



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/06/13
(87) Date publication PCT/PCT Publication Date: 2019/12/26
(85) Entrée phase nationale/National Entry: 2020/12/18
(86) N° demande PCT/PCT Application No.: US 2019/037101
(87) N° publication PCT/PCT Publication No.: 2019/245877
(30) Priorité/Priority: 2018/06/18 (US62/686,274)

(51) Cl.Int./Int.Cl. *C07D 211/06* (2006.01),
A61K 8/49 (2006.01), *A61K 8/67* (2006.01),
A61Q 17/04 (2006.01), *C07D 239/70* (2006.01),
C07D 471/04 (2006.01), *C07D 487/04* (2006.01)
(71) Demandeur/Applicant:
ELC MANAGEMENT LLC, US
(72) Inventeur/Inventor:
BONDA, CRAIG ALAN, US
(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : COMPOSES PHOTOSTABILISANTS, COMPOSITIONS ET PROCÉDES
(54) Title: PHOTOSTABILIZING COMPOUNDS, COMPOSITIONS, AND METHODS




FIG. 1

(57) **Abrégé/Abstract:**

Heterocyclic compounds are provided. In particular, the heteroatom of the heterocyclic compound may be nitrogen. The heterocyclic compounds may demonstrate capacity of stabilizing photoactive compounds. Topical compositions comprising these heterocyclic compounds are also provided. In particular, these topical compositions further comprise photoactive compounds. Methods for stabilizing photoactive compounds are also provided. These methods comprise mixing the photoactive compounds with photostabilizing heterocyclic compounds.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

26 December 2019 (26.12.2019)



(10) International Publication Number

WO 2019/245877 A1

(51) International Patent Classification:

C07D 211/06 (2006.01) *A61K 8/49* (2006.01)*C07D 239/70* (2006.01) *A61K 8/67* (2006.01)*C07D 471/04* (2006.01) *A61Q 17/04* (2006.01)*C07D 487/04* (2006.01)

(21) International Application Number:

PCT/US2019/037101

(22) International Filing Date:

13 June 2019 (13.06.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/686,274 18 June 2018 (18.06.2018) US

(71) Applicant: **ELC MANAGEMENT LLC** [US/US]; 155 Pinelawn Road, Suite 345 South, Melville, New York 11747 (US).

(72) Inventor: **BONDA, Craig Alan**; Os630 Madison Street, Winfield, Illinois 60190 (US).

(74) Agent: **WU, Yonggang** et al.; 155 Pinelawn Road, Suite 345 South, Melville, 11747 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: PHOTOSTABILIZING COMPOUNDS, COMPOSITIONS, AND METHODS




FIG. 1

(57) Abstract: Heterocyclic compounds are provided. In particular, the heteroatom of the heterocyclic compound may be nitrogen. The heterocyclic compounds may demonstrate capacity of stabilizing photoactive compounds. Topical compositions comprising these heterocyclic compounds are also provided. In particular, these topical compositions further comprise photoactive compounds. Methods for stabilizing photoactive compounds are also provided. These methods comprise mixing the photoactive compounds with photostabilizing heterocyclic compounds.

[Continued on next page]

**WO 2019/245877 A1**

WO 2019/245877 A1

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

PHOTOSTABILIZING COMPOUNDS, COMPOSITIONS, AND METHODS

Related Applications

[0001] This application claims priority from provisional patent application serial no. 62/686,274 filed on June 18, 2018.

5

Technical Field

[0002] The disclosure is in the field of compounds that stabilize chemical sunscreens or other compounds that are photoactive, and related compositions and methods.

Background of the Disclosure


[0003] Photoactive compounds are widely used. For example, sunscreens are
10 photoactive compounds. The most widely used UVA and UVB filters in sunscreens are Avobenzene (butyl methoxydibenzoylmethane) and Octinoxate (ethylhexyl methoxycinnamate). While effective in blocking UVA and UVB rays respectively, upon exposure to UV light both Avobenzene and Octinoxate are subject to degradation. Upon exposure to UV light Octinoxate will sometimes form dimers with other Octinoxate
15 molecules. These dimers no longer absorb UVB and UVB efficacy is lost. Octinoxate will also react with the double bond of the dominant form of Avobenzene resulting in the formation of cyclobutane which then forms ring opening structures. The result is loss of UVA efficacy.

[0004] Retinoids are also photoactive compounds. Upon exposure to UV light,
20 retinoids are subject to photoreactions, such as photoisomerization, photopolymerization, photooxidation, and photodegradation. The resulted photodecomposition products do not have the same level of biological activities. The result is loss of biological efficacy.

- [0005] Photostabilizers such as N-cyanodiphenylacrylates such as Octocrylene (2-Cyano-3,3-Diphenyl Acrylic Acid, 2-Ethylhexyl Ester) are known to inhibit the UV-induced photo degradation of Avobenzone. When Avobenzone absorbs a photon of UV light its electron enters a triplet energy state, which can lead to the photo-degradation of the Avobenzone. Octocrylene is able to accept the triplet excited state energy and return the Avobenzone to its original unexcited state. However, when Octoxinate is present, it sometimes will accept the triplet excited state energy from Avobenzone and then react with the double bond found in the dominant form of Avobenzone. Accordingly, Octocrylene is sometimes, but not always, effective for its intended purpose.
- 5 [0006] The problem of solving the instability of photoactive compounds is critical. Sunscreens like Avobenzone and Octinoxate are widely used. Particularly, Avobenzone is one of the only UVA sunscreens approved for global use in sunscreen products. Also, retinoids are highly desired due to their biological benefits and efficacies. Particularly, retinol is an important regulator in epidermal cell growth, normal cell differentiation, and cell maintenance.
- 10 [0007] The disclosure is directed to heterocyclic compounds, compositions comprising those heterocyclic compounds, and related methods for stabilizing photoactive compounds that may include chemical sunscreens, such as Avobenzone or Octinoxate in particular, as well as other unstable compounds such as retinol.
- 15

Summary of the Disclosure

- 20 [0008] The disclosure is directed to heterocyclic compounds having a structure according to Formula I:



Formula I

[0009] In one embodiment, each of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ is independently selected from the group consisting of CR³ and N.

[0010] In one aspect, R³ is selected from the group consisting of H, OH, a straight or
 5 branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkoxy group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, and an aryl group having from about 6 to about 20 carbon atoms. Preferably, R³ is selected from H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an
 10 alkoxy group having from about 1 to about 20 carbon atoms. More preferably, R³ is selected from H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms. Most preferably, R³ is selected from the group consisting of H, and a straight or branched chain alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or
 15 dodecyl.

[0011] In one alternative embodiment, each of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ is independently selected from the group consisting of CH and N.

[0012] In one embodiment, at least one of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ is N. In another embodiment, no more than four of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are N.

20 [0013] In one embodiment, each of R¹ and R² is independently selected from the group consisting of CN, C(=O)OR⁴, C(=O)R⁴, F, CF₃. In one aspect, R¹ and R² are not both CN.

Preferably, each of R¹ and R² is independently selected from the group consisting of CN and C(=O)OR⁴. More preferably, one of R¹ and R² is CN.

[0014] In one aspect, R⁴ is selected from the group consisting of H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, and an aryl group having from about 6 to about 20 carbon atoms. Preferably, R⁴ is a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms. More preferably, R⁴ is a straight or branched chain alkyl group having at least 8, no more than 12 carbon atoms. Most Preferably, R⁴ is a straight or branched chain alkyl group having 8 carbon atoms.

[0015] In one embodiment, examples of the compounds include, but not limit to, Compounds 1-8. Preferably, the compound is Compound 1.

[0016] The disclosure is also directed to compositions comprising at least one heterocyclic compound having a structure according to Formula I.

[0017] In one embodiment, the composition comprises the heterocyclic compounds present in amount ranging from 0.01 to 25% by weight of the total composition. Preferably, the heterocyclic compounds are present in the composition in amount ranging from 0.05 to 15% by weight of the total composition. More preferably, the heterocyclic compounds are present in the composition in amount ranging from 0.1 to 5% by weight of the total composition.

[0018] In one embodiment, the composition further comprises at least one photoactive compound. Preferably, the photoactive compound is selected from the group consisting of a retinoid, a sunscreen, or mixture thereof.

[0019] In one aspect, the photoactive compound is a retinoid. Preferably, the retinoid is retinol.

[0020] In one aspect, the retinoid is present in amount ranging from about 0.0001 to about 20% by weight of the total composition. Preferably, the retinoid is present in amount ranging from about 0.001 to about 10% by weight of the total composition. More preferably, the retinoid is present in amount ranging from about 0.01 to about 8% by weight of the total composition. Most preferably, the retinoid is present in amount ranging from about 0.05 to about 5% by weight of the total composition.

[0021] In one aspect, the photoactive compound is a sunscreen. Preferably, the sunscreen is selected from the group consisting of a UVA chemical sunscreen, a UVB chemical sunscreen, a physical sunscreen, and mixture thereof.

[0022] In one alternative aspect, the sunscreen is a UVA chemical sunscreen. Preferably, the UVA chemical sunscreen is selected from a group consisting of a dibenzoylmethane compound and a dicamphor sulfonic acid derivative. More preferably, the UVA chemical sunscreen is selected from a group consisting of dibenzoylmethane compounds. Examples of the dibenzoylmethane compounds include, but not limit to, 4-methyldibenzoylmethane, 2-methyldibenzoylmethane, 4-isopropyldibenzoylmethane, 4-tert-butyldibenzoylmethane, 2,4-dimethyldibenzoylmethane, 2,5-dimethyldibenzoylmethane, 4,4'-diisopropylbenzoylmethane, 4-tert-butyl-4'-methoxydibenzoylmethane, 4,4'-diisopropylbenzoylmethane, 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane, 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane. Most preferably, the UVA chemical sunscreen is Avobenzene.

[0023] In one alternative aspect, the UVA chemical sunscreen is present in amount ranging from about 0.001 to about 20% by weight of the total composition. Preferably, the UVA chemical sunscreen is present in amount ranging from about 0.005 to about 5% by weight of the total composition. More preferably, UVA chemical sunscreen is present in amount ranging from about 0.005 to about 3% by weight of the total composition.

[0024] In one alternative aspect, the UVA chemical sunscreen is Avobenzene and is present at not greater than about 3% by weight of the total composition.

[0024] In one alternative aspect, the sunscreen is a UVB chemical sunscreen.

Preferably, the UVB chemical sunscreen is selected from the group consisting of an alpha-

5 cyano-beta, beta-diphenyl acrylic acid ester, a benzylidene camphor derivative, a cinnamate derivative, a benzophenone derivative, a menthyl salicylate derivative, an amino benzoic acid derivative, a salicylate derivative, and an ester of 2-phenyl ethanol and benzoic acid. More preferably, the UVB chemical sunscreen is selected from the group consisting of Octocrylene, 4-methylbenzylidene camphor, Octinoxate, Cinoxate, Benzophenone 3, Sulisobenzene, 10 Sulisobenzene Sodium, Homosalate, ethyl hexyl dimethyl PABA, ethyldihydroxypropyl PABA, octyl salicylate, TEA-salicylate, DEA-salicylate, phenylethyl benzoate. More preferably, the UVB chemical sunscreen is selected from the group consisting of Octocrylene, 4-methylbenzylidene camphor, Octinoxate, Benzophenone 3, Homosalate, ethyl hexyl dimethyl PABA, octyl salicylate. Most preferably, the UVB chemical sunscreen is Octinoxate.

15 [0025] In one alternative aspect, the UVB chemical sunscreen is present in amount ranging from about 0.001 to about 45% by weight of the total composition. Preferably, the UVB chemical sunscreen is present in amount ranging from about 0.005 to about 40% by weight of the total composition. More preferably, UVA chemical sunscreen is present in amount ranging from about 0.01 to about 35% by weight of the total composition.

20 [0027] In one aspect, the composition is a sunscreen composition. Preferably, the sunscreen composition has a SPF value ranging from about 1 to about 50. More preferably, the sunscreen composition has a SPF value ranging from about 2 to about 45. Most preferably, the sunscreen composition has a SPF value ranging from about 5 to about 30.

[0026] The disclosure is also directed to methods for stabilizing a photoactive compound, comprising mixing at least one photoactive compound with at least one heterocyclic compound having a structure according to Formula I.

[0027] In one embodiment, the photoactive compound is selected from a group
5 consisting of Avobenzene, Octinoxate, retinol, or mixtures thereof.

Brief Description of The Drawings

[0028] FIG. 1 illustrates an absorption spectrum of Compound 1.

[0029] FIG. 2 illustrates a graph showing inverse fluorescence lifetime vs. Compound
1 concentration used for calculation of the bimolecular quenching rate constant (K_q) for
10 quenching of retinol fluorescence by Compound 1.

[0030] FIG. 3 illustrates a graph showing inverse triplet lifetime vs. Compound 1
concentration used for calculation of the bimolecular quenching rate constant (K_q) for
quenching of *keto*-Avobenzene triplet states by Compound 1.

[0031] FIG. 4 illustrates a graph showing inverse triplet lifetime vs. Compound 1
15 concentration used for calculation of the bimolecular quenching rate constant (K_q) for
quenching of PpIX triplet states by Compound 1.

[0032] FIG. 5 illustrates a graph showing inverse fluorescence lifetime vs. Compound
1 concentration used for calculation of the bimolecular quenching rate constant (K_q) for
quenching of PpIX singlet states by Compound 1.

20 [0033] FIG. 6 illustrates singlet oxygen phosphorescence traces of Pp-MeIX in
absence and presence of various amounts of Compound 1 in air saturated acetonitrile
solutions.

[0034] FIG. 7 illustrates a Stern-Volmer plot of the data illustrated in FIG. 6.

[0035] FIG. 8A illustrates singlet oxygen phosphorescence traces of Pp-MeIX in absence and presence of various amounts of Compound 1 in air saturated acetonitrile solutions.

[0036] FIG. 8B illustrates a Stern-Volmer plot of the data illustrated in FIG. 8A.

5 [0037] FIG. 9A illustrates normalized singlet oxygen phosphorescence traces of Pp-MeIX in air saturated acetonitrile solutions in the absence and presence of Compound 1.

[0038] FIG. 9B illustrates a graph showing inverse singlet oxygen phosphorescence lifetime vs. Compound 1 concentration used for calculation of the bimolecular quenching rate constant (K_q) for singlet oxygen quenching in acetonitrile by Compound 1.

10

Detailed Description

[0039] Photostabilizing compounds are highly desired. In some embodiments, the present disclosure relates to photostabilizing compounds having the capability to stabilize photoactive compounds.

[0040] Each electron in one molecule has two possible spin states. When two electrons
15 of a molecule are at the same molecular orbit and have opposite spin states, these two electrons form an electron pair. When all electrons of a molecule are paired, this molecule is at a singlet state because the electronic energy levels of this molecule would not split when exposed into a magnetic field. When a molecule has only one unpaired electron, this molecule is at a doublet state because the electronic energy levels of this molecule may split into two
20 levels when exposed into a magnetic field. When a molecule has two unpaired electrons whose spin states are parallel to each other, this molecule is at a triplet state because the electronic energy levels of this molecule may split into three levels when exposed into a magnetic field.

[0041] In some embodiments, all electrons of the photoactive compound are paired at the ground state.

[0042] In some embodiments, upon exposure to visible light and/or UV light, the photon absorption of the photoactive compound may cause electron excitation. In some alternative embodiments, upon excitation, an electron of one electron pair may be promoted from the lower energy ground state to a higher energy excited state. The electron pair may be
5 unpaired, with one electron at the excited state and another at the ground state. In one aspect, the excited electron may not change the spin orientation, and may keep the spin orientation opposite to the spin orientation of the other unpaired electron. This excited molecule is at a singlet excited state. In another aspect, the excited electron may change its spin orientation, which may become parallel to the spin orientation of the other unpaired electron. This excited
10 molecule is at a triplet excited state.

[0043] In some embodiments, the photoactive compounds may become less stable upon being excited, subject to photochemical reactions that are mostly irreversible. After undergoing these irreversible reactions, the photoactive compounds generally lose their desired properties and efficacies. Because many photoactive compounds are widely used in the
15 industry due to their great properties and efficacies, it is critical to find a way of stabilizing photoactive compounds.

[0044] In some embodiments, the photostabilizing compounds may stabilize photoactive compounds. In one aspect, the photostabilizing compounds may be capable of directly or indirectly assisting the energy transfer from the excited photoactive compounds. In
20 one alternative aspect, the excited photoactive compounds may be less likely to undergo photochemical reactions as they may more likely get back to their more stable states (i.e., the ground state) before undergoing photochemical reactions due to the co-existing photostabilizing compounds. By lowering the possibility that the photoactive compounds undergo irreversible photochemical reactions after being excited, the photostabilizing
25 compounds may effectively stabilize photoactive compounds.

A. The Compounds

[0045] How photostabilizing compounds assist the energy transfer from the excited photoactive compounds is not well understood.

[0046] In some embodiments, the present disclosure relates to heterocyclic compounds. A heterocyclic compound is one that contains at least a ring made up of more than one kind of atom. Preferably, the heterocyclic compound may be conjugated.

[0047] In one aspect, the heterocyclic compound may be aromatic, non-aromatic, or anti-aromatic. Preferably, the heterocyclic compound may be aromatic.


[0048] In one aspect, the heteroatom of the heterocyclic compound may be nitrogen, oxygen, and/or sulfur. In a heterocyclic compound, a heteroatom is the atom in a ring that is not a carbon atom. Preferably, the heteroatom of the heterocyclic compound may be nitrogen.

[0049] Nitrogen as the heteroatom may affect the properties of heterocyclic compounds in various ways. Nitrogen is more electronegative than carbon is. That is, nitrogen has the higher tendency to attract a bonding pair of electrons than the tendency that carbon has. Also, Nitrogen has a lone pair of electrons that may not form a bond with other atoms.

[0050] In one aspect, the nitrogen's lone pair may be on a p orbital perpendicular to the heterocyclic ring. In this case, nitrogen may act as an electron donor to π orbitals of the heterocyclic system. In another aspect, the nitrogen's lone pair may be on a sp^2 hybrid orbit and lie outside the heterocyclic ring. In this case, nitrogen may act as an electron acceptor of the σ orbitals of the heterocyclic system because it is more electron negative than carbon. The molecular electronic structure of the heterocyclic compound may change dramatically when the number of the nitrogen atom(s) and the position(s) of nitrogen atom(s) change. The photophysical and photochemical properties of the heterocyclic compound may change according to the changes of its molecular electronic structure. By carefully choosing the

number of nitrogen atom(s) on the ring and the position(s) of nitrogen atom(s), desired photophysical and photochemical properties of the compounds may be achieved.

[0051] In some embodiments, the disclosure is related to heterocyclic compounds having the structure according to the Formula I:



Formula I

[0052] In one embodiment, each of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ is independently selected from the group consisting of CR³ and N.

[0053] In one aspect, R³ is selected from the group consisting of:

[0054] (i) H;

10 [0055] (ii) OH;

[0056] (iii) a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, preferably having from about 1 to about 10 carbon atoms, more preferably having from about 1 to about 6 carbon atoms; in one alternative aspect, the alkyl group is a straight or branched chain alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or dodecyl;

[0057] (iv) an alkoxy group having from about 1 to about 20 carbon atoms, preferably having from about 1 to about 12 carbon atoms, more preferably having from about 1 to about 6 carbon atoms, most preferably the alkoxy group is selected from the group consisting of methoxy, ethoxy, propoxy, butoxy;

20

[0058] (v) an alkenyl group having from about 2 to about 20 carbon atoms, preferably having from about 2 to about 12 carbon atoms, more preferably having from about 2 to about 6 carbon atoms, most preferably the alkenyl group is selected from the group consisting of vinyl, allyl, cyclopentenyl, hexenyl;

5 [0059] (vi) an alkynyl group having from about 2 to about 20 carbon atoms, preferably having from about 2 to about 12 carbon atoms, more preferably having from about 2 to about 6 carbon atoms;

[0060] (vii) an aryl group having from about 6 to about 20 carbon atoms, preferably having from about 6 to about 14 carbon atoms, more preferably having from about 6 to about
10 12 carbon atoms.

[0061] In one preferred aspect, R^3 is selected from H; and a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, preferably having from about 1 to about 10 carbon atoms, more preferably having from about 1 to about 6 carbon atoms; in one alternative aspect, the straight or branched chain alkyl is selected from the group consisting of
15 methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or dodecyl.

[0062] In one alternative embodiment, each of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 is independently selected from the group consisting of CH and N.

[0063] In one embodiment, at least one of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 is N.

20 In one embodiment, no more than four of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 are N.

[0064] In one embodiment, each of R^1 and R^2 is independently selected from the group consisting of CN, $C(=O)OR^4$, $C(=O)R^4$, F, CF_3 . In one aspect, R^1 and R^2 are not both CN. Preferably, each of R^1 and R^2 is independently selected from the group consisting of CN and $C(=O)OR^4$. More preferably, one of R^1 and R^2 is CN and another is $C(=O)OR^4$.

25 [0065] In one aspect, R^4 is selected from the group consisting of

[0066] (i) H;

[0067] (ii) a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, preferably having from about 8 to about 12 carbon atoms, more preferably having about 8 carbon atoms; in one alternative aspect, the alkyl group is a straight or

5 branched chain alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or dodecyl, preferably a straight or branched octyl group;

[0068] (iii) an alkenyl group having from about 2 to about 20 carbon atoms, preferably having from about 2 to about 12 carbon atoms, more preferably having from about 2 to about
10 6 carbon atoms, most preferably the alkenyl group is selected from the group consisting of vinyl, allyl, cyclopentenyl, hexenyl;

[0069] (iv) an alkynyl group having from about 2 to about 20 carbon atoms, preferably having from about 2 to about 12 carbon atoms, more preferably having from about 2 to about 6 carbon atoms;


15 [0070] (v) an aryl group having from about 6 to about 20 carbon atoms, preferably having from about 6 to about 14 carbon atoms, more preferably having from about 6 to about 12 carbon atoms.

[0071] In one preferred aspect, R^4 is selected from a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, preferably having from about 1 to about
20 10 carbon atoms; in one alternative aspect, the alkyl group is a straight or branched chain alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or dodecyl, preferably a straight or branched octyl group.


[0072] By carefully selecting R^3 , the photophysical and photochemical properties of
25 the heterocyclic compounds may be further optimized.

[0073] By carefully selecting R^3 and R^4 , the hydrophilicity and/or lipophilicity of the heterocyclic compounds may be optimized. The hydrophilicity and/or lipophilicity of the compounds may play an important role in formulating the compositions comprising the heterocyclic compounds.


5 [0074] Specific, non-limiting examples of heterocyclic compounds are provided:




Compound 1




Compound 2




Compound 3



Compound 4




Compound 5




Compound 6

10



Compound 7



Compound 8

B. The compositions

[0075] In some embodiments, the compositions of the disclosure may be topical

5 compositions. In one aspect, the topical compositions may be in the form of solids, liquids, or gels. In one aspect, the topical compositions may be aqueous based or anhydrous. Aqueous based compositions may be in the form of emulsions, solutions, or dispersions.

[0076] In some embodiments, the compositions comprise at least one compound

10 having the structure according Formulas I. In one aspect, the compound of Formulas I may be present in amounts ranging from about 0.01 to about 25%, preferably about 0.05 to about 15%, more preferably from about 0.1 to about 5% by weight of the total composition.

[0077] In some embodiments, the topical compositions may further comprise certain esters of 2-phenyl ethanol and benzoic acid. One example is phenethyl benzoate, which is sold under the tradename X-Tend 226®, by Ashland.

15 [0078] In some embodiments, the topical compositions may further contain oils, waxes, thickening agents, vitamins, preservatives, antioxidants, botanical extracts, chemical or physical sunscreens or other ingredients.

[0079] In some preferred embodiments, the compositions comprise at least one photoactive compound.


[0080] In one aspect, the photoactive compounds are retinoids and derivatives thereof. Preferably, the compositions comprise retinyl palmitate, retinol, retinoic acid, and/or Vitamin A in the form of beta carotene.

[0081] In one aspect, the retinoid is present in amount ranging from about 0.0001 to about 20% by weight of the total composition. Preferably, the retinoid is present in amount ranging from about 0.001 to about 10% by weight of the total composition. More preferably, the retinoid is present in amount ranging from about 0.01 to about 8% by weight of the total composition. Most preferably, the retinoid is present in amount ranging from about 0.05 to about 5% by weight of the total composition.

[0082] In one aspect, the photoactive compound is sunscreen. Such sunscreens include chemical UVA or UVB sunscreens or physical sunscreens.

1. UVA Chemical Sunscreens

[0083] If desired, the composition may comprise one or more UVA sunscreens. The term "UVA sunscreen" means a chemical compound that blocks UV radiation in the wavelength range of about 320 to 400 nm. Preferred UVA sunscreens are dibenzoylmethane compounds having the general formula:




[0084] wherein R₁ is H, OR and NRR wherein each R is independently H, C₁₋₂₀ straight or branched chain alkyl; R₂ is H or OH; and R₃ is H, C₁₋₂₀ straight or branched chain alkyl.

[0085] Preferred is where R₁ is OR where R is a C₁₋₂₀ straight or branched alkyl, preferably methyl; R₂ is H; and R₃ is a C₁₋₂₀ straight or branched chain alkyl, more preferably, butyl.

[0086] Examples of suitable UVA sunscreen compounds of this general formula

5 include 4-methyldibenzoylmethane, 2-methyldibenzoylmethane, 4-isopropyldibenzoylmethane, 4-tert-butyldibenzoylmethane, 2,4-dimethyldibenzoylmethane, 2,5-dimethyldibenzoylmethane, 4,4'-diisopropylbenzoylmethane, 4-tert-butyl-4'-methoxydibenzoylmethane, 4,4'-diisopropylbenzoylmethane, 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane, 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane, and so on.


10 Particularly preferred is 4-tert-butyl-4'-methoxydibenzoylmethane, also referred to as Avobenzone. Avobenzone is commercially available from Givaudan-Roure or DSM under the trademark Parsol 1789, and Merck & Co. under the tradename Eusolex 9020, and Symrise under the tradename Neo Heliopan 357, and has a structure according to the following formula:



15

[0087] In the preferred embodiment of the disclosure, the composition comprises at least one dibenzoylmethane sunscreen, preferably Avobenzone.


[0088] Other types of UVA sunscreens include dicamphor sulfonic acid derivatives, such as ecamsule, a sunscreen sold by Chimex under the trade name Mexoryl SX, which is
20 terephthalylidene dicamphor sulfonic acid, having the structure according to the following formula:




[0089] The composition may contain from about 0.001-20%, preferably about 0.005-5%, more preferably about 0.005-3% by weight of the composition of UVA sunscreen. In the preferred embodiment of the disclosure the UVA sunscreen is Avobenzone, and it is present at
 5 not greater than about 3% by weight of the total composition.

2. UVB Chemical Sunscreens


[0090] The term "UVB sunscreen" means a compound that blocks UV radiation in the wavelength range of from about 290 to 320 nm. A variety of UVB chemical sunscreens exist including alpha-cyano-beta, beta-diphenyl acrylic acid esters as set forth in U.S. Pat. No.
 10 3,215,724, which is hereby incorporated by reference in its entirety. One particular example of an alpha-cyano-beta, beta-diphenyl acrylic acid ester is Octocrylene, which is 2-ethylhexyl 2-cyano-3,3-diphenylacrylate. In certain cases the composition may contain no more than about 110% by weight of the total composition of octocrylene. Suitable amounts range from about 0.001-10% by weight. Octocrylene may be purchased from BASF under the tradename Uvinul
 15 N-539, from DSM under tradename Parsol 340, and from Symrise under the tradename Neo Heliopan 303, and has a structure according to the following formula:




[0091] Other suitable sunscreens include benzylidene camphor derivatives as set forth in U.S. Pat. No. 3,781,417, which is hereby incorporated by reference in its entirety. Such benzylidene camphor derivatives have the general formula:



5 [0092] wherein R is p-tolyl or styryl, preferably styryl. Particularly preferred is 4-methylbenzylidene camphor, which is a lipid soluble UVB sunscreen compound sold under the tradename Eusolex 6300 by Merck, and Neo Heliopan MBC by Symrise, and Parsol 5000 by DSM, having a structure according to the following formula:




10 [0093] Also suitable are cinnamate derivatives having the general formula:




[0094] wherein R and R₁ are each independently a C₁₋₂₀ straight or branched chain alkyl. Preferred is where R is methyl and R₁ is a branched chain C₁₋₁₀, preferably C₈ alkyl. The preferred compound is ethylhexyl methoxycinnamate, also referred to as Octinoxate or octyl methoxycinnamate. Octinoxate may be purchased from Givaudan Corporation and DSM under

the tradename Parsol MCX, or BASF under the tradename Uvinul MC 80, or Symrise under the tradename Neo Heliopan AV, or Ashland under the tradename Escalol 557, having a structure according to the following structure:




- 5 [0095] Also suitable are mono-, di-, and triethanolamine derivatives of such methoxy cinnamates including diethanolamine methoxycinnamate. Cinoxate, the aromatic ether derivative of the above compound is also acceptable. If present, the Cinoxate should be found at no more than about 3% by weight of the total composition.

- [0096] Also suitable as UVB screening agents are various benzophenone derivatives
10 having the general formula:




- [0097] wherein R through R₉ are each independently H, OH, NaO₃S, SO₃H, SO₃Na, Cl, R", OR" where R" is C₁₋₂₀ straight or branched chain alkyl Examples of such compounds include Benzophenone 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12. Particularly preferred is where
15 the benzophenone derivative is Benzophenone 3 (also referred to as Oxybenzone), Benzophenone 4 (also referred to as Sulisobenzzone), Benzophenone 5 (Sulisobenzzone Sodium), and the like. Most preferred is Benzophenone 3, which may be purchased under the

tradename Uvinul M-40 and NeoHeliopan BB, having the structure according to the following formula:



[0098] Also suitable are certain menthyl salicylate derivatives having the general

5 formula:



[0099] wherein R₁, R₂, R₃, and R₄ are each independently H, OH, NH₂, or C₁₋₂₀

straight or branched chain alkyl. Particularly preferred is where R₁, R₂, and R₃ are methyl and

R₄ is hydroxyl or NH₂, the compound having the name homomenthyl salicylate (also known as


10 Homosalate) or menthyl anthranilate. Menthyl anthranilate is commercially available from

Haarmann & Reimer under the tradename Heliopan. Homosalate is available commercially

from Merck under the tradename Eusolex HMS, and from Symrise under the tradename Neo

Heliopan HMS, and from DSM under the tradename Parsol HMS, having the structure


according to the following formula:



15

[0100] If present, the Homosalate should be present at no more than about 15% by weight of the total composition.


[0101] Various amino benzoic acid derivatives are suitable UVB absorbers including those having the general formula:



5

[0102] wherein R₁, R₂, and R₃ are each independently H, C₁₋₂₀ straight or branched chain alkyl which may be substituted with one or more hydroxy groups. Particularly preferred is wherein R₁ is H or C₁₋₈ straight or branched alkyl, and R₂ and R₃ are H, or C₁₋₈ straight or branched chain alkyl. Particularly preferred are PABA, ethyl hexyl dimethyl PABA (Padimate O), ethyldihydroxypropyl PABA, and the like. If present Padimate O should be found at no more than about 8% by weight of the total composition.

[0103] Salicylate derivatives are also acceptable UVB absorbers. Such compounds have the general formula: wherein R is a straight or branched chain alkyl, including derivatives of the above compound formed from mono-, di-, or triethanolamines. Particular preferred are octyl salicylate, TEA-salicylate, DEA-salicylate, and mixtures thereof. Octyl salicylate has the INCI name Ethylhexyl salicylate, and may be purchased from Ashland under the tradename Escalol 587, and Merck under the tradename Eusolex OS, and has the structure according to the following formula:



[0104] Generally, the amount of the UVB chemical sunscreen present may range from about 0.001-45%, preferably about 0.005-40%, more preferably about 0.01-35% by weight of the total composition.

5 [0105] In one preferred embodiment, the sunscreen may be Avobenzone and/or Octinoxate. It may also be desirable to include one or more other sunscreens in the compositions of the disclosure.

[0106] In one preferred embodiment, the composition may be an oil in water emulsion comprising 5-85% water, 1-40% oil, 0.1-10% Homosalate, 0.1-5% Avobenzone,

10 [0107] If desired, the compositions of the disclosure may be formulated to have a certain SPF (sun protective factor) values ranging from about 1-100, preferably about 4-80, most preferably about 15-60. Calculation of SPF values is well known in the art.

3. Other ingredients:

[0108] The topical composition may contain the following ingredients:


15 Oils

[0109] Suitable oils include silicones, esters, vegetable oils, synthetic oils, including but not limited to those set forth herein. The oils may be volatile or nonvolatile, and are preferably in the form of a pourable liquid at room temperature. If present, the oils may range from about 0.5 to 85%, preferably from about 1-75%, more preferably from about 5-65% by weight of the total composition.

20

[0110] Cyclic and linear volatile silicones are available from various commercial sources including Dow Chemical Corporation and Momentive (formerly General Electric Silicones). The Dow Chemical linear volatile silicones are sold under the trade names Dowsil and Xiameter 244, 245, 344, and 200 fluids. These fluids include hexamethyldisiloxane (viscosity 0.65 centistokes (abbreviated cst)), octamethyltrisiloxane (1.0 cst), decamethyltetrasiloxane (1.5 cst), dodecamethylpentasiloxane (2 cst) and mixtures thereof, with all viscosity measurements being at 25° C.

[0111] Suitable branched volatile silicones include alkyl trimethicones such as methyl trimethicone, a branched volatile silicone having the general formula:



[0112] Methyl trimethicone may be purchased from Shin-Etsu Silicones under the trade name TMF-1.5, having a viscosity of 1.5 centistokes at 25° C.

[0113] Also suitable are various straight or branched chain paraffinic hydrocarbons having 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 carbon atoms, more preferably 8 to 16 carbon atoms. Suitable hydrocarbons include pentane, hexane, heptane, decane, dodecane, tetradecane, tridecane, and C₈₋₂₀ isoparaffins. Suitable C₁₂ isoparaffins are manufactured by Permethyl Corporation under the tradename Permethyl 99A. Various C₁₆ isoparaffins commercially available, such as isohexadecane (having the tradename Permethyl R), are also suitable.

[0114] Also suitable are esters formed by the reaction of a carboxylic acid and an alcohol. The alcohol and the carboxylic acids may both have fatty (C₆₋₃₀) chains. Examples

include hexyl laurate, butyl isostearate, hexadecyl isostearate, cetyl palmitate, isostearyl neopentanoate, stearyl heptanoate, isostearyl isononanoate, stearyl lactate, stearyl octanoate, stearyl stearate, isononyl isononanoate, and so on.

[0115] The ester may also be in the dimer or trimer form. Examples of such esters include diisostearyl malate, neopentyl glycol dioctanoate, dibutyl sebacate, dicetearyl dimer dilinoleate, dicetyl adipate, diisocetyl adipate, diisononyl adipate, diisostearyl dimer dilinoleate, diisostearyl fumarate, diisostearyl malate, dioctyl malate, and so on.

[0116] Examples of other types of esters include those from arachidonic, citric, or behenic acids, such as triarachidin, tributyl citrate, triisostearyl citrate, tri C₁₂₋₁₃ alkyl citrate, tricaprylin, tricaprylyl citrate, tridecyl behenate, trioctyldodecyl citrate, tridecyl behenate; or tridecyl cocoate, tridecyl isononanoate, and so on.

[0117] Synthetic or naturally occurring glyceryl esters of fatty acids, or triglycerides, are also suitable for use in the compositions. Both vegetable and animal sources may be used. Examples of such oils include castor oil, lanolin oil, C₁₀₋₁₈ triglycerides, caprylic/capric/triglycerides, sweet almond oil, apricot kernel oil, sesame oil, camelina sativa oil, tamanu seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, ink oil, olive oil, palm oil, illipe butter, rapeseed oil, soybean oil, grapeseed oil, sunflower seed oil, walnut oil, and the like.

[0118] Also suitable are synthetic or semi-synthetic glyceryl esters, such as fatty acid mono-, di-, and triglycerides which are natural fats or oils that have been modified, for example, mono-, di- or triesters of polyols such as glycerin. In an example, a fatty (C₁₂₋₂₂) carboxylic acid is reacted with one or more repeating glyceryl groups. glyceryl stearate, diglyceryl diisostearate, polyglyceryl-3 isostearate, polyglyceryl-4 isostearate, polyglyceryl-6 ricinoleate, glyceryl dioleate, glyceryl diisostearate, glyceryl tetraisostearate, glyceryl trioctanoate, diglyceryl distearate, glyceryl linoleate, glyceryl myristate, glyceryl isostearate,

PEG castor oils, PEG glyceryl oleates, PEG glyceryl stearates, PEG glyceryl tallowates, and so on.

[0119] Nonvolatile silicone oils, both water soluble and water insoluble, are also suitable for use in the composition. Such silicones preferably have a viscosity ranging from about greater than 5 to 800,000 cst, preferably 20 to 200,000 cst at 25° C. Suitable water insoluble silicones include amine functional silicones such as amodimethicone. Examples include dimethicone, phenyl dimethicone, diphenyl dimethicone, phenyl trimethicone, or trimethylsiloxyphenyl dimethicone. Other examples include alkyl dimethicones such as cetyl dimethicone, stearyl dimethicone, behenyl dimethicone, and the like.

10 Surfactants

[0120] The composition may contain one or more surfactants, especially if in the emulsion form. However, such surfactants may be used if the compositions are anhydrous also, and will assist in dispersing ingredients that have polarity, for example pigments. Such surfactants may be silicone or organic based. The surfactants will aid in the formation of stable emulsions of either the water-in-oil or oil-in-water form. If present, the surfactant may range from about 0.001 to 30%, preferably from about 0.005 to 25%, more preferably from about 0.1 to 20% by weight of the total composition.

[0121] Silicone surfactants may be generically referred to as dimethicone copolyol or alkyl dimethicone copolyol. In some cases the number of repeating ethylene oxide or propylene oxide units in the polymer are also specified, such as a dimethicone copolyol that is also referred to as PEG-15/PPG-10 dimethicone, which refers to a dimethicone having substituents containing 15 ethylene glycol units and 10 propylene glycol units on the siloxane backbone. It is also possible for one or more of the methyl groups in the above general

structure to be substituted with a longer chain alkyl (e.g. ethyl, propyl, butyl, etc.) or an ether such as methyl ether, ethyl ether, propyl ether, butyl ether, and the like.

[0122] Examples of silicone surfactants are those sold by Dow Silicones under the tradename Dowsil 3225C Formulation Aid having the CTFA name cyclotetrasiloxane (and) cyclopentasiloxane (and) PEG/PPG-18 dimethicone; or 5225C Formulation Aid, having the CTFA name cyclopentasiloxane (and) PEG/PPG-18/18 dimethicone; or Dowsil 190 Surfactant having the CTFA name PEG/PPG-18/18 dimethicone; or Dowsil 193 Fluid, Dowsil 5200 having the CTFA name lauryl PEG/PPG-18/18 methicone; or Abil EM 90 having the CTFA name cetyl PEG/PPG-14/14 dimethicone sold by Goldschmidt; or Abil EM 97 having the CTFA name bis-cetyl PEG/PPG-14/14 dimethicone sold by Goldschmidt; or Abil WE 09 having the CTFA name cetyl PEG/PPG-10/1 dimethicone in a mixture also containing polyglyceryl-4 isostearate and hexyl laurate; or KF-6011 sold by Shin-Etsu Silicones having the CTFA name PEG-11 methyl ether dimethicone; KF-6012 sold by Shin-Etsu Silicones having the CTFA name PEG/PPG-20/22 butyl ether dimethicone; or KF-6013 sold by Shin-Etsu Silicones having the CTFA name PEG-9 dimethicone; or KF-6015 sold by Shin-Etsu Silicones having the CTFA name PEG-3 dimethicone; or KF-6016 sold by Shin-Etsu Silicones having the CTFA name PEG-9 methyl ether dimethicone; or KF-6017 sold by Shin-Etsu Silicones having the CTFA name PEG-10 dimethicone; or KF-6038 sold by Shin-Etsu Silicones having the CTFA name lauryl PEG-9 polydimethylsiloxylethyl dimethicone.

[0123] Also suitable are various types of crosslinked silicone surfactants that are often referred to as emulsifying elastomers that contain at least one hydrophilic moiety such as polyoxyalkylenated groups. Polyoxyalkylenated silicone elastomers that may be used in at least one embodiment of the disclosure include those sold by Shin-Etsu Silicones under the names KSG-21, KSG-20, KSG-30, KSG-31, KSG-32, KSG-33; KSG-210 which is dimethicone/PEG-10/15 crosspolymer dispersed in dimethicone; KSG-310 which is PEG-15

lauryl dimethicone crosspolymer; KSG-320 which is PEG-15 lauryl dimethicone crosspolymer dispersed in isododecane; KSG-330 (the former dispersed in triethylhexanoin), KSG-340 which is a mixture of PEG-10 lauryl dimethicone crosspolymer and PEG-15 lauryl dimethicone crosspolymer.

5 [0124] Also suitable are polyglycerolated silicone elastomers like those disclosed in PCT/WO 2004/024798, which is hereby incorporated by reference in its entirety. Such elastomers include Shin-Etsu's KSG series, such as KSG-710 which is dimethicone/polyglycerin-3 crosspolymer dispersed in dimethicone; or lauryl dimethicone/polyglycerin-3 crosspolymer dispersed in a variety of solvent such as
10 isododecane, dimethicone, triethylhexanoin, sold under the Shin-Etsu tradenames KSG-810, KSG-820, KSG-830, or KSG-840. Also suitable are silicones sold by Dow Silicones under the tradenames 9010 and DC9011.

[0125] The composition may comprise one or more nonionic organic surfactants. Suitable nonionic surfactants include alkoxylated alcohols, or ethers, formed by the reaction of
15 an alcohol with an alkylene oxide, usually ethylene or propylene oxide. Preferably the alcohol is either a fatty alcohol having 6 to 30 carbon atoms. Examples of such ingredients include Steareth 2-100, which is formed by the reaction of stearyl alcohol and ethylene oxide and the number of ethylene oxide units ranges from 2 to 100; Beheneth 5-30 which is formed by the reaction of behenyl alcohol and ethylene oxide where the number of repeating ethylene oxide
20 units is 5 to 30; Cetareth 2-100, formed by the reaction of a mixture of cetyl and stearyl alcohol with ethylene oxide, where the number of repeating ethylene oxide units in the molecule is 2 to 100; Ceteth 1-45 which is formed by the reaction of cetyl alcohol and ethylene oxide, and the number of repeating ethylene oxide units is 1 to 45, and so on. All recitations of units include all whole integers between the range.

[0126] Other alkoxyated alcohols are formed by the reaction of fatty acids and mono-, di- or polyhydric alcohols with an alkylene oxide. For example, the reaction products of C₆₋₃₀ fatty carboxylic acids and polyhydric alcohols which are monosaccharides such as glucose, galactose, methyl glucose, and the like, with an alkoxyated alcohol. Examples include

5 polymeric alkylene glycols reacted with glyceryl fatty acid esters such as PEG glyceryl oleates, PEG glyceryl stearate; or PEG polyhydroxyalkanotes such as PEG dipolyhydroxystearate wherein the number of repeating ethylene glycol units ranges from 3 to 1000.

[0127] Other suitable nonionic surfactants include alkoxyated sorbitan and

10 alkoxyated sorbitan derivatives. For example, alkoxylation, in particular ethoxylation of sorbitan provides polyalkoxyated sorbitan derivatives. Esterification of polyalkoxyated sorbitan provides sorbitan esters such as the polysorbates. For example, the polyalkoxyated sorbitan can be esterified with C₆₋₃₀, preferably C₁₂₋₂₂ fatty acids. Examples of such ingredients include Polysorbates 20-85, sorbitan oleate, sorbitan sesquioleate, sorbitan

15 palmitate, sorbitan sesquiisostearate, sorbitan stearate, and so on.

Humectants

[0128] It may also be desirable to include one or more humectants in the composition. If present, such humectants may range from about 0.001 to 25%, preferably from about 0.005 to 20%, more preferably from about 0.1 to 15% by weight of the total composition. Examples

20 of suitable humectants include glycols, sugars, and the like. Suitable glycols are in monomeric or polymeric form and include polyethylene and polypropylene glycols such as PEG 4-200, which are polyethylene glycols having from 4 to 200 repeating ethylene oxide units; as well as C₁₋₆ alkylene glycols such as propylene glycol, butylene glycol, pentylene glycol, and the like. Suitable sugars, some of which are also polyhydric alcohols, are also suitable humectants.

Examples of such sugars include glucose, fructose, honey, hydrogenated honey, inositol, maltose, mannitol, maltitol, sorbitol, sucrose, xylitol, xylose, and so on. Also suitable is urea. Preferably, the humectants used in the composition of the disclosure are C₁₋₆, preferably C₂₋₄ alkylene glycols, most particularly butylene glycol.

5 Botanical Extracts

[0129] It may be desirable to include one or more botanical extracts in the compositions. If so, suggested ranges are from about 0.0001 to 10%, preferably about 0.0005 to 8%, more preferably about 0.001 to 5% by weight of the total composition. Suitable botanical extracts include extracts from plants (herbs, roots, flowers, fruits, seeds) such as

10 flowers, fruits, vegetables, and so on, including yeast ferment extract, *Padina Pavonica* extract, *thermus thermophilis* ferment extract, camelina sativa seed oil, boswellia serrata extract, olive extract, *Aribodopsis Thaliana* extract, *Acacia Dealbata* extract, *Acer Saccharinum* (sugar maple), acidopholus, acorus, aesculus, agaricus, agave, agrimonia, algae, aloe, citrus, brassica, cinnamon, orange, apple, blueberry, cranberry, peach, pear, lemon, lime,

15 pea, seaweed, caffeine, green tea, chamomile, willowbark, mulberry, poppy, and those set forth on pages 1646 through 1660 of the CTFA Cosmetic Ingredient Handbook, Eighth Edition, Volume 2. Further specific examples include, but are not limited to, *Glycyrrhiza Glabra*, *Salix Nigra*, *Macrocystis Pyrifer*, *Pyrus Malus*, *Saxifraga Sarmentosa*, *Vitis Vinifera*, *Morus Nigra*, *Scutellaria Baicalensis*, *Anthemis Nobilis*, *Salvia Sclarea*, *Rosmarinus*

20 *Officinalis*, *Citrus Medica Limonum*, *Panax Ginseng*, *Siegesbeckia Orientalis*, *Fructus Mume*, *Ascophyllum Nodosum*, Bifida Ferment lysate, *Glycine Soja* extract, *Beta Vulgaris*, *Haberlea Rhodopensis*, *Polygonum Cuspidatum*, *Citrus Aurantium Dulcis*, *Vitis Vinifera*, *Selaginella Tamariscina*, *Humulus Lupulus*, *Citrus Reticulata Peel*, *Punica Granatum*,

Asparagopsis, Curcuma Longa, Menyanthes Trifoliata, Helianthus Annuus, Hordeum Vulgare, Cucumis Sativus, Evernia Prunastri, Evernia Furfuracea, and mixtures thereof.

Particulate Materials

- [0130] The compositions of the disclosure may contain particulate materials in the form of pigments, inert particulates, or mixtures thereof. If present, suggested ranges are from about 0.01-75%, preferably about 0.5-70%, more preferably about 0.1-65% by weight of the total composition. In the case where the composition may comprise mixtures of pigments and powders, suitable ranges include about 0.01-75% pigment and 0.1-75% powder, such weights by weight of the total composition.
- [0131] The particulate matter may be colored or non-colored powders. Suitable non-pigmented powders include bismuth oxychloride, titanated mica, fumed silica, spherical silica, polymethylmethacrylate, micronized teflon, boron nitride, acrylate copolymers, aluminum silicate, aluminum starch octenylsuccinate, bentonite, calcium silicate, cellulose, chalk, corn starch, diatomaceous earth, fuller's earth, glyceryl starch, hectorite, hydrated silica, kaolin, magnesium aluminum silicate, magnesium trisilicate, maltodextrin, montmorillonite, microcrystalline cellulose, rice starch, silica, talc, mica, titanium dioxide, zinc laurate, zinc myristate, zinc rosinate, alumina, attapulgite, calcium carbonate, calcium silicate, dextran, kaolin, nylon, silica silylate, silk powder, sericite, soy flour, tin oxide, titanium hydroxide, trimagnesium phosphate, walnut shell powder, or mixtures thereof. The above mentioned powders may be surface treated with lecithin, amino acids, mineral oil, silicone, or various other agents either alone or in combination, which coat the powder surface and render the particles more lipophilic in nature.

[0132] Suitable pigments are organic or inorganic. Organic pigments are generally various aromatic types including azo, indigoid, triphenylmethane, anthroquinone, and xanthine

dyes which are designated as D&C and FD&C blues, browns, greens, oranges, reds, yellows, etc. Organic pigments generally consist of insoluble metallic salts of certified color additives, referred to as the Lakes. Inorganic pigments include iron oxides, ultramarines, chromium, chromium hydroxide colors, and mixtures thereof. Iron oxides of red, blue, yellow, brown,
5 black, and mixtures thereof are suitable.

Vitamins and Antioxidants

[0133] The compositions of the disclosure may contain vitamins and/or coenzymes, as well as antioxidants. If so, 0.001-10%, preferably 0.01-8%, more preferably 0.05-5% by weight of the total composition is suggested. Suitable vitamins include ascorbic acid and
10 derivatives thereof such as ascorbyl palmitate, tetrahexydecyl ascorbate, and so on; the B vitamins such as thiamine, riboflavin, pyridoxin, and so on, as well as coenzymes such as thiamine pyrophosphate, flavin adenin dinucleotide, folic acid, pyridoxal phosphate, tetrahydrofolic acid, and so on. Also suitable is Vitamin E and derivatives thereof such as Vitamin E acetate, nicotinate, or other esters thereof. In addition, Vitamins D and K are
15 suitable.

C. The methods

[0134] In some embodiments, the disclosure is related to methods for stabilizing photoactive compounds, the methods comprise mixing a least one photoactive compound with at least one heterocyclic compound having the structure according to Formula I.

20 [0135] In one aspect, the methods for stabilizing retinoids and derivatives thereof comprise mixing a least one retinoid and/or derivatives thereof with at least one heterocyclic compound having the structure according to Formula I.

[0136] In one aspect, the methods for stabilizing chemical sunscreens comprise mixing a least one chemical sunscreen with at least one heterocyclic compound having the structure according to Formula I.

[0137] The disclosure will be further described in connection with the following


5 examples which are set forth for the purposes of illustration only.

EXPERIMENTAL

Compound Examples

Example 1

Synthesis of Compound 1




10

[0138] 1-Azafluorenone (4.0 gm, 22.1 mmol) and toluene (40 mL) were mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 2.0 mmol), and acetic acid (2.8 mL) were added sequentially at 25-30°C. The reaction mixture was

15 refluxed for approximately 18 hours at 100-115°C. The water was periodically removed from Dean-Stark condenser during the reaction. The reaction was monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 1-azafluorenone by TLC, the reaction mixture was cooled to room temperature. The toluene layer was washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25 mL). The


20 organic layer was evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product was purified by column chromatography by eluting with 10-15% ethyl acetate in hexane to afford pure product (3.5 gm, yield of 44%).

Example 2
Synthesis of Compound 2



[0139] 2,4-Diazafluorenone (4.0 gm, 22.1 mmol) and toluene (40mL) are mixed in a
 5 clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen
 inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99
 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is
 refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from
 Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl
 10 acetate/hexane). After the complete consumption of 2,4-diazafluorenone by TLC, the reaction
 mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL)
 followed by saturated sodium bicarbonate solution (25 mL) and again with water (25mL). The
 organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale
 brown semisolid. The crude product is purified by column chromatography by eluting with 10-
 15 15% ethyl acetate in hexane to afford pure product.


Example 3
Synthesis of Compound 3



[0140] 3,5-Diazafluorenone (4.0 gm, 22.1 mmol) and toluene (40mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 3,5-diazafluorenone by TLC, the reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product is purified by column chromatography by eluting with 10-15% ethyl acetate in hexane to afford pure product.

Example 4


Synthesis of Compound 4



[0141] 2,4,5,7-Tetraazafluorenone (4.0 gm, 22.1 mmol) and toluene (40mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 2,4,5,7-tetraazafluorenone by TLC, the

reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product is purified by column chromatography by eluting with ethyl acetate in hexane to afford pure product.


Example 5
Synthesis of Compound 5



[0142] 4,5-Diazafluorenone (4.0 gm, 22.1 mmol) and toluene (40mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 4,5-diazafluorenone by TLC, the reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product is purified by column chromatography by eluting with 10-15% ethyl acetate in hexane to afford pure product.

Example 6


Synthesis of Compound 6



- 5 [0143] 2,5-Diazafluorenone (4.0 gm, 22.1 mmol) and toluene (40 mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from
- 10 Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 2,5-diazafluorenone by TLC, the reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale
- 15 brown semisolid. The crude product is purified by column chromatography by eluting with 10-15% ethyl acetate in hexane to afford pure product.

Example 7


Synthesis of Compound 7



[0144] 4-Methyl-1-azafluorenone (4.0 gm, 22.1 mmol) and toluene (40 mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexyl cyanoacetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 4-methyl-1-azafluorenone by TLC, the reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25 mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product is purified by column chromatography by eluting with 10-15% ethyl acetate in hexane to afford pure product.

Example 8

Synthesis of Compound 8



15

[0145] 3-Methyl-2-azafluorenone (4.0 gm, 22.1 mmol) and toluene (40mL) are mixed in a clean 250 mL 2 neck round bottom flask equipped with Dean-Stark condenser and nitrogen inlet. 2-ethylhexylcyano acetate (4.4 gm, 22.3 mmol), ammonium acetate (153 mg, 1.99 mmol), and acetic acid (2.8 mL) are added sequentially at 25-30°C. The reaction mixture is refluxed for approximately 18 hours at 100-115°C. The water is periodically removed from Dean-Stark condenser during the reaction. The reaction is monitored by TLC (30% ethyl acetate/hexane). After the complete consumption of 3-methyl-2-azafluorenone by TLC, the

20

reaction mixture is cooled to room temperature. The toluene layer is washed with water (2×25 mL) followed by saturated sodium bicarbonate solution (25 mL) and again with water (25 mL). The organic layer is evaporated under reduced pressure at 45-50°C to obtain crude product as pale brown semisolid. The crude product is purified by column chromatography by
5 eluting with 10-15% ethyl acetate in hexane to afford pure product.

Photophysical and Photochemical Measurements Examples

Example 9

Absorption Spectra of Compound 1

[0146] The acetonitrile solution of Compound 1 was measured by an Agilent 8453
10 spectrometer to record the UV and visible absorption spectra of Compound 1, which is illustrated in FIG. 1.

Example 10


Fluorescence quenching of retinol by Compound 1

[0147] The THF solutions of retinol at 0.2 mM and Compound 1 at various
15 concentration were measured on OB920 TR-SPC (Edinburgh Analytical Instruments) by time-correlated single photon counting (TCSPC) using a pulsed LED (335 nm, 5 MHz) for excitation (PicoQuant) for the corresponding fluorescence lifetimes. The fluorescence signal was collected at 483 nm. The bimolecular quenching rate constant K_q of quenching of retinol fluorescence by Compound 1 was calculated from the slope of the plot in FIG. 2 based on the
20 equation $1/\tau_f = k_0 + k_q [\text{Compound 1}]$ to be $(9.0 \pm 0.3) \times 10^9 \text{ M}^{-1}\text{s}^{-1}$, which showed significant photostablizing capacity of Compound 1.

Example 11

Keto-Avobenzene triplet state quenching by Compound 1

[0148] The deoxygenated acetonitrile solutions of avobenzene at 0.25 mM and Compound 1 at various concentrations were measured for transient absorption on a homebuilt system (Y. Yagci, S. Jockusch and N. J. Turro, *Macromolecules*, 2007, 40, 4481–4485; hereby incorporated by reference in its entirety). *Keto*-Avobenzene was generated by photolysis of *enol*-Avobenzene at 350 nm. Laser flash photolysis of *keto*-Avobenzene was conducted at 266 nm with 5 ns pulse width as shown below:



[0149] The decay traces of the triplet-absorption of *keto*-Avobenzene was monitored at 380 nm. The quenching rate constant was calculated from the slope of the plot in FIG. 3 to be $k_q = (3.5 \pm 0.2) \times 10^9 \text{ M}^{-1}\text{s}^{-1}$, which showed significant photostablizing capacity of Compound 1.

Example 12

PpIX triplet state quenching by Compound 1

[0150] The Argon saturated acetonitrile solutions of Protoporphyrin IX (PpIX) at a concentration optimal for light absorption and Compound 1 at various concentrations were measured for transient absorption on a homebuilt system (Y. Yagci, S. Jockusch and N. J. Turro, *Macromolecules*, 2007, 40, 4481–4485; hereby incorporated by reference in its entirety). Triplet lifetimes were measured at 532 nm (7 ns pulse length) and the transient absorbance was monitored at 630 nm. The quenching rate constant k_q was calculated from the

slope of the plot in FIG. 4 to be $2.7 \pm 0.1 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$ based on the equation $1/\tau_f = k_0 + k_q$ [Compound 1], which showed significant photostabilizing capacity of Compound 1.

Example 13

PpIX singlet excited state quenching by Compound 1

5 [0151] The acetonitrile solutions of PpIX at a concentration optimal for light absorption and Compound 1 at various concentration were measured on OB920 TR-SPC (Edinburgh Analytical Instruments) by time-correlated single photon counting (TCSPC) using a pulsed LED (496 nm) for excitation (PicoQuant) for the corresponding fluorescence lifetimes. The fluorescence signal was collected at 630 nm. The bimolecular quenching rate
10 constant K_q of quenching of PpIX fluorescence by Compound 1 was calculated from the slope of the plot in FIG. 5 based on the equation $1/\tau_f = k_0 + k_q [\text{Compound 1}]$ to be $2.5 \pm 0.1 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$, which showed significant photostabilizing capacity of Compound 1.

Example 14

Singlet oxygen generation quenching by Compound 1 - K_{SV} in acetonitrile

15 [0152] Singlet oxygen phosphorescence measurements of the air saturated acetonitrile solutions of the dimethyl ester derivative of PpIX (Pp-MeIX) at the concentration optimal for light absorption and Compound 1 at various concentrations were performed on a modified Fluorolog-3 spectrometer (HORIBA Jobin Yvon) in conjunction with a NIR sensitive photomultiplier tube (H102330A-45, Hamamatsu). A Spectra Physics GCR-150-30 Nd:YAG
20 laser (532 nm, ca. 5 mJ per pulse, 7 ns) was used for pulsed excitation to collect $^1\text{O}_2$ phosphorescence decay traces at 1270 nm which were stored on a digital oscilloscope (TDS 360 from Tektronics). The singlet oxygen phosphorescence traces were shown in FIG. 6. The Stern-Volmer plots of singlet oxygen phosphorescence were illustrated in FIG. 7. The Stern-

Volmer constant K_{SV} was calculated from the slope of the plot in FIG. 7 to be $57 \pm 2 \text{ M}^{-1}$, which is a direct measurement of the singlet oxygen suppression efficiency of Compound 1.

Compound 1 thus showed significant photostabilizing capacity.

[0153] The structure of Pp-MeIX was disclosed previously (S. Jockusch, C. Bonda and
5 S. Hu, Photochem. PhotoBio. Sci, 2014, 13, 1180–1184; hereby incorporated by reference in its entirety).

Example 15

Singlet oxygen generation quenching by Compound 1 - K_{SV} in CDCl_3

[0154] Singlet oxygen phosphorescence measurements of the air saturated CDCl_3
10 solutions of the dimethyl ester derivative of PpIX (Pp-MeIX) at the concentration optimal for light absorption and Compound 1 at various concentrations were performed on a modified Fluorolog-3 spectrometer (HORIBA Jobin Yvon) in conjunction with a NIR sensitive photomultiplier tube (H102330A-45, Hamamatsu). A Spectra Physics GCR-150-30 Nd:YAG laser (532 nm, ca. 5 mJ per pulse, 7 ns) was used for pulsed excitation to collect $^1\text{O}_2$
15 phosphorescence decay traces at 1270 nm which were stored on a digital oscilloscope (TDS 360 from Tektronics). The singlet oxygen phosphorescence traces were shown in FIG. 8A. The Stern-Volmer plots of singlet oxygen phosphorescence were illustrated in FIG. 8B. The Stern-Volmer constant K_{SV} was calculated from the slope of the plot in FIG. 8B to be $14 \pm 1 \text{ M}^{-1}$, which is a direct measurement of the singlet oxygen suppression efficiency of Compound 1.
20 Compound 1 thus showed significant photostabilizing capacity.

Example 16

Singlet oxygen generation quenching by Compound 1 – k_q in acetonitrile

[0155] Singlet oxygen phosphorescence measurements of the air saturated acetonitrile solutions of the dimethyl ester derivative of PpIX (Pp-MeIX) at the concentration optimal for light absorption and Compound 1 at various concentrations were performed on a modified Fluorolog-3 spectrometer (HORIBA Jobin Yvon) in conjunction with a NIR sensitive photomultiplier tube (H102330A-45, Hamamatsu). A Spectra Physics GCR-150-30 Nd:YAG laser (532 nm, ca. 5 mJ per pulse, 7 ns) was used for pulsed excitation to collect $^1\text{O}_2$ phosphorescence decay traces at 1270 nm which were stored on a digital oscilloscope (TDS 360 from Tektronics). The singlet oxygen phosphorescence traces of absence and presence of Compound 1 (40 mM) were shown in FIG. 9A. The plot of the inverse singlet oxygen phosphorescence lifetime vs. Compound 1 concentration was shown in FIG. 9B. The bimolecular quenching rate constant k_q of singlet oxygen quenching in acetonitrile by Compound 1 was calculated from the slope of the plot in FIG. 9B to be $1.5 \pm 0.1 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, which showed significant photostabilizing capacity of Compound 1.

Composition Examples

Example 17

[0156] An anti-aging cream was prepared as follows:

Ingredient	Wt%
Water	QS100
Shea butter	6.00
Caprylic/capric/myristic/stearic triglyceride	5.50
Methyl trimethicone	5.00
Di-C12-15 alkyl fumarate	4.00
Dimethicone/polysilicone-11	4.00

Butylene glycol	3.00
Steareth-2	2.30
Glyceryl stearate	1.50
Pentylene glycol	1.50
Stearyl alcohol	1.50
Steareth-21	1.20
Glycerin	1.00
Phenoxyethanol	0.50
Acrylamide/sodium acryloyldimethyltaurate copolymer/water/isohexadecane/polysorbate 80	0.50
Fragrance	0.40
Carbomer	0.35
Water/sodium hydroxide	0.28
Cholesterol	0.20
Linoleic acid	0.20
Caffeine	0.20
Dimethicone	0.20
Sodium dehydroacetate	0.10
Tocopherol acetate	0.10
Compound 1	0.50

[0157] The composition was prepared by separately mixing the oil phase ingredients including the Compound 1. The water phase ingredients were combined and emulsified with the oil phase ingredients to form an emulsion.

Example 18

[0158] A formula with stabilized retinol was prepared as follows:

Ingredient	Wt%
Water	QS100
Butylene glycol	1.60
Sodium bisulfite	0.02
Caffeine	0.20
Silica	0.20
Caprylic/capric triglyceride	3.17
Dimethicone	3.00
Cetearyl alcohol	2.00
Tocopheryl acetate	0.50
Tocopherol	0.20
Disodium EDTA	0.10
Sodium hyaluronate	0.10
Cholesterol	0.20
Arachidyl alcohol	1.37
Polysorbate 60	0.03
Behenyl alcohol	0.75
Sodium hydroxide	0.07
Hydroxyethylcellulose	0.30
Retinol	0.30
Stearyl dimethicone	2.25

Caprylyl glycol	0.32
Glycerin	2.50
Shea butter	2.80
Sorbitan olivate	0.80
Sorbitan isostearate	0.03
Cetearyl olivate	1.20
Arachidyl glucoside	0.38
PEG-12 dimethicone/PPG-20 crosspolymer	1.60
Compound 1	0.15

[0159] The composition was prepared by separately combining the water phase and oil phase with the Compound 1. The phases were mixed to emulsify and form a lotion.

Example 19

5 [0160] A sunscreen composition containing Compound 1 was prepared as follows:

Phase	Ingredient	Wt%
A	Deionized water	QS100
B	Glycerin	2.50
B	Triethanolamine	0.60
B	Disodium EDTA	0.10
C	Compound 1	2.00
C	Dimethicone (2 cs)	2.00
C	Glyceryl stearate/PEG 100 stearate	2.50
C	Beeswax	1.00

D	Avobenzone	3.00
D	Homosalate	10.00
D	Octisalate	5.00
D	Octocrylene	4.00
F	Dimethicone/Acrylates dimethicone copolymer	2.50
F	Trimethylsiloxysilicate/dimethicone	2.50
G	Ammonium/Acryloyldimethyltaurate copolymer	0.50
H	Caprylyl glycol/phenoxyethanol/hexylene glycol	1.00

[0161] The Phase A ingredients were charged into a main kettle. Phase B ingredients were added and propeller mixed at medium/high speed until homogeneous. The batch was then heated to a temperature of 67-70° C. In an auxiliary kettle the Phase C ingredients were heated to 65-70° C. and mixed with a propeller at medium speed. The Phase D ingredients were added and mixing at medium speed continued until uniform. The heat was lowered to 63° C. and the Phase E ingredients were added into the vortex with propeller mixing until dispersed. Phases C, D, and E were added into the main batch (A+B) while mixing at high speed. The composition was homogenized at 2000 rpm for 15-20 minutes. When the batch was emulsified and homogeneous, propeller mixing was continued and pre-mixed Phase F ingredients were added and mixed until uniform while cooling the batch to room temperature. Phases G and H were then added and mixed until uniform. The batch was cooled to room temperature.


Other Embodiments

[0162] While the invention has been described in connection with the preferred embodiment, it is not intended to limit the scope of the invention to the particular form set

forth but, on the contrary, it is intended to cover such alternatives, modifications, and equivalents as may be included within the spirit and scope of the invention as defined by the appended claims. Therefore, other embodiments, including those can be easily modified by a person skilled in the art from the present disclosure, are also within the claims.

CLAIMS:

1. A heterocyclic compound having a structure according to Formula I:




Formula I


- wherein each of A1, A2, A3, A4, A5, A6, A7, and A8 is independently selected from the
- 5 group consisting of CR³ and N;
- wherein R³ is selected from the group consisting of H, OH, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkoxy group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, and an aryl group having from
- 10 about 6 to about 20 carbon atoms;
- wherein at least one of A1, A2, A3, A4, A5, A6, A7, and A8 is N;
- wherein no more than four of A1, A2, A3, A4, A5, A6, A7, and A8 are N;
- wherein each of R¹ and R² is independently selected from the group consisting of CN, C(=O)OR⁴, C(=O)R⁴, F, CF₃, with the proviso that R¹ and R² are not both CN;
- 15 wherein R⁴ is selected from the group consisting of H, an alkyl group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, and an aryl group having from about 6 to about 20 carbon atoms.

2. The heterocyclic compound of Claim 1, wherein R^3 is selected from the group consisting of H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, and an alkoxy group having from about 1 to about 20 carbon atoms.
- 5 3. The heterocyclic compound of Claim 2, wherein R^3 is selected from the group consisting of H, and a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms.
4. The heterocyclic compound of Claim 1, wherein R^3 is selected from the group
10 consisting of H, and a straight or branched chain alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, 2-methyl-1-propyl, 2-methyl-2-propyl, pentyl, 2-methyl-2-butyl, hexyl, heptyl, octyl, decyl, or dodecyl.
5. The heterocyclic compound of Claim 1, wherein each of R^1 and R^2 is independently
15 selected from the group consisting of CN and $C(=O)OR^4$.
6. The heterocyclic compound of Claim 5, wherein one of R^1 and R^2 is CN.
7. The heterocyclic compound of Claim 5, wherein R^4 is a straight or branched chain
20 alkyl group having from about 1 to about 20 carbon atoms.
8. The heterocyclic compound of Claim 5, wherein R^4 is a straight or branched chain alkyl group having at least 8, no more than 12 carbon atoms.
- 25 9. The heterocyclic compound of Claim 5, wherein R^4 is a straight or branched chain alkyl group having 8 carbon atoms.


10. The heterocyclic compound of Claim 1, wherein the compound is selected from the group consisting of:




Compound 1




Compound 2




Compound 3




Compound 4




Compound 5



Compound 6




Compound 7




Compound 8

11. The heterocyclic compound of Claim 10, wherein the compound having the structure according to formula:



Compound 1

5 12. A composition comprising at least one heterocyclic compound having the structure according to Formula I:



Formula I

Wherein each of A1, A2, A3, A4, A5, A6, A7, and A8 is independently selected from the
10 group consisting of CR³ and N;

wherein R³ is selected from the group consisting of H, OH, a straight or branched chain alkyl
group having from about 1 to about 20 carbon atoms, an alkoxy group having from about 1 to
about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an
alkynyl group having from about 2 to about 20 carbon atoms, an aryl group having from about
15 6 to about 20 carbon atoms;

wherein at least one of A1, A2, A3, A4, A5, A6, A7, and A8 is N;

wherein no more than four of A1, A2, A3, A4, A5, A6, A7, and A8 are N;

wherein each of R^1 and R^2 is independently selected from the group consisting of CN, $C(=O)OR^4$, $C(=O)R^4$, F, CF_3 , with the proviso that R^1 and R^2 are not both CN.

wherein R^4 is selected from the group consisting of H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, an aryl group having from about 6 to about 20 carbon atoms.

13. The composition of Claim 12, wherein the heterocyclic compounds are present in amount ranging from 0.01 to 25% by weight of the total composition.

10

14. The composition of Claim 12, further comprising at least one photoactive compound.

15. The composition of Claim 14, wherein the photoactive compound is selected from the group consisting of a retinoid, a sunscreen, or mixture thereof.

15

16. The composition of Claim 15, wherein the photoactive compound is a retinoid.

17. The composition of Claim 16, wherein the retinoid is present in amount ranging from about 0.0001 to about 20% by weight of the total composition.

20

18. The composition of Claim 15, wherein the photoactive compound is a sunscreen.

19. The composition of Claim 18, wherein the sunscreen is selected from the group consisting of a UVA chemical sunscreen, a UVB chemical sunscreen, a physical sunscreen, and mixture thereof.

25

20. The composition of Claim 19, wherein the sunscreen is a UVA chemical sunscreen.

21. The composition of Claim 20, wherein the UVA chemical sunscreen is present in amount ranging from about 0.001 to about 20% by weight of the total composition.

5

22. The composition of Claim 19, wherein the sunscreen is a UVB chemical sunscreen.


23. The composition of Claim 22, wherein the UVB chemical sunscreen is present in amount ranging from about 0.001 to about 45% by weight of the total composition.

10

24. The composition of Claim 14, further comprising at least one ingredient selected from the group consisting of oils, surfactants, humectants, botanical extracts, particulate materials, antioxidants, and other vitamins.

15

25. A method for stabilizing a photoactive compound, comprising mixing a least one photoactive compound with at least one heterocyclic compound having the structure according to Formula I:



Formula I

20 Wherein each of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ is independently selected from the group consisting of CR³ and N;

wherein R³ is selected from the group consisting of H, OH, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkoxy group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, an aryl group having from about 6 to about 20 carbon atoms;

wherein at least one of A1, A2, A3, A4, A5, A6, A7, and A8 is N;

wherein no more than four of A1, A2, A3, A4, A5, A6, A7, and A8 are N;

wherein each of R¹ and R² is independently selected from the group consisting of CN,

C(=O)OR⁴, C(=O)R⁴, F, CF₃, with the proviso that R¹ and R² are not both CN.

10 wherein R⁴ is selected from the group consisting of H, a straight or branched chain alkyl group having from about 1 to about 20 carbon atoms, an alkenyl group having from about 2 to about 20 carbon atoms, an alkynyl group having from about 2 to about 20 carbon atoms, an aryl group having from about 6 to about 20 carbon atoms.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5





FIG.6






FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG. 1