

(12) **UK Patent Application** (19) **GB** (11) **2 396 815** (13) **A**

(43) Date of A Publication **07.07.2004**

(21) Application No: **0408783.9**  
(22) Date of Filing: **27.10.1999**  
Date Lodged: **20.04.2004**  
(62) Divided from Application No **9925457.5** under Section 15(4) of the Patents Act 1977

(71) Applicant(s):  
**Phytopharm Plc**  
**(Incorporated in the United Kingdom)**  
**Corpus Christi House, 9 West Street,**  
**GODMANCHESTER, Cambridgeshire,**  
**PE29 2HY, United Kingdom**

(72) Inventor(s):  
**Roelof Marthinus Horak**  
**Vinesh Maharaj**  
**John Hakkinen**

(74) Agent and/or Address for Service:  
**Haseltine Lake & Co**  
**Imperial House, 15-19 Kingsway,**  
**LONDON, WC2B 6UD, United Kingdom**

(51) INT CL<sup>7</sup>:  
**A61K 31/57 31/60 31/704**


(52) UK CL (Edition W ):  
**A5B BJA B23X B23Y B24Y B244 B42Y B421 B423 B58Y**  
**B586**

(56) Documents Cited:  
**None**

(58) Field of Search:  
UK CL (Edition W ) **A5B**  
INT CL<sup>7</sup> **A61K**  
Other: **EPODOC, JAPIO, WPI, CAPLUS, TXTE**

(54) Abstract Title: **A composition comprising a pregnenone derivative and an NSAID**

(57) The composition comprises a non-steroidal anti-inflammatory drug and a compound of the formula wherein: R = an alkyl group containing from one to four carbon atoms;  
R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates or glucose, or a combination thereof;



Wherein: R = an alkyl group containing from one to four atoms;  
R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

The NSAID may be aspirin. The organic ester may be tiglyol or benzoyl. R may be a trisaccharide A particularly preferred compound is  
3-O-[[β-D-thevetopyranosyl-(1→4)-β-D-cymaropyranosyl-(1→4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a derivative thereof.

GASTRIC ACID SECRETION

This invention relates to the reduction of gastric acid secretion in animals including humans; to methods of treating conditions or disorders associated with or exacerbated by gastric acid secretion; to pharmaceutical compositions useful in carrying out such methods; and to the use of certain extracts and compounds in the treatment of human and non-human animals, generally mammals.

Excessive secretion of gastric acid can lead to or aggravate a number of disorders, for example oesophageal reflux disease, e.g. reflux oesophagitis, haemorrhage in and benign ulceration of the stomach and duodenum (including those complicating NSAID therapy). These conditions tend to be more problematical in obese patients and patients with hiatus hernia. The present invention is directed, in part, to the treatment of these conditions and of ancillary indications, e.g. to provide relief of reflux-like symptoms (e.g. heartburn) and/or ulcer-like symptoms (e.g. epigastric pain) associated with acid-related dyspepsia; for general dyspeptic symptoms; and for prophylaxis of acid aspiration.


Non-steroidal anti-inflammatory drugs (NSAIDs), e.g. aspirin (acetylsalicylic acid), are well-known and widely used for their anti-inflammatory and antipyretic properties. A major side effect is their tendency to damage the wall of the stomach; this adverse property is acid-dependent and is generic to the NSAIDs. The present invention is also concerned with means for mitigating these adverse effects.

35

It is known from International Patent Publication

No. WO 98/46243 that extracts of certain plants of the  
genus *Trichocaulon* or *Hoodia* possess appetite  
suppressant properties. This document also discloses  
certain specific compounds which possess appetite  
5 suppressant activity. Among these is the compound 3-O-  
[- $\beta$ -D-thevetopyranosyl-(1-4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -  
O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one; the  
structural formula of this compound is given as formula  
(1) in WO 98/46243. We have found that this compound  
10 is effective in reducing the secretion of gastric acid;  
accordingly, the compound finds application in the  
present invention. Derivatives of this compound are  
also effective in the present invention; such  
derivatives have the general formula

15



25

in which R = alkyl;

30

R<sub>1</sub> = H, alkyl, tigloyl, benzoyl, or any other  
organic ester group;

35

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrate,  
or one or more 2,6-dideoxy carbohydrates, or  
glucose molecules, or combinations thereof;

and in which the dashed lines indicate the optional presence of a double bond in the C4-C5 or C5-C6 positions.

5 Thus, according to a first aspect of the present invention, there is provided a method of reducing gastric acid secretion in an animal, which comprises administering to said animal:


10 (a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or

(b) a compound of the formula

15

20

25



wherein: R = an alkyl group containing from one to four carbon atoms;

30 R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms, or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

35

and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

5            Advantageously, when  $R_1$  is an organic ester group it is tigloyl or benzoyl. In a preferred embodiment,  $R_1$  is an alkyl group having from one to four carbon atoms, tigloyl or benzoyl; and a double bond is present in the C5-C6 position.

10

Advantageously,  $R_2$  is trisaccharide group. The component sugars of said trisaccharide group are preferably 6- deoxy and/or 2,6-dideoxy hexoses. In some embodiments, the terminal hexose moiety is thevetosyl.

15

The compound may be 3-O-[- $\beta$ -D-thevetopyranosyl-(1-4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one.


20

According to a second aspect of the present invention, there is provided a method of treating a disorder of the alimentary system in an animal, which comprises administering to the animal

25

(a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or

(b) a compound of the formula



wherein: R = an alkyl group containing from one to  
15 four carbon atoms;  
R<sub>1</sub> = H, an alkyl group containing from one to  
four carbon atoms, or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates,  
or one or more 2,6-dideoxy carbohydrates, or  
20 glucose, or a combination thereof;

and in which the dashed bonds indicate the optional  
presence of a double bond at the C4-C5 position or the  
C5-C6 position.

25


This method can conveniently be used in the  
treatment of reflux oesophagitis. It is also expected  
to find application in the treatment of epigastric  
pain; dyspepsia; gastric ulceration; and acid  
30 aspiration.

According to a third aspect of the present  
invention, there is provided a method of protecting a  
mammalian gastro-intestinal tract from damage caused by  
35 a non-steroidal anti-inflammatory drug, which comprises  
administering an effective amount of:

(a) an extract of a plant of the genus *Hoodia* or *Trichocaulon*; or

(b) a compound of the formula

5



10

15

20 wherein: R = an alkyl group containing from one to  
four carbon atoms;  
R<sub>1</sub> = H, an alkyl group containing from one to  
four carbon atoms, or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates,  
25 or one or more 2,6-dideoxy carbohydrates, or  
glucose, or a combination thereof;

and in which the dashed bonds indicate the optional  
presence of a double bond at the C4-C5 position or the  
30 C5-C6 position.

According to a fourth aspect, the present  
invention provides a method of protecting a mammalian  
gastro-intestinal tract from damage caused by a non-  
35 steroidal anti-inflammatory drug, which comprises

administering an effective amount of |3-O-[-β-D-thevetopyranosyl-(1-4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a derivative thereof.

5

When a plant extract is used in the invention, it is presently preferred to use an extract from the plant *Hoodia gordonii*. Advantageously, the said extract comprises spray-dried sap of the plant *Hoodia gordonii*.

10


The methods in accordance with this invention are applicable to the treatment of humans.

According to a fifth aspect of the present invention, there is provided a pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a compound of the formula

20

25

30



wherein: R = an alkyl group containing from one to four carbon atoms;

35

R<sub>1</sub> = H, an alkyl group containing from one to



four carbon atoms, or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates,  
or one or more 2,6-dideoxy carbohydrates, or  
glucose, or a combination thereof;

5

and in which the dashed bonds indicate the optional  
presence of a double bond at the C4-C5 position or the  
C5-C6 position.


10

A preferred pharmaceutical composition of this  
type utilises the compound 3-O-[-β-D-thevetopyranosyl-  
(1-4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-  
14β-pregn-5-en-20-one or a derivative thereof. The  
pharmaceutical compositions of this invention are  
15 conveniently prepared in unit dosage form; they may  
also include a pharmaceutically acceptable diluent,  
excipient or carrier.

According to a sixth aspect of the present  
20 invention, there is provided method of treating a  
condition or disorder caused by or exacerbated by  
gastric acid secretion in an animal, which comprises  
administering to the animal a pharmaceutical  
composition as just defined. The animal may be a  
25 human.

According to a seventh aspect, the present  
invention provides the use of a compound of the formula

30



wherein: R = an alkyl group containing from one to  
15 four carbon atoms;  
R<sub>1</sub> = H, an alkyl group containing from one to  
four carbon atoms, or an organic ester group;  
R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates,  
20 or one or more 2,6-dideoxy carbohydrates, or  
glucose, or a combination thereof;

and in which the dashed bonds indicate the  
optional presence of a double bond at the C4-  
C5 position or the C5-C6 position,

25  
in the manufacture of a medicament for the treatment of  
a disorder caused by or exacerbated by gastric acid  
secretion. For such use, the compound may be 3-O-[-β-  
D-thevetopyranosyl-(1-4)-β-D-cymaropyranosyl]-12β-O-  
30 tigloyloxy-14-hydroxy-14β-pregn-5-en-20-one or a  
derivative thereof.

According to an eighth aspect, the present  
invention provides the use of an extract of a plant of  
35 the genus *Hoodia* or *Trichocaulon* in the manufacture of  
a medicament for the treatment of conditions and

disorders caused by or exacerbated by gastric acid secretion. The condition or disorder for which the medicament is intended may be reflux oesophagitis; or gastric or duodenal ulceration; gastro-duodenal  
5 erosion; or epigastric pain.

According to a ninth aspect, the present invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a  
10 medicament for the treatment of conditions and disorders caused by or exacerbated by gastric acid secretion.

According to a tenth aspect, the present invention  
15 provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for the treatment of reflux oesophagitis.

According to an eleventh aspect, the present  
20 invention provides the use of an extract of a plant of the genus *Hoodia* or *Trichocaulon* in the manufacture of a medicament for preventing or reducing gastric damage associated with use of a non-steroidal anti-inflammatory drug.

25 According to a twelfth aspect, the present invention provides the use of 3-O-[- $\beta$ -D-thevetopyranosyl-(1-4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one or a  
30 derivative thereof in the manufacture of a medicament for preventing or reducing gastric damage associated with use of a non-steroidal anti-inflammatory drug.

The extracts of a plant of the genus *Hoodia* or  
35 *Trichocaulon* useful in the present invention may

generally be prepared by extracting sap from the plant and then spray-drying the sap. Alternatively, solvent extraction procedures may be employed. In either case, fractionation of the initial extract, e.g. by column chromatography, may follow in order to generate an extract with enhanced activity.

The extract may be prepared from plant material such as the stems and roots of plants of the genus *Hoodia* or the genus *Trichocaulon*; these genera include succulent plants which grow in the arid regions of southern Africa. Advantageously, the plant extract is obtained from one of the following species: *Trichocaulon piliferum*; *Trichocaulon officinale*; *Hoodia currorii*; *Hoodia gordonii*; and *Hoodia lugardii*.


The plant material may be homogenised in the presence of a suitable solvent, e.g. a methanol/methylene chloride solvent, by means of a device such as a Waring blender. The extraction solution may then be separated from residual plant material by an appropriate separation procedure such as, for example, filtration or centrifugation. The solvent may, for example, be removed by means of a rotary evaporator, preferably in a water bath at a temperature of 60°C. The separated crude extract may then be further extracted with methylene chloride and water before being separated into a methylene chloride extract and a water extract. The methylene chloride extract may have the solvent removed by, for example, a rotary evaporator and the resultant extract may be further purified by way of a methanol/hexane extraction. The methanol/hexane extraction product may then be separated to yield a methanol extract and a hexane extract. The methanol extract may be evaporated

to remove the solvent in order to yield a partially purified active extract.

The partially purified active extract may be  
5 dissolved in methanol, and may be further fractionated  
by column chromatography, employing silica gel as an  
adsorption medium and a chloroform/30% methanol mixture  
as eluent. A plurality of different fractions may be  
obtained, and each may be evaluated, by suitable  
10 bioassaying procedures, to determine their activity.  
High activity fractions may be further fractionated,  
e.g. by column chromatography using silica gel as an  
adsorption medium and a 9:1 chloroform:methanol eluent.  
This process may be repeated using silica gel as an  
15 adsorption medium and a 9:1 ethylacetate:hexane eluent.

Alternatively, the plant product may be compressed  
or macerated to extract sap therefrom, then filtered to  
remove unwanted solids and freeze-dried. The freeze-  
20 dried sap product may then be further purified, if  
desired, for example using chromatographic  
fractionation such as described above.

The derivatives of 3-O-[- $\beta$ -D-thevetopyranosyl-  
25 (1-4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-  
14 $\beta$ -pregn-5-en-20-one which find use in this invention  
are advantageously of the formula:



in which R = alkyl;

R<sub>1</sub> = H, alkyl, tigloyl, benzoyl, or any other organic ester group;

R<sub>2</sub> = H or alkyl;

R<sub>3</sub> = H or OH;

25 and in which (a) the dashed lines indicate the optional presence of a further bond in the C4-C5 or C5-C6 positions; and (b) the carbonyl group is optionally reduced to the group -CHOH.

30 The invention will be illustrated by the following Examples:

Example 1

The ability of the compound 3-O-[-β-D-thevetopyranosyl-  
35 (1-4)-β-D-cymaropyranosyl]-12β-O-tigloyloxy-14-hydroxy-  
14β-pregn-5-en-20-one - hereinafter referred to as

P57AS3 - to prevent aspirin-induced damage to the stomach was investigated in the rat. Three groups each containing eight rats were given aspirin in 0.5% CMC orally at a dose of about 600 mg/kg (1.0 ml of 60  
5 mg/ml). The first group (the control group) was additionally given 0.5% CMC. The second group was given compound P57AS3 at a dose rate of 100mg/kg; and the third group (comparative group) was given  
cimetidine at a dose rate of 20mg/kg.

10

After 5 hours, the animals' stomachs were visually inspected for damage. Damage was assessed on a scale in which each small haemorrhage scored one point; each large haemorrhage scored two points; each small ulcer  
15 \_ scored three points; and each large ulcer scored four points. The results obtained were as follows:

Table 1

Oral treatment	Dose (mg/kg)	Rat no.	Gastric damage score
5 Vehicle (0.5% CMC)	-	1	10
		2	10
		3	16
		4	8
		5	5
		6	3
		7	1
		8	6
		Mean	7.38
10 P57AS3	100	9	0
		10	0
		11	3
		12	0
		13	0
		14	0
		15	2
		16	0
		Mean	0.63
15 Cimetidine	20	17	0
		18	0
		19	4
		20	0
		21	0
		22	0
		23	0
		24	0
		Mean	0.50

20 These results show that compound P57AS3 gave an average reduction in gastric damage of 91.5% compared to 93.2% for cimetidine. In both cases, the statistical



significance of the difference from the control group was  $P < 0.01$ .

Example 2

5

The effect of both the spray dried extract (sap from *Hoodia gordonii*) - referred to hereinafter as P57SD - and the purified compound P57AS3 were examined for effects on gastric acid secretion in the pylorus-  
10 ligated rat. Male Sprague-Dawley derived CD rats, 175-230 g, were used. On the day before the test, rats were outfitted with stainless steel tail cups to prevent coprophagy, housed individually with water allowed ad lib, and fasted overnight (18 hr). In one  
15 test the spray dried sap P57SD (5 or 50 mg/kg), or cimetidine (10 mg/kg) or vehicle (water) was orally administered to the rats immediately before fasting. Water was removed 30 minutes prior to ligation on the  
20 morning of the test. Under isoflurane anesthesia, a midline celiotomy was performed, and the pylorus of each rat was ligated with 4-0 silk.

In a second experiment, purified compound P57AS3 was administered intraduodenally (i.d.) at 1 or 10 mg/kg at  
25 the time of the pylorus ligation. In this experiment with intraduodenal administration of P57AS3, vehicle (DMSO), cimetidine (10 mg/kg) or P57AS3 (1 or 10 mg/kg) was administered immediately after ligation. The incisions were then closed with staples. In the  
30 experiments with subcutaneous administration of P57, vehicle (DMSO), cimetidine (10 mg/kg) or P57, 10 or 30 mg/kg were injected subcutaneously in the nape of the neck immediately after the incisions were closed. Two  
35 hours after ligation, the rats were killed, the contents of each stomach collected and cleaned by

centrifugation (30 minutes at 2000 x g), and the volume of gastric fluid in each stomach was determined. Using a Radiometer pH meter and autotitrator, the pH and acid concentration of each gastric sample was determined by  
5 titration to pH 7 with 0.1N NaOH. Results are expressed as  $\mu\text{eq/hr/100 g}$  (body weight) and % vehicle control.

In the first experiment, gastric acid output was  
10 determined 18 hours after rats were orally administered the spray dried sap P57SD. Results showed that the 5 mg/kg dose of P57SD was without effect, but the 50 mg/kg dose-inhibited gastric acid secretion by 40% (Table 2). In a second experiment, the purified  
15 compound P57AS3 was administered intraduodenally (i.d.) at 1 or 10 mg/kg at the time of the pylorus ligation. The 10 mg/kg i.d. dose of P57AS3 decreased acid secretion by 88%, but the 1 mg/kg i.d. dose had no effect (Table 2). Experiments were then conducted to  
20 determine if systemically administered P57AS3 would also inhibit gastric acid secretion. P57AS3 was administered subcutaneously at the time of ligation in 3 experiments using 10 mg/kg and in one experiment using 30 mg/kg. In one of the experiments with 10  
25 mg/kg, P57AS3 produced a significant decrease in gastric acid secretion, but in the other 2 experiments with 10 mg/kg and the one experiment with 30 mg/kg there was a trend toward decreased acid output but the amount of acid secretion was not significantly  
30 different from vehicle-treated controls (Table 2). In each pylorus ligation experiment, gastric acid secretion was significantly reduced (range 36% to 83% inhibition) in a group of rats treated with cimetidine (10 mg/kg), which was included as a positive control.

Table 2 - Effect of P57 on gastric acid secretion in the pylorus-ligated rat

	Exp.	Active	Dose (mg/kg)	Route	Acid Output % Control mean ± sd
5					
10	I	P57SD	5	po	85.4±34.5
			50	po	*60.0±21.7
	II	P57AS3	1	id	109.0±37.8
			10	id	*12.4±1.1
15	III	P57AS3	10	sc	48.9±29.2
	IV	P57AS3	10	sc	74.0±38.0
	V	P57AS3	10	sc	*57.1±19.8
20	VI	P57AS3	30	sc	63.7±17.3

\*significantly different from vehicle group,  $p \leq 0.05$

## 25 SUMMARY OF RESULTS

### In Vitro Studies

Purified P57AS3 was examined for binding affinity to a variety of receptors, channels, and uptake sites in vitro. At a concentration of 10  $\mu\text{M}$ , In isolated tissue studies, P57AS3 at concentrations  $\leq 10 \mu\text{M}$  had no effect on the rate of spontaneously beating guinea pig right atria or on the force of contraction of electrically stimulated left atria. At concentrations of 1  $\mu\text{M}$  and higher, P57AS3 produced intermittent and transient elevations in the basal tension of electrically stimulated left atria. P57AS3 had no effect on resting tension of isolated guinea pig gallbladder, ileum or trachea, or on CCK octapeptide-induced contractions of the gallbladder (CCK-A receptor mediated), histamine induced

contractions of the guinea pig ileum ( $H_1$  receptor mediated), or acetylcholine-induced contractions of the guinea pig ileum ( $M_3$  receptor mediated). At concentrations  $\geq 1 \mu M$ , P57AS3 noncompetitively inhibited 5 carbachol-induced contractions of the guinea pig trachea ( $M_3$  receptor mediated).

#### In Vivo Studies

The spray dried form of P57 was examined for effects on 10 gastrointestinal motility and both the spray dried and purified forms were examined for effects on gastric acid secretion in the rat. At a dose of 5 mg/kg p.o., spray dried P57 had no effect on gastrointestinal transit or on gastric acid secretion. However, a 50 15 mg/kg dose of spray dried P57 inhibited gastric emptying by 26% and acid output by 40%. Following intraduodenal (id) administration in the rat, purified P57AS3 had no effect on gastric acid secretion at a dose of 1 mg/kg, but a 10 mg/kg dose decreased acid 20 output by 88%. In 2 of 3 experiments at a dose of 10 mg/kg and in one experiment at a dose of 30 mg/kg, subcutaneously (sc) administered purified P57AS3 had no effect on gastric acid output, but in 1 of 3 25 experiments the 10 mg/kg sc dose reduced gastric acid secretion by 43%.


References

1. Hill SJ, and Young JM. Characterization of  
[3H]mepyramine binding to the longitudinal muscle of  
guinea pig small intestine. Mol Pharmacol 1981;  
5 19:379-387
  
2. Eltze M, Mutschler E, and Lambrecht G. Affinity  
profiles of pizotifen, ketotifen and other tricyclic  
antimuscarinics at muscarinic receptor subtypes M1, M2  
10 and M3. Eur J Pharmacol 1992; 211:283-293.
  
3. Bishop LA, Gerskowitch VP, Hull RA, Shankley NP,  
Black JW. Combined dose-ratio analysis of  
cholecystokinin receptor antagonists, devazepide,  
15 lorglumide and loxiglumide in the guinea pig  
gallbladder. Br J Pharmacol 1992; 106:61-66.

CLAIMS

1. A pharmaceutical composition comprising a non-steroidal anti-inflammatory drug and a compound of the formula

5



wherein: R = an alkyl group containing from one to four carbon atoms;

10

R<sub>1</sub> = H, an alkyl group containing from one to four carbon atoms or an organic ester group;

R<sub>2</sub> = H, or one or more 6-deoxy carbohydrates, or one or more 2,6-dideoxy carbohydrates, or glucose, or a combination thereof;

and in which the dashed bonds indicate the optional presence of a double bond at the C4-C5 position or the C5-C6 position.

15

2. A pharmaceutical composition according to claim 1, wherein said organic ester group is tigloyl or benzoyl.

20

3. A pharmaceutical composition according to claim 1, wherein R<sub>1</sub> is an alkyl group having from one to four carbon atoms, tigloyl or benzoyl; and a double bond is present in the C5-C6 position.

25

4. A pharmaceutical composition according to claim 1, wherein R<sub>2</sub> is a trisaccharide group.

5. A pharmaceutical composition according to claim 4, wherein the component sugars of said trisaccharide group are 6-deoxy and/or 2,6-dideoxy hexoses.
6. A pharmaceutical composition according to claim 5, wherein the terminal hexose moiety is thevetosyl.
7. A pharmaceutical composition according to claim 1, wherein said compound is 3-O-[ $\beta$ -D-thevetopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl]-12 $\beta$ -O-tigloyloxy-14-hydroxy-14 $\beta$ -pregn-5-en-20-one or a derivative thereof.
8. A pharmaceutical composition according to any preceding claim, wherein the non-steroidal anti-inflammatory drug is aspirin.
9. A pharmaceutical composition according to any preceding claim, which is prepared in unit dosage form.



INVESTOR IN PEOPLE

Application No: GB0408783.9

Examiner: Mr Jason Bellia

Claims searched: 1-9

Date of search: 5 May 2004

### Patents Act 1977: Search Report under Section 17

#### Documents considered to be relevant:

Category	Relevant to claims	Identity of document and passage or figure of particular reference
		None

#### Categories:

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

#### Field of Search:

Search of GB, EP, WO & US patent documents classified in the following areas of the UKC<sup>W</sup> :

A5B

Worldwide search of patent documents classified in the following areas of the IPC<sup>07</sup>

A61K

The following online and other databases have been used in the preparation of this search report

EPODOC, JAPIO, WPI, CAPLUS, TXTE