




(86) **Date de dépôt PCT/PCT Filing Date:** 2008/11/10
(87) **Date publication PCT/PCT Publication Date:** 2009/05/28
(45) **Date de délivrance/Issue Date:** 2016/07/05
(85) **Entrée phase nationale/National Entry:** 2010/05/20
(86) **N° demande PCT/PCT Application No.:** EP 2008/009451
(87) **N° publication PCT/PCT Publication No.:** 2009/065507
(30) **Priorité/Priority:** 2007/11/23 (IT MI2007A002225)

(51) **Cl.Int./Int.Cl. A61K 9/08** (2006.01),
A61K 9/00 (2006.01)
(72) **Inventeurs/Inventors:**
MAGGI, CARLO ALBERTO, IT;
GIULIANI, SANDRO, IT;
QUARTARA, LAURA, IT
(73) **Propriétaire/Owner:**
ISTITUTO LUSO FARMACO D'ITALIA S.P.A., IT
(74) **Agent:** KIRBY EADES GALE BAKER

(54) **Titre : COMPOSITIONS PHARMACEUTIQUES CONTENANT DES ANTAGONISTES DE LA BRADYKININE ET DE L'ACIDE
HYALURONIQUE, ET LEURS UTILISATIONS**
(54) **Title: PHARMACEUTICAL COMPOSITIONS CONTAINING BRADYKININ ANTAGONISTS AND HYALURONIC ACID, AND USES
THEREOF**

(57) **Abrégé/Abstract:**

Disclosed are pharmaceutical compositions containing, as the active ingredients, a mixture of a hyaluronic acid polymer with a bradykinin B2 receptor antagonist. Said compositions have proved especially effective in the treatment of degenerative joint diseases such as osteoarthritis using intra-articular injections.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 May 2009 (28.05.2009)

PCT

(10) International Publication Number
WO 2009/065507 A2

(51) International Patent Classification: Not classified

(74) Agents: BANFI, Paolo et al.; Bianchetti Bracco Minoja S.r.l., Via Plinio, 63, I-20129 Milano (IT).

(21) International Application Number:
PCT/EP2008/009451

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:
10 November 2008 (10.11.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
MI2007A002225
23 November 2007 (23.11.2007) IT

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): ISTITUTO LUSO FARMACO D'ITALIA S.p.A. [IT/IT]; Via Walter Tobagi, 8, I-20068 Peschiera Borromeo (IT).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): MAGGI, Carlo, Alberto [IT/IT]; Borgo Tegolaio, 15, I-50125 Firenze (IT). GIULIANI, Sandro [IT/IT]; Via Carlo Borrani, 11, I-50012 Bagno a Ripoli (IT). QUARTARA, Laura [IT/IT]; Via delle Cascine, 22, I-50144 Firenze (IT).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING BRADYKININ ANTAGONISTS AND HYALURONIC ACID, AND USES THEREOF

(57) Abstract: Disclosed are pharmaceutical compositions containing, as the active ingredients, a mixture of a hyaluronic acid polymer with a bradykinin B2 receptor antagonist. Said compositions have proved especially effective in the treatment of degenerative joint diseases such as osteoarthritis using intra-articular injections.



WO 2009/065507 A2

**PHARMACEUTICAL COMPOSITIONS CONTAINING BRADYKININ
ANTAGONISTS AND HYALURONIC ACID, AND USES THEREOF**

FIELD OF INVENTION

The invention relates to pharmaceutical compositions containing, as the active ingredients, a mixture of a hyaluronic acid polymer with a bradykinin B2 receptor antagonist. Said compositions have proved especially effective in the treatment of degenerative joint diseases such as osteoarthritis using intra-articular injections.

STATE OF THE ART

Osteoarthritis (OA), also known as degenerative joint disease, is a painful, progressive, degenerative disorder of the joints. The main pathophysiological characteristics of OA are destruction and loss of joint cartilage, hypertrophy, inflammation of the synovial membrane, and consequent swelling of the joint. These effects produce symptoms such as pain, stiffness and loss of function. The high incidence of OA in the elderly population, associated with the increase in average life expectancy, indicates that the number of patients affected by this disorder is likely to increase considerably in the near future. OA patients consider pain reduction to be very important to their quality of life.

No drugs which stop the progress of this disorder are currently available. The existing treatments are mainly designed to reduce the pain symptoms and regain the joint function. Paracetamol and non-steroidal anti-inflammatory drugs (NSAIDs) are widely prescribed for the treatment of pain in osteoarthritis. However, long-term use of said drugs can be accompanied by major adverse effects, especially at gastrointestinal level (ulcers) and in terms of platelet aggregation. Intra-articular injection of corticosteroids reduces the associated inflammation and pain, but they are

rarely used as their effect is short-lived. There is consequently a clear need for new therapeutic agents that reduce the pain and inflammation associated with osteoarthritis.

Bradykinin (BK) is a member of the kinins, a family of small peptides (8-11 aminoacids) which derive from precursors with a high molecular weight (kininogens) following attack by enzymes with peptidase activity (kallikreins). Kinin formation is activated in various circumstances, involving inflammatory, ischaemic and immune processes or bacterial and viral infections.

Two kinin receptors have been pharmacologically characterised: the B1 receptor, which is minimally expressed under normal conditions, but whose expression is induced by the stimuli listed above, and the B2 receptor, which is constitutionally expressed by many cell types. Bradykinin, through stimulation of the B2 receptor, is one of the most important mediators of inflammation and pain, and is involved in the release of pro-inflammatory and hyperalgesic mediators.

It has been demonstrated that bradykinin (BK) participates in the pathophysiology of OA at various levels.

It has long been known that kinins are released into the synovial fluid of OA sufferers. Moreover, in these patients, the B2 receptor has been found in the cells lining the synovial cavity, the fibroblasts, and the endothelial cells of the blood vessels.

Many studies with various preclinical models indicate that BK, when administered by the intra-articular route, induces plasma extravasation and accumulation of neutrophils in the synovial membrane of the rat more effectively than other inflammation mediators such as substance P, histamine, and calcitonin gene-related peptide. Moreover, BK reduces the proteoglycans content in the joint cartilage and generates the release of prostaglandins in

murine OA models.

Some bradykinin B2 receptor antagonists have proved effective in inhibiting inflammatory events and hyperalgesia in various animal synovitis models.

5 After its release, BK excites and sensitises the sensory nerve fibres that innervate the articular capsule.

The clinical significance of BK has been demonstrated in a phase II trial conducted on 58 patients with symptomatic OA of the knee, in which a single intra-articular administration of B2 receptor antagonist icatibant (90 µg/1 ml)
10 reduced the intensity of pain in the knee to a greater extent than the placebo (55 patients). Sanofi-Aventis recently reported that in patients with OA of the knee, intra-articular infiltration of icatibant (3 x 500 µg injections one week apart) induces a strong analgesic response which lasts for up to 3 months after the treatment, and this considerable analgesic effect is obtained with
15 negligible or no side effects.

Many bradykinin B2 receptor antagonists have been described in the literature.

EP370453 describes some compounds with a peptide structure which act as bradykinin antagonists, and said compounds include the one defined as
20 icatibant. Icatibant also forms the subject of patent EP1594520, wherein its use in the prophylaxis and treatment of osteoarthritis is disclosed.

WO03103671 describes a group of very powerful non-peptide bradykinin antagonists. A selection of particularly potent antagonists is reported in WO2006040004, including the compound MEN16132; also these
25 antagonists have proved highly effective in the prophylaxis and treatment of osteoarthritis, especially in intra-articular treatments of the knee.

Hyaluronan (also known as hyaluronic acid or hyaluronate) is a non-sulphated glycosaminoglycan widely distributed in the endothelial,

connective, epithelial and neural tissues. It is one of the major components of extracellular matrix, and significantly contributes to cell proliferation and migration. On average, a person weighing 70 kg has 15 g of hyaluronan in the body, a third of which is exchanged (broken down and synthesised) every day.

5 Hyaluronan is one of the major components of synovial fluid, and increases its viscosity. Together with lubricin, it is one of the main lubricants of the fluid components. Hyaluronan is also an important component of joint cartilage, where it is found as the lining of each cell (chondrocytes).

 Hyaluronan has long been used for treatment of OA of the knee (Puhl
10 W; Scharf P (1997). *Ann Rheum Dis* 56 (7): 637-40.) Said treatment, known as viscosupplementation, consists of a series of injections into the knee joint which are designed to increase the viscosity of the fluids contained in it, lubricating and supporting the joint and consequently producing an analgesic effect. It has also been postulated that hyaluronan has a favourable
15 biochemical effect on the cartilage cells. The first product of this kind to be launched on the market was Hylan G-F 20 (Synvisc®), which has been marketed by Genzyme since 1998. The European Medicines Agency (EMA) extended the approval of Hylan GF-20 as a treatment for OA pain to the hip in 2002, and to the ankle and shoulder in 2007. Other products on the market
20 include Ostenil® (TRB CHEMEDICA), Suplasyn® (MERCKLE RECORDATI) and GO-ON® (ROTTAPHARM).

 A meta-analysis of 18 randomised clinical trials has confirmed the clinical benefits of Hylan G-F 20 and this class of viscosupplements, concluding that viscosupplements are superior to a placebo and that many of
25 them are more effective than steroids in the treatment of pain associated with OA of the knee (Positive Synvisc Data Published by Cochrane Collaboration Source, Press Release 2005, May 05, Genzyme Corp).

 Structurally, hyaluronan is a polymer consisting of repetitive

disaccharide units of N-acetylglucosamine and sodium glucuronate. The one used for the treatment of OA is generally extracted from coxcombs, and fractions of various mean molecular weights, ranging from 0.5 to 10 million daltons, can be isolated. The viscosupplementation effect varies, depending on the molecular weight and density (Gomis et al, Arthritis & Rheumatism, 2004, 50:314-326).

Numerous methods have been proposed for the treatment of osteoarthritis, including the use of BK antagonists or hyaluronic acid.

WO03063799 also proposes the use of pharmaceutical compositions containing a plurality of chondroprotective agents, including matrix metal proteinase (MMP) inhibitors (bradykinin inhibits MMP production), but no mention is made of the use of hyaluronic acid.

However, there is an unmet need for effective treatments for degenerative joint diseases such as osteoarthritis.

15 DISCLOSURE OF THE INVENTION

It has now surprisingly been found that pharmaceutical compositions containing, as the active ingredients:

- a) hyaluronic acid
- b) a bradykinin B2 receptor antagonist

20 show surprising efficacy in the treatment of degenerative joint diseases such as, but not limited to, osteoarthritis.

The present invention relates to pharmaceutical compositions containing a bradykinin B2 receptor antagonist, the following ones being preferred:

- 25 ▪ H-D-Arg-Arg-Pro-Hyp-Gly-Igl-Ser-D-F5F-Igl-Arg-OH (B10056),
- H-Arg-Arg-Pro-Hyp-Gly-Igl-Ser-D-Igl-Oic-Arg-OH (B9430),
- H-D-Arg-Arg-Pro-Hyp-Gly-Thi-Ser-D-Tic-Oic-Arg-OH (Icatibant),
- 4-[2-[[[3-(3-Bromo-2-methyl-imidazo[1.2-a]pyridine-8-yl oxymethyl)-

2,4-dichloro-phenyl]-methyl-carbamoyl]-methyl)-carbamoyl]-vinyl]-
N,N-dimethyl-benzamide, (FR167344)


- 3-(6-Acetylamino-pyridine-3-yl)-N-([2,4-dichloro-3-(2-methyl-
quinoline-8-yl oxymethyl)-phenyl]-methyl-carbamoyl]-methyl)-
5 acrylamide, (FR173657 or FK3657)

- 1-[2,4-Dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-benzene
sulphonyl]-pyrrolidine-2-carboxylic acid [3-(4-carbamido-yl-
benzoylamino)-propyl]-amide, (LF160687, Anatibant)

- 4-(4-[1-[2,4-Dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-
10 benzene sulphonyl]-pyrrolidine-2-carbonyl]-piperazine-1-carbonyl)-
benzamidine, (LF160335)

- 2-[5-(4-Cyano-benzoyl)-1-methyl-1H-pyrrol-2-yl]-N-[2,4-dichloro-3-
(2-methyl-quinoline-8-yl oxymethyl)-phenyl]-N-methyl-acetamide,
or one of the compounds described in WO2006/04004, having general

15 formula (I)



(I)

wherein

- 20 - R is hydrogen or methyl
- W represents a simple bond or an O atom
- $n = 3, 4$

- X is hydrogen or a -NR₁R₂ amino group in which R₁ and R₂ can be independently of one another hydrogen or a group selected from methyl, ethyl n-propyl, isopropyl,
- Y is a quaternary ammonium (-NR₃R₄R₅)⁺A⁻ in which R₃, R₄,
5 R₅ can be independently of one another methyl, ethyl n-propyl, isopropyl, butyl, isobutyl, n-pentyl and A⁻ is an anion of a pharmaceutically acceptable acid;

the pharmacologically acceptable salts, enantiomers and enantiomeric mixtures thereof.

10 For the purposes of this invention, a pharmaceutically acceptable acid is an acid selected from hydrochloric, hydrobromic, phosphoric, carbonic, acetic, sulphuric, trifluoroacetic, methansulphuric, succinic, maleic, malic, malonic, citric, edetic acid; where the anion carries two or more negative charges, A⁻ shall be a fractional value.

15 Among the compounds of general formula (I), the following compound is particularly preferred: (4-(S)-Amino-5-(4-{4-[2,4-dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-benzenesulphonylamino]-tetrahydro-pyran-4-carbonyl}-piperazin-1-yl)-5-oxo-pentyl]-trimethyl-ammonium, in salified
20 form with ions formally deriving from an acid selected from hydrochloric, acetic, sulphuric, trifluoroacetic, methanesulphonic, succinic and edetic acids; chloride dihydrochloride is the compound defined as MEN16132 (MW 871.5).

The hyaluronic acid used has a mean molecular weight between 0.5 and 10 million daltons, and preferably between 4 and 9 million daltons; absolutely preferred is the hyaluronic acid with a mean MW between 5 and 8 million
25 daltons, as is or in the forms of the sodium or potassium salt.

The compositions according to the invention contain an amount of bradykinin antagonist per dose between 5.7×10^{-5} and 2.3×10^{-2} mmols (which, in the case of MEN16132, corresponds to an approximate amount of 0.05 to

20 mg), preferably between 1.1×10^{-4} and 1.1×10^{-2} mmols (which, in the case of MEN16132, corresponds to an approximate amount of 0.1 to 10 mg), and even more preferably between 2.9×10^{-4} and 5.7×10^{-3} mmols (which, in the case of MEN16132, corresponds to an approximate amount of 0.25 to 5 mg).

5 Said compositions also contain hyaluronic acid in the amount of 1 to 100 mg, and preferably 5 to 20 mg, per dose.

Pharmaceutical formulations according to this invention can also contain one or more pharmaceutically acceptable carriers/excipients.

Liquid and semisolid pharmaceutical forms suitable for topical
10 administration, such as solutions, creams, gels or transdermal patches, are preferred; in particular, forms suitable for intra-articular or intrabursal injection, such as solutions, and transdermal application, such as semisolid forms like creams or gels and transdermal patches. The pharmaceutical form
15 can also consist of a form wherein some or all of the components are in a dry form, possibly lyophilised, to be reconstituted with an aqueous solution or other suitable vehicle before use.

Said formulations can be produced by methods well-known in the state of the art using known excipients such as binders, disintegrants, fillers, stabilisers, diluents and colorants. They can also include delayed- or
20 slow-release forms made with suitable polymers known in pharmaceutical technology.

Pharmaceutically acceptable carriers/excipients such as solvents, preservatives such as antioxidants and/or chelating agents and antimicrobials, isotonicity regulators, and buffer systems are preferred for the preparation of
25 liquid forms suitable for injectable use.

Water is preferable as solvent, possibly with co-solvents such as glycols, or polyalcohols such as ethylene glycol.

Preservatives or chelating agents may also be used, sodium edetate and

sodium metabisulphite being preferred, and antimicrobials, benzyl alcohol being preferred.

Sodium chloride or mannitol are particularly preferred as isotonicity regulators.

5 The preferred buffer systems can be the complex of salts for the phosphate and citrate buffer, preferably in the form of sodium or potassium salts.

In this invention, especially in the description of compounds with a peptide structure, the following abbreviations have been used for some
10 non-natural amino acids: Nal = naphthyl-alanine; NMePhe = N-methyl-phenylalanine, Oic = Octahydroindol-2-carboxylic acid, Hyp = Hydroxyproline, Igl = aminoindanecarboxylic acid, Cpg = 1-aminocyclopentanecarboxylic acid, Tic = 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, F5F = pentafluorophenylalanine,

15 Typical examples of formulations according to this invention are:

1. Hyaluronic acid, sodium salt, with a mean molecular weight between 0.5 and 10 million daltons, 5-20 mg, MEN-16132 0.25-5 mg, in saline solution (0.9% NaCl), 0.1 N HCl q.s. to pH 4.5, water q.s. to 1 ml.
- 20 2. Hyaluronic acid, sodium salt, with a mean molecular weight between 0.5 and 10 million daltons, 5-20 mg, MEN-16132 0.25-5 mg, in saline solution (0.9% NaCl), 0.1 N HCl q.s. to pH 6, water q.s. to 1 ml.
3. Hyaluronic acid, sodium salt, with a mean molecular weight
25 between 0.5 and 10 million daltons, 5-20 mg, MEN-16132 0.25-5 mg, in saline solution (0.9% NaCl), phosphate buffer (pH 6-8), water q.s. to 1 ml.
4. Hyaluronic acid, sodium salt, with a mean molecular weight

between 0.5 and 10 million daltons, 5-20 mg, MEN-16132 0.25-5 mg, in saline solution (0.9% NaCl), citrate buffer (pH 6-8), water q.s. to 1 ml.

- 5 5. Extempore preparation obtained by dissolving MEN-16132, 0.25-5 mg, in lyophilised form, with a solution of hyaluronic acid, sodium salt (mean molecular weight between 0.5 and 10 million daltons, 5-20 mg), with phosphate buffer in saline solution (0.9% NaCl), water q.s. to 1 ml.
- 10 6. Hyaluronic acid, sodium salt, with a mean molecular weight between 0.5 and 10 million daltons 5-20 mg, icatibant (MW 1304.5) 0.37-6.5 mg, in saline solution (0.9% NaCl), phosphate buffer (pH 6-8), water q.s. to 1 ml.

The pharmaceutical compositions according to the invention are useful in the prophylaxis and treatment of inflammatory, autoimmune, traumatic and
15 degenerative joint diseases such as osteoarthritis and post-traumatic osteoarthritis, degenerative osteoarthritis (gonarthrosis, spondylarthrosis); spondylosis, synovitis, tenosynovitis, bursitis, contusions, sprains, dislocations and subluxations, and in joint diseases caused by developmental alterations such as osteochondritis and dysplasia.

20 The dose can vary according to age and the patient's general state of health, the nature and gravity of the disease or disorder, and the route and type of administration. In the case of intra-articular use in an adult human patient, the use of the pharmaceutical compositions according to the invention could involve a weekly dose (in a single administration) of bradykinin antagonist
25 amounting to between 2.9×10^{-4} and 5.7×10^{-3} mmols (which, in the case of MEN16132, corresponds to approx. 0.25 to 5 mg) and between 5 and 20 mg of hyaluronic acid.

The following examples illustrate the invention in greater detail:

Example 1

Hyaluronic acid, sodium salt, with a mean molecular weight of 6 million daltons, 10 mg, MEN16132 0.5 mg, in saline solution (0.9% NaCl), 0.1N HCl q.s. to pH 4.5, water q.s. to 1 ml. The solution is placed in pre-filled
5 2.25 ml syringes.

Example 2

Hyaluronic acid, sodium salt, with a mean molecular weight of 6 million daltons, 10 mg, MEN16132 0.5 mg, in saline solution (0.9% NaCl), containing phosphate buffer (Na_2HPO_4 0.16 mg NaH_2PO_4 0.04 mg), water q.s.
10 to 1 ml. The solution is placed in pre-filled 2.25 ml syringes.

Example 3

Hyaluronic acid, sodium salt, with a mean molecular weight of 6 million daltons, 10 mg, MEN16132 0.2 mg, in saline solution (0.9% NaCl), 0.1N HCl q.s. to pH 4.5, water q.s. to 1 ml. The solution is placed in pre-filled
15 2.25 ml syringes.

Example 4

Hyaluronic acid, sodium salt, with a mean molecular weight of 6 million daltons, 10 mg, in saline solution (0.9% NaCl), containing phosphate buffer (Na_2HPO_4 0.16 mg, NaH_2PO_4 0.04 mg), water q.s. to 1 ml. MEN16132,
20 in lyophilised form, is dissolved with the solution described above.

Example 5

Hyaluronic acid, sodium salt, with a mean molecular weight of 6 million daltons, 10 mg, icatibant 0.5 mg, in saline solution (0.9% NaCl), containing phosphate buffer (Na_2HPO_4 0.16 mg, NaH_2PO_4 0.04 mg), water
25 q.s. to 1 ml. The solution is placed in pre-filled 2.25 ml syringes.

Biological activity

The activity of MEN16132, icatibant and hyaluronic acid was measured in an experimental model of osteoarthritis induced by intra-articular injection

of sodium monoiodoacetate (MIA) which inhibits glycolysis in the chondrocytes, thus causing damage and death to those cells with consequent degeneration of the joint surface, in a very similar way to human osteoarthritis.

5 MIA (1 mg/25 μ l) was injected into the intra-articular space of the right knee of the rat, while 25 μ l of saline was administered to the left knee as internal control.

The administration of MIA causes pain, difficulty of walking and inability to bear weight on the paw corresponding to the treated knee; the
10 weight of the body therefore falls mainly on the left paw, to an extent directly proportional to the pain perceived. This effect persists for many weeks. The imbalance in weight on the two weight-bearing paws, evaluated in a non-invasive way with the incapacitance test, measures the pain resulting from the joint damage (osteoarthritis) induced by MIA.

15 The purpose of this test was to evaluate the protective effect of bradykinin B2 receptor antagonists against experimental osteoarthritis, and to establish the effect of co-administration of hyaluronic acid.

Seven days after the treatment with MIA, the compounds under study were injected, and the test was repeated at various times to measure the
20 painkilling effect and its duration.

MEN16132 and icatibant, at the dose of 3 μ g/25 μ l i.a., reduced the pain caused by osteoarthritis by 55% and 40% respectively; the maximal inhibitory effect was observed 3 days after administration of the compounds. No further significant increase in inhibitory effect was observed at the dose of
25 10 μ g/25 μ l i.a.. The antinociceptive effect lasted for over a week with both compounds.

The administration of hyaluronic acid (molecular weight 6 million daltons, 50 μ g/25 μ l i.a.) produced a modest antinociceptive effect, reducing

the pain response by 16%.

Co-administration of MEN16132 or icatibant with hyaluronic acid considerably boosted the antinociceptive response in the rats in which osteoarthritis was induced with MIA. MEN16132 or icatibant (3 µg/25 µl i.a.)
5 and hyaluronic acid (50 µg /25 µl) reduced pain, measured as imbalance of body weight between the untreated paw and the treated paw, by 72% and 65% respectively. On the basis of direct observation of their motor activity, the motor behaviour of the rats treated with MEN16132 or icatibant and hyaluronic acid did not differ from that of the controls not suffering from
10 osteoarthritis. The effect of the two B2 receptor antagonists after a single administration in association with hyaluronic acid also lasted longer, in some cases for over two weeks.

Morphological and histological tests were performed to obtain further confirmation of the surprising efficacy of the association with hyaluronic acid
15 in reducing the symptoms and damage caused by osteoarthritis. 14 days after treatment with the association of MEN16132 or icatibant and hyaluronic acid, a substantial reduction (over 50%) in the damage present on the joint surface of the knee was observed, characterised by reduced loss of chondrocytes, glycosaminoglycan matrix and exposure of subarticular bone, with far superior
20 results to the single administration of B2 antagonists or hyaluronic acid.

Moreover, the association is particularly interesting from the application standpoint, because kinin B2 receptor antagonists achieve the antinociceptive effect slowly but with a long duration, whereas hyaluronic acid with a high molecular weight (6 million daltons) or a salt thereof
25 produces the maximum effect within a few hours of administration. The association disclosed in this invention is therefore rapidly effective and long-lasting due to the complementary and boosting effect of the two classes of compounds.

CLAIMS

1. A pharmaceutical composition comprising, as the active ingredients, a mixture of:

5 a) hyaluronic acid or salts thereof wherein the salts are sodium or potassium salts

b) a bradykinin B2 receptor antagonist

together with a pharmaceutically acceptable carrier or excipient, wherein:

the hyaluronic acid is in polymer form with a mean MW between 0.5 and 10
10 million daltons, and

the bradykinin antagonist is:

- B2 receptor antagonists, wherein the B2 receptor antagonists are:


- H-D-Arg-Arg-Pro-Hyp-Gly-Igl-Ser-D-F5F-Igl-Arg-OH
- H-Arg-Arg-Pro-Hyp-Gly-Igl-Ser-D-Igl-Oic-Arg-OH
- 15 ▪ H-D-Arg-Arg-Pro-Hyp-Gly-Thi-Ser-D-Tic-Oic-Arg-OH
(Icatibant)
- 4-[2-[[[3-(3-Brom-2-methyl-imidazo[1.2-a]pyridine-8-yl
oxymethyl)-2,4-dichloro-phenyl]-methyl-carbamoyl]-methyl)-
carbamoyl]-vinyl]-N,N-dimethyl-benzamide,
- 20 ▪ 3-(6-Acetylamino-pyridine-3-yl)-N-([2,4-dichloro-3-(2-methyl-
quinoline-8-yl oxymethyl)-phenyl]-methyl-carbamoyl)-methyl)-
acrylamide,
- 1-[2,4-Dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-
benzene sulphonyl]-pyrrolidine-2-carboxylic acid [3-(4-
25 carbamidoyl-benzoylamino)-propyl]-amide (Anatibant)
- 4-(4-[1-[2,4-Dichloro-3-(2,4-dimethyl-quinoline-8-yl
oxymethyl)-benzene sulphonyl]-pyrrolidine-2-carbonyl]-
piperazine-1-carbonyl)-benzamidine, or

15

- 2-[5-(4-Cyano-benzoyl)-1-methyl-1H-pyrrol-2-yl]-N-[2,4-dichloro-3-(2-methyl-quinoline-8-yl oxymethyl)-phenyl]-N-methyl-acetamide, or

- B2 receptor antagonists, having general formula (I)

5



(I)

wherein

- R is hydrogen or methyl
- 10 - W represents a simple bond or an O atom
- $n = 3, 4$
- X is hydrogen or a $-NR_1R_2$ amino group in which R_1 and R_2 can be independently of one another hydrogen or methyl, ethyl n-propyl, or isopropyl,
- 15 - Y is a quaternary ammonium $(-NR_3R_4R_5)^+A^-$ in which R_3, R_4, R_5 can be independently of one another methyl, ethyl n-propyl, isopropyl, butyl, isobutyl, or n-pentyl and A^- is an anion of a pharmaceutically acceptable acid;

the pharmacologically acceptable salts, enantiomers and enantiomeric
20 mixtures thereof.

2. The pharmaceutical composition as claimed in claim 1, wherein the bradykinin antagonist is a bradykinin B2 receptor antagonist which is:

- icatibant, or
- a compound of general formula (I) as defined in claim 1.

5 3. The pharmaceutical composition as claimed in claim 1, wherein the bradykinin antagonist is the compound of general formula (I): (4-(S)-amino-5-(4-{4-[2,4-dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-benzenesulphonylamino]-tetrahydro-pyran-4-carbonyl}-piperazin-1-yl-5-oxo-pentyl]-trimethyl-ammonium in salified form with ions formally deriving from
10 an acid wherein the acid is hydrochloric acid, acetic acid, sulphuric acid, trifluoroacetic acid, methanesulphonic acid, succinic acid or edetic acid.

4. The pharmaceutical composition as claimed in claim 3, wherein the compound (4-(S)-amino-5-(4-{4-[2,4-dichloro-3-(2,4-dimethyl-quinoline-8-yl oxymethyl)-benzenesulphonylamino]-tetrahydro-pyran-4-carbonyl}-piperazin-
15 1-yl-5-oxo-pentyl]-trimethyl-ammonium is in the form of chloride dihydrochloride (MEN16132).

5. The pharmaceutical composition as claimed in any one of claims 1-4, wherein the amount of hyaluronic acid per dose is 1-100 mg.

6. The pharmaceutical composition as claimed in claim 5, wherein the
20 amount of hyaluronic acid per dose is 5-20 mg.

7. The pharmaceutical composition as claimed in any one of claims 1-6, wherein the mean MW of the hyaluronic acid ranges between 4 million and 9 million daltons.

8. The pharmaceutical composition as claimed in claim 7, wherein the
25 mean MW of the hyaluronic acid ranges between 5 million and 8 million daltons.

9. The pharmaceutical composition as claimed in any one of claims 1-4, wherein the amount of bradykinin receptor antagonist is between 5.7×10^{-5} and 2.3×10^{-2} mMoles per dose which, in the case of MEN16132, corresponds to a amount of 0.05 to 20 mg per dose.

5 10. The pharmaceutical composition as claimed in claim 9, wherein the amount of bradykinin receptor antagonist is between 1.1×10^{-4} and 1.1×10^{-2} mMoles per dose which, in the case of MEN16132, corresponds to a amount of 0.1 to 10 mg per dose.

11. The pharmaceutical composition as claimed in claim 10, wherein the
10 amount of bradykinin receptor antagonist is between 2.9×10^{-4} and 5.7×10^{-3} mmols per dose which, in the case of MEN16132, substantially corresponds to an amount of 0.25 to 5 mg per dose.

12. The pharmaceutical composition as claimed in any one of claims 1-11,
in the form of an intra-articular or intrabursal injectable solution, or a
15 transdermal form wherein the transdermal form is a cream, gel or patch.

13. The pharmaceutical composition as claimed in claim 12, wherein the
bradykinin antagonist is in a solid form wherein the solid form is crystalline,
amorphous or lyophilised, to be dissolved before use in a solution containing
hyaluronic acid so as to constitute the intra-articular or intrabursal injectable
20 solution.

14. The pharmaceutical composition as claimed in any one of claims 1-13,
also containing a buffer wherein the buffer is phosphate or citrate.

15. The pharmaceutical composition as claimed in any one of claims 1-14,
also containing sodium chloride as isotonicity regulator.

25 16. The pharmaceutical composition as claimed in any one of claims 1-15,
also containing sodium edetate as preservative and chelating agent.

17. The use of hyaluronic acid in association with a bradykinin B2 receptor antagonist for the preparation of the pharmaceutical composition as claimed in claim 1, for the prophylaxis and treatment of inflammatory, autoimmune, traumatic and degenerative joint diseases and in joint diseases caused by developmental alterations.

18. The use as claimed in claim 17 wherein the degenerative joint diseases are osteoarthritis, post-traumatic osteoarthritis, degenerative osteoarthritis, spondylosis, synovitis, tenosynovitis, bursitis, contusions, sprains, dislocations or subluxations.

19. The use as claimed in claim 17 wherein the joint diseases caused by developmental alterations are osteochondritis or dysplasia.

20. The use as claimed in claim 18 wherein the degenerative osteoarthritis is gonarthrititis or spondylarthrititis.

21. The use, as claimed in any one of claims 17-20, of the bradykinin B2 antagonist Men-16132 in association with hyaluronic acid for the preparation of a pharmaceutical composition for the prophylaxis and treatment of inflammatory, autoimmune, traumatic and degenerative joint diseases and in joint diseases caused by developmental alterations.

22. The use as claimed in claim 21 wherein the degenerative joint diseases are osteoarthritis, post-traumatic osteoarthritis, degenerative osteoarthritis, spondylosis, synovitis, tenosynovitis, bursitis, contusions, sprains, dislocations or subluxations.

23. The use as claimed in claim 21 wherein the joint diseases caused by developmental alterations are osteochondritis or dysplasia.

24. The use as claimed in claim 22 wherein the degenerative osteoarthritis is gonarthrititis or spondylarthrititis.

25. The use, as claimed in any one of claims 21-24, of the bradykinin B2 antagonist MEN16132 in association with hyaluronic acid to prepare a pharmaceutical composition for the treatment of osteoarthritis and post-traumatic osteoarthritis.