



(86) Date de dépôt PCT/PCT Filing Date: 2012/12/21  
(87) Date publication PCT/PCT Publication Date: 2013/06/27  
(45) Date de délivrance/Issue Date: 2021/03/09  
(85) Entrée phase nationale/National Entry: 2014/06/17  
(86) N° demande PCT/PCT Application No.: US 2012/071371  
(87) N° publication PCT/PCT Publication No.: 2013/096837  
(30) Priorité/Priority: 2011/12/22 (US61/579,343)

(51) Cl.Int./Int.Cl. *A61K 31/7088* (2006.01),  
*A61K 31/7115* (2006.01), *A61K 31/712* (2006.01),  
*A61K 31/7125* (2006.01), *A61P 35/00* (2006.01),  
*A61P 35/04* (2006.01), *C12N 15/113* (2010.01),  
*C07H 21/00* (2006.01)

(72) Inventeurs/Inventors:  
BENNETT, C. FRANK, US;  
FREIER, SUSAN M., US;  
MARCUSSON, ERIC G., US;  
HSU, SSUCHENG J., US;  
MACLEOD, ROBERT A., US

(73) Propriétaire/Owner:  
IONIS PHARMACEUTICALS, INC., US

(74) Agent: NORTON ROSE FULBRIGHT CANADA

(54) Titre : PROCEDES POUR LA MODULATION D'UNE EXPRESSION D'UN TRANSCRIT 1 D'ADENOCARCINOME ASSOCIE A LA METASTASE (MALAT-1)

(54) Title: METHODS FOR MODULATING METASTASIS-ASSOCIATED-IN-LUNG-ADENOCARCINOMA-TRANSCRIPT-1(MALAT-1) EXPRESSION

(57) Abrégé/Abstract:

The present embodiments provide compounds and methods for reducing expression of Metastasis- Associated-in-Lung-Adenocarcinoma-Transcript-1 (MALAT-1) RNA and/or protein in an animal. Such methods are useful for treating cancer, such as colon cancer, intestinal cancer, lung cancer (e.g. non-small cell lung cancer), liver cancer, and/or prostate cancer. In various aspects, the cancer is a primary cancer.

(74) **Agent(suite/continued):** LLP/S.E.N.C.R.L., S.R.L.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(10) International Publication Number  
**WO 2013/096837 A1**

(43) International Publication Date  
27 June 2013 (27.06.2013)

## (51) International Patent Classification:

C12N 15/11 (2006.01) C07H 21/02 (2006.01)  
A61K 38/28 (2006.01)

## (21) International Application Number:

PCT/US2012/071371

## (22) International Filing Date:

21 December 2012 (21.12.2012)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

61/579,343 22 December 2011 (22.12.2011) US

(71) Applicant: **ISIS PHARMACEUTICALS, INC.** [US/US];  
2855 Gazelle Court, Carlsbad, CA 92010 (US).

## (72) Inventors; and

(71) Applicants : **BENNETT, C., Frank** [US/US]; 2855  
Gazelle Court, Carlsbad, CA 92010 (US). **FREIER,**  
**Susan, M.** [US/US]; 2855 Gazelle Court, Carlsbad, CA  
92010 (US). **MARCUSSON, Eric, G.** [US/US]; 260 King  
Street #727, San Francisco, CA 94107 (US). **HSU,**  
**Ssucheng, J.** [US/US]; 2855 Gazelle Court, Carlsbad, CA  
92010 (US). **MACLEOD, Robert, A.** [CA/US]; 2855  
Gazelle Court, Carlsbad, CA 92010 (US).

(74) Agents: **GINGRICH, Brenden** et al.; Knobbe Martens,  
2040 Main Street, 14th Floor, Irvine, CA 92614 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,  
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,  
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,  
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,  
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,  
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,  
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

## Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: METHODS FOR MODULATING METASTASIS-ASSOCIATED-IN-LUNG-ADENOCARCINOMA-TRANSCRIPT-1(MALAT-1) EXPRESSION

(57) Abstract: The present embodiments provide compounds and methods for reducing expression of Metastasis- Associated-in-Lung-Adenocarcinoma-Transcript-1 (MALAT-1) RNA and/or protein in an animal. Such methods are useful for treating cancer, such as colon cancer, intestinal cancer, lung cancer (e.g. non-small cell lung cancer), liver cancer, and/or prostate cancer. In various aspects, the cancer is a primary cancer.



WO 2013/096837 A1

# METHODS FOR MODULATING METASTASIS-ASSOCIATED-IN-LUNG- ADENOCARCINOMA-TRANSCRIPT-1 (MALAT-1) EXPRESSION

## SEQUENCE LISTING

The present application is being filed along with a Sequence Listing in electronic format. The  
5 Sequence Listing is provided as a file entitled BIOL0181WOSEQ.txt created December 20, 2012, which is  
111 Kb in size and forms part of this description.

## FIELD

The present embodiments relate to the field of cancer biology. More particularly, embodiments  
provided herein are drawn to compounds and methods for reducing expression of Metastasis-Associated-in-  
10 Lung-Adenocarcinoma-Transcript-1 (MALAT-1) RNA and/or protein in an animal. Such methods are  
useful to treat cancer, such as colon cancer, intestinal cancer, lung cancer (e.g. non-small cell lung cancer),  
liver cancer, and/or prostate cancer. In various aspects, the cancer is a primary cancer.

## BACKGROUND

MALAT, also known as noncoding nuclear-enriched abundant transcript 2 (NEAT2) is a large,  
15 infrequently spliced non-coding RNA that is highly conserved amongst mammals. MALAT-1 is expressed  
in the nucleus and positively regulates cell motility by transcriptional and/or post-transcriptional regulation  
of motility-related genes. Additionally, MALAT-1 has been implicated in the regulation of alternative  
splicing. However, the functional role of MALAT-1 in carcinogenesis is largely unknown.

## SUMMARY

20 Embodiments provided herein relate to the discovery that MALAT-1 specific inhibitors can treat  
cancer *in vivo*. Several embodiments are drawn to MALAT-1 specific inhibitors, such as antisense  
compounds, and methods for modulating expression of MALAT-1 RNA and protein using the same. In  
certain embodiments, MALAT-1 specific inhibitors modulate MALAT-1 RNA and/or protein expression or  
activity.

25 Also provided are methods of treating cancer with MALAT-1 specific inhibitors, such as antisense  
compounds. In some embodiments, methods of treating cancer in an animal include administering to the  
animal an antisense compound which reduces expression of MALAT-1. Types of cancers that can be  
treated with the MALAT-1 specific inhibitors provided herein include but are not limited to colon cancer,

intestinal cancer, lung cancer (e.g. non-small cell lung cancer), liver cancer, and/or prostate cancer. In various aspects, the cancer is a primary cancer.

In several embodiments, a method of treating cancer in an animal includes administering to the animal an antisense compound which reduces expression of MALAT-1. In one aspect, the antisense  
5 compound comprises a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 85% complementary to a MALAT-1 nucleic acid.

Several embodiments are directed to the use of a compound including a modified oligonucleotide consisting of 12 to 30 linked nucleosides at least 85% complementary to a MALAT-1 nucleic acid in the manufacture of a medicament for treating cancer.

10 Further embodiments relate to compounds for use in the treatment of cancer including a modified oligonucleotide consisting of 12 to 30 linked nucleosides at least 85% complementary to a MALAT-1 nucleic acid.

In various aspects of any of the aforementioned embodiments, expression of MALAT-1 RNA is reduced; expression of MALAT-1 protein is reduced; the animal is a human; the MALAT-1 nucleic acid is a  
15 human MALAT-1 nucleic acid (e.g. any one of SEQ ID NOs:1-9); the modified oligonucleotide is 100% complementary to a human MALAT-1 nucleic acid (e.g. any one of SEQ ID NOs:1-9); the modified oligonucleotide inhibits cancer growth and/or metastasis; the modified oligonucleotide increases survival of the animal; the cancer is colon cancer, intestinal cancer, lung cancer, liver cancer, or prostate cancer; the cancer is a primary cancer; the expression of MALAT-1 is reduced in cancer cells of the animal compared  
20 to control or untreated animals; the modified oligonucleotide is a single-stranded oligonucleotide; the modified oligonucleotide comprises at least one modified internucleoside linkage such as a phosphorothioate internucleoside linkage; at least one nucleoside comprises a modified sugar; the modified sugar is a bicyclic sugar such as a 4'-CH(CH<sub>3</sub>)-O-2' bridge; the modified oligonucleotide includes at least one tetrahydropyran modified nucleoside wherein a tetrahydropyran ring replaces the furanose ring; the  
25 modified sugar comprises a 2'-O-methoxyethyl group; and/or at least one nucleoside comprises a modified nucleobase such as a 5-methylcytosine.

#### DETAILED DESCRIPTION

It is to be understood that both the foregoing general description and the following detailed description are explanatory only and are not restrictive of the invention as claimed. Herein, the use of the  
30 singular includes the plural unless specifically stated otherwise. As used herein, the use of "or" means "and/or" unless stated otherwise. Additionally, as used herein, the use of "and" means "and/or" unless stated otherwise. Furthermore, the use of the term "including" as well as other forms, such as "includes"

and “included”, is not limiting.

### ***Definitions***

5 Unless specific definitions are provided, the nomenclature utilized in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, and chemical analysis.

Unless otherwise indicated, the following terms have the following meanings:

10 “2’-O-methoxyethyl” (also 2’-MOE and 2’-O(CH<sub>2</sub>)<sub>2</sub>-OCH<sub>3</sub>) refers to an O-methoxy-ethyl modification of the 2’ position of a furanosyl ring. A 2’-O-methoxyethyl modified sugar is a modified sugar.

“2’-MOE nucleoside” (also 2’-O-methoxyethyl nucleoside) means a nucleoside comprising a 2’-MOE modified sugar moiety.

15 “5-methylcytosine” means a cytosine modified with a methyl group attached to the 5’ position. A 5-methylcytosine is a modified nucleobase.

“About” means within  $\pm 7\%$  of a value. For example, if it is stated, “the compounds inhibited MALAT-1 by about 70%”, it is implied that the MALAT-1 levels are inhibited within a range of 63% and 77%.

20 “Active target region” or “target region” means a region to which one or more active antisense compounds is targeted. “Active antisense compounds” means antisense compounds that reduce target nucleic acid levels or protein levels.

“Administering” means providing a pharmaceutical agent to an individual, and includes, but is not limited to administering by a medical professional and self-administering.

25 “Amelioration” or “ameliorate” or “ameliorating” refers to a lessening of at least one indicator, sign, or symptom of an associated disease, disorder, or condition. The severity of indicators may be determined by subjective or objective measures, which are known to those skilled in the art.

“Animal” refers to a human or non-human animal, including, but not limited to, mice, rats, rabbits, dogs, cats, pigs, and non-human primates, including, but not limited to, monkeys and chimpanzees.

“Antisense activity” means any detectable or measurable activity attributable to the hybridization of an antisense compound to its target nucleic acid. In certain embodiments, antisense activity is a decrease in the amount or expression of a target nucleic acid or protein encoded by such target nucleic acid.

“Antisense compound” means an oligomeric compound that is capable of undergoing hybridization to a target nucleic acid through hydrogen bonding. Examples of antisense compounds include single-stranded and double-stranded compounds, such as, antisense oligonucleotides, siRNAs, shRNAs, and miRNAs.

“Antisense inhibition” means reduction of target nucleic acid levels or target protein levels in the presence of an antisense compound complementary to a target nucleic acid compared to target nucleic acid levels or target protein levels in the absence of the antisense compound.

“Antisense oligonucleotide” means a single-stranded oligonucleotide having a nucleobase sequence that permits hybridization to a corresponding region or segment of a target nucleic acid.

“Bicyclic sugar” means a furanosyl ring modified by the bridging of two atoms. A bicyclic sugar is a modified sugar.

“Bicyclic nucleoside” (also BNA) means a nucleoside having a sugar moiety comprising a bridge connecting two carbon atoms of the sugar ring, thereby forming a bicyclic ring system. In certain embodiments, the bridge connects the 4'-carbon and the 2'-carbon of the sugar ring.

“Cap structure” or “terminal cap moiety” means chemical modifications, which have been incorporated at either terminus of an antisense compound.

“cEt” or “constrained ethyl” means a bicyclic nucleoside having a sugar moiety comprising a bridge connecting the 4'-carbon and the 2'-carbon, wherein the bridge has the formula: 4'-CH(CH<sub>3</sub>)-O-2'.

“Constrained ethyl nucleoside” (also cEt nucleoside) means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH(CH<sub>3</sub>)-O-2' bridge.

“Chemically distinct region” refers to a region of an antisense compound that is in some way chemically different than another region of the same antisense compound. For example, a region having 2'-O-methoxyethyl nucleotides is chemically distinct from a region having nucleotides without 2'-O-methoxyethyl modifications.

“Chimeric antisense compound” means an antisense compound that has at least two chemically distinct regions.

“Complementarity” means the capacity for pairing between nucleobases of a first nucleic acid and a second nucleic acid.

5 “Contiguous nucleobases” means nucleobases immediately adjacent to each other.

“Diluent” means an ingredient in a composition that lacks pharmacological activity, but is pharmaceutically necessary or desirable. For example, the diluent in an injected composition may be a liquid, e.g. saline solution.

10 “Effective amount” means the amount of active pharmaceutical agent sufficient to effectuate a desired physiological outcome in an individual in need of the agent. The effective amount may vary among individuals depending on the health and physical condition of the individual to be treated, the taxonomic group of the individuals to be treated, the formulation of the composition, assessment of the individual’s medical condition, and other relevant factors.

15 “Metastasis-Associated-in-Lung-Adenocarcinoma-Transcript-1 (MALAT-1)” means any nucleic acid or protein of MALAT-1. “MALAT-1 nucleic acid” means any nucleic acid encoding MALAT-1. For example, in certain embodiments, a MALAT-1 nucleic acid includes a DNA sequence encoding MALAT-1, an RNA sequence transcribed from DNA encoding MALAT-1 (including genomic DNA comprising introns and exons), including a non-protein encoding (*i.e.* non-coding) RNA sequence, and an mRNA sequence encoding MALAT-1. “MALAT-1 mRNA” means an mRNA encoding a MALAT-1 protein.

20 “MALAT-1 specific inhibitor” refers to any agent capable of specifically inhibiting MALAT-1 RNA and/or MALAT-1 protein expression or activity at the molecular level. For example, MALAT-1 specific inhibitors include nucleic acids (including antisense compounds), peptides, antibodies, small molecules, and other agents capable of inhibiting the expression of MALAT-1 RNA and/or MALAT-1 protein.

25 “Fully complementary” or “100% complementary” means each nucleobase of a first nucleic acid has a complementary nucleobase in a second nucleic acid. In certain embodiments, a first nucleic acid is an antisense compound and a target nucleic acid is a second nucleic acid.

30 “Gapmer” means a chimeric antisense compound in which an internal region having a plurality of nucleosides that support RNase H cleavage is positioned between external regions having one or more nucleosides, wherein the nucleosides comprising the internal region are chemically distinct from the



nucleoside or nucleosides comprising the external regions. The internal region may be referred to as a “gap” and the external regions may be referred to as the “wings.”

“Gap-widened” means a chimeric antisense compound having a gap segment of 12 or more contiguous 2'-deoxyribonucleosides positioned between and immediately adjacent to 5' and 3' wing segments having from one to six nucleosides.

“Hybridization” means the annealing of complementary nucleic acid molecules. In certain embodiments, complementary nucleic acid molecules include an antisense compound and a target nucleic acid.

“Immediately adjacent” means there are no intervening elements between the immediately adjacent elements.

“Individual” means a human or non-human animal selected for treatment or therapy.

“Inhibiting MALAT-1” means reducing expression of MALAT-1 RNA and/or protein levels in the presence of a MALAT-1 specific inhibitor, including a MALAT-1 antisense oligonucleotide, as compared to expression of MALAT-1 RNA and/or protein levels in the absence of a MALAT-1 specific inhibitor, such as a MALAT-1 antisense oligonucleotide.

“Internucleoside linkage” refers to the chemical bond between nucleosides.

“Linked nucleosides” means adjacent nucleosides which are bonded together.

“Mismatch” or “non-complementary nucleobase” refers to the case when a nucleobase of a first nucleic acid is not capable of pairing with the corresponding nucleobase of a second or target nucleic acid.

“Modified internucleoside linkage” refers to a substitution or any change from a naturally occurring internucleoside bond (i.e. a phosphodiester internucleoside bond).

“Modified nucleobase” refers to any nucleobase other than adenine, cytosine, guanine, thymidine, or uracil. An “unmodified nucleobase” means the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C), and uracil (U).

“Modified nucleotide” means a nucleotide having, independently, a modified sugar moiety, modified internucleoside linkage, or modified nucleobase. A “modified nucleoside” means a nucleoside having, independently, a modified sugar moiety or modified nucleobase.

“Modified oligonucleotide” means an oligonucleotide comprising a modified internucleoside linkage, a modified sugar, or a modified nucleobase.

“Modified sugar” refers to a substitution or change from a natural sugar.

“Motif” means the pattern of chemically distinct regions in an antisense compound.

“Naturally occurring internucleoside linkage” means a 3' to 5' phosphodiester linkage.

“Natural sugar moiety” means a sugar found in DNA (2'-H) or RNA (2'-OH).

5 “Nucleic acid” refers to molecules composed of monomeric nucleotides. A nucleic acid includes ribonucleic acids (RNA), deoxyribonucleic acids (DNA), single-stranded nucleic acids, double-stranded nucleic acids, small interfering ribonucleic acids (siRNA), and microRNAs (miRNA).

“Nucleobase” means a heterocyclic moiety capable of pairing with a base of another nucleic acid.

10 “Nucleobase sequence” means the order of contiguous nucleobases independent of any sugar, linkage, or nucleobase modification.

“Nucleoside” means a nucleobase linked to a sugar.

15 “Nucleoside mimetic” includes those structures used to replace the sugar or the sugar and the base and not necessarily the linkage at one or more positions of an oligomeric compound such as for example nucleoside mimetics having morpholino, cyclohexenyl, cyclohexyl, tetrahydropyranyl, bicyclo, or tricyclo sugar mimetics, e.g., non furanose sugar units. Nucleotide mimetic includes those structures used to replace the nucleoside and the linkage at one or more positions of an oligomeric compound such as for example peptide nucleic acids or morpholinos (morpholinos linked by -N(H)-C(=O)-O- or other non-phosphodiester linkage). Sugar surrogate overlaps with the slightly broader term nucleoside mimetic but is intended to indicate replacement of the sugar unit (furanose ring) only. The tetrahydropyranyl rings provided herein are  
20 illustrative of an example of a sugar surrogate wherein the furanose sugar group has been replaced with a tetrahydropyranyl ring system.

“Nucleotide” means a nucleoside having a phosphate group covalently linked to the sugar portion of the nucleoside.

25 “Oligomeric compound” or “oligomer” means a polymer of linked monomeric subunits which is capable of hybridizing to at least a region of a nucleic acid molecule.

“Oligonucleotide” means a polymer of linked nucleosides each of which can be modified or unmodified, independent one from another.

“Parenteral administration” means administration through injection or infusion. Parenteral administration includes subcutaneous administration, intravenous administration, intramuscular

administration, intraarterial administration, intraperitoneal administration, or intracranial administration, e.g., intrathecal or intracerebroventricular administration.

“Peptide” means a molecule formed by linking at least two amino acids by amide bonds. Peptide refers to polypeptides and proteins.

5 “Pharmaceutical composition” means a mixture of substances suitable for administering to an individual. For example, a pharmaceutical composition may comprise one or more active pharmaceutical agents and a sterile aqueous solution.

“Phosphorothioate linkage” means a linkage between nucleosides where the phosphodiester bond is modified by replacing one of the non-bridging oxygen atoms with a sulfur atom. A phosphorothioate  
10 linkage (P=S) is a modified internucleoside linkage.

“Portion” means a defined number of contiguous (i.e., linked) nucleobases of a nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of a target nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of an antisense compound.

“Single-stranded oligonucleotide” means an oligonucleotide which is not hybridized to a  
15 complementary strand.

“Specifically hybridizable” refers to an antisense compound having a sufficient degree of complementarity between an antisense oligonucleotide and a target nucleic acid to induce a desired effect, while exhibiting minimal or no effects on non-target nucleic acids under conditions in which specific binding is desired, i.e., under physiological conditions in the case of *in vivo* assays and therapeutic  
20 treatments.

“Targeting” or “targeted” means the process of design and selection of an antisense compound that will specifically hybridize to a target nucleic acid and induce a desired effect.

“Target nucleic acid,” “target RNA,” and “target RNA transcript” all refer to a nucleic acid capable of being targeted by antisense compounds.

25 “Target segment” means the sequence of nucleotides of a target nucleic acid to which an antisense compound is targeted. “5’ target site” refers to the 5’-most nucleotide of a target segment. “3’ target site” refers to the 3’-most nucleotide of a target segment.

“Therapeutically effective amount” means an amount of a pharmaceutical agent that provides a therapeutic benefit to an individual.

“Treat” or “treating” refers to administering a pharmaceutical composition to effect an alteration or improvement of a disease, disorder, or condition.

“Unmodified nucleotide” means a nucleotide composed of naturally occurring nucleobases, sugar moieties, and internucleoside linkages. In certain embodiments, an unmodified nucleotide is an RNA nucleotide (i.e.  $\beta$ -D-ribonucleosides) or a DNA nucleotide (i.e.  $\beta$ -D-deoxyribonucleoside).

### *Certain Embodiments*

Certain embodiments provided herein relate to methods for decreasing MALAT-1 RNA and/or protein expression in an animal.

Certain embodiments provide methods for the treatment or amelioration of diseases, disorders, and conditions associated with MALAT-1, such as cancer, in an animal in need thereof. In several embodiments, the cancer can be colon cancer, lung cancer (e.g. non-small cell lung cancer), liver cancer, and/or prostate cancer.

Certain embodiments provide for the use of a MALAT-1 specific inhibitor for treating cancer in an animal by administering a MALAT-1 specific inhibitor, such as nucleic acids (including antisense compounds) capable of reducing the levels of MALAT-1 RNA and/or MALAT-1 protein.

Certain embodiments provide for methods of treating cancer in an animal, comprising administering to the animal a therapeutically effective amount of a MALAT-1 specific inhibitor. In certain embodiments, the animal is a human.

In certain embodiments, the MALAT-1 specific inhibitor is an antisense compound. In certain embodiments, the antisense compound is a modified oligonucleotide.

In certain embodiments, the MALAT-1 specific inhibitor is a nucleic acid. In certain embodiments, the nucleic acid is a modified oligonucleotide.

In certain embodiments, the MALAT-1 specific inhibitor is a modified oligonucleotide.

In certain embodiments, the modified oligonucleotide consists of 12 to 30 linked nucleosides.

In certain embodiments, the modified oligonucleotide is a single-stranded oligonucleotide.

In certain embodiments, the modified oligonucleotide consists of 15, 16, 17, 18, 19, or 20 linked nucleosides.

In certain embodiments, the modified oligonucleotide has a nucleobase sequence that is 80%, 85%, 90%, 95%, or 100% complementary to a human MALAT-1 nucleic acid.


In certain embodiments, the modified oligonucleotide comprises at least one modified internucleoside linkage. In certain embodiments, each modified internucleoside linkage is a phosphorothioate internucleoside linkage.

In certain embodiments, at least one nucleoside of the modified oligonucleotide comprises a modified sugar. In certain embodiments, the modified sugar is a bicyclic sugar. In certain embodiments, the bicyclic sugar comprises a 4'-CH(CH<sub>3</sub>)-O-2' bridge.

In certain embodiments, the modified sugar comprises a 2'-O-methoxyethyl group.

In certain embodiments, at least one nucleoside of the modified oligonucleotide comprises a modified nucleobase. In certain embodiments, the modified nucleobase is a 5'-methylcytosine.

In certain embodiments, at least one nucleoside of the modified oligonucleotide comprises at least one tetrahydropyran modified nucleoside wherein a tetrahydropyran ring replaces the furanose ring. In certain embodiments, each of the at least one tetrahydropyran modified nucleoside has the structure:



wherein Bx is an optionally protected heterocyclic base moiety.

In certain embodiments, the modified oligonucleotide of the compound comprises:

- (i) a gap segment consisting of linked deoxynucleosides;
- (ii) a 5' wing segment consisting of linked nucleosides;
- (iii) a 3' wing segment consisting of linked nucleosides, wherein the gap segment is positioned immediately adjacent to and between the 5' wing segment and the 3' wing segment and wherein each nucleoside of each wing segment comprises a modified sugar. In some such embodiments, each cytosine in the modified oligonucleotide is a 5-methylcytosine.

In certain embodiments, the modified oligonucleotide of the compound comprises:

- (i) a gap segment consisting of ten linked deoxynucleosides;
- (ii) a 5' wing segment consisting of five linked nucleosides;

(iii) a 3' wing segment consisting of five linked nucleosides, wherein the gap segment is positioned immediately adjacent to and between the 5' wing segment and the 3' wing segment, wherein each nucleoside of each wing segment comprises a 2'-O-methoxyethyl sugar; and wherein each internucleoside linkage is a phosphorothioate linkage. In some such embodiments, each cytosine in the modified  
 5 oligonucleotide is a 5-methylcytosine.

Several embodiments described herein provide for methods comprising administering to an animal having cancer a therapeutically effective amount of a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 80% complementary to a human MALAT-1 nucleic acid. In certain embodiments, the modified oligonucleotide is at least 90% complementary to a  
 10 human MALAT-1 nucleic acid. In certain embodiments, the modified oligonucleotide is 100% complementary to a human MALAT-1 nucleic acid.

In certain embodiments, the modified oligonucleotide targets a human MALAT-1 nucleic acid which may be selected from, but not limited to, one or more of GENBANK Accession No. EF177381.1 (incorporated herein as SEQ ID NO: 1), GENBANK Accession No. BK001411.1 (incorporated herein as  
 15 SEQ ID NO: 2), GENBANK Accession No. BQ429080.1 (incorporated herein as SEQ ID NO: 3), GENBANK Accession No. BQ428957.1 (incorporated herein as SEQ ID NO: 4), GENBANK Accession No. NT\_033903.7 truncated from nucleobases 10569000 to 10582000 (incorporated herein as SEQ ID NO: 5), GENBANK Accession No. XR\_001309.1 (incorporated herein as SEQ ID NO: 6), or GENBANK Accession No. NR\_002819.2 (incorporated herein as SEQ ID NO: 7), GENBANK Accession No.  
 20 NC\_000011.9 from nucleobases 65265233 to 65273940 (incorporated herein as SEQ ID NO: 8) or the complement thereof, and GENBANK Accession No. AC\_000143.1 from nucleobases 61592326 to 61601033 (incorporated herein as SEQ ID NO: 9) or the complement thereof.

In certain embodiments, the modified oligonucleotide targets a mouse MALAT-1 nucleic acid which may be selected from, but not limited to, one or more of GENBANK Accession No. NR\_002847.2 (incorporated herein as SEQ ID NO: 10), GENBANK Accession No. FJ209304.1 (incorporated herein as  
 25 SEQ ID NO: 11), and the complement of GENBANK Accession No. NT\_082868.4 truncated from nucleobases 2689000 to 2699000 (incorporated herein as SEQ ID NO: 12).

In certain embodiments, antisense compounds may comprise a modified oligonucleotide comprising a nucleobase sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at  
 30 least 98%, or at least 99% complementary to an equal length portion of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12.

In certain embodiments, antisense compounds may comprise a modified oligonucleotide comprising a nucleobase sequence 100% complementary to an equal length portion of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12.

5 In certain embodiments, the nucleobase sequence of the modified oligonucleotide is 100% complementary to a nucleobase sequence of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, or SEQ ID NO:12.

### *Antisense Compounds*

10 Antisense compounds provided herein refer to oligomeric compounds capable of undergoing hybridization to a target nucleic acid through hydrogen bonding. Examples of antisense compounds include single-stranded and double-stranded compounds, such as, antisense oligonucleotides, siRNAs, shRNAs, and miRNAs.

15 In certain embodiments, an antisense compound has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted. In certain such embodiments, an antisense oligonucleotide has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted.

20 In certain embodiments, an antisense compound targeted to a MALAT-1 nucleic acid is 12 to 30 subunits in length. In other words, such antisense compounds are from 12 to 30 linked subunits. In other embodiments, the antisense compound is 8 to 80, 12 to 50, 15 to 30, 18 to 24, 19 to 22, or 20 linked subunits. In certain such embodiments, the antisense compounds are 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 25 77, 78, 79, or 80 linked subunits in length, or a range defined by any two of the above values. In some embodiments the antisense compound is an antisense oligonucleotide, and the linked subunits are nucleosides.

30 In several embodiments an antisense compound targeted to a MALAT-1 nucleic acid can have antisense portions of 10 to 50 nucleobases in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleobases in length, or any range therewithin.

In several embodiments, an antisense compound targeted to a MALAT-1 nucleic acid can have antisense portions of 12 to 30 nucleobases in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleobases in length, or any range therewithin.

5 In some embodiments, an antisense compound targeted to a MALAT-1 nucleic acid can have antisense portions of 12 or 13 to 24 nucleobases in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 or 24 nucleobases in length, or any range therewithin.

10 In some embodiments, an antisense compound targeted to a MALAT-1 nucleic acid can have antisense portions of 19 to 23 nucleobases in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 19, 20, 21, 22 or 23 nucleobases in length, or any range therewithin.

15 In certain embodiments antisense compounds targeted to a MALAT-1 nucleic acid may be shortened or truncated. For example, a single subunit may be deleted from the 5' end (5' truncation), or alternatively from the 3' end (3' truncation). A shortened or truncated antisense compound targeted to a MALAT-1 nucleic acid may have two subunits deleted from the 5' end, or alternatively may have two subunits deleted from the 3' end, of the antisense compound. Alternatively, the deleted nucleosides may be dispersed throughout the antisense compound, for example, in an antisense compound having one nucleoside deleted from the 5' end and one nucleoside deleted from the 3' end.

20 When a single additional subunit is present in a lengthened antisense compound, the additional subunit may be located at the 5' or 3' end of the antisense compound. When two or more additional subunits are present, the added subunits may be adjacent to each other, for example, in an antisense compound having two subunits added to the 5' end (5' addition), or alternatively to the 3' end (3' addition), of the antisense compound. Alternatively, the added subunits may be dispersed throughout the antisense  
25 compound, for example, in an antisense compound having one subunit added to the 5' end and one subunit added to the 3' end.

It is possible to increase or decrease the length of an antisense compound, such as an antisense oligonucleotide, and/or introduce mismatch bases without eliminating activity. For example, in Woolf et al. (Proc. Natl. Acad. Sci. USA 89:7305-7309, 1992), a series of antisense oligonucleotides 13-25 nucleobases  
30 in length were tested for their ability to induce cleavage of a target RNA in an oocyte injection model. Antisense oligonucleotides 25 nucleobases in length with 8 or 11 mismatch bases near the ends of the antisense oligonucleotides were able to direct specific cleavage of the target mRNA, albeit to a lesser extent



than the antisense oligonucleotides that contained no mismatches. Similarly, target specific cleavage was achieved using 13 nucleobase antisense oligonucleotides, including those with 1 or 3 mismatches.

Gautschi et al (J. Natl. Cancer Inst. 93:463-471, March 2001) demonstrated the ability of an oligonucleotide having 100% complementarity to the bcl-2 mRNA and having 3 mismatches to the bcl-xL mRNA to reduce the expression of both bcl-2 and bcl-xL *in vitro* and *in vivo*. Furthermore, this oligonucleotide demonstrated potent anti-tumor activity *in vivo*.

10 Maher and Dolnick (Nuc. Acid. Res. 16:3341-3358,1988) tested a series of tandem 14 nucleobase antisense oligonucleotides, and a 28 and 42 nucleobase antisense oligonucleotides comprised of the sequence of two or three of the tandem antisense oligonucleotides, respectively, for their ability to arrest translation of human DHFR in a rabbit reticulocyte assay. Each of the three 14 nucleobase antisense oligonucleotides alone was able to inhibit translation, albeit at a more modest level than the 28 or 42 nucleobase antisense oligonucleotides.

#### *Certain Antisense Compound Motifs and Mechanisms*

15 In certain embodiments, antisense compounds have chemically modified subunits arranged in patterns, or motifs, to confer to the antisense compounds properties such as enhanced inhibitory activity, increased binding affinity for a target nucleic acid, or resistance to degradation by *in vivo* nucleases. Chimeric antisense compounds typically contain at least one region modified so as to confer increased resistance to nuclease degradation, increased cellular uptake, increased binding affinity for the target nucleic acid, and/or increased inhibitory activity. A second region of a chimeric antisense compound may confer another desired property e.g., serve as a substrate for the cellular endonuclease RNase H, which cleaves the RNA strand of an RNA:DNA duplex.

25 Antisense activity may result from any mechanism involving the hybridization of the antisense compound (e.g., oligonucleotide) with a target nucleic acid, wherein the hybridization ultimately results in a biological effect. In certain embodiments, the amount and/or activity of the target nucleic acid is modulated. In certain embodiments, the amount and/or activity of the target nucleic acid is reduced. In certain embodiments, hybridization of the antisense compound to the target nucleic acid ultimately results in target nucleic acid degradation. In certain embodiments, hybridization of the antisense compound to the target nucleic acid does not result in target nucleic acid degradation. In certain such embodiments, the presence of the antisense compound hybridized with the target nucleic acid (occupancy) results in a modulation of antisense activity. In certain embodiments, antisense compounds having a particular chemical motif or pattern of chemical modifications are particularly suited to exploit one or more mechanisms. In certain 30 embodiments, antisense compounds function through more than one mechanism and/or through mechanisms

that have not been elucidated. Accordingly, the antisense compounds described herein are not limited by particular mechanism.

Antisense mechanisms include, without limitation, RNase H mediated antisense; RNAi mechanisms, which utilize the RISC pathway and include, without limitation, siRNA, ssRNA and microRNA mechanisms; and occupancy based mechanisms. Certain antisense compounds may act through more than one such mechanism and/or through additional mechanisms.

#### *RNase H-Mediated Antisense*

In certain embodiments, antisense activity results at least in part from degradation of target RNA by RNase H. RNase H is a cellular endonuclease that cleaves the RNA strand of an RNA:DNA duplex. It is known in the art that single-stranded antisense compounds which are "DNA-like" elicit RNase H activity in mammalian cells. Accordingly, antisense compounds comprising at least a portion of DNA or DNA-like nucleosides may activate RNase H, resulting in cleavage of the target nucleic acid. In certain embodiments, antisense compounds that utilize RNase H comprise one or more modified nucleosides. In certain embodiments, such antisense compounds comprise at least one block of 1-8 modified nucleosides. In certain such embodiments, the modified nucleosides do not support RNase H activity. In certain embodiments, such antisense compounds are gapmers, as described herein. In certain such embodiments, the gap of the gapmer comprises DNA nucleosides. In certain such embodiments, the gap of the gapmer comprises DNA-like nucleosides. In certain such embodiments, the gap of the gapmer comprises DNA nucleosides and DNA-like nucleosides.

Certain antisense compounds having a gapmer motif are considered chimeric antisense compounds. In a gapmer an internal region having a plurality of nucleotides that supports RNaseH cleavage is positioned between external regions having a plurality of nucleotides that are chemically distinct from the nucleosides of the internal region. In the case of an antisense oligonucleotide having a gapmer motif, the gap segment generally serves as the substrate for endonuclease cleavage, while the wing segments comprise modified nucleosides. In certain embodiments, the regions of a gapmer are differentiated by the types of sugar moieties comprising each distinct region. The types of sugar moieties that are used to differentiate the regions of a gapmer may in some embodiments include  $\beta$ -D-ribonucleosides,  $\beta$ -D-deoxyribonucleosides, 2'-modified nucleosides (such 2'-modified nucleosides may include 2'-MOE and 2'-O-CH<sub>3</sub>, among others), and bicyclic sugar modified nucleosides (such bicyclic sugar modified nucleosides may include those having a constrained ethyl). In certain embodiments, nucleosides in the wings may include several modified sugar moieties, including, for example 2'-MOE and bicyclic sugar moieties such as constrained ethyl or LNA. In certain embodiments, wings may include several modified and unmodified sugar moieties. In certain embodiments, wings may include various combinations of 2'-MOE nucleosides, bicyclic sugar moieties such as constrained ethyl nucleosides or LNA nucleosides, and 2'-deoxynucleosides.

Each distinct region may comprise uniform sugar moieties, variant, or alternating sugar moieties. The wing-gap-wing motif is frequently described as “X-Y-Z”, where “X” represents the length of the 5'-wing, “Y” represents the length of the gap, and “Z” represents the length of the 3'-wing. “X” and “Z” may comprise uniform, variant, or alternating sugar moieties. In certain embodiments, “X” and “Y” may include one or more 2'-deoxynucleosides. “Y” may comprise 2'-deoxynucleosides. As used herein, a gapmer described as “X-Y-Z” has a configuration such that the gap is positioned immediately adjacent to each of the 5'-wing and the 3' wing. Thus, no intervening nucleotides exist between the 5'-wing and gap, or the gap and the 3'-wing. Any of the antisense compounds described herein can have a gapmer motif. In certain embodiments, “X” and “Z” are the same; in other embodiments they are different. In certain embodiments, “Y” is between 8 and 15 nucleosides. X, Y, or Z can be any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30 or more nucleosides.

In certain embodiments, the antisense compound targeted to a MALAT-1 nucleic acid has a gapmer motif in which the gap consists of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 linked nucleosides.

In certain embodiments, the antisense oligonucleotide has a sugar motif described by Formula A as follows:  $(J)_m-(B)_n-(J)_p-(B)_r-(A)_t-(D)_g-(A)_v-(B)_w-(J)_x-(B)_y-(J)_z$

wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;

m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; g is 6-14;

provided that:

at least one of m, n, and r is other than 0;

at least one of w and y is other than 0;

the sum of m, n, p, r, and t is from 2 to 5; and

the sum of v, w, x, y, and z is from 2 to 5.


### *RNAi Compounds*

In certain embodiments, antisense compounds are interfering RNA compounds (RNAi), which include double-stranded RNA compounds (also referred to as short-interfering RNA or siRNA) and single-stranded RNAi compounds (or ssRNA). Such compounds work at least in part through the RISC pathway to degrade and/or sequester a target nucleic acid (thus, include microRNA/microRNA-mimic compounds). In certain embodiments, antisense compounds comprise modifications that make them particularly suited for such mechanisms.

*i. ssRNA compounds*

In certain embodiments, antisense compounds including those particularly suited for use as single-stranded RNAi compounds (ssRNA) comprise a modified 5'-terminal end. In certain such embodiments, the 5'-terminal end comprises a modified phosphate moiety. In certain embodiments, such modified phosphate is stabilized (e.g., resistant to degradation/cleavage compared to unmodified 5'-phosphate). In certain embodiments, such 5'-terminal nucleosides stabilize the 5'-phosphorous moiety. Certain modified 5'-terminal nucleosides may be found in the art, for example in WO/2011/139702.

In certain embodiments, the 5'-nucleoside of an ssRNA compound has Formula IIc:




IIc

wherein:

T<sub>1</sub> is an optionally protected phosphorus moiety;

T<sub>2</sub> is an internucleoside linking group linking the compound of Formula IIc to the oligomeric compound;

A has one of the formulas:



Q<sub>1</sub> and Q<sub>2</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, substituted C<sub>2</sub>-C<sub>6</sub> alkynyl or N(R<sub>3</sub>)(R<sub>4</sub>);

Q<sub>3</sub> is O, S, N(R<sub>5</sub>) or C(R<sub>6</sub>)(R<sub>7</sub>);

each R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> is, independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl or C<sub>1</sub>-C<sub>6</sub> alkoxy;

M<sub>3</sub> is O, S, NR<sub>14</sub>, C(R<sub>15</sub>)(R<sub>16</sub>), C(R<sub>15</sub>)(R<sub>16</sub>)C(R<sub>17</sub>)(R<sub>18</sub>), C(R<sub>15</sub>)=C(R<sub>17</sub>), OC(R<sub>15</sub>)(R<sub>16</sub>) or OC(R<sub>15</sub>)(Bx<sub>2</sub>);

R<sub>14</sub> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub> and R<sub>18</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

Bx<sub>1</sub> is a heterocyclic base moiety;

or if Bx<sub>2</sub> is present then Bx<sub>2</sub> is a heterocyclic base moiety and Bx<sub>1</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

J<sub>4</sub>, J<sub>5</sub>, J<sub>6</sub> and J<sub>7</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

or J<sub>4</sub> forms a bridge with one of J<sub>5</sub> or J<sub>7</sub> wherein said bridge comprises from 1 to 3 linked biradical groups selected from O, S, NR<sub>19</sub>, C(R<sub>20</sub>)(R<sub>21</sub>), C(R<sub>20</sub>)=C(R<sub>21</sub>), C[=C(R<sub>20</sub>)(R<sub>21</sub>)] and C(=O) and the other two of J<sub>5</sub>, J<sub>6</sub> and J<sub>7</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

each R<sub>19</sub>, R<sub>20</sub> and R<sub>21</sub> is, independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, substituted C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl;

G is H, OH, halogen or O-[C(R<sub>8</sub>)(R<sub>9</sub>)]<sub>n</sub>-(C=O)<sub>m</sub>-X<sub>1</sub>]-Z;

each R<sub>8</sub> and R<sub>9</sub> is, independently, H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl or substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

X<sub>1</sub> is O, S or N(E<sub>1</sub>);

Z is H, halogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, substituted C<sub>2</sub>-C<sub>6</sub> alkynyl or N(E<sub>2</sub>)(E<sub>3</sub>);

E<sub>1</sub>, E<sub>2</sub> and E<sub>3</sub> are each, independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl or substituted C<sub>1</sub>-C<sub>6</sub> alkyl;

n is from 1 to about 6;

m is 0 or 1;

j is 0 or 1;

each substituted group comprises one or more optionally protected substituent groups independently selected from halogen,  $\text{OJ}_1$ ,  $\text{N}(\text{J}_1)(\text{J}_2)$ ,  $=\text{NJ}_1$ ,  $\text{SJ}_1$ ,  $\text{N}_3$ ,  $\text{CN}$ ,  $\text{OC}(=\text{X}_2)\text{J}_1$ ,  $\text{OC}(=\text{X}_2)\text{N}(\text{J}_1)(\text{J}_2)$  and  $\text{C}(=\text{X}_2)\text{N}(\text{J}_1)(\text{J}_2)$ ;

5  $\text{X}_2$  is O, S or  $\text{NJ}_3$ ;

each  $\text{J}_1$ ,  $\text{J}_2$  and  $\text{J}_3$  is, independently, H or  $\text{C}_1\text{-C}_6$  alkyl;


when j is 1 then Z is other than halogen or  $\text{N}(\text{E}_2)(\text{E}_3)$ ; and

wherein said oligomeric compound comprises from 8 to 40 monomeric subunits and is hybridizable to at least a portion of a target nucleic acid.

10 In certain embodiments,  $\text{M}_3$  is O,  $\text{CH}=\text{CH}$ ,  $\text{OCH}_2$  or  $\text{OC}(\text{H})(\text{B}_{\text{X}_2})$ . In certain embodiments,  $\text{M}_3$  is O.

In certain embodiments,  $\text{J}_4$ ,  $\text{J}_5$ ,  $\text{J}_6$  and  $\text{J}_7$  are each H. In certain embodiments,  $\text{J}_4$  forms a bridge with one of  $\text{J}_5$  or  $\text{J}_7$ .


In certain embodiments, A has one of the formulas:



15 wherein:

$\text{Q}_1$  and  $\text{Q}_2$  are each, independently, H, halogen,  $\text{C}_1\text{-C}_6$  alkyl, substituted  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_1\text{-C}_6$  alkoxy or substituted  $\text{C}_1\text{-C}_6$  alkoxy. In certain embodiments,  $\text{Q}_1$  and  $\text{Q}_2$  are each H. In certain embodiments,  $\text{Q}_1$  and  $\text{Q}_2$  are each, independently, H or halogen. In certain embodiments,  $\text{Q}_1$  and  $\text{Q}_2$  is H and the other of  $\text{Q}_1$  and  $\text{Q}_2$  is F,  $\text{CH}_3$  or  $\text{OCH}_3$ .

20 In certain embodiments,  $\text{T}_1$  has the formula:




wherein:

$\text{R}_a$  and  $\text{R}_c$  are each, independently, protected hydroxyl, protected thiol,  $\text{C}_1\text{-C}_6$  alkyl, substituted  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_1\text{-C}_6$  alkoxy, substituted  $\text{C}_1\text{-C}_6$  alkoxy, protected amino or substituted amino; and

$R_b$  is O or S. In certain embodiments,  $R_b$  is O and  $R_a$  and  $R_c$  are each, independently,  $OCH_3$ ,  $OCH_2CH_3$  or  $CH(CH_3)_2$ .

In certain embodiments, G is halogen,  $OCH_3$ ,  $OCH_2F$ ,  $OCHF_2$ ,  $OCF_3$ ,  $OCH_2CH_3$ ,  $O(CH_2)_2F$ ,  $OCH_2CHF_2$ ,  $OCH_2CF_3$ ,  $OCH_2-CH=CH_2$ ,  $O(CH_2)_2-OCH_3$ ,  $O(CH_2)_2-SCH_3$ ,  $O(CH_2)_2-OCF_3$ ,  $O(CH_2)_3-$   
 5  $N(R_{10})(R_{11})$ ,  $O(CH_2)_2-ON(R_{10})(R_{11})$ ,  $O(CH_2)_2-O(CH_2)_2-N(R_{10})(R_{11})$ ,  $OCH_2C(=O)-N(R_{10})(R_{11})$ ,  $OCH_2C(=O)-$   
 $N(R_{12})-(CH_2)_2-N(R_{10})(R_{11})$  or  $O(CH_2)_2-N(R_{12})-C(=NR_{13})[N(R_{10})(R_{11})]$  wherein  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$  and  $R_{13}$  are  
 each, independently, H or  $C_1-C_6$  alkyl. In certain embodiments, G is halogen,  $OCH_3$ ,  $OCF_3$ ,  $OCH_2CH_3$ ,  
 $OCH_2CF_3$ ,  $OCH_2-CH=CH_2$ ,  $O(CH_2)_2-OCH_3$ ,  $O(CH_2)_2-O(CH_2)_2-N(CH_3)_2$ ,  $OCH_2C(=O)-N(H)CH_3$ ,  
 $OCH_2C(=O)-N(H)-(CH_2)_2-N(CH_3)_2$  or  $OCH_2-N(H)-C(=NH)NH_2$ . In certain embodiments, G is F,  $OCH_3$  or  
 10  $O(CH_2)_2-OCH_3$ . In certain embodiments, G is  $O(CH_2)_2-OCH_3$ .

In certain embodiments, the 5'-terminal nucleoside has Formula IIc:



IIc

In certain embodiments, antisense compounds, including those particularly suitable for ssRNA  
 15 comprise one or more type of modified sugar moieties and/or naturally occurring sugar moieties arranged  
 along an oligonucleotide or region thereof in a defined pattern or sugar modification motif. Such motifs  
 may include any of the sugar modifications discussed herein and/or other known sugar modifications.

In certain embodiments, the oligonucleotides comprise or consist of a region having uniform sugar  
 modifications. In certain such embodiments, each nucleoside of the region comprises the same RNA-like  
 20 sugar modification. In certain embodiments, each nucleoside of the region is a 2'-F nucleoside. In certain  
 embodiments, each nucleoside of the region is a 2'-OMe nucleoside. In certain embodiments, each  
 nucleoside of the region is a 2'-MOE nucleoside. In certain embodiments, each nucleoside of the region is a  
 cEt nucleoside. In certain embodiments, each nucleoside of the region is an LNA nucleoside. In certain  
 embodiments, the uniform region constitutes all or essentially all of the oligonucleotide. In certain  
 25 embodiments, the region constitutes the entire oligonucleotide except for 1-4 terminal nucleosides.

In certain embodiments, oligonucleotides comprise one or more regions of alternating sugar  
 modifications, wherein the nucleosides alternate between nucleotides having a sugar modification of a first  
 type and nucleotides having a sugar modification of a second type. In certain embodiments, nucleosides of

both types are RNA-like nucleosides. In certain embodiments the alternating nucleosides are selected from: 2'-OMe, 2'-F, 2'-MOE, LNA, and cEt. In certain embodiments, the alternating modifications are 2'-F and 2'-OMe. Such regions may be contiguous or may be interrupted by differently modified nucleosides or conjugated nucleosides.

In certain embodiments, the alternating region of alternating modifications each consist of a single nucleoside (i.e., the pattern is  $(AB)_x A_y$  wherein A is a nucleoside having a sugar modification of a first type and B is a nucleoside having a sugar modification of a second type; x is 1-20 and y is 0 or 1). In certain embodiments, one or more alternating regions in an alternating motif includes more than a single nucleoside of a type. For example, oligonucleotides may include one or more regions of any of the following

nucleoside motifs:

AABBAA;

ABBABB;

AABAAB;

ABBABAABB;

ABABAA;

AABABAB;

ABABAA;

ABBAABBABABAA;

BABBAABBABABAA; or

ABABBAABBABABAA;

wherein A is a nucleoside of a first type and B is a nucleoside of a second type. In certain embodiments, A and B are each selected from 2'-F, 2'-OMe, BNA, and MOE.

In certain embodiments, oligonucleotides having such an alternating motif also comprise a modified 5' terminal nucleoside, such as those of formula IIc or IIe.

In certain embodiments, oligonucleotides comprise a region having a 2-2-3 motif. Such regions comprises the following motif:

$-(A)_2-(B)_x-(A)_2-(C)_y-(A)_3-$

wherein: A is a first type of modified nucleoside;

B and C, are nucleosides that are differently modified than A, however, B and C may have the same

or different modifications as one another;

x and y are from 1 to 15.

In certain embodiments, A is a 2'-OMe modified nucleoside. In certain embodiments, B and C are both 2'-F modified nucleosides. In certain embodiments, A is a 2'-OMe modified nucleoside and B and C are both 2'-F modified nucleosides.

In certain embodiments, oligonucleosides have the following sugar motif:





wherein:

Q is a nucleoside comprising a stabilized phosphate moiety. In certain embodiments, Q is a nucleoside having Formula IIc or IIe;

5 A is a first type of modified nucleoside;

B is a second type of modified nucleoside;

D is a modified nucleoside comprising a modification different from the nucleoside adjacent to it.


Thus, if y is 0, then D must be differently modified than B and if y is 1, then D must be differently modified than A. In certain embodiments, D differs from both A and B.

10 X is 5-15;

Y is 0 or 1;

Z is 0-4.

In certain embodiments, oligonucleosides have the following sugar motif:



15 wherein:

Q is a nucleoside comprising a stabilized phosphate moiety. In certain embodiments, Q is a nucleoside having Formula IIc or IIe;

A is a first type of modified nucleoside;

D is a modified nucleoside comprising a modification different from A.

20 X is 11-30;

Z is 0-4.

In certain embodiments A, B, C, and D in the above motifs are selected from: 2'-OMe, 2'-F, 2'-MOE, LNA, and cEt. In certain embodiments, D represents terminal nucleosides. In certain embodiments, such terminal nucleosides are not designed to hybridize to the target nucleic acid (though one or more might hybridize by chance). In certain embodiments, the nucleobase of each D nucleoside is adenine, regardless of the identity of the nucleobase at the corresponding position of the target nucleic acid. In certain embodiments the nucleobase of each D nucleoside is thymine.

In certain embodiments, antisense compounds, including those particularly suited for use as ssRNA comprise modified internucleoside linkages arranged along the oligonucleotide or region thereof in a defined pattern or modified internucleoside linkage motif. In certain embodiments, oligonucleotides comprise a region having an alternating internucleoside linkage motif. In certain embodiments, oligonucleotides comprise a region of uniformly modified internucleoside linkages. In certain such embodiments, the oligonucleotide comprises a region that is uniformly linked by phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide is uniformly linked by phosphorothioate internucleoside linkages. In certain embodiments, each internucleoside linkage of the

oligonucleotide is selected from phosphodiester and phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate and at least one internucleoside linkage is phosphorothioate.

In certain embodiments, the oligonucleotide comprises at least 6 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 8 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 10 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 6 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 8 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 10 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 12 consecutive phosphorothioate internucleoside linkages. In certain such embodiments, at least one such block is located at the 3' end of the oligonucleotide. In certain such embodiments, at least one such block is located within 3 nucleosides of the 3' end of the oligonucleotide.

Oligonucleotides having any of the various sugar motifs described herein, may have any linkage motif. For example, the oligonucleotides, including but not limited to those described above, may have a linkage motif selected from non-limiting the table below:

5' most linkage	Central region	3'-region
PS	Alternating PO/PS	6 PS
PS	Alternating PO/PS	7 PS
PS	Alternating PO/PS	8 PS

#### ii. siRNA compounds

In certain embodiments, antisense compounds are double-stranded RNAi compounds (siRNA). In such embodiments, one or both strands may comprise any modification motif described above for ssRNA. In certain embodiments, ssRNA compounds may be unmodified RNA. In certain embodiments, siRNA compounds may comprise unmodified RNA nucleosides, but modified internucleoside linkages.

Several embodiments relate to double-stranded compositions wherein each strand comprises a motif defined by the location of one or more modified or unmodified nucleosides. In certain embodiments, compositions are provided comprising a first and a second oligomeric compound that are fully or at least partially hybridized to form a duplex region and further comprising a region that is complementary to and hybridizes to a nucleic acid target. It is suitable that such a composition comprise a first oligomeric compound that is an antisense strand having full or partial complementarity to a nucleic acid target and a

second oligomeric compound that is a sense strand having one or more regions of complementarity to and forming at least one duplex region with the first oligomeric compound.

The compositions of several embodiments modulate gene expression by hybridizing to a nucleic acid target resulting in loss of its normal function. In some embodiments, the target nucleic acid is MALAT-1. In certain embodiment, the degradation of the targeted MALAT-1 is facilitated by an activated RISC complex that is formed with compositions of the invention.

Several embodiments are directed to double-stranded compositions wherein one of the strands is useful in, for example, influencing the preferential loading of the opposite strand into the RISC (or cleavage) complex. The compositions are useful for targeting selected nucleic acid molecules and modulating the expression of one or more genes. In some embodiments, the compositions of the present invention hybridize to a portion of a target RNA resulting in loss of normal function of the target RNA.

Certain embodiments are drawn to double-stranded compositions wherein both the strands comprises a hemimer motif, a fully modified motif, a positionally modified motif or an alternating motif. Each strand of the compositions of the present invention can be modified to fulfil a particular role in for example the siRNA pathway. Using a different motif in each strand or the same motif with different chemical modifications in each strand permits targeting the antisense strand for the RISC complex while inhibiting the incorporation of the sense strand. Within this model, each strand can be independently modified such that it is enhanced for its particular role. The antisense strand can be modified at the 5'-end to enhance its role in one region of the RISC while the 3'-end can be modified differentially to enhance its role in a different region of the RISC.

The double-stranded oligonucleotide molecules can be a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The double-stranded oligonucleotide molecules can be assembled from two separate oligonucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary (i.e. each strand comprises nucleotide sequence that is complementary to nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double-stranded structure, for example wherein the double-stranded region is about 15 to about 30, e.g., about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 base pairs; the antisense strand comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof (e.g., about 15 to about 25 or more nucleotides of the double-stranded oligonucleotide molecule are complementary to the target nucleic acid or a portion thereof). Alternatively, the double-stranded oligonucleotide is assembled from a single

oligonucleotide, where the self-complementary sense and antisense regions of the siRNA are linked by means of a nucleic acid based or non-nucleic acid-based linker(s).

The double-stranded oligonucleotide can be a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. The double-stranded oligonucleotide can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either in vivo or in vitro to generate an active siRNA molecule capable of mediating RNAi.

In certain embodiments, the double-stranded oligonucleotide comprises separate sense and antisense sequences or regions, wherein the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linker molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. In certain embodiments, the double-stranded oligonucleotide comprises nucleotide sequence that is complementary to nucleotide sequence of a target gene. In another embodiment, the double-stranded oligonucleotide interacts with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene.

As used herein, double-stranded oligonucleotides need not be limited to those molecules containing only RNA, but further encompasses chemically modified nucleotides and non-nucleotides. In certain embodiments, the short interfering nucleic acid molecules lack 2'-hydroxy (2'-OH) containing nucleotides. In certain embodiments short interfering nucleic acids optionally do not include any ribonucleotides (e.g., nucleotides having a 2'-OH group). Such double-stranded oligonucleotides that do not require the presence of ribonucleotides within the molecule to support RNAi can however have an attached linker or linkers or other attached or associated groups, moieties, or chains containing one or more nucleotides with 2'-OH groups. Optionally, double-stranded oligonucleotides can comprise ribonucleotides at about 5, 10, 20, 30, 40, or 50% of the nucleotide positions. As used herein, the term siRNA is meant to be equivalent to other terms used to describe nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. In addition, as used herein, the term RNAi is meant to be equivalent to other

terms used to describe sequence specific RNA interference, such as post transcriptional gene silencing, translational inhibition, or epigenetics. For example, double-stranded oligonucleotides can be used to epigenetically silence genes at both the post-transcriptional level and the pre-transcriptional level. In a non-limiting example, epigenetic regulation of gene expression by siRNA molecules of the invention can result from siRNA mediated modification of chromatin structure or methylation pattern to alter gene expression (see, for example, Verdel et al., 2004, Science, 303, 672-676; Pal-Bhadra et al., 2004, Science, 303, 669-672; Allshire, 2002, Science, 297, 1818-1819; Volpe et al., 2002, Science, 297, 1833-1837; Jenuwein, 2002, Science, 297, 2215-2218; and Hall et al., 2002, Science, 297, 2232-2237).

It is contemplated that compounds and compositions of several embodiments provided herein can target MALAT-1 by a dsRNA-mediated gene silencing or RNAi mechanism, including, e.g., "hairpin" or stem-loop double-stranded RNA effector molecules in which a single RNA strand with self-complementary sequences is capable of assuming a double-stranded conformation, or duplex dsRNA effector molecules comprising two separate strands of RNA. In various embodiments, the dsRNA consists entirely of ribonucleotides or consists of a mixture of ribonucleotides and deoxynucleotides, such as the RNA/DNA hybrids disclosed, for example, by WO 00/63364, filed Apr. 19, 2000, or U.S. Ser. No. 60/130,377, filed Apr. 21, 1999. The dsRNA or dsRNA effector molecule may be a single molecule with a region of self-complementarity such that nucleotides in one segment of the molecule base pair with nucleotides in another segment of the molecule. In various embodiments, a dsRNA that consists of a single molecule consists entirely of ribonucleotides or includes a region of ribonucleotides that is complementary to a region of deoxyribonucleotides. Alternatively, the dsRNA may include two different strands that have a region of complementarity to each other.

In various embodiments, both strands consist entirely of ribonucleotides, one strand consists entirely of ribonucleotides and one strand consists entirely of deoxyribonucleotides, or one or both strands contain a mixture of ribonucleotides and deoxyribonucleotides. In certain embodiments, the regions of complementarity are at least 70, 80, 90, 95, 98, or 100% complementary to each other and to a target nucleic acid sequence. In certain embodiments, the region of the dsRNA that is present in a double-stranded conformation includes at least 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 50, 75, 100, 200, 500, 1000, 2000 or 5000 nucleotides or includes all of the nucleotides in a cDNA or other target nucleic acid sequence being represented in the dsRNA. In some embodiments, the dsRNA does not contain any single stranded regions, such as single stranded ends, or the dsRNA is a hairpin. In other embodiments, the dsRNA has one or more single stranded regions or overhangs. In certain embodiments, RNA/DNA hybrids include a DNA strand or region that is an antisense strand or region (e.g., has at least 70, 80, 90, 95, 98, or 100% complementarity to a target nucleic acid) and an RNA strand or region that is a sense strand or region (e.g., has at least 70, 80, 90, 95, 98, or 100% identity to a target nucleic acid), and vice versa.

In various embodiments, the RNA/DNA hybrid is made in vitro using enzymatic or chemical synthetic methods such as those described herein or those described in WO 00/63364, filed Apr. 19, 2000, or U.S. Ser. No. 60/130,377, filed Apr. 21, 1999. In other embodiments, a DNA strand synthesized in vitro is complexed with an RNA strand made in vivo or in vitro before, after, or concurrent with the transformation of the DNA strand into the cell. In yet other embodiments, the dsRNA is a single circular nucleic acid containing a sense and an antisense region, or the dsRNA includes a circular nucleic acid and either a second circular nucleic acid or a linear nucleic acid (see, for example, WO 00/63364, filed Apr. 19, 2000, or U.S. Ser. No. 60/130,377, filed Apr. 21, 1999.) Exemplary circular nucleic acids include lariat structures in which the free 5' phosphoryl group of a nucleotide becomes linked to the 2' hydroxyl group of another nucleotide in a loop back fashion.

In other embodiments, the dsRNA includes one or more modified nucleotides in which the 2' position in the sugar contains a halogen (such as fluorine group) or contains an alkoxy group (such as a methoxy group) which increases the half-life of the dsRNA in vitro or in vivo compared to the corresponding dsRNA in which the corresponding 2' position contains a hydrogen or an hydroxyl group. In yet other embodiments, the dsRNA includes one or more linkages between adjacent nucleotides other than a naturally-occurring phosphodiester linkage. Examples of such linkages include phosphoramidate, phosphorothioate, and phosphorodithioate linkages. The dsRNAs may also be chemically modified nucleic acid molecules as taught in U.S. Pat. No. 6,673,661. In other embodiments, the dsRNA contains one or two capped strands, as disclosed, for example, by WO 00/63364, filed Apr. 19, 2000, or U.S. Ser. No. 60/130,377, filed Apr. 21, 1999.

In other embodiments, the dsRNA can be any of the at least partially dsRNA molecules disclosed in WO 00/63364, as well as any of the dsRNA molecules described in U.S. Provisional Application 60/399,998; and U.S. Provisional Application 60/419,532, and PCT/US2003/033466. Any of the dsRNAs may be expressed in vitro or in vivo using the methods described herein or standard methods, such as those described in WO 00/63364.

### *Occupancy*

In certain embodiments, antisense compounds are not expected to result in cleavage or the target nucleic acid via RNase H or to result in cleavage or sequestration through the RISC pathway. In certain such embodiments, antisense activity may result from occupancy, wherein the presence of the hybridized antisense compound disrupts the activity of the target nucleic acid. In certain such embodiments, the antisense compound may be uniformly modified or may comprise a mix of modifications and/or modified and unmodified nucleosides.

*Target Nucleic Acids, Target Regions and Nucleotide Sequences*

“Targeting” an oligomeric compound to a particular nucleic acid molecule can be a multistep process. The process usually begins with the identification of a target nucleic acid whose function is to be modulated. This target nucleic acid may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In several embodiments provided herein, the target nucleic acid encodes MALAT-1.

Nucleotide sequences that encode human MALAT-1 are target nucleic acids in several embodiments include, without limitation, the following: GENBANK Accession No. EF177381.1 (incorporated herein as SEQ ID NO: 1), GENBANK Accession No. BK001411.1 (incorporated herein as SEQ ID NO: 2), GENBANK Accession No. BQ429080.1 (incorporated herein as SEQ ID NO: 3), GENBANK Accession No. BQ428957.1 (incorporated herein as SEQ ID NO: 4), GENBANK Accession No. NT\_033903.7 truncated from nucleobases 10569000 to 10582000 (incorporated herein as SEQ ID NO: 5), GENBANK Accession No. XR\_001309.1 (incorporated herein as SEQ ID NO: 6), or GENBANK Accession No. NR\_002819.2 (incorporated herein as SEQ ID NO: 7), GENBANK Accession No. NC\_000011.9 from nucleobases 65265233 to 65273940 (incorporated herein as SEQ ID NO: 8), GENBANK Accession No. AC\_000143.1 from nucleobases 61592326 to 61601033 (incorporated herein as SEQ ID NO: 9).

Nucleotide sequences that encode mouse MALAT-1 are target nucleic acids in several embodiments include, without limitation, the following: GENBANK Accession No. NR\_002847.2 (incorporated herein as SEQ ID NO: 10), GENBANK Accession No. FJ209304.1 (incorporated herein as SEQ ID NO: 11), and the complement of GENBANK Accession No. NT\_082868.4 truncated from nucleobases 2689000 to 2699000 (incorporated herein as SEQ ID NO: 12).

It is understood that the sequence set forth in each SEQ ID NO in the Detailed Description and/or Examples contained herein is independent of any modification to a sugar moiety, an internucleoside linkage, or a nucleobase. As such, antisense compounds defined by a SEQ ID NO may comprise, independently, one or more modifications to a sugar moiety, an internucleoside linkage, or a nucleobase. Antisense compounds described by Isis Number (Isis No) indicate a combination of nucleobase sequence and motif.

The targeting process usually also includes determination of at least one target region, segment, or site within the target nucleic acid for the antisense interaction to occur such that the desired effect, e.g., modulation of expression, will result. Within the context of the present embodiments, the term “region” is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or

characteristic. Within regions of target nucleic acids are segments. "Segments" are defined as smaller or sub-portions of regions within a target nucleic acid. "Sites," as used in the present invention, are defined as positions within a target nucleic acid.

5 Targeting includes determination of at least one target segment to which an antisense compound hybridizes, such that a desired effect occurs. In certain embodiments, the desired effect is a reduction in mRNA target nucleic acid levels. In certain embodiments, the desired effect is reduction of levels of protein encoded by the target nucleic acid or a phenotypic change associated with the target nucleic acid.

10 A target region may contain one or more target segments. Multiple target segments within a target region may be overlapping. Alternatively, they may be non-overlapping. In certain embodiments, target segments within a target region are separated by no more than about 300 nucleotides. In certain emodiments, target segments within a target region are separated by a number of nucleotides that is, is about, is no more than, is no more than about, 250, 200, 150, 100, 90, 80, 70, 60, 50, 40, 30, 20, or 10 nucleotides on the target nucleic acid, or is a range defined by any two of the preceeding values. In certain  
15 5 nucleotides on the target nucleic acid. In certain embodiments, target segments are contiguous. Contemplated are target regions defined by a range having a starting nucleic acid that is any of the 5' target sites or 3' target sites listed herein.

Generally, suitable target segments may be found within a 5' UTR, a coding region, a 3' UTR, an intron, an exon, or an exon/intron junction. Target segments containing a start codon or a stop codon  
20 generally are also suitable target segments. A suitable target segment may specifically exclude a certain structurally defined region such as the start codon or stop codon. However, as MALAT-1 transcripts are considered non-coding, suitable target segments may be found throughout the length of the transcript, which is believed to be untranslated.

Nonetheless, target segments including possible MALAT-1 coding transcripts and any structurally  
25 defined regions are still contemplated in several embodiments. For example, a target region may encompass a 3' UTR, a 5' UTR, an exon, an intron, an exon/intron junction, a coding region, a translation initiation region, translation termination region, or other defined nucleic acid region. The structurally defined regions for MALAT-1 can be obtained by accession number from sequence databases such as NCBI. In certain  
30 embodiments, a target region may encompass the sequence from a 5' target site of one target segment within the target region to a 3' target site of another target segment within the same target region.

Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed



mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA transcribed from a gene encoding tyrosinase, regardless of the sequence(s) of such codons. It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively).

The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon. Consequently, the "start codon region" (or "translation initiation codon region") and the "stop codon region" (or "translation termination codon region") are all regions which may be targeted effectively with the antisense compounds of the present invention.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively.

Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA (or corresponding nucleotides on the gene), and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA (or corresponding nucleotides on the gene). The 5' cap site of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap site.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. Targeting splice sites, i.e., intron-exon junctions or exon-intron junctions, may also be particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also possible target sites. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts". It is also known that introns can be effectively targeted using antisense compounds targeted to, for example, DNA or pre-mRNA.

It is also known in the art that alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and exonic sequence.

Upon excision of one or more exon or intron regions, or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

It is also known in the art that variants can be produced through the use of alternative signals to start or stop transcription and that pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites.

The determination of suitable target segments may include a comparison of the sequence of a target nucleic acid to other sequences throughout the genome. For example, the BLAST algorithm may be used to identify regions of similarity amongst different nucleic acids. This comparison can prevent the selection of antisense compound sequences that may hybridize in a non-specific manner to sequences other than a selected target nucleic acid (i.e., non-target or off-target sequences).

There may be variation in activity (e.g., as defined by percent reduction of target nucleic acid levels) of the antisense compounds within an active target region. In certain embodiments, reductions in MALAT-1 mRNA levels are indicative of inhibition of MALAT-1 expression. Reductions in levels of MALAT-1 protein are also indicative of inhibition of target mRNA expression.

## 5 *Hybridization*

In some embodiments, hybridization occurs between an antisense compound disclosed herein and a MALAT-1 nucleic acid. The most common mechanism of hybridization involves hydrogen bonding (e.g., Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding) between complementary nucleobases of the nucleic acid molecules.

10 Hybridization can occur under varying conditions. Stringent conditions are sequence-dependent and are determined by the nature and composition of the nucleic acid molecules to be hybridized.

Methods of determining whether a sequence is specifically hybridizable to a target nucleic acid are well known in the art. In certain embodiments, the antisense compounds provided herein are specifically hybridizable with a MALAT-1 nucleic acid.

## 15 *Complementarity*

An antisense compound and a target nucleic acid are complementary to each other when a sufficient number of nucleobases of the antisense compound can hydrogen bond with the corresponding nucleobases of the target nucleic acid, such that a desired effect will occur (e.g., antisense inhibition of a target nucleic acid, such as a MALAT-1 nucleic acid).

20 Non-complementary nucleobases between an antisense compound and a MALAT-1 nucleic acid may be tolerated provided that the antisense compound remains able to specifically hybridize to a target nucleic acid. Moreover, an antisense compound may hybridize over one or more segments of a MALAT-1 nucleic acid such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure).

25 In certain embodiments, the antisense compounds provided herein, or a specified portion thereof, are, or are at least, 70%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% complementary to a MALAT-1 nucleic acid, a target region, target segment, or specified portion thereof. Percent complementarity of an antisense compound with a target nucleic acid can be determined using routine methods.

30 For example, an antisense compound in which 18 of 20 nucleobases of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90

percent complementarity. In this example, the remaining noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to complementary nucleobases. As such, an antisense compound which is 18 nucleobases in length having 4 (four) noncomplementary nucleobases which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., J. Mol. Biol., 1990, 215, 403-410; Zhang and Madden, Genome Res., 1997, 7, 649-656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482-489).

In certain embodiments, the antisense compounds provided herein, or specified portions thereof, are fully complementary (i.e. 100% complementary) to a target nucleic acid, or specified portion thereof. For example, an antisense compound may be fully complementary to a MALAT-1 nucleic acid, or a target region, or a target segment or target sequence thereof. As used herein, "fully complementary" means each nucleobase of an antisense compound is capable of precise base pairing with the corresponding nucleobases of a target nucleic acid. For example, a 20 nucleobase antisense compound is fully complementary to a target sequence that is 400 nucleobases long, so long as there is a corresponding 20 nucleobase portion of the target nucleic acid that is fully complementary to the antisense compound. Fully complementary can also be used in reference to a specified portion of the first and/or the second nucleic acid. For example, a 20 nucleobase portion of a 30 nucleobase antisense compound can be "fully complementary" to a target sequence that is 400 nucleobases long. The 20 nucleobase portion of the 30 nucleobase oligonucleotide is fully complementary to the target sequence if the target sequence has a corresponding 20 nucleobase portion wherein each nucleobase is complementary to the 20 nucleobase portion of the antisense compound. At the same time, the entire 30 nucleobase antisense compound may or may not be fully complementary to the target sequence, depending on whether the remaining 10 nucleobases of the antisense compound are also complementary to the target sequence.

The location of a non-complementary nucleobase may be at the 5' end or 3' end of the antisense compound. Alternatively, the non-complementary nucleobase or nucleobases may be at an internal position of the antisense compound. When two or more non-complementary nucleobases are present, they may be contiguous (i.e. linked) or non-contiguous. In one embodiment, a non-complementary nucleobase is located in the wing segment of a gapmer antisense oligonucleotide.

In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleobases in length comprise no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a MALAT-1 nucleic acid, or specified portion thereof.

5 In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleobases in length comprise no more than 6, no more than 5, no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a MALAT-1 nucleic acid, or specified portion thereof.

10 The antisense compounds provided herein also include those which are complementary to a portion of a target nucleic acid. As used herein, "portion" refers to a defined number of contiguous (i.e. linked) nucleobases within a region or segment of a target nucleic acid. A "portion" can also refer to a defined number of contiguous nucleobases of an antisense compound. In certain embodiments, the antisense compounds, are complementary to at least an 8 nucleobase portion of a target segment. In certain  
15 embodiments, the antisense compounds are complementary to at least a 12 nucleobase portion of a target segment. In certain embodiments, the antisense compounds are complementary to at least a 15 nucleobase portion of a target segment. Also contemplated are antisense compounds that are complementary to at least a 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleobase portion of a target segment, or a range defined by any two of these values.

### *Identity*

20 The antisense compounds provided herein may also have a defined percent identity to a particular nucleotide sequence, SEQ ID NO, or compound represented by a specific Isis number, or portion thereof. As used herein, an antisense compound is identical to the sequence disclosed herein if it has the same nucleobase pairing ability. For example, a RNA which contains uracil in place of thymidine in a disclosed DNA sequence would be considered identical to the DNA sequence since both uracil and thymidine pair  
25 with adenine. Shortened and lengthened versions of the antisense compounds described herein as well as compounds having non-identical bases relative to the antisense compounds provided herein also are contemplated. The non-identical bases may be adjacent to each other or dispersed throughout the antisense compound. Percent identity of an antisense compound is calculated according to the number of bases that have identical base pairing relative to the sequence to which it is being compared.

30 In certain embodiments, the antisense compounds, or portions thereof, are at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to one or more of the antisense compounds or SEQ ID NOs, or a portion thereof, disclosed herein.

In certain embodiments, a portion of the antisense compound is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

5 In certain embodiments, a portion of the antisense oligonucleotide is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

### *Modifications*

A nucleoside is a base-sugar combination. The nucleobase (also known as base) portion of the nucleoside is normally a heterocyclic base moiety. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to the 2', 3' or 5' hydroxyl moiety of the sugar. Oligonucleotides are formed through the covalent linkage of adjacent nucleosides to one another, to form a linear polymeric oligonucleotide. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside linkages of the oligonucleotide.

15 Modifications to antisense compounds encompass substitutions or changes to internucleoside linkages, sugar moieties, or nucleobases. Modified antisense compounds are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target, increased stability in the presence of nucleases, or increased inhibitory activity.

20 Chemically modified nucleosides may also be employed to increase the binding affinity of a shortened or truncated antisense oligonucleotide for its target nucleic acid. Consequently, comparable results can often be obtained with shorter antisense compounds that have such chemically modified nucleosides.

### *Modified Internucleoside Linkages*

25 The naturally occurring internucleoside linkage of RNA and DNA is a 3' to 5' phosphodiester linkage. Antisense compounds having one or more modified, i.e. non-naturally occurring, internucleoside linkages are often selected over antisense compounds having naturally occurring internucleoside linkages because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for target nucleic acids, and increased stability in the presence of nucleases.

30 Oligonucleotides having modified internucleoside linkages include internucleoside linkages that retain a phosphorus atom as well as internucleoside linkages that do not have a phosphorus atom. Representative phosphorus containing internucleoside linkages include, but are not limited to,

phosphodiester, phosphotriester, methylphosphonate, phosphoramidate, and phosphorothioate. Methods of preparation of phosphorous-containing and non-phosphorous-containing linkages are well known.

In certain embodiments, antisense compounds targeted to a MALAT-1 nucleic acid comprise one or more modified internucleoside linkages. In certain embodiments, the modified internucleoside linkages are phosphorothioate linkages. In certain embodiments, each internucleoside linkage of an antisense compound is a phosphorothioate internucleoside linkage.

### *Modified Sugar Moieties*

Antisense compounds can optionally contain one or more nucleosides wherein the sugar group has been modified. Such sugar modified nucleosides may impart enhanced nuclease stability, increased binding affinity, or some other beneficial biological property to the antisense compounds. In certain embodiments, nucleosides comprise chemically modified ribofuranose ring moieties. Examples of chemically modified ribofuranose rings include without limitation, addition of substituent groups (including 5' and 2' substituent groups, bridging of non-geminal ring atoms to form bicyclic nucleic acids (BNA), replacement of the ribosyl ring oxygen atom with S, N(R), or C(R<sub>1</sub>)(R<sub>2</sub>) (R, R<sub>1</sub> and R<sub>2</sub> are each independently H, C<sub>1</sub>-C<sub>12</sub> alkyl or a protecting group) and combinations thereof. Examples of chemically modified sugars include 2'-F-5'-methyl substituted nucleoside (see PCT International Application WO 2008/101157 Published on 8/21/08 for other disclosed 5',2'-bis substituted nucleosides) or replacement of the ribosyl ring oxygen atom with S with further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on June 16, 2005) or alternatively 5'-substitution of a BNA (see PCT International Application WO 2007/134181 Published on 11/22/07 wherein LNA is substituted with for example a 5'-methyl or a 5'-vinyl group).

Examples of nucleosides having modified sugar moieties include without limitation nucleosides comprising 5'-vinyl, 5'-methyl (R or S), 4'-S, 2'-F, 2'-OCH<sub>3</sub>, 2'-OCH<sub>2</sub>CH<sub>3</sub>, 2'-OCH<sub>2</sub>CH<sub>2</sub>F and 2'-O(CH<sub>2</sub>)<sub>2</sub>OCH<sub>3</sub> substituent groups. The substituent at the 2' position can also be selected from allyl, amino, azido, thio, O-allyl, O-C<sub>1</sub>-C<sub>10</sub> alkyl, OCF<sub>3</sub>, OCH<sub>2</sub>F, O(CH<sub>2</sub>)<sub>2</sub>SCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>2</sub>-O-N(R<sub>m</sub>)(R<sub>n</sub>), O-CH<sub>2</sub>-C(=O)-N(R<sub>m</sub>)(R<sub>n</sub>), and O-CH<sub>2</sub>-C(=O)-N(R<sub>1</sub>)-(CH<sub>2</sub>)<sub>2</sub>-N(R<sub>m</sub>)(R<sub>n</sub>), where each R<sub>1</sub>, R<sub>m</sub> and R<sub>n</sub> is, independently, H or substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl.

As used herein, "bicyclic nucleosides" refer to modified nucleosides comprising a bicyclic sugar moiety. Examples of bicyclic nucleosides include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments, antisense compounds provided herein include one or more bicyclic nucleosides comprising a 4' to 2' bridge. Examples of such 4' to 2' bridged bicyclic nucleosides, include but are not limited to one of the formulae: 4'-(CH<sub>2</sub>)-O-2' (LNA); 4'-(CH<sub>2</sub>)-S-2';

4'-(CH<sub>2</sub>)<sub>2</sub>-O-2' (ENA); 4'-CH(CH<sub>3</sub>)-O-2' (also referred to as constrained ethyl or cEt) and 4'-CH(CH<sub>2</sub>OCH<sub>3</sub>)-O-2' (and analogs thereof see U.S. Patent 7,399,845, issued on July 15, 2008); 4'-C(CH<sub>3</sub>)(CH<sub>3</sub>)-O-2' (and analogs thereof see published International Application WO/2009/006478, published January 8, 2009); 4'-CH<sub>2</sub>-N(OCH<sub>3</sub>)-2' (and analogs thereof see published International Application WO/2008/150729, published December 11, 2008); 4'-CH<sub>2</sub>-O-N(CH<sub>3</sub>)-2' (see published U.S. Patent Application US2004-0171570, published September 2, 2004); 4'-CH<sub>2</sub>-N(R)-O-2', wherein R is H, C<sub>1</sub>-C<sub>12</sub> alkyl, or a protecting group (see U.S. Patent 7,427,672, issued on September 23, 2008); 4'-CH<sub>2</sub>-C(H)(CH<sub>3</sub>)-2' (see Chattopadhyaya *et al.*, *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH<sub>2</sub>-C(=CH<sub>2</sub>)-2' (and analogs thereof see published International Application WO 2008/154401, published on December 8, 2008).

Further reports related to bicyclic nucleosides can also be found in published literature (see for example: Singh *et al.*, *Chem. Commun.*, 1998, 4, 455-456; Koshkin *et al.*, *Tetrahedron*, 1998, 54, 3607-3630; Wahlestedt *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*, 2000, 97, 5633-5638; Kumar *et al.*, *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222; Singh *et al.*, *J. Org. Chem.*, 1998, 63, 10035-10039; Srivastava *et al.*, *J. Am. Chem. Soc.*, 2007, 129(26) 8362-8379; Elayadi *et al.*, *Curr. Opinion Invest. Drugs*, 2001, 2, 558-561; Braasch *et al.*, *Chem. Biol.*, 2001, 8, 1-7; and Orum *et al.*, *Curr. Opinion Mol. Ther.*, 2001, 3, 239-243; U.S. Patent Nos. 6,268,490; 6,525,191; 6,670,461; 6,770,748; 6,794,499; 7,034,133; 7,053,207; 7,399,845; 7,547,684; and 7,696,345; U.S. Patent Publication No. US2008-0039618; US2009-0012281; U.S. Patent Serial Nos. 60/989,574; 61/026,995; 61/026,998; 61/056,564; 61/086,231; 61/097,787; and 61/099,844; Published PCT International applications WO 1994/014226; WO 2004/106356; WO 2005/021570; WO 2007/134181; WO 2008/150729; WO 2008/154401; and WO 2009/006478. Each of the foregoing bicyclic nucleosides can be prepared having one or more stereochemical sugar configurations including for example  $\alpha$ -L-ribofuranose and  $\beta$ -D-ribofuranose (see PCT international application PCT/DK98/00393, published on March 25, 1999 as WO 99/14226).

In certain embodiments, bicyclic sugar moieties of BNA nucleosides include, but are not limited to, compounds having at least one bridge between the 4' and the 2' position of the pentofuranosyl sugar moiety wherein such bridges independently comprises 1 or from 2 to 4 linked groups independently selected from -[C(R<sub>a</sub>)(R<sub>b</sub>)]<sub>n</sub>-, -C(R<sub>a</sub>)=C(R<sub>b</sub>)-, -C(R<sub>a</sub>)=N-, -C(=O)-, -C(=NR<sub>a</sub>)-, -C(=S)-, -O-, -Si(R<sub>a</sub>)<sub>2</sub>-, -S(=O)<sub>x</sub>-, and -N(R<sub>a</sub>)-

wherein:

x is 0, 1, or 2;

n is 1, 2, 3, or 4;



each  $R_a$  and  $R_b$  is, independently, H, a protecting group, hydroxyl,  $C_1$ - $C_{12}$  alkyl, substituted  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl, substituted  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl, substituted  $C_2$ - $C_{12}$  alkynyl,  $C_5$ - $C_{20}$  aryl, substituted  $C_5$ - $C_{20}$  aryl, heterocycle radical, substituted heterocycle radical, heteroaryl, substituted heteroaryl,  $C_5$ - $C_7$  alicyclic radical, substituted  $C_5$ - $C_7$  alicyclic radical, halogen,  $OJ_1$ ,  $NJ_1J_2$ ,  $SJ_1$ ,  $N_3$ ,  $COOJ_1$ , acyl ( $C(=O)$ -H), substituted acyl, CN, sulfonyl ( $S(=O)_2$ - $J_1$ ), or sulfoxyl ( $S(=O)$ - $J_1$ ); and


each  $J_1$  and  $J_2$  is, independently, H,  $C_1$ - $C_{12}$  alkyl, substituted  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl, substituted  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl, substituted  $C_2$ - $C_{12}$  alkynyl,  $C_5$ - $C_{20}$  aryl, substituted  $C_5$ - $C_{20}$  aryl, acyl ( $C(=O)$ -H), substituted acyl, a heterocycle radical, a substituted heterocycle radical,  $C_1$ - $C_{12}$  aminoalkyl, substituted  $C_1$ - $C_{12}$  aminoalkyl or a protecting group.

10 In certain embodiments, the bridge of a bicyclic sugar moiety is  $-[C(R_a)(R_b)]_n-$ ,  $-[C(R_a)(R_b)]_n-O-$ ,  $-C(R_aR_b)-N(R)-O-$  or  $-C(R_aR_b)-O-N(R)-$ . In certain embodiments, the bridge is  $4'$ - $CH_2$ - $2'$ ,  $4'$ -( $CH_2$ ) $_2$ - $2'$ ,  $4'$ -( $CH_2$ ) $_3$ - $2'$ ,  $4'$ - $CH_2$ - $O$ - $2'$ ,  $4'$ -( $CH_2$ ) $_2$ - $O$ - $2'$ ,  $4'$ - $CH_2$ - $O$ - $N(R)$ - $2'$  and  $4'$ - $CH_2$ - $N(R)$ - $O$ - $2'$  wherein each R is, independently, H, a protecting group or  $C_1$ - $C_{12}$  alkyl.

15 In certain embodiments, bicyclic nucleosides are further defined by isomeric configuration. For example, a nucleoside comprising a  $4'$ - $2'$  methylene-oxy bridge, may be in the  $\alpha$ -L configuration or in the  $\beta$ -D configuration. Previously,  $\alpha$ -L-methyleneoxy ( $4'$ - $CH_2$ - $O$ - $2'$ ) BNA's have been incorporated into antisense oligonucleotides that showed antisense activity (Frieden *et al.*, *Nucleic Acids Research*, 2003, 21, 6365-6372).


20 In certain embodiments, bicyclic nucleosides include, but are not limited to, (A)  $\alpha$ -L-methyleneoxy ( $4'$ - $CH_2$ - $O$ - $2'$ ) BNA, (B)  $\beta$ -D-methyleneoxy ( $4'$ - $CH_2$ - $O$ - $2'$ ) BNA, (C) ethyleneoxy ( $4'$ -( $CH_2$ ) $_2$ - $O$ - $2'$ ) BNA, (D) aminooxy ( $4'$ - $CH_2$ - $O$ - $N(R)$ - $2'$ ) BNA, (E) oxyamino ( $4'$ - $CH_2$ - $N(R)$ - $O$ - $2'$ ) BNA, and (F) methyl(methyleneoxy) ( $4'$ - $CH(CH_3)$ - $O$ - $2'$ ) BNA, (G) methylene-thio ( $4'$ - $CH_2$ - $S$ - $2'$ ) BNA, (H) methylene-amino ( $4'$ - $CH_2$ - $N(R)$ - $2'$ ) BNA, (I) methyl carbocyclic ( $4'$ - $CH_2$ - $CH(CH_3)$ - $2'$ ) BNA, (J) propylene carbocyclic ( $4'$ -( $CH_2$ ) $_3$ - $2'$ ) BNA and (K) vinyl BNA as depicted below:

25



wherein Bx is the base moiety and R is independently H, a protecting group, C<sub>1</sub>-C<sub>12</sub> alkyl or C<sub>1</sub>-C<sub>12</sub> alkoxy.

In certain embodiments, bicyclic nucleosides are provided having Formula I:



wherein:

Bx is a heterocyclic base moiety;


-Q<sub>a</sub>-Q<sub>b</sub>-Q<sub>c</sub>- is -CH<sub>2</sub>-N(R<sub>c</sub>)-CH<sub>2</sub>-, -C(=O)-N(R<sub>c</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-O-N(R<sub>c</sub>)-, -CH<sub>2</sub>-N(R<sub>c</sub>)-O- or -N(R<sub>c</sub>)-O-CH<sub>2</sub>;

10

R<sub>c</sub> is C<sub>1</sub>-C<sub>12</sub> alkyl or an amino protecting group; and

T<sub>a</sub> and T<sub>b</sub> are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium.

In certain embodiments, bicyclic nucleosides are provided having Formula II:



5 wherein:


Bx is a heterocyclic base moiety;

T<sub>a</sub> and T<sub>b</sub> are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

10 Z<sub>a</sub> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkynyl, acyl, substituted acyl, substituted amide, thiol or substituted thio.

In one embodiment, each of the substituted groups is, independently, mono or poly substituted with substituent groups independently selected from halogen, oxo, hydroxyl, OJ<sub>c</sub>, NJ<sub>c</sub>J<sub>d</sub>, SJ<sub>c</sub>, N<sub>3</sub>, OC(=X)J<sub>e</sub>, and NJ<sub>c</sub>C(=X)NJ<sub>c</sub>J<sub>d</sub>, wherein each J<sub>c</sub>, J<sub>d</sub> and J<sub>e</sub> is, independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl, or substituted C<sub>1</sub>-C<sub>6</sub> alkyl and X is O or NJ<sub>c</sub>.

15 In certain embodiments, bicyclic nucleosides are provided having Formula III:




wherein:

Bx is a heterocyclic base moiety;

$T_a$  and  $T_b$  are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

$Z_b$  is  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, substituted  $C_1$ - $C_6$  alkyl, substituted  $C_2$ - $C_6$  alkenyl, substituted  $C_2$ - $C_6$  alkynyl or substituted acyl ( $C(=O)-$ ).

5 In certain embodiments, bicyclic nucleosides are provided having Formula IV:



wherein:


$Bx$  is a heterocyclic base moiety;

10  $T_a$  and  $T_b$  are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

$R_d$  is  $C_1$ - $C_6$  alkyl, substituted  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl, substituted  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl or substituted  $C_2$ - $C_6$  alkynyl;

15 each  $q_a$ ,  $q_b$ ,  $q_c$  and  $q_d$  is, independently, H, halogen,  $C_1$ - $C_6$  alkyl, substituted  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl, substituted  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl or substituted  $C_2$ - $C_6$  alkynyl,  $C_1$ - $C_6$  alkoxy, substituted  $C_1$ - $C_6$  alkoxy, acyl, substituted acyl,  $C_1$ - $C_6$  aminoalkyl or substituted  $C_1$ - $C_6$  aminoalkyl;

In certain embodiments, bicyclic nucleosides are provided having Formula V:



wherein:

Bx is a heterocyclic base moiety;

T<sub>a</sub> and T<sub>b</sub> are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

5 q<sub>a</sub>, q<sub>b</sub>, q<sub>c</sub> and q<sub>f</sub> are each, independently, hydrogen, halogen, C<sub>1</sub>-C<sub>12</sub> alkyl, substituted C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, substituted C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, substituted C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>1</sub>-C<sub>12</sub> alkoxy, substituted C<sub>1</sub>-C<sub>12</sub> alkoxy, OJ<sub>j</sub>, SJ<sub>j</sub>, SOJ<sub>j</sub>, SO<sub>2</sub>J<sub>j</sub>, NJ<sub>j</sub>J<sub>k</sub>, N<sub>3</sub>, CN, C(=O)OJ<sub>j</sub>, C(=O)NJ<sub>j</sub>J<sub>k</sub>, C(=O)J<sub>j</sub>, O-C(=O)-NJ<sub>j</sub>J<sub>k</sub>, N(H)C(=NH)NJ<sub>j</sub>J<sub>k</sub>, N(H)C(=O)NJ<sub>j</sub>J<sub>k</sub> or N(H)C(=S)NJ<sub>j</sub>J<sub>k</sub>;


or q<sub>c</sub> and q<sub>f</sub> together are =C(q<sub>g</sub>)(q<sub>h</sub>);

10 q<sub>g</sub> and q<sub>h</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>12</sub> alkyl or substituted C<sub>1</sub>-C<sub>12</sub> alkyl.

The synthesis and preparation of the methyleneoxy (4'-CH<sub>2</sub>-O-2') BNA monomers adenine, cytosine, guanine, 5-methyl-cytosine, thymine and uracil, along with their oligomerization, and nucleic acid recognition properties have been described (Koshkin *et al.*, *Tetrahedron*, 1998, 54, 3607-3630). BNAs and preparation thereof are also described in WO 98/39352 and WO 99/14226.

15 Analogs of methyleneoxy (4'-CH<sub>2</sub>-O-2') BNA and 2'-thio-BNAs, have also been prepared (Kumar *et al.*, *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222). Preparation of locked nucleoside analogs comprising oligodeoxyribonucleotide duplexes as substrates for nucleic acid polymerases has also been described (Wengel *et al.*, WO 99/14226 ). Furthermore, synthesis of 2'-amino-BNA, a novel conformationally restricted high-affinity oligonucleotide analog has been described in the art (Singh *et al.*, *J. Org. Chem.*,  
20 1998, 63, 10035-10039). In addition, 2'-amino- and 2'-methylamino-BNA's have been prepared and the thermal stability of their duplexes with complementary RNA and DNA strands has been previously reported.

In certain embodiments, bicyclic nucleosides are provided having Formula VI:



wherein:

B<sub>x</sub> is a heterocyclic base moiety;

T<sub>a</sub> and T<sub>b</sub> are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

5 each q<sub>i</sub>, q<sub>j</sub>, q<sub>k</sub> and q<sub>l</sub> is, independently, H, halogen, C<sub>1</sub>-C<sub>12</sub> alkyl, substituted C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, substituted C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, substituted C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>1</sub>-C<sub>12</sub> alkoxy, substituted C<sub>1</sub>-C<sub>12</sub> alkoxy, OJ<sub>j</sub>, SJ<sub>j</sub>, SOJ<sub>j</sub>, SO<sub>2</sub>J<sub>j</sub>, NJ<sub>j</sub>J<sub>k</sub>, N<sub>3</sub>, CN, C(=O)OJ<sub>j</sub>, C(=O)NJ<sub>j</sub>J<sub>k</sub>, C(=O)J<sub>j</sub>, O-C(=O)NJ<sub>j</sub>J<sub>k</sub>, N(H)C(=NH)NJ<sub>j</sub>J<sub>k</sub>, N(H)C(=O)NJ<sub>j</sub>J<sub>k</sub> or N(H)C(=S)NJ<sub>j</sub>J<sub>k</sub>; and

10 q<sub>i</sub> and q<sub>j</sub> or q<sub>i</sub> and q<sub>k</sub> together are =C(q<sub>g</sub>)(q<sub>h</sub>), wherein q<sub>g</sub> and q<sub>h</sub> are each, independently, H, halogen, C<sub>1</sub>-C<sub>12</sub> alkyl or substituted C<sub>1</sub>-C<sub>12</sub> alkyl.

One carbocyclic bicyclic nucleoside having a 4'-(CH<sub>2</sub>)<sub>3</sub>-2' bridge and the alkenyl analog bridge 4'-CH=CH-CH<sub>2</sub>-2' have been described (Freier *et al.*, *Nucleic Acids Research*, 1997, 25(22), 4429-4443 and Albaek *et al.*, *J. Org. Chem.*, 2006, 71, 7731-7740). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (Srivastava  
15 *et al.*, *J. Am. Chem. Soc.*, 2007, 129(26), 8362-8379).

As used herein, “4'-2' bicyclic nucleoside” or “4' to 2' bicyclic nucleoside” refers to a bicyclic nucleoside comprising a furanose ring comprising a bridge connecting two carbon atoms of the furanose ring connects the 2' carbon atom and the 4' carbon atom of the sugar ring.

20 As used herein, “monocyclic nucleosides” refer to nucleosides comprising modified sugar moieties that are not bicyclic sugar moieties. In certain embodiments, the sugar moiety, or sugar moiety analogue, of a nucleoside may be modified or substituted at any position.


As used herein, “2'-modified sugar” means a furanosyl sugar modified at the 2' position. In certain embodiments, such modifications include substituents selected from: a halide, including, but not limited to substituted and unsubstituted alkoxy, substituted and unsubstituted thioalkyl, substituted and unsubstituted  
25 amino alkyl, substituted and unsubstituted alkyl, substituted and unsubstituted allyl, and substituted and unsubstituted alkynyl. In certain embodiments, 2' modifications are selected from substituents including, but not limited to: O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>F, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, OCH<sub>2</sub>C(=O)N(H)CH<sub>3</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other 2'-substituent groups can also be selected from: C<sub>1</sub>-C<sub>12</sub> alkyl, substituted alkyl, alkenyl, alkynyl, alkaryl,  
30 aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, F, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA

cleaving group, a reporter group, an intercalator, a group for improving pharmacokinetic properties, or a group for improving the pharmacodynamic properties of an antisense compound, and other substituents having similar properties. In certain embodiments, modified nucleosides comprise a 2'-MOE side chain (Baker *et al.*, *J. Biol. Chem.*, 1997, 272, 11944-12000). Such 2'-MOE substitution have been described as


5 having improved binding affinity compared to unmodified nucleosides and to other modified nucleosides, such as 2'-O-methyl, O-propyl, and O-aminopropyl. Oligonucleotides having the 2'-MOE substituent also have been shown to be antisense inhibitors of gene expression with promising features for *in vivo* use (Martin, *Helv. Chim. Acta*, 1995, 78, 486-504; Altmann *et al.*, *Chimia*, 1996, 50, 168-176; Altmann *et al.*, *Biochem. Soc. Trans.*, 1996, 24, 630-637; and Altmann *et al.*, *Nucleosides Nucleotides*, 1997, 16, 917-926).

10 As used herein, a “modified tetrahydropyran nucleoside” or “modified THP nucleoside” means a nucleoside having a six-membered tetrahydropyran “sugar” substituted in for the pentofuranosyl residue in normal nucleosides (a sugar surrogate). Modified THP nucleosides include, but are not limited to, what is referred to in the art as hexitol nucleic acid (HNA), anitol nucleic acid (ANA), manitol nucleic acid (MNA) (see Leumann, *Bioorg. Med. Chem.*, 2002, 10, 841-854) or fluoro HNA (F-HNA) having a tetrahydropyran

15 ring system as illustrated below:



In certain embodiments, sugar surrogates are selected having Formula VII:



VII

wherein independently for each of said at least one tetrahydropyran nucleoside analog of Formula VII:

Bx is a heterocyclic base moiety;


T<sub>a</sub> and T<sub>b</sub> are each, independently, an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound or one of T<sub>a</sub> and T<sub>b</sub> is an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound and the other of T<sub>a</sub> and T<sub>b</sub> is H, a hydroxyl protecting group, a linked conjugate group or a 5' or 3'-terminal group;

20

5  $q_1, q_2, q_3, q_4, q_5, q_6$  and  $q_7$  are each independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl or substituted C<sub>2</sub>-C<sub>6</sub> alkynyl; and each of R<sub>1</sub> and R<sub>2</sub> is selected from hydrogen, hydroxyl, halogen, substituted or unsubstituted alkoxy, NJ<sub>1</sub>J<sub>2</sub>, SJ<sub>1</sub>, N<sub>3</sub>, OC(=X)J<sub>1</sub>, OC(=X)NJ<sub>1</sub>J<sub>2</sub>, NJ<sub>3</sub>C(=X)NJ<sub>1</sub>J<sub>2</sub> and CN, wherein X is O, S or NJ<sub>1</sub> and each J<sub>1</sub>, J<sub>2</sub> and J<sub>3</sub> is, independently, H or C<sub>1</sub>-C<sub>6</sub> alkyl.

In certain embodiments, the modified THP nucleosides of Formula VII are provided wherein  $q_1, q_2, q_3, q_4, q_5, q_6$  and  $q_7$  are each H. In certain embodiments, at least one of  $q_1, q_2, q_3, q_4, q_5, q_6$  and  $q_7$  is other than H. In certain embodiments, at least one of  $q_1, q_2, q_3, q_4, q_5, q_6$  and  $q_7$  is methyl. In certain embodiments, THP nucleosides of Formula VII are provided wherein one of R<sub>1</sub> and R<sub>2</sub> is fluoro. In certain  
10 embodiments, R<sub>1</sub> is fluoro and R<sub>2</sub> is H; R<sub>1</sub> is methoxy and R<sub>2</sub> is H, and R<sub>1</sub> is methoxyethoxy and R<sub>2</sub> is H.

In certain embodiments, sugar surrogates comprise rings having more than 5 atoms and more than one heteroatom. For example nucleosides comprising morpholino sugar moieties and their use in oligomeric compounds has been reported (see for example: Braasch *et al.*, *Biochemistry*, 2002, 41, 4503-4510; and U.S. Patents 5,698,685; 5,166,315; 5,185,444; and 5,034,506). As used here, the term “morpholino” means a  
15 sugar surrogate having the following formula:




In certain embodiments, morpholinos may be modified, for example by adding or altering various substituent groups from the above morpholino structure. Such sugar surrogates are referred to herein as “modified morpholinos.”

20 Combinations of modifications are also provided without limitation, such as 2'-F-5'-methyl substituted nucleosides (see PCT International Application WO 2008/101157 published on 8/21/08 for other disclosed 5', 2'-bis substituted nucleosides) and replacement of the ribosyl ring oxygen atom with S and further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on June 16, 2005) or alternatively 5'-substitution of a bicyclic nucleic acid (see PCT International Application  
25 WO 2007/134181, published on 11/22/07 wherein a 4'-CH<sub>2</sub>-O-2' bicyclic nucleoside is further substituted at the 5' position with a 5'-methyl or a 5'-vinyl group). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (*see, e.g.*, Srivastava *et al.*, *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).



In certain embodiments, antisense compounds comprise one or more modified cyclohexenyl nucleosides, which is a nucleoside having a six-membered cyclohexenyl in place of the pentofuranosyl residue in naturally occurring nucleosides. Modified cyclohexenyl nucleosides include, but are not limited to those described in the art (see for example commonly owned, published PCT Application WO 2010/036696, published on April 10, 2010, Robeyns *et al.*, *J. Am. Chem. Soc.*, 2008, 130(6), 1979-1984; Horváth *et al.*, *Tetrahedron Letters*, 2007, 48, 3621-3623; Nauwelaerts *et al.*, *J. Am. Chem. Soc.*, 2007, 129(30), 9340-9348; Gu *et al.*, *Nucleosides, Nucleotides & Nucleic Acids*, 2005, 24(5-7), 993-998; Nauwelaerts *et al.*, *Nucleic Acids Research*, 2005, 33(8), 2452-2463; Robeyns *et al.*, *Acta Crystallographica, Section F: Structural Biology and Crystallization Communications*, 2005, F61(6), 585-586; Gu *et al.*, *Tetrahedron*, 2004, 60(9), 2111-2123; Gu *et al.*, *Oligonucleotides*, 2003, 13(6), 479-489; Wang *et al.*, *J. Org. Chem.*, 2003, 68, 4499-4505; Verbeure *et al.*, *Nucleic Acids Research*, 2001, 29(24), 4941-4947; Wang *et al.*, *J. Org. Chem.*, 2001, 66, 8478-82; Wang *et al.*, *Nucleosides, Nucleotides & Nucleic Acids*, 2001, 20(4-7), 785-788; Wang *et al.*, *J. Am. Chem.*, 2000, 122, 8595-8602; Published PCT application, WO 06/047842; and Published PCT Application WO 01/049687. Certain modified cyclohexenyl nucleosides have Formula X.



wherein independently for each of said at least one cyclohexenyl nucleoside analog of Formula X:

Bx is a heterocyclic base moiety;

T<sub>3</sub> and T<sub>4</sub> are each, independently, an internucleoside linking group linking the cyclohexenyl nucleoside analog to an antisense compound or one of T<sub>3</sub> and T<sub>4</sub> is an internucleoside linking group linking the tetrahydropyran nucleoside analog to an antisense compound and the other of T<sub>3</sub> and T<sub>4</sub> is H, a hydroxyl protecting group, a linked conjugate group, or a 5'-or 3'-terminal group; and

q<sub>1</sub>, q<sub>2</sub>, q<sub>3</sub>, q<sub>4</sub>, q<sub>5</sub>, q<sub>6</sub>, q<sub>7</sub>, q<sub>8</sub> and q<sub>9</sub> are each, independently, H, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, substituted C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, substituted C<sub>2</sub>-C<sub>6</sub> alkynyl or other sugar substituent group.

As used herein, "2'-modified" or "2'-substituted" refers to a nucleoside comprising a sugar

comprising a substituent at the 2' position other than H or OH. 2'-modified nucleosides, include, but are not limited to, bicyclic nucleosides wherein the bridge connecting two carbon atoms of the sugar ring connects the 2' carbon and another carbon of the sugar ring; and nucleosides with non-bridging 2'substituents, such as allyl, amino, azido, thio, O-allyl, O-C<sub>1</sub>-C<sub>10</sub> alkyl, -OCF<sub>3</sub>, O-(CH<sub>2</sub>)<sub>2</sub>-O-CH<sub>3</sub>, 2'-O(CH<sub>2</sub>)<sub>2</sub>SCH<sub>3</sub>, O-(CH<sub>2</sub>)<sub>2</sub>-  
5 O-N(R<sub>m</sub>)(R<sub>n</sub>), or O-CH<sub>2</sub>-C(=O)-N(R<sub>m</sub>)(R<sub>n</sub>), where each R<sub>m</sub> and R<sub>n</sub> is, independently, H or substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> alkyl. 2'-modified nucleosides may further comprise other modifications, for example at other positions of the sugar and/or at the nucleobase.

As used herein, "2'-F" refers to a nucleoside comprising a sugar comprising a fluoro group at the 2' position of the sugar ring.

10 As used herein, "2'-OMe" or "2'-OCH<sub>3</sub>" or "2'-O-methyl" each refers to a nucleoside comprising a sugar comprising an -OCH<sub>3</sub> group at the 2' position of the sugar ring.

As used herein, "MOE" or "2'-MOE" or "2'-OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>" or "2'-O-methoxyethyl" each refers to a nucleoside comprising a sugar comprising a -OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> group at the 2' position of the sugar ring.

As used herein, "oligonucleotide" refers to a compound comprising a plurality of linked  
15 nucleosides. In certain embodiments, one or more of the plurality of nucleosides is modified. In certain embodiments, an oligonucleotide comprises one or more ribonucleosides (RNA) and/or deoxyribonucleosides (DNA).

Many other bicyclo and tricyclo sugar surrogate ring systems are also known in the art that can be used to modify nucleosides for incorporation into antisense compounds (see for example review article:  
20 Leumann, *Bioorg. Med. Chem.*, 2002, 10, 841-854). Such ring systems can undergo various additional substitutions to enhance activity.

Methods for the preparations of modified sugars are well known to those skilled in the art. Some representative U.S. patents that teach the preparation of such modified sugars include without limitation, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134;  
25 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,670,633; 5,700,920; 5,792,847 and 6,600,032 and International Application PCT/US2005/019219, filed June 2, 2005 and published as WO 2005/121371 on December 22, 2005.

In nucleotides having modified sugar moieties, the nucleobase moieties (natural, modified or a combination thereof) are maintained for hybridization with an appropriate nucleic acid target.

In certain embodiments, antisense compounds comprise one or more nucleosides having modified sugar moieties. In certain embodiments, the modified sugar moiety is 2'-MOE. In certain embodiments, the 2'-MOE modified nucleosides are arranged in a gapmer motif. In certain embodiments, the modified sugar moiety is a bicyclic nucleoside having a (4'-CH(CH<sub>3</sub>)-O-2') bridging group. In certain embodiments, the (4'-CH(CH<sub>3</sub>)-O-2') modified nucleosides are arranged throughout the wings of a gapmer motif.

#### *Conjugated Antisense compounds*

Antisense compounds may be covalently linked to one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the resulting antisense oligonucleotides. Typical conjugate groups include cholesterol moieties and lipid moieties. Additional conjugate groups include carbohydrates, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes.

Antisense compounds can also be modified to have one or more stabilizing groups that are generally attached to one or both termini of antisense compounds to enhance properties such as, for example, nuclease stability. Included in stabilizing groups are cap structures. These terminal modifications protect the antisense compound having terminal nucleic acid from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap), or at the 3'-terminus (3'-cap), or can be present on both termini. Cap structures are well known in the art and include, for example, inverted deoxy abasic caps. Further 3' and 5'-stabilizing groups that can be used to cap one or both ends of an antisense compound to impart nuclease stability include those disclosed in WO 03/004602 published on January 16, 2003.

In certain embodiments, antisense compounds, including, but not limited to those particularly suited for use as ssRNA, are modified by attachment of one or more conjugate groups. In general, conjugate groups modify one or more properties of the attached oligonucleotide, including but not limited to pharmacodynamics, pharmacokinetics, stability, binding, absorption, cellular distribution, cellular uptake, charge and clearance. Conjugate groups are routinely used in the chemical arts and are linked directly or via an optional conjugate linking moiety or conjugate linking group to a parent compound such as an oligonucleotide. Conjugate groups includes without limitation, intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, thioethers, polyethers, cholesterol, thiocholesterol, cholic acid moieties, folate, lipids, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, adamantane, acridine, fluoresceins, rhodamines, coumarins and dyes. Certain conjugate groups have been described previously, for example: cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Let., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Let., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538),

an aliphatic chain, e.g., do-decan-diol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxysterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

For additional conjugates including those useful for ssRNA and their placement within antisense compounds, see e.g., US Application No.; 61/583,963.

### *Compositions and Methods for Formulating Pharmaceutical Compositions*

Antisense oligonucleotides may be admixed with pharmaceutically acceptable active or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions are dependent upon a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

An antisense compound targeted to a MALAT-1 nucleic acid can be utilized in pharmaceutical compositions by combining the antisense compound with a suitable pharmaceutically acceptable diluent or carrier. A pharmaceutically acceptable diluent includes phosphate-buffered saline (PBS). PBS is a diluent suitable for use in compositions to be delivered parenterally. Accordingly, in one embodiment, employed in the methods described herein is a pharmaceutical composition comprising an antisense compound targeted to a MALAT-1 nucleic acid and a pharmaceutically acceptable diluent. In certain embodiments, the pharmaceutically acceptable diluent is PBS. In certain embodiments, the antisense compound is an antisense oligonucleotide.

### *Treatment of Cancer*

In several embodiments, the antisense compounds provided herein are useful for the treatment of cancer in an animal. Examples of certain types of cancers that can be treated with the antisense compounds described herein include, but are not limited to, colon cancer, intestinal cancer, lung cancer (e.g. non-small cell lung cancer), liver cancer, and/or prostate cancer.

As used herein, the terms "tumor cells," "cancer cells," "malignant cells," and "neoplastic cells" are used interchangeably and do not require a particular distinction as to the extent or degree of transformation

or malignancy relative to a "normal cell." Accordingly, "tumor cells," "cancer cells," and "neoplastic cells" are transformed and/or malignant cells, whereas "normal cells" are not transformed and/or malignant.

The term "treating cancer" refers to performing actions that lead to amelioration of cancer or of the symptoms accompanied therewith to a significant extent. The combination of said actions is encompassed by the term "treatment." Amelioration of a cancer includes but is not limited to reducing in the number of cancer cells in an animal or reducing the number of cancer cells at a specific site in the body of an animal. Said treatment as used herein also includes an entire restoration of the health with respect to the cancers referred to herein. It is to be understood that treatment as used in accordance with embodiments provided herein may not be effective in all subjects to be treated. However, a statistically significant portion of subjects suffering from a cancer referred to herein can be successfully treated. Whether a portion is statistically significant can be determined without by a person of ordinary skill in the art using various well known statistic evaluation tools, e.g., determination of confidence intervals, p-value determination, Student's t-test, Mann-Whitney test, etc.

The term "administration" or "administering" includes routes of introducing a MALAT-1 specific inhibitor to an animal to perform its intended function. An example of a route of administration that can be used includes, but is not limited to parenteral administration, such as subcutaneous, intravenous, intramuscular, intraarterial, intraperitoneal, or intracranial injection or infusion.

When a MALAT-1 specific inhibitor is administered parenterally, such as by subcutaneous or intravenous injection or other injection, it can be in the form of a pyrogen-free, parenterally acceptable aqueous solution or suspension. Suspensions can be formulated using suitable dispersing or wetting agents and suspending agents. Compositions for injection can contain a vehicle such as water, saline (e.g., physiological buffered saline), or other isotonic vehicles such as isotonic sodium chloride solution, Ringer's solution, dextrose solution, or other vehicles known in the art.

As used herein, the term "treatment of cancer" or "treating cancer" can be described by a number of different parameters including, but not limited to, reduction in the size of a tumor in an animal having cancer, reduction in the growth or proliferation of a tumor in an animal having cancer, preventing metastasis or reducing the extent of metastasis, and/or extending the survival of an animal having cancer compared to control. In the context of colon cancer, treating colon cancer can also be measured by a reduction in the number of colon polyps of an animal having colon cancer. In several embodiments, the cancer can be a primary cancer.

Several embodiments are drawn to methods of reducing tumor volume or number in an animal comprising administering a MALAT-1 specific inhibitor to the animal. In various aspects of such embodiments, the MALAT-1 specific inhibitor can be an antisense compound which reduces expression of MALAT-1. It will be understood that any of the MALAT-1 specific inhibitors described herein can be used

in embodiments relating to methods of reducing tumor volume or number in an animal. Furthermore, any antisense compound targeting MALAT-1 as described herein can be used in methods of reducing tumor volume or number in an animal. For example, an antisense compound useful for reducing tumor volume can include a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 85% complementary to a MALAT-1 nucleic acid. In several embodiments, the tumor volume can refer to the volume of a primary tumor.

Various embodiments are drawn to methods of inhibiting tumor growth or proliferation in an animal comprising administering a MALAT-1 specific inhibitor to the animal. In several aspects, the growth or proliferation of a primary tumor can be inhibited by administering a MALAT-1 specific inhibitor to the animal. In various aspects of such embodiments, the MALAT-1 specific inhibitor can be an antisense compound which reduces expression of MALAT-1. It will be understood that any of the MALAT-1 specific inhibitors described herein can be used in embodiments relating to methods of inhibiting tumor growth or proliferation in an animal. Furthermore, any antisense compound targeting MALAT-1 as described herein can be used in methods of inhibiting tumor growth or proliferation in an animal. For example, an antisense compound useful for inhibiting tumor growth or proliferation can include a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 85% complementary to a MALAT-1 nucleic acid.

Certain embodiments are drawn to methods of inhibiting cancer metastasis in an animal comprising administering a MALAT-1 specific inhibitor to the animal. In various aspects of such embodiments, the MALAT-1 specific inhibitor can be an antisense compound which reduces expression of MALAT-1. It will be understood that any of the MALAT-1 specific inhibitors described herein can be used in embodiments relating to methods of inhibiting cancer metastasis in an animal. Furthermore, any antisense compound targeting MALAT-1 as described herein can be used in methods of inhibiting cancer metastasis in an animal. For example, an antisense compound useful for inhibiting cancer metastasis can include a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 85% complementary to a MALAT-1 nucleic acid.

Several embodiments are drawn to methods of increasing survival of an animal having cancer comprising administering a MALAT-1 specific inhibitor to the animal. In various aspects of such embodiments, the MALAT-1 specific inhibitor can be an antisense compound which reduces expression of MALAT-1. It will be understood that any of the MALAT-1 specific inhibitors described herein can be used in embodiments relating to methods of increasing survival of an animal having cancer. Furthermore, any antisense compound targeting MALAT-1 as described herein can be used in methods of increasing survival of an animal having cancer. For example, an antisense compound useful for increasing survival in an

animal having cancer can include a modified oligonucleotide consisting of 12 to 30 linked nucleosides, wherein the modified oligonucleotide is at least 85% complementary to a MALAT-1 nucleic acid.

Certain embodiments are directed to the use of a MALAT-1 specific inhibitor in the manufacture of a medicament for treating cancer. In several aspects, the cancer can be a primary cancer. It will be understood that any of the MALAT-1 specific inhibitors described herein can be used in embodiments relating to use of such inhibitors in the manufacture of a medicament for treating cancer. In some aspects, the MALAT-1 specific inhibitor can comprise a compound including a modified oligonucleotide consisting of 12 to 30 linked nucleosides at least 85% complementary to a MALAT-1 nucleic acid. It will also be understood that MALAT-1 specific inhibitors can be used in the manufacture of a medicament for reducing tumor volume or number, inhibiting tumor growth or proliferation, inhibiting cancer metastasis, and/or increasing survival of an animal having cancer.

Similarly, various embodiments relate to a MALAT-1 specific inhibitor for use in the treatment of cancer. In various aspects the cancer can be a primary cancer. It will be understood that any of the MALAT-1 specific inhibitors described herein can be for use in the treatment of cancer. In some aspects, the MALAT-1 specific inhibitor can comprise a compound including a modified oligonucleotide consisting of 12 to 30 linked nucleosides at least 85% complementary to a MALAT-1 nucleic acid. It will also be understood that MALAT-1 specific inhibitors can be used for reducing tumor volume or number, inhibiting tumor growth or proliferation, inhibiting cancer metastasis, and/or increasing survival of an animal having cancer.

MALAT-1 specific inhibitors of several embodiments can be provided to an administering physician or other health care professional in the form of a kit. The kit is a package which houses a container which contains the MALAT-1 specific inhibitor in a suitable pharmaceutical composition, and instructions for administering the pharmaceutical composition to an animal. The kit can also contain separate doses of a MALAT-1 specific inhibitor for serial or sequential administration. The kit can contain suitable delivery devices, e.g., syringes, and the like, along with instructions for administering the MALAT-1 specific inhibitor. The kit can optionally contain instructions for storage, reconstitution (if applicable), and administration. The kit can include a plurality of containers reflecting the number of administrations to be given to an animal.

#### *Cell culture and antisense compounds treatment*

The effects of antisense compounds on the level, activity or expression of MALAT-1 nucleic acids can be tested *in vitro* in a variety of cell types. Cell types used for such analyses are available from commercial vendors (e.g. American Type Culture Collection, Manassus, VA; Zen-Bio, Inc., Research Triangle Park, NC; Clonetics Corporation, Walkersville, MD) and are cultured according to the vendor's

instructions using commercially available reagents (e.g. Invitrogen Life Technologies, Carlsbad, CA). As an example, b.END cells can be used to test antisense compounds on the activity or expression of MALAT-1 nucleic acids.

*In vitro testing of antisense oligonucleotides*

5 Described herein are methods for treatment of cells with antisense oligonucleotides, which can be modified appropriately for treatment with other antisense compounds.

In general, cells are treated with antisense oligonucleotides when the cells reach approximately 60-80% confluency in culture.

10 One reagent commonly used to introduce antisense oligonucleotides into cultured cells includes the cationic lipid transfection reagent LIPOFECTIN (Invitrogen, Carlsbad, CA). Antisense oligonucleotides are mixed with LIPOFECTIN in OPTI-MEM 1 (Invitrogen, Carlsbad, CA) to achieve the desired final concentration of antisense oligonucleotide and a LIPOFECTIN concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

15 Another reagent used to introduce antisense oligonucleotides into cultured cells includes LIPOFECTAMINE (Invitrogen, Carlsbad, CA). Antisense oligonucleotide is mixed with LIPOFECTAMINE in OPTI-MEM 1 reduced serum medium (Invitrogen, Carlsbad, CA) to achieve the desired concentration of antisense oligonucleotide and a LIPOFECTAMINE concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

20 Another technique used to introduce antisense oligonucleotides into cultured cells includes electroporation.

Cells are treated with antisense oligonucleotides by routine methods. Cells are typically harvested 16-24 hours after antisense oligonucleotide treatment, at which time RNA or protein levels of target nucleic acids are measured by methods known in the art and described herein. In general, when treatments are performed in multiple replicates, the data are presented as the average of the replicate treatments.

25 The concentration of antisense oligonucleotide used varies from cell line to cell line. Methods to determine the optimal antisense oligonucleotide concentration for a particular cell line are well known in the art. Antisense oligonucleotides are typically used at concentrations ranging from 1 nM to 300 nM when transfected with LIPOFECTAMINE. Antisense oligonucleotides are used at higher concentrations ranging from 625 to 20,000 nM when transfected using electroporation.



### ***RNA Isolation***

RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. RNA is prepared using methods well known in the art, for example, using the TRIZOL Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's recommended protocols.

### ***Analysis of inhibition of target levels or expression***

Inhibition of levels or expression of a MALAT-1 nucleic acid can be assayed in a variety of ways known in the art. For example, target nucleic acid levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or quantitative real-time PCR. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. Northern blot analysis is also routine in the art. Quantitative real-time PCR can be conveniently accomplished using the commercially available ABI PRISM 7600, 7700, or 7900 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions.

### ***Quantitative Real-Time PCR Analysis of Target RNA Levels***

Quantitation of target RNA levels may be accomplished by quantitative real-time PCR using the ABI PRISM 7600, 7700, or 7900 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. Methods of quantitative real-time PCR are well known in the art.

Prior to real-time PCR, the isolated RNA is subjected to a reverse transcriptase (RT) reaction, which produces complementary DNA (cDNA) that is then used as the substrate for the real-time PCR amplification. The RT and real-time PCR reactions are performed sequentially in the same sample well. RT and real-time PCR reagents are obtained from Invitrogen (Carlsbad, CA). RT real-time-PCR reactions are carried out by methods well known to those skilled in the art.

Gene (or RNA) target quantities obtained by real time PCR are normalized using either the expression level of a gene whose expression is constant, such as cyclophilin A, or by quantifying total RNA using RIBOGREEN (Invitrogen, Inc. Carlsbad, CA). Cyclophilin A expression is quantified by real time PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RIBOGREEN RNA quantification reagent (Invitrogen, Inc. Eugene, OR). Methods of RNA quantification by RIBOGREEN are taught in Jones, L.J., et al, (Analytical Biochemistry, 1998, 265, 368-374). A CYTOFLUOR 4000 instrument (PE Applied Biosystems) is used to measure RIBOGREEN fluorescence.

Probes and primers are designed to hybridize to a MALAT-1 nucleic acid. Methods for designing real-time PCR probes and primers are well known in the art, and may include the use of software such as PRIMER EXPRESS Software (Applied Biosystems, Foster City, CA).

### *Analysis of Protein Levels*

5 Antisense inhibition of MALAT-1 nucleic acids can be assessed by measuring MALAT-1 protein levels. Protein levels of MALAT-1 can be evaluated or quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), enzyme-linked immunosorbent assay (ELISA), quantitative protein assays, protein activity assays (for example, caspase activity assays), immunohistochemistry, immunocytochemistry or fluorescence-activated cell sorting (FACS). Antibodies  
10 directed to a target can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional monoclonal or polyclonal antibody generation methods well known in the art. Antibodies useful for the detection of MALAT-1 are commercially available

## 15 **EXAMPLES**

Having generally described embodiments drawn to compounds and methods for treating cancer in an animal including administering an antisense compound that targets MALAT-1, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only, and are not intended to be limiting.

20

### **Example 1: Antisense inhibition of murine metastasis associated lung adenocarcinoma transcript 1 (MALAT-1) non-coding RNA in b.END cells**

Antisense oligonucleotides targeted to a murine MALAT-1 nucleic acid were tested for their effects on MALAT-1 RNA *in vitro*. Cultured b.END cells were plated at a density of 4,000 cells per well  
25 and transfected using Cytofectin reagent with 3.125 nM, 6.25 nM, 12.5 nM, 25.0 nM, 50.0 nM, or 100.0 nM concentrations of antisense oligonucleotide, as specified in Table 1. After a treatment period of approximately 16 hours, RNA was isolated from the cells and MALAT-1 RNA levels were measured by quantitative real-time PCR.

ISIS 395251 (CCAGGCTGGTTATGACTCAG; SEQ ID NO: 13), targeting murine MALAT-1  
30 gene sequence, SEQ ID NO: 10 (GENBANK Accession No. NR\_002847.2) at start site 3338; ISIS 399462

(GGGTCAGCTGCCAATGCTAG; SEQ ID NO: 14), targeting SEQ ID NO: 10 at start site 1280; and ISIS 399479 (CGGTGCAAGGCTTAGGAATT; SEQ ID NO: 15) targeting SEQ ID NO: 10 at start site 4004, were three of the antisense oligonucleotides tested in the assay. ISIS 395251 is also cross-reactive with human MALAT-1 gene sequence (GENBANK Accession No. NR\_002819.2; SEQ ID NO: 7) at start site 4897. The antisense oligonucleotides were designed as 5-10-5 MOE gapmers, and are 20 nucleosides in length, wherein the central gap segment comprises ten 2'-deoxynucleosides and is flanked on both sides (in the 5' and 3' directions) by wings comprising 5 nucleosides each. Each nucleoside in the 5' wing segment and each nucleoside in the 3' wing segment has a 2'-MOE modification. The internucleoside linkages throughout the gapmer are phosphorothioate (P=S) linkages. All cytosine residues throughout the gapmer are 5-methylcytosines. The half maximal inhibitory concentration (IC<sub>50</sub>) of each oligonucleotide is also presented in Table 1. As illustrated in Table 1, MALAT-1 RNA levels were significantly reduced in a dose-dependent manner in antisense oligonucleotide treated cells.

Table 1

Dose-dependent inhibition of MALAT-1 RNA in b.END cells

ISIS No	3.125 nM	6.25 nM	12.5 nM	25.0 nM	50.0 nM	100.0 nM	IC <sub>50</sub> (nM)
399479	14	31	55	71	84	91	12.6
399462	20	36	50	68	81	92	12.3
395251	23	45	57	66	85	90	10.1

**Example 2: Effect of antisense inhibition of MALAT-1 in an ApcMin mouse model**

ApcMin (Min, multiple intestinal neoplasia) is a point mutation in the murine homolog of the APC gene. Min/+ mice develop intestinal adenomas and are considered a standard model that mirrors the human condition (Moser, A.R. et al., Proc. Natl. Acad. Sci. USA. 90: 8977, 1993). The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on small intestinal polyps load was examined in ApcMin mice.

*Treatment*

ApcMin mice, 9 weeks in age, were randomly divided into three treatment groups of 4 mice each. The first treatment group was injected with 50 mg/kg of ISIS 399479 (SEQ ID NO: 15), administered subcutaneously 5 days per week for 4 weeks. The second treatment group was injected with 50 mg/kg of

control oligonucleotide, ISIS 141923 (5'- CCTTCCCTGAAGGTTCTCC -3', a 5-10-5 MOE gapmer, designated herein as SEQ ID NO: 16, having no known homology to any mouse gene), administered subcutaneously 5 days per week for 4 weeks. The third group was injected with PBS, administered subcutaneously 5 days per week for 4 weeks. On day 28, the mice were euthanized with isoflurane followed by cervical dislocation. Small intestines, colons, and liver tissue were collected and processed for further analysis.

#### *RNA analysis*

RNA isolation was performed using the Invitrogen PureLink™ Total RNA Purification Kit, according to the manufacturer's protocol. RT-PCR was performed using the Step One Plus system from Applied Biosystems. MALAT-1 RNA expression was measured using primer probe set mMALAT1#2 (forward sequence TGGGTTAGAGAAGGCGTGACTG, designated herein as SEQ ID NO: 17; reverse sequence TCAGCGGCAACTGGGAAA, designated herein as SEQ ID NO: 18; probe sequence CGTTGGCACGACACCTTCAGGGACT, designated herein as SEQ ID NO: 19) and normalized to Cyclophilin mRNA expression. The primer probe set for Cyclophilin was m\_Cyclo24 (forward sequence TCGCCGCTTGCTGCA, designated herein as SEQ ID NO: 20; reverse sequence ATCGGCCGTGATGTCGA, designated herein as SEQ ID NO: 21; probe sequence CCATGGTCAACCCACCGTGTTT, designated herein as SEQ ID NO: 22).

MALAT-1 RNA expression was assessed in the liver and intestines. As shown in Table 2, MALAT-1 RNA expression in mice treated with ISIS 399479 was significantly inhibited compared to the control oligonucleotide-treated group. The mRNA expression levels are expressed as percent inhibition of expression levels compared to that in the PBS control.

**Table 2**

Percent inhibition of MALAT-1 RNA levels (%) compared to the PBS control

	ISIS 399479	ISIS 141923
Liver	98	0
Intestines	94	6

#### *Cell proliferation analysis*

The BrdU cell proliferation assay (Rothaeusler, K. and Baumgarth, N. Curr. Protoc. Cytom. 2007. Chapter 7: Unit 7.31) detects 5-bromo-2'-deoxyuridine (BrdU) incorporated into cellular DNA during cell

proliferation. The quantity of BrdU incorporated into cells is a direct indicator of cell proliferation and was measured using an anti-BrdU antibody (Sigma Aldrich).

BrdU, at a concentration of 50 mg/kg, was injected intraperitoneally for 5 consecutive days starting at day 24. The animals were euthanized on day 28. The entire small intestine from each animal was made in a 'Swiss roll', a standard technique for histological studies of rodent intestine (Moolenbeek, C. and Ruitenber, E.J. Lab Anim. 1981. 15: 57-9). Two sections of the intestine at least 500  $\mu$ m apart were excised from each roll. BrdU positive tumor cells were measured from all the polyps in both sections.

The results are presented in Table 3, as the percentage of BrdU positive cells detected. As shown in Table 3, there was a decrease in the percentage of positive cells in the ISIS 399479-treated mice compared to the control. This result corresponds with the reduction in MALAT-1 expression, measured using ViewRNA software and as shown in Table 4. The data is presented as the signal density of MALAT-1 mRNA expression levels multiplied by the total BrdU positive cells, presented in arbitrary units.

Therefore, treatment of ApcMin mice with antisense oligonucleotides targeting MALAT-1 shows decrease in tumor cell proliferation compared to the control group.

**Table 3**

BrdU positive cells (%) in ApcMin mice

	%
ISIS 399479	23
ISIS 141923	51
PBS	48

**Table 4**

MALAT-1 expression levels in ApcMin mice (signal density of MALAT-1 mRNA expression levels multiplied by the total BrdU positive cells presented as arbitrary units (a.u.))

	a.u.
ISIS 399479	9.6
ISIS 141923	16.2

*Measurement of tumor polyps load*

The entire small intestine from each animal was made in a 'Swiss roll' and processed for paraffin embedding. Two sections of the intestine at least 500  $\mu$ m apart were collected from each roll. The sections were stained with haematoxylin and eosin, and the number of polyps were counted in both sections, and then divided by two. As shown in Table 5, tumor polyps load was decreased in ApcMin mice treated with antisense oligonucleotides targeting MALAT-1 compared to that of the control oligonucleotide group.

**Table 5**

Tumor polyps load in ApcMin mice

	Polyps/animal
ISIS 399479	1.6
ISIS 141923	4.5
PBS	5.8

### Example 3: Effect of antisense inhibition of MALAT-1 in a DEN-induced hepatocellular carcinoma mouse model

Diethyl nitrosamine (DEN) is a standard chemical carcinogen for inducing hepatocellular carcinoma (HCC) in rodents (Park, D.H. et al., Toxicol. Lett. 2009. 191: 321-6). The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on HCC development and progression was examined in this model.

#### Treatment

C57BL/6 male mice, 2 weeks in age, were given 25 mg/ml of DEN via intraperitoneal injection to induce HCC development. Five months after the DEN injection, the mice were randomly divided into three treatment groups. The first treatment group was given 50 mg/kg of ISIS 399479, administered subcutaneously twice a week for 16 weeks. The second treatment group was injected with 50 mg/kg of control oligonucleotide, ISIS 141923, administered subcutaneously twice a week for 16 weeks. The third group was injected with PBS, administered subcutaneously twice a week for 16 weeks. Mice were sacrificed on day 111 (first oligonucleotide treatment counted as day 1). DEN-induced HCCs on the liver surface were counted and collected and processed for further analysis.

#### RNA analysis

RNA was isolated from each HCC using an RNA extraction kit from Qiagen. MALAT-1 RNA expression was measured by qPCR using primer probe set mMALAT1#2 and normalized to cyclophilin mRNA expression.

MALAT-1 RNA expression in mice treated with ISIS 399479 was significantly inhibited by 94% compared to the PBS-treated group.

#### *Measurement of tumor load*

The tumors of the surface of the liver in mice from all groups were counted at the end of experiment. As shown in Table 6, tumor number was significantly decreased in mice treated with antisense oligonucleotides targeting MALAT-1 compared to that of PBS group ( $p=0.017$ ) or the control oligonucleotide group ( $p=0.02$ ).

**Table 6**

**Tumor number in the DEN-induced HCC mouse model**

Treatment groups	Tumors/mouse
ISIS 399479	1.9
ISIS 141923	4.0
PBS	4.3

#### **Example 4: Effect of antisense inhibition of MALAT-1 in a C26 xenograft mouse model**

The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on HCC progression was examined in murine C26 colon cancer xenograft model.

#### *Treatment*

C26 colon carcinoma cells were cultured in RPMI medium containing fetal bovine serum at a final concentration of 10%, and with 5% CO<sub>2</sub> at 37°C. Five million cells were subcutaneously implanted in male CD2F1 mice.

Four days after tumor implantation, the mice were randomly divided into three treatment groups. The first treatment group was injected with 50 mg/kg of ISIS 399462 (SEQ ID NO: 14), administered subcutaneously 5 days per week for 3 weeks. The second treatment group was injected with 50 mg/kg of ISIS 395251 (SEQ ID NO: 13), administered subcutaneously 5 days per week for 3 weeks. The third treatment group was injected with 50 mg/kg of control oligonucleotide, ISIS 347526 (TCTTATGTTTCCGAACCGTT; 5-10-5 MOE gapmer with no known murine or human target) (SEQ ID NO: 23) administered subcutaneously 5 days per week for 3 weeks. Mice were sacrificed on day 26. Liver and tumor tissue were collected and processed for further analysis. The data presented is the average of 2 independent experiments with similar results.

#### *RNA analysis*

RNA extraction and analyses was performed using an RNA extraction kit from Qiagen. MALAT-1 RNA expression was measured using primer probe set mMALAT1 (forward sequence GTAGGTTAAGTTGACGGCCGTTA, designated herein as SEQ ID NO: 24; reverse sequence ATCTTCCCTGTTTCCAACATCATG, designated herein as SEQ ID NO: 25; probe sequence  
 5 AAAAATCCTTCGACTGGCGCATGTACG, designated herein as SEQ ID NO: 26) and normalized to Cyclophilin mRNA expression.

MALAT-1 RNA expression was assessed in the liver. As shown in Table 7, MALAT-1 RNA expression in mice treated with either ISIS 395251 or ISIS 399462 was inhibited compared to the control oligonucleotide-treated group. The mRNA expression levels are expressed as percent inhibition of  
 10 expression levels compared to that in the control group (normalized to 0%).

**Table 7**

Percent inhibition of MALAT-1 RNA levels (%) compared to the PBS control

	% inhibition
ISIS 395251	57
ISIS 399462	71

*Measurement of tumor weight and volume*

15 Tumor volumes were measured on a regular basis throughout the study period, using Vernier calipers. As shown in Table 8, tumor volumes were significantly decreased in mice treated with ISIS 395251 or ISIS 399462.

The weight of the tumor in each mouse from all groups was also assessed on day 26. As shown in Table 9, tumor weight was decreased in mice treated with antisense oligonucleotides targeting MALAT-1  
 20 compared to that of the control oligonucleotide group. These results demonstrate that antisense oligonucleotides targeting MALAT-1 reduced colon cancer growth.

**Table 8**

Tumor volume (mm<sup>3</sup>) in the C26 xenograft model

ISIS No	day 8	day 11	day 14	day 19	day 22	day 25
399462	406	615	960	1365	1802	2275



395251	341	501	877	1480	1635	1641
347526	408	803	1493	2437	2647	3405

**Table 9**

Tumor weight in the C26 xenograft model

ISIS No	Weight (g)
395251	1.6
399462	1.7
347526	2.6

#### 5 **Example 5: Effect of antisense inhibition of MALAT-1 in a Hep3B liver orthotopic mouse model**

An orthotopic xenograft tumor model of hepatocellular carcinoma created by injection of Hep3B cells directly into the liver parenchyma of nude mice is a standard model for studying HCC (Yao, X. et al., Clin. Cancer Res. 2003. 9: 2719). The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on animal survival was examined in the Hep3B liver orthotopic model.

#### 10 *Treatment*

The human HCC cell line Hep3B was purchased from ATCC. Hep3B cells were maintained in MEM media containing fetal bovine serum at a final concentration of 10%, and with 5% CO<sub>2</sub> at 37°C. Exponentially growing Hep3B cells were collected by trypsin-EDTA (Gibco-BRL) treatment and washed once with PBS. The cell pellet was suspended in PBS and kept in ice before intrahepatic injection in mice.

15 Female BALB/c athymic (nu/nu) nude mice, 4-6 weeks in age, were anesthetized with isoflurane. A small transverse incision below the sternum was made to expose the liver. A PBS suspension of  $2 \times 10^6$  Hep3B cells was slowly injected into the upper left lobe of the liver using a 28-gauge needle. The cells were injected at a 30-degree angle into the liver. After injection, a small piece of sterile gauze was placed on the injection site, and light pressure was applied for 1 min to prevent bleeding. The abdomen was then closed  
20 with a 6-0 silk suture. The mice were allowed to recover in a warm cage.

After 10 days, the expression level of alpha-fetoprotein (AFP), which used as a positive marker of tumor growth in the mice, was analyzed. Those mice that tested positive for the marker were then randomly divided into two treatment groups. The first treatment group was injected with 50 mg/kg of ISIS 395251

(SEQ ID NO: 13), administered intraperitoneally 2 days per week for 7 weeks. The second treatment group was injected with 50 mg/kg of control oligonucleotide, ISIS 347526 (SEQ ID NO: 23) administered intraperitoneally 2 days per week for 7 weeks.

#### *Median survival*

Each group was monitored and deaths were recorded. At the end of the study, the median survival of each group was calculated using the statistical formula of Kaplan-Meier and the data is presented in Table 10. As shown in Table 10, the median survival was significantly increased in mice treated with ISIS 395251 compared to the control group. These data demonstrate that antisense oligonucleotides targeting MALAT-1 increased survival of animals having cancer.

**Table 10**

Median survival in the Hep3B liver orthotopic model

ISIS No	days
395251	88
347526	49

#### **Example 6: Effect of antisense inhibition of MALAT-1 in a metastatic EBC-1 xenograft mouse model**

The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on metastasis was examined in EBC-1 lung cancer xenograft mouse model.

#### *Treatment*

The human cell line EBC-1 was purchased from the Health Sciences Foundation, Japan. EBC-1 cells were maintained in RPMI media containing fetal bovine serum at a final concentration of 10%, and with 5% CO<sub>2</sub> at 37°C. Exponentially growing EBC-1 cells were collected by trypsin-EDTA (Gibco-BRL) and washed once with PBS. The cell pellet was suspended in PBS and one million cells were implanted by subcutaneous injection into BALB/c nude mice.

Two weeks after implantation of the EBC-1 human tumor, the mice were randomly divided into two treatment groups. The first treatment group was injected with 50 mg/kg 5 days a week of ISIS 395251 (SEQ ID NO: 13), administered subcutaneously for 5 weeks. The second treatment group was injected with 50 mg/kg 5 days a week of control oligonucleotide, ISIS 347526 (SEQ ID NO: 23) administered subcutaneously for 5 weeks. On week 7, the subcutaneous tumor was surgically removed and the wound closed with a 4-0 suture. Mice were euthanized on week 12 after the start of antisense oligonucleotide treatment. Lung tissue was collected and processed for further analysis. The data presented is the average of 3 independent experiments with similar results.

*Measurement of primary tumor volume and lung tumor multiplicity*

Primary tumor volumes were measured along the course of the study period, using Vernier calipers. As shown in Table 11, at week 7 right before removal of the xenograft, tumor volumes were significantly decreased in mice treated with ISIS 395251 compared to the control group (p=0.00006).

- 5 The tumor multiplicity was also counted using a light microscope. As shown in Table 12, the tumor multiplicity was decreased in mice treated with ISIS 395251 compared to the control. These data demonstrate that antisense oligonucleotides targeting MALAT-1 inhibited metastasis.

**Table 11**

Tumor volume in the EBC-1 xenograft model

ISIS No	Volume (mm <sup>3</sup> )
395251	1755
347526	2838

10

**Table 12**

Tumor multiplicity in the EBC-1 xenograft model

ISIS No	Tumor count
395251	68
347526	167

**Example 7: Effect of antisense inhibition of MALAT-1 in the TRAMP mouse model**

- 15 The transgenic adenocarcinoma of the mouse prostate (TRAMP) model closely mirrors the pathogenesis of human prostate cancer (Hurwitz, A.A. et al., Curr. Protoc. Immunol. 2001. Chapter 20: Unit 20.5). The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on tumor progression was examined in TRAMP mice.

*Treatment*

- 20 TRAMP mice, 23 weeks old, were randomly divided into two treatment groups. The first treatment group was injected with 50 mg/kg of ISIS 395251 (SEQ ID NO: 13), administered subcutaneously 5 days

per week for 3 weeks. The second treatment group was injected with PBS administered subcutaneously 5 days per week for 3 weeks. Mice were sacrificed at the end of 26 weeks. Prostate tissue were collected and processed for further analysis.

#### *RNA analysis*

5 RNA extraction was performed using an RNA extraction kit from Qiagen. MALAT-1 RNA expression was measured using primer probe set mMALAT1 and normalized to Cyclophilin mRNA expression.

MALAT-1 RNA expression was assessed in the tumor. MALAT-1 RNA expression in mice treated with ISIS 395251 was inhibited by 80% compared to the control group.

#### 10 *Measurement of tumor weight*

Tumor tissue was excised from the prostate. Tumor weights were measured on a regular basis throughout the study period, using Vernier calipers. As shown in Table 13, tumor weight was decreased in mice treated with ISIS 395251 compared to the control group.

**Table 13**

15

**Tumor weight in the TRAMP**

	Weight (g)
ISIS 395251	1.0
PBS	9.3

#### **Example 8: Effect of antisense inhibition of MALAT-1 in a patient-derived non-small cell lung cancer xenograft mouse model**

20 Biopsy of tumor mass was done in a non-small cell lung cancer patient (at the University of California, Davis) and this was directly implanted into male NOD.Cg-Prkdc<sup>scid</sup>I12rg<sup>tm1Wjl</sup>/SzJ (NSG; Jackson Laboratories) mice. After 2 *in vivo* passages, the tumor cells from the xenograft were banked at Jackson Laboratories (designated herein as LG-476 P2). The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on tumor progression of this tumor in mice was examined.

#### *Treatment*

25 Two NSG mice were implanted with the LG-476 P2 tumor and monitored three times weekly. Once the tumors reached 1,000 mm<sup>3</sup> in volume, the tumors were harvested and fragmented into 3-5 mm<sup>3</sup> size.

Each fragment was then implanted subcutaneously into the right hind flank of 30 NSG mice. The mice were observed three times a week and once the tumors reached 200-250 mm<sup>3</sup> in size, the mice were randomly divided into two treatment groups. The first treatment group was injected with 50 mg/kg of ISIS 395251 (SEQ ID NO: 13), administered subcutaneously 5 days per week for 3 weeks. The second treatment group was injected with PBS administered subcutaneously 5 days per week for 3 weeks. Mice were euthanized by CO<sub>2</sub> inhalation 24 hrs after the last dose. Tumor tissue were collected and processed for further analysis.

#### *RNA analysis*

RNA extraction was performed using an RNA extraction kit from Qiagen. MALAT-1 RNA expression was measured using primer probe set RTS2736 and normalized to Cyclophilin mRNA expression.

MALAT-1 RNA expression was assessed in the tumor. MALAT-1 RNA expression in mice treated with ISIS 395251 was inhibited by 76% compared to the control group.

#### *Measurement of tumor volume*

Tumor volumes were measured on a regular basis throughout the study period, using Vernier calipers. As shown in Table 14, tumor volumes were decreased in mice treated with ISIS 395251 compared to the control group.

**Table 14**

Tumor volume (mm<sup>3</sup>) on different days in the small cell lung cancer xenograft model

	Day 1	Day 3	Day 6	Day 8	Day 10	Day 13	Day 15	Day 17	Day 20
ISIS 395251	227	376	391	448	529	681	661	747	715
PBS	225	314	384	494	568	696	917	1047	2049

#### **Example 9: Effect of antisense inhibition of MALAT-1 in a Colo201 xenograft mouse model**

The effect of inhibition of MALAT-1 RNA expression with antisense oligonucleotides on tumor progression was examined in the Colo201 xenograft mouse model.

#### *Treatment*

The human colorectal adenocarcinoma cell line Colo201 was purchased from ATCC. Colo201 cells were maintained in RPMI media containing fetal bovine serum at a final concentration of 10%, and with 5%

CO<sub>2</sub> at 37°C. Exponentially growing Colo201 cells were collected by trypsin-EDTA (Gibco-BRL) and washed once with PBS. The cell pellet was suspended in PBS and kept in ice before intrahepatic injection in female BALB/c nude mice.

Four days after tumor implantation, the mice were randomly divided into three treatment groups.

- 5 The first treatment group was injected with 50 mg/kg of ISIS 395251 (SEQ ID NO: 13), administered subcutaneously 5 days per week for 3.5 weeks. The second treatment group was injected with 50 mg/kg of ISIS 347526 (SEQ ID NO: 23), administered subcutaneously 5 days per week for 3.5 weeks. The third group was injected with PBS administered subcutaneously 5 days per week for 3.5 weeks. Mice were sacrificed on day 29. Tumor tissue were collected and processed for further analysis.

#### 10 *RNA analysis*

RNA extraction was performed using an RNA extraction kit from Qiagen. MALAT-1 RNA expression was measured using primer probe set RTS2736 and normalized to Cyclophilin mRNA expression.

- 15 MALAT-1 RNA expression was assessed in the liver. MALAT-1 RNA expression in mice treated with ISIS 395251 was inhibited by 39% compared to the control group.

#### *Measurement of tumor volume*

- 20 Tumor volumes were measured on a regular basis throughout the study period, using Vernier calipers. As shown in Table 15, tumor volumes were decreased in mice treated with ISIS 395251 compared to the control group.

**Table 15**

Tumor volume on different days in the Colo201 cancer xenograft model

	Day 4	Day 7	Day 10	Day 15	Day 18	Day 21	Day 24	Day 29
ISIS 395251	138	132	186	207	241	247	288	341
ISIS 347526	148	156	209	238	354	439	428	476
PBS	159	142	184	240	373	393	404	495

## Claims

1. A single-stranded modified oligonucleotide for use in treating a primary cancer in an animal, wherein the single-stranded modified oligonucleotide consists of 12 to 30 linked nucleosides and has a nucleobase sequence at least 85% complementary over its entire length to a human Metastasis-Associated-in-Lung-Adenocarcinoma-Transcript-1 (MALAT-1) nucleic acid as recited in any one of SEQ ID NOs: 1-9 and wherein the single-stranded modified oligonucleotide is capable of reducing levels of MALAT-1 RNA in the animal.
2. The single-stranded modified oligonucleotide for use according to claim 1, wherein the single-stranded modified oligonucleotide is 100% complementary over its entire length to a human MALAT-1 nucleic acid recited in any one of SEQ ID NOs: 1-9.
3. The single-stranded modified oligonucleotide for use according to claim 1 or 2, wherein the single-stranded modified oligonucleotide is capable of inhibiting growth of the primary cancer in the animal.
4. The single-stranded modified oligonucleotide for use according to any one of claims 1-3, wherein the primary cancer is colon cancer, lung cancer, liver cancer, prostate cancer, or intestinal cancer.
5. The single-stranded modified oligonucleotide for use according to any one of claims 1-4, wherein at least one internucleoside linkage of the single-stranded modified oligonucleotide is a modified internucleoside linkage.
6. The single-stranded modified oligonucleotide for use according to claim 5 wherein the modified internucleoside linkage is a phosphorothioate internucleoside linkage.
7. The single-stranded modified oligonucleotide for use according to any one of claims 1-6, wherein at least one nucleoside of the single-stranded modified oligonucleotide comprises a modified sugar.
8. The single-stranded modified oligonucleotide for use according to claim 7, wherein the modified sugar is a bicyclic sugar comprising a 4'-CH(CH<sub>3</sub>)-O-2', 4'-(CH<sub>2</sub>)-O-2', or 4'-(CH<sub>2</sub>)<sub>2</sub>-O-2' bridge.
9. The single-stranded modified oligonucleotide for use according to claim 7, wherein the modified sugar comprises a 2'-O-methoxyethyl group.
10. The single-stranded modified oligonucleotide for use according to any one of claims 1-9, wherein at least one nucleoside of the single-stranded modified oligonucleotide comprises a modified nucleobase.

11. The single-stranded modified oligonucleotide for use according to claim 10, wherein the single-stranded modified nucleobase is 5-methylcytosine.

12. The single-stranded modified oligonucleotide for use according to claim 11, wherein the single-stranded modified oligonucleotide comprises:

a gap segment consisting of linked deoxynucleosides;

a 5' wing segment consisting of linked nucleosides; and

a 3' wing segment consisting of linked nucleosides, wherein the gap segment is positioned immediately adjacent to and between the 5' wing segment and the 3' wing segment and wherein each nucleoside of each wing segment comprises a modified sugar.

13. Use of a single-stranded modified oligonucleotide in the manufacture of a medicament for treating a primary cancer in an animal, wherein the single-stranded modified oligonucleotide consists of 12 to 30 linked nucleosides and has a nucleobase sequence at least 85% complementary over its entire length to a human Metastasis-Associated-in-Lung-Adenocarcinoma-Transcript-1 (MALAT-1) nucleic acid as recited in any one of SEQ ID NOs: 1-9 and wherein the single-stranded modified oligonucleotide is capable of reducing levels of MALAT-1 RNA in the animal.

14. The use according to claim 13, wherein the single-stranded modified oligonucleotide is 100% complementary over its entire length to a human MALAT-1 nucleic acid recited in any one of SEQ ID NOs: 1-9.

15. The use according to claim 13 or 14, wherein the single-stranded modified oligonucleotide is capable of inhibiting growth of the primary cancer.

16. The use according to any one of claims 13-15, wherein the primary cancer is colon cancer, lung cancer, liver cancer, prostate cancer, or intestinal cancer.

17. The use according to any one of claims 13-16, wherein at least one internucleoside linkage of the single-stranded modified oligonucleotide is a modified internucleoside linkage.

18. The use according to claim 17, wherein the modified internucleoside linkage is a phosphorothioate internucleoside linkage.

19. The use according to any one of claims 13-18, wherein at least one nucleoside of the single-stranded modified oligonucleotide comprises a modified sugar.

20. The use according to claim 19, wherein the modified sugar is a bicyclic sugar comprising a 4'-CH(CH<sub>3</sub>)-O-2', 4'-(CH<sub>2</sub>)-O-2', or 4'-(CH<sub>2</sub>)<sub>2</sub>-O-2' bridge.



21. The use according to claim 19, wherein the modified sugar comprises a 2'-O-methoxyethyl group.

22. The use according to any one of claims 13-21, wherein at least one nucleoside of the single-stranded modified oligonucleotide comprises a modified nucleobase.

23. The use according to claim 22, wherein the modified nucleobase is a 5-methylcytosine.

24. The use according to claim 23, wherein the single-stranded modified oligonucleotide comprises:

gap segment consisting of linked deoxynucleosides;

a 5' wing segment consisting of linked nucleosides; and

a 3' wing segment consisting of linked nucleosides, wherein the gap segment is positioned immediately adjacent to and between the 5' wing segment and the 3' wing segment and wherein each nucleoside of each wing segment comprises a modified sugar.