

**(12) PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 200059664 B2  
(10) Patent No. 779495**

(54) Title  
**Tuberculosis vaccine and diagnostics based on the mycobacterium tuberculosis esat-6 gene family**

(51)<sup>7</sup> International Patent Classification(s)  
**C07K 014/35 C07K 019/00  
A61K 039/04**

(21) Application No: **200059664** (22) Application Date: **2000.07.13**

(87) WIPO No: **WO01/04151**

(30) Priority Data

(31) Number **1999 01020** (32) Date **1999.07.13** (33) Country **DK**  
**60/144011** **1999.07.15** **US**

(43) Publication Date : **2001.01.30**

(43) Publication Journal Date : **2001.04.12**

(44) Accepted Journal Date : **2005.01.27**

(71) Applicant(s)  
**Statens Serum Institut**

(72) Inventor(s)  
**Peter Andersen; Rikke Skjot**

(74) Agent/Attorney  
**F.B. Rice and Co.,139 Rathdowne Street,CARLTON VIC 3053**

(56) Related Art  
**WO 1998/053076**  
**WO 1998/053076**  
**NATURE 1998 VOL. 393 PP 537-544**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number  
WO 01/04151 A2

(51) International Patent Classification<sup>2</sup>: C07K 14/35, 19/00, A61K 39/04

(21) International Application Number: PCT/DK00/00398

(22) International Filing Date: 13 July 2000 (13.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PA 1999 01020 13 July 1999 (13.07.1999) DK  
60/144,011 15 July 1999 (15.07.1999) US

(71) Applicant (for all designated States except US):  
STATENS SERUM INSTITUT [DK/DK]; Artillerivej 5,  
DK-2300 Copenhagen S (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ANDERSEN, Peter [DK/DK]; Sparreholmvej 47, DK-2700 Brønshøj (DK).  
SKJØT, Rikke [DK/DK]; L.A. Rings Vænge 17, DK-2640  
Hedehusenc (DK).

(74) Agent: PLOUGMANN, VINGTOFT & PARTNERS;  
Sankt Annæ Plads 11. P.O. Box 3007, DK-1021 Copenhagen K (DK).

(81) Designated States (national): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (utility model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/04151 A2

(54) Title: TUBERCULOSIS VACCINE AND DIAGNOSTICS BASED ON THE *MYCOBACTERIUM TUBERCULOSIS esat-6* GENE FAMILY

(57) Abstract: This invention relates to a polypeptide fragment which comprises an amino acid sequence encoded by a member of the *esat-6* gene. A member of the *esat-6* gene family is defined as gene encoding a small protein and that two such genes are arranged next to each other on the genome and that at least one of the gene products has an amino acid sequence identity to either Rv3874, Rv3875, or Rv0288 of at least 15 %.

TUBERCULOSIS VACCINE AND DIAGNOSTICS BASED ON THE *MYCOBACTERIUM*  
*TUBERCULOSIS* *esat-6* GENE FAMILY

## FIELD OF THE INVENTION

The present invention relates to a number of immunologically active, novel polypeptide fragments derived from the *Mycobacterium tuberculosis*, vaccines and other immunologic compositions containing the fragments as immunogenic components, and methods of production and use of the polypeptides. The invention also relates to novel nucleic acid fragments derived from *M. tuberculosis* which are useful in the preparation of the polypeptide fragments of the invention or in the diagnosis of infection with *M.*

10 *tuberculosis*.

## BACKGROUND OF THE INVENTION

Human tuberculosis caused by *Mycobacterium tuberculosis* is a severe global health problem responsible for approx. 3 million deaths annually, according to the WHO. The world-wide incidence of new TB cases has been progressively falling for the last decade 15 but during the recent years this trend has markedly changed due to the advent of AIDS and the appearance of multidrug resistant strains of *M. tuberculosis*.

The only vaccine presently available for clinical use is BCG, a vaccine which efficacy remains a matter of controversy. BCG generally induces a high level of acquired resistance in animal models of TB, but several human trials in developing countries have failed to demonstrate significant protection. Notably, BCG is not approved by the FDA for use in the United States because BCG vaccination impairs the specificity of the Tuberculin skin test for diagnosis of TB infection.

25 This makes the development of a new and improved vaccine against TB an urgent matter which has been given a very high priority by the WHO. Many attempts to define protective mycobacterial substances have been made, and from 1950 to 1970 several investigators reported an increased resistance after experimental vaccination. However, the demonstration of a specific long-term protective immune response with the potency of 30 BCG has not yet been achieved.

Immunity to *M. tuberculosis* is characterized by three basic features; 1) Living bacilli efficiently induces a protective immune response; 2) Specifically sensitized T lymphocytes mediate this protection, and 3) The most important mediator molecule seems to be interferon gamma (IFN- $\gamma$ ).

5

*M. tuberculosis* holds, as well as secretes, several proteins of potential relevance for the generation of a new tuberculosis (TB) vaccine. For a number of years a major effort has been put into the identification of new protective antigens for the development of a new vaccine against TB. The search for candidate molecules has primarily focused on

10 proteins released from the dividing bacteria. Several molecules have been identified and characterized from this mycobacterial protein fraction. One low molecular mass protein from culture filtrate, ESAT-6, has been found to be an extraordinary potent IFN- $\gamma$  inducer, when used to stimulate human Peripheral Blood Mononuclear Cells (PBMC) from tuberculosis (TB) patients (Ravn et al. 1999).

15

The total sequencing of the *M. tuberculosis* genome has led to several important findings. A subject of interest was the finding that the potent T-cell antigen ESAT-6 was transcribed together with another low mass protein (CFP10). The genes encoding these two proteins were in other words found next to each other on the mycobacterial genome,

20 located in the same operon and were regulated by the same promoter. The two genes have a sequence identity of approximately 40%. On amino acid level, the sequence identity was approximately 15%. The proteins have approx. same size and pl.

Together with a number of putative Open Reading Frames (ORFs) these two molecules 25 constitute what has been called the *esat-6* gene family (Cole et al. 1998, Berthet et al 1998). All the genes in this family encode low mass proteins, which are placed in operon like structures as are ESAT-6 and CFP10. The family was first described by Cole et al. 1998 with the following words: "The potent T-cell antigen ESAT-6, which is probably secreted in a Sec-independent manner, is encoded by a member of a multigene family.

30 Examination of the genetic context reveals several similarly organized operons that include genes encoding large ATP-hydrolysing membrane proteins that might act as transporters." and was later described by Berthet et al. 1998 as follows: "Several genes sharing weak similarities with *esat-6* have been previously identified during the *M. tuberculosis* genome sequencing project. Although these genes share less than 35% 35 sequence similarity with each other, they were grouped in the *esat-6* gene family since

they all potentially code for small polypeptides of about 100 amino acids. All these genes are organized in operon-like structures and are frequently preceded by genes encoding repetitive proteins of the PE and the PPE families".

5 Diagnosing *M. tuberculosis* in its earliest stage is important for effective treatment of the disease. Current diagnostic assays to determine *M. tuberculosis* infection are expensive and labor intensive. In our part of the world the majority of patients exposed to *M. tuberculosis* receive chest x-rays and attempts are made to culture the bacterium in vitro from sputum samples. X-rays as a diagnostic assay is insensitive and can only identify

10 infections in a very progressed stage. Culturing of *M. tuberculosis* is also not ideal as a diagnostic tool, since the bacteria grows poorly and slowly outside the body, which can produce false negative test results and take weeks before results are obtained. An inexpensive assay, used in third world countries, is the standard tuberculin skin test. It is far from ideal in detecting infection because it cannot distinguish *M. tuberculosis* infected

15 individuals from *M. bovis* BCG vaccinated individuals and therefore cannot be used in areas of the world where patients receive or have received childhood vaccination with bacterial strains related to *M. tuberculosis* (BCG vaccination).

## SUMMARY OF THE INVENTION

20 In the broadest aspect, the present invention relates to a substantially pure polypeptide fragment which comprises an amino acid sequence encoded by a member of the *esat-6* gene family or comprises an amino acid analogue having a sequence identity with a polypeptide fragment encoded by a member of the *esat-6* gene family of at least 70% and at the same time being immunologically equivalent to the polypeptide fragment

25 encoded by a member of the *esat-6* gene family. A member of the *esat-6* gene family is defined as gene encoding a small protein and that two such genes are arranged next to each other on the genome and that at least one of the gene products has an amino acid sequence identity to either Rv3874, Rv3875, or Rv0288 of at least 15%. Presently the following genes are members of the *esat-6* gene family: Rv0287, Rv0288, Rv1036c,

30 Rv1037c, Rv1038c, Rv1197, Rv1198, Rv1792, Rv1793, Rv2346c, Rv2347c, Rv2348c, Rv2653c, Rv2654c, Rv3019c, Rv3020c, Rv3444c, Rv3445c, Rv3619c, Rv3620c, Rv3874, Rv3875, Rv3890c, Rv3891c, Rv3904c, and Rv3905c.

These proteins have an important mycobacteria specific function which may be related to the intracellular habitat of the macrophage phagosome. Furthermore, they show high immunological efficacy, as described in examples 1, 3a and 3b. They are therefore suggested as useful candidates in a vaccine against TB or diagnostic preparation for

5 TB. The genes encoding these proteins are suggested as components in a DNA vaccine against TB.

According to the invention there is provided a fusion polypeptide, which comprises an amino acid sequence chosen from the group consisting of:

10 1) an amino acid sequence analogue having a sequence identity with SEQ ID NO: 7 of at least 70% and at the same time being immunologically equivalent to SEQ ID NO: 7;

2) a T-cell epitope of the amino acid sequence as shown in SEQ ID NO: 7; and

15 3) an amino acid sequence, which has a sequence identity of at least 70% with the T-cell epitope in 2) and at the same time being immunologically equivalent to SEQ ID NO: 7,  
and at least one fusion partner.

20 According to the invention there is also provided a substantially pure polypeptide fragment which comprises an amino acid sequence chosen from the group consisting of:

1) an amino acid sequence analogue having a sequence identity with SEQ ID NO: 7 of at least 70% and at the same time being immunologically equivalent to SEQ ID NO: 7;

25 2) a T-cell epitope of the amino acid sequence as shown in SEQ ID NO: 7; and

3) an amino acid sequence, which has a sequence identity of at least 70% with the T-cell epitope in 2) and at the same time being immunologically equivalent to SEQ ID NO: 7, when used in a pharmaceutical.

According to the invention there is also provided a use of a substantially pure polypeptide chosen from the group consisting of:

35 - the polypeptide fragment according to the invention; and  
- Rv3386

in the preparation of a pharmaceutical composition for the diagnosis of or vaccination against tuberculosis caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis* in a mammal.

According to the invention there is also provided an immunologic composition, which  
5 comprises a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386.

According to the invention there is also provided a vaccine for immunizing an animal,  
10 including a human being, against tuberculosis caused by mycobacteria belonging to the tuberculosis complex, comprising as the effective component a non-pathogenic microorganism, wherein at least one copy of a DNA fragment comprising a DNA sequence encoding a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- 15 - Rv3386

has been incorporated into the genome of the microorganism in a manner allowing the microorganism to express and optionally secrete the polypeptide.

According to the invention there is also provided a composition for diagnosing  
20 tuberculosis in an animal, including a human being, comprising a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386

optionally in combination with a means for detection.

25 According to the invention there is also provided a nucleic acid fragment which  
1) comprises a nucleic acid sequence which has a sequence as disclosed in  
SEQ ID NO: 6,  
2) has a length of at least 10 nucleotides and hybridizes under moderately  
30 stringent conditions with a nucleic acid fragment which has a sequence as  
disclosed in SEQ ID NO: 6 or a sequence complementary thereto,  
with the proviso that SEQ ID NO: 7 disclosed in WO 98/53075 is excluded, when used  
in a pharmaceutical.

According to the invention there is also provided a use of a nucleic acid fragment according to the invention in the preparation of a pharmaceutical composition for the diagnosis of or vaccination against tuberculosis caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis*.

5

According to the invention there is also provided a vaccine comprising a nucleic acid fragment according to the invention, the vaccine effecting *in vivo* expression of antigen by an animal, including a human being, to whom the vaccine has been administered, the amount of expressed antigen being effective to confer substantially increased 10 resistance to infections with mycobacteria of the tuberculosis complex in an animal, including a human being.

According to the invention there is also provided replicable expression vector which comprises a nucleic acid fragment according to the invention.

15

According to the invention there is also provided a transformed cell harbouring at least one vector according to the invention.

According to the invention there is provided a composition for diagnosing tuberculosis 20 in an animal, including a human being, comprising a nucleic acid fragment according to the invention, optionally in combination with a means for detection.

According to the invention there is also provided a method for producing a polypeptide chosen from the group consisting of:

25

- the polypeptide fragment according to the invention; and
- Rv3386,

comprising inserting a nucleic acid fragment, which

- 1) comprises a nucleic acid sequence which has a sequence as disclosed in SEQ ID NO: 6,
- 2) has a length of at least 10 nucleotides and hybridizes under moderately stringent conditions with a nucleic acid fragment which has a sequence as disclosed in SEQ ID NO: 6 or a sequence complementary thereto,

into a vector which is able to replicate in a host cell, introducing the resulting recombinant vector into the host cell, culturing the host cell in a culture medium under 35 conditions sufficient to effect expression of the polypeptide, and recovering the polypeptide from the host cell or culture medium; or

isolating the polypeptide from whole mycobacteria of the tuberculosis complex or from a lysate or fraction thereof ; or  
synthesizing the polypeptide by solid or liquid phase peptide synthesis.

- 5 According to the invention there is also provided a method for producing an immunologic composition according to the invention comprising:  
preparing, synthesizing or isolating a polypeptide chosen from the group consisting of:
  - the polypeptide fragment according to the invention; and
  - Rv3386, and
- 10 solubilizing or dispersing the polypeptide in a medium for a vaccine, and  
optionally adding other *M. tuberculosis* antigens and/or a carrier, vehicle and/or adjuvant substance, or  
cultivating a cell according to the invention, and  
transferring the cells to a medium for a vaccine, and
- 15 optionally adding a carrier, vehicle and/or adjuvant substance.

According to the invention there is also provided a method of diagnosing tuberculosis caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis* into an animal, including a human being, comprising intradermally injecting, in  
20 the animal, a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386

or an immunologic composition according to the invention, a positive skin response at the location of injection being indicative of the animal having tuberculosis, and a  
25 negative skin response at the location of injection being indicative of the animal not having tuberculosis.

According to the invention there is also provided method for immunising an animal, including a human being, against tuberculosis caused by mycobacteria belonging to the  
30 tuberculosis complex, comprising administering to the animal a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386,

the immunologic composition according to the invention, or the vaccine according to  
35 the invention.

According to the invention there is also provided a method for diagnosing ongoing or previous sensitization in an animal or a human being with bacteria belonging to the tuberculosis complex, the method comprising providing a blood sample from the animal or human being, and contacting the sample from the animal with a polypeptide

5 chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386,

a significant release into the extracellular phase of at least one cytokine by mononuclear cells in the blood sample being indicative of the animal being sensitized.

10

According to the invention there is also provided a monoclonal or polyclonal antibody, which is results with a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to the invention; and
- Rv3386

15 in an immunoassay, or a specific binding fragment of said antibody.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

20

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the specification is solely for the purpose of providing a context for the invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the 25 invention as it existed before the priority date of each claim of this application.

## DETAILED DISCLOSURE OF THE INVENTION

In the present specification and claims, the term "polypeptide fragment", or variants 30 thereof, denotes both short peptides with a length of at least two amino acid residues and at most 10 amino acid residues, oligopeptides (11-100 amino acid residues), and longer peptides. The polypeptide fragment may be chemically modified by being glycosylated, by being lipidated, or by comprising prosthetic groups.

In the present context the term "substantially pure polypeptide fragment" means a polypeptide preparation which contains at most 5% by weight of other polypeptide material with which it is natively associated (lower percentages of other polypeptide material are preferred, e. g. at most 4%, at most 3%, at most 2%, at most 1%, and at 5 most 1/2%). It is preferred that the substantially pure polypeptide is at least 96% pure, i. e. that the polypeptide constitutes at least 96% by weight of total polypeptide material present in the preparation, and higher percentages are preferred, such as at least 97%, at least 98%, at least 99%, at least 99,25%, at least 99,5%, and at least 99,75%. It is especially preferred that the polypeptide fragment is in "essentially pure form", i.e. that 10 the polypeptide fragment is essentially free of any other antigen with which it is natively associated, i e. free of any other antigen from bacteria belonging to the tuberculosis complex. This can be accomplished by preparing the polypeptide fragment by means of recombinant methods in a non-mycobacterial host cell as will be described in detail below, or by synthesizing the polypeptide fragment by the well-known 15 methods of solid or liquid phase peptide synthesis, e. g. by the method described by Merrifield or variations thereof.

The "tuberculosis-complex" has its usual meaning, ie. the complex of mycobacteria causing TB which are *Mycobacterium tuberculosis*, *Mycobacterium bovis*, 20 *Mycobacterium bovis* BCG, and *Mycobacterium africanum*.

By the term "virulent *Mycobacterium*" is understood a bacterium capable of causing the tuberculosis disease in a mammal including a human being. Examples of virulent *Mycobacteria* are *M. tuberculosis*, *M. africanum*, and *M. bovis*.

- 5 By "a TB patient" is understood an individual with culture or microscopically proven infection with virulent *Mycobacteria*, and/or an individual clinically diagnosed with TB and who is responsive to anti-TB chemotherapy. Culture, microscopy and clinical diagnosis of TB is well known by the person skilled in the art.
- 10 By the term "PPD positive individual" is understood an individual with a positive Mantoux test or an individual where PPD induces an increase in *in vitro* recall response determined by release of IFN- $\gamma$  of at least 1,000 pg/ml from Peripheral Blood Mononuclear Cells (PBMC) or whole blood, the induction being performed by the addition of 2.5 to 5  $\mu$ g of PPD/ml to a suspension comprising about 1.0 to 2.5  $\times$  10<sup>5</sup> PBMC, the
- 15 release of IFN- $\gamma$  being assessable by determination of IFN- $\gamma$  in supernatant harvested 5 days after the addition of PPD to the suspension compared to the release of IFN- $\gamma$  without the addition of PPD.

By the term "delayed type hypersensitivity reaction" is understood a T-cell mediated inflammatory response elicited after the injection of a polypeptide into or application to the skin, said inflammatory response appearing 72-96 hours after the polypeptide injection or application.

- 20

By the term "IFN- $\gamma$ " is understood interferon-gamma.

25

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations thereof such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

30

The term "sequence identity" indicates a quantitative measure of the degree of homology between two amino acid sequences of equal length or between two nucleotide sequences of equal length. If the two sequences to be compared are not of equal length, they must be aligned to best possible fit. The sequence identity can be calculated as

$\frac{(N_{ref} - N_{dif}) * 100}{N_{ref}}$ , wherein  $N_{dif}$  is the total number of non-identical residues in the two sequences

when aligned and wherein  $N_{ref}$  is the number of residues in one of the sequences. Hence, the DNA sequence AGTCAGTC will have a sequence identity of 75% with the sequence AATCAATC ( $N_{dif}=2$  and  $N_{ref}=8$ ). A gap is counted as non-identity of the specific

5 residue(s), i.e. the DNA sequence AGTGTC will have a sequence identity of 75% with the DNA sequence AGTCAGTC ( $N_{dif}=2$  and  $N_{ref}=8$ ). Sequence identity can alternatively be calculated by the BLAST program e.g. the BLASTP program (Pearson W.R and D.J. Lipman (1988) PNAS USA 85:2444-2448)([www.ncbi.nlm.nih.gov/cgi-bin/BLAST](http://www.ncbi.nlm.nih.gov/cgi-bin/BLAST)). In one aspect of the invention, alignment is performed with the global align algorithm with default 10 parameters as described by X. Huang and W. Miller. Adv. Appl. Math. (1991) 12:337-357, available at [http://www.ch.embnet.org/software/LALIGN\\_form.html](http://www.ch.embnet.org/software/LALIGN_form.html).

A preferred minimum percentage of sequence identity is at least 80%, such as at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, 15 at least 96%, at least 97%, at least 98%, at least 99%, and at least 99.5%.

The *M. tuberculosis* antigens provided herein include variants that are encoded by DNA sequences which are substantially homologous to one or more of the DNA sequences specifically recited herein. Sequence identity as used herein, refers to DNA sequences 20 that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5 % SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50-60°C, 5X SSC overnight or, in the case of cross-species homology at 45°C, 0.5XSSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1 % SDS. Such hybridizing 25 DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

Each polypeptide fragment may thus be characterized by specific amino acid and nucleic 30 acid sequences. It will be understood that such sequences include analogues and variants produced by recombinant methods wherein such nucleic acid and polypeptide sequences have been modified by substitution, insertion, addition and/or deletion of one or more nucleotides in said nucleic acid sequences to cause the substitution, insertion, addition or deletion of one or more amino acid residues in the recombinant polypeptide.

When the term nucleotide is used in the following, for a number of purposes it can be understood as DNA, RNA, PNA or LNA equally. However, as the person skilled in the art will realise, obvious restrictions apply. PNA or LNA may be used instead of DNA. PNA has been shown to exhibit a very dynamic hybridization profile (PNA is described in 5 Nielsen P E et al., 1991, Science 254: 1497-1500). LNA (Locked Nucleic Acids) is a recently introduced oligonucleotide analogue containing bicyclo nucleoside monomers (Koshkin et al., 1998, 54, 3607-3630; Nielsen, N.K. et al. J.Am.Chem.Soc 1998, 120, 5458-5463).

10 The *esat-6* gene family consist of genes, wherein criteria a) through c) below are satisfied:

- a) genes coding for small proteins;
- b) at least two such genes are arranged next to each other on the genome;
- c) at least one of the gene products in criteria b) has an amino acid sequence identity to 15 either Rv3874 (SEQ ID NO: 1), Rv3875 (SEQ ID NO: 2), or Rv0288 (SEQ ID NO: 3) of at least 15%.

One common denominator of the *esat-6* family is the small size of the protein products of the genes. In this context, a small protein is about 80 amino acids, e.g. about 90 amino 20 acids, about 100 amino acids, about 110 amino acids, about 120 amino acids, about 130 amino acids, about 140 amino acids, or about 150 amino acids.

These proteins have an important mycobacteria specific function which may be related to the intracellular habitat of the macrophage phagosome and might suggest that the 25 expression of these molecules are synchronously upregulated in a particular phase of the infection during intracellular growth. This upregulation is a possible explanation of the high antigenicity of the proteins in this family. The amino acid sequence identity in criteria c) is preferably more than 15 %, such as more than 20%, e.g. more than 25%, 30% or even more than 35%.

30

Presently, the following genes satisfy criteria a) through c) above and are thus identified members of the *esat-6* gene family (see Table 1):  
Rv0287, Rv0288 (TB10.4), Rv1036c, Rv1037c, Rv1038c, Rv1197, Rv1198, Rv1792, Rv1793, Rv2346c, Rv2347c, Rv2348c, Rv2653c, Rv2654c, Rv3019c, Rv3020c.

Rv3444c, Rv3445c, Rv3619c, Rv3620c, Rv3874 (CFP10), Rv3875 (ESAT-6), Rv3890c, Rv3891c, Rv3904c, and Rv3905c.

As disclosed in example 1, CFP10, ESAT-6, and TB10.4 (previously named CFP7) are

5 exceedingly good IFN- $\gamma$  inducers, when the purified recombinant antigens are used to stimulate PBMC's from human TB patients. Such an effect (as described in criteria d) ii) below) is an important first test, prior to the determination of whether this protein should be further developed as a component in a vaccine or diagnostic composition.

10 Interestingly, also recombinant Rv1793 and synthetic peptides derived from this protein and from Rv0287, both members of the esat-6 gene family, stimulated T-cell proliferation and IFN- $\gamma$  production in PBMCs from two PPD-positive donors (WO98/53075 and WO98/53076).

15 In one embodiment of the present invention, the protein product of a member of the esat-6 family should further satisfy criteria d):

d) At least one of the properties below should be positive:

i) it induces an *in vitro* response during primary infection with virulent *Mycobacteria*,

20 determined by release of IFN- $\gamma$  of at least 1,500 pg/ml from T-lymphocytes withdrawn from a mouse within 28 days after the mouse has been infected with  $5 \times 10^4$  virulent *Mycobacteria*, the induction being performed by the addition of the polypeptide to a suspension comprising about  $2 \times 10^5$  cells isolated from the spleen, the addition of the polypeptide resulting in a concentration of not more than 20  $\mu$ g per ml suspension, the

25 release of IFN- $\gamma$  being assessable by determination of IFN- $\gamma$  in supernatant harvested 3 days after the addition of the polypeptide to the suspension,

ii) it induces *in vitro* recall response determined by release of IFN- $\gamma$  of at least 500 pg/ml, preferably 1,000 pg/ml from Peripheral Blood Mononuclear Cells (PBMC) or whole blood

30 withdrawn from TB patients 0-6 months after diagnosis, or PPD positive individuals, the induction being performed by the addition of the polypeptide to a suspension comprising about 1.0 to  $2.5 \times 10^5$  PBMC or whole blood cells, the addition of the polypeptide resulting in a concentration of not more than 20  $\mu$ g per ml suspension, the release of IFN- $\gamma$  being assessable by determination of IFN- $\gamma$  in supernatant harvested 5 days after

35 the addition of the polypeptide to the suspension,

iii) it induces a specific antibody response in a TB patient as determined by an ELISA technique or a western blot when the serum is diluted 1:20 in PBS and incubated with the polypeptide in a concentration of at the most 20  $\mu$ g/ml and induces an OD of at least 0.1

5 in ELISA, or a visual response in western blot,

iv) it induces a positive *in vitro* response determined by release of IFN- $\gamma$  of at least 500 pg/ml from Peripheral Blood Mononuclear Cells (PBMC) withdrawn from an individual clinically or subclinically infected with a virulent *Mycobacterium*, the induction being

10 performed by the addition of the polypeptide to a suspension comprising about 1.0 to 2.5  $\times 10^5$  PBMC, the addition of the polypeptide resulting in a concentration of not more than 20  $\mu$ g per ml suspension, the release of IFN- $\gamma$  being assessable by determination of IFN- $\gamma$  in supernatant harvested 5 days after the addition of the polypeptide to the suspension, and preferably does not induce such an IFN- $\gamma$  release in an individual not infected with a

15 virulent *Mycobacterium*,

v) it induces a positive *in vitro* response determined by release of IFN- $\gamma$  of at least 500pg/ml from T cell lines generated from PPD positive individuals, the induction being performed by the addition of the polypeptide to a suspension comprising 1-5  $\times 10^5$

20 cells/ml, the addition of the polypeptide resulting in a concentration not more than 20 $\mu$ g/ml, the release of IFN- $\gamma$  being assessable by determination of the IFN- $\gamma$  in supernatant harvested 3-5 days after the addition of the polypeptide to the suspension.

vi) it induces a positive *in vitro* response determined by T-cell proliferation of at least a

25 stimulation index (SI) of 5 (SI, calculated as mean counts per minute in the presence of antigen divided by the mean counts per minute without antigen) from T cell lines generated from PPD positive individuals, the induction being performed by the addition of the polypeptide to a suspension comprising 1-5  $\times 10^5$  cells/ml, the addition of the polypeptide resulting in a concentration not more than 20 $\mu$ g/ml, the release of IFN- $\gamma$

30 being assessable by determination of the IFN- $\gamma$  in supernatant harvested 3-5 days after the addition of the polypeptide to the suspension.

vii) it induces a positive DTH response determined by intradermal injection or local application patch of at most 100  $\mu$ g of the polypeptide to an individual who is clinically or

subclinically infected with a virulent *Mycobacterium*, a positive response having a diameter of at least 10 mm 72-96 hours after the injection or application,

viii) it induces a positive DTH response determined by intradermal injection or local

5 application patch of at most 100 µg of the polypeptide to an individual who is clinically or subclinically infected with a virulent *Mycobacterium*, a positive response having a diameter of at least 5 mm 72-96 hours after the injection, and preferably does not induce a such response in an individual who has a cleared infection with a virulent *Mycobacterium*.

10

The property described in i) will also be satisfied if the release of IFN- $\gamma$  from reactivated memory T-lymphocytes is 2,000 pg/ml, such as 3,000 pg/ml. In an alternative embodiment of the invention, the immunological effect of the polypeptide could be determined by comparing the IFN- $\gamma$  release as described with the IFN- $\gamma$  release from a

15 similar assay, wherein the polypeptide is not added, a significant increase being indicative of an immunologically effective polypeptide. In a preferred embodiment of the invention, the addition of the polypeptide results in a concentration of not more than 20 µg per ml suspension, such as 15 µg, 10 µg, 5 µg, 3 µg, 2 µg, or 1 µg polypeptide per ml suspension.

20

One example of a mouse strain for property i) is the C57Bl/6j as the animal model. As will be known by a person skilled in the art, due to genetic variation, different strains may react with immune responses of varying strength to the same polypeptide. It is presently unknown which strains of mice will give the best predictability of immunogenic reactivity

25 in which human population. Therefore, it is important to test other mouse strains, such as C3H/HeN, CBA (preferably CBA/J), DBA (preferably DBA/2J), A/J, AKR/N, DBA/1J, FVB/N, SJL/N, 129/SvJ, C3H/HeJ-Lps or BALB mice (preferably BALB/cA, BALB/cJ). It is presently contemplated that also a similar test performed in another animal model such as a guinea pig model or a rat model will have clinical predictability. In order to obtain  
30 good clinical predictability to humans, it is contemplated that a model in any farm animal, such as a cow model, a pig model, a deer model, or any primate model will have clinical predictability and thus serve as an animal model.

It should be noted, moreover, that the tuberculosis disease also affects a number of

35 different animal species such as cows, primates, guinea pigs, badgers, possums, and

deers. A polypeptide which has proven effective in any of the models mentioned above may be of interest for animal treatment even if it is not effective in a human being.

It is proposed to measure the release of IFN- $\gamma$  from reactivated T lymphocytes withdrawn from a mouse within 28 days after the mouse has been infected with virulent *Mycobacteria*. This is due to the fact that when an immune host mounts a protective immune response, the specific T-cells responsible for the early recognition of the infected macrophage stimulate a powerful bactericidal activity through their production of IFN- $\gamma$  (Rook, G.A.W. 1990., Flesch, I. et al. 1987). However, other cytokines could be relevant when monitoring the immunological response to the polypeptide, such as IL-12, TNF- $\alpha$ , IL-4, IL-5, IL-10, IL-6, TGF- $\beta$ . Usually one or more cytokines will be measured utilising for example the PCR technique or ELISA. It will be appreciated by a person skilled in the art that a significant increase or decrease in the amount of any of these cytokines induced by a specific polypeptide can be used in evaluation of the immunological efficacy of the polypeptide. The ability of a polypeptide to induce a IFN- $\gamma$  response is presently believed to be the most relevant correlate of protective immunity as mice with a disruption of the gene coding for IFN- $\gamma$  are unable to control a mycobacterial infection and die very rapidly with widespread dissemination, caseous necrosis and large abcesses (Flynn et al (1993) J.Exp.Med 178: 2249-2254, Cooper et al (1993) J.Exp.Med. 178:2243-2248). A specific model for obtaining information regarding the antigenic targets of a protective immunity in the memory model was originally developed by Lefford (Lefford et al (1973) Immunology 25:703) and has been used extensively in the recent years (Orme et al (1988). Infect.Immun. 140:3589, P.Andersen et al. (1995) J.Immunol.154:3359).

The property described in ii) will also be satisfied if the release of IFN- $\gamma$  from PBMC is determined in PBMC withdrawn from TB patients or PPD positive individuals more than 6 months after diagnosis such as 9 months, 1 year, 2 years, 5 years, or 10 years after diagnosis.

The comments on property i) regarding significant increase in IFN- $\gamma$ , concentration of polypeptide, and other cytokines are equally relevant to property ii).

The property described in iii) will in particular be satisfied, if the ELISA is performed as follows: the polypeptide of interest in the concentration of 1 to 10  $\mu$ g/ml is coated on a 96 wells polystyrene plate (NUNC, Denmark) and after a washing step with phosphate buffer

pH 7.3, containing 0.37 M NaCl and 0.5% Tween-20 the serum or plasma from a TB patient is applied in dilutions from 1:10 to 1:1000 in PBS with 1% Tween-20. Binding of an antibody to the polypeptide is determined by addition of a labeled (e.g. peroxidase labeled) secondary antibody and reaction is thereafter visualized by the use of OPD and 5 H<sub>2</sub>O<sub>2</sub> as described by the manufacturer (DAKO, Denmark). The OD value in each well is determined using an appropriate ELISA reader.

In a preferred embodiment, the western blot is performed as follows: The polypeptide is applied in concentrations from 1-40 µg to a SDS-PAGE and after electrophoresis the 10 polypeptide is transferred to a membrane e.g. nitrocellulose or PVDF. The membrane is thereafter washed in phosphate buffer, pH 7.3, containing 0.37 M NaCl and 0.5% Tween-20 for 30 min. The sera obtained from one or more TB patients are diluted 1:10 to 1:1000 in phosphate buffer pH 7.3 containing 0.37 M NaCl. The membrane is hereafter washed four times five minutes in binding buffer and incubated with peroxidase- or phosphatase- 15 labeled secondary antibody. Reaction is then visualized using the staining method recommended by the manufacturer (DAKO, Denmark).

The property described in iv) will in particular be satisfied if the polypeptide does not induce such an IFN- $\gamma$  release in an individual not infected with a virulent *Mycobacterium*, 20 i.e. an individual who has been BCG vaccinated or infected with *Mycobacterium avium* or sensitised by non-tuberculosis *Mycobacterium* or in an individual cleared of an infection with a virulent *Mycobacterium*, i.e. which does not have any positive culture, microscopical or clinical proven ongoing infection with virulent *Mycobacterium*. The comments on property i) regarding significant increase in IFN- $\gamma$ , concentration of 25 polypeptide, and other cytokines are equally relevant to property iv).

The property described in vii) will in particular be satisfied if the polypeptide does not induce such a response in an individual not infected with a virulent *Mycobacterium*, i.e. an individual who has been BCG vaccinated or infected with *Mycobacterium avium* or 30 sensitised by non-tuberculosis *Mycobacterium*. In a preferred embodiment, the amount of polypeptide intradermally injected or applied is 90 µg, such as 80 µg, 70 µg, 60 µg, 50 µg, 40 µg, or 30 µg. In another embodiment of the invention, the diameter of the positive response is at least 6 mm, such as 7 mm, 8 mm, 9 mm, or 10 mm. In a preferred embodiment, the induration or erythema or both could be determined after administration

of the polypeptide by intradermal injection, patch test or multipuncture. The reaction diameter could be positive after more than 48, such as 72 or 96 hours.

The property described in viii) will in particular be satisfied if the polypeptide does not 5 induce such a response in an individual cleared of an infection with a virulent *Mycobacterium*, i.e. which does not have any positive culture or microscopically proven ongoing infection with virulent *Mycobacterium*. The comments on property vii) regarding the amount of polypeptide intradermally injected or applied and the diameter of the positive response are equally relevant to property viii).

10

One aspect of the present invention relates to a substantially pure polypeptide fragment which comprises an amino acid sequence encoded by a member of the esat-6 gene family having a sequence identity with said polypeptide fragment of at least 70% and at the same time being immunologically equivalent to said polypeptide fragment with the 15 proviso that the substantially pure polypeptide is not selected from the group consisting of Rv0287, Rv0288, Rv1037c, Rv1038c, Rv1197, Rv1198, Rv1792, Rv1793, Rv2347c, Rv2346c, Rv3019c, Rv3619c, Rv3620c, Rv3874, and Rv3875.

In the present context, two polypeptide fragments are immunologically equivalent if they 20 both satisfy property i), property ii), property iii), property iv), property v), property vi), property vii), or property viii).

**Table 1** Tentative list of members of the *esat-6* gene family. Proteins coded by adjacent genes are listed together between the lines (e.g. Rv1036c, Rv1037c and Rv1038c). The percentage of identity to TB10.4, CFP10 and ESAT-6, respectively, was calculated according to the LALIGN algorithm used for definition of criteria c) above.

5 SEQ ID NOs are presented in Table 2 (nucleotide sequences) and Table 3 (protein sequences).

Protein	Size (number of amino acids)	Other names	Percentage of amino acid sequence identity to CFP10 (SEQ ID NO: 1), ESAT-6 (SEQ ID NO: 2), or TB10.4 (SEQ ID NO: 3)
Rv0287	97		18.6 % (TB10.4), 31.0 % (CFP10), 16.5 % (ESAT-6)
Rv0288	96	TB10.4 (previously named CFP7)	18.0 % (CFP10), 21.9 % (ESAT-6)
Rv1036c	100		15.0 % (TB10.4), 31.0 % (CFP10), 15.0 % (ESAT-6)
Rv1037c	94		18.8 % (TB10.4), 14.0 % (CFP10), 22.1 % (ESAT-6)
Rv1038c	98		21.4 % (TB10.4), 18.6 % (CFP10), 9.2 % (ESAT-6)
Rv1197	98		22.4 % (TB10.4), 20.6 % (CFP10), 9.2 % (ESAT-6)
Rv1198	94		18.8 % (TB10.4), 13.0 % (CFP10), 21.1 % (ESAT-6)
Rv1792	98		20.4 % (TB10.4), 19.6 % (CFP10), 11.2 % (ESAT-6)
Rv1793	94		18.0 % (TB10.4), 12.0 % (CFP10), 21.2 % (ESAT-6)
Rv2346c	94		19.8 % (TB10.4), 13.0 % (CFP10), 20 % (ESAT-6)
Rv2347c	98		21.4 % (TB10.4), 18.6 % (CFP10), 10.2 % (ESAT-6)
Rv2348c	108		14.8 % (TB10.4), 13.0 % (CFP10), 13 % (ESAT-6)
Rv2653c	107		18.3 % (TB10.4), 16.5 % (CFP10), 16.7 % (ESAT-6)
Rv2654c	81		21.0 % (TB10.4), 16.0 % (CFP10), 20.0 % (ESAT-6)
Rv3019c	96		84.4 % (TB10.4), 17.0 % (CFP10), 24.0 % (ESAT-6)
Rv3020c	97		17.5 % (TB10.4), 31 % (CFP10), 15.5 % (ESAT-6)
Rv3444c	100		20 % (TB10.4), 15.2 % (CFP10), 22.0 % (ESAT-6)
Rv3445c	125		15.2 % (TB10.4), 12.8 % (CFP10), 15.1 % (ESAT-6)
Rv3619c	94		18.8 % (TB10.4), 14.0 % (CFP10), 22.1 % (ESAT-6)
Rv3620c	98		21.4 % (TB10.4), 19.6 % (CFP10), 10.2 % (ESAT-6)
Rv3874	100	CFP10	18.0 % (TB10.4), 15.0 % (ESAT-6)
Rv3875	95	ESAT-6	21.9 % (TB10.4), 15.0 % (CFP10)
Rv3890c	95		25.8 % (TB10.4), 18.6 % (CFP10), 15.6 % (ESAT-6)
Rv3891c	107		23.4 % (TB10.4), 16.2 % (CFP10), 16.8 % (ESAT-6)
Rv3904c	90		23.2 % (TB10.4), 19.8 % (CFP10), 18.9 % (ESAT-6)
Rv3905c	103		22.3 % (TB10.4), 21.4 % (CFP10), 18.4 % (ESAT-6)

In both immunodiagnostics and vaccine preparation, it is often possible and practical to prepare antigens from segments of a known immunogenic protein or polypeptide. Certain epitopic regions may be used to produce responses similar to those produced by the entire antigenic polypeptide.

5

In order to identify relevant T-cell epitopes which are recognized during an immune response, it is also possible to use a "brute force" method: Since T-cell epitopes are linear, deletion mutants of polypeptides having SEQ ID NOs: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31 will, if constructed systematically, reveal what regions of the 10 polypeptides are essential in immune recognition, e.g. by subjecting these deletion mutants to the IFN- $\gamma$  assay described herein. Another method utilises overlapping oligopeptides (preferably synthetic having a length of e.g. 20 amino acid residues) derived from polypeptides having SEQ ID NOs: 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31. Some of these will give a positive response in the IFN- $\gamma$  assay whereas others 15 will not.

In a preferred embodiment of the invention, the polypeptide fragment of the invention comprises an epitope for a B-cell or T-cell.

20 Although the minimum length of a T-cell epitope has been shown to be at least 6 amino acids, it is normal that such epitopes are constituted of longer stretches of amino acids. Hence, it is preferred that the polypeptide fragment of the invention has a length of at least 7 amino acid residues, such as at least 8, at least 9, at least 10, at least 12, at least 14, at least 16, at least 18, at least 20, at least 22, at least 24, and at least 30 amino acid 25 residues.

In one preferred embodiment, the polypeptide fragment of the invention is free from any signal sequence; this is especially interesting when the polypeptide fragment is produced synthetically but even when the polypeptide fragments are produced recombinantly it is 30 normally acceptable that they are not exported by the host cell to the periplasm or the extracellular space; the polypeptide fragments can be recovered by traditional methods (cf. the discussion below) from the cytoplasm after disruption of the host cells, and if there is need for refolding of the polypeptide fragments, general refolding schemes can be employed, cf. e.g. the disclosure in WO 94/18227 where such a general applicable 35 refolding method is described.

By producing fusion polypeptides, superior characteristics of the polypeptide fragments of the invention can be achieved. For instance, fusion partners which facilitate export of the polypeptide when produced recombinantly, fusion partners which facilitate purification of

- 5 the polypeptide, and fusion partners which enhance the immunogenicity of the polypeptide fragment of the invention are all interesting possibilities. Therefore, the invention also pertains to a fusion polypeptide comprising at least one polypeptide fragment defined above and at least one fusion partner. The fusion partner can, in order to enhance immunogenicity, e.g. be selected from the group consisting of another
- 10 polypeptide fragment as defined above (so as to allow for multiple expression of relevant epitopes), and another polypeptide derived from a bacterium belonging to the tuberculosis complex, such as ESAT-6, TB10.4, CFP10, CFP17, CFP21, CFP25, CFP29, MPB59, MPT59, MPB64, and MPT64 or at least one T-cell epitope of any of these antigens. Other immunogenicity enhancing polypeptides which could serve as fusion partners are
- 15 T-cell epitopes (e.g. derived from the polypeptides ESAT-6, MPB64, MPT64, or MPB59) or other immunogenic epitopes enhancing the immunogenicity of the target gene product, e.g. lymphokines such as IFN- $\gamma$ , IL-2 and IL-12. In order to facilitate expression and/or purification, the fusion partner can e.g. be a bacterial fimbrial protein, e.g. the pilus components pilin and papA; protein A; the ZZ-peptide (ZZ-fusions are marketed by
- 20 Pharmacia in Sweden); the maltose binding protein; glutathione S-transferase;  $\beta$ -galactosidase; or poly-histidine.

Other interesting fusion partners are polypeptides which are lipidated causing that the immunogenic polypeptide is presented in a suitable manner to the immune system. This

- 25 effect is e.g. known from vaccines based on the *Borrelia burgdorferi* OspA polypeptide, wherein the lipidated membrane anchor in the polypeptide confers a self-adjuvating effect to the polypeptide (which is natively lipidated) when isolated from cells producing it. In contrast, the OspA polypeptide is relatively silent immunologically when prepared without the lipidation anchor.

30

Another part of the invention pertains to a nucleic acid fragment in isolated form which

- 1) comprises a nucleic acid sequence which is a member of the esat-6 gene family, and/or

35

2) has a length of at least 10 nucleotides and hybridizes readily under stringent hybridization conditions (as defined in the art, i.e. 5-10°C under the melting point  $T_m$ , cf. Sambrook et al, 1989, pages 11.45-11.49) with a nucleic acid fragment of 1) and/or

5 3) has a length of at least 10 nucleotides and hybridizes readily under stringent hybridization conditions (as defined in the art, i.e. 5-10°C under the melting point  $T_m$ , cf. Sambrook et al, 1989, pages 11.45-11.49) with a nucleic acid fragment which has a nucleotide sequence selected from

10 SEQ SEQ ID NO: 6 or a sequence complementary thereto,  
SEQ ID NO: 12 or a sequence complementary thereto,  
SEQ ID NO: 14 or a sequence complementary thereto,  
SEQ ID NO: 16 or a sequence complementary thereto,  
SEQ ID NO: 18 or a sequence complementary thereto,

15 SEQ ID NO: 20 or a sequence complementary thereto,  
SEQ ID NO: 22 or a sequence complementary thereto,  
SEQ ID NO: 24 or a sequence complementary thereto,  
SEQ ID NO: 26 or a sequence complementary thereto,  
SEQ ID NO: 28 or a sequence complementary thereto, or

20 SEQ ID NO: 30 or a sequence complementary thereto.

It is preferred that the nucleic acid fragment is a DNA fragment.

To provide certainty of the advantages in accordance with the invention, the preferred 25 nucleic acid sequence when employed for hybridization studies or assays includes sequences that are complementary to at least a 10 to 40, or so, nucleotide stretch of the selected sequence. A size of at least 10 nucleotides in length helps to ensure that the fragment will be of sufficient length to form a duplex molecule that is both stable and selective. Molecules having complementary sequences over stretches greater than 10 30 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained.

Hence, the term "subsequence" when used in connection with the nucleic acid fragments 35 of the invention is intended to indicate a continuous stretch of at least 10 nucleotides

which exhibits the above hybridization pattern. Normally, this will require a minimum sequence identity of at least 70% with a subsequence of the hybridization partner having SEQ ID NO: 6, 12, 14, 16, 18, 20, 22, 24, 26, 28 or 30. It is preferred that the nucleic acid fragment is longer than 10 nucleotides, such as at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 70, and at least 80 nucleotides long, and the sequence identity should preferable also be higher than 70%, such as at least 75%, at least 80%, at least 85%, at least 90%, at least 92%, at least 94%, at least 96%, and at least 98%. It is most preferred that the sequence identity is 100%. Such fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, by application of nucleic acid reproduction technology, such as the PCR technology of U.S. Patent 4,603,102, or by introducing selected sequences into recombinant vectors for recombinant production.

It is well known that the same amino acid may be encoded by various codons, the codon usage being related, *inter alia*, to the preference of the organisms in question expressing the nucleotide sequence. Thus, at least one nucleotide or codon of a nucleic acid fragment of the invention may be exchanged by others which, when expressed, result in a polypeptide identical or substantially identical to the polypeptide encoded by the nucleic acid fragment in question. The invention thus allows for variations in the sequence such as substitution, insertion (including introns), addition, deletion and rearrangement of one or more nucleotides, which variations do not have any substantial effect on the polypeptide encoded by the nucleic acid fragment or a subsequence thereof. The term "substitution" is intended to mean the replacement of one or more nucleotides in the full nucleotide sequence with one or more different nucleotides, "addition" is understood to mean the addition of one or more nucleotides at either end of the full nucleotide sequence, "insertion" is intended to mean the introduction of one or more nucleotides within the full nucleotide sequence, "deletion" is intended to indicate that one or more nucleotides have been deleted from the full nucleotide sequence whether at either end of the sequence or at any suitable point within it, and "rearrangement" is intended to mean that two or more nucleotide residues have been exchanged with each other.

The nucleotide sequence to be modified may be of cDNA or genomic origin as discussed above, but may also be of synthetic origin. Furthermore, the sequence may be of mixed cDNA and genomic, mixed cDNA and synthetic or genomic and synthetic origin as discussed above. The sequence may have been modified, e.g. by site-directed mu-

tagenesis, to result in the desired nucleic acid fragment encoding the desired polypeptide. The following discussion focused on modifications of nucleic acid encoding the polypeptide should be understood to encompass also such possibilities, as well as the possibility of building up the nucleic acid by ligation of two or more DNA fragments to

5 obtain the desired nucleic acid fragment, and combinations of the above-mentioned principles.

The nucleotide sequence may be modified using any suitable technique which results in the production of a nucleic acid fragment encoding a polypeptide of the invention.

10

The modification of the nucleotide sequence encoding the amino acid sequence of the polypeptide of the invention should be one which does not impair the immunological function of the resulting polypeptide.

15 A preferred method of preparing variants of the antigens disclosed herein is site-directed mutagenesis. This technique is useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, derived from the antigen sequences, through specific mutagenesis of the underlying nucleic acid. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating

20 one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the nucleic acid. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the nucleotide sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to

25 form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector which includes within its sequence a nucleic acid sequence which encodes the polypeptides of the invention. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically, e.g. by the method of Crea et al. (1978). This primer is then annealed with the single-stranded vector and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment,

35 in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is

formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

5

The preparation of sequence variants of the selected nucleic acid fragments of the invention using site-directed mutagenesis is provided as a means of producing potentially useful species of the genes and is not meant to be limiting, as there are other ways in which sequence variants of the nucleic acid fragments of the invention may be obtained.

10 For example, recombinant vectors encoding the desired genes may be treated with mutagenic agents to obtain sequence variants (see, e.g., a method described by Eichenlaub, 1979) for the mutagenesis of plasmid DNA using hydroxylamine.

15 The invention also relates to a replicable expression vector which comprises a nucleic acid fragment defined above, especially a vector which comprises a nucleic acid fragment encoding a polypeptide fragment of the invention.

20 The vector may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, *i.e.* a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication; examples of such a vector are a plasmid, phage, cosmid, mini-chromosome or virus. Alternatively, the vector may be one which, when introduced in a host cell, is integrated in the host cell genome and replicated together with the 25 chromosome(s) into which it has been integrated.

30 Expression vectors may be constructed to include any of the DNA segments disclosed herein. Such DNA might encode an antigenic protein specific for virulent strains of mycobacteria or even hybridization probes for detecting mycobacteria nucleic acids in samples. Longer or shorter DNA segments could be used, depending on the antigenic protein desired. Epitopic regions of the proteins expressed or encoded by the disclosed DNA could be included as relatively short segments of DNA. A wide variety of expression vectors is possible including, for example, DNA segments encoding reporter gene products useful for identification of heterologous gene products and/or resistance genes 35 such as antibiotic resistance genes which may be useful in identifying transformed cells.

The vector of the invention may be used to transform cells so as to allow propagation of the nucleic acid fragments of the invention or so as to allow expression of the polypeptide fragments of the invention. Hence, the invention also pertains to a transformed cell

5 harbouring at least one such vector according to the invention. Such a transformed cell (which is also a part of the invention) may be any suitable bacterial host cell or any other type of cell such as a unicellular eukaryotic organism, a fungus or yeast, or a cell derived from a multicellular organism, e.g. an animal or a plant. It is especially in cases where glycosylation is desired that a mammalian cell is used, although glycosylation of proteins  
10 is a rare event in prokaryotes. Normally, however, a prokaryotic cell is preferred such as a bacterium belonging to the genera *Mycobacterium*, *Salmonella*, *Pseudomonas*, *Bacillus* and *Escherichia*. It is preferred that the transformed cell is an *E. coli*, *B. subtilis*, or *M. bovis* BCG cell, and it is especially preferred that the transformed cell expresses a  
15 polypeptide according to the invention. The latter opens for the possibility to produce the polypeptide of the invention by simply recovering it from the culture containing the transformed cell. In the most preferred embodiment of this part of the invention the transformed cell is *Mycobacterium bovis* BCG strain: Danish 1331, which is the *Mycobacterium bovis* strain Copenhagen from the Copenhagen BCG Laboratory, Statens Serum Institut, Denmark.

20

The nucleic acid fragments of the invention allow for the recombinant production of the polypeptides fragments of the invention. However, also isolation from the natural source is a way of providing the polypeptide fragments as is peptide synthesis.

25 Therefore, the invention also pertains to a method for the preparation of a polypeptide fragment of the invention, said method comprising inserting a nucleic acid fragment as defined above into a vector which is able to replicate in a host cell, introducing the resulting recombinant vector into the host cell (transformed cells may be selected using various techniques, including screening by differential hybridization, identification of fused  
30 reporter gene products, resistance markers, anti-antigen antibodies and the like), culturing the host cell in a culture medium under conditions sufficient to effect expression of the polypeptide (of course the cell may be cultivated under conditions appropriate to the circumstances, and if DNA is desired, replication conditions are used), and recovering the polypeptide from the host cell or culture medium; or

35

isolating the polypeptide from whole mycobacteria of the tuberculosis complex or from lysates or fractions thereof, e.g. cell wall containing fractions, or

synthesizing the polypeptide by solid or liquid phase peptide synthesis.

5

The medium used to grow the transformed cells may be any conventional medium suitable for the purpose. A suitable vector may be any of the vectors described above, and an appropriate host cell may be any of the cell types listed above. The methods employed to construct the vector and effect introduction thereof into the host cell may be

10 any method known for such purposes within the field of recombinant DNA. In the following, a more detailed description of the possibilities will be given:

In general, of course, prokaryotes are preferred for the initial cloning of nucleic sequences of the invention and constructing the vectors useful in the invention. For ex-

15 ample, in addition to the particular strains mentioned in the more specific disclosure below, one may mention, by way of example, strains such as *E. coli* K12 strain 294 (ATCC No. 31446), *E. coli* B, and *E. coli* X 1776 (ATCC No. 31537). These examples are, of course, intended to be illustrative rather than limiting.

20 Prokaryotes are also preferred for expression. The aforementioned strains, as well as *E. coli* W3110 (F-, lambda-, prototrophic, ATCC No. 273325), bacilli such as *Bacillus subtilis*, or other enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcescens*, and various *Pseudomonas* species may be used. Especially interesting are rapid-growing mycobacteria, e.g. *M. smegmatis*, as these bacteria have a high degree of  
25 resemblance with mycobacteria of the tuberculosis complex and therefore stand a good chance of reducing the need of performing post-translational modifications of the expression product. In one aspect of the invention it is preferred to produce the polypeptide of the invention in a GRAS organism e.g. *Lactococcus*.

30 In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is typically transformed using pBR322, a plasmid derived from an *E. coli* species (see, e.g.,  
35 Bolivar et al., 1977, Gene 2: 95). The pBR322 plasmid contains genes for ampicillin and

tetracycline resistance and thus provides easy means for identifying transformed cells.

The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the microorganism for expression.

- 5 Those promoters most commonly used in recombinant DNA construction include the B-lactamase (penicillinase) and lactose promoter systems (Chang et al., 1978; Itakura et al., 1977; Goeddel et al., 1979) and a tryptophan (trp) promoter system (Goeddel et al., 1979; EPO Appl. Publ. No. 0036776). While these are the most commonly used promoter, other microbial promoters have been discovered and utilized, and details concerning their nucleotide sequences have been published, enabling a skilled worker to ligate them functionally with plasmid vectors (Siebenlist et al., 1980). Certain genes from prokaryotes may be expressed efficiently in *E. coli* from their own promoter sequences, precluding the need for addition of another promoter by artificial means.
- 10
- 15 After the recombinant preparation of the polypeptide according to the invention, the isolation of the polypeptide may for instance be carried out by affinity chromatography (or other conventional biochemical procedures based on chromatography), using a monoclonal antibody which substantially specifically binds the polypeptide according to the invention. Another possibility is to employ the simultaneous electroelution technique
- 20 described by Andersen et al. in J. Immunol. Methods 161: 29-39.

According to the invention, the post-translational modifications may involve lipidation, glycosylation, cleavage, or elongation of the polypeptide.

- 25 In certain aspects, the DNA sequence information provided by this invention allows for the preparation of relatively short DNA (or RNA, PNA, or LNA) sequences having the ability to specifically hybridize to mycobacterial gene sequences. In these aspects, nucleic acid probes of an appropriate length are prepared based on a consideration of the relevant sequence. The ability of such nucleic acid probes to specifically hybridize to
- 30 the mycobacterial gene sequences lends them particular utility in a variety of embodiments. Most importantly, the probes can be used in a variety of diagnostic assays for detecting the presence of pathogenic organisms in a given sample. However, either use is envisioned, including the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructs.

Apart from their use as starting points for the synthesis of polypeptides of the invention and for hybridization probes (useful for direct hybridization assays or as primers in e.g. PCR or other molecular amplification methods), the nucleic acid fragments of the invention may be used for effecting *in vivo* expression of antigens, *i.e.* the nucleic acid

5 fragments may be used in so-called DNA vaccines. Recent research have revealed that a DNA fragment cloned in a vector which is non-replicative in eukaryotic cells may be introduced into an animal (including a human being) by e.g. intramuscular injection or percutaneous administration (the so-called "gene gun" approach). The DNA is taken up by e.g. muscle cells and the gene of interest is expressed by a promoter which is functioning in eukaryotes, *e.g.* a viral promoter, and the gene product thereafter stimulates 10 the immune system. These newly discovered methods are reviewed in Ulmer et al., 1993, which hereby is included by reference.

Hence, the invention also relates to a vaccine comprising a nucleic acid fragment according to the invention, the vaccine effecting *in vivo* expression of antigen by an animal, including a human being, to whom the vaccine has been administered, the amount of expressed antigen being effective to confer substantially increased resistance to infections with mycobacteria of the tuberculosis complex in an animal, including a human being.

20 The efficacy of such a "DNA vaccine" can possibly be enhanced by administering the gene encoding the expression product together with a DNA fragment encoding a polypeptide which has the capability of modulating an immune response. For instance, a gene encoding lymphokine precursors or lymphokines (*e.g.* IFN- $\gamma$ , IL-2, or IL-12) could be 25 administered together with the gene encoding the immunogenic protein, either by administering two separate DNA fragments or by administering both DNA fragments included in the same vector. It also is a possibility to administer DNA fragments comprising a multitude of nucleotide sequences which each encode relevant epitopes of the polypeptides disclosed herein so as to effect a continuous sensitization of the immune 30 system with a broad spectrum of these epitopes.

As explained above, the polypeptide fragments of the invention are excellent candidates for vaccine constituents or for constituents in an immune diagnostic agent.

Thus, another part of the invention pertains to an immunologic composition comprising a polypeptide or fusion polypeptide according to the invention. In order to ensure optimum performance of such an immunologic composition it is preferred that it comprises an immunologically and pharmaceutically acceptable carrier, vehicle or adjuvant.

5

Suitable carriers are selected from the group consisting of a polymer to which the polypeptide(s) is/are bound by hydrophobic non-covalent interaction, such as a plastic, e.g. polystyrene, or a polymer to which the polypeptide(s) is/are covalently bound, such as a polysaccharide, or a polypeptide, e.g. bovine serum albumin, ovalbumin or keyhole limpet haemocyanin. Suitable vehicles are selected from the group consisting of a diluent and a suspending agent. The adjuvant is preferably selected from the group consisting of dimethyldioctadecylammonium bromide (DDA), Quil A, poly I:C, Freund's incomplete adjuvant, IFN- $\gamma$ , IL-2, IL-12, monophosphoryl lipid A (MPL), and muramyl dipeptide (MDP).

15

A preferred immunologic composition according to the present invention comprises at least two different polypeptide fragments, each different polypeptide fragment being a polypeptide or a fusion polypeptide defined above. It is preferred that the immunologic composition comprises between 2-20, such as 3-20 different polypeptide fragments or 20 fusion polypeptides.

Such an immunologic composition may preferably be in the form of a vaccine or in the form of a skin test reagent.

25 In line with the above, the invention therefore also pertain to a method for producing an immunologic composition according to the invention, the method comprising preparing, synthesizing or isolating a polypeptide according to the invention, and solubilizing or dispersing the polypeptide in a medium for a vaccine, and optionally adding other *M. tuberculosis* antigens and/or a carrier, vehicle and/or adjuvant substance.

30

Preparation of vaccines which contain peptide sequences as active ingredients is generally well understood in the art, as exemplified by U.S. Patents 4,608,251; 4,601,903; 4,599,231 and 4,599,230, all incorporated herein by reference. Typically, such vaccines are prepared as injectables either as liquid solutions or suspensions; solid forms suitable

35 for solution in, or suspension in, liquid prior to injection may also be prepared. The

preparation may also be emulsified. The active immunogenic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like, and combinations thereof. In addition, if desired, the vaccine may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, or adjuvants which enhance the effectiveness of the vaccines.

The vaccines are conventionally administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories and, in some cases, oral formulations. For suppositories, traditional binders and carriers may include, for example, polyalkalene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10-95% of active ingredient, preferably 25-70%.

The proteins may be formulated into the vaccine as neutral or salt forms. Pharmaceutically acceptable salts include acid addition salts (formed with the free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups may also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

The vaccines are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immunogenic. The quantity to be administered depends on the subject to be treated, including, e.g., the capacity of the individual's immune system to mount an immune response, and the degree of protection desired. Suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination with a preferred range from about 0.1  $\mu$ g to 1000  $\mu$ g, such as in the range from about 1  $\mu$ g to 300  $\mu$ g, and especially in the range from about 10  $\mu$ g to

50 µg. Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations.

- 5 The manner of application may be varied widely. Any of the conventional methods for administration of a vaccine are applicable. These are believed to include oral application on a solid physiologically acceptable base or in a physiologically acceptable dispersion, parenterally, by injection or the like. The dosage of the vaccine will depend on the route of administration and will vary according to the age of the person to be vaccinated and, to
- 10 a lesser degree, the size of the person to be vaccinated.

Some of the polypeptides of the vaccine are sufficiently immunogenic in a vaccine, but for some of the others the immune response will be enhanced if the vaccine further comprises an adjuvant substance.

15

Various methods of achieving adjuvant effect for the vaccine include use of agents such as aluminum hydroxide or phosphate (alum), commonly used as 0.05 to 0.1 percent solution in phosphate buffered saline, admixture with synthetic polymers of sugars (Carbopol) used as 0.25 percent solution, aggregation of the protein in the vaccine by

- 20 heat treatment with temperatures ranging between 70°C to 101°C for 30 second to 2 minute periods respectively. Aggregation by reactivating with pepsin treated (Fab) antibodies to albumin, mixture with bacterial cells such as *C. parvum* or endotoxins or lipopolysaccharide components of gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with 20

- 25 percent solution of a perfluorocarbon (Fluosol-DA) used as a block substitute may also be employed. According to the invention, DDA (dimethyldioctadecylammonium bromide) is an interesting candidate for an adjuvant, but also Freund's incomplete adjuvants as well as QuilA are interesting possibilities. Further possibilities are monophosphoryl lipid A (MPL), and muramyl dipeptide (MDP).

30

Another highly interesting (and thus, preferred) possibility of achieving adjuvant effect is to employ the technique described in Gosselin *et al.*, 1992 (which is hereby incorporated by reference herein). In brief, the presentation of a relevant antigen such as an antigen of the present invention can be enhanced by conjugating the antigen to antibodies (or

- 35 antigen binding antibody fragments) against the Fc<sub>y</sub> receptors on mono-

cytes/macrophages. Especially conjugates between antigen and anti-Fc<sub>γ</sub>RI have been demonstrated to enhance immunogenicity for the purposes of vaccination.

Other possibilities involve the use of immune modulating substances such as lymphokines (e.g. IFN- $\gamma$ , IL-2 and IL-12) or synthetic IFN- $\gamma$  inducers such as poly I:C in combination with the above-mentioned adjuvants. As discussed in example 3b, it is contemplated that such mixtures of antigen and adjuvant will lead to superior vaccine formulations.

10 In many instances, it will be necessary to have multiple administrations of the vaccine, usually not exceeding six vaccinations, more usually not exceeding four vaccinations and preferably one or more, usually at least about three vaccinations. The vaccinations will normally be at from two to twelve week intervals, more usually from three to five week intervals. Periodic boosters at intervals of 1-5 years, usually three years, will be desirable  
15 to maintain the desired levels of protective immunity. The course of the immunisation may be followed by *in vitro* proliferation assays of PBMC co-cultured with one or more of the polypeptides members used in the vaccine, e.g. co-culture with ESAT-6 or ST-CF, and especially by measuring the levels of IFN- $\gamma$  released from the primed lymphocytes. The assays may be performed using conventional labels, such as radionuclides,  
20 enzymes, fluorescers, and the like. These techniques are well known and may be found in a wide variety of patents, such as U.S. Patent Nos. 3,791,932; 4,174,384 and 3,949,064, as illustrative of these types of assays.

Due to genetic variation, different individuals may react with immune responses of  
25 varying strength to the same polypeptide. Therefore, the vaccine according to the invention may comprise several different polypeptides in order to increase the immune response. The vaccine may comprise two or more polypeptides, where all of the polypeptides are as defined above, or some but not all of the peptides may be derived from a bacterium belonging to the *M. tuberculosis* complex. In the latter example, the  
30 polypeptides not necessarily fulfilling the criteria set forth above for polypeptides may either act due to their own immunogenicity or merely act as adjuvants. Examples of such interesting polypeptides are ESAT-6, TB10.4, and MPT64, but any other substance which can be isolated from mycobacteria are possible candidates.

The vaccine may comprise 1-20, such as 2-20 or even 3-20 different polypeptides, such as 3-10 different polypeptides.

One reason for admixing the polypeptides of the invention with an adjuvant is to effectively activate a cellular immune response. However, this effect can also be achieved in other ways, for instance by expressing the effective antigen in a vaccine in a non-pathogenic microorganism. A well-known example of such a microorganism is *Mycobacterium bovis* BCG.

10 Therefore, another important aspect of the present invention is an improvement of the living BCG vaccine presently available, which is a vaccine for immunizing an animal, including a human being, against TB caused by mycobacteria belonging to the tuberculosis-complex, comprising as the effective component a microorganism, wherein one or more copies of a DNA sequence encoding a polypeptide as defined above has been incorporated into the genome of the microorganism in a manner allowing the microorganism to express and secrete the polypeptide.

In the present context, the term "genome" refers to the chromosome of the microorganisms as well as extrachromosomally DNA or RNA, such as plasmids. It is, however, preferred that the DNA sequence of the present invention has been introduced into the chromosome of the non-pathogenic microorganism, since this will prevent loss of the genetic material introduced.

It is preferred that the non-pathogenic microorganism is a bacterium, e.g. selected from the group consisting of the genera *Mycobacterium*, *Salmonella*, *Pseudomonas* and *Escherichia*. It is especially preferred that the non-pathogenic microorganism is *Mycobacterium bovis* BCG, such as *Mycobacterium bovis* BCG strain: Danish 1331.

The incorporation of one or more copies of a nucleotide sequence encoding the polypeptide according to the invention in a *mycobacterium* from a *M. bovis* BCG strain will enhance the immunogenic effect of the BCG strain. The incorporation of more than one copy of a nucleotide sequence of the invention is contemplated to enhance the immune response even more, and consequently an aspect of the invention is a vaccine wherein at least 2 copies of a DNA sequence encoding a polypeptide is incorporated in the genome of the microorganism, such as at least 5 copies. The copies of DNA sequences may

either be identical encoding identical polypeptides or be variants of the same DNA sequence encoding identical or homologues of a polypeptide, or in another embodiment be different DNA sequences encoding different polypeptides where at least one of the polypeptides is according to the present invention.

5

The living vaccine of the invention can be prepared by cultivating a transformed non-pathogenic cell according to the invention, and transferring these cells to a medium for a vaccine, and optionally adding a carrier, vehicle and/or adjuvant substance.

- 10 The invention also relates to a method of diagnosing TB caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis* in an animal, including a human being, comprising intradermally injecting, in the animal, a polypeptide according to the invention or a skin test reagent described above, a positive skin response at the location of injection being indicative of the animal having TB, and a negative skin
- 15 response at the location of injection being indicative of the animal not having TB. A positive response is a skin reaction having a diameter of at least 5 mm, but larger reactions are preferred, such as at least 1 cm, 1.5 cm, and at least 2 cm in diameter. The composition used as the skin test reagent can be prepared in the same manner as described for the vaccines above.

20

- 25 In line with the disclosure above pertaining to vaccine preparation and use, the invention also pertains to a method for immunising an animal, including a human being, against TB caused by mycobacteria belonging to the tuberculosis complex, comprising administering to the animal the polypeptide of the invention, or a vaccine composition of the invention as described above, or a living vaccine described above. Preferred routes of administration are the parenteral (such as intravenous and intraarterially), intraperitoneal, intramuscular, subcutaneous, intradermal, oral, buccal, sublingual, nasal, rectal or transdermal route.

- 30 30 A number of possible diagnostic assays and methods can be envisaged:

- When diagnosis of previous or ongoing infection with virulent mycobacteria is the aim, a blood sample comprising mononuclear cells (i.e. T-lymphocytes) from a patient could be contacted with a sample of one or more polypeptides of the invention. This contacting
- 35 can be performed *in vitro* and a positive reaction could e.g. be proliferation of the T-cells

or release of cytokines such as  $\gamma$ -interferon into the extracellular phase (e.g. into a culture supernatant); a suitable *in vivo* test would be a skin test as described above. It is also conceivable to contact a serum sample from a subject to contact with a polypeptide of the invention, the demonstration of a binding between antibodies in the serum sample and

5 the polypeptide being indicative of previous or ongoing infection.

The invention therefore also relates to an *in vitro* method for diagnosing ongoing or previous sensitization in an animal or a human being with bacteria belonging to the tuberculosis complex, the method comprising providing a blood sample from the animal

10 or human being, and contacting the sample from the animal with the polypeptide of the invention, a significant release into the extracellular phase of at least one cytokine by mononuclear cells in the blood sample being indicative of the animal being sensitised. By the term "significant release" is herein meant that the release of the cytokine being significantly larger (with a 95% confidence interval as defined by appropriate statistical 15 analysis such as a Student's two-tailed T test) than the cytokine release from a blood sample derived from a patient without the TB diagnosis. Normally, a significant release is at least two times the release observed from such a sample.

Alternatively, a sample of a possibly infected organ may be contacted with an antibody

20 raised against a polypeptide of the invention. The demonstration of the reaction by means of methods well-known in the art between the sample and the antibody will be indicative of an ongoing infection. It is of course also a possibility to demonstrate the presence of anti-mycobacterial antibodies in serum by contacting a serum sample from a subject with at least one of the polypeptide fragments of the invention and using well- 25 known methods for visualizing the reaction between the antibody and antigen.

Also a method of determining the presence of mycobacterial nucleic acids in an animal, including a human being, or in a sample, comprising administering a nucleic acid fragment of the invention to the animal or incubating the sample with the nucleic acid

30 fragment of the invention or a nucleic acid fragment complementary thereto, and detecting the presence of hybridized nucleic acids resulting from the incubation (by using the hybridization assays which are well-known in the art), is also included in the invention. Such a method of diagnosing TB might involve the use of a composition comprising at least a part of a nucleotide sequence as defined above and detecting the presence of 35 nucleotide sequences in a sample from the animal or human being to be tested which

hybridize with the nucleic acid fragment (or a complementary fragment) by the use of PCR technique.

The fact that certain of the disclosed antigens are not present in *M. bovis* BCG but are 5 present in virulent mycobacteria point them out as interesting drug targets; the antigens may constitute receptor molecules or toxins which facilitate the infection by the *mycobacterium*, and if such functionalities are blocked the infectivity of the *mycobacterium* will be diminished.

10 To determine particularly suitable drug targets among the antigens of the invention, the gene encoding at least one of the polypeptides of the invention and the necessary control sequences can be introduced into avirulent strains of mycobacteria (e.g. BCG) so as to determine which of the polypeptides are critical for virulence. Once particular proteins are identified as critical for/contributory to virulence, anti-mycobacterial agents can be 15 designed rationally to inhibit expression of the critical genes or to attack the critical gene products. For instance, antibodies or fragments thereof (such as Fab and (Fab')<sub>2</sub>) fragments can be prepared against such critical polypeptides by methods known in the art and thereafter used as prophylactic or therapeutic agents. Alternatively, small molecules can be screened for their ability to selectively inhibit expression of the critical 20 gene products, e.g. using recombinant expression systems which include the gene's endogenous promoter, or for their ability to directly interfere with the action of the target. These small molecules are then used as therapeutics or as prophylactic agents to inhibit mycobacterial virulence.

25 Alternatively, anti-mycobacterial agents which render a virulent *mycobacterium* avirulent can be operably linked to expression control sequences and used to transform a virulent *mycobacterium*. Such anti-mycobacterial agents inhibit the replication of a specified *mycobacterium* upon transcription or translation of the agent in the *mycobacterium*. Such a "newly avirulent" *mycobacterium* would constitute a superb alternative to the above 30 described modified BCG for vaccine purposes since it would be immunologically very similar to a virulent *mycobacterium* compared to e.g. BCG.

Finally, a monoclonal or polyclonal antibody, which is specifically reacting with a polypeptide of the invention in an immuno assay, or a specific binding fragment of said antibody, is also a part of the invention. The production of such polyclonal antibodies 35

requires that a suitable animal be immunized with the polypeptide and that these antibodies are subsequently isolated, suitably by immune affinity chromatography. The production of monoclonals can be effected by methods well-known in the art, since the present invention provides for adequate amounts of antigen for both immunization and screening of positive hybridomas.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed in Australia before the priority date of each claim of this application.

## REFERENCES

Andersen, P., and I. Heron. 1993. *Journal of Immunological Methods* **161** 29-39

10 Andersen, P., A.B. Andersen, A.L. Sorensen, and S. Nagai. 1995. *J.Immunol.* **154**:3359-3372.

Berthet, F.X., Rasmussen, P.B., Rosenkrands, I., Andersen, P. Gicquel, B. *Microbiology* (1998) **144** 3195-3203

S. T. Cole, et al (1998). *Nature* **393** (6685) 537-544.

Ravn et al (1999). *J. Infect. Dis.* **179**, 637-645

15 WO98/53075

WO98/53076

Rook, G.A.W. (1990). *Res. Microbiol.* **141**:253-256

Flesch, I. et al. (1987). *J. Immunology*, **138**: 4408-4413

Flynn et al (1993) *J.Exp.Med* **178**: 2249-2254,

20 Cooper et al (1993) *J.Exp.Med.* **178**:2243-2248

Orme et al (1988). *Infect.Immun.* **40**:3589,

Ulmer et al., (1993). *Curr. Opin. Invest. Drugs*, **2**: 983-989

Gosselin et al., (1992) *J. Immunol.* **149**: 3477-3481.

Pearson W.R and D.J. Lipman (1988) *PNAS USA* **85**:2444-2448

25 X. Huang and W. Miller. *Adv. Appl. Math.* (1991) **12**:337-357

Nielsen P E et al. (1991. *Science* **254**: 1497-1500

Koshkin et al. (1998). *Tetrahedron* **54**, 3607-3630;

Nielsen, N.K. et al. (1998). *J.Am.Chem.Soc.* **120**, 5458-5463

Lefford et al (1973). *Immunology* **25**:703

30 Crea et al. (1978). *PNAS* **75**(12):5765-5769.

Eichenlaub, (1979). *Journal of Bacteriology*, **138**: 559-566.

Bolivar et al., (1977). *Gene* **2**: 95

Chang et al., (1978). *Nature*, **375**: 515

Itakura et al., (1977). *Science*, **198**: 1056

Goeddel et al., (1979). *Nature*, **281**:544

EPO Appl. Publ. No. 0036776

5 Siebwenlist et al., (1980). *Cell*, **20**: 269

Harboe et al (1998). *Infect. Immun.* **66**, 717-723

Sambrook et. al. (1989)"Molecular cloning: a laboratory manual, 2nd edition Cold Spring Harbor N. Y."

WO 98/44119

10 WO 94/18227

U. S. patent 4,603,102

U. S. patent 4,608,251

U. S. patent 4,601,903

U. S. patent 4,599,231

15 U. S. patent 4,599,230

U. S. patent 3,791,932

U. S. patent 4,174,384

U. S. patent 3,949,064

## FIGURE LEGENDS

### Figure 1.

Human lymphocyte responses to rTB7.3, rTB10.4 and rCFP10. The IFN- $\gamma$  response resulting from stimulation of PBMC's from two human TB patients (circles) and two healthy BCG vaccinated human donors (triangles) with increasing concentrations of rTB7.3 (A), rTB10.4 (B) and rCFP10 (C). All IFN- $\gamma$  analyses were done in duplicates on supernatants pooled from three wells, and have been given as means. The variation on the duplicate wells was always less than 10% of the mean. IFN- $\gamma$  levels below 50 pg/ml were considered negative.

10

100

### Figure 2.

IFN-gamma responses to low mass antigens from *M. tuberculosis* in different groups of donors. 7 healthy non-vaccinated donors, 7 healthy BCG vaccinated donors and 17 TB patients were stimulated with 5  $\mu$ g/ml of ST-CF or recombinant antigens. Individual antigen specific responses are shown as delta values (IFN-gamma release in the antigen stimulated well minus IFN-gamma release in the unstimulated well). ST-CF: Short-term culture filtrate, rTB7.3: Recombinant form of Rv3221c, rTB10.4: Recombinant form of Rv0288, rCFP10: Recombinant form of CFP10, rESAT-6: Recombinant form of ESAT-6.

Figure 3. IFN- $\gamma$  response induced by stimulation of PBMC's from human TB patients with peptide pools of Rv2654c.

Figure 4. IFN- $\gamma$  response induced by stimulation of PBMC's from healthy BCG vaccinated donors with peptide pools of Rv2654c.

## EXAMPLES

### ***Example 1 The effect of CFP10, ESAT-6, and TB10.4 on stimulation of PBMC's from human TB patients.***

The ESAT-6 antigen was identified in the low molecular mass fraction of culture filtrate 5 due to a strong T cell response with high levels of IFN- $\gamma$  released (Andersen et al 1995). This antigen has now in a number of studies been demonstrated to have good stimulatory antigenic properties and is recognized strongly by a high percentage of TB patients as well as different animal species infected with TB. Recently, a few other small 10 proteins have been identified from various mycobacterial extracts and evaluated for their immunological relevance. Recently, a 10 kDa molecule (CFP10) encoded in the same operon as ESAT-6 was identified (Berthet, F.X. 1998).

Two novel low mass *M. tuberculosis* proteins have been identified: TB10.4, and TB7.3 (identical to Rv3221c and not a member of the ESAT-6 gene family). TB10.4 was identified as a novel member of the ESAT-6 family and our data demonstrate that the 15 three members of the ESAT-6 family tested so far (TB10.4, CFP10 and ESAT-6), all are strongly recognized targets by the human immune response against *M. tuberculosis*.

#### *Cloning of the genes encoding CFP10, TB7.3 and TB10.4.*

The gene encoding CFP10 was cloned as described before (Berthet, F.X. 1998). TB7.3 20 (previously named CFP7A) was identified from ST-CF and the corresponding gene was cloned as described (WO98/44119).

The gene encoding TB10.4 (previously named CFP7) was identified by screening a  $\lambda$ gt11 *M. tuberculosis* genome library with the Mab PV-2 and cloned as described previously (WO98/44119).

25

#### *Expression and purification of recombinant TB7.3, TB10.4 and CFP10.*

The histidine-tagged recombinant proteins (rTB7.3, rTB10.4 and rCFP10) were expressed and purified by metal affinity chromatography using a Talon column (Clonetech, Palo Alto, Ca) in the presence of 8M urea, essentially as described by the 30 manufacturer. Purification of the proteins to homogeneity was done by anion exchange chromatography using 1ml Hitrap columns (Pharmacia, Uppsala, Sweden).

Protein concentrations were determined by the BCA - test (Micro BCA Protein Assay Reagent kit, Pierce, Oud-Beijerland, The Netherlands). LPS content in these

preparations, measured by the Limulus Amoebocyte Lysate (LAL) -test, was always below 0.05ng LPS/µg protein.

*Immunological recognition of low mass M. tuberculosis proteins*

5 PBMC were obtained from 17 Danish TB patients diagnosed and treated at the Department of Pulmonary Medicine, University Hospital of Copenhagen, Denmark and from 7 BCG vaccinated and 7 non-vaccinated healthy individuals with no known exposure to *M. tuberculosis*. Blood samples were drawn between 0 and 6 months after diagnosis of tuberculosis, and 2 months to 40 years after BCG vaccination.

10 Separation, culture of PBMC and measurement of IFN-γ in the supernatants was done as described previously by Ravn et al. A dose response study of the three recombinant proteins (rTB7.3, rTB10.4 and rCFP10) was carried out using 0.3 to 10µg antigen /ml culture. Lymphocyte cultures from two Danish TB patients and two healthy Danish BCG vaccinated donors were stimulated with the three antigens. The lymphocyte response

15 after stimulation with TB7.3 was low with IFN-γ releases generally below 1000pg/ml (Figure 1A). Neither IFN-γ nor proliferative responses to this antigen (data not shown), reached more than 20% of the responses seen with ST-CF. For the two other antigens high levels of IFN-γ were induced with increasing antigen concentrations (Figure 1 B and C). Optimal concentrations of the antigens were between 1.25 to 10µg/ml and these

20 concentrations gave responses in the range of 1000-4000pg IFN-γ/ml.

The antigens were investigated in 13-17 TB patients, 4-7 BCG vaccinated and 7 non-vaccinated donors (Figure 2). TB7.3 was recognized but at a low level in both patients and BCG vaccinated donors. Around 40 % (5 out off 13) of the TB patients recognized this molecule at a level significantly above background and for these donors the median

25 response was 659pg IFN-γ/ml versus 4024pg IFN-γ/ml in the same donors for ST-CF. TB10.4 was recognized at a much higher level, by both BCG vaccinated donors (71% responders, median IFN-γ = 3968pg/ml versus 5335pg/ml in the same donors for ST-CF), and TB patients (88% responders, median IFN-γ = 3298pg/ml versus 4707pg/ml in the same donors for ST-CF). In the TB patients, CFP10 induced a pronounced release of

30 IFN-γ (median IFN-γ= 2135pg/ml versus 4755pg/ml in the same donors for ST-CF).

Compared with the pronounced T cell responses to TB10.4, CFP10 and ESAT-6, TB7.3 was a weakly recognized antigen with a very low activity.

Compared to ESAT-6, TB10.4 induced significantly higher levels of IFN-γ in TB patients (P = 0.0017, Wilcoxon Signed Rank Test), whereas T cell responses to CFP10 and

ESAT-6 were similar ( $P = 0.121$ ). Both CFP10 and TB10.4 were recognized by >70% of the TB patients, and interestingly these two potent immunogenic molecules have several points in common with ESAT-6: They have almost identical size and pI (10kDa and 4.5) and show 15 % and 21.9 %, respectively, amino acid sequence identity to ESAT-6, and

5 are members of the *esat-6* gene family as previously defined.

The data presented indicate a striking focusing of the host immune response towards members of the ESAT-6 family, demonstrating that this family contains a number of molecules of potential relevance for future TB vaccines and diagnostics.

10 **Example 2 Cloning of the genes encoding low mass proteins from the ESAT-6 family.**

The genes encoding Rv0287, Rv1036c, Rv1037c, Rv2346c, Rv2348c, Rv2653c, Rv2654c, Rv3020c, Rv3444c, Rv3445c, Rv3890c, Rv3891c, Rv3904c and Rv3905c were cloned into the expression vector pMCT3 (identical to pMCT6, Harboe et al, 1998, 15 except that it only contains six N-terminal histidine residues), by PCR amplification with gene specific primers, for recombinant expression in *E. coli* of the proteins.

For cloning of the proteins, the following gene specific primers were used:

Rv0287:

20 PA0287: 5'- CTGAGATCTATGAGCCTTTGGATGC- 3' (*Bgl*II)  
PB0287: 5'- CTAAGCTTGGATCCTCAGAACCCGGTATAGG - 3' (*Bam*HI)

Rv1036c:

PA1036c: 5'- CTGAGATCTTGATCCCCGGTCGGATGGTG (*Bgl*II).  
PB1036c: 5'- CTCCCATGGGTCAAGGTGATCGAATCAGCCA (*Ncol*)

25 Rv1037c:

PA1037c: 5'- CTGAGATCTATGACCATCAACTATC - 3' (*Bgl*II)  
PB1037c: 5'- CTAAGCTTGGATCCTTAGGCCAGCTGGAGCC - 3' (*Bam*HI)

Rv2346c:

PA2346c: 5'- CTGAGATCTATGACCATCAACTATC - 3' (*Bgl*II)

30 PB2346c: 5'- CTAAGCTTGGATCCTCAGGCCAGCTGGAGCC - 3' (*Bam*HI)

Rv2348c:

PA2348c: 5'- CTGAGATCTGTGCTTTGCCTCTGGTCCG (*Bgl*II)

PB2348c: 5'- CCCAAGCTTCTAGCCGGCCGCCGGAGA (*Hind*III).

Rv2653c:

PA2653c: 5'- CTGAGATCTTGACCCACAAGCGCACTAAA (*Bg*/*II*).

PB2653c: 5'- CTCCCATGGTCACTGTTCGCTGTCGGGTTTC (*Ncol*).

Rv2654c:

PA2654c: 5'- CTGAGATCTATGAGCGGCCACGCGTTGGCT (*Bg*/*II*).

5 PB2654c: 5'- CTCCCATGGTCACGGCGGATCACCCGGTC (*Ncol*).

Rv3020c:

PA3020c: 5'- CTGAGATCTATGAGTTGGATGCCAT (*Bg*/*II*).

PB3020c: 5'- CTCCCATGGTAAAACCCGGTAGCTGGA (*Ncol*).

Rv3444c:

10 PA3444c: 5'- CTGAGATCTATGAACGCAGACCCGTG - 3' (*Bg*/*II*)

PB3444c: 5'- CTAAGCTTGGATCCCTAGCGTGCCAAGCTCC - 3' (*Bam**HI*)

Rv3445c:

PA3445c: 5'- CTGAGATCTATGGTGAACCGGGAAGG - 3' (*Bg*/*II*)

PB3445c: 5'- CTAAGCTTGGATCCCTAGGTCGCCGCCGGC - 3' (*Bam**HI*)

15 Rv3890c:

PA3890c: 5'- CTGAGATCTATGTCAGATCAAATCACG - 3' (*Bg*/*II*)

PB3890c: 5'- CTAAGCTTGGATCCTAGAACACAAGCCCGCG - 3' (*Bam**HI*)

Rv3891c:

PA3891c: 5'- CTGAGATCTATGGCAGACACAATTCAAGG - 3' (*Bg*/*II*)

20 PB3891c: 5'- CTAAGCTTCCGGGTCAAGATCCGTGGCTAGC - 3' (*Sma**I*)

Rv3904c:

PA3904c: 5'- CTGAGATCTATGGATCCGACCGTGTGG - 3' (*Bg*/*II*)

PB3904c: 5'- CTGCCATGGTCACGACCACATACCC - 3' (*Ncol*)

Rv3905c:

25 PA3905c: 5'- CTGAGATCTATGGTGCCGACGACAC - 3' (*Bg*/*II*)

PB3905c: 5'- CTAAGCTTGGATCCTCAGCCACCGCCCACC - 3' (*Bam**HI*)

The primers listed above create the restriction sites indicated after each sequence. The restriction sites are used for the cloning in pMCT3. Where an alternative start codon to

30 ATG is used in the original sequence the primers introduce an ATG codon instead.

PCR reactions contained 10 ng of *M. tuberculosis* chromosomal DNA in 1 x PCR buffer + Mg (Boehringer Manheim) with 400 $\mu$ M dNTP mix (Boehringer Mannheim), 0.4 pM of each primer and 1.5 unit Tag DNA polymerase (Boehringer Mannheim) in 50  $\mu$ l reaction volume. Reactions were initially heated to 94°C for 5 min., run for 30 cycles of the

35 program; 92°C for 1 min., 52°C for 1min. and 72°C for 2min. and terminating with 72°C

for 7min., using PTC-200 thermal cycler (M J Research, Inc.). The PCR products were cloned into the pRC2.1 cloning vector and transformed into One Shot<sup>TM</sup> *E. coli* cells (Invitrogen, Leek, The Netherlands) as described by the manufacturer. Plasmid DNA was digested with the appropriate restriction enzymes (see primer sequence) and cloned into 5 pMCT3 and transformed into *E. coli* XL-1 Blue cells. The correct insert was always confirmed by sequencing. Sequencing of DNA was performed at Statens Serum Institut using the cycle sequencing system in combination with an automated gel reader (model 373A; Applied Biosystems).

10 *Expression and purification of recombinant Rv0287, Rv1036c, Rv1037c, Rv2346c, Rv2348c, Rv2653c, Rv2654c, Rv3020c, Rv3444c, Rv3445c, Rv3890c, Rv3891c, Rv3904c and Rv3905c.*

Expression and metal affinity purification of recombinant protein was undertaken essentially as described by the manufacturers. LB-media containing 100 µg/ml ampicillin 15 and 12.5µg/ml tetracyclin, was inoculated with overnight culture of XL1-Blue cells harbouring recombinant pMCT3 plasmid. The culture was shaken at 37 °C until it reached a density of OD<sub>600</sub> = 0.5. IPTG was hereafter added to a final concentration of 1 mM and the culture was further incubated 2-16 hours. Cells were harvested, resuspended in 1 x sonication buffer + 8 M urea and sonicated 5 X 30 sec. with 30 sec. 20 pausing between the pulses. After centrifugation, the lysate was applied to a column containing 10 ml Talon resin (Clontech, Palo Alto, USA). The column was washed and eluted as described by the manufacturers.

Fractions containing recombinant protein were pooled and to gain homogenous protein 25 preparations the pooled fractions were subjected to either the multielution technique (Andersen and Heron, 1993) or anion exchange on a Hitrap column (Pharmacia, Uppsala, Sweden).

**Table 2** List of nucleotide sequences with their name, Open Reading Frame (ORF) and SEQ ID NOs

Protein	ORF:	SEQ ID NO:
Rv0287	294	4
Rv1036c	339	6
Rv1037c	285	8
Rv2346c	282	10
Rv2348c	327	12
Rv2653c	324	14
Rv2654c	246	16
Rv3020c	294	18
Rv3444c	303	20
Rv3445c	378	22
Rv3890c	288	24
Rv3891c	324	26
Rv3904c	273	28
Rv3905c	312	30

**Table 3** List of proteins with their name, molecular mass (measured in Daltons), their Isoelectric point and their SEQ ID NO's.

Protein	Size (aa)	Molecular Mass (Da)	Isoelectric Point	SEQ ID NO:
CFP10 Rv3874	100	10794	4.41	1
ESAT-6 Rv3875	95	9904	4.3	2
TB10.4 Rv0288	96	10391	4.43	3
Rv0287	97	9778.40	6.3111	5
Rv1036c	112	12996.06	4.60	7
Rv1037c	94	9833.10	4.543	9
Rv2346c	94	9954.01	4.76	11
Rv2348c	108	11396.53	3.89	13
Rv2653c	107	12359.82	8.20	15
Rv2654c	81	7697.71	5.04	17
Rv3020c	97	9842.03	6.14	19
Rv3444c	100	11120.70	6.165	21
Rv3445c	125	13495.10	6.489	23
Rv3890c	95	9920.40	4.176	25
Rv3891c	107	11193.70	4.619	27
Rv3904c	90	9602.90	5.480	29
Rv3905c	103	10460.30	4.641	31

*Synthesis of synthetic peptides*

5 Three of the antigens (Rv3444c, Rv3890c and Rv3905c) were synthesised as synthetic peptides by standard solid-phase methods on an ABIMED peptide synthesiser (ABIMED, Langenfeld, Germany) at Dept. of infectious diseases and Immunohematology/Bloodbank C5-P, Leiden University Medical Centre, Albinusdreef 2, 2333 Leiden, The Netherlands.

10 The peptides covered the following amino acids;

Rv3444c p1: SEQ. ID. NO. 21: amino acid 1-18  
 Rv3444c p2: SEQ. ID. NO. 21: amino acid 11-28  
 Rv3444c p3: SEQ. ID. NO. 21: amino acid 21-38  
 15 Rv3444c p4: SEQ. ID. NO. 21: amino acid 31-48  
 Rv3444c p5: SEQ. ID. NO. 21: amino acid 41-58  
 Rv3444c p6: SEQ. ID. NO. 21: amino acid 51-68

Rv3444c p7: SEQ. ID. NO. 21: amino acid 61-78  
Rv3444c p8: SEQ. ID. NO. 21: amino acid 71-88  
Rv3444c p9: SEQ. ID. NO. 21: amino acid 81-100

5 Rv3890c p1: SEQ. ID. NO. 25: amino acid 1-18  
Rv3890c p2: SEQ. ID. NO. 25: amino acid 11-28  
Rv3890c p3: SEQ. ID. NO. 25: amino acid 21-38  
Rv3890c p4: SEQ. ID. NO. 25: amino acid 31-48  
Rv3890c p5: SEQ. ID. NO. 25: amino acid 41-58  
10 Rv3890c p6: SEQ. ID. NO. 25: amino acid 51-68  
Rv3890c p7: SEQ. ID. NO. 25: amino acid 61-78  
Rv3890c p8: SEQ. ID. NO. 25: amino acid 71-95

Rv3905c p1: SEQ. ID. NO. 31: amino acid 1-18  
15 Rv3905c p2: SEQ. ID. NO. 31: amino acid 11-28  
Rv3905c p3: SEQ. ID. NO. 31: amino acid 21-38  
Rv3905c p4: SEQ. ID. NO. 31: amino acid 31-48  
Rv3905c p5: SEQ. ID. NO. 31: amino acid 41-58  
Rv3905c p6: SEQ. ID. NO. 31: amino acid 51-68  
20 Rv3905c p7: SEQ. ID. NO. 31: amino acid 61-78  
Rv3905c p8: SEQ. ID. NO. 31: amino acid 71-88  
Rv3905c p9: SEQ. ID. NO. 31: amino acid 81-103

***EXAMPLE 3A: Interferon- $\gamma$  induction of T cell lines.***

25 The purified recombinant proteins were screened for the ability to induce a T cell response measured as IFN- $\gamma$  release. The screening involved testing of the IFN- $\gamma$  induction of T cell lines generated from PPD positive donors and / or a measurement of the response in PBMC preparations obtained from TB patients, PPD positive as well as negative healthy donors.

30

**Human donors:** PBMC were obtained from healthy donors with a positive *in vitro* response to PPD.

**T cell line preparation:** T cell lines were prepared by culturing  $1-5 \times 10^6$  freshly isolated PBMC with viable *M. tuberculosis* for 1½ hour at a ratio of 5 bacteria per cell in a total

volume of 1 ml (Donor 1 and 2). After washing, the cells were cultured in RPMI 1640 medium (Gibco, Grand Island, N.Y) supplemented with HEPES, and 10% heat-inactivated NHS. Alternatively, T cell lines were prepared by culturing 1-5 x 10<sup>6</sup> freshly isolated PBMC with 5 µg/ml of ST-CF (Donor 3-5). After 7 days in culture at 37 °C and 5% CO<sub>2</sub>, T cells were supplemented with 30-50 U/well of r-IL-2 (recombinant interleukin-2) (Boehringer Mannheim) for approximately 7 days. Finally, the T cell lines were tested for reactivity against the recombinant antigens and synthetic peptides by stimulating 1-5 x 10<sup>5</sup> cells/ml with 5 µg/ml of PPD and/or ST-CF, recombinant Rv2653c, Rv3891c, Rv3904c and peptide pools (2-9 peptides) of Rv3444c, Rv3890c and Rv3905c, in the presence of 5 x 10<sup>5</sup> autologous antigen-presenting cells/ml (donor 1 and 2) or 1 x 10<sup>8</sup> cells/ml of irradiated (2000 RAD) autologous PBMC (donor 3-5). No antigen (No ag) and PHA were used as negative and positive controls, respectively. The supernatants were harvested after 4 days of culture and stored at -20 °C until the presence of IFN-γ were analysed.

15 Responses obtained with different T cell lines are shown in Table 4, where donor 1 and 2 are based on T cell lines driven by viable *M. tuberculosis* whereas donor 3-5 are generated by stimulation with ST-CF.

20 **Table 4.** Stimulation of T cell lines with recombinant antigen and pools of synthetic peptides. Responses to PHA and PPD or ST-CF are shown for comparison. Results are presented as pg IFN-γ/ml.

Donor	No ag	PHA (1 µg/ml)	PPD (5 µg/ml)	Rv2653c (5µg/ml,1µg/ml)	Rv3444c p1-4 (5µg/ml,1µg/ml)	Rv3444c p5-9 (5µg/ml,1µg/ml)
1	350	3940	3690	1283, 853	132, 602	330, 553
2	325	3845	1824	673, 270	454, 558	1578, 1570

Donor	Rv3890c p1-4 (5µg/ml,1µg/ml)	Rv3890c p5-9 (5µg/ml,1µg/ml)	Rv3904c (5µg/ml,1µg/ml)	Rv3905c p1-4 (5µg/ml,1µg/ml)	Rv3905c p5-9 (5µg/ml,1µg/ml)
1	1167, 872	915, 1109	1827, 1146	1250, 622	332, 778
2	318, 362	522, 242	296, 664	503, 874	817, 422

Donor	No ag	PHA (1 µg/ml)	ST-CF (5 µg/ml)	Rv3891c (5µg/ml,0.5µg/ml)	Rv3444c p1+3 (5µg/ml,0.5µg/ml)	Rv3444c p2 (5µg/ml,0.5µg/ml)
3	136	4467	2425	1260, 606	223, 166	58, 154
4	0	5410	4490	23, 5	14, 12	12, 32
5	0	1996	1175	472, 479	254, 20	26, 33

Donor	Rv3444c p4 (5µg/ml,0.5µg/ml)	Rv3444c p5-6 (5µg/ml,0.5µg/ml)	Rv3444c p7-9 (5µg/ml,0.5µg/ml)	Rv3444c p1-9 (5µg/ml,0.5µg/ml)	Rv3905c p1-9 (5µg/ml,0.5µg/ml)
3	59, 93	700, 682	596, 298	308, 225	262, 116
4	33, 34	109, 69	240, 87	43, 17	452, 25
5	19, 16	119, 148	162, 29	319, 16	407, 26

The results shown in Table 4, regarding the recombinant antigens Rv2653c, Rv3891c and Rv3904c and the peptides covering the antigens Rv3444c, Rv3890c and Rv3905c, indicate that these antigens can induce IFN- $\gamma$  production in T-cell lines generated from 5 PPD positive individuals.

***Example 3B Interferon- $\gamma$  induction in human TB patients and BCG vaccinated***

**Human donors:** PBMC were obtained from healthy BCG vaccinated donors with no 10 known exposure to *M. tuberculosis* and from patients with culture or microscopy proven infection with TB. Blood samples were drawn from the TB patients 0-6 months after diagnosis.

**Lymphocyte preparations and cell culture:** PBMC were freshly isolated by gradient 15 centrifugation of heparinized blood on Lymphoprep (Nycomed, Oslo, Norway) and stored in liquid nitrogen until use. The cells were resuspended in complete RPMI 1640 medium (Gibco, Grand Island, N.Y.) supplemented with 1% penicillin/streptomycin (Gibco BRL, Life Technologies), 1% non-essential-amino acids (FLOW, ICN Biomedicals, CA, USA), and 10% normal human AB0 serum (NHS) from the local blood bank. The number and the 20 viability of the cells were determined by Nigrosin staining. Cultures were established with  $1.25 \times 10^5$  PBMCs in 50 µl in microtitre plates (Nunc, Roskilde, Denmark) and stimulated with ST-CF PDD, Rv0287, Rv1036c, Rv1037c, Rv2653c, Rv3445c, Rv3891c, and Rv3904c. No antigen (No ag) and phytohaemagglutinin (PHA) were used as negative and positive control, respectively. Supernatants for the detection of cytokines were 25 harvested after 5 days of culture, pooled, and stored at -80°C until used.

**Cytokine analysis:** Interferon- $\gamma$  (IFN- $\gamma$ ) was detected with a standard sandwich ELISA technique using a commercially available pair of monoclonal antibodies (Endogen) and used according to the manufacturer's instruction. Recombinant IFN- $\gamma$  (Endogen) was

used as a standard. All data are means of duplicate wells and the variation between wells did not exceed 10 % of the mean. Cytokine levels below 50 pg/ml were considered negative. Responses of 42 individual donors are shown in Table 5 and Table 6.

- 5 As shown in Table 5, marked release of IFN- $\gamma$  is observed after stimulation with several of the recombinant proteins. For 6 donors, stimulation with Rv0287 give rise to high IFN- $\gamma$  responses. Between 40% and 60% of the donors show intermediate IFN- $\gamma$  responses when stimulated with Rv1037c, Rv3891c and Rv3904c, whereas only limited responses are obtained by stimulation with Rv3445c in this experiment.

**Table 5.** Stimulation of PBMCs from 4 healthy non-BCG vaccinated, 4 healthy BCG vaccinated and 6 TB patients with recombinant antigen. Responses to ST-CF and PHA are shown for comparison. Results are given as pg IFN- $\gamma$ /ml.

BCG vaccinated control donors, no known TB exposure

Donor	No ag	PHA (1 $\mu$ g/ml)	ST-CF (5 $\mu$ g/ml)	Rv0287 (10 $\mu$ g/ml)	Rv1037c (10 $\mu$ g/ml)	Rv3445c (10 $\mu$ g/ml)	Rv3891c (10 $\mu$ g/ml)	Rv3904c (10 $\mu$ g/ml)
1	0	8305	622	1459	1800	5	2159	27
2	82	20862	15759	32	30	35	461	50
3	7	17785	16198	380	53	79	610	76
4	912	16198	11350	3020	3137	799	8137	716

5

TB patients

Donor	No ag	PHA (1 $\mu$ g/ml)	ST-CF (5 $\mu$ g/ml)	Rv0287 (10 $\mu$ g/ml)	Rv1037c (10 $\mu$ g/ml)	Rv3445c (10 $\mu$ g/ml)	Rv3891c (10 $\mu$ g/ml)	Rv3904c (10 $\mu$ g/ml)
1	60	12301	11057	2225	799	338	2115	94
2	7	10390	6123	51	44	20	0	522
3	34	11678	8136	1437	665	84	0	528
4	0	13459	7731	17	0	0	0	0
5	21	10143	9513	7869	3135	1646	4116	3018
6	0	10795	10932	8610	1409	421	9	1080

Non-vaccinated control donors, no known TB exposure

Donor	No ag	PHA (1 $\mu$ g/ml)	ST-CF (5 $\mu$ g/ml)	Rv0287 (10 $\mu$ g/ml)	Rv1037c (10 $\mu$ g/ml)	Rv3445c (10 $\mu$ g/ml)	Rv3891c (10 $\mu$ g/ml)	Rv3904c (10 $\mu$ g/ml)
1	61	8379	511	23	115	0	604	269
2	16	11005	1923	12	23	8	615	16
3	0	10190	126	0	0	0	249	0
4	51	10819	1030	0	0	0	n.d.	0

**Table 6.** Stimulation of PBMCs from 9 healthy PPD and/or ST-CF negative, 13 healthy PPD and/or ST-CF positive donors and 6 Tb patients with recombinant antigen. ST-CF, PPD and PHA are shown for comparison.

Results are given in pg IFN- $\gamma$ /ml.

5 Healthy PPD and/or ST-CF negative donors.

Donor	no ag	PHA	PPD	STCF (2.5 $\mu$ g/ml)	Rv1036c (10 $\mu$ g/ml)	Rv1036c (5 $\mu$ g/ml)	Rv1036c (2.5 $\mu$ g/ml)	Rv2653c (5 $\mu$ g/ml)	Rv2653c (2.5 $\mu$ g/ml)	Rv3891c (10 $\mu$ g/ml)	Rv3891c (2.5 $\mu$ g/ml)	Rv3904c (10 $\mu$ g/ml)	Rv3904c (2.5 $\mu$ g/ml)
A	0	3354	113	nd.	nd.	0	37	0	4	nd.	nd.	nd.	nd.
B	0	3803	563	nd.	nd.	15	14	0	50	nd.	nd.	nd.	nd.
C	0	3446	525	nd.	nd.	138	59	97	0	nd.	nd.	nd.	nd.
D	32	1919	nd.	234	137	nd.	148	nd	nd.	nd.	nd.	nd.	nd.
E	0	2889	nd.	178	59	nd.	206	nd.	nd.	nd.	nd.	nd.	nd.
F	42	3998	nd.	175	15	nd.	67	nd.	nd.	nd.	nd.	nd.	nd.
G	44	6269	190	195(5 $\mu$ g)	340	nd.	173	nd.	nd.	30	51	22	41
H	5	2282	n.d.	10 (5 $\mu$ g)	3	nd.	2	nd.	nd.	27	16	nd.	nd.
I	2	10427	n.d.	80 (5 $\mu$ g)	7	nd.	5	nd.	nd.	412	300	nd.	nd.

Table 6. - continued

Healthy PPD and/or ST-CF positive donors.

Donor	no ag	PHA	PPD	STCF (5µg/ml)	Rv1036c (10µg/ml)	Rv1036c (5µg/ml)	Rv1036c (2.5µg/ml)	Rv2653c (5µg/ml)	Rv2653c (2.5µg/ml)	Rv3891c (10µg/ml)	Rv3891c (2.5µg/ml)	Rv3904c (10µg/ml)	Rv3904c (2.5µg/ml)
A	31	6716	2275	nd.	nd.	687	900	1	62	nd.	nd.	nd.	nd.
B	43	4733	6159	nd.	nd.	2244	1108	179	126	nd.	nd.	nd.	nd.
C	7	6165	5808	nd.	nd.	4074	3788	110	30	nd.	nd.	nd.	nd.
D	63	6532	6314	nd.	nd.	1589	1450	2445	235	nd.	nd.	nd.	nd.
E	14	5614	3852	nd.	nd.	390	738	147	448	nd.	nd.	nd.	nd.
F	13	3493	4327	3381	229	nd.	605	nd.	nd.	8	25	42	61
G	12	8164	nd.	738	1774	nd.	2771	nd.	nd.	30	76	25	96
H	5	7378	840	nd.	568	nd.	948	nd.	nd.	15	19	730	102
I	0	5168	n.d.	4241	nd.	0	0	nd.	nd.	575	447	nd.	nd.
J	12	4873	nd.	745	nd.	4	3	nd.	nd.	511	214	nd.	nd.
K	1	4512	nd.	2137	nd.	5	1	nd.	nd.	1903	1105	nd.	nd.
L	75	8047	nd.	2778	812	nd.	235	nd.	nd.	nd.	nd.	nd.	nd.
M	52	6095	nd.	9133	1368	nd.	1223	nd.	nd.	nd.	nd.	nd.	nd.

**Table 6. - continued****Tb patients**

Donor	no ag	PHA	PPD	STCF (5µg/ml)	Rv1036c (10µg/ml)	Rv1036c (2.5µg/ml)	Rv3904c (10µg/ml)	Rv3904c (2.5µg/ml)
A	5	5282	4647	nd.	844 (5µg/ml)	557	nd.	nd.
B	60	7239	nd.	5474	301	595	nd.	nd.
C	44	11014	nd.	11639	384	646	nd.	nd.
D	8	5757	1095	877	624	692	31	88
E	25	7135	7118	5881	362	1035	55	54
F	23	6415	6085	6123	145	237	985	657

5 The results shown in Table 6 regarding the recombinant antigens Rv1036c, Rv2653c, Rv3891c and Rv3904c indicate that these antigens  
 50 can induce IFN- $\gamma$  production in PBMCs from healthy PPD and/or ST-CF positive individuals and /or Tb patients.

## Example 4

### Immunological evaluation of Rv2653c

To evaluate the immunological potential of Rv2653c we characterized the IFN- $\gamma$  response induced, when blood from PPD positive human donors was stimulated with overlapping peptides covering the complete amino acid sequence of Rv2653c.

The synthetic overlapping peptides covering the complete amino acid sequence of Rv2653c were purchased from Mimotopes Pty Ltd. The peptides were synthesized by Fmoc solid phase strategy. No purification steps were performed. Lyophilised peptides were stored dry until reconstitution in PBS.

The peptide sequences were as follows;

#### Rv2653c peptides

- Peptide-1: MTHRKRTKROPAlAAGLNA
- Peptide-2: AIAAGLNApRRNRVGROH
- Peptide-3: RNRVGROHGWPADVPSAE
- Peptide-4: PADVPSAEQRRAQRQRDL
- Peptide-5: RAQRQRDLEAIRRAYAEM
- Peptide-6: IRRAYAEMVATSHEIDDD
- Peptide-7: TSHEIDDDTAELALLSMH
- Peptide-8: ELALLSMHLDDEQRRLEA
- Peptide-9: DEQRRLLEAGHKLGWHPYH
- Peptide-10: MKLGWHPYHFPDEPDSKQ

The above listed synthetic peptides, covering the protein sequence of Rv2653c, were screened in pools of five peptides (Pool I: peptides 1-5, Pool II: peptides 6-10), for their ability to induce a T cell response measured as IFN- $\gamma$  release. The screening involved testing of the IFN- $\gamma$  induction in 24 PBMC preparations obtained from PPD positive human donors (6 TB patients and 18 healthy BCG vaccinated donors).

**Human donors:** PBMC were obtained from PPD positive human donors (6 TB patients with microscopy or culture proven infection and 18 healthy BCG vaccinated donors with no known exposure to TB).

**Lymphocyte preparations and cell culture:** PBMC were freshly isolated by gradient centrifugation of heparinized blood on Lymphoprep (Nycomed, Oslo, Norway) and stored in liquid nitrogen until use. The cells were resuspended in complete RPMI 1640 medium (Gibco BRL, Life Technologies) supplemented with 1% penicillin/streptomycin (Gibco BRL, Life Technologies), 1% non-essential amino acids (FLOW, ICN Biomedicals, CA, USA),

and 10% heat-inactivated normal human AB serum (NHS). The viability and number of the cells were determined by Nigrosin staining. Cell cultures were established with  $1.25 \times 10^5$  PBMCs in 100  $\mu$ l in microtitre plates (Nunc, Roskilde, Denmark) and stimulated with 5  $\mu$ g/ml PPD, or peptide pools I and II of Rv2653c in which the final concentrations of each peptide was 5  $\mu$ g/ml.

"No antigen" was included as negative control (not shown) and PPD was used as positive control. Supernatants for the analysis of secreted cytokines were harvested after 5 days of culture, pooled, and stored at -80 °C until use.

Donor/Antigen	No	PPD	Rv2653c pool I	Rv2653c pool II
PT1	21	15694	275	580
PT2	90	15519	946	3583
PT3	62	1570	30	634
PT4	243	30755	3819	5450
PT5	0	22120	51	734
PT6	0	4174	5	1304
BCG1	0	9601	2010	2014
BCG2	0	3508	1721	2283
BCG3	0	2639	92	490
BCG4	0	9781	142	3475
BCG5	6	21895	629	3404
BCG6	7	8663	6399	730
BCG7	11	6345	1120	2656
BCG8	11	22162	1693	1103
BCG9	0	6170	50	1233
BCG10	0	1387	0	419
BCG11	0	15701	10	1453
BCG12	0	9254	9	3785
BCG13	1	3557	84	1509
BCG14	0	5262	529	1295
BCG15	0	5732	183	1301
BCG16	1	4715	96	1985
BCG17	1	4145	65	2474
BCG18	0	11819	182	2734

As shown in Table 1 stimulation of PBMC's from TB patients (PT1-6) and BCG vaccinated healthy donors (BCG1-18) with peptide pools I and II of Rv2653c resulted in a marked release of IFN- $\gamma$  in both donor groups.

This result makes Rv2653c pool I and II, ideal candidates for vaccination against tuberculosis.

### Rv2654c as a diagnostic antigen.

To evaluate the diagnostic potential of Rv2654c we characterized the immunological response induced, when blood from TB patients and healthy BCG vaccinated donors was stimulated with overlapping peptides covering the complete amino acid sequence of Rv2654c.

The synthetic overlapping peptides covering the complete amino acid sequence of Rv2654c were purchased from Mimotopes Pty Ltd. The peptides were synthesized by Fmoc solid phase strategy. No purification steps were performed. Lyophilised peptides were stored dry until reconstitution in PBS.

.... The peptide sequences were as follows;

#### Rv2654c peptides

Peptide-1: MSGHALAARTLJAAADEL  
 Peptide-2: AADELVGGPPVEASAAAAL  
 Peptide-3: ASAAAALAGDAAGAWRTAA  
 Peptide-4: AWRTAAVELARALIVRAVA  
 Peptide-5: LVRAVAESHGVAAVLFAA  
 Peptide-6: VLFAATAAAAADVDRGDPP

The above listed synthetic peptides, covering the protein sequence of Rv2654c were screened in pools of three peptides (Pool I: peptides 1-3, Pool II: peptides 4-6), for their ability to induce a T cell response measured as IFN- $\gamma$  release. The screening involved testing of the IFN- $\gamma$  induction in PBMC preparations obtained from TB patients and healthy BCG vaccinated donors.

Human donors: PBMC were obtained from healthy BCG vaccinated donors with no known exposure to TB and from TB patients with microscopy or culture proven infection. Blood samples were drawn from TB patients 0-6 months after diagnosis.

**Lymphocyte preparations and cell culture:** PBMC were freshly isolated by gradient centrifugation of heparinized blood on Lymphoprep (Nycomed, Oslo, Norway) and stored in liquid nitrogen until use. The cells were resuspended in complete RPMI 1640 medium (Gibco BRL, Life Technologies) supplemented with 1% penicillin/streptomycin (Gibco BRL, Life Technologies), 1% non-essential-amino acids (FLOW, ICN Biomedicals, CA, USA), and 10% heat-inactivated normal human AB serum (NHS). The viability and number of the

cells were determined by Nigrosin staining. Cell cultures were established with  $1.25 \times 10^5$  PBMCs in 100  $\mu$ l in microtitre plates (Nunc, Roskilde, Denmark) and stimulated with 5  $\mu$ g/ml PPD, 5  $\mu$ g/ml recombinant ESAT-6 or peptide pools I and II of Rv2654c in which the final concentrations of each peptide was 5  $\mu$ g/ml.

"No antigen" was included as negative control (not shown) and PPD was used as positive control. More over the response to the highly responsive and TB specific ESAT-6 was included for comparison. Supernatants for the analysis of secreted cytokines were harvested after 5 days of culture, pooled, and stored at -80 °C until use.

As shown in Figure 1 stimulation of PBMC's from TB patients with peptide pools I and II of Rv2654c resulted in a marked release of IFN- $\gamma$ . In comparison BCG vaccinated donors (Fig. 2) did not respond to either of the pools.

To evaluate the sensitivity of Rv2654c as a diagnostic a cut off value of the "MEAN + 3 SD", based on the response to the BCG vaccinated donors, was chosen. This resulted in a cut off value for peptide pool I on 111 pg IFN- $\gamma$  /ml and for peptide pool II on 67 pg IFN- $\gamma$  /ml.

Using these cut off's 9/24 TB patients recognize pool I and 10/16 donors recognize pool II, with a specificity of 92% and 100% respectively, in the tested population.

This result makes Rv2654c pool I and II, ideal candidates for discriminating between TB infected and BCG vaccinated donors.

**EDITORIAL NOTE**

**APPLICATION NUMBER – 59664/00**

The following Sequence Listing pages **1** to **16** are part of the description. The claims pages follow on pages **51** to **60**.

## SEQUENCE LISTING

&lt;110&gt; Statens Serum Institut

<120> Tuberculosis vaccine and diagnostics  
based on the *Mycobacterium tuberculosis* esat-6 gene family

&lt;130&gt; 23388PC1

&lt;160&gt; 31

&lt;170&gt; FastSEQ for Windows Version 3.0

&lt;210&gt; 1

&lt;211&gt; 100

&lt;212&gt; PRT

<213> *M. Tuberculosis*

&lt;400&gt; 1

Met	Ala	Glu	Met	Lys	Thr	Asp	Ala	Ala	Thr	Leu	Ala	Gln	Glu	Ala	Gly	
1							5			10				15		
Asn	Phe	Glu	Arg	Ile	Ser	Gly	Asp	Leu	Lys	Thr	Gln	Ile	Asp	Gln	Val	
								20		25				30		
Glu	Ser	Thr	Ala	Gly	Ser	Leu	Gln	Gly	Gln	Trp	Arg	Gly	Ala	Ala	Gly	
							35		40		45					
Thr	Ala	Ala	Gln	Ala	Ala	Val	Val	Val	Arg	Phe	Gln	Glu	Ala	Ala	Asn	Lys
							50		55		60					
Gln	Lys	Gln	Glu	Leu	Asp	Glu	Ile	Ser	Thr	Asn	Ile	Arg	Gln	Ala	Gly	
	65						70			75			80			
Val	Gln	Tyr	Ser	Arg	Ala	Asp	Glu	Glu	Gln	Gln	Gln	Ala	Leu	Ser	Ser	
							85			90			95			
Gln	Met	Gly	Phe													
			100													

&lt;210&gt; 2

&lt;211&gt; 95

&lt;212&gt; PRT

<213> *M. Tuberculosis*

&lt;400&gt; 2

Met	Thr	Glu	Gln	Gln	Trp	Asn	Phe	Ala	Gly	Ile	Glu	Ala	Ala	Ala	Ser
1							5			10				15	
Ala	Ile	Gln	Gly	Asn	Val	Thr	Ser	Ile	His	Ser	Leu	Leu	Asp	Glu	Gly
							20		25		30				
Lys	Gln	Ser	Leu	Thr	Lys	Leu	Ala	Ala	Ala	Trp	Gly	Gly	Ser	Gly	Ser
							35		40		45				
Glu	Ala	Tyr	Gln	Gly	Val	Gln	Gln	Lys	Trp	Asp	Ala	Thr	Ala	Thr	Glu
							50		55		60				
Leu	Asn	Asn	Ala	Leu	Gln	Asn	Leu	Ala	Arg	Thr	Ile	Ser	Glu	Ala	Gly
	65						70			75			80		
Gln	Ala	Met	Ala	Ser	Thr	Glu	Gly	Asn	Val	Thr	Gly	Met	Phe	Ala	
							85			90			95		

&lt;210&gt; 3

&lt;211&gt; 96

&lt;212&gt; PRT

<213> *M. Tuberculosis*

<400> 3  
 Met Ser Gln Ile Met Tyr Asn Tyr Pro Ala Met Leu Gly His Ala Gly  
 1 5 10 15  
 Asp Met Ala Gly Tyr Ala Gly Thr Leu Gln Ser Leu Gly Ala Glu Ile  
 20 25 30  
 Ala Val Glu Gln Ala Ala Leu Gln Ser Ala Trp Gln Gly Asp Thr Gly  
 35 40 45  
 Ile Thr Tyr Gln Ala Trp Gln Ala Gln Trp Asn Gln Ala Met Glu Asp  
 50 55 60  
 Leu Val Arg Ala Tyr His Ala Met Ser Ser Thr His Glu Ala Asn Thr  
 65 70 75 80  
 Met Ala Met Met Ala Arg Asp Thr Ala Glu Ala Ala Lys Trp Gly Gly  
 85 90 95

<210> 4

<211> 294

<212> DNA

<213> M Tuberculosis

<220>

<221> CDS

<222> (1) ... (294)

<400> 4

atg agc ctt ttg gat gct cat atc cca cag ttg gtg gcc tcc cag tcg 48  
 Met Ser Leu Leu Asp Ala His Ile Pro Gln Leu Val Ala Ser Gln Ser  
 1 5 10 15

gcg ttt gcc gcc aag gcg ggg ctg atg cggt cac acg atc ggt cag gcc 96  
 Ala Phe Ala Ala Lys Ala Gly Leu Met Arg His Thr Ile Gly Gln Ala  
 20 25 30

gag cag gcg gcg atg tcg gct cag gcg ttt cac cag ggg gag tcg tcg 144  
 Glu Gln Ala Ala Met Ser Ala Gln Ala Phe His Gln Gly Glu Ser Ser  
 35 40 45

gcg gcg ttt cag gcc gcc cat gcc cgg ttt gtg gcg gcg gcc gcc aaa 192  
 Ala Ala Phe Gln Ala Ala His Ala Arg Phe Val Ala Ala Ala Lys  
 50 55 60

gtc aac acc ttg ttg gat gtc gcg cag gcg aat ctg ggt gag gcc gcc 240  
 Val Asn Thr Leu Leu Asp Val Ala Gln Ala Asn Leu Gly Glu Ala Ala  
 65 70 75 80

ggt acc tat gtg gcc gcc gat gct gcg gcc gcg tcg acc tat acc ggg 288  
 Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ser Thr Tyr Thr Gly  
 85 90 95

ttc tga 294  
 Phe \*

<210> 5

<211> 97

<212> PRT

<213> M Tuberculosis

<400> 5  
 Met Ser Leu Leu Asp Ala His Ile Pro Gln Leu Val Ala Ser Gln Ser  
 1 5 10 15  
 Ala Phe Ala Ala Lys Ala Gly Leu Met Arg His Thr Ile Gly Gln Ala  
 20 25 30  
 Glu Gln Ala Ala Met Ser Ala Gln Ala Phe His Gln Gly Glu Ser Ser  
 35 40 45  
 Ala Ala Phe Gln Ala Ala His Ala Arg Phe Val Ala Ala Ala Lys  
 50 55 60  
 Val Asn Thr Leu Leu Asp Val Ala Gln Ala Asn Leu Gly Glu Ala Ala  
 65 70 75 80  
 Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ser Thr Tyr Thr Gly  
 85 90 95  
 Phe

<210> 6  
 <211> 339  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(339)

<400> 6  
 ttg atc ccc ggt cggt atg gtg ctg aac tgg gaa gat ggc ctc aat gcc 48  
 Leu Ile Pro Gly Arg Met Val Leu Asn Trp Glu Asp Gly Leu Asn Ala  
 1 5 10 15  
 ctt gtt gcg gaa ggg att gag gcc atc gtg ttt cgt act tta ggc gat 96  
 Leu Val Ala Glu Gly Ile Glu Ala Ile Val Phe Arg Thr Leu Gly Asp  
 20 25 30  
 cag tgc tgg ttg tgg gag tcg ctg ctg ccc gac gag gtg cgc cga ctg 144  
 Gln Cys Trp Leu Trp Glu Ser Leu Leu Pro Asp Glu Val Arg Arg Leu  
 35 40 45  
 ccc gag gaa ctg gcc cggt gtg gac gca ttg ttg gac gat ccg gcg ttc 192  
 Pro Glu Glu Leu Ala Arg Val Asp Ala Leu Leu Asp Asp Pro Ala Phe  
 50 55 60  
 ttc gcc ccg ttc gtg ccg ttc gac ccg cgc agg ggc ccg ccg tcc 240  
 Phe Ala Pro Phe Val Pro Phe Phe Asp Pro Arg Arg Gly Arg Pro Ser  
 65 70 75 80  
 acg ccg atg gag gtc tat ctg cag ttg atg ttt gtg aag ttc cgc tac 288  
 Thr Pro Met Glu Val Tyr Leu Gln Leu Met Phe Val Lys Phe Arg Tyr  
 85 90 95  
 cggt ctg ggc tat gag tcg ctg tgc ccg gag gtg gct gat tcg atc acc 336  
 Arg Leu Gly Tyr Glu Ser Leu Cys Arg Glu Val Ala Asp Ser Ile Thr  
 100 105 110  
 tga 339  
 <210> 7  
 <211> 112  
 <212> PRT

&lt;213&gt; M Tuberculosis

&lt;400&gt; 7

Met	Ile	Pro	Gly	Arg	Met	Val	Leu	Asn	Trp	Glu	Asp	Gly	Leu	Asn	Ala
1					5				10				15		
Leu	Val	Ala	Glu	Gly	Ile	Glu	Ala	Ile	Val	Phe	Arg	Thr	Leu	Gly	Asp
					20				25				30		
Gln	Cys	Trp	Leu	Trp	Glu	Ser	Leu	Leu	Pro	Asp	Glu	Val	Arg	Arg	Leu
					35				40			45			
Pro	Glu	Glu	Leu	Ala	Arg	Val	Asp	Ala	Leu	Leu	Asp	Asp	Pro	Ala	Phe
					50				55			60			
Phe	Ala	Pro	Phe	Val	Pro	Phe	Phe	Asp	Pro	Arg	Arg	Gly	Arg	Pro	Ser
					65				70			75			80
Thr	Pro	Met	Glu	Val	Tyr	Leu	Gln	Leu	Met	Phe	Val	Lys	Phe	Arg	Tyr
					85				90			95			
Arg	Leu	Gly	Tyr	Glu	Ser	Leu	Cys	Arg	Glu	Val	Ala	Asp	Ser	Ile	Thr
					100				105						110

&lt;210&gt; 8

&lt;211&gt; 285

&lt;212&gt; DNA

&lt;213&gt; M Tuberculosis

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(285)

&lt;400&gt; 8

atg	acc	atc	aac	tat	caa	ttc	ggg	gac	gtc	gac	gct	cac	ggc	gcc	atg
Met	Thr	Ile	Asn	Tyr	Gln	Phe	Gly	Asp	Val	Asp	Ala	His	Gly	Ala	Met
1					5				10			15			

atc	ccg	gct	cag	gcc	ggg	tcg	ctg	gag	gcc	gag	cat	cag	gcc	atc	att
Ile	Arg	Ala	Gln	Ala	Gly	Ser	Leu	Glu	Ala	Glu	His	Gln	Ala	Ile	Ile
						20			25			30			

tct	gat	gtg	ttg	acc	gcg	agt	gac	ttt	tgg	ggc	ggc	gct	ggt	tcg	gcg
Ser	Asp	Val	Leu	Thr	Ala	Ser	Asp	Phe	Trp	Gly	Gly	Ala	Gly	Ser	Ala
					35			40			45				

gcc	tgc	cag	ggg	ttc	att	acc	cag	ctg	ggc	cgt	aac	ttc	cag	gtg	atc
Ala	Cys	Gln	Gly	Phe	Ile	Thr	Gln	Leu	Gly	Arg	Asn	Phe	Gln	Val	Ile
					50			55		60					

tac	gag	cag	gcc	aac	gcc	cac	ggg	cag	aag	gtg	cag	gct	gcc	ggc	aac
Tyr	Glu	Gln	Ala	Asn	Ala	His	Gly	Gln	Lys	Val	Gln	Ala	Ala	Gly	Asn
					65			70		75			80		

aac	atg	gca	caa	acc	gac	agc	gcc	gtc	ggc	tcc	agc	tgg	gcc	taa
Asn	Met	Ala	Gln	Thr	Asp	Ser	Ala	Val	Gly	Ser	Ser	Trp	Ala	*
					85				90					

&lt;210&gt; 9

&lt;211&gt; 94

&lt;212&gt; PRT

&lt;213&gt; M Tuberculosis

&lt;400&gt; 9

Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp Ala His Gly Ala Met  
 1 5 10 15  
 Ile Arg Ala Gln Ala Gly Leu Leu Glu Ala Glu His Gln Ala Ile Val  
 20 25 30  
 Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly Gly Ala Gly Ser Val  
 35 40 45  
 Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg Asn Phe Gln Val Ile  
 50 55 60  
 Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val Gln Ala Ala Gly Asn  
 65 70 75 80  
 Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser Ser Trp Ala  
 85 90

<210> 10  
 <211> 285  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(282)

<400> 10

atg acc atc aac tat cag ttc ggt gat gtc gac gct cat ggc gcc atg 48  
 Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp Ala His Gly Ala Met  
 1 5 10 15

atc cgc gct cag gcc ggg ttg ctg gag gcg gag cat cag gcc atc gtt 96  
 Ile Arg Ala Gln Ala Gly Leu Leu Glu Ala Glu His Gln Ala Ile Val  
 20 25 30

cgt gat gtg ttg gcc gcg ggt gac ttt tgg ggc ggc gcc ggt tcg gtg 144  
 Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly Ala Gly Ser Val  
 35 40 45

gct tgc cag gag ttc att acc cag ttg ggc cgt aac ttc cag gtg atc 192  
 Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg Asn Phe Gln Val Ile  
 50 55 60

taa gag cag gcc aac gcc cac ggg cag aag gtg cag gct gcc ggc aac 240  
 Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val Gln Ala Ala Gly Asn  
 65 70 75 80

aac atg gca caa acc gac agc gcc gtc ggc tcc agc tgg gcc 282  
 Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser Ser Trp Ala  
 85 90

tga 285

<210> 11  
 <211> 94  
 <212> PRT  
 <213> M Tuberculosis

<400> 11

Met Thr Ile Asn Tyr Gln Phe Gly Asp Val Asp Ala His Gly Ala Met  
 1 5 10 15  
 Ile Arg Ala Gln Ala Gly Leu Leu Glu Ala Glu His Gln Ala Ile Val  
 20 25 30

Arg Asp Val Leu Ala Ala Gly Asp Phe Trp Gly Gly Ala Gly Ser Val  
 35 40 45  
 Ala Cys Gln Glu Phe Ile Thr Gln Leu Gly Arg Asn Phe Gln Val Ile  
 50 55 60  
 Tyr Glu Gln Ala Asn Ala His Gly Gln Lys Val Gln Ala Ala Gly Asn  
 65 70 75 80  
 Asn Met Ala Gln Thr Asp Ser Ala Val Gly Ser Ser Trp Ala  
 85 90

<210> 12  
 <211> 327  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(327)

<400> 12  
 gtg ctt ttg cct ctt ggt ccg cct ttg ccg ccc gac gcg gtg gtg gcg 48  
 Val Leu Leu Pro Leu Gly Pro Pro Leu Pro Pro Asp Ala Val Val Ala  
 1 5 10 15  
 aaa cg gct gag tcg gga atg ctc ggc ggg ttg tcg gtt ccg ctc agc 96  
 Lys Arg Ala Glu Ser Gly Met Leu Gly Gly Leu Ser Val Pro Leu Ser  
 20 25 30  
 tgg gga gtg gct gtg cca ccc gat gat tat gac cac tgg gcg cct gcg 144  
 Trp Gly Val Ala Val Pro Pro Asp Asp Tyr Asp His Trp Ala Pro Ala  
 35 40 45  
 ccg gag gac ggc gcc gat gtc gat gtc cag gcg gcc gaa ggg gcg gac 192  
 Pro Glu Asp Gly Ala Asp Val Asp Val Gln Ala Ala Glu Gly Ala Asp  
 50 55 60  
 gca gag gcc gcc atg gac gag tgg gat gag tgg cag gcg tgg aac 240  
 Ala Glu Ala Ala Met Asp Glu Trp Asp Glu Trp Gln Ala Trp Asn  
 65 70 75 80  
 gag tgg gtg gcg gag aac gct gaa ccc cgc ttt gag gtg cca cgg agt 288  
 Glu Trp Val Ala Glu Asn Ala Glu Pro Arg Phe Glu Val Pro Arg Ser  
 85 90 95  
 agc agc agc gtg att ccg cat tct ccg gcg gcc ggc tag 327  
 Ser Ser Ser Val Ile Pro His Ser Pro Ala Ala Gly \*  
 100 105

<210> 13  
 <211> 108  
 <212> PRT  
 <213> M Tuberculosis

<400> 13

Met Leu Leu Pro Leu Gly Pro Pro Leu Pro Pro Asp Ala Val Val Ala  
 1 5 10 15  
 Lys Arg Ala Glu Ser Gly Met Leu Gly Gly Leu Ser Val Pro Leu Ser  
 20 25 30  
 Trp Gly Val Ala Val Pro Pro Asp Asp Tyr Asp His Trp Ala Pro Ala

35	40	45
Pro Glu Asp Gly Ala Asp Val Asp Val Gln Ala Ala Glu Gly Ala Asp		
50	55	60
Ala Glu Ala Ala Ala Met Asp Glu Trp Asp Glu Trp Gln Ala Trp Asn		
65	70	75
Glu Trp Val Ala Glu Asn Ala Glu Pro Arg Phe Glu Val Pro Arg Ser		80
85	90	95
Ser Ser Ser Val Ile Pro His Ser Pro Ala Ala Gly		
100	105	

<210> 14  
 <211> 324  
 <212> DNA  
 <213> M Tuberculosis

<220>															
<221> CDS															
<222> (1)...(324)															
<400> 14															
<table border="0"> <tr> <td>ttg acc cac aag cgc act aaa cgc cag cca gcc atc gcc gca ggg ctc</td> <td></td> <td></td> <td style="text-align: right;">48</td> </tr> <tr> <td>Leu Thr His Lys Arg Thr Lys Arg Gln Pro Ala Ile Ala Ala Gly Leu</td> <td></td> <td></td> <td></td> </tr> <tr> <td>1</td> <td>5</td> <td>10</td> <td>15</td> </tr> </table>				ttg acc cac aag cgc act aaa cgc cag cca gcc atc gcc gca ggg ctc			48	Leu Thr His Lys Arg Thr Lys Arg Gln Pro Ala Ile Ala Ala Gly Leu				1	5	10	15
ttg acc cac aag cgc act aaa cgc cag cca gcc atc gcc gca ggg ctc			48												
Leu Thr His Lys Arg Thr Lys Arg Gln Pro Ala Ile Ala Ala Gly Leu															
1	5	10	15												
<table border="0"> <tr> <td>aac gcc ccg cgt cgg aat cgc gtt ggg cgg caa cat ggt tgg ccg gcc</td> <td></td> <td></td> <td style="text-align: right;">96</td> </tr> <tr> <td>Asn Ala Pro Arg Arg Asn Arg Val Gly Arg Gln His Gly Trp Pro Ala</td> <td></td> <td></td> <td></td> </tr> <tr> <td>20</td> <td>25</td> <td>30</td> <td></td> </tr> </table>				aac gcc ccg cgt cgg aat cgc gtt ggg cgg caa cat ggt tgg ccg gcc			96	Asn Ala Pro Arg Arg Asn Arg Val Gly Arg Gln His Gly Trp Pro Ala				20	25	30	
aac gcc ccg cgt cgg aat cgc gtt ggg cgg caa cat ggt tgg ccg gcc			96												
Asn Ala Pro Arg Arg Asn Arg Val Gly Arg Gln His Gly Trp Pro Ala															
20	25	30													
<table border="0"> <tr> <td>gac gtt ccg tcc gcc gag cag cgc cgc gcc caa cgg cag cgc gac ctc</td> <td></td> <td></td> <td style="text-align: right;">144</td> </tr> <tr> <td>Asp Val Pro Ser Ala Glu Gln Arg Arg Ala Gln Arg Gln Arg Asp Leu</td> <td></td> <td></td> <td></td> </tr> <tr> <td>35</td> <td>40</td> <td>45</td> <td></td> </tr> </table>				gac gtt ccg tcc gcc gag cag cgc cgc gcc caa cgg cag cgc gac ctc			144	Asp Val Pro Ser Ala Glu Gln Arg Arg Ala Gln Arg Gln Arg Asp Leu				35	40	45	
gac gtt ccg tcc gcc gag cag cgc cgc gcc caa cgg cag cgc gac ctc			144												
Asp Val Pro Ser Ala Glu Gln Arg Arg Ala Gln Arg Gln Arg Asp Leu															
35	40	45													
<table border="0"> <tr> <td>gag gct atc cgc cga gcg tac gcc gag atg gtg gcg aca tca cac gaa</td> <td></td> <td></td> <td style="text-align: right;">192</td> </tr> <tr> <td>Glu Ala Ile Arg Arg Ala Tyr Ala Glu Met Val Ala Thr Ser His Glu</td> <td></td> <td></td> <td></td> </tr> <tr> <td>50</td> <td>55</td> <td>60</td> <td></td> </tr> </table>				gag gct atc cgc cga gcg tac gcc gag atg gtg gcg aca tca cac gaa			192	Glu Ala Ile Arg Arg Ala Tyr Ala Glu Met Val Ala Thr Ser His Glu				50	55	60	
gag gct atc cgc cga gcg tac gcc gag atg gtg gcg aca tca cac gaa			192												
Glu Ala Ile Arg Arg Ala Tyr Ala Glu Met Val Ala Thr Ser His Glu															
50	55	60													
<table border="0"> <tr> <td>atc gac gac gac aca gcc gaa ctg gcg ctg ttg tcg atg cat ctc gac</td> <td></td> <td></td> <td style="text-align: right;">240</td> </tr> <tr> <td>Ile Asp Asp Asp Thr Ala Glu Leu Ala Leu Leu Ser Met His Leu Asp</td> <td></td> <td></td> <td></td> </tr> <tr> <td>65</td> <td>70</td> <td>75</td> <td>80</td> </tr> </table>				atc gac gac gac aca gcc gaa ctg gcg ctg ttg tcg atg cat ctc gac			240	Ile Asp Asp Asp Thr Ala Glu Leu Ala Leu Leu Ser Met His Leu Asp				65	70	75	80
atc gac gac gac aca gcc gaa ctg gcg ctg ttg tcg atg cat ctc gac			240												
Ile Asp Asp Asp Thr Ala Glu Leu Ala Leu Leu Ser Met His Leu Asp															
65	70	75	80												
<table border="0"> <tr> <td>gat gag cag cgc cgg ctt gag gcg ggg atg aag ctc ggc tgg cat ccg</td> <td></td> <td></td> <td style="text-align: right;">288</td> </tr> <tr> <td>Asp Glu Gln Arg Leu Glu Ala Gly Met Lys Leu Gly Trp His Pro</td> <td></td> <td></td> <td></td> </tr> <tr> <td>85</td> <td>90</td> <td>95</td> <td></td> </tr> </table>				gat gag cag cgc cgg ctt gag gcg ggg atg aag ctc ggc tgg cat ccg			288	Asp Glu Gln Arg Leu Glu Ala Gly Met Lys Leu Gly Trp His Pro				85	90	95	
gat gag cag cgc cgg ctt gag gcg ggg atg aag ctc ggc tgg cat ccg			288												
Asp Glu Gln Arg Leu Glu Ala Gly Met Lys Leu Gly Trp His Pro															
85	90	95													
<table border="0"> <tr> <td>tat cac ttc ccc gac gaa ccc gac agc aaa cag tga</td> <td></td> <td></td> <td style="text-align: right;">324</td> </tr> <tr> <td>Tyr His Phe Pro Asp Glu Pro Asp Ser Lys Gln *</td> <td></td> <td></td> <td></td> </tr> <tr> <td>100</td> <td>105</td> <td></td> <td></td> </tr> </table>				tat cac ttc ccc gac gaa ccc gac agc aaa cag tga			324	Tyr His Phe Pro Asp Glu Pro Asp Ser Lys Gln *				100	105		
tat cac ttc ccc gac gaa ccc gac agc aaa cag tga			324												
Tyr His Phe Pro Asp Glu Pro Asp Ser Lys Gln *															
100	105														

<210> 15  
 <211> 107  
 <212> PRT  
 <213> M Tuberculosis

<400> 15			
Met Thr His Lys Arg Thr Lys Arg Gln Pro Ala Ile Ala Ala Gly Leu			
1	5	10	15
Asn Ala Pro Arg Arg Asn Arg Val Gly Arg Gln His Gly Trp Pro Ala			
20	25	30	

Asp Val Pro Ser Ala Glu Gln Arg Arg Ala Gln Arg Gln Arg Asp Leu  
 35 40 45  
 Glu Ala Ile Arg Arg Ala Tyr Ala Glu Met Val Ala Thr Ser His Glu  
 50 55 60  
 Ile Asp Asp Asp Thr Ala Glu Leu Ala Leu Ser Met His Leu Asp  
 65 70 75 80  
 Asp Glu Gln Arg Arg Leu Glu Ala Gly Met Lys Leu Gly Trp His Pro  
 85 90 95  
 Tyr His Phe Pro Asp Glu Pro Asp Ser Lys Gln  
 100 105

<210> 16  
 <211> 246  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1) . . . (246)

<400> 16  
 atg agc ggc cac gcg ttg gct gct cggtt acg ctg gcc gac 48  
 Met Ser Gly His Ala Leu Ala Arg Thr Leu Leu Ala Ala Ala Asp  
 1 5 10 15  
 gag ctt gtc ggc ggc ccg cca gtc gag gct tcg gcc gcc gcg ctg gcc 96  
 Glu Leu Val Gly Pro Pro Val Glu Ala Ser Ala Ala Leu Ala  
 20 25 30  
 ggc gac gcc ggc gca tgg cgg acc gcg gcc gtc gag ctt gcg cga 144  
 Gly Asp Ala Ala Gly Ala Trp Arg Thr Ala Ala Val Glu Leu Ala Arg  
 35 40 45  
 gcg ttg gtc cgc gct gtg gcg gag tcg cac ggc gtc gcg gcc gtt ttg 192  
 Ala Leu Val Arg Ala Val Ala Glu Ser His Gly Val Ala Ala Val Leu  
 50 55 60  
 ttc gcc gcg acg gcc gcc gcg gcg gcc gtc gac cgg ggt gat ccg 240  
 Phe Ala Ala Thr Ala Ala Ala Ala Val Asp Arg Gly Asp Pro  
 65 70 75 80  
 ccg tga 246  
 Pro \*

<210> 17  
 <211> 81  
 <212> PRT  
 <213> M Tuberculosis  
 <400> 17  
 Met Ser Gly His Ala Leu Ala Ala Arg Thr Leu Leu Ala Ala Asp  
 1 5 10 15  
 Glu Leu Val Gly Pro Pro Val Glu Ala Ser Ala Ala Leu Ala  
 20 25 30  
 Gly Asp Ala Ala Gly Ala Trp Arg Thr Ala Ala Val Glu Leu Ala Arg  
 35 40 45  
 Ala Leu Val Arg Ala Val Ala Glu Ser His Gly Val Ala Ala Val Leu

50	55	60
Phe Ala Ala Thr Ala Ala Ala Ala Ala Val Asp Arg Gly Asp Pro		
65	70	75
Pro		80

<210> 18  
 <211> 294  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(294)

<400> 18

atg	agt	ttg	ttg	gat	gcc	cat	att	ccg	cag	ttg	atc	gct	tcg	cat	acg	48
Met	Ser	Leu	Leu	Asp	Ala	His	Ile	Pro	Gln	Leu	Ile	Ala	Ser	His	Thr	
1		5						10				15				

gcg	ttt	gcc	gct	aag	gcg	ggg	ttg	atg	cg	cat	acg	atc	ggt	cag	gcc	96
Ala	Phe	Ala	Ala	Lys	Ala	Gly	Leu	Met	Arg	His	Thr	Ile	Gly	Gln	Ala	
20							25					30				

gag	cag	cag	gcg	atg	tcg	g	c	c	g	ttt	cat	cag	gga	gag	tcc	g	144
Glu	Gln	Gln	Ala	Met	Ser	Ala	Gln	Ala	Phe	His	Gln	Gly	Glu	Ser	Ala		
35							40				45						

g	c	g	ttt	c	ag	g	gt	g	c	at	g	cc	g	cc	g	a	192
Ala	Ala	Phe	Gln	Gly	Ala	His	Ala	Arg	Phe	Val	Ala	Ala	Ala	Lys			
50							55			60							

g	t	c	aat	acc	ttt	ctg	gat	atc	g	c	aa	g	cc	g	cc	g	240
Val	Asn	Thr	Leu	Leu	Asp	Ile	Ala	Gln	Ala	Asn	Leu	Gly	Glu	Ala	Ala		
65							70			75		80					

g	gc	ac	t	at	gt	g	cc	g	at	g	cc	g	cc	g	cc	g	288
Gly	Thr	Tyr	Val	Ala	Ala	Asp	Ala	Ala	Ala	Ser	Ser	Tyr	Thr	Gly			
85							90			95							

ttt	tta															294
Phe	Leu															

<210> 19  
 <211> 97  
 <212> PRT  
 <213> M Tuberculosis

<400> 19

Met	Ser	Leu	Leu	Asp	Ala	His	Ile	Pro	Gln	Leu	Ile	Ala	Ser	His	Thr	
1		5						10			15					
Ala	Phe	Ala	Ala	Lys	Ala	Gly	Leu	Met	Arg	His	Thr	Ile	Gly	Gln	Ala	
20							25			30						
Glu	Gln	Gln	Ala	Met	Ser	Ala	Gln	Ala	Phe	His	Gln	Gly	Glu	Ser	Ala	
35							40			45						
Ala	Ala	Phe	Gln	Gly	Ala	His	Ala	Arg	Phe	Val	Ala	Ala	Ala	Lys		
50							55			60						

Val Asn Thr Leu Leu Asp Ile Ala Gln Ala Asn Leu Gly Glu Ala Ala  
 65 70 75 80  
 Gly Thr Tyr Val Ala Ala Asp Ala Ala Ala Ser Ser Tyr Thr Gly  
 85 90 95  
 Phe

<210> 20  
 <211> 303  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(303)

<400> 20

atg aac gca gac ccc gtg ttg tcg tac aac ttt gac gcc atc gaa tac 48  
 Met Asn Ala Asp Pro Val Leu Ser Tyr Asn Phe Asp Ala Ile Glu Tyr  
 1 5 10 15

tcc gtt cgt cag gag atc cac acc acc gcg gcc cgt ttc aac gct gcg 96  
 Ser Val Arg Gln Glu Ile His Thr Thr Ala Ala Arg Phe Asn Ala Ala  
 20 25 30

ctg caa gag ctg agg tcg cag atc gcg ccg ttg cag cag ctc tgg aca 144  
 Leu Gln Glu Leu Arg Ser Gln Ile Ala Pro Leu Gln Gln Leu Trp Thr  
 35 40 45

cgg gaa gcg gcc gcc tac cac gcg gag caa ctc aag tgg cac cag 192  
 Arg Glu Ala Ala Ala Tyr His Ala Glu Gln Leu Lys Trp His Gln  
 50 55 60

gcg gcc agc gcg ctc aac gag atc ctg atc gac ttg gga aac gcg gtt 240  
 Ala Ala Ser Ala Leu Asn Glu Ile Leu Ile Asp Leu Gly Asn Ala Val  
 65 70 75 80

cgc cac ggt gcc gac gac gtg gcg cat gcc gac cgg cgg gcg gct gga 288  
 Arg His Gly Ala Asp Asp Val Ala His Ala Asp Arg Arg Ala Ala Gly  
 85 90 95

gct tgg gca cgc tag 303  
 Ala Trp Ala Arg \*  
 100

<210> 21  
 <211> 100  
 <212> PRT  
 <213> M Tuberculosis

<400> 21

Met Asn Ala Asp Pro Val Leu Ser Tyr Asn Phe Asp Ala Ile Glu Tyr  
 1 5 10 15  
 Ser Val Arg Gln Glu Ile His Thr Thr Ala Ala Arg Phe Asn Ala Ala  
 20 25 30  
 Leu Gln Glu Leu Arg Ser Gln Ile Ala Pro Leu Gln Gln Leu Trp Thr  
 35 40 45  
 Arg Glu Ala Ala Ala Tyr His Ala Glu Gln Leu Lys Trp His Gln

50	55	60														
Ala	Ala	Ser	Ala	Leu	Asn	Glu	Ile	Leu	Ile	Asp	Leu	Gly	Asn	Ala	Val	
65				70				75						80		
Arg	His	Gly	Ala	Asp	Asp	Val	Ala	His	Ala	Asp	Arg	Arg	Arg	Ala	Ala	Gly
				85				90						95		
Ala	Trp	Ala	Arg													
				100												

<210> 22  
 <211> 378  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1) ... (378)

<400> 22	48
ttg gtt gaa ccg gga agg atc gga ggg aac cag acg agg ttg gcg gcg	
Leu Val Glu Pro Gly Arg Ile Gly Gly Asn Gln Thr Arg Leu Ala Ala	
1 5 10 15	
gtc cta ctt gat gtg agc aca ccg aac acg ctg aac gcc gac ttt gac	
Val Leu Leu Asp Val Ser Thr Pro Asn Thr Leu Asn Ala Asp Phe Asp	
20 25 30	
ctg atg cgt tcg gtt ggt atc acg gac gcc cgc aat gag gaa atc	
Leu Met Arg Ser Val Ala Gly Ile Thr Asp Ala Arg Asn Glu Glu Ile	
35 40 45	
cgt gcg atg ctg cag gca ttc atc ggc cgc atg acg ggt gtg ccg ccg	
Arg Ala Met Leu Gln Ala Phe Ile Gly Arg Met Ser Gly Val Pro Pro	
50 55 60	
tcg gtg tgg ggt ggg ctc gcg gct ttc cag gat gtg gtg gat	
Ser Val Trp Gly Gly Leu Ala Ala Arg Phe Gln Asp Val Val Asp	
65 70 75 80	
cgc tgg aac gcc gag tcg acg cgg ctc tac cac gtc ctg cac gcg atc	
Arg Trp Asn Ala Glu Ser Thr Arg Leu Tyr His Val Leu His Ala Ile	
85 90 95	
gcc gac acc atc cgc cac aac gag gcc gcg ctg cgg gaa gcc ggc caa	
Ala Asp Thr Ile Arg His Asn Glu Ala Ala Leu Arg Glu Ala Gly Gln	
100 105 110	
atc cat gcc cgc cac atc gcc gcc ggc ggc gac cta tag	
Ile His Ala Arg His Ile Ala Ala Gly Gly Asp Leu *	
115 120 125	
<210> 23	
<211> 125	
<212> PRT	
<213> M Tuberculosis	
<400> 23	
Met Val Glu Pro Gly Arg Ile Gly Gly Asn Gln Thr Arg Leu Ala Ala	
1 5 10 15	

Val Leu Leu Asp Val Ser Thr Pro Asn Thr Leu Asn Ala Asp Phe Asp  
 20 25 30  
 Leu Met Arg Ser Val Ala Gly Ile Thr Asp Ala Arg Asn Glu Glu Ile  
 35 40 45  
 Arg Ala Met Leu Gln Ala Phe Ile Gly Arg Met Ser Gly Val Pro Pro  
 50 55 60  
 Ser Val Trp Gly Gly Leu Ala Ala Ala Arg Phe Gln Asp Val Val Asp  
 65 70 75 80  
 Arg Trp Asn Ala Glu Ser Thr Arg Leu Tyr His Val Leu His Ala Ile  
 85 90 95  
 Ala Asp Thr Ile Arg His Asn Glu Ala Ala Leu Arg Glu Ala Gly Gln  
 100 105 110  
 Ile His Ala Arg His Ile Ala Ala Gly Gly Asp Leu  
 115 120 125

<210> 24  
 <211> 288  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(288)

<400> 24

atg tca gat caa atc acg tat aac ccg gga gcc gta tcc gac ttc gct 48  
 Met Ser Asp Gln Ile Thr Tyr Asn Pro Gly Ala Val Ser Asp Phe Ala  
 1 5 10 15

tcc gac gtg ggc tcg cgc gcc ggc cag ctc cac atg att tac gaa gac 96  
 Ser Asp Val Gly Ser Arg Ala Gly Gln Leu His Met Ile Tyr Glu Asp  
 20 25 30

acc gcc agc aaa aca aat gcg ctg caa gag ttt ttc gcg ggc cac ggc 144  
 Thr Ala Ser Lys Thr Asn Ala Leu Gln Glu Phe Phe Ala Gly His Gly  
 35 40 45

gcg caa ggg ttt ttc gac gcc cag gcg cag atg ctg tcg ggg ctg cag 192  
 Ala Gln Gly Phe Phe Asp Ala Gln Ala Gln Met Leu Ser Gly Leu Gln  
 50 55 60

ggg ctc att gag acg gtg ggt cag cat ggg act acc acc ggc cac gtg 240  
 Gly Leu Ile Glu Thr Val Gly Gln His Gly Thr Thr Gly His Val  
 65 70 75 80

ctg gac aac gcg atc gga acc gac cag gcc atc gcg ggc ttg ttc taa 288  
 Leu Asp Asn Ala Ile Gly Thr Asp Gln Ala Ile Ala Gly Leu Phe \*  
 85 90 95

<210> 25  
 <211> 95  
 <212> PRT  
 <213> M Tuberculosis

<400> 25

Met Ser Asp Gln Ile Thr Tyr Asn Pro Gly Ala Val Ser Asp Phe Ala  
 1 5 10 15  
 Ser Asp Val Gly Ser Arg Ala Gly Gln Leu His Met Ile Tyr Glu Asp

20	25	30	
Thr Ala Ser Lys Thr Asn Ala Leu Gln Glu Phe Phe Ala Gly His Gly			
35	40	45	
Ala Gln Gly Phe Phe Asp Ala Gln Ala Gln Met Leu Ser Gly Leu Gln			
50	55	60	
Gly Leu Ile Glu Thr Val Gln His Gly Thr Thr Gly His Val			
65	70	75	80
Leu Asp Asn Ala Ile Gly Thr Asp Gln Ala Ile Ala Gly Leu Phe			
85	90	95	

<210> 26  
 <211> 324  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(324)

<400> 26

gtg gca gac aca att cag gta aca ccg cag atg ctg cgc agc acc gcc	48		
Val Ala Asp Thr Ile Gln Val Thr Pro Gln Met Leu Arg Ser Thr Ala			
1	5	10	15

aac gat atc cag gcg aat atg gag caa gcc atg gga atc gcc aag ggc	96		
Asn Asp Ile Gln Ala Asn Met Glu Gln Ala Met Gly Ile Ala Lys Gly			
20	25	30	

tac cta gcc aac cag gaa aac gtc atg aac ccc gcc acc tgg tct ggt	144		
Tyr Leu Ala Asn Gln Glu Asn Val Met Asn Pro Ala Thr Trp Ser Gly			
35	40	45	

acc ggc gtc gtt gct tcg cat atg aca gcc acc gag atc acc aat gaa	192		
Thr Gly Val Val Ala Ser His Met Thr Ala Thr Glu Ile Thr Asn Glu			
50	55	60	

ttg aac aag gtc ctt acc ggg ggc acg cgc ctg gcc gag ggc ctc gtg	240		
Leu Asn Lys Val Leu Thr Gly Gly Thr Arg Leu Ala Glu Gly Leu Val			
65	70	75	80

cag gcc gca gcc ctg atg gag gga cac gag gcg gac tcg cag aca gcg	288		
Gln Ala Ala Ala Leu Met Glu Gly His Glu Ala Asp Ser Gln Thr Ala			
85	90	95	

ttt cag gcg ctg ttc ggc gct agc cac gga tcc tga	324		
Phe Gln Ala Leu Phe Gly Ala Ser His Gly Ser *			
100	105		

<210> 27  
 <211> 107  
 <212> PRT  
 <213> M Tuberculosis

<400> 27

Met Ala Asp Thr Ile Gln Val Thr Pro Gln Met Leu Arg Ser Thr Ala			
1	5	10	15
Asn Asp Ile Gln Ala Asn Met Glu Gln Ala Met Gly Ile Ala Lys Gly			
20	25	30	

Tyr Leu Ala Asn Gln Glu Asn Val Met Asn Pro Ala Thr Trp Ser Gly  
 35 40 45  
 Thr Gly Val Val Ala Ser His Met Thr Ala Thr Glu Ile Thr Asn Glu  
 50 55 60  
 Leu Asn Lys Val Leu Thr Gly Gly Thr Arg Leu Ala Glu Gly Leu Val  
 65 70 75 80  
 Gln Ala Ala Ala Leu Met Glu Gly His Glu Ala Asp Ser Gln Thr Ala  
 85 90 95  
 Phe Gln Ala Leu Phe Gly Ala Ser His Gly Ser  
 100 105

<210> 28  
 <211> 273  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1) . . . (273)

<400> 28  
 gtg gat ccg acc gtg ttg gct gat gcg gtg gcg cgg atg gcc gaa ttc 48  
 Val Asp Pro Thr Val Leu Ala Asp Ala Val Ala Arg Met Ala Glu Phe  
 1 5 10 15

ggt cgc cac gtc gag gag ctg gtc gcc gag att gag tcc ttg gtt acc 96  
 Gly Arg His Val Glu Glu Leu Val Ala Glu Ile Glu Ser Leu Val Thr  
 20 25 30

cgg ctg cat gtg acg tgg acg ggg gag ggc gcg gct cat gct gag 144  
 Arg Leu His Val Thr Trp Thr Gly Glu Gly Ala Ala Ala His Ala Glu  
 35 40 45

gcg caa cga cat tgg gct gcc ggt gag gcg atg atg cgc cag gcg ttg 192  
 Ala Gln Arg His Trp Ala Ala Gly Glu Ala Met Met Arg Gln Ala Leu  
 50 55 60

gcc cag ctc acg gcc ggg cag acg gcg cac gcc aac tac acc ggc 240  
 Ala Gln Leu Thr Ala Ala Gly Gln Ser Ala His Ala Asn Tyr Thr Gly  
 65 70 75 80

gcg atg gcc acg aat ttg ggt atg tgg tcg tga 273  
 Ala Met Ala Thr Asn Leu Gly Met Trp Ser \*  
 85 90

<210> 29  
 <211> 90  
 <212> PRT  
 <213> M Tuberculosis

<400> 29  
 Met Asp Pro Thr Val Leu Ala Asp Ala Val Ala Arg Met Ala Glu Phe  
 1 5 10 15  
 Gly Arg His Val Glu Glu Leu Val Ala Glu Ile Glu Ser Leu Val Thr  
 20 25 30  
 Arg Leu His Val Thr Trp Thr Gly Glu Gly Ala Ala Ala His Ala Glu  
 35 40 45  
 Ala Gln Arg His Trp Ala Ala Gly Glu Ala Met Met Arg Gln Ala Leu

50 55 60  
 Ala Gln Leu Thr Ala Ala Gly Gln Ser Ala His Ala Asn Tyr Thr Gly  
 65 70 75 80  
 Ala Met Ala Thr Asn Leu Gly Met Trp Ser  
 85 90

<210> 30  
 <211> 312  
 <212> DNA  
 <213> M Tuberculosis

<220>  
 <221> CDS  
 <222> (1)...(312)

<400> 30

atg ggt gcc gac gac acg ctg cgc gta gag cct gcg gtg atg cag cag ggt 48  
 Met Gly Ala Asp Asp Thr Leu Arg Val Glu Pro Ala Val Met Gln Gly  
 1 5 10 15

ttc gcc gcg tcg ttg gat gga gcg gcc gag cat ctc gcg gtt caa ctg 96  
 Phe Ala Ala Ser Leu Asp Gly Ala Ala Glu His Leu Ala Val Gln Leu  
 20 25 30

gcc gag ctg gac gct cag gtc ggg cag atg ttg ggc ggg tgg cgc ggg 144  
 Ala Glu Leu Asp Ala Gln Val Gly Gln Met Leu Gly Gly Trp Arg Gly  
 35 40 45

gcg tcg ggc agt gcg tat ggc tcg gcg tgg gag cta tgg cat cgc ggg 192  
 Ala Ser Gly Ser Ala Tyr Gly Ser Ala Trp Glu Leu Trp His Arg Gly  
 50 55 60

gcc ggt gag gtg cag ctg gga ttg tcg atg ctg gcg gcg gcg ata gct 240  
 Ala Gly Glu Val Gln Leu Gly Leu Ser Met Leu Ala Ala Ala Ile Ala  
 65 70 75 80

cac gcc ggt gcg ggt tat caa cac aac gag acc gcg tcg gcg cag gtg 288  
 His Ala Gly Ala Gly Tyr Gln His Asn Glu Thr Ala Ser Ala Gln Val  
 85 90 95

ctt cgt gag gtg ggc ggt ggc tga 312  
 Leu Arg Glu Val Gly Gly \*  
 100

<210> 31  
 <211> 103  
 <212> PRT  
 <213> M Tuberculosis

<400> 31

Met Gly Ala Asp Asp Thr Leu Arg Val Glu Pro Ala Val Met Gln Gly  
 1 5 10 15  
 Phe Ala Ala Ser Leu Asp Gly Ala Ala Glu His Leu Ala Val Gln Leu  
 20 25 30  
 Ala Glu Leu Asp Ala Gln Val Gly Gln Met Leu Gly Gly Trp Arg Gly  
 35 40 45  
 Ala Ser Gly Ser Ala Tyr Gly Ser Ala Trp Glu Leu Trp His Arg Gly  
 50 55 60

Ala Gly Glu Val Gln Leu Gly Leu Ser Met Leu Ala Ala Ala Ile Ala  
65 70 75 80  
His Ala Gly Ala Gly Tyr Gln His Asn Glu Thr Ala Ser Ala Gln Val  
85 90 95  
Leu Arg Glu Val Gly Gly Gly  
100

## THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A fusion polypeptide, which comprises an amino acid sequence chosen from the group consisting of:

5           1) an amino acid sequence analogue having a sequence identity with SEQ ID NO: 7 of at least 70% and at the same time being immunologically equivalent to SEQ ID NO: 7;

10           2) a T-cell epitope of the amino acid sequence as shown in SEQ ID NO: 7; and

10           3) an amino acid sequence, which has a sequence identity of at least 70% with the T-cell epitope in 2) and at the same time being immunologically equivalent to SEQ ID NO: 7,

and at least one fusion partner.

2. The fusion polypeptide according to claim 1, wherein the fusion partner is selected  
15 from the group consisting of

1) a polypeptide fragment, which comprises an amino acid sequence as shown in SEQ ID NOs: 7, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31;

20           2) a polypeptide fragment, which comprises an amino acid sequence analogue having a sequence identity with an amino acid sequence as shown in SEQ ID NOs: 7, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31 of at least 70% and at the same time being immunologically equivalent to said sequence;

25           3) a polypeptide fragment, which comprises a T-cell epitope of amino acid sequence as shown in SEQ ID NOs: 7, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31;

30           4) a polypeptide fragment, which comprises an amino acid sequence, which has a sequence identity of at least 70% with a T-cell epitope of the amino acid sequence as shown in SEQ ID NOs: 7, 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31 and at the same time being immunologically equivalent to the amino acid sequence; and

5) a polypeptide fragment derived from a bacterium belonging to the tuberculosis complex, such as ESAT-6 or at least one T-cell epitope thereof, TB10.4 or at least one T-cell epitope thereof, and MPT59 or at least one T-cell epitope thereof.

35 3. The fusion polypeptide fragment according to either claim 1 or claim 2, wherein the fusion partner is selected from the group consisting of DnaK, GroEL, urease, glutamine synthetase, the proline rich complex, L-alanine dehydrogenase, phosphate binding protein,

Ag 85 complex, HBHA (heparin binding hemagglutinin), MPT51, superoxide dismutase, 19 kDa lipoprotein,  $\alpha$ -crystallin, GroES, and MPT59.

4. The fusion polypeptide according to any one of the preceding claims, which is 5 lipidated so as to allow a self-adjuvating effect of the polypeptide.

5. A substantially pure polypeptide fragment which comprises an amino acid sequence chosen from the group consisting of:

- 1) an amino acid sequence analogue having a sequence identity with SEQ ID 10 NO: 7 of at least 70% and at the same time being immunologically equivalent to SEQ ID NO: 7;
- 2) a T-cell epitope of the amino acid sequence as shown in SEQ ID NO: 7; and
- 3) an amino acid sequence, which has a sequence identity of at least 70% with 15 the T-cell epitope in 2) and at the same time being immunologically equivalent to SEQ ID NO: 7,

when used in a pharmaceutical.

6. The polypeptide fragment according to claim 5 in essentially pure form, when 20 used in a pharmaceutical.

7. The polypeptide fragment according to either claim 5 or claim 6, which has a length of at least 7 amino acid residues, when used in a pharmaceutical.

8. The polypeptide fragment according to any one of claims 5 to 7, which has a length 25 of at least 14 amino acid residues, when used in a pharmaceutical.

9. The polypeptide fragment according to any one of claims 5 to 8, which is free from any signal sequence, when used in a pharmaceutical.

30 10. The fusion polypeptide, according to any one of claims 1 to 4, when used in a pharmaceutical.

11. The polypeptide fragment according to any one of claims 5 to 9, wherein the sequence identity is at least 80%, when used in a pharmaceutical.

35 12. The polypeptide fragment according to any one of claims 5 to 9 and 11, wherein the sequence identity is at least 90%, when used in a pharmaceutical.

13. Use of a substantially pure polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- Rv3386

5

in the preparation of a pharmaceutical composition for the diagnosis of or vaccination against tuberculosis caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis* in a mammal.

10 14. An immunologic composition, which comprises a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- Rv3386.

15

15. The immunologic composition according to claim 14, which further comprises at least one polypeptide fragment comprising an amino acid sequence chosen from the group consisting of:

20

- an amino acid sequence as shown in SEQ ID NO: 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31;

25

- an analogue having a sequence identity with an amino acid sequence as shown in SEQ ID NO: 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31 of at least 70% and at the same time being immunologically equivalent to said amino acid sequence;

30

- a T-cell epitope of a amino acid sequence as shown in SEQ ID NO: 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31; and
- an amino acid sequence which has a sequence identity of at least 70% with a T-cell epitope of an amino acid sequence as shown in SEQ ID NO: 13, 15, 17, 19, 21, 23, 25, 27, 29 or 31 and at the same time being immunologically equivalent to said amino acid sequence.

35

16. The immunologic composition according to either claim 14 or claim 15, which further comprises an immunologically and pharmaceutically acceptable carrier, vehicle or adjuvant.

17. The immunologic composition according to claim 16, wherein the carrier is selected from the group consisting of a polymer to which the polypeptide(s) is/are bound

by hydrophobic non-covalent interaction, a polymer to which the polypeptide(s) is/are covalently bound, and a polypeptide; the vehicle is selected from the group consisting of a diluent and a suspending agent; and the adjuvant is selected from the group consisting of dimethyldioctadecylammonium bromide (DDA), Quil A, poly I:C, Freund's incomplete 5 adjuvant, IFN- $\gamma$ , IL-2, IL-12, monophosphoryl lipid A (MPL), and muramyl dipeptide (MDP).

18. The immunologic composition according to claim 17, wherein the polymer to which the polypeptide(s) is/are bound by hydrophobic non-covalent interaction is a plastic.

10

19. The immunologic composition according to claim 18, wherein the plastic is polystyrene.

15

20. The immunologic composition according to any one of claims 17 to 19, wherein the polymer to which the polypeptide(s) is/are covalently bound is a polysaccharide.

21. The immunologic composition according to any one of claims 17 to 20, wherein the polypeptide is selected from the group consisting of bovine serum albumin, ovalbumin and keyhole limpet hemocyanin.

20

22. The immunologic composition according to any of claims 14 to 21, comprising at least two different polypeptide fragments, each different polypeptide fragment being a polypeptide chosen from the group consisting of:

25

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- Rv3386.

23. The immunologic composition according to claim 22, comprising 3 - 20 different polypeptide fragments.

30

24. The immunologic composition according to any one of claims 14 to 23, which is in the form of a vaccine.

35

25. The immunologic composition according to any one of claims 14 to 23, which is in the form of a skin test reagent.

26. A vaccine for immunizing an animal, including a human being, against tuberculosis caused by mycobacteria belonging to the tuberculosis complex, comprising as the effective component a non-pathogenic microorganism, wherein at least one copy of a DNA fragment comprising a DNA sequence encoding a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- Rv3386

has been incorporated into the genome of the microorganism in a manner allowing the 10 microorganism to express and optionally secrete the polypeptide.

27. The vaccine according to claim 26, wherein the microorganism is a bacterium.

28. The vaccine according to claim 27, wherein the bacterium is selected from the 15 group consisting of the genera *Mycobacterium*, *Salmonella*, *Pseudomonas* and *Escherichia*.

29. The vaccine according to any one of claims 26 to 28, wherein the microorganism is *Mycobacterium bovis* BCG.

20 30. The vaccine according to any one of claims 26 to 29, wherein the microorganism is *Mycobacterium bovis* BCG strain: Danish 1331.

31. The vaccine according to any of claims 26 to 30, wherein at least 2 copies of a DNA fragment encoding a polypeptide chosen from the group consisting of:

25 - the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and

- Rv3386

are incorporated into the genome of the microorganism.

30 32. The vaccine according to claim 31, wherein the number of copies is at least 5.

33. A composition for diagnosing tuberculosis in an animal, including a human being, comprising a polypeptide chosen from the group consisting of:

35 - the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and

- Rv3386

optionally in combination with a means for detection.

34. A nucleic acid fragment which

- 1) comprises a nucleic acid sequence which has a sequence as disclosed in SEQ ID NO: 6,
- 5 2) has a length of at least 10 nucleotides and hybridizes under moderately stringent conditions with a nucleic acid fragment which has a sequence as disclosed in SEQ ID NO: 6 or a sequence complementary thereto,

with the proviso that SEQ ID NO: 7 disclosed in WO 98/53075 is excluded, when used in a pharmaceutical.

10

35. The nucleic acid fragment according to claim 34, which is a DNA fragment, when used in a pharmaceutical

36. Use of the nucleic acid fragment according to either claim 34 or claim 35 in the preparation of a pharmaceutical composition for the diagnosis of or vaccination against 15 tuberculosis caused by *Mycobacterium tuberculosis*, *Mycobacterium africanum* or *Mycobacterium bovis*.

37. A vaccine comprising the nucleic acid fragment according to either claim 34 or 20 claim 35, the vaccine effecting *in vivo* expression of an antigen by an animal, including a human being, to whom the vaccine has been administered, the amount of expressed antigen being effective to confer substantially increased resistance to infections with mycobacteria of the tuberculosis complex in an animal, including a human being.

25 38. A replicable expression vector which comprises the nucleic acid fragment according to either claim 34 or claim 35.

39. The vector according to claim 38, which is selected from the group consisting of a virus, a bacteriophage, a plasmid, a cosmid, and a microchromosome.

30

40. A transformed cell harbouring at least one vector according to either claim 38 or claim 39.

41. The transformed cell according to claim 40, which is a bacterium belonging to the 35 tuberculosis complex.

42. The transformed cell according to either claim 40 or claim 41, which is a *M. tuberculosis bovis* BCG cell.

43. The transformed cell according to any one of claims 40 to 42, which is a *M. tuberculosis bovis* BCG strain: Danish 1331 cell.

44. The transformed cell according to any one of claims 40 to 43, which expresses a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 10 12; and
- Rv3386.

45. A composition for diagnosing tuberculosis in an animal, including a human being, comprising a nucleic acid fragment according to either claim 34 or claim 35, optionally in 15 combination with a means for detection.

46. A method for producing a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- 20 - Rv3386,

comprising inserting a nucleic acid fragment, which

- 1) comprises a nucleic acid sequence which has a sequence as disclosed in SEQ ID NO: 6,
- 2) has a length of at least 10 nucleotides and hybridizes under moderately stringent conditions with a nucleic acid fragment which has a sequence as disclosed in SEQ ID NO: 6 or a sequence complementary thereto, 25 into a vector which is able to replicate in a host cell, introducing the resulting recombinant vector into the host cell, culturing the host cell in a culture medium under conditions sufficient to effect expression of the polypeptide, and recovering the 30 polypeptide from the host cell or culture medium; or

isolating the polypeptide from whole mycobacteria of the tuberculosis complex or from a lysate or fraction thereof; or  
synthesizing the polypeptide by solid or liquid phase peptide synthesis.

35 47. The method according to claim 46, wherein the fraction is a cell-wall containing fraction.

48. A method for producing an immunologic composition according to any one of claims 14 to 25 comprising:

preparing, synthesizing or isolating a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- Rv3386, and

5 solubilizing or dispersing the polypeptide in a medium for a vaccine, and optionally adding other *M. tuberculosis* antigens and/or a carrier, vehicle and/or adjuvant substance, or

10 cultivating a cell according to any one of claims 40 to 44, and transferring the cell to a medium for a vaccine, and optionally adding a carrier, vehicle and/or adjuvant substance.

49. A method of diagnosing tuberculosis caused by *Mycobacterium tuberculosis*,

15 *Mycobacterium africanum* or *Mycobacterium bovis* in an animal, including a human being, comprising intradermally injecting, into the animal, a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- 20 - Rv3386

or an immunologic composition according to any one of claims 14 to 25, a positive skin response at the location of injection being indicative of the animal having tuberculosis, and a negative skin response at the location of injection being indicative of the animal not having tuberculosis.

25

50. A method for immunising an animal, including a human being, against tuberculosis caused by mycobacteria belonging to the tuberculosis complex, comprising administering to the animal a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and
- 30 - Rv3386,

the immunologic composition according to any one of claims 14 to 25, or the vaccine according to any any one of claims 26 to 32 and 37.

35 51. The method according to claim 50, wherein the polypeptide, immunologic composition, or vaccine is administered by a parenteral, intraperitoneal, intramuscular, subcutaneous, intradermal, oral, buccal, sublingual, nasal, rectal or transdermal route.

52. The method according to claim 51, wherein the parenteral route is intravenous or intra-arterial.

5 53. A method for diagnosing ongoing or previous sensitization in an animal or a human being with bacteria belonging to the tuberculosis complex, the method comprising providing a blood sample from the animal or human being, and contacting the sample from the animal with a polypeptide chosen from the group consisting of:

10 - the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and

- Rv3386,

a significant release into the extracellular phase of at least one cytokine by mononuclear cells in the blood sample being indicative of the animal being sensitized.

15 54. A monoclonal or polyclonal antibody, which specifically reacts with a polypeptide chosen from the group consisting of:

- the polypeptide fragment according to any one of the claims 5 to 9, 11 and 12; and

- Rv3386

20 in an immunoassay, or a specific binding fragment of said antibody.

55. A fusion polypeptide according to claim 1, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

25 56. A substantially pure polypeptide fragment according to claim 5, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

30 57. Use of a substantially pure polypeptide according to claim 13, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

35 58. An immunologic composition according to claim 14, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

59. A vaccine according to either claim 26 or claim 37, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

5 60. A composition for diagnosing tuberculosis according to either claim 33 or claim 45, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

61. A nucleic acid fragment according to claim 34, substantially as herein described  
10 with reference to any one or more of the Examples and/or accompanying Figures.

62. Use according to claim 36, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

15 63. A replicable expression vector according to claim 38, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

64. A transformed cell according to claim 40, substantially as herein described with  
20 reference to any one or more of the Examples and/or accompanying Figures.

65. A method according to any one of claims 46, 48, 49, 50 or 53, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

25 66. A monoclonal or polyclonal antibody according to claim 54, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

Dated this eighth day of November 2004

Statens Serum Institut  
Patent Attorneys for the Applicant:  
F B RICE & CO




Fig. 1 a




Fig. 1 b




Fig. 1 c




Fig. 2 a

5/8




Fig. 2 b




Fig. 2 c

11-03-06 50670

## Healthy BCG vaccinated




Figure 3

11. 2.02 59670

TB patients




Figure 4