



(86) Date de dépôt PCT/PCT Filing Date: 2001/11/01
(87) Date publication PCT/PCT Publication Date: 2003/05/08
(85) Entrée phase nationale/National Entry: 2004/04/29
(86) N° demande PCT/PCT Application No.: KR 2001/001845
(87) N° publication PCT/PCT Publication No.: 2003/037085

(51) Cl.Int.⁷/Int.Cl.⁷ A01N 43/38


(71) Demandeur/Applicant:
DONGBU HANNONG CHEMICAL CO., LTD., KR

(72) Inventeurs/Inventors:
KIM, DAE WHANG, KR;
CHANG, HAE SUNG, KR;
KO, YOUNG KWAN, KR;
RYU, JAE WOON, KR;
WOO, JAE CHUN, KR;
KOO, DONG WAN, KR;
KIM, JIN SEOG, KR;
CHUNG, BONG-JIN, KR;
KWON, OH-YEON, KR

(74) Agent: OGILVY RENAULT

(54) Titre : AMIDES N-METHYL-N-2-FLUOROPHENYL D'ACIDE PHENOXYPROPIONIQUE (R) HERBICIDES ACTIFS OPTIQUEMENT

(54) Title: OPTICALLY ACTIVE HERBICIDAL (R)-PHENOXYPROPIONIC ACID-N-METHYL-N-2-FLUOROPHENYL AMIDES



(1)

(57) Abrégé/Abstract:

The present invention relates to optically active herbicidal (R)-phenoxypropionic acid N-methyl-N-2-fluorophenyl amide compounds represented in the following formula (1), a method for preparing thereof, their use to prevent generation of barnyard grass produced from rice and composition as suitable herbicides, (I) wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl; is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 May 2003 (08.05.2003)

PCT

(10) International Publication Number
WO 03/037085 A1

(51) International Patent Classification⁷: **A01N 43/38**

(21) International Application Number: PCT/KR01/01845

(22) International Filing Date:

1 November 2001 (01.11.2001)

(25) Filing Language:

Korean

(26) Publication Language:

English

(71) Applicant (*for all designated States except US*):
DONGBU HANNONG CHEMICAL CO., LTD.
[KR/KR]; 838 Yuksam-dong, Kangnam-ku, Seoul
135-080 (KR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KIM, Dae Whang**
[KR/KR]; 132-304 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **CHANG, Hae
Sung** [KR/KR]; 105-201 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **KO, Young Kwan**
[KR/KR]; 102-1702 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **RYU, Jae Wook**
[KR/KR]; 120-305 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **WOO, Jae Chun**
[KR/KR]; 1-406 Kyungnam Apt., 205 Doma-2dong,
Seo-ku, Daejeon 302-162 (KR). **KOO, Dong Wan**
[KR/KR]; 128-604 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **KIM, Jin Seog**
[KR/KR]; 128-208 Hanbit Apt., 99 Uheun-dong,
Yusung-ku, Daejeon 305-333 (KR). **CHUNG, Bong-Jin**

[KR/KR]; 301, 1092-8 Galsan-dong, Dongan-ku,
Kyunggi-do, Anyang 431-800 (KR). **KWON, Oh-Yeon**
[KR/KR]; 905 ByuksanGrandCore Apt., 9-2 Jungja-dong,
Jangan-ku, Kyunggi-do, Suwon 440-830 (KR).

(74) Agent: **HUH, Sang Hoon**; 16th Fl. Woori Invest-
ment Bldg., 826-20 Yeoksam-dong, Kangnam-ku, Seoul
135-180 (KR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— with international search report


For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: OPTICALLY ACTIVE HERBICIDAL (R)-PHENOXYPROPIONIC ACID-N-METHYL-N-2-FLUOROPHENYL AMIDES



(1)

(57) Abstract: The present invention relates to optically active herbicidal (R)-phenoxypropionic acid N-methyl-N-2-fluorophenyl amide compounds represented in the following formula (1), a method for preparing thereof, their use to prevent generation of barnyard grass produced from rice and composition as suitable herbicides, (I) wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxy carbonyl, or C₁-C₃ alkyl carbonyl; is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.




WO 03/037085 A1

OPTICALLY ACTIVE HERBICIDAL (R)-PHENOXYPROPIONIC ACID-N-METHYL-N-2-FLUOROPHENYL AMIDES

【Technical Field】

The present invention relates to optically active herbicidal (R)-phenoxypropionic acid N-methyl-N-2-fluorophenyl amide compounds represented in the following formula (1), a method for preparing thereof, their use to prevent generation of barnyard grass produced from rice and composition as suitable herbicides,



(1)

wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl;


Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other


substituents when n is 2.

【Background Art】

U.S. Pat. No. 4,130,413 discloses the compounds represented in the following formula (2),





wherein $(R_1)_m$ is hydrogen, halogen, CF_3 , NO_2 , CN or alkyl; A is O, S or NH; R_2

is hydrogen or alkyl; Z is , where R_3 and R_4 may be identical or different and represent hydrogen, C_1 - C_6 alkyl, C_1 - C_6 hydroxyalkyl, C_3 - C_6 cycloalkyl, C_1 - C_4 alkoxy, or phenyl substituted with 1 to 3 substituents chosen from C_1 - C_4 alkyl, C_1 - C_6 alkoxy, halogen and CF_3 .

10

U.S. Pat. No. 4,531,969 discloses the compounds represented in the following formula (3),




wherein R_5 is , where R_6 is hydrogen or halogen; R_7 is


hydrogen or alkyl ; and Z is $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{N} \begin{array}{l} \nearrow \text{R}_3 \\ \searrow \text{R}_4 \end{array} \end{array}$, where R₃ and R₄ may be identical or different and represent hydrogen, C₁-C₆ alkyl, C₁-C₆ hydroxyalkyl, C₃-C₆ cycloalkyl, C₁-C₄ alkoxy, or phenyl substituted with 1 to 3 substituents chosen from C₁-C₄ alkyl, C₁-C₆ alkoxy, halogen and CF₃.

5

U.S. Pat. No. 5,254,527 discloses the compounds represented in the following formula (4),



10

wherein R₅ is , Z is $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{N} \begin{array}{l} \nearrow \text{R}_3 \\ \searrow \text{R}_4 \end{array} \end{array}$, where R₃ and R₄ may be identical or different and represent hydrogen, C₁-C₆ alkyl, C₁-C₆ hydroxyalkyl, C₃-C₆ cycloalkyl, C₁-C₄ alkoxy, or phenyl substituted with 1 to 3 substituents chosen from C₁-C₄ alkyl, C₁-C₆ alkoxy, halogen and CF₃.


15

Even though some compounds of formula (1) of the present invention are disclosed in the above patents, none of the patents teach the synthesis of the compound of formula (1) and have tested the same for herbicidal activity.

20

JP Patent publication 2-11580 discloses the compound represented in the


following formula (5),



5 wherein L is low alkyl, halogen, methoxy, methoxyphenoxy, benzyloxy, methylthio or methylvinyl; and n is an integer of 0 to 2.

JP Patent publication sho 53-40767 and sho 54-112828 also disclose that phenoxypropionic acid amide derivatives have herbicidal activity.

Further, inventors of the present invention disclosed herbicidal
 10 phenoxypropionic acid N-alkyl-N-2-fluorophenyl amide compounds in International Pat. Publication No. WO 2000/05956.



15 Even though many of herbicides for rice have been recently developed and used, barnyard grass among weeds is the biggest problem in rice paddy.

Development of herbicides to control barnyard grass is an urgent to one who is in the field of agriculture. After transplanting young rice, herbicides, developed until now, cannot effectively control the production of barnyard
 20 grass so that it causes a huge damage to harvest. It has been reported that the

amount of rice harvest is decreased by 2% when barnyard grass is produced 1 week per 1 m², decreased by about 10% when produced 5 weeks per 1 m², decreased by about 19% when produced 10 weeks per 1 m² and decreased by about 35% when produced 20 weeks per 1 m².

5 Many different kinds of herbicides have been used for the purpose of controlling barnyard grass that damages in amount of harvest of rice. However, the herbicide with a broader herbicidal activity, environment-friendly property and cost-effectiveness is still in demand.


10 The inventors have intensively studied to provide herbicides to effectively control barnyard grass, and particularly, to find out selective herbicidal activity of phenoxypropionic acid *N*-alkyl-*N*-2-fluorophenyl amide compounds of formula (6). As a result, we completed this invention by finding that some phenoxypropionic acid *N*-alkyl-*N*-2-fluorophenyl amides of formula (6) exist as (R)- or (S)-stereoisomer, and (R)-stereoisomers provide higher
15 stability to rice and better herbicidal activity compared to (S)-stereoisomers or mixtures thereof. This superior activity of (R)-stereoisomers is distinguished from the conventional inventions.

Therefore, an object of the present invention is to provide optically active herbicide compounds which exhibit excellent selectivity toward rice and
20 prevent the production of harmful barnyard grass.

【Disclosure of Invention】

The present invention is to provide optically active herbicide phenoxypropionic acid *N*-methyl-*N*-2-fluorophenyl amides of formula (1) with

an excellent herbicidal activity as well as selective and remarkable stability toward rice,



5


wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl; Y is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

10

The optically active compounds of formula (1) according to the present invention may be specified as the following Table 1.

15

Table 1

 (1)		
R	Y	X
CH ₃	H	H
CH ₂ CH ₃	H	H
CH ₃	H	4-F
CH ₃	H	3-F
CH ₃	H	5-F
CH ₃	H	4-Cl
CH ₃	H	4-F, 5-F
CH ₃	H	4-Br
CH ₃	H	4-CH ₃
CH ₃	H	4-SCH ₃
CH ₃	H	4-CH ₂ CH ₃
CH ₃	H	4-propyl
CH ₃	H	4-isopropyl
CH ₃	H	4-butyl
CH ₃	H	4-isobutyl

R	Y	X
CH ₃	H	4-CO ₂ CH ₃
CH ₃	H	4-OCH ₃
CH ₃	H	4-OEt
CH ₃	H	4-O-isopropyl
CH ₃	H	4-O-allyl
CH ₃	H	4-O-propyl
CH ₃	F	H
CH ₃	F	3-F
CH ₃	F	4-F
CH ₂ CH ₃	F	4-F
CH ₃	F	4-Cl
CH ₃	F	4-Br
CH ₃	F	4-CH ₃
CH ₃	F	4-CH ₂ CH ₃
CH ₃	F	4-propyl
CH ₃	F	4-isopropyl
CH ₃	F	4-cyclopropyl
CH ₃	F	4-butyl
CH ₃	F	4-isobutyl


R	Y	X
CH ₃	F	4-OCH ₃
CH ₃	F	4-OEt
CH ₃	F	4-O-isopropyl
CH ₃	F	4-O-propyl
CH ₃	F	3-F, 5-F
CH ₃	H	5-F
CH ₃	H	5-Cl
CH ₃	H	5-Br
CH ₃	H	5-CH ₃
CH ₃	H	5-SCH ₃
CH ₃	H	5-CH ₂ CH ₃
CH ₃	H	5-propyl
CH ₃	H	5-isopropyl
CH ₃	H	5-cyclopropyl
CH ₃	H	5- butyl
CH ₃	H	5-isobutyl
CH ₃	H	5-OCH ₃
CH ₃	H	4-OEt

R	Y	X
CH ₃	H	5-O-isopropyl
CH ₃	H	5-O-propyl
CH ₃	H	5-O-allyl
CH ₃	F	5-H
CH ₃	F	5-F
CH ₃	F	5-Cl
CH ₃	F	5-Br
CH ₃	F	5-CH ₃
CH ₃	F	5-CH ₂ CH ₃
CH ₃	F	5-propyl
CH ₃	F	5-isopropyl
CH ₃	F	5-cyclopropyl
CH ₃	F	5-n-butyl
CH ₃	F	5-isobutyl
CH ₃	F	5-OCH ₃
CH ₃	F	5-OEt
CH ₃	F	5-O-isopropyl
CH ₃	F	5-O-propyl

The optically active compounds of formula (1) according to this invention may be synthesized by employing a conventional method
 5 represented in the following Scheme 1, reacting a compound of formula (7)

with a compound of formula (8),

Scheme 1




- 5 wherein X' is OH, Cl, Br, or phenoxy; X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl; Y is hydrogen
 10 or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

In the method according to Scheme 1, it is performed preferably to use a binder such as triphenylphosphine and an organic base such as triethylamine or
 15 pyridine at a temperature of 0 to 100°C in an inert solvent such as ethers like tetrahydrofuran, ethyethyl acetate, acetonitrile, toluene, xylene, hexane, methylene chloride, carbon tetrachloride, dichloroethane or the like. After the solvent is evaporated, the crude product is purified by column chromatography.

Another method for preparing the compounds (1) represented in the
 20 following Scheme 2 is an alkylation of a compound of formula (9) to a compound of formula (10),

Scheme 2




wherein X'' is Cl, Br, I, benzenesulfonyloxy, toluenesulfonyloxy, methanesulfonyloxy or low alkyl sulfate; and X is hydrogen, halogen, hydroxy, NH_2 , CO_2H , C_1 - C_6 alkylamino substituted with 1 or 2 of C_1 - C_3 alkyl, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_3 haloalkyl, C_1 - C_3 haloalkoxy, C_2 - C_4 alkoxyalkoxy, C_1 - C_4 alkylthonyl, C_1 - C_4 alkylsulfonyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_2 - C_6 alkenyloxy, C_2 - C_6 alkynyloxy, C_1 - C_3 alkoxy carbonyl, or C_1 - C_3 alkyl carbonyl; Y is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

In scheme 2, it is performed preferably using a strong base which is enough to pull out a hydrogen from amide, NH. Examples of a strong base include NaOH, KOH, LiOH, NaH, n-BuLi, LDA, and the like. The reaction is performed at a temperature of -78 to $50^\circ C$ in an inert solvent such as ethers like ethylether, dioxane or tetrahydrofuran or hydrocarbons like hexane.

Another method for preparing the compounds (1) represented in the following Scheme 3 is a reaction of a compound of formula (11) with a compound of formula (12) in the presence of a base,

Scheme 3



wherein Y' is halogen, alkylsulfonyloxy, haloalkylsulfonyloxy, benzenesulfonyloxy or toluenesulfonyloxy; X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl; Y is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.


In Scheme 3, examples of the base include inorganic bases of alkali metal hydroxides such as sodium hydroxide and potassium hydroxide, alkali metal carbonates such as sodium carbonate and potassium carbonate, alkali metal hydrogen carbonates such as sodium hydrogencarbonate and potassium hydrogencarbonate; and organic bases such as triethylamine, *N,N*-dimethylaniline, pyridine and 1,8-diazabicyclo[5,4,0]undec-7-ene.

A phase transition catalyst such as tetra-*n*-butylammonium bromide or 18-crown-6-[1,4,7,10,13,16-hexaoctacyclooctadecane] may be added to rapidly complete the reaction, if necessary. Further, one or more than two solvents may be used, if deemed necessary. Examples of the inert organic solvent include ketones such as acetone; aromatic hydrocarbons such as toluene, xylene and

chlorobenzene; aliphatic hydrocarbons such as petroleum ether and ligroin; ethers such as diethylether, tetrahydrofuran and dioxane; nitrites such as acetonitrile and propionitrile; and amides such as *N,N*-dimethylformamide, *N,N*-dimethylacetamide, and *N*-methylpyrrolidone. A reaction is carried out at
 5 a temperature of from 0°C to reflux, preferably at 5 to 50°C, for 1 to 24 hour(s) to afford the desired product with high yield.

Another method for preparing the optically active compound (1) represented in the following Scheme 4 is a reaction of a compound of formula (13) with a compound of formula (14) in the presence of a base,

10



wherein Y' is halogen, alkylsulfonyloxy, haloalkylsulfonyloxy, benzenesulfonyloxy or toluenesulfonyloxy; X is hydrogen, halogen, hydroxy, NH₂, CO₂H, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthonyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl; Y is hydrogen or fluoro; and n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

20

In Scheme 4, examples of the base include inorganic bases of alkali

metal hydroxides such as sodium hydroxide and potassium hydroxide, alkali metal carbonates such as sodium carbonate and potassium carbonate, alkali metal hydrogencarbonates such as sodium hydrogencarbonate and potassium hydrogencarbonate; and organic bases such as triethylamine, *N,N*-dimethylaniline, pyridine, picoline, quinoline, and 1,8-diazabicyclo[5,4,0]undec-7-ene.

A phase transition catalyst such as tetra-*n*-butylammonium bromide or 18-crown-6[1,4,7,10,13,16-hexaoctacyclooctadecane] may be used, if necessary. Further, more than one solvent may be used if deemed necessary. Examples of the inert organic solvent include ketones such as acetone and butanone; aromatic hydrocarbons such as benzene, toluene, xylene and chlorobenzene; aliphatic hydrocarbons such as petroleum ether, and ligroin; ethers such as diethylether, tetrahydrofuran and dioxane; nitriles such as acetonitrile or propionitrile; and amides such as *N,N*-dimethylformamide, *N,N*-dimethylacetamide and *N*-methylpyrrolidone. A reaction is carried at a temperature of from 0°C to reflux, preferably at 20 to 100°C for 1 to 24 hour(s) to afford the desired product with high yield.

The present invention will be further illustrated by the following examples. However, they should not be construed as limiting the scope of this invention defined by the appended claims.

【Examples】**Example 1: Preparation of (S)-2-bromo-propionic acid-N-(2-fluorophenyl)-N-methyl amide**

(S)-2-Bromopropionic acid(3.4 g, 0.022 mol) and 2-fluoroaniline(3 g, 0.024 mol) were dissolved in 50 ml of chloroform and cooled to 0°C. Dicyclohexylcarbodiimide(5 g, 0.024 mol) dissolved in 10 ml of chloroform was slowly injected through a syringe. The temperature of the reaction mixture was raised to room temperature and the reaction mixture was stirred for 1 hour. Solid remained during the reaction was filtered out and washed twice with 20 ml of chloroform. The filtrate was concentrated under reduced pressure and the crude product was purified by column chromatography (eluent; ethyl acetate/*n*-hexane=1/3) to afford 5 g of the target product.

¹H-NMR(CDCl₃) : δ1.7(3H, d), 3.24(3H, s), 4.16(0.7H, q), 4.34(0.3H, q), 7.13-7.48(4H, m)

Example 2: Preparation of (R)-2-(4-hydroxyphenoxy)propionic acid-N-(2-fluorophenyl)-N-methyl amide

(S)-2-bromo-propionic acid-N-(2-fluorophenyl)-N-methyl amide (18.2 g, 0.07 mol), hydroquinone (7 g, 0.064 mol), potassium carbonate (10.54 g, 0.076 mol) and tetra-*n*-butylammonium bromide (1 g) were dissolved in 350 ml of acetonitrile and heated at reflux for 6 hours. The reaction mixture was cooled to room temperature and solid remained during the reaction was filtered out. The filtrate was concentrated under reduced pressure and the crude product was

purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 16 g of the target product.

¹H-NMR(CDCl₃) : δ1.42(3H, t), 3.25(3H, s), 4.56(1H, q), 6.5-7.4(8H, m)

5

Example 3: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-(2-fluorophenyl)-*N*-methyl amide

(R)-2-(4-hydroxyphenoxy)propionic acid-*N*-(2-fluorophenyl)-*N*-methyl amide (11.5 g, 0.04 mol), 2,6-dichlorobenzoxazole (6.85 g, 0.036 mol), potassium carbonate (6 g, 0.043 mol) and tetra-*n*-butylammonium bromide (1 g) were dissolved in 300 ml of acetonitrile and heated at reflux for 7 hours. The reaction mixture was cooled to room temperature and solid remained during the reaction was filtered out. The filtrate was concentrated under reduced pressure and the crude product was purified by column chromatography (eluent: ethyl acetate/*n*-hexane=1/3) to afford 12.5 g of the target product.

¹H-NMR(CDCl₃) : δ1.42(3H, t), 3.3(3H, s), 4.62(1H, m), 6.8-7.4(11H, m)

Example 4: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-(2-fluorophenyl)-*N*-methyl amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid (346.7 mg, 1 mmol) was dissolved in 10 ml of tetrahydrofuran. 2-Fluoroaniline(111.12 mg, 1 mmol), triphenylphosphine(393.4 mg, 1.5 mmol), triethylamine(0.15 ml, 1 mmol) and carbon tetrachloride(1 ml) were added sequentially and heated at

reflux for 8 hours. The reaction mixture was cooled to room temperature and acidified with 5% hydrochloric acid, followed by addition of water. The acidified reaction mixture was extracted three times with ethyl acetate. The combined organic layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/4) to afford 200 mg of the target product.

m.p : 132-136 °C

¹H-NMR(CDCl₃) : δ1.7(3H, d), 4.81(1H, q), 7.05-7.45(10H, m), 8.35(1H, m), 8.5(1H, br)

Example 5: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-(2-fluorophenyl)-*N*-methyl Amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid-*N*-(2-fluorophenyl)amide (100 mg, 0.24 mmol) was dissolved in 10 ml of anhydrous tetrahydrofuran and 60% NaH(10 mg, 0.24 mmol) and CH₃I(34 mg, 0.24 mmol) were added sequentially at 0 °C. The reaction mixture was stirred at room temperature for 5 hours. Ice water was poured to the reaction mixture and it was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 75 mg of the target product.

$^1\text{H-NMR}(\text{CDCl}_3)$: δ 1.42(3H, t), 3.3(3H, s), 4.62(1H, m), 6.8-7.4(11H, m)

Example 6: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-(2-fluorophenyl)-*N*-methyl amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid(346.7 mg, 1 mmol) was dissolved in 10 ml of tetrahydrofuran and *N*-methyl-2-fluoroaniline(125 mg, 1 mmol), triphenylphosphine(393.4 mg, 1.5 mmol), triethylamine(0.15 ml, 1 mmol) and carbon tetrachloride(1 ml) were added sequentially and the reaction was heated at reflux for 12 hours. The reaction mixture was cooled to room temperature and acidified with 5% hydrochloric acid, followed by addition of water. The acidified reaction mixture was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 100 mg of the target product.

Example 7: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-methyl-*N*-(2,4,5-trifluorophenyl)amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid (0.693 g, 2 mmol) was dissolved in 15 ml of tetrahydrofuran and *N*-methyl-2,4,5-trifluoroaniline(0.322 g, 2 mmol), triphenylphosphine(0.78g, 2 mmol), triethylamine(0.4 ml) and carbon tetrachloride(2 ml) were added sequentially and then the reaction mixture was heated at reflux for 18 hours. The reaction

mixture was cooled to room temperature and acidified with 5% hydrochloric acid. The acidified reaction mixture was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by
5 column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 250 mg of the target product.

¹H-NMR(CDCl₃) : δ1.42(3H, d), 3.2(3H, s), 4.65(1H, m), 6.6-7.4(9H, m)

10 **Example 8: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-*N*-methyl-*N*-(2,6-difluoro-phenyl)amide**

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid(0.693 g, 2 mmol) and *N*-methyl-2,6-difluoroaniline(0.284 g, 2 mmol) were dissolved in 20 ml of tetrahydrofuran and triphenylphosphine(0.78 g, 2 mmol),
15 triethylamine(0.42 ml) and carbon tetrachloride(2 ml) were added sequentially. The reaction mixture was heated at reflux for 16 hours. The reaction mixture was cooled to room temperature and acidified with 5% hydrochloric acid. The acidified reaction mixture was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and
20 concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 205 mg of the target product.

¹H-NMR(CDCl₃) : δ1.4(3H, d), 3.3(3H, s), 4.62(1H, q), 6.8-7.4(10H, m)

Example 9: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-N-(2,4-difluorophenyl)-N-methyl amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid(0.693 g, 2 mmol) was dissolved in 15 ml of tetrahydrofuran and N-methyl-2,4-difluoroaniline(0.284 g, 2 mmol), triphenylphosphine(0.78 g, 2 mmol), triethylamine(0.42 ml) and carbon tetrachloride(2 ml) were added sequentially. The reaction mixture was heated at reflux for 12 hours. The reaction mixture was cooled to room temperature and acidified with 5% hydrochloric acid, followed by addition of water. The acidified reaction mixture was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 230 mg of the target product.

¹H-NMR(CDCl₃) : δ1.4(3H, d), 3.2(3H, s), 4.6(1H, q), 6.6-7.2(10H, m)

Example 10: Preparation of (R)-2-[4-(6-chloro-2-benzoxazolyloxy)-phenoxy]propionic acid-N-methyl-N-(2,3,6-trifluorophenyl)amide

(R)-2-[4-(6-chloro-2-benzoxazoyloxy)-phenoxy]propionic acid(0.693g, 2 mmol) was added to 6 ml of thionyl chloride and the reaction mixture was heated at reflux for 2 hours. Excess of thionyl chloride was removed under reduced pressure and 3 ml of anhydrous tetrahydrofuran was added to it. A solution of N-methyl-2,3,6-trifluoroaniline(0.32 g, 2 mmol) and triethyl amine(0.42 ml) in anhydrous tetrahydrofuran(10 ml) was added slowly to the


reaction mixture at 0°C. The mixture was stirred at 0°C for 30 minutes and stirred at room temperature for additional 1 hour. After pouring water the reaction mixture was extracted three times with ethyl acetate. The combined organic solvent layer was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography(eluent: ethyl acetate/*n*-hexane=1/2) to afford 240 mg of the target product.

¹H-NMR(CDCl₃) : δ1.45(3H, d), 3.25(3H, s), 4.6(1H, q), 6.7–7.4(9H, m)

Examples 11-16

The compounds represented in the following Table 2 were prepared by the same procedure of example 10 except using of aniline compounds instead of *N*-methyl-2,3,6-trifluoroaniline.

Table 2

			
Classification	X ₁	X ₂	¹ H-NMR(CDCl ₃)
Example 11	H	CH ₃	1.42(3H, t), 2.3(3H, s), 3.25(3H, s), 4.62 (1H, m), 6.8-7.4(10H, m)
Example 12	Cl	H	1.42(3H, t), 3.3(3H, s), 4.62(1H, m), 6.7-7.5(10H, m)
Example 13	H	F	1.42(3H, t), 3.3(3H, s), 4.62(1H, m), 6.5-7.4(10H, m)
Example 14	CH ₃	H	1.42(3H, t), 2.38(3H, s), 3.25(3H, s), 4.62(1H, m), 6.8-7.4(10H, m)
Example 15	OCH ₃	H	1.42(3H,t), 3.25(3H, s), 3.8(3H, s), 4.65(1H, m), 6.65-7.45(10H, m)
Example 16	OCH ₂ CH ₃	H	1.25(3H, t), 1.42(3H, t), 3.25(3H, s), 4.0(2H, q), 4.62(1H, m), 6.65-7.42(10H. m)

Formulation

In order to use the compounds according to the present invention as herbicides, they should be formulated in such a suitable type such as wettable powder, emulsions, granules, dusts, suspensions and solutions by combining a carrier, a surfactant, a dispersing agent or a supplement agent. Many of these may be applied directly or after diluted with suitable media. Formulations can be prepared at spray volume of from hundreds liters to thousands liters per hectare. The formulations contain about 0.1% to 99% by weight of active ingredient(s) and 0.1% to 20% surfactant(s) or 0% to 99.9% solid or liquid diluent(s) are recommended to be added. The formulations will contain these

ingredients in the following approximate proportions shown in Table 3.

Table 3

Formulations	Wt. %		
	Active ingredient	Diluent	Surfactant
Wettable powder	10-90	0-74	1-10
Suspension	3-50	40-95	0-15
Emulsion . solution	3-50	40-95	0-15
Granule	0.1-95	5-99.9	1-15

The proportion of active ingredients depends on the intended use.

- 5 Higher ratio of a surfactant is sometimes desired to active ingredients and is achieved by incorporation into the formulation or tank mixing.

Solid diluents with high absorption are preferred for wettable powder.

- Liquid diluents and solvents are preferred to be stable against phase separation at 0°C. All the formulations may contain a small amount of additives to prevent
10 forming, caking, corrosion and growth of microorganisms.

- According to conventional methods to prepare the composition, solutions can be made only by blending ingredients and fine solids by blending and pulverizing with hammer-mill. Suspensions can be made by wet-milling and granules can be made by spraying the active ingredients on performed
15 granular carrier.

Preparation examples of typical formulations are as follows.

Formulation 1: Wettable Powder

The ingredients are thoroughly blended, re-blended after spraying liquid surfactant on the solid ingredients and hammer-milled until all the solids are essentially under 100 μm .

5	Active ingredient (Compound of Example 3)	20 wt. %
	Dodecylphenol polyethylene glycol ether	2 wt. %
	Sodium ligninsulfonate	4 wt. %
	Sodium silicon aluminate	6 wt. %
	Montmorillonite	68 wt. %

10

Formulation 2: Wettable Powder

The ingredients are blended, hammer-milled until all the solids are under 25 μm and packaged.

	Active ingredient (Compound of Example 3)	80 wt. %
15	Sodium alkyl naphthalenesulfonate	2 wt. %
	Sodium ligninsulfonate	2 wt. %
	Synthetic amorphous silica	3 wt. %
	Kaolinite	13 wt. %

20 Formulation 3: Emulsion

The ingredients are mixed and homogeneously dissolved to give emulsions.

	Active ingredient (Compound of Example 3)	30 wt. %
	Cyclohexanone	20 wt. %

Polyoxyethylene alkylaryl ether	11 wt. %
Calcium alkylbenzenesulfonate	4 wt. %
Methylnaphthalene	35 wt. %

5 Formulation 4: Granule

The ingredients were thoroughly blended. 20 parts by weight of water was added to 100 parts of weight the ingredient mixture. The ingredient mixture was granulated with a size of 14 to 32 mesh by using extrusive granulator and dried.

10	Active ingredient (Compound of Example 3)	5 wt. %
	Sodium laurylalkoholsulfonate	2 wt. %
	Sodium ligninsulfonate	5 wt. %
	Carboxymethyl cellulose	2 wt. %
	Potassium sulfate	16 wt. %
15	Plaster	70 wt. %

The formulations according to this invention were sprayed with diluting to a certain concentration.

20 Utility

The compounds according to the present invention represent high activity as leaf treatment herbicides for rice and especially effective in rice due to an excellent control of barnyard grass.

The active ingredients can be used from 10 g to 4 kg per hectare, preferably from 50 g to 400 g. The amount of the compounds of the present invention depends on the amount and size of weeds and formulations. The herbicides of the present invention can be used as alone or in combination with
5 other herbicides, insecticides or bactericides. Especially it is essential to add one agent selected from the group consisting of bentazon, quinclorac, propanil, simetryn, 2,4-D, fenoxaprop-ethyl, linuron, MCPA, azafenidin, carfentrazone, molinate, thiobencarb, pendimethalin, bensulfuron-methyl, pyrazosulfuron-ethyl, metsulfuron-methyl, thifensulfuron-methyl, tribenuron-methyl,
10 trifluralin, amidosulfuron, bromoxynil, butachlor, mecoprop, metribuzin, bifenox, benfuresate, isoproturon, cyhalofop-butyl, mefenaset, fentrazamide, pyriminobac-methyl, bispyribac sodium, azimsulfuron, cyclosulfamuron, pyanchor, and mixtures thereof.

The herbicidal effect of the compounds of this invention was tested and
15 the examples are as follows.

Experimental Example 1: Leaf Treatment Test

Seeds of rice, wheat, barley, corn, cotton, barnyard grass, common sorghum, large crabgrass and fall panicum were seeded at a pot with a surface
20 area of 600 cm². When barnyard grass grown in a green house kept at 20 – 30 °C had three leaves, wettable powder prepared by mixing 1 part by weight of the active compound, 5 parts by weight of acetone and 1 part by weight of emulsifier and diluted with water was applied directly to the leaves in 2000 L per hectare. The concentration of the spray liquid was so chosen the particular

amounts of the active compound desired. 14 days after the treatment, the degree of damage to the plants was rated in % damage in comparison to the development of untreated control.

	0%	no effect (same as untreated control)
5	20%	slight effect
	70%	herbicidal effect
	100%	total destruction

In the test, the active compound(s) of formula (1) according to the invention exhibited an excellent selectivity toward the plants and herbicidal activity against weeds.

Table 4


Abbr.	Scientific Name	English Name
ORYSA	Oryza sativa L. cv. Dongjin	Rice
ECHCG	Echinochloa crus-galli Beauv. var. caudata Kitagawa	Barnyard grass

Among the compounds of formula (1), herbicidal activity of (R)-2-[4-chloro-2-benzoxazoyloxy]-phenoxy]propionic acid-N-(2-fluorophenyl)-N-methyl amide (Example 3) was compared to the (S)-stereoisomer and racemic mixture thereof and the result is summarized in table 5.


Table 5

Amount of Leaf treatment (g/ha)	(R)-compound		R,S-racemic compound		(S)-compound	
	Rice (4 leaves)	Barnyard grass (4 leaves)	Rice (4 leaves)	Barnyard grass (4 leaves)	Rice (4 leaves)	Barnyard grass (4 leaves)
4000	22.5	100	3.8	100	0.0	100
2000	11.3	100	0.0	100	0.0	100
1000	2.5	100	0.0	100	0.0	100
500	0.0	100	0.0	100	0.0	100
250	0.0	100	0.0	100	0.0	100
125	0.0	100	0.0	100	0.0	100
63	0.0	100	0.0	100	0.0	100
32	0.0	100	0.0	100	0.0	92.5
16	0.0	100	0.0	100	0.0	65.0
8	0.0	98.8	0.0	45.0	0.0	7.5
4	0.0	62.5	0.0	2.5	0.0	0.0
2	0.0	6.3	0.0	0.0	0.0	0.0
1	-	-	-	-	-	-


(R)-Compound



R,S-Racemic compound



(S)-Compound



5

【Industrial Applicability】

As described above, it is noted that optically active (R)-stereoisomers of
 10 the present invention exhibit excellent selectivity toward rice and superior
 herbicidal activity against barnyard grass to racemic mixtures and (S)-
 stereoisomers thereof. Therefore, the optically active compounds of the present
 invention may be very effective in rice farming. Further, it is proved that the
 optically active compounds are very stable for wheat, barley, beans, and corn
 15 and useful to control weeds.

20


PCT/ KR 2001/001845

03 FEBRUARY 2004

CLAIMS

What is claimed is:

1. A herbicidal compound, (R)-phenoxypropionic acid-N-methyl-N-2-fluorophenyl amide of formula (1), having stability toward rice and preventing the generation of barnyard grass:



wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, CN, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthionyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

2. The herbicidal compound according to claim 1, wherein said X is H, F, Cl, Br, CN, CH₃, or OCH₃; Y is H or F; and n=1.

PCT/ KR 2001/001845


03 FEBRUARY 2004

3. The herbicidal compound according to claim 1, wherein said X is H; and Y is H.

4. The herbicidal compound according to claim 1, wherein said X is 5-CH₃; and Y is H.

5. The herbicidal compound according to claim 1, wherein said X is 4,5-F₂; and Y is H.

6. A method of controlling barnyard grass produced while growing rice without inflicting any substantial harm to said rice which comprises applying an effective amount of at least one compound of formula (1):



wherein X is hydrogen, halogen, hydroxy, NH₂, CO₂H, CN, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthionyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxycarbonyl, or C₁-C₃ alkylcarbonyl;


Y is hydrogen or fluoro; and

PCT/ KR 2001/001845

03 FEBRUARY 2004

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

7. A herbicidal composition comprising at least one compound of formula (1) together with at least one member selected from the group consisting of: an agriculturally acceptable carrier, a supplement agent, a surfactant and at least one other herbicidal compound:



wherein X is hydrogen, halogen, hydroxy, NH_2 , CO_2H , CN , $\text{C}_1\text{-C}_6$ alkylamino substituted with 1 or 2 of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_3$ haloalkyl, $\text{C}_1\text{-C}_3$ haloalkoxy, $\text{C}_2\text{-C}_4$ alkoxyalkoxy, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_2\text{-C}_6$ alkenyloxy, $\text{C}_2\text{-C}_6$ alkynyloxy, $\text{C}_1\text{-C}_3$ alkoxycarbonyl, or $\text{C}_1\text{-C}_3$ alkylcarbonyl;


Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

8. A herbicidal compound, (R)-phenoxypropionic acid-N-methyl-N-2-fluorophenyl amide of formula (1), having stability toward rice and preventing the generation of barnyard grass:

PCT/ KR 2001/001845

03 FEBRUARY 2004




wherein X is hydroxy, NH_2 , CO_2H , CN , $\text{C}_1\text{-C}_6$ alkylamino substituted with 1 or 2 of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkoxyalkoxy, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_2\text{-C}_6$ alkenyloxy, $\text{C}_2\text{-C}_6$ alkynyloxy, $\text{C}_1\text{-C}_3$ alkoxycarbonyl, or $\text{C}_1\text{-C}_3$ alkylcarbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

9. A herbicidal compound, (R)-phenoxypropionic acid-N-methyl-N-2-fluorophenyl amide of formula (1), having stability toward rice and preventing the generation of barnyard grass:



wherein X is hydroxy, NH_2 , CO_2H , CN , $\text{C}_1\text{-C}_6$ alkylamino substituted with 1 or 2 of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_1\text{-C}_3$ alkoxycarbonyl,


03 FEBRUARY 2004

or C₁-C₃ alkylcarbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

10. A method of controlling barnyard grass produced while growing rice without inflicting any substantial harm to said rice which comprises applying an effective amount of at least one compound of formula (1):



wherein X is hydroxy, NH₂, CO₂H, CN, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₂-C₄ alkoxyalkoxy, C₁-C₄ alkylthionyl, C₁-C₄ alkylsulfonyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxy carbonyl, or C₁-C₃ alkylcarbonyl;


Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

11. A method of controlling barnyard grass produced while growing rice without inflicting any substantial harm to said rice which comprises applying an effective amount of at least one compound of formula (1):

PCT/ KR 2001/001845

03 FEBRUARY 2004




(1)

wherein X hydroxy, NH_2 , CO_2H , CN , $\text{C}_1\text{-C}_6$ alkylamino substituted with 1 or 2 of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_1\text{-C}_3$ alkoxy carbonyl, or $\text{C}_1\text{-C}_3$ alkylcarbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.

12. A herbicidal composition comprising at least one compound of formula (1) together with at least one member selected from the group consisting of: an agriculturally acceptable carrier, a supplement agent, a surfactant and at least one other herbicidal compound:



(1)

wherein X is hydroxy, NH_2 , CO_2H , CN , $\text{C}_1\text{-C}_6$ alkylamino substituted with 1 or 2 of $\text{C}_1\text{-C}_3$ alkyl, $\text{C}_2\text{-C}_4$ alkoxyalkoxy, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_2\text{-C}_4$ alkoxyalkoxy, $\text{C}_1\text{-C}_4$ alkylthionyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_2\text{-C}_4$ alkoxyalkoxy,


03 FEBRUARY 2004

C₆ alkenyl, C₂-C₆ alkynyl, C₂-C₆ alkenyloxy, C₂-C₆ alkynyloxy, C₁-C₃ alkoxy carbonyl, or C₁-C₃ alkyl carbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.


13. A herbicidal composition comprising at least one compound of formula (1) together with at least one member selected from the group consisting of: an agriculturally acceptable carrier, a supplement agent, a surfactant and at least one other herbicidal compound:



wherein X is hydroxy, NH₂, CO₂H, CN, C₁-C₆ alkylamino substituted with 1 or 2 of C₁-C₃ alkyl, C₁-C₄ alkylthionyl, C₁-C₄ alkylsulfonyl, C₁-C₃ alkoxy carbonyl, or C₁-C₃ alkyl carbonyl;

Y is hydrogen or fluoro; and

n is an integer of 0 to 2, wherein X can be a combination of other substituents when n is 2.



(1)