UK Patent Application (19)GB (11)2490450

(43) Date of Reproduction by UK Office

31.10.2012

1214255.0 (21) Application No:

(22) Date of Filing:

19.01.2011

Date Lodged:

09.08.2012

(30) Priority Data:

(31) 1002278.8

(32) 11.02.2010

(33) GB

(86) International Application Data: PCT/GB2011/050079 En 19.01.2011

(87) International Publication Data: WO2011/098780 En 18.08.2011

(71) Applicant(s):

Johnson Matthey Public Limited Company 5th Floor 25 Farringdon Street, LONDON, EC4A 4AB, **United Kingdom**

(72) Inventor(s):

Alan Cooper Mark Dixon

(74) Agent and/or Address for Service:

Johnson Matthey Catalysts Intellectual Property Department, PO Box 1, Belasis Avenue, BILLINGHAM, Cleveland, TS23 1LB, **United Kingdom**

(51) INT CL:

B01J 31/22 (2006.01) C08G 18/48 (2006.01) C08G 63/84 (2006.01) C08G 18/22 (2006.01) C08G 18/66 (2006.01) C08G 63/85 (2006.01)

(56) Documents Cited by ISA:

WO 2008/155569 A1 WO 2004/050239 A2 US 4502967 A

WO 2007/003966 A1 DE 004242622 A1

(58) Field of Search by ISA: INT CL B01J, C08G Other: EPO -INTERNAL

- (54) Title of the Invention: Glycerol complex composition and method of preparation Abstract Title: Glycerol complex composition and method of preparation
- (57) A composition has an empirical formula M (glycerol) a (X) b, where M represents a metal atom selected from titanium, zirconium, hafnium or aluminium, X is a ligand derived from acetylacetone or a peroxo ion; a is a number between 1 and 2. 5; b is a number in the range from 1 to 2. An alternative composition results from the reaction of a compound of titanium, zirconium, hafnium or aluminium with (a) glycerol and (b) either: (i) acetylacetone or (ii) hydrogen peroxide, an inorganic base and water. The composition is useful in applications requiring water- stable metal chelates, particularly as a catalyst for esterification and polyurethane reactions.