

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number
WO 2006/021000 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2005/029383

(22) International Filing Date: 18 August 2005 (18.08.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/603,347 19 August 2004 (19.08.2004) US

(71) Applicant (for all designated States except US): **AVENTIS PHARMACEUTICALS INC.** [US/US]; 300 Somerset Corporate Boulevard, Bridgewater, New Jersey 08807-2854 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FINK, David Marc** [US/US]; 1 Kensington Court, Lebanon, New Jersey 08833 (US). **CHIANG, Yulin** [US/US]; 24 Vom Eigen Drive, Convent Station, New Jersey 07960 (US). **COLLAR, Nicola Dawn** [GB/US]; 64 Monroe Street, Hoboken, New Jersey 07030 (US).

(74) Agents: **STRUPCZEWSKI, Joseph** et al.; sanofi-aventis, Route 202-206, P. O. Box 6800, Bridgewater, New Jersey 08807-0800 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:



- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED THIENOPYRROLE CARBOXYLIC ACID AMIDES, PYRROLOTHIAZOLE CARBOXYLIC ACID AMIDES, AND RELATED ANALOGS AS INHIBITORS OF CASEIN KINASE I EPSILON



(57) Abstract: The present invention discloses and claims compounds as inhibitors of human casein kinase Iε and methods for using said compounds for treating central nervous system diseases and disorders including mood disorders and sleep disorders. Pharmaceutical compositions comprising said compounds and a method for the preparation are also disclosed and claimed.

5

**SUBSTITUTED THIENOPYRROLE CARBOXYLIC ACID AMIDES,
PYRROLOTHIAZOLE CARBOXYLIC ACID AMIDES, AND RELATED ANALOGS
AS INHIBITORS OF CASEIN KINASE I ϵ**

10

Background of the Invention

1. Field of the Invention

15

This invention relates to a series of substituted 4H-thieno[3,2-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amides, 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amides and related analogs. More specifically the invention relates to 3-arylthio-substituted and 3-heterocyclethio-substituted 4H-thieno[3,2-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amides, 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amides, and related analogs, and to methods of making the compounds of the invention. The compounds of the invention are inhibitors of human casein kinase I ϵ phosphorylation of the human clock protein Period (hPER) and are therefore useful, as pharmaceutical agents, especially in the treatment and/or prevention of diseases and disorders associated with the central nervous system.

25

2. Description of the Art

Rhythmic variations in behavior are displayed by many organisms, ranging from single cells to man. When the rhythm persists under constant conditions, and has a period of about one day, depending little on temperature, the rhythm is called "circadian" (Konopka, R.J. and Benzer, S. (1971) Proc. Nat. Acad. Sci. USA **68**, 2112-2116).

30

Circadian rhythms are generated by endogenous biological pacemakers (circadian clocks) and are present in most living organisms including humans, fungi, insects and bacteria (Dunlap, J.C. (1999) *Cell* **96**, 271-290; Hastings, J.W. et al. *Circadian Rhythms, The Physiology of Biological Timing*. In: Prosser, C.L. ed. *Neural and Integrative Animal Physiology*, New York: Wiley-Liss (1991) 435-546; Allada, R. et al. (1998) *Cell* **93**, 791-804; Kondo et al. (1994) *Science* **266**, 1233-1236; Crosthwaite, S.K. et al. (1997) *Science* **276**, 763-769; Shearman, L.P. et al. (1997) *Neuron*, **19**, 1261-1269). Circadian rhythms are self-sustaining and constant even under conditions of total darkness, but can be synchronized (entrained) to a new day/night regime by environmental signals such as light and temperature cycles (Pittendrigh, C.S. (1993) *Annu. Rev. Physiol.*, **55**, 16-54; Takahashi, J.S. (1995) *Annu. Rev. Neurosci.* **18**, 531-553; Albrecht, U. et al. (1997) *Cell*, **91**, 1055-1064). Circadian clocks are essential for maintaining biological rhythms and regulate a variety of circadian behaviors such as daily fluctuations in behavior, food intake and the sleep/wake cycle, as well as physiological changes such as hormone secretion and fluctuations in body temperature (Hastings, M. (1997) *Trends Neurosci.* **20**, 459-464; Reppert, S.M. and Weaver, D.R. (1997) *Cell* **89**, 487-490).

Genetic and molecular studies in the fruit fly *Drosophila melanogaster* led to elucidation of some of the genes involved in circadian rhythmicity. These studies led to recognition of a pathway that is closely auto-regulated and comprised of a transcription/translation-based negative feed back loop (Dunlap, J.C. (1999) *Cell*, **96**, 271-290; Dunlap, J.C. (1996) *Annu. Rev. Genet.* **30**, 579-601; Hall, J.C. (1996) *Neuron*, **17**, 799-802). The core elements of the circadian oscillator in *Drosophila* consists of two stimulatory proteins dCLOCK/dBMAL (CYCLE) and two inhibitory proteins dPERIOD (dPER) and dTIMELESS (dTIM). dCLOCK and dBMAL heterodimerize forming the transcription factor dCLOCK/dBMAL that promotes expression of two genes termed *Drosophila Period* (*dper*) and *Drosophila Timeless* (*dtim*). Ultimately the mRNAs from these genes are transcribed to afford the proteins dPER and dTIM, respectively. For several hours the protein products dPER and dTIM are synthesized and phosphorylated in the cytoplasm, reach a critical level, and form heterodimers that are translocated into the nucleus. Once in the nucleus dPER and dTIM function as negative regulators of their own transcription, accumulation of dPER and dTIM declines, and activation of *dper* and *dtim* by dCLOCK/dBMAL starts again (Zylka, M.J. et al. (1998) *Neuron* **20**, 1103-1110; Lowrey, P.L. et al. (2000) *Cell* **288**, 483-491). The *dper* gene

has been shown to be a necessary element in controlling circadian rhythms in adult eclosion (the emergence of the adult fly from the pupa) behavior and locomotor activity (Konopka, R.J., & Benzer, S. (1971) Proc. Natl. Acad. Sci. USA, **68**, 2112-2116). Missense mutations of the *per* gene can either shorten (*per^S*) or lengthen (*per^L*) the period of circadian rhythms, while
5 nonsense mutations (*per^o*) cause arrhythmicity in their behaviors (Hall, J.C. (1995) Trends Neurosci. **18**, 230-240).

In mammals, the suprachiasmatic nuclei (SCN) of the anterior hypothalamus are the site of a master biological clock (for review see Panda et al, (2002) Nature **417**, 329 - 335;
10 Reppert, S.M. and Weaver, D.R. (1997) Cell, **89**, 487-490). The SCN clock is entrained to the 24 hour day by the daily light-dark cycle, with light acting through both direct and indirect retina-to-SCN pathways (Klein, D.C. et al. (1991) Suprachiasmatic Nuclei: The Mind's Clock, Oxford University Press, New York). In the SCN of rodents, three *Per* genes have been identified and cloned, and are designated as mouse *Per1* (*mPer1*), *mPer2* and *mPer3*. The
15 protein products of these mammalian genes (mPER1, mPER2, mPER3) share several regions of homology to each other, and each mammalian *Per* gene encodes a protein with a protein dimerization domain designated as PAS (PAS is an acronym for the first three proteins PER, ARNT and SIM found to share this functionally important dimerization domain) that is highly homologous to the PAS domain of insect PER. All *Per* messenger RNAs (mRNAs) and
20 protein levels oscillate during the circadian day and are intimately involved in both positive and negative regulation of the biological clock, but only mPER1 and mPER2 oscillate in response to light (Zylka, M.J. et al. (1998) Neuron **20**, 1103-1110.; Albrecht, U. et al., (1997) Cell **91**, 1055-1064; Shearman, L.P. et al. (1997) Neuron **19**, 1261-1269). The mammalian homolog of the *Drosophila tim* gene was cloned and designated as *mTim*. However, there was
25 no evidence for mPER-mTIM interactions analogous to those observed in *Drosophila*, and it was suggested that PER-PER interactions may have replaced the function of PER-TIM dimers in the molecular workings of the mammalian circadian clock (Zylka, M.J. et al., (1998) Neuron **21**, 1115-1122). Another possibility is that rhythms in PER1 and PER2 form negative feedback loops that regulate the transcriptional activity of the Clock protein (via their PAS
30 domains), which, in turn, drives expression of either or both *Per* genes (Shearman, L.P. et al. (1997) Neuron **19**, 1261-1269).

Understanding the roles of the three *mPer* genes in the mammalian clockwork has been the subject of much investigation. The structural homology of the mPER proteins to dPER led to the expectation that the mPER proteins would function as negative elements in the mammalian feedback loop. PER1 is believed to be involved in the negative regulation of its own transcription in the feedback loop, but recent evidence points to it being involved in the input pathway (Hastings, M.H. et al. (1999) Proc. Natl. Acad. Sci. USA **26**, 15211-15216). PER2 is the most well characterized protein, and mPER2 mutant mice (*mPer2^{Brdm1}*), lacking 87 residues at the carboxyl portion of the PAS dimerization domain, have a shortened circadian cycle in normal light-dark settings, but show arrhythmicity in complete darkness. The mutation also diminishes the oscillating expression of both *mPer1* and *mPer2* in the SCN, indicating that *mPer2* may regulate *mPer1* in vivo (Zheng, B. et al. (1999) Nature **400**, 169-173). PER2 has been shown to have a dual function in the regulation of the "gears" of the central clock (Shearman, L.P. et al. (2000) Science **288**, 1013-1018). In that study, PER2 was shown to bind to cryptochrome (CRY) proteins and translocate to the nucleus where CRY negatively regulated transcription driven by the CLOCK and BMAL1 positive transcriptional complexes. Upon nuclear entry, PER2 initiated the positive arm of the clock by positively regulating BMAL1 transcription by a yet unidentified mechanism. The function of PER3 is poorly understood; however, in *mPer3* knockout mice a subtle effect on circadian activity is observed, and therefore PER3 has been suggested to be involved in the circadian controlled output pathways (Shearman, L.P. et al. (2000) Mol. Cell. Biol. **17**, 6269-6275). It has been reported that mPER proteins interact with each other and that mPER3 can serve as a carrier of mPER1 and mPER2 to bring them into the nucleus which is critical for the generation of circadian signals in the SCN (Kume, K. et al. (1999) Cell **98**, 193-205; Takano, A. et al. (2000), FEBS Letters, **477**, 106-112).

Phosphorylation of the components of the circadian clock has been postulated to regulate the duration of the cycle. The first genetic evidence that a specific protein kinase regulates the *Drosophila* circadian rhythm was the discovery of the novel gene doubletime (*dbt*), encoding a protein serine/threonine kinase (Price J.L. et al. (1998) Cell **94**, 83-95; Kloss B. et al. (1998) Cell **94**, 97-107). Missense mutations in the *dbt* result in an altered circadian rhythm. Null alleles of *dbt* result in hypophosphorylation of dPER and arrhythmia.

The mammalian kinases most closely related to DBT are casein kinase I ϵ (CKI ϵ) and casein kinase I δ (CKI δ). Both kinases have been shown to bind to mPER1, and several studies have shown that CKI ϵ phosphorylates both mouse and human PER1 (Price J.L. et al. (1998) *Cell* **94**, 83-95; Kloss B. et al. (1998) *Cell* **94**, 97-107). In a study with human embryonic kidney 293T cells co-transfected with wild-type hCKI ϵ , hPER1 showed a significant increase in phosphorylation (evidenced by a shift in molecular mass). In this study, the phosphorylated hPER1 had a half-life of approximately twelve hours whereas unphosphorylated hPER1 remained stable in the cell for more than 24 hours, suggesting phosphorylation of hPER1 leads to a decrease in protein stability (Kessler, G.A. et al. (2000) *NeuroReport*, **11**, 951-955). Another study also showed the consequence of PER1 phosphorylation by hCKI ϵ includes both cytoplasmic retention and protein instability (Vielhaber, E. et al. (2000) *Mol. Cell. Biol.* **13**, 4888-4899; Takano, A. et al. (2000) *FEBS Letters* **477**, 106-112).

There has been no biochemical reason to choose between CKI ϵ or CKI δ as a potential regulator in mammals until Lowery et al. [(2000) *Science* **288**, 483-491] found that in the Syrian Golden hamster, semidominant mutations in CKI ϵ (*tau* mutation, Ralph, M.R. and Menaker, M. (1988) *Science* **241**, 1225-1227) caused a shortened circadian day in both heterozygous (22 h) and homozygous (20 h) animals. In this instance, reduced levels of CKI ϵ activity resulted in less PER phosphorylation with presumably higher levels of cytoplasmic PER protein leading to enhanced nuclear entry and altered circadian cycles. More recently, it has been suggested that CKI δ may also be involved in regulating circadian rhythmicity by post-translation modification of mammalian clock proteins hPER1 and hPER2 [Camacho, F. et al., (2001) *FEBS Letters* **489**(2,3), 159-165]. Thus, inhibitors, including small molecule inhibitors, of mammalian or human CKI ϵ and/or CKI δ provide a novel means to phase shift or reset the circadian clock. As discussed below, the alteration of circadian rhythm may find utility for the treatment of sleep or mood disorders.

U.S. patent 6,555,328 B1 discloses screening methods in cells to identify compounds that alter circadian rhythms based on a test compound altering the ability of human casein kinase I ϵ and/or human casein kinase I δ to phosphorylate the human clock proteins hPER1, hPER2 and hPER3. For example, HEK293T cells are co-transfected with hCKI ϵ and Per1 or Per2. For the purpose of evaluating the relevancy of CKI ϵ inhibition and CKI ϵ inhibitors to

circadian biology, a high-throughput cellular assay (33rd Annual Meeting, Soc. for Neurosci., November 8-12, 2003, Abstract numbers 284.1, 284.2, and 284.3) was developed in which circadian rhythm could be monitored in a routine manner. The assay consists of Rat-1 fibroblasts stably expressing a *Mper1-luc* construct, thus enabling the determination of the rhythmic activation of the *Mper1* promoter in living cells by repeatedly estimating luciferase activity by monitoring light-output over several days. The repeated measure format of the assay permits accurate and reproducible assessment of the concentration-dependent effects of CKI ϵ inhibitors on circadian rhythm and provides the nexus for relating CKI ϵ inhibition to circadian period alteration.

Sleep disorders have been classified into four major categories that include primary sleep disorders (dyssomnias and parasomnias), sleep disorders associated with medical/psychiatric disorders and a category of proposed sleep disorders for sleep disorders that cannot be classified due to insufficient data. Primary sleep disorders are thought to arise from abnormalities in the intrinsic systems responsible for sleep-wake generation (homeostatic system) or timing (circadian system). Dyssomnias are disorders in initiating or maintaining sleep and include primary insomnia, hypersomnia (excessive sleepiness), narcolepsy, breathing-related sleep disorder, circadian rhythm sleep disorder, and dyssomnias not otherwise specified. Primary insomnia is characterized by the persistence (>1 month) in difficulty of initiating and maintaining sleep or of non-restorative sleep. Difficulties in sleeping associated with primary insomnia leads to significant distress or impairment, including daytime irritability, loss of attention and concentration, fatigue and malaise, and deterioration of mood and motivation. Circadian rhythm sleep disorders include jet lag syndrome, shift work sleep disorder, advanced sleep phase syndrome and delayed sleep phase syndrome (J. Wagner, M.L. Wagner and W.A. Hening, *Annals of Pharmacotherapy* (1998) **32**, 680-691). Individuals in a forced sleep paradigm demonstrate a greater wakefulness, as a percentage of sleep time, at certain periods of circadian day (Dijk and Lockley, *J. Appl. Physiol.* (2002) **92**, 852-862). It has been generally accepted that with age there is an advance in our circadian rhythm for sleep and often results in less quality sleep (*Am J Physiol Endocrinol Metab.* (2002) **282**, E297-E303). Thus, sleep occurring out of circadian phase may suffer in qualitative and quantitative terms, as further exemplified by alterations in sleep with shift work and jet lag. Disturbance of the human circadian clock can cause sleep disorders and agents that modulate circadian rhythmicity, such as an inhibitor of CKI ϵ and/or

CKI δ , may be useful for the treatment of sleep disorders, and particularly circadian rhythm sleep disorders.

Mood disorders are divided into depressive disorders ("unipolar depression"), bipolar disorders, and two disorders based on etiology that include mood disorder due to a general medical condition and substance-induced mood disorder. Depressive disorders are subclassified as major depressive disorder, dysthymic disorder and depressive disorder not otherwise specified. Bipolar disorders are subclassified as bipolar I disorder and bipolar II disorder. It has been observed that the specifier "seasonal pattern" can be applied to major depressive disorders that are recurrent and to the pattern of major depressive episodes in bipolar I disorder and bipolar II disorder. Prominent anergy, hypersomnia, overeating, weight gain, and a craving for carbohydrates often characterize major depressive episodes that occur in a seasonal pattern. It is unclear whether a seasonal pattern is more likely in major depressive disorder that is recurrent or in bipolar disorders. However, within the bipolar disorders, a seasonal pattern appears to be more likely in bipolar II disorder than in bipolar I disorder. In some individuals the onset of manic or hypomanic episodes may also be linked to a particular season. The winter-type seasonal pattern appears to vary with latitude, age and sex. Prevalence increases with higher latitudes, younger persons are at higher risk for winter depressive episodes, and females comprise 60% to 90% of persons with seasonal pattern. Seasonal affective disorder (SAD), a term commonly used in the literature, is a subtype of mood disorder that in the Diagnostic and Statistical Manual of Mental Disorders IV (DSM-IV) (American Psychiatric Association: "Diagnostic and Statistical Manual of Mental Disorders", Fourth Edition, Text Revision. Washington, DC, American Psychiatric Association, 2000) is denoted by the term "with seasonal pattern" when describing a seasonal pattern of major depressive episodes in bipolar I disorder, bipolar II disorder or recurrent major depressive disorder (E. M. Tam et al., Can. J. Psychiatry (1995) **40**, 457-466). The characteristics and diagnoses of depressive disorders, major depressive disorder, major depressive episode, bipolar I disorder, bipolar II disorder and seasonal effects are described in DSM-IV,

Patients suffering from major depressive disorders, including SAD that is characterized by recurrent depressive episodes typically in winter, have been shown to be positively responsive to light therapy (Kripke, Journal of Affective Disorders (1998) **49**(2), 109-117). The success of bright light treatment for patients with SAD and major depression

resulted in the proposal of several hypotheses to explain the underlying mechanism of action for the therapeutic effect of light. These hypotheses included the “circadian rhythm hypothesis” that suggests the antidepressant effect of bright light could be associated with phase-shifting the circadian pacemaker relative to sleep (E. M. Tam et al., *Can. J. Psychiatry* (1995) **40**, 457-466). In support of the link between light therapy and circadian rhythm, clinically effective light therapy in major depressive disorders causes a concomitant shift in circadian phase and the clinical effectiveness of light therapy appears to depend on the phase-shifting ability of the light therapy (Czeisler et al., *The Journal of Physiology* (2000) 526 (Part 3), 683-694; Terman et al., *Arch. Gen. Psychiatry* (2001) **58**, 69-75). Additionally, light-therapy has been shown to accelerate and augment the effectiveness of the pharmacological treatment of major depressive disorders (Benedetti et al., *J. Clin. Psychiatry* (2003) **64**, 648-653). Thus, inhibition of casein kinase I ϵ and/or casein kinase I δ would be expected to cause a circadian phase shift and such inhibition represents a potential clinically effective mono- or combined therapy for mood disorders.

It should be noted that sleep disturbance is a criterion symptom for many psychiatric disorders (W.V. McCall, *J. Clin. Psychiatry* (2001) **62** (suppl 10), 27-32). Sleep disturbances are a common feature of depressive disorders and insomnia is the sleep disturbance that is frequently reported in depression, occurring in over 90% of depressed patients (M.E. Thase, *J. Clin. Psychiatry* (1999) **60** (suppl 17), 28-31). Accumulating evidence supports a common pathogenesis for primary insomnia and major depressive disorder. It has been hypothesized that corticotrophin releasing factor (CRF) hyperactivity (due to genetic predisposition or possibly early stress) and stress induce a process leading to exaggerated and protracted sleep disturbances, and eventually primary insomnia. Circadian rhythmicity in CRF secretion under nonstressed conditions may play a role in the normal sleep-wake expression (G.S. Richardson and T. Roth, *J. Clin Psychiatry* (2001) **62** (suppl 10), 39-45). Thus, agents that modulate circadian rhythmicity, for example by inhibition of casein kinase I ϵ and/or casein kinase I δ , may be useful for treatment of depressive disorders due to effects on CRF secretion.



All of the references referred to hereinabove are incorporated herein by reference in their entirety.

Thus it is an object of this invention to provide a series of substituted 4H-thieno[3,2-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amides, 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amides and related analogs that are inhibitors of casein kinase I ϵ . This object and other
 5 objects of this invention become apparent from the detailed discussion of the invention that follows.

SUMMARY OF THE INVENTION

The present invention provides substituted 4H-thieno[3,2-b]pyrrole-5-carboxylic acid
 10 amides, 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amides, 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amides, 4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amides and related analogs of formula (I) and formula (II), and the stereoisomers, enantiomers, racemates and tautomers of said compounds and the pharmaceutically acceptable salts thereof, as inhibitors of human casein kinase I ϵ activity, and methods of using the compounds of
 15 formula (I) and formula (II) as pharmaceutical agents for the treatment of diseases and disorders of the central nervous system, such as for example mood disorders including major depressive disorder, bipolar I disorder and bipolar II disorder, and sleep disorders including circadian rhythm sleep disorders such as for example shift work sleep disorder, jet lag syndrome, advanced sleep phase syndrome and delayed sleep phase syndrome. The present
 20 invention also provides methods for making the compounds of formula (I) and formula (II) of the invention.

Accordingly, a broad embodiment of the invention is directed to a compound of formula (I) or formula (II):



wherein

X is S or S(O)_n;

R₁ is H or C₁-C₆alkyl;

R₂ is NR₅R₆;

R₃ is aryl or heterocycle;

R₄ is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl),
5 C₁-C₆alkoxy, aryl-(C₁-C₆alkoxy), heterocycle-(C₁-C₆alkoxy), CF₃, halogen, SH, C₁-C₆alkylthio,
aryl-(C₁-C₆alkylthio), heterocycle-(C₁-C₆alkylthio), NO₂, NH₂, NR₅R₆, aryl-(C₁-
C₆alkylamino), heterocycle-(C₁-C₆alkylamino), or XR₃ wherein X and R₃ are as defined
above;

R₅ is H or C₁-C₆alkyl;

10 R₆ is H or C₁-C₆alkyl;

L is N or CR₇ wherein R₇ is H or C₁-C₆alkyl;

M is S, O or NR₈ wherein R₈ is H, C₁-C₆alkyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl) or
acyl;

n is 1 or 2; or

15 a stereoisomer, an enantiomer, a racemate or a tautomer thereof; or
a pharmaceutically acceptable salt thereof.

Another embodiment of the present invention relates to a method for inhibiting casein
kinase Iε activity in a patient comprising administering to said patient a therapeutically
20 effective amount of a compound of formula (I) or formula (II).

Another embodiment of the present invention relates to a method for treating a patient
suffering from a disease or disorder ameliorated by inhibition of casein kinase Iε activity
comprising administering to said patient a therapeutically effective amount of a compound of
25 formula I or formula II.

A further embodiment of the present invention relates to a process for preparing a
compound of formula (I) or formula (II).

30 DETAILED DESCRIPTION OF THE INVENTION

As used herein, "stereoisomer" is a general term used for all isomers of individual
molecules that differ only in the orientation of their atoms in space. The term stereoisomer

includes mirror image isomers (enantiomers), mixtures of mirror image isomers (racemates, racemic mixtures), geometric (cis/trans or E/Z) isomers, and isomers of compounds with more than one chiral center that are not mirror images of one another (diastereoisomers). The compounds of the present invention may have asymmetric centers and occur as racemates, racemic mixtures, individual diastereoisomers, or enantiomers, or may exist as geometric isomers, with all isomeric forms of said compounds being included in the present invention.

As used herein, "R" and "S" are used as commonly used in organic chemistry to denote specific configuration of a chiral center. The term "R" (rectus) refers to that configuration of a chiral center with a clockwise relationship of group priorities (highest to second lowest) when viewed along the bond toward the lowest priority group. The term "S" (sinister) refers to that configuration of a chiral center with a counterclockwise relationship of group priorities (highest to second lowest) when viewed along the bond toward the lowest priority group. The priority of groups is based upon sequence rules wherein prioritization is first based on atomic number (in order of decreasing atomic number). A listing and discussion of priorities is contained in *Stereochemistry of Organic Compounds*, Ernest L. Eliel, Samuel H. Wilen and Lewis N. Mander, editors, Wiley-Interscience, John Wiley & Sons, Inc., New York, 1994.

In addition to the (R)-(S) system, the older D-L system may also be used herein to denote absolute configuration, especially with reference to amino acids. In this system a Fischer projection formula is oriented so that the number 1 carbon of the main chain is at the top. The prefix "D" is used to represent the absolute configuration of the isomer in which the functional (determining) group is on the right side of the carbon at the chiral center and "L", that of the isomer in which it is on the left.

As used herein, "tautomer" or "tautomerism" refers to the coexistence of two (or more) compounds that differ from each other only in the position of one (or more) mobile atoms and in electron distribution, for example, keto-enol tautomers or tautomerism.

As used herein, "alkyl" refers to a saturated linear or branched chain aliphatic hydrocarbon group having from one to six carbon atoms, and includes methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tert-butyl and the like groups. Included within the meaning of "alkyl" are "alkylene" or "alkylenyl" as are defined herein below.

As used herein "alkylene" or "alkylenyl" refers to a linear or branched, divalent, saturated aliphatic chain of one to six carbons and includes methylenyl, ethylenyl, propylenyl, isopropylenyl, butylenyl, isobutylenyl, t-butylenyl, pentylenyl, isopentylenyl, hexylenyl and the like groups.

5 As used herein "alkenyl" refers to a linear or branched monovalent unsaturated aliphatic chain having from two to six carbon atoms and includes ethenyl (also known as vinyl), 1-methylethenyl, 1-methyl-1-propenyl, 1-butenyl, 1-hexenyl, 2-methyl-2-propenyl, 2,4-hexadienyl, 1-propenyl, 2-propenyl, 2-butenyl, 2-pentenyl, and the like groups.

10 As used herein "alkynyl" is a linear or branched monovalent unsaturated aliphatic having from two to six carbon atoms with at least one triple bond and includes ethynyl, 1-propynyl, 1-butylnyl, 1-hexynyl, 2-propynyl, 2-butylnyl, 2-pentylnyl and the like groups.

15 As used herein, "alkoxy" or "alkyloxy" refers to a monovalent substituent which consists of a linear or branched alkyl chain having from one to six carbon atoms linked through an ether oxygen atom and having its free valence bond from the ether oxygen, and includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, tert-butoxy and the like groups.

20 As used herein, "alkylthio" refers to a monovalent substituent which consists of a linear or branched alkyl chain having from one to six carbon atoms linked through a sulfur atom and having its free valence bond from the sulfur, and includes methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio and the like groups.

25 As used herein "alkenyloxy" refers to a linear or branched monovalent unsaturated aliphatic chain having from two to six carbon atoms linked through an ether oxygen atom and having its free valence bond from the ether oxygen, and includes ethenyloxy (also known as vinyloxy), 1-methylethyloxy, 1-methyl-1-propenyloxy, 1-butyloxy, 1-hexenyloxy, 2-methyl-2-propenyl, 2,4-hexadienyloxy, 1-propenyloxy, 2-propenyloxy, 2-butyloxy, 2-pentyloxy, and the like groups.

30 As used herein "alkynyloxy" refers to a linear or branched monovalent unsaturated aliphatic chain having from two to six carbon atoms with at least one triple bond linked through an ether oxygen atom and having its free valence bond from the ether oxygen, and includes ethynyloxy, 1-propynyloxy, 1-butyloxy, 1-hexynyloxy, 2-propynyloxy, 2-butyloxy, 2-pentyloxy and the like groups.

As used herein the term C₃-C₈cycloalkyl refers to a saturated hydrocarbon ring structure containing from three to eight carbon atoms and includes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

As used herein, "aryl" or "Ar" means any stable monocyclic, bicyclic or tricyclic carbon ring of up to seven members in each ring, wherein at least one ring is aromatic and unsubstituted or substituted with from one to three substituents selected from the group consisting of methylenedioxy, hydroxy, C₁-C₆alkoxy, halogen, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, trifluoromethyl, trifluoromethoxy, -NO₂, -NH₂, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NH-acyl, and -N(C₁-C₆alkyl)acyl. Examples of "aryl" or "Ar" include phenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-bromophenyl, 3-bromophenyl, 4-bromophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-aminophenyl, 3-aminophenyl, 4-aminophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 2,4-dichlorophenyl, 2,3-dichlorophenyl, 3,5-dimethylphenyl, 2-trifluoromethoxyphenyl, 3-trifluoromethoxyphenyl, 4-trifluoromethoxyphenyl, naphthyl, tetrahydronaphthyl and biphenyl.

As used herein, the term "aryl-(C₁-C₆alkyl)" refers to an aryl group as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms and having its free valence bond from one of the alkylene chain carbons. Examples of "aryl-(C₁-C₆alkyl)" include phenylmethyl (benzyl), phenylethyl, p-methoxybenzyl, p-fluorobenzyl, p-chlorobenzyl and the like groups.

As used herein, the term "aryl-(C₁-C₆alkoxy)" refers to an aryl group as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through an ether oxygen atom and having its free valence bond from the ether oxygen. Examples of aryl-(C₁-C₆alkoxy) include phenylmethoxy (benzyloxy), phenylethoxy, and the like groups.

As used herein, the term "aryl-(C₁-C₆alkylamino)" refers to an aryl group as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through a nitrogen atom and having its free valence bond from the nitrogen wherein said nitrogen is optionally substituted by a hydrogen or a C₁-C₆alkyl. Examples of aryl-(C₁-

C₆alkylamino) include phenylmethylamino (benzylamino), phenylethylamino, N-methyl-N-benzylamino and the like groups.

As used herein, the term "aryl-(C₁-C₆alkylthio)" refers to an aryl group as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through a sulfur atom and having its free valence bond from the sulfur. Examples of aryl-(C₁-C₆alkylthio) include phenylmethylthio (benzylthio), phenylethylthio, and the like groups.

As used herein, the term "acyl" refers to a H-(C=O)-, C₁-C₆alkyl-(C=O)-, aryl-(C=O)-, aryl(C₁-C₆alkyl)-(C=O)-, heterocycle-(C=O)-, or heterocycle(C₁-C₆alkyl)-(C=O)- group, wherein alkyl, aryl and heterocycle are as defined herein, and having its free valence bond from the carbonyl (C=O) moiety. Included within the meaning of acyl are acetyl, propionyl, butyryl, isobutyryl, trifluoroacetyl, trichloroacetyl, benzoyl and the like groups.

As used herein, "heterocycle" or "heterocyclic" means a stable 5- to 7-membered monocyclic or stable 8- to 11-membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. The heterocyclic ring may be unsubstituted or substituted with from one to three substituents selected from the group consisting of C₁-C₆alkoxy, hydroxy, halogen, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, trifluoromethyl, trifluoromethoxy, -NO₂, -NH₂, -NH(C₁-C₆alkyl), -N(C₁-C₆alkyl)₂, -NH-acyl, and -N(C₁-C₆alkyl)acyl. Examples of such heterocyclic elements include piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepinyl, azepinyl, pyrrolyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, benzofuranyl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, and oxadiazolyl.

As used herein, the term "heterocycle-(C₁-C₆alkyl)" or "heterocyclic-(C₁-C₆alkyl)" refers to a heterocycle or heterocyclic ring as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms to another carbon atom or to a heteroatom selected from the group consisting of O, N and S. Included within the meaning of
5 heterocycle(C₁-C₆alkyl) or heterocyclic(C₁-C₆alkyl) are 4-pyridinylmethyl, 3-pyridinylmethyl, 2-pyridinylmethyl, 2-furanmethyl, 2-thenyl (2-thiophenemethyl), 5-nitro-2-thenyl, 5-(2-chlorophenyl)-2-furanmethyl, 1-(phenylsulfonyl)-1H-pyrrole-2-methyl and the like groups.

As used herein, the term "heterocycle-(C₁-C₆alkoxy)" or "heterocyclic-(C₁-C₆alkoxy)"
10 refers to a heterocycle or heterocyclic ring as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through an ether oxygen atom and having its free valence bond from the ether oxygen. Included within the meaning of heterocycle(C₁-C₆alkoxy) or heterocyclic(C₁-C₆alkoxy) are 2-thienylmethoxy, 3-thienylmethoxy, 2-furanmethoxy, 3-furanmethoxy, 4-pyridinylmethoxy, 3-pyridinylmethoxy,
15 2-pyridinylmethoxy and the like groups.

As used herein, the term "heterocycle-(C₁-C₆alkylamino)" or "heterocyclic-(C₁-C₆alkylamino)" refers to a heterocycle or heterocyclic ring as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through a nitrogen atom and having its free valence bond from the nitrogen wherein said nitrogen is optionally
20 substituted by a hydrogen or a C₁-C₆alkyl. Included within the meaning of heterocycle(C₁-C₆alkylamino) or heterocyclic(C₁-C₆alkylamino) are 2-thienylmethylanino, 3-thienylmethylanino, 2-furanmethylanino, 3-furanmethylanino, 4-pyridinylmethylanino, 3-pyridinylmethylanino, 2-pyridinylmethylanino and the like groups.

As used herein, the term "heterocycle-(C₁-C₆alkylthio)" or "heterocyclic-(C₁-C₆alkylthio)" refers to a heterocycle or heterocyclic ring as defined above linked by a linear or branched alkylene chain containing one to six carbon atoms linked through a sulfur atom and having its free valence bond from the sulfur. Included within the meaning of heterocycle(C₁-C₆alkylthio) or heterocyclic(C₁-C₆alkylthio) are 2-thienylmethylthio, 3-thienylmethylthio, 2-furanmethylthio, 3-furanmethylthio, 4-pyridinylmethylthio, 3-pyridinylmethylthio, 2-pyridinylmethylthio and the like groups.
30

As used herein, "halogen", "hal" or "halo" refers to a member of the family of fluorine, chlorine, bromine or iodine.

When any variable (e.g., aryl, heterocycle, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, X) occurs more than one time in any constituent or in a compound of formula (I) or formula (II) of this invention, its definition on each occurrence is independent of its definition at every other occurrence unless indicated otherwise. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "treat", "treating" or "treatment" refers to:

- (i) preventing a disease, disorder or condition from occurring in a patient that may be predisposed to the disease, disorder and/or condition, but has not yet been diagnosed as having it;
- (ii) inhibiting a disease, disorder or condition, i.e., arresting its development; or
- (iii) relieving a disease, disorder or condition, i.e., causing regression of the disease, disorder and/or condition.

As used herein, the term "patient" refers to a warm blooded animal such as a mammal which is afflicted with a particular disease, disorder or condition. It is explicitly understood that guinea pigs, dogs, cats, rats, mice, horses, cattle, sheep, and humans are examples of animals within the scope of the meaning of the term.

As used herein, "disease" refers to an illness, sickness or an interruption, cessation or disorder of body functions, systems or organs.

As used herein, "disorder" refers to a disturbance of function, structure or both resulting from a genetic or embryologic failure in development, or from exogenous factors such as poison, injury or disease.

As used herein, "condition" refers to a state of being, health or physical fitness.

As used herein, "prophylaxis" refers to the prevention of disease.

As used herein, the term "sleep disorder", "sleep disorders" or "sleep disturbance" means insomnia.

As used herein, the term "insomnia" means the inability to sleep in the absence of external impediments, such as noise, bright light, etc., during the period when sleep should normally occur and the inability to sleep may vary in degree from restlessness or disturbed slumber to a curtailment of the normal length of sleep or to absolute wakefulness. The term "insomnia" includes primary insomnia, insomnia related to a mental disorder, substance-induced insomnia and circadian rhythm insomnia that is insomnia due to a change in the

normal sleep-wake schedule (shift changes, shift work sleep disorder, jet lag or jet lag syndrome, etc.).

As used herein the term "primary insomnia" means difficulty in initiating sleep, in maintaining sleep or having restorative sleep which is not caused by a mental disorder or due to physiological effects of taking or withdrawing from certain substances (substance-induced insomnia).

As used herein the term "circadian rhythm sleep disorder" includes jet lag or jet lag syndrome, shift work sleep disorder, advanced sleep phase syndrome and delayed sleep phase syndrome.

As used herein the term "effective inhibitory amount of a compound" or "effective casein kinase I ϵ inhibitory amount of a compound" means enough of a compound that becomes bioavailable through the appropriate route of administration to treat a patient afflicted with a disease, disorder or condition amenable to such treatment.

As used herein the term "a therapeutically effective amount" means an amount of a compound which is effective in treating the named disease, disorder or condition.

As used herein, the phrase "lengthening of circadian rhythm period" refers to increasing the interval between seminal events in a process that occurs regularly with a frequency of approximately once every 24 hours.

As used herein, the phrase "shortening of circadian rhythm period" refers to decreasing the interval between seminal events in a process that occurs regularly with a frequency of approximately once every 24 hours.

As used herein, the term "pharmaceutically acceptable salt" is intended to apply to any salt, whether previously known or future discovered, that is used by one skilled in the art that is a non-toxic organic or inorganic addition salt which is suitable for use as a pharmaceutical.

Illustrative bases which form suitable salts include alkali metal or alkaline-earth metal hydroxides such as sodium, potassium, calcium or magnesium hydroxides; ammonia and aliphatic, cyclic or aromatic amines such as methylamine, dimethylamine, triethylamine, diethylamine, isopropyl-diethylamine, pyridine and picoline. Illustrative acids which form suitable salts include inorganic acids such as, for example, hydrochloric, hydrobromic, sulfuric, phosphoric and like acids, and organic carboxylic acids such as, for example, acetic, propionic, glycolic, lactic, pyruvic, malonic, succinic, fumaric, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic and dihydroxymaleic, benzoic, phenylacetic, 4-aminobenzoic, 4-hydroxybenzoic, anthranilic, cinnamic, salicylic, 4-aminosalicylic, 2-phenoxybenzoic, 2-



acetoxybenzoic, mandelic and like acids, and organic sulfonic acids such as methanesulfonic, benzenesulfonic and p-toluenesulfonic acids.

As used herein, "pharmaceutical carrier" or "pharmaceutically acceptable carrier" refers to known pharmaceutical excipients useful in formulating pharmaceutically active compounds for administration, and which are substantially nontoxic and non-sensitizing under conditions of use. The exact proportion of these excipients is determined by the solubility and chemical properties of the active compound, the chosen route of administration as well as standard pharmaceutical practice. In practicing the methods of this invention, the active ingredient is preferably incorporated into a composition containing a pharmaceutical carrier, although the compounds are effective and can be administered, in and of themselves. That said, the proportion of active ingredient can vary from about 1% to about 90% by weight.

Further abbreviations that may appear in this application shall have the following meanings:

Me (methyl), Et (ethyl), Ph (phenyl), Et₃N (triethylamine), p-TsOH (para-toluene sulfonic acid), TsCl (para-toluenesulfonyl chloride), hept (heptane), DMF (dimethylformamide), NMP (1-methyl-2-pyrrolidinone or N-methyl-2-pyrrolidinone), IPA (isopropanol or isopropyl alcohol), DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), DBN (1,5-diazabicyclo[4.3.0]non-5-ene), rt or r.t. (room temperature or ambient temperature), min or min. (minutes), h (hour or hours), UV (ultraviolet), LCMS (liquid chromatography mass spectrometry), t-Boc or Boc (tert-butoxycarbonyl), Bn (benzyl), t-Bu (tertiary butyl), i-Pr (isopropyl), TFA (trifluoroacetic acid), HOAc (acetic acid), EtOAc (ethyl acetate), Et₂O (diethylether), EtOH (ethanol), DIEA (diisopropylethylamine), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride); HOBt (1-hydroxybenzotriazole), g (gram), mg (milligram), µg (microgram), ng (nanogram), mL (milliliter), µL (microliter), L (liter), HPLC (high-performance liquid chromatography), TLC, tlc or Tlc (thin layer chromatography), g/L (grams per liter), SiO₂ (silica gel), L/min (liters per minute), mL/min (milliliters per minute), mmol (millimole), M (molar), mM (millimolar), µM (micromolar), nM (nanomolar), µCi (microCurie), CPM (counts per minute), rpm (revolutions per minute), mm (millimeter), µm (micrometer), µ (micron), nm (nanometer), ppm (parts per million), psi (pounds per square inch), eq. or equiv. (equivalent), R_T (retention time), °C (degrees Celsius), and K (Kelvin).

Accordingly, a broad embodiment of the invention is directed to a compound of formula (I) or formula (II):



wherein X is S or S(O)_n; R₁ is H or C₁-C₆alkyl; R₂ is NR₅R₆; R₃ is aryl or heterocycle; R₄ is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl), C₁-C₆alkoxy, aryl-(C₁-C₆alkoxy), heterocycle-(C₁-C₆alkoxy), CF₃, halogen, SH, C₁-C₆alkylthio, aryl-(C₁-C₆alkylthio), heterocycle-(C₁-C₆alkylthio), NO₂, NH₂, NR₅R₆, aryl-(C₁-C₆alkylamino), heterocycle-(C₁-C₆alkylamino), or XR₃ wherein X and R₃ are as defined above; R₅ is H or C₁-C₆alkyl; R₆ is H or C₁-C₆alkyl; L is N or CR₇ wherein R₇ is H or C₁-C₆alkyl; M is S, O or NR₈ wherein R₈ is H, C₁-C₆alkyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl) or acyl; and n is 1 or 2.

A further embodiment of this invention relates to compounds of formula (I) or formula (II) wherein M and X are each S.

Another embodiment of this invention relates to compounds of formula (I) wherein L is CR₇ and M and X are each S.

A further embodiment of this invention relates to compounds wherein of formula (I) wherein M and X are each S, L is CR₇, and R₇ is H. The following compounds are representative examples within the scope of this embodiment:

6-phenylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
 6-(3-fluorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
 6-(4-chlorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
 6-(2-aminophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
 6-(pyridin-2-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
 6-p-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,

- 6-(thiophen-2-yl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3,5-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(pyridin-4-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-m-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
5 6-o-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2,3-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2,5-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-ethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3-bromo-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
10 6-(3,5-dimethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3-methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-trifluoromethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-fluoro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide, and
15 6-(3-trifluoromethoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide.

Another embodiment of this invention relates to compounds of formula (I) wherein L is N and M and X are each S. The following compounds are representative examples within the scope of this embodiment:

- 20 6-phenylsulfanyl-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide,
6-(3-fluoro-phenylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide, and
6-(pyridin-2-ylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide.

- Another embodiment of this invention relates to compounds of formula (II) wherein L
25 is CR₇ and M and X are each S.

A further embodiment of this invention relates to compounds wherein of formula (II) wherein M and X are each S, L is CR₇, and R₇ is H. The following compounds are representative examples within the scope of this embodiment:

- 4-(pyridin-2-ylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
30 4-(phenylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
6-(3-fluorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,

4-(pyridin-4-ylsulfanyl)-6H-thieno[2,3-b]-pyrrole-5-carboxylic acid amide,
4-(3,5-dichlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(thiophen-2-yl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(3-bromophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
5 4-(3-methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(2-methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(3-chlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide, and
4-(3-methylphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide.

10 Another embodiment of this invention relates to compounds of formula (II) wherein L is N and M and X are each S. The following compounds are representative examples within the scope of this embodiment:

2-methyl-6-phenyl-sulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-methoxyphenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
15 6-(3-fluorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-chlorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-trifluoromethoxy-phenylsulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid
amide,
2,6-bis-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
20 2,6-bis-(3-methoxy-phenylsulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide, and
6-(3-methoxyphenyl-sulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide.

25 Another embodiment of the present invention relates to a method for inhibiting casein kinase Iε activity in a patient that comprises administering to said patient a therapeutically effective amount of a compound of formula (I) or formula (II) that results in a lengthening of circadian rhythm period.


30 Another embodiment of the present invention relates to a method for treating a patient suffering from a disease or disorder ameliorated by inhibition of casein kinase Iε activity that comprises administering to said patient a therapeutically effective amount of a compound of

formula (I) or formula (II) wherein said inhibition of casein kinase I ϵ activity results in a lengthening of circadian rhythm period.

The compounds of the present invention can be prepared by processes analogous to those known in the art. Reaction schemes 1, 2 and 3, and the corresponding descriptive text, describe the preparation of the various compounds of the invention. The disclosed methods and examples are provided for illustration purposes and in no way limit the scope of the present invention. Alternative reagents, reaction conditions, and other combinations and permutations of the steps herein described to arrive at individual compounds are readily apparent to one of ordinary skill in the art. Tables 1, 2, 3 and 4 provide summaries of the example compounds, and biological data for example compounds of the invention is summarized in Table 5.

CHEMICAL SYNTHESIS

Scheme 1



Scheme 1 describes the synthesis of 4H-thieno[3,2-b]pyrroles (M is S), 4H-furo[3,2-b]pyrroles (M is O), and 1,4-dihydropyrrolo[3,2-b]pyrroles (M is NR₈) of formula (I) wherein L is CR₇, and the synthesis of 6H-thieno[2,3-b]pyrroles (M is S), 6H-furo[2,3-b]pyrroles (M is O), and 1,6-dihydropyrrolo[2,3-b]pyrroles (M is NR₈) of formula (II) wherein L is CR₇ from

known or commercially available esters or carboxylic acids **1** and **3**, respectively, wherein R is alkyl or H.

In scheme 1, step a, starting esters **1** or **3**, wherein R is alkyl, are converted to amides **2** or **4**, respectively, by methods well known to one skilled in the art. Thus, treating a mixture of about 7M ammonia and ester **1** or **3** in a suitable polar solvent, such as for example methanol or ethanol, with a chip of lithium hydroxide and heating the resultant mixture in a pressure vessel at about 100 °C for about 16 hours provides, after chromatographic purification as is well known to one skilled in the art, primary amide **2** or **4**, respectively. Alternatively, other reaction conditions well known to one skilled in the art may be employed, such as treating a solution of ester **1** or **3** in a suitable polar solvent, such as for example methanol or ethanol, with about 5M to about 7M ammonia solution for about one day to about three days at ambient temperature, or by heating the solution to about 55 °C for about 10 hours, provides primary amide **2** or **4**, respectively, after isolation by methods well known to one skilled in the art. Alternatively, ester **1** or **3** may be suspended in a mixture of concentrated ammonium hydroxide solution and lithium chloride at ambient temperature for about three to about five days until thin layer chromatographic analysis, or other suitable chromatographic analysis as is well known to one skilled in the art, indicates that the reaction is substantially complete. Primary amides **2** or **4** are isolated from the reaction mixture by methods well known to one skilled in the art. If primary or secondary C₁-C₆alkylamines are employed in place of ammonia or ammonium hydroxide, there is obtained the corresponding secondary and tertiary amides **2** or **4** wherein R₂ is NR₅R₆, R₅ is H or C₁-C₆alkyl and R₆ is C₁-C₆alkyl.

As shown in scheme I, step b, commercially available or known carboxylic acids **1** or **3** (wherein R is H) may be converted to amides **2** or **4**, respectively, by methods well known to one skilled in the art. Where desired, carboxylic acids **1** or **3** (R is H) may also be prepared by hydrolysis of the corresponding esters **1** or **3** (R is alkyl) by methods well known to one skilled in the art. For example, a suitable base, such as for example potassium hydroxide, sodium hydroxide, lithium hydroxide and like bases, is added to a mixture of ester **1** or **3** in a suitable solvent such as for example a mixture of tetrahydrofuran and water. The mixture is heated at about 90 °C to about 110 °C for about 0.5 hour to about 2 hours. The product is recovered as a salt by filtration and the filtrate is concentrated to provide additional material as a residue. The filter cake and residue are combined and acidified by methods well known to one skilled in the art, such as for example acidification with a suitable acid such as acetic acid in a suitable solvent such as methanol, ethanol and like solvents, to provide carboxylic acids **1**

or **3**, respectively, wherein R is H. As shown in scheme I, step b, for example, a solution of carboxylic acid **1** or **3** in a suitable solvent such as dimethylformamide is treated with a base such as diisopropylethylamine, a carbodiimide such as for example (1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride, 1-hydroxybenzotriazole and ammonium chloride.

5 When the reaction is complete as determined by thin layer chromatography, or other suitable chromatographic analysis as is well known to one skilled in the art, the mixture is diluted with a suitable solvent, and the product is isolated and chromatographically purified by methods well known to one skilled in the art to provide primary amides **2** or **4**, respectively, wherein R₂ is NH₂. If primary or secondary C₁-C₆alkylamines are employed in place of ammonium
10 chloride, there is obtained the corresponding secondary and tertiary amides **2** or **4** wherein R₂ is NR₅R₆, R₅ is H or C₁-C₆alkyl and R₆ is C₁-C₆alkyl.

As shown in scheme 1, step c, intermediate amides **2** or **4** are each thioarylated at the 3-position of the amide-bearing pyrrole ring by methods well known to one skilled in the art. For example, a suspension of intermediate amide **2** or **4** in a suitable solvent, such as for
15 example dimethylformamide or NMP, is treated with a suitable base, such as for example sodium hydride or lithium hydride, at ambient temperature, followed by treatment with a suitable diaryldisulfide or diheterocycledisulfide, and then the mixture is stirred at ambient temperature to about 100°C for about 12 hours to about 20 hours. The course of the reaction is followed by thin layer chromatographic analysis or other chromatographic methods as are
20 well known to one skilled in the art. When complete, the reaction is worked-up by extractive methods as are well known to one skilled in the art. The desired 4H-thieno[3,2-b]pyrroles (M is S), 4H-furo[3,2-b]pyrroles (M is O), and 1,4-dihydropyrrolo[3,2-b]pyrroles (M is NR₈) of formula (I), wherein L is CR₇, X is S and R₃ is aryl or heterocycle, and 6H-thieno[2,3-b]pyrroles (M is S), 6H-furo[2,3-b]pyrroles (M is O), and 1,6-dihydropyrrolo[2,3-b]pyrroles
25 (M is NR₈) of formula (II), wherein L is CR₇, X is S and R₃ is aryl or heterocycle, are each isolated and chromatographically purified by methods as are well known to one skilled in the art.

Alternatively, a mixture of the diaryldisulfide or diheterocycledisulfide and about one
30 equivalent of cesium carbonate in a suitable solvent, such as for example dimethylformamide or NMP, is treated with intermediate amide **2** or **4**, and then the mixture is heated at about 80 °C to about 120°C for about one to about six hours. The reaction is monitored by thin layer chromatography or other chromatographic methods as are well known to one skilled in the art.

The desired 4H-thieno[3,2-b]pyrroles (M is S), 4H-furo[3,2-b]pyrroles (M is O) and 1,4-dihydropyrrolo[3,2-b]pyrroles (M is NR₈) of formula (I), wherein L is CR₇, X is S and R₃ is aryl or heterocycle, and 6H-thieno[2,3-b]pyrroles (M is S), 6H-furo[2,3-b]pyrroles (M is O) and 1,6-dihydropyrrolo[2,3-b]pyrroles (M is NR₈) of formula (II), wherein L is CR₇, X is S and R₃ is aryl or heterocycle, are each isolated and chromatographically purified by methods well known to one skilled in the art.

As shown in scheme 1, optional step d, the nitrogen of the pyrrole ring of a compound of formula (I) or formula (II), wherein R₁ is H, is N-alkylated by treating a solution of the compound formula (I) or formula (II) wherein R₁ is H and a suitable solvent, such as for example 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone with a C₁-C₆-dialkylsulfate and a suitable base, such as for example cesium carbonate, at ambient temperature for about 12 hours to about 20 hours. Completion of the reaction is determined by thin layer chromatographic analysis or other chromatographic methods as are well known to one skilled in the art. When complete, the reaction mixture is diluted with water and the compound of formula (I) or formula (II) wherein R₁ is C₁-C₆alkyl is isolated and purified by methods well known to one skilled in the art.

Alternatively, the nitrogen of the pyrrole ring of a scheme 1 compound of formula (I) or formula (II) is alkylated by treating a pyridine solution of a compound of formula (I) or formula (II), wherein R₁ is H, with a C₁-C₆-alkyl halide in the presence of a suitable base such as for example cesium carbonate with heating for about 0.25 hour to about 3 hours. The reaction mixture is cooled, diluted with water or concentrated to dryness, and extracted with ethyl acetate. Concentration and subsequent purification by chromatographic methods as are well known to one skilled in the art provides the scheme 1 compound of formula (I) or formula (II) wherein R₁ is C₁-C₆-alkyl.

Additionally, N-alkylation of the pyrrole ring nitrogen of a compound formula (I) or formula (II) wherein R₁ is H is achieved by other methods that are well known to one skilled in the art, for example by treatment of a compound formula (I) or formula (II) wherein R₁ is H in a suitable polar solvent such as for example dimethylformamide or NMP, with a suitable base, such as for example sodium hydride or potassium t-butoxide, and then a C₁-C₆alkyl halide such as for example, propyl iodide is added. Completion of the reaction is determined

by thin layer chromatographic analysis or other chromatographic methods well known to one skilled in the art. When complete, the reaction mixture is diluted with water and the scheme 1 compound of formula (I) or formula (II) wherein R_1 is C_1 - C_6 alkyl is isolated and purified by methods well known to one skilled in the art.

5


As is well recognized by one skilled in the art, when M is NR_8 and R_8 is H, under the above described conditions N-alkylation may also occur on the aforesaid NR_8 nitrogen to provide scheme 1 compounds of formula (I) or formula (II) wherein R_1 and R_8 are each the same C_1 - C_6 alkyl group. The preparation of starting esters **1** wherein R is ethyl, R_4 and R_7 is each H, M is NR_8 and R_8 is methyl is known by thermolysis of 2-azidoacrylates (also known as 2-azidopropenoic acid esters) as is also described in scheme 2 below (H. Hemetsberger and D. Knittel, *Monatsh. Chem.* (1972) 103(1), 194-204). Starting ester **1** wherein M is NR_8 and R_8 is C_1 - C_6 alkyl is prepared as described and then converted as described in scheme 1 to a compound of formula (I) wherein M is NR_8 , R_8 is C_1 - C_6 alkyl and R_1 is H or C_1 - C_6 alkyl and wherein said R_1 and R_8 C_1 - C_6 alkyl groups may be the same or different. This methodology is also employed to provide similarly substituted esters **3** that are converted as described in scheme 1 to a compound of formula (II) wherein R_8 is C_1 - C_6 alkyl and R_1 is H or C_1 - C_6 alkyl and wherein said R_1 and R_8 C_1 - C_6 alkyl groups may be the same or different.

20 Additionally a compound of formula (I) or formula (II) of scheme 1 wherein R_1 is H or C_1 - C_6 alkyl and X is S, is optionally oxidized to a sulfone or a sulfoxide wherein X is $S(O)_n$ and n is 1 or 2, respectively, by methods well known to one skilled in the art, such as for example, treating a solution of said compound of formula (I) or formula (II) with H_2O_2 and Na_2CO_3 . Alternatively, compound **2** or **4** of scheme 1 is treated with an arylsulfonyl chloride, an arylsulfinyl chloride, a heterocyclesulfonyl chloride or a heterocyclesulfinyl chloride (used in place of the diaryldisulfide or diheterocycledisulfide) as described in step c above, to provide a scheme 1 compound of formula (I) or formula (II) wherein X is $S(O)_n$, n is 1 or 2 and R_3 is aryl or heterocycle.

30 Scheme 2, as shown below, describes the synthesis of 4H-pyrrolo[2,3-d]thiazoles (M is S), 4-H-pyrrolo[2,3-d]oxazoles, (M is O) and 1,4-dihydro-pyrrolo[2,3-d]imidazoles (M is NR_8) of formula (I) wherein L is N, and the synthesis of 6H-pyrrolo[3,2-d]thiazoles (M is S), 6H-pyrrolo[3,2-d]oxazoles (M is O), and 3,4-dihydro-pyrrolo[2,3-d]imidazoles (M is NR_8) of

formula (II) wherein L is N, from known or commercially available starting materials. One skilled in the art readily understands that when L is N, M is NR₈ and R₈ is H, that the imidazole ring can exist in tautomeric forms. In scheme 2, step a, carboxaldehyde **5** or **7** is condensed with a 2-azidoacetate ester **6** wherein R is alkyl, in the presence of a suitable base such as for example potassium hydroxide, sodium hydroxide or like bases, to provide the corresponding 2-azidopropenoic acid ester **8** or **11**, respectively, wherein R is alkyl.

Scheme 2



As shown in scheme 2, step b, thermolysis of 2-azidopropenoic acid ester **8** or **11** is effected by heating a mixture of 2-azidopropenoic acid ester **8** or **11** in a suitable solvent such as for example xylene at about 120 °C to about 140 °C for about 30 to 90 minutes to provide, after chromatographic purification by methods well known to one skilled in the art, the corresponding ester **9** or **12**, respectively, wherein R is alkyl.

As shown in scheme 2, optional step c, ester **9** or **12** obtained from step b may be hydrolyzed by methods well known to one skilled in the art to provide the corresponding carboxylic acid **9** or **12**, respectively, wherein R is H. For example, a suitable base, such as for example potassium hydroxide, sodium hydroxide, lithium hydroxide and like bases, is added to a mixture of ester **9** or **12** and a suitable solvent, such as for example a mixture of tetrahydrofuran and water. The mixture is heated at about 90 °C to about 110 °C for about 0.5 hour to 2 hours. The product is recovered as a salt by filtration and the filtrate is concentrated to provide additional material as a residue. The filter cake and residue are combined and acidified by methods well known to one skilled in the art, such as for example acidification with a suitable acid such as acetic acid in a suitable solvent such as methanol, ethanol and like solvents, to provide carboxylic acid **9** or **12**, respectively, wherein R is H.

As shown in scheme 2, step d, ester **9** or **12**, wherein R is alkyl, is converted to amide **10** or **13**, respectively, as was described above for scheme 1, step a. Alternatively, carboxylic acid **9** or **12**, wherein R is H, is converted to the corresponding amide **10** or **13**, respectively, by methods as are well known to one skilled in the art and as described in scheme 1, step b. For example, a solution of carboxylic acid **9** or **12** in a suitable solvent such as dimethylformamide is treated with a base such as diisopropylethylamine, a carbodiimide such as for example (1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride, 1-hydroxybenzotriazole and ammonium chloride. When the reaction is complete, the mixture is diluted with a suitable solvent, and the product is isolated and chromatographically purified by methods well known to one skilled in the art to afford the corresponding 4H-pyrrolo[2,3-d]thiazole (M is S), 4-H-pyrrolo[2,3-d]oxazole, (M is O) or 1,4-dihydro-pyrrolo[2,3-d]imidazole (M is NR₈) primary amide **10**, or 6H-pyrrolo[3,2-d]thiazole (M is S), 6H-pyrrolo[3,2-d]oxazole (M is O), or 3,4-dihydro-pyrrolo[2,3-d]imidazole (M is NR₈) primary amide **13**, respectively, wherein R₂ is NH₂. If a primary or secondary C₁-C₆alkylamine is employed in place of ammonium chloride, there is obtained the corresponding secondary or

tertiary amide **10** or **13**, respectively, wherein R_2 is NR_5R_6 , R_5 is H or C_1 - C_6 alkyl and R_6 is C_1 - C_6 alkyl.

As shown in scheme 2, step e, intermediate amide **10** or **13** is thioarylated at the 3-position of the amide bearing pyrrole ring by methods analogous to the methods described above for scheme 1, step c, to provide 4H-pyrrolo[2,3-d]thiazole (M is S), 4-H-pyrrolo[2,3-d]oxazole, (M is O) or 1,4-dihydro-pyrrolo[2,3-d]imidazole (M is NR_8) amide of formula (I), or 6H-pyrrolo[3,2-d]thiazole (M is S), 6H-pyrrolo[3,2-d]oxazole (M is O) or 3,4-dihydro-pyrrolo[2,3-d]imidazole (M is NR_8) amide of formula (II), respectively, wherein X is S and R_3 is aryl or heterocycle.


Wherein R_4 is halogen, such as for example Br, in intermediate amide **10** or **13**, thioarylation at the 3-position of the amide-bearing pyrrole ring and displacement of the aforesaid halogen atom may both occur under the conditions described. Concomitant displacement of the aforesaid halogen from intermediate amide **10** or **13** under conditions described for step e above employing a diaryldisulfide or diheterocycledisulfide is thus advantageously utilized to provide a scheme 2 compound of formula (I) or formula (II), wherein R_4 is an arylthio or a heterocyclethio moiety (that is, XR_3) that is identical to the pyrrole ring XR_3 moiety wherein X is S and R_3 is aryl or heterocycle. Additionally, displacement of the aforesaid halogen atom from intermediate amide **10** or **13**, or from an earlier intermediate such as for example an intermediate ester **9** or **12**, with an anion prepared by treating an arylthiol or heterocyclethiol with a suitable base also advantageously provides a compound of formula (I) or formula (II) wherein R_4 is an arylthio or a heterocyclethio moiety that can be the same or different from the XR_3 moiety introduced by thioarylation as described above for scheme 2, step e. Additionally, displacement of the aforesaid halogen from intermediate ester **9** or **12** or from intermediate amide **10** or **13**, with an anion prepared by methods well known to one skilled in the art from a C_1 - C_6 alkyl-OH, an aryl(C_1 - C_6 alkyl)-OH, a heterocycle(C_1 - C_6 alkyl)-OH, a C_1 - C_6 alkyl-SH, an aryl(C_1 - C_6 alkyl)-SH, a heterocycle(C_1 - C_6 alkyl)-SH, a C_1 - C_6 alkyl-NH₂, a (C_1 - C_6 alkyl)₂NH, or an aryl(C_1 - C_6 alkyl)-amine or a heterocycle(C_1 - C_6 alkyl)amine wherein said amine nitrogen is optionally substituted with C_1 - C_6 alkyl, provides, after thioarylation, a scheme 2 compound of formula (I) or formula (II) wherein R_4 is C_1 - C_6 alkoxy, aryl-(C_1 - C_6 alkoxy), heterocycle-(C_1 - C_6 alkoxy), C_1 - C_6 alkylthio, aryl-(C_1 - C_6 alkylthio), heterocycle-(C_1 - C_6 alkylthio), NR_5R_6 wherein R_5 is H or C_1 - C_6 alkyl and

R_6 is C_1 - C_6 alkyl, or aryl(C_1 - C_6 alkyl)amino or heterocycle(C_1 - C_6 alkyl)amino wherein said amine nitrogen is optionally substituted with C_1 - C_6 alkyl.

As shown in scheme 2, optional step f, the nitrogen of the pyrrole ring of a compound of formula (I) or formula (II), wherein R_1 is H, is N-alkylated by methods as described above for scheme 1, optional step d, to provide a scheme 2 compound of formula (I) or formula (II) wherein R_1 is C_1 - C_6 alkyl.

Additionally a compound of formula (I) or formula (II) of scheme 2 wherein R_1 is H or C_1 - C_6 alkyl and X is S, is optionally oxidized to a sulfone or a sulfoxide wherein X is $S(O)_n$ and n is 1 or 2, respectively, by methods well known to one skilled in the art, such as for example, treating a solution of said compound of formula (I) or formula (II) wherein X is S with H_2O_2 and Na_2CO_3 . Alternatively, compound **10** or **13** of scheme 2 is treated with an arylsulfonyl chloride, an arylsulfinyl chloride, a heterocyclesulfonyl chloride or a heterocyclesulfinyl chloride (used in place of the diaryldisulfide or diheterocycledisulfide) as described in step e above, to provide a scheme 2 compound of formula (I) or formula (II) wherein X is $S(O)_n$, n is 1 or 2 and R_3 is aryl or heterocycle.

Scheme 3



As shown in scheme 3, diaryldisulfides are prepared by treating a solution of an arylsulfide in a suitable organic solvent, such as for example methanol, with an aqueous solution of sodium perborate and allowing the mixture to stand for about 12 hours to about 24 hours at ambient temperature. The diaryldisulfide may be isolated and purified by methods as are well known to one skilled in the art. Diheterocycledisulfides such as for example bis(2-pyridinyl)disulfide are prepared in a similar manner. The arylsulfide and heterocyclesulfide are each optionally substituted as is defined above for "aryl" and "heterocycle".

All of the various embodiments of the compounds of this invention as disclosed herein can be used in the method for treating various diseases and disorders as described herein. As

stated herein the compounds used in the method of this invention are capable of inhibiting the effects of casein kinase I ϵ .

One embodiment of this invention provides a method for treating a mood disorder or a sleep disorder. Another embodiment of the present invention provides a method for treating mood disorder wherein the mood disorder is a depressive disorder or a bipolar disorder. A further embodiment of the present invention provides a method for treating a depressive disorder wherein the depressive disorder is major depressive disorder. Another embodiment of the present invention provides a method for treating mood disorder wherein the mood disorder is bipolar disorder and the bipolar disorder is selected from the group from the group consisting of bipolar I disorder and bipolar II disorder. Another embodiment of the present invention provides a method for treating a sleep disorder. A further embodiment of the present invention provides a method for treating sleep disorder wherein the sleep disorder is a circadian rhythm sleep disorder. A further embodiment of the present invention provides a method for treating circadian rhythm sleep disorder wherein the circadian rhythm sleep disorder is selected from the group consisting of shift work sleep disorder, jet lag syndrome, advanced sleep phase syndrome and delayed sleep phase syndrome. One skilled in the art readily appreciates that the diseases and disorders expressly stated herein are not intended to be limiting but rather to illustrate the efficacy of the compounds of the present invention. Thus, it is to be understood that the compounds of the invention may be used to treat any disease or disorder ameliorated by the inhibition of casein kinase I ϵ .

In another embodiment of the present invention, pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a compound of formula (I) or formula (II), or a stereoisomer, an enantiomer, a racemate or a tautomer of said compound; or a pharmaceutically acceptable salt thereof, are prepared in a manner well known to one skilled in the pharmaceutical arts. The carrier or excipient may be a solid, semisolid or liquid material which can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition may be adapted for oral, inhalation, parenteral or topical use, and may be administered to the patient in the form of tablets, capsules, suspensions, syrups, aerosols, inhalants, suppositories, salves, powders, solutions and the like. As used herein, the term "pharmaceutical carrier" means one or more excipients. As described herein, the pharmaceutical compositions of the invention provide inhibition of casein kinase I ϵ and are thus useful for the treatment of diseases or disorders

ameliorated by inhibition of casein kinase I ϵ

In preparing pharmaceutical compositions or formulations of the compounds of the present invention, care should be taken to ensure bioavailability of an effective therapeutic amount of the active compound or compounds by the selected route of administration, including oral, parenteral and subcutaneous routes. For example, effective routes of administration may include subcutaneous, intravenous, transdermal, intranasal, rectal, vaginal and the like routes including release from implants as well as injection of the active ingredient and/or composition directly into the tissue.

For oral administration, the compounds of the present invention can be formulated into solid or liquid preparations, with or without inert diluents or edible carriers, such as capsules, pills, tablets, troches, powders, solutions, suspensions or emulsions. The capsules, pills, tablets, troches and the like may also contain one or more of the following adjuvants: binders such as microcrystalline cellulose, gum tragacanth; excipients such as starch or lactose, disintegrating agents such as alginic acid, corn starch and the like; lubricants such as stearic acid, magnesium stearate or Sterotex \textregistered , (Stokely-Van Camp Inc., Indianapolis, Indiana) glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; and flavoring agents such as peppermint, methyl salicylate or fruit flavoring. When the dosage unit form is a capsule, it may also contain a liquid carrier such as polyethylene glycol or a fatty oil. Materials used should be pharmaceutically pure and nontoxic in the amounts used. Alternatively, the pharmaceutical compositions may be prepared in a form suitable for extended release to provide a therapeutic amount of a compound of formula (I) of the invention in a suitable once daily, once weekly or once monthly form using methods as are well known to one skilled in the art. For example, an erodable polymer containing the active ingredient may be envisaged.

For parenteral administration, the compounds of the present invention may be administered as injectable dosages of a solution or suspension of the compound in a physiologically acceptable diluent with a pharmaceutical carrier which can be a sterile liquid such as water-in-oil or without the addition of a surfactant and other pharmaceutically acceptable excipients. Illustrative oils which can be employed in the preparations are those of petroleum, animal, vegetable or synthetic origin such as, for example, peanut oil, soybean oil

and mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions, ethanol and glycols, such as propylene glycol are preferred liquid carriers, particularly for injectable solutions. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of inert plastic or glass.

5

The solutions or suspensions described above may also include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents, antibacterial agents such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetra-acetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose.

10

The compounds of the present invention can be administered in the form of a cutaneous patch, a depot injection or implant preparation which can be formulated in such a manner as to permit a sustained release of the active ingredient. The active ingredient can be compressed into pellets or small cylinders and implanted subcutaneously or intramuscularly as depot injections or implants. Implants may employ inert materials such as biodegradable polymers and synthetic silicones. Suitable pharmaceutical carriers and formulation techniques are found in standard texts, such as Remington: The Science and Practice of Pharmacy, 19th edition, Volumes 1 and 2, 1995, Mack Publishing Co., Easton, Pennsylvania, U.S.A., which is herein incorporated by reference.

15

20

In the treatment of various diseases, disorders and conditions as described herein, a suitable dosage level is about 0.01 mg/kg per day to about 250 mg/kg per day, preferably about 0.05 mg/kg per day to about 100 mg/kg per day, and especially about 0.05 mg/kg per day to about 40 mg/kg per day. The compounds of the present invention may be administered on a regimen of 1 to 4 times per day and as dictated by the nature of the disease, disorder or condition to be treated.

25

30

EXAMPLES

The following examples are intended to serve for the illustration of the invention in greater detail, without restricting the breadth of the invention in any manner. Tables 1, 2, 3 and 4 provide summaries of the example compounds that are prepared herein.


Unless otherwise noted, all starting materials, reagents and solvents were obtained from commercial suppliers and used without further purification. All reactions were run under inert atmosphere with dry reagents and solvents. Flash chromatography was carried out using silica gel 60 (35–70 μm) according to the literature procedure (Still, W.C.; Kahn, M; Mitra, A. *J. Org. Chem.* **1978** *43*, 2923) or a variation of this method using a commercially available silica gel cartridge (for example Isco Redi Sep). Thin layer chromatography (TLC) was performed on glass-backed, silica gel 60F–254 plates (EM) coated to a thickness of 0.25 mm. The plates were eluted with solvent systems (v/v) as described, and visualized by iodine vapor, UV light, or a staining reagent such as KMnO_4 solution.

^1H NMR spectra were recorded on a Varian Gemini 300, Unity 300, Unity 400, or Unity 500 spectrometers with chemical shifts (δ) reported in ppm relative to tetramethylsilane (0.00 ppm) or chloroform (7.26 ppm) as a reference. Liquid chromatography with mass spectral analysis (LCMS) was recorded on a Micromass LCTAPI LC-TOF (time of flight) Mass Spectrometer and Masslynx Data System. Ionization mode = electrospray (esi), values were determined for the protonated molecular ions ($\text{M}^+ + 1$) using a Synergi 2U HYDRO-RP 20x4 mm column, eluting with 0.1% TFA in water/acetonitrile.

4H-Thieno[3,2-b]pyrrole-5-carboxylic acid ethyl ester and 6H-thieno[2,3-b]pyrrole-5-carboxylic acid ethyl ester were prepared as described in Eras, J.; Galvez, C.; Garcia, F. *Journal of Heterocyclic Chemistry* (1984), 21(1), 215-17. Ethyl esters of 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid, 6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid, and 2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid were prepared in the same fashion as described in WO9940914. 2-Alkylthio-, 2-arylalkylthio- and 2-alkyl-substituted pyrrolo[2,3-d]imidazole-5-carboxylic acid esters can be prepared as described in Shafiee, A. and Hadizadeh, F., *J. of Heterocyclic Chemistry* (1997), 34, 549-550 and in Shafiee, A.; Shahbazi Mojarad, J.; Jalili, M.A.; Adhami, H.R. and Hadizadeh, F. *Journal of Heterocyclic Chemistry*, 39, 367-373. 4-Thiazolecarboxaldehyde, 5-thiazolecarboxaldehyde and 2-methyl-5-thiazolecarboxaldehyde were commercially obtained. 1,4-Dihydro-4-methylpyrrolo[3,2-b]pyrrole-2-carboxylic acid ethyl ester is prepared as described by H. Hemetsberger and D. Knittel, *Monatsh. Chem.* (1972) 103(1), 194-204.

Preparation of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester


2-Azido-3-(2-bromo-4-thiazolyl)propenoic acid ethyl ester



5 Add slowly to a solution of potassium ethoxide (30 ml, 24% w:w, 3 eq. of EtOK) a slurry of 2-bromo-4-thiazolylcarboxyaldehyde (3.87 gm, 30 mol) and 2-azidoacetate ethyl ester (11.5 gm, 3 esq.) in a mixed solvent of ethanol (150ml) DMF (5 ml) and methylene chloride (DCM, 20 ml) at 0 °C over 15-20 minutes. Stir the final mixture overnight (18 hr) at room temperature, quench the reaction with ammonium chloride and remove the ethanol (~50
10 ml) on a rotary evaporator. Extract the aqueous mixture with DCM (3x250ml portions), wash the organic phase with brine and dry over MgSO₄. Filter and concentrate the filtrate, and purify the crude mixture (12.2 gm) by flash chromatography [ISCO, SiO₂, 120 gm cartridge, elute with methanol: DCM (0-5%)] to afford the title compound (3.6 gm, 45%).


LCMS: retention time = 3.68 min, (M⁺) = 302.98

15 2-Bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester




Add dropwise to hot xylene (130 °C, 4 ml) a solution of 2-azido-3-(2-bromo-4-thiazolyl)propenoic acid ethyl ester (60 mg, 0.2 mmol) in DCM (1ml). Heat the mixture for
20 one hour, then cool to room temperature, deposit the mixture on a pad of silica gel and elute with heptane:DCM (50%-100%) to provide the title compound (14 mg).

LCMS: retention time = 3.04 min, (M⁺) = 274.92.

Preparation of 4H-Pyrrolo[2,3-d]thiazole-5-carboxylic acid ethyl ester

Prepare 4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid ethyl ester from 4-thiazolylcarboxyaldehyde in a similar manner as described above for the preparation of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester from 2-bromo-4-thiazolylcarboxyaldehyde.

5


Preparation of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid

Add KOH (1.07 gm, 2 eq) to a mixture of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester (2.6 gm, 9.38 mmol) in THF (15 ml) and water (20 ml), and then heat at 100 °C for 1 hr. Allow to stand overnight at room temperature and collect the crystalline solids by filtration (weight 1.7 gm). Concentrate the aqueous solution in vacuo, and combine the residue with the previously isolated crystalline solid. Acidify with acetic acid in methanol to provide the title compound (2.31 g).

10

LCMS: retention time = 2.35 min, (M^+) = 246.93

15

Preparation of 6H-Pyrrolo[3,2-d]thiazole-5-carboxylic acid


Prepare 6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid by hydrolysis of 6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester in a similar manner as described above for the preparation of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid from 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid ethyl ester.

20

Preparation of Carboxylic Acid Amide Intermediates

4H-Thieno[3,2-b]pyrrole-5-carboxylic acid amide

25




Add to 4H-thieno[3,2-b]pyrrole-5-carboxylic acid ethyl ester (1.74 gm, 8.9 mmol) and 7M ammonia in methanol (100ml) in a steel bomb a chip of lithium hydroxide (0.1gm). Seal the bomb and heat to 100 °C for 16 hrs. Cool to room temperature, and concentrate to remove the volatiles. Purify the crude product via flash chromatography (ISCO, silica cartridge, 40gm, elute with methanol 0-5% in methylene chloride) to provide the title compound (560 mg, 38%) as an off-white solid.

LCMS: retention time = 2.08 min, (M^+) = 166.02


^1H NMR (300 MHz, DMSO- D_6) δ ppm 6.95 (d, $J=5.25$ Hz, 1 H) 7.05 - 7.08 (m, 1 H) 7.11 (br s, 1 H) 7.37 (d, $J=5.25$ Hz, 1 H) 7.68 (br s, 1 H) 11.64 (s, 1 H)

The following amides were also prepared by the above procedure:


6H-Pyrrolo[3,2-d]thiazole-5-carboxylic acid amide (LCMS: retention time = 1.63 min, (M^+ + H) = 168.00)



2-Methyl-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide (LCMS: retention time = 1.28 min, (M^+ + H) = 182)



2-Bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide




Add to a solution of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid (2.53 gm, 10 mmol) in DMF (45 ml), DIEA (diisopropylethylamine, 10 ml, 6 eq), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide HCl, 5.0 gm, 3.5 eq); HOBT (1-hydroxybenzotriazole, 1.91 gm, 14 mmol, 1.4 eq) and NH_4Cl (2.25 gm, 42 mmol). Stir the mixture at room temp for 6 hr and monitor by LC-MS. When complete, dilute the mixture

with ethyl acetate, and wash with water and brine. Collect the solid by filtration (2.21 gm) and purify by chromatography on silica gel (ISCO silica cartridge, 4gm, elute with methanol (10-40%) in methylene chloride) to afford the title compound (550 mg).

LCMS: retention time = 2.11 min, (M^+) = 245.98

5

6H-Pyrrolo[3,2-d]thiazole-5-carboxylic acid amide



Prepare 6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide by amination of 6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid in a similar manner as described above for the preparation of 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide from 2-bromo-6H-pyrrolo[3,2-d]thiazole-5-carboxylic acid.

10


General preparation of diaryldisulfides and diheterocycledisulfides

Add to a solution of the unsubstituted or appropriately substituted arylthiol (17.2 millimole, 1.0 equivalent) and MeOH (30mL), a solution of sodium perborate (22 millimole) and water (20mL) with stirring, and then allow the reaction to stand at rt overnight. Collect the solid by filtration and wash with methanol to give the desired diaryldisulfide. Other disulfides including diheterocycledisulfides (e.g. bis(2-thienyl)disulfide) can be prepared in a similar manner as described for the preparation of the desired diaryldisulfides.

20

Methods for Thioarylation of the Pyrrole Moiety

Method 1: 6-Phenylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide (Ia)



Treat 4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide (75 mg, 0.45 mmol) with NaH (45 mg, 60% in oil, 1.12mmol, 2.5 eq) in N,N-dimethylformamide (1.3 ml) at room temperature under nitrogen for 35 minutes. Add diphenyldisulfide (137 mg, 1.4 eq) and heat


25

the mixture at 70 °C overnight. Dilute the mixture with brine (2 ml) and extract with ethyl acetate. Concentrate the ethyl acetate solution to give an oil and purify by chromatography on silica gel (ISCO silica cartridge, 4 gm, elute with methanol 0-10% in methylene chloride) to afford the title compound (55mg).

5 LCMS: retention time = 3.03 min, ($M^+ + H$) = 275.01

1H NMR (300 MHz, $CDCl_3$) δ ppm 5.94 (s, 1 H) 7.03 (d, $J=5.25$ Hz, 1 H) 7.14 - 7.27 (m, 6 H) 7.79 (br s, 1 H) 10.79 (br s, 1 H)

10 **Method 2: 6-(3-Fluorophenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide (Ib)**




Add 4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide (60 mg, 0.36 mmol) to a mixture of bis(3-fluorophenyl)disulfide, (150 mg, 0.51mmol) and cesium carbonate (120 mg, 1 eq.) in DMF (2.5 ml), and then heat at 95 °C for 3 hrs. Follow the reaction by TLC. When
15 complete, dilute the reaction mixture with ethyl acetate (15ml) and wash with brine (25ml). Dry the organic solution, concentrate to afford a crude oil and purify the oil by chromatography on silica gel (ISCO silica cartridge, 4 gm, elute with methanol 0-10% in methylene chloride) to provide the title compound (81mg).

LCMS: retention time = 3.47 min, ($M^+ + H$) = 293

20 1H NMR (300 MHz, $CDCl_3$) δ ppm 5.67 (s, 1 H), 6.82 - 6.90 (m, 2 H), 6.96 (ddd, $J=7.87$, 1.50, 1.37 Hz, 1 H), 7.03 (d, $J=5.25$ Hz, 1 H), 7.17 - 7.24 (m, 1 H), 7.31 (d, $J=5.25$ Hz, 1 H), 7.70 (s, 1 H), 9.96 (s, 1 H)


25 **Method 3: 4-(Pyridin-2-ylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide (IIa)**



Treat 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide (57 mg, 0.34 mmol) with NaH (19 mg, 0.78 mmol, 2.3 eq) in N,N-dimethylformamide (1ml) at room temperature, under nitrogen for 45 minutes. Add 2,2'-dipyridyldisulfide (106 mg, 1.4 eq) and stir the mixture overnight at ambient temperature. Dilute the mixture with water and extract with ethyl acetate. Concentrate the ethyl acetate solution to give a residue which is purified by flash chromatography (ISCO, silica cartridge, elute with 5% methanol in methylene chloride (+1% 7N ammonia in methanol) to provide the title compound (32 mg, 32%).

LCMS: retention time = 2.53 min, ($M^+ + H$) = 276.022

Method 4: 4-(Phenylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide (IIb)




Treat 6H-thieno[2,3-b] pyrrole-5-carboxylic acid amide (50 mg, 0.30 mmol) in N,N-dimethylformamide (0.5ml) with NaH (29 mg, 60% in oil, 0.75mmol, 2.5 eq) at room temperature, under nitrogen for 45 minutes. Add diphenyl disulfide (92 mg, 0.4 mmol 1.4 eq) and stir the mixture at 60 °C overnight. Increase the temperature to 100 °C for 5 hours and cool to room temperature. Dilute the reaction with water and ethyl acetate* whereupon the title compound crystallizes from solution. Collect the product by filtration and dry under vacuum, to afford the title compound (28 mg).

LCMS: retention time = 3.068 min, ($M^+ + H$) = 275.024

*In some cases (see tables 1 and 2, synthesis method column) where method 4 was employed, the compounds did not crystallize. In this situation, separate the ethyl acetate portion and

concentrate to give a crude residue and purify the residue by flash chromatography (ISCO, silica cartridge, eluted with 10% methanol in methylene chloride (+1% 7N ammonia in methanol) to provide the desired compound.

Method 5: 2,6-Bis-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide (IIq)




Add diphenyldisulfide (93 mg, 0.43 mmol, 1.25 eq) to a mixture of 2-bromo-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide (85 mg, 0.34 mmol) and cesium carbonate (140 mg, 1.25 eq) in DMF (4ml). Heat the mixture at 100 °C for 16 hours. Remove the DMF under reduced pressure and partition the residue between water and ethyl acetate. Wash the organic phase with brine, and then dry and filter the organic phase. Add the filtrate to a flask containing a small amount of silica gel (about 0.5 gm) and evaporate the solvent to provide the crude product adsorbed onto the silica gel. Place the silica gel on top of a small column containing about 4 gm of silica gel and elute with 0-10% of MeOH in DCM to provide 8 mg (6.5%) of the title compound.

LCMS: retention time = 3.30 min, (M^+) = 383.02

Table 1


Compounds of Formula (I)
(M is S)



Example No.	R ₁	R ₂	R ₃ X	R ₄	L	Synthesis Method
Ia	H	NH ₂	C ₆ H ₅ S	H	CH	1
Ib	H	NH ₂	3-F-C ₆ H ₄ S	H	CH	1
Ic	H	NH ₂	4-Cl-C ₆ H ₄ S	H	CH	1
Id	H	NH ₂	2-NH ₂ -C ₆ H ₄ S	H	CH	1
Ie	H	NH ₂	2-pyridinylS	H	CH	1
If	H	NH ₂	4-CH ₃ -C ₆ H ₄ S	H	CH	1
Ig	H	NH ₂	2-thienylS	H	CH	2
Ih	H	NH ₂	3,5-(Cl) ₂ -C ₆ H ₃ S	H	CH	2
Ii	H	NH ₂	4-pyridinylS	H	CH	2
Ij	H	NH ₂	3-CH ₃ -C ₆ H ₄ S	H	CH	2
Ik	H	NH ₂	2-CH ₃ -C ₆ H ₄ S	H	CH	2
Il	H	NH ₂	2,3-(Cl) ₂ -C ₆ H ₃ S	H	CH	2
Im	H	NH ₂	2,5-(Cl) ₂ -C ₆ H ₃ S	H	CH	2
In	H	NH ₂	2-C ₂ H ₅ -C ₆ H ₄ S	H	CH	2
Io	H	NH ₂	3-Br-C ₆ H ₄ S	H	CH	2
Ip	H	NH ₂	3,5-(CH ₃) ₂ -C ₆ H ₃ S	H	CH	2
Iq	H	NH ₂	3-CH ₃ O-C ₆ H ₄ S	H	CH	2
Ir	H	NH ₂	2-CH ₃ O-C ₆ H ₄ S	H	CH	
Is	H	NH ₂	2-CF ₃ -C ₆ H ₄ S	H	CH	2
It	H	NH ₂	2-F-C ₆ H ₄ S	H	CH	2
Iu	H	NH ₂	3-CF ₃ O-C ₆ H ₄ S	H	CH	2
Iv	H	NH ₂	C ₆ H ₅ S	H	N	4
Iw	H	NH ₂	3-F-C ₆ H ₄ S	H	N	4*
Ix	H	NH ₂	2-pyridinylS	H	N	4

*See footnote accompanying method 4.

Table 2
Compounds of Formula (II)
(M is S)



Example No.	R ₁	R ₂	R ₃ X	R ₄	L	Synthesis Method
Ila	H	NH ₂	2-pyridinylS	H	CH	3
Ilb	H	NH ₂	C ₆ H ₅ S	H	CH	4
Ilc	H	NH ₂	3-F-C ₆ H ₄ S	H	CH	4*

IId	H	NH ₂	4-pyridinylS	H	CH	1
Ile	H	NH ₂	3,5-(Cl) ₂ -C ₆ H ₃ S	H	CH	1
IIf	H	NH ₂	2-thienylS	H	CH	1
Ilg	H	NH ₂	3-Br-C ₆ H ₄ S	H	CH	2
Iih	H	NH ₂	3-CH ₃ O-C ₆ H ₄ S	H	CH	2
Ili	H	NH ₂	2-CH ₃ O-C ₆ H ₄ S	H	CH	2
Ilj	H	NH ₂	3-Cl-C ₆ H ₄ S	H	CH	2
Ilk	H	NH ₂	3-CH ₃ -C ₆ H ₄ S	H	CH	2
III	H	NH ₂	C ₆ H ₅ S	CH ₃	N	2
IIm	H	NH ₂	3-CH ₃ O-C ₆ H ₄ S	CH ₃	N	2
IIn	H	NH ₂	3-F-C ₆ H ₄ S	CH ₃	N	2
Ilo	H	NH ₂	3-Cl-C ₆ H ₄ S	CH ₃	N	2
Ilp	H	NH ₂	3-CF ₃ O-C ₆ H ₄ S	CH ₃	N	2
IIq	H	NH ₂	C ₆ H ₅ S	C ₆ H ₅ S	N	5
Ilr	H	NH ₂	3-CH ₃ O-C ₆ H ₄ S	3-CH ₃ O-C ₆ H ₄ S	N	5
IIs	H	NH ₂	C ₆ H ₅ S	H	N	2
IIt	H	NH ₂	3-CH ₃ O-C ₆ H ₄ S	H	N	2

*See footnote accompanying method 4.

Table 3
Spectral Data for Compounds of Formula (I)
(M is S)

Example No.	Compound Name	MS: Obs. Ion Mass* (amu)	MS: Retention time (min)
Ia	6-Phenylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	275.011	3.034
Ib	6-(3-Fluorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	293	3.47
Ic	6-(4-Chlorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	308.967	3.268
Id	6-(2-Aminophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	290.025	2.668
Ie	6-(Pyridin-2-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	276	2.57
If	6-p-Tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	289.024	3.218
Ig	6-(Thiophen-2-yl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	281.02	2.97
Ih	6-(3,5-Dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid	342.96	3.55

	amide		
Ii	6-(Pyridin-4-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	276	1.39
Ij	6-m-Tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	287	2.71
Ik	6-o-Tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	289	2.69
Il	6-(2,3-Dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	342.95	3.57
Im	6-(2,5-Dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	342.95	3.5
In	6-(2-Ethylphenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	303.06	3.45
Io	6-(3-Bromo-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	352.932	3.4
Ip	6-(3,5-Dimethylphenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	303.05	3.47
Iq	6-(3-Methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	305.03	3.17
Ir	6-(2-Methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	305.36	3.13
Is	6-(2-Trifluoromethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	343.01	3.37
It	6-(2-Fluoro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	293.02	3.15
Iu	6-(3-Trifluoromethoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide	359.008	3.47
Iv	6-Phenylsulfanyl-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide	274	3.052
Iw	6-(3-Fluoro-phenylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide	294.013	2.851
Ix	6-(Pyridin-2-ylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide	277.035	2.334

*Ion type is M+H unless otherwise noted.

Table 4
Spectral Data for Compounds of Formula (II)
(M is S)

Example No.	Compound Name NMR (CDCl ₃)	MS: Obs. Ion Mass* (amu)	MS: Retention time (min)
Ila	4-(pyridin-2-ylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	276.022	2.53
Ilb	4-(phenylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	275.024	3.068
Ilc	6-(3-Fluorophenyl-sulfanyl)- 6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	276	3.43
Ild	4-(Pyridin-4-ylsulfanyl)-6H-thieno[2,3-b]-pyrrole-5-carboxylic acid amide	276.1	1.45
Ile	4-(3,5-Dichlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	325.9	2.98
Ilf	4-(Thiophen-2-yl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	264	2.43
Ilg	4-(3-Bromophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	352.9	4.31
Ilh	4-(3-Methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	305	3.44
Ili	4-(2-Methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	305	3.56
Ilj	4-(3-Chlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide 1H NMR (300 MHz,) δ ppm 2.30 (s, 3 H), 5.70 (br s, 1 H), 6.89 (d, J=5.25 Hz, 1 H), 6.94 - 7.06 (m, 4 H), 7.14 (t, J=7.62 Hz, 1 H), 7.87 (br s, 1 H), 10.28 (br s, 1 H)	309.2	3.47
Ilk	4-(3-Methylphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide	289	4.1
III	2-Methyl-6-phenyl-sulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	290.04	2.8
IIm	6-(3-Methoxyphenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide 1H NMR (300 MHz,) δ ppm 2.75 (s, 3 H),	320.05	2.69

	3.71 (s, 3 H), 5.69 (br s, 1 H), 6.64 - 6.68 (m, 1 H), 6.72 - 6.77 (m, 2 H), 7.12 (t, J=7.66 Hz, 1 H), 7.86 (br s, 1 H), 10.35 (br s, 1 H)		
IIn	6-(3-Fluorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	308.04	2.72
Ilo	6-(3-Chlorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide 1H NMR (300 MHz,) δ ppm 2.77 (s, 3 H), 5.72 (br s, 1 H), 7.03 - 7.18 (m, 4 H), 7.76 (br s, 1 H), 10.41 (br s, 1 H)	323.99	2.82
Ilp	6-(3-Trifluoromethoxy-phenylsulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	374.02	3
Iiq	2,6-Bis-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	384.05	3.3
Iir	2,6-Bis-(3-methoxy-phenylsulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	444.08	3.33
Iis	6-Phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide	275.98	2.77
Iit	6-(3-Methoxyphenyl-sulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide 1H NMR (300 MHz,) δ ppm 3.71 (s, 3 H), 5.83 (br s, 1 H), 6.66 - 6.70 (m, 1 H), 6.77 - 6.81 (m, 2 H), 7.13 (t, J=8.00 Hz, 1 H), 7.96 (br s, 1 H), 8.55 (br s, 1 H), 10.63 (br s, 1 H)	306	2.77

*Ion type is M+H unless otherwise noted. For Iic, Iie, Iif observed ion type was (M+H)-NH₃

Biological Examples

Casein Kinase Epsilon ³³P-ATP Filter Plate Assay for Screening CK1 ϵ Inhibitors

Purpose: This assay measures the ability of compounds to inhibit the phosphorylation of the substrate casein by the enzyme casein kinase 1 ϵ using an *in vitro* ³³P-ATP filtration assay. Compounds are tested at five concentrations in duplicate in order to generate IC₅₀ values or % inhibition at a 10 micromolar concentration that are summarized in Table 4.

Materials:

Equipment:

Beckman Biomek 2000 Liquid Handling Robot
Beckman Multimek 96 Automated 96 Channel Pipettor
Millipore Vacuum Manifold Basic Kit # MAVM0960R
Titertek Multidrop Liquid Dispenser
5 Packard TopCount NXT Liquid Scintillation Counter

Plates:

Costar EIA/RIA Plate #9018
Falcon 96 well U bottom Polystyrene Plate #353910
Millipore Multiscreen 96 well Filtration Plates #MAPHNOB50
10 Millipore Multiscreen TopCount Adapter Plates #SE3M203V6

Chemicals:

EGTA from SIGMA #E-3889
Casein (dephosphorylated) from SIGMA #C-4032
ATP from SIGMA #A-7699
15 DTT from Fisher Biotech #BP1725
Trichloroacetic Acid from SIGMA #T-6399
 γ -³³P-ATP 1mCi / 37MBq from Perkin Elmer Life Sciences #NEG-602H

Enzyme:

Casein Kinase 1 ϵ final concentration 0.58mg/ml obtained from fermentation and
20 purification processes as are well known to one skilled in the art. The above are
stored as 100 μ L aliquots at minus 80°C.

Compounds:

Supply compounds for testing as frozen 10mM compound stock dissolved in 100%
DMSO.

Assay Conditions:

Final total assay volume per well is equal to 50 μ L that one prepares as follows:

5 μ L of diluted compound stock (10, 1, 0.1, 0.01 or 0.001 μ M),
5 μ L of dephosphorylated casein final concentration 0.2 μ g/ μ L,
30 20 μ L of CK1 ϵ final concentration 3 ng/ μ L, and
20 μ L of γ -³³P-ATP final concentration 0.02 μ Ci/ μ L mixed with cold ATP (10 μ M final).

Methodology:

1. Prepare 500 mL of fresh assay buffer: 50mM Tris pH 7.5, 10mM MgCl₂, 2 mM DTT and 1 mM EGTA
2. Obtain compounds to be evaluated as 10μL of 10mM stock dissolved in 100% DMSO. Use a Biomek 2000 liquid handling robot, make serial dilutions to yield 10, 1, 0.1, 0.01 and 0.001 μM final compound dilutions added as 5μL additions to Falcon U bottom plates. Typically test 8 compounds per 96 well plate with column 1 and 12 serving as control wells. A routine screening assay will consist of 32 compounds, which equals 4 assay plates.
3. Assay plate maps are set up according to the following pattern CK1εPlateMap.xls

CK1ε Filtration Assay Plate Map												
	1	2	3	4	5	6	7	8	9	10	11	12
A	<div>TOTAL</div>	Compound 1					Compound 5					
B		Compound 1					Compound 5					
C		Compound 2					Compound 6					
D		Compound 2					Compound 6					NonSp.
E		Compound 3					Compound 7					
F		Compound 3					Compound 7					
G		Compound 4					Compound 8					
H		Compound 4					Compound 8					
<div><div><div>1010.10.010.001uM</div><div><div>TOTAL</div><div>Enzyme+Casein+Buffer+33P-ATP</div><div>Test</div></div><div><div>Nonsp.</div><div>Casein+Buffer+33P-ATP</div></div></div><div><div>1010.10.010.001uM</div></div></div>												

4. Add 5 μL of compound as indicated, then add 5μL of dephosphorylated casein (dissolved in distilled H₂O)(0.2μg/μL) and 20μL CK1ε (3ng/μL) to the appropriate wells.
5. Finally add 20μL γ-³³P-ATP (0.02μCi/μL) /10μM cold ATP (equals approximately 2x10⁶ CPM per well).
6. Vortex the Falcon U-Bottom assay plate containing the above 50μL reaction volume and then incubate at room temperature for 2 hours.
7. At the end of 2 hours, stop the reaction by the addition of 65μL of ice cold 2mM cold ATP (made up in assay buffer) to the assay plates using a Beckman Multimek.
8. At the same time add 25 μL 100% ice cold TCA made up in distilled H₂O to a matching number of Millipore MAPH filter plates.
9. Using a handheld 8-channel pipettor, transfer 100μL of the reaction mixture from the Falcon U-Bottom Plate to the Millipore MAPH filter plates presoaked with TCA.

10. Mix the Millipore MAPH filter plates gently and allow to sit at room temperature for at least 30 minutes to precipitate the proteins.
11. After 30 minutes, place the filter plates on a Millipore vacuum manifold and filter at no more than 8mm Hg as the MAPH filters tend to “air lock” at higher vacuum settings.
12. Wash the filter plates sequentially and filter with 2x150 μ L 20% TCA, 2x150 μ L 10% TCA and 2x150 μ L 5% TCA (total of 6 washes per plate/900 μ L per well).
13. Allow the plates to dry overnight at room temperature. The next day add 40 μ L Packard Microscint-20 Scintillation Fluid per well using a Titertek Multidrop dispenser; seal the plates and count for 2 minutes/well in a Packard Topcount NXT Scintillation Counter (to provide CPM values/well).

Calculation:

1. Import Counts Per Minute (CPM) data into a proprietary data calculation and archiving database (Activity Base by IDBS version 5.0).
2. Column 1 for each plate reflects total phosphorylation activity of the enzyme in the absence of any inhibiting compound and thus represents 100%. Column 12 reflects any nonspecific phosphorylation/retained radioactivity activity in the absence of inhibiting compound and enzyme. Typically one observes approximately 1% of Total CPMs that are “nonspecific”.
3. By determining the “total” and “nonspecific” CPMs for each plate, one is able to determine the % inhibition of the enzyme’s ability to phosphorylate the substrate for each concentration of test compound. Use this % inhibition data to calculate an IC₅₀ value (concentration at which a compound is able to inhibit the enzyme activity by 50%) for a compound using a non-linear curve fit program contained with the Activitybase calculation protocol (DG0027-CK1-D-BL).
4. Kinetic studies have determined the K_m value for ATP to be 21 μ M in this assay system.

Casein Kinase 1 δ Streptavidin Affinity Membrane Plate Assay for CKI δ inhibitors

Purpose: To evaluate test compounds for CKI δ activity in Streptavidin Affinity Membrane (SAM) Biotin Capture Plate (Promega V7542)

Supplies and reagents

HEPES Sigma # H3375 MW = 238.3; β -Glycerol phosphate Sigma # G-9891 MW = 216.0; EDTA 0.5M, pH 8.0 GibcoBRL; Sodium orthovanadate ACROS # 205330500 MW = 183.9; DTT (DL-dithiothreitol) Sigma # D-5545 MW = 154.2; Magnesium Chloride ACROS # 41341-5000 MW = 203.3; ATP Sigma # A-7699 MW = 551.1; γ - ^{33}P ATP NEN # NEG602H; Casein Kinase 1 δ Sigma # C4455; Casein Kinase 1 substrate New England Peptide Biotin-RRKDLHDDEEDEAMSITA MW = 2470

Prepare Kinase Buffer (KB, 100 mL) as follows:

10	50 mM HEPES, pH 8.0	5 mL of 1M stock
	10 mM MgCl	1 mL of 1M stock
	10 mM β -glycerophosphate	1 mL of 1M stock
	2.5 mM EDTA	500 μL of 500 mM stock
	1 mM sodium orthovanadate	100 μL of 1M stock
15	1 mM DTT	100 μL of 1 M stock
	water	92.3 mL

Prepare ATP Master Mix as follows:

Prepare 1mL of a 1M ATP solution in water (1M ATP stock).

20 To 12 mL KB:

Add 12 μL of 1M ATP solution, then

Add 12 μL of ^{33}P ATP (10 $\mu\text{Ci}/\mu\text{L}$), NEG602H, Perkin Elmer

Prepare the reaction plate and conduct the assay as follows:

- 25
1. Add 10 μL of KB per well with or without the test compound inhibitor to reaction plate wells
 2. Add 60 μL of KB per well
 3. Add 10 μL of 500 μM Peptide Substrate per well
 4. Bring plate up to 37°C
 - 30 5. Add 10 μL of 1:10 dilution of CK1 δ per well = 0.42 μg or 0.68 units
 6. Initiate the reaction with 10 μL of ATP Master Mix per well
 7. Place the reaction plate in 37°C incubator for 10 min.

8. Stop the reaction with 10 μ L of 1M ATP. Transfer 20 μ L to the SAM Plate and let stand 10 min at room temperature.
9. Wash three times with 100 μ L of 2M NaCl solution, then three times with 100 μ L of 2M NaCl and 1% H_3PO_4 solutions and then three times with 100 μ L of water on a vacuum manifold.
10. Dry the filter plate under a lamp for 30 min.
11. Seal bottom of plate and add 20 μ L of MicroScint 20
12. Read in TOPCOUNT

Cellular Circadian Assay Experimental Procedures

Cell culture: Split *Mper1-luc* Rat-1 fibroblasts (P2C4) cultures every 3-4 days (~10-20% confluence) onto 150 cm² vented polystyrene tissue culture flasks (Falcon # 35-5001) and maintain in growth media [EMEM (Cellgro #10-010-CV); 10% fetal bovine serum (FBS; Gibco #16000-044); and 50 I.U./mL penicillin-streptomycin (Cellgro #30-001-C1)] at 37°C and 5% CO₂.

Stable transfection: Co-transfect Rat-1 fibroblast cultures at 30-50% confluence with vectors containing the Zeocin resistance selectable marker for stable transfection and an mPer-1 promoter-driven luciferase reporter gene. After 24-48 hours, split the cultures onto 96 well plates and maintain in growth media supplemented with 50-100 μ g/mL Zeocin (Invitrogen #45-0430) for 10-14 days. Assess Zeocin-resistant stable transfectants for reporter expression by supplementing growth media with 100 μ M luciferin (Promega #E1603) and assaying luciferase activity on a TopCount scintillation counter (Packard Model #C384V00).

Synchronize Rat-1 clones expressing both Zeocin-resistance and *mPer1*-driven luciferase activity by 50% horse serum [HS (Gibco #16050-122)] serum shock and assess for circadian reporter activity. Select *Mper1-luc* Rat-1 fibroblasts clone P2C4 for compound testing.

Synchronization protocol: Plate *Mper1-luc* Rat-1 fibroblasts (P2C4) (40-50% confluence) onto opaque 96-well tissue culture plates (PerkinElmer #6005680) and maintain in growth media supplemented with 100 μ g/mL Zeocin (Invitrogen #45-0430) until cultures reach 100% confluence (48-72 h). Synchronize cultures with 100 μ L synchronization media [EMEM (Cellgro #10-010-CV); 100 I.U./mL penicillin-streptomycin (Cellgro #30-001-C1); 50% HS (Gibco #16050-122)] for 2 hours at 37°C and 5% CO₂. After synchronization, rinse cultures with 100 μ L EMEM (Cellgro #10-010-CV) for 10 minutes at room temperature. After rinse, replace media with 300 μ L CO₂-independent media [CO2I (Gibco #18045-088); 2mM L-

glutamine (Cellgro #25-005-C1); 100 I.U./mL penicillin-streptomycin (Cellgro #30-001-C1); 100 μ M luciferin (Promega #E1603)]. Add compounds to be tested for circadian effects to CO₂-independent media in 0.3% DMSO (final concentration). Seal cultures immediately with TopSeal-A film (Packard #6005185) and transfer for luciferase activity measurement.

5 **Automated Circadian Reporter Measurement:** After synchronization, maintain assay plates at 37°C in a tissue culture incubator (Forma Scientific Model #3914). Estimate in vivo luciferase activity by measuring relative light output on a TopCount scintillation counter (Packard Model #C384V00). Transfer plates from incubator to reader using an ORCA robotic arm (Beckman Instruments) and SAMI-NT automated scheduling software (Version 3.3; 10 SAGIAN/Beckman Instruments).

Data Analysis: Use Microsoft Excel and XLfit (Version 2.0.9; IDBS) to import, manipulate and graph data. Perform period analysis either by determining the interval between relative light output minima over several days or by Fourier Transform. Both methods produce nearly identical period estimation over a range of circadian periods. Report potency as EC_{Δt+1h}, 15 which is the effective micromolar concentration that induces a 1 hour lengthening of period. Analyze the data by fitting a hyperbolic curve to the data expressed as period change (y-axis) versus the concentration of test compound (x-axis) in XLfit and interpolate the EC_{Δt+1h} from this curve.

20 **Rat Circadian Cycle Assay**

This assay provides a means for assessing the effect of a test compound on circadian cycle in vivo. Use male Wistar rats (Charles River) with a starting body mass of 200-250 g. House each animal individually prior to testing in a controlled environment and maintain a thermoneutral ambient temperature of 24-28 °C under a 12/12 hour (h) light/dark cycle (lights 25 on at 06:00 h), and give standard laboratory chow and water ad libitum. Implant each rat with an intra-abdominal biotlemetry transmitter (Minnimitter-VMFH, series 4000, Sunriver, OR) to monitor core body temperature and general activity. Implant each transmitter as per the manufacturer's recommendations under ketamine/xylazine (78/13 mg kg⁻¹, ip) general anesthesia and allow the animals to recover for 7-10 days. After the recovery period, to 30 establish each animal's internal circadian cycle, place the animals in a constant dark cycle (0/24 h light/dark cycle) and allow the animals to go into free run for 7-10 days prior to test compound administration. During the dosing regimen, animals receive either vehicle or compound (ip, sc, or po) at specific CTs (Circadian Times) over a 48 hour period. Monitor

the animals for 5 to 7 days in a constant dark cycle (0/24 h light/dark cycle) after completion of the dosing regimen. For each experiment, sample abdominal temperature and general activity data at 5-minute intervals. For analysis, use the VitalView and Actiview software supplied by Minimitter. Plot observed abdominal temperatures obtained for each rat on the first day on a horizontal line. Align the line of observed abdominal temperatures below an abscissa line with circadian time (x-axis). Plot observed abdominal temperatures for each successive day as individual lines in a similar manner to provide the ordinate (y-axis, in days). Connect the initial rise of core body temperature that occurs each day with a straight line, which allows the use of multiple days to estimate the circadian phase on any given day for each individual rat. Determine the effect of treatment on phase by using the straight line multi-day estimation of phase before and after dosing. Treatment with an active compound will cause a greater displacement between the straight line connecting the daily initial rise of core body temperature before compound treatment and the straight line connecting the initial rise of core body temperature after compound treatment versus the vehicle control before and after treatment lines. Calculate the difference between those phases projected onto the day prior to dosing for the treated animals. Use ANOVA, together with Students t test, to compare mean body temperature circadian shifts in minutes between groups.



Table 5
Biological Data

Cmpd No.	Casein Kinase I ϵ ³³ P-ATP Filter Plate Assay K _i (nM)	Cell Assay EC _{Δτ+1h} (μM)
Ia	544	
Ib	157.35	
Ic	>10 ⁴	
Id	110	>30
Ie	159	
If	3090	
Ig	915.49	
Ih	70.95	
Ii	1677.81	
Ij	116.26	
Ik	468.28	
Il	324.03	
Im	1266.59	

In	237.97	
Io	78.65	
Ip	356.87	
Iq	37.99	
Ir	331.34	
Is	481.08	
It	425.30	
Iu	562	
Iv	4550.66	
Iw	2025.39	
Ix	1750.71	
Ila	64.04	
Ilb	811	0.874
Ilc	256.88	
Ild	1067.73	
Ile	109.11	
Ilf	1448.84	
Ilg	30.64	2.492
Ilh	24.73	
Ili	-----	
Ilj	112.37	
Ilk	70.56	
Ill	184.63	
IIm	68	
Iln	15.90	0.998
Ilo	9.60	0.662
Ilp	4030	
Ilq	334	
Ilr	333	
Ils	47.00	1.155
Ilt	22.00	1.974

What is claimed is:

1. A compound of formula (I) or formula (II)



wherein

- 5 X is S or S(O)_n;
 R₁ is H or C₁-C₆alkyl;
 R₂ is NR₅R₆;
 R₃ is aryl or heterocycle;
 R₄ is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl),
 10 C₁-C₆alkoxy, aryl-(C₁-C₆alkoxy), heterocycle-(C₁-C₆alkoxy), CF₃, halogen, SH, C₁-C₆alkylthio,
 aryl-(C₁-C₆alkylthio), heterocycle-(C₁-C₆alkylthio), NO₂, NH₂, NR₅R₆, aryl-(C₁-
 C₆alkylamino), heterocycle-(C₁-C₆alkylamino), or XR₃ wherein X and R₃ are as defined
 above;
 R₅ is H or C₁-C₆alkyl;
 15 R₆ is H or C₁-C₆alkyl;
 L is N or CR₇ wherein R₇ is H or C₁-C₆alkyl;
 M is S, O or NR₈ wherein R₈ is H, C₁-C₆alkyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl) or
 acyl;
 n is 1 or 2; or
 20 a stereoisomer, an enantiomer, a racemate or a tautomer thereof; or
 a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1 of formula (I) or formula (II) wherein M
 and X are each S.

25

3. The compound according to claim 2 of formula (I) wherein L is CR₇.
4. The compound according to claim 3 wherein R₇ is H.

5. The compound according to claim 4 selected from the group consisting of:

6-phenylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3-fluorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
5 6-(4-chlorophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-aminophenyl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(pyridin-2-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-p-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(thiophen-2-yl-sulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
10 6-(3,5-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(pyridin-4-ylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-m-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-o-tolylsulfanyl-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2,3-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
15 6-(2,5-dichloro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-ethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3-bromo-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3,5-dimethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(3-methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
20 6-(2-methoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-trifluoromethyl-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide,
6-(2-fluoro-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide, and
6-(3-trifluoromethoxy-phenylsulfanyl)-4H-thieno[3,2-b]pyrrole-5-carboxylic acid amide.

25 6. The compound according to claim 2 of formula (I) wherein L is N.

7. The compound according to claim 6 selected from the group consisting of:

6-phenylsulfanyl-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide,
6-(3-fluoro-phenylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide, and
30 6-(pyridin-2-ylsulfanyl)-4H-pyrrolo[2,3-d]thiazole-5-carboxylic acid amide.

8. The compound according to claim 2 of formula (II) wherein L is CR₇.

9. The compound according to claim 8 wherein R₇ is H.

10. The compound according to claim 9 selected from the group consisting of:

4-(pyridin-2-ylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
5 4-(phenylsulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
6-(3-fluorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(pyridin-4-ylsulfanyl)-6H-thieno[2,3-b]-pyrrole-5-carboxylic acid amide,
4-(3,5-dichlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(thiophen-2-yl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
10 4-(3-bromophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(3-methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(2-methoxyphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide,
4-(3-chlorophenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide, and
4-(3-methylphenyl-sulfanyl)-6H-thieno[2,3-b]pyrrole-5-carboxylic acid amide.



11. The compound according to claim 2 of formula (II) wherein L is N.

12. The compound according to claim 11 selected from the group consisting of:

2-methyl-6-phenyl-sulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
20 6-(3-methoxyphenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-fluorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-chlorophenyl-sulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-(3-trifluoromethoxy-phenylsulfanyl)-2-methyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid
amide,
25 2,6-bis-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
2,6-bis-(3-methoxy-phenylsulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide,
6-phenylsulfanyl-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide, and
6-(3-methoxyphenyl-sulfanyl)-4H-pyrrolo[3,2-d]thiazole-5-carboxylic acid amide.

30 13. A pharmaceutical composition comprising a pharmaceutically acceptable
carrier and a compound of formula (I) or formula (II) according to claim 1, or a stereoisomer,
an enantiomer, a racemate or a tautomer of said compound; or a pharmaceutically acceptable
salt thereof.

14. A method for inhibiting casein kinase I ϵ activity in a patient comprising administering to said patient a therapeutically effective amount of a compound of formula (I) or formula (II),



wherein

X is S or S(O)_n;

R₁ is H or C₁-C₆alkyl;

10 R₂ is NR₅R₆;

R₃ is aryl or heterocycle;

R₄ is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl), C₁-C₆alkoxy, aryl-(C₁-C₆alkoxy), heterocycle-(C₁-C₆alkoxy), CF₃, halogen, SH, C₁-C₆alkylthio, aryl-(C₁-C₆alkylthio), heterocycle-(C₁-C₆alkylthio), NO₂, NH₂, NR₅R₆, aryl-(C₁-

15 C₆alkylamino), heterocycle-(C₁-C₆alkylamino), or XR₃ wherein X and R₃ are as defined above;

R₅ is H or C₁-C₆alkyl;

R₆ is H or C₁-C₆alkyl;

L is N or CR₇ wherein R₇ is H or C₁-C₆alkyl;

20 M is S, O or NR₈ wherein R₈ is H, C₁-C₆alkyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl) or acyl;

n is 1 or 2; or



a stereoisomer, an enantiomer, a racemate or a tautomer thereof; or

a pharmaceutically acceptable salt thereof.

25

15. The method according to claim 14 wherein said inhibition of casein kinase I ϵ activity results in a lengthening of circadian rhythm period.

16. A method for treating a patient suffering from a disease or disorder ameliorated by inhibition of casein kinase I ϵ activity comprising administering to said patient a therapeutically effective amount of a compound of formula (I) or formula (II), or a stereoisomer, an enantiomer, a racemate or a tautomer of said compound, or a pharmaceutically acceptable salt thereof,



wherein

X is S or S(O)_n;

10 R₁ is H or C₁-C₆alkyl;

R₂ is NR₅R₆;

R₃ is aryl or heterocycle;

R₄ is H, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl), C₁-C₆alkoxy, aryl-(C₁-C₆alkoxy), heterocycle-(C₁-C₆alkoxy), CF₃, halogen, SH, C₁-C₆alkylthio,

15 aryl-(C₁-C₆alkylthio), heterocycle-(C₁-C₆alkylthio), NO₂, NH₂, NR₅R₆, aryl-(C₁-C₆alkylamino), heterocycle-(C₁-C₆alkylamino), or XR₃ wherein X and R₃ are as defined above;

R₅ is H or C₁-C₆alkyl;

R₆ is H or C₁-C₆alkyl;

20 L is N or CR₇ wherein R₇ is H or C₁-C₆alkyl;

M is S, O or NR₈ wherein R₈ is H, C₁-C₆alkyl, aryl-(C₁-C₆alkyl), heterocycle-(C₁-C₆alkyl) or acyl;

n is 1 or 2.

25 17. The method according to claim 16 wherein said inhibition of casein kinase I ϵ activity results in a lengthening of circadian rhythm period.

18. The method according to claim 16 wherein the disease or disorder is a mood disorder or a sleep disorder.

19. The method according to claim 18 wherein the disorder is mood disorder.

20. The method according to claim 19 wherein the mood disorder is selected from a depressive disorder or a bipolar disorder.

21. The method according to claim 20 wherein the depressive disorder is major depressive disorder.



22. The method according to claim 20 wherein the bipolar disorder is selected from the group consisting of bipolar I disorder and bipolar II disorder.

23. The method according to claim 18 wherein the disorder is a sleep disorder.

24. The method according to claim 23 wherein the sleep disorder is a circadian rhythm sleep disorder.

25. The method according to claim 24 wherein the circadian rhythm sleep disorder is selected from the group consisting of shift work sleep disorder, jet lag syndrome, advanced sleep phase syndrome and delayed sleep phase syndrome.

26. A process for the preparation of a compound of formula (I) or formula (II)



wherein:

X is S;

R₁ is H or C₁-C₆alkyl;

R_2 is NR_5R_6 ;

R_3 is aryl or heterocycle;

R_4 is XR_3 wherein X and R_3 are as defined above;

R_5 is H or C_1 - C_6 alkyl;



5 R_6 is H or C_1 - C_6 alkyl;

L is N;

M is S, O or NR_8 wherein R_8 is C_1 - C_6 alkyl, aryl- $(C_1$ - C_6 alkyl), heterocycle- $(C_1$ - C_6 alkyl) or acyl;

comprising:

10 reacting a compound of formula **10** or formula **13** wherein R_4 is halogen and R_2 is as defined above



with a diaryldisulfide or a diheterocycledisulfide in the presence of a suitable base in a suitable polar solvent, and

15 isolating the compound of formula (I) or formula (II) as defined above wherein R_4 is XR_3 .

27. The process of claim 26 wherein R_4 of the compound of formula **10** or formula **13** is bromine, the base is cesium carbonate, the polar solvent is dimethylformamide, and said
20 reaction is carried out at a temperature in the range of from about 80 °C to about 120 °C, for a period of time of from about one hour to about 6 hours.