

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

15 July 2021 (15.07.2021)



(10) International Publication Number

WO 2021/139780 A1

(51) International Patent Classification:

C07K 16/28 (2006.01) C07K 14/47 (2006.01)

A61K 39/00 (2006.01) C07K 19/00 (2006.01)

(21) International Application Number:

PCT/CN2021/070907

(22) International Filing Date:

08 January 2021 (08.01.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/CN2020/071336

10 January 2020 (10.01.2020) CN

(71) Applicant: SHANGHAI HENLIUS BIOTECH, INC.

[CN/CN]; Room 303, 304, Block 7, No. 1999 Zhangheng Road, China (Shanghai) Pilot Free Trade Zone, Pudong District, 201210 (CN).

(72) Inventors: YANG, Ming; Room 303, 304, Block 7, No.

1999 Zhangheng Road, China (Shanghai) Pilot Free Trade Zone, Pudong District, 201210 (CN). XU, Wenfeng; Room 303, 304, Block 7, No. 1999 Zhangheng Road, China (Shanghai) Pilot Free Trade Zone, Pudong District, 201210 (CN). JIANG, Wei-Dong; Room 303, 304, Block 7, No. 1999 Zhangheng Road, China (Shanghai) Pilot Free Trade Zone, Pudong District, 201210 (CN). XUE, Jie; Room 303, 304, Block 7, No. 1999 Zhangheng Road, China (Shanghai) Pilot Free Trade Zone, Pudong District, 201210 (CN).

(74) Agent: LECOME INTELLECTUAL PROPERTY

AGENT LTD., 3rd floor, Officebuilding, TaifuHotel, No. 1 Building, No.1 XituchengRoad, Haidian District, Beijing 100081 (CN).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTI-TIGIT ANTIBODIES, MULTISPECIFIC ANTIBODIES COMPRISING THE SAME AND METHODS OF USING THE SAME

(57) Abstract: Provided are anti-TIGIT antibodies that bind to "T cell immunoreceptor with Ig and ITIM domains (TIGIT) ", including multispecific anti-TIGIT antibodies with binding specificity for TIGIT and one or more additional antigen, and methods of using the same. The anti-TIGIT antibodies comprises a single domain antibody that binds to TIGIT. The one or more additional antigen comprises Programmed death protein 1 (PD1).



WO 2021/139780 A1

ANTI-TIGIT ANTIBODIES, MULTISPECIFIC ANTIBODIES COMPRISING THE SAME AND METHODS OF USING THE SAME

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims priority to International Patent Application No. PCT/CN2020/071336, filed January 10, 2020, the contents of which is incorporated by reference in its entirety, and to which priority is claimed.

FIELD

10 The present invention relates to antibodies that bind to T cell immunoreceptor with Ig and ITIM domains (TIGIT), including multispecific anti-TIGIT antibodies with binding specificity for TIGIT and one or more additional antigen, and methods of using the same. In certain embodiments, the one or more additional antigen comprises Programmed cell death protein 1 (PD1).

BACKGROUND

15 T cell immunoreceptor with Ig and ITIM domains (TIGIT) is an immune checkpoint receptor expressed by immune cells such as activated T cells and Natural Killer cells (NK cells) and mediates immunosuppression. The ligands of TIGIT includes PVR (CD155), which has been identified on dendritic cells (DCs), macrophages as well as many human cancer cells and have been
20 shown to downregulate T cell activation and cytokine secretion upon binding to TIGIT. Inhibition of the TIGIT/PVR interaction can mediate potent antitumor activity of immune cells. Given the significant role for TIGIT in immune checkpoint regulation, there remains a need in the art for the development of therapeutic molecules and methods to modulate TIGIT-mediated immune cell regulation for immune therapy and cancer treatment.

25 Programmed cell death protein 1 (PD1) is a key immune checkpoint receptor expressed by activated T and B cells and mediates immunosuppression. It is a member of the CD28 family of receptors, which includes CD28, CTLA-4, ICOS, PD1 and BTLA. Two cell surface glycoprotein ligands for PD1 have been identified, PDL1 and PDL2, which are expressed on antigen-presenting cells as well as many human cancers and have been shown to downregulate T cell activation and
30 cytokine secretion upon binding to PD1 (Freeman et al., 2000; Latchman et al., 2001). Unlike CTLA-4, PD1 primarily functions in peripheral tissues where activated T-cells may encounter the immunosuppressive PDL1 and PDL2 ligands expressed by tumor and/or stromal cells (Flies et al., 2011; Topalian et al., 2012). Inhibition of the PD1/PDL1 interaction mediates potent antitumor activity in preclinical models (U.S. Patent Nos. 8,008,449 and 7,943,743), and the use of antibody
35 inhibitors of the PD1/PDL1 interaction for treating cancer has been studied in clinical trials. Accordingly, there is a need for the development of anticancer therapeutics targeting PD1.

SUMMARY OF THE INVENTION

The present disclosure provides isolated monoclonal antibodies that bind to TIGIT with high affinity, including multispecific antibodies that binds to TIGIT and one or more additional target. In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody that binds to TIGIT. In certain embodiments, the additional target is PD1. This disclosure further provides methods of making and using the antibodies, immunoconjugates and pharmaceutical compositions comprising the antibodies, e.g., for treating diseases and disorders, e.g., cancer. The invention is based, in part, on the discovery of single domain anti-TIGIT antibodies that bind to TIGIT and multispecific antibodies that bind to both TIGIT and PD1, which antibodies can increase an immune response in immune cells and provide improved anti-tumor efficacy.

The present disclosure provides multispecific antibodies that bind to TIGIT and PD1. In certain embodiments, a multispecific antibody disclosed herein comprise: i) a first antigen-binding moiety comprising an anti-TIGIT antibody comprising a single domain antibody that binds to TIGIT; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody that binds to PD1.

In certain embodiments, the single domain antibody comprises a VHH. In certain embodiments, the single domain antibody or the VHH comprises a heavy chain variable region (VH). In certain embodiments, the single domain antibody binds to TIGIT with a K_D of 1×10^{-7} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a K_D of 1×10^{-8} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a K_D of 3×10^{-9} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a K_D of 2×10^{-9} M or less.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference single domain antibody comprising a heavy chain variable region comprising:

a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96,

b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100,

c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104,

d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108,

5 e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112,

10 f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116,

15 g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120,

20 h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124,

i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128,

25 j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132,

30 k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136,

35 l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140,

m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144,

5 n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148,

10 o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 151, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 152,

15 p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 154, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 155, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 156,

20 q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160,

r) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164,

25 s) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168,

30 t) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172,

35 u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176,

v) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180,

5 w) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184,

10 x) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188, or

15 y) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising:

20 a) a heavy chain variable region CDR1 comprising an amino acid sequence of any one of SEQ ID NOs: 94, 98, 102, 106, 110, 114, 118, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186 and 190, or a variant thereof comprising up to about 3 amino acid substitutions;

25 b) a heavy chain variable region CDR2 comprising an amino acid sequence of any one of SEQ ID NOs: 95, 99, 103, 107, 111, 115, 119, 123, 127, 131, 135, 139, 143, 147, 151, 155, 159, 163, 167, 171, 175, 179, 183, 187 and 191, or a variant thereof comprising up to about 3 amino acid substitutions; and

30 c) a heavy chain variable region CDR3 comprising an amino acid sequence of any one of SEQ ID NOs: 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188 and 192, or a variant thereof comprising up to about 3 amino acid substitutions.

35 In certain embodiments, the single domain antibody comprises a heavy chain variable region that comprises a CDR1 domain, a CDR2 domain and a CDR3 domain, wherein the CDR1 domain, the CDR2 domain and the CDR3 domain respectively comprise a CDR1 domain, a CDR2 domain and a CDR3 domain comprised in a reference heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

[illegible]

forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 90% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

[illegible]

amino acid sequence set forth in SEQ ID NO: 181. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 185. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 189. In certain
5 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference single domain antibody comprising a heavy chain variable region comprising:

(a) a heavy chain variable region CDR1 comprising amino acids having the sequence set
10 forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3,

(b) a heavy chain variable region CDR1 comprising amino acids having the sequence set
15 forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7,

(c) a heavy chain variable region CDR1 comprising amino acids having the sequence set
20 forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11,

(d) a heavy chain variable region CDR1 comprising amino acids having the sequence set
forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 15,

(e) a heavy chain variable region CDR1 comprising amino acids having the sequence set
25 forth in SEQ ID NO: 17, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 18, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 19,

(f) a heavy chain variable region CDR1 comprising amino acids having the sequence set
30 forth in SEQ ID NO: 21, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 22, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 23,

(g) a heavy chain variable region CDR1 comprising amino acids having the sequence set
35 forth in SEQ ID NO: 25, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 26, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 27,

(h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 29, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 30, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 31,

5 (i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 33, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 34, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 35,

10 (j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 37, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 38, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 39,

15 (k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 41, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 42, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 43,

20 (l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 45, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 46, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 47,

(m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 49, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 50, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 51,

25 (n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 53, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 54, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 55,

30 (o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 57, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 58, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 59,

35 (p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 61, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 62, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 63,

(q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 65, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 66, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 67,

5 (r) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 69, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 70, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 71,

10 (s) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75,

15 (t) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79, and

20 (u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising:

25 (a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3,

30 (b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7,

(c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11,

35 (d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the

sequence set forth in SEQ ID NO: 50, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 51,

(n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 53, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 54, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 55,

(o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 57, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 58, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 59,

(p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 61, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 62, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 63,

(q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 65, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 66, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 67,

(r) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 69, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 70, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 71,

(s) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75,

(t) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79, and

(u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region having at least about 90% sequence identity to an amino acid sequence selected from the group

consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84.

In certain embodiments, the single domain antibody comprises a humanized framework.

In certain embodiments, the second antigen-binding moiety comprises an anti-PD1 antibody that cross-competes with a reference anti-PD1 antibody comprising:

a) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226);

b) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO: 231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234);

c) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242);

ord) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO: 247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO: 248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

In certain embodiments, the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence

TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226).

In certain embodiments, the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO: 231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234).

In certain embodiments, the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242).

In certain embodiments, the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO: 247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO: 248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

In certain embodiments, the anti-PD1 antibody comprises a humanized antibody.

In certain embodiments, the second antigen binding moiety comprises an anti-PD1 antibody comprising two antibody heavy chains and two antibody light chains. In certain embodiments, the first antigen-binding moiety comprises two or more anti-TIGIT antibodies. In certain embodiments, the first antigen-binding moiety comprises two anti-TIGIT antibodies.

In certain embodiments, the C-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-

terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

In certain embodiments, the first antigen binding moiety is linked to the second antigen binding moiety via a linker. In certain embodiments, the linker is a peptide linker. In certain embodiments, the peptide linker comprises about four to about thirty amino acids. In certain embodiments, the peptide linker comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220.

In certain embodiments, the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM. In certain embodiments, the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc region of IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region comprises a human Fc region. In certain embodiments, the Fc region comprises an IgG1 Fc region. In certain embodiments, the Fc region comprises an IgG4 Fc region. In certain embodiments, the IgG4 Fc region comprises an S228P mutation. In certain embodiments, the multispecific antibody is a bispecific antibody.

In certain embodiments, the multispecific antibody comprises: i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 229, (2) a CDR-H2 comprising the amino acid sequence set forth in SEQ ID NO: 230, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 231; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 232, (2) a CDR-L2 comprising the

amino acid sequence set forth in SEQ ID NO: 233, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 234.

In certain embodiments, the multispecific antibody comprises: i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 237, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 238, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 239; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 240, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 241, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 242.

In certain embodiments, the multispecific antibody comprises: i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 221, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 222, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 223; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 224, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 225, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 226.

In certain embodiments, the multispecific antibody comprises: i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 245, (2) a CDR- H2 comprising the amino acid

sequence set forth in SEQ ID NO: 246, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 247; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 248, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 249, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 250.

In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 253, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 254. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 255, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 256. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 257, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 258. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 259, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 260.

The present disclosure further provides immunoconjugates comprising any multispecific antibodies disclosed herein, linked to a therapeutic agent. In certain embodiments, the therapeutic agent is a cytotoxin. In certain embodiments, the therapeutic agent is a radioactive isotope.

The present disclosure further provides pharmaceutical compositions. In certain embodiments, the pharmaceutical composition comprises a) a multispecific antibody or an immunoconjugate disclosed herein, and b) a pharmaceutically acceptable carrier.

The present disclosure further provides nucleic acids encoding any multispecific antibodies disclosed herein. The present disclosure further provides vectors comprising any nucleic acid disclosed herein. The present disclosure further provides host cells comprising a nucleic acid or a vector disclosed herein.

The present disclosure further provides methods for preparing a multispecific antibody disclosed herein. In certain embodiments, the method comprises expressing a multispecific antibody in a host cell disclosed herein and isolating the multispecific antibody from the host cell.

The present disclosure further provides methods of reducing tumor burden in a subject. In certain embodiments, the method comprises administering to the subject an effective amount of a multispecific antibody, an immunoconjugate, or a pharmaceutical composition disclosed herein. In certain embodiments, the method reduces the number of tumor cells. In certain embodiments, the method reduces tumor size. In certain embodiments, the method eradicates the tumor in the subject.

In certain embodiments, the tumor is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer, blood cancer and a combination thereof.

The present disclosure further provides methods of treating and/or preventing a neoplasm in a subject. In certain embodiments, the method comprises administering to the subject an effective amount of an multispecific antibody, an immunoconjugate, or a pharmaceutical composition disclosed herein.

The present disclosure further provides methods of lengthening survival of a subject having a neoplasm. In certain embodiments, the method comprises administering to the subject an effective amount of an multispecific antibody, an immunoconjugate, or a pharmaceutical composition disclosed herein.

In certain embodiments, the neoplasm is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer, blood cancer and a combination thereof.

The present disclosure further provides multispecific antibodies disclosed herein for use as a medicament. The present disclosure further provides multispecific antibodies disclosed herein for use in treating cancer. The present disclosure further provides pharmaceutical compositions disclosed herein for use as a medicament. The present disclosure further provides pharmaceutical compositions disclosed herein for use in treating cancer. In certain embodiments, the cancer is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer, blood cancer and a combination thereof.

The present disclosure further provides kits comprising an multispecific antibody, an immunoconjugate, a pharmaceutical composition, a nucleic acid, a vector or a host cell disclosed herein. In certain embodiments, the kit comprise a written instruction for treating and/or preventing a neoplasm.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-1C depict whole cell binding of representative VHH bivalent antibodies to human TIGIT determined by flowcytometry assays. Figure 1A shows one assay with comparison to a reference anti-human-TIGIT antibody (Reference Ab 1). Figure 1B shows a different assay with

additional VHH antibodies. Y axis represents Mean Fluorescence Intensity of AlexaFlour 488. X axis represents antibody concentrations in nanomolar. 2B7, 1G1, 1C12, 3G6, 2B10, 3G7, 3G10, 13H11, and 15A5 are VHH clones of anti-human TIGIT. EC50 values were obtained using non-linear regression method by Prism, and values are shown in the table as nanomolar. Figure 1C is a schematic of the structure of a TIGIT VHH bivalent antibody.

Figure 2A-2B depict the potency of representative bivalent antibodies in blocking TIGIT activity determined by luciferase reporter assays. Human TIGIT and NFAT reporter gene stably transfected Jurkat cells were co-cultured with PVR (CD155) stably transfected Raji cells in the presence of representative anti-TIGIT bivalent antibodies and a low concentration of staphylococcal enterotoxin. Y axis represents NFAT luciferase activity in relative luminescence units. X axis represents antibody concentration in nanomolar. Reference Ab 1 is a reference anti-h-TIGIT antibody. 2B7, 1G1, 1C12, 3G6, 2B10, 3G7 and 3F10 are representative clones of anti-h-TIGIT. EC50 values were obtained using non-linear regression method by Prism, and values are shown in the table as nanomolar.

Figures 3A and 3B depict whole cell binding of humanized 1C12 and 1G1 clones to human TIGIT determined by flowcytometry assays. A representative result of 1C12 humanized versions is shown in Figure 3A. 1C12-chimeric is an antibody with Llama VHH sequence of 1C12 clone and human IgG1 CH2 and CH3 domains, 1C12(F-EREF), 1C12 (F-EREW) and 1C12(F-GLEW) are humanized versions of 1C12 clone with difference of mutations in Framework 2. A representative result of 1G1 humanized versions is shown in Figure 3B. 1G1-chimeric is an antibody with Llama VHH sequence of 1G1 clone and human IgG1 CH2 and CH3 domains. 1G1 (F-G-ERES), 1G1 (F-A-ERES), 1G1 (F-A-EREW) and 1G1 (F-A-GLEW) are four different versions of humanized 1G1 clones with different mutations in framework 2. Y axis is values of Mean Fluorescence Intensity of AlexaFlour 488. X axis is values of antibody concentrations in nanomolar.

Figures 4A and 4B depict the potency of humanized 1C12 and 1G1 clones in blocking TIGIT activity determined by luciferase reporter assays. Representative results of 1C12 humanized versions in TIGIT blockade luciferase reporter assay is shown in Figure 4A. All clones are more potent compared to Reference Ab 1, a reference anti-h-TIGIT antibody. Representative results of 1G1 humanized versions are shown in Figure 4B. Y axis represents NFAT luciferase activity (RLU, relative luminescence unite). X axis represents antibody concentrations in nanomolar.

Figure 5 depicts correlations between IC50 values (in nanomolar) of blocking ELISA and EC50 values (in nanomolar) of whole cell binding for representative clones. The clone names are labeled in the figures. Correlations were analyzed using GraphPad Prism.

Figures 6A and 6B depict thermostability tested for representative clones. The VHH antibody samples were heated from 25 to 70 ° C for 60 min. Binding of heated samples to human TIGIT was examined using ELISA or whole cell binding. Figure 6A shows binding of heated

samples of representative clones to h-TIGIT ECD in an ELISA assay with clone names labeled in the figure legends. Y axis represents OD450 obtained from the ELISA assay. X axis represents temperature for the treatment. Figure 6B shows binding of heated samples of representative clones to h-TIGIT stably expressed in Jurkat cells determined by flow cytometry. Y axis represents percentage of binding to h-TIGIT. X axis represents temperature for the treatment.

Figures 7A-7C depict the epitope at which 2A3-Fc binds to human TIGIT determined by Octet binding assay using ForteBio. Recombinant protein of his-tagged human TIGIT ECD (200 nM) was loaded on sensor. Binding was detected by injection of 2A3-Fc at three different concentrations. No additional binding was detected by second injection of 2A3-Fc shown in Figure 7A, however strong binding was detected by injection of three different concentrations of either Reference Ab 2, a reference anti-human-TIGIT antibody, shown in Figure 7B, or Reference Ab 1 shown in Figure 7C, two reference anti-h-TIGIT antibodies. The results indicate that 2A3 clone has a different binding epitope compared to Reference Ab 2 and Reference Ab 1.

Figures 8A-8C depict cross-binding activity of 2A3-Fc to human-, cyno- and mouse-TIGIT determined by ELISA assays. Figure 8A shows binding of 2A3-Fc to recombinant human TIGIT ECD. Reference Ab 2 is an anti-h-TIGIT reference antibody. Both 2A3-Fc and Reference Ab 2 bind to h-TIGIT with similar affinity. Anti-PDL1 did not bind to h-TIGIT. Figure 8B shows binding of 2A3-Fc to recombinant cyno-TIGIT. Both 2A3-Fc and Reference Ab 2 bind to cyno-TIGIT with similar affinity. Anti-PDL1 did not bind to cyno-TIGIT. Figure 8C shows binding of 2A3-Fc to recombinant mouse-TIGIT. Neither 2A3-Fc nor Reference Ab 2 binds to mouse-TIGIT, however an anti-mouse TIGIT reference antibody (Biolegend #142101) bind to mouse-TIGIT with a high affinity. Y axis represents OD450, X axis represents antibody concentrations in μ g/ml.

Figure 9 depicts comparison of potency of representative clone 2A3-Fc and hotspot corrected version of 2A3-LT-Fc in whole cell binding assay and human TIGIT blockade reporter assay. Y axis represents mean fluorescent intensity (MFI) in the upper panel and NFAT luciferase reporter activity in relative luminescence units in the lower panel. X axis represents antibody concentrations in nanomolar. Hotspot corrected version 2A3-LT-Fc had similar potency compared to the parental clone 2A3-Fc.

Figures 10A and 10B depict comparison of affinity of hotspot corrected version 2A3-LT-Fc to a reference anti-h-TIGIT antibody, Reference Ab 2. Figure 10A shows the comparison of potency of hotspot corrected version 2A3-LT-Fc to Reference Ab 2 in whole cell binding. Y axis represents mean fluorescence intensity of AlexaFluor 488 determined by flow cytometry assay using CytoFlex. X axis represents antibody concentrations in nanomolar. Data shown is a representative result from three different experiments. 2A3-LT-Fc had slightly higher affinity to h-TIGIT than Reference Ab 2. Figure 10B shows the comparison of potency of hotspot corrected version 2A3-LT-Fc to Reference Ab 2 in TIGIT blockade NFAT reporter assay. Y axis represents NFAT luciferase

reporter activity in relative luminescence units. X axis represents antibody concentrations in nanomolar. 2A3-LT-Fc had similar potency to that of Reference Ab 2.

Figure 11 depicts *in vitro* antitumor efficacy of 2A3-LT-Fc. Mixed lymphocyte reaction assay (MLR) was performed using TIGIT⁺ T cells and PVR⁺ dendritic cells (DCs). IL-2 secretion from the T cells were measured in culture supernatant after 48 hours of coculture. An anti-PD1 antibody was used as a positive control. An anti-HER2 antibody was used as a negative control.

Figure 12 depicts binding of 2A3-LT-Fc wild type (wt) and Fc mutations (DLE and VLPLL) to human Fc γ RIIIA, human Fc γ RIIB and mouse Fc γ RIV determined using Octet binding assay. Sensors were loaded with recombinant proteins of ECDs of Fc γ RIIIA, human Fc γ RIIB and mouse Fc γ RIV, association and dissociation of 2A3-LT-Fc wt, DLE or VLPLL mutants at 5 different concentrations were detected using ForteBio. Y axis represents association, dissociation & R_{max}. X axis represents time in second. Both DLE and VLPLL mutants had enhanced binding affinity to human Fc γ RIIIA compared to that of wild type, DLE mutant also had enhanced binding affinity to human Fc γ RIIB, however VLPLL mutant reduced binding affinity to human Fc γ RIIB. All versions had similar affinity to mouse Fc γ RIV.

Figures 13A and 13B depict effects of TIGIT mAbs 2A3-LT-Fc wt, and DLE mutant on blocking TIGIT activity and human Fc γ RIIIA-mediated activity, respectively. Figure 13A depicts the effects of TIGIT mAbs 2A3-LT-Fc wt, and DLE mutant on blocking TIGIT activity determined using NFAT luciferase reporter assay. Human TIGIT and NFAT reporter gene stably transfected Jurkat cells were co-cultured with PVR stably transfected Raji cells. 2A3-LT-Fc wt and mutants were added and cultured for 5 hours. The TCR-mediated activity was measured by luciferase activity. Y axis represents NFAT luciferase activity in relative luminescence units. X axis represents antibody concentrations in nanomolar. Figure 13B depicts the effects of TIGIT mAbs 2A3-LT-Fc wt, and DLE mutant on human Fc γ RIIIA-mediated activity determined by Fc γ RIIIA-mediated NFAT luciferase reporter activity. Human Fc γ RIIIA and NFAT stably transfected Jurkat cells were co-cultured with human TIGIT stably transfected 293T cells in the presence of different concentrations of 2A3-LT-Fc wt, and DLE mutant for 5 hours. Luciferase activity was measured and is presented in Y axis as relative luminescence units. X axis represents antibody concentrations in nanomolar.

Figures 14A-14C depict *in vivo* efficacy analyses of the anti-TIGIT antibodies. H-TIGIT Knock-In C57BL/6 mice and MC38 murine colon cancer model were used. MC38 tumor cells were implanted one week before treatment. Treatment started when tumor volume reached about 51 mm³, when drugs were dosed intraperitoneally twice a week for 2.5 weeks at 6 mg/kg for 2A3-LT-Fc antibodies, or 11 mg/kg for reference antibody (equal to 6 mg/kg of 2A3-LT-Fc in mole/kg dosage). Figure 14A depicts tumor growth curves of tumor bearing mice treated with vehicle control, 2A3-Fc wildtype, 2A3-Fc with DLE mutations and Reference Ab 2. The results of individual tumor

volume are shown in Figure 14B. Figure 14C shows that body weight changes were not significant between the groups throughout the study.

Figure 15 depicts various structures of anti-TIGIT/anti-PD1 multispecific antibodies.

Figures 16A-16F depict double binding ability of BsAbs to h-TIGIT and h-PD1 determined by Octet binding assay. Sensors were loaded with BsAbs and then injected with different concentrations of recombinant protein of h-PD1 ECD (his-tagged). After association of BsAbs with h-TIGIT, the sensors were injected again with different concentrations of recombinant protein of h-TIGIT ECD (his-tagged). Binding curves of anti-TIGIT x anti-PD1 (aTIGIT x aPD1)-HCC, HCN, LCC & LCN are shown in (A), (B), (C), & (D) respectively. Binding curves of monoclonal antibody aTIGIT or aPD1 are shown in (E) and (F). All BsAbs had double binding ability to h-PD1 and h-TIGIT. Y axis represents association/disassociation and Rmax. X axis represents time in second.

Figure 17 depicts affinity of BsAbs binding to h-TIGIT single arm determined by whole cell binding assay using flowcytometry. Antibodies were incubated with h-TIGIT transfected Jurkat cells, and binding of antibodies to h-TIGIT was determined using a secondary anti-h-IgG Fc antibody conjugated with Fc AlexaFluor488, which was detected by flow cytometry. Y axis represents mean fluorescent intensity (MFI). X axis represents concentrations of antibodies. BsAb HCN and LCN versions maintained binding affinity to h-TIGIT compared to anti-TIGIT mAb, whereas HCC and LCC versions of BsAbs showed reduced binding affinity to h-TIGIT.

Figure 18 depicts potency of BsAbs in blocking TIGIT determined by TIGIT blockade luciferase reporter assays. Antibodies were cultured with h-TIGIT and NFAT reporter transfected Jurkat cells in the presence of PVR transfected Raji cells and a low concentration of staphylococcal enterotoxin. TCR-mediated NFAT luciferase reporter activity was measured by a chemiluminescent plate reader and is shown in the Y axis in relative luminescence units. HCN version of BsAb was the most potent BsAb in blocking h-TIGIT.

Figure 19 depicts binding affinities of BsAbs to h-PD1 single arm determined by whole cell binding assay using flowcytometry. Antibodies were incubated with h-PD1 transfected Jurkat cells, and binding was determined using a secondary anti-h-IgG Fc antibody conjugated with Fc AlexaFluor488, which was detected by flow cytometry. Y axis represents mean fluorescent intensity (MFI). X axis represents concentrations of antibodies. All BsAbs showed similar binding affinities to h-PD1, with slightly lower affinity compared to monoclonal anti-PD1 parental control.

Figure 20 depicts potency of BsAbs in blocking PD1 single arm determined by PD1 blockade luciferase reporter assay. Antibodies were cultured with h-PD1 and NFAT reporter transfected Jurkat cells in the presence of PDL1 and CD3 transfected CHO cells. TCR-mediated NFAT luciferase activity was measured by a chemiluminescent plate reader and shown in Y axis in

relative luminescence units. All BsAbs maintained blocking ability compared to parental anti-PD1 mAb.

Figure 21 depicts potency of BsAbs in double blocking both TIGIT and PD1 determined using a combination of TIGIT and PD1 blockade assay. mAbs, combination of mAbs and BsAbs were cultured with h-TIGIT and h-PD1 & NFAT reporter transfected Jurkat cells in the presence of PVR and PDL1 transfected CHO cells and a low concentration of staphylococcal enterotoxin. TCR-mediated NFAT luciferase reporter activity was measured by a chemiluminescent plate reader and shown in Y axis in relative luminescence units. The combination of anti-TIGIT (aTIGIT) and anti-PD1 (aPD1) was more potent than either anti-TIGIT or anti-PD1 mAbs. All the BsAbs were more potent than the combination of anti-TIGIT (aTIGIT) and anti-PD1 (aPD1).

Figures 22A and 22B depict in vivo efficacy of BsAb in MC38 syngeneic tumor model using h-PD1 and h-TIGIT double Knock-In C57BL/6 mice. Mice were inoculated with 0.5×10^6 MC38 tumor cells subcutaneously one week before treatment. One week later, when the mean tumor size reached approximately 43 mm^3 , mice were randomly grouped into 6 groups of 7 mice/group and treated with vehicle (PBS), anti-TIGIT-Fc (aTIGIT-Fc), anti-PD1 (aPD1), combination of aTIGIT-Fc+aPD1 and BsAb HCN at the specified dosages by intraperitoneally (IP). Treatment was performed B.I.W. As shown in Figure 22A, tumor volumes were measured biweekly and shown in Y axis in mm^3 , indicating that both aTIGIT-Fc and aPD1 significantly reduced tumor growth. On day 20 of the study, all animals were euthanized, and tumor tissues were excised, weighted. Individual tumor weight (TW) of each study group is plotted on Figure 22B.

Figures 23A and 23B depict multiple mechanisms of action of the anti-TIGIT/anti-PD1 antibodies. Figure 23A shows that the BsAb can block inhibitory signals in PD1 & TIGIT double or single positive CD8⁺ T cells and can block inhibitory signals from PDL1 & PVR double positive tumor cells or PVR single positive tumor cells. Figure 23B shows that the BsAb can block inhibitory signal in NK cells and stimulatory signal in Tregs.

DETAILED DESCRIPTION

The present disclosure provides isolated monoclonal antibodies that bind to TIGIT with high affinity, including multispecific antibodies that binds to TIGIT and one or more additional target. In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody that binds to TIGIT. In certain embodiments, the additional target is PD1. This disclosure further provides methods of making and using the antibodies, immunoconjugates and pharmaceutical compositions comprising the antibodies, e.g., for treating diseases and disorders, e.g., cancer. The invention is based, in part, on the discovery of single domain anti-TIGIT antibodies that bind to TIGIT and multispecific antibodies that bind to both TIGIT and PD1, which antibodies are capable of increasing an immune response in immune cells and provide improved anti-tumor efficacy.

For clarity and not by way of limitation the detailed description of the presently disclosed subject matter is divided into the following subsections:

1. Definitions;
2. Antibodies;
3. Methods of use;
4. Pharmaceutical formulations; and
5. Articles of manufacture.

1. DEFINITIONS

The term “antibody” herein is used in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), single domain antibody and antibody fragments so long as they exhibit the desired antigen-binding activity.

An “antibody fragment” refers to a molecule that comprises an antigen binding portion of an intact full-length antibody that binds to the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂, diabodies, linear antibodies, single-chain antibody molecules (e.g., scFv), multispecific antibodies formed from antibody fragments, a single domain antibody, a VHH, a nanobody, a domain antibody, a bivalent domain antibody, or any other fragment of an antibody that binds to an antigen. A “VHH” refers to a single domain antibody isolated from a camelid animal. In certain embodiments, a VHH comprises a variable region of a heavy chain of a camelid heavy chain antibody. In certain embodiments, a VHH has a size of no more than about 25 kDa. In certain embodiments, a VHH has a size of no more than about 20 kDa. In certain embodiments, a VHH has a size of no more than about 15 kDa.

A “full-length antibody” refers to an antibody comprising two heavy chains and two light chains. The variable regions of the light and heavy chains are responsible for antigen binding. The variable regions of a heavy chain and a light chain may be referred to as “VH” and “VL”, respectively. The variable regions in both chains generally contain three highly variable loops called the complementarity determining regions (CDRs) (light chain (LC) CDRs including LC-CDR1, LC-CDR2, and LC-CDR3, heavy chain (HC) CDRs including HC-CDR1, HC-CDR2, and HC-CDR3). CDR boundaries for the antibodies and antigen-binding fragments disclosed herein may be defined or identified by well-known conventions, e.g., the conventions of Kabat, Chothia, or Al-Lazikani (Al-Lazikani 1997; Chothia 1985; Chothia 1987; Chothia 1989; Kabat 1987; Kabat 1991). The three CDRs of the heavy or light chains are interposed between flanking stretches known as framework regions (FRs), which are more conserved than the CDRs and form a scaffold to support the hypervariable loops. The constant regions of the heavy and light chains are not involved in

antigen binding but exhibit various effector functions. Antibodies are assigned to classes based on the amino acid sequence of the constant region of their heavy chain. The five major classes or isotypes of antibodies are IgA, IgD, IgE, IgG, and IgM, which are characterized by the presence of α , δ , ϵ , γ , and μ heavy chains, respectively. Several of the major antibody classes are divided into subclasses such as IgG1 (γ 1 heavy chain), IgG2 (γ 2 heavy chain), IgG3 (γ 3 heavy chain), IgG4 (γ 4 heavy chain), IgA1 (α 1 heavy chain), or IgA2 (α 2 heavy chain).

An “antibody that cross-competes for binding” with a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition assay by 50% or more. An exemplary competition assay is described in Antibodies, Harlow and Lane (Cold Spring Harbor Press, Cold Spring Harbor, NY).

“Fv” is a minimum antibody fragment which contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region in tight, non-covalent association. From the folding of these two domains emanate six hypervariable loops (3 loops in each of the heavy and light chains) that contribute the amino acid residues to antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) can recognize and bind to an antigen, although sometimes at a lower affinity than the entire binding site.

“Single-chain Fv,” also abbreviated as “sFv” or “scFv,” are antibody fragments that comprise the V_H and V_L antibody domains connected into a single polypeptide chain. In some embodiments, the scFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv, see Plückthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

An “acceptor human framework” or “human framework” for the purposes herein is a framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin framework or a human consensus framework. An acceptor human framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain amino acid sequence changes. In certain embodiments, the number of amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In certain embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence.

“Affinity” refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless

indicated otherwise, as used herein, “binding affinity” refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (KD). Affinity can be measured by common methods known in the art, including those described
 5 herein. Specific illustrative and exemplary embodiments for measuring binding affinity are described in the following.

An “affinity matured” antibody refers to an antibody with one or more alterations in one or more CDRs or hypervariable regions (HVRs), compared to a parent antibody which does not possess such alterations, which alterations provide improved affinity of the antibody for antigen.

10 “T cell immunoreceptor with Ig and ITIM domains” or “TIGIT” as used herein, refers to any native TIGIT polypeptide from any vertebrate source, including mammals such as primates (e.g., humans and cynomolgus monkeys), or any fragment thereof, and may optionally comprise up to one, up to two, up to three, up to four, up to five, up to six, up to seven, up to eight, up to nine or up to ten amino acid substitutions, additions and/or deletions. The term encompasses full-length,
 15 unprocessed TIGIT as well as any form of TIGIT that results from processing in the cell. The term also encompasses naturally occurring variants of TIGIT, e.g., splice variants or allelic variants. A non-limiting example of a human TIGIT amino acid sequence targeted by an anti-TIGIT antibody of the present disclosure is as follows:

```

  1 MRWCLLLIWA QGLRQAPLAS GMMTGTIETT GNISAEKGGG IILQCHLSST TAQVTQVNWE
  20 61 QQDQLLAICN ADLGWHISPS FKDRVAPGPG LGLTLQSLTV NDTGEYFCIY HTYPDGTGTYTG
  121 RIFLEVLESS VAEHGARFQI PLLGAMAATL VVICTAVIVV VALTRKKKAL RIHSVEGDLR
  181 RKSAGQEEWS PSAPSPPGSC VQAEAAPAGL CGEQRGEDCA ELHDYFNVLS YRSLGNCSFF
  241 TETG[SEQ ID NO: 261].
  
```

The term “ECD of TIGIT” refers to the extracellular domain of TIGIT. For example, the
 25 ECD of the exemplary TIGIT protein shown in SEQ ID NO: 261 comprises the following amino acid sequence:

```

  MRWCLLLIWA QGLRQAPLAS GMMTGTIETT GNISAEKGGG IILQCHLSST TAQVTQVNWE
  QQDQLLAICN ADLGWHISPS FKDRVAPGPG LGLTLQSLTV NDTGEYFCIY HTYPDGTGTYTG
  RIFLEVLESS VAEHGARF[SEQ ID NO: 262].
  
```

30 The terms “anti-TIGIT antibody” and “an antibody that binds to TIGIT” refer to an antibody that is capable of binding to TIGIT with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent for targeting TIGIT. In one embodiment, the extent of binding of an anti-TIGIT antibody to an unrelated, non-TIGIT protein is less than about 10% of the binding of the antibody to TIGIT as measured, e.g., by a BIACORE[®] surface plasmon resonance assay. In
 35 certain embodiments, an antibody that binds to TIGIT has a dissociation constant (KD) of < about 1 μM, < about 100 nM, < about 10 nM, < about 1 nM, < about 0.1 nM, < about 0.01 nM, or < about

0.001 nM (e.g., 10^{-8} M or less, e.g., from 10^{-8} M to 10^{-12} M, e.g., from 10^{-9} M to 10^{-10} M). In certain embodiments, an anti-TIGIT antibody binds to an epitope of TIGIT that is conserved among TIGIT from different species. In certain embodiments, an anti-TIGIT antibody binds to an epitope on TIGIT that is in the ECD of the protein.

5 “Programmed cell deathprotein1” or “PD1”, as used herein, refers to any native PD1 polypeptide from any vertebrate source, including mammals such as primates (e.g., humans and cynomolgus monkeys), or any fragment thereof, and may optionally comprise up to one or up to two or up to three amino acid substitutions, additions or deletions. The term encompasses “full-length”, unprocessed PD1 as well as any form of PD1 that results from processing in the cell. The
10 term also encompasses naturally occurring variants of PD1, e.g., splice variants or allelic variants. Non-limiting examples of a human PD1 amino acid sequence targeted by an anti-PD1 antibody of the present disclosure include polypeptides having the NCBI Reference Nos: NP_005009.2, XP_016859782.1, XP_006712636.1, and any variants, modifications and descendants thereof

The terms “anti- PD1 antibody” and “an antibody that binds to PD1” refer to an antibody
15 that is capable of binding PD1 with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting PD1. In certain embodiments, the antibody is any anti-PD1 antibody disclosed in PCT/US2017/050851 (published as International Publication WO 2018/052818 A1), the content of which is incorporated herein by references in its entirety.

The term “chimeric” antibody refers to an antibody in which a portion of the heavy and/or
20 light chain is derived from a particular source or species, while the remainder of the heavy and/or light chain is derived from a different source or species. In certain embodiments, a chimeric antibody disclosed herein comprises a camelid heavy chain variable region and a human Fc region.

As used herein, the term “CDR” or “complementarity determining region” is intended to mean the non-contiguous antigen combining sites within the variable region of a heavy chain and/or
25 a light chain. These particular regions have been described by Kabat et al., J. Biol. Chem. 252:6609-6616 (1977); Kabat et al., U.S. Dept. of Health and Human Services, “Sequences of proteins of immunological interest” (1991); Chothia et al., J. Mol. Biol. 196:901-917 (1987); Al-Lazikani B. et al., J. Mol. Biol., 273: 927-948 (1997); MacCallum et al., J. Mol. Biol. 262:732-745 (1996); Abhinandan and Martin, Mol. Immunol., 45: 3832-3839 (2008); Lefranc M.P. et al., Dev.
30 Comp. Immunol., 27: 55-77 (2003); and Honegger and Plückthun, J. Mol. Biol., 309:657-670 (2001), where the definitions include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of any one of the definitions to refer to a CDR of an antibody or grafted antibodies or variants thereof is intended to be within the scope of the term as defined and used herein. The amino acid residues which encompass the CDRs as defined by each of
35 the above cited references are set forth below in Table 1 as a comparison. CDR prediction algorithms and interfaces are known in the art, including, for example, Abhinandan and Martin,

Mol. Immunol., 45: 3832-3839 (2008); Ehrenmann F. et al., Nucleic Acids Res., 38: D301-D307 (2010); and Adolf-Bryfogle J. et al., Nucleic Acids Res., 43: D432-D438 (2015). The contents of the references cited in this paragraph are incorporated herein by reference in their entireties for use in the present application and for possible inclusion in one or more claims herein.

Table 1: CDR definitions

	Kabat¹	Chothia²	MacCallum³	IMGT⁴	AHo⁵
V _H CDR1	31-35	26-32	30-35	27-38	25-40
V _H CDR2	50-65	53-55	47-58	56-65	58-77
V _H CDR3	95-102	96-101	93-101	105-117	109-137
V _L CDR1	24-34	26-32	30-36	27-38	25-40
V _L CDR2	50-56	50-52	46-55	56-65	58-77
V _L CDR3	89-97	91-96	89-96	105-117	109-137

¹Residue numbering follows the nomenclature of Kabat et al., supra.

²Residue numbering follows the nomenclature of Chothia et al., supra.

³Residue numbering follows the nomenclature of MacCallum et al., supra.

⁴Residue numbering follows the nomenclature of Lefranc et al., supra.

⁵Residue numbering follows the nomenclature of Honegger and Plückthun, supra.

The expression “variable-domain residue-numbering as in Kabat” or “amino-acid-position numbering as in Kabat,” and variations thereof, refers to the numbering system used for heavy-chain variable domains or light-chain variable domains of the compilation of antibodies in Kabat et al., supra. Using this numbering system, the actual linear amino acid sequence may contain fewer or additional amino acids corresponding to a shortening of, or insertion into, a FR or CDR of the variable domain. For example, a heavy-chain variable domain may include a single amino acid insert (residue 52a according to Kabat) after residue 52 of H2 and inserted residues (e.g. residues 82a, 82b, and 82c, etc. according to Kabat) after heavy-chain FR residue 82. The Kabat numbering of residues may be determined for a given antibody by alignment at regions of homology of the sequence of the antibody with a “standard” Kabat numbered sequence.

In certain embodiments, the amino acid residues which encompass the CDRs of a single domain antibody (e.g., a single domain anti-TIGIT antibody disclosed herein) is defined according to the IMGT nomenclature in Lefranc et al., supra. In certain embodiments, the amino acid residues which encompass the CDRs of a full-length antibody (e.g., an anti-PD1 antibody disclosed herein) is defined according to the Kabat nomenclature in Kabat et al., supra. In certain embodiments, the numbering of the residues in an immunoglobulin heavy chain, e.g., in an Fc region, is that of the EU index as in Kabat et al., supra. The “EU index as in Kabat” refers to the residue numbering of the human IgG1 EU antibody.

“Framework” or “FR” refers to residues are those variable-domain residues other than the CDR residues as herein defined.

A “humanized” antibody refers to a chimeric antibody comprising amino acid residues from non-human CDRs/HVRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the HVRs/CDRs correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized antibody optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, e.g., a non-human antibody, refers to an antibody that has undergone humanization.

A “human antibody” is an antibody that possesses an amino-acid sequence corresponding to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies as disclosed herein. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. Human antibodies can be produced using various techniques known in the art, including phage-display libraries. Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991). Also available for the preparation of human monoclonal antibodies are methods described in Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985); Boerner et al., *J. Immunol.*, 147(1):86-95 (1991). See also van Dijk and van de Winkel, *Curr. Opin. Pharmacol.*, 5: 368-74 (2001). Human antibodies can be prepared by administering the antigen to a transgenic animal that has been modified to produce such antibodies in response to antigenic challenge, but whose endogenous loci have been disabled, e.g., immunized xenomice (see, e.g., U.S. Pat. Nos. 6,075,181 and 6,150,584 regarding XENOMOUSE™ technology). See also, for example, Li et al., *Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006) regarding human antibodies generated via a human B-cell hybridoma technology.

“Percent (%) amino acid sequence identity” or “homology” with respect to the polypeptide and antibody sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the polypeptide being compared, after aligning the sequences considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, Megalign (DNASTAR), or MUSCLE software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values

are generated using the sequence comparison computer program MUSCLE (Edgar, R.C., *Nucleic Acids Research* 32(5):1792-1797, 2004; Edgar, R.C., *BMC Bioinformatics* 5(1):113, 2004).

“Homologous” refers to the sequence similarity or sequence identity between two polypeptides or between two nucleic acid molecules. When a position in both of the two compared sequences is occupied by the same base or amino acid monomer subunit, e.g., if a position in each of two DNA molecules is occupied by adenine, then the molecules are homologous at that position. The percent of homology between two sequences is a function of the number of matching or homologous positions shared by the two sequences divided by the number of positions compared times 100. For example, if 6 of 10 of the positions in two sequences are matched or homologous then the two sequences are 60% homologous. By way of example, the DNA sequences ATTGCC and TATGGC share 50% homology. Generally, a comparison is made when two sequences are aligned to give maximum homology.

The term “constant domain” refers to the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable domain, which contains the antigen-binding site. The constant domain contains the C_H1, C_H2 and C_H3 domains (collectively, C_H) of the heavy chain and the C_L domain of the light chain.

The “light chains” of antibodies (e.g., immunoglobulins) from any mammalian species can be assigned to one of two clearly distinct types, called kappa (“κ”) and lambda (“λ”), based on the amino acid sequences of their constant domains.

The “CH1 domain” (also referred to as “C1” or “H1” domain) usually extends from about amino acid 118 to about amino acid 215 (EU numbering system).

“Hinge region” is generally defined as a region in IgG corresponding to Glu216 to Pro230 of human IgG1 (Burton, *Molec. Immunol.* 22:161-206 (1985)). Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the first and last cysteine residues forming inter-heavy chain S-S bonds in the same positions.

The “CH2 domain” of a human IgG Fc region (also referred to as “C2” domain) usually extends from about amino acid 231 to about amino acid 340. The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are interposed between the two CH2 domains of an intact native IgG molecule. It has been speculated that the carbohydrate may provide a substitute for the domain-domain pairing and help stabilize the CH2 domain. Burton, *Molec Immunol.* 22:161-206 (1985).

The “CH3 domain” (also referred to as “C2” domain) comprises the residues between a CH2 domain and the C-terminal of an Fc region (i.e. from about amino acid residue 341 to the C-terminal end of an antibody sequence, typically at amino acid residue 446 or 447 of an IgG).

The term “Fc region” or “fragment crystallizable region” herein is used to define a C-terminal region of an immunoglobulin heavy chain, including native-sequence Fc regions and

variant Fc regions. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy-chain Fc region is usually defined to stretch from an amino acid residue at position Cys226, or from Pro230, to the carboxyl-terminus thereof. The C-terminal lysine (residue 447 according to the EU numbering system) of the Fc region may be removed, for example, during production or purification of the antibody, or by recombinantly engineering the nucleic acid encoding a heavy chain of the antibody. Accordingly, a composition of intact antibodies may comprise antibody populations with all K447 residues removed, antibody populations with no K447 residues removed, and antibody populations having a mixture of antibodies with and without the K447 residue. Suitable native-sequence Fc regions for use in the antibodies described herein include human IgG1, IgG2 (IgG2A, IgG2B), IgG3 and IgG4.

“Fc receptor” or “FcR” describes a receptor that binds the Fc region of an antibody. The preferred FcR is a native human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors, FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibitory receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (See M. Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9: 457-92 (1991); Capel et al., *Immunomethods* 4: 25-34 (1994); and de Haas et al., *J. Lab. Clin. Med.* 126: 330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein.

The term “epitope” as used herein refers to the specific group of atoms or amino acids on an antigen to which an antibody or antigen-binding moiety binds. Two antibodies or antigen-binding moieties may bind the same epitope within an antigen if they exhibit competitive binding for the antigen.

As use herein, the terms “specifically binds,” “specifically recognizing,” and “is specific for” refer to measurable and reproducible interactions, such as binding between a target and an antibody or antibody moiety, which is determinative of the presence of the target in the presence of a heterogeneous population of molecules, including biological molecules. For example, an antibody or antibody moiety that specifically recognizes a target (which can be an epitope) is an antibody or antibody moiety that binds this target with greater affinity, greater avidity, greater readiness, and/or greater duration than its bindings to other targets. In some embodiments, the extent of binding of an antibody to an unrelated target is less than about 10% of the binding of the antibody to the target as measured, e.g., by a radioimmunoassay (RIA). In some embodiments, an antibody that specifically

binds a target has a dissociation constant (K_D) of $\leq 10^{-5}$ M, $\leq 10^{-6}$ M, $\leq 10^{-7}$ M, $\leq 10^{-8}$ M, $\leq 10^{-9}$ M, $\leq 10^{-10}$ M, $\leq 10^{-11}$ M, or $\leq 10^{-12}$ M. In some embodiments, an antibody specifically binds an epitope on a protein that is conserved among the protein from different species. In some embodiments, specific binding can include, but does not require exclusive binding. Binding specificity of the antibody or antigen-binding domain can be determined experimentally by methods known in the art. Such methods comprise, but are not limited to Western blots, ELISA-, RIA-, ECL-, IRMA-, EIA-, BIACORE™-tests and peptide scans.

An “isolated” antibody (or construct) is one that has been identified, separated and/or recovered from a component of its production environment (e.g., natural or recombinant). In certain embodiments, the isolated polypeptide is free or substantially free from association with all other components from its production environment.

An “isolated” nucleic acid molecule encoding a construct, antibody, or antigen-binding fragment thereof described herein is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the environment in which it was produced. In certain embodiments, the isolated nucleic acid is free or substantially free from association with all components associated with the production environment. The isolated nucleic acid molecules encoding the polypeptides and antibodies described herein is in a form other than in the form or setting in which it is found in nature. Isolated nucleic acid molecules therefore are distinguished from nucleic acid encoding the polypeptides and antibodies described herein existing naturally in cells. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location.

The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by

ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term “vector,” as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Certain vectors are capable of directing the expression of nucleic acids to which they are operatively linked. Such vectors are referred to herein as “expression vectors.”

The term “transfected” or “transformed” or “transduced” as used herein refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A “transfected” or “transformed” or “transduced” cell is one which has been transfected, transformed or transduced with exogenous nucleic acid, which cell includes the primary subject cell and its progeny.

The terms “host cell,” “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell and may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

The terms “subject,” “individual,” and “patient” are used interchangeably herein to refer to a mammal, including, but not limited to, human, bovine, horse, feline, canine, rodent, or primate. In some embodiments, the subject is a human.

An “effective amount” of an agent refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result. The specific dose may vary depending on one or more of the particular agent chosen, the dosing regimen to be followed, whether it is administered in combination with other compounds, timing of administration, the tissue to be imaged, and the physical delivery system in which it is carried.

A “therapeutically effective amount” of a substance/molecule of the application, agonist or antagonist may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the substance/molecule, agonist or antagonist to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the substance/molecule, agonist or antagonist are outweighed by the therapeutically beneficial effects. A therapeutically effective amount may be delivered in one or more administrations.

A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, but not necessarily,

since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

As used herein, “treatment” or “treating” is an approach for obtaining beneficial or desired results, including clinical results. For purposes of this application, beneficial or desired clinical results include, but are not limited to, one or more of the following: alleviating one or more symptoms resulting from the disease, diminishing the extent of the disease, stabilizing the disease (e.g., preventing or delaying the worsening of the disease), preventing or delaying the spread (e.g., metastasis) of the disease, preventing or delaying the recurrence of the disease, delaying or slowing the progression of the disease, ameliorating the disease state, providing a remission (partial or total) of the disease, decreasing the dose of one or more other medications required to treat the disease, delaying the progression of the disease, increasing or improving the quality of life, increasing weight gain, and/or prolonging survival. Also encompassed by “treatment” is a reduction of pathological consequence of cancer (such as, for example, tumor volume). The methods of the application contemplate any one or more of these aspects of treatment. “Treatment” does not necessarily mean that the condition being treated will be cured.

It is understood that embodiments of the application described herein include “consisting” and/or “consisting essentially of” embodiments.

As used herein, the term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. In certain embodiments, “about” can mean within 3 or more than 3 standard deviations, per the practice in the art. In certain embodiments, “about” can mean a range of up to 20%, e.g., up to 10%, up to 5%, or up to 1% of a given value. In certain embodiments, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, e.g., within 5-fold or within 2-fold, of a value.

As used herein, the term “modulate” means positively or negatively alter. Exemplary modulations include a about 1%, about 2%, about 5%, about 10%, about 25%, about 50%, about 75%, or about 100% change.

As used herein, the term “increase” means alter positively by at least about 5%. An alteration may be by about 5%, about 10%, about 25%, about 30%, about 50%, about 75%, about 100% or more.

As used herein, the term “reduce” means alter negatively by at least about 5%. An alteration may be by about 5%, about 10%, about 25%, about 30%, about 50%, about 75%, or even by about 100%.

The term “about X-Y” used herein has the same meaning as “about X to about Y.”

As used herein and in the appended claims, the singular forms “a,” “or,” and “the” include plural referents unless the context clearly dictates otherwise.

“Effector functions” refer to those biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity (CDC), Fc receptor binding, antibody-dependent cell-mediated cytotoxicity (ADCC), phagocytosis, down regulation of cell surface receptors (e.g., B cell receptor), and B cell activation.

An “immunoconjugate” refers to an antibody conjugated to one or more heterologous molecule(s), including but not limited to a cytotoxic agent.

The term “pharmaceutical formulation” refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

A “pharmaceutically acceptable carrier,” as used herein, refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer, excipient, stabilizer, or preservative.

The term “variable region” or “variable domain” refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. In certain embodiments, the variable domains of the heavy chain and light chain (VH and VL, respectively) of a native antibody generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three CDRs. (See, e.g., Kindt et al. Kuby Immunology, 61ed., W.H. Freeman and Co., page 91 (2007).) A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody that binds the antigen to screen a library of complementary VL or VH domains, respectively. See, e.g., Portolano et al., J. Immunol. 150:880-887 (1993); Clarkson et al., Nature 352:624-628 (1991).

2. ANTIBODIES

In certain embodiments, the invention is based, in part, on the discovery of single domain antibodies that bind to TIGIT, which can be used in antitumor therapeutics where the antibodies selectively inhibit the TIGIT receptor and induce beneficial immune response of immune cells, e.g., T cells. Accordingly, the present disclosure provides anti-TIGIT antibodies. In certain embodiments, an anti-TIGIT antibody disclosed herein is an antagonist antibody, which inhibits TIGIT receptor functions. In certain embodiments, the anti-TIGIT antibody blocks an interaction between a TIGIT receptor and a ligand. In certain embodiments, the anti-TIGIT antibody blocks an

immune inhibitory signal from a TIGIT receptor. In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody, e.g., a camelid antibody or a VHH antibody. In certain embodiments, the anti-TIGIT antibody has an improved capability of tissue infiltration due to its smaller size compared to traditional antibodies in IgG, Fab and/or scFv forms.

5 The present disclosure further provides anti-PDL1 antibodies and multispecific antibodies that bind to both TIGIT and PD1 (referred to herein as anti-TIGIT/anti-PD1 antibodies. In certain embodiments, an antibody of the present disclosure can be or comprise a monoclonal antibody, including a chimeric, humanized or human antibody. In certain embodiments, the antibody disclosed herein comprises a humanized antibody. In certain embodiments, the antibody comprises
10 an acceptor human framework, e.g., a human immunoglobulin framework or a human consensus framework.

In certain embodiments, an antibody of the present disclosure can be an antibody fragment, e.g., a Fv, Fab, Fab', scFv, diabody, or F(ab')₂ fragment. In certain embodiments, the antibody is a full-length antibody, e.g., an intact IgG 1 antibody, or other antibody class or isotype as defined
15 herein. In certain embodiments, an antibody of the present disclosure can incorporate any of the features, singly or in combination, as described in this application, e.g., Sections 2.1-2.13 detailed herein.

Antibodies of the present disclosure are useful, e.g., for the diagnosis or treatment of a neoplasm or a cancer. In certain embodiments, the neoplasia and cancers whose growth may be
20 inhibited using the antibodies of this disclosure include neoplasia and cancers typically responsive to immunotherapy. In certain embodiments, the neoplasia and cancers include breast cancer (e.g., breast cell carcinoma), ovarian cancer (e.g., ovarian cell carcinoma) and renal cell carcinoma (RCC). Examples of other cancers that may be treated using the methods of this disclosure include melanoma (e.g., metastatic malignant melanoma), prostate cancer, colon cancer, lung cancer, bone
25 cancer, pancreatic cancer, skin cancer, brain tumors, chronic or acute leukemias including acute myeloid leukemia, chronic myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, lymphomas (e.g., Hodgkin's and non-Hodgkin's lymphoma, lymphocytic lymphoma, primary CNS lymphoma, T-cell lymphoma) nasopharyngeal carcinomas, cancer of the head or neck, cutaneous or intraocular malignant melanoma, uterine cancer, rectal cancer, cancer of the anal
30 region, stomach cancer, testicular cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the breast gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, cancer of the bladder,
35 cancer of the kidney or ureter, carcinoma of the breast pelvis, neoplasm of the central nervous system (CNS), tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma,

Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, environmentally induced cancers including those induced by asbestos, e.g., mesothelioma and combinations of said cancers.

2.1 Exemplary anti-TIGIT antibodies

5 The present disclosure provides isolated antibodies that bind to a TIGIT protein. In certain embodiments, an anti-TIGIT antibody of the present disclosure binds to the ECD of TIGIT. In certain embodiments, the anti-TIGIT antibody binds to the ECD of TIGIT that comprises the amino acid sequence set forth in SEQ ID NO: 262. In certain embodiments, the anti-TIGIT antibody binds to the same epitope as an anti-TIGIT antibody, e.g., 2A3, described herein.

10 In certain embodiments, the anti-TIGIT antibody disclosed herein can function as an antagonist of a TIGIT receptor. In certain embodiments, the anti-TIGIT antibody can reduce the activity of the TIGIT receptor by at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 99% or about 99.9%. In certain embodiments, the anti-TIGIT antibody can block the downstream immune inhibitory signaling of the TIGIT
15 receptor. In certain embodiments, the anti-TIGIT antibody increases an immune response and/or an antitumor effect of an immune cell, e.g., a T cell and/or a NK cell. In certain embodiments, treatment using the anti-TIGIT antibody exhibits antitumor efficacy in a subject, whereby reduces tumor growth and/or lengthen the survival of a subject. In certain embodiments, the anti-TIGIT antibody comprising a single domain antibody (e.g., a VHH) has a smaller molecule size compared
20 to a full-length antibody due to the smaller size of a single domain antibody compared to a Fab domain of a full-length antibody, which can result in superior tissue infiltration, e.g., at a tumor site, compared to a full-length antibody. In certain embodiments, treatment using the anti-TIGIT antibody exhibits superior antitumor efficacy compared to treatment using a full-length anti-TIGIT antibody, e.g., Reference Ab 1 having the same amino acid sequences of BMS 22G2 disclosed in
25 U.S. 2016/0176963 A1 and Reference Ab 2 having the same amino acid sequences of Tiragolumab, which sequences are disclosed in U.S. 2017/0088613 A1.

In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody that binds to TIGIT. In certain embodiments, the single domain antibody comprises a VHH. In certain
30 embodiments, the single domain antibody comprises a heavy chain variable region (VH). In certain embodiments, the single domain antibody is linked to a Fc region. In certain embodiments, the single domain antibody is not linked to a Fc region.

In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-7} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-8} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a
35 KD of about 5×10^{-9} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-9} M or less. In certain embodiments, the single domain antibody binds to

TIGIT with a KD of between about 1×10^{-9} M and about 1×10^{-7} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 1×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 2×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 2×10^{-9} M and about 5×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 1×10^{-9} M and about 5×10^{-9} M.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set

forth in SEQ ID NO: 116. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144. In certain embodiments, the single domain antibody cross-competes for

binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 151, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 152. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 154, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 155, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 156. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain

variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising: a) a heavy chain variable region CDR1 comprises an amino acid sequence of any one of SEQ ID NOs: 94, 98, 102, 106, 110, 114, 118, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186 and 190, or a variant thereof comprising up to about 3 amino acid substitutions; b) a heavy chain variable region CDR2 comprises an amino acid sequence of any one of SEQ ID NOs: 95, 99, 103, 107, 111, 115, 119, 123, 127, 131, 135, 139, 143, 147, 151, 155, 159, 163, 167, 171, 175, 179, 183, 187 and 191, or a variant thereof comprising up to about 3 amino acid substitutions; and c) a heavy chain variable region CDR3 comprises an amino acid sequence of any one of SEQ ID NOs: 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188 and 192, or a variant thereof comprising up to about 3 amino acid substitutions.

In certain embodiments, the single domain antibody comprises a heavy chain variable region that comprises a CDR1 domain, a CDR2 domain and a CDR3 domain, wherein the CDR1 domain, the CDR2 domain and the CDR3 domain respectively comprise a CDR1 domain, a CDR2 domain and a CDR3 domain comprised in a reference heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120. In certain embodiments, the single domain antibody comprises a heavy chain variable

[illegible]

sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 5 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 10 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 97. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 101. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 105. In certain 15 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 109. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 113. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 117. In certain 20 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 121. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 125. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 129. In certain 25 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 133. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 137. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 141. In certain 30 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 145. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 149. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 153. In certain 35 embodiments, the single domain antibody comprises a heavy chain variable region comprising the

amino acid sequence set forth in SEQ ID NO: 157. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 161. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 165. In certain
5 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 169. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 173. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 177. In certain
10 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 181. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 185. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 189. In certain
15 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain
20 embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and
25 a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and
30 a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10,
35 and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11. In certain embodiments, the single domain antibody cross-competes for binding to

TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 15. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 17, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 18, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 19. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 21, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 22, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 23. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 25, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 26, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 27. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 29, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 30, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 31. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 33, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 34, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 35. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 37, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 38, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 39. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable

region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 41, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 42, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 43. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 45, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 46, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 47. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 49, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 50, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 51. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 53, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 54, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 55. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 57, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 58, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 59. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 61, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 62, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 63. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 65, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 66, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 67. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 69, a heavy

chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 70, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 71. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:

[illegible]

[illegible]

79. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 4. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 8. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 12. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 16. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 20. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 24. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 28. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 32. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 36. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 40. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 44. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 48. In certain

embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 52. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 56. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 60. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 64. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 68. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 72. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 76. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 80. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84.

In certain embodiments, the single domain antibody comprises a humanized framework. In certain embodiments, the humanized framework comprises a framework sequence of the heavy chain variable region sequences selected from the group consisting of SEQ ID NOs: 85-93. In certain embodiments, the humanized framework comprises a FR2 sequence of the heavy chain variable region sequences selected from the group consisting of SEQ ID NOs: 85-93.

In certain embodiments, the anti-TIGIT antibody does not comprise a Fc region. In certain embodiments, the anti-TIGIT antibody further comprises a Fc region. In certain embodiments, the Fc region comprises a human Fc region. In certain embodiments, the Fc region comprises a Fc region selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM. In certain embodiments, the Fc region comprises a Fc region selected from the group consisting of the Fc regions of IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region comprises an IgG1 Fc region. In certain embodiments, the IgG1 Fc region comprising one or more mutation that enhances an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprises the mutations of L235V, F243L, R292P, Y300L and P396L. In certain embodiments, the IgG1 Fc region comprises the mutations of S239D, A330L and I332E. In certain embodiments, the anti-TIGIT antibody comprises the amino acid sequence set forth in SEQ ID NO: 194

In certain embodiments, the heavy chain variable region is linked to a Fc region via a linker. In certain embodiments, the linker is a peptide linker. In certain embodiments, the peptide linker comprises about four to about thirty amino acids. In certain embodiments, the peptide linker

comprises about four to about fifteen amino acids. In certain embodiments, the peptide linker comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220.

In certain embodiments, the Fc region is capable of dimerization. In certain embodiments, the Fc region can dimerize through one or more disulfide bond. In certain embodiments, the anti-TIGIT antibody comprises two dimerized heavy chains, each of the heavy chain comprising a heavy chain variable region and a Fc region capable of dimerization. In certain embodiments, the anti-TIGIT antibody comprises two disulfide bonds. In certain embodiments, the anti-TIGIT antibody comprises three disulfide bonds. In certain embodiments, the anti-TIGIT antibody does not comprise a light chain.

In certain embodiments, the anti-TIGIT antibody can be a multivalent antibody. In certain embodiments, the anti-TIGIT antibody can be monovalent, bivalent, trivalent, tetravalent, pentavalent, hexavalent, heptavalent or octavalent. In certain embodiments, the anti-TIGIT antibody is monovalent. In certain embodiments, the anti-TIGIT antibody is bivalent. In certain embodiments, the anti-TIGIT antibody is tetravalent.

In certain embodiments, the anti-TIGIT antibody comprises a multispecific antibody, e.g., a bispecific antibody, a full-length immunoglobulin, a single-chain Fv (scFv) fragment, a Fab fragment, a Fab' fragment, a F(ab')₂, an Fv fragment, a disulfide stabilized Fv fragment (dsFv), a (dsFv)₂, a VHH, a Fv-Fc fusion, a scFv-Fc fusion, a scFv-Fv fusion, a diabody, a tribody, a tetrabody or any combination thereof. In certain embodiments, the antibody comprises a multispecific antibody, e.g., a bispecific antibody, which comprises a second antibody moiety that specifically binds to a second antigen.

In certain embodiments, the second antigen is a tumor associated antigen. In certain embodiments, the tumor associated antigen is selected from the group consisting of Her-2, EGFR, PD-L1, c-Met, B Cell Maturation Antigen (BCMA), carbonic anhydrase IX (CAIX), carcinoembryonic antigen (CEA), CD5, CD7, CD10, CD19, CD20, CD22, CD30, CD33, CD34, CD38, CD41, CD44, CD49f, CD56, CD74, CD123, CD133, CD138, CD276 (B7H3), epithelial glycoprotein (EGP2), trophoblast cell-surface antigen 2 (TROP-2), epithelial glycoprotein-40 (EGP-40), epithelial cell adhesion molecule (EpCAM), receptor tyrosine-protein kinases erb-B2,3,4, folate-binding protein (FBP), fetal acetylcholine receptor (AChR), folate receptor- α , Ganglioside G2 (GD2), Ganglioside G3 (GD3), human telomerase reverse transcriptase (hTERT), kinase insert domain receptor (KDR), Lewis A (CA 19.9), Lewis Y (LeY), Glypican-3 (GPC3), L1 cell adhesion molecule (L1CAM), Mucin 16 (Muc-16), Mucin 1 (Muc-1), NG2D ligands, oncofetal antigen (h5T4), prostate stem cell antigen (PSCA), prostate-specific membrane antigen (PSMA), tumor-associated glycoprotein 72 (TAG-72), Claudin18.2 (CLDN18.2), vascular endothelial growth factor R2 (VEGF- R2), Wilms tumor protein (WT-1), type 1 tyrosine-protein kinase transmembrane receptor (ROR1) and any combination thereof.

In certain embodiments, the anti-TIGIT antibody is conjugated to a therapeutic agent or a label. In certain embodiments, the label is selected from the group consisting of a radioisotope, a fluorescent dye and an enzyme.

2.2 Exemplary anti-PD1 antibodies

The present disclosure further provides anti-PD1 antibodies. In certain embodiments, an anti-PD1 antibody disclosed herein binds to a PD1 protein with high affinity. In certain embodiments, the anti-PD1 antibody blocks the interaction between PDL1 and PD1. In certain embodiments, the antibody is any anti-PD1 antibody disclosed in PCT/US2017/050851 (published as International Publication WO 2018/052818 A1), the content of which is incorporated herein by reference in its entirety.

In certain embodiments, the anti-PD1 antibody cross-competes with a reference anti-PD1 antibody that comprises: a) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226); b) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO: 231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234); c) a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242); or d) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence

SSYYYGIDL (SEQ ID NO:247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO:248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

5 In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226). In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO:231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234). In certain
10 20 25 30 35
embodiments, the anti-PD1 antibody comprises a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO:239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO:240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO:242). In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO:247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO:248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%,

95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 227, 235, 243 and 251, and a light chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 228, 236, 244 and 252. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 227, 235, 243 and 251, and a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 228, 236, 244 and 252.

In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 227, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 228. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 235, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 236. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 243, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 244. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 251, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 252.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region and/or the light chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the anti-PD1 antibody comprises a Fc region. In certain embodiments, the Fc region is selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM. In certain embodiments, the Fc region is selected from the group consisting of the Fc regions of IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region comprises a human Fc region. In certain embodiments, the Fc region comprises an IgG1 Fc region. In certain embodiments, the IgG1 Fc region comprising one or more mutation that modifies an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that enhances an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that reduces an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the Fc region comprises an IgG4 Fc region. In certain embodiments, the IgG4 Fc region comprises an S228P mutation.

In certain embodiments, the anti-PD1 antibody comprises a humanized antibody. In certain embodiments, the anti-PD1 antibody is identified by screening of a human phage display library with a polypeptide comprising the ECD of a human PD1 protein.

2.3 Multispecific antibodies

The present disclosure further provides multispecific antibodies, e.g., a bispecific antibody. Multispecific antibodies are monoclonal antibodies that have binding specificities for at least two different epitopes. In certain embodiments, one of the binding specificities is for an epitope present on TIGIT and the other is for any other antigen. In certain embodiments, one of the binding specificities is for an epitope present on PD1 and the other is for any other antigen. In certain embodiments, a multispecific antibody of the present disclosure can bind to an epitope on TIGIT and can bind to an epitope on PD1. In certain embodiments, a multispecific antibody of the present disclosure can comprise a full-length antibody, an antibody fragment and/or any combination thereof.

In certain embodiments, the multispecific antibody binds to TIGIT and PD1. In certain embodiments, the multispecific antibody is a bispecific, anti-TIGIT/anti-PD1 antibodies. In certain embodiments, the multispecific antibody has at least two different binding specificities, see, e.g., U.S. Patent Nos. 5,922,845 and 5,837,243; Zeilder (1999) J. Immunol. 163: 1246-1 252; Somasundaram (1999) Hum. Antibodies 9:47-54; Keler (1997) Cancer Res. 57:4008-401 4. For example, and not by way of limitation, the presently disclosed subject matter provides multispecific antibodies comprising one antigen-binding moiety for a first epitope present on TIGIT and a second antigen-binding moiety for a second epitope present on PDL1. In certain embodiments, the multispecific antibody comprises a first antigen-binding moiety comprising an anti-TIGIT antibody disclosed herein; and a second antigen-binding moiety comprising an anti-PD1 antibody disclosed herein.

In certain embodiments, the anti-TIGIT/anti-PD1 antibody disclosed herein can function as an antagonist of both a TIGIT receptor and PD1/PDL1 signaling. In certain embodiments, the anti-TIGIT/anti-PD1 antibody blocks immune checkpoint inhibition of an immune cell, e.g., a T cell or a NK cell. In certain embodiments, without bound by any theory, the anti-PD1 moiety of the anti-TIGIT/anti-PD1 antibody can guide and/or concentrate the antibody at a tumor site, whereby enhances the antitumor functions of an immune cell at the vicinity of the tumor site and/or reduces the toxicity and side effects of a peripheral immune cells. In certain embodiments, without bound by any theory, the anti-TIGIT/anti-PD1 antibody can ligate a tumor cells expressing PD1 and an immune cell, e.g., a T cell or a NK cell, that expresses TIGIT, whereby brings the immune cell to the vicinity of a tumor site and facilitates tumor eradication. In certain embodiments, treatment using the anti-TIGIT/anti-PD1 antibody exhibits superior antitumor efficacy compared to treatment

using a monospecific anti-TIGIT antibody and/or a monospecific anti-PD1 antibody. In certain embodiments, treatment using the anti-TIGIT/anti-PD1 antibody exhibits superior antitumor efficacy compared to treatment using a combination of a monospecific anti-TIGIT antibody and a monospecific anti-PD1 antibody.

5 In certain embodiments, the anti-TIGIT/anti-PD1 antibody comprises a first antigen-binding moiety comprising an anti-TIGIT antibody comprising a single domain antibody that binds to TIGIT, and a second antigen-binding moiety comprising an anti-PD1 antibody that binds to PD1.

2.3.1 First antigen-binding moiety

10 In certain embodiments, the first antigen-binding moiety of a multispecific anti-TIGIT/anti-PD1 antibody disclosed herein comprises an anti-TIGIT antibody disclosed herein. In certain embodiments, an anti-TIGIT antibody of the present disclosure binds to the ECD of TIGIT. In certain embodiments, the anti-TIGIT antibody binds to the ECD of TIGIT that comprises the amino acid sequence set forth in SEQ ID NO: 262. In certain embodiments, the anti-TIGIT antibody binds
15 to the same epitope as an anti-TIGIT antibody, e.g., 2A3, described herein.

In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody that binds to TIGIT. In certain embodiments, the single domain antibody comprises a VHH. In certain embodiments, the single domain antibody comprises a heavy chain variable region (VH). In certain embodiments, the single domain antibody is linked to a Fc region.

20 In certain embodiments, the anti-TIGIT antibody disclosed herein can function as an antagonist of a TIGIT receptor. In certain embodiments, the anti-TIGIT antibody can reduce the activity of the TIGIT receptor by at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 99% or about 99.9%. In certain embodiments, the anti-TIGIT antibody can block the downstream immune inhibitory signaling of the TIGIT
25 receptor. In certain embodiments, the anti-TIGIT antibody increases an immune response and/or an antitumor effect of an immune cell, e.g., a T cell and/or a NK cell. In certain embodiments, treatment using the anti-TIGIT antibody exhibits antitumor efficacy in a subject, whereby reduces tumor growth and/or lengthen the survival of a subject. In certain embodiments, the anti-TIGIT antibody comprising a single domain antibody (e.g., a VHH) has a smaller molecule size compared
30 to a full-length antibody due to the smaller size of a single domain antibody compared to a Fab domain of a full-length antibody, which can result in superior tissue infiltration, e.g., at a tumor site, compared to a full-length antibody. In certain embodiments, treatment using the anti-TIGIT antibody exhibits superior antitumor efficacy compared to treatment using a full-length anti-TIGIT antibody, e.g., Reference Ab 1 having the same amino acid sequences of BMS 22G2 disclosed in
35 U.S. 2016/0176963 A1 and Reference Ab 2 having the same amino acid sequences of Tiragolumab, which sequences are disclosed in U.S. 2017/0088613 A1.

In certain embodiments, the anti-TIGIT antibody comprises a single domain antibody that binds to TIGIT. In certain embodiments, the single domain antibody comprises a VHH. In certain embodiments, the single domain antibody comprises a heavy chain variable region (VH). In certain embodiments, the single domain antibody is linked to a Fc region. In certain embodiments, the single domain antibody is not linked to a Fc region.

In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-7} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-8} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 5×10^{-9} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of about 1×10^{-9} M or less. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 1×10^{-9} M and about 1×10^{-7} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 1×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 2×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 2×10^{-9} M and about 5×10^{-8} M. In certain embodiments, the single domain antibody binds to TIGIT with a KD of between about 1×10^{-9} M and about 5×10^{-9} M.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set

forth in SEQ ID NO: 108. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136. In certain embodiments, the single domain antibody cross-competes for

binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 151, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 152. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 154, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 155, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 156. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain

variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising: a) a heavy chain variable region CDR1 comprises an amino acid sequence of any one

of SEQ ID NOs: 94, 98, 102, 106, 110, 114, 118, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186 and 190, or a variant thereof comprising up to about 3 amino acid substitutions; b) a heavy chain variable region CDR2 comprises an amino acid sequence of any one of SEQ ID NOs: 95, 99, 103, 107, 111, 115, 119, 123, 127, 131, 135, 139, 143, 147, 151, 155, 159, 163, 167, 171, 175, 179, 183, 187 and 191, or a variant thereof comprising up to about 3 amino acid substitutions; and c) a heavy chain variable region CDR3 comprises an amino acid sequence of any one of SEQ ID NOs: 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188 and 192, or a variant thereof comprising up to about 3 amino acid substitutions.

In certain embodiments, the single domain antibody comprises a heavy chain variable region that comprises a CDR1 domain, a CDR2 domain and a CDR3 domain, wherein the CDR1 domain, the CDR2 domain and the CDR3 domain respectively comprise a CDR1 domain, a CDR2 domain and a CDR3 domain comprised in a reference heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112. In certain embodiments,

the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116. In certain

5 embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120. In certain embodiments, the single domain antibody comprises a heavy chain variable

10 region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a

15 heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in

20 SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino

25 acids having the sequence set forth in SEQ ID NO: 136. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140. In certain embodiments,

30 the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144. In certain

embodiments, the single domain antibody comprises a heavy chain variable region CDR1

35 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a

heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188. In certain embodiments, the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 97. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 101. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 105. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 109. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 113. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 117. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 121. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 125. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 129. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 133. In certain embodiments, the single domain

antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 137. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 141. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 145. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 149. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 153. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 157. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 161. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 165. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 169. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 173. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 177. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 181. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 185. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 189. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region

CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 15. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 17, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 18, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 19. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 21, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 22, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 23. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 25, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 26, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 27. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 29, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 30, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 31. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 33, a heavy

chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 34, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 35. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 37, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 38, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 39. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 41, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 42, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 43. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 45, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 46, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 47. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 49, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 50, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 51. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 53, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 54, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 55. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 57, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 58, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 59. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 61, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO:

62, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 63. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 65, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 66, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 67. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 69, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 70, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 71. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79. In certain embodiments, the single domain antibody cross-competes for binding to TIGIT with a reference anti-TIGIT single domain antibody comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO:

[illegible]

[illegible]

71. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 4. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 8. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 12. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 16. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 20. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 24. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the

amino acid sequence set forth in SEQ ID NO: 28. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 32. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 36. In certain
5 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 40. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 44. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 48. In certain
10 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 52. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 56. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 60. In certain
15 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 64. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 68. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 72. In certain
20 embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 76. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 80. In certain embodiments, the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84.

25 In certain embodiments, the single domain antibody comprises a humanized framework. In certain embodiments, the humanized framework comprises a framework sequence of the heavy chain variable region sequences selected from the group consisting of SEQ ID NOs: 85-93. In certain embodiments, the humanized framework comprises a FR2 sequence of the heavy chain variable region sequences selected from the group consisting of SEQ ID NOs: 85-93.

30 2.3.2 Second antigen-binding moiety

In certain embodiments, the second antigen-binding moiety of a multispecific anti-TIGIT/anti-PD1 antibody disclosed herein comprises an anti-PD1 antibody that binds to PD1. In certain embodiments, the anti-PD1 antibody blocks the interaction between PD1 and PD1. In
35 certain embodiments, the antibody is any anti-PD1 antibody disclosed in PCT/US2017/050851

(published as International Publication WO 2018/052818 A1), the content of which is incorporated herein by references in its entirety.

In certain embodiments, the anti-PD1 antibody cross-competes with a reference anti-PD1 antibody that comprises: a) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226); b) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO: 231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234); c) a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242); or d) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO: 247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO: 248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226).

223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226). In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO: 231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234). In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242). In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO: 247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO: 248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 227, 235, 243 and 251, and a light chain variable region comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 228, 236, 244 and 252. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 227, 235, 243 and 251, and a light chain variable region

comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 228, 236, 244 and 252.

In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 227, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 228. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 235, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 236. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 243, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 244. In certain embodiments, the anti-PD1 antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 251, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 252.

In certain embodiments, any one of the amino acid sequences comprised in the heavy chain variable region and/or the light chain variable region can comprise up to about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9 or about 10 amino acid substitutions, deletions and/or additions. In certain embodiments, the amino acid substitution is a conservative substitution.

In certain embodiments, the anti-PD1 antibody comprises a Fc region. In certain embodiments, the Fc region is selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM. In certain embodiments, the Fc region is selected from the group consisting of the Fc regions of IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region comprises a human Fc region. In certain embodiments, the Fc region comprises an IgG1 Fc region. In certain embodiments, the IgG1 Fc region comprising one or more mutation that modifies an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that enhances an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that reduces an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the Fc region comprises an IgG4 Fc region. In certain embodiments, the IgG4 Fc region comprises an S228P mutation.

In certain embodiments, the anti-PD1 antibody comprises a humanized antibody. In certain embodiments, the anti-PD1 antibody is identified by screening of a human phage display library with a polypeptide comprising the ECD of a human PD1 protein.

2.3.3 Features of multispecific antibodies

In certain embodiments, the anti-TIGIT/anti-PD1 multispecific antibody can be a multivalent antibody. In certain embodiments, the anti-TIGIT/anti-PD1 antibody can be bivalent, trivalent,

tetravalent, pentavalent, hexavalent, heptavalent or octavalent. In certain embodiments, each of the first and the second antigen-binding moieties of the anti-TIGIT/anti-PD1 antibody can be monovalent, bivalent, trivalent, tetravalent, pentavalent, hexavalent, heptavalent or octavalent. In certain embodiments, each of the first and the second antigen-binding moieties is monovalent. In certain embodiments, each of the first and the second antigen-binding moieties is bivalent. In certain embodiments, the anti-TIGIT/anti-PD1 multispecific antibody is bivalent. In certain embodiments, the anti-TIGIT/anti-PD1 multispecific antibody is tetravalent.

In certain embodiments, the second antigen binding moiety comprises an anti-PD1 antibody comprising two antibody heavy chains and two antibody light chains. In certain embodiments, the first antigen-binding moiety comprises two or more anti-TIGIT antibodies. In certain embodiments, the first antigen-binding moiety comprises two anti-TIGIT antibodies. In certain embodiments, the C-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the C-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety. In certain embodiments, the N-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

In certain embodiments, the multispecific antibody comprises i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 229, (2) a CDR-H2 comprising the amino acid sequence set forth in SEQ ID NO: 230, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 231; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 232, (2) a CDR-L2 comprising the

amino acid sequence set forth in SEQ ID NO: 233, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 234.

In certain embodiments, the multispecific antibody comprises i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 237, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 238, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 239; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 240, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 241, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 242.

In certain embodiments, the multispecific antibody comprises i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 221, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 222, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 223; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 224, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 225, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 226.

In certain embodiments, the multispecific antibody comprises i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192; and ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 245, (2) a CDR- H2 comprising the amino acid

sequence set forth in SEQ ID NO: 246, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 247; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 248, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 249, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 250.

In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 253, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 254. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 255, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 256. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 257, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 258. In certain embodiments, the multispecific antibody comprises an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 259, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 260.

In certain embodiments, the first antigen binding moiety is linked to the second antigen binding moiety via a linker. In certain embodiments, the linker is a peptide linker. In certain embodiments, the peptide linker comprises about four to about thirty amino acids. In certain embodiments, the peptide linker comprises about four to about fifteen amino acids. In certain embodiments, the peptide linker comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220.

In certain embodiments, the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM. In certain embodiments, the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc regions of IgG1, IgG2, IgG3 and IgG4. In certain embodiments, the Fc region comprises a human Fc region. In certain embodiments, the Fc region comprises an IgG1 Fc region. In certain embodiments, the IgG1 Fc region comprising one or more mutation that modifies an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that enhances an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the IgG1 Fc region comprising one or more mutation that reduces an antibody-dependent cell-mediated cytotoxicity (ADCC). In certain embodiments, the Fc region comprises an IgG4 Fc region. In certain embodiments, the IgG4 Fc region comprises an S228P mutation.

2.4 Antibody Affinity

In certain embodiments, an antibody or an antigen-binding moiety of a multispecific antibody disclosed herein has a high binding affinity to its target antigen. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of about 1×10^{-7} M or less. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of about 1×10^{-8} M or less. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of about 5×10^{-9} M or less. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of about 1×10^{-9} M or less. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of between about 1×10^{-9} M and about 1×10^{-7} M. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of between about 1×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of between about 2×10^{-9} M and about 1×10^{-8} M. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of between about 2×10^{-9} M and about 5×10^{-8} M. In certain embodiments, the antibody or antigen-binding moiety binds to the target with a K_D of between about 1×10^{-9} M and about 5×10^{-9} M.

The K_D of the antibody or antigen-binding moiety can be determined by methods known in the art. Such methods comprise, but are not limited to Western blots, ELISA-, RIA-, ECL-, IRMA-, EIA-, Octet- BIACORE®-tests and peptide scans.

In certain embodiments, K_D can be measured using a BIACORE® surface plasmon resonance assay. For example, and not by way of limitation, an assay using a BIACORE®-2000 or a BIACORE® 3000 (Biacore, Inc., Piscataway, NJ) is performed at 25°C with immobilized antigen CMS chips at about 10 response units (RU). In certain embodiments, carboxymethylated dextran biosensor chips (CMS, Biacore, Inc.) are activated with N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (about 0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20™) surfactant (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIACORE® Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_D) can be calculated as the ratio k_{off}/k_{on} . See, e.g., Chen et al., J. Mol. Biol. 293:865-881 (1999). If the on-rate exceeds $10^6 \text{ M}^{-1} \text{ s}^{-1}$ by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase

or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCO™ spectrophotometer (ThermoSpectronic) with a stirred cuvette.

2.5 Antibody Fragments

In certain embodiments, an antibody of the present disclosure comprises an antigen-binding fragment or antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, see Hudson et al. Nat. Med. 9: 129-134 (2003). For a review of scFv fragments, see, e.g., Pluckthtin, in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., (Springer- Verlag, New York), pp. 269-315 (1994); see also WO 93/16185; and U.S. Patent Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab)₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Patent No. 5,869,046.

In certain embodiments, an antibody of the present disclosure can be a diabody. Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 404,097; WO 1993/01 161; Hudson et al., Nat. Med. 9:129-134 (2003); and Hollinger et al., Proc. Natl. Acad. Sci. USA 90: 6444-6448 (1993). Triabodies and tetrabodies are also described in Hudson et al., Nat. Med. 9: 129-134 (2003).

In certain embodiments, an antibody of the present disclosure can comprise a singledomain antibody. Singledomain antibodies are antibody fragments that comprise all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, the singledomain antibody is a human single-domain antibody (Domantis, Inc., Waltham, MA; see, e.g., U.S. Patent No. 6,248,516 B1). In certain embodiments, the singledomain antibody is camelid single-domain antibody. In certain embodiments, the singledomain antibody is a VHH. In certain embodiments, the singledomain antibody is humanized.

Antibody fragments can be made by various techniques including, but not limited to, proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g., E. coli or phage), as described herein.

2.6 Chimeric and Humanized Antibodies

In certain embodiments, an antibody, including an antigen-binding moiety of a multispecific antibody, of the present disclosure is a chimeric antibody. Certain chimeric antibodies are described, e.g., in U.S. Patent No. 4,816,567; and Morrison et al., Proc. Natl. Acad. Sci. USA, 81:6851-6855

(1984)). In certain embodiments, a chimeric antibody comprises a non-human variable region (e.g., a variable region derived from mouse) and a human constant region. In certain embodiments, a chimeric antibody is a “class switched” antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-binding fragments thereof.

5 In certain embodiments, an antibody, including an antigen-binding moiety of a multispecific antibody, of the present disclosure can be a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which HVRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or any portion thereof) are derived from human antibody sequences. A humanized antibody optionally can also comprise at least a portion of a human constant region. In certain
10 embodiments, certain FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from which the HVR residues are derived), e.g., to restore or improve antibody specificity or affinity.

15 Humanized antibodies and methods of making them are described, e.g., in Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008), and are further described, e.g., in Riechmann et al., *Nature* 332:323-329 (1988); Queen et al., *Proc. Nat’l Acad. Sci. USA* 86:10029-10033 (1989); US Patent Nos. 5, 821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri et al., *Methods* 36:25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, *Mol. Immunol.* 28:489-498 (1991) (describing “resurfacing”); Dall’Acqua et al., *Methods* 36:43-60 (2005) (describing “FR shuffling”); and Osbourn et al., *Methods* 36:61-68 (2005) and Klimka et al., *Br. J. Cancer*, 83:252-260 (2000) (describing the “guided selection” approach to FR shuffling).
20

Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the “best-fit” method (see, e.g., Sims et al. *J. Immunol.* 151:2296 (1993)); Framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, e.g., Carter et al. *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); and Presta et al. *J. Immunol.*, 151:2623 (1993)); human mature (somatically mutated) framework regions or human germline framework regions (see, e.g., Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008)); and framework regions derived from screening
25 FR libraries (see, e.g., Baca et al., *J. Biol. Chem.* 272:10678-10684 (1997) and Rosok et al., *J. Biol. Chem.* 271:22611-22618 (1996)).
30

2.7 Human Antibodies

35 In certain embodiments, an antibody of the present disclosure can be a human antibody (e.g., human domain antibody, or human DAb). Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk and van de

Winkel, Curr. Opin. Pharmacol. 5: 368-74 (2001), Lonberg, Curr. Opin. Immunol. 20:450-459 (2008), and Chen, Mol. Immunol. 47(4):912-21 (2010). Transgenic mice or rats capable of producing fully human single-domain antibodies (or DAb) are known in the art. See, e.g., US20090307787A1, U.S. Pat. No. 8,754,287, US20150289489A1, US20100122358A1, and
5 WO2004049794.

Human antibodies (e.g., human DABs) may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous immunoglobulin loci,
10 or which are present extrachromosomally or integrated randomly into the animal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of methods for obtaining human antibodies from transgenic animals, see Lonberg, Nat. Biotech. 23:1117-1125 (2005). See also, e.g., U.S. Patent Nos. 6,075,181 and 6,150,584 describing XENOMOUSE™ technology; U.S. Patent No. 5,770,429 describing HuMab® technology; U.S.
15 Patent No. 7,041,870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VelociMouse® technology). Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

Human antibodies (e.g., human DABs) can also be made by hybridoma-based methods.
20 Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described (See, e.g., Kozbor J. Immunol., 133: 3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner et al., J. Immunol., 147: 86 (1991)). Human antibodies generated via human B-cell hybridoma technology are also described in Li et al., Proc.
25 Natl. Acad. Sci. USA, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Patent No. 7,189,826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines) and Ni, Xiandai Mianyixue, 26(4):265-268 (2006) (describing human-human hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers and Brandlein, Histology and Histopathology, 20(3):927-937 (2005) and Vollmers and
30 Brandlein, Methods and Findings in Experimental and Clinical Pharmacology, 27(3):185-91 (2005).

Human antibodies (e.g., human DABs) may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

2.8 Library-Derived Antibodies

The antibody moieties may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are described, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O'Brien et al., ed., Human Press, Totowa, NJ, 2001) and further described, e.g., in the McCafferty et al., *Nature* 348:552-554; Clackson et al., *Nature* 352: 624-628 (1991); Marks et al., *J. Mol. Biol.* 222: 581-597 (1992); Marks and Bradbury, in *Methods in Molecular Biology* 248:161-175 (Lo, ed., Human Press, Totowa, NJ, 2003); Sidhu et al., *J. Mol. Biol.* 338(2): 299-310 (2004); Lee et al., *J. Mol. Biol.* 340(5): 1073-1093 (2004); Fellouse, *Proc. Natl. Acad. Sci. USA* 101(34): 12467-12472 (2004); and Lee et al., *J. Immunol. Methods* 284(1-2): 119-132(2004). Methods for constructing single-domain antibody libraries have been described, for example, see U.S. Pat. NO. 7371849.

In certain phage display methods, repertoires of V_H and V_L genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter et al., *Ann. Rev. Immunol.*, 12: 433-455 (1994). Phage typically displays antibody fragments, either as scFv fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self-antigens without any immunization as described by Griffiths et al., *EMBO J*, 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement in vitro, as described by Hoogenboom and Winter, *J. Mol. Biol.*, 227: 381-388 (1992). Patent publications describing human antibody phage libraries include, for example: US Patent No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/0160598, 2007/0237764, 2007/0292936, and 2009/0002360.

Antibodies or antibody fragments isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

2.9 Antibody Variants

The presently disclosed subject matter further provides amino acid sequence variants of the disclosed antibodies. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino acid sequence variants of an antibody can be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, but are not limited to, deletions from, and/or

insertions into and/or substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final antibody, i.e., modified, possesses the desired characteristics, e.g., antigen-binding.

5

2.9.1 Substitution, Insertion, and Deletion Variants

In certain embodiments, antibody variants having one or more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include the HVRs (or CDRs) and FRs. Conservative substitutions are shown in Table 2 under the heading of “Preferred substitutions.”
 10 More substantial changes are provided in Table 2 under the heading of “exemplary substitutions,” and as further described below in reference to amino acid side chain classes. Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen binding, decreased immunogenicity, or improved ADCC or CDC.

15

Table 2. Amino acid substitutions

Original	Exemplary Substitutions	Preferred
Ala (A)	Val· Leu· Ile	Val
Arg (R)	Lys· Gln· Asn	Lys
Asn (N)	Gln· His· Asp· Lys· Arg	Gln
Asp (D)	Glu· Asn	Glu
Cys (C)	Ser· Ala	Ser
Gln (Q)	Asn· Glu	Asn
Glu (E)	Asn· Gln	Asn
Gly (G)	Ala	Ala
His (H)	Asn· Gln· Lys· Arg	Arg
Ile (I)	Leu· Val· Met· Ala· Phe	Leu
Leu (L)	Norleucine· Ile· Val· Met	Ile
Lys (K)	Arg· Gln· Asn	Arg
Met (M)	Leu· Phe· Ile	Leu
Phe (F)	Trp· Leu· Val· Ile· Ala· Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val· Ser	Ser
Trp (W)	Tyr· Phe	Tyr
Tyr (Y)	Trp· Phe· Thr· Ser	Phe
Val (V)	Ile· Leu· Met· Phe· Ala	Leu

Amino acids may be grouped according to common side-chain properties: (1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: Asp, Glu; (4) basic: His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6)

aromatic: Trp, Tyr, Phe. In certain embodiments, non-conservative substitutions will entail exchanging a member of one of these classes for another class.

In certain embodiments, a type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e.g., a humanized or human antibody).

5 Generally, the resulting variant(s) selected for further study will have modifications (e.g., improvements) in certain biological properties (e.g., increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody, which may be conveniently generated, e.g., using phage display-based affinity maturation techniques such as
10 those described herein. Briefly, one or more HVR (or CDR) residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (e.g. binding affinity).

Alterations (e.g., substitutions) may be made in HVRs (or CDRs), e.g., to improve antibody affinity. Such alterations may be made in HVR (or CDRs) “hotspots,” i.e., residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (see, e.g.,
15 Chowdhury, *Methods Mol. Biol.* 207:179-196 (2008)), and/or SDRs (a-CDRs), with the resulting variant VH or VL being tested for binding affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O’Brien et al., ed., Human Press, Totowa, NJ, (2001)). In certain
20 embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (e.g., error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves HVR (or CDRs) -directed approaches, in which several HVR (or CDRs) residues (e.g., 4-6 residues at a time) are randomized. HVR (or CDRs) residues involved in antigen binding
25 may be specifically identified, e.g., using alanine scanning mutagenesis or modeling. CDR-H3 and CDR-L3 in particular are often targeted.

In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs (or CDRs) so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative substitutions as provided
30 herein) that do not substantially reduce binding affinity may be made in HVRs(or CDRs). Such alterations may be outside of HVR (or CDR) “hotspots” or CDRs. In certain embodiments of the variant VHH sequences provided above, each HVR (or CDR) either is unaltered, or contains no more than one, two or three amino acid substitutions.

A useful method for identification of residues or regions of an antibody that may be targeted
35 for mutagenesis is called “alanine scanning mutagenesis” as described by Cunningham and Wells (1989) *Science*, 244:1081-1085. In this method, a residue or group of target residues (e.g., charged

residues such as Arg, Asp, His, Lys, and Glu) are identified and replaced by a neutral or negatively charged amino acid (e.g., alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of the antibody to an enzyme (e.g., for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

2.9.2 Glycosylation variants

In certain embodiments, the antibody moiety is altered to increase or decrease the extent to which the construct is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

Where the antibody moiety comprises an Fc region (e.g., scFv-Fc), the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the C_H2 domain of the Fc region. See, e.g., Wright et al. TIBTECH 15:26-32 (1997). The oligosaccharide may include various carbohydrates, e.g., mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the "stem" of the biantennary oligosaccharide structure. In certain embodiments, modifications of the oligosaccharide in the antibody moiety may be made in order to create antibody variants with certain improved properties.

In certain embodiments, the antibody moiety has a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. For example, the amount of fucose in such antibody may be from 1% to 80%, from 1% to 65%, from 5% to 65% or from 20% to 40%. The amount of fucose is determined by calculating the average amount of fucose within the sugar chain at Asn297, relative to the sum of all glycostructures attached to Asn 297 (e.g., complex, hybrid and high mannose structures) as measured by MALDI-TOF mass spectrometry, as described in WO 2008/077546, for example. Asn297 refers to the asparagine residue located at about position 297 in the Fc region (EU numbering of Fc region residues); however, Asn297 may also be located

about ± 3 amino acids upstream or downstream of position 297, i.e., between positions 294 and 300, due to minor sequence variations in antibodies. Such fucosylation variants may have improved ADCC function. See, e.g., US Patent Publication Nos. US 2003/0157108 (Presta, L.); US 2004/0093621 (Kyowa Hakko Kogyo Co., Ltd). Examples of publications related to “defucosylated” or “fucose-deficient” antibody variants include: US 2003/0157108; WO 2000/61739; WO 2001/29246; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; WO 2003/085119; WO 2003/084570; WO 2005/035586; WO 2005/035778; WO2005/053742; WO2002/031140; Okazaki et al. J. Mol. Biol. 336:1239-1249 (2004); Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka et al. Arch. Biochem. Biophys. 249:533-545 (1986); US Patent Application No. US 2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams et al.), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (see, e.g., Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004); Kanda, Y. et al., Biotechnol. Bioeng., 94(4):680-688 (2006); and WO2003/085107).

In certain embodiments, the antibody moiety has bisected oligosaccharides, e.g., in which a biantennary oligosaccharide attached to the Fc region of the antibody is bisected by GlcNAc. Such antibody variants may have reduced fucosylation and/or improved ADCC function. Examples of such antibody variants are described, e.g., in WO 2003/011878 (Jean-Mairet et al.); US Patent No. 6,602,684 (Umana et al.); and US 2005/0123546 (Umana et al.). Antibody variants with at least one galactose residue in the oligosaccharide attached to the Fc region are also provided. Such antibody variants may have improved CDC function. Such antibody variants are described, e.g., in WO 1997/30087 (Patel et al.); WO 1998/58964 (Raju, S.); and WO 1999/22764 (Raju, S.).

2.9.3 Fc region variants

In certain embodiments, one or more amino acid modifications may be introduced into the Fc region of the antibody moiety (e.g., scFv-Fc), thereby generating an Fc region variant. The Fc region variant may comprise a human Fc region sequence (e.g., a human IgG1, IgG2, IgG3 or IgG4 Fc region) comprising an amino acid modification (e.g. a substitution) at one or more amino acid positions.

In certain embodiments, the Fc fragment possesses some but not all effector functions, which make it a desirable candidate for applications in which the half-life of the antibody moiety in vivo is important yet certain effector functions (such as complement and ADCC) are unnecessary or deleterious. In vitro and/or in vivo cytotoxicity assays can be conducted to confirm the reduction/depletion of CDC and/or ADCC activities. For example, Fc receptor (FcR) binding assays can be conducted to ensure that the antibody lacks FcγR binding (hence likely lacking ADCC

activity) but retains FcRn binding ability. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 2 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-492 (1991). Non-limiting examples of in vitro assays to assess ADCC activity of a molecule of interest is described in U.S. Patent No. 5,500,362 (see, e.g. Hellstrom, I. et al. *Proc. Nat'l Acad. Sci. USA* 83:7059-7063 (1986)) and Hellstrom, I et al., *Proc. Nat'l Acad. Sci. USA* 82:1499-1502 (1985); 5,821,337 (see Bruggemann, M. et al., *J. Exp. Med.* 166:1351-1361 (1987)). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA; and CytoTox 96® non-radioactive cytotoxicity assay (Promega, Madison, WI). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed in vivo, e.g., in an animal model such as that disclosed in Clynes et al. *Proc. Nat'l Acad. Sci. USA* 95:652-656 (1998). C1q binding assays may also be carried out to confirm that the antibody is unable to bind C1q and hence lacks CDC activity. See, e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., *J. Immunol. Methods* 202:163 (1996); Cragg, M.S. et al., *Blood* 101:1045-1052 (2003); and Cragg, M.S. and M.J. Glennie, *Blood* 103:2738-2743 (2004)). FcRn binding and in vivo clearance/half-life determinations can also be performed using methods known in the art (see, e.g., Petkova, S.B. et al., *Int'l. Immunol.* 18(12):1759-1769 (2006)).

Antibodies with reduced effector function include those with substitution of one or more of Fc region residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

Certain antibody variants with improved or diminished binding to FcRs are described. (See, e.g., U.S. Patent No. 6,737,056; WO 2004/056312, and Shields et al., *J. Biol. Chem.* 9(2): 6591-6604 (2001).)

In certain embodiments, the Fc fragment is an IgG1 Fc fragment. In certain embodiments, the IgG1 Fc fragment comprises a L234A mutation and/or a L235A mutation. In certain embodiments, the Fc fragment is an IgG2 or IgG4 Fc fragment. In certain embodiments, the Fc fragment is an IgG4 Fc fragment comprising a S228P, F234A, and/or a L235A mutation.

In certain embodiments, the antibody moiety comprises an Fc region with one or more amino acid substitutions which improve ADCC, e.g., substitutions at positions 298, 333, and/or 334 of the Fc region (EU numbering of residues).

In certain embodiments, alterations are made in the Fc region that result in altered (i.e., either improved or diminished) C1q binding and/or Complement Dependent Cytotoxicity (CDC), e.g., as described in US Patent No. 6,194,551, WO 99/51642, and Idusogie et al. J. Immunol. 164: 4178-4184 (2000).

5 In certain embodiments, the antibody moiety (e.g., scFv-Fc) variant comprising a variant Fc region comprising one or more amino acid substitutions which alters half-life and/or changes binding to the neonatal Fc receptor (FcRn). Antibodies with increased half-lives and improved binding to the neonatal Fc receptor (FcRn), which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., J. Immunol. 117:587 (1976) and Kim et al., J. Immunol. 24:249 (1994)), are
10 described in US2005/0014934A1 (Hinton et al.). Those antibodies comprise an Fc region with one or more substitutions therein which alters binding of the Fc region to FcRn. Such Fc variants include those with substitutions at one or more of Fc region residues, e.g., substitution of Fc region residue 434 (US Patent No. 7,371,826).

See also Duncan & Winter, Nature 322:738-40 (1988); U.S. Patent No. 5,648,260; U.S.
15 Patent No. 5,624,821; and WO 94/29351 concerning other examples of Fc region variants.

2.9.4 Cysteine engineered antibody variants

In certain embodiments, it may be desirable to create cysteine engineered antibody moieties, e.g., “thioMAbs,” in which one or more residues of an antibody are substituted with cysteine
20 residues. In certain embodiments, the substituted residues occur at accessible sites of the antibody. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate, as described further herein. In certain embodiments, any one or more of the following residues may be substituted with cysteine:
25 A118 (EU numbering) of the heavy chain; and S400 (EU numbering) of the heavy chain Fc region. Cysteine engineered antibody moieties may be generated as described, e.g., in U.S. Patent No. 7,521,541.

2.10 Antibody Derivatives

30 In certain embodiments, the antibody moiety described herein may be further modified to comprise additional nonproteinaceous moieties that are known in the art and readily available. The moieties suitable for derivatization of the antibody include but are not limited to water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol,
35 carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1, 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers

or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer are attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in diagnosis under defined conditions, etc.

In certain embodiments, the antibody moiety may be further modified to comprise one or more biologically active protein, polypeptides or fragments thereof. "Bioactive" or "biologically active", as used herein interchangeably, means showing biological activity in the body to carry out a specific function. For example, it may mean the combination with a particular biomolecule such as protein, DNA, etc., and then promotion or inhibition of the activity of such biomolecule. In certain embodiments, the bioactive protein or fragments thereof include proteins and polypeptides that are administered to patients as the active drug substance for prevention of or treatment of a disease or condition, as well as proteins and polypeptides that are used for diagnostic purposes, such as enzymes used in diagnostic tests or in vitro assays, as well as proteins and polypeptides that are administered to a patient to prevent a disease such as a vaccine.

2.11 Methods of Antibody Production

The antibodies disclosed herein can be produced using any available or known technique in the art. For example, but not by way of limitation, antibodies can be produced using recombinant methods and compositions, e.g., as described in U.S. Patent No. 4,816,567. Detailed procedures to generate antibodies are described in the Examples below.

The presently disclosed subject matter further provides an isolated nucleic acid encoding an antibody disclosed herein. For example, the isolated nucleic acid can encode an amino acid sequence comprising the VL and/or an amino acid sequence comprising the VH of the antibody, e.g., the light and/or heavy chains of the antibody.

In certain embodiments, the nucleic acid can be present in one or more vectors, e.g., expression vectors. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid," which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, where additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into

which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors, expression vectors, are capable of directing the expression of genes to which they are operably linked. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids (vectors). However, the disclosed subject matter is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses) that serve equivalent functions.

Different parts of the antibodies disclosed herein can be constructed in a single, multicistronic expression cassette, in multiple expression cassettes of a single vector, or in multiple vectors. Examples of elements that create polycistronic expression cassette include, but are not limited to, various viral and non-viral Internal Ribosome Entry Sites (IRES, e.g., FGF-1 IRES, FGF-2 IRES, VEGF IRES, IGF-II IRES, NF- κ B IRES, RUNX1 IRES, p53 IRES, hepatitis A IRES, hepatitis C IRES, pestivirus IRES, aphthovirus IRES, picornavirus IRES, poliovirus IRES and encephalomyocarditis virus IRES) and cleavable linkers (e.g., 2A peptides, e.g., P2A, T2A, E2A and F2A peptides). Combinations of retroviral vector and an appropriate packaging line are also suitable, where the capsid proteins will be functional for infecting human cells. Various amphotropic virus-producing cell lines are known, including, but not limited to, PA12 (Miller, et al. (1985) Mol. Cell. Biol. 5:431-437); PA317 (Miller, et al. (1986) Mol. Cell. Biol. 6:2895-2902); and CRIP (Danos, et al. (1988) Proc. Natl. Acad. Sci. USA 85:6460-6464). Non-amphotropic particles are suitable too, e.g., particles pseudotyped with VSVG, RD114 or GALV envelope and any other known in the art.

In certain embodiments, the nucleic acid encoding an antibody of the present disclosure and/or the one or more vectors including the nucleic acid can be introduced into a host cell. In certain embodiments, the introduction of a nucleic acid into a cell can be carried out by any method known in the art including, but not limited to, transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. In certain embodiments, a host cell can include, e.g., has been transformed with: a vector comprising a nucleic acid that encodes an amino acid sequence comprising a single domain antibody and/or the VH of a single domain antibody. In certain embodiments, a host cell can include, e.g., has been transformed with: (1) a vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and an amino acid sequence comprising the VH of the antibody, or (2) a first vector comprising a nucleic acid that encodes an amino acid sequence comprising the VL of the antibody and a second vector comprising a nucleic acid that encodes an amino acid sequence

comprising the VH of the antibody. In certain embodiments, the host cell is eukaryotic, e.g., a Chinese Hamster Ovary (CHO) cell or lymphoid cell (e.g., YO, NSO, Sp20 cell).

In certain embodiments, the methods of making an antibody disclosed herein can include culturing a host cell, in which a nucleic acid encoding the antibody has been introduced, under conditions suitable for expression of the antibody, and optionally recovering the antibody from the host cell and/or host cell culture medium. In certain embodiments, the antibody is recovered from the host cell through chromatography techniques.

For recombinant production of an antibody of the present disclosure, a nucleic acid encoding an antibody, e.g., as described above, can be isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such nucleic acid may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the antibody). Suitable host cells for cloning or expression of antibody-encoding vectors include prokaryotic or eukaryotic cells described herein. For example, antibodies can be produced in bacteria, in particular when glycosylation and Fc effector function are not needed. For expression of antibody fragments and polypeptides in bacteria, see, e.g., U.S. Patent Nos. 5,648,237, 5,789,199, and 5,840,523. (See also Charlton, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ, 2003), pp. 245-254, describing expression of antibody fragments in *E. coli*.) After expression, the antibody may be isolated from the bacterial cell paste in a soluble fraction and can be further purified.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for antibody-encoding vectors, including fungi and yeast strains whose glycosylation pathways have been "humanized," resulting in the production of an antibody with a partially or fully human glycosylation pattern. See Gemgross, *Nat. Biotech.* 22:1409-1414 (2004), and Li et al., *Nat. Biotech.* 24:210-215 (2006). Suitable host cells for the expression of glycosylated antibody can also be derived from multicellular organisms (invertebrates and vertebrates). Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. In certain embodiments, plant cell cultures can be utilized as host cells. See, e.g., US Patent Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIES™ technology for producing antibodies in transgenic plants).

In certain embodiments, vertebrate cells can also be used as hosts. For example, and not by way of limitation, mammalian cell lines that are adapted to grow in suspension can be useful. Non-limiting examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SY40 (COS-7); human embryonic kidney line (293 or 293 cells as described, e.g., in Graham et al., *J Gen Viral.* 36:59 (1977)); baby hamster kidney cells (BHK); mouse sertoli cells (TM4 cells as

described, e.g., in Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV 1); African green monkey kidney cells (VERO-76); human cervical carcinoma cells (HELA); canine kidney cells (MDCK); buffalo rat liver cells (BRL 3A); human lung cells (W138); human liver cells (Hep 02); mouse mammary tumor (MMT 060562); TRI cells, as described, e.g., in Mather et al., *Annals N. Y. Acad. Sci.* 383:44-68 (1982); MRC 5 cells; and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFK CHO cells (Urlaub et al., *Proc. Natl. Acad. Sci. USA* 77:4216 (1980)); and myeloma cell lines such as YO, NSO and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003).

In certain embodiments, techniques for making bispecific and/or multispecific antibodies include, but are not limited to, recombinant expression of two immunoglobulin heavy chain-light chain pairs having the same specificity, where one or two of the heavy chains or the light chains are fused to an antigen binding moiety (e.g., a single domain antibody, e.g., a VHH) having a different specificity, recombinant coexpression of two immunoglobulin heavy chain-light chain pairs having different specificities (see Milstein and Cuello, *Nature* 305: 537 (1983)), PCT Patent Application No. WO 93/08829, and Traunecker et al., *EMBO J* 10: 3655 (1991)), and “knob-in-hole” engineering (see, e.g., U.S. Patent No. 5,731,168). Bispecific antibodies can also be made by engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules (WO 2009/089004A1); cross-linking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bispecific antibodies (see, e.g., Kostelny et al., *J Immunol.*, 148(5): 1547-1553 (1992)); using “diabody” technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g., Gruber et al., *J Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as described, e.g., in Tutt et al. *J Immunol.* 147: 60 (1991).

Bispecific and multispecific molecules of the present disclosure can also be made using chemical techniques (see, e.g., Kranz (1981) *Proc. Natl. Acad. Sci. USA* 78:5807), “polydoma” techniques (see, e.g., U.S. Patent 4,474,893), or recombinant DNA techniques. Bispecific and multispecific molecules of the presently disclosed subject matter can also be prepared by conjugating the constituent binding specificities, e.g., a first epitope and a second epitope binding specificities, using methods known in the art and as described herein. For example, and not by way of limitation, each binding specificity of the bispecific and multispecific molecule can be generated together by recombinant fusion protein techniques, or can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Non-limiting examples of

cross-linking agents include protein A, carbodiimide, N- succinimidyl-S-acetyl-thioacetate (SATA), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC) (see, e.g., Karpovsky (1984) J. Exp. Med. 160:1686; Liu (1985) Proc. Natl. Acad. Sci. USA 82:8648). Other methods include those
5 described by Paulus (Behring Ins. Mitt. (1985) No. 78, 1 18-132; Brennan (1985) Science 229:81-83), Glennie (1987) J Immunol. 139: 2367-2375). When the binding specificities are antibodies (e.g., two humanized antibodies), they can be conjugated via sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In certain embodiments, the hinge region can be modified to contain an odd number of sulfhydryl residues, e.g., one, prior to conjugation.

10 In certain embodiments, both binding specificities of a bispecific antibody can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific and multispecific molecule is a MAb x MAb, MAb x Fab, Fab x F(ab')₂ or ligand x Fab fusion protein. In certain embodiments, a bispecific antibody of the present disclosure can be a single chain molecule, such as a single chain bispecific antibody, a single chain
15 bispecific molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific and multispecific molecules can also be single chain molecules or can comprise at least two single chain molecules. Methods for preparing bi- and multispecific molecules are described, for example, in U.S. Patent No. 5,260,203; U.S. Patent No. 5,455,030; U.S. Patent No. 4,881 ,175; U.S. Patent No. 5,132,405;
20 U.S. Patent No. 5,091 ,513; U.S. Patent No. 5,476,786; U.S. Patent No. 5,013,653; U.S. Patent No. 5,258,498; and U.S. Patent No. 5,482,858. Engineered antibodies with three or more functional antigen binding sites (e.g., epitope binding sites) including "Octopus antibodies," are also included herein (see, e.g., US 2006/0025576A1).

25 In certain embodiments, an animal system can be used to produce an antibody of the present disclosure. One animal system for preparing hybridomas is the murine system.

Hybridoma production in the mouse is a very well-established procedure. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known (see, e.g., Harlow and Lane (1988), Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory Press,
30 Cold Spring Harbor New York).

2.12 Assays

The antibodies of the present disclosure provided herein can be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays
35 known in the art and provided herein.

In certain embodiments, an antibody of the present disclosure can be tested for its antigen binding activity by known methods, such as enzyme-linked immunosorbent assay (ELISA), a radioimmunoassay (RIA), or a Western Blot Assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g.,
5 an antibody) specific for the complex of interest. For example, the antibody can be detected using, e.g., an enzyme-linked antibody or antibody fragment which recognizes and specifically binds to the antibody. Alternatively, the antibody can be detected using any of a variety of other immunoassays. For example, the antibody can be radioactively labeled and used in a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays,
10 Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March 1986, which is incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a Geiger counter or a scintillation counter or by autoradiography.

In certain embodiments, competition assays can be used to identify an antibody that competes with an antibody of the present disclosure, e.g., 1C12, 2A3 or 1G1, for binding to TIGIT.
15 In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by 1C12, 2A3 or 1G1. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in Methods in Molecular Biology vol. 66 (Humana Press, Totowa, NJ).

In a non-limiting example of a competition assay, immobilized TIGIT can be incubated in a
20 solution comprising a first labeled antibody that binds to TIGIT (e.g., 1C12, 2A3 or 1G1) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to TIGIT. The second antibody may be present in a hybridoma supernatant. As a control, immobilized TIGIT is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first
25 antibody to TIGIT, excess unbound antibody is removed, and the amount of label associated with immobilized TIGIT is measured. If the amount of label associated with immobilized TIGIT is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to TIGIT. See Harlow and Lane (1988) Antibodies: A Laboratory Manual ch.14 (Cold Spring Harbor Laboratory, Cold Spring
30 Harbor, NY).

The present disclosure provides assays for identifying anti-TIGIT antibodies thereof having biological activity. Biological activity may include, e.g., activating an immune cell or an immune activation reporter, e.g., a NFAT reporter. Antibodies having such biological activity in vivo and/or in vitro are also provided.

The presently disclosed subject matter further provides immunoconjugates comprising an antibody, disclosed herein, conjugated to one or more detection probe and/or cytotoxic agents, such as chemotherapeutic agents or drugs, growth inhibitory agents, toxins (e.g., protein toxins, enzymatically active toxins of bacterial, fungal, plant, or animal origin, or fragments thereof), or radioactive isotopes. For example, an antibody or antigen-binding portion of the disclosed subject matter can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other binding molecules, such as another antibody, antibody fragment, peptide or binding mimetic.

In certain embodiments, an immunoconjugate is an antibodydrug conjugate (ADC) in which an antibody is conjugated to one or more drugs, including but not limited to a maytansinoid (see U.S. Patent Nos. 5,208,020, 5,416,064 and European Patent EP 0 425 235); an auristatin such as monomethylauristatin drug moieties DE and DF (MMAE and MMAF) (see U.S. Patent Nos. 5,635,483 and 5,780,588, and 7,498,298); a dolastatin; a calicheamicin or derivative thereof (see U.S. Patent Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, and 5,877,296; Hinman et al., *Cancer Res.* 53:3336-3342 (1993); and Lode et al., *Cancer Res.* 58:2925-2928 (1998)); an anthracycline such as daunomycin or doxorubicin (see Kratz et al., *Current Med Chem.* 13:477-523 (2006); Jeffrey et al., *Bioorganic & Med. Chem. Letters* 16:358-362 (2006); Torgov et al., *Bioconj. Chem.* 16:717-721 (2005); Nagy et al., *Proc. Natl. Acad. Sci. USA* 97:829-834 (2000); Dubowchik et al., *Bioorg. & Med. Chem. Letters* 12:1529-1532 (2002); King et al., *J Med. Chem.* 45:4336-4343 (2002); and U.S. Patent No. 6,630,579); methotrexate; vindesine; ataxane such as docetaxel, paclitaxel, larotaxel, tesetaxel, and ortataxel; a trichothecene; and CC1065.

In certain embodiments, an immunoconjugate comprises an antibody as described herein conjugated to an enzymatically active toxin or fragment thereof, including but not limited to diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes.

In certain embodiments, an immunoconjugate comprises an antibody as described herein conjugated to a radioactive atom to form a radioconjugate. A variety of radioactive isotopes are available for the production of radioconjugates. Non-limiting examples include At^{211} , I^{131} , I^{125} , Y^{90} , Re^{186} , Re^{188} , Sm^{153} , Bi^{212} , P^{32} , Pb^{212} and radioactive isotopes of Lu. When the radioconjugate is used for detection, it can include a radioactive atom for scintigraphic studies, for example tc99m or ^{111}In , or a spin label for nuclear magnetic resonance (NMR) imaging (also

known as magnetic resonance imaging, MRI), such as iodine-123, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

Conjugates of an antibody and cytotoxic agent can be made using a variety of bi functional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithio) propionate (SPDP),
5 succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis-(p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine
10 compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science 238: 1098 (1987). Carbon-4-labeled l-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026. The linker can be a "cleavable linker" facilitating release of a cytotoxic drug in the cell. For example, an acid-
15 labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (Chari et al., Cancer Res. 52:127-131 (1992); U.S. Patent No. 5,208,020) can be used.

The immunconjugates or ADCs herein expressly contemplate, but are not limited to, such conjugates prepared with cross-linker reagents including, but not limited to, BMPS, EMCS, GMBS, HBVS, LC-SMCC, MBS, MPBH, SBAP, SIA, SIAB, SMCC, SMPB, SMPH, sulfo-EMCS, sulfo-
20 GMBS, sulfo-KMUS, sulfo-MBS, sulfo-SIAB, sulfo-SMCC, and sulfo-SMPB, and SVSB (succinimidyl-(4-vinylsulfone)benzoate) which are commercially available (e.g., from Pierce Biotechnology, Inc., Rockford, IL., U.S.A).

3. METHODS OF USE

25 The presently disclosed subject matter further provides methods for using the disclosed antibodies, e.g., an anti-TIGIT antibody and an anti-TIGIT/anti-PD1 multispecific antibody. In certain embodiments, the methods are directed to therapeutic uses of the presently disclosed antibodies. In certain embodiments, the methods are directed to diagnostic use of the presently disclosed antibodies.

30

3.1 Treatment methods

The present disclosure provides methods and use of any antibodies disclosed herein (e.g., the anti-TIGIT antibody and/or the anti-TIGIT/anti-PD1 multispecific antibody) for treatment of diseases and disorders or for increasing an immune response. In certain embodiments, the
35 antibody and/or pharmaceutical compositions comprising the same disclosed herein can be administered to subjects (e.g., mammals such as humans) to treat diseases and disorders or to

increases an immune response. In certain embodiments, the diseases and disorders involve immune checkpoint inhibitions and/or abnormal TIGIT and/or PD1 activity. In certain embodiments, the diseases and disorders that can be treated by the antibodies disclosed herein include, but are not limited to, neoplasia, e.g., cancer.

5 In certain embodiments, the present disclosure provides anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies described herein (or fragments thereof) for use in the manufacture of a medicament. In certain embodiments, the present disclosure provides anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies described herein (or fragments thereof) for use in the manufacture of a medicament for treating of cancer. In certain embodiments, the present disclosure provides anti-TIGIT and/or
10 anti-TIGIT/anti-PD1 antibodies described herein (or fragments thereof) for use in treating cancer in a subject. In certain embodiments, the present disclosure provides pharmaceutical compositions comprising an anti-TIGIT and/or anti-TIGIT/anti-PD1 antibody provided herein (or fragments thereof) for use in treating cancer in a subject. In certain embodiments, the cancer can be blood cancers (e.g. leukemias, lymphomas, and myelomas), ovarian cancer, breast cancer, bladder cancer,
15 brain cancer, colon cancer, intestinal cancer, liver cancer, lung cancer, pancreatic cancer, prostate cancer, skin cancer, stomach cancer, glioblastoma, throat cancer, melanoma, neuroblastoma, adenocarcinoma, glioma, soft tissue sarcoma, and various carcinomas (including prostate and small cell lung cancer). Suitable carcinomas further include any known carcinoma in the field of oncology, including, but not limited to, astrocytoma, fibrosarcoma, myxosarcoma, liposarcoma,
20 oligodendroglioma, ependymoma, medulloblastoma, primitive neural ectodermal tumor (PNET), chondrosarcoma, osteogenic sarcoma, pancreatic ductal adenocarcinoma, small and large cell lung adenocarcinomas, chordoma, angiosarcoma, endotheliosarcoma, squamous cell carcinoma, bronchoalveolar carcinoma, epithelial adenocarcinoma, and liver metastases thereof, lymphangiosarcoma, lymphangioendotheliosarcoma, hepatoma, cholangiocarcinoma, synovium,
25 mesothelioma, Ewing's tumor, rhabdomyosarcoma, colon carcinoma, basal cell carcinoma, sweat gland carcinoma, papillary carcinoma, sebaceous gland carcinoma, papillary adenocarcinoma, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, testicular tumor, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic
30 neuroma, oligodendroglioma, meningioma, neuroblastoma, retinoblastoma, leukemia, multiple myeloma, Waldenstrom's macroglobulinemia, breast tumors such as ductal and lobular adenocarcinoma, squamous and adenocarcinomas of the uterine cervix, uterine and ovarian epithelial carcinomas, prostatic adenocarcinomas, transitional squamous cell carcinoma of the bladder, B and T cell lymphomas (nodular and diffuse) plasmacytoma, acute and chronic leukemias,
35 malignant melanoma, soft tissue sarcomas and leiomyosarcomas.

In certain embodiments, the cancer can be melanoma, NSCLC, head and neck cancer, urothelial cancer, breast cancer (e.g., triple-negative breast cancer, TNBC), gastric cancer, cholangiocarcinoma, classical Hodgkin's lymphoma (cHL), Non-Hodgkin lymphoma primary mediastinal B-Cell lymphoma (NHL PMBCL), mesothelioma, ovarian cancer, lung cancer (e.g.,
5 small-cell lung cancer), esophageal cancer, nasopharyngeal carcinoma (NPC), biliary tract cancer, colorectal cancer, cervical cancer or thyroid cancer.

In certain embodiments, the subject to be treated is a mammal (e.g., human, non-human primate, rat, mouse, cow, horse, pig, sheep, goat, dog, cat, etc.). In certain embodiments, the subject is a human. In certain embodiments, the subject is suspected of having or at risk of having a cancer
10 or be diagnosed with a cancer or any other disease having abnormal TIGIT and/or PD1 expression or activity.

Many diagnostic methods for cancer or any other disease exhibiting abnormal TIGIT and/or PD1 activity and the clinical delineation of those diseases are known in the art. Such methods include, but are not limited to, e.g., immunohistochemistry, PCR, fluorescent in situ hybridization
15 (FISH). Additional details regarding diagnostic methods for abnormal TIGIT and/or PD1 activity or expression are described in, e.g., Gupta et al. (2009) *Mod Pathol.* 22(1): 128-133; Lopez-Rios et al. (2013) *J Clin Pathol.* 66(5): 381-385; Ellison et al. (2013) *J Clin Pathol* 66(2): 79-89; and Guha et al. (2013) *PLoS ONE* 8(6): e67782.

Administration can be by any suitable route including, e.g., intravenous, intramuscular, or
20 subcutaneous. In some embodiments, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof) and/or compositions provided herein are administered in combination with a second, third, or fourth agent (including, e.g., an antineoplastic agent, a growth inhibitory agent, a cytotoxic agent, or a chemotherapeutic agent) to treat the diseases or disorders involving abnormal TIGIT/PD1 activity. Such agents include, e.g., docetaxel, gefitinib, FOLFIRI (irinotecan, 5-
25 fluorouracil, and leucovorin), irinotecan, cisplatin, carboplatin, paclitaxel, bevacizumab (anti-VEGF antibody), FOLFOX-4, infusional fluorouracil, leucovorin, and oxaliplatin, afatinib, gemcitabine, capecitabine, pemetrexed, tivantinib, everolimus, CpG-ODN, rapamycin, lenalidomide, vemurafenib, endostatin, lapatinib, PX-866, Imprime PGG, and irlotinibm. In some embodiments, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof) are conjugated to the
30 additional agent.

In certain embodiments, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof) and/or compositions provided herein are administered in combination with one or more additional therapies, such as radiation therapy, surgery, chemotherapy, and/or targeted therapy. In certain embodiments, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof)
35 and/or compositions provided herein are administered in combination with radiation therapy. In certain embodiments, the combination of an anti-TIGIT and/or anti-TIGIT/anti-PD1 antibody (or

fragment thereof) and/or composition provided herein and radiation therapy is used for treating a neoplasm or cancer disclosed herein.

Depending on the indication to be treated and factors relevant to the dosing that a physician of skill in the field would be familiar with, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies or fragments thereof, provided herein will be administered at a dosage that is efficacious for the treatment of that indication while minimizing toxicity and side effects. For the treatment of a cancer, a typical dose can be, for example, in the range of 0.001 to 1000 μg ; however, doses below or above this exemplary range are within the scope of the invention. The daily dose can be about 0.1 μg /kg to about 100 mg/kg of total body weight, about 0.1 μg /kg to about 100 μg /kg of total body weight or about 1 μg /kg to about 100 μg /kg of total body weight. As noted above, therapeutic or prophylactic efficacy can be monitored by periodic assessment of treated patients. For repeated administrations over several days or longer, depending on the condition, the treatment is repeated until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful and are within the scope of the invention. The desired dosage can be delivered by a single bolus administration of the composition, by multiple bolus administrations of the composition, or by continuous infusion administration of the composition.

A pharmaceutical composition comprising the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibody or a fragment thereof can be administered one, two, three, or four times daily. The compositions can also be administered less frequently than daily, for example, six times a week, five times a week, four times a week, three times a week, twice a week, once a week, once every two weeks, once every three weeks, once a month, once every two months, once every three months, or once every six months. The compositions may also be administered in a sustained release formulation, such as in an implant which gradually releases the composition for use over a period of time, and which allows for the composition to be administered less frequently, such as once a month, once every 2-6 months, once every year, or even a single administration. The sustained release devices (such as pellets, nanoparticles, microparticles, nanospheres, microspheres, and the like) may be administered by injection.

The antibody (or a fragment thereof) may be administered in a single daily dose, or the total daily dose may be administered in divided dosages of two, three, or four times daily. The compositions can also be administered less frequently than daily, for example, six times a week, five times a week, four times a week, three times a week, twice a week, once a week, once every two weeks, once every three weeks, once a month, once every two months, once every three months, or once every six months. The antibody (or a fragment thereof) may also be administered in a sustained release formulation, such as in an implant which gradually releases the composition for use over a period of time, and which allows for the composition to be administered less frequently, such as once a month, once every 2-6 months, once every year, or even a single administration. The

sustained release devices (such as pellets, nanoparticles, microparticles, nanospheres, microspheres, and the like) may be administered by injection or surgically implanted in various locations.

Cancer treatments can be evaluated by, e.g., but not limited to, tumor regression, tumor weight or size shrinkage, time to progression, duration of survival, progression free survival, overall response rate, duration of response, quality of life, protein expression and/or activity. Approaches to determining efficacy of the therapy can be employed, including for example, measurement of response through radiological imaging.

In certain embodiments, the efficacy of treatment is measured by the percentage tumor growth inhibition (% TGI), calculated using the equation $100 - (T/C \times 100)$, where T is the mean relative tumor volume of the treated tumor, and C is the mean relative tumor volume of a non-treated tumor. In certain embodiments, the %TGI is about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, or more than 95%.

3.2 Methods of diagnosis and imaging

Labeled anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies, fragments thereof, and derivatives and analogs thereof can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or disorders associated with the expression, aberrant expression and/or activity of TIGIT and/or PD1. For example, the anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof) provided herein can be used in in situ, in vivo, ex vivo, and in vitro diagnostic assays or imaging assays. Methods for detecting expression of a TIGIT polypeptide, comprising (a) assaying the expression of the polypeptide in cells (e.g., tissue) or body fluid of an individual using one or more antibodies of this invention and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed gene expression level compared to the standard expression level is indicative of aberrant expression.

Additional embodiments provided herein include methods of diagnosing a disease or disorder associated with expression or aberrant expression of TIGIT and/or PD1 in an animal (e.g., a mammal such as a human). The methods comprise detecting TIGIT and/or PD1 molecules in the mammal. In certain embodiments, diagnosis comprises: (a) administering an effective amount of a labeled anti-TIGIT and/or anti-TIGIT/anti-PD1 antibody (or fragment thereof) to a mammal (b) waiting for a time interval following the administering for permitting the labeled anti-TIGIT and/or anti-TIGIT/anti-PD1 antibody (or fragment thereof) to preferentially concentrate at sites in the subject where the TIGIT and/or PD1 molecule is expressed (and for unbound labeled molecule to be cleared to background level); (c) determining background level; and (d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with expression or aberrant

expression of TIGIT/PD1. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

Anti-TIGIT and/or anti-TIGIT/anti-PD1 antibodies (or fragments thereof) provided herein can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium ($^{115\text{m}}\text{In}$, $^{113\text{m}}\text{In}$, ^{112}In , ^{111}In), and technetium (^{99}Tc , $^{99\text{m}}\text{Tc}$), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru ; luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

Techniques known in the art may be applied to labeled antibodies (or fragments thereof) provided herein. Such techniques include, but are not limited to, the use of bifunctional conjugating agents (see e.g., U.S. Pat. Nos. 5,756,065; 5,714,631; 5,696,239; 5,652,361; 5,505,931; 5,489,425; 5,435,990; 5,428,139; 5,342,604; 5,274,119; 4,994,560; and 5,808,003).

Alternatively, or additionally, one can measure levels of a TIGIT and/or PD1 polypeptide-encoding nucleic acid or mRNA in the cell, e.g., via fluorescent in situ hybridization using a nucleic acid based probe corresponding to an EGFR-encoding nucleic acid or the complement thereof; (FISH; see WO98/45479 published October, 1998), Southern blotting, Northern blotting, or polymerase chain reaction (PCR) techniques, such as real time quantitative PCR (RT-PCR). One can also study TIGIT and/or PD1 overexpression by measuring shed antigen in a biological fluid such as serum, e.g., using antibody-based assays (see also, e.g., U.S. Patent No. 4,933,294 issued June 12, 1990; WO91/05264 published April 18, 1991; U.S. Patent 5,401,638 issued March 28, 1995; and Sias et al., J. Immunol. Methods 132:73-80 (1990)). Aside from the above assays, various in vivo and ex vivo assays are available to the skilled practitioner. For example, one can expose cells within the body of the mammal to an antibody which is optionally labeled with a detectable label, e.g., a radioactive isotope, and binding of the antibody to the body cells can be evaluated, e.g., by external scanning for radioactivity or by analyzing a sample (e.g., a biopsy or other biological sample) taken from a mammal previously exposed to the antibody.

4. PHARMACEUTICAL FORMULATIONS

The presently disclosed subject matter further provides pharmaceutical formulations containing one or more antibodies disclosed herein, with a pharmaceutically acceptable carrier. In certain embodiments, the pharmaceutical compositions can include a combination of multiple (e.g., two or more) antibodies and/or antigen-binding portions thereof of the presently disclosed subject matter. In certain embodiments, a pharmaceutical composition of the present disclosure can include one or more anti-TIGIT antibodies or one or more of anti-TIGIT/anti-PD1 multispecific antibodies.

In certain embodiments, the disclosed pharmaceutical formulations can be prepared by combining an antibody having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. For example, but not by way of limitation, lyophilized antibody formulations are described in US Patent No. 6,267,958. In certain embodiments, aqueous antibody formulations can include those described in US Patent No. 6,171,586 and WO2006/044908, the latter formulations including a histidine-acetate buffer. In certain embodiments, the antibody can be of a purity greater than about 80%, greater than about 90%, greater than about 91%, greater than about 92%, greater than about 93%, greater than about 94%, greater than about 95%, greater than about 96%, greater than about 97%, greater than about 98%, greater than about 99%, greater than about 99.1%, greater than about 99.2%, greater than about 99.3%, greater than about 99.4%, greater than about 99.5%, greater than about 99.6%, greater than about 99.7%, greater than about 99.8% or greater than about 99.9%.

Pharmaceutically acceptable carriers are generally nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids, antioxidants including ascorbic acid and methionine, preservatives (such as octadecyldimethylbenzyl ammonium chloride, hexamethonium chloride, benzalkonium chloride, benzethonium chloride, phenol, butyl or benzyl alcohol, alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, 3-pentanol, and m-cresol), low molecular weight (less than about 10 residues) polypeptides, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine, monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans, chelating agents such as EDTA, sugars such as sucrose, mannitol, trehalose or sorbitol, salt-forming counter-ions such as sodium, metal complexes (e.g., Zn-protein complexes), and/or non-ionic surfactants such as polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rHuPH20 (HYLENEX®, Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rHuPH20, are described in US Patent

Publication Nos. 2005/0260186 and 2006/0104968. In certain embodiments, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

The carrier can be suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound, i.e., an anti-TIGIT/anti-PD1 antibody or an anti-TIGIT antibody, can be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

Pharmaceutical compositions of the present disclosure also can be administered in combination therapy, i.e., combined with other agents. In certain embodiments, pharmaceutical compositions disclosed herein can also contain more than one active ingredients as necessary for the particular indication being treated, for example, those with complementary activities that do not adversely affect each other. In certain embodiments, the pharmaceutical formulation can include a second active ingredient for treating the same disease treated by the first therapeutic. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended. For example, and not by way of limitation, the formulation of the present disclosure can also contain more than one active ingredients as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, it may be desirable to further provide a second therapeutic useful for treatment of the same disease. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended.

A composition of the present disclosure can be administered by a variety of methods known in the art. The route and/or mode of administration vary depending upon the desired results. The active compounds can be prepared with carriers that protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are described by e.g., Sustained and Controlled Release Drug Delivery Systems, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978. In certain embodiments, the pharmaceutical compositions are manufactured under Good Manufacturing Practice (GMP) conditions of the U.S. Food and Drug Administration.

Sustained-release preparations containing a disclosed antibody can also be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g. films, or microcapsules. In certain embodiments, active ingredients can be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate)

microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

To administer an antibody of the present disclosure by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan et al. (1984) J Neuroimmunol. 7:27).

Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the present disclosure is contemplated. Supplementary active compounds can also be incorporated into the compositions.

Therapeutic compositions typically must be sterile, substantially isotonic, and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

Sterile injectable solutions can be prepared by incorporating one or more disclosed antibodies in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration, e.g., by filtration through sterile filtration membranes. Generally, dispersions are prepared by

incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Therapeutic compositions can also be administered with medical devices known in the art.

For example, a therapeutic composition of the present disclosure can be administered with a needleless hypodermic injection device, such as the devices disclosed in, e.g., U.S. Patent Nos. 5,399,163, 5,383,851, 5,312,335, 5,064,413, 4,941,880, 4,790,824 or 4,596,556. Examples of implants and modules useful in the present disclosure include: U.S. Patent No. 4,487,603, which
5 discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Patent No. 4,486,194, which discloses a therapeutic device for administering medicants through the skin; U.S. Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Patent No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Patent No. 4,439,196,
10 which discloses an osmotic drug delivery system having multi-chamber compartments; and U.S. Patent No. 4,475,196, which discloses an osmotic drug delivery system. Many other such implants, delivery systems, and modules are known.

For the therapeutic compositions, formulations of the present disclosure include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral
15 administration. The formulations can conveniently be presented in unit dosage form and may be prepared by any methods known in the art of pharmacy. The amount of antibody, which can be combined with a carrier material to produce a single dosage form, vary depending upon the subject being treated, and the particular mode of administration. The amount of the antibody which can be combined with a carrier material to produce a single dosage form generally be that amount of the
20 composition which produces a therapeutic effect. Generally, out of one hundred percent, this amount range from about 0.01 percent to about ninety-nine percent of active ingredient, from about 0.1 percent to about 70 percent, or from about 1 percent to about 30 per cent.

Dosage forms for the topical or transdermal administration of compositions of the present disclosure include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and
25 inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

The phrases "parenteral administration" and "administered parenterally" mean modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital,
30 intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

These pharmaceutical compositions can also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of
35 various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride,

and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In certain embodiments, when the antibodies of the present disclosure are administered as pharmaceuticals, to humans and animals, they can be given alone or as a pharmaceutical composition containing, for example, from about 0.01% to about 99.5% (or about 0.1% to about 90%) of an antibody, described herein, in combination with a pharmaceutically acceptable carrier.

5. ARTICLES OF MANUFACTURE

The presently disclosed subject matter further provides articles of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above.

In certain embodiments, the article of manufacture includes a container and a label or package insert on or associated with the container. Non limiting examples of suitable containers include bottles, vials, syringes, IV solution bags, etc. The containers can be formed from a variety of materials such as glass or plastic. The container can hold a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle).

In certain embodiments, at least one active agent in the composition is an antibody of the presently disclosed subject matter. The label or package insert can indicate that the composition is used for treating the condition of choice.

In certain embodiments, the article of manufacture can comprise (a) a first container with a composition contained therein, wherein the composition comprises an antibody of the present disclosure; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. In certain embodiments, the article of manufacture can further comprise a package insert indicating that the compositions can be used to treat a particular condition.

Alternatively, or additionally, the article of manufacture can further an additional container, e.g., a second or third container, including a pharmaceutically acceptable buffer, such as, but not limited to, bacteriostatic water for injection (BWFI), phosphate-buffered saline, Ringer's solution and dextrose solution. The article of manufacture can include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

SEQUENCE TABLE

SEQ ID	Gene Name	Sequence
--------	-----------	----------

NO		
1.	4A11 CDR1	GRPFSNYT
2.	4A11 CDR2	AWPSPST
3.	4A11 CDR3	AADYKSLTQSWLNAALDY
4.	4A11 VHH	QVKLEESGGGEAQPGGSLRLSCTASGRPFSNYTMGWFRR APGKEREFVGLAWPSPSTYVVDVSKGRFTISRDNANTYI LQMNSLKPEDTAIYYCAADYKSLTQSWLNAALDYWGQG TQVTVSS
5.	4B5 CDR1	PRTFSTFH
6.	4B5 CDR2	FNWSSGRT
7.	4B5 CDR3	AAARDRGLHDGTTSDSYLEGSHEY
8.	4B5 VHH	QVKLEESGGRSVLAGGSLRLRCAGTPRTFSTFHIGWFRQA PGKEREFVAAFNWSSGRTYYADSVKGRFTISRNNNGKNMV YLQMTSLTPEDTGLYYCAAARDRGLHDGTTSDSYLEGSH EYEWGQGTQVTVSS
9.	4C5 CDR1	GRSVSTYF
10.	4C5 CDR2	IDRGSTVT
11.	4C5 CDR3	AAKAITRNFIATNDYDY
12.	4C5 VHH	QVQLVDSGGGLVQAGGSLRLSCAVSGRSVSTYFVGWFRQ APGKEREFVAAIDRGSTVTRYDDSVKGRFTISRDNKDTV YLQMNSLKPEDTAVYYCAAIAITRNFIATNDYDYWGQGT QVTVSS
13.	4D5 CDR1	GRAFNEYA
14.	4D5 CDR2	ISSDGRFT
15.	4D5 CDR3	AARDSGSGYYRAQWYDY
16.	4D5 VHH	VDSGGGAVKAGDSLRLVCSAPGRTHGRAFNEYAMAWFR QGPGERESVAAISSDGRFTYYAASVKGRFTISKDNKSA AFLQMNSLKPEDTAVYYCAARDSGSGYYRAQWYDYWG QGTQVTVSS
17.	4D11 CDR1	GSISSINA
18.	4D11 CDR2	ITNSGST
19.	4D11 CDR3	TARRSTWYIS
20.	4D11 VHH	QVQLQESGGGLVQPGGSLRLSCAASGSISSINAMGWYRLA PGKHREFVADITNSGSTNYAASVKGRFNISRDNKDTVYL QMNSLKFEDTAVYYCTARRSTWYISSGRGTQVTVSS

21.	4E5 CDR1	GLTSSDIA
22.	4E5 CDR2	ISSDGRFT
23.	4E5 CDR3	AARDSGSGYYISRAQWYDY
24.	4E5 VHH	QVQLVDSGGGVVEVGASLTLSCTSGLTSSDIAVGWFRQG PGKERESVAAISSDGRFTYYAASVKGRFTISKDNAKSA AFL QMNSLKPEDTAVYYCAARDSGSGYYISRAQWYDYWGQG TQVTVSS
25.	4H6 CDR1	GTIFRLNR
26.	4H6 CDR2	TIWSGRRT
27.	4H6 CDR3	NYRRITPWEASGNY
28.	4H6 VHH	QVQLVESGGGLATAGASLILSCAASGTIFRLNRMGWFRQA PGKERERVAATIWSGRRTHYADSVKGRFTISTDNAKKT VY LRMSSLKPEDTAVYYCNYRRITPWEASGNYWGQGTQVTV SS
29.	4H9 CDR1	GPIARSRS
30.	4H9 CDR2	AAISSDGRFT
31.	4H9 CDR3	AARDSGSGYYISRAQWYDY
32.	4H9 VHH	QVQLVESGGGAVQAGGSLRLSCTASGPIARSRSSTGMGWF RQGPGERESVAAISSDGRFTYYAASVKGRFTISKDNAKS AAFLQMNSLKPEDTAVYYCAARDSGSGYYISRAQWYDYW GQGTQVTVSS
33.	10H5 CDR1	ETTFKSMA
34.	10H5 CDR2	TNYNGGRT
35.	10H5 CDR3	AAKATEGTTFPSRTYEF
36.	10H5 VHH	GGGLVQAGGSLRLACTASDPPFANYETTFKSMAMGWVR HIPGKERELVAATNYNGGRTWYSNSAKARSTISRDN AKN TVYLMSSSLKPEDTAVYYCAAKATEGTTFPSRTYEFW GQ GIQVTVSS
37.	12H7 CDR1	GNFLSVSD
38.	12H7 CDR2	VTEHGRT
39.	12H7 CDR3	KASDVFTDAGAHEAVLIRDY
40.	12H7 VHH	QVQLVDSGGGLVQAGGSLRLSCKVSGNFLSVSDMSWYR QAPGMERDVVATVTEHGRTTYTDSVKGRFTISRDN AHT TYLEMNNLKPEDTAVYFCKASDVFTDAGAHEAVLIRDY W GQGTQVTVSS

41.	13H11 CDR1	GLTFSMYA
42.	13H11 CDR2	ISSDGRFT
43.	13H11 CDR3	AARDSGSGYYSTRAQWYDY
44.	13H11 VHH	QVQLVESGGGLVQAGGSLRLSCAASGLTFSMYAMGWFR QGPGKERESVAAISSDGRFTYYAASVKGRFTISKDNAKSA AFLQMNSLKPEDTAVYYCAARDSGSGYYSTRAQWYDYWG QGTQVTVSS
45.	15A5 CDR1	ERTFSSFA
46.	15A5 CDR2	IDPSGRYI
47.	15A5 CDR3	AARIRGEGYYTRSSFYHY
48.	15A5 VHH	QVQLVESGGGLVQAGGSLRLSCAASERTFSSFAMGWFRQ APGKEREVVAAIDPSGRYIYYKDSVKGRFTMSRDNAKSTV YLQMNSLKPDDTARYYYCAARIRGEGYYTRSSFYHYWGQ GTQVTVSS
49.	2B7 CDR1	GRTFSSYP
50.	2B7 CDR2	ISSDGRFT
51.	2B7 CDR3	AARDSGSGYYSTRAQWYDY
52.	2B7 VHH	QVQLVESGGGLVQAGGSLRLACAASGRTFSSYPMGWFRQ GPGKERESVAAISSDGRFTYYAASVKGRFTISKDNAKSAA FLQMNSLKPEDTAVYYCAARDSGSGYYSTRAQWYDYWGQ GTQVTVSS
53.	2B10 CDR1	SRIFRRYA
54.	2B10 CDR2	ITWSGAST
55.	2B10 CDR3	AADPWGSVIVGTAEYEY
56.	2B10 VHH	QVKLEESGGGLVQTGDSLRLSCAASSRIFRRYAMGWFRQ APGKEREFVAAITWSGASTTYTDSVKGRFTISRDSAENTTY LQMTSLRPEDTAVYYCAADPWGSVIVGTAEYEYWGQGT LTVTVSS
57.	3F10 CDR1	EHTFSNFP
58.	3F10 CDR2	IDSSGRLT
59.	3F10 CDR3	AARTGGVGYYSRSSFYNY
60.	3F10 VHH	QVQLVESGGGLVQAGGSLRLSCASSEHTFSNFPAMGWFRQ APGKERNVVAAIDSSGRLTYADSVKGRFTISKDNAKSTV YLQMNSLKSEDTARYYYCAARTGGVGYYSRSSFYNYWGQ GTLVTVSS

61.	3G6 CDR1	GSIFGISV
62.	3G6 CDR2	LTRAGLT
63.	3G6 CDR3	HANIMESAASTFGRY
64.	3G6 VHH	QVQLVESGGGLVQAGGSLSLSCAASGSIFGISVMGWYRQ APGEQRDLVATLTRAGLT TYGDSVKGRFSISRDSAKNTVY LQMNNLKPEDTAVYYCHANIMESAASTFGRYWGQGTQV TVSS
65.	3G7 CDR1	GRTLSTYT
66.	3G7 CDR2	AWPSPST
67.	3G7 CDR3	AADYKSLTQSWLNAALDY
68.	3G7 VHH	QVQLVESGGGLVQAGDSLRLSCEASGRTLSTYTMGWFRR APGKEREFVGLAWPSPSTYVVD SVKGRFTISRDN AKNTIY LQMNSLKPEDTAIYYCAADYKSLTQSWLNAALDYWGQG TQVTVSS
69.	3H7 CDR1	GSILSAGV
70.	3H7 CDR2	IALDGSTG
71.	3H7 CDR3	NANIRTD MRSAPFDH
72.	3H7 VHH	QVKLEESGGGLVQAGGSLRLSCAASGSILSAGVMRWYRQ APGKQRELVASIALDGSTGYIIDS VKGRFTISRDN AKNIVY LDMRSLEPADTAVYLCNANIRTD MRSAPFDHWGHGTQVT VSS
73.	4C6 CDR1	GRTFSSYP
74.	4C6 CDR2	ISSDGRFT
75.	4C6 CDR3	AVDPTGWGTIEADFRS
76.	4C6 VHH	QVQLVESGGGLVQAGGSLRLACAASGRTFSSYPMGWFRQ GPGKERESVAAISSDGRFTYYAASVKGRFTISKDNAKSAA FLQMNSLKPEDTAVYRCAVDPTGWGTIEADFRSWGQGTQ VTVSS
77.	1C12 CDR1	SRIFSRYG
78.	1C12 CDR2	ISWNGAST
79.	1C12 CDR3	AADPWGAVKLGTAEY EY
80.	1C12 VHH	QVQLVESGGGLVQTGDSLRLSCAASSRIFSRYGMGWFRQ APGKEREFVAAISWNGASTTYTDSVKGRFTISRDSAENTT YLQINSLRPEDTAVYYCAADPWGAVKLGTAEY EYWGQG TQVTVSS

81.	1G1 CDR1	GPSFSSYP
82.	1G1 CDR2	ISSDGRFT
83.	1G1 CDR3	AARDSGSGYYSSRAQWYDY
84.	1G1 VHH	QVQLVESGGGLVQAGDSLRLSCVASGPSFSSYPMGWFRQ GPGKERESVAAISSDGRFTYYAASVKGRFTISKDNAKSAA FLQMNSLKPEDTAVYYCAARDSGSGYYSSRAQWYDYWGQ GTLVTVSS
85.	1G1-F-G-ERES VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ GPGKERESVAAISSDGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWGQ GTLVTVSS
86.	1G1-F-A-ERES VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSDGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWGQ GTLVTVSS
87.	1G1-F-A- EREW VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKEREWVAAISSDGRFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWG QGTLVTVSS
88.	1G1-F-A- GLEW VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKGLEWVAAISSDGRFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWG QGTLVTVSS
89.	1G1-F-A- GREL VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKGRELVAISSDGRFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWG QGTLVTVSS
90.	1G1-F-A-GRES VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKGRESVAAISSDGRFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYYSSRAQWYDYWG QGTLVTVSS
91.	1C12-EREF VHH	QVQLVESGGGVVQPGRSLRLSCAASSRIFSR YGMGWFRQ APGKEREFVAAISWNGASTTYTDSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAADPWGAVKLGTA EYEWGQ GTQVTVSS

92.	1C12-EREW VHH	QVQLVESGGGVVQPGRSLRLSCAASSRIFSRYGMGWFRQ APGKEREWVAAISWNGASTTYTDSVKGRFTISRDNSKNTL YLQMNSLRAEDTAVYYCAADPWGAVKLGTAIEYEWGQ GTQVTVSS
93.	1C12_GLEW VHH	QVQLVESGGGVVQPGRSLRLSCAASSRIFSRYGMGWFRQ APGKGLEWVAAISWNGASTTYTDSVKGRFTISRDNSKNTL YLQMNSLRAEDTAVYYCAADPWGAVKLGTAIEYEWGQ GTQVTVSS
94.	2A3 CDR1	GGSFSSYP
95.	2A3 CDR2	ISSDMRFT
96.	2A3 CDR3	AARDSGVGYYRAQWYDY
97.	2A3 VHH	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDMRFTYYADSVKGRFTISRDNSKNTL YLQMNSLRAEDTAVYYCAARDSGVGYYRAQWYDYWG QGTLVTVSS
98.	1A8 CDR1	GPSFSSYP
99.	1A8 CDR2	ISSRGRFT
100.	1A8 CDR3	AARDSGSGYYRAQWYDY
101.	1A8 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSRGRFTYYADSVKGRFTISRDNSKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYRAQWYDYWGQ GTLVTVSS
102.	1D11 CDR1	GPSFSSSP
103.	1D11 CDR2	ISSMGRFT
104.	1D11 CDR3	AARDSGSGYYRAQWYDY
105.	1D11 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSSPMGWFRQA PGKERESVAAISSMGRFTYYADSVKGRFTISRDNSKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYRAQWYDYWGQ GTLVTVSS
106.	5E8 CDR1	GPSFSSYP
107.	5E8 CDR2	QSSDGRFT
108.	5E8 CDR3	AARDSGSGYYRAQWYDY
109.	5E8 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAQSSDGRFTYYADSVKGRFTISRDNSKNTL YLQMNSLRAEDTAVYYCAARDSGSGYYRAQWYDYWG

		QGTLVTVSS
110.	2A5 CDR1	GPSFSSYP
111.	2A5 CDR2	ISSVGRFT
112.	2A5 CDR3	AARDSGSGYYSRWQWYDY
113.	2A5 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSVGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRWQWYDYWGQ GTLVTVSS
114.	5G10 CDR1	GPRFSSYP
115.	5G10 CDR2	ISSDGRFT
116.	5G10 CDR3	AARDSGSGYYSRAQWYDG
117.	5G10 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPRFSSYPMGWFRQ APGKERESVAAISSDGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRAQWYDYGWQ GTLVTVSS
118.	2A6 CDR1	GPSFSLYP
119.	2A6 CDR2	ISSDRRFT
120.	2A6 CDR3	AARDSGSGYYSRAQWYDY
121.	2A6 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSLYPMGWFRQ APGKERESVAAISSDRRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRAQWYDYWGQ GTLVTVSS
122.	1G1-1C12 CDR1	GPSFSSYP
123.	1G1-1C12 CDR2	ISSDLRFT
124.	1G1-1C12 CDR3	AARDSGSGYYSRKQWYDY
125.	1G1-1C12 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRKQWYDYWGQ GTLVTVSS
126.	5B5 CDR1	GPSFSSYP
127.	5B5 CDR2	ISSDTRFT
128.	5B5 CDR3	AARDSGSGYYSRAQWYDR

129.	5B5 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSDTRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYY SRAQWYDRWGQ GTLVTVSS
130.	6F12 CDR1	GPSFTSYP
131.	6F12 CDR2	ISSDGRFK
132.	6F12 CDR3	AAEDSGSGYY SRAQWYDY
133.	6F12 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFTSYPMGWFRQ APGKERESVAAISSDGRFKYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAAEDSGSGYY SRAQWYDYWG QGTLVTVSS
134.	1C7 CDR1	GPSFSSYP
135.	1C7 CDR2	ISSRGRFT
136.	1C7 CDR3	AARGSGSGYY SRAQWYDY
137.	1C7 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSYPMGWFRQ APGKERESVAAISSRGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARGSGSGYY SRAQWYDYWGQ GTLVTVSS
138.	6E9 CDR1	GPSRSSYP
139.	6E9 CDR2	ISSDGKFT
140.	6E9 CDR3	AARDSGSGYY SRANWYDY
141.	6E9 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSRSSYPMGWFRQ APGKERESVAAISSDGKFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYY SRANWYDYWG QGTLVTVSS
142.	1A10 CDR1	GNSFSSYP
143.	1A10 CDR2	ISSDGRFS
144.	1A10 CDR3	ACRDSGSGYY SRAQWYDY
145.	1A10 VHH	QVQLVESGGGVVQPGRSLRLSCAASGNSFSSYPMGWFRQ APGKERESVAAISSDGRFSYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCACRDSGSGYY SRAQWYDYWGQ GTLVTVSS
146.	6B11 CDR1	GPSFPSYP
147.	6B11 CDR2	ISSRGRFT
148.	6B11 CDR3	AARDSGSGYY SRLQWYDY

149.	6B11 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSPSYPMGWFRQ APGKERESVAAISSRGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYY SRLQWYDYWGQ GTLVTVSS
150.	6D8 CDR1	GPSFSSKP
151.	6D8 CDR2	RSSDGRFT
152.	6D8 CDR3	AARDSGSGRYSRAQWYDY
153.	6D8 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSKPMGWFRQ APGKERESVAARSSDGRFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGRYSRAQWYDYWG QGTLVTVSS
154.	6D12 CDR1	GPSFSTYP
155.	6D12 CDR2	ISSDGVFT
156.	6D12 CDR3	AARDSGSGYY SREQWYDY
157.	6D12 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSTYPMGWFRQ APGKERESVAAISSDGVFTYYADSVKGRFTISRDN SKNTL YLQMNSLRAEDTAVYYCAARDSGSGYY SREQWYDYWG QGTLVTVSS
158.	2C5 CDR1	GPSFSTYP
159.	2C5 CDR2	ISSQGRFT
160.	2C5 CDR3	AARDSGSGYY SRAQWYDY
161.	2C5 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSTYPMGWFRQ APGKERESVAAISSQGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYY SRAQWYDYWGQ GTLVTVSS
162.	7F6 CDR1	GPMFSSYP
163.	7F6 CDR2	ISSDPRFT
164.	7F6 CDR3	AARDSGSGYY SRAQWYDY
165.	7F6 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPMFSSYPMGWFRQ APGKERESVAAISSDPRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYY SRAQWYDYWGQ GTLVTVSS
166.	2D5 CDR1	GPSFSSSP
167.	2D5 CDR2	ISWDGRFT
168.	2D5 CDR3	AARDSGSGYY SRAQWYVY

169.	2D5 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFSSSPMGWFRQA PGKERESVAAISWDGRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRAQWYVYWGQ GTLVTVSS
170.	7B11 CDR1	GPSFLIYP
171.	7B11 CDR2	ISSDGRFW
172.	7B11 CDR3	AARDSGSGYYSRVQWYDY
173.	7B11 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFLIYPMGWFRQA PGKERESVAAISSDGRFWYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGSGYYSRVQWYDYWGQ GTLVTVSS
174.	7D12 CDR1	GPSFLSYP
175.	7D12 CDR2	ISSDGRFS
176.	7D12 CDR3	AARDWGSGYYSRAQWYDY
177.	7D12 VHH	QVQLVESGGGVVQPGRSLRLSCAASGPSFLSYPMGWFRQ APGKERESVAAISSDGRFSYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDWGSGYYSRAQWYDYWGQ GTLVTVSS
178.	2A3ML CDR1	GGSFSSYP
179.	2A3ML CDR2	ISSDLRFT
180.	2A3ML CDR3	AARDSGVGYYSSRAQWYDY
181.	2A3ML VHH	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGVGYYSSRAQWYDYWGQ GTLVTVSS
182.	2A3MI CDR1	GGSFSSYP
183.	2A3MI CDR2	ISSDIRFT
184.	2A3MI CDR3	AARDSGVGYYSSRAQWYDY
185.	2A3MI VHH	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDIRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARDSGVGYYSSRAQWYDYWGQ GTLVTVSS
186.	2A3ML DT CDR1	GGSFSSYP
187.	2A3ML DT	ISSDLRFT

	CDR2	
188.	2A3ML DT CDR3	AARTSGVGYYSRAQWYDY
189.	2A3ML DT VHH	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARTSGVGYYSRAQWYDYWGQ GTLVTVSS
190.	2A3ML DE CDR1	GGSFSSYP
191.	2A3ML DE CDR2	ISSDLRFT
192.	2A3ML DE CDR3	AARESGVGYYSRAQWYDY
193.	2A3ML DE VHH	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARESGVGYYSRAQWYDYWGQ GTLVTVSS
194.	2A3 LTFc	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARTSGVGYYSRAQWYDYWGQ GTLVTVSSEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPK DTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFL YSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSP GK
195.	Exemplary linker	GSGGSGGSGGSG
196.	Exemplary linker	GGGGSGGGGSGGGGS
197.	Exemplary linker	GGGSG
198.	Exemplary linker	GGGSGGGGSG

199.	Exemplary linker	GGSGGGSG
200.	Exemplary linker	GGSGGGSGGGSG
201.	Exemplary linker	GSGGSG
202.	Exemplary linker	GSGGSGGSG
203.	Exemplary linker	GSGSGSG
204.	Exemplary linker	GGGGSGGGGSGGGGSGGG
205.	Exemplary linker	PAPAP
206.	Exemplary linker	PAPAPPAPAPPAPAP
207.	Exemplary linker	IKRTVAA
208.	Exemplary linker	VSSASTK
209.	Exemplary linker	AEAAAKA
210.	Exemplary linker	AEAAAKEAAKA
211.	Exemplary linker	GRPGS GRPGS
212.	Exemplary linker	GRPGS GRPGS GRPGS GRPGS
213.	Exemplary linker	GRGGS GRGGS
214.	Exemplary linker	GRGGS GRGGS GRGGS GRGGS
215.	Exemplary linker	GKPGS GKPGS
216.	Exemplary	GKPGS GKPGS GKPGS GKPGS

	linker	
217.	Exemplary linker	GEPGS GEPGS
218.	Exemplary linker	GEGGS GEGGS GEGGS GEGGS
219.	Exemplary linker	GDPGS GDPGS
220.	Exemplary linker	GDPGS GDPGS GDPGS GDPGS
221.	H1G4CDR-H1	FTFSNYGMS
222.	H1G4CDR-H2	TISGGGSNIY
223.	H1G4CDR-H3	VSYYYGIDF
224.	H1G4CDR-L1	KASQDVTTAVA
225.	H1G4CDR-L2	WASTRHT
226.	H1G4CDR-L3	QQHYTIPWT
227.	H1G4VH	QVQLVESGGGLVKPGGSLRLSCAASG <u>FTFSNYGMS</u> WIRQ APGKGLEWVSTISGGGSNIYADSVKGRFTISRDNKNSL YLQMNSLR AEDTAVYYC <u>VSYYYGIDF</u> WGQGTSVTVSS
228.	H1G4VL	DIQMTQSPSSLSASVGDRVTITC <u>KASQDVTTAVA</u> AWYQQK PGKAPKLLIY <u>WASTRHT</u> GVPSRFSGSGSGTDFTLTISLQP EDFATYYC <u>QQHYTIPWT</u> FGGGTKLEIKR
229.	33B CDR-H1	FRFSNYGMS
230.	33B CDR-H2	TISGGGSNAY
231.	33B CDR-H3	TSYYYGIDF
232.	33B CDR-L1	KASTDVTTAVA
233.	33B CDR-L2	WASLRHT
234.	33B CDR-L3	QQHYGIPWT
235.	33B VH	QVQLVESGGGLVKPGGSLRLSCAASG <u>FRFSNYGMS</u> WIRQ APGKGLEWVSTISGGGSNAYADSVKGRFTISRDNKNS LYLQMNSLR AEDTAVYYC <u>TSYYYGIDF</u> WGQGTSVTVSS
236.	33B VL	DIQMTQSPSSLSASVGDRVTITC <u>KASTDVTTAVA</u> AWYQQK PGKAPKLLIY <u>WASLRHT</u> GVPSRFSGSGSGTDFTLTISLQP EDFATYYC <u>QQHYGIPWT</u> FGGGTKLEIKR
237.	66E CDR-H1	FTFSNYGMS
238.	66E CDR-H2	TISGGGSNIY

239.	66E CDR-H3	VSYYYGIDL
240.	66E CDR-L1	KAKQDVTTAVA
241.	66E CDR-L2	WASTRHT
242.	66E CDR-L3	QQHYWIPWT
243.	66E VH	QVQLVESGGGLVKPGGSLRLSCAASG <u>FTFSNYGMS</u> WIRQ APGKGLEWVSTISGGGSNIYYADSVKGRFTISRDNANKNSL YLQMNSLRAEDTAVYYC <u>VSYYYGIDL</u> WGQGTSTVTVSS
244.	66E VL	DIQMTQSPSSLSASVGDRVTITC <u>KAKQDVTTAVA</u> AWYQQK PGKAPKLLIY <u>WASTRHT</u> GVPSRFSGSGSGTDFTLTISLQP EDFATYYC <u>QQHYWIPWT</u> FGGGTKLEIKR
245.	711D CDR-H1	FTFSNYGMS
246.	711D CDR-H2	TISGGGSNIY
247.	711D CDR-H3	SSYYYGIDL
248.	711D CDR-L1	KASQDVTNAVA
249.	711D CDR-L2	WASTRHT
250.	711D CDR-L3	QQHYTIPWT
251.	711D VH	QVQLVESGGGLVKPGGSLRLSCAASG <u>FTFSNYGMS</u> WIRQ APGKGLEWVSTISGGGSNIYYADSVKGRFTISRDNANKNSL YLQMNSLRAEDTAVYYC <u>SSYYYGIDL</u> WGQGTSTVTVSS
252.	711D VL	DIQMTQSPSSLSASVGDRVTITC <u>KASQDVTNAVA</u> AWYQQK PGKAPKLLIY <u>WASTRHT</u> GVPSRFSGSGSGTDFTLTISLQP EDFATYYC <u>QQHYTIPWT</u> FGGGTKLEIKR
253.	anti-PD1 HC - Linker- 2A3 (HCC heavy chain)	QVQLVESGGGLVKPGGSLRLSCAASGFTFSNYGMSWIRQ APGKGLEWVSTISGGGSNIYYADSVKGRFTISRDNANKNSL YLQMNSLRAEDTAVYYCVSYYYGIDFWGQGTSTVTVSSAS TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVTVPPSSSLGTKTYTCN VDHKPSNTKVDKRVESKYGPCCPPCPAPEFLGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEV HNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLS LSLGKSGSGSGSGSGSGQVQLVESGGGVVQPGRSLRLSC AASGGSFSSYPMGWFRQAPGKERESVAAISSDLRFTYYAD

		SVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAARTS GVGYYSRAQWYDYWGQGTLVTVSS
254.	anti-PD1 LC1 (HCC light chain)	DIQMTQSPSSLSASVGDRVTITCKASQDVTTAVAWYQQKP GKAPKLLIYWASTRHTGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQHYTIPWTFGGGTKLEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE SVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGL SSPVTKSFNRGEC
255.	anti-PD1HC1 (LCC heavy chain)	QVQLVESGGGLVKPGGSLRLSCAASGFTFSNYGMSWIRQ APGKGLEWVSTISGGGSNIYYADSVKGRFTISRDN AKNSL YLQMNSLRAEDTAVYYCVSYYYGIDFWGQGTSVTVSSAS TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTKTYTCN VDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPP KPKDTLMISRTPEVTCVVDVVSQEDPEVQFNWYVDGVEV HNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS FFLYSRLTVDKSRWQEGNVFSCSVMEALHNHYTQKSLS LSLGK
256.	anti-PD1 LC- Linker-2A3 (LCC light chain)	DIQMTQSPSSLSASVGDRVTITCKASQDVTTAVAWYQQKP GKAPKLLIYWASTRHTGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQHYTIPWTFGGGTKLEIKRTVAAPSVFIFPPSD EQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE SVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGL SSPVTKSFNRGECGSGSGSGSGSGQVQLVESGGGVVQP GRSLRLSCAASGGSFSSYPMGWFRQAPGKERESVAAISSD LRFTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVY YCAARTSGVGYYSRAQWYDYWGQGTLVTVSS
257.	2A3-Linker- anti-PD1 HC (HCN heavy chain)	QVQLVESGGGVVQPGSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDN SKNTLY LQMNSLRAEDTAVYYCAARTSGVGYYSRAQWYDYWGQ GTLVTVSSGSGSGSGSGSGQVQLVESGGGLVKPGGSLRL SCAASGFTFSNYGMSWIRQAPGKGLEWVSTISGGGSNIYY ADSVKGRFTISRDN AKNSL YLQMNSLRAEDTAVYYCVSY

		YYGIDFWGQGTSVTVSSASTKGPSVFPLAPCSRSTSESTAA LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVVTVPSSSLGKTKYTCNVDPHKPSNTKVDKRVESKYGP PCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVD VSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPR EPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESN GQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFS CSVMHEALHNHYTQKSLSLGLK
258.	anti-PD1 LC2 (HCN light chain)	DIQMTQSPSSLSASVGDRVTITCKASQDVTTAVAWYQQKP GKAPKLLIYWASTRHTGVPSRFSGSGSGTDFTLTISSLQPE DFATYYCQQHYTIPWTFGGGTKLEIKRTVAAPS VFIFPPSD EQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE SVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGL SSPVTKSFNRGEC
259.	anti-PD1HC2 (LCN heavy chain)	QVQLVESGGGLVKPGGSLRLSCAASGFTFSNYGMSWIRQ APGKGLEWVSTISGGGSNIYYADSVKGRFTISRDNANKNSL YLQMNSLRAEDTAVYYCVSYYYGIDFWGQGTSVTVSSAS TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTKYTCN VDPHKPSNTKVDKRVESKYGP PCPPCPAPEFLGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEV HNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGS FFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLS LSLGLK
260.	2A3-Linker- anti-PD1LC (LCN light chain)	QVQLVESGGGVVQPGRSLRLSCAASGGSFSSYPMGWFRQ APGKERESVAAISSDLRFTYYADSVKGRFTISRDNANKNTLY LQMNSLRAEDTAVYYCAARTSGVGYYSSRAQWYDYWGQ GTLVTVSSGSGSGSGSGSGSDIQMTQSPSSLSASVGDRVTI TCKASQDVTTAVAWYQQKPGKAPKLLIYWASTRHTGVPS RFSGSGSGTDFTLTISSLQPEDFATYYCQQHYTIPWTFGGG TKLEIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPRE AKVQWKVDNALQSGNSQESVTEQDSKDYSLSSSTLTLS

		KADYEKHKVYACEVTHQGLSSPVTKSFNRGEC
261.	Human TIGIT polypeptide	MRWCLLLIWA QGLRQAPLAS GMMTGTIETT GNISAEKGGG IILQCHLSST TAQVTQVNWEQQDQLLAICN ADLGWHISPS FKDRVAPGPG LGLTLQSLTV NDTGEYFCIY HTYPDGTYTGRIFLEVLESS VAEHGARFQI PLLGAMAATL VVICTAVIVV VALTRKKKAL RIHSVEGDLRRKSAGQEEWS PSAPSPPGSC VQAEAAPAGL CGEQRGEDCA ELHDYFNVLS YRSLGNCSSFTETG
262.	ECD of Human TIGIT polypeptide	MRWCLLLIWA QGLRQAPLAS GMMTGTIETT GNISAEKGGG IILQCHLSST TAQVTQVNWEQQDQLLAICN ADLGWHISPS FKDRVAPGPG LGLTLQSLTV NDTGEYFCIY HTYPDGTYTGRIFLEVLESS VAEHGARF

The following examples are merely illustrative of the presently disclosed subject matter and should not be considered as limitations in any way.

5

EXAMPLES

Example 1. Immunization, generation of anti-human TIGIT VHH antibodies & hits identification

Antigen of recombinant human TIGIT extra cellular domain (ECD) protein was purchased from Arco Bio. Immunization of TIGIT was performed using llama under protocols known in the art. The titer of serum antibodies was measured by ELISA assays. After 3 rounds of immunization, a high titer (1:100,000) was observed. Whole blood was then collected, and PBMCs were isolated. RNA was then isolated from the PBMCs.

The VHH antibody genes were amplified by PCR under protocols know in the art, purified by DNA agarose gel, constructed into a phagemid vector pADL-23c (Antibody Design Labs) and transformed to TG1 electrocompetent cells (from Lucigen). Transformed TG1 cells were cultured in Y2T medium. Phages with target VHH displayed were reproduced by adding helper phages and co-culturing overnight. Phages in supernatants of culture were harvested by centrifugation, and panning of binders to human-TIGIT (h-TIGIT) or cynomolgus-TIGIT (cyno-TIGIT) antigen was performed using streptavidin-coupled Dynabeads coated with biotinylated h-TIGIT or cyno-TIGIT ECD. After 3 rounds of panning, binders of h-TIGIT or cyno-TIGIT were eluted, which were used to infect SS320 cells. Colonies of SS320 cells were picked and cultured in Y2T medium, and IPTG was added for secretion of VHH antibodies. Supernatants with VHH antibodies were screened by

ELISA assays using h-TIGIT ECD coated plates. Positive h-TIGIT binders were picked for sequencing. 29 clones with different sequences were selected. Binding ability of VHH antibodies on cyno-TIGIT was also examined by ELISA. The top 21 binders and their CDRs and VHs are shown in the Sequence Table (SEQ ID NOS: 1-84).

The effects of VHH antibody clones on blocking Poliovirus Receptor (PVR, a.k.a. CD155) binding to TIGIT were also determined using blocking ELISA assay. 9 clones with over 90% inhibition of PVR binding to h-TIGIT were further selected (clone name: 2B7, 1G1, 1C12, 3G6, 2B10, 3G7, 3F10, 13H11 and 15A5).

Example 2 – Characterization and selection of TIGIT VHH antibodies

Antibody clones identified from Example 1 were constructed to make bivalent antibody by adding human constant heavy chain2 (CH2) and constant heavy chain 3 (CH3) domains shown in Figure 1C. Constructed bivalent VHH antibodies were expressed in ExpiCHO cells, and proteins in supernatants were harvested and purified by Protein A.

Binding affinities of the bivalent clones to human TIGIT transfected Jurkat cells were confirmed by flowcytometry assays. Jurkat cells stably expressing human TIGIT and NFAT reporter gene were established. Specifically, Jurkat cells were transfected with human TIGIT expression vector by electroporation, and cells stably expressing human TIGIT were selected by 1 µg/ml puromycin during cell culture. Representative antibody clones were incubated at different concentrations with h-TIGIT stably expressing Jurkat cells (0.2x10⁶/ml) in 100 µl/well in 96-well plate in FACS buffer (PBS with 1.5 % FBS) for 30 min. After washing, Alexafluor 488 conjugated anti-human IgG Fc secondary antibody (Alexa Fluor® 488 AffiniPure Goat Anti-Human IgG, Fcγ fragment specific, Jackson labs, 1:500 dilution) was added and incubated for 30 min. After washing, mean fluorescence intensity was measured using CytoFlex (Beckman Coulter) by gating live cell population. Binding affinity was calculated using GraphPad Prism. Representative results are shown in Figures 1A and 1B. All tested antibodies showed high binding affinity to cells expressing human TIGIT compared to Reference Ab 1, a reference anti-human-TIGIT antibody disclosed in U.S. 2016/0176963 A1 as BMS 22G2, which was synthesized in-house based on the disclosed sequences.

Furthermore, Jurkat cells stably expressing human TIGIT were transfected with NFAT reporter gene using electroporation. NFAT reporter expressing cells were selected by 300 µg/ml hygromycin in culture medium. Moreover, Raji cells were stably transfected with human PVR, and transfected cells were selected by hygromycin at 125 µg/ml in culture medium. The antibodies' effects on PVR-mediated suppression of TCR-induced NFAT reporter activity were then determined by co-incubation of Raji cells expressing human PVR and Jurkat cells expressing human TIGIT and NFAT reporter gene in the presence of staphylococcal enterotoxin (SEE, 0.01 ng/ml) for

5 hours. Bright-Glo luciferase assay buffer with substrate(Promega) was added, and luciferase activity was measured by chemiluminescence activity using a plate reader. Potency of antibody in blocking TIGIT activity was calculated using non-linear regression method by GraphPad Prims. Representative results are shown in Figure 2. All tested clones except 3F10 showed similar TIGIT blocking effects compared to the reference antibody Reference Ab 1.

Example 3 – Humanization of anti-TIGIT antibodies

Two representative clones 1G1 and 1C12 were selected for humanization of their framework. Briefly, Igbblast was performed using the sequences of the two clones to search database of human germline genes. Ideal germline sequences were selected, and mutations of framework sequences were made to change the framework sequences from llama to human. For 1C12 clone, human germline IGHV-3-30*10 was used, and three version of humanized 1C12 were made (1C12-EREF, 1C12-EREW and 1C12-GLEW). For 1G1, human germline IGHV-3-30*01 was used, and six versions of humanized 1G1 were made (1G1-F-G-ERES, 1G1-F-A-ERES, 1G1-F-A-EREW, 1G1-F-A-GLEW, 1G1-F-A-GREL and 1G1-F-A-GRES). The constructs were cloned into expressionvectors, and antibody proteins were produced by transient transfection of ExpiCHO and purified by protein A.

The binding affinities of the humanized bivalent antibodies to human TIGIT were determined by whole cell binding to h-TIGIT stably expressed on Jurkat cells. Briefly, antibodies were incubated with h-TIGIT transfected Jurkat cells (0.2×10^6 /well in 100 μ l) for 30 min in FACS buffer. The cells were washed once, then incubated with anti-human IgG Fc AlexaFluor488 (1:500). Mean fluorescence intensity was determined by CytoFlex, and binding affinities of antibodies to h-TIGIT were calculated using non-liner regression by GraphPad Prism 8.0 as shown in Figure 3A. Similar binding affinities were observed for all three versions of 1C12 clones compared to the chimeric parentalclone. For 1G1 clone, two humanized versions, F-G-ERES and F-A-ERES,exhibited similar affinity to h-TIGIT compared to chimeric clone, whereas two versions (F-A-EREW and F-A-GLEW) lost binding affinity to h-TIGIT as shown in Figure 3B.

The activity of humanized antibodies in blocking PVR-mediated suppressive effects on TCR-mediated NFAT reporter activity was determined using a method described previously. Briefly, human TIGIT and NFAT reporter transfected Jurkat cells were incubated with antibodies in different concentrations, in the presence of PVR transfected Raji cells and a low concentration of staphylococcal enterotoxin (SEE, 0.01 ng/ml) for 5 hours. Bright-Glo luciferase assay buffer with substrate (Promega) was added, and luciferase activity was measured. Representative results are shown in Figures 4A and 4B. All humanized versions of 1C12 clone had similar potency in blocking TIGIT compared to the chimeric parentalclone and Reference Ab 1, a reference anti-h-TIGIT antibody. All humanized versions of 1G1 had similar potency compared to the chimeric

parental clone in blocking TIGIT, except two versions 1G1-(F-A-EREW) and 1G1-(F-A-GLEW) lost the blocking effect.

Example 4 – Affinity maturation, selection & modification

Affinity maturation was performed for the 1G1-F-A-ERES clone. Primers for making single mutation of amino acid for each CDR region were designed. A library of mutations was prepared using assembly PCR and cloned into phagemid vectors. Library quality was measured by transformation of TG1 cells and DNA sequencing of clones. Phage production was carried out using helper phages, and phage panning was performed using streptavidin-coupled Dynabeads coated with biotinylated h-TIGIT ECD or cyno-TIGIT ECD. After two round of panning, elution of panning products was used to infect SS320 cells, and colonies were picked and cultured in Y2T medium with IPTG. VHH antibodies in supernatants were examined by ELISA assays. Positive clones against h- & cyno-TIGIT were selected for whole cell binding to h- & cyno-TIGIT in stable cells. PVR blocking ELISA and human TIGIT blockade NFAT reporter assays were also performed for selected clones. EC₅₀ or IC₅₀ values were calculated using GraphPad Prism. The top 25 binders and their CDRs and VHHs are shown in the Sequence Table (SEQ ID NOS: 94-177).

Correlations between EC₅₀/IC₅₀ in whole cell binding and blocking ELISA were graphed using GraphPad Prism, and representative data are shown in Figure 5. 2A3 was identified with the highest affinity and potency.

The thermostability of 12 representative clones was tested by heat treatment at 25 to 70 °C for 60 min, then the binding of the treated samples to human TIGIT was examined using ELISA and whole cell binding flow cytometry assays, the results of which are shown in Figures 6A and 6B. Clone 2A3 exhibited superior thermostability compared to other clones.

Bivalent 2A3 antibody (2A3-Fc) was constructed using human IgG1 CH2 and CH3 domains. The antibody was expressed in ExpiCHO cells and purified by protein A column. Epitope of 2A3-Fc binding to human TIGIT was studied using Octet binding assay in comparison with anti-TIGIT reference antibodies Reference Ab 1 and Reference Ab 2 (a reference anti-human-TIGIT antibody having the same amino acid sequences of Tiragolumab, which was synthesized in-house based on the disclosed sequences in U.S. 2017/0088613A1). As shown in Figures 7A-7C, 2A3 clone bound to a different epitope compared to Reference Ab 1 and Reference Ab 2.

Cross-species binding activity of 2A3-Fc to human, cynomolgus and mouse TIGIT was determined by ELISA assays, results of which are shown in Figures 8A-8C. 2A3-Fc bound to human, cynomolgus monkey but not mouse TIGIT.

Analysis of CDR regions of 2A3 identified two hotspots, a methionine in CDR2 and an aspartic acid in CDR3 followed by serine. Mutations of methionine to leucine and isoleucine, and mutations of aspartic acid to threonine and glutamic acid were carried out. CDRs and VHHs of

the modified antibodies (2A3 ML, 2A3 MI, 2A3 ML_DT (a.k.a 2A3 LT) and 2A3 ML_DE) are shown in the Sequence Table (SEQ ID NOs: 178-193). The modified versions were tested in whole cell binding and NFAT luciferase reporter assays, all of which exhibited similar properties compared to the parental 2A3 clone. The representative data of 2A3-LT-Fc with changes of M to L and D to T (2A3 ML DT) are shown in Figure 9, where similar human TIGIT binding affinity in whole cell binding assay and similar potency in NFAT reporter assay were observed compared to parental 2A3-Fc.

The affinity and potency of 2A3-Fc were compared to a reference anti-TIGIT antibody Reference Ab 2. 2A3-Fc exhibited significantly higher affinity than Reference Ab 2. The EC₅₀ value of 2A3-Fc was 0.32 ± 0.06 nM compared to 0.61 ± 10.13 nM of Reference Ab 2 (n=3) in whole cell binding assays (Figure 10A). 2A3-Fc also exhibited similar potency of blocking TIGIT in NFAT luciferase reporter assays compared to Reference Ab 2. The EC₅₀ value of 2A3-Fc was 0.48 ± 1.018 nM compared to 0.72 ± 0.42 nM of Reference Ab 2 (n=3) (Figure 10B).

Example 5 – In vitro antitumor efficacy study

The anti-TIGIT antibody's antitumor effects were further examined in vitro. In the tumor microenvironment, when TIGIT⁺ T cells are in contact with PVR⁺ dendritic cells (DCs), PVR expressed on the DCs can bind to TIGIT expressed on the T cells and thereby suppresses the T cells' antitumor activity, e.g., antitumor cytokine secretion. Treatment using an effective anti-TIGIT antibody can block the interaction between TIGIT and PVR and thereby enhance the antitumor activity of T cells.

Mixed lymphocyte reaction assay (MLR) was used to model this phenomenon in the tumor microenvironment. CD14⁺ monocytes isolated from PBMCs of one healthy donor were cultured with 50 ng/ml IL-4 and 50 ng/ml GM-CSF in RPMI for 7 days (medium was changed on Day 4) to differentiate into DCs. Mature DCs were obtained by further culturing the DCs with 100 ng/ml LPS in RPMI for 1 day on Day 7. The matured DCs stably expressed CD11c (mature DC biomarker), MHC class II, CD80 (CD28 ligand), and PVR (TIGIT ligand). CD3⁺ T cells were isolated from PBMCs of another healthy donor using ThermoFisher T cell isolation kit and cultured overnight. Next, the CD3⁺ T cells (200,000) were mixed with the mature DCs (10,000) in RPMI 1640 culture medium with 10% FBS-HI, and anti-TIGIT antibody or control antibodies were added at various concentrations. After incubation for 48 hours at 37°C and 5.0% CO₂, IL-2 secretion from the T cells were measured in culture supernatant by PerkinElmer (AlphaLisa human IL-2 kit: Cat#AL221C). An anti-PD1 antibody was used as a positive control. An anti-HER2 antibody was used as a negative control.

As shown in Figure 11, 2A3-LT-Fc enhanced the IL-2 secretion from T cells in a dose dependent manner comparable to the anti-PD1 antibody. In contrast, Reference Ab 2, a

Tiragolumab analog, did not enhance the IL-2 secretion. As IL-2 is an important antitumor cytokine, the results indicate that the anti-TIGIT antibody 2A3-LT-Fc can significantly enhance T cell antitumor activity in the tumor microenvironment, and it exhibited superior antitumor effects compared to the Tiragolumab analog.

5

Example 6 – Fc engineering

The mechanism of action of anti-TIGIT antibodies in enhancing immune function & anti-tumor activity can involve not only blockade effect on TIGIT but also bridging effect between antigen presenting cells (APC) and effective T cells (CD8⁺ T cells) through the binding of the antibody Fc region by FcγRIIIA on APC and the binding of the VHH domain to TIGIT on effective T cells. To take advantage of this mechanism, two different Fc-enhanced versions were made, one with mutations of DLE (S239D, A330L and I332E), one with mutations of VLPLL (L235V, F243L, R292P, Y300L and P396L). The binding of the Fc-mutants to human FcγRIIIA and FcγRIIB was examined by Octet binding assay using recombinant proteins of the ECD of FcγRIIIA and FcγRIIB, the results of which are shown in Figure 12. Both DLE and VLPLL mutants had enhanced binding affinity to human FcγRIIIA. DLE mutant also had enhanced binding affinity to human FcγRIIB, whereas VPVLL mutant had reduced binding affinity to human FcγRIIB.

The TIGIT blockade activity of a Fc-mutant was also examined, where the DLE mutant showed reduced TIGIT blockade function as shown in Figure 13A.

Moreover, when antibody variable region binds to its specific antigen, its Fc region can cross-link the FcγRIIIA and triggers down-stream signaling. Using this mode of action, the effect of Fc-mutants on FcγRIIIA-mediated activity was examined using a human-FcγRIIIA and NFAT reporter transfected Jurkat cells and 293T cells overexpressing TIGIT. In this assay system, the DLE mutant significantly increased human-FcγRIIIA-mediated NFAT reporter activity compared to the wild type Fc version as shown in Figure 13B.

Table 3. Test results shown in Figure 13A

EC50 (nM)	2A3 LT-Fc	DLE
Exp 1	0.58	0.67
Exp 2	0.687	0.92
Exp 3	0.45	1.32
Average	0.57	0.97
SD	0.12	0.3279

Table 4. Test results shown in Figure 13B

EC50 (nM)	2A3 LT-Fc	DLE
-----------	-----------	-----

Exp 1	0.226	0.027
Exp 2	0.175	0.0295
Exp 3	0.144	0.018
Exp 4	0.18	0.02
Average	0.18	0.024
SD	0.03	0.005

Example 7—*In vivo* efficacy study

Using h-TIGIT Knock-In C57BL/6 mice and MC38 murine colon cancer model, efficacies of 2A3-LT-Fc wt and, 2A3-LT-Fc-DLE were studied and compared to Reference Ab 2. MC38 tumor cells were implanted one week before treatment. Treatment started when tumor volume reached about 51 mm³, when drugs were dosed intraperitoneally twice a week for 2.5 weeks at 6 mg/kg for 2A3-LT-Fc antibodies, or 11 mg/kg for reference antibody (equal to 6 mg/kg of 2A3-LT-Fc in mole/kg dosage). On day 16 post-treatment, tumor volumes in multiple mice of the control group reached tumor size limits (2000 mm³). Therefore, day 16 post-treatment was the data end point for the analyses. Average tumor volume in the control group was 1548.76±191mm³ (mean±SEM) on day 16, the treatment with 2A3-LT-Fc wt and 2A3-LT-Fc-DLE significantly reduced tumor growth compared to vehicle control, resulting in respectively 38% and 50% of TGI (tumor growth inhibition) and tumor volume of 985.05±123mm³ and 802.20±126mm³ (P=0.037 and P=0.007 compared to Control by Mann-Whitney test), respectively. The treatment with the reference antibody also reduced tumor growth, but the reduction caused by the antibody was not statistically significant compared to the control group with the TGI of 26%, and tumor volume of 1156.16±195mm³ (Figure 14A, Table 5). The results of individual tumor volume are shown in Figure 14B. These results indicated that both 2A3-LT-Fc and 2A3-LT-Fc-DLE were more efficacious than Reference Ab 2. Body weight changes were not significant between the groups throughout the study (Figure 14C), indicating the treatments were well tolerated.

Table 5. Mean Tumor Volumes on Day 16 in MC38 Syngeneic h-TIGIT Mouse Model

Groups	Treatment	N	TV (mm ³) ^a	P ^b	% TGI (Day 16)
1	Vehicle	8	1548.76±191		
2	2A3-LT-Fc-wt 6 mg/kg	8	985.05±123	0.037	38
3	2A3-LT-Fc-DLE 6 mg/kg	8	802.20±126	0.007	50
4	Reference Ab 2, 11 mg/kg	8	1156.16±195	0.234	26

^a Mean $\pm$ SEM

^b Compared to vehicle control by Mann-Whitney test on day 16 post-treatment

Example 8– Anti-TIGITxanti-PD1 bispecific antibody

Bispecific antibodies (BsAbs) were constructed using humanized, affinity matured anti-TIGIT clone 2A3 and humanized, affinity matured anti-PD1 antibody disclosed in PCT/US2017/050851 (published as International Publication WO 2018/052818 A1). Four different versions were made, where VHH 2A3 was fused to N- or C-terminal of the heavy chain of the anti-PD1 antibody, or N- or C-terminal of the light chain of the anti-PD1 antibody (named as HCN, HCC, LCN, LCC, respectively). The structures of the BsAbs are shown in Figure 15, and the sequences are shown in SEQ ID NOs: 253-260 in the Sequence Table. Production of the BsAbs were made from transient transfection of ExpiCHO and purified by Protein A column.

Double binding of the BsAbs to PD1 or TIGIT was confirmed using octet binding assay. Briefly, sensors were loaded with BsAbs and then injected with different concentrations of recombinant protein of h-PD1 ECD (his-tagged). After association of BsAbs with h-PD1 ECD, the sensors were injected again with different concentrations of recombinant protein of h-TIGIT ECD (his-tagged). As shown in Figures 16A-16F, all the BsAbs bound to both h-TIGIT and h-PD1 with similar affinity compared to parental mAb anti-PD1 and anti-TIGIT (2A3-Fc).

Affinity to TIGIT was further examined in whole cell binding assay using human TIGIT transfected Jurkat cells. Antibodies were incubated with h-TIGIT transfected Jurkat cells, and binding of antibodies to h-TIGIT was determined using a secondary anti-human-IgG Fc antibody conjugated with AlexaFluor488, which was detected by flow cytometry. HCN and LCN versions of the BsAbs kept similar affinity to TIGIT expressed on cells compared to parental mAb 2A3-Fc, whereas both HCC and LCC versions showed reduced affinities to TIGIT compared to 2A3-Fc as shown in Figure 17.

Furthermore, cell based TIGIT blockade luciferase reporter assay was carried out using the BsAbs. Antibodies were cultured with h-TIGIT and NFAT reporter transfected Jurkat cells in the presence of PVR transfected Raji cells and a low concentration of staphylococcal enterotoxin. TCR-mediated NFAT luciferase reporter activity was measured by a chemiluminescent plate reader. HCN version showed similar potency compared to parental anti-TIGIT 2A3-Fc, whereas all other three versions of BsAbs showed reduced potency compared to 2A3-Fc as shown in Figure 18.

The binding affinities of the BsAbs to human PD1 were also determined by whole cell binding assays using human PD1 transfected Jurkat cells. Antibodies were incubated with h-PD1 transfected Jurkat cells, and binding was determined using a secondary anti-h-IgG antibody conjugated with AlexaFluor488, which was detected by flow cytometry. All the BsAbs showed slightly lower affinities than parental anti-PD1 antibody as shown in Figure 19.

The potency of the BsAbs in blocking PD1-mediated suppressive effect was determined by PD1 blockade reporter assay. Antibodies were cultured with h-PD1 and NFAT reporter transfected Jurkat cells in the presence of PDL1 and CD3 transfected CHO cells. TCR-mediated NFAT luciferase activity was measured by a chemiluminescent plate reader. Similar to the affinity, all the BsAbs had slightly lower potency of blocking PD1 than the monoclonal anti-PD1 antibody as shown in Figure 20.

As the potency measured above only indicated the effect of the BsAbs on a single target (either TIGIT or PD1), the potencies of BsAbs toward both targets were examined using a combination blockade assay (double blockade assay). As shown in Figure 21, mAbs, combination of mAbs and BsAbs were cultured with h-TIGIT and h-PD1 & NFAT reporter transfected Jurkat cells in the presence of PVR and PDL1 transfected CHO cells and a low concentration of staphylococcal enterotoxin. TCR-mediated NFAT luciferase reporter activity was measured by a chemiluminescent plate reader and shown in Y axis in relative luminescence units. The combination of anti-TIGIT and anti-PD1 was more potent than either anti-TIGIT or anti-PD1 mAbs. All the BsAbs were more potent compared to the combination of anti-TIGIT (aTIGIT) and anti-PD1 (aPD1), with LCC version being the most potent BsAb followed by HCN and LCN versions.

The developability of four versions of the BsAbs was assessed by SDS PAGE gel, SEC and DLS analysis. Results of SDS PAGE gel showed that LCC and LCN versions had degradation and unpaired light chains. SEC data also showed that the percentage monomers were less than 84% with percentage of low molecular weight higher than 15% for either LCC or LCN versions of the BsAbs, indicating low stability and developability for both BsAbs. In addition, in DLS assay, the Tonset for both LCC and LCN were lower than 60 °C, suggesting low thermostability. DLS results showed that HCC version had much higher radius than other versions of the BsAbs, suggesting aggregation of this form. The summary results are shown in Table 4. HCN version of the BsAb demonstrated superior characteristics than other three versions.

Table 4. summary of results of in vitro assays and developability

EC50 (Mean, nM)	TIGIT WCB	TIGIT Blockade Assay	PD1 WCB	PD1 Blockade Assay	PD1+TIGIT Combination Blockade Assay	SEC Monomer	DLS Radius	DLS Tonset	SDS PAGE
aTIGIT-Fc	0.35	2.3	/	/		95%		60	
aPD1	/	/	0.6	0.73		99%	7-9	58	
aTIGIT-Fc +	/	/	/	/	7.37				

aPD1									
aTIGITxaPD 1-HCN	0.81	2.7	2.24	1.52	1.15	99%	6	64	+++
aTIGITxaPD 1-HCC	4.93	25.9	2.84	3.29	2.61	93.40%	20	65	+++
aTIGITxaPD 1-LCN	0.75	8.7	1.82	4.64	1.23	84.50%	8	60	+
aTIGITxaPD 1-LCC	192	10.3	1.25	1.35	0.48	78%	8	60	+

Based on *in vitro* results and developability assessment, HCN version of the BsAbs was selected for *in vivo* efficacy study using h-TIGIT and h-PD1 double knock-in C57BL/6 mice with MC38 tumor model. MC38 tumor cells were implanted one week before treatment. Treatment started when tumor reached about 43 mm³ volume, where drugs were dosed intraperitoneally twice a week for 3 weeks with dosing volume shown in Figure 22. On Day 18 of the study, the tumor volume (TV, mean $\pm$ SEM) of animals in the control group was 1706.15 $\pm$ 135.86 mm³. Although anti-TIGIT-Fc monotherapy did not show significant anti-tumor efficacy (TGI: 8.40%, P=0.6282) compared with the vehicle control due to low dose volume, all other 3 treatment groups showed significant anti-tumor efficacy resulting in %TGI of 52.56, 61.14, 66.60 in groups treated with anti-PD1 monotherapy, anti-PD1 + anti-TIGIT-Fc combination, and BsAb HCN version, respectively, on Day 18 with TV (mean $\pm$ SEM) of 831.83 $\pm$ 103.75 mm³ (P=0.0012), 689.30 $\pm$ 128.93 mm³ (P=0.0012), and 598.53 $\pm$ 95.42 mm³ (P=0.0006), respectively (Mann-Whitney test). With BsAb HCN, the endpoint tumor weight was also significantly reduced, not only over the vehicle control (P=0.0006, Mann-Whitney test) but also over the monotherapy to anti-PD1 (P<0.03, Mann-Whitney test). Body weight changes were not significant between the groups throughout the study, indicating the treatments were well tolerated.

Table 5. Mean Tumor Volumes on Day 18 in MC38 Syngeneic B-hPD-1/hTIGIT Mouse Model

Groups	Treatment	N	TV (mm ³) ^a	P ^b	P ^c	P ^d	% TGI (Day 18)
1	Vehicle	7	1706.15 $\pm$ 135.86	-	0.0012	0.6282	-
2	aPD1 (0.3 mg/kg)	7	831.86 $\pm$ 103.75	0.0012	-	0.0047	52.56

3	aTIGIT-Fc (1 mg/kg)	6	1566.45 ± 192.41	0.6282	0.0047	-	8.40
4	aPD1+aTIGIT-Fc	7	689.30 ± 128.93	0.0012	0.3829	0.0047	61.14
5	BsAb_HCN	7	598.53 ± 95.42	0.0006	0.2086	0.0012	66.60

^a Mean ± SEM

^b Compared to vehicle control by Mann-Whitney test on day 18 post-treatment

^c Compared to aPD1 by Mann-Whitney test on day 18 post-treatment

^d Compared to aTIGIT-Fc by Mann-Whitney test on day 18 post-treatment

5

Table 6. Mean Tumor Weight of Treatment groups in MC38 Syngeneic B-hPD-1/h-TIGIT Mouse Model

Gps	Treatment	N	TW (g) ^a	% TWI	<i>P</i> ^b	<i>P</i> ^c	<i>P</i> ^d
1	Vehicle	7	2.701 ± 0.129	-	-	0.0006	0.4452
2	aPD1	7	1.322 ± 0.121	51.05	0.0006	-	0.0012
3	aTIGIT-Fc	6	2.465 ± 0.349	8.73	0.4452	0.0012	-
4	aPD1 + aTIGIT-Fc	7	1.128 ± 0.157	58.22	0.0006	0.3829	0.0012
5	BsAb_HCN	7	0.880 ± 0.141	67.42	0.0006	0.0262	0.0012

^a Mean ± SEM

^b Compared to vehicle control by Mann-Whitney test

^c Compared to aPD1 by Mann-Whitney test

^d Compared to 2A3-Fc by Mann-Whitney test

10

15 TIGIT is an immune checkpoint inhibitor similar to PD1 and CTLA4 but has a different expression pattern. It is known that PD1 is expressed at high levels in CD8⁺ T cells but at low levels in Natural Killer (NK) cells and regulatory T cells (Tregs). In contrast, TIGIT is expressed at high levels in CD8⁺ T cells, NK cells and Tregs. Both PD1 and TIGIT are up-regulated significantly in tumor infiltrating lymphocytes in tumor tissues, suggesting that both immune checkpoint inhibitors

contribute to immune suppression for tumor immune escape. Bispecific anti-TIGIT/anti-PD1 antibodies can provide superior therapeutic benefits via mechanisms not available to the monotherapy or the combinatory therapy. For example, as illustrated in Figure 23A, anti-TIGIT/anti-PD1 BsAbs can block inhibitory signal in PD1 & TIGIT double positive and single positive CD8⁺ T cells. Moreover, the BsAbs can block inhibitory signals triggered from PDL1 & PVR double positive tumor cells and PVR single positive tumor cells.

Furthermore, as illustrated in Figure 23B, anti-TIGIT/anti-PD1 BsAbs can block inhibitory signal in NK cells and block stimulatory signal in Tregs. PVR expressed on tumor cells or antigen presenting cells (APC) can also bind to TIGIT on NK cells, which induces inhibitory signal and suppresses NK cells function. Blocking binding of PVR to TIGIT by the BsAbs can reduce the suppressive effect, leading to enhanced NK cell anti-tumor function. PVR on tumor cells and APC can also bind to TIGIT on Treg cells, which induces stimulatory signal and enhances Tregs function, leading to suppression of anti-tumor activity. Blocking TIGIT on Tregs by BsAb can reduce Tregs function and enhance anti-tumor activity.

In addition to the various embodiments depicted and claimed, the disclosed subject matter is also directed to other embodiments having other combinations of the features disclosed and claimed herein. As such, the particular features presented herein can be combined with each other in other manners within the scope of the disclosed subject matter such that the disclosed subject matter includes any suitable combination of the features disclosed herein. The foregoing description of specific embodiments of the disclosed subject matter has been presented for purposes of illustration and description. It is not intended to be exhaustive or to limit the disclosed subject matter to those embodiments disclosed.

It will be apparent to those skilled in the art that various modifications and variations can be made in the compositions and methods of the disclosed subject matter without departing from the spirit or scope of the disclosed subject matter. Thus, it is intended that the disclosed subject matter include modifications and variations that are within the scope of the appended claims and their equivalents.

Various publications, patents and patent applications are cited herein, the contents of which are hereby incorporated by reference in their entireties.

WHAT IS CLAIMED IS:

1. A multispecific antibody that binds to TIGIT and PD1, comprising:
 - i) a first antigen-binding moiety comprising an anti-TIGIT antibody comprising a single domain antibody that binds to TIGIT; and
 - 5 ii) a second antigen-binding moiety comprising an anti-PD1 antibody that binds to PD1.
2. The multispecific antibody of claim 1, wherein the single domain antibody comprises a VHH.
3. The multispecific antibody of claim 1 or 2, wherein the single domain antibody or the VHH comprises a heavy chain variable region (VH).
4. The multispecific antibody of any one of claims 1-3, wherein the single domain antibody
- 10 binds to TIGIT with a K_D of 1×10^{-7} M or less.
5. The multispecific antibody of any one of claims 1-4, wherein the single domain antibody binds to TIGIT with a K_D of 1×10^{-8} M or less.
6. The multispecific antibody of any one of claims 1-5, wherein the single domain antibody binds to TIGIT with a K_D of 3×10^{-9} M or less.
- 15 7. The multispecific antibody of any one of claims 1-6, wherein the single domain antibody binds to TIGIT with a K_D of 2×10^{-9} M or less.
8. The multispecific antibody of any one of claims 1-7, wherein the single domain antibody cross-competes for binding to TIGIT with a reference single domain antibody comprising a heavy chain variable region comprising:
 - 20 a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96,
 - b) a heavy chain variable region CDR1 comprising amino acids having the sequence set
 - 25 forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100,
 - c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids
 - 30 having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104,
 - d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids
 - 35 having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108,

- e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112,
- 5 f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116,
- 10 g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120,
- h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124,
- 15 i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128,
- 20 j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132,
- 25 k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136,
- l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140,
- 30 m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144,
- 35

- n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148,
- 5 o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 151, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 152,
- 10 p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 154, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 155, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 156,
- 15 q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160,
- 20 r) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164,
- 25 s) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168,
- 30 t) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172,
- u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176,
- 35 v) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180,

w) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184,

5 x) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188, or

10 y) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

9. The multispecific antibody of any one of claims 1-8, wherein the single domain antibody comprises a heavy chain variable region comprising:

15 a) a heavy chain variable region CDR1 comprising an amino acid sequence of any one of SEQ ID NOs: 94, 98, 102, 106, 110, 114, 118, 122, 126, 130, 134, 138, 142, 146, 150, 154, 158, 162, 166, 170, 174, 178, 182, 186 and 190, or a variant thereof comprising up to about 3 amino acid substitutions;

20 b) a heavy chain variable region CDR2 comprising an amino acid sequence of any one of SEQ ID NOs: 95, 99, 103, 107, 111, 115, 119, 123, 127, 131, 135, 139, 143, 147, 151, 155, 159, 163, 167, 171, 175, 179, 183, 187 and 191, or a variant thereof comprising up to about 3 amino acid substitutions; and

25 c) a heavy chain variable region CDR3 comprising an amino acid sequence of any one of SEQ ID NOs: 96, 100, 104, 108, 112, 116, 120, 124, 128, 132, 136, 140, 144, 148, 152, 156, 160, 164, 168, 172, 176, 180, 184, 188 and 192, or a variant thereof comprising up to about 3 amino acid substitutions.

10. The multispecific antibody of any one of claims 1-9, wherein the single domain antibody comprises a heavy chain variable region that comprises a CDR1 domain, a CDR2 domain and a CDR3 domain, wherein the CDR1 domain, the CDR2 domain and the CDR3 domain respectively
30 comprise a CDR1 domain, a CDR2 domain and a CDR3 domain comprised in a reference heavy chain variable region comprising the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

11. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody
35 comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the

sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96.

12. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 98, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 99, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 100.

13. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 102, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 103, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 104.

14. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 106, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 107, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 108.

15. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 110, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 111, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 112.

16. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 114, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 115, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 116.

17. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 118, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 119, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 120.

18. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 122, a heavy chain variable region CDR2 comprising amino acids having the

sequence set forth in SEQ ID NO: 123, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 124.

19. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 126, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 127, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 128.

20. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 130, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 131, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 132.

21. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 134, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 135, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 136.

22. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 138, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 139, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 140.

23. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 142, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 143, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 144.

24. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 146, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 147, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 148.

25. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 150, a heavy chain variable region CDR2 comprising amino acids having the

sequence set forth in SEQ ID NO: 151, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 152.

26. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 154, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 155, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 156.

27. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 158, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 159, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 160.

28. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 162, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 163, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 164.

29. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 166, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 167, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 168.

30. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 170, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 171, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 172.

31. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 174, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 175, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 176.

32. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the

sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180.

33. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 182, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 183, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 184.

34. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188.

35. The multispecific antibody of any one of claims 1-10, wherein the single domain antibody comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192.

36. The multispecific antibody of any one of claims 1-35, wherein the single domain antibody comprises a heavy chain variable region comprising an amino acid sequence having at least about 90% sequence identity to the amino acid sequence selected from the group consisting of SEQ ID NOs: 97, 101, 105, 109, 113, 117, 121, 125, 129, 133, 137, 141, 145, 149, 153, 157, 161, 165, 169, 173, 177, 181, 185, 189 and 193.

37. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 97.

38. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 101.

39. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 105.

40. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 109.

41. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:113.
42. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody
5 comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:117.
43. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:121.
- 10 44. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:125.
45. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID
15 NO:129.
46. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:133.
47. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody
20 comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:137.
48. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:141.
- 25 49. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:145.
50. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID
30 NO:149.
51. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:153.
52. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody
35 comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:157.

53. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:161.

54. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:165.

55. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:169.

56. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:173.

57. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:177.

58. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:181.

59. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:185.

60. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:189.

61. The multispecific antibody of any one of claims 1-36, wherein the single domain antibody comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO:193.

62. The multispecific antibody of any one of claims 1-61, wherein the single domain antibody cross-competes for binding to TIGIT with a reference single domain antibody comprising a heavy chain variable region comprising:

(a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3,

(b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having

the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7,

(c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11,

(d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 15,

(e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 17, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 18, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 19,

(f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 21, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 22, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 23,

(g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 25, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 26, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 27,

(h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 29, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 30, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 31,

(i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 33, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 34, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 35,

(j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 37, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 38, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 39,

(k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 41, a heavy chain variable region CDR2 comprising amino acids

having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79, and

(u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

63. The multispecific antibody of any one of claims 1-62, wherein the single domain antibody comprises a heavy chain variable region comprising:

(a) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 1, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 2, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 3,

(b) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 5, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 6, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 7,

(c) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 9, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 10, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 11,

(d) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 13, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 14, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 15,

(e) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 17, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 18, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 19,

(f) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 21, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 22, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 23,

(g) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 25, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 26, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 27,

- (h) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 29, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 30, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 31,
- 5 (i) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 33, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 34, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 35,
- 10 (j) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 37, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 38, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 39,
- 15 (k) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 41, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 42, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 43,
- 20 (l) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 45, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 46, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 47,
- 25 (m) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 49, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 50, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 51,
- 30 (n) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 53, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 54, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 55,
- 35 (o) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 57, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 58, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 59,
- (p) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 61, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 62, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 63,

(q) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 65, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 66, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 67,

(r) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 69, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 70, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 71,

(s) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 73, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 74, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 75,

(t) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 77, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 78, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 79, and

(u) a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 81, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 82, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 83.

64. The multispecific antibody of any one of claims 1-63, wherein the single domain antibody comprises a heavy chain variable region having at least about 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 8, 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80 and 84.

65. The multispecific antibody of any one of claims 1-64, wherein the single domain antibody comprises a humanized framework.

66. The multispecific antibody of any one of claims 1-65, wherein the second antigen-binding moiety comprises an anti-PD1 antibody that cross-competes with a reference anti-PD1 antibody comprising:

a) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226);

b) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO:231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234);

c) a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO:239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO:240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241), and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO:242); or

d) a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the amino acid sequence SSYYYGIDL (SEQ ID NO:247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO:248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

67. The multispecific antibody of any one of claims 1-66, wherein the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 221), (2) a CDR- H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 222), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDF (SEQ ID NO: 223); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTTAVA (SEQ ID NO: 224), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 225), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 226).

68. The multispecific antibody of any one of claims 1-66, wherein the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FRFSNYGMS (SEQ ID NO: 229), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNAY (SEQ ID NO: 230), and (3) a CDR-H3 comprising the amino acid sequence TSYYYGIDF (SEQ ID NO:231); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASTDVTTAVA (SEQ

ID NO: 232), (2) a CDR-L2 comprising the amino acid sequence WASLRHT (SEQ ID NO: 233), and (3) a CDR-L3 comprising the amino acid sequence QQHYGIPWT (SEQ ID NO: 234).

69. The multispecific antibody of any one of claims 1-66, wherein the second antigen-binding moiety comprises a heavy chain variable (VH) domain sequence comprising (1) a CDR-H1

5 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 237), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 238), and (3) a CDR-H3 comprising the amino acid sequence VSYYYGIDL (SEQ ID NO: 239); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KAKQDVTTAVA (SEQ ID NO: 240), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 241),
10 and (3) a CDR-L3 comprising the amino acid sequence QQHYWIPWT (SEQ ID NO: 242).

70. The multispecific antibody of any one of claims 1-66, wherein the second antigen-binding moiety comprises a heavy chain variable domain (VH) sequence comprising (1) a CDR-H1 comprising the amino acid sequence FTFSNYGMS (SEQ ID NO: 245), (2) a CDR-H2 comprising the amino acid sequence TISGGGSNIY (SEQ ID NO: 246), and (3) a CDR-H3 comprising the
15 amino acid sequence SSYYGIDL (SEQ ID NO: 247); and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence KASQDVTNAVA (SEQ ID NO: 248), (2) a CDR-L2 comprising the amino acid sequence WASTRHT (SEQ ID NO: 249), and (3) a CDR-L3 comprising the amino acid sequence QQHYTIPWT (SEQ ID NO: 250).

71. The multispecific antibody of any one of claims 1-70, wherein the anti-PD1 antibody
20 comprises a humanized antibody.

72. The multispecific antibody of any one of claims 1-71, wherein the second antigen binding moiety comprises an anti-PD1 antibody comprising two antibody heavy chains and two antibody light chains.

73. The multispecific antibody of any one of claims 1-72, wherein the first antigen-binding
25 moiety comprises two or more anti-TIGIT antibodies.

74. The multispecific antibody of any one of claims 1-73, wherein the first antigen-binding moiety comprises two anti-TIGIT antibodies.

75. The multispecific antibody of any one of claims 1-74, wherein the C-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding
30 moiety.

76. The multispecific antibody of claim 75, wherein the C-terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

77. The multispecific antibody of any one of claims 1-76, wherein the N-terminus of at least one of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding
35 moiety.

78. The multispecific antibody of claim 77, wherein the N-terminus of each of the two anti-PD1 light chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

79. The multispecific antibody of any one of claims 1-78, wherein the C-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

80. The multispecific antibody of claim 79, wherein the C-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

81. The multispecific antibody of any one of claims 1-80, wherein the N-terminus of at least one of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

82. The multispecific antibody of claim 81, wherein the N-terminus of each of the two anti-PD1 heavy chains is linked to an anti-TIGIT antibody of the first antigen binding moiety.

83. The multispecific antibody of any one of claims 1-82, wherein the first antigen binding moiety is linked to the second antigen binding moiety via a linker.

84. The multispecific antibody of claim 83, wherein the linker is a peptide linker.

85. The multispecific antibody of claim 84, wherein the peptide linker comprises about four to about thirty amino acids.

86. The multispecific antibody of claim 84 or 85, wherein the peptide linker comprises an amino acid sequence selected from the group consisting of SEQ ID NOs: 195-220.

87. The multispecific antibody of any one of claims 1-86, wherein the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc regions of IgG, IgA, IgD, IgE and IgM.

88. The multispecific antibody of any one of claims 1-87, wherein the anti-PD1 antibody of the second antigen-binding moiety comprises an Fc region selected from the group consisting of the Fc region of IgG1, IgG2, IgG3 and IgG4.

89. The multispecific antibody of claim 87 or 88, wherein the Fc region comprises a human Fc region.

90. The multispecific antibody of any one of claims 87-89, wherein the Fc region comprises an IgG1 Fc region.

91. The multispecific antibody of any one of claims 87-89, wherein the Fc region comprises an IgG4 Fc region.

92. The multispecific antibody of claim 91, wherein the IgG4 Fc region comprises an S228P mutation.

93. The multispecific antibody of any one of claims 1-92, wherein the multispecific antibody is a bispecific antibody.

94. The multispecific antibody of any one of claims 1-93, comprising:

i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 94, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 95, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 96; and

ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 229, (2) a CDR-H2 comprising the amino acid sequence set forth in SEQ ID NO: 230, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 231; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 232, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 233, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 234.

95. The multispecific antibody of any one of claims 1-93, comprising:

i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 178, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 179, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 180; and

ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 237, (2) a CDR-H2 comprising the amino acid sequence set forth in SEQ ID NO: 238, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 239; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 240, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 241, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 242.

96. The multispecific antibody of any one of claims 1-93, comprising:

i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 186, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 187, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 188; and

ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid

sequence set forth in SEQ ID NO: 221, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 222, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 223; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 224, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 225, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 226.

97. The multispecific antibody of any one of claims 1-93, comprising:

i) a first antigen-binding moiety comprising a single domain anti-TIGIT antibody that comprises a heavy chain variable region CDR1 comprising amino acids having the sequence set forth in SEQ ID NO: 190, a heavy chain variable region CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 191, and a heavy chain variable region CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 192; and

ii) a second antigen-binding moiety comprising an anti-PD1 antibody comprising a heavy chain variable domain (VH) sequence that comprises (1) a CDR-H1 comprising the amino acid sequence set forth in SEQ ID NO: 245, (2) a CDR- H2 comprising the amino acid sequence set forth in SEQ ID NO: 246, and (3) a CDR-H3 comprising the amino acid sequence set forth in SEQ ID NO: 247; and a light chain variable domain (VL) sequence comprising (1) a CDR-L1 comprising the amino acid sequence set forth in SEQ ID NO: 248, (2) a CDR-L2 comprising the amino acid sequence set forth in SEQ ID NO: 249, and (3) a CDR-L3 comprising the amino acid sequence set forth in SEQ ID NO: 250.

98. The multispecific antibody of any one of claims 1-97, comprising an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 253, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 254.

99. The multispecific antibody of any one of claims 1-97, comprising an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 255, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 256.

100. The multispecific antibody of any one of claims 1-97, comprising an anti-PD1 antibody heavy chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 257, and an anti-PD1 antibody light chain comprising the amino acid sequence set forth in SEQ ID NO: 258.

101. The multispecific antibody of any one of claims 1-97, comprising an anti-PD1 antibody heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 259, and an anti-PD1 antibody light chain linked to an anti-TIGIT antibody comprising the amino acid sequence set forth in SEQ ID NO: 260.

102. An immunoconjugate comprising the multispecific antibody of any one of claims 1-101, linked to a therapeutic agent.
103. The immunoconjugate of claim 102, wherein the therapeutic agent is a cytotoxin.
104. The immunoconjugate of claim 102, wherein the therapeutic agent is a radioactive isotope.
- 5 105. A pharmaceutical composition comprising a) the multispecific antibody of any one of claims 1-101 or the immunoconjugate of any one of claims 102-104, and b) a pharmaceutically acceptable carrier.
106. A nucleic acid encoding the multispecific antibody of any one of claims 1-101.
107. A vector comprising the nucleic acid of claim 106.
- 10 108. A host cell comprising the nucleic acid of claim 106 or the vector of claim 107.
109. A method for preparing a multispecific antibody of any one of claims 1-101 comprising expressing the multispecific antibody in the host cell of claim 108 and isolating the multispecific antibody from the host cell.
110. A method of reducing tumor burden in a subject, the method comprising administering to
15 the subject an effective amount of a multispecific antibody of any one of claims 1-101, an immunoconjugate of any one of claims 102-104, or a pharmaceutical composition of claim 105.
111. The method of claim 110, wherein the method reduces the number of tumor cells.
112. The method of claim 110 or 111, wherein the method reduces tumor size.
113. The method of any one of claims 110-112, wherein the method eradicates the tumor in the
20 subject.
114. The method of any one of claims 110-113, wherein the tumor is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer,
25 blood cancer and a combination thereof.
115. A method of treating and/or preventing a neoplasm, the method comprising administering to the subject an effective amount of a multispecific antibody of any one of claims 1-101, an immunoconjugate of any one of claims 102-104, or a pharmaceutical composition of claim 105.
116. A method of lengthening survival of a subject having a neoplasm, the method comprising
30 administering to the subject an effective amount of a multispecific antibody of any one of claims 1-101, an immunoconjugate of any one of claims 102-104, or a pharmaceutical composition of claim 105.
117. The method of claim 115 or 116, wherein the neoplasm is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon
35 cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic

carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer, blood cancer and a combination thereof.

118. An multispecific antibody of any one of claims 1-101 for use as a medicament.

119. An multispecific antibody of any one of claims 1-101 for use in treating cancer.

5 120. A pharmaceutical composition of claim 105 for use as a medicament.

121. A pharmaceutical composition of claim 105 for use in treating cancer.

122. The multispecific antibody of claim 119 or the pharmaceutical composition of claim 121, wherein the cancer is selected from the group consisting of mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, 10 gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, cholangiocarcinoma, head and neck cancer, blood cancer and a combination thereof.

123. A kit comprising an multispecific antibody of any one of claims 1-101, an immunoconjugate of any one of claims 102-104, a pharmaceutical composition of claim 105, a nucleic acid of claim 106, a vector of claim 107 or a host cell of claim 108.

15 124. The kit of claim 123, further comprising a written instruction for treating and/or preventing a neoplasm.

FIGURE 1A




FIGURE 1B




FIGURE 1C




FIGURE 2A




FIGURE 2B




FIGURE 3A




FIGURE 3B




FIGURE 4A




FIGURE 4B




FIGURE 5




FIGURE 6A




FIGURE 6B




FIGURE 7A




FIGURE 7B




FIGURE 7C




FIGURE 8A




FIGURE 8B




FIGURE 8C




FIGURE 9




FIGURE 10A




FIGURE 10B




FIGURE 11






FIGURE 12

h-FcγRIIIA binding



h-FcγRIIB binding



m-FcγRIV binding




FIGURE 13A




FIGURE 13B




FIGURE 14A




FIGURE 14B




FIGURE 14C




FIGURE 15




FIGURE 16




FIGURE 17




FIGURE 18






FIGURE 19



EC ₅₀ , nM	aPD1	aTIGITxaPD1_HCN	aTIGITxaPD1_HCC	aTIGITxaPD1_LCN	aTIGITxaPD1_LCC
Average (n=4)	0.36	1.56	1.40	1.35	0.78
SD	0.18	0.97	1.26	0.63	0.41

FIGURE 20



EC ₅₀ (nM)	aPD1	aTIGITxaPD1_HCN	aTIGITxaPD1_HCC	aTIGITxaPD1_LCN	aTIGITxaPD1_LCC
Average (n=2-3)	0.73	1.52	3.29	4.64	1.35
SD	0.58	0.49	1.54	1.69	0.26

FIGURE 21




FIGURE 22A




FIGURE 22B




FIGURE 23A





FIGURE 23B



序列表

<110> SHANGHAI HENLIUS BIOTECH, INC.

<120> ANTI-TIGIT ANTIBODIES, MULTISPECIFIC ANTIBODIES COMPRISING THE SAME AND METHODS OF USING THE SAME

<130> SH1903-20P450257

<150> PCT/CN2020/071336

<151> 2020-01-10

<160> 262

<170> SIPOSequenceListing 1.0

<210> 1

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 4A11 CDR1

<400> 1

Gly Arg Pro Phe Ser Asn Tyr Thr

1 5

<210> 2

<211> 7

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(7)

<223> 4A11 CDR2

<400> 2

Ala Trp Pro Ser Pro Ser Thr

1 5

<210> 3

<211> 18

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(18)

<223> 4A11 CDR3

<400> 3

Ala Ala Asp Tyr Lys Ser Leu Thr Gln Ser Trp Leu Asn Ala Ala Leu
1 5 10 15
Asp Tyr

<210> 4

<211> 124

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(124)

<223> 4A11 VHH

<400> 4

Gln Val Lys Leu Glu Glu Ser Gly Gly Gly Glu Ala Gln Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Thr Ala Ser Gly Arg Pro Phe Ser Asn Tyr
20 25 30
Thr Met Gly Trp Phe Arg Arg Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45
Gly Leu Ala Trp Pro Ser Pro Ser Thr Tyr Val Val Asp Ser Val Lys
50 55 60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Ile Tyr Leu
65 70 75 80
Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Ile Tyr Tyr Cys Ala
85 90 95
Ala Asp Tyr Lys Ser Leu Thr Gln Ser Trp Leu Asn Ala Ala Leu Asp
100 105 110
Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 5

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 4B5 CDR1

<400> 5

Pro Arg Thr Phe Ser Thr Phe His
1 5

<210> 6

<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4B5 CDR2

<400> 6
Phe Asn Trp Ser Gly Gly Arg Thr
1 5

<210> 7
<211> 26
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(26)
<223> 4B5 CDR3

<400> 7
Ala Ala Ala Arg Asp Arg Gly Leu His Asp Gly Thr Thr Ser Asp Ser
1 5 10 15
Tyr Leu Glu Gly Ser His Glu Tyr Glu Tyr
20 25

<210> 8
<211> 133
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(133)
<223> 4B5 VHH

<400> 8
Gln Val Lys Leu Glu Glu Ser Gly Gly Arg Ser Val Leu Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Arg Cys Ala Gly Thr Pro Arg Thr Phe Ser Thr Phe
20 25 30
His Ile Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45
Ala Ala Phe Asn Trp Ser Gly Gly Arg Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asn Asn Gly Lys Asn Met Val Tyr
65 70 75 80
Leu Gln Met Thr Ser Leu Thr Pro Glu Asp Thr Gly Leu Tyr Tyr Cys

				85					90					95			
Ala	Ala	Ala	Arg	Asp	Arg	Gly	Leu	His	Asp	Gly	Thr	Thr	Ser	Asp	Ser		
			100					105					110				
Tyr	Leu	Glu	Gly	Ser	His	Glu	Tyr	Glu	Tyr	Trp	Gly	Gln	Gly	Thr	Gln		
		115					120					125					
Val	Thr	Val	Ser	Ser													
	130																

<210> 9
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 4C5 CDR1

<400> 9
 Gly Arg Ser Val Ser Thr Tyr Phe
 1 5

<210> 10
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 4C5 CDR2

<400> 10
 Ile Asp Arg Gly Ser Thr Val Thr
 1 5

<210> 11
 <211> 17
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(17)
 <223> 4C5 CDR3

<400> 11
 Ala Ala Lys Ala Ile Thr Arg Asn Phe Ile Ala Thr Asn Asp Tyr Asp
 1 5 10 15
 Tyr

<210> 12
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 4C5 VHH

<400> 12
Gln Val Gln Leu Val Asp Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Val Ser Gly Arg Ser Val Ser Thr Tyr
20 25 30
Phe Val Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45
Ala Ala Ile Asp Arg Gly Ser Thr Val Thr Arg Tyr Asp Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asp Thr Val Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Lys Ala Ile Thr Arg Asn Phe Ile Ala Thr Asn Asp Tyr Asp
100 105 110
Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 13
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4D5 CDR1

<400> 13
Gly Arg Ala Phe Asn Glu Tyr Ala
1 5

<210> 14
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4D5 CDR2

<400> 14
Ile Ser Ser Asp Gly Arg Phe Thr
1 5

<210> 15
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 4D5 CDR3

<400> 15
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 16
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 4D5 VHH

<400> 16
Val Asp Ser Gly Gly Gly Ala Val Lys Ala Gly Asp Ser Leu Arg Leu
1 5 10 15
Val Cys Ser Ala Pro Gly Arg Thr His Gly Arg Ala Phe Asn Glu Tyr
20 25 30
Ala Met Ala Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125

<210> 17
<211> 8

<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4D11 CDR1

<400> 17
Gly Ser Ile Ser Ser Ile Asn Ala
1 5

<210> 18
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> 4D11 CDR2

<400> 18
Ile Thr Asn Ser Gly Ser Thr
1 5

<210> 19
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> 4D11 CDR3

<400> 19
Thr Ala Arg Arg Ser Thr Trp Tyr Ile Ser
1 5 10

<210> 20
<211> 116
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(116)
<223> 4D11 VHH

<400> 20
 Gln Val Gln Leu Gln Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Ser Ile Ser Ser Ile Asn
 20 25 30
 Ala Met Gly Trp Tyr Arg Leu Ala Pro Gly Lys His Arg Glu Phe Val
 35 40 45
 Ala Asp Ile Thr Asn Ser Gly Ser Thr Asn Tyr Ala Ala Ser Val Lys
 50 55 60
 Gly Arg Phe Asn Ile Ser Arg Asp Asn Ala Lys Asp Thr Val Tyr Leu
 65 70 75 80
 Gln Met Asn Ser Leu Lys Phe Glu Asp Thr Ala Val Tyr Tyr Cys Thr
 85 90 95
 Ala Arg Arg Ser Thr Trp Tyr Ile Ser Ser Gly Arg Gly Thr Gln Val
 100 105 110
 Thr Val Ser Ser
 115

<210> 21
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 4E5 CDR1

<400> 21
 Gly Leu Thr Ser Ser Asp Ile Ala
 1 5

<210> 22
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 4E5 CDR2

<400> 22
 Ile Ser Ser Asp Gly Arg Phe Thr
 1 5

<210> 23
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN
<222> (1)..(18)
<223> 4E5 CDR3

<400> 23
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 24
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 4E5 VHH

<400> 24
Gln Val Gln Leu Val Asp Ser Gly Gly Gly Val Val Glu Val Gly Ala
1 5 10 15
Ser Leu Thr Leu Ser Cys Glu Thr Ser Gly Leu Thr Ser Ser Asp Ile
20 25 30
Ala Val Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125

<210> 25
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4H6 CDR1

<400> 25
Gly Thr Ile Phe Arg Leu Asn Arg
1 5

<210> 26
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4H6 CDR2

<400> 26
Thr Ile Trp Ser Gly Arg Arg Thr
1 5

<210> 27
<211> 14
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(14)
<223> 4H6 CDR3

<400> 27
Asn Tyr Arg Arg Ile Thr Pro Trp Glu Ala Ser Gly Asn Tyr
1 5 10

<210> 28
<211> 121
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(121)
<223> 4H6 VHH

<400> 28
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ala Thr Ala Gly Ala
1 5 10 15
Ser Leu Ile Leu Ser Cys Ala Ala Ser Gly Thr Ile Phe Arg Leu Asn
20 25 30
Arg Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Arg Val
35 40 45
Ala Ala Thr Ile Trp Ser Gly Arg Arg Thr His Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Thr Asp Asn Ala Lys Lys Thr Val Tyr
65 70 75 80
Leu Arg Met Ser Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys

				85					90					95				
Asn	Tyr	Arg	Arg	Ile	Thr	Pro	Trp	Glu	Ala	Ser	Gly	Asn	Tyr	Trp	Gly			
			100					105					110					
Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser										
		115					120											

<210> 29
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 4H9 CDR1

<400> 29
 Gly Pro Ile Ala Arg Ser Arg Ser
 1 5

<210> 30
 <211> 10
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(10)
 <223> 4H9 CDR2

<400> 30
 Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr
 1 5 10

<210> 31
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 4H9 CDR3

<400> 31
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
 1 5 10 15
 Asp Tyr

<210> 32

<211> 127
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(127)
<223> 4H9 VHH

<400> 32
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Ala Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Thr Ala Ser Gly Pro Ile Ala Arg Ser Arg
20 25 30
Ser Thr Gly Met Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu
35 40 45
Ser Val Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala
50 55 60
Ser Val Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala
65 70 75 80
Ala Phe Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr
85 90 95
Tyr Cys Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln
100 105 110
Trp Tyr Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125

<210> 33
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 10H5 CDR1

<400> 33
Glu Thr Thr Phe Lys Ser Met Ala
1 5

<210> 34
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 10H5 CDR2

<400> 34
Thr Asn Tyr Asn Gly Gly Arg Thr
1 5

<210> 35
<211> 17
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(17)
<223> 10H5 CDR3

<400> 35
Ala Ala Lys Ala Thr Glu Gly Thr Thr Phe Pro Ser Arg Thr Tyr Glu
1 5 10 15
Phe

<210> 36
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 10H5 VHH

<400> 36
Gly Gly Gly Leu Val Gln Ala Gly Gly Ser Leu Arg Leu Ala Cys Thr
1 5 10 15
Ala Ser Asp Pro Pro Phe Ala Asn Tyr Glu Thr Thr Phe Lys Ser Met
20 25 30
Ala Met Gly Trp Val Arg His Ile Pro Gly Lys Glu Arg Glu Leu Val
35 40 45
Ala Ala Thr Asn Tyr Asn Gly Gly Arg Thr Trp Tyr Ser Asn Ser Ala
50 55 60
Lys Ala Arg Ser Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Val Tyr
65 70 75 80
Leu Gln Met Ser Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Lys Ala Thr Glu Gly Thr Thr Phe Pro Ser Arg Thr Tyr Glu
100 105 110
Phe Trp Gly Gln Gly Ile Gln Val Thr Val Ser Ser
115 120

<210> 37
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 12H7 CDR1

<400> 37
Gly Asn Phe Leu Ser Val Ser Asp
1 5

<210> 38
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> 12H7 CDR2

<400> 38
Val Thr Glu His Gly Arg Thr
1 5

<210> 39
<211> 20
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(20)
<223> 12H7 CDR3

<400> 39
Lys Ala Ser Asp Val Phe Thr Asp Ala Gly Ala His Glu Ala Val Leu
1 5 10 15
Ile Arg Asp Tyr
20

<210> 40
<211> 126
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(126)
<223> 12H7 VHH

<400> 40
 Gln Val Gln Leu Val Asp Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Lys Val Ser Gly Asn Phe Leu Ser Val Ser
 20 25 30
 Asp Met Ser Trp Tyr Arg Gln Ala Pro Gly Met Glu Arg Asp Val Val
 35 40 45
 Ala Thr Val Thr Glu His Gly Arg Thr Thr Tyr Thr Asp Ser Val Lys
 50 55 60
 Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Glu His Thr Thr Tyr Leu
 65 70 75 80
 Glu Met Asn Asn Leu Lys Pro Glu Asp Thr Ala Val Tyr Phe Cys Lys
 85 90 95
 Ala Ser Asp Val Phe Thr Asp Ala Gly Ala His Glu Ala Val Leu Ile
 100 105 110
 Arg Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
 115 120 125

<210> 41
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 13H11 CDR1

<400> 41
 Gly Leu Thr Phe Ser Met Tyr Ala
 1 5

<210> 42
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 13H11 CDR2

<400> 42
 Ile Ser Ser Asp Gly Arg Phe Thr
 1 5

<210> 43
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN
<222> (1)..(18)
<223> 13H11 CDR3

<400> 43
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 44
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 13H11 VHH

<400> 44
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Leu Thr Phe Ser Met Tyr
20 25 30
Ala Met Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125

<210> 45
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 15A5 CDR1

<400> 45
Glu Arg Thr Phe Ser Ser Phe Ala
1 5

<210> 46
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 15A5 CDR2

<400> 46
Ile Asp Pro Ser Gly Arg Tyr Ile
1 5

<210> 47
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 15A5 CDR3

<400> 47
Ala Ala Arg Ile Arg Gly Glu Gly Tyr Tyr Thr Arg Ser Ser Phe Tyr
1 5 10 15
His Tyr

<210> 48
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 15A5 VHH

<400> 48
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Glu Arg Thr Phe Ser Ser Phe
20 25 30
Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Val Val
35 40 45
Ala Ala Ile Asp Pro Ser Gly Arg Tyr Ile Tyr Tyr Lys Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Met Ser Arg Asp Asn Ala Lys Ser Thr Val Tyr

65					70					75				80
Leu	Gln	Met	Asn	Ser	Leu	Lys	Pro	Asp	Asp	Thr	Ala	Arg	Tyr	Tyr
				85					90					95
Ala	Ala	Arg	Ile	Arg	Gly	Glu	Gly	Tyr	Tyr	Thr	Arg	Ser	Ser	Phe
			100					105					110	
His	Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser		
		115					120					125		

<210> 49
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2B7 CDR1

<400> 49
 Gly Arg Thr Phe Ser Ser Tyr Pro
 1 5

<210> 50
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2B7 CDR2

<400> 50
 Ile Ser Ser Asp Gly Arg Phe Thr
 1 5

<210> 51
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 2B7 CDR3

<400> 51
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
 1 5 10 15
 Asp Tyr

<210> 52
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2B7 VHH

<400> 52
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ala Cys Ala Ala Ser Gly Arg Thr Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120 125

<210> 53
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2B10 CDR1

<400> 53
Ser Arg Ile Phe Arg Arg Tyr Ala
1 5

<210> 54
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2B10 CDR2

<400> 54
Ile Thr Trp Ser Gly Ala Ser Thr
1 5

<210> 55
<211> 17
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(17)
<223> 2B10 CDR3

<400> 55
Ala Ala Asp Pro Trp Gly Ser Val Ile Val Gly Thr Ala Glu Tyr Glu
1 5 10 15
Tyr

<210> 56
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 2B10 VHH

<400> 56
Gln Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Thr Gly Asp
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Arg Ile Phe Arg Arg Tyr
20 25 30
Ala Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45
Ala Ala Ile Thr Trp Ser Gly Ala Ser Thr Thr Tyr Thr Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Ser Ala Glu Asn Thr Thr Tyr
65 70 75 80
Leu Gln Met Thr Ser Leu Arg Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Asp Pro Trp Gly Ser Val Ile Val Gly Thr Ala Glu Tyr Glu
100 105 110
Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 57
<211> 8

<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 3F10 CDR1

<400> 57
Glu His Thr Phe Ser Asn Phe Pro
1 5

<210> 58
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 3F10 CDR2

<400> 58
Ile Asp Ser Ser Gly Arg Leu Thr
1 5

<210> 59
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 3F10 CDR3

<400> 59
Ala Ala Arg Thr Gly Gly Val Gly Tyr Tyr Ser Arg Ser Ser Phe Tyr
1 5 10 15
Asn Tyr

<210> 60
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 3F10 VHH

<400> 60
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ser Ser Glu His Thr Phe Ser Asn Phe
 20 25 30
 Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Asn Val Val
 35 40 45
 Ala Ala Ile Asp Ser Ser Gly Arg Leu Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Thr Val Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Lys Ser Glu Asp Thr Ala Arg Tyr Tyr Cys
 85 90 95
 Ala Ala Arg Thr Gly Gly Val Gly Tyr Tyr Ser Arg Ser Ser Phe Tyr
 100 105 110
 Asn Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120 125

<210> 61
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 3G6 CDR1

<400> 61
 Gly Ser Ile Phe Gly Ile Ser Val
 1 5

<210> 62
 <211> 7
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(7)
 <223> 3G6 CDR2

<400> 62
 Leu Thr Arg Ala Gly Leu Thr
 1 5

<210> 63
 <211> 15
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(15)
<223> 3G6 CDR3

<400> 63
His Ala Asn Ile Met Glu Ser Ala Ala Ser Thr Phe Gly Arg Tyr
1 5 10 15

<210> 64
<211> 121
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(121)
<223> 3G6 VHH

<400> 64
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Ser Leu Ser Cys Ala Ala Ser Gly Ser Ile Phe Gly Ile Ser
20 25 30
Val Met Gly Trp Tyr Arg Gln Ala Pro Gly Glu Gln Arg Asp Leu Val
35 40 45
Ala Thr Leu Thr Arg Ala Gly Leu Thr Thr Tyr Gly Asp Ser Val Lys
50 55 60
Gly Arg Phe Ser Ile Ser Arg Asp Ser Ala Lys Asn Thr Val Tyr Leu
65 70 75 80
Gln Met Asn Asn Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys His
85 90 95
Ala Asn Ile Met Glu Ser Ala Ala Ser Thr Phe Gly Arg Tyr Trp Gly
100 105 110
Gln Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 65
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 3G7 CDR1

<400> 65
Gly Arg Thr Leu Ser Thr Tyr Thr
1 5

<210> 66
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> 3G7 CDR2

<400> 66
Ala Trp Pro Ser Pro Ser Thr
1 5

<210> 67
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 3G7 CDR3

<400> 67
Ala Ala Asp Tyr Lys Ser Leu Thr Gln Ser Trp Leu Asn Ala Ala Leu
1 5 10 15
Asp Tyr

<210> 68
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 3G7 VHH

<400> 68
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Asp
1 5 10 15
Ser Leu Arg Leu Ser Cys Glu Ala Ser Gly Arg Thr Leu Ser Thr Tyr
20 25 30
Thr Met Gly Trp Phe Arg Arg Ala Pro Gly Lys Glu Arg Glu Phe Val
35 40 45
Gly Leu Ala Trp Pro Ser Pro Ser Thr Tyr Val Val Asp Ser Val Lys
50 55 60
Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Ile Tyr Leu

65					70					75				80	
Gln	Met	Asn	Ser	Leu	Lys	Pro	Glu	Asp	Thr	Ala	Ile	Tyr	Tyr	Cys	Ala
				85					90					95	
Ala	Asp	Tyr	Lys	Ser	Leu	Thr	Gln	Ser	Trp	Leu	Asn	Ala	Ala	Leu	Asp
			100					105					110		
Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser				
		115					120								

<210> 69
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 3H7 CDR1

<400> 69
 Gly Ser Ile Leu Ser Ala Gly Val
 1 5

<210> 70
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 3H7 CDR2

<400> 70
 Ile Ala Leu Asp Gly Ser Thr Gly
 1 5

<210> 71
 <211> 15
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(15)
 <223> 3H7 CDR3

<400> 71
 Asn Ala Asn Ile Arg Thr Asp Met Arg Ser Ala Pro Phe Asp His
 1 5 10 15

<210> 72

<211> 122
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(122)
<223> 3H7 VHH

<400> 72
Gln Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Ser Ile Leu Ser Ala Gly
20 25 30
Val Met Arg Trp Tyr Arg Gln Ala Pro Gly Lys Gln Arg Glu Leu Val
35 40 45
Ala Ser Ile Ala Leu Asp Gly Ser Thr Gly Tyr Tyr Ile Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ile Val Tyr
65 70 75 80
Leu Asp Met Arg Ser Leu Glu Pro Ala Asp Thr Ala Val Tyr Leu Cys
85 90 95
Asn Ala Asn Ile Arg Thr Asp Met Arg Ser Ala Pro Phe Asp His Trp
100 105 110
Gly His Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 73
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4C6 CDR1

<400> 73
Gly Arg Thr Phe Ser Ser Tyr Pro
1 5

<210> 74
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 4C6 CDR2

<400> 74
Ile Ser Ser Asp Gly Arg Phe Thr
1 5

<210> 75
<211> 16
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(16)
<223> 4C6 CDR3

<400> 75
Ala Val Asp Pro Thr Gly Trp Gly Thr Ile Glu Ala Asp Phe Arg Ser
1 5 10 15

<210> 76
<211> 123
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(123)
<223> 4C6 VHH

<400> 76
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Gly
1 5 10 15
Ser Leu Arg Leu Ala Cys Ala Ala Ser Gly Arg Thr Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Arg Cys
85 90 95
Ala Val Asp Pro Thr Gly Trp Gly Thr Ile Glu Ala Asp Phe Arg Ser
100 105 110
Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 77
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN
<222> (1)..(8)
<223> 1C12 CDR1

<400> 77
Ser Arg Ile Phe Ser Arg Tyr Gly
1 5

<210> 78
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 1C12 CDR2

<400> 78
Ile Ser Trp Asn Gly Ala Ser Thr
1 5

<210> 79
<211> 17
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(17)
<223> 1C12 CDR3

<400> 79
Ala Ala Asp Pro Trp Gly Ala Val Lys Leu Gly Thr Ala Glu Tyr Glu
1 5 10 15
Tyr

<210> 80
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 1C12 VHH

<400> 80
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Thr Gly Asp

1				5					10					15			
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Ser	Arg	Ile	Phe	Ser	Arg	Tyr		
			20					25					30				
Gly	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Phe	Val		
		35					40					45					
Ala	Ala	Ile	Ser	Trp	Asn	Gly	Ala	Ser	Thr	Thr	Tyr	Thr	Asp	Ser	Val		
	50					55					60						
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Ser	Ala	Glu	Asn	Thr	Thr	Tyr		
65					70				75					80			
Leu	Gln	Ile	Asn	Ser	Leu	Arg	Pro	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys		
			85					90					95				
Ala	Ala	Asp	Pro	Trp	Gly	Ala	Val	Lys	Leu	Gly	Thr	Ala	Glu	Tyr	Glu		
		100					105						110				
Tyr	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser						
		115				120											

<210> 81
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 1G1 CDR1

<400> 81
 Gly Pro Ser Phe Ser Ser Tyr Pro
 1 5

<210> 82
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 1G1 CDR2

<400> 82
 Ile Ser Ser Asp Gly Arg Phe Thr
 1 5

<210> 83
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)

<223> 1G1 CDR3

<400> 83

Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 84

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 1G1 VHH

<400> 84

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Ala Gly Asp
1 5 10 15
Ser Leu Arg Leu Ser Cys Val Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Gly Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Ala Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ala Lys Ser Ala Ala Phe
65 70 75 80
Leu Gln Met Asn Ser Leu Lys Pro Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 85

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 1G1-F-G-ERES VHH

<400> 85

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30

Pro	Met	Gly	Trp	Phe	Arg	Gln	Gly	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
		50				55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75					80
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 86
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)
 <223> 1G1-F-A-ERES VHH

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1			5					10					15		
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr
		20					25					30			
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
		50				55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75					80
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 87
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)
 <223> 1G1-F-A-EREW VHH

<400> 87

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr
			20					25					30		
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Trp	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90				95		
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 88
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)
 <223> 1G1-F-A-GLEW VHH

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr
			20					25					30		
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90				95		
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 89
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)

<223> 1G1-F-A-GREL VHH

<400> 89

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg	
1				5					10					15		
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr	
			20					25					30			
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Arg	Glu	Leu	Val	
		35					40					45				
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	
	50					55					60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	
65					70					75				80		
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85						90				95			
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr	
			100					105					110			
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser				
		115					120					125				

<210> 90

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 1G1-F-A-GRES VHH

<400> 90

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg	
1				5					10					15		
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr	
			20					25					30			
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Arg	Glu	Ser	Val	
		35					40					45				
Ala	Ala	Ile	Ser	Ser	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	
	50					55					60					
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	
65					70					75				80		
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85						90				95			
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr	
			100					105					110			
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser				
		115					120					125				

<210> 91

<211> 124

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(124)
 <223> 1C12-EREF VHH

<400> 91
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Ser Arg Ile Phe Ser Arg Tyr
 20 25 30
 Gly Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Phe Val
 35 40 45
 Ala Ala Ile Ser Trp Asn Gly Ala Ser Thr Thr Tyr Thr Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ala Asp Pro Trp Gly Ala Val Lys Leu Gly Thr Ala Glu Tyr Glu
 100 105 110
 Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
 115 120

<210> 92
 <211> 124
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(124)
 <223> 1C12-EREW VHH

<400> 92
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Ser Arg Ile Phe Ser Arg Tyr
 20 25 30
 Gly Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Trp Val
 35 40 45
 Ala Ala Ile Ser Trp Asn Gly Ala Ser Thr Thr Tyr Thr Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ala Asp Pro Trp Gly Ala Val Lys Leu Gly Thr Ala Glu Tyr Glu
 100 105 110
 Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
 115 120

<210> 93
<211> 124
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(124)
<223> 1C12_GLEW VHH

<400> 93
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Ser Arg Ile Phe Ser Arg Tyr
20 25 30
Gly Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ala Ala Ile Ser Trp Asn Gly Ala Ser Thr Thr Tyr Thr Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Asp Pro Trp Gly Ala Val Lys Leu Gly Thr Ala Glu Tyr Glu
100 105 110
Tyr Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
115 120

<210> 94
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 CDR1

<400> 94
Gly Gly Ser Phe Ser Ser Tyr Pro
1 5

<210> 95
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 CDR2

<400> 95
Ile Ser Ser Asp Met Arg Phe Thr
1 5

<210> 96
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2A3 CDR3

<400> 96
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 97
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A3 VHH

<400> 97
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Gly Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Met Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 98
<211> 8
<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 1A8 CDR1

<400> 98

Gly Pro Ser Phe Ser Ser Tyr Pro

1 5

<210> 99

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 1A8 CDR2

<400> 99

Ile Ser Ser Arg Gly Arg Phe Thr

1 5

<210> 100

<211> 18

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(18)

<223> 1A8 CDR3

<400> 100

Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr

1 5 10 15

Asp Tyr

<210> 101

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 1A8 VHH

<400> 101
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
 20 25 30
 Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
 35 40 45
 Ala Ala Ile Ser Ser Arg Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
 100 105 110
 Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120 125

<210> 102
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 1D11 CDR1

<400> 102
 Gly Pro Ser Phe Ser Ser Ser Pro
 1 5

<210> 103
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 1D11 CDR2

<400> 103
 Ile Ser Ser Met Gly Arg Phe Thr
 1 5

<210> 104
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 1D11 CDR3

<400> 104
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 105
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 1D11 VHH

<400> 105
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Ser
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Met Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 106
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 5E8 CDR1

<400> 106
Gly Pro Ser Phe Ser Ser Tyr Pro

1 5

<210> 107
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 5E8 CDR2

<400> 107
Gln Ser Ser Asp Gly Arg Phe Thr
1 5

<210> 108
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 5E8 CDR3

<400> 108
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 109
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 5E8 VHH

<400> 109
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Gln Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60

Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 110
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2A5 CDR1

<400> 110
 Gly Pro Ser Phe Ser Ser Tyr Pro
 1 5

<210> 111
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2A5 CDR2

<400> 111
 Ile Ser Ser Val Gly Arg Phe Thr
 1 5

<210> 112
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 2A5 CDR3

<400> 112
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Trp Gln Trp Tyr
 1 5 10 15
 Asp Tyr

<210> 113
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A5 VHH

<400> 113
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Val Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Trp Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 114
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 5G10 CDR1

<400> 114
Gly Pro Arg Phe Ser Ser Tyr Pro
1 5

<210> 115
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)

<223> 5G10 CDR2

<400> 115

Ile Ser Ser Asp Gly Arg Phe Thr
1 5

<210> 116

<211> 18

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(18)

<223> 5G10 CDR3

<400> 116

Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Gly

<210> 117

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 5G10 VHH

<400> 117

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Arg Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Gly Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 118

<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A6 CDR1

<400> 118
Gly Pro Ser Phe Ser Leu Tyr Pro
1 5

<210> 119
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A6 CDR2

<400> 119
Ile Ser Ser Asp Arg Arg Phe Thr
1 5

<210> 120
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2A6 CDR3

<400> 120
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 121
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)

<223> 2A6 VHH

<400> 121

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Leu	Tyr
			20					25					30		
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Arg	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75				80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90				95		
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 122

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 1G1-1C12 CDR1

<400> 122

Gly	Pro	Ser	Phe	Ser	Ser	Tyr	Pro
1				5			

<210> 123

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 1G1-1C12 CDR2

<400> 123

Ile	Ser	Ser	Asp	Leu	Arg	Phe	Thr
1				5			

<210> 124

<211> 18

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(18)

<223> 1G1-1C12 CDR3

<400> 124

Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Lys	Gln	Trp	Tyr
1				5				10						15	
Asp	Tyr														

<210> 125

<211> 125

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(125)

<223> 1G1-1C12 VHH

<400> 125

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5				10						15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Tyr
			20					25					30		
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Leu	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75				80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Lys	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 126

<211> 8

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(8)

<223> 5B5 CDR1

<400> 126
Gly Pro Ser Phe Ser Ser Tyr Pro
1 5

<210> 127
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 5B5 CDR2

<400> 127
Ile Ser Ser Asp Thr Arg Phe Thr
1 5

<210> 128
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 5B5 CDR3

<400> 128
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Arg

<210> 129
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 5B5 VHH

<400> 129
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45

Ala	Ala	Ile	Ser	Ser	Asp	Thr	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75						80
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Arg	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 130
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 6F12 CDR1

<400> 130
 Gly Pro Ser Phe Thr Ser Tyr Pro
 1 5

<210> 131
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 6F12 CDR2

<400> 131
 Ile Ser Ser Asp Gly Arg Phe Lys
 1 5

<210> 132
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 6F12 CDR3

<400> 132
 Ala Ala Glu Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr

<221> DOMAIN
<222> (1)..(8)
<223> 1C7 CDR2

<400> 135
Ile Ser Ser Arg Gly Arg Phe Thr
1 5

<210> 136
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 1C7 CDR3

<400> 136
Ala Ala Arg Gly Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 137
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 1C7 VHH

<400> 137
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Arg Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Gly Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 138
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 6E9 CDR1

<400> 138
Gly Pro Ser Arg Ser Ser Tyr Pro
1 5

<210> 139
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 6E9 CDR2

<400> 139
Ile Ser Ser Asp Gly Lys Phe Thr
1 5

<210> 140
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 6E9 CDR3

<400> 140
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Asn Trp Tyr
1 5 10 15
Asp Tyr

<210> 141
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN
<222> (1)..(125)
<223> 6E9 VHH

<400> 141
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Arg Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Lys Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Asn Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 142
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 1A10 CDR1

<400> 142
Gly Asn Ser Phe Ser Ser Tyr Pro
1 5

<210> 143
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 1A10 CDR2

<400> 143
Ile Ser Ser Asp Gly Arg Phe Ser
1 5

<210> 144

<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 1A10 CDR3

<400> 144
Ala Cys Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 145
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 1A10 VHH

<400> 145
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Asn Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Ser Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Cys Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 146
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 6B11 CDR1

<400> 146
Gly Pro Ser Phe Pro Ser Tyr Pro
1 5

<210> 147
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 6B11 CDR2

<400> 147
Ile Ser Ser Arg Gly Arg Phe Thr
1 5

<210> 148
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 6B11 CDR3

<400> 148
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Leu Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 149
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 6B11 VHH

<400> 149
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Pro Ser Tyr
20 25 30

Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Arg	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70					75				80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Leu	Gln	Trp	Tyr
		100					105						110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 150
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 6D8 CDR1

<400> 150
 Gly Pro Ser Phe Ser Ser Lys Pro
 1 5

<210> 151
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 6D8 CDR2

<400> 151
 Arg Ser Ser Asp Gly Arg Phe Thr
 1 5

<210> 152
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 6D8 CDR3

<400> 152
 Ala Ala Arg Asp Ser Gly Ser Gly Arg Tyr Ser Arg Ala Gln Trp Tyr
 1 5 10 15
 Asp Tyr

<210> 153
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)
 <223> 6D8 VHH

<400> 153
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Ser Lys
 20 25 30
 Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
 35 40 45
 Ala Ala Arg Ser Ser Asp Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ala Arg Asp Ser Gly Ser Gly Arg Tyr Ser Arg Ala Gln Trp Tyr
 100 105 110
 Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120 125

<210> 154
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 6D12 CDR1

<400> 154
 Gly Pro Ser Phe Ser Thr Tyr Pro
 1 5

<210> 155
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 6D12 CDR2

<400> 155
Ile Ser Ser Asp Gly Val Phe Thr
1 5

<210> 156
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 6D12 CDR3

<400> 156
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Glu Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 157
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 6D12 VHH

<400> 157
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Thr Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Val Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Glu Gln Trp Tyr
100 105 110

Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 158
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2C5 CDR1

<400> 158
Gly Pro Ser Phe Ser Thr Tyr Pro
1 5

<210> 159
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2C5 CDR2

<400> 159
Ile Ser Ser Gln Gly Arg Phe Thr
1 5

<210> 160
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2C5 CDR3

<400> 160
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 161
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2C5 VHH

<400> 161
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Ser Thr Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Gln Gly Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 162
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 7F6 CDR1

<400> 162
Gly Pro Met Phe Ser Ser Tyr Pro
1 5

<210> 163
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 7F6 CDR2

<400> 163
Ile Ser Ser Asp Pro Arg Phe Thr
1 5

<210> 164
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 7F6 CDR3

<400> 164
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 165
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 7F6 VHH

<400> 165
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Met Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Pro Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 166
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN

<222> (1)..(8)
<223> 2D5 CDR1

<400> 166
Gly Pro Ser Phe Ser Ser Ser Pro
1 5

<210> 167
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2D5 CDR2

<400> 167
Ile Ser Trp Asp Gly Arg Phe Thr
1 5

<210> 168
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2D5 CDR3

<400> 168
Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Val Tyr

<210> 169
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2D5 VHH

<400> 169
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Pro	Ser	Phe	Ser	Ser	Ser
			20					25				30			
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Trp	Asp	Gly	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85					90					95		
Ala	Ala	Arg	Asp	Ser	Gly	Ser	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
		100					105					110			
Val	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 170
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 7B11 CDR1

<400> 170
 Gly Pro Ser Phe Leu Ile Tyr Pro
 1 5

<210> 171
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 7B11 CDR2

<400> 171
 Ile Ser Ser Asp Gly Arg Phe Trp
 1 5

<210> 172
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(18)
 <223> 7B11 CDR3

<400> 172
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Val Gln Trp Tyr
 1 5 10 15
 Asp Tyr

<210> 173
 <211> 125
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(125)
 <223> 7B11 VHH

<400> 173
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Leu Ile Tyr
 20 25 30
 Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
 35 40 45
 Ala Ala Ile Ser Ser Asp Gly Arg Phe Trp Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Ala Arg Asp Ser Gly Ser Gly Tyr Tyr Ser Arg Val Gln Trp Tyr
 100 105 110
 Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120 125

<210> 174
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 7D12 CDR1

<400> 174
 Gly Pro Ser Phe Leu Ser Tyr Pro
 1 5

<210> 175
 <211> 8

<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 7D12 CDR2

<400> 175
Ile Ser Ser Asp Gly Arg Phe Ser
1 5

<210> 176
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 7D12 CDR3

<400> 176
Ala Ala Arg Asp Trp Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 177
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 7D12 VHH

<400> 177
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Pro Ser Phe Leu Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Gly Arg Phe Ser Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Ala Arg Asp Trp Gly Ser Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 178
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 ML CDR1

<400> 178
Gly Gly Ser Phe Ser Ser Tyr Pro
1 5

<210> 179
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 ML CDR2

<400> 179
Ile Ser Ser Asp Leu Arg Phe Thr
1 5

<210> 180
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2A3 ML CDR3

<400> 180
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 181
<211> 125

<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A3 ML VHH

<400> 181
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Gly Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Leu Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 182
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 MI CDR1

<400> 182
Gly Gly Ser Phe Ser Ser Tyr Pro
1 5

<210> 183
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 MI CDR2

<400> 183

Ile Ser Ser Asp Ile Arg Phe Thr
1 5

<210> 184
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2A3 MI CDR3

<400> 184
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 185
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A3 MI VHH

<400> 185
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Gly Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Ile Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Asp Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 186
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 ML DT CDR1

<400> 186
Gly Gly Ser Phe Ser Ser Tyr Pro
1 5

<210> 187
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> 2A3 ML DT CDR2

<400> 187
Ile Ser Ser Asp Leu Arg Phe Thr
1 5

<210> 188
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> 2A3 ML DT CDR3

<400> 188
Ala Ala Arg Thr Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 189
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A3 ML DT VHH

<400> 189

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Gly	Ser	Phe	Ser	Ser	Tyr
			20					25					30		
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val
		35					40					45			
Ala	Ala	Ile	Ser	Ser	Asp	Leu	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90				95		
Ala	Ala	Arg	Thr	Ser	Gly	Val	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
			100					105					110		
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser			
		115					120					125			

<210> 190
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2A3 ML DE CDR1

<400> 190
 Gly Gly Ser Phe Ser Ser Tyr Pro
 1 5

<210> 191
 <211> 8
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(8)
 <223> 2A3 ML DE CDR2

<400> 191
 Ile Ser Ser Asp Leu Arg Phe Thr
 1 5

<210> 192
 <211> 18
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN

<222> (1)..(18)
<223> 2A3 ML DE CDR3

<400> 192
Ala Ala Arg Glu Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
1 5 10 15
Asp Tyr

<210> 193
<211> 125
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(125)
<223> 2A3 ML DE VHH

<400> 193
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Gly Ser Phe Ser Ser Tyr
20 25 30
Pro Met Gly Trp Phe Arg Gln Ala Pro Gly Lys Glu Arg Glu Ser Val
35 40 45
Ala Ala Ile Ser Ser Asp Leu Arg Phe Thr Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Ala Arg Glu Ser Gly Val Gly Tyr Tyr Ser Arg Ala Gln Trp Tyr
100 105 110
Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120 125

<210> 194
<211> 357
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(357)
<223> 2A3 LTFc

<400> 194
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Gly Ser Phe Ser Ser Tyr

<400> 195
Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly
1 5 10

<210> 196
<211> 15
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(15)
<223> Exemplary linker196

<400> 196
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10 15

<210> 197
<211> 5
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(5)
<223> Exemplary linker197

<400> 197
Gly Gly Gly Ser Gly
1 5

<210> 198
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> Exemplary linker198

<400> 198
Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10

<210> 199
<211> 8
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(8)
<223> Exemplary linker199

<400> 199
Gly Gly Ser Gly Gly Gly Ser Gly
1 5

<210> 200
<211> 12
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(12)
<223> Exemplary linker200

<400> 200
Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly
1 5 10

<210> 201
<211> 6
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(6)
<223> Exemplary linker201

<400> 201
Gly Ser Gly Gly Ser Gly
1 5

<210> 202
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> Exemplary linker202

<400> 202
Gly Ser Gly Gly Ser Gly Gly Ser Gly

1

5

<210> 203
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> Exemplary linker203

<400> 203
Gly Ser Gly Ser Gly Ser Gly
1 5

<210> 204
<211> 18
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(18)
<223> Exemplary linker204

<400> 204
Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15
Gly Gly

<210> 205
<211> 5
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(5)
<223> Exemplary linker205

<400> 205
Pro Ala Pro Ala Pro
1 5

<210> 206
<211> 15
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(15)
<223> Exemplary linker206

<400> 206
Pro Ala Pro Ala Pro Pro Ala Pro Ala Pro Pro Ala Pro Ala Pro
1 5 10 15

<210> 207
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> Exemplary linker207

<400> 207
Ile Lys Arg Thr Val Ala Ala
1 5

<210> 208
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> Exemplary linker208

<400> 208
Val Ser Ser Ala Ser Thr Lys
1 5

<210> 209
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> Exemplary linker209

<400> 209
Ala Glu Ala Ala Ala Lys Ala
1 5

<210> 210
<211> 12
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(12)
<223> Exemplary linker210

<400> 210
Ala Glu Ala Ala Ala Lys Glu Ala Ala Ala Lys Ala
1 5 10

<210> 211
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> Exemplary linker211

<400> 211
Gly Arg Pro Gly Ser Gly Arg Pro Gly Ser
1 5 10

<210> 212
<211> 20
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(20)
<223> Exemplary linker212

<400> 212
Gly Arg Pro Gly Ser Gly Arg Pro Gly Ser Gly Arg Pro Gly Ser Gly
1 5 10 15
Arg Pro Gly Ser
20

<210> 213
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN
<222> (1)..(10)
<223> Exemplary linker213

<400> 213
Gly Arg Gly Gly Ser Gly Arg Gly Gly Ser
1 5 10

<210> 214
<211> 20
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(20)
<223> Exemplary linker214

<400> 214
Gly Arg Gly Gly Ser Gly Arg Gly Gly Ser Gly Arg Gly Gly Ser Gly
1 5 10 15
Arg Gly Gly Ser
20

<210> 215
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> Exemplary linker215

<400> 215
Gly Lys Pro Gly Ser Gly Lys Pro Gly Ser
1 5 10

<210> 216
<211> 20
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(20)
<223> Exemplary linker216

<400> 216
Gly Lys Pro Gly Ser Gly Lys Pro Gly Ser Gly Lys Pro Gly Ser Gly

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(20)

<223> Exemplary linker220

<400> 220

Gly	Asp	Pro	Gly	Ser	Gly	Asp	Pro	Gly	Ser	Gly	Asp	Pro	Gly	Ser	Gly
1				5				10					15		
Asp	Pro	Gly	Ser												
			20												

<210> 221

<211> 9

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(9)

<223> H1G4 CDR-H1

<400> 221

Phe	Thr	Phe	Ser	Asn	Tyr	Gly	Met	Ser
1			5					

<210> 222

<211> 10

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(10)

<223> H1G4 CDR-H2

<400> 222

Thr	Ile	Ser	Gly	Gly	Gly	Ser	Asn	Ile	Tyr
1			5					10	

<210> 223

<211> 9

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(9)

<223> H1G4 CDR-H3

<400> 223
Val Ser Tyr Tyr Tyr Gly Ile Asp Phe
1 5

<210> 224
<211> 11
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(11)
<223> H1G4 CDR-L1

<400> 224
Lys Ala Ser Gln Asp Val Thr Thr Ala Val Ala
1 5 10

<210> 225
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> H1G4 CDR-L2

<400> 225
Trp Ala Ser Thr Arg His Thr
1 5

<210> 226
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> H1G4 CDR-L3

<400> 226
Gln Gln His Tyr Thr Ile Pro Trp Thr
1 5

<210> 227
<211> 116
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(116)
 <223> H1G4 VH

<400> 227
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30
 Gly Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Thr Ile Ser Gly Gly Gly Ser Asn Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Val Ser Tyr Tyr Tyr Gly Ile Asp Phe Trp Gly Gln Gly Thr Ser Val
 100 105 110
 Thr Val Ser Ser
 115

<210> 228
 <211> 108
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(108)
 <223> H1G4 VL

<400> 228
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Thr Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Ile Pro Trp
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> 229
 <211> 9

<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 33B CDR-H1

<400> 229
Phe Arg Phe Ser Asn Tyr Gly Met Ser
1 5

<210> 230
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> 33B CDR-H2

<400> 230
Thr Ile Ser Gly Gly Gly Ser Asn Ala Tyr
1 5 10

<210> 231
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 33B CDR-H3

<400> 231
Thr Ser Tyr Tyr Gly Ile Asp Phe
1 5

<210> 232
<211> 11
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(11)
<223> 33B CDR-L1

<400> 232
Lys Ala Ser Thr Asp Val Thr Thr Ala Val Ala
1 5 10

<210> 233
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> 33B CDR-L2

<400> 233
Trp Ala Ser Leu Arg His Thr
1 5

<210> 234
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 33B CDR-L3

<400> 234
Gln Gln His Tyr Gly Ile Pro Trp Thr
1 5

<210> 235
<211> 116
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(116)
<223> 33B VH

<400> 235
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Arg Phe Ser Asn Tyr
20 25 30
Gly Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Thr Ile Ser Gly Gly Gly Ser Asn Ala Tyr Tyr Ala Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Thr Ser Tyr Tyr Tyr Gly Ile Asp Phe Trp Gly Gln Gly Thr Ser Val
 100 105 110
 Thr Val Ser Ser
 115

<210> 236
 <211> 108
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(108)
 <223> 33B VL

<400> 236
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Thr Asp Val Thr Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Trp Ala Ser Leu Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Gly Ile Pro Trp
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> 237
 <211> 9
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(9)
 <223> 66E CDR-H1

<400> 237
 Phe Thr Phe Ser Asn Tyr Gly Met Ser
 1 5

<210> 238
 <211> 10
 <212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(10)

<223> 66E CDR-H2

<400> 238

Thr Ile Ser Gly Gly Gly Ser Asn Ile Tyr
1 5 10

<210> 239

<211> 9

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(9)

<223> 66E CDR-H3

<400> 239

Val Ser Tyr Tyr Tyr Gly Ile Asp Leu
1 5

<210> 240

<211> 11

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(11)

<223> 66E CDR-L1

<400> 240

Lys Ala Lys Gln Asp Val Thr Thr Ala Val Ala
1 5 10

<210> 241

<211> 7

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> DOMAIN

<222> (1)..(7)

<223> 66E CDR-L2

<400> 241

Trp Ala Ser Thr Arg His Thr
1 5

<210> 242
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 66E CDR-L3

<400> 242
Gln Gln His Tyr Trp Ile Pro Trp Thr
1 5

<210> 243
<211> 116
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(116)
<223> 66E VH

<400> 243
Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30
Gly Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45
Ser Thr Ile Ser Gly Gly Gly Ser Asn Ile Tyr Tyr Ala Asp Ser Val
50 55 60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65 70 75 80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Val Ser Tyr Tyr Tyr Gly Ile Asp Leu Trp Gly Gln Gly Thr Ser Val
100 105 110
Thr Val Ser Ser
115

<210> 244
<211> 108
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN

<222> (1)..(108)
<223> 66E VL

<400> 244
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15
Asp Arg Val Thr Ile Thr Cys Lys Ala Lys Gln Asp Val Thr Thr Ala
20 25 30
Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45
Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Trp Ile Pro Trp
85 90 95
Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
100 105

<210> 245
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 711D CDR-H1

<400> 245
Phe Thr Phe Ser Asn Tyr Gly Met Ser
1 5

<210> 246
<211> 10
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(10)
<223> 711D CDR-H2

<400> 246
Thr Ile Ser Gly Gly Gly Ser Asn Ile Tyr
1 5 10

<210> 247
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 711D CDR-H3

<400> 247
Ser Ser Tyr Tyr Tyr Gly Ile Asp Leu
1 5

<210> 248
<211> 11
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(11)
<223> 711D CDR-L1

<400> 248
Lys Ala Ser Gln Asp Val Thr Asn Ala Val Ala
1 5 10

<210> 249
<211> 7
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(7)
<223> 711D CDR-L2

<400> 249
Trp Ala Ser Thr Arg His Thr
1 5

<210> 250
<211> 9
<212> PRT
<213> 人工序列(Artificial Sequence)

<220>
<221> DOMAIN
<222> (1)..(9)
<223> 711D CDR-L3

<400> 250
Gln Gln His Tyr Thr Ile Pro Trp Thr

1

5

<210> 251
 <211> 116
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(116)
 <223> 711D VH

<400> 251
 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30
 Gly Met Ser Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Thr Ile Ser Gly Gly Gly Ser Asn Ile Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ser Ser Tyr Tyr Tyr Gly Ile Asp Leu Trp Gly Gln Gly Thr Ser Val
 100 105 110
 Thr Val Ser Ser
 115

<210> 252
 <211> 108
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(108)
 <223> 711D VL

<400> 252
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Thr Asn Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln His Tyr Thr Ile Pro Trp

				85					90				95
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg		
			100					105					

<210> 253
 <211> 580
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> CHAIN
 <222> (1)..(580)
 <223> anti-PD1 HC -Linker- 2A3 (HCC heavy chain)

<400> 253

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly
1			5					10						15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Tyr
			20					25					30		
Gly	Met	Ser	Trp	Ile	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
			35				40					45			
Ser	Thr	Ile	Ser	Gly	Gly	Gly	Ser	Asn	Ile	Tyr	Tyr	Ala	Asp	Ser	Val
	50					55					60				
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr
65					70				75					80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Val	Ser	Tyr	Tyr	Tyr	Gly	Ile	Asp	Phe	Trp	Gly	Gln	Gly	Thr	Ser	Val
			100					105					110		
Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala
			115				120					125			
Pro	Cys	Ser	Arg	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu
			130			135					140				
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly
145					150					155					160
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser
				165					170					175	
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu
			180					185					190		
Gly	Thr	Lys	Thr	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr
		195					200					205			
Lys	Val	Asp	Lys	Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro
		210				215					220				
Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
225					230					235					240
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
				245					250					255	
Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn
			260					265					270		
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
			275				280					285			
Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
			290			295					300				

Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser
 305 310 315 320
 Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys
 325 330 335
 Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu
 340 345 350
 Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
 355 360 365
 Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
 370 375 380
 Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
 385 390 395 400
 Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly
 405 410 415
 Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
 420 425 430
 Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys Gly Ser Gly Gly Ser
 435 440 445
 Gly Gly Ser Gly Gly Ser Gly Gln Val Gln Leu Val Glu Ser Gly Gly
 450 455 460
 Gly Val Val Gln Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser
 465 470 475 480
 Gly Gly Ser Phe Ser Tyr Pro Met Gly Trp Phe Arg Gln Ala Pro
 485 490 495
 Gly Lys Glu Arg Glu Ser Val Ala Ala Ile Ser Ser Asp Leu Arg Phe
 500 505 510
 Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp
 515 520 525
 Asn Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu
 530 535 540
 Asp Thr Ala Val Tyr Tyr Cys Ala Ala Arg Thr Ser Gly Val Gly Tyr
 545 550 555 560
 Tyr Ser Arg Ala Gln Trp Tyr Asp Tyr Trp Gly Gln Gly Thr Leu Val
 565 570 575
 Thr Val Ser Ser
 580

<210> 254
 <211> 214
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> CHAIN
 <222> (1)..(214)
 <223> anti-PD1 LC1 (HCC light chain)

<400> 254
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Thr Thr Ala
 20 25 30
 Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile

Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly
145					150					155					160
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser
				165					170						175
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu
			180						185						190
Gly	Thr	Lys	Thr	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr
		195					200						205		
Lys	Val	Asp	Lys	Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro
	210					215					220				
Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro
225					230					235					240
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr
				245						250					255
Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn
			260						265						270
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg
		275					280						285		
Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val
		290					295					300			
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser
305					310						315				320
Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys
				325						330					335
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu
			340						345					350	
Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe
		355					360					365			
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu
		370				375						380			
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe
385						390					395				400
Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly
				405						410					415
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr
			420					425						430	
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys					
		435					440								

<210> 256
 <211> 351
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> CHAIN
 <222> (1)..(351)
 <223> anti-PD1 LC-Linker-2A3 (LCC light chain)

<400> 256
 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val Thr Thr Ala

Gln 1	Val	Gln	Leu	Val 5	Glu	Ser	Gly	Gly	Gly 10	Val	Val	Gln	Pro	Gly 15	Arg
Ser	Leu	Arg	Leu 20	Ser	Cys	Ala	Ala	Ser 25	Gly	Gly	Ser	Phe	Ser 30	Ser	Tyr
Pro	Met	Gly 35	Trp	Phe	Arg	Gln	Ala 40	Pro	Gly	Lys	Glu	Arg 45	Glu	Ser	Val
Ala 50	Ala	Ile	Ser	Ser	Asp	Leu 55	Arg	Phe	Thr	Tyr	Tyr 60	Ala	Asp	Ser	Val
Lys 65	Gly	Arg	Phe	Thr	Ile 70	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu 80	Tyr
Leu	Gln	Met	Asn	Ser 85	Leu	Arg	Ala	Glu	Asp 90	Thr	Ala	Val	Tyr	Tyr 95	Cys
Ala	Ala	Arg	Thr 100	Ser	Gly	Val	Gly	Tyr 105	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr
Asp	Tyr	Trp	Gly 115	Gln	Gly	Thr	Leu 120	Val	Thr	Val	Ser	Ser	Gly	Ser	Gly
Gly	Ser	Gly 130	Gly	Ser	Gly	Gly 135	Ser	Gly	Gln	Val	Gln	Leu	Val	Glu	Ser
Gly 145	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala
Ala	Ser	Gly	Phe	Thr 165	Phe	Ser	Asn	Tyr	Gly 170	Met	Ser	Trp	Ile	Arg	Gln
Ala	Pro	Gly	Lys 180	Gly	Leu	Glu	Trp	Val	Ser	Thr	Ile	Ser	Gly	Gly	Gly
Ser	Asn	Ile	Tyr 195	Tyr	Ala	Asp	Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser
Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg
Ala 225	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Val	Ser	Tyr	Tyr	Tyr	Gly	Ile
Asp	Phe	Trp	Gly	Gln 245	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr
Lys	Gly	Pro	Ser 260	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	Ser	Thr	Ser
Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu
Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His
Thr 305	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser
Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr	Tyr	Thr	Cys
Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu
Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Phe	Leu
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
Met 385	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr

Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	
		435					440					445				
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	Ser	Ser	
	450					455					460					
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	
465					470					475					480	
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	
			485						490					495		
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	
			500				505					510				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	
	515					520					525					
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Arg	Leu	Thr	
	530					535					540					
Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	
545					550					555					560	
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	
				565					570						575	
Ser	Leu	Gly	Lys													
			580													

<210> 258
 <211> 214
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> CHAIN
 <222> (1)..(214)
 <223> anti-PD1 LC2 (HCN light chain)

<400>	258															
Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	
1				5				10						15		
Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Gln	Asp	Val	Thr	Thr	Ala	
		20					25					30				
Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	
	35					40					45					
Tyr	Trp	Ala	Ser	Thr	Arg	His	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
	50				55					60						
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
65					70					75					80	
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	His	Tyr	Thr	Ile	Pro	Trp	
			85					90					95			
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	
		100					105					110				
Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	
		115				120						125				
Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	
	130				135					140						
Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	
145					150					155					160	
Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	

				165					170					175			
Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr		
			180					185					190				
Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser		
		195					200					205					
Phe	Asn	Arg	Gly	Glu	Cys												
	210																

<210> 259

<211> 443

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> CHAIN

<222> (1)..(443)

<223> anti-PD1 HC2 (LCN heavy chain)

<400> 259

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Gly		
1			5					10						15			
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Asn	Tyr		
		20					25						30				
Gly	Met	Ser	Trp	Ile	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val		
	35					40						45					
Ser	Thr	Ile	Ser	Gly	Gly	Gly	Ser	Asn	Ile	Tyr	Tyr	Ala	Asp	Ser	Val		
	50				55					60							
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr		
65					70				75						80		
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys		
			85					90					95				
Val	Ser	Tyr	Tyr	Tyr	Gly	Ile	Asp	Phe	Trp	Gly	Gln	Gly	Thr	Ser	Val		
		100					105						110				
Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala		
	115					120						125					
Pro	Cys	Ser	Arg	Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu		
	130				135					140							
Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly		
145				150					155						160		
Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser		
			165				170						175				
Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu		
	180					185						190					
Gly	Thr	Lys	Thr	Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr		
	195				200							205					
Lys	Val	Asp	Lys	Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro		
	210				215						220						
Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro		
225				230					235						240		
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr		
			245					250					255				
Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn		
		260					265						270				

Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	
		275					280					285				
Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	
		290				295					300					
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	
305					310					315					320	
Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	
				325					330					335		
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	
			340				345						350			
Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	
		355				360						365				
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	
		370				375					380					
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	
385					390					395					400	
Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	
				405					410					415		
Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	
			420					425					430			
Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys						
		435				440										

<210> 260

<211> 351

<212> PRT

<213> 人工序列(Artificial Sequence)

<220>

<221> CHAIN

<222> (1)..(351)

<223> 2A3-Linker-anti-PD1 LC (LCN light chain)

<400> 260

Gln	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Val	Val	Gln	Pro	Gly	Arg	
1			5					10					15			
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Gly	Ser	Phe	Ser	Ser	Tyr	
		20					25					30				
Pro	Met	Gly	Trp	Phe	Arg	Gln	Ala	Pro	Gly	Lys	Glu	Arg	Glu	Ser	Val	
		35				40						45				
Ala	Ala	Ile	Ser	Ser	Asp	Leu	Arg	Phe	Thr	Tyr	Tyr	Ala	Asp	Ser	Val	
	50				55					60						
Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ser	Lys	Asn	Thr	Leu	Tyr	
65				70				75						80		
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85					90					95			
Ala	Ala	Arg	Thr	Ser	Gly	Val	Gly	Tyr	Tyr	Ser	Arg	Ala	Gln	Trp	Tyr	
		100					105						110			
Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser	Gly	Ser	Gly	
	115					120						125				
Gly	Ser	Gly	Gly	Ser	Gly	Gly	Ser	Gly	Asp	Ile	Gln	Met	Thr	Gln	Ser	
	130				135				140							
Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	

145					150				155				160		
Lys	Ala	Ser	Gln	Asp	Val	Thr	Thr	Ala	Val	Ala	Trp	Tyr	Gln	Gln	Lys
				165					170					175	
Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr	Trp	Ala	Ser	Thr	Arg	His
			180					185					190		
Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe
		195					200					205			
Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr
	210					215				220					
Cys	Gln	Gln	His	Tyr	Thr	Ile	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr	Lys
225				230						235					240
Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro
			245						250					255	
Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu
		260						265					270		
Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp
	275					280						285			
Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp
	290				295						300				
Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys
305				310						315					320
Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln
			325					330						335	
Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys	
		340					345						350		

<210> 261
 <211> 244
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> PEPTIDE
 <222> (1)..(244)
 <223> Human TIGIT polypeptide

<400> 261

Met	Arg	Trp	Cys	Leu	Leu	Ile	Trp	Ala	Gln	Gly	Leu	Arg	Gln	Ala	
1			5				10						15		
Pro	Leu	Ala	Ser	Gly	Met	Met	Thr	Gly	Thr	Ile	Glu	Thr	Thr	Gly	Asn
		20					25					30			
Ile	Ser	Ala	Glu	Lys	Gly	Gly	Ser	Ile	Ile	Leu	Gln	Cys	His	Leu	Ser
		35			40						45				
Ser	Thr	Thr	Ala	Gln	Val	Thr	Gln	Val	Asn	Trp	Glu	Gln	Gln	Asp	Gln
	50				55					60					
Leu	Leu	Ala	Ile	Cys	Asn	Ala	Asp	Leu	Gly	Trp	His	Ile	Ser	Pro	Ser
65				70					75						80
Phe	Lys	Asp	Arg	Val	Ala	Pro	Gly	Pro	Gly	Leu	Gly	Leu	Thr	Leu	Gln
			85				90						95		
Ser	Leu	Thr	Val	Asn	Asp	Thr	Gly	Glu	Tyr	Phe	Cys	Ile	Tyr	His	Thr
		100					105					110			
Tyr	Pro	Asp	Gly	Thr	Tyr	Thr	Gly	Arg	Ile	Phe	Leu	Glu	Val	Leu	Glu
		115					120					125			

Ser Ser Val Ala Glu His Gly Ala Arg Phe Gln Ile Pro Leu Leu Gly
 130 135 140
 Ala Met Ala Ala Thr Leu Val Val Ile Cys Thr Ala Val Ile Val Val
 145 150 155 160
 Val Ala Leu Thr Arg Lys Lys Lys Ala Leu Arg Ile His Ser Val Glu
 165 170 175
 Gly Asp Leu Arg Arg Lys Ser Ala Gly Gln Glu Glu Trp Ser Pro Ser
 180 185 190
 Ala Pro Ser Pro Pro Gly Ser Cys Val Gln Ala Glu Ala Ala Pro Ala
 195 200 205
 Gly Leu Cys Gly Glu Gln Arg Gly Glu Asp Cys Ala Glu Leu His Asp
 210 215 220
 Tyr Phe Asn Val Leu Ser Tyr Arg Ser Leu Gly Asn Cys Ser Phe Phe
 225 230 235 240
 Thr Glu Thr Gly

<210> 262
 <211> 138
 <212> PRT
 <213> 人工序列(Artificial Sequence)

<220>
 <221> DOMAIN
 <222> (1)..(138)
 <223> ECD of Human TIGIT polypeptide

<400> 262
 Met Arg Trp Cys Leu Leu Leu Ile Trp Ala Gln Gly Leu Arg Gln Ala
 1 5 10 15
 Pro Leu Ala Ser Gly Met Met Thr Gly Thr Ile Glu Thr Thr Gly Asn
 20 25 30
 Ile Ser Ala Glu Lys Gly Gly Ser Ile Ile Leu Gln Cys His Leu Ser
 35 40 45
 Ser Thr Thr Ala Gln Val Thr Gln Val Asn Trp Glu Gln Gln Asp Gln
 50 55 60
 Leu Leu Ala Ile Cys Asn Ala Asp Leu Gly Trp His Ile Ser Pro Ser
 65 70 75 80
 Phe Lys Asp Arg Val Ala Pro Gly Pro Gly Leu Gly Leu Thr Leu Gln
 85 90 95
 Ser Leu Thr Val Asn Asp Thr Gly Glu Tyr Phe Cys Ile Tyr His Thr
 100 105 110
 Tyr Pro Asp Gly Thr Tyr Thr Gly Arg Ile Phe Leu Glu Val Leu Glu
 115 120 125
 Ser Ser Val Ala Glu His Gly Ala Arg Phe
 130 135