



(86) Date de dépôt PCT/PCT Filing Date: 2011/01/14
(87) Date publication PCT/PCT Publication Date: 2011/07/21
(45) Date de délivrance/Issue Date: 2019/05/14
(85) Entrée phase nationale/National Entry: 2012/06/21
(86) N° demande PCT/PCT Application No.: US 2011/021279
(87) N° publication PCT/PCT Publication No.: 2011/088303
(30) Priorité/Priority: 2010/01/15 (US61/295,576)


(51) Cl.Int./Int.Cl. *C07D 409/12* (2006.01),
A61K 31/381 (2006.01), *A61K 31/4436* (2006.01),
A61P 31/00 (2006.01), *C07D 409/14* (2006.01),
C07D 493/04 (2006.01)

(72) Inventeurs/Inventors:
CANALES, EDA, US;
CHONG, LEE S., US;
CLARKE, MICHAEL O'NEIL HANRAHAN, US;
DOERFFLER, EDWARD, US;
LAZERWITH, SCOTT E., US;
LEW, WILLARD, US;

...

(73) Propriétaire/Owner:

(54) Titre : INHIBITEURS DE FLAVIVIRUS
(54) Title: INHIBITORS OF FLAVIVIRIDAE VIRUSES



Formula (I)

(57) Abrégé/Abstract:

Provided are compounds of Formula (I): and pharmaceutically acceptable salts and esters thereof. The compounds, compositions, and methods provided are useful for the treatment of Flaviviridae virus infections, particularly hepatitis C infections.

(72) Inventeurs(suite)/Inventors(continued): MERTZMAN, MICHAEL, US; MORGANELLI, PHILIP ANTHONY, US;
WATKINS, WILLIAM J., US

(73) Propriétaires(suite)/Owners(continued): GILEAD SCIENCES, INC., US

(74) Agent: ROBIC

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 July 2011 (21.07.2011)

PCT

(10) International Publication Number
WO 2011/088303 A1

(51) International Patent Classification:

C07D 409/12 (2006.01) *A61K 31/381* (2006.01)
C07D 409/14 (2006.01) *A61K 31/4436* (2006.01)
C07D 493/04 (2006.01) *A61P 31/00* (2006.01)

(21) International Application Number:

PCT/US2011/021279

(22) International Filing Date:

14 January 2011 (14.01.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/295,576 15 January 2010 (15.01.2010) US

(71) Applicant (for all designated States except US):
GILEAD SCIENCES, INC. [US/US]; 333 Lakeside
 Drive, Foster City, CA 94404 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CANALES, Eda**
 [US/US]; 419 Turner Terrace, Apt. 9, San Mateo, CA
 94401 (US). **CHONG, Lee, S.** [US/US]; 37469 Marsten
 Drive, Newark, CA 94560 (US). **CLARKE, Michael,**
O'neil Hanrahan [US/US]; 545 Seastorm Drive, Red-
 wood City, CA 94065 (US). **DOERFFLER, Edward**
 [US/US]; 2607 Royal Ann Drive, Union City, CA 94587
 (US). **LAZERWITH, Scott, E.** [US/US]; 930 Scott
 Street #1, San Francisco, CA 94115 (US). **LEW, Willard**
 [US/US]; 1437 Overland Drive, San Mateo, CA 94403
 (US). **MERTZMAN, Michael** [US/US]; 2603 Carlmont
 Drive, Belmont, CA 94002 (US). **MORGANELLI,**

Philip, Anthony [US/US]; 3923 Webster Street, Oakland,
 CA 94609 (US). **WATKINS, William, J.** [GB/US];
 14275 Hilltop Way, Saratoga, CA 95070 (US).

(74) Agents: **WARD, John, S.** et al.; Gilead Sciences, Inc.,
 333 Lakeside Drive, Foster City, CA 94404 (US).


(81) Designated States (unless otherwise indicated, for every
 kind of national protection available): AE, AG, AL, AM,
 AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
 CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
 DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
 HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
 KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
 ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
 NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
 SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
 TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
 kind of regional protection available): ARIPO (BW, GH,
 GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
 ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
 TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
 LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
 SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: INHIBITORS OF FLAVIVIRIDAE VIRUSES



Formula (I)

(57) Abstract: Provided are compounds of Formula (I): and pharmaceutically acceptable salts and esters thereof. The compounds, compositions, and methods provided are useful for the treatment of Flaviviridae virus infections, particularly hepatitis C infections.

WO 2011/088303 A1

INHIBITORS OF *FLAVIVIRIDAE* VIRUSES

FIELD OF THE INVENTION


5 The present application includes novel inhibitors of *Flaviviridae* viruses, compositions containing such compounds, therapeutic methods that include the administration of such compounds.

BACKGROUND OF THE INVENTION

10 Viruses comprising the *Flaviviridae* family include at least three distinguishable genera including *pestiviruses*, *flaviviruses*, and *hepaciviruses* (Calisher, *et al.*, J. Gen. Virol., 1993, 70, 37-43). While *pestiviruses* cause many economically important animal diseases such as bovine viral diarrhea virus (BVDV), classical swine fever virus (CSFV, hog cholera) and border disease of sheep (BDV), their importance in human disease is less well characterized (Moennig, V., *et al.*, Adv. Vir. Res. 1992, 48, 53-98). *Flaviviruses* are responsible
15 for important human diseases such as dengue fever and yellow fever while *hepaciviruses* cause hepatitis C virus infections in humans. Other important viral infections caused by the *Flaviviridae* family include West Nile virus (WNV) Japanese encephalitis virus (JEV), tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis, St Louis encephalitis, Omsk hemorrhagic fever virus
20 and Zika virus.

The hepatitis C virus (HCV) is the leading cause of chronic liver disease worldwide (Boyer, N. *et al.* J Hepatol. 32:98-112, 2000) so a significant focus of current antiviral research is directed toward the development of improved methods of treatment of chronic HCV infections in humans (Di Besceglie, A.M. and Bacon, B. R., Scientific American, Oct.: 80-85, (1999); Gordon, C. P., *et al.*, J. Med. Chem. **2005**, 48, 1-20; Maradpour, D.; *et al.*, Nat. Rev. Micro. **2007**, 5(6), 453-463). A

number of HCV treatments are reviewed by Bymock et al. in Antiviral Chemistry & Chemotherapy, 11:2; 79-95 (2000). Virologic cures of patients with chronic HCV infection are difficult to achieve because of the prodigious amount of daily virus



production in chronically infected patients and the high spontaneous mutability of HCV virus (Neumann, et al., *Science* **1998**, 282, 103-7; Fukimoto, et al., *Hepatology*, **1996**, 24, 1351-4; Domingo, et al., *Gene*, **1985**, 40, 1-8; Martell, et al., *J. Virol.* **1992**, 66, 3225-9.


5 Currently, there are primarily two antiviral compounds, ribavirin, a nucleoside analog, and interferon-alpha (α) (IFN), that are used for the treatment of chronic HCV infections in humans. Ribavirin alone is not effective in reducing viral RNA levels, has significant toxicity, and is known to induce anemia. The combination of IFN and ribavirin has been reported to be effective in the management of chronic hepatitis C
10 (Scott, L. J., et al. *Drugs* **2002**, 62, 507-556) but less than half the patients given this treatment show a persistent benefit.

 Combined, infections from the *Flaviviridae* virus family cause significant mortality, morbidity and economic losses throughout the world. Alkynyl substituted thiophenes with anti-*Flaviviridae* virus activity have been disclosed by Chan, et al.,
15 WO 2008058393; Wunberg, et al., WO 2006072347; and Chan, et al., WO 2002100851; but none of these are currently clinically approved antiviral therapeutics. Therefore, there remains a need to develop effective treatments for *Flaviviridae* virus infections.

20

SUMMARY OF THE INVENTION

Provided are compounds of Formula I:



25

Formula (I),

or a pharmaceutically acceptable salt or ester thereof, wherein:

R¹ is selected from the group consisting of optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 -

- 14 membered heteroaryl, optionally substituted 3 -12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted 3-18 membered heterocyclylalkyl and optionally substituted C₆₋₁₈ arylalkyl, wherein, each substituted R¹ is substituted with one or more Q¹;
- 5 each Q¹ is independently selected from the group consisting of halogen, oxo, oxide, -NO₂, -N(=O), -SR¹⁰, -S(O)R¹⁰, -S(O)₂R¹⁰, -S(O)₂NR¹⁰R¹¹, -NR¹⁰C(O)R¹¹, -NR¹⁰C(O)NR¹¹R¹², -NR¹⁰S(O)R¹¹, -NR¹⁰S(O)₂R¹¹, -OP(O)R¹¹R¹², -P(O)R¹¹R¹², -P(O)OR¹¹R¹², -P(O)(OR¹¹)OR¹², -C(O)NR¹¹R¹², optionally substituted C₁₋₆ alkyl, optionally substituted C₂₋₆ alkenyl, optionally substituted C₂₋₆ alkynyl,
- 10 optionally substituted C₃₋₆ cycloalkyl, optionally substituted C₆₋₁₂ arylalkyl, optionally substituted C₆₋₁₂ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted C₁₋₆ alkyloxy, optionally substituted C₂₋₆ alkenyloxy, optionally substituted C₂₋₆ alkynyloxy, optionally substituted C₃₋₆ cycloalkyloxy, optionally substituted C₆₋₁₂ aryloxy, optionally substituted 3 - 14 membered heteroaryloxy, optionally substituted
- 15 4-12 membered heterocyclyloxy, optionally substituted -C(O)C₁₋₆ alkyl, optionally substituted -C(O)C₂₋₆ alkenyl, optionally substituted -C(O)C₂₋₆ alkynyl, optionally substituted -C(O)C₃₋₆ cycloalkyl, optionally substituted -C(O)C₆₋₁₂ aryl, optionally substituted -C(O)-3 - 14 membered heteroaryl, optionally substituted -C(O)C₆₋₁₂ arylalkyl, optionally substituted -3-10 membered
- 20 heterocyclyl, -OH, -NR¹¹R¹², -C(O)OR¹⁰, -CN, -N₃, -C(=NR¹³)NR¹¹R¹², -C(=NR¹³)OR¹⁰, -NR¹⁰C(=NR¹³)NR¹¹R¹², -NR¹¹C(O)OR¹⁰, and -OC(O)NR¹¹R¹²;
- each R¹⁰, R¹¹, and R¹², independently, is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted
- 25 C₆₋₁₄ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, and optionally substituted C₆₋₁₈ arylalkyl;
- or R¹¹ and R¹² taken together with the atoms to which they are attached form a 3 to 10 membered heterocyclyl;
- 30 each R¹³, independently, is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl,
- 35 optionally substituted C₆₋₁₈ arylalkyl, -CN, -C(O)R¹⁴, -CHO and -S(O)₂R¹⁴;

- each R^{14} , independently, is optionally substituted C_{1-12} alkyl;
 wherein, each substituted Q^1 , substituted R^{10} , substituted R^{11} , substituted R^{12} , substituted R^{13} , or substituted R^{14} is independently substituted with one or more Q^6 ;
 R^2 is selected from the group consisting of optionally substituted C_{1-12} alkyl,
 5 optionally substituted C_{2-12} alkenyl, optionally substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3 - 12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, and optionally substituted C_{6-18} arylalkyl;
 10 wherein, each substituted R^2 is substituted with one or more Q^2 ;
 each Q^2 , independently, is selected from the group consisting of halogen, oxo, oxide, $-NO_2$, $-N(=O)$, $-SR^{20}$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)_2NR^{20}R^{21}$, $-NR^{20}C(O)R^{21}$, $-NR^{20}C(O)NR^{21}R^{22}$, $-NR^{20}S(O)R^{21}$, $-NR^{20}S(O)_2R^{21}$, $-OP(O)R^{21}R^{22}$, $-P(O)R^{21}R^{22}$, $-P(O)OR^{21}R^{22}$, $-P(O)(OR^{21})OR^{22}$, $-C(O)NR^{21}R^{22}$, optionally substituted
 15 C_{1-6} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted C_{3-6} cycloalkyl, optionally substituted C_{6-12} arylalkyl, optionally substituted C_{6-12} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted C_{1-6} alkyloxy, optionally substituted C_{2-6} alkenyloxy, optionally substituted C_{2-6} alkynyloxy, optionally substituted C_{3-6} cycloalkyloxy, optionally substituted C_{6-12}
 20 aryloxy, optionally substituted 3 - 14 membered heteroaryloxy, optionally substituted 4-12 membered heterocyclyloxy, optionally substituted $-C(O)C_{1-6}$ alkyl, optionally substituted $-C(O)C_{2-6}$ alkenyl, optionally substituted $-C(O)C_{2-6}$ alkynyl, optionally substituted $-C(O)C_{3-6}$ cycloalkyl, optionally substituted $-C(O)C_{6-12}$ aryl, optionally substituted $-C(O)-3-14$ membered heteroaryl, optionally substituted $-C(O)C_{6-12}$
 25 arylalkyl, optionally substituted 3-10 membered heterocyclyl, $-OH$, $-NR^{21}R^{22}$, $-C(O)OR^{20}$, $-CN$, $-N_3$, $-C(=NR^{23})NR^{21}R^{22}$, $-C(=NR^{23})OR^{20}$, $-NR^{20}C(=NR^{23})NR^{21}R^{22}$, $-NR^{21}C(O)OR^{20}$, and $-OC(O)NR^{21}R^{22}$;
 each R^{20} , R^{21} , and R^{22} , independently, is selected from the group consisting of H, optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, optionally
 30 substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, and optionally substituted C_{6-18} arylalkyl;
 or R^{21} and R^{22} taken together with the atoms to which they are attached form
 35 a 3 to 10 membered heterocyclyl;

- each R^{23} , independently, is selected from the group consisting of H, optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, optionally substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted C_{6-18} arylalkyl, -CN, $-C(O)R^{24}$, -CHO and $-S(O)_2R^{24}$;
- each R^{24} individually is optionally substituted C_{1-12} alkyl;
- wherein, each substituted Q^2 , substituted R^{20} , substituted R^{21} , substituted R^{22} , substituted R^{23} , or substituted R^{24} is independently substituted with one or more Q^6 ;
- 10 Y is $-R^3-L-Het$, $-N(R^4)(R^5)$ or $-R^6=NOR^7$;
- R^3 is selected from the group consisting of optionally substituted C_{1-12} alkylene, C_{2-12} alkenylene, substituted C_{2-12} alkenylene, C_{2-12} alkynylene, substituted C_{2-12} alkynylene, C_{3-12} cycloalkylene, substituted C_{3-12} cycloalkylene, C_{3-12} cycloalkylalkylene, substituted C_{3-12} cycloalkylalkylene, optionally substituted C_{6-14} arylene, optionally substituted 3 - 14 membered heteroarylene, optionally substituted 3 - 12 membered heterocyclylene, optionally substituted 3-18 membered heteroarylalkylene, and optionally substituted C_{6-18} arylalkylene;
- wherein each substituted R^3 is substituted with one or more Q^3 ;
- each Q^3 , independently, is selected from the group consisting of
- 20 halogen, oxo, oxide, $-NO_2$, $-N(=O)$, $-SR^{30}$, $-S(O)R^{30}$, $-S(O)_2R^{30}$, $-S(O)_2NR^{30}R^{31}$, $-NR^{30}C(O)R^{31}$, $-NR^{30}C(O)NR^{31}R^{32}$, $-NR^{30}S(O)R^{31}$, $-NR^{30}S(O)_2R^{31}$, $-OP(O)R^{31}R^{32}$, $-P(O)R^{31}R^{32}$, $-P(O)OR^{31}R^{32}$, $-P(O)(OR^{31})OR^{32}$, $-C(O)NR^{31}R^{32}$, optionally substituted C_{1-6} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted C_{3-6} cycloalkyl, optionally substituted C_{6-12} arylalkyl, optionally substituted C_{6-12} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted C_{1-6} alkyloxy, optionally substituted C_{2-6} alkenyloxy, optionally substituted C_{2-6} alkynyloxy, optionally substituted C_{3-6} cycloalkyloxy, optionally substituted C_{6-12} aryloxy, optionally substituted 3 - 14 membered heteroaryloxy, optionally substituted 4-12 membered heterocycliloxy, optionally substituted $-C(O)C_{1-6}$ alkyl, optionally substituted $-C(O)C_{2-6}$ alkenyl, optionally substituted $-C(O)C_{2-6}$ alkynyl, optionally substituted $-C(O)C_{3-6}$ cycloalkyl, optionally substituted $-C(O)C_{6-12}$ aryl, optionally substituted $-C(O)-3-14$ membered heteroaryl, optionally substituted $-C(O)C_{6-12}$ arylalkyl, optionally substituted 3-10 membered heterocyclyl, -OH, $-NR^{31}R^{32}$, $-C(O)OR^{30}$, -CN, $-N_3$, $-C(=NR^{33})NR^{31}R^{32}$,
- 35 $-C(=NR^{33})OR^{30}$, $-NR^{30}C(=NR^{33})NR^{31}R^{32}$, $-NR^{31}C(O)OR^{30}$, and $-OC(O)NR^{31}R^{32}$;

each R^{30} , R^{31} , and R^{32} , independently is selected from the group consisting of H, optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, optionally substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, and optionally substituted C_{6-18} arylalkyl;

or R^{31} and R^{32} taken together with the atoms to which they are attached form a 3 to 10 membered heterocyclyl;

each R^{33} independently is selected from the group consisting of H, optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, optionally substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted C_{6-18} arylalkyl, -CN, -C(O) R^{34} , -CHO and -S(O) R^{34} ;

each R^{34} individually is optionally substituted C_{1-12} alkyl;

wherein, each substituted Q^3 , substituted R^{30} , substituted R^{31} , substituted R^{32} , substituted R^{33} , or substituted R^{34} is independently substituted with one or more Q^6 ;

L is selected from the group consisting of -OC(O)N(R^4)-, -N(R^4)C(O)O-, -N(R^4)S(O) $_2$ -, -N(R^4)C(O)-, -C(O)-, -C(O)O-, -OC(O)-, -N(R^4)N(R^4)C(O)O-, and -N(R^4)N(R^4)-;

Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl;

wherein, each substituted Het is substituted with one or more Q^4 ;

each Q^4 , independently, is selected from the group consisting of halogen, oxo, oxide, -NO $_2$, -N(=O), -SR ^{40} , -S(O)R ^{40} , -S(O) $_2$ R ^{40} , -S(O) $_2$ NR ^{40} R ^{41} , -NR ^{40} C(O)R ^{41} , -NR ^{40} C(O)NR ^{41} R ^{42} , -NR ^{40} S(O)R ^{41} , -NR ^{40} S(O) $_2$ R ^{41} , -OP(O)R ^{41} R ^{42} , -P(O)R ^{41} R ^{42} , -P(O)OR ^{41} R ^{42} , -P(O)(OR ^{41})OR ^{42} , -C(O)NR ^{41} R ^{42} , optionally substituted C_{1-6} alkyl, optionally substituted C_{2-6} alkenyl, optionally substituted C_{2-6} alkynyl, optionally substituted C_{3-6} cycloalkyl, optionally substituted C_{6-12} arylalkyl, optionally substituted C_{6-12} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted C_{1-6} alkyloxy, optionally substituted C_{2-6} alkenyloxy, optionally substituted C_{2-6} alkynyloxy, optionally substituted C_{3-6} cycloalkyloxy, optionally substituted C_{6-12} aryloxy, optionally substituted 3 - 14 membered heteroaryloxy, optionally substituted 4-12 membered heterocyclyloxy, optionally substituted -C(O) C_{1-6} alkyl, optionally substituted -C(O) C_{2-6} alkenyl, optionally substituted -C(O) C_{2-6} alkynyl, optionally

- substituted -C(O)C₃₋₆ cycloalkyl, optionally substituted -C(O)C₆₋₁₂ aryl, optionally substituted -C(O)- 3 -14 membered heteroaryl, optionally substituted -C(O)C₆₋₁₂ arylalkyl, optionally substituted 3-10 membered heterocyclyl, -OH, -NR⁴¹R⁴², -C(O)OR⁴⁰, -CN, -N₃, -C(=NR⁴³)NR⁴¹R⁴²,
5 -C(=NR⁴³)OR⁴⁰, -NR⁴⁰C(=NR⁴³)NR⁴¹R⁴², -NR⁴¹C(O)OR⁴⁰, and -OC(O)NR⁴¹R⁴²;
each R⁴⁰, R⁴¹, and R⁴², independently is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl,
10 and optionally substituted C₆₋₁₈ arylalkyl;
or R⁴¹ and R⁴² taken together with the atoms to which they are attached form a 3 to 10 membered heterocyclyl;
each R⁴³, independently, is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted C₆₋₁₈ arylalkyl, -CN, -C(O)R⁴⁴, -CHO and -S(O)₂R⁴⁴;
15 each R⁴⁴ individually is optionally substituted C₁₋₁₂ alkyl;
wherein, each substituted Q⁴, substituted R⁴⁰, substituted R⁴¹, substituted R⁴², substituted R⁴³, or substituted R⁴⁴ is independently substituted with one or more Q⁵;
each Q⁵, individually, is selected from the group consisting of halogen, oxo, oxide, -NO₂, -N(=O), -SR⁵⁰, -S(O)R⁵⁰, -S(O)₂R⁵⁰, -S(O)₂NR⁵⁰R⁵¹,
25 -NR⁵⁰C(O)R⁵¹, -NR⁵⁰C(O)NR⁵¹R⁵², -NR⁵⁰S(O)R⁵¹, -NR⁵⁰S(O)₂R⁵¹, -OP(O)R⁵¹R⁵², -P(O)R⁵¹R⁵², -P(O)OR⁵¹R⁵², -P(O)(OR⁵¹)OR⁵², -C(O)NR⁵¹R⁵², optionally substituted C₁₋₆ alkyl, optionally substituted C₂₋₆ alkenyl, optionally substituted C₂₋₆ alkynyl, optionally substituted C₃₋₆ cycloalkyl, optionally substituted C₆₋₁₂ arylalkyl, optionally substituted C₆₋₁₂ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted C₁₋₆ alkyloxy, optionally substituted C₂₋₆ alkenyloxy, optionally substituted C₂₋₆ alkynyloxy, optionally substituted C₃₋₆ cycloalkyloxy, optionally substituted C₆₋₁₂ aryloxy, optionally substituted 3 - 14 membered heteroaryloxy, optionally substituted 4-12 membered heterocyclyloxy, optionally substituted -C(O)C₁₋₆ alkyl, optionally substituted -C(O)C₂₋₆ alkenyl, optionally substituted -C(O)C₂₋₆ alkynyl, optionally substituted -C(O)C₃₋₆ cycloalkyl, optionally substituted -C(O)C₆₋₁₂ aryl, optionally substituted -C(O)C₃₋₆ cycloalkyl, optionally substituted -C(O)C₆₋₁₂ aryl, optionally

- substituted -C(O)- 3 -14 membered heteroaryl, optionally substituted -C(O)C₆₋₁₂ arylalkyl, optionally substituted 3-10 membered heterocyclyl, -OH, -NR⁵¹R⁵², -C(O)OR⁵⁰, -CN, -N₃, -C(=NR⁵³)NR⁵¹R⁵², -C(=NR⁵³)OR⁵⁰, -NR⁵⁰C(=NR⁵³)NR⁵¹R⁵², -NR⁵¹C(O)OR⁵⁰, and -OC(O)NR⁵¹R⁵²;
- 5 each R⁵⁰, R⁵¹, and R⁵², independently is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl,
- 10 and optionally substituted C₆₋₁₈ arylalkyl;
or R⁵¹ and R⁵² taken together with the atoms to which they are attached form a 3 to 10 membered heterocyclyl;
each R⁵³, independently, is selected from the group consisting of H, optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl,
- 15 optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3-12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted C₆₋₁₈ arylalkyl, -CN, -C(O)R⁵⁴, -CHO and -S(O)₂R⁵⁴;
- each R⁵⁴, independently, is optionally substituted C₁₋₁₂ alkyl;
- 20 wherein, each substituted Q⁵, substituted R⁵⁰, substituted R⁵¹, substituted R⁵², substituted R⁵³, or substituted R⁵⁴ is independently substituted with one or more Q⁶;
- each Q⁶, independently, is selected from the group consisting of halogen, oxo, oxide, -NO₂, -N(=O), -SR⁶⁰, -S(O)R⁶⁰, -S(O)₂R⁶⁰, -S(O)₂NR⁶⁰R⁶¹, -NR⁶⁰C(O)R⁶¹, -NR⁶⁰C(O)NR⁶¹R⁶², -NR⁶⁰S(O)R⁶¹, -NR⁶⁰S(O)₂R⁶¹, -OP(O)R⁶¹R⁶²,
- 25 -P(O)R⁶¹R⁶², -P(O)OR⁶¹R⁶², -P(O)(OR⁶¹)OR⁶², -C(O)NR⁶¹R⁶², C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₆ cycloalkyl, C₆₋₁₂ arylalkyl, C₆₋₁₂ aryl, 3 - 14 membered heteroaryl, C₁₋₆ alkyloxy, C₂₋₆ alkenyloxy, C₂₋₆ alkynyloxy, C₃₋₆ cycloalkyloxy, C₆₋₁₂ aryloxy, 3 - 14 membered heteroaryloxy, 4-12 membered heterocyclyloxy, -C(O)C₁₋₆ alkyl, -C(O)C₂₋₆ alkenyl, -C(O)C₂₋₆ alkynyl, -C(O)C₃₋₆ cycloalkyl, -C(O)C₁₋₆
- 30 haloalkyl, -C(O)C₆₋₁₂ aryl, -C(O)- 3-14 membered heteroaryl, -C(O)C₃₋₁₂ arylalkyl, 3-10 membered heterocyclyl, -OH, -NR⁶¹R⁶², -C(O)OR⁶⁰, -CN, -N₃, -C(=NR⁶³)NR⁶¹R⁶², -C(=NR⁶³)OR⁶⁰, -NR⁶⁰C(=NR⁶³)NR⁶¹R⁶², -NR⁶¹C(O)OR⁶⁰, and -OC(O)NR⁶¹R⁶²;
- each R⁶⁰, R⁶¹, and R⁶², independently, is selected from the group consisting of H, C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl, C₃₋₁₂ cycloalkyl, C₁₋₁₂ haloalkyl, C₆₋₁₄ aryl,

3 - 14 membered heteroaryl, 3-12 membered heterocyclyl, 3-18 membered heteroarylalkyl, and C₆₋₁₈ arylalkyl;

or R⁶¹ and R⁶² taken together with the atoms to which they are attached form a 3 to 10 membered heterocyclyl;

5 each R⁶³ independently is selected from the group consisting of H, C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl, C₃₋₁₂ cycloalkyl, C₆₋₁₄ aryl, 3 - 14 membered heteroaryl, 3-12 membered heterocyclyl, 3-18 membered heteroarylalkyl, C₆₋₁₈ arylalkyl, -CN, -C(O)R⁶⁴, -CHO and -S(O)₂R⁶⁴;

each R⁶⁴ individually is C₁₋₁₂ alkyl;

10 each R⁴ is independently H, C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl is optionally substituted with one or more Q¹;


15 each R⁵ is independently C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl is optionally substituted with one or more Q¹;

20 R⁶ is C_{1-C12} alkylyne, C_{3-C12} cycloalkylyne, or 3-12 membered heterocycllyne wherein each C_{1-C12} alkylyne, C_{3-C12} cycloalkylyne, or 3-12 membered heterocycllyne is optionally substituted with one or more Q¹; and

25 R⁷ is selected from the group consisting of optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 3 -14 membered heteroaryl, optionally substituted 3 -12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted 3-18 membered heterocyclalkyl and optionally substituted C₆₋₁₈ arylalkyl;

wherein, each substituted R^7 is substituted with one or more Q^2 .

Also provided is a compound of Formula I:



Formula (I),

5 or a pharmaceutically acceptable salt or ester thereof, wherein:

R^1 is C_{1-12} alkyl;

R^2 is optionally substituted C_{3-12} cycloalkyl;

wherein, each substituted R^2 is substituted with one or more C_{1-6} alkyl;

Y is $-R^3-L-Het$, $-N(R^4)(R^5)$ or $-R^6=NOR^7$;

10 R^3 is selected from the group consisting of C_{1-12} alkylene, C_{3-12} cycloalkylene, and 3-12 membered heterocyclylene, wherein said 3-12 membered heterocyclylene comprises one to four heteroatoms selected from O, S, or N;

L is selected from the group consisting of $-OC(O)N(R^4)-$, $-N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-N(R^4)N(R^4)C(O)O-$, and $-N(R^4)N(R^4)-$;

Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 5-10 membered heteroaryl, wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 5-10 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N;

20 wherein, each substituted Het is substituted with one or more Q^4 ;

each Q^4 , independently, is selected from the group consisting of halogen, oxo, C_{1-6} alkyl, and $-C(O)OR^{40}$;

R⁴⁰ is C₁₋₁₂ alkyl;

each R⁴ is independently selected from the group consisting of H, C₁-C₁₂ alkyl, C₃-C₁₂ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C₁-C₁₂ alkyl, C₃-C₁₂ cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

each R⁵ is independently selected from the group consisting of C₁-C₁₂ alkyl, C₃-C₁₂ cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C₁-C₁₂ alkyl, C₃-C₁₂ cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

each Q¹ is independently selected from the group consisting of C₁₋₆ alkyl, 5-10 membered heteroaryloxy, 4-12 membered heterocyclyloxy, 3-10 membered heterocyclyl, -OH, and -C(O)OR¹⁰, wherein said heteroaryl portion of said 5-10 membered heteroaryloxy, said heterocyclyl portion of said 4-12 membered heterocyclyloxy, and said 3-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

R¹⁰ is selected from the group consisting of H, and C₁₋₁₂ alkyl;

R⁶ is C₃-C₁₂ cycloalkyne;

R⁷ is selected from the group consisting of optionally substituted C₁₋₁₂ alkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 6-11 membered heteroarylalkyl, and optionally substituted C₇₋₁₁ arylalkyl, wherein said heteroaryl portion of said 6-11 membered heteroarylalkyl comprises one to four heteroatoms selected from O, S, or N;

wherein, each substituted R⁷ is substituted with one or more Q²,

each Q^2 , independently, is selected from the group consisting of $-S(O)_2R^{20}$, optionally substituted C_{1-6} alkyl, OH, $-C(O)OR^{20}$, and $-CN$; and


each R^{20} , independently, is selected from the group consisting of H, and C_{1-12} alkyl,

5 wherein:

the term alkyl refers to a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms; and

the term cycloalkyl refers to a saturated or partially unsaturated ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about
10 20 carbon atoms as a polycycle.

In another embodiment, also provided is a compound of compound of Formula I:



Formula (I),

15 or a pharmaceutically acceptable salt or ester thereof, wherein:

R^1 is C_{1-12} alkyl;

R^2 is optionally substituted C_{3-12} cycloalkyl;

wherein, each substituted R^2 is substituted with one or more Q^2 ,

each Q^2 , independently, is selected from the group consisting of $-S(O)_2R^{20}$, optionally substituted C_{1-6} alkyl, OH, $-C(O)OR^{20}$, and $-CN$;

20 each R^{20} , independently, is selected from the group consisting of H, and C_{1-12} alkyl;

Y is $-R^3-L-Het$, $-N(R^4)(R^5)$ or $-R^6=NOR^7$;

R³ is selected from the group consisting of C₁₋₁₂ alkylene, C₃₋₁₂ cycloalkylene, and 3-12 membered heterocyclylene, wherein said 3-12 membered heterocyclylene comprises one to four heteroatoms selected from O, S, or N;

5 **L** is selected from the group consisting of -OC(O)N(R⁴)-, -N(R⁴)C(O)O-, -N(R⁴)S(O)₂-, -N(R⁴)C(O)-, -C(O)-, -C(O)O-, -OC(O)-, -N(R⁴)N(R⁴)C(O)O-, and -N(R⁴)N(R⁴)-;

10 **Het** is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 5-10 membered heteroaryl, wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 5-10 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N;

 wherein, each substituted Het is substituted with one or more Q⁴;

 each Q⁴, independently, is selected from the group consisting of halogen, oxo, C₁₋₆ alkyl, and -C(O)OR⁴⁰;

 R⁴⁰ is C₁₋₁₂ alkyl;

15 each R⁴ is independently selected from the group consisting of H, C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

20 each R⁵ is independently selected from the group consisting of C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

25 each Q¹ is independently selected from the group consisting of C₁₋₆ alkyl, 5-10 membered heteroaryloxy, 4-12 membered heterocyclloxy, 3-10 membered

heterocyclyl, -OH, and -C(O)OR¹⁰, wherein said heteroaryl portion of said 5-10 membered heteroaryloxy, said heterocyclyl portion of said 4-12 membered heterocycliloxy, and said 3-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

5 R¹⁰ is selected from the group consisting of H, and C₁₋₁₂ alkyl;

R⁶ is C₃-C₁₂ cycloalkyl;

10 R⁷ is selected from the group consisting of optionally substituted C₁₋₁₂ alkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 6-11 membered heteroarylalkyl, and optionally substituted C₇₋₁₁ arylalkyl, wherein said heteroaryl portion of said 6-11 membered heteroarylalkyl comprises one to four heteroatoms selected from O, S, or N;

wherein, each substituted R⁷ is substituted with one or more Q²; and
wherein:

15 the term alkyl refers to a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms; and

the term cycloalkyl refers to a saturated or partially unsaturated ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to 20 carbon atoms as a polycycle.

20 In another embodiment, a method for treating *Flaviviridae* viral infections is provided comprising administering an effective amount of a compound of Formula I to a patient in need thereof. The compound of Formula I is administered to a human subject in need thereof, such as a human being who is infected with viruses of the *Flaviviridae* family. In another embodiment, the compound of Formula I is administered to a human subject in need thereof, such as a human being who is
25 infected with a HCV virus. In one embodiment, the treatment results in the reduction of one or more of the in viral loads or clearance of RNA in the patient.

In another embodiment, provided is a method of treating and/or preventing a disease caused by a viral infection wherein the viral infection is caused by a virus

selected from the group consisting of dengue virus, yellow fever virus, West Nile virus, Japanese encephalitis virus, tick-borne encephalitis virus, Junjin virus, Murray Valley encephalitis virus, St Louis encephalitis virus, Omsk hemorrhagic fever virus, bovine viral diarrhoea virus, Zika virus and Hepatitis C virus; by
5 administering to a subject in need thereof a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

In another embodiment, provided is the use of a compound of Formula I for the manufacture of a medicament for the treatment of *Flaviviridae* viral infections. In another aspect of this embodiment, the *Flaviviridae* viral infection is an HCV
10 infection.

In another embodiment, provided is a compound of Formula I for use in treating a *Flaviviridae* viral infection. In another aspect of this embodiment, the *Flaviviridae* viral infection is an acute or chronic HCV infection. In another aspect of this embodiment, the treatment results in the reduction of one or more of the
15 viral loads or clearance of RNA in the patient. In another aspect of this embodiment, the treatment results in the reduction of the HCV viral load or clearance of HCV viral RNA in the patient.

In another embodiment, provided is a method for treating or preventing HCV comprising administering an effective amount of a compound of Formula I to a
20 patient in need thereof. In another embodiment, provided is the use of a compound of the present invention for the manufacture of a medicament for the treatment or prevention of HCV.

In another embodiment, provided is a pharmaceutical composition comprising a compound of Formula I and one or more pharmaceutically acceptable carriers or excipients. The pharmaceutical composition of Formula I may further
25 comprise one or more additional therapeutic agents. The one or more additional therapeutic agent may be, without limitation, selected from: interferons, ribavirin or


its analogs, HCV NS3 protease inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers, and other drugs for treating HCV, or mixtures thereof.

In one embodiment, provided is the use of the compound described herein, for the treatment of a *Flaviviridae* viral infection.

In one embodiment, provided is the use of the compound described herein, for the treatment of HCV.

In one embodiment, provided is the use of the compound described herein, in combination with at least one additional agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, other anti-fibrotic agents, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers and other drugs for treating HCV; or mixtures thereof, for the treatment of HCV.

In another embodiment, provided is a method for the treatment or prevention of the symptoms or effects of an HCV infection in an infected animal which comprises



administering to, *i.e.* treating, said animal with a pharmaceutical combination composition or formulation comprising an effective amount of a Formula I compound, and a second compound having anti-HCV properties.

5 In another embodiment, provided are compounds of Formula I and pharmaceutically acceptable salts thereof and all racemates, enantiomers, diastereomers, tautomers, polymorphs, pseudopolymorphs and amorphous forms thereof.

In another embodiment, provided are processes and novel intermediates disclosed herein which are useful for preparing Formula I compounds.

10 In other embodiments, novel methods for synthesis, analysis, separation, isolation, purification, characterization, and testing of the compounds of Formula I are provided.

The present invention includes combinations of aspects and embodiments, as well as preferences, as herein described throughout the present specification.


DETAILED DESCRIPTION

Reference will now be made in detail to certain embodiments of the invention, examples of which are illustrated in the accompanying structures and formulas. While the invention will be described in conjunction with the enumerated embodiments, it will be understood that they are not intended to limit the invention to those embodiments. On the contrary, the invention is intended to cover all alternatives, modifications, and equivalents, which may be included within the scope of the present invention as defined herein.


25 In one embodiment of Formula I, R¹ is optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, or optionally substituted C₃₋₁₂ cycloalkyl. In another aspect of this embodiment, R¹ is optionally

substituted C₁-C₁₂ alkyl. In another aspect of this embodiment, R¹ is optionally substituted C₃-C₇ secondary or tertiary alkyl. In another aspect of this embodiment, R¹ is prop-2-yl(isopropyl) or 2-methylprop-2-yl(*t*-butyl).


5 In another embodiment of Formula I, R² is optionally substituted C₁₋₁₂ alkyl, optionally substituted C₂₋₁₂ alkenyl, optionally substituted C₂₋₁₂ alkynyl, optionally substituted C₃₋₁₂ cycloalkyl, or optionally substituted C₆₋₁₈ arylalkyl. In another aspect




of this embodiment, R^2 is optionally substituted C_{3-12} cycloalkyl. In another aspect of this embodiment, R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^2 is optionally substituted 4-methylcyclohexyl. In another aspect of this embodiment, R^2 is optionally substituted 4-methylcyclohexenyl. In a preferred aspect of this embodiment, R^2 is




In another preferred aspect of this embodiment, R^2 is



10 In another preferred aspect of this embodiment, R^2 is



In another preferred aspect of this embodiment, R^2 is



15 In another embodiment of Formula I, Y is $-R^3-L-Het$. In another embodiment of Formula I, Y is $-N(R^4)(R^5)$. In another embodiment of Formula I, Y is $-R^6=NOR^7$.

In another embodiment of Formula I, Y is $-R^3-L-Het$, R^1 is optionally substituted C_1-C_{12} alkyl, and R^2 is optionally substituted C_{3-12} cycloalkyl. In another aspect of this embodiment, R^3 is optionally substituted C_{1-12} alkylene, C_{2-12} alkenylene, substituted C_{2-12} alkenylene, C_{2-12} alkynylene, substituted C_{2-12} alkynylene, C_{3-12} cycloalkylene, substituted C_{3-12} cycloalkylene, C_{3-12} cycloalkylalkylene, substituted C_{3-12} cycloalkylalkylene, optionally substituted C_{6-14} arylene, optionally substituted 3 - 14 membered heteroarylene, optionally substituted 3 - 12 membered heterocyclylene, optionally substituted 3-18 membered heteroarylalkylene, or optionally substituted C_{6-18} arylalkylene. In another aspect of this embodiment, R^3 is optionally substituted C_{1-12} alkylene, C_{3-12} cycloalkylene, substituted C_{3-12} cycloalkylene, C_{3-12}

20

25

cycloalkylalkylene, substituted C_{3-12} cycloalkylalkylene, optionally substituted C_{6-14} arylene, or optionally substituted 3-12 membered heterocyclylene. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$, $-N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-N(R^4)N(R^4)C(O)O-$, or $-N(R^4)N(R^4)-$. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$. In another aspect of this embodiment, L is $-N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)S(O)_2-$. In another aspect of this embodiment, L is $-N(R^4)C(O)-$. In another aspect of this embodiment, L is $-C(O)-$. In another aspect of this embodiment, L is $-C(O)O-$. In another aspect of this embodiment, L is $-OC(O)-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)-$.

In another embodiment of Formula I, Y is $-R^3-L-Het$, R^1 is optionally substituted C_{1-12} alkyl, R^2 is optionally substituted C_{3-12} cycloalkyl and R^3 is optionally substituted C_{1-12} alkylene, C_{3-12} cycloalkylene, substituted C_{3-12} cycloalkylene, C_{3-12} cycloalkylalkylene, substituted C_{3-12} cycloalkylalkylene, optionally substituted C_{6-14} arylene, or optionally substituted 3-12 membered heterocyclylene. In another aspect of this embodiment, R^3 is optionally substituted C_{1-6} alkylene. In another aspect of this embodiment, R^3 is C_{4-6} cycloalkylene or substituted C_{4-6} cycloalkylene. In another aspect of this embodiment, R^3 is an optionally substituted 5-6 membered heterocyclylene. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$, $-N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-N(R^4)N(R^4)C(O)O-$, or $-N(R^4)N(R^4)-$. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$. In another aspect of this embodiment, L is $-N(R^4)C(O)O-$. In another aspect of this embodiment, $-N(R^4)S(O)_2-$. In another aspect of this embodiment, L is $-N(R^4)C(O)-$. In another aspect of this embodiment, L is $-C(O)-$. In another aspect of this embodiment, L is $-C(O)O-$. In another aspect of this embodiment, L is $-OC(O)-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)-$. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N.

In another embodiment of Formula I, Y is $-N(R^4)(R^5)$, R^1 is optionally substituted C_1-C_{12} alkyl, and R^2 is optionally substituted C_3-C_{12} cycloalkyl. In another aspect of this embodiment, R^4 is independently H or C_1-C_{12} alkyl wherein C_1-C_{12} alkyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment,

5 R^4 is H or optionally substituted C_1-C_6 alkyl. In another aspect of this embodiment, R^5 is C_1-C_{12} alkyl, C_3-C_{12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl wherein each C_1-C_{12} alkyl, C_3-C_{12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^5 is C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5-10

10 membered heteroaryl or 4-10 membered heterocyclyl wherein each C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5 - 10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q^1 .

In another embodiment of Formula I, Y is $-R^6=NOR^7$, R^1 is optionally substituted C_1-C_{12} alkyl, and R^2 is optionally substituted C_3-C_{12} cycloalkyl. In another aspect of this embodiment, R^6 is C_1-C_{12} alkylyne, C_3-C_{12} cycloalkylyne, or 3-12 membered heterocycliylyne wherein each C_1-C_{12} alkylyne, C_3-C_{12} cycloalkylyne, or 3-12 membered heterocycliylyne is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^6 is C_5-C_6 cycloalkylyne or 4-6 membered heterocycliylyne wherein each C_5-C_6 cycloalkylyne or 4-6 membered heterocycliylyne

20 is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^6 is cyclohexylyne. In another aspect of this embodiment, R^7 is optionally substituted C_{1-12} alkyl, optionally substituted C_{2-12} alkenyl, optionally substituted C_{2-12} alkynyl, optionally substituted C_{3-12} cycloalkyl, optionally substituted C_{6-14} aryl, optionally substituted 3 - 14 membered heteroaryl, optionally substituted 3 - 12 membered heterocyclyl, optionally substituted 3-18 membered heteroarylalkyl, optionally substituted 3-18 membered heterocyclylalkyl or optionally substituted C_{6-18} arylalkyl; wherein, when R^7 is substituted, R^7 is substituted with one or more Q^2 . In another aspect of this embodiment, R^7 is optionally substituted C_1-C_6 alkyl. In another aspect of this embodiment, R^7 is optionally substituted C_7-C_{11} arylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heteroarylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heterocyclalkyl.


25

30


In another embodiment of Formula I, R^1 is optionally substituted C_3-C_7 secondary or tertiary alkyl and R^2 is optionally substituted methylcyclohexyl or

35 optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^1


is prop-2-yl (isopropyl) or 2-methylprop-2-yl (*t*-butyl). In another aspect of this embodiment, R^2 is optionally substituted 4-methylcyclohexyl. In a preferred aspect of this embodiment, R^2 is




5 In another preferred aspect of this embodiment, R^2 is



In another preferred aspect of this embodiment, R^2 is



In another preferred aspect of this embodiment, R^2 is



10

In another embodiment of Formula I, Y is $-R^3-L-Het$, R^1 is optionally substituted C_3-C_7 secondary or tertiary alkyl and R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^3 is optionally substituted C_{1-6} alkylene. In another aspect of this embodiment, R^3 is C_{4-6} cycloalkylene or substituted C_{4-6} cycloalkylene. In another aspect of this embodiment, R^3 is an optionally substituted 5-6 membered heterocyclylene. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$, $-N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-N(R^4)N(R^4)C(O)O-$, or $-N(R^4)N(R^4)-$. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$. In another aspect of this embodiment, L is $-N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)S(O)_2-$. In another aspect of this embodiment, L is $-N(R^4)C(O)-$. In another aspect of this embodiment, L is $-C(O)-$. In another aspect of this embodiment, L is $-C(O)O-$. In another aspect of this embodiment, L is $-OC(O)-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)N(R^4)-$. In another aspect of this embodiment,

15

20


25

Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment,

- 5 Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N.

- In another embodiment of Formula I, Y is $-R^3-N(R^4)C(O)O-Het$, R^1 is
 10 optionally substituted C_3-C_7 secondary or tertiary alkyl and R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^3 is optionally substituted C_{1-6} alkylene. In another aspect of this embodiment, R^3 is C_{4-6} cycloalkylene or substituted C_{4-6} cycloalkylene. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered
 15 heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N.
 20 In aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N.

- In another embodiment of Formula I, Y is $-R^3-C(O)O-Het$, R^1 is optionally substituted C_3-C_7 secondary or tertiary alkyl, R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl and R^3 is an optionally
 25 substituted 5-6 membered heterocyclylene. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment,
 30 Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment $-R^3-C(O)O-Het$ is:




wherein the pyrrolidinyl or piperidinyl ring is optionally substituted.

In another embodiment of Formula I, Y is $-N(R^4)(R^5)$, R^1 is optionally substituted C_3 - C_7 secondary or tertiary alkyl and R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^4 is H or optionally substituted C_1 - C_6 alkyl. In another aspect of this embodiment, R^5 is C_1 - C_{12} alkyl, C_3 - C_{12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl wherein each C_1 - C_{12} alkyl, C_3 - C_{12} cycloalkyl, 3 - 14 membered heteroaryl or 3-12 membered heterocyclyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^5 is C_1 - C_6 alkyl, C_5 - C_6 cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl wherein each C_1 - C_6 alkyl, C_5 - C_6 cycloalkyl, 5 - 10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q^1 .

In another embodiment of Formula I, Y is $-R^6=NOR^7$, R^1 is optionally substituted C_3 - C_7 secondary or tertiary alkyl and R^2 is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl. In another aspect of this embodiment, R^6 is C_5 - C_6 cycloalkyl, or 4-6 membered heterocyclyl wherein each C_5 - C_6 cycloalkyl or 4-6 membered heterocyclyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^6 is cyclohexyl. In another aspect of this embodiment, R^7 is optionally substituted C_1 - C_6 alkyl. In another aspect of this embodiment, R^7 is optionally substituted C_7 - C_{11} arylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heteroarylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heterocyclalkyl.

In another embodiment, compounds of Formula I comprise compounds of Formula II:




Formula II

or pharmaceutically acceptable salts and esters thereof, wherein:


R^2 is optionally substituted 4-methylcyclohexyl or optionally substituted

5 methylcyclohexenyl and the remaining variables are defined as for Formula I.


In one embodiment of Formula II, R^2 is:




In one embodiment of Formula II, R^2 is:



10 In another embodiment, R^2 is



In another embodiment, R^2 is




15 In one embodiment of Formula II, Y is $-R^3-L-Het$. In another aspect of this embodiment, R^3 is optionally substituted C_{1-6} alkylene. In another aspect of this embodiment, R^3 is C_{4-6} cycloalkylene or substituted C_{4-6} cycloalkylene. In another aspect of this embodiment, R^3 is an optionally substituted 5-6 membered heterocyclylene. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$, -
 20 $N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, -
 $N(R^4)N(R^4)C(O)O-$, or $-N(R^4)N(R^4)-$. In another aspect of this embodiment, L is $-OC(O)N(R^4)-$. In another aspect of this embodiment, L is $-N(R^4)C(O)O-$. In another aspect of this embodiment, L is $-N(R^4)S(O)_2-$. In another aspect of this embodiment, L is $-N(R^4)C(O)-$. In another aspect of this embodiment, $-C(O)-$. In another aspect of
 25 this embodiment, $-C(O)O-$. In another aspect of this embodiment, L is $-OC(O)-$. In


another aspect of this embodiment, L is $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)\text{C}(\text{O})\text{O}-$. In another aspect of this embodiment, L is $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)-$. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered

5 heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four


10 hetroatoms selected from O, S, or N. In another aspect of this embodiment, R^2 is:




In another aspect of this embodiment, R^2 is:



In another preferred aspect of this embodiment, R^2 is



In another preferred aspect of this embodiment, R^2 is




In another embodiment of Formula II, Y is $-\text{R}^3\text{-L-Het}$ wherein Het is an


20 optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is optionally substituted pyridinyl. In another aspect of this embodiment, Het is

25 optionally substituted pyridazinyl. In another aspect of this embodiment, Het is


optionally substituted tetrahydro-2*H*-pyranyl. In another aspect of this embodiment, Het is optionally substituted piperidinyl. In another aspect of this embodiment, Het is optionally substituted pyrrolidinyl. In another aspect of this embodiment, Het is optionally substituted tetrahydrothiophenyl. In another aspect of this embodiment, Het is optionally substituted pyrazinyl. In another aspect of this embodiment, Het is optionally substituted 1*H*-tetrazolyl. In another aspect of this embodiment, Het is optionally substituted azetidiny. In another aspect of this embodiment, Het is optionally substituted tetrahydrofuranyl. In another aspect of this embodiment, Het is optionally substituted tetrahydro-2*H*-furo[2,3-*b*]furanyl. In another aspect of this embodiment, Het is optionally substituted thiazolyl. In another aspect of this embodiment, Het is optionally substituted 1*H*-imidazolyl. In another aspect of this embodiment, Het is optionally substituted 4*H*-1,2,4-triazolyl. In another aspect of this embodiment, Het is optionally substituted 1*H*-pyrazolyl. In another aspect of this embodiment, Het is optionally substituted 1,3,4-thiadiazolyl. In another aspect of this embodiment, Het is optionally substituted quinolinyl. In another aspect of this embodiment, Het is optionally substituted [1,2,4]triazolo[4,3-*a*]pyridinyl. In another aspect of this embodiment, Het is optionally substituted thiophenyl. In another aspect of this embodiment, Het is optionally substituted 1,2,4-thiadiazolyl. In another aspect of this embodiment, Het is optionally substituted pyrimidinyl. In another aspect of this embodiment, Het is optionally substituted 1*H*-1,2,3-triazolyl. In another aspect of this embodiment, Het is optionally substituted 1,3,4-oxadiazolyl. In another aspect of this embodiment, Het is optionally substituted imidazo[1,2-*b*]pyridazinyl. In another aspect of this embodiment, R² is:




25 In another aspect of this embodiment, R² is:



In another preferred aspect of this embodiment, R² is




In another preferred aspect of this embodiment, R^2 is




- In another embodiment of Formula II, Y is $-R^3-N(R^4)C(O)O-Het$. In another aspect of this embodiment, R^3 is optionally substituted C_{1-6} alkylene. In another aspect of this embodiment, R^3 is C_{4-6} cycloalkylene or substituted C_{4-6} cycloalkylene.
- 5 In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N.
- 10 In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is optionally substituted pyridinyl. In another aspect of this embodiment, Het is optionally substituted pyridazinyl. In another aspect of this embodiment, Het is optionally substituted tetrahydro-2H-pyranyl. In another aspect of this embodiment, Het is optionally substituted piperidinyl. In another aspect of this embodiment, Het is optionally substituted pyrrolidinyl. In another aspect of this embodiment, Het is optionally substituted tetrahydrothiophenyl. In another aspect of this embodiment, Het is optionally substituted pyrazinyl. In another aspect of this embodiment, Het is optionally substituted 1H-tetrazolyl. In another aspect of this embodiment, Het is optionally substituted azetidyl. In another aspect of this embodiment, Het is optionally substituted tetrahydrofuranyl. In another aspect of this embodiment, Het is optionally substituted tetrahydro-2H-furo[2,3-b]furanyl. In another aspect of this embodiment, Het is optionally substituted thiazoyl. In another aspect of this embodiment, Het is optionally substituted 1H-imidazolyl. In another aspect of this embodiment, Het is optionally substituted 4H-1,2,4-triazolyl. In another aspect of this embodiment, Het is optionally substituted 1H-pyrazolyl. In another aspect of this embodiment, Het is optionally substituted 1,3,4-thiadiazolyl. In another aspect of this embodiment, Het is optionally substituted quinolinyl. In another aspect of this embodiment, Het is optionally substituted [1,2,4]triazolo[4,3-a]pyridinyl. In another aspect of this embodiment, Het is optionally substituted thiophenyl. In
- 25
- 30


- another aspect of this embodiment, Het is optionally substituted 1,2,4-thiadiazolyl. In another aspect of this embodiment, Het is optionally substituted pyrimidinyl. In another aspect of this embodiment, Het is optionally substituted 1*H*-1,2,3-triazolyl. In another aspect of this embodiment, Het is optionally substituted 1,3,4-oxadiazolyl. In
- 5 another aspect of this embodiment, Het is optionally substituted imidazo[1,2-*b*]pyridazinyl. In another aspect of this embodiment, R^2 is:




In another aspect of this embodiment, R^2 is:



- 10 In another preferred aspect of this embodiment, R^2 is




In another preferred aspect of this embodiment, R^2 is




- 15 In another embodiment of Formula II, Y is $-R^3-C(O)O-Het$ and R^3 is an optionally substituted 5-6 membered heterocyclylene. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl wherein said optionally substituted 3-12 membered heterocyclyl or optionally substituted 3-14 membered heteroaryl
- 20 comprises one to four heteroatoms selected from O, S, or N. In another aspect of this embodiment, Het is an optionally substituted 3-12 membered heterocyclyl comprising one or two heteroatoms selected from O, S, or N. In aspect of this embodiment, Het is an optionally substituted 5-10 membered heteroaryl comprising one to four heteroatoms selected from O, S, or N. In another aspect of this
- 25 embodiment, Het is optionally substituted pyridinyl. In another aspect of this


embodiment, Het is optionally substituted pyridazinyl. In another aspect of this
embodiment, Het is optionally substituted tetrahydro-2*H*-pyranyl. In another aspect
of this embodiment, Het is optionally substituted piperidinyl. In another aspect of this
embodiment, Het is optionally substituted pyrrolidinyl. In another aspect of this
5 embodiment, Het is optionally substituted tetrahydrothiophenyl. In another aspect of
this embodiment, Het is optionally substituted pyrazinyl. In another aspect of this
embodiment, Het is optionally substituted 1*H*-tetrazolyl. In another aspect of this
embodiment, Het is optionally substituted azetidiny. In another aspect of this
embodiment, Het is optionally substituted tetrahydrofuranyl. In another aspect of this
10 embodiment, Het is optionally substituted tetrahydro-2*H*-furo[2,3-*b*]furanyl. In
another aspect of this embodiment, Het is optionally substituted thiazoyl. In another
aspect of this embodiment, Het is optionally substituted 1*H*-imidazolyl. In another
aspect of this embodiment, Het is optionally substituted 4*H*-1,2,4-triazolyl. In another
aspect of this embodiment, Het is optionally substituted 1*H*-pyrazolyl. In another
15 aspect of this embodiment, Het is optionally substituted 1,3,4-thiadiazolyl. In another
aspect of this embodiment, Het is optionally substituted quinolinyl. In another aspect
of this embodiment, Het is optionally substituted [1,2,4]triazolo[4,3-*a*]pyridinyl. In
another aspect of this embodiment, Het is optionally substituted thiophenyl. In
another aspect of this embodiment, Het is optionally substituted 1,2,4-thiadiazolyl. In
20 another aspect of this embodiment, Het is optionally substituted pyrimidinyl. In
another aspect of this embodiment, Het is optionally substituted 1*H*-1,2,3-triazolyl. In
another aspect of this embodiment, Het is optionally substituted 1,3,4-oxadiazolyl. In
another aspect of this embodiment, Het is optionally substituted imidazo[1,2-
b]pyridazinyl. In another aspect of this embodiment $-R^3-C(O)O-Het$ is:




wherein the pyrrolidinyl or piperidinyl ring is optionally substituted. In another aspect
of this embodiment, R^2 is:




In another aspect of this embodiment, R^2 is:



In another preferred aspect of this embodiment, R^2 is




In another preferred aspect of this embodiment, R^2 is




5

In another embodiment of Formula II, Y is $-N(R^4)(R^5)$. In another aspect of this embodiment, R^4 is H or optionally substituted C_1-C_6 alkyl. In another aspect of this embodiment, R^5 is C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl wherein each C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5 - 10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^4 is optionally substituted C_1-C_6 alkyl and R^5 is C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl wherein each C_1-C_6 alkyl, C_5-C_6 cycloalkyl, 5 - 10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^2 is:


15



In another aspect of this embodiment, R^2 is:




In another preferred aspect of this embodiment, R^2 is




20


In another preferred aspect of this embodiment, R^2 is




- In another embodiment of Formula II, Y is $-R^6=NOR^7$. In another aspect of this embodiment, R^6 is C_5-C_6 cycloalkylyne, or 4-6 membered heterocyclylyne wherein each C_5-C_6 cycloalkylyne or 4-6 membered heterocyclylyne is optionally substituted with one or more Q^1 . In another aspect of this embodiment, R^6 is cyclohexylyne. In another aspect of this embodiment, R^7 is optionally substituted C_1-C_6 alkyl. In another aspect of this embodiment, R^7 is optionally substituted C_7-C_{11} arylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heteroarylalkyl. In another aspect of this embodiment, R^7 is optionally substituted 6-11 membered heterocyclylalkyl. In another aspect of this embodiment, R^6 is cyclohexylyne and R^7 is optionally substituted C_1-C_6 alkyl. In another aspect of this embodiment, R^6 is cyclohexylyne and R^7 is optionally substituted C_7-C_{11} arylalkyl. In another aspect of this embodiment, R^6 is cyclohexylyne and R^7 is optionally substituted 6-11 membered heteroarylalkyl. In another aspect of this embodiment, R^6 is cyclohexylyne and R^7 is optionally substituted 6-11 membered heterocyclylalkyl. In another aspect of this embodiment, R^2 is:




In another aspect of this embodiment, R^2 is:




- 20 In another preferred aspect of this embodiment, R^2 is









In another preferred aspect of this embodiment, R^2 is





In another embodiment, the compound of Formula I or Formula II is




























Definitions

- 5 Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings. The fact that a particular term or phrase is not specifically defined should not be correlated to indefiniteness or lacking clarity, but rather terms herein are used within their ordinary meaning. When trade names are used herein, applicants intend to independently include the tradename product
- 10 and the active pharmaceutical ingredient(s) of the tradename product.

- The term “treating”, and grammatical equivalents thereof, when used in the context of treating a disease, means slowing or stopping the progression of a disease, or ameliorating at least one symptom of a disease, more preferably ameliorating more than one symptom of a disease. For example, treatment of a hepatitis C virus
- 15 infection can include reducing the HCV viral load in an HCV infected human being, and/or reducing the severity of jaundice present in an HCV infected human being.

- “Alkyl” is hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms. For example, an alkyl group can have 1 to 20 carbon atoms (i.e., C₁-C₂₀ alkyl), 1 to 10 carbon atoms (i.e., C₁-C₁₀ alkyl), or 1 to 6 carbon atoms (i.e., C₁-C₆ alkyl).
- 20 Examples of suitable alkyl groups include, but are not limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (*n*-Pr, *n*-propyl, -CH₂CH₂CH₃), 2-propyl (*i*-Pr, *i*-propyl, -CH(CH₃)CH₂CH₃), 1-butyl (*n*-Bu, *n*-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (*i*-Bu, *i*-butyl, -CH₂CH(CH₃)CH₂CH₃), 2-butyl (*s*-Bu, *s*-butyl, -CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (*t*-Bu, *t*-butyl, -C(CH₃)₃), 1-pentyl (*n*-pentyl, -CH₂CH₂CH₂CH₂CH₃), 2-pentyl
- 25 (-CH(CH₃)CH₂CH₂CH₃), 3-pentyl (-CH(CH₂CH₃)₂), 2-methyl-2-butyl (-C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (-CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (-CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (-CH₂CH(CH₃)CH₂CH₃), 1-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (-CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (-CH(CH₂CH₃)(CH₂CH₂CH₃)), 2-methyl-2-pentyl (-C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-

pentyl $(-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3)$, 4-methyl-2-pentyl $(-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2)$, 3-methyl-3-pentyl $(-\text{C}(\text{CH}_3)(\text{CH}_2\text{CH}_3)_2)$, 2-methyl-3-pentyl $(-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)_2)$, 2,3-dimethyl-2-butyl $(-\text{C}(\text{CH}_3)_2\text{CH}(\text{CH}_3)_2)$, 3,3-dimethyl-2-butyl $(-\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_3)$, and octyl $(-\text{CH}_2)_7\text{CH}_3$.

5 “Alkoxy” means a group having the formula $-\text{O-alkyl}$, in which an alkyl group, as defined above, is attached to the parent molecule via an oxygen atom. The alkyl portion of an alkoxy group can have 1 to 20 carbon atoms (i.e., $\text{C}_1\text{-C}_{20}$ alkoxy), 1 to 12 carbon atoms (i.e., $\text{C}_1\text{-C}_{12}$ alkoxy), or 1 to 6 carbon atoms (i.e., $\text{C}_1\text{-C}_6$ alkoxy).

10 Examples of suitable alkoxy groups include, but are not limited to, methoxy $(-\text{O-CH}_3)$ or $-\text{OMe}$, ethoxy $(-\text{OCH}_2\text{CH}_3)$ or $-\text{OEt}$, t-butoxy $(-\text{O-C}(\text{CH}_3)_3)$ or $-\text{OtBu}$, and the like.

 “Haloalkyl” is an alkyl group, as defined above, in which one or more hydrogen atoms of the alkyl group is replaced with a halogen atom. The alkyl portion of a haloalkyl group can have 1 to 20 carbon atoms (i.e., $\text{C}_1\text{-C}_{20}$ haloalkyl), 1 to 12 carbon atoms (i.e., $\text{C}_1\text{-C}_{12}$ haloalkyl), or 1 to 6 carbon atoms (i.e., $\text{C}_1\text{-C}_6$ alkyl).

15 Examples of suitable haloalkyl groups include, but are not limited to, $-\text{CF}_3$, $-\text{CHF}_2$, $-\text{CFH}_2$, $-\text{CH}_2\text{CF}_3$, and the like.

 “Alkenyl” is a hydrocarbon containing normal, secondary, tertiary, or cyclic carbon atoms with at least one site of unsaturation, i.e. a carbon-carbon, sp^2 double bond. For example, an alkenyl group can have 2 to 20 carbon atoms (i.e., $\text{C}_2\text{-C}_{20}$ alkenyl), 2 to 12 carbon atoms (i.e., $\text{C}_2\text{-C}_{12}$ alkenyl), or 2 to 6 carbon atoms (i.e., $\text{C}_2\text{-C}_6$ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, vinyl $(-\text{CH}=\text{CH}_2)$, allyl $(-\text{CH}_2\text{CH}=\text{CH}_2)$, cyclopentenyl $(-\text{C}_5\text{H}_7)$, and 5-hexenyl $(-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2)$.

25 “Alkynyl” is a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms with at least one site of unsaturation, i.e. a carbon-carbon, sp triple bond. For example, an alkynyl group can have 2 to 20 carbon atoms (i.e., $\text{C}_2\text{-C}_{20}$ alkynyl), 2 to 12 carbon atoms (i.e., $\text{C}_2\text{-C}_{12}$ alkynyl), or 2 to 6 carbon atoms (i.e., $\text{C}_2\text{-C}_6$ alkynyl). Examples of suitable alkynyl groups include, but are not limited to, acetylenic $(-\text{C}\equiv\text{CH})$, propargyl $(-\text{CH}_2\text{C}\equiv\text{CH})$, and the like.

30 “Alkylene” refers to a saturated, branched or straight chain radical or or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, an alkylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkylene radicals include, but are not limited to,

methylene ($-\text{CH}_2-$), 1,1-ethylene ($-\text{CH}(\text{CH}_3)-$), 1,2-ethylene ($-\text{CH}_2\text{CH}_2-$), 1,1-propylene ($-\text{CH}(\text{CH}_2\text{CH}_3)-$), 1,2-propylene ($-\text{CH}_2\text{CH}(\text{CH}_3)-$), 1,3-propylene ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), 1,4-butylene ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$), and the like.

5 "Alkenylene" refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. For example, an alkenylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkenylene radicals include, but are not limited to, 1,2-ethylene ($-\text{CH}=\text{CH}-$).

10 "Alkynylene" refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. For example, an alkynylene group can have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Typical alkynylene radicals include, but are not limited to, acetylene ($-\text{C}\equiv\text{C}-$), propargyl ($-\text{CH}_2\text{C}\equiv\text{C}-$), and 4-pentynyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}-$).

15 "Alkylyne" refers to a saturated, branched or straight chain radical having two radical centers derived by the removal of three hydrogen atoms from two carbon atoms of a parent alkane. For example, an alkylyne group can have 2 to 20 carbon atoms, 2 to 10 carbon atoms, or 2 to 6 carbon atoms. Typical alkylyne radicals include, but are not limited to, 1,2-ethyne ($-\text{CH}_2\text{CH}=\text{CH}_2$), 1,2-propyne ($-\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$), 1,3-propyne ($-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 1,4-butyne ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$), and the like.

20 "Aryl" means a monovalent aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. For example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 12 carbon atoms. Typical aryl groups include, but are not limited to, radicals derived from benzene (e.g., phenyl), substituted benzene, naphthalene, anthracene, biphenyl, and the like.

25 "Arylene" refers to an aryl as defined above having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent aryl. Typical arylene radicals include, but are not limited to, phenylene.


"Arylalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl radical. Typical arylalkyl groups include, but are not limited to, benzyl, 2-

phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. The arylalkyl group can comprise 6 to 20 carbon atoms, e.g., the alkyl moiety is 1 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.


5 "Cycloalkyl" refers to a saturated or partially unsaturated ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic cycloalkyl groups have 3 to 6 ring atoms, still more typically 5 or 6 ring atoms. Bicyclic cycloalkyl groups have 7 to 12 ring atoms, e.g., arranged as a bicyclo (4,5), (5,5), (5,6) or (6,6) system, or 9 or 10 ring atoms arranged as a bicyclo (5,6) or (6,6) system. Cycloalkyl groups include hydrocarbon mono-, bi-, and poly-cyclic rings, whether fused, bridged, or spiro. Non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, bicyclo[3.1.0]hex-6-yl and the like.

"Cycloalkylene" refers to a cycloalkyl as defined above having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent cycloalkyl. Typical cycloalkylene radicals include, but are not limited to, cyclopropylene, cyclobutylene, cyclopentylene and cyclohexylene.


20 "Cycloalkylyne" refers to a cycloalkyl as defined above having two radical centers derived by the removal of three hydrogen atoms from two carbon atoms of a parent cycloalkyl. Two of the hydrogen atoms are removed from the same carbon atom and one hydrogen atom is removed from an alternative carbon atom of the ring. Non-limiting examples of cycloalkylyne radicals include:



cyclobutylyne



cyclopentylyne



cyclohexylyne

25

"Halogen" refers to F, Cl, Br, or I.


As used herein the term "haloalkyl" refers to an alkyl group, as defined herein, that is substituted with at least one halogen. Examples of branched or straight chained "haloalkyl" groups as used herein include, but are not limited to, methyl, ethyl, propyl, isopropyl, n-butyl, and t-butyl substituted independently with one or more halogens, for example, fluoro, chloro, bromo, and iodo. The term "haloalkyl" should be interpreted to include such substituents as perfluoroalkyl groups such as $-\text{CF}_3$.

30

As used herein, the term "haloalkoxy" refers to a group $-OR^a$, where R^a is a haloalkyl group as herein defined. As non-limiting examples, haloalkoxy groups include $-O(CH_2)F$, $-O(CH)F_2$, and $-OCF_3$.


"Heterocycle" or "heterocyclyl" refers to a saturated or partially saturated
5 cyclic group having from 1 to 14 carbon atoms and from 1 to 6 heteroatoms selected from N, S, P, or O, and includes single ring and multiple ring systems including, fused, bridged, and spiro ring systems. "Heterocycle" or "heterocyclyl" as used herein includes by way of example and not limitation those heterocycles described in
10 Paquette, Leo A.; Principles of Modern Heterocyclic Chemistry (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; The Chemistry of Heterocyclic Compounds, A Series of Monographs (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; and J. Am. Chem. Soc. (1960) 82:5566. In one embodiment, the carbon, nitrogen, phosphorous, or sulfur atom(s) of the heterocyclic group may be oxidized to provide for C(=O), N-oxide, phosphinane
15 oxide, sulfinyl, or sulfonyl moieties.

As one example, substituted heterocyclyls include, for example, heterocyclic rings substituted with any of the substituents disclosed herein including oxo groups. A non-limiting example of a carbonyl substituted heterocyclyl is:




20


Examples of heterocycles include by way of example and not limitation dihydropyridyl, tetrahydropyridyl (piperidyl), tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, piperidinyl, 4-piperidonyl, pyrrolidinyl, azetidiny, 2-pyrrolidonyl, tetrahydrofuranyl, decahydroquinolinyl, octahydroisoquinolinyl, pyranly, morpholinyl,
25 and bis-tetrahydrofuranyl:



- “Heterocyclene” or “heterocyclylene” refers to a “heterocycle” or “heterocyclyl” as defined above having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent heterocycle, the removal of two hydrogen atoms from two nitrogen atoms of a parent heterocycle, or the removal of a hydrogen atom from a nitrogen and the removal of a hydrogen atom from a carbon atom of a parent heterocycle. Non-limiting examples of heterocyclene or heterocyclenes are:



- “Heterocyclelyne” or “heterocyclelyne” refers to a “heterocycle” or “heterocyclyl” as defined above having two radical centers derived by the removal of three hydrogen atoms from two carbon atoms of a parent heterocycle or the removal of a hydrogen atom from a nitrogen atom and the removal of two hydrogen atoms from the same carbon atom of a parent heterocycle. Non-limiting examples of heterocyclelyne or “heterocyclelyne radicals include:



"Heteroaryl" refers to a monovalent aromatic heterocyclyl having at least one heteroatom in the ring. Thus, "heteroaryl" refers to an aromatic group of from 1 to 14 carbon atoms and 1 to 6 heteroatoms selected from oxygen, nitrogen, sulfur, or phosphorous. For multiple ring systems, by way of example, the term "heteroaryl" includes fused, bridged, and spiro ring systems having aromatic and non-aromatic rings. . In one embodiment, the carbon, nitrogen, or sulfur ring atom(s) of the heteroaryl group may be oxidized to provide for C(=O), N-oxide, sulfinyl, or sulfonyl moieties.

Examples of heteroaryls include by way of example and not limitation pyridyl, thiazolyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl, indolyl, quinoliny, isoquinoliny, benzimidazolyl, tetrahydroquinoliny, tetrahydroisoquinoliny, 6H-1,2,5-thiadiaziny, 2H,6H-1,5,2-dithiaziny, thienyl, thianthrenyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxathiny, 2H-pyrroly, isothiazolyl, isoxazolyl, pyraziny, pyridaziny, indoliziny, isoindolyl, 3H-indolyl, 1H-indazoly, puriny, 4H-quinoliziny, naphthyridiny, quinoxaliny, quinazoliny, cinnoliny, pteridiny, 4aH-carbazolyl, carbazolyl, β -carboliny, phenanthridiny, acridiny, pyrimidinyl, phenanthroliny, phenaziny, phenothiaziny, furazany, phenoxaziny, isochromany, chromany, imidazolidiny, imidazoliny, pyrazolidiny, pyrazoliny, piperaziny, indoliny, isoindoliny, quinuclidiny, morpholiny, oxazolidiny, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazoliny, and isatinoyl. "Heterocyclylene" refers to a heterocyclyl, as defined herein, derived by replacing a hydrogen atom from a carbon atom or heteroatom of a heterocyclyl, with an open valence. Similarly, "heteroarylene" refers to an aromatic heterocyclylene.

"Heterocyclylalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heterocyclyl radical (i.e., a heterocyclyl-alkylene- moiety). Typical heterocyclyl alkyl groups include, but are not limited to heterocyclyl- CH_2 -, 2-(heterocyclyl)ethan-1-yl, and the like, wherein the "heterocyclyl" portion includes any of the heterocyclyl groups described above, including those described in Principles of Modern Heterocyclic Chemistry. One skilled in the art will also understand that the heterocyclyl group can be attached to the alkyl portion of the heterocyclyl alkyl by means of a carbon-carbon bond or a carbon-heteroatom bond, with the proviso that the resulting group is chemically stable. The heterocyclylalkyl group comprises 2 to 20 carbon atoms and 1-6 heteroatoms, e.g., the alkyl portion of the heterocyclylalkyl

group comprises 1 to 6 carbon atoms and the heterocyclyl moiety comprises 1 to 14 carbon atoms. Examples of heterocyclylalkyls include by way of example and not limitation 5-membered sulfur, oxygen, phosphorus, and/or nitrogen containing heterocycles such as pyrrolidylmethyl, 2-tetrahydrofuranylethan-1-yl, and the like,
 5 6-membered sulfur, oxygen, and/or nitrogen containing heterocycles such as piperidinylmethyl, morpholinylmethyl, piperidinylethyl, tetrahydropyranylethyl, and the like.

"Heteroarylalkyl" refers to an alkyl group, as defined herein, in which a hydrogen atom has been replaced with a heteroaryl group as defined herein. Non-
 10 limiting examples of heteroaryl alkyl include -CH₂-pyridinyl, -CH₂-pyrrolyl, -CH₂-oxazolyl, -CH₂-indolyl, -CH₂-isoindolyl, -CH₂-purinyl, -CH₂-furanyl, -CH₂-thienyl, -CH₂-benzofuranyl, -CH₂-benzothiophenyl, -CH₂-carbazolyl, -CH₂-imidazolyl, -CH₂-thiazolyl, -CH₂-isoxazolyl, -CH₂-pyrazolyl, -CH₂-isothiazolyl, -CH₂-quinolyl, -CH₂-isoquinolyl, -CH₂-pyridazyl, -CH₂-pyrimidyl, -CH₂-pyrazyl, -CH(CH₃)-pyridinyl, -CH(CH₃)-pyrrolyl, -
 15 CH(CH₃)-oxazolyl, -CH(CH₃)-indolyl, -CH(CH₃)-isoindolyl, -CH(CH₃)-purinyl, -CH(CH₃)-furanyl, -CH(CH₃)-thienyl, -CH(CH₃)-benzofuranyl, -CH(CH₃)-benzothiophenyl, -CH(CH₃)-carbazolyl, -CH(CH₃)-imidazolyl, -CH(CH₃)-thiazolyl, -CH(CH₃)-isoxazolyl, -CH(CH₃)-pyrazolyl, -CH(CH₃)-isothiazolyl, -CH(CH₃)-quinolyl, -CH(CH₃)-isoquinolyl, -CH(CH₃)-pyridazyl, -CH(CH₃)-pyrimidyl, -CH(CH₃)-pyrazyl, and the like.

The term "heterocyclyloxy" represents a heterocyclyl group attached to the adjacent atom by an oxygen.

When there is a sulfur atom present, the sulfur atom can be at different
 25 oxidation levels, namely, S, SO, SO₂, or SO₃. All such oxidation levels are within the scope of the present invention.

When there is a phosphorous atom present, the phosphorous atom can be at different oxidation levels, namely, POR^aR^bR^c, PO₂R^aR^b, or PO₃R^aR^b, where R^a, R^b, and R^c each independently is chosen from H, C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl,
 30 C₆₋₁₄ aryl, 3-12 membered heterocycle, 3-18 membered heteroarylalkyl, C₆₋₁₈ arylalkyl, or two taken together (with or without oxygens) form a 5 to 10 membered heterocycle. All such oxidation levels are within the scope of the present invention.

The term "optionally substituted" in reference to a particular moiety of the compound of the Formulae of the invention, for example an "optionally substituted
 35 aryl group", refers to a moiety having none, one, or more substituents.

- The term "substituted" in reference to alkyl, alkylene, aryl, arylalkyl, alkoxy, heterocyclyl, heteroaryl, carbocyclyl, etc., for example, "substituted alkyl", "substituted alkylene", "substituted aryl", "substituted arylalkyl", "substituted heterocyclyl", and "substituted carbocyclyl" means alkyl, alkylene, aryl, arylalkyl, heterocyclyl, carbocyclyl respectively, in which one or more hydrogen atoms are each independently replaced with a non-hydrogen substituent. Divalent groups may also be similarly substituted. Unless otherwise indicated, typical substituents include, but are not limited to, -X, -R^b, -O⁻, =O, -OR^b, -SR^b, -S⁻, -NR^b₂, -N⁺R^b₃, =NR^b, -CX₃, -CN, -OCN, -SCN, -N=C=O, -NCS, -NO, -NO₂, =N₂, -N₃, -NHC(=O)R^b, -OC(=O)R^b, -NHC(=O)NR^b₂, -S(=O)₂⁻, -S(=O)₂OH, -S(=O)₂R^b, -OS(=O)₂OR^b, -S(=O)₂NR^b₂, -S(=O)R^b, -OP(=O)(OR^b)₂, -P(=O)(OR^b)₂, -P(=O)(O⁻)₂, -P(=O)(OH)₂, -P(O)(OR^b)(O⁻), -C(=O)R^b, -C(=O)X, -C(S)R^b, -C(O)OR^b, -C(O)O⁻, -C(S)OR^b, -C(O)SR^b, -C(S)SR^b, -C(O)NR^b₂, -C(S)NR^b₂, -C(=NR^b)NR^b₂, where each X is independently a halogen: F, Cl, Br, or I; and each R^b is independently H, alkyl, aryl, arylalkyl, a heterocycle, or a protecting group or prodrug moiety. Alkylene, alkenylene, and alkynylene groups may also be similarly substituted. Unless otherwise indicated, when the term "substituted" is used in conjunction with groups such as arylalkyl, which have two or more moieties capable of substitution, the substituents can be attached to the aryl moiety, the alkyl moiety, or both.
- Those skilled in the art will recognize that when moieties such as "alkyl", "aryl", "heterocyclyl", etc. are substituted with one or more substituents, they could alternatively be referred to as "alkylene", "arylene", "heterocyclylene", etc. moieties (i.e., indicating that at least one of the hydrogen atoms of the parent "alkyl", "aryl", "heterocyclyl" moieties has been replaced with the indicated substituent(s)). When moieties such as "alkyl", "aryl", "heterocyclyl", etc. are referred to herein as "substituted" or are shown diagrammatically to be substituted (or optionally substituted, e.g., when the number of substituents ranges from zero to a positive integer), then the terms "alkyl", "aryl", "heterocyclyl", etc. are understood to be interchangeable with "alkylene", "arylene", "heterocyclylene", etc.
- As will be appreciated by those skilled in the art, the compounds of the present invention may exist in solvated or hydrated form. The scope of the present invention includes such forms. Again, as will be appreciated by those skilled in the art, the compounds may be capable of esterification. The scope of the present invention includes esters and other physiologically functional derivatives. The scope of the present invention includes prodrug forms of the compound herein described.

“Ester” means any ester of a compound in which any of the --COOH functions of the molecule is replaced by a --C(O)OR function, or in which any of the --OH functions of the molecule are replaced with a --OC(O)R function, in which the R moiety of the ester is any carbon-containing group which forms a stable ester moiety, including but not limited to alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl and substituted derivatives thereof.

The term “prodrug” as used herein refers to any compound that when administered to a biological system generates the drug substance, i.e., active ingredient, as a result of spontaneous chemical reaction(s), enzyme catalyzed chemical reaction(s), photolysis, and/or metabolic chemical reaction(s). A prodrug is thus a covalently modified analog or latent form of a therapeutically active compound. Example of prodrugs include ester moieties, quaternary ammonium moieties, glycol moieties, and the like.

One skilled in the art will recognize that substituents and other moieties of the compounds of Formula I should be selected in order to provide a compound which is sufficiently stable to provide a pharmaceutically useful compound which can be formulated into an acceptably stable pharmaceutical composition. Compounds of Formula I which have such stability are contemplated as falling within the scope of the present invention.

As will be appreciated by those skilled in the art, the compounds of the present invention may contain one or more chiral centers. The scope of the present invention includes such forms. Again, as will be appreciated by those skilled in the art, the compound is capable of esterification. The scope of the present invention includes esters and other physiologically functional derivatives. In addition, the scope of the present invention includes prodrug forms of the compound herein described.

A compound of Formula I-II and its pharmaceutically acceptable salts may exist as different polymorphs or pseudopolymorphs. As used herein, crystalline polymorphism means the ability of a crystalline compound to exist in different crystal structures. Polymorphism generally can occur as a response to changes in temperature, pressure, or both. Polymorphism can also result from variations in the crystallization process. Polymorphs can be distinguished by various physical characteristics known in the art such as x-ray diffraction patterns, solubility, and melting point. The crystalline polymorphism may result from differences in crystal packing (packing polymorphism) or differences in packing between different

conformers of the same molecule (conformational polymorphism). As used herein, crystalline pseudopolymorphism means the ability of a hydrate or solvate of a compound to exist in different crystal structures. The pseudopolymorphs of the instant invention may exist due to differences in crystal packing (packing
5 pseudopolymorphism) or due to differences in packing between different conformers of the same molecule (conformational pseudopolymorphism). The instant invention comprises all polymorphs and pseudopolymorphs of the compounds of Formula I-II and their pharmaceutically acceptable salts.

10 A compound of Formula I-II and its pharmaceutically acceptable salts may also exist as an amorphous solid. As used herein, an amorphous solid is a solid in which there is no long-range order of the positions of the atoms in the solid. This definition applies as well when the crystal size is two nanometers or less. Additives, including solvents, may be used to create the amorphous forms of the instant invention. The instant invention comprises all amorphous forms of the compounds of
15 Formula I-II and their pharmaceutically acceptable salts.

Certain of the compounds described herein contain one or more chiral centers, or may otherwise be capable of existing as multiple stereoisomers. The scope of the present invention includes mixtures of stereoisomers as well as purified enantiomers or enantiomerically/diastereomerically enriched mixtures. Also included
20 within the scope of the invention are the individual isomers of the compounds represented by the formulae of the present invention, as well as any wholly or partially equilibrated mixtures thereof. The present invention also includes the individual isomers of the compounds represented by the formulas above as mixtures with isomers thereof in which one or more chiral centers are inverted.


25 The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to molecules which are superimposable on their mirror image partner.

The term "stereoisomers" refers to compounds which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space.


30 "Diastereomer" refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another. Diastereomers have different physical properties, e.g., melting points, boiling points, spectral properties, and reactivities. Mixtures of diastereomers may separate under high resolution analytical procedures such as electrophoresis and chromatography.

“Enantiomers” refer to stereoisomers of a compound which are non-superimposable mirror images of one another.

“Atropisomers” refer to stereoisomers of a compound resulting from hindered rotation about single bonds where the steric strain barrier to rotation is high enough to allow for the isolation of the individual conformer. Atropisomers display axial
 5 chirality. Atropisomers may be equilibrated thermally and the interconversion barrier may be measured kinetically. Atropisomerism may occur apart from the presence of other forms of chiral isomerism. Thus, as illustrated, the depicted nitrogen atom is planar and compounds of Formula I are capable of existing as atropisomers:



In one embodiment of the present invention, the compounds exist in a conformeric form of Formula Ia:



Stereochemical definitions and conventions used herein generally follow S. P.
 15 Parker, Ed., McGraw-Hill Dictionary of Chemical Terms (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., Stereochemistry of Organic Compounds (1994) John Wiley & Sons, Inc., New York.

Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically active
 20 compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with (-) or l meaning that the compound is levorotatory. A compound prefixed with (+) or d is dextrorotatory.

25 A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of

enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species, devoid of optical activity.

5 The present invention includes a salt or solvate of the compounds herein described, including combinations thereof such as a solvate of a salt. The compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms, and the present invention encompasses all such forms.


Typically, but not absolutely, the salts of the present invention are
10 pharmaceutically acceptable salts. Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention.

Examples of suitable pharmaceutically acceptable salts include inorganic acid addition salts such as chloride, bromide, sulfate, phosphate, and nitrate; organic acid
15 addition salts such as acetate, galactarate, propionate, succinate, lactate, glycolate, malate, tartrate, citrate, maleate, fumarate, methanesulfonate, p-toluenesulfonate, and ascorbate; salts with acidic amino acid such as aspartate and glutamate; alkali metal salts such as sodium salt and potassium salt; alkaline earth metal salts such as magnesium salt and calcium salt; ammonium salt; organic basic salts such as
20 trimethylamine salt, triethylamine salt, pyridine salt, picoline salt, dicyclohexylamine salt, and N,N'-dibenzylethylenediamine salt; and salts with basic amino acid such as lysine salt and arginine salt. The salts may be in some cases hydrates or ethanol solvates.

The modifier "about" used in connection with a quantity is inclusive of the
25 stated value and has the meaning dictated by the context (e.g., includes the degree of error associated with measurement of the particular quantity).

Whenever a compound described herein is substituted with more than one of the same designated group, e.g., "R" or "R¹", then it will be understood that the groups may be the same or different, i.e., each group is independently selected.

30 Wavy lines,



indicate the site of covalent bond attachments to the adjoining substructures, groups, moieties, or atoms.

The compounds of the invention can also exist as tautomeric isomers in certain cases. Although only one delocalized resonance structure may be depicted, all such forms are contemplated within the scope of the invention. For example, enamine tautomers can exist for purine, pyrimidine, imidazole, guanidine, amidine, and tetrazole systems and all their possible tautomeric forms are within the scope of the invention.

Selected substituents comprising the compounds of Formula I-II may be present to a recursive degree. In this context, "recursive substituent" means that a substituent may recite another instance of itself. The multiple recitations may be direct or indirect through a sequence of other substituents. Because of the recursive nature of such substituents, theoretically, a large number of compounds may be present in any given embodiment. One of ordinary skill in the art of medicinal chemistry understands that the total number of such substituents is reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such as ease of synthesis. Recursive substituents may be an intended aspect of the invention. One of ordinary skill in the art of medicinal chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in an embodiment of the invention, they may recite another instance of themselves, 0, 1, 2, 3, or 4 times.

The compounds of Formula I-II also include molecules that incorporate isotopes of the atoms specified in the particular molecules. Non-limiting examples of these isotopes include D, T, ^{14}C , ^{13}C , ^{18}O and ^{15}N .

25 Protecting Groups

In the context of the present invention, protecting groups include prodrug moieties and chemical protecting groups.

Protecting groups are available, commonly known and used, and are optionally used to prevent side reactions with the protected group during synthetic procedures, i.e. routes or methods to prepare the compounds of the invention. For the most part the decision as to which groups to protect, when to do so, and the nature of the chemical protecting group "PG" will be dependent upon the chemistry of the reaction to be protected against (e.g., acidic, basic, oxidative, reductive or other conditions) and the intended direction of the synthesis. The PG groups do not need


to be, and generally are not, the same if the compound is substituted with multiple PG. In general, PG will be used to protect functional groups such as carboxyl, hydroxyl, thio, or amino groups and to thus prevent side reactions or to otherwise facilitate the synthetic efficiency. The order of deprotection to yield free, deprotected groups is dependent upon the intended direction of the synthesis and the reaction conditions to be encountered, and may occur in any order as determined by the artisan.

Various functional groups of the compounds of the invention may be protected. For example, protecting groups for -OH groups (whether hydroxyl, carboxylic acid, phosphonic acid, or other functions) include "ether- or ester-forming groups". Ether- or ester-forming groups are capable of functioning as chemical protecting groups in the synthetic schemes set forth herein. However, some hydroxyl and thio protecting groups are neither ether- nor ester-forming groups, as will be understood by those skilled in the art, and are included with amides, discussed below.

A very large number of hydroxyl protecting groups and amide-forming groups and corresponding chemical cleavage reactions are described in Protective Groups in Organic Synthesis, Theodora W. Greene and Peter G. M. Wuts (John Wiley & Sons, Inc., New York, 1999, ISBN 0-471-16019-9) ("Greene"). See also Kocienski, Philip J.; Protecting Groups (Georg Thieme Verlag Stuttgart, New York, 1994). In particular Chapter 1, Protecting Groups: An Overview, pages 1-20, Chapter 2, Hydroxyl Protecting Groups, pages 21-94, Chapter 3, Diol Protecting Groups, pages 95-117, Chapter 4, Carboxyl Protecting Groups, pages 118-154, Chapter 5, Carbonyl Protecting Groups, pages 155-184. For protecting groups for carboxylic acid, phosphonic acid, phosphonate, sulfonic acid and other protecting groups for acids see Greene as set forth below. Such groups include by way of example and not limitation, esters, amides, hydrazides, and the like.

Ether- and Ester-forming protecting groups

5 Ester-forming groups include: (1) phosphonate ester-forming groups, such as phosphoramidate esters, phosphorothioate esters, phosphonate esters, and phosphon-bis-amidates; (2) carboxyl ester-forming groups, and (3) sulphur ester-forming groups, such as sulphonate, sulfate, and sulfinde.

Metabolites of the Compounds of the Invention

50a

Also falling within the scope of this invention are the in vivo metabolic products of the compounds described herein. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabelled (e.g., C^{14} or ^3H) compound of the invention, administering it parenterally in a detectable dose (e.g., greater than about 0.5 mg/kg) to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur (typically about 30 seconds to 30 hours) and isolating its conversion products from the urine, blood or other biological samples. These products are easily isolated since they are labeled (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, e.g., by MS or NMR analysis. In general, analysis of metabolites is done in the same way as conventional drug metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found in vivo, are useful in diagnostic assays for therapeutic dosing of the compounds of the invention even if they possess no anti-infective activity of their own.

The definitions and substituents for various genus and subgenus of the present compounds are described and illustrated herein. It should be understood by one skilled in the art that any combination of the definitions and substituents described above should not result in an inoperable species or compound. "Inoperable species or compounds" means compound structures that violates relevant scientific principles (such as, for example, a carbon atom connecting to more than four covalent bonds) or compounds too unstable to permit isolation and formulation into pharmaceutically acceptable dosage forms.

Pharmaceutical Formulations

The compounds of this invention are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain

excipients such as those set forth in the Handbook of Pharmaceutical Excipients (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such as EDTA, carbohydrates such as dextrin, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.


While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations of the invention, both for veterinary and for human use, comprise at least one active ingredient, together with one or more acceptable carriers and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

The formulations include those suitable for the foregoing administration routes. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, Pa.). Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water

liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

5 A tablet is made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid



diluent. The tablets may optionally be coated or scored and optionally are formulated so as to provide slow or controlled release of the active ingredient.

For administration to the eye or other external tissues e.g., mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active ingredient(s) in a range between 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

Emulgents and emulsion stabilizers suitable for use in the formulation of the invention include Tween® 60, Span® 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl mono-stearate and sodium lauryl sulfate.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties. The cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids,

isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as
5 white soft paraffin and/or liquid paraffin or other mineral oils are used.

Pharmaceutical formulations according to the present invention comprise one or more compounds of the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. Pharmaceutical formulations containing the active ingredient may be in any form
10 suitable for the intended method of administration. When used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions
15 may contain one or more agents including sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium
20 carbonate, lactose, lactose monohydrate, croscarmellose sodium, povidone, calcium or sodium phosphate; granulating and disintegrating agents, such as maize starch, or alginic acid; binding agents, such as cellulose, microcrystalline cellulose, starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques including
25 microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

Formulations for oral use may be also presented as hard gelatin capsules
30 where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions of the invention contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions.
35 Such excipients include a suspending agent, such as sodium carboxymethylcellulose,

methycellulose, hydroxypropyl methycellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of
5 ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring
10 agents, one or more flavoring agents and one or more sweetening agents, such as sucrose or saccharin.

Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oral suspensions may contain a thickening agent,
15 such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth herein, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules of the invention suitable for preparation of
20 an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

25 The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth, naturally occurring phosphatides, such as soybean lecithin, esters or
30 partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan monooleate, and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may
35 also contain a demulcent, a preservative, a flavoring or a coloring agent.

The pharmaceutical compositions of the invention may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been
5 mentioned herein. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may
10 conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The amount of active ingredient that may be combined with the carrier
15 material to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a time-release formulation intended for oral administration to humans may contain approximately 1 to 1000 mg of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions
20 (weight:weight). The pharmaceutical composition can be prepared to provide easily measurable amounts for administration. For example, an aqueous solution intended for intravenous infusion may contain from about 3 to 500 µg of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur.

25 Formulations suitable for administration to the eye include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent for the active ingredient. The active ingredient is preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% particularly about 1.5% w/w.

30 Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

Formulations suitable for intrapulmonary or nasal administration have a particle size for example in the range of 0.1 to 500 μm (including particle sizes in a range between 0.1 and 500 μm in increments such as 0.5 μm , 1 μm , 30 μm , 35 μm , etc.), which is administered by rapid inhalation through the nasal passage or by inhalation through the mouth so as to reach the alveolar sacs. Suitable formulations include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol or dry powder administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as compounds heretofore used in the treatment or prophylaxis of infections as described herein.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

The formulations are presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

Compounds of the invention can also be formulated to provide controlled release of the active ingredient to allow less frequent dosing or to improve the pharmacokinetic or toxicity profile of the active ingredient. Accordingly, the invention

also provided compositions comprising one or more compounds of the invention formulated for sustained or controlled release.

The effective dose of an active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician using conventional dose escalation studies. The effective dose can be expected to be from about 0.0001 to about 100 mg/kg body weight per day; typically, from about 0.01 to about 10 mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For example, the daily candidate dose for an adult human of approximately 70 kg body weight will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and may take the form of single or multiple doses.

In yet another embodiment, the present application discloses pharmaceutical compositions comprising a compound of Formula I or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

Routes of Administration

One or more compounds of the invention (herein referred to as the active ingredients) are administered by any route appropriate to the condition to be treated. Suitable routes include oral, rectal, nasal, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural), and the like. It will be appreciated that the preferred route may vary with for example the condition of the recipient. An advantage of the compounds of this invention is that they are orally bioavailable and can be dosed orally.

Combination Therapy, Including HCV Combination Therapy

In another embodiment, the compounds of the present invention may be combined with one or more active agent. Non-limiting examples of suitable combinations include combinations of one or more compounds of the present invention with one or more interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B

polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, pharmacokinetic enhancers, and other drugs for treating HCV.

More specifically, one or more compounds of the present invention may be combined with one or more compounds selected from the group consisting of

- 5 1) interferons, *e.g.*, pegylated rIFN-alpha 2b (PEG-Intron), pegylated rIFN-alpha 2a (Pegasys), rIFN-alpha 2b (Intron A), rIFN-alpha 2a (Roferon-A), interferon alpha (MOR-22, OPC-18, Alfaferone, Alfaferone, Multiferon, subalin), interferon alfacon-1 (Infergen), interferon alpha-n1 (Wellferon), interferon alpha-n3 (Alferon), interferon-beta (Avonex, DL-8234), interferon-omega (omega DUROS, Biomed 510),
10 albinterferon alpha-2b (Albupharm), IFN alpha XL, BLX-883 (Locteron), DA-3021, glycosylated interferon alpha-2b (AVI-005), PEG-Infergen, PEGylated interferon lambda (PEGylated IL-29), and belerofon,
- 2) ribavirin and its analogs, *e.g.*, ribavirin (Rebetol, Copegus), and taribavirin (Viramidine),
- 15 3) HCV NS3 protease inhibitors, *e.g.*, boceprevir (SCH-503034, SCH-7), telaprevir (VX-950), VX-813, TMC-435 (TMC435350), ABT-450, BI-201335, BI-1230, MK-7009, SCH-900518, VBY-376, VX-500, GS-9256, GS-9451, BMS-790052, BMS-605339, PHX-1766, AS-101, YH-5258, YH5530, YH5531, and ITMN-191 (R-7227),
- 4) alpha-glucosidase 1 inhibitors, *e.g.*, celgosivir (MX-3253), Miglitol, and UT-
20 231B,
- 5) hepatoprotectants, *e.g.*, emericasan (IDN-6556), ME-3738, GS-9450 (LB-84451), silibinin, and MitoQ,
- 6) nucleoside or nucleotide inhibitors of HCV NS5B polymerase, *e.g.*, R1626, R7128 (R4048), IDX184, IDX-102, PSI-7851, BCX-4678, valopicitabine (NM-283),
25 and MK-0608,
- 7) non-nucleoside inhibitors of HCV NS5B polymerase, *e.g.*, filibuvir (PF-868554), ABT-333, ABT-072, BI-207127, VCH-759, VCH-916, JTK-652, MK-3281, VBY-708, VCH-222, A848837, ANA-598, GL60667, GL59728, A-63890, A-48773, A-48547, BC-2329, VCH-796 (nesbuvir), GSK625433, BILN-1941, XTL-2125, and GS-
30 9190,
- 8) HCV NS5A inhibitors, *e.g.*, AZD-2836 (A-831), AZD-7295 (A-689), and BMS-790052,
- 9) TLR-7 agonists, *e.g.*, imiquimod, 852A, GS-9524, ANA-773, ANA-975, AZD-8848 (DSP-3025), PF-04878691, and SM-360320,
- 35 10) cyclophilin inhibitors, *e.g.*, DEBIO-025, SCY-635, and NIM811,

- 11) HCV IRES inhibitors, e.g., MCI-067,
- 12) pharmacokinetic enhancers, e.g., BAS-100, SPI-452, PF-4194477, TMC-41629, GS-9350, GS-9585, and roxythromycin,
- 13) other drugs for treating HCV, e.g., thymosin alpha 1 (Zadaxin),
- 5 nitazoxanide (Alinia, NTZ), BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), GS-9525, KRN-7000, civacir, GI-5005, XTL-6865, BIT225, PTX-111, ITX2865, TT-033i, ANA 971, NOV-205, tarvacin, EHC-18, VGX-410C, EMZ-702, AVI 4065, BMS-650032, BMS-791325, Bavituximab, MDX-1106 (ONO-4538), Oglufanide, FK-788, and VX-497 (merimepodib);
- 10 14) mevalonate decarboxylase antagonists, e.g., statins, HMGCoA synthase inhibitors (e.g., hymeglusil), squalene synthesis inhibitors (e.g., zaragozic acid);
- 15 15) angiotensin II receptor antagonists, e.g., losartan, irbesartan, olmesartan, candesartan, valsartan, telmisartan, eprosartan;
- 16) angiotensin-converting enzyme inhibitors, e.g., captopril, zofenopril,
- 15 enalapril, ramipril, quinapril, perindopril, lisinopril, benazepril, fosinopril;
- 17) other anti-fibrotic agents, e.g., amiloride and
- 18) endothelin antagonists, e.g. bosentan and ambrisentan.

In yet another embodiment, the present application discloses pharmaceutical compositions comprising a compound of the present invention, or a pharmaceutically acceptable salt thereof, in combination with at least one additional active agent, and a pharmaceutically acceptable carrier or excipient. In yet another embodiment, the present application provides a combination pharmaceutical agent with two or more therapeutic agents in a unitary dosage form. Thus, it is also possible to combine any compound of the invention with one or more other active agents in a unitary dosage form.

The combination therapy may be administered as a simultaneous or sequential regimen. When administered sequentially, the combination may be administered in two or more administrations.

Co-administration of a compound of the invention with one or more other active agents generally refers to simultaneous or sequential administration of a compound of the invention and one or more other active agents, such that therapeutically effective amounts of the compound of the invention and one or more other active agents are both present in the body of the patient.

Co-administration includes administration of unit dosages of the compounds of the invention before or after administration of unit dosages of one or more other

active agents, for example, administration of the compounds of the invention within seconds, minutes, or hours of the administration of one or more other active agents. For example, a unit dose of a compound of the invention can be administered first, followed within seconds or minutes by administration of a unit dose of one or more
5 other active agents. Alternatively, a unit dose of one or more other active agents can be administered first, followed by administration of a unit dose of a compound of the invention within seconds or minutes. In some cases, it may be desirable to administer a unit dose of a compound of the invention first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of one or more other active
10 agents. In other cases, it may be desirable to administer a unit dose of one or more other active agents first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of a compound of the invention.

The combination therapy may provide "synergy" and "synergistic effect", i.e. the effect achieved when the active ingredients used together is greater than the sum
15 of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined formulation; (2) delivered by alternation or in parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are
20 administered or delivered sequentially, e.g., in separate tablets, pills or capsules, or by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together.

25 As will be appreciated by those skilled in the art, when treating a viral infection such as HCV, such treatment may be characterized in a variety of ways and measured by a variety of endpoints. The scope of the present invention is intended to encompass all such characterizations.

30 **Synthetic Examples**

Certain abbreviations and acronyms are used in describing the experimental details. Although most of these would be understood by one skilled in the art, Table 1 contains a list of many of these abbreviations and acronyms.

Table 1. List of abbreviations and acronyms.

Abbreviation	Meaning
Ac	acetyl
ACN	acetonitrile
AIBN	2,2'-azobis(2-methylpropionitrile)
BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
Bn	benzyl
BnBr	benzylbromide
BSA	bis(trimethylsilyl)acetamide
BzCl	benzoyl chloride
CDI	carbonyl diimidazole
DABCO	1,4-diazabicyclo[2.2.2]octane
dba	dibenzylideneacetone
DBN	1,5-diazabicyclo[4.3.0]non-5-ene
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DBU	1,5-diazabicyclo[5.4.0]undec-5-ene
DCA	dichloroacetamide
DCC	dicyclohexylcarbodiimide
DCE	1,2-dichloroethane
DCM	dichloromethane
deg	degrees
DIAD	di-isopropylazodicarboxylate
DIEA	<i>N,N</i> -di-isopropylethylamine
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMTCI	dimethoxytrityl chloride
DMSO	dimethylsulfoxide
DMTr	4, 4'-dimethoxytrityl
DMF	dimethylformamide
EtOAc	ethyl acetate
ES, ESI	electrospray ionization
HMDS	hexamethyldisilazane
HPLC	high pressure liquid chromatography


LC	liquid chromatography
LDA	lithium diisopropylamide
LRMS	low resolution mass spectrum
MCPBA	meta-chloroperoxybenzoic acid
MeCN	acetonitrile
MeOH	methanol
MMTC	mono methoxytrityl chloride
m/z or m/e	mass to charge ratio
MH ⁺	mass plus 1
MH ⁻	mass minus 1
MsOH	methanesulfonic acid
MS or ms	mass spectrum
NBS	N-bromosuccinimide
NMP	N-methylpyrrolidine
Ph	phenyl
rt or r.t.	room temperature
TBAF	tetrabutylammonium fluoride
TES	triethylsilyl
THF	tetrahydrofuran
THP	tetrahydropyran
TMSCl	chlorotrimethylsilane
TMSBr	bromotrimethylsilane
TMSI	iodotrimethylsilane
TMSOTf	(trimethylsilyl)trifluoromethylsulfonate
TEA	triethylamine
TBA	tributylamine
TBAP	tributylammonium pyrophosphate
TBSCl	t-butyldimethylsilyl chloride
TEAB	triethylammonium bicarbonate
TFA	trifluoroacetic acid
TLC or tlc	thin layer chromatography
Tr	triphenylmethyl
Tol	4-methylbenzoyl

Turbo Grignard	1:1 mixture of isopropylmagnesium chloride and lithium chloride
xantphos	4,5-bis(diphenylphosphino)-9,9-dimethylxanthene
δ	parts per million down field from tetramethylsilane

General Schemes


The compounds of this invention may be synthesized by several routes with key bond-forming steps as indicated in Schemes A-C, in which the carboxylate substituent R indicates either a protecting group such as an alkyl ester (where necessary), or the free acid itself. Alkyl ester protecting groups are conveniently removed by saponification with an alkali metal hydroxide in a protic solvent such as water or an alcohol, and may be facilitated by use of ethereal solvent mixtures and/or heating. Alternatively they may be removed by dealkylation through heating with an alkali metal halide in an aprotic solvent. As will be appreciated, substituents on Het may be modified subsequent to other bond-forming steps by, for example, N-oxidation with a typical oxidant such as metachloroperbenzoic acid in a solvent such as dichloromethane, O-dealkylation through treatment with a reagent such as boron tribromide in a solvent such as dichloromethane, or hydrolysis.

Scheme A



The bond between L and Het may be formed by displacement of X on Het, where X is a leaving group such as a halide, sulfinate, sulfonate or phosphate moiety. The reaction is conveniently performed by deprotonation of L-H with a base such as sodium hydride or potassium hexamethyldisilazide, or is facilitated by the presence of a tertiary amine; it can be carried out in a variety of solvents such as THF, dioxane, dichloromethane, NMP, DMF or DMSO and may be accelerated by heating.


Scheme B



The bond between R^3 and L may be formed by nucleophilic displacement of a leaving group X on R^3 . The leaving group may vary widely and includes, but is not limited to, halide, carboxylate, sulfinate, sulfonate or phosphate moieties, and it may be generated from the corresponding alcohol in situ through treatment with reagents such as dialkyl azodicarboxylates. The reaction may also be facilitated by deprotonation of $Het-L-H$ with a base such as sodium hydride or potassium hexamethyldisilazide, or is facilitated by the presence of a tertiary amine; it can be carried out in a variety of solvents such as THF, dioxane, dichloromethane, NMP, DMF or DMSO and may be accelerated by heating.

Scheme C

15




The starting material in Scheme A may be synthesized as depicted in Scheme C. Substituted 3-aminothiophenes II may be generated by reductive amination of $Y-R^3-L-R$ (where Y indicates an aldehyde or ketone and R and R^a depict

optional protecting groups), or by direct alkylation (where Y indicates a leaving group such as a halide, sulfinate, sulfonate or phosphate moiety) of the 3-aminothiophene I (see patent application WO2008/58393). In the latter case the alkylation may be facilitated by deprotonation of the amine with a base such as sodium hydride or potassium hexamethyldisilazide, and can be carried out in a variety of solvents such as THF, dioxane, dichloromethane, NMP, DMF or DMSO and may be accelerated by heating. In cases where R³ is aromatic, the reaction may be catalyzed by Pd (*J. Org. Chem.*, **2000**, 65, 1158-1174). Alternatively II may be generated by coupling of an amine with a 3-iodothiophene IV catalyzed by Pd (*J. Org. Chem.*, **2000**, 65, 1158-1174). The amine II is converted to the amide III by acylation with a carboxylic acid derivative such as an acyl chloride or anhydride in the presence of a base such as pyridine or a tertiary amine in an inert solvent such as dichloromethane. Alternatively IV may be converted to III directly by amidation catalyzed by Cu (*J. Am. Chem. Soc.*, **2002**, 124, 7421-7428).

The starting material for Scheme B may be generated in an analogous fashion, with the leaving group X being generated in a final step by standard methods from the precursor alcohol.

The synthesis of iodothiophene IV is illustrated below for the case where R¹ = tBu, and other variants may be synthesized in analogous fashion:

20




To a solution of 5-(3,3-Dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid (6.2 g, 30 mmol; see patent application US5861421) in THF (100 mL) was added a solution of nBuLi (2.0 M in pentane, 33 mL, 66 mmol) via an addition funnel at -78 °C. After addition, the reaction was stirred at -78 °C for 1 h. A solution of I₂ (7.7 g, 30 mmol) in THF (100 mL) was added slowly (ca. 15 min) to the flask. After a further 10 mins, the reaction was quenched with 1 N HCl (50 mL) and warmed to room temperature. The volatiles were removed in vacuo and the residue was dissolved in ether (500 mL).

The organic solution was washed with 1 M $\text{Na}_2\text{S}_2\text{O}_3$ (100 mL x 2), brine (100 mL) and dried over Na_2SO_4 . After concentrated in vacuo, the residue was purified by silica gel chromatography (EtOAc/hexanes) to give 5-(3,3-Dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid (5.9 g, 65%) as a white solid.

5

Scheme E



10

To a solution of 5-(3,3-Dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid (1.0 g, 3.0 mmol) and DMF (20 μL) in dry dichloromethane (10 mL) was added oxalyl chloride (508 μL , 6.0 mmol) at room temperature. After stirring at room temperature for 90 min, the reaction was concentrated in vacuo to remove volatiles.


15 The residue was dissolved in pyridine (5 mL) and methanol (5 mL) and stirred for 2 h. The volatiles were removed in vacuo and the residue was participated between ether (150 mL) and saturated NH_4Cl solution (50 mL). The organic layer was washed with saturated NH_4Cl solution (50 mL) and dried over Na_2SO_4 . After concentration in vacuo, the residue was purified by silica gel chromatography (EtOAc/hexanes) to give the desired product (835 mg, 80%).


20

Experimentals

Example 31: Compound 31 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester

25





- 5 4-[[5-(3,3-dimethyl-but-1-ynyl)-2-methoxycarbonyl-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester (1.5g, 2.73mmol) was dissolved in ACN (10mL). To the solution was added a solution of lithium hydroxide (253mg, 11.01mmol) in water (10mL). The reaction was stirred at room temperature for 6 hours. The reaction was complete as determined
- 10 by LC/MS. The pH was adjusted to 5 with 1N HCl in water. The product was extracted with ethyl acetate (3x10mL). The combined organics were dried with sodium sulfate, filtered and were concentrated under reduced pressure and 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester (1.35g, 93%) was recovered as a white solid.
- 15

4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tert-butyl ester (1.35g,

2.53mmol) was dissolved in 4N HCl in dioxane (6mL, 24mmol). The reaction was stirred at room temperature for 0.5 hours and found to be complete as determined by LC/MS. The reaction was concentrated under reduced pressure.

5 The HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid (100mg, 0.2mmol) was suspended in ACN (1mL) and a saturated aqueous solution of NaHCO₃ (1mL). After 15 minutes, 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester chloroformate (1.0mmol) was added as a solution in THF (1mL). The reaction was stirred at room temperature for 1 hour. The reaction was found to be complete
10 by LC/MS. Boc protected 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester was purified by HPLC to afford a white solid (76mg, 53%).
LC/MS (m/z): 658 [M+1], 558 [M-99]
Retention time: 2.64min

15 LC: Thermo Finnigan PDA Detector
MS: Thermo Scientific LCQ Fleet
Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm
Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid


20 Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

Boc protected 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester (35mg, 0.05mmol) was dissolved in 4N HCl in dioxane (2mL, 8.0mmol). The reaction was stirred at room temperature for 1 hour. The reaction was complete as
25 determined by LC/MS. The reaction was concentrated under reduced pressure. The HCl salt of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester (31mg, 94%) was found as a white solid.

LC/MS (m/z): 558 [M+1]
30 Retention time: 2.14min
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Synthesis of 4-Hydroxy-piperidine-1-carboxylic acid tert-butyl ester chloroformate


35



To a solution of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester (201mg, 1.0mmol) in THF (2mL) was added a 20% solution of phosgene in toluene (358μL, 1.7mmol). The reaction was stirred at room temperature for 1 hour. The reaction was concentrated under reduced pressure.


Method B

10 Example 32: Compound 32 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-pyran-4-yloxy)carbonylamino]-cyclohexyl]-amino)-thiophene-2-carboxylic acid



15

Scheme 7



5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (5.00g, 10.93mmol) and ammonium acetate (8.42g, 109.30mmol) were dissolved in MeOH (100mL) under an atmosphere of nitrogen. To the reaction was added 4Å molecular sieves, powdered (500mg). After 30 min, sodium triacetoxyborohydride (3.46g, 16.40mmol) was added in 4 portions. The reaction was stirred for 18h until found complete by LC/MS. The reaction was filtered through a pad of celite™ followed by a MeOH wash and was concentrated. 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (3.35g, 67%) was purified by HPLC to afford a white solid.

3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (100mg, 0.22mmol) was suspended in ACN (1 mL) and a saturated aqueous solution of NaHCO₃ (1 mL). After 15 minutes, tetrahydro-pyran-4-ol chloroformate (0.65mmol) as prepared in a similar fashion to 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester chloroformate except that tetrahydro-pyran-4-ol was used instead of hydroxy-piperidine-1-carboxylic acid tert-butyl ester was added as a solution in THF (1 mL). The reaction was stirred at room temperature for 1.5 hours. The reaction was complete by LC/MS. 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-pyran-4-yloxy-carbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid methyl ester was extracted with EtOAc (3x5mL). The combined organics were dried with sodium sulfate, filtered and were concentrated under reduced pressure.


5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-pyran-4-yloxy-carbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid methyl ester was dissolved in ACN (1mL). To the reaction was added a solution of lithium hydroxide (25mg, 1.1mmol) in water (1mL). The reaction was stirred at room temperature for 2.5 hours. The reaction was complete by LC/MS. 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-pyran-4-yloxy-carbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid (50mg, 40% over 2 steps) was purified by HPLC to afford a white solid.

LC/MS (m/z): 448 [M-124]

Retention time: 2.46min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Example 33: Compound 33 - Synthesis of 3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(S)-yl ester



33

5

3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(S)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(S)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 4-(S)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

10

LC/MS (m/z): 544 [M+1]

Retention time: 1.98min

LC: Thermo Finnigan PDA Detector

MS: Thermo Scientific LCQ Fleet

20

Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN


25

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(S)-yl-amino]-thiophene-2-carboxylic acid was synthesized as follows:

a) Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid methyl ester

30

Scheme 8



A mixture of 5-Bromo-thiophene-2-carboxylic acid ethyl ester (7g, 30 mmol),
 5 copper iodide (1.2g, 6 mmol), triethylamine (20 mL) in DMF (100 mL) was degassed
 in a 350mL pressure bottle. Then tris(dibenzylideneacetone)dipalladium(0) (2.1 g, 3
 mmol) and 3,3-dimethyl-but-1-yne (18.3 mL, 150 mmol) were added and heated at
 80 degree for 3 hours. The reaction mixture was filtered on celite and washed with
 ethyl acetate. The solution was diluted with water and extracted twice with ethyl
 10 acetate. The organic phases were combined and washed with water. After drying and
 concentration, the crude residue was purified by flash chromatography to yield 6.9g
 (95%) of 5-(3,3-Dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid ethyl ester as a
 yellow oil.

A solution of 5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid ethyl
 15 ester (6.9g) in THF (100 mL) was added LiOH (1.5N, 100 mL). The mixture was
 stirred at room temperature for 4 hours. Acidified reaction with HCl to pH =2, then
 remove volatiles under vacuo. The resulting beige color solid was collected by
 filtration, washed with water then dried overnight to give 6.2 g of product which was
 used without further purification.

20 To a solution of 5-(3,3-Dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid (6.2 g,
 30 mmol; see patent application US5861421) in THF (100 mL) was added a solution
 of nBuLi (2.0 M in pentane, 33 mL, 66 mmol) via an addition funnel at -78 °C. After
 addition, the reaction was stirred at -78 °C for 1 h. A solution of I₂ (7.7 g, 30 mmol) in
 THF (100 mL) was added slowly (ca. 15 min) to the flask. After a further 10 mins, the
 25 reaction was quenched with 1 N HCl (50 mL) and warmed to room temperature. The
 volatiles were removed in vacuo and the residue was dissolved in ether (500 mL).

The organic solution was washed with 1 M $\text{Na}_2\text{S}_2\text{O}_3$ (100 mL x 2), brine (100 mL) and dried over Na_2SO_4 . After concentrated in vacuo, the residue was purified by silica gel chromatography (EtOAc/hexanes) to give 5-(3,3-Dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid (5.9 g, 65%) as a white solid.


5 To a solution of 5-(3,3-dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid (1.0 g, 3.0 mmol) and DMF (20 μL) in dry dichloromethane (10 mL) was added oxalyl chloride (508 μL , 6.0 mmol) at room temperature. After stirring at room temperature for 90 min, the reaction was concentrated in vacuo to remove volatiles. The residue was dissolved in pyridine (5 mL) and methanol (5 mL) and stirred for 2 h. The
10 volatiles were removed in vacuo and the residue was participated between ether (150 mL) and saturated NH_4Cl solution (50 mL). The organic layer was washed with saturated NH_4Cl solution (50 mL) and dried over Na_2SO_4 . After concentration in vacuo, the residue was purified by silica gel chromatography (EtOAc/hexanes) to give the desired product (835 mg, 80%).

15

b) Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-pyrrolidin-3S-yl-amino]-thiophene-2-carboxylic acid methyl ester

Scheme 9

20



A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid methyl ester (0.5 g, 1.5 mmol), palladium acetate (0.015 g, 0.32 mmol), BINAP (0.009 g, 0.15 mmol), cesium carbonate (1.2 g, 4.5 mmol) and (3S)-Amino-


pyrrolidine-1-carboxylic acid tert-butyl ester (0.252 g, 1.01 mmol) in toluene (8 mL) was heated to 110 °C for 8 h. The reaction was diluted with ethyl acetate filtered through a Celite pad and purified by silica gel chromatography to give 3-[5-(3,3-dimethyl-but-1-ynyl)-2-methoxycarbonyl-thiophen-3S-ylamino]-pyrrolidine-1-carboxylic acid tert-butyl ester in 70% yield.

To a cooled (0 °C) THF (3 mL) solution of 3-[5-(3,3-dimethyl-but-1-ynyl)-2-methoxycarbonyl-thiophen-3S-ylamino]-pyrrolidine-1-carboxylic acid tert-butyl ester (1.0 mmol) was first added KHMDS (1.0 mmol, 0.5 M in toluene), followed by neat *trans*-4-methyl-cyclohexanecarbonyl chloride (0.2 mL, 1.24 mmol). The reaction was warmed slowly to room temperature and quenched with saturated ammonium chloride solution. The reaction mixture was diluted with EtOAc, washed with water, brine, dried over sodium sulfate, filtrated and concentrated. The crude was then diluted with EtOAc (10 mL), treated with 4M HCl in dioxane (0.5 mL) and heated to 50 °C for 30 min. The reaction mixture was cooled to room temperature concentrated and purified by silica gel column chromatography to give the title compound in 70% yield.

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-pyrrolidin-3S-yl-amino]-thiophene-2-carboxylic acid methyl ester (0.3 g, 0.7 mmol) in a 3:2:1 mixture of THF:MeOH:water (5 mL) was treated with lithium hydroxide monohydrate (0.69 g, 1.65 mmol) and heated to 60 °C for 1 hour. The organic volatiles were evaporated under reduced pressure and the crude material was purified by reverse-phase HPLC to afford 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-*trans*-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(S)-yl-amino]-thiophene-2-carboxylic acid in 60% yield.

25

Example 34: Compound 34 - Synthesis of 3-(S)-[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-*trans*-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(R)-yl ester



34


30

3-(*S*)-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 4-(*R*)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (*m/z*): 544 [M+1]
Retention time: 2.18min
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

15

Example 35: Compound 35 - Synthesis of 3-(*S*)-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid hexahydro-furo[2,3-*b*]furan-3-(*R*)-yl ester



20

35

3-(*S*)-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid hexahydro-furo[2,3-*b*]furan-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 4-nitro-benzoic acid hexahydro-furo[2,3-*b*]furan-3-(*R*)-yl ester was used in place of the chloroformate of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

30


LC/MS (m/z): 572.92 [M+1], 595.15 [M+Na⁺]

Retention time: 2.48min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

5

Example 36: Compound 36 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid hexahydro-furo[2,3-b]furan-3-(R)-yl ester



36

10

4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid hexahydro-furo[2,3-b]furan-3-(R)-yl ester was prepared in a similar fashion to 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-pyran-4-yloxy)carbonylamino]-cyclohexyl]-amino]-thiophene-2-carboxylic acid using method B, except the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid methyl ester was used instead of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester, and 4-nitro-benzoic acid hexahydro-furo[2,3-b]furan-3-(R)-yl ester was used in place of the chloroformate of tetrahydro-pyran-4-ol.

15

20


LC/MS (m/z): 587.01 [M+1], 609.20 [M+Na⁺]

Retention time: 2.40min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

25

Example 37: Compound 37 - Synthesis of 3-(R)-{3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxyloxy}-piperidine-1-carboxylic acid tert-butyl ester



37

3-(*R*)-{3-(*S*)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxyloxy}-piperidine-1-carboxylic acid tert-butyl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (*m/z*): 545 [M-98]

Retention time: 2.26min

LC: Thermo Finnigan PDA Detector


MS: Thermo Scientific LCQ Fleet

Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

Example 38: Compound 38 - Synthesis of 3-(*S*)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid piperidin-3-(*R*)-yl ester




38

3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid piperidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 544 [M+1]
Retention time: 2.18min
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

15

Example 39: Compound 39 - Synthesis of 4-{3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxyloxy}-piperidine-1-carboxylic acid tert-butyl ester



20

39

4-{3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxyloxy}-piperidine-1-carboxylic acid tert-butyl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid.

LC/MS (m/z): 644 [M+1], 544 [M-99]
Retention time: 2.56min

30

LC: Thermo Finnigan PDA Detector

MS: Thermo Scientific LCQ Fleet

Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm


Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic

5 acid

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

Example 40: Compound 40 - Synthesis of 3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-
ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-

10 carboxylic acid piperidin-4-yl ester




3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-
 cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid piperidin-4-yl ester was
 15 prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-
 yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid
 piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-
 ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(S)-yl-amino]-thiophene-2-
 carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-
 20 trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid.
 LC/MS (m/z): 544 [M+1]

Retention time: 2.09min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5
 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

25

Example 41: Compound 41 - Synthesis of 3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-
ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-
carboxylic acid tetrahydro-pyran-4-yl ester



- 3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid tetrahydro-pyran-4-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[[4-trans-methyl-cyclohexanecarbonyl]-pyrrolidin-3-(S)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[[4-trans-methyl-cyclohexanecarbonyl]-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and tetrahydro-pyran-4-ol was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 544 [M]

Retention time: 2.57min

- 15 LC: Thermo Finnigan PDA Detector


MS: Thermo Scientific LCQ Fleet

Column: Phenomenex Gemini-rx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid

- 20 Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% CAN

Example 42: Compound 42 - Synthesis of 3-(S)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid tetrahydro-furan-3-(R)-yl ester



25

3-(*S*)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid tetrahydro-furan-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and tetrahydro-furan-3-(*R*)-ol was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (*m/z*): 531 [*M*+1]

Retention time: 2.21min

LC: Thermo Finnigan PDA Detector


MS: Thermo Scientific LCQ Fleet

15 Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid


Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

20 Example 43: Compound 43 - Synthesis of 3-(*S*)-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 2-oxo-pyrrolidin-3-(*R*)-yl ester



43

25 3-(*S*)-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid 2-oxo-pyrrolidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-pyrrolidin-3-(*S*)-yl-amino]-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-



45


5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(piperidin-4-yloxy-carbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl]-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid.

LC/MS (m/z): 572 [M+1]

Retention time: 2.07min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Example 46: Compound 46 - Synthesis of 3-(R)-{4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl]-amino]-cyclohexylcarbamoyloxy}-piperidine-1-carboxylic acid tert-butyl ester



46

3-(R)-{4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl]-amino]-cyclohexylcarbamoyloxy}-piperidine-1-carboxylic acid tert-butyl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl]-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl


salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

5 LC/MS (*m/z*): 670 [M-1]

Retention time: 2.58min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

10 Example 47: Compound 47 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(piperidin-3-(*R*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid




15 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(piperidin-3-(*R*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl]-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

25 LC/MS (*m/z*): 572 [M+1]

Retention time: 2.11min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Example 48: Compound 48 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-furan-3-(*R*)-yloxycarbonylamino]-cyclohexyl]-amino]-thiophene-2-carboxylic acid



5

48

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(tetrahydro-furan-3-(*R*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-[4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and tetrahydro-furan-3-(*R*)-ol was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

15


LC/MS (m/z): 434 [M-124]

Retention time: 2.42min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

20

Example 49: Compound 49 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(5-oxo-pyrrolidin-3-(*S*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid



25

49

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(5-oxo-pyrrolidin-3-(*S*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic

86

acid was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 4-(*S*)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 572 [M+1]

10 Retention time: 2.08min

LC: Thermo Finnigan PDA Detector

MS: Thermo Scientific LCQ Fleet


Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic acid

15 acid

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

Example 50: Compound 50 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(2-oxo-pyrrolidin-3-(*R*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid



50

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(2-oxo-pyrrolidin-3-(*R*)-yloxycarbonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-


carboxylic acid and 3-(*R*)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 447 [M-124]

Retention time: 2.52min

- 5 Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.


Example 51: Compound 51 - Synthesis of Boc protected 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-3-(*R*)-yl ester



Boc protected 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.


- 20 LC/MS (m/z): 558 [M-99]
Retention time: 2.64min
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

- 25 Example 52: Compound 52 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-3-(*R*)-yl ester



- 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that 3-(*R*)-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.
- LC/MS (*m/z*): 558 [*M*+1]
Retention time: 2.15min
Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

- Example 53: Compound 53 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tetrahydro-furan-3-(*R*)-yl ester**



- 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tetrahydro-furan-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl except that tetrahydro-furan-3-(*R*)-ol was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.
- LC/MS (*m/z*): 545 [*M*+1]
Retention time: 2.28min

LC: Thermo Finnigan PDA Detector

MS: Thermo Scientific LCQ Fleet


Column: Phenomenex Gemini-nx 3u C18 110A 30 x 3 mm

Solvents: Acetonitrile with 0.1% trifluoroacetic acid, Water with 0.1% trifluoroacetic

5 acid

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

10 Example 54: Compound 54 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tetrahydro-pyran-4-yl ester




4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid tetrahydro-pyran-4-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl except that tetrahydro-pyran-4-ol was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 559 [M+1]

20 Retention time: 2.35min

Gradient: 0 min-3.1 min 2%-100% ACN, 3.1 min-3.75 min 100% ACN

25 Example 55: Compound 55 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(R)-yl ester




4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl except that 4-(*R*)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (*m/z*): 433 [M-124]

Retention time: 2.54min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Example 56: Compound 56 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(*S*)-yl ester



15

4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 5-oxo-pyrrolidin-3-(*R*)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl except that 4-(*S*)-hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

20

LC/MS (*m/z*): 433 [M-124]


Retention time: 2.54min

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN

25

Example 57: Compound 57 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 2-oxo-pyrrolidin-3-(*S*)-yl ester

30



4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 2-oxo-pyrrolidin-3-(S)-yl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl except that 3-(S)-Hydroxy-pyrrolidin-2-one was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 576 [M+19]

Retention time: 2.16min

LC: Thermo Electron Surveyor HPLC

MS: Finnigan LCQ Advantage MAX Mass Spectrometer


Column: Phenomenex Polar RP 30 mm X 4.6 mm


Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN

Method D

Example 64: Compound 64 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[[3-(hexahydro-furo[2,3-b]furan-3-yloxycarbonylamino)-1-(S)-methyl-propyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid





A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[(3-hydroxy-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid TFA salt (1.40g, 3.23mmol) in THF (30mL) was cooled to 0 °C and triphenylphosphine (2.53g, 9.69mmol) was added followed by phthalimide (0.713g, 4.84mmol). The reaction was stirred until homogenous then DIAD (1.06mL, 5.49mmol) was added. After warming to rt., the reaction was determined to be complete by LC/MS in 1h. The reaction was quenched with CH_3OH (1.0mL) and the solvent removed under reduced pressure. 5-(3,3-Dimethyl-but-1-ynyl)-3-[[3-(1,3-dioxo-1,3-dihydro-isoindole-2-yl)-1-(S)-methyl propyl]-(trans-4-methyl-cyclohexylcarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester (1.67g, 92%) was isolated by silica gel chromatography as an off-white solid.

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[[3-(1,3-dioxo-1,3-dihydro-isoindole-2-yl)-1-(S)-methyl propyl]-(trans-4-methyl-cyclohexylcarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester (168mg, 0.30mmol) in CH_3OH (2mL) and hydrazine

(15 μ L, 0.45mmol) was placed in a preheated 80 °C oil bath and stirred for 2h. The reaction was determined to be complete by LC/MS. Solvent was removed under reduced pressure and the crude material was coevaporated with toluene. 3-[[3-Amino-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester was used as is for the next step without purification.

A mixture of crude 3-[[3-Amino-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester in ACN (3.0mL) was treated with carbonic acid hexahydro-furo[2,3-*b*]furan-3-yl ester 4-nitro-phenyl ester (0.11g, 0.375mmol) followed by diisopropylethyl amine (0.1mL, 0.625mmol) and DMAP (cat). The reaction was stirred for 1h and quenched with CH₃OH (1mL). Solvent was removed under reduced pressure and the crude material was partitioned between EtOAc, and ½ sat NaHCO_{3(aq)}. The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 5-(3,3-Dimethyl-but-1-ynyl)-3-[[3-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester was isolated by reverse phase HPLC and carried forward wet to the next reaction.


A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[[3-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester in THF (2mL) and CH₃OH (1mL) was treated with dissolve LiOH H₂O (0.1g, 2.38mmol) in H₂O (2mL). The reaction was determined to be complete after 5h. The pH was adjusted to 2 with 2N HCl_(aq). After diluting with CH₃OH (3mL), 5-(3,3-Dimethyl-but-1-ynyl)-3-[[3-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid TFA salt (0.41g, 24% 3-steps) was isolated by reverse phase HPLC as a white solid.

LC/MS = 597.20 (M⁺ +Na)

Retention time: 3.76 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

Example 65: Compound 65 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-[1-(S)-methyl-3-(tetrahydrofuran-3-yloxy-carbonylamino)-propyl]-amino]-thiophene-2-carboxylic acid



5

65

A solution of 3-[[3-Amino-1-(S)-methyl-propyl)-(trans-4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (0.08g, 0.146mmol) in CH₃CN (1.5mL) was prepared. 3-(R)-Hydroxy-tetrahydrofuranyl-chloroformate (0.132g, 0.88mmol), which was prepared in a manner similar to method G, was added followed by diisopropylethyl amine (0.25mL, 1.50mmol) and DMAP (cat). After stirring at rt for 30min., the reaction was determined to be complete by LC/MS. The reaction was quenched with H₂O (1mL) and the organics removed under reduced pressure and 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-[1-(S)-methyl-3-(tetrahydrofuran-3-yloxy-carbonylamino)-propyl]-amino]-thiophene-2-carboxylic acid methyl ester was carried forward without purification.

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-[1-(S)-methyl-3-(tetrahydrofuran-3-yloxy-carbonylamino)-propyl]-amino]-thiophene-2-carboxylic acid methyl ester in THF (2mL) and CH₃OH (1mL) was treated with dissolve LiOH·H₂O (0.1g, 2.38mmol) in H₂O (2mL). The reaction was determined to be complete after 1h. The pH was adjusted to 2 with 2NHCl_(aq), the reaction diluted with CH₃OH (3mL) and 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-[1-(S)-methyl-3-(tetrahydrofuran-3-yloxy-carbonylamino)-propyl]-amino]-thiophene-2-carboxylic acid TFA salt (0.046g, 47%) was isolated by reverse phase HPLC as a white solid.

LC/MS = 555.24 (M+ Na⁺)

Retention time: 3.65 min


Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

30

Method E

Example 66: Compound 66 - 5-(3,3-Dimethyl-but-ynyl)-3-[[4-(hexahydro-furo[2,3-
b]furan-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-
amino]-thiophene-2-carboxylic acid

5



10 A solution of 3-[[4-(4-amino-cyclohexyl)-(trans-4-methylcyclohexanecarbonyl)-
amino]-5-(3,3-dimethyl-but-1ynyl)thiophene-2-carboxylic acid methyl ester (0.40g,
0.873mmol) in ACN (5.0mL) was cooled to 0 °C and carbonic acid hexahydro-
furo[2,3-b]furan-3-yl ester 4-nitro-phenyl ester (0.295g, 1.0mmol) was added,
followed by DIEA (391μL, 2.18mmol) and DMAP (cat). The reaction mixture was
15 warmed to rt, and stirred for 1h. The solvent was removed under reduced pressure
and the crude reaction mixture was partitioned between EtOAc and 2N K₂CO_{3(aq)}.

The layers were separated and the organics washed repeatedly with 2N $K_2CO_{3(aq)}$. Both epimers of 5-(3,3-Dimethyl-but-ynyl)-3-[[4-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester were isolated by reverse phase HPLC and

5 carried forward wet into the next reaction.

To each of the wet fractions of 5-(3,3-Dimethyl-but-ynyl)-3-[[4-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid methyl ester were added THF (0.5mL), CH_3OH (0.2mL), and LiOH (0.021g, 0.5mmol). The reactions were

10 allowed to stir at rt for 2h. The reactions were determined to be complete by LC/MS. The reactions were quenched with 2N $HCl_{(aq)}$ to pH = ca. 2. Both epimers of 5-(3,3-Dimethyl-but-ynyl)-3-[[4-(hexahydro-furo[2,3-*b*]furan-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid TFA salt (fast eluting: 89mg, 34%, slow eluting 34mg, 14%) were isolated by

15 reverse phase HPLC as an off-white solid.

LC/MS = 623.20 ($M^+ + Na$)

Retention time: 3.76 min


Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

20

Method F

Example 67: Compound 67 - 3-(S)-{3-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]](trans-4-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxyloxy}-3-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester


25



67

And
97

Example 68: Compound 68 - 3-(S)-[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methylcyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic-acid-3-methyl-pyrrolidin-3-yl-ester



3-(S)-[[5-(3,3-Dimethyl-but-1-ynyl)-2-methoxycarbonyl-thiophene-3-(S)-yl]-
 (trans-4-methyl-cyclohexanecarbonyl)-amino]-pyrrolidine-1-carboxylic acid *tert*-butyl
 ester (3.10g, 5.85mmol) was taken up in CH₂Cl₂ (50mL) and 4N HCl in dioxane
 (25mL, 100mmol) was added in one portion. The reaction was stirred for 2h and was
 5 determined to be complete by LC/MS. Solvents were removed under reduced
 pressure and the crude reaction mixture was co-evaporated with 3 X 25mL toluene.
 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-pyrrolidin-3-(S)-
 yl-amino]-thiophene-carboxylic acid methyl ester was carried forward crude into the
 next step.

10 A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[[trans-4-methyl-
 cyclohexanecarbonyl]-pyrrolidin-3-(S)-yl-amino]-thiophene-carboxylic acid methyl
 ester in THF (25mL) and CH₃OH (10mL) was treated with LiOH·H₂O (1.43g,
 34.2mmol) in H₂O (10mL). The reaction was determined to be complete by LC/MS
 after 2h. The pH was adjusted to 2 with 2N HCl_(aq) and the solvents were removed
 15 under reduced pressure. Co-evaporation with CH₃OH, then EtOAc, and finally
 toluene removed all H₂O. 5-(3,3-Dimethyl-but-1-ynyl)-3-[[trans-4-methyl-
 cyclohexanecarbonyl]-pyrrolidin-3-(S)-yl-amino]-thiophene-carboxylic acid was
 carried forward crude with LiCl salt into the next step.

20 5-(3,3-Dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-
 pyrrolidin-3-(S)-yl-amino]-thiophene-carboxylic acid (150mg) was taken up in ACN
 (2.0mL). The heterogeneous solution was allowed to stir for 5min, then DIEA (250μL,
 2.3mmol) and DMAP (cat) was added sequentially to the solution. The reaction was
 allowed to stir for 5 min then 3-chlorocarbonyloxy-3-methyl-pyrrolidine-1-carboxylic
 acid *tert*-butyl ester (≈200mg) prepared in a manner similar to method G was added
 25 in one portion. The reaction was stirred at rt for 15 min and determined to be
 complete by LC/MS. Solvent was removed under reduced pressure. The reaction
 mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried
 over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 3-{3-[[2-
 Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methyl-
 30 cyclohexanecarbonyl)-3-(S)-amino]-pyrrolidine-1-carboxyloxy}-3-methyl-pyrrolidine-1-
 carboxylic acid *tert*-butyl ester TFA salt (35mg, 20%) and 3-{3-[[2-Carboxy-5-(3,3-
 dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-3-(S)-
 amino]-pyrrolidine-1-carboxyloxy}-3-methyl-pyrrolidine-1-carboxylic acid TFA salt
 (11mg, 7%) were isolated by reverse phase HPLC as white solids.

3-{3-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-3-(*S*)-amino]-pyrrolidine-1-carboxyloxy}-3-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester TFA salt

LC/MS = 544.14 ($M^+ - \text{Boc}$)

5 Retention time: 4.02 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

3-{3-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-3-(*S*)-amino]-pyrrolidine-1-carboxyloxy}-3-methyl-pyrrolidine-1-carboxylic acid TFA salt

10 LC/MS = 544.10 ($M^+ + 1$)

Retention time: 3.36 min


Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20

15 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

Method G

3-Chlorocarbonyloxy-3-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester

20




A solution of 3-hydroxy-3-methylpyrrolidine-carboxylic acid *tert*-butyl ester (200mg, 1.00mmol) in THF (4mL) was slowly treated with phosgene 20% in toluene (0.954mL). Aftering stirring at rt for 16h, solvents were removed under reduced pressure and co-evaporate with CH_2Cl_2 . The crude 3-chlorocarbonyloxy-3-methylpyrrolidine-carboxylic acid *tert*-butyl ester was used directly in the next step.

25

Example 69: Compound 69 - Synthesis of 3-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-3-(*S*)-amino]-pyrrolidine-1-carboxylic acid 1-*tert*-butoxycarbonyl-3-methyl-azetidin-3-yl ester

30




69

5-(3,3-Dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-pyrrolidin-3-(S)-yl-amino]-thiophene-carboxylic acid (150mg) was taken up in ACN (2.0mL). The heterogeneous solution was allowed to stir for 5min, and then DIEA (250μL, 2.3mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-3-methyl-azetidine-1-carboxylic acid *tert*-butyl ester (≈200mg) which was prepared in a similar manner to Method G was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 3-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-3-(S)-amino]-pyrrolidine-1-carboxylic acid 1- *tert*-butoxycarbonyl-3-methyl-azetidin-3-yl ester TFA salt (46mg, 26%)

LC/MS = 630 (M⁺ +1)
Retention time: 3.89 min
Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

20

Example 70: Compound 70 - Synthesis of 3-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methylcyclohexanecarbonyl)-3-(S)-amino]-pyrrolidine-1-carboxylic acid 1,1-dioxo-tetrahydro-12⁶-thiophene-3-yl ester




70

5-(3,3-Dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl]-pyrrolidin-3-(S)-amino]-thiophene-carboxylic acid (200mg) was taken up in ACN (4.0mL). The heterogeneous solution was allowed to stir for 5min, and then DIEA (500 μ L, 4.6mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-1,1-dioxo-tetrahydro-1 λ^6 -thiophene ($\approx$ 200mg), which was prepared in a similar manner to method G, was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 3-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophene-3-yl]-(trans-4-methylcyclohexanecarbonyl)-3-(S)-amino]-pyrrolidine-1-carboxylic acid 1,1-dioxo-tetrahydro-1 λ^6 -thiophene-3-yl ester TFA salt (37mg, 19%)

LC/MS = 454.97 (M⁺ - methylcyclohexylcarbonyl)
Retention time: 3.56 min
Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

20

Example 71: Compound 71 - 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1-tert-butoxycarbonyl-3-methyl-pyrrolidin-3-yl ester




102

71

And

- 5 Example 72: Compound 72 - 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 3-methyl-pyrrolidin-3-yl ester



72

- 10 5-(3,3-Dimethyl-but-1-ynyl)-3-[(trans-4-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid (150mg) was taken up in ACN (2.0mL). The heterogeneous solution was allowed to stir for 5min, and then DIEA (250μL, 2.3mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-3-methylpyrrolidine-carboxylic acid
- 15 *tert*-butyl ester (≈200mg), which was prepared in a similar manner to method G, was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 4-[[2-
- 20 Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1-*tert*-butoxycarbonyl-3-methyl-pyrrolidin-3-yl ester TFA salt (26mg, 18%) and 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 3-methyl-pyrrolidin-3-yl ester TFA salt (8mg, 6%) were isolated by
- 25 reverse phase HPLC as white solids.
- 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1-*tert*-butoxycarbonyl-3-methyl-pyrrolidin-3-yl ester
- LC/MS = 558.19 (M⁺ -Boc)
- 30 Retention time: 4.03 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.


4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 3-methyl-pyrrolidin-3-yl ester

LC/MS = 558.09 ($M^+ + 1$)

Retention time: 3.40 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN

Example 73: Compound 73 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1-tert-butoxycarbonyl-3-methyl-azetidin-3-yl ester



73

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid (150mg) was taken up in ACN (2.0mL). The heterogeneous solution was allowed to stir for 5min, and then DIEA (250 μ L, 2.3mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-3-methylpyrrolidine-carboxylic acid *tert*-butyl ester ($\approx$ 200mg), which was prepared in a similar manner to method G, was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-


piperidine-1-carboxylic acid 1-tert-butoxycarbonyl-3-methyl-azetidin-3-yl ester TFA salt (38mg, 22%) was isolated by reverse phase HPLC as a white solid.

LC/MS = 544.20 (M^+ -Boc)

Retention time: 4.01 min

- 5 Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

Example 74: Compound 74 - Synthesis of 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1,1-dioxo-tetrahydro-1 λ^6 -thiophen-3-yl ester



74

- 5-(3,3-Dimethyl-but-1-ynyl)-3-[[trans-4-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid (200mg) was taken up in ACN (4.0mL). The heterogeneous solution was allowed to stir for 5min, then DIEA (350 μ L, 2.3mmol) and DMAP (cat) were added. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-1,1-dioxo-tetrahydro-1 λ^6 -thiophene ($\approx$ 200mg), which was prepared in a similar fashion as method G, was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid 1,1-dioxo-tetrahydro-1 λ^6 -thiophen-3-yl ester TFA salt (2.0mg, 1%) was isolated

by reverse phase HPLC as a white solid.


LC/MS = 469.10 (M^+ -methylcyclohexycarbonyl)

Retention time: 3.70 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

Example 75: Compound 75 - Synthesis of 3-{4-[[2-carboxy-5-(3,3-dimethyl-but-1-

5 ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-cyclohexylcarbamoyloxy}-3-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester




3-[4-Amino-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid (150mg) was taken up in ACN (2.5mL). The heterogeneous solution was allowed to stir for 5min, then DIEA (250μL, 2.3mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir for 5 min then 3-chlorocarbonyloxy-3-methylpyrrolidine-carboxylic acid *tert*-butyl ester (≈200mg), which was prepared in a manner similar to method G, was added in one portion. The reaction was stirred at rt for 15 min and determined to be complete by LC/MS. Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 3-{4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(trans-4-methyl-cyclohexanecarbonyl)-cyclohexylcarbamoyloxy}-3-methyl-pyrrolidine-1-carboxylic acid *tert*-butyl ester TFA salt (84mg, 47%) was isolated by reverse phase HPLC as a white solid.

LC/MS = 572.21 (M⁺ -Boc)

Retention time: 4.03 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.


Example 76: Compound 76 - 5-(3,3-Dimethyl-but-1-ynyl)-3- {(trans-4-cyclohexanecarbonyl)-[4-(3-methyl-pyrrolidin-3-yloxycarbonylamino)-cyclohexyl]-amino}-thiophene-2-carboxylic acid



A mixture of 3-{4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-
 (trans-4-methyl-cyclohexanecarbonyl)-cyclohexylcarbamoyloxy}-3-methyl-pyrrolidine-
 5 1-carboxylic acid *tert*-butyl ester TFA salt (50mg) in CH₂Cl₂ (2.0mL) and 4N HCl in
 dioxane (0.5mL) was stirred at rt for 2h. The reaction was determined to be complete
 by LC/MS. Solvents were removed under reduced pressure. 5-(3,3-Dimethyl-but-1-
 ynyl)-3- {(trans-4-cyclohexanecarbonyl)-[4-(3-methyl-pyrrolidin-3-
 yloxy-carbonylamino)-cyclohexyl]-amino}-thiophene-2-carboxylic acid TFA salt (32mg,
 10 73%) was isolated by reverse phase HPLC as a white solid.
 LC/MS = 572.19 (M⁺ +1)
 Retention time: 3.17/3.24 min
 Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20
 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

15

Example 77: Compound 77 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[[4-(1,1-
 dioxo-tetrahydro-1λ⁶-thiophen-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-
 methylcyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid



20

3-[4-Amino-cyclohexyl]-(trans-4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-
 dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid (200mg) was taken up in ACN
 (4.0mL). The heterogeneous solution was allowed to stir for 5min, then DIEA (500μL,
 4.6mmol) and DMAP (cat) were added sequentially. The reaction was allowed to stir
 25 for 5 min then 3-chlorocarbonyloxy-1,1-dioxo-tetrahydro-1λ⁶-thiophene (≈200mg),
 which was prepared in a similar manner to method G, was added in one portion. The

reaction was stirred at rt for 15 min and determined to be complete by LC/MS.


Solvent was removed under reduced pressure. The reaction mixture was partitioned between EtOAc and 2N HCl_(aq). The organics were dried over Na₂SO₄, solids filtered and solvent removed under reduced pressure. 5-(3,3-Dimethyl-but-1-ynyl)-3-[[4-(1,1-dioxo-tetrahydro-1λ⁶-thiophen-3-yloxy-carbonylamino)-cyclohexyl]-(trans-4-methylcyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid TFA salt (36mg, 19%)

LC/MS = 606.86 (M⁺ +1)

Retention time: 3.61 min

Gradient: 0 min-0.2min 5% ACN, 0.2 min-3.95 min 5%-100% ACN, 3.95 min-5.20 min 100% ACN, 5.20 min-5.5 min 100%-5% ACN, 5.5 min-6 min 5% ACN.

Example 79: Compound 79 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-[4-(N'-methyl-N'-pyridin-2-yl-hydrazino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid




25 2-Bromopyridine (5 mL, 52.4 mmol) and methylhydrazine (20 mL, 360 mmol) were mixed together in a flask fitted with a water-condenser. After a few minutes a vigorous exothermic reaction occurred with refluxing of the methylhydrazine. When the reaction had subsided, the mixture was left for 16 hours, and the excess methylhydrazine was removed under reduced pressure. The cooled residue was stirred with aqueous sodium hydroxide solution (30 ml of 20%), and the resulting

solution was extracted with ether (3 x 100 ml). The combined extracts were dried (K_2CO_3) and purified by silica gel chromatography 0-3% EtOH/ CH_2Cl_2 to give 4.8 g of N-Methyl-N-pyridin-2-yl-hydrazine.

A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (412 mg, 0.90 mmol) and N-Methyl-N-pyridin-2-yl-hydrazine (222 mg, 1.8 mmol) in DCE (6 mL) was treated with AcOH (200 μ L, 3.0 mmol) followed by $NaBH(OAc)_3$ (300 mg, 4.08 mmol) in two or three portions. After 30 min, $NaHCO_3$ (saturated aqueous solution, 4-8 mL) was added to the mixture, followed by brine (20 mL), and the crude product was extracted with ethyl acetate (2x20 mL). The combined organic layers were concentrated and the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (20 mL), treated with lithium hydroxide (4.5 mmol, 188 mg) and heated to 60 deg C for 2 hours. The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 189 mg (37% yield) of the title compound as its *trans* isomer (TFA salt): MS (m/z): 551.3 $[M-H]^+$; HPLC retention time: 3.39 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) and 71 mg (14% yield) of the title compound as its *cis* isomer (TFA salt): MS (m/z): 551.3 $[M-H]^+$; HPLC retention time: 3.52 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).


Example 80: Compound 80 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-[4-(N'-methyl-N'-pyrazin-2-yl-hydrazino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid



2-Chloropyrazine (1g, 8.3 mmol) and methylhydrazine (1.31 mL, 25 mmol) were mixed together in a flask fitted with a water condenser. After a few minutes an exotherm was observed. After 16 h the excess methylhydrazine was removed under reduced pressure. The cooled residue was stirred with aqueous sodium hydroxide solution (20 ml of 20%), and the resulting solution was extracted with ether (6 x 100 ml). The combined extracts were dried (K_2CO_3) and concentrated to give a light orange solid 370 mg of N-Methyl-N-pyrazin-2-yl-hydrazine.

A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (100 mg, 0.22 mmol) and N-Methyl-N-pyrazin-2-yl-hydrazine (41 mg, 0.33 mmol) in DCE (2 mL) was treated with AcOH (100 μ L, 1.5 mmol) followed by $NaBH(OAc)_3$ (120 mg, 0.5 mmol) in two portions. After 5 h, $NaHCO_3$ (saturated aqueous solution, 2 mL) was added to the mixture, followed by brine (20 mL), and the crude product was extracted with ethyl acetate (2x20 mL). The combined organic layers were concentrated and the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (20 mL), treated with lithium hydroxide (1.1 mmol, 46 mg) and heated to 60 deg C for 1 hours. The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 11 mg of the title compound as its *trans* isomer (TFA salt): MS (m/z): 552.2 $[M-H]^+$; HPLC retention time: 26.05 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient) and 7 mg of the title compound as its *cis* isomer (TFA salt): MS (m/z): 552.2 $[M-H]^+$; HPLC retention time: 24.38 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient).

Example 81: Compound 81 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid




81

2-Chlorothiazole (0.9 g, 7.52 mmol) was placed in a flask fitted with a water condenser and methylhydrazine (1.31 mL, 25 mmol) was added dropwise. An
 5 immediate exotherm was observed and methylhydrazine began to reflux. When the reaction had subsided, the mixture was left for 16 hours, and the excess methylhydrazine was removed under reduced pressure. The cooled residue was stirred with aqueous sodium hydroxide solution (20 ml of 20%), and the resulting solution was extracted with ether (6 x 100 ml). The combined extracts were dried
 10 (K_2CO_3) and concentrated to give 485 mg of N-Methyl-N-thiazol-2-yl-hydrazine.

A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (200 mg, 0.44 mmol) and N-Methyl-N-thiazol-2-yl-hydrazine (112 mg, 0.87 mmol) in DCE (4 mL) was treated with AcOH (200 μ L, 3.0 mmol) followed by $NaBH(OAc)_3$ (120 mg, 0.57
 15 mmol) in two portions. After 5 h, $NaHCO_3$ (saturated aqueous solution, 3 mL) was added to the mixture, followed by brine (20 mL), and the crude product was extracted with ethyl acetate (2x20 mL). The combined organic layers were concentrated and the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (20 mL), treated with lithium hydroxide (2.2 mmol, 92 mg) and heated to 60 deg C for 1 hours.
 20 The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 36 mg of the title compound as its *trans* isomer (TFA salt): MS (m/z): 557.2 $[M-H]^+$; HPLC retention time: 3.43 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) and 8 mg of the title compound
 25 as its *cis* isomer (TFA salt): MS (m/z): 557.2 $[M-H]^+$; HPLC retention time: 3.41 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 82: Compound 82 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-[N'-(6-hydroxy-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl)-(4-methyl-
 30 cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid

Example 83: Compound 83 - [(4-[N'-(6-Chloro-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl)-(4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid



5 3,6-Dichloro-pyridazine (1g, 6.75 mmol) was placed in a flask fitted with a water condenser and methylhydrazine (1.31 mL, 25 mmol) was added dropwise. An immediate exotherm was observed and methylhydrazine begins to reflux. When the reaction had subsided, the mixture was left for 16 hours, and the excess methylhydrazine was removed under reduced pressure. The cooled residue was

10 stirred with aqueous sodium hydroxide solution (20 ml of 20%), and the resulting solution was extracted with ether (6 x 100 ml). The combined extracts were dried (K_2CO_3) and concentrated to give 427 mg of N-(6-Chloro-pyridazin-3-yl)-N-methylhydrazine.

 A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (500 mg, 1.09 mmol) and N-(6-Chloro-pyridazin-3-yl)-N-methyl-hydrazine (260 mg, 1.64 mmol) in DCE (8 mL) was treated with AcOH (0.5 mL, 8.4 mmol) followed by $NaBH(OAc)_3$ (347 mg, 1.64 mmol) in two portions. After 5 h, $NaHCO_3$ (saturated aqueous solution, 10 mL) was added to the mixture, followed by brine (20 mL), and the crude product

20 was extracted with ethyl acetate (2x50 mL). The combined organic layers were concentrated and the crude material was dissolved in 4 mL of acetic acid, treated with NaOAc (893 mg, 10.9 mmol) and heated in a sealed tube at 100 deg C for 16 h. Upon cooling the mixture was diluted with ethyl acetate and neutralized with $NaHCO_3$ (saturated aqueous solution). The combined organic layers were concentrated and


25 the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (20 mL), treated with lithium hydroxide (5.45 mmol, 228 mg) and heated to 60 deg C for 1 hour.

The crude material was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile:water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 16 mg of 3-[[4-[N'-(6-Chloro-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl]-(4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid as the *trans* isomer (TFA salt): MS (m/z): 587.2 [M-H]⁺; HPLC retention time: 26.39 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient); 4 mg of 3-[[4-[N'-(6-Chloro-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl]-(4-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid as the *cis* isomer (TFA salt): MS (m/z): 587.2 [M-H]⁺; HPLC retention time: 24.03 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient); 44 mg of 5-(3,3-Dimethyl-but-1-ynyl)-3-[[4-[N'-(6-hydroxy-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl]-(4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid as its *trans* isomer (TFA salt): MS (m/z): 569.4 [M-H]⁺; HPLC retention time: 26.15 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient), and 71 mg of 5-(3,3-Dimethyl-but-1-ynyl)-3-[[4-[N'-(6-hydroxy-pyridazin-3-yl)-N'-methyl-hydrazino]-cyclohexyl]-(4-methyl-cyclohexanecarbonyl)-amino]-thiophene-2-carboxylic acid as its *cis* isomer (TFA salt): MS (m/z): 569.4 [M-H]⁺; HPLC retention time: 24.67 min (Phenomenex Luna C18, 2-98% acetonitrile:water with 0.1% trifluoroacetic acid, 30 min gradient).

Hydroxylamine Synthesis

25

Hydroxylamine 1: Synthesis of O-pyridin-2-ylmethylhydroxylamine dihydrochloride





2-Chloromethylpyridine hydrochloride (514 mg, 3.13 mmol), N-hydroxyphthalimide (515 mg, 3.18 mmol), and triethylamine (1.3 mL, 9.3 mmol) were mixed in acetonitrile (5 mL) and stirred at 80 °C for 2.5 hours. The reaction mixture was diluted with ethyl acetate and washed sequentially with 1N NaOH_(aq), water, and
 5 brine. The organic phase was dried over MgSO₄, filtered, and concentrated. The residue was recrystallized from ethyl acetate and hexanes to provide N-(pyridine-2-ylmethoxy)phthalimide (364 mg, 1.44 mmol).

Hydrazine monohydrate (53 μL, 1.5 mmol) was added to a solution of N-(pyridine-2-ylmethoxy)phthalimide (356 mg, 1.41 mmol) in ethanol (2 mL) and stirred
 10 at 80 °C for one hour in a sealed tube. The reaction mixture was diluted with ethyl acetate, adsorbed onto silica gel, and purified by SGC (0-20% [8:1 EtOH:NH₄OH_(aq)]/DCM). After concentration of fractions the residue was taken up in diethyl ether and treated with 4N HCl_(Et₂O) to provide O-pyridin-2-ylmethylhydroxylamine dihydrochloride (152 mg, 0.77 mmol) as a white solid.

15


Using the same procedure as described for O-pyridin-2-ylmethylhydroxylamine dihydrochloride, the hydroxylamines shown below were synthesized.



20

Hydroxylamine 6


Example 112: Compound 112 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-((4-*trans*-methyl-cyclohexanecarbonyl)-[4-(pyridin-2-ylmethoxyimino)-cyclohexyl]-amino)-thiophene-2-carboxylic acid



112

- 5 5-(3,3-dimethyl-but-1-ynyl)-3-[(*trans*-4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (264 mg, 0.58 mmol), O-pyridin-2-ylmethylhydroxylamine dihydrochloride (142 mg, 0.72 mmol), and sodium acetate (177 mg, 2.2 mmol) were mixed in 2:1 MeOH:DCM (3 mL) and stirred at 50 °C for 1 hour. Lithium hydroxide monohydrate (240 mg, 5.7 mmol) and water (1 mL) were then added and the reaction mixture continued to stir at 50 °C for 2 hours.
- 10 The reaction mixture was then partitioned between ethyl acetate and 5% citric acid_(aq). The aqueous phase was neutralized with 4N NaOH_(aq) and thrice extracted with ethyl acetate. The combined organic phases were washed with brine, dried over MgSO₄, filtered, and concentrated. The residue was purified by HPLC (Gemini column; 25% acetonitrile:water, 2 min; 25-100% acetonitrile:water, 16 min; 100% acetonitrile, 3 min; both solvents containing 0.1% trifluoroacetic acid). This resulted in 234 mg
- 15 (19% yield over 2 steps) of the title compound as a white powder (TFA salt): MS (*m/z*): 550.2 [M+H]⁺; HPLC retention time: 3.55 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

- 20 Example 113: Compound 113 - 3-(N-(4-Benzoyloximino)cyclohexyl)-4-methylcyclohexanecarboxamido)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid




113

115

Prepared in the same manner as Example 112 using O-benzylhydroxylamine.
MS (m/z): 549.1 [M+H]⁺; HPLC retention time: 5.23 min.


- 5 Example 114: Compound 114 - 3-(N-(4-((1H-1,2,4-Triazol-1-yl)methoxyimino)cyclohexyl)-4-methylcyclohexanecarboxamido)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid



10

Prepared in the same manner as Example 112 using hydroxylamine 2. MS (m/z): 540.1 [M+H]⁺; HPLC retention time: 4.29 min.


- 15 Example 115: Compound 115 - 3-(N-(4-(Carboxymethoxyimino)cyclohexyl)-4-methylcyclohexanecarboxamido)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid




20

Prepared in the same manner as Example 112 using hydroxylamine 5. MS (m/z): 517.0 [M+H]⁺; HPLC retention time: 4.28 min.

Example 116: Compound 116 - 3-(N-(4-(Cyanomethoxyimino)cyclohexyl)-4-methylcyclohexanecarboxamido)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid



5




5-(3,3-Dimethyl-but-1-ynyl)-3-[(*trans*-4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid methyl ester (519 mg, 1.13 mmol) was dissolved in a mixture of 3:2:1 THF:MeOH:H₂O (25 mL). Lithium hydroxide (5 mL, 1.0N aqueous solution) was added and the reaction mixture was stirred at 60 °C for 1 hour. The reaction mixture was then partitioned between ethyl acetate and water. The aqueous phase was neutralized with 5% citric acid_(aq), and thrice extracted with ethyl acetate. The combined organic phases were washed with brine, dried over MgSO₄, filtered, and concentrated to afford 505 mg of 5-(3,3-dimethyl-but-1-ynyl)-3-[(*trans*-4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid which was carried on without further purification.

5-(3,3-Dimethyl-but-1-ynyl)-3-[(*trans*-4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid (505 mg, 1.02 mmol), O-cyanomethylhydroxylamine hydrochloride (174 mg, 1.32 mmol), and sodium acetate (212 mg, 2.6 mmol) were mixed in 2:1 MeOH:DCM (7.5 mL) and stirred at 50 °C for 1 hour. The reaction mixture was then partitioned between ethyl acetate and water. The aqueous phase was thrice extracted with ethyl acetate and the combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated.


The residue was purified by HPLC (Gemini column; 25% acetonitrile:water, 2 min; 25-100% acetonitrile:water, 16 min; 100% acetonitrile, 3 min; both solvents containing 0.1% trifluoroacetic acid). This resulted in 159 mg (28% yield over 2 steps) of the title compound as a white powder: MS (m/z): 495.7 [M-H]⁻; HPLC retention time: 4.68 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 117: Compound 117 - 5-(3,3-Dimethylbut-1-ynyl)-3-(4-methyl-N-(4-(pyridin-4-ylmethoxyimino)cyclohexyl)cyclohexanecarboxamido)thiophene-2-carboxylic acid




Prepared in the same manner as Example 112 using hydroxylamine 3. MS (m/z): 550.1 [M+H]⁺; HPLC retention time: 3.46 min.

Example 118: Compound 118 - 5-(3,3-Dimethylbut-1-ynyl)-3-(4-methyl-N-(4-(pyridin-3-ylmethoxyimino)cyclohexyl)cyclohexanecarboxamido)thiophene-2-carboxylic acid



Prepared in the same manner as Example 112 using hydroxylamine 4. MS (m/z): 550.1 [M+H]⁺; HPLC retention time: 3.47 min.


Example 119: Compound 119 - 5-(3,3-Dimethylbut-1-ynyl)-3-(4-methyl-N-(4-(phenoxymino)cyclohexyl)cyclohexanecarboxamido)thiophene-2-carboxylic acid



Prepared in the same manner as Example 112 using O-phenylhydroxylamine.

5 MS (m/z): 535.0 [M+H]⁺; HPLC retention time: 5.38 min.

Example 120: Compound 120 - 5-(3,3-Dimethylbut-1-ynyl)-3-(N-(4-(methoxyimino)cyclohexyl)-4-methylcyclohexanecarboxamido)thiophene-2-carboxylic acid




10

Prepared in the same manner as Example 112 using O-methylhydroxylamine.

MS (m/z): 473.0 [M+H]⁺; HPLC retention time: 4.79 min.

15


Example 121: Compound 121 - 3-(N-(4-(4-(Methylsulfonyl)benzyloxyimino)cyclohexyl)-4-methylcyclohexanecarboxamido)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid



20

Prepared in the same manner as Example 112 using hydroxylamine **6**. MS (m/z): 627.0 [M+H]⁺; HPLC retention time: 4.75 min.

5 Example 122: Compound 122 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(1,4-dioxo-spiro[4.5]dec-8-yl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid



10 To a solution of 3-tert-butoxycarbonylamino-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (4.3 g, 12.8 mmol) in NMP (64 mL) was added K-OtBu (1.6 g, 14.1 mmol). After 10 min a solution of O-4-nitrobenzoylhydroxylamine (3.0 g, 16.6 mmol) in NMP (16 mL) was added dropwise

and the reaction was allowed to stir for 16 h. The reaction was quenched with LiCl solution (5% in water, 50 mL) and diluted with EtOAc (50 mL). The crude product was washed 3 X 50 mL with 5% LiCl solution, dried over Na₂SO₄, and purified by silica gel chromatography 0-30% EtOAc/hexanes to give 3-(N-tert-butoxycarbonyl-hydrazino)-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (2.6 g, 7.4 mmol).

3-(N-tert-Butoxycarbonyl-hydrazino)-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (2.6 g, 7.4 mmol) in Et₂O (37 mL) was cooled to 0 °C and treated with trifluoroacetic anhydride (1.9 g, 8.9 mmol) for 1 h. The reaction was concentrated, dissolved with neat TFA (4.3 g, 37 mmol) and heated to 60 °C. After 15 min, the reaction was cooled to rt, diluted with toluene (30 mL) and concentrated, this step was repeated three times, to give crystalline 5-(3,3-Dimethyl-but-1-ynyl)-3-[N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic acid methyl ester (2.1 g, 6.0 mmol).


A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic acid methyl ester (2.1 g, 6.0 mmol), 4-methyl-cyclohexanecarbonyl chloride (1.45 g, 9 mmol), DMAP (1.1 g, 9 mmol) and DCE (20 mL) was heated to 40 °C for 16 h. After cooling, the mixture was concentrated and the product was purified by silica gel chromatography 0-30% EtOAc/hexanes to give 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic acid methyl ester (2.74 g, 5.8 mmol).

To a solution of 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic acid methyl ester (1.6 g, 3.4 mmol) in THF (17 mL) was added K-OtBu (4.1 mmol, 1.0 M in THF), followed by MeI (1.0 g, 6.8 mmol). After stirring at rt for 16 h the reaction was diluted with EtOAc washed once with 20 mL of brine and purified by silica gel chromatography 0-20% EtOAc/hexanes to give 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic acid methyl ester. This was then diluted with EtOAc (20 mL) and treated with NaOH (6.8 mL of 2M aq solution) to give 5-(3,3-Dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid.

A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid (300 mg, 0.8 mmol),

1,4-cyclohexadione-mono-ethylene ketal (149 mg, 1 mmol), AcOH (300 mg, 5 mmol) in DCE (2 mL) was treated with $\text{NaBH}(\text{OAc})_3$ (430 mg, 2 mmol) for 16 h. The reaction was quenched with water (10 mL) and extracted with EtOAc. A portion of this material was then purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid) giving 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(1,4-dioxo-spiro[4.5]dec-8-yl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid as its TFA salt: MS (m/z): 517.1 $[\text{M}-\text{H}]^+$; HPLC retention time: 4.73 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 123 (Compound 123) and 124 (Compound 124) - Synthesis of 3-(2-(4-hydroxycyclohexyl)-2-methyl-1-(4-methylcyclohexanecarbonyl)hydrazinyl)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid (123) and 3-(2-(4-hydroxycyclohexyl)-2-methyl-1-(4-methylcyclohexanecarbonyl)hydrazinyl)-5-(3,3-dimethylbut-1-ynyl)thiophene-2-carboxylic acid (124)




A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(1,4-dioxo-spiro[4.5]dec-8-yl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic (500

mg, 1 mmol) in a 1:1 mixture of THF and MeOH (2.5 mL) was treated with 1 M HCl (2.5 mL) and heated for 3 h at 60 °C. The reaction mixture was cooled to room temperature, diluted with EtOAc (10 mL), neutralized with sat. NaHCO₃, washed once with brine (10 mL), dried over Na₂SO₄, and concentrated. The crude material
 5 was dissolved in wet THF (5 mL) and treated with NaBH₄ (38 mg, 1.1 mmol) at 0 °C for 30 min. A portion of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(4-hydroxy-cyclohexyl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50%
 10 acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid) to give 20 mg as its TFA salt: MS (m/z): 475.7 [M-H]⁺; HPLC retention time: 4.33 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).


Additional HPLC purification separated Example 123 (retention time: 27.89 min) from Example 124 (retention time 27.89).

15

Example 125 and 126 - Synthesis of 5-(3,3-dimethylbut-1-ynyl)-3-(2-methyl-1-(4-methylcyclohexanecarbonyl)-2-(4-(pyridin-3-yloxy)cyclohexyl)hydrazinyl)thiophene-2-carboxylic acid (Compound 125) and 5-(3,3-dimethylbut-1-ynyl)-3-(2-methyl-1-(4-methylcyclohexanecarbonyl)-2-(4-(pyridin-3-yloxy)cyclohexyl)hydrazinyl)thiophene-2-carboxylic acid (Compound 126)
 20



125




126

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(4-hydroxy-cyclohexyl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid
 25 (100 mg, 0.2 mmol) and 3-fluoro-pyridine (97 μL, 1 mmol) in DMF (0.6 mL) was treated with sodium hydride (40 mg, 1 mmol, 60% oil dispersion) in two or three

portions. The mixture was stirred until the bubbling slowed, and was sealed and heated at 100 deg C for 8 h. After cooling, ethyl acetate (2-3 mL) was added and the mixture was carefully quenched with citric acid (10% aqueous solution, 2-3 mL). Water was added and the mixture was extracted with ethyl acetate (2x30 mL). The combined organic layers were dried over sodium sulfate, filtered and concentrated. The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-95% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 37 mg of the title compound as a white powder (bis-TFA salt): MS (m/z): 552.1 [M-H]⁺; HPLC retention time: 3.50 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Additional HPLC purification separated Example 125 (retention time: 22.46 min) from Example 126 (retention time 22.62).


Example 127: Compound 127 - Synthesis of 4-[N'-(2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl)]-N-methyl-N'-(4-methyl-cyclohexanecarbonyl)-hydrazino]-piperidine-1-carboxylic acid tert-butyl ester



127


The title compound was synthesized in a manner analogous to Example 122, using 1-N-Boc-4-piperidone in place of 1,4-cyclohexadione-mono-ethylene ketal: MS (m/z): 560.1 [M-H]⁺; HPLC retention time: 5.10 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 128: Compound 128 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-piperidin-4-yl-hydrazino]-thiophene-2-carboxylic acid

**128**

Crude 4-[N'-[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-N-methyl-N'-(4-methyl-cyclohexanecarbonyl)-hydrazino]-piperidine-1-carboxylic acid tert-butyl ester (220 mg, 500 μ mol) (Example 127) was treated with TFA (3 mL, 4.6 mmol) at 60 °C for 10 min. After cooling, toluene (2-3 mL) was added and the mixture was concentrated, this was repeated several times, and a portion of this crude material was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-95% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 20 mg of the title compound as a white powder (bis-TFA salt): MS (m/z): 460.2 [M-H]⁺; HPLC retention time: 3.15 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 129: Compound 129 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(1-methyl-piperidin-4-yl)-hydrazino]-thiophene-2-carboxylic acid


**129**

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-piperidin-4-yl-hydrazino]-thiophene-2-carboxylic acid (100 mg, 225 μ mol) (Example 128), formaldehyde (1.12 mmol), acetic acid (0.5 mL, 8 mmol) in DCE (3 mL) was treated with NaBH(OAc)₃ (36 mg, 168 μ mol) for 16 h. The reaction was quenched with water (10 mL) and extracted with EtOAc. The crude material was then purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents

containing 0.1% trifluoroacetic acid) giving 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(1-methyl-piperidin-4-yl)-hydrazino]-thiophene-2-carboxylic acid as its bis-TFA salt: MS (m/z): 474.2 [M-H]⁺; HPLC retention time: 3.17 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

5

Example 130: Compound 130 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(2-(pyridin-3-yloxy)-ethyl)-hydrazino]-thiophene-2-carboxylic acid




130

10

The title compound was synthesized in a manner analogous to Example 122, using (pyridin-3-yloxy)-acetaldehyde in place of 1,4-cyclohexadione-mono-ethylene ketal: MS (m/z): 498.1 [M-H]⁺; HPLC retention time: 3.31 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

15

Example 131: Compound 131 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(tetrahydro-pyran-4-ylmethyl)-hydrazino]-thiophene-2-carboxylic acid



131


20

To a solution of 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-N'-(2,2,2-trifluoro-acetyl)-hydrazino]-thiophene-2-carboxylic

acid methyl ester (1.6 g, 3.4 mmol) in EtOAc (20 mL) was added NaOH (6.8 mL of 2M aq solution). After 2 h stirring at rt, the reaction was neutralized with 1M HCl aq, extracted with EtOAc, dried over Na₂SO₄, and concentrated to give 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid (1.1g, 3 mmol) as an off white solid.

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid (50 mg, 0.132 mmol), tetrahydro-pyran-4-carbaldehyde (19 mg, 0.172 mmol), AcOH (24 mg, 0.4 mmol) in DCE (2 mL) was treated with NaBH(OAc)₃ (42 mg, 0.2 mmol) for 16 h. The reaction was quenched with water (10 mL), extracted with EtOAc and concentrated. A mixture of the crude material, AcOH (47 mg, 2.37 mmol), dissolved in iPrOH (3 mL) was treated with NaCNBH₃ (13 mg, 0.198 mmol) then heated to 50 °C for 16 h. The product was then purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid) giving 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(tetrahydro-pyran-4-ylmethyl)-hydrazino]-thiophene-2-carboxylic acid as its TFA salt: MS (m/z): 475.1 [M-H]⁺; HPLC retention time: 4.62 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 132: Compound 132 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-(2-(pyridin-2-yloxy)-ethyl)-hydrazino]-thiophene-2-carboxylic acid




132

25

The title compound was synthesized in a manner analogous to Example 131, using (pyridin-2-yloxy)-acetaldehyde in place of tetrahydro-pyran-4-carbaldehyde: MS (m/z): 498.1 [M-H]⁺; HPLC retention time: 29.8 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) 30 min run.

Example 133: Compound 133 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-N'-[2-(pyridin-3-yloxy)-ethyl]-hydrazino]-thiophene-2-carboxylic acid



5


133

The title compound was synthesized from 5-(3,3-dimethyl-but-1-ynyl)-3-[N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid and (pyridin-3-yloxy)-acetaldehyde in a manner similar to that of Example 131. MS (m/z): 484.0 [M-H]⁻; HPLC retention time: 3.44 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

10

Example 134: Compound 134 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(2-hydroxy-ethyl)-N-(4-methyl-cyclohexanecarbonyl)-N'-oxetan-3-yl-hydrazino]-thiophene-2-carboxylic acid


15

**134**

The title compound was synthesized in a manner analogous to Example 131, using 3-(tert-butyl-dimethyl-silanyloxy)-acetaldehyde in place of tetrahydro-pyran-4-carbaldehyde, treating the crude with TBAF (1 mmol, 0.1 M in THF) prior to the second reductive amination in which formaldehyde is replaced with oxetan-3-one: MS (m/z): 461.2 [M-H]⁻; HPLC retention time: 4.38 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

20

Example 135: Compound 135 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-(2-hydroxy-ethyl)-N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid




5

135

The title compound was synthesized in a manner analogous to Example 131, using 3-(tert-butyl-dimethyl-silanyloxy)-acetaldehyde in place of tetrahydro-pyran-4-carbaldehyde and treating the crude with TBAF (1 mmol, 0.1 M in THF) prior to the second reductive amination with formaldehyde: MS (m/z): 422.1 [M-H]⁺; HPLC retention time: 4.06 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

10

Example 136: Compound 136 – Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-dimethyl-N-(4-methyl-cyclohexanecarbonyl)-hydrazino]-thiophene-2-carboxylic acid



15


136

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid methyl ester (0.200 g, 0.574 mmol), LiCl (0.048, 1.14 mmol), N,N-dimethylhydrazine (0.043 mL, 0.57 mmol), Pd₂(dba)₃ (0.003 g, 0.005 mmol), xantphos (0.008 g, 0.005 mmol), and NaOtBu (0.037 g, 0.394 mmol) in toluene (3 mL) was heated to 80 °C for 16h. The reaction was diluted with ethyl acetate filtered through a Celite pad and concentrated the crude material was purified by silica gel chromatography (0-15% EtOAc/hexane) to give 5-(3,3-dimethyl-but-1-ynyl)-3-(N'-dimethyl-hydrazino)-thiophene-2-carboxylic acid methyl ester in 63% yield. This was then dissolved in

20

pyridine (5 mL) and treated with neat *trans*-4-methyl-cyclohexanecarbonyl chloride (0.132 g, 0.825 mmol). The reaction was heated for 16 h at 85 °C and quenched with saturated ammonium chloride solution. The reaction mixture was diluted with EtOAc, washed with water, brine, dried over sodium sulfate, filtered and concentrated. The crude material was then dissolved in a 3:2:1 mixture of THF:MeOH:water (5 mL), treated with lithium hydroxide monohydrate (0.69 g, 1.65 mmol) and heated to 60 °C for 1 hour. The organic volatiles were evaporated under reduced pressure and the crude material was purified by HPLC to afford the title compound. MS (m/z): 456.0 [M+H]⁺; HPLC retention time: 3.91 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 137: Compound 137 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[N'-methyl-N-(4-methyl-cyclohexanecarbonyl)-N'-pyridin-2-yl-hydrazino]-thiophene-2-carboxylic acid




137

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid methyl ester (0.100 g, 0.28 mmol), N, N-diethylenediamine (9 µL, 0.086 mmol), N-methyl-N-pyridin-2-yl-hydrazine (0.105 g, 0.86 mmol), CuI (0.01 g, 0.057 mmol), 4 Å ms (0.114 mg) and K₂CO₃ (0.118 g, 0.861 mmol) in DMF (3 mL) was heated to 80 °C for 16h. The reaction was diluted with ethyl acetate, washed twice with 5% LiCl, concentrated and the crude material was purified by silica gel chromatography (0-20% EtOH in DCM) to give 5-(3,3-dimethyl-but-1-ynyl)-3-(N'-methyl-N'-pyridin-2-yl-hydrazino)-thiophene-2-carboxylic acid methyl ester in 50% yield. This was then dissolved in pyridine (5 mL) and treated with neat *trans*-4-methyl-cyclohexanecarbonyl chloride (0.132 g, 0.825 mmol). The reaction was heated for 16 h at 85 °C and quenched with saturated ammonium chloride solution. The reaction mixture was diluted with EtOAc, washed with water, brine, dried over sodium sulfate,


filtered and concentrated. The crude material was then dissolved in a 3:2:1 mixture of THF:MeOH:water (5 mL), treated with lithium hydroxide monohydrate (0.69 g, 1.65 mmol) and heated to 60 °C for 1 hour. The organic volatiles were evaporated under reduced pressure and the crude material was purified by HPLC to afford the title compound. MS (m/z): 454.1 [M+H]⁺; HPLC retention time: 3.95 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 140: Compound 140 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-methyl-cyclohexanecarbonyl)-[1-(pyridine-3-carbonyl)-piperidin-4-yl]-amino)-thiophene-2-carboxylic acid



A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-((4-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino)-thiophene-2-carboxylic acid methyl ester (63 mg, 0.141 mmol) and pyridine-2-carbonyl chloride HCl salt (33 mg, 0.184 mmol) in DCM (2 mL) was treated with DIEA (124 μ L, 0.7 mmol). After 30 min, NaHCO₃ (saturated aqueous solution, 4-8 mL) was added to the mixture, followed by brine (20 mL), and the crude product was extracted with ethyl acetate (2x20 mL). The combined organic layers were concentrated and the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (5 mL), treated with lithium hydroxide (42 mg, 1 mmol,) and heated to 60 deg C for 2 hours. The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile:water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 40 mg of the title compound as its TFA salt: MS (m/z): 536.9 [M-H]⁺; HPLC retention time: 4.17 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 141: Compound 141 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-[1-(pyrazine-2-carbonyl)-piperidin-4-yl]-amino]-thiophene-2-carboxylic acid




5

141

The title compound was synthesized in a manner analogous to Example 140, using pyrazine-2-carbonyl chloride in place of pyridine-2-carbonyl chloride HCl salt: MS (m/z): 538.0 [M-H]⁺; HPLC retention time: 3.52 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) 30 min run.

10

Example 142: Compound 142 - Synthesis of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-[1-(pyridine-3-carbonyl)-pyrrolidin-3-yl]-amino]-thiophene-2-carboxylic acid



15

142

A mixture of 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-pyrrolidin-3-yl-amino]-thiophene-2-carboxylic acid methyl ester (74 mg, 0.158 mmol) and pyridine-2-carbonyl chloride HCl salt (56 mg, 0.31 mmol) in DCM (2 mL) was treated with DIEA (124 μ L, 0.8 mmol). After 30 min, NaHCO₃ (saturated aqueous solution, 4-8 mL) was added to the mixture, followed by brine (20 mL), and the crude product was extracted with ethyl acetate (2x20 mL). The combined organic layers were concentrated and the crude material was dissolved in a 3:2:1 mixture of THF:MeOH:water (5 mL), treated with lithium hydroxide (42 mg, 1 mmol,) and heated


20

to 60 deg C for 2 hours. The residue was purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid). This resulted in 37 mg of the title compound as its TFA salt: MS (m/z): 522.0 [M-H]⁺; HPLC retention

5 time: 3.58 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid).

Example 164: Compound 164 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-(4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl)-amino)-thiophene-2-carboxylic acid

10



(4-trans-Amino-cyclohexyl)-carbamic acid tert-butyl ester (705mg, 2.8mmol) and triethylamine were dissolved in DCM (30mL) under an atmosphere of nitrogen and cooled to 0°C. After 10 minutes, pyridine-3-sulfonyl chloride (1.00g, 5.6mmol) was added. The reaction was allowed to warm to room temperature and was stirred

15

for 18h. The reaction was found complete by LC/MS. The reaction was washed with water (2x 10mL). The organic layer was dried with sodium sulfate and filtered. [4-trans-(pyridine-3-sulfonylamino)-cyclohexyl]-carbamic acid tert-butyl ester (927mg, 90%) was purified by precipitation into a white solid. LC/MS = 300 (M^+ -55)

5 [4-trans-(Pyridine-3-sulfonylamino)-cyclohexyl]-carbamic acid tert-butyl ester (1.08g, 3.04mmol) was dissolved in DMF (30mL) under an atmosphere of nitrogen. The solution was cooled to 0° C and Cs_2CO_3 (2.97g, 9.12mmol) was added. After 10 minutes, iodomethane (947 μL , 15.19mmol) was added. After 10 minutes, the reaction was allowed to warm to room temperature. The reaction was stirred for 2.5h
10 until found complete by LC/MS. The reaction was quenched with water and extracted with ethyl acetate (2x30mL). The combined organics were washed with a 5% LiCl solution in water, dried with sodium sulfate, filtered and concentrated. {4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl}-carbamic acid tert-butyl ester (741mg, 66%) was purified by silica gel chromatography to afford a white solid.

15 LC/MS = 314 (M^+ -55)

{4-trans-[Methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl}-carbamic acid tert-butyl ester (741mg, 2.01mmol) was dissolved in a 4N solution of HCl in dioxane (2.0mL, 8.00mmol). The reaction was allowed to stir for 30min until complete by LC/MS. The reaction was concentrated and used without purification. LC/MS = 269

20 (M^+)

5-(3,3-Dimethyl-but-1-ynyl)-3-iodo-thiophene-2-carboxylic acid methyl ester (469mg, 1.35mmol), BINAP (126mg, 0.2mmol), $\text{Pd}(\text{OAc})_2$ (55mg, 0.2mmol) and Cs_2CO_3 (2.20g, 6.75mmol) were combined in degassed toluene (15mL) under an atmosphere of argon. To the reaction was added the HCl salt of pyridine-3-sulfonic

25 acid (4-trans-amino-cyclohexyl)-methyl-amide (crude, 2.01mmol max). After 15 minutes, the reaction was heated to 100°C. After 17h, the reaction was complete by LC/MS. The reaction was cooled to room temperature and was filtered. The reaction was concentrated. 5-(3,3-dimethyl-but-1-ynyl)-3-{4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexylamino}-thiophene-2-carboxylic acid methyl ester (281mg, 42%)

30 was purified by silica gel chromatography to afford a yellow solid. LC/MS = 490 (M^+ +1)

5-(3,3-Dimethyl-but-1-ynyl)-3-{4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexylamino}-thiophene-2-carboxylic acid methyl ester (281mg, 0.57mmol) was dissolved in pyridine (5mL) under an atmosphere of nitrogen. To the reaction was

added 4-trans-methyl-cyclohexanecarbonyl chloride (138mg, 0.86mmol). The reaction was heated to 90°C for 20h. The reaction was cooled to room temperature and was diluted with EtOAc. The reaction was washed with water (2x5mL). The organic layer was dried with sodium sulfate, filtered and was concentrated. 5-(3,3-dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-{4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl}-amino)-thiophene-2-carboxylic acid methyl ester (105mg, 30%) was purified by silica gel chromatography to afford a yellow solid. LC/MS = 613 (M⁺)

5-(3,3-Dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-{4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl}-amino)-thiophene-2-carboxylic acid methyl ester (105mg, 0.18mmol) was dissolved in THF (1mL) and MeOH (0.5mL). To the reaction was added a solution of LiOH (20mg, 0.88mmol) in water (1mL). The reaction was stirred for 1h. The reaction was neutralized with 0.88mL of 1N HCl in water. The solution was concentrated. 5-(3,3-dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-{4-trans-[methyl-(pyridine-3-sulfonyl)-amino]-cyclohexyl}-amino)-thiophene-2-carboxylic acid (62mg, 57) was purified by HPLC to afford a white powder.

LC/MS = 597 (M⁺-2) Neg ionization

Retention time: 2.62 min

LC: Thermo Electron Surveyor HPLC


MS: Finnigan LCQ Advantage MAX Mass Spectrometer

Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

Example 165: Compound 165 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-trans-methyl-cyclohexanecarbonyl)-[4-(pyridine-3-sulfonylamino)-cyclohexyl]-amino)-thiophene-2-carboxylic acid



3-[(4-Amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid methyl ester (200mg, 0.44mmol) and TEA (245μL, 1.76mmol) were dissolved in DCM (3mL) under an atmosphere of nitrogen and cooled to 0°C. To the reaction was added pyridine-3-sulfonyl chloride (155mg, 0.88mmol). The reaction was allowed to warm to room temperature and was stirred for 30 minutes. The reaction was concentrated. 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(pyridine-3-sulfonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid methyl ester (50mg, 22%) was purified by silica gel chromatography to afford a yellow solid. LC/MS = 600 (M⁺+1)

5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(pyridine-3-sulfonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid methyl ester (50mg, 0.1mmol) was dissolved in a THF (1mL) and MeOH (1mL). To the reaction was added a solution of LiOH (12mg, 0.5mmol) in water (1mL). The reaction was stirred for 1 hour. The reaction was quenched with 1N HCl in water (0.5mL). 5-(3,3-Dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-[4-(pyridine-3-sulfonylamino)-cyclohexyl]-amino]-thiophene-2-carboxylic acid (23mg, 40%) was purified by HPLC to afford a white powder.

LC/MS = 584 (M⁺-1) Neg ionization

Retention time: 2.52 min

LC: Thermo Electron Surveyor HPLC


MS: Finnigan LCQ Advantage MAX Mass Spectrometer

Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.

- 5 Example 186: Compound 186 - Synthesis of 3-{4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-cyclohexylcarbamoyloxy}-azetidine-1-carboxylic acid tert-butyl ester



10 **186**

- 3-{4-[[2-Carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-methyl-cyclohexanecarbonyl)-amino]-cyclohexylcarbamoyloxy}-azetidine-1-carboxylic acid tert-butyl ester was prepared in a similar fashion to 4-[[2-carboxy-5-(3,3-dimethyl-but-1-ynyl)-thiophen-3-yl]-(4-trans-methyl-cyclohexanecarbonyl)-amino]-piperidine-1-carboxylic acid piperidin-4-yl ester using method A except that the HCl salt of 3-[(4-amino-cyclohexyl)-(4-trans-methyl-cyclohexanecarbonyl)-amino]-5-(3,3-dimethyl-but-1-ynyl)-thiophene-2-carboxylic acid was used instead of the HCl salt of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-trans-methyl-cyclohexanecarbonyl)-piperidin-4-yl-amino]-thiophene-2-carboxylic acid and 3-hydroxy-azetidine-1-carboxylic acid tert-butyl ester was used instead of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester.

LC/MS (m/z): 544 [M-99]

Retention time: 2.52min

LC: Thermo Electron Surveyor HPLC


MS: Finnigan LCQ Advantage MAX Mass Spectrometer

- 25 Column: Phenomenex Polar RP 30 mm X 4.6 mm

Solvents: Acetonitrile with 0.1% formic acid, Water with 0.1% formic acid

Gradient: 0 min-0.1 min 5% ACN, 0.1 min-1.95 min 5%-100% ACN, 1.95 min-3.5 min 100% ACN, 3.5 min-3.55 min 100%-5% ACN, 3.55 min-4 min 5% ACN.


Example 217: Compound 217 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-methyl-cyclohexanecarbonyl)-{4-[N'-methyl-N'-(tetrahydro-pyran-4-yloxy-carbonyl)-hydrazino]-cyclohexyl}-amino)-thiophene-2-carboxylic acid



To a solution of tetrahydro-pyran-4-ol (0.215 mL, 2.27 mmol) and N,N'-disuccinimidyl carbonate (0.87 g, 3.4 mmol) in ACN (7 mL) was added triethylamine (1 mL, 6.81 mmol). After 16 h stirring at rt, the reaction was quenched with LiCl solution (5% in water, 10 mL) and diluted with EtOAc (20 mL). The crude product was washed 2 X 10 mL with 5% LiCl solution, dried over Na₂SO₄ and concentrated. The crude material was used as is for the next step.

A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid (100 mg, 0.23 mmol), N-methyl-hydrazinecarboxylic acid tetrahydro-pyran-4-yl ester (70 mg, ~0.4 mmol), acetic acid (3 drops) in DCE (3 mL) was treated with NaBH(OAc)₃ (12 mg, 56 mmol) for 16 h. The reaction was quenched with water (10 mL) and extracted with EtOAc. The crude material was then purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile:water 13 min, both solvents containing 0.1% trifluoroacetic acid) giving the title compound as a TFA salt: MS (m/z): 603.1 [M-H]⁺; HPLC retention time: 27.4 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) 30 min run.

Example 218: Compound 218 - Synthesis of 5-(3,3-dimethyl-but-1-ynyl)-3-((4-methyl-cyclohexanecarbonyl)-{4-[N'-methyl-N'-(pyridin-2-yloxy-carbonyl)-hydrazino]-cyclohexyl}-amino)-thiophene-2-carboxylic acid



To a solution of carbonic acid di-2-pyridyl ester (300 mg, 1.38 mmol) in MeOH (10 mL) was added methylhydrazine (94 μ L, 1.8 mmol). After 16 h stirring at rt, the reaction was concentrated and the crude material was used as is for the next step.

10 A mixture of 5-(3,3-dimethyl-but-1-ynyl)-3-[(4-methyl-cyclohexanecarbonyl)-(4-oxo-cyclohexyl)-amino]-thiophene-2-carboxylic acid (100 mg, 0.23 mmol), N-methyl-hydrazinecarboxylic acid pyridin-2-yl ester (67 mg, ~0.4 mmol), acetic acid (3 drops) in DCE (3 mL) was treated with $\text{NaBH}(\text{OAc})_3$ (12 mg, 56 mmol) for 16 h. The reaction was quenched with water (10 mL) and extracted with EtOAc. The crude

15 material was then purified by HPLC (Gemini column, 35% acetonitrile:water, 2 min, 35-50% acetonitrile:water, 2 min, 50-100% acetonitrile: water 13 min, both solvents containing 0.1% trifluoroacetic acid) giving the title compound as a TFA salt: MS (m/z): 595.1 $[\text{M}-\text{H}]^+$; HPLC retention time: 28.9 min (2-98% acetonitrile:water with 0.05% trifluoroacetic acid) 30 min run.

20

Biological Examples

Antiviral Activity

Another aspect of the invention relates to methods of inhibiting viral infections,

comprising the step of treating a sample or subject suspected of needing such inhibition with a composition of the invention.

Within the context of the invention samples suspected of containing a virus include natural or man-made materials such as living organisms; tissue or cell
 5 cultures; biological samples such as biological material samples (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which induces a
 10 viral infection, frequently a pathogenic organism such as a tumor virus. Samples can be contained in any medium including water and organic solvent/water mixtures. Samples include living organisms such as humans, and man made materials such as cell cultures.

If desired, the anti-virus activity of a compound of the invention after
 15 application of the composition can be observed by any method including direct and indirect methods of detecting such activity. Quantitative, qualitative, and semiquantitative methods of determining such activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism are
 20 also applicable.

The antiviral activity of a compound of the invention can be measured using standard screening protocols that are known. For example, the antiviral activity of a compound can be measured using the following general protocols.

Cell-based Flavivirus Immunodetection assay

25 BHK21 or A549 cells are trypsinized, counted and diluted to 2×10^5 cells/mL in Hams F-12 media (A549 cells) or RPMI-1640 media (BHK21 cells) supplemented with 2% fetal bovine serum (FBS) and 1% penicillin/streptomycin. 2×10^4 cells are dispensed in a clear 96-well tissue culture plates per well and palced at 37°C , 5% CO_2 overnight. On the next day, the cells are infected with viruses at multiplicity of
 30 infection (MOI) of 0.3 in the presence of varied concentrations of test compounds for

1 hour at 37° C and 5% CO₂ for another 48 hours. The cells are washed once with PBS and fixed with cold methanol for 10 min. After washing twice with PBS, the fixed cells are blocked with PBS containing 1% FBS and 0.05% Tween-20 for 1 hour at room temperature. The primary antibody solution (4G2) is then added at a concentration of 1:20 to 1:100 in PBS containing 1% FBS and 0.05% Tween-20 for 3 hours. The cells are then washed three times with PBS followed by one hour incubation with horseradish peroxidase(HRP)-conjugated anti-mouse IgG (Sigma, 1:2000 dilution). After washing three times with PBS, 50 microliters of 3,3',5,5'-tetramethylbenzidine (TMB) substrate solution (Sigma) is added to each well for two minutes. The reaction is stopped by addition of 0.5 M sulfuric acid. The plates are read at 450 nm absorbance for viral load quantification. After measurement, the cells are washed three times with PBS followed by incubation with propidium iodide for 5 min. The plate is read in a Tecan Safire™ reader (excitation 537 nm, emission 617 nm) for cell number quantification. Dose response curves are plotted from the mean absorbance versus the log of the concentration of test compounds. The EC₅₀ is calculated by non-linear regression analysis. A positive control such as N-nonyl-deoxynojirimycin may be used.

Cell-based Flavivirus cytopathic effect assay

For testing against West Nile virus or Japanese encephalitis virus, BHK21 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in RPMI-1640 media supplemented with 2% FBS and 1% penicillin/streptomycin. For testing against dengue virus, Huh7 cells are trypsinized and diluted to a concentration of 4×10^5 cells/mL in DMEM media supplemented with 5% FBS and 1% penicillin/streptomycin. A 50 microliter of cell suspension (2×10^4 cells) is dispensed per well in a 96-well optical bottom PIT polymer-based plates (Nunc). Cells are grown overnight in culture medium at 37° C, 5% CO₂, and then infected with West Nile virus (e.g. B956 strain) or Japanese encephalitis virus (e.g. Nakayama strain) at MOI = 0.3, or with dengue virus (e.g. DEN-2 NGC strain) at MOI = 1, in the presence of different concentrations of test compounds. The plates containing the virus and the compounds are further incubated at 37°C, 5% CO₂ for 72 hours. At the end of incubation, 100 microliters of CellTiter-Glo™ reagent is added into each well. Contents are mixed for 2 minutes on an orbital shaker to induce cell lysis. The plates are incubated at room temperature for 10 minutes to stabilize luminescent signal.

Luminescence reading is recorded using a plate reader. A positive control such as N-nonyl-deoxyriboflavin may be used.

Antiviral Activity in a Mouse Model of Dengue Infection.

Compounds are tested *in vivo* in a mouse model of dengue virus infection (Schul *et al.* J. Infectious Dis. 2007; 195:665-74). Six to ten week old AG129 mice (B&K Universal Ltd, Hill, UK) are housed in individually ventilated cages. Mice are injected intraperitoneally with 0.4 mL TSV01 dengue virus 2 suspension. Blood samples are taken by retro orbital puncture under isoflurane anaesthesia. Blood samples are collected in tubes containing sodium citrate to a final concentration of 0.4%, and immediately centrifuged for 3 minutes at 6000g to obtain plasma. Plasma (20 microliters) is diluted in 780 microliters RPMI-1640 medium and snap frozen in liquid nitrogen for plaque assay analysis. The remaining plasma is reserved for cytokine and NS1 protein level determination. Mice develop dengue viremia rising over several days, peaking on day 3 post-infection.

For testing of antiviral activity, a compound of the invention is dissolved in vehicle fluid, e.g. 10% ethanol, 30% PEG 300 and 60% D5W (5% dextrose in water; or 6N HCl (1.5 eq):1N NaOH (pH adjusted to 3.5): 100 mM citrate buffer pH 3.5 (0.9% v/v:2.5% v/v: 96.6% v/v). Thirty six 6-10 week old AG129 mice are divided into six groups of six mice each. All mice are infected with dengue virus as described above (day 0). Group 1 is dosed by oral gavage of 200 μ L/mouse with 0.2 mg/kg of a compound of the invention twice a day (once early in the morning and once late in the afternoon) for three consecutive days starting on day 0 (first dose just before dengue infection). Groups 2, 3 and 4 are dosed the same way with 1 mg/kg, 5 mg/kg and 25 mg/kg of the compound, respectively. A positive control may be used, such as (2R,3R,4R,5R)-2-(2-amino-6-hydroxy-purin-9-yl)-5-hydroxymethyl-3-methyl-tetrahydro-furan-3,4-diol, dosed by oral gavage of 200 microliters/mouse the same way as the previous groups. A further group is treated with only vehicle fluid.

On day 3 post-infection approximately 100 microliter blood samples (anti-coagulated with sodium citrate) are taken from the mice by retro-orbital puncture under isoflurane anaesthesia. Plasma is obtained from each blood sample by centrifugation and snap frozen in liquid nitrogen for plaque assay analysis. The

collected plasma samples are analyzed by plaque assay as described in Schul *et al.* Cytokines are also analysed as as described by Schul. NS1 protein levels are analysed using a Platelia™ kit (BioRad Laboratories). An anti-viral effect is indicated by a reduction in cytokine levels and/or NS1 protein levels.

- 5 Typically, reductions in viremia of about 5-100 fold, more typically 10-60 fold, most typically 20-30 fold, are obtained with 5-50 mg/kg bid dosages of the compounds of the invention.

HCV Assay Protocol

- 10 The anti-HCV activity of the compounds of this invention was tested in a human hepatoma Huh-7 cell line harboring a HCV replicon. The assay comprised the following steps:

Step 1: compound preparation and serial dilution.

- 15 Serial dilution was performed in 100% DMSO in a 384-well plate. A solution containing a compound at 225-fold concentration of the starting final serial dilution concentration was prepared in 100% DMSO and 15 uL added to the pre-specified wells in column 3 or 13 of a polypropylene 384-well plate. The rest of the 384-well plate was filled with 10 uL 100% DMSO except for columns 23 and 24, where 10 uL of 500 uM a HCV protease inhibitor (ITMN-191) in 100% DMSO was added. The
20 HCV protease inhibitor was used a control of 100% inhibition of HCV replication. The plate was then placed on a Biomek FX Workstation to start the serial dilution. The serial dilution was performed for ten cycles of 3-fold dilution from column 3 to 12 or from column 13 to 22.

Step 2: cell culture plate preparation and compound addition

- 25 To each well of a black polypropylene 384-well plate, 90 µL of cell media containing 1600 suspended Huh-7 HCV replicon cells was added with a Biotek uFlow Workstation. A volume of 0.4 µL of the compound solution was transferred from the serial dilution plate to the cell culture plate on a Biomek FX Workstation. The DMSO concentration in the final assay condition was 0.44%. The plates were incubated for 3
30 days at 37 °C with 5% CO2 and 85% humidity.

Step 3: detection of cytotoxicity and inhibition of viral replication

- a) Assessment of cytotoxicity: The media in the 384-well cell culture plate was aspirated with a Biotek EL405 plate-washer. A volume of 50 µL of a solution

containing 400 nM Calcein AM in 100% PBS was added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated for 30 minutes at room temperature before the fluorescence signal (emission 490 nm, excitation 520 nm) was measured with a Perkin Elmer Envision Plate Reader.

- 5 b) Assessment of inhibition of viral replication: The calcein-PBS solution in the 384-well cell culture plate was aspirated with a Biotek EL405 plate-washer. A volume of 20 μ L of Dual-Glo luciferase buffer (Promega, Dual-Glo Luciferase Assay Reagent, cat. #E298B) was added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated for 10 minutes at room temperature. A volume of 20 μ L of a
- 10 solution containing 1:100 mixture of Dual-Glo Stop & Glo substrate(Promega, Dual-Glo Luciferase Assay Reagent, cat. #E313B) and Dual-Glo Stop & Glo buffer (Promega, Dual-Glo Luciferase Assay Reagent, cat. #E314B) was then added to each well of the plate with a Biotek uFlow Workstation. The plate was incubated at room temperature for 10 minutes before the luminescence signal was measured with
- 15 a Perkin Elmer Envision Plate Reader.

Step 4: calculation

- The percent cytotoxicity was determined by calcein AM conversion to fluorescent product. The average fluorescent signal from the DMSO control wells were defined as 100% nontoxic. The individual fluorescent signal from testing
- 20 compound treated well was divided by the average signal from DMSO control wells and then multiplied by 100% to get the percent viability. The percent anti-HCV replication activity was determined by the luminescence signal from the testing well compared to DMSO controls wells. The background signal was determined by the average luminescence signal from the HCV protease inhibitor treated wells and was
- 25 subtracted from the signal from the testing wells as well as the DMSO control wells. Following 3-fold serial dilutions, the EC₅₀ and CC₅₀ values were calculated by fitting % inhibition at each concentration to the following equation:

$$\% \text{ inhibition} = 100\% / [(EC_{50}/[I])^b + 1]$$

- Where b is Hill's coefficient. See, for reference, Hill, A. V., *The Possible*
- 30 *Effects of the Aggregation of the Molecules of Hæmoglobin on its Dissociation Curves*, J. Physiol. **40**: iv-vii. (1910).

% inhibition values at a specific concentration, for example 2 μ M, can also be derived from the formula above.

When tested, certain compounds of this invention were found to inhibit viral replication as listed in Table 1:

Table 1

5

Compound	% inhibition at 2 μ M
31	93.1
32	100
33	96.2
34	97.8
35	99.8
36	100
37	99.6
38	99.6
39	98.8
40	98.6
41	99.9
42	99.9
43	96.2
44	100
45	100
46	97.0
47	99.9
48	100
49	99.2
50	99.9
51	85.5
52	93.5
53	99.9
54	99.8
55	87.9
56	92.8
57	78.2

64	100
65	99.9
66	99.9
67	99.0
68	98.5
69	99.5
70	99.1
71	95.1
72	99.7
73	93.1
74	99.7
75	97.5
76	99.4
77	99.9
79	100
80	99.8
81	100
82	100
83	99.9
112	99.9
113	98.1
114	100
115	99.8
116	99.9
117	99.9
118	100
119	97.1
120	99.9
121	99.9
122	99.7
123	99.9
124	99.9
125	99.7

126	98.9
127	98.3
128	99.8
129	100
130	98.7
131	99.9
132	77.4
133	96.7
134	99.0
135	99.8
136	98.9
137	100
140	99.9
141	99.9
142	99.6
164	98.9
165	95.7
186	98.9
217	100
218	100


Preferred compounds according to Table 1 include Examples 32, 36, 38, 41,48, 50, 64, 65, 66, 77, 79, 80, 81, 82, 83,112, 114, 118, 129, 217, and 218.

5 The specific pharmacological responses observed may vary according to and depending on the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with practice of the present invention.

10 Although specific embodiments of the present invention are herein illustrated and described in detail, the invention is not limited thereto. The above detailed descriptions are provided as exemplary of the present invention and should not be construed as constituting any limitation of the invention.

WHAT IS CLAIMED IS:

1. A compound of Formula I:



Formula (I),

or a pharmaceutically acceptable salt or ester thereof, wherein:

R^1 is C_{1-12} alkyl;

R^2 is optionally substituted C_{3-12} cycloalkyl;

wherein, each substituted R^2 is substituted with one or more Q^2 ,

each Q^2 , independently, is selected from the group consisting of $-S(O)_2R^{20}$, optionally substituted C_{1-6} alkyl, OH, $-C(O)OR^{20}$, and $-CN$;

each R^{20} , independently, is selected from the group consisting of H, and C_{1-12} alkyl;

Y is $-R^3-L-Het$, $-N(R^4)(R^5)$ or $-R^6=NOR^7$;

R^3 is selected from the group consisting of C_{1-12} alkylene, C_{3-12} cycloalkylene, and 3-12 membered heterocyclylene, wherein said 3-12 membered heterocyclylene comprises one to four heteroatoms selected from O, S, or N;

L is selected from the group consisting of $-OC(O)N(R^4)-$, $-N(R^4)C(O)O-$, $-N(R^4)S(O)_2-$, $-N(R^4)C(O)-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-N(R^4)N(R^4)C(O)O-$, and $-N(R^4)N(R^4)-$;

Het is an optionally substituted 3-12 membered heterocyclyl or optionally substituted 5-10 membered heteroaryl, wherein said optionally substituted 3-12

membered heterocyclyl or optionally substituted 5-10 membered heteroaryl comprises one to four heteroatoms selected from O, S, or N;

wherein, each substituted Het is substituted with one or more Q⁴;

each Q⁴, independently, is selected from the group consisting of halogen, oxo, C₁₋₆ alkyl, and -C(O)OR⁴⁰;

R⁴⁰ is C₁₋₁₂ alkyl;

each R⁴ is independently selected from the group consisting of H, C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

each R⁵ is independently selected from the group consisting of C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C_{1-C12} alkyl, C_{3-C12} cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q¹, wherein said 5-10 membered heteroaryl and said 4-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

each Q¹ is independently selected from the group consisting of C₁₋₆ alkyl, 5-10 membered heteroaryloxy, 4-12 membered heterocyclyloxy, 3-10 membered heterocyclyl, -OH, and -C(O)OR¹⁰, wherein said heteroaryl portion of said 5-10 membered heteroaryloxy, said heterocyclyl portion of said 4-12 membered heterocyclyloxy, and said 3-10 membered heterocyclyl each comprises one to four heteroatoms selected from O, S, or N;

R¹⁰ is selected from the group consisting of H, and C₁₋₁₂ alkyl;

R⁶ is C_{3-C12} cycloalkylene;

R⁷ is selected from the group consisting of optionally substituted C₁₋₁₂ alkyl, optionally substituted C₆₋₁₄ aryl, optionally substituted 6-11 membered

heteroarylalkyl, and optionally substituted C₇₋₁₁ arylalkyl, wherein said heteroaryl portion of said 6-11 membered heteroarylalkyl comprises one to four heteroatoms selected from O, S, or N;

wherein, each substituted R⁷ is substituted with one or more Q²; and

wherein:


the term alkyl refers to a hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms; and

the term cycloalkyl refers to a saturated or partially unsaturated ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to 20 carbon atoms as a polycycle.

2. The compound of claim 1, wherein R¹ is C₃-C₇ secondary or tertiary alkyl.

3. The compound of claim 1 or 2, wherein R² is optionally substituted methylcyclohexyl or optionally substituted methylcyclohexenyl.

4. The compound of any one of claims 1 to 3, represented by Formula II:




Formula II


or a pharmaceutically acceptable salt or ester thereof, wherein

R² is optionally substituted 4-methylcyclohexyl or optionally substituted 4-methylcyclohexenyl.

5. The compound of any one of claims 1 to 4, wherein R² is:



6. The compound of any one of claims 1 to 4, wherein R² is:



5

7. The compound of any one of claims 1 to 6, wherein Y is -R³-L-Het.

8. The compound of any one of claims 1 to 7, wherein R³ is selected from the group consisting of C₁₋₆ alkylene, C₄₋₆ cycloalkylene, and 5-6 membered heterocyclylene.

10

9. The compound of any one of claims 1 to 8, wherein Het is selected from the group consisting of optionally substituted pyridinyl, optionally substituted pyridazinyl, optionally substituted tetrahydro-2H-pyranyl, optionally substituted piperidinyl, optionally substituted pyrrolidinyl, optionally substituted tetrahydrothiophenyl, optionally substituted pyrazinyl, optionally substituted 1H-tetrazolyl, optionally substituted azetidiny, optionally substituted tetrahydrofuranyl, optionally substituted tetrahydro-2H-furo[2,3-b]furanyl, optionally substituted thiazoyl, optionally substituted 1H-imidazolyl, optionally substituted 4H-1,2,4-triazolyl, optionally substituted 1H-pyrazolyl, optionally substituted 1,3,4-thiadiazolyl, optionally substituted quinolinyl, optionally substituted [1,2,4]triazolo[4,3-a]pyridinyl, optionally substituted thiophenyl, optionally substituted 1,2,4-thiadiazolyl, optionally substituted

15

20

pyrimidinyl, optionally substituted 1*H*-1,2,3-triazolyl, optionally substituted 1,3,4-oxadiazolyl, and optionally substituted imidazo[1,2-*b*]pyridazinyl.

10. The compound of any one of claims 1 to 6, wherein Y is $-N(R^4)(R^5)$.

5

11. The compound of any one of claims 1 to 6 and 10, wherein R^4 is selected from the group consisting of H and optionally substituted C_1 - C_6 alkyl.

12. The compound of any one of claims 1 to 6, 10 and 11, wherein R^5 is selected from the group consisting of C_1 - C_6 alkyl, C_5 - C_6 cycloalkyl, 5-10 membered heteroaryl and 4-10 membered heterocyclyl wherein each C_1 - C_6 alkyl, C_5 - C_6 cycloalkyl, 5-10 membered heteroaryl or 4-10 membered heterocyclyl is optionally substituted with one or more Q^1 .

10


13. The compound of any one of claims 1 to 6, wherein Y is $-R^6=NOR^7$.

15

14. The compound of any one of claims 1 to 6 and 13, wherein R^6 is cyclohexyne.


15. The compound of claim 1, which is:

20




5


153





5




5 16. The compound of claim 1, which is:











or a pharmaceutically acceptable salt or ester thereof.


- 5 17. The compound of claim 1, which is:









18. The compound of claim 1, which is:



5





or a pharmaceutically acceptable salt or ester thereof.

5

19. A pharmaceutical composition, comprising the compound of any one of claims 1 to 18, and a pharmaceutically acceptable carrier or excipient.

10

20. A pharmaceutical composition, comprising a therapeutically effective amount of the compound of any one of claims 1 to 18, and a pharmaceutically acceptable carrier or excipient, for use in the treatment of HCV.


21. The pharmaceutical composition of claim 19 and 20, further comprising at least one additional agent selected from the group consisting of interferons, ribavirin

or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, and mixtures thereof.

22. Use of the compound of any one of claims 1 to 18, for the treatment of a *Flaviviridae* viral infection.

23. Use of the compound of any one of claims 1 to 18, for the treatment of HCV.

24. Use of the compound of any one of claims 1 to 18, in combination with at least one additional agent selected from the group consisting of interferons, ribavirin or its analogs, HCV NS3 protease inhibitors, NS5a inhibitors, alpha-glucosidase 1 inhibitors, hepatoprotectants, mevalonate decarboxylase antagonists, antagonists of the renin-angiotensin system, nucleoside or nucleotide inhibitors of HCV NS5B polymerase, non-nucleoside inhibitors of HCV NS5B polymerase, HCV NS5A inhibitors, TLR-7 agonists, cyclophilin inhibitors, HCV IRES inhibitors, and mixtures thereof, for the treatment of HCV.



Formula (I)