

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
18.10.2006 Bulletin 2006/42

(51) Int Cl.:
A61K 8/02^(2006.01) A61K 8/365^(2006.01)
A61K 8/892^(2006.01)

(21) Application number: **00951710.3**

(86) International application number:
PCT/GB2000/003031

(22) Date of filing: **07.08.2000**

(87) International publication number:
WO 2001/013880 (01.03.2001 Gazette 2001/09)

(54) **TOWELETTE PRODUCT**
KOSMETIKTUCH
ESSUIE-DOIGTS

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: **24.08.1999 US 150481 P**

(43) Date of publication of application:
29.05.2002 Bulletin 2002/22

(73) Proprietors:
• **UNILEVER PLC**
London EC4P 4BQ (GB)
Designated Contracting States:
CY GB IE
• **UNILEVER N.V.**
3013 AL Rotterdam (NL)
Designated Contracting States:
AT BE CH DE DK ES FI FR GR IT LI LU MC NL PT SE

• **SLAVTCHEFF, Craig**
Unilever Home & Pers. Care USA
Trumbull, CT 06611 (US)
• **ZNAIDEN, Alexander**
Nippon Lever B.V.
Haga-gun
Tochigi-ken 321-33 (JP)

(74) Representative: **Acham, Nicholas Clive et al**
Unilever Patent Group
Colworth House
Sharnbrook
Bedford, MK44 1LQ (GB)

(56) References cited:
EP-A- 0 875 233 WO-A1-00/44343
DE-A- 1 467 886 DE-A- 19 903 717
US-A- 3 795 624

(72) Inventors:
• **GOTT, Robert E.**
Unilever Home & Pers. Care USA
Trumbull, CT 06611 (US)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention

[0001] The present invention relates to single use towelettes for cosmetically delivering alpha-hydroxycarboxylic acids, selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof.

10 The Related Art

[0002] Alpha-hydroxycarboxylic acids and their derivatives are widely known as providing treatment for the maintenance of a youthful look. These substances are said to control the development of facial fine lines and wrinkles. However, formulation of these substances has been difficult. Among the problems that have been encountered are compatibility with carrier systems, physical stability and skin irritation. In addition, it is particularly difficult to formulate low pH systems.

[0003] U.S. Patent 5,091,171 (Yu et al.) was one of the first documents to describe the use of alpha-hydroxycarboxylic acids as being effective against the appearance of fine lines and wrinkles. Subsequent thereto a vast amount of literature and many commercial products have been generated based on the efficacy of these materials. Most of the formulations described to date have been of either the cream or lotion type. One problem with these formulations is that they do not always evenly distribute over the applied surfaces. Secondly, any time an active treated surface is followed by cleansing, the active is washed away. Hence, methods of maintaining alpha-hydroxycarboxylic acids on a skin surface are needed which are not susceptible to subsequent cleansing actions. Irritation has also been of great concern because the formulations generally have a low pH.

[0004] WO 96/11572 (Moberg) has utilised a variety of acids, including the alpha-hydroxy substance known as lactic acid, in an aqueous hexylene glycol formulation which can be impregnated onto textiles or refreshing napkins. These were employed in order to overcome the problem of microbial growth on skin and also act as a disinfectant.

[0005] U.S. Patent 4,828,912 (Hossain et al.) and U.S. Patent 4,764,418 (Kuenn et al.) describe virucidal tissue products directed at controlling disease inducing organisms such as viruses and the common cold. Carboxylic acids such as citric, malic, succinic and benzoic acids are formulated with surfactants and a carrier for impregnation onto facial tissues or other non-woven materials.

[0006] None of the above disclosures have addressed the issue of stickiness of the alpha-hydroxycarboxylic acids once they have been deposited onto the skin by the towelette. Neither has there been any discussion with respect to odour control which may arise in low pH systems.

[0007] Accordingly, it is an object of the present invention to provide a product and a method for both cleansing skin and reducing the signs of aging.

[0008] Another object of the present invention is to provide a product and a method for depositing alpha-hydroxycarboxylic acids selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof in a manner which avoids the formation of sticky residues on the skin.

[0009] Yet another object of the present invention is to provide a product for delivering alpha-hydroxycarboxylic acids selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof in a formulation which avoids the formation of foul odours.

[0010] These objects of the present invention will become more apparent from the following summary and detailed discussion which follow.

45 SUMMARY OF THE INVENTION

[0011] According to the present invention there is provided a towelette product which includes:

(a) a water-insoluble substrate;

50 (b) a cosmetic composition impregnated into the substrate including:

(i) an alpha-hydroxycarboxylic acid selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof;

(ii) a silicone microemulsion;

55 (iii) the composition in water having a pH of no higher than about 6.5.

[0012] Silicone microemulsions of the present invention provide both stability to the composition and counteract any stickiness which may occur when the alpha-hydroxycarboxylic acids are deposited onto the skin.

[0013] Surfactants, especially mild surfactants such as those of the amphoteric type may decompose at low pH emitting foul odours. The silicone microemulsions of the present invention have also been found to be useful in counteracting the generation of foul odours.

5 DETAILED DESCRIPTION OF THE INVENTION

[0014] The applicants have now surprisingly found that towelettes impregnated with alpha-hydroxycarboxylic acids selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof, can be delivered to the skin without imparting any stickiness. This is achieved by the use of silicone microemulsions.

10 [0015] Typically creams, lotions and other types of cosmetic vehicles containing the alpha-hydroxycarboxylic acids also contain emollients which are deposited on the skin along with the other actives. These emollients include, for example, esters, hydrocarbons or dimethicone oils. The presence of the emollients obscures any stickiness which may result from evaporation of water or the carrier leaving dried active behind.

15 [0016] Unfortunately, towelette products cannot be impregnated with fluids with too high viscosity as the towelettes will not adequately wet out when the fluid to be impregnated is too thick. However, low viscosity fluids which must be used with towelettes are susceptible to poor emulsion stability and large amounts of emollients are difficult to formulate into these systems. Consequently, actives such as alpha-hydroxycarboxylic acids deposit in a sticky manner onto skin from the low viscosity fluids spread thereon by the towelettes.

20 [0017] A first necessary aspect of the present invention is that of a substrate. Preferably the substrate is a water insoluble substance. By "water insoluble" is meant that the substrate does not dissolve in or readily break apart upon immersion in water. Another advantage of the substrate in combination with the active is that the former helps the active penetrate. The substrate is also much better than a mere liquid or gel formulation enabling for more accurate application to the skin and the avoidance of sensitive areas, such as the eye where inadvertent application of the astringent composition would cause irritation.

25 [0018] A wide variety of materials may be used as the substrate. The following non-limiting characteristics are desirable: (i) sufficient wet strength for use, (ii) sufficient abrasivity, (iii) sufficient loft and porosity, (iv) sufficient thickness, (v) appropriate size, and (vi) non-reactive with components of the impregnating composition.

30 [0019] Nonlimiting examples of suitable substrates which meet the above criteria include nonwoven substrates, woven substrates, hydroentangled substrates, and air entangled substrates. Preferred embodiments employ nonwoven substrates since they are economical and readily available in a variety of materials. By "nonwoven" is meant that the layer is comprised of fibers which are not woven into a fabric but rather are formed into a sheet, particularly a tissue. The fibers can either be random (i.e., randomly aligned) or they can be carded (i.e. combed to be oriented in primarily one direction). Furthermore, the nonwoven substrate can be composed of a combination of layers of random and carded fibers.

35 [0020] Nonwoven substrates may be comprised of a variety of materials both natural and synthetic. By natural is meant that the materials are derived from plants, animals, insects or byproducts. By synthetic is meant that the materials are obtained primarily from various man-made materials or from material that is usually a fibrous web comprising any of the common synthetic or natural textile-length fibers, or mixtures thereof.

40 [0021] Nonlimiting examples of natural materials useful in the present invention are silk fibers, keratin fibers and cellulosic fibers. Nonlimiting examples of keratin fibers include those selected from the group comprising wool fibers, and camel hair fibers. Nonlimiting examples of cellulosic fibers include those selected from the group comprising wood pulp fibers, cotton fibers, hemp fibers, jute fibers, flax fibers, and mixtures thereof. Wood pulp fibers are preferred while all cotton fibers (e.g. cotton pads) are normally avoided.

45 [0022] Nonlimiting examples of synthetic materials useful in the present invention include those selected from the group comprising acetate fibers, acrylic fibers, cellulose ester fibers, modacrylic fibers, polyamide fibers, polyester fibers, polyolefin fibers, polyvinyl alcohol fibers, rayon fibers and mixtures thereof. Examples of some of these synthetic materials include acrylics such as Acrilan®, Creslan®, and the acrylonitrile-based fiber, Orlon®; cellulose ester fibers such as cellulose acetate, Arnel® and Acele®; polyamides such as Nylons (e.g., Nylon 6, Nylon 66, Nylon 610 and the like); polyesters such as Fortrel® Kodel® and the polyethylene terephthalate fibers, Dacron® polyolefins such as polypropylene, polyethylene; polyvinyl acetate fibers and mixtures thereof.

50 [0023] Nonwoven substrates made from natural materials consist of webs or sheets most commonly formed on a fine wire screen from a liquid suspension of the fibers.

[0024] Substrates made from natural materials useful in the present invention can be obtained from a wide variety of commercial sources. Nonlimiting examples of suitable commercially available paper layers useful herein include Airtex® an embossed airlaid cellulosic layer having a base weight of about 85 g/m² (71 gsy), available from James River Corporation, Green Bay, WI; and Walkisoft®, an embossed airlaid cellulosic having a base weight of about 90 g/m² (75 gsy), available from Walkisoft U.S.A., Mount Holly, NC.

55 [0025] Nonwoven substrates made from synthetic materials useful in the present invention can also be obtained from a wide variety of commercial sources. Nonlimiting examples of suitable nonwoven layer materials useful herein include

HEF 40-047, an apertured hydroentangled material containing about 50% rayon and 50% polyester, and having a basis weight of about 51 g/m² (43 grams per square yard (gsy)), available from Veratec, Inc., Walpole, MA; HEF 140-102, an apertured hydroentangled material containing about 50% rayon and 50% polyester, and having a basis weight of about 67 g/m² (56 gsy) available from Veratec, Inc., Walpole, MA; Novenet® 149-191, a thermo-bonded grid patterned material containing about 69% rayon, about 25% polypropylene, and about 6% cotton, and having a basis weight of about 120 g/m² (100 gsy) available from Veratec, Inc., Walpole, MA; HEP Nubtex® 149-801, a nubbed, apertured hydroentangled material, containing about 100% polyester, and having a basis weight of about 84 g/m² (70 gsy), available from Veratec, Inc. Walpole, MA; Keybak® 951V, a dry formed apertured material, containing about 75% rayon, about 25% acrylic fibers, and having 2 basis weight of about 52 g/m² (43 gsy), available from Chicopee Corporation, New Brunswick, NJ; Keybak® 1368, an apertured material, containing about 75% rayon, about 5% polyester, and having a basis weight of about 47 g/m² (39 gsy), available from Chicopee Corporation, New Brunswick, NJ; Duralace® 1236, an apertured, hydroentangled material, containing about 100% rayon, and having a basis weight from about 48 g/m² (40 gsy) to about 138 g/m² (115 gsy), available from Chicopee Corporation, New Brunswick, NJ; Duralace® 5904, an apertured, hydroentangled material, containing about 100% polyester, and having a basis weight from about 48 g/cm² (40 gsy) to about 138 g/m² (115 gsy), available from Chicopee Corporation, New Brunswick, NJ; Sontaro® 8868, a hydroentangled material, containing about 50% cellulose and about 50% polyester, and having a basis weight of about 72 g/m² (60 gsy), available from Dupont Chemical Corp.

[0026] For the purposes of the present invention the most preferred towelettes are non-woven substrates, especially blends of rayon/polyester in ratios of 10:90 to 90:10, preferably ratios of 20:80 to 80:20, optimally 40:60 to 60:40 by weight. The most preferred towelette is a 70:30 rayon/polyester non-woven wipe article.

[0027] The substrate can be made into a wide variety of shapes and forms. Generally the substrate is in single use towelette form. Advantageously, the towelettes are folded in a Z-shaped formation. They may be interleaved with one another but preferably are not interleaved. The Z-fold consists of a center panel flanked by upper and lower wing panels.


[0028] The upper and lower wing panels are substantially of equal width and substantially half a width of the center panel. Each towelette is folded medially in a direction orthogonal to that of the Z-shaped formation. Advantageously the size of the towelette may range in length from 10 to 40 cm, preferably from 15 to 30 cm, optimally from 18 to 24 cm. The width of the towelette may range from 8 to 30 cm, preferably from 10 to 25 cm, optimally from 15 to 20 cm.

[0029] Anywhere from 5 to 100, preferably from 10 to 50 single towelettes may be stored within a dispensing pouch, preferably a moisture impermeable pouch. During storage and between dispensing, the pouch is resealable, usually via an adhesive strip covering a dispensing opening. Single towelette containing pouches may also be employed.

[0030] The substrates of the present invention may optionally comprise two or more layers, each having a different texture and abrasiveness. The differing textures can result from the use of different combinations of materials or from the use of a substrate having a more abrasive side for exfoliation and a softer, absorbent side for gentle cleansing. In addition, separate layers of the substrate can be manufactured to have different colors, thereby helping the user to further distinguish the surfaces.

[0031] A second important element of the present invention is the alpha-hydroxycarboxylic acid selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof. By this term is meant not only the acid form but also salts thereof. Typical cationic counterions to form the salt are the alkali metals, alkaline earth metals, ammonium, C2-C8 trialkanolammonium cation and mixtures thereof.

[0032] Alpha-hydroxyacids are organic carboxylic acids in which one hydroxyl group is attached to the alpha carbon adjacent the carboxy group. The generic structure is as follows:



wherein Ra and Rb are independently an H, F, Cl, Br, alkyl, aralkyl or an aryl group and saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms. In addition Ra and Rb may optionally be substituted by OH, CHO, COOH and alkoxy groups having 1 to 9 carbon atoms. The alpha-hydroxyacids may exist as stereoisomers as D, L, and DL forms when Ra and Rb are not identical.

[0033] Typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, lauryl, stearyl, benzyl and phenyl, etc. The alpha-hydroxyacids of the first group may be subdivided into (1) alkyl alpha-hydroxyacids, (2) aralkyl and aryl alpha-hydroxyacids, (3) polyhydroxy alpha-hydroxyacids, and (4) polycarboxylic alpha-hydroxyacids. The following are representative alpha hydroxyacids in each subgroup:

(1) Alkyl Alpha Hydroxyacids

2-Hydroxyethanoic acid (Glycolic acid, hydroxyacetic acid)

2-Hydroxypropanoic acid (Lactic acid)

2-Methyl 2-hydroxypropanoic acid (Methylactic acid)

2-Hydroxybutanoic acid

2-Hydroxypentanoic acid
 2-Hydroxyhexanoic acid
 2-Hydroxyheptanoic acid
 2-Hydroxyoctanoic acid
 5 2-Hydroxynonanoic acid
 2-Hydroxydecanoic acid
 2-Hydroxyundecanoic acid
 2-Hydroxydodecanoic acid (Alpha hydroxylauric acid)
 2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid)
 10 2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid)
 2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid)
 2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid)

(2) Aralkyl And Aryl Alpha-Hydroxyacids

15 2-Phenyl 2-hydroxyethanoic acid (Mandelic acid)
 2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid)
 3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid)
 2-Phenyl 2-methyl 2-hydroxyethanoic acid (Atrolactic acid)
 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid (4-Hydroxymandelic acid)
 20 2-(4'-Chlorophenyl)-2-hydroxyethanoic acid (4-Chloromandelic acid)
 2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic acid (3-Hydroxy-4-methoxymandelic acid)
 2-(4'-Hydroxy-3'-methoxyphenyl acid)
 3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid [3(2'-Hydroxyphenyl) lactic acid]
 3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid [3-(4'-Hydroxyphenyl) lactic acid]
 25 2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid (3,4-Dihydroxymandelic acid)

(3) Polyhydroxy Alpha-Hydroxyacids

2,3-Dihydroxypropanoic acid (Glyceric acid)
 2,3,4-Trihydroxybutanoic acid, Isomers; erythronic acid, threonic acid)
 30 2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic acid, arabinoic acid, xylonic acid, lyxonic acid)
 2,3,4,5,6-Pentahydroxyhexanoic acid (Isomers; allonic acid, altronic acid, gluconic acid, mannoic acid, gulonic acid,
 idonic acid, galatronic acid, talonic acid)
 2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid, galactoheptonic acid etc.)

(4) Polycarboxylic Alpha-Hydroxyacids

2-Hydroxypropane-1,3-dioic acid (Tartronic acid)
 2-Hydroxybutane, 1,4-dioic acid (Malic acid)
 2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid)
 2-Hydroxy-2-carboxypentane, 1,5-dioic acid (Citric acid)
 40 2,3,4,5-Tetrahydroxyhexane, 1-5, dioic acid (Isomers: saccharic acid, mucic acid)

(5) Lactone Forms

45 **[0034]** The typical lactone forms are gluconolactone, galactonolactone, glucuronolactone, glacturonolactone, gluconolactone, ribonolactone, saccharic acid lactone, pantoyllactone, glucoheptonolactone, mannonolactone, and galactohettonolactone.

[0035] Alpha ketoacids are as follows.

2-Ketoethanoic acid (Glyoxylic acid)
 Methyl 2-ketoethanoate
 50 2-Ketopropanoic acid (Pyruvic acid)
 Methyl 2-ketopropanoate (Methyl pyruvate)
 Ethyl 2-ketopropanoate (Ethyl pyruvate)
 Propyl 2-ketopropanoate (Propyl pyruvate)
 2-Phenyl-2-ketoethanoic acid (Benzoylformic acid)
 55 Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate)
 Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate)
 3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid)
 Methyl 3-phenyl-2-ketopropanoate (Methyl phenylpyruvate)

Ethyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate)

2-Ketobutanoic acid

2-Ketopentanoic acid

2-Ketohexanoic acid

5 2-Ketoheptanoic acid

2-Ketooctanoic acid

2-Ketododecanoic acid

Methyl 2-ketooctanoate

10 II. Dimeric and Polymeric Forms of Hydroxyacids

[0036] When two or more molecules of hydroxycarboxylic acids, either identical or non-identical compounds, are reacted chemically to each other, dimeric or polymeric compounds will be formed. Such dimeric and polymeric compounds may be classified into three groups, namely (a) acyclic ester, (b) cyclic ester and (c) miscellaneous dimer and polymer.

15 **[0037]** Acyclic esters of hydroxycarboxylic acids are

Glycolyl glycollate (Glycolic acid glycollate)

Lactyl lactate (Lactic acid lactate)

Mandelyl mandellate

Atrolactyl atrolactate

20 Phenyllactyl phenyllactate

Benzilyl benzillate

Glycolyl lactate

Lactyl glycollate

Glycolyl glycolyl glycollate

25 Lactyl lactyl lactate

Lactyl glycolyl lactate

Glycolyl glycolyl glycolyl glycollate

Lactyl lactyl lactyl lactate

Glycolyl lactyl glycolyl lactyl glycollate

30 Polyglycolic acid and polylactic acid

[0038] Amounts of the alpha-hydroxycarboxylic acids selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof, may range from about 0.01 to about 20%, preferably from about 0.1 to about 15%, more preferably from about 1 to about 10%, optimally from about 3 to about 8% by weight of the composition which impregnates the substrate.

35 **[0039]** Compositions of the present invention when placed in water will have a pH no higher than about 6.5, preferably from about 6.0 to about 2.0, more preferably from about 5.5 to about 2.5, even more preferably from about 5.0 to about 3.0, optimally from about 4.5 to about 3.5. The compositions may be placed on the towelettes in a dry state and activated by a consumer through moistening with water. In a preferred embodiment compositions of the present invention are pre-prepared fluid compositions of low viscosity. Typical viscosities may range from 0.5 to 100 cps, preferably from about 40 2 to about 20 cps at 200C (Brookfield RVT).

[0040] The amount of impregnating composition relative to the substrate may range from about 20:1 to 1:20, preferably from 10:1 to about 1:10 and optimally from about 2:1 to about 1:2 by weight.

[0041] Typically a humectant is incorporated with compositions of the present invention. Humectants are normally polyols. Representative polyols include glycerin, diglycerin, polyalkylene glycols and more preferably alkylene polyols and their derivatives including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,2-butylene glycol, 1,2,6-hexanetriol, isoprene glycol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. The most preferred is 2-methyl-1,3-propanediol available as MP Diol from the Arco Chemical Company. Amounts of the polyol may range from about 0.5 to about 95%, preferably from about 1 to about 50%, more preferably from about 1.5 to 20%, optimally from about 3 to about 10% by weight of the impregnating composition.

[0042] An essential further element of product according to the present invention is that of a silicone microemulsion. Average particle size of the silicone material in these microemulsions may range from about 0.01 nm to about 500 nm, preferably from about 1 to about 100 nm, optimally from about 5 to about 50 nm. Particle size may be measured by means of a laser light scattering technique, using a 2600 D Particle Sizer from Malvern Instruments.

55 **[0043]** The microemulsions may be prepared by high shear mechanical mixing of the silicone and water, or by emulsifying the insoluble, non-volatile silicone with water and an emulsifier-mixing the silicone into a heated solution of the emulsifier for instance, or by a combination of mechanical and chemical emulsification.

[0044] Any surfactant materials either alone or in admixture may be used as emulsifiers in the preparation of the

silicone emulsions. Preferred emulsifiers include anionic emulsifiers such as alkylarylsulphonates, e.g. sodium dodecylbenzene sulphonate, alkyl sulphates e.g. sodium lauryl sulphate, alkyl ether sulphates e.g. sodium lauryl ether sulphate nEO, where n is from 1 to 20 alkylphenol ether sulphates e.g. octylphenol ether sulphate nEO where n is from 1 to 20, and sulphosuccinates e.g. sodium dioctylsulphosuccinate.

5 [0045] Also suitable are nonionic emulsifiers such as alkylphenol ethoxylates e.g. nonylphenol ethoxylate nEO, where n is from 1 to 50, alcohol ethoxylates e.g. lauryl alcohol nEO, where n is from 1 to 50, ester ethoxylates e.g. polyoxyethylene monostearate where the number of oxyethylene units is from 1 to 30.

10 [0046] Silicones which are particularly preferred for purposes of this invention are dimethiconols, these may be linear or branched. Average number molecular weight may range from about 1,000 to about 1 million, preferably from about 20,000 to about 300,000, optimally from about 40,000 to about 100,000. Microemulsions may be loaded with the silicone at levels ranging from about 1 to about 95%, preferably from about 10 to about 60%, optimally from about 20 to about 40% by weight. Pre-formed microemulsions are available from suppliers such as Dow Corning, General Electric, Union Carbide, Wacker Chemie, Shin Etsu, and Toray Silicone Company. Particularly preferred is a linear dimethiconol microemulsion at 25% silicone with a maximum particle size of 40 nm, pH 6.5-8 and surfactant combination of dodecylbenzene sulphonic acid triethanolamine/Lanreth-24 available from Dow Corning under the trademark DC 2-1870.

15 [0047] Compositions of this invention when in the form of a fluid will usually be provided with a variety of cosmetically acceptable carrier vehicles. Normally the carrier vehicle will be water. Amounts of the carrier vehicle may range from about 0.5 to about 99%, preferably from about 1 to about 80%, more preferably from about 50 to about 70%, optimally from about 65 to 75% by weight of the impregnating composition.

20 [0048] Preservatives can desirably be incorporated into the cosmetic compositions of this invention to protect against the growth of potentially harmful microorganisms. Suitable traditional preservatives for compositions of this invention are alkyl esters of para-hydroxybenzoic acid. Other preservatives which have more recently come into use include hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the preservative challenge test and to provide product stability. Particularly preferred preservatives are phenoxyethanol, methyl paraben, propyl paraben, imidazolidinyl urea, sodium dehydroacetate and benzyl alcohol. The preservatives should be selected having regard for the use of the composition and possible incompatibilities between the preservatives and other ingredients in the composition. Preservatives are preferably employed in amounts ranging from 0.01 % to 2 % by weight of the composition.

25 [0049] Compositions of the present invention may further include herbal extracts. Illustrative extracts include Roman Chamomile, Green Tea, Scullcap, Nettle Root, Swertia Japonica, Fennel and Aloe Vera extracts. Amount of each of the extracts may range from about 0.001 to about 1 %, preferably from about 0.01 to about 0.5% , optimally from about 0.05 to about 0.2% by weight of a composition.

30 [0050] Minor adjunct ingredients may also be present in the compositions. Among these may be vitamins such as Vitamin E Acetate, Vitamin C, Vitamin A Palmitate, Panthenol and any of the Vitamin B complexes. Anti-irritant agents may also be present including those of steviosides, alpha-bisabolol and glycyrrhizinate salts, each vitamin or anti-irritant agent being present in amounts ranging from about 0.001 to about 1.0%, preferably from about 0.01 to about 0.3 % by weight of the composition.

35 [0051] Emulsifiers may also be incorporated into compositions of this invention. These emulsifiers may be anionic, nonionic, cationic, amphoteric and combinations thereof. Useful nonionic type emulsifiers include the C 10-C20 fatty alcohol or acid hydrophobes condensed with from 2 to 100 moles of ethylene oxide or propylene oxide per mole of hydrophobe; C2-C10 alkyl phenols condensed with from 2 to 20 moles of alkylene oxide; mono- and di-fatty acid esters of ethylene glycol; fatty acid monoglyceride; sorbitan, mono- and di-C8-C20 fatty acids; block copolymers (ethylene oxide/propylene oxide); and polyoxyethylene sorbitan as well as combinations thereof. Alkyl polyglycosides and saccharide fatty amides (e.g. methyl gluconamides) are also suitable nonionic emulsifiers. Particularly preferred as the emulsifier is a hydrogenated castor wax alkoxyated with 40 moles ethylene oxide, available commercially as Cremophore RH-400®.

40 [0052] Mild emulsifiers of the anionic and amphoteric type may also be employed. Particularly preferred anionic examples include lauroamphoacetate salts and sarcosinate salts. Preferred amphoteric include cocamidopropylbetaine and dimethylbetaine.

45 [0053] Amounts of the emulsifiers may range from about 0.05 to about 20 % , preferably from about 0.1 to about 5%, optimally from about 0.5 to about 0.8% by weight.

50 [0054] Low pH systems containing fatty acid (C10-C22 alkyl) groups such as lauroamphoacetates have been found to emit foul odours. Although not wishing to be bound by theory, it is believed that hydrolysis of surfactants with fatty acid groups results in cleavage of these groups. Malodorous fatty acids are thereby formed. The applicants have surprisingly found that silicone microemulsions such as dimethiconol microemulsions effectively inhibit odour formation.

55 [0055] These impregnating compositions of the present invention may involve a range of pH although it is preferred to have a relatively low pH, for instance, a pH from about 2 to about 6.5, preferably from about 2.5 to about 4.5.

[0056] Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this

description indicating amounts of material ought to be understood as modified by the word "about".

[0057] The following examples will more fully illustrate the embodiments of this invention. All parts, percentages and proportions referred to herein and in the appended claims are by weight unless otherwise illustrated.

5 EXAMPLES 1-8

[0058] Table I provides a listing of formulations which are suitable for impregnation into a cellulosic substrate forming a towelette. The pH of the resulting composition solutions range from about 2.8 to about 4.0.

10 TABLE I

INGREDIENT	EXAMPLE (WEIGHT %)							
	1	2	3	4*	5	6	7	8
Water	To 100	To 100	To 100	To 100	To 100	To 100	To 100	To 100
Glycolic Acid (70% Active)	11.40	-	-	-	10.90	10.90	1.10	-
Potassium Lactate	-	8.00	-	-	-	0.05	-	4.00
Alpha-Hydroxy-Octanoic Acid	-	-	0.50	-	0.50	-	-	-
Glucarolactone*	-	-	-	2.00	-	-	-	-
Ammonia Solution (30% Active)	3.50	-	-	-	3.00	3.00	0.35	-
Glycerin	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Dimethiconol Microemulsion (25% Active)	0.75	0.75	0.75	0.75	4.00	3.00	2.00	0.10
Disodium Caprylamphodiacetate	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Witch Hazel Extract	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Glydant Plus	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20
PEG-40 Hydrogenated Castor Oil	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Fragrance	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15
Hexylene Glycol	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00
Vitamin E Acetate	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
* comparative example								

40 EXAMPLE 9

[0059] A study was conducted to evaluate the effects of silicones and other substances for stability and reducing stickiness in low pH hydroxycarboxylic acid depositing towelette products. Towelettes were impregnated with a formulation substantially similar to Example 1 except that the dimethiconol microemulsion was replaced with materials reported under Table II at the specified usage level. An amount of 4 grams formulation was impregnated into each towelette (1.8 gram weight; 6 inch by 8 inch size). Stability refers to the phase compatibility of the fluid which impregnated the towelettes. Ratings were 'good' for homogeneous fluids, 'fair, for those with only a small amount of top creaming, and 'poor, for significant phase separation. Stickiness was evaluated by a trained evaluator and measured by sensory finger feel.

TABLE II

Effects of Various Materials on Stability and Stickiness of Low pH Fluids Delivered by Towelettes				
Material	CTFA Name	Usage Level	Stability	Effectiveness on stickiness removal (excellent, very good, good, fair, no effect)
Silicones				
Dow Corning 2501	Dimethicone copolyol (wax)	3%	Good	Fair
D/C 2-1788	Dimethicone emulsion	3%	Poor	Good
D/C 2-1310	Dimethicone emulsion	3%	Poor	Good
D/C FB-50	Dimethicone emulsion	3%	Poor	Good
D/C 1664	Dimethicone emulsion	3%	Poor	Good
D/C 2-1870	Dimethiconol microemulsion	3%	Good	Good
D/C 2-8739	Dimethicone microemulsion	3%	Good	Fair
Silwet L-7087	Organosilicone fluid	3%	Good	No effect
Silwax WS-L	Dimethicone copolyol laurate	3%	Good	No effect
Silube CP-I	Dimethicone copolyol phthalate	3%	Good	Fair
Silwax WS	Orfanommodified dimethicone copolyol	3%	Good	Fair
Pecosil PS-100 Q	Dimethicone copolyol phospahte	3%	Good	No effect
D/C antifoam 1510	Dimethicone emulsion	3%	Poor	Good
Others				
Hetester PHA	Propylene glycol isoceteth-3 acetate	3%	Poor	Fair
Lubragel Oil	Glyceryl polymethacralate, propylene glycol, PVM/MA	3%	Good	No effect
Glycerox HE	PEG-7 glyceryl cocoate	3 %	Good	No effect
HPS-1180	Polysulfonic acid solution	3 %	Good	No effect
		10%	Good	No effect
Lipopeg 4-L	PEG-8 laurate	3%	Good	No effect
Crodesta F-110	Sucrose stearate and surcose distearate	3%	Fair	No effect
SMEC concentrate	OMC emulsion	3%	Good	No effect
Glycerox 767	PEG 6 capric / caprylic glycerides	3%	Good	No effect
Aqua Keep 10 NF	Sodium polyacrylate (SAP)	1 %	Poor	No effect
Arconate HP	Propylene carbonate	3%	Good	No effect
MP diol glycol	2-Methyl 1,3 propanediol	3%	Good	No effect
	2-Methyl 1,3 pentanediol	3%	Good	No effect

(continued)

Others					
5	Emulgade CM	Cetearyl isonanoate, cetereth-20, cetearyl alcohol, glycerin, cetyl palmitate	3%	Good	No effect

10 [0060] Almost all the non-silicone materials had no beneficial effect on ameliorating stickiness of deposited hydroxy-carboxylic acids, Dimethicone emulsions did work well to remove stickiness. However, these emulsions imparted poor stability to the fluid formulations. Only dimethiconol microemulsions retained both good physical stability while also effectively combating stickiness.

15 Claims

1. A towelette product comprising:

20 (a) a water-insoluble substrate;

(b) a cosmetic composition impregnated into the substrate including:

25 i) an alpha-hydroxycarboxylic acid selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof;

ii) a silicone microemulsion;

iii) the composition in water having a pH no higher than about 6.5.

2. A towelette product according to claim 1 wherein the pH of the cosmetic composition ranges from about 2.0 to about 6.0.

30 3. A towelette product according to claim 2 wherein the pH of the cosmetic composition ranges from about 3.5 to about 4.5.

35 4. A towelette product according to any of the preceding claims wherein the silicone microemulsion is a dimethiconol microemulsion.

5. A towelette product according to Claim 4 wherein the silicone material in the microemulsion has a particle size ranging from about 0.01 to about 500 nm.

40 6. A towelette product according to claim 5 wherein the silicone material in the microemulsion has a particle size ranging from about 5 to about 50 nm.

7. A towelette product according to any of the preceding claims wherein the hydroxycarboxylic acid is present in an amount from about 0.1 to about 15% by weight of the composition which impregnates the substrate.

45 8. A towelette product according any of the preceding claims wherein the microemulsion is present in an amount from about 0.1 to about 20% by weight of the composition which impregnates the substrate.

50 9. A towelette product according to any preceding claim, wherein the acid is present as an acid salt, the salt being selected from ammonium, alkali, alkaline earth, C₂ - C₈ trialkanolammonium and mixtures thereof.

Patentansprüche

55 1. Kosmetiktuchprodukt, umfassend:

(a) ein in Wasser unlösliches Substrat;

(b) eine kosmetische Zusammensetzung, die in das Substrat imprägniert ist, welche einschließt:

- i) eine α -Hydroxycarbonsäure, ausgewählt aus der Gruppe, umfassend Glycol-, Milch-, Hydroxyoctansäuren und Gemische davon;
- ii) eine Silikonmikroemulsion;
- iii) wobei die Zusammensetzung in Wasser einen pH-Wert aufweist, der nicht höher als etwa 6,5 ist.

5

2. Kosmetiktuchprodukt nach Anspruch 1, wobei der pH-Wert der kosmetischen Zusammensetzung im Bereich von etwa 2,0 bis etwa 6,0 liegt.

10

3. Kosmetiktuchprodukt nach Anspruch 2, wobei der pH-Wert der kosmetischen Zusammensetzung im Bereich von etwa 3,5 bis etwa 4,5 liegt.

4. Kosmetiktuchprodukt nach einem der vorangehenden Ansprüche, wobei die Silikonmikroemulsion eine Dimethiconolmikroemulsion ist.

15

5. Kosmetiktuchprodukt nach Anspruch 4, wobei das Silikonmaterial in der Mikroemulsion eine Teilchengröße im Bereich von etwa 0,01 bis etwa 500 nm aufweist.

6. Kosmetiktuchprodukt nach Anspruch 5, wobei das Silikonmaterial in der Mikroemulsion eine Teilchengröße im Bereich von etwa 5 bis etwa 50 nm aufweist.

20

7. Kosmetiktuchprodukt nach einem der vorangehenden Ansprüche, wobei die Hydroxycarbonsäure in einer Menge von etwa 0,1 bis etwa 15 Gewichtsprozent der Zusammensetzung, die das Substrat imprägniert, vorliegt.

25

8. Kosmetiktuchprodukt nach einem der vorangehenden Ansprüche, wobei die Mikroemulsion in einer Menge von etwa 0,1 bis etwa 20 Gewichtsprozent der Zusammensetzung, die das Substrat imprägniert, vorliegt.

9. Kosmetiktuchprodukt nach einem vorangehenden Anspruch, wobei die Säure als ein saures Salz vorliegt, wobei das Salz ausgewählt ist aus Ammonium, Alkali, Erdalkali, C_2 - C_8 -Trialkanolammonium und Gemischen davon.

30

Revendications

1. Lingette comprenant :

35

(a) un substrat insoluble dans l'eau ;

(b) une composition cosmétique imprégnée dans le substrat, comprenant :

40

i) un acide α -hydroxycarboxylique choisi dans le groupe comprenant les acides glycolique, lactique, hydroxyoctanoïque et leurs mélanges ;

ii) une microémulsion de silicone ;

iii) la composition dans l'eau ayant un pH de pas plus d'environ 6,5.

2. Lingette selon la revendication 1, dans laquelle le pH de la composition cosmétique se situe dans la plage allant d'environ 2,0 à environ 6,0.

45

3. Lingette selon la revendication 2, dans laquelle le pH de la composition cosmétique se situe dans la plage allant d'environ 3,5 à environ 4,5.

4. Lingette selon l'une quelconque des revendications précédentes, dans laquelle la microémulsion de silicone est une microémulsion de diméthiconol.

50

5. Lingette selon la revendication 4, dans laquelle le matériau de silicone dans la microémulsion a une taille de particule située dans la plage allant d'environ 0,01 à environ 500 nm.

55

6. Lingette selon la revendication 5, dans laquelle le matériau de silicone dans la microémulsion a une taille de particule située dans la plage allant d'environ 5 à environ 50 nm.

7. Lingette selon l'une quelconque des revendications précédentes, dans laquelle l'acide hydroxycarboxylique est

présent en une quantité d'environ 0,1 à environ 15 % en poids de la composition qui imprègne le substrat.

- 5
8. Lingette selon l'une quelconque des revendications précédentes, dans laquelle la microémulsion est présente en une quantité d'environ 0,1 à environ 20 % en poids de la composition qui imprègne le substrat.
9. Lingette selon l'une quelconque des revendications précédentes, dans laquelle l'acide est présent sous la forme d'un sel d'acide, le sel étant choisi parmi l'ammonium, un alcalin, un alcalino-terreux, un trialcanolammonium en C₂ à C₈ et leurs mélanges.
- 10
- 15
- 20
- 25
- 30
- 35
- 40
- 45
- 50
- 55