

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2011380507 B2

(54) Title
Methods for treating hyperuricemia in patients with gout using halofenate or halofenic acid and a second urate-lowering agent

(51) International Patent Classification(s)
A61K 31/19 (2006.01)

(21) Application No: **2011380507** (22) Date of Filing: **2011.11.04**

(87) WIPO No: **WO13/066349**

(43) Publication Date: **2013.05.10**
(44) Accepted Journal Date: **2017.06.15**

(71) Applicant(s)
CYMABAY THERAPEUTICS, INC.;DIATEX, INC.

(72) Inventor(s)
Roberts, Brian K.;Saha, Gopal Chandra;Lavan, Brian Edward;McWherter, Charles A.

(74) Agent / Attorney
AJ PARK, GPO Box 2600, Sydney, NSW, 2001, AU

(56) Related Art
**"NTC01416402 on 2011_10_21: ClinicalTrials.gov Archive" 21 October 2011Retrieved from the internet:<URL: https://clinicaltrials.gov/archive/NCT01416402/2011_10_21>[retrieved on 2 September 2016]
SAHA, G. C. Arthritis and Rheumatism, vol. 63, no. 10, Suppl. S. October 2011, page S1014**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2013/066349 A1

(43) International Publication Date
10 May 2013 (10.05.2013)

(51) International Patent Classification:

A61K 31/19 (2006.01)

(21) International Application Number:

PCT/US2011/059394

(22) International Filing Date:

4 November 2011 (04.11.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant (for all designated States except US): META-BOLEX, INC. [US/US]; 3876 Bay Center Place, Hayward, CA 94545 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ROBERTS, Brian K. [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, CA 94545 (US). SAHA, Gopal Chandra [BD/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, CA 94545 (US). LAVAN, Brian Edward [GB/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, CA 94545 (US). McWHERTER, Charles A. [US/US]; c/o Metabolex, Inc., 3876 Bay Center Place, Hayward, CA 94545 (US).

(74) Agents: NGUYEN, Sam L. et al.; Hamilton, DeSanctis & Cha LLP, 3239 El Camino Real, Suite 220, Palo Alto, CA 94306 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2013/066349 A1

(54) Title: METHODS FOR TREATING HYPERURICEMIA IN PATIENTS WITH GOUT USING HALOFENATE OR HALOFENIC ACID AND A SECOND URATE-LOWERING AGENT

(57) Abstract: Disclosed herein are pharmaceutical compositions, methods and kits for lowering the serum uric acid level of a subject and for the treatment of a condition associated with elevated serum uric acid levels comprising administering a composition comprising a first urate-lowering agent and a second urate-lowering agent. In some aspects the first urate-lowering agent is (-)-halofenate, (-)-halofenic acid, or a pharmaceutically acceptable salts thereof. In some aspects the second urate-lowering agent is an inhibitor of uric acid production, a uricosuric agent, a uricase, or a pharmaceutically acceptable salt thereof.

METHODS FOR TREATING HYPERURICEMIA IN PATIENTS WITH GOUT USING HALOFENATE OR HALOFENIC ACID AND A SECOND URATE- LOWERING AGENT

BACKGROUND

[0001] Conditions associated with elevated serum uric acid levels (hyperuricemia) include disorders of urate crystal deposition such as gout arthropathy and tophi, urolithiasis (urinary tract stones), urate nephropathy, as well as the sequelae of these disorders. Hyperuricemia is associated with an increased risk of developing gout arthropathy, and the risk of gout increases with the degree and duration of the hyperuricemia. In addition to gout arthropathy, chronic hyperuricemia may lead to the deposition of uric acid crystals in the urinary tract, renal parenchyma, and soft tissues, resulting in urolithiasis, urate nephropathy with chronic kidney disease, and soft tissue tophi, respectively. Because of limitations and disadvantages of current uric acid lowering agents, more effective methods, compositions and therapies to lower uric acid are needed. It is an object of the present invention to go somewhat towards meeting this need and/or to provide the public with a useful choice.

SUMMARY

[0002] In a first embodiment, the invention provides an oral dosage composition comprising:

- (a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and
- (b) febuxostat.


[0002a] In a second embodiment, the invention provides a method of treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout, comprising administering an effective amount of an oral dosage composition of the invention.

[0002b] In a third embodiment, the invention provides the use of:

- (a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and
- (b) febuxostat,

in the manufacture of an oral dosage composition for treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout.

[0002c] Also described are methods of lowering the serum uric acid level of a subject with hyperuricemia, the method comprising administering to the subject a first urate-lowering agent and a second urate-lowering agent, wherein the first urate-lowering agent is a compound of Formula (I)



wherein R is selected from the group consisting of hydroxy, lower aralkoxy, di-lower alkylamino-lower alkoxy, lower alkanamido-lower alkoxy, benzamido-lower alkoxy, ureido-lower alkoxy, N'-lower alkyl-ureido-lower alkoxy, carbamoyl-lower alkoxy, halophenoxy-substituted lower alkoxy, carbamoyl-substituted phenoxy, carbonyl-lower alkylamino, N,N-di-lower alkylamino-lower alkylamino, halo-substituted lower alkylamino, hydroxyl- substituted lower alkylamino, lower alkanolyloxy-substituted lower alkylamino, ureido, and lower alkoxy carbonylamino; and each X is independently a halogen; or a pharmaceutically acceptable salt thereof.

[0003] Also disclosed are methods of treating a subject having a condition associated with hyperuricemia, the method comprising administering to the subject a first urate-lowering agent and a second urate-lowering agent, wherein the first urate-lowering agent is a compound of Formula (I). Also disclosed are methods of treating hyperuricemia in a subject with gout comprising administering to the subject a composition comprising a first urate-lowering agent and a second urate-lowering agent, wherein the first urate-lowering agent is a compound of Formula (I).

[0004] Also disclosed are compositions and kits comprising a first urate-lowering agent and a second urate-lowering agent, wherein the first urate-lowering agent is a compound of Formula (I).

[0005] In some aspects, the compound of Formula (I) is (-)-halofenate, (-)-halofenic acid, or a pharmaceutically acceptable salt thereof. In some aspects, the second urate-lowering agent is a

xanthine oxidase inhibitor, an inhibitor of uric acid production, a uricosuric agent or a uricase. In some aspects, the second urate-lowering agent is allopurinol or febuxostat. Other aspects are provided below.

[0006] Currently available uric acid lowering agents and other therapeutic agents in development have limitations in their ability to lower serum uric acid to a desirable level, and their use may be limited by various adverse side effects or toxicities. For example, certain agents including allopurinol and febuxostat, when used to treat hyperuricemia at commonly prescribed doses, often fail to reach the common therapeutic target of serum uric acid levels of 6 mg/dL or less. Advantages of the compositions, methods, and kits disclosed herein over currently available uric acid lowering agents at commonly prescribed doses and treatment methods using such agents may include improved therapeutic benefits; a synergistic effect on lowering uric acid (i.e., an additive or over-additive effect as compared to the effects of single-agent therapies); beneficial effects on other conditions associated with hyperuricemia and urate crystal deposition; and provoking fewer or less intense side effects. In some aspects, the synergistic effect allows dose reduction or dosing interval extension relative to a currently available uric acid lowering agent taken individually at prescribed doses.

DETAILED DESCRIPTION

[0007] As utilized in accordance with the present disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

[0008] “About” when qualifying a number, refers to a range of plus or minus ten percent of that value or number, unless indicated otherwise. Without limiting the application of the doctrine of equivalents as to the scope of the claims, each number should be construed in light of such factors as the number of reported significant digits and the manner or method (e.g. instrumentation, sample preparation, etc.) used to obtain that number.

[0009] “Administering” or “administration” refers to the act of giving a drug, prodrug, or therapeutic agent to a subject. Exemplary routes of administration are discussed below.

[0010] “Acute gout” refers to gout present in a subject with at least one gouty symptom (e.g., podagra or other gouty arthritis, gout flare, gouty attack).

[0011] “Arhalofenate” refers to (−)-halofenate, i.e. (−)-(R)-(4-chloro-phenyl)-(3-trifluoromethyl-phenoxy)-acetic acid 2-acetylamino-ethyl ester.

[0012] “Chronic gout” refers to gout present in a subject having recurrent or prolonged gout flares, tophus formation, chronic inflammatory arthritis, or joint deterioration associated with gout, and includes the periods following recovery from acute gout and between acute gout attacks (i.e. intercritical gout).

[0013] “Composition” or, interchangeably, “formulation” refers to a preparation that contains a mixture of various excipients and key ingredients that provide a relatively stable, desirable, and useful form of a compound or drug.

[0014] The prefixes “d” and “l” or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with (+) or d- meaning that the compound is “dextrorotatory” and with (-) or l- meaning that the compound is “levorotatory”. For a given chemical structure, these isomers or “optical isomers” are identical except that they are mirror images of one another. In describing an optically active compound, the prefixes R and S are used to denote the absolute configuration of the molecule about its chiral center(s). There is no correlation between the nomenclature for the absolute stereochemistry and for the rotation of an enantiomer (i.e., the R- isomer can also be the l- isomer). A specific optical isomer can also be referred to as an “enantiomer,” and a mixture of such isomers is often called an “enantiomeric” or “racemic” mixture. *See, e.g., A. Streitwieser, & C.H. Heathcock, INTRODUCTION TO ORGANIC CHEMISTRY, 2nd Edition, Chapter 7 (MacMillan Publishing Co., U.S.A. 1981).*

The optical rotation $[\alpha]_D$ of (-)-halofenate was measured in methyl alcohol.

[0015] “Elevated serum uric acid level” refers to a serum uric acid level greater than normal and, in patients with gout, generally refers to a serum uric acid level greater than or equal to about 6 mg/dL. In some instances, elevated serum uric acid levels are above the mean level in a given population, such as those of a particular gender or age.

[0016] “Effective amount” refers to an amount required (i) at least partly to attain the desired response in a subject; (ii) to delay or to prevent the onset of a particular condition being treated in a subject; or (iii) or to inhibit or to prevent the progression of a particular condition being treated in a subject. The effective amount for a particular subject varies depending upon the health and physical condition of the subject to be treated, the taxonomic group of individual to be treated, the degree of protection desired, the formulation of the composition, the assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

[0017] “First urate-lowering agent” refers to a compound of any of Formulae (I), (II), (III), or (IV) or a therapeutically acceptable salt or prodrug thereof.

[0018] “Gout” refers to a group of disorders or symptoms most often associated with the build up of uric acid due to an overproduction of uric acid or a reduced ability of the kidney to excrete uric acid. Gout is often characterized by the deposition of urate crystals (uric acid or salts thereof, e.g. monosodium urate) in the joints (gouty arthropathy) or soft tissue (tophi). “Gout” as used herein includes acute gout, chronic gout, moderate gout, refractory gout and severe gout.

[0019] “Gout-associated inflammation” refers to local or systemic inflammation due to immune responses to the deposition of urate crystals.

[0020] “Halofenate” refers to the compound of Formula (III), i.e. (4-chlorophenyl)-(3-trifluoromethylphenoxy)-acetic acid 2-acetylaminooethyl ester (also referred to as the 2-acetamidoethyl ester of 4-chlorophenyl-(3-trifluoromethylphenoxy)-acetic acid. The term halofenate and the corresponding chemical names include both the (+) and (-) enantiomer of compounds of Formula (III) as well as mixtures thereof, unless otherwise specified.

[0021] “Halofenic acid” and “CPTA” refer to the compound of Formula (IV), i.e. 4-chlorophenyl-(3-trifluoromethylphenoxy)-acetic acid [also referred to as 2-(4-chlorophenyl)-2-(3-(trifluoromethyl)phenoxy)acetic acid] as well as its pharmaceutically acceptable salts. The term halofenic acid and the corresponding chemical names include both the (+) and (-) enantiomer of compounds of Formula (II) as well as mixtures thereof, unless otherwise specified.

[0022] “Hyperuricemia” refers to an elevated serum uric acid level (see above).

[0023] “Lower,” when used to describe chemical substituents of compounds of Formulae (I) and (II) such as lower aralkoxy, di-lower alkylamino-lower alkoxy, lower alkanamido, lower alkoxy, benzamido-lower alkoxy, ureido-lower alkoxy, N'-lower alkyl-ureido-lower alkoxy, carbamoyl-lower alkoxy, halophenoxy substituted lower alkoxy, carbonyl-lower alkylamino, N,N-di-lower alkylamino-lower alkylamino, halo substituted lower alkylamino, hydroxy substituted lower alkylamino, lower alkanolylxy substituted lower alkylamino, lower alkoxy carbonyl amino, phenyl-lower alkyl, lower alkanamido-lower alkyl, and benzamido-lower alkyl refers to groups having from one to six carbon atoms. For example, “lower alkoxy” means C₁₋₆alkoxy.

[0024] “Moderate gout” refers to gout present in a subject having at least two gout flares in the past 12 months.

[0025] “Pharmaceutically acceptable” refers to that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and neither biologically nor otherwise undesirable, and includes that which is acceptable for veterinary or human pharmaceutical use.

[0026] “Pharmaceutically acceptable salt” includes pharmaceutically acceptable acid addition salts and pharmaceutically acceptable base addition salts and includes both solvated and unsolvated forms. Representative non-limiting lists of pharmaceutically acceptable salts can be found in S.M. Berge *et al.*, *J. Pharma Sci.*, 66(1), 1-19 (1977), and *Remington: The Science and Practice of Pharmacy*, R. Hendrickson, ed., 21st edition, Lippincott, Williams & Wilkins, Philadelphia, PA, (2005), at p. 732, Table 38-5, both of which are hereby incorporated by reference herein.

[0027] “Pharmaceutically acceptable acid addition salt” refers to salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

[0028] “Pharmaceutically acceptable base addition salt” refers to salts prepared from the addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like.

[0029] “Refractory gout” refers to gout in patients who are unresponsive or poorly responsive, or have experienced or are at an increased risk of experiencing an adverse event, after being administered either (1) one or more second urate-lowering agents but not a first urate-lowering

agent or (2) a first-urate lowering agent but not a second urate-lowering agent. The terms “unresponsive” and “poorly responsive” in this context include (1) no or insignificant lowering of serum uric acid, (2) failure to reach a target serum uric acid level (e.g. as determined by a physician or other medical practitioner), and (3) the persistence of one or more gouty conditions or symptoms such as gout flares, gouty tophus, gouty arthritis, or other associated conditions regardless of any lowering of serum uric acid levels.

[0030] “Second urate-lowering agent” refers to a therapeutic agent that lowers serum uric acid levels that is not a first urate-lowering agent. Second urate-lowering agents include currently available agents (i.e. an agent approved by the FDA or other appropriate regulatory authority as of the filing date of this application) that lower serum uric acid, as well as compounds currently in development or under regulatory review. Examples of second urate-lowering agents are provided below.

[0031] “Subject” and “patient” refer to animals such as mammals, including humans, other primates, domesticated animals (e.g. dogs, cats), farm animals (e.g. horses, cattle, goats, sheep, pigs), rats and mice.

[0032] “Severe gout” refers to gout present in a subject having tophaceous deposits in the joints, skin, or kidneys resulting in chronic arthritis, joint destruction, subcutaneous tophi, or kidney dysfunction, and, in some cases, with subsequent deformity and/or disability.

[0033] “Substantially free from” when used in reference to (−)-halofenate or (−)-halofenic acid (or a salt thereof) being substantially free from the corresponding (+) enantiomer (i.e. (+)-halofenate, (+)-halofenic acid, or a salt thereof) refers to a composition containing a high proportion of a compound’s (−) enantiomer in relation to the (+) enantiomer. In one embodiment, the term means that by weight, the compound included in the composition is at least 85% (−) enantiomer and at most 15% (+) enantiomer. In one embodiment, the term means that by weight, the compound included in the composition is at least 90% (−) enantiomer and at most 10% (+) enantiomer. In other embodiments, the term means that by weight, the compound included in the composition is at least 91% (−) enantiomer and at most 9% (+) enantiomer, at least 92% (−) enantiomer and at most 8% (+) enantiomer, at least 93% (−) enantiomer and at most 7% (+) enantiomer, at least 94% (−) enantiomer and at most 6% (+) enantiomer, at least 95% (−) enantiomer and at most 5% (+) enantiomer, at least 96% (−) enantiomer and at most 4% (+) enantiomer, at least 97% (−) enantiomer and at most 3% (+) enantiomer, at least 98% (−)

enantiomer and at most 2% (+) enantiomer, or at least 99% (-) enantiomer or greater than 99% (-) enantiomer. Other percentages of the (-) and (+) enantiomers may also be provided. These percentages are based upon the amount of the enantiomer relative to the total amount of both enantiomers of the compound in the composition.

[0034] “Therapeutically effective dose,” “therapeutically effective amount,” or, interchangeably, “pharmacologically acceptable dose” and “pharmacologically acceptable amount” mean that a sufficient amount of a therapeutic agent, therapeutic agents, or metabolites thereof will be present in order to achieve a desired result, e.g., lowering uric acid levels to a target goal or treating gout in its various forms or treating conditions associated with hyperuricemia.

[0035] “Treatment” and “treating” of a disease, disorder, condition or symptom refer to (1) preventing or reducing the risk of developing the disease, disorder or condition, i.e., causing the clinical symptoms of the disease, disorder or condition not to develop in a subject who may be exposed to or predisposed to the disease, disorder or condition but who does not yet experience or display symptoms of the disease, disorder or condition (i.e. prophylaxis); (2) inhibiting the disease, disorder or condition, i.e., arresting or reducing the development of the disease, disorder or condition or its clinical symptoms; and (3) relieving the disease, disorder or condition, i.e., causing regression, reversal, or amelioration of the disease, disorder or condition or reducing the number, frequency, duration or severity of its clinical symptoms. The term “management” may be used synonymously.

[0036] “Urate” refers to uric acid (7,9-dihydro-1H-purine-2,6,8(3H)-trione) and ions and salts thereof.


[0036a] The term “comprising” as used in this specification and claims means “consisting at least in part of”. When interpreting statements in this specification, and claims which include the term “comprising”, it is to be understood that other features that are additional to the features prefaced by this term in each statement or claim may also be present. Related terms such as “comprise” and “comprised” are to be interpreted in similar manner.

[0037] This application describes compositions, kits and methods for the treatment of hyperuricemia, that is, for lowering serum uric acid levels. One aspect of the current disclosure provides an oral dosage composition comprising:

(a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and


(b) febuxostat.

[0037a] Also described is a composition comprising a first urate-lowering agent and a second urate-lowering agent, wherein said first urate-lowering agent is a compound of Formula (I)




wherein *R* is selected from the group consisting of a hydroxy, lower aralkoxy, di-lower alkylamino-lower alkoxy, lower alkanamido-lower alkoxy, benzamido-lower alkoxy, ureido-lower alkoxy, N'-lower alkyl-ureido-lower alkoxy, carbamoyl-lower alkoxy, halophenoxy substituted lower alkoxy, carbamoyl substituted phenoxy, carbonyl-lower alkylamino, N,N-di-lower alkylamino-lower alkylamino, halo substituted lower alkylamino, hydroxy substituted lower alkylamino, lower alkanolyl substituted lower alkylamino, ureido, and lower alkoxy carbonyl amine; and each *X* is independently a halogen; or a pharmaceutically acceptable salt thereof.

[0038] In certain aspects, the first urate-lowering agent is a compound of Formula (II)




wherein R² is selected from the group consisting of phenyl-lower alkyl, lower alkanamido-lower alkyl, and benzamido-lower alkyl; and each *X* is independently a halogen, or a pharmaceutically acceptable salt thereof.

[0039] In other aspects, the first urate-lowering agent is a compound of Formula (III), also referred to as halofenate



or a pharmaceutically acceptable salt thereof.

[0040] In other aspects, the first urate-lowering agent is a compound of Formula (IV), also referred to as halofenic acid



(IV)

or a pharmaceutically acceptable salt thereof.

[0041] It should also be noted that any carbon atom with unsatisfied valences in the formulae and examples herein is assumed to have the hydrogen atom to satisfy the valences.

[0042] In certain embodiments the compound is a compound that generates the compound of Formula (IV) or a pharmaceutically acceptable salt thereof via a chemical reaction after being administered, as discussed in more detail below.

[0043] Another aspect provides a method of treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout, comprising administering an effective amount of an oral dosage composition of the invention.

[0043a] Also described are methods of treating a condition associated with an elevated serum uric acid level comprising administering to a subject in need thereof a pharmaceutical composition comprising a first urate-lowering agent, wherein said first urate-lowering agent is a compound of Formulae (I), (II), (III) or (IV) or a pharmaceutically acceptable salt thereof; and a second urate-lowering agent. Also described is a method of lowering the serum uric acid level in a subject comprising administering to a subject in need thereof a pharmaceutical composition comprising a first urate-lowering agent, wherein said first urate-lowering agent is a compound of Formulae (I), (II), (III) or (IV) or a pharmaceutically acceptable salt thereof; and a second urate-lowering agent.

[0044] In certain embodiments, the first urate-lowering agent is (-)-halofenate (i.e. (-)-(R)-(4-chlorophenyl)-(3-trifluoromethylphenoxy)-acetic acid 2-acetylaminooethyl ester, also referred to as arhalofenate). In other embodiments, the first urate-lowering agent is (-)-halofenic acid (i.e. (-)-4-chlorophenyl-(3-trifluoromethylphenoxy) acetic acid) or a pharmaceutically acceptable salt

thereof. In certain embodiments, the (-)-halofenate, (-)-halofenic acid, or pharmaceutically acceptable salt thereof is substantially free from the corresponding (+) enantiomer.

[0045] The enantiomers (stereoisomers) of compounds of Formulae (I), (II), (III) or (IV) and pharmaceutically acceptable salt thereof can be prepared by using reactants or reagents or catalysts in their single enantiomeric form in the process wherever possible or by resolving the mixture of stereoisomers by conventional methods including use of microbial resolution, resolving the diastereomeric salts formed with chiral acids or chiral bases and chromatography using chiral supports. *See, also* U.S. Patent No. 7,199,259 (Daugs), U.S. Patent Nos. 6,646,004; 6,624,194; 6,613,802; and 6,262,118 (each to Luskey et al.), U.S. Patent No. 7,714,131 (Zhu et al.), U.S. Patent No. 7,432,394 (Cheng et al.) and U.S. Publication No. 2010/0093854 (Broggini et al.) each of which are incorporated herein by reference in their entireties.

[0046] The chemical synthesis of racemic mixtures of (3-trihalomethylphenoxy) (4-halophenyl) acetic acid derivatives can also be performed by the methods described in U.S. Patent No. 3,517,050, the teachings of which are incorporated herein by reference. The individual enantiomers can be obtained by resolution of the racemic mixture of enantiomers using conventional means known to and used by those of skill in the art. *See, e.g.*, Jaques, J., et al., in *Enantiomers, Racemates, and Resolutions*, John Wiley and Sons, New York (1981). Other standard methods of resolution known to those skilled in the art, including but not limited to, simple crystallization and chromatographic resolution, can also be used (*see, e.g.*, *Stereochemistry of Carbon Compounds* (1962) E. L. Eliel, McGraw Hill; J. Lochmuller, *Chromatography*, 113, 283-302 (1975)). Additionally, halofenate, halofenic acid, or a pharmaceutically acceptable salt thereof, *i.e.*, the optically pure isomers, can be prepared from the racemic mixture by enzymatic biocatalytic resolution. Enzymatic biocatalytic resolution has been generally described previously (*see, e.g.*, U.S. Patent Nos. 5,057,427 and 5,077,217, the disclosures of which are incorporated herein by reference). Other generic methods of obtaining enantiomers include stereospecific synthesis (*see, e.g.*, A.J. Li et al., *Pharm. Sci.* 86, 1073-77 (1997)).

[0046a] Another aspect provides the use of:

- (a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and
- (b) febuxostat,

in the manufacture of an oral dosage composition for treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout.

[0047] Also described is a composition comprising a pharmaceutically acceptable salt of halofenate or halofenic acid. The neutral forms of the therapeutic agents may be regenerated by contacting the salt with a base or acid and isolating the parent therapeutic agent in the conventional manner. The parent form of the therapeutic agent differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form.

[0048] The second urate-lowering agent may be any other agent (i.e. not a first urate-lowering agent, as defined herein) that lowers serum uric acid levels. These second urate-lowering agents include inhibitors of uric acid production (e.g. xanthine oxidase inhibitors and purine nucleoside phosphorylase inhibitors), uricosuric agents and uricases.

[0049] For example, in some embodiments, the second urate-lowering agent is a xanthine oxidase inhibitor. Xanthine oxidase inhibitors lower the amount of urate in blood by decreasing the synthesis of uric acid. Xanthine oxidase is involved in purine metabolism and inhibiting the enzyme reduces uric acid levels. Xanthine oxidase inhibitors include, but are not limited to: allopurinol, febuxostat, oxypurinol, tisopurine, an inositol and propolis. In some embodiments, the xanthine oxidase inhibitor is allopurinol, febuxostat, oxypurinol, tisopurine, inositol, phytic acid, myo-inositol, kaempferol, myricetin⁴ and quercetin. Allopurinol (1,5-dihydro-4H-pyrazolo [3,4-d]pyrimidin-4-one), a xanthine oxidase inhibitor, is the current first line standard of care for lowering urate levels. Another xanthine oxidase inhibitor, febuxostat (2-(3-cyano-4-isobutoxyphenyl)-4-methyl-1,3-thiazole-5-carboxylic acid), was approved for treatment of gout in February 2009. In one embodiment, halofenate, halofenic acid or a pharmaceutically acceptable salt thereof, is administered before, concurrently or subsequent to administration of allopurinol. In one embodiment, halofenate, halofenic acid or a pharmaceutically acceptable salt thereof is administered before, concurrently or subsequent to administration of febuxostat.

[0050] In other embodiments, the second urate-lowering agent is a purine nucleoside phosphorylase (PNP) inhibitor. Purine nucleoside phosphorylase inhibitors represent a relatively new approach to lowering serum uric acid levels in patient with hyperuricemia, gout, and related conditions. In some embodiments, the PNP inhibitor is forodesine (BCX-1777) (BioCryst

Pharmaceuticals, Inc.). In other embodiments, the PNP inhibitor is BCX-4208 (7-(((3R,4R)-3-hydroxy-4-(hydroxymethyl)pyrrolidin-1-yl)methyl)-3H-pyrrolo[3,2-d]pyrimidin-4(5H)-one) (BioCryst Pharmaceuticals, Inc.). BCX4208 monotherapy administered at 40, 80, 120, 160 and 240 mg/day has been shown to rapidly and significantly reduced serum uric acid in gout patients.

[0051] In some embodiments, the second urate-lowering agent is a uricosuric agent. Uricosuric agents enhance renal excretion of uric acid and generally act by lowering the absorption of uric acid from the kidney proximal tubule back to the blood, e.g., by inhibiting urate transporters, e.g., SLC22A12. Uricosuric agents include, but are not limited to, probenecid, 2-((5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-yl)thio)acetic acid (RDEA594, lesinurad), potassium 4-(2-((5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-yl)thio)acetamido)-3-chlorobenzoate (RDEA806), RDEA684, benzboronarone, sulfapyrazone, amlodipine, atorvastatin, fenofibrate, guaifenesin, losartan, adrenocorticotropic hormone, and cortisone. Probenecid is the most commonly used uricosuric agent in the U.S. and may be given in combination with allopurinol to some gout patients. Benzboronarone and sulfapyrazone are also used as first line uricosuric agents. Guaifenesin, losartan, atorvastatin, amlodipine, adrenocorticotropic hormone (ACTH or corticotropin), fenofibrate and cortisone also have uricosuric effects. In one embodiment, a first urate-lowering agent (e.g. (-)-halofenate, (-)-halofenic acid or a pharmaceutically acceptable salt thereof) is administered before, concurrently or subsequent to administration of a uricosuric agent. In one embodiment, a first urate-lowering agent (e.g. (-)-halofenate, (-)-halofenic acid or a pharmaceutically acceptable salt thereof) is administered before, concurrently or subsequent to administration of probenecid, benzboronarone or sulfapyrazone.

[0052] In some embodiments, the second urate-lowering agent is a uricase enzyme, or a fragment or pegylated derivative thereof. Uricase or urate oxidase enzymes are found in many mammals but not in humans. They can lower uric acid levels by converting uric acid into allantoin, a benign end metabolite which is easily excreted in the urine. Uricase enzymes include, but are not limited to, rasburicase or a pegylated uricase enzyme (PEG-uricase). In some embodiments, the pegylated uricase enzyme is Krystexxa® (PURICASE®; pegloticase) (Savient Pharmaceuticals, Inc.) which is approved in the U.S. for the treatment of chronic gout in adult patients refractory to conventional therapy.

[0053] Also described are methods of treating one or more conditions associated with an elevated serum uric acid level, i.e. hyperuricemia, the methods comprising administering to a subject in need thereof a pharmaceutical composition comprising a first urate-lowering agent, wherein said first urate-lowering agent is a compound of Formulae (I), (II), (III) or (IV) or a pharmaceutically acceptable salt thereof; and a second urate-lowering agent. Conditions associated with hyperuricemia include, but are not limited to gout; acute gout; chronic gout; moderate gout; refractory gout; severe gout; deposition of uric acid crystals in the urinary tract, renal parenchyma, soft tissues, joints, cartilage or bones; urolithiasis; urate nephropathy; tophi; podagra; acute inflammatory gouty arthritis; joint destruction; urinary tract infections; renal impairment; chronic kidney disease; kidney stones; local inflammation; systemic inflammation; immune-related disorders; cardiovascular disease including peripheral vascular disease, coronary artery disease and cerebrovascular disease; insulin resistance; diabetes; fatty liver disease; dementia including vascular dementia; dyslipidemia; preeclampsia; hypertension; obesity; muscle spasm; localized swelling; pain including joint pain, muscle fatigue; and stress feelings.

[0054] A variety of factors increase the risk that a patient will have gout or will experience one or more of its symptoms. In addition to hyperuricemia, these factors include obesity, diabetes, chronic kidney failure, hypertension, use of diuretic drugs and certain other drugs (e.g. salicylates, pyrazinamide, ethambutol, nicotinic acid, cyclosporin, 2-ethylamino-1,3,4-thiadiazole, fructose and cytotoxic agents), overeating or fasting, a high purine diet, a high fructose diet, exposure to lead, consumption of red meat and protein, alcohol intake, and injury or recent surgery. Acute gout can be precipitated by perioperative ketosis in surgical patients, reduced body temperature, e.g., while sleeping, and by dehydration, e.g., by use of diuretic drugs. Genetic risk factors for gout and hyperuricemia have also been identified.

[0055] In various embodiments, the methods described herein may be used to treat any of the aforementioned conditions or disorders. That is, in one embodiment, the condition associated with an elevated serum uric acid level is gout. In some embodiments, the subject has acute gout. In some embodiments, the subject has chronic gout. In some embodiments the subject has moderate gout. In some embodiments the subject has refractory gout. In some embodiments the subject has severe gout. For example, one method provides for the management of hyperuricemia in a subject with gout. Certain methods provide for the treatment or management of hyperuricemia in a subject with gout comprising administering a pharmaceutical composition

comprising a first urate-lowering agent and a second urate-lowering agent. In some embodiments the first urate-lowering agent is (-)-halofenate, (-)-halofenic acid or a pharmaceutically acceptable salt thereof. In certain embodiments, the treatment can be for about four weeks or longer, for about one month or longer, for about 12 weeks or longer, for about three months or longer, for about six months or longer, for about one year or longer, for about two years or longer, for about five years or longer, for about 10 years or longer. In certain embodiments the treatment can be indefinite, e.g. for the remainder of the lifetime of the subject. In certain embodiments the second urate-lowering agent is selected from the group consisting of a uric acid synthesis inhibitor, a uricase, and a uricosuric agent, and pharmaceutically acceptable salts thereof. In certain embodiments the second agent may be allopurinol or febuxostat.

[0056] In various embodiments the methods comprise treating gout. In some embodiments, the methods comprise treating gout by preventing gout flares. In another embodiment the method comprises reducing the number, frequency, duration or severity of one or more gout flares. In another embodiment the method comprises preventing, reducing or reversing uric acid crystal formation. In some embodiments of the methods for treating uric acid crystal formation, the uric acid crystal formation is in one or more of the joints, under skin, and kidney. In some embodiments, the formations include tophaceous deposits. In some embodiments, the subject has uric acid crystal formation determined by aspiration of tophi or by aspiration of synovial fluid of an inflamed joint. In another embodiment the method comprises reducing uric acid burden. In another embodiment the method comprises reducing the size or number of tophi. The size or number of tophi may be assessed by known methods, for example, use of CT scans.

[0057] Also described are methods of lowering the serum uric acid level, treating a subject having a condition associated with an elevated serum uric acid level, and treating hyperuricemia in a subject with gout, in subjects with refractory gout. In certain embodiments, the subject is refractory to allopurinol, 2-((5-bromo-4-(4-cyclopropyl)naphthalen-1-yl)-4H-1,2,4-triazol-3-yl)thio)acetic acid (RDEA594, lesinurad), 2-(3-cyano-4-isobutoxyphenyl)-4-methyl-1,3-thiazole-5-carboxylic acid (febuxostat), or BCX4208. In some embodiments the subject is refractory to allopurinol. For example, in one embodiment, the subject is refractory to allopurinol administered at from 100 mg/day to 800 mg/day (e.g. from 100 mg/day to 300 mg/day) for about one month or longer, about three months or longer, about one year or longer, etc. In some embodiments the subject is refractory to febuxostat. For example, in one embodiment the

subject is refractory to febuxostat administered at from 40 mg/day to 120 mg/day for about one month or longer, about three months or longer, about one year or longer, etc. In certain embodiments the subject has mild or moderate chronic kidney disease (CKD2-3). In other embodiments the subject has severe chronic kidney disease (CKD4). In other embodiments, the subject is on aspirin or diuretic therapy.

[0058] It will be recognized by persons with ordinary skill in the art that patients with gout or at risk of developing gout may be administered agents such as non-steroidal anti-inflammatory drugs (NSAIDS), colchicine, steroids, or similar medicaments to treat or manage gout flares. Accordingly, in certain embodiments of the methods described herein, the subjects may also be administered an agent such as an NSAID, colchicine or a steroid.

[0059] The methods described herein may be accomplished by the administration of a compound that generates the compound of Formula (IV) or a salt thereof via a chemical reaction after being administered. Such compounds include prodrugs of the compound of Formula (IV). Prodrugs of a compound are prepared by modifying functional groups present in the compound in such a way that the modifications may be cleaved *in vivo* to release the parent compound, or an active metabolite. For example, prodrugs include compounds wherein a hydroxy, amino, or sulfhydryl group in a compound is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Certain prodrugs may increase the bioavailability of the compounds of the embodiments when such compounds are administered to a subject (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a certain organ or tissue (e.g., kidneys, adipose tissue, liver, muscles or joints) relative to the parent species. Prodrugs of the compound of Formula (IV) include esters, amides, and carbamates (e.g., N, N-dimethylaminocarbonyl) of the hydroxy functional group of the compound of Formula (IV). The compounds of Formulae (I), (II), and (III) are non-limiting examples of prodrugs of the compound of Formula (IV). Further examples of prodrugs can be found in J. Rautio et al. *Prodrugs: design and clinical applications*, Nat. Rev. Drug Discov., 7, 255-270 (2008); Edward B. Roche, ed., *Bioreversible Carriers in Drug Design*, American Pharmaceutical Association and Pergamon Press, (1987); and T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems*, Vol. 14 of the A.C.S. Symposium Series (1975), each of which are hereby incorporated by reference herein.

[0060] In various embodiments, the compositions, methods, and kits described herein lower serum uric acid levels in a subject by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90% or more, as compared to serum uric acid levels in the subject prior to administering the methods described herein. In various embodiments, serum uric acid levels are decreased about 5% to about 50%, decreased by about 25% to about 75%, or decreased by about 50% to about 99%. Methods to determine serum uric acid levels are well known in the art and are often measured as part of a standard chemistry panel of blood serum samples.

[0061] In some embodiments, the compositions, methods, and kits of the present disclosure lower serum uric acid levels in a subject to about 7 mg/dL or less, to about 6.8 mg/dL or less, to about 6 mg/dL or less, to about 5 mg/dL or less, to about 4 mg/dL or less, or to about 3 mg/dL or less as compared to serum uric acid levels in the subject prior to administering the methods or compositions described herein. In some embodiments, the methods of the present disclosure lower serum uric acid levels in a subject by 0.1, 0.2, 0.3, 0.4, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5 or 10.0 mg/dL, or greater, as compared to serum uric acid levels in the subject prior to administering the methods or compositions described herein. In further embodiments, the methods described herein lower serum uric acid levels by between 0.1 and 10.0 mg/dL, between 0.5 and 6.0 mg/dL, between 1.0 and 4.0 mg/dL or between 1.5 and 2.5 mg/dL. The appropriate serum uric acid level may vary depending on the subject, and may vary for a given subject over time, depending upon the subject's overall medical condition. Similarly, the appropriate serum uric acid level for one group of subjects sharing a common medical condition may be different from that which is appropriate for a different group of subjects sharing a different medical condition. Thus, it may be advisable to reduce the serum uric acid level of a given group of subjects to, for example, below about 5 mg/dL, and to reduce the serum uric acid level of a different group of subjects to, for example, below about 4 mg/dL. In certain embodiments, the methods of the present disclosure decrease a serum uric acid level in the subject by an amount sufficient to result in the disappearance, reduction, amelioration, or the prevention of the onset, of one or more conditions associated with elevated serum uric acid over a certain timeframe, for example about four weeks or longer, about one month or longer, about three months or longer, about one year or longer, about two years or longer, etc. For example, a

method can decrease the serum uric acid level in a subject by an amount sufficient to result in the disappearance or reduction of tophi over about four weeks or longer, about one month or longer, about three months or longer, about one year or longer, about two years or longer, etc.

[0062] In further embodiments, the methods of the present disclosure comprise administering a pharmaceutical composition comprising a first urate-lowering agent and a second therapeutic agent, as described herein, to a subject whose serum uric acid level is at least about 4 mg/dL, at least about 5 mg/dL, at least about 6 mg/dL, at least about 6.8 mg/dL, at least about 7 mg/dL, at least about 8 mg/dL, at least about 9 mg/dL, at least about 10 mg/dL, or at least about 11 mg/dL. Again, the amount of decrease of serum uric acid level that is appropriate may vary depending on the subject, depending upon the subject's overall medical condition. Similarly, the amount of decrease of serum uric acid level that is appropriate for one group of subjects sharing a common medical condition may be different from that which is appropriate for a different group of subjects sharing a different medical condition.

[0063] The therapeutic agents and combinations thereof disclosed herein are contemplated to exhibit therapeutic activity when administered in an amount which can depend on the particular case. The variation in amount can depend, for example, on the subject being treated and the active ingredients chosen. A broad range of doses can be applicable. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, weekly, monthly or other at suitable time intervals or the dose may be proportionally reduced as indicated by the exigencies of the situation. Such dosages are optionally altered depending on a number of variables, not limited to the activity of the one or more active ingredients used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

[0064] As described throughout, the present disclosure contemplates combination therapy and methods of concomitant administration of a first and second urate-lowering agent (wherein these first and second urate-lowering agents are described above). Combination therapy and concomitant administration refer to the administration of the two agents (i.e., a first agent and a second urate-lowering agent, as described above) in any manner in which the pharmacological effects of both are manifested in the subject at the same time. Thus, such administration does not require that a single pharmaceutical composition, the same type of formulation, the same dosage

form, or even the same route of administration be used for administration of both the first and second urate-lowering agents, or that the two agents be administered at the same time. Such administration may be accomplished most conveniently by the same dosage form and the same route of administration, at substantially the same time. For example, a first urate-lowering agent, e.g. halofenate, halofenic acid, or a pharmaceutically acceptable salt thereof, and a second urate-lowering agent, e.g. xanthine oxidase inhibitor (e.g., allopurinol or febuxostat), can be administered to the human subject together in a single oral dosage composition, such as a tablet or capsule, or each agent can be administered in separate oral dosage formulations. One advantage with separate formulations is an added flexibility in dosing, i.e. the dosage of the first and second urate-lowering agents can be changed independently, quickly, and easily. Where separate dosage formulations are used, the first and second urate-lowering agents can be administered at essentially the same time (i.e., simultaneously or concurrently), or at separately staggered times (i.e., sequentially).

[0065] Depending on factors such as the diagnosis, symptoms, and therapeutic goals of a particular subject, a wide range of dosages of the first and second agent can be contemplated. In various embodiments, the first urate-lowering agents may be administered from about 10 mg to about 1000 mg per day and the second urate-lowering agent may be administered from about 10 mg to about 4000 mg per day. For example, halofenate, halofenic acid, or a pharmaceutically acceptable salt thereof may be administered at about 100 mg/day, about 200 mg/day, about 300 mg/day, about 400 mg/day, about 500 mg/day, about 600 mg/day, about 700 mg/day, about 800 mg/day, about 900 mg/day, or about 1000 mg/day.

[0066] As described above, in certain embodiments the second agent is allopurinol. The currently recommended daily dosage of allopurinol is from 100 mg/day to 800 mg/day in increments of 100 mg/day. When administered as a second urate-lowering agent as described herein (i.e. when a first urate-lowering agent is also administered), the dosage range of allopurinol may be within, above, or below the currently recommended daily dosage, as provided above and as appropriate for the subject being treated. By way of non-limiting example, in certain embodiments wherein the first urate-lowering agent is arhalofenate (i.e. (-)-halofenate) and wherein the second urate-lowering agent is allopurinol, the following daily dosages may be administered: arhalofenate 100 mg/day, allopurinol 50 mg/day; arhalofenate 100 mg/day, allopurinol 100 mg/day; arhalofenate 100 mg/day, allopurinol 200 mg/day; arhalofenate 100

mg/day, allopurinol 300 mg/day; arhalofenate 100 mg/day, allopurinol 400 mg/day; arhalofenate 100 mg/day, allopurinol 600 mg/day; arhalofenate 100 mg/day, allopurinol 800 mg/day; arhalofenate 200 mg/day, allopurinol 50 mg/day; arhalofenate 200 mg/day, allopurinol 100 mg/day; arhalofenate 200 mg/day, allopurinol 200 mg/day; arhalofenate 200 mg/day, allopurinol 300 mg/day; arhalofenate 200 mg/day, allopurinol 400 mg/day; arhalofenate 200 mg/day, allopurinol 600 mg/day; arhalofenate 200 mg/day, allopurinol 800 mg/day; arhalofenate 300 mg/day, allopurinol 50 mg/day; arhalofenate 300 mg/day, allopurinol 100 mg/day; arhalofenate 300 mg/day, allopurinol 200 mg/day; arhalofenate 300 mg/day, allopurinol 300 mg/day; arhalofenate 300 mg/day, allopurinol 400 mg/day; arhalofenate 300 mg/day, allopurinol 600 mg/day; arhalofenate 300 mg/day, allopurinol 800 mg/day; arhalofenate 400 mg/day, allopurinol 50 mg/day; arhalofenate 400 mg/day, allopurinol 100 mg/day; arhalofenate 400 mg/day, allopurinol 200 mg/day; arhalofenate 400 mg/day, allopurinol 300 mg/day; arhalofenate 400 mg/day, allopurinol 400 mg/day; arhalofenate 400 mg/day, allopurinol 600 mg/day; arhalofenate 400 mg/day, allopurinol 800 mg/day; arhalofenate 600 mg/day, allopurinol 50 mg/day; arhalofenate 600 mg/day, allopurinol 100 mg/day; arhalofenate 600 mg/day, allopurinol 200 mg/day; arhalofenate 600 mg/day, allopurinol 300 mg/day; arhalofenate 600 mg/day, allopurinol 400 mg/day; arhalofenate 600 mg/day, allopurinol 600 mg/day; arhalofenate 600 mg/day, allopurinol 800 mg/day.

[0067] Also as described above, in certain embodiments the second agent is febuxostat. The currently recommended daily dosage of febuxostat is 40 mg/day or 80 mg/day in the United States, and 40 mg/day, 80 mg/day or 120 mg/day in certain other countries. When administered as a second urate-lowering agent as described herein (i.e. when a first urate-lowering agent is also administered), the dosage range of febuxostat may be within, above, or below the currently recommended daily dosage, as provided above and as appropriate for the subject being treated. By way of non-limiting example, in certain embodiments wherein the first urate-lowering agent is arhalofenate (i.e. (-)-halofenate) and wherein the second urate-lowering agent is febuxostat, the following daily dosages may be administered: arhalofenate 100 mg/day, febuxostat 20 mg/day; arhalofenate 100 mg/day, febuxostat 40 mg/day; arhalofenate 100 mg/day, febuxostat 80 mg/day; arhalofenate 100 mg/day, febuxostat 120 mg/day; arhalofenate 200 mg/day, febuxostat 20 mg/day; arhalofenate 200 mg/day, febuxostat 40 mg/day; arhalofenate 200 mg/day, febuxostat 80 mg/day; arhalofenate 200 mg/day, febuxostat 120 mg/day; arhalofenate

300 mg/day, febuxostat 20 mg/day; arhalofenate 300 mg/day, febuxostat 40 mg/day; arhalofenate 300 mg/day, febuxostat 80 mg/day; arhalofenate 300 mg/day, febuxostat 120 mg/day; arhalofenate 400 mg/day, febuxostat 20 mg/day; arhalofenate 400 mg/day, febuxostat 40 mg/day; arhalofenate 400 mg/day, febuxostat 80 mg/day; arhalofenate 400 mg/day, febuxostat 120 mg/day; arhalofenate 600 mg/day, febuxostat 20 mg/day; arhalofenate 600 mg/day, febuxostat 40 mg/day; arhalofenate 600 mg/day, febuxostat 80 mg/day; arhalofenate 600 mg/day, febuxostat 120 mg/day; arhalofenate 800 mg/day, febuxostat 20 mg/day; arhalofenate 800 mg/day, febuxostat 40 mg/day; arhalofenate 800 mg/day, febuxostat 80 mg/day; arhalofenate 800 mg/day, febuxostat 120 mg/day.

[0068] Other dose ranges within the ranges described above for each of the first urate-lowering agent and the second urate-lowering agent may be readily envisaged. One of skill in the art will appreciate that the dose and dosing regimen may be adjusted when therapeutic agents are used in combination. When such combinations are used, the dose of one or more of the agents may be reduced to a level below the level required for a desired efficacy when the one or more agents are used alone. Similarly, the dosing regimen may be modified, e.g., to synchronize the dosing of the one or more therapeutic agents to facilitate improved patient ease of use and compliance. Alternately, the dosing regimen of the one or more therapeutic agents can be sequential, e.g., to reduce the combined load of the agents at a given time. For example, in certain embodiments, the dose of the second urate-lowering agent (e.g. allopurinol, febuxostat, or the other second urate-lowering agents described herein) can be adjusted to a lower level than that currently recommended when the first urate-lowering agent is and second urate-lowering agents are administered.

[0069] Dose titration or dose escalation protocols may be employed to determine the proper or optimal dose to administer to a subject. For example, dose titration or escalation studies may select for doses that improve efficacy or tolerability. Dose titration or escalation allows for the gradual adjusting of the dose administered until the desired effect is achieved. Dose titration gradually decreases the dosage administered while dose escalation gradually increases the dose administered. Methods of dose titration and escalation are well known in the art. As a non-limiting example, a subject may be administered 200 mg/day halofenate, halofenic acid, or a pharmaceutically acceptable salt thereof every day and measured for serum uric acid levels on a

daily basis. The dosage may be increased or decreased, for example, on a weekly basis. The subject may be monitored for a period of, for example, 2 to 12 weeks to find the desired dose.

[0070] In accordance with the compositions, methods and kits described herein, the first urate-lowering agent and second urate-lowering agent may be administered in any manner in which the pharmacological effects of both are likely to be manifested in the subject at approximately the same time. Such administration does not require that a single pharmaceutical composition, the same type of formulation, the same dosage form, or even the same route of administration be used for administration of both the first and second urate-lowering agents, or that the two agents be administered at the same time. That is, in various embodiments, the first urate-lowering agent and the second urate-lowering agent described herein may be present in a single dosage form (e.g. a single tablet or capsule for oral administration), and in other embodiments the first urate-lowering agent may be present in a first dosage form (e.g. a first tablet or capsule) and the second urate-lowering agent may be present in a second dosage form (e.g. a second tablet or capsule). The dosage forms may include the first and second urate-lowering agents in doses according to the examples provided above. A single dosage form (e.g. a single tablet or capsule) may include a single daily supply of the first and second urate-lowering agent, or a fraction thereof, e.g. one-half of a daily supply, one-third a daily supply, one-fourth a daily supply, etc. For example, the pharmaceutical composition described herein can be in a single tablet comprising 200 mg of arhalofenate and 150 mg of allopurinol. By way of further example, the pharmaceutical composition described herein can be in a single tablet comprising 200 mg of arhalofenate and 40 mg of febuxostat. Other dosage forms within the scope of this disclosure may be readily envisaged.

[0071] In unit dosage form, the formulation may be divided into unit doses containing appropriate quantities of the one or more active ingredients. In some embodiments, the unit dosage is in the form of a package containing discrete quantities of the formulation. Non-limiting examples include packaged tablets or capsules, and powders in vials or ampoules. In some embodiments, aqueous suspension compositions are packaged in single-dose non-reclosable containers. Alternatively, multiple-dose reclosable containers are used, in which case it is typical to include a preservative in the composition. By way of example only, formulations for parenteral injection are presented in unit dosage form, which include, but are not limited to ampoules, or in multi dose containers, with an added preservative. Tablets, troches, pills,

capsules and the like may also contain the components as listed hereafter: a binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain one or more active ingredients, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. In some embodiments, additional ingredients, for example, nonsteroidal anti-inflammatory drugs or colchicine, ingredients for treating other related indications, or inert substances such as artificial coloring agents are added. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the one or more active ingredients may be incorporated into sustained-release preparations and formulations as described herein.

[0072] The pharmaceutical compositions of the present disclosure may be administered once daily (QD), twice daily (BID), three times daily (TID) or four times per day (QID). In one embodiment, the pharmaceutical composition of the present disclosure is administered once daily (QD). In another embodiment, the pharmaceutical composition of the present disclosure is administered twice daily (BID).

[0073] This disclosure also describes articles of manufacture such as kits comprising a composition comprising a first and second urate-lowering agent (wherein these first and second urate-lowering agents are described above). In some embodiments the first urate-lowering agent in the kit is (-)-halofenate (i.e. arhalofenate). In some embodiments the second urate-lowering agent in the kit is allopurinol. In other embodiments the second urate-lowering agent in the kit is febuxostat. The kits can include the compositions packaged for distribution and in quantities sufficient to carry out the methods described herein. Kits may also include instructions (e.g. a package insert, package label, etc.) for using the kit components in one or more methods described herein. For example, a kit may comprise dosage forms of a first urate-lowering agent and a second urate-lowering agent described herein, and instructions for prescribing,

administering or otherwise using the dosage forms to lower serum uric acid levels. In some embodiments a kit is for a subject with hyperuricemia or a condition associated with hyperuricemia (e.g., gout) to use in the self-administration of the pharmaceutical composition, wherein the kit comprises a container housing a plurality of dosage forms containing a first and second urate-lowering agent described herein and instructions for carrying out drug administration therewith. In one embodiment, a kit comprises a first dosage form comprising halofenate, halofenic acid, or a pharmaceutically acceptable salt thereof in one or more of the forms identified above and at least a second dosage form comprising one or more of the forms identified above, in quantities sufficient to carry out the methods of the present disclosure. The second dosage form, and any additional dosage forms (e.g., a third, fourth or fifth dosage form) can comprise any active ingredient disclosed herein for the treatment of a hyperuricemic disorder (e.g., gout). All dosage forms together can comprise a therapeutically effective amount of each compound for the treatment of a condition associated with hyperuricemia (e.g., gout). In some embodiments a kit is for a subject with a condition associated with hyperuricemia (e.g., gout) to use in the self-administration of at least one oral agent, wherein the kit comprises a container housing a plurality of said oral agents and instructions for carrying out drug administration therewith. The embodiments of this application are characterized by the specification and by the features of the Claims of this application as filed, and of corresponding pharmaceutical compositions, methods and uses of these compounds.

Examples

Example 1: Clinical Trial

[0074] This is a randomized, double-blind, placebo-controlled study to evaluate the safety and efficacy of daily oral doses of between 400 to 600 mg of arhalofenate (i.e., (-)-halofenate) in combination with 300 mg oral doses of allopurinol will be conducted in approximately 90 gout patients with inadequate hypouricemic (uric acid lowering) response to allopurinol alone. Approximately 45 of these 90 patients will participate in a allopurinol/oxypurinol serial PK sample collection sub-study.

[0075] The 90 patients will be randomized in total; approximately 30 to each of 3 study arms in the following manner (1:1:1):

- 1) Arhalofenate 400 mg (plus allopurinol 300 mg)
- 2) Arhalofenate 600 mg (plus allopurinol 300 mg)

3) Placebo (plus allopurinol 300 mg)

[0076] Dose/Route/Regimen

[0077] Patients in all treatment groups will take colchicine 0.6 mg once daily by mouth starting at Week -3 through the final study follow-up visit, as background therapy for prophylaxis of gout flares. Patients in all treatment groups will also take allopurinol 300 mg once daily by mouth starting at Week -3 during the run-in and continuing through Week 4. Patients who fail to achieve target serum uric acid lowering on allopurinol alone will be randomized into the study.

[0078] The randomized treatment regimens will be the following (Day 1 through Week 4):

- Treatment Group #1: Arhalofenate 400 mg (plus allopurinol)
- Treatment Group #2: Arhalofenate 600 mg (plus allopurinol)
- Treatment Group #3: Placebo (plus allopurinol)

[0079] Analysis

[0080] All participating patients who will have received treatment in any dose group and for whom the primary PK data are considered to be sufficient and interpretable will be analyzed in the PK analyses. For patients included in the serial PK subset, allopurinol and oxypurinol pharmacokinetics in the presence and absence of arhalofenate at both doses will be determined from the repeat dose (Week 3, Visit 5) plasma concentrations, including the following PK parameters, as appropriate:

- Exposure, or area under the concentration-time curve (AUC0-24, AUC0-last,AUC0-inf)
- Maximum concentration (Cmax)
- Time to reach maximum concentration (Tmax)
- Terminal elimination half-life (t_{1/2})

[0081] Safety

[0082] The interpretation of the safety and tolerability will be made based on the assessment of safety parameters evaluated throughout the study, including clinical laboratory tests, 12-lead ECGs, vital signs, physical examination, concomitant medication review, and AEs (excluding medical events, which are “AEs” captured before dosing on Day 1). The reporting of the safety data is descriptive, and will include all patients receiving at least one dose of arhalofenate or allopurinol. Descriptive analysis will include the incidence and type of AEs by treatment group including tabulation by severity, as well as actual data and changes in laboratory, vital signs, and 12-lead ECG measurements from pre-dosing to all post-dosing time points.

[0083] Pharmacodynamics

[0084] The effects of each of the three treatment groups will be assessed as the absolute and percent change from baseline to each of the selected post-dosing time points for the following endpoints:

- sUA at Week 4 of treatment
- The proportion of patients achieving a sUA < 6 mg/dL
- The proportion of patients achieving a sUA < 5 mg/dL
- The proportion of patients achieving a sUA < 4 mg/dL

Example 2: Clinical Trial

[0085] This study evaluates the safety and efficacy of single oral doses of between 400 to 600 mg of arhalofenate in combination with 80 mg oral doses of febuxostat in approximately 10 to 15 gout patients for the treatment of hyperuricemia in patients with gout.

[0086] In addition to colchicine 0.6 mg daily for flare prophylaxis, all patients will receive febuxostat and arhalofenate in the following order during the Treatment Phase:

- Days 1 through 7: febuxostat 80 mg orally once daily (febuxostat only period)
- Days 8 through 21: febuxostat 80 mg plus arhalofenate 400 mg orally once daily (febuxostat plus arhalofenate 400 mg period)
- Days 22 through 35: febuxostat 80 mg plus arhalofenate 600 mg orally once daily (febuxostat plus arhalofenate 600 mg period)

[0087] The sUA level will be assessed on the last day of each treatment period (Day 7, Day 21, and Day 35); on these days, sUA samples will be collected at four different time points: pre-dose (fasting), 2 hours post-dose, 6 hours post-dose, and 10 hours post-dose (prior to evening meal).

[0088] Dose/Route/Regimen

- Colchicine: 0.6 mg /oral/daily from Day -16 through Day 49
- febuxostat: 80 mg/oral/daily from Day 1 through Day 35
- Arhalofenate: 400 mg/oral daily from Day 8 through Day 21; 600 mg/oral/daily from Day 22 through Day 35

[0089] Duration of Treatment

- Phase 1: Screening Phase: 1 to 4 weeks
- Phase 2: Run-in/Stabilization Phase: \geq 2 weeks
- Phase 3: Treatment Phase: 5 weeks
- Phase 4: Follow-up Phase: 2 weeks

[0090] Analysis

[0091] Safety

[0092] The interpretation of the safety and tolerability will be made based on the assessment of safety parameters evaluated throughout the study, including clinical laboratory tests, 12-lead ECGs, vital signs, physical examination, concomitant medication review, and AEs. The reporting of the safety data is descriptive, and will include all patients receiving at least one dose

of arhalofenate. Descriptive analysis will include the incidence and type of AEs including tabulation by severity, as well as actual data and changes in laboratory, vital signs, and 12-lead ECG measurements. Listings of clinically significant abnormal laboratory data will be presented. The case report form (CRF)-captured ECG and physical exam will be summarized using categorical statistical method.

[0093] Pharmacodynamics

[0094] The effects of each of the febuxostat plus arhalofenate combination treatment periods will be assessed as the change from baseline (Day 1) to end of treatment period for the following endpoints:

- The proportion of patients achieving a sUA < 6 mg/dL
- The proportion of patients achieving a sUA < 5 mg/dL
- The proportion of patients achieving a sUA < 4 mg/dL
- The proportion of patients achieving a sUA < 3 mg/dL
- Absolute and percent change in sUA

[0095] While the foregoing description describes specific embodiments, those with ordinary skill in the art will appreciate that various modifications and alternatives can be developed. Accordingly, the particular embodiments and examples described above are meant to be illustrative only, and not to limit the scope of the invention, which is to be given the full breadth of the appended claims, and any and all equivalents thereof.

[0096] In this specification where reference has been made to patent specifications, other external documents, or other sources of information, this is generally for the purpose of providing a context for discussing the features of the invention. Unless specifically stated otherwise, reference to such external documents is not to be construed as an admission that such documents, or such sources of information, in any jurisdiction, are prior art, or form part of the common general knowledge in the art.

[0097] In the description in this specification reference may be made to subject matter that is not within the scope of the claims of the current application. That subject matter should be readily identifiable by a person skilled in the art and may assist in putting into practice the invention as defined in the claims of this application.

WHAT WE CLAIM IS:

1. An oral dosage composition comprising:
 - (a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and
 - (b) febuxostat.
2. The composition of claim 1, where the compound is (-)-halofenate, substantially free from the corresponding (+)-enantiomer.
3. The composition of claim 1, where the compound is (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer.
4. The composition of any one of claims 1 to 3 where the amounts of the compound and of the febuxostat together are effective for once/day dosing.
5. The composition of any one of claims 1 to 4 where the amount of the compound is between about 10 mg and about 1000 mg.
6. The composition of claim 5 where the amount of the compound is between about 100 mg and about 800 mg.
7. The composition of any one of claims 1 to 6 where the amount of the febuxostat is between about 20 mg and about 120 mg.
8. The composition of claim 7 where the amount of the febuxostat is between about 40 mg and about 120 mg.
9. The composition of any one of claims 1 to 8 that is a tablet, a capsule, a syrup, or a suspension.
10. A method of treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout, comprising administering an effective amount of an oral dosage composition of any one of claims 1 to 9.
11. The method of claim 10 that is a method of treating hyperuricemia in a subject with gout.
12. The method of claim 10 that is a method of lowering serum uric acid in a subject with gout.
13. The method of claim 10 that is a method of reducing the size or number of tophi in a subject with gout.

14. The method of claim 10 that is a method of treating urate crystal formation in a subject with gout.

15. The use of:

(a) a compound that is (-)-halofenate, or (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer; and

(b) febuxostat,

in the manufacture of an oral dosage composition for treating hyperuricemia, or of lowering serum uric acid, or of reducing the size or number of tophi, or of treating urate crystal formation, in a subject with gout.

16. The use of claim 15, where the compound is (-)-halofenate, substantially free from the corresponding (+)-enantiomer.

17. The use of claim 15, where the compound is (-)-halofenic acid or a pharmaceutically acceptable salt thereof, substantially free from the corresponding (+)-enantiomer.

18. The use of any one of claims 15 to 17 where the amounts of the compound and of the febuxostat together are effective for once/day dosing.

19. The use of any one of claims 15 to 18 where the amount of the compound is between about 10 mg and about 1000 mg.

20. The use of claim 19 where the amount of the compound is between about 100 mg and about 800 mg.

21. The use of any one of claims 15 to 20 where the amount of the febuxostat is between about 20 mg and about 120 mg.

22. The use of claim 21 where the amount of the febuxostat is between about 40 mg and about 120 mg.

23. The use of any one of claims 15 to 22 where the oral dosage composition is a tablet, a capsule, a syrup, or a suspension.