



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/05/14
(87) Date publication PCT/PCT Publication Date: 2020/11/26
(85) Entrée phase nationale/National Entry: 2021/11/12
(86) N° demande PCT/PCT Application No.: US 2020/032950
(87) N° publication PCT/PCT Publication No.: 2020/236526
(30) Priorités/Priorities: 2019/05/23 (US62/852,262);
2019/07/22 (US62/877,183); 2019/08/12 (US62/885,746)

(51) Cl.Int./Int.Cl. *C07D 473/18* (2006.01),
C07D 473/34 (2006.01), *C07H 21/02* (2006.01)

(71) Demandeur/Applicant:
STRATOS GENOMICS, INC., US


(72) Inventeurs/Inventors:
O'CONNELL, DYLAN, US;
JACOBS, AARON, US;
VELLUCCI, JESSICA, US;
BANASIK, BRENT, US;
LOPEZ, MATTHEW, US;
GOODMAN, DREW, US;

...

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : ELEMENTS DE COMMANDE DE TRANSLOCATION, CODES RAPPORTEURS, ET AUTRES MOYENS DE
COMMANDE DE TRANSLOCATION DESTINES A ETRE UTILISES DANS LE SEQUENCAGE DE NANOPORES
(54) Title: TRANSLOCATION CONTROL ELEMENTS, REPORTER CODES, AND FURTHER MEANS FOR
TRANSLOCATION CONTROL FOR USE IN NANOPORE SEQUENCING

FIG. 4



(57) **Abrégé/Abstract:**

Phosphoramidate-based monomers are provided for use in the synthesis of expandable polymers for nanopore-based sensing. Such monomers comprising a reporter construct that contain a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code, wherein the ends of the reporter construct are attached to phosphoramidate-nucleoside. Related methods and products are also provided.

(72) **Inventeurs(suite)/Inventors(continued)**: NABAVI, MELUD, US; TABONE, JOHN, US; KOKORIS, MARK STAMATIOS, US; MACHACEK, CARA, US; MERRILL, LACEY, US; CHANDRASEKAR, JAGADEESWARAN, US; KRITZER, SVETLANA, US; CECH, CYNTHIA, US

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

26 November 2020 (26.11.2020)



(10) International Publication Number

WO 2020/236526 A1

(51) International Patent Classification:

C07D 473/18 (2006.01)*C07H 21/02* (2006.01)*C07D 473/34* (2006.01)

(21) International Application Number:

PCT/US2020/032950

(22) International Filing Date:

14 May 2020 (14.05.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/852,262 23 May 2019 (23.05.2019) US

62/877,183 22 July 2019 (22.07.2019) US

62/885,746 12 August 2019 (12.08.2019) US

(71) Applicant: STRATOS GENOMICS, INC. [US/US];

2211 Elliott Avenue, Suite 210, Seattle, Washington 98121 (US).


(72) Inventors: O'CONNELL, Dylan; 807 3rd Ave N., Apt.

46, Seattle, Washington 98109 (US). JACOBS, Aaron;

3500 S. Hudson Street, Seattle, Washington 98118 (US). **VELLUCCI, Jessica**; 1541 Marguerite Ave., Bremerton, Washington 98337 (US). **BANASIK, Brent**; 123 N. 145th St., Seattle, Washington 98133 (US). **LOPEZ, Matthew**; 8316 11th Ave NW, Seattle, Washington 98117 (US). **GOODMAN, Drew**; 1103 East Republican St., Apt. 4, Seattle, Washington 98102 (US). **NABAVI, Melud**; 2040 Waverly Place North #104, Seattle, Washington 98109 (US). **TABONE, John**; 7439 NE 123rd Place, Kirkland, Washington 98034 (US). **KOKORIS, Mark Stamatios**; 10535 Northeast 196th Street, Bothell, Washington 98011 (US). **MACHACEK, Cara**; 5332 8th Avenue NE, Seattle, Washington 98105 (US). **MERRILL, Lacey**; 321 10th Ave. S., #707, Seattle, Washington 98104 (US). **CHANDRASEKAR, Jagadeeswaran**; 937 NW 57th St., Seattle, Washington 98107 (US). **KRITZER, Svetlana**; 1900 1st Avenue, Apt. 607, Seattle, Washington 98101 (US). **CECH, Cynthia**; 7026 119th Place SE, Newcastle, Washington 98056 (US).

(54) Title: TRANSLOCATION CONTROL ELEMENTS, REPORTER CODES, AND FURTHER MEANS FOR TRANSLOCATION CONTROL FOR USE IN NANOPORE SEQUENCING

FIG. 4



(57) Abstract: Phosphoramidate-based monomers are provided for use in the synthesis of expandable polymers for nanopore-based sensing. Such monomers comprising a reporter construct that contain a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code, wherein the ends of the reporter construct are attached to phosphoramidate-nucleoside. Related methods and products are also provided.

[Continued on next page]



WO 2020/236526 A1

WO 2020/236526 A1

(74) **Agent:** HERMANN, Karl, R. et al.; Seed Intellectual Property Law Group LLP, Suite 5400, 701 Fifth Avenue, Seattle, Washington 98104-7064 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *of inventorship (Rule 4.17(iv))*

Published:

— *with international search report (Art. 21(3))*
 — *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

TRANSLOCATION CONTROL ELEMENTS, REPORTER CODES, AND FURTHER MEANS FOR TRANSLOCATION CONTROL FOR USE IN NANOPORE SEQUENCING

BACKGROUND

Technical Field

[0001] The present invention relates generally to new synthetic reporter constructs, more specifically to new nucleotide-free, phosphoramidite-based translocation control elements, reporter codes and other features that generate unique signals when passed through a nanopore, and methods for the manufacture and utilization thereof, particularly in nanopore-based polymer sequencing methods.

Description of the Related Art

[0002] Measurement of biomolecules is a foundation of modern medicine and is broadly used in medical research, and more specifically in diagnostics and therapy, as well in drug development. Nucleic acids encode the necessary information for living things to function and reproduce, and are essentially a blueprint for life. Determining such blueprints is useful in pure research as well as in applied sciences. In medicine, sequencing can be used for diagnosis and to develop treatments for a variety of pathologies, including cancer, heart disease, autoimmune disorders, multiple sclerosis, and obesity. In industry, sequencing can be used to design improved enzymatic processes or synthetic organisms. In biology, this tool can be used to study the health of ecosystems, for example, and thus have a broad range of utility. Similarly, measurement of proteins and other biomolecules has provided markers and understanding of disease and pathogenic propagation.

[0003] An individual's unique DNA sequence provides valuable information concerning their susceptibility to certain diseases. It also provides patients with the opportunity to screen for early detection and/or to receive preventative treatment. Furthermore, given a patient's individual blueprint, clinicians will be able to administer personalized therapy to maximize drug efficacy and/or to minimize the risk of an adverse drug response. Similarly, determining the blueprint of pathogenic organisms can lead to new treatments for infectious diseases and more robust pathogen surveillance. Low cost, whole genome DNA sequencing will provide the foundation for modern medicine. To achieve this goal, sequencing technologies must continue to advance with respect to throughput, accuracy, and read length.

[0004] Over the last decade, a multitude of next generation DNA sequencing technologies have become commercially available and have dramatically reduced the cost of sequencing whole genomes. These include sequencing by synthesis ("SBS") platforms (Illumina, Inc., 454 Life Sciences, Ion Torrent, Pacific Biosciences) and analogous ligation based platforms (Complete Genomics, Life Technologies Corporation). A number of other technologies are being developed that utilize a wide variety of sample processing and detection methods. For example, GnuBio, Inc. (Cambridge, Mass.) uses picoliter reaction vessels to control millions of discrete probe sequencing reactions, whereas Halcyon Molecular (Redwood City, Calif.) was attempting to develop technology for direct DNA measurement using a transmission electron microscope.

[0005] Nanopore based nucleic acid sequencing is a compelling approach that has been widely studied. Kasianowicz et al. (Proc. Natl. Acad. Sci. USA 93: 13770-13773, 1996) characterized single-stranded polynucleotides as they were electrically translocated through an alpha hemolysin nanopore embedded in a lipid bilayer. It was demonstrated that during polynucleotide translocation partial blockage of the nanopore aperture could be measured as a decrease in ionic current. Polynucleotide sequencing in nanopores, however, is burdened by having to resolve tightly spaced bases (0.34 nm) with small signal differences immersed in significant background noise. The measurement challenge of single base resolution in a nanopore is made more demanding due to the rapid translocation rates observed for polynucleotides, which are typically on the order of 1 base per microsecond. Translocation speed can be reduced by adjusting run parameters such as voltage, salt composition, pH, temperature, and viscosity, to name a few. However, such adjustments have been unable to reduce translocation speed to a level that allows for single base resolution.

[0006] Stratos Genomics has developed a method called Sequencing by Expansion ("SBX") that uses a biochemical process to transcribe the sequence of DNA onto a measurable polymer called an "Xpandomer" (Kokoris et al., U.S. Pat. No. 7,939,259, "High Throughput Nucleic Acid Sequencing by Expansion"). The transcribed sequence is encoded along the Xpandomer backbone in high signal-to-noise reporters that are separated by ~10 nm and are designed for high-signal-to-noise, well-differentiated responses. These differences provide significant performance enhancements in sequence read efficiency and accuracy of Xpandomers relative to native DNA. Xpandomers can enable several next generation DNA sequencing detection technologies and are well suited to nanopore sequencing.

[0007] Nanopores have proven to be powerful amplifiers, much like their highly-exploited predecessors, Coulter Counters. However, the current generation of organic

nanopores (such as Hemolysin and MspA), that have been tasked with base recognition of DNA, are transmembrane proteins that do not interact with DNA in nature. They do not have natural functions for controlling DNA translocation. This is a recognized shortcoming that some have attempted to correct by adding functionality with protein motors adjacent to the nanopores. For example, Akeson's group added phi 29 polymerase adjacent to the alpha-hemolysin nanopore so that ss-DNA could be fed into the pore at a controlled rate (see G. M. Cherf et al. "Automated forward and reverse ratcheting of DNA in a nanopore at 5-A precision," Nat Biotech, vol. advance online publication, February 2012). This approach complicates the assay and imposes a separation of the measurement region in the alpha hemolysin from the position control in the polymerase that can introduce additional noise and sequence dependent variation to the measurement.

[0008] In another approach, referred to as translocation control by hybridization (TCH), a nanopore translocation event is paused by using a structure created by hybridization, which disassociates for translocation to proceed (see, e.g. US Patent No. 10,457,979 to McRuer and Kokoris). Akeson et al. (U.S. Pat. No. 6,465,193) first demonstrated this by pausing DNA translocation with sequential hairpin duplexed regions. Translocation stopped at the duplex because it was larger than the alpha-hemolysin nanopore aperture. When the duplex released due to stochastic thermal fluctuation, translocation proceeded to the next duplex. During each pause, the region of DNA located in the nanopore (adjacent to the duplex) could be measured and identified. When applied to nanopore sequencing, this duplexing approach to translocation control suffers from limitations, including incomplete duplex formation, or hybridization fill rate, and the stochastics of duplex dissociation, which can lead to deletions or insertions events. Insertion and deletions that cannot be localized can seriously degrade the data quality.

[0009] While significant advances have been made in this field, commercially viable implementation of translocation control with, for example, Xpandomers, would benefit from improvements that overcome limitations caused by duplexing. The present invention fulfills these needs and provides further related advantages as discussed below.

[0010] All of the subject matter discussed in the Background section is not necessarily prior art and should not be assumed to be prior art merely as a result of its discussion in the Background section. Along these lines, any recognition of problems in the prior art discussed in the Background section or associated with such subject matter should not be treated as prior art unless expressly stated to be prior art. Instead, the discussion of any subject matter in the Background section should be treated as part of the inventor's


approach to the particular problem, which in and of itself may also be inventive.

BRIEF SUMMARY OF THE INVENTION

[0011] In brief, compounds (e.g., XNTPs) including polymeric reporter and linker constructs synthesized from a collection of novel phosphoramidite monomeric units and methods are disclosed for improved nanopore sequencing (for example, generating sequences of higher read length, accuracy, and/or throughput) of polymeric analytes (e.g., Xpandomers).

[0012] In some embodiments, the polymeric constructs may be designed to completely lack nucleotides.

[0013] In one aspect, the present disclosure provides a compound (i.e., an XNTP) having the following structure:



wherein R is OH or H; nucleobase is adenine, cytosine, guanine, thymine, uracil or a nucleobase analog; reporter construct is a polymer having a first end and a second end, and includes, in series from the first end to the second end, a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code; linker A joins the oxygen atom of the alpha phosphoramidate to the first end of the reporter construct; linker B joins the nucleobase to the second end of the reporter construct; and in which the translocation control element is a polymer as described below.

[0014] In one embodiment, the translocation control element is a polymer comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole-propane (compound 35b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 16), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-

nitroindole)-propane (compound 20g), 1,8-O-bis(phosphodiester)-N,N-Diethylpiperazine (compound 26h), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47Gg, 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), or 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52).

[0015] In some embodiments, R is OH.

[0016] In some embodiments, R is H.

[0017] In other embodiments, nucleobase is adenine, cytosine, guanine, thymine, or uracil.

[0018] In other embodiments, nucleobase is a nucleobase analog.

[0019] In other embodiments, the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphodiester)-propane, 1,3-bis-(5-O-phosphodiester-pentylamido)-2-O-phosphodiester-propane, or 1,4,7-O-tris-(phosphodiester)-heptane.

[0020] In other embodiments, the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphodiester)-propane.

[0021] In other embodiments, the translocation control element is a polymer comprising two or more repeat units selected from Table 1A.

[0022] In other embodiments, the translocation control element is a polymer comprising two or more repeat units selected from 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b) and 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b).

[0023] In yet other embodiments, the translocation control element is a polymer comprising the following sequence: [(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))]n1[(1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b))]n2, wherein n1 is from 0 to 6 and n2 is from 6 to 10.

[0024] In other embodiments, the first and second reporter codes are identical.

[0025] In further embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from: hexaethylene glycol (D), ethane (L), triaethylene glycol (X), 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane (compound 31a), 2,3-O-bis(phosphodiester)-1-(1-dimethoxyquinazolinedione)-propane (compound 20c), 2,3-O-bis(phosphodiester)-1-(N9-(3,6-dimethoxycarbazole)-propane (compound 20e), 1,1'-O-bis(phosphodiester)-2,2'-(sulfonylbis(benz-4-yl))-diethanol (compound 26d), 1,1'-O-bis(phosphodiester)-2,2'-bipyridin-4,4'-yl)-dimethanol (compound 26a), 2,3-O-bis(phosphodiester)-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane (compound 20b), 3-(1,2-O-bis(phosphodiester)-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide (compound 20d), 2,3-O-bis(phosphodiester)-1-(N1-(6-Azathymine))-propane (compound 20f), 1,5-O-bis(phosphodiester)-hexahydrofuro[2,6]furan (compound 23), 1,1'-O-bis(phosphodiester)-octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol (compound 26e), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20h), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 5b), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane (compound 31b), and 1,3-O-bis(phosphodiester)-3-O-mPEG4-propane (compound 5a).

[0026] In other embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene

glycol, and any of the compounds set forth in Table 1A.

[0027] In further embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b).

[0028] In more specific embodiments, the first and second reporter codes are polymers comprising a sequence selected from: (i) [(hexaethylene glycol)2(ethane)3(hexaethylene glycol)(triaethylene glycol)], (ii) [(hexaethyleneglycol)2(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))2(ethane)(triaethylene glycol)3], (iii) [(hexaethylene glycol)2(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))3(ethane)2(hexaethylene glycol)(triaethylene glycol)], and (iv) [(triaethylene glycol)2(ethane)(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))6(ethane)7].

[0029] In other embodiments, Linker A and Linker B are polymers comprising two or more repeat units selected from: spermine (Q), hexaethylene glycol (D), 2-(((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((phosphodiester-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-phosphodiester-propane-1,3-diyl dibenzoate (compound 62), 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47g), and 1,1'-O-bis(phosphodiester)-N(p-tolyl)-diethanolamine (compound 26b).

[0030] In other embodiments, Linker A and Linker B are polymers comprising two or more repeat units selected from spermine and any of the compounds set forth in Table 1A.

[0031] In yet other embodiments, Linker A and Linker B comprise a polymerase

enhancement region comprising two repeat units of spermine.

[0032] In further embodiments, Linker A and Linker B comprise a translocation deceleration region comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), and 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b).

[0033] In more specific embodiments, Linker A and Linker B comprise a translocation deceleration region comprising a polymer selected from: (i) [((hexaethylene glycol) (1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))3(hexaethylene glycol)2], (ii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))4(hexaethylene glycol)2], (iii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d))4(hexaethylene glycol)2], and (iv) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b))4(hexaethylene glycol)2].

[0034] In other embodiments, Linker A is joined to the oxygen atom of the alpha phosphoramidate by a linkage comprising a triazole and Linker B is joined to the nucleobase by a linkage comprising a triazole.

[0035] In another aspect, the present invention provides a reporter construct comprising a polymer having a first end and a second end, and including in series from the first end to the second end a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code; and in which the translocation control element is a polymer comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-(Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 16), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 1,8-O-bis(phosphodiester)-N,N-Diethylpiperazine (compound 26h), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound

35a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47Gg, 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), or 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52).

[0036] In some embodiments, the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphosphodiester)-propane, 1,3-bis-(5-O-phosphodiester-pentylamido)-2-O-phosphodiester-propane, or 1,4,7-O-tris-(phosphodiester)-heptane.

[0037] In another embodiment, the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphosphodiester)-propane.

[0038] In other embodiments, the translocation control element is a polymer comprising two or more repeat units selected from Table 1A.

[0039] In yet other embodiments, the translocation control element is a polymer comprising two or more repeat units selected from 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b) and 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b).

[0040] In further embodiments, the translocation control element is a polymer comprising the following sequence: [(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))]n1[(1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b))]n2, wherein n1 is from 0 to 6 and n2 is from 6 to 10.

[0041] In other embodiments, the first and second reporter codes are identical.

[0042] In some embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from: hexaethylene glycol (D), ethane (L), triaethylene glycol (X), 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane

(compound 35b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane (compound 31a), 2,3-O-bis(phosphodiester)-1-(1-dimethoxyquinazolin-2-one)-propane (compound 20c), 2,3-O-bis(phosphodiester)-1-(N9-(3,6-dimethoxycarbazole)-propane (compound 20e), 1,1'-O-bis(phosphodiester)-2,2'-(sulfonylbis(benz-4-yl))-diethanol (compound 26d), 1,1'-O-bis(phosphodiester)-2,2'-bipyridin-4,4'-yl)-dimethanol (compound 26a), 2,3-O-bis(phosphodiester)-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane (compound 20b), 3-(1,2-O-bis(phosphodiester)-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide (compound 20d), 2,3-O-bis(phosphodiester)-1-(N1-(6-Azathymine))-propane (compound 20f), 1,5-O-bis(phosphodiester)-hexahydrofuro[2,6]furan (compound 23), 1,1'-O-bis(phosphodiester)-octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol (compound 26e), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20h), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 5b), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane (compound 31b), and 1,3-O-bis(phosphodiester)-3-O-mPEG4-propane (compound 5a).

[0043] In other embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and any of the compounds set forth in Table 1A.

[0044] In further embodiments, the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b).

[0045] In further embodiments, the first and second reporter codes are polymers comprising a sequence selected from: (i) [(hexaethylene glycol)2(ethane)3(hexaethylene glycol)(triaethylene glycol)], (ii) [(hexaethyleneglycol)2(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))2(ethane)(triaethylene glycol)3], (iii) [(hexaethylene glycol)2(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound

12b))3(ethane)2(hexaethylene glycol)(triaethylene glycol)], and (iv) [(triaethylene glycol)2(ethane)(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))6(ethane)7].

[0046] In another aspect, the present invention provides a symmetrically synthesized report tether (SSRT), in which the symmetrically synthesized reporter tether is a polymer having a first end and a second end, and includes in series from the first end to the second end a first linker, a reporter construct according to any one of the above reporter constructs, and a second linker, in which the first and second linkers are identical and are polymers comprising two or more repeat units selected from: spermine (Q), hexaethylene glycol (D), 2-(((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((phosphodiester-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-phosphodiester-propane-1,3-diyl dibenzoate (compound 62), 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47g), and 1,1'-O-bis(phosphodiester)-N(p-tolyl)-diethanolamine (compound 26b).

[0047] In some embodiments, the symmetrically synthesized reporter tether (SSRT) includes a polymerase enhancement region comprising two repeat units of spermine.

[0048] In other embodiments, the symmetrically synthesized reporter tether (SSRT) includes a translocation deceleration region comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), and 1,3-O-

bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b).

[0049] In other embodiments, the symmetrically synthesized reporter tether (SSRT) includes a translocation deceleration region comprising a polymer selected from: (i) [((hexaethylene glycol) (1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac))-1,2,3-triazole)-propane (compound 35c))3(hexaethylene glycol)2], (ii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac))-1,2,3-triazole)-propane (compound 35c))4(hexaethylene glycol)2], (iii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz))-1,2,3-triazole)-propane (compound 35d))4(hexaethylene glycol)2], and (iv) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b))4(hexaethylene glycol)2].

[0050] In yet other embodiments, the first end and the second end of the symmetrically synthesized reporter tether (SSRT) include a linkage moiety and, in certain embodiments, the linkage moiety is an azido ($-N_3$) group.

[0051] In another aspect, the present invention provides a method for sequencing a target nucleic acid, comprising the steps of: a) providing a daughter strand produced by a template-directed synthesis, the daughter strand comprising a plurality of XNTP subunits coupled in a sequence corresponding to a contiguous nucleotide sequence of all or a portion of the target nucleic acid, wherein the individual XNTP subunits of the daughter strand comprise a reporter construct, a nucleobase residue, and a selectively cleavable bond, and wherein the reporter construct, upon cleavage of the selectively cleavable bond, permits lengthening of the subunits of the daughter strand; b) cleaving the selectively cleavable bonds to yield an Xpandomer of a length longer than the plurality of the subunits of daughter strand, the Xpandomer comprising the reporter constructs for parsing genetic information in a sequence corresponding to the contiguous nucleotide sequence of all or a portion of the target nucleic acid; and c) detecting the reporter constructs of the Xpandomer.

[0052] In some aspects, the reporter constructs for parsing the genetic information comprise a reporter code and a translocation control element, wherein the translocation control element provides translocation control by steric hindrance and pauses translocation of the Xpandomer when passed through a nanopore subjected to a baseline voltage, wherein the translocation control element engages the reporter code within the aperture of the nanopore, wherein the reporter code is sensed by the nanopore.

[0053] In some embodiments, the Xpandomer resumes translocation through the

nanopore by application of a pulse voltage, in which the pulse voltage is sufficient to allow translocation of the translocation control element, while leaving the next reporter construct of the Xpandomer free to engage with the nanopore.

[0054] In other embodiments, the translocation control element of the reporter construct engaged with the nanopore by steric hindrance translocates upon each pulse of the pulsed voltage.

[0055] In some embodiments, the target construct is sensed by the nanopore during the time period between pulses of the pulsed voltage.

[0056] In certain embodiments, the baseline voltage is from about 55mV to about 75mV and the pulse voltage is from about 550mV to about 700mV.

[0057] In some embodiments, the pulse voltage has a duration from about 5 μ s to about 10 μ s and a periodicity from about 0.5ms to 1.5ms.

[0058] In other embodiments, the nanopore is subjected to an alternating current (AC).

[0059] In further embodiments, one or more of the XNTP subunits includes a 2' Fluoroarabinosyl epimer.

[0060] In another aspect, the present disclosure provides a buffer for controlling the rate of translocation of a polymer through a nanopore comprising at least one salt selected from the group consisting of NH₄Cl, MgCl₂, LiCl, KCl, CsCl, NaCl, and CaCl₂.

[0061] In some embodiments, the buffer further comprises at least one solvent selected from the group consisting of 3-methyl-2-oxazolidinone (MOA), DMF, ACN, DMSO, and NMP, wherein the solvent is present in the range from about 1% vol/vol to about 35% vol/vol.

[0062] In other embodiments, the buffer further comprises at least one additive selected from the group consisting of sodium hexanoate (NaHex), EDTA, redox reagents, PEG, glycerol, ficoll, and the like.

[0063] In another aspect, the present disclosure provides a buffer system for controlling the rate of translocation of a polymer through a nanopore detector comprising a cis buffer and a trans buffer, wherein the cis buffer has a first salt concentration and the trans buffer has a second salt concentration, wherein the first salt concentration is lower than the second salt concentration.

BRIEF DESCRIPTION OF THE DRAWINGS

[0064] FIGS. 1A, 1B, 1C and 1D are condensed schematics illustrating the main features of a generalized XNTP and their functions in Sequencing by Expansion (SBX).

[0065] FIG. 2 is a schematic illustrating more details of one embodiment of an XNTP.

[0066] FIG. 3 is a schematic illustrating one embodiment of an Xpandomer passing through a biological nanopore.

[0067] FIG. 4 is a schematic illustrating another embodiment of an Xpandomer passing through a biological nanopore.

[0068] FIG.s 5A – 5D are schematics illustrating alternative embodiments of reporter codes.

[0069] FIG. 6 is a schematic illustrating one embodiment of solid state synthesis of a SSRT reporter construct.

[0070] FIG. 7 illustrates alternative structural embodiments of SSRT reporter constructs.

[0071] FIG.s 8A and 8B are schematics illustrating one embodiment of the cyclization of an SSRT and a dNTP-2c to form an XNTP.

[0072] FIG.s 9A and 9B are schematics illustrating one embodiment of translocation control of an Xpandomer through a nanopore.

[0073] FIG. 10 is a schematic illustrating one embodiment of a biotin derivative.

[0074] FIG. 11 is a schematic illustrating one embodiment of a cleavable extension oligonucleotide.

[0075] FIG. 12 is a schematic illustrating one embodiment of a polymer subjected to ratcheting through a nanopore.

[0076] FIG.s 13A and 13B are representative traces depicting properties of reporter codes.

[0077] FIG.s 14A and 14B are histogram displays of populations of aligned reads of nanopore-derived sequences.

[0078] FIG. 15 is a representative trace showing the sequence of a simple DNA template.

[0079] FIG. 16 is a representative trace showing the sequence of a CAGT repeat DNA template.

[0080] FIG. 17 is a representative trace showing the sequence of a complex DNA

template.

[0081] FIG. 18 is a representative trace showing the sequence of a complex DNA 222mer template.

[0082] FIG. 19 is a histogram display of a population of aligned reads of nanopore-derived sequences.

[0083] Fig. 20A is a schematic illustrating one embodiment of an Xpandomer subjected to ratcheting through a nanopore.

[0084] Fig. 20B is an example of the current measurement for a translocating Xpandomer subjected to ratcheting.

DETAILED DESCRIPTION OF THE INVENTION

[0085] The present invention may be understood more readily by reference to the following detailed description of preferred embodiments of the invention and the Examples included herein. Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs.

[0086] Reference throughout this specification to “one embodiment” or “an embodiment” and variations thereof means that a particular feature, structure, or characteristic described in connection with the embodiment is included in at least one embodiment. Thus, the appearances of the phrases “in one embodiment” or “in an embodiment” in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments.

[0087] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents, i.e., one or more, unless the content and context clearly dictates otherwise. It should also be noted that the conjunctive terms, “and” and “or” are generally employed in the broadest sense to include “and/or” unless the content and context clearly dictates inclusivity or exclusivity as the case may be. Thus, the use of the alternative (e.g., “or”) should be understood to mean either one, both, or any combination thereof of the alternatives. In addition, the composition of “and” and “or” when recited herein as “and/or” is intended to encompass an embodiment that includes all of the associated items or ideas and one or more other alternative embodiments that include fewer than all of

the associated items or ideas.

[0088] Unless the context requires otherwise, throughout the specification and claims that follow, the word “comprise” and synonyms and variants thereof such as “have” and “include”, as well as variations thereof such as “comprises” and “comprising” are to be construed in an open, inclusive sense, e.g., “including, but not limited to.” The term “consisting essentially of” limits the scope of a claim to the specified materials or steps, or to those that do not materially affect the basic and novel characteristics of the claimed invention.

[0089] The abbreviation, “e.g.” is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation “e.g.” is synonymous with the term “for example.” It is also to be understood that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise, the term “X and/or Y” means “X” or “Y” or both “X” and “Y”, and the letter “s” following a noun designates both the plural and singular forms of that noun. In addition, where features or aspects of the invention are described in terms of Markush groups, it is intended, and those skilled in the art will recognize, that the invention embraces and is also thereby described in terms of any individual member and any subgroup of members of the Markush group, and Applicants reserve the right to revise the application or claims to refer specifically to any individual member or any subgroup of members of the Markush group.

[0090] Any headings used within this document are only being utilized to expedite its review by the reader, and should not be construed as limiting the invention or claims in any manner. Thus, the headings and Abstract of the Disclosure provided herein are for convenience only and do not interpret the scope or meaning of the embodiments.

[0091] Where a range of values is provided herein, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0092] For example, any concentration range, percentage range, ratio range, or integer range provided herein is to be understood to include the value of any integer within the

recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated. Also, any number range recited herein relating to any physical feature, such as polymer subunits, size or thickness, are to be understood to include any integer within the recited range, unless otherwise indicated. As used herein, the term "about" means $\pm 20\%$ of the indicated range, value, or structure, unless otherwise indicated.

Sequencing by Expansion

[0093] The "Sequencing by Expansion" (SBX) protocol, developed by Stratos Genomics (see, e.g., Kokoris et al., U.S. Pat. No. 7,939,259, "High Throughput Nucleic Acid Sequencing by Expansion") is based on the polymerization of highly modified, non-natural nucleotide analogs referred to as "XNTPs". In general terms, SBX uses biochemical polymerization to transcribe the sequence of a DNA template onto a measurable polymer called an "Xpandomer". The transcribed sequence is encoded along the Xpandomer backbone in high signal-to-noise reporters that are separated by ~ 10 nm and are designed for high-signal-to-noise, well-differentiated responses. These differences provide significant performance enhancements in sequence read efficiency and accuracy of Xpandomers relative to natural DNA. A generalized overview of the SBX process is depicted in FIGS. 1A, 1B, 1C and 1D.

[0094] XNTPs are expandable, 5' triphosphate modified non-natural nucleotide analogs compatible with template dependent enzymatic polymerization. A highly simplified XNTP is illustrated in FIG. 1A, which emphasizes the unique features of these non-natural substrates: XNTP **100** has two distinct functional regions; namely, a selectively cleavable phosphoramidate bond **110**, linking the 5' α -phosphate **115** to the nucleobase **105**, and a symmetrically synthesized reporter tether (SSRT) **120** that is attached within the nucleoside triphosphoramidate at positions that allow for controlled expansion by cleavage of the phosphoramidate bond. The SSRT includes linkers **125A** and **125B** separated by the selectively cleavable phosphoramidate bond. Each linker attaches to one end of a reporter code **130**. XNTP **100** is illustrated in the "constrained configuration", characteristic of the XNTP substrates and the daughter strand products of template-dependent polymerization. The constrained configuration of polymerized XNTPs is the precursor to the expanded configuration, as found in Xpandomer products. The transition from the constrained configuration to the expanded configuration occurs upon scission of the P--N bond of the phosphoramidate within the primary backbone of the daughter strand.


[0095] Synthesis of an Xpandomer polymer is summarized in FIGS. 1B and 1C. During assembly, the monomeric XNTP substrates **145** (XATP, XCTP, XGTP and XTPP) are polymerized on the extendable terminus of a nascent daughter strand **150** by a process of template-directed polymerization using single-stranded template **140** as a guide. Generally, this process is initiated from a primer and proceeds in the 5' to 3' direction. Generally, a DNA polymerase or other polymerase is used to form the daughter strand, and conditions are selected so that a complimentary copy of the template strand is obtained. After the daughter strand is synthesized, the coupled SSRTs form the constrained Xpandomer that further forms the daughter strand. SSRTs in the daughter strand have the "constrained configuration" of the XNTP substrates. The constrained configuration of the SSRT is the precursor to the expanded configuration, as found the Xpandomer product.

[0096] As shown in FIG. 1C, the transition from the constrained configuration **160** to the expanded configuration **165** results from cleavage of the selectively cleavable phosphoramidate bonds (illustrated for simplicity by the unshaded ovals) within the primary backbone of the daughter strand. In this embodiment, the SSRTs include one or more reporters or reporter codes, **130A**, **130C**, **130G**, or **130T**, specific for the nucleobase to which they are linked, thereby encoding the sequence information of the template. In this manner, the SSRTs provide a means to expand the length of the Xpandomer and lower the linear density of the sequence information of the parent strand.

[0097] FIG. 1D illustrates an Xpandomer **165** translocating through a nanopore **180**, from the cis reservoir **175** to the trans reservoir **185**. Upon passage through the nanopore, each of the reporter codes of the linearized Xpandomer (in this illustration, labeled "G", "C" and "T") generates a distinct and reproducible electronic signal (illustrated by superimposed trace **190**), specific for the nucleobase to which it is linked.

[0098] FIG. 2 depicts the generalized structure of one embodiment of an XNTP in more detail. XNTP **200** includes nucleoside triphosphoramidate **210** with linker arm moieties **220A** and **220B** separated by selectively cleavable phosphoramidate bond **230**. SSRTs are joined to the nucleoside triphosphoramidate at linkage groups **250A** and **250B**, in which a first SSRT end is joined to the heterocycle **260** (represented here by cytosine, though the heterocycle may be any one of the four standard nucleobases, A, C, G, or T) and a second SSRT end is joined to the alpha phosphate **270** of the nucleobase backbone. The skilled artisan will appreciate that many suitable coupling chemistries known in the art may be used to form the final XNTP substrate product, for example, SSRT conjugation may be accomplished through formation of a triazole linkage group.

[0099] In this embodiment, SSRT **275** includes several functional elements, or “features” such as polymerase enhancement regions **280A** and **280B**, reporter codes **285A** and **285B**, and translation control element (TCEs) **290A** and **290B**. In other embodiments, the SSRT includes a single TCE. Each of these features performs a unique function during translocation of the Xpandomer through a nanopore to produce a series of unique and reproducible electronic signal. SSRT **275** is designed for controlling the rate of Xpandomer translocation by the TCE through a combination of sterics and/or electrorepulsion, as discussed further herein. Different reporter codes are sized to block ion flow through a nanopore at different measureable levels. Specific SSRT polymeric sequences can be efficiently synthesized using phosphoramidite chemistry typically used for oligonucleotide synthesis. Reporter codes and other features can be designed by selecting a sequence of specific phosphoramidites from commercially available and/or proprietary libraries. Such libraries include, but are not limited to, polyethylene glycol with lengths of 1 to 12 or more ethylene glycol units and aliphatic polymers with lengths of 1 to 12 or more carbon units. In certain embodiments, the SSRTs include features referred to as “polymerase enhancement regions” at the ends of the SSRTs proximal to the nucleotide triphosphoramidate diester. Polymerase enhancement regions may include positively charged polyamine spacers (e.g., primary, secondary, tertiary, or quarternary amines) or triamine spacers (three secondary amines each separated by three carbons) that facilitate incorporation of XNTP structures by a nucleic acid polymerase. In certain embodiments, the polymerase enhancement region includes two repeat units of spermine, in which the spermine moiety is provided by a phosphoramidite monomer having the following structure (as one of skill in the art will recognize, the trifluoroacetamide protecting groups are removed at the end of SSRT synthesis to expose the amine groups on spermine):




[00100] As used throughout the present disclosure, the term “reporter construct” refers to the element of the SSRT that includes the reporter codes, a symmetrical chemical brancher, and a translocation control element. In certain embodiments, the reporter construct is a polymer that includes, in series, from a first end to a second end, a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code. The term “bearing” refers to a covalent linkage between the symmetrical brancher and

the translocation control element, which produces an advantageous orientation of the translocation control element with respect to the two reporter codes. As discussed further herein and with reference to Figs 6 – 8, the symmetrical chemical brancher can be represented by the letter “Y”, in which the two reporter codes are joined to the arms of the Y, while the translocation control element is joined to the stem of the Y. Thus, the two reporter codes are joined in-line by the brancher, while the brancher bears the translocation control element in a perpendicular orientation with respect to the linear, in-line, SSRT.

[00101] As used throughout the present disclosure, the terms “linker A” and “linker B” refer to the regions of the SSRT that each include a polymerase enhancing region and one or more translocation deceleration features or regions, and, in certain embodiments, a spacer region that includes a polymer of, e.g., PEG6, which can be customized to modulate the length of the SSRT traversed in a nanopore.

[00102] In certain embodiments, an XNTP may be a compound having the following generalized structure:



[00103] In one embodiment, R may be H, for example, when the compounds are used to sequence a DNA template. In another embodiment, R may be OH, for example, when the compounds are used to sequence an RNA template.

[00104] In certain embodiments, nucleobase is adenine, cytosine, guanine, thymine, uracil or a nucleobase analog. As one of skill in the art will appreciate, adenine, cytosine, guanine, thymine, and uracil are naturally occurring nucleobases. As used herein, the term “nucleobase analog” refers to non-naturally occurring nucleobases that are capable of forming Watson and Crick base pair with a complementary nucleobase on an adjacent single-stranded nucleic acid template. Exemplary nucleobase analogs include, but are not limited to, 5-fluorouracil; 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-

acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, 2,6-diaminopurine, 3-nitropyrrole, 8-aza-7-deazaguanine, 8-aza-7-deazainosine, and 8-aza-7-deazaadenine.

[00105] As discussed herein, the reporter construct is a polymer having a first end and a second end, and includes, in series from the first end to the second end, the first reporter code, the symmetrical chemical brancher bearing the translocation control element, and the second reporter code. This series of features reflects the symmetrical structure of the reporter construct (and the entire SSRT, which includes the symmetrical linkers, linker A and linker B), in which the sequences of the two reporter codes are identical and joined, in-line in reverse orientation by the symmetrical chemical brancher. Synthesis of the entire SSRT, including the reporter construct, is discussed further herein with reference to Figs. 6 – 8). Briefly, synthesis proceeds in the 3' to 5' direction, initiating at the 3' end of the TCE. Addition of the symmetrical brancher to the 5' end of the TCE enables simultaneous polymerization of the first and second reporter codes off each arm of the brancher, followed by simultaneous synthesis of linker A and linker B, terminating at the 5' end of the first end and the second end of the SSRT. It has been found that the in-line redundancy provided by two identical reporter codes separated by the symmetrical brancher bearing the translocation control element offers several advantages during nanopore sequencing. For example, Xpandomers can potentially be read by the nanopore when translocated in either direction, i.e., the Xpandomer can be read either “forwards” or “backwards”. This flexibility enables the “ratcheting” method of sequencing, which is discussed further herein, and other methods, such as “flossing” that are based on an AC pattern of voltage application.

[00106] FIG. 3 shows one embodiment of a cleaved Xpandomer in the process of translocating an α -hemolysin nanopore. This biological nanopore is embedded into a lipid bilayer membrane which separates and electrically isolates two reservoirs of electrolytes. A typical electrolyte has 1 molar KCl buffered to a pH of 7.0. When a small voltage, typically

100 mV, is applied across the bilayer, the nanopore constricts the flow of ion current and is the primary resistance in the circuit. Xpandomer reporter codes are designed to give specific ion current blockage levels and sequence information can be read by measuring the sequence of ion current levels as the sequence of reporter codes translocate the nanopore.

[00107] The α -hemolysin nanopore is typically oriented so translocation occurs by entering the vestibule side and exiting the stem side. As shown in FIG. 3, the nanopore is oriented to capture the Xpandomer from the stem side first. In some circumstances, this orientation may cause fewer blockage artifacts than occur when entering vestibule first. However, according to the present invention, the α -hemolysin nanopore may be oriented in either direction. As the Xpandomer translocates, a reporter enters the stem until its translocation control element stops at the stem entrance. The reporter is held in the stem until the TCE is enabled to pass into and through the stem, whereupon translocation proceeds to the next reporter. In this embodiment, TCE passage into the stem is enabled by dissociation of a translocation control moiety from the TCE. Advantageously, the inventors have discovered that TCEs constructed from a novel class of pendant-PEG phosphoramidites provide significantly improved translocation control based on intrinsic physicochemical and steric properties and thus obviate reliance on association and dissociation of translocation control moieties that act in trans.

Novel Compounds for SSRT Feature Design

[00108] Phosphoramidite chemistry, typically used for automated oligonucleotide synthesis, provides an efficient and convenient means to synthesize polymeric SSRTs. However, the ultimate potential of SSRT feature design is significantly limited by the repertoire of phosphoramidite monomers (PPAs) available in commercial libraries. Commercial PPAs are largely based on nucleosidic core structures and therefore do not offer the range of physicochemical properties necessary for the design of a broader array of features that improve the efficiency and accuracy of nanopore reads. To address this shortcoming in the art, the inventors have designed and synthesized a large collection of new PPA monomeric compounds. Significantly, these compounds are not based on nucleosidic core structures, which are well known in the art and, as mentioned, constrain feature design.

[00109] As used herein, the abbreviation “PPA” refers to phosphoramidites that are O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidites. It is readily understood by one of skill in the art, that the term “phosphoramidite” refers to the structure of the monomeric precursor; following in-line polymerization of PPAs into an SSRT, the monomers are converted into

phosphodiester linked oligomeric products.


[00110] Other methods used to make phosphodiester backbones polymers can be used to synthesize SSRTs. Accordingly, the monomers used with these chemistries can also produce SSRTs with non-nucleosidic elements. Additional methods of assembly may involve use of automated or manual assembly strategies done in solution phase or on a solid support. H-phosphonate synthesis and phosphotriester synthesis are examples known in the art. In addition, methods using enzymatic synthesis may be adapted to synthesize SSRTs (e.g., those employed in enzymatic oligonucleotide synthesis). In some embodiments, synthesis of an SSRT may be based on a combination of any of the above synthesis methods.

[00111] What follows is a brief, non-limiting summary of certain principles used to guide PPA monomer design. 1) Phosphate spacing. Compounds were designed that maintained a C3 (3 atom) spacing, which mimics the spacing of a natural nucleotide backbone. Other suitable spacings include, in certain embodiments, from 2 to 20 atom spacing. Unexpectedly, atom spacing was found to influence the rate of nanopore translocation, allowing for fine-tuning of translocation control. 2) Hydrophilicity. Compounds were designed to optimize the hydrophilic properties of SSRT features, as desired for particular functionalities. Several monomer designs were based on PEG, due to its ability to increase water solubility, which is an important property of, e.g., reporter codes. The inventors were able to fine-tune the hydrophilicity of PPA monomers by adjusting the length of the PEG polymer, as well as by terminating the PEG polymer with methyl ether or introducing 1,2,3-triazoles into the polymer, which had the unanticipated effect of further improving water solubility. 3) Steric volume. Several alternative configurations of linear, branched, cyclic, and dendrimeric PPA structures were designed and tested to evaluate the effect of steric volume on current flow through the nanopore. 4) Chirality in the backbone. The nanopore is a chiral environment. Enantiomeric compounds were designed to determine whether key nanopore signal properties were affected in any way. 5) Charge. In addition to back phosphate charge, certain compounds carry either a positive charge, e.g., tertiary amines or a negative charge, e.g., carboxylic acid. 6) Aromaticity. Compounds composed of a wide variety of aromatic hydrocarbons and heteroaromatic structures were incorporated into the backbone to determine if interactions with the nanopore produced desirable signal properties.


[00112] PPA monomeric compounds, in addition to attenuating nanopore signal properties, such as translocation rate control or current level control, also influence physicochemical properties of the Xpandomer. The Xpandomer, as a semi-synthetic

polymer, exhibits properties associated with both natural polymers, e.g., DNA, and synthetic polymers. In some embodiments, certain PPA monomers may enable attenuation of undesirable inter-Xpandomer interactions or interaction between the Xpandomer and certain process elements of the SBX work-flow. For example, it may be possible to reduce Xpandomer self-aggregation, formation of higher order glasses or gels, isolation, passive adsorption to, or interaction with, surfaces of containers, walls of nanochannels or fabrication devices containing the membrane and nanopore.

[00113] One class of compounds that has proven to provide outstanding functionality when incorporated into SSRT features is referred to herein as “pendant PEG”. These structures are based on a molecular core that enables linkage of one or more PEG-containing polymers in a pendant configuration relative to the core. A structural analogy can be drawn between polymers of pendant PEG compounds and a comb, in which the phosphodiester bonds between individual compounds form the base of the comb and the PEG-based polymers form the teeth. Advantageously, several properties of the pendant PEG “teeth” can be customized for particular SSRT features, e.g., one or more of the spacing, length, and composition of the polymeric teeth. Structures 1a, 2a, 3a, and 4a below illustrate four exemplary embodiments of pendant PEG core structures.



In certain non-limiting embodiments, X or X' may represent $-\text{CH}_2\text{O}-[\text{CH}_2\text{CH}_2\text{O}]_m\text{O}-$ in











[00114] Tables 1A-C set forth non-limiting collections of novel phosphoramidite monomeric compounds for use in, e.g., SSRT feature design. Synthetic schemes for each compound are referred to with reference to the relevant Example and specific precursors are

included for each. Analytic data characterizing the purified synthesized compounds are also set forth in Table 1A. These compounds may be used to synthesize any suitable polymeric feature, e.g., SSRT reporter codes, translocation control elements and translocation deceleration features, as described in further detail herein. Table 1A also provides the names of the compounds with reference to the in-line structures they assume following incorporation into synthetic polymers.



Table 1A[illegible]



27

 A. 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane B. 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane C. code: PPA032	<u>12b</u> Ex. 2		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.04 - 1.21 (m, 12 H, N(iPr)₂), 2.57 (m, 2 H, -CH₂-CN) 3.07 - 3.22 (m, 2 H, NH(iPr)) 3.30 (s, 3 H, -OCH₃ (PEG)) 3.44 - 3.76 (m, 23 H, -CH₂O (PEG, CNEt and glycerol)) 3.79 (s, 6 H, Ph-OCH₃) 6.89 (d, J=8.71 Hz, 4 H, Ph-H) 7.20 - 7.53 (m, 9 H, Ph-H) 147.95
 A. 1-O-DMT-3-O-PPA-2S-O-mPEG6-propane B. 1,3-O-bis(phosphodiester)-2S-O-mPEG6-propane C. code: PPA031	<u>12c</u> Ex. 2		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.06 - 1.23 (m, 12 H, N(iPr)₂) 2.52 - 2.65 (m, 2 H, -CH₂-CN) 3.07 - 3.22 (m, 2 H, NH(iPr)₂) 3.31 (s, 3 H, -OCH₃ (PEG)) 3.47 - 3.74 (m, 31 H, -CH₂O (PEG, glycerol, CNEt)) 3.79 (s, 6 H, Ph-OCH₃) 6.80 - 6.94 (m, 4 H, Ph-H) 7.22 - 7.39 (m, 7 H, Ph-H) 7.42 - 7.52 (m, 2 H, Ph-H) 147.98





 DMTrO-PPA A. 1-O-DMT-2-O-PPA-3-O-mPEG2-propane B. 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane C. code: PPA029	<u>16</u> Ex. 3		1H NMR (300 MHz, ACN-d3) δ ppm 1.06 (m, 12 H, N(iPr)2) 2.55 - 2.76 (m, 2 H, -CH2-CN) 3.15 - 3.23 (m, 2 H, N(iPr)2) 3.29 (s, 3 H, -OCH3 (PEG)) 3.41 - 3.82 (m, 14 H, -CH2O (glycerol, PEG, CNEt)) 3.79 (s, 6 H, Ph-OCH3) 4.01 - 4.16 (m, 1 H, CHO (glycerol)) 6.83 - 6.99 (m, 4 H, Ph-H) 7.23 - 7.39 (m, 7 H, Ph-H) 7.45 - 7.54 (m, 2 H, Ph-H)	148.77, 149.26
 DMT-PPA A. 3-O-DMT-2S-O-PPA-1-(N3-Thymine)-propane B. 2,3-O-bis(phosphodiester)-1-(N3-Thymine)-propane C. code: PPA002	<u>20a</u> Ex. 4		Expected 1H NMR δ ppm 1.06 (d, 12 H, N(iPr)2) 2.34 (s, 3 H, C5-CH3) 2.60 (m, 2 H, -CH2CN) 2.83 (m, 2 H, N(iPr)2) 3.17 - 3.63 (m, 4 H, -CH2O (glycerol), -CH2N) 3.81 (s, 6 H, Ph-OCH3) 3.90 (m, 2 H, CNEt) 4.42 (m, 1 H, CHO) 6.89 (m, 4 H, Ph-H) 7.19 (s, 1 H, C6-H) 7.26 - 7.28 (m, 9 H, Ph-H)	149.79, 149.64



A. 3-O-DMT-2S-O-PPA-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane B. 2,3-O-bis(phosphodiester)-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane C. code: PPA006	<u>20b</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.02 - 1.17 (m, 12 H, N(iPr)₂) 2.26 (d, J=3.84 Hz, 3 H, indole-CH₃) 2.43 - 2.54 (m, 2 H, -CH₂CN) 2.89 - 3.14 (m, 2 H, N(iPr)₂) 3.02 (dd, J=8.45, 4.86 Hz, 1 H, CH) 3.52 - 3.71 (m, 4 H, -CH₂O (DMT, CNEt)) 3.75 (s, 3 H, indole-OCH₃) 3.76 (s, 3 H, indole-OCH₃) 3.82 (s, 3 H, Ph-OCH₃) 4.11 - 4.36 (m, 2 H, -CH₂-indole) 6.11 (d, J=1.54 Hz, 1 H, indole-H) 6.40 (d, J=1.54 Hz, 1 H, indole-H) 6.57 (d, J=9.47 Hz, 1 H, indole-H) 6.82 (td, J=5.76, 2.56 Hz, 4 H, Ph-H) 7.19 - 7.33 (m, 7 H, Ph-H) 7.43 - 7.49 (m, 2 H, Ph-H) 149.91, 148.47
---	--------------------------------	---	--




 A. 3-O-DMT-2S-O-PPA-1-(1-dimethoxyquinazolin-2-ylidene)-propane B. 2,3-O-bis(phosphodiester)-1-(1-dimethoxyquinazolin-2-ylidene)-propane C. code: PPA007	<u>20c</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.88 - 1.17 (m, 12 H, N(iPr)₂) 2.43 - 2.53 (m, 2 H, -CH₂CN) 3.25 - 3.39 (m, 2 H, N(iPr)₂) 3.43 - 3.79 (m, 10 H, -CH₂O (DMT, CNEt), Ph-OCH₃) 3.85 - 3.94 (m, 3 H, Ph-OCH₃) 3.99 - 4.16 (m, 1 H, -CH) 4.30 - 4.47 (m, 2 H, -CH₂N) 6.75 - 6.92 (m, 4 H, Ph-H) 7.21 - 7.38 (m, 7 H, Ph-H) 7.40 - 7.54 (m, 4 H, Ph-H)	149.19, 148.83
---	-----------------------------	--	---	---------------------------





 A. 3-(1-O-DMT-2S-O-PPA-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide B. 3-(1,2-O-bis(phosphodiester)-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide C. code: PPA009	<u>20d</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.80 - 0.97 (m, 6 H, camphor-CH₃) 1.14 - 1.29 (m, 16 H, N(iPr)₂, camphor) 1.58 - 2.01 (m, 6 H, camphor) 2.53 (t, J=6.02 Hz, 2 H, -CH₂N) 2.67 - 2.82 (m, 2 H, -CH₂CN) 2.89 - 3.21 (m, 4 H, CH₂O (DMT, CNEt)) 3.25 - 3.42 (m, 2 H, -CH₂S) 3.72 - 3.82 (m, 6 H, Ph-OCH₃) 4.15 - 4.31 (m, 1 H, CH) 6.78 - 6.95 (m, 4 H, Ph-H) 7.23 - 7.40 (m, 7 H, Ph-H) 7.44 - 7.58 (m, 2 H, Ph-H)	148.79, 148.37
---	--------------------------------	---	--	---------------------------



DMTrO O PPA A. 3-O-DMT-2S-O-PPA-1-(N9-(3,6-dimethoxycarbazole)-propane B. 2,3-O-bis(phosphodiester)-1-(N9-(3,6-dimethoxycarbazole)-propane C. code: PPA010	20e Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.84 - 1.04 (m, 12 H, N(iPr)₂) 2.30 - 2.41 (m, 2 H, -CH₂CN) 3.00 - 3.23 (m, 2 H, N(iPr)₂) 3.14 (d, J=4.15 Hz, 1 H, CH) 3.30 - 3.62 (m, 4 H, CH₂O (glycerol, CNEt)) 3.75 (d, J=3.32 Hz, 6 H, carbazole-OCH₃) 3.86 (d, J=1.45 Hz, 6 H, Ph-OCH₃) 4.34 - 4.52 (m, 2 H, CH₂-carbazole) 6.80 (ddd, J=8.81, 5.70, 3.11 Hz, 4 H, Ph-H) 6.99 (dt, J=8.76, 3.08 Hz, 2 H, carbazole) 7.19 - 7.36 (m, 9 H, Ph-H, carbazole) 7.46 (d, J=7.67 Hz, 2 H, Ph-H) 7.57 (s, 2 H, carbazole) 148.67, 148.98
--	-------------------------	---	--





 A. 3-O-DMT-2S-O-PPA-1-(N1-(6-Azathymine))-propane B. 2,3-O-bis(phosphodiester)-1-(N1-(6-Azathymine))-propane C. code: PPA012	<u>20f</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.03 - 1.19 (m, 12 H) 2.03 - 2.20 (m, 2 H, glycerol) 2.47 - 2.65 (m, 2 H, CH₂CN) 3.18 - 3.81 (m, 5 H, CH₂O (glycerol, CNEt)) 3.78 (s, 6 H, Ph-OCH₃) 4.05 - 4.17 (m, 1 H, CH₂O (glycerol)) 4.23 - 4.36 (m, 1 H, CHO (glycerol)) 6.86 - 6.87 (m, 4 H, Ph-H) 7.22 - 7.38 (m, 7 H, Ph-H) 7.40 - 7.56 (m, 2 H, Ph-H)	148.83, 149.06
 A. 3-O-DMT-2S-O-PPA-1-(N1-(2-Me-5-nitroindole))-propane B. 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole))-propane C. code: PPA013	<u>20g</u> Ex. 4		Expected ¹H NMR δ ppm 1.06 (d, 12 H, N(iPr)₂) 2.60 (m, 2 H, -CH₂CN) 2.83 (m, 2 H, N(iPr)₂) 3.35 - 3.63 (m, 5 H, -CH₂O, -CHO (glycerol), -CH₂N) 3.77 (m, 2 H, -CH₂N) 3.81 (s, 6 H, Ph-OCH₃) 3.90 (m, 2 H, CNEt) 4.17 (m, 2 H, CH₂O) 6.70 - 6.89 (m, 8 H, Ph-H) 7.26 - 7.28 (m, 9 H, Ph-H)	148.93, 148.69





 A. 3-O-DMT-2S-O-PPA-1-(N1-(2-Me-5-nitroindole))-propane B. 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole))-propane C. code: PPA015	<u>20h</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.94 - 1.35 (m, 12 H, N(iPr)₂) 2.39 (ddd, J=17.03, 6.02, 5.89 Hz, 2 H, -CH₂CN) 2.50 (d, J=6.14 Hz, 3 H, indole-CH₃) 3.30 - 3.38 (m, 2 H, N(iPr)₂) 3.44 - 3.62 (m, 4 H, CH₂O (glycerol, CNEt)) 3.79 (d, J=4.61 Hz, 6 H, Ph-OCH₃) 4.25 - 4.47 (m, 3 H, -CH₂-indole, -CH) 6.39 - 6.52 (m, 1 H, indole) 6.80 - 6.96 (m, 4 H, Ph-H) 7.30 - 7.43 (m, 7 H, Ph-H) 7.48 - 7.56 (m, 2 H, Ph-H) 7.97 (dt, J=9.15, 1.70 Hz, 1 H, indole) 8.41 (t, J=2.56 Hz, 1 H, indole)	149.01, 149.28
---	------------------------------	--	--	----------------------------





DMT-O- A. 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane B. 2,3-O-PPA O-bis(phosphodiester)-1-(5-benzofuran)-propane C. code: PPA018	<u>20i</u> Ex. 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.09 (dd, J=19.97, 6.91 Hz, 12 H, N(iPr)₂) 2.51 (dt, J=17.92, 6.02 Hz, 2 H, -CH₂CN) 3.10 - 3.73 (m, 6 H, -CH₂ (DMT, CNEt, -NAr)) 3.76 (s, 6 H, Ph-OCH₃) 3.98 - 4.15 (m, 1 H, CHO-PPA) 4.30 - 4.46 (m, 2 H, pyr) 6.50 - 6.86 (m, 1 H, furan) 6.86 - 6.96 (m, 4 H, Ph-H) 7.24 - 7.40 (m, 7 H, Ph-H) 7.44 - 7.57 (m, 2 H, Ph-H) 7.57 - 7.67 (m, 1 H, furan)	149.08
DMTrO- A. 1-O-DMT-2S-O-PPA-3-(4-methylpiperazine-1-yl)-propane B. 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane C. code: PPA043	<u>20j</u> Ex 4		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.98 - 1.26 (m, 12 H, N(iPr)₂) 2.17 (d, J=4.10 Hz, 3 H, NCH₃) 2.22 - 2.47 (m, 6 H, CH₂N) 2.50 - 2.60 (m, 4 H, CH₂N) 2.64 - 2.71 (m, 2 H, CH₂CN) 3.09 - 3.23 (m, 2 H, N(iPr)₂) 3.60 - 3.82 (m, 10 H, HC₂O, Ph-OCH₃) 3.98 - 4.14 (m, 1 H, CH) 6.79 - 6.95 (m, 4 H, Ph-H) 7.23 - 7.40 (m, 7 H, Ph-H) 7.43 - 7.57 (m, 2 H, Ph-H)	148.49, 148.02



 A. 1-O-DMT-2S-O-PPA-3-O-(2-Me-(10-crown-5))-propane B. 1,2-O-bis(phosphodiester)-3-O-(2-Me-(10-crown-5))-propane C. code: PPA021	<u>20k</u> Ex. 4		¹H NMR (300 MHz, CDCl₃) δ ppm 0.82 - 0.94 (m, 1 H) 1.05 (d, J=6.66 Hz, 2 H) 1.13 - 1.30 (m, 15 H, N(iPr)₂) 2.65 (t, J=6.53 Hz, 2 H, CH₂CN) 3.60 - 3.76 (m, 24 H -CH₂O, Ph-OCH₃ (glycerol, crown-ether)) 3.79 - 3.85 (m, 4 H -CH₂O (CNEt, glycerol)) 6.73 - 6.88 (m, 1 H Ph-H) 7.20 - 7.36 (m, 2 H, Ph-H)	148.59
 A. 1-O-DMT-5-O-PPA-hexahydrofuro[2,6]furan B. 1,5-O-bis(phosphodiester)-hexahydrofuro[2,6]furan C. code: PPA014	<u>23</u> Ex. 5		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.09 - 1.25 (m, 12 H) 2.57 - 2.71 (m, 2 H, -CH₂CN) 2.98 - 3.14 (m, 2 H, N(iPr)₂) 3.53 - 3.67 (m, 2 H, CNEt) 3.70 - 4.08 (m, 6 H, isosorbide) 3.79 (s, 6 H, Ph-OCH₃) 4.16 - 4.32 (m, 2 H, -CHO (isosorbide)) 6.77 - 6.89 (m, 4H, Ph-H) 7.27 (d, J=7.17 Hz, 1 H, Ph-H) 7.31 - 7.44 (m, 6 H, Ph-H) 7.47 - 7.62 (m, 2 H, Ph-H)	148.30, 147.73


 A. 1-O-DMT-1'-O-PPA-(2,2'-bipyridin-4,4'-yl)-dimethanol B. 1,1'-O-bis(phosphodiester)-2,2'-bipyridin-4,4'-yl)-dimethanol C. code: PPA005	<u>26a</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.24 (d, J=7.17 Hz, 12 H, N(iPr)₂) 2.71 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.68 - 3.81 (m, 2 H, N(iPr)₂) 3.79 (s, 6 H, Ph-OCH₃) 3.84 - 3.93 (m, 2 H, CNEt) 4.34 (s, 2 H, CH₂O-DMT) 4.79 - 4.95 (m, 2 H, CH₂O-PPA) 6.91 (d, J=8.96 Hz, 4 H, Ph-H) 7.29 (d, J=7.42 Hz, 1 H, Ph-H) 7.33 - 7.45 (m, 8 H, Ph-H) 7.48 - 7.63 (m, 2 H, pyr) 8.37 (s, 1 H, pyr) 8.45 (s, 1 H, pyr) 8.55 - 8.67 (m, 2 H, pyr)	148.96
---	-----------------------------	--	---	---------------



 A. 1-O-DMT-1'-O-PPA-N(p-tolyl)-diethanolamine B. 1,1'-O-bis(phosphodiester)-N(p-tolyl)-diethanolamine C. code: PPA001	<u>26b</u> Ex. 6		¹H NMR (300 MHz, CDCl₃-d) δ ppm 1.19 (dd, J=10.88, 6.78 Hz, 12 H, N(iPr)₂) 2.24 (s, 3 H, Ph-CH₃) 2.57 (t, J=6.53 Hz, 2 H, -CH₂CN) 3.22 - 3.35 (m, 2 H, N(iPr)₂) 3.52 - 3.68 (m, 6 H, -CH₂O) (DMT, PPA, CNEt) 3.70 - 3.83 (m, 4 H, -CH₂N) 3.80 (s, 6 H, Ph-OCH₃) 6.57 (m, J=8.70 Hz, 2 H, toluyl-H) 6.80 (d, J=8.96 Hz, 4 H, Ph-H) 6.99 (m, J=8.45 Hz, 2 H, toluyl-H) 7.18 - 7.35 (m, 7 H, Ph-H) 7.41 - 7.48 (m, 2 H, Ph-H)	148.04
 A. 1-O-DMT-1'-O-PPA-(adamantan-1,3-yl)-dimethanol B. 1,1'-O-bis(phosphodiester)-adamantan-1,3-yl)-dimethanol C. code: PPA003	<u>26c</u> Ex. 6		Expected ¹H NMR δ ppm 0.82 - 1.66 (m, 26 H, adamantane, N(iPr)₂) 2.60 (m, 2 H, -CH₂CN) 3.27 - 3.52 (m, 4 H, CH₂O) 3.81 (s, 6 H, Ph-OCH₃) 3.90 (m, 2 H, CNEt) 6.89 (m, 4 H, Ph-H) 7.26 - 7.28 (m, 9 H, Ph-H)	Expected 148


 DMTrO A. 1-O-DMT-1'-O-PPA-2,2'-(sulfonylbis(benz-4-yl))-diethanol B. 1,1'-O-bis(phosphodiester)-2,2'-(sulfonylbis(benz-4-yl))-diethanol C. code: PPA004	<u>26d</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.16 (dd, J=10.37, 6.84 Hz, 12 H, N(iPr)₂) 2.62 (t, J=6.01 Hz, 2 H, -CH₂CN) 3.32 - 3.40 (m, 2 H, N(iPr)₂) 3.62 (m, 2 H, -CH₂O (CNEt)) 3.78 (s, 6 H, Ph-OCH₃), 3.70 - 3.99 (m, 4 H, -CH₂O (DMT, PPA)) 4.17 - 4.30 (m, 4 H, -CH₂O-sultone) 6.81 - 6.93 (m, 4 H, Ph-H) 7.04 - 7.18 (m, 4 H, sultone) 7.19 - 7.35 (m, 7 H, Ph-H) 7.37 - 7.48 (m, 2 H, Ph-H) 7.82 - 7.88 (m, 4 H, sultone) 148.83
 DMTrO A. 1-O-DMT-1'-O-PPA-(octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol B. 1,1'-O-bis(phosphodiester)-octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol C. code: PPA008	<u>26e</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.10 - 1.27 (m, 12 H, N(iPr)₂) 1.31 - 2.76 (m, 10 H, -CH₂, -CH₂, -CH₂CN) 2.95 - 3.88 (m, 10 H, N-CH₃, CNEt, N(iPr)₂) 3.79 (s, 6 H, Ph-OCH₃) 6.89 (d, J=8.96 Hz, 4 H, Ph-H) 7.21 - 7.38 (m, 7 H, Ph-H) 7.40 - 7.54 (m, 2 H, Ph-H) 147.76



 A. 3,5-bis(DMT-O-ethoxy)- 1-(PPA-O-ethoxy)-1,3,5-triazene-2,4,6-trione B. 3,5-bis(phosphodiester-O-ethoxy)- 1-(phosphodiester-O-ethoxy)-1,3,5-triazene-2,4,6-trione C. code: PPA038	<u>26f</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.88 - 1.14 (m, 12 H, N(iPr)₂) 2.56 (t, J=6.14 Hz, 2 H, -CH₂CN) 3.25 - 4.02 (m, 14 H, Et and N(iPr)₂) 6.73 - 6.81 (m, 8 H, Ph-H) 7.14 - 7.27 (m, 14 H, Ph-H) 7.32 - 7.38 (m, 4 H, Ph-H)	148.13
DMTrO  A. 1-O-DMT-1'-O-PPA-N-methyl-N,N-diethanolamine B. 1,1'-O-bis(phosphodiester)-N-methyl-N,N-diethanolamine C. code: PPA041	<u>26g</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.07 - 1.21 (m, 12 H, N(iPr)₂) 2.21 (s, 3 H, -NCH₃) 2.52 - 2.67 (m, 6 H, -CH₂CN, -CH₂N) 3.04 - 3.11 (m, 2 H, N(iPr)₂) 3.52 - 3.77 (m, 6 H, -CH₂O (CNEt, DMT, PPA)) 3.75 (s, 6 H, Ph-OCH₃) 6.82 - 6.88 (m, 4 H, Ph-H) 7.19 - 7.34 (m, 7 H, Ph-H) 7.41 - 7.46 (m, 2 H, Ph-H)	147.49



DMTrO  A. 1-O-DMT-8-O-PPA-N,N-Diethylpiperazine B. 1,8-O-bis(phosphodiester)-N,N-Diethylpiperazine C. code: PPA044	<u>26h</u> Ex. 6		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.19 (d, J=6.91 Hz, 12 H), 2.41 - 2.57 (m, 12 H, CH₂N) 2.64 - 2.71 (m, 2 H, CH₂CN) 3.10 (t, J=5.76 Hz, 2 H, N(iPr)₂) 3.60 - 3.69 (m, 2 H, CH₂O (CNEt)) 3.71 - 3.84 (m, 4 H, CH₂O (DMT, PPA)) 3.79 (s, 6 H, Ph-OCH₃) 6.81 - 6.98 (m, 4 H, Ph-H) 7.21 - 7.38 (m, 7 H, Ph-H) 7.42 - 7.57 (m, 2 H, Ph-H)	147.79
---	--------------------------------	--	--	---------------


 A. 1-O-DMT-3-O-PPA-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane C. code: PPA026	<u>31a</u> Ex. 7		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.93 - 1.17 (m, 12 H, N(iPr)₂) 2.13 (m, 1 H, CH) 2.57 (q, J=5.63 Hz, 2 H, -CH₂CH₂CN) 2.69 - 2.83 (m, 2 H, -CH₂-triazole) 3.09 (d, J=5.63 Hz, 2 H, N(iPr)₂) 3.24 (s, 3 H, OCH₂ (PEG)) 3.35 - 3.70 (m, 10, CH₂O (PEG, glycerol)) 3.79 (s, 6 H, Ph-OCH₃) 4.37 - 4.42 (m, 2 H, -CH₂-triazole) 6.82 - 6.87 (m, 4 H, Ph-H) 7.19 - 7.32 (m, 7 H, Ph-H) 7.38 - 7.44 (m, 3 H, Ph-H and triazole) 147.16
--	-----------------------------	--	--



 A. 1-O-DMT-3-O-PPA-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane C. code: PPA027	<u>31b</u> Ex. 7		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.90 - 1.17 (m, 12 H, N(iPr)₂) 2.57 (q, J=5.72 Hz, 2 H, -CH₂-CN) 2.67 - 2.84 (m, 2 H, NH(iPr)₂) 2.96 - 3.18 (m, 2 H, O-CH₂ (glycerol)) 3.27 (s, 3 H, -OCH₃ (PEG)) 3.41 - 3.61 (m, 14 H, -CH₂O (PEG, glycerol)) 3.64 - 3.80 (m, 12 H, Ph-OCH₃, -CH₂O (glycerol, PEG)) 4.40 (t, J=4.74 Hz, 2 H, CH₂-triazole (glycerol)) 6.81 - 6.88 (m, 4 H, Ph-H) 7.18 - 7.33 (m, 7 H, Ph-H) 7.37 - 7.53 (m, 3 H, Ph-H, C-CH (triazole))	147.03
--	-----------------------------	--	---	---------------




 A. 1-O-DMT-2-O-PPA-3-(4-(Me-O-PEG1-O-Bz)-1-(1,2,3-triazole))-propane B. 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG1-O-Bz)-1-(1,2,3-triazole))-propane C. code: PPA054	<u>31c</u> Ex. 7		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.05 - 1.11 (m, 12 H, N(iPr)₂) 2.61 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.10 - 3.25 (m, 2 H, N(iPr)₂) 3.49 - 3.87 (m, 12 H, -CH₂O (DMT, CNEt, PEG), Ph-OCH₃) 4.40 - 4.47 (m, 1 H, CH) 4.52 - 4.67 (m, 4 H, -CH₂OBz, -CH₂-triazole) 6.82 - 6.95 (m, 4 H, Ph-H) 7.23 - 7.39 (m, 9 H, Ph-H and OBz) 7.42 - 7.57 (m, 3 H, Ph-H and triazole) 7.59 - 8.06 (m, 3 H, OBz)	149.08, 149.67
--	--------------------------------	--	---	---------------------------



 A. 1-O-DMT-2-O-PPA-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane B. 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane C. code: PPA060	<u>31d</u> Ex. 7		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.08 (d, J=6.66 Hz, 3 H, N(iPr)₂) 1.02 (d, J=6.91 Hz, 4 H, N(iPr)₂) 1.13 (d, J=6.66 Hz, 7 H, N(iPr)₂) 1.91 - 2.02 (m, 2 H, CH₃CN) 2.48 (t, J=6.02 Hz, 1 H, CH₂-CN) 2.62 (t, J=5.89 Hz, 1 H, CH₂-CN) 3.11 - 3.25 (m, 2 H, NH(iPr)₂) 3.52 - 3.67 (m, 7 H, CH₂O (PEG, glycerol)) 3.57 (s, 5 H, CH₂-O, (PEG)) 3.74 - 3.80 (m, 8 H, O-CH₂ (CNEt, Ph-OCH₃) 3.81 (br. s., 1 H) 4.30 - 4.45 (m, 3 H, CH-CH₂-triazole (glycerol)) 4.49 - 4.63 (m, 4 H, triazole-CH₂-PEG, PEG-CH₂-Bz) 6.81 - 6.96 (m, 4 H, Ph-H) 7.23 - 7.39 (m, 8 H, Ph-H) 7.42 - 7.58 (m, 3 H, Ph-H) 7.58 - 7.71 (m, 1 H, Ph-H) 7.98 - 8.07 (m, 1 H, C-CH (triazole)) 149.06, 149.76
--	-----------------------------	--	---



 A. 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG2))-1-(Et-OBz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2))-1-(Et-OBz)-1,2,3-triazole)-propane C. code: PPA062	<u>35a</u> Ex. 8		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.06 - 1.19 (m, 12 H, N(iPr)₂) 2.51 - 2.64 (m, 2 H, -CH₂CN) 3.05 - 3.22 (m, 2 H, N(iPr)₂) 3.50 - 3.73 (m, 13 H, -CH₂O (PEG)) 3.78 (s, 6 H, Ph-OCH₃) 4.56 (s, 2 H, -OCH₂-triazole) 4.61 - 4.77 (m, 4 H, Et-Bz) 6.88 (d, J=8.96 Hz, 4 H, Ph-H) 7.22 - 7.38 (m, 7 H, Ph-H) 7.48 (d, J=7.68 Hz, 3 H, Ph-H) 7.88 - 8.01 (m, 3 H, Ph-H, triazole)	147.98
--	-----------------------------	--	---	---------------




 A. 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole-propane B. 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole-propane C. code: PPA061/081	<u>35b</u> Ex. 8		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.08 - 1.24 (m, 12 H, N(iPr)₂) 1.99 (s, 3 H, OAc) 2.55 - 2.70 (m, 2 H, -CH₂CN) 3.08 - 3.21 (m, 2 H, N(iPr)₂) 3.54 - 3.74 (m, 19 H, CH₂O (glycerol, PEG, CNEt) 3.79 (s, 6 H, Ph-OCH₃) 4.43 (d, J=5.60 Hz, 2 H, -OCH₂-triazole) 4.55 - 4.71 (m, 4 H, EtOAc) 6.85 - 6.99 (m, 4 H, Ph-H) 7.29 - 7.42 (m, 7 H, Ph-H) 7.45 - 7.60 (m, 3 H, Ph-H and triazole)	147.95
--	--------------------------------	--	---	---------------



 A. 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5))-1-(Et-O-Ac)-1,2,3-triazole-propane B. 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5))-1-(Et-O-Ac)-1,2,3-triazole)-propane C. code: PPA083/PPA063	35c Ex. 8		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.05 - 1.20 (m, 12 H, N(iPr)₂) 1.90 - 2.06 (m, 1 H) 1.90 - 2.06 (m, 5 H, OAc) 2.60 (dt, J=8.45, 5.89 Hz, 2 H, CH₂-CN) 3.04 - 3.21 (m, 2 H, NH(iPr)₂) 3.51 - 3.63 (m, 21 H, -OCH₂ (PEG, -CH (glycerol))) 3.65 - 3.74 (m, 5 H, -OCH₃ (Glycerol, CNEt)) 3.78 (s, 6 H, Ph-OCH₃) 4.33 - 4.47 (m, 2 H, PEG-OCH₂-Triazole) 4.51 - 4.66 (m, 4 H, triazole-CH₂-CH₂-Ac) 6.88 (d, J=8.70 Hz, 4 H, Ph-H) 7.22 - 7.38 (m, 7 H, Ph-H) 7.47 (s, 1 H, Ph-H) 7.48 - 7.54 (m, 1 H, C=CH (Triazole))	147.95
---	-------------------------	--	--	---------------



 A. 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane C. code: PPA064	<u>35d</u> Ex. 8		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.08 - 1.19 (m, 12 H, N(iPr)₂) 2.51 - 2.65 (m, 2 H, CH₂-CN) 3.06 - 3.22 (m, 2 H, NH(iPr)₂) 3.49 - 3.72 (m, 36 H, -CH₂O (PEG, glycerol)) 3.79 (s, 7 H, Ph-OCH₃) 4.58 (s, 2 H, PEG-CH₂-triazole) 4.66 - 4.81 (m, 4 H, triazole-CH₂-CH₂-Bz) 6.89 (d, J=8.70 Hz, 4 H, Ph-H) 7.19 - 7.69 (m, 12 H, Ph-H) 7.83 - 8.01 (m, 2 H, Ph-H, C=CH (triazole))	148
 A. 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane C. code: PPA065	<u>35e</u> Ex. 8		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.07 - 1.20 (m, 12 H, N(iPr)₂) 2.52 - 2.67 (m, 2 H, CH₂CN) 3.07 - 3.23 (m, 2 H, N(iPr)₂) 3.54 - 3.80 (m, 22 H, -CH₂O (PEG, Glycerol, CNEt), -OCH₃) 4.60 (s, 2 H, O-CH₂-Triazole) 5.21 (s, 2 H, CH₂-(N)Triazole)) 6.89 (d, J=8.96 Hz, 4 H, Ph-H) 7.23 - 7.55 (m, 9 H, Ph-H)	147.93



 A. 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane C. code: PPA068	37a Ex. 9		¹H NMR (300 MHz, DMSO-d₆) δ ppm 0.98 - 1.13 (m, 12 H, N(iPr)₂) 2.71 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.26 (br. s., 2H, NH(iPr)₂) 3.49 (br. s., 2 H, -CH₂O (glycerol)) 3.52 (d, J=8.70 Hz, 5 H -CH₂O (glycerol, PEG)) 3.62 (s, 7 H, Ph-OCH₃) 3.67 - 3.83 (m, 4 H, -CH₂O (glycerol, PEG)) 4.23 (s, 3 H, PEG-CH₂-triazole) 4.29 - 4.42 (m, 11 H, CH₂-Bz) 4.52 (br. s., 2 H) 4.60 (s, 2 H, triazole-CH₂-C) 6.71 (d, J=8.96 Hz, 4 H, Ph-H) 7.11 - 7.25 (m, 7 H, Ph-H) 7.30 - 7.36 (m, 2 H, Ph-H) 7.39 - 7.54 (m, 13 H, Ph-H) 7.60 - 7.67 (m, 4 H, Ph-H) 7.78 - 7.86 (m, 7 H, Ph-H) 7.90 (s, 1 H, C-CH, triazole) 7.91 (d, J=6.91 Hz, 3 H, Ph-H)	147.93
---	-------------------------	--	---	---------------


 A. 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane C. code: PPA093	<u>37b</u> Ex. 9		¹H NMR (300 MHz, ACN-d₃) δ ppm 0.99 - 1.14 (m, 12 H, N(iPr)₂) 2.03 (s, 9 H, -OAc) 2.45 - 2.61 (m, 2 H, -CH₂CN) 3.00 - 3.17 (m, 2 H, N(iPr)₂) 3.41 - 3.70 (m, 27 H, -CH₂O (PEG, CNEt), -CH (glycerol), -NCH₂ (Erythritol)) 3.73 (s, 6 H, Ph-OCH₃) 4.00 (s, 6 H, -OCH₂ (Erythritol)) 4.51 (d, J=8.71 Hz, 4 H, P-OCH₂ (Glycerol), -OCH₂-Triazole) 6.78 - 6.89 (m, 4 H, Ph-H) 7.12 - 7.47 (m, 9 H, Ph-H) 7.70 (s, 1 H, -CH (Triazole)) 147.96
---	--------------------------------	--	--



 A. 1-O-DMT-3-O-PPA-2S-O-(PEG4-O-Bz)-propane B. 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz))-propane C. code: PPA050	<u>38b</u> Ex. 10		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.05 - 1.22 (m, 12 H, N(iPr)₂ 2.52 - 2.68 (m, 2 H, -CH₂-CN) 3.06 - 3.22 (m, 2 H, N(iPr)₂ 3.53 - 3.77 (m, 21 H, -CH₂O (glycerol, PEG, CNEt)) 3.79 (s, 6 H, Ph-OCH₃) 4.42 (dd, J=5.63, 3.84 Hz, 2 H, -CH₂-OBz) 6.88 (d, J=8.70 Hz, 4 H, Ph-H) 7.23 - 7.38 (m, 10 H, Ph-H) 7.42 - 7.57 (m, 3 H, Ph-H) 7.99 - 8.07 (m, 1 H, Ph-H)	147.98
 A. 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG2)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2))-propane C. code: PPA040	<u>45b</u> Ex. 11		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.07 - 1.23 (m, 12 H, N(iPr)₂ 2.59 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.10 (m, 2 H N(iPr)₂ 3.30 (s, 6 H, -OCH₃ (PEG)) 3.43 - 3.59 (m, 20 H, -CH₂O (PEG)) 3.65 - 3.81 (m, 4 H, -CH₂O (DMT, PPA)) 3.79 (s, 6 H, Ph-OCH₃) 6.88 (d, J=8.96 Hz, 4 H, Ph-H) 7.21 - 7.37 (m, 7 H, Ph-H) 7.40 - 7.55 (m, 2 H, Ph-H)	147.1



 A. 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG4)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG4)-propane C. code: PPA035	<u>45c</u> Ex. 11		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.06 - 1.20 (m, 12 H, N(iPr)₂) 2.57 (t, J=5.89 Hz, 2 H, -CH₂CN) 3.07 (s, 2 H, N(iPr)₂) 3.26 - 3.30 (m, 6 H, OCH₃ (PEG)) 3.44 - 3.59 (m, 36 H, -CH₂O (PEG, Erythritol)) 3.77 (s, 6 H, Ph-OCH₃) 6.85 (d, J=8.96 Hz, 4 H, Ph-H) 7.22-7.33 (m, 7 H, Ph-H) 7.42 - 7.46 (m, 2 H, Ph-H)	147.1
---	-------------------------------	--	---	--------------





 A. 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG1-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG1-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane C. code: PPA070	46e Ex. 11		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.02 - 1.19 (m, 12 H, N(iPr)₂) 2.56 (t, J=5.89 Hz, 2 H, -CH₂CN) 3.07 (s, 2 H, N(iPr)₂) 3.42 - 3.59 (m, 16 H, -CH₂O (PEG)) 3.61 - 3.78 (m, 6 H, -CH₂O (DMT, CNEt, PPA)) 3.76 (s, 6 H, Ph-OCH₃) 4.46 - 4.59 (m, 4 H, -OCH₂-triazole) 4.60 - 4.76 (m, 8 H, Et-OBz) 6.84 (d, J=8.70 Hz, 4 H, Ph-H) 7.17 - 7.34 (m, 7 H, Ph-H) 7.37 - 7.49 (m, 4 H, Ph-H) 7.54 - 7.95 (m, 10 H, Ph-H and triazole)	147.03
---	--------------------------	--	--	---------------

 A. 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane C. code: PPA069	47f Ex. 11		1H NMR (300 MHz, ACN-d₃) δ ppm 1.12 (dd, J=18.31, 6.78 Hz, 12 H, N(iPr)₂) 2.57 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.07 (s, 2 H, N(iPr)₂) 3.43 - 3.59 (m, 24 H, -CH₂O (PEG, Erythritol) 3.62 - 3.73 (m, 2 H, CH₂O (CNEt)) 3.77 (s, 6 H, Ph-OCH₃) 4.55 (s, 4 H, -OCH₂-triazole) 4.63 - 4.80 (m, 8 H, Et-OBz) 6.86 (d, J=8.96 Hz, 4 H, Ph-H) 7.19 - 7.35 (m, 7 H, Ph-H) 7.50 (d, J=7.94 Hz, 2 H, Ph-H) 7.45 - 7.70 (m, 6 H, Ph-H) 7.83 (s, 2 H, triazole) 7.96 (dd, J=8.32, 1.15 Hz, 4 H, Ph-H) 147.12
---	--------------------------	--	--

 A. 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane C. code: PPA075	47g Ex. 11		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.16 (d, J=6.91 Hz, 6 H, N(iPr)) 1.10 (d, J=6.91 Hz, 6 H, N(iPr)) 2.58 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.08 (s, 2 H, N(iPr)₂) 3.43 - 3.80 (m, 34 H, CH₂O (Erythritol, PEG, CNEt) 3.77 (s, 6 H, Ph-OCH₃) 4.57 (s, 4 H, -CH₂-triazole) 4.62 - 4.79 (m, 8 H, EtOBz) 6.87 (d, J=8.96 Hz, 4 H, Ph-H) 7.19 - 7.36 (m, 9 H, Ph-H) 7.39 - 7.80 (m, 10 H, Ph-H) 7.90 - 8.00 (m, 2 H, triazole)	147.05
---	--------------------------	--	--	---------------

 A. 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane C. code: PPA078	<u>47i</u> Ex. 11		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.11 (dd, J=16.90, 6.91 Hz, 12 H, N(iPr)₂) 2.57 (t, J=6.02 Hz, 2 H, CH₂CN) 3.07 (m, 2 H, NH(iPr)₂) 3.42 - 3.79 (m, 37 H, -CH₂ (PEG, glycerol, CNEt), OCH₃) 3.77 (s, 4 H, -CH₂(N)triazole)) 4.47 (s, 4 H, -CH₂-triazole) 4.56 (s, 12 H, -CH₂OBz) 4.88 (s, 4 H) 6.79 - 6.92 (m, 4 H, Ph-H) 7.18 - 7.43 (m, 9 H, Ph-H) 7.48 - 7.69 (m, 18 H, OBz-H) 7.84 (s, 2 H, triazole) 7.99 - 8.01 (m, 12 H, OBz-H)	147.17
---	---------------------------------	--	---	---------------

 A. 1-O-DMT-3-O-PPA-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane B. 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane C. code: PPA056	<u>52</u> Ex. 12		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.09 (d, J=6.66 Hz, 6 H, N(iPr)) 1.18 (d, J=6.66 Hz, 6 H, N(iPr)) 2.64 (t, J=6.02 Hz, 2 H, -CH₂CN) 2.96 - 3.13 (m, 2 H, N(iPr)₂) 3.40 - 3.69 (m, 12 H, -CH₂O (PEG)) 3.72 - 3.88 (m, 6 H, -CH₂O (Erythritol, CNEt)) 3.79 (s, 6 H, Ph-OCH₃) 4.39 - 4.45 (m, 4 H, -OCH₂-triazole) 4.47 - 4.56 (m, 8 H, -CH₂-OBz, -CH₂-triazole) 6.87 (d, J=8.96 Hz, 4 H, Ph-H) 7.22 - 7.36 (m, 7 H, Ph-H) 7.37 - 7.42 (m, 2 H, Ph-H) 7.45 - 7.61 (m, 10 H, Ph-H) 8.02 (d, J=7.42 Hz, 2 H, triazole)	148.11
---	------------------------------	--	--	---------------

 A. 2-((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((PPA-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-DMT-propane-1,3-diyl dibenzoate B. 2-((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((phosphodiester-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-phosphodiester-propane-1,3-diyl dibenzoate C. code: PPA058	<u>62</u> Ex. 13 	¹H NMR (300 MHz, DMSO-d₆) δ ppm 0.98 - 1.13 (m, 12 H, N(iPr)₂) 2.71 (t, J=6.02 Hz, 2 H, -CH₂CN) 3.52 (d, J=8.70 Hz, 4 H, -CH₂(Erythritol)) 3.62 (s, 6 H, Ph-OCH₃) 3.67 - 3.83 (m, 4 H, -CH₂(Erythritol)) 4.23 - 4.42 (m, 12 H, CH₂OBz) 4.52 (br. s., 2 H, -CH₂-triazole) 4.60 (s, 2 H, -CH₂-triazole) 6.71 (d, J=8.96 Hz, 4 H, Ph-H) 7.11 - 7.25 (m, 7 H, Ph-H) 7.30 - 7.36 (m, 2 H, Ph-H) 7.39 - 7.91 (m, 32 H, Ph-H and triazole). 147.76
 DMT PPA A. 2-O-DMT-3-O-PPA-1S,4S-bis(O-mPEG2)-butane B. 2,3-O-bis(phosphodiester-1S,4S-bis(O-mPEG2)-butane C. code: PPA016	<u>67</u> Ex. 14 	¹H NMR (300 MHz, ACN-d₃) δ ppm 1.07 - 1.21 (m, 12 H, N(iPr)₂) 2.50 (td, J=5.89, 3.33 Hz, 1 H, -CH₂CN) 2.62 (t, J=6.14 Hz, 1H, CH₂CN) 3.17 - 3.83 (m, 36 H, -CH₂O (PEG, CNEt), -OCH₃, -CHO) 6.80 - 6.92 (m, 4 H, Ph-H) 7.12 - 7.44 (m, 7 H, Ph-H) 7.52 (d, J=8.19 Hz, 2 H, Ph-H) 147.61, 147.66

















 A. 1-O-DMT-3-O-PPA-2,2-bis(Me-O-levulinate)-propane B. 1,3-O-bis(phosphodiester 2,2-bis(Me-O-levulinate))-propane C. code: PPA036	<u>72</u> Ex. 15		¹H NMR (300 MHz, ACN-d₃) δ ppm 1.07 - 1.24 (m, 12 H, N(iPr)₂) 2.05 - 2.22 (m, 6 H, OAc) 2.43 (td, J=6.66, 3.33 Hz, 4 H, levuliny) 2.52 - 2.65 (m, 2 H, N(iPr)₂) 2.65 - 2.77 (m, 4 H, levuliny) 3.07 - 3.23 (m, 2 H, N(iPr)₂) 3.50 - 3.82 (m, 6 H, -CH₂O (erythritol-PPA, erythritol-DMT, CNEt)) 3.79 (s, 6 H, Ph-OCH₃) 4.02 - 4.18 (m, 4 H, -CH₂O (erythritol-levuliny)) 6.90 (d, J=8.70 Hz, 4 H, Ph-H) 7.21 - 7.38 (m, 7 H, Ph-H) 7.40 - 7.52 (m, 2 H, Ph-H)	147.71
--	-----------------------------	--	--	---------------






Table 1B

Exemplary Novel Phosphoramidite (PPA) Monomers – Group B

Chemical name (code identifier)	Chemical Structure
1-O-DMT-3-O-PPA-2R-mPEG4-propane (PPA106)	
1-O-DMT-3-O-PPA-2S-O-mPEG8-propane (PPA033)	
1-O-DMT-3-O-PPA-2S-O-mPEG10-propane (PPA034)	
1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (PPA079)	
1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG7)-1-(Et-O-Ac)-1,2,3-triazole)-propane (PPA084)	

1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG7)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (PPA094)	
1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (PPA077)	
2-(16-(2-DMT-methoxy)ethyl)-1,4,10,13-tetraoxa-7,16-diazacyclooctadecan-7-yl)ethyl PPA (PPA087)	
1-O-DMT-1'-O-PPA-(N,N)-(4,7,13,16-tetraoxa-1,10-diazacyclooctadecane-1,10-diyl)-bisethylurea (PPA047)	

3-O-DMT-10-O-PPA-3,4,6,7,10,11-hexahydro-2H,9H-benzo[b][1,4,8,11]tetraoxacyclotetradecine (PPA049)	
2-((4-((2-DMT)ethoxy)methyl-1,2,3-triazol-1-yl)methyl)-2-((4-((PPA)ethoxy)methyl)-1,2,3-triazol-1-yl)methyl)propane-1,3-diyl dibenzoate (PPA057)	
1-O-DMT-3-O-PPA-2-(4-methyl-1-(<i>p</i>-methoxy)-1,2,3-triazole)-propane (PPA019)	
1-O-DMT-2S-O-PPA-3-(N-(1-aza-15-crown-5)-propane (PPA048)	
1-O-DMT-2S-O-PPA-3-(3-methoxy-pyridin-2-one-1-yl)-propane (PPA051)	

1-O-DMT-2-O-PPA-3-((N,N-dipyridin-2-yl)amino)-propane (PPA052)	
1-O-DMT-2-O-PPA-3-(pyridazin-3-on-2-yl)-propane (PPA053)	
1-O-DMT-1'-O-PPA-N(butyl) Diethanolamine (PPA045)	
1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (PPA069)	
(PPA046)	

(PPA070)



















Table 1C






Exemplary Novel Phosphoramidite (PPA) Monomers – Group C







Chemical name (code identifier)	Chemical Structure
3-O-DMT-4-O-PPA- 1,2,5,6-tetrakis-(O-mPEG2)-hexane (PPA085)	
1-O-DMT-2S-O-PPA-3-O-(acetoxamido-15-crown-6)-propane (PPA024)	
1-O-DMT-2S-O-PPA-3-O-(2,2-bis(Me-O-mPEG2)-3-O-mPEG2-propane) propane (PPA020)	
1-O-DMT-3-O-PPA-2S-(4-(Me-O-PEG2)-1-(Et-O-Ac)-	



1,2,3-triazole)-propane (PPA092)	
1-O-DMT-3-O-PPA-2S-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (PPA103)	
1-O-DMT-3-O-PPA-2S-(4-(Me-O-PEG7)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (PPA104)	
1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (PPA080)	
1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (PPA082)	

1-O-DMT-3-O-PPA-2S-O-(bis(Et-O-PEG2-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole))-3-O--(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole))-propane (PPA076)	
2,2-bis(DMT-O-Me)-1-O-PPA-butane (PPA039)	
N4-(2-O-DMT-Et)-N10-(2-O-PPA-ethyl)-1,7-dioxo-4,10-diazacyclododecane-4,10-dicarboxamide (PPA086)	
19-DMT-1,4,7,11,14,17-hexaoxacycloicosan-9-yl PPA (PPA028)	

2-(3-(10-(3-(2-DMT-methoxy)ethoxy)propanoyl)-1,7-dioxo-4,10-diazacyclododecan-4-yl)-3-oxopropoxy)ethyl PPA (PPA088)	
3-O-DMT-10-O-PPA-5,8,15,18,23,26-hexaoxa-1,12-diazabicyclo[10.8.8]octacosane-13,20-dione (PPA090)	
1-O-DMT-1'-O-PPA-2-(4,7,13,16-tetraoxa-1,10-diazacyclododecane-1,10-diyl)-diethylamide (PPA017)	
1-O-DMT-1'-O-PPA-(4,7,13,16-tetraoxa-1,10-diazacyclododecane-1,10-diyl)-di-PEG1-amide (PPA030)	
1-O-DMT-22-O-PPA-11,11-bis(Me-O-Bz)-3,6,9,13,16,19-hexaoxadocosane (PPA059)	
1-O-DMT-23-O-PPA-5,5,12,12,19,19-hexakis(Me-O-Bz)hexaoxatricosane (PPA066)	

1-(DMT-O-Me)-4-(PPA-2-O-Et) -1,2,3-triazole (PPA067)	
3-O-DMT-2S-O-PPA-1-(7-dioxaneindole)-propane (PPA011)	
1-O-DMT-2S-O-PPA-3-O-((N,N-bis-mPEG2)acetamide)-propane (PPA025)	
1-O-DMT-2-O-PPA-3-(4-methylpiperazine-1-yl)acetamide (PPA095)	
1-O-DMT-2-O-PPA-3-((4-methylpiperazine-1-yl)-Et-(1,2,3-triazole)-propane) (PPA096)	

1-O-DMT-2-O-PPA-3- ((trimethylamino)acetamide)- propane (PPA097)	
1-O-DMT-1'-O-PPA- N,N(dimethyl) Diethanolamine (PPA098)	
1-O-DMT-1'-O-PPA-N(PEG4- OMe) Diethanolamine (PPA099)	
1-O-DMT-2-(S)-O-PPA-3- (N,N-di(PEG4-OMe)amine) (PPA100)	
1-O-DMT-2(S)-O-PPA-3-(1-H- 3-imidazole)-propane (PPA101)	
1-O-DMT-2-O-PPA-3-((1H- imidazol-4-yl)acetamide)- propane (PPA102)	

1-O-DMT-2-O-PPA-3-((1H-imidazol-4-yl)acetamide)-propane (PPA107)	
1-O-DMT-2-O-PPA-3-((1-Me-imidazol-4-yl)acetamide)-propane (PPA108)	

Translocation Control Elements (TCEs) based on novel phosphoramidite monomers.

[00115] As discussed herein, the TCE feature of an SSRT is designed to stall Xpandomer translocation so as to position the reporter code within the nanopore aperture for measurement. The availability of the new phosphoramidite monomeric compounds of the present invention has enabled design of next-generation TCE structures, which control translocation rate through one or more of steric hindrance, electro-repulsion, and preferential interaction with the nanopore. The resistance of the TCE to the driving force of the ion current when positioned at the pore aperture and the consequent increase in applied voltage (i.e., the voltage pulse) necessary to overcome the arrest and resume translocation, can be customized by modulating various properties of the TCE, (and in some embodiments, the reporter codes and other elements of the SSRT) e.g., the bulk, length, and/or charge density. Importantly, because translocation rate is controlled by properties intrinsic to the TCE, translocation control is relieved of the burden of relying on prior art strategies, which employ, e.g., nucleotide hybridization strategies based on reversible interaction with soluble oligonucleotides.

[00116] In certain embodiments, TCEs are polymers produced by solid-phase synthesis using the phosphoramidite method with suitable monomeric building blocks that terminate with a branched structure (i.e., the “brancher”). Branched phosphoramidites are known in the art and include both symmetrical and asymmetrical branchers, commercially available from,

e.g., Glen Research and ChemGenes. In one embodiment, the TCE brancher is a symmetrical branching CED phosphoramidite, wherein each arm of the brancher is linked to a reporter code. Exemplary symmetrical chemical branchers include 1,2,3 -O-tris-(phosphodiester)-propane, 1,3-bis-(5-O-phosphodiester-pentylamido)-2-O-phosphodiester-propane, and 1,4,7-O-tris-(phosphodiester)-heptane.

[00117] Fig. 4 illustrates, in simplified form, how a TCE arrests Xpandomer translocation through a nanopore. Here, each SSRT of the Xpandomer includes reporter codes **485A** and **485B** that are linked to TCE **490** at the ends of the arms of brancher structure **493** of the TCE. In this embodiment, TCE **490** includes a structure **495** with a larger physical bulk relative to that of the reporter codes. Xpandomer translocation through the barrel of nanopore **450** is arrested when TCE **490**, encounters the pore aperture. In certain embodiments, both the bulk of the TCE and the charge densities of the reporter codes (i.e., the local electric field at the arrest site) contribute to translocation arrest. During the pause, reporter code **495A** is held in the barrel of the nanopore and blocks the flow of current through the pore in a characteristic and detectable manner. To overcome the arrest, a voltage pulse is applied to the system, which forces the TCE to enter and pass through the pore. Translocation then resumes until the next TCE encounters the pore aperture.

[00118] To customize translocation control, several structural properties of the TCE (and in certain embodiments, other features of the SSRT) can be adapted. For example, one or more of the length, bulk, and charge density of the TCE and the spatial positioning of charged elements within the barrel of the nanopore can be modified. In some embodiments, the bulk of the TCE is increased by incorporating one or more pendant PEG phosphoramidites into the polymeric structure. For example, the TCE may incorporate from 2 to 30, from 2 to 20, from 3 to 15, or from 4 to 14 pendant PEG phosphoramidite compounds. In other embodiments, TCEs may include any suitable number and combination of phosphoramidite compounds set forth in Tables 1A - C. For example, a TCE may include from 1 to 10, 2 to 8, or 2 or 3 different phosphoramidite compounds, in any order; in certain embodiments, at least one of the phosphoramidite compounds is a pendant PEG phosphoramidite. In certain embodiments, the length of the entire TCE may include from 2 to 30, from 2 to 20, from 3 to 15, or from 4 to 14 phosphoramidite compounds. In some embodiments, the formula of the TCE may be represented by $(PPA1)_{n1}(PPA2)_{n2}$ wherein PPA1 and/or PPA2 represent a pendant PEG phosphoramidite compound and $n1 = 1$ to 12 and $n2 = 0$ to 10. The inventors have discovered that TCE based on this formula significantly reduce sequencing errors (e.g., insertion or deletion events) and enable single-

pulse transitions between sequential SSRTs. In certain embodiments, the TCE includes a polymer synthesized from phosphoramidite compounds with the following sequence: [(1-O-DMT-3-O-PPA-2S-O-mPEG4-propane (compound 12b))]_{n1}[(1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b)]_{n2}, in which n1 is from 0 to 6 and n2 is from 6 to 10. In other embodiments, the TCE includes one or more phosphoramidite chromophores that can be detected by UV radiation, e.g., benzofuran or triazole-containing PPAs.

[00119] In other embodiments, TCEs may include a brancher structure with more than two arms. In one embodiment, the phosphoramidite brancher may have a terminally branched structure. In this embodiment, the brancher has four arms, two of which are linked to the reporter codes, and two of which contribute to translocation control. In other embodiments, the brancher may be customized to optimize features such as size, polarity, and stability. In one embodiment, the brancher includes an isocyanate trimer.

[00120] In further embodiments, the TCEs of the present invention may include two branchers (i.e., “double brancher” TCEs), in which the branchers are separated by a plurality of unbranched phosphoramidites. The branchers may be symmetrical or asymmetrical structures. The asymmetric structure may be a single enantiomer or racemic. Moreover, a combination of the racemate and/or both enantiomers can be used at different positions in the TCE. In this embodiment, each brancher contributes to a distinct translocation pause event, the first of which may be referred to as a “code pause” that maintains a reporter code in the nanopore, and the second of which may be referred to as a “clock pause” that produces a unique signal indicating that the preceding reporter codes has been “read” by the detection system.

[00121] Table 2 sets forth several exemplary TCE sequences. It is to be emphasized that the present invention is not intended to be limited to these particular embodiments, as the skilled artisan will appreciate that, based on the present disclosure, an extensive library of diverse TCEs can be designed to suit a wide range of experimental requirements. In certain embodiments, any of the TCE sequences set forth below could terminate at the end distal to the brancher with one or more spacer compounds, including C3, benzofuran, or PEG3. The key in Table 2 identifies the compounds in their form as phosphoramidite monomers. It will be readily apparent to one of ordinary skill in the art that the descriptor “phosphoramidite” only applies to the compounds in monomeric form; descriptors that apply to the compounds in multimeric, “in-line”, form are set forth in Table 1A.

TABLE 2
Exemplary Translocation Control Elements (TCEs)

TCE sequence	Key
Y(PPA032) ₆ (PPA061) ₆ Y(PPA032) ₂ (PPA061) ₁₀ Y(PPA032) ₂ (PPA061) ₈ Y(PPA032) ₂ (PPA061) ₆ Y(PPA032) ₅ (PPA061) ₆	Y: symmetrical brancher (e.g., Chemgenese Cat. No. CLP-5215) PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
Y(PPA061) ₈	Y: symmetrical brancher PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
Y(PPA032) ₁₂	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
Y(PPA031) ₂ (PPA032) ₁₀	Y: symmetrical brancher PPA031: 1-O-DMT-3-O-PPA-2s-O-mPEG6-propane PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
Y(PPA040) ₁₂	Y: symmetrical brancher PPA040: 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG2)-propane

Y(PPA032) ₄ (PPA064) ₃	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA064: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane
Y(PPA032) ₄ (PPA068) ₅	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA068: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
Y(PPA032) ₄ (PPA094) ₃	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA094: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG7)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane
Y[(PPA032)(PPA062)] ₆	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA062: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane
Y(PPA044) ₈	Y: symmetrical brancher PPA044: 1-O-DMT-8-O-PPA-N,N-Diethylpiperazine
Y(PPA062) ₁₂	Y: symmetrical brancher PPA062: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane
Y(PPA063) ₆	Y: symmetrical brancher PPA063: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane

Y(PPA065) ₈	Y: symmetrical brancher PPA065: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane
Y(PPA068) ₆	Y: symmetrical brancher PPA068: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
Y(PPA069) ₄	Y: symmetrical brancher PPA069: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane
Y(PPA075) ₂	Y: symmetrical brancher PPA075: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane
Y(PPA078) Y(PPA078) ₃	Y: symmetrical brancher PPA078: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
Y(PPA093) ₅	Y: symmetrical brancher PPA093: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane
Y(PPA094) ₄	Y: symmetrical brancher PPA094: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG7)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane
Y(PPA032) ₆ (PPA060) ₆	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA060: 1-O-DMT-2-O-PPA-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane
Y(PPA032) ₈ (PPA056) ₃	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane

	PPA056: 1-O-DMT-3-O-PPA-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane
Y(PPA032) ₆ (PPA043)(PPA032) ₆ (PPA043) ₂	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA043: 1-O-DMT-2S-O-PPA-3-(4-methylpiperazine-1-yl)-propane
Y(PPA050) ₁₂	Y: symmetrical brancher PPA050: 1-O-DMT-3-O-PPA-2S-O-(PEG4-O-Bz)-propane
Y(PPA031) ₁₂ (PPA018) ₂ Y(PPA031) ₆ (PPA018) ₂	Y: symmetrical brancher PPA031: 1-O-DMT-3-O-PPA-2s-O-mPEG6-propane PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(PPA033) ₁₂ (PPA018) ₂	Y: symmetrical brancher PPA033: 1-O-DMT-3-O-PPA-2S-O-mPEG8-propane PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(PPA033) ₇	Y: symmetrical brancher PPA033: 1-O-DMT-3-O-PPA-2S-O-mPEG8-propane
Y(DTB)X(PPA018) ₄ XD(DTB) Y(DTB)X(PPA018) ₄ XD(DTB)	Y: symmetrical brancher DTB: desthiobiotin X: PEG3 D: PEG6 PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(DTB)X(PPA032) ₄ XD(DTB)(PPA018) ₂	Y: symmetrical brancher DTB: desthiobiotin X: PEG3 D: PEG6 PPA032: PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(PPA032) ₁₂ (PPA018) ₂	Y: symmetrical brancher

Y(PPA029) ₁₂ (PPA018) ₂	PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA029: 1-O-DMT-2-O-PPA-3-O-mPEG2-propane PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(PPA032) ₄ (PPA018) ₂ W Y(PPA018) ₂ W	Y: symmetrical brancher PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane W: SIMA-HEX fluorophore
YL(Fluor)(PPA018) ₂	Y: symmetrical brancher L: C2 spacer Fluor: Fluorescein PPA PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
Y(PPA032) ₆ (Lev)(PPA032) ₄ X (PPA032) (PPA032) (PPA032) (PPA032) (PPA018) (PPA018) N Y(PPA032) ₆ (Lev)DDDX D D D (PPA018) (PPA018) N Y(PPA032) ₈ DD(Lev)(PPA032) ₄ X (PPA032) (PPA032) (PPA032) (PPA032) (PPA018) (PPA018) N	Y: symmetrical brancher Lev: asymmetrical branching CED phosphoramidite (Chemgenes CLP-7169) X: PEG3 D: PEG6 PPA032: PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane PPA018: PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane

Y(PPA032)₈DD(Lev)DDDX D D D (PPA018) (PPA018) N Y(PPA032)₆(Lev)DDD(PPA018)₂X D D D N	
Y(DTB)	Y: symmetrical brancher DTB: desthiobiotin

Methods and Constructs to Reduce the Rate of Insertion and Deletion Events during Xpandomer Translocation

[00122] In other aspects, the present invention provides means of translocation control through Xpandomer modification in combination with discrete translocation deceleration features (referred to herein also as “D-cells”) designed into the SSRT. As disclosed herein, Xpandomers are subjected to several processing steps following synthesis, including an amine modification step. During amine modification, Xpandomers are treated with succinic anhydride, which reacts with the secondary amine groups (and, in certain circumstances, the primary amine group introduced by the acid cleavage step) on the spermine constituents of the polymerase enhancement regions of the SSRT. Succinylation of an amine group results in the introduction of a negatively charged hemi-succinate group. By increasing the degree of succinylation, the degree of negative charge on the spermine-based enhancers is likewise increased. Each spermine phosphoramidite constituent has a net charge of (+3); in a standard modification reaction, the charge of each individual amine moiety is changed from (+1) to (-1). The inventors have discovered conditions that give varying degrees of spermine succinylation such that the net charge of a spermine constituent can be changed from between (+3) to (-5). Under certain conditions, increasing the negative charge of the enhancer regions may be desirable so as to increase the rate of Xpandomer translocation upon application of a voltage pulse (referred to herein as an enhancer electromobility. Notably, this electromobility has been found to reduce the percentage of insertion errors during the sequence read as well as to increase overall sequencing throughput.

[00123] Thus, in certain embodiments, Xpandomer processing includes an amine modification step. Amine (e.g., spermine) modification may be achieved through altering one or more of the succinylation reaction conditions, e.g., the reaction time, temperature, pH, and/or the concentration of succinic anhydride used in the reaction. In other embodiments, Xpandomer processing further includes one or more of a HEPES wash step following the amine modification step in order to achieve more complete amine succinylation.

[00124] In another aspect, the present disclosure provides one or more translocation deceleration features or regions (the terms “features” and “regions” are used interchangeably in this context), which are permanently charged, e.g., tertiary or quarternary amines and/or bulky compounds. Translocation deceleration features may be introduced into the SSRT at a position within or adjacent to the polymerase enhancers. The deceleration features are selected so as not to be altered during the Xpandomer modification (e.g., succinylation)

reaction. Inclusion of one or more deceleration features into a suitable location in the SSRT has been found to reduce the percentage of deletion errors, which arise due to the increased translocation rate resulting from over-modification of the enhancer. Without being bound by theory, it is speculated that the bulk of the deceleration feature creates a “friction”-type of force that reduces the rate of Xpandomer translocation upon encountering the nanopore aperture. Typically, the deceleration features are introduced into the SSRT at a position between the polymerase enhancer and the reporter code (i.e., adjacent to the enhancer).

[00125] The translocation deceleration features of the present invention may incorporate any suitable number and combination of the phosphoramidite compounds set forth in Tables 1A – 1C or commercially available phosphoramidites. In some embodiments, the deceleration features include a combination of from 1 to 4 different monomeric units. In other embodiments, the deceleration features may include 1 or 2 different monomeric units. The entire length of a deceleration feature may be from 1 to 15 monomeric units or, in other embodiments, from 4 to 12 or from 6 to 10 monomeric units. Table 3 sets forth non-limiting examples of alternative translocation deceleration features. The key in Table 3 identifies the compounds in their form as phosphoramidite monomers. It will be readily apparent to one of ordinary skill in the art that the descriptor “phosphoramidite” only applies to the compounds in monomeric form; descriptors that apply to the compounds in multimeric, “in-line”, form are set forth in Table 1A.

TABLE 3
Exemplary Translocation Deceleration Features

D-cell sequence	Key
(PPA058) ₆	PPA058: 2-((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((PPA-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-DMT-propane-1,3-diyl dibenzoate
[D(PPA056)] ₃ DD	D: PEG6 PPA056: 1-O-DMT-3-O-PPA-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane
D(PPA060)D(PPA060)DDDD	D: PEG6 PPA060: 1-O-DMT-2-O-PPA-3-(4-(Me-O-

	PEG3-O-Bz)-1-(1,2,3-triazole))-propane
[D(PPA063)] ₃ DD	D: PEG6
[D(PPA063)] ₄ DD	PPA063: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane
[D(PPA064)] ₃ DD	D: PEG6
[D(PPA064)] ₄ DD	PPA064: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane
[D(PPA068)] ₄ DD	D: PEG6
	PPA068: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
[D(PPA069)] ₃ DD	D: PEG6
	PPA069: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane
[D(PPA078)] ₃ DD	D: PEG6
	PPA078: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
[D(PPA63)] ₃ DD	D: PEG6
	PPA063: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane
[D(PPA093)] ₄ DD	D: PEG6
	PPA093: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane
[D(PPA094)] ₃ DD	D: PEG6
	PPA094: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG7)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane
DDD(PPA063) ₃ DD	D: PEG6
	PPA063: 1-O-DMT-3-O-PPA-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane
D(PPA043)(PPA043)DDDDDD	D: PEG6
D(PPA043) ₂ D(PPA043) ₂ DDDDD	PPA043: 1-O-DMT-2S-O-PPA-3-(4-methylpiperazine-1-yl)-propane
(PPA109)(PPA043) ₂ W(PPA043) ₂ WDD	D: PEG6
	PPA043: 1-O-DMT-2S-O-PPA-3-(4-

	methylpiperazine-1-yl)-propane PPA109:in-line PEG12
DD(PPA033)D(PPA033)D(PPA033)DD DD(PPA033) ₂ D(PPA033) ₂ D(PPA033) ₂ DD	D: PEG6 PPA033: 1-O-DMT-3-O-PPA-2S-O-mPEG8-propane
[D(PPA075)] ₃ DD	D: PEG6 PPA075: 1-O-DMT-3-O-PPA-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane
DD(PPA001)D(PPA001)DDD	D: PEG6 PPA001: 1-O-DMT-1'-O-PPA-N(p-tolyl)-diethanolamine

Reporter Codes

[00126] Each SSRT uses the TCE to position the reporter code within a zone of the nanopore that has high ion current resistance. In alpha hemolysin, this zone is the stem. In this zone, different reporters are sized to block ion flow at different measurable levels. Reporters can be designed by selecting a sequence of specific phosphoramidites from the collection of phosphoramidite monomeric compounds set for in Tables 1A – 1C and/or commercially available libraries. Suitable monomeric compounds are also disclosed in Applicants' US patent no. 10,457,979, which is herein incorporated by reference in its entirety, including PEG3, PEG6, and C2.

[00127] Each constituent monomeric compound contributes to the net current resistance according to its position in the nanopore, its displacement, its charge, its interaction with the nanopore, its chemical and thermal environment and other factors.

[00128] Reporter code design is guided by balancing measurement characteristics including: (i) normalized ion current (I/I_0): where I is ion current and I_0 is the open channel current; (ii) ion current noise: includes multi-state responses, blockages, random spiking, and the like; and/or (iii) release time of the control moiety or the time during which the TCE is otherwise is stalled at the stem entrance.

[00129] Figs 5A – 5D illustrate how different reporter codes can be designed to maintain similar charge densities along the backbone, while providing signature levels of pore blockage due to differential volumes occupied by each unique monomeric constituent.

In these illustrations, a nanopore is depicted in cross-section with the barrel and vestibule portions indicated. The reporter codes are depicted in simplified form with the black circles representing the charged phosphodiester moieties introduced by the phosphoramidite constituents of the codes. Fig. 5A illustrates a linear code, in this embodiment identified a “zero PEG code”, although the invention is not intended to be so limited, e.g., in certain embodiments linear codes may include PEG moieties. Figs 5B, 5C, and 5D illustrate how unique codes can be constructed from repeating units of a single branched monomeric compound showing exemplary PEG-based structures, e.g., a pendant PEG compound. In these embodiments, the branched-moieties of the three different codes occupy different volumes in the pore channel and thus generate unique signals that can be differentiated from each other. In these embodiments, the charge density along the backbone is the same for each reporter code. However, as discussed further herein, in other embodiments, reporter codes can be designed with a gradient of charge density along the backbone.

[00130] Reporter ion current blockage and its duplex release time is also modulated by measurement conditions such as: (i) voltage; (ii) electrolyte; (iii) temperature; (iv) pressure; and/or (v) pH, as described further herein.

[00131] In some embodiments, the TCE associated with the reporter also contributes to the ion current blockage.

[00132] For a given set of measurement conditions reporters can be designed for a minimum and maximum I/I_o levels that define the measurement dynamic range. Other reporters can be designed with different I/I_o levels within the dynamic range. As each reporter is paused in the nanopore, the measured I/I_o level must remain stationary long enough and have low enough noise that the reporter type can be uniquely distinguished. Dynamic range is maximized by selecting a backbone of low impedance molecules (reporter code polymers), typically those with small physical cross-sections and low linear mass densities.

[00133] Table 4 sets forth exemplary reporter codes, though It is to be understood that the present invention contemplates reporter codes incorporating any suitable combination and number of phosphoramidite compounds disclosed herein. The key in Table 4 identifies the compounds in their form as phosphoramidite monomers. It will be readily apparent to one of ordinary skill in the art that the descriptor “phosphoramidite” only applies to the compounds in monomeric form; descriptors that apply to the compounds in multimeric, “in-line”, form are set forth in Table 1A.

TABLE 4
Exemplary Reporter Codes

Code Sequence	Key
XXXXXX	X: Triethyleneglycol (PEG3)
DDDD	D: Hexaethyleneglycol (PEG6)
LLLLLLLLLLLLL	L: ethane (C2)
WW	W: Dodecaethyleneglycol (PEG12)
DDDDDLLL	D: PEG6 L: C2
XXXXLLLL	X: PEG3 L: C2
DDLLEDX	D: PEG6 X: PEG3 L: C2
DDLLELXX	D: PEG6 L: C2
DDLLELLEDX	D: PEG6 X: PEG3 L: C2
DD(PPA032)(PPA032)LDX	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD(PPA032)LL(PPA032) DX	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD(PPA032)(PPA032)LXXX	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DXLLL(PPA032)(PPA032)LXXX	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD((PPA040)(PPA040)LXXX	D: PEG6 X: PEG3 L: C2 PPA040: 1-O-DMT-3-O-PPA-2,2-bis(Me-O-

	mPEG2)-propane
DD(PPA050)(PPA050)LXXX	D: PEG6 X: PEG3 L: C2 PPA050: 1-O-DMT-3-O-PPA-2S-O-(PEG4-O-Bz)-propane
DD(PPA061)(PPA061)LXXX	D: PEG6 X: PEG3 L: C2 PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
DD(PPA032) ₃ LLDX	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD(PPA040) ₃ LLDX	D: PEG6 X: PEG3 L: C2 PPA040: 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG2)-propane
DD(PPA032) ₃ XXLLLL	D: PEG6 X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD(PPA032) ₃ LLLLLLLL	D: PEG6 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DD(PPA050) ₃ LLDX	D: PEG6 X: PEG3 L: C2 PPA050: 1-O-DMT-3-O-PPA-2S-O-(PEG4-O-Bz)-propane

DD(PPA061) ₃ LLDX	D: PEG6 X: PEG3 L: C2 PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
[X(PPA032)] ₄ X	X: PEG3 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
DDLL(PPA031)X(PPA031)X(PPA031)L	D: PEG6 X: PEG3 L: C2 PPA031: 1-O-DMT-3-O-PPA-2s-O-mPEG6-propane
DDLL(PPA033)X(PPA033)X(PPA033)L	D: PEG6 X: PEG3 L: C2 PPA033: 1-O-DMT-3-O-PPA-2S-O-mPEG8-propane
XXL(PPA032) ₆ LLLLLLL	X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
XXL(PPA032) ₆ XXL	X: PEG3 L: C2 PPA032: 1-O-DMT-3-O-PPA-2S-O-mPEG4-propane
XXL(PPA050) ₆ LLLLLLL	X: PEG3 L: C2 PPA050: 1-O-DMT-3-O-PPA-2S-O-(PEG4-O-Bz)-propane
XXL(PPA061) ₆ LLLLLLL	X: PEG3 L: C2 PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
XXL(PPA040) ₆ XLLLL	X: PEG3 L: C2 PPA040: 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG2)-propane

XXL(PPA068) ₆ LLLLLLL	X: PEG3 L: C2 PPA068: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane
XXL(PPA065) ₆ LLLLLLL	X: PEG3 L: C2 PA065: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane
XXL[(PPA050)(PPA061)] ₃ LLLLLLL	X: PEG3 L: C2 50: hydroxyl (Bz) pendant PEG4 (PPA050) PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
XXL(PPA062) ₆ LLLLLLL	X: PEG3 L: C2 PPA062: 1-O-DMT-3-O-PPA-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane
XX(PPA061) ₅ XLLLLLL	X: PEG3 L: C2 PPA061: 1-O-DMT-3-O-PPA-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane
XXX(PPA026)LL(PPA026)XXLL	D: PEG6 X: PEG3 L: C2 PPA026: 1-O-DMT-3-O-PPA-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane
XXL(PPA007) ₄ LLXXLL	X: PEG3 L: C2 PPA007: 3-O-DMT-2S-O-PPA-1-(1-dimethoxyquinazolinedione)-propane
XXL(PPA010) ₄ LLXXLL	X: PEG3 L: C2 PPA010: 3-O-DMT-2S-O-PPA-1-(N9-(3,6-dimethoxycarbazole)-propane
(PPA004) ₄ LLLL	L: C2 PPA004: 1-O-DMT-1'-O-PPA-2,2'-(sulfonylbis(benz-4-yl))- diethanol
XXL(PPA005) ₄ LLXXLL	X: PEG3 L: C2 PPA005: 1-O-DMT-1'-O-PPA-(2,2'-

	bipyridin-4,4'-yl)-dimethanol
XXL(PPA006) ₄ LLXXLL	X: PEG3 L: C2 PPA006: 3-O-DMT-2S-O-PPA-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane
XXL(PPA009) ₄ LLXXLL	X: PEG3 L: C2 PPA009: 3-(1-O-DMT-2S-O-PPA-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide
XXL(PPA012) ₄ LLXXLL	X: PEG3 L: C2 PPA012: 3-O-DMT-2S-O-PPA-1-(N1-(6-Azathymine))-propane
XXL(PPA014) ₄ LLXXLL	X: PEG3 L: C2 PPA014: 1-O-DMT-5-O-PPA-hexahydrofuro[2,6]furan
XXL(PPA008) ₄ LLXXLL	X: PEG3 L: C2 PPA008: 1-O-DMT-1'-O-PPA-(octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol
XXL(PPA015) ₄ LLXXLL	X: PEG3 L: C2 PPA015: 3-O-DMT-2S-O-PPA-1-(N1-(2-Me-5-nitroindole)-propane
XXL(PPA013) ₄ LLXXLL	X: PEG3 L: C2 PPA013: 3-O-DMT-2S-O-PPA-1-(N1-(2-Me-5-nitroindole)-propane
XXL(PPA018) ₄ LLXXLL	X: PEG3 L: C2 PPA018: 3-O-DMT-2S-O-PPA-1-(5-benzofuran)-propane
XXL(PPA023) ₄ LLXXLL	X: PEG3 L: C2

	PPA023: 1-O-DMT-2-O-PPA-3-O-mPEG4-propane
XXL(PPA026) ₄ LLXXLL	X: PEG3 L: C2 PPA026: 1-O-DMT-3-O-PPA-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane
XXL(PPA024) ₄ LLXXLL	X: PEG3 L: C2 PPA024: 1-O-DMT-2S-O-PPA-3-O-(acetoxamido-15-crown-6)-propane
XLL(PPA040) ₂ XXXLLLL	X: PEG3 L: C2 PPA040: 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG2)-propane
XLL(PPA027) ₂ XXXLLLL	X: PEG3 L: C2 PPA027: 1-O-DMT-3-O-PPA-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane
XLL(PPA022) ₂ XXXLLLL	X: PEG3 L: C2 PPA022: 1-O-DMT-2-O-PPA-3-O-mPEG2-propane
XX(PPA022)X(PPA022)X(PPA022)X	X: PEG3 L: C2 PPA022: 1-O-DMT-2-O-PPA-3-O-mPEG2-propane

Synthesis of SSRTs and XNTPs

[00134] As disclosed herein, symmetrically synthesized reporter tethers (SSRTs) are synthesized using standard automated oligonucleotide synthesis protocols. Fig. 6 illustrates SSRT synthesis in simplified form. In step A, a phosphoramidite is immobilized on a solid support; in step B, phosphoramidite coupling is used to polymerize phosphoramidite monomers on the support; in step C, a symmetric brancher is adding to the growing SSRT

structure; in step D, symmetric phosphoramidite branches are polymerized off each arm of the symmetric brancher; in step E, a terminal azido group is added that enables conjugation of the SSRT to a dNTP-2c via a click reaction; in step F, the SSRT is released from the substrate in its final form.

In one embodiment, SSRT synthesis utilizes a four-step iterative process that includes 1) synthesis of SSRT polymers on solid support controlled pore glass beads (reflected in the cartoons of steps A – D). In this step, SSRTs are synthesized one reporter construct at a time at a 1 μ M scale using a MerMade™ 12 Synthesizer (commercially available from BioAutomation). The MerMade™ sequence manager is first prepared followed by preparation of the phosphoramidites (e.g., preparation of 0.067M solution of each phosphoramidite). Suitable coupling times for each phosphoramidite are programmed into the synthesizer. The SSRT synthesis cycle is based on a conventional four step process: detritylation (using a solvent of, e.g., 3% DCA in dichloromethane), monomer coupling (using a solvent of, e.g., 0.25M ETT in acetonitrile), capping (using solvents of, e.g., THF/lutidine/Ac₂O (CAP A) and 16% methylimidazole in THF (CAP B)), and oxidation (using a solvent of, e.g., 0.02M I₂ in THF/pyridine/H₂O). Step 2) functionalization of the 5' end with a manual conversion that displaces the Br with azide (reflected in the cartoon of step E), i.e. “azido modification”. In this step, the synthesis column is washed with 1mL DCM and transferred to a 2mL tube; an azide conversion solution is prepared (100mM sodium iodide and 100mM sodium azide in DMF) and 1.6mL is added to the tube and incubated for 2 hrs. at RT; the support is then rinsed with 1mL DMF and transferred to the column; the column is rinsed with 2mL DMF followed by 3mL ACN and 1mL DCM. Step 3) removal of cyanoethyl protection groups. In this step, a 10% DEA solution is prepared in ACN that may include 0.1M nitromethane; with vacuum, a steady stream of this solution is passed through the column for at least 10'; the column is then rinsed with 2mL ACN followed by 1mL DCM. Step 4) cleavage complete deprotection of the SSRT from the solid support (reflected in the cartoon of step F). In this step, the support is transferred to a 2ml tube and 500 μ L of 30% NH₄OH that may include 100mM nitromethane is added to the tube and incubated for 30' at 55°C; the tube is then chilled for 5' in a freezer; 500 μ L of 40% methylamine is added to the tube and incubated for 1hr at 65°C. The sample is then chilled for 5' in a freezer; the sample is then desalted by draining the column and rinsing with 15mL H₂O; the SSRT is then eluted from the column with 100mM TEAA and quantitated.

Fig. 7 illustrates four exemplary SSRT products that can be used as reporter constructs for the formation of XATP, XCTP, XGTP, and XTTP, each of which is designed

to generate a unique electronic signal when passed through a nanopore. The key illustrates the chemical structures of these particular phosphoramidite as they exist in the final SSRT reporter construct. As used throughout the present disclosure, R represents azido hex, Q represents spermine, D represents PEG6, X represents PEG3, L represents C2 spacer, 4 represents pendant PEG 4, Y represents the symmetric chemical brancher, and 5 represents benzofuran. In this embodiment, R provides the azide conjugation feature, the QQ polymer provides the enhancer feature, the Y444444444444 polymer provides the TCE feature, and the DDLLDX, DDD44LXXX, DDD4444LLDX, and XXL444444LLLLLLL polymers provide the four unique reporter code features.

Fig. 8A summarizes formation of an exemplary XNTP via the cyclization of an SSRT and a dNTP-2c, via a copper catalyzed click reaction. As used throughout the present disclosure, a “dNTP-2c” refers to a cleavable nucleoside triphosphoramidate analog that includes an 1,7-octadiynyl linker conjugated to the heterocycle moiety of the nucleoside and a 5-hexynyl linker conjugated to the alpha phosphoramidate moiety of the triphosphoramidate. This is a three-step process that includes: 1) the click reaction; in one embodiment, the SSRT and dNTP-2c are added to a click reaction solution composed of 0.2mM CuSO₄/0.6mM THPTA/1.2mM NaAsc and incubated for 30'; 2) standard HPLC purification; and 3) desalting using art-recognized methodologies. Further details of the structure and synthesis of the dNTP-2c compounds is disclosed in Applicants' issued U.S. patent no. 10,301,345, which is herein incorporated by reference in its entirety. Fig. 8B illustrates an exemplary XNTP in a detailed chemical structure.

In certain embodiments, the nucleobase of the XNTP may be a non-natural analog, e.g., 7-deazaadenine, 7-deazaguanine, and the like.

Additional means of translocation control

I. Translocation control by reversible binding of translocation control moieties

[00135] This embodiment of translocation control is illustrated in simplified form in Figs 9A and 9B. In this embodiment, the rate of translocation is controlled by the binding and dissociation of soluble translocation binding moieties to the TCEs of the Xpandomer. As shown in Fig. 9A, binding of a first soluble translocation binding moiety to the first TCE element, proximal to a first reporter code of the Xpandomer, forms a “code translocation control complex”. This reversible interaction stalls, pauses, or arrests (for simplicity, these terms are used interchangeably herein) the first reporter code in the nanopore and generates a change in current that is unique to the first code. These code signals are used to identify each

monomeric unit of the Xpandomer. Subsequently, as depicted in Fig. 9B, the first translocation binding moiety dissociates from the first TCE, allowing the first reporter code to pass through, i.e. exit, the nanopore on the opposite side from which it entered the pore. This translocation event is then stalled by the binding of a second translocation binding moiety to the second TCE in the Xpandomer, distal to the first reporter code. This second translocation control complex is referred to herein as the “clock translocation control complex”. This interruption of translocation generates a change in current (i.e. a “clock signal”) signaling complete translocation of the first code region through the nanopore. In certain embodiments, the clock signals of each unit of the Xpandomer may be identical, or nearly indistinguishable, from each other. One of skill in the art will appreciate that the code signals are sufficient in themselves to determine sequence information of the Xpandomer.

[00136] Any suitable set of reversible binding partners may be used for translocation control according to the present invention. In one embodiment, the TCEs include a derivative of biotin, while the translocation control moiety is provided by streptavidin. In this embodiment, the biotin derivative may be engineered to bind streptavidin with lower affinity than natural biotin. One example of a suitable biotin derivative is desthiobiotin (DTB), which is depicted in Fig. 10. In other embodiments, the biotin-SA TCE system can be controlled by, e.g., using other biotin analogs that form weaker biotin-SA complexes, and/or using SA mutants that form weaker complexes.

II. Translocation control through tuning of run conditions

[00137] It has been found that fine tuning of various conditions used in the nanopore-based detection system improves Xpandomer translocation control and the accuracy of code reads. Thus, in other aspects, the present disclosure provides means to improve the rate of polymer translocation through a nanopore by modification of one or more of the following run conditions:

A. Voltage parameters

[00138] The flow of ions from the cis chamber to the trans chamber of the nanopore-based detection systems described herein results from the application of a voltage potential across the membrane that is referred to interchangeably as the “read voltage” or “baseline voltage”. In one embodiment of the present invention, Xpandomer translocation rate is modulated by altering the baseline voltage. In some embodiments, the baseline voltage may be in the range of from about 40mV to about 150mV. In other embodiments, the baseline voltage may be in the range of from about 90mV to about 110mV. In yet other embodiments, the baseline voltage may be in the range of from about 55mV to about 75mV. In some

embodiments, a higher baseline voltage may be desired to capture reporter code reads at a higher rate.

[00139] As discussed herein, Xpandomer translocation is arrested when the TCE proximal to a reporter code encounters the aperture of the pore. The reporter code is maintained in the pore until a voltage pulse is applied that is sufficiently strong to overcome the resistance provided by TCE structure held at the pore. Thus, in another embodiment, Xpandomer translocation rate is modulated by altering the strength of the pulse voltage. In some embodiments, the pulse voltage is in the range of from about 250mV to about 2000mV. In other embodiments, the pulse voltage is in the range of from about 550mV to about 700mV. Likewise, the duration of the voltage pulse can influence the rate of Xpandomer translocation. In some embodiments, the duration of the voltage pulse is in the range of from about 1 μ s to about 50 μ s. In other embodiments, the duration of the voltage pulse is in the range of from about 5 μ s to about 10 μ s. In another embodiment, the periodicity of the pulse voltages may be optimized. In some embodiments, the periodicity is in the range of from about 0.5ms to about 20ms. In yet other embodiments, the periodicity of the pulse voltages is from about 0.5ms to about 1.5ms. The skilled artisan will appreciate that the strength, duration, and periodicity of the optimal voltage pulses will depend upon many factors, e.g., the force of TCE.

B. Salts

[00140] The rate of current flow through the nanopore-based detection systems described herein can be influenced by the salt composition of the buffers that fill the cis and trans chambers of the system. Thus, in certain embodiments, the rate of Xpandomer translocation through the pore can be modulated by salt composition. In these embodiments, salts comprising any suitable mono- or di-valent cation may be utilized. In some embodiments, suitable salts include, but are not limited, to NH₄Cl, MgCl₂, LiCl, KCl, CsCl, NaCl, and CaCl₂. In other embodiments, suitable salts include those in which the anion is acetate. In conditions where a slower current is desired, salts with a lower ionic mobility, e.g., LiCl may be advantageous. In some embodiments, the trans chamber comprises 2M NH₄Cl and a second optional salt with a suitable molarity around 0.2M and the cis chamber comprises NH₄Cl with a suitable molarity in the range from about 0.4M to about 1M and a second optional salt with a suitable molarity in the range from about 0.2M to about 0.8M. In other embodiments, other molarities lying outside of these ranges and/or other combinations of salts may be desirable.

C. Chaotropic agents

[00141] In certain other aspects, the cis chamber of the nanopore-based detection systems of the present invention may include one or more chaotropic agents to improve translocation of individual polymeric analytes, e.g., linearized Xpandomers. Any suitable chaotropic agent may be employed, e.g., urea and/or guanidine hydrochloride (GuCl). In some embodiments, the buffer compositions of the cis chamber include GuCl and/or urea in the range from about 200mM to about 2M.

D. Osmotic gradients

[00142] In other aspects, present invention provides nanopore-based detection systems in which an osmotic gradient is established across the membrane to influence that rate of Xpandomer translocation through the pore. Without being bound by theory, it is hypothesized that a gradient, wherein the concentration of salts and/or other additives is higher in the trans chamber relative to the cis chamber, generates a flow of water towards the nanopore, thereby drawing Xpandomers towards the pore. Under these conditions, an increase in the rate of event frequencies (e.g., code reads) may be observed at a lower run voltage. Thus, in some embodiments, the run conditions include establishment of an osmotic gradient of around 50% across the membrane; e.g., a salt (and/or other additive) concentration of around 1M in the cis chamber and a salt (and/or other additive) concentration of around 2M in the trans chamber. In further embodiments, any other suitable osmotic gradient may be employed.

E. Solvents

[00143] It has been found that certain solvents can enhance Xpandomer solubility and improve the rate translocation through a nanopore. Thus, in certain embodiments, the sample buffers of the present invention include one or more organic solvents. Suitable solvents include, but are not limited to, 3-methyl-2-oxazolidinone (MOA), DMF, ACN, DMSO, and NMP used in the range of from about 1% to about 25%.

F. Buffers, additives, and other run conditions

[00144] Suitable buffers for use in the present invention include, but are not limited to, 20mM-100mM HEPES with a pH of about 7.4 and bis-tris-propane buffers with a molarity in the range of from about 25mM to about 250mM and with a pH in the range of from about 6 to about 10. In other aspects, the buffers of the present invention may include certain detergent additives, such as sodium hexanoate (NaHex), to enhance the rate of Xpandomer translocation. In certain embodiments, the sample buffers of the present invention include around 20mM NaHex. Other suitable additives include, but are not limited to, stabilizers such as EDTA and redox reagents. The viscosity of any of the buffers may also be altered by

additives such as PEG, glycerol, ficoll and the like.

[00145] In other aspects, translocation rate of Xpandomers may be modulated by temperature. In some embodiments, the run temperature may be in the range from about 4°C to about 40°C. In other embodiments, the run temperature may be in the range from about 16°C to about 22°C.

Cleavable Extension Oligonucleotides

[00146] In another aspect, the present disclosure provides cleavable extension oligonucleotides (EO) for Xpandomer synthesis. The cleavable design feature enables the EO to be removed, i.e., cleaved from, the Xpandomer following synthesis and prior to nanopore analysis. This functionality provides advantages when it is undesirable to translocate a polynucleotide sequence through a nanopore. Xpandomer synthesis, processing, and nanopore sequence analysis are carried out as has been described, e.g., in Applicants' PCT patent application no. PCT/US18/67763, which is herein incorporated by reference in its entirety.

[00147] One embodiment of a cleavable extension oligonucleotide is illustrated in simplified form in Fig. 11. The 3' end of the EO is modified to include a cleavable bond, e.g., an acid cleavable phosphoramidate bond, represented here by "P-NH". The nucleobase attached to the EO by the cleavable linker provides a free 3' hydroxyl group for Xpandomer synthesis, the directionality of which is indicated by the dashed arrow. The base moiety of the same nucleobase is modified to provide other features necessary for nanopore translocation, e.g., a leader group also referred to herein as a "pendant leader sequence". Thus, following extension of the EO to form the Xpandomer, acid treatment releases the oligonucleotide primer, while the leader group and other joined features, remain associated with the Xpandomer. Advantageously, the inventors have found that Xpandomer synthesis is unaffected by the addition of a pendant leader sequence on the cleavable extension oligo.

Ratcheting

[00148] One drawback of nanopore-based detection systems practiced in the art is the depletion of current over time resulting from electrolyte exhaustion that occurs during continuous application of DC voltage. For example, where an electrolyte circuit is based on a ferrocyanide-ferricyanide redox couple, each well in a nanopore array has a limited volume and thus contains a limited number of these redox ion species. Under DC voltage, one species converts to the other and will cause a drop in current. To overcome this problem of

current depletion and maintain a more balanced current over time, the present disclosure provides means for detecting polymeric analytes with a nanopore-based detection system that relies, instead, on an alternating current (AC) pattern of voltage application. This pattern is referred to herein as “ratcheting”. A generalized overview of ratcheting is presented in FIG. 12. The top panel of FIG. 12 illustrates an exemplary pattern of voltage application; in this embodiment, a “forward” read voltage of +70mV is applied, in the middle of which the system is subjected to a brief (5 μ s) pulse voltage of +500mV. The “forward” read voltage is then followed by a “reverse” read voltage of -70mV, and this cycle of +70mV forward read/500mV pulse/+70mV forward read /-70mV reverse read is repeated until the entire polymeric analyte has passed through the nanopore. One significant advantage of the ratchet protocol is that the electrolyte distribution is replenished during each forward read-reverse read cycle.

[00149] The bottom panel of FIG. 12 illustrates how the directionality of polymer translocation changes during ratcheting. In this embodiment, each unit of the polymeric analyte includes two identical reporter codes (e.g., **1210A** and **1210B**) separated by a translocation control element, e.g., TCE **1215**. As illustrated in FIG. 12, application of the +70mM forward run voltage results in polymer movement through the pore from the cis side of the membrane to the trans side until reporter code **1210A** is arrested in the pore by the pause in translocation induced by TCE **1215**. The change in current through the pore due to arrested reporter code **1210A** is read as signal “L1+”. Application of the 500mV pulse forces TCE **1215** and reporter code **1210B** through the pore. Following the 5 μ s pulse, the read voltage returns to +70mV, whereupon the next reporter code in series, **1220A**, is arrested in the pore by its corresponding TCE **1225**. The change in current flow through the pore due to reporter code **1220A** is read as signal “L2+”. Next, the voltage is reversed by application of the reverse read voltage of -70mM to the system, resulting in reversal of the direction of polymer translocation. During this period, reporter code **1220A** is pushed back through the pore to the cis side of the membrane. Translocation proceeds until TCE **1215** encounters the pore, whereupon translocation is arrested, positioning reporter code **1210B** in the pore to generate a change in current measured as level “L1-”. Next, the voltage is returned to the +70mM forward run voltage, whereupon the direction of polymer translocation is reversed back to the cis-to-trans direction, until an arrest occurs due to the ITC effect of TCE **1225**, which positions reporter code **1220A** in the pore. The resulting change in current is read as level “L2+”. During this ratcheting protocol, the reporter code characterizing each unit in the polymer is read three times (reporter code “A” gets read twice and reporter code “B” gets

read once). This redundancy provides a means of quality control for the sequence reads and improves the accuracy of the resulting sequence data. Insertion and deletion errors can readily be identified as deviations from the expected pattern, e.g., L1+/L2+/L1-/L2+.

[00150] Although the ratcheting pattern depicted in FIG. 12 shows the pulse voltage being applied during the “middle” of the forward read voltage, other patterns of pulsing are contemplated by the present invention. For example, in some embodiments, the pulse may be applied prior to application of the forward read voltage, while in other embodiments, the pulse may be applied at the end of the forward read voltage, just prior to application of the reverse read voltage.

[00151] In other embodiments, ratcheting provides means for compensating, or correcting, for one or both of current depletion due to pulsing and asymmetry in the resistance of the different reporter codes. For example, in some embodiments, the reverse read voltage can be increased to compensate for the current loss due to the pulses applied during the forward read voltage. The percent increase in the reverse read voltage can also be adjusted to balance the current when the different reporter codes have different intrinsic resistances.

[00152] In other variations of the ratcheting scheme, the sequence of forward, pulse, and reverse voltages can be altered. For example, in one embodiment, a ratcheting cycle could be run as follows: (forward read voltage), (forward read voltage), (reverse read voltage). In another embodiment, a ratcheting cycle could be run as follows: (forward read)_n(reverse read)_n in which “n” represents the total number of monomeric units in the polymer being measured by the nanopore.

[00153] In a related idea, “flossing” was proposed whereby DNA may be read in a nanopore along its full length, stopped and then upon reversing the voltage polarity could be read in the other direction (see, e.g., Kasianowicz, John J. “Nanopores: Flossing with DNA.” *Nature Materials* 3, no. 6 (2004): 355–56. <https://doi.org/10.1038/nmat1143>). This is a less efficient method in an array because the DNA polymers are not captured or stopped synchronously whereas ratcheting is a continuous forward process on these time scales.

SBX-Based Diagnostic Methods and Kits

[00154] In another aspect, the present invention discloses methods and kits for the detection and diagnostics of genetic alterations/mutations in a target sample, which may be a solid tissue or a bodily fluid. The genetic alterations may be either germline or somatic mutations. The invention may be used for detection and diagnostics related to cancer, auto-

immune disease, organ transplant rejection, genetic fetal abnormalities, pathogens, and other suitable conditions.

EXAMPLES

Material and Methods

[00155] The following materials, having the abbreviations as indicated, were obtained from the mentioned sources in the United States, unless otherwise indicated. 2-Phenyl-1,3-dioxan-5-ol, TBDPS-Cl (t-butyldiphenylchlorosilane), DMAP (4-dimethylaminopyridine), (R)-(+)-glycidol, (+)-2,3-O-Isopropylidene-L-threitol, isosorbide, 4,4'-Bis(hydroxymethyl)-2,2'-bipyridine, TBTA (Tris[(1-benzyl-1H-1,2,3-triazol-4-yl)methyl]amine) from TCI America (Portland, OR). NaH (sodium hydride), MeOH (methanol), toluene, THF (tetrahydrofuran), TBAF (tetrabutylammonium fluoride), DCM (dichloromethane), HCl (concentrated hydrochloric acid), DMSO (dimethylsulfoxide), Na ascorbate (sodium ascorbate), sodium bicarbonate, copper sulfate, dimethyl propargylmalonate, lithium borohydride, and acetic acid were obtained from Sigma-Aldrich (St. Louis, MO). DMT-Cl (4,4'-dimethoxytrityl chloride) and PPA-Cl (N,N-diisopropylamino cyanoethyl phosphoramidic chloride) from ChemGenes Corporation (Wilmington, MA). TEA (triethylamine), hexanes, ethyl acetate, EDTA (ethylenediaminetetraacetic acid), diethyl ether from EMD Millipore (Billerica, MA). m-PEG4-Tos was made from m-PEG4-OH (Cat. No. BP-23742). Furo[3,2-c]pyridin-4(5h)-one (Combi-Blocks, San Diego, CA).

[00156] High performance liquid chromatography (HPLC) was performed on a ProStar Helix™ HPLC system from Agilent Technologies, Inc. (Santa Clara, CA) consisting of two pumps (ProStar 210 Solvent Delivery Modules) with 10ml titanium pump heads, a column oven (ProStar 510 Air Oven), a UV detector (ProStar 320 UV/Vis Detector) set at 292nm. The system is controlled by Star Chromatography Workstation Software (version 6.41). The column used was a Cadenza Guard Column System CD-C18 (2.0mm x 5mm) both from Imtakt USA (Portland, OR). The buffers used are Buffer A (100mM triethylammonium acetate, pH 7.0) and Buffer B (100mM triethylammonium acetate, pH 7.0 with 95% by volume acetonitrile). Automated solid phase phosphoramidite synthesis was done on a MerMade™ 12 synthesizer (Bioautomation Corp, Plano, TX). Synthesis solutions for the MerMade™ were purchased from Glen Research (Sterling, VA).

Example 1

Synthesis of DMT Phosphoramidite of racemic 2-(3,6-Dioxaheptyloxy)-1,3-propanediol


Pendant Code 2-glyceryl PEG-2 Phosphoramidite [racemate]

[00157] 2-Phenyl-1,3-dioxan-5-ol (1, 2.7 g, 15 mmol) was dissolved in 30 mL anhydrous THF. Sodium hydride (1.08 g, 27 mmol) was added to generate alkoxide. When the bubbling ceased, mPEG4-Tos (4.94 g, 18 mmol) was dissolved in 10 mL THF and added portion-wise. The reaction was brought to 40°C and incubated with stirring for 3h, then allowed to come down to room temperature overnight. Excess NaH was quenched with 1 mL MeOH, then diluted with water and extracted with DCM. The combined organic layers were concentrated under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 2 in 73% yield.

[00158] Benzylidene protected 2b (3.05 g, 10.8 mmol) was dissolved in 10 mL MeOH and HCl (0.2 mL, 2.3 mmol) was added. The solution was incubated for 20 minutes, then neutralized with sodium bicarbonate (200 mg) and dried under reduced pressure. The residue was resuspended in DCM and purified by flash chromatography to afford diol 3 in 72% yield.

[00159] Diol 3 (1.52 g, 7.8 mmol) was dissolved in 20 mL DCM and TEA (2.17 mL, 15.6 mmol). A solution of DMT-Cl (1.85 g, 5.46 mmol) in 10 mL DCM was added portion-wise over 90 minutes to maximize monotritylation. MeOH (1 mL) was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford the mono-trityl product 4 in 48% yield as well as recovered starting diol.

[00160] Monotrityl 4 (1.84 g, 3.7 mmol) was dissolved in 10 mL DCM and TEA (1.03 mL 7.4 mmol). PPA-Cl (1.05 g, 4.4 mmol) was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 5 (2.03g, 2.9 mmol) was afforded in 79% yield and confirmed by ¹H and ³¹P NMR.



Example 2

Synthesis of DMT Phosphoramidite of enantiomer 2-(3,6-Dioxaheptyloxy)-1,3-propanediol (12b) (Pendant Code 2-glyceryl PEG-2 Phosphoramidite [enantiomer])

[00161] 2,3-Isopropylidene-*sn*-glycerol 6 was dissolved in anhydrous DCM and TEA. DMAP and TBDPS-Cl were added. The reaction was extracted from water with DCM and purified by flash chromatography to afford product 7.

[00162] Silyl ether 7 was dissolved in MeOH and HCl was added. The solution was incubated 20 minutes, then neutralized with sodium bicarbonate and dried under reduced pressure. The residue was resuspended in DCM and purified by flash chromatography to afford diol 8.


[00163] Diol 8 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford the mono-trityl product 9.

[00164] Secondary alcohol 9 was dissolved in anhydrous THF. Sodium hydride was

added to generate alkoxide. When the bubbling ceased, mPEG4-Tos was dissolved in THF and added portion-wise. The reaction was brought to 40°C and incubated with stirring for 3h, then allowed to come down to room temperature overnight. Excess NaH was quenched with 1 mL MeOH, then diluted with water and extracted with DCM. The combined organic layers were dried under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 10.

[00165] mPEG4 ether 10b was resuspended in THF and TBAF was added. The reaction was concentrated under reduced pressure and purified by flash chromatography to afford 11.

[00166] DMT PEG4 alcohol 11b was dissolved in DCM and TEA. PPA-Cl was added and the reaction incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 12 was isolated and confirmed by ¹H and ³¹P NMR.



Example 3

Synthesis of DMT Phosphoramidite of enantiomer 1-(3,6-Dioxaheptyloxy)-2,3-propanediol
(16)


Pendant Code 1-glyceryl PEG-2 Phosphoramidite

[00167] 2,3-Isopropylidene-*sn*-glycerol 6 was dissolved in anhydrous THF. Sodium hydride was added to generate alkoxide. When the bubbling ceased, mPEG2-Tos (Broadpharm Cat. No. BP-2-982) was dissolved in THF and added portion-wise. The reaction was incubated for 24 hours. Excess NaH was quenched with MeOH, then diluted with water and extracted with DCM. The combined organic layers were concentrated under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 13.

[00168] The PEG2 product 13 was dissolved in MeOH and HCl was added. The solution was incubated 20 minutes, then neutralized with sodium bicarbonate and dried under reduced pressure. The residue was resuspended in DCM and purified by flash chromatography to afford diol 14.


[00169] Diol 14 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford the mono-trityl product 15.

[00170] Mono DMT 15 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 16 was isolated and confirmed by ¹H and ³¹P NMR.



Pendant Code 1-glyceryl heterocycle Phosphoramidite

[00173] Secondary alcohol 19 was dissolved in DCM and TEA. PPA-Cl was added and the reaction and incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 20 was isolated and confirmed by ¹H and ³¹P NMR.




Example 5

Synthesis of In-line Code DMT Phosphoramidite of Isosorbide (23)

Bis-secondary alcohol code backbone

[00174] Isosorbide 21 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford monotrityl 22.

[00175] Monotrityl 22 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 23 was isolated and confirmed by ¹H and ³¹P NMR.




Example 6

Synthesis of In-line code DMT Phosphoramidite of Bipyridyl (26)

Bis-Primary alcohol code backbone

[00176] 4,4'-Bis(hydroxymethyl)-2,2'-bipyridine 24 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford monotrityl 25.

[00177] Monotrityl 25 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 26 was isolated and confirmed by ¹H and ³¹P NMR.



Example 7

Synthesis of Pendant Code DMT Phosphoramidite of 2-alkyl triazole


PEG-2 1,3-propane diol (31a). Pendant Triazole PEG Code

[00178] Dimethyl propargylmalonate 27 was added dropwise to a cold suspension of lithium borohydride in diethyl ether. The reaction was warmed to room temperature and incubated overnight. The reaction was quenched with methanol, then water, then acetic acid. The solution was extracted with ether and the combined organic layers were concentrated under reduced pressure. The crude material was purified by flash chromatography to afford diol 28.

[00179] Diol 28 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford DMT alcohol 29.

[00180] DMT alcohol 29 was dissolved in DMSO and azide (Cat. No. BP-20988 Broadpharm) was added. Separately, TBTA was dissolved in DMSO and sodium ascorbate and copper sulfate were combined. The TBTA solution was added to the alkyne/azide solution in portions with stirring. After 45 minutes of incubation, the reaction was quenched with EDTA. The solution was diluted with water and extracted with ethyl acetate, then the organic layers were concentrated under reduced pressure and purified by flash chromatography to afford 30.

[00181] 1,2,3-Triazole 30a was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 31 was isolated and confirmed by ¹H and ³¹P NMR.



Example 8

Synthesis of Pendant PEG-2, PEG-4 and PEG-6 Phosphoramidites [isomerically pure]


(35a)

[00182] 1-O-TBDPS-3-O-DMTr-propane-1,2,3-triol 9 (from Example 2) was dissolved in anhydrous THF. Sodium hydride was added to generate alkoxide. When the bubbling ceased, tosylate (prepared via tosylation of Cat. No. BP-21657, Broadpharm) was added portion-wise. The reaction was incubated with stirring for 48h. Excess NaH was quenched with water, then the solution was transferred to a separatory funnel and extracted with ethyl acetate. The combined organic layers were washed with brine, dried with sodium sulfate, filtered and concentrated under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 32.

[00183] Alkyne 32a was resuspended in THF and TBAF was added. The reaction was concentrated under reduced pressure and purified by flash chromatography to afford 33.

[00184] To a solution of alkyne 33a in DMSO was added 2-azidoethyl acetate. In a separate vial, dissolve sodium ascorbate in water and add DMSO followed by 1M CuSO₄, to prepare catalyst mixture. Add catalyst mixture to solution of alkyne/azide dropwise over 10 minutes. Upon completion, quench with 0.5M EDTA and stir for 15 minutes. Dilute with water and extract with ethyl acetate three times. Wash combined organic extractions with brine and dry over sodium sulfate. The residue was resuspended in toluene and purified by flash chromatography to afford 34a.

[00185] Triazole 34a was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 35 was isolated and confirmed by ¹H and ³¹P NMR.




Example 9

Synthesis of PEG-Based Reporter Codes (37b)

[00186] To a solution of product 33 in DMSO was added 2-(acetoxymethyl)-2-(azidomethyl)propane-1,3-diyl diacetate (prepared by dissolving 2-(Bromomethyl)-2-(hydroxymethyl)-1,3-propanediol in DMF and subsequent addition of NaN₃. The reaction was incubated at 110 °C, concentrated, and purified by flash chromatography. Upon isolation of the product, the residue was reacted with acetic anhydride and purified). In a separate vial, dissolve sodium ascorbate in water and add DMSO followed by 1M CuSO₄, to prepare catalyst mixture. Add catalyst mixture to solution of alkyne/azide dropwise over 10 minutes. Upon completion, quench with 0.5M EDTA and stir for 15 minutes. Dilute with water and extract with ethyl acetate three times. Wash combined organic extractions with brine and dry over sodium sulfate. The residue was resuspended in toluene and purified by flash chromatography to afford 36.

[00187] Primary alcohol 36b was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 37 was isolated and confirmed by ¹H and ³¹P NMR.




Example 10

Synthesis of PEG-Based Reporter Codes (40)

[00188] 1-O-TBDPS-3-O-DMTr-propane-1,2,3-triol **9** (from Example 2) was dissolved in anhydrous THF. Sodium hydride was added to generate alkoxide. When the bubbling ceased, tosylate (prepared by sequential tosylation and silyl-protection of Cat. No.BP-21036, Broadpharm) was dissolved in THF and added portion-wise. The reaction was incubated with stirring overnight. Excess NaH was quenched with water and extracted with DCM. The combined organic layers were dried under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford **38**.

[00189] Bis-silyl ether **38** was resuspended in THF and TBAF was added. The reaction was concentrated under reduced pressure and purified by flash chromatography. The purified material was resuspended in DCM and TEA was added. BzCl was added dropwise. The reaction was stirred at room temperature until complete. The reaction was concentrated under reduced pressure and purified by flash chromatography to afford alcohol **39**.

[00190] Alcohol 39 was dissolved in DCM and TEA. PPA-Cl was added and the reaction and incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 40 was isolated and confirmed by ^1H and ^{31}P NMR.



Example 11

Synthesis of bis-PEG-Based Reporter Codes (45a-d and 47e-i)

[00191] O,O'-Benzylidenepentaerythritol (41, Cat. No. B2682, TCI) was dissolved in anhydrous THF. Sodium hydride was added to generate alkoxide. When the bubbling ceased, tosylate (prepared via tosylation of Cat. No. BP-21397, Broadpharm or Cat. No. BP-21657, Broadpharm) was dissolved in THF and added portion-wise. The reaction was incubated with stirring overnight. Excess NaH was quenched with water and extracted with DCM. The combined organic layers were dried under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 42a-h.

[00192] Products 42a-h were dissolved in MeOH and HCl was added. The reaction was incubated at room temperature overnight, then neutralized with sodium bicarbonate. It was concentrated under reduced pressure and purified by flash chromatography to afford 43a-h.

[00193] Products 43a-h were dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. The reaction was dried under reduced pressure. The residue

was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford 44a-h.

[00194] Products 44a-d were dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography to afford phosphoramidites 45a-d.

[00195] To a solution of product 44 (e-h) in DMSO was added 2-azidoethyl acetate. In a separate vial, dissolve sodium ascorbate in water and add DMSO followed by 1M CuSO₄, to prepare catalyst mixture. Add catalyst mixture to solution of alkyne/azide dropwise over 10 minutes. Upon completion, quench with 0.5M EDTA and stir for 15 minutes. Dilute with water and extract with ethyl acetate three times. Wash combined organic extractions with brine and dry over sodium sulfate. The residue was resuspended in toluene and purified by flash chromatography to afford 46e-h.

[00196] Products 46e-h were dissolved in DCM and TEA. PPA-Cl was added and the reaction and incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography to afford phosphoramidites 47e-h.



Example 12


Synthesis and Testing of PEG-Based Reporter Codes (52)

[00197] 2,2-Bis(bromomethyl)-1,3-propanediol (48, Cat. No. D1808, TCI) was dissolved in DMF and NaN₃ was added. The reaction was incubated at 110 °C, concentrated, and purified by flash chromatography to afford product 49.

[00198] To a solution of 49 in DMSO was added 2-(2-(prop-2-yn-1-yloxy)ethoxy)ethyl benzoate (prepared by benzylation of the commercial alcohol precursor). In a separate vial, dissolve sodium ascorbate in water and add DMSO followed by 1M CuSO₄, to prepare catalyst mixture. Add catalyst mixture to solution of alkyne/azide dropwise over 10 minutes. Upon completion, quench with 0.5M EDTA and stir for 15 minutes. Dilute with water and extract with ethyl acetate three times. Wash combined organic extractions with brine and dry over sodium sulfate. The residue was resuspended in toluene and purified by flash chromatography to afford 50.

[00199] Product 50 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. The reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford 51.

[00200] Product 51 dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography to afford phosphoramidite 52.



Example 13

Synthesis and Testing of PEG-Based Reporter Codes (62)

[00201] 2-(Bromomethyl)-2-(hydroxymethyl)-1,3-propanediol (53, Cat. No. B4057, TCI) was dissolved in DMF and NaN_3 was added. The reaction was incubated at 110 °C, concentrated, and purified by flash chromatography to afford product 54.

[00202] Product 54 was dissolved in DCM and TEA. Benzoyl chloride was added. The solution was incubated overnight at room temperature. The reaction was extracted from water with DCM and purified by flash chromatography to afford product 55, the bis-Bz which was separated from any mono- or tri-protected species and divided.

[00203] A portion of product 55 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. MeOH was added and the reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford the tritylated 56.

[00204] 41 was dissolved in anhydrous THF. Sodium hydride was added to generate

alkoxide. When the bubbling ceased, propargyl bromide was dissolved in THF and added portion-wise. Excess NaH was quenched with 1 mL MeOH, then diluted with water and extracted with DCM. The combined organic layers were dried under reduced pressure. The residue was resuspended in toluene, separated from remaining salts, and purified by flash chromatography to afford 57.


[00205] Product 57 was dissolved in MeOH and HCl was added. The reaction was incubated at room temperature overnight, then neutralized with sodium bicarbonate. It was concentrated under reduced pressure and purified by flash chromatography to afford 58.

[00206] Product 58 was dissolved in DCM and TEA. Benzoyl chloride was added and the reaction was incubated 60 minutes. The reaction was extracted from water with DCM and purified by flash chromatography to afford product 59.

[00207] Products 56 and 59 were dissolved in 9:1 DMSO:H₂O. A solution of TBTA, sodium ascorbate and copper sulfate was added and the reaction was incubated 60 minutes. The reaction was extracted from water with DCM and purified by flash chromatography to afford product 60.

[00208] Products 60 and 55 were dissolved in 9:1 DMSO:H₂O. A solution of TBTA, sodium ascorbate and copper sulfate was added and the reaction was incubated 60 minutes. The reaction was extracted from water with DCM and purified by flash chromatography to afford product 61.

[00209] Product 61 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 62 was isolated and confirmed by ¹H and ³¹P NMR.



Example 14

Synthesis DMT Phosphoramidite of enantiomer (9S,10S)-2,5,8,11,14,17-hexaoxaoctadecane-9,10-diol (Pendant Code C2 Bis-PEG-2 Phosphoramidite [enantiomer]) (67)

[00210] A solution of (+)-2,3-O-Isopropylidene-L-threitol 63 in anhydrous DMF was slowly added to a mixture of NaH in anhydrous DMF (Note: vigorous evolution of H₂ gas). When the bubbling ceased, mPEG2-Tos (Broadpharm Cat. No. BP-20983) was dissolved in DMF and added portion-wise to stir at ambient temperature overnight. The reaction mixture was poured over water, extracted with ethyl acetate and purified by flash chromatography to


afford 64.

[00211] Product 64 was dissolved in MeOH and HCl was added. The solution was incubated 20 minutes, then neutralized with sodium bicarbonate and dried under reduced pressure. The residue was resuspended in ethyl acetate and purified by flash chromatography to afford 65.

[00212] Product 65 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. The reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford 66.

[00213] Product 66 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 67 was confirmed by ¹H and ³¹P NMR.

[00214]



Example 15


Synthesis of 1-O-DMT-3-O-PPA-2,2-bis(Me-O-mPEG4)-propane (72)

[00215] To a solution of pentaerythritol (68, Cat. No. P0039, TCI) in DMF was added p-toluenesulfonic acid. The reaction was neutralized with triethylamine, concentrated and purified by flash chromatography to afford 69.

[00216] Product 69 was added to a stirring solution of EDC-HCl, DMAP and levulinic acid in THF and stirred overnight at ambient temperature. The solution was concentrated and purified by flash chromatography to afford 70.

[00217] Product 70 was dissolved in DCM and TEA. A solution of DMT-Cl in DCM was added portion-wise. The reaction was dried under reduced pressure. The residue was resuspended in toluene and separated from the salts, then purified by flash chromatography to afford 71.

[00218] Product 71 was dissolved in DCM and TEA. PPA-Cl was added and the reaction was incubated 15 minutes. The reaction was dried down under reduced pressure and resuspended in toluene with 1% TEA, then purified by flash chromatography. Phosphoramidite 72 was confirmed by ¹H and ³¹P NMR.



Example 16

Synthesis and Testing of PEG-Based Reporter Codes

[00219] In this Example, reporter codes were synthesized with PEG-based phosphoramidites; notably, these codes do not contain nucleotides. Four exemplary reporter codes are set forth in Table 5.

Table 5

Code designator	sequence
-----------------	----------

L1	DDDLLLL
L2	DXXX3LL3XXLL
L3	DXLL44XXXLLLL
L4	DXXL4444LLXXLL

Key: D = PEG6; X = PEG3; L = C2; 3 = clicked PEG-2; 4 = pendant PEG-4

[00220] The level discrimination (i.e. distinguishable electronic signal) and translocation time of each code was assessed by synthesizing an 100mer Xpandomer copy of a sequence derived from the HIV-2 genome that incorporates XNTPs in which each of the four XNTPs includes a unique code from the group set forth in Table 4. Xpandomer synthesis, processing, and nanopore sequence analysis were carried out as described in Applicants' PCT patent application no. PCT/US18/67763, which is herein incorporated by reference in its entirety. As a control, an Xpandomer copy of the same HIV-2 sequence incorporating different, known codes was sequenced in parallel. A representative trace illustrating the level discrimination and translocation time of each code is shown in Fig. 13A (control – old codes) and 13B (test – new, PEG-based controls).

[00221] To assess the accuracy of the Xpandomer sequence information, sequence data was analyzed by histogram display of the population of sequence reads from the SBX reactions. The analysis software aligns each sequence read to the sequence of the template and trims the extent of the sequence at the end of the reads that does not align with the correct template sequence. Representative histograms of SBX sequencing of the 100mer template are presented in FIG. 14A (control) and FIG. 14B (new, PEG-based codes). Notably, Xpandomers incorporating the nucleotide-free PEG-based codes yielded highly accurate sequence reads of this template.

Example 17Translocation control with pendant PEG-based TCEs – 2 Code Levels

[00222] In this Example, translocation control with a TCE incorporating a pendant PEG phosphoramidite was assessed by using the SBX protocol to sequence a simple 60mer template consisting of TG dinucleotide repeats. Both XATP and XCTP substrates were designed to incorporate the following TCE: Y2222222222255, in which “Y” represents the symmetric phosphoramidite brancher; “2” represents pendant PEG2; and “5” represents benzofuran. The XATP substrate was designed to incorporate the following reporter code: DDDDDDL L L L L, in which “D” represents PEG6 and “L” represents C2. The XCTP substrate was designed to incorporate the following reporter code: DDDDX X 44 X X D L, in which “X” represents PEG3 and “4” represents pendant PEG4.

[00223] To produce Xpandomer copies of the 60mer template, primer extension reactions were conducted using 4pm of an extension oligonucleotide and 250pm of each XNTP. The 10μL extension reaction included the following reagents: 50mM TrisCl, pH 8.84, 200mM NH₄OAc, 20% PEG8K, 5% NMS, 0.75nmol polyphosphate PP-60.20, 2μg SSB, 0.5M urea, 5mM PEM additive (suitable Polymerase Enhancing Molecules are disclosed in Applicants’ pending PCT patent application no. PCT/US18/67763, which is herein incorporated by reference in its entirety), and 1.2μg purified recombinant DNA polymerase (suitable engineered variants of DPO4 polymerase are disclosed in Applicants’ PCT patent application no.s WO2017/087281, PCT/US2018/030972, and PCT/US1864794, which are herein incorporated by reference in their entireties) The extension reaction was run for 30 minutes at 42□ C.

[00224] Xpandomer products of the extension reactions were next sequenced using the SBX protocol. Briefly, the constrained Xpandomer products were cleaved to generate linearized Xpandomers. This was accomplished by first quenching the extension reaction and subjecting the Xpandomers to amine modification with 2M succinic anhydride. The phosphoramidate bonds of the Xpandomers were then cleaved by treating the sample with 11.7M DCl for 30 minutes at 23□ C. Linearized Xpandomers were purified by ethanol precipitation and resuspended in a buffer supplemented with 34% ACN and 15% DMF.

[00225] For sequencing, Xpandomers were added to a sample buffer of 2.8M NH₄Cl, 1.2M GuanCl, 20mM NaHex, 10% DMF, 2mM EDTA, and 20mM HEPES pH 7.4. Protein nanopores were prepared by inserting α-hemolysin into a DPhPE/hexadecane bilayer member

in a buffer containing 2 M NH₄Cl and 100mM HEPES, pH 7.4. This experiment used buffers of 0.4M NH₄Cl, 600mM GuanCl, and 100mM HEPES, pH 7.4 in the cis well and 2M NH₄Cl and 100mM HEPES, pH 7.4 in the trans well of the detection system. The Xpandomer sample was heated to 70° C for 2 minutes, cooled completely, followed by addition of 2μL of the sample to the cis well. The voltage parameters run were as follows: 60mV/300mV/10μs/2ms (read voltage/pulse voltage/pulse voltage duration/pulse frequency). Data were acquired via Labview acquisition software. A representative trace from this run is shown in Fig. 15.

[00226] As shown by the level numbers superimposed above the trace in Fig. 15, all reporter codes were read correctly by the detection system in this experiment, demonstrating 100% accuracy. The absence of deletion or insertion errors validates the efficacy of pendant PEG-based TCEs as structures enabling tight regulation of Xpandomer translocation. Significantly, with the pendant PEG-based TCE used in this experiment, transitions between the two code levels was observed following a single voltage pulse. The ability to transition between sequential reporter codes of the Xpandomer with single voltage pulses is a significant advancement in the art and enables a much higher sequencing throughput.

Example 18

Sequencing by Expansion with pendant PEG-based TCEs– 4 Code Levels

[00227] In this Example, translocation control with a TCE incorporating a pendant PEG phosphoramidite was assessed by using the SBX protocol to sequence a 60mer template consisting of repeats of the sequence, CATG. All XNTP substrates were designed to incorporate the following TCE: Y444444444444455, in which “Y” represents the symmetric phosphoramidite brancher; “4” represents pendant PEG4; and “5” represents benzofuran. The XATP substrate was designed to incorporate the following reporter code: DDDDDDLLDX; the XCTP substrate was designed to incorporate the following reporter code: DDDDDDLLLL; the XTTP substrate was designed to incorporate the following reporter code: DDDDD44LXXX; and the XGTP substrate was designed to incorporate the following reporter code: DDDDXL444444XLLLL, in which “D” represents PEG6, “L” represents C2, “X” represents PEG3 and “4” represents pendant PEG4.

[00228] To produce Xpandomer copies of the 60mer template, primer extension

reactions were conducted using 4pm of an extension oligonucleotide and 1000pm of each XNTP and. The 10 μ L extension reaction included the following reagents: 50mM TrisCl, pH 8.84, 200mM NH₄OAc, 20% PEG8K, 10% NMP, 3nmol polyphosphate PP-60.20, 2 μ g SSB, 1M urea, 10mM PEM additive, and 1.8 μ g purified recombinant DNA polymerase. The extension reaction was run for 30 minutes at 37 $\square$ C.

[00229] Xpandomer products of the extension reactions were next sequenced using the SBX protocol. Briefly, the constrained Xpandomer products were cleaved to generate linearized Xpandomers. This was accomplished by first quenching the extension reaction and subjecting the Xpandomers to amine modification with 2M succinic anhydride. The phosphoramidate bonds of the Xpandomers were then cleaved by treating the sample with 11.7M DCl for 30 minutes at 23 $\square$ C. Linearized Xpandomers were purified by ethanol precipitation and resuspended in a buffer supplemented with 34% ACN and 15% DMF.

[00230] For sequencing, Xpandomers were added to a sample buffer of 0.8M NH₄Cl, 1.2M GuanCl, and 200mM HEPES, pH 7.4. Protein nanopores were prepared by inserting α -hemolysin into a DPhPE/hexadecane bilayer member in a buffer containing 2 M NH₄Cl and 100 mM HEPES, pH 7.4. This experiment used buffers of 0.4M NH₄Cl, 600mM GuanCl, and 100mM HEPES, pH 7.4 in the cis well and 2M NH₄Cl and 100mM HEPES, pH 7.4 in the trans well. The Xpandomer sample was heated to 70 $\square$ C for 2 minutes, cooled completely, followed by addition of 2 μ L of the sample to the cis well. The voltage parameters run were as follows: 70mV/650mV/6 μ s/1.5ms (read voltage/pulse voltage/pulse voltage duration/pulse frequency). Data were acquired via Labview acquisition software. A representative trace from this run is shown in Fig. 16

[00231] The level numbers superimposed above the trace in Fig. 16 correspond to the signals generated by the XCTP code (L1), the XTTP code (L2), the XATP code (L3) and the XGTP code (L4). In this experiment, all reporter codes were read correctly by the detection system, demonstrating 100% accuracy. The absence of deletion and insertion errors underscores the efficacy of pendant PEG-based TCEs in transiently pausing reporter codes in the pore channel to produce accurate signal reads during the “read voltage” and allowing resumption of translocation during the “pulse voltage”. Again, with the pendant PEG-based TCE used in this experiment, transitions between sequential reporter codes was achieved with single voltage pulses. Of note, the same level of accuracy was observed in separate runs of aliquots the of the same Xpandomer sample using the following voltage parameters: 65/700/6/1.5; 65/650/6/1.5; 60/650/6/1.5 and 60/600/6/1.5. It was further observed that the throughput of codes reads can be influenced by various voltage parameters.

Example 19Sequencing by Expansion of a Complex Template with pendant PEG-based TCEs

[00232] In this Example, translocation control with a TCE incorporating a pendant PEG phosphoramidite was assessed by using the SBX protocol to sequence a complex 100mer template. Each XNTP substrate was synthesized with the following TCE: Y22222222222255, in which “Y” represents the symmetric phosphoramidite brancher; “2” represents pendant PEG2; and “5” represents benzofuran. The XNTP substrates were synthesized with the following reporter codes: (XC)DDDDDDLLLL, in which “D” represents PEG6 and “L” represents C2; (XT)DDDDDD44LDX, in which “X” represents PEG3 and “4” represents pendant PEG4; (XA)DDDDXX44XXDL; and (XG)DDDDXXL.

[00233] To produce Xpandomer copies of the 100mer template solid-state primer extension reactions were conducted using 1pmol of XATP and XCTP and 1.5pmol of XGTP and XTTP (solid-state Xpandomer synthesis in which the extension oligo is covalently bound to a chip substrate is described in Applicants’ provisional patent application no. 62/826,805, which is herein incorporated by reference in its entirety). The 50μL extension reaction included the following reagents: 50mM TrisCl, pH 8.84, 200mM NH4OAc, 20% PEG8K, 10% NMP, 15pmol polyphosphate PP-60.20, 10μg SSB, 1M urea, 10mM PEM additive and 9μg purified recombinant DNA polymerase. The extension reaction was run for 30 minutes at 37□ C.

[00234] Xpandomer products were next sequenced using the SBX protocol. Briefly, the constrained Xpandomer products were cleaved to generate linearized Xpandomers. This was accomplished by first quenching the extension reaction and subjecting the Xpandomers to amine modification with succinic anhydride. The phosphoramidate bonds of the Xpandomers were then cleaved by treating the sample with 7.5M DCl for 30 minutes at 23□ C. Linearized Xpandomers were released from the chip substrate by photocleavage of the extension oligonucleotide and recovered in elution buffer supplemented with 15% ACN and 5% DMSO (20% final solvent).

[00235] For sequencing, Xpandomers were added to a sample buffer of 0.8M NH4Cl, 1.2M GuCl, 200mM HEPES; pH 7.4. Protein nanopores were prepared by inserting α-hemolysin into a DPhPE/hexadecane bilayer member in a buffer of 2 M NH4Cl and 100 mM HEPES, pH 7.4. The cis well was perfused with buffer containing 0.4M NH4Cl, 600mM

GuanCl, 100mM HEPES; pH 7.4 and the trans well was perfused with a buffer containing 2M NH₄Cl, 100mM HEPES; pH 7.4. The Xpandomer sample was heated to 70° C for 2 minutes, cooled completely and vortexed, then a 2 µL aliquot was added to the cis well. The voltage parameters were run as follows: 60mV/600mV/6µs/1.5ms (read voltage/pulse voltage/pulse voltage duration/pulse frequency). Data were acquired via Labview acquisition software. A representative trace from this run is shown in Fig. 17.

[00236] As shown in Fig. 17, the sequence of all reporter codes was read correctly by the detection system in this experiment, demonstrating 100% accuracy. The absence of deletion or insertion errors again validates the efficacy of pendant PEG-based TCEs as structures enabling highly reliable sequence reads of a polymeric template. Significantly, runs of homopolymers were accurately sequenced under these conditions (see, e.g., the sequence of 4x L1 codes near the start of the trace shown in Fig. 14), thereby highlighting the accuracy of SBX with pendant PEG-based TCEs.

Example 20

Sequencing by Expansion of a Complex 222mer Template with pendant PEG-based TCEs

[00237] In this Example, translocation control with pendant PEG-based TCEs was assessed by using the SBX protocol to sequence a complex 222mer template. Each XNTP substrate was synthesized to include the following TCE: Y44444444444455, in which “Y” represents the symmetric phosphoramidite brancher; “4” represents pendant PEG-4; and “5” represents benzofuran. The XNTP substrates were synthesized with the following reporter codes: (XC)DDDDDDLLLDX; (XT)DDDDDD44LXXX; (XA)DDDDDD444LLDX; and (XG)DDDDXXL444444XL LLL, in which “D” represents PEG-6, “L” represents C2, “X” represents PEG-3, and “4” represents pendant PEG-4.

[00238] To produce Xpandomer copies of the 222mer template, solid-state primer extension reactions were conducted using 1.25pmol of each XNTP, 10pmol template and 20pmol E-oligo primer (solid-state Xpandomer synthesis in which the extension oligo is covalently bound to a chip substrate is described in Applicants’ provisional patent application no. 62/826,805, which is herein incorporated by reference in its entirety). The 50µL extension reaction included the following reagents: 50mM TrisCl, pH 8.84, 200mM NH₄OAc, 20% PEG8K, 8% NMP, 15nmol polyphosphate PP-60.20, 10µg SSB, 1M urea, 5mM PEM additive and 9µg purified recombinant DNA polymerase. The extension reaction was run for

30 minutes at 37 ° C.

[00239] Xpandomer products were next sequenced using the SBX protocol. Briefly, the constrained Xpandomer products were washed in buffer B.001 (1% Tween-20/3% SDS/5mM HEPES, pH 8.0/100mM NaPO₄/15% DMF) and cleaved to generate linearized Xpandomer by adding 200μl buffer C.001 (7.5M DCl) and incubating for 30 minutes at 23 °C. The sample was then neutralized by adding 1000μl buffer B.001. The Xpandomer sample was then subjected to amine modification by adding 666μmol succinic anhydride and incubating for 5 minutes at 23 °C. The sample was then washed in buffer D.094 (50% ACN) and the Xpandomers were released from the substrate by photocleavage and stored in buffer AG497 (0.8M NH₄Cl/1.2M GuanCl/200mM HEPES, pH 7.4)

[00240] Protein nanopores were prepared by inserting α-hemolysin into a DPhPE/hexadecane bilayer member in a buffer of 2 M NH₄Cl and 100 mM HEPES, pH 7.4. The cis well was perfused with buffer containing 0.4M NH₄Cl, 600mM GuanCl, 100mM HEPES; pH 7.4 and the trans well was perfused with a buffer containing 2M NH₄Cl, 100mM HEPES; pH 7.4. The Xpandomer sample was heated to 70° C for 2 minutes, cooled completely and vortexed, then a 2 μL aliquot was added to the cis well. The voltage parameters were run as follows: 60mV/650mV/6μs/1.0ms (read voltage/pulse voltage/pulse voltage duration/pulse frequency). Data were acquired via Labview acquisition software. A representative trace from this run is shown in Fig. 18.

[00241] As shown in Fig. 18, the sequence of all reporter codes was read correctly by the detection system in this experiment, demonstrating 100% accuracy. The absence of deletion or insertion errors again validates the efficacy of pendant PEG-based TCEs as structures enabling highly reliable sequence reads of a polymeric template. Again, runs of homopolymers were accurately sequenced under these conditions, thus underscoring the accuracy of SBX with pendant PEG-based TCEs.

Example 21Sequencing by Expansion of a Complex 222mer Template with pendant PEG-based TCEs and D-cells

[00242] In this Example, translocation control with pendant PEG-based TCEs and D-cell features was assessed by using the SBX protocol to sequence a complex 222mer template. Each XNTP substrate was synthesized to include the following TCE: Y(32)(32)(32)(32)(32)(32)(61)(61)(61)(61)(61)(61), in which “Y” represents the symmetric phosphoramidite brancher; “32” represents pendant mPEG4 (PPA032); and “61” represents pendant PEG (PPA061). Each XNTP also included the following D-cell feature: D(63)D(63)D(63)DD in which “D” represents PEG6 and “63” represent pendant PEG (PPA063). The XNTP substrates were synthesized with the following reporter codes: (XC)DDLLLX; (XT)LXXX; (XA)DD(32)(32)(32)LLLLLLL; and (XG)XXL(32)(32)(32)(32)(32)(32)LLLLLLL, in which “D” represents PEG-6, “L” represents C2, “X” represents PEG-3, and “32” represents pendant mPEG-4 (PPA032).

[00243] To produce Xpandomer copies of the 222mer template, solid-state primer extension reactions were conducted using 5000pmol of each XNTP, 4pmol template and 20pmol E-oligo primer (solid-state Xpandomer synthesis in which the extension oligo is covalently bound to a chip substrate is described in Applicants’ provisional patent application no. 62/826,805, which is herein incorporated by reference in its entirety). The 50µL extension reaction included the following reagents: 50mM TrisCl, pH 8.84, 200mM NH4OAc, 50mM GuCl/20% PEG8K, 10% NMP, 15nmol polyphosphate PP-60.23, 2.5µg Kod SSB, 0.1M urea, 15mM PEM additive and 13µg purified recombinant DNA polymerase (a variant of DPO4 polymerase). The extension reaction was run for 60 minutes at 37°C.

[00244] Xpandomer products were next sequenced using the SBX protocol. Briefly, the constrained Xpandomer products were washed in buffer B.064 (1% Tween-20/3% SDS/5mM HEPES, pH 8.0/100mM NaPO4/15% DMF) and cleaved to generate linearized Xpandomer by adding 200µl buffer C.001 (7.5M DCl) and incubating for 30 minutes at 23°C. The sample was then neutralized by adding 2000µl buffer B.064 and incubating for 2’ at RT. The Xpandomer sample was then subjected to amine modification by adding 500µmol succinate anhydride in buffer B.065 and incubating for 5 minutes at 23°C. The sample was then washed in buffer D.102 (50% ACN) and the Xpandomers were released from the

substrate by photocleavage and eluted in 60 μ l elution buffer.

[00245] Protein nanopores were prepared by inserting α -hemolysin into a DPhPE/hexadecane bilayer member in a buffer of 2 M NH₄Cl and 100 mM HEPES, pH 7.4. The cis well was perfused with buffer AG242 containing 0.4M NH₄Cl, 600mM GuanCl, 100mM HEPES; pH 7.4, and 5% glycerol and the trans well was perfused with buffer AB080 containing 0.4M NH₄Cl, 600mM GuanCl, 5% ethyl acetate, 10mM HEPES; pH 7.4. The Xpandomer sample was heated to 70° C for 2 minutes, cooled completely and vortexed, then a 2 μ L aliquot was added to the cis well. The voltage parameters were run as follows: 70mV/625mV/6 μ s/1.0ms (read voltage/pulse voltage/pulse voltage duration/pulse frequency). Data were acquired via Labview acquisition software.

[00246] To assess the accuracy of the Xpandomer sequence information, sequence data was analyzed by histogram display of the population of sequence reads from the SBX reaction. The analysis software aligns each sequence read to the sequence of the template and trims the extent of the sequence at the end of the reads that does not align with the correct template sequence. A representative histogram of SBX sequencing of the 222mer template is presented in FIG. 19. As can be seen, the SBX experiment generated highly accurate reads of the 222mer template. Notably, the throughput of this experiment was outstanding. These results suggest that SSRTs incorporating pendant PEG-based TCE and D-cell features provide highly accurate and efficient control of Xpandomer translocation through the nanopore.

Example 22

Ratcheting

[00247] In this example of ratcheting a single hemolysin nanopore is prepared in a lipid bilayer with vestibule on the trans-side and having reagent mix composed of 0.4M NH₄Cl, 600mM GuanCl, 100mM HEPES; pH 7.4 in the cis reservoir and 2M NH₄Cl, 100mM HEPES; pH 7.4 in the trans reservoir. Current passing between Ag/AgCl electrodes located in each reservoir is measured by an Axopatch 200B amplifier and digitized at 100k samples/s. To drive the current through the nanopore, a square wave with 50% duty cycle alternating between +70 mV and -50 mV is applied to the trans reservoir along with a 6 μ s pulse of +600 mV applied between the transition from positive to negative voltage (all voltages referenced to the cis reservoir potential). With this applied pulse train, assuming ideal translocation with no deletions or insertions, both reporters for each XNTP in

incorporated into an Xpandomer are measured, one with +70mV and the other with -50 mV. Having two measurements for each base provides redundancy that can provide higher confidence in matched results and also help identify deletions and insertions in non-homopolymer sequence. Fig. 20A illustrates how the cycles of +70mV/600mV pulse/-50mV influence Xpandomer translocation through the nanopore and results in two measurements for the C code followed by two measurements for the A code (and the 1st measurement for the G code). The pattern code shows how in this non-homopolymer sequence, the 2 reporter measurements for each base can be used to identify insertions and deletions. Insertions and deletions are caused when the Xpandomer does not advance to the next reporter (in the nanopore) or it skips the next reporter, respectively.

[00248] Using an SBX synthesis and purification protocol, an Xpandomer sample generated from a synthetic DNA template of known sequence was introduced to the cis reservoir and measurement proceeded. An example of the current measurement for a translocating Xpandomer is shown in Fig. 20B. The graph is scaled so the four reporter current levels from +70 mV measurements (14, 20, 27 and 35 pA) and the 4 four reporter current levels from -50 mV measurements (-12, -20, -25 and -29pA) are shown as indicated by horizontal dashed lines. The data is aligned to the expected DNA template sequence and indicated by the numeric sequence on the graph. Each of the four numbers refers to a base. The blue sequence number indicates a confirmed base match to the template. The arrows depict errors. The arrows designated with the number 1 are non-homopolymer insertions that can be recognized because of the basecall indicated the Xpandomer did not advance. The arrows designated with the number 2 indicate homopolymer insertions that are not recognized since the basecalls are all the same level.

Example 24


Synthesis of Fluoroarabinosyl XNTP Epimers

[00249] This Example describes the synthesis of 2' fluoro (F) epimers of each XNTP (2' FANA XNTPs). These epimers are based on fluorinated nucleosides, referred to as "fluoroarabinosyl nucleic acids" (FANA). It is predicted that the 2' F epimers will demonstrate increased stability during acid treatment, which is a critical step in the synthetic pathway that produces the linearized Xpandomer product. Below are synthetic schemes for generating each 2' FANA XNTP.

A.

Process for making 2'FANA XTTP

commercial from TCI America




[00250] In the first step, fialuridine (compound 1, available from TCI America) is coupled to 1-8 octadiyne via a Sonogashira reaction (see, e.g., Bag, S., Jana, S., and Kasula, M. (2018). Sonogashira Cross-Coupling: Alkyne-Modified Nucleosides and Their Applications. In Palladium-Catalyzed Modification of Nucleosides, Nucleosides, and Oligonucleotides (pp. 75-146). Elsevier). In the second step, compound 2 is treated with approximately one equivalent of DMTrCL in pyridine to produce compound 3. In the third step, compound 3 is converted to the amidate triphosphate, following the protocol described in U.S. patent no. 10,301,345 to Kokoris et al. entitled, “Phosphoramidate esters and use and synthesis thereof”, which is herein incorporated by reference in its entirety.

B.

Process for making 2'FANA XCTP

commercial from TRC Canada




[00251] In the first step, fialcitabine (compound 5, available from TRC Canada) is coupled to 1-8 octadiyne via a Sonogashira reaction (as described above) to produce compound 6. In the second step, compound 6 is treated with approximately one equivalent of DMTrCL in pyridine to produce compound 7. In the third step, the exocyclic amine of compound 7 is protected by an acetyl group (see, e.g., Fan, Y., Gaffney, B., and Jones, R. (2004). Transient Silylation of the Guanosine O6 and the Amino Groups Facilitates N-Acylation. Organic Letters, 6, 15, 2555-2557.) and subsequently converted to the amidate triphosphate 8, as described in U.S. patent no. 10,301,345 to Kokoris et al.

C.


Process for making 2'FANA XGTP

commercial from Granlen



[00252] In the first step, 7-Deaza-7-iodoguanosine (Compound 9 available from Granlen; CAS: 444020-71-7) is treated with 1 equivalent of 1,3-Dichloro-1,1,3,3-tetramethyldisiloxane to provide compound 10 (see, e.g., Markiewicz, W.T. and Wiewiorowski, M. (1978) A new type of silyl protecting groups in nucleoside chemistry. *Nuc. Acids Res.* 5, s185-ss190). In the second step, compound 10 is converted to compound 11 by using fluorinating agent DAST (see, e.g., Pankiewicz, K., Kreminski, J., Ciszewski, L., Ren, W., and Watanabe, K. (1992). A synthesis of 9-(2-deoxy-2-fluoro-β-D-arabinofuranosyl)adenine and -hypoxanthine. An effect of C3'-endo to C2'-endo conformational shift on the reaction course of 2'-hydroxyl group with DAST. *J. of Organic Chem.* 57, 2, 553-559.) In the third step, the exocyclic amine in compound 11 is protected with a phenoxyacetyl group as described above. In the fourth step, the resulting compound 12 is coupled to 1-8 octadiyne by the Sonogashira reaction described above to afford compound 13. In the fifth step, deprotection of the siloxane group as described above will give compound 14. In the sixth step, treatment of compound 14 with 1 equivalent of DMTrCl in pyridine produces compound 15. In the seventh step, compound 15 is converted to guanosine amidate triphosphate 16 as described in U.S. patent no. 10,301,345 to Kokoris et al.

[00253] This same scheme can be used to synthesize the following adenosine triphosphoramidate analog:




from starting compound 7-Deaza-7-iodoadenosine (available from Granlen, CAS:24386-93-4).

[00254] All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, including but not limited to, U.S. Provisional Patent Application No. 62/852,262 filed on May 23, 2019, U.S. Provisional Patent Application No. 62/877,183 filed on July 22, 2019, and U.S. Provisional Patent Application No. 62/885,746 filed on August 12, 2019, are incorporated herein by reference, in their entirety. Such documents may be incorporated by reference for the purpose of describing and disclosing, for example, materials and methodologies described in the publications, which might be used in connection with the presently described invention. The publications discussed above and throughout the text are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the inventors are not entitled to antedate any referenced publication by virtue of prior invention.

CLAIMS

What is claimed is:

1. A compound having the following structure:



wherein

R is OH or H;

nucleobase is adenine, cytosine, guanine, thymine, uracil or a nucleobase analog;

reporter construct is a polymer having a first end and a second end, and comprising, in series from the first end to the second end, a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code;

linker A joins the oxygen atom of the alpha phosphoramidate to the first end of the reporter construct;

linker B joins the nucleobase to the second end of the reporter construct;

and wherein

the translocation control element is a polymer comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 16), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 1,8-O-bis(phosphodiester)-N,N-Diethylpiperazine (compound 26h), 1,2-O-

bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47g), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), or 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52).

2. The compound of claim 1, wherein R is OH.
3. The compound of claim 1, wherein R is H.
4. The compound of any one of claims 1-3 wherein nucleobase is adenine.
5. The compound of any one of claims 1-3 wherein nucleobase is cytosine.
6. The compound of any one of claims 1-3 wherein nucleobase is guanine.
7. The compound of any one of claims 1-3 wherein nucleobase is thymine.
8. The compound of any one of claims 1-3 wherein nucleobase is uracil.
9. The compound of any one of claims 1-3 wherein nucleobase is a nucleobase analog.

10. The compound of any one of claims 1-9 wherein the symmetrical chemical brancher is selected from 1,2,3-O-tris-(phosphodiester)-propane, 1,3-bis-(5-O-phosphodiester-pentylamido)-2-O-phosphodiester-propane, and 1,4,7-O-tris-(phosphodiester)-heptane.

11. The compound of any one of claims 1-9 wherein the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphodiester)-propane.

12. The compound of any one of claims 1-11, wherein the translocation control element is a polymer comprising two or more repeat units selected from Table 1A.

13. The compound of any one of claims 1-11, wherein the translocation control element is a polymer comprising two or more repeat units selected from 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b) and 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b).

14. The compound of any one of claims 1-11, wherein the translocation control element is a polymer comprising the following sequence: [(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))]n1[(1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b))]n2, wherein n1 is from 0 to 6 and n2 is from 6 to 10.

15. The compound of any one claims 1-14, wherein the first and second reporter codes are identical.

16. The compound of any one claims 1-14, wherein the first and second reporter codes are polymers comprising two or more repeat units selected from: hexaethylene glycol (D), ethane (L), triaethylene glycol (X), 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester)-2-(4-Et-1-

(Et-O-mPEG1)-1,2,3-triazole)-propane (compound 31a), 2,3-O-bis(phosphodiester)-1-(1-dimethoxyquinazolinedione)-propane (compound 20c), 2,3-O-bis(phosphodiester)-1-(N9-(3,6-dimethoxycarbazole)-propane (compound 20e), 1,1'-O-bis(phosphodiester)-2,2'-(sulfonylbis(benz-4-yl))-diethanol (compound 26d), 1,1'-O-bis(phosphodiester)-2,2'-bipyridin-4,4'-yl)-dimethanol (compound 26a), 2,3-O-bis(phosphodiester)-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane (compound 20b), 3-(1,2-O-bis(phosphodiester)-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide (compound 20d), 2,3-O-bis(phosphodiester)-1-(N1-(6-Azathymine))-propane (compound 20f), 1,5-O-bis(phosphodiester)-hexahydrofuro[2,6]furan (compound 23), 1,1'-O-bis(phosphodiester)-octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol (compound 26e), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20h), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 5b), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane (compound 31b), and 1,3-O-bis(phosphodiester)-3-O-mPEG4-propane (compound 5a).

17. The compound of any one of claims 1-14, wherein the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and any of the compounds set forth in Table 1A.

18. The reporter code of any one of claims 1-14, wherein the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b).

19. The compound of any one of claims 1-14, wherein the first and second reporter codes are polymers comprising a sequence selected from: (i) [(hexaethylene glycol)₂(ethane)₃(hexaethylene glycol)(triaethylene glycol)], (ii) [(hexaethyleneglycol)₂(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₂(ethane)(triaethylene glycol)₃], (iii) [(hexaethylene glycol)₂(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₃(ethane)₂(hexaethylene glycol)(triaethylene glycol)], and (iv) [(triaethylene glycol)₂(ethane)(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₆(ethane)₇].

20. The compound of any one of claims 1-19, wherein Linker A and Linker B are polymers comprising two or more repeat units selected from: spermine (Q), hexaethylene glycol (D), 2-(((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((phosphodiester-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-phosphodiester-propane-1,3-diyl dibenzoate (compound 62), 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47g), and 1,1'-O-bis(phosphodiester)-N(p-tolyl)-diethanolamine (compound 26b).

21. The compound of any one of claims 1-19, wherein Linker A and Linker B are polymers comprising two or more repeat units selected from spermine and any of the compounds set forth in Table 1A.

22. The compound of any one of claims 1-19 wherein Linker A and Linker B comprise a polymerase enhancement region comprising two repeat units of spermine.

23. The compound of any one of claims 1-22 wherein Linker A and Linker B comprise a translocation deceleration region comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), and 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b).

24. The compound of any one of claims 1-22 wherein Linker A and Linker B comprise a translocation deceleration region comprising a polymer selected from: (i) [((hexaethylene glycol) (1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))3(hexaethylene glycol)2], (ii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))4(hexaethylene glycol)2], (iii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d))4(hexaethylene glycol)2], and (iv) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b))4(hexaethylene glycol)2].

25. The compound of any one of claims 1-24 wherein Linker A is joined to the oxygen atom of the alpha phosphoramidate by a linkage comprising a triazole.

26. The compound of any one of claims 1-24 wherein Linker B is joined to the nucleobase by a linkage comprising a triazole.

27. A reporter construct comprising a polymer having a first end and a second end, and comprising, in series from the first end to the second end, a first reporter code, a symmetrical chemical brancher bearing a translocation control element, and a second reporter code; and wherein the translocation control element is a polymer comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 16), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 1,8-O-bis(phosphodiester)-N,N-Diethylpiperazine (compound 26h), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-

propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,3-O-bis(phosphodiester-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47Gg, 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), or 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52).

28. The reporter construct of claim 27, wherein the symmetrical chemical brancher is selected from 1,2,3 -O-tris-(phosphosphodiester)-propane, 1,3-bis-(5-O-phosphodiester-pentylamido)-2-O-phosphodiester-propane, and 1,4,7-O-tris-(phosphodiester)-heptane.

29. The reporter construct of claim 27, wherein the symmetrical chemical brancher is 1,2,3 -O-tris-(phosphosphodiester)-propane,

30. The reporter construct of any one of claims 27-29, wherein the translocation control element is a polymer comprising two or more repeat units selected from Table 1A.

31. The reporter construct of any one of claims 27-29, wherein the translocation control element is a polymer comprising two or more repeat units selected from 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b) and 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b).

32. The reporter construct of any one of claims 27-29, wherein the translocation control element is a polymer comprising the following sequence: [(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))]n1[(1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b))]n2, wherein n1 is from 0 to 6 and n2 is from 6 to 10.

33. The reporter construct of any one of claims 27-32, wherein the wherein the first and second reporter codes are identical.

34. The reporter construct of any one of claims 27-32, wherein the wherein the first and second reporter codes are polymers comprising two or more repeat units selected from: hexaethylene glycol (D), ethane (L), triaethylene glycol (X), 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b), 1,3-O-bis(phosphodiester)-2-(4-Me-O-PEG3)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35b), 1,3-O-bis(phosphodiester)-2,2-bis(Me-O-mPEG2)-propane (compound 45b), 1,3-O-bis(phosphodiester)-2S-O-(PEG4-O-Bz)-propane (compound 38b), 1,3-O-bis(phosphodiester)-2s-O-mPEG6-propane (compound 12c), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG3)-1-(Me-acetate)-1,2,3-triazole)-propane (compound 35e), 1,3-O-bis(phosphodiester)-2s-O-(4-(Me-O-PEG2)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35a), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG1)-1,2,3-triazole)-propane (compound 31a), 2,3-O-bis(phosphodiester)-1-(1-dimethoxyquinazolinedione)-propane (compound 20c), 2,3-O-bis(phosphodiester)-1-(N9-(3,6-dimethoxycarbazole)-propane (compound 20e), 1,1'-O-bis(phosphodiester)-2,2'-(sulfonylbis(benz-4-yl))-diethanol (compound 26d), 1,1'-O-bis(phosphodiester)-2,2'-bipyridin-4,4'-yl)-dimethanol (compound 26a), 2,3-O-bis(phosphodiester)-1-(N1-(4,6-dimethoxy-3-Me-indole)-propane (compound 20b), 3-(1,2-O-bis(phosphodiester)-propyl)-8,8-dimethylhexahydro-3H-3a,6-methanobenzo[c]isothiazole 2,2-dioxide (compound 20d), 2,3-O-bis(phosphodiester)-1-(N1-(6-Azathymine))-propane (compound 20f), 1,5-O-bis(phosphodiester)-hexahydrofuro[2,6]furan (compound 23), 1,1'-O-bis(phosphodiester)-octahydro-2,6-dimethyl-3,8:4,7-dimethano-2,6-naphthyridin-4,8-diyl)-dimethanol (compound 26e), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20h), 2,3-O-bis(phosphodiester)-1-(N1-(2-Me-5-nitroindole)-propane (compound 20g), 2,3-O-bis(phosphodiester)-1-(5-benzofuran)-propane (compound 20i), 1,2-O-bis(phosphodiester)-3-O-mPEG2-propane (compound 5b), 1,3-O-bis(phosphodiester)-2-(4-Et-1-(Et-O-mPEG3)-1,2,3-triazole)-propane (compound 31b), and 1,3-O-bis(phosphodiester)-3-O-mPEG4-propane (compound 5a).

35. The reporter construct of any one of claims 27-32, wherein the wherein the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and any of the compounds set forth in Table 1A.

36. The reporter construct of any one of claims 27-32, wherein the wherein the first and second reporter codes are polymers comprising two or more repeat units selected from hexaethylene glycol, ethane, triaethylene glycol, and 1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b).

37. The reporter construct of any one of claims 27-32, wherein the wherein the first and second reporter codes are polymers comprising a sequence selected from: (i) [(hexaethylene glycol)₂(ethane)₃(hexaethylene glycol)(triaethylene glycol)], (ii) [(hexaethyleneglycol)₂(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₂(ethane)(triaethylene glycol)₃], (iii) [(hexaethylene glycol)₂(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₃(ethane)₂(hexaethylene glycol)(triaethylene glycol)], and (iv) [(triaethylene glycol)₂(ethane)(1,3-O-bis(phosphodiester)-2S-O-mPEG4-propane (compound 12b))₆(ethane)₇].

38. A symmetrically synthesized report tether (SSRT), wherein the symmetrically synthesized reporter tether is a polymer having a first end and a second end, and comprising in series from the first end to the second end a first linker, a reporter construct according to any one of claims 27-37, and a second linker, wherein the first and second linkers are identical and are polymers comprising two or more repeat units selected from: spermine (Q), hexaethylene glycol (D), 2-(((4-((3-(benzoyloxy)-2-(((1-(3-(benzoyloxy)-2-((benzoyloxy)methyl)-2-((phosphodiester-oxy)methyl)propyl)-1H-1,2,3-triazol-4-yl)methoxy)methyl)-2-((benzoyloxy)methyl)propoxy)methyl)-1H-1,2,3-triazol-1-yl)methyl)-2-O-phosphodiester-propane-1,3-diyl dibenzoate (compound 62), 1,3-O-bis(phosphodiester-2,2-bis(1-Me-4-(Me-O-PEG2-O-Bz)-1,2,3-triazole)-propane (compound 52), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b), 1,2-O-bis(phosphodiester)-3-(4-(Me-O-PEG3-O-Bz)-1-(1,2,3-triazole))-propane (compound 31d), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG2-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47f), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-O-PEG3-O-Me)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 47i), 1,2-O-bis(phosphodiester)-3-(4-methylpiperazine-1-yl)-propane (compound 20j), 1,3-O-bis(phosphodiester-2,2-bis(4-(Me-

O-PEG3-O-Me)-1-(Et-O-Bz)-1,2,3-triazole)-propane (compound 47g), and 1,1'-O-bis(phosphodiester)-N(p-tolyl)-diethanolamine (compound 26b).

39. The symmetrically synthesized reporter tether (SSRT) of claim 38, wherein the first and second linker comprise a polymerase enhancement regions comprising two repeat units of spermine.

40. The symmetrically synthesized reporter tether (SSRT) of claims 38 or 39 comprising a translocation deceleration region comprising two or more repeat units selected from: 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d), 1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG3)-1-(Et-2,2,2-Tris-(Me-O-Bz))-1,2,3-triazole)-propane (compound 37a), and 1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b).

41. The symmetrically synthesized reporter tether (SSRT) of claims 38 or 39 comprising a translocation deceleration region comprising a polymer selected from: (i) [((hexaethylene glycol) (1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))3(hexaethylene glycol)2], (ii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-O-Ac)-1,2,3-triazole)-propane (compound 35c))4(hexaethylene glycol)2], (iii) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2s-O-(4-(Me-O-PEG7)-1-(Et-OBz)-1,2,3-triazole)-propane (compound 35d))4(hexaethylene glycol)2], and (iv) [((hexaethylene glycol)(1,3-O-bis(phosphodiester-2-(4-(Me-O-PEG5)-1-(Et-2,2,2-Tris-(Me-O-Ac))-1,2,3-triazole)-propane (compound 37b))4(hexaethylene glycol)2].

42. The symmetrically synthesized reporter tether (SSRT) of any one of claims 38-41 wherein the first end and the second end comprise a linkage moiety.

43. The symmetrically synthesized reporter tether (SSRT) of claim 42 wherein the linkage moieties comprise an azido ($-N_3$) group.

44. A method for sequencing a target nucleic acid, comprising: a) providing a daughter strand produced by a template-directed synthesis, the daughter strand comprising a plurality of XNTP subunits of claim 1 coupled in a sequence corresponding to a

contiguous nucleotide sequence of all or a portion of the target nucleic acid, wherein the individual XNTP subunits of the daughter strand comprise a reporter construct, a nucleobase residue, and a selectively cleavable bond, and wherein the reporter construct, upon cleavage of the selectively cleavable bond, permits lengthening of the subunits of the daughter strand; b) cleaving the selectively cleavable bonds to yield an Xpandomer of a length longer than the plurality of the subunits of daughter strand, the Xpandomer comprising the reporter constructs for parsing genetic information in a sequence corresponding to the contiguous nucleotide sequence of all or a portion of the target nucleic acid; and c) detecting the reporter constructs of the Xpandomer.

45. The method of claim 44, wherein the reporter constructs for parsing the genetic information comprise a reporter code and a translocation control element, wherein the translocation control element provides translocation control by steric hindrance and pauses translocation of the Xpandomer when passed through a nanopore subjected to a baseline voltage, wherein the translocation control element engages the reporter code within the aperture of the nanopore, wherein the reporter code is sensed by the nanopore.

46. The method of claim 44, wherein the Xpandomer resumes translocation through the nanopore by application of a pulse voltage, wherein the pulse voltage is sufficient to allow translocation of the translocation control element, while leaving the next reporter construct of the Xpandomer free to engage with the nanopore.

47. The method of claim 45, wherein the translocation control element of the reporter construct engaged with the nanopore by steric hindrance translocates upon each pulse of the pulsed voltage.

48. The method of claim 45, wherein the target construct is sensed by the nanopore during the time period between pulses of the pulsed voltage.

49. The method of claim 44, wherein the baseline voltage is from about 55mV to about 75mV.

50. The method of claim 45, wherein the pulse voltage is from about 550mV to about 700mV.

51. The method of claim 45, wherein the pulse voltage has a duration from about 5 μ s to about 10 μ s.

52. The method of claim 45, wherein periodicity of the pulse voltage is from about 0.5ms to 1.5ms.

53. The method of claim 44, wherein the nanopore is subjected to an alternating current (AC).

54. The method of any one of claims 44-53, wherein one or more of the plurality of XNTP subunits comprises a 2' fluoroarabinosyl epimer.

55. A buffer for controlling the rate of translocation of a polymer through a nanopore comprising at least one salt selected from the group consisting of NH₄Cl, MgCl₂, LiCl, KCl, CsCl, NaCl, and CaCl₂.

56. The buffer of claim 55, further comprising at least one solvent selected from the group consisting of 3-methyl-2-oxazolidinone (MOA), DMF, ACN, DMSO, and NMP, wherein the solvent is present in the range from about 1% vol/vol to about 35% vol/vol.

57. The buffer of claim 55, further comprising at least one additive selected from the group consisting of sodium hexanoate (NaHex), EDTA, redox reagents, PEG, glycerol, ficoll and the like.

58. A buffer system for controlling the rate of translocation of a polymer through a nanopore detector comprising a cis buffer and a trans buffer, wherein the cis buffer comprises a first salt concentration and the trans buffer comprises a second salt concentration, wherein the first salt concentration is lower than the second salt concentration.

1 / 24




FIG. 1A

2 / 24

FIG. 1B






FIG. 1C




3 / 24

FIG. 1D



4 / 24



5 / 24





FIG. 3

6 / 24






FIG. 5A





FIG. 5B




FIG. 5C




FIG. 5D

Fig. 6






Fig. 7


Exemplary
SSRT

Key:







10 / 24






12 / 24



13 / 24


FIG. 10

DTB = Desthiobiotin



14 / 24

FIG. 11






FIG. 13A

FIG. 13B

Control codes

PEG codes

L1 / C	10.8 ms
L2 / T	9.4 ms
L3 / A	7.6 ms
L4 / G	10.2 ms

L1 / C	10.0 ms
L2 / T	11.2 ms
L3 / A	8.0 ms
L4 / G	11.2 ms





FIG. 14A




FIG. 14B






FIG. 15




key:
level 1 - C
level 3 - A

FIG. 16



key:
level 1 - C
level 2 - T
level 3 - A
level 4 - G


FIG. 17



key:
level 1 - C
level 2 - T
level 3 - A
level 4 - G

21 / 24

FIG. 18



22 / 24

Fig. 19





FIG. 20A



Examples:

No errors: C C A A G G
A Insertion: C C A C A A G G
A Deletion: C A G G

24 / 24

FIG. 20B




FIG. 4

