

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2005293703 B2**

(54) Title
A method for the carry-over protection in DNA amplification systems targeting methylation analysis achieved by a modified pre-treatment of nucleic acids

(51) International Patent Classification(s)
C12Q 1/68 (2006.01)

(21) Application No: **2005293703** (22) Date of Filing: **2005.10.11**

(87) WIPO No: **WO06/040187**

(30) Priority Data

(31) Number	(32) Date	(33) Country
04090389.0	2004.10.11	EP
60/618,267	2004.10.12	US

(43) Publication Date: **2006.04.20**

(44) Accepted Journal Date: **2010.02.18**

(71) Applicant(s)
Epigenomics AG

(72) Inventor(s)
Tetzner, Reimo;Dietrich, Dimo

(74) Agent / Attorney
PIZZEYS, Level 14, ANZ Centre 324 Queen Street, Brisbane, QLD, 4000

(56) Related Art
US2003/0022215 A1
Kubareva, E. A. et al., Biochemistry (Moscow), 2001, vol. 66
no.12, pages 1356-1360
WO2003/070986 A1
Grunau, C. et al., Nucleic Acids Research, 2001, vol. 29, no. 13, e65
Thomassin, H. et al, Methods,1999, vol. 19, pages 465-475

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 April 2006 (20.04.2006)

PCT

(10) International Publication Number
WO 2006/040187 A3

(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/EP2005/011133

(22) International Filing Date: 11 October 2005 (11.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04090389.0 11 October 2004 (11.10.2004) EP
60/618,267 12 October 2004 (12.10.2004) US

(71) Applicant (*for all designated States except US*): **EPIGENOMICS AG** [DE/DE]; Kleine Präsidentenstr. 1, 10178 Berlin (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **TETZNER, Reimo** [DE/DE]; Wicherstr. 6, 10439 Berlin (DE). **DIETRICH, Dima** [DE/DE]; Katharinenstr. 27, 10711 Berlin (DE).

(74) Agent: **SCHUBERT, Klemens**; Neue Promenade 5, 10178 Berlin (DE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
5 October 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A METHOD FOR THE CARRY-OVER PROTECTION IN DNA AMPLIFICATION SYSTEMS TARGETING METHYLATION ANALYSIS ACHIEVED BY A MODIFIED PRE-TREATMENT OF NUCLEIC ACIDS

(57) Abstract: Disclosed is a method for the specific amplification of template DNA in the presence of potentially contaminating PCR products from previous amplification experiments. In the first step DNA is contacted with a bisulfite solution, which reacts with unmethylated cytosines but not with methylated cytosines, by sulfonating them. This results in deamination of the cytosine whereby sulfonated uracil is generated. Such sulfonation protects the template nucleic acid from being a target for the enzyme UNG. Any contaminating DNA, which contains unprotected unsulfonated or desulfonated uracils is degraded enzymatically while UNG is active. After UNG treatment and inactivation, the sulfonated uracil bases are converted into uracil by desulfonation. This method is useful for decontamination of nucleic acid samples, or rather for avoiding amplification of 'carry over products' in particular in the context of DNA methylation analysis. Furthermore it can be used as a simplified method of bisulfite treatment in general.

WO 2006/040187 A3

A method for the carry-over protection in DNA amplification systems targeting methylation analysis achieved by a modified pre-treatment of nucleic acids

5

In recent decades in molecular biology studies have focused primarily on genes, the transcription of those genes into RNA, and the translation of the RNA into protein. There has been a more limited analysis of the regulatory mechanisms associated with gene control. Gene regulation, for example, at what stage of development of the individual a gene is activated or inhibited, and the tissue specific nature of this regulation is less understood. However, it can be correlated with a high degree of probability to the extent and nature of methylation of the gene or genome. Specific cell types can be correlated with specific methylation patterns and this has been shown for a number of cases (Adorjan et al. (2002) Tumour class prediction and discovery by microarray-based DNA methylation analysis. Nucleic Acids Res. 30 (5) e21).

20

In higher order eukaryotes DNA is methylated nearly exclusively at cytosines located 5' to guanine in the CpG dinucleotide. This modification has important regulatory effects on gene expression, especially when involving CpG rich areas, known as CpG islands, located in the promoter regions of many genes. While almost all gene-associated islands are protected from methylation on autosomal chromosomes, extensive methylation of CpG islands has been associated with transcriptional inactivation of selected imprinted genes and genes on the inactive X-chromosome of females.

30

The cytosine's modification in form of methylation contains significant information. It is obvious that the identification of 5-methylcytosine in a DNA sequence as opposed to unmethylated cytosine is of greatest importance to analyze its role further. But, because the 5-methylcytosine behaves just as a cytosine for what concerns its hybridization preference (a property relied on for sequence analysis) its position cannot be identified by a normal sequencing reaction.

35

Furthermore, in any amplification, such as a PCR amplification, this relevant epigenetic information, methylated cytosine or unmethylated cytosine, will be completely lost.

5

Several methods are known that solve this problem. Usually genomic DNA is treated with a chemical or enzyme leading to a conversion of the cytosine bases, which consequently allows to differentiate the bases afterwards. The most common methods are a) the use of methylation sensitive restriction enzymes capable of differentiating between methylated and unmethylated DNA and b) the treatment with bisulfite. The use of said enzymes is limited due to the selectivity of the restriction enzyme towards a specific recognition sequence.

10

15

Therefore, the '*bisulfite treatment*', allowing for the specific reaction of bisulfite with cytosine, which, upon subsequent alkaline hydrolysis, is converted to uracil, whereas 5-methylcytosine remains unmodified under these conditions (Shapiro et al. (1970) Nature 227: 1047) is currently the most frequently used method for analyzing DNA for 5-methylcytosine. Uracil corresponds to thymine in its base pairing behavior, that is it hybridizes to adenine; whereas 5-methylcytosine does not change its chemical properties under this treatment and therefore still has the base pairing behavior of a cytosine, that is hybridizing with guanine. Consequently, the original DNA is converted in such a manner that 5-methylcytosine, which originally could not be distinguished from cytosine by its hybridization behavior, can now be detected as the only remaining cytosine using "normal" molecular biological techniques, for example, amplification and hybridization or sequencing. All of these techniques are based on base pairing, which can now be fully exploited. Comparing the sequences of the DNA with and without bisulfite treatment allows an easy identification of those cytosines that have been unmethylated.

20

25

30

35

An overview of the further known methods of detecting 5-methylcytosine may be gathered from the following review arti-

cle: Fraga FM, Esteller M, Biotechniques 2002 Sep;33(3):632, 634, 636-49.

As the use of methylation-specific enzymes is restricted to
5 certain sequences (comprising restriction sites), most methods
are based on a *bisulfite treatment* that is conducted before a
detection or amplifying step (for review: DE 100 29 915, A1
p.2, lines 35-46 or the according translated US application
10/311,661, see also WO 2004/067545). The term 'bisulfite
10 treatment' is meant to comprise treatment with a bisulfite, a
disulfite or a hydrogensulfite solution. As known to the ex-
pert skilled in the art and according to the invention, the
term "bisulfite" is used interchangeably for "hydrogensul-
fite".

15 Several protocols are known in the art. However, all of the
described protocols, comprise of the following steps: The ge-
nomic DNA is isolated, , denatured, converted several hours by
a concentrated bisulfite solution and finally desulfonated and
20 desalted (e.g.: Frommer et al.: A genomic sequencing protocol
that yields a positive display of 5-methylcytosine residues in
individual DNA strands. Proc Natl Acad Sci U S A. 1992 Mar
1;89(5):1827-31).

25 In recent times several technical improvements of the bisul-
fite methods were developed.

The agarose bead method incorporates the DNA to be investi-
gated in an agarose matrix, through which diffusion and rena-
30 turation of the DNA is prevented (bisulfite reacts only on
single-stranded DNA) and all precipitation and purification
steps are replaced by rapid dialysis (Olek A. et al. A modi-
fied and improved method for bisulphite based cytosine methy-
lation analysis, Nucl. Acids Res. 1996, 24, 5064-5066).

35 In the patent application WO 01/98528 (20040152080) a bisul-
fite conversion is described in which the DNA sample is incu-
bated with a bisulfite solution of a concentration range be-
tween 0.1 mol/l to 6 mol/l in presence of a denaturing reagent

and/or solvent and at least one scavenger. In said patent application several suitable denaturing reagents and scavengers are described. The final step is incubation of the solution under alkaline conditions whereby the deaminated nucleic acid is desulfonated.

In the patent application WO 03/038121 (US 20040115663) a method is disclosed in which the DNA to be analysed is bound to a solid surface during the bisulfite treatment. Consequently, purification and washing steps are facilitated.

In the patent application WO 04/067545 a method is disclosed in which the DNA sample is denatured by heat and incubated with a bisulfite solution of a concentration range between 3 mol/l to 6.25 mol/l. Thereby the pH value is between 5.0 and 6.0 and the nucleic acid is deaminated. Finally an incubation of the solution under alkaline conditions takes place, whereby the deaminated nucleic acid is desulfonated.

The understanding in the art that a "bisulfite conversion" usually comprises the step of desulfonation can for example be taken from said application: "According to the invention the term a "bisulfite reaction", "bisulfite treatment" or "bisulfite method" shall mean a reaction for the conversion of a cytosine base, preferably cytosine bases, in a nucleic acid to an uracil base, preferably uracil bases, in the presence of bisulfite ions whereby preferably a 5-methyl-cytosine base, preferably 5-methyl-cytosine bases, is not significantly converted. This reaction for the detection of methylated cytosines is described in detail by Frommer et al., supra and Grigg and Clark (Grigg, G. and Clark, S. , Bioessays 16 (1994) 431-436). The bisulfite reaction contains a deamination step and a desulfonation step, which can be conducted separately or simultaneously (see Figure 1 ; Grigg and Clark, supra). The statement that 5-methyl-cytosine bases are not significantly converted shall only take the fact into account that it cannot be excluded that a small percentage of 5-methyl-cytosine bases is converted to uracil although it is intended to convert only and exclusively the (non-methylated) cytosine bases (Frommer

et al., supra). The expert skilled in the art knows how to perform the bisulfite reaction, e. g. by referring to Frommer et al. , supra or Grigg and Clark, supra who disclose the principal parameters of the bisulfite reaction."

5

Furthermore in said application it is described what the general state of the art is with regard to the different protocols: "From Grunau et al., supra, it is known to the expert in the field what variations of the bisulfite method are possible. In summary, in the deamination step a buffer containing bisulfite ions, optionally chaotropic agents and optionally further reagents as an alcohol or stabilizers as hydroquinone are employed and the pH is in the acidic range. The concentration of bisulfite is between 0.1 and 6 M bisulfite, preferably between 1 M and 5.5 M, the concentration of the chaotropic agent is between 1 and 8 M, whereby preferably guanidinium salts are employed, the pH is in the acidic range, preferably between 4.5 and 6.5, the temperature is between 0 °C and 90 °C, preferably between room temperature (25 °C) and 90 °C, and the reaction time is between 30 min and 24 hours or 48 hours or even longer, but preferably between 1 hour and 24 hours. The desulfonation step is performed by adding an alkaline solution or buffer as e. g. a solution only containing a hydroxide, e. g sodium hydroxide, or a solution containing ethanol, sodium chloride and sodium hydroxide (e. g. 38 % EtOH, 100 mM NaCl, 200 mM NaOH) and incubating at room temperature or elevated temperatures for several min, preferably between 5 min and 60 min."

It is therefore clear that the desulfonation is an inherent feature of all of these methods, in any case a desulfonation takes place before the nucleic acids are used as templates for amplification reactions, in order to provide an ideal template for the polymerase utilized in the following reactions.

35

In the patent application WO 05/038051 improvements for the conversion of unmethylated cytosine to uracil by treatment with a bisulfite reagent are described. According to this method the reaction is carried out in the presence of 10-35 %

by volume, preferentially in the presence of 20-30 % by volume of dioxane, one of its derivatives or a similar aliphatic cyclic ether. The bisulfite reaction can also be carried out in the presence of a n-alkylene glycol compound, particularly in the presence of their dialkyl ethers, and especially in the presence of diethylene glycol dimethyl ether (DME). These compounds can be present in a concentration of 1-35 % by volume, preferentially of 5-25 % by volume. According to this invention the bisulfite conversion is conducted at a temperature in the range of 0-80 °C and that the reaction temperature is increased for 2 to 5 times to a range of 85-100 °C briefly during the course of the conversion (thermospike). It is further preferred that the temperature increases to 85-100 °C, in particular to 90-98 °C during the temperature increase of brief duration.

Subsequent to a bisulfite treatment, usually short, specific fragments of a known gene are amplified and either completely sequenced (Olek A, Walter J. (1997) The pre-implantation ontogeny of the H19 methylation imprint. Nat Genet. 3: 275-6) or individual cytosine positions are detected by a primer extension reaction (Gonzalzo ML and Jones PA. (1997) Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (MS-SNuPE). Nucleic Acids Res. 25 :2529-31, WO 95/00669) or by enzymatic digestion (Xiong Z, Laird PW. (1997) COBRA: a sensitive and quantitative DNA methylation assay. Nucleic Acids Res. 25: 2535-4).

Another technique to detect hypermethylation is the so-called methylation specific PCR (MSP) (Herman JG, Graff JR, Myohanen S, Nelkin BD and Baylin SB. (1996), Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. Proc Natl Acad Sci U S A. 93: 9821-6). The technique is based on the use of primers that differentiate between a methylated and a non-methylated sequence if applied after bisulfite treatment of said DNA sequence. The primer either contains a guanine at the position corresponding to the cytosine in which case it will after bisulfite treatment only bind if the position was

5 methylated. Or the primer contains an adenine at the corresponding cytosine position and therefore only binds to said DNA sequence after bisulfite treatment if the cytosine was unmethylated and has hence been altered by the bisulfite treatment so that it hybridizes to adenine. With the use of these primers, amplicons can be produced specifically depending on the methylation status of a certain cytosine and will as such indicate its methylation state.

10 Another technique is the detection of methylation via a labelled probe, such as used in the so called Taqman PCR, also known as MethyLight (US 6,331,393). With this technique it became feasible to determine the methylation state of single or of several positions directly during PCR, without having to
15 analyze the PCR products in an additional step.

In addition, detection by hybridization has also been described (Olek et al., WO 99/28498).

20 The treatment with bisulfite (or similar chemical agents or enzymes) with the effect of altering the base pairing behaviour of one type of cytosine specifically, either the methylated or the unmethylated, thereby introducing different hybridisation properties, makes the treated DNA more applicable
25 to the conventional methods of molecular biology, especially the polymerase based amplification methods, such as the PCR.

Base excision repair occurs in vivo to repair DNA base damage involving relatively minor disturbances in the helical DNA
30 structure, such as deaminated, oxidized, alkylated or absent bases. Numerous DNA glycosylases are known in the art, and function in vivo during base excision repair to release damaged or modified bases by cleavage of the glycosidic bond that links such bases to the sugar phosphate backbone of DNA (Memisoglu, Samson, Mutation Res. (2000), 451:39-51). All DNA gly-
35 cosylases cleave glycosidic bonds but differ in their base substrate specificity and in their reaction mechanisms.

One widely recognized application of such glycosylases is decontamination in PCR applications. In any such PCR amplification, 2 to the 30 (2^{30}) or more copies of a single template are generated. This very large amount of DNA produced helps in the subsequent analysis, like in DNA sequencing according to the Sanger method, but it can also become a problem when this amount of DNA is handled in an analytical laboratory. Even very small reaction volumes, when inadvertently not kept in a closed vial, can lead to contamination of the whole work environment with a huge number of DNA copies. These DNA copies may be templates for a subsequent amplification experiment performed, and the DNA analysed subsequently may not be the actual sample DNA, but contaminating DNA from a previous experiment. This may also lead to positive negative controls that should not contain any DNA and therefore no amplification should be observed.

In practice, this problem can be so persistent that whole laboratories may move to a new location, because contamination of the work environment makes it impossible to still carry out meaningful PCR based experiments. In a clinical laboratory, however, the concern is also that contaminating DNA may cause false results when performing molecular diagnostics. This would mean that actually contaminating DNA that stems from a previous patient is analysed, instead of the actual sample to be investigated.

Therefore, measures have been implemented to avoid contamination. This involves, for example, a PCR amplification and detection in one tube in a real time PCR experiment. In this case, it is not required that a PCR tube be opened. After use, the tube will be kept closed and discarded and therefore the danger of contamination leading to false results is greatly reduced.

In addition, molecular means exist that reduce the risk of contamination. In a polymerase chain reaction, the enzyme uracil-DNA-glycosylase (UNG) reduces the potential for false positive reactions due to amplicon carryover (see e.g. US

5,035,996 or Thornton CG, Hartley JL, Rashtchian A (1992). Utilizing uracil DNA glycosylase to control carryover contamination in PCR: characterization of residual UDG activity following thermal cycling. *Biotechniques*. 13(2):180-4). The principle of this contamination protection method is that in any amplification instead of dTTP dUTP is provided and incorporated and the resulting amplicon can be distinguished from its template and any future sample DNA by uracil being present instead of thymine. Prior to any subsequent amplification, uracil DNA-glycosylase (UNG) is used to cleave these bases from any contaminating DNA, and therefore only the legitimate template remains intact and can be amplified. This method is considered the standard method of choice in the art and is widely used in DNA based diagnostics. The following is a citation from a publication that summarizes the use of UNG (Longo MC, Berninger MS, Hartley JL (1990). Use of uracil DNA glycosylase to control carry-over contamination in polymerase chain reactions. *Gene*. 1990 Sep 1;93(1):125-8.):

"Polymerase chain reactions (PCRs) synthesize abundant amplification products. Contamination of new PCRs with trace amounts of these products, called carry-over contamination, yields false positive results. Carry-over contamination from some previous PCR can be a significant problem, due both to the abundance of PCR products, and to the ideal structure of the contaminant material for re-amplification. We report that carry-over contamination can be controlled by the following two steps: (i) incorporating dUTP in all PCR products (by substituting dUTP for dTTP, or by incorporating uracil during synthesis of the oligodeoxyribonucleotide primers; and (ii) treating all subsequent fully preassembled starting reactions with uracil DNA glycosylase (UNG), followed by thermal inactivation of UNG. UNG cleaves the uracil base from the phosphodiester backbone of uracil-containing DNA, but has no effect on natural (i.e., thymine-containing) DNA. The resulting apyrimidinic sites block replication by DNA polymerases, and are very labile to acid/base hydrolysis. Because UNG does not react with dUTP, and is also inactivated by heat denaturation prior to the actual PCR, carry-over contamination of PCRs can

be controlled effectively if the contaminants contain uracils in place of thymines."

Another method for carry over protection in PCR has been described by Walder et al (Walder RY, Hayes JR, Walder JA Use of PCR primers containing a 3-terminal ribose residue to prevent cross-contamination of amplified sequences. Nucleic Acids Res 1993 Sep 11;21(18):4339-43.)

It has been described here that carry over protection can be achieved - however not very reproducibly - by using primers consisting of a 3'-end which is characterized as a ribocytidine. After primer extension the amplification product is cleaved specifically at the site of this ribonucleotide by an enzyme known as RNase A. That way the potentially contaminating amplicates are shortened at their ends and cannot serve as templates for said primers in the following amplification procedure. However, the disadvantage inherent to this method is the instability of the primer molecules, containing a ribonucleotide at the 3'-end.

All of the documents cited herein are hereby incorporated by reference in its entirety.

As the existence of uracils is an inherent feature of bisulfite converted DNA and the necessary property relied upon for detecting methylation differences, the method of choice for carry over protection based on uracil-DNA-glycosylase enzyme activity as described above cannot be applied. However, a number of powerful assays for diagnosis are based on PCR performed on bisulfite converted DNA as a template. Therefore for the routinely performance of such assays in a laboratory new methods for carry over prevention need to be developed. There is a great need in the art to provide solutions to the problem of how to achieve a reliable carry over protection when analysing methylation of cytosine positions in DNA from patient samples.

The difficulty of solving the problem for decontamination of bisulfite converted templates is considered a general one, that can not be solved by adaptation of the standard UNG method, as any bisulfite converted DNA will contain uracil as well. It has therefore commonly been argued that, in any uracil-DNA-glycosylase step, the template DNA would be destroyed along with any contaminating DNA.

Surprisingly, the inventors were able to solve this principal problem by inventing the claimed method. The central idea of the method according to the invention is to sulfonate and/or deaminate unmethylated cytosines only without a subsequent desulfonation. After the unmethylated cytosines are converted to C6-sulfonated uracils the reaction mixture is treated with UNG, which degrades all uracil containing DNA and hence every contaminating DNA, but has no effect on the sulfonated uracil containing DNA. If the case may be, a deactivation of the UNG followed by a desulfonation of the sulfonated uracils can be carried out.

The discovery, reported upon for the first time in this application, that sulfonation of uracil at the C6 position protects the uracil from being degraded by UNG allows to find a new and surprisingly easy solution to said problem. One embodiment of this invention therefore comprises a method which provides both a sufficient and reliable differentiation between methylated and unmethylated cytosines, as well as the applicability of the gold standard of carry over protection (based on use of UNG) for common PCR based assays.

Short description of the invention

Disclosed is a method for the specific amplification of template DNA in the presence of potentially contaminating PCR products from previous amplification experiments. This template DNA is usually derived from isolating the genomic DNA to be analysed before the method can be applied. Also, the template nucleic acid used in this method is usually already de-

natured and therefore present in a single stranded modus. In the first step of the method according to the invention the DNA is contacted with a bisulfite solution, which reacts with unmethylated cytosines but not with methylated cytosines, by sulfonating them. This results in a modification of said nucleic acids, which is known as sulfonation. This sulfonation of unmethylated cytosine in aqueous solution results in deamination of the cytosine whereby sulfonated uracil is generated. It has now for the first time been recognized said such sulfonation, which occurs only at the unmethylated cytosine bases a) protects the template nucleic acid from being a target for the enzyme UNG and thereby allows for discrimination of template nucleic acid and potentially contaminating nucleic acids. Any contaminating DNA, which contains unprotected unsulfonated or desulfonated uracils while UNG is active, is subsequently degraded enzymatically and only the template nucleic acid from the sample remains to be amplified in the next step.

After treatment with UNG has been accomplished and UNG activity was terminated the sulfonated uracil bases (which replace the unmethylated cytosines) are converted into uracil by desulfonation. The method is useful for decontamination of nucleic acid samples, or rather for avoiding amplification of 'carry over products' in particular in the context of DNA methylation analysis.

Presently, no method has been reported to decontaminate DNA samples that would be compatible with bisulfite treated DNA employed as the template for an amplification procedure like PCR.

By the provided method according to the invention, it could be achieved to make the most commonly used method based on the glycosylase enzyme UNG, as described above, applicable to DNA methylation analysis:

The present invention solves the problem by describing a method for providing a decontaminated template nucleic acid

for polymerase based amplification reactions suitable for DNA methylation analysis, comprising the following steps:

incubating nucleic acids with a bisulfite reagent solution,
5 whereby the unmethylated cytosines within said nucleic acid are sulfonated, or sulfonated and deaminated, and

mixing the sulfonated or sulfonated and deaminated template nucleic acid with the components required for a polymerase mediated amplification reaction or an amplification based detection assay, and
10

adding to this mixture UNG and incubating the mixture, whereby nucleic acids containing non-sulfonated uracil are degraded,
15 and

terminating UNG activity and desulfonating the template nucleic acid, thereby converting unmethylated deaminated and sulfonated cytosines, i.e. sulfonated uracils into uracils.
20

Subsequently, a polymerase based amplification or amplification based assay is performed, which preferably takes place in the presence of dUTPs instead of dTTPs.

25 Preferably, during the polymerase activity is started during desulfonation step.

Detailed description of the invention

30

The method will typically be carried out performing at least the following steps in the given order:

Firstly, incubating a template nucleic acids with a bisulfite reagent containing solution, whereby the unmethylated cytosines within said nucleic acid are sulfonated, or sulfonated and deaminated, and
35

secondly, mixing the sulfonated, or sulfonated and deaminated, template nucleic acid with the components required for a polymerase mediated amplification reaction or an amplification based detection assay, and thirdly, adding to this mixture UNG units and incubating said mixture, whereby any nucleic acids containing non-sulfonated uracils are degraded, whereas sulfonated uracils essentially remain intact and fourthly terminating UNG activity and desulfonating the template nucleic acid.

The sulfonation takes place at C6 of the base cytosine (see figure 1). Deamination of sulfonated cytosines takes place spontaneously in aqueous solution. Thereby the sulfonated cytosine is converted into a sulfonated uracil.

The method according to invention is based on two essential discoveries. Firstly, we found out that sulfonated nucleic acids are stable up to at least 6 days at 4 °C, e.g. when stored in a common laboratory fridge. This discovery was essential to the method according to the invention because a spontaneous uncontrolled desulfonation of the nucleic acids would render the method unreliable and unstable. Whereas knowing that the nucleic acid's sulfonation pattern, basically resembling the nucleic acids methylation pattern, will remain stable when stored for several days at a lower temperature allows the use of this feature in performing sensitive assays to detect exactly which nucleic acids were methylated within a given sample to which extent.

Secondly, it could be shown in our hands that UNG (uracil-DNA-glycosylase) does not degrade nucleic acids containing sulfonated uracils, in other words sulfonated uracils are no substrate for UNG activity, and therefore protected from degradation with UNG.

It could also be shown that "real time PCR" assays using sulfonated nucleic acids as template performed well in presence of UNG activity, indicating that sulfonated nucleic acids as

derived from the first step of the bisulfite treatment can serve as templates in PCR based assays.

5 Lastly, the question was to be answered whether the desulfonation reaction, which must take place before the nucleic acid can be amplified by a polymerase mediated amplification, can be performed within the PCR reaction.

10 The developed method, according to the invention, was tested successfully for GSTP1 and connexine, see examples.

In the first step, the bisulfite mediated cytosine sulfonation may be initiated according to the first steps of common bisulfite conversion protocols as indicated above, in particular as indicated in WO 05/038051. The reaction may take place both in solution as well as also on DNA bound to a solid phase. Preferably sodium disulfite (= sodium bisulfite/sodium metabisulfite) is used, since it is more soluble in water than sodium sulfite. The disulfite salt disproportionates in aqueous solution to the hydrogen sulfite anions necessary for the cytosine sulfonation. When bisulfite concentration is discussed in more detail, this refers to the concentration of hydrogen sulfite and sulfite anions in the reaction solution. For the method according to the invention, concentration ranges of 0.1 to 6 mol/l are possible. Particularly preferred is a concentration range of 1 to 6 mol/l, and most particularly preferred, 2-4 mol/l. However, when dioxane is used as a denaturing agent, the maximal working concentration of bisulfite is smaller. Dioxane may also be utilized in different concentrations. Preferably, the dioxane concentration amounts to 10 to 35 %, particularly preferred is 20 to 30 %, and most particularly preferred is 22 to 28 %, especially 25 %.

35 In the particularly preferred embodiments with a dioxane concentration of 22-28 %, the final preferred bisulfite concentration amounts to 3.3 to 3.6 mol/l, and in the most particularly preferred embodiment with a dioxane concentration of 25 %, it amounts to 3.5 mol/l (see Examples).

In another preferred embodiment, DME is used as denaturing agent in different concentrations. DME is used in concentrations in the range of 1-35 %, preferable in the range of 5-
5 25 %, and most preferably 10 %.

In a particularly preferred embodiment the bisulfite conversion is carried out in the presence of scavengers. The preferred scavengers are chromane derivatives, e.g., 6-hydroxy-
10 2,5,7,8, -tetramethylchromane 2-carboxylic acid (also known as: Trolox-C™). Further scavengers are listed in the patent application WO 01/98528 (= DE 100 29 915; = US application 10/311,661; incorporated herein in its entirety).

15 The bisulfite conversion can be conducted in a wide temperature range from 0 to 95 °C. However, in a preferred embodiment the reaction temperature lies between 30-70 °C. Particularly preferred is a range between 45-60 °C; most particularly preferred between 50-55 °C.

20 The optimal reaction time of the bisulfite treatment depends on the reaction temperature. The reaction time normally amounts to between 1 and 18 hours (see: Grunau et al. 2001, Nucleic Acids Research; 29(13):E65-5.). The preferred reaction
25 time is 4-6 hours for a reaction temperature of 50 °C.

In a particularly preferred embodiment of the method according to the invention, the bisulfite conversion is conducted at mild reaction temperatures, wherein the reaction temperature
30 is then clearly increased for a short time at least once during the course of the conversion. The temperature increases of short duration are named "thermospikes" below. The "standard" reaction temperature outside the thermospikes is denoted as the basic reaction temperature. The basic reaction temperature
35 amounts to between 0 and 80 °C, preferably between 30-70 °C, most preferably 45°-55°C, as described above. The reaction temperature during a thermospike is increased to over 85 °C by at least one thermospike. The optimal number of thermospikes is a function of the basic reaction temperature. The higher

the optimal number of thermospikes is, the lower is the basic reaction temperature. At least one thermospike is necessary in each case. And, on the other hand, in principle, any number of thermospikes is conceivable.

5

In a particular embodiment the preferred number of thermospikes is between 1 and 10 thermospikes, depending on the basic reaction temperature. Two to five thermospikes are particularly preferred. During the thermospikes the reaction temperature increases preferably to 85 to 100 °C, particularly preferably to 90-98 °C, and most preferably to 94 °C-96 °C. The duration in time of the temperature increases also depends on the volume of the reaction batch.

15 The duration in time of the thermospikes also depends on the volume of the reaction batch. It must be assured that the temperature is increased uniformly throughout the total reaction solution. For a 20 µl reaction batch when using a thermocycler a duration between 15 seconds and 1.5 minutes, especially a duration between 20 and 50 seconds is preferred. In a particular preferred embodiment the duration is 30 seconds. Operating on a volume of 100 µl the preferred range lies between 30 seconds and 5 minutes, especially between 1 and 3 minutes. Particularly preferred are 1.5 minutes. For a volume of 600 µl, a duration of 1 to 6 minutes is preferred, especially between 2 and 4 minutes. Particularly preferred is a duration of 3 minutes. A person skilled in the art will easily be able to determine suitable durations of thermospikes in relation to a variety of reaction volumes.

30

The above described use of thermospikes leads to a significantly better conversion rates in the bisulfite conversion reaction, even when the above-described denaturing solvents are not utilized. According to the invention, a method for bisulfite conversion of DNA is hereby characterized in that the basic reaction temperature amounts to 0 °C to 80 °C and that the reaction temperature is increased for a short time to over 85 °C at least once in the course of the conversion.

35

In the second step, prior to any desulfonation step, units of an enzyme activity, which specifically degrades non-sulfonated uracil containing nucleic acids, are added to said premix. In a preferred embodiment, this degrading enzyme is a DNA-glycosylase or an endonuclease, in particularly UNG (uracil-DNA-glycosylase). The contaminating nucleic acid is characterized in that it contains non-sulfonated uracil bases. The added degrading enzyme is characterized by cleaving the non-sulfonated uracil base from the phosphodiester backbone of non-sulfonated uracil-containing nucleic acids, but has no effect on sulfonated-uracil containing nucleic acid or on thymine containing nucleic acid, that does not contain uracil. The resulting apyrimidinic sites block replication by DNA polymerases, and are very labile to acid/base hydrolysis.

In another preferred embodiment, the first step is carried out as described above. Thereafter, in an intermediate step, the sulfonated and/or deaminated nucleic acid is mixed with components required for a polymerase mediated amplification reaction or an amplification based detection assay. The amplification reaction mix is prepared according to standard protocols. Such an amplification mix, preferably a PCR mix, may contain at least one primer set of two primers and a polymerase. This polymerase preferably is a heat stable enzyme, even more preferred is the use of a thermally activated polymerase for hot start PCR, and most particularly preferred a thermally activated Taq polymerase is used.

The following second step is also carried out as described above. Units of an enzyme activity, which specifically degrades sulfonated-uracil containing nucleic acid, are added to said premix. The sulfonated sample nucleic acid and a set of at least two primer oligonucleotides are incubated with a composition of enzymes, including an enzyme with sulfonated-uracil containing nucleic acid degrading activity and buffers to cleave or degrade any contaminating nucleic acid. The contaminating nucleic acid is characterized in that it contains uracil bases. The added degrading enzyme activity is charac-

terized by cleaving the uracil base from the phosphodiester backbone of non-sulfonated uracil containing nucleic acid, but has no effect on sulfonated uracil containing nucleic acid or on thymine containing nucleic acid, that does not contain uracil. The resulting apyrimidinic sites block replication by DNA polymerases, and are very labile to acid/base hydrolysis. In principle, the enzymatic activity is any enzymatic activity, which causes specifically apyrimidinic sites or one or more nicks adjacent to a non-sulfonated uracil base. In any case this will result in a block of the replication by DNA polymerase.

The primer oligonucleotides will be chosen such that they amplify a fragment of interest. It is particularly preferred that these primers are designed to amplify a nucleic acid fragment of a template nucleic acid sample by means of a polymerase reaction, in particular a polymerase chain reaction, as known in the art. The primer oligonucleotides are therefore designed to anneal to the template nucleic acids to form a double strand, following the Watson-Crick base pairing rules, and the length of these oligonucleotide primers will be selected such that they anneal at approximately the same temperature.

In said second step, an enzyme and the matching buffers are added to achieve cleavage of any present, contaminating amplificates that were generated in any of the preceding experiments. These amplificates will have the property that they comprise uracil bases instead of thymine bases, if generated in a polymerase reaction providing dUTPs instead of dTTPs. Therefore, not the sample nucleic acid at this step would be recognized and degraded by the enzyme, but only nucleic acids that were generated in preceding amplifications, the contaminating DNA that has to be removed before the next round of amplification.

It is particularly preferred that the enzyme employed in this second step is uracil-DNA-glycosylase (UNG). It is further preferred that said non-sulfonated uracil containing nucleic

acid degrading enzyme is thermolabile, in particular the DNA-glycosylase or the Endonuclease are thermolabile, respectively, and most particularly preferred the UNG is thermolabile.

5

In the third step, after enzymatic degradation, the composition of enzymes and buffer is subsequently inactivated, in that it is not capable of substantially cleaving any product of the subsequent amplification step. The non-sulfonated uracil containing nucleic acid degrading enzyme activity is terminated, in particular the DNA-glycosylase activity or the endonuclease activity is terminated, and most particularly preferred the UNG activity is terminated.

10

15

After the composition of non-sulfonated uracil containing nucleic acid degrading enzymes and buffer is inactivated, in that it is not capable of substantially degrading any product of the subsequent amplification step, the fourth step is performed, that is the desulfonation of the template nucleic acids. Prior to the amplification by a polymerase, a fourth step must be conducted, that is the desulfonation of the sulfonated template nucleic. Desulfonation may take place under alkaline conditions (as described in the art). Desulfonation however may also be catalysed by an increase in temperature under pH conditions as they are common to the PCR reaction.

20

25

It is a preferred embodiment of the invention that steps 3 and 4 are conducted simultaneously by a short increase of the incubation temperature of said premix, which results in deactivation of the non-sulfonated uracil containing nucleic acid degrading enzyme on the one hand and in thermal desulfonation of the template nucleic acid, on the other hand. This increase in the incubation temperature can also be suitable to transfer double-stranded DNA into single-stranded form enabling an amplification.

30

35

The sample nucleic acid may now be amplified in the next step using the set of primer oligonucleotides and a polymerase, while any cleaved contaminating DNA is essentially not amplified.

fied. The sample nucleic acid may be amplified, using a set of primer oligonucleotides and a polymerase, while the cleaved or degraded contaminating nucleic acid cannot be amplified. The amplified products may now be analysed and the methylation status in the genomic DNA may be deduced from the presence of an amplified product and/or from the analysis of the sequence within the amplified product.

This amplification may be carried out, in a particularly preferred embodiment of the invention, by means of a polymerase chain reaction, but also by other means of DNA amplification known in the art, like TMA (transcription mediated amplification), isothermal amplifications, rolling circle amplification, ligase chain reaction, and others.

The generated DNA fragments will then be analysed, concerning their presence, the amount, or their sequence properties or a combination thereof.

Therefore one embodiment of the invention is a method for providing a decontaminated template nucleic acid for polymerase based amplification reactions suitable for DNA methylation analysis, which is characterized by firstly, incubating a template nucleic acid with a bisulfite reagent containing solution, whereby the unmethylated cytosines within said nucleic acid are sulfonated and/or deaminated, and secondly, mixing the sulfonated and/or deaminated template nucleic acid with the components required for a polymerase mediated amplification reaction or an amplification based detection assay, and thirdly, adding to this mixture an enzyme with uracil-DNA-glycosylase activity and incubating the mixture, whereby nucleic acids containing non-sulfonated uracils are degraded, and fourthly, terminating the UNG activity, and fifthly, desulfonating the template nucleic acid. In a preferred embodiment of this invention the method is further characterized by a step 4 and 5 taking place simultaneously, by briefly incubating the mixture at an increased temperature, whereby the UNG activity is terminated, whereby desulfonation of the template nucleic acid takes place, and whereby the DNA is trans-

ferred from a double-stranded form into a single-stranded form suitable for amplification.

5 It is a further preferred embodiment of the method according to the invention wherein in a subsequent step 6 the template nucleic acid is amplified.

10 It is further preferred that upon termination of the non-sulfonated uracil containing nucleic acid degrading activity and desulfonation of the template nucleic acid a polymerase based amplification reaction is started and/or an amplification based assay is performed.

15 It is further preferred that the polymerase based amplification reaction is started by a brief incubation at increased temperature (heat activation).

In a preferred embodiment of the method the polymerase is a heat stable polymerase.

20 It is particularly preferred according to the invention that the polymerase mediated amplification or amplification based assay is performed in the presence of dUTPs instead of dTTPs.

25 In one preferred embodiment of the invention, the method is performed by adding an amount of units of the enzyme, which specifically degrades non-sulfonated uracil containing nucleic acids, in the second step that is required to degrade essentially all potential contaminating nucleic acids.

30 It is especially preferred that upon activation of the polymerase enzyme a polymerase based amplification reaction or an amplification based assay is performed.

35 It is further preferred that upon activation of the polymerase enzyme a polymerase based amplification reaction or amplification based assay is performed in the presence of dUTPs instead of dTTPs.

It is further preferred that this assay is a real time assay.

In a particularly preferred embodiment, the sample DNA is obtained from serum or other body fluids of an individual. It is further particularly preferred, that the DNA samples are obtained from cell lines, tissue embedded in paraffin, for example tissue from eyes, intestine, kidneys, brain, heart, prostate, lungs, breast or liver, histological slides, body fluids and all possible combinations thereof. The term body fluids is meant to comprise fluids such as whole blood, blood plasma, blood serum, urine, sputum, ejaculate, semen, tears, sweat, saliva, lymph fluid, bronchial lavage, pleural effusion, peritoneal fluid, meningeal fluid, amniotic fluid, glandular fluid, fine needle aspirates, nipple aspirate fluid, spinal fluid, conjunctival fluid, vaginal fluid, duodenal juice, pancreatic juice, bile, stool and cerebrospinal fluid. It is especially preferred that said body fluids are whole blood, blood plasma, blood serum, urine, stool, ejaculate, bronchial lavage, vaginal fluid and nipple aspirate fluid.

In a particularly preferred embodiment of the invention, the chemical treatment is conducted with a bisulfite (= disulfite, hydrogen sulfite). It is again preferred that the chemical treatment is conducted after embedding the DNA in agarose, or that it is conducted in the presence of a denaturing agent and/or a radical scavenger.

The following methylation detection assays are all preferred embodiments of the invention when performed subsequently to the steps of the method according to the invention:

Methylation Assay Procedures. Various methylation assay procedures are known in the art, and can be used in conjunction with the present invention. These assays allow for determination of the methylation state of one or a plurality of CpG dinucleotides (e.g., CpG islands) within a DNA sequence. Such assays involve, among other techniques, DNA sequencing of bisulfite-treated DNA, and a number of PCR based methylation assays, some of them - known as COBRA, MS-SNuPE, MSP, nested

MSP, HeavyMethyl and MethyLight - are described in more detail now.

5 BISULFITE SEQUENCING. DNA methylation patterns and 5-methylcytosine distribution can be analyzed by sequencing analysis of a previously amplified fragment of the bisulfite treated genomic DNA, as described by Frommer et al. (Frommer et al. Proc. Natl. Acad. Sci. USA 89:1827-1831, 1992). As the bisulfite treated DNA is amplified before sequencing, the amplification procedure according to the invention may be used
10 in combination with this detection method.

COBRA. COBRA analysis is a quantitative methylation assay useful for determining DNA methylation levels at specific gene
15 loci in small amounts of genomic DNA (Xiong & Laird, Nucleic Acids Res. 25:2532-2534, 1997). Briefly, restriction enzyme digestion is used to reveal methylation-dependent sequence differences in PCR products of sodium bisulfite-treated DNA. Methylation-dependent sequence differences are first introduced into the genomic DNA by standard bisulfite treatment according to the procedure described by Frommer et al. (Proc.
20 Natl. Acad. Sci. USA 89:1827-1831, 1992) or as described by Olek et al (Olek A, Oswald J, Walter J. (1996) Nucleic Acids Res. 24: 5064-6). PCR amplification of the bisulfite converted DNA is then performed using methylation unspecific primers
25 followed by restriction endonuclease digestion, gel electrophoresis, and detection using specific, labeled hybridization probes. Methylation levels in the original DNA sample are represented by the relative amounts of digested and undigested PCR product in a linearly quantitative fashion across a wide
30 spectrum of DNA methylation levels. In addition, this technique can be reliably applied to DNA obtained from microdissected paraffin-embedded tissue samples. Typical reagents (e.g., as might be found in a typical COBRA-based kit) for
35 COBRA analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); restriction enzyme and appropriate buffer; gene-hybridization oligo; control hybridization oligo; kinase labeling kit for oligo probe; and radioactive nucleotides. Addi-

tionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kits (e.g., precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

5

Additionally, restriction enzyme digestion of PCR products amplified from bisulfite-converted DNA is also used, in the method described by Sadri & Hornsby (Nucl. Acids Res. 24:5058-5059, 1996)

10

The bisulfite conversion and amplification procedure according to the invention may be used in combination with this detection method.

15

Ms-SNuPE (Methylation-sensitive Single Nucleotide Primer Extension). The Ms-SNuPE technique is a quantitative method for assessing methylation differences at specific CpG sites based on bisulfite treatment of DNA, followed by single-nucleotide primer extension (Gonzalzo & Jones, Nucleic Acids Res.

20

25:2529-2531, 1997). Briefly, genomic DNA is reacted with sodium bisulfite to convert unmethylated cytosine to uracil while leaving 5-methylcytosine unchanged. Amplification of

the desired target sequence is then performed using *PCR primers* specific for bisulfite-converted DNA, and the resulting product is isolated and used as a template for methylation

25

analysis at the CpG site(s) of interest. Small amounts of DNA can be analyzed (e.g., microdissected pathology sections), and it avoids utilization of restriction enzymes for determining the methylation status at CpG sites.

30

Typical reagents (e.g., as might be found in a typical Ms-SNuPE-based kit) for Ms-SNuPE analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); optimized PCR buffers and deoxynucleotides; gel extraction kit; positive control primers; Ms-SNuPE primers for specific gene; reaction buffer (for the Ms-SNuPE reaction); and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kit

35

(e.g., precipitation, ultrafiltration, affinity column);
desulfonation buffer; and DNA recovery components.

5 The bisulfite conversion and amplification procedure according
to the invention may be used in combination with this detec-
tion method.

10 MSP. MSP (methylation-specific PCR) allows for assessing the
methylation status of virtually any group of CpG sites within
a CpG island, independent of the use of methylation-sensitive
restriction enzymes (Herman et al. Proc. Natl. Acad. Sci. USA
93:9821-9826, 1996; US Patent No. 5,786,146). Briefly, DNA is
modified by sodium bisulfite converting all unmethylated, but
not methylated cytosines to uracil, and subsequently amplified
15 with primers specific for methylated versus unmethylated DNA.

MSP primer pairs contain at least one primer, which hybridizes
to a bisulfite treated CpG dinucleotide. Therefore, the se-
quence of said primers comprises at least one CpG dinucleo-
20 tide. MSP primers specific for non-methylated DNA contain a
"T" at the 3' position of the C position in the CpG. Prefera-
bly, therefore, the base sequence of said primers is required
to comprise a sequence having a length of at least 9 nucleo-
tides which hybridizes to the bisulfite converted nucleic acid
25 sequence, wherein the base sequence of said oligomers com-
prises at least one CpG dinucleotide. MSP requires only small
quantities of DNA, is sensitive to 0.1 % methylated alleles of
a given CpG island locus, and can be performed on DNA ex-
tracted from paraffin-embedded samples. Typical reagents
30 (e.g., as might be found in a typical MSP-based kit) for MSP
analysis may include, but are not limited to: methylated and
unmethylated PCR primers for specific gene (or methylation-
altered DNA sequence or CpG island), optimized PCR buffers and
deoxynucleotides, and specific probes.

35 The bisulfite conversion and amplification procedure according
to the invention may be used in combination with this detec-
tion method.

NESTED MSP (Belinsky and Palmisano in US application 20040038245). Considering the apparent conflict of requiring high specificity of the MSP primer to sufficiently differentiate between CG and TG positions but allowing for a mismatch in order to create a unique restriction site it is preferred to use an amended version of MSP, known as nested MSP, as described in WO 02/18649 and US patent application 20040038245 by Belinsky and Palmisano. This method to detect the presence of gene-specific promoter methylation, comprises the steps of: expanding the number of copies of the genetic region of interest by using a polymerase chain reaction to amplify a portion of said region where the promoter methylation resides, thereby generating an amplification product; and using an aliquot of the amplification product generated by the first polymerase chain reaction in a second, methylation-specific, polymerase chain reaction to detect the presence of methylation. In other words a non methylation specific PCR is performed prior to the methylation specific PCR. The bisulfite conversion and amplification procedure according to the invention may be used in combination with this detection method.

HEAVYMETHYL. (WO 02/072880; Cottrell SE et al. Nucleic Acids Res. 2004 Jan 13;32(1):e10) A further preferred embodiment of the method comprises the use of blocker oligonucleotides. In the HeavyMethyl assay blocking probe oligonucleotides are hybridized to the bisulfite treated nucleic acid concurrently with the PCR *primers*. PCR amplification of the nucleic acid is terminated at the 5' position of the blocking probe, such that amplification of a nucleic acid is suppressed where the complementary sequence to the blocking probe is present. The probes may be designed to hybridize to the bisulfite treated nucleic acid in a methylation status specific manner. For example, for detection of methylated nucleic acids within a population of unmethylated nucleic acids, suppression of the amplification of nucleic acids which are unmethylated at the position in question would be carried out by the use of blocking probes comprising a 'CpA' or 'TpA' at the position in question, as opposed to a 'CpG' if the suppression of amplification of methylated nucleic acids is desired.

For PCR methods using blocker oligonucleotides, efficient disruption of polymerase-mediated amplification requires that blocker oligonucleotides not be elongated by the polymerase.

5 Preferably, this is achieved through the use of blockers that are 3'-deoxyoligonucleotides, or oligonucleotides derivatized at the 3' position with other than a "free" hydroxyl group. For example, 3'-O-acetyl oligonucleotides are representative of a preferred class of blocker molecule.

10

Additionally, polymerase-mediated decomposition of the blocker oligonucleotides should be precluded. Preferably, such preclusion comprises either use of a polymerase lacking 5'-3' exonuclease activity, or use of modified blocker oligonucleotides

15

having, for example, thioate bridges at the 5'-termini thereof that render the blocker molecule nuclease-resistant. Particular applications may not require such 5' modifications of the blocker. For example, if the blocker- and primer-

20

binding sites overlap, thereby precluding binding of the primer (e.g., with excess blocker), degradation of the blocker oligonucleotide will be substantially precluded. This is because the polymerase will not extend the primer toward, and through (in the 5'-3' direction) the blocker—a process that normally results in degradation of the hybridized blocker oligonucleotide.

25

A particularly preferred blocker/PCR embodiment, for purposes of the present invention and as implemented herein, comprises the use of peptide nucleic acid (PNA) oligomers as blocking oligonucleotides. Such PNA blocker oligomers are ideally suited, because they are neither decomposed nor extended by the polymerase.

30

Preferably, therefore, the base sequence of said blocking oligonucleotide is required to comprise a sequence having a length of at least 9 nucleotides which hybridizes to the chemically treated nucleic acid sequence, wherein the base sequence of said oligonucleotides comprises at least one CpG, TpG or CpA dinucleotide.

35

The bisulfite conversion and amplification procedure according to the invention may be used in combination with this detection method.

5

Preferably, real-time PCR assays are performed specified by the use of such primers according to the invention. Real-time PCR assays can be performed with methylation specific primers (MSP-real time) as methylation-specific PCR ("MSP"; as described above), or with non-methylation specific primers in
10 presence of methylation specific blockers (HM real-time) ("HEAVYMETHYL", as described above). Real-time PCR may be performed with any suitable detectably labelled labeled probes. For details see below.

15

Both of these methods (MSP or HM) can be combined with the detection method known as MethyLight™ (a fluorescence-based real-time PCR technique) (Eads et al., Cancer Res. 59:2302-2306, 1999), which generally increases the specificity of the
20 signal generated in such an assay. Whenever the real-time probe used is methylation specific in itself, the technology shall be referred to as MethyLight™, a widely used method.

Another assay makes use of the methylation specific probe, the
25 so called "QM" (quantitative methylation) assay. A methylation unspecific, therefore unbiased real-time PCR amplification is performed which is accompanied by the use of two methylation specific probes (MethyLight™) one for the methylated and a second for the unmethylated amplificate. That way two signals
30 are generated which can be used to a) determine the ratio of methylated (CG) to unmethylated (TG) nucleic acids, and at the same time b) the absolute amount of methylated nucleic acids can be determined, when calibrating the assay with a known amount of control DNA.

35

MethyLight™. The MethyLight™ assay is a high-throughput quantitative methylation assay that utilizes fluorescence-based real-time PCR (TaqMan.™) technology that requires no further manipulations after the PCR step (Eads et al., Cancer Res.

59:2302-2306, 1999). Briefly, the MethyLight™ process begins with a mixed sample of genomic DNA that is converted, in a sodium bisulfite reaction, to a mixed pool of methylation-dependent sequence differences according to standard procedures (the bisulfite process converts unmethylated cytosine residues to uracil). Fluorescence-based PCR is then performed either in an "unbiased" (with primers that do not overlap known CpG methylation sites) PCR reaction, or in a "biased" (with PCR primers that overlap known CpG dinucleotides) reaction. Sequence discrimination can occur either at the level of the amplification process or at the level of the fluorescence detection process, or both.

The MethyLight™ assay may be used as a quantitative test for methylation patterns in the genomic DNA sample, wherein sequence discrimination occurs at the level of probe hybridization. In this quantitative version, the PCR reaction provides for unbiased amplification in the presence of a fluorescent probe that overlaps a particular putative methylation site. An unbiased control for the amount of input DNA is provided by a reaction in which neither the primers, nor the probe overlap any CpG dinucleotides. Alternatively, a qualitative test for genomic methylation is achieved by probing of the biased PCR pool with either control oligonucleotides that do not "cover" known methylation sites (a fluorescence-based version of the "MSP" technique), or with oligonucleotides covering potential methylation sites.

The MethyLight™ process can be used with a "TaqMan®" probe in the amplification process. For example, double-stranded genomic DNA is treated with sodium bisulfite and subjected to one of two sets of PCR reactions using TaqMan® probes; e.g., with either biased primers and TaqMan® probe, or unbiased primers and TaqMan® probe. The TaqMan® probe is dual-labeled with fluorescent "reporter" and "quencher" molecules, and is designed to be specific for a relatively high GC content region so that it melts out at about 10 °C higher temperature in the PCR cycle than the forward or reverse primers. This allows the TaqMan® probe to remain fully hybridized during the PCR anneal-

ing/extension step. As the Taq polymerase enzymatically synthesizes a new strand during PCR, it will eventually reach the annealed TaqMan[®] probe. The Taq polymerase 5' to 3' endonuclease activity will then displace the TaqMan[®] probe by digesting it to release the fluorescent reporter molecule for quantitative detection of its now unquenched signal using a real-time fluorescent detection system.

Variations on the TaqMan[™] detection methodology that are also suitable for use with the described invention include the use of dual-probe technology (LightCycler[™]) or fluorescent amplification primers (Sunrise[™] technology). Both these techniques may be adapted in a manner suitable for use with bisulfite treated DNA, and moreover for methylation analysis within CpG dinucleotides.

Typical reagents (e.g., as might be found in a typical MethyLight[™]-based kit) for MethyLight[™] analysis may include, but are not limited to: PCR primers for specific bisulfite sequences, i.e. bisulfite converted genetic regions (or bisulfite converted DNA or *bisulfite converted CpG islands*); probes (e.g. TaqMan[®] or LightCycler[™]) specific for said amplified bisulfite converted sequences; optimized PCR buffers and deoxynucleotides; and a polymerase, such as Taq polymerase.

The bisulfite conversion and amplification procedure according to the invention may be used in combination with this detection method.

The fragments obtained by means of the amplification can carry a directly or indirectly detectable label. Preferred are labels in the form of fluorescence labels, radionuclides, or detachable molecule fragments having a typical mass, which can be detected in a mass spectrometer. Where said labels are mass labels, it is preferred that the labeled amplicates have a single positive or negative net charge, allowing for better detectability in the mass spectrometer. The detection may be carried out and visualized by means of, e.g., matrix assisted

laser desorption/ionization mass spectrometry (MALDI) or using electron spray mass spectrometry (ESI).

Matrix Assisted Laser Desorption/Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas & Hillenkamp, Anal Chem., 60:2299-301, 1988). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionized by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones. MALDI-TOF spectrometry is well suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut & Beck, Current Innovations and Future Trends, 1:147-57, 1995). The sensitivity with respect to nucleic acid analysis is approximately 100-times less than for peptides, and decreases disproportionately with increasing fragment size. Moreover, for nucleic acids having a multiply negatively charged backbone, the ionization process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity between peptides and nucleic acids has not been reduced. This difference in sensitivity can be reduced, however, by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. For example, phosphorothioate nucleic acids, in which the usual phosphates of the backbone are substituted with thiophosphates, can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut & Beck, Nucleic Acids Res. 23: 1367-73, 1995). The coupling of a charge tag to this modified DNA results in an increase in MALDI-TOF sensitivity to the same level as that found for peptides.

The amplicates may also be further detected and/or analysed by means of oligonucleotides constituting all or part of an "array" or "DNA chip" (i.e., an arrangement of different oligonucleotides and/or PNA-oligomers bound to a solid phase).

5 Such an array of different oligonucleotide- and/or PNA-oligomer sequences can be characterized, for example, in that it is arranged on the solid phase in the form of a rectangular or hexagonal lattice. The solid-phase surface may be composed of silicon, glass, polystyrene, aluminum, steel, iron, copper,
10 nickel, silver, or gold. Nitrocellulose as well as plastics such as nylon, which can exist in the form of pellets or also as resin matrices, may also be used. An overview of the Prior Art in oligomer array manufacturing can be gathered from a special edition of Nature Genetics (Nature Genetics Supplement, Volume 21, January 1999, and from the literature cited
15 therein). Fluorescently labeled probes are often used for the scanning of immobilized DNA arrays. The simple attachment of Cy3 and Cy5 dyes to the 5'-OH of the specific probe are particularly suitable for fluorescence labels. The detection of
20 the fluorescence of the hybridized probes may be carried out, for example, via a confocal microscope. Cy3 and Cy5 dyes, besides many others, are commercially available.

The bisulfite conversion and amplification procedure according
25 to the invention may be used in combination with this detection method.

A particular preferred embodiment of the invention is a method for providing a decontaminated nucleic acid for hybridisation
30 on a DNA-Array, preferably an Oligonucleotide-Array, suitable for DNA methylation analysis.

Of course, a particular preferred embodiment of the invention is also a improved method for bisulfite conversion of DNA.
35 Thereby non-methylated cytosines are converted to uracil while methylated cytosines remain unchanged. According to this embodiment, a nucleic acid is incubated with a bisulfite reagent containing solution, whereby the unmethylated cytosines within said nucleic acid are sulfonated and/or deaminated but not yet

desulfonated as this is described above. Afterwards the sulfonated and/or deaminated template nucleic acid is mixed with components required for a polymerase mediated amplification reaction or an amplification based detection assay. Thereafter
5 the template nucleic acid is desulfonated by briefly incubating the mixture at an increased temperature. Subsequently the desulfonated template nucleic acid is amplified. In a particularly preferred variant the polymerase based amplification reaction is started by a brief incubation at increased temperature (heat activation). Simultaneously, this brief incubation
10 at increased temperature serves to desulfonate the sulfonated and/or deaminated template nucleic acid. Furthermore it is particularly preferred that the polymerase is a heat stable polymerase.

15 This particular embodiment has the advantage in comparison to known methods of bisulfite treatment that the purification step after bisulfite treatment becomes dispensable. This is a simplification which results in reduction of costs and handling effort, minimises loss of bisulfite treated DNA and is
20 also time saving. Therefore the use of this embodiment is preferred if DNA samples are treated with bisulfite and subsequently are amplified. This is especially preferred if large amount of samples are analysed. The use of this embodiment is
25 further preferred with regard to sensitive detection methods for DNA methylation analysis like COBRA, MS-SNuPE, MSP, nested MSP, HeavyMethyl and MethyLight.

Furthermore, the invention regards to a test kit for the realisation of the method according to the invention with a component containing bisulfite, for example a reagent or solution containing bisulfite, and a component containing an enzymatic activity. This enzymatic activity specifically degrades DNA containing non-sulfonated uracil. In particular this enzymatic
30 activity is an activity of a DNA-glycosylase and/or an endonuclease, preferentially this enzymatic activity is an uracil-DNA-glycosylase, and more preferentially this enzymatic activity is uracil-N-DNA-Glycosylase (UNG). The added degrading enzyme activity is characterized by its ability to cause spe-

cifically apyrimidinic sites and/or one or more nicks adjacent to a non-sulfonated uracil base. In any case this will result in a block of the replication by DNA polymerase. In a particular test kit the enzymatic activity is characterized by cleav-
5 ing the uracil base from the phosphodiester backbone of non-sulfonated uracil containing nucleic acid, but has no effect on sulfonated uracil containing nucleic acid or on thymine containing nucleic acid, that does not contain uracil. The re-
10 sulting apyrimidinic sites block replication by DNA polymerases, and are very labile to acid/base hydrolysis.

A further test kit comprises one or more of the additional components. This can be:

15 - one or more denaturing reagent and/or solution, for example: dioxane or diethylene glycol dimethyl ether (DME) or any substance, which is suitable as described in WO 05/038051,

20 - one or more scavenger, for example 6-hydroxy-2,5,7,8-tetramethylchromane 2-carboxylic acid or other scavengers as described in WO 01/98528 or WO 05/038051,

25 - one or more primers, which are suitable for the amplification of one or more DNA amplificates, amongst others the primer or primers can be modified, for example with a quencher and/or a label for detection as well known by a person skilled in the art like the dye FAM or the quencher BHQ black hole or dabcy1,

30 - one or more probes, which can be any probe, which can be used to specifically record the amplification of one or more amplificates for example in a real-time-assay, amongst others the probe or probes can be modified, for example with a quencher and/or a label for detection as well known by a person
35 skilled in the art like the dye FAM or the quencher BHQ black hole or dabcy1,

- one or more blockers, which are nucleic acids and can be used to block the binding of a specific primer or the replica-

tion by DNA polymerase, amongst others the blocker or blockers can be modified, for example with a quenscher and/or a label for detection as well known by a person skilled in the art like the dye FAM or the quencher BHQ black hole or dabcy1,

5

- one or more reaction buffers, which are suitable for a bisulfite treatment and/or a PCR reaction,

10

- nucleotides, which can be dATP, dCTP, dUTP and dGTP or any derivative of these nucleotides,

- $MgCl_2$ as a substance or in solution and/or any other magnesium salt, which can be used to carry out a DNA polymerase replication,

15

- DNA polymerase, for example Taq polymerase or any other polymerase with or without proof-reading activity, - dye or quencher, which can be used for the detection of the amplificates as known in the art, for example an intercalating dye like SYBR Green or a dye for linkage to a primer or probe or blocker like the dye FAM or the quencher BHQ black hole or dabcy1, and/or

20

- any reagent, solution, device and/or instruction which is useful for realisation of an assay according to the invention.

25

The methods and test kits disclosed here are preferable used for the diagnosis and/or prognosis of adverse events for patients or individuals, whereby diagnosis means diagnose of a adverse event, a predisposition for a adverse event and/or a progression of a adverse events. These adverse events belong to at least one of the following categories: undesired drug interactions; cancer diseases; CNS malfunctions, damage or disease; symptoms of aggression or behavioral disturbances; clinical, psychological and social consequences of brain damage; psychotic disturbances and personality disorders; dementia and/or associated syndromes; cardiovascular disease, malfunction or damage; malfunction, damage or disease of the gastrointestinal tract; malfunction, damage or disease of the

30

35

respiratory system; lesion, inflammation, infection, immunity and/or convalescence; malfunction, damage or disease of the body as an abnormality in the development process; malfunction, damage or disease of the skin, of the muscles, of the connective tissue or of the bones; endocrine and metabolic malfunction, damage or disease; headaches or sexual malfunction.

The methods and test kits also serve in a particularly preferred manner for distinguishing cell types, tissues or for investigating cell differentiation. They serve in a particularly preferred manner for analysing the response of a patient to a drug treatment.

In another preferred manner the methods and test kits of the invention can also be used to characterize the DNA methylation status in that positions are methylated or non-methylated compared to normal conditions if a single defined disease exists. In a particular preferred manner they can serve for identifying an indication-specific target, wherein a template nucleic acid is treated with bisulfite and UNG enzyme activity, and wherein an indication-specific target is defined as differences in the DNA methylation status of a DNA derived from a diseased tissue in comparison to a DNA derived from a healthy tissue. These tissue samples can originate from diseased or healthy patients or from diseased or healthy adjacent tissue of the same patient.

In a particular preferred manner the indication specific target is a protein, peptide or enzyme, and in particular a per se known modulator of the coded protein, peptide or enzyme is assigned with the specific indication of the diseased tissue. In a particular preferred manner this modulator serves for preparing a pharmaceutical composition with a specific indication, in particular a specific cancer indication.

In a particular preferred manner the enzyme UNG serves as an enzyme for generation of contamination free nucleic acids for methylation analysis.

Brief description of the drawings

5 Figure 1:

Figure 1 describes the complete conversion of unmethylated cytosine to uracil, so called bisulfite conversion.

10 The first step of this reaction takes place when unmethylated cytosine bases are contacted with hydrogensulfite at a pH around 5. The sulfonation takes place at position 6 of the cyclic molecule (C6 position).

15 The second step is the deamination that takes place rather spontaneously in aqueous solution and thereby converts cytosine sulfonate into uracil sulfonate.

The third step is the desulfonation step, which takes place in alkaline conditions, resulting in uracil.

20

Figure 2:

Figure 2 is a plot of real time amplification of methylated DNA of the GSTP1 gene from desulfonated bisulfite converted DNA, according to the state of the art. The Y-axis shows the fluorescence signal measured in channel F2 normalized against channel F1 at each cycle (X-axis). 10 ng, 1 ng respective 25 0,1 ng bisulfite treated methylated DNA were added to the reaction. No signals were determined using the reaction mix containing Uracil-DNA-glycosylase (labeled in open circles) indicating a complete degradation of the bisulfite converted DNA. 30 Amplification occurs only in absence of Uracil-DNA-glycosylase (labeled in rectangles). No template control is marked as solid line.

35 Figure 3:

Figure 3 is a plot of real time amplification of methylated DNA of the GSTP1 gene from bisulfite converted DNA according to the claimed new method without desulfonation. The Y-axis shows the fluorescence signal measured in channel F2 normal-

ized against channel F1 at each cycle (X-axis). 10 ng, 1 ng
respective 0,1 ng bisulfite treated methylated DNA were added
to the reaction. The signals generated from reaction without
Uracil-DNA-glycosylase are labeled with circles. No signifi-
cant difference in amplification was determined from the reac-
tion containing Uracil-DNA-glycosylases (labeled in triangles)
indicating that 6-Sulfon-Uracil containing DNA is not a tem-
plate for UNG. No template control is marked as solid line.

Figure 4:

Figure 4 demonstrates of the efficient degradation of uracil
containing DNA. The plot shows the reamplification of PCR
products containing uracil. The Y-axis shows the fluorescence
signal measured in channel F2 normalized against channel F1 at
each cycle (X-axis). 10^5 copies were added to the reaction. The
signals determined from reaction without Uracil-DNA-
glycosylase are labeled diamonds showing a high efficient
reamplification. The reaction containing Uracil-DNA-
glycosylases results in dramatically increased crossing points
(labeled in stars) indicating a strong degradation of uracil
containing PCR products by UNG. No template control is marked
as solid line.

Figure 5:

Figure 5 shows a correlation plot of the results obtained in
example 3 by the standard workflow and the method according to
the invention (carry over prevention). Every symbol represents
a single sample: quadrates tumor tissues, triangles normal ad-
jacent tissues. The percentage of methylation determined ac-
cording to the standard workflow (x-axis) or to the method ac-
cording to the invention (y-axis) is indicated for each sam-
ple.

The method according to the invention has led only in 2 out of
24 samples to a different methylation percentage as the stan-
dard workflow. This means that although the samples treated
according to the method of the invention were contaminated
with uracil containing TPEF amplicons only DNA of the samples
served as a template for amplification of the TPEF amplicon in

nearly all cases. In case of the said two samples, the differing results occurred presumable because of the low methylation percentage of the DNA (smaller than 0.2 %).

EXAMPLES:

Example 1:

Amplification of methylated DNA of the GSTP1 gene (also known
5 as GST-pi gene) wherein human DNA containing sulfonated
uracils served as template

The use of uracil-DNA-glycosylase is a method well known in
the art to avoid false positive results in polymerase based
10 amplification methods, caused by cross contamination by previ-
ously amplified products (Pang J., Mol Cell Probes. 1992
Jun;6(3):251-6). This method is however not applicable for po-
lymerase based amplification methods which have the purpose to
detect uracil bases within the given template. This is the
15 case in DNA methylation analysis, wherein one way to detect
the difference between methylated and unmethylated cytosines
is to mirror these differences into the difference between cy-
tosine and uracil, which is facilitated by the widely spread
use of common bisulfite conversion methods. These have the ef-
20 fect to convert unmethylated cytosines into uracils whereas
methylated cytosines remain cytosines. Therefore in subsequent
amplification reactions to detect methylation patterns the
template contains uracils.

25 In the following example it was shown that the method accord-
ing to the invention allows Uracil-DNA-glycosylase (UNG) based
technique for carry over prevention of bisulfite converted
DNA, without loss of the critical information, which bases
were unmethylated and which were methylated. To achieve this
30 the following steps were carried out:

Two nucleic acid samples, containing 1,5µg GpGenome™ Universal
Methylated DNA (Chemicon International) diluted in 100µl water
were mixed with 354 µl of bisulfite solution (5.89 mol/l) and
35 146 µl of dioxane containing a radical scavenger (6-hydroxy-
2,5,7,8-tetramethylchromane 2-carboxylic acid, 98.6 mg in
2.5 ml of dioxane). The reaction mixture was denatured for
3 min at 99 °C and subsequently incubated with the following
temperature program for a total of 5 h: 30 min at 50 °C; a

first thermospike (99.9 °C) for 3 min; 1.5 h at 50 °C; a second thermospike (99.9 °C) for 3 min; 3 h at 50 °C. One of the reaction mixtures served as a control whereas the other was treated according to the invention. The reaction mixtures of both the control and the test reaction were subsequently purified by ultrafiltration by means of a Millipore Microcon column. The purification was conducted essentially according to the manufacturer's instructions. For this purpose, the reaction mixture was mixed with 200 µl of water, loaded onto the ultrafiltration membrane, centrifuged for 15 min and subsequently washed with water. The DNA remains on the membrane in this treatment. For the control sample an alkaline desulfonation was performed according to the methods, which are state of the art (see for example US 20040152080, 20040115663, WO 2004/067545). For this purpose, 100 µl of a 0.2 mol/l NaOH was added and incubated for 10 min. For the other sample this desulfonation step was replaced by adding 100µl water. A centrifugation (10 min) was then conducted, followed by a final washing step with water. After this, the DNA was eluted. For this purpose, the membrane was mixed for 10 minutes with 50 µl of warm 1 x TE buffer (50 °C) adjusted to pH 7. The membrane was turned over according to the manufacturer's instructions. Subsequently a repeated centrifugation was conducted, with which the DNA was removed from the membrane.

Subsequently the DNA was stored at 4 °C for 12 h and then used as template in a PCR reaction.

By stopping the chemical reaction after sulfonation all unmethylated cytosines are converted into C6 sulfonated uracils (5,6-Dihydro-6-sulfonyl-uracil) and methylated cytosines remain unchanged. However after a complete desulfonation, as described in the art, all unmethylated cytosines are converted in uracil and methylated cytosines remain unchanged.

As a control for the UNG activity 10^5 copies of a uracil containing PCR product were added to the reaction premix, generated by use of the same primers but under presence of dUTP instead of dTTP. Reamplification took place when UNG was absent,

with the expected efficiency of a crossing point of 22,6. However when UNG was present in the PCR - mix the crossing points reached only a value of 35,8 (Figure 4). This difference demonstrates nicely the efficient degradation of PCR products by UNG resulting from the cleavage of uracils out of the DNA. In this example it was shown that desulfonated conventionally bisulfite treated DNA is also degraded by UNG (Figure 2). However, not desulfonated bisulfite treated DNA does not work as substrate for UNG and serves - even after preincubation with UNG - as working template in an amplification reaction (Figure 3).

In the example the desulfonation of template DNA required for the successful amplification took place during the initial denaturing phase of the PCR reaction at 95 °C. Only precondition for this step is an alkaline pH, such as given in the utilized PCR buffer. Simultaneously the UNG activity is terminated and is hence not capable of cleaving or degrading newly generated PCR product anymore.

In this example three different concentrations (10 ng, 1 ng and 0.1 ng) of each desulfonated and sulfonated (containing 6-sulfonated 5,6-dihydro-uracils) DNA were used as templates in two different Hot-Start PCR reactions.

In one case the reaction mix contained 0.2 Units UNG, in the other case no UNG was added. PCR reactions were performed in the LightCycler in 20µl reaction volume and contained:

10 µl of template DNA (in different concentrations)
2 µl of FastStart LightCycler Mix for Hybridization probes (Roche Diagnostics)
3.5 mM MgCl₂ (Roche Diagnostics)
0.30 µM forward primer (SeqID-1, TIB-MolBiol)
0.30 µM reverse primer (SeqID-2, TIB-MolBiol)
0.15 µM Probe1 (SeqID-3, TIB-MolBiol)
0.15 µM Probe2 (SeqID-4, TIB-MolBiol)
optional 0.2 Unit Uracil-DNA-Glycosylase (Roche Diagnostics)

The temperature-time-profile was programmed as follows:

Pre-incubation (UNG active) 15 min by 25 °C

Activation of polymerase: 20 min by 95 °C

50 temperature cycles: 10 sec by 95 °C

5 30 sec at 56 °C

10 sec at 72 °C

Finally the reaction is cooled down to 35 °C.

10 The primers (Seq ID 1, Seq ID 2) used amplify a 123 bp long
fragment of the GSTP1 gene (Seq ID 5, nt 1184 to nt 1304 in
Genbank Accession X08058). By utilizing sequence specific hy-
bridization probes (SeqID 3, SeqID 4) the amplification rate
was detected in a Real Time PCR. Data interpretation was car-
ried out via the LightCycler Software in channel F2/F1.
15

The crossing point (Cp) was generated automatically by employ-
ing the method "Second Derivative Maximum" (Table 1).

20 The results of the experiment are summarized in table 1. The
reamplification of 10^5 copies of uracil containing amplicons
results in CT of 22,6 without UNG and 35,8 with UNG. The CT
delay of 13 cycles demonstrates the efficient degradation of
uracil containing template by the glycosylase. Also desul-
fonated bisulfite converted DNA was degraded by UNG and no am-
plification was measurable in the reaction with UNG. In the
25 reaction without UNG the 10, 1, and 0,1 ng DNA was detected at
CT of 28,5 / 31,7 / 33,8. In contrast to this, sulfonated DNA,
prepared according to the invention, was amplified in both
cases, without and with Uracil-DNA-Glycosylase with almost the
30 same efficiency and were detected at CT of 29,3 / 32,2 / 34,1
and 29,9 / 32,7 / 34,8 respectively.

Table 1:

DNA	Template DNA in ng	Crossing point of reaction without UNG	Crossing point of reaction with 0,2 Unit UNG added
PCR Amplicons containing Uracil	10 ⁵ copies	22,6	35,8
desulfonated DNA	10	28,5	no signal
	1	31,7	no signal
	0,1	33,8	no signal
sulfonated DNA	10	29,3	29,9
	1	32,2	32,7
	0,1	34,1	34,8

Table 2: Sequences of Oligonucleotides

SeqID	Name	Sequence
Seq ID 1	GSTP1.10F1	GGGAttAtttTTATAAGGtT
Seq ID 2	GSTP1.10R5	TaCTaaaAaCTCTaAaCCCCATC
Seq ID 3	GSTP1.10-fluo1	TTCGtCGtCGtAGTtTTCGtt-Fluo
Seq ID 4	GSTP1.10-red1	red640-tAGTGAGTACGCGCGGtt-PH
Seq ID 5	GSTP1 amplicon	5'GGGAttAtttTTATAAGGtTCGGAGGtCGCGAGGttT TCGtTGGAGTTTCGtCGtCGtAGTtTTCGttAttAGTGA GTACGCGCGGttCGCGTtttCGGGGATGGGGTtTAGAG- tTtttAGtA

5

Fluo=fluoresceine label, red640=LightCycler fluorescence label for channel F2, PH=3'OH-Phosphorylation. Small written t's point to converted cytosines by bisulfite treatment, respectively small a's point to the complementary adenosine bases in the reverse complement synthesized strand.

10

Example 2:

In this experiment the stability of the sulfonated nucleic bases in the presence of UNG activity was analyzed when stored at 4 °C or 40 °C. Again, two nucleic acid samples, containing 1,5 µg GpGenome™ Universal Methylated DNA (Chemicon International) diluted in 100µl water were mixed with 354 µl of bisulfite solution (5.89 mol/l) and 146 µl of dioxane containing a radical scavenger (6-hydroxy-2,5,7,8-tetramethylchromane 2-

20

carboxylic acid, 98.6 mg in 2.5 ml of dioxane). The reaction mixture was denatured for 3 min at 99 °C and subsequently incubated with the following temperature program for a total of 5 h: 30 min at 50 °C; a first thermospike (99.9 °C) for 3 min; 1.5 h at 50 °C; a second thermospike (99.9 °C) for 3 min; 3 h at 50 °C. One of the reaction mixtures served as a control whereas the other was treated according to the invention. The reaction mixtures of both the control and the test reaction were subsequently purified by ultrafiltration by means of a Millipore Microcon column. The purification was conducted essentially according to the manufacturer's instructions. For this purpose, the reaction mixture was mixed with 200 µl of water, loaded onto the ultrafiltration membrane, centrifuged for 15 min and subsequently washed with water. The DNA remains on the membrane in this treatment. For the control sample an alkaline desulfonation was performed according to the methods, which are state of the art (see for example US 20040152080, 20040115663, WO 2004/067545) (named 'desulfonated' in table 3). For this purpose, 100 µl of a 0.2 mol/l NaOH was added and incubated for 10 min. For the other sample this desulfonation step was replaced by adding 100 µl water (named 'sulfonated' in table 3). A centrifugation (10 min) was then conducted, followed by a final washing step with water. After this, the DNA was eluted. For this purpose, the membrane was mixed for 10 minutes with 50 µl of warm 1 x TE buffer (50 °C) adjusted to pH7. The membrane was turned over according to the manufacturer's instructions. Subsequently a repeated centrifugation was conducted, with which the DNA was removed from the membrane.

Subsequently the DNA was divided in aliquots and some of them were stored at 4 °C for 12 h, others at 4 °C for 144 h and then used as template in a PCR reaction. To show the robustness of the method according to the invention we wanted to analyze whether this protecting effect of sulfonation would be stable over a period of time. The PCR reaction was performed under the same conditions.

In addition aliquots were stored at an increased temperature of 40 °C for 22 h and then used as template in a PCR reaction.

By stopping the chemical reaction after sulfonation the unmethylated cytosines were converted into C6 sulfonated uracils (5,6-Dihydro-6-sulfonyl-uracil) and methylated cytosines remained unchanged. However after a complete desulfonation, as described in the art, all unmethylated cytosines would have converted into uracil instead and methylated cytosines remain unchanged.

As a control for the UNG activity 10^5 copies of a uracil containing PCR product were added to the reaction premix, generated by use of the same primers but under presence of dUTP instead of dTTP. Reamplification took place when UNG was absent, with the expected efficiency of a crossing point of 22,6. However, when UNG was present in the PCR - mix the crossing points reached only a value of 35,1 (table 3). This difference demonstrates nicely the efficient degradation of PCR products by UNG resulting from the cleavage of uracils out of the DNA.

In this example it could be shown that bisulfite treated DNA which is not desulfonated according to the invention is stable at 4 °C for a longer period of at least 144 hrs. In addition it could be shown that even storage at 40 °C for a period of 22 hrs does not have a major effect on the UNG protecting effect of sulfonation at C6-uracils.

Table 3:

DNA after 12 h at 4 °C			
	Template DNA in ng	Crossing point of reaction without UNG	Crossing point of reaction <u>with</u> 0,2 Unit UNG added
PCR amplicons containing uracil	10 ⁵ copies	22.6	35.8
desulfonated DNA	0,1	33,8	no signal
sulfonated DNA	0,1	34,1	34,8
DNA after 144 h at 4 °C			
	Template DNA in ng	Crossing point of reaction without UNG	Crossing point of reaction <u>with</u> 0,2 Unit UNG added
PCR amplicons containing uracil	10.00E5 copies	22.6	35.1
desulfonated DNA	10	28,3	no signal
desulfonated DNA	1	30,8	no signal
desulfonated DNA	0,1	33,5	no signal
sulfonated DNA	10	28,7	29,5
sulfonated DNA	1	31,7	31,8
sulfonated DNA	0,1	33,7	34,2
DNA after 22 h at 40 °C			
	Template DNA in ng	Crossing point of reaction without UNG	Crossing point of reaction <u>with</u> 0,2 Unit UNG added
PCR amplicons containing uracil	10.00E5 copies	22.6	35.7
desulfonated DNA	10	29,5	no signal
desulfonated DNA	1	32,5	no signal
desulfonated DNA	0,1	34,9	no signal
sulfonated DNA	10	29,5	30.5
sulfonated DNA	1	32,5	33.3
sulfonated DNA	0,1	34,6	36.1

Example 3:

Comparison of the method according to the invention with the
5 standard workflow by means of the determination of the methylation rate of the TPEF gene (also known as TMEFF2) in colon cancer tissue.

10 1 µg of genomic DNA (200 µl) was extracted from tumours and normal adjacent tissue of 12 patients with colon cancer, respectively. The 24 samples obtained in this way were each divided into 2 x 100 µl DNA. 100 µl of each sample was treated according to standard procedures (bisulfite treatment protocol A, sample set A) or to the method according to the invention
15 (bisulfite protocol B, sample set B). In between the DNA was stored at -20 °C.

Standard workflow:

20 Sample set A:

Measurement of the DNA was performed according to the C3 quantification assay version A and according to the HeavyMethyl assay for the TPEF gene version A. A standard A was generated for calibration.

25

Generation of standard A:

5 tubes each with 2 µg universal methylated DNA were treated with bisulfite according to the bisulfite treatment protocol A and pooled afterwards. The concentration of the DNA in solution was determined by means of UV at 260 nm after the bisulfite reaction.
30

Bisulfite treatment protocol A (standard procedures):

100 µl of the samples (sample set A) containing 0,5 µg DNA diluted in 100 µl water were mixed with 354 µl of bisulfite solution (5.89 mol/l) and 146 µl of dioxane containing a scavenger (6-hydroxy-2,5,7,8-tetramethylchromane 2-carboxylic acid, 98.6 mg in 2.5 ml of dioxane). The reaction mixture was denatured for 3 min at 99 °C and subsequently incubated with the
35

following temperature program for a total of 5 h: 30 min
50 °C; one thermospike 99.9 °C for 3 min; 1.5 h 50 °C; one
thermospike 99.9 °C for 3 min; 3 h 50 °C. The DNA of the reac-
tion mixtures was subsequently purified by ultrafiltration by
5 means of a Millipore Microcon column. The purification was
conducted essentially according to the manufacturer's instruc-
tions. For this purpose, the reaction mixture was mixed with
200 µl of water, loaded onto the ultrafiltration membrane,
centrifuged for 15 min and subsequently washed with water. The
10 DNA remains on the membrane in this treatment. For complete
desulfonation 100 µl of a 0.2 mol/l NaOH solution was added
and incubated for 10 min. A centrifugation for 10 min was then
conducted, followed by a final washing step with water. After
this, the DNA was eluted. For this purpose, the membrane was
15 mixed for 10 minutes with 75 µl of prewarmed 1 x TE buffer
(50 °C) adjusted to pH 8.5. Then the membrane was turned over
and centrifuged according to the manufacturer's instructions
to recover the DNA from the membrane.

C3 quantification assay (EP05075404):

The C3 quantification assay is a quantification assay specific for the total amount of bisulfite converted DNA as described in detail in EP 05075404. The assay amplifies a fragment of DNA that comprises multiple cytosine (but not CpG) positions in the genomic form, which are initially converted to uracil and during amplification replaced by thymine in the bisulfite converted variant. Accordingly the assay does not quantify for unconverted or partially converted bisulfite treated DNA (i.e. wherein the target sequence comprises one or more cytosine positions which have not been converted to thymine). The quantity of DNA in the sample is deduced by comparison of the measured CP (crossing point, which represents the threshold cycle) to a standard curve relating such CP values to DNA amounts. The standard curve is based on measurements of known quantities of bisulfite converted DNA with the according assay.

C3 quantification assay version A:

A 20 µl reaction mixture contained:

- 2 µl of template DNA
- 2 µl of FastStart LightCycler Mix for hybridisation probes (Roche Diagnostics)
- 3.5 mmol/l MgCl₂ (Roche Diagnostics)
- 0.60 µmol/l forward primer (Seq ID-6, TIB-MolBiol)
- 0.60 µmol/l reverse primer (Seq ID-7, TIB-MolBiol)
- 0.2 µmol/l probe1 (Seq ID-8, TIB-MolBiol)

The assay was performed according to the following temperature-time-profile:

- activation 10 min at 95 °C
- 50 cycles: 10 sec at 95 °C
30 sec at 56 °C
10 sec at 72 °C

The used primers (Seq ID-6 and Seq ID-7) amplify a fragment of 123 bp of the GSTP1 gene (Seq ID-9. nucleotide 2273 to nucleotide 2402 of GenBank Accession Number X08058). The detection was carried out during the annealing phase at 56 °C in channel

F1 at 530 nm. The crossing points (CP) were calculated according to the "second derivative maximum" method by means of the LightCycler software.

5 Detection of the methylation rate according to the HeavyMethyl assay for the TPEF gene version A:

A 20 µl reaction mixture contained:

- 2 µl of template DNA
- 2 µl of FastStart LightCycler Mix for hybridization
- 10 probes (Roche Diagnostics)
- 3.5 mmol/l MgCl₂ (Roche Diagnostics)
- 0.30 µmol/l forward primer (Seq ID-10, TIB-MolBiol)
- 0.30 µmol/l reverse primer (Seq ID-11, TIB-MolBiol)
- 4.0 µmol/l blocker (Seq ID-12, TIB-MolBiol)
- 15 • 0.15 µmol/l hybridization probe (Seq ID-13, TIB-MolBiol)
- 0.15 µmol/l hybridization probe (Seq ID-14, TIB-MolBiol)

The assay was performed according to the following temperature-time-profile:

- 20 - activation 10 min at 95 °C
- 50 cycles: 10 sec at 95 °C
- 30 sec at 56 °C
- 10 sec at 72 °C

25 The used primers (Seq ID-10 and Seq ID-11) amplify a fragment of 113 bp of the TPEF gene (Seq ID-15. nucleotide 1102 to nucleotide 1214 of GenBank Accession Number AF242221). The detection was carried out during the annealing phase at 56 °C in channel F2/F1 at 640/530 nm. The crossing points (CP) were
30 calculated according to the "second derivative maximum" method by means of the LightCycler software.

Calculation of DNA amounts from CP:

Both the C3 quantification assay and HeavyMethyl assay for the
35 TPEF gene are Real Time PCR assays using an external standard for calculating the DNA amount of the measured samples. The absolute value (ng) for an unknown concentration is obtained by a comparison of the amplification of DNA in an unknown sample against a standard curve prepared with known concentra-

tions of the same target. The standard samples are amplified in separate capillaries but within the same LightCycler run. The standard curve is the linear regression line through the data points on a plot of crossing points (threshold cycle) versus logarithm of standard sample concentration. The absolute amount of DNA (ng) of the unknown sample matches the data point of the standard curve at which the CP of the unknown sample fits the standard curve.

Table 4: Sequence of Oligonucleotides

SeqID	Name	Sequence
Seq ID-6	C3F	GGAGTGGAGGAAAtTGAGAt
Seq ID-7	C3R	CCACACAaCAaaTaCTCAaAaC
Seq ID-8	C3-TAQ	FAMTGGGTGTTTGTAATTTTGTGTTAGGTT-BHQ1
Seq ID-9	C3-amplicon	GGAGTGGAGGAAAtTGAGAtttAtTGAGGTTACGTAGTTTGttt AAGGTtAAGttTGGGTGttTGtAATttTTGtttTGTGttAGGtTG- ttTttt AGGTGttAGGTGAGtTtTGAGtAttTGtTGTGTGG
Seq ID-10	TPEF -61S	aAAAaAaAAAAaCTCCTCTaCATAC
Seq ID-11	TPEF -62S	GGTtAtTGttTGGGttAAtAAATG
Seq ID-12	TPEF -6B2	aCATACaCCaCaaaTaaaTtaCCaaaAaCATCaaCCaa-PH
Seq ID-13	TPEF -6SF1	tTttttTTttCGGACGtCGtT-Fluo
Seq ID-14	TPEF -6SR1	red640-tCGGtCGATGtTttCGGtAA-PH
Seq ID-15	TPEF -amplicon	GGTtAtTGttTGGGttAAtAAATGGAGttCGtTtTttttTTttCG- GACG TCGtTGttCGGtCGATGtTttCGGtAAtttAttCGCGGCGTATG- tAG AGGAGttTTTtTtTTTt

Fluo=fluoresceine label, red640=LightCycler fluorescence label for channel F2, PH=3'OH-Phosphorylation, FAM=5'-FAM label, BHQ1=BlackHoleQuencher1. Small written t's point to converted

cytosines by bisulfite treatment, respectively small a's point to the complementary adenosine bases in the reverse complement synthesized strand.

5

Method according to the invention:

Sample set B:

Measurement of the DNA was performed according to the C3 quantification assay version B and according to the HeavyMethyl assay for the TPEF gene version B in addition of 10,000 copies of a PCR product of methylated DNA. A standard B (C6 sulfonated uracil containing DNA) was generated for calibration.

15 Generation of standard B (C6 sulfonated uracil containing DNA):

5 tubes each with 2,0 µg universal methylated DNA were treated with bisulfite according to the bisulfite treatment protocol B. The concentration of the DNA in solution was determined by means of UV at 260 nm after the bisulfite reaction.

Generation of PCR products.

10 ng methylated bisulfite converted DNA generated according to standard procedures (bisulfite protocol A) were amplified by means of the HeavyMethyl assay for the TPEF gene version A. The PCR products were purified with the QIAquick PCR Purification Kit and subsequently analysed on a 2 % agarose gel. After this, a serial dilution was carried out with water to a final dilution of 1:10¹⁰. 2 µl of this dilution was reamplified and quantificated according to the HeavyMethyl assay for the TPEF gene version A. The copy number was determined: 2 µl of the said dilution contain 10,000 copies of PCR product.

35 Bisulfite treatment protocol B (protocol for carry over protection):

100 µl of the samples (sample set B) containing 0,5 µg DNA diluted in 100 µl water were mixed with 354 µl of bisulfite solution (5.89 mol/l) and 146 µl of dioxane containing a scavenger (6-hydroxy-2,5,7,8-tetramethylchromane 2-carboxylic acid,

98.6 mg in 2.5 ml of dioxane). The reaction mixture was denatured for 3 min at 99 °C and subsequently incubated with the following temperature program for a total of 5 h: 30 min 50 °C; one thermospike 99.9 °C for 3 min; 1.5 h 50 °C; one thermospike 99.9 °C for 3 min; 3 h 50 °C. The DNA of the reaction mixtures was subsequently purified by ultrafiltration by means of a Millipore Microcon column. The purification was conducted essentially according to the manufacturer's instructions. For this purpose, the reaction mixture was mixed with 200 µl of water, loaded onto the ultrafiltration membrane, centrifuged for 15 min and subsequently washed with water. The DNA remains on the membrane in this treatment. In contrast to the bisulfite treatment protocol A the DNA was not incubated with NaOH, but additionally washed with water. After this, the DNA was eluted. For this purpose, the membrane was mixed for 10 minutes with 75 µl of prewarmed water (50 °C). Then the membrane was turned over and centrifuged according to the manufacturer's instructions to recover the DNA from the membrane.

C3 quantification assay version B:

A 20 µl reaction mixture contained:

- 2 µl of template DNA
- 2 µl PCR product (10,000 copies)
- 2 µl of FastStart LightCycler Mix for hybridisation probes (Roche Diagnostics)
- 3.5 mmol/l MgCl₂ (Roche Diagnostics)
- 0.60 µmol/l forward primer (Seq ID-6, TIB-MolBiol)
- 0.60 µmol/l reverse primer (Seq ID-7, TIB-MolBiol)
- 0.2 µmol/l probe1 (Seq ID-8, TIB-MolBiol)
- 0.2 units uracil-DNA-glycosylase (Roche Diagnostics)

The assay was performed according to the following temperature-time-profile:

- preincubation 10 min at 37 °C
- desulfonation / activation 30 min at 95 °C
- 50 cycles: 10 sec at 95 °C
 30 sec at 56 °C
 10 sec at 72 °C

The used primers (Seq ID-6 and Seq ID-7) amplify a fragment of 123 bp of the GSTP1 gene (Seq ID-9, nucleotide 2273 to nucleotide 2402 of GenBank Accession Number X08058). The detection was carried out during the annealing phase at 56 °C in channel F1 at 530 nm. The crossing points (CP) were calculated according to the "second derivative maximum" method by means of the LightCycler software.

Detection of the methylation rate according to the HeavyMethyl assay for the TPEF gene version B:

A 20 µl reaction mixture contained:

- 2 µl of template DNA
- 2 µl PCR product (10,000 copies)
- 2 µl of FastStart LightCycler Mix for hybridisation probes (Roche Diagnostics)
- 3.5 mmol/l MgCl₂ (Roche Diagnostics)
- 0.30 µmol/l forward primer (Seq ID-10, TIB-MolBiol)
- 0.30 µmol/l reverse primer (Seq ID-11, TIB-MolBiol)
- 4.0 µmol/l blocker (Seq ID-12, TIB-MolBiol)
- 0.15 µmol/l hybridisation probe (Seq ID-13, TIB-MolBiol)
- 0.15 µmol/l hybridisation probe (Seq ID-14, TIB-MolBiol)
- 0.2 units uracil-DNA-glycosylase (Roche Diagnostics)

The assay was performed according to the following temperature-time-profil:

- preincubation 10 min at 37 °C
- desulfonation / activation 30 min at 95 °C
- 50 cycles: 10 sec at 95 °C
30 sec at 56 °C
10 sec at 72 °C

The used primers (Seq ID-10 and Seq ID-11) amplify a fragment of 113 bp of the TPEF gene (Seq ID-15, nucleotide 1102 to nucleotide 1214 of GenBank Accession Number AF242221). The detection was carried out during the annealing phase at 56 °C in channel F2/F1 at 640/530 nm. The crossing points (CP) were

calculated according to the "second derivative maximum" method by means of the LightCycler software.

Calculation of DNA amounts from CP:

5 Both the C3 quantification assay and HeavyMethyl assay for the TPEF gene are Real Time PCR assays using an external standard for calculating the DNA amount of the measured samples. The absolute value (ng) for an unknown concentration is obtained by a comparison of the amplification of DNA in an unknown sam-
10 ple against a standard curve prepared with known concentrations of the same target. The standard samples are amplified in separate capillaries but within the same LightCycler run. The standard curve is the linear regression line through the data points on a plot of crossing points (threshold cycle)
15 versus logarithm of standard sample concentration. The absolute amount of DNA (ng) of the unknown sample matches the data point of the standard curve at which the CP of the unknown sample fits the standard curve.

20 Calculation of the methylation rate from DNA amounts: The results of the study are presented as methylation rates of the promotor region of the TPEF gene. According to the PMR value methode (Eads CA et al. Cancer Res 2001 Apr 15;61(8):3410-8. PMID: 11309301) the methylation rate is equal to the percent-
25 age of methylated copies measured in a sample as proportion of the total DNA measured in the same sample. In table 5 and 6 all CPs received from the C3 and the TPEF assay and the resulting DNA amounts are listed. In the right column the methylation rate (PMR) is shown, which was calculated from the DNA
30 amounts listed in columns before.

Results:

Table 5: Results from the standard workflow. Colon cancer and normal adjacent tissue samples were bisulfite treated with bisulfite treatment protocol A followed by quantification with the C3 quantification assay version A using a calibration curve made by means of standard A. The table shows the crossing points and the calculated DNA amount of 2 replicates. The HeavyMethyl assay for the TPEF gene version A detects only methylated DNA from the promoter region of TPEF gene. The table shows the measured CP values of 2 replicates and the calculated DNA amount. Finally the methylation percentages (PMR) were calculated by the ratio of methylated DNA and total DNA.

Sample	Type	C3 Quantification Assay			HeavyMethyl Assay			PMR
		Version A			TPEF gene			
		CP	CP	ng/PCR	CP	CP	ng/PCR	
		1 st run	2 nd run	mean	1 st run	2 nd run	mean	%
standard A	20 ng	25,86	25,64		26,28	26,55		
standard A	5 ng	28,29	27,72		28,51	28,24		
standard A	5 ng	28,44	27,74		28,51	28,15		
standard A	1,25 ng	30,27	30,2		29,91	30,46		
standard A	1,25 ng	30,23	30,41		29,92	30,21		
standard A	0,31 ng	32,48	32,01		31,46	31,51		
standard A	0,31 ng	31,89	32,18		31,13	31,54		
1	normal	31,49	32,6	0,4	27,27	28,68	7,4	5 %
2	tumor	-	-	0,0	28,22	29,49	4,1	0 %
3	normal	32,74	33,56	0,1	27,84	28,97	5,4	2 %
4	tumor	29,51	30,96	1,5	28,08	29,42	4,4	33 %
5	normal	31,97	33,03	0,3	27,72	29,09	5,6	5 %
6	tumor	27,8	29,65	4,0	27,45	29,16	6,2	65 %
7	normal	32,02	33,47	0,2	27,62	28,93	6,0	4 %
8	tumor	28,5	30,53	2,5	27,56	29,2	5,9	43 %
9	normal	32,47	33,8	0,1	28,06	29,13	4,7	3 %
10	tumor	29,87	31,17	1,2	27,91	29,07	5,1	23 %
11	normal	31,93	33,65	0,2	27,17	29,12	7,2	3 %
12	tumor	32,07	33,27	0,2	27,1	28,29	8,6	3 %
13	normal	31,45	33,41	0,3	27,52	28,61	6,7	5 %
14	tumor	29,87	30,64	1,3	27,83	28,44	6,2	22 %
15	normal	35,97	32,94	0,1	27,8	28,2	6,7	1 %
16	tumor	28,92	29,07	2,8	28,54	28,73	4,4	64 %
17	normal	32,49	34,58	0,1	27,98	29,45	4,6	3 %
18	tumor	27,05	27,93	7,5	27,16	28,08	8,8	84 %
19	normal	31,6	32,78	0,3	27,6	28,95	6,0	6 %
20	tumor	28,78	30,11	2,4	27,59	28,79	6,2	38 %
21	normal	35,23	35,88	0,0	28,07	29,32	4,5	0 %
22	tumor	35	36,93	0,0	27,91	29,08	5,1	0 %
23	normal	31,86	32,72	0,3	27,63	28,31	6,9	4 %
24	tumor	30,01	31,55	1,0	27,47	28,58	6,9	15 %
neg. contr.	-	-	-	-	-	-	-	

Table 6: Results generated by the method according to the invention (carry over prevention). Colon cancer and normal adjacent tissue samples were bisulfite treated with bisulfite treatment protocol B resulting in C6 sulfonated uracil containing DNA. Total DNA was measured with the C3 quantification assay version B using a calibration curve made by means of standard B. The table shows the crossing points and the calculated DNA amount from 2 replicates. Before the measurement of the methylated DNA with the HeavyMethyl Assay for the TPEF gene version B, the reactions were contaminated with 10,000 copies of the TPEF amplicon containing uracil instead of thymine. The table shows the measured CP of 2 replicates and the calculated DNA amount. Finally the methylation percentages (PMR) were calculated by the ratio of methylated DNA and total DNA.

Sample	Type	C3 Quantification Assay Version B			HeavyMethyl Assay for the TPEF gene Version B			PMR %
		CP	CP	ng/PCR	CP	CP	ng/PCR	
		1 st Run	2 nd Run	mean	1 st Run	2 nd Run	mean	
standard B	20 ng	27,73	27,29		27,63	27,31		
standard B	5 ng	29,18	28,58		28,82	28,62		
standard B	5 ng	29,18	28,89		28,76	28,73		
standard B	1,25 ng	31,04	30,71		30,57	30,05		
standard B	1,25 ng	31,08	30,51		30,33	30,01		
standard B	0,31 ng	32,9	32,84		31,87	31,46		
standard B	0,31 ng	32,78	32,65		32,27	31,69		
1	normal	33,84	33,49	0,2	28,13	28,16	9,8	2 %
2	tumor	37,54	-	0,0	28,53	29,14	5,3	0 %
3	normal	34,43	34,72	0,2	29,14	29,68	3,0	5 %
4	tumor	30,46	30,1	1,9	28,23	28,2	9,1	21 %
5	normal	34,17	32,89	0,2	28,7	28,92	5,2	4 %
6	tumor	29,23	28,69	5,6	27,87	27,83	13,1	43 %
7	normal	32,97	32,87	0,3	27,92	28,01	11,7	2 %
8	tumor	30,82	30,63	1,3	28,44	28,78	6,4	21 %
9	normal	33,87	33,55	0,2	28,43	28,59	6,9	3 %
10	tumor	30,97	31,19	1,0	28,58	29,12	5,2	20 %
11	normal	33,14	33,53	0,2	28,11	28,2	9,7	3 %
12	tumor	33,13	33,66	0,2	27,69	28	13,5	2 %
13	normal	33,04	33,57	0,3	28,44	28,86	6,2	4 %
14	tumor	31,08	31,54	0,9	28,03	28,79	8,4	10 %
15	normal	33,09	33,6	0,2	28,03	28,54	9,0	3 %
16	tumor	29,74	29,78	3,0	28,59	28,95	5,5	54 %
17	normal	33,95	33,52	0,2	28,43	28,68	6,7	3 %
18	tumor	28,66	28,69	7,3	28,09	28,09	10,3	70 %
19	normal	33,01	34,08	0,2	28,11	28,2	9,7	3 %
20	tumor	30,03	30,71	1,9	28,19	28,17	9,5	20 %
21	normal	36,24	36,98	0,2	28,92	29,25	4,0	4 %
22	tumor	36,04	35,12	0,1	28,58	28,9	5,6	2 %
23	normal	33,79	33,77	0,2	28,1	28,43	9,0	2 %
24	tumor	31,89	32,2	0,5	28,69	29,09	4,9	10 %
neg. contr.	-	-	-	-	-	-	-	

The results obtained by the standard workflow and the method according are compared in a correlation plot (Figure 5). Every symbol represents a single sample: quadrates tumor tissues, triangles normal adjacent tissues. The percentage of methylation determined according to the standard workflow (x-axis) or to the method according to the invention (y-axis) is indicated for each sample.

The method according to the invention has led only in 2 out of 24 samples to a different methylation percentage as the standard workflow. Although the samples treated according to the method of the invention were contaminated with uracil containing TPEF amplicons only DNA of the samples served as a template for amplification of the TPEF amplicon in nearly all cases. In case of the said two samples, the differing results occurred presumable because of the low methylation percentage of the DNA (smaller than 0.2 %).

Claims

1. A method for providing a decontaminated nucleic acid
suitable for DNA methylation analysis, characterized
by
- a) incubating a nucleic acid with a bisulfite reagent
containing solution, whereby the unmethylated cyto-
sines within said nucleic acid are sulfonated and/or
deaminated but wherein no desulfonation reaction is
carried out, and
- b) adding to this mixture a enzyme, which degrades
non-sulfonated uracil containing nucleic acids and
does not degrade sulfonated uracil containing nucleic
acids, and incubating the mixture, whereby nucleic
acids containing non-sulfonated uracils are degraded.
2. A method of claim 1 for providing a decontaminated
template nucleic acid for polymerase based amplifica-
tion reactions, characterized by
- a) mixing the sulfonated and/or deaminated template
nucleic acid with the components required for a poly-
merase mediated amplification reaction or an amplifi-
cation based detection assay, and
- b) after degradation of non-sulfonated uracil con-
taining nucleic acids, terminating the enzymatic ac-
tivity, which caused the degradation of non-
sulfonated uracil containing nucleic acid, and
- c) desulfonating the template nucleic acid.

3. The method of claim 2 wherein step b) and c) take place simultaneously by briefly incubating the mixture at an increased temperature, whereby the enzymatic activity, which specifically degrades non-sulfonated uracil containing nucleic acids, is terminated and whereby desulfonation of the template nucleic acid takes place.
4. The method of claim 2 wherein in a subsequent step d) the template nucleic acid is amplified.
5. The method of claim 2 wherein the enzyme, which degrades non-sulfonated uracil containing nucleic acids and does not degrade sulfonated uracil containing nucleic acids, is a DNA glycosylase or endonuclease, in particular uracil-DNA-glycosylase.
6. The method of claim 2 wherein upon termination of the enzymatic activity, which causes a degradation of sulfonated uracil containing nucleic acids, and desulfonation of the template nucleic acid a polymerase based amplification reaction is started and/or an amplification based assay is performed.
7. The method of claim 6 wherein the polymerase based amplification reaction is started by a brief incubation at increased temperature (heat activation).
8. The method of claim 6 wherein the polymerase is a heat stable polymerase.
9. The method of claim 6, wherein the polymerase mediated amplification or amplification based assay is performed in the presence of dUTPs instead of dTTPs.

10. A method for a bisulfite treatment and subsequent amplification of a nucleic acid characterized in that

5 a) the nucleic acid is incubated with a bisulfite reagent containing solution, whereby the unmethylated cytosines within said nucleic acid are sulfonated and/or deaminated but wherein no desulfonation reaction is carried out;

10 b) the sulfonated and/or deaminated template nucleic acid is mixed with components required for a polymerase mediated amplification reaction or an amplification based detection assay;

15 c) the sulfonated and/or deaminated template nucleic acid is desulfonated, by a brief incubation of the mixture at an increased temperature.

20 11. The method of claim 10, characterized in that the desulfonated template nucleic acid is amplified, preferably with a polymerase mediated amplification reaction or an amplification based detection assay, wherein a heat stable polymerase is preferably used.

25 12. The method of claim 11, characterized in that the polymerase based amplification reaction is started by a brief incubation at increased temperature (heat activation), whereby simultaneously to the amplification start the desulfonation of the sulfonated and/or
30 deaminated template nucleic acid can take place.

35 13. A kit when used according to the method of any one of claims 1 to 12 comprising a reagent containing bisulfite and an enzymatic activity, which specifically degrades non-sulfonated uracil containing DNA and does not degrade sulfonated uracil containing nucle-

ic, in particular this enzymatic activity is uracil-DNA-glycosylase.



5 14. The kit of claim 13, comprising one or more of the following additional components: denaturing reagent and solution; scavenger; primer, probe, reaction buffer, nucleotides, $MgCl_2$ solution, polymerase and/or dye for the production of amplicates; and/or reagent, solution and instructions, which are useful for
10 conducting an assay according to any one of the preceding claims.

15 15. Use of the kit of claim 13 or 14 for diagnosis and/or prognosis of adverse events for patients or individuals, whereby these adverse events belong to at least one of the following categories: undesired drug interactions; cancer diseases; CNS malfunctions; damage
20 or disease; symptoms of aggression or behavioral disturbances; clinical; psychological and social consequences of brain damages; psychotic disturbances and personality disorders; dementia and/or associated syndromes; cardiovascular disease of the gastrointestinal tract; malfunction, damage or disease of the
25 respiratory system; lesion, inflammation, infection, immunity and/or convalescence; malfunction, damage or disease of the body as an abnormality in the development process; malfunction, damage or disease of the skin, of the muscles, of the connective tissue or of
30 the bones; endocrine and metabolic malfunction, damage or disease; headaches or sexual malfunction.

35 16. Use of the kit of claim 13 or 14 for distinguishing cell types or tissue or for investigating cell differentiation.

17. Use of the method of any one of claims 1 to 12 or of
a kit of claim 13 or 14 for methylation analysis
wherein the DNA methylation status is characterized
in that positions are methylated or non-methylated
compared to normal conditions if a single defined
disease exists.
18. Use of the enzyme uracil-DNA-glycosylase for the gen-
eration of contamination free nucleic acids for me-
thylation analysis.
19. Use of the method of any one of claims 1 to 12 for
methylation analysis, whereby the methylation analy-
sis is performed on samples from patients or individ-
uals to diagnose and/or prognose adverse events for
them, whereby these adverse events belong to at least
one of the following categories: undesired drug inte-
ractions; cancer diseases; CNS malfunctions; damage
or disease; symptoms of aggression or behavioral dis-
turbances; clinical; psychological and social conse-
quences of brain damages; psychotic disturbances and
personality disorders; dementia and/or associated
syndromes; cardiovascular disease of the gastrointes-
tinal tract; malfunction, damage or disease of the
respiratory system; lesion, inflammation , infection,
immunity and/or convalescence; malfunction, damage or
disease of the body as an abnormality in the develop-
ment process; malfunction, damage or disease of the
skin, of the muscles, of the connective tissue or of
the bones; endocrine and metabolic malfunction, dam-
age or disease; headaches or sexual malfunction.
20. Use of the method of any one of claims 1 to 12 for
methylation analysis, whereby the methylation analy-
sis is performed to distinguish cell types or tissues
or to investigate cell differentiation.

Figure 1

**Sulfonation****Hydrolytic
Deamination**

Incubation with Uracil-DNA-Glycosylase


Carry Over Prevention**Desulfonation**

Figure 2




Figure 3




Figure 4




Figure 5

