

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

30 November 2017 (30.11.2017)



WIPO | PCT



(10) International Publication Number

WO 2017/205496 A1

(51) International Patent Classification:

C07F 9/6558 (2006.01) C07F 9/6561 (2006.01)

(21) International Application Number:

PCT/US2017/034235

(22) International Filing Date:

24 May 2017 (24.05.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/341,049	24 May 2016 (24.05.2016)	US
62/357,153	30 June 2016 (30.06.2016)	US
62/357,166	30 June 2016 (30.06.2016)	US
62/357,072	30 June 2016 (30.06.2016)	US
62/356,923	30 June 2016 (30.06.2016)	US
62/508,256	18 May 2017 (18.05.2017)	US

(71) Applicant: SAREPTA THERAPEUTICS, INC.

[US/US]; 215 First Street, Cambridge, MA 02142 (US).

(72) Inventors: CAI, Bao; c/o Sarepta Therapeutics, Inc., 215

First Street, Cambridge, MA 02142 (US). MARTINI, Mitchell; c/o Sarepta Therapeutics, Inc., 215 First Street, Cambridge, MA 02142 (US). SHIMABUKU, Ross; c/o Sarepta Therapeutics, Inc., 215 First Street, Cambridge, MA 02142 (US). THOMAS, Katie; c/o Sarepta Therapeutics, Inc., 215 First Street, Cambridge, MA 02142 (US). FRANK, Diane Elizabeth; c/o Sarepta Therapeutics, Inc., 215 First Street, Cambridge, MA 02142 (US). BESTWICK, Richard K..

(74) Agent: TRINQUE, Brian C. et al.; Lathrop & Gage, LLP,

28 State Street, Boston, MA 02109 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: PROCESSES FOR PREPARING OLIGOMERS

(57) Abstract: Provided herein are processes for preparing an oligomer (e.g., a morpholino oligomer). The synthetic processes described herein may be advantageous to scaling up oligomersynthesis while maintaining overall yield and purity of a synthesized oligomer.



WO 2017/205496 A1

PROCESSES FOR PREPARING OLIGOMERS

RELATED APPLICATIONS

This application claims benefit of United States provisional patent application no. 62/508,256, filed May 18, 2017; United States provisional patent application no. 62/341,049 filed May 24, 2016; United States provisional patent application no. 62/356,923, filed June 30, 2016; United States provisional patent application no. 62/357,072, filed June 30, 2016; United States provisional patent application no. 62/357,153, filed June 30, 2016; and United States provisional patent application no. 62/357,166, filed June 30, 2016; the entire contents of each of which are incorporated herein by reference.

BACKGROUND

Antisense technology provides a means for modulating the expression of one or more specific gene products, including alternative splice products, and is uniquely useful in a number of therapeutic, diagnostic, and research applications. The principle behind antisense technology is that an antisense compound, e.g., an oligonucleotide, which hybridizes to a target nucleic acid, modulates gene expression activities such as transcription, splicing or translation through any one of a number of antisense mechanisms. The sequence specificity of antisense compounds makes them attractive as tools for target validation and gene functionalization, as well as therapeutics to selectively modulate the expression of genes involved in disease.

Duchenne muscular dystrophy (DMD) is caused by a defect in the expression of the protein dystrophin. The gene encoding the protein contains 79 exons spread out over more than 2 million nucleotides of DNA. Any exonic mutation that changes the reading frame of the exon, or introduces a stop codon, or is characterized by removal of an entire out of frame exon or exons, or duplications of one or more exons, has the potential to disrupt production of functional dystrophin, resulting in DMD.

Recent clinical trials testing the safety and efficacy of splice switching oligonucleotides (SSOs) for the treatment of DMD are based on SSO technology to induce alternative splicing of pre-mRNAs by steric blockade of the spliceosome (Cirak et al., 2011; Goemans et al., 2011; Kinali et al., 2009; van Deutekom et al., 2007). However, despite these successes, the pharmacological options available for treating DMD are limited.

Eteplirsen is a phosphorodiamidate morpholino oligomer (PMO) designed to skip exon 51 of the human dystrophin gene in patients with DMD who are amendable to exon


51 skipping to restore the read frame and produce a functional shorter form of the dystrophin protein. Sarepta Therapeutics, Inc., submitted a New Drug Application (NDA) to the United States Food and Drug Administration (FDA) seeking approval for the treatment of DMD in patients amenable to exon 51 skipping. Sarepta's NDA is currently under review by the FDA.

Although significant progress has been made in the field of antisense technology, there remains a need in the art for methods of preparing phosphorodiamidate morpholino oligomers with improved antisense or antigene performance.

SUMMARY

Provided herein are processes for preparing phosphorodiamidate morpholino oligomers (PMOs). The synthetic processes described herein allow for a scaled-up PMO synthesis while maintaining overall yield and purity of a synthesized PMO.


In one aspect, the disclosure provides a process for preparing an oligomeric compound of Formula (B):



(B),

or a pharmaceutically acceptable salt thereof, wherein T, X, R², and n are as defined herein.


In some embodiments, the disclosure provides a process for preparing an oligomeric compound of Formula (A):



(A),

or a pharmaceutically acceptable salt thereof, wherein R^2 and n are as defined herein.


In certain embodiments, provided herein is a process for preparing an oligomeric compound of Formula (E):



or a pharmaceutically acceptable salt thereof.

In yet another embodiment, the oligomeric compound of the disclosure including, for example, some embodiments of an oligomeric compound of Formula (E), is an oligomeric compound of Formula (XII):

10




(XII),

or a pharmaceutically acceptable salt thereof.

For clarity, the structural formulas including, for example, oligomeric compound of Formula (E) and Eteplirsen depicted by Formula (XII), are a continuous structural formula from 5' to 3', and, for the convenience of depicting the entire formula in a compact form in the above structural formulas, Applicants have included various illustration breaks labeled "BREAK A," "BREAK B," "BREAK C," and "BREAK D." As would be understood by the skilled artisan, for example, each indication of "BREAK A" shows a continuation of the illustration of the structural formula at these points. The skilled artisan understands that the same is true for each instance of "BREAK B," "BREAK C," and "BREAK D" in the structural formulas above including Eteplirsen. None of the illustration breaks, however, are intended to indicate, nor would the skilled artisan understand them to mean, an actual discontinuation of the structural formulas above including Eteplirsen.


In another aspect of the disclosure is provided a process for preparing an oligomeric compound of Formula (D):



(D),

or a pharmaceutically acceptable salt thereof, wherein R^1 , R^2 , R^4 - R^7 , X , and n are as defined herein.


5 In some embodiments, the oligomeric compound of Formula (D) is of Formula (G):



(G),

or a pharmaceutically acceptable salt thereof, wherein R^2 , X , and n are as defined herein.


10 In another aspect of the disclosure is provided a process for preparing an oligomeric compound of Formula (F):




(F);

wherein:

each A is, independently for each occurrence, selected from the group consisting of:



5 each R² is, independently for each occurrence, selected from the group consisting of:



n is an integer from 10 to 40; and

R is H or acetyl.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows a representative analytical high performance liquid chromatography (HPLC) chromatogram of a synthesized and deprotected Eteplirsen (AVI-4658) crude drug substance (see Example 4).

Figure 2 shows a representative analytical HPLC chromatogram of a purified Eteplirsen drug substance solution (see Example 5).

Figure 3 shows a representative analytical HPLC chromatogram of a desalted and lyophilized Eteplirsen drug substance (see Example 5).

DETAILED DESCRIPTION

Provided herein are processes for preparing a morpholino oligomer. The a morpholino oligomer described herein displays stronger affinity for DNA and RNA without compromising sequence selectivity, relative to native or unmodified oligonucleotides. In some embodiments, the morpholino oligomer of the disclosure minimizes or prevents cleavage by RNase H. In some embodiments, the morpholino oligomer of the disclosure does not activate RNase H.

The processes described herein are advantageous in an industrial-scale process and can be applied to preparing quantities of a morpholino oligomer in high yield and scale (e.g., about 1 kg, about 1-10 kg, about 2-10 kg, about 5-20 kg, about 10-20 kg, or about 10-50 kg).

Definitions

Listed below are definitions of various terms used to describe this disclosure. These definitions apply to the terms as they are used throughout this specification and claims, unless otherwise limited in specific instances, either individually or as part of a larger group.

“Base-protected” or “base protection” refers to protection of the base-pairing groups, e.g., purine or pyrimidine bases, on the morpholino subunits with protecting groups suitable to prevent reaction or interference of the base-pairing groups during stepwise oligomer synthesis. An example of a base-protected morpholino subunit is the activated C subunit Compound (C) having a CBZ protecting group on the cytosine amino group depicted below.

An “activated phosphoramidate group” is typically a chlorophosphoramidate group, having substitution at nitrogen which is desired in the eventual phosphorodiamidate linkage in the oligomer. An example is (dimethylamino)chlorophosphoramidate, i.e., $\text{-O-P(=O)(NMe}_2\text{)Cl}$.

The term “support-bound” refers to a chemical entity that is covalently linked to a support-medium.

The term “support-medium” refers to any material including, for example, any particle, bead, or surface, upon which an oligomer can be attached or synthesized upon, or can be modified for attachment or synthesis of an oligomer. Representative substrates include, but are not limited to, inorganic supports and organic supports such as glass and modified or functionalized glass, plastics (including acrylics, polystyrene and copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, TEFLON, etc.), polysaccharides, nylon or nitrocellulose, ceramics, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, plastics, optical fiber bundles, and a variety of other polymers. Particularly useful support-medium and solid surfaces for some embodiments are located within a flow cell apparatus. In some embodiments of the processes described herein, the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.

In some embodiments, representative support-medium comprise at least one reactive site for attachment or synthesis of an oligomer. For example, in some embodiments, a support-medium of the disclosure comprises one or more terminal amino or hydroxyl groups capable of forming a chemical bond with an incoming subunit or other activated group for attaching or synthesizing an oligomer.

Some representative support-media that are amenable to the processes described herein include, but are not limited to, the following: controlled pore glass (CPG); oxalyl-controlled pore glass (see, e.g., Alul et al., *Nucleic Acids Research* 1991, 19, 1527); silica-containing particles, such as porous glass beads and silica gel such as that formed by the reaction of trichloro-[3-(4-chloromethyl)phenyl]propylsilane and porous glass beads (see Parr and Grohmann, *Angew. Chem. Internatl. Ed.* 1972, 11, 314, sold under the trademark “PORASIL E” by Waters Associates, Framingham, Mass., USA); a mono ester of 1,4-dihydroxymethylbenzene and silica (see Bayer and Jung, *Tetrahedron Lett.* 1970, 51, 4503, sold under the trademark “BIOPAK” by Waters Associates); TENTAGEL (see, e.g., Wright et al., *Tetrahedron Lett.* 1993, 34, 3373); cross-linked styrene/divinylbenzene copolymer beaded matrix, or POROS, a copolymer of polystyrene/divinylbenzene (available from PerSeptive Biosystems); soluble support-medium such as polyethylene glycol PEG's (see Bonora et al., *Organic Process Research & Development* 2000, 4, 225–231); PEPS support, which is a polyethylene (PE) film with pendant long-chain polystyrene (PS) grafts (see Berg et al., *J. Am. Chem. Soc.* 1989, 111, 8024 and International Patent Application WO

1990/02749); copolymers of dimethylacrylamide cross-linked with N,N'-bisacryloylethylenediamine, including a known amount of N-tertbutoxycarbonyl-beta-alanyl-N'-acryloylhexamethylenediamine (see Atherton et al., *J. Am. Chem. Soc.* 1975, 97, 6584, Atherton et al., *Bioorg. Chem.* 1979, 8, 351, and Atherton et al., *J. Chem. Soc. Perkin I* 1981, 538); glass particles coated with a hydrophobic cross-linked styrene polymer (see Scott et al., *J. Chrom. Sci.* 1971, 9, 577); fluorinated ethylene polymer onto which has been grafted polystyrene (see Kent and Merrifield, *Israel J. Chem.* 1978, 17, 243 and van Rietschoten in *Peptides* 1974, Y. Wolman, Ed., Wiley and Sons, New York, 1975, pp. 113–116); hydroxypropylacrylate-coated polypropylene membranes (Daniels et al., *Tetrahedron Lett.* 1989, 4345); acrylic acid-grafted polyethylene-rods (Geysen et al., *Proc. Natl. Acad. Sci. USA* 1984, 81, 3998); a “tea bag” containing traditionally-used polymer beads (Houghten, *Proc. Natl. Acad. Sci. USA* 1985, 82, 5131); and combinations thereof.

The term “flow cell apparatus” refers to a chamber comprising a surface (e.g., solid surface) across which one or more fluid reagents (e.g., liquid or gas) can be flowed.

The term “deblocking agent” refers to a composition (e.g., a solution) comprising a chemical acid or combination of chemical acids for removing protecting groups. Exemplary chemical acids used in deblocking agents include halogenated acids, e.g., chloroacetic acid, dichloroacetic acid, trichloroacetic acid, fluoroacetic acid, difluoroacetic acid, and trifluoroacetic acid. In some embodiments, a deblocking agent removes one or more trityl groups from, for example, an oligomer, a support-bound oligomer, a support-bound subunit, or other protected nitrogen or oxygen moiety.

The terms “halogen” and “halo” refer to an atom selected from the group consisting of fluorine, chlorine, bromine, and iodine.

The term “capping agent” refers to a composition (e.g., a solution) comprising an acid anhydride (e.g., benzoic anhydride, acetic anhydride, phenoxyacetic anhydride, and the like) useful for blocking a reactive cite of, for example, a support-medium forming a chemical bond with an incoming subunit or other activated group.

The term “cleavage agent” refers to a composition (e.g., a liquid solution or gaseous mixture) comprising a chemical base (e.g., ammonia or 1,8-diazabicycloundec-7-ene) or a combination of chemical bases useful for cleaving, for example, a support-bound oligomer from a support-medium.


The term “deprotecting agent” refers to a composition (e.g., a liquid solution or gaseous mixture) comprising a chemical base (e.g., ammonia, 1,8-diazabicycloundec-7-ene or potassium carbonate) or a combination of chemical bases useful for removing protecting

groups. For example, a deprotecting agent, in some embodiments, can remove the base protection from, for example, a morpholino subunit, morpholino subunits of a morpholino oligomer, or support-bound versions thereof.

The term “solvent” refers to a component of a solution or mixture in which a solute is dissolved. Solvents may be inorganic or organic (e.g., acetic acid, acetone, acetonitrile, acetyl acetone, 2-aminoethanol, aniline, anisole, benzene, benzonitrile, benzyl alcohol, 1-butanol, 2-butanol, i-butanol, 2-butanone, t-butyl alcohol, carbon disulfide, carbon tetrachloride, chlorobenzene, chloroform, cyclohexane, cyclohexanol, cyclohexanone, di-n-butylphthalate, 1,1-dichloroethane, 1,2-dichloroethane, diethylamine, diethylene glycol, diglyme, dimethoxyethane (glyme), N,N-dimethylaniline, dimethylformamide, dimethylphthalate, dimethylsulfoxide, dioxane, ethanol, ether, ethyl acetate, ethyl acetoacetate, ethyl benzoate, ethylene glycol, glycerin, heptane, 1-heptanol, hexane, 1-hexanol, methanol, methyl acetate, methyl t-butyl ether, methylene chloride, 1-octanol, pentane, 1-pentanol, 2-pentanol, 3-pentanol, 2-pentanone, 3-pentanone, 1-propanol, 2-propanol, pyridine, tetrahydrofuran, toluene, water, p-xylene).

The phrases “morpholino oligomer” and “phosphorodiamidate morpholino oligomer” or “PMO” refers to an oligomer having morpholino subunits linked together by phosphorodiamidate linkages, joining the morpholino nitrogen of one subunit to the 5'-exocyclic carbon of an adjacent subunit. Each morpholino subunit comprises a nucleobase-pairing moiety effective to bind, by nucleobase-specific hydrogen bonding, to a nucleobase in a target.

The term “EG3 tail” refers to triethylene glycol moieties conjugated to the oligomer, e.g., at its 3'- or 5'-end. For example, in some embodiments, “EG3 tail” conjugated to the 3' end of an oligomer can be of the structure:



25

In certain embodiments, an oligomeric compound prepared by the methods of the disclosure can have a length from 10 to 50 nucleotides, 10 to 40 nucleotides, 10 to 30 nucleotides, 10 to 25 nucleotides, 10 to 24 nucleotides, 10 to 23 nucleotides, 10 to 22 nucleotides, 10 to 21 nucleotides, 10 to 20 nucleotides, 15 to 30 nucleotides, 15 to 25 nucleotides, 15 to 24 nucleotides, 15 to 23 nucleotides, 15 to 22 nucleotides, 15 to 21 nucleotides, 15 to 20 nucleotides, 20 to 30 nucleotides, 20 to 25 nucleotides, 20 to 24

30

nucleotides, 20 to 23 nucleotides, or 20 to 22 nucleotides in length, including all integers in between these ranges. Preferably, an oligomeric compound prepared by the methods of the disclosure is of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 20 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 21 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 22 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 23 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 24 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 25 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 26 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 27 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 28 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 29 nucleotides in length. In some embodiments, an oligomeric compound prepared by the methods of the disclosure is 30 nucleotides in length.

Accordingly, in some embodiments of oligomeric compounds prepared by the methods of the disclosure including, for example, embodiments of oligomeric compounds of Formula (A), (B), (D), (G), (F), and (S), n is an integer ranging from 10 to 50, 10 to 40, 10 to 30, 10 to 25, 10 to 24, 10 to 23, 10 to 22, 10 to 21, 10 to 20, 15 to 30, 15 to 25, 15 to 24, 15 to 23, 15 to 22, 15 to 21, 15 to 20, 20 to 30, 20 to 25, 20 to 24, 20 to 23, or 20 to 22. In certain embodiments, n is 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40. In some embodiments, n is 10. In some embodiments, n is 11. In some embodiments, n is 12. In some embodiments, n is 15. In some embodiments, n is 18. In some embodiments, n is 20. In some embodiments, n is 21. In some embodiments, n is 22. In some embodiments, n is 23. In some embodiments, n is 24. In some embodiments, n is 25. In some embodiments, n is 26. In some embodiments, n is 27. In some embodiments, n is 28. In some embodiments, n is 29. In some embodiments, n is 30.


The terms “about” or “approximately” are generally understood by persons knowledgeable in the relevant subject area, but in certain circumstances can mean within $\pm 10\%$, or within $\pm 5\%$, of a given value or range.

5 *Processes for Preparing Morpholino Oligomers*

Synthesis is generally prepared, as described herein, on a support-medium. In general a first synthon (e.g., a monomer, such as a morpholino subunit) is first attached to a support-medium, and the oligomer is then synthesized by sequentially coupling subunits to the support-bound synthon. This iterative elongation eventually results in a final oligomeric compound. Suitable support-media can be soluble or insoluble, or may possess variable solubility in different solvents to allow the growing support-bound polymer to be either in or out of solution as desired. Traditional support-media are for the most part insoluble and are routinely placed in reaction vessels while reagents and solvents react with and/or wash the growing chain until the oligomer has reached the target length, after which the oligomer is
10 cleaved from the support, and, if necessary, further worked up to produce the final polymeric compound. More recent approaches have introduced soluble supports including soluble polymer supports to allow precipitating and dissolving the iteratively synthesized product at desired points in the synthesis (Gravert et al., *Chem. Rev.* 1997, 97,489–510).

Provided herein are processes for preparing morpholino oligomers).


20 Thus, in one aspect, provided herein is a process for preparing a compound of Formula (II):



(II);

wherein R¹ is a support-medium;

25 wherein the process comprises contacting a compound of Formula (A1):




(A1);

wherein R¹ is a support-medium and R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

5 with a deblocking agent to form the compound of Formula (II).


In another aspect, provided herein is a process for preparing a compound of Formula (A3):



(A3);

10 wherein R¹ is a support-medium, and R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


wherein the process comprises contacting a compound of Formula (II):



(II);

15 wherein R¹ is a support-medium;

with a compound of Formula (A2):




(A2);

wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

to form the compound of Formula (A3).


In still another aspect, provided herein is a process for preparing a compound of Formula (IV):



(IV);

wherein R^1 is a support-medium;

wherein the process comprises contacting a compound of Formula (A3):




(A3);

wherein R^1 is a support-medium, and R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


with a deblocking agent to form a compound of Formula (IV).


In yet another aspect, provided herein is a process for preparing a compound of Formula (A5):




(A5);

- 5 wherein R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:






wherein the process comprises contacting a compound of Formula (IV):



5 (IV);


wherein R^1 is a support-medium;

with a compound of Formula (A4):




(A4);

10 wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:




to form a compound of Formula (A5).


- 5 In another aspect, provided herein is a process for preparing a compound of Formula (A9):



(A9);


wherein n is an integer from 10 to 40, R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each occurrence, selected from the group consisting of:





wherein the process comprises the sequential steps of:


(a) contacting a compound of Formula (IV):



(IV);


wherein R^1 is a support-medium;

with a compound of Formula (A4):




(A4);

wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:




to form a compound of Formula (A5):



(A5);

wherein R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and


R^4 is selected from the group consisting of:




(b) performing $n-1$ iterations of the sequential steps of:

(b1) contacting the product formed by the immediately prior step with a deblocking agent; and

10 (b2) contacting the compound formed by the immediately prior step with a compound of Formula (A8):




wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:




to form a compound of Formula (A9).

In yet another aspect, provided herein is a process for preparing a compound of Formula (A10):




(A10);


wherein n is an integer from 10 to 40, R^1 is a support-medium, and R^4 is, independently for each occurrence, selected from the group consisting of:



5




wherein the process comprises contacting a compound of Formula (A9):



5


(A9);

wherein n is an integer from 10 to 40, R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each occurrence, selected from the group consisting of:




with a deblocking agent to form a compound of Formula (A10).

5 In still another aspect, provided herein is a process for preparing a compound of Formula (A11):




(A11);


wherein n is an integer from 10 to 40, and R^4 is, for each occurrence independently selected from the group consisting of:



5




wherein the process comprises contacting the compound of Formula (A10):



5


(A10);

wherein n is an integer from 10 to 40, R^1 is a support-medium, and R^4 is, independently for each occurrence, selected from the group consisting of:




with a cleaving agent to form a compound of Formula (A11).


- 5 In another aspect, provided herein is a process for preparing an oligomeric compound of Formula (A):



wherein n is an integer from 10 to 40, and each R^2 is, independently for each occurrence,
 10 selected from the group consisting of:




wherein the process comprises contacting a compound of Formula (A11):



5


(A11);

wherein n is an integer from 10 to 40, and R^4 is, independently for each occurrence, selected from the group consisting of:




with a deprotecting agent to form the oligomeric compound of Formula (A).

- 5 In another aspect, provided herein is a process for preparing an oligomeric compound of Formula (A):




- wherein n is an integer from 10 to 40, and each R^2 is, independently for each occurrence,
 10 selected from the group consisting of:



wherein the process comprises the sequential steps of:


(a) contacting a compound of Formula (A1):



(A1);

wherein R^1 is a support-medium and R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


with a deblocking agent to form the compound of Formula (II):



(II);

wherein R^1 is a support-medium;


(b) contacting the compound of Formula (II) with a compound of Formula (A2):



(A2);

wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


5 to form a compound of Formula (A3):



(A3);

wherein R^1 is a support-medium, and R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


10 (c) contacting the compound of Formula (A3) with a deblocking agent to form a compound of Formula (IV):



(IV);


wherein R^1 is a support-medium;

15 (d) contacting the compound of Formula (IV) with a compound of Formula (A4):




(A4);


wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:




(PC),




(DPG),




(T),




(PA),




(P5mC),



(U),




(I), and



(PG),


to form a compound of Formula (A5):




(A5);

wherein R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and


5 R^4 is selected from the group consisting of:






(e) performing n-1 iterations of the sequential steps of:


- (e1) contacting the product formed by the immediately prior step with a deblocking agent; and
- (e2) contacting the compound formed by the immediately prior step with a compound of Formula (A8):




(A8);

- wherein R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is, independently for each compound of Formula (A8), selected from the group consisting of:






to form a compound of Formula (A9):



5


(A9);

wherein n is an integer from 10 to 40, R¹ is a support-medium, R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is, independently for each occurrence, selected from the group consisting of:




(f) contacting the compound of Formula (A9) with a deblocking agent to form a compound of

5 Formula (A10):




(A10);


wherein n is an integer from 10 to 40, R^1 is a support-medium, and R^4 is, independently for each occurrence, selected from the group consisting of:



5




(g) contacting the compound of Formula (A10) with a cleaving agent to form a compound of Formula (A11):



5

(A11);

wherein n is an integer from 10 to 40, and R^4 is, independently for each occurrence, selected from the group consisting of:



(h) contacting the compound of Formula (A11) with a deprotecting agent to form the
 5 oligomeric compound of Formula (A).

In one embodiment, step (d) or step (e2) further comprises contacting the compound of Formula (IV) or the compound formed by the immediately prior step, respectively, with a capping agent.

In another embodiment, each step is performed in the presence of at least one solvent.

10 In yet another embodiment, the deblocking agent used in each step is a solution comprising a halogenated acid.

In still another embodiment, the deblocking agent used in each step is cyanoacetic acid.

15 In another embodiment, the halogenated acid is selected from the group consisting of chloroacetic acid, dichloroacetic acid, trichloroacetic acid, fluoroacetic acid, difluoroacetic acid, and trifluoroacetic acid.

In another embodiment, the halogenated acid is trifluoroacetic acid.

In yet another embodiment, at least one of steps (a), (c), (e1), and (f) further comprises the step of contacting the deblocked compound of each step with a neutralization agent.

5 In still another embodiment, each of steps (a), (c), (e1), and (f) further comprises the step of contacting the deblocked compound of each step with a neutralization agent.

In another embodiment, the neutralization agent is in a solution comprising dichloromethane and isopropyl alcohol.

In yet another embodiment, the neutralization agent is a monoalkyl, dialkyl, or trialkyl amine.

10 In still another embodiment, the neutralization agent is N,N-diisopropylethylamine.

In another embodiment, the deblocking agent used in each step is a solution comprising 4-cyanopyridine, dichloromethane, trifluoroacetic acid, trifluoroethanol, and water.

15 In yet another embodiment, the capping agent is in a solution comprising ethylmorpholine and methylpyrrolidinone.

In still another embodiment, the capping agent is an acid anhydride.

In another embodiment, the acid anhydride is benzoic anhydride.

In another embodiment, the compounds of Formula (A4) and Formula (A8) are each, independently, in a solution comprising ethylmorpholine and dimethylimidazolidinone.

20 In another embodiment, the cleavage agent comprises dithiothreitol and 1,8-diazabicyclo[5.4.0]undec-7-ene.


In still another embodiment, the cleavage agent is in a solution comprising N-methyl-2-pyrrolidone.

In yet another embodiment, the deprotecting agent comprises NH_3 .

25 In still another embodiment, the deprotecting agent is in an aqueous solution.

In yet another embodiment, the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.

In another embodiment, the compound of Formula (A4) is of Formula (A4a):




(A4a);


wherein:

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

5 R^4 is selected from:



In another embodiment, the compound of Formula (A5) is of Formula (A5a):




(A5a);


wherein:

R^1 is a support-medium


5 R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

R^4 is selected from:





In yet another embodiment, the compound of Formula (A8) is of Formula (A8a):




5


(A8a);

wherein:


R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

10 R⁴ is, independently at each occurrence of the compound of Formula (A8a), selected from the group consisting of:





In still another embodiment, the compound of formula (A9) is of Formula (A9a):



5 (A9a);

wherein:


n is an integer from 10 to 40,

R¹ is a support-medium


R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl

10 and trimethoxytrityl, and

R⁴ is, independently for each occurrence, selected from the group consisting of:




In another embodiment, the compound of Formula (A10) is of Formula (A10a):




(A10a);


wherein:

n is an integer from 10 to 40,

5 R^1 is a support-medium, and R^4 is, independently for each occurrence, selected from the group consisting of:



In another embodiment, the compound of Formula (A11) is of Formula (A11a):




5

(A11a);

wherein:

n is an integer from 10 to 40, and


R^4 is, independently for each occurrence, selected from the group consisting of:



In an embodiment of the oligomeric compound of Formula (A), n is 30, and R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	C	11	A	21	G
2	T	12	A	22	C
3	C	13	G	23	A
4	C	14	G	24	T
5	A	15	A	25	T
6	A	16	A	26	T
7	C	17	G	27	C
8	A	18	A	28	T
9	T	19	T	29	A
10	C	20	G	30	G

wherein the oligomeric compound of Formula (A) is a compound of Formula (E):




or a pharmaceutically acceptable salt thereof.

- 5 Eteplirsen (see e.g., International Patent Application Publication No. WO 2006/000057, incorporated herein by reference in its entirety) has been the subject of clinical studies to test its safety and efficacy, and clinical development is ongoing. Eteplirsen is a phosphorodiamidate morpholino oligomer (PMO). The dystrophin therapeutic “Eteplirsen,” also known as “AVI-4658” is a PMO having the base sequence 5’-
- 10 CTCCAACATCAAGGAAGATGGCATTCTAG-3’ (SEQ ID NO:1). Eteplirsen is registered under CAS Registry Number 1173755-55-9. Chemical names include: RNA, [*P*-deoxy-*P*-(dimethylamino)](2’,3’-dideoxy-2’,3’-imino-2’,3’-seco)(2’a→5’)(C-m5U-C-C-A-A-C-A-m5U-C-A-A-G-G-A-A-G-A-m5U-G-G-C-A-m5U-m5U-m5U-C-m5U-A-G) (SEQ ID NO:2), 5’-[*P*-[4-[[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]carbonyl]-1-piperazinyl]-*N,N*-
- 15 dimethylphosphonamidate] and *P*,2’,3’-trideoxy-*P*-(dimethylamino)-5’-O-{*P*-[4-(10-hydroxy-2,5,8- trioxadecanoyl)piperazin-1-yl]-*N,N*-dimethylphosphonamidoyl}-2’,3’-imino-2’,3’-secocytidylyl-(2’a→5’)-*P*,3’-dideoxy-*P*-(dimethylamino)-2’,3’-imino-2’,3’-secothymidylyl-(2’a→5’)-*P*,2’,3’-trideoxy-*P*-(dimethylamino)-2’,3’-imino-2’,3’-secocytidylyl-(2’a→5’)- *P*,2’,3’-trideoxy-*P*-(dimethylamino)-2’,3’-imino-2’,3’-secocytidylyl-(2’a→5’)-*P*,2’,3’- trideoxy-*P*-
- 20 (dimethylamino)-2’,3’-imino-2’,3’-secoadenylyl-(2’a→5’)-*P*,2’,3’-trideoxy-*P*- (dimethylamino)-2’,3’-imino-2’,3’-secoadenylyl-(2’a→5’)-*P*,2’,3’-trideoxy-*P*- (dimethylamino)-2’,3’-imino-2’,3’-

secocytidylyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-
 (2'a→5')-P,3'-dideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secothymidylyl-(2'a→5')-P,2',3'-
 trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secocytidylyl-(2'a→5')-P,2',3'-trideoxy-P-
 (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-
 5 2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-
 secoguanilyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secoguanilyl-
 (2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,2',3'-
 trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,2',3'-trideoxy-P-
 (dimethylamino)-2',3'-imino-2',3'-secoguanilyl-(2'a→5')-P,2',3'-trideoxy-P-
 10 (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,3'-dideoxy-P- (dimethylamino)-
 2',3'-imino-2',3'-secothymidylyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-
 2',3'-secoguanilyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-
 secoguanilyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secocytidylyl-
 (2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secoadenylyl-(2'a→5')-P,3'-
 15 dideoxy-P- (dimethylamino)-2',3'-imino-2',3'-secothymidylyl-(2'a→5')-P,3'-dideoxy-P-
 (dimethylamino)-2',3'-imino-2',3'-secothymidylyl-(2'a→5')-P,3'-dideoxy-P- (dimethylamino)-
 2',3'-imino-2',3'-secothymidylyl-(2'a→5')-P,2',3'-trideoxy-P- (dimethylamino)-2',3'-imino-
 2',3'-secocytidylyl-(2'a→5')-P,3'-dideoxy-P-(dimethylamino)- 2',3'-imino-2',3'-
 secothymidylyl-(2'a→5')-P,2',3'-trideoxy-P-(dimethylamino)-2',3'-imino- 2',3'-secoadenylyl-
 20 (2'a→5')-2',3'-dideoxy-2',3'-imino-2',3'-secoguanosine.


Eteplirsen has the following structure:



B(1-30):


C-T-C-C-A-A-C-A-T-C-A-A-G-G-A-A-G-A-T-G-G-C-A-T-T-T-C-T-A-G (SEQ ID NO:1)

Eteplirsen can also be depicted by the structure of Formula (XII):




Thus, in one embodiment of the process described above, the oligomeric compound of

5 Formula (A) is a compound of Formula (E):



or a pharmaceutically acceptable salt thereof.

In yet another embodiment, the oligomeric compound of Formula (E) is an oligomeric compound of Formula (XII):



5


(XII),

or a pharmaceutically acceptable salt thereof.

In still another embodiment, R^3 is, at each occurrence, trityl.


In another aspect, provided herein is a process for preparing an oligomeric compound

10 of Formula (S):




(S);

wherein n is an integer from 10 to 40, and each R^2 is, independently for each occurrence, selected from the group consisting of:



wherein the process comprises the sequential steps of:


(a) contacting a compound of Formula (II):



(II);


wherein R^1 is a support-medium;

with a compound of Formula (S2):




(S2);

wherein R^4 is selected from the group consisting of:




wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

to form a compound of Formula (S3):




(S3);

wherein R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is selected from the group consisting of:




- 5 (b) contacting the compound of Formula (S3) with a deblocking agent to form a compound of Formula (IV-S):




(IV-S);

wherein R¹ is a support-medium, R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is selected from the group consisting of:

10




(c) contacting the compound of Formula (IV-S) with a compound of Formula (A4):




(A4);

wherein R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is selected from the group consisting of:



to form a compound of Formula (S5):




5

(S5);

wherein R¹ is a support-medium, R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

R^4 , independently at each occurrence, is selected from the group consisting of:




5

(d) performing n-2 iterations of the sequential steps of:


(d1) contacting the product formed by the immediately prior step with a deblocking agent; and

(d2) contacting the compound formed by the immediately prior step with a compound


10 of Formula (A8):



wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each compound of Formula (A8), selected from the group consisting of:




to form a compound of Formula (S9):




(S9);


wherein n is an integer from 10 to 40, R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is,

5 independently for each occurrence, selected from the group consisting of:






(e) contacting the compound of Formula (S9) with a deblocking agent to form a compound of Formula (S10):




(S10);

wherein n is an integer from 10 to 40, R^1 is a support-medium, and R^4 is, independently for each occurrence, selected from the group consisting of:




(f) contacting the compound of Formula (S10) with a cleaving agent to form a compound of

5 Formula (S11):



(S11);

wherein n is an integer from 10 to 40, and R^4 is, independently for each occurrence, selected from the group consisting of:




(g) contacting the compound of Formula (S11) with a deprotecting agent to form the oligomeric compound of Formula (S).

Processes for Preparing Eteplirsen


10 Provided herein are processes for preparing Eteplirsen.

In another aspect, provided herein is a process for preparing an oligomeric compound of Formula (E):




wherein the process comprises the sequential steps of:

(a) contacting a compound of Formula (I):



wherein R^1 is a support-medium,


with a deblocking agent to form the compound of Formula (II):



(II);


wherein R¹ is a support-medium;

(b) contacting the compound of Formula (II) with compound (B):



(B);


to form a compound of Formula (III):



(III);

wherein R¹ is a support-medium;


(c) contacting the compound of Formula (III) with a deblocking agent to form a compound of Formula (IV):



(IV);


wherein R¹ is a support-medium;

(d) contacting the compound of Formula (IV) with a compound of Formula (C):



(C);

to form a compound of Formula (V):




(V);

wherein R^1 is a support-medium;

(e) contacting the compound of Formula (V) with a deblocking agent to form a compound of


5 Formula (VI):



(VI);


wherein R^1 is a support-medium;

(f) contacting the compound of Formula (VI) with a compound of Formula (F):



(F);

to form a compound of Formula (VII):




(VII);

wherein R¹ is a support-medium;

(g) performing 28 iterations of the sequential steps of:


(g1) contacting the product formed by the immediately prior step with a deblocking agent; and

(g2) contacting the compound formed by the immediately prior step with a compound of Formula (VIII):



(VIII);

- 5 wherein R² is, independently for each compound of Formula (VIII), selected from the group consisting of:




wherein, for each iteration from 1 to 28, R² is:

Iteration No.	R ²	Iteration No.	R ²	Iteration No.	R ²
1	PC	11	DPG	21	PA
2	PC	12	DPG	22	T
3	PA	13	PA	23	T


4	PA	14	PA	24	T
5	PC	15	DPG	25	PC
6	PA	16	PA	26	T
7	T	17	T	27	PA
8	PC	18	DPG	28	DPG
9	PA	19	DPG		
10	PA	20	PC		

to form a compound of Formula (IX):



(IX);


5 wherein R¹ is a support-medium,
wherein R² is, independently for each occurrence, selected from the group consisting
of:



wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


- 5 (h) contacting the compound of Formula (IX) with a deblocking agent to form a compound of Formula (X):



wherein R^1 is a support-medium,

wherein R^2 is, independently for each occurrence, selected from the group consisting


5 of:



wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


(i) contacting the compound of Formula (X) with a cleaving agent to form a compound of Formula (XI):



5

(XI),

wherein R² is, independently for each occurrence, selected from the group consisting of:



wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

5 and

(j) contacting the compound of Formula (XI) with a deprotecting agent to form the oligomeric compound of Formula (E).

In an embodiment, step (d), step (f), step g(2), or combinations thereof further comprises contacting the compound of Formula (IV), Formula (VI), or the compound formed
 10 by the immediately prior step, respectively, with a capping agent.

In certain embodiments, each of step (d), step (f) and step g(2) further comprises contacting the compound of Formula (IV), Formula (VI), or the compound formed by the immediately prior step, respectively, with a capping agent.

In another embodiment, each step is performed in the presence of at least one solvent.

5 In yet another embodiment, the deblocking agent used in each step is a solution comprising a halogenated acid.

In still another embodiment, the deblocking agent used in each step is cyanoacetic acid.

10 In another embodiment, the halogenated acid is selected from the group consisting of chloroacetic acid, dichloroacetic acid, trichloroacetic acid, fluoroacetic acid, difluoroacetic acid, and trifluoroacetic acid.

In yet another embodiment, the halogenated acid is trifluoroacetic acid.

In still another embodiment, at least one of steps (c), (e1), and (f) further comprise the step of contacting the deblocked compound of each step with a neutralization agent.

15 In another embodiment, each of steps (c), (e1), and (f) further comprise the step of contacting the deblocked compound of each step with a neutralization agent.

In yet another embodiment, the neutralization agent is in a solution comprising dichloromethane and isopropyl alcohol.

20 In still another embodiment, the neutralization agent is a monoalkyl, dialkyl, or trialkyl amine.

In another embodiment, the neutralization agent is N,N-diisopropylethylamine.

In yet another embodiment, the deblocking agent used in each step is a solution comprising 4-cyanopyridine, dichloromethane, trifluoroacetic acid, trifluoroethanol, and water.

25 In still another embodiment, the capping agent is in a solution comprising ethylmorpholine and methylpyrrolidinone.

In another embodiment, the capping agent is an acid anhydride.

In yet another embodiment, the acid anhydride is benzoic anhydride.

30 In still another embodiment, the compound of Formula (VIII), compound (C), and compound (F) are each, independently, in a solution comprising ethylmorpholine and dimethylimidazolidinone.

In another embodiment, the cleavage agent comprises dithiothreitol and 1,8-diazabicyclo[5.4.0]undec-7-ene.


In yet another embodiment, the cleavage agent is in a solution comprising N-methyl-2-pyrrolidone.

In still another embodiment, the deprotecting agent comprises NH_3 .

In another embodiment, the deprotecting agent is in an aqueous solution.


5 In yet another embodiment, the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.

In another embodiment, the compound of Formula (C) is of Formula (C1):



(C1).


10 In another embodiment, the compound of Formula (V) is of Formula (Va):



(Va),


wherein R^1 is a support-medium.

In another embodiment, the compound of Formula (F) is of Formula (F1):



(F1).


In another embodiment, the compound of Formula (VII) is of Formula (VIIa):



(VIIa),


wherein R^1 is a support-medium.

In another embodiment, the compound of Formula (VIII) is of Formula (VIIIa):




(VIIIa);

wherein R^2 is, independently for each compound of Formula (VIIIa), selected from the group consisting of:




5



(T), and

(PA).

In another embodiment, the compound of Formula (IX) is of Formula (IXa):




(IXa),

or a pharmaceutically acceptable salt thereof, wherein

 R^1 is a support-medium, and


5 R^2 is, independently at each occurrence, selected from the group consisting of:



wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

In another embodiment, the compound of Formula (X) is of Formula (Xa):




5

(Xa),

or a pharmaceutically acceptable salt thereof, wherein

R^1 is a support-medium, and

R^2 is, independently at each occurrence, selected from the group consisting of:




wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


5

In another embodiment, the compound of Formula (XI) is of Formula (XIa):




(XIa),

or a pharmaceutically acceptable salt thereof, wherein:

 R^2 is, independently at each occurrence, selected from the group consisting of:wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


In another embodiment, the compound of Formula (VI) is of Formula (VIa):



5 (VIa);


wherein R^1 is a support-medium.

In still another embodiment, the oligomeric compound of Formula (E) is an oligomeric compound of Formula (XII):



(XII).

In another aspect, provided herein is a compound of Formula (IX):




(IX),

or a pharmaceutically acceptable salt thereof, wherein:

R^1 is a support-medium, and


5 R^2 is, independently at each occurrence, selected from the group consisting of:



wherein R² is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

In one embodiment, the compound of Formula (IX) is of Formula (IXa):




(IXa),

or a pharmaceutically acceptable salt thereof, wherein


 R^1 is a support-medium, and R^2 is, independently at each occurrence, selected from the group consisting of:

5

wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


In another aspect, provided herein is a compound of Formula (A9):




(A9),


or a pharmaceutically acceptable salt thereof, wherein:

n is an integer from 10 to 40;

5 R¹ is a support-medium;R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl; andR⁴ is, independently at each occurrence, selected from the group consisting of:



In one embodiment, the compound of Formula (A9) is of Formula (A9a):



5

(A9a),

or a pharmaceutically acceptable salt thereof, wherein:


n is an integer from 10 to 40;

R^1 is a support-medium;


R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl

10 and trimethoxytrityl; and

R^4 is, independently at each occurrence, selected from the group consisting of:



In another aspect, provided herein is a compound of Formula (X):




(X);

or a pharmaceutically acceptable salt thereof, wherein


 R^1 is a support-medium, and R^2 is, independently at each occurrence, selected from the group consisting of:

5

wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

In one embodiment, the compound of Formula (X) is of Formula (Xa):




(Xa),

or a pharmaceutically acceptable salt thereof, wherein

R^1 is a support-medium, and


5 R^2 is, independently at each occurrence, selected from the group consisting of:



wherein R² is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

In another aspect, provided herein is a compound of Formula (A10):



(A10),


5

or a pharmaceutically acceptable salt thereof, wherein:


n is an integer from 10 to 40;

R¹ is a support-medium; and

R⁴ is, independently at each occurrence, selected from the group consisting of:



In one embodiment, the compound of Formula (A10) is of Formula (A10a):





(A10a),

or a pharmaceutically acceptable salt thereof, wherein:

n is an integer from 10 to 40;


5

R¹ is a support-medium; andR⁴ is, independently at each occurrence, selected from the group consisting of:



In another embodiment of these compounds, the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.


5 In another aspect, provided herein is a compound of Formula (XI):



(XI),

or a pharmaceutically acceptable salt thereof, wherein:

R^2 is, independently at each occurrence, selected from the group consisting of:




wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2	Position No. 5' to 3'	R^2
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG


5

In one embodiment, the compound of Formula (XI) is of Formula (XIa):




(XIa),

or a pharmaceutically acceptable salt thereof, wherein:

 R^2 is, independently at each occurrence, selected from the group consisting of:wherein R^2 is at each position from 1 to 30 and 5' to 3':

Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²	Position No. 5' to 3'	R ²
1	PC	11	PA	21	DPG
2	T	12	PA	22	PC
3	PC	13	DPG	23	PA
4	PC	14	DPG	24	T
5	PA	15	PA	25	T
6	PA	16	PA	26	T
7	PC	17	DPG	27	PC
8	PA	18	PA	28	T
9	T	19	T	29	PA
10	PC	20	DPG	30	DPG

In another aspect, provided herein is a compound of Formula (A11):



(A11),


5

or a pharmaceutically acceptable salt thereof, wherein:


n is an integer from 10 to 40;

R¹ is a support-medium; and

R⁴ is, independently at each occurrence, selected from the group consisting of:



In one embodiment, the compound of Formula (A11) is of formula (A11a):





(A11a),

or a pharmaceutically acceptable salt thereof, wherein:

n is an integer from 10 to 40;

5


R¹ is a support-medium; andR⁴ is, independently at each occurrence, selected from the group consisting of:



Oligomers

5 Oligomers prepared by the process of the disclosure may contain a variety of nucleotide analog subunits.

In another aspect of the disclosure is provided a process for preparing an oligomeric compound of Formula (D):



10 (D),


or a pharmaceutically acceptable salt thereof, wherein:

R¹ is selected from -O(C₁-C₆alkyl), halo, -O(C₁-C₆alkyl)-O(C₁-C₆alkyl), and -O(C₁-C₆alkyl)-C(O)-NH-C₁-C₆alkyl;

15 R² is H or R¹ and R² join to form a cycloalkyl or heterocyclic ring having from 4 to 7 ring atoms optionally containing an oxygen;

R⁵, R⁶, and R⁷ are each hydrogen, or R⁵ and R⁷ join to form a cycloalkyl ring having from 4 to 7 ring atoms and R⁶ and R⁷ join to form a cycloalkyl ring having from 3 to 4 ring atoms;

R⁴ is, independently at each occurrence, selected from the group consisting of:




R^8 is, independently at each occurrence, selected from the group consisting of H, trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


5 n is 10 to 40; and

X is O or S.

In some embodiments, the oligomeric compound of Formula (D) is of Formula (G):



10 wherein R^2 is, independently at each occurrence, selected from the group consisting of:




n is 10 to 40.

Embodiments of the compounds of Formula D and G include, for example:

1. Locked Nucleic Acids (LNAs)

Modified antisense oligomer compounds may also contain “locked nucleic acid” subunits (LNAs). “LNAs” are a member of a class of modifications called bridged nucleic acid (BNA). BNA is characterized by a covalent linkage that locks the conformation of the ribose ring in a C3'-endo (northern) sugar pucker. For LNA, the bridge is composed of a methylene between the 2'-O and the 4'-C positions. LNA enhances backbone preorganization and base stacking to increase hybridization and thermal stability.


The structures of LNAs can be found, for example, in Singh et al., *Chem. Commun.* 1998, 455; Wengel, *Tetrahedron* 1998, 54, 3607; Wengel, *Acc. Chem. Res.* 1999, 32, 301; Obika et al., *Tetrahedron Lett.* 1997, 38, 8735; Obika et al., *Tetrahedron Lett.* 1998, 39, 5401; and Obika et al., *Bioorg. Med. Chem.* 2008, 16, 9230, which are hereby incorporated by reference in their entirety. A non-limiting example of an LNA oligomer comprising LNA subunits and phosphodiester internucleoside linkages is depicted below:



LNA

Compounds of the disclosure may incorporate one or more LNAs; in some cases, the compounds may be entirely composed of LNAs. Methods for the synthesis of individual LNA nucleoside subunits and their incorporation into oligomers are described, for example, in U.S. Patent Nos. 7,572,582, 7,569,575, 7,084,125, 7,060,809, 7,053,207, 7,034,133, 6,794,499, and 6,670,461, which are hereby incorporated by reference in their entirety. Typical internucleoside linkers include phosphodiester and phosphorothioate moieties; alternatively, non-phosphorous containing linkers may be employed. Further embodiments include an LNA containing compound where each LNA subunit is separated by a DNA subunit. Certain compounds are composed of alternating LNA and DNA subunits where the internucleoside linker is phosphorothioate.


2'O,4'C-ethylene-bridged nucleic acids (ENAs) are another member of the class of BNAs. A non-limiting example of an ENA subunit and phosphodiester internucleoside linkage is depicted below:



ENA oligomers and their preparation are described in Obika et al., *Tetrahedron Lett.* 1997, 38, 8735, which is hereby incorporated by reference in its entirety. Compounds of the disclosure may incorporate one or more ENA subunits.

2. Phosphorothioates

“Phosphorothioates” (or S-oligos) are a variant of native DNA or RNA in which one of the nonbridging oxygens of the phosphodiester internucleoside linkages is replaced by sulfur. A non-limiting example of a phosphorothioate DNA (left), comprising deoxyribose subunits and phosphorothioate internucleoside linkages, and phosphorothioate RNA (right), comprising ribose subunits and phosphorothioate internucleoside linkages, are depicted below:




The sulfurization of the internucleoside bond reduces the action of endo- and exonucleases including 5' to 3' and 3' to 5' DNA Pol 1 exonuclease, nucleases S1 and P1, RNases, serum nucleases and snake venom phosphodiesterase. Phosphorothioates may be made by two principal routes: by the action of a solution of elemental sulfur in carbon disulfide on a hydrogen phosphonate, or by the method of sulfurizing phosphite triesters with either tetraethylthiuram disulfide (TETD) or 3H-1, 2-benzodithiol-3-one 1, 1-dioxide (BDTD) (see, e.g., Iyer et al., *J. Org. Chem.* 1990, 55, 4693-4699, which is hereby incorporated by reference in its entirety). The latter methods avoid the problem of elemental

sulfur's insolubility in most organic solvents and the toxicity of carbon disulfide. The TETD and BDTD methods also yield higher purity phosphorothioates.

3. Tricyclo-DNAs and Tricyclo-Phosphorothioate Nucleotides

Tricyclo-DNAs (tc-DNA) are a class of constrained DNA analogs in which each nucleotide is modified by the introduction of a cyclopropane ring to restrict conformational flexibility of the backbone and to optimize the backbone geometry of the torsion angle γ . Homobasic adenine- and thymine-containing tc-DNAs form extraordinarily stable A-T base pairs with complementary RNAs. Tricyclo-DNAs and their synthesis are described in PCT Publication No. WO 2010/115993, which is hereby incorporated by reference in its entirety. Compounds of the disclosure may incorporate one or more tricyclo-DNA subunits; in some cases, the compounds may be entirely composed of tricyclo-DNA subunits.

Tricyclo-phosphorothioate nucleotides are tricyclo-DNA subunits with phosphorothioate internucleoside linkages. Tricyclo-phosphorothioate nucleotides and their synthesis are described in PCT Publication No. WO 2013/053928, which is hereby incorporated by reference in its entirety. Compounds of the disclosure may incorporate one or more tricyclo-DNA subunits; in some cases, the compounds may be entirely composed of tricyclo-DNA nucleotides. A non-limiting example of a tricyclo-DNA/tricycle subunit and phosphodiester internucleoside linkage is depicted below:




tricyclo-DNA


4. 2' O-Methyl, 2' O-MOE, and 2'-F Oligomers

"2'-O-Me oligomer" molecules comprise subunits that carry a methyl group at the 2'-OH residue of the ribose molecule. 2'-O-Me-RNAs show the same (or similar) behavior as DNA, but are protected against nuclease degradation. 2'-O-Me-RNAs can also be combined with phosphorothioate oligomers (PTOs) for further stabilization. 2'-O-Me oligomers (wherein the 2'-OMe subunits are connected by phosphodiester or phosphorothioate internucleoside linkages) can be synthesized according to routine techniques in the art (see,


e.g., Yoo et al., *Nucleic Acids Res.* 2004, 32, 2008-16, which is hereby incorporated by reference in its entirety). A non-limiting example of a 2' O-Me oligomer comprising 2'-OMe subunits and phosphodiester intersubunit linkages is depicted below:



- 5 2' O-Me oligomers may also comprise a phosphorothioate linkage (2' O-Me phosphorothioate oligomers). 2' O-Methoxyethyl oligomers (2'-O MOE), like 2' O-Me oligomers, comprise subunits that carry a methoxyethyl group at the 2'-OH residue of the ribose molecule and are discussed in Martin et al., *Helv. Chim. Acta* 1995, 78, 486-504, which is hereby incorporated by reference in its entirety. A non-limiting example of a 2' O-
- 10 MOE subunit is depicted below:



- In contrast to the preceding alkylated 2'OH ribose derivatives, 2'-fluoro oligomers comprise subunits that have a fluoro radical at the 2' position in place of the 2'OH. A non-limiting example of a 2'-F oligomer comprising 2'-F subunits and phosphodiester internucleoside linkages is depicted below:
- 15



2'-fluoro oligomers are further described in WO 2004/043977, which is hereby incorporated by reference in its entirety. Compounds of the disclosure may incorporate one or


more 2'-O-Methyl, 2' O-MOE, and 2'-F subunits and may utilize any of the internucleoside linkages described here. In some instances, a compound of the disclosure could be composed of entirely 2'-O-Methyl, 2' O-MOE, or 2'-F subunits. One embodiment of a compound of the disclosure is composed entirely of 2'-O-methyl subunits.

5

5. 2'-O-[2-(N-methylcarbamoyl)ethyl] Oligomers (MCEs)

MCEs are another example of 2'-O modified ribonucleotides useful in the compounds of the disclosure. Here, the 2'-OH is derivatized to a 2-(N-methylcarbamoyl)ethyl moiety to increase nuclease resistance. A non-limiting example of an MCE oligomer comprising MCE subunits and phosphodiester internucleoside linkages is depicted below:


10



MCEs and their synthesis are described in Yamada et al., *J. Org. Chem.* 2011, 76(9), 3042-53, which is hereby incorporated by reference in its entirety. Compounds of the disclosure may incorporate one or more MCE subunits.

15


In another aspect, provided herein is a process for preparing an oligomeric compound of Formula (F):




(F);

wherein:

each A is, independently for each occurrence, selected from the group consisting of:



wherein each R^2 is, independently for each occurrence, selected from the group consisting of:




n is an integer from 10 to 40, and


R is H or acetyl,

wherein the process comprises the sequential steps of:

(a) contacting a compound of Formula (II):



$$(II);$$

wherein R¹ is a support-medium with a compound of Formula (F2):




(F2);

wherein A is selected from the group consisting of:




wherein R⁴ is selected from the group consisting of:




R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and PG is a protecting group;

to form a compound of Formula (F3):




(F3);

wherein R^1 is a support-medium, A is selected from the group consisting of:




wherein R^4 is selected from the group consisting of:



R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and PG is a protecting group;


(b) contacting the compound of Formula (F3) with a deblocking agent to form a compound of Formula (IV-F):



5


(IV-F);

wherein R^1 is a support-medium, A is selected from the group consisting of:




10

wherein R⁴ is selected from the group consisting of:




and PG is a protecting group;


- 5 (c) contacting the compound of Formula (IV-F) with a compound of Formula (F4):




wherein R⁵ is $-\text{OCH}_2\text{CH}_2\text{CN}$ or $-\text{CH}_2\text{C}(=\text{O})\text{OC}(\text{CH}_3)_2\text{CH}_2\text{CN}$, A is selected from the group consisting of:



wherein R^4 is selected from the group consisting of:



R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and PG is a protecting group, to form a compound of Formula (F5):




5


(F5);

wherein R^1 is a support-medium,

each A is, independently at each occurrence, selected from the group consisting of:



each R^4 is, independently at each occurrence, selected from the group consisting of:



R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl,

X is O or S, and


R^5 is $-\text{OCH}_2\text{CH}_2\text{CN}$ or $-\text{CH}_2\text{C}(=\text{O})\text{OC}(\text{CH}_3)_2\text{CH}_2\text{CN}$,

wherein where X is S, then step (d) further comprises a sulfurization step, or where X is O, then step (d) further comprises an oxidation step;

(e) performing n-2 iterations of the sequential steps of:

(e1) contacting the product formed by the immediately prior step with a deblocking agent; and


(e2) contacting the compound formed by the immediately prior step with a compound of Formula (F8):




(F8);


wherein R^5 is $-\text{OCH}_2\text{CH}_2\text{CN}$ or $-\text{CH}_2\text{C}(=\text{O})\text{OC}(\text{CH}_3)_2\text{CH}_2\text{CN}$,

each A is, independently for each iteration of (F8), selected from the group consisting of:




wherein R^4 is, independently for each iteration of (F8), selected from the group consisting of:






R^3 is, independently for each iteration of (F8), selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and PG is a protecting group,

to form a compound of Formula (F9):





(F9);

wherein n is an integer from 10 to 40, R^1 is a support-medium, X is O or S, R^5 is $-OCH_2CH_2CN$ or $-CH_2C(=O)OC(CH_3)_2CH_2CN$, each A, independently at each occurrence, is selected from the group consisting of:




R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each occurrence, selected from the group consisting of:






wherein where X is S, then each iteration further comprises a sulfurization step, or where X is O, then each iteration further comprises an oxidation step; and

- 5 (f) contacting the compound of Formula (F9) with a deblocking agent to form a compound of Formula (F10):





(F10);

wherein n is an integer from 10 to 40, R^1 is a support-medium, X is O or S, R^5 is $-OCH_2CH_2CN$ or $-CH_2C(=O)OC(CH_3)_2CH_2CN$, each A, independently at each occurrence, is selected from the group consisting of:




and R^4 is, independently for each occurrence, selected from the group consisting of:





(g) contacting the compound of Formula (F10) with a cleaving agent to form a compound of Formula (F11):

5




(F11);


wherein n is an integer from 10 to 40,

X is O or S, R^5 is $-\text{OCH}_2\text{CH}_2\text{CN}$ or $-\text{CH}_2\text{C}(=\text{O})\text{OC}(\text{CH}_3)_2\text{CH}_2\text{CN}$, each A,

10 independently at each occurrence, is selected from the group consisting of:






and R^4 is, independently for each occurrence, selected from the group consisting of:





(h) contacting the compound of Formula (F11) with a deprotecting agent to form the oligomeric compound of Formula (F).

The disclosure further provides a compound selected from:










wherein n is an integer from 10 to 40, R^1 is a support-medium, X is O or S, R^5 is $-\text{OCH}_2\text{CH}_2\text{CN}$ or $-\text{CH}_2\text{C}(=\text{O})\text{OC}(\text{CH}_3)_2\text{CH}_2\text{CN}$, each A, independently at each occurrence,

5 is selected from the group consisting of:



R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl
5 and trimethoxytrityl, and R⁴ is, independently for each occurrence, selected from the group
consisting of:





Morpholino Oligomers


5 Important properties of morpholino-based subunits include: 1) the ability to be linked in an oligomeric form by stable, uncharged or positively charged backbone linkages; 2) the ability to support a nucleotide base (e.g. adenine, cytosine, guanine, thymidine, uracil, 5-methyl-cytosine and hypoxanthine) such that the polymer formed can hybridize with a complementary-base target nucleic acid, including target RNA; 3) the ability of the oligomer to be actively or passively transported into mammalian cells; and 4) the ability of the oligomer and oligomer:RNA heteroduplex to resist RNase and RNase H degradation, respectively.

In some embodiments, the antisense oligomers contain base modifications or substitutions. For example, certain nucleobases may be selected to increase the binding affinity of the antisense oligomers described herein. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C, and may be incorporated into the antisense oligomers described herein. In one embodiment, at least one pyrimidine base of the oligomer comprises a 5-substituted pyrimidine base, wherein the pyrimidine base is selected from the group consisting of cytosine, thymine and uracil. In one embodiment, the 5-substituted pyrimidine base is 5-methylcytosine. In another embodiment, at least one purine base of the oligomer comprises hypoxanthine.

Morpholino-based oligomers (including antisense oligomers) are detailed, for example, in U.S. Patent Nos. 5,698,685, 5,217,866, 5,142,047, 5,034,506, 5,166,315, 5,185,444, 5,521,063, 5,506,337, 8,299,206, and 8,076,476; International Patent Application Publication Nos. WO/2009/064471 and WO/2012/043730; and Summerton et al., *Antisense*


Nucleic Acid Drug Dev. 1997, 7, 187-195, each of which are hereby incorporated by reference in their entirety.

In some embodiments, the disclosure provides a process for preparing an oligomeric compound of Formula (B):




(B);

wherein n is an integer from 10 to 40,



X is, independently at each occurrence, selected from H, OH, $-\text{OCH}_2\text{CH}_2\text{CN}$, $\text{N}(\text{R}^{17})_2$,



wherein

R^{17} is, independently for each occurrence, H or C_{1-6} -alkyl;

R^{18} is selected from the group consisting of H, C_{1-6} -alkyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(\text{O})(\text{CH}_2)_q\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$, and $-\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$;

q is 1, 2, 3, 4, or 5;

R^{19} is, independently for each occurrence, H or methyl;

R^{20} is H or C_{1-6} -alkyl;

R^{21} is selected from the group consisting of H, C_{1-6} -alkyl, $\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(\text{O})(\text{CH}_2)_r\text{NR}^{23}\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_3\text{NHC}(=\text{NH})\text{NH}_2$,


$\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{23}\text{C}(=\text{NH})\text{NH}_2$, and $\text{C}(\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_4\text{NH}_2$;

R^{22} is H or C_{1-6} -alkyl;

R^{23} is H or C_{1-6} -alkyl; and


r is 1, 2, 3, 4, or 5;

each R² is, independently for each occurrence, selected from the group consisting of:



5 wherein the process comprises the sequential steps of:


(a) contacting a compound of Formula (B1):



(B1);

wherein

- 10 R¹¹ is selected from the group consisting of halo, CN, and NO₂;
 R¹² is C₁₋₆-alkyl;
 R¹³ is H or C₁₋₆-alkyl; and
 R¹⁴ is a first linker bound to a support-medium,
 R³ is selected from the group consisting of trityl, monomethoxytrityl,
 15 dimethoxytrityl and trimethoxytrityl;
 with a deblocking agent to form the compound of Formula (II-B):



(II-B);

wherein


R^{11} is selected from the group consisting of halo, CN, and NO_2 ;

R^{12} is C_{1-6} -alkyl;

R^{13} is H or C_{1-6} -alkyl; and

R^{14} is a first linker bound to a support-medium;

(b) contacting the compound of Formula (II) with a compound of Formula (III):




(III-B);

wherein


w is selected from O or C;

t is 1, 2, 3, 4, or 5;

R^8 is selected from:




wherein R^4 is selected from the group consisting of:



R^3 is, independently at each occurrence, selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

R^9 and R^{10} are, independently, C_{1-6} -alkyl, or R^9 and R^{10} , together with the atoms to which they are attached, form a cyclic or polycyclic ring comprising 4-10 carbon atoms to form a compound of Formula (B3):



(B3);

wherein

w is selected from O or C;

t is 1, 2, 3, 4, or 5;


R¹¹ is selected from the group consisting of halo, CN, and NO₂;

R¹² is C₁₋₆-alkyl;


R¹³ is H or C₁₋₆-alkyl;

5 R¹⁴ is a first linker bound to a support-medium;

R⁸ is selected from:




wherein R⁴ is selected from the group consisting of:



R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

(c) contacting the compound of Formula (A3) with a deblocking agent to form a compound of Formula (IV-B):



(IV-B);

5 wherein

w is selected from O or C;

t is 1, 2, 3, 4, or 5;


R¹¹ is selected from the group consisting of halo, CN, and NO₂;

R¹² is C₁₋₆-alkyl;


10 R¹³ is H or C₁₋₆-alkyl;

R¹⁴ is a first linker bound to a support-medium; and

R⁸ is selected from:




wherein R⁴ is selected from the group consisting of:




15 (PC),

(DPG),




(d) contacting the compound of Formula (IV-B) with a compound of Formula (B4):



(B4);

wherein

X is, independently at each occurrence, selected from H, OH, -OCH₂CH₂CN,



R¹⁷ is, independently for each occurrence, H or C₁₋₆-alkyl;

R¹⁸ is selected from the group consisting of H, C₁₋₆-alkyl, -C(=NH)NH₂, -C(O)(CH₂)_qNR²⁰C(=NH)NH₂, and -C(O)(CH₂)₂NHC(O)(CH₂)₅NR²⁰C(=NH)NH₂;

q is 1, 2, 3, 4, or 5;

R¹⁹ is, independently for each occurrence, H or methyl;

R²⁰ is H or C₁₋₆-alkyl;

R²¹ is selected from the group consisting of H, C₁₋₆-alkyl, C(=NH)NH₂, C(O)(CH₂)_rNR²³C(=NH)NH₂, C(O)CH(NH₂)(CH₂)₃NHC(=NH)NH₂, C(O)(CH₂)₂NHC(O)(CH₂)₅NR²³C(=NH)NH₂, and C(O)CH(NH₂)(CH₂)₄NH₂;


R²² is H or C₁₋₆-alkyl;

R^{23} is H or C_{1-6} -alkyl; and


r is 1, 2, 3, 4, or 5;

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

5 R^4 is selected from the group consisting of:



to form a compound of Formula (B5):




(B5);

wherein:

w is selected from O or C;

t is 1, 2, 3, 4, or 5;

 R^{11} is selected from the group consisting of halo, CN, and NO_2 ; R^{12} is C_{1-6} -alkyl; R^{13} is H or C_{1-6} -alkyl; R^{14} is a first linker bound to a support-medium;X is, independently at each occurrence, selected from H, OH, $-\text{OCH}_2\text{CH}_2\text{CN}$,

wherein:

 R^{17} is, independently for each occurrence, H or C_{1-6} -alkyl; R^{18} is selected from the group consisting of H, C_{1-6} -alkyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(\text{O})(\text{CH}_2)_q\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$,and $-\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$;

q is 1, 2, 3, 4, or 5;

 R^{19} is, independently for each occurrence, H or methyl; R^{20} is H or C_{1-6} -alkyl;

R^{21} is selected from the group consisting of H, C_{1-6} -alkyl, $C(=NH)NH_2$, $C(O)(CH_2)_rNR^{23}C(=NH)NH_2$, $C(O)CH(NH_2)(CH_2)_3NHC(=NH)NH_2$, $C(O)(CH_2)_2NHC(O)(CH_2)_5NR^{23}C(=NH)NH_2$, and $C(O)CH(NH_2)(CH_2)_4NH_2$;


R^{22} is H or C_{1-6} -alkyl;

5 R^{23} is H or C_{1-6} -alkyl; and


r is 1, 2, 3, 4, or 5;

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl,

R^8 is selected from:



R^4 is, independently at each occurrence, selected from:




(e) performing n-1 iterations of the sequential steps of:


(e1) contacting the product formed by the immediately prior step with a deblocking agent; and


(e2) contacting the compound formed by the immediately prior step with a compound

5 of Formula (B8):



wherein:

X is selected from H, OH, $-\text{OCH}_2\text{CH}_2\text{CN}$, $\text{N}(\text{R}^{17})_2$, , and



wherein

R^{17} is, independently for each occurrence, H or C_{1-6} -alkyl;

R^{18} is selected from the group consisting of H, C_{1-6} -alkyl, $-\text{C}(=\text{NH})\text{NH}_2$,

$-\text{C}(\text{O})(\text{CH}_2)_q\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$,

and $-\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$;

q is 1, 2, 3, 4, or 5;

R^{19} is, independently for each occurrence, H or methyl;

R^{20} is H or C_{1-6} -alkyl;

R^{21} is selected from the group consisting of H, C_{1-6} -alkyl, $\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(\text{O})(\text{CH}_2)_r\text{NR}^{23}\text{C}(=\text{NH})\text{NH}_2$, $\text{C}(\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_3\text{NHC}(=\text{NH})\text{NH}_2$, $\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{23}\text{C}(=\text{NH})\text{NH}_2$, and $\text{C}(\text{O})\text{CH}(\text{NH}_2)(\text{CH}_2)_4\text{NH}_2$;


R^{22} is H or C_{1-6} -alkyl;

R^{23} is H or C_{1-6} -alkyl;


r is 1, 2, 3, 4, or 5;

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

5 R^4 is, independently for each compound of Formula (A8), selected from the group consisting of:



10 to form a compound of Formula (B9):



(B9);


wherein:

w is selected from O or C;

t is 1, 2, 3, 4, or 5;

 R^{11} is selected from the group consisting of halo, CN, and NO_2 ; R^{12} is C_{1-6} -alkyl; R^{13} is H or C_{1-6} -alkyl; R^{14} is a first linker bound to a support-medium;

n is an integer from 10 to 40,

X is, independently at each occurrence, selected from H, OH, $-\text{OCH}_2\text{CH}_2\text{CN}$,

wherein:

 R^{17} is, independently for each occurrence, H or C_{1-6} -alkyl; R^{18} is selected from the group consisting of H, C_{1-6} -alkyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(\text{O})(\text{CH}_2)_q\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$,and $-\text{C}(\text{O})(\text{CH}_2)_2\text{NHC}(\text{O})(\text{CH}_2)_5\text{NR}^{20}\text{C}(=\text{NH})\text{NH}_2$;

q is 1, 2, 3, 4, or 5;

R^{19} is, independently for each occurrence, H or methyl;

R^{20} is H or C_{1-6} -alkyl;

R^{21} is selected from the group consisting of H, C_{1-6} -alkyl, $C(=NH)NH_2$, $C(O)(CH_2)_rNR^{23}C(=NH)NH_2$, $C(O)CH(NH_2)(CH_2)_3NHC(=NH)NH_2$,
 5 $C(O)(CH_2)_2NHC(O)(CH_2)_5NR^{23}C(=NH)NH_2$, and $C(O)CH(NH_2)(CH_2)_4NH_2$;


R^{22} is H or C_{1-6} -alkyl;

R^{23} is H or C_{1-6} -alkyl; and


r is 1, 2, 3, 4, or 5;


R^3 is selected from the group consisting of trityl, monomethoxytrityl,
 10 dimethoxytrityl and trimethoxytrityl,

R^8 is selected from:




R^4 is, independently for each occurrence, selected from the group consisting
 of:





(f) contacting the compound of Formula (A9) with a deblocking agent to form a compound of Formula (B10):



(B10);

wherein:

w is selected from O or C;

t is 1, 2, 3, 4, or 5;

R¹¹ is selected from the group consisting of halo, CN, and NO₂;


R¹² is C₁₋₆-alkyl;

R¹³ is H or C₁₋₆-alkyl;

R¹⁴ is a first linker bound to a support-medium;

n is an integer from 10 to 40,

X is, independently at each occurrence, selected from H, OH, -OCH₂CH₂CN,



wherein:

R^{17} is, independently for each occurrence, H or C₁₋₆-alkyl;

R^{18} is selected from the group consisting of H, C₁₋₆-alkyl, -C(=NH)NH₂,

-C(O)(CH₂)_qNR²⁰C(=NH)NH₂,

and -C(O)(CH₂)₂NHC(O)(CH₂)₅NR²⁰C(=NH)NH₂;

q is 1, 2, 3, 4, or 5;

R^{19} is, independently for each occurrence, H or methyl;

R^{20} is H or C₁₋₆-alkyl;


R^{21} is selected from the group consisting of H, C₁₋₆-alkyl, C(=NH)NH₂, C(O)(CH₂)_rNR²³C(=NH)NH₂, C(O)CH(NH₂)(CH₂)₃NHC(=NH)NH₂, C(O)(CH₂)₂NHC(O)(CH₂)₅NR²³C(=NH)NH₂, and C(O)CH(NH₂)(CH₂)₄NH₂;

R^{22} is H or C₁₋₆-alkyl;

R^{23} is H or C₁₋₆-alkyl; and


r is 1, 2, 3, 4, or 5;

R^8 is selected from:




R^4 is, independently for each occurrence, selected from the group consisting

of:



(g) contacting the compound of Formula (B10) with a cleaving agent to form a compound of

5 Formula (B11):



(B11);

wherein:


w is selected from O or C;

t is 1, 2, 3, 4, or 5;

R¹¹ is selected from the group consisting of halo, CN, and NO₂;

5 n is an integer from 10 to 40,

X is, independently at each occurrence, selected from H, OH, -OCH₂CH₂CN,



wherein:

R¹⁷ is, independently for each occurrence, H or C₁₋₆-alkyl;

10 R¹⁸ is selected from the group consisting of H, C₁₋₆-alkyl, -C(=NH)NH₂,

-C(O)(CH₂)_qNR²⁰C(=NH)NH₂,

and -C(O)(CH₂)₂NHC(O)(CH₂)₅NR²⁰C(=NH)NH₂;

q is 1, 2, 3, 4, or 5;

15 R¹⁹ is, independently for each occurrence, H or methyl;

R²⁰ is H or C₁₋₆-alkyl;


R²¹ is selected from the group consisting of H, C₁₋₆-alkyl, C(=NH)NH₂,
C(O)(CH₂)_rNR²³C(=NH)NH₂, C(O)CH(NH₂)(CH₂)₃NHC(=NH)NH₂,
C(O)(CH₂)₂NHC(O)(CH₂)₅NR²³C(=NH)NH₂, and C(O)CH(NH₂)(CH₂)₄NH₂;

20 R²² is H or C₁₋₆-alkyl;

R²³ is H or C₁₋₆-alkyl; and


r is 1, 2, 3, 4, or 5;

R⁸ is selected from:



25 R⁴ is, independently for each occurrence, selected from the group consisting

of:



(j) contacting the compound of Formula (B11) with a deprotecting agent to form the
 5 oligomeric compound of Formula (B).

In some embodiments, step (d) or step (e2) further comprises contacting the
 compound of Formula (IV-B) or the compound formed by the immediately prior step,
 respectively, with a capping agent.

10 In some embodiments, each step is performed in the presence of at least one solvent.

In some embodiments, the deblocking agent used in each step is a solution comprising
 a halogenated acid. In some embodiments, the deblocking agent used in each step is
 cyanoacetic acid. In some embodiments, the halogenated acid is selected from the group
 consisting of chloroacetic acid, dichloroacetic acid, trichloroacetic acid, fluoroacetic acid,
 15 difluoroacetic acid, and trifluoroacetic acid. In some embodiments, the halogenated acid is
 trifluoroacetic acid.

In some embodiments, at least one of steps (a), (c), (e1), and (f) further comprise the
 step of contacting the deblocked compound of each step with a neutralization agent. In some

embodiments, each of steps (a), (c), (e1), and (f) further comprise the step of contacting the deblocked compound of each step with a neutralization agent. In some embodiments, the neutralization agent is in a solution comprising dichloromethane and isopropyl alcohol. In some embodiments, the neutralization agent is a monoalkyl, dialkyl, or trialkyl amine. In some embodiments, the neutralization agent is N,N-diisopropylethylamine.

In some embodiments, the deblocking agent used in each step is a solution comprising 4-cyanopyridine, dichloromethane, trifluoroacetic acid, trifluoroethanol, and water.

In some embodiments, the capping agent is in a solution comprising ethylmorpholine and methylpyrrolidinone. In some embodiments, the capping agent is an acid anhydride. In some embodiments, the acid anhydride is benzoic anhydride.

In some embodiments, the compounds of Formula (B4) and Formula (B8) are each, independently, in a solution comprising ethylmorpholine and dimethylimidazolidinone.


In some embodiments, the cleavage agent comprises dithiothreitol and 1,8-diazabicyclo[5.4.0]undec-7-ene. In some embodiments, the cleavage agent is in a solution comprising N-methyl-2-pyrrolidone.

In some embodiments, the deprotecting agent comprises NH_3 . In some embodiments, the deprotecting agent is in an aqueous solution.

In some embodiments, the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.


Oligomeric compounds of the disclosure may have asymmetric centers, chiral axes, and chiral planes (as described, for example, in: E. L. Eliel and S. H. Wilen, Stereo-chemistry of Carbon Compounds, John Wiley & Sons, New York, 1994, pages 1119-1190, and March, J. , Advanced Organic Chemistry, 3d. Ed., Chap. 4, John Wiley & Sons, New York (1985)), and may occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers. Oligomeric compounds of the disclosure herein specifically mentioned, without any indication of its stereo-chemistry, are intended to represent all possible isomers and mixtures thereof.

Specifically, without wishing to be bound by any particular theory, oligomeric compounds of the disclosure are prepared, as discussed herein, from activated morpholino subunits including such non-limiting examples such as a compound of Formula (VIII):




(VIII);

wherein R^2 is, independently for each compound of Formula (VIII), selected from the group consisting of:




Each of the above-mentioned compounds of Formula (VIII) may be prepared, for example, from the corresponding beta-D-ribofuranosyl as depicted below:



See Summerton et al., *Antisense Nucleic Acid Drug Dev.* 1997, 7, 187-195. Without being bound by any particular theory, the stereochemistry of the two chiral carbons is retained


under the synthetic conditions such that a number of possible stereoisomers of each morpholino subunit may be produced based on selection of, for example, an alpha-L-ribofuranosyl, alpha-D-ribofuranosyl, beta-L-ribofuranosyl, or beta-D-ribofuranosyl starting material.

- 5 For example, in some embodiments, a compound of Formula (VIII) of the disclosure may be of Formula (VIIIa):



(VIIIa);

- wherein R² is, independently for each compound of Formula (VIIIa), selected from
10 the group consisting of:







- Without being bound by any particular theory, incorporation of 10 to 40 compounds of Formula (VIII), for example, into an oligomeric compound of the disclosure may result in
15 numerous possible stereoisomers.


Without wishing to be bound by any particular theory, oligomeric compounds of the disclosure comprise one or more phosphorous-containing intersubunit linkages, which create a chiral center at each phosphorus, each of which is designated as either an “Sp” or “Rp” configuration as understood in the art. Without wishing to be bound by any particular theory, this chirality creates stereoisomers, which have identical chemical composition but different three-dimensional arrangement of their atoms.

Without wishing to be bound by any particular theory, the configuration of each phosphorous-containing intersubunit linkage occurs randomly during synthesis of, for example, oligomeric compounds of the disclosure. Without wishing to be bound by any particular theory, the synthesis process generates an exponentially large number of stereoisomers of an oligomeric compound of the disclosure because oligomeric compounds of the disclosure are comprised of numerous phosphorous-containing intersubunit linkages – with each phosphorous-containing intersubunit linkage having a random chiral configuration. Specifically, without wishing to be bound by any particular theory, each intersubunit linkage of an additional morpholino subunit doubles the number of stereoisomers of the product, so that a conventional preparation of an oligomeric compound of the disclosure is in fact a highly heterogeneous mixtures of 2^N stereoisomers, where N represents the number of phosphorous-containing intersubunit linkages.

Table 1 depicts various embodiments of morpholino subunits provided in the processes described herein.

Table 1: Various embodiments of morpholino subunits.

A = 	G = 
C = 	T = 




EXAMPLES

Examples have been set forth below for the purpose of illustration and to describe certain specific embodiments of the disclosure. However, the scope of the claims is not to be in any way limited by the examples set forth herein. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art and such changes and modifications including, without limitation, those relating to the chemical structures, substituents, derivatives, formulations or methods of the disclosure may be made without departing from the spirit of the disclosure and the scope of the appended claims. Definitions of the variables in the structures in the schemes herein are commensurate with those of corresponding positions in the formulae presented herein.

Example 1: NCP2 Anchor Synthesis

1. Preparation of Methyl 4-Fluoro-3-Nitrobenzoate (1)




To a 100L flask was charged 12.7kg of 4-fluoro-3-nitrobenzoic acid was added 40kg of methanol and 2.82kg concentrated sulfuric acid. The mixture was stirred at reflux (65° C)

for 36 hours. The reaction mixture was cooled to 0° C. Crystals formed at 38° C. The mixture was held at 0° C for 4 hrs then filtered under nitrogen. The 100L flask was washed and filter cake was washed with 10kg of methanol that had been cooled to 0° C. The solid filter cake was dried on the funnel for 1 hour, transferred to trays, and dried in a vacuum oven at room temperature to a constant weight of 13.695kg methyl 4-fluoro-3-nitrobenzoate (100% yield; HPLC 99%).


2. Preparation of 3-Nitro-4-(2-oxopropyl)benzoic Acid

A. (Z)-Methyl 4-(3-Hydroxy-1-Methoxy-1-Oxobut-2-en-2-yl)-3-Nitrobenzoate (2)




To a 100L flask was charged 3.98kg of methyl 4-fluoro-3-nitrobenzoate (1) from the previous step 9.8kg DMF, 2.81kg methyl acetoacetate. The mixture was stirred and cooled to 0° C. To this was added 3.66kg DBU over about 4 hours while the temperature was maintained at or below 5° C. The mixture was stirred an additional 1 hour. To the reaction flask was added a solution of 8.15kg of citric acid in 37.5kg of purified water while the reaction temperature was maintained at or below 15° C. After the addition, the reaction mixture was stirred an addition 30 minutes then filtered under nitrogen. The wet filter cake was returned to the 100L flask along with 14.8kg of purified water. The slurry was stirred for 10 minutes then filtered. The wet cake was again returned to the 100L flask, slurried with 14.8kg of purified water for 10 minutes, and filtered to crude (Z)-methyl 4-(3-hydroxy-1-methoxy-1-oxobut-2-en-2-yl)-3-nitrobenzoate.

B. 3-Nitro-4-(2-oxopropyl)benzoic Acid



The crude (Z)-methyl 4-(3-hydroxy-1-methoxy-1-oxobut-2-en-2-yl)-3-nitrobenzoate was charged to a 100L reaction flask under nitrogen. To this was added 14.2kg 1,4-dioxane and the stirred. To the mixture was added a solution of 16.655kg concentrated HCl and 13.33kg purified water (6M HCl) over 2 hours while the temperature of the reaction mixture was maintained below 15° C. When the addition was complete, the reaction mixture was heated at reflux (80° C) for 24 hours, cooled to room temperature, and filtered under nitrogen. The solid filter cake was triturated with 14.8kg of purified water, filtered, triturated again with 14.8kg of purified water, and filtered. The solid was returned to the 100L flask with 39.9kg of DCM and refluxed with stirring for 1 hour. 1.5kg of purified water was added to dissolve the remaining solids. The bottom organic layer was split to a pre-warmed 72L flask, then returned to a clean dry 100L flask. The solution was cooled to 0° C, held for 1 hour, then filtered. The solid filter cake was washed twice each with a solution of 9.8kg DCM and 5kg heptane, then dried on the funnel. The solid was transferred to trays and dried to a constant weight of 1.855kg 3-Nitro-4-(2-oxopropyl)benzoic Acid. Overall yield 42% from compound **1**. HPLC 99.45%.

3. Preparation of N-Tritylpiperazine Succinate (NTP)




To a 72L jacketed flask was charged under nitrogen 1.805kg triphenylmethyl chloride and 8.3kg of toluene (TPC solution). The mixture was stirred until the solids dissolved. To a 100L jacketed reaction flask was added under nitrogen 5.61kg piperazine, 19.9kg toluene, and 3.72kg methanol. The mixture was stirred and cooled to 0° C. To this was slowly added in portions the TPC solution over 4 hours while the reaction temperature was maintained at or

below 10° C. The mixture was stirred for 1.5 hours at 10° C, then allowed to warm to 14° C. 32.6kg of purified water was charged to the 72L flask, then transferred to the 100L flask while the internal batch temperature was maintained at 20+/-5° C. The layers were allowed to split and the bottom aqueous layer was separated and stored. The organic layer was
 5 extracted three times with 32kg of purified water each, and the aqueous layers were separated and combined with the stored aqueous solution.

The remaining organic layer was cooled to 18° C and a solution of 847g of succinic acid in 10.87kg of purified water was added slowly in portions to the organic layer. The mixture was stirred for 1.75 hours at 20+/-5° C. The mixture was filtered, and the solids
 10 were washed with 2kg TBME and 2kg of acetone then dried on the funnel. The filter cake was triturated twice with 5.7kg each of acetone and filtered and washed with 1kg of acetone between triturations. The solid was dried on the funnel, then transferred to trays and dried in a vacuum oven at room temperature to a constant weight of 2.32kg of NTP. Yield 80%.

15 4. Preparation of (4-(2-Hydroxypropyl)-3-Nitrophenyl)(4-Tritylpiperazin-1-yl)Methanone


A. Preparation of 1-(2-Nitro-4(4-Tritylpiperazine-1-Carbonyl)Phenyl)Propan-2-one



To a 100L jacketed flask was charged under nitrogen 2kg of 3-Nitro-4-(2-oxopropyl)benzoic Acid (**3**), 18.3 kg DCM, 1.845kg N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDC.HCl). The solution was stirred until a homogenous
 20 mixture was formed. 3.048kg of NTP was added over 30 minutes at room temperature and stirred for 8 hours. 5.44kg of purified water was added to the reaction mixture and stirred for 30 minutes. The layers were allowed to separate and the bottom organic layer containing the product was drained and stored. The aqueous layer was extracted twice with 5.65kg of DCM.
 25 The combined organic layers were washed with a solution of 1.08kg sodium chloride in 4.08kg purified water. The organic layers were dried over 1.068kg of sodium sulfate and filtered. The sodium sulfate was washed with 1.3kg of DCM. The combined organic layers


were slurried with 252g of silica gel and filtered through a filter funnel containing a bed of 252g of silica gel. The silica gel bed was washed with 2kg of DCM. The combined organic layers were evaporated on a rotovap. 4.8kg of THF was added to the residue and then evaporated on the rotovap until 2.5 volumes of the crude 1-(2-nitro-4(4-tritylpiperazine-1-carbonyl)phenyl)propan-2-one in THF was reached.

B. Preparation of (4-(2-Hydroxypropyl)-3-Nitrophenyl)(4-Tritylpiperazin-1-yl)Methanone (**5**)



To a 100L jacketed flask was charged under nitrogen 3600g of **4** from the previous step and 9800g THF. The stirred solution was cooled to $\leq 5^{\circ}\text{C}$. The solution was diluted with 11525g ethanol and 194g of sodium borohydride was added over about 2 hours at $\leq 5^{\circ}\text{C}$. The reaction mixture was stirred an additional 2 hours at $\leq 5^{\circ}\text{C}$. The reaction was quenched with a solution of about 1.1kg ammonium chloride in about 3kg of water by slow addition to maintain the temperature at $\leq 10^{\circ}\text{C}$. The reaction mixture was stirred an additional 30 minutes, filtered to remove inorganics, and recharged to a 100L jacketed flask and extracted with 23kg of DCM. The organic layer was separated and the aqueous was twice more extracted with 4.7kg of DCM each. The combined organic layers were washed with a solution of about 800g of sodium chloride in about 3kg of water, then dried over 2.7kg of sodium sulfate. The suspension was filtered and the filter cake was washed with 2kg of DCM. The combined filtrates were concentrated to 2.0 volumes, diluted with about 360g of ethyl acetate, and evaporated. The crude product was loaded onto a silica gel column of 4kg of silica packed with DCM under nitrogen and eluted with 2.3kg ethyl acetate in 7.2kg of DCM. The combined fractions were evaporated and the residue was taken up in 11.7kg of toluene. The toluene solution was filtered and the filter cake was washed twice with 2kg of toluene each. The filter cake was dried to a constant weight of 2.275kg of compound **5** (46% yield from compound **3**) HPLC 96.99%.

5. Preparation of 2,5-Dioxopyrrolidin-1-yl(1-(2-Nitro-4-(4-triphenylmethylpiperazine-1 Carbonyl)Phenyl)Propan-2-yl) Carbonate (**NCP2 Anchor**)



To a 100L jacketed flask was charged under nitrogen 4.3kg of compound **5** (weight adjusted based on residual toluene by H^1 NMR; all reagents here after were scaled accordingly) and 12.7kg pyridine. To this was charged 3.160 kg of DSC (78.91 weight % by H^1 NMR) while the internal temperature was maintained at $\leq 35^\circ C$. The reaction mixture was aged for about 22 hours at ambience then filtered. The filter cake was washed with 200g of pyridine. In two batches each comprising $\frac{1}{2}$ the filtrate volume, filtrate wash charged slowly to a 100L jacketed flask containing a solution of about 11kg of citric acid in about 50 kg of water and stirred for 30 minutes to allow for solid precipitation. The solid was collected with a filter funnel, washed twice with 4.3kg of water per wash, and dried on the filter funnel under vacuum.

The combined solids were charged to a 100L jacketed flask and dissolved in 28kg of DCM and washed with a solution of 900g of potassium carbonate in 4.3kg of water. After 1 hour, the layers were allowed to separate and the aqueous layer was removed. The organic layer was washed with 10kg of water, separated, and dried over 3.5kg of sodium sulfate. The DCM was filtered, evaporated, and dried under vacuum to 6.16kg of **NCP2 Anchor** (114% yield).


Example 2: Anchor Loaded Resin Synthesis

To a 75L solid phase synthesis reactor with a Teflon stop cock was charged about 52L of NMP and 2300g of aminomethyl polystyrene resin. The resin was stirred in the NMP to swell for about 2 hours then drained. The resin was washed twice with about 4L DCM per wash, then twice with 39L Neutralization Solution per wash, then twice with 39L of DCM per wash. The NCP2 Anchor Solution was slowly added to the stirring resin solution, stirred

for 24 hours at room temperature, and drained. The resin was washed four times with 39L of NMP per wash, and six times with 39L of DCM per wash. The resin was treated and stirred with $\frac{1}{2}$ the DEDC Capping Solution for 30 minutes, drained, and was treated and stirred with the 2nd $\frac{1}{2}$ of the DEDC Capping Solution for 30 minutes and drained. The resin was washed six times with 39L of DCM per wash then dried in an oven to constant weight of 3573.71g of Anchor Loaded Resin.


Example 3: Preparation of Activated EG3 Tail (See Fig. 2)

1. Preparation of Trityl Piperazine Phenyl Carbamate 35




To a cooled suspension of NTP in dichloromethane (6 mL/g NTP) was added a solution of potassium carbonate (3.2 eq) in water (4 mL/g potassium carbonate). To this two-phase mixture was slowly added a solution of phenyl chloroformate (1.03 eq) in dichloromethane (2 g/g phenyl chloroformate). The reaction mixture was warmed to 20° C. Upon reaction completion (1-2 hr), the layers were separated. The organic layer was washed with water, and dried over anhydrous potassium carbonate. The product 35 was isolated by crystallization from acetonitrile. Yield=80%.

2. Preparation of Carbamate Alcohol 36



Sodium hydride (1.2 eq) was suspended in 1-methyl-2-pyrrolidinone (32 mL/g sodium hydride). To this suspension were added triethylene glycol (10.0 eq) and compound 35 (1.0 eq). The resulting slurry was heated to 95° C. Upon reaction completion (1-2 hr), the mixture was cooled to 20° C. To this mixture was added 30% dichloromethane/methyl tert-butyl ether (v:v) and water. The product-containing organic layer was washed successively with aqueous NaOH, aqueous succinic acid, and saturated aqueous sodium chloride. The product 36 was isolated by crystallization from dichloromethane/methyl tert-butyl ether/heptane. Yield=90%.


3. Preparation of EG3 Tail Acid 37



To a solution of compound 36 in tetrahydrofuran (7 mL/g 36) was added succinic anhydride (2.0 eq) and DMAP (0.5 eq). The mixture was heated to 50° C. Upon reaction completion (5 hr), the mixture was cooled to 20° C and adjusted to pH 8.5 with aqueous NaHCO₃. Methyl tert-butyl ether was added, and the product was extracted into the aqueous layer. Dichloromethane was added, and the mixture was adjusted to pH 3 with aqueous citric acid. The product-containing organic layer was washed with a mixture of pH=3 citrate buffer

and saturated aqueous sodium chloride. This dichloromethane solution of 37 was used without isolation in the preparation of compound 38.

4. Preparation of Activated EG3 Tail 38



5

To the solution of compound 37 was added N-hydroxy-5-norbornene-2,3-dicarboxylic acid imide (HONB) (1.02 eq), 4-dimethylaminopyridine (DMAP) (0.34 eq), and then 1-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDC) (1.1 eq). The mixture was heated to 55° C. Upon reaction completion (4-5 hr), the mixture was cooled to 20° C and washed successively with 1:1 0.2 M citric acid/brine and brine. The dichloromethane solution underwent solvent exchange to acetone and then to N,N-dimethylformamide, and the product was isolated by precipitation from acetone/N,N-dimethylformamide into saturated aqueous sodium chloride. The crude product was reslurried several times in water to remove residual N,N-dimethylformamide and salts. Yield=70% of Activated EG3 Tail 38 from compound 36.

15

Example 4: 50L Solid-phase Synthesis of

Eteplirsen [Oligomeric Compound (XII)] Crude Drug Substance

1. Materials


Table 2: Starting Materials

Material Name	Chemical Name	CAS Number	Chemical Formula	MW
Activated A Subunit	Phosphoramidochloridic acid, <i>N,N</i> -dimethyl-, [6-[6-(benzoylamino)-9H-purin-9-yl]-4-(triphenylmethyl)-2-morpholinyl]methyl ester	1155373-30-0	C ₃₈ H ₃₇ ClN ₇ O ₄ P	722.2
Activated C Subunit	Phosphoramidochloridic acid, <i>N,N</i> -dimethyl-, [6-[4-(benzoylamino)-2-oxo-1(2H)-pyrimidinyl]-4-(triphenylmethyl)-2-morpholinyl]methyl ester	1155373-31-1	C ₃₇ H ₃₇ ClN ₅ O ₅ P	698.2

Activated DPG Subunit	Propanoic Acid, 2,2-dimethyl-4-[[[9-[6-[[[chloro(dimethylamino)phosphinyl]oxy]methyl]-4-(triphenylmethyl)-2-morpholinyl]-2-[(2-phenylacetyl)amino]-9H-purin-6-yl]oxy]methyl]phenyl ester	1155309-89-9	C ₅₁ H ₅₃ ClN ₇ O ₇ P	942.2
Activated T Subunit	Phosphoramidochloridic acid, <i>N,N</i> -dimethyl-, [6-(3,4-dihydro-5-methyl-2,4-dioxo-1(2H)-pyrimidinyl)]-4-(triphenylmethyl)-2-morpholinyl]methyl ester	1155373-34-4	C ₃₁ H ₃₄ ClN ₄ O ₅ P	609.1
Activated EG3 Tail	Butanedioic acid, 1-[3a <i>R</i> ,4 <i>S</i> ,7 <i>R</i> ,7a <i>S</i>)-1,3,3a,4,7,7a-hexahydro-1,3-dioxo-4,7-methano-2H-isoindol-2-yl] 4-[2-[2-[2-[[[4-(triphenylmethyl)-1-piperazinyl]carbonyl]oxy]ethoxy]ethoxy]ethyl] ester	1380600-06-5	C ₄₃ H ₄₇ N ₃ O ₁₀	765.9

Chemical Structures of Starting Materials:


A. Activated EG3 Tail



5


Compound (B)

B. Activated C Subunit (for preparation, see U.S. Patent No. 8,067,571)




Compound (C1)

C. Activated A Subunit (for preparation, see U.S. Patent No. 8,067,571)




Compound (D1)

D. Activated DPG Subunit (for preparation, see WO 2009/064471)




Compound (E1)

E. Activated T Subunit (for preparation, see WO 2013/082551)



Compound (F1)

10 F. Anchor Loaded Resin



Formula (I)

wherein

R¹ is a support-medium.

5

Table 3: Description of Solutions for Solid Phase Oligomer Synthesis of Eteplirsen Crude Drug Substance

Solution Name	Solution Composition
NCP2 Anchor Solution	37.5L NMP and 1292g NCP2 Anchor
DEDC Capping Solution	4.16L Diethyl Dicarbonate (DEDC), 3.64L NEM, and 33.8L DCM
CYTFA Solution	2.02 kg 4-cyanopyridine, 158 L DCM, 1.42 L TFA, 39 L TFE, and 2 L purified water
Neutralization Solution	35.3 L IPA, 7.5 L DIPEA, and 106.5 L DCM
Cleavage Solution	1,530.04 g DTT, 6.96 L NMP, and 2.98 L DBU

2. Synthesis of Eteplirsen Crude Drug Substance

10

A. Resin swelling

750 g of Anchor Loaded Resin and 10.5 L of NMP were charged to a 50 L silanized reactor and stirred for 3 hours. The NMP was drained and the Anchor Loaded Resin was washed twice with 5.5L each of DCM and twice with 5.5 L each of 30% TFE/DCM.

B. Cycle 0: EG3 Tail Coupling

15

The Anchor Loaded Resin was washed three times with 5.5 L each of 30% TFE/DCM and drained, washed with 5.5 L of CYTFA solution for 15 minutes and drained, and again washed with 5.5 L of CYTFA Solution for 15 minutes without draining to which 122 mL of 1:1 NEM/DCM was charged and the suspension stirred for 2 minutes and drained. The resin was washed twice with 5.5 L of Neutralization Solution for 5 minutes and drained, then twice

with 5.5 L each of DCM and drained. A solution of 706.2 g of activated EG3 Tail (MW 765.85) and 234 mL of NEM in 3 L of DMI was charged to the resin and stirred for 3 hours at RT and drained. The resin was washed twice with 5.5 L each of Neutralization Solution for 5 minutes per each wash, and once with 5.5 L of DCM and drained. A solution of 374.8 g of benzoic anhydride and 195 mL NEM in 2680 mL NMP was charged and stirred for 15 minutes and drained. The resin was stirred with 5.5 L of Neutralization Solution for 5 minutes, then washed once with 5.5 L of DCM and twice with 5.5 L each of 30% TFE/DCM. The resin was suspended in 5.5 L of 30% TFE/DCM and held for 14 hours.

C. Subunit Coupling Cycles 1-30

i. Pre-coupling treatments

Prior to each coupling cycle as described in Table 4, the resin was: 1) washed with 30% TFE/DCM; 2) a) treated with CYTFA Solution 15 minutes and drained, and b) treated with CYTFA solution for 15 minutes to which was added 1:1 NEM/DCM, stirred, and drained; 3) stirred three times with Neutralization Solution; and 4) washed twice with DCM. See Table 4.

ii. Post Coupling Treatments

After each subunit solution was drained as described in Table 4, the resin was: 1) washed with DCM; and 2) washed two times with 30% TFE/DCM. If the resin was held for a time period prior to the next coupling cycle, the second TFE/DCM wash was not drained and the resin was retained in said TFE/DCM wash solution. See Table 4.

iii. Activated Subunit Coupling Cycles

The coupling cycles were performed as described in Table 4.

iv. Final IPA Washing

After the final coupling step was performed as described in Table 4, the resin was washed 8 times with 19.5 L each of IPA, and dried under vacuum at room temperature for about 63.5 hours to a dried weight of 5,579.8 g.

D. Cleavage

The above resin bound Eteplirsén Crude Drug Substance was divided into two lots, each lot was treated as follows. A 2,789.9g lot of resin was: 1) stirred with 10L of NMP for 2hrs, then the NMP was drained; 2) washed three times with 10L each of 30% TFE/DCM; 3) treated with 10L CYTFA Solution for 15 minutes; and 4) 10L of CYTFA Solution for 15 minutes to which 130ml 1:1 NEM/DCM was then added and stirred for 2 minutes and drained. The resin was treated three times with 10L each of Neutralization Solution, washed six times with 10L of DCM, and eight times with 10L each of NMP. The resin was treated

with a Cleaving Solution of 1530.4g DTT and 2980 DBU in 6.96L NMP for 2 hours to detach the Eteplisen Crude Drug Substance from the resin. The Cleaving solution was drained and retained in a separate vessel. The reactor and resin were washed with 4.97L of NMP which was combined with the Cleaving Solution.

5

Table 4:

Cycle No.: Subunit (SU)	Pre-coupling Treatment				Coupling Cycle		Post-Coupling Treatment	
	1	2	3	4			1	2
	30% TFE/DCM Wash	CYTFA Solution ¹	Neutralization Solution	DCM Wash	Quantity SU (g) NEM (L) DMI (L)	RT Coupling Time (Hrs.)	DCM Wash	30% TFE/DCM Wash
1:C	5.5L	a) 5.5L b) 5.5L, 122ml	3x5.5L	5.5L	536.7g; 195 ml NEM; 3.2L DMI	5	5.5L	2x5.5L
2:T	7.0L	a) 7L b) 7L, 158ml	3x7L	2x7L	468.2g and 195ml NEM 3.2L DMI	4.25	7L	2x7L ²
3:C	8L	a) 8L b) 8L, 182ml	3x8L	2x8L	536.7g; 195ml NEM; 3.4L DMI	4.25	8L	2x8L
4:C	9L	a) 9L b) 9L, 206ml	3x9L	2x9L	536.7g; 195ml NEM; 3.6L DMI	4.25	9L	2x9L ³
5:A	9.5L	a) 9.5L b) 9.5L, 220ml	3x9.5L	2x9.5L	555.2g; 195ml NEM; 3.4L DMI	4.25	9.5L	2x9.5L
6:A	10L	a) 10L b) 10L, 232ml	3x10L	2x10L	555.2g; 195ml NEM; 3.45L DMI	4.25	10L	2x10L ⁴

¹ ml indicates the amount of 1:1 NEM/DCM

² Resin held at this step for ½ day

³ Resin held at this step for ½ day

⁴ Resin held at this stage for 0.4 days

Table 4 Cont.

Cycle No.: Subunit (SU)	Pre-coupling Treatment				Coupling Cycle		Post-Coupling Treatment	
	1	2	3	4			1	2
	30% TFE/DCM Wash	CYTFA Solution	Neutralization Solution	DCM Wash	Quantity SU (g) NEM (L) DMI (L)	RT Coupling Time (Hrs.)	DCM Wash	30% TFE/DCM Wash
7:C	11L	a) 11L b) 11L, 256ml	3x11L	2x11L	536.7g; 195ml NEM; 3.57L DMI	4.25	11L	2x11L
8:A	11L	a) 11L b) 11L, 256ml	3x11L	2x11L	555.2g; 195ml NEM; 3.64L DMI	4.25	11L	2x11L ⁵
9:T	11.5L	a) 11.5L b) 11.5L 268ml	3x11.5L	2x 11.5L	468.2g; 195ml NEM; 3.72L DMI	4.25	11.5L	2x11.5L
10:C	12L	a) 12L b) 12L, 280ml	3x12L	2x12L	536.7g; 195ml NEM; 3.96L DMI	4.25	12L	2x12L ⁶
11:A	13.5L	a) 13.5L b) 13.5L, 204ml	3x13.5L	2x 13.5L	721.7g; 253ml NEM; 4.02L DMI	4.25	13.5L	2x13.5L
12:A	13.5L	a) 13.5L b) 13.5L, 204ml	3x13.5L	2x 13.5L	721.7g; 253ml NEM; 4.02L DMI	4.25	13.5L	2x13.5L ⁷

⁵ Resin held at this stage for 2.5 days⁶ Resin held at this stage for ½ day⁷ Resin held at this stage for 0.4 days

Table 4 Cont.

Cycle No.: Subunit (SU)	Pre-coupling Treatment				Coupling Cycle		Post-Coupling Treatment	
	1	2	3	4			1	2
	30% TFE/DCM Wash	CYTFA Solution	Neutralization Solution	DCM Wash	Quantity SU (g) NEM (L) DMI (L)	RT Coupling Time (Hrs.)	DCM Wash	30% TFE/DCM Wash
13:DP G	14L	a) 14L b) 14L, 216ml	3x14L	2x14L	941.9g; 253ml NEM; 4.02L DMI	4.25	14L	2x14L
14:DP G	14.5L	a) 14.5L b) 14.5L, 228ml	3x14.5L	2x 14.5L	941.9g; 253ml NEM; 4.1L DMI	4.25	14.5L	2x14.5L ⁸
15:A	15.5L	a) 15.5L b) 15.5L, 254ml	3x15.5L	2x 15.5L	721.7g; 253ml NEM; 4.26L DMI	4.25	15.5L	2x15.5L
16:A	15.5L	a) 15.5L b) 15.5L, 254ml	3x15.5L	2x 15.5L	721.7g; 253ml NEM; 4.26L DMI	4.25	15.5L	2x15.5L ⁹
17:DP G	16L	a) 16L b) 16L, 366ml	3x16L	2x16L	941.9g; 253ml NEM; 4.4L DMI	4.75	16L	2x16L
18:A	16.5L	a) 16.5L b) 16.5L, 378ml	3x16.5L	2x 16.5L	721.7g; 253ml NEM; 4.4L DMI	4.25	16.5L	2x16.5L ¹⁰

⁸ Resin held at this stage for 0.4 days⁹ Resin held at this stage for 0.4 days¹⁰ Resin held at this stage for 1.5 days

Table 4 Cont.

Cycle No.: Subunit (SU)	Pre-coupling Treatment				Coupling Cycle		Post-Coupling Treatment	
	1	2	3	4			1	2
	30% TFE/DCM Wash	CYTF A Solution	Neutralization Solution	DCM Wash	Quantity SU (g) NEM (L) DMI (L)	RT Coupling Time (Hrs.)	DCM Wash	30% TFE/DCM Wash
19:T	16.5L	a) 16.5L b) 16.5L, 378ml	3x16.5L	2x 16.5L	608.7g; 253ml NEM; 4.57L DMI	4.25	16.5L	2x16.5L
20:DPG	17L	a) 17L b) 17L, 390ml	3x17L	2x17L	941.9g; 253ml NEM; 4.57L DMI	4.75	17L	2x17L ¹¹
21:DPG	17L	a) 17L b) 17L, 390ml	3x17L	2x17L	1159.2g; 311ml NEM; 4.72L DMI	4.25	17L	2x17L
22:C	17.5L	a) 17.5L b) 17.5L, 402ml	3x17.5L	2x 17.5L	858.7g; 311ml NEM; 4.72L DMI	4.75	17.5L	2x17.5L ¹²
23:A	17.5L	a) 17.5L b) 17.5L, 402ml	3x17.5L	2x 17.5L	888.3g; 311ml NEM; 4.88L DMI	4.25	17.5L	2x17.5L
24:T	18L	a) 18L b) 18L, 414ml	3x18L	2x18L	749.1g; 311ml NEM; 4.95L DMI	4.25	18L	2x18L ¹³

¹¹ Resin held at this stage for 0.3 days¹² Resin held at this stage for 0.4 days¹³ Resin held at this stage for 0.4 days

Table 4 Cont.

Cycle No.: Subunit (SU)	Pre-coupling Treatment				Coupling Cycle		Post-Coupling Treatment	
	1	2	3	4			1	2
	30% TFE/DCM Wash	CYTFA Solution	Neutralization Solution	DCM Wash	Quantity SU (g) NEM (L) DMI (L)	RT Coupling Time (Hrs.)	DCM Wash	30% TFE/DCM Wash
25:T	18L	a) 18L b) 18L, 414ml	3x18L	2x18L	749.1g; 311ml NEM; 5.1L DMI	4.25	18L	2x18L
26:T	18.5L	a) 18.5L b) 18.5L, 426ml	3x18.5L	2x 18.5L	749.1g; 311ml NEM; 5.1L DMI	4.25	18.5 L	2x18.5L ¹⁴
27:C	18.5L	a) 18.5L b) 18.5L, 426ml	3x18.5L	2x 18.5L	858.7g; 311ml NEM; 5.25L DMI	4.25	18.5 L	2x18.5L
28:T	19L	a) 19L b) 19L, 438ml	3x19L	2x19 L	749.1g; 311ml NEM; 5.25L DMI	4.25	19L	2x19L ¹⁵
29:A	19L	a) 19L b) 19L, 438ml	3x19L	2x19 L	888.3g; 311ml NEM; 5.41L DMI	4.25	19L	2x19L
30:DP G	19.5L	a) 19.5L b) 19.5L, 450ml	3x19.5L	2x 19.5L	1159.2g; 311ml NEM; 5.44L DMI	4.75	19.5 L	2x19.5L

E. Deprotection

5 The combined Cleaving Solution and NMP wash were transferred to a pressure vessel to which was added 39.8L of NH_4OH ($\text{NH}_3 \cdot \text{H}_2\text{O}$) that had been chilled to a temperature of -10° to -25° C in a freezer. The pressure vessel was sealed and heated to 45° C for 16hrs then allowed to cool to 25° C. This deprotection solution containing the Eteplirsén crude drug

¹⁴ Resin held at this stage for 0.4 days¹⁵ Resin held at this stage for 0.3 days

substance was diluted 3:1 with purified water and pH adjusted to 3.0 with 2M phosphoric acid, then to pH 8.03 with NH₄OH. HPLC (C18) 73-74% (**Figure 1**).

Table 5. Data of Figure 1

Peak #	Compound Name	Retention Time (min)	Rel. Ret. Time. Product	Area {mAu*min}	Rel.Area %	Plates (USP)
1		2.488	0.381	0.821928	0.18	1105
2		3.047	0.467	17.661449	3.91	4047
3		3.324	0.509	0.818258	0.18	n.a.
4		3.605	0.552	0.465598	0.10	7
5		4.213	0.645	6.558899	1.45	301
6		4.504	0.690	3.324238	0.74	191690
7		5.160	0.790	5.644073	1.25	651
8	AVI-4658	6.531	1.000	332.238891	73.47	2313
9		7.269	1.113	2.063159	0.46	n.a.
10		7.643	1.170	5.556411	1.23	2734
11		8.139	1.246	8.711530	1.93	3572
12		8.382	1.283	4.654783	1.03	1835
13		8.678	1.329	0.562426	0.12	n.a.
14		9.009	1.379	12.031923	2.66	6078
15		9.500	1.455	0.385563	0.09	n.a.
16		9.626	1.474	1.171507	0.26	46084
17		9.898	1.516	0.484362	0.11	21328
18		10.598	1.623	14.589918	3.23	n.a.
19		10.680	1.635	7.520577	1.66	918
20		10.811	1.656	8.604558	1.90	296
21		11.045	1.691	18.351689	4.06	49919

5

Example 5: Purification of Eteplirsén Crude Drug Substance

The deprotection solution from Example 2, part D, containing the Eteplirsén crude drug substance was loaded onto a column of ToyoPearl Super-Q 650S anion exchange resin (Tosoh Bioscience) and eluted with a gradient of 0-35% B over 17 column volume (Buffer A: 10 mM sodium hydroxide; Buffer B: 1 M sodium chloride in 10 mM sodium hydroxide) and fractions of acceptable purity (C18 and SCX HPLC) were pooled to a purified drug product solution. HPLC (**Figure 2**): 97.74% (C18) 94.58% (SCX).

The purified drug substance solution was desalted and lyophilized to 1959g purified Eteplirsén drug substance. Yield 61.4%; HPLC (**Figure 3**): 97.7% (C18) 94.6% (SCX).

15

Table 6. Data of Figure 2


Peak #	Compound Name	Retention Time (min)	Rel. Ret. Time. (Product)	Area {mAu*min}	Area Percent	Plates (USP)
1		6.837	0.750	0.050757	0.058	41574
2		7.405	0.813	0.303271	0.344	841
3		8.086	0.887	1.130007	1.280	13
4		8.615	0.946	2.265128	2.567	761
5	AVI-4658	9.111	1.000	83.468700	94.583	4405
6		10.019	1.100	0.704599	0.798	n.a.
7		11.069	1.215	0.326550	0.370	3044

Table 7. Data of Figure 3


Peak #	Compound Name	Retention Time. (min)	Rel. Ret. Time. (Product)	Area {mAu*min}	Area Percent	Plates (USP)
1		6.866	0.751	0.044399	0.063	608
2		7.794	0.852	0.280589	0.397	n.a.
3		8.188	0.895	0.816793	1.156	209
4		8.644	0.945	1.842896	2.608	1147
5	AVI-4658	9.145	1.000	66.857088	94.622	4664
6		10.058	1.100	0.575793	0.815	n.a.
7		11.103	1.214	0.239454	0.339	4375


5 **Example 6: Additional Oligomers Prepared by the Process of the Disclosure**

The following oligomers of the following sequences were prepared according to the methods of Examples 4 and 5 about:





10 where R² is, at each occurrence, independently selected from the group consisting of:



**Table 8.** Oligomer R² at each position from 1 to n and 5' to 3'

Name	n	R ² 5' to 3' and 1 to n	T
PMO1	22	CAA TGC CAT CCT GGA GTT CCT G	EG3
PMO2	25	GTT GCC TCC GGT TCT GAA GGT GTT C	EG3
AVI-7100	20	CGGT+TAGAAGAC+TCATC+TTT	EG3
AVI-7537	19	GCC+ATGGT+TTT+TTC+TC+AGG	EG3
AVI-7288	23	GAATATTAAC+AI+AC+TGAC+A+AGTC	EG3



In the above Table 8, “+” indicates wherein X is , otherwise X is -N(CH₃)₂.

5 Table 9. Acronyms

Acronym	Name
DBU	1,8-Diazabicycloundec-7-ene
DCM	Dichloromethane
DIPEA	N,N-Diisopropylethylamine
DMI	1,3-Dimethyl-2-imidazolidinone
DTT	Dithiothreitol
IPA	Isopropyl alcohol
MW	Molecular weight
NEM	N-Ethylmorpholine
NMP	N-Methyl-2-pyrrolidone
RT	Room temperature
TFA	2,2,2-Trifluoroacetic acid
TFE	2,2,2-Trifluoroethanol

Incorporation by Reference


The contents of all references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated herein in their entireties. Unless otherwise
5 defined, all technical and scientific terms used herein are accorded the meaning commonly known to one with ordinary skill in the art.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than
10 routine experimentation, many equivalents of the specific embodiments of the disclosure described herein. Such equivalents are intended to be encompassed by the following claims.

CLAIMS


1. A process for preparing an oligomeric compound of Formula (A):



5


(A);

wherein n is an integer from 10 to 40, and each R^2 is, independently for each occurrence, selected from the group consisting of:



- 10 wherein the process comprises the sequential steps of:

(a) contacting a compound of Formula (A1):




(A1);

wherein R^1 is a support-medium and R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

15


with a deblocking agent to form the compound of Formula (II):



(II);

wherein R^1 is a support-medium;


(b) contacting the compound of Formula (II) with a compound of Formula (A2):



(A2);

wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


to form a compound of Formula (A3):



(A3);

wherein R^1 is a support-medium, and R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;


(c) contacting the compound of Formula (A3) with a deblocking agent to form a compound of Formula (IV):



(IV);


wherein R¹ is a support-medium;

(d) contacting the compound of Formula (IV) with a compound of Formula (A4):




(A4);

wherein R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is selected from the group consisting of:




(PC),

(DPG),



to form a compound of Formula (A5):




5

(A5);

wherein R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and


R^4 is selected from:



5 (e) performing n-1 iterations of the sequential steps of:


(e1) contacting the product formed by the immediately prior step with a deblocking agent; and


(e2) contacting the compound formed by the immediately prior step with a compound of Formula (A8):



(A8);


wherein R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each compound of Formula (A8), selected from the group consisting of:







(A9);

wherein n is an integer from 10 to 40, R^1 is a support-medium, R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R^4 is, independently for each occurrence, selected from the group consisting of:






(f) contacting the compound of Formula (A9) with a deblocking agent to form a compound of Formula (A10):



5


(A10);

wherein n is an integer from 10 to 40, R¹ is a support-medium, and R⁴ is, independently for each occurrence, selected from the group consisting of:




(g) contacting the compound of Formula (A10) with a cleaving agent to form a compound of


5 Formula (A11):



(A11);

wherein n is an integer from 10 to 40, and R^4 is, independently for each occurrence, selected from the group consisting of:





(h) contacting the compound of Formula (A11) with a deprotecting agent to form the oligomeric compound of Formula (A).

5

2. The process of claim 1, wherein step (d) or step (e2) further comprises contacting the compound of Formula (IV) or the compound formed by the immediately prior step, respectively, with a capping agent.

10

3. The process of any one of claims 1-3, wherein the deblocking agent used in each step is a solution comprising a halogenated acid.

4. The process of any one of claims 1-3, wherein the deblocking agent used in each step is cyanoacetic acid.

15

5. The process of claim 3, wherein the halogenated acid is selected from the group consisting of chloroacetic acid, dichloroacetic acid, trichloroacetic acid, fluoroacetic acid, difluoroacetic acid, and trifluoroacetic acid.

20

6. The process of claim 5, wherein the halogenated acid is trifluoroacetic acid.

7. The process of any one of claims 1 to 6, wherein at least one of steps (a), (c), (e1), and (f) further comprise the step of contacting the deblocked compound of each step with a neutralization agent.

25

8. The process of any one of claims 1 to 7, wherein each of steps (a), (c), (e1), and (f) further comprise the step of contacting the deblocked compound of each step with a neutralization agent.
- 5 9. The process of claim 7 or 8, wherein the neutralization agent is in a solution comprising dichloromethane and isopropyl alcohol.
10. The process of any one of claims 7 to 9, wherein the neutralization agent is a monoalkyl, dialkyl, or trialkyl amine.
- 10 11. The process of claim 10, wherein the neutralization agent is N,N-diisopropylethylamine.
12. The process of any one of claims 1-11, wherein the deblocking agent used in each
15 step is a solution comprising 4-cyanopyridine, dichloromethane, trifluoroacetic acid, trifluoroethanol, and water.
13. The process of any one of claims 1-12, wherein the capping agent is in a solution comprising ethylmorpholine and methylpyrrolidinone.
- 20 14. The process of claim 13, wherein the capping agent is an acid anhydride.
15. The process of claim 14, wherein the acid anhydride is benzoic anhydride.
- 25 16. The process of any one of claims 1-15, wherein the compounds of Formula (A4) and Formula (A8) are each, independently, in a solution comprising ethylmorpholine and dimethylimidazolidinone.
17. The process of any one of claims 1-16, wherein the cleavage agent comprises
30 dithiothreitol and 1,8-diazabicyclo[5.4.0]undec-7-ene.
18. The process of claim 17, wherein the cleavage agent is in a solution comprising N-methyl-2-pyrrolidone.

19. The process of any one of claims 1-18, wherein the deprotecting agent comprises NH_3 .


20. The process of claim 19, wherein the deprotecting agent is in an aqueous solution.

5

21. The process of any one of claims 1-20, wherein the support-medium comprises polystyrene with 1% crosslinked divinylbenzene.

22. The process of any one of claims 1-21, wherein the compound of Formula (A4) is of Formula (A4a):

10




(A4a);

wherein:


R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl

15 and trimethoxytrityl, and


R^4 is selected from:




(PC),



(DPG),



23. The process of any one of claims 1-22, wherein the compound of Formula (A5) is of Formula (A5a):



(A5a);


wherein:

n is an integer from 10 to 40,


R¹ is a support-medium

R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

R⁴ is selected from:



24. The process of any one of claims 1-23, wherein the compound of Formula (A8) is of Formula (A8a):




(A8a);


wherein:

- R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

R⁴ is, independently at each occurrence of the compound of Formula (A8a), selected from the group consisting of:



- 5 25. The process of any one of claims 1-24, wherein the compound of formula (A9) is of Formula (A9a):



(A9a);


wherein:


n is an integer from 10 to 40,

5 R^1 is a support-medium


R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and

R^4 is, independently for each occurrence, selected from the group consisting of:





26. The process of any one of claims 1-25, wherein the compound of Formula (A10) is of
5 Formula (A10a):




(A10a);

wherein:


n is an integer from 10 to 40,

R¹ is a support-medium, and

R⁴ is, independently for each occurrence, selected from the group consisting of:




- 5 27. The process of any one of claims 1-26, wherein the compound of Formula (A11) is of Formula (A11a):




(A11a);


wherein:

n is an integer from 10 to 40, and

5 R^4 is, independently for each occurrence, selected from the group consisting of:



28. A compound of Formula (B1):



(B1);

or a pharmaceutically acceptable salt thereof

wherein

R^{11} is selected from the group consisting of halo, CN, and NO_2 ;


R^{12} is C_{1-6} -alkyl;

R^{13} is H or C_{1-6} -alkyl; and

R^{14} is a first linker bound to a support-medium,

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl.


29. The compound of claim 28, wherein the compound of Formula (B1) is of Formula (I):



(I),

wherein R^1 is a support-medium.

5 30. A compound of Formula (II-B):




(II-B);

or a pharmaceutically acceptable salt thereof

wherein

- 10 R^{11} is selected from the group consisting of halo, CN, and NO_2 ;
 R^{12} is C_{1-6} -alkyl;
 R^{13} is H or C_{1-6} -alkyl; and
 R^{14} is a first linker bound to a support-medium.


15 31. The compound of claim 30, wherein the compound of Formula (II-B) is Formula (II):



(II);

wherein R^1 is a support-medium.

5 32. A compound of Formula (B3):




(B3);


or a pharmaceutically acceptable salt thereof

wherein

- 10 w is selected from O or C;
 t is 1, 2, 3, 4, or 5;
 R^{11} is selected from the group consisting of halo, CN, and NO_2 ;
 R^{12} is C_{1-6} -alkyl;
 R^{13} is H or C_{1-6} -alkyl;
 15 R^{14} is a first linker bound to a support-medium;
 R^8 is selected from:



wherein R^4 is selected from the group consisting of:




5

and

R^3 is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl.



- 10 33. The compound of claim 32, wherein the compound of Formula (B3) is of Formula (III):



(III);

wherein R^1 is a support-medium.

5 34. A compound selected from:




wherein:

n is an integer from 10 to 40;


R¹ is a support-medium;

5 R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl; and


R⁴ is, independently at each occurrence, selected from the group consisting of:




35. A compound selected from:



;





; and



wherein:

- 5 R¹¹ is selected from the group consisting of halo, CN, and NO₂;
 R¹² is C₁₋₆-alkyl;
 R¹³ is H or C₁₋₆-alkyl;
 R¹⁴ is a first linker bound to a support-medium;
 R¹ is selected from -O(C₁-C₆alkyl), halo, -O(C₁-C₆alkyl)-O(C₁-C₆alkyl), and
 10 -O(C₁-C₆alkyl)-C(O)-NH-C₁-C₆alkyl
 R² is H or R¹ and R² join to form a cycloalkyl or heterocyclic ring having from 4 to 7
 ring atoms optionally containing an oxygen;
 R⁵, R⁶, and R⁷ are each hydrogen, or R⁵ and R⁷ join to form a cycloalkyl ring having
 from 4 to 7 ring atoms and R⁶ and R⁷ join to for a cycloalkyl ring having from 3 to 4
 15 ring atoms,
 R⁹ is -OCH₂CH₂CN or -CH₂C(=O)OC(CH₃)₂CH₂CN;
 R⁴ is, independently at each occurrence, selected from the group consisting of:







R^8 is, independently at each occurrence, selected from the group consisting of H, trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl;

5 n is 10 to 40; and

X is O or S.


36. A compound selected from:








wherein:



- 5 n is an integer from 10 to 40;
- R¹ is a support-medium;
- R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl; and
- R⁴ is, independently at each occurrence, selected from the group consisting of:



5 37. A compound selected from:









wherein n is an integer from 10 to 40, R^1 is a support-medium, X is O or S, R^5 is $-OCH_2CH_2CN$ or $-CH_2C(=O)OC(CH_3)_2CH_2CN$, each A, independently at each occurrence,

5 is selected from the group consisting of:



5 R³ is selected from the group consisting of trityl, monomethoxytrityl, dimethoxytrityl and trimethoxytrityl, and R⁴ is, independently for each occurrence, selected from the group consisting of:











Fig. 1




Fig. 2





Fig. 3



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/034235

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07F9/6558 C07F9/6561
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/082551 A1 (SAREPTA THERAPEUTICS INC [US]) 6 June 2013 (2013-06-06) cited in the application page 47ff.; figures 2-4; example 23 -----	1-37
X	WO 2014/153240 A2 (SAREPTA THERAPEUTICS INC [US]) 25 September 2014 (2014-09-25) page 60, line 1; figure 2B; example 1; compound 34 -----	1-37
X	WO 2011/150408 A2 (AVI BIOPHARMA INC [US]; HANSON GUNNAR J [US]; RUDOLPH ALEXANDER CHARLE) 1 December 2011 (2011-12-01) page 80, line 20 - page 81, line 2; figures 1A, 1B, 4; example 32 ----- -/-	1-37



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

12 July 2017

Date of mailing of the international search report

31/07/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stroeter, Thomas

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/034235

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VIRTA P ET AL: "Solid-supported synthesis of oligomeric bioconjugates", TETRAHE, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 59, no. 28, 7 July 2003 (2003-07-07), pages 5137-5174, XP004434882, ISSN: 0040-4020, DOI: 10.1016/S0040-4020(03)00704-X the whole document -----	1-37

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/034235

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013082551 A1	06-06-2013	AU 2012345638 A1 CA 2857664 A1 EP 2785840 A1 HK 1202891 A1 JP 2015504650 A US 2014329772 A1 WO 2013082551 A1	24-07-2014 06-06-2013 08-10-2014 09-10-2015 16-02-2015 06-11-2014 06-06-2013
WO 2014153240 A2	25-09-2014	AU 2014236160 A1 CA 2903872 A1 CN 105392884 A EA 201591694 A1 EP 2970964 A2 HK 1218138 A1 JP 2016521119 A KR 20150133752 A NZ 631132 A US 2014315977 A1 US 2016040162 A1 WO 2014153240 A2	01-10-2015 25-09-2014 09-03-2016 30-06-2016 20-01-2016 03-02-2017 21-07-2016 30-11-2015 30-06-2017 23-10-2014 11-02-2016 25-09-2014
WO 2011150408 A2	01-12-2011	AU 2011257980 A1 AU 2016234953 A1 CA 2799501 A1 CN 103003288 A EP 2576574 A2 JP 2013530154 A JP 2016000059 A KR 20130118227 A NZ 603606 A TW 201202262 A US 2012065169 A1 US 2015073140 A1 WO 2011150408 A2	10-01-2013 13-10-2016 01-12-2011 27-03-2013 10-04-2013 25-07-2013 07-01-2016 29-10-2013 26-06-2015 16-01-2012 15-03-2012 12-03-2015 01-12-2011