

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Method and reagent for measuring nitroreductase enzyme activity

(51)⁶ International Patent Classification(s)
C09B 57/00 (2006.01) 1/26
C12Q 1/26 (2006.01) 20060101ALI2005100
C12Q 1/32 (2006.01) ^{8BMEP} **C12Q**
C09B 57/00 1/32
20060101AFI2005100 20060101ALI2007072
^{8BMEP} **C12Q** ^{1BMEP}
PCT/GB2005/002057

(21) Application No: 2005250209

(22) Application Date: 2005 .05 .24

(87) WIPO No: W005/118839

(30) Priority Data

(31) Number	(32) Date	(33) Country
0411993.9	2004 .05 .28	GB

(43) Publication Date : 2005 .12 .15

(71) Applicant(s)
GE Healthcare UK Limited

(72) Inventor(s)
Ismail, Rahman, West, Richard Martin

(74) Agent/Attorney
Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
US 2003/0186348 A1 (THOMAS) 2 October 2003

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 December 2005 (15.12.2005)


PCT

(10) International Publication Number
WO 2005/118839 A1

- (51) International Patent Classification⁷: C12Q 1/26, C09B 57/00
- (21) International Application Number: PCT/GB2005/002057
- (22) International Filing Date: 24 May 2005 (24.05.2005)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0411993.9 28 May 2004 (28.05.2004) GB
- (71) Applicant (for all designated States except US): AMERSHAM BIOSCIENCES UK LIMITED [GB/GB]; Amersham Place, Little Chalfont Buckinghamshire HP7 9NA (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): WEST, Richard Martin [GB/GB]; Amersham Biosciences UK Limited, The Maynard Centre, Forest Farm Estate, Whitechurch, Cardiff CF14 7YT (GB). ISMAIL, Rahman [GB/GB]; Amersham Biosciences UK Limited, The Maynard Centre, Forest Farm Estate, Whitechurch, Cardiff CF14 7YT (GB).
- (74) Agents: FRANKS, Barry et al.; GB Healthcare Limited, Amersham Place, Little Chalfont Buckinghamshire HP7 9NA (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KR, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: METHOD AND REAGENT FOR MEASURING NITROREDUCTASE ENZYME ACTIVITY




(I)



(A)

NITRO-SUBSTITUTED CYANINE DYE NITROREDUCTASE SUBSTRATES:
COMPOUNDS (i) AND (ii)



(57) Abstract: Disclosed are nitro-substituted squaraine reporter dyes and methods using such dyes for detecting nitroreductase enzymic activity and nitroreductase gene expression in cellular assays. The dyes are of the structure: formula (I) in which Z¹ and Z² independently represent a phenyl or a naphthyl ring system; X and Y are selected from oxygen, sulphur, -CH=CH- and the group: formula (A); R¹ and R² are selected from C₁-C₄ alkyl, -(CH₂)_n-P, -(CH₂)₂-O-R⁶ and group W; where P is selected from COOR⁷, SO₂- and OH, W is mono- or di-substituted nitrobenzyl, R⁶ is methyl or ethyl, R⁷ is selected from H, C₁-C₄ alkyl and CH₂OC(O)R⁸, where R⁸ is methyl, or t-butyl, n is an integer from 1 to 10, and p is an integer from 1 to 3; R³ and R⁴ are selected from hydrogen, NO₂, halogen, SO₂-, C₁-C₄ alkoxy and -(CH₂)_m-COOR⁷; where R⁷ is hereinbefore defined and m is 0 or an integer from 1 to 5; R⁵ is C₁-C₅ alkyl optionally substituted with COOR⁷, SO₂-, or OH; where R⁷ is hereinbefore defined; and at least one of groups R¹, R², R³ and R⁴ comprises at least one NO₂ group. Also provided are methods for screening for a test agent whose effect upon nitroreductase enzyme activity and nitroreductase gene expression is to be determined.

WO 2005/118839 A1



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LI, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NI, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— with international search report


1

Method and Reagent for Measuring Nitroreductase Enzyme Activity

The present invention relates to the field of enzyme assays. In particular, the invention relates to nitroreductase enzyme assays and new
5 reporter dyes for measuring nitroreductase enzyme activity and nitroreductase gene expression in cell based systems.

Reporter gene technology is widely used to monitor cellular events associated with signal transduction and gene expression. Transcriptional
10 regulation, coupled to the expression of a reporter gene is routinely used to monitor a wide variety of cellular events. To establish a reporter gene assay, the reporter gene is placed under the transcriptional control of a promoter or an enhancer with a minimal promoter. The reporter is inserted into a suitable plasmid vector typically containing a selectable marker that confers resistance
15 to growth suppressing compounds, such as antibiotics. The vector DNA is introduced into cells using standard laboratory procedures. Addition of a suitable agonist will switch on the cell signalling pathway, leading to activation of a transcription factor and gene expression. A review of reporter gene technology is given by Naylor et al, in *Biochem.Pharmacol.*, (1999), 58, 749-
20 757.

A cell-based fluorescent gene reporter system has been described, the assay employing bacterial nitroreductase (NTR) and a cell permeable nitro-
substituted quenched (or non-fluorescent) cyanine dye (shown as Compound
25 (i)), which functions as a substrate for the enzyme (US 2003/0186348, Thomas, N. et al).



Compound (I)

Cellular uptake of the substrate, by passive diffusion across the plasma membrane, was promoted through the use of ethyl ester groups to mask latent polar functionality. Intracellular cleavage of the ester groups by cellular hydrolases results in retention of the substrate inside live cells. Addition of the substrate to a cell that is expressing nitroreductase results in the reduction of the nitro group to the hydroxylamine with a concomitant increase in fluorescence emission. Depending on the structure of the quenched cyanine dye, the fluorescence emission from the product of the NTR reaction may be generated across a wide range of wavelengths, typically 500-900nm. Emission at longer wavelengths is advantageous in avoiding background fluorescence and increasing sensitivity in biological systems.

Wild type nitroreductase expressed from a reporter construct is localised in the cytoplasm of the host cell (Spooner et al, *Int.J.Cancer*, (2001), 93, 123-30). To achieve a maximum signal output from the assay it is desirable to localise the substrate to the same cellular compartment as the reporter enzyme, i.e. within the cytoplasm of the host cell such that the substrate is available for activation by nitroreductase. The masking of hydrophilic groups on, or attached to, the substrate molecule can generate membrane permeable compounds. Furthermore, the masking group can be designed to cleave from the substrate within the cell to generate the substrate intracellularly, preferably within the cytoplasm of the cell. Masking strategies to enable delivery of nitro-substituted cyanine dyes relatively uniformly to the cell cytoplasm have not proved to be entirely successful. A study of the localisation of cell permeant


quenched cyanine dye (Cy-Q) derivatives within cells using fluorescence microscopy, has shown the localisation of some of the substrate to internal cell membranes and organelles, predominantly the mitochondria of the cell.

Accumulation of lipophilic, cationic nitro-substituted cyanine dye substrates

5 within mitochondria is accompanied by an increase in fluorescence of the probe and this accumulation results in an increase in background fluorescence in NTR assays. Thus, there is a need for new and improved reagents for use as NTR substrates that display lower background fluorescence, improved fluorescence signal and cellular distribution.

10

Squarylium (squaraine) dyes are a class of dyes that have overall electrical neutrality; an example is shown as Compound (ii).



Compound (ii)

15

Nitro-substituted squaraine dyes are known from EP 645680 (Bugner D., et al) as near infra-red absorbing additives for use in electrophotographic imaging processes. PCT Application No.WO97/40104 (Hamilton, A.L. et al) discloses squaraine dyes and adducts of squaraine dyes with biological molecules such

20

as peptides, proteins and nucleotides. The dyes may be substituted by electron donating and electron withdrawing substituents, for example nitro; however, the fluorescence properties of the nitro-substituted dyes are not disclosed. The present inventors have now discovered that nitro-group-containing quenched squaraine dyes are effective substrates for nitroreductase through reduction of the nitro group, resulting in a change in an optical property, preferably a change in fluorescence emission, of the squaraine dye.

25

Use of nitro-substituted squaraine dyes in assays for determining nitroreductase activity results in greater sensitivity and lower background fluorescence than in assays that employ conventional NTR substrates.

In a first aspect of the invention, there is provided a method for detecting nitroreductase enzyme activity in a composition comprising:

- i) mixing said composition under conditions to promote nitroreductase activity with a dye molecule; and
- 5 ii) measuring a change in an optical property of said dye molecule wherein said change is a measure of the amount of nitroreductase activity; characterised in that said dye molecule is a squaraine dye comprising at least one NO₂ group.

- 10 In one embodiment, the composition in which nitroreductase enzyme activity is to be detected comprises at least one cell or a cell extract. The cell may be *ex-vivo*, or *in-vivo*. For example, the cell may be cultured under standard laboratory conditions, or the composition may be a live animal cell.

- 15 In another embodiment, the method is conducted in the presence of a test agent whose effect on nitroreductase enzyme activity is to be determined.


In a second aspect of the invention there is provided a method which comprises:

- 20 i) contacting a host cell with a dye molecule wherein said host cell has been transfected with a nucleic acid molecule comprising expression control sequences operably linked to a sequence encoding a nitroreductase; and
- ii) measuring a change in an optical property of said dye molecule wherein said change is a measure of the amount of nitroreductase activity;
- 25 characterised in that said dye molecule is a squaraine dye comprising at least one NO₂ group.

- The optical property that is measured in the dye molecule is suitably the fluorescence emission intensity, such that there is an increase in fluorescence emission as a result of the action of the enzyme upon the dye. For example,
- 30 the composition may be excited at a first wavelength, suitably the excitation maximum of the dye, and the fluorescence emission intensity measured at a second wavelength corresponding to the emission maximum of the product of

the enzyme reaction. Excitation of the dye molecule and measurement of fluorescence emission may also be over a range of wavelengths, so as to maximise emission signal and to distinguish between excitation and emission signals. Alternatively, the measured change in an optical property may be a change in fluorescence lifetime of the dye, before and after the action of the nitroreductase enzyme. The change in fluorescence lifetime may also be used to distinguish the product of the enzyme reaction from the dye molecule used as the substrate. As a further alternative, the change in an optical property may be a change in the absorption maximum of the dye molecule, relative to the absorption maximum of the product. In preferred embodiments, the change in an optical property is an increase in the fluorescence intensity of the dye molecule, whereby the increase is a measure of the amount of nitroreductase activity.

Suitably, the squaraine dye according to the first and second aspects is a compound of formula (I):



(I)

wherein:

R^3 is attached to the Z^1 ring structure and R^4 is attached to the Z^2 ring structure;

Z^1 and Z^2 independently represent a phenyl or a naphthyl ring system;

X and Y are the same or different and are selected from oxygen, sulphur,

$-\text{CH}=\text{CH}-$ and the group:



groups R^1 and R^2 are independently selected from C_1 - C_4 alkyl, $-(CH_2)_n-P$, $-((CH_2)_2-O)_p-R^6$ and the group W; where P is selected from $COOR^7$, SO_3^- and OH, W is mono- or di-substituted nitrobenzyl, R^6 is methyl or ethyl, R^7 is selected from H, C_1 - C_4 alkyl and $CH_2OC(O)R^8$, where R^8 is methyl, or t-butyl, n
 5 is an integer from 1 to 10, and p is an integer from 1 to 3;
 groups R^3 and R^4 are independently selected from hydrogen, NO_2 , halogen, SO_3^- , C_1 - C_4 alkoxy and $-(CH_2)_m-COOR^7$; where R^7 is hereinbefore defined and m is 0 or an integer from 1 to 5;
 R^5 is C_1 - C_6 alkyl optionally substituted with $COOR^7$, SO_3^- , or OH; where R^7 is
 10 hereinbefore defined; and
 at least one of groups R^1 , R^2 , R^3 and R^4 comprises at least one NO_2 group.




Suitably, the squaraine dye of formula (I) may include a counter-ion, which may be positive or negative to balance the formal charge (or charges) on
 15 the dye chromophore or on substituent groups. The nature of the counter-ion is not material to the invention and could be one of many known ions such as H^+ , NH_4^+ , K^+ , Na^+ , trifluoroacetate ($F_3C-CO_2^-$), perchlorate (ClO_4^-), Br^- , or I^- .

Suitably, the at least one nitro group comprised in the dyes of formula
 20 (I), may be attached directly to the Z^1 and Z^2 ring structures. In this embodiment, one or both of groups R^3 and R^4 of the squaraine dye are NO_2 . In an alternative embodiment, one or both of groups R^1 and R^2 of said squaraine dye are the group W, where W is hereinbefore defined. The squaraine dye may optionally be further substituted with one or two nitro groups attached to
 25 the aromatic ring structures.

In preferred embodiments, the squaraine dye employed in the methods according to the invention is permeable to cells. In these embodiments, at least one of groups R^1 , R^2 , R^3 and R^4 comprises a cell membrane
 30 permeabilising group. Membrane permeant compounds can be generated by masking hydrophilic groups to provide more hydrophobic compounds. The masking groups can be designed to be cleaved from the substrate within the cell to generate the derived substrate intracellularly. Since the substrate is

more hydrophilic than the membrane permeant derivative, it is trapped within the cell. Suitable cell membrane permeabilising groups may be selected from acetoxymethyl ester which is readily cleaved by endogenous mammalian intracellular esterases (Jansen, A. B. A. and Russell, T. J., J. Chem. Soc., 2127-2132 (1965) and Daehne, W. et al. J. Med. Chem. 13, 697-612 (1970)) and pivaloyl ester (Madhu et al., J. Ocul. Pharmacol. Ther., (1998), 14, 5, pp 389-399), although other suitable groups will be recognised by those skilled in the art.

10 In one embodiment, one or both of groups R^1 and R^2 of the squaraine dye is the group W, where W is hereinbefore defined. Particular examples of dyes utilised in this embodiment of the methods of the invention are those selected from dyes of formula (II), (III) and (IV):



15 wherein:

X and Y are the same or different and are selected from oxygen, sulphur, $-\text{CH}=\text{CH}-$ and the group:




wherein R^5 is hereinbefore defined;

at least one of groups R^1 and R^2 is the group W; where W is hereinbefore defined;

- 5 any remaining group R^1 or R^2 is selected from C_1 - C_4 alkyl, $-(CH_2)_n-P$ and $-[(CH_2)_2-O]_p-R^6$; where P is selected from $COOR^7$, SO_3^- and OH, R^6 is methyl or ethyl, R^7 is selected from H, C_1 - C_4 alkyl and $CH_2OC(O)R^8$, where R^8 is methyl, or t-butyl, n is an integer from 1 to 10 and p is an integer from 1 to 3; and
- 10 groups R^3 and R^4 are independently selected from hydrogen, halogen, SO_3^- , C_1 - C_4 alkoxy and $-(CH_2)_m-COOR^7$; where R^7 is hereinbefore defined and m is 0 or an integer from 1 to 5.


- In this embodiment, preferably one of groups R^1 and R^2 is selected from group W where W is selected from:
- 15



remaining R^1 or R^2 is selected from methyl and ethyl, or is the group

$-(CH_2)_n-COOR^7$ where R^7 is selected from H, C_1 - C_4 alkyl and $CH_2OC(O)R^8$, where R^8 is methyl, or t-butyl, n is an integer from 1 to 10, preferably 5 or 6. In

- 20 a particularly preferred embodiment, W is the group:



and remaining R^1 or R^2 is hereinbefore defined.

In an alternative embodiment, one or both of groups R^3 and R^4 of the squaraine dye according to formula (II), (III) and (IV) are NO_2 . In this embodiment, X and Y are the same or different and are selected from oxygen, sulphur, $-CH=CH-$ and the group:



5

wherein R^5 is hereinbefore defined;

groups R^1 or R^2 are independently selected from C_1 - C_4 alkyl, $-(CH_2)_n-P$ and $-[(CH_2)_2-O]_p-R^6$; where P is selected from $COOR^7$, SO_3^- and OH, R^6 is methyl or ethyl, R^7 is selected from H, C_1 - C_4 alkyl and $CH_2OC(O)R^8$, where R^8 is

10

methyl, or t-butyl, n is an integer from 1 to 10 and p is an integer from 1 to 3;

at least one of groups R^3 and R^4 is NO_2 ; and

any remaining group R^3 or R^4 is selected from hydrogen, SO_3^- , C_1 - C_4 alkoxy

and $-(CH_2)_m-COOR^7$; where R^7 is selected from H, C_1 - C_4 alkyl and

$CH_2OC(O)R^8$, where R^8 is methyl, or t-butyl, and m is 0 or an integer from 1 to

15

5.

In preferred embodiments X and Y are selected from oxygen, sulphur and



20

where R^5 is methyl.

Preferred C_1 - C_4 alkyl groups are selected from methyl and ethyl. A particularly preferred $-(CH_2)_n-COOR^7$ group is selected from $-(CH_2)_5-COOR^7$, and $-(CH_2)_6-COOR^7$; where R^7 is hereinbefore defined.

25

Halogen atoms are selected from fluorine, chlorine, bromine and iodine.

In a third aspect there is provided a method for screening for a test agent whose effect upon nitroreductase gene expression is to be determined.

The method comprises the steps of: a) performing the method according to the second aspect in the absence and in the presence of said test agent; and b) determining the amount of nitroreductase gene expression in the absence and in the presence of said agent; wherein a difference between the amount of nitroreductase gene expression in the absence and in the presence of said agent is indicative of the effect of said agent on nitroreductase gene expression.

In an alternative aspect, the method for screening for a test agent may be conducted by a) performing the method according to the second aspect in the presence of said agent; and b) comparing the amount of nitroreductase gene expression with a control value for the amount of nitroreductase gene expression in the absence of the test agent. The control values may be stored electronically in a database or other electronic format.

Methods for using a variety of enzyme genes as reporter genes in mammalian cells are well known (for a review see Naylor L.H. (1999) *Biochemical Pharmacology* **58**, 749-757). The reporter gene is chosen to allow the product of the gene to be measurable in the presence of other cellular proteins and is introduced into the cell under the control of a chosen regulatory sequence which is responsive to changes in gene expression in the host cell. Typical regulatory sequences include those responsive to hormones, second messengers and other cellular control and signalling factors. For example, agonist binding to seven transmembrane receptors is known to modulate promoter elements including the cAMP responsive element, NFAT, SRE and AP1; MAP kinase activation leads to modulation of SRE leading to Fos and Jun transcription; DNA damage leads to activation of transcription of DNA repair enzymes and the tumour suppressor gene p53. By selection of an appropriate regulatory sequence the reporter gene can be used to assay the effect of added agents on cellular processes involving the chosen regulatory sequence under study.

For use as a reporter, the nitroreductase gene may be isolated by well known methods, for example by amplification from a cDNA library by use of the polymerase chain reaction (PCR) (Molecular Cloning, A Laboratory Manual 2nd Edition, Cold Spring Harbour Laboratory Press 1989 pp 14.5-14.20). Once
5 isolated, the nitroreductase gene may be inserted into a vector suitable for use with mammalian promoters (Molecular Cloning, A Laboratory Manual 2nd Edition, Cold Spring Harbour Laboratory Press 1989 pp 16.56-16.57) in conjunction with and under the control of the gene regulatory sequence under study. The vector containing the nitroreductase reporter and associated
10 regulatory sequences may then be introduced into the host cell by transfection using well known techniques, for example by use of DEAE-Dextran or Calcium Phosphate (Molecular Cloning, A Laboratory Manual 2nd Edition, Cold Spring Harbour Laboratory Press 1989 pp 16.30-16.46). Other suitable techniques will be well known to those skilled in the art. Nitroreductase has been shown to be
15 retained in cells when expressed in this manner (see Bridgewater et al., Eur. J. Cancer, (1995), 31A, 2362-70).

The methods of the invention may be used with any adherent cell type that can be cultured on standard tissue culture plastic-ware, including cell types
20 derived from any recognised source with respect to species (e.g. human, rodent, simian), tissue source (e.g. brain, liver, lung, heart, kidney skin, muscle) and cell type (e.g. epithelial, endothelial). There are established protocols available for the culture of diverse cell types. (See for example, Freshney, R.I., Culture of Animal Cells: A Manual of Basic Technique, 2nd Edition, Alan R. Liss
25 Inc. 1987). The chosen host cell line is seeded into sterile, tissue culture treated dishes and incubated at 37°C in an humidified atmosphere of 5% CO₂ in a suitable medium, typically Dulbecco's Modified Eagles medium containing 10% foetal calf serum + 2mM L-glutamine. Transfection of the plasmid vector into mammalian cells may be achieved using well known methods, e.g. by the
30 use of cationic lipids, calcium phosphate, and electroporation. It is recommended that transfection efficiencies are optimised for each cell line prior to testing to ensure that reproducible data are obtained. Transient expression of nitroreductase is typically assayed 24-72 hours post transfection. The

prepared nitroreductase gene reporter DNA/transfection reagent complex is added in a dropwise manner to each dish. The contents of the dish are carefully mixed and incubated for a minimum of 4 hours. Overnight incubation at 37°C in a humidified atmosphere of 5% CO₂ is convenient. Following
5 incubation, medium is removed from each dish and the cell monolayer washed with sterile phosphate buffered saline (PBS). Transfected cells may be assayed directly in the transfection dish, or alternatively cells may be detached from each dish pooled to produce a suspension of transfected cells. Transient expression of nitroreductase is typically assayed 24-72 hours post transfection.

10

In a typical adenoviral based NTR gene reporter assay according to the invention, the chosen host cell line is subcultured twenty-four hours prior to viral transduction and incubated overnight at 37°C in a humidified atmosphere of 5% CO₂. Cells are detached with trypsin and cells from each flask pooled to
15 produce a suspension of cells. Cells in suspension are combined with virus at a predetermined multiplicity of infection (MOI) in a sufficient volume of complete medium to cover the base of a suitable tissue culture treated flask and incubated overnight at 37°C in a humidified atmosphere of 5% CO₂. Cells are detached (trypsin) to produce a suspension of transduced cells.

20

Suitably, the nitroreductase containing vector may be used to produce both transient and stable cells for use in gene reporter assays. For stable cell line production, selection with a suitable reagent, such as the antibiotic G418 is necessary. According to this procedure, cells should be seeded at a low
25 density, suitably 100-500 into a suitable dish, and the selection agent added to the medium. The optimum concentration of selection agent will vary according to the cell type and growth rate required and is suitably added at a concentration of between 0.1 mg/ml and 1 mg/ml.

30

For the assay of the effect of a test agent on nitroreductase activity, cells are dispensed in the wells of a microwell plate, suitably a microtitre plate having 24, 96, 384 or higher densities of wells, e.g. 1536 wells. Following overnight incubation at 37°C, medium is removed and the test agent is added in serum

free medium. Wells containing serum free medium only are used as the control. Following incubation, the nitroreductase substrate is added and the fluorescent signal increase is measured over time using a suitable fluorimeter or imaging system.

5

To assay the activity of a test agent to activate cellular responses via the regulatory sequence under study, cells that have been transfected with the nitroreductase reporter are incubated with the test agent, followed by addition of a cell-permeant squaraine dye substrate, such as a squaraine dye comprising at least one NO₂ group. After an appropriate period required for conversion of the dye substrate to a form emitting increased fluorescence, the fluorescence emission from the cells is measured over time at a wavelength appropriate for the chosen squaraine dye, using a suitable fluorimeter or imaging system.

15

Typically, gene reporter assays are performed under "stopped" conditions, e.g. lysis of cells for detection of reporter gene. Thus, the reaction is allowed to proceed for a predetermined time and then terminated with a stop reagent, normally a surfactant. An example of a stop reagent is Triton X-100, which is used to disrupt cell membranes and release the enzymatic activity. In addition, cells may be "fixed" using standard reagents, such as formaldehyde, and the product of the nitroreductase reaction retained within the cell. This allows storage of assay plates until a suitable time for reading is available.

20

25


Where an assay is to be formatted for the determination of the activity of a test agent on nitroreductase activity, the assay may be performed under continuous measurement of the fluorescence of the substrate. In this format, the fluorescence emission intensity of the substrate changes continuously. A time-course of the reaction may be obtained, allowing kinetic studies to be performed in real time. Measurements of emitted fluorescence may be compared with fluorescence measurements from control cells not exposed to the test agent and the effects, if any, of the test agent on gene expression



30

modulated through the regulatory sequence is determined from the ratio of fluorescence in the test cells to the fluorescence in the control cells.

Measurements of changes in fluorescence intensity may be made using instruments incorporating photo-multiplier tubes as detectors, for example an "Ultra" fluorimeter (Tecan), or by means of a charge coupled device (CCD) imager (such as a scanning imager or an area imager) to image all of the wells of a microtitre plate. The LEADseeker™ system features a CCD camera allowing fluorescence imaging of high density microtitre plates in a single pass. Imaging is quantitative and fast, and instrumentation suitable for imaging applications can now simultaneously image the whole of a multiwell plate. Alternatively, cells may be imaged in "live cell" format using an INCell™ 1000 Analyzer or INCell™ 3000 Analyzer. In this format, a suitable cell marker should be introduced into the cell, such as a cytosolic, nuclear or membrane fluorescent label having a fluorescence emission wavelength that is different and distinguishable from the fluorescence emission of the reduced substrate. Suitably, the increase in fluorescence emitted by the substrate is detected at a wavelength in the range 500 nm to 900 nm, preferably 550-780 nm, and, most preferably 630-700 nm. For example, for Compound (1) (Example 1), the fluorescence emission may be monitored at 645 nm with excitation at 630 nm. Alternatively, the dye may be administered *in vivo* to a suitably engineered transgenic animal model. Nitroreductase activity and localisation may then be determined by imaging with a suitable optical system, for example, the eXplore Optix™.

In a further aspect, the present invention provides nitro-substituted squaraine dyes selected from dyes of formula:





wherein:

X and Y are the same or different and are selected from oxygen, sulphur, $-\text{CH}=\text{CH}-$ and the group:




- 5 groups R^1 and R^2 are independently selected from $\text{C}_1\text{-C}_4$ alkyl, $-(\text{CH}_2)_n\text{-P}$, $-\{(\text{CH}_2)_2\text{-O}\}_p\text{-R}^6$ and the group W; where P is selected from COOR^7 , SO_3^- and OH, W is mono- or di-substituted nitrobenzyl, R^6 is methyl or ethyl, R^7 is selected from H, $\text{C}_1\text{-C}_4$ alkyl and $\text{CH}_2\text{OC(O)R}^8$, where R^8 is methyl, or t-butyl, n is an integer from 1 to 10, and p is an integer from 1 to 3;
- 10 groups R^3 and R^4 are independently selected from hydrogen, NO_2 , halogen, SO_3^- , $\text{C}_1\text{-C}_4$ alkoxy and $-(\text{CH}_2)_m\text{-COOR}^7$; where R^7 is selected from H, $\text{C}_1\text{-C}_4$ alkyl and $\text{CH}_2\text{OC(O)R}^8$, where R^8 is methyl, or t-butyl, and m is 0 or an integer from 1 to 5;
- R^5 is $\text{C}_1\text{-C}_6$ alkyl optionally substituted with COOR^7 , SO_3^- , or OH; where R^7 is
- 15 hereinbefore defined; and
- at least one of groups R^1 , R^2 , R^3 and R^4 comprises at least one NO_2 group.

Preferably, X and Y are selected from oxygen, sulphur and



where R⁵ is methyl.

In one embodiment, one of groups R¹ and R² is



5

remaining R¹ or R² is selected from methyl, ethyl and the group $-(CH_2)_n-$
COOR⁷; where R⁷ is hereinbefore defined and n is an integer from 1 to 10,
preferably 5 or 6.

10

In an alternative embodiment, groups R¹ and R² are independently
selected from C₁-C₄ alkyl, $-(CH_2)_n-$ COOR⁷ and $-((CH_2)_2-O)_p-R^8$; where R⁸ is
methyl or ethyl, R⁷ is selected from H, C₁-C₄ alkyl and CH₂OC(O)R⁸, where R⁸
is methyl, or t-butyl, and n and p are hereinbefore defined;
at least one of groups R³ and R⁴ is NO₂; and

15

remaining group R³ or R⁴ is selected from hydrogen, SO₃⁻, C₁-C₄ alkoxy and
 $-(CH_2)_m-$ COOR⁷; where R⁷ is hereinbefore defined and m is 0 or an integer
from 1 to 5.

20

The squaraine dyes are useful as substrates for the detection and/or
measurement of nitroreductase enzyme activity and in particular for measuring
the amount of nitroreductase gene expression in cellular assays.

Examples of dyes according to the first aspect of the invention are as
follows:

25

i) 2-(1-methyl-3,3-dimethyl-2-indolinyliidenemethyl)-4-(1-(3,5-dinitrobenzyl)-
3,3-dimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound
1);

- ii) 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound 2);
- iii) 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-benzindolinyliidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound 3);
- iv) 2-(3-ethyl-6-nitro-2-benzothiazolinyliidenemethyl)-4-(1-(2-(2-methoxyethoxy)ethyl)-3,3-dimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound 4);
- v) 2-(1-ethyl-3,3-dimethyl-5-methoxy-2-indolinyliidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound 5);
- vi) 2-(1-(5-carboxypentyl)-3,3-dimethyl-5-methoxy-2-indolinyliidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound 6);
- vii) 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-3-methyl-1,3-dihydro-2H-indol-2-ylidenemethyl-4-((1-(3,5-dinitrobenzyl)-3,3-dimethyl-3H-indolium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound 7); and
- viii) 2-((3,3-dimethyl-5-sulfo-1-(4-sulfobutyl)-1,3-dihydro-2H-indol-2-ylidene)methyl)-4-((1-methyl-6-nitroquinolinium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound 8)


The present invention is further illustrated with reference to the following Figures and Examples in which:

- Figure 1 shows the molecular structures of two nitro-substituted cyanine type dyes, Compounds (i) and (iii) compared with a nitro-substituted squaraine dye of the invention (Compound (1)) as substrates in a nitroreductase gene reporter assay, as in Example 11;
- Figure 2 illustrates the comparison of two nitro-substituted cyanine dyes, Compounds (i) and (iii) compared with a nitro-substituted squaraine in an NTR gene reporter assay (Compound (1));

Figure 3 is a comparative study of nitro-substituted squaraine dyes (Compounds (1) and 2) as substrates in a nitroreductase gene reporter assay; Figure 4 shows the distribution of Compound (2) in HeLa cells; and Figure 5 shows the evaluation of Compounds (2), (3) and (4) as nitroreductase substrates in live cell NTR assays.

Examples

1. Preparation of 2-(1-methyl-3,3-dimethyl-2-indolinylidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinylidenemethyl)cyclobutenediylum-1,3-diolate (Compound (1))



Compound (1)

- 15 1.1 Preparation of 1-(3,5-Dinitrobenzyl)-2,3,3-trimethyl-3H-indolium iodide

To 2,3,3-trimethylindolenine (1.64g) was added 3,5-dinitrobenzyl iodide (3.71g) and dichlorobenzene (15ml). After heating to 90°C for 6 hours the mixture was allowed to cool and the resultant precipitate removed by filtration.

- 20 The solid was washed with dichlorobenzene (2x10ml) and ether (2x50ml). The material was dried in a vacuum oven to give the product as a yellow solid (2.69g).


MALDI-TOF ($C_{18}H_{18}N_3O_4$ requires M^+ 340) 339, 340.

25

1.2 Preparation of Compound (1)

To 1-(3,5-dinitrobenzyl)-2,3,3-trimethyl-3H-indolium iodide (100mg) was added 3-hydroxy-4-(1,3,3-trimethyl-1,3-dihydroindol-2-ylidenemethyl)cyclobut-3-ene-1,2-dione (54mg), pyridine (2.25ml), acetic acid (2.25ml) and acetic anhydride (0.5ml). The mixture was heated to reflux for 6 hours and the solvent then stripped using rotary evaporation. The residue was partitioned between water and dichloromethane, and the organic phase sequentially washed with dilute aqueous sodium hydrogen carbonate solution and 1M HCl. The solvent was stripped and silica flash column chromatography performed (MeOH/DCM). The resulting material was further purified by reverse phase HPLC (CH₃CN / H₂O / TFA). MALDI-TOF (C₃₄H₃₀N₄O₆ requires M⁺ 590) 591.

15 2. Preparation of 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-indolinylidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinylidenemethyl)cyclobutenediylum-1,3-diolate (Compound (2))



Compound (2)

20


To 1-(3,5-dinitrobenzyl)-2,3,3-trimethyl-3H-indolium iodide (467mg) was added 3,4-dihydroxy-3-cyclobuten-1,2-dione (110mg), 1-(5-carboxypentyl)-2,3,3-trimethyl-3H-indolium iodide (354mg), pyridine (4.5ml), acetic acid (4.5ml) and acetic anhydride (1ml). The mixture was heated to reflux for 3 hours and the solvent then stripped using rotary evaporation. This crude material was

25

subjected to silica flash column chromatography (eluted with MeOH / DCM). Fractions containing product were combined and stripped of solvent. The resulting material was further purified by reverse phase HPLC (CH₃CN / H₂O / TFA).

5

3. Preparation of 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-benzindolinylidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinylidenemethyl)cyclobutenediylium-1,3-diolate (Compound (3))

10
Compound (3)

3.1 Preparation of 3-(5-carboxypentyl)-1,1,2-trimethyl-1H-benzo[e]indolium iodide

15 To 1,1,2-trimethyl-1H-benzo[e]indolenine (16.2g) was added 6-bromohexanoic acid (31.2g) and dichlorobenzene (50ml). The mixture was heated at 110°C for 136hrs, cooled to ambient temperature, chilled upon ice and filtered. The filter cake was washed with dichlorobenzene (50ml), diethyl ether (50ml) and dried at 40°C under a low vacuum to afford the title compound
20 as a beige solid (25.38g).


LCMS (C₂₁H₂₆NO₂ requires M⁺ 324) 324.

25

3.2 Preparation of Compound (3)

To 1-(3,5-dinitrobenzyl)-2,3,3-trimethyl-3H-indolium iodide (132mg) was added 3-(5-carboxypentyl)-1,1,2-trimethyl-1H-benzo[e]indolium iodide (114mg), 3,4-dihydroxy-3-cyclobuten-1,2-dione (32mg), pyridine (4.5ml), acetic acid (4.5ml) and acetic anhydride (1ml). The mixture was heated to 90°C for 4 hours and the solvent then stripped using rotary evaporation. Silica flash column chromatography was performed (EA/MeOH) and the relevant fractions combined and concentrated. The resulting material was further purified by reverse phase HPLC (CH₃CN / H₂O / TFA) to give 1.7mg. MALDI-TOF (C₄₃H₄₀N₄O₈ requires M⁺ 740) 741.

4. Preparation of 2-(3-ethyl-6-nitro-2-benzothiazolinyliidenemethyl)-4-(1-(2-(2-methoxyethoxy)ethyl)-3,3-dimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (Compound (4))



Compound (4)

4.1 Preparation of 1-(2-(2-methoxyethoxy)ethyl)-2,3,3-trimethyl-3H-indolium bromide

To 2,3,3-trimethylindolenine (1.59g) was added 1-bromo-2-(2-methoxyethoxy)ethane (2.75g) and dichlorobenzene (5ml). The mixture was heated to 70°C overnight. The volatiles were stripped and the material purified by HPLC. MALDI-TOF (C₁₆H₂₄NO₂ requires M⁺ 262) 263.

4.2 Preparation of 2-methyl-6-nitrobenzothiazole

2-Methylbenzothiazole (22g) in conc. sulfuric acid (80ml) was cooled to -
5°C. A mixture of conc. sulfuric acid (12ml) in conc. nitric acid (20ml) was
5 added so as to maintain the temperature below 5°C (ca. 1.5 hours). After this
time the mixture was allowed to warm to room temperature and the solution
poured onto ice to give a yellow precipitate. The solid was removed by filtration
and recrystallised from ethanol. After filtration the solid was washed with
ethanol and dried in a vacuum oven to give 18g of the desired product.
10 LCMS ($C_8H_6N_2O_2S$ requires M^+ 194) 195.


4.3 Preparation of 3-ethyl-2-methyl-6-nitrobenzothiazolium iodide

To 2-methyl-6-nitrobenzothiazole (0.36g) was added ethyl iodide (1.5ml)
15 and dichlorobenzene (20ml). The mixture was heated to 120°C for 2 days and
then allowed to cool to room temperature. Ethyl acetate was added and the
resulting precipitate removed by filtration. Drying in a vacuum oven gave the
desired material (80mg).

20 4.4 Preparation of Compound (4)

To 3-ethyl-2-methyl-6-nitrobenzothiazolium iodide (400mg) was added
3,4-dihydroxy-3-cyclobuten-1,2-dione (128mg), 1-(2-(2-methoxyethoxy)ethyl)-
2,3,3-trimethyl-3H-indolium bromide (420mg), pyridine (20ml), acetic acid
25 (18ml) and acetic anhydride (8ml). The mixture was heated to 120°C for 4
hours and then allowed to cool to room temperature. The volatiles were
stripped by rotary evaporation; the residue dissolved in DCM, and silica flash
chromatography performed (DCM / EA / MeOH). The material was further
purified by prep. TLC to give 23mg.
30 LCMS ($C_{30}H_{37}N_3O_6S$ requires M^+ 561) 560.

5. Preparation of 2-(1-ethyl-3,3-dimethyl-5-methoxy-2-indolinylidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinylidenemethyl)cyclobutenediylum-1,3-diolate (Compound (5))



5

5.1 Preparation of 5-methoxy-2,3,3-trimethyl-3H-indole

To 4-methoxyphenyl hydrazine hydrochloride (4.84g) was added 3-methyl-2-butanone (6.4ml) and acetic acid (45ml). The mixture was heated to 100°C for 2.5 hours after which time the solvent was removed by rotary evaporation. Flash column chromatography gave the product (4.66g).

δ H (270MHz; CDCl₃) 1.3 (6H, s), 2.2 (3H, s), 3.8 (3H, s), 6.8 (2H, m), 7.4 (1H, m).

15

5.2 Preparation of 1-ethyl-5-methoxy-2,3,3-trimethyl-3H-indolium iodide

To 5-methoxy-2,3,3-trimethyl-3H-indole (1.9g) was added iodoethane (5ml) and 1,2-dichlorobenzene (10ml). The mixture was heated to 80°C for 4 hours after which time the mixture was allowed to cool and the precipitate removed by filtration and washed sequentially with dichlorobenzene and diethyl ether. Drying in a vacuum oven gave the product (3g).

δ H (270MHz; CDCl₃) 1.6 (3H, t), 1.6 (6H, s), 3.1 (3H, s), 3.9 (3H, s), 4.7 (2H, q), 7.1 (2H, m), 7.7 (1H, m).

25

5.3 Preparation of 1-(3,5-dinitrobenzyl)-5-methoxy-2,3,3-trimethyl-3H-indolium iodide


To 5-methoxy-2,3,3-trimethyl-3H-indole (1.90g) was added 3,5-dinitrobenzyl iodide (4.62g) and 1,2-dichlorobenzene (10ml). The mixture was heated at 75°C for 3 hours, during which time an orange solid separated. The mixture was then cooled in an ice bath and the solid fraction collected by filtration; it was washed sequentially with dichlorobenzene and diethyl ether and dried under vacuum to give the product (2.62g).
8H (270MHz; DMSO-d₆) 1.6 (6H, s), 2.9 (~3H, s), 3.85 (3H, s), 6.1 (2H, s), 7.1 (1H, dd), 7.55 (1H, d), 7.8 (1H, d), 8.65 (2H, s) and 8.8 (1H, s).

5.4 Preparation of Compound (5)

To 1-(3,5-dinitrobenzyl)-5-methoxy-2,3,3-trimethyl-3H-indolium iodide (250mg) was added 3,4-dihydroxy-3-cyclobuten-1,2-dione (55mg), 1-ethyl-5-methoxy-2,3,3-trimethyl-3H-indolium iodide (175mg), pyridine (2.25ml), acetic acid (2.25ml) and acetic anhydride (0.5ml). The mixture was heated to reflux for 5 hours and the solvent then removed using rotary evaporation. The crude material was partitioned between DCM and 1M HCl. The organic layer was further washed with water. Silica flash column chromatography was performed (DCM/MeOH) and the relevant fractions combined and concentrated. The resulting material was further purified by reverse phase HPLC (CH₃CN / H₂O / TFA).
MALDI-TOF (C₃₇H₃₆N₄O₈ requires M+ 664) 665.

6. Preparation of 2-(1-(5-carboxypentyl)-3,3-dimethyl-5-methoxy-2-indolinylidenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinylidenemethyl)cyclobutenediylum-1,3-diolate (Compound (6))

30



6.1 Preparation of 1-(5-carboxypentyl)-5-methoxy-2,3,3-trimethyl-3H-indolium bromide

5

To 5-methoxy-2,3,3-trimethyl-3H-indole (1.9g) was added 6-bromohexanoic acid (3g) and 1,2-dichlorobenzene (10ml). The mixture was heated to 100°C for 3 hours and then allowed to cool to room temperature. Diethyl ether was added and the precipitated material removed by filtration.

10

Drying in a vacuum oven gave the product (3.12g).

δ H (270MHz; CDCl₃) 1.4 (2H, m), 1.5 (6H, s), 1.6 (2H, m), 1.8 (2H, m), 2.2 (2H, m), 2.8 (3H, s), 3.8 (3H, s), 4.4 (2H, m), 7.1 (1H, m), 7.5 (1H, m), 7.9 (1H, m).

6.2 Preparation of Compound (6)

15


To 1-(3,5-dinitrobenzyl)-5-methoxy-2,3,3-trimethyl-3H-indolium iodide (500mg) (see 5.3) was added 3,4-dihydroxy-3-cyclobuten-1,2-dione (114mg), 1-(5-carboxypentyl)-5-methoxy-2,3,3-trimethyl-3H-indolium bromide (385mg), pyridine (4.5ml), acetic acid (4.5ml) and acetic anhydride (1ml). The mixture was heated to 110°C for 4.5 hours and the solvent then removed using rotary evaporation. This crude material was partitioned between DCM and 1M HCl. The organic layer was further washed with water. Silica flash column chromatography was performed (DCM/MeOH) and the relevant fractions combined and concentrated. The resulting material was further purified by

25

reverse phase HPLC (CH₃CN / H₂O / TFA).

MALDI-TOF (C₄₁H₄₂N₄O₁₀ requires M⁺ 750) 751.

7. Preparation of 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-3-methyl-1,3-dihydro-2H-indol-2-ylidenemethyl-4-((1-(3,5-dinitrobenzyl)-3,3-dimethyl-3H-indolium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound (7))



5 Compound (7)

7.1 Preparation of 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-2,3-dimethyl-3H-indolium bromide

10 To 6-(2,3-dimethyl-3H-indol-3-yl)hexanoic acid (100mg) was added 1-bromo-2-(2-methoxyethoxy)ethane (1ml) and the mixture heated to 90°C overnight. On cooling diethyl ether (10ml) was added and the material removed by filtration.

LCMS ($C_{21}H_{32}NO_4$ requires M^+ 362) 363.

15

7.2 Preparation of Compound (7)

20 To 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-2,3-dimethyl-3H-indolium bromide was added squaric acid (44mg), 1-(3,5-dinitrobenzyl)-2,3,3-trimethyl-3H-indolium iodide (177mg), pyridine (4.5ml), acetic acid (4.5ml) and acetic anhydride (1ml). The mixture was heated to 80°C overnight. On cooling preparative HPLC was performed to give the desired material.

LCMS ($C_{43}H_{46}N_4O_{10}$ requires M^+ 778) 779.

8. Preparation of the acetoxymethyl ester derivative of Compound (7)

To 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-3-methyl-1,3-dihydro-2H-indol-2-ylidenemethyl-4-((1-(3,5-dinitrobenzyl)-3,3-dimethyl-3H-indolium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (14mg) was added acetonitrile (3ml), Hunigs base (32μl) and bromomethyl acetate (9ul). After stirring at room temperature for 2 hours, HPLC was performed to give the desired material (8mg).

LCMS (C₄₆H₆₀N₄O₁₂ requires M⁺ 850) 851.


10

9. Preparation of the ethyl ester derivative of Compound (7)

To ethanol (10ml) was added acetyl chloride (1ml) followed by 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-3-methyl-1,3-dihydro-2H-indol-2-ylidenemethyl-4-((1-(3,5-dinitrobenzyl)-3,3-dimethyl-3H-indolium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (2mg). The mixture was stirred at room temperature for 5 hours, after which time the volatiles were stripped *in vacuo*.

LCMS (C₄₅H₅₀N₄O₁₀ requires M⁺ 806) 807.

20

10. Preparation of -2-((3,3-dimethyl-5-sulfo-1-(4-sulfobutyl)-1,3-dihydro-2H-indol-2-ylidene)methyl)-4-((1-methyl-6-nitroquinolinium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound (8))

25

Compound (8)

10.1 Preparation of 1,2-dimethyl-6-nitroquinolinium iodide

2-Methyl-6-nitroquinoline (0.5g, 2.66mmol) and iodomethane (1ml, 16mmol) were heated together in acetonitrile (10ml) at reflux for 48hrs. The mixture was cooled to room temperature and a grey material crystallised out of solution and was filtered off. This was shown to be starting material. The filtrate was diluted with ethyl acetate (200ml) to give a yellow/green precipitate. The product was filtered off, washed with ethyl acetate and then dried under vacuum. The product was obtained as a yellow/green solid (147mg, 16.8%).

LCMS ($C_{11}H_{11}N_2O_2$ requires M^+ 203) single component M^+ 203.

10.2 Preparation of 2,3,3-trimethyl-5-sulfo-1-(4-sulfobutyl)-3H-indolium potassium salt

2,3,3-Trimethylindolenine-5-sulfonate, potassium salt (6g, 21.6mmol) and 1,4-butanedithione (55ml) were heated together, under nitrogen, at 90°C for 24hrs. On cooling the reaction mixture was diluted with ethyl acetate and the resultant solid filtered off, washed with ethyl acetate and dried under vacuum. The product was isolated as a pale pink solid (10.3g). The product was characterised by 1H NMR (CD_3OD).

10.3 Preparation of Compound (8)

1,2-Dimethyl-6-nitro-quinolinium iodide (100mg, 0.30mmol), 3,4-dihydroxy-3-cyclobutene-1,2-dione (34.5mg, 0.30mmol) and 2,3,3-trimethyl-5-sulfo-1-(4-sulfobutyl)-3H-indolium, potassium salt (124mg, 0.30mmol) were heated together in a mixture of pyridine (3ml), acetic acid (3ml) and acetic anhydride (2ml) at 120°C for 1 hr. The reaction mixture is seen to turn to a dark green/blue colour. On cooling the reaction mixture was poured into ethyl acetate to precipitate the products. The products were filtered off and mixture purified by RP HPLC using eluent mixtures of water/acetonitrile/0.1%TFA. The product was obtained as a dark blue solid (14mg).

LCMS ($C_{30}H_{30}N_3O_{10}S_2$ requires M^+ 656) ES^- gives $(M-H)^{-2}$ reconstruction gives M^- at 654.

11. A Comparative Study of a Nitro-substituted Squaraine Dye (Compound (1)) with Nitro-substituted Cyanine Dyes (Compounds (i) and (iii)) as Substrates in a Nitroreductase Gene Reporter Assay

A reporter construct containing the NF- κ B response element upstream of the NTR gene was constructed in pDC511 (AdmaxTM). The reporter was packaged with Ad5 genomic DNA in helper cells, HEK293, and replication incompetent Adenovirus rescued.

HeLa cells were subcultured for twenty-four hours prior to viral transduction and incubated overnight at 37°C in a humidified atmosphere of 5% CO₂ in Dulbecco's Modified Eagles medium containing 10% foetal calf serum + 2mM L-glutamine. After the overnight incubation, the cells were detached from each flask with trypsin, pooled to produce a suspension of cells and the cell concentration determined. The HeLa cell suspension was mixed with virus at a predetermined multiplicity of infection (MOI) in a sufficient minimal volume of complete medium to cover the base of a tissue culture flask; typically 15ml for 10⁶ cells in a T75cm² Costar flask. The cell/virus suspension was returned to the incubator and left overnight at 37°C in a humidified atmosphere of 5% CO₂. The following day the medium was removed from each flask and the cell monolayer rinsed with 5-10ml PBS. The cells were detached with trypsin and pooled to produce a suspension of transduced cells; the concentration of the cell suspension was determined and adjusted to 5.0 x 10⁴ cells per ml. 200 μ l of this cell suspension was dispensed into each well of a 96-well microtitre plate; \approx 10⁴ cells per well. All plates were incubated overnight at 37°C in a humidified atmosphere of 5% CO₂. The overnight medium was replaced with 200 μ l PBS. The PBS was removed from each well and replaced with TNF α agonist in serum free Dulbecco's Modified Eagles medium (100ng/ml, 90 μ l) or control (90 μ l serum free medium) was added to replicate wells. Plates were returned to the incubator at 37°C in an atmosphere of 5% CO₂ for 2 hours.

After this time, 10 μ l of a 10 μ M solutions of Compounds (i) and (iii) (nitro-cyanine dyes) and Compound (1) (nitro-squaraine dye) were dispensed individually into replicate wells and plates returned to 37°C in a humidified atmosphere of 5% CO₂. The fluorescence signal was monitored over time by
5 means of a Tecan "Ultra" fluorimeter. All substrates were measured under identical conditions to avoid instrument artefacts.

Figure 2 compares the performance of a nitro group-containing squaraine dye (Compound (1) with nitro group-containing cyanine dyes
10 (Compounds (i) and (iii)). The signal to background ratio for Compound (1) was 3:1 (compared with 1.3:1 for Compound (i), clearly demonstrating a reduction in background fluorescence combined with a similar increase in fluorescence signal.

15 12. A Comparative Study of Nitro-substituted Squaraine Dyes (Compounds (1) and (2)) as Substrates in a Nitroreductase Gene Reporter Assay

Using the same methodology as that described in Example 11, HeLa cells were transduced with the adenoviral NF- κ B reporter system. At the
20 appropriate time Compounds (1) and (2) were added individually to replicate wells. The fluorescence signal was monitored over time on a Tecan Ultra fluorimeter and data presented in Figure 3.

Compound (2) clearly shows a significant increase in assay signal when
25 compared to Compound (1) in the presence of the agonist, TNF- α . Compound (2) was also capable of detecting basal transcriptional activity in the control sample cells containing the reporter. This basal activity is the difference in signal between cells containing the NF- κ B reporter but no agonist and mock transduced cells. Compound (1) was not sensitive enough to detect this low
30 level activity. The improvements in assay sensitivity are believed to be a direct consequence of the availability of the improved compounds within the cell. Thus, Compound (2) is thought to be available in the cell cytoplasm which is also the same compartment as the expressed reporter protein. Microscopic

imaging of the assay plates have produced cells with intense red cytoplasm staining following NTR expression.

5 The data from Figures 2 and 3 illustrate that the improved properties of the nitro-substituted squaraine dye (Compound (2)) are a result of introducing both the squarylium moiety into the dye and the addition of the hexanoic acid group. The presence of the hexanoic acid group in Compound (iii), nitro-substituted cyanine dye, was not sufficient to alter the cellular localisation of Compound (i), nor its performance in the NTR assay.

10

13. Localisation of Compound (2) in HeLa Cells

HeLa cells were plated at 120,000 per dish and incubated overnight at 37°C. in Dulbecco's Modified Eagles medium containing 10% foetal calf serum + 2Mm L-glutamine. Overnight medium was removed and replaced with 2ml of serum free medium containing 1µM Compound (2). Dishes were returned to the incubator for 2 hours before imaging on a Zeiss Confocal Microscope.

20 Figure 4 shows the uptake and distribution of Compound (2) in HeLa cells. There was no evidence to indicate that Compound (2) was sequestered to organelles. Although there is evidence of background labelling of cell structures, this does not compromise the assay performance.

25 14. Evaluation of Compounds (2), (3) and (4) as Nitroreductase Substrates in live cell NTR assays

Further examples of the utility of nitro-substituted squaraine dyes (Compounds (2), (3) and (4)) as nitroreductase substrates are shown in live cell NTR assays, Figure 5. The data presented in Figure 5 for Compound (3) shows that a squaraine dye substrates that emits at longer wavelength may be obtained by extending the conjugation system of the dye. The presence of the hexanoic group increases the retention properties of the probe within the cell as demonstrated by the fixation of cells. Compounds (2) and (3) show very little

decrease in signal following fixation while Compound (4) has lost almost 50% of the signal post-fixation.

15. Example of shift in absorption maximum after action of NTR upon a substrate

Compound (8) in DMSO (1mMol) (4 μ l) was diluted into PBS buffer (0.01M) (1.76ml). The UV/Vis spectrum of the solution was measured. The substrate has an absorbance maximum at 682nm AU= 0.22. To the solution was added NADH (0.01M in PBS) (200 μ l) and NTR enzyme (446ng/ml, 37 μ l) the mixture was leave to stand at room temperature for 30 mins. After this time, the absorbance spectrum was re-measured. A new absorbance maximum at 621nm, AU=0.17 is observed.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

30

14 Mar 2007


2005250209

2005250209 16 Jun 2009

-33-

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for detecting nitroreductase enzyme activity in a composition comprising:
 - 5 i) mixing said composition under conditions to promote nitroreductase activity with a dye molecule; and
 - ii) measuring a change in an optical property of said dye molecule wherein said change is a measure of the amount of nitroreductase activity; wherein said dye molecule is a squaraine dye comprising at least one NO₂ group.
- 10 2. A method according to claim 1, wherein said composition comprises at least one cell or a cell extract.
3. A method according to claim 1 or claim 2, wherein said method is
15 conducted in the presence of a test agent whose effect on nitroreductase enzyme activity is to be determined.
4. A method which comprises:
 - 20 i) contacting a host cell with a dye molecule wherein said host cell has been transfected with a nucleic acid molecule comprising expression control sequences operably linked to a sequence encoding a nitroreductase; and
 - ii) measuring a change in an optical property of said dye molecule wherein said change is a measure of the amount of nitroreductase activity; wherein said dye is a squaraine dye comprising at least one NO₂ group.
- 25 5. A method according to any one of claims 1 to 4, wherein said change in an optical property is an increase in the fluorescence intensity of the dye molecule, whereby said increase is a measure of the amount of nitroreductase activity.
- 30 6. A method according to any one of claims 1 to 5, wherein said squaraine dye is of formula (I):



(I)

wherein:

- 5 R^3 is attached to the Z^1 ring structure and R^4 is attached to the Z^2 ring structure;
 Z^1 and Z^2 independently represent a phenyl or a naphthyl ring system;
 X and Y are the same or different and are selected from oxygen, sulphur, $-\text{CH}=\text{CH}-$ and the group:



10

groups R^1 and R^2 are independently selected from $\text{C}_1\text{-C}_4$ alkyl, $-(\text{CH}_2)_n\text{-P}$, $-\{(\text{CH}_2)_2\text{-O}\}_p\text{-R}^6$ and the group W; where P is selected from COOR^7 , SO_3^- and OH, W is mono- or di-substituted nitrobenzyl, R^6 is methyl or ethyl, R^7 is selected from H, $\text{C}_1\text{-C}_4$ alkyl and $\text{CH}_2\text{OC(O)R}^8$, where R^8 is methyl, or t-butyl, n

15

is an integer from 1 to 10, and p is an integer from 1 to 3;
 groups R^3 and R^4 are independently selected from hydrogen, NO_2 , halogen, SO_3^- , $\text{C}_1\text{-C}_4$ alkoxy and $-(\text{CH}_2)_m\text{-COOR}^7$; where R^7 is hereinbefore defined and m is 0 or an integer from 1 to 5;

R^5 is $\text{C}_1\text{-C}_6$ alkyl optionally substituted with COOR^7 , SO_3^- , or OH; where R^7 is

20




hereinbefore defined; and
 at least one of groups R^1 , R^2 , R^3 and R^4 comprises at least one NO_2 group.

7. A method according to any one of claims 4 to 6, wherein said dye is, or is rendered, cell permeable.

25

8. A method according to any one of claims 1 to 7, wherein said squaraine dye is a compound selected from dyes of formula:

2005250209 14 Mar 2007



wherein:

X and Y are the same or different and are selected from oxygen, sulphur, -CH=CH- and the group:



5

wherein R⁵ is hereinbefore defined;

at least one of groups R¹ and R² is the group W; where W is hereinbefore defined;

any remaining group R¹ or R² is selected from C₁-C₄ alkyl, -(CH₂)_n-P and

10 -{(CH₂)₂-O}_p-R⁶; where P is selected from COOR⁷, SO₃⁻ and OH, R⁶ is methyl

or ethyl, R⁷ is selected from H, C₁-C₄ alkyl and CH₂OC(O)R⁸, where R⁸ is


methyl, or t-butyl, n is an integer from 1 to 10 and p is an integer from 1 to 3;

and

2005250209 16 Jun 2009

groups R³ and R⁴ are independently selected from hydrogen, halogen, SO₃⁻, C₁-C₄ alkoxy and -(CH₂)_m-COOR⁷; where R⁷ is hereinbefore defined and m is 0 or an integer from 1 to 5.


- 5 9. A method according to claim 8, wherein one of groups R¹ and R² is selected from group W where W is selected from:



remaining R¹ or R² is selected from methyl and ethyl, or is the group -(CH₂)_n-COOR⁷ where R⁷ is selected from H, C₁-C₄ alkyl and CH₂OC(O)R⁸,

- 10 where R⁸ is methyl, or t-butyl, n is an integer from 1 to 10, preferably 5 or 6.


10. A method according to claim 9, wherein W is the group:



and remaining R¹ or R² is hereinbefore defined.



15

11. A method according to any one of claims 1 to 7, wherein said squaraine dye is a compound of formula:



16 Jun 2009

2005250209



wherein:

X and Y are the same or different and are selected from oxygen, sulphur, $-\text{CH}=\text{CH}-$ and the group:



5

wherein R^5 is hereinbefore defined;

groups R^1 or R^2 are independently selected from $\text{C}_1\text{-C}_4$ alkyl, $-(\text{CH}_2)_n\text{-P}$ and $-\{(\text{CH}_2)_2\text{-O}\}_p\text{-R}^6$; where P is selected from COOR^7 , SO_3^- and OH, R^6 is methyl or ethyl, R^7 is selected from H, $\text{C}_1\text{-C}_4$ alkyl and $\text{CH}_2\text{OC}(\text{O})\text{R}^8$, where R^8 is

10 methyl, or t-butyl, n is an integer from 1 to 10 and p is an integer from 1 to 3; at least one of groups R^3 and R^4 is NO_2 ; and

any remaining group R^3 or R^4 is selected from hydrogen, SO_3^- , $\text{C}_1\text{-C}_4$ alkoxy and $-(\text{CH}_2)_m\text{-COOR}^7$; where R^7 is selected from H, $\text{C}_1\text{-C}_4$ alkyl and $\text{CH}_2\text{OC}(\text{O})\text{R}^8$, where R^8 is methyl, or t-butyl, and m is 0 or an integer from 1 to

15 5.

12. A method for screening for a test agent whose effect upon nitroreductase gene expression is to be determined, said method comprising:

a) performing the method according to any one of claims 4 to 11 in the

20 absence and in the presence of said test agent; and

16 Jun 2009

2005250209

5 b) determining the amount of nitroreductase gene expression in the absence and in the presence of said agent; wherein a difference between the amount of nitroreductase gene expression in the absence and in the presence of said agent is indicative of the effect of said agent on nitroreductase gene expression.



10 13. A method of screening for a test agent whose effect upon nitroreductase gene expression is to be determined, said method comprising:

a) performing the method according to any one of claims 4 to 11 in the presence of said agent; and

b) comparing the amount of nitroreductase gene expression with a control value for the amount of nitroreductase gene expression in the absence of the test agent.


15 14. A method according to claim 13, wherein said control value is stored electronically in a database or other electronic format.

15. A compound selected from dyes of formula:



14 Mar 2007

2005250209




wherein:

X and Y are the same or different and are selected from oxygen, sulphur, $-\text{CH}=\text{CH}-$ and the group:



- 5 one of groups R^1 and R^2 is the group:



remaining R^1 or R^2 is selected from methyl, ethyl and the group $-(\text{CH}_2)_n-\text{COOR}^7$; where R^7 is selected from H, C_1 - C_4 alkyl and $\text{CH}_2\text{OC}(\text{O})\text{R}^8$, where R^8 is methyl, or t-butyl and n is an integer from 1 to 10;

- 10 groups R^3 and R^4 are independently selected from hydrogen, NO_2 , halogen, SO_3 , C_1 - C_4 alkoxy and $-(\text{CH}_2)_m-\text{COOR}^7$; where R^7 is selected from H, C_1 - C_4 alkyl and $\text{CH}_2\text{OC}(\text{O})\text{R}^8$, where R^8 is methyl, or t-butyl, and m is 0 or an integer from 1 to 5;

- 15 R^5 is C_1 - C_6 alkyl optionally substituted with COOR^7 , SO_3 , or OH; where R^7 is hereinbefore defined.

16. A compound according to claim 15, wherein said remaining R^1 or R^2 is selected from the group $-(\text{CH}_2)_5-\text{COOR}^7$ and $-(\text{CH}_2)_6-\text{COOR}^7$; where R^7 is hereinbefore defined.

20

17. A compound selected from:

2005250209 16 Jun 2009

- a) 2-(1-methyl-3,3-dimethyl-2-indolinyldenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 1);
- b) 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-indolinyldenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 2);
- c) 2-(1-(5-carboxypentyl)-3,3-dimethyl-2-benzindolinyldenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 3);
- d) 2-(1-ethyl-3,3-dimethyl-5-methoxy-2-indolinyldenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 5);
- e) 2-(1-(5-carboxypentyl)-3,3-dimethyl-5-methoxy-2-indolinyldenemethyl)-4-(1-(3,5-dinitrobenzyl)-3,3-dimethyl-5-methoxy-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 6); and
- f) 3-(5-carboxypentyl)-1-(2-(2-methoxyethoxy)ethyl)-3-methyl-1,3-dihydro-2H-indol-2-ylidenemethyl-4-((1-(3,5-dinitrobenzyl)-3,3-dimethyl-3H-indolium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound 7).
18. A compound selected from:
- a) 2-(3-ethyl-6-nitro-2-benzothiazolinyldenemethyl)-4-(1-(2-(2-methoxyethoxy)ethyl)-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (Compound 4); and
- b) 2-((3,3-dimethyl-5-sulfo-1-(4-sulfobutyl)-1,3-dihydro-2H-indol-2-ylidene)methyl)-4-((1-methyl-6-nitroquinolinium-2-yl)methylene)-3-oxocyclobut-1-en-1-olate (Compound (8)).
19. Use of a compound according to any one of claims 15 to 18 as a substrate for the detection and/or measurement of nitroreductase enzyme activity.
20. Use according to claim 19, wherein said compound is administered *in vivo* and nitroreductase enzyme activity determined by optical imaging.


2005250209 16 Jun 2009

-41-

21. A method according to any one of claims 1 to 14, a compound according to any one of claims 15 to 18, the use of a compound according to claim 19 or claim 20 substantially as hereinbefore described with reference to the Figures and/or Examples.


5

1a



Compound (i)


1b




Compound (iii)

Comparison of two nitro-substituted cyanine type dyes, Compounds (i) and (iii) compared with a nitro-substituted squaraine dye (Compound (1))

2a. Fluorescence emission




2b. Signal:Background




A Comparative Study of Nitro-substituted Squaraine Dyes (Compounds (1) and 2) as Substrates in a Nitroreductase Gene Reporter Assay

3a. Fluorescence emission



3b. Signal:Background




Distribution of Compound (2) in HeLa cells



Evaluation of Compounds (2), (3) and (4) as Nitroreductase Substrates
in live cell NTR assays

5a. Fluorescence emission



5b. Signal:Background

