

(19) **DANMARK**

(10) **DK/EP 2917210 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 D 405/14 (2006.01)** **A 61 K 31/558 (2006.01)** **A 61 P 35/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2019-05-27**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-02-27**
- (86) Europæisk ansøgning nr.: **13853072.0**
- (86) Europæisk indleveringsdag: **2013-11-05**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-09-16**
- (86) International ansøgning nr.: **CN2013086571**
- (87) Internationalt publikationsnr.: **WO2014071832**
- (30) Prioritet: **2012-11-06 US 201261723099 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Shanghai Fochon Pharmaceutical Co. Ltd, Room 512, Building A, No. 1289 Yishan Road, Shanghai 200233, Kina**
Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 555 Zu Chong Zhi Road , Zhangjiang Hi-Tech Park , Pudong New District, Shanghai 201203, Kina
- (72) Opfinder: **WANG, Weibo, 62 Sanders Ranch Road, Moraga, California 94556, USA**
GENG, Meiyu, No. 555 Zu Chong Zhi Road, Zhangjiang Hi-Tech Park, Pudong, Shanghai 201203, Kina
DING, Jian, No. 555 Zu Chong Zhi Road, Zhang Jiang Hi-Tech Park, Pudong, Shanghai 201203, Kina
AI, Jing, No. 555 Zu Chong Zhi Road, Zhang Jiang Hi-Tech Park, Pudong, Shanghai 201203, Kina
TIAN, Qiang, No.565 Tushan Road, Chongqing 400061, Kina
PENG, Xia, No. 555 Zu Chong Zhi Road, Zhang Jiang Hi-Tech Park, Pudong, Shanghai 201203, Kina
ZHANG, Weipeng, No.565 Tushan Road, Chongqing 400061, Kina
LIU, Hongbin, No. 565 Tushan Road, Chongqing 400061, Kina
TAN, Haohan, No. 565 Tushan Road, Chongqing 400061, Kina
CHEN, Ling, No. 565 Tushan Road, Chongqing 400061, Kina
ZHAO, Xingdong, No. 565, Tushan Road, Chongqing 400061, Kina
- (74) Fuldmægtig i Danmark: **Larsen & Birkeholm A/S Skandinavisk Patentbureau, Banegårdspladsen 1, 1570 København V, Danmark**
- (54) Benævnelse: **ALK KINASE-INHIBITORER**
- (56) Fremdragne publikationer:
WO-A1-2005/016894
WO-A1-2005/080393
WO-A1-2008/051547
WO-A1-2009/008371
WO-A1-2010/128659

Fortsættes ...

WO-A2-2007/066187

WO-A2-2008/073687


DESCRIPTION

[0001] Disclosed herein are protein kinase inhibitors, more particularly novel pyrimidine derivatives and pharmaceutical compositions thereof, and their use as pharmaceuticals.



[0002] Anaplastic lymphoma kinase (ALK), a member of the insulin receptor superfamily of receptor tyrosine kinases, has been implicated in oncogenesis in hematopoietic and non-hematopoietic tumors. The aberrant expression of full-length ALK receptor proteins has been reported in neuroblastomas and glioblastomas; and ALK fusion proteins have occurred in anaplastic large cell lymphoma. The study of ALK fusion proteins has also raised the possibility of new therapeutic treatments for patients with ALK-positive malignancies. (Pulford et al., Cell. Mol. Life. Sci. 61:2939-2953 (2004)).

[0003] Because of the emerging disease-related roles of ALK, there is a continuing need for compounds which may be useful for treating and preventing a disease which responds to inhibition of ALK and have at least one advantageous property selected from potency, stability, selectivity, toxicity, pharmacodynamics properties and pharmacokinetics properties as an alternative. In this regard, a novel class of ALK inhibitors is provided herein..

[0004] WO/2005/016894 discloses novel pyrimidine derivatives of formula (I), wherein R is selected from C₆₋₁₀ aryl, C₅₋₁₀ heteroaryl, C₃₋₁₂ cycloalkyl and C₃₋₁₀ heterocycloalkyl; R⁰-R⁶ as described herein; and their use for the manufacture of a medicament for the treatment or prevention of a disease which responds to inhibition of FAK and/or ALK and/or ZAP-70 and/or IGF-IR.




[0005] WO/2008/051547 provides a compound of formula I or II or a pharmaceutically acceptable salt form thereof, wherein R¹, R², R³, R⁴, R⁵, A¹, A², A³, A⁴, and A⁵, are as defined herein. The compounds of formula I or II have ALK and/or c-Met inhibitory activity, and may be used to treat proliferative disorders.



[0006] Disclosed herein are certain novel pyrimidine derivatives and pharmaceutical compositions thereof, and their use as pharmaceuticals.

[0007] In one aspect, the present invention provides at least one compound of formula (I):



and/or at least one pharmaceutically acceptable salt thereof, wherein:

each R¹ is independently selected from:

hydrogen,

halogen,

hydroxyl,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkyl-alkyl,

heterocyclyl,

heterocyclylalkyl

aryl,

arylalkyl,

heteroaryl, and

heteroarylalkyl,

wherein alkyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a}, and wherein aryl and heteroaryl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6b};

each R² is independently selected from: C₁₋₁₀ alkyl and piperidinyl, wherein the C₁₋₁₀ alkyl and piperidinyl are each independently unsubstituted or substituted with at least one substituent,

such as one, two, three or four substituents, independently selected from R^{6a} ;

each R^3 is independently selected from:

hydrogen,

halogen,

-CN,

- NR^7R^8 , and

C_{1-10} alkyl;

wherein alkyl is unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} ;

each R^4 is independently selected from:

hydrogen,

halogen,

-CN,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl, and

C_{3-10} cycloalkyl;

wherein C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, and C_{3-10} cycloalkyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} ;

or R^3 and R^4 together with the carbon atoms to which they are attached form a 5-6 membered ring containing 0, 1, 2 or 3 heteroatoms independently selected from oxygen, sulfur and nitrogen, and optionally substituted with 1-2 R^{6b} groups;

each R^5 is independently selected from:

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-O(CR⁹R¹⁰)_tNR⁷R⁸,

-C(O)R⁷,

-CO₂R⁸,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸,

-OC(O)R⁷,

-CN,

-C(O)NR⁷R⁸,

-NR⁷C(O)R⁸,

-OC(O)NR⁷R⁸,

-NR⁷C(O)OR⁸,

-NR⁷C(O)NR⁷R⁸,

-CR⁷(N-OR⁸),

-CHF₂,

-CF₃,

-OCHF₂, and

-OCF₃;

wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and C₃₋₁₀ cycloalkyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a};

each R^{6a} is independently selected from:

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,


















-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-(CR⁹R¹⁰)_tOR⁸,

-(CR⁹R¹⁰)_tNR⁷R⁸,



$-\text{NR}^7\text{C}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{CR}^7(\text{N}-\text{OR}^8)$,

$-\text{CHF}_2$,

$-\text{CF}_3$,

$-\text{OCHF}_2$, and

$-\text{OCF}_3$;

each R^{6b} is independently selected from:

R^{6a} ,

aryl,

aryl- C_{1-4} alkyl,

heteroaryl, and

heteroaryl- C_{1-4} alkyl;

each R^7 and each R^8 are independently selected from:

hydrogen,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C_{3-10} cycloalkyl,

cycloalkyl- C_{1-10} alkyl;

heterocyclyl,

heterocyclyl- C_{1-10} alkyl,

aryl,

heteroaryl,

aryl- C_{1-10} alkyl, and

heteroaryl- C_{1-10} alkyl;

wherein alkyl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} , and aryl and heteroaryl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6b} ; or

R^7 and R^8 together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0, 1, or 2 additional heteroatoms independently selected from oxygen, sulfur and NR^{11} ,

each R^7 and R^8 may be unsubstituted or substituted on a carbon or nitrogen atom with at least one substituent, such as one, two, or three substituents, independently selected from R^{12} ;

each R^9 and each R^{10} are independently selected from:

hydrogen,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C_{3-10} cycloalkyl,

cycloalkyl- C_{1-10} alkyl,

heterocyclyl,

heterocyclyl- C_{1-10} alkyl,

aryl,

heteroaryl,

aryl- C_{1-10} alkyl, and

heteroaryl- C_{1-10} alkyl; or

R^9 and R^{10} together with the carbon to which they are attached form a ring of 3 to 7 members containing 0, 1, or 2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

each R^{11} is independently selected from:

hydrogen,

C_{1-10} alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl,

heterocyclyl,

heterocyclyl-C₁₋₄ alkyl,

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl,

heteroaryl-C₁₋₄ alkyl,

-S(O)_rR⁷,

-C(O)R⁷,

-CO₂R⁷,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸, and

-C(O)NR⁷R⁸;

each R¹² is independently selected from:

halogen,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkylalkyl,

heterocyclyl,

heterocyclylalkyl,

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl,

heteroaryl-C₁₋₄ alkyl,

-OR⁷,

-NR⁷S(O)_rR⁸,

-S(O)_rR⁷,

-SR⁷,

-S(O)₂OR⁷,

-OS(O)₂R⁷,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-O(CR⁹R¹⁰)_tNR⁷R⁸,

-C(O)R⁷,

-CO₂R⁸,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸,

-OC(O)R⁷,

-CN,

-C(O)NR⁷R⁸,

-NR⁷C(O)R⁸,

-OC(O)NR⁷R⁸,

-NR⁷C(O)OR⁸,

-NR⁷C(O)NR⁷R⁸,

-CHF₂,

-CF₃,

-OCHF₂, and

-OCF₃;

each m is independently selected from 0, 1 and 2;

each n is independently selected from 1, 2, and 3;

p is 2;

each q is independently selected from 0, 1, 2, and 3;

each r is independently selected from 1 and 2;

each t is independently selected from 1, 2, and 3.

[0008] In yet another aspect, the present invention provides pharmaceutical compositions comprising at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof and a pharmaceutically acceptable excipient.

[0009] The present disclosure enables methods for modulating ALK, comprising administering to a system or a subject in need thereof, a therapeutically effective amount of at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof or pharmaceutical compositions thereof, thereby modulating said ALK. The present disclosure also enables at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof or pharmaceutical compositions thereof, and optionally in combination with a second therapeutic agent, for use in treating, ameliorating or preventing a condition which responds to inhibition of ALK. The present disclosure also enables the use of at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a condition mediated by ALK. In particular embodiments, the compounds of the invention may be used alone or in combination with a second therapeutic agent to treat a condition mediated by ALK, wherein said condition is an autoimmune disease, a transplantation disease, an infectious disease or a cell proliferative disorder.

[0010] Furthermore, the invention provides at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof or pharmaceutical compositions thereof, and optionally in combination with a second therapeutic agent, for use in for treating a cell proliferative disorder.

[0011] The present disclosure also enables the use of at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a cell-proliferative disorder. In particular examples, the compounds of the disclosure may be used alone or in combination with a chemotherapeutic agent to treat a cell proliferative disorder, including but not limited to, lymphoma, osteosarcoma, melanoma, or a tumor of breast, renal, prostate, colorectal, thyroid, ovarian, pancreatic, neuronal, lung, uterine or gastrointestinal tumor.

[0012] The present disclosure also enables at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof to be administered to a system comprising cells or tissues, or to a mammalian subject such as a human or animal subject.

BRIEF DESCRIPTION OF THE DRAWING

[0013] Figure 1. Effect of a compound disclosed herein on relative tumor volume.

[0014] As used herein the following definitions are applicable.

[0015] The term "alkyl" refers to both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Unless otherwise specified, "alkyl" refers to C₁-C₁₀ alkyl. For example, C₁-C₁₀, as in "C₁₋₁₀ alkyl" is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, "C₁₋₁₀ alkyl" includes but is not limited to methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, i-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and decyl.

[0016] The term "cycloalkyl" means a saturated aliphatic cyclic hydrocarbon group having the specified number of carbon atoms. Unless otherwise specified, "cycloalkyl" refers to C₃₋₁₀ cycloalkyl. For example, "cycloalkyl" includes but is not limited to cyclopropyl, methylcyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, and cyclohexyl.

[0017] The term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. In some embodiments, one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Thus, "C₂₋₁₀ alkenyl" means an alkenyl radical having from 2 to 10 carbon atoms. Alkenyl groups include but are not limited to ethenyl, propenyl, butenyl, 2-methylbutenyl and cyclohexenyl. The straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

[0018] The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. In some embodiments, up to three carbon-carbon triple bonds may be present. Thus, "C₂₋₁₀ alkynyl" means an alkynyl radical having from 2 to 10 carbon atoms. Alkynyl groups include but are not limited to ethynyl, propynyl, butynyl, and 3-methylbutynyl. The straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.

[0019] The term "alkoxy" refers to either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge such as, for example, methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentyloxy, isopentoxy,

neopentoxo, hexoxy, 2-hexoxy, 3-hexoxy, 3-methylpentoxy, cyclopropanyloxy, and cyclobutyloxy. "Alkoxy" therefore encompasses the definitions of alkyl and cycloalkyl above.

[0020] The term "aryl" encompasses: 5- and 6-membered carbocyclic aromatic rings, for example, benzene; bicyclic ring systems wherein at least one ring is carbocyclic and aromatic, for example, naphthalene, indane, and 1,2,3,4-tetrahydroquinoline; and tricyclic ring systems wherein at least one ring is carbocyclic and aromatic, for example, fluorene. In cases where the aryl substituent is bicyclic or tricyclic and at least one ring is non-aromatic, it is understood that attachment is via the aromatic ring.

[0021] For example, aryl includes 5- and 6-membered carbocyclic aromatic rings fused to a 5- to 7-membered heterocyclic ring containing one or more heteroatoms selected from N, O, and S, provided that the point of attachment is at the carbocyclic aromatic ring. Bivalent radicals formed from substituted benzene derivatives and having the free valences at ring atoms are named as substituted phenylene radicals. Bivalent radicals derived from univalent polycyclic hydrocarbon radicals whose names end in "-yl" by removal of one hydrogen atom from the carbon atom with the free valence are named by adding "-idene" to the name of the corresponding univalent radical, e.g., a naphthyl group with two points of attachment is termed naphthylidene. Aryl, however, does not encompass or overlap in any way with heteroaryl, separately defined below. Hence, if one or more carbocyclic aromatic rings are fused with a heterocyclic aromatic ring, the resulting ring system is heteroaryl, not aryl, as defined herein.

[0022] The term "halogen" (or "halo") refers to fluorine, chlorine, bromine and iodine.

[0023] The term "heteroaryl" refers to

5- to 8-membered aromatic, monocyclic rings containing one or more, for example, from 1 to 4, or, in some embodiments, from 1 to 3, heteroatoms selected from N, O, and S, with the remaining ring atoms being carbon;

8- to 12-membered bicyclic rings containing one or more, for example, from 1 to 4, or, in some embodiments, from 1 to 3, heteroatoms selected from N, O, and S, with the remaining ring atoms being carbon and wherein at least one heteroatom is present in an aromatic ring; and

11- to 14-membered tricyclic rings containing one or more, for example, from 1 to 4, or in some embodiments, from 1 to 3, heteroatoms selected from N, O, and S, with the remaining ring atoms being carbon and wherein at least one heteroatom is present in an aromatic ring.

[0024] When the total number of S and O atoms in the heteroaryl group exceeds 1, those heteroatoms are not adjacent to one another. In some embodiments, the total number of S and O atoms in the heteroaryl group is not more than 2. In some embodiments, the total number of S and O atoms in the heteroaryl group is not more than 1.

[0025] Examples of heteroaryl groups include, but are not limited to, (as numbered from the linkage position assigned priority 1), 2-pyridyl, 3-pyridyl, 4-pyridyl, 2,3-pyrazinyl, 3,4-pyrazinyl, 2,4-pyrimidinyl, 3,5-pyrimidinyl, 1-pyrazolyl, 2,3-pyrazolyl, 2,4-imidazolyl, isoxazolyl, oxazolyl, thiazolyl, thiadiazolyl, tetrazolyl, thienyl, benzothienyl, furyl, benzofuryl, benzoimidazolyl,

indolinyl, pyridizinyl, triazolyl, quinolinyl, pyrazolyl, and 5,6,7,8-tetrahydroisoquinoline.

[0026] Further heteroaryl groups include but are not limited to pyrrolyl, isothiazolyl, triazinyl, pyrazinyl, pyridazinyl, indolyl, benzotriazolyl, quinoxalinyl, and isoquinolinyl. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heteroaryl.

[0027] Bivalent radicals derived from univalent heteroaryl radicals whose names end in "-yl" by removal of one hydrogen atom from the atom with the free valence are named by adding "-idene" to the name of the corresponding univalent radical, e.g., a pyridyl group with two points of attachment is a pyridylidene. Heteroaryl does not encompass or overlap with aryl as defined above.

[0028] In cases where the heteroaryl substituent is bicyclic or tricyclic and at least one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively.

[0029] The term "heterocycle" (and variations thereof such as "heterocyclic", or "heterocyclyl") broadly refers to a single aliphatic ring, usually with 3 to 7 ring atoms, containing at least 2 carbon atoms in addition to 1-3 heteroatoms independently selected from oxygen, sulfur, and nitrogen, as well as combinations comprising at least one of the foregoing heteroatoms. "Heterocycle" also refers to 5- to 7-membered heterocyclic ring containing one or more heteroatoms selected from N, O, and S fused with 5- and 6-membered carbocyclic aromatic ring, provided that the point of attachment is at the heterocyclic ring. The rings may be saturated or have one or more double bonds (i.e. partially unsaturated). The heterocycle can be substituted by oxo. The point of the attachment may be carbon or heteroatom in the heterocyclic ring, provided that attachment results in the creation of a stable structure. When the heterocyclic ring has substituents, it is understood that the substituents may be attached to any atom in the ring, whether a heteroatom or a carbon atom, provided that a stable chemical structure results. Heterocycle does not overlap with heteroaryl.

[0030] Suitable heterocycles include, for example (as numbered from the linkage position assigned priority 1), 1-pyrrolidinyl, 2-pyrrolidinyl, 2,4-imidazolidinyl, 2,3-pyrazolidinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-piperazinyl, 2-piperazinyl, and 2,5-piperazinyl. Morpholinyl groups are also contemplated, including 2-morpholinyl and 3-morpholinyl (numbered wherein the oxygen is assigned priority 1). Substituted heterocycle also includes ring systems substituted with one or more oxo moieties, such as piperidinyl N-oxide, morpholinyl-N-oxide, 1-oxo-1-thiomorpholinyl and 1,1-dioxo-1-thiomorpholinyl.

[0031] As used herein, "arylalkyl" refers to an alkyl moiety substituted by an aryl group. Example arylalkyl groups include benzyl, phenethyl, and naphthylmethyl groups. In some embodiments, arylalkyl groups have from 7 to 20 or 7 to 11 carbon atoms. When used in the phrase "aryl-C₁₋₄ alkyl", the term "C₁₋₄" refers to the alkyl portion of the moiety and does not describe the number of atoms in the aryl portion of the moiety. Likewise, when used in the

phrase "aryl-C₁₋₁₀ alkyl", the term "C₁₋₁₀" refers to the alkyl portion of the moiety and does not describe the number of atoms in the aryl portion of the moiety.

[0032] As used herein, "heterocyclalkyl" refers to alkyl substituted by heterocycl. When used in the phrase "heterocycl-C₁₋₁₀ alkyl", the term "C₁₋₁₀" refers to the alkyl portion of the moiety and does not describe the number of atoms in the heterocycl portion of the moiety.

[0033] As used herein, "cycloalkylalkyl" refers to alkyl substituted by cycloalkyl. When used in the phrase "C₃₋₁₀ cycloalkylalkyl", the term "C₃₋₁₀" refers to the cycloalkyl portion of the moiety and does not describe the number of atoms in the alkyl portion of the moiety. When used in the phrase "C₃₋₇ cycloalkylalkyl", the term "C₃₋₇" refers to the cycloalkyl portion of the moiety and does not describe the number of atoms in the alkyl portion of the moiety. When used in the phrase "C₃₋₈ cycloalkylalkyl", the term "C₃₋₈" refers to the cycloalkyl portion of the moiety and does not describe the number of atoms in the alkyl portion of the moiety. When used in the phrase "cycloalkyl -C₁₋₁₀ alkyl", the term "C₁₋₁₀" refers to the alkyl portion of the moiety and does not describe the number of atoms in the cycloalkyl portion of the moiety.

[0034] As used herein, "heteroarylalkyl" refers to alkyl substituted by heteroaryl. When used in the phrase "heteroaryl -C₁₋₄ alkyl", the term "C₁₋₄" refers to the alkyl portion of the moiety and does not describe the number of atoms in the heteroaryl portion of the moiety. Likewise, when used in the phrase "heteroaryl -C₁₋₁₀ alkyl", the term "C₁₋₁₀" refers to the alkyl portion of the moiety and does not describe the number of atoms in the heteroaryl portion of the moiety.

[0035] For avoidance of doubt, reference, for example, to substitution of alkyl, cycloalkyl, heterocycl, aryl, and/or heteroaryl refers to substitution of each of those groups individually as well as to substitutions of combinations of those groups. That is, if R¹ is arylalkyl, the aryl portion may be unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6b} and the alkyl portion may also be unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a}.

[0036] The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. Salts derived from inorganic bases may be selected, for example, from aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, and zinc salts. Further, for example, the pharmaceutically acceptable salts derived from inorganic bases may be selected from ammonium, calcium, magnesium, potassium, and sodium salts. Salts in the solid form may exist in one or more crystal structures, and may also be in the form of hydrates. Salts derived from pharmaceutically acceptable organic non-toxic bases may be selected, for example, from salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine,

caffeine, choline, N,N'-dibenzylethylene-diamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, and tripropylamine, tromethamine.

[0037] When the compound disclosed herein is basic, salts may be prepared using at least one pharmaceutically acceptable non-toxic acid, selected from inorganic and organic acids. Such acid may be selected, for example, from acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, and p-toluenesulfonic acids. In some embodiments, such acid may be selected, for example, from citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, fumaric, and tartaric acids.

[0038] The terms "co-administration" or "combined administration" or the like as used herein are meant to encompass administration of the selected therapeutic agents to a single patient, and are intended to include treatment regimens in which the agents are not necessarily administered by the same route of administration or at the same time.

[0039] The term "pharmaceutical combination" as used herein refers to a product obtained from mixing or combining active ingredients, and includes both fixed and non-fixed combinations of the active ingredients.

[0040] The term "fixed combination" means that the active ingredients, e.g. a compound of Formula (I) and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage.

[0041] The term "non-fixed combination" means that the active ingredients, e.g. a compound of Formula (I) and a co-agent, are both administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific time limits, wherein such administration provides therapeutically effective levels of the active ingredients in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[0042] The term "therapeutically effective amount" means the amount of the subject compound that will elicit a biological or medical response in a cell, tissue, organ, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

[0043] The term "administration" or "administering" of the subject compound means providing a compound of the disclosure and prodrugs thereof to a subject in need of treatment.


[0044] The term "composition" as used herein is intended to encompass a product comprising

the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. Such term in relation to a pharmaceutical composition is intended to encompass a product comprising the active ingredient (s), and the inert ingredient (s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients.

[0045] By "pharmaceutically acceptable" it is meant compatible with the other ingredients of the formulation and not unacceptably deleterious to the recipient thereof.

[0046] The term "protecting group" or "Pg" refers to a substituent that can be commonly employed to block or protect certain functionality while reacting other functional groups on the compound. For example, an "amino-protecting group" is a substituent attached to an amino group that blocks or protects the amino functionality in the compound. Suitable amino-protecting groups include but are not limited to acetyl, trifluoroacetyl, t-butoxycarbonyl (BOC), benzyloxycarbonyl (CBZ) and 9-fluorenylmethylenoxycarbonyl (Fmoc). Similarly, a "hydroxy-protecting group" refers to a substituent of a hydroxy group that blocks or protects the hydroxy functionality. Suitable protecting groups include but are not limited to acetyl and silyl. A "carboxy-protecting group" refers to a substituent of the carboxy group that blocks or protects the carboxy functionality. Common carboxy-protecting groups include $-\text{CH}_2\text{CH}_2\text{SO}_2\text{Ph}$, cyanoethyl, 2-(trimethylsilyl)ethyl, 2-(trimethylsilyl)ethoxymethyl, 2-(p-toluenesulfonyl)ethyl, 2-(p-nitrophenylsulfonyl)ethyl, 2-(diphenylphosphino)-ethyl, nitroethyl and the like. For a general description of protecting groups and their use, see T. W. Greene, *Protective Groups in Organic Synthesis*, John Wiley & Sons, New York, 1991.

[0047] In one embodiment, disclosed herein is at least one compound of formula (I):



and/or at least one pharmaceutically acceptable salt thereof, wherein

each R^1 is independently selected from:

hydrogen,

halogen,

hydroxyl,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkyl-alkyl,

heterocyclyl,

heterocyclylalkyl

aryl,

arylalkyl,

heteroaryl, and

heteroarylalkyl,

wherein alkyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a}, and wherein aryl and heteroaryl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6b};

each R² is independently selected from C₁₋₁₀ alkyl and piperidinyl, wherein the C₁₋₁₀ alkyl and piperidinyl are each independently unsubstituted or substituted with at least one substituent, such as one, two, three or four substituents, independently selected from R^{6a};

each R³ is independently selected from:

hydrogen,

halogen,

-CN,

-NR⁷R⁸, and

C₁₋₁₀ alkyl;

wherein alkyl is unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a};

each R₄ is independently selected from:

hydrogen,

halogen,

-CN,

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl, and

C₃₋₁₀ cycloalkyl;

wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and C₃₋₁₀ cycloalkyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a};

or R³ and R⁴ together with the carbon atoms to which they are attached form a 5-6 membered ring containing 0, 1, 2 or 3 heteroatoms independently selected from oxygen, sulfur and nitrogen, and optionally substituted with 1-2 R^{6b} groups;

each R⁵ is independently selected from:

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

$-\text{O}(\text{CR}^9\text{R}^{10})_t\text{NR}^7\text{R}^8$,

$-\text{C}(\text{O})\text{R}^7$,

$-\text{CO}_2\text{R}^8$,

$-\text{CO}_2(\text{CR}^9\text{R}^{10})_t\text{CONR}^7\text{R}^8$,

$-\text{OC}(\text{O})\text{R}^7$,

$-\text{CN}$,

$-\text{C}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{R}^8$,

$-\text{OC}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{OR}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{CR}^7(\text{N}-\text{OR}^8)$,

$-\text{CHF}_2$,

$-\text{CF}_3$,

$-\text{OCHF}_2$, and

$-\text{OCF}_3$;

wherein C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, and C_{3-10} cycloalkyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} ;

each R^{6a} is independently selected from:

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C_{3-10} cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-(CR⁹R¹⁰)_tOR⁸,

-(CR⁹R¹⁰)_tNR⁷R⁸,

-(CR⁹R¹⁰)_tSR⁸,

-(CR⁹R¹⁰)_tS(O)_rR⁸,

-(CR⁹R¹⁰)_tCO₂R⁸,

-(CR⁹R¹⁰)_tCONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷CO₂R⁸,

-(CR⁹R¹⁰)_tOCONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷CONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷SO₂NR⁷R⁸,

-O(CR⁹R¹⁰)_tNR⁷R⁸,

-C(O)R⁷,

$-\text{C}(\text{O})(\text{CR}^9\text{R}^{10})_t\text{OR}^8$,

$-\text{C}(\text{O})(\text{CR}^9\text{R}^{10})_t\text{NR}^7\text{R}^8$,

$-\text{C}(\text{O})(\text{CR}^9\text{R}^{10})_t\text{SR}^8$,

$-\text{C}(\text{O})(\text{CR}^9\text{R}^{10})_t\text{S}(\text{O})_r\text{R}^8$,

$-\text{CO}_2\text{R}^8$,

$-\text{CO}_2(\text{CR}^9\text{R}^{10})_t\text{CONR}^7\text{R}^8$,

$-\text{OC}(\text{O})\text{R}^7$,

$-\text{CN}$,

$-\text{C}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{R}^8$,

$-\text{OC}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{OR}^8$,

$-\text{NR}^7\text{C}(\text{O})\text{NR}^7\text{R}^8$,

$-\text{CR}^7(\text{N}-\text{OR}^8)$,

$-\text{CHF}_2$,

$-\text{CF}_3$,

$-\text{OCHF}_2$, and

$-\text{OCF}_3$;

each R^{6b} is independently selected from:

R^{6a} ,

aryl,

aryl- C_{1-4} alkyl,

heteroaryl, and

heteroaryl C_{1-4} alkyl;

each R^7 and each R^8 are independently selected from:

hydrogen,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C_{3-10} cycloalkyl,

cycloalkyl- C_{1-10} alkyl;

heterocyclyl,

heterocyclyl- C_{1-10} alkyl,

aryl,

heteroaryl,

aryl- C_{1-10} alkyl, and

heteroaryl- C_{1-10} alkyl;

wherein alkyl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} , and aryl and heteroaryl are each unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6b} ; or

R^7 and R^8 together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0, 1, or 2 additional heteroatoms independently selected from oxygen, sulfur and NR^{11} ,

each R^7 and R^8 may be unsubstituted or substituted on a carbon or nitrogen atom with at least one substituent, such as one, two, or three substituents, independently selected from R^{12} ;

each R^9 and each R^{10} are independently selected from:

hydrogen,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

cycloalkyl-C₁₋₁₀ alkyl,

heterocyclyl,

heterocyclyl-C₁₋₁₀ alkyl,

aryl,

heteroaryl,

aryl-C₁₋₁₀ alkyl, and

heteroaryl-C₁₋₁₀ alkyl; or

R⁹ and R¹⁰ together with the carbon to which they are attached form a ring of 3 to 7 members containing 0, 1, or 2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

each R¹¹ is independently selected from:

hydrogen,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl,

heterocyclyl,

heterocyclyl-C₁₋₄ alkyl,

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl,

heteroaryl-C₁₋₄ alkyl,

-S(O)_rR⁷,

-C(O)R⁷,

-CO₂R⁷,

$-\text{CO}_2(\text{CR}^9\text{R}^{10})_t\text{CONR}^7\text{R}^8$, and

$-\text{C}(\text{O})\text{NR}^7\text{R}^8$;

each R^{12} is independently selected from:

halogen,

C_{1-10} alkyl,

C_{3-10} cycloalkyl,

C_{3-10} cycloalkylalkyl,

heterocyclyl,

heterocyclylalkyl,

aryl,

aryl- C_{1-4} alkyl,

heteroaryl,

heteroaryl- C_{1-4} alkyl,

$-\text{OR}^7$,

$-\text{NR}^7\text{S}(\text{O})_r\text{R}^8$,

$-\text{S}(\text{O})_r\text{R}^7$,

$-\text{SR}^7$,

$-\text{S}(\text{O})_2\text{OR}^7$,

$-\text{OS}(\text{O})_2\text{R}^7$,

$-\text{S}(\text{O})_r\text{NR}^7\text{R}^8$,

$-\text{NR}^7\text{R}^8$,

$-\text{O}(\text{CR}^9\text{R}^{10})_t\text{NR}^7\text{R}^8$,

$-\text{C}(\text{O})\text{R}^7$,

$-\text{CO}_2\text{R}^8$,

$-\text{CO}_2(\text{CR}^9\text{R}^{10})_t\text{CONR}^7\text{R}^8,$

$-\text{OC}(\text{O})\text{R}^7,$

$-\text{CN},$

$-\text{C}(\text{O})\text{NR}^7\text{R}^8,$

$-\text{NR}^7\text{C}(\text{O})\text{R}^8,$

$-\text{OC}(\text{O})\text{NR}^7\text{R}^8,$

$-\text{NR}^7\text{C}(\text{O})\text{OR}^8,$

$-\text{NR}^7\text{C}(\text{O})\text{NR}^7\text{R}^8,$

$-\text{CHF}_2,$

$-\text{CF}_3,$

$-\text{OCHF}_2,$ and

$-\text{OCF}_3;$

each m is independently selected from 0, 1 and 2; each n is independently selected from 1, 2, and 3; p is 2;

each q is independently selected from 0, 1, 2, and 3; each r is independently selected from 1 and 2; each t is independently selected from 1, 2, and 3.

[0048] In some embodiments, each R^1 is independently selected from hydrogen and C_{1-10} alkyl, wherein alkyl is unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from hydroxyl and C_{1-10} alkoxy.

[0049] In some embodiments, R^1 is hydrogen.

[0050] In some embodiments, each R^1 is independently selected from C_{1-10} alkyl, wherein alkyl is unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from hydroxyl and C_{1-10} alkoxy.

[0051] In some embodiments, R^1 is methyl, hydroxymethyl, or methoxymethyl.

[0052] In some embodiments, each R^2 is independently selected from C_{1-10} alkyl and piperidinyl which are unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} , R^{6a} is as described above.

[0053] In some embodiments, each R^2 is independently selected from methyl and 4-piperidinyl which are unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} , R^{6a} is as described above.

[0054] In some embodiments, each R^2 is independently selected from methyl and 4-piperidinyl, wherein 4-piperidinyl is unsubstituted or substituted with at least one substituent, such as one, two, three, or four substituents, independently selected from R^{6a} , wherein R^{6a} is independently selected from C_{1-10} alkyl, $-C(O)R^7$, $-S(O)_rR^7$, $-(CR^9R^{10})_tCO_2R^8$, $-C(O)(CR^9R^{10})_tNR^7R^8$, $-(CR^9R^{10})_tOR^8$, $-(CR^9R^{10})_tCONR^7R^8$, $-(CR^9R^{10})_tS(O)_rR^8$, $-(CR^9R^{10})_tOR^8$, R^7 , R^8 , R^9 , R^{10} , t and r are as described above.

[0055] In some embodiments, R^3 is hydrogen.

[0056] In some embodiments, each R^4 is independently selected from hydrogen, C_{1-10} alkyl, and halogen.

[0057] In some embodiments, each R^4 is independently selected from halogen.

[0058] In some embodiments, R^4 is chlorine.

[0059] In some embodiments, R^3 and R^4 together with the carbon atoms to which they are attached form a 5-membered ring containing 2 nitrogen atoms, wherein the 5-membered ring is optionally substituted with 1 or 2 C_{1-10} alkyl.

[0060] In some embodiments, R^3 and R^4 together with the carbon atoms to which they are attached form pyrazolo.

[0061] In some embodiments, R^5 is $-S(O)_rR^7$, R^7 and r are as described above.

[0062] In some embodiments, R^5 is $-S(O)_2R^7$, wherein R^7 is selected from C_{1-10} alkyl.

[0063] In some embodiments, R^5 is $-S(O)_2R^7$, wherein R^7 is isopropyl.

[0064] In some embodiments, n is 1.

[0065] In some embodiments, p is 2.

[0066] In some embodiments, m is 1.

[0067] In some embodiments, q is 1.

[0068] Also provided is at least one compound, selected from:

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*S*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*S*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(7-(4-(2-(isopropylsulfonyl)phenylamino)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-6-ylamino)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol;

N-4-(2-(isopropylsulfonyl)phenyl)-*N*-6-(2-(methoxymethyl)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-5-chloro-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*R*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*S*)-5-chloro-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*S*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(S)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(R)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid;

(S)-2-amino-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(S)-2-amino-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(R)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(S)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-*N,N*-dimethylacetamide;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-*N,N*-dimethylacetamide;

(S)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(R)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(S)-5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-

dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(R)-5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

(R)-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(S)-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxyethanone;

(S)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one;

(R)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one;

(S)-methyl 4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxylate;

(S)-ethyl 2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetate;

(S)-2-amino-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(S)-4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxamide;

(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(methylamino)ethanone;

(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(dimethylamino)ethanone;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methylpropan-1-one;

(*R*)-4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)(cyclopropyl)methanone;

(*R*)-5-chloro-*N*-2-(4-(1-(ethylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*R*)-5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(4-(1-(isopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(*R*)-5-chloro-*N*-2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*R*)-1-(4-((*S*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one;

(*R*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one;

(*S*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one;

(*S*)-1-(4-((*S*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one, and pharmaceutically acceptable salts thereof.

[0069] The compounds disclosed herein and their pharmaceutically acceptable salts exhibit valuable pharmacological properties when tested in vitro in cell-free kinase assays and in cellular assays, and are therefore useful as pharmaceuticals.

[0070] In one aspect, compounds of formula (I) may inhibit the tyrosine kinase activity of ALK and the fusion protein of NPM-ALK. This protein tyrosine kinase results from a gene fusion of nucleophosmin (NPM) and ALK, rendering the protein tyrosine kinase activity of ALK ligand independent. NPM-ALK plays a key role in signal transmission in a number of hematopoietic and other human cells leading to hematological and neoplastic diseases, for example in anaplastic large-cell lymphoma (ALCL) and non-Hodgkin's lymphomas (NHL), specifically in ALK+NHL or Alkomas, in inflammatory myofibroblastic tumors (IMT) and neuroblastomas. (Duyster et al. 2001 Oncogene 20, 5623-5637). In addition to NPM-ALK, other gene fusions have been identified in human hematological and neoplastic diseases; for example, TPM3-ALK (a fusion of nonmuscle tropomyosin with ALK).

[0071] The inhibition of ALK tyrosine kinase activity may be demonstrated using known methods, for example using the recombinant kinase domain of the ALK in analogy to the VEGF-R kinase assay described in J. Wood et al. Cancer Res. 60, 2178-2189 (2000). In general, in vitro enzyme assays using GST-ALK protein tyrosine kinase are performed in 96-

well plates as a filter binding assay in 20 mM Tris HCl, pH = 7.5, 3 mM MgCl₂, 10 mM MnCl₂, 1 mM DTT, 0.1 µCi/assay (=30 µl) [γ -³²P]-ATP, 2 µM ATP, 3 µg/mL poly (Glu, Tyr 4:1) Poly-EY (Sigma P-0275), 1% DMSO, 25 ng ALK enzyme. Assays are incubated for 10 min at ambient temperature. Reactions are terminated by adding 50 µl of 125 mM EDTA, and the reaction mixture is transferred onto a MAIP Multiscreen plate (Millipore, Bedford, Mass., USA), previously wet with methanol, and rehydrated for 5 min with H₂O. Following washing (0.5% H₃PO₄), plates are counted in a liquid scintillation counter. IC₅₀ values are calculated by linear regression analysis of the percentage inhibition.

[0072] Compounds of formula (I) may potently inhibit the growth of human NPM-ALK overexpressing murine BaF3 cells (DSMZ Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Germany). The expression of NPM-ALK may be achieved by transfecting the BaF3 cell line with an expression vector pCneo™ (Promega Corp., Madison Wis., USA) coding for NPM-ALK and subsequent selection of G418 resistant cells. Non-transfected BaF3 cells depend on IL-3 for cell survival. In contrast, NPM-ALK expressing BaF3 cells (named BaF3-NPM-ALK hereinafter) can proliferate in the absence of IL-3 because they obtain proliferative signal through NPM-ALK kinase. Putative inhibitors of the NPM-ALK kinase therefore abolish the growth signal and may result in antiproliferative activity. The antiproliferative activity of putative inhibitors of the NPM-ALK kinase can however be overcome by addition of IL-3, which provides growth signals through an NPM-ALK independent mechanism. An analogous cell system using FLT3 kinase has also been described (see, E Weisberg et al. Cancer Cell; 1, 433-443 (2002)).

[0073] The inhibitory activity of the compounds of the invention may be determined as follows. In general, BaF3-NPM-ALK cells (15,000/microtitre plate well) are transferred to 96-well microtitre plates. Test compounds dissolved in dimethyl sulfoxide (DMSO) are added in a series of concentrations (dilution series) in such a manner that the final concentration of DMSO is not greater than 1% (v/v). After the addition, the plates are incubated for two days during which the control cultures without test compound are able to undergo two cell-division cycles. The growth of the BaF3-NPM-ALK cells is measured by means of YOPRO™ staining [T Idziorek et al. J. Immunol. Methods; 185: 249-258 (1995)]: 25 µl of lysis buffer comprising 20 mM sodium citrate, pH 4.0, 26.8 mM sodium chloride, 0.4% NP40, 20 mM EDTA and 20 mM is added to each well. Cell lysis is completed within 60 min at room temperature and total amount of YOPRO™ bound to DNA is determined by measurement using the Cytofluor II 96-well reader (PerSeptive Biosystems) with the following settings: Excitation (nm) 485/20 and Emission (nm) 530/25.

[0074] IC₅₀ values may be determined by a computer-aided system using the formula:

$$IC_{50} = [(ABS_{test} - ABS_{start}) / (ABS_{control} - ABS_{start})] \times 100.$$
 (ABS=absorption)

[0075] The IC₅₀ value in those experiments is given as that concentration of the test compound in question that results in a cell count that is 50% lower than that obtained using the control without inhibitor. The compounds of the disclosure in free form or in pharmaceutically

acceptable salt form, may exhibit valuable pharmacological properties, for example, as indicated by the in vitro tests described in this application. In general, compounds of the disclosure have IC_{50} values from less than 1 nM to 10 μ M. In some examples, compounds of the disclosure have IC_{50} values from less than 1 nM to 500 nM. In other examples, compounds of the disclosure have IC_{50} values from < 1nM to 200 nM.

[0076] The antiproliferative action of the inventive compounds may also be determined in the human KARPAS-299 lymphoma cell line (DSMZ Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany, described in WG Dirks et al. Int. J. Cancer 100, 49-56 (2002)) using the same methodology described above for the BaF3-NPM-ALK cell line. In some embodiments, compounds disclosed herein may exhibit inhibitory activity with an IC_{50} in the range from approximately less than 1nM to 1 μ M. The action of the inventive compounds on autophosphorylation of the ALK may be determined in the human KARPAS-299 lymphoma cell line by means of an immunoblot as described in WG Dirks et al. Int. J. Cancer 100, 49-56 (2002).

[0077] In accordance with the foregoing, the present disclosure provides:

1. (1) a compound disclosed herein for use as a pharmaceutical;
2. (2) a compound disclosed herein for use as an ALK inhibitor, for example, for use in any of the particular indications set forth above;
3. (3) a pharmaceutical composition, e.g. for use in any of the indications herein set forth above, comprising at least one compound disclosed herein as active ingredient together with one or more pharmaceutically acceptable diluents or carriers;
4. (4) at least one compound disclosed herein or at least one pharmaceutical composition comprising thereof for use in treating any particular indication set forth above;
5. (5) the use of a compound disclosed herein for making a medicament for treatment or prevention of a disease or condition in which ALK activation plays a role or is implicated;
6. (6) the compound or pharmaceutical composition as defined above under (4) comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a compound disclosed herein and one or more additional drug substances, said additional drug substance being useful in any of the particular indications set forth above;
7. (7) a composition comprising a therapeutically effective amount of at least one compound disclosed herein and at least one additional drug substance, wherein said additional drug substance is useful in any of the particular indications set forth above;
8. (8) use of a compound disclosed herein for making a medicament for treatment or prevention of a disease which responds to inhibition of the anaplastic lymphoma kinase;
9. (9) the use according to (8), wherein the disease to be treated is selected from anaplastic large cell lymphoma, non-Hodgkin's lymphomas, inflammatory myofibroblastic tumors, neuroblastomas and neoplastic diseases;
10. (10) the use according to (8) or (9), wherein the compound is a pharmaceutically acceptable salt of any one of the examples disclosed herein;
11. (11) at least one compound disclosed herein or at least one pharmaceutical composition

comprising thereof for use in treating a disease which responds to inhibition of the anaplastic lymphoma kinase, such as a disease selected from anaplastic large-cell lymphoma, non Hodgkin's lymphomas, inflammatory myofibroblastic tumors, neuroblastomas and neoplastic diseases, comprising administering an effective amount of at least one compound disclosed herein and/or a pharmaceutically acceptable salt thereof. In some examples, the compounds of the disclosure may be used alone or in combination with a chemotherapeutic agent to treat a cell proliferative disorder, including but not limited to, lymphoma, osteosarcoma, melanoma, or a tumor of breast, renal, prostate, colorectal, thyroid, ovarian, pancreatic, neuronal, lung, uterine or gastrointestinal tumor.

Administration and Pharmaceutical Compositions

[0078] In general, compounds of the disclosure will be administered in therapeutically effective amounts via any of the usual and acceptable modes known in the art, either singly or in combination with one or more therapeutic agents. A therapeutically effective amount may vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the compound used and other factors known to those of ordinary skill in the art. For example, for the treatment of neoplastic diseases and immune system disorders, the required dosage will also vary depending on the mode of administration, the particular condition to be treated and the effect desired.

[0079] In general, satisfactory results are indicated to be obtained systemically at daily dosages of from about 0.001 to about 100 mg/kg per body weight, or particularly, from about 0.03 to 2.5 mg/kg per body weight. An indicated daily dosage in the larger mammal, e.g. humans, may be in the range from about 0.5 mg to about 2000 mg, or more particularly, from about 0.5 mg to about 100 mg, conveniently administered, for example, in divided doses up to four times a day or in retard form. Suitable unit dosage forms for oral administration comprise from ca. 1 to 50 mg active ingredient.

[0080] Compounds of the disclosure may be administered as pharmaceutical compositions by any conventional route; for example, enterally, e.g., orally, e.g., in the form of tablets or capsules; parenterally, e.g., in the form of injectable solutions or suspensions; or topically, e.g., in the form of lotions, gels, ointments or creams, or in a nasal or suppository form.

[0081] Pharmaceutical compositions comprising a compound of the present disclosure in free form or in a pharmaceutically acceptable salt form in association with at least one pharmaceutically acceptable carrier or diluent may be manufactured in a conventional manner by mixing, granulating, coating, dissolving or lyophilizing processes. For example, pharmaceutical compositions comprising a compound of the disclosure in association with at least one pharmaceutical acceptable carrier or diluent may be manufactured in conventional

manner by mixing with a pharmaceutically acceptable carrier or diluent. Unit dosage forms for oral administration contain, for example, from about 0.1 mg to about 500 mg of active substance.

[0082] In one embodiment, the pharmaceutical compositions are solutions of the active ingredient, including suspensions or dispersions, such as isotonic aqueous solutions. In the case of lyophilized compositions comprising the active ingredient alone or together with a carrier such as mannitol, dispersions or suspensions can be made up before use. The pharmaceutical compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. Suitable preservatives include but are not limited to antioxidants such as ascorbic acid, or microbicides, such as sorbic acid or benzoic acid. The solutions or suspensions may further comprise viscosity-increasing agents, including but not limited to, sodium carboxymethylcellulose, carboxymethylcellulose, dextran, polyvinylpyrrolidone, gelatins, or solubilizers, e.g. Tween 80 (polyoxyethylene(20)sorbitan mono-oleate).

[0083] Suspensions in oil may comprise as the oil component the vegetable, synthetic, or semi-synthetic oils customary for injection purposes. Examples include liquid fatty acid esters that contain as the acid component a long-chained fatty acid having from 8 to 22 carbon atoms, or in some embodiments, from 12 to 22 carbon atoms. Suitable liquid fatty acid esters include but are not limited to lauric acid, tridecylic acid, myristic acid, pentadecylic acid, palmitic acid, margaric acid, stearic acid, arachidic acid, behenic acid or corresponding unsaturated acids, for example oleic acid, elaidic acid, erucic acid, brassidic acid and linoleic acid, and if desired, may contain antioxidants, for example vitamin E, 3-carotene or 3,5-di-tert-butyl-hydroxytoluene. The alcohol component of these fatty acid esters may have six carbon atoms and may be monovalent or polyvalent, for example a mono-, di- or trivalent, alcohol. Suitable alcohol components include but are not limited to methanol, ethanol, propanol, butanol or pentanol or isomers thereof; glycol and glycerol.

[0084] Other suitable fatty acid esters include but are not limited ethyl-oleate, isopropyl myristate, isopropyl palmitate, LABRAFIL® M 2375, (polyoxyethylene glycerol), LABRAFIL® M 1944 CS (unsaturated polyglycolized glycerides prepared by alcoholysis of apricot kernel oil and comprising glycerides and polyethylene glycol ester), LABRASOL™ (saturated polyglycolized glycerides prepared by alcoholysis of TCM and comprising glycerides and polyethylene glycol ester; all available from GaKefosse, France), and/or MIGLYOL® 812 (triglyceride of saturated fatty acids of chain length C8 to C12 from Hüls AG, Germany), and vegetable oils such as cottonseed oil, almond oil, olive oil, castor oil, sesame oil, soybean oil, or groundnut oil.

[0085] Pharmaceutical compositions for oral administration may be obtained, for example, by combining the active ingredient with one or more solid carriers, and if desired, granulating a resulting mixture, and processing the mixture or granules by the inclusion of additional excipients, to form tablets or tablet cores.

[0086] Suitable carriers include but are not limited to fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations, and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starches, for example corn, wheat, rice or potato starch, methylcellulose, hydroxypropyl methylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone, and/or, if desired, disintegrators, such as the above-mentioned starches, carboxymethyl starch, crosslinked polyvinylpyrrolidone, alginic acid or a salt thereof, such as sodium alginate. Additional excipients include flow conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol, or derivatives thereof.

[0087] Tablet cores may be provided with suitable, optionally enteric, coatings through the use of, inter alia, concentrated sugar solutions which may comprise gum arable, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable organic solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Dyes or pigments may be added to the tablets or tablet coatings, for example for identification purposes or to indicate different doses of active ingredient.

[0088] Pharmaceutical compositions for oral administration may also include hard capsules comprising gelatin or soft-sealed capsules comprising gelatin and a plasticizer, such as glycerol or sorbitol. The hard capsules may contain the active ingredient in the form of granules, for example in admixture with fillers, such as corn starch, binders, and/or glidants, such as talc or magnesium stearate, and optionally stabilizers. In soft capsules, the active ingredient may be dissolved or suspended in suitable liquid excipients, such as fatty oils, paraffin oil or liquid polyethylene glycols or fatty acid esters of ethylene or propylene glycol, to which stabilizers and detergents, for example of the polyoxyethylene sorbitan fatty acid ester type, may also be added.

[0089] Pharmaceutical compositions suitable for rectal administration are, for example, suppositories comprising a combination of the active ingredient and a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols.

[0090] Pharmaceutical compositions suitable for parenteral administration may comprise aqueous solutions of an active ingredient in water-soluble form, for example of a water-soluble salt, or aqueous injection suspensions that contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if desired, stabilizers. The active ingredient, optionally together with excipients, can also be in the form of a lyophilizate and can be made into a solution before parenteral administration by the addition of suitable solvents. Solutions such as are used, for example, for parenteral administration can also be employed as infusion solutions. The manufacture of injectable preparations is usually carried out under sterile conditions, as is the filling, for example, into ampoules or vials, and the

sealing of the containers.

[0091] The compounds of the disclosure may be administered as the sole active ingredient, or together with other drugs useful against neoplastic diseases or useful in immunomodulating regimens. For example, the compounds of the disclosure may be used in accordance with the disclosure in combination with pharmaceutical compositions effective in various diseases as described above, e.g. with cyclophosphamide, 5-fluorouracil, fludarabine, gemcitabine, cisplatin, carboplatin, vincristine, vinblastine, etoposide, irinotecan, paclitaxel, docetaxel, rituxan, doxorubicine, gefitinib, or imatinib; or also with cyclosporins, rapamycins, ascomycins or their immunosuppressive analogs, e.g. cyclosporin A, cyclosporin G, FK-506, sirolimus or everolimus, corticosteroids, e.g. prednisone, cyclophosphamide, azathioprene, methotrexate, gold salts, sulfasalazine, antimalarials, brequinar, leflunomide, mizoribine, mycophenolic acid, mycophenolate, mofetil, 15-deoxyspergualine, immuno-suppressive monoclonal antibodies, e.g. monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, I CD40, CD45, CD58, CD80, CD86, CD152, CD137, CD154, ICOS, LFA-1, VLA-4 or their ligands, or other immunomodulatory compounds, e.g. CTLA41g.

[0092] The disclosure also provides for a pharmaceutical combinations, e.g. a kit, comprising a) a first agent which is a compound of the disclosure as disclosed herein, in free form or in pharmaceutically acceptable salt form, and b) at least one co-agent. The kit can comprise instructions for its administration.

EXAMPLES

[0093] Various methods may be developed for synthesizing the at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof. Representative methods for synthesizing the at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof are provided in the Examples. It is noted, however, that the at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof may also be synthesized by other synthetic routes that others may devise.

[0094] It will be readily recognized that certain compounds of formula (I) have atoms with linkages to other atoms that confer a particular stereochemistry to the compound (e.g., chiral centers). It is recognized that synthesis of the at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof may result in the creation of mixtures of different stereoisomers (enantiomers, diastereomers). Unless a particular stereochemistry is specified, recitation of a compound is intended to encompass all of the different possible stereoisomers.

[0095] The at least one compound of formula (I) can also be prepared as a pharmaceutically acceptable acid addition salt by, for example, reacting the free base form of the at least one compound with a pharmaceutically acceptable inorganic or organic acid. Alternatively, a pharmaceutically acceptable base addition salt of the at least one compound of formula (I) can

be prepared by, for example, reacting the free acid form of the at least one compound with a pharmaceutically acceptable inorganic or organic base. Inorganic and organic acids and bases suitable for the preparation of the pharmaceutically acceptable salts of compounds of formula (I) are set forth in the definitions section of this Application. Alternatively, the salt forms of the compounds of formula (I) can be prepared using salts of the starting materials or intermediates.

[0096] The free acid or free base forms of the compounds of formula (I) can be prepared from the corresponding base addition salt or acid addition salt form. For example, a compound of formula (I) in an acid addition salt form can be converted to the corresponding free base thereof by treating with a suitable base (e.g., ammonium hydroxide solution, sodium hydroxide, and the like). A compound of formula (I) in a base addition salt form can be converted to the corresponding free acid thereof by, for example, treating with a suitable acid (e.g., hydrochloric acid, etc).

[0097] The N-oxides of the at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof can be prepared by methods known to those of ordinary skill in the art. For example, N-oxides can be prepared by treating an unoxidized form of the compound of formula (I) with an oxidizing agent (e.g., trifluoroperacetic acid, permaleic acid, perbenzoic acid, peracetic acid, meta-chloroperoxybenzoic acid, or the like) in a suitable inert organic solvent (e.g., a halogenated hydrocarbon such as dichloromethane) at approximately 0 to 80 °C. Alternatively, the N-oxides of the compounds of formula (I) can be prepared from the N-oxide of an appropriate starting material.

[0098] Compounds of formula (I) in an unoxidized form can be prepared from N-oxides of compounds of formula (I) by, for example, treating with a reducing agent (e.g., sulfur, sulfur dioxide, triphenyl phosphine, lithium borohydride, sodium borohydride, phosphorus trichloride, tribromide, and the like) in a suitable inert organic solvent (e.g., acetonitrile, ethanol, aqueous dioxane, and the like) at 0 to 80 °C.

[0099] Protected derivatives of the compounds of formula (I) can be made by methods known to those of ordinary skill in the art. A detailed description of the techniques applicable to the creation of protecting groups and their removal can be found in T.W. Greene, *Protecting Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons, Inc. 1999.

[0100] The at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof may be conveniently prepared, or as solvates (e.g. hydrates). Hydrates of the at least one compound of formula I and/or at least one pharmaceutically acceptable salt thereof may be conveniently prepared by recrystallization from an aqueous/organic solvent mixture, using organic solvents such as dioxin, tetrahydrofuran and/or methanol.

[0101] The compounds of formula (I) can also be prepared as their individual stereoisomers by reacting a racemic mixture of the compounds with an optically active resolving agent to form a pair of diastereoisomeric compounds, separating the diastereomers, and recovering the

optically pure enantiomer. While resolution of enantiomers can be carried out using covalent diastereomeric derivatives of compounds, dissociable complexes are preferred (e.g., crystalline diastereoisomeric salts). Diastereomers have distinct physical properties (e.g., melting points, boiling points, solubilities, reactivity, etc.) and can be readily separated by taking advantage of these dissimilarities. The diastereomers can be separated by chromatography or, for example, by separation/resolution techniques based upon differences in solubility. The optically pure enantiomer is then recovered, along with the resolving agent, by any practical means that would not result in racemization. A more detailed description of the techniques applicable to the resolution of stereoisomers of compounds from their racemic mixture can be found in Jean Jacques Andre Collet, Samuel H. Wilen, *Enantiomers, Racemates and Resolutions*, John Wiley & Sons, Inc. (1981).

[0102] As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the Journal of the American Chemical Society or the Journal of Biological Chemistry. Standard single-letter or three-letter abbreviations are generally used to designate amino acid residues, which are assumed to be in the L-configuration unless otherwise noted. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. For example, the following abbreviations may be used in the examples and throughout the specification: g (grams); mg (milligrams); L (liters); mL (milliliters); μ L (microliters); psi (pounds per square inch); M (molar); mM (millimolar); i.v. (intravenous); Hz (Hertz); MHz (megahertz); mol (moles); mmol (millimoles); RT (room temperature); min (minutes); h (hours); mp (melting point); TLC (thin layer chromatography); Rt (retention time); RP (reverse phase); MeOH (methanol); i-PrOH (isopropanol); TEA (triethylamine); TFA (trifluoroacetic acid); TFAA (trifluoroacetic anhydride); THF (tetrahydrofuran); DMSO (dimethyl sulfoxide); EtOAc (ethyl acetate); DME (1,2-dimethoxyethane); DCM (dichloromethane); DCE (dichloroethane); DMF (N,N-dimethylformamide); DMPU (N,N'-dimethylpropyleneurea); CDI (1,1-carbonyldiimidazole); IBCF (isobutyl chloroformate); HOAc (acetic acid); HOSu (N-hydroxysuccinimide); HOBt (1-hydroxybenzotriazole); Et₂O (diethyl ether); EDCI (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride); BOC (tert-butyloxycarbonyl); Fmoc (9-fluorenylmethoxycarbonyl); DCC (dicyclohexylcarbodiimide); CBZ (benzyloxycarbonyl); Ac (acetyl); atm (atmosphere); TMSE (2-(trimethylsilyl)ethyl); TMS (trimethylsilyl); TIPS (triisopropylsilyl); TBS (t-butyldimethylsilyl); DMAP (4-dimethylaminopyridine); Me (methyl); OMe (methoxy); Et (ethyl); tBu (tert-butyl); HPLC (high pressure liquid chromatography); BOP (bis(2-oxo-3-oxazolidinyl)phosphinic chloride); TBAF (tetra-n-butylammonium fluoride); m-CPBA (meta-chloroperoxybenzoic acid).

[0103] References to ether or Et₂O are to diethyl ether; brine refers to a saturated aqueous solution of NaCl. Unless otherwise indicated, all temperatures are expressed in °C (degrees Centigrade). All reactions were conducted under an inert atmosphere at RT unless otherwise noted.

[0104] ¹H NMR spectra were recorded on a Varian Mercury Plus 400. Chemical shifts are expressed in parts per million (ppm). Coupling constants are in units of hertz (Hz). Splitting

patterns describe apparent multiplicities and are designated as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), and br (broad).

[0105] Low-resolution mass spectra (MS) and compound purity data were acquired on a Shimadzu LC/MS single quadrupole system equipped with electrospray ionization (ESI) source, UV detector (220 and 254 nm), and evaporative light scattering detector (ELSD). Thin-layer chromatography was performed on 0.25 mm E. Merck silica gel plates (60F-254), visualized with UV light, 5% ethanolic phosphomolybdic acid, ninhydrin, or p-anisaldehyde solution. Flash column chromatography was performed on silica gel (230-400 mesh, Merck).

Synthetic Schemes

[0106] The at least one compound of formula (I) and/or at least one pharmaceutically acceptable salt thereof may be synthesized according to a variety of reaction schemes. Some illustrative schemes are provided below and in the examples. Other reaction schemes could be readily devised by those skilled in the art in view of the present disclosure.

[0107] In the reactions disclosed below, it may be necessary to protect reactive functional groups, for example hydroxy, amino, imino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice, for example, see T.W. Greene and P. G. M. Wuts in "Protective Groups in Organic Chemistry" John Wiley and Sons, 1991.


[0108] Synthetic methods for preparing the compounds in the present disclosure are illustrated in the following Schemes and Examples. Starting materials are commercially available or may be made according to procedures known in the art or as illustrated herein.

Preparation of Intermediates

Intermediate A

4-Hydroxy-3-methyl-6-mercaptopyrazolo[3,4-d]pyrimidine (A-1)

[0109]




A-1

[0110] This reagent was prepared according to the method described in literature: J. Med. Chem. 1990, 33: 2174-2178.

4-Hydroxy-3-methyl-6-methylmercaptopyrazolo[3,4-d]pyrimidine (A-2)


[0111]



[0112] To a mixture of 4-hydroxy-6-mercapto-3-methylpyrazolo[3,4-d]pyrimidine (A-1) (2.86 g, 11.8 mmol) and NaOH (0.8 g, 20 mmol) in water (20 mL) at 5 °C was added CH₃I (2.0 g, 14.2 mmol). After stirred at ambient temperature for 30 min, the solution was acidified with HOAc to give the crude product of the title compound (A-2), which used in next step without further purification. MS-ESI (m/z): 197.0 (M + H)⁺.

4-chloro-3-methyl-6-methylmercaptopyrazolo[3,4-d]pyrimidine (A-3)


[0113]



[0114] To a mixture of POCl₃ (26 mL) and of dimethylaniline (2 mL) was added crude 4-hydroxy-3-methyl-6-methylmercaptopyrazolo[3,4-d]pyrimidine (A-2) (1.46 g, 7.45 mmol). The solution was heated to reflux for 60 min until all solid dissolved. The excess of POCl₃ was removed under reduced pressure. The residue was poured into ice water with vigorous stirring, it is stirred for 10 min, and the aqueous solution was extracted with ether. The ether layer was washed with cold water, dried with anhydrous Na₂SO₄ and concentrated to give the title compound (A-3) (1.21 g, 5.63 mmol). MS-ESI (m/z): 215.0 (M + H)⁺.

2-(isopropylthio)aniline (A-4)


[0115]



[0116] This reagent was prepared according to the method described in literature: J. Med. Chem. 2002, 45: 2229-2239.

***N*-(2-(isopropylthio)phenyl)-3-methyl-6-(methylthio)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4-amine (A-5)**


[0117]



[0118] A mixture of 4-chloro-3-methyl-6-methylmercaptopyrazolo[3,4-*d*]pyrimidine (A-3) (1.21 g, 5.63 mmol) and 2-(isopropylthio)aniline (A-4) (1.42 g, 8.50 mmol) in isopropanol (56 mL) was stirred at reflux for 3 h. After cooling down to ambient temperature, TEA (0.78 mL, 5.66 mmol) was added to the mixture, and the solution was again heated to reflux for 0.5 h. After cooling down to ambient temperature, the mixture was filtered to give the title compound (A-5) as white solid (1.30 g, 3.77 mmol). MS-ESI (*m/z*): 346 (*M* + *H*)⁺.

***N*-(2-(isopropylsulfonyl)phenyl)-3-methyl-6-(methylsulfonyl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4-amine (Intermediate A)**

[0119]





[0120] To a solution of *N*-(2-(isopropylsulfonyl)phenyl)-6-(methylthio)-1*H*-pyrazolo[3,4-*d*]pyri-

midin-4-amine (A-5) (1.3 g, 3.77 mmol) in DCM (150 mL) was added m-CPBA (5.6 g, 22.6 mmol) at 0 °C. Then the reaction mixture was warmed to ambient temperature and stirred for 4.5 h, washed with saturated NaHSO₃, brine, dried with Na₂SO₄ and concentrated to give the title compound (Intermediate A) as a solid. MS-ESI (m/z): 410 (M + H)⁺.


Intermediate B

2,5-dichloro-N-(2-(isopropylsulfonyl)phenyl)pyrimidin-4-amine (B)

[0121]




[0122] To a mixture of 2-(isopropylthio)aniline (23 g, 0.14 mol) in acetonitrile (250 mL) at RT was added DMAP (8.5 g, 70 mmol) and (Boc)₂O (150 g, 688 mmol). After stirred at ambient temperature for 12 h, the solution was concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (B-1).




[0123] To a mixture of (B-1) (54 g, 0.14 mol) in DCM (500 mL) at 0 °C was added m-CPBA (101 g, 0.44 mol). After stirred at ambient temperature for 12 h, the solution was washed with saturated NaHSO₃, brine, dried with Na₂SO₄ and concentrated to give the compound (B-2) as a solid.

2-(isopropylsulfonyl)aniline (B-3)

[0128]

**2-bromo-1-(2-bromoethoxy)-4-methylbenzene (1a)**


[0129]




[0130] A mixture of 2-bromo-4-methylphenol (5.0 g, 26 mmol), 1,2-dibromoethane (7.0 g, 37 mmol) in water (20 mL) and NaOH (1.1 g, 28 mmol) was stirred at 100 °C for 24 h. The mixture was extracted with DCM (150 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1) to give the title compound (1a).

5-methyl-2,3-dihydrobenzofuran (1b)


[0131]




[0132] To a solution of 2-bromo-1-(2-bromoethoxy)-4-methylbenzene (1a) (1.0 g, 3.4 mmol) in THF (20 mL) at -78 °C under N₂ was added n-BuLi (2.5 M, 11 mL, 27 mmol) and the resulting mixture was stirred at -78 °C for 1 h. The mixture was quenched with saturated NH₄Cl (10 mL), extracted with EtOAc (150 mL), washed with brine, dried and concentrated to give the title compound (1b) as a colorless oil.

5-methyl-7-nitro-2,3-dihydrobenzofuran (1c)**[0133]**

[0134] A mixture of 5-methyl-2,3-dihydrobenzofuran (1b) (2.7 g, 20 mmol) in TFA (50 mL) was cooled to 0 °C, NaNO₂ (2.5 g, 36 mmol) was added thereto. Then it was warmed up to ambient temperature and stirred for 12 h. The mixture was concentrated, poured into water (50 mL), extracted with EtOAc (150 mL), washed with brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (1c).

5-methyl-2,3-dihydrobenzofuran-7-amine (1d)**[0135]**


[0136] A mixture of 5-methyl-7-nitro-2,3-dihydrobenzofuran (1c) (0.30 g, 1.7 mmol) and Pd/C (150 mg, 50%) in THF (5 mL) was introduced H₂. It was stirred at ambient temperature for 12 h. The mixture was filtered through celite and concentrated to give the title compound (1d) as a solid. MS-ESI (m/z): 150 (M + 1)⁺.

4-bromo-5-methyl-2,3-dihydrobenzofuran-7-amine (1e)**[0137]**

[0138] To a mixture of 5-methyl-2,3-dihydrobenzofuran-7-amine (1d) (1.0 g, 6.7 mmol) in DMF (21 mL) at 0 °C was added NBS (1.19 g, 6.7 mmol), it was stirred at 0 °C for 15 min. The mixture was extracted with EtOAc (150 mL), washed with saturated NaHSO₃ (50 mL), water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1) to give the title compound (1e) as a yellow solid. MS-ESI (m/z): 228 (M + 1)⁺.

5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (1f)


[0139]



[0140] A mixture of 4-bromo-5-methyl-2,3-dihydrobenzofuran-7-amine (1e) (0.80 g, 3.5 mmol), pyridin-4-ylboronic acid (0.86 g, 7 mmol), Cs₂CO₃ (2.84 g, 8.7 mmol), Pd(PPh₃)₄ (0.40 g, 0.346 mmol) and water (7 mL) in DMF (35 mL) was stirred at 135 °C under N₂ for 1 h. It was cooled to ambient temperature, poured into water (50 mL), extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1 to 5:1 to 1:1) to give the title compound (1f) as a yellow solid. MS-ESI (m/z): 227 (M + 1)⁺.

5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (1g)

[0141]




[0142] A mixture of 5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (1f) (0.40 g, 1.8 mmol), PtO₂ (120 mg, 35%) and TEA (0.24 mL, 3.6 mmol) in HOAc (30 mL) was introduced H₂

and stirred at ambient temperature for 24 h (88 psi). The mixture was filtered, concentrated, diluted with EtOAc (50 mL). A solution of ammonium hydroxide was added until pH = 10, it was extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated to give the title compound (1g) as a yellow oil. MS-ESI (m/z): 233 (M + 1)⁺.

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-N-6-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (1)

[0143]




[0144] A mixture of 5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (1g) (25 mg, 0.017 mmol), *N*-(2-(isopropylsulfonyl)phenyl)-3-methyl-6-(methylsulfonyl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4-amine (intermediate A) (53 mg, 0.12 mmol) and *p*-TsOH (20 mg, 0.11 mmol) in *i*-PrOH (0.5 mL) was stirred in a sealed tube at 160 °C for 5 h. The mixture was diluted with EtOAc (20 mL). A solution of ammonium hydroxide was added until pH = 10, it was extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (1) as a yellow solid. MS-ESI (m/z): 562 (M + 1)⁺.

Example 2

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-N-6-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (2)

[0145]






[0146] A mixture of of *N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1H-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (1) (37 mg, 0.064 mmol), formaldehyde (5.5 μ L, 0.64 mmol), HOAc (cat.) and NaBH₃CN (10.3 mg, 0.16 mmol) was stirred at ambient temperature for 0.5 h. The mixture was diluted with NaHCO₃ (10 mL) and EtOAc (20 mL), extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (2) as a yellow solid. MS-ESI (*m/z*): 576 (*M* + 1)⁺.

Example 3


(*R*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (3)

[0147]



1-(benzyloxy)-2-bromo-4-methylbenzene (3a)


[0148]



[0149] To a mixture of 2-bromo-4-methylphenol (5.0 g, 27 mmol) and K_2CO_3 (7.4 g, 53 mmol) in DMF (20 mL) was added benzyl bromide (3.5 mL, 29 mmol) at ambient temperature. It was stirred at 70 °C for 12 h. The mixture was poured into water (50 mL), extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (3a).

(S)-1-(2-(benzyloxy)-5-methylphenyl)propan-2-ol (3b)


[0150]



[0151] To a solution of 1-(benzyloxy)-2-bromo-4-methylbenzene (3a) (1.0 g, 3.6 mmol) in THF (20 mL) at -78 °C under N_2 atmosphere was added n -BuLi (2.5 M, 1.6 mL, 3.9 mmol), the resulting mixture was stirred for 30 min and (S)-2-methyloxirane (0.38 mL, 5.4 mmol) and $BF_3 \cdot Et_2O$ (0.67 mL, 5.4 mmol) were added to the mixture. It was stirred at -78 °C for 1.5 h and warmed up to ambient temperature, quenched with water (10 mL), extracted with EtOAc (30 mL), washed with brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (3b).

(S)-2-(2-hydroxypropyl)-4-methylphenol (3c)

[0152]




[0153] The mixture of (S)-1-(2-(benzyloxy)-5-methylphenyl)propan-2-ol (3b) (0.50 g, 1.9

mmol) and Pd/C (100 mg, 20%) in MeOH (10 mL) was introduced H₂ at ambient temperature. It was stirred at ambient temperature for 18 h. The mixture was filtered through Celite and concentrated to give the title compound (3c).

(R)-2,5-dimethyl-2,3-dihydrobenzofuran (3d)


[0154]



[0155] To a mixture of (S)-2-(2-hydroxypropyl)-4-methylphenol (3c) (1.6 g, 8.4 mmol) and triphenylphosphine (2.7 g, 10 mmol) in THF (60 mL) was added DIAD at ambient temperature. It was concentrated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (3d).

(R)-2,5-dimethyl-7-nitro-2,3-dihydrobenzofuran (3e)


[0156]



[0157] A mixture of (R)-2,5-dimethyl-2,3-dihydrobenzofuran (3d) (2.0 g, 13 mmol) in TFA (40 mL) was cooled to 0 °C, NaNO₂ (1.7 g, 23 mmol) was added thereto. It was stirred at ambient temperature for 12 h. The mixture was concentrated, poured into water (50 mL), extracted with EtOAc (150 mL), washed with brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1) to give the title compound (3e).

(R)-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (3f)


[0158]



[0159] The mixture of (*R*)-2,5-dimethyl-7-nitro-2,3-dihydrobenzofuran (3e) (0.84 g, 4.3 mmol) and Pd/C (100 mg, 15%) in THF (10 mL) was introduced H₂ at ambient temperature. It was stirred at ambient temperature for 12 h. The mixture was filtered through Celite and concentrated to give the title compound (3f) as a solid. MS-ESI (*m/z*): 164 (*M* + 1)⁺.

(*R*)-4-bromo-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (3g)


[0160]



[0161] To a mixture of (*R*)-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (3f) (1.3 g, 7.9 mmol) in DMF (16 mL) was added NBS (1.42 g, 7.9 mmol) at 0 °C. It was stirred for 15 min at 0 °C. The mixture was extracted with EtOAc (150 mL), washed with NaHSO₃ (50 mL), water and brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1) to give the title compound (3g) as a solid. MS-ESI (*m/z*): 242 (*M* + 1)⁺.

(*R*)-2,5-dimethyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (3h)

[0162]




[0163] The mixture of (*R*)-4-bromo-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (3g) (1.0 g, 4.3 mmol), pyridin-4-ylboronic acid (1.08 g, 8.77 mmol), Cs₂CO₃ (3.56 g, 10.9 mmol), Pd(PPh₃)₄ (0.506 g, 0.438 mmol) and water (9 mL) in DMF (44 mL) was stirred at 135 °C under N₂ for 1 h. It was cooled to ambient temperature, poured into water (50 mL), extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1 to 5:1 to 1:1)

to give the title compound (3h) as a solid. MS-ESI (m/z): 241 (M + 1)⁺.

(R)-2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (3i)


[0164]



[0165] [The mixture of (R)-2,5-dimethyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (3h) (0.60 g, 2.4 mmol), PtO₂ (240 mg, 35 %) and TEA (0.38 mL, 4.9 mmol) in HOAc (40 mL) was introduced H₂ and stirred for 24 h at ambient temperature (88 psi). The mixture was filtered, concentrated, and diluted with EtOAc (50 mL). A solution of ammonium hydroxide was added until pH = 10, then extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated to give the title compound (3i) as a yellow oil. MS-ESI (m/z): 247 (M + 1)⁺.

(R)-N-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (3)

[0166]




[0167] A mixture of (R)-2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (3i) (44 mg, 0.017 mmol), N-(2-(isopropylsulfonyl)phenyl)-3-methyl-6-(methylsulfonyl)-1H-pyrazolo[3,4-

d]pyrimidin-4-amine (Intermediate A) (83 mg, 0.20 mmol) and *p*-TsOH (32 mg, 0.17 mmol) in *i*-PrOH (0.5 mL) was stirred in a sealed tube at 160 °C for 5 h. The mixture was diluted with EtOAc (20 mL), ammonium hydroxide was added until pH = 10, it was extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (3) as a yellow solid. MS-ESI (*m/z*): 576 (*M* + 1)⁺.

Example 4

(*R*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (4)

[0168]




[0169] A mixture (*R*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (3) (37 mg, 0.064 mmol), formaldehyde (5.5 µL, 0.64 mmol), HOAc (cat.) and NaBH₃CN (10.3 mg, 0.16 mmol) was stirred at ambient temperature for 0.5 h. The mixture was diluted with NaHCO₃ (10 mL) and EtOAc (20 mL), extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (4) as a solid. MS-ESI (*m/z*): 590 (*M* + 1)⁺.

Example 5


(*S*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine (5)

[0170]



(R)-1-(2-(benzyloxy)-5-methylphenyl)propan-2-ol (5a)

[0171]



[0172] The title compound 5a was prepared according to the method for preparation of compound 3b of Example 3 by replacing (S)-2-methyloxirane with (R)-2-methyloxirane.

(R)-2-(2-hydroxypropyl)-4-methylphenol (5b)


[0173]



[0174] The title compound 5b was prepared according to the method for preparation of compound 3c of Example 3 by replacing 3b with 5a.

(S)-2,5-dimethyl-2,3-dihydrobenzofuran (5c)


[0175]



[0176] The title compound 5c was prepared according to the method for preparation of compound 3d of Example 3 by replacing 3c with 5b.

(S)-2,5-dimethyl-7-nitro-2,3-dihydrobenzofuran (5d)


[0177]



[0178] The title compound 5d was prepared according to the method for preparation of compound 3e of Example 3 by replacing 3d with 5c.

(S)-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (5e)


[0179]



[0180] The title compound 5e was prepared according to the method for preparation of compound 3f of Example 3 by replacing 3e with 5d. MS-ESI (m/z): 164 ($M + 1$)⁺.

(S)-4-bromo-2,5-dimethyl-2,3-dihydrobenzofuran-7-amine (5f)


[0181]



[0182] The title compound 5f was prepared according to the method for preparation of compound 3g of Example 3 by replacing 3f with 5e. MS-ESI (m/z): 242 (M + 1)⁺.

(S)-2,5-dimethyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (5g)


[0183]



[0184] The title compound 5g was prepared according to the method for preparation of compound 3h of Example 3 by replacing 3g with 5f. MS-ESI (m/z): 241 (M + 1)⁺.

(S)-2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (5h)

[0185]




[0186] The title compound 5h was prepared according to the method for preparation of compound 3i of Example 3 by replacing 3h with 5g. MS-ESI (m/z): 247 (M + 1)⁺.

(S)-N-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (5)

[0187]






[0188] The title compound (5) was prepared according to the method for preparation of compound 3 of Example 3 by replacing 3i with 5h. MS-ESI (m/z): 576 (M + 1)⁺.

Example 6

(S)-N-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (6)

[0189]




[0190] Prepared according to the method of Example 4 by replacing (R)-N-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (3) with (S)-N-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (5). The title compound (6) was obtained. MS-ESI (m/z): 590 (M + 1)⁺.

Example 7


(7-(4-(2-(isopropylsulfonyl)phenylamino)-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-6-ylamino)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7)

[0191]



1-(allyloxy)-4-methyl-2-nitrobenzene (7a)


[0192]



[0193] A mixture of 4-methyl-2-nitrophenol (5.0 g, 0.033 mol), 3-bromoprop-1-ene (4.2 g, 0.034 mol) and K_2CO_3 (4.6 g, 0.033 mol) in acetone (20 mL) was stirred for 8 h at ambient temperature. The mixture was poured into water (50 mL), extracted with EtOAc (90 mL), washed with brine, dried and concentrated to give the title compound (7a).

2-allyl-4-methyl-6-nitrophenol (7b)

[0194]




7a

7b

[0195] The compound 1-(allyloxy)-4-methyl-2-nitrobenzene (7a) (6.1 g, 0.032 mol) was stirred for 5 h at 200 °C. The mixture was purified by column chromatography on silica gel eluting with hexanes/EtOAc (50:1) to give the title compound (7b) as a yellow oil.

(5-methyl-7-nitro-2,3-dihydrobenzofuran-2-yl)methanol (7c)


[0196]



[0197] A mixture of 2-allyl-4-methyl-6-nitrophenol (7b) (0.50 g, 2.5 mmol) and *m*-CPBA (0.64 g, 2.5 mmol) in CHCl₃ (9 mL) was stirred for 6 h at 85 °C. The mixture was poured into saturated NaHSO₃ (20 mL), extracted with DCM (40 mL), washed with brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (10:1) to give the title compound (7c) as a solid. MS-ESI (*m/z*): 210 (*M* + 1)⁺.

(7-amino-5-methyl-2,3-dihydrobenzofuran-2-yl)methanol (7d)

[0198]




[0199] A mixture of (5-methyl-7-nitro-2,3-dihydrobenzofuran-2-yl)methanol (7c) (2.3 g, 0.011 mol) and Pd/C (230 mg, 10%) in THF (30 mL) was introduced H₂ and stirred at ambient temperature for 12 h. The mixture was filtered through celite and concentrated to give the title compound (7d) as a solid. MS-ESI (*m/z*): 180 (*M* + 1)⁺.

(7-amino-4-bromo-5-methyl-2,3-dihydrobenzofuran-2-yl)methanol (7e)

[0200]






[0201] To a mixture of (7-amino-5-methyl-2,3-dihydrobenzofuran-2-yl)methanol (7d) (0.80 g, 4.5 mmol) in DMF (13 mL) at 0 °C was added NBS (0.08 g, 4.5 mmol), it was stirred at 0 °C for 15 min. The mixture was extracted with EtOAc (150 mL), washed with NaHSO₃ (50 mL), water, brine, dried and concentrated to give the title compound (7e) as a yellow solid. MS-ESI (m/z): 258 (M + 1)⁺.

(7-amino-5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7f)


[0202]



[0203] A mixture of (7-amino-4-bromo-5-methyl-2,3-dihydrobenzofuran-2-yl)methanol (7e) (1.7 g, 6.6 mmol), pyridin-4-ylboronic acid (1.6 g, 1.3 mmol), Cs₂CO₃ (5.3 g, 16 mmol), Pd(PPh₃)₄ (0.76 g, 0.66 mmol) and water (13 mL) in DMF (60 mL) was stirred at 135 °C under N₂ for 1 h. It was cooled to ambient temperature, poured into water (100 mL), extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1 to 5:1 to 1:1) to give the title compound (7f) as a yellow solid. MS-ESI (m/z): 257 (M + 1)⁺.

(7-amino-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7g)


[0204]



[0205] A mixture of (7-amino-5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7f) (0.10 g, 0.39 mmol), PtO₂ (50 mg, 50%) and TEA (0.061 mL, 0.78 mmol) in HOAc (5 mL) was introduced H₂ and stirred at ambient temperature for 24 h (88 psi). The mixture was filtered, concentrated, and diluted with EtOAc (50 mL). A solution of ammonium hydroxide was added until pH = 10, it was extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated to give the title compound (7g) as a yellow oil. MS-ESI (m/z): 262 (M + 1)⁺.

(7-((4-((2-(isopropylsulfonyl)phenyl)amino)-3-methyl-1H-pyrazolo[3,4-d]pyrimidin-6-yl)amino)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7)

[0206]




[0207] A mixture of (7-amino-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (7g) (40 mg, 0.15 mmol), *N*-(2-(isopropylsulfonyl)phenyl)-3-methyl-6-(methylsulfonyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine (Intermediate A) (75 mg, 0.18 mmol) and *p*-TsOH (29 mg, 0.15 mmol) in *i*-PrOH (0.5 mL) was stirred in a sealed tube at 160 °C for 5 h. The mixture was diluted with EtOAc (20 mL), ammonium hydroxide was added until pH = 10, it was extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (7) as a yellow solid. MS-ESI (m/z): 592 (M + 1)⁺.


Example 8

***N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-6-(2-(methoxymethyl)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (8)**

[0208]

**2-(methoxymethyl)-5-methyl-7-nitro-2,3-dihydrobenzofuran (8a)**


[0209]



[0210] A mixture of (5-methyl-7-nitro-2,3-dihydrobenzofuran-2-yl)methanol (0.70 g, 3.3 mmol) in THF was cooled to 0 °C. NaH was added to the mixture at 0 °C. It was stirred for 15 min at 0 °C. MeI was added dropwise to the mixture at 0 °C. It was stirred for 0.5 h at 0 °C. The mixture was poured into water (20 mL), extracted with EtOAc (40 mL), washed with water, brine, dried and concentrated to give the title compound (8a) as a yellow oil. MS-ESI (m/z): 224 (M + 1)⁺.

2-(methoxymethyl)-5-methyl-2,3-dihydrobenzofuran-7-amine (8b)

[0211]




[0212] A mixture of 2-(methoxymethyl)-5-methyl-7-nitro-2,3-dihydrobenzofuran (8a) (0.85 g, 3.8 mmol) and Pd/C (90 mg, 10%) in THF (10 mL) was introduced H₂ and stirred at ambient temperature for 12 h. The mixture was filtered through celite and concentrated to give the title

compound (8b) as oil. MS-ESI (m/z): 194 (M + 1)⁺.

4-bromo-2-(methoxymethyl)-5-methyl-2,3-dihydrobenzofuran-7-amine (8c)


[0213]



[0214] To a mixture of 2-(methoxymethyl)-5-methyl-2,3-dihydrobenzofuran-7-amine (8b) (0.60 g, 3.1 mmol) in DMF (8 mL) at 0 °C was added NBS (0.55 g, 3.1 mmol). It was stirred at 0 °C for 15 min. The mixture was extracted with EtOAc (40 mL), washed with NaHSO₃ (20 mL), water, brine, dried and concentrated to give the title compound (8c) as a yellow oil. MS-ESI (m/z): 272 (M + 1)⁺.

2-(methoxymethyl)-5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (8d)

[0215]




[0216] A mixture of 4-bromo-2-(methoxymethyl)-5-methyl-2,3-dihydrobenzofuran-7-amine (8c) (0.85 g, 3.1 mmol), pyridin-4-ylboronic acid (0.76 g, 6.2 mmol), Cs₂CO₃ (2.5 g, 7.8 mmol), Pd(PPh₃)₄ (0.18 g, 0.16 mmol) and water (6 mL) in DMF (20 mL) was stirred at 135 °C under N₂ for 1 h. It was cooled to ambient temperature, poured into water (50 mL), extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with hexanes/EtOAc (20:1 to 5:1 to 1:1) to give the title compound (8d) as a yellow oil. MS-ESI (m/z): 271 (M + 1)⁺.

2-(methoxymethyl)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (8e)

[0217]


NIL



[0218] A mixture of 2-(methoxymethyl)-5-methyl-4-(pyridin-4-yl)-2,3-dihydrobenzofuran-7-amine (8d) (0.57 g, 2.1 mmol), PtO_2 (300 mg, 50%) and TEA (0.323 mL, 4.20 mmol) in HOAc (15 mL) was introduced H_2 and stirred at ambient temperature for 24 h (88 psi). The mixture was filtered, concentrated, diluted with EtOAc (50 mL), a solution of ammonium hydroxide was added until pH = 10. It was extracted with EtOAc (150 mL), washed with water, brine, dried and concentrated to give the title compound (8e) as a yellow oil. MS-ESI (m/z): 277 ($M + 1$)⁺.

N-4-(2-(isopropylsulfonyl)phenyl)-N-6-(2-(methoxymethyl)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-3-methyl-1H-pyrazolo[3,4-d]pyrimidine-4,6-diamine (8)

[0219]




[0220] A mixture of (7-amino-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol (8e) (100 mg, 0.36 mmol), N-(2-(isopropylsulfonyl)phenyl)-3-methyl-6-(methylsulfonyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine (Intermediate A) (163 mg, 0.400 mmol) and *p*-TsOH (69 mg, 0.36 mmol) in *i*-PrOH (0.5 mL) was stirred in a sealed tube at 160 °C for 5 h. The mixture was diluted with DCM (20 mL), ammonium hydroxide was added until pH = 10, it was extracted with DCM (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (8) as a yellow solid. MS-ESI (m/z): 606 ($M + 1$)⁺.

Example 9

(R)-5-chloro-N-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine (9)

[0221]




[0222] A mixture of 3i (0.8 g, 3.5 mmol), Intermediate B (1.32 g, 3.8 mmol) and TFA (1.0 g, 8.8 mmol) was stirred in a sealed tube at 130 °C for 5 h . The mixture was diluted with EtOAc (20 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (9) as a brown solid. MS-ESI (m/z): 556 (M + 1)⁺.

Example 10

(R)-5-chloro-N-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine (10)

[0223]




[0224] A mixture of Compound 9 (210 mg, 0.38 mmol), formaldehyde (11.4 mg, 0.38 mmol) and NaBH₃CN (57.3 mg, 0.91 mmol) was stirred at ambient temperature for 0.5 h. The mixture was diluted with NaHCO₃ (10 mL) and EtOAc (20 mL), extracted with EtOAc (60 mL), washed with water, brine, dried and concentrated. The residue was purified by column chromatography on silica gel eluting with DCM/MeOH (10:1) to give the title compound (10) as a yellow solid. MS-ESI (m/z): 570 (M + 1)⁺.

Example 11

(S)-5-chloro-N-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine (11)

[0225]




[0226] Prepared according to the method of Example 9 by replacing 3i with 5h. The title compound (11) was obtained. MS-ESI (m/z): 556 (M + 1)⁺.

Example 12

(S)-5-chloro-N-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine (12)

[0227]






12

[0228] Prepared according to the method of Example 10 by replacing Compound 9 with Compound 11. The title compound (12) was obtained. MS-ESI (m/z): 570 ($M + 1$)⁺.

Example 13

5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine (13)

[0229]




13


[0230] Prepared according to the method of Example 9 by replacing 3i with 5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-amine (1g). The title compound (13) was obtained. MS-ESI (m/z): 542 ($M + 1$)⁺.

Example 14

5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine (14)

[0231]





[0232] Prepared according to the method of Example 10 by replacing Compound 9 with Compound 13. The title compound (14) was obtained. MS-ESI (m/z): 556 ($M + 1$)⁺.

Examples 15-58

[0233] Examples 15-58 (Table 1) were derivatized from Compound 9 or Compound 11 via N-alkylation, acylation, or sulfonylation as shown the following scheme using conditions known in the art. The carbon marked with "*" has chirality as set forth in the chemical structures in Table 1.

























Table 1

Example	Name	Structure	M+1
15	(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone		598
16	(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone		598






Example	Name	Structure	M+1
			
17	(S)-5-chloro-N-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine		634
18	(R)-5-chloro-N-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine		634
19	(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid		614
20	(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid		614
21	(S)-2-amino-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-		627

Example	Name	Structure	M+1
	4-yl)piperidin-1-yl)propan-1-one		
22	(S)-2-amino-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one		627
23	(R)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol		614
24	(S)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol		614
25	(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide		613






Example	Name	Structure	M+1
			
26	(<i>R</i>)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide		613
27	(<i>S</i>)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)- <i>N,N</i> -dimethylacetamide		641
28	(<i>R</i>)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)- <i>N,N</i> -dimethylacetamide		641
29	(<i>S</i>)-5-chloro- <i>N</i> -2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)- <i>N</i> -4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine		662
30	(<i>R</i>)-5-chloro- <i>N</i> -2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)- <i>N</i> -4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-		662







Example	Name	Structure	M+1
	diamine		
31	(S)-5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine		614
32	(R)-5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine		614
33	(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol		600
34	(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol		600


Example	Name	Structure	M+1
35	(<i>R</i>)-1-(4-((<i>S</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol		614
36	(<i>S</i>)-1-(4-((<i>S</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol		614
37	(<i>R</i>)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxyethanone		614
38	(<i>S</i>)-1-(4-((<i>R</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one		628
39	(<i>R</i>)-1-(4-((<i>R</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one		628

Example	Name	Structure	M+1
			
40	(S)-methyl 4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxylate		614
41	(S)-ethyl 2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetate		642
42	(S)-2-amino-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone		613
43	(S)-4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxamide		599
44	(S)-1-(4-(7-((5-chloro-4-((2-		627

Example	Name	Structure	M+1
	(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(methylamino)ethanone		
45	(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(dimethylamino)ethanone		641
46	(S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(piperazin-1-yl)methanone		585
47	(S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(4-methylpiperazin-1-yl)methanone		599
48	(S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(morpholino)methanone		586

Example	Name	Structure	M+1
49	(<i>R</i>)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one		612
50	(<i>R</i>)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methylpropan-1-one		626
51	(<i>R</i>)-4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)(cyclopropyl)methanone		624
52	(<i>R</i>)-5-chloro- <i>N</i> -2-(4-(1-(ethylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)- <i>N</i> -4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine		648
53	(<i>R</i>)-5-chloro- <i>N</i> -4-(2-(isopropylsulfonyl)phenyl)- <i>N</i> -2-(4-(-(isopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine		662

Example	Name	Structure	M+1
			
54	(<i>R</i>)-5-chloro- <i>N</i> -2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)- <i>N</i> -4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine		660
55	(<i>R</i>)-1-(4-((<i>S</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one		642
56	(<i>R</i>)-1-(4-((<i>R</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one		642
57	(<i>S</i>)-1-(4-((<i>R</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one		642
58	(<i>S</i>)-1-(4-((<i>S</i>)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-		642

Example	Name	Structure	M+1
	yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one		

Biological Activity

Inhibition Activity of Cell Proliferation

[0234] SU-DHL-1 cells were seeded in 96-well tissue culture plates. On the next day, cells were exposed to various concentrations of compounds and further cultured for 72 h. Cell proliferation was then determined using Cell Counting Kit-8 [(CCK-8), Dojindo] assay. Briefly, add 10 μ L of the CCK-8 solution to each well of the plate. Incubate the plate for 1-4 hours in the incubator. Measure the absorbance at 450 nm using a multi-well spectrophotometer (SpectraMAX 190, Molecular Devices). The inhibition rate (%) was calculated using the following equation: $[1 - (A_{450} \text{ Compound} / A_{450} \text{ Control})] \times 100\%$. IC₅₀ values were calculated according to inhibition curve by four-parameter fit.

Biological Data for Select Compounds

[0235] Select compounds prepared as described above were assayed according to the biological procedures described herein. The results are given in the table 2.

Table 2

Example	SU-DHL-1 IC ₅₀ (nM)
1	25.5
2	18.9
3	61.0
4	10.9
5	12.1
6	34.6
7	298.1
8	52.7

Example	SU-DHL-1 IC ₅₀ (nM)
9	98.3
10	<0.2
11	<4.1
12	4.0
13	<4.1
14	6.6
15	38.2
16	11.6
17	68.5
18	15.9
19	4.9
20	61.9
21	68.6
22	86.0
23	12.3
24	8.3
25	36.3
26	13.7
27	37.2
28	46.6
29	64.7
30	17.4
31	74.5
32	19.3
33	11.8
34	12.3
35	36.9
36	26.2
37	12.5
38	13.6
39	10.3
40	119.6
41	43.2
42	54.6

Example	SU-DHL-1 IC ₅₀ (nM)
43	106.6
44	263.7
45	39.5
46	114.4
47	117.1
48	218.7
49	97.8
50	81.2
51	52.4
52	59.0
53	109.6
54	333
55	35.7
56	26.3
57	32.8
58	141.0
Crizotinib	75.4-153.8
LDK378*	94.6-339.0
*: LDK378 was prepared as example 7 of WO2008073687.	

in vivo antitumor activity

[0236] Male nude mice (6-week old) were housed in a specific pathogen-free room with a 12 h light/dark schedule at 20-25 °C and were fed an autoclaved chow diet and water ad libitum.

[0237] NCI-H2228 cells at a density of 5.1×10^6 in 200 μ L were first implanted sc into the right flank of each nude mice and then allowed to grow to above 230 mm³, the mice were randomly assigned into control and treatment groups (n = 6 per group for compound treated group, n = 12 per group for vehicle group). Vehicle control group were given vehicle alone, and treatment groups were received Compound 4, LDK378 or Crizotinib as indicated doses via oral administration once daily for 21 days. The mice tumor volumes in each group were measured twice per week.


[0238] The results are given in the Table 3 and Figure 1.

Table 3. Effect on relative tumor volume

Group	Mean relative tumor volume (RTV) $\pm$ S.E.						
	D0	D4	D7	D11	D14	D18	D21
Vehicle Control	100.00 $\pm$ 0.00	151.74 $\pm$ 10.44	173.58 $\pm$ 8.81	214.45 $\pm$ 9.56	229.76 $\pm$ 11.06	241.34 $\pm$ 11.04	275.65 $\pm$ 17.77
Compound 4, 10 mg/kg, QD	100.00 $\pm$ 0.00	52.92 $\pm$ 3.59**	36.55 $\pm$ 5.03**	29.20 $\pm$ 5.18**	18.82 $\pm$ 4.53**	9.48 $\pm$ 2.64**	5.28 $\pm$ 2.35**
LDK378, 20 mg/kg, QD	100.00 $\pm$ 0.00	71.66 $\pm$ 4.52**	56.79 $\pm$ 8.60**	49.09 $\pm$ 8.81**	48.58 $\pm$ 7.88**	35.67 $\pm$ 7.17**	31.44 $\pm$ 7.90**
Crizotinib, 50 mg/kg, QD	100.00 $\pm$ 0.00	67.71 $\pm$ 3.20**	54.10 $\pm$ 5.56**	44.71 $\pm$ 5.98**	42.69 $\pm$ 6.12**	41.44 $\pm$ 6.35**	43.69 $\pm$ 6.82**

[0239] The following numbered paragraphs define particular embodiments of the present disclosure:

1. At least one compound of formula (I):



and/or at least one pharmaceutically acceptable salt thereof, wherein:

each R^1 is independently selected from:

halogen,

hydroxyl,

C_{1-10} alkyl,

C_{3-10} cycloalkyl,

C_{3-10} cycloalkyl-alkyl,

heterocyclyl,

heterocyclylalkyl

aryl,

arylalkyl,

heteroaryl, and

heteroarylalkyl,

wherein alkyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent independently selected from R^{6a} , and wherein aryl and heteroaryl are each unsubstituted or substituted with at least one substituent independently selected from R^{6b} ;

each R^2 is independently selected from:

hydrogen,

halogen,

hydroxyl,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl,

C_{3-10} cycloalkyl,

C_{3-10} cycloalkylalkyl,

heterocyclyl,

heterocyclylalkyl

heterocyclylcarbonyl,

aryl,

heteroaryl,

arylalkyl, and

heteroarylalkyl,

wherein alkyl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent independently selected from R^{6a} , and each aryl and heteroaryl is unsubstituted or substituted with at least one substituent independently selected from R^{6b} ;

each R^3 is independently selected from:

hydrogen,

halogen,

-CN,

-NR⁷R⁸, and

C₁₋₁₀ alkyl;

wherein alkyl is unsubstituted or substituted with at least one substituent independently selected from R^{6a};

each R⁴ is independently selected from:

hydrogen,

halogen,

-CN,

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl, and

C₃₋₁₀ cycloalkyl;

wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and C₃₋₁₀ cycloalkyl are each unsubstituted or substituted with at least one substituent independently selected from R^{6a};

or R³ and R⁴ together with the carbon atoms to which they are attached form a 5-6 membered ring containing 0, 1, 2 or 3 heteroatoms independently selected from oxygen, sulfur and nitrogen, and optionally substituted with 1-2 R^{6b} groups;

each R⁵ is independently selected from:

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-O(CR⁹R¹⁰)_tNR⁷R⁸,

-C(O)R⁷,

-CO₂R⁸,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸,

-OC(O)R⁷,

-CN,

-C(O)NR⁷R⁸,

-NR⁷C(O)R⁸,

-OC(O)NR⁷R⁸,

-NR⁷C(O)OR⁸,

-NR⁷C(O)NR⁷R⁸,

-CR⁷(N-OR⁸),

-CHF₂,

-CF₃,

-OCHF₂, and

-OCF₃;

wherein C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl, and C₃₋₁₀ cycloalkyl are each unsubstituted or substituted with at least one substituent independently selected from R^{6a};

each R^{6a} is independently selected from:

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,


-NR⁷R⁸,

-(CR⁹R¹⁰)_tOR⁸,

-(CR⁹R¹⁰)_tNR⁷R⁸,

-(CR⁹R¹⁰)_tSR⁸,

-(CR⁹R¹⁰)_tS(O)_rR⁸,



-CHF₂,

-CF₃,

-OCHF₂, and

-OCF₃;

each R^{6b} is independently selected from:

R^{6a},

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl, and

heteroaryl-C₁₋₄ alkyl;

each R⁷ and each R⁸ are independently selected from:

hydrogen,

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

cycloalkyl-C₁₋₁₀ alkyl;

heterocyclyl,

heterocyclyl-C₁₋₁₀ alkyl,

aryl,

heteroaryl,

aryl-C₁₋₁₀ alkyl, and

heteroaryl-C₁₋₁₀ alkyl;

wherein alkyl, alkenyl, alkynyl, cycloalkyl, and heterocyclyl are each unsubstituted or substituted with at least one substituent independently selected from R^{6a}, and aryl and heteroaryl are each unsubstituted or substituted with at least one substituent

independently selected from R^{6b}; or

R⁷ and R⁸ together with the atom(s) to which they are attached form a heterocyclic ring of 4 to 7 members containing 0, 1, or 2 additional heteroatoms independently selected from oxygen, sulfur and NR¹¹, each R⁷ and R⁸ may be unsubstituted or substituted on a carbon or nitrogen atom with at least one substituent independently selected from R¹²;

each R⁹ and each R¹⁰ are independently selected from:

hydrogen,

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

cycloalkyl-C₁₋₁₀ alkyl,

heterocyclyl,

heterocyclyl-C₁₋₁₀ alkyl,

aryl,

heteroaryl,

aryl-C₁₋₁₀ alkyl, and

heteroaryl-C₁₋₁₀ alkyl; or

R⁹ and R¹⁰ together with the carbon to which they are attached form a ring of 3 to 7 members containing 0, 1, or 2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

each R¹¹ is independently selected from:

hydrogen,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl,

heterocyclyl,

heterocyclyl-C₁₋₄ alkyl,

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl,

heteroaryl-C₁₋₄ alkyl,

-S(O)_rR⁷,

-C(O)R⁷,

-CO₂R⁷,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸, and

-C(O)NR⁷R⁸;

each R¹² is independently selected from:

halogen,

C₁₋₁₀ alkyl,

C₃₋₁₀ cycloalkyl,

C₃₋₁₀ cycloalkylalkyl,

heterocyclyl,

heterocyclylalkyl,

aryl,

aryl-C₁₋₄ alkyl,

heteroaryl,

heteroaryl-C₁₋₄ alkyl,

-OR⁷,

-NR⁷S(O)_rR⁸,

-S(O)_rR⁷,

-SR⁷,

-S(O)₂OR⁷,

-OS(O)₂R⁷,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-O(CR⁹R¹⁰)_tNR⁷R⁸,

-C(O)R⁷,

-CO₂R⁸,

-CO₂(CR⁹R¹⁰)_tCONR⁷R⁸,

-OC(O)R⁷,

-CN,

-C(O)NR⁷R⁸,

-NR⁷C(O)R⁸,

-OC(O)NR⁷R⁸,

-NR⁷C(O)OR⁸,

-NR⁷C(O)NR⁷R⁸,

-CHF₂,

-CF₃,

-OCHF₂, and

-OCF₃;

each m is independently selected from 0, 1 and 2;

each n is independently selected from 1, 2, and 3;

each p is independently selected from 0, 1, 2, and 3;

each q is independently selected from 0, 1, 2, and 3;

each r is independently selected from 1 and 2;

each t is independently selected from 1, 2, and 3.

2. 2. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^1 is hydrogen or C_{1-10} alkyl, wherein alkyl is unsubstituted or substituted with at least one substituent independently selected from hydroxyl and C_{1-10} alkoxy.
3. 3. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein m is 0.
4. 4. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^1 is C_{1-10} alkyl, wherein alkyl is unsubstituted or substituted with at least one substituent independently selected from hydroxyl and C_{1-10} alkoxy.
5. 5. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein m is 1.
6. 6. At least one compound of paragraph 5 and/or at least one pharmaceutically acceptable salt thereof, wherein R^1 is methyl, hydroxymethyl, or methoxymethyl.
7. 7. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein each R^2 is independently selected from C_{1-10} alkyl, heterocyclylcarbonyl, and heterocyclyl, wherein alkyl and heterocyclyl are independently unsubstituted or substituted with at least one substituent independently selected from R^{6a} , and p is selected from 0, 1, 2, and 3.
8. 8. At least one compound of paragraph 7 and/or at least one pharmaceutically acceptable salt thereof, wherein each R^2 is independently selected from C_{1-10} alkyl and piperidinyl, and p is 2.
9. 9. At least one compound of paragraph 8 and/or at least one pharmaceutically acceptable salt thereof, wherein each R^2 is independently selected from methyl and 4-piperidinyl.
10. 10. At least one compound of paragraph 9 and/or at least one pharmaceutically acceptable salt thereof, wherein each R^2 is independently selected from methyl and 4-piperidinyl, wherein 4-piperidinyl is unsubstituted or substituted with at least one substituent independently selected from R^{6a} , wherein R^{6a} is independently selected from C_{1-10} alkyl, $-C(O)R^7$, $-S(O)_rR^7$, $-(CR^9R^{10})_tCO_2R^8$, $-C(O)(CR^9R^{10})_tOR^8$, $-C(O)(CR^9R^{10})_tNR^7R^8$, $-(CR^9R^{10})_tOR^8$, $-(CR^9R^{10})_tCONR^7R^8$, $-(CR^9R^{10})_tS(O)_rR^8$, $-CO_2R^8$, $-C(O)NR^7R^8$, $-(CR^9R^{10})_tOR^8$; wherein R^7 , R^8 , R^9 , R^{10} , t and r are as described in paragraph 1.
11. 11. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^3 is hydrogen.

12. 12. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^4 is independently selected from hydrogen, C_{1-10} alkyl, and halogen.
13. 13. At least one compound of paragraph 12 and/or at least one pharmaceutically acceptable salt thereof, wherein R^4 is independently selected from halogen.
14. 14. At least one compound of paragraph 13 and/or at least one pharmaceutically acceptable salt thereof, wherein R^4 is chlorine.
15. 15. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^3 and R^4 together with the carbon atoms to which they are attached form a 5 membered ring containing 2 nitrogen atoms, wherein the 5-membered ring is optionally substituted with 1-2 C_{1-10} alkyl.
16. 16. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^3 and R^4 together with the carbon atoms to which they are attached form pyrazolo.
17. 17. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein R^5 is $-S(O)_rR^7$.
18. 18. At least one compound of paragraph 17 and/or at least one pharmaceutically acceptable salt thereof, wherein R^5 is $-S(O)_2R^7$, wherein R^7 is C_{1-10} alkyl, and q is 1.
19. 19. At least one compound of paragraph 18 and/or at least one pharmaceutically acceptable salt thereof, wherein R^5 is $-S(O)_2R^7$, wherein R^7 is isopropyl.
20. 20. At least one compound of paragraph 1 and/or at least one pharmaceutically acceptable salt thereof, wherein n is 1.
21. 21. At least one compound selected from:

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*S*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*S*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(7-(4-(2-(isopropylsulfonyl)phenylamino)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-6-ylamino)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-2-yl)methanol;

N-4-(2-(isopropylsulfonyl)phenyl)-*N*-6-(2-(methoxymethyl)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidine-4,6-diamine;

(*R*)-5-chloro-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*R*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*S*)-5-chloro-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*S*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(*S*)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(*R*)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(*S*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*R*)-5-chloro-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(*S*)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid;

(*R*)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetic acid;

(*S*)-2-amino-1-(4-((*S*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(*S*)-2-amino-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(*R*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(*S*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-

yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamide;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-N,N-dimethylacetamide;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-N,N-dimethylacetamide;

(S)-5-chloro-N-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(R)-5-chloro-N-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(S)-5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(R)-5-chloro-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(R)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

(S)-2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

(R)-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(S)-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxyethanone;

(S)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one;

(R)-1-(4-((R)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-one;

(S)-methyl 4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxylate;

(S)-ethyl 2-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetate;

(S)-2-amino-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanone;

(S)-4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidine-1-carboxamide;

(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(methylamino)ethanone;

(S)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(dimethylamino)ethanone;

(S)-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(piperazin-1-yl)methanone;

(S)-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(4-methylpiperazin-1-yl)methanone;

(S)-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)(morpholino)methanone;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-one;

(R)-1-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methylpropan-1-one;

(R)-(4-(7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)(cyclopropyl)methanone;

(R)-5-chloro-*N*-2-(4-(1-(ethylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(R)-5-chloro-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(4-(1-(isopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidine-2,4-diamine;

(R)-5-chloro-*N*-2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidine-2,4-diamine;

(R)-1-(4-((S)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one;

(*R*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one;

(*S*)-1-(4-((*R*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one; (*S*)-1-(4-((*S*)-7-((5-chloro-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-one, and pharmaceutically acceptable salts thereof.

22. 22. A pharmaceutical composition, comprising at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.
23. 23. A method for modulating anaplastic lymphoma kinase, comprising administering to a subject in need thereof a therapeutically effective amount of at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof, or at least one pharmaceutical composition thereof.
24. 24. A method to treat, ameliorate or prevent a condition, which responds to inhibition of anaplastic lymphoma kinase, comprising administering to a subject in need of such treatment an effective amount of at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof, or of at least one pharmaceutical composition thereof, and optionally in combination with a second therapeutic agent.
25. 25. A method for making a medicament for treating a condition, which responds to inhibition of anaplastic lymphoma kinase, comprising including to the medicament at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof.
26. 26. A method for treating a cell proliferative disorder, comprising administering to a subject in need of such treatment an effective amount of at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof or of at least one pharmaceutical composition thereof, and optionally in combination with a second therapeutic agent.
27. 27. A method for making a medicament for treating a cell-proliferative disorder, comprising including in the medicament at least one compound of paragraph 1, and/or at least one pharmaceutically acceptable salt thereof.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- [WO2005016894A](#) [\[0004\]](#)
- [WO2008051547A](#) [\[0005\]](#)
- [WO2008073687A](#) [\[0235\]](#)

Non-patent literature cited in the description

- **PULFORD et al.** Cell. Mol. Life. Sci., 2004, vol. 61, 2939-2953 [\[0002\]](#)
- **T. W. GREENE** Protective Groups in Organic Synthesis John Wiley & Sons 19910000 [\[0046\]](#)
- **DUYSTER et al.** Oncogene, 2001, vol. 20, 5623-5637 [\[0070\]](#)
- **J. WOOD et al.** Cancer Res., 2000, vol. 60, 2178-2189 [\[0071\]](#)
- **E WEISBERG et al.** Cancer Cell, 2002, vol. 1, 433-443 [\[0072\]](#)
- **T IDZIOREK et al.** J. Immunol. Methods, 1995, vol. 185, 249-258 [\[0073\]](#)
- **WG DIRKS et al.** Int. J. Cancer, 2002, vol. 100, 49-56 [\[0076\]](#) [\[0076\]](#)
- **T.W. GREENE** Protecting Groups in Organic Synthesis John Wiley & Sons, Inc. 19990000 [\[0099\]](#)
- **JEAN JACQUES ANDRE COLLET** **SAMUEL H. WILEN** Enantiomers, Racemates and Resolutions John Wiley & Sons, Inc. 19810000 [\[0101\]](#)
- **T.W. GREENE** **G. M. WUTS** Protective Groups in Organic Chemistry John Wiley and Sons 19910000 [\[0107\]](#)
- J. Med. Chem., 1990, vol. 33, 2174-2178 [\[0110\]](#)
- J. Med. Chem., 2002, vol. 45, 2229-2239 [\[0116\]](#)

PATENTKRAV

1. Forbindelse med formelen (I):



eller et farmaceutisk acceptabelt salt deraf, hvor:

hver R¹ uafhængigt er valgt blandt:

hydrogen,
 halogen,
 hydroxyl,
 C₁₋₁₀ alkyl,
 C₃₋₁₀ cycloalkyl,
 C₃₋₁₀ cycloalkyl-alkyl,
 heterocyclyl,
 heterocyclylalkyl,
 aryl,
 arylalkyl,
 heteroaryl og
 hetero arylalkyl,

hvor alkyl, cycloalkyl og heterocyclyl hver er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a}, og hvor aryl og heteroaryl

hver er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6b} ;

hver R^2 uafhængigt er valgt blandt C_{1-10} alkyl og piperidiny, hvor C_{1-10} alkyl og piperidiny hver uafhængigt er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a} ;

hver R^3 uafhængigt er valgt blandt:

hydrogen,

halogen,

-CN,

- NR^7R^8 og

C_{1-10} alkyl;

hvor alkyl er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a} ;

hver R^4 uafhængigt er valgt blandt:

hydrogen,

halogen,

-CN,

C_{1-10} alkyl,

C_{2-10} alkenyl,

C_{2-10} alkynyl og

C_{3-10} cycloalkyl;

hvor C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, og C_{3-10} cycloalkyl hver er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a} ;

eller R^3 og R^4 sammen med kulstofatomerne, til hvilke de er bundet, danner en 5-til 6-leddet ring indeholdende 0, 1, 2 eller 3 heteroatomer uafhængigt valgt blandt oxygen, svovl og nitrogen, og valgfrit substitueret med 1-2 R^{6b} grupper;

hver R^5 uafhængigt er valgt blandt:

- | | |
|----|---|
| | C_{1-10} alkyl, |
| | C_{2-10} alkenyl, |
| 5 | C_{2-10} alkynyl, |
| | C_{3-10} cycloalkyl, |
| | -OR ⁸ , |
| | -NR ⁷ S(O) _r R ⁸ , |
| | -NO ₂ , |
| 10 | halogen, |
| | -S(O) _r R ⁷ , |
| | -SR ⁸ , |
| | -S(O) ₂ OR ⁷ , |
| | -OS(O) ₂ R ⁸ , |
| 15 | -S(O) _r NR ⁷ R ⁸ , |
| | -NR ⁷ R ⁸ , |
| | -O(CR ⁹ R ¹⁰) _i NR ⁷ R ⁸ , |
| | -C(O)R ⁷ , |
| | -CO ₂ R ⁸ , |
| 20 | -CO ₂ (CR ⁹ R ¹⁰) _i CONR ⁷ R ⁸ , |
| | -OC(O)R ⁷ , |
| | -CN, |
| | -C(O)NR ⁷ R ⁸ , |
| | -NR ⁷ C(O)R ⁸ , |
| 25 | -OC(O)NR ⁷ R ⁸ , |
| | -NR ⁷ C(O)OR ⁸ , |
| | -NR ⁷ C(O)NR ⁷ R ⁸ , |
| | -CR ⁷ (N-OR ⁸), |
| | -CHF ₂ , |
| 30 | -CF ₃ , |
| | -OCHF ₂ og |

-OCF₃;

hvor C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₂₋₁₀ alkynyl og C₃₋₁₀ cycloalkyl hver er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a};

hver R^{6a} uafhængigt er valgt blandt:

C₁₋₁₀ alkyl,

C₂₋₁₀ alkenyl,

C₂₋₁₀ alkynyl,

C₃₋₁₀ cycloalkyl,

-OR⁸,

-NR⁷S(O)_rR⁸,

-NO₂,

halogen,

-S(O)_rR⁷,

-SR⁸,

-S(O)₂OR⁷,

-OS(O)₂R⁸,

-S(O)_rNR⁷R⁸,

-NR⁷R⁸,

-(CR⁹R¹⁰)_tOR⁸,

-(CR⁹R¹⁰)_tNR⁷R⁸,

-(CR⁹R¹⁰)_tSR⁸,

-(CR⁹R¹⁰)_tS(O)_rR⁸,

-(CR⁹R¹⁰)_tCO₂R⁸,

-(CR⁹R¹⁰)_tCONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷CO₂R⁸,

-(CR⁹R¹⁰)_tOCONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷CONR⁷R⁸,

-(CR⁹R¹⁰)_tNR⁷SO₂NR⁷R⁸,

- 5
- O(CR⁹R¹⁰)_tNR⁷R⁸,
 - C(O)R⁷,
 - C(O)(CR⁹R¹⁰)_tOR⁸,
 - C(O)(CR⁹R¹⁰)_tNR⁷R⁸,
 - C(O)(CR⁹R¹⁰)_tSR⁸,
 - C(O)(CR⁹R¹⁰)_tS(O)_rR⁸,
 - CO₂R⁸,
 - CO₂(CR⁹R¹⁰)_tCONR⁷R⁸,
 - OC(O)R⁷,
 - 10 -CN,
 - C(O)NR⁷R⁸,
 - NR⁷C(O)R⁸,
 - OC(O)NR⁷R⁸,
 - NR⁷C(O)OR⁸,
 - 15 -NR⁷C(O)NR⁷R⁸,
 - CR⁷(N-OR⁸),
 - CHF₂,
 - CF₃,
 - OCHF₂ og
 - 20 -OCF₃;

hver R^{6b} uafhængigt er valgt blandt:

- 25
- R^{6a},
 - aryl,
 - aryl-C₁₋₄ alkyl,
 - heteroaryl og
 - heteroaryl-C₁₋₄ alkyl;

- 30
- hver R⁷ og hver R⁸ are uafhængigt valgt blandt:

hydrogen,
 C₁₋₁₀ alkyl,
 C₂₋₁₀ alkenyl,
 C₂₋₁₀ alkynyl,
 5 C₃₋₁₀ cycloalkyl,
 cycloalkyl-Ci-io alkyl;
 heterocyclyl,
 heterocyclyl-Ci-io alkyl,
 aryl,
 10 heteroaryl,
 aryl-C₁₋₁₀ alkyl og
 heteroaryl-Ci-io alkyl;

hvor alkyl, alkenyl, alkynyl, cycloalkyl og heterocyclyl hver er ikke-substitueret eller
 15 substitueret med mindst en substituent uafhængigt valgt blandt R^{6a}, og aryl og
 heteroaryl hver er ikke-substitueret eller substitueret med mindst en substituent
 uafhængigt valgt blandt R^{6b}; eller

R⁷ og R⁸ sammen med atomet/atomerne, til hvilke de er bundet, danner en 4- til
 7-leddet heterocyklisk ring indeholdende 0, 1 eller 2 yderligere heteroatomer, som
 20 uafhængigt er valgt blandt oxygen, svovl og NR¹¹, hver R⁷ og R⁸ kan være ikke-
 substitueret eller substitueret på et kulstof- eller nitrogenatom med mindst en
 substituent uafhængigt valgt blandt R¹²;
 hver R⁹ og hver R¹⁰ uafhængigt er valgt blandt:

25 hydrogen,
 C₁₋₁₀ alkyl,
 C₂₋₁₀ alkenyl,
 C₂₋₁₀ alkynyl,
 C₃₋₁₀ cycloalkyl,
 30 cycloalkyl-Ci-io alkyl,
 heterocyclyl,

heterocyclyl-Ci-io alkyl,
 aryl,
 heteroaryl,
 aryl-C₁₋₁₀ alkyl og
 heteroaryl-Ci-io alkyl; eller

5

R⁹ og R¹⁰ sammen med kulstoffet, til hvilket de er bundet, danner en 3- til 7-leddet ring indeholdende 0, 1 eller 2 heteroatomer, som uafhængigt er valgt blandt oxygen, svovl og nitrogen;
 hver R¹¹ uafhængigt er valgt blandt:

10

hydrogen,
 C₁₋₁₀ alkyl,
 C₃₋₁₀ cycloalkyl,
 C₃₋₁₀ cycloalkyl-C₁₋₄ alkyl,
 heterocyclyl,
 heterocyclyl-C₁₋₄ alkyl,
 aryl,
 aryl-C₁₋₄ alkyl,
 heteroaryl,
 heteroaryl-C₁₋₄ alkyl,
 -S(O)_rR⁷,
 -C(O)R⁷,
 -CO₂R⁷,
 -CO₂(CR⁹R¹⁰)_tCONR⁷R⁸ og
 -C(O)NR⁷R⁸;

15

20

25

hver R¹² uafhængigt er valgt blandt:

30

halogen,
 C₁₋₁₀ alkyl,

	C ₃₋₁₀ cycloalkyl,
	C ₃₋₁₀ cycloalkylalkyl,
	heterocyclyl,
	heterocyclylalkyl,
5	aryl,
	aryl-C ₁₋₄ alkyl,
	heteroaryl,
	heteroaryl-C ₁₋₄ alkyl,
	-OR ⁷ ,
10	-NR ⁷ S(O) _r R ⁸ ,
	-S(O) _r R ⁷ ,
	-SR ⁷ ,
	-S(O) ₂ OR ⁷ ,
	-OS(O) ₂ R ⁷ ,
15	-S(O) _r NR ⁷ R ⁸ ,
	-NR ⁷ R ⁸ ,
	-O(CR ⁹ R ¹⁰) _i NR ⁷ R ⁸ ,
	-C(O)R ⁷ ,
	-CO ₂ R ⁸ ,
20	-CO ₂ (CR ⁹ R ¹⁰) _i CONR ⁷ R ⁸ ,
	-OC(O)R ⁷ ,
	-CN,
	-C(O)NR ⁷ R ⁸ ,
	-NR ⁷ C(O)R ⁸ ,
25	-OC(O)NR ⁷ R ⁸ ,
	-NR ⁷ C(O)OR ⁸ ,
	-NR ⁷ C(O)NR ⁷ R ⁸ ,
	-CHF ₂ ,
	-CF ₃ ,
30	-OCHF ₂ og
	-OCF ₃ ;

hver m uafhængigt er valgt blandt 0, 1 og 2;
hver n uafhængigt er valgt blandt 1, 2 og 3;
p er 2;
hver q uafhængigt er valgt blandt 0, 1, 2 og 3;
5 hver r uafhængigt er valgt blandt 1 og 2;
hver t uafhængigt er valgt blandt 1, 2 og 3.

2. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor R^1 er hydrogen eller C_{1-10} alkyl, hvor alkyl er ikke-substitueret eller substitueret med
10 mindst en substituent uafhængigt valgt blandt hydroxyl og C_{1-10} alkoxy.

3. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor m er 0 eller 1; for eksempel hvor m er 1, og R^1 er methyl, hydroxymethyl eller methoxymethyl.

15 4. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor hver R^2 uafhængigt er valgt blandt methyl og 4-piperidiny; for eksempel hvor 4-piperidiny er ikke-substitueret eller substitueret med mindst en substituent uafhængigt valgt blandt R^{6a} , hvor R^{6a} uafhængigt er valgt blandt C_{1-10} alkyl,
20 $-C(O)R^7$, $-S(O)_rR^7$, $-(CR^9R^{10})_tCO_2R^8$, $-C(O)(CR^9R^{10})_tOR^8$, $-C(O)(CR^9R^{10})_tNR^7R^8$,
 $-(CR^9R^{10})_tOR^8$, $-(CR^9R^{10})_tCONR^7R^8$, $-(CR^9R^{10})_tS(O)_rR^8$, $-CO_2R^8$, $-C(O)NR^7R^8$,
 $-(CR^9R^{10})_tOR^8$; hvor R^7 , R^8 , R^9 , R^{10} , t og r er som beskrevet i krav 1.

5. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor R^3
25 er hydrogen.

6. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor R^4 uafhængigt er valgt blandt hydrogen, C_{1-10} alkyl og halogen, for eksempel chlor.

30 7. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor R^3 og R^4 sammen med kulstofatomerne, til hvilke de er bundet, danner en 5-leddet

ring indeholdende 2 nitrogenatomer, for eksempel pyrazolo, hvor den 5-ledede ring valgfrit er substitueret med 1-2 C₁₋₁₀ alkyl.

8. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor R⁵ er -S(O)_qR⁷.

9. Forbindelse ifølge krav 8 eller et farmaceutisk acceptabelt salt deraf, hvor R⁵ er -S(O)₂R⁷, hvor R⁷ er C₁₋₁₀ alkyl, for eksempel isopropyl, og q er 1.

10. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf, hvor n er 1.

11. Forbindelse ifølge krav 1 valgt blandt:

N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;
N-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-*N*-6-(5-methyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;

(*R*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;

(*R*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;

(*S*)-*N*-6-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;

(*S*)-*N*-6-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)-3-methyl-1*H*-pyrazolo[3,4-*d*]pyrimidin-4,6-diamin;

(7-(4-(2-(isopropylsulfonyl)phenylamino)-3-methyl-1*H*-pyrazolo[3,4-
d]pyrimidin-6-ylamino)-5-methyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-
2-yl)methanol;

N-4-(2-(isopropylsulfonyl)phenyl)-*N*-6-(2-(methoxymethyl)-5-methyl-4-
(piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-3-methyl-1*H*-pyrazolo[3,4-
d]pyrimidin-4,6-diamin;

(*R*)-5-chlor-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-
yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;

(*R*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-
dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-
diamin;

(*S*)-5-chlor-*N*-2-(2,5-dimethyl-4-(piperidin-4-yl)-2,3-dihydrobenzofuran-7-
yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;

(*S*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)-2,3-
dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-
diamin;

5-chlor-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(piperidin-4-yl)-
2,3-dihydrobenzofuran-7-yl)pyrimidin-2,4-diamin;

5-chlor-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(5-methyl-4-(1-
methylpiperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)pyrimidin-2,4-diamin;

(*S*)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-
yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanon;

(*R*)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-
yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanon;

(*S*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-
dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-
diamin;

(*R*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)-2,3-
dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-
diamin;

(*S*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)eddikesyre;

(*R*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)eddikesyre;

(*S*)-2-amino-1-(4-((*S*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-on;

(*S*)-2-amino-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-on;

(*R*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(*S*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(*S*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamid;

(*R*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetamid;

(*S*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-*N,N*-dimethylacetamid;

(*R*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-*N,N*-dimethylacetamid;

(*S*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;

(*R*)-5-chlor-*N*-2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-2,3-dihydrobenzofuran-7-yl)-*N*-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;

5 (*S*)-5-chlor-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidin-2,4-diamin;

(*R*)-5-chlor-*N*-4-(2-(isopropylsulfonyl)phenyl)-*N*-2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidin-2,4-diamin;

10 (*R*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

(*S*)-2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanol;

15 (*R*)-1-(4-((*S*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

(*S*)-1-(4-((*S*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-2-ol;

20 (*R*)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxyethanon;

(*S*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-

25 hydroxypropan-1-on;

(*R*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-hydroxypropan-1-on;

30 (*S*)-methyl 4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-carboxylat;

- (S)-ethyl 2-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)acetat;
 (S)-2-amino-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)ethanon;
 5 (S)-4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-carboxamid
 (S)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(methylamino)ethanon;
 10 (S)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-(dimethylamino)ethanon;
 (R)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)propan-1-on;
 15 (R)-1-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methylpropan-1-on;
 20 (R)-(4-(7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)(cyclopropyl)methanon;
 (R)-5-chlor-N-2-(4-(1-(ethylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;
 25 (R)-5-chlor-N-4-(2-(isopropylsulfonyl)phenyl)-N-2-(4-(1-(isopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)pyrimidin-2,4-diamin;
 (R)-5-chlor-N-2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2,5-dimethyl-2,3-dihydrobenzofuran-7-yl)-N-4-(2-(isopropylsulfonyl)phenyl)pyrimidin-2,4-diamin;
 30

(*R*)-1-(4-((*S*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-on;

(*R*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-on;

(*S*)-1-(4-((*R*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-on;

(*S*)-1-(4-((*S*)-7-((5-chlor-4-((2-(isopropylsulfonyl)phenyl)amino)pyrimidin-2-yl)amino)-2,5-dimethyl-2,3-dihydrobenzofuran-4-yl)piperidin-1-yl)-2-methoxypropan-1-on og farmaceutiske acceptable salte deraf.

12. Farmaceutisk sammensætning, omfattende en forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf og mindst et farmaceutisk acceptabelt bærestof.

13. Forbindelse ifølge krav 1 eller et farmaceutisk acceptabelt salt deraf eller en farmaceutisk sammensætning deraf til anvendelse ved

a. modulering af anaplastisk lymfom kinase; eller

b. behandling, forbedring eller forebyggelse af en tilstand, som reagerer på inhibering af anaplastisk lymfom kinase, valgfrit i kombination med et andet terapeutisk middel; eller

c. behandling af en celleproliferativ forstyrrelse, valgfrit i kombination med et andet terapeutisk middel.

DRAWINGS




Figure 1