

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number
WO 00/76517 A1

- (51) International Patent Classification⁷: **A61K 31/47**,
31/495, 31/38, 31/17 King of Prussia, PA 19406 (US). **BENSON, Gregory, M.**
[GB/GB]; Nazeing, Essex (GB).
- (21) International Application Number: PCT/US00/16814 (74) Agents: **SIMON, Soma, G.** et al.; SmithKline Beecham
Corporation, Corporate Intellectual Property, UW2220,
709 Swedeland Road, P.O. Box 1539, King of Prussia, PA
19406-0939 (US).
- (22) International Filing Date: 16 June 2000 (16.06.2000)
- (25) Filing Language: English (81) Designated States (*national*): AE, AL, AU, BA, BB, BG,
BR, CA, CN, CZ, DZ, EE, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK,
MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, TZ,
UA, US, UZ, VN, YU, ZA.
- (26) Publication Language: English
- (30) Priority Data: 60/139,677 16 June 1999 (16.06.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*):
SMITHKLINE BEECHAM CORPORATION
[US/US]; One Franklin Plaza, Philadelphia, PA 19103
(US).
- (72) Inventors; and
(75) Inventors/Applicants (*for US only*): **BRYAN, Deborah**,
L. [US/US]; 341 MacKenzie Drive, West Chester, PA
19380 (US). **GLEASON, John, G.** [CA/US]; 8 Heron Hill
Drive, Downingtown, PA 19335 (US). **WIDDOWSON**,
Katherine, L. [CA/US]; 1047 Old Valley Forge Road,
- Published:**
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: IL-8 RECEPTOR ANTAGONISTS

(57) Abstract: This invention relates to novel compounds of Formula (I), and compositions thereof, useful in the treatment of disease states mediated by the chemokine, Interleukin-8 (IL-8).

WO 00/76517 A1

IL-8 RECEPTOR ANTAGONISTS

FIELD OF THE INVENTION

This invention relates to novel guandine substituted compounds,
5 pharmaceutical compositions, processes for their preparation, and use thereof in
treating IL-8, GRO α , GRO β , GRO γ , ENA-78 and NAP-2 mediated diseases.

BACKGROUND OF THE INVENTION

Many different names have been applied to Interleukin-8 (IL-8), such as
neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil
10 chemotactic factor (MDNCF), neutrophil activating factor (NAF), and T-cell
lymphocyte chemotactic factor. Interleukin-8 is a chemoattractant for neutrophils,
basophils, and a subset of T-cells. It is produced by a majority of nucleated cells
including macrophages, fibroblasts, endothelial and epithelial cells exposed to TNF,
IL-1 α , IL-1 β or LPS, and by neutrophils themselves when exposed to LPS or
15 chemotactic factors such as FMLP. M. Baggiolini et al, J. Clin. Invest. 84, 1045
(1989); J. Schroder et al, J. Immunol. 139, 3474 (1987) and J. Immunol. 144, 2223
(1990) ; Strieter, et al, Science 243, 1467 (1989) and J. Biol. Chem. 264, 10621
(1989); Cassatella et al, J. Immunol. 148, 3216 (1992).

GRO α , GRO β , GRO γ and NAP-2 also belong to the chemokine α family.
20 Like IL-8 these chemokines have also been referred to by different names. For
instance GRO α , β , γ have been referred to as MGSA α , β and γ respectively
(Melanoma Growth Stimulating Activity), see Richmond et al, J. Cell Physiology
129, 375 (1986) and Chang et al, J. Immunol 148, 451 (1992). All of the chemokines
of the α -family which possess the ELR motif directly preceding the CXC motif bind
25 to the IL-8 β receptor.

IL-8, GRO α , GRO β , GRO γ and NAP-2 stimulate a number of functions in
vitro. They have all been shown to have chemoattractant properties for neutrophils,
while IL-8 and GRO α have demonstrated T-lymphocytes, and basophiles
chemotactic activity. In addition IL-8 can induce histamine release from basophils
30 from both normal and atopic individuals GRO- α and IL-8 can in addition, induce

lysozomal enzyme release and respiratory burst from neutrophils. IL-8 has also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils without de novo protein synthesis. This may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many known diseases are characterized by massive neutrophil infiltration. As IL-8, GRO α , GRO β , GRO γ and NAP-2 promote the accumulation and activation of neutrophils, these chemokines have been implicated in a wide range of acute and chronic inflammatory disorders including psoriasis and rheumatoid arthritis, Baggiolini et al, FEBS Lett. **307**, 97 (1992); Miller et al, Crit. Rev. Immunol. **12**, 17 (1992); Oppenheim et al, Annu. Rev. Immunol. **9**, 617 (1991); Seitz et al., J. Clin. Invest. **87**, 463 (1991); Miller et al., Am. Rev. Respir. Dis. **146**, 427 (1992); Donnelly et al., Lancet **341**, 643 (1993). In addition the ELR chemokines (those containing the amino acids ELR motif just prior to the CXC motif) have also been implicated in angiostasis. Strieter et al, Science **258**, 1798 (1992).

In vitro, IL-8, GRO α , GRO β , GRO γ and NAP-2 induce neutrophil shape change, chemotaxis, granule release, and respiratory burst, by binding to and activating receptors of the seven-transmembrane, G-protein-linked family, in particular by binding to IL-8 receptors, most notably the B-receptor. Thomas et al., J. Biol. Chem. **266**, 14839 (1991); and Holmes et al., Science **253**, 1278 (1991). The development of non-peptide small molecule antagonists for members of this receptor family has precedent. For a review see R. Freidinger in: Progress in Drug Research, Vol. 40, pp. 33-98, Birkhauser Verlag, Basel 1993. Hence, the IL-8 receptor represents a promising target for the development of novel anti-inflammatory agents.

Two high affinity human IL-8 receptors (77% homology) have been characterized: IL-8R α , which binds only IL-8 with high affinity, and IL-8R β , which has high affinity for IL-8 as well as for GRO- α , GRO β , GRO γ and NAP-2. See Holmes et al., supra; Murphy et al., Science **253**, 1280 (1991); Lee et al., J. Biol. Chem. **267**, 16283 (1992); LaRosa et al., J. Biol. Chem. **267**, 25402 (1992); and Gayle et al., J. Biol. Chem. **268**, 7283 (1993).

There remains a need for treatment, in this field, for compounds which are capable of binding to the IL-8 α or β receptor. Therefore, conditions associated with

an increase in IL-8 production (which is responsible for chemotaxis of neutrophil and T-cells subsets into the inflammatory site) would benefit by compounds which are inhibitors of IL-8 receptor binding.


SUMMARY OF THE INVENTION

5 This invention provides for a method of treating a chemokine mediated disease, wherein the chemokine is one which binds to an IL-8 α or β receptor and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In particular the chemokine is IL-8.

10 This invention also relates to a method of inhibiting the binding of IL-8 to its receptors in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

Compounds of Formula (I) useful in the present invention are represented by the structure:

15



wherein

Z is cyano, OR₁₁, C(O)NR₁₅R₁₆, R₁₈, C(O)OR₁₁, C(O)R₁₁, or S(O)₂R₁₇;

R is any functional moiety having an ionizable hydrogen and a pK_a of 10 or less;

20 R₁ is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C₁₋₁₀ alkyl; C₁₋₁₀ alkyl; C₂₋₁₀ alkenyl; C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀ alkoxy; azide; (CR₈R₈)_q S(O)_tR₄; hydroxy; hydroxy C₁₋₄alkyl; aryl; aryl C₁₋₄ alkyl; aryloxy; aryl C₁₋₄ alkyloxy; heteroaryl; heteroarylalkyl; heterocyclic; heterocyclic C₁₋₄alkyl; heteroaryl C₁₋₄ alkyloxy; aryl C₂₋₁₀ alkenyl; heteroaryl
25 C₂₋₁₀ alkenyl; heterocyclic C₂₋₁₀ alkenyl; (CR₈R₈)_qNR₄R₅; C₂₋₁₀ alkenyl C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₁₀; S(O)₃H; S(O)₃R₈; (CR₈R₈)_q C(O)R₁₁; C₂₋₁₀ alkenyl C(O)R₁₁; C₂₋₁₀ alkenyl

- $C(O)OR_{11}$; $(CR_8R_8)_q C(O)OR_{12}$; $(CR_8R_8)_q OC(O)R_{11}$;
 $(CR_8R_8)_q NR_4C(O)R_{11}$; $(CR_8R_8)_q NHS(O)_2R_{19}$; $(CR_8R_8)_q S(O)_2NR_4R_5$; or
 two R_1 moieties together may form $O-(CH_2)_sO-$ or a 5 to 6 membered saturated
 or unsaturated ring, and wherein the aryl, heteroaryl and heterocyclic containing
 5 moieties may be optionally substituted;
 q is 0, or an integer having a value of 1 to 10;
 t is 0, or an integer having a value of 1 or 2;
 m is an integer having a value of 1 to 3;
 s is an integer having a value of 1 to 3;
 10 R_4 and R_5 are independently hydrogen, optionally substituted C_{1-4} alkyl, optionally
 substituted aryl, optionally substituted aryl C_{1-4} alkyl, optionally substituted
 heteroaryl, optionally substituted heteroaryl C_{1-4} alkyl, heterocyclic, or
 heterocyclic C_{1-4} alkyl, or R_4 and R_5 together with the nitrogen to which they
 are attached form a 5 to 7 member ring which may optionally comprise an
 15 additional heteroatom selected from oxygen, nitrogen or sulfur;
 HET is an optionally substituted heteroaryl moiety;
 R_6 and R_7 are independently hydrogen or a C_{1-4} alkyl group; or R_6 and R_7 together
 with the nitrogen to which they are attached form a 5 to 7 member ring which
 ring may optionally contain an additional heteroatom selected from oxygen,
 20 nitrogen or sulfur;
 R_8 is independently selected from hydrogen or C_{1-4} alkyl;
 R_{10} is C_{1-10} alkyl $C(O)_2R_8$
 R_{11} is hydrogen, C_{1-4} alkyl, optionally substituted aryl, optionally substituted aryl
 C_{1-4} alkyl, optionally substituted heteroaryl, optionally substituted
 25 heteroaryl C_{1-4} alkyl, optionally substituted heterocyclic, or optionally
 substituted heterocyclic C_{1-4} alkyl;
 R_{12} is hydrogen, C_{1-10} alkyl, optionally substituted aryl or optionally substituted
 arylalkyl;
 R_{13} and R_{14} are independently hydrogen or C_{1-4} alkyl;
 30 v is 0, or an integer having a value of 1 to 4;


R₁₅ and R₁₆ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, optionally substituted heterocyclicC₁₋₄alkyl, or R₁₅ and R₁₆ may together with the nitrogen to which they are attached form a 5 to 7 member ring optionally containing an additional heteroatom selected from oxygen, nitrogen, or sulfur;

R₁₇ is C₁₋₄ alkyl, NR₁₅R₁₆, OR₁₁, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;


R₁₈ is optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;

R₁₉ is C₁₋₄alkyl, aryl, arylalkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclic, or heterocyclicC₁₋₄alkyl, wherein the all of these moieties may be optionally substituted;

R_d is NR₆R₇, alkyl, arylC₁₋₄ alkyl, arylC₂₋₄ alkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroarylC₂₋₄ alkenyl, heterocyclic, heterocyclicC₁₋₄ alkyl, wherein the alkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic, and heterocyclic alkyl rings may be optionally substituted;



the E' containing ring is optionally selected from




a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

5 The compounds of Formula (I) may also be used in association with the veterinary treatment of mammals, other than humans, in need of inhibition of IL-8 or other chemokines which bind to the IL-8 α and β receptors. Chemokine mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section.

10 In compounds of Formula (I), W is suitably



The E' containing ring, denoted by its point of attachment through the asterisk (*) may optionally be present. If it is not present the ring is a phenyl moiety which is substituted by the R_1 terms as shown. The E' ring may be substituted by the R_1 moiety in any ring, saturated or unsaturated, and is shown for purposes herein substituted only in the unsaturated ring(s).

R is suitably any functional moiety which provides an ionizable hydrogen having a pK_a of 10 or less, preferably from about 3 to 9, more preferably from about 3 to 7. Such functional groups include, but are not limited to, hydroxy, carboxylic acid, thiol, SR_2 , OR_2 , $NH-C(O)R_a$, $C(O)NR_6R_7$, a substituted sulfonamides of the

formula: $\text{NHS(O)}_2\text{R}_b$, $\text{S(O)}_2\text{NHR}_c$, $\text{NHC(X}_2\text{)NHR}_b$, or a tetrazolyl; wherein X_2 is oxygen or sulfur, preferably oxygen. Preferably, the functional group is other than a sulfonic acid, either directly or as a substituent group on the aryl, heteroaryl, or heterocyclic moiety ring, such as in SR_2 or OR_2 . More preferably R is OH, SH, or

5 $\text{NHS(O)}_2\text{R}_b$.

Suitably, R_2 is a substituted aryl, heteroaryl, or heterocyclic moiety which ring has the functional moiety providing the ionizable hydrogen having a pK_a of 10 or less.

Suitably, R_6' and R_7' are hydrogen, C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, heterocyclic C_{2-4} alkenyl moiety, all of which may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C_{1-4} alkyl, such as CF_3 ; C_{1-4} alkyl, such as methyl; C_{1-4} alkoxy, such as methoxy; $\text{NR}_9\text{C(O)R}_a$; $\text{C(O)NR}_6\text{R}_7$; $\text{S(O)}_3\text{H}$; or C(O)OC_{1-4} alkyl,

15 provided that only one of R_6' and R_7' are hydrogen, but not both.

Suitably, R_6 and R_7 are independently hydrogen or a C_{1-4} alkyl group, or R_6 and R_7 together with the nitrogen to which they are attached form a 5 to 7 member ring which ring may optionally contain an additional heteroatom which heteroatom is selected from oxygen, nitrogen or sulfur. This heteroring may be

20 optionally substituted as defined herein.

Suitably R_a is an aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, or a heterocyclic C_{1-4} alkyl moiety, all of which may be optionally substituted, as defined herein below.

Suitably, R_b is a NR_6R_7 , alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, heterocyclic C_{2-4} alkenyl moiety, or camphor, all of which may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C_{1-4} alkyl, such as CF_3 ; C_{1-4} alkyl, such as methyl; C_{1-4} alkoxy, such as methoxy; $\text{NR}_9\text{C(O)R}_a$; $\text{C(O)NR}_6\text{R}_7$; $\text{S(O)}_3\text{H}$; or C(O)OC_{1-4} alkyl. R_b is

25

preferably an optionally substituted phenyl, benzyl, or styryl. When R_b is a heteroaryl preferably it is an optionally substituted thiazole, optionally substituted thienyl, or optionally substituted quinolinyl ring.

Suitably R_9 is hydrogen or a C_{1-4} alkyl, preferably hydrogen. Preferably, when the substituent group is $NR_9C(O)R_a$, then R_a is preferably an alkyl group, such as methyl.

Suitably R_c is hydrogen, alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{1-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{1-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, or heterocyclic C_{1-4} alkenyl moiety, all of which may be optionally substituted one to three times independently by halogen, nitro, halosubstituted C_{1-4} alkyl, C_{1-4} alkyl, C_{1-4} alkoxy, $NR_9C(O)R_a$, $C(O)NR_6R_7$, $S(O)_3H$, or $C(O)OC_{1-4}$ alkyl. Preferably, R_c is an optionally substituted phenyl.

When R is an OR_2 or SR_2 moiety it is recognized by one of skill in the art that the aryl ring must, therefore, contain the required ionizable hydrogen. The aryl ring may also be additionally substituted, independently, by one to three groups, which groups may also contain an additional ionizable group, and which include but are not limited to, halogen, nitro, halosubstituted C_{1-4} alkyl, C_{1-4} alkyl, C_{1-4} alkoxy, hydroxy, SH , $C(O)NR_6R_7$, $NH-C(O)R_a$, $NHS(O)_2R_b$, $S(O)_2NR_6R_7$, $C(O)OR_8$, or a tetrazolyl ring.

In compounds of Formula (I), suitably R_1 is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C_{1-10} alkyl, such as CF_3 ; C_{1-10} alkyl, such as methyl, ethyl, isopropyl, or *n*-propyl; C_{2-10} alkenyl; C_{1-10} alkoxy, such as methoxy, or ethoxy; halosubstituted C_{1-10} alkoxy, such as trifluoromethoxy; azide; $(CR_8R_8)_q S(O)_tR_4$, wherein t is 0, 1 or 2; hydroxy; hydroxy C_{1-4} alkyl, such as methanol or ethanol; aryl, such as phenyl or naphthyl; aryl C_{1-4} alkyl, such as benzyl; aryloxy, such as phenoxy; aryl C_{1-4} alkyloxy, such as benzyloxy; heteroaryl; heteroarylalkyl; heteroaryl C_{1-4} alkyloxy; aryl C_{2-10} alkenyl; heteroaryl C_{2-10} alkenyl; heterocyclic C_{2-10} alkenyl; $(CR_8R_8)_q NR_4R_5$; C_{2-10} alkenyl $C(O)NR_4R_5$; $(CR_8R_8)_q C(O)NR_4R_5$; $(CR_8R_8)_q C(O)NR_4R_{10}$; $S(O)_3H$; $S(O)_3R_8$; $(CR_8R_8)_q$

C(O)R₁₁; C₂₋₁₀ alkenyl C(O)R₁₁; C₂₋₁₀ alkenyl C(O)OR₁₁; C(O)R₁₁; (CR₈R₈)_q
 C(O)OR₁₂; (CR₈R₈)_qOC(O)R₁₁; (CR₈R₈)_qNR₄C(O)R₁₁; (CR₈R₈)_q
 NHS(O)₂R₁₉; or (CR₈R₈)_qS(O)₂NR₄R₅; or two R₁ moieties together may form
 O-(CH₂)_sO- or a 5 to 6 membered saturated or unsaturated ring; and s is an integer
 5 having a value of 1 to 3. The aryl, heteroaryl, and heterocyclic containing moieties
 may be optionally substituted as defined herein below.

Suitably, q is 0, or an integer having a value of 1 to 10.

When R₁ forms a dioxybridge, s is preferably 1. When R₁ forms an
 additional saturated or unsaturated ring, it is preferably 6 membered ring resulting in
 10 a naphthylene ring system. These rings may be substituted independently, 1 to 3
 times by the other R₁ moieties as defined above.

Suitably, R₄ and R₅ are independently hydrogen, optionally substituted C₁₋₄
 alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally
 substituted heteroaryl, optionally substituted heteroaryl C₁₋₄alkyl, heterocyclic,
 15 heterocyclicC₁₋₄ alkyl, or R₄ and R₅ together with the nitrogen to which they are
 attached form a 5 to 7 member ring which may optionally comprise an additional
 heteroatom selected from O/N/S.

R₈ is suitably independently selected from hydrogen or C₁₋₄ alkyl.

R₁₀ is suitably C₁₋₁₀ alkyl C(O)₂R₈, such as CH₂C(O)₂H or
 20 CH₂C(O)₂CH₃.

R₁₁ is suitably hydrogen, C₁₋₄ alkyl, aryl, aryl C₁₋₄ alkyl, heteroaryl,
 heteroaryl C₁₋₄alkyl, heterocyclic, or heterocyclic C₁₋₄alkyl.

R₁₂ is suitably hydrogen, C₁₋₁₀ alkyl, optionally substituted aryl or
 optionally substituted arylalkyl.

25 R₁₉ is C₁₋₄alkyl, aryl, arylalkyl, heteroaryl, heteroarylC₁₋₄alkyl,
 heterocyclic, or heterocyclicC₁₋₄alkyl, wherein the all of these moieties may be
 optionally substituted;

Preferably R₁ is halogen, cyano, nitro, CF₃, C(O)NR₄R₅, alkenyl
 C(O)NR₄R₅, C(O)R₄R₁₀, alkenyl C(O)OR₁₂, heteroaryl, heteroarylalkyl,

heteroaryl alkenyl, or $S(O)NR_4R_5$, and preferably R_4 and R_5 are both hydrogen or one is phenyl. A preferred ring substitution for R_1 is in the 4-position of the phenyl ring.

When R is OH , SH or NSO_2R_b than R_1 is preferably mono-substituted in the 3-position, the 4- position or is di substituted in the 3,4- position. The substituent group is suitably an electron withdrawing moiety. Preferably when R is OH , SH or NSO_2R_b , than R_1 is nitro, halogen, cyano, trifluoromethyl group, $C(O)NR_4R_5$.

When R is carboxylic acid, than R_1 is preferably hydrogen, or R_1 is preferably substituted in the 4-position, more preferably substituted by trifluoromethyl or chloro.

In compounds of Formula (I), suitably R_{13} and R_{14} are independently hydrogen, optionally substituted C_{1-4} alkyl which may be straight or branched as defined herein, or one of R_{13} and R_{14} are an optionally substituted aryl; v is 0, or an integer having a value of 1 to 4.

When R_{13} or R_{14} are an optionally substituted alkyl, the alkyl moiety may be substituted one to three times independently by halogen; halosubstituted C_{1-4} alkyl such as trifluoromethyl; hydroxy; hydroxy C_{1-4} alkyl, C_{1-4} alkoxy; such as methoxy, or ethoxy, halosubstituted C_{1-10} alkoxy, $S(O)_tR_4$; aryl; NR_4R_5 ; $NHC(O)R_4$; $C(O)NR_4R_5$; or $C(O)OR_8$.

In compounds of Formula (I), Z is suitably cyano, OR_{11} , $C(O)NR_{15}R_{16}$, R_{18} , $C(O)R_{11}$, $C(O)OR_{11}$, or $S(O)_2R_{17}$.

Suitably R_{15} and R_{16} are independently hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-4} alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C_{1-4} alkyl, optionally substituted heterocyclic, optionally substituted heterocyclic C_{1-4} alkyl, or R_{15} and R_{16} may together with the nitrogen to which they are attached form a 5 to 7 member ring optionally containing an additional heteroatom selected from oxygen, nitrogen, or sulfur.

Suitably, R₁₇ is C₁₋₄ alkyl, NR₁₅R₁₆, OR₁₁, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl.

5 In compounds of Formula (I), R₁₈ is suitably hydrogen, halogen, cyano, optionally substituted C₁₋₄ alkyl, halo substituted C₁₋₄ alkyl, C(O)NR₁₅R₁₆, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl.

10 In compounds of Formula (I), HET is suitably a heteroaryl ring or multi-ring system. If the HET moiety is a multi ring system, the ring containing the heteroatom does not need to be directly attached to the urea moiety. All the rings in this ring system may be optionally substituted as defined herein. Preferably the HET moiety is a pyridyl, which may be 2-, 3- or 4-pyridyl. If the ring is a multi
15 system ring it is preferably benzimidazole, dibenzothiophene, or an indole ring. Other heterocyclic rings of interest include, but are not limited to thiophene, furan, pyrimidine, pyrrole, pyrazole, quinoline, isoquinoline, quinazolinyl, pyridine, oxazole, thiazole, thiadiazole, triazole, imidazole, or benzimidazole.

In compounds of Formula (I), the HET ring may be optionally substituted
20 independently one to three times by Y, i.e. (Y_(n)), wherein n is an integer having a value of 1 to 3. Suitably, Y is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C₁₋₁₀ alkyl; C₁₋₁₀ alkyl; C₂₋₁₀ alkenyl; C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀ alkoxy; azide; (CR₈R₈)_q S(O)_tR₄; hydroxy; hydroxyC₁₋₄alkyl; aryl; aryl C₁₋₄ alkyl; aryloxy; arylC₁₋₄ alkyloxy; heteroaryl;
25 heteroarylalkyl; heteroaryl C₁₋₄ alkyloxy; heterocyclic, heterocyclic C₁₋₄alkyl; aryl C₂₋₁₀ alkenyl; heteroaryl C₂₋₁₀ alkenyl; heterocyclic C₂₋₁₀ alkenyl; (CR₈R₈)_q NR₄R₅; C₂₋₁₀ alkenyl C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₁₀; S(O)₃H; S(O)₃R₈; (CR₈R₈)_q C(O)R₁₁; C₂₋₁₀ alkenyl C(O)R₁₁; C₂₋₁₀ alkenyl C(O)OR₁₁; (CR₈R₈)_q C(O)OR₁₂; (CR₈R₈)_q OC(O) R₁₁; (CR₈R₈)_q

NR₄C(O)R₁₁; (CR₈R₈)_q NHS(O)₂R_d; (CR₈R₈)_q S(O)₂NR₄R₅; or two Y moieties together may form O-(CH₂)_sO- or a 5 to 6 membered saturated or unsaturated ring; wherein the aryl, heteroaryl, heterocyclic containing moieties may be optionally substituted as defined herein.

- 5 When Y forms a dioxybridge, s is preferably 1. When Y forms an additional saturated or unsaturated ring, it is preferably 6 membered ring resulting in a naphthylene ring system. These rings may also be substituted 1 to 3 times by other Y moieties as defined above.

- Suitably, R_d is a NR₆R₇, alkyl, aryl C₁₋₄ alkyl, aryl C₂₋₄ alkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl C₂₋₄ alkenyl, heterocyclic, heterocyclic C₁₋₄ alkyl, or heterocyclic C₂₋₄ alkenyl moiety, wherein the aryl, heteroaryl, and heterocyclic containing moieties may all be optionally substituted as defined herein.

- As used herein, "optionally substituted" unless specifically defined shall mean such groups as halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; hydroxy substituted C₁₋₁₀alkyl; C₁₋₁₀ alkoxy, such as methoxy or ethoxy; S(O)_m' C₁₋₁₀ alkyl, wherein m' is 0, 1 or 2, such as methyl thio, methyl sulfinyl or methyl sulfonyl; amino, mono & di-substituted amino, such as in the NR₄R₅ group; NHC(O)R₄; C(O)NR₄R₅; C(O)OH; S(O)₂NR₄R₅; NHS(O)₂R₂₁, C₁₋₁₀ alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; halosubstituted C₁₋₁₀ alkyl, such as CF₃; an optionally substituted aryl, such as phenyl, or an optionally substituted arylalkyl, such as benzyl or phenethyl, optionally substituted heterocyclic, optionally substituted heterocyclicalkyl, optionally substituted heteroaryl, optionally substituted heteroaryl alkyl, wherein these aryl, heteroaryl, or heterocyclic moieties may be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl; C₁₋₁₀ alkoxy; S(O)_m C₁₋₁₀ alkyl; amino, mono & di-substituted amino, such as in the NR₄R₅ group; C₁₋₁₀ alkyl, or halosubstituted C₁₋₁₀ alkyl, such as CF₃.

R₂₁ is suitably C₁₋₄ alkyl, aryl, aryl C₁₋₄alkyl, heteroaryl, heteroaryl C₁₋₄alkyl, heterocyclic, or heterocyclic C₁₋₄alkyl.

Suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of inorganic and organic acids, such as hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid, ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid, 5 oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid. In addition, pharmaceutically acceptable salts of compounds of Formula (I) may also be formed with a pharmaceutically acceptable cation, for instance, if a substituent group comprises a carboxy moiety. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium 10 cations.

The following terms, as used herein, refer to:

- "halo" - all halogens, that is chloro, fluoro, bromo and iodo.
- "C₁₋₁₀alkyl" or "alkyl" - both straight and branched chain radicals of 1 to 15 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.
- The term "cycloalkyl" is used herein to mean cyclic radicals, preferably of 3 to 8 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, 20 and the like.
- The term "alkenyl" is used herein at all occurrences to mean straight or branched chain radical of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.
- 25 • "aryl" - phenyl and naphthyl;
- "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") - a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, quinoline, 30 isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, thiazole, thiadiazole, triazole, imidazole, or benzimidazole.

- "heterocyclic" (on its own or in any combination, such as "heterocyclicalkyl") - a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydropyran, or imidazolidine.
- The term "arylalkyl" or "heteroarylalkyl" or "heterocyclicalkyl" is used herein to mean C₁₋₁₀ alkyl, as defined above, attached to an aryl, heteroaryl or heterocyclic moiety, as also defined herein, unless otherwise indicated.
- "sulfinyl" - the oxide S (O) of the corresponding sulfide, the term "thio" refers to the sulfide, and the term "sulfonyl" refers to the fully oxidized S(O)₂ moiety.
- The term "wherein two R₁ moieties (or two Y moieties) may together form a 5 or 6 membered saturated or unsaturated ring" is used herein to mean the formation of a naphthylene ring system or a phenyl moiety having attached a 6 membered partially unsaturated ring such as a C₆ cycloalkenyl, i.e hexene, or a C₅ cyloalkenyl moiety, cyclopentene.

It is recognized that the compounds of the present invention may exist as stereoisomers, regioisomers, or diastereoisomers. These compounds may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. All of these compounds are included within the scope of the present invention.


The compounds of Formula (I) may be obtained by applying synthetic procedures, some of which are illustrated in the Schemes below. The synthesis provided for in these Schemes is applicable for the producing compounds of Formula (I) having a variety of different R, R₁, and Aryl groups which are reacted, employing optional substituents which are suitably protected, to achieve compatibility with the reactions outlined herein. Subsequent deprotection, in those cases, then affords compounds of the nature generally disclosed. Once the urea nucleus has been established, further compounds of these formulas may be prepared by applying standard techniques for functional group interconversion, well known in

the art. While the schemes are shown with compounds only of Formula (I) this is merely for illustration purposes only.

Methods of Preparation

The title compounds can be synthesized from the thiouronium salt (2, Scheme 1). R' represents the (R₁₃R₁₄)_v- HET linkage as defined in compounds of Formula (I). Z is also as defined in compounds of Formula (I). For purposes of illustration herein the schemes represent the W term by a substituted phenyl.

Scheme 1




a)Na, EtOH, cyanamide b)EDC·HCl

The thiouronium salt (2, Scheme 1) can be synthesized by reacting sodium cyanamide with a commercially available isothiocyanate 1 (if the isothiocyanate is not commercially available it can be synthesized by reacting the desired amine with thiophosgene in the presence of a base like sodium bicarbonate). The thiouronium salt (2) can then be condensed with the appropriate substituted aniline in the presence of a coupling reagent like EDC·HCl or acidified to form the cyanothiurea and then reacted.

Alternatively 4 can be synthesized by reacting the commercially available (Aldrich Chem. Co.), diphenyl cyanocarboimide (5, Scheme 2) with an amine to

form an intermediate O-phenylisourea 5 which then reacts with the appropriately substituted aniline in the presence of trimethyl aluminum by the method of Atwal. (Atwal, K. S., Tetrahedron Lett, 35, 8085 (1994).). Compounds wherein R' is alkyl may be prepared by heating with the appropriately substituted alkylamine, but without the presence of catalyst.

Scheme 2



a) acetonitrile, heat b) $AlMe_3$


Alternatively the title compound can be synthesized using a protected ortho-substituted aniline (8, Scheme 3, See synthesis as described in US provisional application USSN 60/020655 filed June 27, 1996, Attorney Docket No.: P50467P; WO96/25157 filed 22 August 1996, Widdowson et al. (Attorney Docket No.: P50324-1); and USSN 08/701,299 filed 21 August 1996 (Attorney Docket No.: P50324-2) whose disclosures are incorporated herein by reference in their entirety.

An ortho-substituted aniline (7, Scheme 3) is first protected (ie. tert-butyl dimethyl silyl, allyl, benzyl, mom or other suitable protecting group) by reacting the ortho-

substituted aniline with the appropriate alkyl or silyl halide in the presence of a suitable base (ie. cesium carbonate, potassium carbonate or imidazole) in an aprotic solvent. The protected ortho-substituted aniline may also be synthesized from an ortho-substituted nitrobenzene (9) by reacting it with a protecting group under conditions well known in the art (see Greene, T Protecting Groups in Organic Synthesis, Wiley&Sons, New York, 1981) This protected ortho-substituted nitro compound is then reduced to the corresponding aniline using SnCl_2 in EtOH or alternately H_2/Pd or LiAlH_4 in an aprotic solvent.

10

Scheme 3




a) Alkyl or silyl halide, base b) reducing agent

15

This protected ortho-substituted aniline (8) can then be converted into an isothiocyanate using thiophosgene and then reacted with the anion ZNH^- (formed from reaction of ZNH_2 with a base such as NaH). Z is as defined in compounds of Formula (I). The resulting thioanion can then be alkylated with an alkylating agent like methyl iodide to form a thioimide such as 10 (Scheme 4).

20

Scheme 4




a) ClCSCl, NaHCO₃ b) ZNH⁻ c) MeI

5

The thioimide (10, Scheme 5) can be converted to the title compound 4 by reaction with the amine R'NH₂. This reaction can be accelerated by the addition of metal salt with a high affinity for sulfur such as mercuric oxide or silver acetate or by oxidation of the sulfur with dimethyloxirane to form a better leaving group. Finally the phenol protection is removed by standard methods to form the title compound 4.

10

Scheme 5



15

a) R'NH₂ b) deprotect

Alternately the title compound could be synthesized by reaction of a protected carbodiimide (11, Scheme 6) with the anion NH-Z (formed from reaction of NH₂Z with a base such as NaH) or the neutral species NH₂Z (Z=CN) and a tertiary amine base, such as Hünig's base (diisopropylethylamine), triethylamine, triisopropylethylamine, N,N-dimethylbenzylamine, or N,N-dimethylisopropylamine, under conditions where the nucleophile is present in large excess and the reaction


20

time is kept as short as possible by carefully monitoring the reaction for completion followed by deprotection. Other suitable bases for use herein include, secondary amine, such as pyridine, and amino substituted pyridine derivatives. Suitable solvents for use herein when Z is cyano include various aprotic solvents, such as acetonitrile; halogenated solvents, such as chloroform and methylene chloride; ethyl glycol-dimethyl ether (monoGLYME), dioxane, DMF and DMSO; or mixtures thereof, preferably acetonitrile. It is recognized by the skilled artisan that the limiting feature for use of solvents herein will be the solubility of the cyano derivatized compound. For compounds wherein Z is other than cyano, while aprotic solvents are preferred, it is recognized by the skilled artisan that other suitable solvents, such as protic solvents, i.e. alcohols, may be used.

Preferably the reaction (when Z is cyano) temperature is from about -10° to about 100°, preferably about 10° to about 50°, more preferably around room temperature, i.e. 20 to 30 °C.

The protected R'' moiety may be suitably deprotected using art recognized techniques. Preferably the deprotection is by deallylation catalyzed by palladium (O) when the protecting group is an allyl derivative.

Scheme 6



11


a) ZNH^- ($\text{ZNH}_2 + \text{NaH}$) for $\text{Z}=\text{OR}, \text{COOEt}, \text{CHO}, \text{RNHSO}_2, \text{ArNHSO}_2$;

b) $\text{ZNH}_2 \cdot \text{HCl}$ for $\text{Z}=\text{OH}$ c) ZNH_2 and NR_3 for $\text{Z}=\text{CN}$ d) deprotect

The carbodiimide 11 is prepared from the thiourea (12a, Scheme 7) by treatment with phosgene and a tertiary amine base or from the thiourea (12a) or urea (12b) by reaction with triphenylphosphine, carbon tetrachloride and triethylamine.

The carbodiimide may also be prepared by reaction of the thiourea (12b) with an excess, such as 2 or more equivalents of methanesulfonyl chloride and a tertiary amine base, such as Hünig's base (diisopropylethylamine), triethylamine, triisopropylethylamine, N,N-dimethylbenzylamine, or N,N-dimethylisopropylamine, preferably triethylamine. The reaction may use any halogenated solvent, such as methylene chloride, chloroform, or tetrachloroethylene, etc.; suitable reaction temperatures are from about -30°C to about 80°C, preferably -10°C to about 50°C., more preferably from about 0°C to about room temperature. See Fell and Coppola (Fell, J. B., Coppola, J. B., Syn Communications 25, 43, (1995).

Scheme 7




a) phosgene, Et₃N, b) Ph₃P, CCl₄, Et₃N; c) MsCl, Et₃N

The thiourea or urea derivative may be synthesized as described in the US provisional application USSN 60/020655 and WO96/25157, *supra*. The thiourea (12a, Scheme 8) may also be prepared by reaction of the protected ortho-substituted aniline and two equivalents of an appropriate base such as NaH, KH, calcium hydride, and reacting this anion with a commercially available isothiocyanate (W₁-NCS, wherein W₁ is as defined for compounds of Formula (I)). The reaction may take place in any suitable aprotic solvent or halogenated solvent, preferably dimethyl formamide. Suitable reaction temperatures for this reaction are from about -10° to about 50°.

If the desired isothiocyanate is not commercially available, it may be prepared by reaction of a corresponding aniline with thiophosgene and a suitable base such as sodium bicarbonate.


Scheme 8



5

a) base b) $\text{R}'\text{NCS}$

Another aspect of the present invention are the novel compounds of Formula (II)



- 10 wherein W, and HET are as defined for formula (I), and W contains a protected or unprotected R group (R'') as defined for compounds of Formula (I).

Another aspect of the present invention are the novel compounds of Formula (III)



- 15 wherein W, and HET are as defined for formula (I), and W contains a protected or unprotected R group (R'') as defined for compounds of Formula (I).

Also, the protected (R'') versions of compounds of Formula (I) are contemplated as being within the scope of this invention.

- 20 It is recognized that the guanidine functionality may have a number of different tautomers, such as $\text{W}-\text{N}-\text{C}(=\text{NZ})-\text{N}-\text{HET}$; $\text{ZN}=\text{C}(\text{N}-\text{HET})-\text{N}-\text{W}$; $\text{HET}-\text{N}-\text{C}(=\text{NW})-\text{NZ}$, all of which are within the scope of this invention.

Pharmaceutically acceptable salts of compounds of Formula (I) may be obtained in known manner, for example by treatment thereof with an appropriate amount of acid or base in the presence of a suitable solvent.

METHOD OF TREATMENT

The compounds of Formula (I), or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or
5 caused by excessive or unregulated IL-8 cytokine production by such mammal's cell, such as but not limited to monocytes and/or macrophages, or other chemokines which bind to the IL-8 α or β receptor, also referred to as the type I or type II receptor.

Accordingly, the present invention provides a method of treating a
10 chemokine mediated disease, wherein the chemokine is one which binds to an IL-8 α or β receptor and which method comprises administering an effective amount of a compound of Formula (I), or a pharmaceutically acceptable salt thereof. In particular, the chemokines are IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2.

The compounds of Formula (I) are administered in an amount sufficient to
15 inhibit cytokine function, in particular IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2, such that they are biologically regulated down to normal levels of physiological function, or in some case to subnormal levels, so as to ameliorate the disease state. Abnormal levels of IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 for instance in the context of the present invention, constitute: (i) levels of free IL-8
20 greater than or equal to 1 picogram per mL; (ii) any cell associated IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 above normal physiological levels; or (iii) the presence of IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 above basal levels in cells or tissues in which IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 respectively, is produced.

25 There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. Chemokine mediated diseases include psoriasis, atopic dermatitis, arthritis, asthma, chronic obstructive pulmonary disease, adult respiratory distress syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, stroke, septic shock, endotoxic
30 shock, gram negative sepsis, toxic shock syndrome, cardiac and renal reperfusion injury, glomerulonephritis, thrombosis, graft vs. host reaction, alzheimers disease,

allograft rejections, malaria, restinosis, angiogenesis, atherosclerosis, osteoporosis, gingivitis, undesired hematopoietic stem cells release and diseases caused by respiratory viruses, including but not limited to rhinovirus and influenza virus, herpesviruses, including but not limited to herpes simplex I and II, and hepatitis
5 viruses, including but not limited to Hepatitis B and Hepatitis C virus.

These diseases are primarily characterized by massive neutrophil infiltration, T-cell infiltration, or neovascular growth, and are associated with increased IL-8, GRO α , GRO β , GRO γ or NAP-2 production which is responsible for the chemotaxis of neutrophils into the inflammatory site or the directional growth of endothelial
10 cells. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8, GRO α , GRO β , GRO γ or NAP-2 has the unique property of promoting neutrophil chemotaxis, enzyme release including but not limited to elastase release as well as superoxide production and activation. The α -chemokines but particularly, GRO α , GRO β , GRO γ or NAP-2, working through the IL-8 type I or II receptor can promote
15 the neovascularization of tumors by promoting the directional growth of endothelial cells. Therefore, the inhibition of IL-8 induced chemotaxis or activation would lead to a direct reduction in the neutrophil infiltration.

Recent evidence also implicates the role of chemokines in the treatment of HIV infections, Littleman et al., Nature 381, pp661 (1996) and Koup et al., Nature
20 381, pp 667 (1996).

The present invention also provides for a means of treating, in an acute setting, as well as preventing, in those individuals deemed susceptible to, CNS injuries by the chemokine receptor antagonist compounds of Formula (I).

CNS injuries as defined herein include both open or penetrating head trauma, such
25 as by surgery, or a closed head trauma injury, such as by an injury to the head region. Also included within this definition is ischemic stroke, particularly to the brain area.

Ischemic stroke may be defined as a focal neurologic disorder that results from insufficient blood supply to a particular brain area, usually as a consequence of an embolus, thrombi, or local atheromatous closure of the blood vessel. The role of
30 inflammatory cytokines in this are has been emerging and the present invention provides a

mean for the potential treatment of these injuries. Relatively little treatment, for an acute injury such as these has been available.

TNF- α is a cytokine with proinflammatory actions, including endothelial leukocyte adhesion molecule expression. Leukocytes infiltrate into ischemic brain lesions and hence
5 compounds which inhibit or decrease levels of TNF would be useful for treatment of ischemic brain injury. See Liu et al., Stoke, Vol. 25., No. 7, pp 1481-88 (1994) whose disclosure is incorporated herein by reference.

Models of closed head injuries and treatment with mixed 5-LO/CO agents is discussed in Shohami et al., J. of Vasc & Clinical Physiology and Pharmacology,
10 Vol. 3, No. 2, pp. 99-107 (1992) whose disclosure is incorporated herein by reference. Treatment which reduced edema formation was found to improve functional outcome in those animals treated.

The compounds of Formula (I) are administered in an amount sufficient to inhibit IL-8, binding to the IL-8 alpha or beta receptors, from binding to these
15 receptors, such as evidenced by a reduction in neutrophil chemotaxis and activation. The discovery that the compounds of Formula (I) are inhibitors of IL-8 binding is based upon the effects of the compounds of Formulas (I) in the *in vitro* receptor binding assays which are described herein. The compounds of Formula (I) have been shown, in some instances, to be dual inhibitors of both recombinant type I and
20 type II IL-8 receptors. Preferably the compounds are inhibitors of only one receptor, more preferably Type II.

As used herein, the term "IL-8 mediated disease or disease state" refers to any and all disease states in which IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 plays a role, either by production of IL-8, GRO α , GRO β , GRO γ or NAP-2
25 themselves, or by IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease stated mediated by IL-8.

30 As used herein, the term "chemokine mediated disease or disease state" refers to any and all disease states in which a chemokine which binds to an IL-8 α or β

receptor plays a role, such as but not limited to IL-8, GRO α , GRO β , GRO γ , ENA-78 or NAP-2. This would include a disease state in which, IL-8 plays a role, either by production of IL-8 itself, or by IL-8 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease state in which, for instance, IL-1
5 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease stated mediated by IL-8.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes,
10 but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts, basophils, neutrophils, endothelial cells, brain astrocytes, bone marrow stromal cells,
15 epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- α) and Tumor Necrosis Factor beta (TNF- β).

As used herein, the term "chemokine" refers to any secreted polypeptide that
20 affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response, similar to the term "cytokine" above. A chemokine is primarily secreted through cell transmembranes and causes chemotaxis and activation of specific white blood cells and leukocytes, neutrophils, monocytes, macrophages, T-cells, B-cells, endothelial cells and smooth
25 muscle cells. Examples of chemokines include, but are not limited to, IL-8, GRO- α , GRO- β , GRO γ , ENA-78, NAP-2, IP-10, MIP-1 α , MIP- β , PF4, and MCP 1, 2, and 3.

In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. This invention,
30 therefore, also relates to a pharmaceutical composition comprising an effective, non-

toxic amount of a compound of Formula (I) and a pharmaceutically acceptable carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally, topically, parenterally or by inhalation. The compounds of Formula (I) may be administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg. to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I) externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the Formulation. It may however comprise as much as 10% w/w but preferably will comprise less than 5% w/w, more preferably from 0.1% to 1% w/w of the Formulation.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or a macrogel. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such

as a sorbitan ester or a polyoxyethylene derivative thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

Drops according to the present invention may comprise sterile aqueous or
5 oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining
10 at 98-100 °C. for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily
15 solution include glycerol, diluted alcohol and propylene glycol.

Compounds of formula (I) may be administered parenterally, that is by intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. Appropriate dosage forms for such
20 administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation, that is by intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques.

25 For all methods of use disclosed herein for the compounds of Formula (I), the daily oral dosage regimen will preferably be from about 0.01 to about 80 mg/kg of total body weight. The daily parenteral dosage regimen about 0.001 to about 80 mg/kg of total body weight. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times
30 daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be recognized by one of skill in the art

that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

BIOLOGICAL EXAMPLES

The IL-8, and Gro- α chemokine inhibitory effects of compounds of the present invention were determined by the following *in vitro* assay:

Receptor Binding Assays:

[¹²⁵I] IL-8 (human recombinant) is obtained from Amersham Corp., Arlington Heights, IL, with specific activity 2000 Ci/mmol. Gro- α is obtained from NEN- New England Nuclear. All other chemicals are of analytical grade. High levels of recombinant human IL-8 type α and β receptors were individually expressed in Chinese hamster ovary cells as described previously (Holmes, *et al.*, *Science*, **1991**, 253, 1278). The Chinese hamster ovary membranes were homogenized according to a previously described protocol (Haour, *et al.*, *J Biol Chem.*, 249 pp 2195-2205 (1974)). Except that the homogenization buffer is changed to 10mM Tris-HCL, 1mM MgSO₄, 0.5mM EDTA (ethylene-diaminetetraacetic acid), 1mMPMSF (α -toluenesulphonyl fluoride), 0.5 mg/L Leupeptin, pH 7.5. Membrane protein concentration is determined using Pierce Co. micro-assay kit using bovine serum albumin as a standard. All assays are performed in a 96-well micro plate format. Each reaction mixture contains ¹²⁵I IL-8 (0.25 nM) or ¹²⁵I Gro- α and 0.5 μ g/mL of IL-8R α or 1.0 μ g/mL of IL-8R β membranes in 20 mM Bis-Trispropane and 0.4 mM Tris HCl buffers, pH 8.0, containing 1.2 mM MgSO₄, 0.1

mM EDTA, 25 mM NaCl and 0.03% CHAPS. In addition, drug or compound of interest is added which has been pre-dissolved in DMSO so as to reach a final concentration of between 0.01 nM and 100 μ M. The assay is initiated by addition of ^{125} I-IL-8. After 1 hour at room temperature the plate is harvested using a Tomtec 96-well harvester onto a glass fiber filtermat blocked with 1% polyethylenimine/0.5% BSA and washed 3 times with 25 mM NaCl, 10 mM TrisHCl, 1 mM MgSO₄, 0.5 mM EDTA, 0.03 % CHAPS, pH 7.4. The filter is then dried and counted on the Betaplate liquid scintillation counter. The recombinant IL-8 R α , or Type I, receptor is also referred to herein as the non-permissive receptor and the recombinant IL-8 R β , or Type II, receptor is referred to as the permissive receptor.

Chemotaxis Assay :

The *in vitro* inhibitory properties of these compounds are determined in the neutrophil chemotaxis assay as described in Current Protocols in Immunology, vol I, Suppl 1, Unit 6.12.3., whose disclosure is incorporated herein by reference in its entirety. Neutrophils were isolated from human blood as described in Current Protocols in Immunology Vol I, Suppl 1 Unit 7.23.1, whose disclosure is incorporated herein by reference in its entirety. The chemoattractants IL-8, GRO- α , GRO- β , GRO- γ and NAP-2 are placed in the bottom chamber of a 48 multiwell chamber (Neuro Probe, Cabin John, MD) at a concentration between 0.1 and 100 nM. The two chambers are separated by a 5 μ m polycarbonate filter. When compounds of this invention are tested, they are mixed with the cells (0.001 - 1000 nM) just prior to the addition of the cells to the upper chamber. Incubation is allowed to proceed for between about 45 and 90 min at about 37°C in a humidified incubator with 5% CO₂. At the end of the incubation period, the polycarbonate membrane is removed and the top side washed, the membrane then stained using the Diff Quick staining protocol (Baxter Products, McGaw Park, IL, USA). Cells which have chemotaxed to the chemokine are visually counted using a microscope. Generally, four fields are counted for each sample, these numbers are averaged to give the average number of cells which had migrated. Each sample is tested in triplicate and each compound repeated at least four times. To certain cells (positive

control cells) no compound is added, these cells represent the maximum chemotactic response of the cells. In the case where a negative control (unstimulated) is desired, no chemokine is added to the bottom chamber. The difference between the positive control and the negative control represents the chemotactic activity of the cells.

5 **Elastase Release Assay:**

 The compounds of this invention are tested for their ability to prevent Elastase release from human neutrophils. Neutrophils are isolated from human blood as described in Current Protocols in Immunology Vol I, Suppl 1 Unit 7.23.1. PMNs 0.88×10^6 cells suspended in Ringer's Solution (NaCl 118, KCl 4.56, NaHCO₃ 25, KH₂PO₄ 1.03, Glucose 11.1, HEPES 5 mM, pH 7.4) are placed in
10 each well of a 96 well plate in a volume of 50 ul. To this plate is added the test compound (0.001 - 1000 nM) in a volume of 50 ul, Cytochalasin B in a volume of 50 ul (20ug/ml) and Ringers buffer in a volume of 50 ul. These cells are allowed to warm (37 °C, 5% CO₂, 95% RH) for 5 min before IL-8, GRO α , GRO β , GRO γ or
15 NAP-2 at a final concentration of 0.01 - 1000 nM was added. The reaction is allowed to proceed for 45 min before the 96 well plate is centrifuged (800 xg 5 min) and 100 ul of the supernatant removed. This supernatant is added to a second 96 well plate followed by an artificial elastase substrate (MeOSuc-Ala-Ala-Pro-Val-AMC, Nova Biochem, La Jolla, CA) to a final concentration of 6 ug/ml dissolved in
20 phosphate buffered saline. Immediately, the plate is placed in a fluorescent 96 well plate reader (Cytofluor 2350, Millipore, Bedford, MA) and data collected at 3 min intervals according to the method of Nakajima et al J. Biol Chem 254 4027 (1979). The amount of Elastase released from the PMNs is calculated by measuring the rate of MeOSuc-Ala-Ala-Pro-Val-AMC degradation.

25 **TNF- α in Traumatic Brain Injury Assay**

 The present assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions which follow experimentally induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) were anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and
30 subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia

and surgery without injury, n=18). Animals are sacrificed by decapitation at 1, 6 and 24 hr. post injury, brains removed, and tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA was isolated and Northern blot hybridization is performed and quantitated relative to an TNF- α positive control RNA (macrophage = 100%). A marked increase of TNF- α mRNA expression is observed in LH ($104 \pm 17\%$ of positive control, $p < 0.05$ compared with sham), LC ($105 \pm 21\%$, $p < 0.05$) and LA ($69 \pm 8\%$, $p < 0.01$) in the traumatized hemisphere 1 hr. following injury. An increased TNF- α mRNA expression is also observed in LH ($46 \pm 8\%$, $p < 0.05$), LC ($30 \pm 3\%$, $p < 0.01$) and LA ($32 \pm 3\%$, $p < 0.01$) at 6 hr. which resolves by 24 hr. following injury. In the contralateral hemisphere, expression of TNF- α mRNA is increased in RH ($46 \pm 2\%$, $p < 0.01$), RC ($4 \pm 3\%$) and RA ($22 \pm 8\%$) at 1 hr. and in RH ($28 \pm 11\%$), RC ($7 \pm 5\%$) and RA ($26 \pm 6\%$, $p < 0.05$) at 6 hr. but not at 24 hr. following injury. In sham (surgery without injury) or naive animals, no consistent changes in expression of TNF- α mRNA are observed in any of the 6 brain areas in either hemisphere at any times. These results indicate that following parasagittal fluid-percussion brain injury, the temporal expression of TNF- α mRNA is altered in specific brain regions, including those of the non-traumatized hemisphere. Since TNF- α is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated astrocytes, this post-traumatic alteration in gene expression of TNF- α plays an important role in both the acute and regenerative response to CNS trauma.

CNS Injury model for IL- β mRNA

This assay characterizes the regional expression of interleukin-1 β (IL-1 β) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) are anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia and surgery without injury). Animals are sacrificed at 1, 6 and 24 hr. post injury, brains removed, and


tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA is isolated and Northern blot
5 hybridization was performed and the quantity of brain tissue IL-1 β mRNA is presented as percent relative radioactivity of IL-1 β positive macrophage RNA which was loaded on same gel. At 1 hr. following brain injury, a marked and significant increase in expression of IL-1 β mRNA is observed in LC (20.0 $\pm$ 0.7% of positive control, n=6, p < 0.05 compared with sham animal), LH (24.5 $\pm$ 0.9%, p < 0.05) and
10 LA (21.5 $\pm$ 3.1%, p < 0.05) in the injured hemisphere, which remained elevated up to 6 hr. post injury in the LC (4.0 $\pm$ 0.4%, n=6, p < 0.05) and LH (5.0 $\pm$ 1.3%, p < 0.05). In sham or naive animals, no expression of IL-1 β mRNA is observed in any of the respective brain areas. These results indicate that following TBI, the temporal expression of IL-1 β mRNA is regionally stimulated in specific brain regions. These
15 regional changes in cytokines, such as IL-1 β play a role in the post-traumatic.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

20 The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore the Examples
25 herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is claimed is:

1. A method of treating a chemokine mediated disease state, selected from the group consisting of malaria, restinosis, angiogenesis, atherosclerosis, osteoporosis, gingivitis, undesired hematopoietic stem cells release and diseases caused by respiratory viruses, herpesviruses, and hepatitis viruses, wherein the chemokine binds to an IL-8 α or β receptor in a mammal, which comprises administering to said mammal an effective amount of a compound of the formula:



wherein

Z is cyano, OR_{11} , $\text{C(O)NR}_{15}\text{R}_{16}$, R_{18} , C(O)R_{11} , C(O)OR_{11} , or $\text{S(O)}_2\text{R}_{17}$;

R is any functional moiety having an ionizable hydrogen and a pKa of 10 or less;

R₁ is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C₁₋₁₀ alkyl; C₁₋₁₀ alkyl; C₂₋₁₀ alkenyl; C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀ alkoxy; azide; $(\text{CR}_8\text{R}_8)_q \text{S(O)}_t\text{R}_4$; hydroxy; hydroxy C₁₋₄alkyl; aryl; aryl C₁₋₄ alkyl; aryloxy; aryl C₁₋₄ alkyloxy; heteroaryl; heteroarylalkyl; heterocyclic, heterocyclic C₁₋₄alkyl; heteroaryl C₁₋₄ alkyloxy; aryl C₂₋₁₀ alkenyl; heteroaryl C₂₋₁₀ alkenyl; heterocyclic C₂₋₁₀ alkenyl; $(\text{CR}_8\text{R}_8)_q \text{NR}_4\text{R}_5$; C₂₋₁₀ alkenyl C(O)NR₄R₅; $(\text{CR}_8\text{R}_8)_q \text{C(O)NR}_4\text{R}_5$; $(\text{CR}_8\text{R}_8)_q \text{C(O)NR}_4\text{R}_{10}$; S(O)₃H; S(O)₃R₈; $(\text{CR}_8\text{R}_8)_q \text{C(O)R}_{11}$; C₂₋₁₀ alkenyl C(O)R₁₁; C₂₋₁₀ alkenyl C(O)OR₁₁; $(\text{CR}_8\text{R}_8)_q \text{C(O)OR}_{12}$; $(\text{CR}_8\text{R}_8)_q \text{OC(O)R}_{11}$; $(\text{CR}_8\text{R}_8)_q \text{NR}_4\text{C(O)R}_{11}$; $(\text{CR}_8\text{R}_8)_q \text{NHS(O)}_2\text{R}_{19}$; $(\text{CR}_8\text{R}_8)_q \text{S(O)}_2\text{NR}_4\text{R}_5$; or two R₁ moieties together may form O-(CH₂)_sO- or a 5 to 6 membered saturated or unsaturated ring, and wherein the aryl, heteroaryl and heterocyclic containing moieties may be optionally substituted;

m is an integer having a value of 1 to 3;

q is 0, or an integer having a value of 1 to 10;

s is an integer having a value of 1 to 3;

t is 0, or an integer having a value of 1 or 2;

5 v is 0, or an integer having a value of 1 to 4;

R₄ and R₅ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C₁₋₄alkyl, heterocyclic, heterocyclic C₁₋₄ alkyl, or R₄ and R₅ together with the nitrogen to which they are attached form a 5 to 7 member ring which may optionally comprise an additional heteroatom selected from oxygen, nitrogen or sulfur;

HET is an optionally substituted heteroaryl moiety;

R₆ and R₇ are independently hydrogen or a C₁₋₄ alkyl group; or R₆ and R₇ together with the nitrogen to which they are attached form a 5 to 7 member ring which ring may optionally contain an additional heteroatom selected from oxygen, nitrogen or sulfur;

R₈ is independently selected from hydrogen or C₁₋₄ alkyl;

R₁₀ is C₁₋₁₀ alkyl C(O)₂R₈

R₁₁ is hydrogen, C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;

R₁₂ is hydrogen, C₁₋₁₀ alkyl, optionally substituted aryl or optionally substituted arylalkyl;

25 R₁₃ and R₁₄ are independently hydrogen or C₁₋₄ alkyl;

R₁₅ and R₁₆ are independently hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl, or R₁₅


and R₁₆ may together with the nitrogen to which they are attached form a 5 to 7 member ring optionally containing an additional heteroatom selected from oxygen, nitrogen, or sulfur;

R₁₇ is C₁₋₄ alkyl, NR₁₅R₁₆, OR₁₁, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;


R₁₈ is optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, or optionally substituted heterocyclicC₁₋₄alkyl;

R₁₉ is C₁₋₄alkyl, aryl, arylalkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclic, or heterocyclicC₁₋₄alkyl, wherein the all of these moieties may be optionally substituted;

R_d is NR₆R₇, alkyl, arylC₁₋₄ alkyl, arylC₂₋₄ alkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroarylC₂₋₄ alkenyl, heterocyclic, or heterocyclicC₁₋₄ alkyl, wherein the alkyl, aryl, heteroaryl, and heterocyclic containing moieties may be optionally substituted;



the E' containing ring is optionally selected from



a pharmaceutically acceptable salt thereof.

2. The method according to Claim 1 wherein the ionizable hydrogen has a pKa of 3 to 10.

3. The method according to Claim 2 wherein R is hydroxy, carboxylic acid, thiol, SR₂, OR₂, NH-C(O)R_a, C(O)NR₆R₇, NHS(O)₂R_b, S(O)₂NHR_c, NHC(X₂)NHR_b, or tetrazolyl;

wherein R₂ is a substituted aryl, heteroaryl, or heterocyclic moiety which ring has the functional moiety providing the ionizable hydrogen having a pKa of 10 or less;

R₆' and R₇' are hydrogen, C₁₋₄ alkyl, aryl, arylC₁₋₄alkyl, arylC₂₋₄alkenyl, heteroaryl, heteroarylC₁₋₄alkyl, heteroarylC₂₋₄ alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, or heterocyclic C₂₋₄alkenyl moiety, all of which may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C₁₋₄ alkyl; C₁₋₄ alkyl; C₁₋₄ alkoxy; NR₉C(O)R_a; C(O)NR₆R₇; S(O)₃H; or C(O)OC₁₋₄ alkyl, provided that one of R₆' and R₇' are hydrogen, but not both;

R_a is an aryl, aryl C₁₋₄alkyl, heteroaryl, heteroaryl C₁₋₄alkyl, heterocyclic, or a heterocyclic C₁₋₄alkyl moiety, all of which may be optionally substituted;

R₆ and R₇ are independently hydrogen or a C₁₋₄ alkyl group, or R₆ and R₇ together with the nitrogen to which they are attached form a 5 to 7 member ring

which ring may optionally contain an additional heteroatom which heteroatom is selected from oxygen, nitrogen or sulfur;

R_b is NR_6R_7 , alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, heterocyclic C_{2-4} alkenyl moiety, or camphor, all of which may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C_{1-4} alkyl; C_{1-4} alkyl; C_{1-4} alkoxy; $NR_9C(O)R_a$; $C(O)NR_6R_7$, $S(O)_3H$, or $C(O)OC_{1-4}$ alkyl;

R_9 is hydrogen or C_{1-4} alkyl;

R_c is alkyl, aryl, aryl C_{1-4} alkyl, aryl C_{2-4} alkenyl, heteroaryl, heteroaryl C_{1-4} alkyl, heteroaryl C_{2-4} alkenyl, heterocyclic, heterocyclic C_{1-4} alkyl, or a heterocyclic C_{2-4} alkenyl moiety, all of which may be optionally substituted one to three times independently by halogen, nitro, halosubstituted C_{1-4} alkyl, C_{1-4} alkyl, C_{1-4} alkoxy, $NR_9C(O)R_a$, $C(O)NR_6R_7$, $S(O)_3H$, or $C(O)OC_{1-4}$ alkyl; and

X_2 is oxygen or sulfur.

4. The method according to Claim 3 wherein the R_2 is optionally substituted one to three times by halogen, nitro, halosubstituted C_{1-10} alkyl, C_{1-10} alkyl, C_{1-10} alkoxy, OH, SH, $C(O)NR_6R_7$, $NH-C(O)R_a$, $NHS(O)R_b$, $S(O)NR_6R_7$, $C(O)OR_8$, or a tetrazolyl ring.

5. The method according to Claim 3 wherein R is OH, $-NHS(O)_2R_b$ or $C(O)OH$.

6. The method according to Claim 1 wherein R_1 is halogen, cyano, nitro, CF_3 , $C(O)NR_4R_5$, alkenyl $C(O)NR_4R_5$, $C(O)R_4R_{10}$, alkenyl $C(O)OR_{12}$, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, optionally substituted heteroaryl alkenyl, or $S(O)NR_4R_5$.

7. The method according to Claim 1 wherein HET is substituted by (Y)_n;

n is an integer having a value of 1 to 3;

Y is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted

C₁₋₁₀ alkyl; C₁₋₁₀ alkyl; C₂₋₁₀ alkenyl; C₁₋₁₀ alkoxy; halosubstituted C₁₋₁₀
 5 alkoxy; azide; (CR₈R₈)_q S(O)_tR₄; hydroxy; hydroxyC₁₋₄alkyl; aryl; aryl C₁₋₄
 alkyl; aryloxy; arylC₁₋₄ alkyloxy; heteroaryl; heteroarylalkyl; heteroaryl C₁₋₄
 alkyloxy; heterocyclic, heterocyclic C₁₋₄alkyl; aryl C₂₋₁₀ alkenyl; heteroaryl
 C₂₋₁₀ alkenyl; heterocyclic C₂₋₁₀ alkenyl; (CR₈R₈)_q NR₄R₅; C₂₋₁₀ alkenyl
 C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₅; (CR₈R₈)_q C(O)NR₄R₁₀; S(O)₃H;
 10 S(O)₃R₈; (CR₈R₈)_q C(O)R₁₁; C₂₋₁₀ alkenyl C(O)R₁₁; C₂₋₁₀ alkenyl
 C(O)OR₁₁; (CR₈R₈)_q C(O)OR₁₂; (CR₈R₈)_q OC(O) R₁₁; (CR₈R₈)_q
 NR₄C(O)R₁₁; (CR₈R₈)_q NHS(O)₂R_d; or (CR₈R₈)_q S(O)₂NR₄R₅ or two Y
 moieties together may form O-(CH₂)_sO- or a 5 to 6 membered saturated or
 unsaturated ring; wherein the aryl, heteroaryl, heterocyclic containing moieties
 15 may be optionally substituted as defined herein; and

R_d is NR₆R₇, alkyl, arylC₁₋₄ alkyl, arylC₂₋₄ alkenyl, heteroaryl,
 hetroaryl-C₁₋₄alkyl, heteroarylC₂₋₄ alkenyl, heterocyclic, heterocyclicC₁₋₄
 alkyl, wherein the alkyl the aryl, heteroaryl, heterocyclic containing moieties
 may be optionally substituted.

20

8. The method according to Claim 7 wherein Y is halogen, C₁₋₄ alkoxy,
 optionally substituted aryl, optionally substituted arylalkoxy, methylene dioxy,
 NR₄R₅, thioC₁₋₄alkyl, thioaryl, halosubstituted alkoxy, optionally substituted C₁₋₄
 4alkyl, hydroxy alkyl.

25

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/16814

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :A61K 31/47, 31/495, 31/38, 31/17

US CL :514/311, 369, 445, 586, 597

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/311, 369, 445, 586, 597

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN: COMPOUNDS AND IL-8 ANTAGONISTS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,780,483 A (WIDDOWSON ET AL) 14 JULY 1998 (14/07/98), see entire document.	1-8
A	US 4,405,644 A (KABBE ET AL) 20 SEPTEMBER 1983 (20/09/83), see entire document.	1-8
A	EP 0 344 425 A2 (WARNER-LAMBERT CORP) 06 DECEMBER 1989 (06/12/89), see entire document.	1-8

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
22 AUGUST 2000

Date of mailing of the international search report
07 SEP 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230

Authorized officer *my mas*
RUSSELL TRAVERS
Telephone No. (703) 308-1235