

(19) **DANMARK**

(10) **DK/EP 3810201 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 47/02 (2006.01)** **A 61 K 38/26 (2006.01)** **A 61 K 47/10 (2017.01)**
A 61 P 3/10 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2023-06-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-06-07**
- (86) Europæisk ansøgning nr.: **19739766.4**
- (86) Europæisk indleveringsdag: **2019-06-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-04-28**
- (86) International ansøgning nr.: **US2019037146**
- (87) Internationalt publikationsnr.: **WO2019245893**
- (30) Prioritet: **2018-06-22 US 201862688632 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285, USA**
- (72) Opfinder: **CORVARI, Vincent John, c/o Eli Lilly and Company P.O. Box 6288, Indianapolis, Indiana 46206-6288, USA**
MINIE, Christopher Sears, c/o Eli Lilly and Company P.O. Box 6288, Indianapolis, Indiana 46206-6288, USA
MISHRA, Dinesh Shyandeo, c/o Eli Lilly and Company P.O. Box 6288, Indianapolis, Indiana 46206-6288, USA
QIAN, Ken Kangyi, c/o Eli Lilly and Company P.O. Box 6288, Indianapolis, Indiana 46206-6288, USA
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **GIP/GLP1-agonist-sammensætninger**
- (56) Fremdragne publikationer:
US-A1- 2012 329 708
US-B2- 9 474 780

DESCRIPTION

[0001] The present invention is a pharmaceutical GIP/GLP1 co-agonist peptide composition for subcutaneous injection. The composition comprises tirzepatide, NaCl, and dibasic sodium phosphate. The composition provides commercially acceptable shelf-life stability, in-use stability, and is associated with acceptable patient injection site experience.

[0002] Diabetes mellitus is a chronic disorder characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both. In type 2 diabetes mellitus ("T2D"), the combined effects of impaired insulin secretion and insulin resistance are associated with elevated blood glucose levels. Tirzepatide is a GIP/GLP1 co-agonist peptide useful in the treatment of diabetes. Tirzepatide is useful in the treatment of obesity.

[0003] US9474780 generally describes compositions containing a GIP/GLP1 agonist, administered by parenteral routes. US9474780 describes and claims tirzepatide. There is a desire for compositions of tirzepatide providing acceptable stability and acceptable patient injection site experience.

[0004] The present invention seeks to meet these needs by providing pharmaceutically-acceptable compositions comprising tirzepatide, or a pharmaceutically acceptable salt thereof; NaCl; and dibasic sodium phosphate.

[0005] In an embodiment, the NaCl concentration is from about 6.2 mg/mL to about 9.5 mg/mL. In an embodiment the NaCl concentration is from about 7.0 mg/mL to about 9.0 mg/mL. In an embodiment, the NaCl concentration is about 8.2 mg/mL.

[0006] In an embodiment, the dibasic sodium phosphate concentration is from about 0.67 mg/mL to about 2.68 mg/mL. In an embodiment, the dibasic sodium phosphate is about 1.0 mg/mL to about 3.0 mg/mL. In an embodiment, the dibasic sodium phosphate is about 1.34 mg/mL.

[0007] In an embodiment, the tirzepatide concentration is from about 5 mg/mL to about 30 mg/mL. In an embodiment, the tirzepatide concentration is from about 10 mg/mL to about 30 mg/mL. In an embodiment, the tirzepatide concentration is from about 5 mg/mL to about 30 mg/mL; the NaCl concentration is about 8.2 mg/mL, and dibasic sodium phosphate concentration is from about 0.67 mg/mL to about 2.68 mg/mL. In an embodiment, the tirzepatide concentration is from about 5 mg/mL to about 30 mg/mL; the NaCl concentration is about 8.2 mg/mL, dibasic sodium phosphate concentration is from about 0.67 mg/mL to about 2.68 mg/mL, and the composition is presented in a single use automatic injection apparatus.

[0008] In an embodiment, the tirzepatide concentration is from about 5 mg/mL to about 30 mg/mL. In certain embodiments, the tirzepatide concentration is from about 10 mg/mL to about 30 mg/mL. In an embodiment the tirzepatide concentration is selected from the group

consisting of 5, 10, 15, 20, 25, and 30 mg/mL. In an embodiment, 0.5 mL or less of the composition is administered as a dose. In an embodiment, the tirzepatide concentration is selected from the group consisting of 10, 20, and 30 mg/mL.

[0009] In an embodiment, the tirzepatide composition further comprises a preservative. In an embodiment, the tirzepatide composition comprises tirzepatide, dibasic sodium phosphate, propylene glycol, and a preservative. In an embodiment, the preservative is selected from the group consisting of metacresol and phenol. In an embodiment, the metacresol concentration is from about 2.0 mg/mL to about 4.0 mg/mL. In an embodiment the metacresol concentration is from about 3.0 mg/mL to about 3.5 mg/mL. In an embodiment, the metacresol concentration is about 3.15 mg/mL. In an embodiment, the phenol concentration is from about 3.0 mg/mL to about 7.0 mg/mL. In an embodiment the phenol concentration is from about 4.0 mg/mL to about 6.0 mg/mL. In an embodiment, the phenol concentration is about 5.0 mg/mL.

[0010] In an embodiment, the dose of a tirzepatide composition is administered about once weekly. In an embodiment, the dose of a tirzepatide composition is administered once every seven days.

[0011] In an embodiment, there is provided one of the above-described compositions for use as a medicament.

[0012] In an embodiment, there is provided one of the above-described compositions for use in the treatment of diabetes. In an embodiment, there is provided one of the above-described compositions for use in the treatment of obesity.

[0013] In an embodiment, there is provided one of the above-described compositions for use in providing therapeutic weight loss. In an embodiment, there is provided one of the above-described compositions for use in providing non-therapeutic weight loss.

[0014] According to another aspect of the present invention, there is provided an article of manufacture comprising one of the above-described compositions. In certain embodiments, the article of manufacture is a multi-use vial. In certain embodiments, the article of manufacture is a pre-filled syringe. In certain embodiments, the article of manufacture is an automatic injection apparatus ("auto-injector"). An example of an auto-injector, as contemplated herein, is presented in U.S. Patent 8,734,394.

[0015] As used herein, "tirzepatide" means a GIP/GLP1 co-agonist peptide as described in US 9,474,780 and described by CAS Registry Number: 2023788-19-2. Tirzepatide is described in Example 1 of US 9474,780, with the following sequence:

YX1EGTFTSDYSIX2LDKIAQKAFVQWLIAGGPSSGAPPPS

wherein X₁ is Aib; X₂ is Aib; K at position 20 is chemically modified through conjugation to the epsilon-amino group of the K side-chain with (2-[2-(2-Amino-ethoxy)-ethoxy]-acetyl)₂-(γGlu)₁-CO-(CH₂)₁₈-CO₂H; and the C-terminal amino acid is amidated as a C-terminal primary amide (SEQ ID NO: 1).

[0016] When used herein, "pharmaceutically acceptable salt" is well known to the skilled artisan. In an embodiment is a pharmaceutically acceptable salt that is a tirzepatide trifluoroacetate salt.

[0017] When used herein, the term "does not contain a surfactant" means that the composition contains no added surfactant agents, or contains only a *de minimis* quantity of an added surfactant.

[0018] When used herein the term "propylene glycol" is well known to the skilled artisan. Propylene glycol is also represented by the formula: $C_3H_8O_2$.

[0019] The compositions of the present invention have concentrations of tirzepatide between 5 mg/mL and 30 mg/mL. The compositions of the present invention are likely to have specific concentrations of 5, 10, 15, 20, 25, and 30 mg/mL. Such compositions may be presented in a pre-filled syringe. Such pre-filled syringe may be useful for administering one half milliliter of such composition per patient per dose. A dose of tirzepatide composition may be administered using a dosing schedule determined by a physician.

[0020] The compositions are sterile when first produced. If provided in a multi-use vial or cartridge, an anti-microbial preservative compound or mixture of compounds that is compatible with the other components of the composition may be added at sufficient strength to meet applicable regulatory anti-microbial preservative requirements. Pharmaceutically acceptable preservatives are well-known in the art. (See, e.g., Remington: The Science and Practice of Pharmacy (D.B. Troy, Editor, 21st Edition, Lippincott, Williams & Wilkins, 2006). In an embodiment, the preservative is metacresol. In an embodiment, the preservative is phenol. A composition for single use pre-filled syringe requires no preservative. In an embodiment, the composition does not contain a surfactant.

[0021] The pH of tirzepatide compositions of the present invention is typically 6.5 to 7.5 and it is adjusted using physiologically appropriate acids and bases, as may be required to achieve the desired pH. In an embodiment, the pH target is between 6.7 and 7.3. Patient injection site experience is a consideration for a subcutaneously administered composition. It is desirable to select a composition associated with an acceptable patient injection site experience. For example, NaCl and citrate have been associated with painful stinging at the injection site. (Laurson, T.; Hansen, B.; Fisker, S. Pain perception after subcutaneous injections of media containing different buffers. Basic & Clinical Pharmacology & Toxicology 2006, 98, (2), 218-221.), (Fransson, J.; Espander-Jansson, Local tolerance of subcutaneous injections. Journal of Pharmacy and Pharmacology 1996, 48, (10), 1012-1015.) It is further desirable to match the tonicity (i.e., osmolality) of body fluids at the injection site as closely as possible when administering the compositions because solutions that are not approximately isotonic with body fluids can produce a painful stinging sensation when administered. It is desirable that the compositions be approximately isotonic with body fluids at the sites of injection. The present composition comprising tirzepatide, NaCl, and dibasic sodium phosphate is associated

acceptable patient injection site experience. Likewise, the composition comprising tirzepatide, propylene glycol, and dibasic sodium phosphate is associated with acceptable patient injection site experience.

[0022] In an embodiment, the pH is adjusted using a base to facilitate dissolution in the buffer solution. The addition of an acid to the composition may be required to adjust the pH to the desired pH range. In an embodiment, NaOH is used to facilitate dissolution of tirzepatide in a buffer. In an embodiment, HCl is added to adjust the pH of the composition containing the dissolved tirzepatide to the desired pH range.

[0023] The compositions of the present invention are typically administered subcutaneously. The compositions are typically administered using a pre-filled, disposable pen, reusable pen, or automatic pen injector. The composition may be administered using a multi-use vial or a pump device. In an embodiment, the device is an automatic injection apparatus as claimed by U.S. Patent 8,734,394.

[0024] A composition comprising tirzepatide, NaCl, and dibasic sodium phosphate provides a desired shelf life stability and provides patients with an acceptable injection site experience. Likewise, a composition comprising tirzepatide, propylene glycol, and dibasic sodium phosphate provides a desired shelf life stability and provides patients with an acceptable injection site experience. As used herein, "shelf life stability" is measured under controlled conditions at about 5 degrees Celsius. A composition comprising tirzepatide, NaCl, and dibasic sodium phosphate provides acceptable in-use stability. Likewise, a composition comprising tirzepatide, propylene glycol, and dibasic sodium phosphate provides acceptable in-use stability. As used herein, the term "in-use stability" refers to the stability of the composition measured under controlled conditions at or about 25 degrees Celsius or at or about 40 degrees Celsius.

Example #1 - Composition containing NaCl

[0025] The composition is prepared substantially as described herein. The compositions containing 5, 10, 15, 20, 15, and 30 mg/mL of tirzepatide each contain the ingredients set forth in Table 1. Acid or base is optionally added to attain the desired pH range. Water is added *quantum satis* (q.s.) to one milliliter total final volume.

Table 1. Formulation of Tirzepatide, Phosphate, and NaCl

Ingredient	Concentration (mg/mL)
Tirzepatide	5, 10, 15, 20, 25, and 30
dibasic sodium phosphate*	1.34
NaCl	8.2
*5 mM.phosphate buffer is used	

Reference Example #2 - Composition containing Propylene glycol

[0026] The composition is prepared substantially as described herein. The compositions providing 5, 10, 15, 20, 15, and 30 mg/mL compositions of tirzepatide each contain the ingredients set forth in Table 2. Acid or base is optionally added to attain the desired pH range. Water is added *quantum satis* to one milliliter total final volume.

Table 2. Formulation of Tirzepatide, Phosphate, and Propylene Glycol

Ingredient	Concentration (mg/mL)
Tirzepatide	5, 10, 15, 20, 25, and 30
dibasic sodium phosphate*	1.34
Propylene glycol	15
*5mM phosphate buffer is used	

Size Exclusion Chromatography (SEC) In-Use Stability Study

[0027] This procedure is an isocratic size exclusion HPLC method with UV detection at 214 nm and is designed to determine the relative amounts of tirzepatide monomer and total aggregates. Monomer and aggregates are reported as peak area percent to the total area. The procedure is stability indicating as measured by its ability to resolve known impurities from tirzepatide. This study compares alternate compositions with the compositions of this invention, prepared as shown by Table 3. The stability from this study is shown in Table 4.

Stability study comparing alternate compositions: Table 3

Ingredient	Formulation			
	Control	NaCl	Mannitol	Glycerol
Tirzepatide (mg/mL)	2	2	2	2
Sodium phosphate dibasic 7H ₂ O (mg/mL) **	2.68	2.68	2.68	2.68
NaCl (mg/mL)	----	8.8	----	----
Mannitol (mg/mL)	-----	-----	45	-----
Glycerol (mg/mL)	-----	-----	-----	27
Water (mg/mL)	q.s. to 1 mL	q.s. to 1 mL	q.s. to 1 mL	q.s. to 1 mL
**10 mM phosphate buffer is used				

Table 4. Tirzepatide monomer (% peak area) by size-exclusion chromatography (SEC)

Composition	Temp (°C)	Time (month)			
		0	0.5	1	2
Control (Table 3)	25	98.25	98.15	97.76	96.63
	40		97.62	95.80	93.19
NaCl (Table 3)	25	98.23	98.13	97.95	97.47
	40		97.84	97.28	96.22
Mannitol (Table 3)	25	98.23	97.87	97.39	95.17
	40		96.72	92.71	83.48
Glycerol (Table 3)	25	98.25	98.11	97.77	96.62
	40		97.36	94.51	84.11

Shelf Life Stability Study

RP-HPLC:

[0028] This procedure is a gradient reversed-phase HPLC method with UV detection at 214 nm and is designed to determine the quantity, identity, and purity of tirzepatide in the drug product. Identity is determined by matching the retention time of the main peak with that of the main peak of an external reference standard. Quantity is determined by the comparison of the main peak area with the corresponding peak in the external reference standard. Impurities and related substances are reported as peak area percent to the total peak area. The procedure is stability indicating as judged by its ability to resolve known impurities from tirzepatide. As shown by Table 4, a composition comprising NaCl as a tonicity agent provides acceptable in-use stability.

Size Exclusion Chromatography (SEC) Shelf-Life Stability Study

[0029] The size exclusion stability study methods and RP-HPLC described herein above are applied to compare compositions comprising NaCl as a tonicity and stabilizing agent with compositions comprising propylene glycol as a tonicity and stabilizing agent. This study illustrates the acceptable shelf-life stability of a composition of this invention comprising NaCl agent or comprising propylene glycol as agent. The compositions used for this study are set forth in Table 5. The stability from this study is shown in Table 6.

Table 5. Comparison between NaCl and propylene glycol

Ingredient	Formulation of NaCl as agent	Reference Formulation of Propylene glycol as agent
Tirzepatide (mg/mL)	20	20
Sodium phosphate dibasic 7H ₂ O (mg/mL)*	1.34	1.34
NaCl (mg/mL))	8.8	-----
Propylene glycol (mg/mL)	-----	15
Water	q.s. to mL	q.s. to mL
*5 mM phosphate buffer is used in the study		

Table 6. Comparison of NaCl to Propylene Glycol as an agent in Tirzepatide formulations by determining monomer (% peak area) by SEC and purity by RP-HPLC

Monomer by SEC					
Composition	Temp (°C)	Time (month)			
		0	1	2	3
NaCl (Table 5)	5	99.35	-	-	98.81
	30		97.20	98.03	97.79
Propylene glycol (Table 5)	5	98.51	98.20		
	30		97.94		
Purity by RP-HPLC					
Composition	Temp (°C)	Time (month)			
		0	1	2	3
NaCl (Table 5)	5	92.87	-	-	95.95
	30		91.70	90.40	89.26
Propylene glycol (Table 5)	5	93.38	93.31		
	30		92.04		

Pain upon injection study:

[0030] All compositions are prepared as described by Table 7. Each solution composition vial is held at room temperature about 30 minutes, but not more than four hours. Reconstituted lyophilized compositions are used immediately. All injections are rotated between the 4 quadrants of the abdomen, in the following order; lower left quadrant; lower right quadrant; upper left quadrant; and upper right quadrant. A syringe with a 29 gauge needle is used to administer the composition from a vial. A fold of skin at the injection site is grasped by the subject, and the needle is inserted at about a 45 degree angle. A second person uses a stop

watch to measure the length of time of injection. The subject slowly pushes the plunger of the syringe all the way until 0.5 mL of the composition is injected. The target time of injection is 4 seconds duration, and not more than 5 seconds. The needle is removed from the skin after injection and skin is released from the subject's grasp. The subject immediately assesses the pain after each injection. Pain measurements are assessed using a 100-mm validated visual analog scale (VAS) for pain. The VAS is a well-validated tool to assess injection-site pain (Williamson, A.; Hoggart, Pain: A review of three commonly used pain rating scales. Journal of Clinical Nursing 2005, 14, (7), 798-804). The VAS is presented as a 10-cm (100-mm) line, anchored by verbal descriptors, usually "no pain" and "worst imaginable pain." The subject is asked to mark the 100-mm line to indicate pain intensity at time points and as clinically indicated. A staff member uses a caliper to measure the distance from 0 to the mark that the subject placed on the VAS, and to record the measurement in the source document. Results from this study appear in Table 8. An acceptable patient injection site experience is reflected by a mild pain intensity indication (compared to moderate or severe).

Table 7

Compositions for Pain Intensity Study Ingredient	Formulation of NaCl	Reference Formulation of Propylene glycol
Tirzepatide (mg/mL)	20	20
Sodium phosphate dibasic 7H ₂ O (mg/mL)*	1.34	1.34
NaCl (mg/mL)	8.8	-----
Propylene glycol (mg/mL)	-----	15
Water	<i>q.s.</i> to mL	<i>q.s.</i> to mL
Ingredient	Placebo of NaCl as agent	Placebo of Propylene glycol as agent
Tirzepatide (mg/mL)	0	0
Sodium phosphate dibasic 7H ₂ O (mg/mL)*	1.34	1.34
NaCl (mg/mL))	8.8	-----
Propylene glycol (mg/mL)	-----	15
Water	<i>q.s.</i> to mL	<i>q.s.</i> to mL
Ingredient	Formulation of NaCl as agent-Lyophilized	
Tirzepatide (mg/mL)	20	
Sodium phosphate dibasic 7H ₂ O (mg/mL)*	1.34	
NaCl (mg/mL))	8.8	

Ingredient	Formulation of NaCl as agent-Lyophilized	
Propylene glycol (mg/mL)	-----	
Water: sterile water for injection-reconstitute; swirl to dissolve	q.s. to mL	
*5 mM phosphate buffer is used in the study		

Table 8. VAS pain score.

Composition (per Table 7)	Pain intensity		
	Mild (VAS: 0 mm - 30 mm)	Moderate (VAS: 31 mm - 70 mm)	Severe (VAS: 71 mm -100 mm)
Tirzepatide lyophilized, reconstituted	90%	5%	5%
Reference Tirzepatide composition propylene glycol	100%	0%	0%
Tirzepatide composition NaCl	100%	0%	0%
Placebo composition NaCl	100%	0%	0%
Placebo composition propylene glycol	100%	0%	0%

Sequences

[0031] SEQ ID NO:1

Tirzepatide

YX₁EGTFTSDYSIX₂LDKIAQKAFVQWLIAGGPSSGAPPPS

wherein X₁ is Aib; X₂ is Aib; K at position 20 is chemically modified through conjugation to the epsilon-amino group of the K side-chain with (2-[2-(2-Amino-ethoxy)-ethoxy]-acetyl)₂-(γGlu)₁-CO-(CH₂)₁₈-CO₂H; and the C-terminal amino acid is amidated as a C-terminal primary amide

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US9474780B [0003] [0003] [0015] [0015]
- US8734394B [0014] [0023]

Non-patent literature cited in the description

- CHEMICAL ABSTRACTS, 2023788-19-2 [0015]
- Remington: The Science and Practice of Pharmacy Lippincott, Williams & Wilkins 2006 0000 [0020]
- LAURSEN, T. HANSEN, B. FISHER, S. Pain perception after subcutaneous injections of media containing different buffers Basic & Clinical Pharmacology & Toxicology, 2006, vol. 98, 2218-221 [0021]
- FRANSSON, J. Espander-Jansson, Local tolerance of subcutaneous injections Journal of Pharmacy and Pharmacology, 1996, vol. 48, 101012-1015 [0021]
- WILLIAMSON, A.; HOGGART Pain: A review of three commonly used pain rating scales Journal of Clinical Nursing, 2005, vol. 14, 7798-804 [0030]

P a t e n t k r a v

1. Farmaceutisk sammensætning omfattende tirzepatid eller et farmaceutisk acceptabelt salt deraf; NaCl; og diabasisk natriumphosphat.

5

2. Farmaceutisk sammensætning ifølge krav 1, hvor koncentrationen af tirzepatidet eller et farmaceutisk acceptabelt salt deraf er fra ca. 5 til ca. 30 mg/ml.

10

3. Farmaceutisk sammensætning ifølge krav 2, hvor koncentrationen af tirzepatidet eller et farmaceutisk acceptabelt salt deraf er 5, 10, 15, 20, 25 eller 30 mg/ml.

15

4. Farmaceutisk sammensætning ifølge krav 3, hvor koncentrationen af tirzepatidet eller et farmaceutisk acceptabelt salt deraf er 10, 20 eller 30 mg/ml.

20

5. Farmaceutisk sammensætning ifølge krav 1, hvor koncentrationen af diabasisk natriumphosphat er fra ca. 1,0 mg/ml til ca. 3,0 mg/ml.

6. Farmaceutisk sammensætning ifølge krav 1, hvor den dibasiske natriumphosphatkonzentration er fra ca. 0,67 mg/ml til ca. 2,68 mg/mL.

7. Farmaceutisk sammensætning ifølge krav 5 eller krav 6, hvor den diabasiske natriumphosphatkonzentration er ca. 1,34 mg/ml.

25

8. Farmaceutisk sammensætning ifølge krav 1, hvor koncentrationen af NaCl er fra ca. 6,2 mg/ml til ca. 9,5 mg/ml.

9. Farmaceutisk sammensætning ifølge krav 8, hvor NaCl-koncentrationen er ca. 8,2 mg/ml.

30

10. Farmaceutisk sammensætning ifølge krav 1, hvor koncentrationen af tirzepatid eller farmaceutisk acceptabelt salt deraf er fra ca. 5 mg/ml til ca. 30 mg/ml; dibasisk natriumphosphatkonzentration er fra ca. 0,67 mg/ml til ca. 2,68 mg/ml; og koncentrationen af NaCl er fra ca. 6,2 mg/ml til ca. 9,5 mg/ml.

5

11. Farmaceutisk sammensætning ifølge krav 10, hvor koncentrationen af diabasisk natriumphosphat er ca. 1,34 mg/ml; og NaCl-konzentrationen er ca. 8,2 mg/ml.

10

12. Farmaceutisk sammensætning ifølge krav 11, hvor sammensætningen præsenteres i et automatisk injektionsapparat.

13. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 12, hvor pH af sammensætningen er fra ca. 6,5 til ca. 7,5.

15

14. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 13, yderligere omfattende et eller flere konserveringsmidler.

15. Farmaceutisk sammensætning ifølge krav 14, hvor sammensætningen yderligere omfatter et konserveringsmiddel valgt blandt metacresol og phenol.

20

16. Farmaceutisk sammensætning ifølge krav 15, hvor konserveringsmidlet er metacresol i en koncentration på ca. 2,0 mg/ml til ca. 4,0 mg/ml.

25

17. Farmaceutisk sammensætning ifølge krav 15, hvor konserveringsmidlet er phenol i en koncentration fra ca. 3,0 mg/ml til ca. 7,0 mg/ml.

18. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 17 til anvendelse i behandlingen af diabetes.

30

19. Farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 17 til anvendelse i behandlingen af fedme.