



(51) International Patent Classification:

A61K 31/145 (2006.01) *A61K 31/445* (2006.01)
A61K 31/397 (2006.01) *A61K 31/4465* (2006.01)
A61K 31/40 (2006.01) *C07C 323/30* (2006.01)
A61K 31/4402 (2006.01) *C07D 211/54* (2006.01)
A61K 31/4409 (2006.01)

(21) International Application Number:

PCT/US2016/040637

(22) International Filing Date:

1 July 2016 (01.07.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/187,939 2 July 2015 (02.07.2015) US
62/387,337 23 December 2015 (23.12.2015) US

(71) Applicant: **RAPTOR PHARMACEUTICALS INC.**
[US/US]; 7 Hamilton Landing, Suite 100, Novato, CA
94949 (US).

(72) Inventors: **ZANKEL, Todd, C.**; 6930 California Street,
San Francisco, CA 94121 (US). **UNITT, John**; BioCity,
Pennyfoot Street, Nottingham NG1 1GF (GB). **PHILLIPS,**
Timothy; BioCity, Pennyfoot Street, Nottingham NG1
1GF (GB). **GOURDET, Benoit**; BioCity, Pennyfoot
Street, Nottingham NG1 1GF (GB). **DUFFY, Loma**;
BioCity, Pennyfoot Street, Nottingham NG1 1GF (GB).

(74) Agents: **PLUMMER, Kelly, A.** et al.; Marshall, Gerstein
& Borun LLP, 233 S. Wacker Drive, 6300 Willis Tower,
Chicago, IL 60606-6357 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: ADO-RESISTANT CYSTEAMINE ANALOGS AND USES THEREOF

(57) Abstract: The present invention is directed to ADO-resistant cysteamine analogs for use in methods of treating diseases for which cysteamine is indicated, in particular cystinosis, fatty liver disease, fibrosis, thrombotic diseases, MECP-2 related disorders, inherited mitochondrial diseases, neurological diseases or disorders, inflammation and cancer.



ADO-RESISTANT CYSTEAMINE ANALOGS AND USES THEREOF**FIELD OF THE DISCLOSURE**

[0001] The disclosure relates to materials and methods to treat diseases in which therapy with cysteamine is indicated. In particular, the disclosure provides therapeutic methods involving administration to a patient of a compound as disclosed herein.

BACKGROUND

[0002] Cysteamine (HS-CH₂-CH₂-NH₂) is a small sulfhydryl compound that is able to cross cell membranes easily due to its small size. Cysteamine is currently FDA approved for use in the treatment of cystinosis, an intra-lysosomal cystine storage disorder. In cystinosis, cysteamine acts by converting cystine to cysteine and cysteine-cysteamine mixed disulfide, which are then both able to leave the lysosome through the cysteine and lysine transporters respectively (Gahl *et al.*, N Engl J Med 347(2):111-21, 2002). Within the cytosol, the mixed disulfide can be reduced by its reaction with glutathione and the cysteine released can be used for further GSH synthesis. Treatment with cysteamine has been shown to result in lowering of intracellular cystine levels in circulating leukocytes (Dohil *et al.*, J. Pediatr 148(6):764-9, 2006).


[0003] Cysteamine is converted to hypotaurine by cysteamine dioxygenase (ADO) (Coloso *et al.* (2006) Adv Exp Med Biol 583, 25-36; Dominy *et al.* (2007) J Biol Chem 282, 25189-25198; Richerson *et al.* (1987) Methods Enzymol 143, 410-415) and then ultimately to taurine, the most common amino acid in the body. Cysteamine is also discussed in Prescott *et al.*, (Lancet 2(7778):652, 1979); Prescott *et al.*, (Br Med J 1(6116):856-7, 1978); Mitchell *et al.*, (Clin Pharmacol Ther 16(4):676-84, 1974); de Ferreyra *et al.*, (Toxicol Appl Pharmacol. 48(2):221-8, 1979); and Qiu *et al.*, (World J Gastroenterol. 13:4328-32, 2007). Unfortunately, the sustained concentrations of cysteamine necessary for therapeutic effect are difficult to maintain due to rapid metabolism and clearance of cysteamine from the body, with nearly all administered cysteamine converted to taurine in a matter of hours. These difficulties are transferred to patients in the form of high dosing levels and frequencies, with all of the consequent unpleasant side effects associated with cysteamine (e.g., gastrointestinal distress and body odor). See the package insert for CYSTAGON® (cysteamine bitartrate). International Publication No. WO 2007/079670 and US Patents 8,026,2854 and 8,129,433 disclose enterically coated cysteamine products and a method of reducing dosing frequency of cysteamine.

[0004] Cysteamine is addressed in International Patent Application Nos. WO 2009/070781, and WO 2007/089670, and U.S. Patent Publication Nos. 20110070272, 20090048154, and 20050245433.

SUMMARY

[0005] The present disclosure provides methods of treating a patient suffering from a disease for which treatment with cysteamine is indicated. The methods comprise administering to the patient an effective amount of a composition comprising a compound as disclosed herein. It is contemplated that administration of the composition reduces levels of cystine in patients, which can improve the detrimental effects of elevated cystine levels.

[0006] Suitable compositions comprise a compound of formula I or a disulfide thereof:



wherein:

R^1 and R^2 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^1 and R^2 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^3 and R^4 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

G is selected from the group consisting of $-\text{NR}^5\text{R}^6$ and $-\text{CR}^7\text{R}^8\text{NR}^5\text{R}^6$;

R^5 and R^6 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

R^7 and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^7 and R^8 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^2 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;

R^4 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;

R^2 and R^8 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring; or

R^2 and R^4 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring.

[0007] In some cases, when G is $-\text{NH}_2$, at least one of R^1 , R^2 , R^3 , and R^4 is other than H.

[0008] In some cases, R^5 and R^6 are independently selected from the group consisting of H, methyl, and ethyl. In some cases, R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 5-membered heterocyclic ring.

[0009] In some cases, wherein R^4 is methyl and/or R^3 is methyl. In some cases, R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.


[0010] In some cases, R^2 is methyl and/or R^1 is methyl. In some cases, R^1 and R^2 , taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.

[0011] In some cases, G is $-\text{CR}^7\text{R}^8\text{NR}^5\text{R}^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 6-membered heterocyclic ring. In some cases, R^5 is methyl.


[0012] In some cases, G is $-\text{NR}^5\text{R}^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 4- or 6-membered heterocyclic ring. In some cases, R^5 is H.

[0013] In some cases, R^7 and R^8 are both H.

[0014] A compound of formula I includes, but is not limited to, the following compounds:




[0015] A compound of formula I includes, but is not limited to, the following compounds:



[0016] In some cases, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl. In some cases, R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 are independently selected from the group consisting of H and methyl.

[0017] Suitable compositions comprise a compound of formula II, formula III, or a disulfide thereof:



wherein:

L is a hydrocarbon linking group;

R^9 and R^{10} are independently selected from the group consisting of H, C_{1-5} alkyl, and $CO(C_{1-5}alkyl)$; or

R^9 and R^{10} , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

A is a heterocyclic ring contain one N atom; and

n is 0, 1, 2, or 3.


[0018] In some cases, the compound of formula II is not cysteamine.

[0019] In some cases, the S atom in the compound of formula II or formula III is a distance of about 3.6 Angstroms to about 4.7 Angstroms from the N atom in the compound, such as about 3.8 Angstroms to about 4.4 Angstroms, about 4.0 Angstroms to about 4.2 Angstroms, or about 4.1 Angstroms from the N atom in the compound.

[0020] In some cases, L is a 3-, 4-, 5-, 6-, 7-, or 8-membered cycloalkyl ring or a 6-membered aryl ring. In some cases, L is C_{1-5} alkyl. In some cases, L is substituted with one to four groups selected from halo, C_{1-5} alkyl, C_{3-5} cycloalkyl, and $-CO_2(C_{1-5}alkyl)$.

[0021] In some cases, A is a 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic heterocycloalkyl ring, a 6-, 7-, or 8-membered bicyclic heterocycloalkyl ring, or a 5- or 6-membered heteroaryl ring.

[0022] In some cases, the compound of formula III has a structure IIIa:



wherein R^{11} is selected from the group consisting of H and C_{1-5} alkyl.

[0023] In some cases, A is substituted with one to four groups selected from halo, C_{1-5} alkyl, C_{3-5} cycloalkyl, and $-\text{CO}_2(\text{C}_{1-5}\text{alkyl})$.

[0024] In some cases, the compound of formula II, formula III, or disulfide thereof depletes cystine in a subject in an amount that is at least 70% of the level of depletion of cystine by cysteamine.

[0025] In various embodiments, the disclosure provides a compound as disclosed herein (e.g., a compound as represented by formula I, formula II, or formula III, or a disulfide thereof), wherein the compound or disulfide thereof produces reduced levels of dimethyl sulfide when administered to a subject compared to the level of dimethyl sulfide produced when cysteamine is administered to a subject. In some cases, least 2-fold less dimethyl sulfide is produced when the compound of formula I, formula II, or formula III, or disulfide thereof is administered to a subject.

[0026] In various embodiments, the disclosure provides a compound as disclosed herein (e.g., a compound as represented by formula I, formula II, or formula III, or a disulfide thereof), wherein the compound or disulfide thereof inhibits glutamate-induced excitotoxicity (i.e., provides neuroprotection). In some cases, the compound of formula I, formula II, or formula III, or disulfide thereof demonstrates at least 50% cell survival (expressed as a percent of the cell survival for 100 μM cysteamine), under conditions as described herein.

[0027] In various embodiments, the disclosure provides a method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound as disclosed herein (e.g., a compound as represented by formula I, formula II, or formula III, or a disulfide thereof), wherein the compound or disulfide thereof is resistant to metabolism by cysteamine dioxygenase (ADO). In some cases, less than 20% of the compound of formula I, formula II, or formula III, or disulfide thereof is metabolized by ADO when assayed by consumption of oxygen using an oxygen sensitive fluorescent probe.

[0028] Diseases for which treatment with cysteamine is indicated include, but are not limited to, cystinosis, fatty liver disease, fibrosis, a thrombotic disease, an MECP-2 related

disorder, an inherited mitochondrial disease, a neurological disease or disorder, inflammation and cancer.

[0029] Fatty liver diseases include, but are not limited to, non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), fatty liver disease resulting from hepatitis, fatty liver disease resulting from obesity, fatty liver disease resulting from diabetes, fatty liver disease resulting from insulin resistance, fatty liver disease resulting from hypertriglyceridemia, Abetalipoproteinemia, glycogen storage diseases, Weber-Christian disease, Wolmans disease, acute fatty liver of pregnancy, and lipodystrophy.

[0030] Fibrosis includes, but is not limited to, atherosclerosis, asthma, cardiac fibrosis, organ transplant fibrosis, colloid and hypertrophic scar, muscle fibrosis, pancreatic fibrosis, bone-marrow fibrosis, interstitial liver fibrosis, cirrhosis of liver and gallbladder, scleroderma, pulmonary fibrosis, diffuse parenchymal lung disease, idiopathic interstitial fibrosis, interstitial pneumonitis, desquamative interstitial pneumonia, respiratory bronchiolitis, interstitial lung disease, acute interstitial pneumonitis, nonspecific interstitial pneumonia, cryptogenic organizing pneumonia, lymphocytic interstitial pneumonia, renal fibrosis, and chronic kidney disease, cystic fibrosis and Alport's disease.

[0031] Thrombotic diseases include, but are not limited to, sickle cell disease, deep vein thrombosis, pulmonary embolism, cardiac embolism, hypercoagulable state, thrombophilia, Factor V Leiden, Antithrombin III deficiency, Protein C deficiency, Protein S deficiency, Prothrombin gene mutation (G20210A), Hyperhomocysteinemia, antiphospholipid antibody syndrome (APS), anticardiolipin antibody (ACLA) thrombosis syndrome, or lupus anticoagulant (LA) syndrome.

[0032] Neurological diseases or disorders include, but are not limited to, Huntington's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease spinal muscle atrophy, concussion, stroke, and traumatic brain injury (CTE).

[0033] MECP-2 related diseases include, but are not limited to, Rett syndrome, autism, pervasive development disorder, non-syndromic mental retardation, idiopathic neonatal encephalopathy and idiopathic cerebral palsy.

[0034] Inherited mitochondrial diseases include, but are not limited to, Friedreich's ataxia, Leber's hereditary optic neuropathy (LHON), myoclonic epilepsy and ragged-red fibers, mitochondrial encephalomyopathy, lactic acidosis, and stroke-like syndrome (MELAS), Kearns-Sayre syndrome and subacute necrotizing encephalopathy (Leigh's Syndrome).

[0035] Cancers include, but are not limited to, breast cancer, melanoma, prostate cancer, pancreatic cancer, head and neck cancer, lung cancer, non small-cell lung carcinoma, renal cancer, colorectal cancer, colon cancer, ovarian cancer, liver cancer and gastric cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

[0036] FIG. 1 is a graph showing the relationship between the sulfur-nitrogen distance of a compound as disclosed herein compared to its level of cystine depletion.

DETAILED DESCRIPTION

Definitions

[0037] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a derivative” includes a plurality of such derivatives and reference to “a patient” includes reference to one or more patients and so forth.

[0038] Also, the use of “or” means “and/or” unless stated otherwise. Similarly, “comprise,” “comprises,” “comprising” “include,” “includes,” and “including” are interchangeable and not intended to be limiting.

[0039] It is to be further understood that where descriptions of various embodiments use the term “comprising,” those skilled in the art would understand that in some specific instances, an embodiment can be alternatively described using language “consisting essentially of” or “consisting of.”

[0040] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this disclosure belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice of the disclosed methods and products, the exemplary methods, devices and materials are described herein.

[0041] The documents discussed above and throughout the text are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior disclosure. Each document is incorporated by reference in its entirety with particular attention to the disclosure for which it is cited.

[0042] The following references provide one of skill with a general definition of many of the terms used in this disclosure: Singleton, *et al.*, DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY (2d ed. 1994); THE CAMBRIDGE DICTIONARY OF

SCIENCE AND TECHNOLOGY (Walker ed., 1988); THE GLOSSARY OF GENETICS, 5TH ED., R. Rieger, *et al.* (eds.), Springer Verlag (1991); and Hale and Marham, THE HARPER COLLINS DICTIONARY OF BIOLOGY (1991).

[0043] As used herein, a “therapeutically effective amount” or “effective amount” refers to that amount of the compound sufficient to result in amelioration of symptoms, for example, treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions, typically providing a statistically significant improvement in the treated patient population. When referencing an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When referring to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, including serially or simultaneously. In some embodiments, such as for fatty liver disease, a therapeutically effective amount of the compound ameliorates one or more symptoms, including but not limited to, liver fibrosis, fat content of liver, incidence of or progression of cirrhosis, incidence of hepatocellular carcinoma, increased hepatic aminotransferase levels, such as ALT and AST, increased serum ferritin, elevated levels of gamma-glutamyltransferase (gamma-GT), and elevated levels of plasma insulin, cholesterol and triglyceride. In some embodiments, such as for a neurodegenerative disease, a therapeutically effective amount of the compound increases the level of brain-derived neurotrophic factor (BDNF). In some embodiments, such as for a neurodegenerative disease, a therapeutically effective amount of the compound inhibits tissue transglutaminase. In some embodiments, such as for a neurodegenerative disease, a therapeutically effective amount of the compound increases heat shock DnaJ-containing protein 1b.

[0044] As used herein “a disease for which treatment with cysteamine is indicated” refers to a disease in which increasing the level of cysteamine and/or reducing the level of cystine in a patient is beneficial. Contemplated diseases, include, but are not limited to any of the diseases listed herein, including those in the “Indications, Dosing and Administration” section.

[0045] “Treatment” refers to prophylactic treatment or therapeutic treatment. In certain embodiments, “treatment” refers to administration of a compound or composition to a subject for therapeutic or prophylactic purposes.

[0046] A “therapeutic” treatment is a treatment administered to a subject who exhibits signs or symptoms of pathology for the purpose of diminishing or eliminating those signs or symptoms. The signs or symptoms may be biochemical, cellular, histological, functional or physical, subjective or objective.

[0047] A “prophylactic” treatment is a treatment administered to a subject who does not exhibit signs of a disease or exhibits only early signs of the disease, for the purpose of decreasing the risk of developing pathology. The compounds or compositions of the disclosure may be given as a prophylactic treatment to reduce the likelihood of developing a pathology or to minimize the severity of the pathology, if developed.

[0048] “Diagnostic” means identifying the presence, extent and/or nature of a pathologic condition. Diagnostic methods differ in their specificity and selectivity. While a particular diagnostic method may not provide a definitive diagnosis of a condition, it suffices if the method provides a positive indication that aids in diagnosis.

[0049] “Pharmaceutical composition” refers to a composition suitable for pharmaceutical use in subject animal, including humans and mammals. A pharmaceutical composition comprises a therapeutically effective amount of a compound of the disclosure, optionally another biologically active agent, and optionally a pharmaceutically acceptable excipient, carrier or diluent. In an embodiment, a pharmaceutical composition encompasses a composition comprising the active ingredient(s), and the inert ingredient(s) that make up the carrier, as well as any product that results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present disclosure encompass any composition made by admixing a compound of the disclosure and a pharmaceutically acceptable excipient, carrier or diluent.

[0050] “Pharmaceutically acceptable carrier” refers to any of the standard pharmaceutical carriers, buffers, and the like, such as a phosphate buffered saline solution, 5% aqueous solution of dextrose, and emulsions (e.g., an oil/water or water/oil emulsion). Non-limiting examples of excipients include adjuvants, binders, fillers, diluents, disintegrants, emulsifying agents, wetting agents, lubricants, glidants, sweetening agents, flavoring agents, and coloring agents. Suitable pharmaceutical carriers, excipients and diluents are described in Remington's Pharmaceutical Sciences, 19th Ed. (Mack Publishing Co., Easton, 1995).

Preferred pharmaceutical carriers depend upon the intended mode of administration of the active agent. Typical modes of administration include enteral (*e.g.*, oral) or parenteral (*e.g.*, subcutaneous, intramuscular, intravenous or intraperitoneal injection; or topical, transdermal, or transmucosal administration).


[0051] A “pharmaceutically acceptable salt” is a salt that can be formulated into a compound for pharmaceutical use, including but not limited to metal salts (*e.g.*, sodium, potassium, magnesium, calcium, *etc.*) and salts of ammonia or organic amines.

[0052] As used herein “pharmaceutically acceptable” or “pharmacologically acceptable” is meant a material that is not biologically or otherwise undesirable, *i.e.*, the material may be administered to an individual without causing any undesirable biological effects or without interacting in a deleterious manner with any of the components of the composition in which it is contained or with any components present on or in the body of the individual.

[0053] As used herein, the term “unit dosage form” refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of a compound of the disclosure calculated in an amount sufficient to produce the desired effect, optionally in association with a pharmaceutically acceptable excipient, diluent, carrier or vehicle. The specifications for the novel unit dosage forms of the present disclosure depend on the particular compound employed and the effect to be achieved, and the pharmacodynamics associated with each compound in the host.

[0054] As used herein, the term “subject” encompasses mammals. Examples of mammals include, but are not limited to, any member of the mammalian class: humans, non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. The term does not denote a particular age or gender. In various embodiments the subject is human. In various embodiments, the subject is a child or adolescent.

[0055] In one aspect, a method is provided for treating a patient suffering from a disease for which treatment with cysteamine is indicated. The method comprises administering to the patient an effective amount of a composition comprising a compound of formula I or a disulfide thereof:



wherein:

R^1 and R^2 are independently selected from the group consisting of H and C_{1-5} alkyl; or R^1 and R^2 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^3 and R^4 are independently selected from the group consisting of H and C_{1-5} alkyl; or R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

G is selected from the group consisting of $-NR^5R^6$ and $-CR^7R^8NR^5R^6$;

R^5 and R^6 are independently selected from the group consisting of H and C_{1-5} alkyl; or R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

R^7 and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl; or R^7 and R^8 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^2 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;

R^4 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;

R^2 and R^8 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring; or

R^2 and R^4 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring.

[0056] In some cases, when G is $-NH_2$, at least one of R^1 , R^2 , R^3 , and R^4 is other than H.

[0057] In some cases, R^5 and R^6 are independently selected from the group consisting of H, methyl, and ethyl. In some cases, R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 5-membered heterocyclic ring.

[0058] In some cases, wherein R^4 is methyl and/or R^3 is methyl. In some cases, R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.


[0059] In some cases, R² is methyl and/or R¹ is methyl. In some cases, R¹ and R², taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.

[0060] In some cases, G is $\text{—CR}^7\text{R}^8\text{NR}^5\text{R}^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 6-membered heterocyclic ring. In some cases, R^5 is methyl.


[0061] In some cases, G is $\text{—NR}^5\text{R}^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 4- or 6-membered heterocyclic ring. In some cases, R^5 is H.

[0062] In some cases, R^7 and R^8 are both H.

[0063] A compound of formula I includes, but is not limited to, the following compounds:




[0064] A compound of formula I includes, but is not limited to, the following compounds:



[0065] In some cases, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and R⁸ are independently selected from the group consisting of H and C₁₋₅alkyl. In some cases, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and R⁸ are independently selected from the group consisting of H and methyl.

[0066] In one aspect, a method is provided for treating a patient suffering from a disease for which treatment with cysteamine is indicated. The method comprises administering to the patient an effective amount of a composition comprising a compound of formula II, formula III, or a disulfide thereof:



wherein:

L is a hydrocarbon linking group;

R^9 and R^{10} are independently selected from the group consisting of H, C_{1-5} alkyl, and $\text{CO}(\text{C}_{1-5}\text{alkyl})$; or

R^9 and R^{10} , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

A is a heterocyclic ring contain one N atom; and

n is 0, 1, 2, or 3.


[0067] In some cases, the compound of formula II is not cysteamine.

[0068] In some cases, the S atom in the compound of formula II or formula III is a distance of about 3.6 Angstroms to about 4.7 Angstroms from the N atom in the compound, such as about 3.8 Angstroms to about 4.4 Angstroms, about 4.0 Angstroms to about 4.2 Angstroms, or about 4.1 Angstroms from the N atom in the compound.

[0069] In some cases, L is a 3-, 4-, 5-, 6-, 7-, or 8-membered cycloalkyl ring or a 6-membered aryl ring. In some cases, L is C_{1-5} alkyl. In some cases, L is substituted with one to four groups selected from halo, C_{1-5} alkyl, C_{3-5} cycloalkyl, and $\text{---CO}_2(\text{C}_{1-5}\text{alkyl})$.

[0070] In some cases, A is a 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic heterocycloalkyl ring, a 6-, 7-, or 8-membered bicyclic heterocycloalkyl ring, or a 5- or 6-membered heteroaryl ring.


[0071] In some cases, the compound of formula III has a structure IIIa:




wherein R^{11} is selected from the group consisting of H and C_{1-5} alkyl.

[0072] In some cases, A is substituted with one to four groups selected from halo, C_{1-5} alkyl, C_{3-5} cycloalkyl, and $\text{---CO}_2(\text{C}_{1-5}\text{alkyl})$.


C1CNCCCCC1S






the tables herein, and disulfides thereof.

[0074] Compounds disclosed herein include compounds having the following structure or a disulfide thereof:



wherein R^1 is C_{1-5} alkyl; and R^3 , R^4 , R^5 , R^7 , and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl.

[0075] Compounds disclosed herein include compounds having the following structure or a disulfide thereof:



wherein R^1 , R^3 , R^4 , R^5 , R^6 , and R^7 are independently selected from the group consisting of H and C_{1-5} alkyl.

ADO-Resistant Cysteamine Analogs

[0076] The disclosure provides ADO-resistant cysteamine analogs for use in the methods described herein. An “ADO-resistant cysteamine analog” in the present disclosure refers generally to compounds of formula I, formula II, formula III, or a disulfide thereof. ADO-resistant cysteamine analogs generally demonstrate three properties: (1) the compounds are

resistant to metabolism by ADO, (2) the compounds cleave cystine in vivo, and (3) the compounds clear stored cystine in patient cystinotic fibroblasts.

[0077] As used herein a compound that is “resistant to metabolism by cysteamine dioxygenase” or “resistant to metabolism by ADO” refers to a compound that undergoes less than 50% degradation, for example, less than 40%, less than 30%, less than 25%, less than 20%, less than 15%, less than 10%, less than 8%, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, less than 1.5%, and/or less than 1% degradation when assayed in the presence of ADO under conditions as described herein. Due to the rapid metabolism and clearance of cysteamine from the body due to ADO, the sustained concentrations of cysteamine necessary for therapeutic effect are difficult to maintain. Advantageously, compounds that are resistant to metabolism by ADO are more readily maintained at necessary concentrations for therapeutic effect.

[0078] A compound that cleaves cystine in vivo refers to a compound that converts cystine to cysteine and a mixed disulfide containing cysteine and the compound. Such compounds typically have a reactivity similar to or greater than the reactivity of cysteamine for cystine, such as at least at least 50%, at least 75%, at least 90%, at least 100%, at least 110%, at least 120%, at least 130%, at least 140%, at least 150%, at least 160%, at least 170%, at least 180%, at least 190%, and/or at least 200% of the reactivity of cysteamine for cystine, as determined under conditions as described herein.

[0079] A compound that clears stored cystine in patient cystinotic fibroblasts refers to a compound that facilitates transport of cystine out of lysosomes. Such compounds typically deplete cystine in an amount similar to or greater than the depletion of cystine by cysteamine, such as at least 25%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 100%, at least 105%, and/or at least 110% of the depletion by cysteamine, as determined under conditions as described herein.

[0080] The ADO-resistant cysteamine analogs also include biologically active metabolites or derivatives thereof, and includes salts, esters, amides, alkylate compounds, prodrugs, analogs, phosphorylated compounds, sulfated compounds, or other chemically modified forms thereof (e.g., chemically modified forms prepared by labeling with radionucleotides or enzymes and chemically modified forms prepared by attachment of polymers such as polyethylene glycol). Thus, the compounds of formula I, formula II, or formula III can be

administered in the form of a pharmacologically acceptable salt, ester, amide, prodrug or analog or as a combination thereof.

[0081] Salts, esters, amides, prodrugs and analogs of the active agents may be prepared using standard procedures known to those skilled in the art of synthetic organic chemistry and described, for example, by J. March, "Advanced Organic Chemistry: Reactions, Mechanisms and Structure," 4th Ed. (New York: Wiley-Interscience, 1992). For example, basic addition salts are prepared from the neutral drug using conventional means, involving reaction of one or more of the active agent's free hydroxyl groups with a suitable base. Generally, the neutral form of the drug is dissolved in a polar organic solvent such as methanol or ethanol and the base is added thereto. The resulting salt either precipitates or may be brought out of solution by addition of a less polar solvent. Suitable bases for forming basic addition salts include, but are not limited to, inorganic bases such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine, or the like. Preparation of esters involves functionalization of hydroxyl groups which may be present within the molecular structure of the drug. The esters are typically acyl-substituted derivatives of free alcohol groups, i.e., moieties which are derived from carboxylic acids of the formula $R-COOH$ where R is alkyl, and typically is lower alkyl. Esters can be reconverted to the free acids, if desired, by using conventional hydrogenolysis or hydrolysis procedures. Preparation of amides and prodrugs can be carried out in an analogous manner. Other derivatives and analogs of the active agents may be prepared using standard techniques known to those skilled in the art of synthetic organic chemistry, or may be deduced by reference to the pertinent literature.

Dimethyl Sulfide (DMS) Production

[0082] Compounds that produce reduced levels of dimethyl sulfide when administered to a subject (compared to the level produced when cysteamine is administered) are desirable because unpleasant side effects associated with cysteamine (e.g., halitosis) may be reduced. A compound that produces reduced levels of dimethyl sulfide when administered to a subject generally produces at least 2-fold less dimethyl sulfide, such as at least 3-fold less, 4-fold less, 5-fold less, 6-fold less, 8-fold less, 10-fold less, 15-fold less, and/or 20-fold less dimethyl sulfide, compared to the level of dimethyl sulfide produced when cysteamine is administered to the subject at the same dose and same time after administration under conditions as described herein. For example, the level of dimethyl sulfide can be measured by administering the compound in a dose of about 10 mg/kg to about 500 mg/kg, such as about 25 mg/kg to about 400 mg/kg, about 50 mg/kg to about 300 mg/kg, about 75 mg/kg to

about 200 mg/kg, about 10 mg/kg, about 25 mg/kg, about 50 mg/kg, about 75 mg/kg, about 100 mg/kg, about 150 mg/kg, about 200 mg/kg, about 300 mg/kg, about 400 mg/kg, and/or about 500 mg/kg, and by measuring the dimethyl sulfide level about 10 minutes to 4 hours after administration, such as about 15 minutes to about 2 hours, about 30 minutes to about 1 hour, about 15 minutes, about 30 minutes, about 1 hour, and/or about 2 hours after administration.

Neuroprotection

[0083] Excitotoxicity disorders affect the central nervous and peripheral nervous systems and can lead to progressive neurodegeneration. Excitotoxicity results from excess glutamate being secreted by various cells, including immune cells and neurons, in the brain. Glutamate is the primary excitatory neurotransmitter in the mammalian nervous system. Prolonged glutamate signaling leads to a type of toxicity characterized by elevated mitochondrial activity, gradual glutathione (GSH) depletion, oxidative stress and apoptosis (Shih et al., J Neurosci. 26:10514-523, 2006). Cysteamine is capable of inhibiting glutamate-induced excitotoxicity in St-HdhQ^{111/111} cells. A compound that inhibits glutamate-induced excitotoxicity (i.e., provides neuroprotection) generally provides at least 50% cell survival (expressed as a percent of the cell survival for 100 μ M cysteamine), such as at least 65%, at least 75%, at least 80%, at least 90%, and/or at least 95% cell survival, under conditions as described herein.

Pharmaceutical Formulations

[0084] The disclosure provides compounds useful in the treatment of diseases in which therapy with cysteamine is indicated. To administer compounds of the disclosure to patients or test animals, it is preferable to formulate the compounds in a composition comprising one or more pharmaceutically acceptable carriers. Pharmaceutically or pharmacologically acceptable carriers or vehicles refer to molecular entities and compositions that do not produce allergic, or other adverse reactions when administered using routes well-known in the art, as described below, or are approved by the U.S. Food and Drug Administration or a counterpart foreign regulatory authority as an acceptable additive to orally or parenterally administered pharmaceuticals. Pharmaceutically acceptable carriers include any and all clinically useful solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like.

[0085] Pharmaceutical carriers include pharmaceutically acceptable salts, particularly where a basic or acidic group is present in a compound. For example, when an acidic

substituent, such as --COOH, is present, the ammonium, sodium, potassium, calcium and the like salts, are contemplated for administration. Additionally, where an acid group is present, pharmaceutically acceptable esters of the compound (*e.g.*, methyl, tert-butyl, pivaloyloxymethyl, succinyl, and the like) are contemplated as preferred forms of the compounds, such esters being known in the art for modifying solubility and/or hydrolysis characteristics for use as sustained release or prodrug formulations.

[0086] When a basic group (such as amino or a basic heteroaryl radical, such as pyridyl) is present, then an acidic salt, such as hydrochloride, hydrobromide, acetate, maleate, pamoate, phosphate, methanesulfonate, p-toluenesulfonate, and the like, is contemplated as a form for administration.

[0087] In addition, compounds may form solvates with water or common organic solvents. Such solvates are contemplated as well.

[0088] The compounds may be administered orally, parenterally, transocularly, intranasally, transdermally, transmucosally, by inhalation spray, vaginally, rectally, or by intracranial injection. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intracisternal injection, or infusion techniques. Administration by intravenous, intradermal, intramuscular, intramammary, intraperitoneal, intrathecal, retrobulbar, intrapulmonary injection and or surgical implantation at a particular site is contemplated as well. Generally, compositions for administration by any of the above methods are essentially free of pyrogens, as well as other impurities that could be harmful to the recipient. Further, compositions for administration parenterally are sterile.

[0089] Pharmaceutical compositions of the disclosure containing a compound as disclosed herein as an active ingredient may contain pharmaceutically acceptable carriers or additives depending on the route of administration. Examples of such carriers or additives include water, a pharmaceutically acceptable organic solvent, collagen, polyvinyl alcohol, polyvinylpyrrolidone, a carboxyvinyl polymer, carboxymethylcellulose sodium, polyacrylic sodium, sodium alginate, water-soluble dextran, carboxymethyl starch sodium, pectin, methyl cellulose, ethyl cellulose, xanthan gum, gum Arabic, casein, gelatin, agar, diglycerin, glycerin, propylene glycol, polyethylene glycol, Vaseline, paraffin, stearyl alcohol, stearic acid, human serum albumin (HSA), mannitol, sorbitol, lactose, a pharmaceutically acceptable surfactant and the like. Additives used are chosen from, but not limited to, the above or combinations thereof, as appropriate, depending on the dosage form of the disclosure.

[0090] Formulation of the pharmaceutical composition will vary according to the route of administration selected (*e.g.*, solution, emulsion). An appropriate composition comprising

the compound to be administered can be prepared in a physiologically acceptable vehicle or carrier. For solutions or emulsions, suitable carriers include, for example, aqueous or alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles can include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles can include various additives, preservatives, or fluid, nutrient or electrolyte replenishers.

[0091] A variety of aqueous carriers, *e.g.*, water, buffered water, 0.4% saline, 0.3% glycine, or aqueous suspensions may contain the active compound in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0092] In some embodiments, the compounds of this disclosure can be lyophilized for storage and reconstituted in a suitable carrier prior to use. Any suitable lyophilization and reconstitution techniques can be employed. It is appreciated by those skilled in the art that lyophilization and reconstitution can lead to varying degrees of activity loss and that use levels may have to be adjusted to compensate.

[0093] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active compound in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[0094] In one embodiment, the disclosure provides use of an enterically coated composition. Enteric coatings prolong release until the active agent reaches the intestinal

tract, typically the small intestine. Because of the enteric coatings, delivery to the small intestine is improved thereby improving uptake of the active ingredient while reducing gastric side effects. Exemplary enterically coated products are described in International Publication No. WO 2007/089670 published August 9, 2007, which is incorporated in its entirety herein.

[0095] In some embodiments, the coating material is selected such that the therapeutically active agent is released when the dosage form reaches the small intestine or a region in which the pH is greater than pH 4.5. The coating may be a pH-sensitive materials, which remain intact in the lower pH environs of the stomach, but which disintegrate or dissolve at the pH commonly found in the small intestine of the patient. For example, the enteric coating material begins to dissolve in an aqueous solution at pH between about 4.5 to about 5.5 or between about 5.5 to 6.5. For example, pH-sensitive materials will not undergo significant dissolution until the dosage form has emptied from the stomach. The pH of the small intestine gradually increases from about 4.5 to about 6.5 in the duodenal bulb to about 7.2 in the distal portions of the small intestine. In order to provide predictable dissolution corresponding to the small intestine transit time of about 3 hours (*e.g.*, 2-3 hours) and permit reproducible release therein, the coating should begin to dissolve at the pH range within the small intestine. Therefore, the amount of enteric polymer coating should be sufficient to substantially dissolve during the approximate three hour transit time within the small intestine, such as the proximal and mid-intestine.

[0096] Enteric coatings have been used for many years to arrest the release of the drug from orally ingestible dosage forms. Depending upon the composition and/or thickness, the enteric coatings are resistant to stomach acid for required periods of time before they begin to disintegrate and permit release of the drug in the lower stomach or upper part of the small intestines. Examples of some enteric coatings are disclosed in U.S. Pat. No. 5,225,202 which is incorporated by reference fully herein. As set forth in U.S. Pat. No. 5,225,202, some examples of coating previously employed are beeswax and glyceryl monostearate; beeswax, shellac and cellulose; and cetyl alcohol, mastic and shellac, as well as shellac and stearic acid (U.S. Pat. No. 2,809,918); polyvinyl acetate and ethyl cellulose (U.S. Pat. No. 3,835,221); and neutral copolymer of polymethacrylic acid esters (Eudragit L30D) (F. W. Goodhart *et al.*, *Pharm. Tech.*, pp. 64-71, April 1984); copolymers of methacrylic acid and methacrylic acid methylester (Eudragits), or a neutral copolymer of polymethacrylic acid esters containing metallic stearates (Mehta *et al.*, U.S. Pat. Nos. 4,728,512 and 4,794,001). Such coatings comprise mixtures of fats and fatty acids, shellac and shellac derivatives and the cellulose acid phthlates, *e.g.*, those having a free carboxyl content. See, Remington's at page 1590, and

Zeitova *et al.* (U.S. Pat. No. 4,432,966), for descriptions of suitable enteric coating compositions. Accordingly, increased adsorption in the small intestine due to enteric coatings of product compositions can result in improved efficacy.

[0097] Generally, the enteric coating comprises a polymeric material that prevents product release in the low pH environment of the stomach but that ionizes at a slightly higher pH, typically a pH of 4 or 5, and thus dissolves sufficiently in the small intestines to gradually release the active agent therein. Accordingly, among the most effective enteric coating materials are polyacids having a pKa in the range of about 3 to 5. Suitable enteric coating materials include, but are not limited to, polymerized gelatin, shellac, methacrylic acid copolymer type CNF, cellulose butyrate phthalate, cellulose hydrogen phthalate, cellulose propionate phthalate, polyvinyl acetate phthalate (PVAP), cellulose acetate phthalate (CAP), cellulose acetate trimellitate (CAT), hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate, dioxypopyl methylcellulose succinate, carboxymethyl ethylcellulose (CMEC), hydroxypropyl methylcellulose acetate succinate (HPMCAS), and acrylic acid polymers and copolymers, typically formed from methyl acrylate, ethyl acrylate, methyl methacrylate and/or ethyl methacrylate with copolymers of acrylic and methacrylic acid esters (Eudragit NE, Eudragit RL, Eudragit RS). In one embodiment, the product composition is administered in oral delivery vehicle, including but not limited to, tablet or capsule form. Tablets are manufactured by first enterically coating the product. A method for forming tablets herein is by direct compression of the powders containing the enterically coated product, optionally in combination with diluents, binders, lubricants, disintegrants, colorants, stabilizers or the like. As an alternative to direct compression, compressed tablets can be prepared using wet-granulation or dry-granulation processes. Tablets may also be molded rather than compressed, starting with a moist material containing a suitable water-soluble lubricant.

[0098] In a further embodiment, the product is formulated as a capsule. In one embodiment, the capsule comprises the product and the capsule is then enterically coated. Capsule formulations are prepared using techniques known in the art.

[0099] The preparation of delayed, controlled or sustained/extended release forms of pharmaceutical compositions with the desired pharmacokinetic characteristics is known in the art and can be accomplished by a variety of methods. For example, oral controlled delivery systems include dissolution-controlled release (*e.g.*, encapsulation dissolution control or matrix dissolution control), diffusion-controlled release (reservoir devices or matrix devices), ion exchange resins, osmotic controlled release or gastroretentive systems. Dissolution

controlled release can be obtained, *e.g.*, by slowing the dissolution rate of a drug in the gastrointestinal tract, incorporating the drug in an insoluble polymer, and coating drug particles or granules with polymeric materials of varying thickness. Diffusion controlled release can be obtained, *e.g.*, by controlling diffusion through a polymeric membrane or a polymeric matrix. Osmotically controlled release can be obtained, *e.g.*, by controlling solvent influx across a semipermeable membrane, which in turn carries the drug outside through a laser-drilled orifice. The osmotic and hydrostatic pressure differences on either side of the membrane govern fluid transport. Prolonged gastric retention may be achieved by, *e.g.*, altering density of the formulations, bioadhesion to the stomach lining, or increasing floating time in the stomach. For further detail, see the Handbook of Pharmaceutical Controlled Release Technology, Wise, ed., Marcel Dekker, Inc., New York, NY (2000), incorporated by reference herein in its entirety, *e.g.* Chapter 22 (“An Overview of Controlled Release Systems”).

[0100] The concentration of product in these formulations can vary widely, for example from less than about 0.5%, usually at or at least about 1% to as much as 15 or 20% by weight and are selected primarily based on fluid volumes, manufacturing characteristics, viscosities, etc., in accordance with the particular mode of administration selected. Actual methods for preparing administrable compositions are known or apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Science, 15th ed., Mack Publishing Company, Easton, Pa. (1980).

[0101] Compositions useful for administration may be formulated with uptake or absorption enhancers to increase their efficacy. Such enhancers include, for example, salicylate, glycocholate/linoleate, glycholate, aprotinin, bacitracin, SDS, caprate and the like. See, *e.g.*, Fix (J. Pharm. Sci., 85:1282-1285, 1996) and Oliyai and Stella (Ann. Rev. Pharmacol. Toxicol., 32:521-544, 1993).

[0102] The enterically coated product can comprise various excipients, as is well known in the pharmaceutical art, provided such excipients do not exhibit a destabilizing effect on any components in the composition. Thus, excipients such as binders, bulking agents, diluents, disintegrants, lubricants, fillers, carriers, and the like can be combined with the cysteamine product. Oral delivery vehicles contemplated for use herein include tablets or capsules. For solid compositions, diluents are typically necessary to increase the bulk of a tablet or capsule so that a practical size is provided for compression. Suitable diluents include dicalcium phosphate, calcium sulfate, lactose, cellulose, kaolin, mannitol, sodium chloride, dry starch

and powdered sugar. Binders are used to impart cohesive qualities to a oral delivery vehicle formulation, and thus ensure that a tablet remains intact after compression. Suitable binder materials include, but are not limited to, starch (including corn starch and pregelatinized starch), gelatin, sugars (including sucrose, glucose, dextrose and lactose), polyethylene glycol, waxes, and natural and synthetic gums, e.g., acacia sodium alginate, polyvinylpyrrolidone, cellulosic polymers (including hydroxypropyl cellulose, hydroxypropyl methylcellulose, methyl cellulose, hydroxyethyl cellulose, hypromellose, and the like), and Veegum. Lubricants are used to facilitate oral delivery vehicle manufacture; examples of suitable lubricants include, for example, magnesium stearate, calcium stearate, and stearic acid, and are typically present at no more than approximately 1 weight percent relative to tablet weight. Disintegrants are used to facilitate oral delivery vehicle, (e.g., a tablet) disintegration or "breakup" after administration, and are generally starches, clays, celluloses, algin, gums or crosslinked polymers. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and the like. If desired, flavoring, coloring and/or sweetening agents may be added as well. Other optional components for incorporation into an oral formulation herein include, but are not limited to, preservatives, suspending agents, thickening agents, and the like. Fillers include, for example, insoluble materials such as silicon dioxide, titanium oxide, alumina, talc, kaolin, powdered cellulose, microcrystalline cellulose, and the like, as well as soluble materials such as mannitol, urea, sucrose, lactose, dextrose, sodium chloride, sorbitol, and the like.

[0103] A pharmaceutical composition may also comprise a stabilizing agent such as hydroxypropyl methylcellulose or polyvinylpyrrolidone, as disclosed in U.S. Pat. No. 4,301,146. Other stabilizing agents include, but are not limited to, cellulosic polymers such as hydroxypropyl cellulose, hydroxyethyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose phthalate, microcrystalline cellulose and carboxymethylcellulose sodium; and vinyl polymers and copolymers such as polyvinyl acetate, polyvinylacetate phthalate, vinylacetate crotonic acid copolymer, and ethylene-vinyl acetate copolymers. The stabilizing agent is present in an amount effective to provide the desired stabilizing effect; generally, this means that the ratio of cysteamine product to the stabilizing agent is at least about 1:500 w/w, more commonly about 1:99 w/w.

[0104] The tablet, capsule or other oral delivery system is manufactured by first enterically coating the product. A method for forming tablets herein is by direct compression of the powders containing the enterically coated cysteamine product, optionally in combination with diluents, binders, lubricants, disintegrants, colorants, stabilizers or the like. As an alternative to direct compression, compressed tablets can be prepared using wet-granulation or dry-granulation processes. Tablets may also be molded rather than compressed, starting with a moist material containing a suitable water-soluble lubricant.

[0105] In an alternative embodiment, the enterically coated product is granulated and the granulation is compressed into a tablet or filled into a capsule. Capsule materials may be either hard or soft, and are typically sealed, such as with gelatin bands or the like. Tablets and capsules for oral use will generally include one or more commonly used excipients as discussed herein.

[0106] For administration of the dosage form, i.e., the tablet or capsule comprising the enterically coated product, a total weight in the range of approximately 100 mg to 1000 mg is used. The dosage form is orally administered to a patient suffering from a condition for which treatment with cysteamine would typically be indicated.

Indications, Dosing and Administration

[0107] The disclosure provides methods of treating a patient (e.g., a human patient) suffering from a disease where therapy with cysteamine is indicated, comprising administering to the patient a therapeutically effective amount of a compound as disclosed herein.

[0108] In some embodiments of the methods, the disease is cystinosis. In some embodiments the disease is nephropathic cystinosis. In some embodiments of the methods, the disease is a fatty liver disease. In some embodiments, the fatty liver disease is non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), fatty liver disease resulting from hepatitis, fatty liver disease resulting from obesity, fatty liver disease resulting from diabetes, fatty liver disease resulting from insulin resistance, fatty liver disease resulting from hypertriglyceridemia, Abetalipoproteinemia, glycogen storage diseases, Weber-Christian disease, Wolmans disease, acute fatty liver of pregnancy, and lipodystrophy or other fatty liver disease. The term “fatty liver disease” may include or exclude NASH. In some embodiments of the methods, the disease is a fibrosis. In some embodiments, the fibrosis is atherosclerosis, asthma, cardiac fibrosis, organ transplant fibrosis, colloid and

hypertrophic scar, muscle fibrosis, pancreatic fibrosis, bone-marrow fibrosis, interstitial liver fibrosis, cirrhosis of liver and gallbladder, scleroderma, pulmonary fibrosis, diffuse parenchymal lung disease, idiopathic interstitial fibrosis, interstitial pneumonitis, desquamative interstitial pneumonia, respiratory bronchiolitis, interstitial lung disease, acute interstitial pneumonitis, nonspecific interstitial pneumonia, cryptogenic organizing pneumonia, lymphocytic interstitial pneumonia, renal fibrosis, chronic kidney disease, cystic fibrosis, or Alport's disease. In some embodiments of the methods, the disease is a thrombotic disease. In some embodiments, the thrombotic disease is sickle cell disease, deep vein thrombosis, pulmonary embolism, cardiac embolism, hypercoagulable state, thrombophilia, Factor V Leiden, Antithrombin III deficiency, Protein C deficiency, Protein S deficiency, Prothrombin gene mutation (G20210A), Hyperhomocysteinemia, antiphospholipid antibody syndrome (APS), anticardiolipin antibody (ACLA) thrombosis syndrome, or lupus anticoagulant (LA) syndrome. In some embodiments of the methods, the disease is an MECP-2 related disorder such as Rett syndrome, autism, pervasive development disorder, non-syndromic mental retardation, idiopathic neonatal encephalopathy or idiopathic cerebral palsy. In some embodiments of the methods, the disease is an inherited mitochondrial disease such as Friedreich's ataxia, Leber's hereditary optic neuropathy (LHON), myoclonic epilepsy and ragged-red fibers, mitochondrial encephalomyopathy, lactic acidosis, and stroke-like syndrome (MELAS), Kearns-Sayre syndrome or subacute necrotizing encephalopathy (Leigh's Syndrome). In some embodiments of the methods, the disease is a neurological disease or disorder such as Huntington's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease spinal muscle atrophy, concussion, stroke, or traumatic brain injury (CTE). In some embodiments of the methods, the disease is inflammation. In some embodiments of the methods, the disease is cancer, for example, breast cancer, melanoma, prostate cancer, pancreatic cancer, head and neck cancer, lung cancer, non small-cell lung carcinoma, renal cancer, colorectal cancer, colon cancer, ovarian cancer, liver cancer or gastric cancer.

[0109] As used herein, "renal fibrosis or chronic kidney disease" refers to a progressive disorder of the kidney characterized by excessive deposit(s) of extracellular matrix (ECM) and resulting in glomerular sclerosis and renal tubule-interstitium fibrosis. Excessive deposit of fibrous tissue replaces healthy kidney tissue, damaging kidney structure and impairing kidney function. Exemplary renal fibrosis or chronic kidney disease include, but are not limited to, chronic renal insufficiency (CRI), stage III, IV or V chronic kidney disease,

nephropathy, glomerulosclerosis, glomerulonephritis, diabetes, fibrocystic kidney disease, fibrotic kidney cancer, or renal interstitial fibrosis.

[0110] The compound is administered in a therapeutically effective amount. The amount of compound to be administered is dependent on the age, weight, and general condition of the patient, the severity of the condition being treated, and the judgment of the prescribing physician. Suitable therapeutic amounts are determined by standard methods by those skilled in the art. In some embodiments, the dose is administered either one or two times per day. In some embodiments, the dose is administered multiple times per day. In some embodiments, an effective dosage of compound is within the range of about 0.01 mg to about 1000 mg per kg (mg/kg) of body weight per day. Further, the effective dose may be about: 0.5 mg/kg, 1 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 35 mg/kg, 40 mg/kg, 45 mg/kg, 50 mg/kg, 55 mg/kg, 60 mg/kg, 70 mg/kg, 75 mg/kg, 80 mg/kg, 90 mg/kg, 100 mg/kg, 125 mg/kg, 150 mg/kg, 175 mg/kg, 200 mg/kg, and may increase by 25 mg/kg increments up to 1000 mg/kg, or may range between any two of the foregoing values. In some embodiments, the administration route is oral. In some embodiments, the administration route is transdermal.

[0111] The compound is administered in a therapeutically effective amount; typically, the composition is in unit dosage form. The amount of compound administered is, of course, dependent on the age, weight, and general condition of the patient, the severity of the condition being treated, and the judgment of the prescribing physician. Suitable therapeutic amounts will be known to those skilled in the art and/or are described in the pertinent reference texts and literature. In one aspect, the dose is administered either one time per day or multiple times per day. The product may be administered one, two or three or four times per day. In some embodiments, an effective dosage of product may be within the range of 0.01 mg to 1000 mg per kg (mg/kg) of body weight per day. Further, the effective dose may be 0.5 mg/kg, 1 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 35 mg/kg, 40 mg/kg, 45 mg/kg, 50 mg/kg, 55 mg/kg, 60 mg/kg, 70 mg/kg, 75 mg/kg, 80 mg/kg, 90 mg/kg, 100 mg/kg, 125 mg/kg, 150 mg/kg, 175 mg/kg, 200 mg/kg, 225 mg/kg, 250 mg/kg, 275 mg/kg, 300 mg/kg, 325 mg/kg, 350 mg/kg, 375 mg/kg, 400 mg/kg, 425 mg/kg, 450 mg/kg, 475 mg/kg, 500 mg/kg, 525 mg/kg, 550 mg/kg, 575 mg/kg, 600 mg/kg, 625 mg/kg, 650 mg/kg, 675 mg/kg, 700 mg/kg, 725 mg/kg, 750 mg/kg, 775 mg/kg, 800 mg/kg, 825 mg/kg, 850 mg/kg, 875 mg/kg, 900 mg/kg, 925 mg/kg, 950 mg/kg, 975 mg/kg or 1000 mg/kg, or may range between any two of the foregoing values. In some embodiments, the

dose above may be the total daily dose, or may be the dose administered in one of the one, two or three daily administrations. In some embodiments, the product is administered at a total daily dose of from approximately 0.25 g/m² to 4.0 g/m² body surface area, *e.g.*, at least about 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2 g/m², or up to about 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.2, 2.5, 2.7, 3.0, or 3.5 g/m² or may range between any two of the foregoing values. In some embodiments, the product may be administered at a total daily dose of about 0.5 -2.0 g/m² body surface area, or 1-1.5 g/m² body surface area, or 0.5-1 g/m² body surface area, or about 0.7-0.8 g/m² body surface area, or about 1.35 g/m² body surface area, or about 1.3 to about 1.95 grams/m²/day, or about 0.5 to about 1.5 grams/m²/day, or about 0.5 to about 1.0 grams/m²/day, preferably at a frequency of fewer than four times per day, *e.g.* three, two or one times per day. Salts or esters of the same active ingredient may vary in molecular weight depending on the type and weight of the salt or ester moiety. For administration of the dosage form, *e.g.*, a tablet or capsule or other oral dosage form comprising the enterically coated product, a total weight in the range of approximately 100 mg to 1000 mg is used. In certain embodiments, the amount of active ingredient in a tablet or capsule is approximately 15, 20, 25, 50, 75, 100, 125, 150, 175, 200, 250, 300, 400 or 500 mg. Administration may continue for at least 3 months, 6 months, 9 months, 1 year, 2 years, or more.

[0112] The compositions of the invention can be used in combination with other drugs or therapies for each indication contemplated herein. Combinations with other drugs or therapies that are part of the standard of care for each indication are specifically contemplated herein.

[0113] The compounds and other drugs/therapies can be administered in combination either simultaneously in a single composition or in separate compositions. Alternatively, the administration is sequential. In some embodiments, the patient is pre-dosed with the compound before the administration of the other drug/therapy.

[0114] The effectiveness of a method or composition of the described herein can be assessed, for example, by measuring leukocyte cystine concentrations in subjects affected by cystinosis. Additional measures of the efficacy of the methods of the disclosure include assessing relief of symptoms associated with fatty liver disease including, but not limited to, liver fibrosis, fat content of liver, incidence of or progression of cirrhosis, incidence of hepatocellular carcinoma, elevated hepatic aminotransferase levels, increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), and elevated serum

ferritin. Dosage adjustment and therapy can be made by a medical specialist depending upon, for example, the severity of fatty liver disease and/or the concentration of cystine. For example, treatment of fatty liver disease may result in a reduction in hepatic transaminase of between approximately 10% to 40% compared to levels before treatment. In a related embodiment, treatment results in a reduction in alanine aminotransferase levels in a treated patient to approximately 30%, 20% or 10% above normal ALT levels, or at normal ALT levels (≥ 40 iu/L). In another embodiment, treatment with cysteamine product results in a reduction in aspartate aminotransferase levels in a patient to approximately 30%, 20% or 10% above normal AST levels or back to normal AST levels.

[0115] The methods of the invention also include use of a compound as disclosed herein in preparation of a medicament for treatment of a disease in which administration of cysteamine is indicated, and use of a compound as disclosed herein in preparation of a medicament for administration in combination with another agent for treating a disease where administration of cysteamine is indicated. Further provided are kits comprising a compound as disclosed herein for treatment of a disease in which administration of cysteamine is indicated, optionally another agent for treatment, and instructions for use in treatment of a disease where administration of cysteamine is indicated.

Animal Models

[0116] The compounds disclosed herein can be evaluated in animal models known in the art for the disease indications contemplated herein.

[0117] For example, a number of mouse models which exhibit steatosis/steatohepatitis exist and include genetically altered leptin-deficient (ob/ob) or leptin resistant (db/db) and the dietary methionine/choline deficient (MCD) model. Studies comparing male and female rats of varying strains (Wistar, Sprague-Dawley, Long-Evans) with a mouse strain (C57BL/6) as models for NASH can be carried out. More recently the use of supra-nutritional diets in animals has resulted in a NAFLD model that physiologically more resembles the human phenotype. The medical conditions most commonly associated with NAFLD are obesity, Type II diabetes and dyslipidemia. These conditions can be induced by feeding mice and rats with high fat or sucrose diets. Rats fed with a >70% fat-rich diet for 3 weeks develop pan-lobular steatosis, patchy inflammation, enhanced oxidative stress, and increased plasma insulin concentrations suggesting insulin resistance. NASH mice have been induced through intragastric overfeeding. Mice were fed up to 85% in excess of their standard intake for 9 weeks. The mice became obese with 71% increase in final body weight; they demonstrated

increase white adipose tissue, hyperglycemia, hyperinsulinemia, hyperleptinemia, glucose intolerance and insulin resistance. Of these mice 46% developed increased ALT (121 $\pm$ 27 vs 13 $\pm$ 1 U/L) as well as histologic features suggestive of NASH. The livers of the overfed mice were about twice as large expected, beige in color with microscopic evidence of lipid droplets, cytoplasmic vacuoles and clusters of inflammation.

[0118] Mouse models of NASH are created through specific diets (methionine choline deficient, MCD) or intragastric overfeeding. These mice develop serologic and histologic features of NASH. NASH mice are useful in screening and measuring the effects the compounds disclosed herein on NASH related disease and disorders.

[0119] Animal models for kidney fibrosis are known in the art and described, for example, in Eddy et al., "Investigating mechanisms of chronic kidney disease in mouse models" *Pediatr Nephrol.* 2011 Jun 22.

[0120] Animal models of Huntington's Disease and Parkinson's disease are described in the art and useful to determine the effects of the compounds disclosed herein in subjects suffering from disease. See, e.g., Karpuj et al., "Evidence for a role for transglutaminase in Huntington's disease and the potential therapeutic implications." *Neurochem Int.* (2002) Jan;40(1):31-6, and Bove et al., "Neurotoxin-based models of Parkinson's disease." *Neuroscience.* 2011, Nov 10.

[0121] Additional animal models for other indications are available in the art and are useful to measure the efficacy of the compounds disclosed herein in said disorders.

[0122] While the disclosure has been described in conjunction with specific embodiments thereof, the foregoing description as well as the examples which follow are intended to illustrate and not limit the scope of the disclosure. Other aspects, advantages and modifications within the scope of the disclosure will be apparent to those skilled in the art.

EXAMPLES

General Methods

Cystine Reactivity Assay

[0123] Cystine reactivity was assessed by incubating 3 mM test compound with 100 nM BODIPY® FL L-Cystine (Life Tech.) at 30°C, pH 7.4 in phosphate buffer. The initial rate of fluorescence (excitation λ 485nm and emission λ 535nm) increase was used as a measure of cystine reactivity and hence cysteine formation. Compound cystine reactivity was expressed as a percentage of that of 3 mM cysteamine.

[0124] Under resting conditions, the inherent fluorescence of BODIPY® FL L-cystine was very low due to the spatial proximity of the two BODIPY molecules causing quenching. Under chemical reaction (disruption of the S-S bond), the spatial distance was increased, the quenching lost, and measured fluorescence increased. Time (minutes) vs. fluorescence (relative fluorescence units – RFU) was plotted and linear regression analysis was performed to generate a slope value corresponding to the reaction rate in RFU/min and an R2 value.

ADO Metabolism Assay

[0125] The rate of ADO metabolism was determined by incubating 15 mM test compound with 60 μ g/ml human recombinant ADO and 20 nM MitoXpress® O₂-sensitive fluorescent probe (Luxcel) at 37°C in assay buffer/salt in a sealed assay chamber. The initial rate of fluorescence (excitation λ 380nm and emission λ 650nm) increase was used to measure ADO-dependent oxygen depletion activity. The ability of ADO to metabolize compounds and concomitantly consume oxygen was expressed relative to that of 15 mM cysteamine.

Cystine Depletion Assay

[0126] The ability of test compounds to deplete cystine from cystinotic human fibroblasts (Coriell) was determined by incubating compounds with cells for 60 min at 37°C, 95% (v/v) air/5% (v/v) CO₂ in Minimal Essential Media Eagle. Treated cell samples were subsequently harvested, then homogenized to ensure plasma and lysosomal membrane disruption. Protein was removed by acid precipitation and cystine in the supernatant measured using LC-MS/MS and HILIC chromatography. Protein levels were determined and cystine levels reported as nM per mg/ml protein. Cystine levels after test compound treatment were expressed as a percentage of that of untreated control cells.

Rat Hepatocyte Assay

[0127] Rat hepatocytes were used to model in vivo hepatic clearance of the compounds by the ADO enzyme. Briefly, a Hepatocyte Stability Assay was carried out using cryopreserved rat hepatocytes. The test compound was added to cells (e.g., 10^6 cells) at a concentration of approximately 3 μ M (50 μ L of 10 mM solution). Other concentrations can also be used. Presence of compound was measured at varying timepoints (0, 5, 15, 30, 45, 60, 90, 120 minutes) by LC-MS/MS analysis and intrinsic clearance (CL_{int}) rate was measured. See, e.g., Lubberstedt et al., HepaRG human hepatic cell line utility as a surrogate for primary human hepatocytes in drug metabolism assessment in vitro, *J Pharmacol Toxicol Methods* 63:59-68, 2010; or Zanelli et al., Comparison of cryopreserved HepaRG cells with cryopreserved human hepatocytes for prediction of clearance for 26 drugs, *Drug Metab Dispos* 40:104-110, 2012.

Rodent Pharmacokinetic Assays




[0128] Pharmacokinetic parameters of a test compound in a rodent species (mouse and/or rat) were determined by administering a test compound at 2 and 10 mg free base equivalents per kg via the intravenous and oral gavage route to groups of three animals per route.

[0129] Blood samples were taken at various times after administration, plasma prepared, and submitted to analysis for parent drug using a qualified LC-MS-MS assay. Pharmacokinetics parameters derived from the plasma analytical data were determined using non-compartmental analysis.

Example 1 Activity of N-Substituted Compounds

[0130] Compounds were assayed at a concentration of 50 μ M or 100 μ M according to the general methods described above.

Table 1

Compound	Cysteamine	1a	1b	1c
Structure				
Cystine Reactivity	6543	---	7933	11027
ADO metabolism	111%	4%	9%	2%
Cystine depletion	101% (106%)	85% (94%)	104% (104%)	98%
Rat Hepatocytes	33.7	---	27.0	26.3

CL_{int} (ml/min/mg)				
Rat PK –Half life Bioavailability	0.5h 62%	---	0.3h 15%	0.3h 28%
Mouse PK – Half life Bioavailability	2.4h 33%	---	---	---






() = disulfide efficacy

[0131] As shown in Table 1, compounds 1b and 1c demonstrate increased reactivity with cystine compared to cysteamine, along with similar cystine depletion levels. Compound 1a demonstrates slightly decreased cystine depletion levels as compared to the cystine depletion levels for cysteamine. Advantageously, compounds 1a, 1b, and 1c demonstrate significantly reduced metabolism by ADO compared to cysteamine.

Example 2 Activity of Cyclic N-Substituted Compounds

[0132] Compounds were assayed at a concentration of 50 μ M or 100 μ M according to the general methods described above.

Table 2

Compound	Cysteamine	2a	2b	2c	2d
Structure					
Cystine Reactivity	6543	---	12392	---	7332
ADO metabolism	111%	7%	4%	2%	1%
Cystine depletion	101% (106%)	82%	99%	73%	79%
Rat Hepatocytes CL_{int} (ml/min/mg)	33.7	---	15.7	---	12.6
Rat PK –Half life Bioavailability	0.5h 62%	---	---	---	2.3h 34%
Mouse PK – Half life Bioavailability	2.4h 33%	---	0.4h 33%	7.1h 74%	3.9h 38%





() = disulfide efficacy

[0133] As shown in Table 2, compounds 2b and 2d demonstrate increased reactivity with cystine compared to cysteamine. The compounds also demonstrates similar (compound 2b) or slightly decreased (compounds 2a, 2c, 2d) cystine depletion levels as compared to the cystine depletion levels for cysteamine. Advantageously, compounds 2a, 2b, 2c, and 2d demonstrate significantly reduced metabolism by ADO compared to cysteamine.

Example 3 Activity of Alkyl Chain-Substituted Compounds

[0134] Compounds were assayed at a concentration of 50 μ M or 100 μ M according to the general methods described above.

Table 3

Compound	Cysteamine	3a	3b	3c	3d
Structure					
Cystine Reactivity	6543	2284	---	7733	---
ADO metabolism	111%	14%	---	9%	14%
Cystine depletion	101% (106%)	32%	97%	97%	55%
Rat Hepatocytes CL _{int} (ml/min/mg)	33.7	---	---	29.5	---
Rat PK –Half life Bioavailability	0.5h 62%	---	---	0.8h 23%	---
Mouse PK – Half life Bioavailability	2.4h 33%	---	---	---	---

() = disulfide efficacy

[0135] As shown in Table 3, compound 3c demonstrates increased reactivity with cystine compared to cysteamine. The compounds also demonstrate similar (compounds 3b, 3c) or slightly decreased (compounds 3a, 3d) cystine depletion levels as compared to the cystine depletion levels for cysteamine. Advantageously, compounds 3a, 3c, and 3d demonstrate significantly reduced metabolism by ADO compared to cysteamine.

Example 4

Sulfur-Nitrogen Distances and Activity of Compounds



[0136] The sulfur-nitrogen distances for each of the compounds in Table 4 were calculated using quantum mechanic calculations according to the following procedure. First, the 2D chemical structures were read into Spartan '14 and automatically converted to 3D models. Each 3D structure was then minimized using MMFF (Merck Molecular Force Field) (compound in neutral form). Then, to each minimized MMFF structure (neutral form), quantum mechanics calculations were run as follows: Equilibrium Geometry; Density Functional, B3LYP, 6-31G*; vacuum; in neutral form; no unpaired electrons.









[0137] Cystine depletion levels for the compounds are expressed in Table 4 as a percent relative to the cystine depletion achieved by cysteamine (c.f. cysteamine 100%) at a similar concentration.









[0138] Vortex® scatter plots (generated from Dotmatics®) were used to graphically represent the relationship between the sulfur-nitrogen distance and the extent of cystine depletion for the compounds. As shown in Fig. 1, high levels of cystine depletion are observed for compounds having a sulfur-nitrogen distance of about 3.6 to 4.7 Å, and in particular, about 4.1 Å.








[0139] Unless indicated (by the term “Abs”), relative stereochemistry is shown in Table 4.










Table 4










Compound	Cmpd. No.	S-N Dist. (Å)	Cystine Depletion (%)	Cystine Reactivity Assay (30°C) (%) Mean Value	Cystine Reactivity Assay (RT) (RFU) Mean Value	ADO Metabolism Assay (%) Mean Value
	42694	4.2	20	<50	---	---
	42693	3.1	110	>90	---	---










	42692	3.1	100	>90	---	---
	42691	5.3	---	50-90	---	---
	42690	4.1	---	---	---	---
	42689	4.1	---	---	---	---
	42618	4.7	49	---	---	---
	42617	4	101	50-90	---	<25
	42616	4	79	50-90	---	<25
	42615	4.7	106	<50	---	<25
	42614	4.7	117	---	---	<25






	42613	4.1	109	>90	---	<25
	42612	4.2	98	---	---	---
	42611	5.9	55	>90	---	<25
	42610	3.1	91	>90	---	---
	42609	4.7	17	---	---	---
	42494	4.6	88	50-90	---	<25
	42493	4.1	74	50-90	---	<25
	42492	6.2	---	<50	---	---

	42491	5.1	---	<50	---	<25
	42460	4.7	111	50-90	---	<25
	42458	4.7	91	>90	---	---
	42457	3.8	108	50-90	---	<25
	42391	4.7	60	>90	---	---
	42389	4.5	36	50-90	---	<25
	42388	4.1	97	>90	---	<25

	42345	3.4	50	>90	---	<25
	42344	4.7	38	---	---	---
	42343	3.8	84	---	---	---
	42342	3.8	91	---	---	---
	42341	4.7	81	<50	---	<25
	42338	4.6	---	<50	---	---
	42248	4.1	103	>90	---	<25
	42247	4.1	96	>90	---	<25
	42213	3.7	0	>90	---	>75

	42212	3.3	---	<50	---	---
	42211	4.1	79	>90	---	<25
	42210	4.1	87	>90	---	<25
	42209	4.8	99	50-90	---	<25
	42208	4.1	102	>90	---	<25
	42153	4.6	---	---	---	---
	42152	4.1	126	---	---	---
	42151	3.8	133	---	---	---
	42099	4.1	79	---	---	25-75

	42098	4.1	70	---	---	<25
	42097	4.1	97	---	---	---
	41448	3.1	55	---	---	25-75
	41446	3.7	82	---	---	<25
	41188	4.7	87	50-90	---	<25
	18781	4.1	92	---	---	<25
	17522	6.9	6	---	<4000	<25
	17521	4.7	79	50-90	4000-9000	<25
	17518	4.1	99	---	>9000	<25
	17517	4.1	98	---	>9000	<25
	17515	4.1	97	---	4000-9000	<25
	17333	4.1	104	---	4000-9000	<25



	17330	3.2	---	---	4000-9000	25-75
	17329	5.2	36	---	<4000	<25
	17328	4.1	13	---	<4000	<25
	17287	4.1	---	---	<4000	<25
	1585	4.1	100	>90	4000-9000	>75
	18782	4.1	32	---	<4000	<25









Example 5 Activity of Compounds







[0140] Cystine depletion levels for each of the compounds in Table 5 are expressed as a percent relative to the cystine depletion achieved by cysteamine (c.f. cysteamine 100%) at a similar concentration.










[0141] Unless indicated (by the term “Abs”), relative stereochemistry is shown in Table 5.










Table 5








Compound	Cmpd. No.	Cystine Depletion (%) Mean Value	Cystine Reactivity Assay (30°C) (%) Mean Value	ADO Metabolism Assay (%) Mean Value
	152071	70	---	---
	151272	11	---	---

	151271	27	---	---
	151270	3	---	---
	151182	59	50-90	---
	151181	69	50-90	---
	150731	58	>90	---
	150730	14	>90	<25
	149611	27	>90	---
	149610	15	<50	<25

	149609	50	<50	<25
	149608	55	<50	<25
	149607	17	>90	---
	145696	32	>90	<25
	145695	0	>90	<25
	145694	0	>90	<25
	145693	81	50-90	<25
	145692	75	---	---
	145691	0	---	---

	145611	9	<50	<25
	145610	11	<50	<25
	145609	121	---	<25
	145592	156	>90	---
	145591	61	50-90	<25
	145590	104	>90	<25
	145589	70	50-90	<25
	145588	31	>90	<25
	145587	13	>90	<25

	145586	21	50-90	<25
	145585	11	>90	---
	145584	0	>90	<25
	145566	16	<50	<25
	145565	31	---	---
	145564	36	>90	25-75
	145232	16	<50	<25
	42460	78	50-90	---
	42457	77	50-90	---
	42388	63	>90	<25

	42341	83	<50	---
	42210	89	>90	---
	41446	82	>90	<25
	2415	14	---	---
	2403	19	<50	<25
	1176	10	<50	<25
	1141	0	<50	---
	981	25	<50	<25

	653	23	<50	<25
--	-----	----	-----	-----

Example 6
Dimethyl Sulfide (DMS) Detection in Rat Whole Blood Following Oral Compound Administration

[0142] Male Sprague Dawley (SD) rats (200-250g) were housed in cages under a 12 hour light/dark cycle with free access to food and water. Temperature and humidity were controlled according to UK Home Office regulations. For oral (po) administration, test compounds were formulated in water at a concentration of 30 mg/mL, to provide a dose of 150 mg/kg when administered orally in a 5 mL/kg dosing volume. Following compound or water (vehicle) administration to rats (n=15 per compound), terminal blood samples (>5 mL) were taken by cardiac puncture under CO₂ at defined time-points post oral dosing (pre-dose, 0.25 hour, 0.5 hour, 1 hour and 2 hour). Blood was placed in sealed heparinized Eppendorf tubes on wet ice and stored at 4°C prior to analyzing DMS levels by gas chromatography (GC).

[0143] For each sample of rat blood, 1 mL was added to an evacuated 20 mL GC-Headspace vial containing tetrahydrofuran (THF) internal standard. Headspace vials were sealed with a crimp cap and septum (to prevent the vaporization of volatile sulfur compounds into the open air) and thoroughly vortex mixed. All samples were processed using gas chromatography to measure DMS levels. In short, sulfur containing compounds were thermally liberated into the gas carrier stream and injected (1 µL) into the gas chromatography column (30 m x 0.32 mm DB-624 1.8 µm). Under these conditions, the retention time for dimethyl sulfide (DMS) and THF internal standard were approximately 1.83 minutes and 3.82 minutes respectively.

[0144] For each analyzed blood sample, the ratio of DMS peak area to THF peak area was calculated. For the DMS calibration curve and assay linearity checks, this ratio was graphically plotted against the concentration of reference DMS standards (blank, 60 nM, 180 nM, 360 nM, 600 nM, 1200 nM, 1800 nM and 3000 nM). The concentration of DMS in each

blood sample was calculated from the equation produced by this plot by linear regression analysis. The results are shown in Table 6. The lower limit of detection was 60 nM and a value of <60 indicates no DMS peak was detected in the sample chromatogram.

Advantageously, the tested compounds produced reduced levels of DMS compared to cysteamine, suggesting that these compounds have reduced unpleasant side effects (e.g., halitosis) compared to cysteamine therapy.

Table 6

Compound	Cmpd. No.	Dose (mg/kg) po	Mean DMS levels (nM) $\pm$ STD				
			Pre-Dose	Time 0.25 hr	Time 0.5 hr	Time 1 hr	Time 2 hr
<chem>NCCS</chem>	1585	150	<60	641 $\pm$ 324	1097 $\pm$ 418	1162 $\pm$ 373	1437 $\pm$ 334
<chem>C1CCN(C1)CS</chem>	42341	150	<60	<60	<60	<60	<60
<chem>C1CCN(C1)S</chem>	42210	150	<60	<60	No data	<60	<60
<chem>C1CCN(C1)CS</chem>	41188	150	<60	<60	No data	<60	<60
<chem>C1CCN(C1)CS</chem>	41446	150	<60	<60	No data	<60	<60
<chem>CC(C)(C)N</chem>	18782	150	<60	<60	No data	<60	<60




Example 7


Glutamate-Induced Neuronal Excitotoxicity Assay

[0145] The ability of test compounds to inhibit glutamate-induced excitotoxicity (neuroprotection) in St-HdhQ^{111/111} cells was determined by incubating compounds with cells for 60 min at 33°C, 95% (v:v) air/ 5% (v:v) CO₂ in Dulbecco's Modified Eagle Media. Following this, excitotoxicity was induced by the addition of 6 mM L-glutamate for 24 hours. Cell viability was assessed by measuring ATP levels using a luminescent-based CellTitre Glo assay (Promega). Compound neuroprotection (% cell survival) was expressed as a % of the effect recorded with 100 μ M cysteamine (denoted as 100% cell survival). Advantageously, the tested compounds resulted in high levels of cell survival (e.g., at least 50% cell survival when expressed as a % of the effect for 100 μ M cysteamine), including levels of cell survival highly similar to that of cysteamine (e.g., at least 80% cell survival when expressed as a % of

the effect for 100 μ M cysteamine), suggesting that these compounds have similar neuroprotection levels as compared to cysteamine therapy.

Table 7


Compound / Compound No.	HdhQ ^{III/III} cell % survival (mean $\pm$ STD)	
	Test Conc. 100 μ M	Test Conc. 10 μ M
151182	85 $\pm$ 11	1 $\pm$ 5
151181	90 $\pm$ 9	3 $\pm$ 8
149611	102 $\pm$ 13	35 $\pm$ 13
149607	98 $\pm$ 12	32 $\pm$ 9
145695	74 $\pm$ 10	8 $\pm$ 10
145611	94 $\pm$ 8	24 $\pm$ 13
145610	87 $\pm$ 10	11 $\pm$ 8
145592	88 $\pm$ 8	17 $\pm$ 10
145590	90 $\pm$ 11	14 $\pm$ 7
145588	98 $\pm$ 5	9 $\pm$ 9
145587	97 $\pm$ 7	52 $\pm$ 9
145586	89 $\pm$ 9	9 $\pm$ 6
145585	100 $\pm$ 9	13 $\pm$ 10
 145158	86 $\pm$ 10	3 $\pm$ 7
 145154	93 $\pm$ 9	13 $\pm$ 13
 145153	95 $\pm$ 6	11 $\pm$ 11
42693	77 $\pm$ 9	80 $\pm$ 10
42692	71 $\pm$ 5	82 $\pm$ 4
42617	89 $\pm$ 7	7 $\pm$ 10
42616	56 $\pm$ 7	5 $\pm$ 7
42614	59 $\pm$ 12	2 $\pm$ 6
42610	77 $\pm$ 8	19 $\pm$ 6
42493	87 $\pm$ 10	6 $\pm$ 10
42460	95 $\pm$ 9	3 $\pm$ 5
42457	91 $\pm$ 6	6 $\pm$ 5
42388	91 $\pm$ 5	5 $\pm$ 6
42341	55 $\pm$ 21	1 $\pm$ 10
42248	81 $\pm$ 6	2 $\pm$ 7

42247	84 ± 13	5 ± 8
42211	84 ± 19	2 ± 9
42210	105 ± 8	7 ± 6
41446	101 ± 11	6 ± 9
41188	105 ± 11	20 ± 6
1585	105 ± 13	20 ± 6
145694	15 ± 15	2 ± 10
 145156	16 ± 7	1 ± 6
42494	28 ± 17	0 ± 7

[0146] Numerous modifications and variations in the invention as set forth in the above illustrative examples are expected to occur to those skilled in the art. Consequently only such limitations as appear in the appended claims should be placed on the invention.

WHAT IS CLAIMED:

1. A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula I or a disulfide thereof:



wherein:

R^1 and R^2 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^1 and R^2 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^3 and R^4 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

G is selected from the group consisting of $-NR^5R^6$ and $-CR^7R^8NR^5R^6$;

R^5 and R^6 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

R^7 and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl; or

R^7 and R^8 , taken together with the carbon atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;

R^2 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;

R^4 and R^6 , taken together with the atoms to which they are attached, form a 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered heterocyclic ring;


R^2 and R^8 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring; or

R^2 and R^4 , taken together with the atoms to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered carbocyclic ring;


with the proviso that when G is $-NH_2$, at least one of R^1 , R^2 , R^3 , and R^4 is other than H.

2. The method of claim 1, wherein R^5 and R^6 are independently selected from the group consisting of H, methyl, and ethyl.

3. The method of claim 1, wherein R^5 and R^6 , taken together with the nitrogen atom to which they are attached, form a 5-membered heterocyclic ring.
4. The method of any of the preceding claims, wherein R^4 is methyl.
5. The method of claim 4, wherein R^3 is methyl.
6. The method of any one of claims 1-3, wherein R^3 and R^4 , taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.
7. The method of any one the preceding claims, wherein R^2 is methyl.
8. The method of claim 7, wherein R^1 is methyl.
9. The method of any one of claims 1-6, wherein R^1 and R^2 , taken together with the carbon atom to which they are attached, form a 3-membered carbocyclic ring.
10. The method of claim 1, wherein G is $—CR^7R^8NR^5R^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 6-membered heterocyclic ring.
11. The method of claim 10, wherein R^5 is methyl.
12. The method of claim 1, wherein G is $—NR^5R^6$, and R^2 and R^6 , taken together with the atoms to which they are attached, form a 4- or 6-membered heterocyclic ring.
13. The method of claim 12, wherein R^5 is H.
14. The method of any one the preceding claims, wherein R^7 and R^8 are both H.
15. The method of claim 1, wherein the compound of formula I is selected from the group consisting of:




16. The method of claim 1, wherein the compound of formula I is selected from the group consisting of:



17. The method of claim 16, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 are independently selected from the group consisting of H and C_{1-5} alkyl.

18. The method of claim 16, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and R^8 are independently selected from the group consisting of H and methyl.

19. A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula II, formula III, or a disulfide thereof:



wherein:

L is a hydrocarbon linking group;

R^9 and R^{10} are independently selected from the group consisting of H, C_{1-5} alkyl, and $\text{CO}(C_{1-5}\text{alkyl})$; or

R^9 and R^{10} , taken together with the nitrogen atom to which they are attached, form a 3-, 4-, 5-, 6-, 7-, or 8-membered heterocyclic ring;

A is a heterocyclic ring contain one N atom; and

n is 0, 1, 2, or 3;

wherein the S atom in the compound of formula II or formula III is a distance of about 3.6 Angstroms to about 4.7 Angstroms from the N atom in the compound;

with the proviso that the compound of formula II is not cysteamine.


20. The method of claim 19 wherein L is a 3-, 4-, 5-, 6-, 7-, or 8-membered cycloalkyl ring or a 6-membered aryl ring.

21. The method of claim 19 wherein L is C_{1-5} alkyl.

22. The method of any one of claims 19-21 wherein L is substituted with one to four groups selected from halo, C_{1-5} alkyl, C_{3-5} cycloalkyl, and $\text{---CO}_2(C_{1-5}\text{alkyl})$.

23. The method of claim 19 wherein A is a 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic heterocycloalkyl ring, a 6-, 7-, or 8-membered bicyclic heterocycloalkyl ring, or a 5- or 6-membered heteroaryl ring.

24. The method of claim 19 wherein the compound of formula III has a structure IIIa:



wherein R¹¹ is selected from the group consisting of H and C₁₋₅alkyl.

25. The method of any one of claims 19, 23, or 24 wherein A is substituted with one to four groups selected from halo, C₁₋₅alkyl, C₃₋₅cycloalkyl, and —CO₂(C₁₋₅alkyl).

26. The method of claim 19 wherein the S atom in the compound of formula II or formula III is a distance of about 3.8 Angstroms to about 4.4 Angstroms from the N atom in the compound.

27. The method of claim 26 wherein the S atom in the compound of formula II or formula III is a distance of about 4.0 Angstroms to about 4.2 Angstroms from the N atom in the compound.

28. The method of claim 27 wherein the S atom in the compound of formula II or formula III is a distance of about 4.1 Angstroms from the N atom in the compound.

29. The method of any one of claims 19-28 wherein the compound of formula II, formula III, or disulfide thereof depletes cystine in a subject in an amount that is at least 70% of the level of depletion of cystine by cysteamine.

30. The method of any one of claims 1-29, wherein the compound of formula I, formula II, formula III, or disulfide thereof is resistant to metabolism by cysteamine dioxygenase (ADO).

31. The method of claim 30, wherein less than 20% of the compound of formula I, formula II, formula III, or disulfide thereof is metabolized by ADO when assayed by consumption of oxygen using an oxygen sensitive fluorescent probe.

32. The method of any one of claims 1-31 wherein the disease is selected from the group consisting of cystinosis, fatty liver disease, fibrosis, a thrombotic disease, an MECP-2 related disorder, an inherited mitochondrial disease, a neurological disease or disorder, inflammation and cancer.

33. The method of claim 32 wherein the fatty liver disease is selected from the group consisting of non-alcoholic fatty liver disease (NAFLD), non-alcoholic steatohepatitis (NASH), fatty liver disease resulting from hepatitis, fatty liver disease resulting from obesity, fatty liver disease resulting from diabetes, fatty liver disease resulting from insulin resistance, fatty liver disease resulting from hypertriglyceridemia, Abetalipoproteinemia, glycogen storage diseases, Weber-Christian disease, Wolmans disease, acute fatty liver of pregnancy, and lipodystrophy.

34. The method of claim 32 wherein the fibrosis is selected from the group consisting of atherosclerosis, asthma, cardiac fibrosis, organ transplant fibrosis, colloid and hypertrophic scar, muscle fibrosis, pancreatic fibrosis, bone-marrow fibrosis, interstitial liver fibrosis, cirrhosis of liver and gallbladder, scleroderma, pulmonary fibrosis, diffuse parenchymal lung disease, idiopathic interstitial fibrosis, interstitial pneumonitis, desquamative interstitial pneumonia, respiratory bronchiolitis, interstitial lung disease, acute interstitial pneumonitis, nonspecific interstitial pneumonia, cryptogenic organizing pneumonia, lymphocytic interstitial pneumonia, renal fibrosis, and chronic kidney disease, cystic fibrosis and Alport's disease.

35. The method of claim 32 wherein the thrombotic disease is selected from the group consisting of sickle cell disease, deep vein thrombosis, pulmonary embolism, cardiac embolism, hypercoagulable state, thrombophilia, Factor V Leiden, Antithrombin III deficiency, Protein C deficiency, Protein S deficiency, Prothrombin gene mutation (G20210A), Hyperhomocysteinemia, antiphospholipid antibody syndrome (APS), anticardiolipin antibody (ACLA) thrombosis syndrome, or lupus anticoagulant (LA) syndrome.


36. The method of claim 32 wherein the neurological disease or disorder is selected from the group consisting of Huntington's Disease, Parkinson's Disease, amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's disease spinal muscle atrophy, concussion, stroke, and traumatic brain injury (CTE).

37. The method of claim 32 wherein the MECP-2 related disease is selected from the group consisting of Rett syndrome, autism, pervasive development disorder, non-syndromic mental retardation, idiopathic neonatal encephalopathy and idiopathic cerebral palsy.

38. The method of claim 32 wherein the inherited mitochondrial disease is selected from the group consisting of Friedreich's ataxia, Leber's hereditary optic neuropathy (LHON), myoclonic epilepsy and ragged-red fibers, mitochondrial encephalomyopathy, lactic acidosis, and stroke-like syndrome (MELAS), Kearns-Sayre syndrome and subacute necrotizing encephalopathy (Leigh's Syndrome).

39. The method of claim 32 wherein the cancer is selected from the group consisting of breast cancer, melanoma, prostate cancer, pancreatic cancer, head and neck cancer, lung cancer, non small-cell lung carcinoma, renal cancer, colorectal cancer, colon cancer, ovarian cancer, liver cancer and gastric cancer.


40. A compound having the following structure or a disulfide thereof:



wherein R¹ is C₁₋₅alkyl; and

R³, R⁴, R⁵, R⁷, and R⁸ are independently selected from the group consisting of H and C₁₋₅alkyl.

41. A compound having the following structure or a disulfide thereof:



wherein R¹, R³, R⁴, R⁵, R⁶, and R⁷ are independently selected from the group consisting of H and C₁₋₅alkyl.




FIG. 1

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/040637

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K31/145 A61K31/397 A61K31/40 A61K31/4402 A61K31/4409
A61K31/445 A61K31/4465 C07C323/30 C07D211/54

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K C07C C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, BIOSIS, EMBASE, SCISEARCH

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2009/070781 A1 (UNIV CALIFORNIA [US]; DOHIL RANJAN [US]; SCHNEIDER JERRY [US]) 4 June 2009 (2009-06-04)	1-9, 12, 13, 15-18, 30-33
Y	pages 9-10, paragraph [0025] page 21, paragraph [0051] claims 1, 2, 6, 7 ----- -/--	1-18, 37, 38



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 November 2016

Date of mailing of the international search report

25/11/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Cielen, Elsie

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/040637

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/078335 A1 (UNIV CALIFORNIA [US]) 30 May 2013 (2013-05-30)	1-9,12, 13, 15-18, 30-32, 35,36
Y	page 7, paragraph [0014] pages 22-23, paragraph [0072] page 32, paragraph [0097] page 34, paragraph [0101] pages 35-36, paragraph [0107] claims 1-3, 31, 32, 43, 52-54 -----	1-18,37, 38
X	WO 2013/062544 A1 (SEATTLE CHILDREN S RES INST [US]; EDDY ALLISON A [US]; OKAMURA DARYL M) 2 May 2013 (2013-05-02)	1-9,12, 13,15, 18, 30-32,34
Y	page 2, paragraph [0007] page 4, paragraph [0014] pages 18-19, paragraph [0060] claims 1, 9 -----	1-18,37, 38
X	WO 2005/014524 A2 (WISCONSIN ALUMNI RES FOUND [US]; FAHL WILLIAM E [US]; PEEBLES DANIEL D) 17 February 2005 (2005-02-17) page 1, paragraph 3 page 8, paragraph 2-4 page 10, paragraph 2 page 29, paragraph 3 claims 129, 130 figure 7A -----	1-18, 30-32,40
X	WO 2005/039548 A2 (PROBIODRUG AG [DE]; SCHILLING STEPHAN [DE]; HOFFMANN TORSTEN [DE]; NIE) 6 May 2005 (2005-05-06) page 39, lines 9-15 page 44, lines 7-19 page 46, lines 1-5 table 5 page 50, line 31 - page 51, line 8 claims 1, 5, 6, 11, 13 -----	1-18, 30-32, 36,39,40
Y	WO 2015/069888 A2 (RAPTOR PHARMACEUTICALS INC [US]) 14 May 2015 (2015-05-14) page 16, paragraph [0064] pages 17-18, paragraph [0068] page 28, paragraph [0099] claims 1, 2, 7 -----	1-18,38
Y	WO 2012/055826 A1 (UNIV AIX MARSEILLE II [FR]; INST NAT SANTE RECH MED [FR]; INST CURIE []) 3 May 2012 (2012-05-03) page 10, lines 4-13 claims 1, 2 -----	1-18,37
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/040637

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 2016/046524 A1 (NOVABIOTICS LTD [GB]) 31 March 2016 (2016-03-31) page 4, line 31 - page 5, line 9 page 6, lines 6-15 page 17, lines 20-27 claims 1, 5 -----	1,2,7, 30-32,34
X	DATABASE WPI Week 199126 Thomson Scientific, London, GB; AN 1991-190158 XP002764300, & JP H03 118319 A (GREEN CROSS CORP) 20 May 1991 (1991-05-20) abstract -----	1-9,12, 13, 15-18, 30-32
X	APFFEL C A ET AL: "TUMOR REJECTION N EXPERIMENTAL ANIMALS TREATED WITH RADIOPROTECTIVE THIOLS", CANCER RESEARCH, AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 35, no. 2, 1 February 1975 (1975-02-01), pages 429-437, XP001033861, ISSN: 0008-5472 abstract page 429, column 2, paragraph 3 - page 430, column 1, paragraph 2 tables 1, 2 -----	1-11, 14-18, 30-32,39
X	WO 93/06832 A1 (THOENE JESS G [US]) 15 April 1993 (1993-04-15) page 8, paragraph 1-2 claim 4 -----	1-11, 14-18, 30-32
X	WO 2008/128919 A2 (UCB PHARMA SA [BE]; CELANIRE SYLVAIN [FR]; DENONNE FREDERIC [BE]; VALA) 30 October 2008 (2008-10-30) page 13, lines 13-35 page 26, lines 11-16 page 31, line 25 page 33, line 7 example 5; compound a58 page 54, lines 14-27 claim 18 ----- -/--	41

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/040637

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BERNIE CHOUDARY ET AL: "Synthesis of the Pleuromutilin Antibiotic SB-268091: A New Practical and Efficient Synthesis of Quinuclidine-4-thiol", ORGANIC PROCESS RESEARCH AND DEVELOPMENT, vol. 16, no. 12, 21 December 2012 (2012-12-21), pages 1927-1939, XP55320266, US ISSN: 1083-6160, DOI: 10.1021/op300263w figure scheme 2; compound 4 figure scheme 4; compound 12 -----	40
A	MACDONALD I R ET AL: "Thioesters for the in vitro evaluation of agents to image brain cholinesterases", JOURNAL OF ENZYME INHIBITION AND MEDICINAL CHEMISTRY 2013 INFORMA HEALTHCARE GBR, vol. 28, no. 3, June 2013 (2013-06), pages 447-455, XP008182281, ISSN: 1475-6366 abstract page 452; figure scheme 1a -----	40
A	RAO V S: "A convenient preparation of 1-mercapto-3-azidocyclobutane from 1,1-cyclobutanedicarboxylic acid", SYNTHETIC COMMUNICATIONS 1993 US, vol. 23, no. 20, 1993, pages 2915-2920, XP008182285, ISSN: 0039-7911 figure scheme 1; compounds 6, 9 -----	41
A	JP 2001 131067 A (SANKYO CO; UBE INDUSTRIES) 15 May 2001 (2001-05-15) abstract -----	1,3,32, 35

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2016/040637

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-18, 40, 41(completely); 30-39(partially)
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 12, 13(completely); 1-9, 15-18, 30-39(partially)

A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula I or a disulfide thereof wherein G is -NR⁵R⁶ with the proviso that when G is -NH₂ , at least one of R¹ , R² , R³ , and R⁴ is other than H.

2. claims: 10, 11, 14, 40, 41(completely); 1-9, 15-18, 30-39(partially)

A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula I or a disulfide thereof wherein G is -CR⁷R⁸NR⁵R⁶. A compound as defined in claims 40 or 41.

3. claims: 20-22(completely); 19, 26-39(partially)

A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula II, in as far as not comprised in inventions 1 or 2.

4. claims: 23-25(completely); 19, 26-39(partially)

A method of treating a patient suffering from a disease for which treatment with cysteamine is indicated comprising administering to the patient an effective amount of a composition comprising a compound of formula III or IIIa, in as far as not comprised in inventions 1 or 2.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/040637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2009070781	A1	04-06-2009	AR 069500 A1 27-01-2010
		AU 2008329628 A1 04-06-2009	
		BR PI0819690 A2 07-10-2014	
		CA 2706768 A1 04-06-2009	
		CN 101932238 A 29-12-2010	
		CN 104825430 A 12-08-2015	
		DK 2214480 T3 08-07-2013	
		EP 2214480 A1 11-08-2010	
		EP 2636404 A2 11-09-2013	
		ES 2417179 T3 06-08-2013	
		HK 1141945 A1 09-08-2013	
		HR P20130590 T1 31-08-2013	
		JP 5583022 B2 03-09-2014	
		JP 2011505375 A 24-02-2011	
		JP 2014237670 A 18-12-2014	
		KR 20100091219 A 18-08-2010	
		MY 156316 A 11-02-2016	
		NZ 585732 A 24-02-2012	
		PT 2214480 E 23-07-2013	
		RU 2010126631 A 10-01-2012	
		SG 10201403200R A 30-10-2014	
		TW 200927093 A 01-07-2009	
		TW 201513855 A 16-04-2015	
		US 2010303870 A1 02-12-2010	
		US 2012015005 A1 19-01-2012	
		US 2013183351 A1 18-07-2013	
		US 2016143865 A1 26-05-2016	
		WO 2009070781 A1 04-06-2009	
		ZA 201003670 B 31-08-2011	
WO 2013078335	A1	30-05-2013	AU 2012340670 A1 17-04-2014
			CA 2851387 A1 30-05-2013
			CN 103974698 A 06-08-2014
			EP 2782564 A1 01-10-2014
			HK 1202440 A1 02-10-2015
			JP 2014533701 A 15-12-2014
			KR 20140097132 A 06-08-2014
			RU 2014125282 A 27-12-2015
			SG 11201402472Q A 27-06-2014
			US 2014322315 A1 30-10-2014
			WO 2013078335 A1 30-05-2013
WO 2013062544	A1	02-05-2013	AU 2011379972 A1 15-05-2014
			CA 2853484 A1 02-05-2013
			US 2014275279 A1 18-09-2014
			WO 2013062544 A1 02-05-2013
WO 2005014524	A2	17-02-2005	US 2005101676 A1 12-05-2005
			WO 2005014524 A2 17-02-2005
WO 2005039548	A2	06-05-2005	AU 2004283461 A1 06-05-2005
			BR PI0415409 A 05-12-2006
			CA 2542419 A1 06-05-2005
			CN 1867324 A 22-11-2006
			CN 102302781 A 04-01-2012
			DK 1675578 T3 26-08-2013
			EA 200600618 A1 27-10-2006
			EP 1675578 A2 05-07-2006

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/040637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		EP 2289498 A1	02-03-2011
		IL 174372 A	27-09-2011
		JP 5707014 B2	22-04-2015
		JP 2007508347 A	05-04-2007
		JP 2015057400 A	26-03-2015
		KR 20070017477 A	12-02-2007
		KR 20120007079 A	19-01-2012
		KR 20120125379 A	14-11-2012
		MX PA06003998 A	27-06-2006
		NZ 546322 A	28-11-2008
		US 2006189523 A1	24-08-2006
		US 2009149394 A1	11-06-2009
		WO 2005039548 A2	06-05-2005
WO 2015069888	A2	14-05-2015	
		CA 2928442 A1	14-05-2015
		CN 105873579 A	17-08-2016
		EA 201690936 A1	31-08-2016
		EP 3065725 A2	14-09-2016
		KR 20160070154 A	17-06-2016
		TW 201605434 A	16-02-2016
		US 2015125526 A1	07-05-2015
		WO 2015069888 A2	14-05-2015
WO 2012055826	A1	03-05-2012	
		AU 2011322715 A1	23-05-2013
		CA 2815582 A1	03-05-2012
		CN 103260612 A	21-08-2013
		EA 201390616 A1	30-09-2013
		EP 2632444 A1	04-09-2013
		JP 2013540789 A	07-11-2013
		KR 20140041387 A	04-04-2014
		NZ 609640 A	26-06-2015
		SG 189519 A1	31-05-2013
		US 2013316961 A1	28-11-2013
		US 2015265554 A1	24-09-2015
		WO 2012055826 A1	03-05-2012
WO 2016046524	A1	31-03-2016	NONE
JP H03118319	A	20-05-1991	NONE
WO 9306832	A1	15-04-1993	
		AT 173625 T	15-12-1998
		AU 675412 B2	06-02-1997
		BR 9206571 A	17-10-1995
		CA 2120140 A1	15-04-1993
		DE 69227696 D1	07-01-1999
		DE 69227696 T2	20-05-1999
		EP 0607252 A1	27-07-1994
		JP H06511008 A	08-12-1994
		OA 9917 A	15-09-1994
		PT 100914 A	29-10-1993
		WO 9306832 A1	15-04-1993
WO 2008128919	A2	30-10-2008	
		AU 2008240832 A1	30-10-2008
		BR PI0810009 A2	14-10-2014
		CA 2682539 A1	30-10-2008
		CN 101663290 A	03-03-2010
		EA 200901378 A1	30-04-2010
		EP 2146980 A2	27-01-2010

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/040637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		JP 2010524881 A	22-07-2010
		US 2010305116 A1	02-12-2010
		WO 2008128919 A2	30-10-2008

JP 2001131067	A	15-05-2001	NONE
