

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
06 June 2019 (06.06.2019)



(10) International Publication Number
WO 2019/106609 A1

(51) International Patent Classification:

A61K 47/68 (2017.01) C07K 16/24 (2006.01)
A61P 19/02 (2006.01)

MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/IB2018/059482

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

(22) International Filing Date:

29 November 2018 (29.11.2018)

Published:

— with international search report (Art. 21(3))
— with sequence listing part of description (Rule 5.2(a))
— in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/593,776 01 December 2017 (01.12.2017) US
62/595,054 05 December 2017 (05.12.2017) US

(71) Applicant: **ABBVIE INC.** [US/US]; 1 North Waukegan Road, North Chicago, Illinois 60064 (US).

(72) Inventors: **HOBSON, Adrian D.**; 25 Holman Street, Shrewsbury, Massachusetts 01545 (US). **MCPHERSON, Michael J.**; 287 Pillsbury Road, Ashby, Massachusetts 01431 (US). **WAEGELL, Wendy**; 7 Rice Corner Road, Brookfield, Massachusetts 01506 (US). **GOESS, Christian**; 112 Mashapaug Road, Sturbridge, Massachusetts 01566 (US). **HERNANDEZ, Axel, Jr.**; 14 Laurie Lane, Charlton, Massachusetts 01507 (US). **WANG, Lu**; 14 Stratton Way, Northborough, Massachusetts 01532 (US). **WANG, Lu**; 12 Nipmuck Drive, Westborough, Massachusetts 01581 (US). **MARVIN, Christopher C.**; 242 Braxton Way, Grayslake, Illinois 60030 (US). **SANTORA, Ling C.**; 7 Thoreau Circle, Shrewsbury, Massachusetts 01545 (US).

(74) Agent: **HERRITT, Danielle**; Womble Bond Dickinson (US) LLP, Two International Place, Suite 2310, Boston, Massachusetts 02110 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,

(54) Title: GLUCOCORTICOID RECEPTOR AGONIST AND IMMUNOCONJUGATES THEREOF

(57) Abstract: Provided herein are glucocorticoid receptor agonist immunoconjugates, glucocorticoid receptor agonists, and methods of using the same.



WO 2019/106609 A1

GLUCOCORTICOID RECEPTOR AGONIST AND IMMUNOCONJUGATES THEREOF**RELATED APPLICATIONS**

[0001] This application claims priority to US Provisional Application Number 62/593,776, filed December 1, 2017 and US Provisional Application Number 62/595,054, filed December 5, 2017, each of which is incorporated by reference in its entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on November 28, 2018, is named A103017_1490WO_SL.txt and is 14,758 bytes in size.

TECHNICAL FIELD


[0003] Tumor Necrosis Factor α (TNF α) plays a central role in the pathophysiology of several human disorders, and anti-TNF α agents have clinically validated therapeutic utility in the treatment of autoimmune and inflammatory disorders, such as rheumatoid arthritis, psoriasis and inflammatory bowel disease. Despite their success in the clinic, anti-TNF α biologics are still limited in the maximal efficacy they can achieve in patients, necessitating the identification and development of more potent and effective therapeutics. Patients treated with anti-TNF α biologics may also develop an immunogenic response to the therapeutic thus limiting its effectiveness. Therefore anti-TNF α therapies with lower immunogenicity and high efficacy would be useful for further controlling disease.

[0004] Synthetic glucocorticoid receptor agonists are a potent class of small molecules used in the treatment of inflammatory disorders, but their utility in the chronic treatment of disease is limited due to severe side effects. There is a need to develop therapeutics with enhanced efficacy and longer duration of action compared to anti-TNF antibodies and with minimal unwanted effects.


SUMMARY

[0005] The present disclosure provides glucocorticoid receptor agonist immunoconjugates useful for treating autoimmune diseases.


[0006] In one aspect, the present disclosure provides an antibody drug conjugate comprising:
(a) an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4; and (b) a glucocorticoid receptor agonist comprising a radical represented by the formula:



and wherein the antibody is conjugated to the glucocorticoid receptor agonist via a linker represented by the formula:




[0007] In one embodiment, the present disclosure provides an antibody drug conjugate according to the formula:




wherein A is the antibody and n is an integer from 1-10.

[0008] In one aspect, the present disclosure provides an antibody drug conjugate comprising:


(a) an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4; and (b) a glucocorticoid receptor agonist comprising a radical represented by the formula:



and wherein the antibody is conjugated to the glucocorticoid receptor agonist via a linker represented by the formula:




[0009] In one embodiment, the present disclosure provides an antibody drug conjugate according to the formula:



wherein A is the antibody and n is an integer from 1-10.


[0010] In one embodiment, the present disclosure provides the antibody drug conjugate of any preceding embodiment, wherein the drug loading is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10. In one embodiment, the present disclosure provides the antibody drug conjugate of any preceding embodiment, wherein the drug loading is 4, for example, n in the foregoing antibody drug conjugate formula equals 4. In one embodiment, the present disclosure provides the antibody drug conjugate of any preceding embodiment, wherein the drug loading is 2, for example, n in the foregoing antibody drug conjugate formula equals 2.

[0011] In one embodiment, the present disclosure provides a method of making the antibody drug conjugate of any preceding embodiment, comprising the step of conjugating the antibody to the glucocorticoid receptor agonist. In one embodiment, the present disclosure provides the method of the preceding embodiment, further comprising the step of introducing a PO₄ moiety on the glucocorticoid receptor agonist before conjugating the antibody to the glucocorticoid receptor agonist. In one embodiment, the present disclosure provides the method of any preceding embodiment, wherein the conjugating comprises partially reducing the antibody, and alkylating the partially reduced antibody with a compound according to the formula:




[0012] In one embodiment, the present disclosure provides a pharmaceutical composition comprising the antibody drug conjugate of any preceding embodiment and a pharmaceutically acceptable carrier. In one embodiment, the present disclosure provides the pharmaceutical composition of any of the preceding embodiments, comprising a drug to antibody ratio (DAR) of 1-10.

[0013] In one preferred embodiment, the present disclosure provides an antibody drug conjugate according to the formula:




wherein A is adalimumab and n is 4.

[0014] In one preferred embodiment, the present disclosure provides an antibody drug conjugate according to the formula:




wherein A is an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4, and n is 4.

[0015] In another preferred embodiment, the present disclosure provides an antibody drug conjugate according to the formula:



[0016] wherein A is adalimumab and n is 2. In another preferred embodiment, the present disclosure provides an antibody drug conjugate according to the formula:



wherein A is an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4, and n is 2.

[0017] In one preferred embodiment, the present disclosure provides the pharmaceutical composition of any preceding embodiment, comprising a drug to antibody ratio (DAR) of 2.0.

[0018] In one preferred embodiment, the present disclosure provides the pharmaceutical composition of any preceding embodiment, comprising a drug to antibody ratio (DAR) of 4.0.

[0019] In one embodiment, the present disclosure provides a method of treating a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis in a subject, comprising administering an effective amount of the antibody drug conjugate of any preceding embodiment or the pharmaceutical composition of any preceding embodiment to the subject.

[0020] In one embodiment, the present disclosure provides the antibody drug conjugate of any preceding embodiment or the pharmaceutical composition of any preceding embodiment for use in the treatment of a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis.

[0021] In one embodiment, the present disclosure provides use of the antibody drug conjugate of any preceding embodiment or the pharmaceutical composition of any preceding embodiment for preparation of a medicament for treating a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis.

[0022] In one embodiment, the present disclosure provides a kit comprising: (a) a container comprising the antibody drug conjugate of any preceding embodiment or the pharmaceutical composition of any preceding embodiment; and (b) a label or package insert on or associated with the one or more containers, wherein the label or package insert indicates that the antibody drug conjugate or pharmaceutical composition is used for treating a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis,

plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis.

[0023] In one embodiment, the present disclosure provides a method of delivering a glucocorticoid receptor agonist to a TNF α -expressing cell, comprising the step of contacting the cell with the antibody drug conjugate of any preceding embodiment. In one embodiment, the present disclosure provides a method of determining anti-inflammatory activity of an antibody drug conjugate comprising: (a) contacting a TNF α -expressing cell with the antibody drug conjugate of any preceding embodiment; and (b) determining release of pro-inflammatory cytokines from the cell as compared to a control cell.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024] Fig. 1 provides a chromatographic resolution of BrAc-Gly-Glu-glucocorticoid receptor modulator (GRM)-PO₄, as performed and described in Example 7. As shown, the ADC is a heterogeneous ADC mixture containing ADCs with two drug linker molecules attached, and ADCs with four drug linker molecules attached.

[0025] Fig. 2 sets forth deconvoluted MS data of adalimumab conjugated with BrAc-Gly-Glu-glucocorticosteroid-PO₄. As shown, conjugation was achieved.


[0026] Fig. 3 provides a graph demonstrating the efficacy of a high and low dose of ADC1 compared to anti-TNF α mAb (high dose) or vehicle in a mouse collagen-induced arthritis (CIA) model of arthritis, as performed and described in Example 7. As shown, a single dose of anti-TNF α glucocorticosteroid ADC1 exhibited an extended duration of action through amelioration of paw swelling for ~28 days compared to anti-TNF α mAb or vehicles alone.

[0027] Fig. 4 is a graph of concentration (ug/ml) for closed and open ring ADC in cynomolgus monkeys over time, as performed and described in Example 7. As shown, the ring closed form is susceptible to the reverse Michael reaction and subsequent loss of linker-drug *in vivo*.

DETAILED DESCRIPTION

[0028] Provided herein are glucocorticoid receptor agonist immunoconjugates, glucocorticoid receptor agonists, and methods of making and using the same.

[0029] Provided herein is an antibody drug conjugate according to the formula:



wherein A is adalumimab and n is 4. As demonstrated in Example 7 below, this ADC (i.e., ADC4 below) demonstrates *in vitro* activity, stability in plasma, and minimal aggregation.

[0030] Also provided are methods of making and methods of using ADC4.

I. Definitions

[0031] To facilitate an understanding of the present disclosure, a number of terms and phrases are defined below.

[0032] The term "anti-TNF α protein" refers to proteins that are capable of (i) binding to TNF α and (ii) inhibiting binding of soluble TNF α to cell surface TNF receptors (p55 and/or p75) and/or lysing surface TNF α or TNF α receptor expressing cells *in vitro* in the presence of complement. In some embodiments, the anti-TNF antibody, can bind to TNF alpha on the surface of a cell and become internalized. For example, US 2014/0294813, which is herein incorporated by reference in its entirety, discloses anti-TNF antibodies that exhibit cellular internalization upon binding to cell surface human TNF. Anti-TNF α proteins include, for example, anti-TNF α antibodies (e.g., adalimumab, infliximab, and golimumab). Anti-TNF α antibodies are actively internalized upon binding to transmembrane TNF on monocyte-derived DCs and rapidly enter the lysosomes where they are degraded. (Deora et.al. MABS, 2017, Vol. 9, No. 4, 680-694).

[0033] The term "antibody", as used herein, is intended to refer to immunoglobulin molecules comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

[0034] The term "anti-TNF α antibody" or "an antibody that binds to TNF α " refers to an antibody that is capable of binding TNF α , e.g., with sufficient affinity such that the antibody is useful as a therapeutic agent

in targeting TNF α . The extent of binding of an anti-TNF α antibody to an unrelated, non-TNF α protein can be less than about 10% of the binding of the antibody to TNF α as measured, e.g., by a radioimmunoassay (RIA). In certain embodiments, an antibody that binds to TNF α has a dissociation constant (K_d) of ≤ 1 μ M, ≤ 100 nM, ≤ 10 nM, ≤ 1 nM, or ≤ 0.1 nM.

[0035] The term "immunoconjugate," "conjugate," "antibody-drug conjugate," or "ADC" as used herein refers to a compound or a derivative thereof that is conjugated to a protein such as a cell binding agent (e.g., an anti-TNF α antibody). Such immunoconjugates can be defined by a generic formula: (SM-L-Q)_n-A, wherein SM = radical derived from a small-molecule glucocorticoid receptor agonist, e.g., a glucocorticosteroid, L = linker, Q = heterobifunctional group or is absent, and A = a protein (e.g., an antibody), and n = 1-10. Immunoconjugates can also be defined by the generic formula in reverse order: A-(Q-L-SM)_n.

[0036] In the present disclosure, the term "linker" refers to a chemical moiety capable of linking the anti-TNF α protein (e.g. antibody) to a glucocorticosteroid. Linkers may be susceptible to cleavage (a "cleavable linker") thereby facilitating release of the glucocorticosteroid. For example, such cleavable linkers may be susceptible to peptidase-induced cleavage, at conditions under which the glucocorticosteroid and/or the antibody remains active.

[0037] In particular, the cleavable linker component disclosed herein comprises a peptide comprising two to three amino acid residues (a dipeptide or tripeptide) and specifically to dipeptides and tripeptides selected from the group consisting of alanine-alanine (Ala-Ala), glycine-glutamic acid (Gly-Glu), glutamic acid-alanine-alanine (Glu-Ala-Ala), and glycine-lysine (Gly-Lys). The peptide allows for cleavage of the linker by a protease, thereby facilitating release of the glucocorticosteroid upon exposure to intracellular proteases, such as lysosomal enzymes (Doronina et al. (2003) *Nat. Biotechnol.* 21:778-784).

[0038] In the present disclosure, the term "glucocorticosteroid" refers to a naturally-occurring or synthetic steroid hormone that interacts with glucocorticoid receptors, and specific glucocorticosteroids are disclosed in detail herein. A "radical of a glucocorticosteroid" is derived by the removal of one or more hydrogen atoms from a parent glucocorticosteroid. The removal of hydrogen atom(s) facilitates the attachment of the parent glucocorticosteroid to a linker. In the present disclosure, the hydrogen atom is removed from any suitable -NH₂ group of the parent glucocorticosteroid. In particular, the "radical of a glucocorticosteroid" is a monovalent radical derived from the removal of one hydrogen atom from a parent glucocorticosteroid.

[0039] In the present disclosure, the term "heterobifunctional group" refers to a chemical moiety that connects the linker and the anti-TNF α protein (e.g. antibody). Heterobifunctional groups are characterized as having different reactive groups at either end of the chemical moiety.

[0040] The term "drug antibody ratio" or "DAR" refers to the number of SMs (e.g., radical derived from a small-molecule glucocorticoid receptor agonist, e.g., a glucocorticosteroid) linked to A (e.g., an antibody). Thus, in the immunoconjugate having the generic formula (SM-L-Q)_n-A, the DAR is defined by drug loading per antibody drug conjugate, for example, "n".

[0041] When referring to a compound having formula (SM-L-Q)_n-A representing an individual immunoconjugate, the term "compound DAR" refers to the number of SMs linked to the individual A (e.g., drug loading or n as an integer of 1 to 10).

[0042] When referring to a compound having formula (SM-L-Q)_n-A representing a population of immunoconjugates, the term "population DAR" refers to the average number of SMs linked to the As (e.g., drug loading or n as an integer or fraction of 1 to 10 ± 0.5, ± 0.4, ± 0.3, ± 0.2, ± 0.1).

[0043] The term "subject" refers to humans, non-human primates, , and the like, which is to be the recipient of a particular treatment. Typically, the terms "subject" and "patient" are used interchangeably herein in reference to a human subject.

[0044] The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of the active ingredient to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered. The formulation can be sterile.

[0045] An "effective amount" of an immunoconjugate as disclosed herein is an amount sufficient to carry out a specifically stated purpose. An "effective amount" can be determined in relation to the stated purpose.

[0046] The term "therapeutically effective amount" refers to an amount of an immunoconjugate effective to "treat" a disease or disorder in a subject or mammal. A "prophylactically effective amount" refers to an amount effective to achieve the desired prophylactic result.

[0047] Terms such as "treating" or "treatment" or "to treat" or "alleviating" or "to alleviate" refer to therapeutic measures that cure, slow down, lessen one or more symptoms of, and/or slow or halt progression of a diagnosed pathologic condition or disorder ("therapeutic treatment"). Thus, those in need of therapeutic treatment include those already diagnosed with or suspected of having the disorder. Prophylactic or preventative measures refer to measures that prevent the development of a targeted pathological condition or disorder ("prophylactic treatment"). Thus, those in need of prophylactic treatment include those prone to have the disorder and those in whom the disorder is to be prevented.

II. Proteins for linkage to glucocorticoid receptor agonists

[0048] The present disclosure provides immunoconjugates containing glucocorticoid receptor agonists linked to proteins, for example, antibodies. In some embodiments, the antibody is human, humanized, chimeric, or murine. In some embodiments, the protein, e.g., antibody, can bind to a target on the surface of a cell and become internalized.

[0049] The present disclosure also provides immunoconjugates containing glucocorticoid receptor agonists linked to anti-TNF α proteins. In certain embodiments, the anti-TNF α proteins are antibodies. In certain embodiments, the anti-TNF α proteins are antibodies that bind to TNF α (e.g., soluble TNF α and/or membrane bound TNF α). In certain embodiments, the anti-TNF α proteins are soluble TNF receptor

proteins, e.g., soluble TNF receptor proteins fused to a heavy chain constant. In some embodiments, the anti-TNF α protein, e.g., anti-TNF α antibody, binds to TNF α on the surface of a cell and become internalized. For example, US patent application publication no. 2014/0294813, incorporated herein by reference, discloses anti-TNF α proteins that exhibit cellular internalization upon binding to cell surface human TNF α .

[0050] In certain embodiments, the antibodies bind to human and/or mouse TNF α .

[0051] The full-length amino acid sequence for membrane bound human TNF α is:

[0052] MSTESMIRDVELAEEALPKKTGGPQGSRRCLFLSLFSFLIVAGATTLFCLLHFGVIGPQRE
EFPRDLISLISPLAQAVRSSSRTPSDKPVAVHVVANPQAEGQLQWLNRRANALLANGVELRDNQLVVP
SEGLYLIYSQVLFKGGQCPSTHVLLTHTISRIVSYQTKVNLLSAIKSPCQRETPEGAEAKPWYEPIY
LGGVFQLEKGDRLSAEINRPDYLDFAESGQVYFGIIAL (SEQ ID NO:1). Soluble human TNF α
contains amino acids 77-233 of SEQ ID NO:1. The full-length amino acid sequence for membrane bound
murine TNF α is:

MSTESMIRDVELAEEALPQKMGGFQNSRRCLCLSLFSFLLVAGATTLFCLLNFGVIGPQRDEKFPNG
LPLISSMAQTLTLRSSSQNSSDKPVAVHVVANHQVEEQLEWLSQRANALLANGMDLKNQLVVPAD
GLYLVYSQVLFKGGQCPDYVLLTHTVSRFAISYQEKVNLLSAVKSPCPKDTPEGAELKPWYEPIYL
GGVFQLEKGDQLSAEVNLPKYLDFAESGQVYFGVIAL (SEQ ID NO:2). Soluble murine TNF α
contains amino acids 80-235 of SEQ ID NO:2.

[0053] In some embodiments, the anti-TNF α antibody binds to human TNF α .

[0054] In some embodiments, the anti-TNF α antibody binds to murine TNF α .

[0055] In certain embodiments, the anti-TNF α antibody has one or more of the following effects: neutralizes human TNF α cytotoxicity in an *in vitro* L929 assay with an IC₅₀ of 1X10⁻⁷ M or less; blocks the interaction of TNF α with p55 and p75 cell surface receptors; and/or lyses surface TNF expressing cells *in vitro* in the presence of complement.

[0056] In certain embodiments, the anti-TNF α antibody does not bind to TNF-beta.

[0057] Anti-TNF α antibodies include, for example, adalimumab, which is a recombinant human antibody. Amino acid sequences corresponding to the CDR and variable regions of adalimumab are described in U.S. Patent No. 6,258,562 in reference to antibody D2E7, *i.e.*, SEQ ID Nos: 1 to 8.

[0058] The international nonproprietary name (INN) adalimumab is provided in the WHO INN listing site: Year 2000, List 44 (*WHO Drug Information* (2000) Vol. 14(3)).

[0059] In certain embodiments, an anti-TNF α antibody comprises sequences of adalimumab, e.g., the complementarity-determining regions (CDRs), the variable heavy domain (VH), and/or the variable light domain (VL). Exemplary sequences are provided in Table 1.

Table 1: Exemplary adalimumab antibody region sequences

Antibody Region	Amino Acid Sequence
Heavy chain	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKG LEWVSAITWNSGHIDYADSVEGRFTISRDNKNSLYLQMNSLRAE DTAVYYCAKVSYLSTASSLDYWGQGTLVTVSSASTKGPSVFPLAP SSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKEPKSCDK THTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFCSCVMHEALHNYHTQKSLSLSPGK (SEQ ID NO: 3)
Light chain	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKPGKAPKL LIYAASTLQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNR APYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFY PREAKVQWKVDNALQSGNSQESVTEQDSKSTYLSLSTLTKAD YEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:4)
Heavy chain variable region	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKG LEWVSAITWNSGHIDYADSVEGRFTISRDNKNSLYLQMNSLRAE DTAVYYCAKVSYLSTASSLDYWGQGTLVTVSS (SEQ ID NO:5)
Light chain variable region	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKPGKAPKL LIYAASTLQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNR APYTFGQGTKVEIK (SEQ ID NO:6)
VH-CDR1	DYAMH (SEQ ID NO:7) or GFTFDDYAMH (SEQ ID NO:8)
VH-CDR2	AITWNSGHIDYADSVEG (SEQ ID NO:9)
VH-CDR3	VSYLSTASS (SEQ ID NO:10) or VSYLSTASSLDY (SEQ ID NO:11)
VL-CDR1	RASQGIRNYLA (SEQ ID NO:12)
VL-CDR2	AASTLQS (SEQ ID NO:13)
VL-CDR3	QRYNRAPYT (SEQ ID NO:14)

[0060] In certain embodiments, the anti-TNF α antibody comprises the CDRs of SEQ ID NOs: 3 and 4. In some embodiments, the CDRs comprise SEQ ID NOs: 7 or 8, 9, 10 or 11, 12, 13, and 14. In certain embodiments, the anti-TNF α antibody comprises the heavy chain of SEQ ID NO:3 and/or the light chain of SEQ ID NO:4.

[0061] The present disclosure further embraces variants and equivalents which are substantially homologous to anti-TNF α antibodies set forth herein. These can contain, for example, conservative substitution mutations, i.e., the substitution of one or more amino acids by similar amino acids. For example, conservative substitution refers to the substitution of an amino acid with another within the same general class such as, for example, one acidic amino acid with another acidic amino acid, one basic amino acid with another basic amino acid or one neutral amino acid by another neutral amino acid. What is intended by a conservative amino acid substitution is well known in the art.

[0062] The isolated anti-TNF α antibodies described herein can be produced by any suitable method known in the art. Such methods range from direct protein synthetic methods to constructing a DNA sequence encoding isolated polypeptide sequences and expressing those sequences in a suitable transformed host. In some embodiments, a DNA sequence is constructed using recombinant technology by isolating or synthesizing a DNA sequence encoding a wild-type protein of interest. Optionally, the sequence can be mutagenized by site-specific mutagenesis to provide functional analogs thereof. See, e.g., Zoeller et al., *Proc. Nat'l. Acad. Sci. USA* 81:5662-5066 (1984) and U.S. Pat. No. 4,588,585.

[0063] In some embodiments a DNA sequence encoding an antibody of interest would be constructed by chemical synthesis using an oligonucleotide synthesizer. Such oligonucleotides can be designed based on the amino acid sequence of the desired polypeptide and selecting those codons that are favored in the host cell in which the recombinant polypeptide of interest will be produced. Standard methods can be applied to synthesize an isolated polynucleotide sequence encoding an isolated polypeptide of interest.


[0064] In certain embodiments, recombinant expression vectors are used to amplify and express DNA encoding anti-TNF α antibodies. A wide variety of expression host/vector combinations can be employed. Useful expression vectors for eukaryotic hosts, include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Useful expression vectors for bacterial hosts include known bacterial plasmids, such as plasmids from *Escherichia coli*, including pCR 1, pBR322, pMB9 and their derivatives, wider host range plasmids, such as M13 and filamentous single-stranded DNA phages.

[0065] Suitable host cells for expression of anti-TNF α antibodies) include prokaryotes, yeast, insect or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include established cell lines of mammalian origin. Cell-free translation systems could also be employed. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, N.Y., 1985). Additional information regarding methods of protein production, including antibody production, can be found, e.g., in U.S. Patent Publication No. 2008/0187954, U.S. Patent Nos. 6,413,746 and 6,660,501, and International Patent Publication No. WO 04009823.

III. Immunoconjugates containing glucocorticoid receptor agonists

[0066] Immunoconjugates containing glucocorticoid receptor agonists are also provided. In some embodiments, an immunoconjugate binds to Fc gamma receptor. In some embodiments, an immunoconjugate is active in the GRE transmembrane TNF α reporter assay (as used herein the "GRE transmembrane TNF α reporter assay" refers to the assay used below Example 7 below). In some embodiments, an immunoconjugate shows reduced immunogenicity (reduced anti-drug immune response (ADA)) as compared to the protein in the immunoconjugate (e.g., the antibody) alone.

[0067] In one embodiment, disclosed herein is a compound having Formula **I-a**:



wherein:

A is an anti-tumor necrosis factor (TNF) α antibody, an anti-TNF α monoclonal antibody, or adalimumab;

L is a linker;


Q is a heterobifunctional group; or

Q is absent;

n is 1-10; and


SM is a monovalent radical of a glucocorticosteroid having any one of:

(1) Formula **II-a**:




II-a;

(2) Formula **II-b**:




II-b;

(3) Formula **II-c**:



II-c; or

(4) Formula **II-d**:




II-d.

[0068] In another embodiment, disclosed herein is a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, or **II-d**, wherein L


is a cleavable linker comprising a dipeptide or tripeptide, Q is a heterobifunctional group or Q is absent and n is 1-10. In particular, L comprises a dipeptide or tripeptide selected from the group consisting of alanine-alanine (Ala-Ala), glycine-glutamic acid (Gly-Glu), glutamic acid-alanine-alanine (Glu-Ala-Ala), and glycine-lysine (Gly-Lys).

[0069] In another embodiment, disclosed herein is a compound having Formula I-a, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae II-a, II-b, II-c, and II-d, wherein Q is a heterobifunctional group represented by:




wherein m is 0 or 1.

[0070] In another embodiment, m is 0, and Q is represented by:





[0071] In another embodiment, m is 1, and Q is represented by:



[0072] In another embodiment, disclosed herein is a compound having Formula I-a, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae II-a, II-b, II-c, and II-d, wherein L-Q- is any one of the chemical structures of Table 2:

Table 2





[0073] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a** wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein n is 2-8. In another embodiment, n is 1-5. In another embodiment, n is 2-5. In another embodiment, n is 1. In another embodiment, n is 2. In another embodiment n is 3. In another embodiment, n is 4. In another embodiment, n is 5. In another embodiment, n is 6. In another embodiment, n is 7. In another embodiment, n is 8.

[0074] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A is an antibody.

[0075] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein the antibody is murine, chimeric, humanized, or human.

[0076] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A competitively inhibits binding of an antibody selected from the group consisting of adalimumab, infliximab, certolizumab pegol, and golimumab to TNF α .

[0077] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A binds to the same TNF α epitope as an antibody selected from

the group consisting of adalimumab, infliximab, certolizumab pegol, afelimomab, nerelimomab, ozoralizumab, placulumab, and golimumab.

[0078] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A comprises the variable heavy chain CDR1, CDR2, and CDR3 sequences of SEQ ID NO:7 or 8, SEQ ID NO:9, and SEQ ID NO: 10 or 11, respectively, and the variable light chain CDR1, CDR2, and CDR3 sequences of SEQ ID NO:12, SEQ ID NO:13, and SEQ ID NO:14, respectively.

[0079] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A is an anti-TNF α antibody comprising a heavy chain variable region set forth as SEQ ID NO: 5 and a light chain variable region set forth as SEQ ID NO: 6.

[0080] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A is an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4,.

[0081] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A blocks the interaction of TNF α with p55 and p75 cell surface receptors.


[0082] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A lyses surface TNF expressing cells *in vitro* in the presence of complement.


[0083] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A is etanercept.



[0084] In another embodiment, disclosed herein is a compound having Formula **I-a**, e.g., a compound having Formula **I-a**, wherein SM is a monovalent radical of a glucocorticosteroid having any one of Formulae **II-a**, **II-b**, **II-c**, and **II-d**, wherein A is adalimumab.


[0085] In another embodiment, disclosed herein is a compound having Formula **I-a**, which is any one of the chemical structures of Table 3:

Table 3










herein n is 1-5 and A is adalimumab.

[0086] In another embodiment, disclosed herein is an antibody drug conjugate according to the formula:




wherein A is adalimumab and n is 4. As demonstrated in Example 7 below, this ADC (i.e., ADC4 below) demonstrates *in vitro* activity, stability in plasma, and minimal aggregation.

IV. Methods of making immunoconjugates and synthetic intermediates

[0087] The general synthesis of various immunoconjugates of the disclosure involves reacting a NH₂-functionalized small molecule (SM) of any of Formulae III-a, III-b, III-c, or III-d below with a linker


portion and functionalizing the resulting compound to give a bromoacetamide-functionalized intermediate. The bromoacetamide-functionalized intermediate is then reacted with HS-A, wherein HS-A is an antibody, for example adilumimab, having a limited number of reduced interchain disulfide bonds.

[0088] (1) Formula **III-a**:




III-a;

[0089] (2) Formula **III-b**:




III-b;

[0090] (3) Formula **III-c**:




III-c; or

[0091] (4) Formula **III-d**:



III-d.

[0092] In another embodiment, disclosed herein is a method of making a compound having Formula **IV-a**:



wherein:

A is adalimumab;


L is a linker;

n is 1-10; and

SM is a radical of a glucocorticosteroid having any one of Formulae **III-a- III-d**;

the method comprising:

a) conjugating a compound having Formula **V**:




with an anti-tumor necrosis factor (TNF) α protein or a protein; and

b) isolating the compound having Formula **IV-a**.

[0093] In some embodiments, the disclosed method further comprises purifying the compound of Formula **IV-a**. In certain embodiments, anionic exchange chromatography (AEC) is used, which (due to charged moieties on the peptide portion of the linker in some embodiments and/or the phosphate group on the SM in some embodiments) can provide efficient separation of different DAR species.

[0094] In some embodiments, the disclosed method requires fewer synthetic steps than methods relying on standard maleimide-based linker chemistry. In particular, methods relying on standard maleimide-based linker chemistry can require a subsequent succinimide ring-opening hydrolysis step that is performed after purification of the appropriate DAR species. As such, in certain embodiments, the disclosed method significantly shortens the conjugation protocol with respect to standard maleimide-based linker chemistry.

[0095] In another embodiment, disclosed herein is a method of making a compound having Formula **VI-a**:




wherein:

A is adalimumab; and

n is 1-10,

the method comprising:

a) conjugating a compound of Formula **VII-a**:



VII-a,

with partially-reduced adalimumab; and

b) isolating, e.g., by chromatography, the compound having Formula **VI-a**.

[0096] In another embodiment, disclosed herein is a method of making a compound having Formula **IV-a** or Formula **VI-a**, wherein n is 1-7. In another embodiment, n is 1-5. In another embodiment, n is 2-4. In another embodiment, n is 1. In another embodiment, n is 2. In another embodiment, n is 3. In another embodiment, n is 4. In another embodiment, n is 5. In another embodiment, n is 6. In another embodiment, n is 7. In another embodiment, n is 8.

[0097] In another embodiment, disclosed herein is a compound having Formula **IV-a** or **VI-a**, wherein:
A is adalimumab; and
n is 1-10.

[0098] In another embodiment, disclosed herein is a compound having Formula **IV-a** or **VI-a**, wherein n is 1-7. In another embodiment, n is 1-5. In another embodiment, n is 2-4. In another embodiment, n is 1. In another embodiment, n is 2. In another embodiment, n is 3. In another embodiment, n is 4. In another embodiment, n is 5. In another embodiment, n is 6. In another embodiment, n is 7. In another embodiment, n is 8.

[0099] Also provided herein are synthetic intermediates that are useful for the preparation of immunoconjugates.

[00100] In one embodiment, the synthetic intermediate disclosed herein is a compound having any one of Formulae **V** or **VII-a**.

VI. Methods of use and pharmaceutical compositions

[00101] Provided herein are conjugates having Formula **I-a** (e.g., having the formulas shown in Table 3) that can be used *in vitro* or *in vivo*. Accordingly, also provided are compositions, e.g., pharmaceutical compositions for certain *in vivo* uses, comprising a conjugate or a glucocorticoid receptor agonist having the desired degree of purity in a physiologically acceptable carrier, excipient or stabilizer (Remington's

Pharmaceutical Sciences (1990) Mack Publishing Co., Easton, PA). Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations employed.

[00102] The compositions (e.g., pharmaceutical compositions) to be used for *in vivo* administration can be sterile, which can be accomplished by filtration through, e.g., sterile filtration membranes. The compositions (e.g., pharmaceutical compositions) to be used for *in vivo* administration can comprise a preservative.

[00103] Antibody drug conjugates and/or pharmaceutical compositions comprising antibody drug conjugates described herein can be useful in lysing a cell expressing surface TNF α (*in vitro* or *in vivo*) and/or for the treatment of diseases or disorders characterized by increased TNF α (e.g., increased TNF α in synovial fluid). In some embodiments, the antibody drug conjugates and/or compositions are useful in inhibiting cytokine release (*in vitro* or *in vivo*) and/or for the treatment of autoimmune or inflammatory diseases. In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of Crohn's disease (e.g., moderate to severely active Crohn's disease involving the ileum and/or the ascending colon and/or the maintenance of clinical remission of moderate to severely active Crohn's disease involving the ileum and/or the ascending colon for up to 3 months). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of ulcerative colitis (e.g., for the induction of remission in patients with active, moderate to severe ulcerative colitis). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of rheumatoid arthritis (RA). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of juvenile idiopathic arthritis (JA). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of psoriatic arthritis (PsA). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of a spondyloarthropathy such as ankylosing spondylitis (AS) or axial spondyloarthritis (axSpA). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of adult Crohn's disease (CD). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of pediatric Crohn's disease. In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of ulcerative colitis (UC). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of plaque psoriasis (Ps). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of hidradenitis suppurativa (HS). In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of uveitis. In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of Behcets disease. In some embodiments, the antibody drug conjugates and/or compositions are used for the treatment of psoriasis, including plaque psoriasis. Some embodiments comprise use of drug conjugates and/or pharmaceutical compositions for the preparation of a medicament for treating the diseases or disorders described herein.

[00104] Some embodiments comprise methods of delivering a glucocorticoid receptor agonist to a TNF α -expressing cell. Such methods can include a step of contacting a TNF α -expressing cell with an

antibody drug conjugate as described herein. Some embodiments comprise an *in vitro* method of delivering a glucocorticoid receptor agonist to a TNF α -expressing cell.

[00105] Also provided are methods of determining anti-inflammatory activity of an antibody drug conjugate. Such methods can include a step of contacting a TNF α -expressing cell with an antibody drug conjugate as described herein. Some embodiments comprise contacting a TNF α -expressing cell with an antibody drug conjugate as described herein and determining reduced release of pro-inflammatory cytokines from the cell as compared to a control cell. Some embodiments comprise an *in vitro* method of determining anti-inflammatory activity of an antibody drug conjugate.

[00106] Some embodiments comprise screening methods (*e.g. in vitro* methods) that include contacting, directly or indirectly, cells (*e.g.*, TNF α -expressing cells) with an antibody drug conjugate and determining if the antibody drug conjugate modulates an activity or function of the cells, as reflected for example by changes in cell morphology or viability, expression of a marker, differentiation or de-differentiation, cell respiration, mitochondrial activity, membrane integrity, maturation, proliferation, viability, apoptosis or cell death. One example of a direct interaction is physical interaction, while an indirect interaction includes, for example, the action of a composition upon an intermediary molecule that, in turn, acts upon the referenced entity (*e.g.*, cell or cell culture).

VII. Articles of Manufacture

[00107] The disclosure also includes pharmaceutical packs and kits comprising one or more containers, wherein a container can comprise one or more doses of an antibody drug conjugate or composition as described herein. In certain embodiments, the pack or kit contains a unit dosage, meaning a predetermined amount of a composition or antibody drug conjugate, with or without one or more additional agents.

[00108] The kit can comprise one or multiple containers and a label or package insert in, on or associated with the container(s), indicating that the enclosed composition is used for treating the disease condition of choice. Suitable containers include, for example, bottles, vials, syringes, etc. The containers can be formed from a variety of materials such as glass or plastic. The container(s) can comprise a sterile access port, for example, the container may be an intravenous solution bag or a vial having a stopper that can be pierced by a hypodermic injection needle.

[00109] In some embodiments the kit can contain a means by which to administer the antibody and any optional components to a patient, *e.g.*, one or more needles or syringes (pre-filled or empty), an eye dropper, pipette, or other such like apparatus, from which the formulation may be injected or introduced into the subject or applied to a diseased area of the body. The kits of the disclosure will also typically include a means for containing the vials, or the like, and other components in close confinement for commercial sale, such as, *e.g.*, blow-molded plastic containers into which the desired vials and other apparatus are placed and retained.

Examples

[00110] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this disclosure.

[00111] Starting materials are commercially available, may be prepared by the procedures described herein, by literature procedures, or by procedures that would be well known to one skilled in the art of organic chemistry. Reagent/reactant names given are as named on the commercial bottle or as generated by IUPAC conventions, CambridgeSoft® ChemDraw Ultra 12.0, CambridgeSoft® Chemistry E-Notebook 11, or AutoNom 2000. It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this disclosure.

Analytical Methods for Compound Synthesis and Characterization

[00112] Analytical data is included within the procedures below, in the illustrations of the general procedures, or in the tables of examples. Unless otherwise stated, all ¹H and ¹³C NMR data were collected on a Varian Mercury Plus 400 MHz or a Bruker AVIII 300 MHz instrument; chemical shifts are quoted in parts per million (ppm). HPLC analytical data are either detailed within the experimental or referenced to the table of LC/MS and HPLC conditions, using the method provided in Table 4.

Table 4

Method	Conditions
a	The gradient was 1-90% B in 3.4 min, 90-100% B in 0.45 min, 100-1% B in 0.01 min, and then held at 1% B for 0.65 min (0.8 mL/min flow rate). Mobile phase A was 0.0375% TFA in H ₂ O, mobile phase B was 0.018% TFA in MeCN. The column used for the chromatography was a 2.0 X 50 mm Phenomenex Luna-C18 column (5 μm particles). Detection methods are DAD and ELSD as well as positive electrospray ionization.


[00113] Abbreviations used in the examples that follow are:

ACTH	Adrenocorticotropin Hormone	HIC	Hydrophobic Interaction Chromatography
BrAc	Bromo acetamide	HPLC	High performance liquid chromatography
CV	Column volumes	LCMS	Liquid chromatography mass spectrometry
DAD	Diode array	MeCN	Acetonitrile
DAR	Drug to antibody ratio	MEM	Minimal Essential Media
DBP	Dibutyl phthalate	MeOH	Methanol
DCM	Dichloromethane	MS	Mass spectrometry

DMA	Dimethyl acetamide	NEAA	Non-essential amino acids
DMF	Dimethyl formamide	NMR	Nuclear magnetic resonance
DMSO	Dimethyl sulfoxide	PBS	Phosphate buffer saline
DTT	Dithiothreitol	PE	Petroleum ether
EEDQ	2-Ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline	P1NP	Procollagen type 1 amino-terminal propeptide
ELSD	Evaporative light scattering detector	rt	Room temperature
Eq	Equivalents	RPMI	Roswell Park Memorial Institute
EtOAc	Ethyl acetate	SEC	Size exclusion chromatography
FBS	Fetal Bovine Serum	TCEP	(tris(2-carboxyethyl)phosphine)
Fmoc	9-Fluorenylmethoxycarbonyl	TFA	Trifluoro acetic acid


[00114] Example 1: Synthesis of (2S,6aS,6bR,7S,8aS,8bS,10S,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-2,6b-difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one

[00115] Step 1: Synthesis of 4-(bromomethyl)benzaldehyde



[00116] Diisobutylaluminum hydride (153 mL, 153 mmol, 1 M in toluene) was added drop-wise to a 0 °C solution of 4-(bromomethyl)benzonitrile (20 g, 102 mmol) in toluene (400 mL) over 1 hour. Two additional vials were set up as described above. All three reaction mixtures were combined. To the mixture was added 10% aqueous HCl (1.5 L). The mixture was extracted with DCM (3 X 500 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluted with PE:EtOAc = 10:1) to obtain the title compound (50 g, yield 82%) as white solid. ¹H NMR (400MHz, CDCl₃) δ 10.02 (s, 1H), 7.91 - 7.82 (m, 2H), 7.56 (d, J=7.9 Hz, 2H), 4.55 - 4.45 (m, 2H).


[00117] Step 2: Synthesis of 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline



[00118] To a solution of 3-bromoaniline (40 g, 233 mmol) in 1,4-dioxane (480 mL) was added 4,4,4',5,5,5'-tetramethyl-2,2'-bi(1,3,2-dioxaborolane) (94 g, 372 mmol), potassium acetate (45.6 g, 465 mmol), 2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl (8.07 g, 13.95 mmol), and tris(dibenzylideneacetone)dipalladium(0) (8.52 g, 9.30 mmol). The resulting mixture was heated at 80°C for


4 hours under nitrogen. An additional vial was set up as described above. The two reaction mixtures were combined and concentrated and the residue was purified by column chromatography on silica gel (eluted with PE:EtOAc = 10:1) to obtain the title compound (60 g, yield 55.4%) as a light yellow solid. ¹H NMR (400MHz, CDCl₃) δ 7.23 - 7.13 (m, 3H), 6.80 (d, *J*=7.5 Hz, 1H), 3.82 - 3.38 (m, 2H), 1.34 (s, 12H).

[00119] Step 3: Synthesis of tert-butyl (3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl) carbamate




[00120] 3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (Example 1, Step 2) (30 g, 137 mmol) and di-tert-butyl dicarbonate (38.9 g, 178 mmol) were mixed in toluene (600 mL) at 100 °C for 24 hours. Another vial was set up as described above. The two reaction mixtures were combined and the mixture was evaporated, dissolved in EtOAc (1.5 L), washed with 0.1 N HCl (3 X 2 L) and brine (3 L), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give the title compound (50 g, yield 57%) as a red solid. ¹H NMR (400MHz, CDCl₃) δ 7.63 (br m, 2H), 7.48 (d, *J*=7.1 Hz, 1H), 7.37 - 7.28 (m, 1H), 1.52 (s, 9H), 1.34 (s, 12H).

[00121] Step 4: Synthesis of tert-butyl (3-(4-formylbenzyl)phenyl)carbamate




[00122] A mixture of 4-(bromomethyl)benzaldehyde (Example 1, Step 1) (24.94 g, 125 mmol), 1,1'-bis(diphenylphosphino) ferrocenedichloro palladium(II) DCM complex (13.75 g, 18.80 mmol), tert-butyl (3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl) carbamate (from Example 1, Step 3) (20 g, 62.7 mmol) and potassium carbonate (43.3 g, 313 mmol) in tetrahydrofuran (400 mL) was heated to 80 °C for 12 hours. Another vial was set up as described above. The two reaction mixtures were combined and diluted with water (500 mL). The aqueous mixture was extracted with EtOAc (3 X 500 mL). The organic layers were combined and dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluted with PE:EtOAc = 10:1) to obtain the title compound (15 g, yield 38.4%) as white solid. ¹H NMR (400MHz, CDCl₃) δ 9.95 (s, 1H), 7.78 (d, *J*=7.9 Hz, 2H), 7.33 (d, *J*=7.9 Hz, 2H), 7.27 - 7.13 (m, 3H), 6.82 (d, *J*=7.1 Hz, 1H), 6.47 (br. s., 1H), 4.00 (s, 2H), 1.48 (s, 9H).

[00123] Step 5: Synthesis of (6*S*,8*S*,9*R*,10*S*,11*S*,13*S*,14*S*,16*R*,17*S*)-6,9-difluoro-11,16,17-trihydroxy-17-(2-hydroxyacetyl)-10,13-dimethyl-6,7,8,9,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[*a*]phenanthren-3-one



[00124] (2S,6aS,6bR,7S,8aS,8bS,11aR,12aS,12bS)-2,6b-Difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a,10,10-tetramethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one (20 g, 44.2 mmol) was suspended in 40% aqueous HBF₄ (440 mL) and the mixture was stirred at 25 °C for 48 hours. After the reaction was complete, 2 L of water was added and the solid was collected by filtration. This solid was washed with water (1 L) and then MeOH (200 mL) to give the title compound (11 g, yield 60.3%) as a white solid. ¹H NMR (400MHz, DMSO-d₆) δ 7.25 (d, *J*=10.1 Hz, 1H), 6.28 (d, *J*=10.1 Hz, 1H), 6.10 (s, 1H), 5.73 - 5.50 (m, 1H), 5.39 (br. s., 1H), 4.85 - 4.60 (m, 2H), 4.50 (d, *J*=19.4 Hz, 1H), 4.20 - 4.04 (m, 2H), 2.46 - 2.06 (m, 6H), 1.87 - 1.75 (m, 1H), 1.56 - 1.30 (m, 6H), 0.83 (s, 3H).

[00125] Step 6: Synthesis of (2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-2,6b-difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one.




[00126] A suspension of (6S,8S,9R,10S,11S,13S,14S,16R,17S)-6,9-difluoro-11,16,17-trihydroxy-17-(2-hydroxyacetyl)-10,13-dimethyl-6,7,8,9,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthren-3-one (Example 1, Step 5) (4.4 g, 10.67 mmol) and MgSO₄ (6.42 g, 53.3 mmol) in MeCN (100 mL) was stirred at 20 °C for 1 hour. A solution of tert-butyl (3-(4-formylbenzyl)phenyl)carbamate (Example 1, Step 4) (3.65 g, 11.74 mmol) in MeCN (100 mL) was added in one portion. Trifluoromethanesulfonic acid (9.01 mL, 53.3 mmol) was added dropwise while maintaining an internal temperature below 25 °C using an ice bath. After the addition, the mixture was stirred at 20 °C for 2 hours. Three additional vials were set up as described above. All four reaction mixtures were combined and concentrated and the residue was purified by Prep HPLC to give the title compound (4.5 g, yield 14.2%) as yellow solid. LCMS (Method a, Table 4) R_t = 2.65 min; MS *m/z* = 606.2 (M+H)⁺; ¹H NMR (400MHz, DMSO-d₆) δ 7.44 - 7.17 (m, 5H), 6.89 (t, *J*=7.7 Hz, 1H), 6.44 - 6.25 (m, 4H), 6.13 (br. s., 1H), 5.79 - 5.52 (m, 2H), 5.44 (s, 1H), 5.17 - 4.89 (m, 3H), 4.51 (d, *J*=19.4 Hz, 1H), 4.25 - 4.05 (m, 2H), 3.73 (s, 2H), 3.17 (br. s., 1H), 2.75 - 2.55 (m, 1H), 2.36 - 1.97 (m, 3H), 1.76 - 1.64 (m, 3H), 1.59 - 1.39 (m, 4H), 0.94 - 0.78 (m, 3H). Prep HPLC Method: Instrument: Gilson 281 semi-preparative HPLC system; Mobile phase: A:

Formic Acid/H₂O=0.01% v/v; B: MeCN; Column: Luna C18 150*25 5 micron; Flow rate: 25 mL/min;
Monitor wavelength: 220 and 254nm.

Time	0.0	10.5	10.6	10.7	13.7	13.8	15.0
B %	15	35	35	100	100	10	10


[00127] Example 2: Synthesis of (6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one.



[00128] Example 2 was synthesized in a similar procedure to Example 1 using (8S,9S,10R,11S,13S,14S,16R,17S)-11,16,17-trihydroxy-17-(2-hydroxyacetyl)-10,13-dimethyl-6,7,8,9,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthren-3-one.

¹H NMR (400MHz, DMSO-d₆) δ 7.36 (d, *J*=7.9 Hz, 2H), 7.31 (d, *J*=10.1 Hz, 1H), 7.20 (d, *J*=7.9 Hz, 2H), 6.89 (t, *J*=7.9 Hz, 1H), 6.39 - 6.28 (m, 3H), 6.16 (dd, *J*=1.5, 9.9 Hz, 1H), 5.93 (s, 1H), 5.39 (s, 1H), 5.08 (t, *J*=5.7 Hz, 1H), 4.98 - 4.87 (m, 3H), 4.78 (d, *J*=3.1 Hz, 1H), 4.49 (dd, *J*=6.2, 19.4 Hz, 1H), 4.29 (br. s., 1H), 4.17 (dd, *J*=5.5, 19.6 Hz, 1H), 3.74 (s, 2H), 2.61 - 2.53 (m, 1H), 2.36 - 2.26 (m, 1H), 2.11 (d, *J*=11.0 Hz, 1H), 2.07 (s, 1H), 2.02 (d, *J*=12.8 Hz, 1H), 1.83 - 1.54 (m, 5H), 1.39 (s, 3H), 1.16 - 0.96 (m, 2H), 0.85 (s, 3H). LCMS (Method a, Table 4) R_t = 2.365 min; m/z = 570.2 (M+H)⁺.

[00129] Example 3: Synthesis of (6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-6b-fluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one.




[00130] Example 3 was synthesized in a similar procedure to Example 1 using (8S,9R,10S,11S,13S,14S,16R,17S)-9-fluoro-11,16,17-trihydroxy-17-(2-hydroxyacetyl)-10,13-dimethyl-6,7,8,9,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthren-3-one.

[00131] ¹H NMR (400MHz, DMSO-d₆) δ 7.37 - 7.26 (m, 3H), 7.21 (d, *J*=7.9 Hz, 2H), 6.89 (t, *J*=7.7 Hz, 1H), 6.43 - 6.30 (m, 3H), 6.23 (d, *J*=10.1 Hz, 1H), 6.04 (s, 1H), 5.75 (s, 1H), 5.44 (s, 2H), 5.09 (t, *J*=5.7 Hz, 1H), 4.93 (br. s., 3H), 4.50 (dd, *J*=6.2, 19.4 Hz, 1H), 4.28 - 4.09 (m, 2H), 3.74 (s, 2H), 2.73 - 2.54 (m, 2H), 2.35 (d, *J*=13.2 Hz, 1H), 2.25 - 2.12 (m, 1H), 2.05 (d, *J*=15.0 Hz, 1H), 1.92 - 1.77 (m, 1H), 1.74 - 1.58 (m,

3H), 1.50 (s, 3H), 1.45 - 1.30 (m, 1H), 0.87 (s, 3H). LCMS (Method a, Table 4) $R_t = 2.68$ min; $m/z = 588.1$ (M+H)⁺

[00132] Example 4: Synthesis of (S)-4-(2-(2-bromoacetamido)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-7-hydroxy-6a,8a-dimethyl-4-oxo-8b-(2-(phosphonoxy)acetyl)-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoic acid

[00133] Step 1: Synthesis of (S)-2-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-(tert-butoxy)-5-oxopentanoic acid.




[00134] A mixture of 2-chlorotrityl chloride resin (30 g, 92 mmol), triethylamine (46.4 g, 458 mmol) and (S)-2-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)-5-(tert-butoxy)-5-oxopentanoic acid (25.5 g, 60 mmol) in dry DCM (200 mL) was bubbled with N₂ at 20 °C for 8 hours. The mixture was filtered and the resin was washed with DCM (2 X 200 mL), MeOH (2 X 200 mL), and DMF (2 X 200 mL). The resin was added a solution of piperidine:DMF (1:4, 400 mL) and the mixture was bubbled with N₂ for 8 minutes and then filtered. This operation was repeated five times to give complete removal of the Fmoc protecting group. The resin was washed with DMF (5 X 500 mL) to afford resin bound (S)-2-amino-5-(tert-butoxy)-5-oxopentanoic acid. A mixture of 2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetic acid (13.38 g, 45.0 mmol), *N,N*-diisopropylethylamine (7.86 mL, 45 mmol), hydroxybenzotriazole (6.89 g, 45 mmol), 2-(6-chloro-1H-benzo[d][1,2,3]triazol-1-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate (V) (18.62 g, 45.0 mmol) in DMF (200 mL) was stirred at 20 °C for 30 min. To the mixture was added the resin bound (S)-2-amino-5-(tert-butoxy)-5-oxopentanoic acid and the resulting mixture was bubbled with N₂ at 25 °C for 1.5 hours. The mixture was filtered and the resin was washed with DMF (4 X 500 mL), and DCM (2 X 500 mL). To the mixture was added 1% TFA / DCM (5 X 500 mL) and bubbled with N₂ for 5 minutes. The mixture was filtered and the filtrate was added to saturated solution of NaHCO₃ (200 mL) directly. The combined mixture was separated, and the organic phase was washed with saturated citric acid water solution (4 X 400 mL) and brine (2 X 300 mL). The final organic solution was dried over Na₂SO₄ (20 g), filtered, concentrated under reduced pressure to afford the title compound (10 g, yield 20%) as a light yellow solid.

[00135] ¹H NMR: (CDCl₃, 400 MHz) $\delta = 7.75$ (d, $J = 7.5$ Hz, 2H), 7.59 (br d, $J = 7.5$ Hz, 2H), 7.41 - 7.36 (m, 2H), 7.30 (t, $J = 7.0$ Hz, 2H), 5.82 (br s, 1H), 4.57 (br d, $J = 4.8$ Hz, 1H), 4.38 (br d, $J = 7.5$ Hz, 2H), 4.27 - 4.15 (m, 1H), 4.06 - 3.83 (m, 2H), 2.50 - 2.29 (m, 2H), 2.26 - 2.13 (m, 1H), 2.06 - 2.02 (m, 1H), 1.43 (s, 9H).

[00136] Step 2: Synthesis of tert-butyl (S)-4-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-7-hydroxy-


8b-(2-hydroxyacetyl)-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate



[00137] To a solution of (S)-2-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-(tert-butoxy)-5-oxopentanoic acid (Example 4, step 1) (424 mg, 0.878 mmol) in DMF (3.5 mL) was added (6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one (Example 2) (500 mg, 0.878 mmol) and triethylamine (0.3 mL, 2.63 mmol) at 25 °C. The solution was cooled to 0 °C and then 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (1.12 g, 1.755 mmol) was added. The reaction mixture was stirred for 12 hours at 25 °C. LCMS showed the reaction was complete. Fourteen additional vials were set up as described above. All fifteen reaction mixtures were combined. The mixture was purified by reverse phase column to afford the title compound (5 g, yield 38.4%) as a yellow solid. Reverse phase column method: Instrument: Shimadzu LC-8A preparative HPLC; Column: Phenomenex Luna C18 200*40 mm*10 mm; Mobile phase: A for H₂O (0.05% TFA) and B for MeCN; Gradient: B from 30% to 100% in 30 min; Flow rate: 60 mL/min; Wavelength: 220 & 254 nm.

[00138] LCMS (Method a, Table 4) R_t = 1.34 min; m/z 1016.6 (M+H-18)⁺.


[00139] **Step 3:** Synthesis of tert-butyl (S)-4-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-((3-(4-(((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate.



[00140] To a solution of tert-butyl (S)-4-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-((3-(4-(((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate (Example 4, step 2) (400 mg, 0.387 mmol) in DMF (2.5 mL) was added 1H-tetrazole (271 mg, 3.87 mmol) and di-tert-butyl diethylphosphoramidite (1.16 g, 4.64 mmol). The

reaction was stirred at rt for 2.5 hours then cooled to 0 °C. Hydrogen peroxide (241 mg, 2.127 mmol) was added to the resulting mixture allowed to warm to rt and stirred for 1 hour after which time LCMS showed the reaction was complete. Eleven additional vials were set up as described above. All twelve reaction mixtures were combined. The mixture was purified by reverse phase column to afford the title compound (4.4 g, yield 64.2%) as a yellow solid. Reverse phase column method: Instrument: Shimadzu LC-8A preparative HPLC; Column: Phenomenex Luna C18 200*40mm*10 mm; Mobile phase: A for H₂O and B for MeCN; Gradient: B from 50% to 100% in 30min; Flow rate: 60 mL/min; Wavelength: 220 & 254 nm. LCMS (Method a, Table 4) R_t = 1.41 min; m/z 1226.7 (M+H)⁺.


[00141] Step 4: Synthesis of tert-butyl (S)-4-(2-aminoacetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate.



[00142] To a solution of tert-butyl (S)-4-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate (Example 4, Step 3) (1.1 g, 0.897 mmol) in MeCN (6 mL) was added piperidine (0.75 mL, 7.58 mmol) at 25°C. The reaction was stirred at rt for 20 minutes after which time LCMS showed the reaction was complete. Three additional vials were set up as described above. All four reaction mixtures were combined. The mixture was concentrated to afford a residue, which was treated with PE (10 mL) under stirring for 2 hours. The resulting solid was collected by filtration, and dried under reduced pressure to afford the title compound (3.8 g, yield 90%) as a yellow solid.

[00143] LCMS (Method a, Table 4) R_t = 1.16 min; m/z 1004.6 (M+H)⁺.


[00144] Step 5: Synthesis of tert-butyl (S)-4-(2-(2-bromoacetamido)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate.



[00145] To a solution of 2-bromoacetic acid (97 mg, 0.697 mmol) in DMF (2.5 mL) was added EEDQ (172 mg, 0.697 mmol) at rt. The mixture was stirred at rt for 1 hour. tert-Butyl (S)-4-(2-aminoacetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate (Example 4, Step 4) (350 mg, 0.349 mmol) was added and the resulting solution was stirred for 2.5 hours after which time LCMS showed the reaction was complete. Seven additional vials were set up as described above. All eight reaction mixtures were combined. The reaction was diluted with DCM (100 mL), washed with aqueous HBr (1 M, 2 X 80 mL), aqueous NaHCO₃ (60 mL), and brine (60 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (2 g, yield 63.7%) as yellow oil.

[00146] LCMS (Method a, Table 4) R_t = 1.30 min; m/z 1124.2, 1125.9 (M+H)⁺.

[00147] Step 6: Synthesis of (S)-4-(2-(2-bromoacetamido)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-7-hydroxy-6a,8a-dimethyl-4-oxo-8b-(2-(phosphonoxy)acetyl)-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoic acid




[00148] To a solution of tert-butyl (S)-4-(2-(2-bromoacetamido)acetamido)-5-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-5-oxopentanoate (Example 4, Step 5) (2 g, 1.778 mmol) in DCM (16 mL) was added TFA (8 mL, 104 mmol) and the resulting mixture was stirred at rt for 40 minutes after which time LCMS showed the reaction was complete. The solvent was removed under reduced pressure. The resulting residue was purified by Prep HPLC. The mobile phase was lyophilized directly to afford the title compound (640 mg, yield 35.3%) as yellow solid. Prep HPLC method: Instrument: Shimadzu LC-8A preparative HPLC; Column: Phenomenex Luna C18 200*40 mm*10 μm; Mobile phase: A for H₂O (0.09% TFA) and B for MeCN; Gradient: B from 30% to 40% in 20 min; Flow rate: 60mL/min; Wavelength: 220 & 254 nm.

[00149] ^1H NMR: (DMSO- d_6 , 400 MHz) δ = 9.88 (s, 1H), 8.52 (s, 1H), 8.24 (br d, J = 8.4 Hz, 1H), 7.46 (br d, J = 7.9 Hz, 1H), 7.42 (s, 1H), 7.36 (br d, J = 7.9 Hz, 2H), 7.30 (br d, J = 9.7 Hz, 1H), 7.23 - 7.17 (m, 3H), 6.90 (br d, J = 6.8 Hz, 1H), 6.16 (br d, J = 10.4 Hz, 1H), 5.93 (s, 1H), 5.47 (s, 1H), 4.96 - 4.85 (m, 3H), 4.58 (br dd, J = 7.9, 18.7 Hz, 1H), 4.38 (br d, J = 5.3 Hz, 1H), 4.29 (br s, 1H), 3.93 (s, 2H), 3.89 (s, 2H), 3.80 (br s, 2H), 2.30 - 2.22 (m, 2H), 2.16 - 1.91 (m, 4H), 1.85 - 1.62 (m, 6H), 1.39 (s, 3H), 1.00 (br s, 2H), 0.87 (s, 3H). LCMS (Method a, Table 4) R_t = 2.86 min; m/z 956.0, 958.0 (M+H) $^+$.

[00150] **Example 5:** Synthesis of 2-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-((S)-6-amino-2-(2-(2-bromoacetamido)acetamido)hexanamido)benzyl)phenyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-1,2,4,6a,6b,7,8,8a,11a,12,12a,12b-dodecahydro-8bH-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-8b-yl)-2-oxoethyl dihydrogen phosphate.

[00151] **Step 1:** Synthesis of tert-butyl ((S)-5-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-2,6b-difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate.




[00152] To a solution of N^2 -(((9H-fluoren-9-yl)methoxy)carbonyl)glycyl)- N^6 -(tert-butoxycarbonyl)-L-lysine (5.58 g, 8.26 mmol) in DMF (60 mL) at 0 °C was added 2,4,6-tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (10.51 g, 16.51 mmol) and triethylamine (3.45 mL, 24.77 mmol). The resulting mixture was stirred at rt for 1 hour (2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-aminobenzyl)phenyl)-2,6b-difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-1,2,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-4H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-4-one (Example 1, Step 6) (5 g, 8.26 mmol) was added. The resulting mixture was stirred for 5 hours at rt after which time LCMS showed the reaction was complete. Six additional vials were set up as described above. All seven reaction mixtures were combined. The reaction was purified by reverse phase column to afford the title compound (24 g, yield 24.62%) as a white solid. Reverse phase column method: Instrument: Shimadzu LC-8A prep HPLC; Column: Phenomenex Luna C18 200*40 mm*10 mm; Mobile phase: A for H₂O (0.05% TFA) and B for MeCN; Gradient: B from 30% to 100% in 30 min; Flow rate: 60 mL/min; Wavelength: 220 & 254 nm.

[00153] LCMS (Method a, Table 4) R_t = 1.29 min; m/z 1095.6 (M+H-18) $^+$.

[00154] **Step 2:** Synthesis of tert-butyl ((S)-5-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-


((di-tert-butoxyphosphoryl)oxy)acetyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate.



[00155] To a solution of tert-butyl ((S)-5-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-2,6b-difluoro-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate (Example 5, Step 1) (3 g, 2.69 mmol) in DMF (30 mL) was added 1H-tetrazole (1.888 g, 26.9 mmol) and di-tert-butyl diethylphosphoramidite (8.06 g, 32.3 mmol) and the reaction was stirred at rt for 3.5 hours. Hydrogen peroxide (224 mg, 1.97 mmol) was added to the reaction and stirred for 0.5 hours after which time LCMS showed the reaction was complete. Six additional vials were set up as described above. All seven reaction mixtures were combined. The reaction was purified by reverse phase column to afford the title compound (10 g, purity: 78%, yield 37.1%) as a white solid. Reverse phase column method: Instrument: Shimadzu LC-8A prep HPLC; Column: Phenomenex Luna C18 200*40mm*10 mm; Mobile phase: A for H₂O and B for MeCN; Gradient: B from 50% to 100% in 30 min; Flow rate: 60 mL/min; Wavelength: 220 & 254 nm.

[00156] LCMS (Method a, Table 4) R_t = 1.42 min; m/z 1305.7 (M+H)⁺.

[00157] Step 3: Synthesis of tert-butyl ((S)-5-(2-aminoacetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate.




[00158] To a solution of tert-butyl ((S)-5-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)acetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate (Example 5, Step 2) (2.5 g, 1.969 mmol) in MeCN (10 mL) was added piperidine (2 mL, 1.969 mmol) and the reaction stirred at rt for 1 hour after which time LCMS showed the reaction was complete. Three additional vials were set up as described above. All four reaction mixtures were combined. The reaction was concentrated

to afford a crude product, which was stirred in PE (30 mL) for 2 hours. The resulting solid was collected by filtration, and dried under reduced pressure to afford the title compound (7 g, purity: 83%, yield 70.4%) as a yellow solid.

[00159] LCMS (Method a, Table 4) $R_t = 1.17$ min; m/z 1083.5 (M+H)⁺.


[00160] **Step 4:** Synthesis of tert-butyl ((S)-5-(2-(2-bromoacetamido)acetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate.



[00161] To a solution of 2-bromoacetic acid (0.929 g, 6.68 mmol) in DMF (35 mL) was added 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (1.653 g, 6.68 mmol) and the resulting mixture stirred at rt for 1 hour. The product from Example 5, Step 3 (3.5 g, 3.34 mmol) was added and the resulting mixture stirred at rt for 2 hours. LCMS showed the reaction was completed. The reaction was diluted with DCM (100 mL), washed with aqueous HBr (1 M, 2 X 80 mL), aqueous NaHCO₃ (60 mL) and brine (60 mL). The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (2 g, yield 51.2%) as yellow oil.

[00162] LCMS (Method a, Table 4) $R_t = 1.32$ min; m/z 1205.5 (M+H)⁺.

[00163] **Step 5:** Synthesis of 2-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-10-(4-(3-((S)-6-amino-2-(2-(2-bromoacetamido)acetamido)hexanamido)benzyl)phenyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-1,2,4,6a,6b,7,8,8a,11a,12,12a,12b-dodecahydro-8bH-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-8b-yl)-2-oxoethyl dihydrogen phosphate.




[00164] To a solution of tert-butyl ((S)-5-(2-aminoacetamido)-6-((3-(4-((2S,6aS,6bR,7S,8aS,8bS,10R,11aR,12aS,12bS)-8b-(2-((di-tert-butoxyphosphoryl)oxy)acetyl)-2,6b-difluoro-7-hydroxy-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-6-oxohexyl)carbamate (Example 5, Step 3) (2 g, 1.661 mmol) in DCM (10 mL) was added TFA (5 mL, 64.9 mmol) and the reaction stirred at rt for 40 min after which time LCMS showed the reaction was complete. The solvent was removed under reduced pressure and the crude product purified by Prep HPLC. The mobile phase was lyophilized directly

to afford the title compound (550 mg, purity: 96.9%, yield 32.3%) as an off-white solid. Prep-HPLC method: Instrument: Shimadzu LC-8A prep HPLC; Column: Phenomenex Luna C18 200*40 mm*10 mm; Mobile phase: A for H₂O (0.09% TFA) and B for MeCN; Gradient: B from 30% to 40% in 20 min; Flow rate: 60 mL/min; Wavelength: 220 & 254 nm.

[00165] ¹H NMR: (DMSO-d₆, 400 MHz) δ ppm 0.90 (s, 3 H) 1.19 - 1.41 (m, 2 H) 1.43 - 1.62 (m, 7 H) 1.64 - 1.77 (m, 3 H) 1.84 (br d, *J*=14.55 Hz, 1 H) 1.95 - 2.07 (m, 1 H) 2.18 - 2.36 (m, 3 H) 2.65 - 2.78 (m, 3 H) 3.71 - 3.86 (m, 3 H) 3.89 (s, 2 H) 3.93 (s, 2 H) 4.20 (br d, *J*=9.48 Hz, 1 H) 4.33 - 4.41 (m, 1 H) 4.59 (br dd, *J*=18.41, 8.05 Hz, 1 H) 4.81 (br dd, *J*=18.52, 8.60 Hz, 1 H) 4.94 (d, *J*=4.63 Hz, 1 H) 5.50 (s, 1 H) 5.54 - 5.76 (m, 1 H) 6.13 (s, 1 H) 6.29 (dd, *J*=10.14, 1.32 Hz, 1 H) 6.95 (d, *J*=7.72 Hz, 1 H) 7.15 - 7.28 (m, 4 H) 7.30 - 7.41 (m, 3 H) 7.51 (br d, *J*=7.94 Hz, 1 H) 7.72 (br s, 3 H) 8.21 (br d, *J*=7.72 Hz, 1 H) 8.54 (t, *J*=5.62 Hz, 1 H) 9.93 (br d, *J*=2.65 Hz, 1 H) LCMS (Method a, Table 4) R_t = 2.31 min.

Example 6: Synthesis of (S)-2-((2-(2-bromoacetamido)ethyl)amino)-N-(((S)-1-((3-(4-((6aR,6bS,7S,8aS,8bS,10R,11aR,12aS,12bS)-7-hydroxy-8b-(2-hydroxyacetyl)-6a,8a-dimethyl-4-oxo-2,4,6a,6b,7,8,8a,8b,11a,12,12a,12b-dodecahydro-1H-naphtho[2',1':4,5]indeno[1,2-d][1,3]dioxol-10-yl)benzyl)phenyl)amino)-1-oxopropan-2-yl)propanamide.

[00166] Example 6 product may be synthesized from coupling of N-(2-(((9H-fluoren-9-yl)methoxy)carbonyl)amino)ethyl)-N-(tert-butoxycarbonyl)-L-alanyl-L-alanine (the product of steps **S1** and **S2** below) to the amino product of Example 2, followed by steps **S4-S6**: (1) Fmoc deprotection, (2) coupling with 2-bromoacetic acid, and (3) Boc deprotection. Fmoc = Fluorenylmethyloxycarbonyl; Boc = tertbutoxycarbonyl.



[00167] General Cysteine Conjugation Protocol

[00168] An approximate 5-20 mg/mL solution of the desired antibody was prepared in PBS buffer, pH 6 - 7.4. A reducing agent of choice, such as TCEP, was diluted or dissolved in solvents like H₂O, DMSO,

DMA or DMF to give a solution with concentration range between 1 to 25 mM. Antibodies (anti-hTNF hIgG1 (D2E7) or anti-mTNF mIgG2a (8C11; McRae BL et al. *J Crohns Colitis* 10 (1): 69-76 (2016)) were then partially reduced by adding about 2-3.5 equivalents of reducing agent, briefly mixing, and incubating overnight at 0 – 4 °C. Tris buffer, pH 8-8.5 (20-50 mM) was then added, followed by the linker-drug in DMSO or DMA (less than 15% total) and the mixture was incubated for 2 - 3 hours at rt. Excess linker-drug and organic solvent were then removed by purification. Purified ADC samples were then analyzed by SEC, HIC and reduced mass spectrometry.

[00169] ADC Analytical Procedures

[00170] ADCs were profiled by either anionic exchange chromatography (AEC) or hydrophobic interaction chromatography (HIC) to determine the degree of conjugation and purity of ADC.

[00171] **AEC.** Approximately 20 µg of ADC was loaded onto an Ultimate 3000 Dual LC system (Thermo Scientific) equipped with a 4 X 250 mm Propac™ WAX-10 column (Tosoh Bioscience, cat. 054999). Column was equilibrated with 100% buffer A and eluted using a linear gradient from 100% buffer A to 100% buffer B over 18 min at 1.0 mL/min, where buffer A is 20 mM MES, pH 6.7 and buffer B is 20 mM MES, 500 sodium chloride, pH 6.7.

[00172] **HIC.** Approximately 20 mg of the ADC was loaded onto an Ultimate 3000 Dual LC system (Thermo Scientific) equipped with a 4.6 X 35 mm butyl-NPR column (Tosoh Bioscience, cat. 14947). Column was equilibrated in 100% buffer A and eluted using a linear gradient from 100% buffer A to 100% buffer B over 12 min at 0.8 mL/min, where buffer A is 25 mM sodium phosphate, 1.5 M ammonium sulfate, pH 7.0 and buffer B is 25 mM sodium phosphate, 25% isopropanol, pH 7.0.

[00173] **SEC.** Size distributions of the ADCs were profiled by size exclusion SEC using an Ultimate 3000 Dual LC system (Thermo Scientific) equipped with a 7.8 X 300 mm TSK-gel 3000SW_{XL} column (Tosoh Bioscience, cat. 08541). Approximately 20 µg of ADC was loaded onto the column and eluted over 17 min using an isocratic gradient of 100 mM sodium sulfate, 100 mM sodium phosphate, pH 6.8 at a flow rate of 1.0 mL/min.

[00174] **MS.** Reduced samples (10 µL) were injected to an Agilent 6550 QToF LC/MS system through a temperature controlled (5 °C) CTC autosampler. Sample elution was achieved on a Waters C-4, 3.5 µm, 300 Å, 2.1 x 50 mm i.d. HPLC column. The mobile phases were: A: 0.1% formic acid in water, and B: 0.1% formic acid in MeCN; the flow rate was 0.45 mL/min, and the column compartment was maintained at 40°C.

[00175] The HPLC gradient is as set forth in Table 5:

Table 6




Time (min)	Mobile Phase A (%)	Mobile Phase B (%)
0	75	25
14	5	95
16	0	100
18	0	100

Figure 2 shows a deconvoluted mass spectrum of the purified ADC. This ADC has 4 drug linker molecules conjugated to each antibody. The peak on the left has molecular weight of 24284.74 Da, which is the result of one drug linker attached to single light chain. The peak on the right has molecular weight of 51513.96 Da, which is the result of one drug linker attached to single heavy chain. ADCs in Table 7 and 8 were prepared according to the procedure described above.

Table 7: Anti-mouse TNF α antibody drug conjugates. X refers to the anti-murine TNF α antibody 8C11

ID NO.	Structure	DAR
ADC1		4
ADC2		4
ADC3		4

Table 8: Anti-human TNF α antibody drug conjugates. X refers to the anti-human TNF α antibody adalimumab

Corporate ID	Structure	DAR
ADC4		4
ADC5		2
ADC6		4

Generation of human and mouse transmembrane TNF-alpha GRE reporter cell lines

[00180] In order to create a parental cell line, K562 cells were seeded onto a 6 well dish (Costar: 3516) with 2 mL of complete growth medium (RPMI, 10% FBS, 1% L-glutamine, 1% Na Pyruvate and 1% MEM NEAA) at 500,000 cells per well for 24 hours at 37°, 5% CO₂. The next day, 1.5 mg of pGL4.36[Luc2P/MMTV/Hygro] (Promega: E316), 1.5 ug pGI4.75 [hRLuc/CMV] (Promega: E639A), and 3 uL of PLUS reagent (Invitrogen: 10964-021) were diluted into 244 mL Opti-MEM (Gibco: 31985-070) and incubated at rt for 15 minutes. The pGL4.36[luc2P/MMTV/Hygro] vector contains MMTV LTR (Murine Mammary Tumor Virus Long Terminal Repeat) that drives the transcription of the luciferase reporter gene *luc2P* in response to activation of several nuclear receptors such as glucocorticoid receptor and androgen

receptor. The pGL4.75[*hRluc*/CMV] Vector encodes the luciferase reporter gene *hRluc* (*Renilla reniformis*) and is designed for high expression and reduced anomalous transcription.

[00181] After incubation, diluted DNA solution was pre-incubated with 1:1 Lipofectamine LTX solution (Invitrogen: 94756) (13.2 mL + 256.8 mL Opti-MEM) and incubated at room temperature for 25 minutes to form DNA-Lipofectamine LTX complexes. After incubation, 500 μ L of DNA-Lipofectamine complexes were added directly to the well containing cells. K562 cells were transfected for 24 h at 37 °C, 5% CO₂. After incubation, cells were washed with 3 mL of PBS and selected with complete growth medium containing 125 mg/mL of hygromycin B (Invitrogen: 10687-010) for two weeks. "K562 pGL4.36[Luc2P/MMTV/Hygro]_pGL4.75[hRLuc/CMV]" cells were produced.

[00182] In order to create a murine transmembrane TNF-alpha GRE reporter cell line, the parental cells, K562 pGL4.36[Luc2P/MMTV/Hygro]_pGL4.75[hRLuc/CMV], were seeded onto 6 well dish (Costar: 3516) with 2 mL of complete growth medium (RPMI, 10%FBS, 1%L-glutamine, 1% Na Pyruvate and 1% MEM NEAA) at 500,000 cells per well for 24 h at 37°, 5% CO₂. The next day, 3 mg of mFL_TNFa DNA (Origene: MC208048), which encodes untagged mouse TNF, and 3 mL of PLUS reagent (Invitrogen: 10964-021) were diluted into 244 mL Opti-MEM (Gibco: 31985-070) and incubated at rt for 15 minutes. After incubation, diluted DNA solution was pre-incubated with 1:1 Lipofectamine LTX solution (Invitrogen: 94756) (13.2 mL + 256.8 mL Opti-MEM) and incubated at rt for 25 minutes to form DNA-Lipofectamine LTX complexes. After incubation, 500 μ L of DNA-Lipofectamine complexes were added directly to the well containing cells. The parental K562 pGL4.36[Luc2P/MMTV/Hygro]_pGL4.75[hRLuc/CMV] cells were transfected for 24 h at 37°C, 5% CO₂. After incubation, cells were washed with 3 mL of PBS and selected with complete growth medium containing 125 mg /mL of hygromycin B (Invitrogen: 10687-010) and 250 mg /mL G418 (Gibco: 10131-027) for two weeks. "K562 mouse FL-TNFa GRE (pGL4.36[luc2P/MMTV/Hygro])" cells were produced.

[00183] In order to create a human transmembrane TNF-alpha GRE reporter cell line, the parental cells, K562 pGL4.36[Luc2P/MMTV/Hygro]_pGL4.75[hRLuc/CMV], were transfected with the plasmid hTNF delta 1-12 C-Myc pcDNA3.1(-) plasmid construct. This plasmid is pcDNA 3.1 (ThermoFisher cat# V79020) encoding tace resistant transmembrane TNF (i.e., SEQ ID NO:1 lacking amino acids 77-88). (See Perez C et al. *Cell* 63 (2): 251-8 (1990) discussing tace resistant transmembrane TNF.) These cell lines were then used in the TNF-alpha reporter assays described in the subsequent examples.

Activity of anti-TNF-alpha immunoconjugates in GRE transmembrane TNF-alpha reporter assays

[00184] K562 parental GRE (pGL4.36[luc2P/MMTV/Hygro]) cells and K562 mFL-TNF-a or hTNF delta 1-12 GRE (pGL4.36[luc2P/MMTV/Hygro]) cells were plated onto 96 well tissue culture treated white plates (Costar: 3917) at 50,000 cells per well in 50 μ L of assay medium (RPMI, 1% CSFBS, 1% L-glutamine, 1% Na Pyruvate and 1% MEAA). The cells were treated with 25 μ L of 3X serial diluted murine or human anti-TNF-a antibody drug conjugates in assay medium, steroid compound, or media alone and

incubated for 48 hours at 37 °C, 5% CO₂. After 48 hours of incubation, cells were treated with 75 µL of Dual-Glo Luciferase Assay System (Promega-E2920) for 10 min and analyzed for luminescence using the Microbeta (PerkinElmer). Data were analyzed using a four parameter curve fit to generate EC₅₀ values. % maximum activation was normalized to 100 nM dexamethasone. The results using the murine TNF-alpha cell line are shown in Table 9 below, and the results using the human TNF-alpha cell line are shown in Table 10 below. In Table 9 below, the antibody in the ADC is the anti-murine TNFα antibody 8C11. In Table 10 below, the antibody in the ADC is the anti-human TNFα antibody adalimumab. Percent (%) monomer was determined by SEC as previously described (see ADC analytical procedures).

Table 9: *In vitro* activity of anti-murine TNFα antibody drug conjugate in mouse transmembrane TNFα GRE reporter assay

ADC	% Monomer	mTNFα GRE EC ₅₀ (µg/mL)	mTNFα GRE (% max)	K562 GRE EC ₅₀ (µg/mL)	K562 GRE (% max)
ADC1	99.9	0.06	150	>50	63
ADC2	99.3	0.39	164	>50	72
ADC3	100	0.04	104	3.9	84

Table 10: *In vitro* activity of anti-human TNFα antibody drug conjugate in human transmembrane TNFα GRE reporter assay

ADC	% Monomer	hTNFα GRE EC ₅₀ (µg/mL)	hTNFα GRE (% max)	K562 GRE EC ₅₀ (µg/mL)	K562 GRE (% max)
ADC4	99.9	0.03	118	>50	63
ADC5	100	0.03	126	>50	28
ADC6	100	0.05	97	>50	83

Activity of anti-hTNF alpha immunoconjugates in Lipopolysacharride Stimulated Human PBMC Cytokine Release Assay

[00185] Primary human peripheral blood mononuclear cells (PBMCs) were purchased from Biological Specialty Corporation (cat# 214-00-10), washed in 50 mL PBS, re-suspended in FBS with 5% DMSO, aliquoted and cryopreserved in liquid nitrogen until use. The PBMCs were thawed, re-suspended in RPMI

media supplemented with 2% FBS, and 1% Penicillin-Streptomycin, and plated into a cell assay plate (Costar #3799). Then the cells were incubated varying concentration of anti-TNF ADC at 37 °C and 5% CO₂ for 4 hours. Cells were then stimulated with 100 ng/mL LPS overnight. On the following day, plate was spun for five minutes at 1000 rpm, and 100 µL of supernatant media was directly transferred to an additional 96-well plate and analyzed for IL-6 (MSD, #K151AKB) and IL-1 beta (MSD, #K151AGB) concentrations. The dose response data were fitted to a sigmoidal curve using nonlinear regression, and the IC₅₀ values calculated with the aid of GraphPad 5.0 (GraphPad Software, Inc.). The results shown in Table 11 demonstrate that the anti-TNF ADC has potent activity in inhibiting the release of pro-inflammatory cytokines IL-6 and IL-1beta from activated primary immune cells.

Table 11. In vitro activity of anti-TNF steroid ADC in stimulated PBMC cytokine release assay

ADC	IL-1 beta IC ₅₀ (ng/mL)	IL-6 IC ₅₀ (ng/ml)
ADC4	44	265

Activity of anti- mTNF-alpha immunoconjugate in contact hypersensitivity model

[00186] Anti-TNF α steroid ADC was evaluated in an acute contact hypersensitivity model, an elicitation of acute skin inflammation using delayed-type hypersensitivity (DTH) response (T-cell driven) via application of a sensitizing agent (fluorescein isothiocyanate (FITC)). The efficacy of anti-TNF α steroid ADCs was measured by the ability to reduce ear swelling. The steroid biomarkers corticosterone and procollagen type 1 N-terminal propeptide (P1NP) were included as readouts to assess the putative impact of anti-TNF α steroid ADC treatment on the Hypothalamus-Pituitary-Adrenal (HPA) axis and bone turnover respectively.

[00187] *Ear Swelling*

[00188] On day 0 mice were placed under general anesthesia and the abdomens were shaved. Using a micropipettor, mice were sensitized by epicutaneous application of 400 µL of FITC solution (1.5% solution in 1:1 acetone:DBP) on the abdomen. Six days later mice were dosed with vehicle or therapeutic agent 1 hour prior to ear challenge with FITC. For ear challenge, mice were placed under general anesthesia and were challenged with 20 µL FITC applied onto right ear. Twenty-four hours after challenge mice were placed under general anesthesia and their ear thickness is measured by caliper. Difference between challenged and unchallenged ears was calculated. Seventy-two hours after ear challenge, mice were injected with ACTH at 1 mpk IP, and terminally bled at 30 min post-ACTH. Plasma was collected and analyzed P1NP, corticosterone, free steroid, and large molecule levels.

[00189] *Quantification of released free steroid and endogenous corticosterone*

[00190] Calibration curve of steroid was prepared in mouse plasma with final concentrations from 0.03 nM to 0.1 μ M at 8 different concentration levels. Corticosterone calibration curve ranging from 0.3 nM to 1 μ M final corticosterone concentrations was prepared in 70 mg/mL bovine serum albumin solution in PBS buffer. A solution of 160 μ L MeCN with 0.1% formic acid was added to 40 μ L study plasma samples or calibration standards. Supernatants were diluted with distilled water and 30 μ L final sample solution was injected for LC/MS analysis.

[00191] Quantification of released free steroid and corticosterone was conducted on an AB Sciex 5500 triple quadrupole mass spectrometer connected to a Shimadzu AC20 HPLC system interfaced with an electrospray ionization source operating in positive mode. A Waters XBridge BEH C18, 2.1x30mm, 3.5 μ m column was used for chromatography separation. The mobile phase A was 0.1% formic acid in Milli Q HPLC water, and mobile phase B was 0.1% formic acid in MeCN. A linear gradient from 2% of mobile phase B to 98% mobile phase B was applied from 0.6 to 1.2 min. The total run time was 2.6 min at a flow rate of 0.8 mL/min. The mass spectrometer was operated in positive MRM mode at source temperature of 700 C.

[00192] *Quantification of plasma PINP*

[00193] Quantification of plasma PINP was conducted on a LCMS platform based on protein trypsin digestion. Plasma samples were partially precipitated and fully reduced by adding MeCN/0.1M ammonium bicarbonate/DTT mixture. Supernatant was collected and alkylated by adding iodoacetic acid. The alkylated proteins were digested by trypsin and resulting tryptic peptides were analyzed by LCMS. Calibration curve were generated by using synthetic tryptic peptide spiked into horse serum (non-interfering surrogate matrix). Stable isotope labeled flanking peptide (3-6 amino acids extension on both termini of the tryptic peptide) was used as internal standard added in the MeCN/DTT protein precipitation mixture to normalize both digestion efficiency and LCMS injection.

[00194] A Columnex Chromenta BB-C18, 2.1x150 mm, 5 μ m column was used for chromatography separation. The mobile phase A was 0.1% formic acid in Milli Q HPLC water and mobile phase B was 0.1% formic acid in MeCN. A linear gradient from 2% of mobile phase B to 65% mobile phase B was applied from 0.6 to 3 min. The total run time was 8 min at a flow rate of 0.45 mL/min. An AB Sciex 4000Q trap mass spectrometer was used in positive MRM mode to quantify PINP peptides, at source temperature of 700 C.

[00195] *Results*

[00196] The results are shown in Table 12 below:

Table 12: Comparison of anti-mTNF alpha steroid ADC activity on ear swelling and steroid biomarkers in CHS model of inflammation

ADC	Ear swelling (% inhib @ 10mpk ± SEM)	PINP (% inhib. @ 10mpk ± SEM)	Corticosterone (% inhib @ 10mpk ± SEM)
ADC1	82.8 ± 2.3	32.6 ± 3.8	4.7 ± 10.5

Activity of anti- mTNF-alpha immunoconjugates in collagen-induced arthritis

[00197] The ability of anti-mTNF α steroid ADC (ADC1) to impact disease was assessed in the collagen-induced arthritis (CIA) model of arthritis.

[00198] In these experiments, male DBA/1J mice were obtained from Jackson Labs (Bar Harbor, ME). Mice were used at 6 to 8 weeks of age. All animals were maintained at constant temperature and humidity under a 12 hour light/dark cycle and fed with rodent chow (Lab Diet 5010 PharmaServ, Framingham, MA) and water ad libitum. AbbVie is AAALAC (Association for Assessment and Accreditation of Laboratory Animal Care) accredited, and all procedures were approved by the Institutional Animal Care and Use Committee (IACUC) and monitored by an attending veterinarian. Body weight and condition were monitored, and animals were euthanized if exhibiting >20% weight loss.

[00199] The male DBA/J mice were immunized intradermally (i.d.) at the base of the tail with 100 μ L of emulsion containing 100 μ g of type II bovine collagen (MD Biosciences) dissolved in 0.1 N acetic acid and 200 μ g of heat-inactivated Mycobacterium tuberculosis H37Ra (Complete Freund's Adjuvant, Difco, Laurence, KS). Twenty-one days after immunization with collagen, mice were boosted IP with 1 mg of Zymosan A (Sigma, St. Louis, MO) in PBS. Following the boost, mice were monitored 3 to 5 times per week for arthritis. Rear paws were evaluated for paw swelling using Dyer spring calipers (Dyer 310-115)

[00200] Mice were enrolled between days 24 and 28 at the first clinical signs of disease and distributed into groups of equivalent arthritic severity. Early therapeutic treatment began at the time of enrollment.

[00201] Animals were dosed once intraperitoneal (i.p.) with anti-mTNF mAb (high dose) or anti-mTNF steroid ADC (high and low dose - mpk) in 0.9% saline. Blood was collected for antibody exposure by tail nick at 24 and 72 hours after dose. Paws were collected at the terminal time-point for histopathology. Blood was collected at the terminal time-point by cardiac puncture for complete blood counts (Sysmex XT-2000iV). Statistical significance was determined by ANOVA.

[00202] The results are shown in Fig. 3 and demonstrate that a single dose of anti-TNF α steroid ADC can exhibit an extended duration of action through amelioration of paw swelling for ~28 days compared to anti-TNF α mAb or vehicle alone.

ADC Stability in Plasma

[00203] Although hydrolysis has been employed to increase *in vivo* stability of maleimide-based linkers, it generally requires exposure to basic pH, which may lead to modifications in the antibody (e.g.,

deamidation), increased heterogeneity, decreased yield and the like. (Shen et al., *Nature Biotechnology* 30:184–189 (2012) (hydrolyzing maleimides avoids premature release of the drug and systemic exposure to the drug); Strop et al., *Chemistry & Biology* 20(2):161-167 (2013) (a similar study); Tumey et al., *Bioconjugate Chem* 25(10):1871-1880 (2014) (use of a proximal PEG chain to enable ring hydrolysis under basic conditions); Lyon et al., *Nature Biotechnology* 32:1059-1062 (2014) (processes for facilitating the creation of hydrolyzed succinimide); Christie et al., *J Control Release* 220(PtB):660-70 (28 Dec 2015) (use of N-aryl maleimides to facilitate succinimide ring hydrolysis under basic conditions); Dovgan et al., *Scientific Reports* 6:1 (2016)(use of 2-(maleimidomethyl)-1,3-dioxane for facilitating ring hydrolysis under mildly basic conditions for a prolonged period of time); and *J Pharm Sci* 2013: 102 (6) 1712-1723 (asparagine deamidation dependence on buffer type, pH and temperature). In contrast, typical conditions for conjugation using bromo acetamide are complete within a few hours, compared to multiple days for maleimide (conjugation and subsequent ring hydrolysis at basic pH). In addition, the 2-mercaptoacetamide formed during reaction of cysteine with the bromo acetamide is not susceptible to the reverse Michael reaction and affords a stable attachment of the linker to the antibody as demonstrated in Table 13 below for ADC4.

Table 13


% SM released from ADC in plasma after 4d @ 37°C				
PBS1X buffer	Mouse	Rat	Cyno	Human
BLQ	BLQ	BLQ	BLQ	BLQ

BLQ- Below level of quantification (<1.0nM)

ADC stability in Solution

[00204] To assess long-term stability, a biologic is subjected to an accelerated stress test. The protocol for this test is 100 mg/mL of the biologic in 15 mM histidine at 40 °C for 21 days. When ADC4 was subjected to this test the percent increase in aggregate was <5% compared to ADC203 of US Patent Application Publication no.2018/012600, published May 10, 2018, which saw an increase in aggregation of 18% (Table 14). This demonstrates the improved properties imparted to ADC4 by the Gly-Glu linker with the phosphate prodrug of the payload. ADC203 of US2018/0126000 is:

203




where n=4 and A refers to the anti-human TNF α antibody adalimumab.

Table 14

ADC	Monomer Loss at 5°C for 21 Days	Monomer Loss at 25°C for 21 Days	Monomer Loss at 40°C for 21 Days
ADC203	1.43%	2.26%	17.56%
ADC4	<0.5%	<0.5%	3.46%

[00205] An additional benefit of the phosphate prodrug is it enables the use of anion exchange chromatography for the DAR purification. This results in improved peak resolution compared to hydrophobic interaction chromatography resulting in higher yields of DAR purified ADC.

[00206] In formulation buffer the hydrolyzed succinimide ring of ADC203 is in equilibrium with the ring closed form. The ring closed form is susceptible to the reverse Michael reaction and subsequent loss of linker-drug *in vivo* in cynomolgus monkeys (Fig. 4).



[00207] Under nominal liquid storage conditions the succinimide attachment in the open conformation will reform the closed conformation greater than 5% at 5 °C and greater than 15% at 25 °C after 6 months (Table 15).

Table 15

Time/Temp	% Closed Succinimide Ring	
	Light Chain	Heavy Chain
3 Months/ 5°C	2.2	4.6
6 Months/ 5°C	3.2	6.9
3 Months/ 25°C	10.1	13.3
6 Months/ 25°C	16.6	24.1
3 Months/ 40°C	26.2	33.7
3 Months/ 40°C	26.8	39.0

[00208] It is to be appreciated that the Detailed Description section, and not the Summary and Abstract sections, is intended to be used to interpret the claims. The Summary and Abstract sections sets forth one or more, but not all, exemplary embodiments of the present disclosure as contemplated by the inventor(s), and thus, are not intended to limit the present disclosure and the appended claims in any way.

[00209] The present disclosure has been described above with the aid of functional building blocks illustrating the implementation of specified functions and relationships thereof. The boundaries of these functional building blocks have been arbitrarily defined herein for the convenience of the description. Alternate boundaries can be defined so long as the specified functions and relationships thereof are appropriately performed.

[00210] The foregoing description of the specific embodiments will so fully reveal the general nature of the disclosure that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present disclosure. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.


[00211] The breadth and scope of the present disclosure should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents.

INCORPORATION BY REFERENCE


[00212] All publications, including patents and published applications, referred to in the Detailed Description are incorporated by reference herein in their entirety.

WHAT IS CLAIMED IS:


1. An antibody drug conjugate comprising:
 - (a) an anti-TNF α antibody comprising a heavy chain set forth as SEQ ID NO: 3 and a light chain set forth as SEQ ID NO: 4; and
 - (b) a glucocorticoid receptor agonist comprising a radical represented by the formula:



wherein the antibody is conjugated to the glucocorticoid receptor agonist via a linker represented by the formula:



2. The antibody drug conjugate of claim 1, according to the formula:



wherein A is the antibody and n is an integer from 1-10.

3. The antibody drug conjugate of claim 2, wherein n is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.
4. The antibody drug conjugate of claim 2, wherein n is 4.
5. The antibody drug conjugate of claim 2, wherein n is 2.

6. A pharmaceutical composition comprising the antibody drug conjugate of claim 4 and a pharmaceutically acceptable carrier.

7. A method of treating a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis in a subject, comprising administering an effective amount of the antibody drug conjugate of claim 4 to the subject.


8. A kit comprising:

(a) a container comprising the antibody drug conjugate of claim 4 or the pharmaceutical composition of claim 6; and

(b) a label or package insert on or associated with the one or more containers, wherein the label or package insert indicates that the antibody drug conjugate or pharmaceutical composition is used for treating a condition selected from rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, plaque psoriasis, ulcerative colitis, adult Crohn's disease, pediatric Crohn's disease, uveitis, hidradenitis suppurativa, and juvenile idiopathic arthritis.


9. A method of delivering a glucocorticoid receptor agonist to a TNF α -expressing cell, comprising the step of contacting the cell with the antibody drug conjugate of claim 4.

10. An antibody drug conjugate according to the formula:



wherein A is adalimumab and n is 4.

11. An antibody drug conjugate according to the formula:



wherein A is adalimumab and n is 2.




FIG. 1





FIG. 2



* p < 0.05 VS. VEHICLE BY 2-way ANOVA

DAYS OF DISEASE

FIG. 3




FIG. 4

INTERNATIONAL SEARCH REPORT

International application No PCT/IB2018/059482

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K47/68 A61P19/02 C07K16/24 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols) A61K A61P C07K				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A	PATEL TORAL ET AL: "Adalimumab: efficacy and safety in psoriasis and rheumatoid arthritis", DERMATOLOGIC THERAPY, WILEY-BLACKWELL PUBLISHING, INC, US, vol. 17, no. 5, 1 January 2004 (2004-01-01), pages 427-431, XP002477489, ISSN: 1396-0296, DOI: 10.1111/J.1396-0296.2004.04045.X the whole document -----	1-11		
A	WO 2015/153401 A1 (MERCK SHARP & DOHME [US]; AMBRX INC [US]; GARBACCIO ROBERT M [US]; KER) 8 October 2015 (2015-10-08) the whole document claims 6-11 ----- -/--	1-11		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.				
* Special categories of cited documents : <table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none; vertical-align: top;"> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td style="width: 50%; border: none; vertical-align: top;"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
22 February 2019	14/03/2019			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Mooren, Nicolai			

INTERNATIONAL SEARCH REPORT

International application No PCT/IB2018/059482

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	WO 2017/210471 A1 (ABBVIE INC [US]) 7 December 2017 (2017-12-07) the whole document Compound no. 210 in Table 12 on page 351 -----	1-11

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/IB2018/059482
--

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2015153401 A1	08-10-2015	EP 3125943 A1	08-02-2017
		US 2017182181 A1	29-06-2017
		WO 2015153401 A1	08-10-2015

WO 2017210471 A1	07-12-2017	AU 2017274442 A1	13-12-2018
		CA 3025377 A1	07-12-2017
		DO P2018000261 A	31-12-2018
		SG 11201810678W A	28-12-2018
		TW 201801749 A	16-01-2018
		US 2018126000 A1	10-05-2018
		UY 37269 A	02-01-2018
		WO 2017210471 A1	07-12-2017
