

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199534772 B2
(10) Patent No. 707004

(54) Title
Modified human C3 proteins

(51)⁶ International Patent Classification(s)
C12N 015/12 C07K 014/47
A61K 038/17 C12N 009/64
A61K 038/48 C12N 015/57
A61K 047/48

(21) Application No: **199534772**

(22) Application Date: **1995.09.08**

(87) WIPO No: **WO96/07738**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9418147	1994.09.08	GB
9509102	1995.05.04	GB

(43) Publication Date : **1996.03.27**

(43) Publication Journal Date : **1996.05.16**

(44) Accepted Journal Date : **1999.07.01**

(71) Applicant(s)
Imutran Limited

(72) Inventor(s)
Richard Alexander Harrison; Timothy Charles Farries

(74) Agent/Attorney
F B RICE and CO,605 Darling Street,BALMAIN NSW 2041

(56) Related Art
US 4661347

OPI DATE 27/03/96 APPLN. ID 34772/95
AOJP DATE 16/05/96 PCT NUMBER PCT/GB95/02121



AU9534772

(51) International Patent Classification ⁶ : C12N 15/12, 15/57, C07K 14/47, C12N 9/64, A61K 38/48, 38/17, 47/48		A2	(11) International Publication Number: WO 96/07738
			(43) International Publication Date: 14 March 1996 (14.03.96)
(21) International Application Number: PCT/GB95/02121		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 8 September 1995 (08.09.95)			
(30) Priority Data: 9418147.6 8 September 1994 (08.09.94) GB 9509102.1 4 May 1995 (04.05.95) GB			
(71) Applicant (for all designated States except US): IMUTRAN LIMITED [GB/GB]; 21 Holborn Viaduct, London EC1A 2DY (GB).		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(72) Inventors; and (75) Inventors/Applicants (for US only): HARRISON, Richard, Alexander [GB/GB]; MRC Centre, Hills Road, Cambridge CB2 2QH (GB). FARRIES, Timothy, Charles [GB/GB]; MRC Centre, Hills Road, Cambridge CB2 2QH (GB).			
(74) Agents: WHITE, Martin, Paul et al.; Kilburn & Strode, 30 John Street, London WC1N 2DD (GB).			
(54) Title: MODIFIED HUMAN C3 PROTEINS			
(57) Abstract Native complement pathway proteins modified such that the protein is capable of forming a stable C3 convertase. Preferably the modified protein is a modified human C3 protein. DNA sequences encoding such proteins are also provided, together with DNA constructs. Conjugates comprising such proteins and a specific binding moiety, for example an antibody, are also described, as are uses of such proteins and/or conjugates in therapy.			

MODIFIED PROTEINS

5 The present invention relates to novel modified proteins capable of forming stable C3 convertases, DNA sequences encoding such proteins and the use of such proteins as therapeutic agents, particularly for use in depleting levels of complement pathway proteins.

10 The complement system functions in the immune response of humans and other vertebrates, being of major importance in the effector functions such as phagocytosis, cytolysis and recruitment of cells that induce local inflammatory responses [15]. These properties are desirable for elimination of invading pathogens, such as bacteria, but undesirable when triggered to act against host tissues
15 (e.g. in post-ischemic reperfusion injury [3]) or against foreign therapeutic material (e.g. hyperacute rejection of xenografts [7]). There have been attempts to abrogate these undesirable properties by exploiting derivatives of complement regulatory proteins whose normal function is
20 to suppress complement activation [10, 18].

The complement system comprises proteins both on the surface of cells, (receptors and regulators) as well as in the fluid-phase (blood plasma and other extracellular
25 environments). The critical step for the generation of responses is the proteolytic conversion of C3 to the fragments C3b and C3a. C3a is an anaphylatoxin that, like C5a, attracts mast cells to the site of challenge, resulting in local release of histamine, vasodilation and
30 other inflammatory effects. The nascent C3b has an ability to bind to surfaces around its site of generation. This C3b then focuses attack by the cytolytic complement components (C5-C9).

Surface-bound C3b, and its degradation products, also function as ligands for C3 receptors mediating, for example, phagocytosis [15]. There are two distinct pathways of complement activation that both result in conversion of C3 to C3b and subsequent responses. The classical pathway is commonly triggered by complexes of antibody with antigen, initiating an enzyme cascade involving the proteins C1q, C1r, C1s, C2 and C4. The alternative pathway depends on an activation loop involving C3 itself and requiring factors B and D.

Conversion of C3 to C3b (or C3i) produces a product that can combine with factor B, giving C3bB (or C3iB). These complexes are acted upon by factor D to generate C3bBb, which is a C3 convertase capable of cleaving more C3 to C3b, leading to more C3bBb and even more C3 conversion. Under certain circumstances the C3bBb complex is stabilised by association with the positive regulator properdin (P). However, this positive-feedback loop is normally limited to a slow tick-over by regulatory proteins, notably factor H and factor I.

Factor H (and structurally related cell-associated molecules) (i) displaces B and Bb from C3b, and (ii) acts as a cofactor for factor I which cleaves C3b into iC3b thereby preventing any recombination with factor B to form more C3 convertases. The pathway is "fired" into amplified generation of C3b in the presence of surfaces, such as many bacterial cell walls, that bind nascent C3b and impede its regulation by factors H and I. Nascent C3b is also able to bind to endogenous cells. Endogenous cell surfaces normally exposed to complement are therefore additionally protected by membrane-bound regulators such as MCP, DAF and CR1 acting in a similar manner to factor

H.

There are a few rare naturally occurring conditions where the normal fluid-phase regulation cannot occur and spontaneous C3 conversion ultimately results in generalised depletion of C3 from the circulation:- (i) genetic deficiencies of factor H or I [13], (ii) the presence of antibodies (nephritic factors) that bind to C3bBb and impede dissociation [4], and (iii) contact with a protein in cobra venom, called cobra venom factor (CVF), that combines with factor B and forms a C3 convertase enzyme which does not contain C3b and is not affected by factors H and I [14]. These illustrate the normal physiological importance of down-regulation of complement in the absence of specific activation.

There are also circumstances where specific activation occurs, but is unwanted, particularly when it is directed against tissues of the host (e.g. tissue damaged by ischemia or surgery) or against foreign material deliberately given for therapeutic purposes (such as a xenograft, artificial organ or a dialysis membrane). The complement activation results in undesirable attack and further damage, so in these cases it would be beneficial to block or inhibit the activation and response.

Existing approaches to preventing complement-mediated damage have targeted the use of down-regulatory proteins (CR1, MCP, DAF and factors H and I) to inhibit complement activation. Complement inhibitors like factor I, factor H and soluble derivatives of the membrane-bound proteins CR1, DAF, MCP do suppress the fluid-phase amplification loop of the alternative pathway. Therefore there have been attempts to use these molecules, particularly CR1

(which seems to be the most potent) to reduce complement-mediated damage in models of physiological situations [10, 18].

5 Factor H is endogenously present in blood plasma in high concentrations (typically 0.3-0.5 mg/ml [15]), so even though increased levels of inhibitors do dampen-down fluid-phase reactions, their potency is weak so large amounts of purified proteins would have to be
10 administered *in vivo* (e.g probably in excess of 5 mg/Kg body weight of soluble CR1). In addition, the alternative pathway is activated by surfaces where the effect of factor H is already impeded. While this does not necessarily concomitantly reduce the activities of other
15 inhibitors, the same factors suggest that they are unlikely to be completely or universally effective.

Cobra Venom Factor (CVF) has the property of generating a stable C3 convertase which can be used experimentally to
20 deplete complement in animals *in vivo*, and in other samples (e.g. human blood plasma) *in vitro*. CVF is potent (e.g. 40 μ g/Kg can destroy the complement activity of a mouse [16]). However, there are disadvantages that make it unsuitable for therapeutic use in humans.

25 Firstly, it is obtained from cobra venom (a difficult source to obtain and dangerous to handle) and must therefore be carefully purified from the venom neurotoxins. There is also the obvious difficulty in
30 obtaining supplies. This problem cannot readily be overcome by cloning and expressing the gene *ex vivo*, because there are post-translational modifications that occur in the snake (specific proteolytic processing) that may be difficult (or impossible) to reproduce *in vitro*.

In addition, the enzymes and digestion conditions required for this processing are currently unknown. Secondly, the protein is of foreign origin (to humans) and therefore immunogenic. This precludes its repeated therapeutic use, as would be required to deplete a patient over many weeks (e.g. to allow xenograft survival).

Although CVF has some structural and functional homologies with human C3 [17], it also has major differences in both respects (e.g. chain structure, site of biosynthesis, insensitivity to complement regulators, formation of a stable C3 convertase). It is not derived from the cobra equivalent of C3 which is known, having been cloned and sequenced, and which in gross structure and function resembles human C3 more closely than does CVF [8].

CVF is a venom-specific product of an animal of great evolutionary distance from homo sapiens. It is therefore not practicable to use genetic manipulation to modify this protein into a product that can be used non-immunogenically in humans.

We have now devised an alternative strategy which relies on by-passing the physiological regulation and, instead of inhibiting complement activation, causes the system to be super-activated. This has two applications. Firstly, it can be used *in vivo* to activate complement until one or more components are exhausted, resulting in loss of ability to produce local responses to any subsequent challenge (such as a xenograft). Secondly, the unregulated super-activation can be deliberately localised to a particular target (e.g. a virus or a

virally-infected cell) to increase the sensitivity of that target to complement-mediated destructive responses.

5 The term "regulators of complement activation" is used herein to include all proteins that act to inhibit amplification of C3 conversion, and is not intended to be restricted in meaning to those proteins whose genes are located in the RCA genetic locus. It does not however include "up-regulators" such as properdin. "C3
10 conversion" is defined as the proteolytic conversion of C3 into C3b and C3a, unless otherwise indicated, and "C3 convertase" (or simply "convertase") is defined as an enzyme (typically a complex of two or more protein components; for example C3bBb, C3iBb, CVFBb or C4b2a)
15 that catalyses this reaction.

Thus, in a first aspect the invention provides a native complement pathway protein modified such that the protein is capable of forming a stable C3 convertase.
20

By "native" is meant naturally occurring, ie is obtainable in nature. Thus, the definition encompasses any naturally occurring complement pathway protein modified as defined above. It is not intended to be
25 restricted to species specific proteins. In other words, a modified human protein could be used as a stable C3 convertase in other mammalian species, for example. Typically, modified complement pathway proteins from the same species will be used.

30 Modification of the C3 DNA coding sequence, for example using site directed mutagenesis, can produce a variant of C3 that is resistant to complement regulatory proteins while retaining positive functional properties (cleavage

to C3b by C3 convertase) and features of structural integrity (correct chain structure, and presence of a thiolester bond). The invention described herein relates to genetically-modified forms of native complement proteins, for example human C3, whose C3b fragment acquires the property of being resistant to physiological complement regulation. Because of this resistance, these molecules can generate stabilised forms of the corresponding C3 convertase that produce amplified conversion of C3 to C3b, and later degradation products, in physiological environments (e.g. in vivo).

In a preferred embodiment the invention provides a modified human C3 protein which is resistant to cleavage by factor I.

This can be achieved by modifying residues of the protein at proteolytic sites.

A particularly preferred embodiment of the invention relates to a modified human C3 protein wherein the protein is modified by replacement of either Arg-1303, Arg-1320 or both by another amino acid. The other amino acid may be Tyrosine, Cystine, Tryptophan, Glutamine, Glutamic acid or Glycine. Arg-1303 is preferably replaced by Glutamic acid or Glycine (less preferably by Glutamine). Arg-1320 is preferably replaced by Glutamine.

Other strategies for producing suitable modified proteins of the invention include:

i) Reduced susceptibility to the inhibitory actions of factor H and related proteins (eg. MCP, DAF, CR1). For

example, in human C3 residues 767-776 and 1209-1271 have been implicated in factor H binding [20,24], and substitution of one or more of these residues or other residues also associated with the action of these proteins, could reduce the binding of one or more of these regulatory proteins.

ii) Reduced rate of dissociation of C3bBb. Mutations can be introduced which would strengthen the interaction between C3b and Bb. This would result in both a reduction in spontaneous decomposition of the enzyme, and diminish the effectiveness of factor H (and related regulators) in displacing Bb from C3b.

These mutations are desirable to reduce the rates of both the spontaneous and the factor H-mediated decomposition of C3bBb. Even in the absence of factor H, the fluid phase C3bBb complex has a half-life of only about 10 mins at 37°C in the presence of properdin [6].

iii) Human C3 residues 752-761 are implicated in binding factor B. It is a highly conserved region in C3, and a closely related sequence is found in C4. As C4 binds the factor B homolog C2, the strong similarity of this region between C3 and C4, together with its high conservation in C3, further supports its role in C3 as a factor B binding site. Thus, changes in this region could have effects on B affinity and on the stability of C3bBb.

iv) Resistance to other regulators of complement activation such as CR1, DAF and MCP would also be desirable. The mode of action of these regulators are all similar to factor H, so additional mutagenesis would not necessarily be required. Similarly, some pathogenic

organisms express their own inhibitors of complement activation that are often structurally and functionally homologous to factor H (e.g. Vaccinia virus secretory peptide [1]). These molecules protect the invaders against immune responses, and it would be advantageous to be able to attack them with targeted C3 convertase enzymes resistant to these defences.

5
10
15
20
v) Mutations that increase the stabilisation of the C3 convertase by properdin. The activity of properdin is to stabilise the C3bBb complex, retarding spontaneous and factor H-dependent dissociation. This stabilisation is ineffective in the fluid-phase, but seems to be more important in amplifying the process once it has already started on a suitable activating surface [5]. Increasing its activity (by increasing its affinity) may upset the balance in the fluid-phase, and thereby promote spontaneous C3 conversion. This should be particularly useful in combination with the other modifications described above.

25
30
vi) Mutations that prevent the C3bBb from possessing C5 convertase activity. When used to deplete active C3 from the circulation an undesirable side-effect could be the generation of large amounts of anaphylactic peptides. The most potent of these is C5a, which is cleaved from C5 by some C3 convertase enzymes. This reaction probably depends on the affinity of the convertase for another molecule of C3b [11], and so may be subject to suppression by mutations to the C3 that remove this interaction.

vii) Improved activity of the C3 convertase. The active site of the C3bBb C3 convertase enzyme resides in the Bb

portion. The C3b component presumably functions to impose an active conformation on Bb and/or to bind and orientate the substrate to be acted upon by Bb. This is not known, but in either case there may be scope for enhancing the activity of the convertase through mutations in C3.

viii) Expression in a functional form. Wild-type C3 requires conversion to C3b before it can combine into a new C3 convertase complex. When used in vivo, a requirement for conversion to C3b (or C3i) would delay the action of the modified C3. It would therefore be desirable to either administer the protein in a form capable of immediate convertase formation, or to administer pre-formed convertase complexes. It is therefore advantageous to generate a functionally C3b-like reagent ex-vivo. This could be achieved in vitro (e.g. by proteolysis).

ix) Modifications to the native protein which serve to introduce new cleavage sites such that peptide regions required for factor B binding are retained but those required exclusively for factor H binding can be specifically removed. For example, sites can be introduced such that the C3b-like form of the modified C3 can be further cleaved into a form that still binds factor B but is less susceptible to inactivation by factors H and I.

x) Modifications in other regions which may affect the C3b interaction with factor B and/or factor H.

The invention is based on reversing the traditional approach by promoting C3 conversion to deplete C3 and

thereby disable the system. An additional application of the invention is the potential to promote C3 conversion at a particular site, and thereby recruit the complement-dependent effector mechanisms to attack a specific target.

Therefore the ultimate effect will be to increase the amount of C3 conversion when the modified protein is administered into a physiological medium (e.g. blood) containing regulators of complement activation. This activity can then be used either to deplete that medium of native C3, or to localise the C3 conversion at a desired target.

The analogue of C3 whose C3b-fragment is resistant to the actions of factor I (e.g. the derivative described in example 1) would bind factor B, which would then be cleaved by factor D and eventually dissociate in an inactive form. In the absence of inactivation by factor I, the modified C3b would be able to repeatedly bind new molecules of factor B and thereby promote its inactivation. Therefore another potential application of modifications described in this invention would be the inactivation of the alternative pathway by consumption of factor B activity. An analogous approach could also be used to modify C4 to promote the consumption of C2, and thereby disable the classical pathway of complement activation.

The invention includes any other protease used in an analogous manner to the C3bBb enzyme which leads to cleavage of C3 to C3b, despite the presence of regulators of complement activation.

The invention also includes DNA sequences which code for a protein of the invention as well as DNA constructs comprising such DNA sequences.

5 "DNA sequences" include all other nucleic acid sequences which, by virtue of the degeneracy of genetic code, also code for the given amino acid sequence or which are substantially homologous to this sequence. These sequences are thus also included within the scope of the
10 invention.

Nucleic acid sequences which are "substantially homologous" are also within the scope of the present invention. "Substantial homology" may be assessed either
15 at the nucleic acid level or at the amino acid level. At the nucleic acid level, sequences having substantial homology may be regarded as those which hybridise to the nucleic acid sequences of the invention under stringent conditions (for example, at 35 to 65°C in a salt solution
20 of about 0.9M). At the amino acid level, a protein sequence may be regarded as substantially homologous to another protein sequence if a significant number of the constituent amino acids exhibit homology. At least 55%, 60%, 70%, 80%, 90%, 95% or even 99%, in increasing order
25 of preference, of the amino acids may be homologous.

As discussed above the proteins of the invention can be used to achieve localised complement activation effects. One way of ensuring this is to conjugate the protein to
30 a moiety which will bind at the desired target. Thus, in another aspect the invention provides a conjugate comprising a protein of the invention linked to a specific binding moiety, for example a specific binding protein. An example of such a protein would be an

antibody or an antigen binding fragment thereof.

5 The proteins of the invention are intended to be administered to a subject to elicit a desired therapeutic effect. To that end therefore the invention also provides:

- a) A protein of the invention for use in therapy;
- 10 b) The use of a protein or a conjugate of the invention in the manufacture of a medicament for use in depleting levels of complement pathway protein, and in particular for use in preventing rejection of foreign matter;
- 15 c) A pharmaceutical formulation comprising one or more proteins or conjugates of the invention together with one or more pharmaceutically acceptable carriers and/or excipients; and
- 20 d) A method of reducing complement pathway protein in a mammal which comprises administering to the mammal a protein of the invention, preferably in the form of a pharmaceutical formulation.
- 25 Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per dose. Such a unit may contain as a minimum, for example, 1mg of active ingredient, and preferably 2-3mg. The upper limit which such a unit dose
- 30 can contain will depend on many factors such as the condition being treated, the route of administration and the age, weight and condition of the patient as well as economic considerations. As an example a unit dose form can contain as much as 10mg or even 100mg of active

ingredient.

5 The proteins of the invention could be used in vivo to disable the complement system. Circumstances where this may be desirable include the following:-

10 (a) In order to prevent complement-mediated destruction or damage to a transplant, particularly a xenograft (material transplanted from a different species of animal), and especially a discordant xenograft (where the donor and recipient species are discordantly related). The recipient would be decomplemented prior to the operation and maintained in this state until the transplant had either been accommodated or been replaced
15 by a more compatible organ.

The initial treatment could be made within several days before transplantation. Additional decomplementation could be required at times of rejection crisis. The
20 treatments may be accompanied by the use of anti-histamine reagents to control the general inflammatory responses (e.g. vasodilation) likely to result from the generation of C3a and/or C5a.

25 Decomplementation may also be beneficial in the use of artificial organs or tissues (e.g. artificial kidney dialysis membranes) which activate the complement system. As described above, the protein may be given either as the unactivated form, a functionally C3b-like form or a
30 preformed active C3 convertase (like C3bBb). These may be administered by any route whereby the active convertase will encounter the circulating C3 (e.g. intravenously, subcutaneously etc.).

Another alternative would be an ex vivo treatment, for example by transfusing the circulation through a matrix bearing the active convertase. This could have the advantage of allowing anaphylactic peptides (C3a and C5a) and other low molecular weight inflammatory mediators (e.g. histamine and nitric oxide) to be removed (e.g. by dialysis) prior to the decompemented blood (or plasma) being returned to the patient.

(b) To prevent complement-mediated damage resulting from major surgery. The patient would be decompemented, as above, preferably before the operation (but if necessary afterwards) and kept in this state until the danger of additional internal injury due to complement-dependent immune attack had diminished.

(c) To minimise complement-mediated damage resulting from non-surgical injury. In these cases the decompementation must be performed after the initial injury, but the formulations and methods of administration are likely to be otherwise similar to those described above. This may be particularly useful when the recovery involves reperfusion of an ischemic tissue by the circulation (e.g. myocardial ischemia, frostbite, burns etc.).

(d) To minimise complement-mediated damage resulting from antibody-antigen interactions. Complement-mediated defensive responses are particularly undesirable in autoimmune diseases which may include glomerulonephritis, haemolytic anaemia, myasthenia gravis and type II collagen-induced arthritis. Disabling the complement system during severe episodes of disease may alleviate the condition.

- (e) To make a specific pathogenic target more susceptible to complement-mediated immune mechanisms. In this approach, the aim is not to use the super-active C3 convertase to produce generalised depletion of C3, but instead to use the convertase locally to concentrate the C3 conversion at a desired target. The target may be a pathogenic organism, such as a bacteria, virus or other parasite, or a deleterious host cell or tissue, such as a tumour cell or a virally-infected cell. The C3 convertase could be localised to the target either by local administration (e.g. direct injection, possibly in a medium that retards its dispersion into the general circulation), or by combining with a targeting moiety, e.g. an antibody. Thus the modified protein could be linked to a specific immunoglobulin either by chemical cross-linking of the proteins, or by joining the DNA coding sequences and expressing (and purifying) the fusion protein (e.g. in the case of IgG, either the heavy or the light chain could be attached to C3 and co-expressed with C3, or both chains could be combined within one complete fusion polypeptide), or by incorporation of specific coding sequences (eg. for "leucine zipper"-like domains) to the DNA of both fusion partners (eg. modified C3 and specific antibody) such that the expressed products, when mixed together, self-associate to form stable conjugates. The fusion protein could then be administered locally or into the general circulation.
- 30 Liposomes (bearing the antibody on the surface with the modified protein either on the surface or inside the liposome) and/or virions (e.g. engineered to express the proteins on their surface) could also be used for co-delivery of antibody and modified protein. This

strategy could be used directly, alone or in combination with other treatments, at any stage in the disease process. It may be particularly appropriate for use in eliminating any cancerous cells left in the circulation after surgical removal of a tumour. The antibody-modified protein conjugates could also be used *ex vivo* to eliminate pathogenic tissue. For example to kill leukaemic cells from an extracted bone-marrow and then returning the remaining healthy cells to the patient.

Alternatively lymphocytes that do not match the MHC types of the recipient could be eliminated from a bone marrow prior to transplantation. Also the modified protein could be linked to an antigen, and this combination could be used, either *in vivo* or *ex vivo*, to attack lymphocytes of undesirable reactivities (e.g. against transplant or self tissue).

The same technology would be applicable to treating other species, using either a human modified protein derivative, or a similar analogue tailor-made for that species.

Preferred features of each aspect of the invention are as for each other aspect *mutatis mutandis*.

The invention will now be described by way of the following examples, which should not be construed as in any way limiting the invention. The examples refer to the accompanying drawings in which:

Figure 1: shows the predicted protein sequence of human C3 as encoded in PC3;
(using the standard one letter amino acid code)

Figure 2: shows the cDNA sequence in PC3;
(using the standard one letter deoxynucleotide code
for the sense strand, written 5'-3').

5 Figure 3: shows a visualisation of modified proteins
of the invention.

10 Figure 4: shows the effect of various mutations to
human C3 which replace Arg 1303 or Arg 1320 on
factor I-mediated cleavage at these sites.

N.B.

1. [35S]-biosynthetically labelled samples.
2. Reactions performed at normal ionic strength.
3. Immunoprecipitated with anti-C3.
- 15 4. SDS-PAGE under reducing conditions.
5. Autoradiography.

20 Figure 5: shows enhanced resistance of human C3
incorporating the Arg 1303 -> Gln 1303 mutation to
inactivation by factors I and H.

25 Figure 6: shows an analysis of the cleavage of a C3
convertase mutated at amino acid residues 752-754
and 758-760.

30 This is a photograph of a Western Blot developed
from a 7.5% polyacrylamide SDS-PAGE gel (reducing
conditions), after electrophoretic transfer onto
nitrocellulose, probing with a sheep anti-human C3
antibody, and development with horse-radish-
peroxidase-coupled anti-sheep Immunoglobulin
antibody and Enhance ChemiLuminescence (method and
detection reagents from Amersham, U.K.) recorded on
X-ray film. The cleavage reactions and detection

procedure were performed as described in Example 4 with reference to the results shown in Figure 3.

Key:

- 5 Tracks 1-4: wild-type C3 (expressed in COS cells)
Tracks 5-8: Mutant C3 (residues 752-754 changed to Gly-Ser-Gly and residues 758-760 also being changed to Gly-Ser-Gly) (expressed in COS cells)
- 10 Tracks 1,5: no addition
Tracks 2,6: + CVFBb
Tracks 3,7: + factors H + I
Tracks 4,8: + CVFBb + factors H + I
- 15 The bands indicated by arrows are:
A: C3 alpha-chain
B: C3 alpha'-chain
C: C3 beta chain
D: 68 kDa cleavage product of C3 alpha'-chain
20 E: IgG heavy chain

Figure 7: shows an analysis of the cleavage of radiolabelled factor B by factor D, in the presence of wild-type and mutant C3's (C3i's)

25 A photograph of the autoradiograph of the SDS-PAGE gel is shown. All samples contained factor D and ¹²⁵I-labelled factor B, and were incubated for 3 hours at 37°C.

30 The samples in the numbered tracks also included:
1. Buffer alone
2. 1/125 wild-type C3
3. 1/25 wild-type C3

4. 1/5 wild-type C3
5. 1/25 mutant C3 (residues 1427 Gln, 1431 Asp and
1433 Gln)
6. 1/5 mutant C3
5 7. undiluted mutant C3

The bands indicated by arrows are:

- A. Uncleaved ^{125}I -labelled factor B (93 kDa)
B. 60 kDa cleavage product ("Bb")
10 C. 33 kDa cleavage product ("Ba")

Figure 8: shows an SDS-PAGE study illustrating the formation of a conjugate between C3i and IgG.

- 15 This is a Coomassie stain of a 4% acrylamide SDS-PAGE gel run under non-reducing conditions. The numbered tracks contain samples of:
1. PDP-IgG—
2. C3i
20 3. PDP-IgG + C3i reaction mixture

Indicated by arrows are:

- A. Probably C3i-IgG conjugate (350 kDa)
B. C3i (200 kDa)
25 C. IgG (150 kDa)

Figure 9: demonstrates that conjugate targets C3 convertase activity against sheep erythrocytes.

- (This graph shows the % lysed sheep erythrocytes after coating with dilutions of either the C3i-IgG conjugate, PDP-IgG or C3i followed by washing, generation of C3 convertases with properdin and factors B and D, and finally development of lysis by NGPS in CFD/EDTA, as described in the methods. Only
30

the conjugate produces lysis, and this lysis is dose dependent.)

5 The following standard methods and definitions are applicable to all the examples.

10 All complement components referred to are of human origin, unless otherwise specified, using standard terminology for all proteins and their derived fragments (e.g. as contained in reference [15]). In addition the term "C3i" refers to any molecular form of C3 without an intact thiolester bond, but retaining the C3a polypeptide on the alpha chain.

15 The human C3 cDNA and coding sequence are numbered as shown in FIGURE 2, using the numbering used in the EMBL nucleotide data base (derived from reference [2]). The sequence shown is that of our construct ('PC3'), which lacks the first 11 nucleotides of the 5' untranslated region reported in reference [2], and hence the first base is numbered 12. The putative initiation codon is nucleotides number 61-63, the codon for the amino-terminal serine residue of the beta-chain is nucleotides 127-129, and the codon for the amino-terminal
20 serine residue of the alpha-chain is nucleotides 2074-2076.

30 The protein sequence is numbered according to the precursor sequence as shown in FIGURE 1, which is a predicted translation of the DNA sequence in Appendix 1 (amino acids 1-22 are expected to comprise a signal sequence that is removed during biosynthesis, and amino acids 668-671 are expected to be removed when the precursor is cleaved into the alpha and beta chains).

The following abbreviations have the following meanings;
CVF cobra venom factor; ELISA, Enzyme-linked
immunoabsorbant assay; *E. coli*, *Escherichia coli*; kb,
kilobase; HSV-1, herpes simplex virus type 1; PBS,
phosphate-buffered saline. COS-1 is a cell line derived
from monkey kidney cells. The following are restriction
endonucleases:- AflIII, DraI, DraIII, EcoRI, EcoRV,
HindIII, NaeI, NheI, XbaI.

Standard methods

Methods for standard molecular biological procedures such
as plasmid isolations, agarose gel electrophoresis, and
DNA ligations can be found in reference [21]. Double
stranded DNA was sequenced using the 'Sequenase version
2.0' kit supplied by 'United States Biochemicals'. C3
expression was measured by an ELISA assay using plastic
plates pre-coated with affinity-purified polyclonal sheep
anti-human C3 to which samples of culture supernatant
were added. Bound C3 was detected with a monoclonal rat
antibody to C3 conjugated to alkaline phosphatase, and
the chromogenic substrate, p-nitrophenol phosphate.
Assays were calibrated with purified human plasma C3.

Methods for purification of complement proteins and CVF,
and for the preparation of affinity purified anti-C3
antibodies used in the analysis can be found in reference
[28]. Equivalent reagents can also be purchased from
Sigma chemical company LTD.

C3 cDNA coding sequence

Our C3 cDNA coding sequence was constructed from two
segments isolated from a random-primed human liver cDNA

library carried in the vector pGEM4 (Promega). Five oligodeoxynucleotides, corresponding to known segments in the human C3 coding sequence, were radiolabeled with T4 polynucleotide kinase and [γ -32P]ATP and used to probe filter transfers of the library from agarose plates. Two clones containing inserts of approximately 4 kb were isolated. Restriction endonuclease digestion, hybridisation to specific oligodeoxynucleotide probes and partial sequence analysis demonstrated that one of these ('A13') included the 5'-end of the 5.1kb message, whereas the other ('B44') extended to the 3'-end.

These inserts therefore overlapped by approximately 3 kb, including a unique *EcoRI* restriction enzyme site. The incomplete 5' section of A13 was cut out with *EcoRI* and *NheI*, and replaced with the complete segment isolated from B44 by digestion with *EcoRI* and *XbaI*. Both pieces were purified by gel electrophoresis in low-melting point agarose before ligating together with T4 DNA ligase to produce a vector ('PGC3') containing 5.1 kb of DNA encoding the entire C3 precursor protein.

Linker sequences 5' to the C3 coding region contained two ATG's which are potential false translation start sites. These were therefore removed by gapped-plasmid mutagenesis, as described in the method of example 1, using an oligodeoxynucleotide PL-ATC-3 (tagggagacc ggaagcttgc cctctccctc tgtccctctg t) that deleted approximately 50 base pairs of linker/adaptor DNA, without altering the C3 coding sequence. This mutated vector, 7.7kb containing 5.1kb of C3 cDNA sequence plus 2.6 kb of sequence from the pGEM4 vector (Promega) is referred to as PC3.

The C3 coding region of the PGC3 plasmid was completely sequenced and revealed only four differences from a previously published human C3 ("S" allele) cDNA sequence [2].

5 (i) the changes C2481->G, and C2805->T do not alter the coding;

(ii) T1001->C encodes the previously described HAV 4-1-
10 (Leucine314->Proline) polymorphic form [20]; and

(iii) G2716->A encodes Valine886->Isoleucine, that has not been previously reported in human C3, although Ile is found in this position in mouse and rat C3.

15 Our sequence includes start and stop codons, with a complete signal sequence and should, therefore, encode functional C3.

20 Levels of up to 1.7 μ g/ml expressed wild type C3 in culture supernatants of COS-1 cells (transfected using lipofectamine and the pcDNA3 (Invitrogen) expression vector) have been detected by ELISA. No detectable C3 was produced by cells transfected with pcDNA3 vector alone. Furthermore, analysis of the expressed product by
25 cleavage reactions followed by immunoprecipitation, SDS-PAGE and immunoblotting demonstrated that:

(i) the primary translation product had been correctly processed into the mature two-chain form;

30 (ii) this product was, like native C3, cleavable to C3b by C3 convertase (CVFBb); and

(iii) the expressed protein was, like native C3, not

cleavable by factor H plus I, but became cleavable after conversion to C3b by C3 convertase enzyme. This confirms that our starting plasmid can be translated into functional C3.

5

For an alternative description of a construction and expression of a C3 coding sequence see reference [25].

EXAMPLE 1: Production of C3 that has the arginine residues at both factor I cleavage sites (amino acid positions 1303 and 1320) converted to glutamine residues to prevent cleavage of the C3b fragment by factor I.

a) Mutagenesis

15

Mutagenic oligodeoxynucleotides used were QRI1 (caactgcccagccaaagctccaagatcacc), QRI2 (gccagcctcctgcaatcagaagagaccaag), and AFL4149 (taataaattcgaccttaaggtcaccataaaac), as well as the corresponding antisense oligodeoxynucleotides QRI1n (ggatgatcttggagctttggctgggcagttg), QRI2n (cttgggtctcttctgattgcaggaggctggc) and AFL4149n (gttttatggtgaccttaaggtcgaatttatta).

QRI1 and QRI1n specify the replacement of arginine for glutamine at the factor I cleavage site at amino acid residue 1303 in the C3 precursor sequence (by changing G3968C3969 to AA in the cDNA sequence), and QRI2 and QRI2n effect the same substitution at the factor cleavage site at amino acid residue 1320 (by changing nucleotide G4019 to A).

30

AFL4149 and AFL4149n introduce a cleavage site for the restriction endonuclease *Afl*III at position 4149 in the

cDNA sequence (by changing C4149 to T) without altering the encoded amino acid sequence. These two primers were used as markers, allowing successful mutagenesis to be identified on the basis of cleavage of the DNA product by *AflIII*.

Mutagenesis was effected using the 'gapped plasmid' method. A batch of PGC3 ('UPGC3'), enriched in uridine in place of thymidine, was prepared by growth in *E. Coli* strain TJ236 in the presence of 0.25 $\mu\text{g/ml}$ uridine. This plasmid was digested with *SmaI* and the 7.2kb product ('US1') agarose gel purified to remove a 0.5kb fragment from the C3 sequence (residues 1463-1947). The other component of the gapped plasmid ('DN2') was prepared by digesting PGC3 with *DraIII* plus *NaeI* and purifying the 5.1kb piece twice by agarose gel electrophoresis. 200ng DN2 was mixed with approximately 500ng US1 in 50 μl H₂O, heated to 100°C and cooled slowly to below 50°C, before adding 20 μl to 25 μl of 2XT7 buffer (100mM Tris/HCl/pH 7.4/ 14mM MgCl₂, 100mM NaCl, 2mM dithiothreitol, and 1mM each of ATP, dATP, dCTP, dTTP and dGTP) plus 10nmol of each 5'-phosphorylated mutagenic primer (one reaction used QRI1, QRI2 plus AFL4149, another reaction used QRI1n, QRI2n plus AFL4149n). The mixtures were reheated to 70°C for 5 min and cooled slowly (over 30-60 min) to 20°C. At 0°C, 10 units of T7 DNA polymerase plus 80 units T4 DNA ligase are added. The mixture (total volume 50 μl) was incubated first at 0°C, for 5 min, then at room temperature for 5 min, and finally at 37°C for 3 hours. 1 μl of each mixture was used to transform 100 μl supercompetent XL1 *E. Coli* (Stratagene) according to the manufacturer's instructions.

Ampicillin resistant colonies were screened for *AflIII*

cleavage, and successful mutants were grown up in 100ml cultures from which the plasmids were isolated and sequenced (using a sequencing primer C3pa-3876, cttcatggtgttccaagcct, matching nucleotides 3876-3895 of C3 cDNA) to characterise mutations at the factor I cleavage sites.

For an alternative protocol for "gapped plasmid" mutagenesis see references [26,27].

b) Transfer of mutant DNA to eukaryotic expression vector

The C3 coding fragments from mutant plasmids were excised by double digestion with *HindIII* and *NaeI*. *DraI* was also included to incapacitate the residual plasmid. The C3 coding sequence was agarose gel purified and ligated into pcDNA3 vector (Invitrogen) that had been linearised with *HindIII* and *EcoRV* enzymes and dephosphorylated with calf intestinal phosphorylase. Ligation mixtures were used to transform supercompetent XL1 *E. coli*, which were then plated onto culture plates containing ampicillin.

A random selection (three or four) of ampicillin resistant colonies were grown up in 2-3ml cultures and small scale isolation of the plasmid DNA. The plasmids containing the correct insert were identified by digestion of the plasmid DNA with restriction endonucleases *EcoRI*, *HindIII* and *AflII*. The corresponding colonies grown up in 100ml cultures and the plasmids purified by the standard procedure. These mutants were originally constructed from PGC3 and so retained the two ATG's 5' to the coding region. This region (plus the 5' 3kb of the C3 coding sequence) was therefore excised with

*Hind*III plus *Eco*RI and replaced by ligation of the same segment cut out of PC3. These reconstructed vectors were prepared by the standard procedure and used for transfection of COS cells.

5

c) Expression of wild-type and mutant C3's

Mutants and wild-type C3 were transiently expressed from plasmids transfected into COS-1 cells using lipofectamine® (GIBCO) according to the manufacturer's instructions. Typically, $1-1.5 \times 10^5$ cells per well of a standard 6 well culture plate were transfected with 2-4 µg of plasmid using 9 µl of lipofectamine reagent. Supernatants were assayed for C3 secretion, and typical yields of 0.3-1.7 µg per ml supernatant were obtained 3-6 days after transfection.

Results

20 a) Generation of mutants

The following mutants, named according to the mutagenic oligodeoxynucleotide sequences that have been incorporated, have so far been isolated:-

25 (i) 3 mutants with both QRI1 and QRI2 mutations plus AFL4149: C3M-26, C3M-58 and C3M-61;

(ii) 1 mutant with QRI1 and QRI2 but without AFL4149: C3M-8; and

30

(iii) 1 mutant with QRI2 and AFL4149, but without QRI1: C3M-51 (used in example 3)

b) Validation that functional effects were due to the

mutations specifically introduced at the factor I cleavage sites

5 Sequencing has confirmed the absence of other alterations in 178-350 bases around the mutated region of each mutant. The sequence of one mutant produced by this procedure, C3M-51 (see example 3), has been analysed throughout the entire 'gap' (bases 2463-5067) used in mutagenesis, and no other deviations from the wild-type
10 sequence were found.

Furthermore, representative sequencing of a total of 2922 bases from all mutants have not revealed any single point mutations that could have been caused by polymerase-mediated errors. The expressed mutants all displayed the
15 two-chain structure and cleavage by C3 convertases characteristic of native C3. In summary, the mutants used are unlikely to contain any unwanted changes although they have not been completely re-sequenced.

20 EXAMPLE 2: Production of C3 that has the arginine residue at one factor I cleavage site (amino acid position 1303) converted to a glutamine residue

25 The procedure of Example 1 was followed except that only mutagenic oligodeoxynucleotides AFL4149 plus QRI1 or AFL4149n plus QRI1n (i.e. no QRI2 or QRI2n), were used in mutagenesis.

30 Results

a) Mutants obtained

2 mutants with QRI1 and AFL4149 but without QRI2 were isolated: -C3M-I23,27. The mutant C3M-I23 was expressed,

as described in Example 1.

5 This protein was cleavable by CVFBb. The C3b-like product was relatively (compared to the wild-type) resistant to cleavage at position 1303 by factors I and H, but could still be cleaved at position 1320. This C3b derivative is therefore partially resistant to factor I.

10 EXAMPLE 3: Production of C3 that has the arginine residue at one factor I cleavage site (amino acid position 1320) converted to a glutamine residue

15 The procedure of Example 1 was followed except that only mutagenic oligodeoxynucleotides AFL4149 plus QRI2 or AFL4149n plus QRI2n (i.e. no QRI1 or QRI1n), were used in mutagenesis. In addition, the method used in example 1 also yielded one mutant with QRI2 and AFL4149, but without QRI1. —

20 Results

a) Mutants obtained

25 3 mutants with QRI2 and AFL4149 but without QRI1 were isolated: -C3M-51, C3M-Q2, C3M-Q13. The mutant C3M-51 was expressed, as described in Example 1. This protein was cleavable by CVFBb. The C3b-like product was not readily cleaved at position 1320 by factors I and H, but it could still be cleaved at position 1303. This C3b derivative is therefore partially resistant to factor I.

30 EXAMPLE 4 Analysis of the functional effects of mutations.

Supernatants (100-400 µl) from transfected COS cells were

incubated at 37°C for 2h with:

COS cells were transfected with pcDNA3 carrying inserts of:-

5

- 1) the unmutated C3 sequence;
- 2) mutant C3M-I23 (encoding Arg¹³⁰³->Gln);
- 3) mutant C3M-26 (encoding Arg¹³⁰³->Gln, Arg¹³²⁰->Gln); and
- 10 4) mutant C3M-51 (encoding Arg¹³²⁰->Gln).

200 µl of the culture supernatants, taken 3 days after transfection, were pretreated with 2 mM phenylmethanesulphonyl fluoride (0°C, 15 min) and then
15 incubated at 37°C for 2 hours with the following:

- A) no addition;
- 20 B) preformed C3 convertase, CVFBb (10 µl from 200 µl containing 6.6 µg CVF, 100 µg factor B and 1.4 µg factor D in phosphate-buffered saline (PBS) containing 10 mM MgCl₂, preincubated at 37°C, 15 mins);
- C) factors H (5 µg) and I (1 µg); and
- 25 D) CVFBb plus factors H and I.

These were then immunoprecipitated by adding 0.6 µg affinity-purified sheep anti-human C3 immunoglobulin at
30 room temperature and after 1 hour adding 20 µl a 5% suspension of washed formalin-fixed Group C *Streptococcus* sp. cells (protein G) (Sigma). After 45 min at room temperature the particles were washed once in PBS, 5 mM NaN₃, and once in 20 mM Tris/HCl, 137 mM NaCl, 0.1% (v/v)

Tween 20, pH 7.6 before eluting in 1% SDS/2% 2-mercaptoethanol (90-100°C, 5 min). These eluates were separated by SDS-PAGE, electroblotted onto nitrocellulose and the C3 bands detected by probing with affinity-purified sheep anti-human C3 immunoglobulin followed by horse radish peroxidase-coupled donkey anti-sheep immunoglobulin (Sigma) and detection using the "Enhanced Chemiluminescence" substrates supplied by Amersham. A photograph of a 2 minute exposure to X-ray film is shown. The visible C3-derived bands are indicated by labelled arrows, and the individual samples (1-4, A-D) are those just described. (The prominent band of about 50 kDa (between the 46 and 68 kDa bands) present in all samples is the heavy chain of the IgG used in the immunoprecipitation and detected by the horse radish peroxidase-coupled donkey anti-sheep immunoglobulin.).

Results (see Figure 3)

1. All the untreated samples (1-A, 2-A, 3-A, 4-A) contain bands of the correct migration for alpha and beta chains of C3, indicating that all the mutants are expressed, and post-translationally processed correctly. The presence of 43 or 46 kDa bands in these samples indicates the presence of some factor H + factor I-like activity in the culture medium. Spontaneous hydrolysis of C3 during the 3 day biosynthetic period produces C3i which is cleaved by this activity. In the unmutated C3 this generates bands of 43 kDa and 75 kDa (the 75 kDa band is invisible because (i) it is hidden by the 75 kDa beta chain, and (ii) the antibody used to develop the western blot has very little activity towards this portion of the C3 alpha chain:- its presence was subsequently confirmed by reprobing with a rat monoclonal

antibody, "Clone-3", that is specific for this region). The addition of factors H and I without CVFBb (1-C, 2-C, 3-C, 4-C), did not cleave the remaining C3 indicating that this represented active C3 (thiolester intact).

5

2. The unmutated C3 (1) is cleaved by CVFBb and the C3b product is further cleaved by endogenous enzymes in 1-B or added factors H and I in 1-D. The 43 kDa band indicates cleavage at Arg¹³²⁰, and the 68 kDa band (visible in longer exposures) indicates cleavage at Arg¹³⁰³.

10

3. The mutant C3M-I23 (Arg¹³⁰³->Gln) was cleavable by CVFBb and the product was relatively resistant to endogenous factor H and I-like activity (2-B), with distinct amounts of alpha' chain (C3b) persisting, but was still cleavable when extra factor H and I were added (2-D). The 43 kDa product indicates cleavage at Arg¹³²⁰, (a faint band at 71 kDa representing the other fragment of the alpha' chain could be seen in longer exposures) but no 68 kDa band was present, showing that this mutant is resistant to cleavage at the mutated Gln¹³⁰³.

15

20

4. The mutant C3M-26 (Arg¹³⁰³->Gln, Arg¹³²⁰->Gln) was cleavable by CVFBb and the C3b-like product (alpha') was resistant to endogenous factor H and I-like activity (3-B). It was also very resistant to the additional factors H and I (3-D) in comparison with the unmutated C3 (1) and other mutants (2 and 4). There was a small amount of 46 kDa product indicating some cleavage at the mutated Gln¹³⁰³ (the accompanying 68 kDa fragment was also visible on longer exposures). There was little or no detectable 43 kDa that would correspond to any cleavage at Gln¹³²⁰. Therefore the Arg->Gln mutation at position 1303 is less

25

30

effective than that at position 1320 at preventing cleavage by factor I. (This slow residual cleavage might also be occurring in the mutant C3M-I23 (Arg¹³⁰³->Gln), but the 46 kDa intermediate is probably being rapidly processed to 43 kDa by further cleavage at the unmutated Arg¹³²⁰.)

5. The mutant C3M-51 (Arg¹³²⁰->Gln) was cleavable by CVFBb and the product was cleaved by endogenous factor H and I-like activity (4-B), and by additional factor H and I (4-D). The 46 kDa product (and faint 68 kDa band) indicates cleavage at Arg¹³⁰³. However, the absence of a 43 kDa band indicates that it is not cleaved at the mutated Gln¹³²⁰.

Example 5 Comparison of various amino acid substitutions at position 1303

1. Introduction

The previous examples described mutations of arg 1303 and arg 1320 to glutamine residues. Both mutations imparted resistance to cleavage at those positions by factor I. However, there was a small but detectable degree of cleavage at gln 1303. Therefore a number of other amino acid substitutions at this position have been made and tested. Cleavage occurs, in decreasing order of efficacy when residue 1303 is: Arg > Tyr > [Cys or Trp] > Gln > [Glu or Gly]. These results are unexpected because (i) all known naturally occurring human factor I-mediated cleavages occur C-terminal to arginine residues, so it would have been deduced that the enzyme had a requirement for arginine; and (ii) if it did cleave at other residues

one would predict that they would have to be electrostatically similar to arg, i.e. a basic residue (lys or his), (e.g. trypsin selectively cleaves C-terminal to arg, lys or his), so one could not have predicted cleavage of the tyrosine substitution.

Therefore substitution of arg 1303 with glycine or glutamic acid is preferred for the purpose of creating a derivative of C3 resistant to inactivation by factor I.

2. Methods

2.1 Mutagenesis: the degenerate mutagenic primer used was:

caactgcccagc(gt)(ag)(cg)agctccaagatcacc (letters in brackets indicate mixture of bases at that position). Mutants were constructed either by the gapped-plasmid method (as described in the earlier examples), or by the "megaprimer method" (V. Picard et al, Nuc Acid Res 22:2587-91, (1994)), in which the upstream primer was caccaggaactgaatctagatgtgtccctc and the downstream primer was gttttatggtgaccttaaggtcgaatttatta. All mutations were performed on templates in which the C3-encoding DNA had already been mutated such that amino acid residue 1320 was glutamine, and a restriction site for AflII had been introduced at position 4149 (as described in the earlier examples) and were confirmed by DNA sequencing.

2.2 Expression: mutants were expressed in COS cells using the pcDNA3 vector as described in the earlier examples, biosynthetically labelled with [³⁵S]methionine in serum-free medium.

2.3 Assay: the supernatants were treated with CVFBb (formed by reaction of CVF with factors B and D in magnesium-containing buffer) and factors H and I followed by immunoprecipitation with anti-C3 and separation by SDS-polyacrylamide gel electrophoresis performed under reducing conditions (as described in the earlier examples). The gel was fixed, treated with Amersham "Amplify" reagent, dried and exposed to autoradiography film to yield the result shown in the figure.

3. Results

Factor I-mediated cleavage at position 1303 (site 1), without cleavage at 1320 (site 2) (where this has been mutated to glutamine) produces bands of 46 and 68 kDa. It can be seen that cleavage occurs in the order: arg(R) > tyr(Y) > cys(C) and trp(W) > gln(Q) > gly(G) and glu(E). The wild-type (arginine at both positions) is cleaved at both positions to produce fragments of 43 (too small to be visible on this gel) and 68 kDa.

4. Figure

The results are shown in Figure 4. The residues at site 1 (position 1303) and site 2 (1320) are indicated above the respective tracks.

Example 6 Demonstration of enhanced resistance to inactivation by factors I and H after mutation of arg 1303 to gln

1. Introduction

The earlier examples demonstrated that conversion of either arg 1303 or arg 1320 to glutamine made that site resistant to cleavage by factor I. Mutation of both sites makes a molecule that is resistant to cleavage at either site. Here, we further demonstrate that mutation of arg 1303 to gln alone (without alteration to arg 1320) results in a considerable resistance, compared to the wild-type, to functional inactivation by factors I and H.

2. Method

2.1 Expression: The preparation of the arg 1303->gln mutation was described in an earlier example. This was transfected into CHO (a common laboratory cell line derived from chinese hamster ovary cells) by the calcium phosphate method, and stable transfectants selected on the basis of resistance to G418 ("Geneticin" available from Sigma). Cell culture supernatants were collected, and the expressed C3 was partially purified by sodium sulphate precipitation (10-20% (w/v) fraction), and ion-exchange chromatography on Q-sepharose and mono-Q

sepharose (A W Dodds *Methods Enzymol* 223: 46 (1993)).

2.2 Assay: Sheep erythrocytes were coated with SO16 monoclonal antibody (R A Harrison and P J Lachmann *Handbook of Experimental Immunology* 4th Edition chpt. 39 (1986)) and 4.4 ml of a 5% (v/v) suspension was then incubated with approximately 10 μ g C2, 24 μ g C4 and 1 μ g C1 (purified human components) for 10 min at 37°C in CFD (R A Harrison and P J Lachman *supra*). 0.8 ml of this mixture was then incubated for 105 min with 0.25 ml containing the semi-purified mutant or wild-type C3 and EDTA to a final concentration of 12.5 mM. The cells were then washed in CFD and used in CFD containing 0.1% (w/v) gelatin (CFD-gel). Radioligand binding with [125 I]-labelled clone 4 monoclonal anti-C3 antibody was used to confirm that similar amounts of wild-type or mutant C3b were deposited.

For the assay, 40 μ l of a 5% suspension of cells was diluted in 250 μ l CFD-gel and 50 μ l aliquots were incubated with 50 μ l CFD-gel containing dilutions of factors I and H to final concentrations of 100, 10, 1 and 0 μ g/ml each, at 37°C for 30 min. 0.9 ml of CFD was then added, the cells pelleted by centrifugation and washed twice more with 1 ml of CFD each time. The cells were then resuspended in 100 μ l CFD-gel containing 100 μ g/ml factor B, 100 μ g/ml properdin, 1 μ g/ml factor D and 0.3 mM NiCl_2 . After 10 minutes at 37°C, 0.9 ml of CFD containing 10 mM EDTA and 2% (v/v) normal guinea-pig serum. After a further 30 min at 37°C, unlysed cells were pelleted by centrifugation, and the degree of lysis determined by measuring the absorbance of the supernatant at 412 nm. The absorbance equivalent to 100% lysis was determined from an aliquot of cells lysed in water, and

hence the percentage lysis was calculated.

5 This assay measures the ability of deposited C3b to form
a functional C3bBbP convertase. Conversion to iC3b
prevents convertase formation and subsequent lysis in
serum/EDTA.

10 3. Results

The result shown in the figure indicates that more than
ten times as much factor I and factor H are required to
abrogate the hemolytic activity of the arg 1303->gln
mutant, when compared to the wild-type. This mutation is
15 therefore advantageous for the creation of a derivative
of C3 whose C3b product is resistant to inactivation by
factors H and I. The effect could either be due to the
greater resistance to cleavage at position 1303 (when arg
is mutated to gln), or to greater resistance to cleavage
20 at position 1320 when cleavage can first take place at
position 1303.

4. Figure

25 The results are shown in Figure 5. The x-axis indicates
the concentration of factors H and I. Q1 represents the
arg 1303->gln mutation. % lysis is measured as described
in the methods.

Discussion

The essential features of Human C3, with respect to modified variants described herein are as follows:-

5

(i) The molecule has a functionally C3b-like derivative in that it can combine with functionally active human factor B, which can then be cleaved by human factor D to form an enzyme capable of cleaving human C3.

10

(ii) The amino acid sequences of derivatives are more homologous to C3 from humans than to C3 from any other species for which a sequence is presently known, or to any other presently known protein sequence. Structural features of C3 present in wild-type protein, but not necessarily in modified derivatives, include the following:-

15

(a) The DNA coding sequence and translated protein sequence for the variant of human C3 used in the examples of the invention described herein are given in Figures 2 and 1 respectively. This protein sequence differs from the published sequence [2] at just two amino acids (details are given in the examples). It is assumed that many more variations are compatible with C3 function, even though most will not be present in the population.

20

25

(b) The primary translation product is proteolytically processed into two disulphide-linked chains, alpha (residues 672-1663) and beta (residues 23-667), with removal of the signal sequence (residues 1-22).

30

(c) The mature protein contains a thiolester bond between residues Cys1010 and Gln1013.

(d) C3 convertases cleave C3 to remove C3a (residues 672-748). This reaction is followed by breakage of the thiolester bond.

5 (e) In the presence of factor H, factor I cleaves C3b between residues Arg1303 and Ser1304, and between Arg1320 and Ser1321.

Modifications made to the native C3 molecule

10

Replacement of Arg1303 by Gln

15 This modification is at one site of cleavage of C3b by factor I. The effect is to reduce the rate of cleavage by factor I at this position. The change to glutamine was selected to take away the positive charge of the arginine, which is likely to be important for the serine protease activity of factor I, while retaining a hydrophilic character and a similar side-chain size that
20 should minimise any disruptions to the tertiary protein structure. Evidence supporting this presumption is that the mutation did not prevent processing into a two-chain structure, formation of a thiolester or cleavage of C3 by C3 convertase. Mutation of Arg1303 to another amino acid
25 can achieve a similar or even a superior effect, as demonstrated in Example 5.

30 It may also be possible to reduce this cleavage by mutating Ser1304 (the other side of the cleavage site) or other residues involved in the enzyme-substrate interaction.

Replacement of Arg1320 by Gln

5 This modification is at the other site of cleavage of C3b
by factor I. The effect is to drastically reduce
(virtually abolish) the rate of cleavage by factor I at
this position. The change to glutamine was made on the
same criteria described above, and this mutation also did
not prevent processing into a two-chain structure,
10 formation of a thiolester or cleavage of C3 by C3
convertase. Again, mutation to another amino acid may
achieve the same effect, as may mutation of Ser1321 or
other residues involved in the enzyme-substrate
interaction.

15 When in combination the two mutations, Arg1303-Gln and
Arg1320-Gln, protect the C3b from inactivation and hence
maintain its ability to form part of an active C3bBb
convertase. Other mutations (including combinations of
mutations) that abolish both cleavage reactions could
20 also be used (for example Arg 1303 Glu or Arg 1303 Gly
could be used in combination with Arg 1320 Gln).

EXAMPLE 7 Various mutations that reduce the interaction
of C3b/C3i with factor H

7.1 Introduction

5

10

15

20

25

30

Other laboratories have produced evidence based either on the effects of synthetic peptides (Ganu, V.S. and Muller-Eberhard, H.J., 1985, *Complement* 2:27; Becherer, J.D. et al., 1992, *Biochemistry* 31: 1787-1794), or limited mutagenesis (Taniguchi-Sidle, A. and Isenman, D.E., 1994 *J. Immunol.* 153: 5285-5302) to suggest that the residues 752-761 in the primary sequence of the C3 transcript (see figure 1) could be involved in the interaction with factor H. However, other published evidence suggests that only residues 767-776 are involved in the interaction with factor H, whereas residues 752-761 are important for the interaction with factor B (Fishelson, 1991, *Mol. Immunol.* 28:545-552). We surmised that more extensive mutagenesis of this region might reduce the affinity for factor H and therefore be desirable for the objective of creating a C3 derivative that is resistant to factor H. Furthermore, we guessed that the important residues to mutate could be the prominent acidic residues (aspartic and glutamic acids) and that it would be desirable to change them to neutral residues less likely to mediate strong interactions. In this example we changed residue 752-754 from Asp-Glu-Asp to Gly-Ser-Gly, in combination with changing residues 758-760 from Glu-Glu-Asn to Gly-Ser-Gly. The product displayed reduced cleavage characteristics consistent with a reduction in the susceptibility to factor H. This provides evidence that C3 can be modified to reduce the binding of factor H, and hence the susceptibility to factors H and I. These modifications are desirable for the creation of a

C3 convertase that is stable under physiological conditions.

7.2 Method

5

The methods of mutagenesis, expression and analysis have been described in the earlier examples. The mutagenic oligonucleotide that was synthesised had the sequence:
agtaacctgggttcgggcatcattgcaggatcgggcatcgtttcc.

10

7.3 Results

The results of cleavage reactions are shown in Figure 6. These indicate that:

15

1. Addition of CVFBb⁻ to wild-type C3 results in elimination of the alpha chain (track 2) because the C3b that is formed is susceptible to the low concentrations of factor I and H in the culture supernatant. C3i that has been formed during expression or this subsequent incubation has been broken down to iC3i in the same way. Addition of exogenous factors I and H (tracks 3 and 4) are therefore no different from tracks 1 and 2 respectively, because the medium itself contains sufficient factor H and I activity to effect complete cleavage.

20

25

30

2. In contrast, treatment of the mutant C3 with CVFBb (track 6) does not result in disappearance of the alpha chain. There is some generation of alpha', corresponding to C3b, but some or all of this remains, indicating that the persistence of alpha chain is not merely the result of a failure of cleavage by CVFBb. The remaining uncleaved alpha chain in track 2 may therefore represent

C3i that has not been cleaved by the endogenous activities of factors H and I, although it is also possible that some of this represents native C3 persisting if the mutant has acquired a partial resistance to CVFBb. Addition of high concentrations of exogenous factors H and I (track 7 and 8) does produce depletion of alpha and alpha' chains, indicating that (i) the mutant is not completely resistant to these factors, and (ii) the alpha chain uncleaved by CVFBb in track 2 is predominantly derived from C3i (which is cleavable by factors H and I but not by CVFBb) rather than from native C3 (which is cleavable by CVFBb but not by factors H and I). Still not all the alpha chain is cleaved, even in track 8, probably because of the resistance to factors H and I.

Therefore mutation of residues 752-754, and residues 758-760 can generate a C3 molecule that can still be cleaved by C3 convertases, but is partially resistant to the actions of factors H and I. In view of other published data, this is most probably because the mutations have modified a region that is involved in the interaction with factor H and hence have resulted in a reduced affinity for factor H.

EXAMPLE 8 A site in C3 that can be mutated to modify the interaction of C3i with factor B

8.1 Introduction

The previous examples have demonstrated that mutations to C3 can modulate the interactions with factors H and I. In order to discover other sites in C3 that might

interact with factor B, we compared the known sequences of C3 molecules from different species, as well as with available sequences for C4 and other homologous proteins. We identified the region corresponding to residues 1427-1433 of human C3 that might be involved in C3 and C4 specific functions. This could include interaction with factor B (or its homologue, C2, in the case of C4), but not necessarily because other potential functions include thiolester formation, conversion into C3b (or C4b form), interaction with substrate C3 and/or C5 in convertase activity and interaction with factor I and its cofactors. Therefore selected residues were mutated to the corresponding residues (based on sequence alignments) found in another homologous protein, in this case human C5. Thus residue 1427 was changed from an Arg to a Gln, residue 1431 from a Lys to Asp, and residue 1433 from a Glu to a Gln. The resulting mutant was found to be susceptible to cleavage by C3 convertase (CVFBb) and the C3b product was cleavable by factors H and I. However, this mutant did not support the conversion of factor B to Bb plus Ba, which is dependent on the binding of factor B to C3i (or C3b). Therefore we have evidence that mutation of this region has diminished the interaction with factor B. Whilst this is undesirable for the generation of a super-active C3 convertase, it does provide an indication that other modifications to this region of C3 will also alter the interaction with factor B, and some of these will probably increase the affinity. As a consequence such mutations may also increase the stability and activity of the bimolecular convertase enzyme, C3bBb (or C3iBb).

8.2 Methods

The alignments shown in Table 1 overleaf illustrate why we considered that this region was a candidate for mutagenesis. We surmised that characters of certain residues were well conserved in C3 and C4 but distinctly different in the other proteins. Residues 1427, 1431 and 1433 were selected because their charged nature might be indicative of groups involved in protein-protein interactions. The changes were made to the corresponding residues in human C5 because these displayed very different electrostatic properties, but within the context of some other conserved residues that might indicate a similar local structure.

Table 1

Alignments of sequences of C3 and related molecules for region of residues 1427-1435 of human C3

Protein	Species	RESIDUE (Human)								
		1427	1428	1429	1430	1431	1432	1433	1434	1435
C3	Human	R	Y	I	S	K	Y	E	L	D
	Mouse	R	Y	I	S	K	Y	E	M	N
	Rat	R	Y	I	S	K	Y	E	M	D
	G. pig	R	Y	I	S	K	Y	E	L	D
	Rabbit	R	Y	I	S	K	Y	E	L	N
	Cobra	R	Y	I	S	K	F	E	I	D
	Xenopus	K	Y	I	S	K	Y	E	V	N
	Trout	R	Y	I	E	K	F	E	M	D
C4	Human	R	Y	V	S	H	F	E	T	E
	Mouse	R	Y	V	S	H	F	E	T	D
Slp	Mouse	R	Y	V	S	H	F	E	T	D
C3/C4-like	Hagfish	N	Y	I	V	Q	Y	E	I	R
	Lamprey	K	Y	I	S	N	Y	E	I	T
C5	Human	Q	L	F	T	D	Y	Q	I	K
	Mouse	Q	L	L	T	D	Y	Q	I	K
A2M	Human	P	T	V	K	M	L	E	R	S
	Mouse	P	S	V	K	R	L	Q	D	Q
	Rat	P	T	V	K	M	L	E	R	S
PZP	Human	P	T	V	K	M	L	E	R	S
Murinoglobulin	Mouse	P	T	V	K	K	L	E	R	L
A1M	Rat	P	S	V	K	K	L	Q	D	Q
A1M	G. Hamster	P	T	V	K	K	L	E	R	S
A1I3	Rat	P	T	V	K	K	L	E	R	L

SUBSTITUTE SHEET (RULE 26)

The methods of mutagenesis, expression and analysis of C3 cleavage reactions were as described in the earlier examples (Examples 1-4). The mutagenic oligonucleotide was synthesised with the sequence:

5 tgggtgttgaccaatacatctccgactatcagctggacaa.

Assay for turnover of factor B.

The expressed product was purified from the COS cell medium by affinity purification on a column of Clone-3-Sepharose as described in Example 9. This method results in considerable conversion of the thiolester broken form, C3i. Wild-type C3 was isolated by the same procedure. Dilutions of the wild-type C3 (1/5, 1/25 and 1/125) were run on an SDS-PAGE gel (reducing conditions) along with the mutant C3, and silver staining indicated that the mutant was present at a concentration equivalent to slightly less than the 1/25 but much more than the 1/125 dilution of wild-type. The same dilutions were used in the assay of factor B turnover. 5 μ l of these C3's were incubated with 25 μ l of CFD-G containing 5 μ g/ml factor D and approximately 1.6 μ g/ml of ¹²⁵I-labelled factor B (approx. 1000-2000 dpm/ μ l) for 3h at 37°C. The samples were then analysed by SDS-PAGE (reducing conditions) with autoradiography of the dried gel. The results are shown in Fig. 7.

8.3 Results

As shown in Fig. 7, distinct cleavage of factor B occurs even at a 1/125 dilution of the wild-type C3 (C3i). In contrast, no significant cleavage was observed in the presence of the mutant C3, even undiluted which should be at a concentration higher than the 1/125 sample of the wild-type.

This mutant therefore appears to have an impaired ability to support the cleavage of factor B, most likely due to a reduction in its binding affinity for factor B. Therefore this is a region of C3 that can be mutated to modulate the interaction between C3i (or C3b) and factor B and perhaps also the stability of the convertase (C3iBb or C3bBb).

10 EXAMPLE 9 Purification of expressed mutant C3 molecules

9.1 Introduction

This example demonstrates how the mutant C3 molecules may be isolated from an expression medium, such as the culture medium of transfected eukaryotic cells. By simple affinity purification the C3 molecules are obtained in sufficient purity for functional tests and for conjugation to antibody by the method described in Example 10. Although elution from an antibody is accompanied by hydrolysis of a considerable proportion of the internal thiolester, the C3i product is still a suitable precursor for the generation of an active C3 convertase, as well as for the production of C3i-antibody conjugates. This approach is also likely to be useful as part of the preparation required for *in vivo* use.

9.2 Method

30 Affinity-purification on Clone-3-Sepharose.
Clone-3 is a rat monoclonal antibody that is specific for C3 and its derivatives, including C3b and C3i (Lachmann, P.J. *et al.*, 1980, *J. Immunol.* 41:503-515). Other monoclonal antibodies against C3 are available, and in

some cases have been successfully used to isolate C3 from small quantities of human plasma (Dodds, A.W., 1993, *Methods Enzymol.* 223:46-61) and are therefore also likely to be applicable for the isolation of molecules expressed *ex vivo*. The IgG fraction was coupled to Sepharose CL-4B using cyanogen bromide (methodology may be found in Harrison and Lachmann, 1986, *Handbook of Experimental Immunology*, 4th edn., Ed.s Weir, Herzenberg, Blackwell and Herzenberg; Blackwell, Oxford). Culture supernatants were either passed directly through a column of this resin (re-circulated), or first concentrated by precipitation with 25% (w/v) Na₂SO₄, and resolubilization and dialysis into PBS, 5 mM NaN₃. The column is then washed successively with (i) PBS, 5 mM NaN₃ and (ii) PBS containing 1 M NaCl. Bound C3 elutes with 50 mM Na borate buffer, pH 10.5, and is immediately neutralised by collection of 0.9 ml fractions into 0.1 ml 1 M Tris/HCl pH 7. The material is then dialysed into PBS, 5 mM NaN₃.

Preparation of C3 bearing a "His-Tag"

A "His-Tag" is a string of histidine residues that displays affinity for columns bearing Nickel ions. This method has been employed to aid the isolation of expressed proteins. We thought that this could be useful for the isolation of expressed mutant C3 molecules so we have used insertion mutagenesis to generate a plasmid encoding C3 with a tail of 6 histidine residues at the carboxy terminus (immediately carboxy-terminal to residue 1663). This location for the his tag was selected so as to minimise interference with the synthesis, folding, processing and disulphide bond formation of the nascent C3. Residue 1661 is a cysteine residue that is involved in a disulphide bond to a residue earlier in the sequence (probably Cys 1537; Dolmer, K. and Sottrup-Jensen, L.,

1993, FEBS-Lett 315: 85-90) and therefore it seemed prudent to make the insertion beyond this structural feature. The mutation was introduced using the "gapped-plasmid" technique used in Example 1, using the mutagenic oligonucleotide synthesised with the sequence:

5 tgggtgcccccaaccatcatcatcatcattgaccacaccccc.

Incorporation of the correct sequence was confirmed by DNA sequencing. This DNA sequence may now be transferred to an expression vector. After transfection of eukaryotic cells, it should be possible to isolate the expressed C3 by affinity for a column bearing Nickel ions, or by any other matrix with specific affinity for the "His-Tag".

10

15

9.3 Results

A number of mutant C3 have been purified on the Clone-3-Sepharose, including those described in Examples 1 and 2 expressed in CHO cells. The products retained the ability to support the cleavage of factor B by factor D. The same method was used to isolate the mutant described in Example B2, expressed in COS cells. Silver-staining of SDS-PAGE gels indicated that the isolated products were not 100% pure, but often appeared to be greater than or equal to 50% pure. This comes from starting materials generally containing less than 10µg/ml C3 in 10% (v/v) fetal calf serum plus other cellular proteins. In addition the C3's were not degraded during isolation, and endogenous factor H and I activity appeared to have been removed.

20

25

30

Purification by virtue of the "His-Tag" involves milder elution conditions from a column bearing Nickel ions.

For example, EDTA has been used. Application of this method to C3 should therefore allow isolation without rupture of the internal thiolester bond.

5

EXAMPLE 10 Conjugation of C3i to antibody and use to target C3 convertase activity against a particular cell

10 10.1 Introduction

One aspect of the invention is that stable C3 convertases derived from mutant C3 molecules will cause enhanced C3 conversion which, if localised at a particular target site, will promote complement-dependent attack of that target. The favoured approach for targeting the response is to couple the mutant C3 molecule, as either the C3i or C3b derivative, to an antibody specific for the desired target. In this example we demonstrate a working methodology for formation of such conjugates, which is applicable to mutant C3i or C3b molecules and can be used on material affinity-purified from an expression system, even if the thiolester of C3 has been broken in the process. By coupling C3i to an antibody that specifically binds to sheep erythrocytes, we further show that the the conjugate fixes C3i to the erythrocyte surface such that a convertase, C3iBbP, can be formed that initiates lysis of these cells when other complement components are supplied in the form of normal guinea-pig serum (in EDTA to prevent de-novo formation of C3 convertases). Hence conjugation to antibody can be used to target a C3i molecule to initiate complement-dependent attack of a particular cell type. This example uses wild-type C3i, from human plasma, that forma a C3

convertase *in vitro*. *In vivo*, wild-type C3i and C3b are broken down by factor H and I. Therefore a mutant C3, constructed according to the plans in this patent to be resistant to factors H and I and therefore forming a stable C3 convertase, would be advantageous in a physiological context.

10.2 Method

10 (i) Generation and purification of C3i-antibody conjugate

The antibody used was the IgG fraction isolated from a polyclonal rabbit anti-sheep erythrocyte antiserum. 1.1 mg was incubated with 75 nmol of SPDP in conjugation buffer, pH 7.5 (20 mM KH_2PO_4 , 60 mM Na_2HPO_4 , 0.12 M NaCl) for 2h at room temperature. The PDP-IgG was purified by gel-filtration on a Superose-6 column (Pharmacia) (in a phosphate buffer, pH 7.4, containing 0.5 M NaCl). Reduction of a sample with dithiothreitol was used to estimate 4 PDP groups coupled per molecule of IgG. C3i was prepared by treatment of purified C3 with 0.1 M methylamine, pH 7.2 (2h at 37°C). Excess methylamine was removed by gel-filtration followed by dialysis into conjugation buffer. 18 nmole of C3i was mixed with 1.7 nmoles of PDP-IgG in 1.26 ml conjugation buffer and incubated for 1 day at room temperature followed by 1.5 days at 4°C. Figure 8 shows a Coomassie Blue stained SDS-PAGE gel of the conjugation reaction mixture showing the appearance of a species of approximately 350 kDa that was not present in either PDP-IgG or C3i. This species was partially purified by gel-filtration on the Superose-6 column in a phosphate buffer, pH 7.4, containing 0.5 M NaCl and then dialysed into PBS. It eluted before the C3, in a volume from which a molecular weight of 300-400

kDa could be estimated by calibration with globular molecular weight standards. Concentrations of conjugate, free antibody and uncoupled C3 were estimated from a Coomassie-stained SDS-PAGE gel (non-reducing conditions).
5 Two-dimensional SDS-PAGE (first dimension unreduced, second dimension reduced) revealed a pattern compatible with a 1:1 conjugate between IgG and C3i.

(ii) Demonstration that the C3-antibody conjugate can be
10 used to target convertase activity against a particular cell.

20µl of dilutions of the conjugate (0 (no conjugate), 1/100, 1/50, 1/10) were incubated with 100µl of approximately 1% (v/v) sheep erythrocytes (prewashed in
15 CFD) for 1 hour at 37°C. Parallel incubations were performed with equivalent amounts of PDP-IgG (no C3) and C3 alone. The cells were then washed 4 times in CFD and resuspended to 100µl in CFD-G. 50µl of this were lysed with 150 µl H₂O, followed by addition of 800 µl of CFD
20 containing 10 mM EDTA and 2% (v/v) NGPS. The other 50µl of conjugate-coated cells were incubated for 15 min at 37°C with 50µl of CFD-G containing 190µg/ml factor B, 2µg/ml factor D, 20µg/ml properdin and 0.6 mM NiCl₂, followed by lysis with 900µl of CFD containing 10mM EDTA
25 and 2% (v/v) NGPS. After 30 min at 37°C, the cells were pelleted by centrifugation (2000 X g, about 3 min) and the optical absorbance of the supernatant was measured at 412 nm. Using the H₂O-treated samples as 100% lysis, and a buffer blank devoid of cells, the % lysis was
30 calculated, as shown in Fig. 9. The conjugate produced dose-dependent lysis, whereas neither the PDP-IgG nor the C3i alone generated any lysis significantly above that observed in the absence of any such treatment.

10.3 Summary of Results

The method used has proved successful for coupling C3i to IgG as shown by:

5

1. The formation of a band of appropriate size (about 350 kDa) for a 1:1 C3:IgG conjugate shown by SDS-PAGE in Fig. 8.

10

2. Two-dimensional SDS-PAGE (first dimension non-reduced, second dimension reduced) indicated that this species contained both IgG and C3i.

15

3. The elution characteristic of this species on gel-filtration is again consistent with a molecule of about 350 kDa.

20

4. The conjugate displays a haemolytic activity that is not displayed by either PDP-IgG or C3i (Fig. 9).

The haemolytic assay (Fig. 9) further demonstrates that:

25

1. The specific anti-sheep erythrocyte antibody has localised the C3i to the target cell (sheep erythrocyte) membrane, preventing it from being removed by washing (in contrast to free C3i).

30

2. The conjugate retains the activity of the C3i in that it is still able to form a C3 convertase by reaction with properdin and factors B and D.

3. This convertase can initiate complement-dependent attack of the target, in this case by activating the lytic pathway (C5-9) to lyse the erythrocyte.

Additional data from other laboratories show that cobra venom factor can be coupled to an antibody and that these conjugates can target complement activation against a particular cell type (Vogel, 1988, *Targeted. Diagn. Ther.*, 1:191-224; Muller, B. and Muller-Ruchholtz, W., 1987, *Leuk. Res.* 11:461-468; Parker, C.J., White, V.F. and Falk, R.J., 1986, *Complement* 3:223-235; Petrella, E.C. et al, 1987, *J. Immunol. Methods* 104:159-172). These data support the contention that C3 modified so that it is capable of forming a stable C3 convertase, like cobra venom factor, could be used to target complement-mediated responses, as outlined in this invention.

15

EXAMPLE 11 Demonstration that mutant C3 molecules induce factor B turnover in normal human serum

20 11.1 Introduction

A major purpose of the invention described herein is the consumptive depletion of complement activity from biological fluids. The invention describes methods for the manufacture of C3 molecules that are resistant to down-regulation by factors H and I. In this state they will bind factor B and generate active C3 convertases. The activity of these convertases is demonstrated by the haemolytic assay employed in Example 6. Such a convertase will therefore consume C3. If the convertase is unstable, it will dissociate without much C3 conversion. However this will allow binding of fresh factor B, and its conversion to Bb and Ba. Thus the mutant C3 will promote the consumption of factor B,

leading ultimately to the disablement of the alternative pathway, and its inability to amplify classical pathway stimulation. If a stable C3 convertase is formed, turnover of factor B will be reduced, but consumption of C3 will be increased. Both situations can therefore be desirable. In this example we demonstrate that mutant C3 molecules that are modified to make them resistant to factor I, but without any modification to modify the stability of the convertase, promote accelerated turnover of factor B in human serum. Wild-type C3, in contrast, causes no significant turn-over, presumably because wild-type C3i is rapidly degraded by factors H and I.

11.2 Method

The Mutants prepared are as follows:

Q1R2 Arg1303 changed to Gln (Example 2)

Q1Q2 Arg1303 changed to Gln, plus Arg1320 changed to Gln
(Example 1)

E1Q2 Arg1303 changed to Glu, plus Arg1320 changed to Gln
(Example 5)

These mutants were all expressed in CHO cells and then purified by precipitation with Na_2SO_4 , followed by affinity purification on Clone-3-Sepharose, as described in Example B3. Wild-type C3 (R1R2) was similarly isolated. By SDS-PAGE with silver-staining, the concentration of Q1 was between 1/5 and 1/25 of the wild-type, the concentration of Q1Q2 was about that of 1/5 wild-type, and the concentration of E1Q2 was between 1/25 and 1/125 of wild-type. All preparations probably contained a majority of thiolester-broken molecules (C3i).

10 μ l of these C3 preparations were incubated with 10 μ l of a solution of 20% (v/v) normal human serum in PBS containing 1 mM MgCl₂ and approximately 300 ng ¹²⁵I-labelled factor B (approx. 2-300,000 dpm) for 1 hour at 37°C. 5 μ l was then analysed by SDS-PAGE (reducing conditions). The dried gel was exposed to autoradiography film to indicate the positions of the bands corresponding to the intact factor B and its cleavage products. These were then excised and counted to accurately determine the degree of cleavage. The value obtained in buffer alone was subtracted as background (encompassing not only background cleavage, but also degradation products and other impurities present in the radioligand preparation).

11.3 Results

The resulting degrees of factor B cleavage are shown below:

20	1/25 Wild-type	1.49%
	1/5 Wild-type	2.74%
	Q1R2	6.19%
	Q1Q2	7.41%
25	E1Q2	6.42%

Therefore the factor I resistant mutants all produce greater levels of factor B cleavage than equivalent amounts of wild-type C3 (C3i). With larger doses or longer incubations, complete incapacitation of the alternative pathway should result.

The abbreviations used in the foregoing examples include: CFD, complement fixation diluent (defined in Harrison and

5 Lachmann, 1986, *Handbook of Experimental Immunology*, 4th edn., Ed.s Weir, Herzenberg, Blackwell and Herzenberg; Blackwell, Oxford); CFD-G, CFD containing 0.1% (w/v) gelatin; PBS, phosphate-buffered saline; NGPS, normal guinea-pig serum; SDS-PAGE, SDS-polyacrylamide gel electrophoresis; SPDP, N-Succinimidyl-3-[2-pyridyldithio]propionate.

REFERENCES:

1. Bergmann, M. & Fruton, J.S. (1941) *Adv. Enzymol.*, 1:63-98.
- 5 2. de Bruijn, M.H. & Fey, G.H. (1985), *Proc. Natl. Acad. Sci. U.S.A.* 82:708-712
3. Crawford-MH et al. (1988) *Circulation*. 78:1449-58
4. Daha, M.R. & van Es, L.A. (1982) *Immunol.* 43:33-38.
5. Farries, TC; Lachmann, PJ & Harrison, RA (1988)
- 10 *Biochem. J.* 252:47-54
6. Farries, TC; Lachmann, PJ & Harrison, RA (1988)
- Biochem. J.* 253:667-75
7. Forty, J; Hasan, R; Cary, N; White, DJ & Wallwork, J
- (1992) *Transplant. Proc.* 24:488-9
- 15 8. Fritzinger, D.C. et al. (1992) *J. Immunol.* 149:3554-3562
9. Harrison, R.A. & Lachmann, P.J. (1980) *Mol. Immunol.* 17:9-20.
10. Kalli, K.R., Hsu, P. & Fearon, D.T. (1994) *Springer*
- 20 *Semin. Immunopathol.* 15:417-431.
11. Kinoshita, T; Takata, Y; Kozono, H; Takeda, J; Hong, KS
- & Inoue, K (1988) *J. Immunol.* 141:3895-901
12. McNearney, TA; Odell, C; Holers, VM; Spear, PG;
- Atkinson, JP (1987) *J. Exp. Med.* 166:1525-35
- 25 13. Nicol, P.A.E. & Lachmann, P.J. (1973) *Immunol.* 24:259-275
14. Pangburn, MK & Muller-Eberhard, HJ (1984) *Springer*
- Semin. Immunopathol.* 7:163-92
15. Rother, K. & Till, G.O. (eds) (1988) "The complement
- 30 *System"* (Springer-Verlag Berlin Heidelberg, Germany)
16. Van den Berg, C.W., Aerts, P.C. & Van Dijk, H. (1991)
- J. Immunol. Methods* 136:287-294.
17. Vogel, CW; Smith, CA & Muller-Eberhard, HJ (1984) *J. Immunol.* 133:3235-41

18. Weisman, HF et al. (1990) *Science* **249**:146-51.
19. Wu, R. (ed.) (1993) *Methods Enzymol.* **217**: ch.s 12-14
(Academic Press, San Diego, U.S.A.)
20. Botto, M, Fong, K.Y., So, A.K., Koch, C. & Walport, M.J.
5 (1990) *J. Exp. Med.* **172**:1011-7
21. Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989)
"Molecular Cloning. A Laboratory Manual" second
edition (Cold Spring Harbor Laboratory Press)
22. Fishelson, Z. (1991) *Mol. Immunol.* **28**:545-52.
- 10 23. Taniguchi-Sidle, A & Isenman, D.E. (1993) *Mol.*
Immunol. **30**:54.
24. Lambris, J.D., Avila, D., Becherer, J.D. &
Muller, Eberhard, H.J. (1988) *J. Biol. Chem.*
263:12147-50.
- 15 25. Taniguchi-Sidle, A. and Isenman, D.E. (1992) *J.*
Biol. Chem. **267**:635-643.
26. Hofer, B. and Kuhlein, B. (1993) *Methods Enzymol.*
217:173-189. —
27. Morinaga, Y., Franceschini, T., Inouye, S. and
20 Inouye, M. (1984) *Bio-technology* **2**:636-639.
28. Harrison, R.A. and Lachmann, P.J. (1986) "Handbook
of Experimental Immunology" (eds Weir, Herzenberg,
Blackwell and Herzenberg; Blackwell, Oxford) 4th ed.,
29. Kotwal, G., J., and Moss, B., *Nature* (1988) **335**
25 (**6186**):176-8.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A modified human C3 protein which is capable of forming a stable C3 convertase wherein said modified protein is selected from the group consisting of:
 - 5 (a) a C3 protein in which either Arg-1303, Arg-1320, or both is replaced with another amino acid;
 - (b) a C3 protein which has reduced susceptibility to Factor H and/or Factor I relative to native human C3 convertase, said protein having one or more amino acid changes relative to native human C3 convertase in the
 - 10 region corresponding to amino acid residues 752-754 and/or residues 758-780 of native human C3 convertase; and
 - (c) a C3 protein having amino acid changes relative to native human C3 convertase at amino acid residues 1427, 1431 and/or 1433 of native human C3 convertase.
- 15 2. A protein according to claim 1 wherein the protein is modified by replacement of either Arg-1303, Arg-1320, or both by another amino acid.
3. A protein according to claim 2 wherein Arg-1303, Arg-1320, or both are replaced by glutamine, tyrosine, cystine, tryptophan, glutamic acid or glycine.
- 20 4. A protein according to claim 3, wherein Arg-1320 is replaced by glutamine.
5. A protein according to claim 3 or 4, wherein Arg-1303 is replaced by glutamic acid, glycine or glutamine.
6. A protein according to any preceding claim which has reduced
 - 25 susceptibility to Factor H and/or Factor I relative to native human C3 convertase, said protein having one or more amino acid changes relative to native human C3 convertase in the region corresponding to amino residues 752-754 and/or residues 758-780 of native human C3 convertase.
 7. A protein according to claim 6 wherein the one or more amino acid changes are changes from acidic amino acid residues to neutral amino acid residues.
 8. A protein according to claim 6 or 7 wherein the amino acid residue changes are changes from Asp-Glu-Asp to Gly-Ser-Gly.
 9. A protein according to any preceding claim having amino acid changes
 - 35 relative to native human C3 convertase at amino acid residues corresponding to residues 1427, 1431 and/or 1433 of native human C3 convertase.



10. An isolated DNA molecule having a DNA sequence coding for a protein according to any one of claims 1 to 9.

11. A DNA construct or vector including a DNA sequence according to claim 10.

5 12. A protein according to any one of claims 1 to 9 when used in therapy.

13. A conjugate comprising a protein according to any one of claims 1 to 9 linked to a specific binding moiety.

14. A conjugate according to claim 13 wherein the specific binding moiety is a specific binding protein.

10 15. A conjugate according to claim 14 wherein the specific binding protein is an antibody or antigen binding fragment thereof.

16. The use of a protein according to any one of claims 1 to 9 or of a conjugate according to any one of claims 13 to 15 in the manufacture of a medicament for use in depleting levels of complement pathway protein in a subject.

15 17. The use according to claim 16 wherein the medicament is for use in preventing rejection of foreign matter.

18. The use according to claim 16 wherein the medicament is for use in localising and/or amplifying endogenous complement protein conversion and deposition at a specific site.

20 19. A pharmaceutical formulation comprising one or more proteins according to any one of claims 1 to 9 or a conjugate according to any one of claims 13 to 16 together with one or more pharmaceutically acceptable carriers or excipients.

25 20. A pharmaceutical formulation according to claim 19 which is for use in depleting levels of complement pathway protein.

21. A pharmaceutical formulation according to claim 19 which is for use in preventing rejection of foreign matter.

30 22. A pharmaceutical formulation according to claim 19 which is for use in localising and/or amplifying complement protein conversion and deposition at a specific site.

23. A method of reducing complement pathway protein in a mammal, the method comprising administering to the mammal an effective amount of a protein according to any one of claims 1 to 9.



24. A method according to claim 23 wherein the protein is administered in the form of a pharmaceutical formulation according to claim 19.

5

Dated this thirtieth day of April 1999

IMUTRAN LIMITED

Patent Attorneys for the Applicant:

F B RICE & CO

RECEIVED
APR 30 1999
F B RICE & CO



1/13

FIGURE 1

10	20	30	40	50	60
MGPTSGPSLL	LLLLTHLPLA	LGSPMYSIIT	PNILRLESEE	TMVLEAHDAQ	GDVPVTVTVH
70	80	90	100	110	120
DFPGKKLVLS	SEKTVLTPAT	NHMGNVFTFI	PANREFKSEK	GRNKFVTVQA	TFGTQVVEKV
130	140	150	160	170	180
VLVSLQSGYL	FIQTDKTIYT	PGSTVLYRIF	TVNHKLLPVG	RTVMVNIENP	EGIPVKQDSL
190	200	210	220	230	240
SSQNQLGVLP	LSWDIPELVN	MGQWKIRAYY	ENSPQQVFST	EFEVKEYVLP	SFEVIVEPTE
250	260	270	280	290	300
KFYIYNEKG	LEVITITARFL	YGKKVEGTAF	VIFGIQDGEQ	RISLPESLKR	IPIEDGSGEV
310	320	330	340	350	360
VLSRKVLLDG	VQNPRAEDLV	GKSLYVSATV	ILHSGSDMVQ	AERSGPIIVT	SPYQIHFTKT
370	380	390	400	410	420
PKYFKPGMPF	DLMVFVTNPD	GSPAYRVFVA	VQGEDTVQSL	TQGDGVAKLS	INTHPSQKPL
430	440	450	460	470	480
SITVRTKKQE	LSEAEQATRT	MQALPYSTVG	NSNNYLHLSV	LRTELRPGET	LVNFNLLRMD
490	500	510	520	530	540
RAHEAKIRYY	TYLIMNKGRL	LKAGRQVREP	GQDLVVLPLS	ITTDIFIPSR	LVAYYTLIGA
550	560	570	580	590	600
SGQREVADS	VWVDVKDSCV	GSLVVKSGQS	EDRQPVPGQQ	MTLKIEGDHG	ARVVLVAVDK
610	620	630	640	650	660
GVFVLNKKNK	LTQSKIWDVV	EKADIGCTPG	SGKDYAGVFS	DAGLTFTSSS	GQQTAAQRAEL
670	680	690	700	710	720
QCPQPAARRR	RSVQLTEKRM	DKVGKYPKEL	RKCCEDGMRE	NPMRFSCQRR	TRFISLGEAC
730	740	750	760	770	780
KKVFLDCCNY	ITELRRQHAR	ASHLGLARSN	LDEDIIAEEN	IVSRSEFPES	WLWNVEDLKE
790	800	810	820	830	840
PPKNGISTKL	MNIFLKDSIT	TWEILAVSMS	DKKGICVADP	FEVTVMQDFF	IDLRLPYSVV
850	860	870	880	890	900
RNEQVEIRAV	LYNYRQNQEL	KVRVELLHNP	AFCSLATTKR	RHQQTITIPP	KSSLSVPYVI
910	920	930	940	950	960
VPLKTGLQEV	EVKAAVYHHF	ISDGVKSLK	VVPEGIRMNK	TVAVRTLDP	RLGREGVQKE
970	980	990	1000	1010	1020
DIPPADLSQ	VPDTESETRI	LLQGTPVAQM	TEDAVIDAERL	KHLIVTPSGC	GEQNMIGMTP
1030	1040	1050	1060	1070	1080
TVIAVHYLDE	TEQWEKFGLE	KRQGALELIK	KGYTQQLAFR	QPSSAFAAFV	KRAPSTWLTA
1090	1100	1110	1120	1130	1140
YVVKVFS LAV	NLIAIDSQVL	CGAVKWLILE	KQKPDGVFQE	DAPVIHQEMI	GGLRNNNEKD
1150	1160	1170	1180	1190	1200
MALTAFLVIS	LQEAKDICEE	QVNSLPGSIT	KAGDFLEANY	MNLQRSYTVA	IAGYALAQMG

SUBSTITUTE SHEET (RULE 26)

2/13

1-2

1210	1220	1230	1240	1250	1260
RLKGPLLKNF	LTTAKDKNRW	EDPGKQLYNV	EATSYALLAL	LQLKDFDFVP	PVVRWLNEQR
1270	1280	1290	1300	1310	1320
YYGGGYGSTQ	ATFMVFAQAL	QYQKDAPDHQ	ELNLDVSLQL	PSRSSKITHR	IHWESASLLR
1330	1340	1350	1360	1370	1380
SEETKENEGF	TVTAEGKGQG	TLSVVTMYHA	KAKDQLTCNK	FDLKVTIKPA	PETEKRPQDA
1390	1400	1410	1420	1430	1440
KNTMILEICT	RYRGDQDATM	SILDISMMTG	FAPDTDDLKQ	LANGVDRIYS	KYELDKAFSD
1450	1460	1470	1480	1490	1500
RNTLIIYLDK	VSHSEDDCLA	FKVHQYFNVE	LIQPGAVKVY	AYYNLEESCT	RFYHPEKEDG
1510	1520	1530	1540	1550	1560
KLNLKCRDEL	CRCAEENCFI	QKSDDKVTLE	ERLDKACEPG	VDYVYKTRLV	KVQLSNDFDE
1570	1580	1590	1600	1610	1620
YIMAIEQTIK	SGSDEVQVGQ	QRTFISPIKC	REALKLEKK	HYLMWGLSSD	FWGEKPNLSY
1630	1640	1650	1660		
IIGKDTWVEH	WPEEDECODE	ENQKQCQDLG	AFTESMVVFG	CPN	

3/13

FIGURE 2

	cctctccct	ctgtccctct	gtccctctga	ccctgcactg	tcccagcacc	
	12	20	30	40	50	60
atgggaccca	cctcaggtcc	cagcctgctg	ctcctgctac	taaccacact	ccccctggct	
70	80	90	100	110	120	
ctggggagtc	ccatgtactc	tatcatcacc	cccaacatct	tgcggtctgga	gagcgaggag	
130	140	150	160	170	180	
accatgggtgc	tggaggccca	cgacgcgcaa	ggggatgttc	cagtcactgt	tactgtccac	
190	200	210	220	230	240	
gacttcccag	gcaaaaaact	agtgtgtgtcc	agtgagaaga	ctgtgctgac	ccctgccacc	
250	260	270	280	290	300	
aaccacatgg	gcaacgtcac	cttcacgatc	ccagccaaca	gggagttcaa	gtcagaaaag	
310	320	330	340	350	360	
gggcgcaaca	agttcgtgac	cgtgcaggcc	accttcggga	cccaagtggg	ggagaagggtg	
370	380	390	400	410	420	
gtgctgggtca	gcctgcagag	cgggtacctc	ttcatccaga	cagacaagac	catctacacc	
430	440	450	460	470	480	
cctgggtcca	cagttctcta	tcggatcttc	accgtcaacc	acaagctgct	acccgtgggc	
490	500	510	520	530	540	
cggacgggtca	tgggtcaacat	tgagaaccgc	gaaggcatcc	cgggtcaagca	ggactccttg	
550	560	570	580	590	600	
tcttctcaga	accagcttgg	cgtcttgccc	ttgtcttggg	acattccgga	actcgtcaac	
610	620	630	640	650	660	
atggggcagt	ggaagatccg	agcctactat	gaaaactcac	cacagcaggt	cttctccact	
670	680	690	700	710	720	
gagtttgagg	tgaaggagta	cgtgctgccc	agtttcgagg	tcatagtgga	gcctacagag	
730	740	750	760	770	780	
aaattctact	acatctataa	cgagaagggc	ctggagggtca	ccatcacgcg	caggttcctc	
790	800	810	820	830	840	
tacgggaaga	aagtggaggg	aactgccttt	gtcatcttcg	ggatccagga	tggcgaacag	
850	860	870	880	890	900	
aggatttccc	tgctgaatc	cctcaagcgc	attccgattg	aggatggctc	gggggaggtt	
910	920	930	940	950	960	
gtgctgagcc	ggaaggtact	gctggacggg	gtgcagaacc	cccagacaga	agacctgggtg	
970	980	990	1000	1010	1020	
gggaagtctt	tgtacgtgtc	tgccaccgtc	atcttgact	caggcagtga	catgggtgcag	
1030	1040	1050	1060	1070	1080	
gcagagcgca	gcgggatccc	catcgtgacc	tctccctacc	agatccactt	caccaagaca	
1090	1100	1110	1120	1130	1140	
cccaagtact	tcaaaccagg	aatgcccttt	gacctcatgg	tgttcgtgac	gaaccctgat	
1150	1160	1170	1180	1190	1200	
ggctctccag	cctaccgagt	ccccgtggca	gtccagggcg	aggacactgt	gcagtctcta	
1210	1220	1230	1240	1250	1260	

acccagggag 1270	atggcggtggc 1280	caaactcagc 1290	atcaacacac 1300	accccagcca 1310	gaagcccttg 1320
agcatcacgg 1330	tgcgcacgaa 1340	gaagcaggag 1350	ctctcggagg 1360	cagagcaggc 1370	taccaggacc 1380
atgcaggctc 1390	tgccctacag 1400	caccgtgggc 1410	aactccaaca 1420	attacctgca 1430	tctctcagtg 1440
ctacgtacag 1450	agctcagacc 1460	cggggagacc 1470	ctcaacgtca 1480	acttcctcct 1490	gcgaatggac 1500
cgcgcccacg 1510	aggccaagat 1520	ccgctactac 1530	acctacctga 1540	tcatgaacaa 1550	gggcaggctg 1560
ttgaaggcgg 1570	gacgccagggt 1580	gcgagagccc 1590	ggccaggacc 1600	tgggtgggtgct 1610	gccccgtgcc 1620
atcaccaccg 1630	acttcatccc 1640	ttccttccgc 1650	ctgggtggcgt 1660	actacacgct 1670	gacgggtgcc 1680
agcggccaga 1690	gggaggtggt 1700	ggccgactcc 1710	gtgtgggtgg 1720	acgtcaagga 1730	ctcctgcgtg 1740
ggctcgctgg 1750	tggtaaaaag 1760	cggccagtca 1770	gaagaccggc 1780	agcctgtacc 1790	tgggcagcag 1800
atgaccctga 1810	agatagaggg 1820	tgaccacggg 1830	gcccgggtgg 1840	tactgggtggc 1850	cgtggacaag 1860
ggcgtgttcg 1870	tgctgaataa 1880	gaagaacaaa 1890	ctgacgcaga 1900	gtaagatctg 1910	ggacgtgggtg 1920
gagaaggcag 1930	acatcggtcg 1940	caccccgggc 1950	agtgggaagg 1960	attacgccgg 1970	tgtcttctcc 1980
gacgcagggc 1990	tgaccttcac 2000	gagcagcagt 2010	ggccagcaga 2020	ccgcccagag 2030	ggcagaactt 2040
cagtgccgcg 2050	agccagccgc 2060	ccgccgacgc 2070	cgttccgtgc 2080	agctcacgga 2090	gaagcgaatg 2100
gacaaagtctg 2110	gcaagtaccc 2120	caaggagctg 2130	cgcaagtgtg 2140	gcgaggacgg 2150	catgcgggag 2160
aaccccatga 2170	ggttctcgtg 2180	ccagcgccgg 2190	acccgtttca 2200	tctccctggg 2210	cgaggcgtgc 2220
aagaaggtct 2230	tcctggactg 2240	ctgcaactac 2250	atcacagagc 2260	tgcgggcgga 2270	gcacgcgcgg 2280
gccagccacc 2290	tgggcctggc 2300	caggagtaac 2310	ctggatgagg 2320	acatcattgc 2330	agaagagaac 2340
atcgtttccc 2350	gaagtgagtt 2360	cccagagagc 2370	tggctgtgga 2380	acgttgagga 2390	cttgaaagag 2400
ccaccgaaaa 2410	atggaatctc 2420	tacgaagctc 2430	atgaatatat 2440	ttttgaaaga 2450	ctccatcacc 2460
acgtgggaga 2470	ttctggctgt 2480	gagcatgtcg 2490	gacaagaaag 2500	ggatctgtgt 2510	ggcagacccc 2520
ttcgaggtca 2530	cagtaatgca 2540	ggacttcttc 2550	atcgacctgc 2560	ggctacccta 2570	ctctgttgtt 2580

2-3

cgaaacgagc 2590	aggtggaaat 2600	ccgagccgtt 2610	ctctacaatt 2620	accggcagaa 2630	ccaagagctc 2640
aaggtgaggg 2650	tggaactact 2660	ccacaatcca 2670	gccttctgca 2680	gcctggccac 2690	caccaagagg 2700
cgtcaccagc 2710	agaccataac 2720	catccccccc 2730	aagtcctcgt 2740	tgtccgttcc 2750	atatgtcatc 2760
gtgccgctaa 2770	agaccggcct 2780	gcaggaagtg 2790	gaagtcaagg 2800	ctgctgtcta 2810	ccatcatttc 2820
atcagtgaag 2830	gtgtcaggaa 2840	gtccctgaag 2850	gtcgtgccgg 2860	aaggaatcag 2870	aatgaacaaa 2880
actgtggctg 2890	ttcgcaccct 2900	ggatccagaa 2910	cgctggggcc 2920	gtgaaggagt 2930	gcagaaagag 2940
gacatcccac 2950	ctgcagacct 2960	cagtgaacaa 2970	gtcccggaac 2980	ccgagtctga 2990	gaccagaatt 3000
ctcctgcaag 3010	ggaccccagt 3020	ggcccagatg 3030	acagaggatg 3040	ccgtcgacgc 3050	ggaacggctg 3060
aagcacctca 3070	ttgtgacccc 3080	ctcgggctgc 3090	ggggaacaga 3100	acatgatcgg 3110	catgacgccc 3120
acggtcacgc 3130	ctgtgcatta 3140	cctggatgaa 3150	acggagcagt 3160	gggagaagtt 3170	cggcctagag 3180
aagcggcagg 3190	gggccttgga 3200	gctcatcaag 3210	aaggggtaca 3220	cccagcagct 3230	ggccttcaga 3240
caaccacagc 3250	ctgcctttgc 3260	ggccttcgtg 3270	aaacgggcac 3280	ccagcacctg 3290	gctgaccgcc 3300
tacgtggctc 3310	aggtcttctc 3320	tctggctgtc 3330	aacctcatcg 3340	ccatcgactc 3350	ccaagtcttc 3360
tgcggggctg 3370	ttaaatggct 3380	gatcctggag 3390	aagcagaagc 3400	ccgacggggg 3410	cttccaggag 3420
gatgcgcccc 3430	tgatacacca 3440	agaaatgatt 3450	ggtggattac 3460	ggaacaacaa 3470	cgagaaagae 3480
atggccctca 3490	cggcctttgt 3500	tctcatctcg 3510	ctgcaggagg 3520	ctaaagatat 3530	ttgcgaggag 3540
caggtcaaca 3550	gcctgccagg 3560	cagcatcact 3570	aaagcaggag 3580	acttccttga 3590	agccaactac 3600
atgaacctac 3610	agagatccta 3620	cactgtggcc 3630	attgctggct 3640	atgctctggc 3650	ccagatgggc 3660
aggctgaagg 3670	ggcctcttct 3680	taacaaattt 3690	ctgaccacag 3700	ccaaagataa 3710	gaaccgctgg 3720
gaggaccctg 3730	gtaagcagct 3740	ctacaacgtg 3750	gaggccacat 3760	cctatgccct 3770	cttggcccta 3780
ctgcagctaa 3790	aagactttga 3800	ctttgtgcct 3810	cccgtcgtgc 3820	gttggctcaa 3830	tgaacagaga 3840
tactacgggtg 3850	gtggctatgg 3860	ctctaccag 3870	gccaccttca 3880	tgggtgtcca 3890	agccttggct 3900


6/13

2-4

caataccaaa 3910	aggacgcccc 3920	tgaccaccag 3930	gaactgaacc 3940	ttgatgtgtc 3950	cctccaactg 3960
cccagccgca 3970	gctccaagat 3980	caccacccgt 3990	atccactggg 4000	aatctgccag 4010	cctcctgcga 4020
tcagaagaga 4030	ccaaggaaaa 4040	tgagggtttc 4050	acagtcacag 4060	ctgaaggaaa 4070	aggccaaggc 4080
accttgctcg 4090	tggtgacaat 4100	gtaccatgct 4110	aaggccaaag 4120	atcaactcac 4130	ctgtaataaa 4140
ttcgacctca 4150	aggtcaccat 4160	aaaaccagca 4170	ccggaaacag 4180	aaaagaggcc 4190	tcaggatgcc 4200
aagaacacta 4210	tgatccttga 4220	gatctgtacc 4230	aggtaccggg 4240	gagaccagga 4250	tgccactatg 4260
tctatattgg 4270	acatatccat 4280	gatgactggc 4290	tttgctccag 4300	acacagatga 4310	cctgaagcag 4320
ctggccaatg 4330	gtgttgacag 4340	atacatctcc 4350	aagtatgagc 4360	tggacaaagc 4370	cttctccgat 4380
aggaacaccc 4390	tcatcatcta 4400	cctggacaag 4410	gtctcacact 4420	ctgaggatga 4430	ctgtctagct 4440
ttcaaagttc 4450	accaataactt 4460	taatgtagag 4470	cttatccagc 4480	ctggagcagt 4490	caaggtctac 4500
gcctattaca 4510	acctggagga 4520	aagctgtacc 4530	cggttctacc 4540	atccggaaaa 4550	ggaggatgga 4560
aagctgaaca 4570	agctctgccg 4580	tgatgaactg 4590	tgccgctgtg 4600	ctgaggagaa 4610	ttgcttcata 4620
caaaagtcgg 4630	atgacaaggt 4640	caccctggaa 4650	gaacggctgg 4660	acaaggcctg 4670	tgagccagga 4680
gtggactatg 4690	tgtacaagac 4700	ccgactggtc 4710	aagggttcagc 4720	tgtccaatga 4730	ctttgacgag 4740
tacatcatgg 4750	ccattgagca 4760	gaccatcaag 4770	tcaggctcgg 4780	atgagggtgca 4790	ggttggacag 4800
cagcgcacgt 4810	tcatcagccc 4820	catcaagtgc 4830	agagaagccc 4840	tgaagctgga 4850	ggagaagaaa 4860
cactacctca 4870	tgtggggtct 4880	ctcctccgat 4890	ttctggggag 4900	agaagcccaa 4910	cctcagctac 4920
atcatcgagg 4930	aggacacttg 4940	ggtggagcac 4950	tgccctgagg 4960	aggacgaatg 4970	ccaagacgaa 4980
gagaaccaga 4990	aacaatgcca 5000	ggacctcggc 5010	gccttcaccg 5020	agagcatggg 5030	tgtctttggg 5040
tgccccaact 5050	gaccacaccc 5060	ccattcc			


7/13

FIGURE 3




8/13

FIGURE 4




9/13

FIGURE 5




10/13

FIGURE 6



11/13


FIGURE 7



12/13

FIGURE 8

1 - 2 - 3



13/13

FIGURE 9

