

(19) **DANMARK**

(10) **DK/EP 3442602 T3**



(12)

## Oversættelse af europæisk patentskrift

Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **A 61 K 48/00 (2006.01)** **C 12 N 15/11 (2006.01)** **C 12 N 15/113 (2010.01)**
- (45) Oversættelsen bekendtgjort den: **2025-05-19**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2025-02-19**
- (86) Europæisk ansøgning nr.: **17783236.7**
- (86) Europæisk indleveringsdag: **2017-04-14**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-02-20**
- (86) International ansøgning nr.: **US2017027636**
- (87) Internationalt publikationsnr.: **WO2017181015**
- (30) Prioritet: **2016-04-15 US 201662323163 P** **2017-03-17 US 201762473253 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Research Institute at Nationwide Children's Hospital, 700 Children's Drive, Room W172, Columbus, Ohio 43205, USA**
- (72) Opfinder: **RODINO-KLAPAC, Louise, 4912 Bixby Ridge Drive E, Groveport, OH 43125, USA**  
**MENDELL, Jerry, R., 8176 Copperfield Drive, Columbus, OH 43235, USA**  
**HELLER, Kristin, N., The Research Institute At Nationwide, Children's Hospital, 700 Children's Drive, Columbus, OH 43205, USA**
- (74) Fuldmægtig i Danmark: **RWS Group, RWS Compass House, Vanwall Business Park, Vanwall Road, Maidenhead, Berkshire, SL6 4UB, Storbritannien**
- (54) Benævnelse: **ADENO-ASSOCIERET VIRUSVEKTORS FRIGØRELSE AF MIKRODYSTROFIN TIL BEHANDLING AF MUSKELDYSTROFI**
- (56) Fremdragne publikationer:  
**WO-A1-2015/110449**  
**WO-A1-2015/197232**  
**WO-A1-2017/165859**  
**US-A1- 2008 249 052**  
**US-A1- 2010 003 218**  
**US-A1- 2014 234 255**  
**US-A1- 2014 256 802**  
**TAEYOUNG KOO ET AL: "Long-term functional adeno-associated virus-microdystrophin expression in the dystrophic CXMDj dog", THE JOURNAL OF GENE MEDICINE, vol. 13, no. 9, 1 September 2011 (2011-09-01), pages 497 - 506, XP055152320, ISSN: 1099-498X, DOI: 10.1002/jgm.1602**  
**LOUISE R RODINO-KLAPAC ET AL: "Persistent Expression of FLAG-tagged Micro dystrophin in Nonhuman Primates Following Intramuscular and Vascular Delivery", MOLECULAR THERAPY, vol. 18, no. 1, 1 January 2010 (2010-01-01), pages 109 - 117, XP055631567, ISSN: 1525-0016, DOI: 10.1038/mt.2009.254**

Fortsættes ...

**LG CHICOINE ET AL: "Plasmapheresis Eliminates the Negative Impact of AAV Antibodies on Microdystrophin Gene Expression Following Vascular Delivery", MOLECULAR THERAPY, vol. 22, no. 2, 1 February 2014 (2014-02-01), pages 338 - 347, XP055367763, ISSN: 1525-0016, DOI: 10.1038/mt.2013.244**

**HELLER ET AL: "379. MicroRNA-29 and Micro-Dystrophin Combinatorial Therapy Suppresses Fibrosis and Restores Function to mdx/utrn+/- Mice", MOLECULAR THERAPY, vol. 24, no. 1, 14 December 2016 (2016-12-14), pages S151, XP055433385, DOI: 10.1016/S1525-0016(16)33188-4**

**KRISTIN N. HELLER ET AL: "MicroRNA-29 overexpression by adeno-associated virus suppresses fibrosis and restores muscle function in combination with micro-dystrophin", JCI INSIGHT, vol. 2, no. 9, 4 May 2017 (2017-05-04), pages 1 - 13, XP055597289, DOI: 10.1172/jci.insight.93309**

# DESCRIPTION

Description

## FIELD OF INVENTION

**[0001]** The invention provides gene therapy vectors, such as adeno-associated virus (AAV) vectors, expressing a miniaturized human micro-dystrophin gene, such as for use in methods to reduce and prevent fibrosis in subjects suffering from muscular dystrophy.

## BACKGROUND

**[0002]** The importance of muscle mass and strength for daily activities, such as locomotion and breathing, and for whole body metabolism is unequivocal. Deficits in muscle function produce muscular dystrophies (MDs) that are characterized by muscle weakness and wasting and have serious impacts on quality of life. The most well-characterized MDs result from mutations in genes encoding members of the dystrophin-associated protein complex (DAPC). These MDs result from membrane fragility associated with the loss of sarcolemmal-cytoskeleton tethering by the DAPC. Duchenne Muscular Dystrophy (DMD) is one of the most devastating muscle diseases affecting 1 in 5000 newborn males.

**[0003]** This application includes two translational approaches to develop treatment for DMD. Fibrotic infiltration is profound in DMD and is a significant impediment to any potential therapy. It is also important to consider that gene replacement alone is hampered by the severity of fibrosis, already present in very young children with DMD. In fact, muscle biopsies at the usual age of diagnosis, between 4-5 years old, show very significant levels of fibrosis.

**[0004]** DMD is caused by mutations in the DMD gene leading to reductions in mRNA and the absence of dystrophin, a 427 kD sarcolemmal protein associated with the dystrophin-associated protein complex (DAPC) (Hoffman et al., Cell 51(6):919-28, 1987). The DAPC is composed of multiple proteins at the muscle sarcolemma that form a structural link between the extra-cellular matrix (ECM) and the cytoskeleton via dystrophin, an actin binding protein, and alpha-dystroglycan, a laminin-binding protein. These structural links act to stabilize the muscle cell membrane during contraction and protect against contraction-induced damage. With dystrophin loss, membrane fragility results in sarcolemmal tears and an influx of calcium, triggering calcium-activated proteases and segmental fiber necrosis (Straub et al., Curr Opin. Neurol. 10(2): 168-75, 1997). This uncontrolled cycle of muscle degeneration and regeneration ultimately exhausts the muscle stem cell population (Sacco et al., Cell, 2010. 143(7): p. 1059-

71; Wallace et al., *Annu Rev Physiol*, 2009. 71: p. 37-57), resulting in progressive muscle weakness, endomysial inflammation, and fibrotic scarring.

**[0005]** Without membrane stabilization from dystrophin or a micro-dystrophin, DMD will manifest uncontrolled cycles of tissue injury and repair and ultimately replace lost muscle fibers with fibrotic scar tissue through connective tissue proliferation. Fibrosis is characterized by the excessive deposits of ECM matrix proteins, including collagen and elastin. ECM proteins are primarily produced from cytokines such as TGF $\beta$  that is released by activated fibroblasts responding to stress and inflammation. Although the primary pathological feature of DMD is myofiber degeneration and necrosis, fibrosis as a pathological consequence has equal repercussions. The over-production of fibrotic tissue restricts muscle regeneration and contributes to progressive muscle weakness in the DMD patient. In one study, the presence of fibrosis on initial DMD muscle biopsies was highly correlated with poor motor outcome at a 10-year follow-up (Desguerre et al., *J Neuropathol Exp Neurol*, 2009. 68(7): p. 762-7). These results point to fibrosis as a major contributor to DMD muscle dysfunction and highlight the need to develop therapies that reduce fibrotic tissue. Most anti-fibrotic therapies that have been tested in mdx mice act to block fibrotic cytokine signaling through inhibition of the TGF $\beta$  pathway. MicroRNAs (miRNAs) are single-stranded RNAs of ~22 nucleotides that mediate gene silencing at the post-transcriptional level by pairing with bases within the 3' UTR of mRNA, inhibiting translation or promoting mRNA degradation. A seed sequence of 7 bp at the 5' end of the miRNA targets the mRNA; additional recognition is provided by the remainder of the targeted sequence, as well as its secondary structure. MiRNAs play an important role in muscle disease pathology and exhibit expression profiles that are uniquely dependent on the type of muscular dystrophy in question (Eisenberg et al. *Proc Natl Acad Sci U S A*, 2007. 104(43): p. 17016-21). A growing body of evidence suggests that miRNAs are involved in the fibrotic process in many organs including heart, liver, kidney, and lung (Jiang et al., *Proc Natl Acad Sci U S A*, 2007. 104(43): p. 17016-21). Recently, the down-regulation of miR-29 was shown to contribute to cardiac fibrosis (Cacchiarelli et al., *Cell Metab*, 2010. 12(4): p. 341-51) and reduced expression of miR-29 was genetically linked with human DMD patient muscles (Eisenberg et al. *Proc Natl Acad Sci U S A*, 2007. 104(43): p. 17016-2). The miR-29 family consists of three family members expressed from two bicistronic miRNA clusters. MiR-29a is coexpressed with miR-29b (miR-29b-1); miR-29c is coexpressed with a second copy of miR-29b (miR-29b-2). The miR-29 family shares a conserved seed sequence and miR-29a and miR-29b each differ by only one base from miR-29c. Furthermore, electroporation of miR-29 plasmid (a cluster of miR-29a and miR-29b-1) into mdx mouse muscle reduced the expression levels of ECM components, collagen and elastin, and strongly decreased collagen deposition in muscle sections within 25 days post-treatment (Cacchiarelli et al., *Cell Metab*, 2010. 12(4): p. 341-51).

**[0006]** Adeno-associated virus (AAV) is a replication-deficient parvovirus, the single-stranded DNA genome of which is about 4.7 kb in length including 145 nucleotide inverted terminal repeat (ITRs). There are multiple serotypes of AAV. The nucleotide sequences of the genomes of the AAV serotypes are known. For example, the nucleotide sequence of the AAV serotype 2 (AAV2) genome is presented in Srivastava et al., *J Virol*, 45: 555-564 (1983) as corrected by

Ruffing et al., J Gen Virol, 75: 3385-3392 (1994). As other examples, the complete genome of AAV-1 is provided in GenBank Accession No. NC\_002077; the complete genome of AAV-3 is provided in GenBank Accession No. NC\_1829; the complete genome of AAV-4 is provided in GenBank Accession No. NC\_001829; the AAV-5 genome is provided in GenBank Accession No. AF085716; the complete genome of AAV-6 is provided in GenBank Accession No. NC\_001862; at least portions of AAV-7 and AAV-8 genomes are provided in GenBank Accession Nos. AX753246 and AX753249, respectively (see also U.S. Patent Nos. 7,282,199 and 7,790,449 relating to AAV-8); the AAV-9 genome is provided in Gao et al., J. Virol., 78: 6381-6388 (2004); the AAV-10 genome is provided in Mol. Ther., 13(1): 67-76 (2006); and the AAV-11 genome is provided in Virology, 330(2): 375-383 (2004). The AAVrh74 serotype is described in Rodino-Klapac et al. J. Trans. Med. 5: 45 (2007). *Cis*-acting sequences directing viral DNA replication (rep), encapsidation/packaging and host cell chromosome integration are contained within the ITRs. Three AAV promoters (named p5, p19, and p40 for their relative map locations) drive the expression of the two AAV internal open reading frames encoding rep and cap genes. The two rep promoters (p5 and p19), coupled with the differential splicing of the single AAV intron (e.g., at AAV2 nucleotides 2107 and 2227), result in the production of four rep proteins (rep 78, rep 68, rep 52, and rep 40) from the rep gene. Rep proteins possess multiple enzymatic properties that are ultimately responsible for replicating the viral genome. The cap gene is expressed from the p40 promoter and it encodes the three capsid proteins VP1, VP2, and VP3. Alternative splicing and non-consensus translational start sites are responsible for the production of the three related capsid proteins. A single consensus polyadenylation site is located at map position 95 of the AAV genome. The life cycle and genetics of AAV are reviewed in Muzyczka, Current Topics in Microbiology and Immunology, 158: 97-129 (1992).

**[0007]** AAV possesses unique features that make it attractive as a vector for delivering foreign DNA to cells, for example, in gene therapy. AAV infection of cells in culture is noncytopathic, and natural infection of humans and other animals is silent and asymptomatic. Moreover, AAV infects many mammalian cells allowing the possibility of targeting many different tissues *in vivo*. Moreover, AAV transduces slowly dividing and non-dividing cells, and can persist essentially for the lifetime of those cells as a transcriptionally active nuclear episome (extrachromosomal element). The AAV proviral genome is infectious as cloned DNA in plasmids which makes construction of recombinant genomes feasible. Furthermore, because the signals directing AAV replication, genome encapsidation and integration are contained within the ITRs of the AAV genome, some or all of the internal approximately 4.3 kb of the genome (encoding replication and structural capsid proteins, rep-cap) may be replaced with foreign DNA such as a gene cassette containing a promoter, a DNA of interest and a polyadenylation signal. The rep and cap proteins may be provided *in trans*. Another significant feature of AAV is that it is an extremely stable and hearty virus. It easily withstands the conditions used to inactivate adenovirus (56° to 65°C for several hours), making cold preservation of AAV less critical. AAV may even be lyophilized. Finally, AAV-infected cells are not resistant to superinfection.

**[0008]** Multiple studies have demonstrated long-term (> 1.5 years) recombinant AAV-mediated protein expression in muscle. See, Clark et al., Hum Gene Ther, 8: 659-669 (1997); Kessler et al., Proc Nat. Acad Sc. USA, 93: 14082-14087 (1996); and Xiao et al., J Virol, 70: 8098-8108

(1996). See also, Chao et al., Mol Ther, 2:619-623 (2000) and Chao et al., Mol Ther, 4:217-222 (2001). Moreover, because muscle is highly vascularized, recombinant AAV transduction has resulted in the appearance of transgene products in the systemic circulation following intramuscular injection as described in Herzog et al., Proc Natl Acad Sci USA, 94: 5804-5809 (1997) and Murphy et al., Proc Natl Acad Sci USA, 94: 13921-13926 (1997). Moreover, Lewis et al., J Virol, 76: 8769-8775 (2002) demonstrated that skeletal myofibers possess the necessary cellular factors for correct antibody glycosylation, folding, and secretion, indicating that muscle is capable of stable expression of secreted protein therapeutics.

**[0009]** Functional improvement in patients suffering from DMD and other muscular dystrophies require both gene restoration and reduction of fibrosis. There is a need for methods of reducing fibrosis that may be paired with gene restoration methods for more effective treatments of DMD and other muscular dystrophies. miR29 is a potential gene regulator and an ideal candidate for reducing muscle fibrosis.

## SUMMARY OF INVENTION

**[0010]** In one aspect, the invention provides a recombinant AAVrh74 vector comprising a muscle-specific control element operably linked to the nucleotide sequence of SEQ ID NO: 7.

**[0011]** In one aspect, the invention provides a composition comprising the recombinant AAVrh74 vector of the invention, and a pharmaceutically acceptable carrier.

**[0012]** In one aspect, the invention provides the recombinant AAVrh74 vector of the invention or the composition of the invention for use in the treatment of muscular dystrophy.

**[0013]** In one aspect, the invention provides the recombinant AAVrh74 vector of the invention or the composition of the invention for use in reducing or preventing fibrosis in a subject suffering from muscular dystrophy.

**[0014]** In one aspect, the invention provides the recombinant AAVrh74 vector of the invention or the composition of the invention for use in increasing muscular force or muscle mass in a subject suffering from muscular dystrophy.

**[0015]** In some embodiments, the subject is suffering from Duchenne muscular dystrophy.

**[0016]** In one aspect, the invention provides an in vitro method of producing a functional micro-dystrophin protein comprising infecting a host cell with a recombinant AAVrh74 vector of the invention and expressing a functional micro-dystrophin protein in the host cell.

**[0017]** Combination therapies and approaches for reducing and preventing fibrosis using gene therapy vectors may deliver miR-29 to suppress fibrosis along with micro-dystrophin to address the gene defect observed in DMD. As shown in Examples 5-7, the combination treatment

resulted in a greater reduction in fibrosis, increased muscle size and increased muscle force.

**[0018]** For example the muscle-specific control element is human skeletal actin gene element, cardiac actin gene element, myocyte-specific enhancer binding factor MEF, muscle creatine kinase (MCK), tMCK (truncated MCK), myosin heavy chain (MHC), C5-12 (synthetic promoter), murine creatine kinase enhancer element, skeletal fast-twitch troponin C gene element, slow-twitch cardiac troponin C gene element, the slow-twitch troponin I gene element, hypoxia-inducible nuclear factors, steroid-inducible element or glucocorticoid response element (GRE).

**[0019]** For example, any of the rAAV vectors of the invention are operably linked to the muscle-specific control element comprising the MCK enhancer nucleotide sequence of SEQ ID NO: 10 and/or the MCK promoter sequence of SEQ ID NO: 11.

**[0020]** The invention also provides for pharmaceutical compositions (or sometimes referred to herein as simply "compositions") comprising any of the rAAV vectors of the invention.

**[0021]** The invention also provides for viral particles comprising any of the recombinant AAV vectors of the invention.

**[0022]** For example, any of the rAAV of the invention may be administered to subjects suffering from muscular dystrophy to reduce fibrosis, and in particular reduces fibrosis in skeletal muscle or in cardiac muscle of the subject. These methods may further comprise the step of administering a rAAV expressing micro-dystrophin.

**[0023]** "Fibrosis" refers to the excessive or unregulated deposition of extracellular matrix (ECM) components and abnormal repair processes in tissues upon injury including skeletal muscle, cardiac muscle, liver, lung, kidney, and pancreas. The ECM components that are deposited include fibronectin and collagen, e.g. collagen 1, collagen 2 or collagen 3.

**[0024]** For example, any of the rAAV of the invention may be administered to subjects suffering from muscular dystrophy to prevent fibrosis, e.g. administered before fibrosis is observed in the subject. The rAAV of the invention are administered to the subject suffering from muscular dystrophy in order to prevent new fibrosis in these subjects. These methods may further comprise the step of administering a rAAV expressing micro-dystrophin.

The terms "combination therapy" and "combination treatment" refer to administration of a rAAV vector expressing miR-29 and a rAAV vector expressing micro-dystrophin.

**[0025]** The subject may be suffering from muscular dystrophy such as DMD, Becker muscular dystrophy or any other dystrophin-associated muscular dystrophy. In addition, the subject may be suffering from dystrophinopathy.

**[0026]** The invention provides for recombinant AAV vectors comprising a nucleotide sequence encoding a micro-dystrophin protein. The invention provides for a rAAV comprising a) a muscle-specific control element operably linked to the nucleotide sequence of SEQ ID NO: 7,

or b) the nucleotide sequence of SEQ ID NO: 9.

**[0027]** An exemplary rAAV expressing micro-dystrophin of the invention is the pAAV.mck.micro-dystrophin which comprises the nucleotide sequence of SEQ ID NO: 9 and shown in Figure 10 and 11. This rAAV vector comprises the MCK promoter, a chimeric intron sequence, the coding sequence for the micro-dystrophin gene, polyA, ampicillin resistance and the pGEX plasmid backbone with pBR322 origin of replication. The recombinant AAV vectors of the invention are AAVrh.74.

**[0028]** Described are rAAV vectors expressing micro-dystrophin comprising a nucleotide sequence that has at least 65%, at least 70%, at least 75%, at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, or 89%, more typically at least 90%, 91%, 92%, 93%, or 94% and even more typically at least 95%, 96%, 97%, 98% or 99% sequence identity to SEQ ID NO: 7, and encodes a functional micro-dystrophin protein.

**[0029]** The invention provides for rAAV vectors expressing micro-dystrophin comprising a nucleotide sequence that hybridizes under stringent conditions to the nucleic acid sequence of SEQ ID NOS: 7, or compliments thereof, and encodes a functional micro-dystrophin protein.

**[0030]** The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Hybridization stringency is principally determined by temperature, ionic strength, and the concentration of denaturing agents such as formamide. Examples of stringent conditions for hybridization and washing are 0.015 M sodium chloride, 0.0015 M sodium citrate at 65-68°C or 0.015 M sodium chloride, 0.0015M sodium citrate, and 50% formamide at 42°C. See Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, (Cold Spring Harbor, N.Y. 1989). More stringent conditions (such as higher temperature, lower ionic strength, higher formamide, or other denaturing agent) may also be used, however, the rate of hybridization will be affected. In instances wherein hybridization of deoxyoligonucleotides is concerned, additional exemplary stringent hybridization conditions include washing in 6x SSC 0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos).

**[0031]** Other agents may be included in the hybridization and washing buffers for the purpose of reducing non-specific and/or background hybridization. Examples are 0.1% bovine serum albumin, 0.1% polyvinyl-pyrrolidone, 0.1% sodium pyrophosphate, 0.1% sodium dodecylsulfate, NaDodSO<sub>4</sub>, (SDS), ficoll, Denhardt's solution, sonicated salmon sperm DNA (or other non-complementary DNA), and dextran sulfate, although other suitable agents can also be used. The concentration and types of these additives can be changed without substantially affecting the stringency of the hybridization conditions. Hybridization experiments are usually carried out at pH 6.8-7.4, however, at typical ionic strength conditions, the rate of hybridization is nearly independent of pH. See Anderson et al., *Nucleic Acid Hybridisation: A Practical Approach*, Ch. 4, IRL Press Limited (Oxford, England). Hybridization conditions can be adjusted by one skilled in the art in order to accommodate these variables and allow DNAs of



different sequence relatedness to form hybrids.

**[0032]** In another aspect, the rAAV vectors expressing micro-dystrophin comprises the coding sequence of the micro-dystrophin gene operably linked to a muscle-specific control element. For example, the muscle-specific control element is human skeletal actin gene element, cardiac actin gene element, myocyte-specific enhancer binding factor MEF, muscle creatine kinase (MCK), tMCK (truncated MCK), myosin heavy chain (MHC), C5-12 (synthetic promoter), murine creatine kinase enhancer element, skeletal fast-twitch troponin C gene element, slow-twitch cardiac troponin C gene element, the slow-twitch troponin I gene element, hypoxia-inducible nuclear factors, steroid-inducible element or glucocorticoid response element (GRE).

**[0033]** In addition, the invention provides for rAAV vectors expressing micro-dystrophin comprising a muscle-specific control element comprising the nucleotide sequence of SEQ ID NO: 10 or SEQ ID NO: 11.

**[0034]** The invention also provides for pharmaceutical compositions (or sometimes referred to herein as simply "compositions") comprising any of the rAAV vectors of the invention.

**[0035]** The invention also provides for viral particles comprising any of the recombinant AAV vectors of the invention.

**[0036]** The invention also provides for in vitro methods of producing a functional micro-dystrophin protein comprising infecting a host cell with a recombinant AAV vector expressing micro-dystrophin of the invention and expressing a functional micro-dystrophin protein in the host cell.

**[0037]** Reducing fibrosis in a subject in need may comprise administering a therapeutically effective amount of any rAAV vector of the invention expressing micro-dystrophin. For example, any of the rAAV of the invention are administered to subjects suffering from muscular dystrophy or dystrophinopathy to reduce fibrosis, and in particular reduces fibrosis in skeletal muscle or in cardiac muscle of the subject.

**[0038]** Preventing fibrosis in a subject in need may comprise administering a therapeutically effective amount of the any recombinant AAV vector of the invention expressing micro-dystrophin. For example, any of the rAAV of the invention are administered to subjects suffering from muscular dystrophy or dystrophinopathy to prevent fibrosis, e.g. the rAAV of the invention expressing micro-dystrophin are administered before fibrosis is observed in the subject. In addition, the rAAV of the invention expressing micro-dystrophin are administered to a subject at risk of developing fibrosis, such as those suffering or diagnosed with dystrophinopathy or muscular dystrophy, e.g. DMD or Becker muscular dystrophy. The rAAV of the invention are administered to the subject suffering from dystrophinopathy or dystrophinopathy muscular dystrophy in order to prevent new fibrosis in these subjects.

**[0039]** The terms "combination therapy" and "combination treatment" refer to administration of

a rAAV vector expressing miR-29 and an rAAV vector expressing micro-dystrophin.

**[0040]** When administering an rAAV vector expressing miR-29 and an rAAV vector expressing the micro-dystrophin protein, these rAAV vectors may be administered concurrently, or administered consecutively with the rAAV vector expressing miR29 administered immediately before the rAAV expressing the micro-dystrophin protein, or administered consecutively with the rAAV vector expressing miR29 is administered immediately after the rAAV expressing the micro-dystrophin protein. Alternatively, the AAV vector expressing the micro-dystrophin protein may be administered within about 1-5 hours or 5-12 hours or 12 to 15 hours or 15 to 24 hours after administering the rAAV expressing miR-29 or the AAV vector expressing the micro-dystrophin protein may be administered within about 1-5 hours or 5-12 hours or 12 to 15 hours or 15 to 24 hours before administering the rAAV expressing miR-29. Alternatively, the AAV vector expressing the micro-dystrophin protein may be administered within about 1 or 6 or 12 or 24 hours after administering the rAAV expressing miR-29 or the AAV vector expressing the micro-dystrophin protein may be administered within about 1 or 6 or 12 or 24 hours before administering the rAAV expressing miR-29.

**[0041]** Any of the AAV vectors of the invention may be administered to patients diagnosed with dystrophinopathy or muscular dystrophy, such as DMD or Becker Muscular dystrophy, before fibrosis is observed in the subject or before the muscle force has been reduced in the subject or before the muscle mass has been reduced in the subject.

**[0042]** Any of the rAAV of the invention may be administered to a subject suffering from dystrophinopathy or muscular dystrophy, such as DMD or Becker Muscular dystrophy, who already has developed fibrosis, in order to prevent new fibrosis in these subjects. The invention also provides for administering any of the rAAV of the invention to the patient suffering from muscular dystrophy who already has reduced muscle force or has reduced muscle mass in order to protect the muscle from further injury.

**[0043]** The rAAV vector may be, for example, administered by intramuscular injection or intravenous injection.

**[0044]** The rAAV vector or composition may be administered systemically. For examples, the rAAV vector or composition is parentally administration by injection, infusion or implantation.

**[0045]** In another embodiment, the invention provides for composition comprising the rAAV vectors expressing micro-dystrophin or comprising both a rAAV vector expressing miR-29 and the rAAV vector expressing micro-dystrophin for reducing fibrosis in a subject in need. In addition, the invention provides for compositions comprising the rAAV vectors expressing micro-dystrophin or comprising both a rAAV vector expressing miR-29 and the rAAV vector expressing micro-dystrophin for preventing fibrosis in a patient suffering from dystrophinopathy or muscular dystrophy, such as DMD or Becker Muscular dystrophy.

**[0046]** The invention also provides for compositions comprising the rAAV vectors expressing

micro-dystrophin protein or comprising both a rAAV vector expressing miR-29 and the rAAV vector expressing micro-dystrophin protein for increasing muscular force and/or muscle mass in a subject suffering from dystrophinopathy or muscular dystrophy, such as DMD or Becker Muscular dystrophy.

**[0047]** In a further embodiment, the invention provides for compositions comprising the rAAV vectors expressing micro-dystrophin protein or comprising both a rAAV vector expressing miR-29 and the rAAV vector expressing micro-dystrophin protein for treatment of dystrophinopathy or muscular dystrophy, such as DMD or Becker Muscular dystrophy.

**[0048]** The compositions of the invention are formulated for intramuscular injection or intravenous injection. The composition of the invention is also formulated for systemic administration, such as parentally administration by injection, infusion or implantation. In addition, any of the compositions are formulated for administration to a subject suffering from dystrophinopathy or muscular dystrophy, such as DMD, Becker muscular dystrophy or any other dystrophin associated muscular dystrophy.

**[0049]** The invention also provides for administering any of the rAAV of the invention to the patient suffering from muscular dystrophy who already has reduced muscle force or has reduced muscle mass in order to protect the muscle from further injury.

**[0050]** In any of the uses of the invention, the medicament is formulated for intramuscular injection. In addition, any of the medicaments may be prepared for administration to a subject suffering from muscular dystrophy such as DMD or any other dystrophin associated muscular dystrophy.

**[0051]** In addition, any of the medicaments of the invention may be a combination therapy in which the rAAV vectors expressing miR-29 and rAAV vectors expressing micro-dystrophin are administered concurrently, or administered consecutively with the rAAV vector expressing miR29 administered immediately before the rAAV expressing micro-dystrophin, or administered consecutively with the rAAV vector expressing miR29 administered immediately after the rAAV expressing micro-dystrophin. Alternatively, the medicament comprises administration of the AAV vector expressing micro-dystrophin administered within about 1-5 hours after administering the rAAV expressing miR-29 or the medicament comprises the AAV vector expressing micro-dystrophin administered within about 1-5 hours before administering the rAAV expressing miR-29.

#### **BRIEF DESCRIPTION OF DRAWING**

**[0052]**

**Figure 1** provide a schematic of rAAV vector scAAVCrh.74.CMV.miR29c and the nucleotide sequence of the miR-29c in a natural miR-30 backbone and the nucleotide sequence of the

predicted hairpin structure.

**Figure 2A-CD** illustrates that injection of miR-29c into muscle reduces collagen throughout the muscle and restores miR-29c expression.

**Figure 3A-3C** demonstrates that injection of miR-29c improves absolute muscle force (panel A) and specific muscle force (panel B) but does not protect against contraction-induced damage (panel C).

**Figure 4A-4C** displays the number of muscle fibers expression micro-dystrophin to measure of efficacy of transgene delivery.

**Figure 5A-5C** demonstrates that co-delivery of miR-29c with micro-dystrophin reduces collagen expression (panel A) and fibrosis-induced dystrophin expression.

**Figure 6A-6D** illustrates that intramuscular injection of miR-29c /micro-dystrophin inhibits extracellular matrix (ECM) in mdx/utrn<sup>+/-</sup> mice as measured by collagen 1 alpha (panel A), collagen 3 alpha (panel B), fibronectin (panel C) and TGF- $\beta$  (panel D).

**Figure 7A-7C** demonstrates the intramuscular injection of miR-29c increased absolute force (panel A), normalized specific force (panel B) and added protection from contraction-induced damage (panel C) in the muscle.

**Figure 8** illustrates that the miR-29c/ $\mu$ -dys combination increases muscle size in mice treated at 3 months of age. Sections of treated and untreated mdx/utrn<sup>+/-</sup> gastrocnemius muscles stained with picrosirius Red to stain for collagen are shown. Fibrotic areas are pink and intact muscle is in green. On the macroscopic level, miR-29c/ $\mu$ -dys combination decreases fibrosis and increases total cross sectional area.

**Figure 9A-F** demonstrates that treatment with miR-29c co-delivered with micro-dystrophin increased muscle hypertrophy and hyperplasia as shown by an increase in the overall weight of the injected gastroc compared to either one injected alone (panel A), an increase in the an increase in average fiber size (panel B), an increase in cross-sectional area of the muscle (panel D; uninjected: 24.6 vs. miR-29c: 26.3 vs. micro-dys: 26.6 vs. micro-dys/miR-29c: 33.1) and an increase in the number of muscle fibers (panel E) but the number of muscle fibers per unit area was not affected (panel F). Panel C compares mdx/utrn<sup>+/-</sup> controls with miR-29c/ $\mu$ -dys treated mdx/utrn<sup>+/-</sup>, the average diameter increased from 25.96 to 30.97 $\mu$ m

**Figure 10A-G** demonstrates that early treatment of AAV.miR-29c/micro-dystrophin combination therapy is more effective at reducing fibrosis and ECM expression. Panel A shows picrosirius red staining of wild-type, uninjected, AAV.miR-29c, AAV.micro-dystrophin, and AAV.miR-29c/AAV.micro-dystrophin of mice injected at 4-5 wks of age taken out twelve weeks post-injection. Panel B provides quantification of picrosirius red staining showing co-treated muscle had a 51.1% reduction in collagen compared to uninjected GAS muscle. Panel C demonstrates that qRT-PCR confirms an increase in miR-29c transcript levels in the treated cohorts. Semi-quantitative qRT-PCR shows a significant reduction in collagen I and III (panels

d, e), fbn (panel f) and TGF- $\beta$ 1 (panel g) levels in the AAV.miR-29c/AAV.micro-dystrophin treated muscle compared to the contralateral limb and each of the single therapies Error bars, SEM for n=5 (scAAVrh.74.CMV.miR-29c), n=5 (scAAVrh.74.CMV.miR-29c/ssAAVrh.74.MCK.micro-dystrophin), n=6 (ssAAVrh.74.MCK.micro-dystrophin), n=9(mdx/utrn<sup>+/-</sup> mice). 1-way ANOVA (\*p<0.05, \*\* p<0.01, \*\*\* p<0.001)

**Figure 11** demonstrates early combination therapy restores force and protects against contraction-induced damage. Measurement of absolute (panel A) and normalized specific force (panel b) following tetanic contraction in all three treatment injected GAS muscles were significantly increased compared to untreated mdx/utrn<sup>+/-</sup> muscle (panel C). Muscles were then assessed for loss of force following repetitive eccentric contractions. Only mice co-treated with miR-29c/micro-dystrophin and micro-dystrophin alone showed a protection from loss of force compared with untreated mdx/utrn<sup>+/-</sup> muscles (blue). Two-way analysis of variance demonstrates significance in decay curves Error bars, SEM for n=5 (rAAVrh.74.CMV.miR-29c), n=6 (rAAVrh.74.CMV.miR-29c/rAAVrh.74.MCK.micro-dystrophin), n=5 (rAAVrh.74.MCK.micro-dystrophin), n=15 (mdx/utrn<sup>+/-</sup> mice). 1-way ANOVA (\*p<0.05,\*\*p<0.01, \*\*\* p<0.001, \*\*\*\*p<0.0001).

**Figure 12** illustrates miR-29c/micro-dystrophin combination treatment increases muscle size in mice treated at 1 month of age. Treated and untreated mdx/utrn<sup>+/-</sup> GAS muscles were sectioned and staining with picrosirius Red to stain for collagen. Fibrotic areas are pink and intact muscle is in green. On the macroscopic level, miR-29c/micro-dystrophin combination decreases fibrosis and increases total cross sectional area.

**Figure 13A - 13G** demonstrates that early treatment (at 4-5 weeks) of AAV.MCK.miR-29c/micro-dystrophin combination therapy is more effective at reducing fibrosis and ECM expression. Panel A provide picrosirius red staining of uninjected and AAV.MCK.miR-29c/AAV.MCK.micro-dystrophin of mice injected at 4-5wks of age taken out twelve weeks post-injection. Original magnification, x20 Panel B provides quantification of picrosirius red staining demonstrating co-treated muscle had a 50.9% reduction in collagen compared to untreated GAS muscle Panel C provides qRT-PCR confirming an increase in miR-29c transcript levels in the treated cohort. Semi-quantitative qRT-PCR shows a significant reduction in Collagen 1A (Col1A; panel D) and Collagen 3A (Col3A; panel E), Fibronectin (Fbn; panel F) and Tgf $\beta$ 1 (panel G) levels in the AAV.MCK.miR-29c/AAV.micro-dystrophin treated muscle compared to the contralateral limb therapies. (\*p<0.05,\*\*\*\*p<0.0001).

**Figure 14A - 14G** demonstrates that late treatment (treatment at 12 weeks) with AAV.MCK.miR-29c/micro-dystrophin combination therapy is effective at reducing fibrosis and ECM expression. Panel A provides picrosirius red staining of untreated, AAV.MCK.miR-29c and AAV.MCK.miR-29c/AAV.micro-dystrophin twelve weeks post-injection. Original magnification, x20. Panel B provides quantification of picrosirius red staining which demonstrates that co-treated muscle had a 30.3 % reduction in collagen compared to untreated GAS muscle. Panel C provides qRT-PCR confirming an increase in miR-29c transcript levels in the treated cohorts. Semi-quantitative qRT-PCR demonstrated a significant reduction in Collagen 1A (Col1A; panel

D), Collagen 3A (Col3A; panel E), Fibronectin (Fbn; Panel F) and Tgf $\beta$ 1 (panel G) levels in the AAV.miR-29c/AAV.micro-dystrophin treated muscle compared to the contralateral limb. One-way ANOVA. All data represent mean  $\pm$  SEM. (\*\*  $p < 0.01$ , \*\*\*\* $p < 0.0001$ ).

**Figure 15A-15C** demonstrates that early combination therapy (treatment at 4-5 weeks) restored force and protected against contraction-induced damage. Measurement of absolute (panel A) and normalized specific force (panel B) following tetanic contraction MCK.miR-29c/micro-dystrophin injected GAS muscles were significantly increased compared to untreated mdx/utrn<sup>+/-</sup> muscle. (C) Muscles were then assessed for loss of force following repetitive eccentric contractions. Mice co-treated with miR-29c/micro-dystrophin and micro-dystrophin alone showed protection from loss of force compared with untreated mdx/utrn<sup>+/-</sup> muscles (red). Two-way ANOVA. All data represent mean  $\pm$  SEM (\*\*\*\* $p < 0.0001$ ).

**Figure 16A - 16C** demonstrates that late combination therapy restored force and protected against contraction-induced damage. Measurement of absolute (panel A) and normalized specific force (panel B) following tetanic contraction rAAV.MCK.miR-29c and rAAV expressing micro-dystrophin injected GAS muscles were significantly increased compared to untreated mdx/utrn<sup>+/-</sup> muscle. In Panel C, muscles were then assessed for loss of force following repetitive eccentric contractions. Mice co-treated with rAAV.MCK.miR-29c/rAAV expressing micro-dystrophin showed a protection from loss of force compared with untreated mdx/utrn<sup>+/-</sup> muscles (red). Two-way ANOVA. All data represent mean  $\pm$  SEM (\*\* $p < 0.01$ , \*\*\*\* $p < 0.0001$ ).

**Figure 17A-17D** demonstrates that combination treatment increases muscle hypertrophy 3 months post injection. Panel A demonstrates that rAAV. MCK.miR-29c co-delivered with rAAV expressing micro-dystrophin failed to increase the overall weight of the injected GAS. Panel B demonstrates that rAAV.MCK.miR-29c/rAAV expressing micro-dystrophin combination treatment induced an increase in average fiber size. Comparing mdx/utrn<sup>+/-</sup> controls with miR-29c/micro-dystrophin treated mdx/utrn<sup>+/-</sup>, the average diameter increased from 28.96 to 36.03 $\mu$ m. Panel C shows that co-delivery produced a shift towards wild-type fiber size distribution. Panel D provided the number of muscle fibers per mm<sup>2</sup> in the miR-29c/micro-dystrophin combination treatment was significantly less than untreated mice and wild-type (\*\* $p < 0.01$ , \*\*\*\* $p < 0.0001$ ).

**Figure 18A-18B** provides the nucleic acid sequence (SEQ ID NO: 1 pAAV.CMV.Mir29C) of an exemplary rAAV vector comprising the mature guide strand of miR-29c (nucleotides 1257-1284) and the natural mi-30 backbone (nucleotides 1088-1375). The construct also comprises the CMV promoter (nucleotides 120-526), two EF1a introns at nucleotides 927-1087 and 1380-1854 and a polA at nucleotides 1896-2091.

**Figure 19** provides a schematic of the rAAV vector pAAV.MCK.micro-dystrophin.

**Figure 20A-D** provides the nucleic acid sequence (SEQ ID NO: 9; pAAV.MCK.micro-dystrophin) of an exemplary rAAV vector expressing micro-dystrophin.

**Figure 21A-D** provides the nucleotide sequence of the human micro-dystrophin nucleotide

sequence (SEQ ID NO: 7)

**Figure 22** provides the nucleotide sequence (SEQ ID NO: 12 pAAV.MCK.Mir29C) of an exemplary rAAV vector comprising the mature guide strand of miR-29c (nucleotides 1487-1512) and the natural mi-30 backbone (nucleotides 1088-1375). The construct also comprises the MCK enhancer (nucleotides 190-395), MCK promoter (nucleotides 396-753), two EF1a introns at nucleotides 1155-1315 and 1609-2083 and a polA at nucleotides 2094-2148.

## DETAILED DESCRIPTION

**[0053]** Muscle biopsies taken at the earliest age of diagnosis of DMD reveal prominent connective tissue proliferation. Muscle fibrosis is deleterious in multiple ways. It reduces normal transit of endomysial nutrients through connective tissue barriers, reduces the blood flow and deprives muscle of vascular-derived nutritional constituents, and functionally contributes to early loss of ambulation through limb contractures. Over time, treatment challenges multiply as a result of marked fibrosis in muscle. This can be observed in muscle biopsies comparing connective tissue proliferation at successive time points. The process continues to exacerbate leading to loss of ambulation and accelerating out of control, especially in wheelchair-dependent patients.

**[0054]** Without a parallel approach to reduce fibrosis it is unlikely that the benefits of exon skipping, stop-codon read-through, or gene replacement therapies can ever be fully achieved. Even small molecules or protein replacement strategies are likely to fail without an approach to reduce muscle fibrosis. Previous work in aged *mdx* mice with existing fibrosis treated with AAV.micro-dystrophin demonstrated that we could not achieve full functional restoration (*Human molecular genetics* **22**, 4929-4937 (2013)). It is also known that progression of DMD cardiomyopathy is accompanied by scarring and fibrosis in the ventricular wall. Micro-RNA delivery is particularly innovative because of lack of immune barriers and relative ease of delivery. MicroRNAs are small (~200bp) and can therefore be packaged in AAV along with a therapeutic cassette to correct or bypass the genetic defect.

**[0055]** As used herein, the term "AAV" is a standard abbreviation for adeno-associated virus. Adeno-associated virus is a single-stranded DNA parvovirus that grows only in cells in which certain functions are provided by a co-infecting helper virus. There are currently thirteen serotypes of AAV that have been characterized. General information and reviews of AAV can be found in, for example, Carter, 1989, Handbook of Parvoviruses, Vol. 1, pp. 169-228, and Berns, 1990, Virology, pp. 1743-1764, Raven Press, (New York). However, it is fully expected that these same principles will be applicable to additional AAV serotypes since it is well known that the various serotypes are quite closely related, both structurally and functionally, even at the genetic level. (See, for example, Blacklowe, 1988, pp. 165-174 of Parvoviruses and Human Disease, J. R. Pattison, ed.; and Rose, Comprehensive Virology 3:1-61 (1974)). For example,

all AAV serotypes apparently exhibit very similar replication properties mediated by homologous rep genes; and all bear three related capsid proteins such as those expressed in AAV2. The degree of relatedness is further suggested by heteroduplex analysis which reveals extensive cross-hybridization between serotypes along the length of the genome; and the presence of analogous self-annealing segments at the termini that correspond to "inverted terminal repeat sequences" (ITRs). The similar infectivity patterns also suggest that the replication functions in each serotype are under similar regulatory control.

**[0056]** An "AAV vector" as used herein refers to a vector comprising one or more polynucleotides of interest (or transgenes) that are flanked by AAV terminal repeat sequences (ITRs). Such AAV vectors can be replicated and packaged into infectious viral particles when present in a host cell that has been transfected with a vector encoding and expressing rep and cap gene products.

**[0057]** An "AAV virion" or "AAV viral particle" or "AAV vector particle" refers to a viral particle composed of at least one AAV capsid protein and an encapsidated polynucleotide AAV vector. If the particle comprises a heterologous polynucleotide (i.e. a polynucleotide other than a wild-type AAV genome such as a transgene to be delivered to a mammalian cell), it is typically referred to as an "AAV vector particle" or simply an "AAV vector". Thus, production of AAV vector particle necessarily includes production of AAV vector, as such a vector is contained within an AAV vector particle.

## AAV

**[0058]** Recombinant AAV genomes of the invention comprise nucleic acid molecule of the invention and one or more AAV ITRs flanking a nucleic acid molecule. AAV DNA in the rAAV genomes may be from any AAV serotype for which a recombinant virus can be derived including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12 and AAV-13. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692. Other types of rAAV variants, for example rAAV with capsid mutations, are also contemplated. See, for example, Marsic et al., Molecular Therapy, 22(11): 1900-1909 (2014). As noted in the Background section above, the nucleotide sequences of the genomes of various AAV serotypes are known in the art. To promote skeletal muscle specific expression, AAV1, AAV6, AAV8 or AAVrh.74 may be used.

**[0059]** DNA plasmids may comprise rAAV genomes of the invention. The DNA plasmids are transferred to cells permissible for infection with a helper virus of AAV (e.g., adenovirus, E1-deleted adenovirus or herpes virus) for assembly of the rAAV genome into infectious viral particles. Techniques to produce rAAV particles, in which an AAV genome to be packaged, rep and cap genes, and helper virus functions are provided to a cell, are standard in the art. Production of rAAV requires that the following components are present within a single cell (denoted herein as a packaging cell): a rAAV genome, AAV rep and cap genes separate from (i.e., not in) the rAAV genome, and helper virus functions. The AAV rep and cap genes may be



from any AAV serotype for which recombinant virus can be derived and may be from a different AAV serotype than the rAAV genome ITRs, including, but not limited to, AAV serotypes AAV-1, AAV-2, AAV-3, AAV-4, AAV-5, AAV-6, AAV-7, AAVrh.74, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12 and AAV-13. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692.

**[0060]** A method of generating a packaging cell is to create a cell line that stably expresses all the necessary components for AAV particle production. For example, a plasmid (or multiple plasmids) comprising a rAAV genome lacking AAV rep and cap genes, AAV rep and cap genes separate from the rAAV genome, and a selectable marker, such as a neomycin resistance gene, are integrated into the genome of a cell. AAV genomes have been introduced into bacterial plasmids by procedures such as GC tailing (Samulski et al., 1982, Proc. Natl. Acad. Sci. USA, 79:2077-2081), addition of synthetic linkers containing restriction endonuclease cleavage sites (Laughlin et al., 1983, Gene, 23:65-73) or by direct, blunt-end ligation (Senapathy & Carter, 1984, J. Biol. Chem., 259:4661-4666). The packaging cell line is then infected with a helper virus such as adenovirus. The advantages of this method are that the cells are selectable and are suitable for large-scale production of rAAV. Other examples of suitable methods employ adenovirus or baculovirus rather than plasmids to introduce rAAV genomes and/or rep and cap genes into packaging cells.

**[0061]** General principles of rAAV production are reviewed in, for example, Carter, 1992, Current Opinions in Biotechnology, 1533-539; and Muzyczka, 1992, Curr. Topics in Microbial. and Immunol., 158:97-129). Various approaches are described in Ratschin et al., Mol. Cell. Biol. 4:2072 (1984); Hermonat et al., Proc. Natl. Acad. Sci. USA, 81:6466 (1984); Tratschin et al., Mol. Cell. Biol. 5:3251 (1985); McLaughlin et al., J. Virol., 62:1963 (1988); and Lebkowski et al., 1988 Mol. Cell. Biol., 7:349 (1988). Samulski et al. (1989, J. Virol., 63:3822-3828); U.S. Patent No. 5,173,414; WO 95/13365 and corresponding U.S. Patent No. 5,658,776 ; WO 95/13392; WO 96/17947; PCT/US98/18600; WO 97/09441 (PCT/US96/14423); WO 97/08298 (PCT/US96/13872); WO 97/21825 (PCT/US96/20777); WO 97/06243 (PCT/FR96/01064); WO 99/11764; Perrin et al. (1995) Vaccine 13:1244-1250; Paul et al. (1993) Human Gene Therapy 4:609-615; Clark et al. (1996) Gene Therapy 3:1124-1132; U.S. Patent. No. 5,786,211; U.S. Patent No. 5,871,982; and U.S. Patent. No. 6,258,595.

**[0062]** Described are packaging cells that produce infectious rAAV. In one embodiment packaging cells may be stably transformed cancer cells such as HeLa cells, 293 cells and PerC.6 cells (a cognate 293 line). In another embodiment, packaging cells are cells that are not transformed cancer cells, such as low passage 293 cells (human fetal kidney cells transformed with E1 of adenovirus), MRC-5 cells (human fetal fibroblasts), WI-38 cells (human fetal fibroblasts), Vero cells (monkey kidney cells) and FRhL-2 cells (rhesus fetal lung cells).

**[0063]** Recombinant AAV (*i.e.*, infectious encapsidated rAAV particles) of the invention comprise a rAAV genome. In exemplary embodiments, the genomes of both rAAV lack AAV rep and cap DNA, that is, there is no AAV rep or cap DNA between the ITRs of the genomes. Examples of rAAV that may be constructed to comprise the nucleic acid molecules of the invention are set out in International Patent Application No. PCT/US2012/047999 (WO

2013/016352).

**[0064]** The rAAV may be purified by methods standard in the art such as by column chromatography or cesium chloride gradients. Methods for purifying rAAV vectors from helper virus are known in the art and include methods disclosed in, for example, Clark et al., Hum. Gene Ther., 10(6): 1031-1039 (1999); Schenpp and Clark, Methods Mol. Med., 69 427-443 (2002); U.S. Patent No. 6,566,118 and WO 98/09657.

**[0065]** In another embodiment, the invention contemplates compositions comprising rAAV of the present invention. Compositions of the invention comprise rAAV and a pharmaceutically acceptable carrier. The compositions may also comprise other ingredients such as diluents and adjuvants. Acceptable carriers, diluents and adjuvants are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counter ions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

**[0066]** Titers of rAAV to be administered in methods described herein will vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of rAAV may range from about  $1 \times 10^6$ , about  $1 \times 10^7$ , about  $1 \times 10^8$ , about  $1 \times 10^9$ , about  $1 \times 10^{10}$ , about  $1 \times 10^{11}$ , about  $1 \times 10^{12}$ , about  $1 \times 10^{13}$  to about  $1 \times 10^{14}$  or more DNase resistant particles (DRP) per ml. Dosages may also be expressed in units of viral genomes (vg).

**[0067]** Methods of transducing a target cell with rAAV, *in vivo* or *in vitro*, are described. The *in vivo* methods comprise the step of administering an effective dose, or effective multiple doses, of a composition comprising a rAAV of the invention to an animal (including a human being) in need thereof. If the dose is administered prior to development of a disorder/disease, the administration is prophylactic. If the dose is administered after the development of a disorder/disease, the administration is therapeutic. In embodiments of the invention, an effective dose is a dose that alleviates (eliminates or reduces) at least one symptom associated with the disorder/disease state being treated, that slows or prevents progression to a disorder/disease state, that slows or prevents progression of a disorder/disease state, that diminishes the extent of disease, that results in remission (partial or total) of disease, and/or that prolongs survival. An example of a disease contemplated for prevention or treatment with methods of the invention is FSHD.

**[0068]** Combination therapies are also contemplated by the invention. Combination as used herein includes both simultaneous treatment and sequential treatments. Combinations of methods of the invention with standard medical treatments (e.g., corticosteroids) are

specifically contemplated, as are combinations with novel therapies.

**[0069]** Administration of an effective dose of the compositions may be by routes standard in the art including, but not limited to, intramuscular, parenteral, intravenous, oral, buccal, nasal, pulmonary, intracranial, intraosseous, intraocular, rectal, or vaginal. Route(s) of administration and serotype(s) of AAV components of the rAAV (in particular, the AAV ITRs and capsid protein) of the invention may be chosen and/or matched by those skilled in the art taking into account the infection and/or disease state being treated and the target cells/tissue(s) that are to express the miR-29 miRNA and/or micro-dystrophin.

**[0070]** The invention provides for local administration and systemic administration of an effective dose of rAAV and compositions of the invention including combination therapy of the invention. For example, systemic administration is administration into the circulatory system so that the entire body is affected. Systemic administration includes enteral administration such as absorption through the gastrointestinal tract and parental administration through injection, infusion or implantation.

**[0071]** In particular, actual administration of rAAV of the present invention may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration according to the invention includes, but is not limited to, injection into muscle, the bloodstream and/or directly into the liver. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV (although compositions that degrade DNA should be avoided in the normal manner with rAAV). Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the invention. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

**[0072]** The dose of rAAV to be administered in methods disclosed herein will vary depending, for example, on the particular rAAV, the mode of administration, the treatment goal, the individual, and the cell type(s) being targeted, and may be determined by methods standard in the art. Titers of each rAAV administered may range from about  $1 \times 10^6$ , about  $1 \times 10^7$ , about  $1 \times 10^8$ , about  $1 \times 10^9$ , about  $1 \times 10^{10}$ , about  $1 \times 10^{11}$ , about  $1 \times 10^{12}$ , about  $1 \times 10^{13}$ , about  $1 \times 10^{14}$ , or to about  $1 \times 10^{15}$  or more DNase resistant particles (DRP) per ml. Dosages may also be expressed in units of viral genomes (vg) (i.e.,  $1 \times 10^7$  vg,  $1 \times 10^8$  vg,  $1 \times 10^9$  vg,  $1 \times 10^{10}$  vg,  $1 \times 10^{11}$  vg,  $1 \times 10^{12}$  vg,  $1 \times 10^{13}$  vg,  $1 \times 10^{14}$  vg,  $1 \times 10^{15}$  respectively). Dosages may also be expressed in units of viral genomes (vg) per kilogram (kg) of bodyweight (i.e.,  $1 \times 10^{10}$  vg/kg,  $1 \times 10^{11}$  vg/kg,  $1 \times 10^{12}$  vg/kg,  $1 \times 10^{13}$  vg/kg,  $1 \times 10^{14}$  vg/kg,  $1 \times 10^{15}$  vg/kg respectively). Methods

for titering AAV are described in Clark et al., Hum. Gene Ther., 10: 1031-1039 (1999).

**[0073]** In particular, actual administration of rAAV of the present invention may be accomplished by using any physical method that will transport the rAAV recombinant vector into the target tissue of an animal. Administration according to the invention includes, but is not limited to, injection into muscle, the bloodstream and/or directly into the liver. Simply resuspending a rAAV in phosphate buffered saline has been demonstrated to be sufficient to provide a vehicle useful for muscle tissue expression, and there are no known restrictions on the carriers or other components that can be co-administered with the rAAV (although compositions that degrade DNA should be avoided in the normal manner with rAAV). Capsid proteins of a rAAV may be modified so that the rAAV is targeted to a particular target tissue of interest such as muscle. See, for example, WO 02/053703. Pharmaceutical compositions can be prepared as injectable formulations or as topical formulations to be delivered to the muscles by transdermal transport. Numerous formulations for both intramuscular injection and transdermal transport have been previously developed and can be used in the practice of the invention. The rAAV can be used with any pharmaceutically acceptable carrier for ease of administration and handling.

**[0074]** For purposes of intramuscular injection, solutions in an adjuvant such as sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions. Such aqueous solutions can be buffered, if desired, and the liquid diluent first rendered isotonic with saline or glucose. Solutions of rAAV as a free acid (DNA contains acidic phosphate groups) or a pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. A dispersion of rAAV can also be prepared in glycerol, liquid polyethylene glycols and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

**[0075]** The pharmaceutical carriers, diluents or excipients suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating actions of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of a dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by use of agents delaying absorption, for example,

aluminum monostearate and gelatin.

**[0076]** Sterile injectable solutions are prepared by incorporating rAAV in the required amount in the appropriate solvent with various other ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique that yield a powder of the active ingredient plus any additional desired ingredient from the previously sterile-filtered solution thereof.

**[0077]** Transduction with rAAV may also be carried out *in vitro*. In one embodiment, desired target muscle cells are removed from the subject, transduced with rAAV and reintroduced into the subject. Alternatively, syngeneic or xenogeneic muscle cells can be used where those cells will not generate an inappropriate immune response in the subject.

**[0078]** Suitable methods for the transduction and reintroduction of transduced cells into a subject are known in the art. In one embodiment, cells can be transduced *in vitro* by combining rAAV with muscle cells, e.g., in appropriate media, and screening for those cells harboring the DNA of interest using conventional techniques such as Southern blots and/or PCR, or by using selectable markers. Transduced cells can then be formulated into pharmaceutical compositions, and the composition introduced into the subject by various techniques, such as by intramuscular, intravenous, subcutaneous and intraperitoneal injection, or by injection into smooth and cardiac muscle, using e.g., a catheter.

**[0079]** Transduction of cells with rAAV of the invention results in sustained expression of micro-dystrophin. Described are methods of administering/delivering rAAV which express of miR-29 and or micro-dystrophin to an animal, preferably a human being. These methods include transducing tissues (including, but not limited to, tissues such as muscle, organs such as liver and brain, and glands such as salivary glands) with one or more rAAV of the present invention. Transduction may be carried out with gene cassettes comprising tissue specific control elements. For example, one embodiment provides methods of transducing muscle cells and muscle tissues directed by muscle specific control elements, including, but not limited to, those derived from the actin and myosin gene families, such as from the myoD gene family [See Weintraub et al., Science, 251: 761-766 (1991)], the myocyte-specific enhancer binding factor MEF-2 [Cserjesi and Olson, Mol Cell Biol 11: 4854-4862 (1991)], control elements derived from the human skeletal actin gene [Muscat et al., Mol Cell Biol, 7: 4089-4099 (1987)], the cardiac actin gene, muscle creatine kinase sequence elements [See Johnson et al., Mol Cell Biol, 9:3393-3399 (1989)] and the murine creatine kinase enhancer (mCK) element, control elements derived from the skeletal fast-twitch troponin C gene, the slow-twitch cardiac troponin C gene and the slow-twitch troponin I gene: hypoxia-inducible nuclear factors (Semenza et al., Proc Natl Acad Sci USA, 88: 5680-5684 (1991)), steroid-inducible elements and promoters including the glucocorticoid response element (GRE) (See Mader and White, Proc. Natl. Acad. Sci. USA 90: 5603-5607 (1993)), and other control elements.

**[0080]** Muscle tissue is an attractive target for *in vivo* DNA delivery, because it is not a vital organ and is easy to access. Described is sustained expression of miRNAs from transduced myofibers.

**[0081]** By "muscle cell" or "muscle tissue" is meant a cell or group of cells derived from muscle of any kind (for example, skeletal muscle and smooth muscle, e.g. from the digestive tract, urinary bladder, blood vessels or cardiac tissue). Such muscle cells may be differentiated or undifferentiated, such as myoblasts, myocytes, myotubes, cardiomyocytes and cardiomyoblasts.

**[0082]** The term "transduction" is used to refer to the administration/delivery of the miR29 guide strand or the coding region of the micro-dystrophin to a recipient cell either *in vivo* or *in vitro*, via a replication-deficient rAAV resulting in expression of a miR29 or micro-dystrophin by the recipient cell.

**[0083]** Thus, described are methods of administering an effective dose (or doses, administered essentially simultaneously or doses given at intervals) of rAAV that encode miR29 and/or micro-dystrophin to a patient in need thereof.

## EXAMPLES

### Example 1

#### Confirmation of Duchenne Muscular Dystrophy Models

**[0084]** The mdx mouse provides a convenient, yet incomplete, animal model to study DMD pathogenesis. This model is a cross of the *mdx* mouse with a heterozygous knockout of the utrophin gene (*mdx:utrn+/-*), which presents with increased fibrosis and more faithfully recapitulates the pathology of human DMD. Mdx mice have a nonsense mutation in exon 23 of DMD that results in a relatively mild phenotype and a near-normal life span. By 3 weeks of age, the diaphragm and limb muscle of mdx mice develop signs of endomysial inflammation. These symptoms subside in the limb muscle after the mice reach adulthood while the inflammation in the diaphragm muscle continues to progressively worsen. In mdx mice lacking telomerase, muscular dystrophy progressively worsens with age; mdx mice lacking utrophin (DKO) have a phenotype more characteristic of human DMD with early onset muscle weakness, severe fibrosis, and premature death. Utrophin, an autosomal paralog of the dystrophin, shares a high degree of sequence homology that may compensate for the lack of dystrophin in the mdx mouse in the double KO (dystrophin plus utrophin); a severe phenotype with early death is observed. The premature death in the DKO mouse precludes progression of inflammation and

fibrosis, but the *mdx:utrn*<sup>+/-</sup> mouse presents a model with similarities to the human disease exhibiting a striking degree of fibrosis, and a longer survival than the DKO, providing a better model for our proposed translational studies. A recent report confirms the use of the *mdx:utrn*<sup>+/-</sup> mouse as an ideal model to study fibrosis in the context of DMD. In the present study, increased fibrosis as measured by Sirius red staining was accompanied by increased collagen transcript levels and decreased mir29c levels.

## Example 2

### Delivery of miR29 to DMD Mice Reduces Fibrosis

**[0085]** Preliminary studies have demonstrated that there is a significant increase in Sirius Red staining for collagen and a decrease in miR-29c levels in human DMD patients and the *mdx/utrn*<sup>+/-</sup> mouse. Gene delivery of miR-29 using muscle specific AAV vectors is potentially safe and efficient. To generate the rAAV vector, referred to herein as rAAVrh.74.CMV.miR29c, the 22 nucleotide miR29c sequence (target strand SEQ ID NO: 3 and guide strand SEQ ID NO: 4) was cloned into a miR-30 scaffold driven by a CMV promoter. The expression cassette (SEQ ID NO: 2) was cloned into a self-complementary AAV plasmid and packaged using AAVrh.74, a serotype known to express well in muscle. The miR-29c cDNA was synthesized using a custom primer containing the miR-29c target (sense) strand, miR-30 stem loop and miR-29c guide (antisense) strand in the miR-30 backbone. Three bases of the miR-29c sequence were modified. This sequence was then cloned into a self-complementary AAV ITR containing plasmid driven by the CMV promoter and poly A sequence.

**[0086]** As shown in Figure 1, the pAAV.CMV.miR29C plasmid contains the mir29c cDNA in a miR-30 stem loop backbone flanked by AAV2 inverted terminal repeat sequences (ITR). It is this sequence that was encapsidated into AAVrh.74 virions. In addition, a few nucleotides with in the miR-29c target sequence were changed to mimic Watson-crick pairing at this site as in shRNA-miR(luc). According to ShRNA-luc design, the hairpin should be perfectly complementary throughout its length. Plus, the more changes to the passenger strand, the more likely the elimination of any endogenous mechanism that regulates miR-29 processing that could recognize the miRNA via the stem. The 19<sup>th</sup> base of the guide strand was modified to a cytosine to mimic the nucleotide that precedes the cleavage site in natural mi-29c sequence and the corresponding base on the other strand was changed to preserve pairing.

**[0087]** The gene therapy vector scrAAVrh.74.CMV.miR29c ( $1 \times 10^{11}$  vgs) was injected into the quadriceps muscle of 3 month old *mdx/utrn*<sup>+/-</sup> mice. Quadriceps muscle was analyzed 3 months post-injection by Sirius Red staining and analyzed by NIH ImageJ software as described in Nevo et al. (PloS One, 6: e18049 (2011)). MiR29c, collagen and elastin levels were quantified by RT-PCR. Delivery of miR-29c to young *mdx/utrn*<sup>+/-</sup> mice significantly increases mir-29c

levels and a significant reduction in Sirius red staining in the quadriceps muscle of 6 month old mdx/utrn<sup>+/-</sup> mice (3 months post injection). There was a reduction in collagen and elastin levels in the treated muscles when evaluated by RT-PCR.

**[0088]** Demonstration of increased fibrosis and decreased miR29 expression in the mdx/utrn<sup>+/-</sup> mice and dystrophin-deficient patients validates the mouse model as being representative of the human disease. Initial results using AAV-delivered miR29 as an anti-fibrotic therapy suggest that there is significant beneficial effect with reduction in Sirius Red staining and collagen and elastin levels, which are key contributors in fibrosis.

### Example 3

#### Injection of MiR-29c Reduces Collagen and Restores miR-29c

**[0089]** To determine whether rAAVrh.74.CMV.MiR-29c could reduce fibrosis, 12-week-old mdx/utrn<sup>+/-</sup> mice received an intramuscular injection of rAAVrh.74.CMV.MiR-29c at  $5 \times 10^{11}$  vgs to the left gastrocnemius (GAS) muscle. The mice were analyzed at 12 weeks post injection. Picrosirius red staining revealed a significant decrease in collagen staining throughout the GAS muscles (Fig. 2a) compared to the untreated contralateral mdx/utrn<sup>+/-</sup> GAS muscle. Quantification of the picrosirius red staining shows that treated muscle had a 18.3% reduction in collagen compared to the untreated muscle (treated-  $23.3\% \pm 1.3$  vs. untreated-  $29.5\% \pm 0.7$ ) (Fig2b). To confirm overexpression of miR-29c in treated muscle, total RNA was extracted from the GAS muscle from 24 week old WT, miR-29c treated and mdx/utrn<sup>+/-</sup> mice and subjected to quantitative reverse-transcription -PCR (qRT-PCR) analysis for miR-29c expression. The results showed that miR-29c was significantly increased in the GAS muscle of the treated mice compared to untreated mice (Fig. 2d).

### Example 4

#### MiR-29c Improves Absolute and Specific Muscle Force but does not protect against Contraction-Induced Damage

**[0090]** Knowing that fibrosis can impact muscle function, we wanted to test whether reducing fibrosis by increasing expression of MiR-29c could protect mdx/utrn<sup>+/-</sup> muscle from contraction-induced injury and increase overall force. The functional properties of the gastrocnemius muscle from mdx/utrn<sup>+/-</sup> mice treated with rAAVrh.74.CMV.MiR-29c were assessed. Twelve weeks post injection, the GAS was isolated to perform *in vivo* force measurements.



**[0091]** The GAS procedure follows the protocol listed in Hakim et al., (Methods Mol Biol. 709: 75-89, 2011) for analyzing transverse abdominal muscle physiology but adapted for the GAS. Briefly, mice were anesthetized using ketamine/xylazine mixture. The hind limb skin was removed to expose the GAS muscle and the Achilles tendon. The distal tendon was dissected out and a double square knot was tied around the tendon with 4-0 suture as close to the muscle as possible, another second double square knot is tied right next to the first knot and then tendon is cut. The exposed muscle was constantly dampened with saline. Mice were then transferred to a thermal controlled platform and maintained at 37°. The knee was secured to the platform with a needle through the patella tendon, the tendon suture to the level arm of the force transducer (Aurora Scientific, Aurora, ON, Canada), and the foot was secured with tape. The GAS muscle contractions were elicited by stimulating the sciatic nerve via bipolar platinum electrodes. Once the muscle was stabilized, the optimal length was determined by incremental stretching the muscle until the maximum twitch force was achieved. After a 3-minute rest period, the GAS was stimulated at 50,100,150, and 200 Hz, allowing a 1-minute rest period between each stimulus to determine maximum tetanic force. Muscle length was measured. Following a 5-minute rest, the susceptibility of the GAS muscle to contraction-induced damage was assessed. After 500 ms of stimulation, the muscle was lengthened by 10% of the optimal length. This consisted of stimulating the muscle at 150Hz for 700ms. After the stimulation, the muscle was returned to the optimal length. The cycle was repeated every minute for a total of 5 cycles. Specific force was calculated by dividing the maximum tetanic force by the GAS muscle cross sectional area. After the eccentric contractions, the mice were then euthanized and the GAS muscle was dissected out, weighed and frozen for analysis.

**[0092]** Each GAS was subjected to a series of repeated eccentric contraction. By comparing the force ratio of each contraction versus the first contraction revealed that after the fifth contraction untreated muscle decayed to  $0.56 \pm 0.05$  versus treated  $0.50 \pm 0.04$  ( $p \leq 0.0001$ ). The injected group showed a slight decrease in the degree of protection compared to WT controls, that decayed to  $0.92 \pm 0.02$  (Fig. 3c). This data shows that reducing fibrosis by increasing expression of miR-29c leads to increase in both absolute and specific force but does not significantly protect muscle from contraction-induced injury.

**[0093]** rAAVrh.74.MiR-29c treated GAS muscle showed significant improvement in absolute force when compared to untreated mdx/utrn<sup>+/-</sup> GAS muscle (rAAV.miR-29c-  $2277 \pm 161.7$  vs. mdx/utrn<sup>+/-</sup> untreated-  $1722 \pm 145.7$ ; Fig. 3a), and also normalized specific force in rAAVrh.74.miR-29c treated GAS muscle specific improvement when compared to untreated GAS muscle (rAAV.miR-29c-  $204.7 \pm 11.7$  vs. mdx/utrn<sup>+/-</sup> untreated-  $151.6 \pm 14.5$ ; Fig. 3b). Force was still significantly reduced when compared to wild-type controls (rAAV.miR-29c-  $204.7 \pm 11.7$  vs. wild-type-  $312.0 \pm 34.1$ ).

## Example 5

### Co-delivery with Micro-Dystrophin Further Reduces fibrosis

**[0094]** To determine whether miR-29c/micro-dystrophin combined gene therapy approach would be more beneficial at reducing fibrosis, 12-week-old mdx/utrn<sup>+/-</sup> mice received an intramuscular injection of rAAVrh.74.CMV.MiR-29c at  $5 \times 10^{11}$  vgs to the left gastrocnemius muscle. The following gene therapy vectors were administered by intramuscular injection (IM) into the left gastrocnemius (GAS) muscle of 3 month old mdx/utrn<sup>+/-</sup> mice, a DMD mouse model: scAAVrh.74.CMV.miR-29c alone, co-delivered with rAAVrh.74.MCK.micro-dystrophin, and rAAVrh.74.MCK.micro-dystrophin alone.

**[0095]** The pAAV.MCK.micro-dystrophin plasmid contains the human micro-dystrophin cDNA expression cassette flanked by AAV2 inverted terminal repeat sequences (ITR) as shown in Fig. 10. It is this sequence that was encapsidated into AAV rh.74 virions. The pAAV.MCK.micro-dystrophin plasmid was constructed by inserting the MCK expression cassette driving a codon optimized human micro-dystrophin cDNA sequence into the AAV cloning vector as described in Rodino-Klapac et al. (Mol Ther. 2010 Jan;18(1):109-17). A MCK promoter/enhancer sequence was used to drive muscle-specific gene expression and is composed of the mouse MCK core enhancer (206 bp) fused to the 351 bp MCK core promoter (proximal). After the core promoter, the 53 bp endogenous mouse MCK Exon1 (untranslated) is present for efficient transcription initiation, followed by the SV40 late 16S/19S splice signals (97 bp) and a small 5'UTR (61 bp). The intron and 5' UTR are derived from plasmid pCMV $\beta$  (Clontech). The micro-dystrophin cassette has a consensus Kozak immediately in front of the ATG start and a small 53 bp synthetic polyA signal for mRNA termination. The human micro-dystrophin cassette contains the (R4-R23/ $\Delta$ 71-78) domains. The complementary DNA was codon optimized for human usage and synthesized by GenScript (Piscataway, NJ).

**[0096]** The mice were analyzed at 12 and 24 weeks post injection. First, the number of muscle fibers expressing micro-dystrophin was used to assess the efficacy of transgene delivery and to make sure we had similar levels of micro-dystrophin expressed in each group. We found that micro-dystrophin was not different between cohorts treated with micro-dystrophin alone ( $71.85 \pm 2.25\%$ ) compared with miR-29c/micro-dystrophin combination therapy ( $75.03 \pm 1.91\%$ ) (Fig.4).

**[0097]** GAS muscle was analyzed 12 months post-injection to assess collagen accumulation by Sirius Red staining and subsequent quantification with ImageJ. Additional outcomes included miR-29c and collagen transcript levels, force measurements in the GAS muscle, fiber diameter measurements and western blot analysis for proteins involved in muscle regeneration (MyoD, Myogenin). The amount of fibrosis was analyzed by picosirius red staining, which revealed a significant decrease in collagen staining throughout the GAS muscles in all treated groups (Fig. 5a) compared to the untreated contralateral mdx/utrn<sup>+/-</sup> GAS muscle or micro-dystrophin alone. Quantification of the picosirius red staining shows that co-treated muscle had a 40.8% reduction in collagen compared to the untreated muscle (treated-  $17.47\% \pm 0.75$  vs. untreated-  $29.5\% \pm 0.7$ ) (Fig. 5b). To confirm expression of miR-29c, qRT-PCR was performed on the GAS

muscle and all treatment groups had an increase in miR-29c compared to untreated muscle (Fig. 5c).

**[0098]** Analogous to DMD tissue, a significant reduction in miR-29c levels in mdx/utrn<sup>+/−</sup> muscle was observed which correlated with increased fibrosis measured by picrosirius red staining. Following 3 months of treatment with scAAV.miR-29c alone, there was a significant reduction in fibrosis (treated-23.5%±1.3 vs. untreated-27.8% ±0.6) in the GAS muscle. When co-delivered with micro-dystrophin, further reduction in collagen (41%) was observed by picrosirius red staining (combination treatment: 17.47%±0.75 vs. untreated: 29.5% ±0.7) (p<0.0001) (Fig. 5b). To confirm expression of miR-29c, qRT-PCR was performed on the GAS muscle and all treatment groups had an increase in miR-29c compared to untreated muscle (Fig. 5b).

**[0099]** At 24 weeks post-injection, the results were similar to those observed 12 weeks post injection. There was a 47% reduction in collagen by picrosirius red staining compared to the untreated muscle (combination treatment: 16.5±1.23 vs. untreated: 31.07±0.93; p<0.0001) and a coincident increase in miR-29c transcript level.

**[0100]** To further validate reduction of collagen observed by picrosirius red staining, qRT-PCR was performed on the muscle to quantify transcript levels of Col1A, Col3A and also another ECM component, fibronectin (Fbn). qRT-PCR analysis detected a decrease in Col1A and Col3A following each treatment, however only the cohort treated with both micro-dystrophin and miR-29c showed significant reduction (Fig. 6a and 6b). The analysis revealed that Fbn was significantly reduced only in the co-treated cohort (Fig. 6c).

**[0101]** TGF-β1 has been previously shown to be up regulated in dystrophic muscle, likely playing a role in the initiation of the fibrotic cascade. TGF-β1 is a known pro-fibrotic cytokine that down regulates miR-29c and is responsible for conversion of myoblasts to myofibroblasts with an increase in collagen and muscle fibrogenesis. qRT-PCR analysis shows that co-treated muscle had significantly lower levels of TGF-β1 compared to uninjected muscle and either treatment alone (Fig. 6d). At 6 months post injection, co-treated muscle continued to show reduced Col1A, Col3A, Fbn and TGF-β1 levels, whereas only slight reductions in Col1A mRNA levels in the miR-29 and the micro-dystrophin only groups were observed

**[0102]** An increase in specific and absolute force was observed in the muscle treated with miR-29c alone compared to the untreated limb, which when combined with micro-dystrophin led to absolute and specific force that were not significantly different than wild-type. We also observed a significant increase in gastroc weight in those muscles that were co-treated.

**[0103]** Initial results using rAAV.miR-29c as an anti-fibrotic therapy suggest that there is beneficial effect with reduction in collagen levels, a key contributor in fibrosis. Moreover, when combined with micro-dystrophin to improve membrane stability, miR29 up regulation normalized muscle force.

**Example 6****Further Increase in Absolute Force and Added Protection from Contraction-induced Damage**

[0104] Knowing that miR-29-treated muscle had a modest but significant increase in absolute and specific force, the combination therapy of miR-29c overexpression and micro-dystrophin gene replacement impact on muscle function was investigated. Twelve weeks post injection, we isolated the GAS for which we performed in vivo force measurements. The rAAVrh.74.MiR-29c vector described above in Example 2 and a rAAV

[0105] Co-treated rAAVrh.74.MiR-29c and rAAV expressing Micro-Dys treated GAS muscle showed significant improvement in absolute force when compared to untreated mdx/utrn<sup>+/-</sup> GAS muscle (co-treated- 3582.4 ± 79.4 nM vs. mdx/utrn<sup>+/-</sup> untreated- 1722 ± 145.7 nM vs. wild-type- 3005 ± 167.3 nM) (Fig. 7), and also normalized specific force in rAAVrh.74.miR-29c/micro-dys treated GAS muscle specific improvement when compared to untreated GAS muscle (co-treated mice- 244.2 ± 6.6 nM/mm<sup>2</sup> vs. mdx/utrn<sup>+/-</sup> untreated- 151.6 ± 14.5 nM/mm<sup>2</sup> vs. 312.0 ± 34.1 nM/mm<sup>2</sup>) (Fig 7). Both absolute and specific force was not significantly different from wild-type controls.

[0106] Each GAS was subjected to a series of repeated eccentric contraction. By comparing the force ratio of each contraction versus the first contraction revealed that after the fifth contraction untreated muscle decayed to 0.54 ± 0.06 versus co-treated 0.66 ± 0.04 ( $p \leq 0.0001$ ), which can be contributed to the micro-dystrophin since the micro-dystrophin alone also decayed to 0.66 ± 0.04. The treated group was still significantly lower than wild-type that decayed to 0.92 ± 0.02 (Fig. 7c). Similar findings were seen at 24 weeks post injection. This data shows that reducing fibrosis and gene replacement leads to increase in both absolute and specific force and significantly protects muscle from contraction-induced injury.

**Example 7****Combination treatment increases muscle hypertrophy and hyperplasia**

[0107] MiR-29c co-delivered with micro-dystrophin increased the overall weight of the injected gastroc compared to either one injected alone at three months of age (Fig. 8, Fig. 9a). To investigate the source of increased muscle mass, myofiber diameters are measured. miR-29c/μ-dys combination treatment demonstrated an increase in average fiber size. Comparing

mdx/utrn<sup>+/-</sup> controls with miR-29c/ $\mu$ -dys treated mdx/utrn<sup>+/-</sup>, the average diameter increased from 25.96 to 30.97 $\mu$ m (Fig. 9b). The co-delivery produced a shift towards wild-type fiber size distribution (Fig. 9c). Although the average fiber size was increased does not explain the ~30% increase in gross muscle weight. Total cross-sectional area of the muscle was also measured. Gastroc muscles from all groups were full slide scanned and the total area was measured. Muscles co-treated with micro-dys/miR-29c had a significant increase in cross sectional area compared to untreated and either treatment alone (uninjected: 24.6 vs. miR-29c: 26.3 vs. micro-dys: 26.6 vs. micro-dys/miR-29c: 33.1) (Fig. 8, Fig. 9d).

**[0108]** miR-29c has been reported it to play a role in the myoD/Pax7/myogenin pathway and it was hypothesized that miR-29c may be impacting regeneration and activation of satellite cells (muscle stem cells) to differentiate in myogenic lineage. To test this, the total number of muscle fibers from the full slide scanned images was counted. An increased number of muscle fibers following miR-29c/ $\mu$ -dys combination treatment (Fig. 9e). Finally, given that muscle fiber diameters in mdx/utrn<sup>+/-</sup> mice are heterogeneous with many small fibers and some hypertrophic fibers, it was determined whether the number of fibers per unit area (cells/mm<sup>2</sup>) was affected with treatment. miR-29c/ $\mu$ -dys combination treatment was not different than wild-type (Fig. 9f).

### Example 8

#### Early Treatment With Combination Prevents Fibrosis

**[0109]** In view of the potential importance of combinatorial miR-29c and micro-dystrophin as a prophylactic therapy for DMD, a cohort of younger mdx/utrn<sup>+/-</sup> mice were treated at 4 weeks of age. Using the same paradigm as for other groups as described herein, the following treatments were compared for efficacy for prevention of fibrosis by intramuscular injection of GAS: scAAVrh.74.CMV.miR-29c alone, ssAAVrh74.MCK.micro-dystrophin + scAAVrh.74.CMV.miR-29c combination therapy, or ssAAVrh74.MCK.micro-dystrophin alone at the same dose. The mice were necropsied 12 weeks post injection. A significant decrease in collagen staining throughout the GAS muscles in all treated groups compared to the untreated contralateral mdx/utrn<sup>+/-</sup> GAS muscle was observed (Fig. 10A). Quantification of the picosirius red staining showed that muscle co-treated with micro-dystrophin/miR-29c had a 51% reduction in collagen compared to the untreated muscle (treated- 11.32% $\pm$  1.18 vs. untreated- 23.15%  $\pm$  0.90) ( $p < 0.0001$ ) (Fig. 10) and qRT-PCR confirmed Col1A, Col3A, Fbn and TGF- $\beta$ 1 reduction following combinatorial therapy (Fig. 10D and E).

### Example 9

#### Early Combination Therapy Restores Force and Protects from Contraction-Induced

## Damage Better than Late Treatment

[0110] *In vivo* force measurement was carried out on the GAS of the mice treated early with the combination therapy as described in Example 8. In 4-week-old mdx/utrn<sup>+/-</sup> mice, co-treatment using miR-29c/micro-dystrophin showed significant improvement in absolute force when compared to untreated mdx/utrn<sup>+/-</sup> mice and there was no difference from wild type (co-treated: 2908± 129.5 mN vs. untreated: 1639.4± 116.9 mN vs. wild-type: 3369.73± 154.1 mN). Specific force was also normalized to wild type levels following combinatorial therapy (co-treated 338.9 ± 22.34 mN/mm<sup>2</sup> vs. untreated 184.3 ± 13.42 mN/mm<sup>2</sup> vs. WT 364.3 ± 7.79 mN/mm<sup>2</sup>) (Fig. 11A and B and 12).

[0111] Next, each GAS was subjected to a series of repeat eccentric contractions. By comparing the force ratio of each contraction by the fifth contraction, untreated muscle decayed to 0.53± 0.04 versus co-treated 0.82± 0.04 ( $p \leq 0.0001$ ). The combinatorial treatment group was slightly lower than wild type but not significantly different, which decayed to 0.93 ± 0.01 (Fig. 11C). These data show that reducing fibrosis and gene replacement lead to increase in both absolute and specific force and significantly protects muscle from contraction-induced injury.

[0112] These experiments suggest that gene replacement should be started in the newborn period. Efforts are clearly moving in the direction of identifying DMD and other muscular dystrophies in the newborn period. The Ohio Newborn Screening Study illustrates the potential for identification of DMD in newborns using CK 7 Neurol. as a biomarker (>2000 U/L) with DNA confirmation on the same dried blood spot (Mendell et al., Ann. Neurol. 71: 304-313, 2012). This methodology is now being extended to other states in the USA (PPMD May 16, 2016: Next Steps with Newborn Screening) and in other countries, particularly the UK (UK National Screening Committee) and China (Perkin Elmer<sup>™</sup> launches screening in China).

[0113] miR-29 has also shown promise as a treatment modality for cardiac, pulmonary, and liver fibrosis. Myocardial infarction in mice and humans is associated with miR-29 down-regulation. Rooij et al. (Proc. Natl. Acad. Sci. USA 105:13027-13032, 2008) demonstrated that exposing fibroblasts to a miR-29b mimic decreased collagen transcripts providing a path for clinical translation for cardiac fibrosis. Subsequent studies showed that in a bleomycin-induced pulmonary fibrosis mouse model, attenuation of fibrosis could be achieved using the Sleeping Beauty (SB) transposon system-based delivery of miR-29b.<sup>14</sup> Currently, a miR-29b mimic is in a clinical Phase 1 Safety-Tolerability local intradermal trial in healthy volunteers (miRagen Therapeutics<sup>™</sup> MRG-201). Compared to miR-29 oligonucleotide delivery that would require repeated administration related to the half-life of the oligonucleotides, AAV gene therapy could potentially provide a path for single-delivery gene transfer.

## Example 10

## Treatment with Muscle Specific Expression of miR-29 and Micro-dystrophin Reduced Fibrosis and ECM Expression

**[0114]** AAV vectors comprising the miR29c sequence and a muscle specific promoter MCK were also generated and tested as a combination therapy with AAV vectors expressing micro-dystrophin. To generate the rAAV vector, referred to herein as rAAV.MCK.miR29c, the 22 nucleotide miR29c sequence (target strand SEQ ID NO: 3 and guide strand SEQ ID NO: 4) was cloned into a miR-30 scaffold driven by a MCK promoter (SEQ ID NO: 11). The expression cassette (SEQ ID NO: 12) was cloned into a single stranded AAV plasmid and packaged using AAVrh74, a serotype known to express well in muscle. The miR-29c cDNA was synthesized using a custom primer containing the miR-29c target (sense) strand, miR-30 stem loop and miR-29c guide (antisense) strand in the miR-30 backbone. Three bases of the miR-29c sequence were modified. This sequence was then cloned into a single stranded AAV ITR containing plasmid driven by the MCK promoter and polyA sequence.

**[0115]** The pAAV.MCK.miR29C plasmid contains the mir29c cDNA in a miR-30 stem loop backbone flanked by AAV2 inverted terminal repeat sequences (ITR). It is this sequence that was encapsidated into AAVrh74 virions. In addition, a few nucleotides within the miR-29c target sequence were changed to mimic Watson-crick pairing at this site as in shRNA-miR(luc). According to ShRNA-luc design, the hairpin should be perfectly complementary throughout its length. Plus, the more changes to the passenger strand, the more likely the elimination of any endogenous mechanism that regulates miR-29 processing that could recognize the miRNA via the stem. The 19<sup>th</sup> base of the guide strand was modified to a cytosine to mimic the nucleotide that precedes the cleavage site in natural mi-29c sequence and the corresponding base on the other strand was changed to preserve pairing.

**[0116]** Early treatment of AAV.MCK.miR-29c/micro-dystrophin combination therapy was more effective at reducing fibrosis and ECM expression. 4-5-week-old mdx/utrn<sup>+/-</sup> mice received an intramuscular injection of rAAVrh.74.MCK.MiR-29c and rAAVrh74.MCK.micro-dystrophin at  $5 \times 10^{11}$  vgs to the left gastrocnemius muscle as described in Example 5. The muscles were harvested twelve weeks post injection. Picrosirius red staining of muscle harvested from uninjected and mice injected with combination therapy of rAAV.MCK.miR-29c/rAAV.MCK.micro-dystrophin showed co-treated muscle had a 50.9% reduction in collagen compared to untreated GAS muscle (See Fig. 13a and 13b). qRT-PCR confirmed an increase in miR-29c transcript levels in the treated cohort (Fig. 13c). Semi-quantitative qRT-PCR showed a significant reduction in Collagen A1 and Collagen 3A (Fig. 13d, e), Fibronectin (Fig. 13f) and Tgf $\beta$ 1 (Fig. 13g) levels in the AAV.MCK.miR-29c/AAV.micro-dystrophin treated muscle compared to the contralateral limb therapies. (\*p<0.05, \*\*\*\*p<0.0001). Late treatment of AAV.MCK.miR-29c/micro-dystrophin combination therapy is effective at reducing fibrosis and ECM expression. Three month old mdx/utrn<sup>+/-</sup> mice received an intramuscular injection of

rAAVrh.74.MCK.MiR-29c and rAAVrh.74.MCK.micro-dystrophin at  $5 \times 10^{11}$  vgs to the left gastrocnemius muscle as described in Example 5. The muscles were harvested twelve weeks post injection. Picrosirius red staining of untreated, AAV.MCK.miR-29c and AAV.MCK.miR-29c/AAV.micro-dystrophin treated muscle showed co-treated muscle had a 30.3 % reduction in collagen compared to untreated GAS muscle (See Fig. 14a and 14b) qRT-PCR confirmed an increase in miR-29c transcript levels in the treated cohorts (Fig. 14c). Semi-quantitative qRT-PCR shows a significant reduction in Collagen 1A and Collagen 3A (Fig. 14d, e), Fibronectin (Fig. 14f) and Tgf $\beta$ 1 (Fig. 14G) levels in the AAV.miR-29c/AAV.micro-dystrophin treated muscle compared to the contralateral limb. One-way ANOVA. All data represent mean  $\pm$  SEM. (\*\*  $p < 0.01$ , \*\*\*\*  $p < 0.0001$ ).

### Example 11

#### Early Combination Therapy Restores Force and Protects from Contraction-Induced Damage Better than Late Treatment

[0117] *In vivo* force measurement was carried out on the GAS of the mice treated early with the muscle-specific expression of miR-29 and micro-dystrophin. as described in Examples 8 and 9. In 4-week-old mdx/utrn<sup>+/-</sup> mice, co-treatment using rAAV.MCK.miR-29c/and rAAV expressing micro-dystrophin showed significant improvement in absolute force when compared to untreated mdx/utrn<sup>+/-</sup> mice and there was no difference from wild type (Fig. 15a). Specific force was also normalized to wild type levels following combination therapy (Fig. 15b).

[0118] Muscles were then assessed for loss of force following repetitive eccentric contractions as described in Example 9. Mice co-treated with rAAV.MCK.miR-29c/rAAV.MCK.micro-dystrophin and rAAV.MCK.micro-dystrophin alone showed a protection from loss of force compared with untreated mdx/utrn<sup>+/-</sup> muscles (Fig. 15c).

[0119] In 12-week-old mdx/utrn<sup>+/-</sup> mice, co-treatment using rAAV.MCK.miR-29c/and rAAV expressing micro-dystrophin restored force and protected against contraction-induced damage. Measurement of absolute (Fig. 16a) and normalized specific force (Fig. 16b) following tetanic contraction rAAV.MCK.miR-29c and rAAV expressing micro-dystrophin injected GAS muscles were significantly increased compared to untreated mdx/utrn<sup>+/-</sup> muscle. Subsequently, muscles were assessed for loss of force following repetitive eccentric contractions as described in Example 9. Mice co-treated with MCK.miR-29c/micro-dystrophin showed a protection from loss of force compared with untreated mdx/utrn<sup>+/-</sup> muscles (Fig. 16c). These data show that reducing fibrosis and gene replacement lead to increase in both absolute and specific force and significantly protects muscle from contraction-induced injury.

### Example 12



### Early Combination treatment increases muscle hypertrophy and hyperplasia

**[0120]** Co-delivery of rAAV.MCK.miR-29 with rAAV expressing micro-dystrophin did not increase overall weight of the injected gastroc compared to either one injected alone at three months post-injection (Fig. 17a). Myofiber diameters were also measured. miR-29c/micro-dystrophin combination treatment demonstrated an increase in average fiber size. Comparing mdx/utrn<sup>+/-</sup> controls with miR-29c/micro-dystrophin treated mdx/utrn<sup>+/-</sup>, the average diameter increased from 28.96 to 36.03µm (Fig. 17b). The co-delivery produced a shift towards wild-type fiber size distribution (Fig. 17c). The number of muscle fibers per mm<sup>2</sup> in the miR-29c/micro-dystrophin combination treatment was significantly less than untreated mice and wild-type (Fig. 17d; \*\*\*p<0.01, \*\*\*\*p<0.0001).

### REFERENCES

#### **[0121]**

1. Hoffman, E.P., Brown, R.H., Jr. & Kunkel, L.M. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* 51, 919-928 (1987).
2. Straub, V. & Campbell, K.P. Muscular dystrophies and the dystrophin-glycoprotein complex. *Curr Opin Neurol* 10, 168-175 (1997).
3. Sacco, A., et al. Short telomeres and stem cell exhaustion model Duchenne muscular dystrophy in mdx/mTR mice. *Cell* 143, 1059-1071 (2010).
4. Wallace, G.Q. & McNally, E.M. Mechanisms of muscle degeneration, regeneration, and repair in the muscular dystrophies. *Annu Rev Physiol* 71, 37-57 (2009).
5. Zhou, L. & Lu, H. Targeting fibrosis in Duchenne muscular dystrophy. *J Neuropathol Exp Neurol* 69, 771-776 (2010).
6. Desguerre, I., et al. Endomysial fibrosis in Duchenne muscular dystrophy: a marker of poor outcome associated with macrophage alternative activation. *J Neuropathol Exp Neurol* 68, 762-773 (2009).
7. Kim, J., et al. microRNA-directed cleavage of ATHB 15 mRNA regulates vascular development in Arabidopsis inflorescence stems. *Plant J* 42, 84-94 (2005).
8. Ambros, V. MicroRNA pathways in flies and worms: growth, death, fat, stress, and timing. *Cell* 113, 673-676 (2003).
9. Eisenberg, I., et al. Distinctive patterns of microRNA expression in primary muscular disorders. *Proc Natl Acad Sci U S A* 104, 17016-17021 (2007).
10. Jiang, X., Tsitsiou, E., Herrick, S.E. & Lindsay, M.A. MicroRNAs and the regulation of fibrosis. *FEBS J* 277, 2015-2021 (2010).
11. van Rooij, E., et al. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci U S A* 105, 13027-13032 (2008).

12. 12. Cacchiarelli, D., et al. MicroRNAs involved in molecular circuitries relevant for the Duchenne muscular dystrophy pathogenesis are controlled by the dystrophin/nNOS pathway. *Cell Metab* 12, 341-351 (2010).
13. 13. DiPrimio, N., McPhee, S.W. & Samulski, R.J. Adeno-associated virus for the treatment of muscle diseases: toward clinical trials. *Curr Opin Mol Ther* 12, 553-560 (2010).
14. 14. Mendell, J.R., et al. Sustained alpha-sarcoglycan gene expression after gene transfer in limb-girdle muscular dystrophy, type 2D. *Ann Neurol* 68, 629-638 (2010).
15. 15. Mendell, J.R., et al. Limb-girdle muscular dystrophy type 2D gene therapy restores alpha-sarcoglycan and associated proteins. *Ann Neurol* 66, 290-297 (2009).
16. 16. Mendell, J.R., et al. A phase 1/2a follistatin gene therapy trial for becker muscular dystrophy. *Molecular therapy : the journal of the American Society of Gene Therapy* 23, 192-201 (2015).
17. 17. Carnwath, J.W. & Shotton, D.M. Muscular dystrophy in the mdx mouse: histopathology of the soleus and extensor digitorum longus muscles. *J Neural Sci* 80, 39-54 (1987).
18. 18. Coulton, G.R., Morgan, J.E., Partridge, T.A. & Sloper, J.C. The mdx mouse skeletal muscle myopathy: I. A histological, morphometric and biochemical investigation. *Neuropathol Appl Neurobiol* 14, 53-70 (1988).
19. 19. Cullen, M.J. & Jaros, E. Ultrastructure of the skeletal muscle in the X chromosome-linked dystrophic (mdx) mouse. Comparison with Duchenne muscular dystrophy. *Acta Neuropathol* 77, 69-81 (1988).
20. 20. Dupont-Versteegden, E.E. & McCarter, R.J. Differential expression of muscular dystrophy in diaphragm versus hindlimb muscles of mdx mice. *Muscle Nerve* 15, 1105-1110 (1992).
21. 21. Stedman, H.H., et al. The mdx mouse diaphragm reproduces the degenerative changes of Duchenne muscular dystrophy. *Nature* 352, 536-539 (1991).
22. 22. Deconinck, A.E., et al. Utrophin-dystrophin-deficient mice as a model for Duchenne muscular dystrophy. *Cell* 90, 717-727 (1997).
23. 23. Grady, R.M., et al. Skeletal and cardiac myopathies in mice lacking utrophin and dystrophin: a model for Duchenne muscular dystrophy. *Cell* 90, 729-738 (1997).
24. 24. Love, D.R., et al. An autosomal transcript in skeletal muscle with homology to dystrophin. *Nature* 339, 55-58 (1989).
25. 25. Tinsley, J.M., et al. Primary structure of dystrophin-related protein. *Nature* 360, 591-593 (1992).
26. 26. Tinsley, J., et al. Expression of full-length utrophin prevents muscular dystrophy in mdx mice. *Nat Med* 4, 1441-1444 (1998).
27. 27. Squire, S., et al. Prevention of pathology in mdx mice by expression of utrophin: analysis using an inducible transgenic expression system. *Hum Mol Genet* 11, 3333-3344 (2002).
28. 28. Rafael, J.A., Tinsley, J.M., Potter, A.C., Deconinck, A.E. & Davies, K.E. Skeletal muscle-specific expression of a utrophin transgene rescues utrophin-dystrophin deficient mice. *Nat Genet* 19, 79-82 (1998).
29. 29. Zhou, L., et al. Haploinsufficiency of utrophin gene worsens skeletal muscle

- inflammation and fibrosis in mdx mice. *J Neural Sci* 264, 106-111 (2008).
30. 30. Gutpell, K.M., Hrinivich, W.T. & Hoffman, L.M. Skeletal Muscle Fibrosis in the mdx/utrn+/- Mouse Validates Its Suitability as a Murine Model of Duchenne Muscular Dystrophy. *PloS one* 10, e0117306 (2015).
  31. 31. Rodino-Klapac, L.R., et al. Micro-dystrophin and follistatin co-delivery restores muscle function in aged DMD model. *Human molecular genetics* 22, 4929-4937 (2013).
  32. 32. Cushing, L., et al. MIR-29 is a Major Regulator of Genes Associated with Pulmonary Fibrosis. *Am J Respir Cell Mol Biol* (2010).
  33. 33. Roderburg, C., et al. Micro-RNA profiling reveals a role for miR-29 in human and murine liver fibrosis. *Hepatology* 53, 209-218 (2011).
  34. 34. Nevo, Y., et al. The Ras antagonist, farnesylthiosalicylic acid (FTS), decreases fibrosis and improves muscle strength in dy/dy mouse model of muscular dystrophy. *PloS one* 6, e18049 (2011).
  35. 35. Rodino-Klapac, L.R., et al. A translational approach for limb vascular delivery of the micro-dystrophin gene without high volume or high pressure for treatment of Duchenne muscular dystrophy. *J Transl Med* 5, 45 (2007).
  36. 36. Mulieri, L.A., Hasenfuss, G., Ittleman, F., Blanchard, E.M. & Alpert, N.R. Protection of human left ventricular myocardium from cutting injury with 2,3-butanedione monoxime. *Circ Res* 65, 1441-1449 (1989).
  37. 37. Rodino-Klapac, L.R., et al. Persistent expression of FLAG-tagged micro dystrophin in nonhuman primates following intramuscular and vascular delivery. *Molecular therapy : the journal of the American Society of Gene Therapy* 18, 109-117 (2010).
  38. 38. Grose, W.E., et al. Homologous recombination mediates functional recovery of dysferlin deficiency following AAV5 gene transfer. *PloS one* 7, e39233 (2012).
  39. 39. Liu, M., et al. Adeno-associated virus-mediated microdystrophin expression protects young mdx muscle from contraction-induced injury. *Mol Ther* 11, 245-256 (2005).

## REFERENCES CITED IN THE DESCRIPTION

### Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [US7282199B](#) [0006]
- [US7790449B](#) [0006]
- [WO0183692A](#) [0058] [0059]
- [US5173414A](#) [0061]
- [WO9513365A](#) [0061]
- [US5658776A](#) [0061]
- [WO9513392A](#) [0061]
- [WO9617947A](#) [0061]
- [US9818600W](#) [0061]
- [WO9709441A](#) [0061]
- [US9614423W](#) [0061]
- [WO9708298A](#) [0061]
- [US9613872W](#) [0061]
- [WO9721825A](#) [0061]
- [US9620777W](#) [0061]
- [WO9706243A](#) [0061]
- [FR9601064W](#) [0061]
- [WO9911764A](#) [0061]
- [US5786211A](#) [0061]
- [US5871982A](#) [0061]
- [US6258595B](#) [0061]
- [US2012047999W](#) [0063]
- [WO2013016352A](#) [0063]
- [US6566118B](#) [0064]
- [WO9809657A](#) [0064]
- [WO02053703A](#) [0071] [0073]

#### Non-patent literature cited in the description

- **HOFFMAN et al.**Cell, 1987, vol. 51, 6919-28 [0004]
- **STRAUB et al.**Curr Opin. Neurol., 1997, vol. 10, 2168-75 [0004]
- **SACCO et al.**Cell, 2010, vol. 143, 71059-71 [0004]
- **WALLACE et al.**Annu Rev Physiol, 2009, vol. 71, 37-57 [0004]
- **DESGUERRE et al.**J Neuropathol Exp Neurol, 2009, vol. 68, 7762-7 [0005]
- **EISENBERG et al.**Proc Natl Acad Sci U S A, 2007, vol. 104, 4317016-21 [0005]
- **JIANG et al.**Proc Natl Acad Sci U S A, 2007, vol. 104, 4317016-21 [0005]
- **CACCHIARELLI et al.**Cell Metab, 2010, vol. 12, 4341-51 [0005] [0005]
- **EISENBERG et al.**Proc Natl Acad Sci U S A, 2007, vol. 104, 4317016-2 [0005]
- **SRIVASTAVA et al.**J Virol, 1983, vol. 45, 555-564 [0006]

- **RUFFING et al.**J Gen Virol, 1994, vol. 75, 3385-3392 [0006]
- **GAO et al.**J. Virol., 2004, vol. 78, 6381-6388 [0006]
- **Mol. Ther.**, 2006, vol. 13, 167-76 [0006]
- **Virology**, 2004, vol. 330, 2375-383 [0006]
- **RODINO-KLAPAC et al.**J. Trans. Med., 2007, vol. 5, 45- [0006]
- **MUZYCZKA**Current Topics in Microbiology and Immunology, 1992, vol. 158, 97-129 [0006]
- **CLARK et al.**Hum Gene Ther, 1997, vol. 8, 659-669 [0008]
- **KESSLER et al.**Proc Nat. Acad Sc. USA, 1996, vol. 93, 14082-14087 [0008]
- **XIAO et al.**J Virol, 1996, vol. 70, 8098-8108 [0008]
- **CHAO et al.**Mol Ther, 2000, vol. 2, 619-623 [0008]
- **CHAO et al.**Mol Ther, 2001, vol. 4, 217-222 [0008]
- **HERZOG et al.**Proc Natl Acad Sci USA, 1997, vol. 94, 5804-5809 [0008]
- **MURPHY et al.**Proc Natl Acad Sci USA, 1997, vol. 94, 13921-13926 [0008]
- **LEWIS et al.**J Virol, 2002, vol. 76, 8769-8775 [0008]
- **SAMBROOK et al.**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory19890000 [0030]
- **ANDERSON et al.**Nucleic Acid Hybridisation: A Practical ApproachIRL Press Limited [0031]
- **CARTER**Handbook of Parvoviruses19890000vol. 1, 169-228 [0055]
- **BERNS**VirologyRaven Press19900000 [0055]
- **BLACKLOWE**Parvoviruses and Human Disease19880000165-174 [0055]
- **ROSE**Comprehensive Virology, 1974, vol. 3, 1-61 [0055]
- **MARSIC et al.**Molecular Therapy, 2014, vol. 22, 111900-1909 [0058]
- **SAMULSKI et al.**Proc. Natl. Acad. S6. USA, 1982, vol. 79, 2077-2081 [0060]
- **LAUGHLIN et al.**Gene, 1983, vol. 23, 65-73 [0060]
- **SENAPATHY**CARTERJ. Biol. Chem., 1984, vol. 259, 4661-4666 [0060]
- **CARTER**Current Opinions in Biotechnology, 1992, 1533-539 [0061]
- **MUZYCZKA**Curr. Topics in Microbial. and Immunol., 1992, vol. 158, 97-129 [0061]
- **RATSCHIN et al.**Mol. Cell. Biol., 1984, vol. 4, 2072- [0061]
- **HERMONAT et al.**Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 6466- [0061]
- **TRATSCHIN et al.**Mo1. Cell. Biol., 1985, vol. 5, 3251- [0061]
- **MCLAUGHLIN et al.**J. Virol., 1988, vol. 62, 1963- [0061]
- **LEBKOWSKI et al.**Mol. Cell. Biol., 1988, vol. 7, 349- [0061]
- **SAMULSKI et al.**J. Virol., 1989, vol. 63, 3822-3828 [0061]
- **PERRIN et al.**Vaccine, 1995, vol. 13, 1244-1250 [0061]
- **PAUL et al.**Human Gene Therapy, 1993, vol. 4, 609-615 [0061]
- **CLARK et al.**Gene Therapy, 1996, vol. 3, 1124-1132 [0061]
- **CLARK et al.**Hum. Gene Ther., 1999, vol. 10, 61031-1039 [0064]
- **SCHENPPCLARK**Methods Mol. Med., 2002, vol. 69, 427-443 [0064]
- **CLARK et al.**Hum. Gene Ther., 1999, vol. 10, 1031-1039 [0072]
- **WEINTRAUB et al.**Science, 1991, vol. 251, 761-766 [0079]
- **CSERJESIOLSON**Mol Cell Biol, 1991, vol. 11, 4854-4862 [0079]
- **MUSCAT et al.**Mol Cell Biol, 1987, vol. 7, 4089-4099 [0079]

- **JOHNSON et al.** Mol Cell Biol, 1989, vol. 9, 3393-3399 [0079]
- **SEMENZA et al.** Proc Natl Acad Sci USA, 1991, vol. 88, 5680-5684 [0079]
- **MADERWHITE** Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 5603-5607 [0079]
- **NEVO et al.** PloS One, 2011, vol. 6, e18049- [0087]
- **HAKIM et al.** Methods Mol Biol., 2011, vol. 709, 75-89 [0091]
- **RODINO-KLAPAC et al.** Mol Ther., 2010, vol. 18, 1109-17 [0095]
- **MENDELL et al.** Ann. Neurol., 2012, vol. 71, 304-313 [0112]
- **ROOIJ et al.** Proc. Natl. Acad. Sci. USA, 2008, vol. 105, 13027-13032 [0113]
- **HOFFMAN, E.P.BROWN, R.H., JR.KUNKEL, L.M** Dystrophin: the protein product of the Duchenne muscular dystrophy locus Cell, 1987, vol. 51, 919-928 [0121]
- **STRAUB, V.CAMPBELL, K.P.** Muscular dystrophies and the dystrophin-glycoprotein complex Curr Opin Neurol, 1997, vol. 10, 168-175 [0121]
- **SACCO, A. et al.** Short telomeres and stem cell exhaustion model Duchenne muscular dystrophy in mdx/mTR mice Cell, 2010, vol. 143, 1059-1071 [0121]
- **WALLACE, G.Q.MCNALLY, E.M.** Mechanisms of muscle degeneration, regeneration, and repair in the muscular dystrophies Annu Rev Physiol, 2009, vol. 71, 37-57 [0121]
- **ZHOU, L.LU, H.** Targeting fibrosis in Duchenne muscular dystrophy J Neuropathol Exp Neurol, 2010, vol. 69, 771-776 [0121]
- **DESGUERRE, I. et al.** Endomysial fibrosis in Duchenne muscular dystrophy: a marker of poor outcome associated with macrophage alternative activation J Neuropathol Exp Neurol, 2009, vol. 68, 762-773 [0121]
- **KIM, J. et al.** microRNA-directed cleavage of ATHB 15 mRNA regulates vascular development in Arabidopsis inflorescence stems Plant J, 2005, vol. 42, 84-94 [0121]
- **AMBROS, V.** MicroRNA pathways in flies and worms: growth, death, fat, stress, and timing Cell, 2003, vol. 113, 673-676 [0121]
- **EISENBERG, I. et al.** Distinctive patterns of microRNA expression in primary muscular disorders Proc Natl Acad Sci U S A, 2007, vol. 104, 17016-17021 [0121]
- **JIANG, X.TSITSIOU, E.HERRICK, S.E.LINDSAY, M.A.** MicroRNAs and the regulation of fibrosis FEBS J, 2010, vol. 277, 2015-2021 [0121]
- **VAN ROOIJ, E. et al.** Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis Proc Natl Acad Sci U S A, 2008, vol. 105, 13027-13032 [0121]
- **CACCHIARELLI, D. et al.** MicroRNAs involved in molecular circuitries relevant for the Duchenne muscular dystrophy pathogenesis are controlled by the dystrophin/nNOS pathway Cell Metab, 2010, vol. 12, 341-351 [0121]
- **DIPRIMIO, N.MCPHEE, S.W.SAMULSKI, R.J** Adeno-associated virus for the treatment of muscle diseases: toward clinical trials Curr Opin Mol Ther, 2010, vol. 12, 553-560 [0121]
- **MENDELL, J.R. et al.** Sustained alpha-sarcoglycan gene expression after gene transfer in limb-girdle muscular dystrophy, type 2D Ann Neurol, 2010, vol. 68, 629-638 [0121]
- **MENDELL, J.R. et al.** Limb-girdle muscular dystrophy type 2D gene therapy restores alpha-sarcoglycan and associated proteins Ann Neurol, 2009, vol. 66, 290-297 [0121]
- **MENDELL, J.R. et al.** A phase 1/2a follistatin gene therapy trial for becker muscular dystrophy Molecular therapy : the journal of the American Society of Gene Therapy, 2015, vol. 23, 192-201 [0121]

- **CARNWATH, J.W.SHOTTON, D.M.**Muscular dystrophy in the mdx mouse: histopathology of the soleus and extensor digitorum longus muscles*J Neural Sci*, 1987, vol. 80, 39-54 [\[0121\]](#)
- **COULTON, G.R.MORGAN, J.E.PARTRIDGE, T.A.SLOPER, J.C**The mdx mouse skeletal muscle myopathy: I. A histological, morphometric and biochemical investigation*Neuropathol Appl Neurobiol*, 1988, vol. 14, 53-70 [\[0121\]](#)
- **CULLEN, M.J.JAROS, E.**Ultrastructure of the skeletal muscle in the X chromosome-linked dystrophic (mdx) mouse. Comparison with Duchenne muscular dystrophy*Acta Neuropathol*, 1988, vol. 77, 69-81 [\[0121\]](#)
- **DUPONT-VERSTEEGDEN, E.E.MCCARTER, R.J.**Differential expression of muscular dystrophy in diaphragm versus hindlimb muscles of mdx mice*Muscle Nerve*, 1992, vol. 15, 1105-1110 [\[0121\]](#)
- **STEDMAN, H.H. et al.**The mdx mouse diaphragm reproduces the degenerative changes of Duchenne muscular dystrophy*Nature*, 1991, vol. 352, 536-539 [\[0121\]](#)
- **DECONINCK, A.E. et al.**Utrophin-dystrophin-deficient mice as a model for Duchenne muscular dystrophy*Cell*, 1997, vol. 90, 717-727 [\[0121\]](#)
- **GRADY, R.M. et al.**Skeletal and cardiac myopathies in mice lacking utrophin and dystrophin: a model for Duchenne muscular dystrophy*Cell*, 1997, vol. 90, 729-738 [\[0121\]](#)
- **LOVE, D.R. et al.**An autosomal transcript in skeletal muscle with homology to dystrophin*Nature*, 1989, vol. 339, 55-58 [\[0121\]](#)
- **TINSLEY, J.M. et al.**Primary structure of dystrophin-related protein*Nature*, 1992, vol. 360, 591-593 [\[0121\]](#)
- **TINSLEY, J. et al.**Expression of full-length utrophin prevents muscular dystrophy in mdx mice*Nat Med*, 1998, vol. 4, 1441-1444 [\[0121\]](#)
- **SQUIRE, S. et al.**Prevention of pathology in mdx mice by expression of utrophin: analysis using an inducible transgenic expression system*Hum Mol Genet*, 2002, vol. 11, 3333-3344 [\[0121\]](#)
- **RAFAEL, J.A.TINSLEY, J.M.POTTER, A.C.DECONINCK, A.E.DAVIES, K.E**Skeletal muscle-specific expression of a utrophin transgene rescues utrophin-dystrophin deficient mice*Nat Genet*, 1998, vol. 19, 79-82 [\[0121\]](#)
- **ZHOU, L. et al.**Haploinsufficiency of utrophin gene worsens skeletal muscle inflammation and fibrosis in mdx mice*J Neural Sci*, 2008, vol. 264, 106-111 [\[0121\]](#)
- **GUTPELL, K.M.HRINIVICH, W.T.HOFFMAN, L.M.**Skeletal Muscle Fibrosis in the mdx/utrn+/- Mouse Validates Its Suitability as a Murine Model of Duchenne Muscular Dystrophy*PloS one*, 2015, vol. 10, e0117306- [\[0121\]](#)
- **RODINO-KLAPAC, L.R. et al.**Micro-dystrophin and follistatin co-delivery restores muscle function in aged DMD model*Human molecular genetics*, 2013, vol. 22, 4929-4937 [\[0121\]](#)
- **CUSHING, L. et al.**MIR-29 is a Major Regulator of Genes Associated with Pulmonary Fibrosis*Am J Respir Cell Mol Biol*, 2010, [\[0121\]](#)
- **RODERBURG, C. et al.**Micro-RNA profiling reveals a role for miR-29 in human and murine liver fibrosis*Hepatology*, 2011, vol. 53, 209-218 [\[0121\]](#)
- **NEVO, Y. et al.**The Ras antagonist, farnesylthiosalicylic acid (FTS), decreases fibrosis

and improves muscle strength in dy/dy mouse model of muscular dystrophyPloS one, 2011, vol. 6, e18049- [0121]

- **RODINO-KLAPAC, L.R. et al.**A translational approach for limb vascular delivery of the micro-dystrophin gene without high volume or high pressure for treatment of Duchenne muscular dystrophyJ Transl Med, 2007, vol. 5, 45- [0121]
- **MULIERI, L.A.HASENFUSS, G.ITTLEMAN, F.BLANCHARD, E.M.ALPERT, N.R.**Protection of human left ventricular myocardium from cutting injury with 2,3-butanedione monoximeCirc Res, 1989, vol. 65, 1441-1449 [0121]
- **RODINO-KLAPAC, L.R. et al.**Persistent expression of FLAG-tagged micro dystrophin in nonhuman primates following intramuscular and vascular deliveryMolecular therapy : the journal of the American Society of Gene Therapy, 2010, vol. 18, 109-117 [0121]
- **GROSE, W.E. et al.**Homologous recombination mediates functional recovery of dysferlin deficiency following AAV5 gene transferPloS one, 2012, vol. 7, e39233- [0121]
- **LIU, M. et al.**Adeno-associated virus-mediated microdystrophin expression protects young mdx muscle from contraction-induced injuryMol Ther, 2005, vol. 11, 245-256 [0121]



## Patentkrav

1. Rekombinant AAVrh74-vektor, der omfatter et muskelspecifikt kontrolelement, som er funktionsmæssigt forbundet med nukleotidsekvensen ifølge SEQ ID NO: 7.  
5
2. Rekombinant AAVrh74-vektor ifølge krav 1, hvor det muskelspecifikke kontrolelement er afledt af muskelcreatinkinase (MCK).  
10
3. Rekombinant AAVrh74-vektor ifølge krav 2, hvor det muskelspecifikke kontrolelement omfatter nukleotidsekvensen ifølge SEQ ID NO: 10 eller SEQ ID NO: 11.
- 15 4. Sammensætning, der omfatter den rekombinante AAVrh74-vektor ifølge et hvilket som helst af kravene 1-3 og en farmaceutisk acceptabel bærer.
5. Rekombinant AAVrh74-vektor ifølge et hvilket som helst af  
20 kravene 1-3 eller sammensætning ifølge krav 4 til anvendelse ved behandling af muskeldystrofi.
6. Rekombinant AAVrh74-vektor ifølge et hvilket som helst af kravene 1-3 eller sammensætning ifølge krav 4 til anvendelse ved  
25 reducering eller forebyggelse af fibrose hos et individ, der lider af muskeldystrofi.
7. Rekombinant AAVrh74-vektor ifølge et hvilket som helst af kravene 1-3 eller sammensætning ifølge krav 4 til anvendelse til  
30 forøgelse af muskelkraft eller muskelmasse hos et individ, der lider af muskeldystrofi.
8. Rekombinant AAVrh74-vektor eller sammensætning til anvendelse ifølge et hvilket som helst af kravene 5-7, hvor den  
35 rekombinante AAVrh74-vektor eller sammensætningen indgives, inden fibrose observeres hos individet, eller inden muskelkraften reduceres hos individet, eller inden muskelmassen reduceres hos individet.

9. Rekombinant AAVrh74-vektor eller sammensætning til anvendelse ifølge et hvilket som helst af kravene 5-8, hvor individet lider af Duchennes muskeldystrofi.

5

10. In vitro-fremgangsmåde til fremstilling af et funktionelt mikrodystrofinprotein, hvilken fremgangsmåde omfatter inficering af en værtscelle med en rekombinant AAVrh74-vektor ifølge et hvilket som helst af kravene 1-3 og ekspresion af et

10

funktionelt mikrodystrofinprotein i værtscellen.



Figure 2A




Figure 2B  
Sirius RedFigure 2C  
miR-29c

Figure 3A




Figure 3B




Figure 3C




Figure 4C




Figure 4B




Figure 4A




Figure 5A




Figure 5B

Sirius Red




Figure 5C

miR-29c




Figure 6A

Col1A




Figure 6B

Col3A




Figure 6C

FBN




Figure 6D


TGF- $\beta$ 



Figure 7A




Figure 7B




Figure 7C




Figure 8




Figure 9A

Gas Weight




Figure 9B

Fiber Diameters




Figure 9C

Histogram of all diameters




Figure 9D

Total Area




Figure 9E

total number of fibers




Figure 9F

Muscle Fibers/mm²




Figure 10A




Figure 10D




Figure 10G




Figure 10C




Figure 10F




Figure 10B




Figure 10E




Figure 11




Figure 12




Figure 13A







Figure 13D




Figure 13G




Figure 13C




Figure 13F




Figure 13B




Figure 13E




Figure 14A






Figure 14D

*Col1A*




Figure 14C

miR-29c levels




Figure 14B

Sirius Red




Figure 14G

*Tgfb1*




Figure 14F

*Fbn*




Figure 14E

*Col3A*




Figure 15A




Figure 15B




Figure 15C




Figure 16A




Figure 16B




Figure 16C




Figure 17A




Figure 17B




Figure 17C




Figure 17D




Figure 18A

pAAV.CMV.miR29C Sequence (SEQ ID NO: 1)

Main features:

CMV promoter- 120-526

EF1a Intron- 927-1087, 1380-1854

miR-29c-1257-1284

shRNA-miR29-c with primary seed sequence- 1088-1375

PolyA- 1896-2091

CAGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCGGGCGTCGGGCGACCTTTGGTCGCCCG  
GCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGGGTTAACTCGTTACATAACTTACGGTAAATGGCCCG  
CCTGGCTGACCGCCCAACGACCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGA  
CTTTCCATTGACGTCAATGGGTGGAGTATTACGGTAACTGCCACCTTGGCAGTACATCAAGTGTATCATATGCC  
AAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTATGGGAC  
TTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGG  
GCGTGGATAGCGGTTTGACTCACGGGGATTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGTGTTTGCAC  
CAAAATCAACGGGACTTTCCAAATGTCGTAACAACCTCCGCCCATTGACGCAATGGGCGGTAGGCGTGTACGG  
TGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCTGGAGACGCCATCCACGCTGTTTTGACC  
TCCATAGAAGACACCGGGACCGATCCAGCCTCCGGACTCTAGAGGATCCGGTACTCGAGGAAGTGAACCAACAGA  
AAGTTAACTGGTAAGTTTAGTCTTTTGTCTTTATTTCAGTCCCGGATCCGGTGGTGGTGCAATCAAAGAAGT  
CTCCTCAGTGGATGTTGCCTTTACTTCTAGGCTGTACGGAAGTGTTACTTCTGCTCTAAAAGCTGCGGAATTGTAC  
CCGGGGCCGATCCACCGGCTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAGTGGCGTGTGTGGTTCCCG  
CGGGCGGCGACGGGGCCCGTGGCTCCAGCGCACATGTTGGGCGAGGCGGGGCTGCGAGCGCGGCCACCGAG  
AATCGGACGGGGGTAGTCTCAAGCTGGCCGGCCTGTTGAATGAGGCTTCACTTTACAGAATCGTTGCCTGC  
ACATCTTGGAAACACTTGTGGGATTACTTCTTCAGGTTAACCCACAGAAAGGCTCGAGAAGGTATATTGCTGTTG  
ACAGTGAGCGCAACCGATTTCAAATGGTGCTAGAGTGAAGCCACAGATGTCTAGCACCATTTGAAATCGGTTATG  
CCTACTGCCTCGGAATTCAGGGGCTACTTTAGGAGCAATTATCTTGTTTACTAAAACTGAATACCTTGCTATCTCTT  
TGATACATTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCCGTGTATCGCCCCGCCCTGGGCGGCAAGGCTG  
GCCCCGTGGGACCAAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAG  
GACGCGGCGCTCGGGAGAGCGGGCGGGTGAGTACCCACACAAAGGAAAAGGGCTTTCCGCTCTCAGCCGTCG  
CTTCATGTGACTCCACGGAGTACCGGGCGCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTC  
TTTAGGTTGGGGGGAGGGGTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGAAGTTAGGCCAG  
CTTGGCACTTGATGTAATTCCTTGGAATTTGCCCTTTTGAAGTTTGGATCTTGGTTTATTCTCAAGCCTCAGACAG  
TGGTTCAAAGTTTTTTCTTCCATTTAGGTTGTCGTGAAAAGCTAGCGCTACCGGACTCAGATCTCGAGCTCAAGCT  
GCGGGGATCCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGAAAAAATGCT  
TTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAGTTAACAACAACAATT  
GCATTCATTTTATGTTTCAAGTTCAAGGGGAGGTTGGGAGGTTTTTCACTAGTAGCATGGCTACGTAGATAAGT  
AGCATGGCGGGTTAATCATTAACTACAAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTC  
GCTCACTGAGGCCGGGCGACCAAGGTCGCCGACGCCGGGCTTGGCCGGGCGGCTCAGTGAGCGAGCGAG  
CGCGCCAGCTGGCGTAATAGCGAAGAGGCGCCGACCGATCGCCCTCCCAACAGTTGCGCAGCCTGAATGGCGAA  
TGGAATTCAGACGATTGAGCGTCAAAATGTAGGTATTTCCATGAGCGTTTTCTGTTGCAATGGCTGGCGGTAA  
TATTGTTCTGGATATTACCAGCAAGGCCGATAGTTTGAAGTTCTTCTACTCAGGCAAGTGATGTTATTACTAATCAA  
GAAGTATTGCGACAACGGTTAATTTGCGTGATGGACAGACTCTTTACTCGGTGGCCTCACTGATTATAAAAAACAC  
TTCTCAGGATTCTGGCGTACCGTTCTGTCTAAAATCCCTTAATCGGCCTCCTGTTAGCTCCGCTCTGATTCTAA  
CGAGGAAAGCACGTTATACGTGCTCGTCAAAGCAAC

Figure 18B

CATAGTACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTG  
 CCAGCGCCCTAGCGCCCGCTCCTTTGCTTTCTTCCCTTCTTTCTCGCCACGTTGCGCGGCTTTCCCGTCAAGCTCT  
 AAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAACTTGATTAGGGTGAT  
 GGTTACGTAAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTG  
 GACTCTTGTTCCAACTGGAACAACACTCAACCCTATCTCGGTCTATTCTTTGATTATAAGGGATTTTGCCGATT  
 CGGCCTATTGGTTAAAAATGAGCTGATTTAACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTACAATT  
 TAAATATTTGCTTATACAATCTTCCTGTTTTGGGGCTTTCTGATTATCAACCGGGGTACATATGATTGACATGCTA  
 GTTTTACGATTACCGTTCATCGATTCTCTGTTTGCTCCAGACTCTCAGGCAATGACCTGATAGCCTTTGTAGAGACC  
 TCTCAAAAATAGCTACCTCTCCGGCATGAATTTATCAGCTAGAACGGTTGAATATCATATTGATGGTGATTGACT  
 GTCTCCGGCCTTTCTACCCGTTTGAATCTTTACCTACACATTACTCAGGCATTGCATTTAAATATATGAGGGTCT  
 AAAAAATTTTATCCTTGCGTTGAAATAAGGGCTTCTCCCGCAAAAGTATTACAGGGTCATAATGTTTTGGTACAAC  
 CGATTTAGCTTTATGCTCTGAGGCTTTATTGCTTAATTTTGCTAATCTTTGCCTTGCCTGTATGATTTATTGGATGTT  
 GGAATTCCTGATGCGGTATTTTCTCTTACGCATCTGTGCGGTATTTACACCGCATATGGTGCACCTCAGTACAA  
 TCTGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGCT  
 GCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCAGAGGTTTTACCGTCATCA  
 CCGAAACGCGCGAGACGAAAGGGCCTCGTGATACGCCTATTTTATAGGTTAATGTCATGATAATAATGGTTTCTT  
 AGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTCTAAATACATTCAAATATG  
 TATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTATCAACATT  
 TCCGTGTCGCCCTTATTCCCTTTTTGCGGCATTTTGCTTCTGTTTTGCTCACCCAGAAACGCTGGTGAAGTAA  
 AAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGA  
 GTTTTCGCCCGAAGAAGCTTTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATT  
 GACGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACA  
 GAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAACACTGCG  
 GCCAACTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGACAACATGGGGGATCATGTA  
 ACTCGCCTTGATCGTTGGGAACCGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGTA  
 GCAATGGCAACAACGTTGCGCAAACTATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACT  
 GGATGGAGGCGGATAAAGTTGACGAGCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGTTTATTGCTGATAAAT  
 CTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAG  
 TTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGA  
 TTAAGCATTGGTAAGTGTGACACCAAGTTTACTCATATATACTTTAGATTGATTTAAACTTCATTTTAAATTTAAAA  
 GGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCA  
 GACCCCGTAGAAAAGATCAAAGGATCTTCTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAACAAAAAA  
 ACCACCGCTACCAGCGGTGGTTTGTGTCGGGATCAAGAGCTACCAACTCTTTTCCGAAGGTAAGTGGCTTCAGC  
 AGAGCGCAGATACCAAACTACTGTCCTTCTAGTGAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGC  
 CTACATACCTCGCTCTGCTAATCCTGTTACCACTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGA  
 CTAAGACGATAGTTACCGGATAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTG  
 AGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGA  
 AAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAAC  
 GCCTGGTATCTTTATAGTCTGTGCGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTGTCAGGGGG  
 GCGGAGCCTATGGAAAACGCCAGCAACGCGGCCCTTTTACGGTTCCTGGCCTTTTGTGGCCTTTTGTGCACATG  
 TTCTTCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCGCTCGCCGAG  
 CCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCG  
 CGCGTTGGCCGATTCAATG



Figure 19




Figure 20A

SEQ ID NO: 9Main features:MCK promoterChimeric intron sequenceHuman micro-dystrophin sequencePoly A tailAmpicillin resistancepGEX plasmid backbone with pBR322 origin or replication

GCCCAATACGCAAACCGCCTCTCCCCGCGCGTTGGCCGATTCATTAATGCAGCTGGC  
 GCGCTCGCTCGCTCACTGAGGCCGCCGGGCAAAGCCCCGGGCGTCGGGCGACCTTT  
 GGTCGCCCCGGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCAT  
 CACTAGGGGTTCTTGTAGTTAATGATTAACCCGCCATGCTAATTATCTACGTAGCC  
 ATGTCTAGACAGCCACTATGGGTCTAGGCTGCCCATGTAAGGAGGCAAGGCCTGGG  
 GACACCCGAGATGCCTGGTTATAATTAACCCAGACATGTGGCTGCTCCCCCCCCCA  
 ACACCTGCTGCCTGAGCCTCACCCCCACCCCGGTGCCTGGGICTTAGGCTCTGTACA  
 CCATGGAGGAGAAGCTCGCTCTAAAAATAACCTGTCCCTGGTGGGCTGTGGGGGA  
 CTGAGGGCAGGCTGTAACAGGCTTGGGGGCCAGGGCTTATACGTGCCTGGGACTCC  
 CAAAGTATTACTGTTCCATGTTCCCGGCGAAGGGCCAGCTGTCCCCCGCCAGCTAGA  
 CTCAGCACTTAGTTTAGGAACCAAGTGAGCAAGTCAGCCCTTGGGGCAGCCATACA  
 AGGCCATGGGGCTGGGCAAGCTGCACGCCTGGGTCCGGGGTGGGCACGGTGCCCGG  
 GCAACGAGCTGAAAGCTCATCTGCTCTCAGGGGGCCCTCCCTGGGGACAGCCCCTCC  
 TGGCTAGTCACACCCTGTAGGCTCCTCTATATAACCCAGGGGCACAGGGGCTGCCCC  
 CGGGTCACCACCACCTCCACAGCACAGACAGACACTCAGGAGCCAGCCAGCCAGGT  
 AAGTTTAGTCTTTTTGTCTTTTATTTACAGGTCCCGGATCCGGTGGTGGTGCAAATCAA  
 AGAACTGCTCCTCAGTGGATGTTGCCTTTACTTCTAGGCCTGTACGGAAGTGTTACTT  
 CTGCTCTAAAAGCTGCGGAATTGTACCCGCGGCCGCCACCATGCTGTGGTGGGAGG  
 AGGTGGAGGATTGTTATGAAAGGGAGGACGTGCAGAAGAAGACTTTTACCAAGTGG  
 GTGAACGCTCAGTTCAGCAAATTTGGGAAGCAGCACATCGAGAATCTGTTTTCCGAC  
 CTGCAGGATGGGAGACGGCTGCTGGATCTGCTGGAAGGACTGACTGGCCAGAAGCT  
 GCCCAAAGAGAAGGGGAGCACTAGGGTGCACGCCCTGAACAACGTGAACAAAGCT  
 CTGAGAGTGCTGCAGAACAACAACGTGGATCTGGTGAATATTGGCAGTACTGATAT  
 CGTGGACGGGAACCACAACTGACACTGGGCCTGATCTGGAACATTATTCTGCACTG  
 GCAGGTGAAAAATGTGATGAAGAACATCATGGCCGGGCTGCAGCAGACCAATTCCG  
 AGAAGATCCTGCTGTCTTGGGTGCGGCAGAGCACCCGCAACTATCCCCAGGTGAAC  
 GTGATTAACCTCACTACATCCTGGAGCGACGGGCTGGCCCTGAATGCTCTGATTCAC  
 AGCCACAGGCCTGATCTGTTGACTGGAATAGCGTGGTGTGCCAGCAGTCTGCCACA  
 CAGCGCCTGGAACATGCCTTCAATATCGCTCGGTACCAGCTGGGGATCGAAAAACT  
 GCTGGACCCAGAGGATGTGGACACTACATACCCAGATAAAAAGTCTATTCTGATGT  
 ACATTACTAGCCTGTTCCAGGTGCTGCCACAGCAGGTGTCTATTGAAGCCATTACAG  
 AGGTGGAAATGCTGCCCCGCCCCCCCCAAAGTGACTAAAGAGGAGCATTTTCAGCTG  
 CATCATCAGATGCATTACAGCCAGCAGATTACCGTGAGCCTGGCTCAGGGATATGA  
 GCGCACCAAGTAGTCCAAAACACGGTTCAAGTCCTACGCTTATAC

Figure 20B

CCAGGCTGCCTACGTGACAACTAGCGACCCTACTAGATCCCCCTTCCATCCCAGCA  
 CCTGGAGGCCCCAGAGGACAAGAGCTTTGGGTCCAGCCTGATGG  
 AAAGCGAGGTGAATCTGGATCGGTACCAGACAGCCCTGGAGGAGGTGCTGAGCTGG  
 CTGCTGAGTGCTGAAGACACACTGCAGGCCAGGGCGAAATTTCCAATGACGTGGA  
 AGTGGTGAAGGATCAGTTCCACACACACGAGGGCTATATGATGGACCTGACAGCTC  
 ACCAGGGGCGCGTGGGCAATATCCTGCAGCTGGGCTCTAAACTGATCGGCACCGGG  
 AAAGTGAAGTGAAGGACGAGGAAACAGAAGTGCAGGAGCAGATGAACCTGCTGAACA  
 GCGCTGGGAGTGTCTGAGAGTGGCTAGTATGGAGAAGCAGTCCAACCTGCACCGG  
 GTGCTGATGGACCTGCAGAACCAGAACTGAAAGAGCTGAACGACTGGCTGACAAA  
 GACTGAGGAACGCACAAGGAAGATGGAGGAGGAGCCACTGGGACCCGACCTGGAG  
 GATCTGAAGAGACAGGTGCAGCAGCATAAGGTGCTGCAGGAGGATCTGGAACAGG  
 AGCAGGTGCGGGTGAAGTCCCTGACACATATGGTGGTGGTGGTGGACGAATCTAGT  
 GGAGATCACGCCACCGCCGCCCTGGAGGAACAGCTGAAGGTGCTGGGGGACCGGTG  
 GCGCAACATTTGCCGGTGGACCGAGGACAGGTGGGTGCTGCTGCAGGACATCCTGC  
 TGAAATGGCAGAGGCTGACCGAGGAGCAGTGTCTGTTLAGTGTCTGGCTGAGCGAG  
 AAAGAGGACGCGCTGAACAAGATCCACACAACCGGCTTTAAGGATCAGAACGAAT  
 GCTGTCTAGCCTGCAGAACTGGCTGTGCTGAAGGCCGATCTGGAGAAAAAGAAGC  
 AGAGCATGGGCAAACCTGTATAGCCTGAAACAGGACCTGCTGAGCACCTGAAGAAC  
 AAGAGCGTGACCCAGAAGACAGAAGCCTGGCTGGATAACTTTGCCCGCTGCTGGGA  
 CAACCTGGTGCAGAACTGGAGAAAAGTACAGCTCAGATCTCTCAGGCTGTGACCA  
 CAACCCAGCCTAGCCTGACCCAGACAACCGTGATGGAAACCGTGACCACCGTGACA  
 ACCCGCGAACAGATCCTGGTGAACATGCCAGGAAGAGCTGCCACCTCCACCTCC  
 CCAGAAGAAGAGAACCCTGGAGCGGCTGCAGGAGCTGCAGGAAGCCACTGACGAA  
 CTGGACCTGAAGCTGAGGCAGGCCGAAGTGATTAAGGGGTCTTGGCAGCCTGTGGG  
 CGATCTGCTGATTGATTCCCTGCAGGACCACCTGGAAAAGGTGAAGGCTCTGAGAG  
 GCGAAATTGCTCCACTGAAGGAGAACGTGAGTCATGTGAACGATCTGGCTAGACAG  
 CTGACAACACTGGGCATCCAGCTGAGCCCATAAATCTGAGCACACTGGAGGACCT  
 GAATACCAGGTGGAAGCTGCTGCAGGTGGCTGTGGAAAGACCGGGTGGCGCAGCTGC  
 ATGAGGCCCATCGCGACTTCGGACCAGCCAGCCAGCACTTTCTGAGCACATCCGTGC  
 AGGGGCCCTGGGAGAGGGGCCATTTCTCCCAACAAGGTGCCCTACTATATTAATCACG  
 AGACCCAGACCACTTGTTGGGACCATCCCAAGATGACAGAACTGTACCAGTCCCTG  
 GCCGATCTGAACAACGTGAGGTTTACGCGCTTACAGAACCGCTATGAAGCTGAGACG  
 GCTGCAGAAGGCCCTGTGCCTGGATCTGCTGTCCCTGTCCGCCGCTGCGATGCCCT  
 GGATCAGCATAATCTGAAGCAGAACGATCAGCCAATGGATATCCTGCAGATCATCA  
 ACTGCCTGACCACTATCTACGACAGGCTGGAGCAGGAGCACACAACCTGGTGAAC  
 GTGCCCTCTGTCCGTGGATATGTGCCTGAAGTGGCTGCTGAACGTGTATGACACTGGG  
 CGCACCGGCCCGGATCAGAGTGCTGAGTTTTAAAGTGGGATTATCTCCCTGTGTAAG  
 GCGCACCTGGAGGACAAGTACAGGTACCTGTTCAAGCAGGTGGCTAGTAGCACTGG  
 ATTTTGTGACCAGCGCCGCTGGGACTGCTGCTGCATGATAGTATCCAGATTCCCTAG  
 ACAGCTGGGAGAGGTGGCTAGTTTCGGAGGATCTAACATCGAACCCAGCGTGCGCA  
 GCTGTTTCCAGTTTGCCAATAACAAACCTGAAATCGAGGCTGCTCTGTTCCTGGATT  
 GGATGCGCCTGGAACACAGAGCATGGTGTGGCTGCCTGTGCTGCACAGAGTGG

Figure 20C

CTGCCGCCGAAACTGCCAAGCACCAGGCTAAATGCAACATCTGCAAGGAATGTCCC  
ATTATCGGCTTTTCGCTACAGGAGTCTGAAACATTTTAACTACGAT  
ATTTGCCAGAGCTGCTTCTTTTCCGGAAGAGTGGCCAAAGGACACAAGATGCACTAC  
CCTATGGTGGAATATTGCACCCCAACTACATCTGGCGAAGATGTGCGCGATTTTGGC  
AAGGTGCTGAAGAATAAGTTTCCGACTAAGAGGTACTTCGCCAAGCACCCCCGCAT  
GGGGTATCTGCCAGTGCAGACAGTGTCTGGAAGGAGACAATATGGAGACCGATACAA  
TGTGAGCGGGCGCAATAAAAGATCTTTATTTTCATTAGATCTGTGTGTTGGTTTTTG  
TGTGTCTAGAGCATGGCTACGTAGATAAGTAGCATGGCGGGTTAATCATTAACTACA  
AGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTCACTG  
AGGCCGGGCGACCAAAGGTGCCCCGACGCCCGGGCTTTGCCCGGGCGGCCTCAGTG  
AGCGAGCGAGCGCGCCAGCTGGCGTAATAGCGAAGAGGCCCCGCACCGATCGCCCTT  
CCCAACAGTTGCGCAGCCTGAATGGCGAATGGAAGTTCCAGACGATTGAGCGTCAA  
AATGTAGGTATTTCCATGAGCGTTTTTCTGTGCAATGGCTGGCGGTAATATTGTTT  
TGGATATTACCAGCAAGGCCGATAGTTTGAGTTCTTCTACTCAGGCAAGTGATGTTA  
TTACTAATCAAAGAAGTATTGCGACAACGGTTAATTTGCGTGATGGACAGACTCTTT  
TACTCGGTGGCTCACTGATTATAAAAAACACTTCTCAGGATTCTGGCGTACCGTTTCT  
GTCTAAAATCCCTTTAATCGGCCTCCTGTTTAGCTCCCGCTCTGATTCTAACGAGGAA  
AGCACGTTATACGTGCTCGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGCGCATT  
AAGCGCGGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCC  
TAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCCCTTCTCGCCACGTTCGCCGGCTTTCCC  
CGTCAAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGATTTACGGCAC  
CTCGACCCCAAAAAAAGTTGATTAGGGTGATGGTTACAGTAGTGGGCCATCGCCCTGA  
TAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGAAGTCTGT  
TCCAAACTGGAACAACACTCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGAT  
TTTGCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGC  
GAATTTTAACAAAATATTAACGTTTACAATTTAAATATTTGCTTATACAATCTTCCTG  
TTTTTGGGGCTTTTCTGATTATCAACCGGGGTACATATGATTGACATGCTAGTTTTAC  
GATTACCGTTCATCGATTCTCTTGTGTTTGCTCCAGACTCTCAGGCAATGACCTGATAGC  
CTTTGTAGAGACCTCTCAAAAATAGCTACCCTCTCCGGCATGAATTTATCAGCTAGA  
ACGGTTGAATATCATATTGATGGTGATTGACTGTCTCCGGCCTTTCTCACCCGTTTG  
AATCTTTACCTACACATTACTCAGGCATTGCATTTAAAATATATGAGGGTTCTAAAA  
ATTTTTATCCTTGCGTTGAAATAAAGGCTTCTCCCGCAAAGTATTACAGGGTCATA  
ATGTTTTTGGTACAACCGATTTAGCTTTATGCTCTGAGGCTTTATTGCTTAATTTTGCT  
AATTCTTTGCCTTGCCGTGATGATTTATTGGATGTTGGAAGTTCCTGATGCGGTATTT  
TCTCCTTACGCATCTGTGCGGTATTTACACCCGCATATGGTGCACTCTCAGTACAATC  
TGCTCTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCG  
CCCTGACGGGCTTGCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCC  
GGGAGCTGCATGTGTCAGAGGTTTTACCGGTCATCACCGAAACGCGCGAGACGAAA  
GGGCCTCGTGATACGCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTAG  
ACGTCAGGTGGCACTTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCT  
AAATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAAT  
AATATTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTT

Figure 20D

ATTCCCTTTTTTGCGGCATTTTTGCCTTCCTGTTTTTTGCTCACCCAGAAACGCTGGTGA  
AAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTAC  
ATCGAACTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGT  
TTTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTG  
ACGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTG  
AGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTA  
TGCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACG  
ATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAAC  
TCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTG  
ACACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACTATTAACTGGCGAA  
CTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAAAGTT  
GCAGGACCACCTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCT  
GGAGCCGGTGAGCGTGCGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAA  
GCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACG  
AAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAG  
ACCAAGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAG  
GATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTT  
TTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCC  
TTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGT  
GGTTTGTTGCCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTAACTGGCTTCAG  
CAGAGCGCAGATACCAAATACTGTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTT  
CAAGAACTCTGTAGCACCGCGTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGC  
TGCTGCCAGTGGCGATAAGTCGTGCTTACCGGGTTGGACTCAAGACGATAGTTACC  
GGATAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCGTGACACAGCCCAGCTTG  
AGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCC  
ACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAA  
CAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCT  
GTCCGGTTTTCGCCACCTCTGACTTGAGCGTTCGATTTTTGTGATGCTCGTCAGGGGGG  
CGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGC  
TGGCCTTTTGCTCATATGTTCTTTTCTGCGTTATCCCCTGATTCTGTGGATAACCGTA  
TTACCGGGTTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGC  
GAGTCAGTGAGCGACCAAGCGGAAGAGC

Figure 21A

SEQ ID NO: 7 : human micro-dystrophin gene

ATGCTGTGGTGGGAGGAGGTGGAGGATTGTTATGAAAGGGAGGACGTGCAGAAGA  
AGACTTTTACCAAGTGGGTGAACGCTCAGTTCAGCAAATTTGGGAAGCAGCACATC  
GAGAATCTGTTTTCCGACCTGCAGGATGGGAGACGGCTGCTGGATCTGCTGGAAGG  
ACTGACTGGCCAGAAGCTGCCCCAAAGAGAAGGGGAGCACTAGGGTGCACGCCCTGA  
ACAACGTGAACAAAGCTCTGAGAGTGTGCAGAACAACAACGTGGATCTGGTGAAT  
ATTGGCAGTACTGATATCGTGGACGGGAACCACAACTGACACTGGGCCTGATCTG  
GAACATTATTCTGCACTGGCAGGTGAAAAATGTGATGAAGAACATCATGGCCGGGC  
TGCAGCAGACCAATTCCGAGAAGATCCTGCTGTCTTGGGTGCGGCAGAGCACCCGC  
AACTATCCCCAGGTGAACGTGATTAACCTTCACTACATCCTGGAGCGACGGGCTGGCC  
CTGAATGCTCTGATTCACAGCCACAGGCCTGATCTGTTCGACTGGAATAGCGTGGTG  
TGCCAGCAGTCTGCCACACAGCGCCTGGAACATGCCTTCAATATCGCTCGGTACCAG  
CTGGGGATCGAAAACTGCTGGACCCAGAGGATGTGGACACTACATACCCAGATAA  
AAAGTCTATTCTGATGTACATTACTAGCCTGTTCCAGGTGCTGCCACAGCAGGTGTC  
TATTGAAGCCATTCAGGAGGTGGAAATGCTGCCCCGCCCCCCCCAAAGTGAATAAG  
AGGAGCATTTTCAGCTGCATCATCAGATGCATTACAGCCAGCAGATTACCGTGAGCC  
TGGCTCAGGGATATGAGCGCACCAAGTAGTCCAAAACACGGTTCAAGTCTACGCTT  
ATACCCAGGCTGCCTACGTGACAACCTAGCGACCCTACTAGATCCCCCTTTCCATCCC  
AGCACCTGGAGGGCCCCAGAGGACAAGAGCTTTGGGTCCAGCCTGATGGAAAGCGAG  
GTGAATCTGGATCGGTACCAGACAGCCCTGGAGGAGGTGCTGAGCTGGCTGCTGAG  
TGCTGAAGACACACTGCAGGCCCAGGGCGAAATTTCCAATGACGTGGAAGTGGTGA  
AGGATCAGTTCCACACACACGAGGGCTATATGATGGACCTGACAGCTCACCAGGGG  
CGCGTGGGCAATATCCTGCAGCTGGGCTCTAAACTGATCGGCACCGGGAAACTGAG  
TGAGGACGAGGAAACAGAAGTGCAGGAGCAGATGAACCTGCTGAACAGCCGCTGG  
GAGTGTCTGAGAGTGGCTAGTATGGAGAAGCAGTCCAACCTGCACCGGGTGCTGAT  
GGACCTGCAGAACCAGAACTGAAAGAGCTGAACGACTGGCTGACAAAGACTGAG  
GAACGCACAAGGAAGATGGAGGAGGAGCCACTGGGACCCGACCTGGAGGATCTGA  
AGAGACAGGTGCAGCAGCATAAGGTGCTGCAGGAGGATCTGGAACAGGAGCAGGT  
GCGGGTGAACCTCCCTGACACATATGGTGGT

Figure 21B

GGTGGTGGACGAATCTAGTGGAGATCACGCCACCGCCGCCCTGGAGGAACAGCTGA  
AGGTGCTGGGGGACCGGTGGGCCAACATTTGCCGGTGGACCGAGGACAGGTGGGTG  
CTGCTGCAGGACATCCTGCTGAAATGGCAGAGGCTGACCGAGGAGCAGTGTCTGTTT  
AGTGCTTGGCTGAGCGAGAAAGAGGACGCCGTGAACAAGATCCACACAACCGGCTT  
TAAGGATCAGAACGAAATGCTGTCTAGCCTGCAGAACTGGCTGTGCTGAAGGCCG  
ATCTGGAGAAAAAGAAGCAGAGCATGGGCAAACCTGTATAGCCTGAAACAGGACCTG  
CTGAGCACCTTGAAGAACAAGAGCGTGACCCAGAAGACAGAAGCCTGGCTGGATA  
ACTTTGCCCCGCTGCTGGGACAACCTGGTGCAGAACTGGAGAAAAGTACAGCTCAG  
ATCTCTCAGGCTGTGACCACAACCCAGCCTAGCCTGACCCAGACAACCGTGATGGA  
AACCGTGACCACCGTGACAACCCGCGAACAGATCCTGGTGAAACATGCCCAGGAAG  
AGCTGCCACCTCCACCTCCCCAGAAGAAGAGAACCCTGGAGCGGCTGCAGGAGCTG  
CAGGAAGCCACTGACGAACTGGACCTGAAGCTGAGGCAGGCCGAAGTGATTAAGGG  
GTCTTGGCAGCCTGTGGGCGATCTGCTGATTGATTCCCTGCAGGACCACCTGGAAAA  
GGTGAAGGCTCTGAGAGGCGAAATTGCTCCACTGAAGGAGAACGTGAGTCATGTGA  
ACGATCTGGCTAGACAGCTGACAACACTGGGCATCCAGCTGAGCCCATACAATCTG  
AGCACACTGGAGGACCTGAATACCAGGTGGAAGCTGCTGCAGGTGGCTGTGGAAGA  
CCGGGTGCGGCAGCTGCATGAGGCCCATCGCGACTTCGGACCAGCCAGCCAGCACT  
TTCTGAGCACATCCGTGCAGGGGGCCCTGGGAGAGGGCCATTTCTCCCAACAAGGTG  
CCCTACTATATTAATCACGAGACCCAGACCACTTGTGGGACCATCCCAAGATGACA  
GAACTGTACCAGTCCCTGGCCGATCTGAACAACGTGAGGTTTAGCGCTTACAGAACC  
GCTATGAAGCTGAGACGGCTGCAGAAGGCCCTGTGCCTGGATCTGCTGTCCCTGTCC  
GCCGCCTGCGATGCCCTGGATCAGCATAATCTGAAGCAGAACGATCAGCCAATGGA  
TATCCTGCAGATCATCAACTGCCTGACCACTATCTACGACAGGCTGGAGCAGGAGC  
ACAACAACCTGGTGAACGTGCCTCTGTGCGTGGATATGTGCCTGAACTGGCTGCTGA  
ACGTGTATGACACTGGGCGCACCGGCCGGATCAGAGTGCTGAGTTTTAAACTGGG  
ATTATCTCCCTGTGTAAGGCCACCTGGAGGACAAGTACAGGTACCTGTTCAAGCAG  
GTGGCTAGTAGCACTGGATTTTGTGACCAGCGCCGCCTGGGACTGCTGCTGCATGAT  
AGTATCCAGATTCCCTAGACAGCTGGGAGAGG

Figure 21C

TGGCTAGTTTCGGAGGATCTAACATCGAACCCAGCGTGCGCAGCTGTTTCCAGTTTG  
CCAATAACAAACCTGAAATCGAGGCTGCTCTGTTCCCTGGATTGGATGCGCCTGGAAC  
CACAGAGCATGGTGTGGCTGCCTGTGCTGCACAGAGTGGCTGCCGCCGAAACTGCC  
AAGCACCAGGCTAAATGCAACATCTGCAAGGAATGTCCCATTATCGGCTTTCGCTAC  
AGGAGTCTGAAACATTTTAACTACGATATTTGCCAGAGCTGCTTCTTTTCCGGAAGA  
GTGGCCAAAGGACACAAGATGCACTACCCTATGGTGGAATATTGCACCCCAACTAC  
ATCTGGCGAAGATGTGCGCGATTTTGCCAAGGTGCTGAAGAATAAGTTTCGGACTAA  
GAGGTACTTCGCCAAGCACCCCCGCATGGGGTATCTGCCAGTGCAGACAGTGCTGG  
AAGGAGACAATATGGAGACCGATACAATGTGAGC



Figure 22A

**pAAV.MCK.miR29C Sequence (SEQ ID NO: 12)**Main features:MCK enhancer- 190-395 (SEQ ID NO: 10)MCK promoter- 396-753 (SEQ ID NO: 11)EF1a Intron- 1155-1315, 1609-2083miR-29c-1487-1512shRNA-miR29-c with primary seed sequence- 1316-1608SV40 PolyA- 2094-2146

CTGNNNNNNNGCGCGCTCGCTCGCTCACTGAGGCCGCCCGGGCAAAGCCCAGGGCGTC  
 GGGCGACCTTTGGTCGCCCCGGCCTCAGTGAGCGAGCGAGCGCGCAGAGAGGGAGTG  
 GCCAACTCCATCACTAGGGGTTCTTGTAGTTAATGATTAACCCGCCATGCTAATTA  
 TCTACGTAGCCATGTCTAGACAGCCACTATGGGTCTAGGCTGCCCATGTAAGGAGGC  
AAGGCCTGGGGACACCCGAGATGCCTGGTTATAATTAACCCAGACATGTGGCTGCTC  
CCCCCCCCAACACCTGCTGCCTGAGCCTCACCCCCACCCCGGTGCCTGGGTCTTAG  
GCTCTGTACACCATGGAGGAGAAGCTCGCTCTAAAAATAACCCTGTCCCTGGTGGGC  
TGTGGGGGACTGAGGGCAGGCTGTAACAGGCTTGGGGGCCAGGGCTTATACGTGCC  
TGGGACTCCCAAAGTATTACTGTTCCATGTTCCCGGCGAAGGGCCAGCTGTCCCCG  
CCAGCTAGACTCAGCACTTAGTTTAGGAACCAGTGAGCAAGTCAGCCCTTGGGGCA  
GCCCATACAAGGCCATGGGGCTGGGCAAGCTGCACGCCTGGGTCCGGGGTGGGCAC  
GGTGCCCGGGCAACGAGCTGAAAGCTCATCTGCTCTCAGGGGCCCCCTCCCTGGGGA  
CAGCCCCCTCTGGCTAGTCACACCCTGTAGGCTCCTCTATATAACCCAGGGGCACAG  
GGGCTGCCCCCGGGTCAACCACCTCCACAGCACAGACAGACACTCAGGAGCCAG  
CCAGCCAGGTAAGTTTAGTCTTTTTGTCTTTTATTTAGGTCCCGGATCCGGTGGTGG  
TGCAAATCAAAGAACTGCTCCTCAGTGATGTTGCCTTTACTTCTAGGCCTGTACGG  
AAGTGTTACTTCTGCTCTAAAAGCTGCGGAATTGTACCCGCCTAGAGGATCCGGTAC  
TCGAGGAAGTGA AAAACCAGAAAGTTAACTGGTAAGTTTAGTCTTTTTGTCTTTAT  
TTCAGGTCCCGGATCCGGTGGTGGTGCAAATCAAAGAACTGCTCCTCAGTGATGTT  
GCCTTTACTTCTAGGCCTGTACGGAAGTGTTACTTCTGCTCTAAAAGCTGCGGAATT  
GTACCCGGGGCCGATCCACCGGTCTTTTTCGCAACGGGTTTGCCGCCAGAACACAGG  
TAAGTGCCGTGTGTGGTTCCCGCGGGCGGGCAGCGGGCCCCGTGCGTCCAGCGCAC  
ATGTTCCGGCAGGCGGGGCCTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGT  
CTCAAGCTGGCCGGCCTGTTTGAATGAGGCTTCAGTACTTTACAGAATCGTTGCCTG  
CACATCTTGGAACACTTGCTGGGATTACTTCTTCAGGTTAACCCAACAGAAGGCTC  
GAGAAGGTATATTGCTGTTGACAGTGAGCGCAACCGATTTCAAATGGTGCTAGAGT  
GAAGCCACAGATGTCTAGCACCATTTGAAATCGGTTATGCCTACTGCCTCGGAATTC  
AAGGGGCTACTTTAGGAGCAATTATCTTGTTTACTAAACTGAATACCTTGCTATCT  
CTTTGATACATTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCCGTGTATCGCC  
CCGCCCTGGGCGGCAAGGCTGGCCCGGTGCGCACCAAGTTGCGTGAGCGGAAAGATG  
GCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCGCTCGGGAG  
AGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTTTCCGTCCTCAGCCGTC  
GCTTCATGTGACTCCACGGAGTACCGGGCGCCGTCCAGGCACCTCGATTAGTT

Figure 22B

CTCGAGCTTTTGGAGTACGTCGTCTTTAGGTTGGGGGGAGGGGTTTTATGCGATGGA  
 GTTCCCCACACTGAGTGGGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTA  
 ATTCTCCTTGGAATTTGCCCTTTTTGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAG  
 ACAGTGGTTCAAAGTTTTTTTTCTTCCATTTACAGGTGTCGTGAAAAGCTAGTGC GGCC  
GCAATAAAAAGATCTTTATTTTCATTAGATCTGTGTGTTGGTTTTTTGTGTGTCTAGAC  
 ATGGCTACGTAGATAATTAGCATGGCGGGTAAATCATTAACTACAAGGAACCCCTAG  
 TGATGGAGTTGGCCACTCCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGAC  
 CAAAGGTCGCCCCGACGCCCGGGCTTTGCCCGGGCGGCCCTCAGTGAGCGAGCGAGCG  
 CGCNNNNNNCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAAC  
 AGTTGCGCAGCCTGAATGGCGAATGGAAGTTCCAGACGATTGAGCGTCAAAATGTA  
 GGTATTTCCATGAGCGTTTTTCTGTGTGCAATGGCTGGCGGTAATATTGTTCTGGATA  
 TTACCAGCAAGGCCGATAGTTTGAGTTCTTCTACTCAGGCAAGTGATGTTATTACTA  
 ATCAAAGAAGTATTGCGACAACGGTTAATTTGCGTGATGGACAGACTCTTTTACTCG  
 GTGGCCTCACTGATTATAAAAACACTTCTCAGGATTCTGGCGTACCGTTCCTGTCTA  
 AAATCCCTTTAATCGGCCTCCTGTTTAGCTCCCGCTCTGATTCTAACGAGGAAAGCA  
 CGTTATACGTGCTCGTCAAAGCAACCATAGTACGCGCCCTGTAGCGGCGCATTAAGC  
 GCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGC  
 GCCCGCTCCTTTGCTTTCTTCCCTTCCCTTCTCGCCACGTTGCGCGGCTTTCCCCGTC  
 AAGCTCTAAATCGGGGGCTCCCTTTAGGGTTCCGATTTAGTGATTTACGGCACCTCG  
 ACCCCAAAAAAGCTTGATTAGGGTGATGGTTCACGTAGTGGGCCATCGCCCTGATAGA  
 CGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGAAGTCTTGTTCCA  
 AACTGGAACAACACTCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTG  
 CCGATTTGCGCCTATTGGTTAAAAAATGAGCTGATTTAACAAAAATTTAACGCGAAT  
 TTTAACAAAAATATTAACGTTTACAATTTAAATATTTGCTTATACAATCTTCCTGTTTT  
 TGGGGCTTTTCTGATTATCAACCGGGGTACATATGATTGACATGCTAGTTTTACGATT  
 ACCGTTTCATCGATTCTCTTGTGTTGCTCCAGACTCTCAGGCAATGACCTGATAGCCTTT  
 GTAGAGACCTCTCAAAAATAGCTACCCTCTCCGGCATGAATTTATCAGCTAGAACGG  
 TTGAATATCATATTGATGGTGATTTGACTGTCTCCGGCCTTTCTCACCCGTTTGAATC  
 TTTACCTACACATTACTCAGGCATTGCATTTAAAATATATGAGGGTTCTAAAAATTTT  
 TATCCTTTCGTTGAAATAAAGGCTTCTCCCGCAAAAGTATTACAGGGTCATAATGTT  
 TTTGGTACAACCGATTTAGCTTTATGCTCTGAGGCTTTATTGCTTAATTTTGCTAATT  
 CTTTGCTTGCCTGTATGATTTATTGGATGTTGGAAGTTCCTGATGCGGTATTTTCTC  
 CTTACGCATCTGTGCGGTATTTACACCCGCATATGGTGCACTCTCAGTACAATCTGCT  
 CTGATGCCGCATAGTTAAGCCAGCCCCGACACCCGCCAACACCCGCTGACGCGCCCT  
 GACGGGCTTGCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGA  
 GCTGCATGTGTCAGAGGTTTTACCGTCATACCGAAACGCGCGAGACGAAAGGGC  
 CTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTTAGACGT  
 CAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTCTAAAT  
 ACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATA  
 TTGAAAAAGGAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATCCCTTTTTT  
 GCGGCATTTTGCCTTCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTAAAAGAT  
 GCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGG  
 TAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTT

Figure 22C

TTCCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTATTGA  
CGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGA  
GTA CTACCCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTAT  
GCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGA  
TCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAATC  
GCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGAC  
ACCACGATGCCTGTAGCAATGGCAACAACGTTGCGCAAACTATTA ACTGGCGAACT  
ACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGC  
AGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGG  
AGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGC  
CCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGA  
AATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAAGCATTGGTA ACTGTCAGAC  
CAAGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAGGA  
TCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTC  
GTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTT  
TTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGT  
TTGTTTGCCGGATCAAGAGCTACCAACTCTTTTTCCGAAGGTA ACTGGCTTCAGCAG  
AGCGCAGATACCAAATACTGTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAA  
GAACTCTGTAGCACCGCGTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGC  
TGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGA  
TAAGGCGCAGCGGTTCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTTGGAGC  
GAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACG  
CTTCCCGAAGGGAGAAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGAACAG  
GAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTATAGTCCTGTC  
GGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGG  
AGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGG  
CCTTTTGCTCACATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTAC  
CGGGTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGT  
CAGTGAGCGACCAAGCGGAAGAGC

## SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

