



(86) Date de dépôt PCT/PCT Filing Date: 2007/12/21
(87) Date publication PCT/PCT Publication Date: 2008/03/07
(45) Date de délivrance/Issue Date: 2015/04/21
(85) Entrée phase nationale/National Entry: 2009/06/19
(86) N° demande PCT/PCT Application No.: EP 2007/064423
(87) N° publication PCT/PCT Publication No.: 2008/077924
(30) Priorité/Priority: 2006/12/22 (EP06127149.0)


(51) Cl.Int./Int.Cl. *A01N 43/40* (2006.01),
A01N 51/00 (2006.01), *A01P 3/00* (2006.01),
A01P 7/04 (2006.01)

(72) Inventeurs/Inventors:
HUNGENBERG, HEIKE, DE;
THIELERT, WOLFGANG, DE;
BUSCHERMOEHLE, ALEXANDER, DE

(73) Propriétaire/Owner:
BAYER CROPSCIENCE AG, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : COMPOSITION PESTICIDE COMPRENANT UN DERIVE DE 2-PYRDILMETHYLBENZAMIDE ET UN
COMPOSE INSECTICIDE
(54) Title: PESTICIDAL COMPOSITION COMPRISING A 2-PYRDILMETHYLBENZAMIDE DERIVATIVE AND AN
INSECTICIDE COMPOUND



(57) Abrégé/Abstract:

A composition comprising at least a pyridylmethylbenzamide derivative of general formula (I) (A) and an insecticide compound (B) in a (A) / (B) weight ratio of from 1/1000 to 1000/1; a composition further comprising an additional fungicidal compound and a method for preventively or curatively combating the pests and diseases of crops by using this composition.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2008 (03.07.2008)

PCT

(10) International Publication Number
WO 2008/077924 A3

(51) International Patent Classification:

A01N 43/40 (2006.01) *A01P 7/04* (2006.01)
A01N 51/00 (2006.01) *A01P 3/00* (2006.01)

(21) International Application Number:

PCT/EP2007/064423

(22) International Filing Date:

21 December 2007 (21.12.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

06127149.0 22 December 2006 (22.12.2006) EP

(71) Applicant (for all designated States except US): **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Strasse 50, 40789 Monheim am Rhein (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HUNGENBERG, Heike** [DE/DE]; Louveciennesstrasse 2a, 40764 Langenfeld (DE). **THIELERT, Wolfgang** [DE/DE]; Buschweg 69, 51519 Odenthal (DE). **BUSCHERMOEHLE, Alexander** [DE/DE]; Schachstrasse 9, 50735 Köln (DE).

(74) Agent: **BAYER CROPSCIENCE AG**; Business Planning and Administration, Law and Patents, Patents and Licensing, Alfred-Nobel-Strasse 50, 40790 Monheim (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

7 May 2009

(54) Title: PESTICIDAL COMPOSITION COMPRISING A 2-PYRIDILMETHYLBENZAMIDE DERIVATIVE AND AN INSECTICIDE COMPOUND



(I)

(57) Abstract: A composition comprising at least a pyridylmethylbenzamide derivative of general formula (I) (A) and an insecticide compound (B) in a (A) / (B) weight ratio of from 1/1000 to 1000/1; a composition further comprising an additional fungicidal compound and a method for preventively or curatively combating the pests and diseases of crops by using this composition.

**PESTICIDAL COMPOSITION COMPRISING A 2-PYRIDYMETHYLBENZAMIDE DERIVATIVE AND AN
INSECTICIDE COMPOUND**


DESCRIPTION

- 5 The present invention relates to novel pesticidal compositions comprising a 2-pyridylmethylbenzamide derivative and an insecticide compound. The present invention also relates to a method of combating or controlling pests and diseases by applying at a locus infested or liable to be infested such a composition.
- 10 International patent application WO-99/42447 discloses fluopicolide among other 2-pyridylmethylbenzamide derivatives, as well as their use as fungicide. The possibility of combining one or more of these 2-pyridylmethylbenzamide derivatives with known fungicidal products to develop a fungicidal activity is disclosed. No mention is made of any association of fluopicolide with an insecticide active ingredient.
- 15 International patent application WO-2004/091299 discloses compositions comprising 2-pyridylmethylbenzamide derivatives according to the present invention in mixture with a chloronitrile fungicide derivative and their use as fungicide. No mention is made of mixtures comprising fenamidone with an insecticide active ingredient.
- In international patent application WO-2004/080181 there are generically disclosed numerous
20 mixtures of some phthalamide insecticide compounds with known fungicide substances. The association of these insecticide compounds with fluopicolide has not been tested for in connection with their combined fungicide and insecticide activity. Such an association does not form part of the present invention.
- 25 It is always of high-interest in agriculture to use novel pesticidal mixtures showing a broader scope of activity and a fungicide or insecticide synergistic effect in order notably to avoid or to control the development of resistant strains to the active ingredients or to the mixtures of known active ingredients used by the farmer while minimising the doses of chemical products spread in the environment and reducing the cost of the treatment.
- 30 We have now found some novel pesticidal compositions which possess the above mentioned characteristics.

Accordingly, the present invention relates to a composition comprising :

A) a 2-pyridylmethylbenzamide derivative of general formula (I)

2



wherein :

- R¹ may be a hydrogen atom, an optionally substituted alkyl group or an optionally substituted acyl group;


- R² may be a hydrogen atom or an optionally substituted alkyl group;

- R³ and R⁴ may be chosen independently from each other as being a halogen atom, a hydroxyl group, a cyano group, a nitro group, -SF₅, a trialkylsilyl group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which E may be an alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl or a heterocyclyl group each of which may optionally be substituted;

- p represents 0, 1, 2, 3 or 4;

- q represents 0, 1, 2, 3 or 4; and its agriculturally acceptable optical and/or geometric isomers, tautomers and addition salts with an acid or a base; and

B) an insecticide compound; in a (A) / (B) weight ratio of from 1/1000 to 1000/1; provided that insecticide B compound is different from compounds of general formula (II)



wherein :

- Hal represents a chlorine atom, a bromine atom or a iodine atom;

- R represents hydrogen or methyl and * may represent a carbon atom in R- or S-configuration.

In the context of the present invention :

- the term halogen means bromine, chlorine, iodine or fluorine;

30725-978

3

- the term alkyl means a linear or branched saturated hydrocarbon group containing from 1 to 6 carbon atoms;

- the term alkenyl means a linear or branched hydrocarbon group containing from 2 to 6 carbon atoms and an unsaturation in the form of double bond;

5 - the term alkynyl means a linear or branched hydrocarbon group containing from 2 to 6 carbon atoms and an unsaturation in the form of a triple bond;

- the term alkoxy means a linear or branched alkyloxy group containing from 1 to 6 carbon atoms;

10 - the term acyl means a formyl group or linear or branched alkoxy carbonyl group containing from 2 to 6 carbon atoms;

- the term cycloalkyl means a saturated cyclic hydrocarbon group containing from 3 to 8 carbon atoms;

- the term aryl means a phenyl or naphthyl group;


15 - the term heterocyclyl means saturated, partially saturated, unsaturated or aromatic cyclic group containing from 3 to 8 atoms, which may be a carbon atom, a nitrogen atom, a sulphur atom or an oxygen atom. Examples of such heterocyclyl may be pyridyl, pyridinyl, quinolyl, furyl, thienyl, pyrrolyl, oxazoliny;

20 - the term "optionally substituted " means that the group thus termed may be substituted with one or more groups which may be halogen, alkyl, alkoxy, hydroxyl, nitro, amin, cyano or acyl.

30725-978

3a

In one composition aspect, the invention relates to a composition comprising: (A) 2,6-dichloro-N-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl}benzamide (fluopicolide) of formula (Ia):



- 5 or agriculturally acceptable optical and/or geometric isomers, tautomers or addition salts with an acid or a base; and (B) an insecticide compound selected from the group consisting of: (b1) an acetylcholine receptor agonist/antagonist, (b2) an acetylcholinesterase (AChE) inhibitor, (b3) a sodium channel modulator/voltage-gated sodium channel blocker, (b5) a GABA-gated chloride channel antagonist, (b16) an inhibitor of lipid synthesis, (b17) a carboxamide, and
- 10 (b20) rynaxapyr, in a (A) / (B) weight ratio of from 1/1000 to 1000/1.

The composition according to the present invention provides a synergistic effect. This synergistic effect allows a reduction of the chemical substances spread into the environment and a reduction of the cost of the pesticidal treatment.

15 In the context of the present invention, the term “synergistic effect” is defined by Colby according to the article entitled “Calculation of the synergistic and antagonistic responses of herbicide combinations” Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = X + Y - \frac{XY}{100}$$

20 in which E represents the expected percentage of inhibition of the pest for the combination of the two pesticides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the pest by the compound (a) at a defined dose (equal to x), y is the percentage of inhibition observed for the pest by the compound (b) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The latter article also mentions the formula :

$$E = X + Y + Z - \frac{XYZ}{100}$$

in which E represents the expected percentage of inhibition of the pest for the combination of the three pesticides at defined doses (for example equal to x, y and z respectively), x is the percentage of inhibition observed for the pest by the compound (a) at a defined dose (equal to x), y is the percentage of inhibition observed for the pest by the compound (b) at a defined dose (equal to y) and z is the percentage of inhibition observed for the pest by the compound (c) at a defined dose (equal to z). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The composition according to the present invention comprises a 2-pyridylmethylbenzamide derivative of general formula (I).

Preferably, the present invention relates to a composition comprising a 2-pyridylmethylbenzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards R^1 and R^2 , R^1 and R^2 may be chosen independently from each other as being a hydrogen atom or an optionally substituted alkyl group. More preferably, R^1 and R^2 may be chosen independently from each other as being a hydrogen atom, a methyl group or an ethyl group. Even more preferably, R^1 and R^2 may be both hydrogen atoms.

- as regards R^3 and R^4 , R^3 and R^4 may be chosen independently from each other as being a halogen atom, a hydroxyl group, a nitro group, an optionally substituted amino group, an acyl group, or a group E, OE or SE, in which E may be a alkyl, a cycloalkyl, a phenyl or a heterocyclyl group, each of which may optionally be substituted. More preferably, R^3 and R^4 may be chosen independently from each other as being a halogen atom, a nitro group or a halogenoalkyl group. Even more preferably R^3 and R^4 may be chosen independently from each other as being a chlorine atom, a nitro group or a trifluoromethyl group.

- as regards p, p may be 1 or 2. More preferably, p may be 2.

- as regards q, q may be 1 or 2. More preferably, q may be 2;

and its agriculturally acceptable possible tautomers and addition salts with an acid or a base.

More preferably, the 2-pyridylmethylbenzamide derivative of general formula (I) present in the composition of the present invention is :

- a compound (Ia) which is 2,6-dichloro-N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide, also known as fluopicolide; or

- a compound (Ib) which is N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]-2-fluoro-6-nitrobenzamide; or

- a compound (Ic) which is N-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]-2-methyl-6-nitrobenzamide;

and its agriculturally acceptable possible tautomers and addition salts with an acid or a base.

- 5 The composition according to the present invention comprises an insecticide compound (b). Suitable insecticide are chosen in the following groups :

B1) acetylcholine receptor agonists/antagonists such as chloronicotinyis/neonicotinoids, nicotine, bensultap or cartap. Suitable examples of chloronicotinyis/neonicotinoids include acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, nithiazine, thiacloprid, thiamethoxam, imidaclothiz and (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine;

B2) acetylcholinesterase (AChE) inhibitors such as carbamates and organophosphates. Suitable examples of carbamates include alanycarb, aldicarb, aldoxycarb, allyxycarb, aminocarb, bendiocarb, benfuracarb, bufencarb, butacarb, butocarboxim, butoxycarboxim, carbaryl, carbofuran, carbosulfan, chloethocarb, dimetilan, ethiofencarb, fenobucarb, fenothiocarb, formetanate, furathiocarb, isoprocarb, metam-sodium, methiocarb, methomyl, metolcarb, oxamyl, phosphocarb, pirimicarb, promecarb, propoxur, thiodicarb, thiofanox, triazamate, trimethacarb, XMC and xylylcarb. Suitable examples of organophosphates include acephate, azamethiphos, azinphos (-methyl, -ethyl), bromophos-ethyl, bromfenvinfos (-methyl), butathiofos, cadusafos, carbophenothion, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos (-methyl/-ethyl), coumaphos, cyanofenphos, cyanophos, demeton-S-methyl, demeton-S-methylsulphon, dialifos, diazinon, dichlofenthion, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, dioxabenzofos, disulfoton, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fensulfothion, fenthion, flupyrazofos, fonofos, formothion, fosmethilan, fosthiazate, heptenophos, iodofenphos, iprobenfos, isazofos, isofenphos, isopropyl O-salicylate, isoxathion, malathion, mecarbam, methacrifos, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion (-methyl/-ethyl), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, pirimiphos (-methyl/-ethyl), profenofos, propaphos, propetamphos, prothiofos, prothoate, pyraclofos, pyridaphenthion, pyridathion, quinalphos, sebufos, sulfotep, sulprofos, tebupirimfos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, triclofon and vamidothion;

B3) sodium channel modulators/voltage-gated sodium channel blockers such as pyrethroids and oxadiazines. Suitable examples of pyrethroids include acrinathrin, allethrin (d-cis-trans, d-trans), beta-cyfluthrin, bifenthrin, bioallethrin, bioallethrin-S-cyclopentyl-isomer,

bioethanomethrin, biopermethrin, bioresmethrin, chlovaporthrin, cis-cypermethrin, cis-resmethrin, cis-permethrin, clocythrin, cycloprothrin, cyfluthrin, cyhalothrin, cypermethrin (alpha-, beta-, theta-, zeta-), cyphenothrin, DDT, deltamethrin, empenthrin (1R-isomer), esfenvalerate, etofenprox, fenfluthrin, fenpropathrin, fenpyrithrin, fenvalerate, flubrocycythrinate, flucythrinate, 5 flufenprox, flumethrin, fluvalinate, fubfenprox, gamma-cyhalothrin, imiprothrin, kadethrin, lambda-cyhalothrin, metofluthrin, permethrin (cis-, trans-), phenothrin (1R-trans isomer), prallethrin, profluthrin, protrifenbute, pyresmethrin, resmethrin, RU 15525, silafluofen, tau-fluvalinate, tefluthrin, terallethrin, tetramethrin (1R-isomer), traloccythrin, tralomethrin, transfluthrin, ZXI 8901 and pyrethrins (pyrethrum). Suitable example of oxadiazines includes 10 indoxacarb;

B4) acetylcholine receptor modulators such as spinosyns. Suitable example of spinosyns includes spinosad;

B5) GABA-gated chloride channel antagonists such as cyclodiene organochlorines and fiproles. Suitable examples of cyclodiene organochlorines include camphechlor, chlordane, 15 endosulfan, gamma-HCH, HCH, heptachlor, lindane and methoxychlor. Suitable examples of fiproles include acetoprole, ethiprole, fipronil and vaniliprole;

B6) chloride channel activators such as mectins. Suitable examples of mectins include abamectin, avermectin, emamectin, emamectin-benzoate, ivermectin, lepimectin, milbemectin and milbemycin;

20 B7) juvenile hormone mimetics such as diofenolan, epofenonane, fenoxycarb, hydroprene, kinoprene, methoprene, pyriproxifen, triprene;

B8) ecdysone agonists/disruptors such as diacylhydrazines. Suitable examples of diacylhydrazines include chromafenozide, halofenozide, methoxyfenozide and tebufenozide;

B9) inhibitors of chitinbiosynthesis such as benzoylureas, buprofezin and cyromazine. 25 Suitable examples of benzoylureas include bistrifluron, chlofluazuron, diflubenzuron, fluazuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, penfluron, teflubenzuron and triflumuron;

B10) inhibitors of oxidative phosphorylation, ATP disruptors such as organotins and diafenthiuron. Suitable examples of organotins include azocyclotin, cyhexatin and fenbutatin oxide;

30 B11) decouplers of oxidative phosphorylation by disruption of the H⁺ proton gradient such as pyrroles and dinitrophenols. Suitable example of pyrroles includes chlorfenapyr. Suitable examples of dinitrophenols include binapacryl, dinobuton, dinocap and DNOC;

B12) site I electron transport inhibitors such as METIs, hydramethylnone and dicofol. Suitable examples of METIs include fenazaquin, fenpyroximate, pyrimidifen, pyridaben, 35 tebufenpyrad, tolfenpyrad;

B13) site II electron transport inhibitors such as rotenone;

B14) site III electron transport inhibitors such as acequinocyl and fluacrypyrim;

B15) microbial disrupters of the intestinal membrane of insects such as *Bacillus thuringiensis* strains;

5 B16) inhibitors of lipid synthesis such as tetronic acids and tetramic acids. Suitable examples of tetronic acids include spirodiclofen, spiromesifen and spirotetramat. Suitable example of tetramic acids includes cis-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl carbonate (alias: carbonic acid, 3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl ester (CAS Reg. No.: 382608-10-8);

10 B17) carboxamides such as flonicamid;

B18) octopaminergic agonists such as amitraz;

B19) inhibitors of the magnesium-stimulated ATPase such as propargite;

B20) ryanodin receptor agonists such as phthalamides or rynaxapyr. Suitable example of phthalamides includes N²-[1,1-dimethyl-2-(methylsulphonyl)ethyl]-3-iodo-N¹-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide (i.e. flubendiamide, CAS reg. No.: 272451-65-7);

B21) nereistoxin analogues such as thiocyclam hydrogen oxalate and thiosultap-sodium;

B22) biologics, hormones or pheromones such as azadirachtin, *Bacillus spec.*, *Beauveria spec.*, codlemone, *Metarrhizium spec.*, *Paecilomyces spec.*, *thuringiensis* and *Verticillium spec.*;

20 B23) active compounds having unknown or non-specified mechanisms of action such as fumigants, selective feeding inhibitors, mite growth inhibitors, amidoflumet; bendlothiaz, benzoximate, bifenazate, bromopropylate, buprofezin, chinomethioat, chlordimeform, chlorobenzilate, chloropicrin, clothiazoben, cycloprene, cyflumetofen, dicyclanil, fenoxacrim, fentrifanil, flubenzimine, flufenerim, flutenzin, gossyplure, hydramethylnone, japonilure, metoxadiazon, petroleum, piperonyl butoxide, potassium oleate, pyrafluprole, pyridalyl, pyriprole, sulfluramid, tetradifon, tetrasul, triarathene, verbutin, furthermore the compound 3-methylphenyl propylcarbamate (Tsumacide Z), the compound 3-(5-chloro-3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane-3-carbonitrile (CAS reg. No. 185982-80-3) and the corresponding 3-endo isomer (CAS reg. No. 185984-60-5) (cf. WO 96/37494, WO 98/25923), and also preparations

30 comprising insecticidal effective plant extracts, nematodes, fungi or viruses. Suitable examples of fumigants include aluminium phosphide, methyl bromide and sulphuryl fluoride. Suitable examples of selective feeding inhibitors include cryolite, flonicamid and pymetrozine. Suitable examples of mite growth inhibitors include clofentezine, etoxazole and hexythiazox.

Preferably, the insecticide compound (b) is chosen as being abamectin, acephate, acetamiprid, acrinathrin, aldicarb, alpha-cypermethrin, beta-cyfluthrin, bifenthrin, carbaryl, carbofuran, chlorfenapyr, chlorfluazuron, chlorpyrifos-E, clothianidin, cyfluthrin, cypermethrin, cyromazine, deltamethrin, diflubenzuron, dinotefuran, emamectin benzoate, ethiprole, fenpyroximate, fipronil, flonicamid, flubendiamide, flufenoxuron, gamma-cyhalothrin, hexaflumuron, imidacloprid, indoxacarb, L-cyhalothrin, lepimectin, lufenuron, methamidophos, methiocarb, methomyl, methoxyfenozide, milbemycin, nitenpyram, novaluron, profenofos, pymetrozine, rynaxapyr, spinosad, spirotetramate, spiromesifen, spirotetramate, tebufenozide, tebufenozide, tebufenpyrad, tebufenpyrad, tebupirimphos, teflubenzuron, tefluthrin, thiacloprid, thiamethoxam, thiodicarb, triazophos triflumuron, imidaclothiz and (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine. More preferably, the insecticide compound (b) is chosen as being abamectin, acetamiprid, aldicarb, beta-cyfluthrin, carbofuran, chlorpyrifos-E, clothianidin, cypermethrin, cyromazine, deltamethrin, diflubenzuron, emamectin benzoate, ethiprole, fipronil, gamma-cyhalothrin, imidacloprid, L-cyhalothrin, lufenuron, methiocarb, methoxyfenozide, pymetrozine, rynaxapyr, spinosad, spirotetramate, spiromesifen, spirotetramate, tebufenozide, tebufenpyrad, tefluthrin, thiacloprid, thiamethoxam, thiodicarb, triflumuron, imidaclothiz and (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine. Even more preferably, the insecticide compound (b) is chosen as being abamectin, aldicarb, beta-cyfluthrin, chlorpyrifos-E, clothianidin, cyromazine, deltamethrin, diflubenzuron, emamectin benzoate, fipronil, gamma-cyhalothrin, imidacloprid, L-cyhalothrin, methiocarb, pymetrozine, rynaxapyr, spinosad, spirotetramate, spiromesifen, spirotetramate, tebufenozide, tebufenpyrad, tefluthrin, thiamethoxam, thiodicarb, imidaclothiz, (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine.

Non limitative examples of suitable mixtures according to the present invention may include mixtures of fluopicolide with abamectin, fluopicolide with acephate, fluopicolide with acetamiprid, fluopicolide with acrinathrin, fluopicolide with aldicarb, fluopicolide with alpha-cypermethrin, fluopicolide with beta-cyfluthrin, fluopicolide with bifenthrin, fluopicolide with carbaryl, fluopicolide with carbofuran, fluopicolide with chlorfenapyr, fluopicolide with chlorfluazuron, fluopicolide with chlorpyrifos-E, fluopicolide with clothianidin, fluopicolide with cyfluthrin, fluopicolide with cypermethrin, fluopicolide with cyromazine, fluopicolide with deltamethrin, fluopicolide with diflubenzuron, fluopicolide with dinotefuran, fluopicolide with emamectin benzoate, fluopicolide with ethiprole, fluopicolide with fenpyroximate, fluopicolide with fipronil, fluopicolide with flonicamid, fluopicolide with flubendiamide, fluopicolide with flufenoxuron, fluopicolide with gamma-cyhalothrin, fluopicolide with hexaflumuron, fluopicolide with

imidacloprid, fluopicolide with indoxacarb, fluopicolide with L-cyhalothrin, fluopicolide with
lepimectin, fluopicolide with lufenuron, fluopicolide with methamidophos, fluopicolide with
methiocarb, fluopicolide with methomyl, fluopicolide with methoxyfenozide, fluopicolide with
milbemycin, fluopicolide with nitenpyram, fluopicolide with novaluron, fluopicolide with
5 profenofos, fluopicolide with pymetrozine, fluopicolide with rynaxapyr, fluopicolide with
spinosad, fluopicolide with spirotetramate, fluopicolide with spiromesifen, fluopicolide with
spirotetramate, fluopicolide with tebufenozide, fluopicolide with tebufenozide, fluopicolide with
tebufenpyrad, fluopicolide with tebufenpyrad, fluopicolide with tebupirimphos, fluopicolide with
teflubenzuron, fluopicolide with tefluthrin, fluopicolide with thiacloprid, fluopicolide with
10 thiamethoxam, fluopicolide with thiodicarb, fluopicolide with triazophos, fluopicolide with
triflumuron, fluopicolide with imidaclothiz, fluopicolide with (2E)-1-[(2-chloro-1,3-thiazol-5-
yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine, compound (Ib) with abamectin,
compound (Ib) with acephate, compound (Ib) with acetamiprid, compound (Ib) with acrinathrin,
compound (Ib) with aldicarb, compound (Ib) with alpha-cypermethrin, compound (Ib) with beta-
15 cyfluthrin, compound (Ib) with bifenthrin, compound (Ib) with carbaryl, compound (Ib) with
carbofuran, compound (Ib) with chlorfenapyr, compound (Ib) with chlorfluazuron, compound (Ib)
with chlorpyrifos-E, compound (Ib) with clothianidin, compound (Ib) with cyfluthrin, compound
(Ib) with cypermethrin, compound (Ib) with cyromazine, compound (Ib) with deltamethrin,
compound (Ib) with diflubenzuron, compound (Ib) with diflubenzuron, compound (Ib) with
20 dinotefuran, compound (Ib) with emamectin benzoate, compound (Ib) with ethiprole, compound
(Ib) with fenpyroximate, compound (Ib) with fipronil, compound (Ib) with flonicamid, compound
(Ib) with flubendiamide, compound (Ib) with flufenoxuron, compound (Ib) with gamma-
cyhalothrin, compound (Ib) with hexaflumuron, compound (Ib) with imidacloprid, compound (Ib)
with indoxacarb, compound (Ib) with L-cyhalothrin, compound (Ib) with lepimectin, compound
25 (Ib) with lufenuron, compound (Ib) with methamidophos, compound (Ib) with methiocarb,
compound (Ib) with methomyl, compound (Ib) with methoxyfenozide, compound (Ib) with
milbemycin, compound (Ib) with nitenpyram, compound (Ib) with novaluron, compound (Ib) with
profenofos, compound (Ib) with pymetrozine, compound (Ib) with rynaxapyr, compound (Ib) with
spinosad, compound (Ib) with spirotetramate, compound (Ib) with spiromesifen, compound (Ib)
30 with spirotetramate, compound (Ib) with tebufenozide, compound (Ib) with tebufenozide,
compound (Ib) with tebufenpyrad, compound (Ib) with tebufenpyrad, compound (Ib) with
tebupirimphos, compound (Ib) with teflubenzuron, compound (Ib) with tefluthrin, compound (Ib)
with thiacloprid, compound (Ib) with thiamethoxam, compound (Ib) with thiodicarb, compound
(Ib) with triazophos, compound (Ib) with triflumuron, compound (Ib) with imidaclothiz, compound
35 (Ib) with (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine,

compound (Ic) with abamectin, compound (Ic) with acephate, compound (Ic) with acetamiprid,
compound (Ic) with acrinathrin, compound (Ic) with aldicarb, compound (Ic) with alpha-
cypermethrin, compound (Ic) with beta-cyfluthrin, compound (Ic) with bifenthrin, compound (Ic)
with carbaryl, compound (Ic) with carbofuran, compound (Ic) with chlorfenapyr, compound (Ic)
5 with chlorfluazuron, compound (Ic) with chlorpyrifos-E, compound (Ic) with clothianidin,
compound (Ic) with cyfluthrin, compound (Ic) with cypermethrin, compound (Ic) with cyromazine,
compound (Ic) with deltamethrin, compound (Ic) with diflubenzuron, compound (Ic) with
diflubenzuron, compound (Ic) with dinotefuran, compound (Ic) with emamectin benzoate,
compound (Ic) with ethiprole, compound (Ic) with fenpyroximate, compound (Ic) with fipronil,
10 compound (Ic) with flonicamid, compound (Ic) with flubendiamide, compound (Ic) with
flufenoxuron, compound (Ic) with gamma-cyhalothrin, compound (Ic) with hexaflumuron,
compound (Ic) with imidacloprid, compound (Ic) with indoxacarb, compound (Ic) with L-
cyhalothrin, compound (Ic) with lepimectin, compound (Ic) with lufenuron, compound (Ic) with
methamidophos, compound (Ic) with methiocarb, compound (Ic) with methomyl, compound (Ic)
15 with methoxyfenozide, compound (Ic) with milbemycin, compound (Ic) with nitenpyram,
compound (Ic) with novaluron, compound (Ic) with profenofos, compound (Ic) with pymetrozine,
compound (Ic) with rynaxapyr, compound (Ic) with spinosad, compound (Ic) with spirotetrafen,
compound (Ic) with spiromesifen, compound (Ic) with spirotetramate, compound (Ic) with
tebufenozide, compound (Ic) with tebufenozide, compound (Ic) with tebufenpyrad, compound
20 (Ic) with tebufenpyrad, compound (Ic) with tebupirimphos, compound (Ic) with teflubenzuron,
compound (Ic) with tefluthrin, compound (Ic) with thiacloprid, compound (Ic) with thiamethoxam,
compound (Ic) with thiodicarb, compound (Ic) with triazophos, compound (Ic) with triflumuron,
compound (Ic) with imidaclothiz and compound (Ic) with (2E)-1-[(2-chloro-1,3-thiazol-5-
yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine.

25

The composition according to the present invention comprises a compound of general formula (I) (a) and an insecticide compound (b) in a (a) / (b) weight ratio of from 1/1000 to 1000/1. Preferably, (a) / (b) weight ratio is of from 1/125 to 125/1. Even more preferably, (a) / (b) weight ratio is of from 1/25 to 25/1.

30 Furthermore in the combinations according to the invention the compounds (I) (a) and (b) are present in a synergistically effective weight ratio of A:B in a range of 100:1 to 1:50, 100:1 to 1:20, 50:1 to 1:100, 50:1 to 1:20, 20:1 to 1:100, 20:1 to 1:50. Further ratios of A:B which can be used according to the present invention with increasing preference in the order given are: 95:1 to 1:95, 95:1 to 1:90, 95:1 to 1:85, 95:1 to 1:80, 95:1 to 1:75, 95:1 to 1:70, 95:1 to 1:65, 95:1 to 1:60, 95:1 to 1:55, 95:1 to 1:50, 95:1 to 1:45, 95:1 to 1:40, 95:1 to 1:35, 95:1 to 1:30, 95:1 to

35

1:25, 95:1 to 1:20, 95:1 to 1:15, 95:1 to 1:10, 95:1 to 1:5, 95:1 to 1:4, 95:1 to 1:3, 95:1 to 1:2,
 90:1 to 1:90, 90:1 to 1:95, 90:1 to 1:85, 90:1 to 1:80, 90:1 to 1:75, 90:1 to 1:70, 90:1 to 1:65,
 90:1 to 1:60, 90:1 to 1:55, 90:1 to 1:50, 90:1 to 1:45, 90:1 to 1:40, 90:1 to 1:35, 90:1 to 1:30,
 90:1 to 1:25, 90:1 to 1:20, 90:1 to 1:15, 90:1 to 1:10, 90:1 to 1:5, 90:1 to 1:4, 90:1 to 1:3, 90:1 to
 5 1:2, 85:1 to 1:85, 85:1 to 1:95, 85:1 to 1:90, 85:1 to 1:80, 85:1 to 1:75, 85:1 to 1:70, 85:1 to
 1:65, 85:1 to 1:60, 85:1 to 1:55, 85:1 to 1:50, 85:1 to 1:45, 85:1 to 1:40, 85:1 to 1:35, 85:1 to
 1:30, 85:1 to 1:25, 85:1 to 1:20, 85:1 to 1:15, 85:1 to 1:10, 85:1 to 1:5, 85:1 to 1:4, 85:1 to 1:3,
 85:1 to 1:2, 80:1 to 1:80, 80:1 to 1:95, 80:1 to 1:90, 80:1 to 1:85, 80:1 to 1:75, 80:1 to 1:70, 80:1
 to 1:65, 80:1 to 1:60, 80:1 to 1:55, 80:1 to 1:50, 80:1 to 1:45, 80:1 to 1:40, 80:1 to 1:35, 80:1 to
 10 1:30, 80:1 to 1:25, 80:1 to 1:20, 80:1 to 1:15, 80:1 to 1:10, 80:1 to 1:5, 80:1 to 1:4, 80:1 to 1:3,
 80:1 to 1:2, 75:1 to 1:75, 75:1 to 1:95, 75:1 to 1:90, 75:1 to 1:85, 75:1 to 1:80, 75:1 to 1:70, 75:1
 to 1:65, 75:1 to 1:60, 75:1 to 1:55, 75:1 to 1:50, 75:1 to 1:45, 75:1 to 1:40, 75:1 to 1:35, 75:1 to
 1:30, 75:1 to 1:25, 75:1 to 1:20, 75:1 to 1:15, 75:1 to 1:10, 75:1 to 1:5, 75:1 to 1:4, 75:1 to 1:3,
 75:1 to 1:2, 70:1 to 1:70, 70:1 to 1:95, 70:1 to 1:90, 70:1 to 1:85, 70:1 to 1:80, 70:1 to 1:75, 70:1
 15 to 1:65, 70:1 to 1:60, 70:1 to 1:55, 70:1 to 1:50, 70:1 to 1:45, 70:1 to 1:40, 70:1 to 1:35, 70:1 to
 1:30, 70:1 to 1:25, 70:1 to 1:20, 70:1 to 1:15, 70:1 to 1:10, 70:1 to 1:5, 70:1 to 1:4, 70:1 to 1:3,
 70:1 to 1:2, 65:1 to 1:65, 65:1 to 1:95, 65:1 to 1:90, 65:1 to 1:85, 65:1 to 1:80, 65:1 to 1:75, 65:1
 to 1:70, 65:1 to 1:60, 65:1 to 1:55, 65:1 to 1:50, 65:1 to 1:45, 65:1 to 1:40, 65:1 to 1:35, 65:1 to
 1:30, 65:1 to 1:25, 65:1 to 1:20, 65:1 to 1:15, 65:1 to 1:10, 65:1 to 1:5, 65:1 to 1:4, 65:1 to 1:3,
 20 65:1 to 1:2, 60:1 to 1:60, 60:1 to 1:95, 60:1 to 1:90, 60:1 to 1:85, 60:1 to 1:80, 60:1 to 1:75, 60:1
 to 1:70, 60:1 to 1:65, 60:1 to 1:55, 60:1 to 1:50, 60:1 to 1:45, 60:1 to 1:40, 60:1 to 1:35, 60:1 to
 1:30, 60:1 to 1:25, 60:1 to 1:20, 60:1 to 1:15, 60:1 to 1:10, 60:1 to 1:5, 60:1 to 1:4, 60:1 to 1:3,
 60:1 to 1:2, 55:1 to 1:55, 55:1 to 1:95, 55:1 to 1:90, 55:1 to 1:85, 55:1 to 1:80, 55:1 to 1:75, 55:1
 to 1:70, 55:1 to 1:65, 55:1 to 1:60, 55:1 to 1:50, 55:1 to 1:45, 55:1 to 1:40, 55:1 to 1:35, 55:1 to
 25 1:30, 55:1 to 1:25, 55:1 to 1:20, 55:1 to 1:15, 55:1 to 1:10, 55:1 to 1:5, 55:1 to 1:4, 55:1 to 1:3,
 55:1 to 1:2, 50:1 to 1:95, 50:1 to 1:90, 50:1 to 1:85, 50:1 to 1:80, 50:1 to 1:75, 50:1 to 1:70, 50:1
 to 1:65, 50:1 to 1:60, 50:1 to 1:55, 50:1 to 1:45, 50:1 to 1:40, 50:1 to 1:35, 50:1 to 1:30, 50:1 to
 1:25, 50:1 to 1:20, 50:1 to 1:15, 50:1 to 1:10, 50:1 to 1:5, 50:1 to 1:4, 50:1 to 1:3, 50:1 to 1:2,
 45:1 to 1:45, 45:1 to 1:95, 45:1 to 1:90, 45:1 to 1:85, 45:1 to 1:80, 45:1 to 1:75, 45:1 to 1:70,
 30 45:1 to 1:65, 45:1 to 1:60, 45:1 to 1:55, 45:1 to 1:50, 45:1 to 1:40, 45:1 to 1:35, 45:1 to 1:30,
 45:1 to 1:25, 45:1 to 1:20, 45:1 to 1:15, 45:1 to 1:10, 45:1 to 1:5, 45:1 to 1:4, 45:1 to 1:3, 45:1 to
 1:2, 40:1 to 1:40, 40:1 to 1:95, 40:1 to 1:90, 40:1 to 1:85, 40:1 to 1:80, 40:1 to 1:75, 40:1 to
 1:70, 40:1 to 1:65, 40:1 to 1:60, 40:1 to 1:55, 40:1 to 1:50, 40:1 to 1:45, 40:1 to 1:35, 40:1 to
 1:30, 40:1 to 1:25, 40:1 to 1:20, 40:1 to 1:15, 40:1 to 1:10, 40:1 to 1:5, 40:1 to 1:4, 40:1 to 1:3,
 35 40:1 to 1:2, 35:1 to 1:35, 35:1 to 1:95, 35:1 to 1:90, 35:1 to 1:85, 35:1 to 1:80, 35:1 to 1:75, 35:1

to 1:70, 35:1 to 1:65, 35:1 to 1:60, 35:1 to 1:55, 35:1 to 1:50, 35:1 to 1:45, 35:1 to 1:40, 35:1 to 1:30, 35:1 to 1:25, 35:1 to 1:20, 35:1 to 1:15, 35:1 to 1:10, 35:1 to 1:5, 35:1 to 1:4, 35:1 to 1:3, 35:1 to 1:2, 30:1 to 1:30, 30:1 to 1:95, 30:1 to 1:90, 30:1 to 1:85, 30:1 to 1:80, 30:1 to 1:75, 30:1 to 1:70, 30:1 to 1:65, 30:1 to 1:60, 30:1 to 1:55, 30:1 to 1:50, 30:1 to 1:45, 30:1 to 1:40, 30:1 to 1:35, 30:1 to 1:25, 30:1 to 1:20, 30:1 to 1:15, 30:1 to 1:10, 30:1 to 1:5, 30:1 to 1:4, 30:1 to 1:3, 30:1 to 1:2, 25:1 to 1:25, 25:1 to 1:95, 25:1 to 1:90, 25:1 to 1:85, 25:1 to 1:80, 25:1 to 1:75, 25:1 to 1:70, 25:1 to 1:65, 25:1 to 1:60, 25:1 to 1:55, 25:1 to 1:50, 25:1 to 1:45, 25:1 to 1:40, 25:1 to 1:35, 25:1 to 1:30, 25:1 to 1:20, 25:1 to 1:15, 25:1 to 1:10, 25:1 to 1:5, 25:1 to 1:4, 25:1 to 1:3, 25:1 to 1:2, 20:1 to 1:95, 20:1 to 1:90, 20:1 to 1:85, 20:1 to 1:80, 20:1 to 1:75, 20:1 to 1:70, 20:1 to 1:65, 20:1 to 1:60, 20:1 to 1:55, 20:1 to 1:50, 20:1 to 1:45, 20:1 to 1:40, 20:1 to 1:35, 20:1 to 1:30, 20:1 to 1:25, 20:1 to 1:15, 20:1 to 1:10, 20:1 to 1:5, 20:1 to 1:4, 20:1 to 1:3, 20:1 to 1:2, 15:1 to 1:15, 15:1 to 1:95, 15:1 to 1:90, 15:1 to 1:85, 15:1 to 1:80, 15:1 to 1:75, 15:1 to 1:70, 15:1 to 1:65, 15:1 to 1:60, 15:1 to 1:55, 15:1 to 1:50, 15:1 to 1:45, 15:1 to 1:40, 15:1 to 1:35, 15:1 to 1:30, 15:1 to 1:25, 15:1 to 1:20, 15:1 to 1:10, 15:1 to 1:5, 15:1 to 1:4, 15:1 to 1:3, 15:1 to 1:2, 10:1 to 1:10, 10:1 to 1:95, 10:1 to 1:90, 10:1 to 1:85, 10:1 to 1:80, 10:1 to 1:75, 10:1 to 1:70, 10:1 to 1:65, 10:1 to 1:60, 10:1 to 1:55, 10:1 to 1:50, 10:1 to 1:45, 10:1 to 1:40, 10:1 to 1:35, 10:1 to 1:30, 10:1 to 1:25, 10:1 to 1:20, 10:1 to 1:15, 10:1 to 1:5, 10:1 to 1:4, 10:1 to 1:3, 10:1 to 1:2, 5:1 to 1:5, 5:1 to 1:95, 5:1 to 1:90, 5:1 to 1:85, 5:1 to 1:80, 5:1 to 1:75, 5:1 to 1:70, 5:1 to 1:65, 5:1 to 1:60, 5:1 to 1:55, 5:1 to 1:50, 5:1 to 1:45, 5:1 to 1:40, 5:1 to 1:35, 5:1 to 1:30, 5:1 to 1:25, 5:1 to 1:20, 5:1 to 1:15, 5:1 to 1:10, 5:1 to 1:4, 5:1 to 1:3, 5:1 to 1:2, 4:1 to 1:4, 4:1 to 1:95, 4:1 to 1:90, 4:1 to 1:85, 4:1 to 1:80, 4:1 to 1:75, 4:1 to 1:70, 4:1 to 1:65, 4:1 to 1:60, 4:1 to 1:55, 4:1 to 1:50, 4:1 to 1:45, 4:1 to 1:40, 4:1 to 1:35, 4:1 to 1:30, 4:1 to 1:25, 4:1 to 1:20, 4:1 to 1:15, 4:1 to 1:10, 4:1 to 1:5, 4:1 to 1:3, 4:1 to 1:2, 3:1 to 1:3, 3:1 to 1:95, 3:1 to 1:90, 3:1 to 1:85, 3:1 to 1:80, 3:1 to 1:75, 3:1 to 1:70, 3:1 to 1:65, 3:1 to 1:60, 3:1 to 1:55, 3:1 to 1:50, 3:1 to 1:45, 3:1 to 1:40, 3:1 to 1:35, 3:1 to 1:30, 3:1 to 1:25, 3:1 to 1:20, 3:1 to 1:15, 3:1 to 1:10, 3:1 to 1:5, 3:1 to 1:4, 3:1 to 1:2, 2:1 to 1:2, 2:1 to 1:95, 2:1 to 1:90, 2:1 to 1:85, 2:1 to 1:80, 2:1 to 1:75, 2:1 to 1:70, 2:1 to 1:65, 2:1 to 1:60, 2:1 to 1:55, 2:1 to 1:50, 2:1 to 1:45, 2:1 to 1:40, 2:1 to 1:35, 2:1 to 1:30, 2:1 to 1:25, 2:1 to 1:20, 2:1 to 1:15, 2:1 to 1:10, 2:1 to 1:5, 2:1 to 1:4, 2:1 to 1:3.

30

The composition of the present invention may further comprise at least one other different fungicide active ingredient (c).

Examples of suitable fungicide mixing partners may be selected in the following lists :

C1) a compound capable to inhibit the nucleic acid synthesis like benalaxyl, benalaxyl-M, bupirimate, chiralaxyl, clozylacon, dimethirimol, ethirimol, furalaxyl, hymexazol, mefenoxam, metalaxyl, metalaxyl-M, ofurace, oxadixyl, oxolinic acid ;

5 C2) a compound capable to inhibit the mitosis and cell division like benomyl, carbendazim, diethofencarb, ethaboxam, fuberidazole, pencycuron, thiabendazole thiophanate-methyl, zoxamide;

C3) a compound capable to inhibit the respiration for example

as CI-respiration inhibitor like diflumetorim;

10 as CII-respiration inhibitor like boscalid, carboxin, fenfuram, flutolanil, furametpyr, furmecyclox, mepronil, oxycarboxine, penthiopyrad, thifluzamide;

as CIII-respiration inhibitor like amisulbrom, azoxystrobin, cyazofamid, dimoxystrobin, enestrobin, famoxadone, fenamidone, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, trifloxystrobin;

15 C4) a compound capable of to act as an uncoupler like dinocap, fluazinam, meptyldinocap;

C5) a compound capable to inhibit ATP production like fentin acetate, fentin chloride, fentin hydroxide, silthiofam;

20 C6) a compound capable to inhibit AA and protein biosynthesis like andoprim, blasticidin-S, cyprodinil, kasugamycin, kasugamycin hydrochloride hydrate, mepaniprim, pyrimethanil;

C7) a compound capable to inhibit the signal transduction like fenpiclonil, fludioxonil, quinoxyfen;

25 C8) a compound capable to inhibit lipid and membrane synthesis like biphenyl, chlozolate, edifenphos, etridiazole, iodocarb, iprobenfos, iprodione, isoprothiolane, procymidone, propamocarb, propamocarb hydrochloride, pyrazophos, tolclofos-methyl, vinclozolin ;

30 C9) a compound capable to inhibit ergosterol biosynthesis like aldimorph, azaconazole, bitertanol, bromuconazole, cyproconazole, diclobutrazole, difenoconazole, diniconazole, diniconazole-M, dodemorph, dodemorph acetate, epoxiconazole, etaconazole, fenarimol, fenbuconazole, fenhexamid, fenpropidin, fenpropimorph, fluquinconazole, flurprimidol, flusilazole, flutriafol, furconazole, furconazole-cis, hexaconazole, imazalil, imazalil sulfate, imibenconazole, ipconazole, metconazole, myclobutanil, naftifine, nuarimol, oxpoconazole, paclobutrazol, pefurazoate, penconazole, prochloraz, propiconazole, prothioconazole, pyributicarb, pyrifenox, simeconazole, spiroxamine, tebuconazole, terbinafine, tetraconazole,

triadimefon, triadimenol, tridemorph, triflumizole, triforine, triticonazole, uniconazole, viniconazole, voriconazole;

C10) a compound capable to inhibit cell wall synthesis like bentiavalicarb, bialaphos, dimethomorph, flumorph, iprovalicarb, mandipropamid, polyoxins, polyoxorim, validamycin A;

5 C11) a compound capable to inhibit melanine biosynthesis like carpropamid, diclocymet, fenoxanil, phthalide, pyroquilon, tricyclazole;

C12) a compound capable to induce a host defence like acibenzolar-S-methyl, probenazole, tiadinil;

10 C13) a compound capable to have a multisite action like Bordeaux mixture, captafol, captan, chlorothalonil, copper naphthenate, copper oxide, copper oxychloride, copper preparations such as copper hydroxide, copper sulphate, dichlofluanid, dithianon, dodine, dodine free base, ferbam, fluorofolpet, folpet, guazatine, guazatine acetate, iminoctadine, iminoctadine albesilate, iminoctadine triacetate, mancopper, mancozeb, maneb, metiram, metiram zinc, oxine-copper, propineb, sulphur and sulphur preparations including calcium
15 polysulphide, thiram, tolylfluanid, zineb, ziram;

C14) a compound selected in the following list: (2E)-2-(2-([6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl]oxy}phenyl)-2-(methoxyimino)-N-methylacetamide, (2E)-2-(2-([([1E)-1-(3-([E)-1-fluoro-2-phenylvinyl]oxy}phenyl)ethylidene]amino}oxy)methyl]phenyl)-2-(methoxyimino)-N-methylacetamide, 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol,
20 1-[(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl-1H-imidazole-1-carboxylate, 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine, 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)nicotinamide, 2-phenylphenol and salts, 3,4,5-trichloropyridine-2,6-dicarbonitrile, 3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide, 3-[5-(4-chlorophenyl)-2,3-dimethylisoxazolidin-3-yl]pyridine, 5-chloro-6-(2,4,6-trifluorophenyl)-N-
25 [(1R)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidin-7-amine, 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine, 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, 8-hydroxyquinoline sulfate, benthiazole, bethoxazin, capsimycin, carvone, chinomethionat, cufraneb, cyflufenamid, cymoxanil, dazomet, debacarb, dichlorophen, diclomezine, dicloran,
30 difenzoquat, difenzoquat methylsulphate, diphenylamine, ferimzone, flumetover, fluopicolide, fluoroimide, flusulfamide, fosetyl-aluminium, fosetyl-calcium, fosetyl-sodium, hexachlorobenzene, irumamycin, isotianil, methasulfocarb, methyl (2E)-2-(2-([cyclopropyl]-(4-methoxyphenyl)imino)methyl}thio)methyl]phenyl)-3-methoxyacrylate, methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate, methyl isothiocyanate, metrafenone,
35 mildiomycin, N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1,3-dimethyl-1H-

pyrazole-4-carboxamide, N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-(formylamino)-2-hydroxybenzamide, N-(4-chloro-2-nitrophenyl)-N-ethyl-4-methylbenzenesulfonamide, N-(4-chlorobenzyl)-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, N-[(4-chlorophenyl)(cyano)methyl]-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloronicotinamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloronicotinamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodonicotinamide, N-[2-(4-{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl)ethyl]-N-(methylsulfonyl)valinamide, N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide, N-{2-[1,1'-bi(cyclopropyl)-2-yl]phenyl}-3-(difluoromethyl)-, 1-methyl-1H-pyrazole-4-carboxamide, N-{2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethyl}-2-(trifluoromethyl)benzamide, natamycin, N-ethyl-N-methyl-N'-(2-methyl-5-(trifluoromethyl)-4-[3-(trimethylsilyl)propoxy]phenyl)imidoformamide, N-ethyl-N-methyl-N'-(2-methyl-5-(difluoromethyl)-4-[3-(trimethylsilyl)propoxy]phenyl)imidoformamide, nickel dimethyldithiocarbamate, nitrothal-isopropyl, O-{1-[(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl}1H-imidazole-1-carbothioate, octhilinone, oxamocarb, oxyfenthiin, pentachlorophenol and salts, phosphorous acid and its salts, piperalin, propamocarb fosetyl, propanosine-sodium, proquinazid, pyribencarb, pyrrolnitrine, quintozone, tecloftalam, tecnazene, triazoxide, trichlamide, valiphenal, zarilamid.

20

Preferably, fungicidal active ingredient (c) is selected from N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, benalaxyl, benalaxyl-M, benthiavalicarb, carboxin, chlorothalonil, cyazofamid, cymoxanil, dimetomorph, fluazinam, fludioxonil, fluquinconazole, fluoxastrobin, flutriafol, fosetyl-aluminium, hexaconazole, hymexazole, ipconazole, mancozeb, mandipropamid, maneb, mefenoxam, metiram, metalaxyl, metalaxyl-M, peconazole, penthiopyrad, phosphorous acid, propamocarb.HCl, propineb, prothioconazole, tebuconazole, thiram, triadimenol, trifloxystrobin and triticonazole.

Where the third active ingredient (c) as defined above is present in the composition, this compound may be present in an amount of (a) : (b) : (c) weight ratio of from 1 : 0.001 : 0.001 to 1 : 1000 : 1000; the ratios of compound (a) and compound (c) varying independently from each other. Preferably, the (a) : (b) : (c) weight ratio may be of from 1 : 0.01 : 0.01 to 1 : 100 : 100. More preferably, the (a) : (b) : (c) weight ratio may be of from 1 : 0.05 : 0.05 to 1 : 80 : 80.

Following compositions may be cited to illustrate in a non-limited manner the present invention :

fluopicolide with N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide and clothianidin, fluopicolide with benalaxyl and clothianidin, fluopicolide with benalaxyl-M and clothianidin, fluopicolide with benthiavalicarb and clothianidin, fluopicolide with carboxin and clothianidin, fluopicolide with chlorothalonil and clothianidin, fluopicolide with cyazofamid and clothianidin, fluopicolide with cymoxanil and clothianidin, fluopicolide with dimetomorph and clothianidin, fluopicolide with fluazinam and clothianidin, fluopicolide with fludioxonil and clothianidin, fluopicolide with fluquinconazole and clothianidin, fluopicolide with fluoxastrobin and clothianidin, fluopicolide with flutriafol and clothianidin, fluopicolide with fosetyl-aluminium and clothianidin, fluopicolide with hexaconazole and clothianidin, fluopicolide with hymexazole and clothianidin, fluopicolide with ipconazole and clothianidin, fluopicolide with mancozeb and clothianidin, fluopicolide with mandipropamid and clothianidin, fluopicolide with maneb and clothianidin, fluopicolide with mefenoxam and clothianidin, fluopicolide with metiram and clothianidin, fluopicolide with metalaxyl and clothianidin, fluopicolide with metalaxyl-M and clothianidin, fluopicolide with peconazole and clothianidin, fluopicolide with penthiopyrad and clothianidin, fluopicolide with phosphorous acid and clothianidin, fluopicolide with propamocarb.HCl and clothianidin, fluopicolide with propineb and clothianidin, fluopicolide with prothioconazole and clothianidin, fluopicolide with tebuconazole and clothianidin, fluopicolide with thiram and clothianidin, fluopicolide with triadimenol and clothianidin, fluopicolide with trifloxystrobin and clothianidin, fluopicolide with triticonazole and clothianidin, fluopicolide with N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide and imidacloprid, fluopicolide with benalaxyl and imidacloprid, fluopicolide with benalaxyl-M and imidacloprid, fluopicolide with benthiavalicarb and imidacloprid, fluopicolide with carboxin and imidacloprid, fluopicolide with chlorothalonil and imidacloprid, fluopicolide with cyazofamid and imidacloprid, fluopicolide with cymoxanil and imidacloprid, fluopicolide with dimetomorph and imidacloprid, fluopicolide with fluazinam and imidacloprid, fluopicolide with fludioxonil and imidacloprid, fluopicolide with fluquinconazole and imidacloprid, fluopicolide with fluoxastrobin and imidacloprid, fluopicolide with flutriafol and imidacloprid, fluopicolide with fosetyl-aluminium and imidacloprid, fluopicolide with hexaconazole and imidacloprid, fluopicolide with hymexazole and imidacloprid, fluopicolide with ipconazole and imidacloprid, fluopicolide with mancozeb and imidacloprid, fluopicolide with mandipropamid and imidacloprid, fluopicolide with maneb and imidacloprid, fluopicolide with mefenoxam and imidacloprid, fluopicolide with metiram and imidacloprid, fluopicolide with metalaxyl and imidacloprid, fluopicolide with metalaxyl-M and imidacloprid, fluopicolide with peconazole and imidacloprid, fluopicolide with penthiopyrad and imidacloprid, fluopicolide with phosphorous acid and imidacloprid, fluopicolide with

propamocarb.HCl and imidacloprid, fluopicolide with propineb and imidacloprid, fluopicolide with prothioconazole and imidacloprid, fluopicolide with tebuconazole and imidacloprid, fluopicolide with thiram and imidacloprid, fluopicolide with triadimenol and imidacloprid, fluopicolide with trifloxystrobin and imidacloprid, fluopicolide with triticonazole and imidacloprid, fluopicolide with
5 N-[2-(1,3-dimethylbutyl)phenyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide and thiametoxam, fluopicolide with benalaxyl and thiametoxam, fluopicolide with benalaxyl-M and thiametoxam, fluopicolide with benthiavalicarb and thiametoxam, fluopicolide with carboxin and thiametoxam, fluopicolide with chlorothalonil and thiametoxam, fluopicolide with cyazofamid and thiametoxam, fluopicolide with cymoxanil and thiametoxam, fluopicolide with dimetomorph and
10 thiametoxam, fluopicolide with fluazinam and thiametoxam, fluopicolide with fludioxonil and thiametoxam, fluopicolide with fluquinconazole and thiametoxam, fluopicolide with fluoxastrobin and thiametoxam, fluopicolide with flutriafol and thiametoxam, fluopicolide with fosetyl-aluminium and thiametoxam, fluopicolide with hexaconazole and thiametoxam, fluopicolide with hymexazole and thiametoxam, fluopicolide with ipconazole and thiametoxam, fluopicolide with
15 mancozeb and thiametoxam, fluopicolide with mandipropamid and thiametoxam, fluopicolide with maneb and thiametoxam, fluopicolide with mefenoxam and thiametoxam, fluopicolide with metiram and thiametoxam, fluopicolide with metalaxyl and thiametoxam, fluopicolide with metalaxyl-M and thiametoxam, fluopicolide with peconazole and thiametoxam, fluopicolide with penthiopyrad and thiametoxam, fluopicolide with phosphorous acid and thiametoxam,
20 fluopicolide with propamocarb.HCl and thiametoxam, fluopicolide with propineb and thiametoxam, fluopicolide with prothioconazole and thiametoxam, fluopicolide with tebuconazole and thiametoxam, fluopicolide with thiram and thiametoxam, fluopicolide with triadimenol and thiametoxam, fluopicolide with trifloxystrobin and thiametoxam and fluopicolide with triticonazole and thiametoxam.

25

The composition according to the present invention may further comprise an other additional component such as an agriculturally acceptable support, carrier or filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably
30 to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid. Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

The composition may also comprise other additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before they are applied to the crop.

The pesticidal compositions of the present invention can be used to curatively or preventively control phytopathogenic fungi of crops but also to curatively or preventively control insects.

Thus, according to a further aspect of the present invention, there is provided a method for preventively or curatively controlling phytopathogenic fungi of crops but also to curatively or preventively control insects characterised in that an effective and non-phytotoxic amount of a composition as hereinbefore defined is applied via seed treatment, foliar application, stem application or drench/drip application (chemigation) to the seed, the plant and/or to the fruit of the plant or to soil and/or to inert substrate (e.g. inorganic substrates (e.g. sand, rockwool, glasswool, expanded minerals (e.g. perlite, vermiculite, zeolite, expanded clay)), Pumice, Pyroclastic materials/tuff, synthetic organic substrates (e.g. Polyurethane), organic substrates (e.g. peat, composts, tree waste products (e.g. coir, wood fibre/chips, tree bark)) and/or to a liquid substrate (e.g. floating hydroponic systems, Nutrient Film Technique, Aeroponics) in which the plant is growing or in which it is desired to grow.

The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the pests and/or diseases present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the pests and diseases to be combated or controlled, the type of crop, the climatic conditions and the compounds included in the composition according to the invention.

This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

Among the plants that can be protected by the method according to the present invention, mention may be made of cotton; flax; vine; fruit or vegetable crops such as *Rosaceae* sp. (for instance pip fruit such as apples and pears, but also stone fruit such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp., *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantains), *Rubiaceae* sp., *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruit); *Solanaceae* sp. (for instance tomatoes), *Liliaceae*

sp., *Asteraceae* sp. (for instance lettuces), *Umbelliferae* sp., *Cruciferae* sp., *Chenopodiaceae* sp., *Cucurbitaceae* sp., *Papilionaceae* sp. (for instance peas), *Rosaceae* sp. (for instance strawberries); major crops such as *Graminae* sp. (for instance maize, lawn or cereals such as wheat, rice, barley and triticales), *Asteraceae* sp. (for instance sunflower), *Cruciferae* sp. (for instance colza), *Fabaceae* sp. (for instance peanuts), *Papilionaceae* sp. (for instance soybean), *Solanaceae* sp. (for instance potatoes), *Chenopodiaceae* sp. (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

The method of treatment according to the invention can be used in the treatment of genetically modified organisms (GMOs), e.g. plants or seeds. Genetically modified plants (or transgenic plants) are plants in which a heterologous gene has been stably integrated into the genome. The expression "heterologous gene" essentially means a gene which is provided or assembled outside the plant and when introduced in the nuclear, chloroplastic or mitochondrial genome gives the transformed plant new or improved agronomic or other properties by expressing a protein or polypeptide of interest or by downregulating or silencing other gene(s) which are present in the plant (using for example, antisense technology, co suppression technology or RNA interference – RNAi - technology). A heterologous gene that is located in the genome is also called a transgene. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the active compounds and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, bigger fruits, larger plant height, greener leaf color, earlier flowering, higher quality and/or a higher nutritional value of the harvested products, higher sugar concentration within the fruits, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

At certain application rates, the active compound combinations according to the invention may also have a strengthening effect in plants. Accordingly, they are also suitable for mobilizing the defense system of the plant against attack by unwanted phytopathogenic fungi and/ or microorganisms and/or viruses. This may, if appropriate, be one of the reasons of the enhanced activity of the combinations according to the invention, for example against fungi. Plant-

strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances or combinations of substances which are capable of stimulating the defense system of plants in such a way that, when subsequently inoculated with unwanted phytopathogenic fungi and/ or microorganisms and/or viruses, the treated plants display a substantial degree of resistance to these unwanted phytopathogenic fungi and/ or microorganisms and/or viruses. In the present case, unwanted phytopathogenic fungi and/ or microorganisms and/or viruses are to be understood as meaning phytopathogenic fungi, bacteria and viruses. Thus, the substances according to the invention can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period of time within which protection is effected generally extends from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active compounds.

Plants and plant cultivars which are preferably to be treated according to the invention include all plants which have genetic material which impart particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

Plants and plant cultivars which are also preferably to be treated according to the invention are resistant against one or more biotic stresses, i.e. said plants show a better defense against animal and microbial pests, such as against nematodes, insects, mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

Plants and plant cultivars which may also be treated according to the invention are those plants which are resistant to one or more abiotic stresses. Abiotic stress conditions may include, for example, drought, cold temperature exposure, heat exposure, osmotic stress, flooding, increased soil salinity, increased mineral exposure, ozon exposure, high light exposure, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients, shade avoidance.

Plants and plant cultivars which may also be treated according to the invention, are those plants characterized by enhanced yield characteristics. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can furthermore be affected by improved plant architecture (under stress and non-stress conditions), including but not limited to, early flowering, flowering control for hybrid seed production, seedling vigor, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content, protein content, oil content

and composition, nutritional value, reduction in anti-nutritional compounds, improved processability and better storage stability.

Plants that may be treated according to the invention are hybrid plants that already express the characteristic of heterosis or hybrid vigor which results in generally higher yield, vigor, health and resistance towards biotic and abiotic stress factors. Such plants are typically made by crossing an inbred male-sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). Hybrid seed is typically harvested from the male sterile plants and sold to growers. Male sterile plants can sometimes (e.g. in corn) be produced by detasseling, i.e. the mechanical removal of the male reproductive organs (or males flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants it is typically useful to ensure that male fertility in the hybrid plants is fully restored. This can be accomplished by ensuring that the male parents have appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic determinants responsible for male-sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described in Brassica species (WO 1992/005251, WO 1995/009910, WO 1998/27806, WO 2005/002324, WO 2006/021972 and US 6,229,072). However, genetic determinants for male sterility can also be located in the nuclear genome. Male sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 1989/10396 in which, for example, a ribonuclease such as barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar (e.g. WO 1991/002069).

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated according to the invention are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

Herbicide-tolerant plants are for example glyphosate-tolerant plants, i.e. plants made tolerant to the herbicide glyphosate or salts thereof. Plants can be made tolerant to glyphosate through different means. For example, glyphosate-tolerant plants can be obtained by transforming the plant with a gene encoding the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are the AroA gene (mutant CT7) of the bacterium *Salmonella typhimurium* (Comai et al., Science (1983), 221, 370-371), the CP4 gene of the

bacterium *Agrobacterium* sp. (Barry et al., Curr. Topics Plant Physiol. (1992), 7, 139-145), the genes encoding a *Petunia* EPSPS (Shah et al., Science (1986), 233, 478-481), a *Tomato* EPSPS (Gasser et al., J. Biol. Chem. (1988), 263, 4280-4289), or an *Eleusine* EPSPS (WO 2001/66704). It can also be a mutated EPSPS as described in for example EP-A 0837944, WO 2000/066746, WO 2000/066747 or WO 2002/026995. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate oxido-reductase enzyme as described in US 5,776,760 and US 5,463,175. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate acetyl transferase enzyme as described in for example WO 2002/036782, WO 2003/092360, WO 2005/012515 and WO 2007/024782. Glyphosate-tolerant plants can also be obtained by selecting plants containing naturally-occurring mutations of the above-mentioned genes, as described in for example WO 2001/024615 or WO 2003/013226.

Other herbicide resistant plants are for example plants that are made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition. One such efficient detoxifying enzyme is an enzyme encoding a phosphinothricin acetyltransferase (such as the bar or pat protein from *Streptomyces* species). Plants expressing an exogenous phosphinothricin acetyltransferase are for example described in US 5,561,236; US 5,648,477; US 5,646,024; US 5,273,894; US 5,637,489; US 5,276,268; US 5,739,082; US 5,908,810 and US 7,112,665.

Further herbicide-tolerant plants are also plants that are made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvate dioxygenase (HPPD). Hydroxyphenylpyruvate dioxygenases are enzymes that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is transformed into homogentisate. Plants tolerant to HPPD-inhibitors can be transformed with a gene encoding a naturally-occurring resistant HPPD enzyme, or a gene encoding a mutated HPPD enzyme as described in WO 1996/038567, WO 1999/024585 and WO 1999/024586. Tolerance to HPPD-inhibitors can also be obtained by transforming plants with genes encoding certain enzymes enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD-inhibitor. Such plants and genes are described in WO 1999/034008 and WO 2002/36787. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an enzyme prephenate dehydrogenase in addition to a gene encoding an HPPD-tolerant enzyme, as described in WO 2004/024928.

Still further herbicide resistant plants are plants that are made tolerant to acetolactate synthase (ALS) inhibitors. Known ALS-inhibitors include, for example, sulfonylurea, imidazolinone,

triazolopyrimidines, pyrimidinyloxy(thio)benzoates, and/or sulfonylaminocarbonyltriazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxyacid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides, as described for example in Tranel and Wright, *Weed Science* (2002), 50, 700-712, but also, in US 5,605,011, US 5,378,824, US 5,141,870, and US 5,013,659. The production of sulfonylurea-tolerant plants and imidazolinone-tolerant plants is described in US 5,605,011; US 5,013,659; US 5,141,870; US 5,767,361; US 5,731,180; US 5,304,732; US 4,761,373; US 5,331,107; US 5,928,937; and US 5,378,824; and international publication WO 1996/033270. Other imidazolinone-tolerant plants are also described in for example WO 2004/040012, WO 2004/106529, WO 2005/020673, WO 2005/093093, WO 2006/007373, WO 2006/015376, WO 2006/024351, and WO 2006/060634. Further sulfonylurea- and imidazolinone-tolerant plants are also described in for example WO 2007/024782.

Other plants tolerant to imidazolinone and/or sulfonylurea can be obtained by induced mutagenesis, selection in cell cultures in the presence of the herbicide or mutation breeding as described for example for soybeans in US 5,084,082, for rice in WO 1997/41218, for sugar beet in US 5,773,702 and WO 1999/057965, for lettuce in US 5,198,599, or for sunflower in WO 2001/065922.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are insect-resistant transgenic plants, i.e. plants made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such insect resistance. An "insect-resistant transgenic plant", as used herein, includes any plant containing at least one transgene comprising a coding sequence encoding:

1) an insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins listed by Crickmore et al., *Microbiology and Molecular Biology Reviews* (1998), 62, 807-813, updated by Crickmore et al. (2005) at the *Bacillus thuringiensis* toxin nomenclature, online at:

http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/), or insecticidal portions thereof, e.g., proteins of the Cry protein classes Cry1Ab, Cry1Ac, Cry1F, Cry2Ab, Cry3Aa, or Cry3Bb or insecticidal portions thereof; or

2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein from *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cry34 and Cry35 crystal proteins (Moellenbeck et al., *Nat. Biotechnol.* (2001), 19, 668-72; Schnepf et al., *Applied Environm. Microbiol.* (2006), 71, 1765-1774); or

3) a hybrid insecticidal protein comprising parts of different insecticidal crystal proteins from *Bacillus thuringiensis*, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2) above, e.g., the Cry1A.105 protein produced by corn event MON98034 (WO 2007/027777); or

5 4) a protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in corn events MON863 or MON88017, or the Cry3A protein in corn event
10 MIR604;

5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal (VIP) proteins listed at: http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html, e.g., proteins from the VIP3Aa protein class; or

15 6) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B. cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins (WO 1994/21795); or

7) a hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the
20 proteins in 2) above; or

8) a protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding
25 an insecticidal protein), such as the VIP3Aa protein in cotton event COT102.

Of course, an insect-resistant transgenic plant, as used herein, also includes any plant comprising a combination of genes encoding the proteins of any one of the above classes 1 to 8. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the above classes 1 to 8, to expand the range of target insect species
30 affected when using different proteins directed at different target insect species, or to delay insect resistance development to the plants by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering)
35 which may also be treated according to the invention are tolerant to abiotic stresses. Such

plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress tolerance plants include:

plants which contain a transgene capable of reducing the expression and/or the activity of poly(ADP-ribose)polymerase (PARP) gene in the plant cells or plants as described in WO
5 2000/004173 or WO2006/045633 or PCT/EP07/004142.

plants which contain a stress tolerance enhancing transgene capable of reducing the expression and/or the activity of the PARP encoding genes of the plants or plants cells, as described e.g. in WO 2004/090140.

plants which contain a stress tolerance enhancing transgene coding for a plant-functional
10 enzyme of the nicotinamide adenine dinucleotide salvage synthesis pathway including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenylyl transferase, nicotinamide adenine dinucleotide synthetase or nicotine amide phosphoribosyltransferase as described e.g. in WO2006/032469 or WO 2006/133827 or PCT/EP07/002433.

15 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention show altered quantity, quality and/or storage-stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as :

1) transgenic plants which synthesize a modified starch, which in its physical-chemical
20 characteristics, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the side chain distribution, the viscosity behaviour, the gelling strength, the starch grain size and/or the starch grain morphology, is changed in comparison with the synthesised starch in wild type plant cells or plants, so that this is better suited for special applications. Said transgenic plants synthesizing a modified starch are
25 disclosed, for example, in EP 0571427, WO 1995/004826, EP 0719338, WO 1996/15248, WO 1996/19581, WO 1996/27674, WO 1997/11188, WO 1997/26362, WO 1997/32985, WO 1997/42328, WO 1997/44472, WO 1997/45545, WO 1998/27212, WO 1998/40503, WO99/58688, WO 1999/58690, WO 1999/58654, WO 2000/008184, WO 2000/008185, WO 2000/008175, WO 2000/28052, WO 2000/77229, WO 2001/12782, WO 2001/12826, WO
30 2002/101059, WO 2003/071860, WO 2004/056999, WO 2005/030942, WO 2005/030941, WO 2005/095632, WO 2005/095617, WO 2005/095619, WO 2005/095618, WO 2005/123927, WO 2006/018319, WO 2006/103107, WO 2006/108702, WO 2007/009823, WO 2000/22140, WO 2006/063862, WO 2006/072603, WO 2002/034923, EP 06090134.5, EP 06090228.5, EP 06090227.7, EP 07090007.1, EP 07090009.7, WO 2001/14569, WO 2002/79410, WO
35 2003/33540, WO 2004/078983, WO 2001/19975, WO 1995/26407, WO 1996/34968, WO

1998/20145, WO 1999/12950, WO 1999/66050, WO 1999/53072, US 6,734,341, WO 2000/11192, WO 1998/22604, WO 1998/32326, WO 2001/98509, WO 2001/98509, WO 2005/002359, US 5,824,790, US 6,013,861, WO 1994/004693, WO 1994/009144, WO 1994/11520, WO 1995/35026, WO 1997/20936.

5 2) transgenic plants which synthesize non starch carbohydrate polymers or which synthesize non starch carbohydrate polymers with altered properties in comparison to wild type plants without genetic modification. Examples are plants producing polyfructose, especially of the inulin and levan-type, as disclosed in EP 0663956, WO 1996/001904, WO 1996/021023, WO 1998/039460, and WO 1999/024593, plants producing alpha 1,4 glucans as disclosed in WO
10 1995/031553, US 2002/031826, US 6,284,479, US 5,712,107, WO 1997/047806, WO 1997/047807, WO 1997/047808 and WO 2000/014249, plants producing alpha-1,6 branched alpha-1,4-glucans, as disclosed in WO 2000/73422, plants producing alternan, as disclosed in WO 2000/047727, EP 06077301.7, US 5,908,975 and EP 0728213,

3) transgenic plants which produce hyaluronan, as for example disclosed in WO 2006/032538,
15 WO 2007/039314, WO 2007/039315, WO 2007/039316, JP 2006/304779, and WO 2005/012529.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as cotton plants, with altered fiber characteristics. Such plants can be obtained by genetic transformation,
20 or by selection of plants contain a mutation imparting such altered fiber characteristics and include:

Plants, such as cotton plants, containing an altered form of cellulose synthase genes as described in WO 1998/000549

Plants, such as cotton plants, containing an altered form of rsw2 or rsw3 homologous nucleic
25 acids as described in WO2004/053219

Plants, such as cotton plants, with increased expression of sucrose phosphate synthase as described in WO 2001/017333

Plants, such as cotton plants, with increased expression of sucrose synthase as described in WO02/45485

30 Plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fiber cell is altered, e.g. through downregulation of fiberselective β 1,3-glucanase as described in WO2005/017157

Plants, such as cotton plants, having fibers with altered reactivity, e.g. through the expression of N-acetylglucosaminetransferase gene including nodC and chitinsynthase genes as described in
35 WO2006/136351

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be obtained by genetic transformation or by selection of plants contain a mutation imparting such

5 altered oil characteristics and include:

Plants, such as oilseed rape plants, producing oil having a high oleic acid content as described e.g. in US 5,969,169, US 5,840,946 or US 6,323,392 or US 6,063,947

Plants such as oilseed rape plants, producing oil having a low linolenic acid content as described in US 6,270,828, US 6,169,190 or US 5,965,755

10 Plant such as oilseed rape plants, producing oil having a low level of saturated fatty acids as described e.g. in US 5,434,283

Particularly useful transgenic plants which may be treated according to the invention are plants which comprise one or more genes which encode one or more toxins, such as the following which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans),
 15 KnockOut® (for example maize), BiteGard® (for example maize), Bt-Xtra® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucotn® (cotton), Nucotn 33B®(cotton), NatureGard® (for example maize), Protecta® and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for
 20 example maize, cotton, soya bean), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned include the varieties sold under the name Clearfield® (for example maize).

25 Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or combination of transformation events, that are listed for example in the databases from various national or regional regulatory agencies (see for example http://gmoinfo.jrc.it/gmp_browse.aspx and <http://www.agbios.com/dbase.php>).

30

Among the diseases of plants or crops that can be controlled by the method according to the present invention, mention may be made of :

Powdery mildew diseases such as :

Blumeria diseases, caused for example by *Blumeria graminis*;

35 Leveillula diseases, caused for example by *Leveillula taurica*

Podosphaera diseases, caused for example by *Podosphaera leucotricha*;

Sphaerotheca diseases, caused for example by *Sphaerotheca fuliginea* or *Sphaerotheca pannosa*;

Uncinula diseases, caused for example by *Uncinula necator*;

5 Rust diseases such as :

Gymnosporangium diseases, caused for example by *Gymnosporangium sabinae*;

Hemileia diseases, caused for example by *Hemileia vastatrix*;

Phakopsora diseases, caused for example by *Phakopsora pachyrhizi* or *Phakopsora meibomia*;

10 Puccinia diseases, caused for example by *Puccinia recondita*;

Uromyces diseases, caused for example by *Uromyces appendiculatus*;

Oomycete diseases such as :

Bremia diseases, caused for example by *Bremia lactucae*;

Peronospora diseases, caused for example by *Peronospora pisi* or *P. brassicae*;

15 Phytophthora diseases, caused for example by *Phytophthora infestans*;

Plasmopara diseases, caused for example by *Plasmopara viticola*;

Pseudoperonospora diseases, caused for example by *Pseudoperonospora humuli* or *Pseudoperonospora cubensis*;

Pythium diseases, caused for example by *Pythium ultimum*;

20 Leafspot, leaf blotch and leaf blight diseases such as :

Alternaria diseases, caused for example by *Alternaria solani*;

Cercospora diseases, caused for example by *Cercospora beticola*;

Cladosporium diseases, caused for example by *Cladosporium cucumerinum*;

Cochliobolus diseases, caused for example by *Cochliobolus sativus*;

25 Colletotrichum diseases, caused for example by *Colletotrichum lindemuthianum*;

Cycloconium diseases, caused for example by *Cycloconium oleaginum*;

Diaporthe diseases, caused for example by *Diaporthe citri*;

Diplocarpon diseases, caused for example by *Diplocarpon rosae*

Elsinoe diseases, caused for example by *Elsinoe fawcettii*;

30 Gloeosporium diseases, caused for example by *Gloeosporium laeticolor*;

Glomerella diseases, caused for example by *Glomerella cingulata*;

Guignardia diseases, caused for example by *Guignardia bidwelli*;

Leptosphaeria diseases, caused for example by *Leptosphaeria maculans*; *Leptosphaeria nodorum*;

35 Magnaporthe diseases, caused for example by *Magnaporthe grisea*;

Mycosphaerella diseases, caused for example by *Mycosphaerella graminicola*; *Mycosphaerella arachidicola*; *Mycosphaerella fijiensis*;

Phaeosphaeria diseases, caused for example by *Phaeosphaeria nodorum*;

Pyrenophora diseases, caused for example by *Pyrenophora teres*;

5 Ramularia diseases, caused for example by *Ramularia collo-cygni*;

Rhynchosporium diseases, caused for example by *Rhynchosporium secalis*;

Septoria diseases, caused for example by *Septoria apii* or *Septoria lycopersici*;

Typhula diseases, caused for example by *Typhula incarnata*;

Venturia diseases, caused for example by *Venturia inaequalis*;

10 Root and stem diseases such as :

Corticium diseases, caused for example by *Corticium graminearum*;

Fusarium diseases, caused for example by *Fusarium oxysporum*;

Gaeumannomyces diseases, caused for example by *Gaeumannomyces graminis*;

Rhizoctonia diseases, caused for example by *Rhizoctonia solani*;

15 Tapesia diseases, caused for example by *Tapesia acuformis*;

Thielaviopsis diseases, caused for example by *Thielaviopsis basicola*;

Ear and panicle diseases such as :

Alternaria diseases, caused for example by *Alternaria spp.*;

Aspergillus diseases, caused for example by *Aspergillus flavus*;

20 Cladosporium diseases, caused for example by *Cladosporium spp.*;

Claviceps diseases, caused for example by *Claviceps purpurea*;

Fusarium diseases, caused for example by *Fusarium culmorum*;

Gibberella diseases, caused for example by *Gibberella zeae*;

Monographella diseases, caused for example by *Monographella nivalis*;

25 Smut and bunt diseases such as :

Sphacelotheca diseases, caused for example by *Sphacelotheca reiliana*;

Tilletia diseases, caused for example by *Tilletia caries*;

Urocystis diseases, caused for example by *Urocystis occulta*;

Ustilago diseases, caused for example by *Ustilago nuda*;

30 Fruit rot and mould diseases such as :

Aspergillus diseases, caused for example by *Aspergillus flavus*;

Botrytis diseases, caused for example by *Botrytis cinerea*;

Penicillium diseases, caused for example by *Penicillium expansum*;

Sclerotinia diseases, caused for example by *Sclerotinia sclerotiorum*;

35 Verticillium diseases, caused for example by *Verticillium albo-atrum*;

Seed and soilborne decay, mould, wilt, rot and damping-off diseases :

Fusarium diseases, caused for example by *Fusarium culmorum*;

Phytophthora diseases, caused for example by *Phytophthora cactorum*;

Pythium diseases, caused for example by *Pythium ultimum*;

5 Rhizoctonia diseases, caused for example by *Rhizoctonia solani*;

Sclerotium diseases, caused for example by *Sclerotium rolfsii*;

Microdochium diseases, caused for example by *Microdochium nivale*;

Canker, broom and dieback diseases such as :

Nectria diseases, caused for example by *Nectria galligena*;

10 Blight diseases such as :

Monilinia diseases, caused for example by *Monilinia laxa*;

Leaf blister or leaf curl diseases such as :

Taphrina diseases, caused for example by *Taphrina deformans*;

Decline diseases of wooden plants such as :

15 Esca diseases, caused for example by *Phaemoniella clamydospora*;

Diseases of flowers and Seeds such as :

Botrytis diseases, caused for example by *Botrytis cinerea*;

Diseases of tubers such as :

Rhizoctonia diseases, caused for example by *Rhizoctonia solani*;

20 Helminthosporium diseases, caused for example by *Helminthosporium solani*.

Furthermore the treatments according to the invention can be able to reduce the contents of mycotoxins in the harvested crops and therefore in foods and animal feed stuff made therefrom.

Especially but not exclusively the following mycotoxins can be specified:

25 Deoxynivalenole (DON), Nivalenole, 15-Ac-DON, 3-Ac-DON, T2- und HT2- Toxins, Fumonisin, Zearalenone Moniliformine, Fusarine, Diacetoxyscirpenole (DAS), Beauvericine, Enniatine, Fusaroproliferine, Fusarenole, Ochratoxines, Patuline, Ergocalciferol und Aflatoxines, which are caused for example by the following fungal diseases: *Fusarium spec.*, like *Fusarium acuminatum*, *F. avenaceum*, *F. crookwellense*, *F. culmorum*, *F. graminearum* (30 *Gibberella zeae*), *F. equiseti*, *F. fujikuroi*, *F. musarum*, *F. oxysporum*, *F. proliferatum*, *F. poae*, *F. pseudograminearum*, *F. sambucinum*, *F. scirpi*, *F. semitectum*, *F. solani*, *F. sporotrichoides*, *F. langsethiae*, *F. subglutinans*, *F. tricinctum*, *F. verticillioides* and others but also by *Aspergillus spec.*, *Penicillium spec.*, *Claviceps purpurea*, *Stachybotrys spec.* and others.

The composition according to the present invention is well tolerated by plants, have favourable homeotherm toxicity and are environmentally friendly; it is suitable for protecting plants and plant organs, for increasing harvest yields, for improving the quality of the harvested material and for controlling animal pests, in particular insects, arachnids and nematodes encountered in
 5 agriculture, in forests, in gardens and leisure facilities, in the protection of stored products and materials and in the hygiene sector. It is preferably used as crop protection agents. It is active against normally sensitive and resistant species and against all or some stages of development. Among the animal pests that can also be controlled by the method according to the present invention, mention may be made of :

- 10 Pest from the order of the Isopoda, for example *Oniscus asellus*, *Armadillidium vulgare* and *Porcellio scaber*;
- Pest from the order of the Diplopoda, for example *Blaniulus guttulatus*;
- Pest from the order of the Chilopoda, for example *Geophilus carpophagus* and *Scutigera* spp.;
- Pest from the order of the Symphyla, for example *Scutigera* spp.;
- 15 Pest from the order of the Thysanura, for example *Lepisma saccharina*;
- Pest from the order of the Collembola, for example *Onychiurus armatus*;
- Pest from the order of the Orthoptera, for example *Acheta domesticus*, *Gryllotalpa* spp., *Locusta migratoria migratorioides*, *Melanoplus* spp. and *Schistocerca gregaria*;
- Pest from the order of the Blattaria, for example *Blatta orientalis*, *Periplaneta americana*,
 20 *Leucophaea maderae* and *Blattella germanica*;
- Pest from the order of the Dermaptera, for example *Forficula auricularia*;
- Pest from the order of the Isoptera, for example *Reticulitermes* spp.;
- Pest from the order of the Phthiraptera, for example *Pediculus humanus corporis*, *Haematopinus* spp., *Linognathus* spp., *Trichodectes* spp., *Damalinia* spp.;
- 25 Pest from the order of the Thysanoptera, for example *Hercinothrips femoralis*, *Thrips tabaci*, *Thrips palmi*, *Frankliniella accidentalis*;
- Pest from the order of the Heteroptera, for example *Eurygaster* spp., *Dysdercus intermedius*, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus* and *Triatoma* spp.;
- Pest from the order of the Homoptera, for example *Aleurodes brassicae*, *Bemisia tabaci*,
 30 *Trialeurodes vaporariorum*, *Aphis gossypii*, *Brevicoryne brassicae*, *Cryptomyzus ribis*, *Aphis fabae*, *Aphis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*, *Phylloxera vastatrix*, *Pemphigus* spp., *Macrosiphum avenae*, *Myzus* spp., *Phorodon humuli*, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelis bilobatus*, *Nephotettix cincticeps*, *Lecanium corni*, *Saissetia oleae*, *Laodelphax striatellus*, *Nilaparvata lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp. and *Psylla* spp.;

Pest from the order of the Lepidoptera, for example *Pectinophora gossypiella*, *Bupalus piniarius*, *Cheimatobia brumata*, *Lithocolletis blancardella*, *Hyponomeuta padella*, *Plutella xylostella*, *Malacosoma neustria*, *Euproctis chrysorrhoea*, *Lymantria* spp., *Bucculatrix thurberiella*, *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp., *Earias insulana*, *Heliothis* spp.,
 5 *Mamestra brassicae*, *Panolis flammea*, *Spodoptera* spp., *Trichoplusia ni*, *Carpocapsa pomonella*, *Pieris* spp., *Chilo* spp., *Pyrausta nubilalis*, *Ephestia kuehniella*, *Galleria mellonella*, *Tineola bisselliella*, *Tinea pellionella*, *Hofmannophila pseudospretella*, *Cacoecia podana*, *Capua reticulana*, *Choristoneura fumiferana*, *Clysia ambiguella*, *Homona magnanima*, *Tortrix viridana*, *Cnaphalocerus* spp. and *Oulema oryzae*;

10 Pest from the order of the Coleoptera, for example *Anobium punctatum*, *Rhizopertha dominica*, *Bruchidius obtectus*, *Acanthoscelides obtectus*, *Hylotrupes bajulus*, *Agelastica alni*, *Leptinotarsa decemlineata*, *Phaedon cochleariae*, *Diabrotica* spp., *Psylliodes chrysocephala*, *Epilachna varivestis*, *Atomaria* spp., *Oryzaephilus surinamensis*, *Anthonomus* spp., *Sitophilus* spp., *Otiorrhynchus sulcatus*, *Cosmopolites sordidus*, *Ceuthorrhynchus assimilis*, *Hypera postica*,
 15 *Dermestes* spp., *Trogoderma* spp., *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*, *Ptinus* spp., *Niptus hololeucus*, *Gibbium psyllodes*, *Tribolium* spp., *Tenebrio molitor*, *Agriotes* spp., *Conoderus* spp., *Melolontha melolontha*, *Amphimallon solstitialis*, *Costelytra zealandica* and *Lissorhoptrus oryzophilus*;

Pest from the order of the Hymenoptera, for example *Diprion* spp., *Hoplocampa* spp., *Lasius* spp.,
 20 *Monomorium pharaonis* and *Vespa* spp.;

Pest from the order of the Diptera, for example *Aedes* spp., *Anopheles* spp., *Culex* spp., *Drosophila melanogaster*, *Musca* spp., *Fannia* spp., *Calliphora erythrocephala*, *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp., *Gastrophilus* spp., *Hyppobosca* spp., *Stomoxys* spp., *Oestrus* spp., *Hypoderma* spp., *Tabanus* spp., *Tannia* spp., *Bibio hortulanus*, *Oscinella frit*, *Phorbia* spp., *Pegomyia*
 25 *hyoscyami*, *Ceratitis capitata*, *Dacus oleae*, *Tipula paludosa*, *Hylemyia* spp. and *Liriomyza* spp.;

Pest from the order of the Siphonaptera, for example *Xenopsylla cheopis* and *Ceratophyllus* spp.;

Pest from the class of the Arachnida, for example *Scorpio maurus*, *Latrodectus mactans*, *Acarus siro*, *Argas* spp., *Ornithodoros* spp., *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptruta oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes*
 30 spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp., *Tetranychus* spp., *Hemitarsonemus* spp. and *Brevipalpus* spp.;

The plant-parasitic nematodes such as *Pratylenchus* spp., *Radopholus similis*, *Ditylenchus dipsaci*, *Tylenchulus semipenetrans*, *Heterodera* spp., *Globodera* spp., *Meloidogyne* spp., *Aphelenchoides* spp., *Longidorus* spp., *Xiphinema* spp., *Trichodorus* spp. and *Bursaphelenchus*
 35 spp.

The composition according to the present invention may also be used against pests and diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 800 g/ha, preferably between 50 and 300 g/ha for applications in foliar treatment. If a drench/drip application is possible, the dose can be lower, especially in artificial substrates like rockwool or perlite. The dose of active substance applied is generally and advantageously between 2 and 200 g per 100 kg of seed, preferably between 3 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to adapt the application doses according to the nature of the crop to be treated.

The composition according to the present invention may also be used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into whose genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

Biological Examples

Formula for the efficacy of the combination of two compounds

The expected efficacy of a given combination of two compounds is calculated as follows (see Colby, S.R., „Calculating Synergistic and antagonistic Responses of Herbicide Combinations“, Weeds 15, pp. 20-22, 1967):

If

X is the efficacy expressed in % mortality of the untreated control for test compound A at a concentration of m ppm respectively m g/ha,

Y is the efficacy expressed in % mortality of the untreated control for test compound B at a concentration of n ppm respectively n g/ha,

E is the efficacy expressed in % mortality of the untreated control using the mixture of A and B at m and n ppm respectively m and n g /ha,

$$E = X + Y - \frac{X \times Y}{100}$$

If the observed insecticidal efficacy of the combination is higher than the one calculated as „E“, then the combination of the two compounds is more than additive, i.e., there is a synergistic effect.

Example A

Plutella xylostella - test

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycoether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with larvae of the diamond back moth (*Plutella xylostella*) as long as the leaves are still moist.

After the specified period of time, the mortality in % is determined. 100 % means that all the caterpillars have been killed; 0 % means that none of the caterpillars have been killed.

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table A plant damaging insects Plutella xylostella - test		
Active Ingredient	Concentration in ppm	Efficacy in % after 3d
Fluopicolide	200	0
Clothianidin	4	0
Fluopicolide + Clothianidin (5 : 1) according to the invention	200 + 4	obs.* cal.** 30 0

* obs. = observed insecticidal efficacy

** cal. = efficacy calculated with Colby-formula

Example B**Spodoptera exigua - test**

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycolether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with larvae of the beet army worm (*Spodoptera exigua*) as long as the leaves are still moist.

After the specified period of time, the mortality in % is determined. 100 % means that all the caterpillars have been killed; 0 % means that none of the caterpillars have been killed.

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table A plant damaging insects Spodoptera exigua - test		
Active Ingredient	Concentration in ppm	Efficacy in % after 3d
Fluopicolide	500	0
Imidacloprid	500	40
Fluopicolide + Imidacloprid (1 : 1) according to the invention	500 + 500	obs.* cal.** 65 40

* obs. = observed insecticidal efficacy

** cal. = efficacy calculated with Colby-formula

Example B**Spodoptera exigua - test**

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycoether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with larvae of the beet army worm (*Spodoptera exigua*) as long as the leaves are still moist.

After the specified period of time, the mortality in % is determined. 100 % means that all the caterpillars have been killed; 0 % means that none of the caterpillars have been killed.

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table B plant damaging insects Spodoptera exigua - test		
Active Ingredient	Concentration in ppm	Efficacy in % after 3d
Fluopicolide	500	0
Imidacloprid	500	40
Fluopicolide + Imidacloprid (1 : 1) according to the invention	500 + 500	obs.* cal.** 65 40

*obs. = observed insecticidal efficacy

** cal. = efficacy calculated with Colby-formula

Example C**Spodoptera frugiperda - test**

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycolether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

10 Cabbage leaves (*Brassica oleracea*) are treated by being dipped into the preparation of the active compound of the desired concentration and are infested with larvae of the fall army worm (*Spodoptera frugiperda*) as long as the leaves are still moist.

After the specified period of time, the mortality in % is determined. 100 % means that all the caterpillars have been killed; 0 % means that none of the caterpillars have been killed.

According to the present application in this test e.g. the following combinations show a synergistic effect in comparison to the single compounds:

Table C plant damaging insects Spodoptera frugiperda - test		
Active Ingredient	Concentration in ppm	Efficacy in % after 4d
Fluopicolide	500	0
Thiacloprid	20	25
Fluopicolide + Thiacloprid (25 : 1) according to the invention	500 + 20	obs.* cal.** 50 25

* obs. = observed insecticidal efficacy

** cal. = efficacy calculated with Colby-formula

Table D plant damaging insects Spodoptera frugiperda - test		
Active Ingredient	Concentration in ppm	Efficacy in % after 6d
Fluopicolide	200	0

Thiamethoxam	100	0
Fluopicolide + Thiamethoxam (2 : 1) according to the invention	200 + 100	obs.* cal.** 25 0

* obs. = observed insecticidal efficacy

** cal. = efficacy calculated with Colby-formula


30725-978

44

CLAIMS:

1. A composition comprising:

(A) 2,6-dichloro-N-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl} benzamide (fluopicolide) of formula (Ia):



5

or agriculturally acceptable optical and/or geometric isomers, tautomers or addition salts with an acid or a base; and

(B) an insecticide compound selected from the group consisting of:

(b1) an acetylcholine receptor agonist/antagonist,

10 (b2) an acetylcholinesterase (AChE) inhibitor,

(b3) a sodium channel modulator/voltage-gated sodium channel blocker,

(b5) a GABA-gated chloride channel antagonist,

(b16) an inhibitor of lipid synthesis,

(b17) a carboxamide, and

15 (b20) rynaxapyr,

in a (A) / (B) weight ratio of from 1/1000 to 1000/1.

2. The composition according to claim 1, wherein:

(b1) is a chloronicotinyl, a neonicotinoid, nicotine, bensultap or cartap,

30725-978

45

(b2) is a carbamate or an organophosphate,

(b3) is a pyrethroid or an oxadiazine,

(b5) is a cyclodiene organochlorine or a fiprole,

(b16) is a tetrionic acid, or a tetramic acid, and

5 (b17) is flonicamid.

3. A composition according to claim 1, wherein the insecticide compound (B) is selected from the group consisting of acephate, acetoprole, acrinathrin, alanycarb, aldicarb, aldoxycarb, allethrin (d-cis-trans, d-trans), allyxycarb, aminocarb, acetamiprid, azamethiphos, azinphos (-methyl, -ethyl), bendiocarb, benfuracarb, bensultap, beta-cyfluthrin, bifenthrin,
 10 bioallethrin, bioallethrin-S-cyclopentyl-isomer, bioethanomethrin, biopermethrin, bioresmethrin, bromophos-ethyl, bromfenvinfos (-methyl), bufencarb, butacarb, butocarboxim, butoxycarboxim, butathiofos, carbaryl, carbofuran, carbonic acid, carbosulfan, cadusafos, camphechlor, carbophenothion, cartap, chloethocarb, chlordane, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos (-methyl/-ethyl), chlovaporthrin, cis-
 15 cypermethrin, cis-resmethrin, cis-permethrin, clocythrins, clothianidin, coumaphos, cyanofenphos, cyanophos, cycloprothrin, cyfluthrin, cyhalothrin, cypermethrin (alpha-, beta-, theta-, zeta-), cyphenothrin, DDT, deltamethrin, demeton-S-methyl, demeton-S-methylsulphon, dialifos, diazinon, dichlofenthion, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, dimetilan, dinotefuran, dioxabenzofos, disulfoton, empenthrin (1R-isomer),
 20 endosulfan, EPN, esfenvalerate, ethiofencarb, ethion, ethiprole, ethoprophos, etofenprox, etrimfos, famphur, fenamiphos, fenfluthrin, fenitrothion, fenobucarb, fenothiocarb, fenpropathrin, fenpyrithrin, fensulfothion, fenthion, fenvalerate, fipronil, flonicamid, flubrocycythrinate, flubrocycythrinate, flucythrinate, flufenprox, flumethrin, flupyrazofos, fluvalinate, fonofos, formetanate, formothion, fosmethilan, fosthiazate, fubfenprox,
 25 furathiocarb, gamma-cyhalothrin, gamma-HCH, HCH, heptachlor, heptenophos, imidacloprid, imidaclothiz, imiprothrin, indoxacarb, iodofenphos, iprobenfos, isazofos, isofenphos, isoprocab, isopropyl O-salicylate, isoxathion, kadethrin, lambda-cyhalothrin, lindane, malathion, mecarbam, metam-sodium, methacrifos, methamidophos, methidathion,

30725-978

46

methiocarb, methomyl, methoxychlor, metofluthrin, metolcarb, mevinphos, monocrotophos, naled, nicotine, nitenpyram, nithiazine, omethoate, oxamyl, oxydemeton-methyl, parathion (-methyl/-ethyl), permethrin (cis-, trans-), phenothrin (1R-trans isomer), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, pirimicarb, pirimiphos (-methyl/-ethyl), prallethrin, profenofos, profluthrin, promecarb, propaphos, propetamphos, propoxur, prothiofos, prothoate, protrifenbute, pyraclofos, pyresmethrin, pyrethrins (pyrethrum), pyridaphenthion, pyridathion, quinalphos, resmethrin, RU 15525, rynaxapyr, sebufos, silafluofen, spiroadiclofen, spiromesifen, spirotetramate, sulfotep, sulprofos, tau-fluvalinate, tebupirimfos, tefluthrin, temephos, terallethrin, terbufos, tetrachlorvinphos, tetramethrin (1R-isomer), thiacloprid, thiodicarb, thiofanox, traloccythrin, tralomethrin, transfluthrin, trimethacarb, thiamethoxam, thiometon, triazamate, triazophos, triclofon, vamidothion, vaniliprole, XMC, xylylcarb, ZXI 8901 and (2E)-1-[(2-chloro-1,3-thiazol-5-yl)methyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine.

4. The composition according to claim 1, wherein the insecticide compound (B) is selected from the group consisting of acephate, acetamiprid, acrinathrin, aldicarb, alpha-cypermethrin, beta-cyfluthrin, bifenthrin, carbaryl, carbofuran, chlorpyrifos-E, clothianidin, cyfluthrin, cypermethrin, deltamethrin, dinotefuran, ethiprole, fipronil, flonicamid, gamma-cyhalothrin, imidacloprid, indoxacarb, L-cyhalothrin, methamidophos, methiocarb, methomyl, nitenpyram, profenofos, rynaxapyr, spinosad, spiroadiclofen, spiromesifen, spirotetramate, tebupirimphos, tefluthrin, thiacloprid, thiamethoxam, thiodicarb and triazophos.

5. The composition according to claim 1, wherein the insecticide compound (B) is selected from the group consisting of clothianidin, imidacloprid, thiacloprid and thiamethoxam.

6. The composition according to claim 1, wherein the insecticide compound (B) is methiocarb.

7. The composition according to claim 1, wherein the insecticide compound (B) is selected from the group consisting of deltamethrin and tefluthrin.

30725-978

47

8. The composition according to claim 1, wherein the insecticide compound (B) is fipronil.
9. The composition according to claim 1, wherein the insecticide compound (B) is spirotetramate.
- 5 10. The composition according to claim 1, wherein the insecticide compound (B) is flonicamid.
11. The composition according to claim 1, wherein the insecticide compound (B) is rynaxapyr.
12. The composition according to any one of claims 1 to 11, further comprising a
10 fungicide compound (C).
13. The composition according to claim 12, wherein the compounds (A), (B) and (C) are present in a weight ratio of (A):(B):(C) of from 1:0.001:0.001 to 1:1000:1000.
14. The composition according to claim 12 or 13, wherein the fungicide compound (C) is selected from the group consisting of 5-fluoro-1,3-dimethyl-N-[2-(1,3-
15 dimethyl)-but-2-ol-yl]-1H-pyrazole-4-carboxamide, benalaxyl, benalaxyl-M, benthiavalicarb, carboxin, chlorothalonil, cyazofamid, cymoxanil, dimetomorph, fluazinam, fludioxonil, fluquinconazole, fluoxastrobin, flutriafol, fosetyl-aluminium, hexaconazole, hymexazole, ipconazole, mancozeb, mandipropamid, maneb, mefenoxam, metiram, metalaxyl, metalaxyl-M, peconazole, penthiopyrad, phosphorous acid, propamocarb.HCl, propineb,
20 prothioconazole, tebuconazole, thiram, triadimenol, trifloxystrobin and triticonazole.
15. The composition according to any one of claims 1 to 14, further comprising an agriculturally acceptable support, carrier, filler, surfactant or a mixture thereof.
16. A method for preventively or curatively controlling phytopathogenic fungi of plants and also curatively or preventively controlling insects, comprising applying a
25 non-phytotoxic amount of the composition according to any one of claims 1 to 15, via seed treatment, foliar application, stem application or drench/drip application (chemigation) to seed

30725-978

48

of the plants, the plants and/or fruit of the plants, or to soil and/or to inert substrate, Pumice, Pyroclastic materials/tuff, synthetic organic substrates, organic substrates and/or to a liquid substrate in which the plants are growing or in which it is desired to grow the plants.

