



(86) Date de dépôt PCT/PCT Filing Date: 2006/12/08
 (87) Date publication PCT/PCT Publication Date: 2007/06/14
 (45) Date de délivrance/Issue Date: 2017/02/07
 (85) Entrée phase nationale/National Entry: 2008/06/06
 (86) N° demande PCT/PCT Application No.: AU 2006/001869
 (87) N° publication PCT/PCT Publication No.: 2007/065226
 (30) Priorité/Priority: 2005/12/09 (AU2005906947)

(51) Cl.Int./Int.Cl. *A61K 38/17* (2006.01),
A61K 38/04 (2006.01), *A61P 9/00* (2006.01),
A61P 9/04 (2006.01)
 (72) Inventeur/Inventor:
 DUGGAN, KAREN ANNETTE, AU
 (73) Propriétaire/Owner:
 VECTUS BIOSYSTEMS LIMITED, AU
 (74) Agent: BLAKE, CASSELS & GRAYDON LLP

(54) Titre : FRAGMENTS DE VIP ET METHODES D'UTILISATION
 (54) Title: VIP FRAGMENTS AND METHODS OF USE

(57) **Abrégé/Abstract:**

The invention relates to composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments, and the use of those compositions in the treatment of fibrosis, hypertension and other disorders.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 June 2007 (14.06.2007)

PCT

(10) International Publication Number
WO 2007/065226 A1

(51) International Patent Classification:

A61K 38/17 (2006.01) A61P 9/00 (2006.01)
A61K 38/04 (2006.01) A61P 9/04 (2006.01)

(21) International Application Number:

PCT/AU2006/001869

(22) International Filing Date:

8 December 2006 (08.12.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2005906947 9 December 2005 (09.12.2005) AU

(71) Applicant (for all designated States except US): **VECTUS BIOSYSTEMS LIMITED** [AU/AU]; 3-11 Primrose Avenue, Rosebery, NSW 2018 (AU).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **DUGGAN, Karen, Annette** [AU/AU]; 3 Soudan Street, Randwick, NSW 2031 (AU).(74) Agent: **SHELSTON IP**; 60 Margaret Street, Sydney, NSW 2000 (AU).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VIP FRAGMENTS AND METHODS OF USE

(57) Abstract: The invention relates to composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments, and the use of those compositions in the treatment of fibrosis, hypertension and other disorders.



WO 2007/065226 A1

“VIP fragments and methods of use ”

TECHNICAL FIELD

This invention relates to compositions and methods for therapeutic or prophylactic treatment of myocardial fibrosis or associated conditions. In particular this invention
5 concerns compositions comprising certain active fragments of VIP and their use in the treatment of myocardial fibrosis.

BACKGROUND

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common
10 general knowledge in the field.

In myocardial fibrosis, heart muscle is replaced by fibrous or scar tissue. This can interfere with the flexibility of the heart muscle. It can lead to a decrease in function and, eventually, to overt heart failure. Use of VIP and certain fragments of VIP in therapeutic and prophylactic treatment of myocardial fibrosis and related conditions was
15 described in PCT/AU2005/000835.

Conventional view of structure/function relationship with respect to VIP activity is that the N-terminal amino acid residues (1-5) are important and necessary for signal delivery once VIP binds to its receptor. Further, there are certain key amino acid residues throughout the VIP molecule, distal to the N-terminus, that are important for
20 receptor binding. This would suggest that fragments of VIP lacking either the N-terminal residues or significant portions that encompass the receptor binding residues would not be fully functional. Earlier studies with two VIP fragments, described in PCT/AU2005/000835, indicate that such fragments of VIP could be active, at least in the treatment of myocardial fibrosis.

25 It is an object of the present invention to identify additional fragments of VIP that are active in the treatment of conditions such as myocardial fibrosis, that would overcome certain disadvantages associated with using the entire VIP molecule or large portions thereof, or at least provide a useful alternative.

SUMMARY OF THE INVENTION

Despite the currently prevailing view, it was surprisingly found that activity of VIP and its fragments in the treatment and/or prevention of myocardial fibrosis is not curtailed by either deletion of the N-terminal residues of VIP or the majority of amino acid residues responsible for receptor binding.

In a broad aspect, the invention relates to compositions for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, such as hypertension, comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments. Most preferably, the (VIP) fragments have as a core sequence amino acid residues 6 to 10 of native VIP, or conservative substitutions thereof. However, they may also have as a core sequence amino acid residues 16 to 25 of native VIP, or conservative substitutions thereof.

Accordingly, in a first aspect the invention provides a composition for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, the composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6 to 10 of native VIP, or conservative substitutions thereof.

The preferred active fragments of VIP are VIP(4-16), VIP(6-16), VIP(6-10), VIP(6-12), VIP(6-20) and VIP(6-24), with the proviso that fragments VIP (1-12) and VIP (6-28) are not included. However, the use of VIP fragments, including VIP (1-12) and VIP (6-28) for the treatment of myocardial fibrosis and other associated conditions such as hypertension are encompassed by the present invention

The pharmaceutically effective amount of an active VIP fragment will vary according to the patient and/or with the severity of the disease or condition. These variables can be ascertained by one skilled in the art by routine experimentation. An appropriate dosage range, as a starting point, can be derived from dosages administered in the animal models described herein, or with reference to PCT/AU2005/000835.

Accordingly, in a second aspect the invention provides a composition for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, the composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 16 to 25 of native VIP, or conservative substitutions thereof.

Accordingly, in a third aspect the invention provides a composition for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, the composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments selected from the group
5 consisting of VIP(4-16), VIP(6-16), VIP(6-10), VIP(6-12), VIP(10-28), VIP(16-28), VIP(6-20) and VIP(6-24), or conservative substitutions thereof.

Accordingly, in a fourth aspect the invention provides a composition for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, the composition comprising a pharmaceutically effective amount of one or more
10 functional vasoactive intestinal peptide (VIP) fragments selected from the group listed in Table 1, or conservative substitutions thereof.

The compositions of the invention may be administered in conjunction with a pharmaceutically acceptable carrier, which may be any of those known in the art or devised hereafter and suitable for the intended use. As well as carriers, the
15 pharmaceutical composition of the invention may include other ingredients, including dyes, preservatives, buffers and anti-oxidants, for example. The compositions of the present invention may also include other active agents useful in the treatment of cardiovascular conditions.

The pharmaceutical composition of the invention may take any suitable form, but
20 is preferably suitable for administration by oral, intravenous, intramuscular or subcuticular routes. Other methods of administration such as patches, snuffs, nasal sprays and the like will be clear to those skilled in the art.

According to a fifth aspect the invention provides a method of therapeutic treatment of myocardial fibrosis or an associated condition in a subject, the method
25 including administering to the subject with myocardial fibrosis or an associated condition, a composition according to the invention.

The compositions of the invention may be used to prevent or slow down progression of established myocardial fibrosis, as well as to reduce the degree of established fibrosis.

30 According to a sixth aspect the invention provides a method of prophylactic treatment of myocardial fibrosis or an associated condition in a subject, the method including administering to the subject at risk of developing myocardial fibrosis or an associated condition, a composition according to the invention.

With respect to prophylactic treatment it will be understood that such a treatment would benefit particularly subjects who are at risk of developing myocardial fibrosis or an associated condition. As an example of subjects in the risk category are those having hypertension, diabetes, myocarditis, ischaemic heart disease, drugs such as daunorubicin and others which are used in cancer chemotherapy, genetic predisposition, other conditions such as Conn's Syndrome and Pheochromocytoma, high salt diet and the like. The prophylactic treatment may be used to prevent or slow down the development of fibrosis in the at risk group. High proportion of subjects may already have signs of early heart failure on echocardiography. For example such signs are present in nearly 80% of patients with hypertension. The incidence in diabetics is even higher.

According to a seventh aspect the invention provides a method of prophylactic or therapeutic treatment of congestive cardiac failure in a subject, the method including administering to the subject at risk of developing congestive cardiac failure, a pharmaceutical composition according to the invention.

According to an eighth aspect the invention provides a method of reducing the levels, inhibiting or reducing the production, of pro-fibrotic mediators in a subject, the method including administering to the subject a composition according to the invention.

According to a ninth aspect the invention provides a method of reducing collagen formation or enhancing collagen degradation in the cardiac muscle of a subject, the method including administering to the subject a composition according to the invention.

According to a tenth aspect the invention provides a method of therapeutic or prophylactic treatment of hypertension, the method including administering to the subject a composition according to the invention.

According to an eleventh aspect the invention provides a method of lowering blood pressure in a subject, the method including administering to a subject a composition according to the invention.

It will be apparent to one skilled in the art that the pattern of use of the compositions of the invention may need to be altered for optimum effect. It may be necessary to take into account the nature of the disease or condition as well as its severity.

The associated conditions, which may be subject to prevention or treatment by the compositions of the invention, may include left ventricular hypertrophy, diastolic

dysfunction, myocarditis, cardiomyopathy, left ventricular dysfunction and congestive cardiac failure (for which myocardial fibrosis may be an underlying pathology). The associated condition may also include conditions which give rise to generation of profibrotic mediators or conditions which predispose a subject to myocardial fibrosis, such as for example hypertension and/or high salt intake, diseases such as diabetes and the like.

Further aspect of the invention includes the use of a composition of the invention in the manufacture of a medicament for the prophylactic or therapeutic treatment of myocardial fibrosis, or an associated condition.

Yet another aspect of the invention provides the use of a composition of the invention in the manufacture of a medicament for the prophylactic or therapeutic treatment of congestive cardiac failure.

In another aspect, the invention relates to a composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments with the proviso that fragments VIP (1-12) and VIP (6-28) are not included.

In another aspect, the invention relates to a composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6-10 or 16-25 of native VIP, or conservative substitutions thereof. Preferably, fragments VIP (1-12) and VIP (6-28) are not included.

Preferably, the VIP fragment is one or more fragments selected from the group consisting of: VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28), VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

In another aspect, the invention relates to a composition for prophylactic or therapeutic treatment of myocardial fibrosis or an associated condition, the composition

comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments wherein the VIP fragment is one or more fragments selected from the group consisting of: VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

In another aspect, the invention relates to a composition for prophylactic or therapeutic treatment of hypertension, the composition comprising a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments wherein the VIP fragment is one or more fragments selected from the group consisting of: VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27) VIP(6-28) VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

Preferably, compositions according to the present invention are administered in conjunction with a pharmaceutically acceptable carrier. They may preferably be administered in conjunction with one or more other active agents useful in the treatment of cardiovascular conditions. They may, for preference, be formulated for administration by oral, intravenous, intramuscular or subcuticular routes.

In another aspect, the invention relates to method of therapeutic or prophylactic treatment of myocardial fibrosis or an associated condition in a subject, the method including administering to a subject a composition including a pharmaceutically

effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6-10 or 16-25 of native VIP, or conservative substitutions thereof.

In another aspect, the invention relates to a method of therapeutic or prophylactic treatment of myocardial fibrosis or an associated condition in a subject, the method including administering to a subject a composition according to the present invention.

In another aspect, the invention relates to a method of therapeutic or prophylactic treatment of myocardial fibrosis or an associated condition in a subject, the method including administering to a subject a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

Preferably, the method is used to prevent or slow down progression of established myocardial fibrosis, or alternatively, to reduce the degree of established fibrosis.

Preferably the associated condition is hypertension, diabetes, myocarditis, ischaemic heart disease, left ventricular hypertrophy, diastolic dysfunction, myocarditis, cardiomyopathy, left ventricular dysfunction, congestive cardiac failure (for which myocardial fibrosis may be an underlying pathology), treatment with drugs such as daunorubicin and others which are used in cancer chemotherapy, genetic predisposition, other conditions such as Conn's Syndrome and Pheochromocytoma, high salt diet and the like.

Preferably the associated condition includes conditions which give rise to generation of profibrotic mediators or conditions which predispose a subject to myocardial fibrosis.

Preferably the prophylactic treatment is used to prevent or slow down the development of fibrosis in the at risk group.

Preferably the condition is hypertension and/or high salt intake, diseases such as diabetes and the like. Most preferably, the condition is hypertension.

In another aspect, the invention relates to method of therapeutic or prophylactic treatment of myocardial fibrosis by reducing hypertension, the method including
5 administering to a subject in need or at risk a composition including a VIP fragment.

In another aspect, the invention relates to a method of therapeutic or prophylactic treatment of myocardial fibrosis by reducing hypertension, the method including administering an associated condition a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12),
10 VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19),
15 VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

Preferably, the method is used to prevent or slow down progression of established
20 myocardial fibrosis.

Preferably the method is used to reduce the degree of established fibrosis.

In another aspect, the invention relates to method of lowering blood pressure comprising administering to a subject in need thereof a composition including a pharmaceutically effective amount of one or more of vasoactive intestinal peptide (VIP)
25 and/or active fragment thereof with a pharmaceutically acceptable carrier.

In another aspect, the invention relates to a method of lowering blood pressure comprising administering to a subject in need thereof a composition including a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6-10 or 16-25 of
30 native VIP, or conservative substitutions thereof.

In another aspect, the invention relates to a method of lowering blood pressure, the method including administering to a subject a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12),

VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23),
5 VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28),VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof.

10 In another aspect, the invention relates to a method of treating hypertension comprising administering to a subject in need thereof a composition including a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6-10 or 16-25 of native VIP, or conservative substitutions thereof.

15 In another aspect, the invention relates to a method of treating hypertension comprising administering to a subject in need thereof a composition including a pharmaceutically effective amount of one or more functional vasoactive intestinal peptide (VIP) fragments having as a core sequence amino acid residues 6-10 or 16-25 of native VIP, or conservative substitutions thereof.

20 In another aspect, the invention relates to a method of therapeutic or prophylactic treatment of hypertension, the method including administering to a subject a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24),
25 VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28),VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26),
30 VIP(16-27), VIP(16-28) or conservative substitutions thereof.

Preferably, the subject has myocardial fibrosis or an associated condition.

In another aspect, the invention relates to a method of reducing the levels, inhibiting or reducing the production of pro-fibrotic mediators in a subject, the method including administering to the subject a composition according to the present invention.

In another aspect, the invention relates to a method of reducing collagen formation
5 or enhancing collagen degradation in the cardiac muscle of a subject, the method including administering to the subject a composition according to the invention

In another aspect, the invention relates to the use of a composition according to the invention for the manufacture of a medicament for the treatment or prophylaxis of myocardial fibrosis or an associated condition

10 Preferably, the use is to prevent or slow down progression of established myocardial fibrosis. Alternatively, the use is to prevent or slow down the development of fibrosis in an at risk group. Also preferably, the use is to reduce the degree of established fibrosis.

Preferably, the associated condition is hypertension, diabetes, myocarditis,
15 ischaemic heart disease, left ventricular hypertrophy, diastolic dysfunction, myocarditis, cardiomyopathy, left ventricular dysfunction, congestive cardiac failure (for which myocardial fibrosis may be an underlying pathology), treatment with drugs such as daunorubicin and others which are used in cancer chemotherapy, genetic predisposition, other conditions such as Conn's Syndrome and Pheochromocytoma, high salt diet and
20 the like.

Preferably, the associated condition includes conditions which give rise to generation of profibrotic mediators or conditions which predispose a subject to myocardial fibrosis. Preferably, the condition is hypertension and/or high salt intake, diseases such as diabetes and the like.

25 In another aspect, the invention relates to the use of a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-
30 15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-

26), VIP(6-27), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof in the manufacture of a medicament for the therapeutic or prophylactic treatment of myocardial fibrosis or an associated condition.

In another aspect, the invention relates to the use of a composition including one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28) or conservative substitutions thereof in the manufacture of a medicament for the therapeutic or prophylactic treatment of myocardial fibrosis by reducing hypertension.

In another aspect, the invention relates to the use of VIP or a fragment thereof in the manufacture of a medicament for lowering blood pressure in a subject. Preferably, the VIP fragment is one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20), VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-28), VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16), VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26), VIP(16-27), VIP(16-28).

In another aspect, the invention relates to the use of a VIP in the manufacture of a medicament for the therapeutic or prophylactic treatment of hypertension. Preferably, the VIP fragment is one or more VIP fragments selected from the group consisting of VIP(4-10), VIP(4-11), VIP(4-12), VIP(4-13), VIP(4-14), VIP(4-15), VIP(4-16), VIP(4-17), VIP(4-18), VIP(4-19), VIP(4-20), VIP(4-21), VIP(4-22), VIP(4-23), VIP(4-24), VIP(4-25), VIP(4-26), VIP(4-27), VIP(4-28) VIP(5-10), VIP(5-11), VIP(5-12), VIP(5-

13), VIP(5-14), VIP(5-15), VIP(5-16), VIP(5-17), VIP(5-18), VIP(5-19), VIP(5-20),
VIP(5-21), VIP(5-22), VIP(5-23), VIP(5-24), VIP(5-25), VIP(5-26), VIP(5-27), VIP(5-
28),VIP(6-10), VIP(6-11), VIP(6-12), VIP(6-13), VIP(6-14), VIP(6-15), VIP(6-16),
VIP(6-17), VIP(6-18), VIP(6-19), VIP(6-20), VIP(6-21), VIP(6-22), VIP(6-23), VIP(6-
5 24), VIP(6-25), VIP(6-26), VIP(6-27), VIP(6-28), VIP(10-28), VIP(16-25), VIP(16-26),
VIP(16-27), VIP(16-28).

BRIEF DESCRIPTION OF THE FIGURES

Figure 1: Comparison of the degree of myocardial fibrosis in WKY rats on a high salt
10 diet after 4 week infusion of vehicle , VIP(6-16) or VIP(4-16).

Figure 2: Comparison of myocardial fibrosis in rats aged 14 weeks (zero time) then on
a high salt diet for 4 weeks and receiving either vehicle infusion or VIP(6-16) infusion
(5pmol/kg/min) for 4 weeks).

Figure 3: Myocardial fibrosis in 18 week old WKY rats after 4 weeks on a high salt diet
15 and treated with VIP(6-16), VIP(10-28), VIP(16-28), VIP(4-16) and VIP(1-12) or
vehicle Control. All peptides were infused at a dose of 5pmol/kg/min for 4 weeks.

Figure 4: Myocardial fibrosis in 18 week old WKY rats after 4 weeks on a high salt
diet and treated with native VIP, VIP(6-28), VIP(6-16), or vehicle Control. All peptides
were infused at a dose of 5pmol/kg/min for 4 weeks.

20 **Figure 5:** Myocardial fibrosis in 18 week old WKY rats after 4 weeks on a high salt
diet and treated with native VIP, VIP(6-28), VIP(10-28), VIP(16-28), VIP(4-16) and
VIP(1-12) or vehicle Control. All peptides were infused at a dose of 5pmol/kg/min for 4
weeks.

Figure 6: Systolic blood pressure (SBP) in SHR after 4 weeks on a high salt diet and
25 treated with VIP(1-28), VIP(4-10), VIP(4-12), VIP(4-16), VIP(4-20), VIP(4-24) and
VIP(4-28) or control. All peptides were infused at a dose of 5pmol/kg/min for 4 weeks.

Figure 7: Systolic blood pressure (SBP) in SHR after 4 weeks on a high salt diet and
treated with VIP(1-28), VIP(6-10), VIP(6-12), VIP(6-16), VIP(6-20), VIP(6-24) and
VIP(6-28) or control. All peptides were infused at a dose of 5pmol/kg/min for 4 weeks.

30 **Figure 8:** Systolic blood pressure (SBP) in SHR after 4 weeks on a high salt diet and
treated with VIP(1-28), VIP(16-20), VIP(16-24) and VIP(16-28) or control. All peptides
were infused at a dose of 5pmol/kg/min for 4 weeks.

Figure 9: Systolic blood pressure (SBP) in SHR after 4 weeks on a high salt diet and treated with VIP(1-28), VIP(1-12), and VIP(10-28) or control. All peptides were infused at a dose of 5pmol/kg/min for 4 weeks.

5

DESCRIPTION OF THE PREFERRED EMBODIMENT

It has now been found, quite surprisingly in view of the well accepted views held in this field, that VIP fragments lacking amino acids and motifs thought to be important for their function are nevertheless useful therapeutic agents to reverse or delay onset of myocardial fibrosis, or prevent onset of fibrosis in subjects at risk of developing myocardial fibrosis. Thus, VIP fragments are also useful in the treatment of congestive cardiac failure.

The use of the pharmaceutical compositions of the invention in the treatment of myocardial fibrosis or associated conditions represents a new class of therapeutic agents for these conditions. Existing treatments for myocardial fibrosis or associated conditions usually target one, or at the most two, of the known causative mechanisms in myocardial fibrosis. Without wishing to be bound by any particular mechanism of action, it is believed that the pharmaceutical preparations of the invention may target virtually all the currently known promoters of myocardial fibrosis.

On the basis of the present studies, and not wishing to be bound by theory, it is postulated that VIP fragments act as a major regulator to prevent the development of fibrosis and that the depletion of VIP may unleash the synthesis of a number of profibrotic mediators, thereby causing myocardial injury. The VIP fragments of the present invention seem to be able to act in much the same way as the native VIP but are more suited for therapeutic applications due to smaller size and hence increased stability and ease of manufacture.

Further, and unexpectedly, a number of fragments lacking certain amino acid residues and domains commonly thought to be important for its function have been shown to be active in the present studies. Particularly useful fragments are VIP(6-16), VIP(16-28), VIP(10-28), VIP(6-12), VIP(6-10), VIP(6-20), VIP(6-24) and VIP(4-16). Further, fragments listed in Table 1 are also expected to be useful in the methods of the present invention. Fragments VIP(1-12) and VIP(6-28) have been described earlier and

are not included in the present invention.

5

Other useful VIP fragments are listed in Table 1. All the sequences relate to VIP
10 and fragments of human origin but due to the very high level of amino acid
conservation, VIP and fragments thereof derived from other mammalian species are also
contemplated and encompassed by the present invention.

The present invention also contemplates pharmaceutical compositions which
include active VIP fragments. Such compositions may include any type of dosage form
15 such as tablets, capsules, powders, liquid formulations, delayed or sustained release,
patches, snuffs, nasal sprays and the like. The formulations may additionally include
other ingredients such as dyes, preservatives, buffers and anti-oxidants, for example.
The physical form and content of the pharmaceutical formulations contemplated are
conventional preparations that can be formulated by those skilled in the pharmaceutical
20 formulation field and are based on well established principles and compositions
described in, for example, Remington: The Science and Practice of Pharmacy, 19th
Edition, 1995; British Pharmacopoeia 2000 and similar formulation texts and manuals.
The compositions of the present invention may also include other active agents useful in
the treatment of cardiovascular conditions.

25 The route and frequency of administration of the compositions of the present
invention will depend on the treatment requirements and the nature of the molecule to be
administered. Thus the formulations may be suitably prepared for administration by
intravenous, intramuscular or subcuticular injection. VIP fragments may also be suitable
for mucosal administration such as oral, sublingual, nasal and the like. These parameters
30 are easily established by those skilled in the art.

The pharmaceutical compositions of the invention have been shown to be effective
in preventing or slowing down progression of established myocardial fibrosis, as well as
in reducing the degree (reversal) of established fibrosis and thus important in therapeutic

applications. The compositions of the present invention are also useful for prophylactic or therapeutic treatment of congestive cardiac failure. These are important findings with respect to the range and severity of conditions which can be treated with the compositions of the present invention.

5 Further, the compositions of the present invention may be used prophylactically in subjects at risk of developing myocardial fibrosis or an associated condition. As an example of subjects in the risk category are those having hypertension, diabetes, myocarditis, ischaemic heart disease, drugs such as daunorubicin and others which are used in cancer chemotherapy, genetic predisposition, other conditions such as Conn's
10 Syndrome and Pheochromocytoma, high salt diet and the like. The term "prophylactic" as used in the context of the present invention is intended *inter alia* to encompass treatments used to prevent or slow down the development of fibrosis in the at risk group. High proportion of subjects which may be given prophylactic treatment may already have signs of early heart failure on echocardiography.

15 The term "associated condition" as used in the context of the present invention and in reference to myocardial fibrosis is intended to encompass, without limitation, left ventricular hypertrophy, diastolic dysfunction, myocarditis, cardiomyopathy, left ventricular dysfunction and congestive cardiac failure (for which myocardial fibrosis may be an underlying pathology). The associated condition may also include conditions
20 which give rise to generation of profibrotic mediators or conditions which predispose a subject to myocardial fibrosis, such as for example hypertension and/or high salt intake, diseases such as diabetes and the like.

By conserving the VIP content of the cardiac muscle in a subject with, or at risk of developing, myocardial fibrosis or associated condition, through the use of the
25 compositions of the present invention significant therapeutic benefits can be achieved including reduction of fibrosis, reduction in the level, production or activity of profibrotic mediators, reduction in progression of fibrosis, reduction in collagen formation or enhancing collagen degradation in the cardiac muscle.

The invention will now be described more particularly with reference to non-
30 limiting examples.

EXPERIMENTAL

All general methodology and techniques have been described in detail in PCT/AU2005/000835.

Example 1. – Effect of VIP fragment infusion on fibrosis in animal models of fibrosis

5 Four animal models of myocardial fibrosis were used (animals obtained from Australian Animal Resources, Perth, Western Australia, Australia)

- i) WKY rat fed a high salt diet
- ii) WKY rat fed a high salt diet and given L-NAME (ω -monomethyl-nitro-L-arginine, Sigma Chemical Co.) 10 mg/kg/day in the drinking water
- 10 iii) WKY rats with diabetes induced by streptozotocin injection 60 mg/kg
- iv) SHR rat fed a 2.2% salt diet for 4 weeks

In each model the rats were randomised to VIP(4-16) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln), VIP(6-16) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln), VIP(4-20) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-
15 Met-Ala-Val-Lys-), or VIP(4-24) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn), VIP(16-28) (Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(10-28) (Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(6-12) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg), VIP(6-10) (Phe-Thr-Asp-Asn-Tyr), VIP(6-20) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-
20 Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-), or VIP(6-24) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn) (All obtained from or synthesised by Auspep, Australia) or vehicle control (Hartman's Solution, Baxter) infusion for 4 weeks via Alzet minipump (n=6-8 each experimental group). The dose of each peptide was 5 pmol/kg/min.

25 In models i) and ii) 14 week old WKY rats were commenced on high salt diet or high salt diet plus L-NAME. They underwent operative insertion of an Alzet minipump for infusion of VIP or vehicle. After 4 weeks the rats were anaesthetised and the hearts harvested. Myocardial fibrosis levels were quantitated.

In model iii) rats are injected with streptozotocin 60mg/kg at 14 weeks of age.
30 After 8 weeks the diabetic rats are randomised to peptide infusion (5pmol/kg/min) or no treatment. The peptides are administered as above. After a further 4 weeks the rats are anaesthetised and the hearts harvested as above.

In model iv) 14 week old SHR rats are commenced on a 2.2% salt diet. They undergo operative insertion of an Alzet minipump for peptide or vehicle infusion. After 4 weeks the rats are anaesthetised and the hearts harvested. Myocardial fibrosis levels are quantitated as above.

5 The degree of myocardial fibrosis is quantitated by two methods in known manner (refer Ye VZC, Hodge G, Yong JLC & Duggan KA (2002) Early myocardial fibrosis is associated with depletion of vasoactive intestinal peptide in the heart *Exp. Physiol* 87:539-546 Ye VZC, Hodge G, Yong JLC & Duggan KA (2003) Myocardial VIP and myocardial fibrosis induced by nitric oxide synthase inhibition in the rat *Acta Physiol. Scand.* 179:353-360. Ye VZC, Hodge G, Yong JLC & Duggan KA (2004) Vasopeptidase inhibition reverses myocardial VIP depletion and decreases myocardial fibrosis in salt sensitive hypertension *Europ. J. Pharmacol.* 485:235-242).

15 A significant improvement was found in the hearts of the treated rats in that the myocardial fibrosis index was significantly lower in the treated rats compared to the control groups (Figure 1).

Example 2 - Effect of VIP fragment treatment on regression of fibrosis in animal models of fibrosis.

To determine whether VIP (6-16), VIP(6-20) and VIP(6-24) infusion caused regression of existing fibrosis as well as prevented progression of fibrosis two groups of studies were performed. The degree of myocardial fibrosis in untreated 14 week old WKY rats was compared with the degree of fibrosis in 18 week old WKY rats after 4 weeks treatment with a high salt diet or a high salt diet plus L-NAME (10mg/kg/day) and either VIP (6-16), VIP(6-20) and VIP(6-24) (5pmol/kg/min) or vehicle control infusion (see Figure 2)

25 Preliminary results are shown in Figure 2. The data clearly show regression of myocardial fibrosis following VIP(6-16) infusion. The regression of fibrosis may be due to either reduction in collagen formation or enhancement of its degradation. Although not wishing to be bound by any particular mechanism of action, the regression of myocardial fibrosis is likely to be due to the action of collagenases, which play a part in collagen resorption.

30 The importance of the present invention to health care will be immediately apparent to one skilled in the art upon reading this disclosure. Although the capacity to

treat cardiac failure has improved significantly with the advent of angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers, as well as the realisation that aldosterone antagonists and beta blockers improve outcome in later stage disease, the addition of the pharmaceutical preparations of the invention, which act to prevent the progression of the underlying lesion (fibrosis), or even reverse fibrosis, has the capacity to prevent the escalation of mild to severe disease and hence to substantially reduce the health care burden. The overall size of certain VIP fragments and their activity makes them ideally suitable as targets for drug development.

10 EXAMPLE 3 - Effect of VIP fragment treatment on blood pressure in animal models of hypertension.

To determine whether VIP and fragments were effective in controlling blood pressure, Spontaneous Hypertensive Rats (SHR) were acclimated to blood pressure measurement by tail cuff plethysmography and then they were placed on a 2.2% salt diet for 4 weeks.

The rats were randomized to Control (Hartman's Solution) vehicle infusion, VIP(1-28) (His-Ser-Asp-Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(4-10) (Ala-Val-Phe-Thr-Asp-Asn-Tyr), VIP(4-12) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg), VIP(4-16) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln), VIP(4-20) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys), VIP(4-24) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn), VIP(4-28) (Ala-Val-Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(6-10) (Phe-Thr-Asp-Asn-Tyr), VIP(6-12) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg), VIP(6-16) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln), VIP(6-20) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys), VIP(6-24) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn), VIP(6-28) (Phe-Thr-Asp-Asn-Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(16-20) (Gln-Met-Ala-Val-Lys), VIP(16-24) (Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn), VIP(16-28) (Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(1-12) (His-Ser-Asp-Ala-Val-Phe-Thr-

Asp-Asn-Tyr-Thr-Arg) and VIP(10-28) (Tyr-Thr-Arg-Leu-Arg-Lys-Gln-Met-Ala-Val-Lys-Lys-Tyr-Leu-Asn-Ser-Ile-Leu-Asn), VIP(4-10) (Ala-Val-Phe-Thr-Asp-Asn-Tyr).

Each peptide was infused at 5pmol/kg/min.

Blood pressure was measured by tail cuff plethysmography twice weekly for 4
5 weeks.

The resultant systolic blood pressure (SBP) measurements are shown in Figures 6-9. Several of the fragments showed a reduction in SPB relative to the control solution. VIP itself was not seen to have any significant effect in reducing SBP, but a number of fragments, notably VIP(6-28), VIP(4-24), VIP(4-28), and more particularly VIP(4-16),
10 VIP(4-12), VIP(4-20), VIP(6-20) and VIP(6-24) achieved significant reductions in SBP. Figure 6 illustrates to some extent the difficulty in predicting VIP fragment activity from structure. VIP(4-16) is less efficacious than either VIP(4-12), which has 4AA less, or VIP(4-20), which is 4AA longer.

VIP(6-28) was disclosed in PCT/AU2005/000835 as being effective for
15 preventing myocardial fibrosis.

The scope of the claims should not be limited by the preferred embodiments set forth in the examples but should be given the broadest interpretation consistent with the description as a whole.

Amino acid position No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33				
VIP 6-10						P h e	T h r	A s p	A s n	T y r																											
VIP 6-11						P h e	T h r	A s p	A s n	T h r																											
VIP 6-12						P h e	T h r	A s p	A s n	T h r	A r g																										
VIP 6-13						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u																								
VIP 6-14						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g																							
VIP 6-15						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g	L y s																						
VIP 6-16						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g	L y s	G l n																					
VIP 6-17						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g	L y s	G l n	M e t																				
VIP 6-18						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g	L y s	G l n	M e t	A l a																			
VIP 6-19						P h e	T h r	A s p	A s n	T h r	A r g	A r g	L e u	A r g	L y s	G l n	M e t	A l a	V a l																		

Amino acid position No.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33		
VIP 16-25																	Gln	Met	Ala	Val	Lys	Lys	Tyr	Leu	Asn	Ser									
VIP 16-26																	Gln	Met	Ala	Val	Lys	Lys	Tyr	Leu	Asn	Ser	Ile								
VIP 16-27																	Gln	Met	Ala	Val	Lys	Lys	Tyr	Leu	Asn	Ser	Ile	Leu							
VIP 16-28																	Gln	Met	Ala	Val	Lys	Lys	Tyr	Leu	Asn	Ser	Ile	Leu	Asn						

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A composition comprising one or more vasoactive intestinal peptide (VIP) fragments selected from the group consisting of: VIP(4-12) (SEQ ID NO: 31), VIP(4-16) (SEQ ID NO: 35), VIP(4-20) (SEQ ID NO: 39) and VIP(4-24) (SEQ ID NO: 43).
2. The composition according to claim 1 for prophylactic or therapeutic treatment of myocardial fibrosis and/or hypertension.
3. The composition according to claim 1 for prophylactic or therapeutic treatment of hypertension in a patient with myocardial fibrosis.
4. The composition according to any one of claims 1 to 3 wherein the composition is formulated for administration by oral, intravenous, intramuscular or subcuticular routes.
5. Use of the composition according to claim 1 for the manufacture of a medicament for the therapeutic or prophylactic treatment of myocardial fibrosis and/or hypertension.
6. The use according to claim 5 wherein the medicament prevents or slows down progression of established myocardial fibrosis.
7. The use according to claim 5 wherein the medicament reduces the degree of established fibrosis.
8. The use according to claim 5 wherein the medicament prevents or slows down the development of fibrosis.
9. Use of the composition according to claim 1 for the manufacture of a medicament for the therapeutic or prophylactic treatment of myocardial fibrosis by reducing hypertension.

10. The use according to claim 9 wherein the medicament prevents or slows down progression of established myocardial fibrosis.
11. The use according to claim 9 wherein the medicament reduces the degree of established fibrosis.
12. Use of the composition according to claim 1 for the manufacture of a medicament for lowering blood pressure.
13. Use of the composition according to claim 1 for the manufacture of a medicament for treating hypertension in a subject.
14. The use according to claim 13 wherein the subject has myocardial fibrosis.
15. The use according to any one of claims 5 to 14 wherein the medicament reduces the levels, inhibits or reduces the production of pro-fibrotic mediators.
16. The use according to any one of claims 5 to 15 wherein the medicament reduces collagen formation or enhances collagen degradation in the cardiac muscle.
17. The use according to any one of claims 5 to 16, wherein the medicament is for use with one or more other active agents useful in the treatment of cardiovascular conditions.
18. The use according to any one of claims 5 to 17, wherein the medicament is formulated for administration by oral, intravenous, intramuscular or subcuticular routes.
19. Use of the composition according to claim 1 for the therapeutic or prophylactic treatment of myocardial fibrosis and/or hypertension.
20. The use according to claim 19 to prevent or slow down progression of established myocardial fibrosis.

21. The use according to claim 19 to reduce the degree of established fibrosis.
22. The use according to claim 19 to prevent or slow down the development of fibrosis.
23. Use of the composition according to claim 1 for the therapeutic or prophylactic treatment of myocardial fibrosis by reducing hypertension.
24. The use according to claim 23 to prevent or slow down progression of established myocardial fibrosis.
25. The use according to claim 23 to reduce the degree of established fibrosis.
26. Use of the composition according to claim 1 for lowering blood pressure.
27. Use of the composition according to claim 1 for treating hypertension in a subject.
28. The use according to claim 27 wherein the subject has myocardial fibrosis.
29. The use according to any one of claims 19 to 28 to reduce the levels or to inhibit or reduce the production of pro-fibrotic mediators.
30. The use according to any one of claims 19 to 29 to reduce collagen formation or enhance collagen degradation in the cardiac muscle.
31. The use according to any one of claims 19 to 30, wherein the use is in conjunction with one or more other active agents useful in the treatment of cardiovascular conditions.
32. The use according to any one of claims 19 to 31, wherein the composition is formulated for administration by oral, intravenous, intramuscular or subcuticular routes.




Figure 1:




Figure 2:




Figure 3




Figure 4






Figure 5




Systolic blood pressure (SBP) in SHR on 2.2% salt diet after 4 weeks treatment. Peptides were infused at 5pmol/kg/min. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Figure 6




Systolic blood pressure (SBP) in SHR on 2.2% salt diet after 4 weeks treatment. Peptides were infused at 5pmol/kg/min. * $p < 0.01$, ** $p < 0.0005$

Figure 7



Systolic blood pressure after 4 weeks treatment in SHR on 2.2% salt diet. Peptides were infused at 5 pmol/kg/min.

Figure 8



Systolic blood pressure (SBP) after 4 weeks treatment in SHR on 2.2% salt diet. Peptides were infused at 5pmol/kg/min

Figure 9