



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/09/11
(87) Date publication PCT/PCT Publication Date: 2021/03/18
(85) Entrée phase nationale/National Entry: 2022/03/11
(86) N° demande PCT/PCT Application No.: US 2020/050386
(87) N° publication PCT/PCT Publication No.: 2021/050862
(30) Priorités/Priorities: 2019/09/13 (US62/900,141);
2019/11/18 (US62/936,951)

(51) Cl.Int./Int.Cl. *A61K 35/17* (2015.01),
A61K 39/00 (2006.01), *A61K 39/395* (2006.01),
A61P 35/00 (2006.01), *A61P 35/02* (2006.01),
C07K 14/705 (2006.01)
(71) Demandeurs/Applicants:
MEMORIAL SLOAN-KETTERING CANCER CENTER,
US;
TRI-INSTITUTIONAL THERAPEUTICS DISCOVERY
INSTITUTE, INC., US;
MEMORIAL HOSPITAL FOR CANCER AND ALLIED
DISEASES, US;
SLOAN-KETTERING INSTITUTE FOR CANCER
RESEARCH, US
(72) Inventeurs/Inventors:

(54) Titre : RECEPTEURS DE RECONNAISSANCE D'ANTIGENE CIBLANT CD371 ET LEURS UTILISATIONS
(54) Title: ANTIGEN RECOGNIZING RECEPTORS TARGETING CD371 AND USES THEREOF

Phage-based CD371-targeted CAR s

Et.B10L3H_MT_h28Z (B10 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L3H, and a CD28Z-based signaling domain)

Et.B10L4H_MT_hBBZ (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a 4-1BBZ-based signaling domain)

Et.B10L4H_MT_h28Z (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.B10H3L_MT_h28Z (B10, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.B10H4L_MT_h28Z (B10, Heavy-Light scFv orientation with a 19-amino acid G4S linker, H4L, and a CD28Z-based signaling domain)

Et.C3H3L_MT_h28Z (C3, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.C3L3H_MT_h28Z (C3 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.D6L3H_MT_h28Z (D6 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)




FIG. 1

(57) **Abrégé/Abstract:**

The presently disclosed subject matter provides for antigen-recognizing receptors that specifically target CD371 and cells comprising such CD371 -targeted antigen-recognizing receptors. The presently disclosed subject matter further provides uses of the CD371 -targeted antigen-recognizing receptors for treatment. The presently disclosed subject matter provides methods and compositions for immunotherapies. It relates to antigen-recognizing receptors (e.g., chimeric antigen receptors (CARs) or T-cell receptors (TCRs)) that specifically target CD371, cells comprising such receptors, and methods of using such cells for treatments.

(72) **Inventeurs(suite)/Inventors(continued):** DANIYAN, ANTHONY, US; BRENTJENS, RENIER J., US; LORENZ, IVO C., US;
POHL, MARY ANN, US

(74) **Agent:** SMART & BIGGAR LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
18 March 2021 (18.03.2021)



(10) International Publication Number
WO 2021/050862 A1

(51) International Patent Classification:

A61K 35/17 (2015.01) *A61P 35/00* (2006.01)
A61K 39/00 (2006.01) *A61P 35/02* (2006.01)
A61K 39/395 (2006.01) *C07K 14/705* (2006.01)

(21) International Application Number:

PCT/US2020/050386

(22) International Filing Date:

11 September 2020 (11.09.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/900,141 13 September 2019 (13.09.2019) US
62/936,951 18 November 2019 (18.11.2019) US

(71) Applicants: **MEMORIAL SLOAN-KETTERING CANCER CENTER** [US/US]; 1275 York Avenue, New York, NY 10065 (US). **TRI-INSTITUTIONAL THER-**

APEUTICS DISCOVERY INSTITUTE, INC. [US/US]; 413 East 69th Street, Box 300, New York, NY 10021 (US).

(72) **Inventors:** **DANIYAN, Anthony**; 303 East 60th Street Apt. 12C, New York, NY 10022 (US). **BRENTJENS, Renier, J.**; 1275 York Avenue, New York, NY 10065 (US). **LORENZ, Ivo, C.**; 413 East 69th Street, Box 300, New York, NY 10021 (US). **POHL, Mary, Ann**; 31-01 Vernon Blvd., Apt. 301, Astoria, NY 11106 (US).

(74) **Agent:** **LENDARIS, Steven, P.** et al.; Baker Botts LLP, 30 Rockefeller Plaza, New York, NY 10112-4498 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW,

(54) Title: ANTIGEN RECOGNIZING RECEPTORS TARGETING CD371 AND USES THEREOF

Phage-based CD371-targeted CAR s

Et.B10L3H_MT_h28Z (B10 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L3H, and a CD28Z-based signaling domain)

Et.B10L4H_MT_hBBZ (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a 4-1BBZ-based signaling domain)

Et.B10L4H_MT_h28Z (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.B10H3L_MT_h28Z (B10, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.B10H4L_MT_h28Z (B10, Heavy-Light scFv orientation with a 19-amino acid G4S linker, H4L, and a CD28Z-based signaling domain)

Et.C3H3L_MT_h28Z (C3, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.C3L3H_MT_h28Z (C3 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.D6L3H_MT_h28Z (D6 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)




FIG. 1

(57) **Abstract:** The presently disclosed subject matter provides for antigen-recognizing receptors that specifically target CD371 and cells comprising such CD371 -targeted antigen-recognizing receptors. The presently disclosed subject matter further provides uses of the CD371 -targeted antigen-recognizing receptors for treatment. The presently disclosed subject matter provides methods and compositions for immunotherapies. It relates to antigen-recognizing receptors (e.g., chimeric antigen receptors (CARs) or T-cell receptors (TCRs)) that specifically target CD371, cells comprising such receptors, and methods of using such cells for treatments.

[Continued on next page]



WO 2021/050862 A1

WO 2021/050862 A1

SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

ANTIGEN RECOGNIZING RECEPTORS TARGETING CD371 AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application claims priority to United States Provisional Application
5 No. 62/900,141 filed September 13, 2019 and United States Provisional Application No. 62/936,951 filed November 18, 2019, the contents of each of which are incorporated by reference in their entireties herein, and priority to each of which is claimed.

SEQUENCE LISTING

The present application contains a Sequence Listing which has been submitted in
10 ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on September 11, 2020, is named 0727341147_ST25 and is 211,921 bytes in size.

1. INTRODUCTION

The presently disclosed subject matter provides methods and compositions for
15 immunotherapies. It relates to antigen-recognizing receptors (e.g., chimeric antigen receptors (CARs) or T-cell receptors (TCRs)) that specifically target CD371, cells comprising such receptors, and methods of using such cells for treatments.

2. BACKGROUND OF THE INVENTION

Cell-based immunotherapy is a therapy with curative potential for the treatment of
20 cancer. T cells and other immune cells may be modified to target tumor antigens through the introduction of genetic material coding for artificial or synthetic receptors for antigen, termed Chimeric Antigen Receptors (CARs), specific to selected antigens. Targeted T cell therapy using CARs has shown recent clinical success in treating hematologic malignancies.

25 Acute myeloid leukemia (AML) is the most common type of adult acute leukemia. It is characterized by the accumulation of immature myeloid cells in the bone marrow that results in dysfunction of hematopoiesis. Chemotherapy and hematopoietic stem cell transplantation (HSCT) are standard treatment of AML. However, a majority of patients eventually relapse and succumb to the disease.

30 AML is the most common acute leukemia in adults. The standard induction chemotherapy regimens have not changed substantially over the past 40 years and the overall survival remains very poor. Frequent recurring abnormalities involving genes coding for epigenetic modifiers have been identified. The development of CAR therapy

for AML is hampered by the lack of suitable targets. Accordingly, there are needs for novel therapeutic strategies to design CARs targeting antigens that are highly expressed in AML cells and limited expression in normal tissues for treating AML, and for strategies capable of inducing potent cancer eradication with minimal toxicity and immunogenicity.

3. SUMMARY OF THE INVENTION

The presently disclosed subject matter provides antigen-recognizing receptors that specifically target CD371 and cells comprising such CD371-targeted antigen-recognizing receptors. The presently disclosed subject matter further provides uses of the CD371-targeted antigen-recognizing receptors for treatment.

The presently disclosed subject matter provides an antigen-recognizing receptor, comprising an extracellular antigen-binding domain, a transmembrane domain, and an intracellular signaling domain, wherein the extracellular antigen-binding domain specifically binds to CD371. In certain embodiments, the extracellular antigen-binding domain is a single-chain variable fragment (scFv). In certain embodiments, the extracellular antigen-binding domain is a human scFv. In certain embodiments, the extracellular antigen-binding domain is a Fab, which is optionally crosslinked. In certain embodiments, the extracellular antigen-binding domain is a F(ab)₂. In certain embodiments, one or more of the scFv, Fab and F(ab)₂ are comprised in a fusion protein with a heterologous sequence to form the extracellular antigen-binding domain.

In certain embodiments, the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30 or a conservative modification thereof;

(b) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36 or a conservative modification thereof;

(c) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 42 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 43 or a conservative modification thereof;

(d) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 462 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48 or a conservative modification thereof;

(e) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54 or a conservative modification thereof; or

(f) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain comprises: a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30.

In certain embodiments, the extracellular antigen-binding domain comprises:

(a) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32 or a

conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33 or a conservative modification thereof;

(b) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37 or a conservative modification thereof; a light chain variable
5 region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38 or a conservative modification thereof; and a light chain variable region CDR3 comprising SEQ ID NO: 39 or a conservative modification thereof;

(c) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43 or a conservative modification thereof; a light chain variable
10 region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45 or a conservative modification thereof;

(d) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 49 or a conservative modification thereof; a light chain variable
15 region CDR2 comprising SEQ ID NO: 50 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51 or a conservative modification thereof;

(e) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 55 or a conservative modification thereof; a light chain variable
20 region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57 or a conservative modification thereof;
or

(f) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 61 or a conservative modification thereof; a light chain variable
25 region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain comprises: a
30 heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30.

In certain embodiments, the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33;

(b) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39;

(c) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45;

(d) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 46; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 49; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 50; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51;

(e) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 55; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57; or

(f) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 61; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63.

In certain embodiments, the extracellular antigen-binding domain comprises: a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33.

In certain embodiments, the extracellular antigen-binding domain comprises a heavy chain variable region comprising an amino acid sequence that is at least about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, or SEQ ID NO: 11. In certain embodiments, the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7,

SEQ ID NO: 9, or SEQ ID NO: 11. In certain embodiments, the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1.

In certain embodiments, the extracellular antigen-binding domain comprises a
 5 light chain variable region comprising an amino acid sequence that is at least about 80%,
 about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about
 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%,
 about 96%, about 97%, about 98% or about 99% homologous or identical to the amino
 acid sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8,
 10 SEQ ID NO: 10, or SEQ ID NO: 12. In certain embodiments, the extracellular antigen-
 binding domain comprises a light chain variable region comprising the amino acid
 sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ
 ID NO: 10, or SEQ ID NO: 12. In certain embodiments, the extracellular antigen-binding
 domain comprises a light chain variable region comprising the amino acid sequence set
 15 forth in SEQ ID NO: 2.

In certain embodiments, the extracellular antigen-binding domain comprises: (a) a
 heavy chain variable region comprising an amino acid sequence that is at least about
 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%,
 about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about
 20 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the
 amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID
 NO: 7, SEQ ID NO: 9, or SEQ ID NO: 11; and (b) a light chain variable region
 comprising an amino acid sequence that is at least about 80%, about 81%, about 82%,
 about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about
 25 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%,
 about 98% or about 99% homologous or identical to the amino acid sequence set forth in
 in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, or
 SEQ ID NO: 12. In certain embodiments, the extracellular antigen-binding domain
 comprises: (a) a heavy chain variable region comprising the amino acid sequence set forth
 30 in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, or SEQ
 ID NO: 11; and (b) a light chain variable region comprising the amino acid sequence set
 forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10,
 or SEQ ID NO: 12.

In certain embodiments, the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2;

5 (b) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 3, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 4;

(c) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 5, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 6;

(d) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 7, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 8;

15 (e) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 9, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 10; or

(f) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 11, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 12.

20 In certain embodiments, the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1; and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2.

In certain embodiments, the extracellular antigen-binding domain comprises a linker between a heavy chain variable region and a light chain variable region of the extracellular antigen-binding domain. In certain embodiments, the linker has the amino acid sequence set forth in SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, or SEQ ID NO: 94. In certain embodiments, the linker has the amino acid sequence set forth in SEQ ID NO: 13 or SEQ ID NO: 14.

30 In certain embodiments, the extracellular antigen-binding domain comprises a signal peptide that is covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain embodiments, the extracellular antigen-binding domain

comprises a heavy chain variable region and a light chain variable region, which are positioned from the N- to the C-terminus: V_H-V_L.

In certain embodiments, the extracellular antigen-binding domain binds to CD371 with a low binding affinity. In certain embodiments, the extracellular antigen-binding domain binds to CD371 with a dissociation constant (K_d) of 1×10^{-8} M or more.

In certain embodiments, the transmembrane domain comprises a CD28 polypeptide. In certain embodiments, the intracellular signaling domain comprises a CD3 ζ polypeptide. In certain embodiments, the intracellular signaling domain further comprises at least one co-stimulatory signaling region. In certain embodiments, the at least one co-stimulatory signaling region comprises a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, or a combination thereof. In certain embodiments, the at least one co-stimulatory signaling region comprises a CD28 polypeptide or a 4-1BB polypeptide.

In certain embodiments, the antigen-recognizing receptor is a chimeric antigen receptor (CAR), a T-cell Receptor (TCR), or a T-cell like fusion protein. In certain embodiments, the antigen-recognizing receptor is a CAR.

In certain embodiments, the antigen-recognizing receptor is recombinantly expressed. In certain embodiments, the antigen-recognizing receptor is expressed from a vector. In certain embodiments, the vector is a γ -retroviral vector.

The presently disclosed subject matter provides cells comprising a presently disclosed antigen-recognizing receptor. In certain embodiments, the cell is transduced with the antigen-recognizing receptor. In certain embodiment, the antigen-recognizing receptor is constitutively expressed on the surface of the cell.

In certain embodiments, the cell is engineered to express a cytokine or a fragment thereof. In certain embodiments, the cell further comprises an exogenous polypeptide of the cytokine or fragment thereof. In certain embodiments, the cell further comprises a nucleic acid molecule encoding the cytokine or fragment thereof. In certain embodiments, the cytokine is selected from the group consisting of IL-18, IL-33, IL-36, and combinations thereof. In certain embodiments, the cytokine is IL-18.

In certain embodiments, the cell is an immunoresponsive cell. In certain embodiments, the cell is a cell of the lymphoid lineage or a cell of the myeloid lineage. In certain embodiments, the cell is selected from the group consisting of a T cell, a Natural Killer (NK) cell, and a stem cell from which lymphoid cells may be

differentiated. In certain embodiments, the cell is a T cell. In certain embodiments, the T cell is a cytotoxic T lymphocyte (CTL) or a regulatory T cell. In certain embodiments, the stem cell is a pluripotent stem cell. In certain embodiments, the pluripotent stem cell is an embryoid stem cell or an induced pluripotent stem cell.

5 The presently disclosed subject matter further provides nucleic acid molecules that encode a presently disclosed antigen-recognizing receptor. In certain embodiments, the nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, or SEQ ID NO: 27. In certain embodiments, the nucleic acid molecule comprises the nucleotide sequence set
10 forth in SEQ ID NO: 22. The presently disclosed subject matter further provides vectors comprising the presently disclosed nucleic acid molecules. In certain embodiments, the vector is a viral vector. In certain embodiments, the vector is a γ -retroviral vector.

 In addition, the presently disclosed subject matter provides host cells expressing the nucleic acid molecule disclosed herein. In certain embodiments, the host cell is a T
15 cell.

 The presently disclosed subject matter further provides compositions comprising the cells disclosed herein. In certain embodiments, the composition is a pharmaceutical composition further comprising a pharmaceutically acceptable carrier.

 The presently disclosed subject matter further provides methods of reducing tumor
20 burden in a subject. In certain embodiments, the method comprises administering an effective amount of presently disclosed cells or composition to the subject. In certain embodiments, the method reduces the number of tumor cells, reduces tumor size, and/or eradicates the tumor in the subject. The presently disclosed subject matter further provides methods of increasing or lengthening survival of a subject having a tumor or
25 neoplasm. In certain embodiments, the method comprises administering an effective amount of presently disclosed cells or composition to the subject. The presently disclosed subject matter further provides methods of treating and/or preventing a tumor or neoplasm in a subject. In certain embodiments, the method comprises administering to the subject an effective amount of the presently disclosed cells or composition. In certain
30 embodiments, the tumor or neoplasm is selected from the group consisting of acute myeloid leukemia (AML), multiple myeloma, Non-Hodgkin Lymphoma, Hodgkin Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, myelodysplastic

syndrome (MDS), and chronic myelogenous leukemia (CML). In certain embodiments, the tumor or neoplasm is acute myeloid leukemia (AML).

The presently disclosed subject matter further provides methods for producing a presently disclosed cell comprising a CD371-targeted antigen-recognizing receptor. In certain embodiments, the method comprises introducing into the cell a nucleic acid molecule that encodes the antigen-recognizing receptor.

In addition, the presently disclosed subject matter provides kits for reducing tumor burden in a subject, treating and/or preventing a tumor or neoplasm in a subject, and/or increasing or lengthening survival of a subject having a tumor or neoplasm. In certain embodiments, the kit comprises the cell described herein. In certain embodiments, the kit further comprises written instructions for using the cell for reducing tumor burden in a subject, treating and/or preventing a tumor or neoplasm in a subject, and/or increasing or lengthening survival of a subject having a tumor or neoplasm.

4. BRIEF DESCRIPTION OF THE FIGURES

The following Detailed Description, given by way of example, but not intended to limit the invention to specific embodiments described, may be understood in conjunction with the accompanying drawings.

Figure 1 depicts structures of antigen-recognizing receptors in accordance with the presently disclosed subject matter.

Figure 2 depicts detection of CD371-targeted CAR on the surface of transduced human T cells. Antibodies against EGFRt (CETUXIMAB-APC) and Myc-tag (9B11-PE) detected surface expression of EGFRt and human anti-human CAR in B10-based CAR constructs. Controls included non-transduced human T cells which did not express EGFRt or MYC-tag, and non-MYC tag containing CAR constructs Etah19h28z (an anti-human CD19 CAR T cells) and EtC1HVh28z (anti-CD371 CAR T cells derived from a mouse anti-human CD371 antibody 107537).

Figure 3 depicts tumor cell lysis activities of CD371-targeted CAR T cells. 4 day rested (4dR) human CD371-targeted CAR T cells were cocultured with CD33⁺/CD371⁺ U937 cells expressing GFP and firefly luciferase (U937gL) at different effector:tumor (E:T) ratios. Bioluminescence was measured 24 hours later and plotted as a percentage of signal detected in a coculture of a non-functional CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL. Et.C1HVh28z represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7, and

EtM195Mth28Z represents a CD33-targeted CAR T cell derived from a murine anti-CD33 antibody, M195 (*see Zhao et al., Haematologica* (2010):95:71–78 (2010)).

Figure 4 depicts tumor cell lysis activities of CD371-targeted CAR T cells in 24 hour killing assays. Healthy donor-derived (referred to as “donor C”) CD371-targeted CAR T cells were cocultured with CD33⁺/CD371⁺ U937 cells expressing GFP and firefly luciferase (U937gLuc) at different effector:tumor (E:T) ratios. Bioluminescence was measured 24 hours later and plotted as a percentage of signal detected in a coculture of a non-functional CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gLuc. Et.C1HVh28z represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7, and EtM195Mth28z represents a CD33-targeted CAR T cell derived from murine anti-human CD33 antibody, M195.

Figure 5 depicts tumor cell lysis activities of CD371-targeted CAR T cells in 24 hour killing assays. Healthy donor-derived (referred to as “donor D”) CD371-targeted CAR T cells were cocultured with CD33⁺/CD371⁺ U937 cells expressing GFP and firefly luciferase (U937gLuc) at different effector:tumor (E:T ratios). Bioluminescence was measured 24 hours later and plotted as a percentage of signal detected in a coculture of a non-functional CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gLuc. Et.C1HVh28z represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7, and EtM195Mth28z represents a CD33-targeted CAR T cell derived from a murine anti-human CD33 antibody, M195.

Figure 6 depicts activities of CD371-targeted CAR T cells in a recursive stimulation assay. Healthy donor-derived (referred to as “donor C”) CD371-targeted CAR T cells were cocultured with CD33⁺/CD371⁺ U937gLuc cell at an E:T ratio of 1:12.5 and at a concentration of 30,000 CAR positive/ml. CAR T cells were counted approximately every 4-5 days and characterized by flow cytometry, and the starting number of tumor cells were added back into the culture (indicated by arrows). Et.B10LHdel represents a non-functional CAR T cells (lacking signaling domains); Et.C1HVh28z represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7, and EtM195Mth28z represents a CD33-targeted CAR T cell derived from a murine anti-human CD33 antibody, M195.

Figure 7 depicts activities of CD371-targeted CAR T cells in a recursive stimulation assay. Healthy donor-derived (referred to as “donor D”) CD371-targeted

CAR T cells were cocultured with CD33⁺/CD371⁺ U937gL cell at an E:T ratio of 1:12.5 and at a concentration of 30,000 CAR positive/ml. CAR T cells were counted approximately every 4-5 days and characterized by flow cytometry, and the starting number of tumor cells were added back into the culture (indicated by arrows).

- 5 Et.B10LHdel represents a non-functional CAR T cells (lacking signaling domains); Et.C1HVh28z represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7, and EtM195Mth28z represents a CD33-targeted CAR T cell derived from a murine anti-human CD33 antibody, M195.

Figures 8A and 8B depict *in vitro* anti-tumor activities of CD371-targeted CAR T
10 cells (second generation B10HL-based CAR T cells and B10LH-based CAR T cells) in a U937gL 24-hour killing assay from different healthy donors. Healthy human donor-derived (referred to as “Donor A” and “Donor “B) CD371-targeted CAR T cells were cocultured with CD371-positive U937 cells expressing GFP and firefly luciferase (U937gL) at different effector:tumor ratios. Bioluminescence was measured 24 hours
15 later and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL. Figure 8A shows the results for Donor A. Figure 8B shows the results for Donor B.

Figures 9A and 9B depict *in vitro* anti-tumor activities of CD371-targeted CAR T
20 cells (second generation B10HL-based CAR T cells and B10LH-based CAR T cells) in HL60gL 24-hour killing assay from different healthy donors. Healthy human donor-derived (referred to as “Donor A” and “Donor “B”) CD371-targeted CAR T cells were cocultured with CD371-positive HL60 cells expressing GFP and firefly luciferase (HL60gL) at different effector:tumor ratios. Bioluminescence was measured 24 hours
25 later and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and HL60gL. Figure 9A shows the results for Donor A. Figure 9B shows the results for Donor B.

Figures 10A and 10B depicts generation of human CD371 (hCD371) CRISPR
30 knockout cell lines. Human CD371 was knocked out of HL60gL (Fig. 10A) and U937gL (Fig. 10B) with CRISPR. The knockout was confirmed by flow cytometry using anti-human CD371 APC-conjugated antibody. Figure 10A shows the knockout of HL60gL. Figure 10B shows the knockout of U937gL.

Figures 11A and 11B depict cytotoxicity activities of CD371-targeted CAR T cells (second generation B10-based CAR T cells, i.e., B10-HL- and B10LH-based CAR T cells) against antigen-negative U937gL in a 24-hour killing assay from different healthy donors. Healthy human donor-derived (referred to as “Donor A” and “Donor “B”)

5 CD371-targeted CAR T cells were cocultured with CD371-negative U937 cells expressing RFP and cypridina luciferase (U937RFPcyp.371KO) at different effector:tumor ratios. Bioluminescence was measured 24 hours later and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and
10 U937RFPcyp.371KO. Figure 11A shows the results for Donor A. Figure 11B shows the results for Donor B.

Figures 12A and 12B depict cytotoxicity activities of CD371-targeted CAR T cells (second generation B10-based CAR T cells, i.e., B10-HL- and B10LH-based CAR T cells) against antigen-negative HL60gL in a 24-hour killing assay from different healthy
15 donors. Healthy human donor-derived (referred to as “Donor A” and “Donor “B”)

CD371-targeted CAR T cells were cocultured with CD371-negative HL60 cells expressing GFP and firefly luciferase (HL60L.371KO) at different effector:tumor ratios. Bioluminescence was measured 24 hours later and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on
20 B10L4H, but without a CD28 or CD3 zeta signaling domain) and HL60gL.371KO. Figure 12A shows the results for Donor A. Figure 12B shows the results for Donor B.

Figures 13A and 13B depict interferon gamma (IFN- γ) secretion of CD371-targeted CAR T cells (second generation B10-based CAR T cells, i.e., B10-HL- and B10LH-based CAR T cells). Human CD371-targeted CAR T cells were cocultured alone,
25 with CD371-negative U937 cells, or CD371+ U937cells at an effector:tumor ratio (E:T) of 1:1 (4.0×10^4 : 4.0×10^4 cells in 200 μ l). 24 hours later, supernatant was collected and IFN- γ was measured utilizing a bead-based multiplex assay. Figure 13A shows the results for Donor A. Figure 13B shows the results for Donor B.

Figures 14A and 14B depict interleukin-2 (IL-2) secretion of CD371-targeted CAR
30 T cells (second generation B10-based CAR T cells, i.e., B10-HL- and B10LH-based CAR T cells). Human CD371-targeted CAR T cells were cocultured alone, with CD371-negative U937 cells, or CD371+ U937 cells at an effector:tumor ratio (E:T) of 1:1 (4.0×10^4 : 4.0×10^4 cells in 200 μ l). 24 hours later, supernatant was collected and IL-2 was

measured using a bead-based multiplex assay. Figure 14A shows the results for Donor A. Figure 14B shows the results for Donor B.

Figures 15A and 15B depict tumor necrosis factor alpha (TNF- α) secretion of CD371-targeted CAR T cells (second generation B10-based CAR T cells, i.e., B10-HL- and B10LH-based CAR T cells). Human CD371-targeted CAR T cells were cocultured alone, with CD371-negative U937 cells, or CD371+ U937 cells at an effector:tumor ratio (E:T) of 1:1 (4.0×10^4 : 4.0×10^4 cells in 200 μ l). 24 hours later, supernatant was collected and TNF- α was measured using a bead-based multiplex assay. Figure 15A shows the results for Donor A. Figure 15B shows the results for Donor B.

Figures 16A and 16B depicts cell proliferation of B10-based CAR T cells in a recursive stimulation assay. CD371-targeted CAR T cells were generated from two healthy donors (Donor A and Donor B). CAR T cells were cocultured with CD371+ U937gL at an E:T ratio of 1:5 and at a concentration of 50,000 CAR positive/ml. Approximately every 5 days, CAR T cells were counted and characterized by flow cytometry. The starting number of tumor cells were added back into the culture (indicated at arrows). Figure 16A shows the results for Donor A. Figure 16B shows the results for Donor B.

Figures 17 illustrates a murine xenograft model of AML. NCG mice were inoculated with 5×10^4 U937gL AML FAB-M5 cell lines via tail vein. CD371-targeted CAR T cells were administered to the mice three days later. Tumor kinetics measured non-invasively with bioluminescent imaging approximately every 5 days, and survival was monitored.

Figure 18 depicts the survival of AML cell-line Xenografted mice (Donor 1) treated with human CD371-targeted CAR T cells. NCG mice were inoculated with 5×10^4 U937gL tumor cells and treated with approximately 1.25×10^6 human CD371-targeted CAR T cells three days later. Survival of mice were monitored.

Figure 19 depicts the survival of AML cell-line Xenografted mice (Donor 2) treated with human CD371-targeted CAR T cells. NCG mice were inoculated with U937gL tumor cells and treated with approximately 1.25×10^6 human CD371 targeted CAR T cells three days later. Survival of mice were monitored.

Figure 20 depicts *in vivo* activity of B10-based CAR T cells. NCG mice were inoculated with 5×10^4 U937gL tumor cells and treated with various doses of CD371-targeted CAR T cells (1.25×10^5 , 2.5×10^5 , 5.0×10^5 , and 1.0×10^6) three days later.

Survival of the mice was monitored within 55 days (Figure 20A) and 60 days (Figure 20B). Figure 21 depicts binding of scFvs to HEK293 cells expressing human CD371.

Figure 22 depicts the *in vitro* cytotoxicity of second-generation B10HL-based CAR T cells against CD371-positive targets. The B10HL-based CAR T cells have G4S linkers of varying lengths. Healthy human donor-derived (including Donors A & B) CD371-targeted CAR T cells were cocultured with antigen-positive U937 cells expressing GFP and firefly luciferase (U937gL) at different effector:tumor ratios. Bioluminescence was measured 24 hours later, and was plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL.

Figure 23 depicts *in vivo* activities of B10-based CAR T cells. NCG mice were inoculated with 5×10^4 U937gL tumor cells, and were treated with 5×10^5 of CAR T cells. Survival of the mice was monitored. B10H3L, B10H5L, and IL33-secreting B10H4L human CD371-targeted CAR T cells outperformed B10H4L-based CAR T cells *in vivo*.

Figure 24 depicts *in vivo* activities of B10-based CAR T cells. NCG mice were inoculated with 5×10^4 U937gL tumor cells, and were treated with 2.5×10^5 of CAR T cells. Survival of the mice was monitored. IL18- and IL33-secreting B10H4L Human CD371-targeted CAR T cells outperformed B10H4L-based CAR T cells *in vivo*.

Figure 25 depicts *in vivo* activities of B10-based CAR T cells. NCG mice were inoculated with 5×10^4 U937gL tumor cells, and were treated with 1.0×10^5 of CAR T cells. Survival of the mice was monitored. At a low dosage of 1.0×10^5 , IL18-secreting B10H4L human CD371-targeted CAR T cells outperformed all other constructs *in vivo*.

5. DETAILED DESCRIPTION OF THE INVENTION

The presently disclosed subject matter provides antigen-recognizing receptors (e.g., chimeric antigen receptors (CARs) or T-cell receptors (TCRs)) that specifically target CD371. The presently disclosed subject matter further provides cells comprising such receptors. The cells can be immunoresponsive cells, e.g., genetically modified immunoresponsive cells (e.g., T cells or NK cells). The presently disclosed subject matter also provides methods of using such cells for treatments, e.g., for treating and/or preventing a tumor or neoplasm (e.g., AML).

Non-limiting embodiments of the present disclosure are described by the present specification and Examples.

For purposes of clarity of disclosure and not by way of limitation, the detailed description is divided into the following subsections:

- 5.1. Definitions;
- 5.2. CD371;
- 5.3. Antigen-Recognizing Receptors;
- 5.4. Cells;
- 5.5. Compositions and Vectors;
- 5.6. Polypeptides;
- 5.7. Formulations and Administration;
- 5.8. Methods of Treatment; and
- 5.9. Kits

5.1. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag (1991); and Hale & Marham, The Harper Collins Dictionary of Biology (1991). As used herein, the following terms have the meanings ascribed to them below, unless specified otherwise.

As used herein, the term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system. For example, “about” can mean within 3 or more than 3 standard deviations, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value.

By “immunoresponsive cell” is meant a cell that functions in an immune response or a progenitor, or progeny thereof. In certain embodiments, the immunoresponsive cell is a cell of lymphoid lineage. Non-limiting examples of cells of lymphoid lineage include

T cells, Natural Killer (NK) cells, B cells, and stem cells from which lymphoid cells may be differentiated. In certain embodiments, the immunoresponsive cell is a cell of myeloid lineage.

By “activates an immunoresponsive cell” is meant induction of signal transduction or changes in protein expression in the cell resulting in initiation of an immune response. For example, when CD3 Chains cluster in response to ligand binding and immunoreceptor tyrosine-based inhibition motifs (ITAMs) a signal transduction cascade is produced. In certain embodiments, when an endogenous TCR or an exogenous CAR binds to an antigen, a formation of an immunological synapse occurs that includes clustering of many molecules near the bound receptor (e.g. CD4 or CD8, CD3 $\gamma/\delta/\epsilon/\zeta$, etc.). This clustering of membrane bound signaling molecules allows for ITAM motifs contained within the CD3 chains to become phosphorylated. This phosphorylation in turn initiates a T cell activation pathway ultimately activating transcription factors, such as NF- κ B and AP-1. These transcription factors induce global gene expression of the T cell to increase IL-2 production for proliferation and expression of master regulator T cell proteins in order to initiate a T cell mediated immune response.

By “stimulates an immunoresponsive cell” is meant a signal that results in a robust and sustained immune response. In various embodiments, this occurs after immune cell (e.g., T-cell) activation or concomitantly mediated through receptors including, but not limited to, CD28, CD137 (4-1BB), OX40, CD40 and ICOS. Receiving multiple stimulatory signals can be important to mount a robust and long-term T cell mediated immune response. T cells can quickly become inhibited and unresponsive to antigen. While the effects of these co-stimulatory signals may vary, they generally result in increased gene expression in order to generate long lived, proliferative, and anti-apoptotic T cells that robustly respond to antigen for complete and sustained eradication.

The term “antigen-recognizing receptor” as used herein refers to a receptor that is capable of recognizing a target antigen (e.g., CD371). In certain embodiments, the antigen-recognizing receptor is capable of activating an immune or immunoresponsive cell (e.g., a T cell) upon its binding to the target antigen.

As used herein, the term “antibody” means not only intact antibody molecules, but also fragments of antibody molecules that retain immunogen-binding ability. Such fragments are also well known in the art and are regularly employed both *in vitro* and *in vivo*. Accordingly, as used herein, the term “antibody” means not only intact

immunoglobulin molecules but also the well-known active fragments F(ab')₂, and Fab. F(ab')₂, and Fab fragments that lack the Fe fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding of an intact antibody (Wahl et al., . *Nucl Med* (1983);24:316-325). As used herein, include whole native

5 antibodies, bispecific antibodies; chimeric antibodies; Fab, Fab', single chain V region fragments (scFv), fusion polypeptides, and unconventional antibodies. In certain embodiments, an antibody is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V_H) and a heavy chain constant

10 (C_H) region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as V_L) and a light chain constant C_L region. The light chain constant region is comprised of one domain, C_L. The V_H and V_L regions can be further sub-divided into regions of hypervariability, termed complementarity determining regions (CDR),

15 interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the

20 immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells) and the first component (C1q) of the classical complement system.

As used herein, "CDRs" are defined as the complementarity determining region amino acid sequences of an antibody which are the hypervariable regions of immunoglobulin heavy and light chains. *See, e.g.*, Kabat et al., Sequences of Proteins of

25 Immunological Interest, 4th U. S. Department of Health and Human Services, National Institutes of Health (1987), or IMGT numbering system (Lefranc, *The Immunologist* (1999);7:132-136; Lefranc *et al.*, *Dev. Comp. Immunol.* (2003);27:55-77). Generally, antibodies comprise three heavy chain and three light chain CDRs or CDR regions in the variable region. CDRs provide the majority of contact residues for the binding of the

30 antibody to the antigen or epitope. In certain embodiments, the CDRs regions are delineated using the IMGT numbering system. In certain embodiments, the CDR regions are delineated using the IMGT numbering system accessible at http://www.imgt.org/IMGT_vquest/input.

As used herein, the term “single-chain variable fragment” or “scFv” is a fusion protein of the variable regions of the heavy (V_H) and light chains (V_L) of an immunoglobulin (e.g., mouse or human) covalently linked to form a V_H::V_L heterodimer. The heavy (V_H) and light chains (V_L) are either joined directly or joined by a peptide-
 5 encoding linker (e.g., 10, 15, 20, 25 amino acids), which connects the N-terminus of the V_H with the C-terminus of the V_L, or the C-terminus of the V_H with the N-terminus of the V_L. The linker is usually rich in glycine for flexibility, as well as serine or threonine for solubility. The linker can link the heavy chain variable region and the light chain variable region of the extracellular antigen-binding domain. Non-limiting examples of linkers are
 10 disclosed in Shen et al., Anal. Chem. 80(6):1910-1917 (2008) and WO 2014/087010, the contents of which are hereby incorporated by reference in their entireties. In certain embodiments, the linker is a G4S linker.

In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 13, which is provided below:

15 GGGGSGGGGSGGGGSGGGGS [SEQ ID NO:13]

In certain embodiments, the linker comprise the amino acid sequence set forth in SEQ ID NO: 14, which is provided below:

GGGGSGGGGSGGGGSGGGGS [SEQ ID NO: 14]

In certain embodiments, the linker comprises the amino acid sequence set forth in
 20 SEQ ID NO: 91, which is provided below:

GGGGSGGGGSGGGGSGGGGSGGGGS [SEQ ID NO: 91]

In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 92, which is provided below:

GGGGSGGGGSGGGGSGGGGSGGGGSGGGGS [SEQ ID NO: 92]

25 In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 93, which is provided below:

GGGGS [SEQ ID NO: 93]

In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 94, which is provided below:

30 GGGGSGGGGS [SEQ ID NO: 94]

Despite removal of the constant regions and the introduction of a linker, scFv proteins retain the specificity of the original immunoglobulin. Single chain Fv polypeptide antibodies can be expressed from a nucleic acid comprising V_H - and V_L -encoding sequences as described by Huston, et al. *Proc. Nat. Acad. Sci. USA*,

(1988);85:5879-5883; U.S. Patent Nos. 5,091,513, 5,132,405 and 4,956,778; and U.S. Patent Publication Nos. 20050196754 and 20050196754. Antagonistic scFvs having inhibitory activity have been described (see, e.g., Zhao et al., *Hybridoma* (Larchmt) (2008);27(6):455-51; Peter et al., *J Cachexia Sarcopenia Muscle* (2012);August 12; Shieh et al., *J Immunol* (2009);183(4):2277-85; Giomarelli et al., *Thromb Haemost* (2007);97(6):955-63; Fife et al., *J Clin Invest* (2006);116(8):2252-61; Brocks et al., *Immunotechnology* 1997 3(3):173-84; Moosmayer et al., *Ther Immunol* 1995 2(10:31-40). Agonistic scFvs having stimulatory activity have been described (Peter et al., *J Biol Chem* (2003);278(38):36740-7; Xie et al., *Nat Biotech* 1997 15(8):768-71; Ledbetter et al., *Crit Rev Immunol* (1997);17(5-6):427-55; Ho et al., *Biochim Biophys Acta* (2003);1638(3):257-66).

The term “chimeric antigen receptor” or “CAR” as used herein refers to a molecule comprising an extracellular antigen-binding domain that is fused to an intracellular signaling domain that is capable of activating or stimulating an immunoresponsive cell, and a transmembrane domain. In certain embodiments, the extracellular antigen-binding domain of a CAR comprises a scFv. The scFv can be derived from fusing the variable heavy and light regions of an antibody. Alternatively or additionally, the scFv may be derived from Fab’s (instead of from an antibody, e.g., obtained from Fab libraries). In certain embodiments, the scFv is fused to the transmembrane domain and then to the intracellular signaling domain. By “substantially identical” or “substantially homologous” is meant a polypeptide or nucleic acid molecule exhibiting at least about 50% homologous or identical to a reference amino acid sequence (for example, any of the amino acid sequences described herein) or a reference nucleic acid sequence (for example, any of the nucleic acid sequences described herein). In certain embodiments, such a sequence is at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 99%, or at least about 100% homologous or identical to the sequence of the amino acid or nucleic acid used for comparison.

Sequence identity can be measured by using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, BLAST, BESTFIT, GAP, or PILEUP/PRETTYBOX programs). Such software matches identical or similar sequences by assigning degrees of homology to various

substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. In an exemplary approach to determining the
5 degree of identity, a BLAST program may be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence.

An “effective amount” is an amount sufficient to affect a beneficial or desired clinical result upon treatment. An effective amount can be administered to a subject in one or more doses. In certain embodiments, an effective amount can be an amount that is
10 sufficient to palliate, ameliorate, stabilize, reverse or slow the progression of the disease, or otherwise reduce the pathological consequences of the disease. The effective amount can be determined by a physician on a case-by-case basis and is within the skill of one in the art. Several factors are typically taken into account when determining an appropriate dosage to achieve an effective amount. These factors include age, sex and weight of the
15 subject, the condition being treated, the severity of the condition and the form and effective concentration of the cells administered.

As used herein, the term “endogenous” refers to a nucleic acid molecule or polypeptide that is normally expressed in a cell or tissue.

As used herein, the term “exogenous” refers to a nucleic acid molecule or
20 polypeptide that is not endogenously present in a cell. The term “exogenous” would therefore encompass any recombinant nucleic acid molecule or polypeptide expressed in a cell, such as foreign, heterologous, and over-expressed nucleic acid molecules and polypeptides. By “exogenous” nucleic acid is meant a nucleic acid not present in a native wild-type cell; for example, an exogenous nucleic acid may vary from an endogenous
25 counterpart by sequence, by position/location, or both. For clarity, an exogenous nucleic acid may have the same or different sequence relative to its native endogenous counterpart; it may be introduced by genetic engineering into the cell itself or a progenitor thereof, and may optionally be linked to alternative control sequences, such as a non-native promoter or secretory sequence.

30 By a “heterologous nucleic acid molecule or polypeptide” is meant a nucleic acid molecule (*e.g.*, a cDNA, DNA or RNA molecule) or polypeptide that is not normally present in a cell or sample obtained from a cell. This nucleic acid may be from another

organism, or it may be, for example, an mRNA molecule that is not normally expressed in a cell or sample.

By “modulate” is meant positively or negatively alter. Exemplary modulations include a about 1%, about 2%, about 5%, about 10%, about 25%, about 50%, about 75%,
5 or about 100% change.

By “increase” is meant to alter positively by at least about 5%. An alteration may be by about 5%, about 10%, about 25%, about 30%, about 50%, about 75%, about 100% or more.

By “reduce” is meant to alter negatively by at least about 5%. An alteration may
10 be by about 5%, about 10%, about 25%, about 30%, about 50%, about 75%, or even by about 100%.

The terms “isolated,” “purified,” or “biologically pure” refer to material that is free to varying degrees from components which normally accompany it as found in its native state. “Isolate” denotes a degree of separation from original source or
15 surroundings. “Purify” denotes a degree of separation that is higher than isolation. A “purified” or “biologically pure” protein is sufficiently free of other materials such that any impurities do not materially affect the biological properties of the protein or cause other adverse consequences. That is, a nucleic acid or peptide is purified if it is substantially free of cellular material, viral material, or culture medium when produced by
20 recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Purity and homogeneity are typically determined using analytical chemistry techniques, for example, polyacrylamide gel electrophoresis or high-performance liquid chromatography. The term “purified” can denote that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. For a protein that
25 can be subjected to modifications, for example, phosphorylation or glycosylation, different modifications may give rise to different isolated proteins, which can be separately purified.

By “isolated cell” is meant a cell that is separated from the molecular and/or cellular components that naturally accompany the cell.

30 The term “antigen-binding domain” as used herein refers to a domain capable of specifically binding a particular antigenic determinant or set of antigenic determinants present on a cell.

By “neoplasm” is meant a disease characterized by the pathological proliferation of a cell or tissue and its subsequent migration to or invasion of other tissues or organs. Neoplastic growth is typically uncontrolled and progressive, and occurs under conditions that would not elicit, or would cause cessation of, multiplication of normal cells.

- 5 Neoplasm can affect a variety of cell types, tissues, or organs, including but not limited to an organ selected from the group consisting of bladder, bone, brain, breast, cartilage, glia, esophagus, fallopian tube, gallbladder, heart, intestines, kidney, liver, lung, lymph node, nervous tissue, ovaries, pancreas, prostate, skeletal muscle, skin, spinal cord, spleen, stomach, testes, thymus, thyroid, trachea, urogenital tract, ureter, urethra, uterus, and
- 10 vagina, or a tissue or cell type thereof. Neoplasia include cancers, such as sarcomas, carcinomas, or plasmacytomas (malignant tumor of the plasma cells). The neoplasia can a primary tumor or primary cancer. In addition, the neoplasm can be in metastatic status.

By “receptor” is meant a polypeptide, or portion thereof, present on a cell membrane that selectively binds one or more ligand.

- 15 By “recognize” is meant selectively binds to a target. A T cell that recognizes a tumor can expresses a receptor (*e.g.*, a TCR or CAR) that binds to a tumor antigen.

By “reference” or “control” is meant a standard of comparison. For example, the level of scFv-antigen binding by a cell expressing a CAR and an scFv may be compared to the level of scFv-antigen binding in a corresponding cell expressing CAR alone.

- 20 By “secreted” is meant a polypeptide that is released from a cell via the secretory pathway through the endoplasmic reticulum, Golgi apparatus, and as a vesicle that transiently fuses at the cell plasma membrane, releasing the proteins outside of the cell.

- By “signal sequence” or “leader sequence” is meant a peptide sequence (*e.g.*, 5, 10, 15, 20, 25 or 30 amino acids) present at the N-terminus of newly synthesized proteins
- 25 that directs their entry to the secretory pathway

- By “specifically binds” or “specifically binds to” or “specifically target” is meant a polypeptide or a fragment thereof that recognizes and/or binds to a biological molecule of interest (*e.g.*, a polypeptide, *e.g.*, a CD371 polypeptide), but which does not substantially recognize and/or bind other molecules in a sample, for example, a biological
- 30 sample, which naturally includes a presently disclosed polypeptide (*e.g.*, a CD371 polypeptide).

The terms “comprises”, “comprising”, and are intended to have the broad meaning ascribed to them in U.S. Patent Law and can mean “includes”, “including” and the like.

As used herein, “treatment” refers to clinical intervention in an attempt to alter the disease course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Therapeutic effects of treatment include, without limitation, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastases, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis. By preventing progression of a disease or disorder, a treatment can prevent deterioration due to a disorder in an affected or diagnosed subject or a subject suspected of having the disorder, but also a treatment may prevent the onset of the disorder or a symptom of the disorder in a subject at risk for the disorder or suspected of having the disorder.

An “individual” or “subject” herein is a vertebrate, such as a human or non-human animal, for example, a mammal. Mammals include, but are not limited to, humans, primates, farm animals, sport animals, rodents and pets. Non-limiting examples of non-human animal subjects include rodents such as mice, rats, hamsters, and guinea pigs; rabbits; dogs; cats; sheep; pigs; goats; cattle; horses; and non-human primates such as apes and monkeys. The term “immunocompromised” as used herein refers to a subject who has an immunodeficiency. The subject is very vulnerable to opportunistic infections, infections caused by organisms that usually do not cause disease in a person with a healthy immune system but can affect people with a poorly functioning or suppressed immune system.

Other aspects of the presently disclosed subject matter are described in the following disclosure and are within the ambit of the presently disclosed subject matter.

5.2. CD371

CD371 (CEC12A), also known as DCAL-2, MICL or CLL-1, is a 30 kD C-type lectin transmembrane glycoprotein. It is expressed on monocytes, granulocytes, natural killer (NK) cells, and basophils. CD371 is an immunoinhibitory receptor that recruits Src homology phosphatases SHP-1 and SHP-2 to its phosphorylated cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) (Sancho et al., *Annu Rev Immunol* (2012); 30:491-529; Yan et al., *Front Immunol* (2015);6:408; Lahoud et al., *J Immunol* (2011);187:842). CD371 has been implicated as a negative regulatory uric acid crystals (monosodium urate, MSU) receptor that controls autoimmunity and inflammatory disease (Neumann et al., *Immunity* (2014);40:389-99). CD371 is a negative regulator of

granulocyte and monocyte function (Marshall et al., *J Biol Chem* (2004);279(15):14792-802; Pyz et al., *Eur J Immunol* (2008);38(4):1157-63).

In certain embodiments, CD371 is a human CD371 comprising or consisting of the amino acid sequence with a NCBI Reference No: NP_612210.4 (SEQ ID NO: 15), or
5 a fragment thereof.

SEQ ID NO: 15 is provided below:

10 MSEEVTYADL QFQNSSEMEK IPEIGKFGEK APPAPSHVWR PAALFLTLLC LLLIIGLGVL
ASMFHVTLKI EMKMKMNLQN ISEELQRNIS LQLMSNMINIS NKIRNLSTTL QTIATKLCRE
LYSKEQEHKC KPCPRRWIWH KDSCYFLSDD VQWQESKMA CAAQNASLLK INNKNALFTI
KSQSRSDYDW LGLSPEEDST RGMVRDNIIN SSAWVIRNAP DLNNMYCGYI NRLVQYYHC
TYKKRMICEK MANPVQLGST YFREA [SEQ ID NO: 15]

In certain embodiments, the CD371 comprises or consists of an amino acid sequence that is at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99%, at
15 least about 100% identical to the amino acid sequence set forth in SEQ ID NO: 15 or a fragment thereof.

5.3. *Antigen-Recognizing Receptors*

The presently disclosed antigen-recognizing receptors specifically target or binds to CD371. In certain embodiments, the antigen-recognizing receptor is a chimeric
20 antigen receptor (CAR). In certain embodiments, the antigen-recognizing receptor is a T-cell receptor (TCR). In certain embodiments, the antigen-recognizing receptor is a TCR like fusion molecule.

The presently disclosed subject matter also provides nucleic acid molecules that encode the presently disclosed antigen-recognizing receptors. In certain embodiments,
25 the nucleic acid molecule comprises a nucleotide sequence that encodes a polypeptide of a CD371-targeted antigen recognizing receptor disclosed herein.

5.3.1. T-Cell Receptor (TCR)

In certain embodiments, the antigen-recognizing receptor is a TCR. A TCR is a disulfide-linked heterodimeric protein consisting of two variable chains expressed as part
30 of a complex with the invariant CD3 chain molecules. A TCR found on the surface of T cells is responsible for recognizing antigens as peptides bound to major histocompatibility complex (MHC) molecules. In certain embodiments, a TCR comprises an alpha chain and a beta chain (encoded by TRA and TRB, respectively). In certain embodiments, a TCR comprises a gamma chain and a delta chain (encoded by TRG and TRD,
35 respectively).

Each chain of a TCR is composed of two extracellular domains: Variable (V) region and a Constant (C) region. The Constant region is proximal to the cell membrane, followed by a transmembrane region and a short cytoplasmic tail. The Variable region binds to the peptide/MHC complex. The variable domain of both chains each has three complementarity determining regions (CDRs).

In certain embodiments, a TCR can form a receptor complex with three dimeric signaling modules CD3 δ / ϵ , CD3 γ / ϵ and CD247 ζ / ζ or ζ / η . When a TCR complex engages with its antigen and MHC (peptide/MHC), the T cell expressing the TCR complex is activated.

In certain embodiments, the TCR is an endogenous TCR. In certain embodiments, the antigen-recognizing receptor is naturally occurring TCR.

In certain embodiments, the antigen-recognizing receptor is an exogenous TCR. In certain embodiments, the antigen-recognizing receptor is a recombinant TCR. In certain embodiments, the antigen-recognizing receptor is a non-naturally occurring TCR.

In certain embodiments, the non-naturally occurring TCR differs from any naturally occurring TCR by at least one amino acid residue. In certain embodiments, the non-naturally occurring TCR differs from any naturally occurring TCR by at least about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 20, about 25, about 30, about 40, about 50, about 60, about 70, about 80, about 90, about 100 or more amino acid residues. In certain embodiments, the non-naturally occurring TCR is modified from a naturally occurring TCR by at least one amino acid residue. In certain embodiments, the non-naturally occurring TCR is modified from a naturally occurring TCR by at least about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 20, about 25, about 30, about 40, about 50, about 60, about 70, about 80, about 90, about 100 or more amino acid residues.

5.3.2. Chimeric Antigen Receptor (CAR)

In certain embodiments, the antigen-recognizing receptor is a CAR. CARs are engineered receptors, which graft or confer a specificity of interest onto an immune effector cell. CARs can be used to graft the specificity of a monoclonal antibody onto a T cell; with transfer of their coding sequence facilitated by retroviral vectors.

There are three generations of CARs. "First generation" CARs are typically composed of an extracellular antigen-binding domain (*e.g.*, an scFv), which is fused to a

transmembrane domain, which is fused to cytoplasmic/intracellular signaling domain.

“First generation” CARs can provide *de novo* antigen recognition and cause activation of both CD4⁺ and CD8⁺ T cells through their CD3ζ chain signaling domain in a single fusion molecule, independent of HLA-mediated antigen presentation. “Second generation” CARs add intracellular signaling domains from various co-stimulatory molecules (*e.g.*, CD28, 4-1BB, ICOS, OX40) to the cytoplasmic tail of the CAR to provide additional signals to the T cell. “Second generation” CARs comprise those that provide both co-stimulation (*e.g.*, CD28 or 4-1BB) and activation (CD3ζ). “Third generation” CARs comprise those that provide multiple co-stimulation (*e.g.*, CD28 and 4-1BB) and activation (CD3ζ). In certain embodiments, the antigen-recognizing receptor is a first-generation CAR. In certain embodiments, the antigen-recognizing receptor is a CAR that does not comprise an intracellular signaling domain of a co-stimulatory molecule or a fragment thereof. In certain embodiments, the antigen-recognizing receptor is a second-generation CAR.

In certain embodiments, the CAR comprises an extracellular antigen-binding domain that specifically binds to CD371, a transmembrane domain, and an intracellular signaling domain.

5.3.2.1. Extracellular Antigen-Binding Domain of A CAR

In certain embodiments, the extracellular antigen-binding domain is an scFv. In certain embodiments, the scFv is a human scFv. In certain embodiments, the scFv is a humanized scFv. In certain embodiments, the scFv is a murine scFv. In certain embodiments, the scFv is identified by screening scFv phage library with an antigen-Fc fusion protein.

In certain embodiments, the extracellular antigen-binding domain is a Fab. In certain embodiments, the Fab is crosslinked. In certain embodiments, the extracellular antigen-binding domain is a F(ab)₂.

Any of the foregoing molecules may be comprised in a fusion protein with a heterologous sequence to form the extracellular antigen-binding domain. In certain non-limiting embodiments, the extracellular antigen-binding domain of the CAR (embodied, for example, an scFv or an analog thereof) binds to CD371 (*e.g.*, human CD371) with a dissociation constant (K_d) of about 1×10^{-6} M or less, *e.g.*, about 1×10^{-7} M or less, about 1×10^{-8} M or less, about 1×10^{-9} M or less, about 1×10^{-10} M or less, or about 1×10^{-11} M or less. In certain embodiments, the extracellular antigen-binding domain of the CAR

binds to CD371 (e.g., human CD371) with a K_d of about 1×10^{-7} M or less. In certain embodiments, the extracellular antigen-binding domain of the CAR binds to CD371 (e.g., human CD371) with a K_d of about 1×10^{-8} M or less. In certain embodiments, the extracellular antigen-binding domain of the CAR binds to CD371 (e.g., human CD371) with a K_d of about 1.5×10^{-8} M or about 1×10^{-8} M. In certain embodiments, the extracellular antigen-binding domain of the CAR binds to CD371 (e.g., human CD371) with a K_d of between about 1×10^{-8} M and about 1×10^{-7} M.

In certain embodiments, the extracellular antigen-binding domain of the CAR binds to CD371 (e.g., human CD371) with a low binding affinity. In certain embodiments, the extracellular antigen-binding domain of the CAR binds to CD371 (e.g., human CD371) with a K_d of about 1.5×10^{-8} M or more, about 1×10^{-8} M or more, about 1×10^{-7} M or more, or about 1×10^{-6} M or more.

Binding of the extracellular antigen-binding domain of the CAR can be confirmed by, for example, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), FACS analysis, bioassay (e.g., growth inhibition), or Western Blot assay. Each of these assays generally detect the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody, or a scFv) specific for the complex of interest. For example, the scFv can be radioactively labeled and used in a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a γ counter or a scintillation counter or by autoradiography. In certain embodiments, the CD371-targeted extracellular antigen-binding domain is labeled with a fluorescent marker. Non-limiting examples of fluorescent markers include green fluorescent protein (GFP), blue fluorescent protein (e.g., EBFP, EBFP2, Azurite, and mKalama1), cyan fluorescent protein (e.g., ECFP, Cerulean, and CyPet), and yellow fluorescent protein (e.g., YFP, Citrine, Venus, and YPet). In one embodiment, the CD371-targeted human scFv is labeled with GFP.

In certain embodiments, the CDRs are identified according to the IMGT numbering system.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about

80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 1. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to SEQ ID NO: 1. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 1. SEQ ID NO: 1 is provided in Table 1 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 2. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to SEQ ID NO: 2. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 2. SEQ ID NO: 2 is provided in Table 1 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29 or a conservative modification thereof, and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30 or a conservative modification thereof. SEQ ID NOs: 28-30 are provided in Table 1.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33 or a conservative modification thereof. SEQ ID NOs: 31-33 are provided in Table 1.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29 or a conservative modification thereof,
 5 a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32 or a conservative modification, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33 or a
 10 conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, a
 15 V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 1, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 2. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 107, which is provided in Table 1. In certain embodiments, the scFv is
 25 designated as “B10H4L”.
 30

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is

positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 16. In certain embodiments, the scFv is designated as “B10L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 16 is set forth in SEQ ID NO: 22. SEQ ID NOS: 16 and 22 are provided in Table 1 below.

Table 1

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFSDYQ [SEQ ID NO: 28]	IQGGGGST [SEQ ID NO: 29]	AREMWRGDYYSGMDV [SEQ ID NO: 30]
V _L	QSVLDSYNNENN [SEQ ID NO: 31]	WAS [SEQ ID NO: 32]	QQYTSEPIT [SEQ ID NO: 33]
Full V _H	EVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPGKLEWVSGIQGGGGSTY YADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSGMDVWGQGT TVSS [SEQ ID NO: 1]		
Full V _L	DIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNENNLA WYQQKPGQP PKLLIYWAST RESGVPDRFSGSGSGTDFTLT ISSLQAEDVAVYYCQQYTSEPITFGQGTKVEIK [SEQ ID NO: 2]		
V _L -V _H scFv	DIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNENNLA WYQQKPGQP PKLLIYWAST RESGVPDRFSGSGSGTDFTLT ISSLQAEDVAVYYCQQYTSEPITFGQGTKVEIKGGGGS GGGSGGGSGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPGKGL EWVSGIQGGGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGD YYSGMDVWGQGT TVTVSS [SEQ ID NO: 16]		
DNA for V _L -V _H scFv	GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCAC CATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCTATAACAATGAGAACAATTTAG CTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCATCTACC CGGGAATCCGGGGTCCCTGACCGATTAGTGGCAGCGGTCTGGGACAGATTTCACTCT CACCATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGCAATATACCA GCGAACCTATCACGTTTCGGCCAAGGTACCAAGGTGGAAATCAAAGGTGGTGGTGGTTCA GGTGGTGGTGGTCTGGCGGCGGCTCCGGTGGTGGTGGATCCGAGGTGCAGCTGTTGGA GTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTG GATTCACCTTTAGCGACTATCAGATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTG GAGTGGGTGTCAGGCATTAGGGTGGCGGTGGTAGCACATATTACGCAGACTCCGTGAA GGGCCGGTTCACCATCTCCCGTGACAATCCCAAGAACACGCTGTATCTGCAATGAACA GCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGAC TACTACTCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTCACCGTCTCCTCA [SEQ ID NO: 22]		
V _H -V _L scFv	EVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPGKLEWVSGIQGGGGSTY YADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSGMDVWGQGT TVSSGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNE NLA WYQQKPGQP PKLLIYWAST RESGVPDRFSGSGSGTDFTLT ISSLQAEDVAVYYCQ QYTSEPITFGQGTKVEIK [SEQ ID NO: 107]		

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or

identical to the amino sequence set forth in SEQ ID NO: 3. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 3. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 3. SEQ ID NO: 3 is provided in Table 2 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 4. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 4. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 4. SEQ ID NO: 4 is provided in Table 2 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 3, as shown in Table 2. In certain embodiments, the anti-CD371 scFv comprises a V_L comprising the amino acid sequence set forth in SEQ ID NO: 4. In certain embodiments, the anti-CD371 scFv comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 3 and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 4.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35 or a conservative modification thereof,

and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36 or a conservative modification thereof. SEQ ID NOs: 34-36 are provided in Table 2.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in
 5 SEQ ID NO: 37 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39 or a conservative modification thereof. SEQ ID NOs: 37-39 are provided in Table 2.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 10 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35 or a conservative modification thereof, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set
 15 forth in SEQ ID NO: 37 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 20 (e.g., a scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38, and a V_L CDR3 comprising the
 25 amino acid sequence set forth in SEQ ID NO: 39.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 3, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 4. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the
 30 linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H)

is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 108, which is provided in Table 2. In certain embodiments, the scFv is designated as “C3H4L”.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 17. In certain embodiments, the scFv is designated as “C3L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 17 is set forth in SEQ ID NO: 23. SEQ ID NOS: 17 and 23 are provided in Table 2 below.

Table 2

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFTSYA [SEQ ID NO: 34]	IDGSGGGT [SEQ ID NO: 35]	ARAYYDIL [SEQ ID NO: 36]
V _L	QSVLSSYNNENN [SEQ ID NO: 37]	AAS [SEQ ID NO: 38]	QQYYSEPYT [SEQ ID NO: 39]
Full V _H	EVQLLESGLVQPGGSLRLSCAASGFTFTSYAMSWVRQAPGKLEWVSGIDGSGGGTNYA DSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARAYYDILTGYPDGMDVWGQGTTVTVSS [SEQ ID NO: 3]		
Full V _L	DIVMTQSPDSLAVSLGERATINCKSSQSVLSSYNNENNLAWEYQQKPGQPPKLLIYAASRE SGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYYSEPYTFGQGTKVEIK [SEQ ID NO: 4]		
V _L -V _H scFv	DIVMTQSPDSLAVSLGERATINCKSSQSVLSSYNNENNLAWEYQQKPGQPPKLLIYAASRE SGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYYSEPYTFGQGTKVEIKGGGSGGGG SGGGSGGGGSEVQLLESGLVQPGGSLRLSCAASGFTFTSYAMSWVRQAPGKLEWVSGI DSGGGTNYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARAYYDILTGYPDGMD VWGQGTTVTVSS [SEQ ID NO: 17]		
DNA for V _L -V _H scFv	GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCA TCAACTGCAAGTCCAGCCAGAGTGTTTTAAGCAGCTATAACAATGAGAACAATTTAGCTTG GTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGCCGCATCTACCCGGGAA TCCGGGGTCCCTGACCGATTCACTGGCAGCGGTCTGGGACAGATTTCACTCTCACCATCA GCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGCAATATTTATAGCGAACCTTA TACGTTTCGGCCAAGGTACCAAGGTGGAAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGT TCTGGCGGCGGCTCCGGTGGTGGTGGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCT TGGTACAGCCTGGGGGGTCCCTGCGACTCTCTGTGCAGCCTCTGGATTACCTTTACCAG CTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTGAGGCATT GACGGTAGCGGTGGTGGCACAAATTACGCAGACTCCGTGAAGGGCCGGTTACCATCTCCC GTGACAATTCAGAACACGCTGTATCTGCAATGAACAGCCTGCGTGCCGAGGACACGGC		

	TGTGTATTACTGTGCGAGAGCGTATTACGATATTTTGACTGGTTACCCCGTGGACGGTATG GACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA [SEQ ID NO: 23]
V _H -V _L scFv	EVQLLESQGGGLVQPGGSLRLSCAASGFTFTSYAMSWVRQAPGKGLEWVSGIDGSGGGTNYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARAYYDILTGYFVDGMDVWGQGT TTVT VSSGGGSGGGGSGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLSSYNNENNL AWYQQKPGQPPKLLIYAASSTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQYYSE PYTFGQGTKVEIK [SEQ ID NO: 108]

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 5. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 5.

10 In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 5. SEQ ID NO: 5 is provided in Table 3 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 6. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 6. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 6. SEQ ID NO: 6 is provided in Table 3 below.

25 In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41 or a conservative modification thereof,

and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42 or a conservative modification thereof. SEQ ID NOS: 40-42 are provided in Table 3.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in
 5 SEQ ID NO: 43 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45 or a conservative modification thereof. SEQ ID NOS: 43-45 are provided in Table 3.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 10 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41 or a conservative modification thereof, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set
 15 forth in SEQ ID NO: 43 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 20 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44, and a V_L CDR3
 25 comprising the amino acid sequence set forth in SEQ ID NO: 45.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 5, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 6. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the
 30 linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H)

is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 109, which is provided in Table 3. In certain embodiments, the scFv is designated as “D6H4L”.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 18. In certain embodiments, the scFv is designated as “D6L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 18 is set forth in SEQ ID NO: 24. SEQ ID NOS: 18 and 24 are provided in Table 3 below.

Table 3

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFTDYA [SEQ ID NO: 40]	IDSGGGST [SEQ ID NO: 41]	ALELGATTVY [SEQ ID NO: 42]
V _L	QSVLRSSNNKNN [SEQ ID NO: 43]	AAS [SEQ ID NO: 44]	QQYYREPLT [SEQ ID NO: 45]
Full V _H	EVQLLESGGGLVQPGGSLRLSCAASGFTFTDYAMSWVRQAPGKGLEWVSDIDSGGGSTDYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCALELGATTVYWGQGT LVTVSS [SEQ ID NO: 5]		
Full V _L	DIVMTQSPDSLAVSLGERATINCKSSQSVLRSSNNKNNLAWYQQKPGQP PKLLIYAASTRE SGVPDRFSGSGSGTDFTLTIS SLQAEDVAVYYCQQYYREPLTFGQGTKVEIK [SEQ ID NO: 6]		
V _L -V _H scFv	DIVMTQSPDSLAVSLGERATINCKSSQSVLRSSNNKNNLAWYQQKPGQP PKLLIYAASTRE SGVPDRFSGSGSGTDFTLTIS SLQAEDVAVYYCQQYYREPLTFGQGTKVEIKGGGGSGGGG SGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFTDYAMSWVRQAPGKGLEWVSDI DSGGGSTDYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCALELGATTVYWGQGT LVTVSS [SEQ ID NO: 18]		
DNA for V _L -V _H scFv	GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCA TCAACTGCAAGTCCAGCCAGAGTGTTTTACGCAGCAGCAACAATAAAAACAATTTAGCTTG GTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTACGCCGCATCTACCCGGGAA TCCGGGGTCCCTGACCGATTCACTGGCAGCGGTCTGGGACAGATTTCACTCTCACCATCA GCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGCAATATTTATCGCGAACCTCT GACGTTTCGGCCAAGGTACCAAGGTGGAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGT TCTGGCGGCGGCTCCGGTGGTGGTGGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCT TGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTACCGA CTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTGAGACATT GACGGTAGCGGTGGTAGCACAGACTACGCAGACTCCGTGAAGGGCCGGTTACCATCTCCC GTGACAATTC AAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGC		

	TGTGTATTACTGTGCGCTAGAGCTGGGAGCTACTACCGTCTACTGGGGCCAGGGAACCTGTGTCACCGTCTCCTCA [SEQ ID NO: 24]
V _H -V _L scFv	EVQLLESQGGGLVQPGGSLRLSCAASGFTFTDYAMSWVRQAPGKGLEWVSDIDGSGGSTDYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCALELGATTVYWGQGT LVT VSSGGGGS GGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLRSSNNKNNLAWYQQKPG QPPKLLIYAASSTRESGVPDRFSGSGSTDFTLTISSLQAEDVAVYYCQYYREPLTFGQGT KVEIK [SEQ ID NO: 109]

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 7. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 7.

10 In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 7. SEQ ID NO: 7 is provided in Table 4 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 8. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 8. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 8. SEQ ID NO: 8 is provided in Table 4 below.

25 In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 46 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47 or a conservative modification thereof,

and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48 or a conservative modification thereof. SEQ ID NOs: 46-48 are provided in Table 4.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in
 5 SEQ ID NO: 49 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 50 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51 or a conservative modification thereof. SEQ ID NOs: 49-51 are provided in Table 4.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 10 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 46 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47 or a conservative modification thereof, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set
 15 forth in SEQ ID NO: 49 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 50 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 20 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 46, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 49, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 50, and a V_L CDR3
 25 comprising the amino acid sequence set forth in SEQ ID NO: 51.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 7, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 8. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the
 30 linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H)

is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 110, which is provided in Table 4. In certain embodiments, the scFv is designated as “A11H4L”.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 19. In certain embodiments, the scFv is designated as “A11L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 19 is set forth in SEQ ID NO: 25. SEQ ID NOS: 19 and 25 are provided in Table 4 below.

Table 4

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFTSTQ [SEQ ID NO: 46]	ISGYGGST [SEQ ID NO: 47]	AKDTEVSGDAFDI [SEQ ID NO: 48]
V _L	QSVDDSN [SEQ ID NO: 49]	GAS [SEQ ID NO: 50]	QQYRSWPIT [SEQ ID NO: 51]
Full V _H	EVQLLESGGGLVQPGGSLRLSCAASGFTFTSTQMSWVRQAPGKGLEWVSEISGYGGSTYYA DSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKDTEVSGDAFDIWGQGTMTVSS [SEQ ID NO: 7]		
Full V _L	EIVLTQSPGTLSPGERATLSCRASQSVDDSNLAWYQQKPGQAPRLLIYGASSRATGIPD RFSGSGSGTDFTLTISRLEPEDFAVYYCQQYRSWPITFGQGTKVEIK [SEQ ID NO: 8]		
V _L -V _H scFv	EIVLTQSPGTLSPGERATLSCRASQSVDDSNLAWYQQKPGQAPRLLIYGASSRATGIPD RFSGSGSGTDFTLTISRLEPEDFAVYYCQQYRSWPITFGQGTKVEIKGGGGSGGGSGGGG GGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFTSTQMSWVRQAPGKGLEWVSEISGYGG STYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKDTEVSGDAFDIWGQGTMTV VSS [SEQ ID NO: 19]		
DNA for V _L -V _H scFv	GAAATTGTGTTGACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAACGTGCCACCC TCTCCTGCCGTGCCAGTCAGAGTGTTGACAGCAGCAATTTAGCCTGGTATCAGCAGAAACC TGGCCAGGCTCCCCGACTCCTCATCTATGGCGCATCTAGCCGTGCCACTGGTATCCCAGAC CGTTTCAGTGGCAGTGGGTCTGGGACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTG AAGATTTTGCAGTGTATTACTGTCTCAGCAGTATCGCAGCTGGCCTATCACGTTTCGGCCAAGG TACCAAGGTGGAAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGTCTGGCGGCGGCTCC GGTGGTGGTGGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGG GGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTACCAGCACCCAGATGAGCTG GGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCTAGAGATTAGCGGTTATGGTGGT AGCACATACTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAATTCGAAGA ACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGC		

	AAAAGACACGGAGGTTTCGGGAGATGCTTTTGATATCTGGGGCCAAGGGACAATGGTCACC GTCTCTTCA [SEQ ID NO: 25]
V _H -V _L scFv	EVQLLESQGGGLVQPGGSLRLSCAASGFTFTSTQMSWVRQAPGKGLEWVSEISGYGGSTYYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKDTEVSGDAFDIWGGQTMVTVSSGG GGSGGGGSGGGSGGGSEIVLTQSPGTLSPGERATLSCRASQSVDSNLAQYQQKPGQA PRLIYGAASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQYRSWPITFGQGTKV EIK [SEQ ID NO: 110]

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 9. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 9.

10 In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 9. SEQ ID NO: 9 is provided in Table 5 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 10. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 10. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 10. SEQ ID NO: 10 is provided in Table 5 below.

25 In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53 or a conservative modification thereof,

and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54 or a conservative modification thereof. SEQ ID NOs: 52-54 are provided in Table 5.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in
 5 SEQ ID NO: 55 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57 or a conservative modification thereof. SEQ ID NOs: 55-57 are provided in Table 5.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 10 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53 or a conservative modification thereof, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set
 15 forth in SEQ ID NO: 55 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 20 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 55, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56, and a V_L CDR3
 25 comprising the amino acid sequence set forth in SEQ ID NO: 57.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 9, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 10. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the
 30 linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H)

is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 111, which is provided in Table 5. In certain embodiments, the scFv is designated as “E4H4L”.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 20. In certain embodiments, the scFv is designated as “E4L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 20 is set forth in SEQ ID NO: 26. SEQ ID NOS: 20 and 26 are provided in Table 5 below.

Table 5

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFTSY [SEQ ID NO: 52]	ISGSGDST [SEQ ID NO: 53]	AREAGGDYDSGAFDI [SEQ ID NO: 54]
V _L	QSVLYSGNNKNY [SEQ ID NO: 55]	GAS [SEQ ID NO: 56]	QQYDYAPFT [SEQ ID NO: 57]
Full V _H	EVQLLES GGG LVQ PGGSLRLSCAASGFTFTSYMSWVRQAPGKGLEWVSGISGSGDSTSYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREAGGDYDSGAFDIWGQGTMTVSS [SEQ ID NO: 9]		
Full V _L	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSGNNKNYLAWYQQKPGQPPKLLIYGASTRE SGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYDYAPFTFGQGTKVEIK [SEQ ID NO: 10]		
V _L -V _H scFv	DIVMTQSPDSLAVSLGERATINCKSSQSVLYSGNNKNYLAWYQQKPGQPPKLLIYGASTRE SGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYDYAPFTFGQGTKVEIKGGGGSGGGG SGGGSGGGGSEVQLLES GGG LVQ PGGSLRLSCAASGFTFTSYMSWVRQAPGKGLEWVSGI SSGSDSTSYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREAGGDYDSGAFDIW GQGTMTVSS [SEQ ID NO: 20]		
DNA for V _L -V _H scFv	GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCA TCAACTGCAAGTCCAGCCAGAGTGTTTTATATAGCGGCAACAATAAAACTATTTAGCTTG GTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGGCGCATCTACCCGGGAA TCCGGGGTCCCTGACCGATTCACTGGCAGCGGTCTGGGACAGATTTCACTCTCACCATCA GCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATGACTATGCCCTTT TACGTTTCGGCCAAGGTACCAAGGTGGAAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGT TCTGGCGGCGGCTCCGGTGGTGGTGGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCT TGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTACCAG CTATTATATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGCATT AGCGGTAGCGGTGACAGCACAAGCTACGCAGACTCCGTGAAGGGCCGGTTACCATCTCCC GTGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGC		

	TGTGTATTACTGTGCGAGAGAGGCAGGTGGTGACTACGATAGTGGTGCTTTTGATATCTGG GGCCAAGGGACAATGGTCACCGTCTCTTCA [SEQ ID NO: 26]
V _H -V _L scFv	EVQLLESQGGGLVQPGGSLRLSCAASGFTFTSYMSWVRQAPGKGLEWVSGISGSGDSTSYA DSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREAGGDYDSGAFDIWGQGTMTVTVSS GGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLYSGNNKNYLAWY QQKPGQPPKLLIYGASTRESGVPRDFSGSGSGTDFTLTISLQAEDVAVYYCQDYDYAPFT FGQGTKVEIK [SEQ ID NO: 111]

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is at least about 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 11. For example, the

5 extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth

10 in SEQ ID NO: 11. In certain embodiments, the extracellular antigen-binding domain comprises a V_H comprising the amino sequence set forth in SEQ ID NO: 11. SEQ ID NO: 11 is provided in Table 6 below.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is at least about

15 80% (e.g., at least about 85%, at least about 90%, or at least about 95%) homologous or identical to the amino sequence set forth in SEQ ID NO: 12. For example, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L comprising an amino acid sequence that is about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%,

20 about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino sequence set forth in SEQ ID NO: 12. In certain embodiments, the extracellular antigen-binding domain comprises a V_L comprising the amino sequence set forth in SEQ ID NO: 12. SEQ ID NO: 12 is provided in Table 6 below.

25 In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59 or a conservative modification thereof,

and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60 or a conservative modification thereof. SEQ ID NOs: 58-60 are provided in Table 6.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_L CDR1 comprising the amino acid sequence set forth in
 5 SEQ ID NO: 61 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63 or a conservative modification thereof. SEQ ID NOs: 61-63 are provided in Table 6.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 10 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58 or a conservative modification thereof, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59 or a conservative modification thereof, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60 or a conservative modification thereof, a V_L CDR1 comprising the amino acid sequence set
 15 forth in SEQ ID NO: 61 or a conservative modification thereof, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62 or a conservative modification thereof, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63 or a conservative modification thereof.

In certain embodiments, the extracellular antigen-binding domain of the CAR
 20 (e.g., an scFv) comprises a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59, a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60, a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 61, a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62, and a V_L CDR3
 25 comprising the amino acid sequence set forth in SEQ ID NO: 63.

In certain embodiments, the extracellular antigen-binding domain of the CAR (e.g., an scFv) comprises a V_H comprising the amino acid sequence set forth in SEQ ID NO: 11, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 12. In certain embodiments, the V_H and V_L are linked via a linker. In certain embodiments, the
 30 linker comprises the amino acid sequence set forth in SEQ ID NO: 13.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H)

is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the scFv comprises the amino acid sequence set forth in SEQ ID NO: 112, which is provided in Table 6. In certain embodiments, the scFv is designated as “E8H4L”.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, scFv comprises the amino acid sequence set forth in SEQ ID NO: 21. In certain embodiments, the scFv is designated as “E8L4H”. An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 21 is set forth in SEQ ID NO: 27. SEQ ID NOS: 21 and 27 are provided in Table 6 below.

Table 6

Antigen	A CD371 polypeptide consisting of the amino acid sequence of SEQ ID NO: 15 or a fragment thereof		
CDRs	1	2	3
V _H	GFTFSSYA [SEQ ID NO: 58]	IDGEGGYT [SEQ ID NO: 59]	AREGVDDYDILTGYYPYGMDV [SEQ ID NO: 60]
V _L	QSVLDSSNNKNY [SEQ ID NO: 61]	DAS [SEQ ID NO: 62]	QQGTSSPLT [SEQ ID NO: 63]
Full V _H	EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSEIDGEGGYTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREGVDDYDILTGYYPYGMDVWGQGTITVTVSS [SEQ ID NO: 11]		
Full V _L	DIVMTQSPDSLAVSLGERATINCKSSQSVLDSSNNKNYLAWYQQKPGQPPKLLIYDASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQGTSSPLTFGQGTKVEIK [SEQ ID NO: 12]		
V _L -V _H scFv	DIVMTQSPDSLAVSLGERATINCKSSQSVLDSSNNKNYLAWYQQKPGQPPKLLIYDASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQGTSSPLTFGQGTKVEIKGGGGSGGGSGGGSGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKLEWVSEIDGEGGYTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREGVDDYDILTGYYPYGMDVWGQGTITVTVSS [SEQ ID NO: 21]		
DNA for V _L -V _H scFv	GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCAGCAACAATAAAAACATATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGACGCATCTACCCGGGAATCCGGGGTCCCTGACCGATTCAAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGTCAGCAAGGCACCAGCAGCCCTCTGACGTTCCGCCAAGGTACCAAGGTGGAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGTCTGGCGCGGCTCCGGTGGTGGTGGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTGAGAGATTGACGGTGAGGGTGGTTATACAAATTACGCAGACTCCGTGAAGGGCCGGTTACCATCTCCCGTGACAATTC AAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCCGTGTATTACT		

	GTGCGAGAGAAGGGGTAGATTACGATATTTTGACTGGTTATTATCCTTACGGTATGGACGTC TGGGGCCAAGGGACCACGGTCACCGTCTCCTCA [SEQ ID NO: 27]
V _H -V _L scFv	EVQLLESQGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSEIDEGGYTNYAD SVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREGVVDYDILTGYYPYGMVDVWGQGT TVTVSSGGGGSGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSSNNK NYLA WYQQKPGQPPKLLIYDASTRESGVPRFSGSGSGTDFTLTISSLQAEDVAVYYCQQTSSPL TFGQGTKVEIK [SEQ ID NO: 112]

As used herein, the term “a conservative sequence modification” refers to an amino acid modification that does not significantly affect or alter the binding characteristics of the presently disclosed mesothelin-targeted CAR (*e.g.*, the extracellular antigen-binding domain of the CAR) comprising the amino acid sequence. Conservative modifications can include amino acid substitutions, additions and deletions. Modifications can be introduced into the extracellular antigen-binding domain of the presently disclosed CAR by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Amino acids can be classified into groups according to their physicochemical properties such as charge and polarity. Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid within the same group. For example, amino acids can be classified by charge: positively-charged amino acids include lysine, arginine, histidine, negatively-charged amino acids include aspartic acid, glutamic acid, neutral charge amino acids include alanine, asparagine, cysteine, glutamine, glycine, isoleucine, leucine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine. In addition, amino acids can be classified by polarity: polar amino acids include arginine (basic polar), asparagine, aspartic acid (acidic polar), glutamic acid (acidic polar), glutamine, histidine (basic polar), lysine (basic polar), serine, threonine, and tyrosine; non-polar amino acids include alanine, cysteine, glycine, isoleucine, leucine, methionine, phenylalanine, proline, tryptophan, and valine. Thus, one or more amino acid residues within a CDR region can be replaced with other amino acid residues from the same group and the altered antibody can be tested for retained function (*i.e.*, the functions set forth in (c) through (l) above) using the functional assays described herein. In certain embodiments, no more than one, no more than two, no more than three, no more than four, no more than five residues within a specified sequence or a CDR region are altered.

The V_H and/or V_L amino acid sequences having at least about 80%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% (*e.g.*, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about

97%, about 98%, or about 99%) homology or identity to a specific sequence (*e.g.*, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12) may contain substitutions (*e.g.*, conservative substitutions), insertions, or deletions relative to the specified sequence(s), but retain the ability to bind to a target antigen (*e.g.*, mesothelin). In certain embodiments, a total of 1 to 10 amino acids are substituted, inserted and/or deleted in a specific sequence (*e.g.*, SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, or SEQ ID NO: 12). In certain embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (*e.g.*, in the FRs) of the extracellular antigen-binding domain. In certain embodiments, the extracellular antigen-binding domain comprises V_H and/or V_L sequence selected from SEQ ID NOs: 1-12, including post-translational modifications of that sequence (SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12).

As used herein, the percent homology between two amino acid sequences is equivalent to the percent identity between the two sequences. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions × 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm.

The percent homology between two amino acid sequences can be determined using the algorithm of E. Meyers and W. Miller (Comput. Appl. Biosci., 4:11-17 (1988)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent homology between two amino acid sequences can be determined using the Needleman and Wunsch (J. Mol. Biol. 48:444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at www.gcg.com), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

Additionally or alternatively, the amino acids sequences of the presently disclosed subject matter can further be used as a “query sequence” to perform a search against

public databases to, for example, identify related sequences. Such searches can be performed using the XBLAST program (version 2.0) of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the specified sequences (*e.g.*, heavy and light chain variable region sequences of scFv m903, m904, m905, m906, and m900) disclosed herein. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) Nucleic Acids Res. 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used.

In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with a reference antibody or an antigen-binding fragment thereof comprising the V_H CDR1, CDR2, and CDR3 sequences and the V_L CDR1, CDR2, and CDR3 sequences of, for example, any one of the presently disclosed scFvs (*e.g.*, V10, C3, D6, A11, E4, and D8). In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with a reference antibody or an antigen-binding portion thereof comprising the V_H and V_L sequences of, for example, any one of the presently disclosed scFvs (*e.g.*, V10, C3, D6, A11, E4, and D8).

In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with a reference antibody or an antigen-binding portion thereof comprising the V_H CDR1, CDR2, and CDR3 sequences and the V_L CDR1, CDR2, and CDR3 sequences of scFv B10. For example, the extracellular antigen-binding domain of a presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with a reference antibody or an antigen-binding portion thereof comprising a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28, a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a V_L CDR2 comprising amino acids having the sequence set forth in SEQ ID NO: 32; and a V_L CDR3 comprising amino acids having the sequence set forth in SEQ ID NO: 33. In certain embodiments, the extracellular antigen-binding domain of a

presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with a reference antibody or an antigen-binding portion thereof comprising the V_H and V_L sequences of scFv B10. For example, the extracellular antigen-binding domain of a presently disclosed CAR cross-competes for binding to CD371 (*e.g.*, human CD371) with
 5 a reference antibody or an antigen-binding portion thereof comprising a V_H comprising amino acids having the sequence set forth in SEQ ID NO: 1, and a V_L comprising amino acids having the sequence set forth in SEQ ID NO: 2.

In certain embodiments, the extracellular antigen-binding domain binds to the same epitope on CD371 (*e.g.*, human CD371) as the reference antibody or antigen-
 10 binding portion thereof. For example, the extracellular antigen-binding domain of a presently disclosed CAR binds to the same epitope on CD371 (*e.g.*, human CD371) as a reference antibody or an antigen-binding portion thereof comprising the V_H CDR1, CDR2, and CDR3 sequences and the V_L CDR1, CDR2, and CDR3 sequences of, for example, any one of the presently disclosed scFvs (*e.g.*, B10, C3, D6, A11, E4, and E8).
 15 In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR binds to the same epitope on EMR2 (*e.g.*, human EMR2) as a reference antibody or an antigen-binding portion thereof comprising the V_H and V_L sequences of, for example, any one of the presently disclosed scFvs (*e.g.*, B10, C3, D6, A11, E4, and E8).

In certain embodiments, the extracellular antigen-binding domain of a presently
 20 disclosed CAR binds to the same epitope on CD371 (*e.g.*, human CD371) as a reference antibody or an antigen-binding fragment thereof comprising the V_H CDR1, CDR2, and CDR3 sequences and the V_L CDR1, CDR2, and CDR3 sequences of scFv B10. For example, the extracellular antigen-binding domain of a presently disclosed CAR binds to the same epitope on CD371 (*e.g.*, human CD371) as a reference antibody or an antigen-
 25 binding fragment thereof comprising a V_H CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a V_H CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a V_L CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a V_L CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a V_L
 30 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33. In certain embodiments, the extracellular antigen-binding domain of a presently disclosed CAR binds to the same or substantially the same epitope on CD371 (*e.g.*, human CD371) as a reference antibody or an antigen-binding fragment thereof comprising the V_H and V_L

sequences of scFv B10. For example, the extracellular antigen-binding domain of a presently disclosed CAR binds to the same epitope on CD371 (*e.g.*, human CD371) as a reference antibody or an antigen-binding fragment thereof comprising a V_H comprising the amino acid sequence set forth in SEQ ID NO: 1, and a V_L comprising the amino acid sequence set forth in SEQ ID NO: 2.

Extracellular antigen-binding domains that cross-compete or compete with the reference antibody or antigen-binding portions thereof for binding to CD371 (*e.g.*, human CD371) can be identified by using routine methods known in the art, including, but not limited to, ELISAs, radioimmunoassays (RIAs), Biacore, flow cytometry, Western blotting, and any other suitable quantitative or qualitative antibody-binding assays. Competition ELISA is described in Morris, "Epitope Mapping of Protein Antigens by Competition ELISA", *The Protein Protocols Handbook* (1996), pp 595-600, edited by J. Walker, which is incorporated by reference in its entirety. In certain embodiments, the antibody-binding assay comprises measuring an initial binding of a reference antibody to a CD371 polypeptide, admixing the reference antibody with a test extracellular antigen-binding domain, measuring a second binding of the reference antibody to the CD371 polypeptide in the presence of the test extracellular antigen-binding domain, and comparing the initial binding with the second binding of the reference antibody, wherein a decreased second binding of the reference antibody to the CD371 polypeptide in comparison to the initial binding indicates that the test extracellular antigen-binding domain cross-competes with the reference antibody for binding to CD371, *e.g.*, one that recognizes the same or substantially the same epitope, an overlapping epitope, or an adjacent epitope. In certain embodiments, the reference antibody is labeled, *e.g.*, with a fluorochrome, biotin, or peroxidase. In certain embodiments, the CD371 polypeptide is expressed in cells, *e.g.*, in a flow cytometry test. In certain embodiments, the CD371 polypeptide is immobilized onto a surface, including a Biacore chip (*e.g.*, in a Biacore test), or other media suitable for surface plasmon resonance analysis. The binding of the reference antibody in the presence of a completely irrelevant antibody (that does not bind to CD371) can serve as the control high value. The control low value can be obtained by incubating a labeled reference antibody with an unlabeled reference antibody, where competition and reduced binding of the labeled reference antibody would occur. In certain embodiments, a test extracellular antigen-binding domain that reduces the binding of the reference antibody to a CD371 polypeptide by at least about 20%, at least about

30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% is considered to be an extracellular antigen-binding domain that cross-competes with the reference antibody for binding to CD371. In certain embodiments, the assays are performed at room temperature.

In certain embodiments, the antibody-binding assay comprises measuring an initial binding of a test extracellular antigen-binding domain to a CD371 polypeptide, admixing the test extracellular antigen-binding domain with a reference antibody, measuring a second binding of the test extracellular antigen-binding domain to the CD371 polypeptide in the presence of the reference antibody, and comparing the initial binding with the second binding of the test extracellular antigen-binding domain, where a decreased second binding of the test extracellular antigen-binding domain to the CD371 polypeptide in comparison to the initial binding indicates that the test extracellular antigen-binding domain cross-competes with the reference antibody for binding to CD371, *e.g.*, one that recognizes the same or substantially the same epitope, an overlapping epitope, or an adjacent epitope. In certain embodiments, the test extracellular antigen-binding domain is labeled, *e.g.*, with a fluorochrome, biotin, or peroxidase. In certain embodiments, the CD371 polypeptide is expressed in cells, *e.g.*, in a flow cytometry test. In certain embodiments, the CD371 polypeptide is immobilized onto a surface, including a Biacore chip (*e.g.*, in a Biacore test), or other media suitable for surface plasmon resonance analysis. The binding of the test extracellular antigen-binding domain in the presence of a completely irrelevant antibody (that does not bind to CD371) can serve as the control high value. The control low value can be obtained by incubating a labeled test extracellular antigen-binding domain with an unlabeled test extracellular antigen-binding domain, where competition and reduced binding of the labeled test extracellular antigen-binding domain would occur. In certain embodiments, a test extracellular antigen-binding domain, whose binding to a CD371 polypeptide is decreased by at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% in the presence of a reference antibody, is considered to be an extracellular antigen-binding domain that cross-competes with the reference antibody for binding to CD371. In certain embodiments, the assays are performed at room temperature.

In certain non-limiting embodiments, the extracellular antigen-binding domain of the presently disclosed CAR comprises a linker connecting the heavy chain variable region and light chain variable region of the extracellular antigen-binding domain. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 13. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 91. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 92. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 93. In certain embodiments, the linker comprises the amino acid sequence set forth in SEQ ID NO: 94.

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a heavy chain variable region (V_H) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_H - V_L .

In certain embodiments, the variable regions within the extracellular antigen-binding domain of the CAR have to be linked one after another such that at the N-terminus of the extracellular antigen-binding domain, a light chain variable region (V_L) is positioned. In certain embodiments, if the extracellular antigen-binding domain of the CAR is an scFv, the variable regions are positioned from the N- to the C-terminus: V_L - V_H .

In addition, the extracellular antigen-binding domain can comprise a leader or a signal peptide that directs the nascent protein into the endoplasmic reticulum. Signal peptide or leader can be essential if the CAR is to be glycosylated and anchored in the cell membrane. The signal sequence or leader can be a peptide sequence (about 5, about 10, about 15, about 20, about 25, or about 30 amino acids long) present at the N-terminus of newly synthesized proteins that directs their entry to the secretory pathway. In certain embodiments, the signal peptide is covalently joined to the 5' terminus of the extracellular antigen-binding domain. In certain embodiments, the signal peptide comprises a CD8 polypeptide, e.g., the CAR comprises a truncated CD8 signal peptide.

5.3.2.2. Transmembrane Domain of a CAR

In certain non-limiting embodiments, the transmembrane domain of the CAR comprises a hydrophobic alpha helix that spans at least a portion of the membrane. Different transmembrane domains result in different receptor stability. After antigen recognition, receptors cluster and a signal are transmitted to the cell. In accordance with

5 the presently disclosed subject matter, the transmembrane domain of the CAR can comprise a native or modified transmembrane domain of CD8 or a fragment thereof, a native or modified transmembrane domain of CD28 or a fragment thereof, a native or modified transmembrane domain of CD3 ζ or a fragment thereof, a native or modified transmembrane domain of CD4 or a fragment thereof, a native or modified

10 transmembrane domain of 4-1BB or a fragment thereof, a native or modified transmembrane domain of OX40 or a fragment thereof, a native or modified transmembrane domain of ICOS or a fragment thereof, a native or modified transmembrane domain of CD84 or a fragment thereof, a native or modified transmembrane domain of CD166 or a fragment thereof, a native or modified

15 transmembrane domain of CD8a or a fragment thereof, a native or modified transmembrane domain of CD8b or a fragment thereof, a native or modified transmembrane domain of ICAM-1 or a fragment thereof, a native or modified transmembrane domain of CTLA-4 or a fragment thereof, a native or modified transmembrane domain of CD27 or a fragment thereof, a native or modified

20 transmembrane domain of CD40 or a fragment thereof, NKGD2 or a fragment thereof, or a combination thereof.

In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide (e.g., a transmembrane domain of CD8 or a fragment thereof).

In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide (e.g., a transmembrane domain of CD8 or a fragment thereof). In certain

25 polypeptide (e.g., a transmembrane domain of CD8 or a fragment thereof). In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide (e.g., a transmembrane domain of human CD8 or a fragment thereof). In certain embodiments, the CD8 polypeptide comprises or consists of an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or

30 about 100% homologous or identical to the amino acid sequence having a NCBI Reference No: NP_001139345.1 (SEQ ID NO: 64) or a fragments thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In certain embodiments, the CD8 polypeptide comprises or consists of an

amino acid sequence that is a consecutive portion of SEQ ID NO: 64, which is at least 20, or at least 30, or at least 40, or at least 50, and up to 235 amino acids in length.

Alternatively or additionally, in non-limiting various embodiments, the CD8 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 235, 1 to 50, 50 to 100, 100 to 150, 150 to 200, 137 to 209 or 200 to 235 of SEQ ID NO: 64. In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide comprising or consisting of amino acids 137 to 209 of SEQ ID NO: 64. SEQ ID NO: 64 is provided below.

10 MALPVTALLLPLALLLHAARPSQFRVSPLDRTWNLGETVELKQCQVLLSNPTSGCSWLFQPRGAAASPTFLLY
LSQNKPKAAEGLDTQRFSGKRLGDTFVLTLSDFRRENEGYYFCSALSNSIMYFSHFVPVFLPAKPTTTPAPR
PPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVITLYCNHRNRRRVCK
CPRPVVKS GDKPSLSARYV [SEQ ID NO: 64]

In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide (e.g., a transmembrane domain of mouse CD8 or a fragment thereof). In certain embodiments, the CD8 polypeptide comprises or consists of an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino acid sequence having a NCBI Reference No: AAA92533.1 (SEQ ID NO: 65) or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In certain embodiments, the CD8 polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 65, which is at least about 20, or at least about 30, or at least about 40, or at least about 50, or at least about 60, or at least about 70, or at least about 100, or at least about 200, and up to 247 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD8 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 247, 1 to 50, 50 to 100, 100 to 150, 150 to 200, 151 to 219, or 200 to 247 of SEQ ID NO: 65. In certain embodiments, the transmembrane domain of the CAR comprises a CD8 polypeptide comprising or consisting of amino acids 151 to 219 of SEQ ID NO: 65. SEQ ID NO: 65 is provided below.

30 1 MASPLTRFLS LNLLLMGESI ILGSGEAKPQ APELRIFPKK MDAELGQKVD LVCEVLGSVS
61 QGCSWLFQNS SSKLPQPTFV VYMASSHNI TWDEKLNSSK LFSVRDNTN KYVLTLNKF
121 KENEGYYFCS VISNSVMYFS SVVPVLQKVN STTKPVLRT PSPVHPTGTS QPQRPEDCRP
181 RGSVKGTGLD FACDIYIWAP LAGICVAPLL SLIITLICYH RSRKRVCKCP RPLVRQEGKP
241 RPSEKIV [SEQ ID NO: 65]

35 In certain embodiments, the transmembrane domain of a presently disclosed CAR comprises a CD28 polypeptide (e.g., a transmembrane domain of CD28 or a fragment thereof).

In certain embodiments, the transmembrane domain of the CAR comprises a CD28 polypeptide (e.g., a transmembrane domain of human CD28 or a fragment thereof). In certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100% homologous or identical to the amino acid sequence having a NCBI Reference No: NP_006130 (SEQ ID No: 66) or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 66 which is at least 20, or at least 30, or at least 40, or at least 50, and up to 220 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 220, 1 to 50, 50 to 100, 100 to 150, 150 to 200, 153 to 179, or 200 to 220 of SEQ ID NO: 66. In certain embodiments, the transmembrane domain of the CAR comprises a CD28 polypeptide comprising or consisting of amino acids 153 to 179 of SEQ ID NO: 66. SEQ ID NO: 66 is provided below:

```

1  MLRLLLLALNL FPSIQVTGNK ILVKQSPMLV AYDNAVNLSC KYSYNLFSRE FRASLHKGLD
61  SAVEVCVVYG NYSQQLQVYS KTGFNCDGKL GNESVTFYLQ NLYVNQTDIY FCKIEVMYPP
121 PYLDNEKSNG TIIHVKGKHL CPSPLFPGPS KFEWVLVVVG GVLACYSLLV TVAFIIFWVR
181 SKRSRL LHSD YMNMTFRRPG PTRKHYQPYA PPRDFAAYRS [SEQ ID NO: 66]

```

An exemplary nucleotide sequence encoding amino acid 153 to 179 of SEQ ID NO: 66 is set forth in SEQ ID NO: 67, which is provided below.

```

TTTTGGGTGCTGGTGGTGGTGGTGGAGTCTGCTTGTCTATAGCTTGCTAGTAACAGTGGCCTTTATTATT
TTCTGGGTG [SEQ ID NO: 67]

```

In certain embodiments, the transmembrane domain of the CAR comprises a CD28 polypeptide (e.g., a transmembrane domain of mouse CD28 or a fragment thereof). In certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or 100% homologous or identical to the amino acid sequence having a NCBI Reference No: NP_031668.3 (SEQ ID No: 68) or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 66 which is at least 20, or at least 30, or at least 40, or at least 50, and up to 218 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 220, 1 to

50, 50 to 100, 100 to 150, 150 to 200, 151 to 177, or 200 to 218 of SEQ ID NO: 68. In certain embodiments, the transmembrane domain of the CAR comprises a CD28 polypeptide comprising or consisting of amino acids 151 to 177 of SEQ ID NO: 68. SEQ ID NO: 68 is provided below:

5 1 MTLRLLFLAL NFFSVQVTEN KILVKQSPLL VVDSNEVSLS CRYSYNLLAK EFRASLYKGV
61 NSDVEVCVGN GNFTYQPQFR SNAEFNCDGD FDNETVTFRL WNLHVNHTDI YFCKIEFMYP
121 PPYLDNERSN GTIIHIKEKH LCHTQSSPKL FWALVVVAGV LFCYGLLVTV ALCVIWTNSR
181 RNRLQLQSDYM NMTPRRPGLT RKPYPYAPA RDAAYRP [SEQ ID NO: 68]

10 In certain non-limiting embodiments, the CAR further comprises a spacer region that links the extracellular antigen-binding domain to the transmembrane domain. The spacer region can be flexible enough to allow the antigen binding domain to orient in different directions to facilitate antigen recognition while preserving the activating activity of the CAR.

15 In certain embodiments, the hinge/spacer region of the CAR comprises a native or modified hinge region of CD8 or a fragment thereof, a native or modified hinge region of CD28 or a fragment thereof, a native or modified hinge region of CD3 ζ or a fragment thereof, a native or modified hinge region of CD40 or a fragment thereof, a native or modified hinge region of 4-1BB or a fragment thereof, a native or modified hinge region of OX40 or a fragment thereof, a native or modified hinge region of CD84 or a fragment thereof, a native or modified hinge region of CD166 or a fragment thereof, a native or modified hinge region of CD8a or a fragment thereof, a native or modified hinge region of CD8b or a fragment thereof, a native or modified hinge region of ICOS or a fragment thereof, a native or modified hinge region of ICAM-1 or a fragment thereof, a native or modified hinge region of CTLA-4 or a fragment thereof, a native or modified hinge region of CD27 or a fragment thereof, a native or modified hinge region of CD40 or a fragment thereof, a native or modified hinge region of NKGD2 or a fragment thereof, a synthetic polypeptide (not based on a protein associated with the immune response), or a combination thereof. The hinge/spacer region can be the hinge region from IgG1, or the CH₂CH₃ region of immunoglobulin and portions of CD3, a portion of a CD28 polypeptide (e.g., a portion of SEQ ID NO: 66 or 68), a portion of a CD8 polypeptide (e.g., a portion of SEQ ID NO: 64 or 65), a variation of any of the foregoing which is at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 100% homologous or identical thereto, or a synthetic spacer sequence.

5.3.2.3. Intracellular Signaling Domain of a CAR

In certain embodiments, the CAR comprises an intracellular signaling domain. In certain non-limiting embodiments, the intracellular signaling domain of the CAR comprises a CD3 ζ polypeptide. CD3 ζ can activate or stimulate a cell (*e.g.*, a cell of the lymphoid lineage, *e.g.*, a T cell). Wild type (“native”) CD3 ζ comprises three functional immunoreceptor tyrosine-based activation motifs (ITAMs), three functional basic-rich stretch (BRS) regions (BRS1, BRS2 and BRS3). CD3 ζ transmits an activation signal to the cell (*e.g.*, a cell of the lymphoid lineage, *e.g.*, a T cell) after antigen is bound. The intracellular signaling domain of the CD3 ζ -chain is the primary transmitter of signals from endogenous TCRs.

In certain embodiments, the intracellular signaling domain of the CAR comprises a native CD3 ζ . In certain embodiments, the CD3 ζ polypeptide comprises or consists of an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100% homologous or identical to the amino acid sequence having a NCBI Reference No: NP_932170 (SEQ ID NO: 69) or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In certain non-limiting embodiments, the CD3 ζ polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 69, which is at least 20, or at least 30, or at least 40, or at least 50, and up to 164 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD3 ζ polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 164, 1 to 50, 50 to 100, 52 to 164, 100 to 150, or 150 to 164 of SEQ ID NO: 69. In certain embodiments, the intracellular signaling domain of the CAR comprises a CD3 ζ polypeptide comprising or consisting of amino acids 52 to 164 of SEQ ID NO: 69. SEQ ID NO: 69 is provided below:

```

1  MKWKALFTAA ILQAQLPITE AQSFGLLDPK LCYLLDGILF IYGVILTALF LRVKFSRSAD
61  APAYQQGQNQ LYNELNLGRR EEDVDLDRR GRDPEMGGKP QRRKNPQEGLYNELQKDKMA
121 EAYSEIGMKG ERRRGKGHDG LYQGLSTATK DTYDALHMQA LPPR [SEQ ID NO: 69]

```

In certain embodiments, the intracellular signaling domain of the CAR comprises a CD3 polypeptide comprising or consisting of the amino acid sequence set forth in SEQ ID NO: 89. SEQ ID NO: 89 is provided below.

```

RVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYS
EIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 89]

```

An exemplary nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 89 is set forth in SEQ ID NO: 90, which is as provided below.

AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGGTACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTC
 AATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAGATGGGGGAAAG
 CCGAGAAGGAAGAACCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCCTACAGT
 GAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCC
 5 ACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCTCGC [SEQ ID NO: 90]

In certain non-limiting embodiments, the intracellular signaling domain of the CAR further comprises at least a co-stimulatory signaling region. In certain embodiments, the co-stimulatory signaling region comprises at least one co-stimulatory molecule or a fragment thereof. In certain embodiments, the co-stimulatory signaling
 10 region comprises an intracellular domain of at least one co-stimulatory molecule or a fragment thereof.

As used herein, a “co-stimulatory molecule” refers to a cell surface molecule other than antigen receptor or its ligand that can provide an efficient response of lymphocytes to an antigen. In certain embodiments, a co-stimulatory molecule can provide optimal
 15 lymphocyte activation. Non-limiting examples of co-stimulatory molecules include CD28, 4-1BB, OX40, ICOS, DAP-10, CD27, CD40, NKGD2, CD2, FN14, HVEM, LTBR, CD28H, TNFR1, TNFR2, BAFF-R, BCMA, TACI, TROY, RANK, CD40, CD27, CD30, EDAR, XEDAR, GITR, DR6, and NGFR, and combinations thereof. The co-stimulatory molecule can bind to a co-stimulatory ligand, which is a protein expressed
 20 on cell surface that upon binding to its receptor produces a co-stimulatory response, *i.e.*, an intracellular response that effects the stimulation provided when an antigen-recognizing receptor (e.g., a chimeric antigen receptor (CAR)) binds to its target antigen. As one example, a 4-1BB ligand (*i.e.*, 4-1BBL) may bind to 4-1BB for providing an intracellular signal that in combination with a CAR signal induces an effector cell
 25 function of the CAR⁺ T cell.

In certain embodiments, the intracellular signaling domain of the CAR comprises a co-stimulatory signaling region that comprises a CD28 polypeptide, e.g., an intracellular domain of CD28 or a fragment thereof. The CD28 polypeptide can comprise or have an amino acid sequence that is at least about 80%, at least about 85%, at least about 90%, at
 30 least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99%, at least about 100% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 66 or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence
 35 that is a consecutive portion of SEQ ID NO: 66, which is at least 20, or at least 30, or at

least 40, or at least 50, and up to 220 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 220, 1 to 50, 50 to 100, 100 to 150, 114 to 220, 150 to 200, 180 to 220, or 200 to 220 of SEQ ID NO: 66. In certain embodiments, the intracellular signaling domain of the CAR comprises a co-stimulatory signaling region that comprises a CD28 polypeptide comprising or consisting of an amino acid sequence of amino acids 180 to 220 of SEQ ID NO: 66.

An exemplary nucleic acid sequence encoding amino acids 180 to 220 of SEQ ID NO: 66 is set forth in SEQ ID NO: 70, which is provided below.

10 AGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGCCACCCGC
AAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCCTATCGCTCC [SEQ ID NO: 70]

In certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99%, at least about 100% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 68 or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three conservative amino acid substitutions. In non-limiting certain embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 68 which is at least about 20, or at least about 30, or at least about 40, or at least about 50, and up to 218 amino acids in length. Alternatively or additionally, in non-limiting various embodiments, the CD28 polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 218, 1 to 50, 50 to 100, 100 to 150, 150 to 218, 178 to 218, or 200 to 218 of SEQ ID NO: 68. In certain embodiments, the co-stimulatory signaling region of a presently disclosed CAR comprises a CD28 polypeptide that comprises or consists of the amino acids 178 to 218 of SEQ ID NO: 68.

In certain embodiments, the intracellular signaling domain of the CAR comprises a co-stimulatory signaling region that comprises a 4-1BB polypeptide, e.g., an intracellular domain of 4-1BB or a fragment thereof. The 4-1BB polypeptide can comprise or consists of an amino acid sequence that is at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or at least about 99%, at least about 100% homologous or identical to the amino acid sequence having a NCBI Ref. No.: NP_001552 (SEQ ID NO: 71) or a fragment thereof, and/or may optionally comprise up to one or up to two or up to three

conservative amino acid substitutions. In non-limiting certain embodiments, the 4-1BB polypeptide comprises or consists of an amino acid sequence that is a consecutive portion of SEQ ID NO: 71, which is at least 20, or at least 30, or at least 40, or at least 50, or at least 100, or at least 150, or at least 150, and up to 255 amino acids in length.

- 5 Alternatively or additionally, in non-limiting various embodiments, the 4-1BB polypeptide comprises or consists of an amino acid sequence of amino acids 1 to 255, 1 to 50, 50 to 100, 100 to 150, 150 to 200, or 200 to 255 of SEQ ID NO: 71. In certain embodiments, the intracellular signaling domain of the CAR comprises a co-stimulatory signaling region that comprises a 4-1BB polypeptide comprising or consisting of an amino acid sequence of amino acids 214 to 255 of SEQ ID NO: 71. SEQ ID NO: 71 is provided below.

15
1 MGNSCYNIVA TLLLVLNFER TRSLQDPCSN CPAGTFCDNN RNQICSPCPP NSFSSAGGQR
61 TCDICRQCKG VERTRKECSS TSNAECDCTP GFHCLGAGCS MCEQDCKQGQ ELTKKGCKDC
121 CFGTFNDQKR GICRPWTNCS LDGKSVLVNG TKERDVVCGP SPADLSPGAS SVTPPAPARE
181 PGHSPQIISF FLALTSTALL FLLFFLTLRF SVVKRGRKKL LYIFKQPFMR PVQTTQEEDG
241 CSCRFPPEEEE GGCEL [SEQ ID NO: 71]

An exemplary nucleic acid sequence encoding amino acids 214 to 255 of SEQ ID NO: 71 is set forth in SEQ ID NO: 72, which is provided below.

20
AAACGGGGCAGAAAGAACTCCTGTATATATTCAAACAACCATTTATGAGACCAGTACAACTACT
CAAGAGGAAGATGGCTGTAGCTGCCGATTTCCAGAAGAAGAAGAGGAGGATGTGAACTG [SEQ ID NO: 72]

- 25 In certain embodiments, the intracellular signaling domain of the CAR comprises a co-stimulatory signaling region that comprises intracellular domains of two or more co-stimulatory molecules or portions thereof, e.g., an intracellular domain of CD28 or a fragment thereof and an intracellular domain of 4-1BB or a fragment thereof, or an intracellular domain of CD28 or a fragment thereof and an intracellular domain of OX40 or a fragment thereof.

5.3.2.4. Exemplified CARs

- 30 In certain embodiments, the CAR is a CD371-targeted CAR. In certain embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31, 35 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a

transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain of human CD28 or a fragment thereof), and (c) an intracellular signaling domain comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a fragment thereof).

- 5 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the V_H and V_L are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, the CAR is designed as “Et.B10L3H_MT_h28Z”. In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 73, which is provided below.

10 MLLLVTSLLLCELPHPAFLLIIPKVCNGIGIGEFKDSLSINATNIKHFKNCTSSISGDLHILPVAFR
GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLG
LRLSKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR
DCVSCRNVSRGECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPKIPSIATGMVGALLLLLVVALGIG
15 LFMGSGEGRGSLLTCGDVEENPGPMALPVTALLPLALLLHADIVMTQSPDSLAVSLGERATINCKSSQSVL
DSYNNENNLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYTSEPI
TFGQGTKEIKGGGSGGGGSGGGGSEVQLLESGLLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPGKGLE
WVSGIQGGGSGSTYYADSVKGRFTISRDNKNTLYLQMNSLRRAEDTAVYYCAREMWRGDYYSGMDVWGQGTTV
TVSSEQKLISEEDLAAAIEMVPPPYLDNEKSNGTIIHVKGKHLCPSPFPGPSKPFWVLVVGVLACYSL
20 LVTVAFIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQL
YNELNLGRREEYDVLDKRRGRDPGEGKPPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKGDGLYQG
LSTATKDTYDALHMQALPPR [SEQ ID NO: 73]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 73 is set forth in SEQ ID NO: 74, which is provided below.

25 ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCTCCTGATCCCA
CGCAAAGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAA
CACTTCAAAAAGTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTCTAGGGGTGACTCCTTC
ACACATACTCCTCCTCTGGACCCACAGGAAGTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTG
CTGATTCAAGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG
30 ACCAAGCAACATGGTCAGTTTTCTCTTGCAGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
AAGGAGATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
AAAAAAGTGTGGGACCTCCGGTCAGAAAACAAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCGAGGGCTGCTGGGGCCCGGAGCCAGGGACTGCGTCTCT
TGCCGGAATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
35 GTGGAGAACTCTGAGTGCATACAGTGCCACCCAGAGTGCCCTGCCTCAGGCCATGAACATCACCTGCACAGGA
CGGGGACCAGACAACTGTATCCAGTGTGCCCCTACATTGACGGCCCCCTGCGTCAAGACCTGCCCGGCA
GGAGTCATGGGAGAAAAACAACACCTGGTCTGGAGGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
CCAAACTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC

ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGACATCGTGATGACCCAGTCTCCAGACTCC
 CTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCTATAAC
 5 AATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCATCT
 ACCCGGGAATCCGGGGTCCCTGACCGATTCACTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGC
 AGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGAGCAATATACCAGCGAACCTATCACGTTTCGGCCAA
 GGTACCAAGGTGGAATCAAAGGCGGGGGTGGTAGTGGCGGTGGAGGTAGCGGAGGTGGCGGGTCTGAGGTG
 CAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGA
 10 TTCACCTTTAGCGACTATCAGATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGC
 ATTCAGGGTGGCGGTGGTAGCACATATTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAAT
 TCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGA
 GAGATGTGGCGTGGGGACTACTACTCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTCACCGTCTCCTCA
 GAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCTTACCTAGAC
 15 AATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGA
 CCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCCTGGCTTGCTATAGCTTGCTAGTAACAGTG
 GCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCC
 CGCCGCCCCGGGCCCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCCTATCGCTCC
 AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGCTACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTC
 20 AATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAGATGGGGGAAAG
 CCGAGAAGGAAGAACCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCTACAGT
 GAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCC
 ACCAAGGACACCTACGACGCCCTTACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO: 74]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 25 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
 30 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 of human CD28 or a portion thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 35 a 4-1BB polypeptide (e.g., an intracellular domain of human 4-1BB or a portion thereof).
 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 the amino acid sequence set forth in SEQ ID NO: 13. In certain embodiments, the V_H

and V_L are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, the CAR is designed as “Et.B10L4H_MT_hBBZ”. In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 75, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFKNCTSI SGDLHLIPVAFR
 5 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGFSLAVVSLNITSLG
 LRSLKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR
 DCVSCRNVSRGRECVDKCNLLEGEPPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
 TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGTGPGLEGCPNGPKIPSIATGMVGALLLLLVVALGIG
 LFMGSGEGRSLLTCGDVEENPGPMALPVTALLLPLALLLHADIVMTQSPDSLAVSLGERATINCKSSQSVL
 10 DSYNNENNLAWYQQKPGQPPKLLIYWASTRESGVDPDRFSGSGSGTDFTLTITSSSLQAEDVAVYYCQQYTSEPI
 TFGQGTKVEIKGGGGSGGGSGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPG
 KGLEWVSGIQGGGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSGMDVWGQ
 GTTVTVSSEQKLISEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSPFPGPSKPFWVLVVGGVLA
 CYSLLVTVAFIIFVWKGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAYQQ
 15 GQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHD
 GLYQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 75]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 75 is set forth in SEQ ID NO: 76, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
 20 CGCAAAGTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAA
 CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTGT
 CTGATTACAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGCAGTCGTCAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 25 AAGGAGATAAGTATGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 AAAAACTGTTTGGGACCTCCGGTCAGAAAACCAAAATTTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGGAGCCAGGGACTGCGTCTCT
 TGCCGGAATGTGAGCCGAGGACGGGAATGCGTGACAAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGCATACAGTGCCACCCAGAGTGCCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 30 CGGGGACCAGACAACTGTATCCAGTGTGCCACTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCA
 GGAGTCATGGGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 35 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGACATCGTGATGACCCAGTCTCCAGACTCC
 CTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCTATAAC
 AATGAGAACAAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCATCT
 ACCCGGGAATCCGGGGTCCCTGACCGATTAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGC
 AGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATACCAGCGAACCTATCACGTTCCGGCCAA

GGTACCAAGGTGGAAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGTCTGGCGGCGGCTCCGGTGGTGGT
 GGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGT
 GCAGCCTCTGGATTACCTTTAGCGACTATCAGATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAG
 TGGGTGTCAGGCATTGAGGGTGGCGGTGGTAGCACATATTACGCAGACTCCGTGAAGGGCCGGTTCACCATC
 5 TCCCGTGACAATTCCAAGAACACGCTGTATCTGCAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTAT
 TACTGTGCGAGAGAGATGTGGCGTGGGGACTACTACTCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTC
 ACCGTCTCCTCAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCT
 CCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCC
 CTATTTCCCGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCTGGCTTGCTATAGCTTG
 10 CTAGTAACAGTGGCCTTTATTATTTCTGGGTGAAACGGGGCAGAAAAGAACTCCTGTATATATTCAAACAA
 CCATTTATGAGACCAGTACAACTACTCAAGAGGAAGATGGCTGTAGCTGCCGATTTCCAGAAGAAGAAGAA
 GGAGGATGTGAACTGAGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGCTACCAGCAGGGCCAGAACCAG
 CTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCT
 GAGATGGGGGGAAAGCCGAGAAGGAAGAACCTCAGGAAGGCCGTGTACAATGAACTGCAGAAAGATAAGATG
 15 GCGGAGGCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAG
 GGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ
 ID NO: 76]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 20 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
 25 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 of human CD28 or a portion thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof).
 30 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 the amino acid sequence set forth in SEQ ID NO: 13. In certain embodiments, the V_H
 and V_L are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, the
 CAR is designed as “Et.B10L4H_MT_h28Z”. In certain embodiments, the CAR
 comprises the amino acid sequence set forth in SEQ ID NO: 77, which is provided below.

35 MLLLVTSLLLCELPHPAFLLIIPKVCNGIGIGEFKDSLSINATNIKHFKNCTSIISGDLHILPVAFR
 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLG
 LRSLKEISDGDVLIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR

DCVSCRNVSRGRECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
 TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPKIPSIATGMVGALLLLLVVALGIG
 LFMGSGEGRSLLTTCGDVEENPGPMALPVTALLLPLALLLHADIVMTQSPDSLAVSLGERATINCKSSQSVL
 DSYNNENNLAWYQQKPGQPPKLLIYWASTRESGVDPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYTSEPI
 5 TFGQGTKEIKGGGSGGGGSGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSWVRQAPG
 KGLEWVSGIQGGGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSGMDVWGQ
 GTTVTVSSEQKLI SEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSP LFPGPSKPFWVLVVGGLA
 CYSLLVTVAFIIFWVRSKRSRL LHS DYMNMTPRRPGPTRKHYPYAPPRDFAAYRSRVKFSRSADAPAYQQG
 QNQLYNELNLGRREEYDVL DKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKGH DG
 10 LYQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 77]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 77
 is set forth in SEQ ID NO: 78, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
 CGCAAAGTGTGTAAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAA
 15 CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACCTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTG
 CTGATT CAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGCACTCGTCAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 AAGGAGATAAGTGATGGAGATGTGATAATTT CAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 20 AAAAAACTGTTTGGGACCTCCGGTCAGAAAACAAAAATTTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGGAGCCCAGGGACTGCGTCTCT
 TGCCGGAATGT CAGCCGAGGCAGGGAATGCGTGACAAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGATACAGTGCCACCCAGAGTGCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 CGGGGACCAGACAACTGTATCCAGTGTGCCCACTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCCGCA
 25 GGAGTCATGGGAGAAAACAACACCCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGACATCGTGATGACCCAGTCTCCAGACTCC
 30 CTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCTATAAC
 AATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCATCT
 ACCCGGAATCCGGGGTCCCTGACCGATT CAGTGGCAGCGGTCTGGGACAGATTTCACTCTCACCATCAGC
 AGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGT CAGCAATATACCAGCGAACCTATCACGTT CGGCCAA
 GGTACCAAGGTGGAATCAAAGGTGGTGGTGGTTCAGGTGGTGGTGGTCTGGCGGCGGCTCCGGTGGTGGT
 35 GGATCCGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGT
 GCAGCCTCTGGATTACCTTTAGCGACTATCAGATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAG
 TGGGTGT CAGGCATT CAGGGTGGCGGTGGTAGACATATTACGCAGACTCCGTGAAGGGCCGGTT CACCATC
 TCCCGTGACAATTCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTAT
 TACTGTGCGAGAGAGATGTGGCGTGGGGACTACTACTCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTC
 40 ACCGTCTCCTCAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCT

CCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCC
 CTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCCCTGGCTTGCTATAGCTTG
 CTAGTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGA CTACATG
 AACATGACTCCCCGCCGCCCGGGCCCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCA
 5 GCCTATCGCTCCAGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGGTACCAGCAGGGCCAGAACCAGCTC
 TATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAG
 ATGGGGGGAAAGCCGAGAAGGAAGAACCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCG
 GAGGCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGT
 CTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID
 10 NO: 78]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
 15 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 20 of human CD28 or a portion thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof).
 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the V_H
 25 and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the
 CAR is designed as “Et.B10H3L_MT_h28Z”. In certain embodiments, the CAR
 comprises the amino acid sequence set forth in SEQ ID NO: 79, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPKVCNGIGIGEFKDSLSINATNIKHFNCTSSISGDLHILPVAFR
 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLG
 30 LRLSLKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR
 DCVSCRNVSRGRECVDKCNLLEGEPPREFVENSEICQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
 TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCP TNGPKIPSIATGMVGALLLLLVVALGIG
 LFMGSGEGRGSLLTCGDVEENPGPMALPVTALLPLALLLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFS
 DYQMSWVRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWR
 35 GDYYSGMDVWGQGT TTVTVSSGGGGSGGGSGGGSGDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNEN
 NLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTIS SLQAEDVAVYYCQQTSEPI TFGQGTK
 VEIKEQKLI SEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSP LFPGPSKPFWVLVVGGVLACYSL

LVTVAFIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQL
 YNELNLGRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKGDGLYQG
 LSTATKDTYDALHMQALPPR [SEQ ID NO: 79]

- An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 79
 5 is set forth in SEQ ID NO: 80, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
 CGCAAAGTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTTAA
 CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTTG
 10 CTGATTACAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGCAGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 AAGGAGATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 AAAAACTGTTTGGGACCTCCGGTCAGAAAACCAAAATTTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGAGGCCAGGGACTGCGTCTCT
 15 TGCCGGAATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGCATACAGTGCCACCCAGAGTGCCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 CGGGGACCAGACAACTGTATCCAGTGTGCCCCTACTACATTGACGGCCCCCTGCGTCAAGACCTGCCCCGCA
 GGAGTCATGGGAGAAAACAACACCCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 20 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGC
 TTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCGACTATCAGATG
 AGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGCATTACAGGTGGCGGTGGTAGCACA
 25 TATTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAATTCCAAGAACACGCTGTATCTGCAA
 ATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGACTACTAC
 TCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTCAACCTCTCCTCAGGCGGGGGTGGTAGTGGCGGTGGA
 GGTAGCGGAGGTGGCGGGTCTGACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAG
 CGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGCTATAACAATGAGAACAATTTAGCTTGG
 30 TATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGGGCATCTACCCGGGAATCCGGGGTCCCT
 GACCGATTACAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTG
 GCAGTTTATTACTGTGAGCAATATACCAGCGAACCTATCACGTTTCGGCCAAGGTACCAAGGTGGAAATCAAA
 GAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCTTACCTAGAC
 AATGAGAAGAGCAATGGAACCATTTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCATTTCCCGGA
 35 CCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCTGGCTTGCTATAGCTTGCTAGTAACAGTG
 GCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGAATGACATGAACATGACTCCC
 CGCCGCCCCGGGCCCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCGAGCCTATCGCTCC
 AGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGCTACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTC
 AATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAGATGGGGGAAAG
 40 CCGAGAAGGAAGAACCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCTACAGT

GAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCC
ACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO: 80]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
5 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
10 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
of human CD28 or a portion thereof), and (c) an intracellular signaling domain
comprising (i) a CD3ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof).

15 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
the amino acid sequence set forth in SEQ ID NO: 13. In certain embodiments, the V_H
and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the
CAR is designated as “Et.B10H4L_MT_h28Z”. In certain embodiments, the CAR
comprises the amino acid sequence set forth in SEQ ID NO: 81, which is provided below.

20 MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFKNCTSIGDLHILPVAFRGDSFTH
TPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHQQFSLAVVSLNITSLGLRSLKE
ISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPRDCVSCR
NVSRGRECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCPAGV
MGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPGPKIPSIATGMVGALLLLLVVALGIGLFMGSG
25 EGRGSLTTCGDVEENPGPMALPVTALLLPALLLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSW
VRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDNKNTLYLQMNSLR AEDTAVYYCAREMWRGDYYSG
MDVWGQGTITVTVSSGGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNENNL
AWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTITSSSLQAEDVAVYYCQQYTSEPITFGQGTKVE
IKEQKLI SEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSPLPFGPSKPFVVLVVGGVLACYSLLV
30 TVAFIIIFWVRSKRSLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQLYN
ELNLGRREEYDVLDRGRDPGEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKGHGGLYQGLS
TATKDTYDALHMQALPPR [SEQ ID NO: 81]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 81
is set forth in SEQ ID NO: 82, which is provided below.

35 ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCACGCAAA
GTGTGTAAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAACACTTC

In certain embodiments, the CAR is a CD371-targeted CAR. In certain embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35, and a

V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36, and (ii) a V_L that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39; (b) a

5 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain of human CD28 or a portion thereof), and (c) an intracellular signaling domain comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof).

In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of

10 the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the V_H and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the CAR is designated as "Et.C3H3L_MT_h28Z". In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 83, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFKNCTSSISGDLHILPVAFR

15 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLG

LRLSLKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR

DCVSCRNVSRGREGVDKCNLLEGEPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK

TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPKIPSIATGMVGALLLLLVVALGIG

LFMGSGEGRGSLLTCGDVEENPGPMALPVTALLPLALLLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFT

20 SYAMSWVRQAPGKGLEWVSGIDGSGGGTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARAYYD

ILTGYPVDGMDVWQGTTTVTVSSGGGSGGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLSSYN

NENNLAWYQQKPGQPPLKLLIYAASSTRESGVPDRFSGSGSGTDFTLTITSSQLAEDVAVYYCQQYYSEPYTFGQ

GTKVEIKEQKLISEEDLAAAEVMPYPYLDNEKSNGTIIHVKGKHLCPSPLPFGPSKPFWVLVVGGVLAC

YSLLVTVAFIIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRSRVKFSRSADAPAYQQGQ

25 NQLYNELNLGRREEYDVLDRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKHDGL

YQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 83]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 83 is set forth in SEQ ID NO: 84, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA

30 CGCAAAGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAA

CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC

ACACATACTCCTCCTCTGGACCCACAGGAACCTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTGTG

CTGATTACAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG

ACCAAGCAACATGGTCAGTTTTCTCTTGACAGTCGTCAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC

35 AAGGAGATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG

AAAAAACTGTTTGGGACCTCCGGTCAGAAAACAAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC

ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGAGCCCAGGGACTGCGTCTCT

TGCCGGAATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGCATACAGTGCCACCCAGAGTGCCCTGCCCTCAGGCCATGAACATCACCTGCACAGGA
 CGGGGACCAGACAACCTGTATCCAGTGTGCCCACTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCA
 GGAGTCATGGGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 5 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGC
 TTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACACCTTTACCAGCTATGCCATG
 10 AGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGCATTGACGGTAGCGGTGGTGGCACA
 AATTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAATTCCAAGAACACGCTGTATCTGCAA
 ATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGCGTATTACGATATTTTACTGGT
 TACCCCGTGACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCAGGCGGGGGTGGTAGT
 GGCGGTGGAGGTAGCGGAGGTGGCGGGTCTGACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCT
 15 CTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAAGCAGCTATAACAATGAGAACAAT
 TTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGCCGCATCTACCCGGGAATCC
 GGGGTCCCTGACCGATTTCAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCT
 GAAGATGTGGCAGTTTATTACTGTGAGCAATATTATAGCGAACCTTATACGTTTCGGCCAAGGTACCAAGGTG
 GAAATCAAAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCT
 20 TACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCTA
 TTTCCCGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCCCTGGCTTGCTATAGCTTGCTA
 GTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAAC
 ATGACTCCCCGCCGCCCGGGCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCC
 TATCGCTCCAGAGTGAAGTTTCAGCAGGAGCGCAGACGCCCCCGGTACCAGCAGGGCCAGAACCAGCTCTAT
 25 AACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAGATG
 GGGGGAAGCCGAGAAGGAAGAACCTCAGGAAGGCCGTACAATGAACTGCAGAAAGATAAGATGGCGGAG
 GCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTC
 AGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO:
 84]

30 In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 34, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36, and (ii) a V_L
 35 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37,
 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain

of human CD28 or a portion thereof), and (c) an intracellular signaling domain comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof). In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the V_H and V_L are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, the CAR is designated as "Et.C3L3H_MT_h28Z". In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 85, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFNCTSIISGDLHILPVAFR
 10 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGFSLAVVSLNITSLG
 LRSLEISDGDVLIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR
 DCVSCRNVSRGRECVDKCNLLEGEPPREFVENSEICIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
 TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPKIPSIATGMVGALLLLLVVALGIG
 LFMGSGEGRGSLLTTCGDVEENPGPMALPVTALLPLALLLHADIVMTQSPDSLAVSLGERATINCKSSQSVL
 15 SSYNNENNLAWYQQKPGQPPKLLIYAASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQYYSEPY
 TFGQGTKEIKGGGSGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFTSYAMSWVRQAPGKGLE
 WVSGIDGSGGGTNYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARAYYDILTGYPDGMDVWGQG
 TTVTVSSEQKLISEEDLAAAEVMPYPYLDNEKSNGTIIHVKGKHLCPSPFPGPSKPFVWL VVVGVLAC
 YSLLVTVAFIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRSRVKFSRSADAPAYQQGQ
 20 NQLYNELNLGRREEYDVLDRGRDPEMGGKPRRKNPQEGLYNELQDKMAEAYSEIGMKGERRRGKGHDGL
 YQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 85]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 85 is set forth in SEQ ID NO: 86, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
 25 CGCAAAGTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAA
 CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACCTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTTG
 CTGATTACAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGACAGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 30 AAGGAGATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 AAAAACTGTTTGGGACCTCCGGTCAGAAAACAAAAATTTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGAGGCCAGGGACTGCGTCTCT
 TGCCGGAATGTGAGCCGAGGCAGGGAATGCGTGACAAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGATACAGTGCCACCCAGAGTGCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 35 CGGGGACCAGACAACTGTATCCAGTGTGCCCCTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCCGCA
 GGAGTCATGGGAGAAAACAACACCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT

TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGACATCGTGATGACCCAGTCTCCAGACTCC
 CTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAAGCAGCTATAAC
 AATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGCCGCATCT
 5 ACCCGGAATCCGGGGTCCCTGACCGATTAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGC
 AGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATTATAGCGAACCTTATACGTTCCGGCCAA
 GGTACCAAGGTGGAATCAAAGGCGGGGGTGGTAGTGGCGGTGGAGGTAGCGGAGGTGGCGGGTCTGAGGTG
 CAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGA
 TTCACCTTTACCAGCTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGC
 10 ATTGACGGTAGCGGTGGTGGCACAATATTACGCAGACTCCGTGAAGGGCCGGTTACCATCTCCCGTGACAAT
 TCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGA
 GCGTATTACGATATTTTACTGGTTACCCCGTGACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACC
 GTCTCCTCAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCT
 TACCTAGACAATGAGAAGAGCAATGGAACCATTTATCCATGTGAAAGGGAACACCTTTGTCCAAGTCCCCTA
 15 TTTCCCGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCCTGGCTTGCTATAGCTTGCTA
 GTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAAC
 ATGACTCCCCGCCGCCCGGGCCACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCC
 TATCGCTCCAGAGTGAAGTTCAGCAGGAGCGCAGACGCCCCCGCTACCAGCAGGGCCAGAACCAGCTCTAT
 AACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCTGAGATG
 20 GGGGGAAGCCGAGAAGGAAGAACCCCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAG
 GCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTC
 AGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO:
 86]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 25 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 40, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43,
 30 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 of human CD28 or a portion thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 35 a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a portion thereof).
 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 the amino acid sequence set forth in SEQ ID NO: 14. In certain embodiments, the V_H

and V_L are positioned from the N- to the C-terminus: V_L-V_H. In certain embodiments, the CAQR is designated as “Et.D6L3H_MT_h28Z”. In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 87, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFNCTSI SGDLHLIPVAFR
 5 GDSFTHTPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRKQHGQFSLAVVSLNITSLG
 LRSLKEISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPR
 DCVSCRNVSRGRECVDKCNLLEGEPPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVK
 TCPAGVMGENNTLVWKYADAGHVCHLCHPNCTYGTGPGLEGCPNGPKIPSIATGMVGALLLLLVVALGIG
 LFMGSGEGRGSLLTGCDVEENPGPMALPVTALLLPLALLLHADIVMTQSPDSLAVSLGERATINCKSSQSVL
 10 RSSNNKNNLAWYQQKPGQPPKLLIYAASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYYREPL
 TFGQGTKVEIKGGGGSGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTFTDYAMSWVRQAPGKGLE
 WVSDIDGSGGSTDYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCALELGATTVYWGQGLTVTSSE
 QKLISEEDLAAAEVMPYPYLDNEKSNGTIIHVKGKHLCPSPLPFGPSKPFWVLVVGGVLACYSLLVTVVA
 FIIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAAYRSRVKFSRSADAPAYQQGQNQLYNELN
 15 LGRREEYDVLDRKRRDPEMGGKPRRKNPQEGLYNELQDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTAT
 KDTYDALHMQALPPR [SEQ ID NO: 87]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 87 is set forth in SEQ ID NO: 88, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
 20 CGCAAAGTGTGTAAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAA
 CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTG
 CTGATTACAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGACGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 25 AAGGAGATAAGTATGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 AAAAACTGTTTGGGACCTCCGGTCAAGAAACCAAAATTTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGGAGCCAGGGACTGCGTCTCT
 TGCCGGAATGTGAGCCGAGGACGGGAATGCGTGACAAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGCATACAGTGCCACCCAGAGTGCCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 30 CGGGGACCAGACAACTGTATCCAGTGTGCCACTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCA
 GGAGTCATGGGAGAAAACAACACCCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 35 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGACATCGTGATGACCCAGTCTCCAGACTCC
 CTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTACGCAGCAGCAAC
 AATAAAAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACGCCGCATCT
 ACCCGGGAATCCGGGGTCCCTGACCGATTGAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACCATCAGC
 AGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATTATCGCGAACCTCTGACGTTTCGGCCAA

GGTACCAAGGTGGAAATCAAAGGCGGGGGTGGTAGTGGCGGTGGAGGTAGCGGAGGTGGCGGGTCTGAGGTG
 CAGCTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGA
 TTCACCTTTACCGACTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGAC
 ATTGACGGTAGCGGTGGTAGCACAGACTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAAT
 5 TCCAAGAACACGCTGTATCTGCAAATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGCTA
 GAGCTGGGAGCTACTACCGTCTACTGGGGCCAGGGAACCCTGGTCACCGTCTCCTCAGAACAGAACTGATC
 TCTGAAGAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCTTACCTAGACAATGAGAAGAGCAAT
 GGAACCATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTT
 TGGGTGCTGGTGGTGGTTGGTGGAGTCTTGGCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATTATTTTC
 10 TGGGTGAGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGGCC
 ACCCGCAAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCCTATCGCTCCAGAGTGAAGTTCAGC
 AGGAGCGCAGACGCCCCCGGTACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGA
 GAGGAGTACGATGTTTTGGACAAGAGACGTGGCCGGGACCCTGAGATGGGGGGAAAGCCGAGAAGGAAGAAC
 CCTCAGGAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCTACAGTGAGATTGGGATGAAA
 15 GGCGAGCGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCCACCAAGGACACCTAC
 GACGCCCTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO: 88]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 20 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
 25 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 of human CD28 or a fragment thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a fragment thereof).
 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 30 the amino acid sequence set forth in SEQ ID NO: 93. In certain embodiments, the V_H
 and V_L are positioned from the N- to the C-terminus: V_H - V_L . In certain embodiments, the
 CAR is designed as “Et.B10H1L_MT_h28Z”. In certain embodiments, the CAR
 comprises the amino acid sequence set forth in SEQ ID NO: 95, which is provided below.
 MLLLVTSLLLCELPHPAFLLIIPKVCNGIGIGEFKDSLSINATNIKHFNCTSSISGDLHILPVAFRGDSFTH
 35 TPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLGLRSLKE
 ISDGDVLIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPRDCVSCR
 NVSRGRECVDKCNLLEGEPPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCPAGV

MGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPKIPSIATGMVGALLLLLVALGIGLFMGS
 EGRGSLTTCGDVEENPGPMALPVTALLLPALLLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSW
 VRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDNKNTLYLQMNLSRAEDTAVYYCAREMWRGDYYSG
 MDVWGQGTTVTVSSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNENNLAWEYQQKPGQPPKLL
 5 IYWASTRESGVDPDRFSGSGGTDFLTITISLQAEDVAVYYCQQTSEPIITFGQGTKEIKEQKLISEEDLAA
 AIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSPLEFPGPSKPFVWLTVVGGVLACYSLLVTVAFIIFWVRSKRS
 RLLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVL
 KRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKHDGLYQGLSTATKDTYDALHMQA
 LPPR [SEQ ID NO: 95]

10 An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 95
 is set forth in SEQ ID NO: 96, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACACACCCAGCATTCCTCCTGATCCCACGCAAA
 GTGTGTAAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAACACTTC
 AAAAAGTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTCACACAT
 15 ACTCCTCCTCTGGACCCACAGGAAGTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTGTCTGATT
 CAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGGACCAAG
 CAACATGGTCAGTTTTCTCTTGCAGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTCAAGGAG
 ATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGGAAAAA
 CTGTTTGGGACCTCCGGTCAGAAAACAAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCCACAGGC
 20 CAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGAGCCAGGGACTGCGTCTCTTGCCTGG
 AATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTTGTGGAG
 AACTCTGAGTGATACAGTGCCACCCAGAGTGCTGCCTCAGGCCATGAACATCACCTGCACAGGACGGGGA
 CCAGACAAGTGTATCCAGTGTGCCCCTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCAGGAGTC
 ATGGGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCATCCAAAC
 25 TGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCAAGATCCCGTCCATCGCC
 ACTGGGATGGTGGGGGCCCTCCTCTTGTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGTTCGGT
 GAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGTCCATGGCTCTCCAGTGACT
 GCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTA
 CAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCGACTATCAGATGAGCTGG
 30 GTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTGAGGCATTGAGGTGGCGGTGGTAGCACATATTAC
 GCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAATTCCAAGAACACGCTGTATCTGCAAATGAAC
 AGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGACTACTACTCCGGT
 ATGGACGTCTGGGGCCAGGGGACCACGGTCACCGTCTCCTCAGGCGGGGTGGTAGTGACATCGTGATGACC
 CAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTT
 35 TTAGACAGCTATAACAATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTC
 ATTTACTGGGCATCTACCCGGGAATCCGGGGTCCCTGACCGATTGAGTGGCAGCGGGTCTGGGACAGATTTT
 ACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGAGCAATATACCAGCGAACCT
 ATCACGTTCCGCCAAGGTACCAAGGTGAAATCAAAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCC
 GCAATTGAAGTTATGTATCCTCCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAA
 40 GGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGT

GGAGTCCTGGCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGC
 AGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGCCACCCGCAAGCATTACCAGCCC
 TATGCCCCACCACGCGACTTCGCAGCCTATCGCTCCAGAGTGAAAGTTCAGCAGGAGCGCAGACGCCCCCGCG
 TACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGAC
 5 AAGAGACGTGGCCGGGACCCTGAGATGGGGGAAAGCCGAGAAGGAAGAACCCCTCAGGAAGGCCTGTACAAT
 GAACTGCAGAAAGATAAGATGGCGGAGGCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAG
 GGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCC
 CTGCCCCCTCGCTAG [SEQ ID NO: 96]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 10 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
 V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
 that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
 15 a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
 transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
 of human CD28 or a fragment thereof), and (c) an intracellular signaling domain
 comprising (i) a CD3ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
 20 a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a fragment thereof).
 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
 the amino acid sequence set forth in SEQ ID NO: 94. In certain embodiments, the V_H
 and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the
 CAR is designed as “Et.B10H2L_MT_h28Z”. In certain embodiments, the CAR
 25 comprises the amino acid sequence set forth in SEQ ID NO: 97, which is provided below.
 MLLLVTSLLLCELPHPAFLLI PRKVCNGIGIGEFKDSL SINATNIKHFNCTSI SGDLHILPVAFRGDSFTH
 TPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGR TKQHGFSLAVVSLNITSLGLRSLKE
 ISDGDV IISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPRDCVSCR
 NVSRGRECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCPAGV
 30 MGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCP TNGPKIPSIATGMVGALLLLLVVALGIGLFMGSG
 EGRGSLLT CGDVEENPGPMALPVTALLLP LALLLHAEVQLLES GGGGLVQP GGSRLRLS CAASGFTFS DYQMSW
 VRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSG
 MDVWGQGT TVTVSSGGGGSGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYNNENNLA WYQQKPGQ
 PPKLLIYWASTRESGVPDRFSGSGSGTDFTLTITSSLQAEDVAVYYCQQYTSEPITFGQGT KVEIKEQKLISE
 35 EDLAAAEVMPYPYLDNEKSNGTIIHVKGKHLCPSP LFPGPSKPFVVLVVVGGVLACYSLLVTVAFIIFVW
 RSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAA YRSRVKFSRSADAPAYQQGQNQLYNELNLGRREE

YDVLDKRRGRDPPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDLGLYQGLSTATKDTYDA
LHMQUALPPR [SEQ ID NO: 97]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 97
is set forth in SEQ ID NO: 98, which is provided below.

5 ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACACACCCCAGCATTCTCCTGATCCCACGCAAA
GTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAACACTTC
AAAACTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTCACACAT
ACTCCTCCTCTGGACCCACAGGAAGTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTGTCTGATT
CAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGGACCAAG
10 CAACATGGTCAGTTTTCTCTTGCAGTCGTCAGCCTGAACATAACATCCTTGGGATTACGCTCCCTCAAGGAG
ATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGGAAAAAA
CTGTTTGGGACCTCCGGTCAGAAAACAAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCCACAGGC
CAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGGAGCCCAGGGACTGCGTCTCTTGGCCGG
AATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTTGTGGAG
15 AACTCTGAGTGACATACAGTGCCACCCAGAGTGCCTGCCTCAGGCCATGAACATCACCTGCACAGGACGGGGA
CCAGACAAGTGTATCCAGTGTGCCCCTACATTGACGGCCCCCTGCGTCAAGACCTGCCGGCAGGAGTCT
ATGGGAGAAAACAACACCCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCATCCAAAC
TGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCAAGATCCCGTCCATCGCC
ACTGGGATGGTGGGGGCCCTCCTCTTGTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGTTCGGGT
20 GAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCAGTGACT
GCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTA
CAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCGACTATCAGATGAGCTGG
GTCCGCCAGGCTCCAGGGAAGGGCTGGAGTGGGTGTGAGGCATTGAGGGTGGCGGTGGTAGCACATATTAC
GCAGACTCCGTGAAGGGCCGGTTACCATCTCCCGTGACAATTCCAAGAACACGCTGTATCTGCAAATGAAC
25 AGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGACTACTACTCCGGT
ATGGACGTCTGGGGCCAGGGGACCACGGTCACCGTCTCCTCAGGCGGGGTGGTAGTGGAGGTGGCGGGTCT
GACATCGTGATGACCCAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAG
TCCAGCCAGAGTGTTTTAGACAGCTATAACAATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAG
CCTCCTAAGCTGCTCATTTACTGGGCATCTACCCGGGAATCCGGGGTCCCTGACCGATTCACTGGCAGCGGG
30 TCTGGGACAGATTTCACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGAGCAA
TATACCAGCGAACCTATCACGTTTCGGCCAAGGTACCAAGGTGGAAATCAAAGAACAGAACTGATCTCTGAA
GAAGACCTGGCGGCCGCAATTGAAGTTATGTATCCTCCTCCTTACCTAGACAATGAGAAGAGCAATGGAACC
ATTATCCATGTGAAAGGGAAACACCTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTG
CTGGTGGTGGTGGTGGAGTCTGGCTTGTCTATAGCTTGTAGTAACAGTGGCCTTTATTATTTTCTGGGTG
35 AGGAGTAAGAGGAGCAGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGGCCACCCGC
AAGCATTACCAGCCCTATGCCCCACCACGCGACTTCGCAGCCTATCGCTCCAGAGTGAAGTTCAGCAGGAGC
GCAGACGCCCCCGGTACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAG
TACGATGTTTTGGACAAGAGACGTGGCCGGGACCTGAGATGGGGGAAAGCCGAGAAGGAAGAACCTCAG
GAAGGCCTGTACAATGAACTGCAGAAAGATAAGATGGCGGAGGCCCTACAGTGAGATTGGGATGAAAGGCGAG

CGCCGGAGGGGCAAGGGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCC
CTTCACATGCAGGCCCTGCCCCCTCGCTAG [SEQ ID NO: 98]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
5 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a
V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L
that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31,
a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L
10 CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a
transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain
of human CD28 or a fragment thereof), and (c) an intracellular signaling domain
comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising
a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a fragment thereof).
15 In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of
the amino acid sequence set forth in SEQ ID NO: 91. In certain embodiments, the V_H
and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the
CAR is designed as “Et.B10H5L_MT_h28Z”. In certain embodiments, the CAR
comprises the amino acid sequence set forth in SEQ ID NO: 99, which is provided below.

20 MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFKNCTISGDLHILPVAFRGDSFTH
TPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGQFSLAVVSLNITSLGLRSLKE
ISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPRDCVSCR
NVSRGRECVDKCNLLEGEPRFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCPAGV
MGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPGPKIPSIATGMVGALLLLLVALGIGLFMGSG
25 EGRGSLTTCGDVEENPGPMALPVTALLLPALLLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSW
VRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDNKNTLYLQMNSLR AEDTAVYYCAREMWRGDYYSG
MDVWGQGTITVTVSSGGGGSGGGSGGGSGGGSGGGSDIVMTQSPDSLAVSLGERATINCKSSQSVLDSYN
NENNLAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGTDFTLTITSSSLQAEDVAVYYCQQYTSEPITFGQ
GTKVEIKEQKLISEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSPFPGPSKPFVVLVVGGV LAC
30 YSLLVTVAFIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAYRSRVKFSRSADAPAYQQGQ
NQLYNELNLGRREEYDVLDKRRGRDP EMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRRGKGH DGL
YQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 99]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 99
is set forth in SEQ ID NO: 100, which is provided below.

35 ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCA
CGCAAAGTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAG

CACTTCAAAAACCTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTC
 ACACATACTCCTCCTCTGGACCCACAGGAACTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTTG
 CTGATTTCAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCCTAGAAATCATACGCGGCAGG
 ACCAAGCAACATGGTCAGTTTTCTCTTGTCAGTCGTGAGCCTGAACATAACATCCTTGGGATTACGCTCCCTC
 5 AAGGAGATAAGTGATGGAGATGTGATAATTTTCAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGG
 AAAAACTGTTTGGGACCTCCGGTCAGAAAACAAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCC
 ACAGGCCAGGTCTGCCATGCCTTGTGCTCCCCCGAGGGCTGCTGGGGCCCGGAGCCCAGGGACTGCGTCTCT
 TGCCGGAATGTCAGCCGAGGCAGGGAATGCGTGGACAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTT
 GTGGAGAACTCTGAGTGATACAGTGCCACCCAGAGTGCCCTGCCTCAGGCCATGAACATCACCTGCACAGGA
 10 CGGGGACCAGACAACCTGTATCCAGTGTCGCCACTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCGGCA
 GGAGTCATGGGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCAT
 CCAAACCTGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCC
 ATCGCCACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGT
 TCCGGTGAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGGTCCCATGGCTCTCCCA
 15 GTGACTGCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGC
 TTGGTACAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCGACTATCAGATG
 AGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTCAGGCATTCAGGGTGGCGGTGGTAGCACA
 TATTACGCAGACTCCGTGAAGGGCCGGTTCACCATCTCCCGTGACAATTCCAAGAACACGCTGTATCTGCAA
 ATGAACAGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGACTACTAC
 20 TCCGGTATGGACGTCTGGGGCCAGGGGACCACGGTCACCGTCTCCTCAGGTGGTGGTGGTTCAGGTGGTGGT
 GGTTCTGGAGGGGGCGGTTCTGGCGGCGGCTCCGGTGGTGGTGGATCCGACATCGTGATGACCCAGTCTCCA
 GACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTTTTAGACAGC
 TATAACAATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTCATTTACTGG
 GCATCTACCCGGGAATCCGGGGTCCCTGACCGATTTCAGTGGCAGCGGGTCTGGGACAGATTTCACTCTCACC
 25 ATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTGAGCAATATACCAGCGAACCTATCACGTTT
 GGCCAAGGTACCAAGGTGGAATCAAAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCCGCAATTGAA
 GTTATGTATCCTCCTCCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAAGGGAAACAC
 CTTTGTCCAAGTCCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGTGGAGTCTTG
 GCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGCAGGCTCCTG
 30 CACAGTGACTACATGAACATGACTCCCCGCCGCCCGGGCCACCCGCAAGCATTACCAGCCCTATGCCCCA
 CCACGCGACTTCGCAGCCTATCGCTCCAGAGTGAAGTTTCAGCAGGAGCGCAGACGCCCCCGCGTACCAGCAG
 GGCCAGAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGACAAGAGACGT
 GGCCGGGACCTGAGATGGGGGGAAAGCCGAGAAGGAAGAACCTCAGGAAGGCCTGTACAATGAACTGCAG
 AAAGATAAGATGGCGGAGGCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAGGGGCACGAT
 35 GGCCTTTACCAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCCCTGCCCCCT
 CGCTAG [SEQ ID NO: 100]

In certain embodiments, the CAR is a CD371-targeted CAR. In certain
 embodiments, the CAR comprises (a) an extracellular antigen-binding domain comprising
 (i) a V_H that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID
 40 NO: 28, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29, and a

V_H CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30, and (ii) a V_L that comprises a CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31, a CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32, and a V_L CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33; (b) a transmembrane domain comprising a CD28 polypeptide (e.g., a transmembrane domain of human CD28 or a fragment thereof), and (c) an intracellular signaling domain comprising (i) a CD3 ζ polypeptide, and (ii) a co-stimulatory signaling region comprising a CD28 polypeptide (e.g., an intracellular domain of human CD28 or a fragment thereof). In certain embodiments, the V_H and V_L are linked via a linker comprising or consisting of the amino acid sequence set forth in SEQ ID NO: 92. In certain embodiments, the V_H and V_L are positioned from the N- to the C-terminus: V_H-V_L. In certain embodiments, the CAR is designed as "Et.B10H6L_MT_h28Z". In certain embodiments, the CAR comprises the amino acid sequence set forth in SEQ ID NO: 101, which is provided below.

MLLLVTSLLLCELPHPAFLLIIPRKVCNGIGIGEFKDSLSINATNIKHFKNCTSI SGDLHILPVAFRGDSFTH
 TPPLDPQELDILKTVKEITGFLLIQAWPENRTDLHAFENLEIIRGRTKQHGGFSLAVVSLNITSLGLRSLKE
 ISDGDVIIISGNKNLCYANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEPRDCVSCR
 NVSRGRECVDKCNLLEGEPPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCAPGV
 MGENNTLVWKYADAGHVCHLCHPNCTYGCTGPGLEGCPNTPGPKIPSIATGMVGALLLLLVVALGIGLFMGSG
 EGRGSLLTCDGVEENPGPMALPVTALLLPALALLHAEVQLLESGGGLVQPGGSLRLSCAASGFTFSDYQMSW
 VRQAPGKGLEWVSGIQGGGGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREMWRGDYYSG
 MDVWGQGTITVTVSSGGGGSGGGGGSGGGGGSGGGGGSGGGGGSDIVMTQSPDSLAVSLGERATINCKSSQSV
 LDSYNNENNLAWYQQKPGQPPKLLIYWASTRESGPVDRFSGSGSGTDFTLTISSLQAEDVAVYYCQQYTSEP
 ITFGQGTKEIKEQKLI SEEDLAAAIEVMYPPPYLDNEKSNGTIIHVKGKHLCPSPFPGPSKPFVVLVVG
 GVLACYSLLVTVAFIIFWVRSKRSRLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRSRVKFSRSADAPA
 YQQGQNQLYNELNLGRREEYDVLDRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGK
 GHDGLYQGLSTATKDTYDALHMQALPPR [SEQ ID NO: 101]

An exemplary nucleic acid sequence the amino acid sequence of SEQ ID NO: 101 is set forth in SEQ ID NO: 102, which is provided below.

ATGCTTCTCCTGGTGACAAGCCTTCTGCTCTGTGAGTTACCACACCCAGCATTCCTCCTGATCCCACGCAAA
 GTGTGTAACGGAATAGGTATTGGTGAATTTAAAGACTCACTCTCCATAAATGCTACGAATATTAAACACTTC
 AAAAAGTGCACCTCCATCAGTGGCGATCTCCACATCCTGCCGGTGGCATTTAGGGGTGACTCCTTCACACAT
 ACTCCTCCTCTGGACCCACAGGAAGTGGATATTCTGAAAACCGTAAAGGAAATCACAGGGTTTTTGTCTGATT
 CAGGCTTGGCCTGAAAACAGGACGGACCTCCATGCCTTTGAGAACCTAGAAATCATACGCGGCAGGACCAAG
 CAACATGGTCAGTTTTCTCTTGCAGTCGTCAGCCTGAACATAACATCCTTGGGATTACGCTCCCTCAAGGAG
 ATAAGTGATGGAGATGTGATAATTTAGGAAACAAAAATTTGTGCTATGCAAATACAATAAACTGGAAAAAA
 CTGTTTGGGACCTCCGGTCAGAAAACCAAAATTATAAGCAACAGAGGTGAAAACAGCTGCAAGGCCACAGGC

CAGGTCTGCCATGCCTTGTGCTCCCCGAGGGCTGCTGGGGCCCGGAGCCCAGGGACTGCGTCTCTTGCCGG
 AATGTGAGCCGAGGCAGGGAATGCGTGACAAAGTGCAACCTTCTGGAGGGTGAGCCAAGGGAGTTTGTGGAG
 AACTCTGAGTGATACAGTGCCACCCAGAGTGCCCTGCCTCAGGCCATGAACATCACCTGCACAGGACGGGGA
 CCAGACAACTGTATCCAGTGCTGCCCCTACATTGACGGCCCCCACTGCGTCAAGACCTGCCCCGAGGAGTC
 5 ATGGGAGAAAACAACACCCTGGTCTGGAAGTACGCAGACGCCGGCCATGTGTGCCACCTGTGCCATCCAAAC
 TGCACCTACGGATGCACTGGGCCAGGTCTTGAAGGCTGTCCAACGAATGGGCCTAAGATCCCGTCCATCGCC
 ACTGGGATGGTGGGGGCCCTCCTCTTGCTGCTGGTGGTGGCCCTGGGGATCGGCCTCTTCATGGGTTCCGGT
 GAGGGACGGGGGTCACTGCTCACCTGCGGAGATGTAGAAGAGAATCCCGTCCCATGGCTCTCCAGTGACT
 GCCCTACTGCTTCCCCTAGCGCTTCTCCTGCATGCAGAGGTGCAGCTGTTGGAGTCTGGGGGAGGCTTGGTA
 10 CAGCCTGGGGGGTCCCTGCGACTCTCCTGTGCAGCCTCTGGATTACCTTTAGCGACTATCAGATGAGCTGG
 GTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTGTGAGGCATTGAGGTGGCGGTGGTAGCACATATTAC
 GCAGACTCCGTGAAGGGCCGGTTCACCATCTCCGTGACAATTCGAAGAACACGCTGTATCTGCAAATGAAC
 AGCCTGCGTGCCGAGGACACGGCTGTGTATTACTGTGCGAGAGAGATGTGGCGTGGGGACTACTACTCCGGT
 ATGGACGTCTGGGGCCAGGGGACCACGGTCAACCTCTCCTCAGGTGGTGGTGGTTCAGGTGGTGGTGGTCT
 15 GGTGGGGGCGGTAGTGAGGGGGCGGTTCTGGCGGCGGCTCCGGTGGTGGTGGATCCGACATCGTGATGACC
 CAGTCTCCAGACTCCCTGGCTGTGTCTCTGGGCGAGCGTGCCACCATCAACTGCAAGTCCAGCCAGAGTGTT
 TTAGACAGCTATAACAATGAGAACAATTTAGCTTGGTATCAGCAGAAACCAGGACAGCCTCCTAAGCTGCTC
 ATTTACTGGGCATCTACCCGGGAATCCGGGGTCCCTGACCGATTGAGTGGCAGCGGGTCTGGGACAGATTTT
 ACTCTCACCATCAGCAGCCTGCAGGCTGAAGATGTGGCAGTTTATTACTGTCAGCAATATACCAGCGAACCT
 20 ATCACGTTTCGGCCAAGGTACCAAGGTGGAATCAAAGAACAGAACTGATCTCTGAAGAAGACCTGGCGGCC
 GCAATTGAAGTTATGTATCCTCCTTACCTAGACAATGAGAAGAGCAATGGAACCATTATCCATGTGAAA
 GGGAAACACCTTTGTCCAAGTCCCTATTTCCCGGACCTTCTAAGCCCTTTTGGGTGCTGGTGGTGGTGGT
 GGAGTCTGGCTTGCTATAGCTTGCTAGTAACAGTGGCCTTTATTATTTTCTGGGTGAGGAGTAAGAGGAGC
 AGGCTCCTGCACAGTGACTACATGAACATGACTCCCCGCGCCCCGGGGCCACCCGCAAGCATTACCAGCCC
 25 TATGCCCCACCACGCGACTTCGCAGCCTATCGCTCCAGAGTGAAGTTGAGCAGGAGCGCAGACGCCCCCGCG
 TACCAGCAGGGCCAGAACCAGCTCTATAACGAGCTCAATCTAGGACGAAGAGAGGAGTACGATGTTTTGGAC
 AAGAGACGTGGCCGGGACCTGAGATGGGGGGAAAGCCGAGAAGGAAGAACCTCAGGAAGGCCTGTACAAT
 GAACTGCAGAAAGATAAGATGGCGGAGGCCTACAGTGAGATTGGGATGAAAGGCGAGCGCCGGAGGGGCAAG
 GGGCACGATGGCCTTTACCAGGGTCTCAGTACAGCCACCAAGGACACCTACGACGCCCTTCACATGCAGGCC
 30 CTGCCCCCTCGCTAG [SEQ ID NO: 102]

In certain embodiments, a presently disclosed CAR further comprises an inducible promoter, for expressing nucleic acid sequences in human cells. Promoters for use in expressing CAR genes can be a constitutive promoter, such as ubiquitin C (UbiC) promoter.

35 5.3.3. TCR like Fusion Molecules

In certain embodiments, the antigen-recognizing receptor is a TCR like fusion molecule. Non-limiting examples of TCR fusion molecules include HLA-Independent TCR-based Chimeric Antigen Receptor (also known as “HIT-CAR”, e.g., those disclosed in International Patent Application No. PCT/US19/017525, which is incorporated by

reference in its entirety), and T cell receptor fusion constructs (TRuCs) (e.g., those disclosed in Baeuerle et al., “Synthetic TRuC receptors engaging the complete T cell receptor for potent anti-tumor response,” *Nature Communications* volume 10, Article number: 2087 (2019), which is incorporated by reference in its entirety).

5 In certain embodiments, the TCR like fusion molecule comprises an antigen binding chain that comprises an extracellular antigen-binding domain and a constant domain, wherein the TCR like fusion molecule binds to an antigen in an HLA-independent manner. In certain embodiments, the constant domain comprises a T cell receptor constant region selected from the group consisting of a native or modified TRAC peptide, 10 a native or modified TRBC peptide, a native or modified TRDC peptide, a native or modified TRGC peptide and any variants or functional fragments thereof. In certain embodiments, the constant domain comprises a native or modified TRAC peptide. In certain embodiments, the constant domain comprises a native or modified TRBC peptide. In certain embodiments, the constant domain is capable of forming a homodimer or a 15 heterodimer with another constant domain. In certain embodiments, the antigen binding chain is capable of associating with a CD3 ζ polypeptide. In certain embodiments, the antigen binding chain, upon binding to an antigen, is capable of activating the CD3 ζ polypeptide associated to the antigen binding chain. In certain embodiments, the activation of the CD3 ζ polypeptide is capable of activating an immunoresponsive cell. In 20 certain embodiments, the TCR like fusion molecule is capable of integrating with a CD3 complex and providing HLA-independent antigen recognition. In certain embodiments, the TCR like fusion molecule replaces an endogenous TCR in a CD3/TCR complex. In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule is capable of dimerizing with another extracellular antigen-binding domain. In 25 certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises a ligand for a cell-surface receptor, a receptor for a cell surface ligand, an antigen binding portion of an antibody or a fragment thereof or an antigen binding portion of a TCR. In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises one or two immunoglobulin variable 30 region(s). In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises a heavy chain variable region (V_H) of an antibody. In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises a light chain variable region (V_L) of an antibody. In certain

embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule is capable of dimerizing with another extracellular antigen-binding domain. In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises a V_H of an antibody, wherein the V_H is capable of dimerizing with another
 5 extracellular antigen-binding domain comprising a V_L of the antibody and form a fragment variable (Fv). In certain embodiments, the extracellular antigen-binding domain of the TCR like fusion molecule comprises a V_L of an antibody, wherein the V_L is capable of dimerizing with another extracellular antigen-binding domain comprising a V_H of the antibody and form a fragment variable (Fv).

10 The TCR like fusion molecule can bind to a tumor antigen or a pathogen antigen. In certain embodiments, the TCR like fusion molecule binds to a tumor antigen.

5.4. Cells

The presently disclosed subject matter provides cells comprising a presently disclosed CD371-targeted antigen-recognizing receptor (e.g., one disclosed in Section
 15 5.3). In certain embodiments, the cell is selected from the group consisting of cells of lymphoid lineage and cells of myeloid lineage. In certain embodiments, the cell is an immunoresponsive cell. In certain embodiments, the immunoresponsive cell is a cell of lymphoid lineage.

In certain embodiments, the cell is a cell of the lymphoid lineage. Cells of the
 20 lymphoid lineage can provide production of antibodies, regulation of cellular immune system, detection of foreign agents in the blood, detection of cells foreign to the host, and the like. Non-limiting examples of cells of the lymphoid lineage include T cells, Natural Killer (NK) cells, B cells, dendritic cells, stem cells from which lymphoid cells may be differentiated. In certain embodiments, the stem cell is a pluripotent stem cell (e.g.,
 25 embryonic stem cell).

In certain embodiments, the cell is a T cell. T cells can be lymphocytes that mature in the thymus and are chiefly responsible for cell-mediated immunity. T cells are involved in the adaptive immune system. The T cells of the presently disclosed subject matter can be any type of T cells, including, but not limited to, helper T cells, cytotoxic T
 30 cells, memory T cells (including central memory T cells, stem-cell-like memory T cells (or stem-like memory T cells), and two types of effector memory T cells: e.g., TEM cells and TEMRA cells, Regulatory T cells (also known as suppressor T cells), tumor-infiltrating lymphocyte (TIL), Natural killer T cells, Mucosal associated invariant T cells,

and $\gamma\delta$ T cells. Cytotoxic T cells (CTL or killer T cells) are a subset of T lymphocytes capable of inducing the death of infected somatic or tumor cells. A patient's own T cells may be genetically modified to target specific antigens through the introduction of an antigen-recognizing receptor, *e.g.*, a CAR or a TCR. In certain embodiments, the

5 immunoresponsive cell is a T cell. The T cell can be a $CD4^+$ T cell or a $CD8^+$ T cell. In certain embodiments, the T cell is a $CD4^+$ T cell. In certain embodiments, the T cell is a $CD8^+$ T cell.

In certain embodiments, the cell is a NK cell. Natural killer (NK) cells can be lymphocytes that are part of cell-mediated immunity and act during the innate immune

10 response. NK cells do not require prior activation in order to perform their cytotoxic effect on target cells.

Types of human lymphocytes of the presently disclosed subject matter include, without limitation, peripheral donor lymphocytes. *e.g.*, those disclosed in Sadelain et al., *Nat Rev Cancer* (2003); 3:35-45 (disclosing peripheral donor lymphocytes genetically

15 modified to express CARs), in Morgan, R.A., *et al.* 2006 *Science* 314:126-129 (disclosing peripheral donor lymphocytes genetically modified to express a full-length tumor antigen-recognizing T cell receptor complex comprising the α and β heterodimer), in Panelli et al., *J Immunol* (2000);164:495-504; Panelli et al., *J Immunol* (2000);164:4382-4392 (disclosing lymphocyte cultures derived from tumor infiltrating lymphocytes (TILs) in

20 tumor biopsies), and in Dupont et al., *Cancer Res* (2005);65:5417-5427; Papanicolaou et al., *Blood* (2003);102:2498-2505 (disclosing selectively *in vitro*-expanded antigen-specific peripheral blood leukocytes employing artificial antigen-presenting cells (AAPCs) or pulsed dendritic cells).

The cells (*e.g.*, T cells) can be autologous, non-autologous (*e.g.*, allogeneic), or

25 derived *in vitro* from engineered progenitor or stem cells.

The cells of the presently disclosed subject matter can be cells of the myeloid lineage. Non-limiting examples of cells of the myeloid lineage include monocytes, macrophages, neutrophils, dendritic cells, basophils, neutrophils, eosinophils, megakaryocytes, mast cell, erythrocyte, thrombocytes, and stem cells from which

30 myeloid cells may be differentiated. In certain embodiments, the stem cell is a pluripotent stem cell (*e.g.*, an embryonic stem cell or an induced pluripotent stem cell).

In certain embodiments, the presently disclosed cells are capable of modulating the tumor microenvironment. Tumors have a microenvironment that is hostile to the host

immune response involving a series of mechanisms by malignant cells to protect themselves from immune recognition and elimination. This “hostile tumor microenvironment” comprises a variety of immune suppressive factors including infiltrating regulatory CD4⁺ T cells (Tregs), myeloid derived suppressor cells (MDSCs), tumor associated macrophages (TAMs), immune suppressive cytokines including TGF-β, and expression of ligands targeted to immune suppressive receptors expressed by activated T cells (CTLA-4 and PD-1). These mechanisms of immune suppression play a role in the maintenance of tolerance and suppressing inappropriate immune responses, however within the tumor microenvironment these mechanisms prevent an effective anti-tumor immune response. Collectively these immune suppressive factors can induce either marked anergy or apoptosis of adoptively transferred CAR modified T cells upon encounter with targeted tumor cells.

In certain embodiments, the cells can be transduced with the presently disclosed CD371-targeted antigen-recognizing receptor such that the cells express the antigen-recognizing receptor.

In certain embodiments, the cell further comprises a soluble single-chain variable fragment (scFv) that binds a polypeptide that has immunosuppressive activity or immunostimulatory activity. In certain embodiments, immunosuppressive activity refers to induction of signal transduction or changes in protein expression in a cell (e.g., an activated immunoresponsive cell) resulting in a decrease in an immune response. Polypeptides known to suppress or decrease an immune response via their binding include CD47, PD-1, CTLA-4, and their corresponding ligands, including SIRPα, PD-L1, PD-L2, B7-1, and B7-2. Such polypeptides are present in the tumor microenvironment and inhibit immune responses to neoplastic cells. In various embodiments, inhibiting, blocking, or antagonizing the interaction of immunosuppressive polypeptides and/or their ligands enhances the immune response of the immunoresponsive cell.

In certain embodiments, immunostimulatory activity refers to induction of signal transduction or changes in protein expression in a cell (e.g., an activated immunoresponsive cell) resulting in an increase in an immune response.

Immunostimulatory activity may include pro-inflammatory activity. Polypeptides known to stimulate or increase an immune response via their binding include CD28, OX-40, 4-1BB, and their corresponding ligands, including B7-1, B7-2, OX-40L, and 4-1BBL. Such polypeptides are present in the tumor microenvironment and activate immune responses

to neoplastic cells. In various embodiments, promoting, stimulating, or agonizing pro-inflammatory polypeptides and/or their ligands enhances the immune response of the immunoresponsive cell.

Cells comprising an antigen-recognizing receptor (e.g., a CAR) and a soluble scFv that binds a polypeptide that has immunosuppressive activity or immunostimulatory activity are disclosed in International Patent Publication No. WO 2014/134165, which is incorporated by reference in its entirety.

In certain embodiments, the cell further comprises an exogenous CD40L. Cells comprising an antigen-recognizing receptor (e.g., a CAR) and an exogenous CD40L are disclosed in International Patent Publication No. WO 2014/134165.

Furthermore, in certain embodiments, the cell is engineered to express IL-18. In certain embodiments, the cell further comprises an exogenous IL-18 polypeptide. In certain embodiments, the exogenous IL-18 polypeptide comprises or consists of the amino acid sequence set forth in SEQ ID NO: 103, which is provided below.

MGYRMQLLSICIALSLALVTNSGYFGKLESKLSVIRNLNDQVLFIDQGNRPLFEDMTDSDCRDNAP
RTIFIISMYKDSQPRGMAVTISVKCEKISTLSCENKIISFKEMNPPDNIKDTKSDIIFQRSVPG
HDNKMQFESSYEGYFLACEKERDLFKLILKKEDELGDRSIMFTVQNE [SEQ ID NO: 103]

In certain embodiments, the cells further comprise a nucleic acid molecule encoding an IL-18 polypeptide. In certain embodiments, the nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 104, which is provided below.

ATGGGTACAGGATGCAACTCCTGTCTTGCATTGCACTAAGTCTTGCCTTGTACAAACAGTGG
CTACTTTGGCAAGCTTGAATCTAAATTATCAGTCATAAGAAATTTGAATGACCAAGTTCTCTTCA
TTGACCAAGGAAATCGGCCTCTATTTGAAGATATGACTGATTCTGACTGTAGAGATAATGCACCC
CGGACCATATTTATTATAAGTATGTATAAAGATAGCCAGCCTAGAGGTATGGCTGTAACATCTC
TGTGAAGTGTGAGAAAATTTCAACTCTCTCCTGTGAGAACAAAATTATTTCTTTAAGGAAATGA
ATCCTCCTGATAACATCAAGGATACAAAAAGTGACATCATATTCTTTCAGAGAAGTGTCCCAGGA
CATGATAATAAGATGCAATTTGAATCTTCATCATACGAAGGATACTTTCTAGCTTGTGAAAAAGA
GAGAGACCTTTTTAACTCATTTTGAAAAAGAGGATGAATTGGGGGATAGATCTATAATGTTCA
CTGTTCAAACGAAGACTAG [SEQ ID NO: 104]

Alternatively, in certain embodiments, the cell further comprises a modified promoter/enhancer at an IL-18 gene locus, which can increase IL-18 gene expression, e.g., a constitutive or inducible promoter is placed to drive IL-18 gene expression.

Cells comprising an antigen-recognizing receptor (e.g., a CAR) and engineered to express IL-18, e.g., comprising an exogenous IL-18 polypeptide or a modified promoter/enhancer at an IL-18 gene locus are disclosed in International Patent Publication No. WO2018/027155, which is incorporated by reference in its entirety.

5 Additionally or alternatively, the cell is engineered to express IL-33. In certain embodiments, the cell further comprises an exogenous IL-33 polypeptide. In certain embodiments, the exogenous IL-33 polypeptide comprises or consists of the amino acid sequence set forth in SEQ ID NO: 105, which is provided below.

10 MYRMQLLSICIALSLALVTNSSITGISPITEYLASLSTYNDQSITFALEDESYEIYVEDLKKDEKK
DKVLLSYYESQHPSPNESGDGVDGKMLMVTLSPTKDFWLHANNKEHSVELHKCEKPLPDQAFFVLH
NMHSNCVSFECKTDPGVFIGVKDNHLALIKVDSSENLCTENILFKLSET [SEQ ID NO: 105]

15 In certain embodiments, the cells further comprise a nucleic acid molecule encoding an IL-33 polypeptide. In certain embodiments, the nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 106, which is provided below.

20 ATGTACAGGATGCAACTCCTGTCTTGCATTGCACTAAGTCTTGCCTTGTACAAACAGTAGTAT
CACAGGAATTTACCTATTACAGAGTATCTTGCTTCTCTAAGCACATACAATGATCAATCCATTA
CTTTTGCTTTGGAGGATGAAAGTTATGAGATATATGTTGAAGACTTGAAAAAGATGAAAAGAA
GATAAGGTGTTACTGAGTTACTATGAGTCTCAACACCCCTCAAATGAATCAGGTGACGGTGTGTA
TGGTAAGATGTTAATGGTAACCCCTGAGTCTACAAAAGACTTCTGGTTGCATGCCAACACAAGG
AACACTCTGTGGAGCTCCATAAGTGTGAAAAACCACTGCCAGACCAGGCCTTCTTTGTCCTTCAT
AATATGCACTCCAACGTGTTTCATTTGAATGCAAGACTGATCCTGGAGTGTATATAGGTGTAA
GGATAATCATCTTGCTCTGATTAAAGTAGACTCTTCTGAGAATTTGTGTACTGAAAATATCTTGT
TTAAGCTCTCTGAAACTTAG [SEQ ID NO: 106]

25 Alternatively, in certain embodiments, the cell further comprises a modified promoter/enhancer at an IL-33 gene locus, which can increase IL-33 gene expression, e.g., a constitutive or inducible promoter placed to drive IL-33 gene expression. Cells comprising an antigen-recognizing receptor (e.g., a CAR) and engineered to express IL-33, e.g., comprising an exogenous IL-33 polypeptide or a modified promoter/enhancer at
30 an IL-33 gene locus are disclosed in International Patent Publication No. WO2019/099479, which is incorporated by reference in its entirety.

Additionally or alternatively, the cell is engineered to express IL-36. In certain embodiments, the cell further comprises an exogenous IL-36 polypeptide. In certain embodiments, the cell further comprises a modified promoter/enhancer at an IL-36 gene

locus, which can increase IL-36 gene expression, e.g., a constitutive or inducible promoter placed to drive IL-36 gene expression. Cells comprising an antigen-recognizing receptor (e.g., a CAR) and engineered to express IL-36, e.g., comprising an exogenous IL-36 polypeptide or a modified promoter/enhancer at an IL-36 gene locus are disclosed in International Patent Publication No. WO2019/099483, which is incorporated by reference in its entirety.

5.5. *Compositions and Vectors*

The presently disclosed subject matter provides compositions comprising a presently disclosed CD371-targeted antigen-recognizing receptor (e.g., one disclosed in Section 5.3). Also provided are cells comprising such compositions.

In certain embodiments, the presently disclosed CD371-targeted antigen-recognizing receptor is operably linked to a promoter.

Furthermore, the present discloses subject matter provides nucleic acid compositions comprising a polynucleotide encoding a presently disclosed CD371-targeted antigen-recognizing receptor (e.g., one disclosed in Section 5.3). Also provided are cells comprising such nucleic acid compositions.

In certain embodiments, the nucleic acid composition further comprises a promoter that is operably linked to the presently disclosed CD371-targeted antigen-recognizing receptor.

In certain embodiments, the promoter is endogenous or exogenous. In certain embodiments, the exogenous promoter is selected from an elongation factor (EF)-1 promoter, a cytomegalovirus immediate-early promoter (CMV) promoter, a simian virus 40 early promoter (SV40) promoter, a phosphoglycerate kinase (PGK) promoter, and a metallothionein promoter. In certain embodiments, the promoter is an inducible promoter. In certain embodiment, the inducible promoter is selected from a NFAT transcriptional response element (TRE) promoter, a CD69 promoter, a CD25 promoter, and an IL-2 promoter.

The compositions and nucleic acid compositions can be administered to subjects or and/delivered into cells by art-known methods or as described herein. Genetic modification of a cell (e.g., a T cell or a NK cell) can be accomplished by transducing a substantially homogeneous cell composition with a recombinant DNA construct. In certain embodiments, a retroviral vector (e.g., gamma-retroviral vector or lentiviral vector) is employed for the introduction of the DNA construct into the cell. For example,

a polynucleotide encoding an antigen-recognizing receptor can be cloned into a retroviral vector and expression can be driven from its endogenous promoter, from the retroviral long terminal repeat, or from a promoter specific for a target cell type of interest. Non-viral vectors may be used as well.

5 For initial genetic modification of a cell to include a presently disclosed CD371-targeted antigen-recognizing receptor (*e.g.*, a CAR or a TCR), a retroviral vector can be employed for transduction, however any other suitable viral vector or non-viral delivery system can be used. The antigen-recognizing receptor can be constructed in a single, multicistronic expression cassette, in multiple expression cassettes of a single vector, or in
10 multiple vectors. Examples of elements that create polycistronic expression cassette include, but is not limited to, various viral and non-viral Internal Ribosome Entry Sites (IRES, *e.g.*, FGF-1 IRES, FGF-2 IRES, VEGF IRES, IGF-II IRES, NF- κ B IRES, RUNX1 IRES, p53 IRES, hepatitis A IRES, hepatitis C IRES, pestivirus IRES, aphthovirus IRES, picornavirus IRES, poliovirus IRES and encephalomyocarditis virus
15 IRES) and cleavable linkers (*e.g.*, 2A peptides, *e.g.*, P2A, T2A, E2A and F2A peptides). Combinations of retroviral vector and an appropriate packaging line are also suitable, where the capsid proteins will be functional for infecting human cells. Various amphotropic virus-producing cell lines are known, including, but not limited to, PA12 (Miller *et al.*, (1985) *Mol Cell Biol* (1985);5:431-437); PA317 (Miller, *et al.*, *Mol Cell*
20 *Biol* (1986); 6:2895-2902); and CRIP (Danos *et al.*, *Proc Natl Acad Sci USA* (1988);85:6460-6464). Non-amphotropic particles are suitable too, *e.g.*, particles pseudotyped with VSVG, RD114 or GALV envelope and any other known in the art.

Possible methods of transduction also include direct co-culture of the cells with producer cells (Bregni *et al.*, *Blood* (1992);80:1418-1422), or culturing with viral
25 supernatant alone or concentrated vector stocks with or without appropriate growth factors and polycations (Xu *et al.*, *Exp Hemat* (1994); 22:223-230; and Hughes *et al.* *J Clin Invest* (1992); 89:1817).

Other transducing viral vectors can be used to modify a cell. In certain embodiments, the chosen vector exhibits high efficiency of infection and stable
30 integration and expression (*see, e.g.*, Cayouette *et al.*, *Human Gene Therapy* 8:423-430, 1997; Kido *et al.*, *Current Eye Research* 15:833-844, 1996; Bloomer *et al.*, *Journal of Virology* 71:6641-6649, 1997; Naldini *et al.*, *Science* 272:263-267, 1996; and Miyoshi *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 94:10319, 1997). Other viral vectors that can be used

include, for example, adenoviral, lentiviral, and adena-associated viral vectors, vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus (also see, for example, the vectors of Miller, *Human Gene Thera* (1990);15-14; Friedman, *Science* 244:1275-1281, 1989; Eglitis et al., *BioTechniques* (1988);6:608-614; Tolstoshev et al.,
 5 *Cur Opin Biotechnol* (1990); 1:55-61; Sharp, *The Lancet* (1991);337:1277-78; Cornetta et al., *Nucleic Acid Research and Molecular Biology* 36:311-22, 1987; Anderson, *Science* (1984);226:401-409; Moen, *Blood Cells* 17:407-16, 1991; Miller et al., *Biotechnol* (1989);7:980-90; LeGal La Salle et al., *Science* (1993);259:988-90; and Johnson, *Chest* (1995)107:77S- 83S). Retroviral vectors are particularly well developed and have been
 10 used in clinical settings (Rosenberg et al., *N Engl J Med* (1990);323:370, 1990; Anderson et al., U.S. Patent. No. 5,399,346).

Non-viral approaches can also be employed for genetic modification of a cell. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner et al., *Proc Natl Acad Sci U.S.A.*
 15 (1987);84:7413; Ono et al., *Neurosci Lett* (1990);17:259; Brigham et al., *Am J Med Sci* (1989);298:278; Staubinger et al., *Methods in Enzymol* (1983);101:512, Wu et al., *J Biol Chem* (1988);263:14621; Wu et al., *J Biol Chem* (1989);264:16985), or by micro-injection under surgical conditions (Wolff et al., *Science* (1990);247:1465). Other non-viral means for gene transfer include transfection *in vitro* using calcium phosphate,
 20 DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transplantation of normal genes into the affected tissues of a subject can also be accomplished by transferring a normal nucleic acid into a cultivatable cell type *ex vivo* (e.g., an autologous or heterologous primary cell or progeny thereof), after which the cell (or its descendants) are injected into a targeted
 25 tissue or are injected systemically. Recombinant receptors can also be derived or obtained using transposases or targeted nucleases (e.g. Zinc finger nucleases, meganucleases, or TALE nucleases, CRISPR). Transient expression may be obtained by RNA electroporation.

Any targeted genome editing methods can also be used to deliver a presently
 30 disclosed antigen-recognizing receptor to a cell or a subject. In certain embodiments, a CRISPR system is used to deliver a presently disclosed antigen-recognizing receptor disclosed herein. In certain embodiments, zinc-finger nucleases are used to deliver the

antigen-recognizing receptor. In certain embodiments, a TALEN system is used to deliver a presently disclosed antigen-recognizing receptor.

Clustered regularly-interspaced short palindromic repeats (CRISPR) system is a genome editing tool discovered in prokaryotic cells. When utilized for genome editing, the system includes Cas9 (a protein able to modify DNA utilizing crRNA as its guide), CRISPR RNA (crRNA, contains the RNA used by Cas9 to guide it to the correct section of host DNA along with a region that binds to tracrRNA (generally in a hairpin loop form) forming an active complex with Cas9), trans-activating crRNA (tracrRNA, binds to crRNA and forms an active complex with Cas9), and an optional section of DNA repair template (DNA that guides the cellular repair process allowing insertion of a specific DNA sequence). CRISPR/Cas9 often employs a plasmid to transfect the target cells. The crRNA needs to be designed for each application as this is the sequence that Cas9 uses to identify and directly bind to the target DNA in a cell. The repair template carrying CAR expression cassette need also be designed for each application, as it must overlap with the sequences on either side of the cut and code for the insertion sequence. Multiple crRNA's and the tracrRNA can be packaged together to form a single-guide RNA (sgRNA). This sgRNA can be joined together with the Cas9 gene and made into a plasmid in order to be transfected into cells.

A zinc-finger nuclease (ZFN) is an artificial restriction enzyme, which is generated by combining a zinc finger DNA-binding domain with a DNA-cleavage domain. A zinc finger domain can be engineered to target specific DNA sequences which allows a zinc-finger nuclease to target desired sequences within genomes. The DNA-binding domains of individual ZFNs typically contain a plurality of individual zinc finger repeats and can each recognize a plurality of basepairs. The most common method to generate new zinc-finger domain is to combine smaller zinc-finger "modules" of known specificity. The most common cleavage domain in ZFNs is the non-specific cleavage domain from the type IIs restriction endonuclease FokI. Using the endogenous homologous recombination (HR) machinery and a homologous DNA template carrying CAR expression cassette, ZFNs can be used to insert the CAR expression cassette into genome. When the targeted sequence is cleaved by ZFNs, the HR machinery searches for homology between the damaged chromosome and the homologous DNA template, and then copies the sequence of the template between the two broken ends of the chromosome, whereby the homologous DNA template is integrated into the genome.

Transcription activator-like effector nucleases (TALEN) are restriction enzymes that can be engineered to cut specific sequences of DNA. TALEN system operates on almost the same principle as ZFNs. They are generated by combining a transcription activator-like effectors DNA-binding domain with a DNA cleavage domain.

5 Transcription activator-like effectors (TALEs) are composed of 33-34 amino acid repeating motifs with two variable positions that have a strong recognition for specific nucleotides. By assembling arrays of these TALEs, the TALE DNA-binding domain can be engineered to bind desired DNA sequence, and thereby guide the nuclease to cut at specific locations in genome. cDNA expression for use in polynucleotide therapy methods
10 can be directed from any suitable promoter (e.g., the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element or intron (e.g. the elongation factor 1a enhancer/promoter/intron structure). For example, if desired, enhancers known to preferentially direct gene expression in specific cell types can be used to direct the
15 expression of a nucleic acid. The enhancers used can include, without limitation, those that are characterized as tissue- or cell-specific enhancers. Alternatively, if a genomic clone is used as a therapeutic construct, regulation can be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

20 Methods for delivering the genome editing agents/systems can vary depending on the need. In certain embodiments, the components of a selected genome editing method are delivered as DNA constructs in one or more plasmids. In certain embodiments, the components are delivered via viral vectors. Common delivery methods include but is not limited to, electroporation, microinjection, gene gun, impalefection, hydrostatic pressure,
25 continuous infusion, sonication, magnetofection, adeno-associated viruses, envelope protein pseudotyping of viral vectors, replication-competent vectors cis and trans-acting elements, herpes simplex virus, and chemical vehicles (e.g., oligonucleotides, lipoplexes, polymersomes, polyplexes, dendrimers, inorganic Nanoparticles, and cell-penetrating peptides).

30 **5.6. Polypeptides**

The presently disclosed subject matter provides methods for optimizing an amino acid sequence or a nucleic acid sequence by producing an alteration in the sequence. Such alterations may include certain mutations, deletions, insertions, or post-translational

modifications. The presently disclosed subject matter further includes analogs of any naturally-occurring polypeptides disclosed herein (including, but not limited to, CD371, CD8, CD28, 4-1BB, and CD3ζ.). Analogs can differ from a naturally-occurring polypeptide disclosed herein by amino acid sequence differences, by post-translational modifications, or by both. Analogs can exhibit at least about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or more homologous or identical to all or part of a naturally-occurring amino acid sequence of the presently disclosed subject matter. The length of sequence comparison is at least 5, 10, 15 or 20 amino acid residues, *e.g.*, at least 25, 50, or 75 amino acid residues, or more than 100 amino acid residues. Again, in an exemplary approach to determining the degree of identity, a BLAST program may be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence. Modifications include *in vivo* and *in vitro* chemical derivatization of polypeptides, *e.g.*, acetylation, carboxylation, phosphorylation, or glycosylation; such modifications may occur during polypeptide synthesis or processing or following treatment with isolated modifying enzymes. Analogs can also differ from the naturally-occurring polypeptides by alterations in primary sequence. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethanemethylsulfate or by site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory Manual* (2d ed.), CSH Press, 1989, or Ausubel et al., *supra*). Also included are cyclized peptides, molecules, and analogs which contain residues other than L-amino acids, *e.g.*, D-amino acids or non-naturally occurring or synthetic amino acids, *e.g.*, β or γ amino acids.

In addition to full-length polypeptides, the presently disclosed subject matter also provides fragments of any of the polypeptides disclosed herein. As used herein, the term “a fragment” means at least 5, 10, 13, or 15 amino acids. In certain embodiments, a fragment comprises at least 20 contiguous amino acids, at least 30 contiguous amino acids, or at least 50 contiguous amino acids. In certain embodiments, a fragment comprises at least 60 to 80, 100, 200, 300 or more contiguous amino acids. Fragments can be generated by methods known to those skilled in the art or may result from normal protein processing (*e.g.*, removal of amino acids from the nascent polypeptide that are not required for biological activity or removal of amino acids by alternative mRNA splicing or alternative protein processing events).

5.7. *Formulations and Administration*

The presently disclosed subject matter also provides compositions comprising the presently disclosed cells. Compositions comprising the presently disclosed cells can be conveniently provided as sterile liquid preparations, *e.g.*, isotonic aqueous solutions, suspensions, emulsions, dispersions, or viscous compositions, which may be buffered to a selected pH. Liquid preparations are normally easier to prepare than gels, other viscous compositions, and solid compositions. Additionally, liquid compositions are somewhat more convenient to administer, especially by injection. Viscous compositions, on the other hand, can be formulated within the appropriate viscosity range to provide longer contact periods with specific tissues. Liquid or viscous compositions can comprise carriers, which can be a solvent or dispersing medium containing, for example, water, saline, phosphate buffered saline, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like) and suitable mixtures thereof.

Sterile injectable solutions can be prepared by incorporating the genetically modified cells in the required amount of the appropriate solvent with various amounts of the other ingredients, as desired. Such compositions may be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose, dextrose, or the like. The compositions can also be lyophilized. The compositions can contain auxiliary substances such as wetting, dispersing, or emulsifying agents (*e.g.*, methylcellulose), pH buffering agents, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard texts, such as “REMINGTON’S PHARMACEUTICAL SCIENCE”, 17th edition, 1985, incorporated herein by reference, may be consulted to prepare suitable preparations, without undue experimentation.

Various additives which enhance the stability and sterility of the compositions, including antimicrobial preservatives, antioxidants, chelating agents, and buffers, can be added. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin. According to the presently disclosed subject matter, however, any vehicle, diluent, or additive used would have to be compatible with the genetically modified cells.

The compositions can be isotonic, *i.e.*, they can have the same osmotic pressure as blood and lacrimal fluid. The desired isotonicity of the compositions may be accomplished using sodium chloride, or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol or other inorganic or organic solutes. Sodium chloride can be particularly for buffers containing sodium ions.

Viscosity of the compositions, if desired, can be maintained at the selected level using a pharmaceutically acceptable thickening agent. For example, methylcellulose is readily and economically available and is easy to work with. Other suitable thickening agents include, for example, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, carbomer, and the like. The concentration of the thickener can depend upon the agent selected. The important point is to use an amount that will achieve the selected viscosity. Obviously, the choice of suitable carriers and other additives will depend on the exact route of administration and the nature of the particular dosage form, *e.g.*, liquid dosage form (*e.g.*, whether the composition is to be formulated into a solution, a suspension, gel or another liquid form, such as a time release form or liquid-filled form).

Compositions comprising the presently disclosed cells can be provided systemically or directly to a subject for inducing and/or enhancing an immune response to an antigen and/or treating and/or preventing a neoplasia. In certain embodiments, the presently disclosed cells or compositions comprising thereof are directly injected into an organ of interest (*e.g.*, an organ affected by a neoplasia). Alternatively, the presently disclosed cells or compositions comprising thereof are provided indirectly to the organ of interest, for example, by administration into the circulatory system (*e.g.*, the tumor vasculature). Expansion and differentiation agents can be provided prior to, during or after administration of the cells or compositions to increase production of cells (*e.g.*, T cells or NK cells) *in vitro* or *in vivo*.

The presently disclosed cells can be administered in any physiologically acceptable vehicle, normally intravascularly, although they may also be introduced into bone or other convenient site where the cells may find an appropriate site for regeneration and differentiation (*e.g.*, thymus).

The quantity of cells to be administered can vary for the subject being treated. In certain embodiments, between about 10^4 and about 10^{10} , between about 10^4 and about 10^7 , between about 10^5 and about 10^7 , between about 10^5 and about 10^9 , or between about 10^6 and about 10^8 of the presently disclosed cells are administered to a subject. More

effective cells may be administered in even smaller numbers. Usually, at least about 1×10^5 cells will be administered, eventually reaching about 1×10^{10} or more. In certain embodiments, at least about 1×10^5 , 5×10^5 , 1×10^6 , about 5×10^6 , about 1×10^7 , about 5×10^7 , about 1×10^8 , or about 5×10^8 of the presently disclosed cells are administered to a subject.

5 In certain embodiments, about 1×10^6 of the presently disclosed cells are administered to a subject. The precise determination of what would be considered an effective dose can be based on factors individual to each subject, including their size, age, sex, weight, and condition of the particular subject. Dosages can be readily ascertained by those skilled in the art from this disclosure and the knowledge in the art.

10 The presently disclosed cells can comprise a purified population of cells. Those skilled in the art can readily determine the percentage of the presently disclosed cells in a population using various well-known methods, such as fluorescence activated cell sorting (FACS). Suitable ranges of purity in populations comprising the presently disclosed immunoresponsive cells are about 50% to about 55%, about 5% to about 60%, and about
15 65% to about 70%. In certain embodiments, the purity is about 70% to about 75%, about 75% to about 80%, or about 80% to about 85%. In certain embodiments, the purity is about 85% to about 90%, about 90% to about 95%, and about 95% to about 100%. Dosages can be readily adjusted by those skilled in the art (*e.g.*, a decrease in purity may require an increase in dosage). The cells can be introduced by injection, catheter, or the
20 like.

The skilled artisan can readily determine the amount of cells and optional additives, vehicles, and/or carrier in compositions and to be administered in methods. Typically, any additives (in addition to the active cell(s) and/or agent(s)) are present in an amount of 0.001 to 50% (weight) solution in phosphate buffered saline, and the active
25 ingredient is present in the order of micrograms to milligrams, such as about 0.0001 to about 5 wt %, about 0.0001 to about 1 wt %, about 0.0001 to about 0.05 wt% or about 0.001 to about 20 wt %, about 0.01 to about 10 wt %, or about 0.05 to about 5 wt %. For any composition to be administered to an animal or human, the followings can be determined: toxicity such as by determining the lethal dose (LD) and LD50 in a suitable
30 animal model *e.g.*, rodent such as mouse; the dosage of the composition(s), concentration of components therein and timing of administering the composition(s), which elicit a suitable response. Such determinations do not require undue experimentation from the

knowledge of the skilled artisan, this disclosure and the documents cited herein. And, the time for sequential administrations can be ascertained without undue experimentation.

In certain embodiments, the composition is a pharmaceutical composition comprising the presently disclosed cells and a pharmaceutically acceptable carrier.

5 Administration of the compositions can be autologous or heterologous. For example, cells can be obtained from one subject, and administered to the same subject or a different, compatible subject. Peripheral blood derived cells or their progeny (*e.g.*, *in vivo*, *ex vivo* or *in vitro* derived) can be administered. When administering a presently disclosed composition (*e.g.*, a pharmaceutical composition comprising presently disclosed
10 cells), it can be formulated in a unit dosage injectable form (solution, suspension, emulsion).

The presently disclosed cells and compositions can be administered by any method known in the art including, but not limited to, oral administration, intravenous administration, subcutaneous administration, intranodal administration, intratumoral
15 administration, intrathecal administration, intrapleural administration, intraosseous administration, intraperitoneal administration, pleural administration, and direct administration to the subject.

5.8. *Methods of Treatment*

The presently disclosed subject cells and compositions comprising thereof can be
20 used for treating and/or preventing a tumor or neoplasm. Such cells can be administered to a subject (*e.g.*, a human subject) in need thereof for treatment and/or prevention of a tumor or neoplasm (*e.g.*, acute myeloid leukemia (AML), multiple myeloma, Non-Hodgkin's Lymphoma, Hodgkin's Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma). In certain embodiments, the cell is a T cell. The T cell can be a CD4⁺ T
25 cell or a CD8⁺ T cell. In certain embodiments, the T cell is a CD4⁺ T cell.

The presently disclosed subject matter provides methods for inducing and/or increasing an immune response in a subject in need thereof. The presently disclosed cells and compositions comprising thereof can be used in a therapy or medicament. The presently disclosed subject matter provides various methods of using the cells (*e.g.*, T
30 cells) or compositions comprising thereof. For example, the presently disclosed cells and compositions comprising thereof can be used for reducing tumor burden in a subject. The presently disclosed cell can reduce the number of tumor cells, reduce tumor size, and/or eradicate the tumor in the subject. The presently disclosed cells and compositions

comprising thereof can be used for treating and/or preventing a tumor or neoplasm in a subject. The presently disclosed cells and compositions comprising thereof can be used for prolonging the survival of a subject suffering from a tumor or neoplasm. Such methods comprise administering the presently disclosed cells or a composition (*e.g.*, a pharmaceutical composition) comprising thereof to achieve the desired effect, *e.g.*, palliation of an existing condition or prevention of recurrence. For treatment, the amount administered is an amount effective in producing the desired effect. An effective amount can be provided in one or a series of administrations. An effective amount can be provided in a bolus or by continuous perfusion.

The presently disclosed subject matter provides various methods of using the cells (*e.g.*, T cells) or compositions comprising thereof. For example, the presently disclosed subject matter provides methods of reducing tumor burden in a subject. In certain embodiments, the method of reducing tumor burden comprises administering the presently disclosed cells or a composition comprising thereof to the subject. The presently disclosed cell can reduce the number of tumor cells, reduce tumor size, and/or eradicate the tumor in the subject.

The tumor or neoplasm can be a solid tumor. Non-limiting examples of solid tumors include mesothelioma, lung cancer, pancreatic cancer, ovarian cancer, breast cancer, colon cancer, pleural tumor, glioblastoma, esophageal cancer, gastric cancer, synovial sarcoma, thymic carcinoma, endometrial carcinoma, stomach cancer, and cholangiocarcinoma.

The presently disclosed subject matter also provides methods of increasing or lengthening survival of a subject having a tumor or neoplasm. In certain embodiments, the method of increasing or lengthening survival of a subject having a tumor or neoplasm comprises administering the presently disclosed immunoresponsive cells or a composition comprising thereof to the subject. The method can reduce or eradicate tumor burden in the subject. Additionally, the presently disclosed subject matter provides methods for increasing an immune response in a subject, comprising administering the presently disclosed cell or a composition comprising thereof to the subject. The presently disclosed subject matter further provides methods for treating and/or preventing a tumor or neoplasm in a subject, comprising administering the presently disclosed cells or a composition comprising thereof to the subject.

Non-limiting examples of neoplasms or tumors include acute myeloid leukemia (AML), multiple myeloma, Chronic Lymphocytic Leukemia (CLL), lymphoma (Hodgkin's lymphoma, non-Hodgkin's lymphoma), glioblastoma, myelodysplastic syndrome (MDS), and chronic myelogenous leukemia (CML), bone cancer, intestinal cancer, liver cancer, skin cancer, cancer of the head or neck, melanoma (cutaneous or intraocular malignant melanoma), renal cancer (*e.g.* clear cell carcinoma), throat cancer, prostate cancer (*e.g.* hormone refractory prostate adenocarcinoma), blood cancers (*e.g.* leukemias, lymphomas, and myelomas), uterine cancer, rectal cancer, cancer of the anal region, bladder cancer, brain cancer, stomach cancer, testicular cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, leukemias (*e.g.*, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute monocytic leukemia, acute erythroleukemia, chronic leukemia, chronic myelocytic leukemia, , polycythemia vera, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, lymphocytic lymphoma, cancer of the bladder, cancer of the kidney or ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally induced cancers including those induced by asbestos, include Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors such as sarcomas and carcinomas (*e.g.*, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, hepatoma, nile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, salivary gland cancer, uterine cancer, testicular cancer, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma,

hemangioblastoma, acoustic neuroma, oligodendroglioma, schwannoma, meningioma, melanoma, neuroblastoma, and retinoblastoma).

In certain embodiments, the tumor or neoplasm is selected from the group consisting of acute myeloid leukemia (AML), multiple myeloma, Chronic Lymphocytic Leukemia (CLL), Hodgkin's lymphoma, non-Hodgkin's lymphoma, glioblastoma, myelodysplastic syndrome (MDS), and chronic myelogenous leukemia (CML). In certain
5 embodiments, the tumor or neoplasm is AML.

The subjects can have an advanced form of disease, in which case the treatment objective can include mitigation or reversal of disease progression, and/or amelioration of
10 side effects. The subjects can have a history of the condition, for which they have already been treated, in which case the therapeutic objective will typically include a decrease or delay in the risk of recurrence.

As a consequence of surface expression of a presently disclosed CD371-targeted antigen-recognizing receptor, adoptively transferred cells (e.g., immunoresponsive cells, e.g., T cells or NK cells) are endowed with augmented and selective cytolytic activity at
15 the tumor site. Furthermore, subsequent to their localization to tumor or viral infection and their proliferation, the cells turn the tumor or viral infection site into a highly conducive environment for a wide range of immune cells involved in the physiological anti-tumor or antiviral response (tumor infiltrating lymphocytes, NK-, NKT- cells, dendritic cells, and macrophages).
20

Further modification can be introduced to the presently disclosed cells (e.g., T cells) to avert or minimize the risks of immunological complications (known as "malignant T-cell transformation"), e.g., graft versus-host disease (GvHD), or when healthy tissues express the same target antigens as the tumor cells, leading to outcomes
25 similar to GvHD. A potential solution to this problem is engineering a suicide gene into the presently disclosed cells. Suitable suicide genes include, but are not limited to, Herpes simplex virus thymidine kinase (hsv-tk), inducible Caspase 9 Suicide gene (iCasp-9), and a truncated human epidermal growth factor receptor (EGFRt) polypeptide. In certain embodiments, the suicide gene is an EGFRt polypeptide. The EGFRt polypeptide
30 can enable T cell elimination by administering anti-EGFR monoclonal antibody (e.g., cetuximab). EGFRt can be covalently joined to the upstream of the antigen-recognizing receptor (e.g., CAR). The suicide gene can be included within the vector comprising nucleic acids encoding a presently disclosed antigen-recognizing receptor (e.g., CAR). In

this way, administration of a prodrug designed to activate the suicide gene (*e.g.*, a prodrug (*e.g.*, AP1903 that can activate iCasp-9) during malignant T-cell transformation (*e.g.*, GVHD) triggers apoptosis in the suicide gene-activated cells expressing the antigen-recognizing receptor (*e.g.*, CAR). The incorporation of a suicide gene into the a presently disclosed antigen-recognizing receptor (*e.g.*, CAR) gives an added level of safety with the ability to eliminate the majority of receptor-expressing cells within a very short time period. A presently disclosed cell (*e.g.*, a T cell) incorporated with a suicide gene can be pre-emptively eliminated at a given timepoint post the cell infusion, or eradicated at the earliest signs of toxicity.

5.9. Kits

The presently disclosed subject matter provides kits for inducing and/or enhancing an immune response in a subject, treating and/or preventing a tumor or neoplasm in a subject, reducing tumor burden in a subject, and/or increasing or lengthening survival of a subject having a tumor or neoplasm in a subject. In certain embodiments, the kit comprises the presently disclosed cells or a composition comprising thereof. In certain embodiments, the kit comprises a sterile container; such containers can be boxes, ampules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. Such containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments. In certain non-limiting embodiments, the kit includes a nucleic acid molecule encoding a presently disclosed CD371-targeted antigen-recognizing receptor (*e.g.*, a CAR or a TCR).

If desired, the cells and/or nucleic acid molecules are provided together with instructions for administering the cells or nucleic acid molecules to a subject having or at risk of developing a tumor or neoplasm. The instructions generally include information about the use of the composition for the treatment and/or prevention of a tumor or neoplasm. In certain embodiments, the instructions include at least one of the following: description of the therapeutic agent; dosage schedule and administration for treatment or prevention of a tumor or neoplasm; precautions; warnings; indications; counter-indications; over-dosage information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container.

6. EXAMPLES

The practice of the present disclosure employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are well within the purview of the skilled artisan. Such techniques are explained fully in the literature, such as, “Molecular Cloning: A Laboratory Manual”, second edition (Sambrook, 1989); “Oligonucleotide Synthesis” (Gait, 1984); “Animal Cell Culture” (Freshney, 1987); “Methods in Enzymology” “Handbook of Experimental Immunology” (Weir, 1996); “Gene Transfer Vectors for Mammalian Cells” (Miller and Calos, 1987); “Current Protocols in Molecular Biology” (Ausubel, 1987); “PCR: The Polymerase Chain Reaction”, (Mullis, 1994); “Current Protocols in Immunology” (Coligan, 1991). These techniques are applicable to the production of the polynucleotides and polypeptides disclosed herein, and, as such, may be considered in making and practicing the presently disclosed subject matter. Particularly useful techniques for particular embodiments will be discussed in the sections that follow.

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the presently disclosed cells and compositions, and are not intended to limit the scope of what the inventors regard as their invention.

Example 1 – Generation of CD371-targeted CARs

The following six CD371-targeted CARs were generated:

“Et.B10L3H_MT_h28z”, “Et.B10L4H_MT_hBBz”, “Et.B10L4H_MT_h28z”, “Et.B10H3L_MT_h28z”, “Et.B10H4L_MT_h28z”, “Et.C3H3L_MT_h28z”, “Et.C3L3H_MT_h28z”, and “Et.D6L3H_MT_h28z”. The structures of the CARs are illustrated in Figure 1.

As shown in Figure 2, the CARs were detected by EGFR and Myc-TAG on the surface of transduced human T cells. Antibodies against EGFRt (CETUXIMAB-APC) and Myc-tag (9B11-PE) were used to detect surface expression of EGFRt and human anti-human CAR in B10-based constructs. Controls included non-transduced human T cells not expressing EGFRt or Myc-tag, and non-Myc Tag containing CAR constructs Etah19h28Z (an anti-human CD19 CAR T cell) and EtC1HVh28Z (an anti-CD371 CAR T cell derived from a mouse anti-human CD371 antibody 1075.7)

Example 2 – In vitro cytotoxicity activities of CD371-targeted CAR T cells

T cells were transfected with a number of CD371-targeted CARs (i.e., “Et.C1HVh28Z”, “Et.B10LH_MT_h28Z”, “Et.B10LHg4s_MT_h28Z”, “Et.B10HL_MT_h28Z”, “Et.C3HL_MT_h28Z”, “Et.C3LH_MT_h28Z”, and “Et.D6LH_MT_h28Z”). Et.C1HVh28Z was used as a positive control and is derived from a mouse anti-human CD371 antibody 1075.7. “Et.B10LHg4s_MT_hDEL” is a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain). These CD371-targeted CAR T cells were cocultured with CD371+ U937 cells expressing GFP and firefly luciferase (U937gL) at different effector:tumor ratios. 24 hours later, bioluminescence was measured and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL. The results are shown in Figure 3. As shown in Figure 3, CD371-targeted CAR T cells were able to lyse cells expressing CD371.

Example 3 – In vitro cytotoxicity activities of CD371-targeted CAR T cells

T cells were transfected with a number of CD371-targeted CARs (i.e., “Et.C1HVh28Z”, “Et.B10L4H_MT_h28Z”, “Et.B10L3H_MT_h28Z”, “Et.B10L4H_MT_hBBZ”, “Et.B10H3L_MT_h28Z”, “Et.C3HL_MT_h28Z”, “Et.C3LH_MT_h28Z”, and “Et.M195_MT_h28Z”). Et.C1HVh28Z was used as a positive control and is derived from a mouse anti-human antibody 1075.7). “Et.M195MTh28Z” represents a CD33-targeted CAR T cell derived from the murine anti-human CD33 antibody, M195. “Et.M195MTh28Z” used as a positive control as U937 cells also express CD33. Healthy human donor-derived (i.e., Donor C and Donor D) CD371-targeted CAR T cells were cocultured with CD33+/CD371+ U937 cells expressing GFP and firefly luciferase (U937gL) at different effector:tumor ratios. 24 hours later, bioluminescence was measured and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL. The results are shown in Figure 4 (for Donor C) and Figure 5 (for Donor D). As shown in Figures 4 and 5, B10-based CARs (both HL/LH) T cells outperformed C3-based CAR T cells.

Example 4 – Antigen-stimulation of CD371-targeted CAR T cells

Healthy human donor-derived (i.e., Donor C and Donor D) CD371-targeted CAR T cells were cocultured with CD33+/CD371+ U937gL at an E:T ratio of 1:12.5 and at a

concentration of 30,000 CAR+/mL. Roughly every 4-5 days, CAR T cells were counted and characterized by flow cytometry, and the starting number of tumor cells were added back into the culture (indicated by arrows). The results are shown in Figure 6 (for Donor C) and Figure 7 (for Donor D). “Et.B10LHdel” represents a non-functional CAR T cell (lacks signaling domains). “Et.C1HVh28Z” represents a CD371-targeted CAR derived from a mouse anti-human CD371 antibody 1075.7), and “EtM195MTh28Z” or “M19528” represents a CD33-targeted CAR T cell derived from the murine anti-human CD33 antibody, M195. As shown in Figures 6 and 7, B10-based CAR T cells outperformed C3-based CAR T cells in a long-term repeated stimulation assay.

10 ***Example 5 – In vitro cytotoxicity activities of CD371-targeted CAR T cells***

Healthy human donor-derived (i.e., Donor A and Donor B) CD371-targeted CAR T cells were cocultured with CD371-positive U937 cells or HL60 cells expressing GFP and firefly luciferase (U937gL or HL60gL) at different effector:tumor ratios.

Bioluminescence was measured 24 hours after coculture, and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937gL. The results are shown in Figures 8A and 8B (for U937gL cells) and Figures 9A and 9B (for HL 60 cells). As shown in Figures 8A-8B and 9A-9B, B10HL-based CD371-targeted CAR T cells outperformed B10LH-based CD371-targeted CAR T cells.

20 ***Example 6 – Antigen-specific cytotoxicity activities of CD371-targeted CAR T cells***

Whether the cytotoxicity activities of the CD371-targeted CAR T cells are specific to CD371 was assessed. Human CD371 (hCD371) CRISPR knockout cells lines were generated. Human CD371 was knocked out of HL60gL and U937gL with CRISPR, and knockout was confirmed by flow cytometry using anti-human CD371 APC-conjugated antibody. See Figure 10.

Healthy human donor-derived (i.e., Donor A and Donor B) CD371-targeted CAR T cells were cocultured with antigen-negative U937 cells or HL60 expressing RFP and cypridina luciferase (U937RFPcyp.371KO or HLgL.371KO) at different effector:tumor ratios. 24 hours later, bioluminescence was measured and plotted as a percentage of signal detected in a coculture of a non-functional CD371-targeted CAR T cells (based on B10L4H, but without a CD28 or CD3 zeta signaling domain) and U937RFPcyp.371KO. The results are shown in Figures 11A-11B (for U937 cells) and Figures 12A-12B (for HL60 cells). As shown in Figures 11A-11B and 12A-12B, both B10HL-based and

B10LH-based CD371-targeted CAR T cells showed no significant cytotoxicity against CD371-negative U937 cells, and showed limited cytotoxicity against CD371-negative HL60 cells.

Example 7 – Cytokines secretion by the CD371-targeted CAR T cells

5 The ability for producing cytokines by the CD371-targeted CAR T cells was assessed. CD371-targeted CAR T cells were cocultured alone, with CD371-negative tumor cells, or CD371-positive U937 cells at an effector:tumor ratio of 1:1 (40,000:40,000 cells in 200 μ L). 24 hours later, supernatant was collected from the culture and the secretion levels of IFN- γ , IL-2, and TNF- α were measured utilizing a
10 bead-based multiplex assay. The results are shown in Figures 13A-13B (for IFN- γ), Figures 14A-14B (for IL-2), and Figures 15A-15B (for TNF- α). As shown in Figures 13A-13B, Figures 14A-14B, and Figures 15A-15B, B10HL-based CAQR T cells produced more cytokines in an antigen-dependent manner compared to B10LH-based CAR T cells.

15 ***Example 8 – Proliferation Capacity of CD371-targeted CAR T cells***

CAR T cells were generated from two healthy donors (i.e., Donor A and Donor B). CD371-targeted CAR T cells were cocultured with CD371+ U937gL at an E:T ratio of 1:5 and at a concentration of 50,000 CAR+/mL. CAR T cells were counted approximately every 5 days, and characterized by flow cytometry. The starting number of
20 tumor cells were added back into the culture at the time point indicated by arrows. The results are shown in Figure 16. As shown in Figure 16, B10HL-based and B10-LH-based CD371-targeted CAR T cells showed near-comparative expansion capacities *in vitro*.

Example 9 – In vivo Anti-tumor activities of CD371-targeted CAR T cells

In vivo anti-tumor activities of the B10-based CD371-targeted CAR T cells were
25 assessed. Figure 17 shows the xenograft model of AML. NCG mice were inoculated with 5.0×10^4 U937gL tumor cells, AML FAB-M5 cell line via tail vein, and treated with approximately 1.25×10^6 CAR T cells three days later. Mice survival was monitored. The results are shown in Figure 18 (for Donor 1) and Figure 19 (for Donor 2). As shown in Figures 18 and 19, B10HL-based CAR T cells showed superior survival in an AML
30 xenografted mice as compared to B10LH-based CAR T cells.

Next, whether the *in vivo* activities are dose-dependent was assessed. NCG mice were inoculated with 5.0×10^4 U937gL tumor cells, and treated with various doses of CAR T cells (1.25×10^5 , 2.5×10^5 , 5.0×10^5 , and 1.0×10^6) three days later. The results

are shown in Figure 20. As shown in Figure 20, B10HL-based CAR T cells outperformed B10LH-based CAR T cells in a dose-response treatment model.

Example 10: B10 scFvs Binding to CD371-expressing Cell Lines

The binding of B10 (also referred to as “1B10”) scFvs both orientations (V_H - V_L or V_L - V_H) to CD371-expressing cells was assessed. The results are shown in Figure 21. As shown in Figure 21, binding to cells was detected for the B10 scFv in the V_L - V_H orientation. However, binding of the B10 scFv in the V_H - V_L orientation was not observed by flow cytometry, suggesting a lower affinity and thus a requirement for bivalent binding.

Example 11: B10 scFvs Binding to Recombinant CD371 in Solution

For scFv affinity measurements, biotinylated CD371 was captured with streptavidin, and soluble scFv was used as analyte. Table 8 shows dissociation constants (K_D), on-rates (k_{on}) and off-rates (k_{off}) for the different scFv formats. Consistent with the flow cytometry results, weak binding of the 1B10 scFv in the V_H - V_L orientation was observed but a dissociation constant could not be calculated by any curve fit method.

Thus, it is surprising that the CARs comprising the lower binding affinity scFv outperformed *in vitro* and *in vivo* than CARs comprising the higher binding affinity scFv, as shown in Examples 5, 7 and 9.

Table 8. Binding affinity of B10 scFvs in V_H - V_L or V_L - V_H orientation to soluble CD371

Format	K_D (nM)	K_{on} (1/Ms)	K_{off} (1/s)
V_L - V_H scFv	16	5.99×10^5	9.52×10^{-3}
V_H - V_L scFv	Weak binding	NA	NA

Example 12: In vivo and in vitro activities of B10HL-based second-generation CAR T cells

B10HL-based second-generation CAR T cells having a truncated EGFR (Et), a Myc-tag (MT), and G4S linkers of varying lengths were successfully generated:

- Et.B10H4L_MT_h28Z is described in Example 1.
- Et.B10H1L_MT_h28Z comprised a B10 based scFv (designated as B10H1L) having V_H - V_L orientation and a 5 amino acids long G4S linker (GGGGS [SEQ ID NO: 93]), and a CD28Z-based signaling domain.

- Et.B10H2L_MT_h28Z comprised a B10 based scFv (designated as B10H2L) having V_H-V_L orientation and a 10 amino acids long G4S linker (GGGGSGGGGS [SEQ ID NO: 94]), and a CD28Z-based signaling domain.
- Et.B10H5L_MT_h28Z comprised a B10 based scFv (designated as B10H5L) having V_H-V_L orientation and a 24 amino acids long G4S linker (GGGGSGGGGS GGGGSGGGSGGGGS [SEQ ID NO: 91]), and a CD28Z-based signaling domain.
- Et.B10H6L_MT_h28Z comprised a B10 based scFv (designated as B10H6L) having V_H-V_L orientation and a 29-amino acid G4S linker (GGGGSGGGGS GGGGSGGGSGGGSGGGGS [SEQ ID NO: 92]), and a CD28Z-based signaling domain.

Additional B10H4L-based second-generation CAR T cells incorporating either constitutive IL18 (designated as “Et.B10H4Lmt28ZpIL18”) or IL33 (designated as “Et.B10H4Lmt28ZpIL33”) secretion were generated. T cells comprising

Et.B10H4Lmt28ZpIL18 comprise the CAR designated as “Et.B10H4L_MT_h28Z” and an exogenous IL-18 polypeptide (e.g., an exogenous IL-18 polypeptide comprising the amino acid sequence set forth in SEQ ID NO 103). T cells comprising Et.B10H4Lmt28ZpIL33 comprise the CAR designated as “Et.B10H4L_MT_h28Z” and an exogenous IL-33 polypeptide (e.g., an exogenous IL-33 polypeptide comprising the amino acid sequence set forth in SEQ ID NO 105).

As shown in Figure 22, CAR T cells having V_H-V_L oriented scFvs had superior cytotoxicity against U937 cells than CAR T cells having V_L-V_H oriented scFvs. To assess the *in vivo* activity of these B10 based CAR T cells, NCG mice were inoculated with 5×10^4 U937gL tumor cells, and were treated with various doses of CAR T cells. Survival of the treated mice were monitored recorded. As shown in Figure 23, at the dose of 5×10^5 CAR T cells, B10H3L, B10H5L, and IL33-secreting B10H4L based human CD371-targeted CAR T cells outperformed B10H4L-based CAR T cells. As shown in Figure 24, at the dose of 2.5×10^5 CAR T cells, IL18- and IL33-secreting B10H4L based human CD371-targeted CAR T cells outperformed B10H4L-based CAR T cells. As shown in Figure 25, at the low dose of 1.0×10^5 CAR T cells, IL18-secreting B10H4L human CD371-targeted CAR T cells outperform other CAR T cells.

Embodiments of the presently disclosed subject matter

From the foregoing description, it will be apparent that variations and modifications may be made to the presently disclosed subject matter to adopt it to various
5 usages and conditions. Such embodiments are also within the scope of the following claims.

The recitation of a listing of elements in any definition of a variable herein includes definitions of that variable as any single element or combination (or sub-combination) of listed elements. The recitation of an embodiment herein includes that
10 embodiment as any single embodiment or in combination with any other embodiments or portions thereof.

All patents and publications mentioned in this specification are herein incorporated by reference to the same extent as if each independent patent and publication was specifically and individually indicated to be incorporated by reference.

WHAT IS CLAIMED IS:

1. An antigen-recognizing receptor, comprising an extracellular antigen-binding domain, a transmembrane domain, and an intracellular signaling domain, wherein the extracellular antigen-binding domain specifically binds to CD371.
2. The antigen-recognizing receptor of claim 1, wherein the extracellular antigen-binding domain is a single-chain variable fragment (scFv).
3. The antigen-recognizing receptor of claim 2, wherein the extracellular antigen-binding domain is a human scFv.
4. The antigen-recognizing receptor of claim 1, wherein the extracellular antigen-binding domain is a Fab, which is optionally crosslinked.
5. The antigen-recognizing receptor of claim 1, wherein the extracellular antigen-binding domain is a F(ab)₂.
6. The antigen-recognizing receptor of any one of claims 2-5, wherein one or more of the scFv, Fab and F(ab)₂ are comprised in a fusion protein with a heterologous sequence to form the extracellular antigen-binding domain.
7. The antigen-recognizing receptor of any one of claims 1-6, wherein the extracellular antigen-binding domain comprises:
 - (a) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30 or a conservative modification thereof;
 - (b) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36 or a conservative modification thereof;

(c) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 42 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 43 or a conservative modification thereof;

(d) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 462 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 48 or a conservative modification thereof;

(e) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54 or a conservative modification thereof; or

(f) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58 or a conservative modification thereof; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59 or a conservative modification thereof; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60 or a conservative modification thereof;

8. The antigen-recognizing receptor of any one of claims 1-7, wherein the extracellular antigen-binding domain comprises: a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; and a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30.

9. The antigen-recognizing receptor of any one of claims 1-8, wherein the extracellular antigen-binding domain comprises:

(a) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33 or a conservative modification thereof;

(b) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38 or a conservative modification thereof; and a light chain variable region CDR3 comprising SEQ ID NO: 39 or a conservative modification thereof;

(c) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45 or a conservative modification thereof;

(d) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 49 or a conservative modification thereof; a light chain variable region CDR2 comprising SEQ ID NO: 50 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51 or a conservative modification thereof;

(e) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 55 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57 or a conservative modification thereof;
or

(f) a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 61 or a conservative modification thereof; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62 or a conservative modification thereof; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63 or a conservative modification thereof;

10. The antigen-recognizing receptor of any one of claims 1-9, wherein the extracellular antigen-binding domain comprises: a light chain variable region CDR1

comprising the amino acid sequence set forth in SEQ ID NO: 31; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33.

11. The antigen-recognizing receptor of any one of claims 1-10, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33;

(b) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 34; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 35; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 36; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 37; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 38; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 39;

(c) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 40; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 41; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 42; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 43; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 44; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 45;

(d) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 46; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 47; a heavy chain variable region CDR3 comprising

the amino acid sequence set forth in SEQ ID NO: 48; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 49; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 50; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 51;

(e) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 52; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 53; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 54; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 55; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 56; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 57; or

(f) a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 58; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 59; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 60; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 61; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 62; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 63.

12. The antigen-recognizing receptor of any one of claims 1-11, wherein the extracellular antigen-binding domain comprises: a heavy chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 28; a heavy chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 29; a heavy chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 30; a light chain variable region CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 31; a light chain variable region CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 32; and a light chain variable region CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 33.

13. The antigen-recognizing receptor of any one of claims 1-12, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising

an amino acid sequence that is at least about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, or SEQ ID NO: 11.

14. The antigen-recognizing receptor of any one of claims 1-13, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, or SEQ ID NO:11.

15. The antigen-recognizing receptor of any one of claims 1-14, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising amino acids having a sequence set forth in SEQ ID NO: 1.

16. The antigen-recognizing receptor of any one of claims 1-15, wherein the extracellular antigen-binding domain comprises a light chain variable region comprising an amino acid sequence that is at least about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, or SEQ ID NO: 12.

17. The antigen-recognizing receptor of any one of claims 1-16, wherein the extracellular antigen-binding domain comprises a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, or SEQ ID NO: 12.

18. The antigen-recognizing receptor of any one of claims 1-17, wherein the extracellular antigen-binding domain comprises a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2.

19. The antigen-recognizing receptor of any one of claims 1-18, wherein the extracellular antigen-binding domain comprises: (a) a heavy chain variable region

comprising an amino acid sequence that is at least about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the amino acid sequence selected set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, or SEQ ID NO: 11; and (b) a light chain variable region comprising an amino acid sequence that is at least about 80%, about 81%, about 82%, about 83%, about 84%, about 85%, about 86%, about 87%, about 88%, about 89%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99% homologous or identical to the amino acid sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, or SEQ ID NO: 12.

20. The antigen-recognizing receptor of any one of claims 1-19, wherein the extracellular antigen-binding domain comprises: (a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, or SEQ ID NO: 11; and (b) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, or SEQ ID NO: 12.

21. The antigen-recognizing receptor of claim 14, wherein the extracellular antigen-binding domain comprises:

(a) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2;

(b) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 3, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 4;

(c) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 5, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 6;

(d) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 7, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 8;

(e) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 9, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 10; or

(f) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 11, and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 12.

22. The antigen-recognizing receptor of any one of claims 1-21, wherein the extracellular antigen-binding domain comprises a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 1; and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 2.

23. The antigen-recognizing receptor of any one of claims 1-22, wherein the extracellular antigen-binding domain comprises a linker between a heavy chain variable region and a light chain variable region of the extracellular antigen-binding domain.

24. The antigen-recognizing receptor of claim 23, wherein the linker has the amino acid sequence set forth in SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 91, SEQ ID NO: 92, SEQ ID NO: 93, or SEQ ID NO: 94.

25. The antigen-recognizing receptor of claim 23, wherein the linker has the amino acid sequence set forth in SEQ ID NO: 13 or SEQ ID NO: 14.

26. The antigen-recognizing receptor of any one of claims 1-25, wherein the extracellular antigen-binding domain comprises a signal peptide that is covalently joined to the 5' terminus of the extracellular antigen-binding domain.

27. The antigen-recognizing receptor of any one of claims 1-26, wherein the extracellular antigen-binding domain comprises a heavy chain variable region and a light chain variable region, which are positioned from the N- to the C-terminus: V_H-V_L.

28. The antigen-recognizing receptor of any one of claims 1-26, wherein the extracellular antigen-binding domain binds to CD371 with a low binding affinity.

29. The antigen-recognizing receptor of any one of claims 1-26, wherein the extracellular antigen-binding domain binds to CD371 with a dissociation constant (K_d) of 1×10^{-8} M or more.
30. The antigen-recognizing receptor of any one of claims 1-29, wherein the transmembrane domain comprises a CD8 polypeptide, a CD28 polypeptide, a CD3 ζ polypeptide, a CD4 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a CTLA-4 polypeptide, a PD-1 polypeptide, a LAG-3 polypeptide, a 2B4 polypeptide, a BTLA polypeptide, or a combination thereof.
31. The antigen-recognizing receptor of claim 30, wherein the transmembrane domain comprises a CD28 polypeptide.
32. The antigen-recognizing receptor of any one of claims 1-31, wherein the intracellular signaling domain comprises a CD3 ζ polypeptide.
33. The antigen-recognizing receptor of any one of claims 1-32, wherein the intracellular signaling domain further comprises at least one co-stimulatory signaling region.
34. The antigen-recognizing receptor of claim 33, wherein the at least one co-stimulatory signaling region comprises a CD28 polypeptide, a 4-1BB polypeptide, an OX40 polypeptide, an ICOS polypeptide, a DAP-10 polypeptide, or a combination thereof.
35. The antigen-recognizing receptor of claim 33 or 34, wherein the at least one co-stimulatory signaling region comprises a CD28 polypeptide or a 4-1BB polypeptide.
36. The antigen-recognizing receptor of any one of claims 1-35, wherein the antigen-recognizing receptor is a chimeric antigen receptor (CAR), a T-cell Receptor (TCR), or a T-cell like fusion protein.
37. The antigen-recognizing receptor of any one of claims 1-36, wherein the antigen-recognizing receptor is a CAR.
38. The antigen-recognizing receptor of any one of claims 1-37, wherein the antigen-recognizing receptor is recombinantly expressed.

39. The antigen-recognizing receptor of any one of claims 1-38, wherein the antigen-recognizing receptor is expressed from a vector.
40. The antigen-recognizing receptor of claim 39, wherein the vector is a γ -retroviral vector.
41. A cell comprising the antigen-recognizing receptor of any one of claims 1-40.
42. The cell of claim 41, wherein the cell is transduced with the antigen-recognizing receptor.
43. The cell of claim 41 or 42, wherein the antigen-recognizing receptor is constitutively expressed on the surface of the cell.
44. The cell of any one of claims 41-43, wherein the cell is engineered to express a cytokine or a fragment thereof.
45. The cell of claim 44, wherein the cell comprises an exogenous polypeptide of the cytokine or fragment thereof.
46. The cell of claim 44 or 45, wherein the cell comprises a nucleic acid molecule encoding the cytokine or fragment thereof.
47. The cell of any one of claims 44-46, wherein the cytokine is selected from the group consisting of IL-18, IL-33, IL-36, and combinations thereof.
48. The cell of any one of claims 44-47, wherein the cytokine is IL-18.
49. The cell of any one of claims 41-48, wherein the cell is an immunoresponsive cell.
50. The cell of any one of claims 41-49, wherein the cell is a cell of the lymphoid lineage or a cell of the myeloid lineage.
51. The cell of any one of claims 41-50, wherein the cell is selected from the group consisting of a T cell, a Natural Killer (NK) cell, and a stem cell from which lymphoid cells may be differentiated.
52. The cell of any one of claims 41-51, wherein the cell is a T cell.

53. The cell of claim 51 or 52, wherein the T cell is a cytotoxic T lymphocyte (CTL) or a regulatory T cell.
54. The cell of claim 53, wherein the stem cell is a pluripotent stem cell.
55. The cell of claim 54, wherein the pluripotent stem cell is an embryoid stem cell or an induced pluripotent stem cell.
56. A nucleic acid molecule encoding the antigen-recognizing receptor of any one of claims 1-40.
57. The nucleic acid molecule of claim 56, wherein the nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, or SEQ ID NO: 27.
58. The nucleic acid molecule of claim 56 or 57, wherein the nucleic acid molecule comprises the nucleotide sequence set forth in SEQ ID NO: 22.
59. A vector comprising the nucleic acid molecule of any one of claims 56-58.
60. The vector of claim 59, wherein the vector is a γ -retroviral vector.
61. A host cell expressing the nucleic acid molecule of any one of claims 56-58.
62. The host cell of claim 61, wherein the host cell is a T cell.
63. A composition comprising the cell of any one of claims 41-55.
64. The composition of claim 63, which is a pharmaceutical composition further comprising a pharmaceutically acceptable carrier.
65. A method of reducing tumor burden in a subject, comprising administering an effective amount of the cell of any one of claims 41-55 or the composition of claim 63 or 64 to the subject.
66. The method of claim 65, wherein the method reduces the number of tumor cells, reduces tumor size, and/or eradicates the tumor in the subject

67. A method of increasing or lengthening survival of a subject having a tumor or neoplasm, comprising administering an effective amount of the cell of any one of claims 41-55 or the composition of claim 63 or 64.

68. A method of treating and/or preventing a tumor or neoplasm in a subject, comprising administering an effective amount of the cell of any one of claims 41-55 or the composition of claim 63 or 64.

69. The method of any one of claims 65-68, wherein the tumor or neoplasm is selected from the group consisting of acute myeloid leukemia (AML), multiple myeloma, Non-Hodgkin's Lymphoma, Hodgkin's Lymphoma, Chronic Lymphocytic Leukemia (CLL), glioblastoma, myelodysplastic syndrome (MDS), and chronic myelogenous leukemia (CML).

70. The method of any one of claims 65-69, wherein the tumor or neoplasm is acute myeloid leukemia (AML).

71. A method for producing a cell comprising an antigen-recognizing receptor of any one of claims 1-40, comprising introducing into the cell a nucleic acid molecule that encodes the antigen-recognizing receptor.

72. A kit for reducing tumor burden in a subject, treating and/or preventing a tumor or neoplasm in a subject, and/or increasing or lengthening survival of a subject having a tumor or neoplasm, comprising the cell of any one of claims 41-55.

73. The kit of claim 72, wherein the kit further comprises written instructions for using the cell for reducing tumor burden in a subject, treating and/or preventing a tumor or neoplasm in a subject, and/or increasing or lengthening survival of a subject having a tumor or neoplasm.

Phage-based CD371-targetd CAR s

Et.B10L3H_MT_h28Z (B10 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L3H, and a CD28Z-based signaling domain)

Et.B10L4H_MT_hBBZ (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a 4-1BBZ-based signaling domain)

Et.B10L4H_MT_h28Z (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.B10H3L_MT_h28Z (B10, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.B10H4L_MT_h28Z (B10, Heavy-Light scFv orientation with a 19-amino acid G4S linker, H4L, and a CD28Z-based signaling domain)

Et.C3H3L_MT_h28Z (C3, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.C3L3H_MT_h28Z (C3 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.D6L3H_MT_h28Z (D6 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)




FIG. 1

CAR Constructs with EGFR and MYC-tag




FIG. 2

CAR Constructs with EGFR and MYC-tag





FIG. 2, Continued



Effector:Tumor Ratio

Dunnnett's multiple comparisons test				Significant?		Summary		Adjusted P Value	
				Mean Diff.		95.00% CI of diff.			
1_1									
4dR Et.C1.H.vh28Zvs. 4dR Et.B.LH.MT.h28Z				14.75		2.322 to 27.18		Yes *	
4dR Et.C1.H.vh28Zvs. 4dR Et.B.LH.g4s.MT.h28Z				17.25		4.822 to 29.68		Yes **	
4dR Et.C1.H.vh28Zvs. 4dR Et.B.LH.MT.h28Z				3.750		-8.678 to 16.18		No ns	
4dR Et.C1.H.vh28Zvs. 4dR Et.C.LH.MT.h28Z				41.50		29.07 to 53.93		Yes ****	
4dR Et.C1.H.vh28Zvs. 4dR Et.C.LH.MT.h28Z				27.00		14.57 to 39.43		Yes ****	
4dR Et.C1.H.vh28Zvs. 4dR Et.D.LH.MT.h28Z				22.25		9.822 to 34.68		Yes ****	

FIG. 3




FIG. 4




FIG. 5

Donor C CAR T Cell Recursive Stimulation w/ Tumor at E:T of 1:12.5




FIG. 6

8/26




FIG. 7

9/26




FIG. 8A




FIG. 8B




FIG. 9A






FIG. 9B

Generation of human CD371 CRISPR Knockout Cell Lines



U937RFPcyp.371KO Killing at 24 hours (Donor A)




FIG. 11A

U937RFPcyp.371KO Killing at 24 hours (Donor B)




FIG. 11B




FIG. 12A




FIG. 12B

14/26

Interferon-gamma Secretion at 24 hours (Donor A)




FIG. 13A

Interferon-gamma Secretion at 24 hours (Donor B)




FIG. 13B

Interleukin-2 Secretion at 24 hours (Donor A)




FIG. 14A

Interleukin-2 Secretion at 24 hours (Donor B)




FIG. 14B

16/26

TNF-alpha Secretion at 24 hours (Donor A)




FIG. 15A

TNF-alpha Secretion at 24 hours (Donor B)




FIG. 15B

Donor A CAR T Cell Recursive Stimulation w/ Tumor at E:T of 1:5




FIG. 16A

Donor B CAR T Cell Recursive Stimulation w/ Tumor at E:T of 1:5




FIG. 16B

Xenograft Model of AML




FIG. 17




FIG. 18

5e4 U937gL (day 0) -> 1.25E6 CAR+ T cells (day 3)




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23

5e4 U937gL (day 0) -> 2e5 CAR+ T cells (day 3)




FIG. 24




FIG. 25

Phage-based CD371-targeted CARs

Et.B10L3H_MT_h28Z (B10 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L3H, and a CD28Z-based signaling domain)

Et.B10L4H_MT_hBBZ (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a 4-1BBZ-based signaling domain)

Et.B10L4H_MT_h28Z (B10 Light-Heavy scFv orientation with a 19-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.B10H3L_MT_h28Z (B10, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.B10H4L_MT_h28Z (B10, Heavy-Light scFv orientation with a 19-amino acid G4S linker, H4L, and a CD28Z-based signaling domain)

Et.C3H3L_MT_h28Z (C3, Heavy-Light scFv orientation with a 15-amino acid G4S linker, H3L, and a CD28Z-based signaling domain)

Et.C3L3H_MT_h28Z (C3 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)

Et.D6L3H_MT_h28Z (D6 Light-Heavy scFv orientation with a 15-amino acid G4S linker, L4H, and a CD28Z-based signaling domain)




FIG. 1