

(19) **DANMARK**

(10) **DK/EP 2056832 T3**



(12) **Oversættelse af  
europæisk patentskrift**


Patent- og  
Varemærkestyrelsen

- 
- (51) Int.Cl.: **A 61 K 31/519 (2006.01)** **A 61 K 9/20 (2006.01)** **A 61 K 9/48 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2017-05-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-03-22**
- (86) Europæisk ansøgning nr.: **07794121.9**
- (86) Europæisk indleveringsdag: **2007-08-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2009-05-13**
- (86) International ansøgning nr.: **SE2007000736**
- (87) Internationalt publikationsnr.: **WO2008024045**
- (30) Prioritet: **2006-08-21 US 823083 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **AstraZeneca AB, , 151 85 Södertälje, Sverige**
- (72) Opfinder: **BANKS, Simon, AstraZeneca R & D Charnwood, Bakewell Road, Loughborough Leicestershire LE11 5RH, Storbritannien**
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **SAMMENSÆTNINGER EGNET TIL ORAL ADMINISTRATION, DER OMFATTER ET TRIAZOL[4,5-D]PYRIMIDINDERIVAT**
- (56) Fremdragne publikationer:  
**WO-A1-00/34283**  
**WO-A1-01/36421**  
**WO-A1-01/92262**  
**WO-A1-2004/037263**  
**WO-A1-2004/052342**  
**WO-A2-2004/024127**  
**WO-A2-2005/113006**  
**US-A1- 2004 001 885**



## DESCRIPTION

[0001] The present invention relates to pharmaceutical compositions and more particularly to a pharmaceutical composition containing the compound of formula (I):



[0002] The compound of formula (I) is conventionally named {1*S*-[1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ (1*S*\*,2*R*\*)*S*]}-3-(7-[[2-(3,4-difluorophenyl)cyclopropyl]amino]-5-(propylthio)-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-3-yl)-5-(2-hydroxyethoxy)cyclopentane-1,2-diol and hereinafter will be referred to as the 'Agent'.

[0003] The Agent is disclosed as an ADP-receptor antagonist in International Patent Application number PCT/SE99/02256 (publication number WO00/34283) and International Patent Application number PCT/SE01/01239 (publication number WO01/92262). It has been found that adenosine 5'-diphosphate (ADP) acts as a key mediator of thrombosis. ADP-induced platelet aggregation is mediated by the P<sub>2</sub>T receptor subtype located on the platelet membrane. The P<sub>2</sub>T receptor (also known as P2Y<sub>ADP</sub> or P2T<sub>AC</sub>) is primarily involved in mediating platelet aggregation/activation and is a G-protein coupled receptor which is as yet uncloned. The pharmacological characteristics of this receptor have been described, for example, in the references by Humphries et al., Br. J. Pharmacology (1994), 113, 1057-1063, and Fagura et al., Br. J. Pharmacology (1998) 124, 157-164. It has been shown that antagonists at this receptor offer significant improvements over other anti-thrombotic agents (see J. Med. Chem. (1999) 42, 213).

[0004] The pharmaceutical compositions of the present invention are suitable for oral administration. One of the qualities that is desirable in a pharmaceutical composition suitable for oral administration is bioavailability. The bioavailability of a drug is the relative amount of an administered dose that reaches the systemic circulation in an unchanged form. Therefore bioavailability is important in determining the therapeutically active concentration at the site of action. Both drug release from the formulation and the stability of the formulation will affect its bioavailability. It is therefore important that the drug formulation should release substantially all of the drug. (see Aulton ME, *Pharmaceutics - The Science of Dosage Form Design*, 2nd Edition, 2002, Churchill Livingstone.). Bioavailability can be measured using tests known in the art, for example using a standard United States Pharmacopoeia (USP) dissolution apparatus and a standard 'bio-relevant' dissolution medium, for example FaSSIF (Pharm. Res., 17:439-444, 2000).

[0005] There are pharmaceutical compositions containing the Agent that retain some of the Agent and hence reduce its bioavailability.

[0006] We have now discovered a novel pharmaceutical composition of the Agent that has advantageous properties and which solves one or more of the problems associated with formulation of the Agent. In a first aspect we have discovered a pharmaceutical composition that is suitable for oral administration and that releases substantially all of the Agent. In one aspect the pharmaceutical composition releases at least 90% of the Agent. In another aspect the pharmaceutical composition releases at least 95% of the Agent. In yet another aspect the pharmaceutical composition releases at least 97% of the Agent.

[0007] Accordingly, the invention is a pharmaceutical composition comprising:

the Agent;

a filler which is a mixture of mannitol and dibasic calcium phosphate dihydrate,

a binder which is hydroxypropyl cellulose,

a disintegrant which is sodium starch glycolate,

and one or more lubricants.

**[0008]** A 'soluble' filler is a filler that is substantially soluble in water at ambient temperature. An 'insoluble' filler is a filler that has low or slow solubility in water at ambient temperature.

**[0009]** The pharmaceutical composition contains at least one 'soluble' filler which is mannitol.

**[0010]** The pharmaceutical composition contains one or more 'insoluble' filler which is dibasic calcium phosphate dihydrate.

**[0011]** In one aspect the pharmaceutical composition contains one or more 'soluble' fillers. In another aspect, the pharmaceutical composition contains one 'soluble' filler.

**[0012]** In one aspect the pharmaceutical composition contains one or more 'insoluble' fillers. In another aspect, the pharmaceutical composition contains one 'insoluble' filler.

**[0013]** In one aspect the pharmaceutical composition contains one or more binders. In another aspect, the pharmaceutical composition contains one binder.

**[0014]** In one aspect the pharmaceutical composition contains one or more disintegrants. In another aspect, the pharmaceutical composition contains one disintegrant.

**[0015]** In one aspect the pharmaceutical composition contains one or more lubricants. In another aspect, the pharmaceutical composition contains one lubricant.

**[0016]** In another aspect, the filler is a mixture of mannitol and dibasic calcium phosphate dihydrate.

**[0017]** The 'soluble' filler is mannitol.

**[0018]** The 'insoluble' filler is dibasic calcium phosphate dihydrate.

**[0019]** The binder is hydroxypropyl cellulose.

**[0020]** The disintegrant is sodium starch glycollate.

**[0021]** Additional conventional excipients, which may be added, include preservatives, stabilisers, anti-oxidants, silica flow conditioners, antiadherents or glidants.

**[0022]** Other suitable lubricants and additional excipients which may be used are described in Handbook of Pharmaceutical Excipients, 2nd Edition, American Pharmaceutical Association; The Theory and Practice of Industrial Pharmacy, 2nd Edition, Lachman, Leon, 1976; Pharmaceutical Dosage Forms: Tablets Volume 1, 2nd Edition, Lieberman, Hebert A., et al, 1989; Modern Pharmaceutics, Banker, Gilbert and Rhodes, Christopher T, 1979; and Remington's Pharmaceutical Sciences, 15th Edition, 1975.

**[0023]** Suitable lubricants include, for example, magnesium stearate, stearic acid, palmitic acid, calcium stearate, carnauba wax, hydrogenated vegetable oils, mineral oil, polyethylene glycols and sodium stearyl fumarate.

**[0024]** In one aspect, the lubricant is selected from magnesium stearate and sodium stearyl fumarate. In another aspect, the lubricant is magnesium stearate.

**[0025]** In one aspect, the pharmaceutical composition contains from 1 to 50% by weight of the Agent. In particular it contains 20 to 45% by weight of the Agent.

**[0026]** In another aspect, the pharmaceutical composition contains from 1 to 90% by weight of filler. In particular, it contains 20 to 70% by weight of filler.

**[0027]** In another aspect, the pharmaceutical composition contains from 1-70% by weight of 'soluble' filler. In particular, it contains 20 to 45% by weight of 'soluble' filler.

**[0028]** In another aspect, the pharmaceutical composition contains from 1 to 30% by weight of 'insoluble' filler. In particular it contains 10 to 30% by weight of 'insoluble' filler.

**[0029]** In another aspect, the pharmaceutical composition contains from 2 to 8% by weight of binder. In particular, it contains 3 to 6% by weight of binder.

**[0030]** In another aspect, the pharmaceutical composition contains from 2 to 6% by weight of disintegrant.

**[0031]** It will be appreciated that a particular excipient may act as both a binder and a filler, or as a binder, a filler and a disintegrant. Typically the combined amount of filler, binder and disintegrant comprises, for example, 50 to 90% by weight of the composition.

**[0032]** Typically one or more lubricants will be present in an amount 0.5 to 3%, and especially 0.5 to 1% by weight.

**[0033]** The invention relates to a pharmaceutical composition comprising the Agent, mannitol, dibasic calcium phosphate dihydrate, hydroxypropylcellulose, sodium starch glycolate and one or more lubricants.

**[0034]** In another aspect the invention relates to a pharmaceutical composition comprising: the Agent in an amount of 20 to 45% by weight; mannitol in an amount of 20 to 45% by weight; dibasic calcium phosphate dihydrate in an amount of 10 to 30% by weight; hydroxypropylcellulose in an amount of 3 to 6% by weight; sodium starch glycolate in an amount of 2 to 6% by weight; and one or more lubricants in an amount of 0.5 to 3% by weight.

**[0035]** It is desirable that the physical properties of these compositions are stable on storage, as changes in for instance, disintegration times, dissolution rates or tablet hardness among others can affect product performance. It is possible that decreases in dissolution rate on storage under International Council for Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use (ICH) stability testing conditions, used to assign product shelf life, can reduce the bioavailability of the Agent. Physical property stability can be measured by USP methodologies for disintegration times and dissolution testing.

**[0036]** It is desirable that the compositions are chemically stable as degradation by oxidation, hydrolysis, isomerisation, photolysis, polymerisation or any other method of degradation, either as a result of mixing with excipients or by any other method, could lead to a reduction in bioavailability. Chemical stability can be measured by a suitable, stability indicating chromatographic method for determining degradation products (see Aulton ME, *Pharmaceutics - The Science of Dosage Form Design*, 2nd Edition, 2002, Churchill Livingstone.).

**[0037]** In another aspect we have discovered a pharmaceutical composition that is suitable for oral administration that releases substantially all of the Agent and has a desirable stability profile.

**[0038]** In one aspect the invention relates to a pharmaceutical composition prepared by wet granulation.

**[0039]** Granulation is a process by which primary particles (powders) are made to adhere to form larger, multiparticulate entities called granules. Granulation normally commences after initial dry mixing of the powdered ingredients so that a fairly uniform distribution of ingredients through the mix is achieved. Granulation methods can be divided into two types, wet granulation methods that utilize a liquid to form the granules and dry methods that do not.

**[0040]** In dry granulation methods, primary powder particles are aggregated under pressure (or compaction). There are two main processes: a large tablet (also known as a slug) is produced with a heavy duty tablet press or the powder particles are compressed between two rollers to produce a sheet or 'ribbon' of material (process known as roller compaction). In both cases, the compacted material is milled using a suitable milling technique to produce granular material. The granules can then be compressed in a standard tablet press to produce tablets.

**[0041]** Wet granulation involves massing the primary powder particles using a granulating fluid. The fluid contains a solvent, which can be removed by drying, and is non-toxic. The granulating fluid can be used alone or more typically with a binding agent (binder) to ensure article adhesion in the dry state. Binding agents can be added to the system as a binder solution (as part of

the granulating fluid) or as dry material mixed with the primary powder particles. There are three main types of wet granulator, shear granulators (such as planetary mixers), high shear mixer granulators (such as Fielder or Diosna) and Fluid Bed Granulators (such as Aeromatic or Glatt).

**[0042]** In another aspect we have discovered a pharmaceutical composition prepared by a wet granulation process that is suitable for oral administration that releases substantially all of the Agent and a desirable stability profile.

**[0043]** In another aspect the invention relates to a pharmaceutical composition prepared by a wet granulation process comprising the Agent, mannitol, dibasic calcium phosphate dihydrate, hydroxypropylcellulose, sodium starch glycolate and one or more lubricants.

**[0044]** In another aspect the invention relates to a pharmaceutical composition prepared by high shear wet granulation.

**[0045]** High shear wet granulation is a process that involves intensive dry mixing of primary powders and subsequent addition of granulating fluid, which results in the formation of granules. The granulating fluid contains a volatile solvent (usually water) and may also include a binder; ensuring particle adhesion (binders may also be added dry as powders to the bulk of the formulation to be granulated). Granules possess major advantages compared to powders, which they are composed of, in terms of improved flow properties, reduced risk of segregation, increased homogeneity. (Information taken from Aulton ME, *Pharmaceutics - The Science of Dosage Form Design*, 2nd Edition, 2002, Churchill Livingstone.)

**[0046]** In one aspect the pharmaceutical composition is in a solid dosage form, such as a tablet or capsule. In another aspect the pharmaceutical composition is in the form of a tablet.

**[0047]** In another aspect the invention relates to a pharmaceutical composition prepared by a high shear wet granulation process comprising the Agent, mannitol, dibasic calcium phosphate dihydrate, hydroxypropylcellulose, sodium starch glycolate and one or more lubricants.

**[0048]** The Agent exists in an amorphous form and in four different substantially crystalline forms (see International Patent Application number PCT/SE01/01239 (publication number WO01/92262)). In another aspect the invention relates to a pharmaceutical composition as hereinabove define in which the Agent is in a crystalline form.

**[0049]** In yet another aspect the invention relates to a pharmaceutical composition comprising the Agent substantially as Polymorph II.

**[0050]** In yet another aspect the invention relates to a pharmaceutical composition comprising the Agent substantially as Polymorph III.

**[0051]** Compositions of the invention, which are of particular interest include, for example, the specific embodiments set out hereinafter in the accompanying Example.

**[0052]** It will be appreciated that modifications of the wet granulation techniques, including the order of addition of the components and their screening and blending prior to compression into tablets, may be carried out according to principles well known in the art.

**[0053]** A further aspect of the present invention comprises a method of preparing a pharmaceutical composition, which comprises admixing the Agent with:

a filler which is a mixture of mannitol and dibasic calcium phosphate dihydrate;

a binder which is hydroxyl propyl cellulose,

a disintegrant which is sodium starch glycolate,

and one or more lubricants.

## Example 1

[0054]

Ingredient	Quantity per unit dose Unit dose (mg)	Quantity (%w/w or w/v)
The Agent	90.00	30.00
Mannitol	126.00	42.00
Dibasic calcium phosphate dihydrate	63.00	21.00
Hydroxypropyl cellulose	9.00	3.00
Sodium starch glycolate	9.00	3.00
Magnesium stearate	3.00	1.00
Core tablet weight	300.000	100.00

[0055] A high shear wet granulator (Fielder GP1 with 10L bowl) was used to dry mix the Agent, mannitol, dibasic calcium phosphate dihydrate, hydroxypropyl cellulose and sodium starch glycolate in amounts to give 2.5 kg of total formulation, for 4 minutes. Water was added via a pressure pot at approximately 50g/min to approx. 25% w/w. The total mixing time was approximately 10 minutes.

[0056] The fluid bed was dried using a Glatt GPCG1 at 60°C to a product temperature of 42°C. The resulting granule was milled by Quadro Comil 197, the milled granules were blended with magnesium stearate and tablets were compressed from the blend.

## REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [WOSE9902256A](#) [0003]
- [WO0034283A](#) [0003]
- [WOSE0101239A](#) [0003] [0048]
- [WO0192262A](#) [0003] [0048]

### Non-patent literature cited in the description

- **HUMPHRIES et al.** Br. J. Pharmacology, 1994, vol. 113, 1057-1063 [0003]
- **FAGURA et al.** Br. J. Pharmacology, 1998, vol. 124, 157-164 [0003]
- J. Med. Chem., 1999, vol. 42, 213- [0003]
- **AULTON ME** Pharmaceuticals - The Science of Dosage Form Design Churchill Livingstone 20020000 [0004] [0036] [0045]
- Pharm. Res., 2000, vol. 17, 439-444 [0004]
- Handbook of Pharmaceutical Excipients American Pharmaceutical Association [0022]
- **LEON** The Theory and Practice of Industrial Pharmacy Lachman 19760000 [0022]
- **HEBERT A. et al.** Pharmaceutical Dosage Forms Lieberman 19890000 vol. 1, [0022]
- **BANKER, GILBERT RHODES, CHRISTOPHER T** Modern Pharmaceutics, 1979, [0022]
- Remington's Pharmaceutical Sciences 19750000 [0022]

## Patentkrav

1. Farmaceutisk sammensætning, der omfatter:  
{1S-[1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ (1S\*,2R\*),5 $\beta$ ]}-3-(7-{[2-(3,4-  
5 difluorphenyl)cyclopropyl]-amino}-5-(propylthio)-3H-1,2,3-  
triazol[4,5-d]pyrimidin-3-yl)-5-(2-hydroxyethoxy)-cyclopentan-  
1,2-diol;  
et fyldstof, som er en blanding af mannitol og dibasisk  
calciumphosphatdihydrat;  
10 et bindemiddel, som er hydroxypropylcellulose;  
et opløsningsmiddel, som er natriumstivelsesglycolat; og  
et eller flere smøremidler.
  
2. Farmaceutisk sammensætning ifølge krav 1, hvor  
15 smøremidlet er udvalgt blandt magnesiumstearat og  
natriumstearylfumarat.
  
3. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor  
{1S-[1 $\alpha$ ,  
2 $\alpha$ ,3 $\beta$ (1S\*,2R\*),5 $\beta$ ]}-3-(7-{[2-(3,4-  
20 difluorphenyl)cyclopropyl]-amino}-5-(propylthio)-3H-1,2,3-  
triazol[4,5-d]pyrimidin-3-yl)-5-(2-hydroxyethoxy)-cyclopentan-  
1,2-diol er til stede i en mængde på 20 til 45 vægt-%.
  
4. Farmaceutisk sammensætning ifølge et hvilket som helst af  
25 kravene 1 til 3, hvor fyldstoffet er til stede i en mængde på  
20 til 70 vægt-%.
  
5. Farmaceutisk sammensætning ifølge et hvilket som helst af  
kravene 1 til 4, hvor bindemidlet er til stede i en mængde på  
30 3 til 6 vægt-%.
  
6. Farmaceutisk sammensætning ifølge et hvilket som helst af  
kravene 1 til 5, hvor opløsningsmidlet er til stede i en  
mængde på 2 til 6 vægt-%.  
35
  
7. Farmaceutisk sammensætning ifølge et hvilket som helst af  
kravene 1 til 6, hvor smøremidlet er til stede i en mængde på  
0,5 til 1 vægt-%.

8. Farmaceutisk sammensætning ifølge krav 1, der omfatter:  
{1S-[1 $\alpha$ ,2 $\alpha$ ,3 $\beta$ (1S\*,2R\*),5 $\beta$ ]}-3-(7-{[2-(3,4-difluorphenyl)cyclopropyl]amino}-5-(propylthio)-3H-1,2,3-triazol[4,5-d]pyrimidin-3-yl)-5-(2-hydroxyethoxy)-cyclopentan-1,2-diol i en mængde på 20 til 45 vægt-%;  
mannitol i en mængde på 20 til 45 vægt-%;  
dibasisk calciumphosphatdihydrat i en mængde på 10 til 30 vægt-%;
- 10 hydroxypropylcellulose i en mængde på 3 til 6 vægt-%;  
natriumstivelsesglycolat i en mængde på 2 til 6 vægt-%; og  
et eller flere smøremidler i en mængde på 0,5 til 3 vægt-%.
9. Farmaceutisk sammensætning ifølge krav 1 eller 2, hvor  
15 smøremidlet er magnesiumstearat.