

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3668972 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 12 N 9/22 (2006.01)** **A 61 K 38/46 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2025-01-13**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-10-09**
- (86) Europæisk ansøgning nr.: **18846060.4**
- (86) Europæisk indleveringsdag: **2018-08-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-06-24**
- (86) International ansøgning nr.: **US2018047084**
- (87) Internationalt publikationsnr.: **WO2019036719**
- (30) Prioritet: **2017-08-18 US 201762547220 P** **2017-12-28 US 201762611166 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Neutrolis, Inc., 1 Broadway, Cambridge, MA 02142, USA**
- (72) Opfinder: **FUCHS, Tobias A., c/o Neutrolis Inc., 700 Main Street, North, Cambridge, MA 02139, USA**
GÖBEL, Josephine (formerly BITTERLING, Josephine), c/o Neutrolis Inc., 700 Main Street, North, Cambridge, MA 02139, USA
ENGLERT, Hanna, c/o Neutrolis Inc., 700 Main Street, North, Cambridge, MA 02139, USA
JIMÉNEZ-ALCÁZAR, Miguel, c/o Neutrolis Inc., 700 Main Street, North, Cambridge, MA 02139, USA
- (74) Fuldmægtig i Danmark: **Cosmovici Intellectual Property, Paul Cosmovici, Rue du Commerce 4, 1204 Geneva, Schweiz**
- (54) Benævnelse: **KONSTRUERED DNASE-ENZYMER OG ANVENDELSE I TERAPI**
- (56) Fremdragne publikationer:
WO-A2-2011/053982
US-A1- 2009 010 966
US-A1- 2016 376 366
ANDREW WILBER ET AL: "Deoxyribonuclease 1-like III is an inducible macrophage barrier to liposomal transfection", MOLECULAR THERAPY, NO LONGER PUBLISHED BY ELSEVIER, vol. 6, no. 1, 1 July 2002 (2002-07-01), pages 35 - 42, XP002641392, ISSN: 1525-0016, DOI: 10.1006/MTHE.2002.0625
JIMÉNEZ-ALCÁZAR M ET AL: "Impaired DNase1-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies", JOURNAL OF THROMBOSIS AND HAEMOSTASIS, vol. 13, no. 5, 13 April 2015 (2015-04-13), pages 732 - 742, XP055782172, Retrieved from the Internet <URL:http://www.ncbi.nlm.nih.gov/pubmed/25418346> DOI: 10.1111/jth.12796
SISIRAK VANJA ET AL: "Digestion of Chromatin in Apoptotic Cell Microparticles Prevents Autoimmunity", CELL, ELSEVIER, AMSTERDAM NL, vol. 166, no. 1, 9 June 2016 (2016-06-09), pages 88 - 101, XP029627879, ISSN: 0092-8674, DOI: 10.1016/J.CELL.2016.05.034
SHIOKAWA DAISUKE ET AL: "Identification of two functional nuclear localization signals in DNase [gamma]

Fortsættes ...

and their roles in its apoptotic DNase activity", *BIOCHEMICAL JOURNAL*, vol. 376, no. 2, 1 December 2003 (2003-12-01), pages 377 - 381, XP055782937, ISSN: 0264-6021, Retrieved from the Internet <URL:<http://portlandpress.com/biochemj/article-pdf/376/2/377/712183/bj3760377.pdf>> DOI: 10.1042/bj20030820

JIMÉNEZ-ALCÁZAR MIGUEL ET AL: "Host DNases prevent vascular occlusion by neutrophil extracellular traps", *SCIENCE*, vol. 358, no. 6367, 1 December 2017 (2017-12-01), US, pages 1202 - 1206, XP055783049, ISSN: 0036-8075, DOI: 10.1126/science.aam8897

MONA SAFFARZADEH ET AL: "Fighting against the dark side of neutrophil extracellular traps in disease : manoeuvres for host protection", *CURRENT OPINION IN HEMATOLOGY*, vol. 20, no. 1, 1 January 2013 (2013-01-01), US, pages 3 - 9, XP055471240, ISSN: 1065-6251, DOI: 10.1097/MOH.0b013e32835a0025

JORCH SELINA K ET AL: "An emerging role for neutrophil extracellular traps in noninfectious disease", *NATURE MEDICINE*, vol. 23, no. 3, 1 March 2017 (2017-03-01), New York, pages 279 - 287, XP055785209, ISSN: 1078-8956, Retrieved from the Internet <URL:<http://www.nature.com/articles/nm.4294>> DOI: 10.1038/nm.4294

DABROWSKA D ET AL: "New Aspects of the Biology of Neutrophil Extracellular Traps.", *SCANDINAVIAN JOURNAL OF IMMUNOLOGY* DEC 2016, vol. 84, no. 6, December 2016 (2016-12-01), pages 317 - 322, ISSN: 1365-3083

BRINKMANN VOLKER ET AL: "Neutrophil extracellular traps kill bacteria", *SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE*, US, vol. 303, no. 5663, 5 March 2004 (2004-03-05), pages 1532 - 1535, XP002469080, ISSN: 0036-8075, DOI: 10.1126/SCIENCE.1092385

KOBAYASHI, Y ET AL.: "Synchronous growth of *Pichia Pastoris* for a high rate production of DnaseI at microquantities", *THE 10TH APCCHE CONGRESS THE ASIAN PACIFIC CONFEDERATION OF CHEMICAL ENGINEERING : CONFERENCE PROCEEDINGS : OCTOBER 17-21, 2004 KITAKYUSHU, JAPAN*, no. 833, 1 January 2004 (2004-01-01), pages 1 - 6, XP055679026, DOI: 10.11491/apcche.2004.0.953.0

HAKKIM, ABDUL ET AL.: "Impairment of neutrophil extracellular trap degradation is associated with lupus nephritis", *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES*, vol. 107, no. 21, May 2010 (2010-05-01), pages 9813 - 9818, XP0002592735, DOI: 10.1073/pnas.0909927107

PICCOLO, F ET AL.: "Intrapleural Tissue Plasminogen Activator and Deoxyribonuclease for Pleural Infection. An Effective and Safe Alternative to Surgery", *ANNALS ATS*, vol. 11, no. 9, November 2014 (2014-11-01), pages 1419 - 1425, XP055678828, DOI: 10.1513/AnnalsATS.201407-329OC

PERINI ET AL: "Topical application of Acheflan on rat skin injury accelerates wound healing: a histopathological, immunohistochemical and biochemical study", *BMC COMPLEMENTARY MEDICINE AND THERAPIES*, vol. 15, 203, 30 June 2015 (2015-06-30), XP055678836, DOI: 10.1186/s12906-015-0745x

ANDREW WILBUR ET AL: "Deoxyribonuclease I-like III Is an Inducible Macrophage Barrier to Liposomal Transfection", *MOLECULAR THERAPY*, vol. 6, no. 1, 2002, pages 35 - 42, XP002641392, DOI: 10.1006/mthe.2002.0625

BERNTSSON RP. ET AL.: "Structural insight into DNA binding and oligomerization of the multifunctional Cox protein of bacteriophage P2.", *NUCLEIC ACIDS RESEARCH*, vol. 42, no. 4, 23 November 2013 (2013-11-23), pages 2725 - 2735, XP055678852, DOI: 10.1093/nar/gkt1119

JIMÉNEZ-ALCÁZAR M ET AL: "Impaired DNase1-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies", *JOURNAL OF THROMBOSIS AND HAEMOSTASIS*, vol. 13, no. 5, 13 April 2015 (2015-04-13), pages 732 - 742, XP055782172, Retrieved from the Internet <URL:<http://www.ncbi.nlm.nih.gov/pubmed/25418346>> DOI: 10.1111/jth.12796

JIMÉNEZ-ALCÁZAR MIGUEL ET AL: "Host DNases prevent vascular occlusion by neutrophil extracellular traps", *SCIENCE*, vol. 358, no. 6367, 1 December 2017 (2017-12-01), US, pages 1202 - 1206, XP055783049, ISSN: 0036-8075, DOI: 10.1126/science.aam8897

DESCRIPTION

Description

PRIORITY

[0001] This Application claims the benefit of, and priority to, US Application Nos. 62/611,166 filed December 28, 2017 and 62/547,220 filed August 18, 2017.

BACKGROUND OF THE INVENTION

[0002] Inflammation is an essential host response to control invading microbes and heal damaged tissues. Uncontrolled and persistent inflammation causes tissue injury in a plethora of inflammatory disorders. Neutrophils are the predominant leukocytes in acute inflammation. During infections neutrophils generate neutrophil extracellular traps (NETs), lattices of DNA-filaments decorated with toxic histones and enzymes that immobilize and neutralize bacteria. The relevance of NETs in host defense is illustrated by the fact that extracellular DNases serve as virulence factors in several pathogenic bacteria. However, inappropriately released NETs may harm host cells due to their cytotoxic, proinflammatory, and prothrombotic activity. Indeed, NETs are frequently associated with inflammatory or ischemic organ damage and the therapeutic infusion of DNases limits host injury in various animal models.

[0003] Extracellular trap (ET) formation is not limited to neutrophils as other leukocytes, namely monocytes, macrophages, basophils, eosinophils, and mast cells release ETs as well. Furthermore, cancer cells, including acute promyelocytic leukemia (APL) cells, and injured endothelial cells can expose DNA-filaments, which have ET-like features.

[0004] How the host degrades NETs in vivo to limit tissue damage during episodes of inflammation is poorly understood. DNase1 (D1) in serum has been shown to digest the DNA-backbone of NETs in vitro. Other extracellular DNases have been identified, including DNase1-like 3 (D1L3) in circulation, which can digest extracellular microparticle-associated chromatin. Moreover, it was shown that impaired DNase1 mediated degradation of NETs (Jiménez-Alcázar et al. (2015), Journal of Thrombosis and Haemostasis, vol. 13(5):732-742.

[0005] D1 and D1L3 form along with DNase1-like 1 (D1L1) and DNase1-like 2 (D1L2) the DNase1-protein family, a group of homologous secreted DNase enzymes that are expressed in humans and provide drug candidates for NET-associated diseases. However, the physical, enzymatic, and pharmacokinetic properties of these enzymes are not ideal for therapy.

[0006] The present disclosure provides engineered DNases, including DNase1-like 3 and DNase1, for therapy, including for preventing or treating vascular occlusions that can result from release of NETs during acute inflammatory episodes.

SUMMARY OF THE INVENTION

[0007] The invention is defined by the appended claims.

The present disclosure provides engineered DNase proteins (including DNase1-like 3 and DNase1) that are useful for treating conditions characterized by neutrophil extracellular trap (NET) accumulation and/or release. The disclosure provides compositions and methods for preventing or treating vascular occlusion involving NETs. As demonstrated herein, NETs participate in a non-canonical mechanism for vascular occlusion, which is not dependent on fibrin and platelets.

[0008] The disclosure provides DNase1-like 3 (D1L3) variants engineered to have physical, pharmacodynamic, and/or enzymatic properties more suitable for therapy, for example, to reduce or prevent NET accumulation in a subject. The D1L3 enzyme may have advantages in manufacturing, providing for production of the recombinant enzyme suitable for use in therapy. The disclosure provides a recombinant D1L3 variant comprising one or more amino acid alterations resulting in one or more glycosylations, inactivation of a nuclear localization signal, and/or deletion of all or part of a C-terminal tail.

[0009] The D1L3 protein variant may have increased protein stability, increased resistance towards inhibition by proteases, increased bioavailability, and substantially the same or better DNA and/or chromatin and/or NET-degrading activity (in vitro or in vivo) as compared to wild-type D1L3 protein. The D1L3 variant may comprise one or more of the following properties relative to wild type D1L3 (e.g., SEQ ID NO:2): the same or substantially the same (or higher) protein-free DNA (naked DNA) degradation activity, the same or substantially the same (or higher) chromosomal DNA (chromatin) degradation activity, protease resistance, an increased circulatory half-life, and higher production levels with in vitro expression systems (e.g. Chinese hamster ovary cells and/or *Pichia pastoris*).

[0010] The D1L3 variant may include one or a plurality of block substitutions from human DNase1 (SEQ ID NO: 1), described herein as building block substitutions.

[0011] The disclosure provides DNase1 (D1) variants engineered to have physical, pharmacodynamic, and/or enzymatic properties more suitable for therapy, for example, to reduce or prevent NET accumulation in a subject. The D1 enzyme may have advantages in manufacturing, providing for production of the recombinant enzyme suitable for use in therapy. The engineered D1 variant may comprise an amino acid sequence that is at least 80% identical to the mature enzyme of amino acid sequence of SEQ ID NO: 1, with one or more amino acid substitutions, additions (including insertions), or deletions resulting in one or more

of a mutated DNA binding site, a mutated chromatin binding site, a mutated actin binding site, a mutated glycosylation site, addition of a nuclear localization signal (e.g., having similarity or identity to NLS1 or NLS2 of D1L3), and/or a C-terminal domain similar to or identical to the C-terminal tail of D1L3.

[0012] The D1 variant may be engineered to comprise one or more of the following characteristics relative to the wild type enzyme: the same or higher protein-free DNA (naked DNA) degradation activity, higher chromosomal DNA (chromatin) degradation activity, similar or improved protease resistance, actin resistance, similar or improved penetration from blood into urine and/or bile, and higher production levels in vitro expression systems (e.g. Chinese hamster ovary cells and/or *Pichia pastoris*).

[0013] D1 variants described herein may have a combination of point mutations, including addition of cationic residues, and/or may comprise one or more block substitutions from D1L3 (SEQ ID NO:2). Such block substitutions are described herein, and termed "building block substitutions."

[0014] The disclosure provides pharmaceutical compositions comprising the D1L3 and/or D1 variants, or polynucleotides or vectors encoding the same, and a pharmaceutically acceptable carrier. The composition may be formulated for parenteral or pulmonary administration. The composition may be formulated for intravenous, intraarterial, intraperitoneal, intraarticular, intramuscular, topical, or subcutaneous administration. The composition may comprise both D1L3 and D1, which are each optionally engineered variants described herein. The composition may comprise a chimeric protein containing block sequences from both D1 and D1L3. Exemplary block substitutions between D1 and D1L3 are shown in FIG 31.

[0015] The disclosure provides a process for DNase enzyme engineering, by producing chimeric sequences. For example, in these aspects the method comprises providing a protein-protein alignment of donor and recipient DNase enzymes; identifying variable amino acid sequences for transfer ("building block"). The variable amino acid or amino acids are flanked by one or more conserved amino acids in the donor and recipient DNase enzymes (upstream and downstream of the building block). These building blocks can be swapped between recipient and donor proteins. The chimeric enzyme is then produced recombinantly.

[0016] The disclosure provides a method of making a pharmaceutical composition for reducing or preventing neutrophil extracellular trap (NET) accumulation. The method employs a genetically modified mouse deficient in D1 and D1L3 activity, and heterologous expression of G-CSF polynucleotide (e.g., in hepatocyte cells) or inducing a sustained endogenous G-CSF expression (e.g. via repetitive administration of microbial compounds). This mouse model accumulates NETs and rapidly develops NET-related vascular occlusions. The method thus comprises administering a candidate NET inhibitor or candidate DNase enzyme (including a D1L3 or D1 variant in accordance with this disclosure) to the genetically-modified mouse, and selecting a NET inhibitor or DNase enzyme that reduces the accumulation of NETs. The selected inhibitor or enzyme is formulated for administration to a human patient.

[0017] The disclosure provides a method for treating a subject in need of neutrophil extracellular trap (NET) degradation. The method comprises administering a therapeutically effective amount of a D1L3 and/or D1, and/or variants thereof according to this disclosure. The D1L3 or D1 variant may be administered as pharmaceutical compositions comprising the recombinant protein, or the encoding DNA or RNA, or vectors comprising the same.

[0018] The disclosed method pertains to the treatment of diseases or conditions characterized by the presence or accumulation of NETs. Such diseases or conditions include, but are not limited to, diseases associated with chronic neutrophilia (e.g., an increase in the number of neutrophils), neutrophil aggregation and leukostasis, thrombosis and vascular occlusion (e.g. sickle cell disease), ischemia-reperfusion injury (e.g. midgut volvulus, testicular torsion, limb ischemia reperfusion, vital organ ischemia-reperfusion, organ transplantation), surgical and traumatic tissue injury, an acute or chronic inflammatory reaction or disease, an autoimmune disease (e.g. systemic lupus erythematosus (SLE), lupus nephritis, rheumatoid arthritis, vasculitis, systemic sclerosis), cardiovascular disease (e.g., myocardial infarction, stroke, atherosclerosis, venous thromboembolism, including thrombolytic therapy), metabolic disease (e.g., diabetes), systemic inflammation (e.g., systemic inflammatory response syndrome (SIRS), sepsis, septic shock, disseminated intravascular coagulation (DIC), and thrombotic microangiopathy (TMA)), inflammatory diseases of the respiratory tract (e.g. cystic fibrosis, chronic obstructive pulmonary disease (COPD), acute lung injury (ALI), smoke induced lung injury, transfusion induced lung injury (TRALI), acute respiratory distress syndrome (ARDS), and asthma, atelectasis, bronchitis, empyema), renal inflammatory diseases (acute and chronic kidney diseases, including acute kidney injury (AKI) and chronic kidney disease (CKD), inflammatory diseases related to transplanted tissue (e.g. graft-versus-host disease) and cancer (e.g. leukemia, tumor metastasis, and solid tumors).

[0019] The subject may have or may be at risk of a ductal occlusion in a ductal system. Nonlimiting examples of a ductal system or an organ or tissue containing a ductal system include bile duct, tear duct, lactiferous duct, cystic duct, hepatic duct, ejaculatory duct, parotid duct, submandibular duct, major sublingual duct, bartholin's duct, cerebral aqueduct, pancreas, mammary gland, vas deferens, ureter, urinary bladder, gallbladder, and liver. As such, the disclosed enzymes are useful for treating a subject who has pancreatitis, cholangitis, conjunctivitis, mastitis, dry eye disease, an obstruction of the vas deferens, or renal disease.

[0020] The subject may have or may be at risk of NETs accumulating on endothelial surfaces (e.g., surgical adhesions), the skin (e.g., wounds/scarring, ulcers), or in synovial joints (e.g., gout, arthritis). For instance, NETs may contribute to surgical adhesions, e.g., after an invasive medical procedure. The disclosed enzymes can be administered during surgery to prevent or inhibit the formation of surgical adhesions. In some instances, D1, D1L3 (including variants thereof), or the combination of D1 and D1L3 (or variants thereof) as outlined herein can be administered topically to the skin to prevent or treat wounds and/or scarring. In other instances, D1, D1L3, or variants, or the combinations thereof as outlined herein can be administered to synovial joints to prevent or treat gout and arthritis.

[0021] The subject may have or may be at risk of a vascular occlusion comprising NETs.

[0022] The present method pertains to the treatment of diseases that are treatable with DNase enzyme, including D1 and streptodornase. Such diseases or conditions include thrombosis, stroke, sepsis, lung injury, atherosclerosis, viral infection, sickle cell disease, myocardial infarction, ear infection, wound healing, liver injury, endocarditis, liver infection, pancreatitis, primary graft dysfunction, limb ischemia reperfusion, kidney injury, blood clotting, alum-induced inflammation, hepatorenal injury, pleural exudations, hemotorax, intrabiliary blood clots, post pneumatic anemia, ulcers, otolaryngological conditions, oral infections, minor injuries, sinusitis, post-operative rhinoplasties, infertility, bladder catheter, wound cleaning, skin reaction test, pneumococcal meningitis, gout, leg ulcers, cystic fibrosis, Kartegener's syndrome, asthma, lobar atelectasis, chronic bronchitis, bronchiectasis, lupus, primary ciliary dyskinesia, bronchiolitis, empyema, pleural infections, cancer, dry eyes disease, lower respiratory tract infections, chronic hematomas, Alzheimer's disease, and obstructive pulmonary disease.

[0023] The disclosure further provides a method for recombinant production of D1 or D1L3, or variants thereof, using a non-mammalian expression system, such as *Pichia pastoris*. The *Pichia pastoris* may encode the DNase enzyme with the native signal peptide allowing for secretion from host cells.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024]

FIG. 1A - FIG. 1F illustrate that DNase1 and DNase1I3 in circulation degrade NETs in vitro, and characterize DNA-degrading activity of sera from WT, DNase1^{-/-} (D1^{-/-}), DNase1I3^{-/-} (D1L3^{-/-}), DNase1/DNase1I3^{-/-} (D1/D1L3^{-/-}) mice. FIG. 1A provides detection of DNase1 (D1), DNase1I3 (D1L3), and total DNase activity by the zymographic assays DPZ and SRED. FIG. 1B provide images and FIG. 1C provide quantification of DNA-stainings of NETs generated in vitro after incubation with sera from indicated genotypes (N=6). Scale bar: 50 μ m. FIG. 1C shows DPZ and SRED analysis of sera from D1/D1L3^{-/-} mice stably expressing a plasmid with DNase1 (D1), DNase1I3 (D1L3), or a control plasmid (Ctrl). FIG. 1E shows images and FIG. 1F shows quantification of DNA-staining of NETs generated in vitro after incubation with buffer or sera from D1/D1L3^{-/-} mice expressing D1, D1L3, or Ctrl (N=5). Scale bar: 50 μ m. Images are representative of two or more independent experiments. Statistics: (FIG. 1C and FIG. 1F) one-way ANOVA followed by Bonferroni's multiple comparisons post test; § P < 0.001 vs. all other groups.

FIG. 2A-FIG. 2I illustrate that DNase1 or DNase1I3 is required to tolerate chronic neutrophilia. Chronic neutrophilia was induced by injection of a G-CSF-expression plasmid (CSF3). Controls received an empty plasmid (Ctrl). FIG. 2A shows blood neutrophil count of WT mice and controls after 2 weeks (N=3-6). FIG. 2B shows NET-like structures (arrow) in DNA-stainings of blood smears from WT mice after 1 week (N=3). Scale bar: 20 μ m. FIG. 2C shows survival of

WT (N=7), DNase1^{-/-} (D1^{-/-}, N=6), DNase1I3^{-/-} (D1L3^{-/-}, N=6), DNase1/DNase1I3^{-/-} mice (D1/D1L3^{-/-}, N=6) and controls (Ctrl, N=4). FIG. 2D shows survival of D1/D1L3^{-/-} mice co-expressing CSF3 with DNase1 (CSF3/D1, N=5), DNase1I3 (CSF3/D1L3, N=6), and controls (CSF3/Ctrl, N=4). FIGS. 2E-2I characterizes mortality during chronic neutrophilia (N= 4). FIG. 2E shows change in peripheral body temperature. FIG. 2F provides photographs of plasma and urine. FIG. 2G provides concentration of hemoglobin in blood. FIG. 2H provides images of blood smears. Arrows point to schistocytes. Scale bar: 20 μ m. FIG. 2I shows LDH concentration in plasma. UT: untreated mice (N=5-6). Images are representative of three or more mice. Statistics: (FIGS. 2A, 2H, 2I) one-way and (FIG. 2E) two-way ANOVA followed by Bonferroni's multiple comparisons post test, (FIG. 2B) student's t-test, (FIGS. 2C, 2D) Log-rank test; # P < 0.01, § P < 0.001 vs. all other groups or baseline (BL).

FIG. 3A-FIG. 3G illustrate that DNase1 and DNase1I3 prevent vascular occlusion by NET-clots during chronic neutrophilia. Histological analysis of DNase1/DNase1I3^{-/-} mice (D1/D1L3^{-/-}) co-expressing CSF3 with DNase1 (CSF3/D1, N=4), DNase1I3 (CSF3/D1L3, N=4), or a control plasmid (CSF3/Ctrl, N=4). FIG. 3A provides hematoxylin and eosin stainings (H&E) of lungs. Scale bars: 500 μ m (overview), 25 μ m (detail). FIG. 3B shows quantification of blood vessels in lungs occluded by hematoxylin-positive clots per field of view (FOV). Untreated WT mice (UTWT, N=4), untreated D1/D1L3^{-/-} mice (UT; N=4). FIG. 3C shows immunostaining of occluded blood vessels for the neutrophil-marker myeloperoxidase (MPO) and chromatin. FIGS. 3D and 3E show immunostaining for von Willebrand factor (vWF), fibrin, and DNA. Three types of NET-clots were detected (α , vWF+/fibrin-; β , vWF+/fibrin+; γ , vWF-/fibrin-). FIG. 3F provides quantification of vWF and fibrin in NET-clots. Data shown as mean $\pm$ SD, n = 108. FIG. 3G shows survival of D1/D1L3^{-/-} mice expressing CSF3 treated with IgG (N=4), anti-platelet-IgG (N=5), and dabigatran (N=5). Scale bars: 50 μ m. Images are representative of three or more mice. (FIGS. 3C, 3D, 3E) dotted line indicates vessel wall. Statistics: (FIG. 3B) one-way ANOVA followed by Bonferroni's multiple comparisons post test; § P < 0.001 vs. all other groups, (FIG. 3G) Log-rank test; P > 0.05 Anti-Platelet-IgG or Dabigatran vs. IgG.

FIG. 4A-FIG. 4I illustrate that DNase1 and DNase1I3 protect against host injury in septicemia. WT mice (N=5) and mice with a combined deficiency in DNase1 and DNase1I3 (D1/D1L3^{-/-}) expressing DNase1 (D1, N=7), DNase1I3 (D1L3, N=8), or a control plasmid (Ctrl, N=11) were treated with lipopolysaccharide and heat-killed *E. coli* to induce septicemia. FIG. 4A shows survival time of septic mice. FIG. 4B shows concentration of hemoglobin in blood. FIG. 4C shows photographs of plasma and urine. FIG. 4D provides LDH concentration in plasma. FIG. 4E shows quantification of schistocytes in blood per field of view (FOV). FIG. 4F shows quantification of occluded blood vessels in lungs per FOV. FIG. 4G shows hematoxylin and eosin stainings (H&E) of lungs. Arrowheads point to occluded blood vessels. Scale bars: 500 μ m. FIGS. 4H and 4I show H&E staining of partially and fully occluded blood vessel. Arrows point to NETs covering the intercellular space. Inserts are overviews. Scale bars: 50 μ m. Statistics: (FIG. 4A) Log-rank test; # P < 0.01 vs. all other groups, (FIGS. 4B-4F) one-way ANOVA followed by Bonferroni's multiple comparisons post test; § P < 0.001, # P < 0.01.

FIG. 5A-FIG. 5E show a murine model of chronic neutrophilia. Four-week old WT mice were injected with a CSF3-expression plasmid (CSF3) or with an empty control plasmid (Ctrl). FIG.

5A shows G-CSF levels in plasma (N=3-6). BL: baseline, before injection; Ctrl: mice expressing the control plasmid without CSF3 for 2 weeks (N=4-5). FIG. 5B shows immunostaining for neutrophil elastase of lungs, kidneys, and livers. Scale bars: 200 μ m. FIG. 5C shows photographs and weights of spleens (N=5-6). Scale bar: 1 cm. FIG. 5D shows body weight of mice at indicated times (N=3-5). FIG. 5E shows neutrophil counts in whole blood and plasma levels of ALT, AST, BUN, and creatinine 2 weeks after the injection. Images are representative of three or more mice. Statistics: (FIGS. 5A, 5C) one-way and (FIG. 5D) two-way ANOVA followed by Bonferroni's multiple comparisons post test, and (FIG. 5E) Student's t-test; # $P < 0.01$, § $P < 0.001$ vs. (FIG. 5A) BL, or (FIG. 5C) UT.

FIG. 6A-FIG. 6D show that DNase1 and DNase1I3 prevent the formation of NET-clots and organ injury in liver and kidneys during chronic neutrophilia. Analysis of DNase1/DNase1I3-/- mice (D1/D1L3-/-) co-expressing CSF3 a control plasmid (CSF3/Ctrl, N=4), with DNase1 (CSF3/D1, N=4), or with DNase1I3 (CSF3/D1L3, N=4). Controls include untreated WT mice (UTWT, N=4) and untreated D1/D1L3-/- mice (UT; N=4). FIG. 6A shows plasma levels of alanine and aspartate aminotransferases (ALT, AST). FIG. 6B shows plasma levels of blood urea nitrogen (BUN) and creatinine. FIG. 6C provides quantification of intravascular hematoxylin-rich clots in livers per section and in kidneys per field of view (FOV). FIG. 6D provides H&E stainings of livers and kidneys. Scale bars: 500 μ m (overviews), 50 μ m (detail). Images are representative of three or more mice. Statistics: (FIGS. 6A, 6B, 6C) one-way ANOVA followed by Bonferroni's multiple comparisons post test; # $P < 0.01$, § $P < 0.001$ vs. all other groups.

FIG. 7A-FIG. 7C show hematoxylin-positive clots are composed of NETs. Analysis of lungs from DNase1/DNase1I3-/- mice (D1/D1L3-/-) with chronic neutrophilia (D1/D1L3-/- + CSF3/Ctrl) shown in FIGS. 3A-3G. FIG. 7A provides staining with hematoxylin and eosin (H&E), for DNA, and for chromatin of consecutive lung sections. Scale bars: 200 μ m. FIG. 7B provides immunostaining of occluded blood vessel for chromatin, CRAMP (cathelin-related antimicrobial peptide), and DNA. Scale bars: 50 μ m. FIG. 7C provides immunostaining of occluded blood vessel for chromatin, citrullinated histone 3 (citH3), and DNA. Scale bars: 50 μ m. Images are representative of three or more mice.

FIG. 8A and **FIG. 8B** show that human NETs form clots in vitro. FIG. 8A provides a photograph of NET-clot (arrow) generated by PMA-activated human neutrophils in vitro. Untreated neutrophils (UT) or neutrophils activated in the presence of recombinant human DNase1 (PMA + D1) served as controls. Scale bars: 5 mm. FIG. 8B provides hematoxylin and eosin stainings (H&E) and immunostaining for chromatin and the neutrophil-marker myeloperoxidase (MPO) of in vitro generated NET-clots. Scale bars: 200 μ m. Images are representative of three or more independent experiments.

FIG. 9A and **FIG. 9B** show NET-degradation correlates with disease activity in STEC-HUS. FIG. 9A shows images and FIG. 9B shows quantification of DNA-staining of NETs generated in vitro after incubation with buffer, plasma from normal healthy donors (N=7), or plasma collected in the acute disease state (Acute) and in remission (Rem.) of 11 patients with STEC-HUS. Scale bar: 50 μ m. Statistics: one-way ANOVA followed by Bonferroni's multiple

comparisons post test; § $P < 0.001$ vs. all other groups.

FIG. 10 provides a table of descriptions of human tissue from various patients examined, such as patient number, cause of death as indicated in the autopsy report (ARDS, acute respiratory distress syndrome). Sex and age of the patient. Type of analyzed tissue. The tissues was analyzed in a three step procedure: first, identification of hematoxylin-rich clots / aggregates in hematoxylin & eosin stainings; second, identification of NETs in hematoxylin-rich clots by immunostaining for chromatin & myeloperoxidase (MPO); third, identification of the localization of the NET-clots within the lung architecture.

FIG. 11A-FIG. 11D show human NETs form clots in situ. Lung tissue collected at autopsy from two patients with sepsis (Patient #6 and #7 in FIG. 10). FIG. 11A shows H&E staining of two hematoxylin-rich intravascular clots in lung tissue from patient #6. FIG. 11B shows immunostaining of consecutive section for chromatin and myeloperoxidase (MPO) of left blood vessel shown in panel A. FIG. 11C shows H&E staining blood vessel in lung tissue from patient #7 shows hematoxylin-rich intravascular clots. FIG. 11D shows immunostaining of consecutive section for chromatin and MPO. Arrows point to clots of NETs. Scale bar: 50 μm .

FIG. 12 shows NET-clots occlude blood vessels in septic mice. Immunostaining of fully occluded blood vessel in septic DNase1/DNase1l3^{-/-} mice for DNA, chromatin and myeloperoxidase (MPO). DNA and chromatin stainings of NETs (arrows) appear dimmer than nuclear stainings (asterisks) and cover the intercellular space. Scale bars: 10 μm .

FIG. 13A-FIG. 13F shows that DNase1 and DNase1l3 degrade NETs by distinct mechanisms. FIG. 13A provides DNA-stainings of PFA-fixed NETs generated in vitro after incubation with sera from WT, DNase1^{-/-} (D1^{-/-}), DNase1l3^{-/-} (D1l3^{-/-}), DNase1/DNase1l3^{-/-} mice (D1/D1l3^{-/-}). Data are representative of ≥ 3 independent experiments. Scale bar: 50 μm . FIG. 13B shows the concentration of cell-free DNA fragments released from PFA-fixed NETs after incubation with sera from WT, D1^{-/-}, D1l3^{-/-}, and D1/D1l3^{-/-} mice. FIG. 13C shows DNA-stainings of non-fixed NETs generated in vitro after incubation with vehicle, recombinant murine DNase1 (rmD1), or recombinant murine DNase1l3 (rmD1l3). FIG. 13D shows the concentration of cell-free DNA fragments released from non-fixed NETs generated in vitro after incubation with vehicle, rmD1, or rmD1l3. FIG. 13E shows DNA-stainings of PFA-fixed NETs generated in vitro after incubation with vehicle, rmD1, or rmD1l3. FIG. 13F shows the concentration of cell-free DNA fragments released from PFA-fixed NETs generated in vitro after incubation with vehicle, rmD1, or rmD1l3. Statistics by one-way ANOVA followed by Bonferroni's multiple comparisons post test; § $P < 0.0001$.

FIG. 14A-FIG. 14G provide DNase1 and DNase1l3 activity in the circulation of humans and mice. FIG. 14A-FIG. 14E show DNase activity in human and murine plasma quantified by SRED. FIG. 14A illustrates supplementation of plasma from normal healthy donors (NHD) with polyclonal antibodies against human DNase1 (α -hDNase1), but not with IgG, blocks the DNA degrading activity in a concentration-dependent manner (*: $p < 0.05$ vs. IgG). FIG. 14B shows plasma supplemented with 200 $\mu\text{g}/\text{ml}$ α -hDNase1 and heparin, an inhibitor of DNase1l3, blocks residual DNase activity in plasma supplemented with 200 $\mu\text{g}/\text{ml}$ α -hDNase1 (*: $p < 0.05$ vs. NHD). FIG. 14C-FIG. 14E provide a comparison of DNase activity in serial dilutions of plasma

from mice and NHD. FIG. 14C illustrates that total DNase activity of WT mice is approximately 10-times higher than in NHD. FIG. 14D shows DNase1 activity in plasma from DNase1I3^{-/-} mice is approximately 10-fold higher than human plasma supplemented with 500 µg/ml heparin. (E DNase1I3 activity in plasma from DNase1^{-/-} mice is approximately 10-fold higher than human plasma supplemented with 200 µg/ml α-hDNase1. (F, G) In vitro NET-degradation is illustrated in FIG. 14F and FIG. 14G. FIG. 14F shows activated neutrophils incubated with plasma from NHD alone or supplemented with 200 µg/ml α-hDNase1, plasma from TMA patients, and plasma from WT, DNase1^{-/-}, and DKO mice (Scale bars: 200 µm). FIG. 14G shows cell-free DNA in supernatants of activated neutrophils incubated with plasma from indicated sources (§: $p < 0.001$ vs. NHD). In vitro NET-degradation by human plasma is dependent on DNase1 as DNase1I3 in human plasma is not sufficient to degrade NETs.

FIG. 15A-FIG. 15C show a lethal course of chronic neutrophilia in an alternative strain of DNase1/DNase1I3^{-/-} mice. An alternative strain of DNase1^{-/-} mice (see Methods of Example 1) was crossed with DNase1I3^{-/-} mice to generate mice with a combined deficiency. FIG. 15A provides survival curves of neutrophilia in WT (N=9), DNase1^{-/-} (D1^{-/-}, N=5), and DNase1/DNase1I3^{-/-} mice (D1/D1L3^{-/-}, N=5). Chronic neutrophilia was induced by injection of a CSF3-expression plasmid (CSF3). FIG. 15B provides hematoxylin and eosin stainings (H&E) of lung sections. Arrow points to hematoxylin positive clots in D1/D1L3^{-/-} mice. Scale bar: 500 µm. FIG. 15C provides survival of D1/D1L3^{-/-} mice coexpressing CSF3 with a control plasmid (CSF3/Ctrl, N=6), with DNase1 (CSF3/D1, N=5), or with DNase1I3 (CSF3/D1L3, N=6). Statistics: Log-rank test; § $P < 0.001$.

FIG. 16A-FIG. 16B show the screening approach to identify hyperactive variants of DNase1 and DNase1L3 in vitro. FIG. 16A shows a library of DNases included wild-type DNase1L3 (I, D1L3), wild-type DNase1 (II, D1), and variants thereof. D1 and D1L3 variants were designed by transferring amino acids from D1L3 to D1 (III, D1^V) and D1 to D1L3 (IV, D1L3^V). DNase activity was tested using the degradation of high-molecular weight (HMW) DNA, dsDNA and chromatin, as a substrate. In FIG. 16B, zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. D1, 0.0005 ng). D1 degrades dsDNA approximately 100-fold more efficiently than D1L3. Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. D1L3 degrades chromatin approximately 100-fold more efficiently than D1. At a ratio of 10:1 or 1:1 (m/m), D1 and D1L3 show synergistic effects in the degradation of chromatin.

FIG. 17 shows an amino acid sequence alignment of human DNase1 (SEQ ID NO: 1) and human DNase1L3 (SEQ ID NO: 2). The signal peptide, conserved amino acids, additional arginine- and lysine-residues in variable regions of D1L3, and the C-terminal tail of D1L3 are highlighted. Sequence rulers indicate the amino acid position.

FIG. 18 shows a list of arginine- and lysine-residues that are present in DNase1L3 only. The corresponding amino acid in DNase1 and amino acid substitution in DNase1 are indicated. Numbers in brackets are the amino acid position in the mature protein without signal peptide.

Arginine- and lysine-residues that are located in the C-terminal tail of D1L3 are shown in lanes 29-37. This application tested 22 of the 37 possible DNase1 variants.

FIG. 19 shows the characterization of DNase1 variants with single arginine and lysine amino acid substitutions listed in FIG. 18. Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. Amino acid substitutions that cause an increase in chromatin degrading activity are highlighted in blue. Samples without such effect are shown in light blue.

FIG. 20 shows the amino acid sequence alignment of human DNase1 (SEQ ID NO: 1), mouse DNase1 (SEQ ID NO: 3), rat DNase1 (SEQ ID NO: 4), and human DNase1L3 (SEQ ID NO: 2). Sequence rulers indicate the amino acid position. The signal peptides are highlighted in grey. Cationic amino acid residues (arginine: R, lysine: K) that are present in mouse/rat DNase1 and human DNase1L3, but not in human DNase1, are highlighted. Four such R/K-residues are shown in mouse DNase1 (Q31R, Q49K, Q79R, W209R). Rat DNase1 contains two additional R/K-residues (Q177K, G262K).

FIG. 21A-FIG. 21B shows a comparison between murine and human DNase1. (FIG. 21A) Zymography show dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Murine DNase1 (mD1) and human DNase1 (D1) show similar levels of dsDNA degrading activity. Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. Murine DNase1 (mD1) shows more chromatin degrading activity than human DNase1 (D1). (FIG. 21B) Gene therapy with human DNase1 provides only partial protection from lethal vascular occlusion by NETs. Chronic neutrophilia with concomitant intravascular NET-formation was induced as by hepatic expression of *Csf3*, which encodes with G-CSF. The data show the survival curve of *Csf3*-expressing *Dnase1*^{-/-} *Dnase1*^{1/3}^{-/-} mice. Animals were injected with *Csf3* and an empty control vector (*Ctrl*, N = 6), with a mixture of *Csf3* and human DNase1 (*hDnase1*, N = 12), and with *Csf3* and murine DNase1 (*mDnase1*, N = 5). *P*-values were calculated using log-rank test.

FIG. 22A-FIG. 22B shows the characterization of rodent-like human DNase1-variants in vivo and in vitro. (FIG. 20A) Gene therapy with rodent-like DNase1-variants provides full protection from lethal vascular occlusion by NETs. Chronic neutrophilia with concomitant intravascular NET-formation was induced as by hepatic expression of *Csf3*, which encodes with G-CSF. The data show the survival curve of *Csf3*-expressing *Dnase1*^{-/-} *Dnase1*^{1/3}^{-/-} mice. Animals were injected with *Csf3* and an empty control vector (*Ctrl*, N = 6), with a mixture of *Csf3* and human DNase1 (*hDnase1*, N = 12) as shown in FIG. 21. A third group of *Dnase1*^{-/-} *Dnase1*^{1/3}^{-/-} mice received an injection with *Csf3* and the gene encoding with murine-like DNase1-variant, which contains 4 amino acid mutations Q31R/Q49K/Q79R/W209R (Q31R/Q49K/Q79R/W209R, N = 6). A fourth group of *Dnase1*^{-/-} *Dnase1*^{1/3}^{-/-} mice received an injection with *Csf3* and the gene

encoding with rat-like DNase1-variant, which contains 6 amino acid mutations Q31R, Q49K, Q79R, Q177K, W209R, and G262K (Q31R/Q49K/Q79R/Q177K/W209R/G262K, N = 6). *P*-values were calculated using log-rank test. (FIG. 22B) Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. The murine-like variant of human DNase1 (Q31R/Q49K/Q79R/W209R), and the rat-like variant of human DNase1 (Q31R/Q49K/Q79R/Q177K/W209R/G262K) show similar dsDNA but more chromatin degrading activity than wild-type human DNase1.

FIG. 23 shows the amino acid sequence alignment of human DNase1 (D1, SEQ ID NO: 1) and human DNase1L3 (D1L3, SEQ ID NO: 2). Sequence rulers demark the amino acid positions. Amino acids of D1 with published mutations are highlighted. Human D1L3 contains three mutations that are associated with a gain-of-function in human DNase1 (Q31R/T227K/A136F) and four mutations that are not linked to increased activity.

FIG. 24A-24B show the characterization of DNase1 variants (D1^V) with the mutations Q31R, T227K, and/or A136F. (FIG. 24A) Culture supernatants of transfected HEK293 cells were analyzed. HEK cells transfected with wild type DNase1 (D1) and DNase1L3 (D1L3) served as controls. Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. The D1^V that features three amino acid mutations Q31R/T227K/A136F has similar dsDNA degrading activity, but a greater capacity to degrade chromatin than D1 and D1^V featuring Q31R, T227K, A136F, or Q31R/T227K. (FIG. 24B) Gene therapy with a hyperactive D1^V provides full partial protection from lethal vascular occlusion by NETs. Chronic neutrophilia with concomitant intravascular NET-formation was induced as by hepatic expression of *Csf3*, which encodes with G-CSF. The data show the survival curve of *Csf3*-expressing *Dnase1*^{-/-}*Dnase1l3*^{-/-} mice. Animals were injected with *Csf3* and an empty control vector (*Ctrl*, N = 6), with a mixture of *Csf3* and human DNase1 (*hDnase1*, N = 12) as shown in FIG. 21. In addition, a group of *Dnase1*^{-/-}*Dnase1l3*^{-/-} mice received an injection with *Csf3* and the gene encoding with a human DNase1-variant, which contains the triple amino acid mutation Q31R/T227K/A136F (Q31R/T227K/A136F, N = 6). *P*-values were calculated using log-rank test.

FIG. 25 shows a comparison of wild-type DNase1 (D1) and the DNase1-variants (D1^V) featuring the mutations Q31R/T227K/A136F. Purified proteins were analyzed at indicated amounts. Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. 0 ng). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular

weight DNA that correlates with chromatin degrading activity. The DNase1-variant featuring the mutations Q31R/T227K/A136F degrades chromatin approximately 10-20-fold more efficiently than wild-type DNase 1.

FIG. 26 shows a comparison of DNase1 variants (D1^V) with the amino acid mutations A164K, A136F, and A164K/A136F with wild-type DNase1 (D1) and wild-type (D1L3). Culture supernatants of transfected HEK cells were analyzed. Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. DNase1-variants featuring A136F/A164K degrades chromatin most efficiently.

FIG. 27A-FIG. 27B shows development of a DNase1 variant with the mutation A226_T227insK. (FIG. 26A) Amino acid sequence alignment of human DNase1 (SEQ ID NO: 1) and human DNase1L3 (SEQ ID NO: 2). Sequence rulers demark the amino acid positions. Critical arginine residues in DNase1L3 are highlighted. Shared amino acids between DNase1 and DNase1L3 are highlighted and serve as anchors to replace the amino acids sequence ATP of DNase1 with VKKS of DNase1L3. (FIG. 26B) Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). The gel was loaded with wild-type DNase1 (D1), DNase1 variants (D1^V) that feature the mutations A226_T227insK, T227K, and A226_P228delinsVKKS. A226_T227insK, but not A226_P228delinsVKKS, is associated with a reduced dsDNA degrading activity.

FIG. 28 shows the structure of human DNase1 [4AWN]. Highlighted are the N-terminal beta-sheet and the C-terminal beta-sheet and the motive S272/D273/H274, which is conserved among D1-protein family members.

FIG. 29 shows the concept of building block engineering of homologous proteins. The technology transfers single or multiple variable amino acids, which are flanked by conserved single or multiple variable amino acids, between a donor and recipient protein.

FIG. 30 shows an amino acid sequence alignment of DNase1 and DNase1L3 of mouse (SEQ ID NOS: 3 and 34), rat (SEQ ID NOS: 4 and 35), chimpanzee (SEQ ID NOS: 36 and 37), and human (SEQ ID NOS: 1 and 2). The N-terminal signal peptide and conserved amino acids are highlighted. Variable amino acids are not highlighted and serve as Building Blocks that can be transferred from DNase1 to DNase1L3 and vice versa. Abbreviations: AA, amino acid.

FIG. 31A-FIG. 31B show lists of Building Blocks in human DNase1 (D1) and human DNase1L3 (D1L3). FIG. 31A shows amino acids that are conserved in D1 and D1L3, which serve as N- and C-anchors, respectively. Building blocks are variable amino acids in D1 and D1L3. Mutations that transfer Building Blocks from D1L3 to D1 are shown. FIG. 31B shows N- and C-anchors in D1L3. Mutations that transfer Building Blocks from D1 to D1L3 are listed. AA: amino acid.

FIG. 32 shows an application of the building block engineering of homologous proteins. The application uses as an initial screening step, the transfer of clusters of building blocks between a homologous donor and recipient protein. Additional optional steps are the transfer of individual building blocks, followed by the transfer of individual amino acids. In a final step (not shown), multiple amino acids, building blocks, and building block clusters may be combined to degenerate a chimeric enzyme.

FIG. 33 shows characterization of DNase1 variants (D1^V) featuring building blocks from DNase1L3 (D1L3). Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. Building block substitutions that cause an increase in chromatin degrading activity are highlighted in dark shade. Samples without such effect are shown in light shade. A DNase1 variant featuring the combination of building blocks 11, 12-14, 26, 41-48, and 49 shows similar chromatin degrading activity than wild-type DNase1L3.

FIG. 34A-FIG. 34B shows the development of a DNase1L3 variant without the C-terminal tail of wild-type DNase1L3. **FIG. 34A** shows an amino acid sequence alignment of human DNase1 (SEQ ID NO: 1) and human DNase1L3 (SEQ ID NO: 2). Sequence rulers demark the amino acid positions. The C-terminal tail of DNase1L3 is highlighted and its nuclear localization signal is boxed. Shared amino acids between DNase1 and DNase1L3 are indicated. Amino acids located after the conserved SDH motive were exchanged the building block cluster 60-62 between DNase1 and DNase1L3. **FIG. 34B** shows zymography showing dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. The gels were loaded with wild-type DNase1 (D1), DNase1 variant (D1^V) that features the BB cluster 60-62 from D1, DNase1L3 variant (D1L3^V) that features the BB cluster 60-62 from D1, and a D1L3^V with two mutations (K301A, K303A) were made to inactivate the NLS2. D1L3^V with BB cluster 60-62 from D1 showed increased chromatin degrading activity compared to wild-type D1L3.

FIG. 35A-FIG. 35B shows the development of glycosylated DNase1L3 variants (D1L3^V). **FIG. 35A** shows the amino acid sequence alignment of human DNase1 (SEQ ID NO: 1) and human DNase1L3 (SEQ ID NO: 2). Sequence rulers demark the amino acid positions. The glycosylation motives N40-X-T42 and N128-X-T130 of DNase1 are highlighted. The sites are part of building block (BB) 4 and building block cluster (BB) 23-25. Relevant shared amino acids between DNase1 and DNase1L3 are highlighted. (**FIG. 35B**) Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. Ctrl).

Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. The gels were loaded with wild-type DNase1L3 (D1L3), DNase1L3 variant (D1L3^V) that features BB 4 from D1, and DNase1L3 variant (D1L3^V) that features the BB cluster 23-25 from D1. A Western Blot (WB) with antibodies against D1L3 detected that D1L3^V showed increased amounts of larger proteins. Both glycosylated DNase1L3 variants (D1L3^V) retained their capacity to degrade chromatin.

FIG. 36A-FIG. 36F shows the therapeutic effect of DNase1L3 (SEQ ID NO: 2) and a variant thereof (SEQ ID NO: 17) against NETs in vivo. FIG. 36A shows an H&E staining of a mouse heart with a hematoxylin-rich and dense clot of NETs within a cardiac chamber. FIG. 36B shows a time line starting with the induction of NETs, leading to vascular occlusion by NETs and death. Hematuria and hypothermia are external signs for in vivo NET-clots. (FIG. 36C) Outline of the experimental design: *Csf3* was expressed in *Dnase1^{-/-}Dnase1l3^{-/-}* to induce vascular clots of NETs. Mice that showed hematuria and hypothermia were randomized by an injection of vehicle (N = 3), dornase alpha (D1, N = 3), purified DNase1L3 (D1L3, N = 3), or a purified DNase1L3 variant (SEQ ID NO: 17, D1L3^V, N = 3). After 30 minutes lungs and serum were collected. The therapy with D1L3 or D1L3^V, but not saline or D1, significantly reduced vascular occlusion by NETs in lungs (FIG. 36D) and increased the levels of DNA in serum (FIG. 36E). FIG. 36F shows oligo-nucleosomes in the serum from mice receiving the D1L3 therapy. Mice receiving D1L3^V showed smaller mono- and di-nucleosomes. DNA isolates from serum of mice receiving D1 therapy showed faint DNA smears of various sizes, whereas no DNA could be isolated from serum of mice injected with vehicle.

FIG. 37A-FIG. 37C shows the therapeutic effect of a DNase1L3 variant (SEQ ID NO: 17) in a rat model of ischemia-reperfusion (IR) injury. (FIG: 37A) Outline of the experimental design: One testicle of male wild-type rats was subjected to testicular torsion. The procedure causes an ischemic tissue injury. Mice were randomized. A de-torsion of the testicle was performed, followed by an injection of vehicle (N = 6), dornase alpha (D1, N = 6), or a purified DNase1L3 variant (SEQ ID NO: 17, D1L3^V, N = 6). The untreated and the ischemic testicle were collected after 7 days. FIG. 37B shows that the therapy with the D1L3^V, but not saline or D1, significantly reduced atrophy in the ischemic testicle. FIG. 37C quantified the macroscopic tissue injury. All untreated testicles are brownish and vascularized, indicating healthy tissue (Ctrl, Score: 0). Testicular torsion followed by reperfusion damaged tissue, which appeared whitish (IR, Score: 4). Therapy with the D1L3^V, but not saline or D1, substantially reduced the tissue injury and 4/6 testicles showed no signs of injury. Scale bar: 1cm.

FIG. 38A-FIG. 38C shows the expression levels of DNase1, DNase1L3, and variants thereof. (FIG. 38A) Stable pools of CHO cells producing wild-type DNase1 (D1), a variant thereof (D1^V, SEQ ID NO: 7), wild-type D1L3, and a variant thereof (D1L3^V, SEQ ID NO: 17) were cultured in bioreactors. Samples were collected periodically and analyzed by Western Blot. Expression of D1 and D1^V was detected, but only low levels of D1L3 and D1L3^V. (FIG. 38B) Robust

expression of D1L3 and D1L3^V was achieved in *Pichia pastoris*. (FIG. 38C) Zymography showed dsDNA degrading activity as dark circles. The dsDNA degrading activity correlates with the diameter. Samples without activity show the loading well as small black spot (e.g. D1L3, 0.0005 ng). Agarose gel electrophoresis (AGE) of DNA isolated from digested chromatin shows a shift from high-molecular weight DNA to lower or low-molecular weight DNA that correlates with chromatin degrading activity. D1L3^V degrades chromatin approximately 10-fold more efficiently than D1L3.

FIG. 39 illustrates certain DNase variants in accordance with this disclosure. The amino acid mutations in DNase1 and DNase1L3 with signal peptide as well as the mature proteins without signal peptide are listed. Mutations that were generated using Building Block Engineering are marked with "+" and the Building Blocks are indicated in brackets.

DETAILED DESCRIPTION OF THE INVENTION

[0025] The present disclosure provides engineered DNase proteins, members of the DNase1-protein family (including DNase1-like 3 and DNase1) that are useful for treating conditions characterized by neutrophil extracellular trap (NET) accumulation and/or release. The disclosure further provides compositions and methods for preventing or treating vascular occlusion involving NETs. As demonstrated herein, NETs participate in a non-canonical mechanism for vascular occlusion, which is not dependent on fibrin and platelets.

[0026] As used herein, the term "neutrophil extracellular trap" or "NET" refers to any extracellular trap ("ET") comprising extracellular DNA formed by cells such as, but not limited to, neutrophils, monocytes, macrophages, basophils, eosinophils, mast cells, cancer cells, injured cells (e.g., injured endothelial cells), and the like. Unless the context indicates otherwise, the terms NET and ET are used interchangeably herein.

[0027] There are two extracellular DNase enzymes found in the circulation of humans and mice, DNase1 (D1) and DNase1-like 3 (D1L3). These enzymes can be deficient in their ability and mechanism to degrade NETs in vitro and in vivo, and are found at relatively low concentrations in humans and mice. The natural enzymes do not have ideal physical, enzymatic, and/or pharmacodynamic properties for recombinant therapy, and particularly for systemic therapy.

[0028] D1 and D1L3 belong, along with DNase1-like 1 (D1L1) and DNase1-like 2 (D1L2), to the DNase1-protein family. D1 and D1L3 are expressed in a variety of species including, humans, primates, and rodents. In humans and mice, D1 and D1L3 show a protein similarity of 49-52%. However, despite their homology, D1 and D1L3 differ in cellular origin, sensitivity towards inhibitors, and substrate affinity. D1 preferentially cleaves protein-free DNA (e.g. bacterial DNA, plasmid DNA), whereas D1L3 targets chromatin, the complex of DNA and histones, which is

commonly found in the nucleus of eukaryotic cells. D1 activity is inhibited upon binding to monomeric actin and sensitive to physiological salt concentrations. In addition, D1 is glycosylated at N40 (corresponds to N18 in the mature enzyme without signal peptide) and N128 (N106), which makes the enzyme resistant to inactivation by serum proteases. In contrast, D1L3 lacks glycosylation and actin-binding sites, which causes its susceptibility towards several proteases and resistance towards actin, respectively.

[0029] The disclosure provides D1L3 variants engineered to have physical, pharmacodynamic, and/or enzymatic properties more suitable for therapy, for example, to reduce or prevent NET accumulation in a subject. The disclosure provides a recombinant D1L3 protein variant comprising: one or more glycosylations, one or more amino acid alterations resulting in increased substrate affinity, inactivation of a nuclear localization signal, deletion of all or part of a C-terminal tail, pegylation, and fusion to a carrier protein or moiety. The D1L3 variant may have increased protein stability, increased protease-resistance, increased bioavailability, increased half-life in circulation, and/or substantially the same or better DNA and/or chromatin and/or NET-degrading activity, higher production levels with in vitro expression systems (e.g. Chinese hamster ovary cells and/or *Pichia pastoris*) as compared to wild-type D1L3 protein of SEQ ID NO:2.

[0030] The D1L3 variant may comprise an amino acid sequence with at least 80% sequence identity to the enzyme defined by the amino acid sequence of SEQ ID NO:2, with one or more amino acid modifications with respect to SEQ ID NO:2 as described herein. The D1L3 variant may comprise an amino acid sequence with at least 90%, or at least 95%, or at least 97%, or at least 98% sequence identity with the DNase enzyme defined by SEQ ID NO:2.

[0031] As used herein, when referring to sequence identity with wild-type DNase enzymes, and unless stated otherwise, sequences refer to mature enzymes lacking the signal peptide. Further, unless stated otherwise, amino acid positions are numbered with respect to the full translated DNase sequence, including signal peptide, for clarity. Accordingly, for example, reference to sequence identity to the enzyme of SEQ ID NO:2 refers to a percent identity with the mature enzyme having M21 at the N-terminus. Similarly, reference to sequence identity to the enzyme of SEQ ID NO:1 (human D1) refers to a percent identity with the mature enzyme having L23 at the N-terminus.

[0032] The similarity of nucleotide and amino acid sequences, i.e. the percentage of sequence identity, can be determined via sequence alignments as known in the art. Such alignments can be carried out with several art-known algorithms, such as with the mathematical algorithm of Karlin and Altschul (Karlin & Altschul (1993) Proc. Natl. Acad. Sci. USA 90: 5873-5877), with hmalign (HMMER package, <http://hmmer.wustl.edu/>) or with the CLUSTAL algorithm (Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994) Nucleic Acids Res. 22, 4673-80). Exemplary algorithms are incorporated into the BLASTN and BLASTP programs of Altschul et al (1990) J. Mol. Biol. 215: 403-410. When utilizing BLAST programs, the default parameters of the respective programs are used.

[0033] The recombinant D1L3 protein variant may comprise one or more glycosylation consensus sequences, e.g., NX(T/S). The human wild type enzyme (SEQ ID NO:2) does not contain a consensus glycosylation sequence. However, engineering a glycosylation consensus sequence into D1L3 can provide for glycosylation (and protease resistance) without substantial impact on the enzyme activity. The glycosylation consensus sequence may include a D38N substitution and an N40T or N40S substitution, and/or an A127N substitution and a V129T or V129S substitution from the wild type sequence (e.g., SEQ ID NO:2). The recombinant D1L3 enzyme may have one, two, three, or four glycosylation sites. The consensus sequence can be introduced into the protein at a location that does not affect its enzymatic activity, and may be located near portions of the enzyme that are important for maintaining structure of the enzyme (e.g., such as within amino acids 25 to 289, and within amino acids 24 to 150).

[0034] The disclosure provides a DNase enzyme comprising an amino acid sequence that is at least 80% identical to the amino acid sequence of the enzyme defined by SEQ ID NO:2, wherein the enzyme has a substitution of asparagine at the position corresponding to position 38 and/or position 127 of SEQ ID NO:2, and the N38 and/or N127 is glycosylated. The substitution at position 38 and/or 127 may be made via a building block substitution from SEQ ID NO: 1 as described herein.

[0035] D1L3 variants having one or more glycosylations may exhibit increased resistance to serum protease degradation and/or an increased serum half-life compared to wild-type D1L3 protein (SEQ ID NO:2).

[0036] D1L3 protein variant may comprise an inactivation of a nuclear localization signal (NLS). D1L3 features two nuclear localization sites (NLS1, NLS2), which both target the enzyme to the nucleus during apoptosis. Indeed, D1L3 is required for fragmentation of nuclear DNA in apoptotic and necrotic cells *in vivo*. NLS1 is located near the N-terminus (about amino acid positions 80 to 96 with respect to SEQ ID NO: 2). NLS2 (amino acid positions 291 to 304 with respect to SEQ ID NO: 2) is embedded within a C-terminal tail (amino acid positions 283 to 305 with respect to SEQ ID NO: 2) that is unique to D1L3 and not present in D1. The C-terminal tail is thought to be required for D1L3 to degrade different substrates of extracellular DNA, namely lipid-encapsulated DNA and chromatin within apoptotic bodies. Lipids encapsulate DNA during transfections. In brief, cDNA, protein-free DNA, and cationic lipids form a complex that penetrates through the plasma membrane of target cells. D1L3 interferes with transfections and the C-terminal tail is critical for this function. Physiological substrates of D1L3 are apoptotic bodies, extracellular lipid vesicles filled with chromatin from apoptotic cells. The C-terminal tail enables D1L3 to penetrate through lipid membranes of apoptotic bodies and degrade the chromatin load. The C-terminal tail is also thought to be required for degradation of lipid-free, extracellular chromatin by D1L3. Thus, the C-terminal tail, a molecular feature unique to D1L3, appears to be responsible for the distinct substrate affinities of D1 and D1L3.

[0037] Inactivation of one or more NLSs enhances the availability and/or activity of the enzyme in the circulation. The NLS inactivation may be by deletion of all or part of NLS1 and/or NLS2.

The NLS may be inactivated by substitution and/or deletion of amino acids within NLS1 and/or NLS2. NLS2 may be deleted, entirely or partially. The D1L3 protein variant may comprise a deletion of all or part of the C-terminal tail, and/or substitution of one or more amino acids in the C-terminal tail.

[0038] The D1L3 variant may contain one or more, e.g., 1, 2, 3, 4, 5, or more amino acid substitutions, additions (e.g., insertions), or deletions in the NLS1 domain. The NLS2 domain or a portion thereof may be deleted. The D1L3 protein variant may contain one or more, e.g., 1, 2, 3, 4, 5, or more amino acid substitutions, additions, or deletions in the C-terminal tail domain. The C-terminal tail domain or a portion thereof may be deleted. The D1L3 protein variant may contain one or more, e.g., 1, 2, 3, 4, 5, or more amino acid substitutions, additions, or deletions in the NLS2 domain which is located in the C-terminal tail domain. The NLS2 domain or a portion thereof can be deleted in the D1L3 protein variant.

[0039] The disclosure provides a protein engineering technology that is based on a transfer of a single amino acid or multiple-adjacent amino acids, termed "building block", between two members of a protein family, such as DNase1-protein family members to generate enzymatically active variants of DNase1-protein family members (including variant of D1 and D1L3). A "building block" is defined by amino acids that are variable between two or more members of the DNase1-protein family. These variable amino acids are flanked by amino acids that are conserved between two or more members of the DNase1-protein family ("anchors"). The variable single amino acid or multiple contiguous amino acids ("building blocks") are exchanged between members of the DNase1-protein family by implanting them between conserved single amino acid or multiple contiguous amino acids ("anchors").

[0040] This approach is referred to herein as "building-block protein engineering." Where three or more amino acids are transferred, up to 1/3 of the amino acids may be further substituted. For example, where three or six amino acids are transferred as a building block, one or up to two residues may be further substituted, respectively. Four or more amino acids may be transferred as a building block substitution, and up to 25% of the transferred amino acids are further substituted, e.g., with conservative or non-conservative amino acid modifications. For example, where four, eight, or twelve amino acids are transferred, one, two, or three amino acids (respectively) may be further substituted in the building block substitution.

[0041] The D1L3 variant may further comprise at least one building block substitution from D1. FIG. 31 illustrates 62 building block substitutions between D1 and D1L3.

[0042] Exemplary building block substitutions from D1 include one or more of: substitution of 1-20 (MSRELAPLLLLLSIHSALA) of SEQ ID NO:2 with 1-22 (MRGMKLLGALLALAALLQGAVS) from SEQ ID NO:1; substitution of 21-25 (MRICS) of SEQ ID NO:2 with 23-27 (LKIAA) from SEQ ID NO:1; substitution of 28-30 (VRS) of SEQ ID NO:2 with 30-32 (IQT) from SEQ ID NO:1; substitution of 33-34 (ES) of SEQ ID NO:2 with 35-36 (ET) from SEQ ID NO:1; substitution of 36-45 (QEDKNAMDVI) of SEQ ID NO:2 with 38-47 (MSNATLVSYI) from SEQ ID NO:1; substitution of 47-51 (KVIK) of SEQ ID NO:2 with 49-53

(QILS) from SEQ ID NO:1; substitution of 52 (C) of SEQ ID NO:2 with 54 (Y) from SEQ ID NO:1; substitution of 54-58 (IILVM) of SEQ ID NO:2 with 56-60 (IALVQ) from SEQ ID NO:1; substitution of 60-61 (IK) of SEQ ID NO:2 with 62-63 (VR) from SEQ ID NO:1; substitution of 63-70 (SNNRICPI) of SEQ ID NO:2 with 65-72 (SHLTAVGK) from SEQ ID NO:1; substitution of 72-74 (MEK) of SEQ ID NO:2 with 74-76 (LDN) from SEQ ID NO:1; substitution of 77-84 (RNSRRGIT) of SEQ ID NO:2 with 79-84 (QDAPDT) from SEQ ID NO:1; substitution of 86 (N) of SEQ ID NO:2 with 86 (H) from SEQ ID NO:1; substitution of 88-89 (VI) of SEQ ID NO:2 with 88-89 (VV) from SEQ ID NO:1; substitution of 91-92 (SR) of SEQ ID NO:2 with 91-92 (EP) from SEQ ID NO:1; substitution of 96-97 (NT) of SEQ ID NO:2 with 96-97 (NS) from SEQ ID NO:1; substitution of 101 (Q) of SEQ ID NO:2 with 101 (R) from SEQ ID NO:1; substitution of 103 (A) of SEQ ID NO:2 with 103 (L) from SEQ ID NO:1; substitution of 105 (L) of SEQ ID NO:2 with 105 (V) from SEQ ID NO:1; substitution of 107-110 (KEKL) of SEQ ID NO:2 with 107-110 (RPDQ) from SEQ ID NO:1; substitution of 113-116 (VKRS) of SEQ ID NO:2 with 113-116 (AVDS) from SEQ ID NO:1; substitution of 118 (H) of SEQ ID NO:2 with 118 (Y) from SEQ ID NO:1; substitution of 120 (H) of SEQ ID NO:2 with 120 (D) from SEQ ID NO:1; substitution of 122-127 (YQDGDA) of SEQ ID NO:2 with 122-128 (GCEPCGN) from SEQ ID NO:1; substitution of 129 (V) of SEQ ID NO:2 with 130 (T) from SEQ ID NO:1; substitution of 131 (S) of SEQ ID NO:2 with 132 (N) from SEQ ID NO:1; substitution of 135-136 (FV) of SEQ ID NO:2 with 136-137 (AIV) from SEQ ID NO:1; substitution of 138 (W) of SEQ ID NO:2 with 139 (R) from SEQ ID NO:1; substitution of 140-143 (QSPH) of SEQ ID NO:2 with 141-144 (FSRF) from SEQ ID NO:1; substitution of 145-148 (AVKD) of SEQ ID NO:2 with 146-149 (EVRE) from SEQ ID NO:1; substitution of 150 (V) of SEQ ID NO:2 with 151 (A) from SEQ ID NO:1; substitution of 152 (I) of SEQ ID NO:2 with 153 (V) from SEQ ID NO:1; substitution of 156-157 (TT) of SEQ ID NO:2 with 157-158 (AA) from SEQ ID NO:1; substitution of 159-161 (ETS) of SEQ ID NO:2 with 160-162 (GDA) from SEQ ID NO:1; substitution of 163 (K) of SEQ ID NO:2 with 164 (A) from SEQ ID NO:1; substitution of 167 (E) of SEQ ID NO:2 with 168 (A) from SEQ ID NO:1; substitution of 169-170 (VE) of SEQ ID NO:2 with 170-171 (YD) from SEQ ID NO:1; substitution of 173 (T) of SEQ ID NO:2 with 174 (L) from SEQ ID NO:1; substitution of 176-178 (KHR) of SEQ ID NO:2 with 177-179 (QEK) from SEQ ID NO:1; substitution of 180-181 (KA) of SEQ ID NO:2 with 181-182 (GL) from SEQ ID NO:1; substitution of 183-186 (NFIF) of SEQ ID NO:2 with 184-187 (DVML) from SEQ ID NO:1; substitution of 198-201 (PKKA) of SEQ ID NO:2 with 199-202 (RPSQ) from SEQ ID NO:1; substitution of 203-204 (KN) of SEQ ID NO:2 with 204-205 (SS) from SEQ ID NO:1; substitution of 208 (R) of SEQ ID NO:2 with 209 (W) from SEQ ID NO:1; substitution of 210 (D) of SEQ ID NO:2 with 211 (S) from SEQ ID NO:1; substitution of 212 (R) of SEQ ID NO:2 with 213 (T) from SEQ ID NO:1; substitution of 214 (V) of SEQ ID NO:2 with 215 (Q) from SEQ ID NO:1; substitution of 218 (G) of SEQ ID NO:2 with 219 (P) from SEQ ID NO:1; substitution of 220-221 (QE) of SEQ ID NO:2 with 221-222 (SA) from SEQ ID NO:1; substitution of 225-228 (VKKS) of SEQ ID NO:2 with 226-228 (ATP) from SEQ ID NO:1; substitution of 230 (N) of SEQ ID NO:2 with 230 (H) from SEQ ID NO:1; substitution of 238-240 (LRG) of SEQ ID NO:2 with 238-240 (VAG) from SEQ ID NO:1; substitution of 241-246 (QEIVSS) of SEQ ID NO:2 with 241-246 (MLLRGA) from SEQ ID NO:1; substitution of 250 (K) of SEQ ID NO:2 with 250 (D) from SEQ ID NO:1; substitution of 252-254 (NSV) of SEQ ID NO:2 with 252-254 (ALP) from SEQ ID NO:1; substitution of 256 (D) of SEQ ID NO:2 with 256 (N) from SEQ ID NO:1; substitution of 259-260 (KA) of SEQ ID NO:2 with 259-260 (AA) from

SEQ ID NO:1; substitution of 262 (K) of SEQ ID NO:2 with 262 (G) from SEQ ID NO:1; substitution of 264-267 (TEEE) of SEQ ID NO:2 with 264-267 (SDQL) from SEQ ID NO:1; substitution of 269-271 (LDV) of SEQ ID NO:2 with 269-271 (QAI) from SEQ ID NO:1; substitution of 275 (F) from SEQ ID NO:2 with 275 (Y) from SEQ ID NO:1; substitution of 279-280 (FK) of SEQ ID NO:2 with 279-280 (VM) from SEQ ID NO: 1; and substitution of 282-305 (QSSRAFTNSKKSVTLRKKTskRS) of SEQ ID NO:2 with 282 (K) from SEQ ID NO: 1.

[0043] These building block substitutions from D1 result in variants of D1L3 which feature one or more of the following mutations: M1_A20delinsMRGMKLLGALLALAALLQGAVS, M21_S25delinsLKIAA, V28_S30delinsIQT, E33_S34delinsET, Q36_I45delinsMSNATLVSYI, K47_K50delinsQILS, C52Y, I54_M58delinsIALVQ, I60_K61delinsVR, S63_I70delinsSHLTAVGK, M72_K74delinsLDN, R77_T84delinsQDAPDT, N86H, V88_I89delinsVV, S91_R92delinsEP, N96_T97delinsNS, Q101R, A103L, L105V, K107_L110delinsRPDQ, V113_S116delinsAVDS, H118Y, H120D, Y122_A127delinsGCEPCGN, V129T, S131N, I35F_I36VdelinsAI, W138R, Q140_H143delinsFSRF, A145_D148delinsAVKD, V150A, I152A, T156_T157delinsAA, E159_S161delinsGDA, K163A, E167A, V169_E170delinsYD, T173L, K176_R178delinsQE, K180_A181delinsGL, N183_F186delinsDVML, P198_A201delinsRPSQ, K203_N204delinsSS, R208W, D210S, R212T, V214Q, G218P, Q220_E221delinsSA, V225_S228delinsATP, N230H, L238_R239delinsVA, Q241_S246delinsMLLRGA, K250D, N252_V254delinsALP, D256N, K259_A260delinsAA, K262G, T264_E267delinsSDQL, L269_V271delinsQAI, F275Y, F279_K280delinsVM, Q282_S305delinsK.

[0044] The term "delins" refers to a deletion between and including two indicated amino acids, with an insertion of an amino acid or sequence of amino acids at the site of the deletion. For example, the notation E91_P92delinsSR means that the amino acids from E91 to P92 are deleted and the amino acids SR are inserted at the site of the deletion (e.g., the resulting amino acid sequence will have S91 and R92).

[0045] The term "ins" refers to an insertion of amino acids between two indicated amino acids. For example, the notation E91_P92insSR means that the amino acids SR are inserted between E91 and P92, resulting in the sequence E91, S92, R93, and P94.

[0046] Multiple mutations in one enzyme are separated by "/" e.g., T227K/A136F or R199_Q202delinsPKKA/5204_S205delinsKN/W209R/S211D/T213R/Q215V/P219G/S221_A222delinsQE.

[0047] The D1L3 variant may comprise one or more adjacent building block substitutions. For example, the D1L3 variant may comprise with respect to SEQ ID NO:2 the mutation F275Y/F279_K280delinsVM/Q282_S305delinsK (SEQ ID NO: 17). The D1L3 variant may comprise with respect to SEQ ID NO:2 the mutation Q36_I45delinsMSNATLVSYI (SEQ ID NO: 19). The D1L3 variant may comprise with respect to SEQ ID NO:2 the mutation Y122_A127delinsGCEPCGN/V129T/S131N (SEQ ID NO: 20).

[0048] The D1L3 variant may comprise 2 or 3 substitutions selected from the mutations with

respect to SEQ ID NO:2: Q36_I45delinsMSNATLVSYI, Y122_A127delinsGCEPCGN/V129T/S131N, F275Y/F279_K280delinsVM/Q282_S305delinsK.

[0049] Thus, the disclosure provides a DNase enzyme comprising an amino acid sequence that is at least 80% identical to the amino acid sequence of the enzyme defined by SEQ ID NO:2, and comprises at least one building block substitution from the amino acid sequence of SEQ ID NO: 1. Building block substitutions between SEQ ID NOS: 1 and 2 are illustrated and numbered in **FIG. 31**. For example, the enzyme comprises the mutation F275Y/F279_K280delinsVM/ Q282_S305delinsK, and which optionally comprises the amino acid sequence of SEQ ID NO: 17. The enzyme may comprise the mutations Q36_I45delinsMSNATLVSYI, Y122_A127delinsGCEPCGN, V129T, or S131N, and optionally has the amino acid sequence of SEQ ID NO: 19 or SEQ ID NO: 20.

[0050] Exemplary D1L3 variants include those comprising the amino acid sequence of SEQ ID Nos: 17 to 20. The disclosure also includes derivatives having from 1 to 10 (e.g., 1 to 5) amino acid insertions, deletions, or substitutions with respect to any one of SEQ ID NOS: 17 to 20. The D1L3 variant may thus comprise one or more additional block amino acid substitutions from a homologous DNase (e.g., from D1). Such block substitutions may replace at least 2, at least 3, at least 4, or at least 5 amino acids. The block substitutions may replace from 2 to 20 amino acids, such as from 3 to 15 amino acids, or from 3 to 10 amino acids, with the comparable building block from the homologous DNase.

[0051] The engineered variants of D1L3 protein may comprise one or more additional amino acid substitutions, additions (insertions), deletions, or truncations in the amino acid sequence of human D1L3 (SEQ ID NO: 2). Amino acid substitutions may include conservative and/or non-conservative substitutions.

[0052] For example, "conservative substitutions" may be made, for instance, on the basis of similarity in polarity, charge, size, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the amino acid residues involved. The 20 naturally occurring amino acids can be grouped into the following six standard amino acid groups: (1) hydrophobic: Met, Ala, Val, Leu, Ile; (2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln; (3) acidic: Asp, Glu; (4) basic: His, Lys, Arg; (5) residues that influence chain orientation: Gly, Pro; and (6) aromatic: Trp, Tyr, Phe. "Conservative substitutions" are defined as exchanges of an amino acid by another amino acid listed within the same group of the six standard amino acid groups shown above. For example, the exchange of Asp by Glu retains one negative charge in the so modified polypeptide. In addition, glycine and proline may be substituted for one another based on their ability to disrupt α -helices. As used herein, "non-conservative substitutions" are defined as exchanges of an amino acid by another amino acid listed in a different group of the six standard amino acid groups (1) to (6) shown above.

[0053] The D1L3 protein variant may comprise an N-terminal or C-terminal fusion to a half-life extending moiety, such as albumin, transferrin, an Fc, or elastin-like protein. See US 9,458,218. The D1L3 may be dimerized by an immunoglobulin hinge region. For example, the

engineered enzymes described herein may also include an Fc-fusion domain (e.g. a hinge and CH2 domains and CH3 domains of an immunoglobulin). In other cases, the engineered D1L3 protein is fused to albumin, e.g., human albumin or a fragment thereof. See WO 2015/066550; US 9,221,896. Albumin can be joined to at the N-terminus or the C-terminus of the engineered D1 protein, and may optionally comprise an amino acid linker. D1L3 and D1 may be together dimerized by an Fc hinge region, creating a dimeric molecule with synergistic functional properties for degrading NETs.

[0054] The recombinant D1L3 protein variant may alternatively or additionally comprise one or more polyethylene glycol (PEG) moieties, which may be conjugated at one or more of positions or the C-terminus. The native amino acid at that position may be substituted with an amino acid having a side chain suitable for crosslinking with hydrophilic moieties, to facilitate linkage of the hydrophilic moiety to the peptide. An amino acid modified to comprise a hydrophilic group may be added to the peptide at the C-terminus. The PEG chain(s) may have a molecular weight in the range of about 500 to about 40,000 Daltons. The PEG chain(s) may have a molecular weight in the range of about 500 to about 5,000 Daltons. Alternatively, the PEG chain(s) have a molecular weight of about 10,000 to about 20,000 Daltons.

[0055] The D1L3 protein variant may have increased protein stability, increased serum availability, and substantially the same or better NET-degrading activity (as determined in vitro or in vivo) as compared to wild-type D1L3 protein. The D1L3 variant may comprise one or more of the following properties relative to wild type D1L3: higher protein-free DNA (naked DNA) degradation activity, the same or substantially the same (e.g., at least 50%) chromosomal DNA degradation activity, protease resistance, an increased half-life, and higher production levels with in vitro expression systems (e.g. Chinese hamster ovary cells and/or *Pichia pastoris*).

[0056] DNA fragmentation can be measured using methods known to those skilled in the art such as gel electrophoresis and DNase activity assays. Methods for quantifying NETs, NET degradation, and DNA fragmentation activity are described in, for example, Hakkim et al., Proc. Natl. Acad. Sci USA, 2010, 107(21):9813-9818; Napirei et al. Biochem J, 2005, 389: 355-364. Fuchs et al., J Cell Biol, 2007, 176(2):231-241, Sisirak et al. Cell, 2016, 166:88-101, and Jiménez-Alcázar et al. J Thromb Haemost, 2015, 13, 732-742.

[0057] The disclosure provides DNase1 (D1) variants engineered to have physical, pharmacodynamic, and/or enzymatic properties more suitable for therapy, for example, to reduce or prevent NET accumulation in a subject. The variant may be more amenable to recombinant expression, thereby providing considerable manufacturing advantages. The engineered D1 variant may comprise an amino acid sequence that is at least 80% identical to the mature enzyme defined by the amino acid sequence of SEQ ID NO:1, with one or more amino acid substitutions, additions, or deletions resulting in one or more of a mutated DNA binding site, addition of a chromatin binding site, a mutated actin binding site, mutation of a glycosylation site, addition of a nuclear localization signal (e.g., having similarity or identity to NLS1 or NLS2 of D1L3), and/or a C-terminal domain similar to the C-terminal tail of D1L3. The

D1 variant comprises an amino acid sequence with at least 80% identity to the amino acid sequence of the enzyme defined by SEQ ID NO:1, with one or more amino acid modifications with respect to SEQ ID NO:1 as described herein. The D1L3 variant may comprise an amino acid sequence with at least 90%, or at least 95%, or at least 97%, or at least 98% identity with the enzyme of SEQ ID NO:1.

[0058] The D1 variant may comprise one or more amino acid substitutions, additions, deletions, or truncations in the amino acid sequence of human D1 (SEQ ID NO: 1). The amino acid substitutions may include conservative and/or non-conservative substitutions as described. For example, the D1 variant may have from 1 to 20 (or from 1 to 10, or from 1 to 5) amino acid substitutions, deletions, or insertions with respect to the enzyme of SEQ ID NO:1.

[0059] The D1 variant may comprise one or more additional block amino acid substitutions from a homologous DNase (e.g., from D1L3). Such block substitutions may each replace from 1 to 24 amino acids, such as from 3 to 15 amino acids, or from 3 to 10 amino acids, from the homologous DNase. The D1 variant may comprise from 2 to 5 block substitutions from the homologous DNase (e.g., D1L3), e.g. 2 or 3 contiguous building block substitutions are transferred.

[0060] The D1 variant may be engineered to comprise one or more of the following characteristics relative to the wild type enzyme: the substantially the same or higher protein-free DNA (naked DNA) degradation activity (e.g., at least 50% or better), higher chromosomal DNA degradation activity, similar or improved protease resistance, actin resistance, an increased serum half-life, improved penetration from blood into urine or bile, or a combination of such properties.

[0061] The engineered D1 enzyme may comprise an amino acid sequence comprising at least 50% (or at least 70%, 80%, 90%) sequence identity to the C-terminal domain of wild-type D1L3 protein (e.g., amino acids 283 to 305 of SEQ ID NO:2).

[0062] The D1 enzyme may comprise an amino acid sequence having at least 50% or at least 75% sequence identity to the NLS1 of wild-type D1L3 (amino acids 35 to 51 of SEQ ID NO:2), a nuclear localization signal 2 (NLS2) having substantial sequence identity to the NLS2 of wild-type D1L3 protein (amino acids 296 to 304 of SEQ ID NO:2).

[0063] In various aspects, the recombinant D1 variant comprises one or more building block substitutions selected from: substitution of 1-22 (MRGMKLLGALLALAALLQGAVS) from SEQ ID NO:1 with 1-20 (MSRELAPLLL LLLSIHSALA) from SEQ ID NO:2; substitution of 23-27 (LKIAA) from SEQ ID NO:1 with 21-25 (MRICS) from SEQ ID NO:2; substitution of 30-32 (IQT) from SEQ ID NO: 1 with 28-30 (VRS) from SEQ ID NO:2; substitution of 35-36 (ET) from SEQ ID NO:1 with 33-34 (ES) from SEQ ID NO:2; substitution of 38-47 (MSNATLVSYI) from SEQ ID NO:1 with 36-45 (QEDKNAMDVI) from SEQ ID NO:2; substitution of 49-52 (QILS) from SEQ ID NO:1 with 47-50 (KVIK) from SEQ ID NO:2; substitution of 54 (Y) from SEQ ID NO:1 with 52 (C) from SEQ ID NO:2; substitution of 56-60 (IALVQ) from SEQ ID NO:1 with 54-58 (III,VM)

from SEQ ID NO:2; substitution of 62-63 (VR) from SEQ ID NO:1 with 60-61 (IK) from SEQ ID NO:2; substitution of 65-72 (SHLTAVGK) from SEQ ID NO:1 with 63-70 (SNNRICPI) from SEQ ID NO:2; substitution of 74-76 (LDN) from SEQ ID NO:1 with 72-74 (MEK) from SEQ ID NO:2; substitution of 79-84 (QDAPDT) from SEQ ID NO:1 with 77-84 (RNSRRGIT) from SEQ ID NO:2; substitution of 86 (H) from SEQ ID NO:1 with 86 (N) from SEQ ID NO:2; substitution of 88-89 (VV) from SEQ ID NO:1 with 88-89 (VI) from SEQ ID NO:2; substitution of 91-92 (EP) from SEQ ID NO:1 with 91-92 (SR) from SEQ ID NO:2; substitution of 96-97 (NS) from SEQ ID NO:1 with 96-97 (NT) from SEQ ID NO:2; substitution of 101 (R) from SEQ ID NO:1 with 101 (Q) from SEQ ID NO:2; substitution of 103 (L) from SEQ ID NO: 1 with 103 (A) from SEQ ID NO:2; substitution of 105 (V) from SEQ ID NO:1 with 105 (L) from SEQ ID NO:2; substitution of 107-110 (RPDQ) from SEQ ID NO:1 with 107-110 (KEKL) from SEQ ID NO:2; substitution of 113-116 (AVDS) from SEQ ID NO:1 with 113-116 (VKRS) from SEQ ID NO:2; substitution of 118 (Y) from SEQ ID NO:1 with 118 (H) from SEQ ID NO:2; substitution of 120 (D) from SEQ ID NO:1 with 120 (H) from SEQ ID NO:2; substitution of 122-128 (GCEPCGN) from SEQ ID NO:1 with 122-127 (YQDGDA) from SEQ ID NO:2; substitution of 130 (TF) from SEQ ID NO:1 with 129 (V) from SEQ ID NO:2; substitution of 132 (N) from SEQ ID NO:1 with 131 (S) from SEQ ID NO:2; substitution of 136-137 (AI) from SEQ ID NO:1 with 135-136 (FV) from SEQ ID NO:2; substitution of 139 (R) from SEQ ID NO:1 with 138 (W) from SEQ ID NO:2; substitution of 141-144 (FSRF) from SEQ ID NO:1 with 140-143 (QSPH) from SEQ ID NO:2; substitution of 146-149 (EVRE) from SEQ ID NO:1 with 145-148 (AVKD) from SEQ ID NO:2; substitution of 151 (A) from SEQ ID NO:1 with 150 (V) from SEQ ID NO:2; substitution of 153 (V) from SEQ ID NO:1 with 152 (I) from SEQ ID NO:2; substitution of 157-158 (AA) from SEQ ID NO:1 with 156-157 (TT) from SEQ ID NO:2; substitution of 160-162 (GDA) from SEQ ID NO:1 with 159-161 (ETS) from SEQ ID NO:2; substitution of 164 (A) from SEQ ID NO:1 with 163 (K) from SEQ ID NO:2; substitution of 168 (A) from SEQ ID NO:1 with 167 (E) from SEQ ID NO:2; substitution of 170-171 (YD) from SEQ ID NO:1 with 169-171 (VE) from SEQ ID NO:2; substitution of 174 (L) from SEQ ID NO:1 with 173 (T) from SEQ ID NO:2; substitution of 177-179 (QEK) from SEQ ID NO:1 with 176-178 (KHR) from SEQ ID NO:2; substitution of 181-182 (GL) from SEQ ID NO:1 with 180-181 (KA) from SEQ ID NO:2; substitution of 184-187 (DVML) from SEQ ID NO:1 with 183-187 (NFIF) from SEQ ID NO:2; substitution of 199-202 (RPSQ) from SEQ ID NO:1 with 198-201 (PKKA) from SEQ ID NO:2; substitution of 204-205 (SS) from SEQ ID NO:1 with 203-204 (KN) from SEQ ID NO:2; substitution of 209 (W) from SEQ ID NO:1 with 208 (R) from SEQ ID NO:2; substitution of 211 (S) from SEQ ID NO:1 with 210 (D) from SEQ ID NO:2; substitution of 213 (T) from SEQ ID NO:1 with 212 (R) from SEQ ID NO:2; substitution of 215 (Q) from SEQ ID NO:1 with 214 (V) from SEQ ID NO:2; substitution of 219 (P) from SEQ ID NO:1 with 218 (G) from SEQ ID NO:2; substitution of 221-222 (SA) from SEQ ID NO:1 with 220-221 (QE) from SEQ ID NO:2; substitution of 226-228 (ATP) from SEQ ID NO:1 with 225-228 (VKKS) from SEQ ID NO:2; substitution of 230 (H) from SEQ ID NO:1 with 230(N) from SEQ ID NO:2; substitution of 238-239 (VA) from SEQ ID NO:1 with 238-239 (LR) from SEQ ID NO:2; substitution of 241-246 (MLLRGA) from SEQ ID NO:1 with 241-246 (QEIVSS) from SEQ ID NO:2; substitution of 250-251 (D) from SEQ ID NO:1 with 250-251 (K) from SEQ ID NO:2; substitution of 252-254 (ALP) from SEQ ID NO:1 with 252-254 (NSV) from SEQ ID NO:2; substitution of 256 (N) from SEQ ID NO:1 with 256 (D) from SEQ ID NO:2; substitution of 259-260 (AA) from SEQ ID NO:1 with 259-260 (KA) from SEQ ID NO:2; substitution of 262 (G) from SEQ ID NO:1 with 262 (K)

from SEQ ID NO:2; substitution of 264-267 (SDQ) from SEQ ID NO:1 with 264-267 (TEEE) from SEQ ID NO:2; substitution of 269-271 (QAI) from SEQ ID NO:1 with 269-271 (LDV) from SEQ ID NO:2; substitution of 275 (Y) from SEQ ID NO:1 with 275 (F) from SEQ ID NO:2; substitution of 279-280 (VM) from SEQ ID NO:1 with 279-280 (FK) from SEQ ID NO:2; and substitution of 282 (K) from SEQ ID NO:1 with 282-305 (QSSRAFTNSKKS SVTLRKTKSKRS) from SEQ ID NO:2.

[0064] These building block substitutions from D1L3 results in variants of D1, which feature one or more of the following mutations: 1M_S22delinsMSRELAPLLLLLSIHSALA, L23_A27delinsMRICS, I30_T32delinsVRS, E35_T36delinsES, M38_I47delinsQEDKNAMDVI, Q49_S52delinsKVIK, Y54C, I56_Q60delinsIILVM, V62_R63delinsIK, S65_K72delinsSNNRICPI, L74_N76delinsMEK, Q79_T84delinsRNSRRGIT, H86N, V88_V89delinsVI, E91_P92delinsSR, N96_S97delinsNT, R101Q, L103A, V105L, R107_Q110delinsKEKL, A113_S116delinsVKRS, Y118H, D120H, G122_N128delinsYQDGDA, T130S, N132S, A136_I137delinsFV, R139W, F141_F144delinsQSPH, E146_E149delinsAVKD, A151V, V153I, A157_A158delinsTT, G160_A162delinsETS, A164K, A168E, Y170_D171delinsVE, L174T, Q177_K179delinsKHR, G181_L182delinsKA, D184_L187delinsNFIF, R199_Q202delinsPKKA, S204_S205delinsKN, W209R, S211D, T213R, Q215V, P219G, S221_A222delinsQE, A226_P228delinsVKKS, H230N, V238_A239delinsLR, M241_A246delinsQEIVSS, D250K, A252_P254delinsNSV, N256D, A259_A260delinsKA, G262K, S264_L267delinsTEEE, Q269_I271delinsLDV, Y275F, V279_M280delinsFK, K282delinsQSSRAFTNSKKS SVTLRKTKSKRS.

[0065] A recombinant D1 variant may comprise the following modifications with respect to SEQ ID NO:1: Y275F, V279_M280delinsFK, and K282delinsQSSRAFTNSKKS SVTLRKTKSKRS (SEQ ID NO: 16). Such variants comprise a C-terminal tail, and display higher affinity for DNA-lipid complexes.

[0066] A recombinant D1 variant may comprise the following amino acid modifications with respect to SEQ ID NO:1: Q79_T84delinsRNSRRGIT (e.g., SEQ ID NO: 10); H86N, and/or, V88_V89delinsVI, and/or E91_P92delinsSR (e.g. SEQ ID NO: 11, which includes all three); A136_I137delinsFV (e.g., SEQ ID NO: 12); R199_Q202delinsPKKA, and/or S204_S205delinsKN, and/or W209R, and/or S211D, and/or T213R, and/or Q215V, and/or P219G, and/or S221_A222delinsQE (e.g., SEQ ID NO: 13, which includes all eight); and A226_P228delinsVKKS (e.g., SEQ ID NO: 14).

[0067] Where the D1 variant comprises the substitution H86N, the variant comprises at least one addition modification, such as a building block substitution or other point mutation described herein (such as the addition of a cationic amino acid or substitution that results in resistance to actin).

[0068] The D1 variant may comprise at least two adjacent, contiguous, building block substitutions that result in mutations with respect to SEQ ID NO: 1 of: H86N/V88_V89delinsVI (e.g., SEQ ID NO: 11); R199_Q202delinsPKKA/S204_S205delinsKN/W209R/S211D/T213R/Q215V/P219G/S221_A22

2delinsQE (e.g., SEQ ID NO: 13);
Y275F/V279_M280delinsFK/K282delinsQSSRAFTNSKKSVTLRKKT KSKRS (SEQ ID NO: 16).

[0069] The disclosure provides a DNase enzyme comprising an amino acid sequence that is at least 80% identical to the amino acid sequence of the enzyme defined by SEQ ID NO:1, and comprising at least one building block substitution from the amino acid sequence of SEQ ID NO:2, where the DNase enzyme has increased chromatin-degrading activity as compared to the DNase enzyme of SEQ ID NO:1. The enzyme may comprise at least two adjacent building block substitutions from SEQ ID NO:2. Building block substitutions between D1 (SEQ ID NO:1) and D1L3 (SEQ ID NO:2) are illustrated in **FIG. 31**.

[0070] The enzyme may comprise the following modification with respect to SEQ ID NO:1: Q79_T84delinsRNSRRGIT, and optionally comprises the amino acid sequence of SEQ ID NO: 10.

[0071] The enzyme may comprise the following modification with respect to SEQ ID NO:1: H86N/V88_V89delinsVI/E91_P92delinsSR, and optionally comprises the amino acid sequence of SEQ ID NO: 11.

[0072] The enzyme may comprise the following modifications with respect to SEQ ID NO:1: A136_I137delinsFV, and optionally comprises the amino acid sequence of SEQ ID NO: 12.

[0073] The enzyme may comprise two or more of the following modifications with respect to SEQ ID NO:1: R199_Q202delinsPKKA, and/or S204_S205delinsKN, and/or W209R, and/or S211D, and/or T213R, and/or Q215V, and/or P219G, and/or S221_A222delinsQE, and optionally comprises the amino acid sequence of SEQ ID NO: 13 (which comprises all eight).

[0074] The enzyme may comprise the following modifications with respect to SEQ ID NO:1: A226_P228delinsVKKS, and optionally comprises the amino acid sequence of SEQ ID NO: 14.

[0075] The enzyme may comprise two or more of the following modifications with respect to SEQ ID NO:1: Q79_T84delinsRNSRRGIT, and/or H86N, and/or V88_V89delinsVI, and/or E91_P92delinsSR, and/or A136_I137delinsFV, and/or R199_Q202delinsPKKA, and/or S204_S205delinsKN, and/or W209R, and/or S211D, and/or T213R, and/or Q215V, and/or P219G, and/or S221_A222delinsQE, and/or A226_P228delinsVKKS, and optionally comprises the amino acid sequence of SEQ ID NO: 15.

[0076] The enzyme may comprise two or more of the following modification with respect to SEQ ID NO:1: Y275F, and/or V279_M280delinsFK, and/or K282delinsQSSRAFTNSKKSVTLRKKT KSKRS, and optionally comprises the amino acid sequence of SEQ ID NO: 16.

[0077] Exemplary D1 variants include those comprising the amino acid sequence of SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID

NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, and SEQ ID NO:16. The disclosure includes derivatives having from 1 to 10 (e.g., 1 to 5) amino acid insertions, deletions, or substitutions with respect to any one of SEQ ID NOS: 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16. The D1 variant may comprise one or more additional block amino acid substitutions from a homologous DNase (e.g., from D1L3). Such block substitutions may replace from 2 to 20 amino acids, such as from 3 to 15 amino acids, or from 3 to 10 amino acids of the D1 variant, with building block amino acids from the homologous DNase (e.g., D1L3, SEQ ID NO:2). Building block substitutions between D1 and D1L3 are illustrated in **FIG. 31**. The DNase enzyme may comprise an amino acid sequence that is at least 95% or at least 97% identical to SEQ ID NO:15.

[0078] The enzyme may exhibit one or more properties selected from: similar or higher affinity for protein-free DNA compared to the enzyme of SEQ ID NO:1, a higher chromatin degradation activity compared to the enzyme of SEQ ID NO:1, an enzymatic activity that is resistant to inhibition by actin compared to the enzyme of SEQ ID NO:1, and a combination thereof.

[0079] The D1 variant may comprise an N-terminal or C-terminal fusion to a half-life extending moiety, such as albumin, transferrin, an Fc, or elastin-like protein, as described for D1L3. The D1 enzyme may be dimerized by an immunoglobulin hinge region.

[0080] The D1 variant may exhibit one or more properties selected from: the same or higher protein-free DNA (naked DNA) degradation activity, higher chromosomal DNA (chromatin) degradation activity, similar or improved protease resistance, actin resistance, similar or improved penetration from blood into urine and/or bile, and higher production levels in vitro expression systems (e.g. Chinese hamster ovary cells and/or *Pichia pastoris*), and a combination thereof.

[0081] The DNase enzyme may comprise an amino acid sequence that is at least 80% identical to the amino acid sequence of the enzyme defined by SEQ ID NO:1, and comprises the substitution, insertion, or addition of one or more arginine and/or lysine residues, wherein the DNase enzyme has increased chromatin-degrading activity as compared to the enzyme of SEQ ID NO:1. The one or more arginine and/or lysine residues may be amino acid substitutions selected from substitutions at positions corresponding to positions 31, 41, 49, 52, 68, 76, 79, 81, 92, 109, 114, 115, 164, 177, 181, 200, 201, 204, 209, 213, 227, 239, 250, 259, 262, and 280 of SEQ ID NO:1. Exemplary modifications at these positions include Q31R, A41K, Q49R, S52K, T68R, N76K, Q79R, D80_A81insR, A81R, P92R, D109K, V114K, D115R, A164K, Q177K, G181K, P200K, S201K, S204R, W209R, T213R, A226_T227insK, T227K, A239R, D250K, A259K, G262K, and K280M. The enzyme may comprise substitution and/or insertion of at least two, or at least three, or at least four, or at least five, or at least six arginine and/or lysine residues. The DNase enzyme may comprise an arginine or lysine at one or more positions corresponding to positions 31, 49, 76, 79, 92, 109, 164, 177, 200, 201, 204, 209, 213, and 262 of SEQ ID NO:1, which are optionally selected from Q31R, Q49R, N76K, Q79R, P92R, D109K, A164K, Q177K, P200K, S201K, S204R, W209R, T213R, W209R, and G262K. These positions/substitutions can independently provide better activity, including activity for

degrading chromatin.

[0082] If the enzyme comprises a substitution of Q31R or T227K, the enzyme comprises at least two amino acid modifications selected from Q31R, A41K, Q49R, S52K, T68R, N76K, Q79R, D80_A81insR, A81R, P92R, D109K, V114K, D115R, A164K, Q177K, G181K, P200K, S201K, S204R, W209R, T213R, A226_T227insK, T227K, A239R, D250K, A259K, G262K, and K280M. The amino acid substitutions may include Q31R and T227K with respect to SEQ ID NO:1.

[0083] The DNase enzyme may have an arginine or lysine at positions corresponding to positions 31, 49, 79, and 209 of SEQ ID NO:1, and may optionally comprise the following amino acid substitutions: Q31R, Q49K, Q79R, and W209R. These positions harbor an arginine or lysine in murine D1, but not wild-type human D1, and can transfer chromatin-degrading activity to human D1.

[0084] The DNase enzyme may have an arginine or lysine at positions corresponding to positions 31, 49, 79, 177, 209, and 262 of SEQ ID NO:1, and may optionally comprise the amino acid substitutions Q31R, Q49K, Q79R, A136F, Q177K, W209R, G262K. Four of these positions harbor an arginine or lysine in rat D1, but not wild-type human D1, and together can transfer chromatin degrading activity to human D1.

[0085] The DNase enzyme may have a lysine or arginine residue at positions corresponding to positions 31 and 227 of SEQ ID NO:1.

[0086] The DNase enzyme may have a lysine or arginine at the position corresponding to position 164 of SEQ ID NO:1, which is optionally lysine (A164K).

[0087] The engineered D1 protein may contain one or more amino acid modifications (e.g., substitutions, additions, and deletions) in the actin binding site (e.g. substitution of A136, such as A136F), and thereby hindering or preventing actin binding. As such, the engineered D1 protein can be actin-resistant or substantially actin-resistant. See Ulmer et al., PNAS USA Vol. 93, pp 8225-8229 (1996). Thus, the DNase enzyme may comprise an amino acid substitution at position 136 with respect to SEQ ID NO:1, and which is optionally A136F. Mutations to D1 providing for actin resistance, such as mutations at position 136 (e.g., A136F, or substitution of another bulky, hydrophobic, cyclic or aromatic amino acid at this position), may be combined with any other mutation described herein, including addition of cationic residues or building block substitutions.

[0088] The D1 variant may be resistant to actin-inhibition and capable of degrading chromatin. For example, the D1 variant may comprise the following amino acid substitutions: Q31R, T227K, and A136F with respect to SEQ ID NO:1. The DNase enzyme may comprise the following amino acid substitutions: A164K and A136F.

[0089] The DNase enzyme may comprise a glycosylation consensus sequence that comprises

a N40S substitution and/or an N128S substitution with respect to SEQ ID NO:1. For example, the enzyme comprises the amino acid substitutions: Q31R, N40S, A136F, N128S, and T227K with respect to SEQ ID NO:1.

[0090] The D1 variant may contain one or more amino acid modifications (e.g., substitutions, additions, insertions, and deletions) in the glycosylation sites, and thereby increasing affinity towards lipid encapsulated DNA. As such, the engineered D1 protein can degrade lipid encapsulated DNA in transfection reactions. See Wilber et al., Mol Ther 6, 35-42 (2002).

[0091] The D1 variant may be glycosylated, or may be non-glycosylated particularly where long circulating half-life is not needed. The amino acid alterations described herein can be made in the context of glycosylated or non-glycosylated variants.

[0092] The D1 variant may be resistant to actin-inhibition and capable of degrading chromatin within lipid vesicles. Exemplary D1 variants comprise one or more amino acid modifications selected from: Q31R, T227K, A136F, N40D or N40A, and N128D or N128S, with amino acids numbered according to SEQ ID NO:1. Such variants may lack glycosylation sites, which are beneficial from a manufacturing perspective.

[0093] The D1 variant may be resistant to actin-inhibition and capable of degrading chromatin within lipid vesicles, and comprises amino acid modifications Q31R, T227K, A136F, N40D, and N128D with respect to SEQ ID NO:1.

[0094] Other modifications to D1 can be as described in Pan et al. J Biol Chem, 1998, 273, 11701-11708 and in US 2014/0199329.

[0095] The engineered D1 variant can have an increased or extended serum half-life compared to the wild-type D1 protein (SEQ ID NO:1). The engineered D1 may comprise an Fc-fusion domain (e.g. a hinge region, and/or one or more CH2 domains and CH3 domains of an immunoglobulin). The engineered D1 may be fused to albumin, e.g., human albumin or a fragment thereof. Albumin can be joined to at the N-terminus or the C-terminus of the engineered D1 enzyme, optionally through an amino acid linker.

[0096] The disclosure provides pharmaceutical compositions comprising the recombinant D1L3 and/or D1 variants, or polynucleotides or vectors encoding the same, and a pharmaceutically acceptable carrier. The recombinant D1L3 protein variant may be formulated for parenteral or pulmonary administration, for intravenous, intraarterial, intraperitoneal, intraarticular, intramuscular, topical, or subcutaneous administration, or other route described herein. The composition may comprise both D1L3 and D1, which are each optionally engineered variants described herein. The recombinant D1L3 variant and the recombinant D1 variant may be formulated for parenteral or pulmonary administration.

[0097] The term "pharmaceutically acceptable carrier" includes, but is not limited to, any carrier that does not interfere with the effectiveness of the biological activity of the ingredients

and that is not toxic to the patient to whom it is administered. Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Such carriers can be formulated by conventional methods and can be administered to the subject at a suitable dose. Preferably, the compositions are sterile. These compositions may also contain adjuvants such as preservative, emulsifying agents and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents.

[0098] Routes of administration include, for example: intradermal, intramuscular, intraperitoneal, intraarticular, intravenous, subcutaneous, intraarterial, oral, sublingual, pulmonary, or transdermal. The administering may be effected orally or by parenteral injection or infusion. Alternatively, the route of administration may be topical, including as eye drops or mouth wash.

[0099] The disclosure also provides a method of making a pharmaceutical composition for reducing or preventing neutrophil extracellular trap (NET) accumulation. The method may employ a genetically modified mouse deficient in D1 and D1L3 activity, and heterologous expression of a G-CSF polynucleotide (e.g., in hepatocyte cells) or induction of a sustained endogenous G-CSF expression (e.g., via repetitive administration of microbial compounds). This mouse model accumulates NETs and rapidly develops NET-related vascular occlusions. the method may comprise administering a candidate NET inhibitor or candidate DNase enzyme (including a D1L3 or D1 variant in accordance with this disclosure) to the genetically-modified mouse, and selecting a NET inhibitor or DNase enzyme that reduces the accumulation of NETs. The selected inhibitor or enzyme is formulated (as described) for administration to a human patient.

[0100] One skilled in the art recognizes standard methods for generating double knockout *Dnase1^{-/-}Dnase1/3^{-/-}* mice. Detailed descriptions can be found in, for example, European Application No. EP 17152528.0.

[0101] Method of treatments, disclosed herein are not part of the invention.

The disclosure provides a method for treating a subject in need of neutrophil extracellular trap (NET) degradation. The method comprises administering a therapeutically effective amount of D1L3 and/or D1 (which may include variants described herein, or one or more wild type enzymes, including for D1L3 and D1 combination therapy) according to this disclosure. The D1L3 or D1 may be administered as pharmaceutical compositions comprising the recombinant protein, or in some embodiments comprising the encoding DNA or RNA or vectors comprising the same.

[0102] The method may comprise administering a therapeutically effective amount of a D1L3 or variant thereof and a D1 or variant thereof. The D1L3 or variant thereof, and the D1 or variant thereof, may be administered in a single pharmaceutical composition or separate pharmaceutical compositions. The ratio of D1L3 or variant thereof to D1 or variant thereof may

be in the range of 100:1 to 1:100 by mass, or 10: 1 to 1:10 by mass, and is optionally about 1:1.

[0103] The subject may exhibit impaired NET degradation and/or exhibits pathological NET accumulation. The subject may have a chronic or acute inflammatory disorder or an acute or chronic infection.

[0104] The disclosed method pertains to the treatment of diseases or conditions characterized by the presence or accumulation of NETs. Such diseases or conditions include, but are not limited to, diseases associated with chronic neutrophilia (e.g., an increase in the number of neutrophils), neutrophil aggregation and leukostasis, thrombosis and vascular occlusion (e.g. sickle cell disease), ischemia-reperfusion injury (e.g. midgut volvulus, testicular torsion, limb ischemia reperfusion, vital organ ischemia-reperfusion, organ transplantation), surgical and traumatic tissue injury, an acute or chronic inflammatory reaction or disease, an autoimmune disease (e.g. systemic lupus erythematosus (SLE), lupus nephritis, rheumatoid arthritis, vasculitis, systemic sclerosis), cardiovascular disease (e.g., myocardial infarction, stroke, atherosclerosis, venous thromboembolism, including thrombolytic therapy), metabolic disease (e.g., diabetes), systemic inflammation (e.g., systemic inflammatory response syndrome (SIRS), sepsis, septic shock, disseminated intravascular coagulation (DIC), and thrombotic microangiopathy (TMA)), inflammatory diseases of the respiratory tract (e.g. cystic fibrosis, chronic obstructive pulmonary disease (COPD), acute lung injury (ALI), smoke induced lung injury, transfusion induced lung injury (TRALI), acute respiratory distress syndrome (ARDS), and asthma, atelectasis, bronchitis, empyema), renal inflammatory diseases (acute and chronic kidney diseases, including acute kidney injury (AKI) and chronic kidney disease (CKD), inflammatory diseases related to transplanted tissue (e.g. graft-versus-host disease) and cancer (e.g. leukemia, tumor metastasis, and solid tumors).

[0105] The subject may have or may be at risk of NETs occluding ductural systems. The herein disclosed enzymes can be administered to a subject to treat pancreatitis, cholangitis, conjunctivitis, mastitis, dry eye disease, obstructions of vas deferens, or renal diseases. In such situation, the method may comprise administering a therapeutically effective amount of a D1L3 or variant thereof, and/or a therapeutically effective amount of D1 or a variant thereof. For example, the subject may be administered an enzyme comprising an amino acid sequence having 80% or more sequence identity with the enzyme defined by SEQ ID NO:2, and an enzyme comprising an amino acid sequence having 80% or more sequence identity with the enzyme defined by SEQ ID NO:1. The enzymes may comprise the amino acid sequence of the enzyme defined by SEQ ID NO:2 and the amino acid sequence of the enzyme defined by SEQ ID NO:1.

[0106] The ductal system may be bile duct, tear duct, lactiferous duct, cystic duct, hepatic duct, ejaculatory duct, parotid duct, submandibular duct, major sublingual duct, Bartholin's duct, cerebral aqueduct, pancreas, mammary gland, vas deferens, ureter, urinary bladder, gallbladder, and liver. For example, the subject may have pancreatitis, cholangitis (e.g., primary sclerosing cholangitis), conjunctivitis, mastitis, dry eye disease, an obstruction of the

vas deferens, or renal disease. The DNase enzyme may be administered by intravenous, intraarterial, or intraperitoneal administration. The DNase when applied, for example, intravenously, will be present in enzymatically active form in various ductal systems, such as in bile fluid.

[0107] The subject may have or may be at risk of NETs accumulating on endothelial surfaces (e.g. surgical adhesions), the skin (e.g. wounds/scarring), or in synovial joints (e.g. gout and arthritis). The herein disclosed enzymes can be administered to a subject to treat a condition characterized by an accumulation of NETs on an endothelial surface such as, but not limited to, a surgical adhesion. The herein disclosed enzymes can be administered to a subject to treat a condition characterized by an accumulation of NETs on skin such as, but not limited to, wounds and scars. The herein disclosed enzymes can be administered to a subject to treat a condition characterized by an accumulation of NETs in a synovial joint such as, but not limited to, gout and arthritis.

[0108] The subject may have or may be at risk of a vascular occlusion comprising NETs. In such situation, the method comprises administering a therapeutically effective amount of a D1L3 or variant thereof, and/or a therapeutically effective amount of D1 or a variant thereof. For example, the subject may be administered an enzyme comprising an amino acid sequence having 80% or more sequence identity with the enzyme defined by SEQ ID NO:2. The enzyme may comprise the amino acid sequence of the enzyme defined by SEQ ID NO:2. The subject may be further administered an enzyme comprising an amino acid sequence having 80% or more sequence identity with the enzyme defined by SEQ ID NO:1, such as the enzyme defined by SEQ ID NO:1.

[0109] The subject may have a condition relating to NETs as described in WO 2016/118476 and US 9,642,822.

[0110] The subject may have a disease that is or has been treated with wild-type DNases, including D1 and streptodornase. Such diseases or conditions include thrombosis, stroke, sepsis, lung injury, atherosclerosis, viral infection, sickle cell disease, myocardial infarction, ear infection, wound healing, liver injury, endocarditis, liver infection, pancreatitis, primary graft dysfunction, limb ischemia reperfusion, kidney injury, blood clotting, alum-induced inflammation, hepatorenal injury, pleural exudations, hemotorax, intrabiliary blood clots, post pneumatic anemia, ulcers, otolaryngological conditions, oral infections, minor injuries, sinusitis, post-operative rhinoplasties, infertility, bladder catheter, wound cleaning, skin reaction test, pneumococcal meningitis, gout, leg ulcers, cystic fibrosis, Kartegener's syndrome, asthma, lobar atelectasis, chronic bronchitis, bronchiectasis, lupus, primary ciliary dyskinesia, bronchiolitis, empyema, pleural infections, cancer, dry eyes disease, lower respiratory tract infections, chronic hematomas, Alzheimer's disease, and obstructive pulmonary disease.

[0111] The disclosure further pertains to the treatment of diseases or conditions characterized by deficiency of D1, deficiency of D1L3, and deficiency of D1 and D1L3. In some cases, the subject has a mutation in the *Dnase1* and/or the *Dnase1/3* gene. Such subjects can also have

an autoimmune disease (e.g., SLE, systemic sclerosis) or an inflammatory disease. In some cases, the subject has an acquired inhibitor of D1 (e.g., anti-DNase1-antibody and actin) and/or the D1L3 (e.g., anti-DNase1L3-antibody). Such subjects can also have an autoimmune disease (e.g., SLE, systemic sclerosis) or an inflammatory disease (e.g., sepsis and ischemia-reperfusion injury).

[0112] The subject may have cystic fibrosis, and the DNase composition is administered by pulmonary delivery.

[0113] For example, where the subject is at risk of, or exhibits symptoms of, NET-related vascular occlusion, the protein or composition (e.g., comprising D1L3, D1, or a variant of D1L3 or D1, including variants described herein) may be administered intravenously, intramuscularly, subcutaneously, or intraarterially.

[0114] The subject may be monitored for the NET-degrading activity of blood, plasma, or serum. For example, the subject may be monitored for about one week to about four weeks, to reduce the risk of vascular or ductal occlusion by NETs, or NET-related damage to organs. The subject may receive from one to four administrations over the course of from one week to one month (or 1-2 weeks), as needed to prevent or reduce intravascular NET accumulation (including NET-related intravascular occlusions) during an acute infection or inflammatory event. Alternatively, the subject may receive continuous infusion over the course of from one hour to one month (or 1-2 weeks). In some cases, the subject may receive at least one, e.g., 1, 2, 3, 4 or more infusions daily, for example (without limitation), in the case of critical illness such as sepsis, stroke, or myocardial infarction.

[0115] The treatment of disease or conditions relating to NETs may involve administration of a D1 variant, a D1L3 variant, or a combination comprising a D1 variant and a D1L3 variant. Two or more of variants of the D1 protein described herein may be administered. Alternatively, two or more of variants of the D1L3 protein described herein may administered.

[0116] The present D1 protein variants and D1L3 protein variants can be administered as a prophylactic therapy, such as before the onset of one or more symptoms of the disease or condition.

[0117] As used herein, "treatment" or "treating" or "treated" refers to therapeutic treatment wherein the object is to slow (lessen) an undesired physiological condition, disorder or disease, or to obtain beneficial or desired clinical results. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent of the condition, disorder or disease; stabilization (i.e., not worsening) of the state of the condition, disorder or disease; delay in onset or slowing of the progression of the condition, disorder or disease; amelioration of the condition, disorder or disease state; and remission (whether partial or total), whether detectable or undetectable, or enhancement or improvement of the condition, disorder or disease. Treatment includes eliciting a clinically significant response without excessive levels of side effects. "Treatment" or "treating" or "treated" refers to prophylactic measures, wherein

the object is to delay onset of or reduce severity of an undesired physiological condition, disorder or disease, such as, for example is a person who is predisposed to a disease (e.g., an individual who carries a genetic marker for a disease such as lupus).

[0118] Therapy is used to provide a positive therapeutic response with respect to a disease or condition. By "positive therapeutic response" is intended an improvement in the disease or condition, and/or an improvement in the symptoms associated with the disease or condition. For example, a positive therapeutic response would refer to one or more of the following improvements in the disease: (1) a reduction in neutrophil extracellular traps; (2) a reduction in intravascular clots; (3) a reduction in ductural clots; (4) a reduction of accumulation of NETs in synovial joints; (5) an increase in the degradation of neutrophil extracellular traps; (6) an increase in degradation of extracellular protein-free DNA; (7) an increase in degradation of extracellular chromosomal DNA or protein-bound DNA; (8) an increase in the degradation of neutrophil extracellular traps in the presence of autoantibodies; (9) a reduction in tissue/organ inflammation; (10) a reduction in tissue/organ injury; (11) a reduction in tissue/organ atrophy; (12) an increased patient survival rate; and (13) some relief from one or more symptoms associated with the disease or condition.

[0119] Positive therapeutic responses in any given disease or condition can be determined by standardized response criteria specific to that disease or condition. Clinical response can be assessed for changes in vascular occlusion and the presence of NETs using screening techniques such as magnetic resonance imaging (MRI) scan, ultrasound scan, histology, and counting of NETs in the circulation. In addition to these positive therapeutic responses, the subject undergoing therapy may experience the beneficial effect of an improvement in the symptoms associated with the disease.

[0120] Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. Parenteral compositions may be formulated in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0121] The disclosure further provides an expression vector, comprising a nucleic acid encoding a wild type or D1L3 or D1 variant described herein, which can find use for genetic therapy. Genetic therapy can be useful for chronic illness such as a genetic deficiency of D1L3 and/or D1 (such as a deleterious mutation). The expression vector may comprise DNA or RNA. The expression vector may be a mammalian expression vector.

[0122] Expression vectors may comprise a nucleic acid encoding the protein variants operably linked to an expression control region, or complement thereof, that is functional in a host cell

(e.g., prokaryotic, eukaryotic, or mammalian cell). The expression control region is capable of driving expression of the operably linked blocking and/or stimulating agent encoding nucleic acid such that the blocking and/or stimulating agent is produced in a human cell transformed with the expression vector.

[0123] Expression control regions are regulatory polynucleotides (sometimes referred to herein as elements), such as promoters and enhancers, that influence expression of an operably linked nucleic acid. An expression control region of an expression vector disclosed herein is capable of expressing operably linked encoding nucleic acid in a human cell. The expression control region may confer regulatable expression to an operably linked nucleic acid. A signal (sometimes referred to as a stimulus) can increase or decrease expression of a nucleic acid operably linked to such an expression control region. Such expression control regions that increase expression in response to a signal are often referred to as inducible. Such expression control regions that decrease expression in response to a signal are often referred to as repressible. Typically, the amount of increase or decrease conferred by such elements is proportional to the amount of signal present; the greater the amount of signal, the greater the increase or decrease in expression.

[0124] Expression systems functional in human cells are well known in the art, and include viral systems. Generally, a promoter functional in a human cell is any DNA sequence capable of binding mammalian RNA polymerase and initiating the downstream (3') transcription of a coding sequence into mRNA. A promoter will have a transcription initiating region, which is usually placed proximal to the 5' end of the coding sequence, and typically a TATA box located 25-30 base pairs upstream of the transcription initiation site. The TATA box is thought to direct RNA polymerase II to begin RNA synthesis at the correct site. A promoter will also typically contain an upstream promoter element (enhancer element), typically located within 100 to 200 base pairs upstream of the TATA box. An upstream promoter element determines the rate at which transcription is initiated and can act in either orientation. Of particular use as promoters are the promoters from mammalian viral genes, since the viral genes are often highly expressed and have a broad host range. Examples include the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter, herpes simplex virus promoter, and the CMV promoter.

[0125] Where appropriate, gene delivery agents such as, e.g., integration sequences can also be employed. Numerous integration sequences are known in the art (see, e.g., Nunes-Duby et al., *Nucleic Acids Res.* 26:391-406, 1998; Sadwoski, J. *Bacteriol.*, 165:341-357, 1986; Bestor, *Cell*, 122(3):322-325, 2005; Plasterk et al., *TIG* 15:326-332, 1999; Kootstra et al., *Ann. Rev. Pharm. Toxicol.*, 43:413-439, 2003). These include recombinases and transposases. Examples include Cre (Sternberg and Hamilton, *J. Mol. Biol.*, 150:467-486, 1981), lambda (Nash, *Nature*, 247, 543-545, 1974), Flp (Broach, et al., *Cell*, 29:227-234, 1982), R (Matsuzaki, et al., *J. Bacteriology*, 172:610-618, 1990), cpC31 (see, e.g., Groth et al., *J. Mol. Biol.* 335:667-678, 2004), sleeping beauty, transposases of the manner family (Plasterk *et al.*, *supra*), and components for integrating viruses such as AAV, retroviruses, and antiviruses having components that provide for virus integration such as the LTR sequences of retroviruses or

lentivirus and the ITR sequences of AAV (Kootstra et al., *Ann. Rev. Pharm. Toxicol.*, 43:413-439, 2003). In addition, direct and targeted genetic integration strategies may be used to insert nucleic acid sequences encoding the chimeric fusion proteins including CRISPR/CAS9, zinc finger, TALEN, and meganuclease gene-editing technologies.

Example 1: Host DNases prevent vascular occlusion by neutrophil extracellular traps

[0126] Platelet and fibrin clots occlude blood vessels in hemostasis and thrombosis. Here we report a non-canonical mechanism for vascular occlusion based on neutrophil extracellular traps (NETs), antimicrobial DNA-fibers released by activated neutrophils. We show that two DNases, DNase1 and DNase1-like 3, degrade NETs in circulation during episodes of murine inflammation. In the absence of both DNases, intravascular NETs form clots that obstruct blood vessels and cause organ damage. Vascular occlusions in patients with severe bacterial infections are associated with a defect to degrade NETs *ex vivo* and the formation of intravascular NET-clots. DNase1 and DNase1-like 3 are independently expressed and thus provide dual host protection against deleterious effects of NETs in inflammation.

[0127] Inflammation is an essential host response to control invading microbes and heal damaged tissues (1). Uncontrolled and persistent inflammation causes tissue injury in a plethora of inflammatory disorders. Neutrophils are the predominant leukocytes in acute inflammation. During infections neutrophils generate extracellular traps (NETs), lattices of DNA-filaments decorated with toxic histones and enzymes that immobilize and neutralize bacteria (2). The relevance of NETs in host defense is illustrated by the fact that extracellular DNases serve as virulence factors in several pathogenic bacteria (3, 4). However, inappropriately released NETs may harm host cells due to their cytotoxic, proinflammatory, and prothrombotic activity (5-7). Indeed, NETs are frequently associated with inflammatory or ischemic organ damage and the therapeutic infusion of DNases limits host injury in various animal models (8, 9).

[0128] How the host degrades NETs *in vivo* to limit tissue damage during episodes of inflammation is poorly understood. Earlier work has shown that DNase1 in serum digests the DNA-backbone of NETs *in vitro* (10). We analyzed serum from wild type mice by zymography and detected two enzymatically active DNases, DNase1 and DNase1-like 3 (DNase1I3) (FIG. 1A). Both enzymes are members of the DNase1 protein family, but differ in their origin and substrate affinity. DNase1 is expressed by non-hematopoietic tissues and preferentially cleaves protein-free DNA (11,12). DNase1I3, also known as DNase γ , is secreted by immune cells and targets DNA-protein-complexes, such as nucleosomes (11,13). We generated mice that lacked DNA-degrading activity in serum due to a combined deficiency of DNase1 and DNase1I3 (FIG. 1A). *In vitro* generated NETs remained intact after exposure to DNase1/DNase1I3^{-/-} sera, whereas sera from wild type, DNase1^{-/-}, and DNase1I3^{-/-} mice degraded NETs (FIG. 1B and FIG. 1C). We used a hepatocyte-specific expression plasmid in conjunction with hydrodynamic gene delivery to stably express the cDNA of DNase1 or DNase1I3 in the liver of DNase1/DNase1I3^{-/-} mice. Given that both enzymes contain a secretory protein signal

sequence (11), this approach restored the activity of DNase1 or DNase1I3 in circulation (FIG. 1D) and the capacity of sera from DNase1/DNase1I3^{-/-} mice to degrade NETs (FIG. 1E and FIG. 1F). Taken together, these data show that two independently expressed host enzymes, DNase1 and DNase1I3, degrade NETs in vitro.

[0129] To test the requirement of DNase1 and DNase1I3 for NET-degradation in vivo, we aimed to chronically stimulate wild type, DNase1^{-/-}, DNase1I3^{-/-}, and DNase1/DNase1I3^{-/-} mice with the granulocyte-colony stimulating factor (G-CSF). G-CSF triggers neutrophilia, a hallmark of acute inflammation, and stimulates a subpopulation of neutrophils to spontaneously release NETs ex vivo (14). We cloned the cDNA of CSF3, which encodes G-CSF, into the hepatocyte-specific expression plasmid. Hydrodynamic injection of wild type mice with the CSF3-plasmid resulted in chronically elevated levels of G-CSF in plasma (FIG. 5A). Consequently, the neutrophil blood count steadily increased and spontaneously formed NETs were detected in blood smears (FIG. 2A and FIG. 2B). We furthermore observed an increased number of resident neutrophils in vital organs and splenomegaly (FIG. 5B and FIG. 5C). Importantly, CSF3-injected wild type mice grew normally, did not develop organ injury, and did not show macroscopic signs of distress or abnormal behavior (FIG. 5D and FIG. 5E). Collectively, these data suggest that chronic neutrophilia with concomitant NET formation is well tolerated in wild type mice.

[0130] Next, we stably expressed CSF3 in the liver of DNase1^{-/-}, DNase1I3^{-/-}, and DNase1/DNase1I3^{-/-} mice. Mice with a single deficiency in DNase1 or DNase1I3 did not show signs of distress (not shown), whereas all mice with a combined deficiency died within 6 days after CSF3-injection (FIG. 2C). DNase1/DNase1I3^{-/-} mice receiving the control plasmid lacking CSF3 survived without showing any abnormalities (FIG. 2C, not shown). We co-expressed DNase1 or DNase1I3 with CSF3 in DNase1/DNase1I3^{-/-} mice to induce neutrophilia and NETs and simultaneously restore DNase1 or DNase1I3 in circulation. Expression of either DNase was sufficient for DNase1/DNase1I3^{-/-} mice to survive without showing any signs of distress (FIG. 2D). DNase1/DNase1I3^{-/-} mice co-expressing CSF3 with a control plasmid lacking *DNase1* and *DNase1I3* died within 5 days after gene delivery (FIG. 2D). The mortality in these mice was preceded by a rapidly progressing hypothermia, which was evidenced as a strong decrease in peripheral body temperature within 8 hours before exitus (FIG. 2E). Hypothermia was accompanied with hemolytic anemia shown by reddish plasma and urine and reduced blood hemoglobin (FIG. 2F and FIG. 2G). Abundant schistocytes in blood smears indicated that the hemolytic anemia was caused by erythrocyte fragmentation (FIG. 2H). Furthermore, we detected elevated plasma levels of lactate dehydrogenase (LDH), liver transaminases, and the renal injury markers blood urea nitrogen and creatinine, which indicated multiple organ damage (FIG. 2I, FIG. 6A, and FIG. 6B). Coexpression of *DNase1* or *DNase1I3* with CSF3 maintained the body temperature and integrity of erythrocytes and organs (FIG. 2E to FIG. 2I; FIG. 6A and FIG. 6B). In conclusion, these data indicate that either DNase1 or DNase1I3 is required to prevent host injury during chronic neutrophilia.

[0131] The histological analysis of DNase1/DNase1I3^{-/-} mice with chronic neutrophilia showed intravascular hematoxylin-positive clots with entrapped erythrocytes that fully or partially

occluded blood vessels in lungs, liver, and kidneys (FIG. 3A and B; FIG. 6C and FIG. 6D). Expression of DNase1 or DNase1I3 in circulation prevented these vascular occlusions. The hematoxylin-positive clots showed an abundant light-violet staining pattern that was sporadically speckled with a dark-violet staining of individual leukocyte nuclei, suggesting that decondensed DNA is a main clot component (FIG. 3A). Given that nuclear breakdown and unfolding of tightly packed chromatin is a hallmark of NET-formation (15), we stained the hematoxylin-positive clots for NET-markers. We observed a robust staining with fluorescent double-stranded DNA-intercalating dyes and antibodies against chromatin (FIG. 7A). The co-localization of decondensed chromatin with the neutrophil granule-derived enzyme myeloperoxidase, antimicrobial cathelicidin peptides, and the NET-surrogate marker citrullinated histones confirmed that the clots were composed of NETs (FIG. 3C, FIG. 7B and FIG. 7C). To identify components of canonical thrombi, we stained NET-clots for fibrin and von Willebrand factor (vWF), a protein stored in the secretory vesicles of platelets and the vascular endothelium. NET-clots were very heterogeneous in their vWF and fibrin content (FIG. 3D and FIG. 3E). Cross-sections of NET-clots were covered on average with 46% of vWF, while 9% of NET-clots did not stain for vWF (FIG. 3E and FIG. 3F). Fibrin was largely absent and detected in less than 23% of the occluded vessels (FIG. 3E). These data are in line with findings that NETs serve as a fibrin-independent scaffold to immobilize platelets and erythrocytes *in vitro* (6). The absence of vWF and fibrin in some clots suggests that NETs may be sufficient for vascular occlusion. To corroborate this notion, we aimed to generate NET-clots from pure neutrophils *in vitro*. We isolated neutrophils from blood and induced NET-formation, while exposing the cells to shear forces to mimic blood flow. In this setup, we observed macroscopically visible and DNase-sensitive clots (FIG. 8A), which resembled the appearance of NET-clots within the murine vasculature (FIG. 8B). Now, we depleted platelets from the circulation and pharmacologically inhibited thrombin to block fibrin formation in CSF3-expressing DNase1/DNase1I3^{-/-} mice. Unlike DNase1- or DNase1I3- expression, neither anti-thrombotic treatment could prevent mortality in these animals (FIG. 3H). Collectively, the data suggest that clots of NETs are sufficient to obstruct blood vessels during chronic neutrophilia in DNase1/DNase1I3^{-/-} mice.

[0132] The formation of NET-clots in DNase1/DNase1I3^{-/-} mice is associated with features of infection-induced thrombotic microangiopathies (TMAs) and disseminated intravascular coagulation in patients, including schistocytes, hemolytic anemia, and organ failure due to vascular occlusions. We analyzed plasma from TMA patients with hemolytic-uremic syndrome due to an infection with Shiga toxin-producing *Escherichia coli* [STEC-HUS, (16)]. Sepsis and septic shock was a frequent complication in these patients (17). NETs generated *in vitro* remained intact after exposure to patient plasma collected in the acute disease state, whereas plasma from patients in remission degraded NETs (FIG. 9A and FIG. 9B). The data indicate an acquired and temporary defect in NET-degradation, thus extending previous reports (18,19). Of note, the STEC-HUS patients were effectively treated with a regimen that included infusion of plasma from healthy donors (17), a source of DNases, which potentially restored NET-degradation.

[0133] Large aggregates of NETs are reportedly formed in the synovial fluid and pancreatic

ducts of patients (21,22), but have not yet been described in other tissues. Therefore, we aimed to identify intravascular aggregates of NETs in patients with severe inflammatory diseases. We screened lung tissue collected at autopsy from patients with acute respiratory distress syndrome and/or sepsis (FIG. 10). We detected hematoxylin-positive clots in blood vessels of two septic patients (FIG. 11A and FIG. 11C). In both cases, clots were comprised of chromatin and MPO (FIG. 11B and FIG. 11D), indicating that NETs can form intravascular clots in human sepsis.

[0134] Septicemia is a potent and rapid trigger of intravascular NET-formation in mice (7). We therefore hypothesized that a defect in NET-degradation may aggravate the disease. Indeed, we observed that mice with a combined deficiency of DNase1 and DNase1I3, but not wild type mice, were highly susceptible to low doses lipopolysaccharide and heat-killed *E. coli* (FIG. 4A). Similar to neutrophilic DNase1/DNase1I3^{-/-} mice, blood analysis of septic DNase1/DNase1I3^{-/-} mice showed hemolytic anemia and hematuria (FIG. 4B and FIG. 4C), along with increased levels of plasma LDH and schistocytes in blood smears (FIG. 4D and FIG. 4E). Furthermore, we detected abundant partially or fully occluded blood vessels in the lung (FIG. 4F and FIG. 4G). Detailed analysis of partially occluded vessels revealed clots of NETs within the vascular lumen (FIG. 4H). In fully occluded vessels the NET-clots were congested with entrapped erythrocytes and leukocytes (FIG. 4I; FIG. 12A and FIG. 12B). Importantly, hepatic expression of *DNase1* or *DNase1I3* in DNase1/DNase1I3^{-/-} mice prevented vascular occlusion and restored the wild type phenotype. Taken together, these data indicate that circulating DNase1 or DNase1I3 prevent the formation of NET-clots and host injury in septicemia.

[0135] In summary, while platelets and fibrin form hemostatic clots and pathological thrombi (23), our data introduces NET-clots as a non-canonical mechanism for vascular occlusion in inflammatory states. Similar to fibrin strands, NETs are ultra-large and stable molecules (6). At high concentrations, such as found in chronic neutrophilia or septicemia, intravascular NETs may form clots, which are sufficient in size to obstruct blood vessels and thus cause damage to erythrocytes and organs. To maintain blood and tissue integrity during inflammation, the host independently expresses DNase1 and DNase1I3 as a dual protection system against intravascular NETs. However, acquired and genetic defects in these host factors may delay the degradation of NETs and thus precipitate disease. Acquired defects may involve DNase1-inhibition by monomeric actin externalized from damaged tissue and inactivation of DNase1I3 by serum proteases (11). Mutations in DNase1 and DNase1I3 have been identified in patients and are associated with systemic lupus erythematosus (SLE), a systemic autoimmune disease (24,25). Interestingly, DNase1 and DNase1I3-deficient mice spontaneously develop SLE-like disease with age (13,26). NETs are composed of prominent autoantigens and neutrophils from SLE patients have an increased capacity to release NETs (27). A reduced clearance capacity may increase the half-life NETs and thus promote the autoimmune disease (10,27). In conclusion, defects in host DNases may bring forward the deleterious effects of NETs in a plethora of inflammatory diseases.

[0136] Consequently, DNase1- and DNase1I3-mediated NET-degradation provides attractive therapeutic opportunity for patients suffering from inflammatory conditions. To test whether

DNase1 and DNase1I3 degrade NETs by the same or via distinct mechanisms, we cross-linked proteins in NETs using fixative. Fixed NETs were efficiently degraded by DNase1I3^{-/-} sera, but were resistant to degradation by DNase1^{-/-} sera (FIG. 13A and FIG. 13B). In line with these results, exposure of NETs to recombinant DNase1 or DNase1I3 showed that both DNases degrade naive NETs (FIG. 13C and FIG. 13D), whereas fixed NETs are resistant to DNase1I3 activity (FIG. 13E and FIG. 13F). Collectively, these in vitro data suggest that DNase1 and DNase1I3 degrade NETs via distinct mechanisms.

[0137] Finally, we compared the activity of DNase1 and DNase1I3 in the circulation of mice to humans. To discriminate DNase1 and DNase1I3 activity, we generated antibodies against human DNase1 (α -hDNase1), which block DNase1 activity in plasma from normal healthy donors (NHD) (FIG. 14A). Heparin is a known inhibitor of DNase1I3 and was used to block the DNA-degrading activity by DNase1I3 (FIG. 14B). Murine plasma showed approximately 10-fold higher total DNase activity (FIG. 14C), DNase1 activity (FIG. 14D), and DNase1I3 activity (FIG. 14E) than human plasma. Furthermore, DNase1I3 in murine plasma, but not in human plasma, degrades NETs efficiently (FIG. 14F and FIG. 14G). The data suggest that the concentration of DNase1I3 in human circulation is not sufficient to degrade NETs. Therefore, enabling DNase1I3-mediated NET-degradation in patients a novel and promising therapeutic strategy for inflammatory diseases.

Materials and Methods

Patient plasma

[0138] Citrated plasma samples were obtained from patients during the STEC-HUS outbreak 2011 in Germany (16). The criteria for inclusion in our study were positive identification of STEC/enterohemorrhagic *E. coli* or bloody diarrhea, platelet count $\leq 150 \times 10^9$ /L or decreased by $\geq 25\%$ in 1 week, evidence of hemolysis (LDH above the normal limit, haptoglobin below the normal limit, or the presence of schistocytes) and acute kidney injury stage ≥ 1 .

Human tissue

[0139] Human tissues were fixed at room temperature overnight in 5% phosphate-buffered formalin. All tissues were collected at autopsy and originally submitted for diagnostic purposes to us and studied in accordance with national ethical principles. The cause of death of the patients is indicated in FIG. 10.

Mice

[0140] All mice were on a C57BL/6 genetic background. We crossed previously described DNase1^{-/-} and DNase1-like 3^{-/-} mice (26,28) to generate DNase1/DNase1I3^{-/-} mice. This strain of DNase1^{-/-} mice has been recently reported to contain an off-target mutation in the *DNase1* overlapping gene *Trap1/Hsp75*, which encodes a mitochondrial chaperone (29). We therefore included an alternative DNase1^{-/-} strain (30) to generate DNase1/DNase1I3^{-/-} mice for control experiments (FIG. 15A-FIG. 15C)

Detection of DNase1 by denaturing polyacrylamide gel electrophoresis zymography (DPZ)

[0141] We performed DPZ, as previously described with modifications (19). In brief, sodium dodecyl sulfate (SDS)-polyacrylamide gels were prepared with 4% (v/v) stacking gels without DNA and 10% (v/v) resolving gels containing 200 µg/ml of salmon testes DNA (Sigma- Aldrich, Germany). For the detection of DNase1, 0.5 µl of murine serum were mixed with 14.5 µl of water and 5 µl SDS gel-loading buffer (BioRad, Germany). The mixture was boiled for 5 minutes and loaded onto the gels. Electrophoresis was carried out at 120 V using Tris/glycine electrophoresis buffer (25 mM Tris, 192 mM glycine, 0.1% (w/v) SDS, pH 8.7). After electrophoresis, proteins were refolded by incubating the gels overnight at 37°C in a refolding buffer containing 5% (w/v) milk powder, 10 mM Tris/HCl pH 7.8, 3 mM CaCl₂, 3 mM MgCl₂, 100 U/mL penicillin, and 100 µg/mL streptomycin. The gels were transferred to a refolding buffer without milk powder and incubated for additional 24 hours at 37°C. DNA was labeled fluorescently with 1x SYBR Safe (Thermo Scientific, Germany) and fluorescent images of gels were recorded using a fluorescence scanner (Molecular Imager FX, BioRad, Germany).

Detection of DNase1I3 by DPZ

[0142] For the detection of DNase1I3, 2 µl of serum were mixed with 12 µl of water, 5 µl SDS gelloading buffer, and 1 µl of beta-mercaptoethanol (BME, Sigma-Aldrich). BME reduces the disulfide bridges of DNase1, which causes its inactivation (11). The mixture was boiled for 5 minutes and loaded onto the gels. Electrophoresis was carried out as described for DNase1. SDS was removed by washing the gels with 10 mM Tris/HCl pH 7.8 for 30 minutes at 50°C twice. The proteins were refolded by incubating the gels for 48 hours at 37°C in a refolding buffer containing 10 mM Tris/HCl pH 7.8, 1 mM BME, 100 U/mL penicillin and 100 µg/mL streptomycin. The gels were then incubated for additional 48 hours at 37°C in refolding buffer supplemented with 3 mM CaCl₂ and 3 mM MnCl₂. The addition of Mn²⁺ is required to enable efficient degradation of protein-free DNA by DNase1I3 (11). DNA was labeled and imaged as described for DNase1.

Detection of total DNase activity by single radial enzyme diffusion (SRED) assay

[0143] To measure total DNase activity, we dissolved 55 µg/ml DNA from salmon testes in a buffer containing Mn^{2+} (20 mM Tris-HCl pH 7.8, 10 mM $MnCl_2$, 2 mM $CaCl_2$, and 2x SYBR Safe). The DNA solution was heated at 50°C for 10 minutes and mixed with an equal volume of 2% ultra-pure agarose (Thermo Scientific). The mixture was poured into plastic trays and stored at room temperature until solidification. Two µl of murine serum were loaded into wells of 1.0 mm diameter. Gels were incubated for 4 hours at 37°C in a humid chamber. The DNA fluorescence of the gels was recorded with a fluorescence scanner.

In vitro NET-degradation assay

[0144] NET-degradation was analyzed as previously described (19). Purified human neutrophils in serum-free DMEM were seeded to sterile 96-well plates (Falcon, BD Technologies, Germany) coated with 0.001% polylysine (Sigma) at a concentration of 5×10^4 cells per well. To induce NET-formation, neutrophils were activated with 100 nM phorbol 12-myristate 13- acetate (PMA; Sigma-Aldrich) for 4 h at 37°C with 5% CO_2 and humidity. We added phosphate buffered saline (PBS) and stored the 96-well plates overnight at 4 °C. NETs were washed with PBS, treated for 5 minutes with PBS containing 0.5% Triton X-100, and washed with PBS again. NETs were incubated with 10% murine serum or 10% citrated human plasma supplemented with 10 µM PPACK (Santa Cruz, Heidelberg, Germany) in HBSS with divalent cations (HBSS+; Thermo Scientific). NETs were allowed to be degraded for 3 hours at 37°C with 5% CO_2 and humidity. We discarded supernatants and stopped NET-degradation by adding 2% PFA in PBS for 1 hour at room temperature. PFA was discarded and non-degraded NETs were labeled fluorescently by adding 2 µM of the fluorescent DNA dye SytoxGreen (Thermo Scientific) in PBS. Images of fluorescently stained NETs were acquired with an inverted fluorescence microscope (Axiovert 200M, Zeiss, Germany). We quantified nondegraded NETs by recording the fluorescence intensity with a plate reader (Excitation: 485 nm; Emission: 535 nm) or by measuring the area coverage of NETs in microscopy pictures using ImageJ software (NIH, USA).

Preparation of in vivo expression vectors

[0145] We used the pLIVE plasmid (Mirus Bio, USA) to express proteins in mice. The vector enables a long-lasting and hepatocyte-specific expression of proteins. We generated pLIVE plasmids with murine *DNase1*, *DNase1I3*, and *CSF3*. For murine *DNase1*, a PCR of the cDNA (Genbank Accession Number NM010061) was performed using the pair of primers DNase1-F 5'-GTCGACATGCGGTACACAGG (SEQ ID NO:27) and DNase1-R 5'-CTCGAGTCAGATTTTCTGAGTGTCA (SEQ ID NO:28) containing Sall and XhoI restriction sites. For *DNase1I3*, a PCR of the cDNA (Genbank Accession Number AF047355) was performed using the pair of primers DNase1I3-F 5'-GAAGTCCCAGGAATTCAAAGATGT (SEQ ID NO:29) and DNase1I3-R 5'-GCGTGATACCCGGGAGCGATTG (SEQ ID NO:30) containing

BamHI and SacI restriction sites. Both cDNAs were cloned using the T4 ligase (New England Biolabs, Germany) into the multi-cloning site (MCS) of the pLIVE vector, which was predigested with the appropriate enzymes. The pLIVE vector containing DNase1I3 was subjected to site-directed mutagenesis with the pair of primers mutDNase1I3-F 5'-AGTCGACTCCCGGCCACCATGTCCCTGCA (SEQ ID NO:31) and its complementary mutDNase1I3-R 5'-TGCAGGGACATGGTGGCCGGGAGTCGACT (SEQ ID NO:32) in order to match the consensus Kozak sequence and optimize protein expression. The generation of the pLIVE vector containing the cDNA of CSF3 was outsourced (Eurofins Genomics, Germany). The cDNA of CSF3 (Genbank Accession Number BC120761) was inserted in the MCS between restriction sites Sall and XhoI. Sequence of all the generated vectors was confirmed by double stranded DNA sequencing. As a control we used the parental pLIVE plasmid without additional inserts. All plasmids were purified using PureLink HiPure Plasmid Maxiprep Kit and potential contaminations of endotoxin were removed using High Capacity Endotoxin Removal Spin Columns (both Thermo Scientific).

***In vitro* expression of murine DNase1 and DNase1I3**

[0146] Plasmids containing the cDNA for DNase1 or DNase1I3, described elsewhere were transfected into Human Embryonic Kidney (HEK) cells with Lipofectamine 3000 (Thermo Scientific) in serum-free conditions with DMEM media (Thermo Scientific) supplemented with 10% KnockOut serum replacement (Thermo Scientific). After 72 hours, supernatants were collected, centrifuged at 500 xg for 10 minutes, sterile filtered, and concentrated by ultracentrifugation with 3K columns (Amicon Ultra, Millipore, Darmstadt, Germany).

***In vivo* gene expression**

[0147] The pLIVE-plasmids containing *DNase1*, *DNase1I3*, *CSF3*, or empty control plasmids were administered to mice via hydrodynamic tail vein injection. In brief, 50 µg of plasmid were diluted in 0.9% saline in a volume equivalent to 10% of the body mass of the mouse. Mice were anaesthetized with isoflurane and the plasmid solution was then injected intravenously over 5 to 8 seconds via the tail vein. In rare cases, mice did not fully recover from the injection within the first 24 hours and these animals were excluded from the study. For coexpression studies, 50 µg of the CSF3-plasmid were mixed with 50 µg of the empty control plasmid, or the plasmids containing *DNase1* and *DNase1I3*. The solution containing both plasmids was administered via hydrodynamic tail vein injection.

Chronic neutrophilia

[0148] Female mice at 4 weeks or 8-12 weeks of age were injected with 50 µg of the pLIVE-plasmid containing *CSF3* to induce G-CSF overexpression and neutrophilia. Mice were

monitored every 8 hours during the first week after injection, and daily afterwards. Temperature was measured in the perianal area by a contactless infrared-thermometer (Etekcity, Germany). For survival studies, mice were euthanized and scored as "non-surviving" if the animals showed signs of distress (no spontaneous movement, closed eyes, occasional gasping). In all cases these signs of distress were accompanied with a rapidly progressing and severe hypothermia, defined as decrease in body temperature of $\geq 4^{\circ}\text{C}$ compared to the body temperature before the plasmid injection, and with hematuria. Non-hypothermic mice did not show any signs of distress and were euthanized at the end of the experiment. For organ, blood, and urine collection, we analyzed four groups of three D1/D1L3-/- mice. Mice in each group were injected with a mixture of CSF3 with empty plasma, *DNase1*, or *DNase1/3*. Each group of mice was euthanized for biosample collection, when the first animal of the group showed signs of distress, severe hypothermia, and hematuria. This time point was defined as "exitus" and occurred within 3 to 6 days after the injection.

Platelet depletion

[0149] Mice received an intraperitoneal injection of 2 $\mu\text{g/g}$ platelet-depleting antibody or isotype control antibody (both Emfret, Germany). This treatment depletes > 95% of platelets from circulation. The treatment was started 24 hours after the injection of CSF3 and repeated every 48 hours until completion of the experiment.

Thrombin inhibition

[0150] Powdered dabigatran etexilate (Pradaxa, Boehringer Ingelheim, Germany) was mixed with normal chow powder (Altromin, Germany) at a dose of 40 mg/g. Pellets were prepared by mixing the powder with distilled water and allowed to dry at room temperature. The feeding of dabigatran was started 24 hours after the injection of CSF3. WT mice fed with the dabigatran diet for 1 day showed a 6.48 ± 1.19 -fold (Mean $\pm$ SD, N=4) increased activated partial thromboplastin time when compared to mice receiving normal chow without dabigatran (Student t-test; $P < 0.001$).

Preparation of bacteria for septicemia

[0151] *Escherichia coli* (XEN 14, Perkin Elmer) was grown overnight in lysogeny broth media containing 50 $\mu\text{g/ml}$ kanamycin. Bacteria were pelleted by centrifuging at $4000 \times g$ for 10 minutes, washed with and resuspended in PBS. Aliquots of 1.5×10^9 bacteria/ml were incubated at 70°C for 15 minutes to heat-kill the bacteria. Aliquots stored at -20°C until further use.

Septicemia

[0152] Male mice at 8-15 weeks of age received three daily intraperitoneal injections of 1 µg/g of LPS from *Salmonella enterica* serotype *thyphimurium* (Sigma-Aldrich) in 0.9% saline. Along with the third LPS injection, mice received an intravenous injection of 1.5×10^7 heat-killed *E.coli/g*. The shown survival time indicates the time after the injection of *E.coli*. Blood and organs were collected at the time of euthanasia. Insufficient biosamples were obtained from two animals (1× D1/D1L3-/- + Ctrl, 1× D1/D1L3-/- + D1L3) to perform the complete analysis shown in FIG. 4A-FIG 4I. Mice were euthanized and scored as "non-surviving" if the animals showed signs of severe distress (irresponsiveness to touch). All non-surviving mice showed hematuria and paleness of extremities. All surviving mice were euthanized and scored as "surviving" 24 hours after the intravenous injection of heat-killed *E.coli*.

Murine blood, plasma, and tissue collection

[0153] Blood was collected by submandibular puncture or by the retroorbital sinus. Blood was collected into 200 µl serum tubes (Monovette; Sarstedt, Germany) and 200 µl EDTA tubes (GK 150, KABE labortechnik, Germany). Plasma was obtained by collecting the supernatant of blood after centrifugation at $3000 \times g$ for 15 minutes. Serum was obtained by allowing the blood to clot for 1 hour at room temperature followed by a centrifugation for 15 minutes at $3000 \times g$. Serum and plasma samples were stored in aliquots at -20°C until further use. For organ analysis, mice were perfused by intracardiac infusion of PBS. Organs were collected and fixed for 24 hours in 4% PFA at 4°C. Fixed organs were embedded in paraffin.

Analysis of blood and plasma

[0154] LDH, AST, ALT, creatinine and BUN in plasma were quantified by using standardized kits (Biotron Diagnostics, CA, USA) following the manufacturer instructions. Mouse G-CSF was quantified with a Quantikine ELISA Kit (R&D, United Kingdom). Hemoglobin in EDTA blood was quantified by an automated hemocytometer (Idexx ProCyt Dx Hematology Analyzer, Netherlands). To quantify neutrophils by FACS, whole blood was incubated on ice for 15 minutes with 0.2 µg of phycoerythrin-labelled anti-mouse CD11b (M1/70, Biolegend, Germany) and 0.5 µg of Alexa Fluor 488 anti-mouse Ly6G (1AB, Biolegend, Germany). The blood was then diluted with 0.5 ml PBS and analyzed with a FACSCalibur (BD Bioscience, USA). Blood smears of EDTA-anticoagulated blood were prepared on polylysine-coated slides (Hecht Assistant, Germany). After air-drying were incubated for 1 minute in methanol supplemented with 1µM SytoxGreen on dry ice or stained with Giemsa using commercial kit (In Vitro Diagnostikum, Germany).

Immunofluorescence stainings

[0155] Paraffin-embedded sections were de-waxed, rehydrated, and subjected to antigen retrieval for 25 minutes at 100°C in citrate buffer (10 mM sodium citrate, 0.1% Tween, pH 6). Thereafter, sections were blocked for 30 minutes with 2.5% normal goat serum (Vector, United Kingdom) followed by incubation with a mouse-on-mouse blocking kit (Vector) for one hour. The sections were then incubated over night at 4°C with 2 µg/ml of the primary antibody against MPO (A0398, Agilent, Germany), CRAMP (PA-CRLP-100, Innovagen, Sweden), citrullinated histone 3 (ab5103, Abcam, United Kingdom), fibrin [clone 59D8], or the complex of histone H2A, H2B, and DNA to detect chromatin (2). Sections were incubated with anti-rabbit and anti-mouse IgG antibodies conjugated with AlexaFluor488 or AlexaFluor555 (all Thermo Scientific) for 1 hour. After washing, DNA was labeled with 1 µg/ml DAPI for 2 minutes. Autofluorescence was quenched by a 25-minute incubation with Sudan Black (0.1% in 70% ethanol), and sections were mounted with Fluoromount G (Southern Biotech, USA). Images of fluorescently labeled sections were acquired with an inverted fluorescence microscope (Axiovert 200M, Zeiss, Germany) or a confocal microscope (TCS SP5, Leica, Germany). Images of NET-clots were quantified using software (ImageJ). A vWF or fibrin staining area of less than 3% of NET-clots was considered negative.

Immunohistochemistry

[0156] After antigen retrieval, deparaffinized sections were blocked for 30 minutes with 2.5% normal horse serum (Vector). The sections were incubated over night at 4°C with 2 µg/ml of the primary antibody against neutrophil elastase (ab68672, Abcam). After washing, sections were stained using the anti-rabbit IgG-AP kit, (ImmPRESS, Vector) according to manufacturer's instruction. Sections were counterstained with hemalum (Merck, Germany) and mounted with Neo-Mount media (Merck). Images of stained sections were acquired with an inverted microscope (Axiovert 200M, Zeiss).

Generation of NET-clots in vitro

[0157] Purified human neutrophils were seeded to tissue culture plates at 3×10^7 cells / ml in DMEM supplemented with 2% BSA. The cells were activated with 0.1 µM PMA for 4 hours at 37°C, 5% CO₂ with humidity and rotating conditions (300 rpm). We used unstimulated neutrophils and neutrophils activated with PMA in the presence 10 U/ml recombinant human DNase1 (dornase alpha, Roche, Germany) as controls. NET-clots were fixed with 2% PFA overnight at 4°C. Fixed NET-clots were embedded in paraffin and sections were stained for NETs as described above.

Statistical evaluation

[0158] Data are shown as mean $\pm$ standard deviation. Statistical analysis was performed using Prism Software (GraphPad, USA) and included Student's t-test, one-way and two-way ANOVA followed by Bonferroni's multiple comparison test, and Log-rank test. Results were considered significant at $P < 0.05$.

Movies

[0159] Chronic neutrophilia is tolerated in wild type mice. A movie generated shows the wild type mice one week after the injection of a CSF3-expression plasmid (CSF3) or a control plasmid lacking CSF3 (Ctrl). Both animals show normal behavior and no signs of distress.

[0160] Chronic neutrophilia causes distress in mice lacking DNase1 and DNase1I3. The movie shows DNase1/DNase1I3^{-/-} mice co-expressing CSF3 with a control plasmid (CSF3/Ctrl), with DNase1 (CSF3/D1), or with DNase1I3 (CSF3/D1L3). Mice expressing circulatory DNases, DNase1 or DNase1I3, show normal behavior. Mice lacking circulatory DNases (CSF3/Ctrl) show signs of distress (no spontaneous movement, closed eyes, occasional gasping).

References

[0161]

1. V. Kumar, A. K. Abbas, N. Fausto, J. C. Aster, Acute and chronic inflammation. In Robbins & Cotran Pathologic Basis of Disease. (Elsevier Health Sciences, 2009), chap. 2.
2. V. Brinkmann, U. Reichard, C. Goosmann, B. Fauler, Y. Uhlemann, D. S. Weiss, Y. Weinrauch, A. Zychlinsky, Neutrophil extracellular traps kill bacteria. *Science* 303, 1532-1535 (2004).
3. P. Li, M. Li, M. R. Lindberg, M. J. Kennett, N. Xiong, Y. Wang, PALM is essential for antibacterial innate immunity mediated by neutrophil extracellular traps. *J. Exp. Med.* 207, 1853-1862 (2010).
4. V. Thammavongsa, D. M. Missiakas, O. Schneewind, Staphylococcus aureus degrades neutrophil extracellular traps to promote immune cell death. *Science*. 342, 863-866 (2013).
5. S. R. Clark, A. C. Ma, S. A. Tavener, B. McDonald, Z. Goodarzi, M. M. Kelly, K. D. Patel, S. Chakrabarti, E. McAvoy, G. D. Sinclair, E. M. Keys, E. Allen-Vercoe, R. Devinney, C. J. Doig, F. H. Green, P. Kubes, Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood. *Nat. Med.* 13, 463-469 (2007).
6. T. A. Fuchs, A. Brill, D. Duerschmied, D. Schatzberg, M. Monestier, D. D. Myers, Jr., S. K. Wroblewski, T. W. Wakefield, J. H. Hartwig, D. D. Wagner, Extracellular DNA traps promote thrombosis. *Proc. Natl. Acad. Sci. U. S. A.* 107, 15880-15885 (2010).
7. A. Warnatsch, M. Ioannou, Q. Wang, V. Papayannopoulos, Neutrophil extracellular traps license macrophages for cytokine production in atherosclerosis. *Science* 349, 316-

- 320 (2015).
8. 8. B. Engelmann, S. Massberg, Thrombosis as an intravascular effector of innate immunity. *Nat. Rev. Immunol.* 13, 34-45 (2013).
 9. 9. S. K. Jorch, P. Kubes, An emerging role for neutrophil extracellular traps in noninfectious disease. *Nat. Med.* 23, 279-287 (2017).
 10. 10. A. Hakkim, B.G. Furnrohr, K. Amann, B. Laube, U.A. Abed, V. Brinkmann, M. Herrmann, R.E. Voll, A. Zychlinsky. Impairment of neutrophil extracellular trap degradation is associated with lupus nephritis. *Proc. Natl. Acad. Sci. U. S. A.* 107, 9813-9818 (2010).
 11. 11. M. Napirei, S. Ludwig, J. Mezhrab, T. Klockl, H. G. Mannherz, Murine serum nucleases contrasting effects of plasmin and heparin on the activities of DNase1 and DNase1-like 3 (DNase1I3). *FEBS J.* 276, 1059-1073 (2009).
 12. 12. M. Napirei, A. Ricken, D. Eulitz, H. Knoop, H. G. Mannherz, Expression pattern of the deoxyribonuclease 1 gene: lessons from the Dnase1 knockout mouse. *Biochem. J.* 380, 929-937 (2004).
 13. 13. V. Sisirak, B. Sally, V. D'Agati, W. Martinez-Ortiz, Z. B. Ozcakar, J. David, A. Rashidfarrokhi, A. Yeste, C. Panea, A. S. Chida, M. Bogunovic, Ivanov, II, F. J. Quintana, I. Sanz, K. B. Elkon, M. Tekin, F. Yalcinkaya, T. J. Cardozo, R. M. Clancy, J. P. Buyon, B. Reizis, Digestion of Chromatin in Apoptotic Cell Microparticles Prevents Autoimmunity. *Cell* 166, 88-101 (2016).
 14. 14. M. Demers, D. S. Krause, D. Schatzberg, K. Martinod, J. R. Voorhees, T. A. Fuchs, D. T. Scadden, D. D. Wagner, Cancers predispose neutrophils to release extracellular DNA traps that contribute to cancer-associated thrombosis. *Proc. Natl. Acad. Sci. U. S. A.* 109, 13076-13081 (2012).
 15. 15. T. A. Fuchs, U. Abed, C. Goosmann, R. Hurwitz, I. Schulze, V. Wahn, Y. Weinrauch, V. Brinkmann, A. Zychlinsky, Novel cell death program leads to neutrophil extracellular traps. *J. Cell Biol.* 176, 231-241 (2007).
 16. 16. C. Frank, D. Werber, J. P. Cramer, M. Askar, M. Faber, M. an der Heiden, H. Bernard, A. Fruth, R. Prager, A. Spode, M. Wadl, A. Zoufaly, S. Jordan, M. J. Kemper, P. Follin, L. Muller, L. A. King, B. Rosner, U. Buchholz, K. Stark, G. Krause, HUS Investigation Team, Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. *N. Engl. J. Med.* 365, 1771-1780 (2011).
 17. 17. S. A. Braune, D. Wichmann, M. C. von Heinz, A. Nierhaus, H. Becker, T. N. Meyer, G. P. Meyer, M. Muller-Schulz, J. Fricke, A. de Weerth, W. W. Hoepker, J. Fiehler, T. Magnus, C. Gerloff, U. Panzer, R. A. Stahl, K. Wegscheider, S. Kluge. Clinical features of critically ill patients with Shiga toxin-induced hemolytic uremic syndrome. *Crit. Care. Med.* 41, 1702-1710 (2013).
 18. 18. M. Jimenez-Alcazar, M. Napirei, R. Panda, E. C. Kohler, J. A. Kremer Hovinga, H. G. Mannherz, S. Peine, T. Renne, B. Lammle, T. A. Fuchs, Impaired DNase1-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies. *J. Thromb. Haemost.* 13, 732-742 (2015).
 19. 19. J. Leffler, Z. Prohaszka, B. Mikes, G. Sinkovits, K. Ciacma, P. Farkas, M. Reti, K. Kelen, G. S. Reusz, A. J. Szabo, M. Martin, A. M. Blom, Decreased neutrophil extracellular trap degradation in shiga toxin-associated haemolytic uraemic syndrome. *J.*

- Innate Immun. 9, 12-21 (2017).
20. 20. M. Leppkes, C. Maueroeder, S. Hirth, S. Nowecki, C. Gunther, U. Billmeier, S. Paulus, M. Biermann, L. E. Munoz, M. Hoffmann, D. Wildner, A. L. Croxford, A. Waisman, K. Mowen, D. E. Jenne, V. Krenn, J. Mayerle, M. M. Lerch, G. Schett, S. Wirtz, M. F. Neurath, M. Herrmann, C. Becker, Externalized decondensed neutrophil chromatin occludes pancreatic ducts and drives pancreatitis. *Nat. Commun.* 7, 10973 (2016).
 21. 21. C. Schauer, C. Janko, L. E. Munoz, Y. Zhao, D. Kienhofer, B. Frey, M. Lell, B. Manger, J. Rech, E. Naschberger, R. Holmdahl, V. Krenn, T. Harrer, I. Jeremic, R. Bilyy, G. Schett, M. Hoffmann, M. Herrmann, Aggregated neutrophil extracellular traps limit inflammation by degrading cytokines and chemokines. *Nat. Med.* 20, 511- 517 (2014).
 22. 22. V. Kumar, A. K. Abbas, N. Fausto, J. C. Aster, Hemodynamic disorders, thromboembolic disease, and shock. In *Robbins & Cotran Pathologic Basis of Disease*. (Elsevier Health Sciences, 2009), chap. 4.
 23. 23. M. Al-Mayouf, A. Sunker, R. Abdwani, S.A. Arawi, F. Almurshedi, N. Alhashmi, A. Al Sonbul, W. Sewairi, A. Qari, E. Abdallah, E., M. Al-Owain, S. Al Motywee, H. Al-Rayes, M. Hashem, H. Khalak, L. Al-Jebali, F. S. Alkuraya. Loss-of-function variant in DNASE1L3 causes a familial form of systemic lupus erythematosus. *Nat. Genet.* 43, 1186-1188 (2011).
 24. 24. K. Yasutomo, T. Horiuchi, S. Kagami, H. Tsukamoto, C. Hashimura, M. Urushihara, Y. Kuroda. Mutation of DNASE1 in people with systemic lupus erythematosus. *Nat. Genet.* 28, 313-314 (2001).
 25. 25. M. Napirei, H. Karsunky, B. Zevnik, H. Stephan, H. G. Mannherz, T. Moroy, Features of systemic lupus erythematosus in Dnase1-deficient mice. *Nat. Genet.* 25, 177-181 (2000).
 26. 26. S. Gupta, M. J. Kaplan, The role of neutrophils and NETosis in autoimmune and renal diseases. *Nat. Rev. Nephrol.* 12, 402-413 (2016).
 27. 27. R. Mizuta, S. Araki, M. Furukawa, Y. Furukawa, S. Ebara, D. Shiokawa, K. Hayashi, S. Tanuma, D. Kitamura, DNase gamma is the effector endonuclease for internucleosomal DNA fragmentation in necrosis. *PloS one* 8, e80223 (2013).
 28. 28. J. Rossaint, J. M. Herter, H. Van Aken, M. Napirei, Y. Doring, C. Weber, O. Soehnlein, A. Zarbock, Synchronized integrin engagement and chemokine activation is crucial in neutrophil extracellular trap-mediated sterile inflammation. *Blood* 123, 2573-2584 (2014).
 29. 29. A. Bradley, K. Anastasiadis, A. Ayadi, J. F. Battey, C. Bell, M. C. Birling, J. Bottomley, S. D. Brown, A. Burger, C. J. Bult, W. Bushell, F. S. Collins, C. Desaintes, B. Doe, A. Economides, J. T. Eppig, R. H. Finnell, C. Fletcher, M. Fray, D. Friendewey, R. H. Friedel, F. G. Grosveld, J. Hansen, Y. Herault, G. Hicks, A. Horlein, R. Houghton, M. Hrabe de Angelis, D. Huylebroeck, V. Iyer, P. J. de Jong, J. A. Kadin, C. Kaloff, K. Kennedy, M. Koutsourakis, K. C. Lloyd, S. Marschall, J. Mason, C. McKerlie, M. P. McLeod, H. von Melchner, M. Moore, A. O. Mujica, A. Nagy, M. Nefedov, L. M. Nutter, G. Pavlovic, J. L. Peterson, J. Pollock, R. Ramirez- Solis, D. E. Rancourt, M. Raspa, J. E. Remacle, M. Ringwald, B. Rosen, N. Rosenthal, J. Rossant, P. Ruiz Noppinger, E. Ryder, J. Z. Schick, F. Schnutgen, P. Schofield, C. Seisenberger, M. Selloum, E. M. Simpson, W. C. Skarnes, D. Smedley, W. L. Stanford, A. F. Stewart, K. Stone, K. Swan, H. Tadepally, L. Teboul, G.

P. Tocchini- Valentini, D. Valenzuela, A. P. West, K. Yamamura, Y. Yoshinaga, W. Wurst, The mammalian gene function resource: the International Knockout Mouse Consortium. Mamm. Genome 23, 580-586 (2012).

Example 2: Development and use of DNase1L3 and variants for therapy

Introduction

[0162] Uncontrolled inflammation causes a plethora of conditions (7). Neutrophil extracellular traps (NETs) are DNA filaments decorated with toxic proteins (2). NETs protect against microbial infection, but excessive NET formation triggers and exacerbates inflammatory and autoimmune diseases (3). DNase1 (D1), a DNA-degrading enzyme found in blood, has been widely used to disarm NETs in vitro and in vivo. Indeed, infusion of recombinant human D1, an FDA-approved drug (dornase alpha) for cystic fibrosis (CF), prevents the pathological effects of NETs in various disease models (3), supporting the use of D1 for prophylactic therapy.

DNase1-like 3 is novel drug candidate for NET associated diseases

[0163] In Example 1, we found that in addition to D1, DNase1-like 3 (D1L3) also degrades NETs. Both enzymes maintain blood and tissue homeostasis during inflammatory responses. In brief, we showed that NETs are generated within blood vessels during neutrophilia, a hallmark of inflammation, under sterile as well as infectious conditions. D1 and D1L3 cleave NETs and thus prevent the aggregation of DNA-filaments of NETs to vascular clots. In the absence of D1 and D1L3, clots of NETs obstruct blood flow and cause organ damage and death. NET-clots are formed independently of platelets or fibrin and are therefore resistant to conventional anti-thrombotic therapy. Thus, we uncovered clots of NETs as a novel mechanism for vascular occlusion and D1L3 a new prophylactic therapy against NETs (see **Example 1**).

[0164] D1 and D1L3 belong, along with DNase1-like 1 (D1L1) and DNase1-like 2 (D1L2), to the DNase1-protein family (4). D1 and D1L3 are expressed in a variety of species including, humans, primates, and rodents. D1 is predominantly expressed in the gastrointestinal tract and exocrine glands (5), whereas hematopoietic cells, namely macrophages and dendritic cells produce D1L3 (6, 7). We found both DNases at low concentrations in serum of mice and humans (Example 1, **FIG. 14**), but the origin of D1 and D1L3 in circulation is unknown (7, 8). D1 preferentially cleaves protein-free DNA (e.g. bacterial DNA, plasmid DNA), whereas D1L3 targets chromatin, the complex of DNA and histones, which is commonly found in the nucleus of eukaryotic cells (7, 9, 10). D1 activity is inhibited upon binding to monomeric actin and sensitive to physiological salt concentrations (11-13). In addition, D1 is glycosylated at N40

(corresponds to N18 in the mature enzyme without signal peptide) and N128 (N106), which makes the enzyme resistant to inactivation by serum proteases (9). In contrast, D1L3 lacks glycosylation and actin-binding sites, which causes its susceptibility towards several proteases and resistance towards actin, respectively (9).

[0165] D1 was first discovered in body fluids of the gastrointestinal tract and urine, taking part in the digestion of DNA from food intake. We observed in the course of this disclosure that systemically administered DNase1 (e.g. through intraperitoneal or intravenous injection) has a relatively short half-life in circulation and is secreted in an enzymatically active form into the urine and, unexpectedly into bile. Its secretion into the bile fluid opens up the opportunity to target extracellular DNA and/or NETs in the biliary duct via systemic DNase1 therapy.

[0166] D1L3 was first described in the context of programmed cell death (14). D1L3 features two nuclear localization sites (NLS1, NLS2). NLS2 is embedded within a C-terminal tail that is unique to D1L3 and not present in D1. Both NLS target the enzyme to the nucleus during apoptosis (14). Indeed, intracellular D1L3 is required for fragmentation of nuclear DNA within apoptotic and necrotic cells *in vivo* (15). D1L3 requires its C-terminal tail to degrade extracellular DNA, namely lipid-encapsulated DNA, such as found in transfections, and chromatin within apoptotic bodies (7, 10). In transfections, cDNA and cationic lipids form a complex that penetrates through the plasma membrane of target cells. D1L3 interferes with transfections and the C-terminal tail is critical for this function (10). Apoptotic bodies, lipid vesicles filled with chromatin from apoptotic cells, are physiological substrates of D1L3 (7). Here, the C-terminal tail enables D1L3 to penetrate through lipid membranes of apoptotic bodies and degrade the chromatin load. Importantly, the C-terminal tail is also required for the degradation of lipid-free extracellular chromatin by D1L3 (7). The C-terminal tail is believed to be critical for the functions of D1L3 in the extracellular space.

[0167] While hD1 is effectively targeting NETs in animal models, mainly mice, it has shown limited or no therapeutic effect in clinical trials of CF (16) and emphysema (17), respectively. NETs are formed in CF (18), indicating that wild-type hD1 is not optimal for targeting NETs in patients. Hyperactive variants have been generated to circumvent the therapeutic limitations of D1, but their clinical manufacturing is challenging (12). Notably, NET-DNA contains histones and is structurally organized like chromatin (2). Thus, we speculated that the efficient degradation of chromatin is required for an optimal therapy against NETs.

[0168] A combinational therapy of hD1 and tissue plasminogen activator (tPA, alteplase) of emphysema patients has been described (17). The data are in line with our hypothesis because tPA enables D1 to degrade chromatin *in vitro* (19). US 9,642,822, further discloses combination therapy with tPA and D1 for therapeutic targeting of NETs (20). However, treatment with tPA is commonly associated with an increased risk of bleeding due to its fibrinolytic activity (17). D1L3 may provide a safer strategy for targeting NETs without putting patients at risk for bleeding, but a therapy with D1L3 protein has not been explored.

Engineering of DNase1 and DNase1-like 3 variants through amino acid transfer

[0169] In this Example we generated variants of D1 and D1L3 that are engineered for therapy against extracellular chromatin (including NETs). To design variants, we hypothesized that the capacity of human D1L3 to degrade chromatin is encoded by individual amino acids and/or amino acid sequences, which are absent in human D1. Furthermore, we speculated that transferring these individual amino acids and/or an amino acid sequence from human D1L3 to D1 generates hyperactive D1 variants with increased chromatin-degrading activity. Similarly, enzymatic properties (e.g. glycosylation) can be transferred from D1 to D1L3. To test our hypothesis, we employed conventional amino acid mutations and developed a novel building-block (BB) technology to engineer variants of DNases from the DNase1-protein family.

[0170] To produce variants of D1 and D1L3, we used in vitro expression systems [e.g. Chinese hamster ovary (CHO) cells, human embryonic kidney (HEK) cells, *Pichia pastoris*]. Enzymatic activity was characterized using the degradation of high-molecular weight (HMW)-DNA as a readout (**FIG. 16**). We chose two types of HMW-DNA: (a) double-stranded DNA (dsDNA) that was isolated from salmon testes. The degradation of HMW-dsDNA was analyzed by zymography; (b) chromatin of purified nuclei from HEK293 cells. To analyse its degradation, HMW-chromatin was first incubated with the DNase variants, followed by DNA isolation and visualization via agarose gel electrophoresis (AGE). Chromatin degradation was detected by low molecular weight (LMW)-DNA. In this setup, wild-type human D1 (SEQ ID NO: 1) has approximately 100-fold higher activity to degrade dsDNA than wild-type D1L3 (SEQ ID NO: 2), whereas wild-type D1L3 degrades chromatin with approximately 100-fold more efficiency than wild-type D1 (**FIG. 16**). We found synergistic effects of D1 and D1L3 in degrading chromatin, but not dsDNA. If D1 and D1L3 are mixed at a ratio of 10: 1 or 1: 1, we observed increased chromatin degradation, compared to each D1 or D1L3 alone (**FIG. 16**). Thus, we screened for D1 and/or D1L3 variants, which show increased chromatin and/or dsDNA degrading activity.

[0171] Selected D1 and D1L3 variants were further tested to degrade intravascular NETs, using an in vivo-expression system for members of the D1-protein family. See PCT/EP2018/051444. The system is based on two components, (a) an expression plasmid that enables the stable in vivo transfection of hepatocytes with cDNA of D1, D1L3, and variants thereof in recipient mice, and (b) *Dnase1^{-/-}Dnase1I3^{-/-}* deficient mice, which are characterized by the absence of DNase activity in serum. In brief, the cDNA of candidate DNases is cloned into the hepatic expression vector. Next, the vector is co-expressed in *Dnase1^{-/-}Dnase1I3^{-/-}* deficient mice along with the cDNA of murine *Csf3*, which encodes for the granulocyte colony-stimulating factor (G-CSF). The expression of *Csf3* induces neutrophilia with concomitant intravascular NET-formation, which in turn causes the aggregation of undigested NETs within blood vessels and death in *Dnase1^{-/-}Dnase1I3^{-/-}* mice. To identify drug candidates, we screened for D1, D1L3, and variants thereof that efficiently degrade NETs and thus prevent morbidity and mortality in this model.

Results

Engineering DNase1 variants through conventional amino acid substitutions

[0172] Our central hypothesis is that the capacity of human D1L3 to degrade chromatin is mediated by amino acids that are not present in D1. Sequence alignment showed that 44% of the amino acids in human D1 and human D1L3 are identical (**FIG. 17**). We exclusively mutated only variable amino acids (56% non-shared amino acids) to generate D1 and DL3 variants.

[0173] D1L3 is a cationic protein with an isoelectric point of 9.35, whereas D1 and the other members are more basic proteins. We therefore speculated that the cationic nature of D1L3 enables the enzyme to degrade chromatin. Through sequence alignment, we identified 37 arginine or lysine residues in the variable amino acids, which are absent in D1 (**FIG. 17**). These amino acid residues are located at the amino acid positions 29 (position 9 with the signal peptide), 39 (9), 47 (7), 50 (30), 66 (46), 74 (54), 77 (57), 80 (60), 81 (61), 92 (72), 109 (89), 114 (94), 115 (95), 163 (143), 176 (156), 180 (160), 199 (179), 200 (180), 203 (183), 208 (188), 212 (192), 226 (206), 227 (207), 239 (219), 250 (230), 259 (239), 262 (242), 280 (260), 285 (265), 291 (271), 292 (272), 297 (277), 298 (278), 299 (279), 301 (281), 303 (283), 304 (284) with respect to SEQ ID NO: 2.

[0174] Twenty-six of these additional arginine or lysine residues in D1L3 correspond to the following amino acid positions in D1 (SEQ ID NO: 1): 31 (position 9 with the signal peptide), 41 (19), 49 (40), 52 (30), 68 (46), 76 (54), 79 (57), 81 (59), 92 (70), 109 (87), 114 (92), 115 (93), 164 (142), 177 (155), 181 (159), 200 (178), 201 (179), 204 (182), 209 (187), 213 (191), 227 (205), 239 (217), 250 (228), 259 (237), 262 (240), and 280 (260). In addition, D1L3 features 3 sites of insertions of arginine or lysine residues: R80, K226 and a C-terminal tail (after amino acid position 282 in SEQ ID NO: 2), which features 3 arginine and 6 lysine residues.

[0175] We aimed to transfer the arginine and lysine residues that are part of the core body of D1L3 to D1 by introducing the following mutations into wild-type D1 (**FIG. 18**): Q31R, A41K, Q49R, S52K, T68R, N76K, Q79R, D80_A81insR, A81R, P92R, D109K, V114K, D115R, A164K, Q177K, G181K, P200K, S201K, S204R, W209R, T213R, A226_T227insK, T227K, A239R, D250K, A259K, G262K, and K280M with respect to SEQ ID NO:1, thus generating 28 variants of D1.

[0176] In this Example, we were able to generate D1 variants that contain 23 of the 28 amino acid substitutions. We expressed the cDNA of these D1-variants in HEK293 cells and analyzed the dsDNA and chromatin degrading activity in the culture supernatants. We observed that the following amino acid modifications partially increased the chromatin degrading activity, while no major differences in dsDNA degradation were observed (**FIG. 19**): Q31R, Q49K, N76K, Q79R, P92R, D109K, A164K, Q177K, P200K, S201K, S204R, T213R. Interestingly, mouse and rat D1 (SEQ ID NO: 4), which share 79% and 78% of the amino acids of human D1, respectively, feature several of the analyzed amino acid substitutions, including Q31R, Q49K, and Q79K

(FIG. 20). Analysis of supernatants from HEK293 cells expressing murine D1 and human D1 showed that murine D1 has an increased activity to degrade chromatin, when compared to human D1 (FIG. 21). We injected *Dnase1^{-/-}Dnase1/3^{-/-}* mice with a mixture of *Csf3* and murine D1 (*mDnase1*, SEQ ID NO: 3), human D1 (*hDnase1*, SEQ ID NO: 1), or an empty control plasmid (*Ctrl*). Mice receiving the control plasmid died within 7 days, whereas the expression of murine D1 provided a full protection for at least 28 days (FIG. 21). Unexpectedly, the expression of human D1 showed only a partial protection from death and more than half of the animals died within 14 days (FIG. 21), indicating that human D1 is less efficient in degrading intravascular NETs than murine D1. We now mutated Q31R, Q49K, Q79R, and W209R in human D1 to generate a murine-like D1-variant (SEQ ID NO: 5). A total of six mutations (Q31R, Q49K, Q79R, Q177K, W209R, G262K) were made to generate a rat-like D1-variant (SEQ ID NO: 6). Both rodent-like D1 variants were cloned into the hepatocyte expression vector and delivered along with *Csf3* into *Dnase1^{-/-}Dnase1/3^{-/-}* mice. We observed that gene therapy with the murine- and rat-like variant of D1 provided a full protection against NET-mediated vascular occlusions and death (FIG. 22). Importantly, in vitro expression of the rodent-like human D1-variants showed an increased activity to degrade chromatin (FIG. 22). Collectively, the data illustrate that (a) human D1 has a limited capacity to degrade NETs in vivo; (b) transferring multiple cationic amino acids from human D1L3 to human D1 generates hyperactive variants.

[0177] Several attempts have been made to generate hyperactive variants of D1 (6, 12, 13, 21-29). We found 184 mutations, which altered 82 of the 282 amino acids of wild-type human D1 (FIG. 23). Alignment of amino acid sequences of wild-type human D1 and human D1L3 revealed three gain-of-function mutations in human D1 that are already present in wild-type human D1L3 (FIG. 23): Q31R (corresponds to Q9R in D1 without signal peptide), T227K (T205K), and A136F (A114F). While the mutations Q31R and T227K may alter the DNA-binding properties of hD1 and thus cause hyperactivity, A136F confers resistance to inhibition by monomeric actin (21-24, 26, 29).

[0178] We designed a new D1-variant that carries the combination of Q31R/T227K/A136F (SEQ ID NO: 7) and compared its enzymatic activity to wild-type D1 (SEQ ID NO: 1), as well as to D1-variants carrying single mutations in Q31R, T227K, A136F, or a combined mutation in Q31R and T227K (SEQ ID NO: 8). We characterized the dsDNA and chromatin-degrading activity in the culture supernatants of transfected HEK293 cells. We observed no major differences in dsDNA-degrading activity with D1-variants carrying single or combinational mutations of Q31R and T227K (FIG. 24). Interestingly, chromatin was partially and fully degraded in A136F mutants and the combinational mutation of Q31R/T227K/A136F, respectively (FIG. 24). Next, we tested whether the Q31R/T227K/A136F variant of D1 is capable of preventing NET-mediated lethal vascular occlusions in neutrophilic *Dnase1^{-/-}Dnase1/3^{-/-}*. All mice receiving a control plasmid without DNases died, whereas all mice expressing the D1-variant (SEQ ID NO: 7) survived until the end of the experiment without showing signs of distress (FIG. 24). Using purified proteins, we observed that the triple amino acid substitution increases the chromatin degrading activity approximately 10-20 fold

compared to wild-type D1 without detectable differences in dsDNA degrading activity (**FIG. 25**). Chromatin from nuclei and NETs contains actin (30, 31). The data suggest that chromatin degradation by D1 variants requires cationic amino acid substitutions as well as mutations that confer actin resistance - two features which are present in wild-type D1L3. To further corroborate this notion, we generated a D1 variant which features the combination of the mutations A164K and A136F (SEQ ID NO: 9). Indeed, we observed a strong increase in chromatin degrading activity in the D1 variant with both mutations (**FIG. 26**), supporting our hypothesis.

Development of building block engineering of DNase1-protein family members

[0179] D1L3 features three sites that contain additional amino acids: the C-terminal tail starting after Q282 (NH₂-SSRAFTBSKKSVTLRKKTKSKRS-COOH) (SEQ ID NO:33), and at two sites within the enzyme at S79/R80 and at K226 (**FIG. 17**). The 23 amino acids of the C-terminal tail of D1L3 have been attached to the C-terminus of D1 (10, 28). We observed that the insertion of an arginine-residue at position 226 of DNase 1 (A226_T227insK) generated a D1-variant with reduced enzymatic activity to degrade dsDNA, while no such effect was observed with the substitution T227K (**FIG. 27**). Thus, an insertion of a K/R-residue goes along with a risk of reducing D1 function. The insertion of a charged amino acid may influence the local protein structure. Given that D1 is a globular enzyme that comprises one amino acid chain [**FIG. 28, (27)**], it is conceivable that such local alteration may render the whole enzyme inactive. Indeed, numerous non-conservative mutations throughout the D1 amino acid sequence inactivate the enzyme (12, 25, 26). We hypothesized that the transfer of local protein structures by implanting not only single arginine and lysine residues but also the neighboring amino acids sequences reduces the risk of inactivation. We searched within D1 and D1L3 for conserved amino acids in the vicinity of A226_T227insK that can be used as anchors for the insertion. We identified a D223/T224/T225 motif and a conserved T229 in D1 as N-terminal and C-terminal anchors, respectively (**FIG. 27**). We replaced 3 amino acids within D1 (ATP) with 4 amino acids, including K226, from D1L3 (VKKS) *in silico*. Expression of the cDNA of the new D1-variant (A226_P228delinsVKKS) in HEK239 cells revealed a functionally active enzyme with a similar dsDNA-degrading activity, when compared to wild-type D1 (**FIG. 27**). The data suggest that the variable amino acids between conserved amino acids are interchangeable between D1 and D1L3.

[0180] We conceptualized a building block-technology to transfer enzymatic properties from one member of the D1-protein family to another. The following cardinal steps characterize the technology (**FIG. 29**):

1. (1) Provide protein-protein alignment of donor and recipient DNase
2. (2) Identify variable amino acid or amino acid sequence for transfer (building block)
3. (3) Identify conserved amino acids in donor and recipient DNase that are located up and downstream of building block (anchors), respectively.

4. (4) Replace cDNA encoding for building block between C- and N-anchors in recipient DNase, with cDNA between the anchors in donor DNase.
5. (5) Synthesize cDNA of chimeric DNase, followed by in vitro/in vivo expression into a recipient organism that is preferably deficient in both donor and recipient DNase (e.g. CHO cells or *Dnase1^{-/-}Dnase1l3^{-/-}* mice).

Engineering DNase1 variants through building block technology

[0181] A multiple-species alignment of D1 and D1L3 from human, mouse, rat, and chimpanzee (**FIG. 30**), showed that N- and C-terminal anchors are conserved among these species. These anchor amino acids or amino acid sequences flank 62 building blocks of variable amino acids and amino acid sequences, which include the amino acid sequence in D1 (ATP) and D1L3 (VKKS) from the above-mentioned experiments as building blocks #49 (**FIG. 31**).

[0182] The transfer of these building blocks from D1L3 into D1 generates D1-variants with the following mutations (**FIG. 31**): 1M_S22delinsMSRELAPLLLLLSIHSALA, L23_A27delinsMRICS, I30_T32delinsVRS, E35_T36delinsES, M38_I47delinsQEDKNAMDVI, Q49_S52delinsKVIK, Y54C, I56_Q60delinsIILVM, V62_R63delinsIK, S65_K72delinsSNNRICPI, L74_N76delinsMEK, Q79_T84delinsRNSRRGIT, H86N, V88_V89delinsVI, E91_P92delinsSR, N96_S97delinsNT, R101Q, L103A, V105L, R107_Q110delinsKEKL, A113_S116delinsVKRS, Y118H, D120H, G122_N128delinsYQDGDA, T130S, N132S, A136_I137delinsFV, R139W, F141_F144delinsQSPH, E146_E149delinsAVKD, A151V, V153I, A157_A158delinsTT, G160_A162delinsETS, A164K, A168E, Y170_D171delinsVE, L174T, Q177_K179delinsKHR, G181_L182delinsKA, D184_L187delinsNFIF, R199_Q202delinsPKKA, S204_S205delinsKN, W209R, S211D, T213R, Q215V, P219G, S221_A222delinsQE, A226_P228delinsVKKS, H230N, V238_A239delinsLR, M241_A246delinsQEIVSS, D250K, A252_P254delinsNSV, N256D, A259_A260delinsKA, G262K, S264_L267delinsTEEE, Q269_I271delinsLDV, Y275F, V279_M280delinsFK, and K282delinsQSSRAFTNSKKSVTLRKKTCKSKRS.

[0183] The following D1L3-variants are generated if the building blocks are transferred from D1 to D1L3 (**FIG. 31**): M1_A20delinsMRGMKLLGALLALAALLQGAVS, M21_S25delinsLKIAA, V28_S30delinsIQT, E33_S34delinsET, Q36_I45delinsMSNATLVSYI, K47_K50delinsQILS, C52Y, I54_M58delinsIALVQ, I60_K61delinsVR, S63_I70delinsSHLTAVGK, M72_K74delinsLDN, R77_T84delinsQDAPDT, N86H, V88_I89delinsVV, S91_R92delinsEP, N96_T97delinsNS, Q101R, A103L, L105V, K107_L110delinsRPDQ, V113_S116delinsAVDS, H118Y, H120D, Y122_A127delinsGCEPCGN, V129T, S131N, I35F_136VdelinsAI, W138R, Q140_H143delinsFSRF, A145_D148delinsAVKD, V150A, I152A, T156_T157delinsAA, E159_S161delinsGDA, K163A, E167A, V169_E170delinsYD, T173L, K176_R178delinsQEK, K180_A181delinsGL, N183_F186delinsDVML, P198_A201delinsRPSQ, K203_N204delinsSS, R208W, D210S, R212T, V214Q, G218P, Q220_E221delinsSA, V225_S228delinsATP, N230H, L238_R239delinsVA, Q241_S246delinsMLLRGA, K250D, N252_V254delinsALP, D256N,

K259_A260delinsAA, K262G, T264_E267delinsSDQL, L269_V271delinsQAI, F275Y, F279_K280delinsVM, and Q282_S305delinsK.

[0184] Next, we conceptualized a sequential approach to engineer D1-variants with D1L3 activity that starts with the transfer of multiple adjacent building blocks (clusters), continues with the transfer of individual building blocks, and ends with a transfer of individual amino acids or the combination of multiple building blocks into new chimeric enzymes (**FIG. 32**). This approach reduces the number of D1-D1L3-chimera in the initial screening.

[0185] To test our method, we designed a total of 19 D1-variants comprising either individual building blocks or clusters of building block cluster from D1L3 (**FIG. 31**). These D1-variants feature the following amino acid mutations: 1M_S22delinsMSRELAPLLLLLSIHSALA, L23_A27delinsMRICS/I30_T32delinsVRS/E35_T36delinsES, M38_I47delinsQEDKNAMDVI, Q49_S52delinsKVIK/Y54C/I56_Q60delinsIILVM, V62_R63delinsIK/S65_K72delinsSNNRICPI/L74_N76delinsMEK, Q79_T84delinsRNSRRGIT, H86N/V88_V89delinsVI/E91_P92delinsSR, N96_S97delinsNT/R101Q/L103A/V105L, R107_Q110delinsKEKL/A113_S116delinsVKRS/Y118H/D120H, G122_N128delinsYQDGDA/T130S/N132S, A136_I137delinsFV, R139W/F141_F144delinsQSPH/E146_E149delinsAVKD/A151V/V153VA157_A158delinsTT/ G160_A162delinsETS/A164K, A168E/Y170_D171delinsVE/L174T, Q177_K179delinsKHR/G181_L182delinsKA/D184_L187delinsNFIF, R199_Q202delinsPKKA/S204_S205delinsKN/W209R/S211D/T213R/Q215V/P219G/S221_A222delinsQE, A226_P228delinsVKKS, H230N/V238_A239delinsLR/M241_A246delinsQEIVSS/D250K/A252_P254delinsNSV, N256D/A259_A260delinsKA/G262K/S264_L267delinsTEEE/Q269_I271delinsLDV, and Y275F/V279_M280delinsFK/K282delinsQSSRAFTNSKKSVTLRKKTCSKRS. Next, we cloned the cDNA into an expression vector, which was transfected into HEK293 cells. Analysis of the cell supernatants showed dsDNA degradation by all samples (**FIG. 33**). Furthermore, we observed that the transfer of building blocks (BB) 11 (see SEQ ID NO: 10), BB 12-14 (see SEQ ID NO: 11), BB 26 (see SEQ ID NO: 12), BB 41-48 (see SEQ ID NO: 13), and BB 49 (see SEQ ID NO: 14) from D1L3 to D1 resulted in enzymes with increased chromatin degrading activity. All these chimeric enzymes exhibited the same or more activity to degrade dsDNA substrates than wild-type D1. The building blocks 11 (RNSRRGI in SEQ ID NO: 2) and 49 (VKKS in SEQ ID NO: 2) from D1L3 contain R80/R81 and K227, respectively, which are not present in D1. The D1L3-BB cluster 41-48 (PKKAWKNIRLRTDPRFVWLIGDQE from SEQ ID NO:2) features 5 additional arginine and lysine residues than its counterpart in D1 (RPSQWSSIRLWTSPTFQWLIPDSA of SEQ ID NO:1). These additional cationic amino acids may be responsible for the hyperactivity. The D1-building blocks 12-14 and 26 contain the amino acid sequences H86 to R95 and A136 to V138 in SEQ ID NO: 1, which includes amino acid residues that are required for binding of the D1-inhibitor actin. Thus, replacement of these amino acid sequences with the respective building blocks from D1L3, which do not interact with actin, likely generates actin-resistant variants of D1. We now combined BB 11, 14, 26, 41-19 in one novel D1-variant (SEQ ID NO: 15). We observed that the combination of these gain-of-function BBs increased the chromatin degrading of the D1 variant to levels of wild-type D1L3

(FIG. 33). Thus, the BB technology provides a robust method to generate hyperactive D1 variants.

Engineering DNaseI-like 3 variants through building block technology

[0186] The BB cluster 60 to 62 of D1L3 includes the amino acid from its C-terminal tail (FIG. 33). The tail starts after L281 (FIG. 34) and comprises the amino acid sequence QSSRAFTBSKKSVTLRKKTCSKRS (SEQ ID NO:33), which includes a bi-partite NLS. D1L3 requires the C-terminal tail to degrade DNA within lipid vesicles including transfection reagents and apoptotic bodies (7, 10, 28). Furthermore, the C-terminal tail is required for degradation of lipid- and cell-free chromatin by D1L3 (7). Both findings are based on a D1L3-variant that lacks the amino acids after Q282 and is thus inactive in degrading these substrates. Wilbert et al. attached the C-terminal extension of D1L3 to the C-terminus of D1 and thus transferred the capacity to degrade DNA within lipid vesicles to D1 (10, 28). Here, we employed our BB technology to transfer the C-terminus from D1L3 to D1 and vice versa. We used the conserved S272/D273/H274 motif, which is located in front of the terminal beta-sheet in D1 (FIG. 28), as a N-anchor and exchanged the terminal 3 building blocks between D1 and D1L3 (FIG. 34). This approach generated a D1-variant that features the C-terminal extension of D1L3 due to the mutations Y275F/V279_M280delinsFK/K282delinsQSSRAFTNSKKSVTLRKKTCSKRS (SEQ ID NO: 16) and a truncated D1L3-variant due to the mutations F275Y/F279_K280delinsVM/Q282_S305delinsK (SEQ ID NO: 17). We expressed both variants and the wild-type enzymes in HEK293 cells and characterized the DNase activity in the supernatant. Unexpectedly, we observed that the D1-variant with the C-terminus of D1L3 did not degrade chromatin, while the D1L3-variant lacking the C-terminus of the wild-type enzyme was still able to cleave chromatin (FIG. 34). In fact, we observed an increased chromatin degrading activity with this D1L3 variant. To corroborate this finding, we tested whether the truncated variant of D1L3 in vivo and observed that its expression of protected from NET-mediated mortality of neutrophilic *Dnase1^{-/-}Dnase1/3^{-/-}* mice (FIG. 34). In conclusion, the deletion of the C-terminal does not inhibit, but rather increases the function of D1L3 to degrade lipid-free and cell-free chromatin.

[0187] Serum proteases, in particular plasmin, degrade D1L3, but not D1 (9). The sensitivity of D1L3 towards protease activity would limit its use in clinical practice. Glycosylated proteins display an increased protection against proteases (32). Using software analysis, we could not identify a known consensus sequence for glycosylation in human and murine D1L3. Software analysis of human D1 revealed 2 glycosylation sites, as observed both in silico and in vitro (9). We hypothesized that glycosylated variants of D1L3 may increase the half-life in circulation due to an improved resistance towards serum proteases. A glycosylation site requires a minimal consensus sequence of N-X-T/S. D1 contain two known N-glycosylation sites, N40 (N18 without signal peptide) and N128 [(N106 without signal peptide, (9)]. The amino acids of the D1-glycosylation sites N40-X-T42 and N128-X-T130 correspond to the D38-X-N40 and A127-X-V129 in D1L3. Thus, a glycosylated version of D1L3 needs to feature the amino acids

mutations D38N, N40T, A127N, and V129T (see SEQ ID NO: 18). Instead of using conventional point mutations to insert these amino acid substitutions, we used BB technology to transfer the D1-glycosylation sites N40-X-T42, which lies within BB4 (MSNATLVSY in SEQ ID NO: 1), and N128-X-T130, which lies within the BB cluster 23-25 (GCEPCGN DTFN in SEQ ID NO: 1), from D1 to D1L3 (**FIG. 35**). Thus, we generated two new D1L3 variants, which feature the mutations Q36_I45delinsMSNATLVSYI (SEQ ID NO: 19) and Y122_A127delinsGCEPCGN/V129T/S131N (SEQ ID NO: 20), respectively. We expressed these D1L3 variants along with the WT enzyme in HEK293 cells. Next, we analyzed the supernatant by Western Blot using a polyclonal antibody targeting an epitope in the C-terminal tail of the WT enzyme to detect the WT enzyme as well as the D1L3 variants. As expected the D1L3 variants showed bands of a higher molecular weight presumably to glycosylation, when compared to the WT enzyme (**FIG. 35**). D1L3 variant with glycosylation at N40 or N128 retained their chromatin degrading activity (**FIG. 35**). In conclusion, the data suggest that BB technology can be used to transfer the glycosylation sites from D1 to D1L3, which may generate D1L3 variants with increased protease-resistance and half-life in patients.

Therapy with DNaseI-like 3 and a variant thereof

[0188] Finally, we tested the therapeutic effects of D1L3 in vivo. Histological analysis of vascular clots of NETs show dense aggregates of chromatin (**FIG. 36A**), indicating that histone-free cleavage sites for D1 are rare in vivo. Thus, we hypothesized that an acute therapy with wild-type D1L3 is more effective in degrading intravascular NETs than wild-type D1. To test our hypothesis, we expressed *Csf3* in *Dnase1^{-/-}Dnase1I3^{-/-}* deficient mice to induce vascular clots of NETs. Importantly, preceding exitus, mice with vascular occlusions by NETs develop hematuria and hypothermia, which can be detected non-invasively. Therefore, the animal model allows identifying and hence selecting mice, which have formed NETs, to test drug candidates for an acute therapy against NETs (**FIG. 36B**). Here, we tested purified wild-type human D1 (dornase alpha; SEQ ID NO: 1), wild-type human D1L3 (SEQ ID NO: 2), and a hyperactive D1L3 variant [D1L3^V; (SEQ ID NO: 17)] for an acute therapy against NETs. We injected *Csf3* into *Dnase1^{-/-}Dnase1I3^{-/-}* deficient mice. After 72 hours, the first individual animals started to develop hypothermia and hematuria. We used a drop in body temperature of 4°C or more and the presence of red urine as inclusion criteria. Mice, which developed these phenotypes, were randomized into the different treatment groups. Animals received a single dose, intravenous injection via the tail vein of 1 mg/kg of D1, D1L3, or vehicle. After 30 minutes, the animals were euthanized to collect biosamples (blood and organs). Quantification of clots of NETs in the vasculature of lungs showed that NETs occluded blood vessels in mice receiving vehicle or D1 therapy (**FIG. 36D**). Importantly, mice injected with D1L3 and D1L3^V showed no or few clot of NETs (**FIG. 36D**). In patients, circulating DNA has the size of DNA in mono-nucleosomes [180 base pairs, (33)], indicating that it is a degradation product of high-molecular weight DNA in chromatin. Circulating DNA is commonly used as a surrogate marker of intravascular NET formation, but can also derive from damaged and necrotic tissue (34). We quantified DNA in serum using a fluorescent DNA probe. We observed that the injection of

D1L3 and D1L3^V, but not D1 or vehicle, caused a strong increase in levels of circulating DNA (**FIG. 36E**). Isolation and visualization of circulating DNA showed multimers 180 base pair fragments, indicating the presence of mono- and oligo-nucleosomes in serum from mice receiving the D1L3 therapy (**FIG. 36F**). Mice receiving D1L3^V showed only mono- and di-nucleosomes. The data confirm the hyperactivity of D1L3^V in vivo. DNA isolates from serum of mice receiving D1 therapy showed faint DNA smears of various sizes, whereas no DNA could be isolated from serum of mice injected with vehicle (**FIG. 36F**). Thus, D1L3 may be used, not only for prophylaxis, but also for acute therapy of diseases that associated with NETs.

[0189] We further explored the therapeutic application of D1L3^V in a model of ischemia-reperfusion injury (**FIG. 37A**). In brief, one testicle of male wild-type rats was subjected to testicular torsion as previously described by us (35). The procedure causes an ischemic tissue injury. After randomization and de-torsion of the testicle, the rats received three daily injections of vehicle dornase alpha (D1, 1mg/kg), or a purified DNase1L3 variant (SEQ ID NO: 17, D1L3^V, 1mg/kg). After 7 days the ischemic testicle and untreated testicle were collected. The therapy with the D1L3^V, but not saline or D1, significantly reduced the testicular atrophy (**FIG. 37B**). Moreover, therapy with the D1L3^V, but not saline or D1, substantially reduced the tissue injury and majority of animals no signs of injury (**FIG. 37C**). Collectively, the data identify D1L3 as a potent therapy against ischemia-reperfusion injury.

Drug development of DNase1 variants, DNase1-like 3 and variants

[0190] Nearly 70% of all protein therapeutics are produced using Chinese Hamster Ovary (CHO) (36). Indeed, wild-type DNase1 (dornase alpha) is produced in CHO cells. Despite significant advantages, cell line development and large-scale production in CHO still remains a huge challenge due a considerable degree of variability and no reliable methods for predicting or modeling cell growth characteristics (36). Importantly, CHO cells were not able to stably produce hyperactive variants of D1, which prevented their clinical manufacturing (12). The manufacturing properties of D1L3 are unknown. Thus, we developed stable CHO cell lines producing wild-type D1 (SEQ ID NO: 1), a hyperactive D1 variant (SEQ ID NO: 7), wild-type D1L3 (SEQ ID NO: 2) and a hyperactive D1L3 variant (SEQ ID NO: 17). We cultured the cell lines in bioreactors and observed a stable protein expression with wild-type D1 and the D1 variant, with the latter having strongly increased expression levels (**FIG. 38**), illustrating the feasibility of producing a hyperactive D1 variant in CHO. Unexpectedly, we observed only minor protein levels of wild-type D1L3 and the D1L3 variant (**FIG. 38**), pointing towards a major challenge for clinical manufacturing D1L3 and variants thereof.

[0191] Due to high cell densities in large-scale bioreactors, cells compete for resources and undergo cellular stress leading to increased cell death called apoptosis, a cell death program used by multi-cellular organisms, including mammals (37). D1L3 targets DNA in apoptotic bodies (7). We speculated that the low levels of D1L3 are due to scavenging of the enzyme by

these apoptotic vesicles. To overcome these limitations, we tested the production of D1, D1L3 and their variants using the microbial expression system, *Pichia pastoris*, an established, FDA approved and Generally Recognized As Safe (GRAS) organism with protein secretory abilities (38, 39). A key limitation of this approach is that proteins produced by *P. pastoris* are often hyperglycosylated and may result in enhanced immunogenicity (40). The composition of D1L3 and its variant are non-glycosylated and optimal for expression in *P. pastoris*. We mutated the glycosylation sites in D1 and its variant at N40S and N128S respectively to generate deglycosylated proteins (SEQ ID NO: 21, 23). Protein expression in *P. pastoris* is generally achieved by coupling proteins of interest to secretory signal peptides of various origins. We omitted this step to avoid the generation of immunogenic chimeric proteins. We speculated that the native signal peptide of D1 and/or D1L3 is recognized by *P. pastoris*. After codon-optimization was performed, we expressed the cDNA in *P. pastoris* (SEQ ID NO: 22, 24-26). We achieved expression of D1 and the D1 variant (not shown). Importantly, this approach furthermore enabled the expression of D1L3 and the D1L3 variant, with the latter showing increased expression levels (**FIG. 38**). We purified the proteins and confirmed that the D1L3 has increased activity (**FIG. 38**). In conclusion, the data suggest that using non-mammalian expression systems, such as *P. pastoris*, enables clinical manufacturing of D1L3 and variants thereof.

Discussion

[0192] Two DNA-degrading enzymes have been widely used therapeutically in experimental models and patients. Streptodornase, a secreted DNase from *Streptococcus spp.*, has been tested in disease models of skin homograft, bladder clots, wound debridement, meningitis. DNase1 (including pancreatic DNase1) has been used therapeutically in experimental models of cancer, cystic fibrosis, lupus, thrombosis, stroke, sepsis, lung injury, atherosclerosis, viral infection, sickle cell disease, myocardial infarction, ear infection, wound healing, liver injury, endocarditis, liver infection, pancreatitis, primary graft dysfunction, limb ischemia reperfusion, kidney injury, blood clotting, asthma, alum-induced inflammation, hepatorenal injury. Streptodornase has also been used to treat patients with pleural exudations, hemotorax, intrabiliary blood clots, post pneumatic anemia, ulcers, otolaryngological conditions, oral infections, minor injuries, sinusitis, postoperative rhinoplasties, infertility, bladder catheter, wound cleaning, skin reaction test. Importantly, DNase1 has been tested therapeutically in patients with pneumococcal meningitis, gout, leg ulcers, cystic fibrosis, Kartegener's syndrome, asthma, lobar atelectasis, chronic bronchitis, bronchiectasis, lupus, primary ciliary dyskinesia, bronchiolitis, empyema, pleural infections, cancer, dry eyes disease, lower respiratory tract infections, chronic hematomas, Alzheimer's disease, and obstructive pulmonary disease. In this example, we identified D1L3, variants thereof, as well as D1 variants as novel candidates for a DNase therapy (**FIG. 39**).

Methods

Detection of chromatin degrading activity

[0193] To measure chromatin degrading activity, we mixed samples with nuclei isolated from HEK293 (1,500,000 - 3,000,000 per reaction), 20 mM Tris-HCl pH 7.4, 2 mM MgCl₂, 2 mM CaCl₂, 50 mM NaCl and protease inhibitors. After incubating the samples for 1 hour at 37°C, they were heated up to 65°C for 10 minutes. We proceeded with DNA isolation using the QIAamp DNA Blood Mini Kit (Qiagen) according to the manufacturer's protocol. The eluent was then loaded onto an agarose gel and the DNA fragments were separated by gel electrophoresis. The DNA fluorescence of the gels was recorded with a fluorescence scanner.

[0194] Further methods are described in Example 1.

References**[0195]**

1. 1. V. Kumar, A. K. Abbas, N. Fausto, J. C. Aster, in Robbins & Cotran Pathologic Basis of Disease. (Elsevier Health Sciences, 2009), chap. 2.
2. 2. V. Brinkmann et al., Neutrophil extracellular traps kill bacteria. Science (New York, N.Y.) 303, 1532-1535 (2004).
3. 3. T. A. Fuchs, A. Hakkim, C. F. Urban, in Inflammation: Fundamental Mechanisms, K. Ley, Ed. (World Scientific, 2018), chap. 6.
4. 4. D. Shiokawa, S. Tanuma, Characterization of human DNase I family endonucleases and activation of DNase gamma during apoptosis. Biochemistry 40, 143-152 (2001).
5. 5. M. Napirei, A. Ricken, D. Eulitz, H. Knoop, H. G. Mannherz, Expression pattern of the deoxyribonuclease 1 gene: lessons from the Dnase1 knockout mouse. Biochem J 380, 929-937 (2004).
6. 6. W. F. Baron et al., Cloning and characterization of an actin-resistant DNase I-like endonuclease secreted by macrophages. Gene 215, 291-301 (1998).
7. 7. V. Sisirak et al., Digestion of Chromatin in Apoptotic Cell Microparticles Prevents Autoimmunity. Cell 166, 88-101 (2016).
8. 8. M. Jimenez-Alcazar et al., Impaired DNaseI-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies. Journal of thrombosis and haemostasis : JTH 13, 732-742 (2015).
9. 9. M. Napirei, S. Ludwig, J. Mezrhah, T. Klockl, H. G. Mannherz, Murine serum nucleases--contrasting effects of plasmin and heparin on the activities of DNase1 and DNase1-like 3 (DNase1I3). FEBS J 276, 1059-1073 (2009).
10. 10. A. Wilber, M. Lu, M. C. Schneider, Deoxyribonuclease I-like III is an inducible macrophage barrier to liposomal transfection. Mol Ther 6, 35-42 (2002).
11. 11. W. Kabsch, H. G. Mannherz, D. Suck, E. F. Pai, K. C. Holmes, Atomic structure of the actin:DNase I complex. Nature 347, 37-44 (1990).

12. 12. C. Lam et al., Taming hyperactive hDNase I: Stable inducible expression of a hyperactive salt- and actin-resistant variant of human deoxyribonuclease I in CHO cells. *Biotechnol Prog* 33, 523-533 (2017).
13. 13. C. Q. Pan et al., Improved potency of hyperactive and actin-resistant human DNase I variants for treatment of cystic fibrosis and systemic lupus erythematosus. *J Biol Chem* 273, 18374-18381 (1998).
14. 14. D. Shiokawa, Y. Shika, S. Tanuma, Identification of two functional nuclear localization signals in DNase gamma and their roles in its apoptotic DNase activity. *Biochem J* 376, 377-381 (2003).
15. 15. R. Mizuta et al., DNase gamma is the effector endonuclease for internucleosomal DNA fragmentation in necrosis. *PLoS One* 8, e80223 (2013).
16. 16. H. J. Fuchs et al., Effect of aerosolized recombinant human DNase on exacerbations of respiratory symptoms and on pulmonary function in patients with cystic fibrosis. The Pulmozyme Study Group. *N Engl J Med* 331, 637-642 (1994).
17. 17. N. M. Rahman et al., Intrapleural use of tissue plasminogen activator and DNase in pleural infection. *N Engl J Med* 365, 518-526 (2011).
18. 18. V. Papayannopoulos, D. Staab, A. Zychlinsky, Neutrophil elastase enhances sputum solubilization in cystic fibrosis patients receiving DNase therapy. *PLoS One* 6, e28526 (2011).
19. 19. M. Napirei, S. Wulf, H. G. Mannherz, Chromatin breakdown during necrosis by serum Dnase1 and the plasminogen system. *Arthritis Rheum* 50, 1873-1883 (2004).
20. 20. D. D. Wagner et al. (Children's Medical Center Corporation, Boston, MA (US), United States, 2017), vol. US 9,642,822 B2.
21. 21. R. A. Lazarus, C. Q. Pan. (Genentech Inc., United States, 2002), vol. US 6,391,607 B1.
22. 22. R. A. Lazarus, C. Q. Pan. (Genentech Inc., 2008), vol. US 7,407,785 B2.
23. 23. R. A. Lazarus, S. Shak, J. S. Ulmer. (Genentech Inc., 2001), vol. US 2001/0041360 A1.
24. 24. C. Q. Pan, R. A. Lazarus, Hyperactivity of human DNase I variants. Dependence on the number of positively charged residues and concentration, length, and environment of DNA. *J Biol Chem* 273, 11701-11708 (1998).
25. 25. C. Q. Pan, D. V. Sinicropi, R. A. Lazarus, Engineered properties and assays for human DNase I mutants. *Methods Mol Biol* 160, 309-321 (2001).
26. 26. C. Q. Pan, J. S. Ulmer, A. Herzka, R. A. Lazarus, Mutational analysis of human DNase I at the DNA binding interface: implications for DNA recognition, catalysis, and metal ion dependence. *Protein Sci* 7, 628-636 (1998).
27. 27. G. Parsiegla, C. Noguere, L. Santell, R. A. Lazarus, Y. Bourne, The structure of human DNase I bound to magnesium and phosphate ions points to a catalytic mechanism common to members of the DNase I-like superfamily. *Biochemistry* 51, 10250-10258 (2012).
28. 28. M. C. Schneider, A. Wilber. (United States, 2004), vol. US 2004/0138156.
29. 29. J. S. Ulmer et al., Engineering actin-resistant human DNase I for treatment of cystic fibrosis. *Proceedings of the National Academy of Sciences of the United States of America* 93, 8225-8229 (1996).

30. 30. B. T. Bettinger, D. M. Gilbert, D. C. Amberg, Actin up in the nucleus. *Nat Rev Mol Cell Biol* 5, 410-415 (2004).
31. 31. C. F. Urban et al., Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex involved in host defense against *Candida albicans*. *PLoS Pathog* 5, e1000639 (2009).
32. 32. D. Russell, N. J. Oldham, B. G. Davis, Site-selective chemical protein glycosylation protects from autolysis and proteolytic degradation. *Carbohydr Res* 344, 1508-1514 (2009).
33. 33. T. A. Fuchs, J. A. Kremer Hovinga, D. Schatzberg, D. D. Wagner, B. Lammle, Circulating DNA and myeloperoxidase indicate disease activity in patients with thrombotic microangiopathies. *Blood* 120, 1157-1164 (2012).
34. 34. M. Jimenez-Alcazar, N. Kim, T. A. Fuchs, Circulating Extracellular DNA: Cause or Consequence of Thrombosis? *Semin Thromb Hemost* 43, 553-561 (2017).
35. 35. M. Boettcher et al., Degradation of Extracellular DNA by DNase1 Significantly Reduces Testicular Damage After Testicular Torsion in Rats. *Urology* 109, 223 e221-223 e227 (2017).
36. 36. A. D. Bandaranayake, S. C. Almo, Recent advances in mammalian protein production. *FEBS letters* 588, 253-260 (2014).
37. 37. Y. K. Han, Y. G. Kim, J. Y. Kim, G. M. Lee, Hyperosmotic stress induces autophagy and apoptosis in recombinant Chinese hamster ovary cell culture. *Biotechnology and bioengineering* 105, 1187-1192 (2010).
38. 38. J. L. Cereghino, J. M. Cregg, Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*. *FEMS microbiology reviews* 24, 45-66 (2000).
39. 39. M. Ahmad, M. Hirz, H. Pichler, H. Schwab, Protein expression in *Pichia pastoris*: recent achievements and perspectives for heterologous protein production. *Applied microbiology and biotechnology* 98, 5301-5317 (2014).
40. 40. R. Daly, M. T. Hearn, Expression of heterologous proteins in *Pichia pastoris*: a useful experimental tool in protein engineering and production. *Journal of molecular recognition* : JMR 18, 119-138 (2005).

Sequences of wild-type DNases

[0196]

SEQ ID NO: 1

Human DNase1 (P24855), Amino acid sequence (Signal Peptide; Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEVDRDS
HLTAVGKLLDNLNQDAPDPTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
SYVRPSQWSSIRLWTSPTTFQWLIPDSADTTATPTHCAVDRIIVAGMLLRGAVVPDSALPFNFQAA
YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 2Human DNase1L3 (Q13609), Amino acid sequence (Signal Peptide: Mature Protein):

MSRELAPLLLLLSIHSALAMRICSFNVRSGESKQEDKNAMDVIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDYQDGDADV
 SREPFVWFQSPHTAVKDFVLIPLHTTPETSVKEIDELVEVYTDVKHRWKAENFIIFMGDFNAGCS
 YVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEIIVSSVVPKSNVDFQKA
 YKLTETEEALDVSDHFPVEFKLQSSRAFTNSKKSVTLRKKTKSKRS

SEQ ID NO: 3Murine DNase1 (P49183): Amino acid sequence (Signal Peptide: Mature Protein):

MRYTGLMGILLTLVNLLQLAGTLRIA AFNIRTFGETKMSNATLSVYFVKILSRDYDIAVIEVRDS
 HLVAVGKLLDELNRDKPDITYRVVSEPLGRKSYKEQYLFVYRPDQVSILDSYQYDDGCEPCGNDT
 FSREPAIVKFFSPYTEVQEF AIVPLHAAPTEAVSEIDALYDVYLDVWQKWGLEDIMFMGDFNAGC
 SYVTSSQWSSIRLRTSPIFQWLIPDSADTTVTSTHCA YDRIVVAGALLQAAVVPNSAVPFDQAE
 YGLSNQLAEAISDHYPVEVTLRKI

SEQ ID NO: 4Rat DNase1 (P21704), Amino acid sequence (Signal Peptide: Mature Protein):

MRYTGLMGILLTLVNLLQLAATLRIA AFNIRTFGDTKMSNATLSSYIVKILSRDYDIAVVQEV RDT
 HLVAVGKLLDELNRDIPDNYRYIISEPLGRKSYKEQYLFVYRPSQVSVLDSYHYDDGCEPCGNDT
 FSREPAIVKFFSPYTEVREF AIVPLHSAPTEAVSEIDALYDVYLDVRQKWGLEDIMFMGDFNAGC
 SYVTSSQWSSIRLRTSPIFQWLIPDSADTTATSTHCA YDRIVVAGALLQAAVVPSSAVPFDQAE
 YRLTNQMAE AISDHYPVEVTLRKT

SEQ ID NO: 34Murine DNase1L3 (O55070): Amino acid sequence (Signal Peptide: Mature Protein):

MSLHPASPRLASLLLFILALHDTLALRLCSFNVRSGASKKENHEAMDIIVKIIKRCDLILLMEI
 KDSNNNICPMLMEKLNNSRRSTTYNYVISSRLGRNTYKEQYAFVYKEKLVSVKTKYHYHDYQDG
 DTDVFSREPFVWFHSPFTAVKDFVIVPLHTTPETSVKEIDELVDVYTDVRSQWK TENFIIFMGDF
 NAGCSYVPKKAWQNIRLRTDPKFVWLIGDQEDTTVKKSTSCAYDRIVLCGQEIIVNSVVP RSSGVF
 DFQKAYDLSEEEALDVSDHFPVEFKLQSSRAFTNNRKS VSLKKRKKGNRS

SEQ ID NO: 35Rat DNase1L3 (O89107): Amino acid sequence (Signal Peptide: Mature Protein):

MSLYPASPYLASLLLFILALHGALSRLCSFNVRSGESKKENHNAMDIIVKIIKRCDLILLMEI
 KDSNNNICPMLMEKLNNSRRSTTYNYVISSRLGRNTYKEQYAFLYKEKLVSVKAKYLYHDYQDG
 DTDVFSREPFVWFQAPFTA AKDFVIVPLHTTPETSVKEIDELADVYTDVRRRWKAENFIIFMGDF
 NAGCSYVPKKAWKNIRLRTDPNFVWLIGDQEDTTVKKSTSCAYDRIVLRGQEIIVNSVVP RSSGVF
 DFQKAYELSEEEALDVSDHFPVEFKLQSSRAFTNSRKS VSLKKKKKGSRS

SEQ ID NO: 36Chimpanzee DNase1 (H2QAH1): Amino acid sequence (Signal Peptide: Mature Protein):

MRSMKLLGALLAALLQGA VSLKIAAFNIQTGETKMSNATLSYIVQILSRDYDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT

FNREPAIVRFFSRFTEVREFAIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLRTSPAQWLIPDSADTTATPTHCAVDRIIVAGMLLQGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 37

Chimpanzee DNase1L3 (H2QMU7): Amino acid sequence (Signal Peptide; Mature Protein):

MSRELTPLLLLLSIHSTLALRICSFNVRSEGESKQEDQNAMDVIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDYQDGDADV
 SREPFVWFQSPHTAVKDFVIIPLHTTPETSVKEIDELVEVYTDVKHRWKAENFI FMGDFNAGCS
 YVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEI VSSVVPKSN SVDFDQKA
 YKLTEEEALDVSDHFPVEFKLQSSRAFTNSKKSVTLRKKTKSKRS

SEQ ID NO: 38

Human DNase1-like 1 (NM_006730): Amino acid sequence (Signal Peptide; Mature Protein):

MHYPTALLFLILANGAQAFRICAFNAQRLTLAKVAREQVMDTLVRILARCDIMVLQEVVDSSGSA
 IPLLLRELNRFDGSGPYSTLSSPQLGRSTYMETVYFYRSHKTQVLSSYVYNDEDDVFAREFFVA
 QFSLPSNVLP SLVLVPLHTTPKAVEKELNALYDVFLVSVQHWQSKDVILLGDFNADCASLTKKRL
 DKLELRTEPGFHWVIADGEDTTVRASTHCTYDRVVLHGERCRSLLHTAAAFDFTSFQLTEEEAL
 NISDHYPVEVELKLSQAHSVQPLSLTVLLLLLSLLSPQLCPAA

SEQ ID NO: 39

Human DNase1-like 2 (NM_001374): Amino acid sequence (Signal Peptide; Mature Protein):

MGGPRALLAALWALEAAGTAALRIGAFNIQSFGDSKVSDPACGSI IAKILAGYDLALVQEVDPD
 LSAVSALMEQINSVSEHEYSFVSSQPLGRDQYKEMYLFVYRKDAVSVDVLYLPDPEDVFSREPF
 VVKFSAPGTGERAPPLPSRRALTPPPLPAAAQNLVLIPLHAAPHQAVAEIDALYDVYLDVIDKWG
 TDDMLFLGDFNADCSYVRAQDWAAIRLSSEVFkwLIPDSADTTVGNSDCAYDRIVACGARLRS
 LKPQSATVHDFQEEFGLDQTQALAI SDHFPVEVTLKFHR

C-terminal tail of human DNase1L3**SEQ ID NO: 33**

[0197] Amino acid sequence:

SSRAFTNSKKSVTLRKKTKSKRS

Sequences of DNase1 variants

[0198]

SEQ ID NO: 5

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIRTFGETKMSNATLVSYIVKILSRydIALVQEVrDS
 HLTAVGKLLDNLNRDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLRTSPTFQWLIPDSADTTATPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 6**Amino acid sequence (Signal Peptide: Mature Protein):**

MRGMKLLGALLALAALLQGAVSLKIAAFNIRTFGETKMSNATLVSYIVKILSRydIALVQEVrDS
 HLTAVGKLLDNLNRDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLRTSPTFQWLIPDSADTTATPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQAA
 YKLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 7**Amino acid sequence (Signal Peptide: Mature Protein):**

MRGMKLLGALLALAALLQGAVSLKIAAFNIRTFGETKMSNATLVSYIVQILSRydIALVQEVrDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPFIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTAKPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 8**Amino acid sequence (Signal Peptide: Mature Protein):**

MRGMKLLGALLALAALLQGAVSLKIAAFNIRTFGETKMSNATLVSYIVQILSRydIALVQEVrDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTAKPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 9**Amino acid sequence (Signal Peptide: Mature Protein):**

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTTFGETKMSNATLVSYIVQILSRydIALVQEVrDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPFIVRFFSRFTEVREFAIIVPLHAAPGDAVKEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTATPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 10**Amino acid sequence (Signal Peptide: Mature Protein):**

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTTFGETKMSNATLVSYIVQILSRydIALVQEVrDS
 HLTAVGKLLDNLNRNSRRGITYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGN
 DTFNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNA
 GCSYVRPSQWSSIRLWTSPTFQWLIPDSADTTATPTHCAyDRIVVAGMLLRGAVVPDSALPFNFQ

AAAYGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 11

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYNYVISSRLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTATPTHCAYDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 12

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPFVVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTATPTHCAYDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 13

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVPPKAWKNIRLRTDPRFVWLIGDQEDTTATPTHCAYDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 14

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLIPDSADTTVKKSTHCAYDRIVVAGMLLRGAVVPDSALPFNFQAA
 AYGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 15

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNRNSRRGITNYVISSRLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT
 DTFNREPFVVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 GCSYVPPKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTHCAYDRIVVAGMLLRGAVVPDSALPFNF
 QAAYGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 16

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFGETKMSNATLVSYIVQILSRDIALVQEV RDS
 HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGNDT

FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDVMLMGDFNAGC
 SYVRPSQWSSIRLWTSPTFQWLI PDSADTTATPTHCA YDRIVVAGMLLRGAVVPDSALPFNFQAA
 YGLSDQLAQAI SDHFPVEFKLQSSRAFTNSKKS VTLRKKTKSKRS

Sequences of DNaseII3 variants

[0199]

SEQ ID NO: 17

Amino acid sequence (Signal Peptide: Mature Protein):

MSRELAPLLLLLLSIHSALAMRICSFNVRSGESKQEDKNAMDVIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRRNSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDYQDGDADV
 SREPFVWFQSPHTAVKDFV I I PLHTTPETSVKEIDELVEVYTDVKHRWKAENFI FMGDFNAGCS
 YVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEIVSSVVPKSN SVDFDQKA
 YKLTEEEALDVSDHYPVEVMLK

SEQ ID NO: 18

Amino acid sequence (Signal Peptide: Mature Protein):

MSRELAPLLLLLLSIHSALAMRICSFNVRSGESKQENKTAMDVIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRRNSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDYQDGDNDTF
 SREPFVWFQSPHTAVKDFV I I PLHTTPETSVKEIDELVEVYTDVKHRWKAENFI FMGDFNAGCS
 YVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEIVSSVVPKSN SVDFDQKA
 YKLTEEEALDVSDHFPVEFKLQSSRAFTNSKKS VTLRKKTKSKRS

SEQ ID NO: 19

Amino acid sequence (Signal Peptide: Mature Protein):

MSRELAPLLLLLLSIHSALAMRICSFNVRSGESKM NATLVSYIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRRNSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDYQDGDADV
 SREPFVWFQSPHTAVKDFV I I PLHTTPETSVKEIDELVEVYTDVKHRWKAENFI FMGDFNAGCS
 YVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEIVSSVVPKSN SVDFDQKA
 YKLTEEEALDVSDHFPVEFKLQSSRAFTNSKKS VTLRKKTKSKRS

SEQ ID NO: 20

Amino acid sequence (Signal Peptide: Mature Protein):

MSRELAPLLLLLLSIHSALAMRICSFNVRSGESKQEDKNAMDVIVKVIKRCDIILVMEIKDSNN
 RICPILMEKLNRRNSRRGITYNVVISSRLGRNTYKEQYAFLYKEKLVSVKRSYHYHDGCEPCGNDT
 FNREPFVWFQSPHTAVKDFV I I PLHTTPETSVKEIDELVEVYTDVKHRWKAENFI FMGDFNAGC
 SYVPKKAWKNIRLRTDPRFVWLIGDQEDTTVKKSTNCAYDRIVLRGQEIVSSVVPKSN SVDFDQK
 AYKLTEEEALDVSDHFPVEFKLQSSRAFTNSKKS VTLRKKTKSKRS

Sequences used in *Pichia pastoris*

[0200]

SEQ ID NO: 21

Amino acid sequence (Signal Peptide: Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIQTFTGETKMSSATLVSYIVQILSRDYDIALVQEVDRS
HLTAVGKLLDNLNQDAPDTHYVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGSDT
FNREPAIVRFFSRFTEVREFAIIVPLHAAPGDAVAEIDALYDVYLDVQEKWGLEDMMLMGDFNAGC
SYVRPSQWSSIRLWTSPTFQWLIPDSADTTATPTHCAYDRIVVAGMLLRGAVVPDSALPFNFQAA
YGLSDQLAQAI SDHYPVEVMLK

SEQ ID NO: 22

Codon-optimized cDNA of SEQ ID NO: 21 (Start codon, Signal peptide):

ATGAGAGGTATGAAGCTCCTTGGTGGCTCCTGGCTCTCGCTGCTCTCCTCCAAGGAGCAGATTTG
TCTAAAGATTGCTGCTTTCAATATTCAAACCTTTGGCGAGACTAAAATGTCTCTCGGCAACATTAG
TATCTTACATTGTTTACAGATCCTTTCTCGTTACGACATAGCTTTGGTCCAGGAGGTTAGAGACTCC
CACTTGACTGCCGTCGGTAAGCTGCTAGACAATCTTAACCAAGACGCACCTGACACCTACCACTA
CGTTGTTAGTGAGCCTCTCGGTAGAACTCCTACAAGGAGAGATACCTATTTGTCTACCGTCCAG
ATCAGGTGTCTGCTGTGGACTCTTACTACTATGACGACGGTTGTGAACCTTGTGGTTCAGACACC
TTCAACAGAGAACCAGCTATCGTTAGATTTTTTCTCCAGATTACCCGAGGTCAGAGAGTTCCGCAT
CGTTCCATTGCACGCTGCGCCTGGAGATGCAGTGGCCGAAATTGACGCTCTCTATGATGTCTACC
TGGACGTTTCAGGAAAAATGGGGTCTAGAAGATGTTATGCTGATGGGAGACTTCAACGCTGGTTGC
TCTTACGTTAGGCCATCTCAATGGTCAAGCATCAGACTATGGACTTCCCCAACGTTCCAATGGCT
TATTCCTGACTCCGCTGATACTACCGCCACTCCCCTCATTGTGCATATGACAGAATTGTTGTCTG
CTGGTATGCTACTGCGTGGAGCTGTTCGTACCAGATTCTGCTCTGCCATTTAACTTCCAAGCCGCA
TATGGTCTTTCTGACCAACTGGCTCAAGCCATCTCTGATCACTACCCTGTGGAAGTTATGCTTAA

SEQ ID NO: 23

Amino acid sequence (Signal Peptide; Mature Protein):

MRGMKLLGALLALAALLQGAVSLKIAAFNIRTFGETKMSSATLVSYIVQILSRDYIALVQEV RDS
HLTAVGKLLDNLNQDAPDTYHYVVSEPLGRNSYKERYLFVYRPDQVSAVDSYYYDDGCEPCGSDT
FNREPFIVRFFSRFTEVREFAI VPLHAAPGDAVAEIDALYDVYLDVQEKWGLE DVMLMGDFNAGC
SYVRPSQWSSI RLWTSPTFQWLIPDSADTTAKPTHCA YDRIVVAGMLLRGAVVPDSALPFNFQAA
YGLSDOLAQAISDHYPVEVMLK

SEQ ID NO: 24

Codon-optimized cDNA of SEQ ID NO: 23 (Start codon, Signal peptide):

ATGAGAGGATATGAAGCTCCTTGGTGCCCTCCTGGCTCTCGCTGCTCTCCTCCAAGGAGCAGTTTC
TCTAAAGATTGCTGCTTTCAATATTAGAACCTTTGGCGAGACTAAAATGTCTCTCGGCAACATTAG
TATCTTACATTGTTTACAGATCCTTTCTCGTTACGACATAGCTTTGGTCCAGGAGGTTAGAGACTCC
CACTTGACTGCCGTCCGTAAGCTGCTAGACAATCTTAACCAAGACGCACCTGACACCTACCACTA
CGTTGTTAGTGAGGCTCTCGGTAGAACTCCTAGCAAGGAGAGATAGCTATTTGCTCTACGCTCAG

CGTTGTTAGTGAAGCTCTCGGTAGAACTCTTACAGGAGAGATACCTATTGTCTACCGTCCAG
 ATCAGGTGTCTGCTGTGGACTCTTACTACTATGACGACGGTTGTGAACCTTGTGGTTTCAGACACC
 TTCAACAGAGAACCATTTCATCGTTAGATTTTTTCTCCAGATTCACCGAGGTCAGAGAGTTTCGCCAT
 CGTTCCATTGCACGCTGCGCCTGGAGATGCAGTGGCCGAAATTGACGCTCTCTATGATGTCTACC
 TGGACGTTTCAGGAAAAATGGGGTCTAGAAGATGTTATGCTGATGGGAGACTTCAACGCTGGTTGC
 TCTTACGTTAGGCCATCTCAATGGTCAAGCATCAGACTATGGACTTCCCCAACGTTCCAATGGCT
 TATTCCTGACTCCGCTGATACTACCGCCAAGCCACTCATTGTGCATATGACAGAATTGTTGTGC
 CTGGTATGCTACTGCGTGGAGCTGTGCTACCAGATTCTGCTCTGCCATTTAACTTCCAAGCCGCA
 TATGGTCTTTCTGACCAACTGGCTCAAGCCATCTCTGATCACTACCCTGTGGAAGTTATGCTTAA
 G

SEQ ID NO: 25

Codon-optimized cDNA of SEQ ID NO: 2 (Start codon, Signal peptide):

ATGTCTAGAGAGCTTGCTCCACTCCTCCTACTCCTTCTTAGTATCCACTCTGCACTCGCCATGAG
 AATTTGTAGCTTCAATGTAAGATCCTTCGGTGAATCTAAGCAGGAGGACAAAAACGCTATGGACG
 TTATTGTGAAGGTCATTAAGAGATGTGATATCATTTCTAGTTATGGAGATCAAGGACTCTAACAAC
 AGAATTTGCCCCATCCTTATGGAAAAGCTGAATAGAACTCTAGAAGAGGAATTACTTACAACATA
 CGTTATCTCTTCTAGGCTGGGTAGAAACACTTACAAGGAGCAATATGCATTTCTATAACAAGGAAA
 AGCTAGTTTCCGTTAAGCGTTCTTACCCTATCATGACTACCAAGACGGCGATGCTGACGTTTTTC
 TCCAGAGAACCATTTCGTTGTTTGGTTTTCAGTCTCCTCACACCGCTGTTAAGGACTTCGTGATTAT
 CCCACTACACACTACGCCAGAAACCTCTGTCAAGGAAATAGATGAACTTGTTGAAGTTTACACCG
 ACGTGAAGCACAGATGGAAGGCCGAGAATTTCAATTTTCATGGGTGATTTCAACGCCGGATGCTCA
 TATGTCCCTAAAAAGGCTTGGA AAAACATCCGTTTGAGAACCGATCCTAGATTTGTCTGGCTCAT
 CGGTGACCAAGAGGACACCACAGTCAAAAAGTCTACCAACTGTGCTTACGACAGAATTGTTCTGC
 GTGGTCAGGAGATTGTTTCATCTGTTGTCCCAAAGTCCAACCTCCGTCTTTGATTTCCAAAAGGCT
 TACAACTGACTGAGGAGGAAGCTTTAGACGTGTCCGACCACTTCCCTGTAGAGTTTAAGCTGCA
 ATCCTCCAGAGCATTCACTAACTCTAAAAAGTCAGTCACTTTGCCTAAAAAGACTAAGTCTAAGA
 GATCG

SEQ ID NO: 26

Codon-optimized cDNA of SEQ ID NO: 17 (Start codon, Signal peptide):

ATGTCTAGAGAGCTTGCTCCACTCCTCCTACTCCTTCTTAGTATCCACTCTGCACTCGCCATGAG
 AATTTGTAGCTTCAATGTAAGATCCTTCGGTGAATCTAAGCAGGAGGACAAAAACGCTATGGACG
 TTATTGTGAAGGTCATTAAGAGATGTGATATCATTTCTAGTTATGGAGATCAAGGACTCTAACAAC
 AGAATTTGCCCCATCCTTATGGAAAAGCTGAATAGAACTCTAGAAGAGGAATTACTTACAACATA
 CGTTATCTCTTCTAGGCTGGGTAGAAACACTTACAAGGAGCAATATGCATTTCTATAACAAGGAAA
 AGCTAGTTTCCGTTAAGCGTTCTTACCCTATCATGACTACCAAGACGGCGATGCTGACGTTTTTC
 TCCAGAGAACCATTTCGTTGTTTGGTTTTCAGTCTCCTCACACCGCTGTTAAGGACTTCGTGATTAT
 CCCACTACACACTACGCCAGAAACCTCTGTCAAGGAAATAGATGAACTTGTTGAAGTTTACACCG
 ACGTGAAGCACAGATGGAAGGCCGAGAATTTCAATTTTCATGGGTGATTTCAACGCCGGATGCTCA
 TATGTCCCTAAAAAGGCTTGGA AAAACATCCGTTTGAGAACCGATCCTAGATTTGTCTGGCTCAT
 CGGTGACCAAGAGGACACCACAGTCAAAAAGTCTACCAACTGTGCTTACGACAGAATTGTTCTGC

GTGGTCAGGAGATTGTTTCATCTGTTGTCCCAAAGTCCAACCTCCGTCTTTGATTTCCAAAAGGCT
TACAAACTGACTGAGGAGGAAGCTTTAGACGTGTCCGACCACTACCCTGTAGAGGTCATGTTGAA
G

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US62611166B \[0001\]](#)
- [US62547220B \[0001\]](#)
- [US9458218B \[0053\]](#)
- [WO2015066550A \[0053\]](#)
- [US9221896B \[0053\]](#)
- [US20140199329A \[0094\]](#)
- [EP17152528 \[0100\]](#)
- [WO2016118476A \[0109\]](#)
- [US9642822B \[0109\] \[0168\]](#)
- [EP2018051444W \[0171\]](#)
- [US9642822B2 \[0195\]](#)
- [US6391607B1 \[0195\]](#)
- [US7407785B2 \[0195\]](#)
- [US20010041360A1 \[0195\]](#)
- [US20040138156A \[0195\]](#)

Non-patent literature cited in the description

- **JIMÉNEZ-ALCÁZAR et al.**Journal of Thrombosis and Haemostasis, 2015, vol. 13, 5732-742 [0004]
- **KARLINALTSCHUL**Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 5873-5877 [0032]
- **THOMPSON, J. D.HIGGINS, D. G.GIBSON, T. J.**Nucleic Acids Res., 1994, vol. 22, 4673-80 [0032]
- **ALTSCHUL et al.**J. Mol. Biol., 1990, vol. 215, 403-410 [0032]
- **HAKKIM et al.**Proc. Natl. Acad. Sci USA, 2010, vol. 107, 219813-9818 [0056]
- **NAPIREI et al.**Biochem J, 2005, vol. 389, 355-364 [0056]
- **FUCHS et al.**J Cell Biol, 2007, vol. 176, 2231-241 [0056]
- **SISIRAK et al.**Cell, 2016, vol. 166, 88-101 [0056]
- **JIMÉNEZ-ALCÁZAR et al.**J Thromb Haemost, 2015, vol. 13, 732-742 [0056]
- **ULMER et al.**PNAS USA, 1996, vol. 93, 8225-8229 [0087]
- **WILBER et al.**Mol Ther, 2002, vol. 6, 35-42 [0090]
- **PAN et al.**J Biol Chem, 1998, vol. 273, 11701-11708 [0094]
- **NUNES-DUBY et al.**Nucleic Acids Res, 1998, vol. 26, 391-406 [0125]
- **SADWOSKI**J. Bacteriol., 1986, vol. 165, 341-357 [0125]
- **BESTOR**Cell, 2005, vol. 122, 3322-325 [0125]
- **PLASTERK et al.**, TIG, 1999, vol. 15, 326-332 [0125]
- **KOOTSTRA et al.**Ann. Rev. Pharm. Toxicol., 2003, vol. 43, 413-439 [0125] [0125]
- **STERNBERGHAMILTON**J. Mol. Biol., 1981, vol. 150, 467-486 [0125]
- **NASH**Nature, 1974, vol. 247, 543-545 [0125]
- **BROACH et al.**Cell, 1982, vol. 29, 227-234 [0125]
- **MATSUZAKI et al.**J. Bacteriology, 1990, vol. 172, 610-618 [0125]
- **GROTH et al.**J. Mol. Biol., 2004, vol. 335, 667-678 [0125]
- Acute and chronic inflammation.V. **KUMARA. K. ABBASN. FAUSTOJ. C. ASTER**Robbins & Cotran Pathologic Basis of Disease.Elsevier Health Sciences20090000 [0161]
- **V. BRINKMANNU. REICHARD C. GOOSMANNB. FAULERY. UHLEMANND. S. WEISSY. WEINRAUCHA. ZYCHLINSKY**Neutrophil extracellular traps kill bacteria.Science, 2004, vol. 303, 1532-1535 [0161]
- **P. LIM. LIM. R. LINDBERGM. J. KENNETTN. XIONGY. WANG**PALM is essential for antibacterial innate immunity mediated by neutrophil extracellular traps.J. Exp. Med., 2010, vol. 207, 1853-1862 [0161]
- **V. THAMMAVONGSAD. M. MISSIAKASO. SCHNEEWIND**Staphylococcus aureus degrades neutrophil extracellular traps to promote immune cell death.Science, 2013, vol. 342, 863-866 [0161]
- **S. R. CLARKA. C. MAS. A. TAVENERB. MCDONALDZ. GOODARZIM. M. KELLYK. D. PATELS. CHAKRABARTIE. MCAVOYG. D. SINCLAIR**Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood.Nat. Med., 2007, vol. 13, 463-469 [0161]
- **T. A. FUCHSA. BRILLD. DUERSCHMIEDD. SCHATZBERGM. MONESTIERD. D. MYERS, JR.S. K. WROBLESKIT. W. WAKEFIELDJ. H. HARTWIGD. D. WAGNER**Extracellular DNA traps promote thrombosis.Proc. Natl. Acad. Sci. U. S. A., 2010, vol. 107, 15880-15885 [0161]

- **A. WARNATSCHM. IOANNOUQ. WANGV. PAPAYANNOPOULOS**Neutrophil extracellular traps license macrophages for cytokine production in atherosclerosis.Science, 2015, vol. 349, 316-320 [\[0161\]](#)
- **B. ENGELMANN S. MASSBERG**Thrombosis as an intravascular effector of innate immunity.Nat. Rev. Immunol, 2013, vol. 13, 34-45 [\[0161\]](#)
- **S. K. JORCHP. KUBES**An emerging role for neutrophil extracellular traps in noninfectious disease.Nat. Med., 2017, vol. 23, 279-287 [\[0161\]](#)
- **A. HAKKIMB.G. FURNROHRK. AMANNB. LAUBEU.A. ABEDV. BRINKMANNM. HERRMANNR.E. VOLLA. ZYCHLINSKY**Impairment of neutrophil extracellular trap degradation is associated with lupus nephritis.Proc. Natl. Acad. Sci. U. S. A., 2010, vol. 107, 9813-9818 [\[0161\]](#)
- **M. NAPIREIS. LUDWIGJ. MEZRHABT. KLOCKLH. G. MANNHERZ**Murine serum nucleasescontrasting effects of plasmin and heparin on the activities of DNase1 and DNase1-like 3 (DNase1l3).FEBS J., 2009, vol. 276, 1059-1073 [\[0161\]](#)
- **M. NAPIREIA. RICKEND. EULITZH. KNOOPH. G. MANNHERZ**Expression pattern of the deoxyribonuclease 1 gene: lessons from the Dnase1 knockout mouse.Biochem. J., 2004, vol. 380, 929-937 [\[0161\]](#)
- **V. SISIRAKB. SALLYV. D'AGATI W. MARTINEZ-ORTIZZ. B. OZCAKARJ. DAVIDA. RASHIDFARROKHIA. YESTEC. PANEAA. S. CHIDA**Digestion of Chromatin in Apoptotic Cell Microparticles Prevents Autoimmunity.Cell, 2016, vol. 166, 88-101 [\[0161\]](#)
- **M. DEMERSD. S. KRAUSED. SCHATZBERGK. MARTINODJ. R. VOORHEEST. A. FUCHSD. T. SCADDEND. D. WAGNER**Cancers predispose neutrophils to release extracellular DNA traps that contribute to cancer-associated thrombosis.Proc. Natl. Acad. Sci. U. S. A., 2012, vol. 109, 13076-13081 [\[0161\]](#)
- **T. A. FUCHSU. ABEDC. GOOSMANNR. HURWITZI. SCHULZE V. WAHNY. WEINRAUCHV. BRINKMANNA. ZYCHLINSKY**Novel cell death program leads to neutrophil extracellular traps.J. Cell Biol., 2007, vol. 176, 231-241 [\[0161\]](#)
- **C. FRANKD. WERBERJ. P. CRAMERM. ASKARM. FABERM. AN DER HEIDENH. BERNARDA. FRUTHR. PRAGERA. SPODEHUS**Investigation Team, Epidemic profile of Shiga-toxin-producing Escherichia coli O104:H4 outbreak in Germany.N. Engl. J. Med., 2011, vol. 365, 1771-1780 [\[0161\]](#)
- **S. A. BRAUNED. WICHMANNM. C. VON HEINZA. NIERHAUSH. BECKERT. N. MEYERG. P. MEYERM. MULLER-SCHULZJ. FRICKEA. DE WEERTH**Clinical features of critically ill patients with Shiga toxin-induced hemolytic uremic syndrome.Crit. Care. Med., 2013, vol. 41, 1702-1710 [\[0161\]](#)
- **M. JIMENEZ-ALCAZARM. NAPIREIR. PANDAE. C. KOHLERJ. A. KREMER HOVINGAH. G. MANNHERZS. PEINET. RENNEB. LAMMLET. A. FUCHS**Impaired DNase1-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies.J. Thromb. Haemost., 2015, vol. 13, 732-742 [\[0161\]](#)
- **J. LEFFLERZ. PROHASZKAB. MIKESG. SINKOVITSK. CIACMAP. FARKASM. RETIK. KELENG. S. REUSZA. J. SZABO**Decreased neutrophil extracellular trap degradation in shiga toxin-associated haemolytic uraemic syndrome.J. Innate Immun., 2017, vol. 9, 12-21 [\[0161\]](#)
- **M. LEPPKESC. MAUERODERS. HIRTHS. NOWECKIC. GUNTHERU. BILLMEIERS.**

- PAULUSM. BIERMANNL. E. MUNOZM. HOFFMANNExternalized decondensed neutrophil chromatin occludes pancreatic ducts and drives pancreatitis.Nat. Commun., 2016, vol. 7, 10973- [0161]
- C. SCHAUERC. JANKOL. E. MUNOZY. ZHAOD. KIENHOFERB. FREYM. LELLB. MANGERJ. RECHE. NASCHBERGERAggregated neutrophil extracellular traps limit inflammation by degrading cytokines and chemokines.Nat. Med., 2014, vol. 20, 511-517 [0161]
 - Hemodynamic disorders, thromboembolic disease, and shock.V. KUMARA. K. ABBASN. FAUSTOJ. C. ASTERRobbins & Cotran Pathologic Basis of DiseaseElsevier Health Sciences20090000 [0161]
 - M. AL-MAYOUFA. SUNKERR. ABDWANIS.A. ABRAWIF. ALMURSHEDIN. ALHASHMIA. AL SONBULW. SEWAIRIA. QARIE. ABDALLAHLoss-of-function variant in DNASE1L3 causes a familial form of systemic lupus erythematosus.Nat. Genet, 2011, vol. 43, 1186-1188 [0161]
 - K. YASUTOMOT. HORIUCHIS. KAGAMIH. TSUKAMOTOC. HASHIMURAM. URUSHIHARAY. KURODA.Mutation of DNASE1 in people with systemic lupus erythematosus.Nat. Genet., 2001, vol. 28, 313-314 [0161]
 - M. NAPIREIH. KARSUNKYB. ZEVIKH. STEPHANH. G. MANNHERZT. MOROYFeatures of systemic lupus erythematosus in Dnase1-deficient mice.Nat. Genet., 2000, vol. 25, 177-181 [0161]
 - S. GUPTAM. J. KAPLANThe role of neutrophils and NETosis in autoimmune and renal diseases.Nat. Rev. Nephrol., 2016, vol. 12, 402-413 [0161]
 - R. MIZUTAS. ARAKIM. FURUKAWAY. FURUKAWAS. EBARAD. SHIOKAWAK. HAYASHIS. TANUMAD. KITAMURADNase gamma is the effector endonuclease for internucleosomal DNA fragmentation in necrosisPloS one, 2013, vol. 8, e80223- [0161]
 - J. ROSSAJTJ. M. HERTERH. VAN AKENM. NAPIREIY. DORINGC. WEBERO. SOEHNLEINA. ZARBOCKSynchronized integrin engagement and chemokine activation is crucial in neutrophil extracellular trap-mediated sterile inflammation.Blood, 2014, vol. 123, 2573-2584 [0161]
 - A. BRADLEYK. ANASTASSIADISA. AYADIJ. F. BATTEYC. BELLM. C. BIRLINGJ. BOTTOMLEYS. D. BROWNA. BURGERC. J. BULTThe mammalian gene function resource: the International Knockout Mouse Consortium.Mamm. Genome, 2012, vol. 23, 580-586 [0161]
 - V. KUMARA. K. ABBASN. FAUSTOJ. C. ASTERRobbins & Cotran Pathologic Basis of Disease.Elsevier Health Sciences20090000 [0195]
 - V. BRINKMANN et al.Neutrophil extracellular traps kill bacteria.Science (New York, N.Y.), 2004, vol. 303, 1532-1535 [0195]
 - T. A. FUCHSA. HAKKIMC. F. URBANInflammation: Fundamental Mechanisms,World Scientific20180000 [0195]
 - D. SHIOKAWAS. TANUMACHaracterization of human DNase I family endonucleases and activation of DNase gamma during apoptosisBiochemistry, 2001, vol. 40, 143-152 [0195]
 - M. NAPIREIA. RICKEND. EULITZH. KNOOPH. G. MANNHERZExpression pattern of the deoxyribonuclease 1 gene: lessons from the Dnase1 knockout mouse.Biochem J, 2004,

vol. 380, 929-937 [0195]

- **W. F. BARON et al.** Cloning and characterization of an actin-resistant DNase I-like endonuclease secreted by macrophages. *Gene*, 1998, vol. 215, 291-301 [0195]
- **V. SISIRAK et al.** Digestion of Chromatin in Apoptotic Cell Microparticles Prevents Autoimmunity. *Cell*, 2016, vol. 166, 88-101 [0195]
- **M. JIMENEZ-ALCAZAR et al.** Impaired DNase1-mediated degradation of neutrophil extracellular traps is associated with acute thrombotic microangiopathies. *Journal of thrombosis and haemostasis : JTH*, 2015, vol. 13, 732-742 [0195]
- **M. NAPIREIS. LUDWIGJ. MEZRHABT. KLOCKLH. G. MANNHERZ** Murine serum nucleases--contrasting effects of plasmin and heparin on the activities of DNase1 and DNase1-like 3 (DNase1I3). *FEBS J*, 2009, vol. 276, 1059-1073 [0195]
- **A. WILBERM. LUM. C. SCHNEIDER** Deoxyribonuclease I-like III is an inducible macrophage barrier to liposomal transfection. *Mol Ther*, 2002, vol. 6, 35-42 [0195]
- **W. KABSCHH. G. MANNHERZD. SUCKE. F. PAIK. C. HOLMES** Atomic structure of the actin:DNase I complex. *Nature*, 1990, vol. 347, 37-44 [0195]
- **C. LAM et al.** Taming hyperactive hDNase I: Stable inducible expression of a hyperactive salt- and actin-resistant variant of human deoxyribonuclease I in CHO cells. *Biotechnol Prog*, 2017, vol. 33, 523-533 [0195]
- **C. Q. PAN et al.** Improved potency of hyperactive and actin-resistant human DNase I variants for treatment of cystic fibrosis and systemic lupus erythematosus. *J Biol Chem*, 1998, vol. 273, 18374-18381 [0195]
- **D. SHIOKAWAY. SHIKAS. TANUMA** Identification of two functional nuclear localization signals in DNase gamma and their roles in its apoptotic DNase activity. *Biochem J*, 2003, vol. 376, 377-381 [0195]
- **R. MIZUTA et al.** DNase gamma is the effector endonuclease for internucleosomal DNA fragmentation in necrosis. *PLoS One*, 2013, vol. 8, e80223- [0195]
- **H. J. FUCHS et al.** Effect of aerosolized recombinant human DNase on exacerbations of respiratory symptoms and on pulmonary function in patients with cystic fibrosis. The Pulmozyme Study Group. *N Engl J Med*, 1994, vol. 331, 637-642 [0195]
- **N. M. RAHMAN et al.** Intrapleural use of tissue plasminogen activator and DNase in pleural infection. *N Engl J Med*, 2011, vol. 365, 518-526 [0195]
- **V. PAPAYANNOPOULOSD. STAABA. ZYCHLINSKY** Neutrophil elastase enhances sputum solubilization in cystic fibrosis patients receiving DNase therapy. *PLoS One*, 2011, vol. 6, e28526- [0195]
- **M. NAPIREIS. WULFH. G. MANNHERZ** Chromatin breakdown during necrosis by serum Dnase1 and the plasminogen system. *Arthritis Rheum*, 2004, vol. 50, 1873-1883 [0195]
- **C. Q. PANR. A. LAZARUS** Hyperactivity of human DNase I variants. Dependence on the number of positively charged residues and concentration, length, and environment of DNA. *J Biol Chem*, 1998, vol. 273, 11701-11708 [0195]
- **C. Q. PAND. V. SINICROPIR. A. LAZARUS** Engineered properties and assays for human DNase I mutants. *Methods Mol Biol*, 2001, vol. 160, 309-321 [0195]
- **C. Q. PANJ. S. ULMERA. HERZKAR. A. LAZARUS** Mutational analysis of human DNase I at the DNA binding interface: implications for DNA recognition, catalysis, and metal ion dependence. *Protein Sci*, 1998, vol. 7, 628-636 [0195]

- **G. PARSIEGLAC. NOGUEREL. SANTELLR. A. LAZARUSY. BOURNE**The structure of human DNase I bound to magnesium and phosphate ions points to a catalytic mechanism common to members of the DNase I-like superfamily.Biochemistry, 2012, vol. 51, 10250-10258 [0195]
- **J. S. ULMER et al.**Engineering actin-resistant human DNase I for treatment of cystic fibrosis.Proceedings of the National Academy of Sciences of the United States of America, 1996, vol. 93, 8225-8229 [0195]
- **B. T. BETTINGERD. M. GILBERTD. C. AMBERG**Actin up in the nucleus.Nat Rev Mol Cell Biol, 2004, vol. 5, 410-415 [0195]
- **C. F. URBAN et al.**Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex involved in host defense against *Candida albicans*.PLoS Pathog, 2009, vol. 5, e1000639- [0195]
- **D. RUSSELLN. J. OLDHAMB. G. DAVIS**Site-selective chemical protein glycosylation protects from autolysis and proteolytic degradation.Carbohydr Res, 2009, vol. 344, 1508-1514 [0195]
- **T. A. FUCHSJ. A. KREMER HOVINGAD. SCHATZBERGD. D. WAGNERB. LAMMLE**Circulating DNA and myeloperoxidase indicate disease activity in patients with thrombotic microangiopathies.Blood, 2012, vol. 120, 1157-1164 [0195]
- **M. JIMENEZ-ALCAZARN. KIMT. A. FUCHS**Circulating Extracellular DNA: Cause or Consequence of Thrombosis?Semin Thromb Hemost, 2017, vol. 43, 553-561 [0195]
- **M. BOETTCHER et al.**Degradation of Extracellular DNA by DNaseI Significantly Reduces Testicular Damage After Testicular Torsion in Rats.Urology, 2017, vol. 109, 223-e221-223e227- [0195]
- **A. D. BANDARANAYAKES. C. ALMO**Recent advances in mammalian protein production.FEBS letters, 2014, vol. 588, 253-260 [0195]
- **Y. K. HANY. G. KIMJ. Y. KIMG. M. LEE**Hyperosmotic stress induces autophagy and apoptosis in recombinant Chinese hamster ovary cell culture.Biotechnology and bioengineering, 2010, vol. 105, 1187-1192 [0195]
- **J. L. CEREGHINOJ. M. CREGG**Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*.FEMS microbiology reviews, 2000, vol. 24, 45-66 [0195]
- **M. AHMADM. HIRZH. PICHLERH. SCHWAB**Protein expression in *Pichia pastoris*: recent achievements and perspectives for heterologous protein production.Applied microbiology and biotechnology, 2014, vol. 98, 5301-5317 [0195]
- **R. DALYM. T. HEARN**Expression of heterologous proteins in *Pichia pastoris*: a useful experimental tool in protein engineering and production.Journal of molecular recognition : JMR, 2005, vol. 18, 119-138 [0195]

PATENTKRAV

1. DNase-1-lignende 3 (D1L3) variant til anvendelse i behandling af sygdomme eller tilstande
 5 **kendetegnet ved** tilstedeværelse eller akkumulation af neutrofile, ekstracellulære fælder (NET), hvor D1L3-varianten omfatter en aminosyresekvens, der har mindst 90 % sekvensidentitet med det modne enzym defineret af SEQ ID NO: 2 uden signalpeptidet, forudsat D1L3-varianten har en deletion af hele eller en del af den C-terminale hale defineret af SEQ ID NO: 33.
- 10 2. D1L3-variant til anvendelse ifølge krav 1, hvor D1L3-varianten kondenseres til en halveringstidsforlængende del, hvor den halveringstidsforlængende del vælges fra gruppen bestående af albumin, transferrin, et Fc eller et elastinlignende protein.
- 15 3. D1L3-variant til anvendelse ifølge krav 1, hvor D1L3-varianten kondenseres til albumin.
4. D1L3-variant til anvendelse ifølge krav 1, hvor varianten omfatter et inaktiveret nukleart lokaliseringssignal (NLS).
- 20 5. D1L3-variant til anvendelse ifølge krav 4, hvor et NLS inaktiveres ved deletion af hele eller en del af NLS1 og/eller NLS2 fra SEQ ID NO: 2, hvor NLS1 omfatter aminosyrerne 35-51 fra SEQ ID NO: 2 og NLS2 omfatter aminosyrerne 296-305 fra SEQ ID NO: 2.
- 25 6. D1L3-variant til anvendelse ifølge krav 5, hvor NLS2 af SEQ ID NO: 2 er deleteret.
7. D1L3-variant til anvendelse ifølge krav 1, og som omfatter deletion af hele den C-terminale hale defineret af SEQ ID NO: 33.
8. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor sygdommen eller tilstanden, der skal behandles, er en kronisk eller akut, inflammatorisk lidelse.
- 30 9. D1L3-variant til anvendelse ifølge krav 8, hvor sygdommen eller tilstanden, der skal behandles, er en akut eller kronisk infektion.
- 35 10. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 9, hvor sygdommen eller tilstanden, der skal behandles, er en vaskulær okklusion omfattende NET.
11. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 9, hvor sygdommen

eller tilstanden, der skal behandles, er kronisk neutrofil, neutrofil aggregering og leukostase, thrombose og vaskulær okklusion, iskæmireperfusionslæsion, kirurgisk og traumatisk vævslæsion, en akut eller kronisk inflammatorisk reaktion eller sygdom, en autoimmun sygdom, kardiovaskulær sygdom, metabolisk sygdom, systemisk inflammation, inflammatorisk sygdom i luftvejene, inflammatoriske nyresygdomme, inflammatoriske sygdomme relateret til transplanteret væv og cancer.

12. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 11, hvor D1L3-varianten administreres intravenøst, intraarterielt, intraartikulært, intramuskulært, subkutant, topisk eller ved pulmonal administration.

13. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor individet har eller har en risiko for at få en gangokklusion i et gangsystem.

14. D1L3-variant til anvendelse ifølge krav 13, hvor gangsystemet er cerebral akvædukt, galdegang, tårekanal, mælkegang, cystisk kanal, leverkanal, ejakulationskanal, parotis kanal, submandikulær kanal, større sublingual kanal, Bartholins kanal, cerebral akvædukt, bugspytkirtel, brystkirtel, sædleder, urinleder, urinblære, galdeblære og lever.

15. D1L3-variant til anvendelse ifølge krav 14, hvor sygdommen eller tilstanden, der skal behandles, er pancreatitis, cholangitis, conjunctivitis, mastitis, tør øjensygdom, obstruktion af sædleder eller nyresygdom.

16. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor sygdommen eller tilstanden, der skal behandles, er systemisk lupus erythematosus (SLE).

17. D1L3-variant til anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor behandlingen omfatter den supplerende administration af DNase1 eller en variant deraf.

DRAWINGS

Drawing

Fig. 1A




Fig. 1B




Fig. 1C




Fig. 1D




Fig. 1E




Fig. 1F




Fig. 2A




Fig. 2B



Fig. 2C




Fig. 2D




Fig. 2E




Fig. 2F




Fig. 2G




Fig. 2H




Fig. 2I




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 3D





Fig. 3E

Fig. 3F

Fig. 3G

Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D




Fig. 4E




Fig. 4F




Fig. 4G




Fig. 4H



Fig. 4I




Fig. 5A







Fig. 5B



Parameter (Units)	Ctl	CSF3	P-Value
N	3	4	-
Neutrophils (K/ μ l)	0.5 $\pm$ 0.1	19.2 $\pm$ 5.2	< 0.001
ALT (U/l)	50 $\pm$ 22	37 $\pm$ 13	> 0.05
AST (U/l)	80 $\pm$ 25	58 $\pm$ 8	> 0.05
BUN (mg/ml)	21 $\pm$ 2	21 $\pm$ 2	> 0.05
Creatinine (mg/ml)	0.4 $\pm$ 0.1	0.3 $\pm$ 0.1	> 0.05

Fig. 5C

Fig. 5D

Fig. 5E

Fig. 6A




Fig. 6B




Fig. 6C





Fig. 6D




Fig. 7A

D1/D13^{+/+} + CSF3/Ctrl
Lung, Overview




Fig. 7B

D1/D13^{+/+} + CSF3/Ctrl
Occluded blood vessels, Detail




Fig. 7C




Fig. 8A




Fig. 8B

Fig. 9A




Fig. 9B




FIG. 10

#	Cause of death	Sex	Age	Tissue	Hematoxylin-rich clots	Chromatin & MPO-positive	Localization
1	ARDS	M	84	Lung	Yes	Yes	Bronchus
2	Septic cardiovascular failure + ARDS	F	63	Lung	No	Not applicable	Not applicable
3	ARDS	F	67	Lung	No	Not applicable	Not applicable
4	Septic multi-organ failure + ARDS	M	72	Lung	Yes	Yes	Bronchus
5	Septic cardiovascular failure + ARDS	F	75	Lung	No	Not applicable	Not applicable
6	Septic multi-organ failure	M	60	Lung	Yes	Yes	Blood vessel
7	Septic cardiovascular failure	F	76	Lung	Yes	Yes	Blood vessel
8	ARDS	F	68	Lung	No	Not applicable	Not applicable

Fig. 11A




Fig. 11B




Fig. 11C




Fig. 11D




FIG. 12




Fig. 13A




Fig. 13B




Fig. 13C




Fig. 13D




Fig. 13E




Fig. 13F




Fig. 14A




Fig. 14B




Fig. 14C




Fig. 14D




Fig. 14E





Fig. 14F




Fig. 14G

Fig. 15A




Fig. 15B




Fig. 15C




Fig. 16A

In vitro screening approach for hyperactive D1 and D1L3 variants




Fig. 16B Readout system: dsDNA & chromatin degradation




FIG. 17

Sequence Ruler	MRGKLLGALLAALLOGAVSLKIAAFNIOTFGETKMSNATLVSYIQIILSRVDIALVQEVDRSHLTAVGKHLDNLNQD	10 20 30 40 50 60 70
Sequence Ruler	MSRELAPLLILLLSIHSAAMRIGCSFNVSEGESKQEDNANDVIYVIRCDYILVMEIKDSNNICPILMEILHN	1 10 20 30 40 50 60 70
Sequence Ruler	APDTYHYVVEPIGRNSYKERYLFVYRPDOVSADVSTYYDGGCEPCGNDTFNHEPAIVRFFSRFTVEVREAIIVPLHAA	80 90 100 110 120 130 140 150
Sequence Ruler	SIGITYNYVVISLGRNTYKEQVAFLYKEILVSVSYHYHDYQDQDADVFSRPFVWVQSPHTAVKDEVIPLMTT	80 90 100 110 120 130 140 150
Sequence Ruler	PQDAVAEIDALYDYVLDVQEKWGLEDVMLMGDFNAGCSYVRPSONSSIRLWTSFTFOWLIPDSADTTA-TPTHCAVDRIY	160 170 180 190 200 210 220 230
Sequence Ruler	PETSVIEIDELVEVYTOVHRVDAENFIHQDFNAGCSYVPAMNIRLTDPFVWIGDOEDTIVSTNCAYDRIY	160 170 180 190 200 210 220 230
Sequence Ruler	VAGMLLRGAVVPSALPENEQAYGLSDQLAQASDNYFVEVMVK	240 250 260 270 280
Sequence Ruler	LGQSEIVSSVVPNSVFDQAYLITEEALDVSDHCFVEFLQSSAFPTNSGVTLTSSSS	240 250 260 270 280 290 300
Signal peptide		Conserved amino acids
		C-terminal tail in D1L3

FIG. 18

#	DNase1	DNase1	Mutation in DNase1	Analyzed in this application
1	R29 (R9)	Q31 (Q9)	Q31R (Q9R)	+
2	K39 (R19)	A41 (A19)	A41K (A19K)	-
3	K47 (K27)	Q49 (Q27)	Q49K (Q27K)	+
4	K50 (K30)	S52 (S30)	S52K (S30K)	-
5	R66 (R46)	T68 (T46)	T68R (T46R)	-
6	K74 (K54)	N76 (N54)	N76K (N54K)	+
7	R77 (R57)	Q79 (Q57)	Q79R (Q57R)	+
8	R80 (R60)	-	D80_A81insR (D60_A61insR)	+
9	R81 (R61)	A81 (A59)	A81R (A61R)	+
10	R92 (R72)	P92 (P70)	P92R (P70R)	+
11	K109 (K89)	D109 (D87)	D109K (D87K)	+
12	K114 (K94)	V114 (V92)	V114K (V92)	+
13	R115 (K95)	D115 (D93)	D115R (D92R)	+
14	K163 (K143)	A164 (A142)	A164K (A142K)	+
15	K176 (K156)	Q177 (Q155)	Q177K (Q115K)	+
16	K180 (K160)	G181 (G159)	G181K (G159K)	+
17	K199 (K179)	P200 (P178)	P200K (P178K)	-
18	K200 (K180)	S201 (S179)	S201K (S179K)	+
19	K203 (K183)	S204 (S182)	S204K (S182K)	+
20	R208 (R188)	W209 (W187)	W209R (W187R)	+
21	R212 (R192)	T213 (T191)	T213R (T191R)	+
22	K226 (K206)	-	A226_T227insK (A204_T205insK)	+
23	K227 (K207)	T227 (T205)	T227K (T205K)	+
24	R239 (R219)	A239 (A217)	A239R (A217R)	+
25	K250 (K230)	D250 (D228)	D250K (D228K)	+
26	K259 (K239)	A259 (A237)	A259K (A237K)	-
27	K262 (K262)	G262 (G240)	G262K (G240K)	-
28	K280 (K260)	M280 (M258)	M280K (M258K)	+
29	R285 (K265)	-	-	-
30	K291 (K271)	-	-	-
31	K292 (K272)	-	-	-
32	R297 (K277)	-	-	-
33	K298 (K278)	-	-	-
34	K299 (K279)	-	-	-
35	K301 (K281)	-	-	-
36	K303 (K283)	-	-	-
37	R304 (K284)	-	-	-

FIG. 20

Human DNase1L3 Sequence Ruler	10	20	30	40	50	60	70
Rat DNase1	MSRELAFLLLLSHSAAMRICSNVAFSGESKOEDKNAMDVIVIKRCDILVMEIKDSNNRIPCILMEKLNEN						
Mouse DNase1	MRYTGLNGILLTLVNLGLQAAALIRIAAFNITFGDTKMSNATLSYIVILSRDYIAVVOEVRDTHLVAVGKLLDELND						
Human DNase1	MRYTGLNGILLTLVNLGLQAAALIRIAAFNITFGDTKMSNATLSYIVILSRDYIAVVOEVRDTHLVAVGKLLDELND						
Sequence Ruler	10	20	30	40	50	60	70
Human DNase1L3 Sequence Ruler	80	90	100	110	120	130	140
Rat DNase1	I--PDNYRYIISEPLGRKSYKEQYLFVYRPSQVSVLDQSYHYDDGCEPCGNDTSREPAIVKFFSPYTEVREFAIVPLHSA						
Mouse DNase1	K--PDYRYVVSEPLGRKSYKEQYLFVYRPSQVSVLDQSYHYDDGCEPCGNDTSREPAIVKFFSPYTEVREFAIVPLHSA						
Human DNase1	A--PDYHYVVSEPLGRNSYKERYLFVYRPSQVSAVSYHYDDGCEPCGNDTSREPAIVKFFSRFTTEVREFAIVPLHAA						
Sequence Ruler	80	90	100	110	120	130	140
Human DNase1L3 Sequence Ruler	160	170	180	190	200	210	220
Rat DNase1	PETSVKEIDELVEVYTDVHRWKAENFIMGDFNAGCSYVPPKKAWNIRLTDPRFVWLDGQEDITVKKSTNCAYDRIV						
Mouse DNase1	PTEAVSEIDALYDVLVDVOKWGLEDMFGDFNAGCSYVTSOWSSIRLTSPIFQWLIPDSADTTA-TSTHCAVDRIY						
Human DNase1	PTEAVSEIDALYDVLVDVOKWGLEDMFGDFNAGCSYVTSOWSSIRLTSPIFQWLIPDSADTTA-TSTHCAVDRIY						
Sequence Ruler	160	170	180	190	200	210	220
Human DNase1L3 Sequence Ruler	240	250	260	270	280	290	300
Rat DNase1	LRGQELVSSVVPKSNVDFDKAYLTTEEALDVSDHFPVEFKLOSSRAFTNSKKSVTLRKKTKSKRS						
Mouse DNase1	VAGALLQAAVVPSSAVPFPDFAEYLTNOMAEAI SDHYPVEVTLRKT--						
Human DNase1	VAGALLQAAVVPNSAVPFPDFAEYGLSNOLAEAI SDHYPVEVTLRKT--						
Sequence Ruler	240	250	260	270	280	290	300

Signal peptide

K/R residues in D1L3, present in murine or rat D1, absent in human D1

Fig. 21A




Fig. 21B




FIG. 22




Fig. 22A




FIG. 23




Fig. 24A




Fig. 24B




FIG. 25




FIG. 26




Fig. 27A




Fig. 27B




FIG. 28




FIG. 29


BUILDING BLOCK-ENGINEERING OF HOMOLOGOUS PROTEINS**I. AMINO ACID SEQUENCE ALIGNMENT OF HOMOLOGOUS ENZYMES****II. EXCISION OF CDNA CODING FOR VARIABLE AMINO ACIDS IN DONOR ENZYME****III. DELETION OF CDNA CODING FOR VARIABLE AMINO ACIDS IN RECIPIENT ENZYME****IV. TRANSFER OF CDNA FROM DONOR TO RECIPIENT ENZYME**

FIG. 30

MOUSE (P49183) RAT (P21704) CHIMPANZEE (H2QAH1) HUMAN (P24855) Sequence Ruler	1 10 20 30 40 50 60 70 MRYS-GLMGILLTL--VNLQGLAQLRIAARNIRTFGETKMSNATLSVYFKILSRDYIAVQEVRSLSHLVAVGKLLDEL MRYS-GLMGILLTL--VNLQGLAQLRIAARNIRTFGETKMSNATLSVYFKILSRDYIAVQEVRSLSHLVAVGKLLDEL MRSM-KLLGALLAL--AALQGAVALKIAARNIQTFGETKMSNATLSVYFKILSRDYIALVQEVRSLSHLTAVGKLLDNL MRSM-KLLGALLAL--AALQGAVALKIAARNIQTFGETKMSNATLSVYFKILSRDYIALVQEVRSLSHLTAVGKLLDNL
MOUSE (O65070) RAT (O68107) CHIMPANZEE (H2QAH1) HUMAN (Q13609) Sequence Ruler	1 10 20 30 40 50 60 70 MSLHPASPRLASLLFICALHDTLALRLCSFNVRSPGASKKENHEAMDIIKIIKRCDIILLMEIKDSNNNICPMLMEKL MSLHPASPRLASLLFICALHDTLALRLCSFNVRSPGASKKENHEAMDIIKIIKRCDIILLMEIKDSNNNICPMLMEKL MSLHPASPRLASLLFICALHDTLALRLCSFNVRSPGASKKENHEAMDIIKIIKRCDIILLMEIKDSNNNICPMLMEKL MSLHPASPRLASLLFICALHDTLALRLCSFNVRSPGASKKENHEAMDIIKIIKRCDIILLMEIKDSNNNICPMLMEKL
MOUSE (P49183) RAT (P21704) CHIMPANZEE (H2QAH1) HUMAN (P24855) Sequence Ruler	80 90 100 110 120 130 140 150 NRDK--PDTVRVVSSEPLGRKSYKEQYLFVVRPDQVSLDSYQXDGCEPCGNDTSREPAIVKFEFSPYTEVQEFATVPL NRDK--PDTVRVVSSEPLGRKSYKEQYLFVVRPDQVSLDSYQXDGCEPCGNDTSREPAIVKFEFSPYTEVQEFATVPL NRDI--PDTVRVVSSEPLGRKSYKEQYLFVVRPDQVSLDSYQXDGCEPCGNDTSREPAIVKFEFSPYTEVQEFATVPL NRDI--PDTVRVVSSEPLGRKSYKEQYLFVVRPDQVSLDSYQXDGCEPCGNDTSREPAIVKFEFSPYTEVQEFATVPL
MOUSE (O65070) RAT (O68107) CHIMPANZEE (H2QAH1) HUMAN (Q13609) Sequence Ruler	80 90 100 110 120 130 140 150 NGNSRRSTTNNYVVSRLGRNTYKQYAFVKEKLVSVKTKYHXYQYQ-DGDTDFVSRPFVWVWFSPFTAVKDFVIVPL NGNSRRSTTNNYVVSRLGRNTYKQYAFVKEKLVSVKTKYHXYQYQ-DGDTDFVSRPFVWVWFSPFTAVKDFVIVPL NGNSRRSTTNNYVVSRLGRNTYKQYAFVKEKLVSVKTKYHXYQYQ-DGDTDFVSRPFVWVWFSPFTAVKDFVIVPL NGNSRRSTTNNYVVSRLGRNTYKQYAFVKEKLVSVKTKYHXYQYQ-DGDTDFVSRPFVWVWFSPFTAVKDFVIVPL

Signal Peptide

Conserved AA = Anchors

Variable AA = Building Blocks

FIG. 30 (continued)

Mouse (P49183) Rat (P21704) Chimpanzee (H20AH1) Human (P24855) Sequence Ruler	HAARTEA VSEIDAL YDYLDVWQKWGLEIDIMFGDFNAGCSVTSQWSSIRLRISPIFQWLPDSADTTVT-STHCAYD HSA RTEA VSEIDAL YDYLDVWQKWGLEIDIMFGDFNAGCSVTSQWSSIRLRISPIFQWLPDSADTTAT-STHCAYD HAARGDAVAEIDAL YDYLDVQEKWGLEIDVLMGDFNAGCSVTRPSQWSSIRLRISPAEQWLPDSADTTAT-PTHCAVD HAARGDAVAEIDAL YDYLDVQEKWGLEIDVLMGDFNAGCSVTRPSQWSSIRLRISPTFQWLPDSADTTAT-PTHCAVD	160 170 180 190 200 210 220 230
	HTTRET SVKEIDELVDVYTDVRRQWKTENFIFMGDFNAGCSVTPKKAWONIRLRIDPKFVMLGGQEDTTVKKSTSCAYD HTTRET SVKEIDELADVYTDVRRQWKAENFIFMGDFNAGCSVTPKKAWKNIRLRIDPNFVMLGGQEDTTVKKSTSCAYD HTTRET SVKEIDELVEYTDVRRQWKAENFIFMGDFNAGCSVTPKKAWKNIRLRIDPNFVMLGGQEDTTVKKSTSCAYD HTTRET SVKEIDELVEYTDVRRQWKAENFIFMGDFNAGCSVTPKKAWKNIRLRIDPREVVMGGQEDTTVKKSTSCAYD	160 170 180 190 200 210 220 230
	RI VVAGALLQAAVVPNSAVPFDQAEYGSNQLAEAI SDHYRVEVTLRKI RI VVAGALLQAAVVPNSAVPFDQAEYRINQMAEAI SDHYRVEVTLRKI RI VVAGMLLQCAVVPDIALPFDQAAVGSQDLAQAI SDHYRVEVMLK RI VVAGMLLQCAVVPDIALPFDQAAVGSQDLAQAI SDHYRVEVMLK	160 170 180 190 200 210 220 230
	RI VLRGQEI VNSVVPKSGVFDQKAYDSEEEALDVSDHFRVEFKQSSRAFTNNSKSVLKKRKGNSRS RI VLRGQEI VNSVVPKSGVFDQKAYDSEEEALDVSDHFRVEFKQSSRAFTNNSKSVLKKRKGNSRS RI VLRGQEI VNSVVPKSNVFDQKAYKTEEEALDVSDHFRVEFKQSSRAFTNNSKSVTLRKKTKSKRS RI VLRGQEI VNSVVPKSNVFDQKAYKTEEEALDVSDHFRVEFKQSSRAFTNNSKSVTLRKKTKSKRS	240 250 260 270 280 290 300
Signal Peptide	Conserved AA = Anchors	Variable AA = Building Blocks

Fig. 31A

Example of Building Block Engineering: human DNase1 (D1) variants with Building Blocks from human DNase1L3 (D1L3)

BB #	D1 N-Anchors: Conserved AA in D1	D1 C-Anchors: Conserved AA in D1	D1 Building Blocks: Variable AA in D1	D1L3 Building Blocks: Variable AA in D1L3	Building Block Mutations of D1 variants	BB Cluster
0	N/A	N/A	1-22 (MRGMKLL GALLAALQGAVS)	1-20 (NSRELAPLLLLLSHSA)	M1_S22delinsRELAPLLL LLLSHSA	BB ₀
1	Signal peptide	F28/N29	23-27 (LKIAA)	21-25 (MRICS)	L23_A27delinsMRICS	BB ₁₋₃
2	F28/N29	F33/G34	30-32 (HOT)	28-30 (VRS)	I30_T32delinsVRS	
3	F33/G34	K37	35-38 (ET)	33-34 (ES)	E35_T36delinsES	
4	K37	V48	38-47 (MSNATLVSYI)	36-45 (QEDKNAMDIVI)	M38_I47delinsQEDKNAMDIVI	BB ₄
5	V48	R53	49-52 (QLS)	47-50 (KVIK)	Q49_S52delinsKVIK	BB ₅₋₇
6	R53	D55	54 (Y)	52 (C)	Y54C	
7	D55	E61	56-60 (IALVQ)	54-58 (ILVM)	I56_Q60delinsILVM	
8	E61	D64	62-63 (VR)	60-61 (IK)	V62_R63delinsIK	BB ₈₋₁₀
9	D64	L73	65-72 (SHLTAVGK)	63-70 (SNNRICPI)	S65_K72delinsSNNRICPI	
10	L73	L77/N78	74-78 (LDN)	72-74 (MEK)	L74_N76delinsMEK	
11	L77/N78	Y65	79-84 (QDAPOI)	77-84 (RNSRRGII)	Q79_I84delinsRNSRRGII	BB ₁₁
12	Y65	Y87	86-91	86 (N)	HPEN	BB ₁₂₋₁₄
13	Y87	S90	92-99 (LV)	88-90 (VLI)	V92_V99delinsVLI	
14	S90	L93/G94/R95	100-103 (RISG)	91-92 (EP)	E91_F92delinsISR	
15	L93/G94/R95	Y98/K99/E100	96-97 (NS)	95-97 (NT)	N96_S97delinsNT	BB ₁₅₋₁₈
16	Y98/K99/E100	Y102	101 (R)	101 (Q)	R101Q	
17	Y102	F104	103 (L)	103 (A)	L103A	
18	F104	Y106	105 (V)	105 (L)	Y105L	BB ₁₉₋₂₂
19	Y106	V111/S112	107-110 (RPDO)	107-110 (KEKL)	R107_Q110delinsKEKL	
20	Y111/S112	Y117	113-116 (AVDS)	113-116 (VKRS)	A113_S116delinsVKRS	
21	Y117	Y119	118 (Y)	118 (H)	Y118H	BB ₂₃₋₂₅
22	Y119	D121	120 (D)	120 (H)	D120H	
23	D121	D128	122-128 (GCEPCGN)	122-127 (YQDGD)	G122_N128delinsYQDGD	
24	D128	F131	130 (T)	129 (V)	T130S	BB ₂₆₋₂₈
25	F131	R133/E134/P135	132 (N)	131 (S)	N132S	
26	R133/E134/P135	V138	136-137 (AI)	135 (E) (V)	A136_I137delinsV	
27	V138	F140	139 (R)	138 (W)	R139W	BB ₂₉₋₃₄
28	F140	T145	141-144 (FSRF)	140-143 (QSPH)	F141_F144delinsQSPH	
29	T145	F150	148-149 (EVRE)	145-148 (AVKD)	E146_E149delinsAVKD	
30	F150	I152	151 (A)	150 (V)	A151V	BB ₃₁₋₃₄
31	I152	P154/L155/H156	153 (V)	152 (I)	V153I	
32	P154/L155/H156	P159	157-158 (AA)	156-157 (TT)	A157_A158delinsTT	
33	P159	V183	160-162 (GDA)	159-161 (ETS)	G160_A162delinsETS	BB ₃₅₋₃₈
34	V183	E165/I166/D167	164 (A)	163 (K)	A164K	

Fig. 31A (continued)

BB #	D1 N-Ancor: Conserved AA in D1	D1 C-Ancor: Conserved AA in D1	D1 Building Block: Variable AA in D1	D1L3 Building Block: Variable AA in D1L3	Building Block Mutations of D1 variants	BB
35	E165/I166/D167	L168	168 (A)	167 (E)	A168E	BB ₃₅₋₃₇
36	L169	V172/V173	170-171 (YD)	169-171 (VE)	Y170 D171delinsVE	
37	V172/V173	D175/V176	174 (L)	173 (I)	L174I	
38	D175/V176	W180	177-179 (QEK)	176-178 (KHR)	Q177 K178delinsKHR	
39	W180	E183	181-182 (GL)	180-181 (KA)	G181 L182delinsKA	BB ₃₉₋₄₀
40	E183	M187-GDFNAGCSY-V198	184-187 (DVML)	183-187 (NFI)	D184_L187delinsNFI	
41	M187-GDFNAGCSY-V198	V203	199-202 (RPSQ)	198-201 (PKA)	R199 D202delinsPKA	
42	V203	D206/R207/D208	204-205 (SS)	203-204 (K)	S204 S205delinsKA	
43	S206/H207/L208	T210	209 (R)	208 (R)	R209	
44	T210	P212	211 (S)	210 (D)	S211D	BB ₄₄₋₄₇
45	E242	E244	243 (L)	242 (K)	L243K	
46	F214	R216/D217/S218	245 (Q)	244 (V)	Q245V	
47	R216/D217/S218	R229	249 (P)	248 (G)	P249G	
48	D220	D227/D228/D229	251-252 (SA)	250-251 (Q)	S251 A252delinsQ	
49	D227/D228/D229	L229	253-254 (AP)	252-253 (KAS)	A253 R254delinsKAS	BB ₄₉
50	T229	C231-AYDRI-V237	230 (H)	230 (N)	H230N	
51	C231-AYDRI-V237	G240	238-239 (VA)	238-239 (LR)	V238 A239delinsLR	BB ₅₁₋₅₄
52	G240	V247N/248/P248	241-246 (MLRGA)	241-248 (QEVSS)	M241 A248delinsQEVSS	
53	V247N/248/P248	S251	250 (D)	250 (K)	D250K	
54	S251	F255	252-254 (ALP)	252-254 (NSV)	A252 P254delinsNSV	
55	F255	F257/Q258	256 (N)	256 (D)	N256D	
56	F257/Q258	Y261	259-260 (AA)	259-260 (KA)	A259 A260delinsKA	
57	Y261	L263	262 (G)	262 (K)	G262K	BB ₅₇₋₅₉
58	L263	A268	264-267 (SDQL)	264-267 (TEEE)	S264 L267delinsTEEE	
59	A268	S272/D273/H274	268-271 (QAI)	269-271 (LDV)	Q269 I271delinsLDV	
60	S272/D273/H274	P276/V277/E278	275 (Y)	275 (F)	Y275F	
61	P276/V277/E278	L281	279-280 (VM)	279-280 (FK)	V279 M280delinsFK	
62	L281	-	282 (K)	282-305 (QSSRAFTNSKK SVTLRKTKSKRS)	K282delinsQSSRAFTNSKKSVTLRKTKSKRS	BB ₆₂

Fig. 31B

Example of Building Block Engineering: human DNase1L3 (D1L3) variants with Building Blocks from human DNase1 (D1)

BB #	D1L3 N-Anter: Conserved AA in D1	D1L3 C-Anter: Conserved AA in D1	D1L3 Building Blocks: Variable AA in D1L3	D1 Building Blocks: Variable AA in D1	Building Block Mutations of D1L3 variants	BB Cluster
0	N/A	N/A	1-20 (MSRELAPILLILLSIHSA)	1-22 (NRGAKLL GALLAALLOGAVS)	M1_A20delinsMRGKLLGALLALA	BB ₀
1	Signal peptide	F26N27	21-28 (MRICS)	23-27 (LKAA)	ALLOGAVS	
2	F26N27	F31G32	28-30 (VRS)	30-32 (IOT)	M21_S25delinsLKAA V28_S30delinsIOT	BB _{1,3}
3	F33G34	K35	33-34 (ES)	35-36 (ET)	E33_S34delinsET	
4	F38	Y48	37-38 (GTPSAKQV)	39-47 (GTPSAKQV)	Y38_A39delinsGTPSAKQV	BB ₄
5	V46	R51	47-50 (KVIK)	49-52 (QILS)	K47_K50delinsQILS	
6	R51	D53	52 (C)	54 (Y)	C52Y	BB _{5,7}
7	D53	E59	54-58 (ILVM)	59-60 (ALVQ)	I54_M58delinsALVQ	
8	E59	D62	60-61 (IK)	62-63 (VRI)	I60_K61delinsVRI	
9	D62	L71	63-70 (SNRICPI)	65-72 (SHLTAYGK)	S63_I70delinsSHLTAYGK	BB _{8,10}
10	L71	L75N76	72-74 (MEK)	74-76 (LON)	M72_K74delinsLON	
11	L75N76	Y85	77-84 (RNSRRGIT)	79-84 (QDAPDT)	R77_T84delinsQDAPDT	BB ₁₁
12	Y85	Y87	86 (N)	88 (H)	N86H	
13	Y87	S90	88-89 (VI)	88-89 (VV)	V88_I89delinsVV	
14	S90	L93/G94/R95	91-92 (SR)	91-92 (EP)	S91_R92delinsEP	BB _{12,14}
15	L93/G94/R95	Y98/K99/E100	96-97 (NT)	96-97 (NS)	N96_I97delinsNS	
16	Y98/K99/E100	Y102	101 (Q)	101 (R)	Q101R	
17	Y102	F104	103 (A)	103 (L)	A103L	BB _{15,18}
18	F104	Y106	105 (L)	105 (V)	L105V	
19	Y106	V111/S112	107-110 (KEKL)	107-110 (RPDQ)	K107_L110delinsRPDQ	
20	Y111/S112	Y117	113-116 (VKRS)	113-118 (AVDS)	V113_S116delinsAVDS	BB _{19,22}
21	Y117	Y119	118 (H)	118 (Y)	H118Y	
22	Y119	D121	120 (H)	120 (D)	H120D	
23	D121	G128	122-127 (TDDGDA)	123-128 (TDDGDA)	Y122_A123delinsTDDGDA	BB _{23,24}
24	D128	F130	129 (V)	130 (I)	V129I	
25	F130	R132/E133/P134	131 (S)	132 (H)	S131H	
26	R132/E133/P134	V137	135-138 (FV)	138-137 (AI)	I35F_I36VdelinsAI	BB ₂₅
27	V137	F139	138 (W)	139 (R)	W138R	
28	F139	T144	140-143 (QSPH)	141-144 (FSRF)	Q140_H143delinsFSRF	
29	T144	F149	145-148 (AVKD)	146-149 (EVRE)	A145_D148delinsAVKD	
30	F149	I151	150 (V)	151 (A)	V150A	
31	I151	P153/L154/M155	152 (I)	153 (V)	I152A	
32	P153/L154/M155	P157	156-157 (TT)	157-158 (AA)	T156_T157delinsAA	BB _{27,34}
33	P157	V162	159-161 (ETS)	160-162 (GDA)	E159_S161delinsGDA	
34	V162	E164/I165/D166	163 (K)	164 (A)	K163A	

Fig. 31B (continued)

BB #	D1L3 N-Anchor: Conserved AA in D1L3	D1L3 C-Anchors: Conserved AA in D1L3	D1L3 Building Blocks: Variable AA in D1L3	D1 Building Block: Variable AA in D1	Building Block Mutations of D1L3 variants	BB Cluster
35	E164/I165/D166	L168	167 (E)	168 (A)	E167A	BB ₃₅₋₃₇
36	L188	V171/Y172	169-170 (VE)	170-171 (VD)	V169 E170delinsYD	
37	V171/Y172	D174/V175	173 (T)	174 (L)	T173L	
38	D174/V175	W179	176-178 (KHR)	177-179 (QEK)	K176 R178delinsQEK	BB ₃₈₋₄₀
39	W179	E182	180-181 (KA)	181-182 (GL)	K180 A181delinsGL	
40	E182	M187-GDFNAGCSY- V197	183-186 (NFF)	184-187 (DVML)	N183 F186delinsDVML	
41	M187-GDFNAGCSY-V197	W202	198-201 (PKKA)	199-202 (RPSQ)	198P 201AdeinsRPSQ	
42	W202	I205/R206/L207	203-204 (KN)	204-205 (SS)	K203 N204delinsSS	
43	I205/R206/L207	T209	208 (R)	208 (W)	R208W	
44	T209	P211	210 (D)	211 (S)	D210S	BB ₄₄₋₄₆
45	P211	F213	212 (R)	213 (T)	R212T	
46	F213	W215/L216/I217	214 (V)	215 (Q)	V214Q	
47	W215/L216/I217	D219	218 (G)	219 (P)	G218P	
48	D219	D222/I223/I224	220-221 (QE)	221-222 (SA)	Q220 E221delinsSA	
49	D222/I223/I224	T229	225-228 (VRKS)	228-229 (ATP)	V225 S228delinsATP	BB ₄₉
50	T229	C231-AYDRI-V237	230 (N)	230 (H)	N230H	
51	C231-AYDRI-V237	G240	238-239 (LR)	238-239 (VA)	L238 R239delinsVA	BB ₅₁₋₅₄
52	G240	V247/V248/P249	241-246 (QEVSS)	241-246 (MLIRGA)	Q241 S246delinsMLIRGA	
53	V247/V248/P249	S251	250 (K)	250 (D)	K250D	
54	S251	F255	252-254 (NSV)	252-254 (ALP)	N252 V254delinsALP	
55	F255	F257/Q258	258 (D)	258 (N)	D258N	
56	F257/Q258	Y261	259-260 (KA)	259-260 (AA)	K258 A260delinsAA	BB ₅₅₋₅₉
57	Y261	L263	262 (K)	262 (G)	K262G	
58	L263	A268	264-267 (TEEE)	264-267 (SDQL)	T264 E267delinsSDQL	
59	A268	S272/D273/H274	269-271 (LDV)	269-271 (QAI)	L269 V271delinsQAI	
60	S272/D273/H274	P284/V285/Q286	276 (F)	276 (T)	P279 S285delinsY59	
61	P284/V285/Q286	L281	279-280 (FA)	279-280 (YH)	Q282 S285delinsK	
62	L281		282 (S)	282 (K)		BB ₆₂

FIG. 32


BUILDING BLOCK CLUSTER-ENGINEERING**I. AMINO ACID SEQUENCE ALIGNMENT OF HOMOLOGOUS ENZYMES****II. TRANSFER OF BUILDING BLOCK CLUSTERS****III. TRANSFER OF INDIVIDUAL BUILDING BLOCKS (OPTIONAL)****IV. TRANSFER OF INDIVIDUAL AMINO ACID RESIDUES (OPTIONAL)**

FIG. 33




Fig. 34A




Fig. 34B




Fig. 36A

NET-clot in cardiac chamber




Fig. 36B

NET-mediated death




Fig. 36C Experimental design: Acute anti-NET therapy




Fig. 36D

Fig. 36E

Fig. 36F

Fig. 37A Experimental design: Ischemia-reperfusion injury therapy




Fig. 37B




Fig. 37C




Fig. 38A

Chinese Hamster Ovary
(CHO) cells




Fig. 38B

Pichia pastoris




Fig. 38C




FIG. 39

SEQ ID NO	Description	Mutation (Mature protein without signal peptide)	Building Block Engineering	Parental SEQ ID NO
5	DNase1 variant	Q31R/Q48K/Q79K/W209K (Q36Q27K/Q57K/W187K)	-	1
6	DNase1 variant	Q31R/Q48K/Q79K/Q177K/W209K/G282K (Q36Q27K/Q57K/Q155K/W187K/G240K)	-	1
7	DNase1 variant	Q31R/A136F/T227K (Q36R/A114F/T205K)	-	1
9	DNase1 variant	A136F/A164K (A114F/A142K)	-	1
10	DNase1 variant	Q79_T64delinsRNSRRGIT (Q57_T62delinsRNSRRGIT)	+ (BB ₁₁)	1
11	DNase1 variant	H86N/V98_V89delinsVVE91_P92delinsSR (H84N/V86_V87delinsVVE91_P70delinsSR)	+ (BB ₁₂₋₁₄)	1
12	DNase1 variant	A136_I137delinsFV (A114_I115delinsFV)	+ (BB ₂₀)	1
13	DNase1 variant	R189_Q202delinsPKKA/S304_S203delinsKNW209R/S211D/TZ13R/Q215V/ P219Q/S221_A222delinsQE (R177_Q180delinsPKKA/S182_S183delinsKNW187R/S189D/T191R/Q193V/ P197G/S189_A200delinsQE)	+ (BB ₁₋₄₀)	1
14	DNase1 variant	A228_P228delinsVKKS (A204_P206delinsVKKS)	+ (BB ₄₀)	1
15	DNase1 variant	Q79_T64delinsRNSRRGIT/H86N/V98_V89delinsVVE91_P92delinsSR/ A136_I137delinsFV/ R189_Q202delinsPKKA/S304_S203delinsKNW209R/S211D/ TZ13R/Q215V/P219Q/S221_A222delinsQE/A226_P228delinsVKKS (Q57_T62delinsRNSRRGIT/H84N/V86_V87delinsVVE91_P70delinsSR/ A114_I115delinsFV/R177_Q180delinsPKKA/S182_S183delinsKNW187R/S189D/ T191R/Q193V/P197G/S189_A200delinsQE/ A204_P206delinsVKKS)	+ (BB ₁₁ , BB ₁₂₋₁₄ , BB ₂₀ , BB ₄₁₋₄₅ , BB ₄₆)	1
17	DNase1L3 variant	F275Y/F279_K280delinsVM/Q282_S305delinsK (F255Y/F259_K280delinsVM/Q282_S285delinsK)	+ (BB ₄₀₋₄₂)	2
19	DNase1L3 variant	Q36_I45delinsMSHATLV8YI (Q16_I25delinsMSHATLV8YI)	+ (BB ₄)	2
20	DNase1L3 variant	Y122_A127delinsGCEPCGNV120T/S131N (Y102_A107delinsGCEPCGNV108T/S111N)	+ (BB ₂₋₂₃)	2