



(12)

Rettet oversættelse af
europæisk patentskriftPatent- og
Varemærkestyrelsen

(51) Int.Cl.: **C 07 K 16/28 (2006.01)** **A 61 K 39/00 (2006.01)** **A 61 K 47/68 (2017.01)**

(45) Oversættelsen bekendtgjort den: **2024-09-23**

(80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2023-09-06**

(86) Europæisk ansøgning nr.: **18163942.8**

(86) Europæisk indleveringsdag: **2012-05-24**

(87) Den europæiske ansøgnings publiceringsdag: **2018-12-19**

(30) Prioritet: **2011-05-27 US 201161490732 P** **2012-05-15 US 201261647196 P**

(62) Stamansøgningsnr: **12723206.4**

(84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**

(73) Patenthaver: **Glaxo Group Limited, GSK Medicines Research Centre , Gunnels Wood Road, Stevenage SG1 2NY, Storbritannien**

(72) Opfinder: **ALGATE, Paul, 580 Kalmia Place NW, Issaquah, WA, USA**
Clegg, Stephanie Jane, GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien
CRAIGEN, Jennifer L., GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien
Hamblin, Paul Andrew, GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien
LEWIS, Alan Peter, GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien
Parmar, Radha Shah, GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien
MAYES, Patrick, GlaxoSmithKline, 1250 S Collegeville Road, Collegeville, PA 19426-0989, USA
Wattam, Trevor Anthony Kenneth, GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY, Storbritannien

(74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**

(54) Benævnelse: **BCMA (CD269/TNFRSF17)-BINDENDE PROTEINER**

(56) Fremdragne publikationer:
WO-A1-2012/066058
WO-A2-2010/104949
WO-A2-2011/108008
RYAN MAUREEN C ET AL: "Antibody targeting of B-cell maturation antigen on malignant plasma cells", MOLECULAR CANCER THERAPEUTICS, AMERICAN ASSOCIATION OF CANCER RESEARCH, US, vol. 6, no. 11, 1 November 2007 (2007-11-01), pages 3009-3018, XP002581270, ISSN: 1535-7163
YAMANE-OHNUKI N ET AL: "Establishment of FUT8 knockout chinese hanster ovary cells: an ideal host cell line for producing completely defucosylated antibodies with enhanced antibody-dependent cellular

cytotoxicity", BIOTECHNOLOGY AND BIOENGINEERING, WILEY & SONS, HOBOKEN, NJ, US, vol. 87, no. 5, 6 August 2004 (2004-08-06) , pages 614-622, XP002983758, ISSN: 0006-3592, DOI: 10.1002/BIT.20151
C. PETERS ET AL: "Antibody-drug conjugates as novel anti-cancer therapeutics", BIOSCIENCE REPORTS, vol. 35, no. 4, 12 June 2015 (2015-06-12), pages e00225-e00225, XP055301629, US ISSN: 0144-8463, DOI: 10.1042/BSR20150089
RYAN MAUREEN C ET AL: "Antibody targeting of B-cell maturation antigen on malignant plasma cells", MOLECULAR CANCER THERAPEUTICS, AMERICAN ASSOCIATION OF CANCER RESEARCH, US, vol. 6, no. 11, 1 November 2007 (2007-11-01), pages 3009 - 3018, XP002581270, ISSN: 1535-7163
YAMANE-OHNUKI N ET AL: "Establishment of FUT8 knockout Chinese hamster ovary cells: an ideal host cell line for producing completely defucosylated antibodies with enhanced antibody-dependent cellular cytotoxicity", BIOTECHNOLOGY AND BIOENGINEERING, WILEY & SONS, HOBOKEN, NJ, US, vol. 87, no. 5, 6 August 2004 (2004-08-06), pages 614 - 622, XP002983758, ISSN: 0006-3592, DOI: 10.1002/BIT.20151
C. PETERS ET AL: "Antibody-drug conjugates as novel anti-cancer therapeutics", BIOSCIENCE REPORTS, vol. 35, no. 4, 12 June 2015 (2015-06-12), US, pages e00225 - e00225, XP055301629, ISSN: 0144-8463, DOI: 10.1042/BSR20150089

DESCRIPTION

Field of the invention

[0001] The present invention relates to antigen binding proteins and fragments thereof that specifically bind B cell maturation antigen (BCMA) and in particular human BCMA (hBCMA).

[0002] The present invention also concerns methods of treating diseases or disorders with said antigen binding fragments, pharmaceutical compositions comprising said antigen binding fragments and methods of manufacture. Other embodiments of the present invention will be apparent from the description below.

Background of the invention

[0003] BCMA (CD269 or TNFRSF17) is a member of the TNF receptor superfamily. It is a non-glycosylated integral membrane receptor for the ligands BAFF and APRIL. BCMA's ligands can also bind additional receptors: TACI (Transmembrane Activator and Calcium modulator and cyclophilin ligand Interactor), which binds APRIL and BAFF; as well as BAFF-R (BAFF Receptor or BR3), which shows restricted but high affinity for BAFF. Together, these receptors and their corresponding ligands regulate different aspects of humoral immunity, B-cell development and homeostasis.

[0004] BCMA's expression is typically restricted to the B-cell lineage and is reported to increase in terminal B-cell differentiation. BCMA is expressed by human plasma blasts, plasma cells from tonsils, spleen and bone marrow, but also by tonsillar memory B cells and by germinal centre B cells, which have a TACI-BAFFR low phenotype (Darce et al, 2007). BCMA is virtually absent on naive and memory B-cells (Novak et al., 2004a and b). The BCMA antigen is expressed on the cell surface so is accessible to the antibody, but is also expressed in the golgi. As suggested by its expression profile, BCMA signalling, typically linked with B-cell survival and proliferation, is important in the late stages of B-cell differentiation, as well as the survival of long lived bone marrow plasma cells (O'Connor et al., 2004) and plasmablasts (Avery et al., 2003). Furthermore, as BCMA binds APRIL with high affinity, the BCMA-APRIL signalling axis is suggested to predominate at the later stages of B-cell differentiation, perhaps being the most physiologically relevant interaction.

[0005] Multiple Myeloma (MM) is a clonal B-cell malignancy that occurs in multiple sites within the bone marrow before spreading to the circulation; either de novo, or as a progression from monoclonal gammopathy of undetermined significance (MGUS). It is commonly characterised by increases in paraprotein and osteoclast activity, as well as hypercalcaemia, cytopenia, renal dysfunction, hyperviscosity and peripheral neuropathy. Decreases in both normal antibody levels and numbers of neutrophils are also common, leading to a life threatening susceptibility to infection. BCMA has been implicated in the growth and survival of myeloma cell lines in vitro (Novak et al., 2004a and b; Moreaux et al., 2004).

[0006] BCMA expression (both transcript and protein) is reported to correlate with disease progression in MM. Using Affymetrix microarrays, it was demonstrated that the *TACI* and *BCMA* genes were overexpressed in Multiple Myeloma Cells (MMC) compared with their normal counterparts (Moreaux et al, 2004). Gene expression analysis has been used to compare human myeloma cells with purified plasma cells from patients with MGUS and from normal bone marrow as well as with primary tumour cells from B-cell lineage leukaemias (Bellucci et al, 2005). The BCMA gene was highly expressed in all myeloma samples. Although purified plasma cells from patients with MGUS had lower expression of BCMA, there was no significant difference when compared with the expression found in normal plasma cells or myeloma cells. In contrast, BCMA expression was significantly lower in B-cell Chronic Lymphocytic Leukaemia (CLL), pre-B Acute Lymphocytic Leukaemia (ALL) and T-cellALL (T-ALL). Mouse models that transgenically over-express BAFF or APRIL have a significant increase in B-cell lymphomas (Batten et al., 2004 - BAFF; Pinalles et al., 2004 - APRIL). In humans, excess BAFF and APRIL have been detected in the sera and micro-environments of patients with a number of B-cell malignancies, as well as other B-cell disorders.

[0007] An anti-BCMA IgG1 antibody SG1 has been shown to block the APRIL-dependent activation of NF κ B. This antibody apparently demonstrated cytotoxicity against multiple myeloma cells as both a naked antibody, or as antibody-drug conjugates with mono methyl auristatin F (MMAF). Enhanced ADCC (100-fold) was achieved through chimerisation with a human Fc region comprising mutations that enhance Fc γ RIIIA binding (Ryan Maureen et al: "Antibody targeting of B-cell maturation antigen on malignant plasma cells", Molecular Cancer Therapeutics, American Association of Cancer Research, US vol. 6, no. 11, 1 November 2007 (2007-11-01), pages 3009-3018, ISSN: 1535-7163) Chimeric (humanised) anti-BCMA IgG1 antibodies, including afucosylated versions thereof have also been disclosed previously and purported to have ADCC activity (WO 2010/104949 A2)

Brief Description of Figures

[0008]

Figure 1: FMAT Binding Assay - Figure showing the results of the FMAT assay for CA8 antibody binding to human and cyno BCMA expressing HEK293 cells. Human chimeric CA8 binds well to human and cyno BCMA expressing cells.

Figure 2: ELISA Binding Assay - Figure showing the ELISA results for CA8 antibodies binding to human and cyno BCMA recombinant proteins. This clearly shows that human chimeric CA8 antibodies bind to human and cyno BCMA proteins equally.

Figure 3: BiaCore Binding Assay - Figure showing the binding of CA8 to BCMA-Fc, TACI-Fc and BAFF-R-Fc proteins in the Biacore experiment. CA8 chimera antibody does not bind to TACI or BAFF-R proteins.

Figure 4: Cell binding assay - Figure showing binding of murine S307118G03, S3222110D07, S332121 F02 and S332126E04 to H929 multiple

myeloma cells and S3322110D07, S332121 F02 and S332126E04 to the BCMA transfected ARH77 cells as determined by FACS.

Multiple myeloma cell line H929 or ARH77-hBCMA 10B5 BCMA expressing transfectant cells were stained with either murine anti BCMA antibodies (solid histogram) or murine IgG2a isotype control (open histograms). Cells were analysed by FACS to detect antibody bound to the cells.

Figure 5: Cell binding assay - Figure showing binding of chimeric CA8 to a panel of multiple myeloma cell lines as determined by FACS. Binding to H929, OPM-2, JJN-3 and U266 was tested by flow cytometry and mean fluorescence intensity (MFI) values measured to determine binding. Synagis was used as an irrelevant isotype control.

Figure 6: Cell binding assay - Figure showing binding curves of humanised CA8 variants to BCMA transfected ARH77 cells (A) and multiple myeloma H929 cells (B) as determined by FACS.

Humanised variants J6M0, J6M1, J6M2, J9M0, J9M1 and J9M2 were tested by flow cytometry and mean fluorescence intensity (MFI) values measured to determine binding compared to the CA8 chimera.

Figure 7: Ligand neutralisation assays -

(A and B) Figure showing the ability of CA8 and J6M0 to neutralise binding of recombinant BAFF or APRIL to recombinant BCMA coated on an ELISA plate. OD values were used to calculate the antibody mediated inhibition of the maximal signal achieved by the relevant ligand alone binding to recombinant BCMA. Data is reported as percentage inhibition of the maximal signal. Antibodies tested were chimeric CA8 and humanised CA8 version J6M0 in both wild type and afucosylated (Potelligent) form.

(A) Neutralisation of BAFF ligand binding; (B)- Neutralisation APRIL ligand binding.

(C) - Figure showing the ability of J6M0 BCMA antibody in inhibition of BAFF or APRIL induced phosphorylation of NFKappaB in H929 cells. H-929 cells were washed 3 times to remove any sBCMA and resuspended in serum free medium. J6M0 potelligent antibody was added to a 96 well plate to give a final well concentrations up to 100ug/ml along with BAFF or APRIL ligand to give a final well concentration of 0.6 or 0.2 ug/ml respectively. H-929 cells were then plated at 7.5x104cells/well in serum free medium. 30 minutes later the cells were lysed and phosphorylated NFkappaB levels measured using a MSD pNFkappaB assay. MSD reader 502819. This is data from one independent experiments. Each data point is the mean/sd of two replicates.

Figure 8: ADCC assay - Figure showing ADCC activity of chimeric CA8 and defucosylated (Fc enhanced) CA8 with target cells expressing BCMA. Human NK cells were incubated with europium labelled ARH77 10B5 BCMA transfected target cells in the presence of varying concentrations of antibody. Europium release from the target cells was measured and specific lysis calculated. (A) ADCC dose response curves of chimeric CA8 compared to isotype control. (B) ADCC dose response curves for chimeric CA8 and defucosylated chimeric CA8 (Fc enhanced), against the BCMA expressing cell line ARH77 10B5.

Figure 9: ADCC assay - Figure showing ADCC assay on CA8 humanised antibodies using ARH77 BCMA expressing target cells.

Human PBMC were incubated with europium labelled ARH77 BCMA transfected target cells in the presence of a range of concentrations of the J5, J6, J7, J8 or J9 series of humanised CA8 antibodies. Europium release from the target cells was measured and specific lysis calculated. EC50 values are shown in ug/ml.

Figure 10: ADCC assay - Figure showing ADCC activity of chimeric, S332121 F02 (A), S33221 10D07 (B) S307118G03 (C) and humanised S307118G03 H3L0 (D) against ARH7710B5 target cells with purified NK cells as effector cells. Human NK target cells were incubated with europium labelled ARH77 10B5 BCMA transfected target cells in the presence of varying concentrations of antibody. Europium release from the target cells was measured and specific lysis calculated.

Figure 11: Viability assay dose response curves - Figure showing dose response curves in a cell viability assay for chimeric CA8 antibody, chimeric CA8-vcMMAE and chimeric CA8-mcMMAF antibody-drug conjugates in human multiple myeloma cell lines (A) NCI-H929 (B) U266-B1 (C) JJN3 and (D) OPM2. Antibody was added to the cells and the number of viable cells after 96 hours measured using CellTiterGlo. Data points represent the mean of triplicate CellTiterGlo measurements. Error bars represent standard error.

Figure 12: Impact of CA8 chimeric antibody on cell cycle.

1. (A) Cell cycle histograms of NCI-H929 cells treated with unconjugated chimeric CA8, chimeric CA8-vcMMAE ADC or chimeric CA8-mcMMAF ADC at 50ng/mL for the timepoints indicated. Paclitaxel (100nM) was used as a positive control for G2/M cell cycle arrest and cell death. Control human IgG1 was used as a negative control. Cell cycle analysis was carried out at the times shown on the graphs.
2. (B) Quantification of the 4N DNA cell population indicative of G2/M arrest and (C) sub-2N DNA cell population indicative of cell death for each of the treatments indicated. Cells were seeded in 12-well plates (2×10^5 cells per well in 1mL of RPMI + 10% FBS). Antibody or ADC was added 6 hours after cell seeding.

Figure 13: Impact of chimeric CA8 on phospho-histone H3.

Chimeric CA8 ADC treatment results in increased phospho-Histone H3 staining of NCI-H929 cells. (A,B) Dot plots of cells stained with propidium iodide to measure DNA content (FL3-H) x-axis and anti-phospho-Histone H3 (Thr11) antibody (FL1-H) y-axis after treatment with either Control IgG (A) or chimeric CA8-mcMMAF (B). (C) Quantification of phospho-Histone H3 positive NCI-H929 cells after a 48 hour treatment with the indicated concentrations of chimeric CA8 ADCs. Paclitaxel (100nM) was used as a positive control for mitotic arrest and control chimera IgG1 was used as a negative control. Cells were seeded in 12-well plates (2×10^5 cells per well in 1mL of RPMI + 10% FBS). Antibody or ADC was added 6 hours after cell seeding.

Figure 14: Impact of chimeric CA8 on Annexin-V.

Chimeric CA8 ADC treatment results in increased Annexin-V staining of NCI-H929 cells. (A) Histograms of Annexin-V-FITC (FL1-H; top panels) and Live cell propidium iodide staining (FL3-H; bottom panels) after treatment with increasing concentrations of chimeric CA8 ADCs (B) Quantification of Annexin-V positive NCI-H929 cells after a 96 hour treatment with the indicated concentrations of chimeric CA8 ADCs. Paclitaxel (100nM) was used as a positive control for apoptosis and control chimera IgG1 was used as a negative control. Cells were seeded in 12-well plates (2×10^5 cells per well in 1mL of RPMI + 10% FBS). Antibody or ADC was added 6 hours after cell seeding.

Figure 15: Viability assay dose response curves - Figure showing dose response curves for the unconjugated (Naked) and vcMMAE and mcmMAF antibody-drug conjugates of chimeric CA8 or humanized J6M0 antibodies. Antibody drug conjugates were tested against human multiple myeloma cell lines NCI-H929 and OPM2.

Figure 16: Viability assay dose response curves - Figure showing dose response curves for the unconjugated antibodies, vcMMAE and mcmMAF antibody-drug conjugates of murine anti-BCMA antibodies S332121F02, S322110D07, S332126E04 and S307118G03 in human multiple myeloma cell lines NCI-H929 and U266-B1.

Figure 17 ADCC activity of ADC J6M0 molecules - Figure showing ADCC assay on J6M0 antibodies using ARH77 BCMA expressing target cells. Human PBMC were incubated with europium labelled ARH77 BCMA transfected target cells in the presence of a range of concentrations of J6M0 WT and potelligent BCMA antibodies conjugated to MMAE, MMAF, or unconjugated. Europium release was monitored on the Victor 2 1420 multilabel reader.

Figure 18 ADCC dose response curves of CA8 J6M0 Potelligent against a panel of 5 multiple myeloma lines - Human PBMC were incubated with multiple myeloma target cells in the presence of varying concentrations of CA8 J6M0 potelligent antibody at an E:T ratio of 50:1 for 18 hours. The percentage of target cells remaining in the effector plus target mixture was then measured by FACS using a fluorescently labelled anti-CD138 antibody to detect the target cells and the percent cytotoxicity calculated. A) Example dose response curves for CA8 J6M0 potelligent against the five multiple myeloma cell lines tested. Each data point is from a singlicate value.

Figure 19 Effect of dose escalation of J6M0 and drug conjugated J6M0 on the growth and establishment of NCI-H929 cells in CB.17 SCID mice Calculated tumour volumes of NCI-H929 tumours in CB17 SCID mice following twice weekly intraperitoneal dosing of either 50 or 100ug J6M0 anti-BCMA or IgG1 isotype control unconjugated, or conjugated to MMAE or MMAF for 2 weeks. Data points represent mean tumour volume of n=5 per group

Figure 20- Determination of soluble BCMA levels in serum from healthy volunteers and myeloma patients. Serum samples were collected from MM patient samples were from a variety of stages (progressive disease, remission, relapsed, newly diagnosed, and others). The samples shown in the figure are those from serum diluted 1/500 prior to the assay.

A Human BCMATNFRSF17 sandwich ELISA kit from R& D Systems which measures soluble human BCMA levels was used to detect BCMA following the standard protocol provided with the kit.

Summary of the Invention

[0009] The invention is defined by the appended claims.

[0010] Any references in the description to methods of treatment refer to the compounds, pharmaceutical compositions and medicaments of the present invention for use in a method for treatment of the human (or animal) body by therapy (or for diagnosis).

Detailed Description of the Invention

[0011] The present invention provides an anti B-Cell Maturation Antigen (CD269) antibody comprising a heavy chain variable region of SEQ. ID. NO:23 and a light chain variable region of SEQ. ID. NO:31.

[0012] Further provided is an anti B-Cell Maturation Antigen (CD269) antibody which comprises a heavy chain of SEQ. ID. NO:55 and a light chain of SEQ. ID. NO:63. In one aspect the anti B-Cell Maturation Antigen (CD269) antibody of the invention has enhanced binding to FcγRIIIA or has enhanced FcγRIIIA mediated effector function, for example in one aspect of the invention as herein provided the antibody is defucosylated.

[0013] Herein disclosed are antigen binding proteins which bind to membrane bound targets and wherein the antigen binding protein is capable of internalisation. In a further embodiment there is provided an immunoconjugate comprising the antigen binding protein of the present invention and a cytotoxic agent. In a further embodiment the antigen binding protein has ADCC effector function for example the antigen binding protein has enhanced ADCC effector function.

[0014] In one such embodiment there is provided antigen binding proteins or fragments thereof which specifically bind to BCMA, for example which specifically binds human BCMA (hBCMA) and which inhibit the binding of BAFF and/or APRIL to the BCMA receptor.

[0015] In a further embodiment the antigen binding proteins or fragments specifically bind to BCMA and inhibit the binding of BAFF and/or APRIL to BCMA wherein the antigen binding proteins or fragments thereof have the ability to bind to FcγRIIIA and mediate FcγRIIIA mediated effector functions, or have enhanced FcγRIIIA mediated effector function. In one embodiment of the invention as herein provided the antigen binding proteins are capable of internalisation.

[0016] In one aspect of the invention there is provided an antigen binding protein according to the invention as herein described which binds to non-membrane bound BCMA, for example to serum BCMA.

[0017] In one aspect of the invention the antigen binding protein has enhanced effector function. In another aspect the antigen binding protein is conjugated to a cytotoxic agent. In yet a further embodiment the antigen binding protein has both enhanced effector function and is conjugated to a cytotoxic agent.

[0018] The antigen binding proteins of the present invention may comprise heavy chain variable regions and light chain variable regions of the invention which may be formatted into the structure of a natural antibody or functional fragment or equivalent thereof. An antigen binding protein of the invention may therefore comprise the VH regions of the invention formatted into a full length antibody, a (Fab')2 fragment, a Fab fragment, or equivalent thereof (such as scFV, bi- tri- or tetra-bodies, Tandabs etc.), when paired with an appropriate light chain. The antibody may be an IgG1, IgG2, IgG3, or IgG4; or IgM; IgA, IgE or IgD or a modified variant thereof. The constant domain of the antibody heavy chain may be selected accordingly. The light chain constant domain may be a kappa or lambda constant domain. Furthermore, the antigen binding protein may comprise modifications of all classes e.g. IgG dimers, Fc mutants that no longer bind Fc receptors or mediate C1q binding. The antigen binding protein may also be a chimeric antibody of the type described in WO86/01533 which comprises an antigen binding region and a non-immunoglobulin region.

[0019] The constant region is selected according to any functionality required e.g. an IgG1 may demonstrate lytic ability through binding to complement and/or will mediate ADCC (antibody dependent cell cytotoxicity).

[0020] In one aspect the antigen binding protein is an antibody or antigen binding fragment thereof comprising one or more CDR's according to the invention described herein, or one or both of the heavy or light chain variable domains according to the invention described herein. In one embodiment the antigen binding protein binds primate BCMA. In one such embodiment the antigen binding protein additionally binds non-human primate BCMA, for example cynomolgus macaque monkey BCMA.

[0021] In another aspect the antigen binding protein is selected from the group consisting of a dAb, Fab, Fab', F(ab')₂, Fv, diabody, triabody, tetrabody, minibody, and a minibody.

[0022] In one aspect of the present invention the antigen binding protein is a humanised or chimaeric antibody, in a further aspect the antibody is humanised.

[0023] In one aspect the antibody is a monoclonal antibody.

[0024] In another aspect the antigen binding protein binds to human BCMA with high affinity for example when measured by Biacore the antigen binding protein binds to human BCMA with an affinity of 20nM or less or an affinity of 15nM or less or an affinity of 5nM or less or an affinity of 1000 pM or less or an affinity of 500pM or less or an affinity of 400pM or less, or 300pM or less or for example about 120pM. In a further embodiment the antigen binding protein binds to human BCMA when measured by Biacore of between about 100pM and about 500pM or between about 100pM and about 400pM, or between about 100pM and about 300pM. In one embodiment of the present invention the antigen binding protein binds BCMA with an affinity of less than 150pm.

[0025] In one such embodiment, this is measured by Biacore, for example as set out in Example 4.

[0026] In another aspect the antigen binding protein binds to human BCMA and neutralises the binding of the ligands BAFF and/or APRIL to the BCMA receptor in a cell neutralisation assay wherein the antigen binding protein has an IC50 of between about 1 nM and about 500nM, or between about 1 nM and about 100nM, or between about 1 nM and about 50nM, or between about 1 nM and about 25nM, or between about 5nM and about 15nM. In a further embodiment of the present invention the antigen binding protein binds BCMA and neutralises BCMA in a cell neutralisation assay wherein the antigen binding protein has an IC50 of about 10nM.

[0027] In one such embodiment, this is measured by a cell neutralisation assay, for example as set out in Example 4.6.

[0028] The antigen binding proteins, for example antibodies of the present invention may be produced by transfection of a host cell with an expression vector comprising the coding sequence for the antigen binding protein of the invention. An expression vector or recombinant plasmid is produced by placing these coding sequences for the antigen binding protein in operative association with conventional regulatory control sequences capable of controlling the replication and expression in, and/or secretion from, a host cell. Regulatory sequences include promoter sequences, e.g., CMV promoter, and signal sequences which can be derived from other known antibodies. Similarly, a second expression vector can be produced having a DNA sequence which encodes a complementary antigen binding protein light or heavy chain. In certain embodiments this second expression vector is identical to the first except insofar as the coding sequences and selectable markers are concerned, so to ensure as far as possible that each polypeptide chain is functionally expressed. Alternatively, the heavy and light chain coding sequences for the antigen binding protein may reside on a single vector.

[0029] A selected host cell is co-transfected by conventional techniques with both the first and second vectors (or simply transfected by a single vector) to create the transfected host cell of the invention comprising both the recombinant or synthetic light and heavy chains. The transfected cell is then cultured by conventional techniques to produce the engineered antigen binding protein of the invention. The antigen binding protein which includes the association of both the recombinant heavy chain and/or light chain is screened from culture by appropriate assay, such as ELISA or RIA. Similar conventional techniques may be employed to construct other antigen binding proteins.

[0030] Suitable vectors for the cloning and subcloning steps employed in the methods and construction of the compositions of this invention may be selected by one of skill in the art. For example, the conventional pUC series of cloning vectors may be used. One vector, pUC19, is commercially available from supply houses, such as Amersham (Buckinghamshire, United Kingdom) or Pharmacia (Uppsala, Sweden). Additionally, any vector which is capable of replicating readily, has an abundance of cloning sites and selectable genes (e.g., antibiotic resistance), and is easily manipulated may be used for cloning. Thus, the selection of the cloning vector is not a limiting factor in this invention.

[0031] The expression vectors may also be characterized by genes suitable for amplifying expression of the heterologous DNA sequences, e.g., the mammalian dihydrofolate reductase gene (DHFR). Other vector sequences include a poly A signal sequence, such as from bovine growth hormone (BGH) and the betaglobin promoter sequence (betaglopro). The expression vectors useful herein may be synthesized by techniques well known to those skilled in this art.

[0032] The components of such vectors, e.g. replicons, selection genes, enhancers, promoters, signal sequences and the like, may be obtained from commercial or natural sources or synthesized by known procedures for use in directing the expression and/or secretion of the product of the recombinant DNA in a selected host. Other appropriate expression vectors of which numerous types are known in the art for mammalian, bacterial, insect, yeast, and fungal expression may also be selected for this purpose.

[0033] The present invention also encompasses a cell line transfected with a recombinant plasmid containing the coding sequences of the antigen binding proteins of the present invention. Host cells useful for the cloning and other manipulations of these cloning vectors are also conventional. However, cells from various strains of *E. Coli* may be used for replication of the cloning vectors and other steps in the construction of antigen binding proteins of this invention.

[0034] Suitable host cells or cell lines for the expression of the antigen binding proteins of the invention include mammalian cells such as NS0, Sp2/0, CHO (e.g. DG44), COS, HEK, a fibroblast cell (e.g., 3T3), and myeloma cells, for example it may be expressed in a CHO or a myeloma cell. Human cells may be used, thus enabling the molecule to be modified with human glycosylation patterns. Alternatively, other eukaryotic cell lines may be employed. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. See, e.g., Sambrook et al., cited above.

[0035] Bacterial cells may prove useful as host cells suitable for the expression of the recombinant Fabs or other embodiments of the present invention (see, e.g., Plückthun, A., Immunol. Rev., 130:151-188 (1992)). However, due to the tendency of proteins expressed in bacterial cells to be in an unfolded or improperly folded form or in a non-glycosylated form, any recombinant Fab produced in a bacterial cell would have to be screened for retention of antigen binding ability. If the molecule expressed by the bacterial cell was produced in a properly folded form, that bacterial cell would be a desirable host, or in alternative embodiments the molecule may express in the bacterial host and then be subsequently re-folded. For example, various strains of *E. Coli* used for expression are well-known as host cells in the field of biotechnology. Various strains of *B. Subtilis*, *Streptomyces*, other bacilli and the like may also be employed in this method.

[0036] Where desired, strains of yeast cells known to those skilled in the art are also available as host cells, as well as insect cells, e.g. *Drosophila* and *Lepidoptera* and viral expression systems. See, e.g. Miller et al., Genetic Engineering, 8:277-298, Plenum Press (1986) and references cited therein.

[0037] The general methods by which the vectors may be constructed, the transfection methods required to produce the host cells of the invention, and culture methods necessary to produce the antigen binding protein of the invention from such host cell may all be conventional techniques. Typically, the culture method of the present invention is a serum-free culture method, usually by culturing cells serum-free in suspension. Likewise, once produced, the antigen binding proteins of the invention may be purified from the cell culture contents according to standard procedures of the art, including ammonium peroxide precipitation, affinity columns, column chromatography, gel electrophoresis and the like. Such techniques are within the skill of the art and do not limit this invention. For example, preparations of altered antibodies are described in WO 99/58679 and WO 96/16990.

[0038] Yet another method of expression of the antigen binding proteins may utilize expression in a transgenic animal, such as described in U. S. Patent No. 4,873,316. This relates to an expression system using the animal's casein promoter which when transgenically incorporated into a mammal permits the female to produce the desired recombinant protein in its milk.

[0039] In a further embodiment of the invention there is provided a method of producing an antibody of the invention which method comprises the step of culturing a host cell transformed or transfected with a vector encoding the light and/or heavy chain of the antibody of the invention and recovering the antibody thereby produced.

[0040] In accordance with the present invention there is provided a method of producing an anti-BCMA antibody of the present invention which binds to and neutralises the activity of human BCMA which method comprises the steps of:

providing a first vector encoding a heavy chain of the antibody;

providing a second vector encoding a light chain of the antibody;

transforming a mammalian host cell (e.g. CHO) with said first and second vectors;

culturing the host cell of step (c) under conditions conducive to the secretion of the antibody from said host cell into said culture media;

recovering the secreted antibody of step (d).

[0041] Once expressed by the desired method, the antibody is then examined for in vitro activity by use of an appropriate assay. Presently conventional ELISA assay formats are employed to assess qualitative and quantitative binding of the antibody to BCMA. Additionally, other in vitro assays may also be used to verify neutralizing efficacy prior to subsequent human clinical studies performed to evaluate the persistence of the antibody in the body despite the usual clearance mechanisms.

[0042] The dose and duration of treatment relates to the relative duration of the molecules of the present invention in the human circulation, and can be adjusted by one of skill in the art depending upon the condition being treated and the general health of the patient. It is envisaged that repeated dosing (e.g. once a week or once every two weeks or once every 3 weeks) over an extended time period (e.g. four to six months) maybe required to achieve maximal therapeutic efficacy.

[0043] In one embodiment of the present invention there is provided a recombinant transformed, transfected or transduced host cell comprising at least one expression cassette, for example where the expression cassette comprises a polynucleotide encoding a heavy chain of an antigen binding

protein according to the invention described herein and further comprises a polynucleotide encoding a light chain of an antigen binding protein according to the invention described herein or where there are two expression cassettes and the 1st encodes the light chain and the second encodes the heavy chain. For example in one embodiment the first expression cassette comprises a polynucleotide encoding a heavy chain of an antigen binding protein comprising a constant region or antigen binding fragment thereof which is linked to a constant region according to the invention described herein and further comprises a second cassette comprising a polynucleotide encoding a light chain of an antigen binding protein comprising a constant region or antigen binding fragment thereof which is linked to a constant region according to the invention described herein for example the first expression cassette comprises a polynucleotide encoding a heavy chain selected from SEQ. ID. NO:56, and a second expression cassette comprising a polynucleotide encoding a light chain selected from SEQ. ID. NO: 64

[0044] In another embodiment of the invention there is provided a stably transformed host cell comprising a vector comprising one or more expression cassettes encoding a heavy chain and/or a light chain of the antibody comprising a constant region or antigen binding fragment thereof which is linked to a constant region as described herein. For example such host cells may comprise a first vector encoding the light chain and a second vector encoding the heavy chain, for example the first vector encodes a heavy chain selected from SEQ. ID. NO: 55, and a second vector encoding a light chain for example the light chain of SEQ ID NO: 63.

[0045] In one such example the first vector encodes a heavy chain selected from SEQ. ID. NO: 55 and a second vector encoding a light chain for example the light chain of SEQ ID NO: 63.

[0046] In another embodiment of the present invention there is provided a host cell according to the invention described herein wherein the cell is eukaryotic, for example where the cell is mammalian. Examples of such cell lines include CHO or NS0.

[0047] In another embodiment of the present invention there is provided a method for the production of an antibody comprising a constant region or antigen binding fragment thereof which is linked to a constant region according to the invention described herein which method comprises the step of culturing a host cell in a culture media, for example serum- free culture media.

[0048] In another embodiment of the present invention there is provided a method according to the invention described herein wherein said antibody is further purified to at least 95% or greater (e.g. 98% or greater) with respect to said antibody containing serum- free culture media.

[0049] In yet another embodiment there is provided a pharmaceutical composition comprising an antigen binding protein and a pharmaceutically acceptable carrier.

[0050] In another embodiment of the present invention there is provided a kit-of-parts comprising the composition according to the invention described herein described together with instructions for use.

[0051] The mode of administration of the therapeutic agent of the invention may be any suitable route which delivers the agent to the host. The antigen binding proteins, and pharmaceutical compositions of the invention are particularly useful for parenteral administration, i.e., subcutaneously (s.c.), intrathecally, intraperitoneally, intramuscularly (i.m.) or intravenously (i.v.). In one such embodiment the antigen binding proteins of the present invention are administered intravenously or subcutaneously.

[0052] Therapeutic agents of the invention may be prepared as pharmaceutical compositions containing an effective amount of the antigen binding protein of the invention as an active ingredient in a pharmaceutically acceptable carrier. In one embodiment the prophylactic agent of the invention is an aqueous suspension or solution containing the antigen binding protein in a form ready for injection. In one embodiment the suspension or solution is buffered at physiological pH. In one embodiment the compositions for parenteral administration will comprise a solution of the antigen binding protein of the invention or a cocktail thereof dissolved in a pharmaceutically acceptable carrier. In one embodiment the carrier is an aqueous carrier. A variety of aqueous carriers may be employed, e.g., 0.9% saline, 0.3% glycine, and the like. These solutions may be made sterile and generally free of particulate matter. These solutions may be sterilized by conventional, well known sterilization techniques (e.g., filtration). The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, etc. The concentration of the antigen binding protein of the invention in such pharmaceutical formulation can vary widely, i.e., from less than about 0.5%, usually at or at least about 1% to as much as about 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, etc., according to the particular mode of administration selected.

[0053] Thus, a pharmaceutical composition of the invention for intravenous infusion could be made up to contain about 250 ml of sterile Ringer's solution, and about 1 to about 30 or 5 mg to about 25 mg of an antigen binding protein of the invention per ml of Ringer's solution. Actual methods for preparing parenterally administrable compositions are well known or will be apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Science, 15th ed., Mack Publishing Company, Easton, Pennsylvania. For the preparation of intravenously administrable antigen binding protein formulations of the invention see Lasmar U and Parkins D "The formulation of Biopharmaceutical products", Pharma. Sci.Tech.today, page 129-137, Vol.3 (3rd April 2000); Wang, W "Instability, stabilisation and formulation of liquid protein pharmaceuticals", Int. J. Pharm 185 (1999) 129-188; Stability of Protein Pharmaceuticals Part A and B ed Ahern T.J., Manning M.C., New York, NY: Plenum Press (1992); Akers,M.J. "Excipient-Drug interactions in Parenteral Formulations", J.Pharm Sci 91 (2002) 2283-2300; Imamura, K et al "Effects of types of sugar on stabilization of Protein in the dried state", J Pharm Sci 92 (2003) 266-274; Izutsu, Kkojima, S. "Excipient crystallinity and its protein-structure-stabilizing effect during freeze-drying", J Pharm. Pharmacol, 54 (2002) 1033-1039; Johnson, R, "Mannitol-sucrose mixtures-versatile formulations for protein peroxidise19g19n", J. Pharm. Sci, 91 (2002) 914-922, and Ha,E Wang W, Wang Y.j. "Peroxide formation in polysorbate 80 and protein stability", J. Pharm Sci, 91, 2252-2264,(2002)..

[0054] In one embodiment the therapeutic agent of the invention, when in a pharmaceutical preparation, is present in unit dose forms. The appropriate therapeutically effective dose will be determined readily by those of skill in the art. Suitable doses may be calculated for patients according to their weight, for example suitable doses may be in the range of about 0.1 to about 20mg/kg, for example about 1 to about 20mg/kg, for example about 10 to about 20mg/kg or for example about 1 to about 15mg/kg, for example about 10 to about 15mg/kg. To effectively treat conditions such as Multiple myeloma, SLE or IPT in a human, suitable doses may be within the range of about 0.1 to about 1000 mg, for example

about 0.1 to about 500mg, for example about 500mg, for example about 0.1 to about 100mg, or about 0.1 to about 80mg, or about 0.1 to about 60mg, or about 0.1 to about 40mg, or for example about 1 to about 100mg, or about 1 to about 50mg, of an antigen binding protein of this invention, which may be administered parenterally, for example subcutaneously, intravenously or intramuscularly. Such dose may, if necessary, be repeated at appropriate time intervals selected as appropriate by a physician.

[0055] The antigen binding proteins described herein can be lyophilized for storage and reconstituted in a suitable carrier prior to use. This technique has been shown to be effective with conventional immunoglobulins and art-known peroxidise and reconstitution techniques can be employed.

[0056] In another aspect of the invention there is provided an antigen binding protein as herein described for use in a medicament.

[0057] In one aspect of the disclosure there is provided an antigen binding protein according to the invention as herein described for use in the treatment of rheumatoid arthritis, Type 1 Diabetes Mellitus, multiple sclerosis or psoriasis wherein said method comprises the step of administering to said patient a therapeutically effective amount of the antigen binding protein as described herein.

[0058] In one embodiment of the present disclosure, methods are provided for treating cancer in a human comprising administering to said human an antigen binding protein that specifically binds to BCMA. In some instances the antigen binding protein is part of an immunoconjugate.

[0059] In another aspect of the present invention there is provided an antigen binding protein according to the invention as herein described for use in the treatment of a B-cell mediated or plasma cell mediated disease or antibody mediated disease or disorder selected from Multiple Myeloma (MM), chronic lymphocytic leukemia (CLL), Non-secretory multiple myeloma, Smoldering multiple myeloma, Monoclonal gammopathy of undetermined significance (MGUS), Solitary plasmacytoma (Bone, Extramedullary), Lymphoplasmacytic lymphoma (LPL), Waldenstrom's Macroglobulinemia, Plasma cell leukemia, Primary Amyloidosis (AL), Heavy chain disease, Systemic lupus erythematosus (SLE), POEMS syndrome / osteosclerotic myeloma, Type I and II cryoglobulinemia, Light chain deposition disease, Goodpasture's syndrome, Idiopathic thrombocytopenic purpura (ITP), Acute glomerulonephritis, Pemphigus and Pemphigoid disorders, and Epidermolysis bullosa acquisita; or any Non-Hodgkin's Lymphoma B-cell leukemia or Hodgkin's lymphoma (HL) with BCMA expression or any diseases in which patients develop neutralising antibodies to recombinant protein replacement therapy wherein said method comprises the step of administering to said patient a therapeutically effective amount of the antigen binding protein as described herein.

[0060] B-cell disorders can be divided into defects of B-cell development/immunoglobulin production (immunodeficiencies) and excessive/uncontrolled proliferation (lymphomas, leukemias). As used herein, B-cell disorder refers to both types of diseases, and methods are provided for treating B-cell disorders with an antigen binding protein.

[0061] In a particular aspect, the disease or disorder is selected from the group consisting of Multiple Myeloma (MM), Chronic Lymphocytic Leukaemia (CLL), Solitary Plasmacytoma (Bone, Extramedullary), Waldenstrom's Macroglobulinemia.

[0062] In one aspect of the present invention the disease is Multiple Myeloma, Smoldering Multiple Myeloma (SMM) or Solitary Plasmacytoma (Bone, Extramedullary).

[0063] In one aspect of the present invention the disease is Multiple Myeloma.

[0064] In one aspect of the present disclosure the disease is Systemic lupus erythematosus (SLE)

[0065] In one aspect of the present disclosure the disease is Idiopathic thrombocytopenic purpura (ITP) Use of the antigen binding protein as described herein in the manufacture of a medicament for the treatment of diseases and disorders as described herein is also provided.

[0066] In another aspect of the disclosure there is provided the use of the antigen binding protein as described herein for use in the treatment or prophylaxis of an antibody mediated or plasma cell mediated disease or disorder selected from rheumatoid arthritis, Type 1 Diabetes Mellitus, multiple sclerosis or psoriasis.

[0067] In another aspect of the disclosure there is provided the use of the antigen binding protein as described herein for use in the treatment or prophylaxis of an antibody mediated or plasma cell mediated disease or disorder selected from Multiple Myeloma (MM), chronic lymphocytic leukemia (CLL), Monoclonal gammopathy of undetermined significance (MGUS), Smoldering multiple myeloma (SMM), Solitary Plasmacytoma (Bone, Extramedullary), Waldenstrom's Macroglobulinemia , Primary Amyloidosis (AL), Heavy chain disease, Systemic lupus erythematosus (SLE), POEMS syndrome / osteosclerotic myeloma, Type I and II cryoglobulinemia, Light chain deposition disease, Goodpastures syndrome, Idiopathic thrombocytopenic purpura (ITP), Acute glomerulonephritis, Pemphigus and Pemphigoid disorders and Epidermolysis bullosa acquisita, any Non-Hodgkin Lymphoma and Leukemia with BCMA expression or any diseases in which patients develop neutralising antibodies to recombinant protein replacement therapy wherein said method comprises the step of administering to said patient a therapeutically effective amount of the antigen binding protein as described herein.

[0068] In one aspect, the invention provides a pharmaceutical composition comprising an antigen binding protein of the present invention or a functional fragment thereof and a pharmaceutically acceptable carrier for treatment or prophylaxis of rheumatoid arthritis, Type 1 Diabetes Mellitus, multiple sclerosis or psoriasis or an antibody mediated or plasma cell mediated disease or disorder selected from Multiple Myeloma (MM), chronic lymphocytic leukemia (CLL), Monoclonal gammopathy of undetermined significance (MGUS), Smoldering multiple myeloma (SMM), Solitary Plasmacytoma (Bone, Extramedullary), Waldenstrom's Macroglobulinemia , Primary Amyloidosis (AL), Heavy chain disease, Systemic lupus erythematosus (SLE), POEMS syndrome / osteosclerotic myeloma, Type I and II cryoglobulinemia, Light chain deposition disease, Goodpastures syndrome, Idiopathic thrombocytopenic purpura (ITP), Acute glomerulonephritis, Pemphigus and Pemphigoid disorders and Epidermolysis bullosa acquisita, any Non-Hodgkin Lymphoma and Leukemia with BCMA expression or any diseases in which patients develop neutralising antibodies to recombinant protein replacement therapy wherein said method comprises the step of administering to said patient a therapeutically effective amount

of the antigen binding protein as described herein.

[0069] In another embodiment of the present disclosure there is provided a method of treating a human patient afflicted with rheumatoid arthritis, Type 1 Diabetes Mellitus, multiple sclerosis or psoriasis or an antibody mediated or plasma cell mediated disorder or disease which method comprises the step of administering a therapeutically effective amount of the antigen binding protein according to the invention as described herein.

[0070] In another aspect of the present disclosure there is provided an antigen binding protein according to the invention as herein described for use in the treatment of an antibody mediated or plasma cell mediated disease or disorder selected from Multiple Myeloma (MM), Chronic Lymphocytic Leukaemia (CLL)Monoclonal gammopathy of undetermined significance (MGUS), Smoldering multiple myeloma (SMM), Solitary Plasmacytoma (Bone, Extramedullary), Waldenstrom's Macroglobulinemia , Primary Amyloidosis (AL), Heavy chain disease, Systemic lupus erythematosus (SLE), POEMS syndrome / osteosclerotic myeloma, Type I and II cryoglobulinemia, Light chain deposition disease, Goodpastures syndrome, Idiopathic thrombocytopenic purpura (ITP), Acute glomerulonephritis, Pemphigus and Pemphigoid disorders and Epidermolysis bullosa acquisita, any Non-Hodgkin Lymphoma and Leukemia with BCMA expression or any diseases in which patients develop neutralising antibodies to recombinant protein replacement therapy wherein said method comprises the step of administering a pharmaceutical composition comprising an antigen binding protein according to the invention herein in combination with a pharmaceutically acceptable carrier.

[0071] In a further embodiment there is provided a method of treating a human patient afflicted with Multiple Myeloma (MM).

Definitions

[0072] As used herein, the terms "cancer," "neoplasm," and "tumor" are used interchangeably and, in either the singular or plural form, refer to cells that have undergone a malignant transformation that makes them pathological to the host organism. Primary cancer cells can be readily distinguished from noncancerous cells by well-established techniques, particularly histological examination. The definition of a cancer cell, as used herein, includes not only a primary cancer cell, but any cell derived from a cancer cell ancestor. This includes metastasized cancer cells, and *in vitro* cultures and cell lines derived from cancer cells. When referring to a type of cancer that normally manifests as a solid tumor, a "clinically detectable" tumor is one that is detectable on the basis of tumor mass; e.g., by procedures such as computed tomography (CT) scan, magnetic resonance imaging (MRI), X-ray, ultrasound or palpation on physical examination, and/or which is detectable because of the expression of one or more cancer-specific antigens in a sample obtainable from a patient. Tumors may be a hematopoietic (or hematologic or hematological or blood-related) cancer, for example, cancers derived from blood cells or immune cells, which may be referred to as "liquid tumors." Specific examples of clinical conditions based on hematologic tumors include leukemias such as chronic myelocytic leukemia, acute myelocytic leukemia, chronic lymphocytic leukemia and acute lymphocytic leukemia; plasma cell malignancies such as multiple myeloma, MGUS and Waldenstrom's macroglobulinemia; lymphomas such as non-Hodgkin's lymphoma, Hodgkin's lymphoma; and the like.

[0073] The cancer may be any cancer in which an abnormal number of blast cells or unwanted cell proliferation is present or that is diagnosed as a hematological cancer, including both lymphoid and myeloid malignancies. Myeloid malignancies include, but are not limited to, acute myeloid (or myelocytic or myelogenous or myeloblastic) leukemia (undifferentiated or differentiated), acute promyeloid (or promyelocytic or promyelogenous or promyeloblastic) leukemia, acute myelomonocytic (or myelomonoblastic) leukemia, acute monocytic (or monoblastic) leukemia, erythroleukemia and megakaryocytic (or megakaryoblastic) leukemia. These leukemias may be referred together as acute myeloid (or myelocytic or myelogenous) leukemia (AML). Myeloid malignancies also include myeloproliferative disorders (MPD) which include, but are not limited to, chronic myelogenous (or myeloid) leukemia (CML), chronic myelomonocytic leukemia (CMML), essential thrombocythemia (or thrombocytosis), and polycythemia vera (PCV). Myeloid malignancies also include myelodysplasia (or myelodysplastic syndrome or MDS), which may be referred to as refractory anemia (RA), refractory anemia with excess blasts (RAEB), and refractory anemia with excess blasts in transformation (RAEBT); as well as myelofibrosis (MFS) with or without agnogenic myeloid metaplasia.

[0074] Hematopoietic cancers also include lymphoid malignancies, which may affect the lymph nodes, spleens, bone marrow, peripheral blood, and/or extranodal sites. Lymphoid cancers include B-cell malignancies, which include, but are not limited to, B-cell non-Hodgkin's lymphomas (B-NHLs). B-NHLs may be indolent (or low-grade), intermediate-grade (or aggressive) or high-grade (very aggressive). Indolent B-cell lymphomas include follicular lymphoma (FL); small lymphocytic lymphoma (SLL); marginal zone lymphoma (MZL) including nodal MZL, extranodal MZL, splenic MZL and splenic MZL with villous lymphocytes; lymphoplasmacytic lymphoma (LPL); and mucosaassociated-lymphoid tissue (MALT or extranodal marginal zone) lymphoma. Intermediate-grade B-NHLs include mantle cell lymphoma (MCL) with or without leukemic involvement, diffuse large cell lymphoma (DLBCL), follicular large cell (or grade 3 or grade 3B) lymphoma, and primary mediastinal lymphoma (PML). High-grade B-NHLs include Burkitt's lymphoma (BL), Burkitt-like lymphoma, small non-cleaved cell lymphoma (SNCL) and lymphoblastic lymphoma. Other B-NHLs include immunoblastic lymphoma (or immunocytoma), primary effusion lymphoma, HIV associated (or AIDS related) lymphomas, and post-transplant lymphoproliferative disorder (PTLD) or lymphoma. B-cell malignancies also include, but are not limited to, chronic lymphocytic leukemia (CLL), prolymphocytic leukemia (PLL), Waldenstrom's macroglobulinemia (WM), hairy cell leukemia (HCL), large granular lymphocyte (LGL) leukemia, acute lymphoid (or lymphocytic or lymphoblastic) leukemia, and Castleman's disease. NHL may also include T-cell non-Hodgkin's lymphoma s(T-NHLs), which include, but are not limited to T-cell non-Hodgkin's lymphoma not otherwise specified (NOS), peripheral T-cell lymphoma (PTCL), anaplastic large cell lymphoma (ALCL), angioimmunoblastic lymphoid disorder (AILD), nasal natural killer (NK) cell / T-cell lymphoma, gamma/delta lymphoma, cutaneous T cell lymphoma, mycosis fungoides, and Sezary syndrome.

[0075] Hematopoietic cancers also include Hodgkin's lymphoma (or disease) including classical Hodgkin's lymphoma, nodular sclerosing Hodgkin's lymphoma, mixed cellularity Hodgkin's lymphoma, lymphocyte predominant (LP) Hodgkin's lymphoma, nodular LP Hodgkin's lymphoma, and lymphocyte depleted Hodgkin's lymphoma. Hematopoietic cancers also include plasma cell diseases or cancers such as multiple myeloma (MM) including smoldering MM, monoclonal gammopathy of undetermined (or unknown or unclear) significance (MGUS), plasmacytoma (bone, extramedullary), lymphoplasmacytic lymphoma (LPL), Waldenstrom's Macroglobulinemia, plasma cell leukemia, and primary amyloidosis (AL). Hematopoietic cancers may also include other cancers of additional hematopoietic cells, including polymorphonuclear leukocytes (or neutrophils), basophils, eosinophils, dendritic cells, platelets, erythrocytes and natural killer cells. Tissues which include hematopoietic cells referred herein to as "hematopoietic cell tissues" include bone marrow; peripheral blood; thymus; and peripheral lymphoid tissues, such as spleen, lymph nodes,

lymphoid tissues associated with mucosa (such as the gut-associated lymphoid tissues), tonsils, Peyer's patches and appendix, and lymphoid tissues associated with other mucosa, for example, the bronchial linings.

[0076] The term "antigen binding protein" as used herein refers to antibodies, antibody fragments and other protein constructs which are capable of binding to and neutralising human BCMA.

[0077] The terms Fv, Fc, Fd, Fab, or F(ab)2 are used with their standard meanings (see, e.g., Harlow et al., *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory, (1988)).

[0078] The term "antibody" is used herein in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies), polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies)

[0079] The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogenous antibodies i.e. the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific being directed against a single antigenic binding site. Furthermore, in contrast to polyclonal antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

[0080] A "chimeric antibody" refers to a type of engineered antibody in which a portion of the heavy and/ or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular donor antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (US Patent No. 4, 816,567 and Morrison et al. Proc. Natl. Acad. Sci. USA 81:6851-6855) (1984)).

[0081] A "humanised antibody" refers to a type of engineered antibody having its CDRs derived from a non-human donor immunoglobulin, the remaining immunoglobulin-derived parts of the molecule being derived from one (or more) human immunoglobulin(s). In addition, framework support residues may be altered to preserve binding affinity (see, e.g., Queen et al., Proc. Natl Acad Sci USA, 86:10029-10032 (1989), Hodgson et al., Bio/Technology, 9:421 (1991)). A suitable human acceptor antibody may be one selected from a conventional database, e.g., the KABAT® database, Los Alamos database, and Swiss Protein database, by homology to the nucleotide and amino acid sequences of the donor antibody. A human antibody characterized by a homology to the framework regions of the donor antibody (on an amino acid basis) may be suitable to provide a heavy chain constant region and/or a heavy chain variable framework region for insertion of the donor CDRs. A suitable acceptor antibody capable of donating light chain constant or variable framework regions may be selected in a similar manner. It should be noted that the acceptor antibody heavy and light chains are not required to originate from the same acceptor antibody. The prior art describes several ways of producing such humanised antibodies - see for example EP-A-0239400 and EP-A-054951.

[0082] For nucleic acids, the term "substantial identity" indicates that two nucleic acids, or designated sequences thereof, when optimally aligned and compared, are identical, with appropriate nucleotide insertions or deletions, in at least about 80% of the nucleotides, at least about 90% to about 95%, or at least about 98% to about 99.5% of the nucleotides. Alternatively, substantial identity exists when the segments will hybridize under selective hybridization conditions, to the complement of the strand. "Identity," means, for polynucleotides and polypeptides, as the case may be, the comparison calculated using an algorithm provided in (1) and (2) below:

1. (1) Identity for polynucleotides is calculated by multiplying the total number of nucleotides in a given sequence by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in said sequence, or:

$$nn \leq xn - (xn \cdot y),$$

wherein nn is the number of nucleotide alterations, xn is the total number of nucleotides in a given sequence, y is 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and • is the symbol for the multiplication operator, and wherein any non-integer product of xn and y is rounded down to the nearest integer prior to subtracting it from xn. Alterations of a polynucleotide sequence encoding a polypeptide may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

2. (2) Identity for polypeptides is calculated by multiplying the total number of amino acids by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids, or:

$$na \leq xa - (xa \cdot y),$$

wherein na is the number of amino acid alterations, xa is the total number of amino acids in the sequence, y is 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and • is the symbol for the multiplication operator, and wherein any non-integer product of xa and y is rounded down to the nearest integer prior to subtracting it from xa

[0083] For nucleotide and amino acid sequences, the term "identical" indicates the degree of identity between two nucleic acid or amino acid sequences when optimally aligned and compared with appropriate insertions or deletions.

[0084] "Isolated" means altered "by the hand of man" from its natural state, has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", including but not limited to when such polynucleotide or polypeptide is introduced back into a cell, even if the cell is of the same species or type as that from which the polynucleotide or polypeptide was separated.

[0085] Throughout the present specification and the accompanying claims the term "comprising" and "comprises" incorporates "consisting of" and "consists of". That is, these words are intended to convey the possible inclusion of other elements or integers not specifically recited, where the context allows.

[0086] The term "specifically binds" as used throughout the present specification in relation to antigen binding proteins of the invention means that the antigen binding protein binds human BCMA (hBCMA) with no or insignificant binding to other human proteins. The term however does not exclude the fact that antigen binding proteins of the invention may also be cross-reactive with other forms of BCMA, for example primate BCMA. For example in one embodiment the antigen binding protein does not bind to TAC1 or BAFF-R.

[0087] The term "inhibits" as used throughout the present specification in relation to antigen binding proteins of the invention means that the biological activity of BCMA is reduced in the presence of the antigen binding proteins of the present invention in comparison to the activity of BCMA in the absence of such antigen binding proteins. Inhibition may be due but not limited to one or more of blocking ligand binding, preventing the ligand activating the receptor, and/ or down regulating the BCMA. Inhibits can also refer to an antigen binding protein binding to BCMA and causing cell apoptosis or ADCC. The antibodies of the invention may neutralise the activity of the BCMA ligands BAFF and/or APRIL binding to BCMA. Levels of neutralisation can be measured in several ways, for example by use of the assays as set out in the examples below, for example in 4.4 in an H929 cell NFkB signalling assay. The BCMA ligands BAFF and APRIL are able to induce NFkB signalling and downstream events following binding to BCMA. The neutralisation of BCMA in this assay is measured by assessing the ability of anti-BCMA monoclonal antibodies to inhibit BAFF or APRIL driven NFkB induction.

[0088] If an antibody or antigen binding fragment thereof is capable of neutralisation then this is indicative of inhibition of the interaction between human BAFF or APRIL and BCMA. Antibodies which are considered to have neutralising activity against human BCMA would have an IC50 of less than 30 micrograms/ml, or less than 20 micrograms/ml, or less than 10 micrograms/ml, or less than 5 micrograms/ml or less than 1 micrograms/ml or less than 0.1 micrograms/ml in the H929 stimulation assay as set out in Example 4.4

[0089] "CDRs" are defined as the complementarity determining region amino acid sequences of an antibody which are the hypervariable domains of immunoglobulin heavy and light chains. There are three heavy chain and three light chain CDRs (or CDR regions) in the variable portion of an immunoglobulin. Thus, "CDRs" as used herein may refer to all three heavy chain CDRs, or all three light chain CDRs (or both all heavy and all light chain CDRs, if appropriate).

[0090] CDRs provide the majority of contact residues for the binding of the antibody to the antigen or epitope. CDRs of interest in this invention are derived from donor antibody variable heavy and light chain sequences, and include analogs of the naturally occurring CDRs, which analogs also share or retain the same antigen binding specificity and/or neutralizing ability as the donor antibody from which they were derived.

[0091] The CDR sequences of antibodies can be determined by the Kabat numbering system (Kabat et al; (Sequences of proteins of Immunological Interest NIH, 1987), alternatively they can be determined using the Chothia numbering system (Al-Lazikani et al., (1997) JMB 273,927-948), the contact definition method (MacCallum R.M., and Martin A.C.R. and Thornton J.M, (1996), Journal of Molecular Biology, 262 (5), 732-745) or any other established method for numbering the residues in an antibody and determining CDRs known to the skilled man in the art

[0092] Other numbering conventions for CDR sequences available to a skilled person include "AbM" (University of Bath) and "contact" (University College London) methods. The minimum overlapping region using at least two of the Kabat, Chothia, AbM and contact methods can be determined to provide the "minimum binding unit". The minimum binding unit may be a sub-portion of a CDR.

[0093] Table A below represents one definition using each numbering convention for each CDR or binding unit. The Kabat numbering scheme is used in Table X to number the variable domain amino acid sequence. It should be noted that some of the CDR definitions may vary depending on the individual publication used.

Table A

	Kabat CDR	Chothia CDR	AbM CDR	Contact CDR	Minimum binding unit
H1	31-35/35A/35B	26-32/33/34	26-35/35A/35B	30-35/35A/35B	31-32
H2	50-65	52-56	50-58	47-58	52-56
H3	95-102	95-102	95-102	93-101	95-101
L1	24-34	24-34	24-34	30-36	30-34
L2	50-56	50-56	50-56	46-55	50-55
L3	89-97	89-97	89-97	89-96	89-96

[0094] Throughout this specification, amino acid residues in antibody sequences are numbered according to the Kabat scheme. Similarly, the terms "CDR", "CDRL1", "CDRL2", "CDRL3", "CDRH1", "CDRH2", "CDRH3" follow the Kabat numbering system as set forth in Kabat et al; Sequences of proteins of Immunological Interest NIH, 1987.

[0095] The terms "Variant" refers to at least one, two or three amino acid changes in the sequence. These amino acid changes may be deletion, substitution or addition but are preferably substitution. In one such embodiment the substitutions are conservative substitutions.

[0096] In an alternative embodiment the variant sequence contains at least one substitution whilst retaining the canonical of the antigen binding protein.

[0097] The complementarity determining regions (CDRs) L1, L2, L3, H1 and H2 tend to structurally exhibit one of a finite number of main chain conformations. The particular canonical structure class of a CDR is defined by both the length of the CDR and by the loop packing, determined by residues located at key positions in both the CDRs and the framework regions (structurally determining residues or SDRs). Martin and Thornton (1996; J Mol Biol 263:800-815) have generated an automatic method to define the "key residue" canonical templates. Cluster analysis is used to define the canonical classes for sets of CDRs, and canonical templates are then identified by analysing buried hydrophobics, hydrogen-bonding residues, and e.g. conserved glycines. The CDRs of antibody sequences can be assigned to canonical classes by comparing the sequences to the

key residue templates and scoring each template using identity or similarity matrices.

[0098] The terms "VH" and "VL" are used herein to refer to the heavy chain variable domain and light chain variable domain respectively of an antibody.

[0099] As used herein the term "domain" refers to a folded protein structure which has tertiary structure independent of the rest of the protein. Generally, domains are responsible for discrete functional properties of proteins and in many cases may be added, removed or transferred to other proteins without loss of function of the remainder of the protein and/or of the domain. An "antibody single variable domain" is a folded polypeptide domain comprising sequences characteristic of antibody variable domains. It therefore includes complete antibody variable domains and modified variable domains, for example, in which one or more loops have been replaced by sequences which are not characteristic of antibody variable domains, or antibody variable domains which have been truncated or comprise N- or C-terminal extensions, as well as folded fragments of variable domains which retain at least the binding activity and specificity of the full-length domain.

[0100] The phrase "immunoglobulin single variable domain" refers to an antibody variable domain (VH, VHH, VL) that specifically binds an antigen or epitope independently of a different V region or domain. An immunoglobulin single variable domain can be present in a format (e.g., homo- or hetero-multimer) with other, different variable regions or variable domains where the other regions or domains are not required for antigen binding by the single immunoglobulin variable domain (i.e., where the immunoglobulin single variable domain binds antigen independently of the additional variable domains). A "domain antibody" or "dAb" is the same as an "immunoglobulin single variable domain" which is capable of binding to an antigen as the term is used herein. An immunoglobulin single variable domain may be a human antibody variable domain, but also includes single antibody variable domains from other species such as rodent (for example, as disclosed in WO 00/29004), nurse shark and Camelid VHH dAbs. Camelid VHH are immunoglobulin single variable domain polypeptides that are derived from species including camel, llama, alpaca, dromedary, and guanaco, which produce heavy chain antibodies naturally devoid of light chains. Such VHH domains may be humanised according to standard techniques available in the art, and such domains are still considered to be "domain antibodies" according to the invention. As used herein "VH" includes camelid VHH domains. NARV are another type of immunoglobulin single variable domain which were identified in cartilaginous fish including the nurse shark. These domains are also known as Novel Antigen Receptor variable region (commonly abbreviated to V(NAR) or NARV). For further details see Mol. Immunol. 44, 656-665 (2006) and US20050043519A.

[0101] The term "Epitope-binding domain" refers to a domain that specifically binds an antigen or epitope independently of a different V region or domain, this may be a domain antibody (dAb), for example a human, camelid or shark immunoglobulin single variable domain or it may be a domain which is a derivative of a scaffold selected from the group consisting of CTLA-4 (E vibody); lipocalin; Protein A derived molecules such as Z-domain of Protein A (Affibody, SpA), A-domain (Avimer/Maxibody); Heat shock proteins such as GroEL and GroES; 30eroxidise30g (trans-body); ankyrin repeat protein (DARPin); peptide aptamer; C-type lectin domain (Tetranectin); human γ -crystallin and human ubiquitin (affilins); PDZ domains; scorpion toxinkunitz type domains of human protease inhibitors; and fibronectin (adnectin); which has been subjected to protein engineering in order to obtain binding to a ligand other than the natural ligand.

CTLA-4 (Cytotoxic T Lymphocyte-associated Antigen 4) is a CD28-family receptor expressed on mainly CD4+ T-cells. Its extracellular domain has a variable domain-like Ig fold. Loops corresponding to CDRs of antibodies can be substituted with heterologous sequence to confer different binding properties. CTLA-4 molecules engineered to have different binding specificities are also known as E vibodies. For further details see Journal of Immunological Methods 248 (1-2), 31-45 (2001)

Lipocalins are a family of extracellular proteins which transport small hydrophobic molecules such as steroids, bilins, retinoids and lipids. They have a rigid β -sheet secondary structure with a number of loops at the open end of the conical structure which can be engineered to bind to different target antigens. Anticalins are between 160-180 amino acids in size, and are derived from lipocalins. For further details see Biochim Biophys Acta 1482: 337-350 (2000), US7250297B1 and US20070224633

An affibody is a scaffold derived from Protein A of *Staphylococcus aureus* which can be engineered to bind to antigen. The domain consists of a three-helical bundle of approximately 58 amino acids. Libraries have been generated by randomisation of surface residues. For further details see Protein Eng. Des. Sel. 17, 455-462 (2004) and EP1641818A1

Avimers are multidomain proteins derived from the A-domain scaffold family. The native domains of approximately 35 amino acids adopt a defined disulphide bonded structure. Diversity is generated by shuffling of the natural variation exhibited by the family of A-domains. For further details see Nature Biotechnology 23(12), 1556 - 1561 (2005) and Expert Opinion on Investigational Drugs 16(6), 909-917 (June 2007)

A Transferrin is a monomeric serum transport glycoprotein. Transferrins can be engineered to bind different target antigens by insertion of peptide sequences in a permissive surface loop. Examples of engineered transferrins scaffolds include the Trans-body. For further details see J. Biol. Chem 274, 24066-24073 (1999).

[0102] Designed Ankyrin Repeat Proteins (DARPins) are derived from Ankyrin which is a family of proteins that mediate attachment of integral membrane proteins to the cytoskeleton. A single ankyrin repeat is a 33 residue motif consisting of two α -helices and a β -turn. They can be engineered to bind different target antigens by randomising residues in the first α -helix and a β -turn of each repeat. Their binding interface can be increased by increasing the number of modules (a method of affinity maturation). For further details see J. Mol. Biol. 332, 489-503 (2003), PNAS 100(4), 1700-1705 (2003) and J. Mol. Biol. 369, 1015-1028 (2007) and US20040132028A1.

[0103] Fibronectin is a scaffold which can be engineered to bind to antigen. Adnectins consists of a backbone of the natural amino acid sequence of the 10th domain of the 15 repeating units of human fibronectin type III (FN3). Three loops at one end of the β -sandwich can be engineered to enable an Adnectin to specifically recognize a therapeutic target of interest. For further details see Protein Eng. Des. Sel. 18, 435-444 (2005), US20080139791, WO2005056764 and US6818418B1.

[0104] Peptide aptamers are combinatorial recognition molecules that consist of a constant scaffold protein, typically thioredoxin (TrxA) which

contains a constrained variable peptide loop inserted at the active site. For further details see Expert Opin. Biol. Ther. 5, 783-797 (2005).

[0105] Microbodies are derived from naturally occurring microproteins of 25-50 amino acids in length which contain 3-4 cysteine bridges - examples of microproteins include KalataB1 and conotoxin and knottins. The microproteins have a loop which can be engineered to include upto 25 amino acids without affecting the overall fold of the microprotein. For further details of engineered knottin domains, see WO2008098796.

[0106] Other epitope binding domains include proteins which have been used as a scaffold to engineer different target antigen binding properties include human γ -crystallin and human ubiquitin (affilins), kunitz type domains of human protease inhibitors, PDZ-domains of the Ras-binding protein AF-6, scorpion toxins (charybdotoxin), C-type lectin domain (tetranectins) are reviewed in Chapter 7 - Non-Antibody Scaffolds from Handbook of Therapeutic Antibodies (2007, edited by Stefan Dubel) and Protein Science 15:14-27 (2006). Epitope binding domains of the present invention could be derived from any of these alternative protein domains.

[0107] As used herein, the term "antigen-binding site" refers to a site on a protein which is capable of specifically binding to antigen, this may be a single domain, for example an epitope-binding domain, or it may be paired VH/VL domains as can be found on a standard antibody. In some embodiments of the invention single-chain Fv (ScFv) domains can provide antigen-binding sites.

[0108] The terms "mAbdAb" and dAbmAb" are used herein to refer to antigen-binding proteins of the present invention. The two terms can be used interchangeably, and are intended to have the same meaning as used herein.

[0109] The term "antigen binding protein" as used herein refers to antibodies, antibody fragments for example a domain antibody (dAb), ScFv, Fab, Fab2, and other protein constructs. Antigen binding molecules may comprise at least one Ig variable domain, for example antibodies, domain antibodies (dAbs), Fab, Fab', F(ab')2, Fv, ScFv, diabodies, mAbdAbs, affibodies, heteroconjugate antibodies or bispecific antibodies. In one embodiment the antigen binding molecule is an antibody. In another embodiment the antigen binding molecule is a dAb, i.e. an immunoglobulin single variable domain such as a VH, VHH or VL that specifically binds an antigen or epitope independently of a different V region or domain. Antigen binding molecules may be capable of binding to two targets, i.e. they may be dual targeting proteins. Antigen binding molecules may be a combination of antibodies and antigen binding fragments such as for example, one or more domain antibodies and/or one or more ScFvs linked to a monoclonal antibody. Antigen binding molecules may also comprise a non-Ig domain for example a domain which is a derivative of a scaffold selected from the group consisting of CTLA-4 (Evibody); lipocalin; Protein A derived molecules such as Z-domain of Protein A (Affibody, SpA); A-domain (Avimer/Maxibody); Heat shock proteins such as GroEl and GroES; 32eroxidise32g (trans-body); ankyrin repeat protein (DARPin); peptide aptamer; C-type lectin domain (Tetranectin); human γ -crystallin and human ubiquitin (affilins); PDZ domains; scorpion toxin/kunitz type domains of human protease inhibitors; and fibronectin (adnectin); which has been subjected to protein engineering in order to obtain binding to OSM. As used herein "antigen binding protein" will be capable of antagonising and/or neutralising human OSM. In addition, an antigen binding protein may inhibit and/or block OSM activity by binding to OSM and preventing a natural ligand from binding and/or activating the gp130 receptor.

[0110] The term "Effector Function" as used herein is meant to refer to one or more of Antibody dependant cell mediated cytotoxic activity (ADCC) , Complement-dependant cytotoxic activity (CDC) mediated responses, Fc-mediated phagocytosis and antibody recycling via the FcRn receptor. For IgG antibodies, effector functionalities including ADCC and ADCP are mediated by the interaction of the heavy chain constant region with a family of Fc receptors present on the surface of immune cells. In humans these include Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16). Interaction between the antigen binding protein bound to antigen and the formation of the Fc/ Fc complex induces a range of effects including cytotoxicity, immune cell activation, phagocytosis and release of inflammatory cytokines.

[0111] The interaction between the constant region of an antigen binding protein and various Fc receptors (FcR) is believed to mediate the effector functions of the antigen binding protein. Significant biological effects can be a consequence of effector functionality, in particular, antibody-dependent cellular cytotoxicity (ADCC), fixation of complement (complement dependent cytotoxicity or CDC), and halflife/clearance of the antigen binding protein. Usually, the ability to mediate effector function requires binding of the antigen binding protein to an antigen and not all antigen binding proteins will mediate every effector function.

[0112] Effector function can be measured in a number of ways including for example via binding of the Fc γ RIII to Natural Killer cells or via Fc γ RI to monocytes/macrophages to measure for ADCC effector function. For example an antigen binding protein of the present invention can be assessed for ADCC effector function in a Natural Killer cell assay. Examples of such assays can be found in Shields et al, 2001 The Journal of Biological Chemistry, Vol. 276, p6591-6604; Chappel et al, 1993 The Journal of Biological Chemistry, Vol 268, p25124-25131; Lazar et al, 2006 PNAS, 103, 4005-4010.

[0113] Examples of assays to determine CDC function include that described in 1995 J Imm Meth 184:29-38.

[0114] Some isotypes of human constant regions, in particular IgG4 and IgG2 isotypes, essentially lack the functions of a) activation of complement by the classical pathway; and b) antibody-dependent cellular cytotoxicity. Various modifications to the heavy chain constant region of antigen binding proteins may be carried out depending on the desired effector property. IgG1 constant regions containing specific mutations have separately been described to reduce binding to Fc receptors and therefore reduce ADCC and CDC (Duncan et al. Nature 1988, 332; 563-564; Lund et al. J. Immunol. 1991, 147; 2657-2662; Chappel et al. PNAS 1991, 88; 9036-9040; Burton and Woof, Adv. Immunol. 1992, 51:1-84; Morgan et al., Immunology 1995, 86; 319-324; Hezareh et al., J. Virol. 2001, 75 (24); 12161-12168).

[0115] In one embodiment of the present invention there is provided an antigen binding protein comprising a constant region such that the antigen binding protein has reduced ADCC and/or complement activation or effector functionality. In one such embodiment the heavy chain constant region may comprise a naturally disabled constant region of IgG2 or IgG4 isotype or a mutated IgG1 constant region. Examples of suitable modifications are described in EP0307434. One example comprises the substitutions of alanine residues at positions 235 and 237 (EU index numbering).

[0116] Human IgG1 constant regions containing specific mutations or altered glycosylation on residue Asn297 have also been described to enhance binding to Fc receptors. In some cases these mutations have also been shown to enhance ADCC and CDC (Lazar et al. PNAS 2006, 103; 4005-

4010; Shields et al. J Biol Chem 2001, 276; 6591-6604; Nechansky et al. Mol Immunol, 2007, 44; 1815-1817).

[0117] In one embodiment of the present invention, such mutations are in one or more of positions selected from 239, 332 and 330 (IgG1), or the equivalent positions in other IgG isotypes. Examples of suitable mutations are S239D and I332E and A330L. In one embodiment the antigen binding protein of the invention herein described is mutated at positions 239 and 332, for example S239D and I332E or in a further embodiment it is mutated at three or more positions selected from 239 and 332 and 330, for example S239D and I332E and A330L. (EU index numbering).

[0118] In an alternative embodiment of the present invention, there is provided an antigen binding protein comprising a heavy chain constant region with an altered glycosylation profile such that the antigen binding protein has enhanced effector function. For example, wherein the antigen binding protein has enhanced ADCC or enhanced CDC or wherein it has both enhanced ADCC and CDC effector function. Examples of suitable methodologies to produce antigen binding proteins with an altered glycosylation profile are described in WO2003011878, WO2006014679 and EP1229125, all of which can be applied to the antigen binding proteins of the present invention.

[0119] The present invention also provides a method for the production of an antigen binding protein according to the invention comprising the steps of:

1. a) culturing a recombinant host cell comprising an expression vector comprising the isolated nucleic acid as described herein, wherein the FUT8 gene encoding alpha-1,6-fucosyltransferase has been inactivated in the recombinant host cell; and
2. b) recovering the antigen binding protein.

[0120] Such methods for the production of antigen binding proteins can be performed, for example, using the POTESSION™ technology system available from BioWa, Inc. (Princeton, NJ) in which CHOK1SV cells lacking a functional copy of the FUT8 gene produce monoclonal antibodies having enhanced antibody dependent cell mediated cytotoxicity (ADCC) activity that is increased relative to an identical monoclonal antibody produced in a cell with a functional FUT8 gene. Aspects of the POTESSION™ technology system are described in US7214775, US6946292, WO0061739 and WO0231240. Those of ordinary skill in the art will also recognize other appropriate systems.

[0121] In one embodiment of the present invention there is provided an antigen binding protein comprising a chimaeric heavy chain constant region for example an antigen binding protein comprising a chimaeric heavy chain constant region with at least one CH2 domain from IgG3 such that the antigen binding protein has enhanced effector function, for example wherein it has enhanced ADCC or enhanced CDC, or enhanced ADCC and CDC functions. In one such embodiment, the antigen binding protein may comprise one CH2 domain from IgG3 or both CH2 domains may be from IgG3.

[0122] Also provided is a method of producing an antigen binding protein according to the invention comprising the steps of:

1. a) culturing a recombinant host cell comprising an expression vector comprising an isolated nucleic acid as described herein wherein the expression vector comprises a nucleic acid sequence encoding an Fc domain having both IgG1 and IgG3 Fc domain amino acid residues; and
2. b) recovering the antigen binding protein.

[0123] Such methods for the production of antigen binding proteins can be performed, for example, using the COMPLEGENT™ technology system available from BioWa, Inc. (Princeton, NJ) and Kyowa Hakko Kogyo (now, Kyowa Hakko Kirin Co., Ltd.) Co., Ltd. In which a recombinant host cell comprising an expression vector in which a nucleic acid sequence encoding a chimeric Fc domain having both IgG1 and IgG3 Fc domain amino acid residues is expressed to produce an antigen binding protein having enhanced complement dependent cytotoxicity (CDC) activity that is increased relative to an otherwise identical antigen binding protein lacking such a chimeric Fc domain. Aspects of the COMPLEGENT™ technology system are described in WO2007011041 and US20070148165. In an alternative embodiment CDC activity may be increased by introducing sequence specific mutations into the Fc region of an IgG chain. Those of ordinary skill in the art will also recognize other appropriate systems.

[0124] It will be apparent to those skilled in the art that such modifications may not only be used alone but may be used in combination with each other in order to further enhance effector function.

[0125] In one such embodiment of the present invention there is provided an antigen binding protein comprising a heavy chain constant region which comprises a mutated and chimaeric heavy chain constant region for example wherein an antigen binding protein comprising at least one CH2 domain from IgG3 and one CH2 domain from IgG1, wherein the IgG1 CH2 domain has one or more mutations at positions selected from 239 and 332 and 330 (for example the mutations may be selected from S239D and I332E and A330L) such that the antigen binding protein has enhanced effector function, for example wherein it has one or more of the following functions, enhanced ADCC or enhanced CDC, for example wherein it has enhanced ADCC and enhanced CDC. In one embodiment the IgG1 CH2 domain has the mutations S239D and I332E.

[0126] In an alternative embodiment of the present invention there is provided an antigen binding protein comprising a chimaeric heavy chain constant region and which has an altered glycosylation profile. In one such embodiment the heavy chain constant region comprises at least one CH2 domain from IgG3 and one CH2 domain from IgG1 and has an altered glycosylation profile such that the ratio of fucose to mannose is 0.8:3 or less, for example wherein the antigen binding protein is defucosylated so that said antigen binding protein has an enhanced effector function in comparison with an equivalent antigen binding protein with an immunoglobulin heavy chain constant region lacking said mutations and altered glycosylation profile, for example wherein it has one or more of the following functions, enhanced ADCC or enhanced CDC, for example wherein it has enhanced ADCC and enhanced CDC. In an alternative embodiment the antigen binding protein has at least one IgG3 CH2 domain and at least one heavy chain constant domain from IgG1 wherein both IgG CH2 domains are mutated in accordance with the limitations described herein.

[0127] In one aspect of the invention there is provided a method of producing an antigen binding protein according to the invention described herein comprising the steps of:

1. a) culturing a recombinant host cell containing an expression vector containing an isolated nucleic acid as described herein, said expression vector further comprising a Fc nucleic acid sequence encoding a chimeric Fc domain having both IgG1 and IgG3 Fc domain amino acid residues, and wherein the FUT8 gene encoding alpha-1,6-fucosyltransferase has been inactivated in the recombinant host cell, and
2. b) recovering the antigen binding protein.

[0128] Such methods for the production of antigen binding proteins can be performed, for example, using the ACCRETAMAB™ technology system available from BioWa, Inc. (Princeton, NJ) which combines the POTELLIGENT™ and COMPLEGENT™ technology systems to produce an antigen binding protein having both ADCC and CDC enhanced activity that is increased relative to an otherwise identical monoclonal antibody lacking a chimeric Fc domain and which has fucose on the oligosaccharide

[0129] In yet another embodiment of the present invention there is provided an antigen binding protein comprising a mutated and chimeric heavy chain constant region wherein said antigen binding protein has an altered glycosylation profile such that the antigen binding protein has enhanced effector function, for example wherein it has one or more of the following functions, enhanced ADCC or enhanced CDC. In one embodiment the mutations are selected from positions 239 and 332 and 330, for example the mutations are selected from S239D and I332E and A330L. In a further embodiment the heavy chain constant region comprises at least one CH2 domain from IgG3 and one Ch2 domain from IgG1. In one embodiment the heavy chain constant region has an altered glycosylation profile such that the ratio of fucose to mannose is 0.8:3 or less for example the antigen binding protein is defucosylated, so that said antigen binding protein has an enhanced effector function in comparison with an equivalent non-chimeraic antigen binding protein or with an immunoglobulin heavy chain constant region lacking said mutations and altered glycosylation profile.

Immunoconjugates

[0130] Also provided is an immunoconjugate (interchangeably referred to as "antibody-drug conjugates," or "ADCs") comprising an antigen binding protein according to the invention as herein described including, but not limited to, an antibody conjugated to one or more cytotoxic agents, such as a chemotherapeutic agent, a drug, a growth inhibitory agent, a toxin (e.g., a protein toxin, an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0131] Immunoconjugates have been used for the local delivery of cytotoxic agents, i.e., drugs that kill or inhibit the growth or proliferation of cells, in the treatment of cancer (Lambert, J. (2005) *Curr. Opinion in Pharmacology* 5:543-549; Wu et al. (2005) *Nature Biotechnology* 23(9):1137-1146; Payne, G. (2003) i 3:207-212; Syrigos and Epenetos (1999) *Anticancer Research* 19:605-614; Niculescu-Duvaz and Springer (1997) *Adv. Drug Deliv. Rev.* 26:151-172; U.S. Pat. No. 4,975,278). Immunoconjugates allow for the targeted delivery of a drug moiety to a tumor, and intracellular accumulation therein, where systemic administration of unconjugated drugs may result in unacceptable levels of toxicity to normal cells as well as the tumor cells sought to be eliminated (Baldwin et al., *Lancet* (Mar. 15, 1986) pp. 603-05; Thorpe (1985) "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review," in *Monoclonal Antibodies '84: Biological And Clinical Applications* (A. Pinchera et al., eds) pp. 475-506. Both polyclonal antibodies and monoclonal antibodies have been reported as useful in these strategies (Rowland et al., (1986) *Cancer Immunol. Immunother.* 21:183-87). Drugs used in these methods include daunomycin, doxorubicin, methotrexate, and vindesine (Rowland et al., (1986) *supra*). Toxins used in antibody-toxin conjugates include bacterial toxins such as diphtheria toxin, plant toxins such as ricin, small molecule toxins such as geldanamycin (Mandler et al (2000) *J. Nat. Cancer Inst.* 92(19):1573-1581; Mandler et al (2000) *Bioorganic & Med. Chem. Letters* 10:1025-1028; Mandler et al (2002) *Bioconjugate Chem.* 13:786-791), maytansinoids (EP 1391213; Liu et al., (1996) *Proc. Natl. Acad. Sci. USA* 93:8618-8623), and calicheamicin (Lode et al (1998) *Cancer Res.* 58:2928; Hinman et al (1993) *Cancer Res.* 53:3336-3342).

[0132] In one embodiment, the present invention includes immunoconjugates having the following general structure:

$$\text{ABP} - (\text{Linker})_n - \text{Ctx})_m$$

Wherein ABP is an antigen binding protein



Linker is either absent or any a cleavable or non-cleavable linker described herein


Ctx is any cytotoxic agent described herein

n is 0, 1, 2, or 3 and

m is 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10.

[0133] Examples of antibodies linked by an MC linker with auristatins such as MMAE and MMAF are depicted in the following structures:





[0134] In certain embodiments, an immunoconjugate comprises an antigen binding protein, including but not limited to, an antibody and a chemotherapeutic agent or other toxin. Chemotherapeutic agents useful in the generation of immunoconjugates are described herein. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the trichothecenes. See, e.g., WO 93/21232 published Oct. 28, 1993. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{211}At , ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re .

[0135] Antigen binding proteins of the present invention may also be conjugated to one or more toxins, including, but not limited to, a calicheamicin, maytansinoids, dolastatins, aurostatins, a trichothecene, and CC1065, and the derivatives of these toxins that have toxin activity. Suitable cytotoxic agents include, but are not limited to, an auristatin including dovaline-valine-dolaisoleanine-dolaproinephenylalanine (MMAF) and monomethyl auristatin E (MMAE) as well as ester forms of MMAE, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a duocarmycin, a taxane, including paclitaxel and docetaxel, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid. Specific cytotoxic agents include topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, calicheamicin, maytansine, DM-1, DM-4, netropsin. Other suitable cytotoxic agents include anti-tubulin agents, such as an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin derivative, a cryptophycin, a maytansinoid, a combretastatin, or a dolastatin. Antitubulin agent include dimethylvaline-valine-dolaisoleuine-dolaproine-phenylalanine-p-phenylenediamine (AFP), MMAF, MMAE, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcimid, estramustine, cemadotin, discodermolide, maytansine, DM-1, DM-4 or eleutherobin.

[0136] Antibody drug conjugates were produced by conjugating the small molecule anti-tubulin agent monomethylauristatin E (MMAE) or monomethylauristatin F (MMAF) to the antibodies. In the case of MMAE the linker consists of a thiol-reactive maleimide, a caproyl spacer, the dipeptide valine-citrulline, and p-aminobenzoyloxycarbonyl, a self-immolative fragmenting group. In the case of MMAF a protease-resistant maleimidocaproyl linker is used. The conjugation process leads to heterogeneity in drug-antibody attachment, varying in both the number of drugs bound to each antibody molecule (mole ratio [MR]), and the site of attachment. The most prevalent species is the material with an MR = 4; less prevalent are materials with MR of 0, 2, 6, and 8. The overall average drug-to-antibody MR is approximately 4.

Production of Immunoconjugates

[0137] The points of attachment are cysteines produced by mild reduction of the interchain disulfides of the antibody which is carried out whilst antibodies are immobilised on Protein G affinity resin (thus enabling the use of large reagent excesses without intermediate purifications). While immobilized, a large excess of TCEP will fully reduce the interchain disulfides but has no impact upon the binding of the antibody to the resin.

[0138] The number of thiols per antibody generated by this procedure depends upon the source and isotype of the antibodies. For example, human (and mouse-human chimeric) IgG1s have 4 reducible disulfides, and thus generate 8 thiols upon full reduction, whereas murine IgG1s have 5 reducible disulfides and produce 10 thiols. If ADCs with the maximal drug loading (e.g., 10 drugs per antibody for the murine IgG1s) are desired, then the maleimido-drug-linker can simply be added to the immobilized antibodies in sufficient excess to ensure complete conjugation. However, ADCs with fewer drugs per antibody can also be prepared from fully reduced antibodies by including a biologically inert capping agent such as N-ethyl maleimide (NEM) which occupies some of the available thiols on the antibody. When the maleimido-drug-linker and the capping agent are added simultaneously to the fully reduced antibody and in large excess (at least 3-fold), the two maleimide electrophiles compete for the limiting number of available thiols. In this fashion, the drug loading is determined by the relative thiol reaction rates of the drug-linker and capping agent, and thus can be considered to be under kinetic control. The relative reaction rates of maleimido-drug-linkers do vary significantly, and thus the molar ratio of drug-linker to NEM present in a reaction mix must be determined empirically to arrive at a panel of ADCs with a desired level of drug loading. The mole fraction of the drug linkers SGD-1006 (vcMMAE) and SGD-1269 (mcMMAF) in NEM mixtures which yield ADCs with approximately 4 drugs per antibody are summarized in Table 2 for common human and murine IgG isotypes.

Auristatins and Dolastatins


[0139] In some embodiments, the immunoconjugate comprises an antigen binding protein or antibody conjugated to dolastatins or dolostatin peptidic analogs and derivatives, the auristatins (U.S. Pat. Nos. 5,635,483; 5,780,588). Dolastatins and auristatins have been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke et al. (2001) *Antimicrob. Agents and Chemother.* 45(12):3580-3584) and have anticancer (U.S. Pat. No. 5,663,149) and antifungal activity (Pettit et al. (1998) *Antimicrob. Agents Chemother.* 42:2961-2965). The dolastatin or auristatin (which are pentapeptide derivatives of dolastatins) drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl) terminus of the peptidic drug moiety (WO 02/088172).

[0140] Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, disclosed in "Monomethylvaline Compounds Capable of Conjugation to Ligands." U.S. Patent No. 7,498,298. As used herein, the abbreviation "MMAE" refers to monomethyl auristatin E. As used herein the abbreviation "MMAF" refers to dovaline-valine-dolaisoleuine-dolaproine-phenylalanine.


[0141] Typically, peptide-based drug moieties can be prepared by forming a peptide bond between two or more amino acids and/or peptide fragments. Such peptide bonds can be prepared, for example, according to the liquid phase synthesis method (see E. Schroder and K. Lubke, "The

Peptides," volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry. The auristatin/dolastatin drug moieties may be prepared according to the methods of: U.S. Pat. No. 5,635,483; U.S. Pat. No. 5,780,588; Pettit et al. (1989) J. Am. Chem. Soc. 111:5463-5465; Pettit et al. (1998) Anti-Cancer Drug Design 13:243-277; Pettit, G. R., et al. Synthesis, 1996, 719-725; and Pettit et al. (1996) J. Chem. Soc. Perkin Trans. 15:859-863. See also Doronina (2003) Nat Biotechnol 21(7):778-784; "Monomethylvaline Compounds Capable of Conjugation to Ligands," U.S. Patent No. 7,498,298, filed Nov. 5, 2004, (disclosing, e.g., linkers and methods of preparing monomethylvaline compounds such as MMAE and MMAF conjugated to linkers). Biologically active organic compounds which act as cytotoxic agents, specifically pentapeptides, are disclosed in US Patent Nos. 6,884,869; 7,498,298; 7,098,308; 7,256,257; and 7,423,116.. Monoclonal antibodies linked with MMAE and MMAF as well as various derivatives of auristatins and methods of making them are described in US Patent NO. 7,964,566.

[0142] Examples of auristatins include MMAE and MMAF the structures of which are shown below:



MMAE



Maytansine and Maytansinoids

[0143] Maytansinoids are mitotic inhibitors which act by inhibiting tubulin polymerization. Maytansine was first isolated from the east African shrub Maytenus serra (U.S. Pat. No. 3,896,111). Subsequently, it was discovered that certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (U.S. Pat. No. 4,151,042). Highly cytotoxic maytansinoid drugs can be prepared from ansamitocin precursors produced by fermentation of microorganisms such as *Actinosynema*. Methods for isolating ansamitocins are described in US Patent No. 6,573,074. Synthetic maytansinol and derivatives and analogues thereof are disclosed, for example, in U.S. Pat. Nos. 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,364,866; 4,424,219; 4,450,254; 4,362,663; and 4,371,533.

[0144] Antibody-maytansinoid conjugates are prepared by chemically linking an antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. See, e.g., U.S. Pat. No. 5,208,020. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, although even one molecule of toxin/antibody would be expected to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources. Suitable maytansinoids are disclosed, for example, in U.S. Pat. No. 5,208,020 and in the other patents and nonpatent publications referred to hereinabove. Maytansinoids are maytansinol and maytansinol analogues modified in the aromatic ring or at other positions of the maytansinol molecule, such as various maytansinol esters. Methods for preparing maytansinoids for linkage with antibodies are disclosed in US Patent Nos. 6,570,024 and 6,884,874.

Calicheamicin

[0145] The calicheamicin family of antibiotics is capable of producing double-stranded DNA breaks at subpicomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Pat. Nos. 5,712,374, 5,714,586, 5,739,116, 5,767,285, 5,770,701, 5,770,710, 5,773,001, 5,877,296 (all to American Cyanamid Company). Structural analogues of calicheamicin which may be used include, but are not limited to, .gamma.11, .alpha.21, .alpha.31, N-acetyl-.gamma.11, PSAG and .theta.11 (Hinman et al., Cancer Research 53:3336-3342 (1993), Lode et al., Cancer Research 58:2925-2928 (1998) and the aforementioned U.S. patents to American Cyanamid). Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

Other Cytotoxic Agents

[0146] Other antitumor agents that can be conjugated to the antibodies include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Pat. Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Pat. No. 5,877,296).

[0147] Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the trichothecenes. See, for example, WO 93/21232 published Oct. 28, 1993.

[0148] The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase).

[0149] For selective destruction of the tumor, the antibody may comprise a highly radioactive atom. A variety of radioactive isotopes are available for the production of radioconjugated antibodies. Examples include At211, I131, I125, Y90, Re186, Re188, Sm153, Bi212, P32, Pb212 and radioactive isotopes of Lu. When the conjugate is used for detection, it may comprise a radioactive atom for scintigraphic studies, for example tc99m or I123, or a spin label for nuclear magnetic resonance (NMR) imaging (also known as magnetic resonance imaging, mri), such as iodine-123 again, iodine-131, indium-111, fluorine-19, carbon-13, nitrogen-15, oxygen-17, gadolinium, manganese or iron.

[0150] The radio- or other labels may be incorporated in the conjugate in known ways. For example, the peptide may be biosynthesized or may be synthesized by chemical amino acid synthesis using suitable amino acid precursors involving, for example, fluorine-19 in place of hydrogen. Labels such as tc99m or I123, Re186, Re188 and In111 can be attached via a cysteine residue in the peptide. Yttrium-90 can be attached via a lysine residue. The IODOGEN method (Fraker et al. (1978) *Biochem. Biophys. Res. Commun.* 80: 49-57) can be used to incorporate iodine-123. "Monoclonal Antibodies in Immunoscintigraphy" (Chatal, CRC Press 1989) describes other methods in detail.

Preparation of ADCs

[0151] In antibody drug conjugates, the antibody can be conjugated directly to the cytotoxic agent or via a linker. Suitable linkers include, for example, cleavable and non-cleavable linkers. A cleavable linker is typically susceptible to cleavage under intracellular conditions. Suitable cleavable linkers include, for example, a peptide linker cleavable by an intracellular protease, such as lysosomal protease or an endosomal protease. In exemplary embodiments, the linker can be a dipeptide linker, such as a valine-citrulline (val-cit) or a phenylalanine-lysine (phe-lys) linker. Other suitable linkers include linkers hydrolyzable at a pH of less than 5.5, such as a hydrazone linker. Additional suitable cleavable linkers include disulfide linkers.

[0152] Bristol-Myers Squibb has described particular lysosomal enzyme-cleavable antitumor drug conjugates. See, for example, U.S. Pat. No. 6,214,345. Seattle Genetics has published applications U.S. Pat. Appl. 2003/0096743 and U.S. Pat. Appl. 2003/0130189, which describe p-aminobenzylethers in drug delivery agents. The linkers described in these applications are limited to aminobenzyl ether compositions.

[0153] Conjugates of the antigen binding protein and cytotoxic agent may be made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridylidithio)propionate (SPDP), succinimidyl-4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCl), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis(p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene).

[0154] Additionally the linker may be composed of one or more linker components. Exemplary linker components include 6-maleimidocaproyl ("MC"), maleimidopropanoyl ("MP"), valine-citrulline ("val-cit"), alanine-phenylalanine ("ala-phe"), p-aminobenzylloxycarbonyl ("PAB"), N-Succinimidyl 4-(2-pyridylthio)pentanoate ("SPP"), N-Succinimidyl 4-(N-maleimidomethyl)cyclohexane-1 carboxylate ("SMCC"), and N-Succinimidyl (4-iodoacetyl)aminobenzoate ("SIAB"). Additional linker components are known in the art and some are described herein. See also "Monomethylvaline Compounds Capable of Conjugation to Ligands," U.S. Patent No. US7,498,298, filed Nov. 5, 2004.

[0155] Linkers may also comprise amino acids and/or amino acid analogs. Amino acid linker components include a dipeptide, a tripeptide, a tetrapeptide or a pentapeptide. Exemplary dipeptides include: valine-citrulline (vc or val-cit), alanine-phenylalanine (af or ala-phe). Exemplary tripeptides include: glycine-valine-citrulline (gly-val-cit) and glycine-glycine-glycine (gly-gly-gly). Amino acid residues which comprise an amino acid linker component include those occurring naturally, as well as minor amino acids and non-naturally occurring amino acid analogs, such as citrulline. Amino acid linker components can be designed and optimized in their selectivity for enzymatic cleavage by a particular enzyme, for example, a tumor-associated protease, cathepsin B, C and D, or a plasmin protease.

[0156] Antigen binding proteins and antibodies may be made reactive for conjugation with linker reagents. Nucleophilic groups on antibodies include, but are not limited to: (i) N-terminal amine groups, (ii) side chain amine groups, e.g., lysine, (iii) side chain thiol groups, e.g. cysteine, and (iv) sugar hydroxyl or amino groups where the antibody is glycosylated. Amine, thiol, and hydroxyl groups are nucleophilic and capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) active esters such as NHS esters, HOBt esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups. Certain antibodies have reducible interchain disulfides, i.e. cysteine bridges. Antibodies may be made reactive for conjugation with linker reagents by treatment with a reducing agent such as DTT (dithiothreitol). Each cysteine bridge will thus form, theoretically, two reactive thiol nucleophiles. Additional nucleophilic groups can be introduced into antibodies through the reaction of lysines with 2-iminothiolane (Traut's reagent) resulting in conversion of an amine into a thiol. Reactive thiol groups may be introduced into the antibody (or fragment thereof) by introducing one, two, three, four, or more cysteine residues (e.g., preparing mutant antibodies comprising one or more non-native cysteine amino acid residues).

[0157] Antigen binding proteins and antibodies may also be modified to introduce electrophilic moieties, which can react with nucleophilic substituents on the linker reagent or drug. The sugars of glycosylated antibodies may be oxidized, e.g. with periodate oxidizing reagents, to form aldehyde or ketone groups which may react with the amine group of linker reagents or drug moieties. The resulting imine Schiff base groups may form a stable linkage, or may be reduced, e.g., by borohydride reagents to form stable amine linkages. In one embodiment, reaction of the carbohydrate portion of a glycosylated antibody with either glucose oxidase or sodium meta-periodate may yield carbonyl (aldehyde and ketone) groups in the protein that can react with appropriate groups on the drug (Hermanson, *Bioconjugate Techniques*). In another embodiment, proteins containing N-terminal serine or threonine residues can react with sodium meta-periodate, resulting in production of an aldehyde in place of the first amino acid (Geoghegan & Stroh, (1992) *Bioconjugate Chem.* 3:138-146; U.S. Pat. No. 5,362,852). Such aldehydes can be reacted with a drug moiety or linker nucleophile.

[0158] Nucleophilic groups on a drug moiety include, but are not limited to: amine, thiol, hydroxyl, hydrazide, oxime, hydrazine, thiosemicarbazone, hydrazine carboxylate, and arylhydrazide groups capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker

reagents including: (i) active esters such as NHS esters, HOBT esters, haloformates, and acid halides; (ii) alkyl and benzyl halides such as haloacetamides; (iii) aldehydes, ketones, carboxyl, and maleimide groups.

[0159] In some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (e.g., within a lysosome or endosome or caveolea). The linker can be, e.g., a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. Typically, the peptidyl linker is at least two amino acids long or at least three amino acids long. Cleaving agents can include cathepsins B and D and plasmin, all of which are known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells (see, e.g., Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123). Peptidyl linkers may be cleavable by enzymes that are present cells. For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a Phe-Leu or a Gly-Phe-Leu-Gly (SEQ ID NO:50) linker). Other such linkers are described, e.g., in U.S. Pat. No. 6,214,345. In specific embodiments, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker or a Phe-Lys linker (see, e.g., U.S. Pat. No. 6,214,345, which describes the synthesis of doxorubicin with the val-cit linker). One advantage of using intracellular proteolytic release of the therapeutic agent is that the agent is typically attenuated when conjugated and the serum stabilities of the conjugates are typically high.

[0160] In other embodiments, the cleavable linker is pH-sensitive, i.e., sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acidlabile linker that is hydrolyzable in the lysosome (e.g., a hydrazone, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like) can be used. (See, e.g., U.S. Pat. Nos. 5,122,368; 5,824,805; 5,622,929; Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123; Neville et al., 1989, *Biol. Chem.* 264:14653-14661.) Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome. In certain embodiments, the hydrolyzable linker is a thioether linker (such as, e.g., a thioether attached to the therapeutic agent via an acylhydrazone bond (see, e.g., U.S. Pat. No. 5,622,929)).

[0161] In yet other embodiments, the linker is cleavable under reducing conditions (e.g., a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-5-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridylthio)propionate), SPDB (N-succinimidyl-3-(2-pyridylthio)butyrate) and SMPT (N-succinimidyl-oxy carbonyl-alphamethyl-alpha-(2-pyridylthio)toluene)-, SPDB and SMPT (See, e.g., Thorpe et al., 1987, *Cancer Res.* 47:5924-5931; Wawrzynczak et al., In *Immunoc conjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer* (C. W. Vogel ed., Oxford U. Press, 1987. See also U.S. Pat. No. 4,880,935.)

[0162] In yet other specific embodiments, the linker is a malonate linker (Johnson et al., 1995, *Anticancer Res.* 15:1387-93), a maleimidobenzoyl linker (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10):1299-1304), or a 3'-N-amide analog (Lau et al., 1995, *Bioorg-Med-Chem.* 3(10):1305-12).

[0163] Typically, the linker is not substantially sensitive to the extracellular environment. As used herein, "not substantially sensitive to the extracellular environment," in the context of a linker, means that no more than about 20%, typically no more than about 15%, more typically no more than about 10%, and even more typically no more than about 5%, no more than about 3%, or no more than about 1% of the linkers, in a sample of ADC or ADC derivative, are cleaved when the ADC or ADC derivative present in an extracellular environment (e.g., in plasma). Whether a linker is not substantially sensitive to the extracellular environment can be determined, for example, by incubating independently with plasma both (a) the ADC or ADC derivative (the "ADC sample") and (b) an equal molar amount of unconjugated antibody or therapeutic agent (the "control sample") for a predetermined time period (e.g., 2, 4, 8, 16, or 24 hours) and then comparing the amount of unconjugated antibody or therapeutic agent present in the ADC sample with that present in control sample, as measured, for example, by high performance liquid chromatography.

[0164] In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (i.e., in the milieu of the linker-therapeutic agent moiety of the ADC or ADC derivative as described herein). In yet other embodiments, the linker promotes cellular internalization when conjugated to both the therapeutic agent and the antigen binding protein or antibody or derivative thereof (i.e., in the milieu of the ADC or ADC derivative as described herein).

[0165] A variety of linkers that can be used with the present compositions and methods are described in WO 2004010957 entitled "Drug Conjugates and Their Use for Treating Cancer, An Autoimmune Disease or an Infectious Disease" filed Jul. 31, 2003, and U.S. Provisional Application No. 60/400,403, entitled "Drug Conjugates and their use for treating cancer, an autoimmune disease or an infectious disease", filed Jul. 31, 2002.

[0166] Alternatively, a fusion protein comprising the antigen binding protein and cytotoxic agent may be made, e.g., by recombinant techniques or peptide synthesis. The length of DNA may comprise respective regions encoding the two portions of the conjugate either adjacent one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

[0167] In yet another embodiment, the antibody may be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pre-targeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionucleotide).

[0168] The term "Non Human antibody or antibody fragment thereof" as used herein is meant to refer to antibodies or fragments thereof which originate from any species other than human wherein human includes chimeric antibodies.

[0169] The term "donor antibody" refers to an antibody (monoclonal, and/or recombinant) which contributes the amino acid sequences of its variable domains, CDRs, or other functional fragments or analogs thereof to a first immunoglobulin partner, so as to provide the altered immunoglobulin coding region and resulting expressed altered antibody with the antigenic specificity and neutralizing activity characteristic of the donor antibody.

[0170] The term "acceptor antibody" refers to an antibody (monoclonal and/or recombinant) heterologous to the donor antibody, which contributes all (or any portion, but preferably all) of the amino acid sequences encoding its heavy and/or light chain framework regions and/or its heavy and/or light chain constant regions to the first immunoglobulin partner. The human antibody is the acceptor antibody. The term "Human acceptor sequence" as used herein is meant to refer to a framework of an antibody or antibody fragment thereof comprising the amino acid sequence of a VH or VL

framework derived from a human antibody or antibody fragment thereof or a human consensus sequence framework into which CDR's from a non-human species may be incorporated.

[0171] The term "incorporation" of CDR's or hypervariable regions as used herein encompasses any means by which the non-human CDR's are situated with the human acceptor framework. It will be appreciated that this can be achieved in various ways, for example, nucleic acids encoding the desired amino acid sequence can be generated by mutating nucleic acids encoding the non-human variable domain sequence so that the framework residues thereof are changed to human acceptor framework residues, or by mutating nucleic acid encoding the human variable domain sequence so that the CDR's are changed to non-human residues, or by synthesizing nucleic acids encoding the desired sequence. In one embodiment the final sequence is generated in silico.

[0172] The present invention is illustrated by the following examples.

Examples

Example 1 Monoclonal Antibody Generation and Selection

1.1 Immunisation strategies

[0173] The anti human BCMA mAb murine parental CA8 was identified from hybridomas derived from mice immunized with full length human BCMA. A BALB/c mouse was immunized i.p. with 25 µg of recombinant (rBCMA) protein combined with CFA. The mouse was boosted three times at one-month intervals with 25 µg of full length rBCMA protein + 10 µg monophosphoryl lipid A-stable emulsion (MPL-SE) (Corixa Corporation, Seattle, WA) and given a pre-fusion boost of 30 µg rBCMA protein i.v. 3 days prior to fusion. Hybridomas were either generated and cloned using the ClonaCell-HY hybridoma cloning kit (StemCell Technologies, Vancouver, BC) or using a conventional method. In the conventional method, B cells from the spleens of the immunized animals were fused with Sp2/0 myeloma cells in the presence of PEG (Sigma-Aldrich, St. Louis, MO). After overnight recovery, fused cells were plated at limiting dilution in 96-well plates and subjected to hypoxanthine-aminopterinthymidine selection. Hybridoma culture supernatants were examined for the presence of anti-BCMA antibodies by ELISA and flow cytometry.

[0174] The anti human BCMA mAb murine parental S307118G03 was identified from hybridomas derived from SJL mice immunized with recombinant human BCMA/TNFRSF17-Fc chimera (R&D 193-Fc) using the RIMMS method (Rapid immunisation multiple sites). At Day 0, 5µg protein per mouse was emulsified in AS02a adjuvant at 2 sites on back (over haunches and over shoulders) and subjacent to the major lymph nodes at 4 sites on front. On day 6 and day 11 2.5µg protein per mouse in RIBI adjuvant was injected subjacent to the major lymph nodes at 4 sites on front. On day 14 the animals were sacrificed. The lymph nodes and spleen were excised, disrupted and a PEG1500 induced somatic cell fusion performed using a 3:1 ratio with mouse myeloma cells X63 AG8 653.GFP.Bcl-2.11 (BioCat 112754; R17209/58). The fusion was plated out into 10 × 96 well plates and screened directly from these.

[0175] The anti human BCMA mAb murine parental S336105A07 was identified from hybridomas derived from identical immunisations. The lymph nodes and spleen were excised at day 14, disrupted, and a Cytopulse electrofusion was performed using a 1:1 ratio with mouse myeloma cells X63 AG8 653.GFP.Bcl-2.11 (BioCat 112754; R17209/58). The fusion was plated out into omnitrays containing semi solid medium prior to picking into 10 × 96 well plates and was screened directly from these 5 days later.

[0176] The anti human BCMA murine parental mAbs S332121F02 and S332126E04 were identified from hybridomas derived from SJL mice immunized with recombinant Fc fusion of the extracellular domain of human BCMA (4-53)BCMA using the RIMMS method (Rapid immunisation). At Day 0, 5µg protein per mouse was emulsified in AS02a adjuvant at 2 sites on back (over haunches and over shoulders) and subjacent to the major lymph nodes at 4 sites on front. On day 6 5µg recombinant cyno BCMA-Fc protein per mouse in RIBI adjuvant was injected subjacent to the major lymph nodes at 4 sites on front. On day 11 2.5µg recombinant human BCMA-Fc and 2.5µg recombinant cyno BCMA-Fc per mouse in RIBI adjuvant was injected subjacent to the major lymph nodes at 4 sites on front. On day 14 the animals were sacrificed and cells treated as for S307118G03.

[0177] The anti human BCMA murine parental mAb S322110D07 was identified from hybridomas derived from SJL mice immunised with recombinant Fc fusion of the extracellular domain of human BCMA (4-53) in complex with recombinant human April (R&D 5860-AP/CF) premixed at 1:1 molar ratio. The mice were immunized i.p. with 5µg April/Cyno BCMA-Fc complex in PBS, suspended in RIBI adjuvant, 100ul dose per mouse and boosted 3 times at 3-4 week intervals with 2.5µg April/Cyno BCMA-Fc complex in PBS, suspended in RIBI adjuvant, 100ul dose per mouse injected via intraperitoneal route and given a pre-fusion boost of the same immunogen 1 day prior to fusion and treated as for S307118G03.

[0178] The anti human BCMA mAb murine parental S335115G01 and S335122F05 were identified from hybridomas derived from SJL mice immunized with a mixture of recombinant Fc fusion of the extracellular domain of human BCMA (4-53) and recombinant Fc fusion of the extracellular domain of cyno BCMA (4-52) using the RIMMS method (Rapid immunisation multiple sites). At Day 0, 2, 5µg of each protein per mouse was emulsified in AS02a adjuvant and injected at 2 sites on the back (over haunches and over shoulders) and subjacent to the major lymph nodes at 4 sites on front. On day 6 and day 11 2.5µg of each protein per mouse in RIBI adjuvant was injected subjacent to the major lymph nodes at 4 sites on front. On day 14 the animals were sacrificed. The lymph nodes and spleen were excised, disrupted and a Cytopulse electrofusion was performed using a 1:1 ratio with mouse myeloma cells X63 AG8 653.GFP.Bcl-2.11 (BioCat 112754; R17209/58). The fusion was plated out into omnitrays containing semi solid medium prior to picking into 32 × 96 well plates and was screened directly from these 5 days later.

Example 2 Humanisation.

2.1 Cloning of CA8 Hybridoma Variable Regions

[0179] Total RNA was extracted from CA8 hybridoma cells, heavy and light variable domain cDNA sequence was then generated by reverse transcription and polymerase chain reaction (RT-PCR). The forward primer for RT-PCR was a mixture of degenerate primers specific for murine immunoglobulin gene leader-sequences and the reverse primer was specific for the antibody constant regions. Reverse primers specific for IgG1, IgG2a and IgG2b were used in this case as the isotype was unknown. To design the primers, DNA multiple sequence alignments of the leader sequences of the mouse V_H and V_K genes were generated.

2.2 Cloning of chimeric CA8

[0180] The DNA expression constructs encoding the chimeric antibody were prepared *de novo* by build-up of overlapping oligonucleotides including restriction sites for cloning into mammalian expression vectors as well as a human signal sequence. *HindIII* and *SpeI* restriction sites were introduced to frame the VH domain containing the signal sequence for cloning into mammalian expression vectors containing the human $\gamma 1$ constant region. *HindIII* and *BsIWI* restriction sites were introduced to frame the VL domain containing the signal sequence for cloning into mammalian expression vector containing the human kappa constant region.

2.3 Cloning of the humanised CA8 variants

[0181] The DNA expression constructs encoding the humanised antibody variants were prepared *de novo* by build-up of overlapping oligonucleotides including restriction sites for cloning into mammalian expression vectors as well as a human signal sequence. *HindIII* and *SpeI* restriction sites were introduced to frame the VH domain containing the signal sequence for cloning into mammalian expression vectors containing the human $\gamma 1$ constant region. *HindIII* and *BsI* restriction sites were introduced to frame the VL domain containing the signal sequence for cloning into mammalian expression vector containing the human kappa constant region.

2.4 Expression of the recombinant CA8 antibodies (including antibody quantification)

[0182] Expression plasmids encoding the heavy and light chains respectively were transiently co-transfected into HEK 293 6E cells and expressed at small scale to produce antibody. Antibodies were quantified by ELISA. ELISA plates were coated with anti human IgG (Sigma I3382) at 1mg/ml and blocked with blocking solution (4% BSA in Tris buffered saline). Various dilutions of the tissue culture supernatants were added and the plate was incubated for 1 hour at room temperature. Dilutions of a known standard antibody were also added to the plate. The plate was washed in TBST and binding was detected by the addition of a peroxidase labelled anti human kappa light chain antibody (Sigma A7164) at a dilution of 1/1000 in blocking solution. The plate was incubated for 1 hour at room temp before washing in TBST. The plate was developed by addition of OPD substrate (Sigma P9187) and colour development stopped by addition of 2M H₂SO₄. Absorbance was measured at 490nm and a standard curve plotted using data for the known standard dilutions. The standard curve was used to estimate the concentration of antibody in the tissue culture supernatants. Larger scale antibody preparations were purified using protein A and concentrations were measured using a Nanodrop (Thermo Scientific).

Table 1. Design of CA8 variable heavy and light humanised variants

Humanised VH	Template	Backmutations (Kabat#)
J0	Straight graft of CA8 VH CDRs onto IGHV1_69 + JH1 minigene	None
J1	J0	G27Y. S30T
J2	J1	A93T
J3	J2	A24G. K73T
J4	J3	M4RI. V67A.
J5	J3	N99D
J6	J0	N99D
J7	J1	N99D
J8	J2	N99D
J9	J4	N99D
M0	Straight graft of CA8 VL CDRs onto IGKV1_39 + JK2 minigene	None
M1	M0	F71Y
M2	M1	M4L. K45E.

2.5 Defucosylated antibody production

[0183] To generate defucosylated antibodies the heavy and light chains respectively were co-transfected into CHO DG44 MS705 BioWa cells and expressed at scale to produce antibody. Briefly, 30 μ g DNA was linearised overnight with Not1, the DNA was ethanol precipitated and re-dissolved in TE buffer. From culture, 2.4X10⁷ BioWa DG44 cells were obtained and washed in 14ml of warmed PBS-sucrose. The cells were spun and the pellet resuspended in 1.6ml of PBS-sucrose. Half (0.8ml) of aforementioned cells, suspended in PBS-sucrose, were added to a BioRad cuvette with the

30 μ g of linearised DNA (in 50 μ l TE buffer). A BioRad GenePulser was programmed to 380V with a capacitance of 25 μ F and the cuvette was entered for electroporation. The resulting 850 μ l of electroporated cells and DNA were added to (80ml) warmed SFM512 medium (including phenol red, 2XHT (nucleosides), glutamax and Gibco supplement4). Finally, the resulting 80ml of cell suspension was transferred (150 μ l/well) to each well of one of 4 X 96-well plates. After 48 hours, the medium was changed to nucleoside free by removing approximately 130 μ l of conditioned and replacing with 150 μ l of fresh selection medium SFM512 medium (including phenol red and glutamax). Every 3-4 days, 130-150 μ l of conditioned medium was removed and replaced with fresh, selection medium. Wells were monitored for colour change and assayed for IgG concentration as discussed previously.

2.6 Additional antibodies - Cloning of Hybridoma Variable Regions

[0184] Total RNA was extracted from S307118G03, S332121F02, S332126E04, S322110D07, S336105A07, S335115G01 and S335122F05 hybridoma cells. Heavy and light variable domain cDNA sequence was then generated by reverse transcription and polymerase chain reaction (RT-PCR). The forward primer for RT-PCR was a mixture of degenerate primers specific for murine immunoglobulin gene leader-sequences and the reverse primer was specific for the antibody constant regions, in this case isotype IgG2a. Primers were designed based on a strategy described by Jones and Bendig (Bio/Technology 9:88, 1991). RT-PCR was carried out for both V-region sequences to enable subsequent verification of the correct V-region sequences. DNA sequence data was obtained for the V-region products generated by the RT-PCR.

2.7 Additional antibodies - Cloning of the chimeras

[0185] The DNA expression constructs encoding the chimeric antibodies were prepared de novo by infusion advantage PCR cloning (Clonetech) of the V-gene PCR products into mammalian expression vectors. This cloning method enabled fusion the murine variable regions to human IgG1 H chain and kappa L chain constant regions.

2.8 S307118G03 - Cloning of the humanized variants

[0186] Cloning was carried out as for paragraph 2.3.


2.9 S307118G03 Expression of the recombinant antibodies

[0187] Expression plasmids encoding the relevant heavy and light chains (listed in Table 8 below) were transiently co-transfected into HEK 293 6E cells and expressed at small scale to produce antibody. The antibodies were Protein A purified from the supernatants and quantified using the Nanodrop spectrophotometer. 8 below) were transiently co-transfected into HEK 293 6E cells and expressed at small scale to produce antibody. The antibodies were Protein A purified from the supernatants and quantified using the Nanodrop spectrophotometer.

Example 3 Conjugation of antibodies to vcMMAE and mcMMAF to form antibody drug conjugates (ADC)

[0188]

Table B Chemical structures of drug-linkers



[0189] Gammabind Plus Protein G Sepharose (GE Healthcare) resin slurry (75 μ L) was added to each well of a deep well (2 mL capacity) filter plate. The antibodies to be conjugated were grouped by species and isotype and up to 0.5 mg of each antibody transferred to each well of the plate. Each antibody was transferred to two separate wells to facilitate the preparation of two conjugates, with the drug-linkers SGD-1006 and SGD-1269.

The filter plate was then shaken at 1200 RPM for 2 hours at 5 °C to bind the antibodies to the resin. The filter plate was then centrifuged at 500x g for 3 minutes to ensure complete pulldown of all fluids and resin to the bottom of the each well.

[0190] The bound antibodies were then reduced by adding 500 μ L of 10 mM TCEP in 100 mM KPO₄, 150 mM NaCl, pH 7, 1 mM EDTA and shaking for 30 minutes at 22 °C. Following reduction, the plate was again centrifuged to remove the TCEP solution and subsequently washed with PBS + 1 mM EDTA, 1 mL per well. The wash solution was removed by centrifugation and the process repeated 3 times for a total of 4 washes. The bound and reduced antibodies were then conjugated using a mixture of NEM and drug linker prepared in accordance with the mole fractions indicated in Table 2.

Table 2.

Antibody (species / isotype)	Reducible Disulfides	SGD-1006 mole fraction	SGD-1269 mole fraction
Human IgG1*	4	0.675	0.688
Murine IgG1	5	0.500	0.586
Murine IgG2a	5	0.500	0.586
Murine IgG2b	6	0.463	0.481

* also for murine / human IgG1 chimerics

[0191] Separate mixtures of NEM and drug linker were thus prepared for each antibody species / isotype using 10 mM DMSO stock solutions of SGD-1006, SGD-1269 (See Table B) and NEM. When mixed at the appropriate ratio the total maleimide concentration was therefore still 10 mM, and this value was used to calculate the volume of maleimide solution to be added to each well. For example for a murine IgG1 with 5 reducible disulfides (10 available thiols when reduced) 0.5 mg of antibody at 150 kD is 3.33 nmol corresponding to 33.3 nmol of thiol. A 3-fold excess is therefore 100 nmol of total maleimide or 10 μ L of the 10 mM drug linker / NEM mix. For the SGD-1269 conjugate this mix would then be prepared with 5.86 μ L of SGD-1269 and 4.14 μ L of NEM. The maleimide mix would then be diluted into 500 μ L of PBS prior to addition to the immobilized reduced antibody. In practice, since multiple antibodies of each isotype were conjugated simultaneously a single SGD-1269 / NEM mixed solution for each isotype was prepared by multiplying the number of wells containing that isotype by 10 μ L per well then diluting into a volume of PBS equal to 500 μ L times the number of wells. In like fashion a total of eight drug-linker / NEM mixes were prepared-four with SGD-1006 and four with SGD-1269-and diluted into PBS. These mixes were then added to the reduced antibodies (500 μ L per well) and the plate was shaken for 30 minutes at 22 °C. The plate was then centrifuged as above to remove the excess reaction solution, and subsequently washed 4 times with PBS as before.

[0192] The bound ADCs were then eluted by adding 200 μ L of 50 mM glycine pH 2.5 to each well and shaking the plate for 3 minutes at 1200 RPM. While shaking 20 μ L of neutralization buffer (1M potassium phosphate, pH 7.4, 500 mM NaCl, 0.2% Tween-20) was added to each well of a 1 mL collection plate. The ADCs were then eluted into the collection plate by spinning at 1500x g for 6 minutes. The collection plate was then shaken briefly to ensure complete mixing of the neutralization buffer.

[0193] The concentration of each ADC was then determined with an absorbance plate reader by transferring the solutions into a UV assay plate (Costar model 3635, Corning) and measuring the optical density at 280 nm. An average IgG extinction coefficient of 1.45 mL mg⁻¹ cm⁻¹ was used to provide an adequate estimation of ADC concentration across the panel. To confirm successful conjugation, a reversed phase protein HPLC method (described below) was used to estimate the drug loading of the isotype controls. For the plate containing the humanization variants of CA8 this method was used to estimate the loading of all ADCs directly.

[0194] The reversed phase protein chromatography method for determining drug loading employs the PLRP-S polymeric stationary phase (Agilent Technologies). Since the antibodies were fully reduced during the conjugation process all of the antibody subunits elute from the column as single polypeptide chains allowing the subpopulations of light and heavy chain species with varying levels of drug loading to be evaluated separately. Thus, the analysis of these data allow for the calculation of the average light chain drug loading and the average heavy chain drug loading as independent factors which can then be combined to determine average antibody drug loading with the basic knowledge that each antibody is comprised of two light and two heavy chains. The chromatographic conditions were as follows: A PLRP-S column, 1000 Å, 50 × 2.1 mm, 8 μ m particle size (Agilent Technologies) with water + 0.05% TFA as mobile phase A and acetonitrile + 0.01% TFA as mobile phase B; elution with a linear gradient of 27% B to 42% B in 12.5 minutes.

[0195] Anti-BCMA antibodies were conjugated with SGD-1006 and SGD-1269 in three separate batches over a period of seven months. In the first batch a total of 29 antibodies were conjugated (resulting in 58 ADCs). The drug loading of each isotype control determined by PLRP chromatography and the data are summarized in Table 3.

Table 3.

Isotype	SGD-1006 loading	SGD-1269 loading
clgG1 (control P)	4.23	4.35
clgG1 (control M)	4.42	4.41
mlgG1	4.26	4.04
mlgG2a	4.51	4.57
mlgG2b	4.39	4.18

[0196] For the second batch an additional 25 antibodies were conjugated (resulting in 50 ADCs). The drug loading of each isotype control was again determined by PLRP chromatography and the data are summarized in Table 4.

Table 4.

Isotype	SGD-1006 loading	SGD-1269 loading
clgG1	3.96	3.78
mlgG1	3.95	3.32
mlgG2a	4.53	3.60
mlgG2b	4.32	3.49

[0197] In the third batch 30 antibodies were conjugated (resulting in 60 ADCs), including 13 humanized variants of CA8. In this final batch, the drug loading of all ADCs were determined and are summarized in the following two plate maps. (Table 5 & 6)

Table 5.

	drug loading										
	1	2	3	4	5	6	7	8	9	10	
A	3.7	4.0	3.6	3.8	3.8	3.5	3.9	2.8	3.8	3.8	
B	3.7	3.6	3.5	3.7	4.0	3.4	3.7	3.3	3.8	3.9	
C	3.6	3.8	3.5	3.7	3.6	3.3	3.8	4.7	3.8	3.7	
D	3.4	3.6	3.6	3.9	3.9	3.4	3.2	4.8	3.8	3.9	
E	3.9		3.8	3.9	3.4	3.6		3.3	3.7	3.4	
F	3.7			4.0	3.6	3.5			3.8	3.7	
G	3.6			3.6		3.4			3.7		
H				3.6					3.6		
	SGD-1006 (vc-MMAE) ADCs					SGD-1269 (mc-MMAF) ADCs					
	3.7	3.8	3.6	3.8	3.4	3.7	3.8		3.7		
	4.1%	5.1%	3.4%	4.8%	2.8%	8.5%	24.1%		3.4%		

Table 6.

	mlgG1	mlgG2a	mlgG2b	humanized	mlgG1	mlgG2a	mlgG2b	humamed			
	1	2	3	4	5	6	7	8	9	10	
A	control	control	control	CA8 J6MO	CA8 J8M2	control	control	control	CA8 J6MO	CA8 J8M2	
B	5336106D07	5336105A07	S336107G05	CAS J6M1	CA8 J9M0	5336106D07	5336105A07	S336107G08	CAS J6M1	CAS J9M0	
C	5335115G03	5335122F05	5336104A09	CA8 J6M2	CA8 J9M1	5335115G03	5335122F05	5336104A09	CA8 J6M2	CA8 J9M1	
D	5335115G01	5335128A12	5335107H11	CA8 J7M0	CA8 J9M2	5335115G01	5335128A12	5335107H11	CA8 J7M0	CA8 J9M2	
E	5335106E08		5335119E11	CA8 J7M1	CA8 Fc ENH	S335106EOS		5335119E11	CA8 T7M1	CA8 Fc ENH	
F	5335132E01			CA8 J7M2	GRITS28785	S335132E01			CA8 J7M2	GRITS28785	
G	5341106G02			CAS J8M0		S341106G02			CAS J8M0		
H				CA8 J8M1					CA8 J8M1		
	SGD-1 006 (vc-MMAE) ADCs					SGD-12 269 (mc-MMAF) ADCs					

[0198] Mean drug loading and %CV are indicated for each isotype series at the bottom. An uncharacteristically large variability in drug loading was observed for the SGD-1269 ADCs prepared with mlgG2b antibodies; the reason for this is unclear. Also, the Fc-enhanced CA8 antibodies yielded somewhat lower drug loading levels than the other CA8 human variants; to address this, additional Fc-enhanced CA8 was conjugated in a solution-phase reaction to better match the drug loading achieved for the other antibodies.

Example 4 - Binding Data

4.1 FMAT binding assay to show binding of Chimeric CA8 to cells expressing human or cyano BCMA.

[0199] Cryopreserved transfected human, cyano BCMA and mock transfected HEK293 cells were recovered from LN2 storage. Assay wells were prepared with human chimeric CA8 antibody, at a range of different concentrations, mixed with human BCMA HEK293, cyano BCMA HEK293 and mock transfected cells respectively. Anti-human IgG FMAT Blue secondary conjugate was added for detection of human chimeric CA8. The assay plates were left for a minimum of 90 minutes before the result was read on the ABI8200 (FMAT) plate reader.

[0200] This showed that the CA8 antibody in chimeric form binds well to both human and cyno BCMA proteins expressed on HEK293 cells.

[0201] Results are shown in Figure 1.

4.2 ELISA experiment showing binding of chimeric CA8 to recombinant BCMA protein

[0202] Chimeric CA8 antibodies were tested for binding to human BCMA and cyno BCMA expressed as Fc fusions. Human BCMA-Fc and cyno BCMA-Fc were coated to ELISA plates and the plates were blocked using BSA to reduce non specific binding. CA8 chimeric antibodies were added in a concentration range from 5ug/ml to 0.1ug/ml to the human and cyno BCMA coated ELISA plates. Any bound human chimeric CA8 antibody was detected using anti-human IgG HRP conjugated secondary antibody as appropriate. HRP substrate (TMB) was added to develop the ELISA. This showed that CA8 antibody binds to recombinant human and cyno BCMA in an ELISA assay.

[0203] Results are shown in Figure 2.

4.3 Biacore experiment to show CA8 antibody binding to BCMA and TACI proteins to determine cross reactivity with TACI protein.

[0204] CA8 chimera antibody was injected and captured on protein A. (A protein A derivitised sensorchip was used). Residual protein A binding was blocked with an injection of a high concentration of human IgG solution. BCMA-Fc, TACI-Fc or BAFF-R-Fc solutions were then tested for binding to the antibody. The 3 proteins were injected in sequence and binding events were measured. The surface was regenerated between injection of each protein.

[0205] Sensorgrams were analysed in the Biaevaluation program. Double reference subtraction was done to remove instrument noise and any non-specific binding from the sensorgram curves.

[0206] This showed that CA8 was specific for binding to BCMA binding and not to TACI and BAFFR.

[0207] Binding of the CA8 antibody to BCMA-Fc, TACI-Fc and BAFF-R-Fc was plotted out as shown in Figure 3.

4.4 Cell binding and neutralisation data

4.4.1 Binding of murine anti BCMA antibodies to multiple myeloma cells and BCMA expressing cells

[0208] Multiple myeloma cell line H929 and ARH77-hBCMA 10B5 BCMA expressing transfectant cells were stained with murine S332211D07, S3332121 F02 or S332126E04 or murine isotype control at 5 µg/mL. Multiple myeloma cell line H929 was stained with murine S307118G03. Cells were incubated for 20 mins at room temperature (RT) and then washed with FACS buffer (PBS + 0.5% BSA +0.1% sodium azide) to remove unbound antibody. Cells were incubated with a secondary PE labelled antimouse IgG antibody for 15 minutes at RT and then washed with FACS buffer to remove unbound antibody. Cells were analysed by FACS to detect antibody bound to the cells.

[0209] The results (Figure 4) showed that all 4 murine antibodies bound to the H929 multiple myeloma cell line and the three antibodies tested on ARH77 BCMA transfected cells bound to these.

4.4.2 Binding curve of chimeric CA8 to multiple myeloma cells as determined by FACS

[0210] A panel of multiple myeloma cell lines were used to determine the binding of chimeric CA8. Cell lines H929, OPM-2, JJN-3 and U266 were stained with either chimeric CA8 or irrelevant antibody (Synagis) at varying concentrations for 20 minutes at RT. Cells were then washed with FACS buffer (PBS + 0.5% BSA + 0.1% sodium azide) to remove unbound antibody. Cells were incubated with a secondary PE labelled anti-human IgG antibody for 15 minutes at RT and then washed with FACS buffer to remove unbound antibody. Cells were analysed by FACS and mean fluorescence intensity (MFI) values measured to determine binding.

[0211] Results showed that chimeric CA8 bound to multiple myeloma cell lines H929, OPM-2, JJN-3 & U266 in a dose dependent manner (Figure 5).

4.4.3 Binding of humanised CA8 to BCMA transfected cells as determined by FACS

[0212] ARH77-hBCMA 10B5 BCMA expressing transfectant cells or H929 cells were stained with either chimeric CA8 or humanised variants of CA8 designated J6M0, J6M1, J6M2, J9M0, J9M1, J9M2 at varying concentrations for 20 minutes at RT. Cells were then washed with FACS buffer (PBS + 0.5% BSA + 0.1% sodium azide) to remove unbound antibody. Cells were incubated with a secondary PE labelled anti-human IgG antibody for 15 minutes at RT and then washed with FACS buffer to remove unbound antibody. Cells were analysed by FACS and mean fluorescence intensity (MFI) values measured to determine binding.

[0213] Results showed that chimeric CA8 and all antibodies tested apart from J9M2 bound to ARH77-hBCMA 10B5 BCMA expressing transfectant

cells and H929 cells in a dose dependent manner (Figure 6).

4.5 Demonstration of ability of CA8 and the humanised version J6MO to neutralise binding of BAFF or APRIL to recombinant BCMA.

[0214] The aim of this assay was to assess the ability of antibody CA8, and humanised version J6M0 in both wild type and afucosylated (Potelligent) form, at various concentrations, to neutralise the binding ability of either BCMA ligand, BAFF or APRIL.

[0215] 96 well flat bottomed plates were coated overnight with 1 μ g/mL solution of recombinant human BCMA Fc 4-53 in PBS. Following a wash step using 0.05% TWEEN20, plates were blocked with 2% Bovine Serum Albumin solution in PBS for 1 hour at room temperature. Plates were washed as before and 40 μ L of each antibody (murine IgG, murine CA8, and chimeric CA8), starting at 10 μ g/mL, titrated at 1 in 2 in duplicate was added to the relevant wells and incubated for 1 hour at room temperature. 40 μ L of 2% BSA was added to the relevant control wells. 10 μ L of either recombinant human BAFF (2149-BF/CF, R&D Systems) or recombinant human APRIL (5860-AP/CF, R&D Systems) was added at 30ng/mL and 750ng/mL respectively, giving a final concentration of 6ng/mL and 150ng/mL respectively in each well. Equivalent volume of 2% BSA was added to the relevant control wells. Plates were allowed to incubate for 2 hours at room temperature, after which they were washed as before. Biotinylated anti-human ligand (BAFF BAF124 or APRIL BAF884, R&D Systems) was added to the relevant wells at 50ng/mL and incubated for 1 hour. Following a wash step, 50 μ L of a 1:4000 dilution of Streptavidin-HRP (Amersham RPN4401) was added to each well and incubated for 30 minutes at room temperature. The wash process was repeated again followed by the addition of 100 μ L of Tetramethylbenzidine substrate solution (T8665, Sigma) into each well. Plates were incubated for 20-25 minutes at room temperature, wrapped in foil. The reaction was stopped with the addition of 100 μ L of 1M H₂SO₄. Optical density was determined at 450nm using Spectromax reader. See Figure 7A and B.

[0216] In a plate based assay for neutralisation of binding of BAFF or APRIL to BCMA, the EC50 values calculated for chimeric CA8 were 0.695 μ g/mL and 0.773 μ g/mL respectively. The values for the humanised J6M0 were 0.776ng/ml and 0.630ng/ml. The values for the J6M0 potelligent version were 0.748 and 0.616ng/ml respectively.

4.6 Effect of chimerised CA8 and humanised J6MO BCMA antibody on BAFF or APRIL induced phosphorylation of NF κ B in H929 cells.

[0217] In one set of experiments, H-929 cells were plated at 75,000cells/well in a 96 well plate in serum free medium. The chimeric CA8 antibody was added 24 hours later to give final well concentrations up to 200 μ g/ml. Ten minutes later, BAFF or APRIL ligand were added to the cells to give final well concentrations of 0.6 or 0.3 μ g/ml respectively. After 30 minutes the cells were lysed and phosphorylated NfkappaB levels measured using a MSD pNFkappaB assay.

[0218] The chimeric BCMA antibody CA8 neutralised both BAFF and APRIL induced NfkappaB cell signalling in H-929 cells. It was particularly potent at neutralising BAFF induced NfkappaB cell signalling in this cell type with a mean IC50 of 10nM, compared to 257nM for APRIL induced NfkappaB cell signalling.

Meaned data for 2 experiments

[0219] IC50s were 10nM for BAFF induced NfkappaB neutralisation and 257nM for APRIL induced NfkappaB neutralisation (mean of 2 independent experiments) are shown in Table 7.

Table 7

	BAFF induced IC50		APRIL induced IC50	
	ug/ml	nM	ug/ml	nM
BCMA antibody CA8	1.5	10	38.5	256.7

[0220] A further set of experiments were carried out to aim to understand why there was such a discrepancy between the potency in neutralisation of APRIL and BAFF in the cell based system. Following the discovery of the soluble form of BCMA the experimental design was changed to include a step where the H929 cells were washed prior to the assay to reduce the interference from the antibody binding to soluble BCMA. H-929 cells were washed 3 times to remove any sBCMA and resuspended in serum free medium. J6M0 potelligent antibody was added to a 96 well plate to give a final well concentrations up to 100 μ g/ml along with BAFF or APRIL ligand to give a final well concentration of 0.6 or 0.2 μ g/ml respectively. H-929 cells were then plated at 7.5 \times 104cells/well in serum free medium. 30 minutes later the cells were lysed and phosphorylated NF κ B levels measured using a MSD pNFkappaB assay. This is data from one experiment. Each data point is the mean/sd of two replicates. The data from this experiment is shown in Figure 7c. The IC50s for inhibition of BAFF and APRIL signalling were determined as 0.91 μ g/ml and 2.43 μ g/ml respectively.

4.7 ProteOn analysis of anti-BCMA CA8 chimeric and humanised constructs

[0221] The initial screen of CA8 chimeric and humanised variants was carried out on the ProteOn XPR36 (Biorad). The method was as follows; Protein A was immobilised on a GLC chip (Biorad, Cat No: 176-5011) by primary amine coupling, CA8 variants were then captured on this surface and recombinant human BCMA (in house or commercial US Biological, B0410) materials (run 2 only) passed over at 256, 64, 16, 4, 1nM with a 0nM injection (i.e. buffer alone) used to double reference the binding curves, the buffer used is the HBS-EP buffer. 50mM NaOH was used to regenerate the capture surface. The data was fitted to the 1:1 model using the analysis software inherent to the ProteOn XPR36.. Run 1 corresponds to the first screen of humanised CA8 variants (J0 to J5 series) and run 2 to the second screen of humanised CA8 variants (J5 to J9 series). Both runs were carried out at 25°C.

[0222] The data obtained from run1 are set out in Table 8 and data from run 2 are set in Table 9. Several molecules in the Run 2 (Table 09) failed to give affinity values measurable by ProteOn, this was due to the off-rate being beyond the sensitivity of the machine in this assay, this does however indicate that all these molecules bind tightly to recombinant human BCMA. From Run 1 the data indicates that some constructs did not show any binding to recombinant cyno BCMA.

Table 8: Run 1-Kinetics analyses of anti-BCMA molecules against Recombinant Human BCMA

Sample name	Human in house BCMA			Cyno in house BCMA		
	ka	kd	KD (nM)	ka	kd	KD (nM)
CA8 humanised J5M0	2.16E+05	1.88E-05	0.087	3.25E+05	8.14E-06	0.025
CA8 humanised J5M2	2.67E+05	3.21E-05	0.12	4.30E+05	4.70E-05	0.109
CA8 humanised J5M1	2.97E+05	4.32E-05	0.145	4.81E+05	5.41E-05	0.112
CA8 humanised J4M1	2.54E+05	7.04E-05	0.278	3.50E+05	7.10E-05	0.203
CA8 humanised J4M2	2.51E+05	7.06E-05	0.281	3.44E+05	6.15E-05	0.179
CA8 humanised J0M2	2.25E+05	6.97E-05	0.31	3.26E+05	1.84E-04	0.563
CA8 humanised J3M2	2.66E+05	9.64E-05	0.362	3.69E+05	5.87E-05	0.159
CA8 humanised J0M1	2.31E+05	8.60E-05	0.373	3.32E+05	1.67E-04	0.503
CA8 humanised J0M0	2.45E+05	1.06E-04	0.435	3.58E+05	2.32E-04	0.648
CA8 humanised J3M1	2.85E+05	1.25E-04	0.438	4.04E+05	7.93E-05	0.196
CA8 humanised J2M2	2.05E+05	9.87E-05	0.482	2.98E+05	3.17E-05	0.106
CA8 Chimera	2.41E+05	1.25E-04	0.519	3.82E+05	1.74E-04	0.457
CA8 humanised J2M1	2.04E+05	1.72E-04	0.842	2.96E+05	6.46E-05	0.218
CA8 humanised J4M0	2.42E+05	2.20E-04	0.906	3.34E+05	2.89E-04	0.866
CA8 humanised J1M2	2.15E+05	2.46E-04	114	3.19E+05	9.67E-05	0.303
CA8 humanised J3M0	2.08E+05	2.85E-04	1.37	2.93E+05	1.54E-04	0.526
CA8 humanised J1M1	2.27E+05	3.43E-04	1.51	3.33E+05	1.47E-04	0.442
CA8 humanised J2M0	1.95E+05	3.77E-04	194	2.81E+05	1.51E-04	0.538
CA8 humanised J1M0	1.78E+05	5.02E-04	2.82	2.47E+05	2.10E-04	0.849
S307118G03 Chimera	4.75E+05	1.95E-03	4.11	No	Analysable	Binding
S307118G03 humanised H3L1	4.69E+05	2.28E-03	4.86	No	Analysable	Binding
S307118G03 humanised H3L0	2.86E+05	1.52E-03	5.31	No	Analysable	Binding
S307118G03 humanised H2L0	3.78E+05	2.41E-03	6.36	No	Analysable	Binding
S307118G03 humanised H2L1	3.38E+05	2.15E-03	6.37	No	Analysable	Binding
S307118G03 humanised H4L1	No	Analysable	Binding	No	Analysable	Binding

Table 9: Run 2-Kinetics analyses of anti-BCMA molecules against Recombinant Human BCMA

For antibodies J8M0, J9M0, J8M1, J9M2, J7M2, J5M0, J7M1, J7M0, J8M2, J9M1, J5M2, J5M1 the off rate was beyond the sensitivity of the assay hence no data shown.									
Sample Name	Human in house BCMA			commercial human BCMA			Cyno in house BCMA		
	ka	kd	KD (nM)	ka	kd	KD (nM)	ka	kd	KD (nM)
CA8 Chimera	2.51E+05	1.03E-04	0.412	7.05E+05	9.79E-05	0.139	5.89E+04	1.21E-04	2.060
CA8 humanised J6M1	2.17E+05	2.70E-05	0.124	5.92E+05	3.75E-05	0.063	4.88E+04	2.58E-04	5.300
CA8 humanised J6M0	2.40E+05	7.40E-05	0.308	6.23E+05	5.37E-05	0.086	5.64E+04	3.18E-04	5.630
CA8 humanised J6M2	2.01E+05	4.06E-05	0.202	5.63E+05	3.97E-05	0.071	4.41E+04	3.02E-04	6.860
S307118G03 H5L0	No Analysable Binding			V weak signal			No	Analysable	Binding
S307118G03 H5L1	No Analysable Binding			V weak signal			No	Analysable	Binding
S307118G03Chimera	4.79E+05	1.65E-03	3.44	1.55E+06	1.48E-03	0.956	No	Analysable	Binding

4.8 BIACore analysis of anti-BCMA CA8 chimeric and humanised constructs (J7 to J9 series)

[0223] Protein A was immobilised on a CM5 chip (GE Healthcare, Cat No: BR-1005-30) by primary amine coupling and this surface was then used to capture the antibody molecules. Recombinant human BCMA (US Biological, B0410) was used as analyte at 256nM, 64nM, 16nM, 4nM and 1nM. Regeneration of the capture surface was carried out using 50mM NaOH. All binding curves were double referenced with a buffer injection (i.e. OnM) and the data was fitted to the using the 1:1 model inherent to T100 evaluation software. The run was carried out at 37°C, using HBS-EP as the running buffer.

[0224] The results showed the molecules tested with the exception of J9M2 bind to recombinant human BCMA, with similar affinity as the chimeric

molecule. Data generated from this experiment are presented in table 10.

Table 10: Kinetics analysis of anti-BCMA humanised molecules against Recombinant Human BCMA

Sample name	Human commercial BCMA			Cyno in house BCMA		
	ka	kd	KD (nM)	ka	kd	KD (nM)
CA8 humanised J9M1	1.96E+07	3.50E-04	0.018	6.77E+05	2.99E-04	0.442
CA8 humanised J9M0	4.95E+06	1.74E-04	0.035	7.03E+05	3.24E-04	0.46
CA8 Chimera	3.27E+07	1.18E-03	0.036	1.15E+06	3.49E-04	0.305
CA8 humanised J8M1	2.66E+06	1.34E-04	0.05	2.82E+05	3.62E-04	1.284
CA8 humanised J8M0	2.44E+06	1.26E-04	0.052	3.89E+05	4.18E-04	1.076
CA8 humanised J7M1	2.35E+06	1.31E-04	0.056	3.70E+05	3.91E-04	1.057
CA8 humanised J8M2	2.63E+06	1.50E-04	0.057	3.83E+05	5.06E-04	1.324
CA8 humanised J7M2	2.37E+06	1.35E-04	0.057	3.46E+05	4.47E-04	1.293
CA8 humanised J7M0	2.36E+06	1.51E-04	0.064	3.21 E+05	3.67E-04	1.143
CA8 humanised J9M2	No	Analysable	Binding	4.88E+05	2.52E-04	0.515

4.9 BIACore analysis of anti-BCMA CA8 chimeric and humanised constructs J6M0 and J9M0

[0225] Protein A was immobilised on a CM5 chip (GE Healthcare, Cat No: BR-1005-30) by primary amine coupling and this surface was then used to capture the antibody molecules. Recombinant human BCMA (US Biological, B0410) was used as analyte at 256nM, 64nM, 16nM, 4nM and 1nM. Regeneration of the capture surface was carried out using 50mM NaOH. All binding curves were double referenced with a buffer injection (i.e. 0nM) the data was fitted to the using the 1:1 model inherent to T100 evaluation software. The run was carried out at 25°C and 37°C for experiment 1 and only 37°C for experiment 2 using HBS-EP as the running buffer.

[0226] The both runs identified J9M0 as the best molecule in term of overall affinity to human BCMA. Data generated from this experiment are presented in table 11.

Table 11 Kinetics analyses of anti-BCMA humanised molecules against Human BCMA

Sample	Human commercial BCMA								
	25°C			37°C					
	Experiment 1			Experiment 1			Experiment 2		
Sample	ka	kd	KD (nM)	ka	kd	KD (nM)	ka	kd	KD (nM)
J9M0	1.59E+ 06	3.38E-05	0.021	3.75E+ 06	1.58E-04	0.042	3.62E+ 06	1.89E-04	0.052
J6M0	1.01E+ 06	1.22E-04	0.121	2.12E+ 06	1.48E-03	0.698	3.78E+ 06	1.88E-03	0.498
Chimera CA8	1.88E+ 06	2.63E-04	0.140	1.72E+ 07	8.72E-04	0.051	1.88E+ 07	1.04E-03	0.055

4.10. ProteOn analysis of new anti-BCMA chimeric constructs

[0227] The initial screen of the new chimeric variants from the second batch of hybridomas was carried out on the ProteOn XPR36 (Biorad). The method was as follows; Protein A was immobilised on a GLM chip (Biorad, Cat No: 176-5012) by primary amine coupling, anti-BCMA variants were then captured on this surface and recombinant human BCMA (in house material) passed over at 256, 64, 16, 4, 1nM with a 0nM injection (i.e. buffer alone) used to double reference the binding curves, the buffer used is the HBS-EP buffer. Regeneration of the capture surface was carried out using 50mM NaOH. The data was fitted to the 1:1 model using the analysis software inherent to the ProteOn XPR36. The run was carried out at 25°C.

[0228] Data generated from this experiment are presented in table 12.

Table 12: Kinetics analyses of anti-BCMA humanised molecules against Human BCMA

Sample name	In house human BCMA		
	ka	kd	KD (nM)
S332110D07	3.11E+05	3.77E-03	12.100
S332121F02	3.73E+05	6.45E-03	17.300

Example 5 Cell Killing Assays.

5.1 ADCC potencies of chimeric CA8 and defucosylated chimeric CA8 version in ARH77 cells expressing BCMA

[0229] Human natural killer (NK) cells were incubated with europium labelled ARH77 BCMA transfected target cells (10B5) in the presence of

varying concentrations of antibody at an E:T ratio of 5:1 for 2 hours. Europium release from the target cells was measured and specific lysis calculated.

[0230] Result: Chimeric CA8 and defucosylated chimeric CA8 killed BCMA expressing target cells via ADCC. The defucosylated chimeric antibody showed more potent ADCC activity, as measured by a higher percent lysis achieved with all the target cells tested and a ten-fold lower EC₅₀ on the high BCMA expressing target cell line 10B5, compared to the parent chimeric antibody. See Figure 8A and 8B.

5.2 ADCC activity of CA8 humanised antibodies using ARH77 BCMA expressing target cells and PBMC as effectors

[0231] Human PBMC were incubated with europium labelled ARH77 BCMA transfected target cells (10B5) in the presence of varying concentrations of humanised versions of CA8 antibody (5ug/ml to 0.005ug/ml) at an E:T ratio of 5:1 for 2 hours. Europium release from the target cells was measured and specific lysis calculated.

Result:

[0232] Result: All the J5, J6, J7 J8 & J9 series of humanised variants of CA8 showed ADCC activity against the ARH77 high BCMA expressing cell line 10B5 in a dose dependent manner. ADCC was at a similar level as that found in the experiments using chimeric CA8 molecule. See Figure 9.

5.3 ADCC potencies of chimeric S322110F02, S322110D07 and S307118G03 and humanised S307118G03 H3L0 against ARH77 10B5 cells expressing BCMA with purified NK cells as effector cells

[0233] Human natural killer (NK) target cells were incubated with europium labelled ARH77 BCMA transfected target cells (10B5) in the presence of varying concentrations of antibody at an E:T ratio of 5:1 for 2 hours. Europium release from the target cells was measured and specific lysis calculated. Result: all 4 antibodies tested showed ADCC activity against ARH77 10B5 cells. See Figure 10.

5.4 Antibody-Drug Conjugate (ADC) activity of Chimeric CA8 ADCs.

[0234] Measuring ADC activity of chimeric CA8 antibody, chimeric CA8-mcMMAF antibody drug conjugates and chimeric CA8-vcMMAE antibody drug conjugates against human multiple myeloma cell lines. Multiple Myeloma cell lines were treated with chimeric CA8 antibody-drug conjugates to determine the ADC concentrations required for growth inhibition and death.

[0235] The antibody drug conjugates tested were added to wells containing multiple myeloma cells at concentrations ranging from 1ug/ml to 5ng/ml. The plates were incubated at 37°C for 96 hours at which point viable cells were quantitated using Cell titre Glo. The unconjugated chimeric CA8 antibody showed no significant growth inhibitory activity at the antibody concentrations that were tested. The chimeric CA8-mcMMAF antibody-drug conjugate showed greater growth inhibitory activity than the chimeric CA8-vcMMAE antibody-drug conjugate in all 4 of the multiple myeloma cell lines that were tested. See Figure 11 and Table 13

Table 13 IC₅₀ values represented in ng/mL for the chimeric CA8-vcMMAE and the chimeric CA8-mcMMAF antibody-drug conjugates in 4 different multiple myeloma cell lines

Multiple Myeloma cell lines	IC ₅₀ (ng/mL)	
	CA8 chimera - vcMMAE	CA8 chimera - mcMMAF
NCI-H929	29.5	8.8
U266-B1	18.9	9.7
JJN3	21.8	12.4
OPM2	92.7	58.1

5.5 Measuring cell cycle arrest activity of chimeric CA8 antibody, chimeric CA8-mcMMAF antibody drug conjugates and chimeric CA8-vcMMAE antibody drug conjugates against human multiple myeloma cell line H929.

[0236] To determine the mechanism that chimeric CA8 Antibody Drug Conjugates (ADC's) cause growth inhibition in multiple myeloma cells, the cell cycle of NCI-H929 cells was monitored by measuring cellular DNA content through fixed cell propidium iodide staining at multiple timepoints following chimeric CA8 antibody and chimeric CA8 ADC treatment.

[0237] At the chimeric CA8 ADC concentration tested (50ng/mL), the chimeric CA8-mcMMAF ADC caused significant G2/M cell cycle arrest (4N DNA content) which peaked at 48 hours. At the later timepoints 48, 72 and 96 hours, treatment with the chimeric CA8-mcMMAF ADC resulted in accumulation of a cell population with sub-2N DNA content, which is representative of cell death. At the 50ng/mL concentration tested the chimeric CA8-vcMMAE ADC had no significant effect on G2/M cell cycle arrest or sub-G1 accumulation. See Figure 12.

5.6 Phospho-Histone-H3 (Thr11) staining as a marker for chimeric CA8-mcMMAF antibody drug conjugate and chimeric CA8-vcMMAE antibody drug conjugate induced mitotic arrest.

[0238] To determine if the accumulation of cells with 4N DNA content is a specific result of mitotic arrest induced by the chimeric CA8 ADCs NCI-H929 cells were stained with an anti-phospho-Histone H3 antibody following treatment with increasing concentrations of unconjugated chimeric CA8, chimeric CA8-vcMMAE or chimeric CA8-mcMMAF for 48 hours.

[0239] Treatment with chimeric CA8 ADCs resulted in a dose-dependent accumulation of NCI-H929 cells that stained positive for 65eroxidi-Histone H3 (Thr11), a specific marker of mitotic cells. The chimeric CA8-mcMMAF ADC caused accumulation of 65eroxidi-Histone H3 positive cells at lower concentrations than the chimeric CA8-vcMMAE ADC. See Figure 13.

5.7 Measuring apoptosis in NCI-H929 cells in response to chimeric CA8 ADCs by staining for Annexin V.

[0240] To determine if the accumulation of cells with sub-2N DNA content is a specific result of apoptosis induced by the chimeric CA8 ADCs, NCI-H929 cells were stained with an anti-Annexin-V antibody following treatment with increasing concentrations of unconjugated chimeric CA8, chimeric CA8-vcMMAE or chimeric CA8-mcMMAF for 48 hours. Treatment with chimeric CA8 ADCs resulted in a dose-dependent accumulation of NCI-H929 cells that stained positive for Annexin-V, a specific marker of apoptosis. The chimeric CA8-mcMMAF ADC caused accumulation of Annexin-V positive cells at lower concentrations than the chimeric CA8-vcMMAE ADC. See Figure 14.

5.8 Antibody-Drug Conjugate (ADC) activity of humanised variants of CA8 anti-BCMA antibody-drug conjugates.

Cells were plated in 96-well plates (4,000 cells per well in 100uL of RPMI + 10% FBS)

[0241] Naked antibody or ADC was added 6 hours after cell seeding and plates were incubated for 144 hours. Growth inhibition in the presence of the antibodies or ADCs was measured at 144 hours using Cell Titre glo. Data points represent the mean of triplicate CellTiterGlo measurements. Error bars represent standard error.

[0242] Multiple Myeloma cell lines NCI-H929 and OPM2 were treated with humanized CA8 anti-BCMA antibody-drug conjugates to determine the ADC concentrations required for growth inhibition and death. The mcMMAF and vcMMAE antibody-drug conjugate forms of these antibodies showed significant growth inhibitory activity comparable to that found with the CA8 chimera. Variant J6M0 showed higher potency than the chimera and data is shown in figure 15 in H929 cells and OPM2 cells. The mcMMAF antibody-drug conjugate showed greater growth inhibitory activity than the vcMMAE antibody-drug conjugate for all antibodies in both cell lines tested. Results for all humanized variants are shown in Table 14.

Table 14. IC₅₀ values represented in ng/mL for the anti BCMA antibody-drug conjugates in NCI-H929 and U266-B1 cells

	NCI-H929 mcMMAF	vcMMAE	OPM2 mcMMAF	vcMMAE
	Average IC ₅₀ (ng/mL)			
CA8 chimera	11.64	37.96	57.04	80.01
CA8 J6M0	5.97	27.67	87.22	121.2
CA8 J6M1	14.6	51.89	205.6	239.9
CA8 J6M2	9.5	39.71	112.9	144.7
CA8 J7M0	18.97	52.25	93.27	127.1
CA8 J7M1	17.87	43.97	95.35	107.5
CA8 J7M2	31.63	55.13	102.6	115.9
CA8 J8M0	15.67	59.94	89.95	132
CA8 J8M1	17.04	46.55	82.96	115.8
CA8 J8M2	15.08	55.98	72.63	124.5
CA8 J9M0	14.95	48.5	58.6	109.8
CA8 J9M1	15.19	55.1	55.88	115
CA8 J9M2	20.87	55.77	80.35	111.7

5.9 Antibody-Drug Conjugate (ADC) activity of other murine anti-BCMA antibody-drug conjugates.

Cells were plated in 96-well plates (4,000 cells per well in 100uL of RPMI + 10% FBS)

[0243] Antibody or ADC was added 6 hours after cell seeding and plates were incubated for 144 hours. Growth inhibition in the presence of the ADCs was measured at 144 hours using Cell Titre glo. The mean of triplicate CellTiterGlo measurements are shown. Table 15a and 15b are from experiments carried out at different times on different series of antibodies. Multiple Myeloma cell lines NCI-H929 and U266-B1 were used for antibodies in Table 15a.

[0244] The mcMMAF and vcMMAE antibody-drug conjugate forms of murine antibodies S322110D07, S332121 F02 and S332136E04 showed

significant growth inhibitory activity. The mcMMAF antibody-drug conjugate showed greater growth inhibitory activity than the vcMMAE antibody-drug conjugate in all of the murine anti-BCMA antibodies tested where activity was seen. IC50 figures are shown in Table 15a. See Figure 16 for dose response curves for these three antibodies and also S107118G03. Error bars represent standard error. NCI-H929, U266-B1, JJN3 and OPM2 cells for antibodies in Table 15b were treated with a different series of murine anti-BCMA antibody-drug conjugates to determine the ADC concentrations required for growth inhibition and death. IC50 figures are shown in Table 15b. All 5 antibodies shown on the table had significant ADC activity.

Table 15a. IC₅₀ values represented in ng/mL for the anti BCMA antibody-drug conjugates in NCI-H929 and U266-B1 cells

Antibody	IC ₅₀ (ng/mL)		-U266-B1			
	NCI-H929	-vcMMAE	-mcMMAF	-vcMMAE	-mcMMAF	
S322110D07 mlgG1	28.4		6.7	53.3		33.3
S332121F02 mlgG1	24.5		7	- 2.3		2.5
S332126E04 mlgG1	46.8		9.7	- 27.1		10.6

Table 15b IC₅₀ values represented in ng/mL for the anti BCMA antibody-drug conjugates in NCI-H929, U266-B1, JJN3 and OPM2 cells

	NCI-H929		U266B1		JJN3		OPM2
Average IC50 (ng/mL)	vcMMAE	mcMMAF	vcMMAE	mcMMAF	vcMMAE	mcMMAF	vcMMAE
S335115G01	14.9	4.2	38.8	18.5	73.9	45.8	162.4
S336105A07	17.8	5.1	21.4	9.3	54.2	23.2	95.5
S335122F05	10.9	4.2	21.1	14.1	29.5	25.5	98.4
S335106E08	19.2	7.9	36.8	32.6	189.8	214.1	243.9
S335128A12	86.3	28.3	101.8	104.1	>500	>500	>500

5.10 ADCC potency of conjugated, afucosylated J6M0 (Potelligent)

[0245] Afucosylated J6M0 conjugated to MMAE or MMAF was tested in ADCC assays using BCMA transfectants to ensure that its ADCC activity was not compromised by the conjugation. Europium labelled ARH77-10B5 cells were incubated with various J6M0 WT and Potelligent BCMA antibodies at concentrations up to 1 0000ng/ml for 30 minutes prior to the addition of PBMCs (PBMC: target cell ratio 50:1). Two hours later an aliquot of cell media was sampled and mixed with enhancement solution. After 30 minutes on a plate shaker, europium release was monitored on the Victor 2 1420 multi-label reader. Datapoints represent means of triplicate values. This data is representative of 2 experiments.

[0246] There were no significant differences in ADCC potency between the unconjugated and ADC forms of J6M0 Potelligent. In the same experiment a wild type version of J6M0 was included to show how the potency compares to the afucosylated version. As expected, defucosylation resulted in a lower EC50 and higher maximal lysis. No lysis was observed with the Fc disabled form of J6M0. (Figure 17)

5.11 ADCC potency of afucosylated J6M0 on MM cell lines

[0247] Human PBMC were incubated with multiple myeloma target cells at an E:T ratio of 50:1 in presence of varying concentrations of afucosylated (Potelligent) J6M0. The percentage of target cells remaining in the effector + target cell mixture after 18 hours was measured by FACS using a fluorescently labelled anti-CD138 antibody to detect the target cells and the percent lysis calculated. This is representative of several experiments.

[0248] J6M0 Potelligent antibody showed ADCC activity against all five multiple myeloma target cell lines tested. This was important to test since earlier studies were carried out using transfected cells. Results are shown in Figure 18. Full dataset with multiple donors is shown in Table 16. The potencies were all in a similar range as those found with the transfectants. The ADCC activity was not directly related to BCMA surface expression on these cell lines.

Table 16 EC₅₀ values generated on 13 independent assays using 11 donors (designated A-K) across the five multiple myeloma cell lines.

Donor	EC ₅₀ (ng/ml)				
	H929	RPMI 8226	JJN-3	OPM-2	U266
A	1.43	NA	1.64	NA	NA
R	057	NA	NA	NA	NA
C	0.73	NA	1.01	NA	NA
C	1.81	NA	NA	NA	NA
A	2.05	NA	NA	NA	NA
D	NA	4.09	NA	NA	NA
F	NA	NA	14.4	NA	NA
F	2.18	NA	NA	NA	NA
G	NA	NA	26.3	NA	NA
H	4.79	NA	111.3	NA	NA
I	NA	NA	40.1	NA	NA

Donor	EC ₅₀ (ng/ml)				
	H929	RPMI 8226	JJN-3	OPM-2	U266
J	2.19	20.4	4.89	NA	NA
K	ND	ND	4.52	4.15	9.04

Example 6. Xenograft data

[0249] 6.1 Murine xenografts of human MM cell lines were tested to ensure that antibody potency detected in vitro can also be demonstrated in vivo. The cell line selected for xenograft studies was NCI-H929 which is sensitive to ADC and ADCC killing in vitro. Studies were carried out in immunocompromised CB.17 SCID mice which lack T and B cells but maintain NK cells to allow for ADCC activity. However it should be noted that although human IgG1 can engage murine Fc receptors, the Potelligent enhancement does not improve the affinity as it does with human Fc receptors.

6.2 Impact of unconjugated and MMAE or MMAF conjugated J6M0 on NCI-H929 tumour growth.

[0250] In order to independently analyze both the ADCC and ADC activities of J6M0 we tested J6M0 antibody in the presence and absence of MMAF or MMAE conjugation. By testing the unconjugated J6M0, any anti-tumour effects could be attributed to some combination of ADDC and functional inhibitory activity. Mice with NCI-H929 tumours that had reached a volume of 200 mm³ on average were treated with a human IgG1 control or the J6M0 antibody (unconjugated, MMAE or MMAF) twice weekly at a dose of 50 ug or 100ug, for 2 weeks. Results from this study show that a 100 ug dose of the J6M0-MMAF conjugate resulted in elimination of tumours in those mice which have completed the dosing. The J6M0-MMAF mice were maintained for 40 days after the last dose with no recurrence of tumour occurring. These results from this experiment demonstrate that MMAF conjugation had increased anti-tumour activity over both unconjugated J6M0 antibody and J6M0-MMAE conjugate. See Figure 19.

Example 7 Evaluation of Soluble BCMA Levels from MM Patient Serum

[0251] 7.1 It is currently unknown whether BCMA is present extracellularly and can be detected in the blood. In this work, we determined the serum level of human BCMA from MM patients. Serum samples from 54 MM and plasma cell dyscrasia patients and 20 normal control samples were analyzed by ELISA. Human Subject Approval was obtained from Western Institutional Review Board.

7.2 Assessment of Serum Human BCMA Levels

[0252] Blood, from patients and normal controls in the clinic, were collected in serum collection tubes. MM patient samples were from a variety of stages (progressive disease, remission, relapsed, newly diagnosed, and others). The Blood samples were spun at 10,000 rpm for 10 minutes and serum transferred into sterile micro-centrifuge plastic tubes.

[0253] A Human BCMA/TNFRSF17 ELISA kit from R& D Systems (catalog # DY193E) which measures soluble human BCMA levels was used to detect BCMA following the standard protocol supplied with the kit.

[0254] Briefly, 96 well micro-plates were coated with 100ul per well capture antibody and incubated overnight at 4oC. The plates were washed three times with wash buffer (0.05% Tween 20 in PBS, pH 7.2) and blocked with 300ul of 1% BSA in PBS at room temperature for 2 hours. The plates were washed three times with washing buffer. 100ul of serum sample or standard was added into each well and incubated for 2 hours at room temperature. The plates were washed three times with washing buffer and then 100ul of the detection antibody was added to each well and incubated 2 hours at room temperature. 100ul of Streptavidin-HRP was added in each well after washing plates three times and incubated in dark room for 20 minutes. The plates were washed three times and added 50ul stop solution and then determined by micro-plate reader with 570nM wavelength.

[0255] A series of assays were carried out in order to determine the serum dilution factor appropriate for the levels of BCMA which were present. A dilution factor of 1:500 was found to be suitable for the majority of samples and is the dilution factor used in the data shown in Figure 20. The full data set is shown in Table 17.

[0256] Patient and normal control serum samples diluted and run in triplicates had BCMA levels determined. The serum levels of BCMA were significantly elevated in the sera from MM patients compared with normal controls in this study. When the disease subset was divided further there was a trend towards elevated serum levels of BCMA in the sera from progressing MM patients compared with those in remission.. This is the first report identifying serum BCMA in any human disease and suggests that these levels may be a novel biomarker for monitoring disease status and therapeutic response of MM patients and for other patients with plasma cell mediated diseases.

Table 17. Figures represent serum concentration of soluble BCMA in ng/ml calculated from samples diluted at 1/50, 1/500 and 1/5000. P values were calculated using the one tailed T-Test and 95% significance values are below the table.

1:5000	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
Mean	14.130	500.804	154.762	151.201	94.457	84.912	22.838

1-5000	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
1-500 Triplicate	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
Mean	15.901	215.877	81.135	43.294	97.584	53.894	22.838
1-500 Single	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
Mean	16.620	207.028	61.576	42.796	71.372	40.623	14.099
1-50 Trial 1	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
Mean	25.568	129.544	41.983	40.507	65.120	42.067	51.650
1-50 Trial 2	Normal	Myeloma: Progressive	Myeloma: Stable	Myeloma: Remission	Myeloma: Other	MGUS	Other Plasma Cell Dyscrasias
Mean	17.160	119.220	34.567	34.264	54.780	26.333	51.650

P-Values (One Tailed T-Test, 95% Significance)

~1-500 Single

Normal vs Progressive: p=.0010*

Progressive vs Remission: p=.0146*

~1-500 Triplicate

Normal vs Progressive: p=.0004*

Progressive vs Remission: p=.0091*

~1-50 Trial 1

Normal vs Progressive: p=.0171*

Progressive vs Remission: p=.0777

~1-50 Trial 2

Normal vs Progressive: p=.0184*

Progressive vs Remission: p=.0876

* shows significance

Sequence Summary (Table C)

Description	Amino acid sequence	Polynucleotide sequence
CA8 CDRH1	SEQ.I.D.NO:1	n/a
CA8 CDRH2	SEQ.I.D.NO:2	n/a
CA8 CDRH3	SEQ.I.D.NO:3	n/a
CA8 CDRL1	SEQ.I.D.NO:4	n/a
CA8 CDRL2	SEQ.I.D.NO:5	n/a
CA8 CDRL3	SEQ.I.D.NO:6	n/a
CA8 V _H domain (murine)	SEQ.I.D.NO:7	SEQ.ID.NO:8
CA8 V _L domain (murine)	SEQ.I.D.NO:9	SEQ.I.D.NO:10
CA8 Humanised V _H J0	SEQ.I.D.NO:11	SEQ.ID.NO:12
CA8 Humanised V _H J1	SEQ.I.D.NO:13	SEQ.I.D.NO:14
CA8 Humanised V _H J2	SEQ.I.D.NO:15	SEQ.I.D.NO:16
CA8 Humanised V _H J3	SEQ.I.D.NO:17	SEQ.I.D.NO:18
CA8 Humanised V _H J4	SEQ.I.D.NO:19	SEQ.I.D.NO:20
CA8 Humanised V _H J5	SEQ.LD.NO:21	SEQ.I.D.NO:22
CA8 Humanised V _H J6	SEQ.I.D.NO:23	SEQ.I.D.NO:24
CA8 Humanised V _H J7	SEQ.I.D.NO:25	SEQ.ID.NO:26
CA8 Humanised V _H J8	SEQ.I.D.NO:27	SEQ.I.D.NO:28
CA8 Humanised V _H J9	SEQ.I.D.NO:29	SEQ.I.D.NO:30
CA8 Humanised V _L M0	SEQ.I.D. NO:31	SEQ.I.D.NO:32
CA8 Humanised V _L M1	SEQ.I.D. NO:33	SEQ.I.D.NO:34
CA8 Humanised V _L M2	SEQ.I.D. NO:35	SEQ.ID.NO:36
Human BCMA	SEQ.I.D.NO:37	SEQ.I.D.NO:38
CD33-hBCMA ECD (1-53) TEV-Fc		
Human BCMA	SEQ.I.D.NO:39	SEQ.I.D.NO:40
CD33-hBCMA ECD (4-53) TEV-Fc		
Cyno BCMA	SEQ.I.D.NO:41	SEQ.I.D.NO:42

Description	Amino acid sequence	Polynucleotide sequence
CD33 cyano BCMA ECD (4-52) TEV-Fc		
CA8 J0 Humanised heavy chain	SEQ.I.D.NO:43	SEQ.I.D.NO:44
CA8 J1 Humanised heavy chain	SEQ.I.D.NO:45	SEQ.I.D.NO:46
CA8 J2 Humanised heavy chain	SEQ.I.D.NO:47	SEQ.I.D.NO:48
CA8 J3 Humanised heavy chain	SEQ.I.D.NO:49	SEQ.I.D.NO:50
CA8 J4 Humanised heavy chain	SEQ.I.D.NO:51	SEQ.I.D.NO:52
CA8 J5 Humanised heavy chain	SEQ.I.D.NO:53	SEQ.ID.NO:S4
CA8 J6 Humanised heavy chain	SEQ.I.D.NO:55	SEQ.I.D.NO:56
CA8 J7 Humanised heavy chain	SEQ.I.D.NO:57	SEQ.ID.NO:S8
CA8 J8 Humanised heavy chain	SEQ.I.D.NO:59	SEQ.ID.NO:60
CA8 J9 Humanised heavy chain	SEQ.I.D.NO:61	SEQ.I.D.NO:62
CA8 M0 Humanised light chain	SEQ.I.D.NO:63	SEQ.I.D.NO:64
CA8 M1 Humanised light chain	SEQ.I.D.NO:65	SEQ.ID.NO:66
CA8 M2 Humanised light chain	SEQ.I.D.NO:67	SEQ.ID.NO:68
S307118G03 V _H domain (murine)	SEQ.ID.NO:69	SEQ.ID.NO:70
S307118G03 V _L domain (murine)	SEQ.ID.NO:71	SEQ.I.D.NO:72
S307118G03 heavy chain (chimeric)	SEQ.I.D.NO:73	SEQ.ID.NO:74
S307118G03 light chain(chimeric)	SEQ.I.D.NO:75	SEQ.I.D.NO:76
S307118G03 Humanised V _H H0	SEQ.ID.NO:77	SEQ.ID.NO:78
S307118G03 Humanised V _H H1	SEQ.ID.NO:79	SEQ.I.D.NO:80
S307118G03 humanised V _H H2	SEQ.ID.NO:81	SEQ.I.D.NO:82
S307118G03 humanised V _H H3	SEQ.ID.NO:83	SEQ.I.D.NO:84
S307118G03 humanised V _H H4	SEQ.ID.NO:85	SEQ.I.D.NO:86
S307118G03 humanised V _H H5	SEQ.ID.NO:87	SEQ.I.D.NO:88
S307118G03 humanised V _L L0	SEQ.I.D.NO:89	SEQ.I.D.NO:90
S307118G03 humanised V _L L1	SEQ.I.D.NO:91	SEQ.I.D.NO:92
S307118G03 CDRH1	SEQ.ID.NO:93	
S307118G03 CDRH2	SEQ.ID.NO:94	
S307118G03 CDRH3	SEQ.I.D.NO:95	
S307118G03 CDRL1	SEQ.ID.NO:96	
S307118G03 CDRL2	SEQ.ID.NO:97	
S307118G03 CDRL3	SEQ.ID.NO:98	
S307118G03 humanised H5 CDRH3	SEQ.ID.NO:99	
S307118G03 H0 Humanised heavy chain	SEQ.I.D.NO:100	SEQ.I.D.NO:101
S307118G03 H1 humanised heavy chain	SEQ.I.D.NO:102	SEQ.ID.NO:103
S307118G03 H2 humanised heavy chain	SEQ.I.D.NO:104	SEQ.I.D.NO:105
S307118G03 H3 humanised heavy chain	SEQ.I.D.NO:106	SEQ.I.D.NO:107
S307118G03 H4 humanised heavy chain	SEQ.I.D.NO:108	SEQ.I.D.NO:109
S307118G03 H5 humanised heavy chain	SEQ.I.D.NO:110	SEQ.ID.NO:111
S307118G03 L0 humanised light chain	SEQ.I.D.NO:112	SEQ.ID.NO:113
S307118G03 L1 humanised light chain	SEQ.I.D.NO:114	SEQ.ID.NO:115
S332121F02 murine variable heavy chain	SEQ.I.D.NO:116	SEQ.ID.NO:117
S332121F02 chimeric variable heavy chain	SEQ.I.D.NO:118	SEQ.ID.NO:119
S332121F02 murine variable light chain	SEQ.I.D.NO:120	SEQ.ID.NO:121
S332121F02 chimeric variable light chain	SEQ.I.D.NO:122	SEQ.ID.NO:123
S322110D07 murine variable heavy chain	SEQ.I.D.NO:124	SEQ.ID.NO:125
S322110D07 chimeric heavy chain	SEQ.I.D.NO:126	SEQ.I.D.NO:127
S322110D07 murine variable light chain	SEQ.I.D.NO:128	SEQ.I.D.NO:129
S322110D07 chimeric light chain	SEQ.I.D.NO:130	SEQ.ID.NO:131
S332126E04 murine variable heavy chain	SEQ.I.D.NO:132	SEQ.ID.NO:133
S332126E04 Chimeric heavy chain	SEQ.I.D.NO:134	SEQ.I.D.NO:135
S332126E04 murine variable light chain	SEQ.I.D.NO:136	SEQ.ID.NO:137
S332126E04 Chimeric light chain	SEQ.I.D.NO:138	SEQ.ID.NO:139
S336105A07 murine variable heavy chain	SEQ.I.D.NO:140	SEQ.ID.NO:141

Description	Amino acid sequence	Polynucleotide sequence
S336105A07 Chimeric heavy chain	SEQ.I.D.NO:142	SEQ.ID.NO:143
S336105A07 murine variable light chain	SEQ.I.D.NO:144	SEQ.I.D.NO:145
S336105A07 chimeric light chain	SEQ.I.D.NO:146	SEQ.ID.NO:147
S335115G01 murine variable heavy chain	SEQ.I.D.NO:148	SEQ.I.D.NO:149
S335115G01 Chimeric heavy chain	SEQ.I.D.NO:150	SEQ.I.D.NO:151
S335115G01 murine variable light chain	SEQ.I.D.NO:152	SEQ.I.D.NO:153
S335115G01 Chimeric light chain	SEQ.I.D.NO:154	SEQ.I.D.NO:155
S335122F05 murine variable heavy chain	SEQ.I.D.NO:156	SEQ.I.D.NO:158
S335122F05 Chimeric heavy chain	SEQ.I.D.NO:158	SEQ.I.D.NO:159
S335122F05 murine variable light chain	SEQ.I.D.NO:160	SEQ.ID.NO:161
S335122F05 Chimeric light chain	SEQ.I.D.NO:162	SEQ.ID.NO:163
S332121F02 CDRH1	SEQ.I.D.NO: 164	
S332121F02 CDRH2	SEQ.I.D.NO: 165	
S332121F02 CDRH3	SEQ.I.D.NO: 166	
S332121F02 CDRL1	SEQ.I.D.NO: 167	
S332121F02 CDRL2	SEQ.I.D.NO: 168	
S332121F02 CDRL3	SEQ.I.D.NO: 169	
S322110D07 CDRH1	SEQ.I.D.NO: 170	
S322110D07 CDRH2	SEQ.I.D.NO: 171	
S322110D07 CDRH3	SEQ.I.D.NO: 172	
S322110D07CDRL1	SEQ.I.D.NO: 173	
S322110D07 CDRL2	SEQ.I.D.NO: 174	
S322110D07 CDRL3	SEQ.I.D.NO: 175	
S332126E04CDRH1	SEQ.I.D.NO: 176	
S332126E04 CDRH2	SEQ.I.D.NO: 177	
S332126E04 CDRH3	SEQ.I.D.NO: 178	
S332126E04 CDRL1	SEQ.I.D.NO: 179	
S332126E04 CDRL2	SEQ.I.D.NO: 180	
S332126E04 CDRL3	SEQ.I.D.NO: 181	
S336105A07 CDRH1	SEQ.I.D.NO: 182	
S336105A07 CDRH2	SEQ.I.D.NO: 183	
S336105A07 CDRH3	SEQ.I.D.NO: 184	
S336105A07 CDRL1	SEQ.I.D.NO: 185	
S336105A07 CDRL2	SEQ.I.D.NO: 186	
S336105A07 CDRL3	SEQ.I.D.NO: 187	
S335115G01 CDRH1	SEQ.I.D.NO: 188	
S335115G01 CDRH2	SEQ.I.D.NO: 189	
S335115G01 CDRH3	SEQ.I.D.NO: 190	
S335115G01 CDRL1	SEQ.I.D.NO: 191	
S335115G01 CDRL2	SEQ.I.D.NO: 192	
S335115G01 CDRL3	SEQ.I.D.NO: 193	
S335122F05 CDRH1	SEQ.I.D.NO: 194	
S335122F05 CDRH2	SEQ.I.D.NO: 195	
S335122F05 CDRH3	SEQ.I.D.NO: 196	
S335122F05 CDRL1	SEQ.I.D.NO: 197	
S335122F05 CDRL2	SEQ.I.D.NO: 198	
S335122F05 CDRL3	SEQ.I.D.NO: 199	

SEQUENCE LISTING

[0257]

SEQ ID 1 - CA8 CDRH1

NYWMH

SEQ ID 2 - CA8 CDRH2
ATYRGHSDTYYNQKFKG

SEQ ID 3 - CA8 CDRH3
GAIYNGYDVLDN

SEQ ID 4 - CA8 CDRL1
SASQDISNYLN

SEQ ID 5 - CA8 CDRL2
YTSNLHS

SEQ ID 6 - CA8 CDRL3
QQYRKLPWT

SEQ ID 7 - CA8 V_H domain (murine)
EVQLQQSGAVLARPGASVKMSCKGSGYTFNTYWMHWVKQRPGQGLEWIGATYRGHSDTYYNQF
KGKAALKTAVTSTSTAYMELSLTNEDSAVYYCTRGAIYNGYDVLDNWGQGTLTVSS

SEQ ID 8 - CA8 V_H domain (murine) (Polynucleotide)
GAGGTGCAGCTGCAGCAGAGCGGCCGCGTGGCTGGCCAGGCCGGAGCTAGCGTGAAGATGAG
CTGCAAGGGCAGCGGCTACACCTTCACCAACTCTGGATGCACTGGGTGAAACAGAGGCCCGG
CCAGGGACTGGAGTGGATCGGCCGACCTACAGGGGCCACAGCGACACCTACTAACACAGAA
GTTCAAGGGCAAGGCAAGCTGACCGCCGCTGACCTCAACCAGCACCGCCCTACATGGAACGTGAG
CAGCCTGACCAACGAGGACAGCGCCGTTACTACTGACCCAGGGGCCCATCTAACACGGCTA
CGACGTGCTGGACATTGGGCCAGGGAACACTAGTGACCGTGTCCAGC

SEQ ID 9 - CA8 V_L domain (murine)
DIQLTQTTSSLSASLGDRTIISCSASQDISNYLNWYQQKPDGTVELVIYTSNLHSGVPSRFSGS
TDYSLTIGYLEPEDVATYYCQQYRKLPWTFGGGSKLEIKR

SEQ ID 10 - CA8 V_L domain (murine) (Polynucleotide)
GATATCAGCTGACCCAGACCAACAGCAGCCTGAGCGCCTCCCTGGGCACAGGGTACCAATT
AGCTGCAGCGCAGGCCAGGACATCAGCAACTACCTGAACCTGGTACCGCAGAACGCCGACGGC
ACCGTGGAGCTGATCTACTACACCTCCAACCTGCACAGCGCCGTGCCCAGCGAGGTTCTCG

GCAGCGGCAGCGGCCACCGACTACAGCCTGACCATCGGCTATCTGGAGCCCGAGGACGTCGCA
CCTACTACTGCCAGCAGTACAGGAAGCTGCCCTGGACCTTCGGCGAGGCTCTAAGCTGGAGA
TTAACCGT

SEQ ID 11 - CA8 Humanised V_H J0
QVQLVQSGAEVKKPSSVKVSCKASGYTFNTYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSLRSEDTAVYYCARGAIYNGYDVLDNWGQGTLTVSS

SEQ ID 12 - CA8 Humanised V_H J0 (Polynucleotide)
CAGGTGCAGCTGGTCCAGAGCGGCCGCGAAGTGAAGAAGGCCGGCAGCTCCGTGAAGATGAG
CTGCAAGGGCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCG
GACAGGGCTGGAGTGGATGGGCCACCTACAGGGGCCACAGCGACACCTACTAACACAGAA
AGTTCAAGGGCGGGTGACCATCACCCTGACAAAGAGCACCGCCTACATGGAACGTGA
GCACCTCAGGAGCGAGGACACCGCTGTGTTACTGCACCCAGGGGCCATCTAACACGGCTA
ACGACGTGCTGGACAACTGGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 13 - CA8 Humanised V_H J1
QVQLVQSGAEVKKPSSVKVSCKASGYTFNTYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSLRSEDTAVYYCARGAIYNGYDVLDNWGQGTLTVSS

SEQ ID 14 - CA8 Humanised V_H J1 (Polynucleotide)
CAGGTGCAGCTGGTCCAGAGCGGCCGCGAAGTGAAGAAGGCCGGCAGCTCCGTGAAGATGAG
CTGCAAGGGCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCG
ACAGGGCTGGAGTGGATGGGCCACCTACAGGGGCCACAGCGACACCTACTAACACAGAA
GTTCAAGGGCGGGTGACCATCACCCTGACAAAGAGCACCGCCTACATGGAACGTGAG
CACCTCAGGAGCGAGGACACCGCTGTGTTACTGCACCCAGGGGCCATCTAACACGGCTA
CGACGTGCTGGACAACTGGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 15 - CA8 Humanised V_H J2
QVQLVQSGAEVKKPSSVKVSCKASGYTFNTYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSLRSEDTAVYYCARGAIYNGYDVLDNWGQGTLTVSS

SEQ ID 16 - CA8 Humanised V_H J2 (Polynucleotide)
CAGGTGCAGCTGGTCCAGAGCGGCCGCGAAGTGAAGAAGGCCGGCAGCTCCGTGAAGATGAG
CTGCAAGGGCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCG
ACAGGGCTGGAGTGGATGGGCCACCTACAGGGGCCACAGCGACACCTACTAACACAGAA
GTTCAAGGGCGGGTGACCATCACCCTGACAAAGAGCACCGCCTACATGGAACGTGAG
CACCTCAGGAGCGAGGACACCGCTGTGTTACTGCACCCAGGGGCCATCTAACACGGCTA
CGACGTGCTGGACAACTGGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 17 - CA8 Humanised V_H J3
QVQLVQSGAEVKKPSSVKVSCKASGYTFNTYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSLRSEDTAVYYCARGAIYNGYDVLDNWGQGTLTVSS

SEQ ID 18 - CA8 Humanised V_H J3 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGGCAGCGGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGCAGCACAGCAGCACCGCTACATGGAACGTAG
CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACAACGGCTA
CGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 19 - CA8 Humanised V_H J4

QVQLVQSGAEVKPGSSVKVSKGSYFTFTNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
KGRATLTADTSTSTAYMELSSLRSEDTAVYYCTRGAIYNGYDVLNDNWQGTLTVSS

SEQ ID 20 - CA8 Humanised V_H J4 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGGCAGCGGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGCAGCACAGCAGCACCGCTACATGGAACGTAG
CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACAACGGCTA
CGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 21 - CA8 Humanised V_H J5

QVQLVQSGAEVKPGSSVKVSKGSYFTFTNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
FKGRVTITADTSTSTAYMELSSLRSEDTAVYYCTRGAIYDGYDVLNDNWQGTLTVSS

SEQ ID 22 - CA8 Humanised V_H J5 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGGCAGCGGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGCAGCACAGCAGCACCGCTACATGGAACGTAG
CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACGACGGCTA
CGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 23 - CA8 Humanised V_H J6

QVQLVQSGAEVKPGSSVKVSKASGGTFSNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARQAYDGYDVLNDNWQGTLTVSS

SEQ ID 24 - CA8 Humanised V_H J6 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGCCAGCGGCCACCTTCAGCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
GACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
AGTCAAGGGCCGGGTGACCATCACCGCGACAGAGCACAGCAGCACCGCTACATGGAACGTAG
GCAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACGACGGCTA
ACGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 25 - CA8 Humanised V_H J7

QVQLVQSGAEVKPGSSVKVSKASGYFTFTNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARQAYDGYDVLNDNWQGTLTVSS

SEQ ID 26 - CA8 Humanised V_H J7 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGCCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGAGCACAGCAGCACCGCTACATGGAACGTAG
CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACGACGGCTA
CGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 27 - CA8 Humanised V_H J8

QVQLVQSGAEVKPGSSVKVSKASGYFTFTNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCTRGAIYDGYDVLNDNWQGTLTVSS

SEQ ID 28 - CA8 Humanised V_H J8 (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGCCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGAGCACAGCAGCACCGCTACATGGAACGTAG
CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCACCAGGGGCCATCTACGACGGCTA
CGACGTGCTGGACAACTGGGGCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 29 - CA8 Humanised V_H J9

QVQLVQSGAEVKPGSSVKVSKGSYFTFTNYWMHWVRQAPGQGLEWIGATYRGHSDTYYNQFK
KGRATLTADTSTSTAYMELSSLRSEDTAVYYCTRGAIYDGYDVLNDNWQGTLTVSS

SEQ ID 30 - CA8 Humanised V_H J9 Polnucleotide

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAAGAAGCCGGCAGCTCGTAAAGTGAG
CTGCAAGGCCAGCGGCCACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGCCCGG
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
GTTCAAGGGCCGGGTGACCATCACCGCGACAGAGCACAGCAGCACCGCTACATGGAACGTAG

CAGCCTCAGGAGCGAGGACACCGCTGTATTACTGTCACCAGGGGCGCCATCTACGACGGCTA
CGACGTGCTGGACAACCTGGGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 31 - CA8 Humanised V_L M0
DIQMTQSPSSLSASVGDRVTITCSASQDISQNLNWYQQKPGKAPKLIYYSNLSLHSGVPSRFSGSGS
GTDFTLTISLQPEDFATYYCQQYRKLPWTFGQGTKLEIKR

SEQ ID 32 - CA8 Humanised V_L M0 (Polynucleotide)
GACATCCAGATGACCCAGAGGCCCTAGCTCACTGAGGCCAGCGTGGGCGACAGGGTGACCTT
ACCTGCTTCCCCAGCCAGGACATCAGCAACTCTGAACCTGGTACCCAGCAGAACCCGGGCAAG
GCCCCCAAGCTGCTGATCTACTACACCTTCAACCTGCACCTGGCGTGGCCCAAGGTTCAAGCG
GAAGCGGCAAGGGCACCGATTTCACCTGACCATCTCAGCGTGCAGCGGAGGACTTCGCCA
CTTCACTACTGCGCACAGCAGTACAGGAAGCTCCCTGGACTTTCGGCCAGGGCACCAACTGGAGAT
CAAGCGT

SEQ ID 33 - CA8 Humanised V_L M1
DIQMTQSPSSLSASVGDRVITTCASQDINSYLNWYQQKPGKAPKLIYTSNLHSGVPSRFGSGS
GTDYTLTISLQPEDFATYYCQQYRKLPWTFGQGTKLEIKR

SEQ ID 34 - CA8 Humanised V_L M1 (Polynucleotide)
GACATCCAGATGACCCAGAGCCCTAGCTACTGAGGCCAGCGTGGGCGACAGGGTGACCTT
ACCTGCTTCCCCAGCCAGGGCATCAGCAACTCTGAACCTGGTACCCAGCAGAACGCCGGCAAG
GCCCTTCAAGCTGTGATCTACACCTCCACCTGCACCTGGCGTGGCCCAAGGTTCAAGC
GAAGGCCAGGGCAGCATTACACCTGACCATCTCAGCCTGCAAGCCCAAGGACTTCGCCA
CCTACTACTGTCAGCAGTACAGGAAGCTCCCTGGACTTTCGGCCAGGGCACCAAACTGGAGAT
CAAGCGT

SEQ ID 35 - CA8 Humanised V_L M2
DIQLTQSPSSLASVGDRVITICCSAQSDISNYLNWYQQKPGKAPELVYYTSNLHSGVPSRFSGSGSC
TDYTLTISSLQQPEDFATYYCQQYRKLPWTFGQGTKLEIKR

SEQ ID 36 - CA8 Humanised V_L M2 (Polynucleotide)
GACATCCAGCTGACCAGGCCAGCTAGCTACTGAGGCCAGCGTGGCGACAGGGTGACCTT
ACCTGCTTCCCAGCCAGGACATCAGCACTACCTGAACTGTGACCCAGGAGCCCGGCAAG
GCCCGCAGCTGGTGTACTACACTCCCAACCTGCACTCCGGCAGTGGCCAGGTTTCAGC
GGAAGGCGCCAGGGCACCGATTACACCCCTGACCATCTCCAGCTGAGCCGAGGACTTCGCC
ACCTACTACTGCCAGCAGTACAGGAAGCTCCCTGGACTTTCGGCCAGGGACCAAAGTGGAG
TCAAGCGT

SEQ ID 37 - Human BCMA CD33-hBCMA ECD (1-53) TEV-Fc
MPLLLLPLWAGALAMLQMGQCSQNEYDSSLHACIPCQLRCSSNTPPLTCQRYCNAVSNTVSVKG
TNSENGLYFQGDPKSCDKTHCTPPCQPAPELLGGPSVLFPPPKDSDLMISRTPVEVTCVVVDVSHEDP
EVKFNWYVVDGVEVHNAKTTPKEEPCQYNSTYRVSVLTVLHQDWLKEYKCKVSNKALPAPIEKTSK
AKQKPPREPVQYTLPPSRDELTKNQVSLTCLVKGFPYPSDIAWEWESNGQPNENNYKTPPVLDSDGSFF
LYSKLTVDSRWRQGQNVFSCSMHEALHNHYTQKSLSSPQGK

SEQ ID 38 - Human BCMA CD33-hBCMA ECD (1-53) TEV-Fc (Polynucleotide)
ATGCCGCTGCTGACTCTGCTGCCCTGCTGGGGCAGGGCGCTAGCTATGCTGCAGATGGCC
GGCCAGTCAGCAGCAGAACGGAGTACTTCGACAGCCTGCTGACGCCGTCATCCCTGCCAGCTG
AGATGAGCAGCAGCAACACACCTCTGACCTCCGAGAAGATCTGCAACGCCAGCTGACCAACA
GCTGGAAGGGACCAACTCCGGAGAGAACCTCTTCAAGGGGATCCCCAAATCTTGTGACAA
AACTCACATGCCCCAGCTGGCCAGCACCTGAACTCTGGGGGAGCCGTGAGCTCTCCCTTC
CCCCCAAAACCAAGGACACCCCTATGATCTCCGGACCCCTGAGGTACATGGCTGGTGGTGGT
GACGTGAGCCACGAAGACCCCTGAGGTCAAGTCAACTGGTACCTGGGACGGCTGGAGGTGCA
AATGCCAAGACAAGCCGGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCTCC
ACCGTCTGACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCAACAAAGCC
TCCCAGCCCCATGAGAAAACCTCTCCAAAGCCAAGGGCAGCCCCGAGACCCACAGGTG
ACACCTCTGCCCCCATCCGGAGTGAAGCTGCAAGAACCCAGGTGACCTGCTGACCTGGTCA
AAGGCTTCTATCCCGACGACATCCGGTGGAGGAGACAGAATGGGAGCCGGAGAACAACT
ACAAAGACCCCTCCGGTGGACTCCGAGCAGGGCTCTTCTCTACAGCAAGCTCACCGT
GGACAAGGAGCAGGTGGCAGCAGGGGAACGCTTCTCATGCTCGTGTGATGCATGAGGCTCTGCA
CAACCACTACAGCAGAACAGGCTCTCCGTCTGGTAA

SEQ ID 39- Human BCMA CD33-hBCMA ECD (4-53) TEV-Fc
MPLLLLLPLWAGALAMAGQCSQNEYFDSSLHACIPCQLRCSSNTPLTCQRYCNASVTSVKGTNS
GENLYFQGDPKSCDKTHCPCPAPAEELLGGPSVFLFPPPKPDLMISRTRPEVTVVWVDSHDEPEVK
FNWYVVGVEVHNAKTTPREEQYNSNTYRVSVLTVLHQDWLNGKEYKCVSNKALPAPIEKTSKAKG
QPRREPQVYTLPLPSRDELTKNQSVLCLVKGKFYPSDIAWEWNSQGPNENNYKTTPPVLDSDGSFFLYS
K1/TD/KDSRWOOGNIVESCSVWMEAI HNHYTOKSI SISPGK

SEQ ID 40 - Human BCMA CD33-hBCMA ECD (4-53) TEV-Fc (Polynucleotide)
ATGCCGCTGCTGACTGCTGCCCTGCTGGGGCAGGGCGCTAGCTATGGCGGCCAGTGC
AGCCAGAACGAGTACTCGACAGCCCTGCTCACCCCTGCATCCTGGCAGCTGAGATGCAGC
AGCAACACACTCTCTGAGCTGCCAGAGATCTGCAACGCCAGCGTGACCAACAGCGTGAAGG
GCACCAACTCGGAGAGAACCTGTACTTCAAGGGATCCAAATCTTGACAAAAACTCACAC
ATGCCAACCGTGCCCCAGCACCTGAACTCCTGGGGGACCGTCAGTCTTCTCTTCCCCCCCCAAA
CCCAAGGACACCCCTCATGATCTCCGGACCCCTGAGGTCACATGGCTGGTGGGGACGTCAGC

UHUGAAGGAAUUCUUGGGTCAAGTTCAGUUGGTTGGTACGGTGGAUUGGUUGTGGAGGGTGUATTAATGUUAG
ACAAAGCCGGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCTCACCGTCTG
CACCAAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAGGTCTCCAAACAAAGCCCTCCAGCC
CCCATCGAGAAAACCATCTCAAAGCCAAGGGCAGCCCGAGAGGCCACAGGTGTACACCCCTG
CCCCCATCCGGATGAGCTGACCAAGAACAGGTAGCCTGCACCTGCTCAAAGGCTCT
ATCCCAGCAGCACATGCCGTGGAGTGGAGAGCAATGGCAGCCGAGAACAACTACAAGACCA
CGCCTCCGTGCTGGACTCGACGGCTCTCTCTACAGCAAGCTACCGTGGACAAGAG
CAGGTGGCAGCAGGGAACGTCTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCAACTAC
ACCCAGAAGAGCCTCTCCGTCTCCGGTAA

SEQ ID 41- Cynomolgous BCMA CD33 cyto BCMA ECD (4-52) TEV-Fc
MPLLLLLLPLWAGALAMARQCSQNEYFDSSLHDCKPCQLRCSSPPLTCQRYCNASMTNSVKGMNS
GENLYFQGDPKSCDKTHTCPPCPAPEELLGGPSVFLFPKPKDTLMISRTPEVTCVVVDVSHEDPEVK
FNWYVVDGVEVHNAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAIEKTISAKG
QPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYS
KLTVDKSRWQQGNVFSCSVVMHEALHNHYTQKSLSLSPGK

SEQ ID 42 - Cynomolgous BCMA CD33 cyto BCMA ECD (4-52) TEV-Fc (Polynucleotide)
ATGCCGCTGCTGCTACTGCTGCCCTGCTGGGCAGGGCGCTAGCTATGGCCAGACAGTGC
AGCCAGAACGAGTACTTCGACAGCCTGCTGCACGACTGCAAGCCCTGCCAGCTGAGATGCAGC
AGCACACCTCTCTGACCTGCCAGAGATACTGCAACGCCAGCATGACCAACAGCGTAAGGGCA
TGAACCTCGGAGAGAACCTGTACTTCAAGGGGATCCCAAATCTTGTACAAAAACTCACACATGC
CCACCGTGGCCAGCACCTGAACTCTGGGGGGACCGTCACTTCTCTCTCCCTCCAAAACCCA
AGGACACCCCTCATGATCTCCCGACCCCTGAGGTACATGCGTGGTGGACGTGAGCCACG
AAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGATAATGCCAAGACAA
GCCCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTACGGTCTCTACCGTCTGCACCA
GGACTGCTGAATGCCAAGGAGTACAAGTGCAGGCTTCAACAAAGCCCTCCAGCCCCCATC
GAGAAAACATCTCAAAGGCAAGGGCAGCCCCAGAGGCCACAGGTGTACACCCCTGCCCCA
TCCCGGGATGAGCTGACCAAGAACAGGTCACTGCCCTGGCTGCTGCTGAAAGGCTTATCCA
GGACATGCCGTGGAGTGGAGGAGCAATGGCCAGGGAGAACAACTACAAGACACGCC
CCGTGCTGGACTCCGACGGCTCTCTCTACAGCAAGCTACCGTGGACAAGAGCAGGTG
GCAGCAGGGAAAGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCAACTACACCGAG
AAGAGCCTCTCCCTGTCTCCGGTAA

SEQ ID 43- CA8 J0 Humanised heavy chain
QVQLVQSGAEVKKPSSVKVSKASGGTFSNYWMHWRQAPGQGLEWMGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARGAIYNGYDVLNDNWQGTLTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVWSNSGALTSGVHTFPVQLQSSGLYSLSSVTVPSLSS

GTQYICNVNHPSTNTVKVDPKSCDKTHTCPPCPAPEELLGGPSVFLFPKPKDTLMISRTPEVTC
VVVDVSHEDPEVKFNWYVVDGVEVHNAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
ALPAIEKTISAKGQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVVMHEALHNHYTQKSLSLSPGK

SEQ ID 44 - CA8 J0 Humanised heavy chain (Polynucleotide)
CAGGTGCAGCTGGCCAGGGCGCCAGACTGCAACTACTGGATGCACTGGGTGAGGCAGGCC
GACAGGGCTGGAGTGGATGGCGCCACCTACAGGGCCACAGCACACCTACTACAACAGAG
AGTTCAAGGGCCGGTGAACCATACCGCCGACAAGAGCACAGCACGCCCTACATGGAAC
GCAGCCTCAGGAGCGAGGACCCCTGTGTATTACTGCGCCAGGGGCCATCTACAACGGCT
ACGACGTGCTGGACAACACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCCAGCAC
GCCCGAGCGTCTCCCTGGCCCCAGCAGCAAGAGCACAGCGCCAGCGCC
GGCTGCTGGTGAAGGACTACTCCCCAACCGGTGACCGTGTGGAACAGCGGAGCC
ACACAGCGCGTGCACACCTCCCCGCCGTGCTGAGAGCACAGCGCC
GTGCTGGAGCTGGCCAGCGAGCTGGCACAGGCCACAGCTACATGTAACGTGAAC
CCAGCAACACCAAGGTGACAAGAAGGTGGAGCCAAGAGCTGTGACAAGACCCAC
CCCCCTCTGGCTGGCCAGCTGCTGGAGGCCAGCGTGTGTTCTGTTCCCCCAAGCCT
AAGGACACCCCTGATGATCAGCAGAACCCCCAGGGTGAACCTGTGTTGGTGGATGTGAG
GAGGACCCCTGAGGTGAAGTCACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAG
AACGCCAGGGAGGAGCAGTACAACAGCACCTACGGCGTGTGCTGCTGACCGTGT
CAGGATGGCTGAACGGCAAGGGTACAAGTGTGAAGGTGTCACAAAGGCCCTGCC
TCGAGAAAACCATCAGCAAGGCCAGGGCAGCCAGAGAGGCCAGGTGACACCTGCC
CTAGCAGAGATGAGCTGACCAAGAACAGGTGTCCCTGACCTGCCCTGGTAAGGGCTTACCC
CAGCGACATGCCGTGGAGTGGAGAGCAACGCCAGGCCAGGAACAACTACAAGACCA
CCCTGTGCTGGACACGGATGGCACTTCTCTGTACAGCAAGCTACCGTGGACAAGAG
TGCAGCAGGGCAACGTGTGAGCTGCTCCGTGATGCACGAGGCCCTGCACAATCA
CAGAAGAGCCTGAGCCTGCTCCCTGGCAAG

SEQ ID 45- CA8 J1 Humanised heavy chain
QVQLVQSGAEVKKPSSVKVSKASGYTFTNYWMHWRQAPGQGLEWMGATYRGHSDTYYNQK
FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARGAIYNGYDVLNDNWQGTLTVSSASTKGPSVF
PLAPSSKSTSGGTAALGCLVKDYFPEPVTVWSNSGALTSGVHTFPVQLQSSGLYSLSSVTVPSLSS
GTQYICNVNHPSTNTVKVDPKSCDKTHTCPPCPAPEELLGGPSVFLFPKPKDTLMISRTPEVTC
VVVDVSHEDPEVKFNWYVVDGVEVHNAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
ALPAIEKTISAKGQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVVMHEALHNHYTQKSLSLSPGK

SEQ ID 46 - CA8 J1 Humanised heavy chain (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAGAGAAGCCGGAGCTCCGTGAAAGTGAG
 CTGCAAGGCCAGCGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGGCCCCGG
 ACAGGGCCTGGAGTGGATGGCGGCCACCTACAGGGGCCACAGCAGCACCTACTACAAACAGAA
 GTTCAAGGGCCGGGTGACCATCACCGCGACAAGAGCACCAGCAGCACCGCTACATGGAACGTAG
 CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCGCCAGGGCGCATCTACAACGGCTA
 CGACGTGCTGGACAACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCCAGCACCAAGG
 GCCCCAGCGTGTCCCCCTGGCCCGCAGCAGCACAGCAGCGGCCACAGCGCCCTG
 GGCTGCGCTGGTAAGGACTACTTCCCAGAACCCTGGTACAGCGGCCGTACAGCGTGA
 ACCAGCGCGTGCACACCTCCCCGCCGTGTGAGAGCAGCGGCCGTACAGCGTGA
 GTGGTACCGTGCACAGCAGCGTGTGAGAGCAGCTGAGAGCAGCGGCCGTACAGCGTGA
 CCCAGCAACACCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCTGTGACAAGACCCACACTGC
 CCCCCCTGCCCTGCCCGAGCTCTGGAGGCCAGCGTGTCTGTGTTCCCCCAAGCCT
 AAGGACACCCGTATGATCAGCGAGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
 GAGGACCCCTGAGGTGAAAGTCACTGGTACGTGACGGCGTGGAGGTGACAATGCAAGACC
 AAGCCCAGGGAGGAGCAGTACAACAGCACCTACCGGGTGGTGTCCGTGACCGTGTGAC
 CAGGATTGGCTGAACGGCAAGGAGTACAAGTGTAAGGTGTCACAAAGGCCCTGCTGCC
 TCGAGAAAACCATCAGCAAGGCCAGGGCAGCCAGAGAGGCCAGGTGTACACCCCTGCC
 CTAGCAGAGATGAGCTGACCAAGAACAGGTGTCCCTGACCTGCGCTGGTGAAGGCC
 CACCGACATCCCCGTGGAGTGGAGAGCAACGCCAGCCAGAACACTACAAGACCAC
 CCCTGTGCTGGACACCGATGGCACCTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGA
 TGGCAGCAGGGCAACGTGTTCACTGCTGCCGTGATGCAAGGCCCTGACAATCACTACACC
 CAGAAGAGCCTGAGCCTGTCCCCCTGGCAAG

SEQ ID 47 - CA8 J2 Humanised heavy chain

QVQLVQSGAEVKKPGSSVKVKSGYFTFTNWMHWVRQAPGQGLEWMGATYRGHSDTYYNQK
 FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCTRGAIYNGYDVLNDWQGTLTVSSASTKGPSVF
 PLAPSSKSTSGGTAALGCLVKDVFPEPVTSWNSGALTSGVHTFPALVLQSSGLYLSVVTPVSSL
 GTQTYICNVNHPKSNVKDKVKEPKSCDKTHTCPPCPAPEELLGGPSVFLFPKPKDTLMSRTPVTC
 VVVDVSHEDPEVKFNWYDVGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
 ALPAPIEKTIKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
 PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVHEALHNHYTQKSLSLSPGK

SEQ ID 48 - CA8 J2 Humanised heavy chain (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAGAGAAGCCGGAGCTCCGTGAAAGTGAG
 CTGCAAGGCCAGCGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGGCCCCGG
 ACAGGGCCTGGAGTGGATGGCGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
 GTTCAAGGGCGGGTACCATCACCGCGACAAGAGCACAGCACCGCTACATGGAACGTAG
 CAGCCTCAGGAGCAGGACACCGCTGTATTACTGACCCAGGGGCCATCTACACGGCTA
 CGACGTGCTGGACAACTGGGCCAGGGCACACTAGTGACCGTGTGAGGCCAGCACCAAGG
 GCCCCAGCGTGTCCCCCTGGCCCGCAGCAGCAAGAGCACAGCGGCCAGCGGCCCTG
 GGCTGCTGGTGAAGGACTACTTCCCAGAACCCTGGTACCGTGTCTGGACAGCGAGGCC
 ACCAGCGCGTGACACCTTCCCAGCGTGTGAGAGCACAGCGGCCCTGACAGCGTGA
 GTGGTACCGTGCACAGCAGCGTGGACCCAGCTACATCTGTAACGTGAACCCACAAG
 CCCAGCAACACCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCTGTGACAAGACCCAC
 CCCCCCTGCCCTGCCCGAGCTGCTGGAGGCCAGCGTGTCTGGTACAGCGTGTGAC
 AAGGACACCCGTATGATCAGCGAGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCC
 GAGGACCCCTGAGGTGAAGTCAACTGGTACGTGAGCGCCGGTGGAGGTGACAATGCC
 AAGCCCAGGGAGGAGCAGTACAACAGCACCTACGGGTGGTGTGCTGACCGTGTGAC
 CAGGATTGGCTGAACGGCAAGGAGTACAAGTGTAAGGTGTCACAAAGGCCCTGCC
 TCAGAGAAAACCATCAGCAAGGCCAGGCCAGAGAGGCCAGGTGTACACCCCTGCC
 CTAGCAGAGATGAGCTGACCAAGAACCCAGGTGTCCCTGACCTGCTGGTAAGGGCTTCA
 CAGCGACATGCCCTGGAGTGGAGAGCAACGCCAGGGGAGAACACTACAAGACCAC
 CCCTGTGCTGGACACCGATGGCACCTCTCTGTACAGCAAGCTGACCGTGGACAAGCAGA
 TGGCAGCAGGGCAACCTGTTCACTGCTGCCGTGATGCAAGGCCCTGACAATCACTACACC
 CAGAAGAGCCTGAGCCTGTCCCCCTGGCAAG

SEQ ID 49- CA8 J3 Humanised heavy chain

QVQLVQSGAEVKKPGSSVKVKSGYFTFTNWMHWVRQAPGQGLEWMGATYRGHSDTYYNQK
 FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCTRGAIYNGYDVLNDWQGTLTVSSASTKGPSVF
 PLAPSSKSTSGGTAALGCLVKDVFPEPVTSWNSGALTSGVHTFPALVLQSSGLYLSVVTPVSSL
 GTQTYICNVNHPKSNVKDKVKEPKSCDKTHTCPPCPAPEELLGGPSVFLFPKPKDTLMSRTPVTC
 VVVDVSHEDPEVKFNWYDVGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
 ALPAPIEKTIKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
 PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVHEALHNHYTQKSLSLSPGK

SEQ ID 50 - CA8 J3 Humanised heavy chain (Polynucleotide)

CAGGTGCAGCTGGTCCAGAGCGGCCGAAGTGAGAGAAGCCGGAGCTCCGTGAAAGTGAG
 CTGCAAGGCCAGCGCTACACCTTCACCAACTACTGGATGCACTGGGTGAGGCAGGGCCCCGG
 ACAGGGCCTGGAGTGGATGGCGGCCACCTACAGGGGCCACAGCAGCACCTACTACAACAGAA
 GTTCAAGGGCGGGTACCATCACCGCGACAGCACAGCACCGCTACATGGAACGTAG
 CAGCCTCAGGAGCAGGAGCACCGCTGTATTACTGACCCAGGGGCCATCTACACGGCTA
 CGACGTGCTGGACAACTGGGCCAGGGCACACTAGTGACCGTGTGAGCGGCCAGCACCAAGG
 GCCCCAGCGTGTCCCCCTGGCCCGCAGCAGCAAGAGCACAGGCCAGGCCACAGCGCCCTG
 GGCTGCTGGTGAAGGACTACTTCCCAGAACCCTGGTACCGTGTCTGGACAGCGAGGCC

ACCAAGCGGGCGTGCACACCTCCCCGCCGTGCTGCAGAGCAGCGGGCTGTACAGCCTGAGCAGC
GTGGTGAACCGTGCACCGCAGCAGCAGCTGGCACCCAGACCTACATCTGTAACGTGAACCCAAG
CCCAAGCAACACCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCCTGTGACAAGACCCACACCTC
CCCCCTGCCCTGCCCTGCCAGCTCTGGAGGCCAGCGTGTCTGTGTTCCCCCAAGCCT
AAGGACACCCCTGATGATCAGCAGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
GAGGACCCCTGAGGTGAAGTCAACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAGACC

AAGCCCAGGGAGGAGCAGTACAACAGCACCTACCGGTGGTGTCCGTGCTGACCGTGTGAC
CAGGATTGGCTGAACGCCAGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCCCTGCC
TCAGAGAAAACCATCAGCAAGGCCAGGGCCAGCCAGAGGCCAGGTGTACACCCCTGCC
CTAGCAGAGATGAGCTGACCAAGAACCAGGTGTCCTGACCTGCCCTGGTAAGGGCTCTACCC
CAGCGACATGCCCTGGAGGTGGAGAGCAACGCCAGCCAGAGAAACAACTAAGACCAC
CCCTGTGCTGGACAGCAGTGGACCTTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGA
TGGCAGCAGGGCAACGTGTTCACTGCTCCGTGATGCAACGAGGCCCTGACAATCACTACACC
CAGAAGAGCCTGAGCCTGCCCCCTGGCAAG

SEQ ID 51 - CA8 J4 Humanised heavy chain
QVQLVQSGAEVKPGSSVKVSKGSYFTFTNWMHWRQAPGQGLEWIGATYRGHSDTYYNQF
KGRATLTADTSTSTAYMELSSLRSEDTAVYYCTRGAIYDGYDVLNDWQGTLTVSSASTKGPSVF
LAPLSSKSTSGGTAALGCLVKDYFPEPVTVSNWSALTSVHFTPAVLQSSGLYLSVVTPSSLG
TQTYICNVNHPKPSNTKVDKVKEPKSCDKTHTCPCCPAPELLGGPSVFLFPKPKDLMISRTPEVTC
VVVDVSHEDPEVKFNWYDGVVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKA
LPAPIEKTIKAKGQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTP
PVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPKG

SEQ ID 52 - CA8 J4 Humanised heavy chain (Polynucleotide)
CAGGTGCAGCTGGCCAGCGGGCGCGAAGTGAAGAAGGCCGGAGCTCCGTGAAAGTGAG
CTCGAAGGGCAGCGGCTACACCTTACCAACTCTGGATGCACTGGGTGAGGCCAGGCC
ACAGGGCCTGGAGGTGGATCGGCCACCTACAGGGGCCACAGCGCACCTACTACAACAGGAA
GTTCAAGGGCCGGCGACCCCTACCGCGACAGAGCACCGACCGCCTACATGAACTGAG
CAGCCTCAGGAGCAGGACACCCTGTGTTACTGCACCAAGGGCGCCATCTACAACGCGTA
CGAGGTGCTGGACAACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGG
GCCCGAGGTGTTCCCCCTGGCCCGACAGCAAGAGCACAGCGGCCAGCGCC
GGCTGCGCTGGTGAAGGACTACTCCCCGAACCGGTGACCGTGTCCGTGAAACAGCGGAGCC
ACAGCGGGCGTGCACACCTTCCCCCGCGTGTGCAAGAGCACGCCGTGACCGTGTGAGCAGC
GTGGTGAACCGTGCACGAGCTGGCACCCAGCCTACATCTGTAACGTGAACCC
CCAGCAACACCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCTGTGACAAGACCCAC
CCCCCTGCCCTGCCCGAGCTCTGGAGGCCAGCGTGTGTTCCCCCAAGCCT
AAGGACACCCCTGATGATCAGCAGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
GAGGACCCCTGAGGTGAAGTCAACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAGACC
AAGCCCAGGGAGGAGCAGTACAACAGCACCTACGGGTGGTGTCCGTGACCGTGTGAC
CAGGATTGGCTGAACGCCAGGGACACTAGTGACCGTGTCCAGCGCAGCACCAAGG
TCGAGAAAACCATCAGCAAGGCCAGGGCCAGCCAGAGGCCAGGTGTACACCCCTGCC
CTAGCAGAGATGAGCTGACCAAGAACCAGGTGTCCTGACCTGCCCTGGTAAGGGCTCTACCC
CAGCGACATGCCCTGGAGGTGGAGAGCAACGCCAGCCAGAGAAACAACTAAGACCAC
CCCTGTGCTGGACAGCAGTGGCAGCTTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGA
TGGCAGCAGGGCAACGTGTTCACTGCTCCGTGATGCAACGAGGCCCTGACAATCACTACACC
CAGAAGAGCCTGAGCCTGCCCCCTGGCAAG

SEQ ID 53 - CA8 J5 Humanised heavy chain
QVQLVQSGAEVKPGSSVKVSKGSYFTFTNWMHWRQAPGQGLEWIGATYRGHSDTYYNQK
FKGRVTITADTSTSTAYMELSSLRSEDTAVYYCTRGAIYDGYDVLNDWQGTLTVSSASTKGPSVF
LAPLSSKSTSGGTAALGCLVKDYFPEPVTVSNWSALTSVHFTPAVLQSSGLYLSVVTPSSLG
TQTYICNVNHPKPSNTKVDKVKEPKSCDKTHTCPCCPAPELLGGPSVFLFPKPKDLMISRTPEVTC
VVVDVSHEDPEVKFNWYDGVVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
LPAPIEKTIKAKGQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTP
PVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPKG

SEQ ID 54 - CA8 J5 Humanised heavy chain (Polynucleotide)
CAGGTGCAGCTGGCCAGCGGGCGCGAAGTGAAGAAGGCCGGAGCTCCGTGAAAGTGAG
CTCGAAGGGCAGCGGCTACACCTTACCAACTCTGGATGCACTGGGTGAGGCCAGGCC
ACAGGGCCTGGAGGTGGATGGGCCACCTACAGGGGCCACAGCGCACCCACTACAACAGGAA
GTTCAAGGGCCGGTGACCATCACCGCCACAGCAGCACAGCACCGCCTACATGAACTGAG
CAGCCTCAGGAGCAGGACACCCTGTGTTACTGCACCAAGGGCGCCATCTAGCACGGCTA
CGAGGTGCTGGACAACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGG
GCCCGAGGTGTTCCCCCTGGCCCGAGCGAAGAGCACAGCGGCCAGCGCC
GGCTGCGCTGGTGAAGGACTACTCCCCGAACCGGTGACCGTGTCCGTGAAACAGCGGAGCC
ACAGCGGGCGTGCACACCTTCCCCGCCGTGCTGCAGAGCACGCCGTGACCGTGTGAGCAGC
GTGGTGAACCGTGCACGAGCTGGCACCCAGCAGCTACATCTGTAACGTGAACCC
CCAGCAACACCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCTGTGACAAGACCCAC
CCCCCTGCCCTGCCCGAGCTGCTGGAGGCCAGCGTGTGTTCCCCCAAGCCT
AAGGACACCCCTGATGATCAGCAGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
GAGGACCCCTGAGGTGAAGTCAACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAGACC
AAGCCCAGGGAGGAGCAGTACAACAGCACCTACGGGTGGTGTCCGTGACCGTGTGAC
CAGGATTGGCTGAACGCCAGGGAGGTGACCTGTGTGGTGGATGTGAGCCAC

TCGAGAAAACCATCAGCAAGGCCAAGGGCCAGCCCAGAGAGCCCCAGGTGTACACCTGCCCC
 CTAGCAGAGATGAGCTGACCAAGAACCGGTGTCCTGACCTGCGCTGTGAAGGGCTTCTACCC
 CAGCGACATCGCCGTGGAGTGGGAGAGCAACGCCAGCCCCAGAAACAACACTACAAGACCACCC
 CCCTGTCTGGACACGGATGGCACTTCTCTGTACAGCAAGCTCACCGTGGACAAGAGCAGA
 TGCAGCAGGGCAACGTGTTCAGCTGCTCCGTATGCACGAGGCCCTGCACAATCACTACACC
 CAGAAGAGCCTGAGCCTGCCCCGGCAAG

SEQ ID 55 - CA8 J6 Humanised heavy chain
 QVQLVQSGAEVKPGSSVKVSKASGYTFTNYWMHWVRQAPGQGLEWMGATYRGHSDTYYNQK
 FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARGAIYDGYDVLNDWQGTLTVSSASTKGPSVF
 PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYLSLSVTVPSSSL

GTQTYICNVNHPNSNTVKDKVKEPKSCDKTHTCPCTCAPEELLGGPSVFLFPKPKDTLMISRTPEVCT
 VVVDVSHEDPEVKFNWYDGVVEHNAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
 ALPAPIEKTSKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
 PPVLDSDGSFFYLSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK

SEQ ID 56 - CA8 J6 Humanised heavy chain (Polynucleotide)
 CAGGTGAGCTGGCCAGAGCGGGCCGAAGTGAAGAAGGCCGGCAGCTCCGTAAAGTGAG
 CTGCAAGGCCAGCGCCGGCACCTCAGCAACTACTGGATGCACTGGGTGAGCAGGCCAGGGCCCG
 GACAGGGCTGGAGTGGATGGCCACCTACAGGGCCACAGCGACACCTACTACACCCAGA
 AGTCAAGGGCCGGGTGACCATCACCGCCGACAAGAGCAGCAGCACCGCCTACATGGAACCTGA
 GCAGCCTCAGGAGCAGGACACGGCTGTATTACTGCGCCAGGGCGCCATCTACGACGGCT
 ACCACGTGCTGACCAACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGG
 GCCCCAGGGTGTCCCCCTGGCCCCCAGCAGCAAGAGCACAGCGCCAGCGCCCTG
 GGCTGCGCTGGTAAGGACTACTCCCCGAACCGGTGACCGTGTCCCTGAAACAGCGAGCCCTG
 ACCAGCGCGTGCACACCTCCCCGCGTGTGACAGCAGCGCCCTGTACAGCCTGAGCAGC
 GTGGTGACCGTGCAGCAGCACCCAGGACACCTACATCTGAAACGCTGACAAGACCCACACCTG
 CCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCAAGAGCTGTGACAAGACCCACACCTG
 CCCCCCTGCCCTGCCCGAGCTGGAGGCCAGCGTGTCCCTGTTCCCCCAAGCCT
 AAGGACACCCCTGATGATCAGCGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
 GAGGACCCCTGAGGTGAAGTCAACTGGTACCGTGAAGCCGGCTGGAGGGTGCACATGCAAGACCC
 AAGCCCAGGGAGGAGCAGTACAACAGCACCTACGGGGTGGTGTGGTCCGTCAGCGTGCAC
 CAGGATTGGCTGAAGGCAAGGAGTACAAGTGTAAAGGTGCTAACACAGGCCCTGCCCT
 TCGAGAAAACCATCAGCAAGGCCAGGGCACAGGCCAGAGGCCAGGTGTACACCCCTGCCCT
 CTAGCAGAGATGAGCTGACCAAGAACGGTGTCCCTGACCTGCGCTGGTAAGGGCTTCTACCC
 CACCGACATCCCCGTGAGTGGAGAGCAACGCCAGGCCAGAGAACACTACAAGACCACCC
 CCTGTGCTGGACAGCAGTGGCAGCTTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGA
 TGGCAGCAGGGCAACGTGTTCAGCTGCTCCGTATGCACGAGGCCCTGCACAATCACTACACC
 CAGAAGAGCCTGAGCCTGCCCCGGCAAG

SEQ ID 57 - CA8 J7 Humanised heavy chain
 QVQLVQSGAEVKPGSSVKVSKASGYTFTNYWMHWVRQAPGQGLEWMGATYRGHSDTYYNQK
 FKGRVTITADKSTSTAYMELSSLRSEDTAVYYCARGAIYDGYDVLNDWQGTLTVSSASTKGPSVF
 PLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYLSLSVTVPSSSL
 GTQTYICNVNHPNSNTVKDKVKEPKSCDKTHTCPCTCAPEELLGGPSVFLFPKPKDTLMISRTPEVCT
 VVVDVSHEDPEVKFNWYDGVVEHNAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNK
 ALPAPIEKTSKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT
 PPVLDSDGSFFYLSKLTVDKSRWQQGNVFSCSVMEALHNHYTQKSLSLSPGK

SEQ ID 58 - CA8 J7 Humanised heavy chain (Polynucleotide)
 CAGGTGAGCTGGCCAGAGCGGGCCGAAGTGAAGAAGGCCGGCAGCTCCGTAAAGTGAG
 CTGCAAGGCCAGCGCCGGTACACCTCAGCAACTACTGGATGCACTGGGTGAGGAGGCCAGGGCCCG
 ACAGGGCTGGAGTGGATGGCCACCTACAGGGCCACAGCGACACCTACTACACCCAGAA
 GTTCAAGGGCCGGGTGACCATCACCGCCGACAAGAGCAGCACAGCGCCCTACATGGAACCTGA
 CAGCCTCAGGAGCAGGACACCGCTGTATTACTGCGCCAGGGCGCCATCTACGACGGCTA
 CGACGTGCTGACCAACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGG
 GCCCCAGCGTGTCCCCCTGGCCCCCAGCAGCAAGAGCACAGCGCCAGCGCCCTG
 GGCTGCGCTGGTAAGGACTACTCCCCGAACCGGTGACCGTGTCCCTGAAACAGCGGAGCCCTG
 ACCAGCGCGTGCACACCTCCCCGCGTGTGACAGCAGCGCCCTGTACAGCCTGAGCAGC
 GTGGTGACCGTGCAGCAGCACCTGGGACCCAGACCTACATCTGAAACGTAACGCAAGACCC
 CCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCAAGAGCTGTGACAAGACCCACACCTG
 CCCCCCTGCCCTGCCCGAGCTGGAGGCCAGCGTGTCCCTGTTCCCCCAAGCCT
 AAGGACACCCCTGATGATCAGCGAACCCCCGAGGTGACCTGTGTGGTGGATGTGAGCCAC
 GAGGACCCCTGAGGTGAAGTCACTGGTACGTGGAGCCGGTGGAGGTGACAATGCCAAGACCC
 AACGCCAGGGAGGAGCAGTACAACAGCACCTACGGGGTGGTGTCCGTGACCGTGTG
 CAGGATTGGCTGAAGGCAAGGAGTACAAGTGTAAAGGTGCTAACACAGGCCCTGCCCT
 TCGAGAAAACCATCAGCAAGGCCAGGGCACAGGCCAGAGGCCAGGTGTACACCCCTGCCCT
 CTAGCAGAGATGAGCTGACCAAGAACCAAGGTGTCCCTGACCTGCGCTGGTAAGGGCTTCTACCC
 CACCGACATCCCCGTGAGTGGAGAGCAACGCCAGGCCAGAGAACACTACAAGACCACCC
 CCTGTGCTGGACAGCAGTGGCAGCTTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGA
 TGGCAGCAGGGCAACGTGTTCAGCTGCTCCGTATGCACGAGGCCCTGCACAATCACTACACC
 CAGAAGAGCCTGAGCCTGCCCCGGCAAG

SEQ ID 59 - CA8 J8 Humanised heavy chain
 QVQLVQSGAEVKPGSSVKVSKASGYTFTNYWMHWVRQAPGQGLEWMGATYRGHSDTYYNQK

FKGRVTITADKSTSTAYMELSSLRSEDTAVYCYTRGAIYDGYDVLDNWGGQTLVTVSASTKGPSVF
PLAPSSKSTSGGTAAALGCLVKDVFPEPVTSVNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSL
GTQTYICNVNHPKPSNTVKDKVKEPKSCDKTHTCPGPAPELLGGPSVFLFPKPKDTLMSRTPEVTC
VVVDVSHEDPEVKFNWVYDGVEVHNATAKTPPREQYNTYRVSVLTVLHQDWLNGKEYKCKVNSK
ALPAPIEKTSKAGQPRPEQVYTLPPSDERLTKNQVSLTICLVKFGYPSDIAWEVSENGQOPENNYKTT
PPVLDSDGSFLYSLKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

SEQ ID 60 - CA8 J8 Humanised heavy chain (Polynucleotide)
CAGGTGCAGCTGGTCAGAGCGGCCGAAGTGAAGAAGCCCGCAGCTCCGTGAAAGTGAG
CTGCAAGGCCAGGGCTACACCTTACCAACTACTGGATGCACTGGGTGAGGCCAGGCCCCCG
ACAGGGCCTGGAGTGGATGGGCCACCTACAGGGGCCACAGCGCACCTACTACAAAGGAGA
GTTCAAGGGGGGGTGGACCATACGGCCGACAGAGCAGGCCACAGGGCCACTACATGGAACTGAG
CAGCCTCAGGAGCGAGGACACCGCTGTGATTACTGCAACCGAGGGGCCACTACAGCACGGCTA
CGACGTGCTGGACAACTGGGCCAGGGCACACTAGTGCACCGTGTCCAGCGCCAGCACCAGG
GCCCCAGCGTGTCCCCCTGGCCCCCAGCAGCAAGAGCACCAGCGGGGCCACAGCCGCCCTG

GGCTGCCTGGTAAGGACTACTTCCCGAACCGGTGACCGTGTCTGGAACAGCGGAGCCCTG
ACCAGCGCGCTGCACACCTTCCCGCGTGTGAGCAGCAGCGGCGCTGACAGCTGAGCAGC
GTGGTACCGTGCCAGCAGCAGCAGCTGGGACCCAGACTACATCTGTAACTGTGAAACCACAAG
CCCAGCAACCCAAGGTGGACAAGAAGGTGGAGGCCAAGAGCTGTGACAAGACCACCTGC
CCCCCTGCCCCGCCCCGAGCTGCTGGAGGCCCAAGCGTGTCTGGTCCCCCAAGCT
AAGGACACCTGTGATCAGCAGAACCCCCGAGGTGACCTGTGGTGTGGATGTGAGCCAC
GAGGGACCTCTGAGGTGAACTTCAACTGGTACGTGGACCGCGTGGAGGCTGACATGCAAGACC
AAGCCCAGGGAGGAGCAGTACAACAGCACCTACCGGGTGTCTCGTGCACCGTGTGCAAC
CAGGATTGGTCGAACCGAACGGAGTACAAGTGTGACAGGCTGCTGGTGTGCAAC
TCGAGAAAACCATCAGCAAGGGCAAGGGCAGGCCAGAGGAGCCCCAGGTGTACACCTGGCCCTA
CTAGCAGAGATGAGCTGACCAAGAACCGGTGCTCTGACCTGCTGGTGAAGGGCTTCTACCC
CAGCGACATGCCGTGGAGGGAGGCAACGGCAGCCCGAGAACAACTACAAGACCACCC
CCCTGTGCTGGACAGCGATGGCAGCTCTCTCTGTACAGCAAGCTGACCGTGGACAAGCGAG
TGGCAGCAGGGCAACGTGTTAGCTGCTCCGTGATGCACTGAGGCCCTGCAACATCACTACCC
CAAGAAGGCCCTGAGCTGTCCTGGCAAG

SEQ ID 61 - CA8 J9 Humanised heavy chain
QVLVQSGAEVKPGSSVKSCKGSGYFTFTNVMHWRQAPGQGLEWIGATYRGHSDTYYNQKF
KGRATLTADTSTAYMELRSEDTAVYCTRGAIYDGYDVLNDWGGQGLTVVSSASTKGPSVFF
LAPSSKSTSGGTAALGCLVKDYPFPEVPTWSNSGALTSGVHTFPAVLQSSGGLSLLSVTPVSSSLG
TQTYCINVNHKPSNTVKDKKVEPKSCDKTHTCPGCPAPELGGPSVFLFPPKPKDLMISRTEPVTC
VVDVSHEDPEVKFNWYDVGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKAL
LPAPIEKTSKAKGGQPREPQVYTLPPSRDTELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTP
PVLDSDGSFFLYSLKLTVDKSRWQQGNVFCSVVMHEALHNHYTQKSLSLSPGK

SEQ ID 63 - CA8 M0 Humanised light chain
DIQMTQSPSSLSASVGDRVTITCSASQDISNLYNWWQQKPGKAPKLLIYITSNLHSGVPSSRFSGSGS
GTDFTLTISSLQPEDFATYYCQQYRKLPWTFGQGTKEIERTVAAPSVFIFPPSDEQLKSGTASVVC
NNFYPREAKWVQVKVDNALQSGNSQESVTEQDSKDTSTYSLSSSTLTSKADYEHKVYACEVTHQGLS
SPVTKSFNRGEC

SEQ ID 64 - CA8 M0 Humanised light chain (Polynucleotide)
GACATCCAGATGCCAGGCCAGGGCTAGCTACTGAGGCCAGCGTGGCGACAGGGTGACCTT
ACCTGCTTCCCCAGCCAGGACATCAGCAACTCTGAACTGGTACCCAGCAGAACGCCGGCAAG
GCCCCCAAGCTGCTGATCTACACCTCCAACTCTGCACTTGGCGTGGCCACAGGTTACGGC
GAAGCCGACCCGCAACGGATTCTACCTGACCATCTCGACCTGGCCAGCCGACGGGACTCTGGCA

CCTACTACTGCCAGCAGTACAGGAAGCTCCCCTGGACTTCGGCCAGGGCACCAAACGGAGAT
 CAAAGCCTACGGTGGCCGCCCCCAGGGTGTCTGCTGAACAAACTCTACCCCGGGAGGCAAGGTGAG
 CGGCACCCGCCAGCGTGGTGTCTGCTGAACAAACTCTACCCCGGGAGGCAAGGTGAG
 GAAGGTGGACAATGCCCTGAGCGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAACACA
 AGGACTCCACCTACAGCCTGAGCGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAACACA
 AGGTGTACGCCGTGAGGTGACCCACCAGGGCTGTCACCCCGTACCAAGAGCTTCACC
 GGGCGAGTGC

SEQ ID 65 - CA8 M1 Humanised light chain
 DIQMTQSPSSLASVGDRVTITCSASQDISNYLNWYQQKPGKAPKLIYIYTSLHSGVPSRFSGSGS
 GTDYLTISSLQPEDFATYYCQQYRKLWPWTFGQGTLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLL
 NNFPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTLKADYEHKVVYACEVTHQGLS
 SPVTKSFNRGEC

SEQ ID 66 - CA8 M1 Humanised light chain (Polynucleotide)
 GACATCCAGATGACCCAGGCCCTAGCTACTGAGCGCAGCGTGGGCACAGGGTACATT
 ACCTGCTCCGCCAGGCCAGGACATCAGCAACTACCTGAACTGGTACCCAGCAGAACCGCCGCAAG
 GCCCCCCAAGCTGCTGATCTACTACACCTCAACCTGCACCCGGCTGCCCAGCAGGTTACCG
 GAAGCGGCAGCGGCCACCGATTACACCTGACCCATCTCCAGCCTGAGCAGCCGAGGACTTCGCCA
 CCTACTACTGCCAGCAGTACAGGAAGCTCCCCTGGACTTCGGCCAGGGCACCAAACGGAGAT
 CAAGCGTACGGTGGCCGCCCCCAGCGTGTCTATCTCCCCCAGCGATGAGCAGCTGAAGAG
 CGGCACCCGCCAGCGTGGTGTCTGCTGAACAACTCTACCCCGGGAGGCAAGGTGAG
 GAAGGTGGACAATGCCCTGAGCGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAACACA
 AGGACTCCACCTACAGCCTGAGCGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAACACA
 AGGTGTACGCCGTGAGGTGACCCACCAGGGCTGTCACCCCGTACCAAGAGCTTCACC
 GGGCGAGTGC

SEQ ID 67 - CA8 M2 Humanised light chain
 DIQLTQSPSSLASVGDRVTITCSASQDISNYLNWYQQKPGKAPELVIYIYTSLHSGVPSRFSGSGS
 TDYLTISLQPEDFATYYCQQYRKLWPWTFGQGTLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLL
 NFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTLKADYEHKVVYACEVTHQGLS
 PVTKSFNRGEC

SEQ ID 68 - CA8 M2 Humanised light chain (Polynucleotide)
 GACATCCAGCTGCCAGGCCAGGCCCTAGCTACTGAGCGCAGCGTGGGCACAGGGTACATT
 ACCTGCTCCGCCAGGCCAGGACATCAGCAACTACCTGAACTGGTACCCAGCAGAACCGCCGCAAG
 GCCCCCCAGCTGCTGATCTACTACACCTCAACCTGCACCCGGCTGCCCAGCAGGTTACCG
 GAAGCGGCAGGCCAGCGATTACACCTGACCCATCTCCAGCCTGAGCAGCCGAGGACTTCGCC
 ACCTACTACTGCCAGCAGTACAGGAAGCTCCCCTGGACTTCGGCCAGGGCACCAAACGGAGA
 TCAAGCGTACGGTGGCCGCCCCCAGCGTGTCTGCTGAACAACTCTACCCCGGGAGGCAAGGTGAG
 CGGGCACCGCCAGCGTGGTGTCTGCTGAACAACTCTACCCCGGGAGGCAAGGTGAG
 GGAAAGGTGGACAATGCCCTGAGCGCACCCAGGGAGGAGCTGACCCGAGCAGGAGC
 AAGGACTCCACCTACAGCCTGAGCGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAACAC
 AAGGTGTACGCCGTGAGGTGACCCACCAGGGCTGTCACCCCGTACCAAGAGAGCTTCACC
 CGGGCGAGTGC

SEQ ID 69 - S307118G03 mouse variable heavy
 EVLQQSGPELVKGASVKISCKASGYTFDTYMKWVQSHGKSLEWIGEIPNNGITYNQFKGK
 ATLTVDKSSSTAYMELRSLTSEDSAVYYCANGYEFVYWGQGTIVTVA
 ATLTVDKSSSTAYMELRSLTSEDSAVYYCANGYEFVYWGQGTIVTVA

SEQ ID 70 - S307118G03 mouse variable heavy (DNA sequence)
 GAGGTCCAGTGCACAACTCTGGACCTGAGCTGGTGAAGCCTGGGCTTCAGTGAAGATATCCT
 GTAAAGGCTTCTGGATACACATTACTGACTACTACATGAAGTGGTGAAGCAGAGCCATGGAAA
 GAGCCTGAGTGGATTGGAGAGATTATCTCTAAATAATGGTGTATTACCTACAAACCAGAAAGTTCA
 AGGGCAAGGCCACATTGACTGTAGACAAGTCTCCAGCACGCCATGGAGCTCCGAGC
 GACATCTGAGGACTCTGAGCTTACTGTGAAATGGTACGAGTTACTGGGCAAG
 GGACTCTGGTCACTGCTCTGCA

SEQ ID 71 - S307118G03 mouse variable light
 DIQMTQTASSLSSLQKPDGTVKLIYIYTSSLHSGVPSRFSGSGS
 TDYSLTISNLEPEDIATYYCQQYSLPWTGGGTLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLL

SEQ ID 72 - S307118G03 mouse variable light (DNA sequence)
 GATATCCAGATGACACAGACTGCATCCTCCCTGCTGCTCTGGAGACAGAGTCACCATCA
 GTTCAGTGCAGCTGGGATTAGCAATTATTTAACTGGTACACTCAGCAGAACCCAGATGGAACT
 GTTAAACTCTGATCTTACACATCAAGTTACACTCAGGAGTCCATCAAGGTTAGTGGCAG
 TGCGTGGGAGAGATTATCTCTACCATCAGCAACCTGGAAGATACTTCA
 ATTGTAGCAGTATAGTAAGCTCCGTGGACGTCGGTGGAGGCAACAGCTGGAAATCAAAC
 G

SEQ ID 73 - S307118G03 chimeric heavy chain
 EVLQQSGPELVKGASVKISCKASGYTFDTYMKWVQSHGKSLEWIGEIPNNGITYNQFKGK
 ATLTVDKSSSTAYMELRSLTSEDSAVYYCANGYEFVYWGQGTIVTVA
 ATLTVDKSSSTAYMELRSLTSEDSAVYYCANGYEFVYWGQGTIVTVA
 GGTAAALGCLVKDYFPEPVTVWSNGALSGVHVTFFAVLQSSGLYSLSSVTVPSSSLGTQTYICNVN
 HKPSNTKVKDKVEPKSCDKTHTCPVCPAPELLGGPSVFLPPKPKDLMISRTPEVTCVVVDVSHED
 PEVKFNVYWDGVEVHNAAKTPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIKTIS
 KAKGQPREPQVYTLPPSRDELTQNQVSLTCLVKGFYPSDIAVEWESNGQPEENYKTTPPVLDSDGSF
 FLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

SEQ ID 74 - S307118G03 chimeric heavy chain (DNA sequence)

GAGGTCCAGTTGCAACAACTGGACCTGAGCTGGTAAGCCTGGGCTCAGTGAAGATATCCT
GTAAGGCTCTGGATACACATTCACTGACTACTACATGAAGTGGGTGAAGCAGAGCCATGGAA
GAGCCTGAGTGGATTGGAGAGATTATCCTAATAATGGGTGTTACCTACAACCAGAAGTTCA
AGGCCAAGGCCACATTGACTGTAGACAAGTCTCCAGCACACCCCTACATGGAGCTCCGAGCCT
GACATCTGAGGACTCTGAGCTATTACTGTCAAATGGTTACGAGTTACTGGGCAAG
GGACTCTGGTCACTGTCTCTGCAGGCCAAAACACAGCAGCCCAGCGTGTCCCCCTGGCCCCAG
CAGCAAGAGCACGGCCAGCAGGCCCTGGCTGGTGAAGGACTACTTCCCCGA
ACCGGTGACCGTGTCTGGAACAGCGGAGCCCTGACCGCGCTGCACACCTCCCCGCGT
GCTCAGAGCACGGCTGTACAGCCTGAGCAGCGTGTGACCGTGTCCCCAGCAGCAGCTGG
GCACCCAGACCTACATGTAACTGTAACGTAAACCACAAGGCCAGCAACACAAGGTGACAAGAAGGT
GGACCCAAAGAGCTGTGACAAGACCCACACTGCCCTGCCCTGCCCTGGAGCTGCTGG
AGGCCCCAGCGTGTCTGTGTTCCCCCAAGCTAAGGACACCCGTATGATCAGCAGAAACCCC
GAGGTGACCTGTGTTGGTGGATGTGAGGCCACGAGGACCCCTGAGGTGAGTTCAACTGGTAC
GTGGACGGCGTGGAGGTGACAATGCCAAGACCAAGGCCAGGGAGGAGCAGTACAACAGCACC
TACCGGGTGGTGTCTGTGACCGTGTGACCCAGATTGGCTGAACGGCAAGGAGTACAAG
TGTAAAGGTGTCACAAGGCCCTGCCCTGCCCTATCAGGAAACCATCAGCAAGGCCAAAGGGCC
AGGCCAGAGGCCAGGTGTACACCCCTGCCCTAGCAGAGTGAAGCTGACCAAGAACAGG
TGTCCCTGACTGCCCTGTGAAAGGCTTCAACCCAGGACATGCCCTGGAGTGGAGAGCA
ACGGCCAGCCCAGAACACTACAAGACCAACCCCCCTGTGCTGGACAGCGATGGCAGCTTCT
TCCGTACAGCAAGCTGACCGTGGACAAGAGCAGATGGCAGCAGGGCAACGTGTTAGCTGCT
CCGTGATGCACGAGGCCCTGACAATCACTACACCCAGAACAGGCCCTGAGCTGTCCCCGGCA
AG

SEQ ID 75 - S307118G03 chimeric light chain

DIQMTQTASSLSASLGDRVITSCASQGISNYLNWYQQKPDGTVKLIYIYTSSLHSGVPSRFSGSGSG
TDYSLTISNLEPEDIATYYCQQYSKLPWTFGGGTKLELKRTVAAPSVFIFPPSDEQLKSGTASVVCNN
NFYPREAKVQWKVDNALQSGNSQESVTEQDSKDDSTYLSSTLTLKADYEKHKVYACEVTHQGLSS
PVTKSFNRGEC

SEQ ID 76 - S307118G03 chimeric light chain (DNA sequence)

GATATCCAGATGACACAGACTGCATCCTCCCTGTCTCCCTCTGGAGACAGAGTCACCATCA
GTTGCAGTCAAGTCAGGGCATTAGCAATTATTTAACTGGTATCACGAGAAACAGATGGAACCT
GTTAAACTCTGTATTACACATCAAGTAACTCAGGTTACACTCAGGAGTCCCACATCAAGGTTCACTGGCAG
TGGGTCTGGACAGATTCTCTACCCATCAGCAACCTGGAAACCTGAAGATATTGCAACTTACT
ATTGTCAGCAGTATAGTAACTGGTCTGGAGCTGGTGGAGGCAAGCTGGAGCTGAAACG
TACGGTGCCCCCAGCTGTTCATCTTCCCCCAGCGATGAGCAGCTGAAGAGCGGCAC
CGCCAGCGTGGTGTCTGTGAAACAATTCTACCCCGGGAGGCCAGGTGAGCTGGAGG
GGACAATGCCCTGAGCGGGCAACAGCAGGAGACCGTGAACGAGCAGGACACAAGGACT
CCACCTACAGCCTGAGCAGCACCTGACCCCTGAGCAAGGCCACTACGAGAAGCACAAGGACT
ACGCCCTGTGAGGTGACCCACCAGGGCCTGTCCAGCCCCGTGACCAAGAGCTCAACCGGGCG
AGTGC

SEQ ID 77 - S307118G03 humanised H0 variable heavy

QVQLVQSGAEVKKPGSSVKVSCKASGGTFSYYMKWVRQAPGQGLEWMGEIYPNNGGITYNQKFK
GRVTITADKSTSTAYMELSSLRSEDTAVYYCARGYEFVYWGQGTLVTSS

SEQ ID 78 - S307118G03 humanised H0 variable heavy (DNA sequence)

CAGGTGCAGCTGGTGCAGAGCGGGCGCGAAGTGAAGAAGGCCGGTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGCGGGCACCTTCAGCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
CCAGGGACTGGAGTGGATGGCGAGATCTACCCAAACACGGGGCATCACCTACAACCCAGAA
GTTCAAGGGCAGGGTGACCATCACCGCCGACAAAAGCACCAGCACCGCTACATGGAACTGAG
CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCAGGGCTACGAGTTCGTGTATTG
GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 79 - S307118G03 humanised H1 variable heavy

QVQLVQSGAEVKKPGSSVKVSCKASGYFTDYYMKWVRQAPGQGLEWMGEIYPNNGGITYNQKFK
GRVTITADKSTSTAYMELSSLRSEDTAVYYCARGYEFVYWGQGTLVTSS

SEQ ID 80 - S307118G03 humanised H1 variable heavy (DNA sequence)

CAGGTGCAGCTGGTGCAGAGCGGGCGCGAAGTGAAGAAGGCCGGTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGCGTACACCTTCACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
CCAGGGACTGGAGTGGATGGCGAGATCTACCCAAACACGGGGCATCACCTACAACCCAGAA
GTTCAAGGGCAGGGTGACCATCACCGCCGACAAAAGCACCAGCACCGCTACATGGAACTGAG
CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCAGGGCTACGAGTTCGTGTATTG
GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 81 - S307118G03 humanised H2 variable heavy

QVQLVQSGAEVKKPGSSVKVSCKASGYFTDYYMKWVRQAPGQGLEWMGEIYPNNGGITYNQKFK
GRVTITADKSTSTAYMELSSLRSEDTAVYYCARGYEFVYWGQGTLVTSS

SEQ ID 82 - S307118G03 humanised H2 variable heavy (DNA sequence)

CAGGTGCAGCTGGTGCAGAGCGGGCGCGAAGTGAAGAAGGCCGGTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGCGTACACCTTCACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
CCAGGGACTGGAGTGGATGGCGAGATCTACCCAAACACGGGGCATCACCTACAACCCAGAA
GTTCAAGGGCAGGGTGTACCATCACCGCCGACAAAAGCACCAGCACCGCTACATGGAACTGAG
CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCAGGGCTACGAGTTCGTGTATTG

GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 83 - S307118G03 humanised H3 variable heavy
 QVQLVQSGAEVKKPGSSVKVSKASGYTFTDYYMKVWRQAPGQGLEWIGEIYPNNGGITYNQKFKG
 RATLTVDKSTSTAYMELSSLRSEDTAVYYCANGYEFVYWGQGTLVTVSS

SEQ ID 84 - S307118G03 humanised H3 variable heavy (DNA sequence)
 CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
 CTGCAAGGCTAGCGGTACACCTTACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
 CCAGGGACTGGAGTGGATAGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
 GTTCAAGGGCAGGGCAGCCCTCACCGTCGACAAAAGCACCAGCACCGCTACATGGAACGTGAG
 CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCACGGCTACGAGTCGTATTG
 GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 85 - S307118G03 humanised H4 variable heavy
 QVQLVQSGAEVKKPGSSVKVSKASGYTFTDYYMKVWRQAPGQGLEWIGEIYPNNGGITYNQKFKG
 GRVTITADKSTSTAYMELSSLRSEDTAVYYCAGYEFVYWGQGTLVTVSS

SEQ ID 86 - S307118G03 humanised H4 variable heavy (DNA sequence)
 CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
 CTGCAAGGCTAGCGGTACACCTTACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
 CCAGGGACTGGAGTGGATAGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
 GTTCAAGGGCAGGGTGACCATCACCGCGACAAAAGCACCAGCACCGCTACATGGAACGTGAG

CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCACGGCTACGAGTCGTATTG
 GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 87 - S307118G03 humanised H5 variable heavy
 QVQLVQSGAEVKKPGSSVKVSKASGYTFTDYYMKVWRQAPGQGLEWIGEIYPNNGGITYNQKFKG
 RATLTVDKSTSTAYMELSSLRSEDTAVYYCANGYEFDYWGQGTLVTVSS

SEQ ID 88 - S307118G03 humanised H5 variable heavy (DNA sequence)
 CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
 CTGCAAGGCTAGCGGTACACCTTACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCCGG
 CCAGGGACTGGAGTGGATAGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
 GTTCAAGGGCAGGGCAGCCCTCACCGTCGACAAAAGCACCAGCACCGCTACATGGAACGTGAG
 CAGCCTGAGGAGCGAGGACACCGCCGTGTACTACTCGGCCACGGCTACGAGTCGACTATTG
 GGGCCAGGGCACACTAGTGACCGTGTCCAGC

SEQ ID 89 - S307118G03 humanised L0 variable light
 DIQMTQSPSSLSASVGDRTVITCSASQGISNYLNWYQQKPGKAPKLIYYTSSLHSGVPSRFSGS
 GTDFTLTSSLQPEDFATYYCQQYSKLPWTFGQGQTKLEIKR

SEQ ID 90 - S307118G03 humanised L0 variable light (DNA sequence)
 GACATCCAGATGACCCAGAGCCCCCTCAAGCCTGAGCGCCAGCGTGGCGACAGGGTACTATC
 ACCTGCAGCGCCTCCAGGGCATCAGCAACTACCTGACTGGTACCGAGCAGAACGCCCCGGCAAG
 GCCCTAAGCTGCTGATCTACTACACCAGCAGCGTGCACAGCGGCGTGCCCAGCAGGTTCTCC
 GGCAGCGGAGCGGAACCGACTTCACCCCTGACCATTAGCAGCCTCCAGCCCCAGGAACCTCGCC
 ACCTACTACTGCCAGCACTACAGCAAGCTGCCCTGACCTTCGGCCAGGGACCAAACGGAG
 ATCAAGCGT

SEQ ID 91 - S307118G03 humanised L1 variable light
 DIQMTQSPSSLSASVGDRTVITCSASQGISNYLNWYQQKPGKAPKLIYYTSSLHSGVPSRFSGS
 GTDFTLTSSLQPEDFATYYCQQYSKLPWTFGQGQTKLEIKR

SEQ ID 92 - S307118G03 humanised L1 variable light (DNA sequence)
 GACATCCAGATGACCCAGAGCCCCCTCAAGCCTGAGCGCCAGCGTGGCGACAGGGTACTATC
 ACCTGCAGCGCCTCCAGGGCATCAGCAACTACCTGACTGGTACCGAGCAGAACGCCCCGGCAAG
 GCCCTAAGCTGCTGATCTACTACACCAGCAGCGTGCACAGCGGCGTGCCCAGCAGGTTCTCC
 GGCAGCGGAGCGGAACCGACTACCCCTGACCATTAGCAGCCTCCAGCCCCAGGAACCTCGCC
 ACCTACTACTGCCAGCACTACAGCAAGCTGCCCTGACCTTCGGCCAGGGACCAAACGGAG
 ATCAAGCGT

SEQ ID 93 - S307118G03 CDRH1
 DYYMK

SEQ ID 94 - S307118G03 CDRH2
 EIYPNNGGITYNQKFKG

SEQ ID 95 - S307118G03 CDRH3
 GYEFVY

SEQ ID 96 - S307118G03 CDRL1
 SASQGISNYLN

SEQ ID 97 - S307118G03 CDRL2
 YTSSLHS

SEQ ID 98 - S307118G03 CDRL3
 QQYSKLPWT

SEQ ID 99 - S307118G03 humanised H5 CDRH3
GYEFDY

SEQ ID 100 - S307118G03 humanised H0 heavy chain
QVQLVQSGAEVKPGSSVKVSKASGGTFSDDYMKWVWRQAPGGQLEWMGEIYPNNGGITYNQKFKF
GRVTITADKSTSTAYMELSSRLSEDATVYYCARGYEFVYWGQGLTVVSSASTKGPSVFLAPSSKS
TSGGTAALGCKLVDKPFPEPVTSWNSGALTSGVHTFPVALQSSGLSLSVTPVSSSLGTQYICN
VNHKPSNTKVDKVKVEPKSCDKTHTCPCCPAELLGGPSVFLFPKPKDLMISRTPTEVTCVVVDVSH
EDPEVKFNWYVGDEVHNAKTFPKREEQYNNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK
TISAKGQGPREQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPVLDSD
GSFFLYSKLTVDKRSWQQGNVFCSVMHEALHNHYTQKSLSLSPGK

SEQ ID 101 - S307118G03 humanised H0 heavy chain (polynucleotide)
CAGGTGCAGCTGGTCAGAGCGGCCGAGCTGAAAGAGCCCGCTCAGGGTGAAGGTQAG
CTGCAAGGCTAGGGCGGCCACCTTCAGCGACTACTCATGAGTGGGGTGAAGGGCAGGCCCCCG
CAGGAGCTGGAGTGGATGGCGAGATCTACCCCCAACACGGGGGATCACCTACAGGAA
GTTCAAGGGCAGGGTGGACCATCACCCGGCAGAACAGGAGCAGGGCTACATGGAACTGAG
CAGCCTGAGGAGCGAGGACCCGGCTGTACTACTGCGCAGGGCTACGAGTTCTGTATTG
GGCCAGGGCAGACTAGTGGACCTGGTCAGGCCAGCACAAAGGGCCCGAGCTGGTTCCTCCCT
GGCCCCCAGCAGCAAGAGCACCGGGGGCACAGCGGCCCTGGGCTGCCTGGTGAAGGACT
ACTTCCCCGAACCGGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGGGCTGCACACCT
TCCCCGGCGTGTGAGAGCAGGGCTGTACAGCCTGAGCAGCTGGTACCGTGGCCAGCA
GCAGCCTGGCAGCCAGCCTACATGTAACTGTAACGTGAACCAACAAGCCAGCAACACCAAGGTGG
ACAAGAAGGTGGAGCCAAAGAGCTGTGACAAAGACCCACACCTGCCCCCTGCCCCCTGCCCCG
AGCTGCTGGAGGCCCAAGCGTGTCTGGTGTGGAGTGTGAGCCACGGGACCCCTGAGGTGAAGT
GCAGAACCCCGAGGTGACCTGTGTGGTGTGGAGTGTGAGCCACGGGACCCCTGAGGTGAAGT
TCAACTGGTACGGACGGCTGGAGGTGCAATGCAAGGACCAAGCCAGGGAGGAGCAGT
ACAACAGCACCTACGGGTGGTGTCCGTGCTGACCGTGTGACCAGGATTGGCTGAACGGCA
AGGAGTACAAGTGTAAAGTGTCAACAAGGCCCTGCCCTGCCCTATCGAGAAAACCATCAGCAA
GGCCAAGGGCAGGCCAGAGGCCCAAGGTGACACCCCTGCCCTAGCAGAGATGAGCTGAC
CAAGAACCCAGGTGCTCTGGTGTGGAGGGCTTACCCCGACGCACTGCCGGTGA
GTGGGAGAGCAACGGCCAGGGCAGAGAACACTAACAGAACCCCTGTGCTGGACAGCGA
TGGCAGCTTCTCTGTACAGCAAGCTGGACCCCTGGACAAGAGCAGATGGCAGGGCAAGT
GTTCACTGCTCCGTGTGACAGCAGGCCCTGCAACATCTACACCCAGAAGAGGCTGAGCCCTG
TCCCCCTGGCAAG

SEQ ID 102 - S307118G03 humanised H1 heavy chain
QVQLVQSGAEVVKPGSSVKVSCAKSAGYFTTDYMKWVWRQAPGQGLEWMGEIYPPNNGITYNQKFK
GRVTITADKSTSTAYMELSSLRSEDTAVYYCARGYEFVYWWGQTLVTVSSASTKGPSPVFLPAPSSKS
TSGGTAAALGCKLVKDPEPVEPVWSNNSGALTSGVHTFPALVQSSGLYSLSSVTVPPSSGLTQTYICN
VNHKPSNTKVDDKKEPKSCDKHTCPCCPAPELLGGPSVFLFPKKPKDTLMISRTEPTEVTCVVVDVSH
EDPEPCKFNWYDGVEVHNAKTPREEQYNSTRYVVSPTVHLQDWLNGKEYKCKVSNKALPAPIEK
TISAKGQQPREQPVYTLPPSRSRDELTQNVSSTCLVKGFYPSDIAVEWNSGQPNENYKTPPPVLDSD
GSFFLYSKLTVDKSRWVQQGNVFCSVHMHEALHNHYTQKSLSLSPGK

SEQ ID 103 - S307118G03 humanised H1 heavy chain (DNA sequence)
CAGGTGCAGCTGGTCAGAGCGCGCGCAAGTGAAGAAGCCCGCTCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGGCTACACCTTACCGACTACTACATGAACTGGGTGAGGCCAGGGCCCGG
CCAGGACTGGATGGTGGATGGCGAGATCTACCCCAACACGGGGGCTACCATCACACAGAA
GTTCAAGGGCAGGGTGACCATCACCGCGACAAAGGACCCAGCAGCCCTACATGGAACTAG
CAGCGCTAGGGAGCGAGGACACCCGCGTGTACTACTGCGCCAGGGCTACAGGTGTTGATTG
GGGCCAGGGCACACTAGTGAACCTGTCCAGGCCAGCACAAGGCCAGGGCTTCAGGTGTTCCCT
GGCCCCAGCACAGAACAGAGCACAGCGCCGGCACAGCGCCCTGGCTGCCAGGTGAGGACT
ACTTCCCCGAAACCGGTGACCGTGTCTGGAAACAGCGGAGCCCTGACCAGCGGTGACACCT
TCCCCGCGTGTGAGACAGCGGCTGTACAGCCTGAGCAGCGTGGTACCGTGCAGCA
GCAGCCTGGCACCCAGACCTACATGTAACGTGAACCAACAAGCCAGCAACACCAAGGTGG
ACAAGAAGGTGGAGCCAAAGAGTGTGACAAGACCCCACACTGCCCCCTGCCCTCCCCCG
AGCTGCTGGAGGCCACGGTGTCTCTGTTCCCCCAAAGCTAAGGACACCTGTATGATCA
GCAGAACCCCGAGGTGACCTGTGTTGGTGGATGTGAGGCCAGAGGACCTGAGGTGAAGT
TCAACTGGTACGTGAGCGCGTGGAGGTGCAACATGCCAAGAACCAAGCCGGAGGAGCAGT
AACACAGCACCTACGGGGTGTGTCGTCAGCGTGCACAGGATTGGCTGAACGGCA
AGGAGTACAAGTGAAGGTGTCACAAAGCCCTGCCTGCCCTATCGAGAAAACCATCAGCA
GGCCAAGGGCCAGGCCAGAGAGCCCCAGGTGACACCTGCCCTAGCGAGAGTGAACGCTGAC

CAAGAACCCAGGTGCTCTGACCTGCCGTGAAGGGCTCTACCCCGAGCAGACATGCCGTGGA
GTGGGAGAGCAACGGCAGCCGAGAACACTACAAGGACCCCCCTGTGTCGGACAGCGA
TGGCAGCTTCTCTGTACAGCAAGCTGACCGTGGACAGAGCAGATGGCAGCAGGGCAACGT
GTTCAAGTGTGCTGGTGTGACAGGCCCTGCAACATCAACCCAGAAGAGCTGAGCGTGA
TCCCCCTGGCAAG

SEQ ID 104 - S307118G03 humanised H2 heavy chain
QVQLVQSGAEVKPGSSVKVSKCAGSTFTDYYMKWVVRQAPGQGLEWMGEIYPNNGITYNQKFK
GRVTITADKSTSTAYMELSSLRSEDTAVYYCANGYEFVYWGQGTLTVTSSASTKGPSVFLPLA
TSGGTAALGCLVKDYYFPEPVTVWSNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPVPS
VNHKPSNTKVDKKVVEPKSCDKTHCPCCPAPELLGGPSVFLPPKPKDTLMSIRTP
EVTCVVDVSH

EUPEVKFNVYVUDGVVEVHNAAKIKPREEQYNSIYRUVSVLVLMQDWLNGKEYKCKVSNKALPAPIEK
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPVLDSD
GSFFLYSKLTVDKSRWQQGNVFSCSVMEHALHNHYTQKSLSLSPGK

SEQ ID 105 - S307118G03 humanised H2 heavy chain (DNA sequence)
CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGGCTACACCTCACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCG
CCAGGGACTGGAGTGGATGGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
GTTCAAGGGCAGGGTGACCATCACCGCGACAAAAGCACCAGCACCGCTACATGAACTGAG
CAGCCTGAGGAGCAGGACACC CGGTGACTACTCGCACAACGGCTACGAGTTCTGTATTG
GGGCCAGGGCACACTAGTGAACCGTGTCCAGCGCCAGCACCAAGGGCCAGCGTGTCCCCCT
GGCCCCAGCAGAGCACCAGCGGCCAGACGGCCCTGGCTGGCTGGCTGGTGAAGGACT
ACTTCCCGAACCGGTGACCGTGTCTGGAAACAGCGGCCCTGACCCAGCGCCGTGACACCT
TCCCGCCGTGCTGAGCAGCAGCGGCCCTGACCGCTGAGCAGCGTGTGACCGTGTCCCCAGCA
GCAGCCTGAGGAGCAGGGCACACTACATCTGAACCTGAAACCAAGGGCAGCAACCAAGGTGG
ACAAGAAGGTGAGCCAAAGAGCTGTGACAAGACCCACCTGCCCTGGCTGGCTGGCTGG
AGCTGCTGGAGGCCAGCGTGTCCCTGTTCCCTGAGCTGAGGACAGCTGAGGACCTGATGATCA
GCAGAACCCCCAGGGTGAACCTGTTCCCTGAGCTGAGGACAGCTGAGGACCTGAGGAG
TCAACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAGACCAAGCCAGGGAGGAGCAGT
ACAACAGCACCTACCGGGTGGTGTGCTGACCCAGGATTGGCTGAACGGCA
AGGAGTACAAGTGAAGGTGTCACAAGGCCCTGCTGCCCTATCGAGAAAACCATCGCAA
GGCCAAGGGCCAGCCAGAGGCCAGGGTGTACCCCTGCCCTAGCAGAGATGAGCTGAC
CAAGAACCGAGGTGCTCCCTGACCTGCCCTGGTAAGGGCTTCTACCCAGCAGACATGCCGTGGA
GTGGGAGGAGCAACGGCAGGCCAGAACAACTACAAGACCAAGCCCCCTGTGCTGGACAGCAGT
TGCAGCTGCTCCGTGATGCAAGGCCCTGCACAATCACTACACCCAGAAGGCCCTGAGCCTG
TCCCCTGGCAAG

SEQ ID 106 - S307118G03 humanised H3 heavy chain
QVQLVQSGAEVKPKGSVVKSCASGYFTDYMVKWRQAPGQGLEWIGEIYPNNGGITYNQKFKG
RATLTVDKSTSTAYMELSLRSEDTAVYYCANGYEFVYWGQGTLTVSSASTKGPSVFLPAPSSKST
SGGTAAALGCLVKDVFPEPVTVWSNNGALTSGVHTFPAVLQSSGLYLSVVTVPSLGLTQTYICNV
NHKPSNTKVDKKVEPKSCDKTHTCPCTGACCTGCCCTGGTAAGGGCTTCTACCCAGCAGACATGCCGTGGA
DPEVKFNWYVUDGVVEVHNAAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT
SKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPVLDSD
SFLLYSLKLTVDKSRWQQGNVFSCSVMEHALHNHYTQKSLSLSPGK

SEQ ID 107 - S307118G03 humanised H3 heavy chain (DNA sequence)
CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGGCTACACCTCACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCG
CCAGGGACTGGAGTGGATAGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
GTTCAAGGGCAGGGCACCCCTCACCGCGACAAAAGCACCAGCACCGCTACATGAACTGAG
CAGCCTGAGGAGCAGGACACC CGGTGACTACTCGCACAACGGCTACGAGTTCTGTATTG
GGGCCAGGGCACACTAGTGAACCGTGTCCAGGCCAGCAGGCCAGGGCCAGCGTGTCCCCCT
GGCCCCAGCAGAGCAGGCCAGCAGGCCAGCAGGCCAGGCCAGCGTGTGAGGACT
ACTTCCCGAACCGGTGACCTGCTGGAAACAGCGGCCCTGACCCAGCAGCGTGTGACACCT
TCCCGCCGTGCTGAGCAGCAGGCCCTGACCGCTGAGCAGCGTGTGACCGTGTGAGGAG
GCACCCCTGGCACCCAGACCTACATCTGAACCTGAAACCAAGGCCAGCAACCAAGGTGG
ACAAGAAGGTGAGCCAAAGAGCTGTGACAAGACCCACCTGCCCTGGCTGGCTGG
AGCTGCTGGAGGCCAGCGTGTCCCTGTTCCCTGAGCTGAGGACAGCTGAGGACCTGATGATCA
GCAGAACCCCCAGGGTGAACCTGTTCCCTGAGCTGAGGACAGCTGAGGACCCCTGAGGAG
TCAACTGGTACGTGGACGGCGTGGAGGTGACAATGCCAAGACCAAGCCAGGGAGGAGCAGT
ACAACAGCACCTACCGGGTGGTGTGCTGACCCAGGATTGGCTGAACGGCA
AGGAGTACAAGTGAAGGTGTCACAAGGCCCTGCCCTATCGAGAAAACCATCGCAA
GGCCAAGGGCCAGCCAGAGGCCAGGGCTGACCCCTGCCCTAGCAGAGATGAGCTGAC
CAAGAACCGAGGTGCTCCCTGACCTGCCCTGGTAAGGGCTTCTACCCAGCAGACATGCCGTGGA
GTGGGAGGAGCAACGGCAGGCCAGAACAACTACAAGACCAAGCCCCCTGTGCTGGACAGCGA
TGCAGCTTCTCCGTACAGCAAGCTGACCGTGGACAAGAGCAGATGGCAGCAGGGCAACGT
GTTCACTGCTCCGTGATGCAAGGCCCTGCACAATCACTACACCCAGAAGGCCCTGAGCCTG
TCCCCTGGCAAG

SEQ ID 108 - S307118G03 humanised H4 heavy chain
QVQLVQSGAEVKPKGSVVKSCASGYFTDYMVKWRQAPGQGLEWIGEIYPNNGGITYNQKFKG
GRVTITADKSTSTAYMELSLRSEDTAVYYCADGYEFVYWGQGTLTVSSASTKGPSVFLPAPSSKST
TSGGTAAALGCLVKDVFPEPVTVWSNNGALTSGVHTFPAVLQSSGLYLSVVTVPSLGLTQTYICNV
VNHKPSNTKVDKKVEPKSCDKTHTCPCTGACCTGCCCTGGTAAGGGCTTCTACCCAGCAGACATGCCGTGGA
DPEVKFNWYVUDGVVEVHNAAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT
TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPVLDSD
GSFFLYSLKLTVDKSRWQQGNVFSCSVMEHALHNHYTQKSLSLSPGK

SEQ ID 109 - S307118G03 humanised H4 heavy chain (DNA sequence)
CAGGTGCAGCTGGCAGAGCGGCCGAAGTGAAGAAGCCGGCTCCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGGCTACACCTCACCGACTACTACATGAAGTGGGTGAGGCAGGGCCCG
CCAGGGACTGGAGTGGATGGCGAGATCTACCCAAACAACGGGGCATCACCTACAACCAGAA
GTTCAAGGGCAGGGTGAACCTCACCGCGACAAAAGCACCAGCACCGCTACATGAACTGAG
CAGCCTGAGGAGCAGGACACC CGGTGACTACTCGCAGCAGGCCAGCTACGAGTTCTGTATTG
GGCCCCAGGGCACACTAGTGAACCTGAGGAGCAGGCCAGCGTGTGAGGAGCTGAGGAG

SEQ ID 110 - S307118G03 humanised H5 heavy chain
QVQLVQSGAEVKKPGSSVKVSCKASGYTFTDYYMKWVRQAPGQGLEWIGEYIYPNNNGITYNQKFKG
RATLTVDKSTSTAYMELSSLRSEDTAYVYCANGYEFDWVQGQTLVTVSSASTKGPSWFPLAPSSKST
SGGTAALCGKVKDVFPEPVTWSNWSALGTSVHFTPAVLQSSGGLYSLSSVTVPSNNSLGTQYTCINV
NHKPSNTVKDKVKEPKSCDCKTHCPCPAFELLGGPSVFLFPKPKDTLMSIRTPETVCVVDSHE
DPEVKFNVWYDGVHVEVNAKTKPREQYNNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT
SAKAGQPREPVQVYTLPPRSDELTKNQVSLTCLVKGYFPPSVDIAVEWESNGQPNENYYKTPPVLDSDG
SFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

SEQ ID 111 - S307118G03 humanised H5 heavy chain (DNA sequence)
CAGGTGCAGCTGGTCAGACGGCGCCGGAAGTGAAGAACGCCGGCTCAGCGTGAAGGTGAG
CTGCAAGGCTAGCGGTCACAGCTTCACCGACTACTACATGAGTGGGGTGAAGCGAGGGCCCG
CCAGGGACTGGAGTAGGGAGATCAGGGACAAACCGGGGCACTACCTAACACCGAGAA
GTTCAAGGGCAGGGGACCCCTACCGCTGCACAAAAGCACCGACCCGCTACATGGGAACGTAG
CAGCTGAGGAGCGAGGACACC CGCGTGTACTACTGCGCCAAAGGCTACGAGTTGCACTATTG

GGGGCCAGGGCACACTAGTGCACCGTGTCCAGCGCCAGCACCAAGGGCCCCAGCGTGTCCCCCT
GGCCCCCAGCAGCAAGAGCACCAGCGCGGCACAGCGGCCCTGGCTGCCTGGTGAAGGACT
ACTTCCCCGAAACCGGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGCACACCT
TCCCCCCCAGTGTGAGCAGCGGCGCTGTACAGCCTGAGCAGCGTGTGACCCTGCCCCAGCA
GCAGCGTGGCACCCAGACTACATCTGTAACTGTAAGCTGAACACAAAGCCAGCACAACAAAGCTGG
ACAAGAAGGGTGGGAGGCCCCAGGGACTGTGACAAAGACCCACACCTGCCCCCTGGCTGGCCCC
AGCTGCTGGGAGGCCCCAGGGTGTCTGGCTGGGAGGCCCCAGGCTAAAGGACACCTGTATGATCA
GCAGAACCCCCGGAGGTGACCTGTGTGTGTGGATGTGAGCACCAGGGACCTGAGGTGAACT
TCAACTGGTACGTGGACGGCGTGGAGGTGCAACATGCCAACAGCCAGGGAGGAGCAGT
ACAACAGCACCTACGGGGTGGTGTCCGTGCTGACCGTGTGCAACAGGATTGGCTGAACGGCA
AGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCCTGCCCTATCGAGAAAACCATCAGCAA
GGCCAAGGGCAGGCCAGAGAGGCCAGGTGACACCCCTGCCCTAGCGAGATGAGCTGAC
CAAGAACAGGTGCTCTGACCTGGCTGGTGAAGGGCTTACCCCCAGCGACATGCCGTGGA
GTGGGAGGAGCACGGGCCAGCCCCAGAGAACACTACAAAGACCCACCCCCCTGTGCTGGACAGCGA
TGGCAGCTTCTCTGGTACAGCAAGGCTGACCGTGGACAAAGAGCAGATGGCAGCAGGGCAACGT
GTTGCGTGTGGCTGGTGAAGGGCCCTGCACAATCACTACACCCAGAAGAGGCTGACGCCCTG
TCCCCCTGGCAAG

SEQ ID 112 - S307118G03 humanised L0 light chain
DIQMTQSPSSLSASVGDRVTITCSASQGINSYLNWYQQKPGKAPKLLIYYTSSLHSGVPSRFGSGS
GTDFTLTISSQLPQEDFATYCQQYSKLPWTFGQQGTLKLEIRTVAAEPSVFIFPPSDEQLKSGTASVVCLL
NNFYFPRKEAVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSLTLSKADYEHKVYACEVTHQGLS
SPVTKSENPIGEC

SEQ ID 113 - S307118G03 humanised L0 light chain (DNA sequence)
GACATCCAGATGACCCAGAGGCCCTCAAGGCCCTAGGCCAGCGTGGCGCAGAGGGTACTATC
ACCTGCAGGCCCTCCAGGGCATCAGCAACTACCTGAACTGGTACCCAGCAGAAAGCCCGCAAG
GCCCTAAGCTGCTGATCTACTACACCAGCAGCCTGCACAGCGCGTGCAGCAGGGTCTCC
GGCAGCGGCAGCGGAACCGACTTACCCCTGACCATTAGCAGCCTCCAGCCCGAGGGACTTCGC
ACCTACTACTGCCAGCAGTACAGCAGGCTGCCCTGGACCTTCCGGCAGGGCACCAAACGGAG
ATCAAGCGTACGGTGGCCGCCCGCCAGCGTGTCTACCTCTCCCGCCAGCGATGAGCAGCTGAAG
AGCGGCACCCGCCAGCGTGTGTCTGCTGAACAACTCTACCCCGGGAGGCAAGGTGCA
TGGAAAGGTGACAATGCCCTGCAGAGCGGCAACAGCCAGGAAGCGCTGACCCAGCAGGACAG
CAAGGGACTCCACCTACGCCCTGAGCAGCACCCCTGACCCCTGAGCAAGGGCAGTACGAGAAGCA
CAAGGTGTAGCCCTGTAGGGTGAACCCACCGGGCTGTCCAGCCCCGTGACCAAGAGCTTCAA
CGGGGGCAGGTGCG

SEQ ID 114 - S307118G03 humanised L1 light chain
DIQMTQSPSSLSASVGDRVTITCSASQGISNYLNWYQQPGKAPKLLIYTSSLHSGVPSRFSGSGS
GTDYTLTISQLQPEDFATYYCQQYSKLPWTFGGQGTLEIKRTVAAPSVFIFPPSDEQLKSGTASVCLL

NNFPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTKADYEKHKVYACEVTHQGLS
SPVTKSFNRGEC

SEQ ID 115 - S307118G03 humanised L1 light chain (DNA sequence)
GACATCCAGATGACCCAGGCCCTCAAGCTGAGGCCAGGGCTGGCGACAGGGTGACTATC
ACCTGAGCGCCCTCCAGGGCATCAGCAACTACCTGAACTGGTACCCAGCAGAAAGCCCGGCAAG
GCCCTAAGCTGCTGATCTACACCCAGCAGCCTGCACAGGGCGTGGCCACAGGGTTCTCC
GGCAGCGGCAGCGGAACCGCATCACCCCTGACCATTAAGCAGCTCCAGCCGAGGACTTCGC
ACCTACTGCGCAGCTACAGCAAGCTGCCCTGGACCTTGGCCAGGGCACAAACTGGAG
ATCAAGCGTACGGTGGCCGGCCCGGAGCTGTTCTATCCCCCCCCAGCGATGAGCAGCTGAAG
AGCGGCACGGCAGCGTGGTGTCTGTAACAACTTCAACCCGGGGAGGCCAAGGTGCA
TGGAGGTGGACAATGCCCTGAGAGCGGCAACAGCAGGGAGAGCGTGAACCAGCAGGACAG
CAAGGACTCCACCTACACCCCTGAGCAGCACCTGACCCCTGAGCAAGGCCACTACGAGAAGCA
CAAGGTGACCCCTGTGAGGTGACCCACCAAGGGCTGTCCACCCCCGTGACCAAGAGCTCAA
CCGGGGCGAGTGC

SEQ ID 116 - S332121F02 murine variable heavy chain
EVQLQQSGPVLVKPGASVKMSCEASGYTFTDYYMNWKQSHGKTLIEWGVINPYNGGTDYNQKFK
GKATLTVDKSSSTAYMELNSLTSEDSAVYYCARSVYDYPFDWGGQGLTVTVSS

SEQ ID 117 S332121F02 murine variable heavy chain (DNA sequence)
GAGGTGCAGCTGCAGCAGAGCGGCCGCTGCTGGTGAAAGCCTGGAGCAGCGTGAATATGAG
CTGCGAAGCCAGCGCTCACCTTACCGACTACTACATGAACTGGGTGAAAGCAGGCCACGG
CAAGACCTCCGGAGTGGATGCCGCTGATCACACCTTACAAACGGGGCACCGACTACAAACAGAA
GTTCAAGGGCAAGGGCACTCTGACCTGGACAAAGGCTCAGCACCCCTACATGGAACTGAA
CAGCCTCACCTCTGAGGACAGCGCCGCTATTACTGCGCCAGGACGCTGACGACTACCCCTTC
GACTACTGGGCCAGGGCACACTAGTGACCGTGCCAGC

SEQ ID 118 - S332121F02 chimeric heavy chain
EVQLQQSGPVLVKGAVMSCEASGYTFTDYYMNWKQSHGKTLIEWGVINPYNGGTQDYNQKFK
GKATLTVDKSSSTAYMELNSLTSEDAVYYCARSVYDQPFDVWQGQLTVTSSASTKGPSVFLPLAPS
SKSTSGGTAAALGCLVKDGYPEPVTVWSNSGALTSGVHTFPAVLQSSGLYSLSSVTPVSSSLGTQTY
CNVNHPKSNTKVDKVKVEPKSCDKTHTCPCPAPELGGPSVFLPPKPKDLMISRTPEVTCVVVDV
SHEDPEVKFNWYVGVEVHNAKTPKREPPQYNSRVRVSVLTVLHQDWLNGKEYKCKVSNKALPAPI
EKTISKAGQGPREPQVYTLPPSREDELTKNQVSLSCLVKGVPYPSDIAVEWESNGQPNENYKTTTPVLD
SDGSFFLYSKLTVDKSRWQQGNVFSCSVMEHEALHNHYTQKSLSLSPGK

SEQ ID 119 - S332121F02 chimeric heavy chain (DNA sequence)
GAGGTGCGAGCTGCAAGCAGCGGCCCTGCTGGTAAGGCTTGGAGGCCAGCGTGAAGAAATGAG
CTGCGAAGCCAGCGGCCACACCTTACCCGACTACTAATGACTGGTTGAAGCAGAGGCCACGG
CAAGACCTGAGTGTGATCGGCCGTGATCACCCCTACACCGGGGCCAGCAGTACAACCCGAA

GTTCAGGGCAAGGCCACTCTGACCGTGGACAAAGAGCTTCAAGCCGCTACATGGAACTGAA
CAGCCTCACCTCTGAGGACAGCGCCGCTATTACTGCGCCAGGAGCGTGTACGACTACCCCTTC
GACTACTGGGCCAGGGCACACTAGTGACCGTGTCCAGCGCCAGCACCAAGGGCCCGACGCT
GTTCCCCCTGCCCCCAACCGAACAGGACCCAGCGGCCACAGGCCCCCTGGCTGGCTGG
TGAAAGACTTCCCCGAAACCGTGACCGTGTCTGGACACGGGAGCCCTGACCGCGCC
TGACACCTTCCCCGGCTGCGAGCAGCGGCCGCTACAGCTGAGCAGCTGGTGACCG
TCCCCAGCAGCAGCCTGGCACCCAGACCTACATCTGTAACGTGAACCCACAGGCAACAC
CAAGGTTGACAAGAAGGTGGAGCCAAAGCTGTGACAAGACCCACACTGCCCCCTGGC
TGCCCCCGAGCTGCTGGAGGCCAGCGTGTCTGTGTTCCCCCAAGCTAAGGACACCT
GATGATCAGCAGAACCCCCGAGGTGAGCTGTGCTGGTGTGAGCCACGAGGACCTGA
GGTGAAGTCACTGGTACGTGGACGGCGTGGAGGTGACAACTGCAAGAACAGGCAAGGG
GGAGCAGTACACAACCTGGGTTGGTGTGGTGTGACCCAGGATTGGCT
GAACGGCAAGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCTGCCCCATCGAGAAAACC
ATCAGCAAGGCCAAGGGCAGCCAGCCAGAGAGCCCCAGGTGTACACCTGCCCCTAGCAGAGAT
GAGCTGACCAAGAACCCAGGTGCTCTGACCTGCTGGTGAAGGGCTTACCCACCGACATC
GCCGTGGAGTGGGGAGAGCACGGCCAGGGAGAACAACTAACAGACCAACCCCCCTGTGCTG
GAACAGCGATGGCAGCTTCTCTGTACAGCAGGCTGACCCAGGAGCAGATGGCAGCAG
GGCACCTGTGTCAGCTGCTCGTATGCGAGGCCCTGACAACTACACCCAGAAGAGCC
TGAGCTGCCCCCTGGCAAG

SEQ ID 120 - S332121F02 murine variable light chain
DIVLTQSPASLAVSLGQRATISCRASESVIHGTHLMHWYQQKPGQPPKLLIYASNLLESGVPARFSG
SGSETDFTLNIHPVEEDAATYFCQQSIEDPRTFGGGTKLEIK

SEQ ID 121 - S332121F02 murine variable light chain (DNA sequence)
GACATCGTCTGACCCAGAGCCCCGCGCAGGCCCTGGCCGTGAGCCCTGGGCCAGAGGGCCACAATC
AGCTGCAGGCCCTCTGAGCTCTGAGCATCCACGGGCCACCCACTGTGACTGTGTTATCAGCG
AAAGCCGGCCAGCCTCCAAAGCTGCTGATCTACGCCGCCAGAACCTTGAGAGCCGGCTGCC
GCTAGTTTCAGCGGAAGCGCAGCGAGACCCGACTTCACCTCTGAAACATCCCCCTGAGGAG
GAAAGCCGCCGCACTTCTGCCAGCAGAGCATCGAGGACCCAGGACCTTCGGCGGGGCC
ACCAAGCTCAGGATTAAGCGT

SEQ ID 122 - S332121F02 chimeric light chain
MGWSCLILFLVATATGVHSIDVLTLQSPASLAVSLGQRATISCRASESEVSVIHGTLMHWYQQKPGQQPK
LLIYASNLSEGVGPARFSGSGSEEDTDLNIIHPEVEEDAATYFCQCSIEDPRTFGGTKEIKRTVAAPS
VFIFPPSDDEQLKSGTASVWCLNNFYPREAKVQWVBDNALQSGNSQESVTEQDSKDTSTYSSLSTLT
SKADYKEHKVYACEVTHQGLSPVTKSFNRGEC

SEQ ID 123 - S332121F02 chimeric light chain (DNA sequence)
ATGGGCTGGCTCTGCATCATCTGTTCTGGTGCCACCGCCACCGGCGTGCACAGCGACATC
GTCTCAGGCCAGAGCCCCCGCAGCCTGCGCCCTGAGCCTGCGGCCAGAGGGCCACAATCAGCTG
CAGGGCTCTGTAGCTGGCTGAGCATCACGGCACCCACCTGTGACTTGTATCAGCAGAACGC
CGGCCAGCCTCTTCAAGCTCTGTATCAGCCGCCAGAACCTGGAGAGGCCGGCTGCCCTAG

GTTCAGCGGAAGCGGCAGCGAGACCGACTTCACCCCTGAACATCCACCCCGTGGAGGGAGGAAGA
CGCGGAGACCTCTCTGCGCAGCAGACATCGAGGACCCAGGACCTTCGGGGGGCACCAA
GCTCGAGATTAAAGCGTACGGTGGCGCCCCCAGCGTGTTCATCTTCCCCCCCAGCGATGAGCA
GCTGAAGAGCGGCACCGCCACCGTGTGTCTGCTGAACAACTTCTACCCCCGGAGGCCAA
GGTCAGTGGAAAGGTGGACAATGCCCTCAGAGCGGCAACAGCCAGGAGCGTGACCGAGC

AGGACAGCAAGGACTCCACCCCTACAGCTGAGCAGCACCCGTGACCTGAGCAAGGCCAGTACAG
AGAAGCACAAGGTGTACCCCTGTGAGGTGACCCACCCAGGGCTGTCCAGCCCCGTGACCAAGA
GCTCAACCGGGCGAGTG

SEQ ID 124 - S322110D07 murine variable heavy chain
EVQLQQSGPELVKPGTSVKIPCKTSGYIFTDYSIDWVKQSHGKSLWIGDIDPNYGDPIYNHKFKGKA
TLTVDRSSSTAYMELRSLTSEDTAVYFCARRATGTDWFAFWQGQTLTVSS

SEQ ID 125 - S322110D07 murine variable heavy chain (DNA sequence)
GAGGTGCAGCTGCAGCAGAGCGGGCCCGAGCTGGTGAACCCGGCACCGCGTGAAGATCCC
CTCGAACACCTCTGGCTACATCTTACCGACTACAGCATCGACTGGTGAAGCAGAGCCACGGC
AACTCTGGAGTGGATGGGACATCGACCCCAACTACGGCGACCCCATCTACAACCCAAAGT
TCAAGGGCAAGGCCACCCCTGACCGTGGACAGGAGCAGCAGCACCCCTACATGGAACCTAGGA
GCCTGACCAGCGAGGGACACCGCCGTGTATTTGCCAGGAGGGGCCACCGGACTGATTGGT
TCGCTCTCTGGGGCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGGGCCAGCG

SEQ ID 126 - S322110D07 chimeric heavy chain
EVQLQQSGPELVKPGTSVKIPCKTSGYIFTDYSIDWVKQSHGKSLWIGDIDPNYGDPIYNHKFKGKA
TLTVDRSSSTAYMELRSLTSEDTAVYFCARRATGTDWFAFWQGQTLTVSS
STSGGTAALGCLVYDFFPEPVWSNSALTSVHFTPAVLQSSGLYLSLSSVTPVSSSLGTQTYIC
NVNHPNSNTKDKVPPKSCDKTHTCPCTPAPELLGGPSVFLFPKPKDLMISRTPEVTCVVVDVS
HEDPEVKFNWYDGEVHNAAKTPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISAKQQPREEPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWEESNGQPENNYKTPPVLD
DGSFFYSLKTVDKSRWQQGNFSCSVMHEALHNHYTQKSLSLSPGK

SEQ ID 127 - S322110D07 chimeric heavy chain (DNA sequence)
GAGGTGCAGCTGCAGCAGAGCGGGCCCGAGCTGGTGAACCCGGCACCGCGTGAAGATCCC
CTCGAACACCTCTGGCTACATCTTACCGACTACAGCATCGACTGGTGAAGCAGAGCCACGGC
AACTCTGGAGTGGATGGGACATCGACCCCAACTACGGCGACCCCATCTACAACCCAAAGT
TCAAGGGCAAGGCCACCCCTGACCGTGGACAGGAGCAGCAGCACCCCTACATGGAACCTAGGA
GCCTGACCAGCGAGGGACACCCCGGTGTATTTGCCAGGAGGGGCCACCGGACTGATTGGT
TCGCTCTCTGGGGCAGGGCACACTAGTGACCGTGTCCAGCGCAGCACCAAGGGCCAGCG
TGTTCCCCCTGGCCCCCAGCAGCAAGCACCAGCGGGCGCACAGCCGCCCTGGCTGCGCTG
GTGAAGGACTACTTCCCCAACCGGTGACCCCTGTCTGGACAGCGGAGGCCCTGACCCAGCG
GTGACACCTTCCCCCGCCGTGCGAGCAGCGGCCCTGTACAGCCTGAGCGCTGGTGACC
GTGCCCCAGCAGCCTGGCACCCAGACCTACATGTAACCTGAACCCAAGGCCACGAAAC
ACCAAGGTGACAAGAGTGGAGCCAAAGAGCTGTGACAAGACCCACACCTGCCCCCTGC
CCTGCCCCCGAGCTGCTGGAGGGCCAGCGTGTCTGTGCCCCCAAGGCTAAGGACACC
CTGATGATCAGCAGAAACCCCGAGGTGACCTGTGAGCCAGGACAGCGGCCCTAGAGAC
GAGGTGAAGTCACTGGTACCGTGGAGGGGTGACAGCAGCGGCCCTGTACAGCCTGAGCG
GAGGAGCAGTACAACAGCAGCCTGGTGTGCTGACCGTGTGCTGACCCAGGATTGG
CTGAACGGCAAGGTACAAGGTGAAAGGTGCTGACCGTGTGCTGACCGTGTGCTGCCCC
CCATCAGCAAGGCCAGGGCCAGCCCAGAGGCCAGGTGACCCCTGCCCCCTAGAGAG
ATGAGCTGACCAAGAACCCAGGTGCTCCGTGACCTGCTGGTGAAGGGCTTCTACCCAGC
CGCCGTGGAGTGGAGAGCACGCCAGCCGAGAACACTACAAGACCAACCCCCCTGTG
GGACAGCGATGCCAGCTTCTGTACACCAAGCTGACCGTGGACAGAGCAGATGGCAGCA
GGGCAACGTGTTGAGCTGCTCCGTGATGCAAGGGCCCTGACAATCACTACACCCAGAAG
CCTGAGCTGCTGCCCTGGCAAG

SEQ ID 128 - S322110D07 murine variable light chain
DIQMTQSPASLSVSVGETVTICRASENIYNNLAWYQQKQGKSPQLVYATIADGVPSRFSGSGSG
TQYSLKINSLQSGDFGTYYCQHFWTPLTFGAGTKLEKLR

SEQ ID 129 - S322110D07 murine variable light chain (DNA sequence)
GACATCCAGATGACCCAGAGCCCCGCTAGCCTCAGCGTGTCCCGAGACCGTGACCATC
ACCTGAGGGCAGCGAGAACATCTACAACACCTGGCTGGTACAGCAGAACGGCAA
AGCCCCCAGCTCTGGTGTACGGCCACCTCTGGCGACGGCGTCCCCAGCAGGTTCTCT
GGAAGCGGCAGCGGCACCCAGTACAGCCTGAAGATCAACAGCCTGAGAGCGGGGACTTCG
CACCTACTACTGCCAGCACTCTGGGGACTCCCTGACCTTCGGAGCCGGCACCAAGCTGG
GCTGAAGCGT

SEQ ID 130 - S322110D07 chimeric light chain
DIQMTQSPASLSVSVGETVTICRASENIYNNLAWYQQKQGKSPQLVYATIADGVPSRFSGSGSG
TQYSLKINSLQSGDFGTYYCQHFWTPLTFGAGTKLEKLR
NNFYPREAKVQWVKDNAQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGL
SPVTKSFNRGEC

SEQ ID 131 - S322110D07 chimeric light chain (DNA sequence)
GACATCCAGATGACCCAGAGCCCCGCTAGCCTCAGCGTGTCCCGAGACCGTGACCATC
ACCTGAGGGCAGCGAGAACATCTACAACACCTGGCTGGTACAGCAGAACGGCAA
AGCCCCCAGCTCTGGTGTACGGCCACCTCTGGCGACGGCGTCCCCAGCAGGTTCTCT
GGAAGCGGCAGCGGCACCCAGTACAGCCTGAAGATCAACAGCCTGAGAGCGGGGACTTCG
CACCTACTACTGCCAGCACTCTGGGGACTCCCTGACCTTCGGAGCCGGCACCAAGCTGG
GCTGAAGCGTACGGTGGGGCCCCCAGCGTGTCTGGTGTCTGGTGTGACCAAGCAGCTGG
GAGCGGGCACCGCCAGCGTGTGCTGACCAACTTCTACCCCGGGAGGGCAAGGTGCA
GTGAAGGTGACAATGCCCTGAGCGAGCACAGCCAGGAGAGCGTACCGAGCAGGACA
GCAAGGACTCCACCTACAGCCTGAGCGAGCACCCGTGACCTGAGCAAGCCGACTACAGGAGA

ACAAGGTGTACGCCGTGAGGTGACCCACCAGGGCTGTCCAGCCCCGTGACCAAGAGCTCA
ACCGGGGGAGTGC

SEQ ID 132 - S332126E04 murine variable heavy chain
QVQLQQPGAEVKGASVVLSCASGYFTNYWMHW/KQRPGQGLEWIGIHPNSGNTNYNEKFKS
KATLTVDKSSSTAYMQLSSLTSEDSAVYYCARGIYDYPFAYWGGTLLTVSS

SEQ ID 133 - S332126E04 murine variable heavy chain (DNA sequence)
CAGGTGAGCTCCAGCAGCCGGAGCCGAAGCTGGTAAGGCCGGAGCCAGCGTCAAAGTGCC
TCCAAGGCCAGGGCTACACCTCACCAACTACTGGATGCACTGGGTGAAGCAGAGGCCGGC
CAGGGCTGTGAGGGATCGGCATCATCCACCCAAACAGCGGGAGCACCAACTACAAGGAG
TTCAAGAGCAAGGGCACCCGTGACCGTGACAGACAGCAGCACTGCCTACATGCAGCTGAGC
AGCCTGACCAAGCGAGGACAGCGCTGTGACTACTGCAGCCAGGGCATCAGACTACCCCTC
GCCTATTGGGGCAGGGCACACTAGTGACCGTGTCCAGCGCCAGCACCAAGGGCCAGCGT
EKTISKAKGQPQREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD
SDGSFFLYSKLTVDSRWWQGNVFCSVMIHEALHNHTQKSLSLSPKG

SEQ ID 135 - S332126E04 Chimeric heavy chain (DNA sequence)
CAGGTGAGCTCCAGCAGCCGGAGCCGAAGCTGGTAAGGCCGGAGCCAGCGTCAAAGTGCC
TCCAAGGCCAGGGCTACACCTCACCAACTACTGGATGCACTGGGTGAAGCAGAGGCCGGC
CAGGGCTGTGAGGGATCGGCATCATCCACCCAAACAGCGGGAGCACCAACTACAAGGAG
TTCAAGAGCAAGGGCACCCGTGACCGTGACAGACAGCAGCACTGCCTACATGCAGCTGAGC
AGCCTGACCAAGCGAGGACAGCGCTGTGACTACTGCAGCCAGGGCATCAGACTACCCCTC
GCCTATTGGGGCAGGGCACACTAGTGACCGTGTCCAGCGCCAGCACCAAGGGCCAGCGT
TTCCCCCTGGCCCCAGCAGCAAGAGCACCAAGCGGGCCAGCGCCAGCGCCAGCGTGG
GAAGGACTACTCCCGAACCGTGACCGTGCTCTGGAACAGGGAGCCCTGACCCAGCGCGT
GCACACCTTCCCCCGTGTGACAGCAGCGGGCTGTACAGCCTGAGCAGCGTGGTACCGT
GCCAGCAGCAGCTGGCACCCAGCCTACATCTGAACTGAAACCACAGCCAGCAACAC
CAAGGTGACAAAGAGTGAGGCAAGAGCTGTGACAAAGACCCACACTCCCCCTGCC
TGCCCCCGAGCTGGAGGCCCCAGCGTGTCTGTTCCCCCAAGCCTAAGGACACCC
GATGATCAGCAGAACCCCGAGGTGACCTGTGTGGTGGATGTGAGCCACGAGGACCC
GGTGAAGTCAACTGTGAGCAGCTGGAGGGTGACACAATGCCAAGGCCAACGG
GGAGCAGTACAACAGCACCTACCGGGTGTGCTGACCGTGTGACCCAGGATTGGCT
GAACCGAACGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCCCTACGAGAAA
ATCAAGCAAGGCAAGGCCAGCCCAGAGGCCAACGCTGACCCCTGCCCTAGCAGAGAT
GAGCTGACCAAGAACCCAGGTGTCCTGACCTGCTGTGAAAGGCCCTACCCAGCGACATC
GCCGTTGGAGTGGAGGCAACGGCCAGGCCAGAACAACTACAAGACCAACCCCC
GACAGCGATGGCAGCTTCCCTGACAGCAAGCTGACCGTGGACAAGAGCAGATGGCAGCAG
GCCAACCTGTTCAAGCTGCTGATGACAGCAGGCCCTGCAACATCACTACACCCAGAGGCC
TGAGCCTGTCCCCCTGGCAAG

SEQ ID 136 - S332126E04 murine variable light chain
DIVLTQSPASLAVSLQQRATISCRASESVIHGTHLMHWYQQKPGQPPKLIYASNLESGVPARFSG
SGSETDFTLNIHPVEEEDAATYFCQQSIEDPYTFGGTLEIKR
VAAAPSVFIFPPSDEQLKSGTASVV
CLNNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSLTLKADYEHK
VYACEVTHQ
GLSPVTKSFNRGEC

SEQ ID 137 - S332126E04 murine variable light chain (DNA sequence)
GACATCGTGTGACCCAGTCTCCGCTAGCCTGGCGTGTCTGGCCAGAGGGCCAAATC
AGCTGAGGGCCAGCAGAGCGTCAAGCATTACGGCACCCACCTGATGCACTGGTACCGAG
AACCCCGGCCAGCCTCCAAGCTCTGATCTACGCCGAGCAACCTGGAAAGCGGAGT
GCCAGGTTCAGCGGCAGCGCTCCGAGACCGACTTACCCCTGAAACATCCACCC
GAGGACGCGGCCACTACTCTGCAAGCAGACATGAGGACCCCTACACCTTCGGCGGCC
ACCAAGCTGGAGATCAAGCGTSEQ ID 138 - S332126E04 Chimeric light chain

DIVLTQSPASLAVSLQQRATISCRASESVIHGTHLMHWYQQKPGQPPKLIYASNLESGVPARFSG
SGSETDFTLNIHPVEEEDAATYFCQQSIEDPYTFGGTLEIKR
VAAAPSVFIFPPSDEQLKSGTASVV
CLNNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSLTLKADYEHK
VYACEVTHQ
GLSPVTKSFNRGEC

SEQ ID 139 - S332126E04 Chimeric light chain (DNA sequence)
GACATCGTGTGACCCAGTCTCCGCTAGCCTGGCGTGTCTGGCCAGAGGGCCAAATC
AGCTGAGGGCCAGCAGAGCGTCAAGCATTACGGCACCCACCTGATGCACTGGTACCGAG
AACCCCGGCCAGCCTCCAAGCTCTGATCTACGCCGAGCAACCTGGAAAGCGGAGT
GCCAGGTTCAGCGGCAGCGCTCCGAGACCGACTTACCCCTGAAACATCCACCC
GAGGACGCGGCCACTACTCTGCAAGCAGACATGAGGACCCCTACACCTTCGGCGGCC
ACCAAGCTGGAGATCAAGCGT

GAGGACGCGGCCACTACTCTGCAAGCAGACATGAGGACCCCTACACCTTCGGCGGCC
ACCAAGCTGGAGATCAAGCGTACGGTGCCGCCAGCGTGTCTCATCTCC
GAGCAGCTGAAGCGGGCACCGCAGCGTGTGTCTGCTGAACAACTCTACCC
GCCAAGGTGCAAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
CGAGCAGGACAGCAAGGACTCCACCTACAGCGTGAAGCAGCACCC
CTACGAGAAGCACAAGGTGACGCGCTGTGAGGTGACCCACCAGGGCTGT
CAAGAGCTTCAACCGGGCGAGGTG

SEQ ID 140 - S336105A07 murine variable heavy chain
EVKLQSGGGLVQPGGSLKSCAASGIDF5RYWMSWRRAPGKLEWIGEINPDRSTINYAPSLKD
FIISRDNAKNTLYLQMSKVRSEDTALYYCAVYYDYE GAMDYWQGQGTSVTSS

SEQ ID 141 - S336105A07 murine variable heavy chain (DNA sequence)

GAGGTGAAGCTCTCCAGCTGGAGGTGGCTGGTCAGCCTGGAGGATCCCTGAAACTCTCT
GTCAAGGCCAGGAATCGATTTAGTAGATACTGGATGAGTTGGGCTGGCGGGCTCAGGGAA
AGGACTAGAATGGATTGGAGAAATTAAATCCAGATAGGAGTACAATCAACTATGCACCATCTCTAA
AGGATAAATTCACTCATCTCCAGAGACACAGCCAAAATACGCTGTACCTGCAAATGAGCAAAGTG
AGACTGAGGACACAGCCCTTATTACTGTGAGTTCTACTATGATTACGAGGGTGTATGGA
CTACTGGGTCAAGGAACCTCACTCAGCTACCGTCTCTCAGCAGGAAACACAGGCCACCGTGT
CCCCTGGGCCCCCAGCAGCAGGACCCAGCGCCACAGCGCCCTGGGCTGCCGTGAA
GGACTTCTCCCGCCGCTGCTGAGAGCCACCGTGTGAGGAAACAGCCACACTGCCCTG
CACGAGCAGCCTGGGACCCAGACCTACATCTGAAACGCTGACAGGAAACCCACCTGCCCTG
GTGAGCAGAAGGTTGGAGGCAAGAGCTGTGACAAAGACCCACACTGCCCTG
CCCAGCTGCTGGAGGCCCCAGCGTGTCTGTTCTGCTCCCTGCTGACAGGAAACCCCTGATG
ATCAGCAGAACCCCGAGGTGACCTGTGAGGTTGGATGTGAGCCAGGACCCCTGAGGTG
AAGTTCACTGGTACGGGACCCGGTGGAGGTGCAAAATGCCAAGGCAAGCCAGGGAG
CACTACACAGCACCTACGGGTTGCTCCGTGCTGACCGTGTGACCCAGGATTGGCTGAAC
GGCAAGGAGTACAAGTGTAAAGGTGTCACAAGGCCCCGCTGCTGCCCTATCGAGAAAACCATCA
GCAAGGCCAGGGCCAGCCAGAGAGGCCCCAGGTGTACACCCCTGCCCTAGCAGAGATGAGC
TGACCAAGAACCCAGGTGCTCTGACCTGCCCTGGTAAGGGCTTCAACCCCTAGCAGACATGCCGT

SEQ ID 142 - S336105A07 Chimeric heavy chain

EVKLLQSGGLVQPGGSLKLSCAASGIDFSRYMSWRRAPGKGLEWIGEINPDRSTINYAPSLDK
FISRDNAKNTLYQMSKVRSEDATLYYCAVFFYDYEGAMDYWGQGTSVTVSSAKTAPSVFPLAPS
SKSTSGGTAALGLVLDYFFPEPVTSVMSGALTSGVHTPAVLQSSGLYSLSSVTVPSSSLGTQTYI
CNVNHKPSNTVKDKVKEPKSCDKTHTCPPCPAPELGGPSVFLPPPKPDLMISRTPEVTCVVDV
SHEDPEVKFNWVVDGVEVHNNAKTKPREEQYNSTYRVSVSLTVLHQDWLNGKEYKCKVSNKALPAPI
EKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNQPENNYKTPPVLD
SDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPKG

SEQ ID 143 - S336105A07 Chimeric heavy chain (DNA sequence)

GAGGTGAAGCTCTCCAGCTGGAGGTGGCTGGTCAGCCTGGAGGATCCCTGAAACTCTCT
GTCAAGGCCAGGAATCGATTTAGTAGATACTGGATGAGTTGGGCTGGCGGGCTCAGGGAA
AGGACTAGAATGGATTGGAGAAATTAAATCCAGATAGGAGTACAATCAACTATGCACCATCTCTAA
AGGATAAATTCACTCATCTCCAGAGACACAGCCAAAATACGCTGTACCTGCAAATGAGCAAAGTG
AGACTGAGGACACAGCCCTTATTACTGTGAGTTCTACTATGATTACGAGGGTGTATGGA
CTACTGGGTCAAGGAACCTCACTCAGCTACCGTCTCTCAGCAGGAAACACAGGCCACCGTGT
CCCCTGGGCCCCCAGCAGCAGGACCCAGCGCCACAGCGCCCTGGGCTGCCGTGAA
GGACTTCTCCCGCCGCTGCTGAGAGCCACCGTGTGAGGAAACAGCCACACTGCCCTG
CACGAGCAGCCTGGGACCCAGACCTACATCTGAAACGCTGACAGGAAACCCACCTGCCCTG
GTGAGCAGAAGGTTGGAGGCAAGAGCTGTGACAAAGACCCACACTGCCCTG
CCCAGCTGCTGGAGGCCCCAGCGTGTCTGTTCTGCTCCCTGCTGACAGGAAACCCCTGATG
ATCAGCAGAACCCCGAGGTGACCTGTGAGGTTGGATGTGAGCCAGGACCCCTGAGGTG
AAGTTCACTGGTACGGGACCCGGTGGAGGTGCAAAATGCCAAGGCAAGCCAGGGAG
CACTACACAGCACCTACGGGTTGCTCCGTGCTGACCGTGTGACCCAGGATTGGCTGAAC
GGCAAGGAGTACAAGTGTAAAGGTGTCACAAGGCCCCGCTGCTGCCCTATCGAGAAAACCATCA
GCAAGGCCAGGGCCAGCCAGAGAGGCCCCAGGTGTACACCCCTGCCCTAGCAGAGATGAGC
TGACCAAGAACCCAGGTGCTCTGACCTGCCCTGGTAAGGGCTTCAACCCCTAGCAGACATGCCGT

GGAGTGGAGGACAACGGCCAGGCCAGGAGAACAAACTACAAGACCAACCCCCCTGTGCTGGACAG
CGATGGCAGCTCTCCGTACAGCAAGGCTGACCGTGGACAAGGAGCAGATGGCAGCAGGCCA
CGTGTTCAGCTGCTCGTGATGCCACGAGGCCCTGACAATCACTACACCCAGAAGAGCCTGAGC
CTGCTCCCTGGCAAG

SEQ ID 144 - S336105A07 murine variable light chain

DIVMTQSQKFMSTSVDGRSVTCKASQNVDTNAWVYQQKPGQSPKALIYSASYRFSGVPDRFTGSG
SGTDFLTISNVQSEDLAEYFCQQYNSFPFTFGSGTKEIKRTVAAPSVIFPPSDEQLKSGTASVCL
LNNFYPREAKVQVMVDNALQSGNSQESVTEQDSKDSYLSSTLTSKADYEKHKVYACEVTHQGL
SSPVTKSFRGEC

SEQ ID 145 - S336105A07 murine variable light chain (DNA sequence)

GACATTGTGATGACCCAGTCTCAAAATTCACTGTCCACATCAGTAGGAGACAGGGTCAGCGTCAC
CTCAAGGCCAGTCTCAGAATGTGATTAATGTGAGCTGGTATCAACAAAAACAGGGCAATCTC
CTAAAGCACTGATTACTCGCCTACCCGGTTCACTGGAGTCCCTGATCGCTTCACAGGCAGT
GGATCTGGGACAGATTCTACCTCACCATCAGCAATGTGAGCTGAAAGACTTGGCAGAGTATT
CTGTGAGCAATATAACGCTTCTACCTCACGGTGTGCTGAGGAGGAAAGTTGAAATAAAACGTA
CGGTGGCCGCCAGGGTGTCTGAGCAACACTTCTACCCGGGAGGCCAGGTGAGTGGAAAGGTG
ACAATGCCCTGAGCGCAAGGCCAGGAGAGCTGACCGAGCAGCACAGGACTCCA
CCTACACCTGAGCGCACCCCTGACCTGAGCAAGGCCAGTACAGAGAACACAAGGTGTACG
CCTGTGAGGTGACCCACCAAGGGCTGTGAGCAAGGCCAGTACAGAGAACACAAGGTGTACG
GC

SEQ ID 146 - S336105A07 chimeric light chain

DIVMTQSQKFMSTSVDGRSVTCKASQNVDTNAWVYQQKPGQSPKALIYSASYRFSGVPDRFTGSG
SGTDFLTISNVQSEDLAEYFCQQYNSFPFTFGSGTKEIKRTVAAPSVIFPPSDEQLKSGTASVCL
LNNFYPREAKVQVMVDNALQSGNSQESVTEQDSKDSYLSSTLTSKADYEKHKVYACEVTHQGL
SSPVTKSFRGEC

SEQ ID 147 - S336105A07 chimeric light chain (DNA sequence)

GACATTGTGATGACCCAGTCTCAAAATTCACTGTCCACATCAGTAGGAGACAGGGTCAGCGTCAC
CTGCAAGGCCAGTCTCAGAATGTGATTAATGTGAGCTGGTATCAACAAAAACAGGGCAATCTC
CTAAAGCACTGATTACTCGCCTACCCGGTTCACTGGAGTCCCTGATCGCTTCACAGGCAGT
GGATCTGGGACAGATTCTACCTCACCATCAGCAATGTGAGCTGAAAGACTTGGCAGAGTATT
CTGTGAGCAATATAACGCTTCTACCTCACGGTGTGCTGAGGAGGAAAGTTGAAATAAAACGTA
CGGTGGCCGCCAGGGTGTCTGAGCAACACTTCTACCCGGGAGGCCAGGTGAGTGGAAAGGTG
ACAATGCCCTGAGCGCAAGGCCAGGAGAGCTGACCGAGCAGCACAGGACTCCA
CCTACACCTGAGCGCACCCCTGACCTGAGCAAGGCCAGTACAGAGAACACAAGGTGTACG
CCTGTGAGGTGACCCACCAAGGGCTGTGAGCAAGGCCAGTACAGAGAACACAAGGTGTACG
GC

SEQ ID 148 - S335115G01 murine variable heavy chain

PVQLQPGTELVRPGTSVKLSCKASGYTFTSYWMHWKQRPGQGLEWIGVIDPSDSYTNYNQKF
GKATLTVDTSSSTAYMQLSSLSEDSAVYYCARQVFDYPMDYWGQGTSVTSS

SEQ ID 149 - S335115G01 murine variable heavy chain (DNA sequence)

CCGGTCAACTGAGCAGCCTGGACTGAGCTGGTGGAGGCTGGGACTTCAGTGAAGTTGTCC
TGCAGGGCTTCTGCTACACCTTACCAAGCTACTGGATGCACTGGGTAAGGAGGAGGCCCTGGAC
AAGGCCCTGAGCTGGAGCTGAGTGTAGACACATCCTCCAGCACAGCCTACATGCAGCTCAGCAGCCT
AGGGCAAGGCCACATTGACTGTAGACACATCCTCCAGCACAGCCTACATGCAGCTCAGCAGCCT
GACATCTGAGGACTCTGCGGTCTTACTGTGCAAGACAGGTGTTGACTATCCTATGGACTACT

GGGGTCAAGGAACCTCAGTCACCGTCTCTCA

SEQ ID 150 - S335115G01 Chimeric heavy chain

PVQLQQPGTELVPRGTSVKLSCKASGYTFTSHWMHWKQRPQGQGLEWIGVIDPDSYTNQKFK
GKATLTVDTSSSTAYMLSSLTSEDAVYYCARQVFDPMDYWGQGTLTVSSASTKGPSVFLAP
SSKGTAALGCLVKDYFPEPVTWSNNSALTSVHFTPAVLQSSGLYSLSSVTVPSSSLGTQ

YICVNHHPSNTKVDKKVEPKSCDKTHTCPCPAPELGGPSVLFPPPKDLMISRTPEVTCVV
VSHEDPEVKFNWYWDGVEVHNAKTPREEQYNSTYRVSVLVLHQDWLNKEYKCKVSNKALP
IEKTSKAGQQPREPVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV
SDGSFFLYSKLTVDKSRWQQGNVFSCVMHEALHNHYTQKSLSLSPGK

SEQ ID 151 - S335115G01 Chimeric heavy chain (DNA sequence)

CCGGTCCAACCTGAGCCTGGGACTGAGCTGGTGGAGGCTGGGACTTCAGTGAAGTTGTCC
TGCAAGGCTTCTGGTACACCTTACCAAGGCACTGGATGCACGGTAAAGCAGAGGCTGGAC
AAGGCCCTTGTGGATCGGAGTGATGATCTTGATAGTTACTAACTACAATCAAAGTTCA
AGGGCAAGGCCACATTGACTGTAGACACATCCTCCAGCACAGCCTACATGCAGCTCAGC
GACATCTGAGGACTCTGCGTCTATTACTGTGAAGACAGGTGTTGACTATCCTATGGACTACT
GGGTCAGGAAAGGAACTACTGTGACCCAGCAGCACCAAGGGCCCCAGGGTGTCCCC
TGGGCTCCAGCAGCAGAGCACAGCCGGGCAAGGGCCCTGGCTGGCTGGTGAAGGAC
TACTTCCCAGAACCCGGTACCGTGTCTGGAAACAGCGGAGGCCCTGACCCAGGGCTGCACACC
TTCCCCGGCGTGTGAGCAGCAGCGGCGTGTACAGCCTGAGCAGCGTGGTACCCGTC
AGCAGCCTGGGACCCAGCTACATGTAACTGTGAACGTGAACCACAAGCCAGCAACACCAAGGTG
GACAAGAAGGTGAGGAGCCAAGAGCTGTGACAAAGGACCCACACCTGCCCTGCCCTGCC
GAGCTGTGGGAGGGGGAGGGGGAGGGGGAGGGGGAGGGGGAGGGGGAGGGGGAGGG
AGCAGAACCCCGAGGTGACCTGTGTTGGTGGATGTGAGCCACGGAGGACCTGAGGTGAAG
TTCAACTGGTACGGTGGAGGGTGTGAGCTGACAAATGCCAAGGACCAAGGCCAGGGAGGG
TACAACAGCACCTACGGGTGGTGTGCTGACCGTGTGACCCAGGATTGGCTGAACGGC
AAGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCCCTGCCCTATCGAGAAAACCATCAGCA
AGGCCAACGGGAGGCCAGAGGAGGCCAGGTGACCCCTGCCCTAGCAGAGATGAGCTGA
CCAAGAACAGGTGCCCCGACCTGGCTGGAGACAGACTCACCCCTCAACATCCATCTGGAGGG
AGTGGAGAACGACGGCCAGCCCAGAACAACTACAAGAACCCCCCTGTGCTGGACAGCG
ATGGCACCTCTCTGTACAGCAAGCTGACCGTGGACAAGAGCAGATGGCAGCAGGGCAACG
TGTTCAGCTGCTCCGTGATGACGAGGCCCTGACAAATCACTACACCCAGAACAGGCTGAGCT
GTCCCCCTGGCAAG

SEQ ID 152 - S335115G01 murine variable light chain

DIVLTQSPASLAVSLQQRATISCRASESVSIHGTHLMHWYQQKPGQPPKLLIYASNLESGVPARFSG
SGSETDFTLNIPVVEEADATYFCQQSIEDPWTFGGGTKEIKR

SEQ ID 153 - S335115G01 murine variable light chain (DNA sequence)

GACATTGTGCTGACCCAATCTCAGCTTCTGGCTGTCTCTAGGGCAGAGGGCCACCATCT
CCTCGAGGCCAGTGAAGGTGCACTGTTATCATGCTACTCATTAATGCACTGGTACCAACAGAAA
CCAGGACGCCACCAAACCTCCATCTATGCTGACATCCAACTAGAAATCTGGAGTCCCTGCC
GGTTCACTGGCAGTGGGTCTGAGACAGACTCACCCCTAACATCCATCTGGAGGGAGGGAG
TGTGCAACCTATTCTGTGAGCAAAGTATTGAGGATCCGGTGGACGTTGGAGGGCACAAG
CTGAAATCAATGTCACCGTGGCCGCCAGCGTGTGTCTGCTGAACAACTTCTACCCCGGGAGGCCAAG
TGAAGAGCGGCCACCGCCAGCGTGTGTCTGCTGAACAACTTCTACCCCGGGAGGCCAAG
TGCAGTGGAAAGGTGACAATGCCCTGAGCGGGCACAGCCAGGAGAGCGTGACCGAGCAG
GACAGCAAGGACTCACGCTGAGCAGCACCCCTGACCCCTGAGCAAGGCCACTACAGAG
AAGCACAAGGTGACGCTGTGAGGTGACCCACCGGGCTGTCCAGCCCCGTGACCAAGAGC
TTAACACGGGGCGAGTGC

SEQ ID 156 - S335122F05 murine variable heavy chain

QVQLQQSGAELVPRPGASVTLSCAKASGYTFTDYEMHWKQTPVHGLEWIGAIDPETGGTAYNQKFKG
KAITLADKSSSTAYMELRSLTSEDAVYYCTRSIYDYYFDYWGQGTTLVSSAKTPPSVFLAPSSK

SEQ ID 157 - S335122F05 murine variable heavy chain (DNA sequence)

CAGGTTCAACTGAGCAGCTGGGCTGAGCTGGTGGAGGCTGGGCTTCAGTGAAGCTGTCC
TGCAAGGCTTGGGCTACACATTACTGACTATGAAATGCACTGGGTGAAGCAGACACCTGTGC
ATGCCCTGGAAATGGATTGGAGCTATTGATCTGAAACTGGTGTACTGCCCTACATCAGAAGTTC
AAGGGCAAGGCCACTGACTGAGCAGACAACTCCAGCACAGCCTACATGGAGCTCCGAGCC
TGACATCTGAGGACTCTGCCGTCTATTACTGTACAAGATGATTATGATTACTACTTGTACT
GGGGCCAAGGCCACACTCTCACAGTCTCCTCA

SEQ ID 158 - S335122F05 Chimeric heavy chain

QVQLQQSGAELVPRPGASVTLSCAKASGYTFTDYEMHWKQTPVHGLEWIGAIDPETGGTAYNQKFKG
KAITLADKSSSTAYMELRSLTSEDAVYYCTRSIYDYYFDYWGQGTTLVSSAKTPPSVFLAPSSK
STGGTAALGCLVKDYFPEPVTWSNNSALTSVHFTPAVLQSSGLYSLSSVTVPSSSLGTQTYIC
NVNHHPSNTKVDKKVEPKSCDKTHTCPCPAPELGGPSVFLPPPKDLMISRTPEVTCVVVDVS

EDPEVKFNYYVDGVEVHNAKTPREEQVNSTYRVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE
KTISAKGQPRQEPVYTLPPSRLDETLKNQVSLTCLVKGFPSDIAWEVESNGQPENNYKTPPVLD
DSGFFSYLSKLTKDVRSPWQQNGVFSCVSMEHALHNHTQKSLSPGK

SEQ ID 159 - S335122F05 Chimeric heavy chain (DNA sequence)
CAGGTTCACTGAGCAGTCTGGGCTGAGCTGGTGAAGCTGGGCTTCACTGACGCTGTCC
TGCAGGCTGGGACATCTGGGACTACACATTACTGACTATAAGTGCAGTGGGCTGAGCACACTGTG
ATGGCCTGGAATGGATTGGAGCTATTGATCCTGAAACTGGTGGTACTGGCTTCAACATCAGAAGTTC
AAGGGCAAGGGCATACTGACTGAGCAGACAACTTCAGCAGCACAGCTACATGGAGCTCGCAGC
TGACATCTGAGGACTCTGGCTCTATTACTGACAAGATGATTATTGATTACTACTTGACTACT
GGGGCCAAGGCACCACTCTCACAGTCTCCTCAGCAGGAAACGACACCCCCCAGCGTGTCCCCCT
GGCCCCCAGCAGCAGAGGACCCAGCGGGCACAGCGGCCCTGGCTGCTGGTGAAGGACT
ACTTCCCCGACCCGGTACCGTGTCTGGAAACAGCGGAGGCCCTGACAGCGGGCTGCAACACCT
TCCCCCGCTGGCTCAGAGCACGGGCTGTACAGCTGAGCAGCTGGTCAAGCTGGCAGC
GGAGCCTGGGCCACCCAGACCTACATGTCAGCTGAAACCAAGCCGACACAAAGCTGG
ACAAGAAGGTGGAGCCCCAAGACTGTGACAAGGACCCACACCTGCCCCCTGGCTGCCCCGG
AGCTGCTGGAGGCCCCAGCGTGTCTGGTCCCCCTGGCTGACAAGGCTAAGGACACCTGATGATCA
GCAGAACCCCCGAGGTGACCTGTGTTGGTGTGAGCCACGGGACCCCTGAGGTGAAGT
TCAACTGGTACGTGGACGGCTGGAGGTGCAATGCAAGGCAAGCCAGGGAGGAGCAGT

ACAACAGCACCTACGGGTTGGTGTCCGTGTCAGCGTGTGACCCAGGATTGGCTGAACGGCA
AGGAGTACAAGTGTAAAGGTGTCACAAAGGCCCTGCCTGCCCTATCGAGAAAACCATCAGCA
GGCAAGGGCCAGCCAGAGAGGCCCTGGTACACCCCTGCCCTTAGCAGAGATGAGCTGAC
CAAGAACAGGTGCTCTGACCTGCTGGTAAGGGCTTACCCCAAGGGACATCGCCGTGGA
GTGGAGGAGCAACGGCCAGCCCCAGAGAACAACTACAAGGACACCCCCCTGTGCTGGACAGCGA
TGGCAGCTTCTCTGTACAGCAGGCTGACCCCTGGACAAGGAGCAGATGGCAGCAGGGCAACGT
GTTGAGCTGCTGGTGTGACAGGAGGACCCCTGACAACTACACTACACCCAGAAGAGGCTGAGCCTG
TCCCCCTGGCAAG

SEQ ID 160 - S335122F05 murine variable light chain
DIVLTQSPASLAVSLGQRATISCRASESVSIIHGTHLMHWYQQKPGQQPKLLIYASNLESGVPARFSG
GGSETDFTLNIIHPVEEEDGATYFCQQSIEYPRTFGGGTKLEINR

SEQ ID 161 - S335122F05 murine variable light chain (DNA sequence)
GACATTGTGCTGACCAATCTCCAGCTCTTGGCTGTCTAGGGCAGAGGGCACCATCT
CCTGCAGAGCCAGTGAAAGTGTCACTATTCTATGGTACTCATTTAATGCACGGTACCAACAGAAA
CCAGGACGACGCCACCCAAAATCTCTCATCTATGGCATCCAACTAGAACCTGGAGTCCCTGCCA
GGTTCAGTGGCGGTGGGCTGAGACAGACTTCACCCCTCAACATCCATCTGTGGAGGGAGGAGG
ATGGTCACCACTTCTGTGACGAAAGTATTGAGTATCCTCGGACGTTGGTGGAGGCCACCA
CTGGAAATCAATCGT

SEQ ID 162 - S335122F05 Chimeric light chain
DIVLTQSPASLAVSLQRATISCRASESVIHGTHLMHWYQQKPGQPPKLLIYASNLESGVPARFSG
GGSETDFTNLINHPVEEEDGATYFCQQSIEYPTRFGGTKLEINRTVAAPSVFIFPPSDEQLKSGTASV
CLNNPFYPREAKVQVBDNALQSGNSQESVTEQDSKDSTYSLSSLTLSKADYEKHKVYACEVTHQ
GLSSPVTKSFNRGEC

SEQ ID 163 - S335122F05 Chimeric light chain (DNA sequence)
GACATTGTGCTGACCCAACTCCAGCTCTTGGCTGTCTAGGGCAGAGGGCCACCATCT
CTCTGGAGAGGAGCTGAAAGATGTCGATTCTGGCTGATCTTAACTGGTACAGGAAACAGAAA
CCAGGAGACGCCACCCAAACCTCTCATCTGGTGCATCCAACCTAGAACTGGAGTCCTGCCA
GGTTCACTGGGGTGGGCTGAGACAGACTTCACCCCTAACATCATCTGTGGAGGAGGAGG
ATGGTCAACCTTCTGCAAGAAAGTATTGAGTATCCTGGAGCTTGGTGGAGGCCACCAA
GCTGGAAATCAATCGTAGGGGGCCCGCCCGAGCGTGTTCATCTTCCCCCCCAGCGATGAGCA
GCTGAAGAGCGGCACCGCCAGCGTGGTGTCTGCTGTAACAACCTTACCCCCGGAGGCCAA
GGTGCAGTGGAAAGGTGGACATGCCCCGCAAGCGCAGACAGCCAGGAGAGCGTACCGAGC
AGGACAGCAGGAACTGACCTACAGGCTGAGCAGCAGCCCTGACCCCTGAGCAAGGCCGACTACGG
AAAGCAACAGGTACGGCTGTAGGGTACCCCTGAGGAGGCCACAGGGCTGTCCAGGCCGTCGACCAAGA
GCTTCACCGGGCGAGTC

SEQ.I.D.NO: 164 - S332121F02 CDRH1
DYYNM

SEQ.I.D.NO: 165 - S332121F02 CDRH2
VVDYNGCSTDYNGKES

SEQ.I.D.NO: 166 - S332121F02 CDRH3

SEQ.I.D.NO: 167 - S332121F02 CDRL1

SEQ.I.D.NO: 168 - S332121F02 CDRL2

SEQ.I.D.NO: 169 - S332121F02 CDRL3

SEQ ID NO: 170 - S322110D07 GDRH1

SEQ.I.D.NO: 171 - S322110D07 CDRH2
DIDPNYGDPIYNHKFKG

SEQ.I.D.NO: 172 - S322110D07 CDRH3
RATGTDWFAF

SEQ.I.D.NO: 173 - S322110D07CDRL1
RASENIYNNLA

SEQ.I.D.NO: 174 - S322110D07 CDRL2
AATILAD

SEQ.I.D.NO: 175 - S322110D07 CDRL3
QHFWGTPLT

SEQ.I.D.NO: 176 - S332126E04CDRH1
NYWMH

SEQ.I.D.NO: 177 - S332126E04 CDRH2
IIHPNSGSTNYNEKFKS

SEQ.I.D.NO: 178 - S332126E04 CDRH3
GIYDYPFAY

SEQ.I.D.NO: 179 - S332126E04 CDRL1
RASESVSIHGTHLMH

SEQ.I.D.NO: 180 - S332126E04 CDRL2
AASNLES

SEQ.I.D.NO: 181 - S332126E04 CDRL3
QQSIEDPYT

SEQ.I.D.NO: 182 - S336105A07 CDRH1
RYWMS

SEQ.I.D.NO: 183 - S336105A07 CDRH2
EINPDRSTINYAPSLKD

SEQ.I.D.NO: 184 - S336105A07 CDRH3
FYYDYEGAMDY

SEQ.I.D.NO: 185 - S336105A07 CDRL1
KASQNVDTNVA

SEQ.I.D.NO: 186 - S336105A07 CDRL2
SASYRFS

SEQ.I.D.NO: 187 - S336105A07 CDRL3
QQYNSFPFT

SEQ.I.D.NO: 188 - S335115G01 CDRH1
SYVMH

SEQ.I.D.NO: 189 - S335115G01 CDRH2
VIDPSDSYTNYNQKFKG

SEQ.I.D.NO: 190 - S335115G01 CDRH3
QVFDYPMDY

SEQ.I.D.NO: 191 - S335115G01 CDRL1
RASESVSIHGTHLMH

SEQ.I.D.NO: 192 - S335115G01 CDRL2
AASNLES

SEQ.I.D.NO: 193 - S335115G01 CDRL3
QQSIEDPWT

SEQ.I.D.NO: 194 - S335122F05 CDRH1
DYEMH

SEQ.I.D.NO: 195 - S335122F05 CDRH2
AIDPETGGTAYNQKFKG

SEQ.I.D.NO: 196 - S335122F05 CDRH3
SIYDYYFDY

SEQ.I.D.NO: 197 - S335122F05 CDRL1
RASESVSIHGTHLMH

SEQ.I.D.NO: 198 - S335122F05 CDRL2
AASNLES

SEQ.I.D.NO: 199 - S335122F05 CDRL3
QQSIEYPRT

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO2010104949A2 [0002]
- WO8501533A [0018]
- WO9358679A [0037]
- WO93163930A [0037]
- US4873316A [0038]
- US4816567A [0039]
- EP0239400A [00811]
- EP054951A [00811]
- WO0029004A [0100]
- US20050042519A [0100]
- US7260297B1 [0101]
- US20070224633A [0101]
- EP1641815A1 [0101]
- US20040132023A1 [0102]
- US20080139791A [0103]
- WO2005056794A [0103]
- US6818416B1 [0103]
- WO2008098736A [0105]
- EP0397434A [0110]
- WO2003011676A [0110]
- WO2006014579A [0110]
- EP1229125A [0118]
- US7214775B [0120]
- US6946282B [0120]
- WO0061739A [0120]
- WO20031230A [0120]
- WO2007011041A [0123]
- US20070146165A [0123]
- US4975276A [0131]
- EP1381213A [0131]
- WO9321232A [0134] (Q147)
- US5635463A [0138] (Q1411)
- US5780568A [0138] (Q1411)
- US5662149A [0139]
- WO20081172A [0139]
- US7496286B [0140] (Q1411) (Q1411) (Q154)
- US6884869B [0141]
- US7292302B [01411]
- US7256257B [01411]
- US7423119B [01411]
- US7964566B [01411]
- US3896111A [0143]
- US4151042A [0143]

- [US6573074B](#) [01431]
- [US4137220A](#) [01431]
- [US4248870A](#) [01431]
- [US4256746A](#) [01431]
- [US4260608A](#) [01431]
- [US4265811A](#) [01431]
- [US4294757A](#) [01431]
- [US4307016A](#) [01431]
- [US4308266A](#) [01431]
- [US4308269A](#) [01431]
- [US4309428A](#) [01431]
- [US4313945A](#) [01431]
- [US4316929A](#) [01431]
- [US4317821A](#) [01431]
- [US4322348A](#) [01431]
- [US4331596A](#) [01431]
- [US4361660A](#) [01431]
- [US4364866A](#) [01431]
- [US4424219A](#) [01431]
- [US4450254A](#) [01431]
- [US4362663A](#) [01431]
- [US4371533A](#) [01431]
- [US5208020A](#) [01441] [01441]
- [US6570024B](#) [01441]
- [US6684874B](#) [01441]
- [US5712274A](#) [01461]
- [US5714580A](#) [01461]
- [US5739119A](#) [01461]
- [US5767285A](#) [01461]
- [US5770701A](#) [01461]
- [US57707105773001A](#) [01461]
- [US5677296A](#) [01461] [01461]
- [US5653394A](#) [01461]
- [US5770710A](#) [01461]
- [US6214345B](#) [01521] [01521] [01521]
- [US200300966743A](#) [01521]
- [US20030130189A](#) [01521]
- [US53922852A](#) [01571]
- [US5122368A](#) [01591]
- [US5624606A](#) [01591]
- [US5622929A](#) [01591] [01591]
- [US48802935A](#) [01611]
- [WO2004010957A](#) [01661]
- [US40040302](#) [01661]

Non-patent literature cited in the description

- Antibody targeting of B-cell maturation antigen on malignant plasma cells **RYAN MAUREEN et al.** Molecular Cancer Therapeutics American Association of Cancer Research 20071101 vol. 6, 3009-3018 [00071]
- **PLÜCKTHUN, A.** Immunol. Rev., 1992, vol. 130, 151-188 [00081]
- **MILLER et al.** Genetic Engineering Plenum Press 19860000 vol. 8, 277-298 [00081]
- **REMINGTON'S PHARMACEUTICAL SCIENCE** Mack Publishing Company [00081]
- **LASMAR UPARKINS D** The formulation of Biopharmaceutical products Pharma. Sci. Tech. today, 2000, vol. 3, 129-137 [00081]
- **WANG, W** in stability, stabilisation and formulation of liquid protein pharmaceuticals Int. J. Pharm., 1999, vol. 185, 129-188 [00081]
- Stability of Protein Pharmaceuticals Part A and B Plenum Press 19920000 [00081]
- **AKERS, M.J.** Excipient-Drug interactions in Parenteral Formulations J. Pharm Sci, 2002, vol. 91, 2283-2300 [00081]
- **IMAMURA, K et al.** Effects of types of sugar on stabilization of Protein in the dried state J. Pharm Sci, 2003, vol. 92, 266-274 [00081]
- **IZUTSU, KKOJIMA, S.** Excipient crystallinity and its protein-structure-stabilizing effect during freeze-drying J. Pharm. Pharmacol., 2002, vol. 54, 1033-1039 [00081]
- **JOHNSON, R** Mannitol-sucrose mixtures-versatile formulations for protein peroxidise 19g19n J. Pharm. Sci, 2002, vol. 91, 914-922 [00081]
- **HA, EWANG WWANG Y.J.** Peroxide formation in polysorbate 80 and protein stability J. Pharm Sci, 2002, vol. 91, 2252-2264 [00081]
- **HARLOW et al.** Antibodies A Laboratory Manual Cold Spring Harbor Laboratory 19880000 [00071]
- **MORRISON et al.** Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 6851-6855 [00081]
- **QUEEN et al.** Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 10029-10032 [00081]
- **HODGSON et al.** Bio/Technology, 1991, vol. 9, 421- [00081]

- **KABAT et al.** Sequences of proteins of Immunological Interest NIH, 1987, [\[Q0081\]](#) [\[Q0084\]](#)
- **AL-LAZIKANI et al.** JMB, 1997, vol. 273, 927-948 [\[Q0081\]](#)
- **MACCALLUM R.M.MARTIN A.C.R.THORNTON J.M** Journal of Molecular Biology, 1996, vol. 262, 5732-745 [\[Q0081\]](#)
- **MARTINTHORNTON J** Mol Biol, 1996, vol. 263, 800-815 [\[Q0081\]](#)
- Mol. Immunol., 2006, vol. 44, 656-665 [\[Q1001\]](#)
- Journal of Immunological Methods, 2001, vol. 248, 1-231-45 [\[Q1001\]](#)
- Biochim Biophys Acta, 2000, vol. 1482, 337-350 [\[Q1001\]](#)
- Protein Eng. Des. Sel., 2004, vol. 17, 455-462 [\[Q1001\]](#)
- Nature Biotechnology, 2005, vol. 23, 121556-1561 [\[Q1001\]](#)
- Expert Opinion on Investigational Drugs, 2007, vol. 16, 6909-917 [\[Q1001\]](#)
- J. Biol. Chem, 1999, vol. 274, 24066-24073 [\[Q1001\]](#)
- J. Mol. Biol., 2003, vol. 332, 489-503 [\[Q1002\]](#)
- PNAS, 2003, vol. 100, 41700-1705 [\[Q1002\]](#)
- J. Mol. Biol., 2007, vol. 369, 1015-1028 [\[Q1002\]](#)
- Protein Eng. Des. Sel., 2005, vol. 18, 435-444 [\[Q1003\]](#)
- Expert Opin. Biol. Ther., 2005, vol. 5, 783-797 [\[Q1004\]](#)
- Non-Antibody Scaffolds from Handbook of Therapeutic Antibodies 20070000 [\[Q1005\]](#)
- Protein Science, 2006, vol. 15, 14-27 [\[Q1006\]](#)
- **SHIELDS et al.** The Journal of Biological Chemistry, 2001, vol. 276, 6591-6604 [\[Q112\]](#)
- **CHAPPEL et al.** The Journal of Biological Chemistry, 1993, vol. 268, 25124-25131 [\[Q112\]](#)
- **LAZAR et al.** PNAS, 2006, vol. 103, 4005-4010 [\[Q116\]](#) [\[Q116\]](#)
- J Imm Meth, 1995, vol. 184, 29-38 [\[Q113\]](#)
- **DUNCAN et al.** Nature, 1988, vol. 332, 563-564 [\[Q114\]](#)
- **LUND et al.** J. Immunol., 1991, vol. 147, 2657-2662 [\[Q114\]](#)
- **CHAPPEL et al.** PNAS, 1991, vol. 88, 9036-9040 [\[Q114\]](#)
- **BURTONWOOF** Adv. Immunol., 1992, vol. 51, 1-84 [\[Q114\]](#)
- **MORGAN et al.** Immunology, 1995, vol. 86, 319-324 [\[Q114\]](#)
- **HEZAREH et al.** J. Virol., 2001, vol. 75, 2412161-12168 [\[Q114\]](#)
- **SHIELDS et al.** J Biol Chem, 2001, vol. 276, 6591-6604 [\[Q116\]](#)
- **NECHANSKY et al.** Mol Immunol, 2007, vol. 44, 1815-1817 [\[Q116\]](#)
- **LAMBERT, J.** Curr. Opinion in Pharmacology, 2005, vol. 5, 543-549 [\[Q131\]](#)
- **WU et al.** Nature Biotechnology, 2005, vol. 23, 91137-1146 [\[Q131\]](#)
- **SYRIGOSEPENETOS** Anticancer Research, 1999, vol. 19, 605-614 [\[Q131\]](#)
- **NICULESCU-DUVAZSPRINGER** Adv. Drug Deliv. Rev., 1997, vol. 26, 151-172 [\[Q131\]](#)
- **BALDWIN et al.** Lancet, 1986, 603-05 [\[Q131\]](#)
- Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review **THORPE et al.** Monoclonal Antibodies '84: Biological And Clinical Applications 19850000475-506 [\[Q131\]](#)
- **ROWLAND et al.** Cancer Immunol. Immunother., 1986, vol. 21, 183-87 [\[Q131\]](#)
- **MANDLER et al.** J. Nat. Cancer Inst., 2000, vol. 92, 191573-1581 [\[Q131\]](#)
- **MANDLERET** Bioorganic & Med. Chem. Letters, 2000, vol. 10, 1025-1028 [\[Q131\]](#)
- **MANDLER et al.** Bioconjugate Chem., 2002, vol. 13, 786-791 [\[Q131\]](#)
- **LIU et al.** Proc. Natl. Acad. Sci. USA, 1996, vol. 93, 8618-8623 [\[Q131\]](#)
- **LODE et al.** Cancer Res., 1998, vol. 58, 2928- [\[Q131\]](#)
- **HINMAN et al.** Cancer Res., 1993, vol. 53, 3336-3342 [\[Q131\]](#)
- **WOYKE et al.** Antimicrob. Agents and Chemother, 2001, vol. 45, 123580-3584 [\[Q138\]](#)
- **PETTIT et al.** Antimicrob. Agents Chemother., 1998, vol. 42, 2961-2965 [\[Q138\]](#)
- **E. SCHRODERK. LUBKE** The Peptides Academic Press 19650000 vol. 1, 76-136 [\[Q141\]](#)
- **PETTIT et al.** J. Am. Chem. Soc., 1989, vol. 111, 5463-5465 [\[Q141\]](#)
- **PETTIT et al.** Anti-Cancer Drug Design, 1998, vol. 13, 243-277 [\[Q141\]](#)
- **PETTIT, G. R. et al.** Synthesis, 1996, 719-725 [\[Q141\]](#)
- **PETTIT et al.** J. Chem. Soc. Perkin Trans., 1996, vol. 15, 859-863 [\[Q141\]](#)
- **DORONINA** Nat Biotechnol, 2003, vol. 21, 7778-784 [\[Q141\]](#)
- Monomethylvaline Compounds Capable of Conjugation to Ligands, [\[Q141\]](#)
- **HINMAN et al.** Cancer Research, 1993, vol. 53, 3336-3342 [\[Q141\]](#)
- **LODE et al.** Cancer Research, 1998, vol. 58, 2925-2928 [\[Q145\]](#)
- **FRAKER et al.** Biochem. Biophys. Res. Commun., 1978, vol. 80, 49-57 [\[Q150\]](#)
- **CHATAL** Monoclonal Antibodies in Immunoscintigraphy CRC Press 19890000 [\[Q150\]](#)
- **GEOGHEGAN & STROH** Bioconjugate Chem., 1992, vol. 3, 138-146 [\[Q157\]](#)
- **DUBOWCHIKWALKER** Pharm. Therapeutics, 1999, vol. 83, 67-123 [\[Q158\]](#) [\[Q158\]](#)
- **NEVILLE et al.** Biol. Chem., 1989, vol. 264, 14653-14661 [\[Q160\]](#)
- **THORPE et al.** Cancer Res., 1987, vol. 47, 5924-5931 [\[Q161\]](#)
- **WAWRZYNCKA et al.** Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer Oxford U. Press 19870000 [\[Q161\]](#)
- **JOHNSON et al.** Anticancer Res., 1995, vol. 15, 1387-93 [\[Q162\]](#)
- **LAU et al.** Bioorg-Med-Chem., 1995, vol. 3, 101299-1304 [\[Q162\]](#)
- **LAU et al.** Bioorg-Med-Chem., 1995, vol. 3, 101305-12 [\[Q162\]](#)

Patentkrav

1. Anti-B-cellemodningsantigen-(CD269)-antistof omfattende en tungkæde variabel region af SEQ. ID. NO:23 og en letkæde variabel 5 region af SEQ. ID. NO:31.
2. Anti-B-cellemodningsantigen-(CD269)-antistof ifølge krav 1 som omfatter en tungkæde af SEQ. ID. NO:55 og en letkæde af SEQ. ID. NO:63.
- 10 3. Anti-B-cellemodningsantigen-(CD269)-antistof ifølge et hvilket som helst foregående krav, hvor antistoffet er et monoklonalt antistof.
4. Anti-B-cellemodningsantigen-(CD269)-antistof ifølge et hvilket som 15 helst foregående krav, hvor antistoffet har forbedret binding til Fc_YRIIIA eller har forbedret Fc_YRIIIA-medieret effektorfunktion.
5. Anti-B-cellemodningsantigen-(CD269)-antistof ifølge krav 4, hvor antistoffet er defucosyleret.
- 20 6. Nukleinsyresekvenser der koder for både de tungkæde variable og letkæde variable sekvenser af antistoffet ifølge et hvilket som helst foregående krav.
- 25 7. Nukleinsyresekvenser omfattende SEQ ID NO 56 der koder for tungkæden; og SEQ ID NO: 64 der koder for letkæden af antistoffet ifølge et hvilket som helst foregående krav.
- 30 8. Ekspressionsvektor omfattende nukleinsyresekvenserne ifølge krav 6 eller krav 7.

9. Rekombinant transformerede eller transfekterede værtscelle omfattende nukleinsyresekvenserne ifølge krav 6 eller krav 7, eller ekspressionsvektoren ifølge krav 8, hvor værtscellen omfatter nukleinsyresekvenser der koder for både tung- og letkæde.

5

10. Fremgangsmåde til fremstillingen af et antistof, hvilken fremgangsmåde omfatter at dyrke værtscellen ifølge krav 9 under betingelser egnet til ekspression af nævnte nukleinsyresekvenser eller vektor(er), hvorved antistoffet fremstilles.

10

11. Antistof som fremstillet med fremgangsmåden ifølge krav 10.

12. Immunokonjugat omfattende antistoffet ifølge et hvilket som helst af kravene 1 til 5 eller krav 11 og et cytotoxiske middel.

15

13. Immunokonjugat ifølge krav 12, hvor det cytotoxiske middel er monomethyl-auristatin E (MMAE) eller monomethyl-auristatin F (MMAF).

20

14. Immunokonjugat af krav 13, hvor det cytotoxiske middel er monomethyl-auristatin F (MMAF).

15. Immunokonjugat ifølge et hvilket som helst af kravene 12-14, hvor antistoffet er forbundet til det cytotoxiske middel via en linker.

25

16. Immunokonjugat af krav 15, hvor linkeren er valgt fra 6-maleimidocaproyl (MC), maleimidopropanoyl (MP), valin-citrullin (val-cit), alanin-phenylalanin (ala-phe), p- aminobenzylloxycarbonyl (PAB), N-Succinimidyl 4-(2-pyridylthio)pentanoat (SPP), N-succinimidyl 4-(N-maleimidomethyl)cyclohexan-1 carboxylat (SMCC), og N-Succinimidyl (4-iodo-acetyl) aminobenzoat (SIAB).

30

17. Immunokonjugat af krav 16, hvor linkeren er 6-maleimidocaproyl

(MC).

18. Immunokonjugat omfattende et anti-B-cellemodningsantigen-(CD269)-antistof og et cytotoxiske middel, hvor anti-B-
5 cellemodningsantigen-(CD269)-antistoffet har forbedret binding til FcyRIIIA eller har forbedret FcyRIIIA-medieret effektorfunktion og omfatter tungkædeaminosyresekvensen i henhold til SEQ ID NO: 55 og letkædeaminosyresekvensen i henhold til SEQ ID NO:63, og hvor antistoffet er defucosyleret og hvor det cytotoxiske middel er
10 monomethyl-auristatin F (MMAF) og hvor nævnte MMAF er forbundet til nævnte antistof via en 6-maleimidocaproyl(MC)-linker.

19. Farmaceutisk sammensætning omfattende antistoffet ifølge et hvilket som helst af kravene 1-5 eller krav 11 eller immunokonjugatet
15 ifølge et hvilket som helst af kravene 12-18 og en farmaceutisk acceptabel bærer.

20. Antistof ifølge et hvilket som helst af kravene 1-5 og 11, et immunokonjugat ifølge et hvilket som helst af kravene 12-18, eller en farmaceutisk sammensætning ifølge krav 19 til anvendelse i behandling af en menneskelig patient ramt af en B-cellesygdom eller -lidelse valgt fra multipel myelom (MM), kronisk lymfocytisk leukæmi (CLL), Non-sekretorisk multipel myelom, ulmende (*Smoldering*) multipel myelom, monoklonal gammopathi af ubestemt signifikans (MGUS), solitært plasmacytom (knogle, ekstramedullært), lymfoplasmacytisk lymfom (LPL), Waldenströms makroglobulinæmie, plasmacelleleukæmi, primær amyloidose (AL), tungkædesygdom, systemisk lupus erythematosus (SLE), POEMS syndrom/osteosklerotisk myeloma, Type I og II kryoglobulinæmie, letkædeaflejringssygdom, Goodpasture-syndrom, 25 idiopatisk trombocytopenisk purpura (ITP), akut glomerulonefritis, pemphigus- og pemphigoidlidelser, og epidermolysis bullosa acquisita; eller enhver non-Hodgkins lymfom B-celleleukæmi eller Hodgkins

lymfom (HL).

21. Antistoffet, immunokonjugatet eller den farmaceutiske sammensætning til anvendelse ifølge krav 20, hvor B-5 cellesygdommen er multipel myelom (MM).

DRAWINGS

Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:





Figure 6:

A- Binding to ARH-77 10B5 cells



B - Binding to H929 cells






Figure 7:

(A)



(B)



(C)





Figure 8:

(A)



(B)





Figure 9:


J5 variants



■ J5M0
▲ J5M1
● J5M2

EC50 | J5M0 0.2087 J5M1 0.3045 J5M2 0.1822


J6 variants



■ J6M0
▲ J6M1
▼ J6M2

EC50 | J6M0 0.09187 J6M1 0.04860 J6M2 0.1364


J8 variants



■ J8M0
▲ J8M1
● J8M2

EC50 | J8M0 0.07106 J8M1 0.1734 J8M2 0.06510


J7 variants



■ J7M0
▲ J7M1
● J7M2

EC50 | J7M0 0.05386 J7M1 0.05761 J7M2 0.1506

J9 variants




■ J9M0
▲ J9M1
● J9M2


EC50 | J9M0 0.04738 J9M1 0.03225 J9M2 0.1170

Figure 10:


A



B



C



D




Figure 11:




Figure 12:




Figure 13:




Figure 14:



A**B**

Figure 15:




Figure 16:




Figure 17:





Figure 18:

A



B




Figure 19:





Figure 20:



SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

