



(86) Date de dépôt PCT/PCT Filing Date: 2005/11/17
(87) Date publication PCT/PCT Publication Date: 2006/05/26
(85) Entrée phase nationale/National Entry: 2007/05/14
(86) N° demande PCT/PCT Application No.: US 2005/041511
(87) N° publication PCT/PCT Publication No.: 2006/055625
(30) Priorité/Priority: 2004/11/18 (US60/628,930)

(51) Cl.Int./Int.Cl. *C07D 307/79* (2006.01),
A61K 31/34 (2006.01), *A61P 3/00* (2006.01),
C07C 311/02 (2006.01), *C07D 307/91* (2006.01)


(71) Demandeur/Applicant:
THE INSTITUTES FOR PHARMACEUTICAL
DISCOVERY, LLC, US

(72) Inventeurs/Inventors:
VAN ZANDT, MICHAEL C., US;
FANG, HAIQUAN, US;
HU, SHAOJING, US;
WHITEHOUSE, DARREN, US

(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : ACIDES CARBOXYLIQUES A SUBSTITUTION PHENYLE

(54) Title: HETEROCYCLYLBIHENYL DERIVATES AS PROTEIN TYROSINE PHOSPHATASE INHIBITORS



(57) Abrégé/Abstract:

Disclosed are compounds and pharmaceutically acceptable salts of formula (A): which are useful in the treatment of metabolic disorders related to insulin resistance, leptin resistance, or hyperglycemia. Compounds of the invention include inhibitors of Protein tyrosine phosphatases, in particular Protein tyrosine phosphatase-1B (PTP-1B), that are useful in the treatment of diabetes and other PTP mediated diseases, such as cancer, neurodegenerative diseases and the like. Also disclosed are pharmaceutical compositions comprising compounds of the invention and methods of treating the aforementioned conditions using such compounds.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 May 2006 (26.05.2006)

PCT

(10) International Publication Number
WO 2006/055625 A3

(51) International Patent Classification:

C07D 307/79 (2006.01) *A61K 31/34* (2006.01)
C07D 307/91 (2006.01) *A61P 3/00* (2006.01)
C07C 311/02 (2006.01)

(21) International Application Number:

PCT/US2005/041511

(22) International Filing Date:

17 November 2005 (17.11.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/628,930 18 November 2004 (18.11.2004) US

(71) Applicant (for all designated States except US): **THE INSTITUTES FOR PHARMACEUTICAL DISCOVERY, LLC** [US/US]; 23 Business Park Drive, Branford, Connecticut 06405 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VAN ZANDT, Michael, C.** [US/US]; 91 Golden Hill Drive, Guilford, Connecticut 06437 (US). **FANG, Haiquan** [US/US]; 304 County Rd., Madison, Connecticut 06443 (US). **HU, Shaojing** [US/US]; 900 Mix Ave., Unit 121, Hamden, Connecticut 06514 (US). **WHITEHOUSE, Darren** [US/US]; 40 Fishing Brook Rd., Westbrook, Connecticut 06498 (US).

(74) Agent: **SARUSSI, Steven, J.**; McDonnell Boehnen Hulbert & Berghoff LLP, 300 South Wacker Drive, Chicago, Illinois 60606 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

30 November 2006

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLYLBIHENYL DERIVATES AS PROTEIN TYROSINE PHOSPHATASE INHIBITORS



(A)

aforementioned conditions using such compounds.

(57) Abstract: Disclosed are compounds and pharmaceutically acceptable salts of formula (A): which are useful in the treatment of metabolic disorders related to insulin resistance, leptin resistance, or hyperglycemia. Compounds of the invention include inhibitors of Protein tyrosine phosphatases, in particular Protein tyrosine phosphatase-1B (PTP-1B), that are useful in the treatment of diabetes and other PTP mediated diseases, such as cancer, neurodegenerative diseases and the like. Also disclosed are pharmaceutical compositions comprising compounds of the invention and methods of treating the

WO 2006/055625 A3

Phenyl Substituted Carboxylic Acids

BACKGROUND OF THE INVENTION

This application claims priority from U.S. Provisional Application Serial No. 60/628930, which was filed November 18, 2004, the disclosure of which is incorporated herein by reference in its entirety.

Field of the Invention

The invention relates to phenyl substituted carboxylic acids and more specifically to such compounds that are useful in the treatment of syndrome X (consisting of such abnormalities as obesity, dyslipidemia, hypercoagulation, hypertension, insulin resistance and leading to heart disease and diabetes), obesity, diabetes, immunological disease, bleeding disorders and/or cancer. More specifically, it relates to such compounds that are capable of inhibiting Protein tyrosine phosphatases (PTPs), in particular Protein tyrosine phosphatase-1B (PTP-1B) which is a negative regulator of the insulin and leptin signaling pathway and improves insulin sensitivity.

Description of the Related Art

This invention relates to a class of heterocycle substituted carboxylic acids that are inhibitors of various PTPs, in particular PTP-1B.

Protein tyrosine phosphatases are a large family of transmembrane or intracellular enzymes that dephosphorylate substrates involved in a variety of regulatory processes (Fischer et al., 1991, Science 253:401-406). Protein tyrosine phosphatase-1B (PTP-1B) is an approximately 50 kd intracellular protein, which is present in abundant amounts in various human tissues (Charbonneau et al., 1989, Proc. Natl. Acad. Sci. USA 86:5252-5256; Goldstein, 1993, Receptor 3:1-15).

Determining which proteins are substrates of PTP-1B has been of considerable interest. One substrate which has aroused

especial interest is the insulin receptor. The binding of insulin to its receptor results in autophosphorylation of the domain. This causes activation of the insulin receptor tyrosine kinase, which phosphorylates the various insulin receptor substrate (IRS) proteins that propagate the insulin signaling event further downstream to mediate insulin's various biological effects.

Seely et al., 1996, Diabetes 45:1379-1385 ("Seely") studied the relationship of PTP-1B and the insulin receptor in vitro. Seely constructed a glutathione S-transferase (GST) fusion protein of PTP-1B that had a point mutation in the PTP-1B catalytic domain. Although catalytically inactive, this fusion protein was able to bind to the insulin receptor, as demonstrated by its ability to precipitate the insulin receptor from purified receptor preparations and from whole cell lysates derived from cells expressing the insulin receptor.

Ahmad et al., 1995, J. Biol. Chem. 270:20503-20508 used osmotic loading to introduce PTP-1B neutralizing antibodies into rat KRC-7 hepatoma cells. The presence of the antibody in the cells resulted in an increase of 42% and 38%, respectively, in insulin stimulated DNA synthesis and phosphatidylinositol 3' kinase activity. Insulin receptor autophosphorylation and insulin receptor substrate-1 tyrosine phosphorylation were increased 2.2 and 2.0-fold, respectively, in the antibody-loaded cells. The antibody-loaded cells also showed a 57% increase in insulin stimulated insulin receptor kinase activity toward exogenous peptide substrates.

Kennedy et al., 1999, Science 283: 1544-1548 showed that protein tyrosine phosphatase PTP-1B is a negative regulator of the insulin signaling pathway, indicating that inhibitors of this enzyme are beneficial in the treatment of Type 2 diabetes, which appears to involve a defect in an early process in insulin signal transduction rather than a structural defect in

the insulin receptor itself. (J. M. Olefsky, W. T. Garvey, R. R. Henry, D. Brillon, S. Matthai and G. R. Freidenberg, G. R. (1988).) Cellular mechanisms of insulin resistance in non-insulin-dependent (Type II) diabetes. (Am. J. Med. 85: Suppl. 5A, 86-105.) A drug that improved insulin sensitivity would have several advantages over traditional therapy of NIDDM using sulfonylureas, which do not alleviate insulin resistance but instead compensate by increasing insulin secretion.

10 Ragab et al (2003, J. Biol. Chem 278(42), 40923-32) showed that PTP 1B is involved in regulating platelet aggregation. Hence, inhibition of PTP 1B can be predicted to have an effect on bleeding disorder, and cardiovascular disease.

Romsicki et al., (2003, Arch Biochem. Biophys 414(1), 40-50) showed that TC PTP is structurally and functionally very similar. A PTP 1B inhibitor is very likely to also inhibit TC PTP. A knockout of the TC PTP gene produces a phenotype with impaired immune function. (You-Ten et al., 1997, J. Exp. Med. 186(5), 683-93). Hence, inhibitors of PTP 1B can be predict to inhibit TC PTP and modulate immune response.

20 It has also been demonstrated that PT-P1B is a negative regulator of leptin signaling (Kaszua et al. MolCell. Endocrinology, 195:109-118, 2002). PTP-1B deficient mice show enhanced potency for exogenous leptin to suppress food intake (Cheng, et al. Developmental Cell 2:497-503, 2002). Thus, inhibitors of PTP-1B augment the beneficial effects of leptin on food intake, body weight regulation and metabolism, in normal individuals and leptin resistant individuals.


Therefore, inhibitors of PTPs, and inhibitors of PTP-1B in particular, are useful in controlling or treating obesity, syndrome X, Type 2 diabetes, in improving glucose tolerance, and in improving insulin sensitivity in patients in need thereof. Such compounds are also useful in treating or controlling other PTP mediated diseases, such as the treatment

of cancer, neurodegenerative diseases, immunological disorders, bleeding and cardiovascular disorders, and the like.

SUMMARY OF THE INVENTION

In a broad aspect, the invention encompasses the compounds of formula (A) shown below, pharmaceutical compositions containing the compounds and methods employing such compounds or compositions in the treatment of diabetes and/or cancer.

The invention provides compounds of formula A:



(A)

- 10 and pharmaceutically acceptable salts thereof, wherein
 R₁ is H, C₁-C₆ alkyl, phenyl(C₁-C₆)alkyl, or C₂-C₆ alkenyl;
 L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -(C₁-C₄)alkyl-
 N(R₁₀)C(O)-, -C(O)N(R₁₀)-(C₁-C₄)alkyl-, -N(R₁₀)C(O)-(C₁-
 C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-(C₁-C₆)alkyl-, -CO-
 15 , -SO₂-, or -(C₁-C₆)alkyl-O-;
 L₃ is absent, a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)
 alkyl-, -alkenyl-, -phenyl-;
 L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -C(O)N(R₉)-(C₁-
 C₄)alkyl-, -N(R₉)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-
 20 C(O)N(R₉)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-N(R₉)C(O)-(C₁-C₄)
 alkyl-, -N(R₉)-(C₁-C₆)alkyl-, -N(R₉)-(C₁-C₆)alkyl- wherein
 -(C₁-C₆)alkyl- is optionally substituted with phenyl, -
 (C₁-C₄)alkyl-N(R₉)-(C₁-C₄)alkyl-, -SO₂N(R₉)-, -SO₂N(R₉)-
 (C₁-C₄)alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -N(R₉)SO₂-, -(C₁-C₄)
 25 alkyl-, -(C₂-C₆)alkenyl-, -N(R₉)C(O)-, -C(O)-(C₁-C₄)
 alkyl-, -S-(C₁-C₄)alkyl-, -(C₁-C₆)alkyl-S-, -or -(C₁-C₄)
 alkyl-S-(C₁-C₄)alkyl-, wherein each alkyl and alkenyl is
 optionally substituted with one or more R₁₁ groups,
 R₉ and R₁₀ are independently H, C₁-C₆ alkyl, C₁-C₆
 30 alkoxycarbonyl, -SO₂-aryl, heteroarylalkyl,

arylalkyl, wherein the aryl or heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl, or haloalkoxy;

R₁₁ at each occurrence is independently N₁₂R₁₃, -

N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂ and R₁₃ are independently H or C₁-C₆ alkyl,

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H,

arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO₂, NH₂, CN, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-aryl, NHC(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄)alkyl-aryl, -NHSO₂-aryl, -N(C₁-C₄alkyl)SO₂aryl, wherein the aryl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, haloalkyl, haloalkoxy;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;

Q is H, aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl, -heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl,

heterocycloalkylcarbonyl, $-C(O)NH_2$, $-C(O)NH(C_1-C_6)alkyl$, $-C(O)N(C_1-C_6)alkyl(C_1-C_6)alkyl$, or $-SO_2-aryl$, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C_1-C_4 alkyl, C_1-C_4 alkoxy, NO_2 , OH , NH_2 , $NH(C_1-C_6)alkyl$, $N(C_1-C_6)alkyl(C_1-C_6)alkyl$, haloalkyl or haloalkoxy;

and

Z is absent, H , $-NHC(O)aryl$, $-N(C_1-C_4 alkyl)C(O)aryl$, or $aryl$ (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, haloalkyl, haloalkoxy, or NO_2 , or

Z is $-NHC(O)-(C_1-C_4)alkyl-(C_3-C_7)cycloalkyl$, $-N(C_1-C_4)alkylC(O)-(C_1-C_4)alkyl-(C_3-C_7)cycloalkyl$.

The compounds of formula A bind to PTPs, and in particular to PTP-1B. The interaction with the enzyme, specifically PTP-1B, preferably results in inhibition of the enzyme.

The invention also includes intermediates that are useful in making the compounds of the invention.

The invention also provides pharmaceutical compositions comprising a compound or salt of formula A and at least one pharmaceutically acceptable carrier, solvent, adjuvant or diluent.

The invention further provides methods of treating disease such as diabetes, syndrome X, cancer, immunological disease, bleeding disorders, or cardiovascular disease in a patient in need of such treatment, comprising administering to the patient a compound or pharmaceutically acceptable salt of formula A, or a pharmaceutical composition comprising a compound or salt of formula A.

In another aspect, the invention provides a method for inhibiting protein tyrosine phosphatases, preferably PTP-1B,

comprising administering a therapeutically effective amount of a compound of formula A.

In another aspect, the invention provides a method for treating metabolic disorders related to insulin resistance or hyperglycemia, comprising administering to a patient in need of such treatment a therapeutically effective amount of a compound of formula A.


The invention also provides the use of a compound or salt according to formula A for the manufacture of a medicament for use in treating diabetes or cancer or other diseases related to PTP.

The invention also provides methods of preparing the compounds of the invention and the intermediates used in those methods.

The invention also provides methods and compositions for combination therapy of Type I and Type II diabetes. In these embodiments, the invention provides formulations and pharmaceutical compositions, as well as methods for treating Type I and Type II diabetes with the compounds of formula A plus additional compounds and medicaments as disclosed in more detail below. In these embodiments, the methods of the invention can comprise treatment methods for Type I and Type II diabetes where the compounds of formula A are formulated with a therapeutically-effective amount of said additional compounds and medicaments. In alternative embodiments, treatment methods of the invention for Type I and Type II diabetes comprise administration of the inventive compounds of formula A as disclosed herein concomitantly, simultaneously or together with a therapeutically-effective amount of said additional compounds and medicaments.

DETAILED DESCRIPTION OF THE INVENTION

A preferred class of compounds of formula A are compounds of formula I:



(I)

Preferred compounds of formula I are compounds of formula I-a, wherein,

Q is H, phenyl, -phenyl-O-phenyl, -phenyl-carbonyl-phenyl, -phenyl-(C₁-C₄)alkyl-phenyl, -phenyl-pyridyl, -phenyl-pyrimidyl, -phenyl-benzofuranyl, -phenyl-indolyl, -phenyl-piperidinyl, -phenyl-pyrrolidinyl, -phenyl-piperazinyl, indoliziny, benzofuranyl, indolyl, dibenzofuranyl, -benzothienyl-(C₁-C₄)alkyl-phenyl, -indolyl-(C₁-C₄)alkyl-phenyl, benzofuranyl-(C₁-C₄)alkyl-phenyl, piperidinyl, pyrrolidinyl, tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or imidazo[2,1-b]thiazol-3-one, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, or phenyl-(C₁-C₆)alkyl-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, phenyl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, phenyl(C₂-C₆)alkanoyl, C₁-C₆ alkoxycarbonyl, phenyl(C₁-C₆)alkoxycarbonyl, heteroarylcarbonyl, heteroaryl, heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, or phenyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, or NO₂.

5 Preferred compounds of formula I are those wherein L₂ and L₃ are in a 1 to 3 positional relationship on the A ring.

Preferred are also compounds wherein L₂ and L₅ are para to each other on the bridging phenylene.

10 Further preferred are compounds wherein L₂ and L₅ are meta to each other on the bridging phenylene.

A preferred class of compounds of formula I-a are compounds of formula I-b, wherein,

15 the A ring is phenyl, pyrido[1,2-a]indolyl, furanyl, thienyl, benzofuranyl, dibenzofuranyl, indolyl, thiazolyl, thiazolidinyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

20 R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H, phenyl(C₁-C₆)alkoxy, phenyl(C₁-C₆)alkyl, halogen, (C₁-C₆)alkyl, OH, alkoxy, CN, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-phenyl, NHC(O)-(C₁-C₄) alkyl-phenyl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-phenyl, N(C₁-C₄)alkyl-phenyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂phenyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy.

30

A preferred class of compounds of formula I-b are compounds of formula I-c, wherein,

R₁ is H, C₁-C₆ alkyl, benzyl, or allyl;

L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -(C₁-C₄)alkyl-N(R₁₀)C(O)-, -C(O)N(R₁₀)-(C₁-C₄)alkyl-, -N(R₁₀)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-(C₁-C₆)alkyl-, or -(C₁-C₆)alkyl-O-;

5 L₃ is absent, a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-, -alkenyl-, or -phenyl-;

L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -C(O)N(R₉)-(C₁-C₄)alkyl-, -N(R₉)C(O)-(C₁-C₄)alkyl-, -N(R₉)-(C₁-C₆)alkyl-, -N(R₉)-(C₁-C₆)alkyl- wherein -(C₁-C₆)alkyl- is
 10 optionally substituted with phenyl, -(C₁-C₄)alkyl-N(R₉), -(C₁-C₄)alkyl-N(R₉)-(C₁-C₄)alkyl-, -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄)alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -N(R₉)SO₂-, -(C₁-C₄)alkyl-, -(C₂-C₆)alkenyl-, -N(R₉)-(C₁-C₆)alkyl- wherein -(C₁-C₆)alkyl- is optionally substituted with phenyl,
 15 -N(R₉)C(O)-, -C(O)-(C₁-C₄)alkyl-, -S-(C₁-C₄)alkyl-, or -(C₁-C₄)alkyl-S-(C₁-C₄)alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups,

R₉ and R₁₀ are independently is H, C₁-C₆ alkyl, C₁-C₆
 20 alkoxycarbonyl, -SO₂phenyl, -C₁-C₆alkyl-furanyl, -C₁-C₆alkyl-tetrazolyl, -C₁-C₆alkyl thienyl, -C₁-C₆alkyl pyrrolyl, -C₁-C₆alkyl pyridyl, benzyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups
 25 that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, C₁-C₄ haloalkyl, or C₁-C₄ haloalkoxy

R₁₁ at each occurrence is independently -N₁₂R₁₃, -
 30 N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂ and R₁₃ are independently H or C₁-C₆ alkyl.

A preferred class of compounds of formula I-c are compounds of formula I-d, wherein,

L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-(C₁-C₆)alkyl-, or -(C₁-C₆)alkyl-O-;

5 the A ring is phenyl, pyrido[1,2-a]indolyl, furanyl, thienyl, indolyl, thiazolyl, thiazolidinyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or
10 N(C₁-C₆)alkyl(C₁-C₆)alkyl;

Z is phenyl, optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, or NO₂.

15 A preferred class of compounds of formulas I-c or I-d are compounds of formula I-e, wherein,

R₂₂ and R₂₃ are both H;

L₅ is a bond, -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄) alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-,
20 -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl, -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, or -N(R₉)SO₂-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups; and
Q is phenyl, -phenyl-O-phenyl, indoliziny, benzofuranyl, indolyl, dibenzofuranyl, -benzothienyl-(C₁-C₄)alkyl-phenyl, -indolyl-(C₁-C₄)alkyl-phenyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or benzofuranyl-(C₁-C₄) alkyl-phenyl, wherein the
25 aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently
30 alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, or phenyl-(C₁-C₆)alkyl-.

A preferred class of compounds of formula I-e are compounds of formula I-f, wherein,
 R₉ is H, C₁-C₆ alkyl, C₁-C₆ alkoxy, -SO₂phenyl, -C₁-C₆ alkyl-furanyl, -C₁-C₆ alkyl-tetrazolyl, -C₁-C₆- alkyl thienyl, -C₁-C₆- alkyl pyrrolyl, -C₁-C₆- alkyl pyridyl, benzyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, C₁-C₂ haloalkyl (preferably CF₃), or C₁-C₄ haloalkoxy (preferably OCF₃); and
 L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl-, -(C₁-C₄) alkyl-, -alkenyl-, or -phenyl-.

15 A preferred class of compounds of formula I-f are compounds of formula I-g, wherein,
 L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -O-(C₁-C₆)alkyl-, or - (C₁-C₆)alkyl-O-;
 the A ring is phenyl, furanyl, indolyl, thiazolyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₂ haloalkyl, C₁-C₂ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;
 Q is phenyl, -phenyl-O-phenyl, benzofuranyl, indolyl, 1,2,3,4-tetrahydroquinoliny, 1,2,3,4-tetrahydroisoquinoliny, dibenzofuranyl, or benzofuranyl-CH₂-phenyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkoxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, CF₃, OCF₃, NR₆R₇, phenyl, or phenyl-(C₁-C₆)alkyl-; wherein R₆ and R₇ are independently H, C₁-C₆ alkyl, benzyl, C₂-C₆ alkanoyl, phenyl(C₁-C₆)alkanoyl, C₁-C₆ alkoxy, or -SO₂-phenyl, wherein the cyclic

groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃; and

5 Z is phenyl, optionally substituted with 1, 2, or 3 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, CF₃, OCF₃, or NO₂.

A preferred class of compounds of formula I-e are
10 compounds of formula I-h, wherein,
L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄) alkyl-;
L₅ is -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄) alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-,
15 -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl, -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, or -N(R₉)SO₂-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups;
R₁ is H or C₁-C₆ alkyl;
R₂₁ is H; and
R₂₂ is H, phenyl(C₁-C₆)alkoxy, benzyl, halogen, (C₁-C₆)alkyl, OH,
20 C₁-C₆ alkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, CF₃ or OCF₃.

25

A preferred class of compounds of formula I-h are
compounds of formula I-i, wherein,
the A ring is phenyl, indolyl, or thiazolyl, each of which is
optionally substituted with 1, or 2 groups that are
30 independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, CF₃, OCF₃, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;

Q is phenyl, -phenyl-O-phenyl, 1,2,3,4-tetrahydroquinolinyl,
 1,2,3,4-tetrahydroisoquinolinyl, benzofuranyl,
 dibenzofuranyl, or benzofuranyl-CH₂-phenyl, wherein the
 aforementioned cyclic groups are optionally substituted
 with 1, 2, or 3 groups that are independently C₁-C₆
 alkoxy, carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, CF₃,
 OCF₃, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl.


Preferred compounds of formula I also include compounds
 wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted with 1 or
 2 R₁₁ groups.

Preferred compounds of formula I also include compounds
 wherein L₅ is -(C₂-C₆) alkenyl-, optionally substituted with 1
 or 2 R₁₁ groups.

Preferred compounds of formula I also include compounds
 wherein L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, wherein the
 alkyl portion of L₅ is optionally substituted with 1 or 2 R₁₁
 groups.

Preferred compounds of formula I also include compounds
 wherein L₅ is -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein the alkyl
 portion of L₅ is optionally substituted with 1 or 2 R₁₁ groups.

A preferred class of compounds of formula I-c are
 compounds of formula II,



(II)

wherein

n is 0, 1, 2, 3, or 4; and

each R₁₀ is independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl

5 R₁ is H, C₁-C₆ alkyl, or benzyl;

Q is H, phenyl, -phenyl-O-phenyl, -phenyl-(C₁-C₄)alkyl-phenyl, -phenyl-pyridyl, -phenyl-pyrimidyl, -phenyl-benzofuranyl, -phenyl-indolyl, -phenyl-piperidinyl, -phenyl-pyrrolidinyl, -phenyl-piperazinyl, indoliziny, benzofuranyl, indolyl, dibenzofuranyl, -benzothienyl-(C₁-C₄)alkyl-phenyl, -indolyl-(C₁-C₄)alkyl-phenyl, benzofuranyl-(C₁-C₄)alkyl-phenyl, piperidinyl, pyrrolidinyl, tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or imidazo[2,1-b]thiazol-3-one, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxy carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, or phenyl-(C₁-C₆)alkyl-;

20 L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -(C₁-C₄)alkyl-N(R₁₀)C(O)-, -C(O)N(R₁₀)-(C₁-C₄)alkyl-, -N(R₁₀)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-(C₁-C₆)alkyl-, or -(C₁-C₆)alkyl-O-;

R₉ and R₁₀ are independently is H, C₁-C₆ alkyl, -SO₂phenyl,

25 -CH₂-furanyl, -CH₂-tetrazolyl, benzyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, C₁-C₄ haloalkyl, or C₁-C₄ haloalkoxy.

30 Preferably, Q-L₃- is in a meta position on the phenylene ring.

Preferred compounds of formula II include compounds of formula II-a, wherein

L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, or -(C₁-C₄)alkyl-; and

- 5 R₂₀ and R₂₁ are independently selected from H, phenyl(C₁-C₆)alkoxy, phenyl(C₁-C₆)alkyl, halogen, (C₁-C₆)alkyl, OH, C₁-C₆ alkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are
- 10 independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, C₁-C₂ haloalkyl, or C₁-C₂ haloalkoxy.

Preferred compounds of formula II-a include compounds of formula II-b, wherein

- 15 R₁ is H, or C₁-C₆ alkyl,
- Q is H, phenyl, -phenyl-O-phenyl, indoliziny, benzofuranyl, indolyl, dibenzofuranyl, -benzothienyl-(C₁-C₄)alkyl-phenyl, -indolyl-(C₁-C₄)alkyl-phenyl, benzofuranyl-(C₁-C₄)alkyl-phenyl, piperidinyl, pyrrolidinyl,
- 20 tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or imidazo[2,1-b]thiazol-3-one, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, or
- 25 phenyl-(C₁-C₆)alkyl-.

Preferred compounds of formula II also include compounds wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted with 1 or


- 30 2 R₁₁ groups.

Preferred compounds of formula II also include compounds wherein L_5 is $-(C_2-C_6)$ alkenyl-, optionally substituted with 1 or 2 R_{11} groups.

5 Preferred compounds of formula II also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

10 Preferred compounds of formula II also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

Preferred class of compounds of formula II-b include
15 compounds of formula III,



wherein

20 n is 0, 1, 2, 3, or 4; and
each R_{10} is independently, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)$ alkyl, or $N(C_1-C_6)$ alkyl (C_1-C_6) alkyl.

25 Preferred compounds of formula III include compounds of formula III-a, wherein
 L_3 is a bond or $-(C_1-C_4)$ alkyl-; and
 L_5 is a bond, $-O-(C_1-C_6)$ alkyl-, $-(C_1-C_6)$ alkyl-O-, $-C(O)N(R_9)-(C_1-C_4)$ alkyl-,
30 $-N(R_9)C(O)-(C_1-C_4)$ alkyl-, $-N(R_9)-(C_1-C_6)$ alkyl-,
 $-(C_1-C_4)$ alkyl- $N(R_9)$, $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-,
 $-SO_2N(R_9)-$, $-SO_2N(R_9)-(C_1-C_4)$ alkyl-, $-N(R_9)SO_2-(C_1-C_4)$ alkyl-

, -N(R₉)SO₂-, -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl, -N(R₉)C(O)-, -C(O)-(C₁-C₄) alkyl-, -S-(C₁-C₄) alkyl-, or -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups.

5

Preferred compounds of formula III-a include compounds of formula III-b, wherein

L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl, -C(O)-(C₁-C₄) alkyl-, -S-(C₁-C₄) alkyl-, or -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups.

Preferred compounds of formula III-b include compounds of formula III-c, wherein

R₁ and R₂₁ are both H; and

R₂₂ is H, phenyl(C₁-C₆)alkoxy, benzyl, halogen, (C₁-C₆)alkyl, OH, C₁-C₆ alkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, C₁-C₂ haloalkyl, or C₁-C₂ haloalkoxy.

Preferred compounds of formula III-c include compounds of formula III-d, wherein

L₅ is -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl, or -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups.

30

Preferred compounds of formula III-c include compounds of formula III-e, wherein

L₅ is -(C₁-C₄)alkyl- or -C(O)-(C₁-C₄) alkyl-. In another aspect, L₅ is -CH₂-.

Preferred compounds of formula III-c include compounds of
 5 formula III-f, wherein
 L₅ is -S-(C₁-C₄)alkyl- or -(C₁-C₄)alkyl-S-(C₁-C₄) alkyl-. In another aspect, L₅ is -S-(C₁-C₂)alkyl-.

Preferred compounds of formula III-a include compounds of
 10 formula III-g, wherein
 L₅ is -C(O)N(R₉)-(C₁-C₄) alkyl-, -N(R₉)C(O)-(C₁-C₄) alkyl-, -N(R₉)-(C₁-C₆) alkyl-, -(C₁-C₄) alkyl-N(R₉), -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄) alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -N(R₉)SO₂-, -N(R₉)C(O)-, -(C₁-C₄)
 15 alkyl-, or -(C₂-C₆) alkenyl, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups,
 R₉ is H, C₁-C₆ alkyl, -SO₂phenyl, -C₁-C₆ alkyl-furanyl, -C₁-C₆ alkyl-tetrazolyl, -C₁-C₆- alkyl thienyl, -C₁-C₆-
 20 alkyl pyrrolyl, -C₁-C₆- alkyl pyridyl, benzyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-
 25 C₆)alkyl(C₁-C₆)alkyl, C₁-C₄ haloalkyl, or C₁-C₄ haloalkoxy.

Preferred compounds of formula III-g include compounds of
 formula III-h, wherein
 30 R₁ and R₂₁ are both H; and
 R₂₂ is H, phenyl(C₁-C₆)alkoxy, benzyl, halogen, (C₁-C₆)alkyl, OH, C₁-C₆ alkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl, wherein the phenyl group is optionally

substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, C₁-C₂ haloalkyl, or C₁-C₂ haloalkoxy.

5 Preferred compounds of formula III-h include compounds of formula III-i, wherein

R₉ is H, C₁-C₆ alkyl, -SO₂phenyl, -C₁-C₄ alkyl-furanyl, -C₁-C₄ alkyl-tetrazolyl, -C₁-C₄- alkyl thienyl, -C₁-C₄- alkyl pyrrolyl, -C₁-C₄- alkyl pyridyl, benzyl, wherein the aryl
10 and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

15 Preferred compounds of formula III-i include compounds of formula III-j, wherein

L₅ is -C(O)N(R₉)-(C₁-C₄) alkyl-, -N(R₉)C(O)-(C₁-C₄) alkyl-,
-N(R₉)C(O)-, -(C₁-C₄)-alkyl-S-(C₁-C₄) alkyl-, -(C₁-C₄)
alkyl-N(R₉)-(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl-, -(C₁-C₄)
20 alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups.

Preferred compounds of formula III-j include compounds of formula III-k, wherein

25 R₉ is H, C₁-C₆ alkyl, or benzyl, wherein phenyl group is optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

30 Preferred compounds of formula III-j include compounds of formula III-l, wherein

R₉ is H, -SO₂phenyl, -C₁-C₄ alkyl-furanyl, -C₁-C₄ alkyl-tetrazolyl, -C₁-C₄- alkyl thienyl, -C₁-C₄- alkyl pyrrolyl, -C₁-C₄- alkyl pyridyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

Preferred compounds of formula III-i include compounds of formula III-m, wherein

L₅ is -N(R₉)-(C₁-C₆) alkyl-, -(C₁-C₄) alkyl-N(R₉), or -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-.

Preferred compounds of formula III-m include compounds of formula III-n, wherein

R₉ is H, C₁-C₆ alkyl, or benzyl, wherein phenyl group is optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

Preferred compounds of formula III-m include compounds of formula III-o, wherein

R₉ is H, -SO₂phenyl, -C₁-C₄ alkyl-furanyl, -C₁-C₄ alkyl-tetrazolyl, -C₁-C₄- alkyl thienyl, -C₁-C₄- alkyl pyrrolyl, -C₁-C₄- alkyl pyridyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

Preferred compounds of formula III-i include compounds of formula III-p, wherein

L₅ is -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl-, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups.

5

Preferred compounds of formula III-p include compounds of formula III-q, wherein

R₉ is H, C₁-C₆ alkyl, or benzyl, wherein phenyl group is optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

10

Preferred compounds of formula III-p include compounds of formula III-r, wherein

15

R₉ is H, -SO₂phenyl, -C₁-C₄ alkyl-furanyl, -C₁-C₄ alkyl-tetrazolyl, -C₁-C₄-alkyl thienyl, -C₁-C₄-alkyl pyrrolyl, -C₁-C₄-alkyl pyridyl, wherein the aryl and heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, CF₃, or OCF₃.

20

Preferred compounds of formula III-a include compounds of formula III-s, wherein

25

R₁ and R₂₁ are both H; and

R₂₂ is H, phenyl(C₁-C₆)alkoxy, benzyl, halogen, (C₁-C₆)alkyl, OH, C₁-C₆ alkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, CF₃, or OCF₃.

30


Preferred compounds of formula III also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-, optionally substituted with 1 or 2 R_{11} groups.

5 Preferred compounds of formula III also include compounds wherein L_5 is $-(C_2-C_6)$ alkenyl-, optionally substituted with 1 or 2 R_{11} groups.

10 Preferred compounds of formula III also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

15 Preferred compounds of formula III also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

Preferred compound of formula III-s include compounds of formula IV,



(IV)

wherein

n is 0, 1, 2, 3, or 4; and

each R_{10} is independently, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy,

25 C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$.

Preferably, the dibenzofuranyl group is in a meta position on the phenylene ring.


Preferred compounds of formula IV also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-, optionally substituted with 1 or 2 R_{11} groups.

5 Preferred compounds of formula IV also include compounds wherein L_5 is $-(C_2-C_6)$ alkenyl-, optionally substituted with 1 or 2 R_{11} groups.

10 Preferred compounds of formula IV also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

15 Preferred compounds of formula IV also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

Preferred compound of formula III-s include compounds of formula V,



wherein

n is 0, 1, 2, 3, or 4; and

each R_{10} is independently, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy,

25 C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$.

Preferably, the benzyl substituted benzofuranyl methylene moiety is in a meta position on the phenylene ring.


Preferred compounds of formula V also include compounds
5 wherein L_5 is $-(C_1-C_4)$ alkyl-, optionally substituted with 1 or 2 R_{11} groups.

Preferred compounds of formula V also include compounds
10 wherein L_5 is $-(C_2-C_6)$ alkenyl-, optionally substituted with 1 or 2 R_{11} groups.

Preferred compounds of formula V also include compounds
15 wherein L_5 is $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

Preferred compounds of formula V also include compounds
20 wherein L_5 is $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

Preferred compound of formula III-s include compounds of
formula VI,



(VI)

wherein

n is 0, 1, 2, 3, or 4;

each R_{10} is independently, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy,

C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$,

30 or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and

R₅ is alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, or phenyl.

5 Preferably, the benzofuranyl methylene moiety is in a meta position on the phenylene ring.


Preferred compounds of formula VI also include compounds wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted with 1 or
10 2 R₁₁ groups.

Preferred compounds of formula VI also include compounds wherein L₅ is -(C₂-C₆) alkenyl-, optionally substituted with 1 or 2 R₁₁ groups.

15 Preferred compounds of formula VI also include compounds wherein L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, wherein the alkyl portion of L₅ is optionally substituted with 1 or 2 R₁₁ groups.

20 Preferred compounds of formula VI also include compounds wherein L₅ is -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein the alkyl portion of L₅ is optionally substituted with 1 or 2 R₁₁ groups.

25 Preferred compound of formula III-s include compounds of formula VII,



(VII)

wherein

n is 0, 1, 2, 3, or 4; and

each R₁₀ is independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy,
C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl,
5 or N(C₁-C₆)alkyl(C₁-C₆)alkyl.

Preferably, the indolyl group is in a meta position on the phenylene ring.


10 Preferred compounds of formula VII also include compounds wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted with 1 or 2 R₁₁ groups.

Preferred compounds of formula VII also include compounds
15 wherein L₅ is -(C₂-C₆) alkenyl-, optionally substituted with 1 or 2 R₁₁ groups.

Preferred compounds of formula VII also include compounds wherein L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, wherein the
20 alkyl portion of L₅ is optionally substituted with 1 or 2 R₁₁ groups.

Preferred compounds of formula VII also include compounds wherein L₅ is -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein the alkyl
25 portion of L₅ is optionally substituted with 1 or 2 R₁₁ groups.

Preferred compound of formula III-s include compounds of formula VIII,



(VIII)

wherein

n is 0, 1, 2, 3, or 4; and

each R₁₀ is independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy,

5 C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or
N(C₁-C₆)alkyl(C₁-C₆)alkyl.

Preferably, the diphenylether group is in a meta position
on the phenylene ring.

10

Preferred compounds of formula VIII also include
compounds wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted
with 1 or 2 R₁₁ groups.

15

Preferred compounds of formula VIII also include
compounds wherein L₅ is -(C₂-C₆) alkenyl-, optionally
substituted with 1 or 2 R₁₁ groups.

20


Preferred compounds of formula VIII also include
compounds wherein L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-,
wherein the alkyl portion of L₅ is optionally substituted with
1 or 2 R₁₁ groups.

25

Preferred compounds of formula VIII also include
compounds wherein L₅ is -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein
the alkyl portion of L₅ is optionally substituted with 1 or 2
R₁₁ groups.

30

A preferred class of compounds of formula A are compounds
of formula X,



Preferred compounds of formula X are compounds of formula

5 X-a, wherein,

Q is H, phenyl, -phenyl-O-phenyl, -phenyl-carbonyl-phenyl, -phenyl-(C₁-C₄)alkyl-phenyl, -phenyl-pyridyl, -phenyl-pyrimidyl, -phenyl-benzofuranyl, -phenyl-indolyl, -phenyl-piperidinyl, -phenyl-pyrrolidinyl, -phenyl-piperazinyl, indoliziny, benzofuranyl, indolyl, dibenzofuranyl, -benzothienyl-(C₁-C₄)alkyl-phenyl, -indolyl-(C₁-C₄)alkyl-phenyl, benzofuranyl-(C₁-C₄) alkyl-phenyl, piperidinyl, pyrrolidinyl, tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or imidazo[2,1-b]thiazol-3-one, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxy, carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, or phenyl-(C₁-C₆)alkyl-; wherein

20 R₆ and R₇ are independently H, C₁-C₆ alkyl, phenyl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, phenyl(C₂-C₆)alkanoyl, C₁-C₆ alkoxy, carbonyl, phenyl(C₁-C₆)alkoxy, carbonyl, heteroaryl, heterocycloalkyl, carbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, or phenyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, or NO₂.

5

A preferred class of compounds of formula X-a are compounds of formula X-b, wherein,

the A ring is phenyl, -phenyl-O-phenyl, pyrido[1,2-a]indolyl, furanyl, thienyl, benzofuranyl, dibenzofuranyl, indolyl, thiazolyl, thiazolidinyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H, phenyl(C₁-C₆)alkoxy, phenyl(C₁-C₆)alkyl, halogen, (C₁-C₆)alkyl, OH, alkoxy, CN, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-phenyl, NHC(O)-(C₁-C₄)alkyl-phenyl, N(C₁-C₄alkyl)C(O)-(C₁-C₄)alkyl-phenyl, N(C₁-C₄)alkyl-phenyl, -NHSO₂-phenyl, -N(C₁-C₄alkyl)SO₂phenyl, wherein the phenyl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy.

25

A preferred class of compounds of formula X-b are compounds of formula X-c, wherein,

R₁ is H, C₁-C₆ alkyl, benzyl, or allyl;

L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-, -(C₁-C₄)alkyl-

N(R₁₀)C(O)-, -C(O)N(R₁₀)-(C₁-C₄)alkyl-, -N(R₁₀)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-(C₁-C₆)alkyl-, or -(C₁-C₆)alkyl-O-;

L₃ is absent, a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-, -alkenyl-, or -phenyl-;

L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -C(O)N(R₉)-(C₁-C₄)alkyl-, -N(R₉)C(O)-(C₁-C₄)alkyl-, -N(R₉)-(C₁-C₆)alkyl-,
 5 , -(C₁-C₄)alkyl-N(R₉), -(C₁-C₄)alkyl-N(R₉)-(C₁-C₄)alkyl-,
 -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄)alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-,
 , -N(R₉)SO₂-, -(C₁-C₄)alkyl-, -(C₂-C₆)alkenyl-, -N(R₉)-(C₁-C₆)alkyl- wherein -(C₁-C₆)alkyl- is optionally
 substituted with phenyl, -N(R₉)C(O)-, -C(O)-(C₁-C₄)alkyl-,
 10 -S-(C₁-C₄)alkyl-, or -(C₁-C₄)alkyl-S-(C₁-C₄)alkyl-,
 wherein each alkyl and alkenyl is optionally substituted
 with one or more R₁₁ groups,

R₉ and R₁₀ are independently is H, C₁-C₆alkyl, -SO₂phenyl,
 -C₁-C₆alkyl-furanyl, -C₁-C₆alkyl-tetrazolyl, -C₁-C₆-
 15 alkyl thienyl, -C₁-C₆-alkyl pyrrolyl, -C₁-C₆-alkyl
 pyridyl, benzyl, wherein the aryl and heteroaryl
 groups are optionally substituted with 1, 2, 3, or 4
 groups that are independently C₁-C₄alkyl, C₁-C₄
 alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-
 20 C₆)alkyl(C₁-C₆)alkyl, C₁-C₄haloalkyl, or C₁-C₄
 haloalkoxy,

R₁₁ at each occurrence is independently N₁₂R₁₃, -
 N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂
 and R₁₃ are independently H or C₁-C₆alkyl.

25 A preferred class of compounds of formula X-c are
 compounds of formula X-d, wherein,

L₂ is a bond;

L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, or -(C₁-C₄)
 30 alkyl-;

the A ring is phenyl; and

Z is absent.

A preferred class of compounds of formulas X-c or X-d are compounds of formula X-e, wherein,

R₂₁ and R₂₃ are both H;

R₂₂ is OH or phenyl(C₁-C₆)alkoxy;

5 L₅ is -N(R₉)C(O)-(C₁-C₄) alkyl-, or -N(R₉)C(O)-, and

Q is phenyl, benzofuranyl, indolyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, or dibenzofuranyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups
10 that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, or phenyl.

A preferred class of compounds of formula X-e are compounds of formula I-f, wherein R₉ is H.

15 Preferred compounds of formula X-e are compounds of formula X-f, wherein Q is benzofuran, optionally substituted with C₁-C₆ alkyl.


Preferred compounds of formula X also include compounds
20 wherein L₅ is -(C₁-C₄) alkyl-, optionally substituted with 1 or 2 R₁₁ groups.

Preferred compounds of formula X also include compounds wherein L₅ is -(C₂-C₆) alkenyl-, optionally substituted with 1
25 or 2 R₁₁ groups.

Preferred compounds of formula X also include compounds wherein L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, wherein the alkyl portion of L₅ is optionally substituted with 1 or 2 R₁₁
30 groups.

Preferred compounds of formula X also include compounds wherein L_5 is $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein the alkyl portion of L_5 is optionally substituted with 1 or 2 R_{11} groups.

5 In another aspect, the invention provides compounds of Formula XI:



XI

and pharmaceutically acceptable salts thereof, wherein:

10 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1-C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1-C_3 alkoxy or C_1-C_2 acyl, or with one phenoxy;

15 R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1-C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1-C_3 alkoxy or C_1-C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1-C_6 alkyl;

20 R_{20} and R_{21} are independently hydrogen or halogen;

L_5 is $-(C_1-C_4)$ alkyl-N(R_9)- (C_1-C_4) alkyl-, $-(C_2-C_6)$ alkenyl-, or $-(C_1-C_4)$ alkyl-S- (C_1-C_4) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or two R_{11} groups,

25 R_9 is H, C_1-C_6 alkyl, C_1-C_6 alkoxycarbonyl;

R_{11} at each occurrence is independently $N_{12}R_{13}$, -

$N(R_{12})C(O)R_{13}$, $N(R_{12})CO_2R_{13}$, or $-C(O)NR_{12}R_{13}$, wherein R_{12} and R_{13} are independently H or C_1-C_6 alkyl,

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and n is 0, 1, or 2.

5

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl.

In particular aspects of Formula XI, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of
10 Formula XI, n is 1 and R_{10} is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another preferred aspect of Formula XI, n is 1 and R_{10} is fluoro, methyl, methoxy, or trifluoromethyl.

15 Preferred compounds of Formula XI include those where R_{20} is hydrogen and R_{21} is hydrogen. Also preferred are compounds wherein R_{20} is hydrogen and R_{21} is halogen. More preferably R_{21} is fluoro. In another aspect, both of R_{20} and R_{21} are halogen, preferably fluoro.

20 Other preferred compounds of Formula XI include those where R_{28} is dibenzofuranyl or adamantanyl and R_{29} is hydrogen, halogen, or *t*-butyl. Other preferred compounds of Formula XI include those where R_{29} is dibenzofuranyl or adamantanyl and R_{28} is hydrogen, halogen, or *t*-butyl.

25 Where one of R_{28} and R_{29} is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably dibenzofuran-4-yl. Each of these preferred R_{28} and R_{29} groups is optionally substituted with from 1-4, more preferably 1-3,
30 and most preferably 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_4 alkoxycarbonyl, C_1 - C_6 alkoxy, halogen, haloalkyl, haloalkoxy, and NR_6R_7 , where R_6 and R_7 are independently H, C_1 - C_6 alkyl, C_1 - C_6 alkanoyl, C_1 - C_6 alkoxycarbonyl, piperidinyl,

pyrrolidinylcarbonyl, $-C(O)NH_2$, $-C(O)NH(C_1-C_6)alkyl$, or $-C(O)N(C_1-C_6)alkyl(C_1-C_6)alkyl$.

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R_{28} is hydrogen, R_{29} is dibenzothiophen-4-yl.

5 In a particular aspect of Formual XI, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In a preferred aspect, R_{20} is hydrogen and R_{21} is halogen, more preferably fluoro. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro. In yet another
10 preferred aspect, both of R_{20} and R_{21} are hydrogen.


In another particular aspect of Formual XI, R_{29} is hydrogen; R_{28} is dibenzofuran-4-yl. Preferably, within this aspect both of R_{20} and R_{21} are halogen, preferably fluoro. More preferably, within this aspect, R_{20} is hydrogen and R_{21} is
15 fluoro.

Preferred compounds of formula XI include compounds wherein L_5 is $-(C_1-C_4)alkyl-N(R_9)-(C_1-C_4)alkyl-$, wherein R_9 is C_1-C_6 alkoxycarbonyl.

Preferred compounds of formula XI include compounds
20 wherein L_5 is $-(C_2-C_6)alkenyl-$, optionally substituted with one R_{11} group.

Preferred compounds of formula XI include compounds wherein L_5 is $-(C_1-C_4)alkyl-S-(C_1-C_4)alkyl-$.

25 In another aspect, the invention provides compounds of Formula XII:



XII

and pharmaceutically acceptable salts thereof, wherein

R₂₇ is C₁-C₆ alkoxy;

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)

dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,

5 or (vii) phenyl substituted with one or two C₁-C₃ alkoxy
or C₁-C₂ acyl, or with one phenoxy;

R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)

dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,

or (vii) phenyl substituted with one or two C₁-C₃ alkoxy

10 or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆
alkyl;

R₂₀ and R₂₁ are independently hydrogen or halogen;

R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

15 each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy,
C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl,
or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

n is 0, 1, or 2.

A preferred dibenzothiophenyl group is dibenzothiophen-4-
20 yl.

Preferred compounds of Formula XII include those where R₂₀
is hydrogen and R₂₁ is hydrogen. Also preferred are compounds
where R₂₀ is hydrogen and R₂₁ is halogen. More preferably R₂₁ is
fluoro. In another aspect, both of R₂₀ and R₂₁ are halogen,
25 preferably fluoro.

In particular aspects of Formula XII, n is 0 or 1. In a
preferred aspect, n is 0. In another preferred aspect of
Formula XII, n is 1 and R₁₀ is chloro, fluoro, methyl, ethyl,
methoxy, ethoxy, trifluoromethyl, or amino. In still another
30 preferred aspect of Formula XII, n is 1 and R₁₀ is fluoro,
methyl, methoxy, or trifluoromethyl.

Other preferred compounds of Formula XII include those
where R₂₈ is dibenzofuranyl or adamantanyl and R₂₉ is hydrogen,

halogen, or t-butyl. Other preferred compounds of Formula XII include those where R₂₉ is dibenzofuranyl or adamantanyl and R₂₈ is hydrogen, halogen, or t-butyl.

Where one of R₂₈ and R₂₉ is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably dibenzofuran-4-yl. Each of these preferred R₂₈ and R₂₉ groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, and NR₆R₇, where R₆ and R₇ are independently H, C₁-C₆ alkyl, C₁-C₆ alkanoyl, C₁-C₆ alkoxy, piperidinyl, pyrrolidinyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, or -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl.

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R₂₈ is hydrogen, R₂₉ is dibenzothiophen-4-yl, and each of R₃₀ and R₃₁ are hydrogen.

In a particular aspect of Formual XII, R₂₈ is hydrogen; R₂₉ is dibenzofuran-4-yl; and each of R₃₀ and R₃₁ is hydrogen. In a preferred aspect, R₂₀ is hydrogen and R₂₁ is halogen, more preferably fluoro. In another preferred aspect, both of R₂₀ and R₂₁ are halogen, preferably fluoro. In yet another preferred aspect, both of R₂₀ and R₂₁ are hydrogen.


In another particular aspect of Formual XII, R₂₉ is hydrogen; R₂₈ is dibenzofuran-4-yl; and each of R₃₀ and R₃₁ is hydrogen. Preferably, within this aspect both of R₂₀ and R₂₁ are halogen, preferably fluoro. More preferably, within this aspect, R₂₀ is hydrogen and R₂₁ is fluoro.

Still other preferred compounds of Formula XII include those where one of R₂₈ and R₂₉ is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XII include those where R₃₀ and R₃₁ are (i) both hydrogen, (ii) hydrogen and methyl, or

(iii) both methyl. More preferred compounds of Formula XII are those where R_{30} and R_{31} are both hydrogen.

In another aspect, the invention provides compounds of
5 Formula XIII:



XIII

and pharmaceutically acceptable salts thereof, wherein
 R_{27} is C_1 - C_6 alkoxy;

10 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)
15 dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1 - C_6 alkyl;

20 R_{20} and R_{21} are independently hydrogen or halogen;

R_{30} and R_{31} are independently hydrogen or C_1 - C_2 alkyl;

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and

25 n is 0, 1, or 2.

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R_{28} is hydrogen, R_{29} is dibenzothiophen-4-yl, and each of R_{30} and R_{31} are hydrogen.

In particular aspects of Formula XIII, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of Formula XIII, n is 1 and R_{10} is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another preferred aspect of Formula XIII, n is 1 and R_{10} is fluoro, methyl, methoxy, or trifluoromethyl.

Preferred compounds of Formula XIII include those where R_{20} is hydrogen and R_{21} is hydrogen. Also preferred are compounds wherein R_{20} is hydrogen and R_{21} is halogen. More preferably R_{21} is fluoro. In another aspect, both of R_{20} and R_{21} are halogen, preferably fluoro.

Other preferred compounds of Formula XIII include those where R_{28} is dibenzofuranyl or adamantanyl and R_{29} is hydrogen, halogen, or *t*-butyl. Other preferred compounds of Formula XIII include those where R_{29} is dibenzofuranyl or adamantanyl and R_{28} is hydrogen, halogen, or *t*-butyl.

Where one of R_{28} and R_{29} is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably dibenzofuran-4-yl. Each of these preferred R_{28} and R_{29} groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_4 alkoxy, carbonyl, C_1 - C_6 alkoxy, halogen, haloalkyl, haloalkoxy, and NR_6R_7 , where R_6 and R_7 are independently H, C_1 - C_6 alkyl, C_1 - C_6 alkanoyl, C_1 - C_6 alkoxy, carbonyl, piperidinyl, pyrrolidinyl, carbonyl, $-C(O)NH_2$, $-C(O)NH(C_1-C_6)alkyl$, or $-C(O)N(C_1-C_6)alkyl(C_1-C_6)alkyl$.

In a particular aspect of Formula XIII, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In a preferred aspect, R_{20} is hydrogen and R_{21} is halogen, more preferably fluoro. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro. In yet another preferred aspect, both of R_{20} and R_{21} are hydrogen.


In another particular aspect of Formual XIII, R_{29} is hydrogen; R_{28} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro.

5 In still another particular aspect of Formual XIII, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is methyl.

Still otherpreferred compounds of Formula XIII include those where one of R_{28} and R_{29} is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XIII include those where R_{30} and R_{31} are (i) both hydrogen, (ii) hydrogen and methyl, or (iii) both methyl. More preferred compounds of Formula XIII are those where R_{30} and R_{31} are both hydrogen.

15 In another aspect, the invention provides compounds of Formula XIV:



XIV

and pharmaceutically acceptable salts thereof, wherein

20 R_{27} is C_1 - C_6 alkoxy;

R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

25 R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;

R₂₀ and R₂₁ are independently hydrogen or halogen;

R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

5 each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and
n is 0, 1, or 2.

10 A preferred dibenzothiophenyl group is dibenzothiophen-4-yl.

Preferred compounds of Formula XIV include those where R₂₀ is hydrogen and R₂₁ is hydrogen. Also preferred are compounds where R₂₀ is hydrogen and R₂₁ is halogen. More preferably R₂₁ is fluoro. In another aspect, both of R₂₀ and R₂₁ are halogen,
15 preferably fluoro.

In particular aspects of Formula XIV, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of Formula XIV, n is 1 and R₁₀ is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another
20 preferred aspect of Formula XIV, n is 1 and R₁₀ is fluoro, methyl, methoxy, or trifluoromethyl.

Other preferred compounds of Formula XIV include those where R₂₈ is dibenzofuranyl or adamantanyl and R₂₉ is hydrogen, halogen, or t-butyl. Other preferred compounds of Formula XIV
25 include those where R₂₉ is dibenzofuranyl or adamantanyl and R₂₈ is hydrogen, halogen, or t-butyl.

Where one of R₂₈ and R₂₉ is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably
30 dibenzofuran-4-yl. Each of these preferred R₂₈ and R₂₉ groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C₁-C₆ alkyl, C₁-C₄ alkoxycarbonyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy,

and NR_6R_7 , where R_6 and R_7 are independently H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkanoyl, $\text{C}_1\text{-C}_6$ alkoxy carbonyl, piperidinyl, pyrrolidinyl carbonyl, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6)\text{alkyl}$, or $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6)\text{alkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$.

5 A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R_{28} is hydrogen, R_{29} is dibenzothiophen-4-yl, and each of R_{30} and R_{31} are hydrogen.


In a particular aspect of Formual XIV, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In a
10 preferred aspect, R_{20} is hydrogen and R_{21} is halogen, more preferably fluoro. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro. In yet another preferred aspect, both of R_{20} and R_{21} are hydrogen.

In another particular aspect of Formual XIV, R_{29} is
15 hydrogen; R_{28} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. Preferably, within this aspect both of R_{20} and R_{21} are halogen, preferably fluoro. More preferably, within this aspect, R_{20} is hydrogen and R_{21} is fluoro.

Still other preferred compounds of Formula XIV include
20 those where one of R_{28} and R_{29} is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XIV include those where R_{30} and R_{31} are (i) both hydrogen, (ii) hydrogen and methyl, or (iii) both methyl. More preferred compounds of Formula XIV are
25 those where R_{30} and R_{31} are both hydrogen.

In another aspect, the invention provides compounds of Formula XV:



XV

and pharmaceutically acceptable salts thereof, wherein
 R_{27} is C_1 - C_6 alkoxy;

- 5 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

- 10 R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1 - C_6 alkyl;

- 15 R_{20} and R_{21} are independently hydrogen or halogen;

R_{30} and R_{31} are independently hydrogen or C_1 - C_2 alkyl;

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and

- 20 n is 0, 1, or 2.

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R_{28} is hydrogen, R_{29} is dibenzothiophen-4-yl, and each of R_{30} and R_{31} are hydrogen.

- 25 In particular aspects of Formula XV, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of Formula XV, n is 1 and R_{10} is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another

preferred aspect of Formula XV, n is 1 and R₁₀ is fluoro, methyl, methoxy, or trifluoromethyl.

Preferred compounds of Formula XV include those where R₂₀ is hydrogen and R₂₁ is hydrogen. Also preferred are compounds wherein R₂₀ is hydrogen and R₂₁ is halogen. More preferably R₂₁ is fluoro. In another aspect, both of R₂₀ and R₂₁ are halogen, preferably fluoro.

Other preferred compounds of Formula XV include those where R₂₈ is dibenzofuranyl or adamantanyl and R₂₉ is hydrogen, halogen, or t-butyl. Other preferred compounds of Formula XV include those where R₂₉ is dibenzofuranyl or adamantanyl and R₂₈ is hydrogen, halogen, or t-butyl.

Where one of R₂₈ and R₂₉ is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably dibenzofuran-4-yl. Each of these preferred R₂₈ and R₂₉ groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C₁-C₆ alkyl, C₁-C₄ alkoxycarbonyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, and NR₆R₇, where R₆ and R₇ are independently H, C₁-C₆ alkyl, C₁-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, piperidinyl, pyrrolidinylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, or -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl.

In a particular aspect of Formual XV, R₂₈ is hydrogen; R₂₉ is dibenzofuran-4-yl; and each of R₃₀ and R₃₁ is hydrogen. In a preferred aspect, R₂₀ is hydrogen and R₂₁ is halogen, more preferably fluoro. In another preferred aspect, both of R₂₀ and R₂₁ are halogen, preferably fluoro. In yet another preferred aspect, both of R₂₀ and R₂₁ are hydrogen.


In another particular aspect of Formual XV, R₂₉ is hydrogen; R₂₈ is dibenzofuran-4-yl; and each of R₃₀ and R₃₁ is hydrogen. In another preferred aspect, both of R₂₀ and R₂₁ are halogen, preferably fluoro.

In still another particular aspect of Formual XV, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is methyl.

Still otherpreferred compounds of Formula XV include those
5 where one of R_{28} and R_{29} is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XV include those where
 R_{30} and R_{31} are (i) both hydrogen, (ii) hydrogen and methyl, or
(iii) both methyl. More preferred compounds of Formula XV are
10 those where R_{30} and R_{31} are both hydrogen.

In another aspect, the invention provides compounds of
15 Formula XVI:



XVI

and pharmaceutically acceptable salts thereof, wherein
p is 0 or 1;

20 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)
25 dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1 - C_6 alkyl;

R_{20} and R_{21} are independently hydrogen or halogen;

R_{30} and R_{31} are independently hydrogen or C_1 - C_2 alkyl;

5 each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and

n is 0, 1, or 2.

10 A preferred dibenzothiophenyl group is dibenzothiophen-4-yl.

Preferred compounds of Formula XVI include those where R_{20} is hydrogen and R_{21} is hydrogen. Also preferred are compounds where R_{20} is hydrogen and R_{21} is halogen. More preferably R_{21} is fluoro. In another aspect, both of R_{20} and R_{21} are halogen,
15 preferably fluoro.

In particular aspects of Formula XVI, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of Formula XVI, n is 1 and R_{10} is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another
20 preferred aspect of Formula XVI, n is 1 and R_{10} is fluoro, methyl, methoxy, or trifluoromethyl.

Other preferred compounds of Formula XVI include those where R_{28} is dibenzofuranyl or adamantanyl and R_{29} is hydrogen, halogen, or *t*-butyl. Other preferred compounds of Formula XVI
25 include those where R_{29} is dibenzofuranyl or adamantanyl and R_{28} is hydrogen, halogen, or *t*-butyl.

Where one of R_{28} and R_{29} is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably
30 dibenzofuran-4-yl. Each of these preferred R_{28} and R_{29} groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_4 alkoxycarbonyl, C_1 - C_6 alkoxy, halogen, haloalkyl, haloalkoxy,

and NR_6R_7 , where R_6 and R_7 are independently H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkanoyl, $\text{C}_1\text{-C}_6$ alkoxy carbonyl, piperidinyl, pyrrolidinyl carbonyl, $-\text{C}(\text{O})\text{NH}_2$, $-\text{C}(\text{O})\text{NH}(\text{C}_1\text{-C}_6)\text{alkyl}$, or $-\text{C}(\text{O})\text{N}(\text{C}_1\text{-C}_6)\text{alkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$.

5 A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R_{28} is hydrogen, R_{29} is dibenzothiophen-4-yl, and each of R_{30} and R_{31} are hydrogen.


In a particular aspect of Formual XVI, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In a
10 preferred aspect, R_{20} is hydrogen and R_{21} is halogen, more preferably fluoro. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro. In yet another preferred aspect, both of R_{20} and R_{21} are hydrogen.

In another particular aspect of Formual XVI, R_{29} is
15 hydrogen; R_{28} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. Preferably, within this aspect both of R_{20} and R_{21} are halogen, preferably fluoro. More preferably, within this aspect, R_{20} is hydrogen and R_{21} is fluoro.

Still other preferred compounds of Formula XVI include
20 those where one of R_{28} and R_{29} is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XVI include those where R_{30} and R_{31} are (i) both hydrogen, (ii) hydrogen and methyl, or (iii) both methyl. More preferred compounds of Formula XIV are
25 those where R_{30} and R_{31} are both hydrogen.

In another aspect, the invention provides compounds of Formula XVII:



XVII

and pharmaceutically acceptable salts thereof, wherein
p is 0 or 1;

- 5 R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

- 10 R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;

- 15 R₂₀ and R₂₁ are independently hydrogen or halogen;

R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

- 20 n is 0, 1, or 2.

A preferred dibenzothiophenyl group is dibenzothiophen-4-yl. Preferably, R₂₈ is hydrogen, R₂₉ is dibenzothiophen-4-yl, and each of R₃₀ and R₃₁ are hydrogen.

- 25 In particular aspects of Formula XVII, n is 0 or 1. In a preferred aspect, n is 0. In another preferred aspect of Formula XVII, n is 1 and R₁₀ is chloro, fluoro, methyl, ethyl, methoxy, ethoxy, trifluoromethyl, or amino. In still another

preferred aspect of Formula XVII, n is 1 and R_{10} is fluoro, methyl, methoxy, or trifluoromethyl.

Preferred compounds of Formula XVII include those where R_{20} is hydrogen and R_{21} is hydrogen. Also preferred are
5 compounds wherein R_{20} is hydrogen and R_{21} is halogen. More preferably R_{21} is fluoro. In another aspect, both of R_{20} and R_{21} are halogen, preferably fluoro.

Other preferred compounds of Formula XVII include those where R_{28} is dibenzofuranyl or adamantanyl and R_{29} is hydrogen,
10 halogen, or *t*-butyl. Other preferred compounds of Formula XVII include those where R_{29} is dibenzofuranyl or adamantanyl and R_{28} is hydrogen, halogen, or *t*-butyl.

Where one of R_{28} and R_{29} is hydrogen, the other is preferably adamantanyl or dibenzofuranyl, more preferably
15 dibenzofuran-3-yl or dibenzofuran-4-yl, most preferably dibenzofuran-4-yl. Each of these preferred R_{28} and R_{29} groups is optionally substituted with from 1-4, more preferably 1-3, and most preferably 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_4 alkoxy, carbonyl, C_1 - C_6 alkoxy, halogen, haloalkyl, haloalkoxy,
20 and NR_6R_7 , where R_6 and R_7 are independently H, C_1 - C_6 alkyl, C_1 - C_6 alkanoyl, C_1 - C_6 alkoxy, carbonyl, piperidinyl, pyrrolidinyl, carbonyl, $-C(O)NH_2$, $-C(O)NH(C_1-C_6)alkyl$, or $-C(O)N(C_1-C_6)alkyl(C_1-C_6)alkyl$.

In a particular aspect of Formula XVII, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen.
25 In a preferred aspect, R_{20} is hydrogen and R_{21} is halogen, more preferably fluoro. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro. In yet another preferred aspect, both of R_{20} and R_{21} are hydrogen.


30 In another particular aspect of Formula XVII, R_{29} is hydrogen; R_{28} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is hydrogen. In another preferred aspect, both of R_{20} and R_{21} are halogen, preferably fluoro.

In still another particular aspect of Formual XVII, R_{28} is hydrogen; R_{29} is dibenzofuran-4-yl; and each of R_{30} and R_{31} is methyl.

Still otherpreferred compounds of Formula XVII include
 5 those where one of R_{28} and R_{29} is 3,4-dimethoxyphenyl, 3-methoxyphenyl, 3-formylphenyl, or 3-acetylphenyl.

Preferred compounds of Formula XVII include those
 where R_{30} and R_{31} are (i) both hydrogen, (ii) hydrogen and
 methyl, or (iii) both methyl. More preferred compounds of
 10 Formula XVII are those where R_{30} and R_{31} are both hydrogen.


In another aspect, the invention provides a method for
 preparing a compound of formula (I)



15 or a pharmaceutically acceptable salt thereof, wherein
 L_2 is a bond; and A, Z, L_3 , L_5 , Q, R_1 , R_{20} , R_{21} , R_{22} , and R_{23} are
 as defined in claim 1;

comprising:


treating a compound of formula



20

wherein X is Cl, Br, I, or OSO_2CF_3 ,

with a metal catalyst, a base, and a compound of formula




wherein R_A is H or $(\text{C}_1\text{-C}_6)\text{alkyl}$, and


L is alkylene,

25

to provide a compound of formula



In another aspect, the invention provides a method for preparing a compound of formula (I)




(I),

or a pharmaceutically acceptable salt thereof, wherein L_2 is a bond; and A, Z, L_3 , L_5 , Q, R_1 , R_{20} , R_{21} , R_{22} , and R_{23} are as defined in claim 1;


comprising:

treating a compound of formula



wherein X is Cl, Br, I, or OSO_2CF_3 ,


with a metal catalyst, a base, and a compound of formula




wherein R_A is H or (C_1-C_6) alkyl, and

L is alkylene,

to provide a compound of formula



In another aspect, the invention provides a method for preparing a compound of formula (I)




(I),

- 5 or a pharmaceutically acceptable salt thereof, wherein L_5 is $-O-(C_1-C_6)\text{alkyl}-$, $-(C_1-C_6)\text{alkyl}-O-$, $-N(R_9)-(C_1-C_6)\text{alkyl}-$, $-N(R_9)-(C_1-C_6)\text{alkyl}-$ wherein $-(C_1-C_6)\text{alkyl}-$ is optionally substituted with phenyl, $-(C_1-C_4)\text{alkyl}-N(R_9)-(C_1-C_4)\text{alkyl}-$, $-S-(C_1-C_4)\text{alkyl}-$, $-(C_1-C_6)\text{alkyl}-S-$, or $-(C_1-C_4)\text{alkyl}-S-$
 10 $(C_1-C_4)\text{alkyl}-$, wherein each alkyl is optionally substituted with one or more R_{11} groups,
 R_9 is H, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-SO_2\text{-aryl}$, heteroarylalkyl, arylalkyl, wherein the aryl or heteroaryl groups are optionally substituted with 1,
 15 2, 3, or 4 groups that are independently C_1-C_4 alkyl, C_1-C_4 alkoxy, halogen, OH, NO_2 , NH_2 , $NH(C_1-C_6)\text{alkyl}$, $N(C_1-C_6)\text{alkyl}(C_1-C_6)\text{alkyl}$, haloalkyl, or haloalkoxy;
 R_{11} at each occurrence is independently $N_{12}R_{13}$, $-N(R_{12})C(O)R_{13}$, $N(R_{12})CO_2R_{13}$, or $-C(O)NR_{12}R_{13}$, wherein R_{12} and R_{13} are
 20 independently H or C_1-C_6 alkyl; and
 A , Z , L_2 , L_3 , Q , R_1 , R_{20} , R_{21} , R_{22} , and R_{23} are as defined in claim 1;

comprising:


treating a compound of formula



wherein R is $(CH_2)_nOH$, $(CH_2)_nSH$, or $(CH_2)_nNH_2$,

n is 0, 1, 2, 3, or 4,

with a base and a compound of formula




wherein X is Cl, Br, I, or $\text{OSO}_2\text{R}'$,

5 R' is methyl, para-methylphenyl, or CF_3 , and

m is 0, 1, 2, 3, or 4,

to provide a compound of formula (I).

In another aspect, the invention provides a compound of formula (XV)



(XV)

wherein X is Cl, Br, I, or OSO_2CF_3 ;

R₁ is H, C₁-C₆ alkyl, phenyl(C₁-C₆)alkyl, or C₂-C₆ alkenyl;

15 L₃ is absent, a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-, -alkenyl-, or -phenyl-;

L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -C(O)N(R₉)-(C₁-C₄)alkyl-, -N(R₉)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₉)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-N(R₉)C(O)-(C₁-C₄)alkyl-, -N(R₉)-(C₁-C₆)alkyl-, -N(R₉)-(C₁-C₆)alkyl- wherein
20 -(C₁-C₆)alkyl- is optionally substituted with phenyl, -(C₁-C₄)alkyl-N(R₉)-(C₁-C₄)alkyl-, -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄)alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -N(R₉)SO₂-, -(C₁-C₄)alkyl-, -(C₂-C₆)alkenyl-, -N(R₉)C(O)-, -C(O)-(C₁-C₄)alkyl-, -S-(C₁-C₄)alkyl-, -(C₁-C₆)alkyl-S-, or -(C₁-C₄)alkyl-S-(C₁-C₄)alkyl-, wherein each alkyl and alkenyl is
25 optionally substituted with one or more R₁₁ groups,
R₉ and R₁₀ are independently H, C₁-C₆ alkyl, C₁-C₆

alkoxycarbonyl, -SO₂-aryl, heteroarylalkyl,


arylalkyl, wherein the aryl or heteroaryl groups are

optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl, or haloalkoxy;

5 R₁₁ at each occurrence is independently N₁₂R₁₃, -N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂ and R₁₃ are independently H or C₁-C₆ alkyl; and

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H, arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO₂,
 10 NH₂, CN, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-aryl, NHC(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄)alkyl-aryl, -NHSO₂-aryl, or -N(C₁-C₄alkyl)SO₂aryl, wherein the aryl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are
 15 independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, haloalkyl, haloalkoxy.

In another aspect, the invention provides a compound of formula (XVI) or formula (XVII)



20 wherein R_A is H or (C₁-C₆)alkyl; and

L is alkylene;

L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-, -alkenyl-, or -phenyl-;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which
 25 is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;

Q is H, aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-
 30 aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl, -heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl,


wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxy carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxy carbonyl, aryl C₁-C₆ alkoxy carbonyl, heteroaryl carbonyl, heteroaryl, heterocycloalkyl carbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂, or

Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, or -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

In another aspect, the invention provides a compound of formula (XVIII)



(XVIII),

wherein X is Cl, Br, I, or OSO₂CF₃;

L₃ is a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄)alkyl-, -alkenyl-, or -phenyl-;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄

haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;


Q is H, aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl, -heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl, heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂, or

Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, or -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

In another aspect, the invention provides a compound of formula (XIX) or formula (XX)



wherein R_A is H or (C₁-C₆)alkyl; and

L is alkylene;

R₁ is H, C₁-C₆ alkyl, phenyl(C₁-C₆)alkyl, or C₂-C₆ alkenyl;


L₅ is a bond, -O-(C₁-C₆)alkyl-, -(C₁-C₆)alkyl-O-, -C(O)N(R₉)-(C₁-C₄) alkyl-, -N(R₉)C(O)-(C₁-C₄) alkyl-, -(C₁-C₄) alkyl-C(O)N(R₉)-(C₁-C₄) alkyl-, -(C₁-C₄) alkyl-N(R₉)C(O)-(C₁-C₄) alkyl-, -N(R₉)-(C₁-C₆) alkyl-, -N(R₉)-(C₁-C₆) alkyl- wherein -(C₁-C₆) alkyl- is optionally substituted with phenyl, -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -SO₂N(R₉)-, -SO₂N(R₉)-(C₁-C₄) alkyl-, -N(R₉)SO₂-(C₁-C₄)alkyl-, -N(R₉)SO₂-, -(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl-, -N(R₉)C(O)-, -C(O)-(C₁-C₄) alkyl-, -S-(C₁-C₄) alkyl-, -(C₁-C₆)alkyl-S-, or -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R₁₁ groups, R₉ and R₁₀ are independently H, C₁-C₆ alkyl, C₁-C₆

alkoxycarbonyl, -SO₂-aryl, heteroarylalkyl, arylalkyl, wherein the aryl or heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C₁-C₄ alkyl, C₁-C₄ alkoxy, halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl, or haloalkoxy;

R₁₁ at each occurrence is independently N₁₂R₁₃, -N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂ and R₁₃ are independently H or C₁-C₆ alkyl; and

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H, arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO₂, NH₂, CN, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-aryl, NHC(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄)alkyl-aryl, -NHSO₂-aryl, or -N(C₁-C₄alkyl)SO₂aryl, wherein the aryl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, haloalkyl, or haloalkoxy.

In another aspect, the invention provides a compound of formula (XXI)



(XXI),

- 5 wherein R is $(\text{CH}_2)_n\text{OH}$, $(\text{CH}_2)_n\text{SH}$, $(\text{CH}_2)_n\text{NH}_2$, $(\text{CH}_2)_n\text{CHO}$, or $(\text{CH}_2)_n\text{-X}$;
 X is Cl, Br, I, or $\text{OSO}_2\text{R}'$;
 R' is methyl, para-methylphenyl, or CF_3 ;
 n is 0, 1, 2, 3, or 4;
 L₂ is a bond or $-\text{C}(\text{O})\text{NR}_{10}-$, $-\text{N}(\text{R}_{10})\text{C}(\text{O})-$,
 10 $-(\text{C}_1-\text{C}_4)\text{alkyl}-\text{N}(\text{R}_{10})\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{N}(\text{R}_{10})-(\text{C}_1-\text{C}_4)\text{alkyl}-$,
 $-\text{N}(\text{R}_{10})\text{C}(\text{O})-(\text{C}_1-\text{C}_4)\text{alkyl}-$, $-(\text{C}_1-\text{C}_4)\text{alkyl}-\text{C}(\text{O})\text{N}(\text{R}_{10})-$, $-\text{O}-$
 $(\text{C}_1-\text{C}_6)\text{alkyl}-$, $-\text{CO}-$, $-\text{SO}_2-$, or $-(\text{C}_1-\text{C}_6)\text{alkyl}-\text{O}-$;
 L₃ is absent, a bond, $-(\text{C}_1-\text{C}_4)\text{alkyl}-\text{O}-$, $-\text{O}-(\text{C}_1-\text{C}_4)\text{alkyl}$,
 $-(\text{C}_1-\text{C}_4)\text{alkyl}-$, $-\text{alkenyl}-$, or $-\text{phenyl}-$;
 15 the A ring is aryl, heteroaryl, heterocycloalkyl, each of
 which is optionally substituted with 1, 2, or 3 groups
 that are independently, halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy,
 C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, NO_2 , NH_2 , $\text{NH}(\text{C}_1-\text{C}_6)\text{alkyl}$,
 or $\text{N}(\text{C}_1-\text{C}_6)\text{alkyl}(\text{C}_1-\text{C}_6)\text{alkyl}$;
 20 R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H,
 arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO_2 ,
 NH_2 , CN, $\text{NH}(\text{C}_1-\text{C}_6)\text{alkyl}$, $\text{N}(\text{C}_1-\text{C}_6)\text{alkyl}(\text{C}_1-\text{C}_6)\text{alkyl}$, NH-aryl ,
 $\text{NHC}(\text{O})-(\text{C}_1-\text{C}_4)\text{alkyl-aryl}$, $\text{N}(\text{C}_1-\text{C}_4\text{alkyl})\text{C}(\text{O})-(\text{C}_1-\text{C}_4)$
 alkyl-aryl , $\text{N}(\text{C}_1-\text{C}_4)\text{alkyl-aryl}$, $-\text{NHSO}_2\text{-aryl}$, or $-\text{N}(\text{C}_1-$
 25 $\text{C}_4\text{alkyl})\text{SO}_2\text{aryl}$, wherein the aryl group is optionally
 substituted with 1, 2, 3, 4, or 5 groups that are
 independently C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, OH, NO_2 ,
 haloalkyl, or haloalkoxy;
 Q is H, aryl, $-\text{aryl-carbonyl-aryl}$, $-\text{aryl-O-aryl}$, $-\text{aryl-alkyl-}$
 30 aryl , $-\text{aryl-heteroaryl}$, $-\text{aryl-heterocycloalkyl}$,

-heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl, heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂; or

Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, or -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

In another aspect, the invention provides a compound of formula (XXI)

wherein R is (CH₂)_nOH, (CH₂)_nSH, (CH₂)_nNH₂, (CH₂)_nCHO, or (CH₂)_n-X;

X is Cl, Br, I, or OSO₂R';

R' is methyl, para-methylphenyl, or CF₃;

n is 0, 1, 2, 3, or 4;

L₂ is a bond;

L₃ is a bond;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy,

C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;

R₂₀, R₂₁, R₂₂, and R₂₃ are H;

Q is aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl, -heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl, wherein the
5
aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxy, carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen,
10
haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxy, carbonyl, aryl C₁-C₆ alkoxy, carbonyl, heteroaryl, carbonyl, heteroaryl, heterocycloalkyl, carbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the
15
cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and
20

Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl(phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂; or
25

Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, or -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

In another aspect, the invention provides a compound of formula (XXI)

30 wherein R is (CH₂)_nOH, (CH₂)_nSH, (CH₂)_nNH₂, (CH₂)_nCHO, or (CH₂)_n-X;

X is Cl, Br, I, or OSO₂R';

R' is methyl, para-methylphenyl, or CF₃;

n is 0, 1, 2, 3, or 4;

L₂ is a bond;

L₃ is a bond,;

the A ring is phenyl;

R₂₀, R₂₁, R₂₂, and R₂₃ are H;

- 5 Q is -heteroaryl optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein
- R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-
- 10 C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl, heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the
- 15 cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and
- Z is H.

- In another aspect, the invention provides a compound of
- 20 formula (XXI)
- wherein R is (CH₂)_nOH, (CH₂)_nSH, (CH₂)_nNH₂, (CH₂)_nCHO, or (CH₂)_n-X;
- X is Cl, Br, I, or OSO₂R';
- R' is methyl, para-methylphenyl, or CF₃;
- n is 0, 1, 2, 3, or 4;

- 25 L₂ is a bond;
- L₃ is a bond,;
- the A ring is phenyl;
- R₂₀, R₂₁, R₂₂, and R₂₃ are H;
- Q is dibenzofuranyl, benzofuranyl, or indolyl, wherein each is
- 30 optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H or C₁-C₆ alkyl; and
Z is H.

In another aspect, the invention provides a method of
5 treating diabetes comprising administering a pharmaceutically
acceptable amount of a compound of formula A to a patient in
need of such treatment.

In yet another aspect, the invention provides a
10 pharmaceutical composition comprising a compound of formula A
and at least one pharmaceutically acceptable solvent, carrier,
adjuvant or excipient.

In another aspect, the invention provides a method of
treating diabetes, comprising administering to a patient in
15 need of such treatment a pharmaceutically acceptable amount of
a compounds of formula A.

In another aspect, the invention encompasses a method of
treating diabetes comprising administering to a patient in need
thereof, a pharmaceutically acceptable amount of a compound or
20 salt of formula A or a pharmaceutical composition comprising a
compound or salt of formula A.

In another aspect, the invention encompasses a method of
inhibiting TPT-1B comprising administering to a patient in need
thereof, a pharmaceutically acceptable amount of a compound or
25 salt of formula A or a pharmaceutical composition comprising a
compound or salt of formula A.

In another aspect, the invention encompasses a method of
treating cancer or neurodegenerative diseases comprising
administering to a patient in need thereof, a pharmaceutically
30 acceptable amount of a compound or salt of formula A or a
pharmaceutical composition comprising a compound or salt of
formula A.

In another aspect, the invention provides a method of treating syndrome X (consisting of such abnormalities as obesity, dyslipidemia, hypercoagulation, hypertension, insulin resistance and leading to heart disease and diabetes), obesity, 5 diabetes, immunological disease, bleeding disorders and/or cancer comprising administering a pharmaceutically acceptable amount of a compound of formula A to a patient in need of such treatment.

As noted above, the compounds of the invention bind to and 10 preferably, inhibit PTP-1B. As a result that are useful in the treatment of various diseases, including controlling or treating Type 2 diabetes, improving glucose tolerance, and in improving insulin sensitivity in patients in need thereof. The compounds are also useful in treating or controlling other PTP- 15 1B mediated diseases, such as the treatment of cancer, neurodegenerative diseases and the like.

In another aspect, the invention provides the use of a compound or salt of formula I for the manufacture of a medicament for treating cancer, neurodegenerative diseases 20 diabetes, syndrome X, immunological disease, bleeding disorders, or cardiovascular diseases in a patient in need of such treatment.

In another aspect, the invention provides the use of a compound or a salt of formula I for the manufacture of a 25 medicament for inhibiting PTP-1B in a patient in need thereof.

In another aspect, the invention provides the use of a pharmaceutical composition for the manufacture of a medicament comprising a compound of formula I and at least one pharmaceutically acceptable solvent, carrier, adjuvant or 30 excipient.

The term "alkoxy" represents an alkyl group of indicated number of carbon atoms attached to the parent molecular moiety

through an oxygen bridge. Examples of alkoxy groups include, for example, methoxy, ethoxy, propoxy and isopropoxy.

As used herein, the term "alkyl" includes those alkyl groups of a designed number of carbon atoms. Alkyl groups may be straight, or branched. Examples of "alkyl" include methyl, ethyl, propyl, isopropyl, butyl, iso-, sec- and tert-butyl, pentyl, hexyl, heptyl, 3-ethylbutyl, and the like.

The term "alkylene" means a divalent group derived from a straight or branched chain hydrocarbon of from 2 to 10 carbon atoms. Representative examples of alkylene include, but are not limited to, $-\text{CH}_2\text{CH}_2-$, $-\text{C}(\text{CH}_3)_2\text{C}(\text{CH}_3)_2-$, $-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2-$.

The term "aryl" refers to an aromatic hydrocarbon ring system containing at least one aromatic ring. The aromatic ring may optionally be fused or otherwise attached to other aromatic hydrocarbon rings or non-aromatic hydrocarbon rings. Examples of aryl groups include, for example, phenyl, naphthyl, 1,2,3,4-tetrahydronaphthalene and biphenyl. Preferred examples of aryl groups include phenyl, naphthyl, and anthracenyl. More preferred aryl groups are phenyl and naphthyl. Most preferred is phenyl.

The term "cycloalkyl" refers to a C_3 - C_8 cyclic hydrocarbon. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

The terms "halogen" or "halo" indicate fluorine, chlorine, bromine, and iodine.

The term "heterocycloalkyl" refers to a ring or ring system containing at least one heteroatom selected from nitrogen, oxygen, and sulfur, wherein said heteroatom is in a non-aromatic ring. The heterocycloalkyl ring is optionally fused to or otherwise attached to other heterocycloalkyl rings and/or non-aromatic hydrocarbon rings and/or phenyl rings.

Preferred heterocycloalkyl groups have from 3 to 7 members. Examples of heterocycloalkyl groups include, for example, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, piperazinyl, morpholinyl, piperidinyl, tetrahydrofuranyl, pyrrolidinyl, pyridinonyl, and pyrazolidinyl. Preferred heterocycloalkyl groups include piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, pyridinonyl, dihydropyrrolidinyl, and pyrrolidinonyl.

The term "heteroaryl" refers to an aromatic ring system containing at least one heteroatom selected from nitrogen, oxygen, and sulfur. The heteroaryl ring may be fused or otherwise attached to one or more heteroaryl rings, aromatic or non-aromatic hydrocarbon rings or heterocycloalkyl rings. Examples of heteroaryl groups include, for example, pyridine, furan, thienyl, 5,6,7,8-tetrahydroisoquinoline and pyrimidine. Preferred examples of heteroaryl groups include thienyl, benzothienyl, pyridyl, quinolyl, pyrazolyl, pyrimidyl, imidazolyl, benzimidazolyl, furanyl, benzofuranyl, dibenzofuranyl, thiazolyl, benzothiazolyl, isoxazolyl, oxadiazolyl, isothiazolyl, benzisothiazolyl, triazolyl, pyrrolyl, indolyl, pyrazolyl, and benzopyrazolyl.

The compounds of this invention may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. These compounds can be, for example, racemates, chiral non-racemic or diastereomers. In these situations, the single enantiomers, i.e., optically active forms, can be obtained by asymmetric synthesis or by resolution of the racemates. Resolution of the racemates can be accomplished, for example, by conventional methods such as crystallization in the presence of a resolving agent; chromatography, using, for example a chiral HPLC column; or derivatizing the racemic mixture with a resolving reagent to generate diastereomers, separating the diastereomers via

chromatography, and removing the resolving agent to generate the original compound in enantiomerically enriched form. Any of the above procedures can be repeated to increase the enantiomeric purity of a compound.

5 When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless otherwise specified, it is intended that the compounds include the cis, trans, Z- and E- configurations. Likewise, all tautomeric forms are also intended to be included.

10 The compounds of general Formula A may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes
15 percutaneous, subcutaneous, intravascular (e.g., intravenous), intramuscular, or intrathecal injection or infusion techniques and the like. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula A and a pharmaceutically acceptable carrier. One or more compounds of
20 general Formula A may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants, and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula A may be in a form suitable for
25 oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of
30 pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preservative agents in order to provide pharmaceutically

elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques. In some cases such coatings may be prepared by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules, wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Formulations for oral use may also be presented as lozenges.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropyl-methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain

aliphatic alcohols, for example heptadecaethylenoxycetanol, or
condensation products of ethylene oxide with partial esters
derived from fatty acids and a hexitol such as polyoxyethylene
sorbitol monooleate, or condensation products of ethylene oxide
5 with partial esters derived from fatty acids and hexitol
anhydrides, for example polyethylene sorbitan monooleate. The
aqueous suspensions may also contain one or more preservatives,
for example ethyl, or n-propyl p-hydroxybenzoate, one or more
coloring agents, one or more flavoring agents, and one or more
10 sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the
active ingredients in a vegetable oil, for example arachis oil,
olive oil, sesame oil or coconut oil, or in a mineral oil such
as liquid paraffin. The oily suspensions may contain a
15 thickening agent, for example beeswax, hard paraffin or cetyl
alcohol. Sweetening agents and flavoring agents may be added
to provide palatable oral preparations. These compositions may
be preserved by the addition of an anti-oxidant such as
ascorbic acid.

20 Dispersible powders and granules suitable for preparation
of an aqueous suspension by the addition of water provide the
active ingredient in admixture with a dispersing or wetting
agent, suspending agent and one or more preservatives.
Suitable dispersing or wetting agents or suspending agents are
25 exemplified by those already mentioned above. Additional
excipients, for example sweetening, flavoring and coloring
agents, may also be present.

Pharmaceutical compositions of the invention may also be
in the form of oil-in-water emulsions. The oily phase may be a
30 vegetable oil or a mineral oil or mixtures of these. Suitable
emulsifying agents may be naturally-occurring gums, for example
gum acacia or gum tragacanth, naturally-occurring phosphatides,
for example soy bean, lecithin, and esters or partial esters

derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening
5 and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

10 The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable
15 preparation may also be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.
20 In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

25 The compounds of general Formula A may also be administered in the form of suppositories, e.g., for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal
30 temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Compounds of general Formula A may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

For disorders of the eye or other external tissues, e.g., mouth and skin, the formulations are preferably applied as a topical gel, spray, ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active ingredients may be employed with either paraffinic or a water-miscible ointment base.

Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to

the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane. The transdermal patch may include the compound in a suitable solvent system with an adhesive system, such as an acrylic emulsion, and a polyester patch. The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others. The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required. Alternatively, high

melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

Formulations suitable for topical administration to the eye also include eye drops wherein the active ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10% and particularly about 1.5% w/w. For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient. The daily dose can be administered in one to four doses per day. In the case of skin conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

For administration to non-human animals, the composition may also be added to the animal feed or drinking water. It may be convenient to formulate the animal feed and drinking water compositions so that the animal takes in a therapeutically appropriate quantity of the composition along with its diet. It may also be convenient to present the composition as a premix for addition to the feed or drinking water. Preferred non-human animals include domesticated animals.

As noted above, the invention also provides methods and compositions for combination therapy of Type I and Type II diabetes. In one such aspect, the invention provides methods of using compounds of formula A in combination with one or more angiotensin converting enzyme (ACE) inhibitors for improving

the cardiovascular risk profile in patients experiencing or subject to Syndrome X or type II diabetes (non-insulin-dependent diabetes mellitus), preferably in human type II diabetics. These methods may also be characterized as the
5 reduction of risk factors for heart disease, stroke or heart attack in a type II diabetic.

These methods include the reduction of hyperlipidemia in a patients experiencing or subject to Syndrome X or type II diabetes. These methods include methods lowering low density
10 lipoprotein (LDL) blood levels and to increase high density lipoprotein (HDL) blood levels. The methods herein may further be characterized as useful for inhibiting, preventing or reducing atherosclerosis in a type II diabetics, or for reducing the risk factors thereof.

15 These methods also include the lowering of free fatty acid blood levels and triglyceride levels in type II diabetics.

Among the ACE inhibitors which may be utilized with the invention described herein are quinapril, ramipril, verapamil, captopril, diltiazem, clonidine, hydrochlorthiazide,
20 benazepril, prazosin, fosinopril, lisinopril, atenolol, enalapril, perindopril, perindopril tert-butylamine, trandolapril and moexipril, or a pharmaceutically acceptable salt form of one or more of these compounds.

The invention also provides methods of using PTPase
25 inhibitors of formula A for improving the cardiovascular or cerebrovascular risk profile in patients experiencing or subject to type II diabetes (non-insulin-dependent diabetes mellitus), preferably in human type II diabetics or a patient experiencing or subject to Syndrome X. These methods may also
30 be characterized as the reduction of risk factors for heart disease, stroke or heart attack in a type II diabetic or a patient experiencing or subject to Syndrome X.

The invention also provides methods of using a pharmacological combination of one or more PTPase inhibiting agents, one or more biguanide agents, and, optionally one or more sulfonylurea agents for treatment of type II diabetes or Syndrome X in a patient in need of such treatment. Also provided are methods of using these agents to treat or inhibit metabolic disorders mediated by insulin resistance or hyperglycemia in a patient in need thereof. Further included in this invention is a method of modulating blood glucose levels in a patient in need thereof.

Each of these methods comprises administering to a patient in need thereof pharmaceutically effective amounts of:

- a) a PTPase inhibiting agent of formula I; and
- b) a biguanide agent; and
- 15 c) optionally, a sulfonylurea agent.

Biguanide agents useful with this invention include metformin and its pharmaceutically acceptable salt forms. Sulfonylurea agents useful for the methods and combinations of this invention may be selected from the group of glyburide, glyburide, glipizide, glimepiride, chlorpropamide, tolbutamide, 20 or tolazamide, or a pharmaceutically acceptable salt form of these agents.

This invention also provides pharmaceutical compositions and methods of using PTPase inhibitors of formula A in combination with one or more alpha-glucosidase inhibitors, such as miglitol or acarbose, for improving the cardiovascular risk profile in patients experiencing or subject to Syndrome X or type II diabetes (non-insulin-dependent diabetes mellitus), preferably in human type II diabetics. These methods may also be characterized as the reduction of risk factors for heart disease, stroke or heart attack in a patient in such need.

These methods include the reduction of hyperlipidemia in type II diabetics, including methods in type II diabetics for

lowering low density lipoprotein (LDL) blood levels and to increase high density lipoprotein (HDL) blood levels. The methods herein may further be characterized as useful for inhibiting, preventing or reducing atherosclerosis in a type II diabetic or a patient experiencing or subject to Syndrome X, or the risk factors of either.

These methods also include the lowering free fatty acid blood levels and triglyceride levels in type II diabetics, or a patient experiencing or subject to Syndrome X.

Among the alpha-glucosidase inhibitors which may be utilized with the invention described herein are miglitol or acarbose, or a pharmaceutically acceptable salt form of one or more of these compounds.

This invention further provides methods for using a PTPase inhibitor of the invention and a sulfonylurea agent for the management of Syndrome X or type 2 diabetes and for improving the cardiovascular risk profile in patients experiencing or subject to those maladies. These methods may also be characterized as the reduction of risk factors in such patients for heart disease, stroke or heart attack in a type II diabetic. Such methods include the reduction of hyperlipidemia in a patients experiencing or subject to Syndrome X or type II diabetes and include methods for lowering low density lipoprotein (LDL) blood levels, high density lipoprotein (HDL) blood levels, and overall blood lipoprotein levels. The methods herein may further be characterized as inhibiting, preventing or reducing atherosclerosis in patients subject to or experiencing Syndrome X or type II diabetes, or the risk factors thereof. Such methods further include the lowering of free fatty acid blood levels and triglyceride levels in such patients.

Representative sulfonylurea agents include glipizide, glyburide (glibenclamide), chlorpropamide, tolbutamide,

tolazamide and glimepiride, or the pharmaceutically acceptable salt forms thereof.

In addition, the invention provides combinations of a PTPase inhibitor of the invention and at least one
5 thiazolidinedione agents. Such combinations are useful for treatment, inhibition or maintenance of Syndrome X or type II diabetes in patients in need of such treatment. Accordingly, methods of using such combinations are provided by the invention. Thus, the invention provides methods of using these
10 agents to treat or inhibit metabolic disorders mediated by insulin resistance or hyperglycemia in patients in need thereof. Further included in this invention are methods of modulating blood glucose levels in a patient in need thereof.

Each of these methods comprises administering to a patient
15 in need thereof pharmaceutically effective amounts of:

a) a thiazolidinedione agent, such as selected from the group of pioglitazone and rosiglitazone, or a pharmaceutically acceptable salt form of these agents; and

b) a compound of formula A.

20 The invention also provides pharmaceutical compositions and methods of using PTPase inhibitors in combination with one or more antilipemic agents. Such methods and compositions are useful for improving the cardiovascular risk profile in patients experiencing or subject to type II diabetes (non-
25 insulin-dependent diabetes mellitus), preferably in type II diabetics or Syndrome X. These methods also include reducing the risk factors for heart disease, stroke or heart attack in a type II diabetic or a patient experiencing or subject to Syndrome X. Such methods further include the reduction of
30 hyperlipidemia in type II diabetics, including such methods in type II diabetics for lowering low density lipoprotein (LDL) blood levels and to increase high density lipoprotein (HDL) blood levels. These compositions and methods are also useful

for inhibiting, preventing or reducing atherosclerosis in a type II diabetic or a patient experiencing or subject to Syndrome X, or the risk factors thereof. In this aspect, the compositions and methods are useful for lowering of free fatty acid blood levels and triglyceride levels in type II diabetics, or patients experiencing or subject to Syndrome X.

Representative antilipemic or agents, also known as antihyperlipidemic agents, suitable for use in the invention are bile acid sequestrants, fibric acid derivatives, HMG-CoA reductase inhibitors and nicotinic acid compounds. Bile acid sequestrant agents useful with this invention include colestipol and colesvelam, and their pharmaceutically acceptable salt forms. Fibric acid derivatives which may be used with the present invention include clifofibrate, gemfibrozil and fenofibrate. HMG-CoA reductase inhibitors, also known as statins, useful with this invention include cerivastatin, fluvastatin, atorvastatin, lovastatin, pravastatin and simvastatin, or the pharmaceutically acceptable salt forms thereof. Niacin is an example of a nicotinic acid compound which may be used with the methods of this invention. Also useful are lipase inhibiting agents, such as orlistat.

This invention also provides pharmaceutical compositions that are a combination of a compound of Formula A and an aldose reductase inhibitor (ARI). Such combinations are useful in methods for treating, inhibiting or preventing type II diabetes, or its related and associated symptoms, disorders and maladies. These methods comprise administering to a patient in need of such therapy a pharmaceutically effective amount of a composition comprising a combination of pharmaceutically effective amounts of a compound of formula A and an ARI. These compositions and methods are useful for the treatment, prevention or inhibition of diabetic neuropathy, diabetic

nephropathy, retinopathy, keratopathy, diabetic uveitis, cataracts.

Representative suitable ARIs are disclosed in U.S. Patent Nos. 6,420,426 and 6,214,991.

5 Combinations of the compounds of Formula A and an ARI are also useful for inhibition or reduction of risk factors for heart disease, stroke or heart attack in a type II diabetic. Therefore, in this aspect the invention is useful for reducing hyperlipidemia and/or low density lipoprotein (LDL) blood
10 levels in type II diabetics. Also included in this aspect are methods for inhibiting, preventing or reducing atherosclerosis or the risk factors thereof in type II diabetics. This aspect includes lowering of free fatty acid blood levels and triglyceride levels.

15 This invention also provides methods of using a compound of formula A and insulin(s) for the management of type I or type II diabetes. Accordingly, the invention provides for combination therapy, i.e., where a compound of Formula A is administered in combination with insulin. Such combination
20 therapy encompasses simultaneous or sequential administration of the compound of Formula A and insulin. The insulins useful in this aspect include both naturally occurring and synthetic insulins.

Insulins useful with the methods and combinations of this
25 invention include rapid acting insulins, intermediate acting insulins, long acting insulins and combinations of intermediate and rapid acting insulins.

Rapid acting commercially available insulin products include HUMALOG® Brand Lispro Injection (rDNA origin); HUMULIN®
30 Regular Human Injection, USP [rDNA origin]; HUMULIN® Regular U-500 Concentrated Human Injection, USP [rDNA origin]; REGULAR ILETIN® II (insulin injection, USP, purified pork) available from Eli Lilly and Co.; and the NOVALIN® Human Insulin

Injection and VENOSULIN® BR Buffered Regular Human Injection, each available from Novo Nordisk Pharmaceuticals.

Commercially available intermediate acting insulins useful with this invention include, but are not limited to, the
5 HUMULIN® L brand LENTE® human insulin [rDNA origin] zinc suspension, HUMULIN® N NPH human insulin [rDNA origin] isophane suspension, LENTE® ILETIN.RTM. II insulin zinc suspension, USP, purified pork, and NPH ILETIN® II isophane insulin suspension, USP, purified pork, available from Eli Lilly and Company,
10 LANTUS® insulin glargine [rDNA origin] injection, available from Aventis Pharmaceuticals, and the NOVOLIN L Lente® human insulin zinc suspension (recombinant DNA origin), and NOVOLIN® N NPH human insulin isophane suspension (recombinant DNA origin) products available from Novo Nordisk Pharmaceuticals,
15 Inc, Princeton N.J.

Also useful with the methods and formulations of this invention are intermediate and rapid acting insulin combinations, such as the HUMALOG® Mix 75/25 (75% Insulin Lispro Protamine Suspension and 25% Insulin Lispro Injection),
20 HUMULIN® 50/50 (50% Human Insulin Isophane Suspension and 50% Human Insulin Injection) and HUMULIN® 70/30 (70% Human Insulin Isophane Suspension and 30% Human Insulin Injection), each available from Eli Lilly and Company. Also useful are the NOVALIN® 70/30 (70% NPH, Human Insulin Isophane Suspension and
25 30% Regular, Human Insulin Injection) line of combination products available from Novo Nordisk Pharmaceuticals.


A commercially available long acting insulin for use with this invention is the HUMULIN® U Ultralente® human insulin [rDNA origin] extended zinc suspension, available from Eli
30 Lilly and Company.

Also useful in the methods of this invention are inhaled insulin products, such as the EXUBERA® inhaled insulin product developed by Pfizer Inc. and Aventis SA.

Each of these insulin products can be administered as directed by a medical professional using administrations, dosages and regimens known in the art, such as those published for each product in the Physicians' Desk Reference, 55 Edition, 5 2001, published by Medical Economics Company, Inc. at Montvale, N.J., the relevant sections of which are incorporated herein by reference. In this aspect, the invention includes, for example, methods for improving the cardiovascular and cerebrovascular risk profiles in patients experiencing or subject to type I or 10 type II diabetes (non-insulin-dependent diabetes mellitus), preferably in human type II diabetics. These methods may also be characterized as the inhibition or reduction of risk factors for heart disease, stroke or heart attack in a type II diabetic.

15 The compounds of the present invention may be prepared by use of known chemical reactions and procedures. Representative methods for synthesizing compounds of the invention are presented below. It is understood that the nature of the substituents required for the desired target compound often 20 determines the preferred method of synthesis. All variable groups of these methods are as described in the generic description if they are not specifically defined below.

Methods of Preparation



Scheme 1

A method for preparing compounds of the invention is illustrated in scheme 1. Certain compounds of the invention are prepared from a substituted 4-bromobenzenesulfonylchloride or 4-bromobenzeneacid chloride as illustrated in scheme 1.

Treatment with the desired amino acid ester in the presence of a base, such as pyridine or triethylamine, gives the corresponding sulfonamide or amide. Activation of the aryl bromide by treatment with bis(pinacolato)diboron and a palladium catalyst give the boronic ester, which is subsequently coupled to a variety of aryl or heteroaryl bromides or iodides using a palladium catalyst. For some examples, the desired aryl or heteroaryl bromide may need to be prepared separately. In general the preparation of these intermediates can be accomplished using methods known in the art.

Once the L₃-Q group is in place, the sulfonamide or amine nitrogen can be alkylated with the desired side chain alkyl halide. This is usually done with a base, such as cesium carbonate, or sodium hydride. Finally, the ester intermediate is hydrolyzed to give the target compound.

Those having skill in the art will recognize that the starting materials and reaction conditions may be varied, the sequence of the reactions altered, and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples. In some cases, protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general, the need for such protecting groups as well as the conditions necessary to attach and remove such groups will be apparent to those skilled in the art of organic synthesis.

The disclosures of all articles and references mentioned in this application, including patents, are incorporated herein by reference in their entirety.


Structures were named using Name Pro IUPAC Naming Software, version 5.09, available from Advanced Chemical Development, Inc., 90 Adelaide Street West, Toronto, Ontario, M5H 3V9, Canada or with ChemDraw v. 6.02, which is available from Cambridgesoft.com in Cambridge, MA.

CHEMISTRY EXAMPLES

The preparation of intermediates and compounds of the invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures and compounds described in them. In all cases, unless otherwise specified, the column chromatography is performed using a silica gel solid phase.

Example 1. Preparation of 4-Dibenzofuran-4-yl-phenyl-boronic acid

Step 1: (4-Dibenzofuran-4-yl-phenyl)-trimethyl-silane




5

A solution of dibenzofuran-4-yl-boronic acid (20.0 g, 94.3 mmol), (4-bromo-phenyl)-trimethyl-silane (21.62 g, 94.3 mmol), K_2CO_3 (39.1 g, 3 equiv., 283 mmol) in toluene (100 mL), ethanol (60 mL) and water (30 mL) was purged with nitrogen for 5 min (bubbled into solution) and treated with $Pd(PPh_3)_4$ (3.59 g, 2.9 mmol). After heating to 80 °C for 4 h, the solution was cooled to room temperature, poured into water (300 mL) and extracted with ethyl acetate (300 mL). The organic phase was washed with sat'd aq NaCl, dried over anhydrous $MgSO_4$, filtered and concentrated in vacuo. Purification by flash column chromatography (5-20% ethyl acetate in heptane) afforded (4-dibenzofuran-4-yl-phenyl)-trimethyl-silane as a colorless oil (28.9 g, 96%).

10

15



20


Step 2: 4-Dibenzofuran-4-yl-phenyl-boronic acid

25

A solution of (4-dibenzofuran-4-yl-phenyl)-trimethyl-silane (28.6 g, 90.2 mmol) in dichloromethane (350 mL, 0.26 M) was cooled to -78 °C and carefully treated with borontribromide (135 mL, 1.5 equiv., 135 mmol). After the addition was complete, the solution was warmed to room temperature and

stirred for 3 h. Next, the reaction mixture was re-cooled to -78 °C, treated with dry methanol (30 mL), slowly warmed to room temperature and stirred for 1.5 h. Next, the solution was re-cooled to -78 °C, carefully quenched with 10% aq HCl (50 mL), warmed to room temperature and stirred for 1 h (solids form). The resulting solution was poured into water (500 mL) and extracted with ethyl acetate (3 X 700 mL). The combined organic layers were washed with sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. The crude product was suspended in a 10% ethyl acetate in heptane solution, filtered and washed with the same solution (5 X 60 mL) to give 4-dibenzofuran-4-yl-phenyl-boronic acid as a white solid (20.2 g, 77%).

Example 2. 5-Benzyl-5-{2-[4'-(2-benzylbenzofuran-3-yl)-biphen-4-yl]-2-oxoethyl}-2,2-dimethyl-[1,3]-dioxane-4,6-dione




A solution of 5-{2-[4'-(2-benzylbenzofuran-3-yl)-biphen-4-yl]-2-oxoethyl}-2,2-dimethyl-[1,3]dioxane-4,6-dione (200 mg, 0.37 mmol) in THF/DMF (5: 1; 6 mL) was added dropwise to a stirred suspension of sodium hydride (95%, 10.2 mg, 0.40 mmol) in anhydrous THF (5 mL) at room temperature. The clear solution was stirred at room temperature for 30 mins and then a solution of benzyl bromide (76 mg, 0.44 mmol) in THF (5 mL) was added dropwise, followed by the addition of tetra-n-butylammonium iodide (5 mg) as a solid. The reaction mixture was warmed to 60°C for 4 hrs (TLC control), cooled to room temperature and

then water (10 mL) was added cautiously. The reaction mixture was extracted with diethyl ether (3 x 15 mL). The combined extract was washed with water (2 x 10 mL), brine (3 x 10 mL), dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*.

5 Purification of the product by trituration and filtration from MeOH afforded the title compound as a pale yellow solid (210 mg). $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): δ 8.12 (2H, d, $J = 8$ Hz, Ar-H), 7.88 (4H, d, $J = 6$ Hz, Ar-H), 7.53-7.69 (4H, m, Ar-H), 7.26 (1H, m, Ar-H), 7.14 (1H, m, Ar-H), 4.26 (2H, s, PhCH_2), 4.16
10 (2H, CH_2CO), 3.38 (2H, s, PhCH_2), 1.98 (3H, s, Me), 0.78 (3H, s, Me).

Example 3. 1-(4-Bromophenyl)-1H-indole.




15

A solution of 1H-indole (3.0 g, 25.6 mmol), 4-fluorobromobenzene (4.48 g, 25.6 mmol), potassium fluoride (40% wt on alumina; 3.0 g) and 18-crown-6 (690 mg, 2.56 mmol) in anhydrous DMSO (30 mL) was heated at 150°C for 24 hours, and
20 then cooled to room temperature. The reaction mixture was poured into water (50 mL) and extracted with diethyl ether (3 x 50 mL). The combined organic extract was washed with water (2 x 30 mL), brine (3 x 30 mL), dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by
25 flash column chromatography, using 20 % ethyl acetate/hexane as eluent, afforded the title compound as a pale yellow solid (5.5 g, 76%).

Example 4. 4'-Indol-1-yl-biphenyl-4-carbaldehyde.

To a stirred solution of the bromide (from example 3) (7.77 g, 28.6 mmol) and *tetrakis*-(triphenylphosphine)palladium(0) (1.8 g, 1.45 mmol) in toluene (100 mL) was added a solution of 4-formylphenylboronic acid (5.21 g, 34.5 mmol) in ethanol (20 mL) and 2N sodium carbonate (28.6 mL, 57.2 mmol). The resulting suspension was stirred at 90°C for 4 hrs (TLC control). The reaction was cooled, diluted with water (50 mL) and extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. The resulting brown solid was redissolved in tetrahydrofuran (50 mL). 2N Hydrochloric acid (10 mL) was added and the resulting solution was stirred at room temperature for 1 hour, and then diluted with water (50 mL) and extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 20% ethyl acetate in heptane as eluent, afforded the title compound as a white solid (8.02 g, 94 %), ¹H NMR (CDCl₃, 300 MHz) δ 10.1 (1H, s, CHO), 8.01 (2H, d, J = 8 Hz, Ar-H), 7.70 (5H, m, Ar-H), 7.62 (2H, d, J = 8 Hz, Ar-H), 7.39 (1H, d, J = 3.5 Hz, Ar-H), 7.22 (3H, m, Ar-H), 6.74 (1H, d, J = 3.5 Hz, Ar-H).

Example 5. (4'-Indol-1-yl-biphen-4-yl)methanol.



Sodium borohydride (783 mg, 20.6 mmol) was added portion-wise to a stirred solution of aldehyde (prepared in example 4) (3.06 g, 10.3 mmol) in a mixture of anhydrous THF and ethanol


(1:1; 100 mL) at room temperature. The reaction mixture was stirred for 10 minutes at room temperature (TLC control), poured into water (50 mL) and acidified to pH 4 with 2N hydrochloric acid, and then extracted with diethyl ether (3 x 5 20 mL). The combined extract was washed with 0.5 N hydrochloric acid (2 x 10 mL), water and finally brine. The ethereal solution was dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 30 % ethyl acetate/hexane as 10 eluent, afforded the title compound as a white solid (2.80 g, 91 %); ¹H NMR (CDCl₃, 300 MHz) δ 7.55 – 7.78 (7H, m, Ar-H), 7.52 (2H, d, J = 8 Hz, Ar-H), 7.41 (1H, d, J = 3.5 Hz, Ar-H), 7.22 (3H, m, Ar-H), 6.72 (1H, d, J = 3.5 Hz, Ar-H), 4.79 (2H, d, J = 5.5 Hz, CH₂O).

15

Example 6. Methanesulfonic acid, (4'-Indol-1-yl-biphen-4-yl)methyl ester.

Methanesulfonyl chloride (194 mg, 131 μL, 1.7 mmol) was 20 added dropwise to a cooled (0°C) solution of alcohol (prepared in example 5) (620 mg, 1.54 mmol) and triethylamine (311 mg, 0.43 mL, 3.08 mmol) in anhydrous methylene chloride (10 mL). The clear reaction mixture was stirred at 0°C for 2-4 hrs (TLC control), then poured into water (50 mL), and extracted with 25 diethyl ether (3 x 30 mL). The combined extract was washed with 0.5 N hydrochloric acid (2 x 10 mL), water and finally brine. The ethereal solution was dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. The crude mesylate was used in the subsequent alkylation step without further 30 purification.

Example 7. 4'-Dibenzofuran-4-yl-biphenyl-4-carbaldehyde



A solution of dibenzofuran-4-boronic acid (1.0 g, 4.7 mmol) in ethanol (10 mL) was added to a stirred solution of 1-bromo-4-iodobenzene (1.33 g, 4.7 mmol) and *tetrakis-*
5 (triphenylphosphine)palladium(0) (271 mg, 5 mol%) in toluene (40 mL). 2N sodium carbonate (4.7 mL, 9.4 mmol) was added and the reaction was heated to 90°C (oil bath temp.) for 2-3 hrs until complete (TLC control).

10 The reaction mixture was cooled to room temperature and partitioned between water and diethyl ether. The phases were separated, the aqueous phase being further extracted with diethyl ether (2 x 20 mL). The combined extract was washed with water and brine. The ethereal solution was dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to yield 4-
15 (4-bromophenyl)-dibenzofuran as a yellow solid, which was used immediately without further purification.

A solution of 4-formylphenylboronic acid (0.9 g, 5.64 mmol) in ethanol (10 mL) was added to a stirred solution of the crude 4-(4-bromophenyl)-dibenzofuran (from the previous
20 reaction) in toluene (40 mL). *tetrakis-* (Triphenylphosphine)palladium(0) (270 mg, 5 mol%) and 2N sodium carbonate (4.7 mL, 9.4 mmol) were added and the reaction was heated to 100°C (oil bath temp.) for 2-3 hrs until complete (TLC control). The reaction mixture was cooled to room
25 temperature and partitioned between water and ethyl acetate. The phases were separated, the aqueous phase being further extracted with ethyl acetate (2 x 20 mL). The combined extract was washed with 0.5 N hydrochloric acid, water and brine and

then dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 10-20% ethyl acetate in hexane as eluent, afforded the title compound as a white solid (1.51g).

5

Example 8. (4'-Dibenzofuran-4-yl-biphenyl-4-yl)methanol

Sodium borohydride (322 mg, 8.4 mmol) was added portion-wise to a stirred solution of aldehyde (prepared in example 7) (1.48 g, 4.2 mmol) in a mixture of anhydrous THF and ethanol (1:2; 50mL) at room temperature. The reaction mixture was stirred for 5-10 minutes at room temperature (TLC control), poured into water (50 mL) and acidified to pH 4 with 2N hydrochloric acid, and then extracted with diethyl ether (3 x 30 mL). The combined extract was washed with 0.5 N hydrochloric acid (2 x 10 mL), water and finally brine. The ethereal solution was dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 30 % ethyl acetate/hexane as eluent, afforded the title compound as a white solid (1.40 g).

20

Example 9. Methanesulfonic acid, 4'-dibenzofuran-4-yl-biphenyl-4-yl-methyl ester

Methanesulfonyl chloride (490 mg, 330 μL , 4.3 mmol) was added dropwise to a cooled (0°C) solution of alcohol (prepared in example 8) (1.38 g, 3.9 mmol) and triethylamine (800 mg, 1.1 mL, 7.9 mmol) in anhydrous methylene chloride (50 mL). The clear reaction mixture was stirred at 0°C for 2-4 hrs (TLC control), then poured into water (50 mL), and extracted with diethyl ether (3 x 30 mL). The combined extract was washed with 0.5 N hydrochloric acid (2 x 10 mL), water and finally brine. The ethereal solution was dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. The crude mesylate was

30


used in the subsequent alkylation step without further purification.

Example 10. Trifluoro-methanesulfonic acid 4-(2-benzyl-
5 benzofuran-3-yl)-phenyl ester.

To a stirred solution of the known phenol, 4-(2-benzylbenzofuran-3-yl)-phenol, (7.2 g, 24 mmol) in anhydrous methylene chloride (100 mL), was added triethylamine (4.86 g, 6.7 mL, 48 mmol) and then *N*-phenyltrifluoromethanesulfonimide
10 (9.4g, 26.4 mmol) portionwise as a solid. The resulting solution was stirred for 2 hours at room temperature and then diluted with water, extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*.

15 Purification of the product by flash column chromatography, using 10% ethyl acetate in heptane as eluent, afforded the title compound as a white solid (9.35 g, 90%).

Example 11. 4'-(2-Benzylbenzofuran-3-yl)biphenyl-4-
20 carbaldehyde.




To a stirred solution of the triflate (from example 10) (9.35 g, 21.6 mmol) and *tetrakis*-(triphenylphosphine)palladium(0) (750 mg, 0.65 mmol) in toluene
25 (70 mL) was added a solution of 4-formylphenylboronic acid (4.06g, 27.05 mmol) in ethanol (20 mL) and 2N sodium carbonate

(21.6 mL, 43.2 mmol). The resulting suspension was stirred at 100°C for 4 hrs (TLC control). The reaction was cooled, diluted with water (50 mL) and extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. The resulting brown solid was redissolved in tetrahydrofuran (50 mL). 2N Hydrochloric acid (10 mL) was added and the resulting solution was stirred at room temperature for 1 hour, and then diluted with water (50 mL) and extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 20% ethyl acetate in heptane as eluent, afforded the title compound as a white solid (7.34g, 88%).

Example 12. 4'-(2-Benzylbenzofuran-3-yl)biphenyl-4-methanol

To a solution of 4'-(2-benzylbenzofuran-3-yl)biphenyl-4-carbaldehyde (5.0g, 12.9 mmol) in ethanol (100 mL) and tetrahydrofuran (25 mL) was added sodium borohydride (980 mg, 25.8 mmol) as a solid in 3 portions. The reaction was stirred at room temperature for 1 hour (TLC control) and then poured into water (100 mL) and extracted with diethyl ether (3 x 100 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to afford the title compound as a white solid (5.02g, 99%).

Example 13. 2-Benzyl-3-(4'-bromomethylbiphen-4-yl)benzofuran



To a solution of 4'-(2-benzylbenzofuran-3-yl)biphenyl-4-
 5 methanol (5.01g, 12.7 mmol) in anhydrous acetonitrile (75 mL)
 was added dibromotriphenylphosphorane (5.45g, 12.7 mmol) as a
 solid portionwise over 15 mins. The reaction was stirred for 2
 hours (TLC control) and then poured into water (100 mL) and
 extracted with diethyl ether (3 x 100 mL). The combined extract
 10 was washed with water, brine, dried over anhydrous MgSO_4 ,
 filtered and concentrated *in vacuo* to afford the title compound
 as an off-white solid (4.98g, 87%): ^1H NMR (CDCl_3 , 300 MHz): δ
 7.70 (2H, m, Ar-H), 7.63 (4H, m, Ar-H), 7.50 (3H, m, Ar-H),
 7.30 (4H, m, Ar-H), 7.25 (4H, m, Ar-H), 4.57 (2H, s, CH_2Br),
 15 4.26 (2H, PhCH_2).

Example 14. 4'-Bromo-biphenyl-4-carboxylic acid methyl ester.

A mixture of methyl 4-iodobenzoate, 9.38g (35.8 mmol), 4-
 bromophenylboronic acid 7.18g (35.8 mmol), $\text{Pd}(\text{PPh}_3)_4$, 2.07g
 20 (1.79 mmol), in 180mL of toluene and 100mL of ethanol was
 heated to obtain a clear solution. To the solution was added
 30mL of 4.0M aq. Na_2CO_3 . The reaction mixture refluxed for 4h
 at 80 °C. The mixture was cooled to room temperature and
 diluted with 300mL ethyl acetate. The organic layer was washed
 25 with 2x300mL portions of water, 2x300mL portions of sat. aq.
 NaCl , and dried (MgSO_4). After the solution was concentrated,

the residue was purified by column chromatography (eluted with 7% EtOAc-Heptane) to afford the desired product in 7.8g (78%) as a white solid. ¹H NMR (CDCl₃) 8.10 (d, 2H, J = 9.0Hz), 7.62 (d, 2H, J = 9.0Hz), 7.59 (d, 2H, J = 9.3Hz), 7.48 (d, 2H, J = 9.3 Hz), 3.95 (s, 3H).

Example 15. (4'-Bromo-biphenyl-4-yl)-methanol.


A solution of 4'-Bromo-biphenyl-4-carboxylic acid methyl ester, 7.8g (27.9 mmol) in 150mL of tetrahydrofuran was cooled to 0 °C via ice-water bath. Lithiumaluminum-hydride, 1.1g (27.9 mmol) was added to the solution in one portion. The reaction mixture stirred at 0 °C for 1h. The mixture was slowly quenched with 10mL of isopropyl alcohol, then with 10mL of water. The aqueous mixture was extracted with 3x50mL portions of ethyl acetate. The organic layers were combined, washed with sat. aq. NaCl, and dried (MgSO₄). The solution was concentrated to afford the desired product in 7.01g (100%) as a white solid. The material was taken to the next step without further purification.

Example 16. 4'-Bromo-4-bromomethyl-biphenyl.

A solution of (4'-bromo-biphenyl-4-yl)-methanol, 7.01g (27.9 mmol) and dibromo-triphenylphosphorane 11.8g (27.9 mmol) in 150mL of methylene chloride stirred at room temperature for 2h. The solution was diluted with 100mL of water and extracted with 2x200mL portions of diethyl ether. The organic layers were combined, washed with sat. aq. NaCl, and dried (MgSO₄). After the solution was concentrated, the residue was purified through a short plug of silica gel (eluted with 50% EtOAc-Heptane) to afford the desired product in 9.1g (100%) as a white solid. The material was taken to the next step without further purification.

Example 17. (4-Dibenzofuran-4-yl-phenyl)boronic acid


Step 1: (4-Dibenzofuran-4-yl-phenyl)-trimethyl-silane



5

A solution of dibenzofuran-4-yl-boronic acid (20.0 g, 94.3 mmol), (4-bromo-phenyl)-trimethyl-silane (21.62 g, 94.3 mmol), K_2CO_3 (39.1 g, 3 equiv., 283 mmol) in toluene (100 mL), ethanol (60 mL) and water (30 mL) was purged with nitrogen for 5 min (bubbled into solution) and treated with $Pd(PPh_3)_4$ (3.59 g, 2.9 mmol). After heating to 80 °C for 4 h, the solution was cooled to room temperature, poured into water (300 mL) and extracted with ethyl acetate (300 mL). The organic phase was washed with sat'd aq NaCl, dried over anhydrous $MgSO_4$, filtered and concentrated in vacuo. Purification by flash column chromatography (5-20% ethyl acetate in heptane) afforded (4-dibenzofuran-4-yl-phenyl)-trimethyl-silane as a colorless oil (28.9 g, 96%).

20 Step 2: 4-Dibenzofuran-4-yl-phenyl-boronic acid



A solution of (4-dibenzofuran-4-yl-phenyl)-trimethyl-silane (28.6 g, 90.2 mmol) in dichloromethane (350 mL, 0.26 M) was cooled to -78 °C and carefully treated with borontribromide (135 mL, 1.5 equiv., 135 mmol). After the addition was complete, the solution was warmed to room temperature and stirred for 3 h. Next, the reaction mixture was re-cooled to -78 °C, treated

25

with dry methanol (30 mL), slowly warmed to room temperature and stirred for 1.5 h. Next, the solution was re-cooled to -78 °C, carefully quenched with 10% aq HCl (50 mL), warmed to room temperature and stirred for 1 h (solids form). The resulting
5 solution was poured into water (500 mL) and extracted with ethyl acetate (3 X 700 mL). The combined organic layers were washed with sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. The crude product was suspended in
10 a 10% ethyl acetate in heptane solution, filtered and washed with the same solution (5 X 60 mL) to give 4-dibenzofuran-4-yl-phenyl-boronic acid as a white solid (20.2 g, 77%).

Example 18. Preparation of [[4'-(2-Benzyl-benzofuran-3-yl)-
15 biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-acetic acid

Step 1: Preparation of Tert-Butyl-[(4-bromobenzenesulfonyl)-(3-trifluoromethylbenzyl)amino]acetate.

20 3-Trifluoromethylbenzyl bromide (313 mg, 1.31 mmol) was added dropwise to a stirred suspension of *tert*-butyl-(4-bromobenzenesulfonylamino)acetate (404 mg, 1.16 mmol) and cesium carbonate (768 mg, 2.36 mmol) in anhyd DMF (5 mL). The resultant reaction mixture was stirred at room temperature for
25 24 h, diluted with ethyl acetate (20 mL), washed with sat'd aq LiCl (3 x 10 mL), sat'd aq NaCl, dried over anhyd MgSO₄, filtered and concentrated in vacuo. Purification by flash column chromatography (10% ethyl acetate in heptane) afforded the title compound as a colorless oil (293 mg).


Step 2: Preparation of Tert-Butyl-{[4'-(2-benzylbenzofuran-3-yl)biphenyl-4-sulfonyl]-(3-trifluoromethylbenzyl)amino}acetate.

5 A solution of 2-benzyl-3-[4'-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)phenyl]-benzofuran (268 mg 0.653 mmol) in anhyd DMSO (5 mL) was added to a stirred suspension of tert-Butyl-[(4-bromobenzenesulfonyl)-(3-trifluoromethylbenzyl)amino]-acetate (296 mg, 0.58 mmol) and
10 tripotassium phosphate (0.37 g, 1.76 mmol) in anhyd DMSO (5 mL). [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II)-DCM complex (60 mg, 0.07 mmol) was added as a solid, and the resulting suspension was heated to 80 °C for 25 h. The
15 reaction mixture was cooled to room temperature, diluted with ethyl acetate, filtered through celite, and washed with sat'd aq LiCl (3 x 10 mL), sat'd aq NaCl (1 x 10 mL), dried over anhyd MgSO₄, filtered and concentrated in vacuo. Purification by flash column chromatography (heptane) afforded the title
20 compound as a white solid (95 mg).

Step 3: [[4'-(2-Benzyl-benzofuran-3-yl)biphenyl-4-sulfonyl]-(3-trifluoromethylbenzyl)-amino]acetic acid.


Tert-Butyl-{[4'-(2-benzylbenzofuran-3-yl)biphenyl-4-sulfonyl]-(3-trifluoromethylbenzyl)amino}acetate (90 mg) was
25 dissolved in methylene chloride (4 mL). TFA (1 mL) was added and the reaction mixture was stirred at room temperature for 16 h, and then concentrated in vacuo. Purification by flash column chromatography (5% methanol in methylene chloride) afforded the
30 title compound as an off white solid (45 mg).

Example 19. Preparation of N-[4'-(2-Butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-yl]-oxalamic acid



5

Step 1: (4-Bromo-phenyl)-(2-butyl-benzofuran-3-yl)-methanone




10

A solution of 2-n-butylbenzofurane (19.8 g, 114 mmol) and 4-bromobenzoyl chloride (25.0 g, 114 mmol) in dry dichloromethane (300 mL, 0.4 M) was cooled to 0 °C and treated with AlCl₃ (16.6 g, 1.1 equiv., 125.4 mmol) in 3 portions. After the additions were complete, the solution was stirred for 3 h and carefully added to ice water. After separation, the aqueous layer was extracted with dichloromethane (2 X 200 mL) and the combined organic layers were washed with water, sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by flash column chromatography (1-2% ethyl acetate in heptane) afforded (4-bromo-phenyl)-(2-butyl-benzofuran-3-yl)-methanone (14.6 g, 36%).


15

20




Step 2: 3-(4-Bromo-benzyl)-2-butyl-benzofuran

A solution of (4-bromo-phenyl)-(2-butyl-benzofuran-3-yl)-
5 methanone (2.25 g, 6.32 mmol) in ethanol (20 mL, 0.3 M) was
cooled to 0 °C and treated with NaBH₄ (0.263 g, 1.1 equiv, 6.95
mmol). After stirring for 1 h, the mixture was poured into a
50% ether in water solution (200 mL). After separation, the
aqueous layer was extracted with ether (50 mL) and the combined
10 organic layers were washed with water, sat'd aq NaCl, dried
over anhydrous MgSO₄, filtered and concentrated in vacuo.
The resulting alcohol was subsequently dissolved in dry
dichloromethane (50 mL), cooled to 0 °C and treated with
triethylsilane (2.0 mL, 2.0 equiv., 12.64 mmol) dropwise via
15 syringe. After stirring an additional 5 min, trifluoroacetic
acid (2.43 mL, 5.0 equiv., 31.6 mmol) was added over 2 min and
the mixture was stirred for 3 h. Once complete, the solution
was washed with water, sat'd aq NaCl, dried over anhydrous
MgSO₄, filtered and concentrated in vacuo. Purification by
20 flash column chromatography (0-2% ethyl acetate in heptane)
afforded 3-(4-bromo-benzyl)-2-butyl-benzofuran as a pale yellow
oil (1.34 g, 63%).



Step 3: 2-Butyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzyl]-benzofuran

A solution of 3-(4-bromo-benzyl)-2-butyl-benzofuran (14.03 g, 41.5 mmol), bis(pinacolato)diborane (11.60 g, 1.1 equiv., 45.7 mmol), potassium acetate (12.2 g, 3.0 equiv., 125 mmol) in DMSO (100 mL, 0.4 M) was treated with $\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ (4.15 g, 0.1 equiv., 4.15 mmol) and heated to 80 °C. After complete by TLC, the solution was cooled to room temperature, diluted with water (150 mL) and filtered through celite (washed with ether, 500 mL). After separation, the aqueous layer was extracted with ether (2 X 150 mL). The combined organic layers were washed with water, sat'd aq NaCl, dried over anhydrous MgSO_4 , filtered and concentrated in vacuo. Purification by flash column chromatography (2-5% ethyl acetate in heptane) afforded 2-butyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzyl]-benzofuran as a pale yellow oil (11.2 g, 69%).




Step 4: 2-Butyl-3-[3'-nitro-4'-(3-phenyl-propoxy)-biphenyl-4-ylmethyl]-benzofuran

A solution of 2-butyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-benzyl]-benzofuran (100 mg, 0.256 mmol), 4-bromo-2-nitro-1-(3-phenylpropylox)-benzene (95 mg, 1.1 equiv., 0.282 mmol) and K_2CO_3 (136 mg, 3.5 equiv., 0.987 mmol) in water (1 mL) and DMF (2 mL) was treated with $\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ (23 mg, 0.1 equiv., 0.0282 mmol) and heated to 120 °C. After complete (by TLC) the solution was cooled to room temperature, acidified to pH < 4 with 10% aq HCl and

diluted with water (20 mL). After separation, the aqueous layer was extracted with ether (3 X 20 mL) and the combined organic layers were washed with water, sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo.


- 5 Purification by prep thin layer chromatography afforded 2-butyl-3-[3'-nitro-4'-(3-phenyl-propoxy)-biphenyl-4-ylmethyl]-benzofuran (53 mg, 41%).



- 10 *Step 5: 4'-(2-Butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-ylamine*


- A solution of 2-butyl-3-[3'-nitro-4'-(3-phenyl-propoxy)-biphenyl-4-ylmethyl]-benzofuran (53 mg, 0.105 mmol) in ethanol
 15 (1 mL) and acetic acid (1 mL) was treated with Fe (26.4 mg, 4.5 equiv., 0.472 mmol) and heated to 120 oC for 3 h. After cooling to room temperature, the solution was poured into a 20% aq NaOH / ice water solution (Ph > 8) and extracted with ether
 20 (3 X 20 mL). The combined organic layers were washed with water, sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by prep thin layer chromatography (25% ethyl acetate in heptane) afforded 4'-(2-butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-ylamine (14.5 mg, 29%).

25



Step 6: *N*-[4'-(2-Butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-yl]-oxalamic acid ethyl ester

A solution of 4'-(2-butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-ylamine (129 mg, 0.264 mmol) and diisopropylethylamine (0.115 mL, 2.5 equiv., 0.66 mmol) in dichloromethane (5 mL) was treated with a solution of ethyl chlorooxoacetate (44 mg, 1.2 equiv., 0.317 mmol) in dichloromethane (1 mL). After stirring for 2 h, the solution was diluted with water and extracted with dichloromethane (2 X 15 mL). The combined organic layers were washed with water, sat'd aq NaCl, dried over anhydrous MgSO₄, filtered and concentrated in vacuo. Purification by prep thin layer chromatography (25% ethyl acetate in heptane) afforded *N*-[4'-(2-butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-yl]-oxalamic acid ethyl ester (120 mg, 77%).



Step 7: *N*-[4'-(2-Butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-yl]-oxalamic acid

A solution of *N*-[4'-(2-butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-biphenyl-3-yl]-oxalamic acid ethyl ester (120 mg, 0.204 mmol) in ethanol (3 mL) was treated with aq 1 N NaOH (0.3 mL, 1.5 equiv., 0.306 mmol) and stirred at room temperature. After stirring 1 h, the solution was acidified to pH < 4 with 10% HCl, concentrated and purified by prep thin layer chromatography (10% methanol in dichloromethane) afforded *N*-[4'-(2-Butyl-benzofuran-3-ylmethyl)-4-(3-phenyl-propoxy)-

biphenyl-3-yl]-oxalamic acid (32 mg, 28%). R_f 0.39 (10% methanol in dichloromethane), ^1H NMR (DMSO- d_6 , 300 MHz) δ 10.26 (s, 1 H), 8.58 (d, J = 2.1 Hz, 1 H), 7.43-7.05 (m, 16 H), 4.07 (t, J = 6.0 Hz, 2 H), 3.95 (s, 2 H), 2.82-2.77 (m, 4 H), 2.13-1.97 (m, 2 H), 1.67-1.59 (m, 2 H), 1.37-1.24 (m, 2 H), 0.86 (t, J = 7.5 Hz, 3 H).

Example 20. Preparation of 4-[4'-(2-benzylbenzofuran-3-yl)biphen-4-yl]-4-oxobutyric acid

10 *Step 1. 5-[2-(4-Bromophenyl)-2-oxoethyl]-2,2-dimethyl-[1,3]dioxane-4,6-dione*

A solution of Meldrum's acid (5.0 g, 35 mmol) in anhydrous THF (25 mL) was added cautiously to a stirred suspension of sodium hydride (95%, 960 mg, 38 mmol) in anhydrous THF (25 mL). The resulting solution was stirred at room temperature for 1 hr. A solution of 2,4'-dibromoacetophenone (11.6 g, 42 mmol) in anhydrous THF (25 mL) was added dropwise, and the resultant solution was stirred at room temperature for 16-24 hrs (TLC control). The reaction mixture was poured into water (50 mL), acidified to pH 2-3 with 0.5N hydrochloric acid and extracted with ethyl acetate (3 x 50 mL). The combined extract was washed with water, brine, dried over anhydrous $MgSO_4$, filtered and concentrated *in vacuo*. Trituration and filtration from MeOH afforded the title compound as a white solid (6.56 g).

Step 2. 2-Benzyl-3-[4-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)phenyl]-benzofuran

30 A solution of bis-(pinacolato)diboron (2.64 g, 10.41 mmol) in anhydrous DMSO (20 mL) was added to a stirred suspension of the known triflate, trifluoromethanesulfonic acid-4-(2-benzylbenzofuran-3-yl)phenyl ester, (4.09 g, 9.47 mmol) and potassium acetate (3.71 g, 37.9 mmol) in anhydrous

DMSO (20 mL). [1,1'-*bis*-(Diphenylphosphino)-ferrocene]dichloropalladium(II)-DCM complex (770 mg, 0.95 mmol) was added as a solid, and the resulting suspension was heated to 80°C for 4 hrs. The reaction mixture was cooled to room temperature, diluted with diethyl ether (150 mL), washed with water (2 x 50 mL), brine (3 x 50 mL), dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 10 % ethyl acetate/heptane as eluent, afforded the title compound as a white solid (2.96 g).


Step 3. 5-{2-[4'-(2-Benzylbenzofuran-3-yl)-biphen-4-yl]-2-oxoethyl}-2,2-dimethyl-[1,3]-dioxane-4,6-dione

A solution of 2-benzyl-3-[4'-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolan-2-yl)phenyl]-benzofuran (500 mg, 1.22 mmol) in anhydrous DMSO (5 mL) was added to a stirred suspension of 5-[2-(4-bromophenyl)-2-oxoethyl]-2,2-dimethyl-[1,3]dioxane-4,6-dione (436 mg, 1.22 mmol) and tripotassium phosphate (1.04 g, 4.88 mmol) in anhydrous DMSO (5 mL). [1,1'-*bis*-(Diphenylphosphino)ferrocene]dichloropalladium(II)-DCM complex (100 mg, 0.12 mmol) was added as a solid, and the resulting suspension was heated to 80°C for 2 hrs. The reaction mixture was cooled to room temperature, diluted with diethyl ether (150 mL), washed with water (2 x 50 mL), brine (3 x 50 mL), dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 50-60 % ethyl acetate/heptane as eluent, afforded the title compound as an off-white solid (502 mg): ¹H-NMR (DMSO-d₆, 300 MHz): δ 8.12 (2H, d, J = 8 Hz, Ar-H), 7.92 (4H, d, J = 6 Hz, Ar-H), 7.53-7.69 (4H, m, Ar-H), 7.26 (7H, m, Ar-H), 4.82 (1H, s, CH), 4.26 (2H, s, PhCH₂), 3.86 (2H, CH₂CO), 1.84 (3H, s, Me), 1.74 (3H, s, Me).


Step 4. 4-[4'-(2-benzylbenzofuran-3-yl)biphen-4-yl]-4-oxobutyric acid

2N Hydrochloric acid (1 mL) was added to a stirred solution of 5-{2-[4'-(2-benzyl-benzofuran-3-yl)-biphen-4-yl]-2-oxoethyl}-2,2-dimethyl-[1,3]-dioxane-4,6-dione (200 mg, 0.36 mmol) in THF (10 mL), and the resultant solution was heated at 70°C for 6 hrs and then cooled to room temperature and concentrated *in vacuo*. The resulting solid was redissolved in DMSO (10 mL), and heated to 150°C for 3 hrs before being cooled to room temperature, and diluted with water (20 mL), and extracted with ethyl acetate (3 x 20 mL). The combined extract was washed with water (2 x 10 mL), brine (3 x 10 mL), dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by trituration and filtration from MeOH afforded the title compound as a white solid (105 mg, 62%). Rf: 0.20 (5% methanol in dichloromethane); ¹H-NMR (CDCl₃, 300 MHz): δ 8.10 (2H, d, J = 8 Hz, Ar-H), 7.80 (5H, m, Ar-H), 7.62 (1H, d, J = 8 Hz, Ar-H), 7.48 (1H, d, J = 8 Hz, Ar-H), 7.29 (6H, m, Ar-H), 4.26 (2H, s, PhCH₂), 3.86 (2H, t, J = 6 Hz), 2.86 (2H, t, J = 6 Hz); ESI-LCMS e/z calcd for C₃₁H₂₄O₄: 460.527, found 461 (M+H)⁺.

Example 21. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid




Step 1. 3-(4-bromophenylmethylsulfanyl)-propanoic acid methyl ester



A solution of 3-thiolpropanoic acid methyl ester (1.2 g, 10 mmol) and 4-bromobenzyl-bromide (2.5 g, 10 mmol) in DMF (20 mL) was cooled to 0 °C and treated with Cs₂CO₃ (3.9 g, 12 mmol). After stirred for 2 h, the reaction was quenched with 5% HCl (25 mL) and diluted with ethyl acetate (50 mL). After separation, the aqueous layer extracted with ethyl acetate (3 x 15 mL). The combined organic layers were washed with sat. aq NaCl, dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-5% ethyl acetate in heptane) provided 3-(4-bromophenylmethylsulfanyl)-propanoic acid methyl ester (2.5 g, 87%) as white solid. ¹H NMR (CDCl₃), 7.43 (d, *J* = 8 Hz, 2 H), 7.20 (d, *J* = 8 Hz, 2 H), 3.68 (s, 5 H), 2.67 (m, 2 H), 2.56 (t, *J* = 6 Hz, 2 H).

Step 2. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid methyl ester




20

A solution of 3-(4-bromophenylmethylsulfanyl)-propanoic acid methyl ester (289 mg, 1 mmol), 4-(4-dibenzofuranyl)benzeneboronic acid (302 mg, 1.05 mmol) and Pd(PPh₃)₄ (52 mg, 5mol%) in toluene (10 mL) and ethanol (3.0 mL) was treated with 2 M K₂CO₃ (1.5 mL). The reaction mixture was heated to reflux for 2 h, cooled to room temperature,

diluted with ethyl acetate (100 mL). The organic layer was washed successively with 2% aq HCl and sat. aq NaCl, dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-10% ethyl acetate in heptane) gave 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid methyl ester (330 mg, 72%) as white solid. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.81 (s, 2 H), 3.71 (s, 3 H), 2.76 (t, *J* = 6 Hz, 2 H), 2.56 (t, *J* = 6 Hz, 2 H).
LCMS 475 (M⁺ + 23).

Step 3. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid



15

A solution of 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid methyl ester (210 mg, 0.46 mmol) in THF (2 mL) and methanol (2 mL) was cooled to 0 °C and treated with 2 N KOH (1.0 mL). After stirring at room temperature for 1 h the solution was acidified with 10% HCl to pH 2 and diluted with ethyl acetate (25 mL). After being separated, the aqueous layer was extracted with ethyl acetate (3 x 15 mL) and the combined organic layers were dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-5% methanol in dichloromethane) provided 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid (180 mg, 90%) as white solid. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8

Hz, 2 H), 7.94 (d, $J = 8$ Hz, 2 H), 7.78 (d, $J = 8$ Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.82 (s, 2 H), 3.82 (s, 2 H), 2.76 (m, 2 H), 2.56 (t, $J = 6$ Hz, 2 H). LCMS 462 ($M^+ + 23$).

5 Example 22. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2-methylpropanoic acid
3-(4-bromophenylmethylsulfanyl)-2-methylpropanoic acid methyl ester

10 A solution of 3-bromo-2-methylpropanoic acid methyl ester (0.366 g, 2 mmol) and 4-bromobenzylthiol (0.402 g, 2 mmol) in DMF (5 mL) was cooled to 0 °C and treated with K_2CO_3 (414 g, 3 mmol). After stirring for 2 h, the reaction was quenched with 5% HCl (15 mL) and diluted with ethyl acetate (50 mL). After
15 separation, the aqueous layer extracted with ethyl acetate (3 x 15 mL). The combined organic layers were washed with sat. aq NaCl, dried over $MgSO_4$ and concentrated. Purification by flash column chromatography (5% ethyl acetate in heptane) provided 3-(4-bromophenylmethylsulfanyl)-2-methylpropanoic acid
20 methyl ester (430 mg, 71%) as white solid. ^1H NMR ($CDCl_3$), 7.43 (d, $J = 8$ Hz, 2 H), 7.20 (d, $J = 8$ Hz, 2 H), 3.69 (s, 3 H), 3.65 (s, 2 H), 2.67 (m, 2 H), 2.45 (q, $J = 6$ Hz, 1 H), 1.21 (d, $J = 7$ Hz, 3 H).

25 Step 1. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2-methylpropanoic acid methyl ester

A solution of 3-(4-bromophenylmethylsulfanyl)-2-methylpropanoic acid methyl ester (0.427 g, 1 mmol), 4-(4-dibenzofuranyl)benzeneboronic acid
30 (0.302 g, 1.05 mmol) and $Pd(PPh_3)_4$ (0.052g, 5 mol%) in toluene (10 mL) and ethanol (3.0 mL) was treated with 2 M K_2CO_3 (1.5 mL). The reaction mixture was heated to reflux for 2 h, cooled to room temperature, diluted with ethyl acetate (100 mL). The


organic layer was washed successively with 2% aq HCl and sat. aq NaCl, dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-10% ethyl acetate in heptane) gave 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid methyl ester (312 mg, 67%) as an oil. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.79 (s, 2 H), 3.71 (s, 3 H), 2.76 (m, 2 H), 2.56 (q, *J* = 6 Hz, 1 H). LCMS 480 (M⁺ + 23).

10


Step 2. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2-methylpropanoic acid

A solution of 2-*tert*-butoxycarbonylamino-3-(4'-dibenzofuran-4-yl-biphen-4-yl)-propanoic acid methyl ester (0.28 g, 0.6 mmol) in THF (2 mL) and methanol (2 mL) was cooled to 0 °C and treated with 2 N KOH (1.0 mL). After stirring at room temperature for 1 h the solution was acidified with 10% HCl to pH 2 and diluted with ethyl acetate (25 mL). After being separated, the aqueous layer was extracted with ethyl acetate (3 x 15 mL) and the combined organic layers were dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-5% methanol in dichloromethane) provided 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid (245 mg, 91%) as white solid. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.81 (s, 2 H), 2.76 (m, 2 H), 2.56 (q, *J* = 6 Hz, 1 H). LCMS 476 (M⁺ + 23).

Example 23. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2, 2-dimethylpropanoic acid




Step 1. 3-(4-bromophenylmethylsulfanyl)-2, 2-dimethylpropanoic acid methyl ester



A solution of 3-bromo-2,2-dimethylpropanoic acid methyl ester (0.39 g, 2 mmol) and 4-bromobenzylthiol (0.402 g, 2 mmol) in DMF (5 mL) was cooled to 0 °C and treated with K₂CO₃ (0.414 g, 3 mmol). After stirring for 2 h, the reaction was quenched with 5% HCl (15 mL) and diluted with ethyl acetate (50 mL). After separation, the aqueous layer extracted with ethyl acetate (3 x 15 mL). The combined organic layers were washed with sat. aq NaCl, dried over MgSO₄ and concentrated. Purification by flash column chromatography (5% ethyl acetate in heptane) provided 3-(4-bromophenylmethylsulfanyl)-2, 2-dimethylpropanoic acid methyl ester (260 mg, 35%) as white solid. ¹H NMR (CDCl₃), 7.43 (d, *J* = 8 Hz, 2 H), 7.18 (d, *J* = 8 Hz, 2 H), 3.69 (s, 5 H), 2.63 (s, 2 H), 1.21 (s, 6 H).

Step 2. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2-dimethylpropanoic acid methyl ester




5

A solution of 3-(4-bromophenylmethylsulfanyl)-2-methylpropanoic acid methyl ester (0.260 g, 0.82 mmol), 4-(4-dibenzofuranyl)benzeneboronic acid (0.236 g, 0.86 mmol) and Pd(PPh₃)₄ (0.052g, 5 mol%) in toluene (10 mL) and ethanol (3.0 mL) was treated with 2 M K₂CO₃ (1.5 mL). The reaction mixture was heated to reflux for 2 h, cooled to room temperature, diluted with ethyl acetate (100 mL). The organic layer was washed successively with 2% aq HCl and sat. aq NaCl, dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-10% ethyl acetate in heptane) gave 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid methyl ester (280 mg, 67%) as an oil. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.79 (s, 2 H), 3.71 (s, 3 H), 2.73 (s, 2 H), 1.27 (s, 6 H). LCMS 504 (M⁺ + 23).

20

Step 3. 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-2-dimethylpropanoic acid



5

A solution of 2-*tert*-butoxycarbonylamino-3-(4'-dibenzofuran-4-yl-biphen-4-yl)-propanoic acid methyl ester (0.125 g, 0.26 mmol) in THF (2 mL) and methanol (2 mL) was cooled to 0 °C and treated with 2 N KOH (1.0 mL). After stirring at room temperature for 1 h the solution was acidified with 10% HCl to pH 2 and diluted with ethyl acetate (25 mL). After being separated, the aqueous layer was extracted with ethyl acetate (3 x 15 mL) and the combined organic layers were dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-5% methanol in dichloromethane) provided 3-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-propanoic acid (80 mg, 67%) as white solid. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.82 (s, 2 H), 2.76 (s, 2 H), 1.32 (s, 6 H). LCMS 490 (M⁺ + 23).

Example 24. 2-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-3-methylbutanoic acid

(4-bromophen-yl-4-methylsulfanyl)-3-methylbutanoic acid

ethyl ester

A solution of 2-bromo-3-methylpropanoic acid ethyl ester (0.418 g, 2 mmol) and 4-bromobenzylthiol (0.402 g, 2 mmol) in

DMF (5 mL) was cooled to 0 °C and treated with K₂CO₃ (0.414 g, 3 mmol). After stirring for 2 h, the reaction was quenched with 5% HCl (15 mL) and diluted with ethyl acetate (50 mL).

After separation, the aqueous layer extracted with ethyl

5 acetate (3 x 15 mL). The combined organic layers were washed with sat. aq NaCl, dried over MgSO₄ and concentrated.

Purification by flash column chromatography (5% ethyl acetate in heptane) provided (4-bromophen-yl-4-methylsulfanyl)-3-methylbutanoic acid ethyl ester (660 mg, 90%) as white solid.

10 ¹H NMR (CDCl₃), 7.43 (d, *J* = 8 Hz, 2 H), 7.18 (d, *J* = 8 Hz, 2 H), 4.17 (q, *J* = 6 Hz, 2 H), 3.75 (s, 2 H), 2.85 (d, *J* = 9 Hz, 1 H), 2.04 (m, 1 H), 1.29 (t, *J* = 6 Hz, 3 H), 1.02 (d, *J* = 6 Hz, 3 H), 0.98 (d, *J* = 6 Hz, 3 H).

15 Step 1. 2-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-3-methylbutanoic acid methyl ester

A solution of (4-bromophen-yl-4-methylsulfanyl)-3-methylbutanoic acid methyl ester (0.331 g, 1 mmol), 4-(4-dibenzofuranyl)benzeneboronic acid (0.305 g, 1.05 mmol) and

20 Pd(PPh₃)₄ (0.052 g, 5%mol) in toluene (10 mL) and ethanol (3.0 mL) was treated with 2 M K₂CO₃ (1.5 mL). The reaction mixture was heated to reflux for 2 h, cooled to room temperature, diluted with ethyl acetate (100 mL). The organic layer was washed successively with 2% aq HCl and sat. aq NaCl, dried over

25 MgSO₄ and concentrated. Purification by flash column chromatography (2-10% ethyl acetate in heptane) gave 2-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-3-methylbutanoic acid methyl ester (336 mg, 70%) as an oil. ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 4.17 (q, *J* = 6 Hz, 2 H), 3.85 (s, 2 H), 2.95 (d, *J* = 9 Hz, 1 H), 2.04 (m, 1 H), 1.29 (t, *J* = 6 Hz, 3 H), 1.05 (d, *J* = 6 Hz, 3 H), 0.99 (d, *J* = 6 Hz, 3 H). LCMS 517 (M⁺ + 23).

Step 2. 2-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-3-methylbutanoic acid

A solution of 2-*tert*-butoxycarbonylamino-3-(4'-dibenzofuran-4-yl-biphen-4-yl)-propanoic acid methyl ester (0.28 g, 0.566 mmol) in THF (2 mL) and methanol (2 mL) was cooled to 0 °C and treated with 2 N KOH (1.0 mL). After stirring at room temperature for 1 h the solution was acidified with 10% HCl to pH 2 and diluted with ethyl acetate (25 mL). After being
separated, the aqueous layer was extracted with ethyl acetate (3 x 15 mL) and the combined organic layers were dried over MgSO₄ and concentrated. Purification by flash column chromatography (2-5% methanol in dichloromethane) provided 2-(4'-dibenzofuran-4-yl-biphen-4-ylmethylsulfanyl)-3-methylbutanoic acid (195 mg, 74%) as white solid. 0.195g (yield 73.8 %). ¹H NMR (CDCl₃), 8.01 (d, *J* = 8 Hz, 2 H), 7.94 (d, *J* = 8 Hz, 2 H), 7.78 (d, *J* = 8 Hz, 2 H), 7.66 (m, 4 H), 7.42 (m, 5 H), 3.85 (s, 2 H), 2.95 (d, *J* = 9 Hz, 1 H), 2.12 (m, 1 H), 1.05 (t, *J* = 6 Hz, 6 H). LCMS 489 (M⁺ + 23).

20

The following compounds are prepared essentially according to the procedures described in the schemes, charts, examples and preparations set forth herein.

25 Example 25. N-[3-Benzoyloxy-4'-(2-butyl-benzofuran-3-ylmethyl)-biphenyl-4-yl]-oxalamic acid.

R_f 0.19 (10% methanol in dichloromethane), ¹H NMR (DMSO-d₆, 300 MHz) δ 9.78 (s, 1 H), 8.31 (s, 1 H), 7.48-7.08 (m, 16 H), 5.25 (s, 2 H), 4.01 (s, 2 H), 2.83 (t, *J* = 7.5 Hz, 2 H), 1.68-1.58 (m 2 H), 1.37-1.27 (m, 2 H), 0.87 (t, *J* = 7.2 Hz, 3 H); ESI-LCMS *m/z* calcd for C₃₄H₃₁NO₅: 533; found 532 (M - 1)⁺.

30

Example 26. N-[3-Benzyloxy-4'-(2-butyl-benzofuran-3-ylmethyl)-biphenyl-4-yl]-malonamic acid.

R_f 0.32 (10% methanol in dichloromethane), ^1H NMR ($CDCl_3$, 300 MHz) δ 8.54 (s, 1 H), 8.41 (s, 1 H), 7.48-7.01 (m, 16 H), 5.16 (s, 1 H), 4.02 (s, 1 H), 3.47 (s, 1 H), 1.77-1.67 (m, 2 H), 1.45-1.33 (m, 2 H), 0.93 (t, $J = 7.2$ Hz, 3 H); ESI-LCMS m/z calcd for $C_{35}H_{33}NO_5$: 547; found 548 ($M + 1$)⁺.


Example 27. N-[4-Benzyloxy-4'-(2-butyl-benzofuran-3-ylmethyl)-biphenyl-3-yl]-malonamic acid.

R_f 0.34 (10% methanol in dichloromethane), ^1H NMR ($CDCl_3$, 300 MHz) δ 8.54 (s, 1 H), 8.34 (s, 1 H), 7.48-7.01 (m, 16 H), 5.16 (s, 2 H), 4.02 (s, 2 H), 3.50 (s, 2 H), 2.79 (t, $J = 7.5$ Hz, 2 H), 1.75-1.67 (m, 2 H), 1.45-1.37 (m, 2 H), 0.93 (t, $J = 6.9$ Hz, 3 H); ESI-LCMS m/z calcd for $C_{35}H_{33}NO_5$: 547; found 548 ($M + 1$)⁺.

Example 28. N-[4'-(2-Butyl-benzofuran-3-ylmethyl)-3-hydroxybiphenyl-4-yl]-malonamic acid.

R_f 0.53 (20% methanol in dichloromethane), ^1H NMR ($CDCl_3$, 300 MHz) δ 9.22 (br s, 1 H), 7.87 (br s, 1 H), 7.40-6.99 (m, 11 H), 3.95 (s, 2 H), 3.56 (s, 2 H), 2.72 (t, $J = 7.5$ Hz, 2 H), 1.68-1.49 (m, 2 H), 1.36-1.28 (m, 2 H), 0.86 (t, $J = 6.9$ Hz, 3 H); ESI-LCMS m/z calcd for $C_{28}H_{27}NO_5$: 457; found 458 ($M + 1$)⁺.

Example 29. N-[4'-(2-Butyl-benzofuran-3-ylmethyl)-4-hydroxybiphenyl-3-yl]-malonamic acid.




R_f 0.53 (20% methanol in dichloromethane), ¹H NMR (CDCl₃, 300 MHz) δ 9.22 (br s, 1 H), 7.39–6.99 (m, 11 H), 3.95 (s, 2 H), 3.56 (s, 2 H), 2.72 (t, J = 7.5 Hz, 2 H), 1.68–1.49 (m, 2 H), 1.36–1.28 (m, 2 H), 0.86 (t, J = 6.9 Hz, 3 H); ESI-LCMS m/z calcd for C₂₈H₂₇NO₅: 457; found 458 (M + 1)⁺.

Example 30. [4'-(2-Benzyl-benzofuran-3-yl)-3-fluoro-biphenyl-4-sulfonylamino]-acetic acid

Isolated as a white solid. R_f 0.42 (20% Methanol-80% Methylene Chloride); ¹H NMR (DMSO-d₆) 8.35 (br. s, 1H), 7.95–7.54 (m, 9H), 7.31–7.21 (m, 6H), 4.27 (s, 2H), 3.73 (s, 2H); LCMS m/z calcd for C₂₉H₂₂FO₅ S : 515.5 found 516.3 (M+1).

Example 31. {[4'-(2-Benzyl-benzofuran-3-yl)-3-fluoro-biphenyl-4-sulfonyl]-methyl-amino}-acetic acid




Isolated as an off-white solid . R_f 0.32 (10% Methanol-90% Methylene Chloride); ¹H NMR (CDCl₃) 7.98–7.24 (m, 15H), 4.26 (s, 2H), 4.21 (s, 2H), 3.04 (s, 3H); LCMS m/z calcd for C₃₀H₂₄FO₅ S : 529.57 found 530.3 (M+1).

Example 32. [[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-nitro-benzyl)-amino]-acetic acid

Isolated as a beige solid. R_f 0.50 (10% Methanol-90% Methylene Chloride); ¹H NMR (DMSO-d₆) 8.10–7.00 (m, 21H), 4.63 (s, 2H), 4.27 (s, 2H), 3.64 (s, 2H).

Example 33. [[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-acetic acid



5

Isolated as a white foam. , R_f 0.61 (50% Ethyl Acetate-50% Heptane); ^1H NMR (CDCl_3) 7.94 (d, $J=7.5\text{Hz}$, 2H), 7.75-7.02 (m, 19H), 4.57 (s, 2H), 4.23 (s, 2H), 4.02 (s, 2H).

10 Example 34. [[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-methoxy-benzyl)-amino]-acetic acid

Isolated as a white foam. R_f 0.62 (20% Methanol-80% Ethyl Acetate); ^1H NMR ($\text{DMSO}-d_6$) 8.00-7.87 (m, 6H), 7.70-7.55 (4H), 7.36-7.17 (7H), 6.81-6.71 (m, 4H), 4.53 (s, 2H), 4.28 (s, 2H),
 15 3.65 (s, 3H), 3.58 (s, 2H).

Example 35. [[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-fluoro-benzyl)-amino]-acetic acid


Isolated as an off-white solid. R_f 0.38 (10% Methanol-90% Methylene Chloride); ^1H NMR ($\text{DMSO}-d_6$) 7.92-7.90 (m, 6H), 7.70-7.56 (m, 4H), 7.35-7.23 (m, 8H), 7.07 (t, $J=9.0\text{ Hz}$, 3H), 4.52 (s, 2H), 4.29 (s, 2H), 3.75 (s, 2H).

25 Example 36. [[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-acetic acid

Isolated as an off-white solid. R_f 0.43 (10% Methanol-90% Methylene Chloride); ^1H NMR ($\text{DMSO}-d_6$) 7.94-7.90 (m, 6H), 7.71-

7.57 (m, 8H), 7.35-7.26 (m, 7H), 4.58 (s, 2H), 4.30 (s, 2H), 3.88 (s, 2H).

Example 37. 2-[[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-4-methyl-pentanoic acid




Isolated as white foam. R_f 0.31 (10% Methanol-90% Methylene Chloride); ^1H NMR (DMSO- d_6) 7.96-7.88 (m, 6H), 7.78-7.57 (m, 8H), 7.34-7.23 (m, 7H), 4.85 (d, $J=17.1$ Hz, 1H), 4.53 (d, $J=16.8$ Hz, 1H), 4.47-4.42 (m, 1H), 4.29 (s, 2H), 1.43-1.24 (m, 3H), 0.83 (d, $J=6.0$ Hz, 3H), 0.46 (d, $J=6.3$ Hz, 3H).

Example 38. 2-[[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-butyric acid

Isolated as a white foam. R_f 0.37 (10% Methanol-90% Methylene Chloride); ^1H NMR (DMSO- d_6) 7.93-7.85 (m, 6H), 7.74-7.54 (m, 7H), 7.34-7.23 (m, 7H), 4.76 (d, $J=17.1$ Hz, 1H), 4.53 (d, $J=16.8$ Hz, 1H), 4.34 (m, 1H), 4.32 (s, 2H), 1.78 (m, 1H), 1.47 (m, 1H), 0.75 (t, $J=6.9$ Hz, 3H).

Example 39. [(4'-Dibenzofuran-4-yl-biphenyl-4-sulfonyl)-(3-trifluoromethyl-benzyl)-amino]-acetic acid




Isolated as a white foam. R_f 0.24 (10% Methanol-90%

5 Methylene Chloride); ^1H NMR (DMSO- d_6) 12.8 (br s, 1H), 8.29 (s, 1H), 8.21-7.40 (m, 18H), 4.57 (s, 2 H), 3.99 (s, 2H); LCMS m/z calcd for $\text{C}_{34}\text{H}_{24}\text{F}_3\text{NO}_5\text{S}$: 615.6 found 616.3 (M+1).

Example 40. 2-[[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-sulfonyl]-(3-trifluoromethyl-benzyl)-amino]-propionic acid

10 Isolated as a white foam. R_f 0.24 (10% Methanol-90% Methylene Chloride); ^1H NMR (DMSO- d_6) 7.01-6.91 (m, 6H), 6.79-8.71 (m, 8H), 6.41-6.31 (m, 7H), 4.00 (s, 2H), 3.94-3.83 (m, 2H), 3.69 (d, J = 16.8 Hz, 1H), 0.46 (d, J = 7.5Hz, 3H); LCMS m/z 15 calcd for $\text{C}_{38}\text{H}_{30}\text{F}_3\text{NO}_5\text{S}$: 669.7 found 670.3 (M+1).

Example 41. [(2-Phenoxy-[1,1';4',1'']terphenyl-4'''-sulfonyl)-(3-trifluoromethyl-benzyl)-amino]-acetic acid



20


Isolated as a white foam. R_f 0.22 (10% Methanol-90% Methylene Chloride); ^1H NMR (DMSO- d_6 with TFA) 7.77-7.83 (m,

21H), 4.43 (s, 2H), 3.86 (s, 2H); LCMS m/z calcd for $C_{34}H_{26}F_3NO_5$
S : 617.6 found 618.3 (M+1).

Example 42. [(4-Propyl-[1,1';4',1'']terphenyl-4''-sulfonyl)-
5 (3-trifluoromethyl-benzyl)-amino]-acetic acid


Isolated as a white foam. R_f 0.30 (10% Methanol-90%
Methylene Chloride); ^1H NMR (DMSO- d_6) 7.93 (d, $J=8.1$ Hz, 2H),
7.76-7.63 (m, 6H), 7.55 (d, $J=8.4$ Hz, 2H), 7.49-7.41 (m, 4H),
7.28 (d, $J=8.1$ Hz, 2H), 4.58 (s, 2H), 3.98 (s, 2H), 2.66 (t,
10 $J=7.5$ Hz, 2H), 1.70 (m, 2H), 0.99 (t, $J=7.5$ Hz, 3H); LCMS m/z
calcd for $C_{31}H_{28}F_3NO_4$ S : 567.62 found 566.3 (M+1).

Example 43. [(4'-Dibenzofuran-4-yl-biphenyl-4-sulfonyl)-(3-
fluoro-benzyl)-amino]-acetic acid




Isolated as an off-white solid. R_f 0.48 (20% Methanol-80%
Methylene Chloride); ^1H NMR (DMSO- d_6) 12.80 (br s, 1H), 8.19-
7.09 (m, 19H), 4.49 (s, 2H), 3.95 (s, 2H).

Example 44. 2-(3'-Dibenzofuran-4-yl-biphen-3-
20 ylmethylsulfanyl)-propionic acid.




Step 1. 4-(3-Bromophenyl)-dibenzofuran



A solution of dibenzofuran-4-boronic acid (1.0 g, 4.7 mmol) in ethanol (10 mL) was added to a stirred solution of 1-bromo-3-iodobenzene (1.33 g, 4.7 mmol) and tetrakis-(triphenylphosphine)palladium(0) (271 mg, 5 mol%) in toluene (40 mL). 2N sodium carbonate (4.7 mL, 9.4 mmol) was added and the reaction was heated to 90°C (oil bath temp.) for 2-3 hrs until complete (TLC control).

The reaction mixture was cooled to room temperature and partitioned between water and diethyl ether. The phases were separated, the aqueous phase being further extracted with diethyl ether (2 x 20 mL). The combined extract was washed with water and brine. The ethereal solution was dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo* to yield 4-(3-bromophenyl)-dibenzofuran as a yellow solid. Trituration of the solid with methanol, followed by vacuum filtration (with methanol washes), yields the title compound, 4-(3-bromophenyl)-dibenzofuran, as a pale yellow solid (1.32 g, 87%).


Step 2. Methyl-3-dibenzofuran-4-ylbiphen-3-yl-carboxylate



A solution of 3-methoxycarbonyl-phenylboronic acid (0.62 g, 3.4 mmol) in methanol (10 mL) was added to a stirred solution of 4-(3-bromophenyl)-dibenzofuran (1.0 g, 3.1 mmol) in toluene (40

mL). *tetrakis*-(Triphenylphosphine)palladium(0) (180 mg, 5 mol%) and 2N sodium carbonate (3.1 mL, 6.2 mmol) were added and the reaction was heated to 90°C (oil bath temp.) for 2-3 hrs until complete (TLC control). The reaction mixture was cooled to room temperature and partitioned between water and ethyl acetate. The phases were separated, the aqueous phase being further extracted with ethyl acetate (2 x 20 mL). The combined extract was washed with 0.5 N hydrochloric acid, water and brine and then dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 10-20% ethyl acetate in hexane as eluent, afforded the title compound; methyl-3-dibenzofuran-4-ylbiphen-3-yl-carboxylate has a white solid (1.10 g, 94%).


Step 3. (3'-Dibenzofuran-4-ylbiphen-3-yl)methanol



Lithium aluminium hydride (3.5 mL, 1.0 M solution in THF, 3.5 mmol) was added dropwise to a stirred solution of ester (prepared in the previous step) (900 mg, 2.38 mmol) in anhydrous THF (15mL) at room temperature. The reaction mixture was stirred for 30 minutes at room temperature (TLC control), and then cooled to 0°C, and quenched with water (0.15 mL), 2N sodium hydroxide (0.15 mL) and water (0.45 mL). The reaction mixture was diluted with diethyl ether (30 mL) and then filtered through celite. The ethereal solution was dried over anhydrous MgSO₄, filtered and concentrated *in vacuo*, affording the title compound; (3'-dibenzofuran-4-ylbiphen-3-yl)methanol,

as a white solid (820 mg, 98%), that did not require further purification.

Step 4. 4-(3'-Bromomethyl-biphen-3-yl)-dibenzofuran




5

Dibromotriphenylphosphorane (630 mg, 1.5 mmol) was added to a stirred solution of alcohol; (3'-dibenzofuran-4-ylbiphen-3-yl)methanol (260 mg, 0.75 mmol) in dichloromethane (15 mL).

The reaction mixture was stirred for 60 minutes at room
10 temperature (TLC control), and then diluted with diethyl ether (30 mL) and washed with water (2x) and brine. The ethereal solution was dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 10-20% ethyl acetate in hexane as
15 eluent, afforded the title compound; 4-(3'-bromomethyl-biphen-3-yl)-dibenzofuran, as a colorless oil (266 mg, 86%).

Step 5. Methyl-2-(3'-dibenzofuran-4-yl-biphen-3-ylmethylsulfanyl)-propionate.




20

Ethyl-2-mercaptopropionate (31 mg, 0.29 mmol) was added to a stirred suspension of 4-(3'-bromomethyl-biphen-3-yl)-

dibenzofuran (100 mg, 0.24 mmol) and cesium carbonate (198 mg, 0.6 mmol) in 5ml DMF. diluted with diethyl ether (30 mL) and washed with water (2x) and brine (3x). The ethereal solution was dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 30 % ethyl acetate in hexane as eluent, afforded the title compound; methyl-2-(3'-dibenzofuran-4-yl-biphen-3-ylmethylsulfanyl)-propionate, as a colorless oil (105 mg, 94%).

Step 6. 2-(3'-Dibenzofuran-4-yl-biphen-3-ylmethylsulfanyl)-propionic acid.







2N Sodium hydroxide solution (0.33 mL, 0.66 mmol) was added dropwise to a stirred solution of methyl-2-(3'-dibenzofuran-4-yl-biphen-3-ylmethylsulfanyl)-propionate (100 mg, 0.22 mmol) in tetrahydrofuran (5 mL) and methanol (1 mL). The clear reaction mixture was stirred at room temperature until the reaction was complete (TLC control), and then diluted with water (5 mL), and acidified to pH 3 with 2N hydrochloric acid. The reaction mixture was extracted with ethyl acetate (2 x 20 mL). The combined extract was washed with water, brine, dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo*. Purification of the product by flash column chromatography, using 5 - 10% methanol in methylene chloride as eluent, afforded the title compound has a white solid (86 mg, 89%); Rf:

0.40 (10% methanol in dichloromethane); ^1H NMR (DMSO- d_6):

8.19 (2H, m, Ar-H), 8.02 (2H, d, $J = 9$ Hz, Ar-H), 7.86 (2H, d, $J = 9$ Hz, Ar-H), 7.75 (4H, m, Ar-H), 7.29 - 7.41 (5H, m, Ar-H), 3.96 (2H, s, PhCH_2S), 3.36 (1H, q, $J = 7$ Hz, SCHMe), 1.36 (3H, d, $J = 7$ Hz, CHMe); ESI-LCMS m/z calcd for $\text{C}_{28}\text{H}_{22}\text{O}_3\text{S}$ 438.54, found 439 ($\text{M}+\text{H}$) $^+$.


Example 45

The following compounds are prepared essentially according to the procedures described in the above schemes and examples.

Cmpd. No.	Structure	Name
1		4-tert-butoxycarbonylamino-5-(4'-dibenzofuran-4-yl-biphenyl-4-yl)-pent-2-enoic acid
2		3-(4'-Dibenzofuran-4-yl-biphenyl-4-yl)-acrylic acid
3		3-(3'-Dibenzofuran-4-yl-biphenyl-4-ylmethylsulfanyl)-propionic acid
4		(3'-Dibenzofuran-4-yl-biphenyl-3-ylmethylsulfanyl)-acetic acid


Example 46

The following compounds are prepared essentially according to the procedures described in the above schemes and examples.



A.





































OR





























































B.

5





































Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
9	10	bond		H		H	H
11	12	bond		F		H	H
13	14	CO		H		H	H
15	16	CO		F		H	H
17	18	SO ₂		H		H	H
19	20	SO ₂		F		H	H
21	22	-CH ₂ O-		H		H	H
23	24	-CH ₂ O-		F		H	H
25	26	bond		H		H	H
27	28	bond		F		H	H
29	30	CO		H		H	H
31	32	CO		F		H	H





























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
33	34	SO ₂		H		H	H
35	36	SO ₂		F		H	H
37	38	-CH ₂ O-		H		H	H
39	40	-CH ₂ O-		F		H	H
41	42	bond		H	t-Bu		H
43	44	bond		F	t-Bu		H
45	46	CO		H	t-Bu		H
47	48	CO		F	t-Bu		H
49	50	SO ₂		H	t-Bu		H
51	52	SO ₂		F	t-Bu		H
53	54	-CH ₂ O-		H	t-Bu		H
55	56	-CH ₂ O-		F	t-Bu		H
59	60	bond		F	H		H
61	62	CO		H	H		H
63	64	CO		F	H		H
65	66	SO ₂		H	H		H
67	68	SO ₂		F	H		H
69	70	-CH ₂ O-		H	H		H





























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
71	72	-CH ₂ O-		F	H		H
73	74	bond		H	F		H
75	76	bond		F	F		H
77	78	CO		H	F		H
79	80	CO		F	F		H
81	82	SO ₂		H	F		H
83	84	SO ₂		F	F		H
85	86	-CH ₂ O-		H	F		H
87	88	-CH ₂ O-		F	F		H
89	90	bond		H	F		H
91	92	bond		F	F		H
93	94	CO		H	F		H
95	96	CO		F	F		H
97	98	SO ₂		H	F		H
99	100	SO ₂		F	F		H
101	102	-CH ₂ O-		H	F		H
103	104	-CH ₂ O-		F	F		H






























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
105	106	bond		H	t-Bu		H
107	108	bond		F	t-Bu		H
109	110	CO		H	t-Bu		H
111	112	CO		F	t-Bu		H
113	114	SO ₂		H	t-Bu		H
115	116	SO ₂		F	t-Bu		H
117	118	-CH ₂ O-		H	t-Bu		H
119	120	-CH ₂ O-		F	t-Bu		H
121	122	bond		H	H	t-Bu	
123	124	bond		F	H	t-Bu	
125	126	CO		H	H	t-Bu	
127	128	CO		F	H	t-Bu	
129	130	SO ₂		H	H	t-Bu	
131	132	SO ₂		F	H	t-Bu	
133	134	-CH ₂ O-		H	H	t-Bu	
135	136	-CH ₂ O-		F	H	t-Bu	
137	138	bond		H	F	t-Bu	


































Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
139	140	bond		F	F	t-Bu	
141	142	CO		H	F	t-Bu	
143	144	CO		F	F	t-Bu	
145	146	SO ₂		H	F	t-Bu	
147	148	SO ₂		F	F	t-Bu	
149	150	-CH ₂ O-		H	F	t-Bu	
151	152	-CH ₂ O-		F	F	t-Bu	
153	154	bond		H		H	H
155	156	bond		F		H	H
157	158	CO		H		H	H
159	160	CO		F		H	H
161	162	SO ₂		H		H	H
163	164	SO ₂		F		H	H
165	166	-CH ₂ O-		H		H	H
167	168	-CH ₂ O-		F		H	H
169	170	bond		H		H	H
171	172	bond		F		H	H



































Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
173	174	CO		H		H	H
175	176	CO		F		H	H
177	178	SO ₂		H		H	H
179	180	SO ₂		F		H	H
181	182	-CH ₂ O-		H		H	H
183	184	-CH ₂ O-		F		H	H
185	186	bond		H	t-Bu		H
187	188	bond		F	t-Bu		H
189	190	CO		H	t-Bu		H
191	192	CO		F	t-Bu		H
193	194	SO ₂		H	t-Bu		H
195	196	SO ₂		F	t-Bu		H
197	198	-CH ₂ O-		H	t-Bu		H
199	200	-CH ₂ O-		F	t-Bu		H
203	204	bond		F	H		H
205	206	CO		H	H		H
207	208	CO		F	H		H
209	210	SO ₂		H	H		H































Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
211	212	SO ₂		F	H		H
213	214	-CH ₂ O-		H	H		H
215	216	-CH ₂ O-		F	H		H
217	218	bond		H	F		H
219	220	bond		F	F		H
221	222	CO		H	F		H
223	224	CO		F	F		H
225	226	SO ₂		H	F		H
227	228	SO ₂		F	F		H
229	230	-CH ₂ O-		H	F		H
231	232	-CH ₂ O-		F	F		H
233	234	bond		H	F		H
235	236	bond		F	F		H
237	238	CO		H	F		H
239	240	CO		F	F		H
241	242	SO ₂		H	F		H
243	244	SO ₂		F	F		H
































Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
245	246	-CH ₂ O-		H	F		H
247	248	-CH ₂ O-		F	F		H
249	250	bond		H	t-Bu		H
251	252	bond		F	t-Bu		H
253	254	CO		H	t-Bu		H
255	256	CO		F	t-Bu		H
257	258	SO ₂		H	t-Bu		H
259	260	SO ₂		F	t-Bu		H
261	262	-CH ₂ O-		H	t-Bu		H
263	264	-CH ₂ O-		F	t-Bu		H
265	266	bond		H	H	t-Bu	
267	268	bond		F	H	t-Bu	
269	270	CO		H	H	t-Bu	
271	272	CO		F	H	t-Bu	
273	274	SO ₂		H	H	t-Bu	
275	276	SO ₂		F	H	t-Bu	
277	278	-CH ₂ O-		H	H	t-Bu	
























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
279	280	-CH ₂ O-		F	H	t-Bu	
281	282	bond		H	F	t-Bu	
283	284	bond		F	F	t-Bu	
285	286	CO		H	F	t-Bu	
287	288	CO		F	F	t-Bu	
289	290	SO ₂		H	F	t-Bu	
291	292	SO ₂		F	F	t-Bu	
293	294	-CH ₂ O-		H	F	t-Bu	
295	296	-CH ₂ O-		F	F	t-Bu	
297	298	bond		H		H	H
299	300	bond		F		H	H
301	302	CO		H		H	H
303	304	CO		F		H	H
305	306	SO ₂		H		H	H
307	308	SO ₂		F		H	H
309	310	-CH ₂ O-		H		H	H
311	312	-CH ₂ O-		F		H	H



























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
313	314	bond		H		H	H
315	316	bond		F		H	H
317	318	CO		H		H	H
319	320	CO		F		H	H
321	322	SO ₂		H		H	H
323	324	SO ₂		F		H	H
325	326	-CH ₂ O-		H		H	H
327	328	-CH ₂ O-		F		H	H
329	330	bond		H	t-Bu		H
331	332	bond		F	t-Bu		H
333	334	CO		H	t-Bu		H
335	336	CO		F	t-Bu		H
337	338	SO ₂		H	t-Bu		H
339	340	SO ₂		F	t-Bu		H
341	342	-CH ₂ O-		H	t-Bu		H
343	344	-CH ₂ O-		F	t-Bu		H
347	348	bond		F	H		H
349	350	CO		H	H		H



























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
351	352	CO		F	H		H
353	354	SO ₂		H	H		H
355	356	SO ₂		F	H		H
357	358	-CH ₂ O-		H	H		H
359	360	-CH ₂ O-		F	H		H
361	362	bond		H	F		H
363	364	bond		F	F		H
365	366	CO		H	F		H
367	368	CO		F	F		H
369	370	SO ₂		H	F		H
371	372	SO ₂		F	F		H
373	374	-CH ₂ O-		H	F		H
375	376	-CH ₂ O-		F	F		H
377	378	bond		H	F		H
379	380	bond		F	F		H
381	382	CO		H	F		H
383	384	CO		F	F		H
385	386	SO ₂		H	F		H



























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
387	388	SO ₂		F	F		H
389	390	-CH ₂ O-		H	F		H
391	392	-CH ₂ O-		F	F		H
393	394	bond		H	t-Bu		H
395	396	bond		F	t-Bu		H
397	398	CO		H	t-Bu		H
399	400	CO		F	t-Bu		H
401	402	SO ₂		H	t-Bu		H
403	404	SO ₂		F	t-Bu		H
405	406	-CH ₂ O-		H	t-Bu		H
407	408	-CH ₂ O-		F	t-Bu		H
409	410	bond		H	H	t-Bu	
411	412	bond		F	H	t-Bu	
413	414	CO		H	H	t-Bu	
415	416	CO		F	H	t-Bu	
417	418	SO ₂		H	H	t-Bu	
419	420	SO ₂		F	H	t-Bu	



























Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
421	422	-CH ₂ O-		H	H	t-Bu	
423	424	-CH ₂ O-		F	H	t-Bu	
425	426	bond		H	F	t-Bu	
427	428	bond		F	F	t-Bu	
429	430	CO		H	F	t-Bu	
431	432	CO		F	F	t-Bu	
433	434	SO ₂		H	F	t-Bu	
435	436	SO ₂		F	F	t-Bu	
437	438	-CH ₂ O-		H	F	t-Bu	
439	440	-CH ₂ O-		F	F	t-Bu	
441	442	bond		H		H	H
443	444	bond		F		H	H
445	446	CO		H		H	H
447	448	CO		F		H	H
449	450	SO ₂		H		H	H
451	452	SO ₂		F		H	H





Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
453	454	-CH ₂ O-		H		H	H
455	456	-CH ₂ O-		F		H	H
457	458	bond		H		H	H
459	460	bond		F		H	H
461	462	CO		H		H	H
463	464	CO		F		H	H
465	466	SO ₂		H		H	H
467	468	SO ₂		F		H	H
469	470	-CH ₂ O-		H		H	H
471	472	-CH ₂ O-		F		H	H
473	474	bond		H	t-Bu		H
475	476	bond		F	t-Bu		H
477	478	CO		H	t-Bu		H

Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
479	480	CO		F	t-Bu		H
481	482	SO ₂		H	t-Bu		H
483	484	SO ₂		F	t-Bu		H
485	486	-CH ₂ O-		H	t-Bu		H
487	488	-CH ₂ O-		F	t-Bu		H
491	492	bond		F	H		H
493	494	CO		H	H		H
495	496	CO		F	H		H
497	498	SO ₂		H	H		H
499	500	SO ₂		F	H		H
501	502	-CH ₂ O-		H	H		H
503	504	-CH ₂ O-		F	H		H
505	506	bond		H	F		H

Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
507	508	bond		F	F		H
509	510	CO		H	F		H
511	512	CO		F	F		H
513	514	SO ₂		H	F		H
515	516	SO ₂		F	F		H
517	518	-CH ₂ O-		H	F		H
519	520	-CH ₂ O-		F	F		H
521	522	bond		H	F		H
523	524	bond		F	F		H
525	526	CO		H	F		H
527	528	CO		F	F		H
529	530	SO ₂		H	F		H
531	532	SO ₂		F	F		H


Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
533	534	-CH ₂ O-		H	F		H
535	536	-CH ₂ O-		F	F		H
537	538	bond		H	t-Bu		H
539	540	bond		F	t-Bu		H
541	542	CO		H	t-Bu		H
543	544	CO		F	t-Bu		H
545	546	SO ₂		H	t-Bu		H
547	548	SO ₂		F	t-Bu		H
549	550	-CH ₂ O-		H	t-Bu		H
551	552	-CH ₂ O-		F	t-Bu		H
553	554	bond		H	H	t-Bu	
555	556	bond		F	H	t-Bu	
557	558	CO		H	H	t-Bu	









Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
559	560	CO		F	H	t-Bu	
561	562	SO ₂		H	H	t-Bu	
563	564	SO ₂		F	H	t-Bu	
565	566	-CH ₂ O-		H	H	t-Bu	
567	568	-CH ₂ O-		F	H	t-Bu	
569	570	bond		H	F	t-Bu	
571	572	bond		F	F	t-Bu	
573	574	CO		H	F	t-Bu	
575	576	CO		F	F	t-Bu	
577	578	SO ₂		H	F	t-Bu	
579	580	SO ₂		F	F	t-Bu	
581	582	-CH ₂ O-		H	F	t-Bu	
583	584	-CH ₂ O-		F	F	t-Bu	

Cmpd No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
A	B						
585	586	bond		H	H	H	
587	588	bond		H	H		H

Additional compounds of formula I, which can be prepared essentially according to the methods and procedures set forth above, include the following.

5



Cmpd. No.		X	R ₁	R ₂	R ₃	R ₄	R ₅
57	58	bond		H	H		H
201	202	bond		H	H		H
345	346	bond		H	H		H
489	490	bond		H	H		H

10

BIOLOGY EXAMPLES

Example 1. Method for measuring PTP-1B activity

The test compounds are evaluated for their in vitro inhibitory activity against recombinant human PTP1B with

phosphotyrosyl dodecapeptide TRDI(P)YETD(P)Y(P)YRK. This corresponds to the 1142-1153 insulin receptor kinase regulatory domain, phosphorylated on the 1146, 1150 and 1151 tyrosine residues; IR-triphosphopeptide as a source of substrate.


5 Enzyme reaction progression is monitored via the release of inorganic phosphate as detected by the malachite green - ammonium molybdate method for the phosphopeptide.

Preferred compounds of the invention exhibit IC_{50} values of less than 10 μM ; more preferred compounds of the invention exhibit IC_{50} values of less than 1 μM . Particularly preferred
10 compounds exhibit IC_{50} values of less than 300 nM.

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which
15 it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the invention and that modifications may be made therein without departing from the spirit or scope of the invention as set forth in the claims. To particularly point out and distinctly
20 claim the subject matter regarded as invention, the following claims conclude this specification.

What is claimed is:

1. A compound of the formula:



5

or a pharmaceutically acceptable salt thereof, wherein

R_1 is H, C_1 - C_6 alkyl, phenyl(C_1 - C_6)alkyl, or C_2 - C_6 alkenyl;

L_2 is a bond or $-C(O)NR_{10}-$, $-N(R_{10})C(O)-$, $-(C_1-C_4)$ alkyl-

10 $N(R_{10})C(O)-$, $-C(O)N(R_{10})-(C_1-C_4)$ alkyl-, $-N(R_{10})C(O)-(C_1-C_4)$ alkyl-, $-(C_1-C_4)$ alkyl- $C(O)N(R_{10})-$, $-O-(C_1-C_6)$ alkyl-, $-CO-$, $-SO_2-$, or $-(C_1-C_6)$ alkyl- $O-$;

L_3 is absent, a bond, $-(C_1-C_4)$ alkyl- $O-$, $-O-(C_1-C_4)$ alkyl, $-(C_1-C_4)$ alkyl-, $-alkenyl-$, $-phenyl-$;

15 L_5 is a bond, $-O-(C_1-C_6)$ alkyl-, $-(C_1-C_6)$ alkyl- $O-$, $-C(O)N(R_9)-(C_1-C_4)$ alkyl-, $-N(R_9)C(O)-(C_1-C_4)$ alkyl-, $-(C_1-C_4)$ alkyl- $C(O)N(R_9)-(C_1-C_4)$ alkyl-, $-(C_1-C_4)$ alkyl- $N(R_9)C(O)-(C_1-C_4)$ alkyl-, $-N(R_9)-(C_1-C_6)$ alkyl-, $-N(R_9)-(C_1-C_6)$ alkyl- wherein $-(C_1-C_6)$ alkyl- is optionally substituted with phenyl, $-(C_1-C_4)$ alkyl- $N(R_9)-(C_1-C_4)$ alkyl-, $-SO_2N(R_9)-$, $-SO_2N(R_9)-(C_1-C_4)$ alkyl-, $-N(R_9)SO_2-(C_1-C_4)$ alkyl-, $-N(R_9)SO_2-$, $-(C_1-C_4)$ alkyl-, $-(C_2-C_6)$ alkenyl-, $-N(R_9)C(O)-$, $-C(O)-(C_1-C_4)$ alkyl-, $-S-(C_1-C_4)$ alkyl-, $-(C_1-C_6)$ alkyl- $S-$, or $-(C_1-C_4)$ alkyl- $S-(C_1-C_4)$ alkyl-, wherein each alkyl and alkenyl is optionally substituted with one or more R_{11} groups,

25 R_9 and R_{10} are independently H, C_1 - C_6 alkyl, C_1 - C_6 alkoxycarbonyl, $-SO_2$ -aryl, heteroarylalkyl, arylalkyl, wherein the aryl or heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C_1 - C_4 alkyl, C_1 - C_4 alkoxy,

halogen, OH, NO₂, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl, or haloalkoxy;

R₁₁ at each occurrence is independently N₁₂R₁₃, -

N(R₁₂)C(O)R₁₃, N(R₁₂)CO₂R₁₃, or -C(O)NR₁₂R₁₃, wherein R₁₂

and R₁₃ are independently H or C₁-C₆ alkyl,

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H,

arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO₂,

NH₂, CN, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-aryl,

NHC(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-

aryl, N(C₁-C₄)alkyl-aryl, -NHSO₂-aryl, -N(C₁-

C₄alkyl)SO₂aryl, wherein the aryl group is optionally

substituted with 1, 2, 3, 4, or 5 groups that are

independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂,

haloalkyl, haloalkoxy;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which

is optionally substituted with 1, 2, or 3 groups that are

independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄

haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or

N(C₁-C₆)alkyl(C₁-C₆)alkyl;

Q is H, aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-

aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl,

-heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl,

wherein the aforementioned cyclic groups are optionally

substituted with 1, 2, 3, 4, or 5 groups that are

independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy,

halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-

C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-

C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl,

aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl,

heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -

C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the


cyclic groups are optionally substituted with 1, 2, 3, or

4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy;

and

- 5 Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂, or
- 10 Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

2. A compound according to claim 1, of the formula:



15

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

20 R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;

25

R₂₀ and R₂₁ are independently hydrogen or halogen;

L₅ is -(C₁-C₄) alkyl-N(R₉)-(C₁-C₄) alkyl-, -(C₂-C₆) alkenyl-, or -(C₁-C₄) alkyl-S-(C₁-C₄) alkyl-, wherein each alkyl and

alkenyl is optionally substituted with one or two R_{11} groups,

R_9 is H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, carbonyl;


R_{11} at each occurrence is independently $N_{12}R_{13}$, -

5 $N(R_{12})C(O)R_{13}$, $N(R_{12})CO_2R_{13}$, or $-C(O)NR_{12}R_{13}$, wherein R_{12} and R_{13} are independently H or C_1 - C_6 alkyl,

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and

10 n is 0, 1, or 2.

3. A compound according to claim 2, of the formula:



R_{27} is C_1 - C_6 alkoxy;

15 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

20 R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1 - C_6 alkyl;


25 R_{20} and R_{21} are independently hydrogen or halogen;

R_{30} and R_{31} are independently hydrogen or C_1 - C_2 alkyl;

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and n is 0, 1, or 2.

5

4. A compound according to claim 2, of the formula:



R_{27} is C_1 - C_6 alkoxy;

10 R_{28} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

15 R_{29} is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C_1 - C_4 alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C_1 - C_3 alkoxy or C_1 - C_2 acyl, or with one phenoxy;

provided that at least one of R_{28} and R_{29} is hydrogen or C_1 - C_6 alkyl;


R_{20} and R_{21} are independently hydrogen or halogen;

20 R_{30} and R_{31} are independently hydrogen or C_1 - C_2 alkyl;

each R_{10} is independently H, halogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, NO_2 , NH_2 , $NH(C_1-C_6)alkyl$, or $N(C_1-C_6)alkyl(C_1-C_6)alkyl$; and n is 0, 1, or 2.

25

5. A compound according to claim 2, of the formula:



and pharmaceutically acceptable salts thereof, wherein

R₂₇ is C₁-C₆ alkoxy;

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)

5 dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,
or (vii) phenyl substituted with one or two C₁-C₃ alkoxy
or C₁-C₂ acyl, or with one phenoxy;

R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv)

10 dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,
or (vii) phenyl substituted with one or two C₁-C₃ alkoxy
or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆
alkyl;

R₂₀ and R₂₁ are independently hydrogen or halogen;


15 R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy,
C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl,
or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

n is 0, 1, or 2.

20

6. A compound according to claim 2, of the formula:



R₂₇ is C₁-C₆ alkoxy;

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

5 R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;


R₂₀ and R₂₁ are independently hydrogen or halogen;

R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and

n is 0, 1, or 2.

7. A compound according to claim 2, of the formula:



20 p is 0 or 1;

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

25 R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl, or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;


provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;

R₂₀ and R₂₁ are independently hydrogen or halogen;

R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

5 each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and
n is 0, 1, or 2.

10 8. A compound according to claim 2, of the formula:



p is 0 or 1;

R₂₈ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,
15 or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

R₂₉ is (i) hydrogen, (ii) halogen, (iii) adamantanyl, (iv) dibenzofuranyl, (v) C₁-C₄ alkyl, (vi) dibenzothiophenyl,
20 or (vii) phenyl substituted with one or two C₁-C₃ alkoxy or C₁-C₂ acyl, or with one phenoxy;

provided that at least one of R₂₈ and R₂₉ is hydrogen or C₁-C₆ alkyl;

R₂₀ and R₂₁ are independently hydrogen or halogen;


R₃₀ and R₃₁ are independently hydrogen or C₁-C₂ alkyl;

25 each R₁₀ is independently H, halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl; and
n is 0, 1, or 2.

9. A compound according to claim 1, selected from the group consisting of:

4-tert-Butoxycarbonylamino-5-(4'-dibenzofuran-4-yl-biphenyl-4-yl)-pent-2-enoic acid;
 3-(4'-Dibenzofuran-4-yl-biphenyl-4-yl)-acrylic acid;
 3-(3'-Dibenzofuran-4-yl-biphenyl-4-ylmethylsulfanyl)-propionic acid; and
 (3'-Dibenzofuran-4-yl-biphenyl-3-ylmethylsulfanyl)-acetic acid.

10. A method for preparing a compound of formula I




or a pharmaceutically acceptable salt thereof, wherein
 L_5 is $-O-(C_1-C_6)\text{alkyl}-$, $-(C_1-C_6)\text{alkyl}-O-$, $-N(R_9)-(C_1-C_6)\text{alkyl}-$, $-N(R_9)-(C_1-C_6)\text{alkyl}-$ wherein $-(C_1-C_6)\text{alkyl}-$ is optionally substituted with phenyl, $-(C_1-C_4)\text{alkyl}-N(R_9)-(C_1-C_4)\text{alkyl}-$, $-S-(C_1-C_4)\text{alkyl}-$, $-(C_1-C_6)\text{alkyl}-S-$, or $-(C_1-C_4)\text{alkyl}-S-(C_1-C_4)\text{alkyl}-$, wherein each alkyl is optionally substituted with one or more R_{11} groups,
 R_9 is H, C_1-C_6 alkyl, C_1-C_6 alkoxycarbonyl, $-SO_2\text{-aryl}$, heteroarylalkyl, arylalkyl, wherein the aryl or heteroaryl groups are optionally substituted with 1, 2, 3, or 4 groups that are independently C_1-C_4 alkyl, C_1-C_4 alkoxy, halogen, OH, NO_2 , NH_2 , $NH(C_1-C_6)\text{alkyl}$, $N(C_1-C_6)\text{alkyl}(C_1-C_6)\text{alkyl}$, haloalkyl, or haloalkoxy;
 R_{11} at each occurrence is independently $N_{12}R_{13}$, $-N(R_{12})C(O)R_{13}$, $N(R_{12})CO_2R_{13}$, or $-C(O)NR_{12}R_{13}$, wherein R_{12} and R_{13} are independently H or C_1-C_6 alkyl; and

A, Z, L₂, L₃, Q, R₁, R₂₀, R₂₁, R₂₂, and R₂₃ are as defined in claim 1;

comprising:

treating a compound of formula




5

wherein R is (CH₂)_nOH, (CH₂)_nSH, or (CH₂)_nNH₂,

n is 0, 1, 2, 3, or 4,

with a base and a compound of formula



10


wherein X is Cl, Br, I, or OSO₂R',

R' is methyl, para-methylphenyl, or CF₃, and

m is 0, 1, 2, 3, or 4,

to provide a compound of formula (I).

15 11. A compound of formula (XXI)



(XXI),

wherein R is (CH₂)_nOH, (CH₂)_nSH, (CH₂)_nNH₂, (CH₂)_nCHO, or (CH₂)_n-X;

X is Cl, Br, I, or OSO₂R';

20 R' is methyl, para-methylphenyl, or CF₃;

n is 0, 1, 2, 3, or 4;

L₂ is a bond or -C(O)NR₁₀-, -N(R₁₀)C(O)-,

- (C₁-C₄)alkyl-N(R₁₀)C(O)-, -C(O)N(R₁₀)-(C₁-C₄)alkyl-,

-N(R₁₀)C(O)-(C₁-C₄)alkyl-, -(C₁-C₄)alkyl-C(O)N(R₁₀)-, -O-

25 (C₁-C₆)alkyl-, -CO-, -SO₂-, or -(C₁-C₆)alkyl-O-;

L₃ is absent, a bond, -(C₁-C₄)alkyl-O-, -O-(C₁-C₄)alkyl, -(C₁-C₄) alkyl-, -alkenyl-, or -phenyl-;

the A ring is aryl, heteroaryl, heterocycloalkyl, each of which is optionally substituted with 1, 2, or 3 groups that are independently, halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, NO₂, NH₂, NH(C₁-C₆)alkyl, or N(C₁-C₆)alkyl(C₁-C₆)alkyl;

R₂₀, R₂₁, R₂₂, and R₂₃ are independently selected from H, arylalkoxy, arylalkyl, halogen, alkyl, OH, alkoxy, NO₂, NH₂, CN, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, NH-aryl, NHC(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄ alkyl)C(O)-(C₁-C₄) alkyl-aryl, N(C₁-C₄)alkyl-aryl, -NHSO₂-aryl, or -N(C₁-C₄alkyl)SO₂aryl, wherein the aryl group is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, OH, NO₂, haloalkyl, or haloalkoxy;

Q is H, aryl, -aryl-carbonyl-aryl, -aryl-O-aryl, -aryl-alkyl-aryl, -aryl-heteroaryl, -aryl-heterocycloalkyl, -heteroaryl, -heteroaryl-alkyl-aryl, or -heterocycloalkyl, wherein the aforementioned cyclic groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxycarbonyl, aryl C₁-C₆ alkoxycarbonyl, heteroarylcarbonyl, heteroaryl, heterocycloalkylcarbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is absent, H, -NHC(O)aryl, -N(C₁-C₄ alkyl)C(O)aryl, or aryl (phenyl), wherein the aryl groups are optionally substituted with 1, 2, 3, 4, or 5 groups that are independently C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, or NO₂; or

Z is -NHC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl, or -N(C₁-C₄)alkylC(O)-(C₁-C₄)alkyl-(C₃-C₇)cycloalkyl.

12. A compound according to claim 11, wherein

10 L₂ is a bond;

L₃ is a bond,;

the A ring is phenyl;

R₂₀, R₂₁, R₂₂, and R₂₃ are H;

Q is -heteroaryl optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxy, carbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein

R₆ and R₇ are independently H, C₁-C₆ alkyl, aryl(C₁-C₆)alkyl, C₂-C₆ alkanoyl, aryl C₂-C₆ alkanoyl, C₁-C₆ alkoxy, carbonyl, aryl C₁-C₆ alkoxy, carbonyl, heteroaryl, heterocycloalkyl, carbonyl, -C(O)NH₂, -C(O)NH(C₁-C₆)alkyl, -C(O)N(C₁-C₆)alkyl(C₁-C₆)alkyl, or -SO₂-aryl, wherein the cyclic groups are optionally substituted with 1, 2, 3, or 4 groups that are independently halogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, NO₂, OH, NH₂, NH(C₁-C₆)alkyl, N(C₁-C₆)alkyl(C₁-C₆)alkyl, haloalkyl or haloalkoxy; and

Z is H.

13. In A compound according to claim 11

30 wherein

L₂ is a bond;

L₃ is a bond,;

the A ring is phenyl;

R₂₀, R₂₁, R₂₂, and R₂₃ are H;

Q is dibenzofuranyl, benzofuranyl, or indolyl, wherein each is optionally substituted with 1, 2, 3, 4, or 5 groups that are independently alkoxycarbonyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, halogen, haloalkyl, haloalkoxy, NR₆R₇, phenyl, phenyl-(C₁-C₆)alkyl-, or phenyloxy-; wherein R₆ and R₇ are independently H or C₁-C₆ alkyl; and Z is H.

10 14. A method of treating syndrome X, obesity, diabetes, immunological disease, bleeding disorders or cancer comprising administering a pharmaceutically acceptable amount of a compound of claim 1 to a patient in need of such treatment.

15 15. A pharmaceutical composition comprising a compound of claim 1 and at least one pharmaceutically acceptable solvent, carrier, adjuvant or excipient.

20 16. Use of a compound or salt of formula I for the manufacture of a medicament for treating diabetes in a patient in need of such treatment.


25 17. Use of a compound or salt of formula I for the manufacture of a medicament for treating syndrome X, obesity, immunological disease, bleeding disorders or cancer in a patient in need of such treatment.

30 18. Use of a compound or a salt of formula I for the manufacture of a medicament for inhibiting PTP-1B in a patient in need thereof.


19. Use of a pharmaceutical composition for the manufacture of a medicament wherein the pharmaceutical composition comprises a

compound of formula I and at least one pharmaceutically acceptable solvent, carrier, adjuvant or excipient.

20. A compound according to claim 1 which is 2-(3'-
5 Dibenzofuran-4-yl-biphen-3-ylmethylsulfanyl)-propionic acid.
21. A compound according to claim 1 which is 4-tert-
Butoxycarbonylamino-5-(4'-dibenzofuran-4-yl-biphenyl-4-yl)-
pent-2-enoic acid.
- 10 22. A compound according to claim 1 which is 3-(4'-
Dibenzofuran-4-yl-biphenyl-4-yl)-acrylic acid.
23. A compound according to claim 1 which is 3-(3'-
Dibenzofuran-4-yl-biphenyl-4-ylmethylsulfanyl)-propionic acid.
24. A compound according to claim 1 which is (3'-Dibenzofuran-
15 4-yl-biphenyl-3-ylmethylsulfanyl)-acetic acid.
25. A compound according to claim 1 which has Formula A or B:






































A






























B

where the substituents R_1 , R_2 , R_3 , R_4 , R_5 , and X are defined in
20 the table below




































X	R_1	R_2	R_3	R_4	R_5
bond		H		H	H
bond		F		H	H
CO		H		H	H




































X	R ₁	R ₂	R ₃	R ₄	R ₅
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H



































X	R ₁	R ₂	R ₃	R ₄	R ₅
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		F	H		H
CO		H	H		H
CO		F	H		H
SO ₂		H	H		H
SO ₂		F	H		H
-CH ₂ O-		H	H		H
-CH ₂ O-		F	H		H
bond		H	F		H
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	F		H


































X	R ₁	R ₂	R ₃	R ₄	R ₅
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		H	H	t-Bu	
bond		F	H	t-Bu	

X	R ₁	R ₂	R ₃	R ₄	R ₅
CO		H	H	t-Bu	
CO		F	H	t-Bu	
SO ₂		H	H	t-Bu	
SO ₂		F	H	t-Bu	
-CH ₂ O-		H	H	t-Bu	
-CH ₂ O-		F	H	t-Bu	
bond		H	F	t-Bu	
bond		F	F	t-Bu	
CO		H	F	t-Bu	
CO		F	F	t-Bu	
SO ₂		H	F	t-Bu	
SO ₂		F	F	t-Bu	
-CH ₂ O-		H	F	t-Bu	
-CH ₂ O-		F	F	t-Bu	
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H




































X	R ₁	R ₂	R ₃	R ₄	R ₅
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H





























X	R ₁	R ₂	R ₃	R ₄	R ₅
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		F	H		H
CO		H	H		H
CO		F	H		H
SO ₂		H	H		H
SO ₂		F	H		H
-CH ₂ O-		H	H		H
-CH ₂ O-		F	H		H
bond		H	F		H
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	F		H

X	R ₁	R ₂	R ₃	R ₄	R ₅
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		H	H	t-Bu	
bond		F	H	t-Bu	

























X	R ₁	R ₂	R ₃	R ₄	R ₅
CO		H	H	t-Bu	
CO		F	H	t-Bu	
SO ₂		H	H	t-Bu	
SO ₂		F	H	t-Bu	
-CH ₂ O-		H	H	t-Bu	
-CH ₂ O-		F	H	t-Bu	
bond		H	F	t-Bu	
bond		F	F	t-Bu	
CO		H	F	t-Bu	
CO		F	F	t-Bu	
SO ₂		H	F	t-Bu	
SO ₂		F	F	t-Bu	
-CH ₂ O-		H	F	t-Bu	
-CH ₂ O-		F	F	t-Bu	
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H





























X	R ₁	R ₂	R ₃	R ₄	R ₅
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H





























X	R ₁	R ₂	R ₃	R ₄	R ₅
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		F	H		H
CO		H	H		H
CO		F	H		H
SO ₂		H	H		H
SO ₂		F	H		H
-CH ₂ O-		H	H		H
-CH ₂ O-		F	H		H
bond		H	F		H
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	F		H



























X	R ₁	R ₂	R ₃	R ₄	R ₅
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		H	H	t-Bu	
bond		F	H	t-Bu	



























X	R ₁	R ₂	R ₃	R ₄	R ₅
CO		H	H	t-Bu	
CO		F	H	t-Bu	
SO ₂		H	H	t-Bu	
SO ₂		F	H	t-Bu	
-CH ₂ O-		H	H	t-Bu	
-CH ₂ O-		F	H	t-Bu	
bond		H	F	t-Bu	
bond		F	F	t-Bu	
CO		H	F	t-Bu	
CO		F	F	t-Bu	
SO ₂		H	F	t-Bu	
SO ₂		F	F	t-Bu	
-CH ₂ O-		H	F	t-Bu	
-CH ₂ O-		F	F	t-Bu	
bond		H		H	H
bond		F		H	H
CO		H		H	H

X	R ₁	R ₂	R ₃	R ₄	R ₅
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H		H	H
bond		F		H	H
CO		H		H	H
CO		F		H	H
SO ₂		H		H	H
SO ₂		F		H	H
-CH ₂ O-		H		H	H
-CH ₂ O-		F		H	H
bond		H	t-Bu		H



X	R ₁	R ₂	R ₃	R ₄	R ₅
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		F	H		H
CO		H	H		H
CO		F	H		H
SO ₂		H	H		H
SO ₂		F	H		H
-CH ₂ O-		H	H		H
-CH ₂ O-		F	H		H

X	R ₁	R ₂	R ₃	R ₄	R ₅
bond		H	F		H
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	F		H
bond		F	F		H
CO		H	F		H
CO		F	F		H
SO ₂		H	F		H
SO ₂		F	F		H

X	R ₁	R ₂	R ₃	R ₄	R ₅
-CH ₂ O-		H	F		H
-CH ₂ O-		F	F		H
bond		H	t-Bu		H
bond		F	t-Bu		H
CO		H	t-Bu		H
CO		F	t-Bu		H
SO ₂		H	t-Bu		H
SO ₂		F	t-Bu		H
-CH ₂ O-		H	t-Bu		H
-CH ₂ O-		F	t-Bu		H
bond		H	H	t-Bu	
bond		F	H	t-Bu	
CO		H	H	t-Bu	
CO		F	H	t-Bu	

X	R ₁	R ₂	R ₃	R ₄	R ₅
SO ₂		H	H	t-Bu	
SO ₂		F	H	t-Bu	
-CH ₂ O-		H	H	t-Bu	
-CH ₂ O-		F	H	t-Bu	
bond		H	F	t-Bu	
bond		F	F	t-Bu	
CO		H	F	t-Bu	
CO		F	F	t-Bu	
SO ₂		H	F	t-Bu	
SO ₂		F	F	t-Bu	
-CH ₂ O-		H	F	t-Bu	
-CH ₂ O-		F	F	t-Bu	
bond		H	H	H	

or

bond		H	H		H.
------	---	---	---	---	----

