



(12) UK Patent (19) GB (11) 2 250 743 (13) B

(54) Title of Invention

Heterocyclic sulphonamide leukotriene antagonist

(51) INT CL^s; C07D 209/18, A61K 31/40

(21) Application No
9126353.3

(22) Date of filing
11.12.1991

(30) Priority Data

(31) **9027018**

(32) **12.12.1990**

(33) **GB**

(43) Application published
17.06.1992

(45) Patent published
15.02.1995

(72) Inventor(s)
Martin Paul Edwards
John David Sherwood

(73) Proprietor(s)
Zeneca Limited

(Incorporated in the United Kingdom)

Imperial Chemical House
9 Millbank
London
SW1P 3JF
United Kingdom

(74) Agent and/or
Address for Service
Paul Millington Denerley
Zeneca Pharmaceuticals
Intellectual Property Group
Intellectual Property
Department
Mereside
Alderley Park
Macclesfield
SK10 4TG
United Kingdom

(52) Domestic classification
(Edition N)
C2C CSC CVM CVW C1175
C1343 C213 C22Y C220 C225
C25Y C250 C251 C30Y C303
C34Y C340 C36Y C364 C385
C510 C52Y C662 C699 C80Y
C802
U1S S1321 S2415 S2416

(56) Documents cited
EP0199543 A2

(58) Field of search

As for published application
2250743 A *viz*:
UK CL(Edition K) C2C CSC
INT CL^s C07D
Online databases : CAS
ONLINE
updated as appropriate







Fig. 2.





4/5



5/5

Fig. 5.





Fig. 6.



HETEROCYCLIC SULPHONAMIDE LEUKOTRIENE ANTAGONIST

The present invention relates to heterocyclic compounds. More particularly, it relates to a new physical form of a heterocyclic amide derivative, to a process for the preparation of this physical form, and to a pharmaceutical composition containing it.

Before a compound in the solid state can be formulated in a pharmaceutical composition, a physical form of the compound is sought which is physically stable and can be prepared substantially free of other physical forms. This latter requirement is important because different physical forms can have markedly different bioavailabilities.

European patent application publication number EP-A2-0199543 discloses certain heterocyclic amide derivatives which antagonise the pharmacological actions of one or more of the arachidonic acid metabolites known as leukotrienes. One of these heterocyclic amide derivatives is N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide, hereinafter referred to as compound 1. Example 105 in EP-A2-0199543 discloses the preparation of compound 1 and its isolation in the solid state. The material obtained had a melting point of 138-140°C.

It has been found that compound 1 can be isolated in the solid state as a material having a range of different physical properties, depending upon the method of isolation. This ability is due to the fact that compound 1 can exist in more than one physical form, and mixture of these forms can be isolated.

A method of preparing a physical form of compound 1 substantially free of other physical forms has now been found, and this form substantially free of other physical forms is provided as the present invention.

Accordingly, the present invention provides a new physical form of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide substantially free of other physical forms, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at $2\theta=8.1$, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} .

The new physical form of compound 1, which is referred to hereinafter as form X, is physically stable and can be prepared substantially free of other physical forms.

Where reference is made in this specification to form X substantially free of other physical forms, it preferably means that at least 90% by weight of the compound 1 present is in that physical form.

The advantageous nature of form X may be demonstrated by comparing its properties with those of other physical forms of compound 1.

One other physical form of compound 1 is a monohydrate of compound 1 which is crystalline, has an infra-red spectrum (0.5% in KBr) having sharp peaks at 3560, 1690, 1660, 1540, 1440, 1165, 880 and 858 cm^{-1} , and an X-ray powder diffraction pattern having peaks at $2\theta = 10.0$, 11.2, 14.6, 19.8 and 23.0°. It is referred to hereinafter as form B.

Form B may be prepared substantially free of other physical forms by crystallisation from hot aqueous acetone. In particular, it may be prepared by dissolving a source of compound 1 in aqueous acetone at an elevated temperature, adding more water, and allowing the resultant mixture to cool. The crystalline product may be dried at an elevated temperature, for example at about 60 °C or below. If it is desired to start from an impure source of compound 1, it has

been found advantageous to triturate this impure source with hot toluene/ethyl acetate prior to the crystallisation.

Form B has been found to be more difficult to prepare substantially free of other physical forms than form X. In particular, it has been found that after form B has been crystallised and filtered off, it is difficult to remove the residual organic solvent without losing some of the water of crystallisation.

Another physical form of compound 1 is amorphous, has an infra-red spectrum (0.5% in KBr) having sharp peaks at 1690, 1530, 1490, 1420, 1155, 1060, 862 and 550 cm^{-1} . This amorphous form of compound 1 is referred to hereinafter as form A. Because form A is amorphous, it is also characterised by an X-ray powder diffraction pattern having no discernable peaks.

Form A may be prepared substantially free of other physical forms by dehydrating form B, prepared as described above, at a temperature of about 120 °C under vacuum.

Form A has been found to be physically unstable in the presence of chlorofluorocarbon aerosol propellants, and is therefore unsuitable for use in an aerosol formulation. In particular, when suspended in a chlorofluorocarbon aerosol propellant, it has been found to change to a new, crystalline form, the crystals of which have been found to grow beyond a size suitable for use in a metered dose inhaler.

Form X has been found to be stable in the presence of chlorofluorocarbon aerosol propellants.

Each of the forms X, A, and B may conveniently be characterised for example by their X-ray powder diffraction pattern alone, or their infra-red pattern alone.

In this specification, infra-red spectra were determined using a 0.5% dispersion of sample material in a potassium bromide disc over the wave number range 4000 to 400cm⁻¹. Examples of infra-red spectra for each of forms X, A and B are given in Figures 1, 2 and 3 hereinafter.

X-ray powder diffraction spectra were determined using 2g of sample material mounted in a Philips standard deep pack holder ('Philips' a Trade or Service Mark) over the scanning range of 4-40° 20 counting for 4 seconds per point at 0.02° intervals to produce a trace of spacings against intensity for this range. Examples of X-ray powder diffraction spectra for each of forms X, A and B are given in Figures 4, 5 and 6 hereinafter.

The melting point of each of the forms X, A and B generally depend upon their level of purity. Typically, form X has been found to have a melting point of above 190°C, for example about 200°C; form A between 115°C and 140°C, for example about 124 to 132°C; and form B about 140 to 160°C, for example from 145 to 155°C. Form B has been observed to lose water at a temperature of from about 60°C, and may not show a sharp melting point.

According to another aspect, the invention provides a process for the preparation of form X substantially free of other physical forms, which comprises dissolving a source of compound 1 in hot aqueous acetone, reducing the volume of the resultant solution by evaporation, adding toluene and further reducing the volume by evaporation.

If it is desired to employ material which is a relatively impure source of compound 1, such material may advantageously be triturated with hot toluene/ethyl acetate prior to the crystallisation step.

As stated previously, compound 1 possesses leukotriene antagonist properties. Thus it antagonises the actions of one or more of the arachidonic acid metabolites known as leukotriens, for

example, C₄, D₄ and/or E₄, which are known to be powerful spasmogens (particularly in the lung), to increase vascular permeability and have been implicated in the pathogenesis of asthma and inflammation (see J. L. Marx, Science, 1982, 215, 1380-1383) as well as of endotoxic shock (see J. A. Cook, et al., J. Pharmacol. Exp. Ther., 1985, 235, 470) and traumatic shock (see C. Denzlinger, et al., Science, 1985, 230, 330). Compound 1 is thus useful in the treatment of diseases in which leukotrienes are implicated and in which antagonism of their action is desired. Such diseases include, for example, allergic pulmonary disorders such as asthma, hay fever and allergic rhinitis and certain inflammatory diseases such as bronchitis, ectopic and atopic eczema, psoriasis, as well as vasospastic cardiovascular disease, and endotoxic and traumatic shock conditions.

Form X may be administered by itself, for example by inhalation as a micronised powder, or in a pharmaceutical composition.

According to another aspect, the invention provides a pharmaceutical composition, which comprises form X substantially free of other physical forms and a pharmaceutically acceptable carrier.

The pharmaceutical composition may be formulated in a conventional manner, and may typically be in the form of tablets, capsules or suspensions for oral administration; in the form of suppositories for rectal administration; in the form of suspensions for parenteral administration; in the form of suspensions for inhalation administration by metered dose inhaler or nebuliser; and in the form of powders together with pharmaceutically acceptable inert solid diluents such as lactose for administration by inhalation.

As stated hereinbefore, form X is especially suitable for use in a metered dose inhaler. According to a preferred aspect, therefore, the invention provides a pharmaceutical composition suitable for administration by a metered dose inhaler, which comprises form X substantially free of other physical forms and a pharmaceutically acceptable propellant.

The non toxic pharmaceutically acceptable propellant may be any of those conventionally used in metered dose inhalers. Examples of propellants include chlorofluorocarbons, for example trichlorofluoromethane, dichlorodifluoromethane and dichlorotetrafluoroethane; and 1,1,1,2-tetrafluoroethane.

The pharmaceutical composition suitable for administration by a metered dose inhaler may further comprise one or more other ingredients conventionally used in metered dose inhalers, for example oleic acid, sorbitan trioleate or lecithin.

The mean mass aerodynamic diameter of form X used in the aerosol composition is conveniently less than 50 microns, preferably in the range of from 1 to 50 microns, more preferably 1 to 10 microns, especially 1 to 5 microns.

The amount of form X present in the aerosol composition is conveniently from 0.01 to 10% by weight, preferably from 0.03 to 2%, more preferably from 0.05 to 1%.

The amount of form X administered to a patient will depend upon the weight of the patient and the severity of the disorder being treated, and the route of administration. Conveniently, a unit aerosol dose will comprise from 0.01 to 2.0 mg of form X, preferably from 0.02 to 1.0 mg, more preferably from 0.05 to 0.6 mg. Administration may take place from 1 to 8 times per day, preferably from 1 to 4 times per day. A typical daily dose for a 70 kg patient will be from 0.01 to 16 mg, preferably 0.02 to 4 mg.

It has been found that micronised form X has very poor flow characteristics, and tends to adhere to many different types of surface. These properties are disadvantageous, because a powder with poor flow characteristics is difficult to handle on a manufacturing scale. Furthermore, such a powder is unsuitable for use in a dry powder inhaler, especially one that utilises volumetric metering.

A flowable state of form X has now been found, and is provided as a further feature of the invention.

Accordingly, the invention provides form X in the form of soft pellets consisting of an agglomeration of particles, at least 90% of which have a diameter of less than 10 microns, which pellets have a diameter in the range of from 50 to 900 microns.

The diameter of the pellets may be, for example in the range of from 75 to 250 microns, such as from 100 to 200 microns, or from 200 to 500 microns, such as from 250 to 400 microns, depending upon their intended use.

It has been found that form X in the form of soft pellets as defined above has good flow characteristics. Furthermore, it has been found that the pellets readily break down to the original particles when exposed to shear forces. These properties are particularly desirable in a powder intended for use in a dry powder inhaler, especially one that utilises volumetric metering. Thus, in a dry powder inhaler which utilises volumetric metering, it is important for powder to be able to flow freely from a powder reservoir into a dosing hole. It is also important, in any dry powder inhaler, that any agglomerates of particles are broken up before leaving the device, otherwise they will be too large to be able to penetrate into the lungs.

According to another aspect, the invention provides a process for obtaining form X in the form of soft pellets, which comprises extruding micronised form X through a sieve having apertures with a diameter in the range of from 150 to 700 microns, rolling the extruded material, and then screening the rolled material.

The micronised form X is preferably extruded through a sieve having apertures with a diameter in the range of from 175 to 600 microns. The extrusion may conveniently be performed by passing the surface of a blade across powder on the surface of the sieve.

The function of the rolling step in the process is to strengthen the agglomerates of particles formed in the extrusion step, and to mould them into a spherical shape. The rolling may conveniently be effected by allowing the agglomerates to tumble in a rotating vessel, preferably a cylindrical vessel.

The screening step removes over and undersized pellets. It is conveniently performed using two sieves with apertures defining the upper and lower desired pellet diameters.

The following Examples illustrate the invention.

Example 1

Preparation of Form X

a) Preparation of an impure source of compound 1

Methyl 3-methoxy-4-(1-methyl-5-nitroindol-3-ylmethyl)-benzoate (prepared as described in Example 4 of EP-A2-0199543) was converted into the free acid by treatment with aqueous sodium hydroxide. The free acid was then converted into the acid chloride by treatment with thionyl chloride in dichloromethane. The acid chloride was then reacted o-toluenesulphonamide in dichloromethane in the presence of 2.2 equivalents of 4-dimethylaminopyridine to afford the dimethylaminopyridine salt of 4-(1-methyl-5-nitroindol-3-ylmethyl)-3-methoxybenzoyl-2-methylbenzenesulphonamide.

A solution of the dimethylaminopyridine salt of 4-(1-methyl-5-nitroindol-3-ylmethyl)-3-methoxybenzoyl-2-methylbenzenesulphonamide (30 g) in 2-methoxyethanol (130 ml) and concentrated sodium hydroxide liquor (3.2 ml) was charged to a nitrogen-purged flask containing 10% palladium on charcoal (3.3 g of a 60.9% water-wet paste). The mixture was then stirred under a hydrogen atmosphere at a pressure of 3 bar for 2.5 hours. The mixture was then filtered through diatomaceous earth, and washed through with 2-methoxyethanol (37.5 ml). To the combined liquors was added cyclopentyl chloroformate (9.2 ml), and the mixture allowed to stir under an

atmosphere of nitrogen overnight. The temperature was then adjusted to 30-33°C, and 0.8M hydrochloric acid (68 ml) was added over 20 minutes with vigorous agitation. The mixture was then cooled to 15-20 °C and stirred for one hour. The crystalline product was then filtered off, washed with water and dried at 50 °C. It was then used in the next step.

b) Trituration of impure compound 1

60 g (0.101 gmol) of the product of step a), toluene 240 ml (4 volumes) and ethyl acetate 150 ml (2.5 volumes) were slowly heated to reflux and 30 ml (0.5 volumes) of distillate were collected to remove most of the released water. The mixture was heated under reflux for one hour (88-90 °C) and was then cooled to 10-15 °C. After stirring for three hours at 10-15 °C, the solid was filtered through a glass sinter and washed with a 2:1 mixture of toluene (80 ml) and ethyl acetate (40 ml). The product was then dried to constant weight on the sinter to afford 53.2 g of dry compound 1 (yield 91.5%).

c) Preparation of form X

The product of step b) (30.0 g, 0.0521 g mol) was dissolved in acetone (150 ml) and water (4.7 ml) by gentle heating to reflux, and the solution screened through a sintered glass funnel. The filtrate was heated to boiling and 90 ml distillate collected. Toluene (120 ml) was added and a further 75 ml distillate collected. More toluene (120 ml) was added and an additional 75 ml distillate collected. After heating for a further hour at reflux, the mixture was cooled to 15-20 °C, and the product collected and washed with toluene (2x30 ml). The yield after drying on the sinter funnel was 29.5 g (98.3%).

Preparation of Form B

30.0 g of the product of step b), 210 ml acetone and 12 ml water were charged to a 500 ml reaction flask. The mixture was then heated under reflux for 15 minutes, and was then screened at 45-50 °C through a diatomaceous earth pad on a glass sinter directly into a 500 ml reaction flask. The flask and sinter were washed with a mixture of acetone (60 ml) and water (3 ml). The combined liquids were then stirred in a water bath at approximately 40 °C and water (120 ml) was added over five minutes. The mixture oiled out at first, but then rapidly crystallised. The mixture was then cooled to 20 °C over one hour, stirred for two hours at 15-20 °C and then filtered. The product was washed with water (60 ml), dried as far as possible on the sinter and then dried in an air oven at 60 °C (max.). The yield of form B was 30.0 g (97%).

It has been found that the drying conditions for form B must be strictly controlled in order to prevent loss of water.

Preparation of Form A

Form B prepared as described above (15.0 g) was placed in a 500 ml round bottomed flask which was then evacuated on a rotary evaporator at 20 mbara. The flask and contents were then immersed in an oil bath preheated to 118 °C, and slowly rotated at this temperature for 6 hours. The mass was broken up on cooling to afford form A as a white powder.

Example 2

Pharmaceutical Composition of Form X suitable for administration by a metered dose inhaler

Active ingredient	0.02g
Oleic acid	0.01g
Trichloromonofluoromethane (chlorofluorocarbon 11)	4.97g
Dichlorodifluoromethane (chlorofluorocarbon 12)	10.5g
Dichlorotetrafluoroethane (chlorofluorocarbon 114)	4.5g

Oleic acid, and then micronised active ingredient were added to a tank containing the trichloromonofluoromethane. The mixture was then homogenised and transferred into an aerosol canister. A valve was then crimped onto the aerosol canister, and then the remaining chlorofluorocarbons were introduced into the canister through the valve.

Comparison of the Stability of Form X in an Aerosol Formulation with that of Form A

Compositions comprising form X or form A were prepared as described in Example 2 and stored at various temperatures and humidity levels for 1, 3 and 6 months. Samples of the materials were then sprayed onto a microscope slide and examined under a microscope. The results are summarised in Tables 1 and 2.

Samples from each stability test were also studied by cascade impaction. The results are summarised in Tables 3 and 4.

The results of these comparative tests demonstrate the superiority of form X over form A in an aerosol formulation.

Example 3

Pharmaceutical Composition of Form X suitable for administration by a metered dose inhaler.

Active ingredient	0.02g
Sorbitan trioleate	0.04g
Trichloromonofluoromethane (chlorofluorocarbon 11)	7.0g
Dichlorodifluoromethane (chlorofluorocarbon 12)	13.0g

The composition was prepared by a method analogous to that described in Example 2.

Table 1

Microscopic Observations of Metered Dose Inhaler Formulation of Form X

Storage conditions	Particle Characteristics
Initial	Highly crystalline irregularly shaped particles which are <5 μ m; some agglomerates (up to 12.5 μ m) of these particles
5 °C	
1 month	NC
3 months	NC
6 months	NC
RT	
1 month	NC
3 months	NC
6 months	NC
30 °C/80%RH	
1 month	NC
3 months	NC
6 months	NC
40 °C	
1 month	NC
3 months	NC
6 months	NC

NC = no change

RH = relative humidity

RT = room temperature

Table 2

Microscopic Observations of Metered Dose Inhaler Formulation of Form A

Storage conditions	Particle Characteristics
Initial	needles
4 °C	
1 month	needles
3 months	needles
6 months	needles
RT	
1 month	needles
3 months	needles
6 months	needles
30 °C/80%RH	
1 month	needles
3 months	needles
6 months	needles, rods (many; up to 20x600µm)
40 °C	
1 month	needles, rods (only a few; up to 6x173µm)
3 months	needles, rods (more; up to 15x289µm)
6 months	needles, rods (many; up to 20x150µm)

RH = relative humidity

RT = room temperature

Table 3

Summary of Cascade Impaction Results for Metered Dose
Inhaler Product Containing Form X

Storage Condition	MMAD (μm)	GSD
Initial	3.94	1.67
5 °C		
1 month	3.60	1.95
3 months	3.60	1.64
6 months	3.16	1.53
RT		
1 month	3.64	1.78
3 months	3.53	1.66
6 months	3.67	1.62
30 °C/80%RH		
1 month	3.74	1.89
3 months	4.25	1.75
6 months	3.98	1.66
40 °C		
1 month	3.73	1.89
3 months	3.76	1.64
6 months	3.67	1.62

MMAD = mass median aerodynamic diameter

GSD = geometric standard deviation

RT = room temperature

RH = relative humidity

Table 4

Summary of Cascade Impaction Results for Metered Dose
Inhaler Product Containing Form A

Storage Condition	MMAD (μm)	GSD
Initial	2.00	1.82
4 °C		
1 month	1.45	1.95
3 months	1.32	1.74
6 months	1.70	1.81
RT		
1 month	1.70	1.87
3 months	1.53	1.77
6 months	1.80	1.85
30 °C/80%RH		
1 month	1.76	1.77
3 months	1.57	1.80
6 months	>>9.0	----
40 °C		
1 month	1.41	1.76
3 months	1.70	1.84
6 months	>>9.0	----

MMAD = mass median aerodynamic diameter

GSD = geometric standard deviation

RT = room temperature

RH = relative humidity

Example 4

Tablet Formulation of Form X

Granulating Material	mg/tablet
Active ingredient	20
Croscarmellose sodium	5
Polyvinylpyrrolidone	4
Microcrystalline cellulose	74
Lactose	50
Purified water for injection	98

Final Blend Material

Dried milled granulation	153
Lactose	20
Croscarmellose sodium	5
Microcrystalline cellulose	20
Magnesium stearate	2

The ingredients of the granulating material are wet granulated. The granules are then dried and milled. The dried milled granulation is then blended with the other ingredients of the final blend mixture and compressed into tablets.

Example 5

Treatment of micronised form X to improve its flow characteristics

Form X was micronised to produce a powder consisting of at least 98% by weight of particles having a diameter of less than 10 microns.

30g of the powder was then placed in one heap on a brass sieve having an aperture size between 210 and 500 microns. The powder was then

extruded through the apertures of the sieve using a stainless steel palette knife. The extrudate thus formed was then placed into a screw topped glass jar.

The glass jar was then placed on to a set of rollers, which were rotated at 100rpm for between 8 and 20 minutes. The soft pellets thus formed were then sieved through an 850 micron sieve and then a 150 micron sieve, the fraction retained on the 150 micron sieve being the required product.

The soft pellets thus produced were free flowing and relatively dust free. However the pellets when sheared, for example in a twin impinger, broke back down to the powder's original particle size distribution. This indicates that the pellets are suitable for use in a multidose dry powder inhaler that utilises volumetric metering to measure out the doses.

CLAIMS

1. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide substantially free of other physical forms, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at $2\theta=8.1$, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} .
2. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide when at least 90% by weight free of other physical forms, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at $2\theta=8.1$, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} .
3. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at $2\theta=8.1$, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} , and a melting point of above 190°C.
4. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at $2\theta=8.1$, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} , and having a melting point of about 200 C.

5. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methylindol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide having an infra-red spectrum (0.5% in KBr) substantially as shown in Figure 1 herein.
6. A physical form of N-[4-[5-cyclopentyloxycarbonyl)amino-1-methylindol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide having an X-ray powder diffraction pattern substantially as shown in figure 4 herein.
7. A process for the preparation of a physical form of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide as defined in any one of claims 1 to 6, which comprises dissolving a source of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide in hot aqueous acetone, reducing the volume of the resultant solution by evaporation, adding toluene and further reducing the volume by evaporation.
8. A process as claimed in claim 7 wherein the source of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is obtained by trituration of an impure source of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide with a hot mixture of toluene and ethyl acetate.

9. A pharmaceutical composition, which comprises a physical form of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide as claimed in any one of claims 1 to 6 and a pharmaceutically acceptable carrier.
10. A pharmaceutical composition as claimed in claim 9, which composition is suitable for administration by a metered dose inhaler, and comprises a pharmaceutically acceptable propellant.
11. A pharmaceutical composition as claimed in claim 10, in which the propellant comprises trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethane or 1,1,1,2-tetrafluoroethane.
12. A pharmaceutical composition as claimed in either claim 10 or claim 11 which further comprises one or more ingredients selected from oleic acid, sorbitan trioleate and lecithin.
13. A pharmaceutical composition as claimed in any one of claims 10 to 12 in which the mean mass aerodynamic diameter of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is in the range of from 1 to 50 microns.
14. A pharmaceutical composition as claimed in claim 13 in which the mean mass aerodynamic diameter of N-[4-[5-(cyclopentyl-oxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is in the range of from 1 to 10 microns.
15. A pharmaceutical composition as claimed in claim 14 in which the mean mass aerodynamic diameter of N-[4-[5-(cyclopentyloxy-carbonyl-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is in the range of from 1 to 5 microns.

16. A pharmaceutical composition as claimed in any one of claims 10 to 15 in which N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is present in the range 0.01 to 10% by weight.
17. A pharmaceutical composition as claimed in claim 16 in which N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is present in the range 0.03 to 2% by weight.
18. A pharmaceutical composition as claimed in claim 17 in which N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide is present in the range 0.05 to 1% by weight.
19. A unit aerosol dose of a pharmaceutical composition as claimed in any one of claims 10 to 18 which comprises 0.01 to 2.0mg of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide.
20. A unit aerosol dose of a pharmaceutical composition as claimed in claim 19 which comprises 0.02 to 1.0mg of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide.
21. A unit aerosol dose of a pharmaceutical composition as claimed in claim 20 which comprises 0.05 to 0.6mg of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide.
22. A pharmaceutical composition as claimed in any one of claims 10 to 21 in combination with a metered dose inhaler.

23. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide as claimed in any one of claims 1 to 6, which is in the form of soft pellets consisting of an agglomeration of particles, at least 90% of which have a diameter of less than 10 microns, which pellets have a diameter in the range of from 50 to 900 microns.

24. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide substantially free of other physical forms, which form is crystalline, has an X-ray powder diffraction pattern with specific peaks occurring at 20=8.1, 13.7, 16.4, 20.5 and 23.7° and an infra-red spectrum (0.5% in KBr) having sharp peaks at 3370, 1670, 1525, 1490, 1280, 890, 870 and 550 cm^{-1} , which is in the form of soft pellets consisting of an agglomeration of particles, at least 90% of which have a diameter of less than 10 microns, which pellets have a diameter in the range of from 50 to 900 microns.

25. A physical form as claimed in either claim 23 or claim 24 wherein the diameter of the pellets is in the range of from 75 to 250 microns.

26. A physical form as claimed in claim 25 wherein the diameter of the pellets is in the range of from 100 to 200 microns.

27. A physical form as claimed in either claim 23 or claim 24 wherein the diameter of the pellets is in the range of from 200 to 500 microns.

28. A physical form as claimed in claim 27 wherein the diameter of the pellets is in the range of from 250 to 400 microns.

29. A physical form as claimed in any one of claims 23 to 28 in combination with a dry powder inhaler.

30. A combination as claimed in claim 29 wherein the dry powder inhaler is one that utilises volumetric metering.

31. A process for obtaining material as claimed in any one of claims 23 to 28, which comprises extruding micronised particles of a physical form of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide as claimed in any one of claims 1 to 6 through a sieve having apertures with a diameter in the range of from 150 to 700 microns, rolling the extruded material, and then screening the rolled material.

32. A process for the preparation of N-[4-[5-(cyclopentyloxycarbonyl)amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide substantially as described in Example 1 (c) herein.

33. A pharmaceutical composition as claimed in claim 9 substantially as herein described with reference to any example thereof.

34. A physical form of N-[4-[5-(cyclopentyloxycarbonyl)-amino-1-methyl-indol-3-ylmethyl]-3-methoxybenzoyl]-2-methylbenzenesulphonamide substantially as herein described with reference to Example 1(c).

35. A physical form of a compound as claimed in any one of claims 1 to 6 when prepared by the process of claim 7.

36. A physical form as claimed in any one of claims 1 to 6 in micronised form consisting of at least 98% by weight of particles having a diameter of less than 10 microns.

37. The use of a physical form as claimed in any one of claims 1 to 6 for the manufacture of a medicament for treatment of asthma, hay fever or allergic rhinitis.

38. The use of a physical form as claimed in any one of claims 1 to 6 for the manufacture of a medicament for antagonising the action of leukotrienes.

FC36080AA

25JUL94

REGISTER ENTRY FOR GB2250743

Form 1 Application No GB9126353.3 filing date 11.12.1991

Priority claimed:

12.12.1990 in United Kingdom - doc: 9027018

Title HETEROCYCLIC COMPOUNDS

Applicant/Proprietor

IMPERIAL CHEMICAL INDUSTRIES PLC, Incorporated in the United Kingdom,
Imperial Chemical House, Millbank, LONDON, SW1P 3JF, United Kingdom

[ADP No. 00000935003]

Inventors

MARTIN PAUL EDWARDS, Mereside, Alderley Park, MACCLESFIELD, Cheshire, SK10
4TG, United Kingdom [ADP No. 06028781001]JOHN DAVID SHERWOOD, Hurdsfield Industrial Estate, Macclesfield Works,
Hurdsfield Road, MACCLESFIELD, Cheshire, United Kingdom

[ADP No. 06028799001]

Classified to

C2C U1S
C07D A61K

Address for Service

MARTIN ALEXANDER HAY, IMPERIAL CHEMICAL INDUSTRIES PLC, ICI Group Patents,
Group Patents Services Dept, PO Box 6, Shire Park, Bessemer Road, WELWYN
GARDEN CITY, Hertfordshire, AL7 1HD, United Kingdom [ADP No. 05770417001]

Publication No GB2250743 dated 17.06.1992

Examination requested 04.11.1992

Patent Granted with effect from 15.02.1995/(Section 25(1)) with title
HETEROCYCLIC SULPHONAMIDE LEUKOTRIENE ANTAGONIST

16.06.1993 Application under Section 32 filed on 30.04.1993

Entry Type 8.1 Staff ID. KH1 Auth ID. F21

14.07.1993 *ZENECA LIMITED* Incorporated in the United Kingdom, Imperial
Chemical House, 9 Millbank, LONDON, SW1P 3JF, United Kingdom
[ADP No. 06254007001]registered as Applicant/Proprietor in place of
IMPERIAL CHEMICAL INDUSTRIES PLC, Incorporated in the United
Kingdom, Imperial Chemical House, Millbank, LONDON, SW1P 3JF,
United Kingdom [ADP No. 00000935003]by virtue of deed of assignment dated 30.12.1992. Certified copy
filed on GB2260996

Entry Type 8.4 Staff ID. RD Auth ID. F21

20.07.1994 Notification of change of Address For Service name of

MARTIN ALEXANDER HAY, IMPERIAL CHEMICAL INDUSTRIES PLC, ICI Group
Patents, Group Patents Services Dept, PO Box 6, Shire Park, Bessemer
Road, WELWYN GARDEN CITY, Hertfordshire, AL7 1HD, United Kingdom
[ADP No. 05770417001]

to

PAUL MILLINGTON DENERLEY, Imperial Chemical Industries PLC, ICI Group Patents, Group Patents Services Dept, PO Box 6, Shire Park, Bessemer Road, WELWYN GARDEN CITY, Hertfordshire, AL7 1HD, United Kingdom [ADP No. 01030618001]
dated 14.07.1994. Official evidence filed on GB9126353.3
Entry Type 7.2 Staff ID. 5AI1 Auth ID. F51

12.12.1994 Notification of change of Address For Service address of
PAUL MILLINGTON DENERLEY, Imperial Chemical Industries PLC, ICI Group Patents, Group Patents Services Dept, PO Box 6, Shire Park, Bessemer Road, WELWYN GARDEN CITY, Hertfordshire, AL7 1HD, United Kingdom [ADP No. 01030618001]
to
PAUL MILLINGTON DENERLEY, ZENECA PHARMACEUTICALS, Intellectual Property Group, Intellectual Property Department, Mereside, Alderley Park, MACCLESFIELD, SK10 4TG, United Kingdom [ADP No. 01030618002]
dated 28.11.1994. Written notification filed on GB9126353.3
Entry Type 7.3 Staff ID. AGT1 Auth ID. AO

***** END OF REGISTER ENTRY *****

OA80-01
FG

OPTICS - PATENTS

12/03/97 08:25:01
PAGE: 1

RENEWAL DETAILS

PUBLICATION NUMBER

GB2250743

PROPRIETOR(S)

Zeneca Limited, Incorporated in the United Kingdom, Imperial
Chemical House, 9 Millbank, LONDON, SW1P 3JF, United Kingdom

PLC 1

DATE FILED 11.12.1991

DATE GRANTED 15.02.1995

DATE NEXT RENEWAL DUE 11.12.1997

DATE NOT IN FORCE

DATE OF LAST RENEWAL 02.12.1996

YEAR OF LAST RENEWAL 06

STATUS PATENT IN FORCE

***** END OF REPORT *****