



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2017/12/15
(87) Date publication PCT/PCT Publication Date: 2018/06/21
(85) Entrée phase nationale/National Entry: 2020/06/10
(86) N° demande PCT/PCT Application No.: IB 2017/058023
(87) N° publication PCT/PCT Publication No.: 2018/109746
(30) Priorité/Priority: 2016/12/15 (US62/434,957)

(51) Cl.Int./Int.Cl. *G01N 33/02* (2006.01),
G01N 33/04 (2006.01), *G01N 33/08* (2006.01),
G01N 33/12 (2006.01), *G01N 33/50* (2006.01),
G01N 33/53 (2006.01), *G01N 33/543* (2006.01)
(71) Demandeur/Applicant:
BIOMERICA, INC., US
(72) Inventeurs/Inventors:
LADERMAN, ELISABETH, US;
IRANI-COHEN, ZACKARY, US
(74) Agent: BCF LLP

(54) Titre : COMPOSITIONS, DISPOSITIFS ET METHODES PERMETTANT DE TESTER UNE SENSIBILITE LIEE AU TROUBLE DEFICITAIRE DE L'ATTENTION AVEC OU SANS HYPERACTIVITE (TDA/TDAH)
(54) Title: COMPOSITIONS, DEVICES, AND METHODS OF ATTENTION DEFICIT DISORDER/ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADD/ADHD) SENSITIVITY TESTING

Distribution of ELISA Signal Scores by Diagnosis

Sex=FEMALE Food= Wheat




Figure 2C

(57) **Abrégé/Abstract:**

Contemplated test kits, diagnostic apparatus and methods using same for food sensitivity are based on rational-based selection of food preparations with established discriminatory p-value. Kits and diagnostic apparatus include those with a minimum number of food preparations that have an average discriminatory p-value of ≤ 0.07 as determined by their raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In further contemplated aspects, compositions and methods for food sensitivity are also stratified by gender to further enhance predictive value.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
21 June 2018 (21.06.2018)



(10) International Publication Number
WO 2018/109746 A1

(51) International Patent Classification:

G01N 33/02 (2006.01) G01N 33/50 (2006.01)
G01N 33/04 (2006.01) G01N 33/53 (2006.01)
G01N 33/08 (2006.01) G01N 33/543 (2006.01)
G01N 33/12 (2006.01)

(21) International Application Number:

PCT/IB2017/058023

(22) International Filing Date:

15 December 2017 (15.12.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/434,957 15 December 2016 (15.12.2016) US

(71) Applicant: **BIOMERICA, INC.** [US/US]; 17571 Von Karman Avenue, Irvine, CA 92614 (US).

(72) Inventors: **LADERMAN, Elisabeth**; 17571 Von Karman Ave, Irvine, California 92614 (US). **IRANI-COHEN, Zackary**; 17571 Von Karman Ave, Irvine, California 92614 (US).

(74) Agent: **HERRITT, Danielle L.** et al.; Womble Bond Dickinson (US) LLP, P.O. Box 7037, Atlanta, GA 30357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(54) Title: COMPOSITIONS, DEVICES, AND METHODS OF ATTENTION DEFICIT DISORDER/ ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADD/ADHD) SENSITIVITY TESTING

Distribution of ELISA Signal Scores by Diagnosis

Sex=FEMALE Food= Wheat




Figure 2C

(57) Abstract: Contemplated test kits, diagnostic apparatus and methods using same for food sensitivity are based on rational-based selection of food preparations with established discriminatory p-value. Kits and diagnostic apparatus include those with a minimum number of food preparations that have an average discriminatory p-value of ≤ 0.07 as determined by their raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In further contemplated aspects, compositions and methods for food sensitivity are also stratified by gender to further enhance predictive value.

[Continued on next page]



WO 2018/109746 A1

WO 2018/109746 A1

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

**COMPOSITIONS, DEVICES, AND METHODS OF ATTENTION DEFICIT DISORDER/
ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADD/ADHD) SENSITIVITY
TESTING**

Related Applications

[0001] This application claims priority to U.S. provisional patent application with the serial number, 62/434,957, filed December 15, 2016, which is incorporated by reference herein in its entirety.

Field

[0002] The field of the disclosure is related to the identification of food items that trigger and/or exacerbate ADD/ADHD, particularly as it relates to the testing, identification and possible elimination of selected food items as trigger foods for patients diagnosed with or suspected to have ADD/ADHD.

Background

[0003] The background description includes information that may be useful in understanding the present disclosure. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0004] Food sensitivity, especially as it relates to ADD/ADHD (a type of neurodevelopmental psychiatric disorder), often presents with significant problems of attention and/or hyperactivity, and underlying causes of ADD/ADHD are not well understood in the medical community. Most typically, ADD/ADHD is diagnosed by an assessment of a person's childhood behavioral and mental development. Unfortunately, treatment of ADD/ADHD is often less than effective and may present new difficulties due to neurochemical modulatory effects. Elimination of one or more food items has also shown promise in at least reducing incidence and/or severity of the symptoms. However, ADD/ADHD is often quite diverse with respect to dietary items triggering symptoms, and no standardized test to help identify trigger food items with a reasonable degree of certainty is known, leaving such patients often to trial-and-error.

[0005] All publications identified herein are incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be

incorporated by reference. Where a definition or use of a term in an incorporated reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply.

[0006] Accordingly, there is still a need for compositions, devices, and methods of food sensitivity testing, especially for identification and possible elimination of trigger foods for patients identified with or suspected of having ADD/ADHD.

Summary

[0007] The subject matter described herein provides systems and methods for identifying of food items that trigger and/or exacerbate ADD/ADHD symptomology in patients diagnosed with or suspected to have ADD/ADHD, or which alleviate ADD/ADHD symptomology when removed. One aspect of the disclosure is a test kit with for identifying food items that trigger and/or exacerbate ADD/ADHD symptomology, or which alleviate ADD/ADHD symptomology when removed, in patients diagnosed with or suspected of having ADD/ADHD. The test kit includes a plurality of distinct food preparations coupled to individually addressable respective solid carriers. The plurality of distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In some embodiments, the average discriminatory p-value is determined by a process, which includes comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or suspected of having ADD/ADHD.

[0008] Another aspect of the embodiments described herein includes a method of identifying food items that trigger and/or exacerbate ADD/ADHD symptomology, or which alleviate ADD/ADHD symptomology when removed, in patients diagnosed with or suspected of having ADD/ADHD. The method includes a step of contacting a food preparation with a bodily fluid of a patient that is diagnosed with or suspected to have ADD/ADHD. The bodily fluid is associated with gender identification. In certain embodiments, the step of contacting is performed under conditions that allow IgG from the bodily fluid to bind to at least one component of the food preparation. The method continues with a step of measuring IgG bound to the at least one component of the food preparation to obtain a signal, and then comparing the signal to a gender-stratified reference value for the food preparation using the gender identification to obtain a result. Then, the method also includes a step of updating or generating a report using the result.

[0009] Another aspect of the embodiments described herein includes a method of generating a test for identifying of food items that trigger and/or exacerbate ADD/ADHD symptomology, or which alleviate ADD/ADHD symptomology when removed, in patients diagnosed with or suspected to have ADD/ADHD. The method includes a step of obtaining test results for a plurality of distinct food preparations. The test results are based on bodily fluids of patients diagnosed with or suspected to have ADD/ADHD and bodily fluids of a control group not diagnosed with or not suspected to have ADD/ADHD. The method also includes a step of stratifying the test results by gender for each of the distinct food preparations. Then the method continues with a step of assigning for a predetermined percentile rank a different cutoff value for male and female patients for each of the distinct food preparations.

[0010] Still another aspect of the embodiments described herein includes a use of a plurality of distinct food preparations coupled to individually addressable respective solid carriers in a diagnosis of ADD/ADHD. The plurality of distinct food preparations are selected based on their average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0011] In one aspect, the disclosure features a test kit for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising one or more distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier; wherein the distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value, or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In some embodiments, the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD. In certain embodiments, the one or more distinct food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least three food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In other embodiments, the one or more distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least five food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least six food preparations prepared from food items of Table 1 or

selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least seven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the one or more distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least nine food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least ten food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least eleven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In yet other embodiments, the one or more distinct food preparations includes at least 12 food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least 13 food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In certain embodiments, the one or more distinct food preparations includes at least 14 or more food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

[0012] In some embodiments, the one or more food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-

37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

[0013] In some embodiments of the disclosure, the one or more distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2. In other embodiments, the one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0014] In other embodiments of the disclosure, the one or more distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value. In other embodiments, the one or more distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.

[0015] In other embodiments of the disclosure, the FDR multiplicity adjusted p-value is adjusted for at least one of age and gender. In another embodiment, the FDR multiplicity adjusted p-value is adjusted for age and gender.

[0016] In other embodiments of the disclosure, at least 50% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 55% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 60% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 65% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 70% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 75% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 80% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 85% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

FDR multiplicity adjusted p-value. In other embodiments, at least 95% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 95% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, all of the one or more distinct food preparations, when adjusted for a single gender, have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0017] In certain embodiments of the disclosure, the one or more distinct food preparations are crude filtered aqueous extracts. In other embodiments, the one or more distinct food preparations are processed aqueous extracts. In yet other embodiments, the one or more distinct food preparations are crude filtered aqueous extracts or processed aqueous extracts. In certain other embodiments, the one or more distinct food preparations are crude filtered aqueous extracts or processed aqueous extracts.

[0018] In certain embodiments of the disclosure, the solid carrier is a well of a multiwell plate, a bead, an electrical, a chemical sensor, a microchip or an adsorptive film.

[0019] In certain other embodiments, the disclosure features a method of testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising contacting a test kit (or diagnostic apparatus) of the disclosure with a bodily fluid of a patient that is diagnosed with or suspected of having ADD/ADHD, wherein the step of contacting is performed under conditions that allow IgG from the bodily fluid to bind to at least one component of the food preparation; measuring IgG bound to the at least one component of the food preparation to obtain a signal; and updating or generating a report using the signal. In some embodiments, the bodily fluid of the patient is selected from the group consisting of whole blood, plasma, serum, saliva, urine and a fecal suspension. In other embodiments, the step of contacting the test kit is performed with a plurality of distinct food preparations.

[0020] In certain embodiments, the plurality of distinct food preparations is prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In yet other embodiments, the plurality of distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food

preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

[0021] In other embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value. In still other embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value. In certain embodiments, all of the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0022] In some embodiments, the food preparation is immobilized on a solid surface. In other embodiments the food preparation is immobilized on a solid surface, optionally in an addressable manner.

[0023] In another embodiment, the step of measuring IgG bound to the at least one component of the food preparation is performed via immunoassay test. In yet another embodiment, the method comprises comparing the signal to a gender-stratified reference value for the food preparation using gender identification to obtain a result, wherein the gender-stratified reference value for the food preparation is at least a 90th percentile value.

[0024] In another aspect, the disclosure features a method of generating a test for food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising obtaining test results for a plurality of distinct food preparations, wherein the test results are based on bodily fluids of patients diagnosed with or suspected of having ADD/ADHD and bodily fluids of a control group not diagnosed with or not suspected of having ADD/ADHD; stratifying the test results by gender for each of the distinct food preparations; assigning for a predetermined percentile rank a different cutoff value for male and female patients for each of the distinct food preparations; selecting a plurality of distinct food preparations that each have a discriminatory p-value of ≤ 0.07 as determined by raw p-value or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value; and generating a test consisting essentially of the selected food preparations for food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD.

[0025] In some embodiments, the test result is an ELISA result.

[0026] In other embodiments, the plurality of distinct food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least three food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least five food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least six food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least seven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another

embodiment, the plurality of distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least nine food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least ten food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eleven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least twelve food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least thirteen food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least fourteen or more food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

[0027] In another embodiment, the plurality of distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2,

at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2. In yet another embodiment, the plurality of distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0028] In other embodiments, the plurality of different food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value. In another embodiment, the plurality of different food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.

[0029] In certain embodiments, the bodily fluid of the patient is selected from the group consisting of whole blood, plasma, serum, saliva, urine or a fecal suspension.

[0030] In other embodiments, the predetermined percentile rank is an at least 90th percentile rank.

[0031] In yet another embodiment, the cutoff value for male and female patients has a difference of at least 10% (abs).

[0032] In a further embodiment, the method further comprises a step of normalizing the result to a patient's total IgG. In another embodiment, the method further comprises a step of normalizing the result to a global mean of the patient's food specific IgG results.

[0033] In other embodiments, the method further comprises a step of identifying a subset of patients, wherein the subset of patients' sensitivities to the food preparations underlies ADD/ADHD by raw p-value or an average discriminatory p-value of ≤ 0.01 . In other embodiments, the method further comprises a step of determining numbers of the food preparations, wherein the numbers of the food preparations can be used to confirm ADD/ADHD by raw p-value or an average discriminatory p-value of ≤ 0.01 .

[0034] In certain aspects, the disclosure features a use of a plurality of distinct food preparations coupled to individually addressable respective solid carriers for the diagnosis of ADD/ADHD,

wherein the plurality of distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In certain embodiments, the plurality of food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least three food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least five food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least six food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least seven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least nine food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least ten food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eleven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least twelve food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least thirteen food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least fourteen or more food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

[0035] In other embodiments, the plurality of food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations

from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2. In yet another embodiment, the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0036] In certain embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value. In other embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.

[0037] In certain other embodiments, the FDR multiplicity adjusted p-value is adjusted for at least one of age and gender. In other embodiments, the FDR multiplicity adjusted p-value is adjusted for age and gender.

[0038] In other embodiments, at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 55% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 60% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 65% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet another embodiment, at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 75% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 80% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 85% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 90% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 95% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet another embodiment, all of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0039] In another embodiment, the plurality of distinct food preparations is crude filtered aqueous extracts. In other embodiments, the plurality of distinct food preparations is processed aqueous extracts. In yet other embodiments, the plurality of distinct food preparations is crude filtered aqueous extracts and processed aqueous extracts.

[0040] In another embodiment, the solid carrier is a well of a multiwell plate, a bead, an electrical sensor, a chemical sensor, a microchip, or an adsorptive film.

[0041] In other embodiments, the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD headaches with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD headaches.

[0042] In yet another embodiment, the test result is an ELISA result derived from a process that includes separately contacting each distinct food preparation with the bodily fluid of each patient.

[0043] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier; wherein the plurality of distinct food preparations consists essentially of food preparations that each have a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, the discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD.

[0044] In some embodiments, the plurality of food preparations includes at least two food preparations selected from food items of Table 1 or selected from foods items 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least three food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least five food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least six food preparations

prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least seven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least nine food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least ten food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eleven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least twelve food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least thirteen food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least fourteen or more food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

[0045] In other embodiments, the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2,

at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2. In yet other embodiments, the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0046] In some embodiments, the plurality of distinct food preparations consists essentially of food preparations that each has a discriminatory p-value of ≤ 0.05 as determined by raw p-value or a discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

[0047] In other embodiments, the FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.

[0048] In yet other embodiments, at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 55% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 60% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 65% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 75% of the plurality of distinct food preparations, when

adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 80% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 85% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 90% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 95% or more of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0049] In other embodiments, the plurality of distinct food preparations is a crude aqueous extract. In other embodiments, the plurality of distinct food preparations is a processed aqueous extract. In other embodiments, the plurality of distinct food preparations is a crude aqueous extract or a processed aqueous extract.

[0050] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier; wherein the plurality of distinct food preparations consists essentially of food preparations having an average discriminatory p-value of ≤ 0.07 as determined by raw p-value, or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In some embodiments, the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or suspected of having ADD/ADHD.

[0051] In other embodiments, the plurality of food preparations includes at least two food preparations selected from food items of Table 1 or selected from foods items 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least three food

preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least five food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least six food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least seven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least nine food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least ten food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least eleven food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least twelve food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least thirteen food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2. In another embodiment, the plurality of distinct food preparations includes at least fourteen or more food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

[0052] In other embodiments, the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2,

at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2. In another embodiment, the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0053] In certain other embodiment, the plurality of distinct food preparations consists essentially of food preparations having an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

[0054] In yet other embodiments, the FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.

[0055] In yet other embodiments, at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 55% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 60% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 65% of the plurality of distinct food preparations, when

adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet another embodiment, at least 75% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In certain embodiments, at least 80% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 85% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0056] In yet another embodiment, the plurality of distinct food preparations is a crude aqueous extract. In another embodiment, the plurality of distinct food preparations is a processed aqueous extract. In yet another embodiment, the plurality of distinct food preparations is a crude aqueous extract or a processed aqueous extract.

[0057] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier; wherein at least 50% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0058] In one embodiment, at least 50% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 55% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet another embodiment, at least 60% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 65% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 70% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 75% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In certain embodiments, at least 80% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 85% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 90% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 95% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0059] In some embodiments, the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-

37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

[0060] In some embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

[0061] In other embodiments, the FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.

[0062] In yet other embodiments, at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In other embodiments, at least 55% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 60% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In another embodiment, at least 65% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 75% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 80% of the plurality of distinct food preparations, when

adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 85% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 90% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In yet other embodiments, at least 95% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[0063] In some embodiments, the plurality of distinct food preparations is crude aqueous. In other embodiments, the plurality of distinct food preparations is processed aqueous extracts. In some embodiments, the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

[0064] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface, wherein the plurality of food preparations consists essentially of trigger foods for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In certain embodiments, the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-

37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods 1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2. In other embodiments, the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0065] In some embodiments, the plurality of distinct food preparations is crude aqueous. In other embodiments, the plurality of distinct food preparations is processed aqueous extracts. In some embodiments, the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

[0066] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface, wherein at least 50% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In another embodiment, at least 55% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In another embodiment, at least 60% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least 65% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In yet another embodiment, at least 70% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In further embodiments, at least 75% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In another embodiment, at least 80% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In certain other embodiments, at least 85% of the plurality of food

preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least 90% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In yet other embodiments, at least 95% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

[0067] In some embodiments, the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods 1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2. In yet another embodiment, the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0068] In some embodiments, the plurality of distinct food preparations is crude aqueous. In other embodiments, the plurality of distinct food preparations is processed aqueous extracts. In some embodiments, the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

[0069] In certain aspects, the disclosure features a detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface, wherein at least seven food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least eight food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least nine food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least ten food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least eleven food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least twelve food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least thirteen food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample. In other embodiments, at least fourteen food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

[0070] In other embodiments, the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from

foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods 1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2. In yet other embodiments, the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0071] In some embodiments, the plurality of distinct food preparations is crude aqueous. In other embodiments, the plurality of distinct food preparations is processed aqueous extracts. In some embodiments, the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

[0072] In certain aspects, the disclosure features a test kit with one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0073] In certain aspects, the disclosure features methods using one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0074] In certain aspects, the disclosure features uses using one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0075] In certain aspects, the disclosure features a detection apparatus with one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

[0076] Various objects, features, aspects and advantages of the embodiments described herein will become more apparent from the following detailed description of the embodiments, along with the accompanying drawing figures in which like numerals represent like components.

Brief Description of The Drawings

[0077] **Table 1** shows a list of food items from which food preparations can be prepared.

[0078] **Table 2** shows statistical data of foods ranked according to 2-tailed FDR multiplicity-adjusted p-values.

[0079] **Table 3** shows statistical data of ELISA score by food and gender.

[0080] **Table 4** shows cutoff values of foods for a predetermined percentile rank.

[0081] **Figure 1A** illustrates ELISA signal score of male ADD/ADHD patients and control tested with cantaloupe.

[0082] **Figure 1B** illustrates a distribution of percentage of male ADD/ADHD subjects exceeding the 90th and 95th percentile tested with cantaloupe.

[0083] **Figure 1C** illustrates a signal distribution in women along with the 95th percentile cutoff as determined from the female control population tested with cantaloupe.

[0084] **Figure 1D** illustrates a distribution of percentage of female ADD/ADHD subjects exceeding the 90th and 95th percentile tested with cantaloupe.

[0085] **Figure 2A** illustrates ELISA signal score of male ADD/ADHD patients and control tested with wheat.

[0086] **Figure 2B** illustrates a distribution of percentage of male ADD/ADHD subjects exceeding the 90th and 95th percentile tested with wheat.

[0087] **Figure 2C** illustrates a signal distribution in women along with the 95th percentile cutoff as determined from the female control population tested with wheat.

[0088] **Figure 2D** illustrates a distribution of percentage of female ADD/ADHD subjects exceeding the 90th and 95th percentile tested with wheat.

[0089] **Figure 3A** illustrates ELISA signal score of male ADD/ADHD patients and control tested with tomato.

[0090] **Figure 3B** illustrates a distribution of percentage of male ADD/ADHD subjects exceeding the 90th and 95th percentile tested with tomato.

[0091] **Figure 3C** illustrates a signal distribution in women along with the 95th percentile cutoff as determined from the female control population tested with tomato.

[0092] **Figure 3D** illustrates a distribution of percentage of female ADD/ADHD subjects exceeding the 90th and 95th percentile tested with tomato.

[0093] **Figure 4A** illustrates ELISA signal score of male ADD/ADHD patients and control tested with cucumber.

[0094] **Figure 4B** illustrates a distribution of percentage of male ADD/ADHD subjects exceeding the 90th and 95th percentile tested with cucumber.

[0095] **Figure 4C** illustrates a signal distribution in women along with the 95th percentile cutoff as determined from the female control population tested with cucumber.

[0096] **Figure 4D** illustrates a distribution of percentage of female ADD/ADHD subjects exceeding the 90th and 95th percentile tested with cucumber.

[0097] **Figure 5A** illustrates distributions of ADD/ADHD subjects by number of foods that were identified as trigger foods at the 90th percentile.

[0098] **Figure 5B** illustrates distributions of ADD/ADHD subjects by number of foods that were identified as trigger foods at the 95th percentile.

[0099] **Table 5A** shows raw data of ADD/ADHD patients and control with number of positive results based on the 90th percentile.

[00100] **Table 5B** shows raw data of ADD/ADHD patients and control with number of positive results based on the 95th percentile.

[00101] **Table 6A** shows statistical data summarizing the raw data of ADD/ADHD patient populations shown in Table 5A.

[00102] **Table 6B** shows statistical data summarizing the raw data of ADD/ADHD patient populations shown in Table 5B.

[00103] **Table 7A** shows statistical data summarizing the raw data of control populations shown in Table 5A.

[00104] **Table 7B** shows statistical data summarizing the raw data of control populations shown in Table 5B.

[00105] **Table 8A** shows statistical data summarizing the raw data of ADD/ADHD patient populations shown in Table 5A transformed by logarithmic transformation.

[00106] **Table 8B** shows statistical data summarizing the raw data of ADD/ADHD patient populations shown in Table 5B transformed by logarithmic transformation.

[00107] **Table 9A** shows statistical data summarizing the raw data of control populations shown in Table 5A transformed by logarithmic transformation.

[00108] **Table 9B** shows statistical data summarizing the raw data of control populations shown in Table 5B transformed by logarithmic transformation.

[00109] **Table 10A** shows statistical data of an independent T-test to compare the geometric mean number of positive foods between the ADD/ADHD and non-ADD/ADHD samples based on the 90th percentile.

[00110] **Table 10B** shows statistical data of an independent T-test to compare the geometric mean number of positive foods between the ADD/ADHD and non-ADD/ADHD samples based on the 95th percentile.

[00111] **Table 11A** shows statistical data of a Mann-Whitney test to compare the geometric mean number of positive foods between the ADD/ADHD and non-ADD/ADHD samples based on the 90th percentile.

[00112] **Table 11B** shows statistical data of a Mann-Whitney test to compare the geometric mean number of positive foods between the ADD/ADHD and non-ADD/ADHD samples based on the 95th percentile.

[00113] **Figure 6A** illustrates a box and whisker plot of data shown in Table 5A.

[00114] **Figure 6B** illustrates a notched box and whisker plot of data shown in Table 5A.

[00115] **Figure 6C** illustrates a box and whisker plot of data shown in Table 5B.

[00116] **Figure 6D** illustrates a notched box and whisker plot of data shown in Table 5B.

[00117] **Table 12A** shows statistical data of a Receiver Operating Characteristic (ROC) curve analysis of data shown in Tables 5A-11A.

[00118] **Table 12B** shows statistical data of a Receiver Operating Characteristic (ROC) curve analysis of data shown in Tables 5B-11B.

[00119] **Figure 7A** illustrates the ROC curve corresponding to the statistical data shown in Table 12A.

[00120] **Figure 7B** illustrates the ROC curve corresponding to the statistical data shown in Table 12B.

[00121] **Table 13A** shows a statistical data of performance metrics in predicting ADD/ADHD status among female patients from number of positive foods based on the 90th percentile.

[00122] **Table 13B** shows a statistical data of performance metrics in predicting ADD/ADHD status among male patients from number of positive foods based on the 90th percentile.

[00123] **Table 14A** shows a statistical data of performance metrics in predicting ADD/ADHD status among female patients from number of positive foods based on the 95th percentile.

[00124] **Table 14B** shows a statistical data of performance metrics in predicting ADD/ADHD status among male patients from number of positive foods based on the 95th percentile.

Detailed Description

[00125] The inventors have discovered that food preparations used in food tests to identify trigger foods in patients diagnosed with or suspected of having ADD/ADHD are not equally well

predictive and/or associated with ADD/ADHD symptoms. Indeed, various experiments have revealed that among a wide variety of food items, certain food items are highly predictive/associated with ADD/ADHD whereas others have no statistically significant association with ADD/ADHD.

[00126] Even more unexpectedly, the inventors discovered that in addition to the high variability of food items, gender variability with respect to response in a test plays a substantial role in the determination of association of a food item with ADD/ADHD. Consequently, based on the inventors' findings and further contemplations, test kits, detection apparatus or devices, array, chip or test panel and methods of using same are now presented with substantially higher predictive power in the choice of food items (i.e., trigger foods) that could be eliminated for reduction of ADD/ADHD signs and symptoms.

[00127] The following discussion provides many example embodiments of the inventive subject matter. Although each embodiment represents a single combination of inventive elements, the inventive subject matter is considered to include all possible combinations of the disclosed elements. Thus if one embodiment comprises elements A, B, and C, and a second embodiment comprises elements B and D, then the inventive subject matter is also considered to include other remaining combinations of A, B, C, or D, even if not explicitly disclosed.

[00128] In some embodiments, the numbers expressing quantities or ranges, used to describe and claim certain embodiments of the disclosure are to be understood as being modified in some instances by the term "about." Accordingly, in some embodiments, the numerical parameters set forth in the written description and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by a particular embodiment. In some embodiments, the numerical parameters should be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of some embodiments of the disclosure are approximations, the numerical values set forth in the specific examples are reported as precisely as practicable. The numerical values presented in some embodiments of the disclosure may contain certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Unless the context dictates the contrary, all ranges set forth herein should be interpreted as being inclusive of their endpoints and open-ended ranges should be interpreted to include only commercially practical values. Similarly, all lists of values should be considered as inclusive of intermediate values unless the context indicates the contrary.

[00129] As used in the description herein and throughout the claims that follow, the meaning of “a,” “an,” and “the” includes plural reference unless the context clearly dictates otherwise. Also, as used in the description herein, the meaning of “in” includes “in” and “on” unless the context clearly dictates otherwise.

[00130] The term “average discriminatory p-value”, as used herein, generally refers to an average of all the p-values with a particular probability (e.g., ≤ 0.05) as determined by raw p-value or with a particular probability (e.g., ≤ 0.1) as determined by FDR multiplicity adjusted p-value that were identified by analytical methods described herein, e.g., the analytical methods that produced the results summarized in Table 2 (for example, and in particular, foods 1-37 in Table 2). In one embodiment, average discriminatory p-value refers to an average of all the p-values with a particular probability (e.g., ≤ 0.05) as determined by raw p-value or with a particular probability (e.g., ≤ 0.1) as determined by FDR multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.1 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.09 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.08 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.075 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.07 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.065 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.05 as determined by raw p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.1 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.09 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.095 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by

analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.09 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.085 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.08 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.075 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.07 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.065 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein. In another embodiment, average discriminatory p-value refers to an average of all the p-values that are ≤ 0.06 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value that were identified by analytical methods described herein.

[00131] The term “food preparation”, as used herein, refers to a solubilized aqueous extraction of a specific food item (e.g., cantaloupe, wheat, milk, etc.). In certain embodiments, the specific food item is solubilized using a blender or similar apparatus, in the presence of a buffer and the food item is processed until the structure of the food item is broken down into a homogenous liquid suspension or solution.

[00132] The term “individually addressable”, as used herein, refers to a portion of a solid carrier (e.g. an ELISA well, etc.), wherein a food preparation is immobilized (or coupled, etc.) to said portion of the solid carrier in a manner that separates said food preparation from other food preparations immobilized to the solid carrier, and that allows for the detection of an immunoglobulin (e.g., an IgG or other binding molecule) capable of binding to said food preparation (or a component thereof).

[00133] The term “one component of the food preparation” or “a component of the food preparation”, as used herein, refers to any portion of a food preparation (e.g., a protein(s), a lipid(s),

a sugar(s), etc.) that is antigenic (i.e., capable of inducing and/or eliciting an immune response in a subject or patient).

[00134] The term “trigger food”, as used herein, broadly refers to a food preparation, or a component thereof, which will result in a significantly elevated immune response in a subject (e.g., a patient) exposed to the food preparation, or a component thereof, wherein the elevated immune response is highly correlated to the presence of a disease symptom(s), and which potentially may trigger and/or exacerbate a disease symptom(s); and/or which may potentially result in an alleviation or reduction of symptoms by removing the food preparation, or component thereof. In one embodiment, a trigger food may be characterized by a p-value of about ≤ 0.15 as determined by raw p-value. In another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.10 as determined by raw p-value. In another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.075 as determined by raw p-value. In yet another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.05 as determined by raw p-value. In other embodiments, a trigger food may be characterized by a p-value of about ≤ 0.10 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.09 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.08 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In yet another embodiment, a trigger food may be characterized by a p-value of about ≤ 0.07 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In another embodiment, a trigger food may be characterized as one of the food preparations listed in Table 1 and / or Table 2. In another embodiment, a trigger food may be characterized as one of the food preparations listed in Table 2. In another embodiment, a trigger food may be characterized as one of the food preparations with a rank number of 1-37 (i.e., also referred to herein as “foods 1-37 of Table 2”) as described in Table 2.

[00135] All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided with respect to certain embodiments herein is intended merely to better illuminate the disclosure and does not pose a limitation on the scope of any embodiments of the disclosure otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of any embodiments of the disclosure.

[00136] Groupings of alternative elements or embodiments of the disclosure disclosed herein are not to be construed as limitations. Each group member can be referred to and claimed individually or in any combination with other members of the group or other elements found herein. One or more members of a group can be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is herein deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

[00137] In one aspect, the inventors therefore contemplate a test kit, detection apparatus or device, array, chip or test panel that is suitable for identifying food items that trigger and/or exacerbate ADD/ADHD symptomology, or which alleviate ADD/ADHD symptomology when removed, in patients where the patient is diagnosed with or suspected of having ADD/ADHD. In some embodiments, such test kit, detection device or apparatus, array, chip or panel will include one or more (e.g., a plurality) of distinct food preparations (e.g., raw or processed extract, e.g., an aqueous extract with optional co-solvent, which may or may not be filtered) that are coupled to individually addressable respective solid carriers (e.g., in a form of an array or a micro well plate), wherein the distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In certain embodiments, the plurality of distinct food preparations consists essentially of food preparations that each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. In certain embodiments, at least 50% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

[00138] In some embodiments, the numbers expressing quantities of ingredients, properties such as concentration, reaction conditions, and so forth, used to describe and claim certain embodiments of the disclosure are to be understood as being **modified in some instances by the term “about.”** Accordingly, in some embodiments, the numerical parameters set forth in the written description and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by a particular embodiment. In some embodiments, the numerical parameters should be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of some embodiments of the disclosure are approximations, the numerical values set forth in the specific examples are reported as precisely as practicable. The numerical values

presented in some embodiments of the disclosure may contain certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Moreover, and unless the context dictates the contrary, all ranges set forth herein should be interpreted as being inclusive of their endpoints and open-ended ranges should be interpreted to include only commercially practical values. Similarly, all lists of values should be considered as inclusive of intermediate values unless the context indicates the contrary.

[00139] While not limiting to the inventive subject matter, food preparations will typically be drawn from foods generally known or suspected to trigger and/or exacerbate signs or symptoms of ADD/ADHD, or which alleviate ADD/ADHD symptomology when removed. Particularly suitable food preparations may be identified by the experimental procedures outlined below. Thus, it should be appreciated that the food items (e.g., trigger foods) need not be limited to the items described herein, but that all items are contemplated that can be identified by the methods presented herein. Therefore, in certain embodiments, exemplary food preparations include at least two, at least four, at least eight, or at least 12 food preparations prepared from foods 1-37 of Table 2. In other embodiments, exemplary food preparations are selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food

preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, at least 37 food preparations from foods 1-37 of Table 2. In other embodiments, exemplary food preparations are selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper. Still further especially contemplated food items and food additives from which food preparations can be prepared are listed in **Table 1**.

[00140] Using bodily fluids from patients diagnosed with or suspected of having ADD/ADHD and healthy control group individuals (*i.e.*, those not diagnosed with or not suspected of having ADD/ADHD), numerous additional food items (*i.e.*, trigger foods) may be identified. In one embodiment, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.15** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.14** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.13** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.12** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.11** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.10** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.09** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.08** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.07** as determined by raw p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.06** as determined by raw p-value. In yet other embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.05** as determined by raw p-value. In certain embodiments, identified food items (*i.e.*, trigger foods) will have high discriminatory power and as such have a **p-value of ≤ 0.10** as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In other embodiments, identified food items (*i.e.*, trigger foods) will

have high discriminatory power and as such have a p-value of ≤ 0.09 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In other embodiments, identified food items (i.e., trigger foods) will have high discriminatory power and as such have a p-value of ≤ 0.08 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In yet other embodiments, identified food items (i.e., trigger foods) will have high discriminatory power and as such have a p-value of ≤ 0.07 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value.

[00141] In certain embodiments, such identified food preparations will have high discriminatory power and, as such, will have a p-value of ≤ 0.15 , ≤ 0.10 , or even ≤ 0.05 as determined by raw p-value, and/or a p-value of ≤ 0.10 , ≤ 0.08 , or even ≤ 0.07 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In one embodiment, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.15 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.14 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.13 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.12 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.11 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.10 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.09 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.08 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.07 as determined by raw p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.06 as determined by raw p-value. In yet other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.05 as determined by raw p-value. In certain embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.10 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.09 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value. In other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.08 as determined by False

Discovery Rate (FDR) multiplicity adjusted p-value. In yet other embodiments, identified food preparations will have high discriminatory power and as such have a p-value of ≤ 0.07 as determined by False Discovery Rate (FDR) multiplicity adjusted p-value.

[00142] Therefore, where a test kit, detection device or apparatus, array, chip, or panel has multiple (i.e., a plurality of) food preparations, it is contemplated that in certain embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value. In other embodiments, the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value. In other embodiments, it should be appreciated that in certain embodiments, the FDR multiplicity adjusted p-value may be adjusted for at least one of age and gender, while in yet other embodiments, the FDR multiplicity adjusted p-value may be adjusted for both age and gender. On the other hand, where a test kit, detection device or apparatus, array, chip or panel is stratified for use with a single gender, it is also contemplated that in a test kit, detection device or apparatus, array, chip or panel at least 50%, or at least 55%, or at least 60%, or at least 65%, or at least 70%, or at last 75%, or at least 80%, or at least 85%, or at least 90%, or at least 95%, or all of the plurality of distinct food preparations, when adjusted for a single gender, have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value. Furthermore, it should be appreciated that other stratifications (e.g., dietary preference, ethnicity, place of residence, genetic predisposition or family history, etc.) are also contemplated, and the person of ordinary skill in the art (PHOSITA) will be readily appraised of the appropriate choice of stratification.

[00143] The recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided with respect to certain embodiments herein is intended merely to better illuminate certain embodiments of the disclosure and does not pose a limitation on the scope of the any embodiments of the disclosure otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the disclosure.

[00144] Of course, it should be noted that the particular format of the test kit, detection device or apparatus, array, chip, or panel may vary considerably and contemplated formats include micro well plates, dip sticks, membrane-bound arrays, etc. Consequently, the solid carrier to which the food preparations are coupled may include wells of a multiwell plate, a (*e.g.*, color-coded or magnetic) bead, or an adsorptive film (*e.g.*, nitrocellulose or micro/nanoporous polymeric film), or an electrical sensor, (*e.g.*, a printed copper sensor or microchip).

[00145] Consequently, the inventors also contemplate a method of identifying food items that trigger and/or exacerbate ADD/ADHD symptomology in patients that are diagnosed with or suspected to have ADD/ADHD. Most typically, such methods will include a step of contacting a food preparation with a bodily fluid (*e.g.*, including, but not limited to, whole blood, plasma, serum, saliva, urine, or a fecal suspension) of a patient that is diagnosed with or suspected to have ADD/ADHD, and wherein the bodily fluid is associated with a gender identification. As noted before, the step of contacting is performed, in certain embodiments, under conditions that allow an immunoglobulin, *e.g.*, IgG (or IgE or IgA or IgM or IgD) from the bodily fluid to bind to at least one component of the food preparation, and the IgG bound to the component(s) of the food preparation are then quantified/measured to obtain a signal. In some embodiments, the signal is then compared against a gender-stratified reference value (*e.g.*, at least a 90th percentile value) for the food preparation using the gender identification to obtain a result, which is then used to update or generate a report (*e.g.*, written medical report; oral report of results from doctor to patient; written or oral directive from physician based on results).

[00146] In certain embodiments, such methods will not be limited to a single food preparation, but will employ multiple different food preparations (*i.e.*, a plurality of distinct food preparations). As noted before, suitable food preparations can be identified using various methods as described herein. In certain embodiments, the food preparations include foods 1-37, of Table 2, and/or items of Table 1. As also noted herein, in some embodiments, at least some, or all of the different food preparations have an average discriminatory p-value of ≤ 0.07 (or ≤ 0.065 , or ≤ 0.06 , or ≤ 0.055 , or ≤ 0.05 , or ≤ 0.045 , or ≤ 0.04 , or ≤ 0.035 , or ≤ 0.03 , or ≤ 0.025) as determined by raw p-value, and/or or an average discriminatory p-value of ≤ 0.10 (or ≤ 0.095 , or ≤ 0.09 , or ≤ 0.085 , or ≤ 0.08 , or ≤ 0.075 , or ≤ 0.07) as determined by FDR multiplicity adjusted p-value.

[00147] While in certain embodiments food preparations are prepared from single food items as crude extracts, or crude filtered extracts, it is contemplated that food preparations can be prepared from mixtures of a plurality of food items (*e.g.*, a mixture of citrus comprising lemon, orange, and a

grapefruit, a mixture of yeast comprising baker's yeast and brewer's yeast, a mixture of rice comprising a brown rice and white rice, a mixture of sugars comprising honey, malt, and cane sugar. In some embodiments, it is also contemplated that food preparations can be prepared from purified food antigens or recombinant food antigens.

[00148] As provided in certain embodiments, the food preparation is immobilized on a solid surface (typically in an addressable manner), accordingly, it is contemplated that the step of measuring the immunoglobulin (e.g., IgG or other type of antibody) bound to the component of the food preparation is performed via an ELISA test. Exemplary solid surfaces include, but are not limited to, wells in a multiwell plate, such that each food preparation may be isolated to a separate microwell. In certain embodiments, the food preparation will be coupled to, or immobilized on, the solid surface. In other embodiments, the food preparation(s) will be coupled to a molecular tag that allows for binding to human immunoglobulins (e.g., IgG) in solution. See, e.g., Example 3.

[00149] Viewed from a different perspective, the inventors also contemplate a method of generating a test for identifying food items that trigger and/or exacerbate ADD/ADHD in patients diagnosed with or suspected of having ADD/ADHD. Thus, in certain embodiments, the method is for identifying triggering food items (i.e., trigger foods) among already diagnosed or suspected ADD/ADHD patients. Such test will typically include a step of obtaining one or more test results (e.g., ELISA) for various distinct food preparations, wherein the test results are based on bodily fluids (e.g., including, but not limited to, whole blood, plasma, serum, saliva, urine, fecal suspension) of patients diagnosed with or suspected to have ADD/ADHD and bodily fluids of a control group not diagnosed with or not suspected to have ADD/ADHD. In other embodiments, the method is for identifying triggering food items (i.e., trigger foods) among patients only suspected of having ADD/ADHD. In certain embodiments, the test results are then stratified by gender for each of the distinct food preparations, a different cutoff value for male and female patients for each of the distinct food preparations (e.g., cutoff value for male and female patients has a difference of at least 10% (abs)) is assigned for a predetermined percentile rank (e.g., 90th or 95th percentile).

[00150] As noted herein, and while not limiting to the inventive subject matter, it is contemplated that in certain embodiments, the plurality of distinct food preparations include at least two (or six, or ten, or 15) food preparations prepared from food items selected from the group consisting of foods 1-37 of Table 2, and/or items of Table 1. In other embodiments, the plurality of distinct food preparations are selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food

preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, at least 37 food preparations from foods 1-37 of Table 2. On the other hand, where new food items are tested, it should be appreciated that the distinct food preparations include a food preparation prepared from a food items other than foods 1-37 of Table 2. Regardless of the particular choice of food items, in certain embodiments, the plurality of distinct food preparations have an average discriminatory p-value of ≤ 0.07 (or ≤ 0.05 , or ≤ 0.025) as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 (or ≤ 0.08 , or ≤ 0.07) as determined by FDR multiplicity adjusted p-value. In other embodiments, the plurality of distinct food preparations have an average discriminatory p-value as determined by raw p-value selected from the group consisting of about ≤ 0.07 , about ≤ 0.065 , about ≤ 0.06 , about ≤ 0.055 , about ≤ 0.05 , about ≤ 0.045 , about ≤ 0.04 , about ≤ 0.035 , about ≤ 0.03 , about ≤ 0.025 , and about ≤ 0.02 . In yet other embodiments, the plurality of distinct food preparations have an average discriminatory p-value as determined by FDR multiplicity adjusted p-value selected from the group consisting of about ≤ 0.10 , about ≤ 0.095 , about ≤ 0.09 , about ≤ 0.085 , about ≤ 0.08 , about ≤ 0.075 , about ≤ 0.07 , about ≤ 0.065 , about $\leq$

0.06, about ≤ 0.055 , and about ≤ 0.05 . Exemplary aspects and protocols, and considerations are provided in the experimental description below.

[00151] Thus, it should be appreciated that by having a high-confidence test system as described herein, the rate of false-positive and false negatives can be significantly reduced, and especially where the test systems and methods are gender stratified or adjusted for gender differences as shown below. Such advantages have heretofore not been realized and it is expected that the systems and methods presented herein will substantially increase the predictive power of food sensitivity tests for patients diagnosed with or suspected of having ADD/ADHD.

Examples

[00152] Example 1: General Protocol for food preparation generation: Commercially available food extracts (available from Biomerica Inc., 17571 Von Karman Ave, Irvine, CA 92614) prepared from the edible portion of the respective raw foods were used to prepare ELISA plates following the manufacturer's instructions.

[00153] For some food extracts, the inventors expect that food extracts prepared with certain specific procedures to generate the food extracts, provides more superior results in detecting elevated immunoglobulin (e.g., IgG) reactivity in ADD/ADHD patients compared to commercially available food extracts. For example, in certain embodiments related to grains and nuts, a three-step procedure of generating food extracts may be used. The first step is a defatting step. In this step, lipids from grains and nuts are extracted by contacting the flour of grains and nuts with a non-polar solvent and collecting residue. Then, the defatted grain or nut flour are extracted by contacting the flour with elevated pH to obtain a mixture and removing the solid from the mixture to obtain the liquid extract. Once the liquid extract is generated, the liquid extract is stabilized by adding an aqueous formulation. In certain embodiments, the aqueous formulation includes a sugar alcohol, a metal chelating agent, protease inhibitor, mineral salt, and buffer component 20-50 mM of buffer from 4-9 pH. This formulation allowed for long term storage at -70 °C and multiple freeze-thaws without a loss of activity.

[00154] For another example, in certain embodiments related to meats and fish, a two-step procedure of generating food extracts may be used. The first step is an extraction step. In this step, extracts from raw, uncooked meats or fish are generated by emulsifying the raw, uncooked meats or fish in an aqueous buffer formulation in a high impact pressure processor. Then, solid materials are removed to obtain liquid extract. Once the liquid extract is generated, the liquid extract is stabilized

by adding an aqueous formulation. In certain embodiments, the aqueous formulation includes a sugar alcohol, a metal chelating agent, protease inhibitor, mineral salt, and buffer component 20-50 mM of buffer from 4-9 pH. This formulation allowed for long term storage at -70 °C and multiple freeze-thaws without a loss of activity.

[00155] For still another example, in certain embodiments related to fruits and vegetables, a two-step procedure of generating food extracts may be used. The first step is an extraction step. In this step, liquid extracts from fruits or vegetables are generated using an extractor (*e.g.*, masticating juicer, etc.) to pulverize foods and extract juice. Then, solid materials are removed to obtain liquid extract. Once the liquid extract is generated, the liquid extract is stabilized by adding an aqueous formulation. In certain embodiments, the aqueous formulation includes a sugar alcohol, a metal chelating agent, protease inhibitor, mineral salt, and buffer component 20-50 mM of buffer from 4-9 pH. This formulation allowed for long term storage at -70 °C and multiple freeze-thaws without a loss of activity.

[00156] Example 2: Blocking of ELISA plates: To optimize signal to noise, plates will be blocked with a blocking buffer. In one embodiment, the blocking buffer includes 20-50 mM of a phosphate buffer (pH 4-9), bovine serum albumin (BSA) and a polyvinyl alcohol (PVA).

[00157] Example 3: ELISA preparation and sample testing: Food antigen preparations (*i.e.*, food preparations) were immobilized onto respective microtiter wells following the manufacturer's instructions. For the assays, the food antigens (*i.e.*, food preparations) were allowed to react with antibodies present in the patients' serum (*e.g.*, IgG), and excess serum proteins were removed by a wash step. For detection of IgG antibody binding, enzyme labeled anti-IgG antibody conjugate was allowed to react with antigen-antibody complex. A color was developed by the addition of a substrate that reacts with the coupled enzyme. The color intensity was measured and is directly proportional to the concentration of IgG antibody specific to a particular food antigen.

[00158] Example 4: Methodology to determine ranked food list in order of ability of ELISA signals to distinguish ADD/ADHD from control subjects: Out of an initial selection (*e.g.*, 100 food items, or 150 food items, or even more), samples can be eliminated prior to analysis due to low consumption in an intended, or target population. In addition, specific food items can be used as being representative of the larger more generic food group, especially where prior testing has established a correlation among different species within a generic group (*e.g.* in both genders, but also suitable for correlation for a single gender). For example, in one embodiment, green pepper could be dropped in favor of chili pepper as representative of the "pepper" food group, or sweet

potato could be dropped in favor of potato as representative of the “potato” food group. In other embodiments, the final list foods will be less than 50 food items. In another embodiment, the final list of foods will be equal to or less than of 40 food items. In other embodiments, the final list of foods is selected from the group consisting of less than 50 food items, less than 49 food items, less than 48 food items, less than 47 food items, less than 46 food items, less than 45 food items, less than 44 food items, less than 43 food items, less than 42 food items, less than 41 food items, less than 40 food items, less than 39 food items, less than 38 food items, less than 37 food items, less than 36 food items, less than 35 food items, less than 34 food items, less than 33 food items, less than 32 food items, less than 31 food items, less than 30 food items, less than 29 food items, less than 28 food items, less than 27 food items, less than 26 food items, less than 25 food items, less than 24 food items, less than 23 food items, less than 22 food items, less than 21 food items, less than 20 food items less than 19 food items, less than 18 food items, less than 17 food items, less than 16 food items, less than 15 food items, less than 14 food items, less than 13 food items, less than 12 food items, less than 11 food items, less than 10 food items, less than 9 food items, less than 8 food items, less than 7 food items, less than 6 food items, less than 5 food items, less than 4 food items, less than 3 food items, and less than 2 food items.

[00159] Since the foods ultimately selected for the food panel will not be specific for a particular gender, a gender-neutral food list is necessary. Since the observed sample will be at least initially imbalanced by gender (*e.g.*, Controls: 38.6% female, ADD/ADHD: 67.1% female), differences in ELISA signal magnitude strictly due to gender will be removed by modeling signal scores against gender using a two-sample t-test and storing the residuals for further analysis. For each of the tested foods (*i.e.*, food preparations), residual signal scores will be compared between ADD/ADHD and controls using a permutation test on a two-sample t-test with a relative high number of resamplings (*e.g.*, in certain embodiments >1,000 resamplings; in other embodiments >10,000 resamplings; in yet other embodiments >50,000 resamplings). The Satterthwaite approximation can then be used for the denominator degrees of freedom to account for lack of homogeneity of variances, and the 2-tailed permuted p-value will represent the raw p-value for each food. False Discovery Rates (FDR) among the comparisons, will be adjusted by any acceptable statistical procedures (*e.g.*, Benjamini-Hochberg, Family-wise Error Rate (FWER), Per Comparison Error Rate (PCER), etc.).

[00160] Foods (*i.e.*, food preparations) were then ranked according to their 2-tailed FDR multiplicity-adjusted p-values. Foods with adjusted p-values equal to or lower than the desired FDR threshold are deemed to have significantly higher signal scores among ADD/ADHD than control subjects and therefore deemed candidates for inclusion into a food panel. A typical result

that is representative of the outcome of the statistical procedure is provided in **Table 2**. Here the ranking of foods is according to 2-tailed permutation T-test p-values with FDR adjustment.

[00161] Based on earlier experiments, the inventors contemplate that even for the same food preparation tested, the ELISA score for at least several food items (i.e., food preparations) will vary dramatically, and exemplary raw data are provided in **Table 3**. As should be readily appreciated, data unstratified by gender will therefore lose significant explanatory power where the same cutoff value is applied to raw data for male and female data. To overcome such disadvantage, the inventors therefore contemplate stratification of the data by gender as described below.

[00162] Example 5: Statistical Method for Cutpoint Selection for each Food: The determination of what ELISA signal scores would constitute a “positive” response can be made by summarizing the distribution of signal scores among the Control subjects. For each food (i.e., food preparations), ADD/ADHD subjects who have observed scores greater than or equal to selected quantiles of the Control subject distribution will be deemed “positive”. To attenuate the influence of any one subject on cutpoint determination, each food-specific and gender-specific dataset will be bootstrap resampled 1000 times. Within each bootstrap replicate, the 90th and 95th percentiles of the Control signal scores will be determined. Each ADD/ADHD subject in the bootstrap sample will be compared to the 90th and 95th percentiles to determine whether he/she had a “positive” response. The final 90th and 95th percentile-based cutpoints for each food and gender will be computed as the average 90th and 95th percentiles across the 1000 samples. The number of foods for which each ADD/ADHD subject will be rated as “positive” was computed by pooling data across foods. Using such method, the inventors will be now able to identify cutoff values for a predetermined percentile rank that in most cases was substantially different as can be taken from **Table 4**.

[00163] Typical examples for the gender difference in IgG response in blood with respect to cantaloupe is shown in **Figures 1A-1D**, where Figure 1A shows the signal distribution in men along with the 95th percentile cutoff as determined from the male control population. Figure 1B shows the distribution of percentage of male ADD/ADHD subjects exceeding the 90th and 95th percentile, while Figure 1C shows the signal distribution in women along with the 95th percentile cutoff as determined from the female control population. Figure 1D shows the distribution of percentage of female ADD/ADHD subjects exceeding the 90th and 95th percentile. In the same fashion, **Figures 2A-2D** exemplarily depict the differential response to wheat, **Figures 3A-3D** exemplarily depict the differential response to tomato, and **Figures 4A-4D** exemplarily depict the differential response to cucumber. **Figures 5A-5B** show the distribution of ADD/ADHD subjects

by number of foods that were identified as trigger foods at the 90th percentile (5A) and 95th percentile (5B). Inventors contemplate that regardless of the particular food items (i.e., food preparations), male and female responses will be notably distinct.

[00164] It should be noted that nothing in the art have provided any predictable food groups related to ADD/ADHD that is gender-stratified. Thus, a discovery of food items (i.e., food preparations) that show distinct responses by gender is a surprising result, which could not be obviously expected in view of all previously available arts. In other words, selection of food items based on gender stratification provides an unexpected technical effect such that statistical significances for particular food items as triggering food among male or female ADD/ADHD patients have been significantly improved.

[00165] Example 6: Normalization of IgG Response Data: While the raw data of the patient's IgG response results can be used to compare strength of response among given foods, it is also contemplated that the IgG response results of a patient are normalized and indexed to generate unit-less numbers for comparison of relative strength of response to a given food. For example, one or more of a patient's food specific IgG results (*e.g.*, IgG specific to orange and IgG specific to malt) can be normalized to the patient's total IgG. The normalized value of the patient's IgG specific to orange can be 0.1 and the normalized value of the patient's IgG specific to malt can be 0.3. In this scenario, the relative strength of the patient's response to malt is three times higher compared to orange. Then, the patient's sensitivity to malt and orange can be indexed as such.

[00166] In other examples, one or more of a patient's food specific IgG results (*e.g.*, IgG specific to shrimp and IgG specific to pork) can be normalized to the global mean of that patient's food specific IgG results. The global means of the patient's food specific IgG can be measured by total amount of the patient's food specific IgG. In this scenario, the patient's specific IgG to shrimp can be normalized to the mean of patient's total food specific IgG (*e.g.*, mean of IgG levels to shrimp, pork, Dungeness crab, chicken, peas, etc.) . However, it is also contemplated that the global means of the patient's food specific IgG can be measured by the patient's IgG levels to a specific type of food via multiple tests. If the patient has been tested for his sensitivity to shrimp five times and to pork seven times previously, the patient's new IgG values to shrimp or to pork are normalized to the mean of five-times test results to shrimp or the mean of seven-times test results to pork. The normalized value of the patient's IgG specific to shrimp can be 6.0 and the normalized value of the patient's IgG specific to pork can be 1.0. In this scenario, the patient has six times higher sensitivity to shrimp at this time compared to his average sensitivity to shrimp, but substantially similar

sensitivity to pork. Then, the patient's sensitivity to shrimp and pork can be indexed based on such comparison.

[00167] Example 7: Methodology to determine the subset of ADD/ADHD patients with food sensitivities that underlie ADD/ADHD: While it is suspected that food sensitivities plays a substantial role in signs and symptoms of ADD/ADHD, some ADD/ADHD patients may not have food sensitivities that underlie ADD/ADHD. Those patients would not be benefit from dietary intervention to treat signs and symptoms of ADD/ADHD. To determine the subset of such patients, body fluid samples of ADD/ADHD patients and non- ADD/ADHD patients can be tested with ELISA test using test devices with up to 37 food samples.

[00168] **Table 5A** and **Table 5B** provide exemplary raw data. As should be readily appreciated, the data indicate number of positive results out of 37 sample foods based on 90th percentile value (**Table 5A**) or 95th percentile value (**Table 5B**). The first column is ADD/ADHD (n=137); second column is non-ADD/ADHD (n=132) by ICD-10 code. Average and median number of positive foods was computed for ADD/ADHD and non-ADD/ADHD patients. From the raw data shown in **Table 5A** and **Table 5B**, average and standard deviation of the number of positive foods (i.e., trigger foods) was computed for ADD/ADHD and non-ADD/ADHD patients. Additionally, the number and percentage of patients with zero positive foods was calculated for both ADD/ADHD and non-ADD/ADHD. The number and percentage of patients with zero positive foods in the ADD/ADHD population is approximately 40% lower than the percentage of patients with zero positive foods in the non- ADD/ADHD population (24.1% vs. 38.6%, respectively) based on 90th percentile value (**Table 5A**), and the percentage of patients in the ADD/ADHD population with zero positive foods is also significantly lower (i.e. approximately 40% lower) than that seen in the non-ADD/ADHD population (35.8 % vs. 59.1%, respectively) based on 95th percentile value (**Table 5B**). Thus, it can be easily appreciated that the ADD/ADHD patient having sensitivity to zero positive foods is unlikely to have food sensitivities underlying their signs and symptoms of ADD/ADHD.

[00169] **Table 6A** and **Table 7A** show exemplary statistical data summarizing the raw data of two patient populations shown in Table 5A. The statistical data includes normality, arithmetic mean, median, percentiles and 95% confidence interval (CI) for the mean and median representing number of positive foods (i.e., trigger foods) in the ADD/ADHD population and the non-ADD/ADHD population. **Table 6B** and **Table 7B** show exemplary statistical data summarizing the raw data of two patient populations shown in Table 5B. The statistical data includes normality,

arithmetic mean, median, percentiles and 95% confidence interval (CI) for the mean and median representing number of positive foods in the ADD/ADHD population and the non-ADD/ADHD population.

[00170] **Table 8A** and **Table 9A** show exemplary statistical data summarizing the raw data of two patient populations shown in Table 5A. In Tables 8A and 9A, the raw data was transformed by logarithmic transformation to improve the data interpretation. **Table 8B** and **Table 9B** show another exemplary statistical data summarizing the raw data of two patient populations shown in Table 5B. In Tables 8B and 9B, the raw data was transformed by logarithmic transformation to improve the data interpretation.

[00171] **Table 10A** and **Table 11A** show exemplary statistical data of an independent T-test (Table 10A, logarithmically transformed data) and a Mann-Whitney test (Table 11A) to compare the geometric mean number of positive foods (i.e., trigger foods) between the ADD/ADHD and non-ADD/ADHD samples. The data shown in Table 10A and Table 11A indicate statistically significant differences in the geometric mean of positive number of foods between the ADD/ADHD population and the non-ADD/ADHD population. In both statistical tests, it is shown that the number of positive responses with 37 food samples is significantly higher in the ADD/ADHD population than in the non-ADD/ADHD population with an average discriminatory p-value of ≤ 0.0001 . These statistical data is also illustrated as a box and whisker plot in **Figure 6A**, and a notched box and whisker plot in **Figure 6B**.

[00172] **Table 10B** and **Table 11B** show exemplary statistical data of an independent T-test (Table 10B, logarithmically transformed data) and a Mann-Whitney test (Table 11B) to compare the geometric mean number of positive foods (i.e., trigger foods) between the ADD/ADHD and non-ADD/ADHD samples. The data shown in Table 10B and Table 11B indicate statistically significant differences in the geometric mean of positive number of foods between the ADD/ADHD population and the non-ADD/ADHD population. In both statistical tests, it is shown that the number of positive responses with 37 food samples is significantly higher in the ADD/ADHD population than in the non-ADD/ADHD population with an average discriminatory p-value of ≤ 0.0001 . These statistical data is also illustrated as a box and whisker plot in **Figure 6C**, and a notched box and whisker plot in **Figure 6D**.

[00173] **Table 12A** shows exemplary statistical data of a Receiver Operating Characteristic (ROC) curve analysis of data shown in Tables 5A-11A to determine the diagnostic power of the test used in Table 5 at discriminating ADD/ADHD from non- ADD/ADHD subjects. When a cutoff

criterion of more than 2 positive foods (i.e., trigger foods) is used, the test yields a data with 53.3% sensitivity and 76.5% specificity, with an area under the curve (AUROC) of 0.660. The p-value for the ROC is significant at a p-value of <0.0001 . **Figure 7A** illustrates the ROC curve corresponding to the statistical data shown in Table 12A. Because the statistical difference between the ADD/ADHD population and the non-ADD/ADHD population is significant when the test results are cut off to a positive number of 2, the number of foods for which a patient tests positive could be used as a confirmation of the primary clinical diagnosis of ADD/ADHD, and whether it is likely that food sensitivities underlies on the patient's signs and symptoms of ADD/ADHD. Therefore, the above test can be used as another 'rule in' test to add to currently available clinical criteria for diagnosis for ADD/ADHD.

[00174] As shown in Tables 5A-12A, and Figure 7A, based on 90th percentile data, the number of positive foods (i.e., trigger foods) seen in ADD/ADHD vs. non-ADD/ADHD subjects is significantly different whether the geometric mean or median of the data is compared. The number of positive foods that a person has is indicative of the presence of ADD/ADHD in subjects. The test has discriminatory power to detect ADD/ADHD with 53.3% sensitivity and 76.5% specificity. Additionally, the absolute number and percentage of subjects with 0 positive foods is also very different in ADD/ADHD vs. non-ADD/ADHD subjects, with a far lower percentage of ADD/ADHD subjects (24.1%) having 0 positive foods than non-ADD/ADHD subjects (38.6%). The data suggests a subset of ADD/ADHD patients may have ADD/ADHD due to other factors than diet, and may not benefit from dietary restriction.

[00175] **Table 12B** shows exemplary statistical data of a Receiver Operating Characteristic (ROC) curve analysis of data shown in Tables 5B-11B to determine the diagnostic power of the test used in Table 5 at discriminating ADD/ADHD from non-ADD/ADHD subjects. When a cutoff criterion of more than 1 positive foods (i.e., trigger foods) is used, the test yields a data with 47.4% sensitivity and 81.1% specificity, with an area under the curve (AUROC) of 0.659. The p-value for the ROC is significant at a p-value of <0.0001 . **Figure 7B** illustrates the ROC curve corresponding to the statistical data shown in Table 12B. Because the statistical difference between the ADD/ADHD population and the non-ADD/ADHD population is significant when the test results are cut off to positive number of >1 , the number of foods that a patient tests positive could be used as a confirmation of the primary clinical diagnosis of ADD/ADHD, and whether it is likely that food sensitivities underlies on the patient's signs and symptoms of ADD/ADHD. Therefore, the above test can be used as another 'rule in' test to add to currently available clinical criteria for diagnosis for ADD/ADHD.

[00176] As shown in Tables 5B-12B, and Figure 7B, based on 95th percentile data, the number of positive foods (i.e., trigger foods) seen in ADD/ADHD vs. non-ADD/ADHD subjects is significantly different whether the geometric mean or median of the data is compared. The number of positive foods that a person has is indicative of the presence of ADD/ADHD in subjects. The test has discriminatory power to detect ADD/ADHD with 47.4% sensitivity and 81.1% specificity. Additionally, the absolute number and percentage of subjects with 0 positive foods is also very different in ADD/ADHD vs. non-ADD/ADHD subjects, with a far lower percentage of ADD/ADHD subjects (35.8%) having 0 positive foods than non-ADD/ADHD subjects (59.1%). The data suggests a subset of ADD/ADHD patients may have ADD/ADHD due to other factors than diet, and may not benefit from dietary restriction.

[00177] Example 8: Method for determining distribution of per-person number of foods declared “positive”: To determine the distribution of number of “positive” foods (i.e., trigger foods) per person and measure the diagnostic performance, the analysis will be performed with 37 food items from Table 2, which shows most positive responses to ADD/ADHD patients. To attenuate the influence of any one subject on this analysis, each food-specific and gender-specific dataset will be bootstrap resampled 1000 times. Then, for each food item in the bootstrap sample, sex-specific cutpoint will be determined using the 90th and 95th percentiles of the control population. Once the sex-specific cutpoints are determined, the sex-specific cutpoints will be compared with the observed ELISA signal scores for both control and ADD/ADHD subjects. In this comparison, if the observed signal is equal or more than the cutpoint value, then it will be determined “positive” food, and if the observed signal is less than the cutpoint value, then it will be determined “negative” food.

[00178] Once all food items (i.e., food preparations) were determined either positive or negative, the results of the 74 (37 foods x 2 cutpoints) calls for each subject will be saved within each bootstrap replicate. Then, for each subject, 37 calls will be summed using 90th percentile as cutpoint to get “Number of Positive Foods (90th),” and the rest of 37 calls will be summed using 95th percentile to get “Number of Positive Foods (95th).” Then, within each replicate, “Number of Positive Foods (90th)” and “Number of Positive Foods (95th)” will be summarized across subjects to get descriptive statistics for each replicate as follows: 1) overall means equals to the mean of means, 2) overall standard deviation equals to the mean of standard deviations, 3) overall medial equals to the mean of medians, 4) overall minimum equals to the minimum of minimums, and 5) overall maximum equals to maximum of maximum. In this analysis, to avoid non-integer “Number of Positive Foods” when computing frequency distribution and histogram, the authors will pretend that the 1000 repetitions of the same original dataset were actually 999 sets of new subjects of the

same size added to the original sample. Once the summarization of data is done, frequency distributions and histograms will be generated for both “Number of Positive Foods (90th)” and “Number of Positive Foods (95th)” for both genders and for both ADD/ADHD subjects and control subjects using programs “a_pos_foods.sas, a_pos_foods_by_dx.sas”.

[00179] Example 9: Method for measuring diagnostic performance: To measure diagnostic performance for each food items (i.e., food preparations) for each subject, we will use data of “Number of Positive Foods (90th)” and “Number of Positive Foods (95th)” for each subject within each bootstrap replicate described above. In this analysis, the cutpoint was set to 1. Thus, if a subject has one or more “Number of Positive Foods (90th)”, then the subject will be called “Has ADD/ADHD.” If a subject has less than one “Number of Positive Foods (90th)”, then the subject will be called “Does Not Have ADD/ADHD.” When all calls were made, the calls were compared with actual diagnosis to determine whether a call was a True Positive (TP), True Negative (TN), False Positive (FP), or False Negative (FN). The comparisons will be summarized across subjects to get the performance metrics of sensitivity, specificity, positive predictive value, and negative predictive value for both “Number of Positive Foods (90th)” and “Number of Positive Foods(95th)” when the cutpoint is set to 1 for each method. Each (sensitivity, 1-specificity) pair becomes a point on the ROC curve for this replicate.

[00180] To increase the accuracy, the analysis above will be repeated by incrementing cutpoint from 2 up to 37, and repeated for each of the 1000 bootstrap replicates. Then the performance metrics across the 1000 bootstrap replicates will be summarized by calculating averages using a program “t_pos_foods_by_dx.sas”. The results of diagnostic performance for female and male are shown in **Tables 13A and 13B** (90th percentile) and **Tables 14 A and 14B** (95th percentile).

[00181] Of course, it should be appreciated that certain variations in the food preparations may be made without altering the inventive subject matter presented herein. For example, where the food item was yellow onion, that item should be understood to also include other onion varieties that were demonstrated to have equivalent activity in the tests. Indeed, the inventors have noted that for each tested food preparation, certain other related food preparations also tested in the same or equivalent manner (data not shown). Thus, it should be appreciated that each tested and claimed food preparation will have equivalent related preparations with demonstrated equal or equivalent reactions in the test.

[00182] It should be apparent to those skilled in the art that many more modifications besides those already described are possible without departing from the inventive concepts herein. The

inventive subject matter, therefore, is not to be restricted except in the spirit of the appended claims. Moreover, in interpreting both the specification and the claims, all terms should be interpreted in the broadest possible manner consistent with the context. In particular, the terms “comprises” and “comprising” should be interpreted as referring to elements, components, or steps in a non-exclusive manner, indicating that the referenced elements, components, or steps may be present, or utilized, or combined with other elements, components, or steps that are not expressly referenced. Where the specification claims refers to at least one of something selected from the group consisting of A, B, C and N, the text should be interpreted as requiring only one element from the group, not A plus N, or B plus N, etc.

Abalone	Cured Cheese	Onion	Walnut, black
Adlay	Cuttlefish	Orange	Watermelon
Almond	Duck	Oyster	Welch Onion
American Cheese	Durian	Papaya	Wheat
Apple	Eel	Paprika	Wheat bran
Artichoke	Egg White (separate)	Parsley	Yeast (<i>S. cerevisiae</i>)
Asparagus	Egg Yolk (separate)	Peach	Yogurt
Avocado	Egg, white/yolk (comb.)	Peanut	
Baby Bok Choy	Eggplant	Pear	FOOD ADDITIVES
Bamboo shoots	Garlic	Pepper, Black	Arabic Gum
Banana	Ginger	Pineapple	Carboxymethyl Cellulose
Barley, whole grain	Gluten - Gliadin	Pinto bean	Carrageneenan
Beef	Goat's milk	Plum	FD&C Blue #1
Beets	Grape, white/concord	Pork	FD&C Red #3
Beta-lactoglobulin	Grapefruit	Potato	FD&C Red #40
Blueberry	Grass Carp	Rabbit	FD&C Yellow #5
Broccoli	Green Onion	Rice	FD&C Yellow #6
Buckwheat	Green pea	Roquefort Cheese	Gelatin
Butter	Green pepper	Rye	Guar Gum
Cabbage	Guava	Saccharine	Maltodextrin
Cane sugar	Hair Tail	Safflower seed	Pectin
Cantaloupe	Hake	Salmon	Whey
Caraway	Halibut	Sardine	Xanthan Gum
Carrot	Hazelnut	Scallop	
Casein	Honey	Sesame	
Cashew	Kelp	Shark fin	
Cauliflower	Kidney bean	Sheep's milk	
Celery	Kiwi Fruit	Shrimp	
Chard	Lamb	Sole	
Cheddar Cheese	Leek	Soybean	
Chick Peas	Lemon	Spinach	
Chicken	Lentils	Squashes	
Chili pepper	Lettuce, Iceberg	Squid	
Chocolate	Lima bean	Strawberry	
Cinnamon	Lobster	String bean	
Clam	Longan	Sunflower seed	
Cocoa Bean	Mackerel	Sweet potato	
Coconut	Malt	Swiss cheese	
Codfish	Mango	Taro	
Coffee	Marjoram	Tea, black	
Cola nut	Millet	Tobacco	
Corn	Mung bean	Tomato	
Cottage cheese	Mushroom	Trout	
Cow's milk	Mustard seed	Tuna	
Crab	Oat	Turkey	
Cucumber	Olive	Vanilla	

Table 1

**Ranking of Foods according to 2-tailed Permutation T-test
p-values with FDR adjustment**

<i>Rank</i>	<i>Food</i>	<i>Raw p-value</i>	<i>FDR Multiplicity-adj p-value</i>
1	Cantaloupe	0.0000	0.0017
2	Wheat	0.0000	0.0017
3	Tomato	0.0001	0.0029
4	Cucumber	0.0001	0.0030
5	Squashes	0.0003	0.0046
6	Almond	0.0003	0.0046
7	Egg	0.0004	0.0048
8	Cauliflower	0.0005	0.0048
9	Pinto_Bean	0.0005	0.0048
10	Broccoli	0.0006	0.0048
11	Orange	0.0008	0.0063
12	Butter	0.0011	0.0077
13	Corn	0.0012	0.0077
14	Lettuce	0.0013	0.0081
15	Rye	0.0016	0.0092
16	Peach	0.0026	0.0139
17	Green_Pea	0.0031	0.0156
18	Carrot	0.0035	0.0168
19	Tea	0.0051	0.0233
20	Mustard	0.0058	0.0249
21	Strawberry	0.0062	0.0253
22	Celery	0.0093	0.0351
23	Green_Pepper	0.0094	0.0351
24	Garlic	0.0101	0.0358
25	Oat	0.0104	0.0358
26	Onion	0.0120	0.0396
27	Banana	0.0139	0.0443
28	Eggplant	0.0156	0.0479
29	Cabbage	0.0162	0.0480
30	Safflower	0.0168	0.0480
31	Olive	0.0175	0.0487
32	Cottage_Ch_	0.0190	0.0512

<i>Rank</i>	<i>Food</i>	<i>Raw p-value</i>	<i>FDR Multiplicity-adj p-value</i>
33	Grapefruit	0.0208	0.0541
34	Walnut_Black	0.0243	0.0615
35	Cow_Milk	0.0256	0.0629
36	Soybean	0.0300	0.0717
37	Chili_Pepper	0.0376	0.0874
38	Tobacco	0.0481	0.1088
39	Malt	0.0507	0.1118
40	Cinnamon	0.0545	0.1171
41	Cane_Sugar	0.0619	0.1299
42	Potato	0.0665	0.1362
43	Millet	0.0704	0.1407
44	Sunflower_Sd	0.0807	0.1577
45	Crab	0.0850	0.1625
46	Lima_Bean	0.1080	0.2020
47	Chocolate	0.1114	0.2038
48	Spinach	0.1205	0.2159
49	Buck_Wheat	0.1283	0.2241
50	Honey	0.1303	0.2241
51	Pineapple	0.1376	0.2274
52	Yeast_Brewer	0.1382	0.2274
53	Rice	0.1401	0.2274
54	Apple	0.1962	0.3125
55	Mushroom	0.2001	0.3128
56	Yeast_Baker	0.2551	0.3918
57	Avocado	0.2707	0.4075
58	Cheddar_Ch_	0.2803	0.4075
59	Sesame	0.2821	0.4075
60	Clam	0.2843	0.4075
61	Pork	0.2982	0.4204
62	Blueberry	0.3534	0.4903
63	Swiss_Ch_	0.3654	0.4987
64	Barley	0.3841	0.5162
65	String_Bean	0.4313	0.5707
66	Peanut	0.4626	0.6028

<i>Rank</i>	<i>Food</i>	<i>Raw p-value</i>	<i>FDR Multiplicity-adj p-value</i>
67	Scallop	0.4997	0.6414
68	Lemon	0.5191	0.6565
69	Goat_Milk	0.5378	0.6624
70	Oyster	0.5392	0.6624
71	Lobster	0.5621	0.6809
72	Chicken	0.6743	0.8055
73	Amer__Cheese	0.7096	0.8312
74	Coffee	0.7152	0.8312
75	Beef	0.7616	0.8733
76	Codfish	0.8330	0.9426
77	Salmon	0.8779	0.9646
78	Cashew	0.8950	0.9646
79	Sardine	0.8972	0.9646
80	Parsley	0.9073	0.9646
81	Turkey	0.9086	0.9646
82	Grape	0.9376	0.9780
83	Yogurt	0.9439	0.9780
84	Cola_Nut	0.9742	0.9931
85	Sweet_Pot_	0.9840	0.9931
86	Trout	0.9931	0.9931

Table 2

Basic Descriptive Statistics of ELISA Score by Food and Gender
Comparing ADD/ADHD to Control

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
FEMALE	Almond	Control	51	4.057	1.630	1.270	8.090
		ADHD	92	6.654	7.307	0.700	49.320
		Diff (1-2)	—	-2.597	5.950	—	—
	Amer__Cheese	Control	51	32.961	66.654	0.100	400.00
		ADHD	92	32.901	33.849	1.210	148.00
		Diff (1-2)	—	0.060	48.113	—	—
	Apple	Control	51	4.792	6.600	0.100	44.160
		ADHD	92	5.578	8.466	0.110	57.720
		Diff (1-2)	—	-0.785	7.855	—	—
	Avocado	Control	51	2.144	1.129	0.100	5.560
		ADHD	92	3.177	4.055	0.100	33.660
		Diff (1-2)	—	-1.032	3.326	—	—
	Banana	Control	51	3.042	2.909	0.100	17.210
		ADHD	92	6.176	11.816	0.100	105.04
		Diff (1-2)	—	-3.134	9.649	—	—
	Barley	Control	51	3.867	3.919	0.100	25.110
		ADHD	92	9.954	19.858	0.100	133.34
		Diff (1-2)	—	-6.087	16.123	—	—
	Beef	Control	51	9.007	12.047	1.030	81.660
		ADHD	92	12.865	40.258	1.270	380.80
		Diff (1-2)	—	-3.858	33.128	—	—
	Blueberry	Control	51	3.533	4.712	0.100	26.460
		ADHD	92	4.293	5.077	0.860	34.650
		Diff (1-2)	—	-0.760	4.951	—	—
	Broccoli	Control	51	5.211	4.424	0.110	29.600
		ADHD	92	9.398	10.582	2.570	91.250
		Diff (1-2)	—	-4.187	8.900	—	—
	Buck_Wheat	Control	51	5.151	4.281	0.100	26.450
		ADHD	92	6.364	7.554	0.870	60.670
		Diff (1-2)	—	-1.213	6.583	—	—
	Butter	Control	51	17.809	24.982	0.100	150.93
		ADHD	92	23.143	21.663	0.740	104.00

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		Diff (1-2)	—	-5.334	22.895	—	—
	Cabbage	Control	51	5.038	6.005	0.350	37.840
		ADHD	92	8.829	14.699	0.740	121.38
		Diff (1-2)	—	-3.791	12.338	—	—
	Cane_Sugar	Control	51	15.190	10.152	3.460	50.450
		ADHD	92	25.370	25.407	4.050	116.64
		Diff (1-2)	—	-10.180	21.287	—	—
	Cantaloupe	Control	51	4.707	2.368	1.150	12.760
		ADHD	92	11.342	20.203	1.070	134.48
		Diff (1-2)	—	-6.635	16.291	—	—
	Carrot	Control	51	2.702	1.549	0.100	6.940
		ADHD	92	6.653	14.010	0.100	89.210
		Diff (1-2)	—	-3.952	11.293	—	—
	Cashew	Control	51	8.620	13.756	0.100	81.890
		ADHD	92	9.483	19.123	0.610	111.06
		Diff (1-2)	—	-0.863	17.410	—	—
	Cauliflower	Control	51	4.203	2.424	0.430	11.770
		ADHD	92	7.444	11.577	1.010	94.290
		Diff (1-2)	—	-3.241	9.412	—	—
	Celery	Control	51	7.815	5.560	2.060	32.830
		ADHD	92	11.498	13.759	2.010	86.600
		Diff (1-2)	—	-3.683	11.539	—	—
	Cheddar_Ch_	Control	51	25.261	59.384	1.530	400.00
		ADHD	92	27.840	36.629	0.100	189.39
		Diff (1-2)	—	-2.579	46.005	—	—
	Chicken	Control	51	14.077	8.350	2.690	50.000
		ADHD	92	16.458	13.609	2.270	79.130
		Diff (1-2)	—	-2.381	12.010	—	—
	Chili_Pepper	Control	51	7.281	6.348	0.570	32.360
		ADHD	92	10.595	16.487	0.870	111.31
		Diff (1-2)	—	-3.314	13.774	—	—
	Chocolate	Control	51	13.516	6.136	3.410	30.540
		ADHD	92	18.291	15.883	3.950	114.88
		Diff (1-2)	—	-4.775	13.272	—	—

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Cinnamon	Control	51	8.315	6.347	1.490	38.800
		ADHD	92	12.609	15.262	0.490	111.50
		Diff (1-2)	—	-4.294	12.830	—	—
	Clam	Control	51	36.890	57.603	7.450	400.00
		ADHD	92	31.984	32.189	3.640	213.69
		Diff (1-2)	—	4.906	42.957	—	—
	Codfish	Control	51	27.484	34.270	6.170	203.91
		ADHD	92	22.764	36.339	1.270	342.06
		Diff (1-2)	—	4.720	35.619	—	—
	Coffee	Control	51	34.003	55.076	6.730	400.00
		ADHD	92	33.999	41.052	3.850	219.51
		Diff (1-2)	—	0.004	46.512	—	—
	Cola_Nut	Control	51	11.928	5.390	2.630	27.260
		ADHD	92	13.835	7.148	1.830	33.990
		Diff (1-2)	—	-1.907	6.579	—	—
	Corn	Control	51	7.351	5.171	2.080	27.010
		ADHD	92	19.225	32.677	0.530	188.39
		Diff (1-2)	—	-11.874	26.431	—	—
	Cottage_Ch_	Control	51	83.139	107.442	2.120	400.00
		ADHD	92	105.535	116.561	2.210	400.00
		Diff (1-2)	—	-22.396	113.411	—	—
	Cow_Milk	Control	51	65.188	90.937	1.800	400.00
		ADHD	92	83.361	88.266	1.100	400.00
		Diff (1-2)	—	-18.173	89.222	—	—
	Crab	Control	51	30.367	21.673	3.770	114.37
		ADHD	92	25.751	27.638	3.460	213.52
		Diff (1-2)	—	4.616	25.682	—	—
	Cucumber	Control	51	6.560	5.111	1.270	26.500
		ADHD	92	21.432	56.304	1.600	353.35
		Diff (1-2)	—	-14.873	45.334	—	—
	Egg	Control	51	70.569	97.119	5.110	400.00
		ADHD	92	109.862	120.655	3.070	400.00
		Diff (1-2)	—	-39.293	112.872	—	—
	Eggplant	Control	51	3.980	4.120	0.100	26.500

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		ADHD	92	9.399	23.449	0.210	155.52
		Diff (1-2)	—	-5.419	18.997	—	—
	Garlic	Control	51	8.615	4.650	0.100	23.410
		ADHD	92	13.875	12.722	2.830	65.170
		Diff (1-2)	—	-5.259	10.589	—	—
	Goat_Milk	Control	51	12.632	28.034	0.100	149.14
		ADHD	92	12.332	15.717	0.100	88.430
		Diff (1-2)	—	0.300	20.931	—	—
	Grape	Control	51	13.069	6.123	2.350	44.190
		ADHD	92	14.913	9.885	5.840	63.770
		Diff (1-2)	—	-1.844	8.738	—	—
	Grapefruit	Control	51	3.128	1.808	0.100	8.040
		ADHD	92	5.639	8.356	0.630	54.840
		Diff (1-2)	—	-2.511	6.799	—	—
	Green_Pea	Control	51	5.735	5.596	0.100	27.010
		ADHD	92	9.805	13.113	0.100	65.560
		Diff (1-2)	—	-4.070	11.049	—	—
	Green_Pepper	Control	51	4.001	2.220	1.150	11.460
		ADHD	92	6.259	8.363	0.320	54.580
		Diff (1-2)	—	-2.258	6.848	—	—
	Halibut	Control	51	11.307	6.425	0.850	35.370
		ADHD	92	12.371	15.568	2.650	98.220
		Diff (1-2)	—	-1.064	13.079	—	—
	Honey	Control	51	8.233	3.766	0.530	22.800
		ADHD	92	10.606	7.379	2.570	48.440
		Diff (1-2)	—	-2.373	6.338	—	—
	Lemon	Control	51	2.559	1.273	0.100	6.270
		ADHD	92	3.103	2.731	0.110	17.390
		Diff (1-2)	—	-0.543	2.321	—	—
	Lettuce	Control	51	9.676	5.181	3.730	27.090
		ADHD	92	17.674	23.129	2.720	121.69
		Diff (1-2)	—	-7.998	18.836	—	—
	Lima_Bean	Control	51	4.955	4.159	0.100	25.010
		ADHD	92	6.487	5.863	0.880	32.820

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		Diff (1-2)	—	-1.531	5.321	—	—
	Lobster	Control	51	7.009	4.347	1.180	23.980
		ADHD	92	9.874	16.089	1.160	118.16
		Diff (1-2)	—	-2.865	13.182	—	—
	Malt	Control	51	13.807	7.087	1.920	31.200
		ADHD	92	17.057	12.192	2.810	98.820
		Diff (1-2)	—	-3.250	10.665	—	—
	Millet	Control	51	3.882	6.158	0.100	45.890
		ADHD	92	3.253	1.777	0.210	8.500
		Diff (1-2)	—	0.629	3.935	—	—
	Mushroom	Control	51	22.699	22.937	1.490	106.22
		ADHD	92	29.321	27.815	2.700	131.64
		Diff (1-2)	—	-6.622	26.189	—	—
	Mustard	Control	51	5.399	2.689	0.910	11.660
		ADHD	92	7.816	6.990	0.120	46.630
		Diff (1-2)	—	-2.417	5.839	—	—
	Oat	Control	51	11.120	9.562	0.100	49.540
		ADHD	92	24.648	35.440	1.210	228.72
		Diff (1-2)	—	-13.529	29.035	—	—
	Olive	Control	51	15.480	19.299	3.050	111.23
		ADHD	92	30.122	61.991	3.280	400.00
		Diff (1-2)	—	-14.642	51.110	—	—
	Onion	Control	51	9.739	10.258	1.170	70.460
		ADHD	92	27.033	65.323	1.800	400.00
		Diff (1-2)	—	-17.293	52.833	—	—
	Orange	Control	51	23.728	28.880	4.170	149.43
		ADHD	92	47.326	75.864	3.210	400.00
		Diff (1-2)	—	-23.598	63.326	—	—
	Oyster	Control	51	44.126	34.721	9.620	168.93
		ADHD	92	51.409	66.452	5.780	400.00
		Diff (1-2)	—	-7.283	57.249	—	—
	Parsley	Control	51	21.958	46.256	5.990	342.33
		ADHD	92	19.005	15.536	5.350	76.780
		Diff (1-2)	—	2.953	30.241	—	—

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Peach	Control	51	6.506	7.491	0.100	34.650
		ADHD	92	23.107	66.936	0.210	400.00
		Diff (1-2)	—	-16.601	53.958	—	—
	Peanut	Control	51	5.445	4.274	0.100	24.230
		ADHD	92	6.205	10.593	0.210	85.730
		Diff (1-2)	—	-0.759	8.882	—	—
	Pineapple	Control	51	8.460	18.977	0.100	122.86
		ADHD	92	17.163	45.432	0.640	400.00
		Diff (1-2)	—	-8.703	38.208	—	—
	Pinto_Bean	Control	51	9.830	9.653	0.210	47.920
		ADHD	92	17.675	26.357	1.480	139.23
		Diff (1-2)	—	-7.845	21.940	—	—
	Pork	Control	51	15.096	8.745	4.800	44.260
		ADHD	92	14.611	15.147	2.000	111.43
		Diff (1-2)	—	0.485	13.236	—	—
	Potato	Control	51	8.664	2.240	4.900	14.010
		ADHD	92	13.359	17.362	3.120	118.58
		Diff (1-2)	—	-4.696	14.012	—	—
	Rice	Control	51	18.985	14.969	4.900	73.100
		ADHD	92	26.191	27.478	3.730	189.04
		Diff (1-2)	—	-7.206	23.807	—	—
	Rye	Control	51	4.185	2.647	0.230	17.990
		ADHD	92	6.593	7.838	0.100	54.320
		Diff (1-2)	—	-2.409	6.491	—	—
	Safflower	Control	51	6.557	5.364	1.620	36.650
		ADHD	92	10.126	19.454	1.180	178.13
		Diff (1-2)	—	-3.569	15.951	—	—
	Salmon	Control	51	13.155	11.632	3.480	68.370
		ADHD	92	15.231	41.326	2.200	400.00
		Diff (1-2)	—	-2.076	33.914	—	—
	Sardine	Control	51	29.733	14.098	12.950	76.730
		ADHD	92	30.082	19.193	6.570	112.43
		Diff (1-2)	—	-0.349	17.556	—	—
	Scallop	Control	51	53.504	22.302	15.620	107.71

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Sesame	ADHD	92	54.016	42.424	10.920	283.14
		Diff (1-2)	—	-0.511	36.578	—	—
		Control	51	91.740	91.167	6.640	400.00
	Shrimp	ADHD	92	91.778	112.395	4.760	400.00
		Diff (1-2)	—	-0.039	105.358	—	—
		Control	51	31.906	31.340	5.360	151.14
	Sole	ADHD	92	17.770	18.162	2.540	103.29
		Diff (1-2)	—	14.136	23.689	—	—
		Control	51	5.010	3.858	0.230	29.090
	Soybean	ADHD	92	5.034	3.886	0.210	32.900
		Diff (1-2)	—	-0.024	3.876	—	—
		Control	51	14.277	10.254	4.150	51.570
	Spinach	ADHD	92	21.073	24.100	2.750	115.37
		Diff (1-2)	—	-6.796	20.301	—	—
		Control	51	20.913	15.581	3.290	66.870
	Squashes	ADHD	92	19.725	19.295	5.350	124.21
		Diff (1-2)	—	1.188	18.065	—	—
		Control	51	5.696	2.997	2.050	13.840
	Strawberry	ADHD	92	13.463	27.966	0.620	210.70
		Diff (1-2)	—	-7.767	22.538	—	—
		Control	51	4.585	4.755	0.110	27.900
	String_Bean	ADHD	92	15.284	50.795	0.100	331.49
		Diff (1-2)	—	-10.699	40.905	—	—
		Control	51	34.495	21.114	12.540	94.210
	Sunflower_Sd	ADHD	92	36.872	27.247	7.860	184.36
		Diff (1-2)	—	-2.377	25.243	—	—
		Control	51	7.402	4.309	1.490	21.170
	Sweet_Pot_	ADHD	92	8.995	6.411	1.790	33.590
		Diff (1-2)	—	-1.593	5.754	—	—
		Control	51	13.319	8.693	4.460	53.650
	Swiss_Ch_	ADHD	92	15.735	14.552	2.520	94.140
		Diff (1-2)	—	-2.416	12.785	—	—
		Control	51	37.893	78.801	1.490	400.00
		ADHD	92	38.949	59.178	1.470	386.11

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
Tea		Diff (1-2)	—	-1.056	66.799	—	—
		Control	51	19.459	7.609	8.930	38.010
		ADHD	92	26.526	16.518	9.070	118.42
Tobacco		Diff (1-2)	—	-7.066	14.022	—	—
		Control	51	28.550	13.486	7.880	65.660
		ADHD	92	38.636	32.339	9.270	196.29
Tomato		Diff (1-2)	—	-10.085	27.193	—	—
		Control	51	7.412	5.926	1.920	30.760
		ADHD	92	26.083	69.738	0.990	400.00
Trout		Diff (1-2)	—	-18.670	56.136	—	—
		Control	51	15.255	16.016	3.000	93.130
		ADHD	92	15.496	41.328	2.570	400.00
Tuna		Diff (1-2)	—	-0.241	34.544	—	—
		Control	51	8.130	6.362	3.050	33.880
		ADHD	92	5.577	5.696	0.850	46.670
Turkey		Diff (1-2)	—	2.553	5.941	—	—
		Control	51	11.859	5.301	4.490	28.920
		ADHD	92	13.482	12.293	2.670	87.480
Walnut_Bl		Diff (1-2)	—	-1.623	10.368	—	—
		Control	51	19.796	13.830	5.670	79.530
		ADHD	92	24.730	20.952	4.530	99.510
Wheat		Diff (1-2)	—	-4.934	18.739	—	—
		Control	51	14.031	16.565	3.200	116.33
		ADHD	92	26.488	47.247	2.110	400.00
Yeast_Baker		Diff (1-2)	—	-12.457	39.217	—	—
		Control	51	6.905	4.322	2.230	24.960
		ADHD	92	14.363	32.026	1.360	230.36
Yeast_Brewer		Diff (1-2)	—	-7.458	25.857	—	—
		Control	51	9.946	8.060	1.490	37.540
		ADHD	92	17.777	29.355	1.790	177.52
Yogurt		Diff (1-2)	—	-7.830	24.066	—	—
		Control	51	19.256	34.792	0.100	223.20
		ADHD	92	16.536	16.856	1.380	101.94
		Diff (1-2)	—	2.720	24.751	—	—

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
MALE	Almond	Control	81	4.955	2.457	1.600	14.840
		ADHD	45	7.873	9.885	0.750	50.700
		Diff (1-2)	—	-2.917	6.210	—	—
	Amer__Cheese	Control	81	33.623	47.729	1.710	234.20
		ADHD	45	41.001	45.930	1.440	238.18
		Diff (1-2)	—	-7.378	47.099	—	—
	Apple	Control	81	4.768	4.226	0.990	30.110
		ADHD	45	14.584	59.891	0.100	400.00
		Diff (1-2)	—	-9.816	35.837	—	—
	Avocado	Control	81	2.950	2.086	0.200	15.510
		ADHD	45	2.762	2.450	0.360	14.930
		Diff (1-2)	—	0.188	2.222	—	—
	Banana	Control	81	4.016	5.531	0.800	48.430
		ADHD	45	5.730	10.042	0.620	57.670
		Diff (1-2)	—	-1.714	7.451	—	—
	Barley	Control	81	9.009	35.683	1.080	324.19
		ADHD	45	9.982	21.334	0.240	142.39
		Diff (1-2)	—	-0.974	31.352	—	—
	Beef	Control	81	10.820	19.739	2.370	162.33
		ADHD	45	8.968	10.461	1.180	64.970
		Diff (1-2)	—	1.853	17.035	—	—
	Blueberry	Control	81	3.790	2.258	0.880	12.560
		ADHD	45	3.964	4.805	0.100	33.490
		Diff (1-2)	—	-0.174	3.388	—	—
	Broccoli	Control	81	7.175	5.133	2.100	30.730
		ADHD	45	10.317	13.071	2.140	84.710
		Diff (1-2)	—	-3.142	8.810	—	—
	Buck_Wheat	Control	81	5.548	3.014	1.670	23.700
		ADHD	45	6.486	4.736	0.610	25.060
		Diff (1-2)	—	-0.938	3.718	—	—
	Butter	Control	81	16.652	19.178	1.550	93.140
		ADHD	45	33.273	31.469	0.120	117.94
		Diff (1-2)	—	-16.621	24.263	—	—
	Cabbage	Control	81	5.952	10.811	0.980	94.740

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		ADHD	45	13.046	34.923	1.780	230.48
		Diff (1-2)	—	-7.094	22.543	—	—
	Cane_Sugar	Control	81	23.047	28.025	3.900	170.78
		ADHD	45	30.327	54.839	3.250	360.99
		Diff (1-2)	—	-7.280	39.672	—	—
	Cantaloupe	Control	81	5.879	4.368	1.960	29.570
		ADHD	45	20.911	61.974	0.360	400.00
		Diff (1-2)	—	-15.032	37.083	—	—
	Carrot	Control	81	4.016	3.787	1.180	27.680
		ADHD	45	14.156	41.503	0.480	224.76
		Diff (1-2)	—	-10.140	24.909	—	—
	Cashew	Control	81	9.724	11.603	1.020	59.200
		ADHD	45	9.675	13.714	0.100	78.260
		Diff (1-2)	—	0.050	12.393	—	—
	Cauliflower	Control	81	4.865	3.698	1.510	24.160
		ADHD	45	11.647	30.698	1.800	205.18
		Diff (1-2)	—	-6.781	18.526	—	—
	Celery	Control	81	8.967	5.476	2.950	34.790
		ADHD	45	15.251	26.609	2.990	155.83
		Diff (1-2)	—	-6.283	16.450	—	—
	Cheddar_Ch_	Control	81	26.696	45.931	1.690	283.73
		ADHD	45	43.425	70.376	0.600	355.84
		Diff (1-2)	—	-16.729	55.843	—	—
	Chicken	Control	81	16.053	12.550	2.940	76.880
		ADHD	45	14.770	10.368	4.290	45.240
		Diff (1-2)	—	1.283	11.822	—	—
	Chili_Pepper	Control	81	7.835	5.613	1.570	38.040
		ADHD	45	10.254	11.413	1.800	54.050
		Diff (1-2)	—	-2.419	8.157	—	—
	Chocolate	Control	81	16.623	11.019	3.010	59.470
		ADHD	45	16.970	12.146	4.580	49.160
		Diff (1-2)	—	-0.347	11.432	—	—
	Cinnamon	Control	81	9.850	7.037	1.640	40.480
		ADHD	45	9.821	6.585	1.810	32.670

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		Diff (1-2)	—	0.029	6.880	—	—
	Clam	Control	81	33.566	20.277	3.190	98.480
		ADHD	45	26.397	25.701	2.780	132.61
		Diff (1-2)	—	7.169	22.353	—	—
	Codfish	Control	81	25.075	33.649	6.490	277.17
		ADHD	45	29.288	58.990	3.900	400.00
		Diff (1-2)	—	-4.212	44.331	—	—
	Coffee	Control	81	30.318	43.408	4.320	356.95
		ADHD	45	33.387	40.314	3.970	237.41
		Diff (1-2)	—	-3.069	42.336	—	—
	Cola_Nut	Control	81	15.243	8.049	4.080	38.820
		ADHD	45	14.139	8.193	4.790	36.670
		Diff (1-2)	—	1.105	8.101	—	—
	Corn	Control	81	9.923	12.544	2.360	95.510
		ADHD	45	11.735	10.231	2.160	53.180
		Diff (1-2)	—	-1.812	11.775	—	—
	Cottage_Ch_	Control	81	76.631	102.972	1.210	400.00
		ADHD	45	123.448	125.633	2.290	400.00
		Diff (1-2)	—	-46.817	111.542	—	—
	Cow_Milk	Control	81	60.822	83.166	1.770	400.00
		ADHD	45	93.019	93.827	2.040	400.00
		Diff (1-2)	—	-32.197	87.098	—	—
	Crab	Control	81	32.448	37.288	4.770	299.11
		ADHD	45	25.297	20.928	5.170	126.85
		Diff (1-2)	—	7.151	32.442	—	—
	Cucumber	Control	81	8.752	8.584	1.880	61.860
		ADHD	45	27.909	68.764	2.530	400.00
		Diff (1-2)	—	-19.157	41.538	—	—
	Egg	Control	81	62.505	92.408	3.780	400.00
		ADHD	45	120.210	128.233	2.650	400.00
		Diff (1-2)	—	-57.705	106.508	—	—
	Eggplant	Control	81	5.045	5.910	1.370	48.790
		ADHD	45	8.238	16.933	1.180	90.580
		Diff (1-2)	—	-3.193	11.148	—	—

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Garlic	Control	81	11.918	9.606	3.040	52.160
		ADHD	45	15.901	20.797	0.230	116.36
		Diff (1-2)	—	-3.983	14.595	—	—
	Goat_Milk	Control	81	11.177	16.325	0.500	96.690
		ADHD	45	14.943	17.272	0.600	90.320
		Diff (1-2)	—	-3.767	16.667	—	—
	Grape	Control	81	15.645	5.750	8.060	47.250
		ADHD	45	14.363	11.715	5.650	65.050
		Diff (1-2)	—	1.282	8.368	—	—
	Grapefruit	Control	81	4.255	3.961	0.810	32.910
		ADHD	45	11.103	44.847	0.120	304.26
		Diff (1-2)	—	-6.848	26.903	—	—
	Green_Pea	Control	81	7.020	6.334	1.020	35.200
		ADHD	45	12.287	18.687	1.320	87.570
		Diff (1-2)	—	-5.267	12.239	—	—
	Green_Pepper	Control	81	4.715	3.714	1.660	32.330
		ADHD	45	7.840	13.389	1.390	76.860
		Diff (1-2)	—	-3.124	8.515	—	—
	Halibut	Control	81	14.289	15.877	4.410	135.74
		ADHD	45	9.302	6.073	1.440	30.940
		Diff (1-2)	—	4.987	13.256	—	—
	Honey	Control	81	10.351	5.111	2.730	29.820
		ADHD	45	11.212	10.312	4.380	64.910
		Diff (1-2)	—	-0.861	7.388	—	—
	Lemon	Control	81	3.051	2.461	0.200	20.660
		ADHD	45	3.049	4.172	0.100	28.260
		Diff (1-2)	—	0.002	3.175	—	—
	Lettuce	Control	81	12.815	7.663	3.730	39.970
		ADHD	45	22.311	39.582	4.060	245.35
		Diff (1-2)	—	-9.497	24.369	—	—
	Lima_Bean	Control	81	6.294	5.249	1.550	35.110
		ADHD	45	8.261	10.797	1.440	64.200
		Diff (1-2)	—	-1.967	7.690	—	—
	Lobster	Control	81	9.454	6.640	1.310	41.980

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Malt	ADHD	45	8.113	5.548	1.920	27.520
		Diff (1-2)	—	1.341	6.274	—	—
		Control	81	15.173	8.267	2.550	51.290
	Millet	ADHD	45	16.614	8.000	4.680	42.770
		Diff (1-2)	—	-1.441	8.173	—	—
		Control	81	4.065	4.304	1.440	40.360
	Mushroom	ADHD	45	3.069	1.661	1.090	10.770
		Diff (1-2)	—	0.996	3.596	—	—
		Control	81	27.234	27.375	2.820	118.76
	Mustard	ADHD	45	31.311	37.013	2.170	165.04
		Diff (1-2)	—	-4.076	31.139	—	—
		Control	81	6.992	4.300	1.950	30.770
	Oat	ADHD	45	9.070	6.716	2.350	27.670
		Diff (1-2)	—	-2.078	5.285	—	—
		Control	81	18.201	20.144	1.180	88.430
	Olive	ADHD	45	24.258	44.906	0.240	289.44
		Diff (1-2)	—	-6.057	31.263	—	—
		Control	81	17.589	31.696	3.550	281.30
	Onion	ADHD	45	32.810	71.753	3.760	400.00
		Diff (1-2)	—	-15.221	49.750	—	—
		Control	81	13.450	23.822	2.270	210.93
	Orange	ADHD	45	21.552	43.799	2.260	288.97
		Diff (1-2)	—	-8.101	32.355	—	—
		Control	81	26.423	37.325	2.820	314.77
	Oyster	ADHD	45	52.303	86.202	6.980	400.00
		Diff (1-2)	—	-25.880	59.460	—	—
		Control	81	49.593	42.026	7.660	250.39
	Parsley	ADHD	45	51.189	40.645	2.990	145.00
		Diff (1-2)	—	-1.596	41.542	—	—
		Control	81	17.745	7.651	5.300	59.620
	Peach	ADHD	45	18.454	19.955	7.580	120.17
		Diff (1-2)	—	-0.710	13.381	—	—
		Control	81	10.413	10.155	1.910	53.130
		ADHD	45	29.666	69.796	1.710	390.11

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		Diff (1-2)	—	-19.253	42.369	—	—
	Peanut	Control	81	5.730	9.912	1.220	89.270
		ADHD	45	7.088	9.960	0.210	61.860
		Diff (1-2)	—	-1.358	9.929	—	—
	Pineapple	Control	81	12.433	44.326	1.660	400.00
		ADHD	45	24.370	71.590	1.680	400.00
		Diff (1-2)	—	-11.937	55.554	—	—
	Pinto_Bean	Control	81	9.371	6.088	2.000	33.950
		ADHD	45	18.432	47.574	0.840	326.50
		Diff (1-2)	—	-9.061	28.758	—	—
	Pork	Control	81	16.675	14.641	4.200	89.420
		ADHD	45	13.623	14.044	3.990	95.400
		Diff (1-2)	—	3.052	14.432	—	—
	Potato	Control	81	12.243	8.339	4.920	75.770
		ADHD	45	19.614	42.471	3.490	238.41
		Diff (1-2)	—	-7.372	26.171	—	—
	Rice	Control	81	24.230	16.518	4.810	79.620
		ADHD	45	26.083	23.639	4.400	104.65
		Diff (1-2)	—	-1.852	19.347	—	—
	Rye	Control	81	5.122	3.376	1.570	23.490
		ADHD	45	8.482	9.991	0.100	42.730
		Diff (1-2)	—	-3.360	6.540	—	—
	Safflower	Control	81	7.553	4.019	2.450	27.490
		ADHD	45	11.774	16.638	2.070	99.350
		Diff (1-2)	—	-4.221	10.423	—	—
	Salmon	Control	81	16.308	15.973	0.100	136.52
		ADHD	45	11.480	12.387	0.370	66.780
		Diff (1-2)	—	4.827	14.800	—	—
	Sardine	Control	81	33.100	14.613	7.840	87.490
		ADHD	45	34.280	34.250	3.530	223.07
		Diff (1-2)	—	-1.181	23.537	—	—
	Scallop	Control	81	50.308	23.098	11.060	116.33
		ADHD	45	55.161	42.723	6.980	162.20
		Diff (1-2)	—	-4.854	31.494	—	—

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
	Sesame	Control	81	73.448	87.621	3.430	400.00
		ADHD	45	98.254	111.756	2.880	400.00
		Diff (1-2)	—	-24.806	96.876	—	—
	Shrimp	Control	81	34.185	40.052	2.930	272.28
		ADHD	45	19.372	14.056	4.160	57.840
		Diff (1-2)	—	14.812	33.242	—	—
	Sole	Control	81	5.290	2.521	2.240	20.370
		ADHD	45	4.138	1.985	0.370	10.540
		Diff (1-2)	—	1.152	2.344	—	—
	Soybean	Control	81	16.814	12.312	3.480	81.380
		ADHD	45	19.752	19.790	4.900	110.99
		Diff (1-2)	—	-2.938	15.387	—	—
	Spinach	Control	81	14.620	6.503	5.380	40.130
		ADHD	45	22.821	33.693	4.590	232.80
		Diff (1-2)	—	-8.201	20.739	—	—
	Squashes	Control	81	7.200	4.790	2.260	24.680
		ADHD	45	19.699	52.168	2.460	332.85
		Diff (1-2)	—	-12.499	31.313	—	—
	Strawberry	Control	81	5.072	4.417	1.000	29.160
		ADHD	45	15.664	59.370	0.730	400.00
		Diff (1-2)	—	-10.591	35.543	—	—
	String_Bean	Control	81	37.257	22.322	7.890	146.17
		ADHD	45	45.077	59.336	7.050	400.00
		Diff (1-2)	—	-7.820	39.633	—	—
	Sunflower_Sd	Control	81	8.566	5.303	2.450	31.260
		ADHD	45	11.785	17.504	3.250	119.64
		Diff (1-2)	—	-3.220	11.264	—	—
	Sweet_Pot_	Control	81	17.535	13.698	4.100	74.660
		ADHD	45	16.360	17.323	3.100	91.230
		Diff (1-2)	—	1.175	15.084	—	—
	Swiss_Ch_	Control	81	35.608	58.963	2.010	299.50
		ADHD	45	54.448	81.068	2.030	357.47
		Diff (1-2)	—	-18.840	67.638	—	—
	Tea	Control	81	23.966	9.868	7.620	46.400

Sex	Food	Diagnosis	N	ELISA Score			
				Mean	SD	Min	Max
		ADHD	45	28.329	23.408	8.760	159.87
		Diff (1-2)	—	-4.363	16.039	—	—
	Tobacco	Control	81	36.231	21.642	8.830	125.93
		ADHD	45	42.184	34.190	8.580	211.54
		Diff (1-2)	—	-5.953	26.776	—	—
	Tomato	Control	81	9.199	6.995	2.320	40.930
		ADHD	45	30.259	70.799	2.530	387.96
		Diff (1-2)	—	-21.059	42.546	—	—
	Trout	Control	81	14.686	9.991	3.220	83.960
		ADHD	45	13.454	14.761	3.330	85.230
		Diff (1-2)	—	1.233	11.904	—	—
	Tuna	Control	81	8.305	6.512	2.110	39.020
		ADHD	45	12.132	25.126	2.120	122.92
		Diff (1-2)	—	-3.827	15.855	—	—
	Turkey	Control	81	14.012	11.117	4.080	65.180
		ADHD	45	13.009	8.626	4.040	36.080
		Diff (1-2)	—	1.003	10.302	—	—
	Walnut_Bl	Control	81	20.821	10.403	5.680	58.470
		ADHD	45	28.055	32.043	7.160	149.93
		Diff (1-2)	—	-7.234	20.836	—	—
	Wheat	Control	81	13.359	10.034	3.240	71.930
		ADHD	45	43.230	74.407	4.300	400.00
		Diff (1-2)	—	-29.871	45.050	—	—
	Yeast_Baker	Control	81	12.471	20.370	2.070	123.35
		ADHD	45	12.101	16.201	1.390	77.180
		Diff (1-2)	—	0.371	18.995	—	—
	Yeast_Brewer	Control	81	15.903	21.144	2.640	130.89
		ADHD	45	18.229	26.775	2.030	121.54
		Diff (1-2)	—	-2.326	23.298	—	—
	Yogurt	Control	81	15.650	16.294	3.000	73.200
		ADHD	45	18.662	17.105	1.280	76.030
		Diff (1-2)	—	-3.012	16.587	—	—

Table 3

Upper Quantiles of ELISA Signal Scores among Control Subjects as Candidates for Test Cutpoints in Determining "Positive" or "Negative"

Top 26 Foods Ranked by Descending order of Discriminatory Ability using Permutation Test
ADD/ADHD Subjects vs. Controls

<i>Food Ranking</i>	<i>Food</i>	<i>Sex</i>	<i>Cutpoint</i>	
			<i>90th percentile</i>	<i>95th percentile</i>
1	Cantaloupe	FEMALE	7.774	9.885
		MALE	10.247	14.980
2	Wheat	FEMALE	22.676	40.002
		MALE	23.891	30.922
3	Tomato	FEMALE	12.757	20.137
		MALE	16.321	23.825
4	Cucumber	FEMALE	12.269	17.960
		MALE	16.150	22.827
5	Squashes	FEMALE	10.267	11.963
		MALE	13.574	18.485
6	Almond	FEMALE	6.392	7.269
		MALE	8.212	10.337
7	Egg	FEMALE	188.62	321.65
		MALE	145.49	298.13
8	Cauliflower	FEMALE	7.594	8.954
		MALE	7.799	12.288
9	Pinto_Bean	FEMALE	20.059	32.885
		MALE	17.813	21.226
10	Broccoli	FEMALE	9.069	12.037
		MALE	13.200	18.126
11	Orange	FEMALE	59.347	90.581
		MALE	47.525	68.415
12	Butter	FEMALE	40.002	67.588
		MALE	43.433	60.984
13	Corn	FEMALE	13.161	19.357
		MALE	17.173	28.120
14	Lettuce	FEMALE	16.940	21.280
		MALE	23.239	30.306
15	Rye	FEMALE	6.198	8.944

<i>Food Ranking</i>	<i>Food</i>	<i>Sex</i>	<i>Cutpoint</i>	
			<i>90th percentile</i>	<i>95th percentile</i>
16	Peach	MALE	8.399	11.978
		FEMALE	16.434	25.262
17	Green_Pea	MALE	22.310	32.987
		FEMALE	13.221	18.595
18	Carrot	MALE	14.660	21.095
		FEMALE	5.055	6.010
19	Tea	MALE	6.600	9.684
		FEMALE	29.837	33.245
20	Mustard	MALE	37.540	42.112
		FEMALE	9.607	10.799
21	Strawberry	MALE	11.762	14.581
		FEMALE	7.900	14.127
22	Celery	MALE	9.448	14.125
		FEMALE	14.210	19.974
23	Green_Pepper	MALE	15.168	19.728
		FEMALE	6.912	8.275
24	Garlic	MALE	7.092	9.348
		FEMALE	15.016	17.352
25	Oat	MALE	24.205	34.240
		FEMALE	22.743	27.596
26	Onion	MALE	51.597	63.749
		FEMALE	18.635	26.190
27	Banana	MALE	26.971	32.220
		FEMALE	5.276	8.990
28	Eggplant	MALE	6.632	10.083
		FEMALE	7.193	11.328
29	Cabbage	MALE	7.631	12.926
		FEMALE	8.484	15.011
30	Safflower	MALE	9.596	16.710
		FEMALE	10.328	16.067
31	Olive	MALE	11.820	14.861
		FEMALE	27.849	55.153
32	Cottage_Ch_	MALE	28.220	44.995
		FEMALE	244.09	354.86

<i>Food Ranking</i>	<i>Food</i>	<i>Sex</i>	<i>Cutpoint</i>	
			<i>90th percentile</i>	<i>95th percentile</i>
33	Grapefruit	MALE	225.78	341.05
		FEMALE	5.731	6.772
34	Walnut_Black	MALE	7.127	10.165
		FEMALE	38.596	49.729
35	Cow_Milk	MALE	35.760	41.973
		FEMALE	155.08	281.94
36	Soybean	MALE	149.48	244.73
		FEMALE	27.522	38.061
37	Chili_Pepper	MALE	31.249	41.905
		FEMALE	15.566	21.521
		MALE	14.808	18.731

Table 4

ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
BRH1227838	10
BRH1227840	2
BRH1227841	13
BRH1227842	2
BRH1227843	6
BRH1227844	13
BRH1227846	13
BRH1243697	0
BRH1243698	8
BRH1243699	2
KH16-13285	0
KH16-14189	1
KH16-14190	1
KH16-14586	0
KH16-14587	0
KH16-16347	10
KH16-21226	3
KH17-4120	3
KH17-4757	1
KH17-5055	0
KH17-5056	0
DLS17-012889	3
KH17-00912	9
KH17-1238	20
KH17-2452	10
KH17-2453	20
KH17-2454	0
KH17-2676	4
KH17-4038	1
KH17-4039	10
DLS16-69541	0
160904AAC0045	3
160904AAC0046	20
160904AAC0047	5
160904AAC0048	2
160904AAC0049	10
160904AAC0050	0
160904AAC0051	5

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
BRH1244994	0
BRH1244995	0
BRH1244996	1
BRH1244997	1
BRH1244998	2
BRH1244999	0
BRH1245000	2
BRH1245001	0
BRH1245002	1
BRH1245004	0
BRH1245007	0
BRH1245008	1
BRH1245009	1
BRH1245010	5
BRH1245011	2
BRH1245014	0
BRH1245015	1
BRH1245018	1
BRH1245019	1
BRH1245022	11
BRH1245023	1
BRH1245024	2
BRH1245026	2
BRH1245029	1
BRH1245030	0
BRH1245031	2
BRH1245032	1
BRH1245033	2
BRH1245035	0
BRH1245037	0
BRH1245038	0
BRH1245039	9
BRH1245040	5
BRH1245041	3
BRH1267328	12
BRH1267329	2
BRH1267330	0
BRH1267332	0

WO 2018/109746

PCT/IB2017/058023

ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
160904AAC0021	9
160904AAC0023	1
160904AAC0025	10
160904AAC0026	6
160904AAC0028	3
160904AAC0037	4
160904AAC0038	0
160904AAC0039	8
160904AAC0040	9
160904AAC0042	3
160904AAC0012	1
160904AAC0013	1
160904AAC0014	2
160904AAC0015	7
160904AAC0016	20
160904AAC0017	1
160904AAC0018	0
160904AAC0019	4
160904AAC0020	5
160880AAB0022	2
160880AAB0024	6
160880AAB0025	0
160880AAB0026	18
160880AAB0028	11
160880AAB0029	0
160904AAC0009	0
160904AAC0010	6
160904AAC0001	0
160904AAC0003	0
160904AAC0004	7
160904AAC0005	2
160880AAB0031	3
160880AAB0032	2
160880AAB0033	3
160880AAB0034	20
160880AAB0002	13
160880AAB0003	9
160880AAB0004	0
160880AAB0006	0

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
BRH1267333	0
BRH1267334	9
BRH1267335	0
BRH1267337	2
BRH1267338	0
BRH1267339	2
BRH1267340	6
BRH1267341	0
BRH1267343	5
BRH1267345	0
BRH1267346	0
BRH1267347	0
BRH1267349	1
BRH1244900	1
BRH1244901	4
BRH1244902	0
BRH1244903	0
BRH1244904	1
BRH1244905	0
BRH1244906	5
BRH1244907	0
BRH1244908	2
BRH1244909	1
BRH1244910	2
BRH1244911	0
BRH1244912	2
BRH1244913	0
BRH1244914	3
BRH1244915	0
BRH1244916	12
BRH1244917	7
BRH1244918	0
BRH1244920	3
BRH1244921	1
BRH1244922	12
BRH1244923	1
BRH1244924	1
BRH1244925	1
BRH1244926	8

ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
160880AAB0007	3
160880AAB0009	3
160880AAB0010	8
160880AAB0011	6
160880AAB0013	1
160880AAB0014	0
160880AAB0015	0
160880AAB0016	6
160880AAB0017	0
160880AAB0018	0
11782002	2
11827629	1
11828836	0
12018682	0
12035395	0
BRH1227836	4
BRH1227837	1
BRH1227839	0
BRH1227845	13
KH16-13576	3
KH16-16348	1
KH16-18260	0
KH16-18261	6
KH16-18262	0
KH16-19420	3
KH17-4040	0
KH17-4613	5
KH17-4614	5
DLS17-012888	0
KH16-19421	3
KH17-00072	0
KH17-00073	8
KH17-1237	1
160904AAC0043	7
160904AAC0044	0
160904AAC0052	3
160904AAC0022	8
160904AAC0024	6
160904AAC0027	18

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
BRH1244928	1
BRH1244929	3
BRH1244931	1
BRH1244932	2
BRH1244933	3
BRH1244934	3
BRH1244938	3
BRH1244939	0
BRH1244940	0
BRH1244941	1
BRH1244942	4
BRH1244943	2
BRH1244944	8
BRH1244945	1
BRH1244946	6
BRH1244947	0
BRH1244948	0
BRH1244949	2
BRH1244950	0
BRH1244951	0
BRH1244952	0
BRH1244953	1
BRH1244954	0
BRH1244956	19
BRH1244959	1
BRH1244960	0
BRH1244961	0
BRH1244962	1
BRH1244963	0
BRH1244964	4
BRH1244965	1
BRH1244967	0
BRH1244969	1
BRH1244970	3
BRH1244971	0
BRH1244972	1
BRH1244973	3
BRH1244974	0
BRH1244975	0

ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
160904AAC0029	1
160904AAC0030	7
160904AAC0031	1
160904AAC0032	1
160904AAC0033	6
160904AAC0034	3
160904AAC0035	16
160904AAC0036	3
160904AAC0041	4
160904AAC0011	1
160880AAB0021	1
160880AAB0023	2
160880AAB0027	4
160904AAC0008	0
160904AAC0002	3
160880AAB0030	19
160880AAB0008	3
160880AAB0012	1
11827735	2
11940652	2
12080010	3

No of Observations	137
Average Number	4.5
Median Number	3

# of Patients w/ 0 Pos Results	33
% Subjects w/ 0 pos results	24.1

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 90th Percentile
BRH1244976	0
BRH1244977	0
BRH1244979	0
BRH1244980	1
BRH1244981	1
BRH1244982	1
BRH1244983	0
BRH1244985	0
BRH1244987	0
BRH1244988	5
BRH1244991	0
BRH1244992	1
BRH1267320	1
BRH1267322	7
BRH1267323	0
BRH1267325	3

No of Observations	132
Average Number	2.0
Median Number	1

# of Patients w/ 0 Pos Results	51
% Subjects w/ 0 pos results	38.6

Table 5A

ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
BRH1227838	5
BRH1227840	1
BRH1227841	7
BRH1227842	2
BRH1227843	2
BRH1227844	5
BRH1227846	7
BRH1243697	0
BRH1243698	3
BRH1243699	0
KH16-13285	0
KH16-14189	0
KH16-14190	1
KH16-14586	0
KH16-14587	0
KH16-16347	6
KH16-21226	1
KH17-4120	2
KH17-4757	1
KH17-5055	0
KH17-5056	0
DLS17-012889	1
KH17-00912	7
KH17-1238	20
KH17-2452	7
KH17-2453	20
KH17-2454	0
KH17-2676	3
KH17-4038	0
KH17-4039	4
DLS16-69541	0
160904AAC0045	2
160904AAC0046	16
160904AAC0047	4
160904AAC0048	0
160904AAC0049	7
160904AAC0050	0
160904AAC0051	2

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
BRH1244994	0
BRH1244995	0
BRH1244996	0
BRH1244997	0
BRH1244998	1
BRH1244999	0
BRH1245000	0
BRH1245001	0
BRH1245002	1
BRH1245004	0
BRH1245007	0
BRH1245008	0
BRH1245009	0
BRH1245010	3
BRH1245011	2
BRH1245014	0
BRH1245015	0
BRH1245018	1
BRH1245019	1
BRH1245022	6
BRH1245023	1
BRH1245024	1
BRH1245026	2
BRH1245029	1
BRH1245030	0
BRH1245031	1
BRH1245032	0
BRH1245033	0
BRH1245035	0
BRH1245037	0
BRH1245038	0
BRH1245039	7
BRH1245040	3
BRH1245041	1
BRH1267328	4
BRH1267329	0
BRH1267330	0
BRH1267332	0

ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
160904AAC0021	6
160904AAC0023	0
160904AAC0025	8
160904AAC0026	5
160904AAC0028	3
160904AAC0037	0
160904AAC0038	0
160904AAC0039	3
160904AAC0040	5
160904AAC0042	1
160904AAC0012	0
160904AAC0013	1
160904AAC0014	2
160904AAC0015	5
160904AAC0016	20
160904AAC0017	0
160904AAC0018	0
160904AAC0019	2
160904AAC0020	2
160880AAB0022	0
160880AAB0024	4
160880AAB0025	0
160880AAB0026	13
160880AAB0028	10
160880AAB0029	0
160904AAC0009	0
160904AAC0010	5
160904AAC0001	0
160904AAC0003	0
160904AAC0004	3
160904AAC0005	1
160880AAB0031	1
160880AAB0032	0
160880AAB0033	0
160880AAB0034	17
160880AAB0002	6
160880AAB0003	4
160880AAB0004	0
160880AAB0006	0

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
BRH1267333	0
BRH1267334	4
BRH1267335	0
BRH1267337	1
BRH1267338	0
BRH1267339	0
BRH1267340	6
BRH1267341	0
BRH1267343	2
BRH1267345	0
BRH1267346	0
BRH1267347	0
BRH1267349	1
BRH1244900	1
BRH1244901	1
BRH1244902	0
BRH1244903	0
BRH1244904	1
BRH1244905	0
BRH1244906	3
BRH1244907	0
BRH1244908	2
BRH1244909	1
BRH1244910	0
BRH1244911	0
BRH1244912	0
BRH1244913	0
BRH1244914	1
BRH1244915	0
BRH1244916	6
BRH1244917	3
BRH1244918	0
BRH1244920	3
BRH1244921	0
BRH1244922	9
BRH1244923	0
BRH1244924	1
BRH1244925	1
BRH1244926	7

ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
160880AAB0007	3
160880AAB0009	1
160880AAB0010	4
160880AAB0011	1
160880AAB0013	1
160880AAB0014	0
160880AAB0015	0
160880AAB0016	3
160880AAB0017	0
160880AAB0018	0
11782002	1
11827629	1
11828836	0
12018682	0
12035395	0
BRH1227836	2
BRH1227837	0
BRH1227839	0
BRH1227845	4
KH16-13576	2
KH16-16348	0
KH16-18260	0
KH16-18261	3
KH16-18262	0
KH16-19420	2
KH17-4040	0
KH17-4613	5
KH17-4614	2
DLS17-012888	0
KH16-19421	3
KH17-00072	0
KH17-00073	3
KH17-1237	1
160904AAC0043	4
160904AAC0044	0
160904AAC0052	2
160904AAC0022	7
160904AAC0024	0
160904AAC0027	16

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
BRH1244928	1
BRH1244929	2
BRH1244931	1
BRH1244932	1
BRH1244933	2
BRH1244934	2
BRH1244938	1
BRH1244939	0
BRH1244940	0
BRH1244941	0
BRH1244942	3
BRH1244943	0
BRH1244944	1
BRH1244945	1
BRH1244946	4
BRH1244947	0
BRH1244948	0
BRH1244949	0
BRH1244950	0
BRH1244951	0
BRH1244952	0
BRH1244953	1
BRH1244954	0
BRH1244956	10
BRH1244959	0
BRH1244960	0
BRH1244961	0
BRH1244962	0
BRH1244963	0
BRH1244964	3
BRH1244965	0
BRH1244967	0
BRH1244969	1
BRH1244970	1
BRH1244971	0
BRH1244972	1
BRH1244973	0
BRH1244974	0
BRH1244975	0

ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
160904AAC0029	1
160904AAC0030	3
160904AAC0031	1
160904AAC0032	1
160904AAC0033	3
160904AAC0034	2
160904AAC0035	16
160904AAC0036	1
160904AAC0041	1
160904AAC0011	1
160880AAB0021	0
160880AAB0023	0
160880AAB0027	4
160904AAC0008	0
160904AAC0002	3
160880AAB0030	19
160880AAB0008	2
160880AAB0012	1
11827735	2
11940652	1
12080010	3

No of Observations	137
Average Number	2.9
Median Number	1

# of Patients w/ 0 Pos Results	49
% Subjects w/ 0 pos results	35.8

NON-ADHD POPULATION	
Sample ID	# of Positive Results Based on 95th Percentile
BRH1244976	0
BRH1244977	0
BRH1244979	0
BRH1244980	0
BRH1244981	0
BRH1244982	0
BRH1244983	0
BRH1244985	0
BRH1244987	0
BRH1244988	3
BRH1244991	0
BRH1244992	0
BRH1267320	0
BRH1267322	0
BRH1267323	0
BRH1267325	1

No of Observations	132
Average Number	1.0
Median Number	0

# of Patients w/ 0 Pos Results	78
% Subjects w/ 0 pos results	59.1

Table 5B

Summary statistics		
Variable	ADHD_90th_percentile ADHD_90th_percentile	
Sample size	137	
Lowest value	0.0000	
Highest value	20.0000	
Arithmetic mean	4.4818	
95% CI for the mean	3.6002 to 5.3633	
Median	3.0000	
95% CI for the median	2.0000 to 3.0000	
Variance	27.2221	
Standard deviation	5.2175	
Relative standard deviation	1.1642 (116.42%)	
Standard error of the mean	0.4458	
Coefficient of Skewness	1.5369 (P<0.0001)	
Coefficient of Kurtosis	1.8249 (P=0.0040)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.0000	
5	0.0000	0.0000 to 0.0000
10	0.0000	0.0000 to 0.0000
25	1.0000	0.0000 to 1.0000
75	6.2500	5.0000 to 8.9203
90	12.6000	9.5756 to 18.0000
95	18.0000	13.0000 to 20.0000
97.5	20.0000	

Table 6A

Summary statistics		
Variable	ADHD_95th_percentile	
Sample size	137	
Lowest value	0.0000	
Highest value	20.0000	
Arithmetic mean	2.9124	
95% CI for the mean	2.1586 to 3.6662	
Median	1.0000	
95% CI for the median	1.0000 to 2.0000	
Variance	19.9040	
Standard deviation	4.4614	
Relative standard deviation	1.5319 (153.19%)	
Standard error of the mean	0.3812	
Coefficient of Skewness	2.4667 (P<0.0001)	
Coefficient of Kurtosis	6.0423 (P<0.0001)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.0000	
5	0.0000	0.0000 to 0.0000
10	0.0000	0.0000 to 0.0000
25	0.0000	0.0000 to 0.0000
75	4.0000	3.0000 to 5.0000
90	7.0000	5.0000 to 16.0000
95	16.0000	7.0000 to 20.0000
97.5	19.0750	

Table 6B

Summary statistics		
Variable	Non_ADHD_90th_percentile Non_ADHD 90th percentile	
Sample size	132	
Lowest value	0.0000	
Highest value	19.0000	
Arithmetic mean	1.9621	
95% CI for the mean	1.4352 to 2.4890	
Median	1.0000	
95% CI for the median	1.0000 to 1.0000	
Variance	9.3650	
Standard deviation	3.0602	
Relative standard deviation	1.5597 (155.97%)	
Standard error of the mean	0.2664	
Coefficient of Skewness	2.7058 (P<0.0001)	
Coefficient of Kurtosis	8.9629 (P<0.0001)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.0000	
5	0.0000	0.0000 to 0.0000
10	0.0000	0.0000 to 0.0000
25	0.0000	0.0000 to 0.0000
75	2.0000	2.0000 to 3.0000
90	5.3000	4.0000 to 8.6297
95	8.9000	6.0000 to 12.0000
97.5	12.0000	

Table 7A

Summary statistics		
Variable	Non_ADHD_95th_percentile	
Sample size	132	
Lowest value	0.0000	
Highest value	10.0000	
Arithmetic mean	0.9848	
95% CI for the mean	0.6707 to 1.2990	
Median	0.0000	
95% CI for the median	0.0000 to 0.0000	
Variance	3.3280	
Standard deviation	1.8243	
Relative standard deviation	1.8524 (185.24%)	
Standard error of the mean	0.1588	
Coefficient of Skewness	2.7344 (P<0.0001)	
Coefficient of Kurtosis	8.2646 (P<0.0001)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.0000	
5	0.0000	0.0000 to 0.0000
10	0.0000	0.0000 to 0.0000
25	0.0000	0.0000 to 0.0000
75	1.0000	1.0000 to 2.0000
90	3.0000	2.0000 to 5.2594
95	5.8000	3.0000 to 7.6200
97.5	7.0000	

Table 7B

Summary statistics		
Variable	ADHD_90th_percentile_1 ADHD_90th_percentile_1	
Back-transformed after logarithmic transformation.		
Sample size	132	
Lowest value	0.1000	
Highest value	19.0000	
Geometric mean	0.6626	
95% CI for the mean	0.4997 to 0.8786	
Median	1.0000	
95% CI for the median	1.0000 to 1.0000	
Coefficient of Skewness	0.03362 (P=0.8699)	
Coefficient of Kurtosis	-1.3824 (P<0.0001)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.10000	
5	0.10000	0.10000 to 0.10000
10	0.10000	0.10000 to 0.10000
25	0.10000	0.10000 to 0.10000
75	2.0000	2.0000 to 3.0000
90	5.2811	4.0000 to 8.6159
95	8.8946	6.0000 to 12.0000
97.5	12.0000	

Table 8A

Summary statistics		
Variable	ADHD_95th_percentile_1	
Back-transformed after logarithmic transformation.		
Sample size	137	
Lowest value	0.1000	
Highest value	20.0000	
Geometric mean	0.8838	
95% CI for the mean	0.6542 to 1.1940	
Median	1.0000	
95% CI for the median	1.0000 to 2.0000	
Coefficient of Skewness	-0.09449 (P=0.6402)	
Coefficient of Kurtosis	-1.4081 (P<0.0001)	
D'Agostino-Pearson test for Normal distribution	reject Normality (P<0.0001)	
Percentiles		95% Confidence interval
2.5	0.10000	
5	0.10000	0.10000 to 0.10000
10	0.10000	0.10000 to 0.10000
25	0.10000	0.10000 to 0.10000
75	4.0000	3.0000 to 5.0000
90	7.0000	5.0000 to 16.0000
95	16.0000	7.0000 to 20.0000
97.5	19.0732	

Table 8B

Summary statistics

Variable	Non_ADHD_90th_percentile_1	
Back-transformed after logarithmic transformation.		
Sample size		132
Lowest value		0.1000
Highest value		19.0000
Geometric mean		0.6626
95% CI for the mean		0.4997 to 0.8786
Median		1.0000
95% CI for the median		1.0000 to 1.0000
Coefficient of Skewness		0.03362 (P=0.8699)
Coefficient of Kurtosis		-1.3824 (P<0.0001)
D'Agostino-Pearson test for Normal distribution		reject Normality (P<0.0001)
Percentiles		95% Confidence interval
2.5	0.10000	
5	0.10000	0.10000 to 0.10000
10	0.10000	0.10000 to 0.10000
25	0.10000	0.10000 to 0.10000
75	2.0000	2.0000 to 3.0000
90	5.2811	4.0000 to 8.6159
95	8.8946	6.0000 to 12.0000
97.5	12.0000	

Table 9A

Summary statistics		
Variable	Non_ADHD_95th_percentile_1	
Back-transformed after logarithmic transformation.		
Sample size		132
Lowest value		0.1000
Highest value		10.0000
Geometric mean		0.3258
95% CI for the mean		0.2518 to 0.4216
Median		0.10000
95% CI for the median		0.10000 to 0.10000
Coefficient of Skewness		0.6874 (P=0.0020)
Coefficient of Kurtosis		-1.1003 (P<0.0001)
D'Agostino-Pearson test for Normal distribution		reject Normality (P<0.0001)
Percentiles		95% Confidence interval
2.5	0.10000	
5	0.10000	0.10000 to 0.10000
10	0.10000	0.10000 to 0.10000
25	0.10000	0.10000 to 0.10000
75	1.0000	1.0000 to 2.0000
90	3.0000	2.0000 to 5.1635
95	5.7616	3.0000 to 7.5672
97.5	7.0000	

Table 9B

Independent samples t-test		
Sample 1		
Variable	ADHD_90th_percentile_1	
Sample 2		
Variable	Non_ADHD_90th_percentile_1	
Back-transformed after logarithmic transformation		
	Sample 1	Sample 2
Sample size	137	132
Geometric mean	1.6449	0.6626
95% CI for the mean	1.2191 to 2.2194	0.4997 to 0.8786
Variance of Logs	0.5930	0.5065
F-test for equal variances	P = 0.364	
T-test (assuming equal variances)		
Difference on Log-transformed scale		
Difference	-0.3949	
Standard Error	0.09049	
95% CI of difference	-0.5730 to -0.2167	
Test statistic t	-4.363	
Degrees of Freedom (DF)	267	
Two-tailed probability	P < 0.0001	
Back-transformed results		
Ratio of geometric means	0.4028	
95% CI of ratio	0.2673 to 0.6072	

Table 10A

Independent samples t-test		
Sample 1		
Variable	ADHD_95th_percentile_1	
Sample 2		
Variable	Non_ADHD_95th_percentile_1	
Back-transformed after logarithmic transformation.		
	Sample 1	Sample 2
Sample size	137	132
Geometric mean	0.8838	0.3258
95% CI for the mean	0.6542 to 1.1940	0.2518 to 0.4216
Variance of Logs	0.5978	0.4228
F-test for equal variances	P = 0.047	
T-test (assuming equal variances)		
Difference on Log-transformed scale		
Difference	-0.4334	
Standard Error	0.08726	
95% CI of difference	-0.6052 to -0.2616	
Test statistic t	-4.967	
Degrees of Freedom (DF)	267	
Two-tailed probability	P < 0.0001	
Back-transformed results		
Ratio of geometric means	0.3686	
95% CI of ratio	0.2482 to 0.5475	

Table 10B

Mann-Whitney test (independent samples)		
Sample 1		
Variable	ADHD_90th_percentile ADHD 90th percentile	
Sample 2		
Variable	Non_ADHD_90th_percentile Non_ADHD 90th percentile	
	Sample 1	Sample 2
Sample size	137	132
Lowest value	0.0000	0.0000
Highest value	20.0000	19.0000
Median	3.0000	1.0000
95% CI for the median	2.0000 to 3.0000	1.0000 to 1.0000
Interquartile range	1.0000 to 6.2500	0.0000 to 2.0000
Mann-Whitney test (independent samples)		
Average rank of first group	156.1679	
Average rank of second group	113.0303	
Mann-Whitney U	6142.00	
Test statistic Z (corrected for ties)	4.637	
Two-tailed probability	P < 0.0001	

Table 11A

Mann-Whitney test (independent samples)		
Sample 1		
Variable	ADHD_95th_percentile	
Sample 2		
Variable	Non_ADHD_95th_percentile	
	Sample 1	Sample 2
Sample size	137	132
Lowest value	0.0000	0.0000
Highest value	20.0000	10.0000
Median	1.0000	0.0000
95% CI for the median	1.0000 to 2.0000	0.0000 to 0.0000
Interquartile range	0.0000 to 4.0000	0.0000 to 1.0000

Mann-Whitney test (independent samples)	
Average rank of first group	156.0401
Average rank of second group	113.1629
Mann-Whitney U	6159.50
Test statistic Z (corrected for ties)	4.792
Two-tailed probability	P < 0.0001

Table 11B

ROC curve	
Variable	ADHD_Test_90th
Classification variable	ADHD_Test_90th
	Diagnosis__1_ADHD_0_Non_ADHD_
	Diagnosis(1_ADHD_0_Non_ADHD)
Sample size	269
Positive group ^a	137 (50.93%)
Negative group ^b	132 (49.07%)
^a Diagnosis__1_ADHD_0_Non_ADHD_ = 1	
^b Diagnosis__1_ADHD_0_Non_ADHD_ = 0	
Disease prevalence (%)	unknown
Area under the ROC curve (AUC)	
Area under the ROC curve (AUC)	0.660
Standard Error ^a	0.0326
95% Confidence interval ^b	0.600 to 0.717
z statistic	4.925
Significance level P (Area=0.5)	<0.0001
^a DeLong et al., 1988	
^b Binomial exact	
Youden index	
Youden index J	0.2980
95% Confidence interval ^a	0.1810 to 0.4018
Associated criterion	>2
95% Confidence interval ^a	>1 to >5
Sensitivity	53.28
Specificity	76.52
^a BC _a bootstrap confidence interval (1000 iterations; random number seed: 978).	

Table 12A

ROC curve	
Variable	ADHD_Test_95th
Classification variable	ADHD_Test_95th Diagnosis__1_ADHD_0_Non_ADHD_ Diagnosis(1_ADHD_0_Non_ADHD)_
Sample size	269
Positive group ^a	137 (50.93%)
Negative group ^a	132 (49.07%)
^a Diagnosis__1_ADHD_0_Non_ADHD_ = 1 ^b Diagnosis__1_ADHD_0_Non_ADHD_ = 0	
Disease prevalence (%)	unknown
Area under the ROC curve (AUC)	
Area under the ROC curve (AUC)	0.659
Standard Error ^a	0.0312
95% Confidence interval ^a	0.599 to 0.716
z statistic	5.117
Significance level P (Area=0.5)	<0.0001
^a DeLong et al., 1988 ^b Binomial exact	
Youden index	
Youden index J	0.2851
95% Confidence interval ^a	0.1724 to 0.3749
Associated criterion	>1
95% Confidence interval ^a	>0 to >2
Sensitivity	47.45
Specificity	81.06
^a BC _a bootstrap confidence interval (1000 iterations; random number seed: 978).	

Table 12B

Performance Metrics in Predicting ADD/ADHD Status from Number of Positive Foods**Using 90th Percentile of ELISA Signal to determine Positive**

<i>Sex</i>	<i>No. of Positive Foods as Cutoff</i>	<i>Sensitivity</i>	<i>Specificity</i>	<i>Positive Predictive Value</i>	<i>Negative Predictive Value</i>	<i>Overall Percent Agreement</i>
FEMALE	1	0.85	0.26	0.67	0.48	0.64
	2	0.75	0.40	0.70	0.48	0.63
	3	0.67	0.55	0.73	0.48	0.63
	4	0.60	0.66	0.76	0.48	0.63
	5	0.56	0.72	0.78	0.47	0.62
	6	0.52	0.76	0.79	0.46	0.60
	7	0.48	0.77	0.79	0.45	0.59
	8	0.45	0.80	0.80	0.44	0.57
	9	0.42	0.82	0.81	0.44	0.56
	10	0.38	0.85	0.82	0.43	0.55
	11	0.35	0.88	0.83	0.43	0.53
	12	0.33	0.89	0.83	0.42	0.53
	13	0.30	0.90	0.84	0.42	0.52
	14	0.28	0.91	0.84	0.41	0.50
	15	0.25	0.91	0.83	0.40	0.49
	16	0.23	0.92	0.83	0.40	0.48
	17	0.21	0.93	0.83	0.39	0.47
	18	0.20	0.94	0.85	0.39	0.46
	19	0.18	0.94	0.86	0.39	0.45
	20	0.16	0.96	0.88	0.39	0.45
	21	0.15	0.97	0.89	0.39	0.44
	22	0.13	0.97	0.90	0.38	0.43
	23	0.12	0.98	0.93	0.38	0.43
	24	0.10	1.00	1.00	0.38	0.42
	25	0.09	1.00	1.00	0.38	0.41
	26	0.08	1.00	1.00	0.38	0.41
	27	0.07	1.00	1.00	0.38	0.40
	28	0.07	1.00	1.00	0.37	0.40
	29	0.07	1.00	1.00	0.37	0.40
	30	0.07	1.00	1.00	0.37	0.40

31	0.06	1.00	1.00	0.37	0.40
32	0.05	1.00	1.00	0.37	0.39
33	0.05	1.00	1.00	0.37	0.39
34	0.04	1.00	1.00	0.37	0.38
35	0.04	1.00	1.00	0.36	0.38
36	0.02	1.00	1.00	0.36	0.37
37	0.02	1.00	1.00	0.36	0.37

Table 13A

Performance Metrics in Predicting ADD/ADHD Status from Number of Positive Foods**Using 90th Percentile of ELISA Signal to determine Positive**

Sex	No. of Positive Foods as Cutoff	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Overall Percent Agreement
MALE	1	0.89	0.25	0.40	0.80	0.48
	2	0.77	0.44	0.43	0.78	0.56
	3	0.65	0.57	0.45	0.74	0.60
	4	0.57	0.66	0.48	0.73	0.63
	5	0.50	0.73	0.52	0.73	0.65
	6	0.44	0.79	0.54	0.72	0.67
	7	0.37	0.83	0.55	0.70	0.67
	8	0.33	0.85	0.56	0.70	0.66
	9	0.29	0.87	0.55	0.69	0.66
	10	0.27	0.88	0.56	0.69	0.66
	11	0.26	0.90	0.58	0.68	0.67
	12	0.24	0.91	0.60	0.68	0.67
	13	0.23	0.92	0.60	0.68	0.67
	14	0.21	0.92	0.58	0.68	0.67
	15	0.18	0.92	0.57	0.67	0.66
	16	0.15	0.93	0.56	0.67	0.65
	17	0.12	0.94	0.50	0.66	0.65
	18	0.11	0.94	0.50	0.66	0.64
	19	0.10	0.94	0.50	0.65	0.64
	20	0.09	0.95	0.50	0.65	0.64
	21	0.08	0.96	0.50	0.65	0.64
	22	0.08	0.96	0.50	0.65	0.64
	23	0.07	0.96	0.50	0.65	0.65
	24	0.07	0.97	0.60	0.65	0.65
	25	0.07	0.98	0.67	0.65	0.65
	26	0.07	0.98	0.67	0.65	0.65
	27	0.07	0.98	0.67	0.65	0.65
	28	0.07	0.98	0.67	0.65	0.65

<i>Sex</i>	<i>No. of Positive Foods as Cutoff</i>	<i>Sensitivity</i>	<i>Specificity</i>	<i>Positive Predictive Value</i>	<i>Negative Predictive Value</i>	<i>Overall Percent Agreement</i>
	29	0.04	0.98	0.67	0.65	0.65
	30	0.04	0.98	0.67	0.65	0.65
	31	0.04	0.98	0.67	0.65	0.65
	32	0.04	1.00	0.67	0.65	0.65
	33	0.03	1.00	1.00	0.65	0.65
	34	0.00	1.00	1.00	0.65	0.65
	35	0.00	1.00	1.00	0.64	0.64
	36	0.00	1.00	0.00	0.64	0.64
	37	0.00	1.00	0.00	0.64	0.64

Table 13B

Performance Metrics in Predicting ADD/ADHD Status from Number of Positive Foods**Using 95th Percentile of ELISA Signal to determine Positive**

Sex	No. of Positive Foods as Cutoff	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Overall Percent Agreement
FEMALE	1	0.73	0.41	0.69	0.46	0.62
	2	0.61	0.58	0.72	0.45	0.60
	3	0.53	0.72	0.77	0.46	0.60
	4	0.47	0.79	0.81	0.46	0.59
	5	0.43	0.82	0.81	0.45	0.57
	6	0.39	0.84	0.81	0.43	0.55
	7	0.35	0.85	0.81	0.42	0.53
	8	0.31	0.88	0.82	0.41	0.51
	9	0.27	0.91	0.84	0.41	0.49
	10	0.23	0.93	0.85	0.40	0.48
	11	0.20	0.94	0.86	0.39	0.47
	12	0.18	0.94	0.86	0.39	0.45
	13	0.16	0.97	0.88	0.39	0.44
	14	0.14	0.97	0.89	0.38	0.44
	15	0.12	0.97	0.91	0.38	0.43
	16	0.11	1.00	1.00	0.38	0.42
	17	0.10	1.00	1.00	0.38	0.42
	18	0.09	1.00	1.00	0.38	0.41
	19	0.08	1.00	1.00	0.38	0.41
	20	0.07	1.00	1.00	0.37	0.40
	21	0.07	1.00	1.00	0.37	0.40
	22	0.07	1.00	1.00	0.37	0.40
	23	0.07	1.00	1.00	0.37	0.40
	24	0.07	1.00	1.00	0.37	0.40
	25	0.06	1.00	1.00	0.37	0.40
	26	0.06	1.00	1.00	0.37	0.39
	27	0.05	1.00	1.00	0.37	0.39
	28	0.05	1.00	1.00	0.37	0.39
	29	0.05	1.00	1.00	0.37	0.38
	30	0.04	1.00	1.00	0.37	0.38

<i>Sex</i>	<i>No. of Positive Foods as Cutoff</i>	<i>Sensitivity</i>	<i>Specificity</i>	<i>Positive Predictive Value</i>	<i>Negative Predictive Value</i>	<i>Overall Percent Agreement</i>
	31	0.04	1.00	1.00	0.36	0.38
	32	0.03	1.00	1.00	0.36	0.38
	33	0.03	1.00	1.00	0.36	0.38
	34	0.02	1.00	1.00	0.36	0.37
	35	0.02	1.00	1.00	0.36	0.37
	36	0.02	1.00	1.00	0.36	0.36
	37	0.02	1.00	1.00	0.36	0.36

Table 14A

Performance Metrics in Predicting ADD/ADHD Status from Number of Positive Foods
Using 95th Percentile of ELISA Signal to determine Positive

Sex	No. of Positive Foods as Cutoff	Sensitivity	Specificity	Positive Predictive Value	Negative Predictive Value	Overall Percent Agreement
MALE	1	0.80	0.43	0.44	0.80	0.57
	2	0.65	0.63	0.49	0.76	0.64
	3	0.50	0.73	0.50	0.72	0.65
	4	0.41	0.82	0.56	0.71	0.67
	5	0.33	0.87	0.59	0.70	0.68
	6	0.28	0.90	0.60	0.69	0.68
	7	0.24	0.91	0.60	0.69	0.67
	8	0.21	0.92	0.62	0.68	0.67
	9	0.18	0.94	0.60	0.67	0.67
	10	0.15	0.94	0.57	0.67	0.66
	11	0.12	0.94	0.57	0.66	0.65
	12	0.10	0.95	0.50	0.66	0.65
	13	0.08	0.96	0.50	0.65	0.64
	14	0.07	0.96	0.50	0.65	0.64
	15	0.07	0.96	0.50	0.65	0.65
	16	0.07	0.97	0.50	0.65	0.65
	17	0.07	0.98	0.60	0.65	0.65
	18	0.07	0.98	0.67	0.65	0.65
	19	0.07	0.98	0.67	0.65	0.65
	20	0.07	0.98	0.67	0.65	0.65
	21	0.07	0.98	0.67	0.65	0.66
	22	0.07	0.98	0.75	0.65	0.66
	23	0.07	1.00	1.00	0.65	0.66
	24	0.07	1.00	1.00	0.66	0.66
	25	0.07	1.00	1.00	0.66	0.66
	26	0.06	1.00	1.00	0.65	0.66
	27	0.04	1.00	1.00	0.65	0.66
	28	0.04	1.00	1.00	0.65	0.66
	29	0.04	1.00	1.00	0.65	0.66
	30	0.04	1.00	1.00	0.65	0.66

<i>Sex</i>	<i>No. of Positive Foods as Cutoff</i>	<i>Sensitivity</i>	<i>Specificity</i>	<i>Positive Predictive Value</i>	<i>Negative Predictive Value</i>	<i>Overall Percent Agreement</i>
	31	0.03	1.00	1.00	0.65	0.65
	32	0.00	1.00	1.00	0.65	0.65
	33	0.00	1.00	1.00	0.64	0.64
	34	0.00	1.00	1.00	0.64	0.64
	35	0.00	1.00	1.00	0.64	0.64
	36	0.00	1.00	.	0.64	0.64
	37	0.00	1.00	.	0.64	0.64

Table 14B

CLAIMS

What is claimed is:

1. A test kit for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
 - one or more distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier;
 - wherein the distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value, or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
2. The test kit of claim 1, wherein the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD.
3. The test kit of claim 1, wherein the one or more distinct food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
4. The test kit of claim 1, wherein the one or more distinct food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
5. The test kit of claim 1, wherein the one or more distinct food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
6. The test kit of claim 1, wherein the one or more distinct food preparations includes at least 12 food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
7. The test kit of claim 1, wherein the one or more distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food

preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

8. The test kit of any one of claims 1-7 wherein the one or more distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.
9. The test kit of any one of claims 1-7 wherein the one or more distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.

10. The test kit of any one of claims 1-9 wherein FDR multiplicity adjusted p-value is adjusted for at least one of age and gender.
11. The test kit of any one of claims 1-9 wherein FDR multiplicity adjusted p-value is adjusted for age and gender.
12. The test kit of any one of claims 1-11 wherein at least 50% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
13. The test kit of any one of the claims 1-11 wherein at least 70% of the one or more distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
14. The test kit of any one of the claims 1-13 wherein all of the one or more distinct food preparations, when adjusted for a single gender, have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
15. The test kit of any one of the claims 1-14 wherein the one or more distinct food preparations are crude filtered aqueous extracts.
16. The test kit of any one of the claims 1-15 wherein the one or more distinct food preparations are processed aqueous extracts.
17. The test kit of any one of the claims 1-16 wherein the solid carrier is a well of a multiwell plate, a bead, an electrical, a chemical sensor, a microchip or an adsorptive film.
18. A method of testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
 - contacting the test kit of claim 1 with a bodily fluid of a patient that is diagnosed with or suspected of having ADD/ADHD,

wherein the step of contacting is performed under conditions that allow IgG from the bodily fluid to bind to at least one component of the food preparation; measuring IgG bound to the at least one component of the food preparation to obtain a signal; and updating or generating a report using the signal.

19. The method of claim 18 wherein the bodily fluid of the patient is selected from the group consisting of whole blood, plasma, serum, saliva, urine and a fecal suspension.
20. The method of claim 18 or claim 19 wherein the step of contacting the test kit is performed with a plurality of distinct food preparations.
21. The method of claim 20 wherein the plurality of distinct food preparations is prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
22. The method of claim 20 wherein the plurality of distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food

preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

23. The method of claim 20 wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
24. The method of claim 20 wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.
25. The method of claim 20 wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.
26. The method of claim 20 wherein all of the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
27. The method of claim 18 wherein the food preparation is immobilized on a solid surface, optionally in an addressable manner.
28. The method of any of the claims 18-26 wherein the food preparation is immobilized on a solid surface, optionally in an addressable manner.
29. The method of any of the claims 18-28 wherein the step of measuring IgG bound to the at least one component of the food preparation is performed via immunoassay test.

30. The method of any of the claims 18-29, further comprising comparing the signal to a gender-stratified reference value for the food preparation using gender identification to obtain a result, wherein the gender-stratified reference value for the food preparation is at least a 90th percentile value.
31. A method of generating a test for food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
- obtaining test results for a plurality of distinct food preparations, wherein the test results are based on bodily fluids of patients diagnosed with or suspected of having ADD/ADHD and bodily fluids of a control group not diagnosed with or not suspected of having ADD/ADHD;
 - stratifying the test results by gender for each of the distinct food preparations;
 - assigning for a predetermined percentile rank a different cutoff value for male and female patients for each of the distinct food preparations;
 - selecting a plurality of distinct food preparations that each have a discriminatory p-value of ≤ 0.07 as determined by raw p-value or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value; and
 - generating a test consisting essentially of the selected food preparations for food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD.
32. The method of claim 31 wherein the test result is an ELISA result.
33. The method of claim 31 or claim 32 wherein the plurality of distinct food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
34. The method of any of claim 31 or claim 32 wherein the plurality of distinct food preparations includes at least six food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
35. The method of any of claim 31 or 32 wherein the plurality of distinct food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from

foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

36. The method of any of claims 31-35 wherein the plurality of different food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.
37. The method of any of claims 31-35 wherein the plurality of different food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.

38. The method of any of claims 31-37 wherein the bodily fluid of the patient is selected from the group consisting of whole blood, plasma, serum, saliva, urine or a fecal suspension.
39. The method of any of claims 31-38 wherein the predetermined percentile rank is an at least 90th percentile rank.
40. The method of any of claims 31-39 wherein the cutoff value for male and female patients has a difference of at least 10% (abs).
41. The method of any of claims 18-40, further comprising a step of normalizing the result to a **patient's total IgG**.
42. The method of any of claims 18-40, further comprising a step of normalizing the result to a **global mean of the patient's food specific IgG results**.
43. The method of any of claims 18-40, further comprising a step of identifying a subset of **patients, wherein the subset of patients' sensitivities to the food preparations underlies ADD/ADHD by raw p-value or an average discriminatory p-value of ≤ 0.01** .
44. The method of any of claims 18-40, further comprising a step of determining numbers of the food preparations, wherein the numbers of the food preparations can be used to confirm ADD/ADHD by raw p-value or an average discriminatory p-value of ≤ 0.01 .
45. Use of a plurality of distinct food preparations coupled to individually addressable respective solid carriers for the diagnosis of ADD/ADHD, wherein the plurality of distinct food preparations have an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
46. Use of claim 45 wherein the plurality of food preparations includes at least two food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
47. Use of claim 45 wherein the plurality of food preparations includes at least four food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.

48. Use of claim 45 wherein the plurality of food preparations includes at least eight food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
49. Use of claim 45 wherein the plurality of food preparations includes at least 12 food preparations prepared from food items of Table 1 or selected from foods 1-37 of Table 2.
50. Use of claim 45 wherein the plurality of food preparations is selected from the group consisting of at least 1 food preparation from foods 1-37 of Table 2, at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

51. Use of any one of claims 45-50, wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.
52. Use of any one of claims 45-50 wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.025 as determined by raw p-value or an average discriminatory p-value of ≤ 0.07 as determined by FDR multiplicity adjusted p-value.
53. Use of any one of claims 45-52 wherein FDR multiplicity adjusted p-value is adjusted for at least one of age and gender.
54. Use of any one of claims 45-52 wherein FDR multiplicity adjusted p-value is adjusted for age and gender.
55. Use of any one of claims 45-54 wherein at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
56. Use of any one of the claims 45-54 wherein at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
57. Use of any one of the claims 45-54 wherein all of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
58. Use of any one of the claims 45-57 wherein the plurality of distinct food preparations is crude filtered aqueous extracts.
59. Use of any one of the claims 45-58 wherein the plurality of distinct food preparations is processed aqueous extracts.

60. Use of any one of the claims 45-59 wherein the solid carrier is a well of a multiwell plate, a bead, an electrical sensor, a chemical sensor, a microchip, or an adsorptive film.
61. Use of any one of claims 45-59, wherein the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD headaches with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD headaches.
62. The method of claim 31, wherein the test result is an ELISA result derived from a process that includes separately contacting each distinct food preparation with the bodily fluid of each patient.
63. A detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier;
wherein the plurality of distinct food preparations consists essentially of food preparations that each have a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
64. The detection apparatus of claim 63, wherein the discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or not suspected of having ADD/ADHD.
65. The detection apparatus of claim 63, wherein the plurality of food preparations includes at least two food preparations selected from food items of Table 1 or selected from foods items 1-37 of Table 2.
66. The detection apparatus of claim 63, wherein the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at

least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

67. The detection apparatus of claim 63, wherein the plurality of distinct food preparations consists essentially of food preparations that each has a discriminatory p-value of ≤ 0.05 as determined by raw p-value or a discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

68. The detection apparatus of claim 63, wherein FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.
69. The detection apparatus of claim 63, wherein at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
70. The detection apparatus of claim 63, wherein the plurality of distinct food preparations is a crude aqueous extract or a processed aqueous extract.
71. A detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier;
wherein the plurality of distinct food preparations consists essentially of food preparations having an average discriminatory p-value of ≤ 0.07 as determined by raw p-value, or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
72. The detection apparatus of claim 71, wherein the average discriminatory p-value is determined by a process comprising comparing assay values of a first patient test cohort that is diagnosed with or suspected of having ADD/ADHD with assay values of a second patient test cohort that is not diagnosed with or suspected of having ADD/ADHD.
73. The detection apparatus of claim 71, wherein the plurality of food preparations includes at least two food preparations selected from food items of Table 1 or selected from foods items 1-37 of Table 2.
74. The detection apparatus of claim 71, wherein the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at

least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37 of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

75. The detection apparatus of claim 71, wherein the plurality of distinct food preparations consists essentially of food preparations having an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

76. The detection apparatus of claim 71, wherein FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.
77. The detection apparatus of claim 71, wherein at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
78. The detection apparatus of claim 71, wherein at least 60% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
79. The detection apparatus of claim 71, wherein at least 65% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
80. The detection apparatus of claim 71, wherein at least 70% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
81. The detection apparatus of claim 71, wherein at least 75% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
82. The detection apparatus of claim 71, wherein at least 80% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value

- of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
83. The detection apparatus of claim 71, wherein at least 85% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
84. The detection apparatus of claim 71, wherein the plurality of distinct food preparations is a crude aqueous extract or a processed aqueous extract.
85. A detection apparatus for testing food sensitivity in a patient diagnosed with or suspected of having ADD/ADHD, comprising:
a plurality of distinct food preparations, wherein each food preparation is independently coupled to an individually addressable solid carrier;
wherein at least 50% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
86. The detection apparatus of claim 85, wherein at least 55% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
87. The detection apparatus of claim 85, wherein at least 60% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
88. The detection apparatus of claim 85, wherein at least 65% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

89. The detection apparatus of claim 85, wherein at least 70% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
90. The detection apparatus of claim 85, wherein at least 75% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
91. The detection apparatus of claim 85, wherein at least 80% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
92. The detection apparatus of claim 85, wherein at least 85% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
93. The detection apparatus of claim 85, wherein at least 90% of the plurality of distinct food preparations each has a discriminatory p-value of ≤ 0.07 as determined by raw p-value, or a discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.
94. The detection apparatus of claim 85, wherein the plurality of food preparations is selected from the group consisting of at least 2 food preparations from foods 1-37 of Table 2, at least 3 food preparations from foods 1-37 of Table 2, at least 4 food preparations from foods 1-37 of Table 2, at least 5 food preparations from foods 1-37 of Table 2, at least 6 food preparations from foods 1-37 of Table 2, at least 7 food preparations from foods 1-37 of Table 2, at least 8 food preparations from foods 1-37 of Table 2, at least 9 food preparations from foods 1-37 of Table 2, at least 10 food preparations from foods 1-37 of Table 2, at least 11 food preparations from foods 1-37 of Table 2, at least 12 food preparations from foods 1-37 of Table 2, at least 13 food preparations from foods 1-37 of Table 2, at least 14 food preparations from foods 1-37 of Table 2, at least 15 food preparations from foods 1-37 of Table 2, at least 16 food preparations from foods 1-37 of Table 2, at least 17 food preparations from foods 1-37 of Table 2, at least 18 food preparations from foods 1-37 of Table 2, at least 19 food preparations from foods 1-37

of Table 2, at least 20 food preparations from foods 1-37 of Table 2, at least 21 food preparations from foods 1-37 of Table 2, at least 22 food preparations from foods 1-37 of Table 2, at least 23 food preparations from foods 1-37 of Table 2, at least 24 food preparations from foods 1-37 of Table 2, at least 25 food preparations from foods 1-37 of Table 2, at least 26 food preparations from foods 1-37 of Table 2, at least 27 food preparations from foods 1-37 of Table 2, at least 28 food preparations from foods 1-37 of Table 2, at least 29 food preparations from foods 1-37 of Table 2, at least 30 food preparations from foods 1-37 of Table 2, at least 31 food preparations from foods 1-37 of Table 2, at least 32 food preparations from foods 1-37 of Table 2, at least 33 food preparations from foods 1-37 of Table 2, at least 34 food preparations from foods 1-37 of Table 2, at least 35 food preparations from foods 1-37 of Table 2, at least 36 food preparations from foods 1-37 of Table 2, and at least 37 food preparations from foods 1-37 of Table 2.

95. The detection apparatus of claim 85, wherein the plurality of distinct food preparations has an average discriminatory p-value of ≤ 0.05 as determined by raw p-value or an average discriminatory p-value of ≤ 0.08 as determined by FDR multiplicity adjusted p-value.

96. The detection apparatus of claim 85, wherein FDR multiplicity adjusted p-value is adjusted for at least one of age or gender.

97. The detection apparatus of claim 85, wherein at least 50% of the plurality of distinct food preparations, when adjusted for a single gender, has an average discriminatory p-value of ≤ 0.07 as determined by raw p-value or an average discriminatory p-value of ≤ 0.10 as determined by FDR multiplicity adjusted p-value.

98. The detection apparatus of claim 85, wherein the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

99. A detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising:

a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface,

wherein the plurality of food preparations consists essentially of trigger foods for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

100. The detection apparatus of claim 99, wherein the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods 1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2.

101. The detection apparatus of claim 99, wherein the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

102. The detection apparatus of claim 99, wherein the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.
103. A detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising:
- a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface,
- wherein at least 50% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
104. The detection apparatus of claim 103, wherein at least 55% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
105. The detection apparatus of claim 103, wherein at least 60% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
106. The detection apparatus of claim 103, wherein at least 65% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
107. The detection apparatus of claim 103, wherein at least 70% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
108. The detection apparatus of claim 103, wherein at least 75% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
109. The detection apparatus of claim 103, wherein at least 80% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

110. The detection apparatus of claim 103, wherein at least 85% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
111. The detection apparatus of claim 103, wherein at least 90% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
112. The detection apparatus of claim 103, wherein at least 95% of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
113. The detection apparatus of claim 103, wherein the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods 1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at

least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2.

114. The detection apparatus of claim 103, wherein the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

115. The detection apparatus of claim 103, wherein the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

116. A detection apparatus for testing food sensitivity in a biological sample from a patient diagnosed with or suspected of having ADD/ADHD, comprising:

a solid surface with a plurality of food preparations, wherein each food preparation is independently coupled to the solid surface,

wherein at least seven food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

117. The detection apparatus of claim 116, wherein at least eight food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

118. The detection apparatus of claim 116, wherein at least nine food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

119. The detection apparatus of claim 116, wherein at least ten food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.

120. The detection apparatus of claim 116, wherein at least eleven food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
121. The detection apparatus of claim 116, wherein at least twelve food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
122. The detection apparatus of claim 116, wherein at least thirteen food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
123. The detection apparatus of claim 116, wherein at least fourteen food preparations of the plurality of food preparations is a trigger food for ADD/ADHD capable of reacting with an immunoglobulin from the biological sample.
124. The detection apparatus of claim 116, wherein the trigger foods is selected from the group consisting of at least 2 food items from foods 1-37 of Table 2, at least 3 food items from foods 1-37 of Table 2, at least 4 food items from foods 1-37 of Table 2, at least 5 food items from foods 1-37 of Table 2, at least 6 food items from foods 1-37 of Table 2, at least 7 food items from foods 1-37 of Table 2, at least 8 food items from foods 1-37 of Table 2, at least 9 food items from foods 1-37 of Table 2, at least 10 food items from foods 1-37 of Table 2, at least 11 food items from foods 1-37 of Table 2, at least 12 food items from foods 1-37 of Table 2, at least 13 food items from foods 1-37 of Table 2, at least 14 food items from foods 1-37 of Table 2, at least 15 food items from foods 1-37 of Table 2, at least 16 food items from foods 1-37 of Table 2, at least 17 food items from foods 1-37 of Table 2, at least 18 food items from foods 1-37 of Table 2, at least 19 food items from foods 1-37 of Table 2, at least 20 food items from foods 1-37 of Table 2, at least 21 food items from foods 1-37 of Table 2, at least 22 food items from foods 1-37 of Table 2, at least 23 food items from foods 1-37 of Table 2, at least 24 food items from foods 1-37 of Table 2, at least 25 food items from foods 1-37 of Table 2, at least 26 food items from foods 1-37 of Table 2, at least 27 food items from foods 1-37 of Table 2, at least 28 food items from foods 1-37 of Table 2, at least 29 food items from foods 1-37 of Table 2, at least 30 food items from foods

1-37 of Table 2, at least 31 food items from foods 1-37 of Table 2, at least 32 food items from foods 1-37 of Table 2, at least 33 food items from foods 1-37 of Table 2, at least 34 food items from foods 1-37 of Table 2, at least 35 food items from foods 1-37 of Table 2, at least 36 food items from foods 1-37 of Table 2, and at least 37 food items from foods 1-37 of Table 2.

125. The detection apparatus of claim 116, wherein the trigger foods is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

126. The detection apparatus of claim 116, wherein the plurality of distinct food preparations is crude aqueous extracts or processed aqueous extracts.

127. The test kit of claim 1, wherein the one or more distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

128. The method of claim 20 wherein the plurality of distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

129. The method of any of claim 31 or 32 wherein the plurality of distinct food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

130. Use of claim 45 wherein the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.
131. The detection apparatus of claim 63, wherein the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.
132. The detection apparatus of claim 71, wherein the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.
133. The detection apparatus of claim 85, wherein the plurality of food preparations is selected from the group consisting of cantaloupe, wheat, tomato, cucumber, squashes, almond, egg, cauliflower, pinto bean, broccoli, orange, butter, corn, lettuce, rye, peach, green pea, carrot, tea, mustard, strawberry, celery, green pepper, garlic, oat, onion, banana, eggplant, cabbage, safflower, olive, cottage cheese, grapefruit, walnut blk, cow milk, soybean, and chili pepper.

Distribution of ELISA Signal Scores by Diagnosis
Sex=MALE Food=Cantaloupe




Figure 1A

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control Cutpoint
across 1000 Bootstrapped Samples*

Sex=MALE Food=Cantaloupe




Figure 1B

Distribution of ELISA Signal Scores by Diagnosis
Sex=FEMALE Food=Cantaloupe




Figure 1C

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control
Cutpoint across 1000 Bootstrapped Samples*

Sex=FEMALE Food=Cantaloupe




Figure 1D

Distribution of ELISA Signal Scores by Diagnosis
Sex=MALE Food=Wheat




Figure 2A

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control
Cutpoint across 1000 Bootstrapped Samples*

Sex=MALE Food= Wheat





Figure 2B

*Distribution of ELISA Signal Scores by Diagnosis**Sex=FEMALE Food= Wheat***Figure 2C**

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control
Cutpoint across 1000 Bootstrapped Samples*

Sex=FEMALE Food= Wheat




Figure 2D

Distribution of ELISA Signal Scores by Diagnosis
Sex=MALE Food=Tomato




Figure 3A

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control
Cutpoint across 1000 Bootstrapped Samples*

Sex=MALE Food=Tomato





Figure 3B

*Distribution of ELISA Signal Scores by Diagnosis**Sex=FEMALE Food=Tomato***Figure 3C**

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control Cutpoint
across 1000 Bootstrapped Samples*

Sex=FEMALE Food=Tomato





Figure 3D

*Distribution of ELISA Signal Scores by Diagnosis**Sex=MALE Food=Cucumber***Figure 4A**

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control
Cutpoint across 1000 Bootstrapped Samples
Sex=MALE Food=Cucumber*




Figure 4B

Distribution of ELISA Signal Scores by Diagnosis
Sex=FEMALE Food=Cucumber




Figure 4C

*Distribution of Percentage of ADD/ADHD Subjects with Signals $\geq$ Control Cutpoint
across 1000 Bootstrapped Samples
Sex=FEMALE Food=Cucumber*




Figure 4D

*Distribution of ADD/ADHD Subjects by Number of Foods in which they were rated as "Positive"
by Sex*

90th Percentile as Cutpoint




Figure 5A

***Distribution of ADD/ADHD Subjects by Number of Foods in which they were rated as "Positive"
by Sex***

95th Percentile as Cutpoint









Figure 5B


**Figure 6A**

**Figure 6B**

**Figure 6C**

**Figure 6D**

**Figure 7A**

**Figure 7B**

Distribution of ELISA Signal Scores by Diagnosis

Sex=FEMALE Food= Wheat




Figure 2C