

A METHOD OF INCREASING THE EFFECT OF AN ACTIVATED-POTENTIATED FORM OF AN ANTIBODY

FIELD

The present invention relates to a method of increasing the effect of an activated-potentiated form of an antibody and a pharmaceutical formulation comprising an activated-potentiated form of an antibody to an endogenous biological molecule and an activated-potentiated form of an antibody to endothelial NO-synthase.

BACKGROUND

Nitric oxide (NO) is a gaseous molecule that has been shown to act in the signaling of different biological processes. Endothelium-derived NO is a key molecule in regulation of vascular tone and its association with vascular disease has long been recognized. NO inhibits many processes known to be involved in the formation of atherosclerotic plaque, including monocyte adhesion, platelet aggregation and vascular smooth muscle cell proliferation. Another important role of endothelial NO is the protection of the vascular wall from the oxidative stress induced by its own metabolic products and by the oxidation products of lipids and lipoproteins. Endothelial dysfunction occurs at very early stages of atherosclerosis. It is therefore possible that deficiency in local NO availability could be a final common pathway that accelerates atherogenesis in humans. In addition to its role in the vascular endothelium, NO availability has been shown to modulate metabolism of lipoproteins. Negative correlation has been reported between plasma concentrations of NO metabolic products and plasma total and Low Density Lipoprotein [LDL] cholesterol levels while High Density Lipoprotein [HDL] improves vascular function in hypercholesterolaemic subjects. The loss of NO has considerable effect on the development of the disease. Diabetes mellitus is associated with increased rates of morbidity and mortality caused primarily by the accelerated development of atherosclerotic disease. Moreover, reports show that diabetics have impaired lung functions. It has been proposed that insulin resistance

leads to airway inflammation. Habib et al., *Nitric Oxide Measurement From Blood To Lungs, Is There A Link?* Pak J Physiol 2007; 3(1).

Nitric oxide is synthesized by the endothelium from L-arginine by nitric oxide synthase (NO synthase). NO synthase occurs in different isoforms, including a constitutive form (cNOS) and an inducible form (iNOS). The constitutive form is present in normal endothelial cells, neurons and some other tissues.

The therapeutic effect of an extremely diluted form (or ultra-low form) of antibodies potentized by homeopathic technology (activated potentiated form) has been discovered by Dr. Oleg I. Epshtein. U.S. Patent No. 7,700,096 discloses a homeopathically potentized form of antibodies to endothelial NO-synthase. The homeopathically potentized form of antibodies to endothelial NO-synthase is marketed in the Russian Federation and other countries under the name Impaza®.

There is a continuing need for a method of increasing the effect of an activated-potentiated form of an antibody.

SUMMARY

In accordance with one aspect, the present invention provides a method of increasing the effect of an activated-potentiated form of an antibody to an endogenous biological molecule, said method comprising combining said endogenous biological molecule with an activated-potentiated form of an antibody to endothelial NO-synthase. Preferably, the method aspect of the invention includes administering said combination to a patient in need of treatment with said activated-potentiated form of an antibody.

In one variant, said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to S-100 protein. In another variant, said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to prostate specific antigen. In another variant, said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to insulin receptor. In another variant, said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to antiotensin receptor II.

In accordance with another aspect, the invention provides a pharmaceutical composition comprising a) an activated-potentiated form of an antibody to an endogenous biological molecule, and b) an activated-potentiated form of an antibody to NO synthase. Preferably, the pharmaceutical composition pharmaceutically acceptable solid carrier. Preferably, the activated-potentiated form of an antibody to endothelial NO synthase contains a mixture of C12, C30, and C200 homeopathic dilutions impregnated onto the solid carrier. The activated-potentiated form of an antibody to an endogenous biological molecule could be a monoclonal, monoclonal, or natural antibody. Preferably, the antibody to human insulin receptor is a polyclonal antibody.

It is contemplated that the pharmaceutical composition comprises an activated-potentiated form of an antibody to an endogenous biological molecule prepared by successive centesimal dilutions coupled with shaking of every dilution. It is also contemplated that the antibody to endothelial NO-synthase is monoclonal, polyclonal or natural antibody. It is particularly preferred that the antibody to endothelial NO-synthase is a polyclonal antibody. It is contemplated that the activated-potentiated form of an antibody to endothelial NO-synthase is prepared by successive centesimal dilutions coupled with shaking of every dilution.

DESCRIPTION OF THE FIGURES

Figure 1 - Illustrates the effect of tested preparations on blood plasma glucose level of rats with streptozotocin-induced diabetes mellitus

Figure 2 - Illustrates the effect of tested preparations on day 14 of injection on indicators of area under concentration-time curve (AUC) in the glucose tolerance test in rats with streptozotocin-induced diabetes mellitus.

Figure 3 - Illustrates the effect of tested preparations on blood plasma glucose level of rats with spontaneous non-insulin-dependent diabetes mellitus.

Figure 4 - Illustrates the effect of tested preparations on day 28 of injection on indicators of area under concentration-time curve (AUC) in glucose tolerance test in rats with spontaneous non-insulin-dependent diabetes mellitus.

DETAILED DESCRIPTION

The invention is defined with reference to the appended claims. With respect to the claims, the glossary that follows provides the relevant definitions.

The term "antibody" as used herein shall mean an immunoglobulin that specifically binds to, and is thereby defined as complementary with, a particular spatial and polar organization of another molecule. Antibodies as recited in the claims may include a complete immunoglobulin or fragment thereof, may be natural, polyclonal or monoclonal, and may include various classes and isotypes, such as IgA, IgD, IgE, IgG1, IgG2a, IgG2b and IgG3, IgM, etc. Fragments thereof may include Fab, Fv and F(ab')², Fab', and the like. The singular "antibody" includes plural "antibodies."

The term "activated-potentiated form" or "potentiated form" respectively, with respect to antibodies recited herein is used to denote a product of homeopathic potentization of any initial solution of antibodies. "Homeopathic potentization" denotes the use of methods of homeopathy to impart homeopathic potency to an initial solution of relevant substance. Although not so limited, "homeopathic potentization" may involve, for example, repeated consecutive dilutions combined with external treatment, particularly vertical (mechanical) shaking. In other words, an initial solution of antibody is subjected to consecutive repeated dilution and multiple vertical shaking of each obtained solution in accordance with homeopathic technology. The preferred concentration of the initial solution of antibody in the solvent, preferably water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml. The preferred procedure for preparing each component, i.e. antibody solution, is the use of the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture) of antibodies diluted 100¹², 100³⁰ and 100²⁰⁰ times, respectively, which is equivalent to centesimal homeopathic dilutions (C12, C30, and C200) or the use of the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution of antibodies diluted 100¹², 100³⁰ and 100⁵⁰ times, respectively, which is equivalent to centesimal homeopathic dilutions (C12, C30 and C50). Examples of homeopathic

potentization are described in U.S. Patent. Nos. 7,572,441 and 7,582,294, which are incorporated herein by reference in their entirety and for the purpose stated. While the term "activated-potentiated form" is used in the claims, the term "ultra-low doses" is used in the examples. The term "ultra-low doses" became a term of art in the field of art created by study and use of homeopathically diluted and potentized form of substance. The term "ultra-low dose" or "ultra-low doses" is meant as fully supportive and primarily synonymous with the term 'activated-potentiated" form used in the claims.

In other words, an antibody is in the "activated-potentiated" or "potentiated" form when three factors are present. First, the "activated-potentiated" form of the antibody is a product of a preparation process well accepted in the homeopathic art. Second, the "activated-potentiated" form of antibody must have biological activity determined by methods well accepted in modern pharmacology. And third, the biological activity exhibited by the "activated potentiated" form of the antibody cannot be explained by the presence of the molecular form of the antibody in the final product of the homeopathic process.

For example, the activated potentiated form of antibodies may be prepared by subjecting an initial, isolated antibody in a molecular form to consecutive multiple dilutions coupled with an external impact, such as mechanical shaking. The external treatment in the course of concentration reduction may also be accomplished, for example, by exposure to ultrasonic, electromagnetic, or other physical factors. V. Schwabe "Homeopathic medicines", M., 1967, U.S. Patents Nos. 7,229,648 and 4,311,897, which are incorporated by reference in their entirety and for the purpose stated, describe such processes that are well-accepted methods of homeopathic potentiation in the homeopathic art. This procedure gives rise to a uniform decrease in molecular concentration of the initial molecular form of the antibody. This procedure is repeated until the desired homeopathic potency is obtained. For the individual antibody, the required homeopathic potency can be determined by subjecting the intermediate dilutions to biological testing in the desired pharmacological model. Although not so limited, 'homeopathic potentization" may involve, for example, repeated consecutive dilutions combined with external treatment, particularly vertical (mechanical) shaking. In other words, an initial solution of antibody is subjected to consecutive

repeated dilution and multiple vertical shaking of each obtained solution in accordance with homeopathic technology. The preferred concentration of the initial solution of antibody in the solvent, preferably, water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml. The preferred procedure for preparing each component, i.e. antibody solution, is the use of the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture) of antibodies diluted 100¹², 100³⁰ and 100²⁰⁰ times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30 and C200 or the mixture of three aqueous or aqueous-alcohol dilutions of the primary matrix solution (mother tincture) of antibodies diluted 100¹², 100³⁰ and 100⁵⁰ times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30 and C50. Examples of how to obtain the desired potency are also provided, for example, in U.S. Patent Nos. 7,229,648 and 4,311,897, which are incorporated by reference for the purpose stated. The procedure applicable to the "activated-potentiated" form of the antibodies described herein is described in more detail below.

There has been a considerable amount of controversy regarding homeopathic treatment of human subjects. While the present invention relies on accepted homeopathic processes to obtain the "activated-potentiated" form of antibodies, it does not rely solely on homeopathy in human subjects for evidence of activity. It has been surprisingly discovered by the inventor of the present application and amply demonstrated in the accepted pharmacological models that the solvent ultimately obtained from consecutive multiple dilution of a starting molecular form of an antibody has definitive activity unrelated to the presence of the traces of the molecular form of the antibody in the target dilution. The "activated-potentiated" form of the antibody provided herein are tested for biological activity in well accepted pharmacological models of activity, either in appropriate *in vitro* experiments, or *in vivo* in suitable animal models. The experiments provided further below provide evidence of biological activity in such models. Human clinical studies also provide evidence that the activity observed in the animal model is well translated to human therapy. Human studies have also provided evidence of availability of the "activated potentiated" forms described herein to treat

specified human diseases or disorders well accepted as pathological conditions in the medical science.

Also, the claimed "activated-potentiated" form of antibody encompasses only solutions or solid preparations the biological activity of which cannot be explained by the presence of the molecular form of the antibody remaining from the initial, starting solution. In other words, while it is contemplated that the "activated-potentiated" form of the antibody may contain traces of the initial molecular form of the antibody, one skilled in the art could not attribute the observed biological activity in the accepted pharmacological models to the remaining molecular form of the antibody with any degree of plausibility due to the extremely low concentrations of the molecular form of the antibody remaining after the consecutive dilutions. While the invention is not limited by any specific theory, the biological activity of the "activated-potentiated" form of the antibodies of the present invention is not attributable to the initial molecular form of the antibody. Preferred is the "activated-potentiated" form of antibody in liquid or solid form in which the concentration of the molecular form of the antibody is below the limit of detection of the accepted analytical techniques, such as capillary electrophoresis and High Performance Liquid Chromatography. Particularly preferred is the "activated-potentiated" form of antibody in liquid or solid form in which the concentration of the molecular form of the antibody is below the Avogadro number. In the pharmacology of molecular forms of therapeutic substances, it is common practice to create a dose-response curve in which the level of pharmacological response is plotted against the concentration of the active drug administered to the subject or tested in vitro. The minimal level of the drug which produces any detectable response is known as a threshold dose. It is specifically contemplated and preferred that the "activated-potentiated" form of the antibodies contains molecular antibody, if any, at a concentration below the threshold dose for the molecular form of the antibody in the given biological model.

The combination pharmaceutical composition in accordance with this aspect of the invention may be in the liquid form or in solid form. Each of the activated potentiated

forms of the antibodies included in the pharmaceutical composition is prepared from an initial molecular form of the antibody via a process accepted in homeopathic art. The starting antibodies may be monoclonal, or polyclonal antibodies prepared in accordance with known processes, for example, as described in *Immunotechniques*, G. Frimel, M., "Meditsyna", 1987, p. 9-33; "Hum. Antibodies. Monoclonal and recombinant antibodies, 30 years after" by Laffly E., Sodoyer R. - 2005 - Vol. 14. - N 1-2. P.33-55, both incorporated herein by reference.

Monoclonal antibodies may be obtained, e.g., by means of hybridoma technology. The initial stage of the process includes immunization based on the principles already developed in the course of polyclonal antisera preparation. Further stages of work involve the production of hybrid cells generating clones of antibodies with identical specificity. Their separate isolation is performed using the same methods as in the case of polyclonal antisera preparation.

Polyclonal antibodies may be obtained via active immunization of animals. For this purpose, for example, suitable animals (e.g. rabbits) receive a series of injections of the appropriate antigen, for example, NO synthase. The animals' immune system generates corresponding antibodies, which are collected from the animals in a known manner. This procedure enables preparation of a monospecific antibody-rich serum.

If desired, the serum containing antibodies may be purified, for example by using affine chromatography, fractionation by salt precipitation, or ion-exchange chromatography. The resulting purified, antibody-enriched serum may be used as a starting material for the preparation of the activated-potentiated form of the antibodies. The preferred concentration of the resulting initial solution of antibody in the solvent, preferably water or a water-ethyl alcohol mixture, ranges from about 0.5 to about 5.0 mg/ml.

The preferred procedure for preparing each component of the combination drug according to the present invention is the use of the mixture of three aqueous-alcohol dilutions of the primary matrix solution of antibodies diluted 100^{12} , 100^{30} and 100^{50} times, respectively, which is equivalent to centesimal homeopathic dilutions C12, C30, and C50 or diluted 100^{12} , 100^{30} and 100^{200} times, respectively, which is equivalent to centesimal

homeopathic dilutions C12, C30 and C200. To prepare a solid dosage form, a solid carrier is treated with the desired dilution obtained via the homeopathic process. To obtain a solid unit dosage form of the combination of the invention, the carrier mass is impregnated with each of the dilutions. Both orders of impregnation are suitable to prepare the desired combination dosage form.

In a preferred embodiment, the starting material for the preparation of the activated potentiated form that comprise the combination of the invention is polyclonal, animal-raised antibody to the corresponding antigen, namely, NO synthase and endogenous biological molecule. To obtain the activated-potentiated form of polyclonal antibodies to NO synthase, the desired antigen may be injected as immunogen into a laboratory animal, preferably, rabbits. Polyclonal antibodies to NO synthase may be obtained using the whole molecule of bovine NO synthase of the following sequence:

SEQ ID NO: 1

Met	Gly	Asn	Leu	Lys	Ser	Val	Gly	Gin	Glu	Pro	Gly	Pro	Pro	Cys
1				5					10					15
Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Cys	Gly	Lys	Gin	Gly
16				20					25					30
Pro	Ala	Ser	Pro	Ala	Pro	Glu	Pro	Ser	Arg	Ala	Pro	Ala	Pro	Ala
31				35					40					45
Thr	Pro	His	Ala	Pro	Asp	His	Ser	Pro	Ala	Pro	Asn	Ser	Pro	Thr
46				50					55					60
Leu	Thr	Arg	Pro	Pro	Glu	Gly	Pro	Lys	Phe	Pro	Arg	Val	Lys	Asn
61				65					70					75
Trp	Glu	Leu	GLys	er	Ile	Thr	Tyr	Asp	Thr	Leu	Cys	Ala	Gin	Ser
76				80					85					90
Gin	Gin	Asp	Gly	Pro	Cys	Thr	Pro	Arg	Cys	Cys	Leu	GLys	er	Leu
91				95					100					105
Val	Leu	Pro	Arg	Lys	Leu	Gin	Thr	Arg	Pro	Ser	Pro	Gly	Pro	Pro
106				110					115					120
Pro	Ala	Glu	Gin	Leu	Leu	Ser	Gin	Ala	Arg	Asp	Phe	Ile	Asn	Gin
121				125					130					135
Tyr	Tyr	Ser	Ser	Ile	Lys	Arg	Ser	GLys	er	Gin	Ala	His	Glu	Glu
136				140					145					150
Arg	Leu	Gin	Glu	Val	Glu	Ala	Glu	Val	Ala	Ser	Thr	Gly	Thr	Tyr
151				155					160					165
His	Leu	Arg	Glu	Ser	Glu	Leu	Val	Phe	Gly	Ala	Lys	Gin	Ala	Trp
166				170					175					180

Arg	Asn	Ala	Pro	Arg	Cys	Val	Gly	Arg	H	e	Gin	Trp	Gly	Lys	Leu
181				185					190						195
Gin	Val	Phe	Asp	Ala	Arg	Asp	Cys	Ser	Ser	Ala	Gin	Glu	Met	Phe	
196				200					205						210
Thr	Tyr	lie	Cys	Asn	His	Ile	Lys	Tyr	Ala	Thr	Asn	Arg	Gly	Asn	
211				215					220						225
Leu	Arg	Ser	Ala	Ile	Thr	Val	Phe	Pro	Gin	Arg	Ala	Pro	Gly	Arg	
226				230					235						240
Gly	Asp	Phe	Arg	Ile	Trp	Asn	Ser	Gin	Leu	Val	Arg	Tyr	Ala	Gly	
241				245					250						255
Tyr	Arg	Gin	Gin	Asp	GLys	er	Val	Arg	Gly	Asp	Pro	Ala	Asn	Val	
256				260					265						270
Glu	Ile	Thr	Glu	Leu	Cys	H	e	Gin	His	Gly	Trp	Thr	Pro	Gly	Asn
271				275					280						285
Gly	Arg	Phe	Asp	Val	Leu	Pro	Leu	Leu	Leu	Gin	Ala	Pro	Asp	Glu	
286				290					295						300
Ala	Pro	Glu	Leu	Phe	Val	Leu	Pro	Pro	Glu	Leu	Val	Leu	Glu	Val	
301				305					310						315
Pro	Leu	Glu	His	Pro	Thr	Leu	Glu	Trp	Phe	Ala	Ala	Leu	Gly	Leu	
316				320					325						330
Arg	Trp	Tyr	Ala	Leu	Pro	Ala	Val	Ser	Asn	Met	Leu	Leu	Glu	Ile	
331				335					340						345
Gly	Gly	Leu	Glu	Phe	Ser	Ala	Ala	Pro	Phe	Ser	Gly	Trp	Tyr	Met	
346				350					355						360
Ser	Thr	Glu	H	e	Gly	Thr	Arg	Asn	Leu	Cys	Asp	Pro	His	Arg	Tyr
361				365					370						375
Asn	Ile	Leu	Glu	Asp	Val	Ala	Val	Cys	Met	Asp	Leu	Asp	Thr	Arg	
376				380					385						390
Thr	Thr	Ser	Ser	Leu	Trp	Lys	Asp	Lys	Ala	Ala	Val	Glu	Ile	Asn	
391				395					400						405
Leu	Ala	Val	Leu	His	Ser	Phe	Gin	Leu	Ala	Lys	Val	Thr	Ile	Val	
406				410					415						420
Asp	His	His	Ala	Ala	Thr	Val	Ser	Phe	Met	Lys	His	Leu	Asp	Asn	
421				425					430						435
Glu	Gin	Lys	Ala	Arg	Gly	Gly	Cys	Pro	Ala	Asp	Trp	Ala	Trp	Ile	
436				440					445						450
Val	Pro	Pro	Ile	Ser	GLys	er	Leu	Thr	Pro	Val	Phe	His	Gin	Glu	
451				455					460						465
Met	Val	Asn	Tyr	Ile	Leu	Ser	Pro	Ala	Phe	Arg	Tyr	Gin	Pro	Asp	
466				470					475						480
Pro	Trp	Lys	GLy	Ser	Ala	Thr	Lys	Gly	Ala	Gly	H	e	Thr	Arg	Lys
481				485					490						495
Lys	Thr	Phe	Lys	Glu	Val	Ala	Asn	Ala	Val	Lys	Ile	Ser	Ala	Ser	
496				500					505						510
Leu	Met	Gly	Thr	Leu	Met	Ala	Lys	Arg	Val	Lys	Ala	Thr	Ile	Leu	
511				515					510						525

Tyr	Ala	Ser	Glu	Thr	Gly	Arg	Ala	Gin	Ser	Tyr	Ala	Gin	Gin	Leu
526				530					535					540
Gly	Arg	Leu	Phe	Arg	Lys	Ala	Phe	Asp	Pro	Arg	Val	Leu	Cys	Met
541				545					550					555
Asp	Glu	Tyr	Asp	Val	Val	Ser	Leu	Glu	His	Glu	Ala	Leu	Val	Leu
556				560					565					570
Val	Val	Thr	Ser	Thr	Phe	Gly	Asn	Gly	Asp	Pro	Pro	Glu	Asn	Gly
571				575					580					585
Glu	Ser	Phe	Ala	Ala	Ala	Leu	Met	Glu	Met	Ser	Gly	Pro	Tyr	Asn
586				590					595					600
Ser	Ser	Pro	Arg	Pro	Glu	Gin	His	Lys	Ser	Tyr	Lys	He	Arg	Phe
601				605					610					615
Asn	Ser	Val	Ser	Cys	Ser	Asp	Pro	Leu	Val	Ser	Ser	Trp	Arg	Arg
616				620					625					630
Lys	Arg	Lys	Glu	Ser	Ser	Asn	Thr	Asp	Ser	Ala	Gly	Ala	Leu	Gly
631				635					640					645
Thr	Leu	Arg	Phe	Cys	Val	Phe	Gly	Leu	GLy	Ser	Arg	Ala	Tyr	Pro
646				650					655					660
His	Phe	Cys	Ala	Phe	Ala	Arg	Ala	Val	Asp	Thr	Arg	Leu	Glu	Glu
661				665					670					675
Leu	Gly	Gly	Glu	Arg	Leu	Leu	Gin	Leu	Gly	Gin	Gly	Asp	Glu	Leu
676				680					685					690
Cys	Gly	Gin	Glu	Glu	Ala	Phe	Arg	Gly	Trp	Ala	Lys	Ala	Ala	Phe
691				695					700					705
Gin	Ala	Ser	Cys	Glu	Thr	Phe	Cys	Val	Gly	Glu	Glu	Ala	Lys	Ala
706				710					715					720
Ala	Ala	Gin	Asp	Ile	Phe	Ser	Pro	Lys	Arg	Ser	Trp	Lys	Arg	Gin
721				725					730					735
Arg	Tyr	Arg	Leu	Ser	Thr	Gin	Ala	Glu	Gly	Leu	Gin	Leu	Leu	Pro
736				740					745					750
Gly	Leu	lie	His	Val	His	Arg	Arg	Lys	Met	Phe	Gin	Ala	Thr	Val
751				755					760					765
Leu	Ser	Val	Glu	Asn	Leu	Gin	Ser	Ser	Lys	Ser	Thr	Arg	Ala	Thr
766				770					775					780
Ile	Leu	Val	Arg	Leu	Asp	Thr	Ala	Gly	Gin	Glu	Gly	Leu	Gin	Tyr
781				785					790					795
Gin	Pro	Gly	Asp	His	Ile	Gly	Ile	Cys	Pro	Pro	Asn	Arg	Pro	Gly
796				800					805					810
Leu	Val	Glu	Ala	Leu	Leu	Ser	Arg	Val	Glu	Asp	Pro	Pro	Pro	Pro
811				815					820					825
Thr	Glu	Ser	Val	Ala	Val	Glu	Gin	Leu	Glu	Lys	GLys	er	Pro	Gly
826				830					835					840
Gly	Pro	Pro	Pro	Ser	Trp	Val	Arg	Asp	Pro	Arg	Leu	Pro	Pro	Cys
841				845					850					855
Thr	Leu	Arg	Gin	Ala	Leu	Thr	Phe	Phe	Leu	Asp	Ile	Thr	Ser	Pro
856				860					865					870

Pro	Ser	Pro	Arg	Leu	Leu	Arg	Leu	Leu	Ser	Thr	Leu	Ala	Glu	Glu
871				875					880					885
Pro	Ser	Glu	Gin	Gin	Glu	Leu	Glu	Thr	Leu	Ser	Gin	Asp	Pro	Arg
886				890					895					900
Arg	Tyr	Glu	Glu	Trp	Lys	Trp	Phe	Arg	Cys	Pro	Thr	Leu	Leu	Glu
901				905					910					915
Val	Leu	Glu	Gin	Phe	Pro	Ser	Val	Ala	Leu	Pro	Ala	Pro	Leu	Leu
916				920					925					930
Leu	Thr	Gin	Leu	Pro	Leu	Leu	Gin	Pro	Arg	Tyr	Tyr	Ser	Val	Ser
931				935					940					945
Ser	Ala	Pro	Asn	Ala	His	Pro	Gly	Glu	Val	His	Leu	Thr	Val	Ala
946				950					955					960
Val	Leu	Ala	Tyr	Arg	Thr	Gin	Asp	Gly	Leu	Gly	Pro	Leu	His	Tyr
961				965					970					975
Gly	Val	Cys	Ser	Thr	Trp	Leu	Ser	Gin	Leu	Lys	Thr	Gly	Asp	Pro
976				980					985					990
Val	Pro	Cys	Phe	lie	Arg	Gly	Ala	Pro	Ser	Phe	Arg	Leu	Pro	Pro
991				995					1000					1005
Asp	Pro	Tyr	Val	Pro	Cys	lie	Leu	Val	Gly	Pro	Gly	Thr	Gly	lie
1006				1010					1015					1020
Ala	Pro	Phe	Arg	Gly	Phe	Trp	Gin	Glu	Arg	Leu	His	Asp	lie	Glu
1021				1025					1030					1035
Ser	Lys	Gly	Leu	Gin	Pro	Ala	Pro	Met	Thr	Leu	Val	Phe	Gly	Cys
1036				1040					1045					1050
Arg	Cys	Ser	Gin	Leu	Asp	His	Leu	Tyr	Arg	Asp	Glu	Val	Gin	Asp
1051				1055					1060					1065
Ala	Gin	Glu	Arg	Gly	Val	Phe	Gly	Arg	Val	Leu	Thr	Ala	Phe	Ser
1066				1070					1075					1080
Arg	Glu	Pro	Asp	Ser	Pro	Lys	Thr	Tyr	Val	Gin	Asp	lie	Leu	Arg
1081				1085					1090					1095
Thr	Glu	Leu	Ala	Ala	Glu	Val	His	Arg	Val	Leu	Cys	Leu	Glu	Arg
1096				1100					1105					1110
Gly	His	Met	Phe	Val	Cys	Gly	Asp	Val	Thr	Met	Ala	Thr	Ser	Val
1111				1115					1120					1125
Leu	Gin	Thr	Val	Gin	Arg	lie	Leu	Ala	Thr	Glu	Gly	Asp	Met	Glu
1126				1130					1135					1140
Leu	Asp	Glu	Ala	Gly	Asp	Val	lie	Gly	Val	Leu	Arg	Asp	Gin	Gin
1141				1145					1150					1155
Arg	Tyr	His	Glu	Asp	lie	Phe	Gly	Leu	Thr	Leu	Arg	Thr	Gin	Glu
1156				1160					1165					1170
Val	Thr	Ser	Arg	lie	Arg	Thr	Gin	Ser	Phe	Ser	Leu	Gin	Glu	Arg
1171				1175					1180					1185
His	Leu	Arg	Gly	Ala	Val	Pro	Trp	Ala	Phe	Asp	Pro	Pro	Gly	Pro
1186				1190					1195					1200
Asp	Thr	Pro	Gly	Pro										
1201				1205										

Polyclonal antibodies to NO synthase may be obtained using the whole molecule of human NO synthase of the following sequence:

SEQ ID NO:2

Met	Gly	Asn	Leu	Lys	Ser	Val	Ala	Gin	Glu	Pro	Gly	Pro	Pro	Cys
1														15
Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Gly	Leu	Cys	Gly	Lys	Gin	Gly
16														30
Pro	Ala	Thr	Pro	Ala	Pro	Glu	Pro	Ser	Arg	Ala	Pro	Ala	Ser	Leu
31														45
Leu	Pro	Pro	Ala	Pro	Glu	His	Ser	Pro	Pro	Ser	Ser	Pro	Leu	Thr
46														60
Gin	Pro	Pro	Glu	Gly	Pro	Lys	Phe	Pro	Arg	Val	Lys	Asn	Trp	Glu
61														75
Val	GLys	er	lie	Thr	Tyr	Asp	Thr	Leu	Ser	Ala	Gin	Ala	Gin	Gin
76														90
Asp	Gly	Pro	Cys	Thr	Pro	Arg	Arg	Cys	Leu	GLys	er	Leu	Val	Phe
91														105
Pro	Arg	Lys	Leu	Gin	Gly	Arg	Pro	Ser	Pro	Gly	Pro	Pro	Ala	Pro
106														120
Glu	Gin	Leu	Leu	Ser	Gin	Ala	Arg	Asp	Phe	lie	Asn	Gin	Tyr	Tyr
121														135
Ser	Ser	lie	Lys	Arg	Ser	GLys	er	Gin	Ala	His	Glu	Gin	Arg	Leu
136														150
Gin	Glu	Val	Glu	Ala	Glu	Val	Ala	Ala	Thr	Gly	Thr	Tyr	Gin	Leu

151	155	160	165											
Arg	Glu	Ser	Glu	Leu	Val	Phe	Gly	Ala	Lys	Gin	Ala	Trp	Arg	Asn
166	170	175	180											
Ala	Pro	Arg	Cys	Val	Gly	Arg	lie	Gin	Trp	Gly	Lys	Leu	Gin	Val
181	185	190	195											
Phe	Asp	Ala	Arg	Asp	Cys	Arg	Ser	Ala	Gin	Glu	Met	Phe	Thr	Tyr
196	200	205	210											
lie	Cys	Asn	His	lie	Lys	Tyr	Ala	Thr	Asn	Arg	Gly	Asn	Leu	Arg
211	215	220	225											
Ser	Ala	lie	Thr	Val	Phe	Pro	Gin	Arg	Cys	Pro	Gly	Arg	Gly	Asp
226	230	235	240											
Phe	Arg	lie	Trp	Asn	Ser	Gin	Leu	Val	Arg	Tyr	Ala	Gly	Tyr	Arg
241	245	250	255											
Gin	Gin	Asp	GLy	Ser	Val	Arg	Gly	Asp	Pro	Ala	Asn	Val	Glu	lie
256	260	265	270											
Thr	Glu	Leu	Cys	lie	Gin	His	Gly	Trp	Thr	Pro	Gly	Asn	Gly	Arg
271	275	280	285											
Phe	Asp	Val	Leu	Pro	Leu	Leu	Leu	Gin	Ala	Pro	Asp	Glu	Pro	Pro
286	290	295	300											
Glu	Leu	Phe	Leu	Leu	Pro	Pro	Glu	Leu	Val	Leu	Glu	Val	Pro	Leu
301	305	310	315											
Glu	His	Pro	Thr	Leu	Glu	Trp	Phe	Ala	Ala	Leu	Gly	Leu	Arg	Trp
316	320	325	330											
Tyr	Ala	Leu	Pro	Ala	Val	Ser	Asn	Met	Leu	Leu	Glu	lie	Gly	Gly
331	335	340	345											

Leu Glu Phe Pro Ala Ala Pro Phe Ser Gly Trp Tyr Met Ser Thr
 346 350 355 360

Glu lie Gly Thr Arg Asn Leu Cys Asp Pro His Arg Tyr Asn lie
 361 365 370 375

Leu Glu Asp Val Ala Val Cys Met Asp Leu Asp Thr Arg Thr Thr
 376 380 385 390

Ser Ser Leu Trp Lys Asp Lys Ala Ala Val Glu lie Asn Val Ala
 391 395 400 405

Val Leu His Ser Tyr Gin Leu Ala Lys Val Thr lie Val Asp His
 406 410 415 420

His Ala Ala Thr Ala Ser Phe Met Lys His Leu Glu Asn Glu Gin
 421 425 430 435

Lys Ala Arg Gly Gly Cys Pro Ala Asp Trp Ala Trp lie Val Pro
 436 440 445 450

Pro lie Ser GLys er Leu Thr Pro Val Phe His Gin Glu Met Val
 451 455 460 465

Asn Tyr Phe Leu Ser Pro Ala Phe Arg Tyr Gin Pro Asp Pro Trp
 466 470 475 480

Lys Gly Ser Ala Ala Lys Gly Thr Gly lie Thr Arg Lys Lys Thr
 481 485 490 495

Phe Lys Glu Val Ala Asn Ala Val Lys lie Ser Ala Ser Leu Met
 496 500 505 510

Gly Thr Val Met Ala Lys Arg Val Lys Ala Thr lie Leu Tyr Gly
 511 515 510 525

Ser Glu Thr Gly Arg Ala Gin Ser Tyr Ala Gin Gin Leu Gly Arg

526	530	535	540											
Leu	Phe	Arg	Lys	Ala	Phe	Asp	Pro	Arg	Val	Leu	Cys	Met	Asp	Glu
541	545	550	555											
Tyr	Asp	Val	Val	Ser	Leu	Glu	His	Glu	Thr	Leu	Val	Leu	Val	Val
556	560	565	570											
Thr	Ser	Thr	Phe	Gly	Asn	Gly	Asp	Pro	Pro	Glu	Asn	Gly	Glu	Ser
571	575	580	585											
Phe	Ala	Ala	Ala	Leu	Met	Glu	Met	Ser	Gly	Pro	Tyr	Asn	Ser	Ser
586	590	595	600											
Pro	Arg	Pro	Glu	Gin	His	Lys	Ser	Tyr.	Lys	lie	Arg	Phe	Asn	Ser
601	605	610	615											
lie	Ser	Cys	Ser	Asp	Pro	Leu	Val	Ser	Ser	Trp	Arg	Arg	Lys	Arg
616	620	625	630											
Lys	Glu	Ser	Ser	Asn	Thr	Asp	Ser	Ala	Gly	Ala	Leu	Gly	Thr	Leu
631	635	640	645											
Arg	Phe	Cys	Val	Phe	Gly	Leu	GLys	er	Arg	Ala	Tyr	Pro	His	Phe
646	650	655	660											
Cys	Ala	Phe	Ala	Arg	Ala	Val	Asp	Thr	Arg	Leu	Glu	Glu	Leu	Gly
661	665	670	675											
Gly	Glu	Arg	Leu	Leu	Gin	Leu	Gly	Gin	Gly	Asp	Glu	Leu	Cys	Gly
676	680	685	690											
Gin	Glu	Glu	Ala	Phe	Arg	Gly	Trp	Ala	Gin	Ala	Ala	Phe	Gin	Ala
691	695	700	705											
Ala	Cys	Glu	Thr	Phe	Cys	Val	Gly	Glu	Asp	Ala	Lys	Ala	Ala	Ala
706	710	715	720											

Arg Asp lie Phe Ser Pro Lys Arg Ser Trp Lys Arg Gin Arg Tyr
721 **725** **730** **735**
 Arg Leu Ser Ala Gin Ala Glu Gly Leu Gin Leu Leu Pro Gly Leu
736 **740** **745** **750**
 Ile His Val His Arg Arg Lys Met Phe Gin Ala Thr Ile Arg Ser
751 **755** **760** **765**
 Val Glu Asn Leu Gin Ser Ser Lys Ser Thr Arg Ala Thr Ile Leu
766 **770** **775** **780**
 Val Arg Leu Asp Thr Gly Gly Gin Glu Gly Leu Gin Tyr Gin Pro
781 **785** **790** **795**
 Gly Asp His Ile Gly Val Cys Pro Pro Asn Arg Pro Gly Leu Val
796 **800** **805** **810**
 Glu Ala Leu Leu Ser Arg Val Glu Asp Pro Pro Ala Pro Thr Glu
811 **815** **820** **825**
 Pro Val Ala Val Glu Gin Leu Glu Lys Gly Ser Pro Gly Gly Pro
826 **830** **835** **840**
 Pro Pro Gly Trp Val Arg Asp Pro Arg Leu Pro Pro Cys Thr Leu
841 **845** **850** **855**
 Arg Gin Ala Leu Thr Phe Phe Leu Asp Ile Thr Ser Pro Pro Ser
856 **860** **865** **870**
 Pro Gin Leu Leu Arg Leu Leu Ser Thr Leu Ala Glu Glu Pro Arg
871 **875** **880** **885**
 Glu Gin Gin Glu Leu Glu Ala Leu Ser Gin Asp Pro Arg Arg Tyr
886 **890** **895** **900**
 Glu Glu Trp Lys Trp Phe Arg Cys Pro Thr Leu Leu Glu Val Leu

901	905	910	915											
Glu	Gin	Phe	Pro	Ser	Val	Ala	Leu	Pro	Ala	Pro	Leu	Leu	Leu	Thr
916	920	925	930											
Gin	Leu	Pro	Leu	Leu	Gin	Pro	Arg	Tyr	Tyr	Ser	Val	Ser	Ser	Ala
931	935	940	945											
Pro	Ser	Thr	His	Pro	Gly	Glu	lie	His	Leu	Thr	Val	Ala	Val	Leu
946	950	955	960											
Ala	Tyr	Arg	Thr	Gin	Asp	Gly	Leu	Gly	Pro	Leu	His	Tyr	Gly	Val
961	965	970	975											
Cys	Ser	Thr	Trp	Leu	Ser	Gin	Leu	Lys	Pro	Gly	Asp	Pro	Val	Pro
976	980	985	990											
Cys	Phe	lie	Arg	Gly	Ala	Pro	Ser	Phe	Arg	Leu	Pro	Pro	Asp	Pro
991	995	1000	1005											
Ser	Leu	Pro	Cys	lie	Leu	Val	Gly	Pro	Gly	Thr	Gly	lie	Ala	Pro
1006	1010	1015	1020											
Phe	Arg	Gly	Phe	Trp	Gin	Glu	Arg	Leu	His	Asp	lie	Glu	Ser	Lys
1021	1025	1030	1035											
Gly	Leu	Gin	Pro	Thr	Pro	Met	Thr	Leu	Val	Phe	Gly	Cys	Arg	Cys
1036	1040	1045	1050											
Ser	Gin	Leu	Asp	His	Leu	Tyr	Arg	Asp	Glu	Val	Gin	Asn	Ala	Gin
1051	1055	1060	1065											
Gin	Arg	Gly	Val	Phe	Gly	Arg	Val	Leu	Thr	Ala	Phe	Ser	Arg	Glu
1066	1070	1075	1080											
Pro	Asp	Asn	Pro	Lys	Thr	Tyr	Val	Gin	Asp	lie	Leu	Arg	Thr	Glu
1081	1085	1090	1095											

Leu	Ala	Ala	Glu	Val	His	Arg	Val	Leu	Cys	Leu	Glu	Arg	Gly	His
1096			1100					1105						1110
Met	Phe	Val	Cys	Gly	Asp	Val	Thr	Met	Ala	Thr	Asn	Val	Leu	Gin
1111			1115					1120						1125
Thr	Val	Gin	Arg	lie	Leu	Ala	Thr	Glu	Gly	Asp	Met	Glu	Leu	Asp
1126			1130					1135						1140
Glu	Ala	Gly	Asp	Val	lie	Gly	Val	Leu	Arg	Asp	Gin	Gin	Arg	Tyr
1141			1145					1150						1155
His	Glu	Asp	lie	Phe	Gly	Leu	Thr	Leu	Arg	Thr	Gin	Glu	Val	Thr
1156			1160					1165						1170
Ser	Arg	lie	Arg	Thr	Gin	Ser	Phe	Ser	Leu	Gin	Glu	Arg	Gin	Leu
1171			1175					1180						1185
Arg	Gly	Ala	Val	Pro	Trp	Ala	Phe	Asp	Pro	Pro	Gly	Ser	Asp	Thr
1186			1190					1195						1200
Asn	Ser	Pro												
1201	1203													

To obtain polyclonal antibodies to NO synthase, it is also possible to use a fragment of NO synthase, selected, for example, from the following sequences:

SEQ ID NO: 3

Pro Trp Ala Phe

1192 1195

SEQ ID NO: 4

Gly Ala Val Pro

1189 1192

SEQ ID NO: 5

Arg
1185

His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp Pro Pro Gly Pro
1186 1190 1195 1200

Asp Thr Pro Gly Pro
1201 1205

SEQ ID NO: 6

Ala Phe Asp Pro Pro Gly Pro
1194 1195 1200

Asp Thr Pro Gly Pro
1201 1205

SEQ ID NO: 7

His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp
1186 1190 1195 1196

SEQ ID NO: 8

His Leu Arg Gly Ala Val Pro Trp Ala Phe Asp Pro Pro Gly Pro
1186 1190 1195 1200

Asp Thr Pro Gly Pro
1201 1205

The exemplary procedure for preparation of the starting polyclonal antibodies to NO synthase may be described as follows. In 7-9 days before blood sampling, 1-3 intravenous injections of the desired antigen are made to the rabbits to increase the level of polyclonal antibodies in the rabbit blood stream. Upon immunization, blood samples are taken to test the antibody level. Typically, the maximum level of immune reaction of the soluble antigen is achieved within 40 to 60 days after the first injection of the antigen. Upon completion of the first immunization cycle, rabbits have a 30-day

rehabilitation period, after which re-immunization is performed with another 1-3 intravenous injections.

To obtain antiserum containing the desired antibodies, the immunized rabbits' blood is collected from rabbits and placed in a 50ml centrifuge tube. Product clots formed on the tube sides are removed with a wooden spatula, and a rod is placed into the clot in the tube center. The blood is then placed in a refrigerator for one night at the temperature of about 40°C. On the following day, the clot on the spatula is removed, and the remaining liquid is centrifuged for 10 min at 13,000 rotations per minute. Supernatant fluid is the target antiserum. The obtained antiserum is typically yellow. 20% of NaN₃ (weight concentration) is added in the antiserum to a final concentration of 0.02% and stored before use in frozen state at the temperature of -20°C or without NaN₃ at the temperature of -70°C. To separate the target antibodies to gamma interferon from the antiserum, the following solid phase absorption sequence is suitable:

10 ml of the antiserum of rabbits is diluted twofold with 0.15 M NaCl, after which 6.26g Na₂SO₄ is added, mixed and incubated for 12-16 hours at 4°C. The sediment is removed by centrifugation, diluted in 10ml of phosphate buffer and dialyzed against the same buffer during one night at ambient temperature. After the sediment is removed, the solution is applied to a DEAE-cellulose column balanced by phosphate buffer. The antibody fraction is determined by measuring the optical density of the eluate at 280 nm.

The isolated crude antibodies are purified using affine chromatography method by attaching the obtained antibodies to NO synthase located on the insoluble matrix of the chromatography media, with subsequent elution by concentrated aqueous salt solutions.

The resulting buffer solution is used as the initial solution for the homeopathic dilution process used to prepare the activated-potentiated form of the antibodies. The preferred concentration of the initial matrix solution of the antigen-purified polyclonal rabbit antibodies to NO synthase is 0.5 to 5.0 mg/ml, preferably, 2.0 to 3.0 mg/ml.

The polyclonal antibodies to endogenous biological molecule may also be obtained by a similar methodology to the methodology described for endothelial NO synthase antibodies using an adjuvant.

The resulting buffer solution is used as the initial solution for the homeopathic dilution process used to prepare the activated potentiated form of the antibodies.

The activated potentiated form of each component of the combination may be prepared from an initial solution by homeopathic potentization, preferably using the method of proportional concentration decrease by serial dilution of 1 part of each preceding solution (beginning with the initial solution) in 9 parts (for decimal dilution), or in 99 parts (for centesimal dilution), or in 999 parts (for millesimal dilution) of a neutral solvent, starting with a concentration of the initial solution of antibody in the solvent, preferably, water or a water-ethyl alcohol mixture, in the range from about 0.5 to about 5.0 mg/ml, coupled with external impact. Preferably, the external impact involves multiple vertical shaking (dynamization) of each dilution. Preferably, separate containers are used for each subsequent dilution up to the required potency level, or the dilution factor. This method is well-accepted in the homeopathic art. See, e.g. V. Schwabe *"Homeopathic medicines"*, M., 1967, p. 14-29, incorporated herein by reference for the purpose stated.

For example, to prepare a 12-centesimal dilution (denoted C12), one part of the initial matrix solution of antibodies to NO synthase with the concentration of 3.0 mg/ml is diluted in 99 parts of neutral aqueous or aqueous-alcohol solvent (preferably, 15%-ethyl alcohol) and then vertically shaked many times (10 and more) to create the 1st centesimal dilution (denoted as C1). The 2nd centesimal dilution (C2) is prepared from the 1st centesimal dilution C1. This procedure is repeated 11 times to prepare the 12th centesimal dilution C12. Thus, the 12th centesimal dilution C12 represents a solution obtained by 12 serial dilutions of one part of the initial matrix solution of antibodies with the concentration of 3.0 mg/ml in 99 parts of a neutral solvent in different containers, which is equivalent to the centesimal homeopathic dilution C12. Similar procedures with the relevant dilution factor are performed to obtain the desired dilutions. The intermediate dilutions may be tested in a desired biological model to check activity. The preferred activated potentiated forms for antibodies comprising the combination of the invention is a C12, C30 and C200 dilutions for each activated-potentiated form. When using the mixture of various homeopathic dilutions (primarily centesimal) of the active substance

as biologically active liquid component, each component of the composition (e.g., C12, C30, C50, C200) is prepared separately according to the above-described procedure until the next-to-last dilution is obtained (e.g., until C11, C29, and C199 respectively), and then one part of each component is added in one container according to the mixture composition and mixed with the required quantity of the solvent (e.g. with 97 parts for centesimal dilution).

It is possible to use the active substance as mixture of various homeopathic dilutions, e.g. decimal and/or centesimal (D20, C30, C100 or C12, C30, C50 or C12, C30, C200, etc.), the efficiency of which is determined experimentally by testing the dilution in a suitable biological model, for example, in models described in the examples herein.

In the course of potentiation and concentration decrease, the vertical shaking may be substituted for external exposure to ultrasound, electromagnetic field or any similar external impact procedure accepted in the homeopathic art.

The solid unit dosage form of the pharmaceutical composition of the invention may be prepared by using impregnating a solid, pharmaceutically acceptable carrier with the mixture of the activated potentiated form aqueous or aqueous-alcohol solutions of active components that are mixed, primarily in 1:1 ratio and used in liquid dosage form. Alternatively, the carrier may be impregnated consecutively with each requisite dilution.

Preferably, the pharmaceutical composition in the solid unit dosage form is prepared from granules of the pharmaceutically acceptable carrier which was previously saturated with the aqueous or aqueous-alcoholic dilutions of the activated potentiated form of antibodies. The solid dosage form may be in any form known in the pharmaceutical art, including a tablet, a capsule, a lozenge, and others. As an inactive pharmaceutical ingredients one can use glucose, sucrose, maltose, amylose, isomaltose, isomalt and other mono- oligo- and polysaccharides used in manufacturing of pharmaceuticals as well as technological mixtures of the above mentioned inactive pharmaceutical ingredients with other pharmaceutically acceptable excipients, for example isomalt, crospovidone, sodium cyclamate, sodium saccharine, anhydrous citric acid etc), including lubricants, disintegrants, binders and coloring agents. The preferred

carriers are lactose and isomalt. The pharmaceutical dosage form may further include standard pharmaceutical excipients, for example, microcrystalline cellulose, magnesium stearate and citric acid.

The example of preparation of the solid unit dosage form is set forth below. To prepare the solid oral form, 100-300 μm granules of lactose are impregnated with aqueous or aqueous-alcoholic solutions of the activated potentiated form of antibodies to histamine, activated-potentiated form of antibodies to NO synthase and the activated potentiated form of antibodies to an endogenous biological molecule in the ratio of 1 kg of antibody solution to 5 or 10 kg of lactose (1:5 to 1:10). To effect impregnation, the lactose granules are exposed to saturation irrigation in the fluidized boiling bed in a boiling bed plant (e.g. "Huttlin Pilotlab" by Hiittlin GmbH) with subsequent drying via heated air flow at a temperature below 40°C. The estimated quantity of the dried granules (10 to 34 weight parts) saturated with the activated potentiated form of antibodies is placed in the mixer, and mixed with 25 to 45 weight parts of "non-saturated" pure lactose (used for the purposes of cost reduction and simplification and acceleration of the technological process without decreasing the treatment efficiency), together with 0.1 to 1 weight parts of magnesium stearate, and 3 to 10 weight parts of microcrystalline cellulose. The obtained tablet mass is uniformly mixed, and tableted by direct dry pressing (e.g., in a Korsch - XL 400 tablet press) to form 150 to 500 mg round pills, preferably, 300 mg. After tableting, 300 mg pills are obtained that are saturated with aqueous-alcohol solution (3.0-6.0 mg/pill) of the combination of the activated-potentiated form of antibodies. Each component of the combination used to impregnate the carrier is in the form of a mixture of centesimal homeopathic dilutions, preferably, C12, C30 and C200.

While the invention is not limited to any specific theory, it is believed that the activated-potentiated form of the antibodies described herein do not contain the molecular form of the antibody in an amount sufficient to have biological activity attributed to such molecular form. The biological activity of the combination drug (combination, pharmaceutical composition) of the invention is amply demonstrated in the appended examples.

Preferably, for the purpose of treatment, the combination of the invention is administered from once daily to four times daily, preferably twice daily, each administration including one or two combination unit dosage forms.

The invention is further illustrated with reference to the appended non-limiting examples.

EXAMPLES

Example 1.

Study of effect of a complex preparation containing ultralow doses (ULD) of activated - potentiated forms of polyclonal affinity purified rabbit antibodies to brain-specific protein S-100 (anti-S100) and endothelial NO-synthase (anti-eNOS), obtained by super-dilution of initial matrix solution (concentration: 2.5 mg/ml) (100^{12} , 100^{30} , 100^{200} times), equivalent to a blend of centesimal homeopathic dilutions C12, C30, C200 (ratio: 1:1) (ULD of anti-S100+anti-eNOS), as well as its components - activated - potentiated form of polyclonal affinity purified rabbit antibodies to ultralow doses (ULD) of brain-specific protein S-100 (anti-S100), purified on antigen, obtained by super-dilution of initial matrix solution (100^{12} , 100^{30} , 100^{200} times), equivalent to a blend of centesimal homeopathic dilution C12, C30, C200, and activated - potentiated form of polyclonal rabbit antibodies to ultralow dose of endothelial NO-synthase (ULD of anti-eNOS), obtained by super-dilution of initial matrix solution (100^{12} , 100^{30} , 100^{200} times), equivalent to a blend of centesimal homeopathic dilution C12, C30, C200 *in vitro* on binding of standard ligand [³H]pentazocine to human recombinant σ 1 receptor was evaluated using radioligand method. Potentiated distilled water (blend of homeopathic dilutions C12+C30+C200) was used as test preparations control.

Sigma-1 (σ 1) receptor - an intracellular one which is localized in the cells of central nervous system, the cells of the most of peripheral tissues and immune component cells. Receptors exhibit a unique ability to be translocated which is caused by many psychotropic medications. The dynamics of sigma-1 receptors is directly linked to various influences which are performed by preparations acting to the sigma-1 receptors. These effects include the regulation of activity channels, ecocytosis, signal transferring,

remodeling of the plasma membrane (formation of rafts) and lipid transportation / metabolism. All this can contribute to the plasticity of neurons in a brain. There is evidence that the sigma-1 receptors have a modulating effect on all the major neuromediator systems: noradrenergic, serotonergic, dopaminergic, cholinergic systems and NMDA- adjustable glutamate effects. Sigma-1 receptor plays an important role in the pathophysiology of neurodegenerative diseases (e.g., Alzheimer's disease, Parkinson), psychiatric and affective disorders and stroke; and it also takes part in the processes of learning and memory. In this regard, the ability of drugs to influence the efficiency of interaction of ligands with sigma-1 receptor indicates on the presence of neuroprotective, anti-ischemic, anxiolytic, antidepressant and anti astenic components in the spectrum of its pharmacological activity that allows the consideration of these drugs as effective preparations particularly for the treatment of cerebrovascular diseases.

During the test (to measure total binding) 20 μ l of complex preparation of ULD of anti-S100+anti-eNOS or 10 μ l of ULD of AB to S100 or 10 μ l of ULD of AB to NOS were transferred in the incubation medium. Thus, the quantity of ULD of anti-S100+anti-eNOS, transferred into the test basin when testing the complex preparation was identical to that of ULD of AB to S100 and ULD of AB to NOS tested as monopreparations, which allow comparing the efficiency of the preparation to its separate components. 20 μ l and 10 μ l of potentiated water were transferred in the incubation medium.

Further, 160 μ l (about 200 μ g of protein) of Jurkat cell line membranes homogenate (human leukemic T-lymphocyte line), and finally, 20 μ l of tritium-labeled radioligand [3 H]pentazocine (15 nm) were transferred.

In order to measure non-specific binding, 20 μ l of non-labeled ligand- haloperidol (10 μ M) were transferred in the incubation medium instead of the preparations or potentiated water.

Radioactivity was measured using a scintillometer (Topcount, Packard) and scintillation blend (Microscint 0, Packard) following the incubation within 120 minutes at 22°C in 50 mM Tris-HCl buffer (pH = 7,4) and filtration using fiberglass filters (GF/B,

Packard). Specific binding (during the test or control) was calculated as a difference between total (during the test or control) and non-specific binding.

Results are represented as percentage of specific binding inhibition in control (distilled water was used as control) (Table 1).

Table 1.

Test group	Quantity per test basin	% of radioligand specific binding in control			% of radioligand binding inhibition in control
		1 st test	2 nd test	Average	
ULD of anti-S100+anti-eNOS	20 µl	48,4	35,5	42,0	58,0
ULD of anti-S100	10 µl	67,3	63,1	65,2	34,8
ULD of anti-eNOS	10 µl	147,5	161,1	154,3	-54,3
Potentiated water	20 µl	98,1	75,8	86,9	13,1
Potentiated water	10 µl	140,1	106,2	123,2	-23,2

Effect of the preparations and potentiated water on binding of standard ligand [³H]pentazocine to human recombinant σ 1 receptor

Note: % of specific binding in control = (specific binding during the test/ specific binding in control)* 100%;

% of specific binding inhibition in control = 100% - (specific binding during the test/ specific binding in control) * 100%.

The outcomes reflecting inhibition above 50% represent significant effects of the tested compounds; inhibition from 25% to 50% confirm mild to moderate effects; inhibition less than 25% is considered to be insignificant effect of the tested compound and is within background level.

Therefore, the conditions of this test model showed that the complex preparation of ULD of anti-S100+anti-eNOS is more efficient than its separate components (ULD of anti-S100 and ULD of anti-eNOS) in inhibiting the binding of standard radioligand [³H]pentazocine to human recombinant σ 1 receptor; ULD of anti-S100, transferred into the test basin, namely 10 μ l, inhibit the binding of standard radioligand [³H]pentazocine to human recombinant σ 1 receptor, but the effect intensity is inferior to that of the complex preparation of ULD of anti-S100+anti-eNOS; ULD of anti-eNOS, transferred into the test basin, namely 10 μ l, had no effect on the binding of standard radioligand [³H]pentazocine to human recombinant σ 1 receptor; potentiated water, transferred into the test basin, namely 10 μ l or 20 μ l, had no effect on the binding of standard radioligand [³H]pentazocine to human recombinant σ 1 receptor.

Example 2.

Alzheimer's disease (AD) is a neurodegenerative disease that is characterized by lowering of cognitive functions, memory deterioration, confused consciousness, and emotional changes. Although the main cause of this pathology is nowadays considered the accumulation of beta amyloid which leads to the formation of beta-amyloid plaques and neurofibrillary tangles in brain tissues; AD is also accompanied by a deficiency of cholinergic system. This is the basis of a most common way of modeling of AD in animals with the help of antagonist of cholinergic system of scopolamine. Injection of scopolamine into experimental animals (usually rats or mice) interrupts the ability to learn and leads to deterioration of memory.

Various methods were used to assess cognitive functions of rats and mice, including Morris water maze. The essence of this test is that the animals are released

into a container with cloudy water from different points are forced to look for a hidden fixed platform. The advantage of this method is that it allows the researcher to monitor the process of animal training (the formation of ideas about the spatial alignment of the platform no matter where the animal was placed in the water) so as to assess the memory strength (for this the test is conducted when the platform is removed).

The effectiveness in rats with Scopolamine amnesia of the combination pharmaceutical composition of the present invention containing activated-potentiated forms of polyclonal affinity purified on antigen of rabbit brain-specific proteins S-100 (anti-S100) and to endothelial NO-syntheses (anti-eNOS) in ultra low doses (ULD) obtained by super dilution of storage stock solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, equivalent to centesimal homeopathic dilutions C12, C30, C200 (ULD anti-S100 + anti-eNOS) is studied.

In a study of the effectiveness of the drug ULD anti-S100 + anti-eNOS in rats with scopolamine amnesia (a model of Alzheimer's disease) 48 male rats of the Rj: Wistar (Han) line (weight 180-280g) were used. During 4 days the rats were subdermally injecting with normal saline (n = 12, intact) or scopolamine in doze of 0.5 mg / kg (n = 36) (scopolamine-induced amnesia). Rats with scopolamine-induced amnesia were divided into three groups and administered with distilled water (7.5 ml / kg, n = 12, control group 1), or ULD anti-S100 (7.5 ml / kg, n = 12, group 2) or ULD anti-S100 + anti-eNOS (7.5 ml / kg, n = 12, group 3) intragastrically for 9 days (4 days prior to the injection of scopolamine, 4 days against the background of scopolamine and 1 day after the last scopolamine injection).

The training session in the Morris water maze was conducted within 4 days of the scopolamine injection through 60 minutes after administration of tested drugs and 30 minutes after administration of scopolamine (4 sequential tests at interval of 60 seconds). Morris' maze is a round reservoir (diameter - 150 cm, height - 45 cm) at 30 cm filled with water (26-28 ° C). At 18 cm from the edge of the container there is hidden platform (diameter - 15 cm) buried on 1.5 cm below the water level. Cloudy water made by adding a non-toxic dye (e.g., milk powder) makes the platform invisible. For each test

the animal was placed in a maze in one of the initial points that are equidistant from the hidden platform and the animal was allowed to find the platform. If the animal could not find the platform within 120 seconds, the animal was put on the platform and left for 60 seconds and the test was restarted. During the four tests in random order the animals began to walk through the maze twice from each starting point. The tests were recorded on videotape and then analyzed for distance overcomes searching the platform in each trial and the latent period of searching for the platform. On day 5 the test was performed: the platform was removed from the maze and rats were given free float for 60 seconds. The time spent in the place where the platform used to be was recorded.

The administration of scopolamine significantly worsened the ability of animals to learn. In the control group the time spent by animals searching for platforms and the distance that animals swam searching for the platform, significantly increased (Table 2, 3). The test shows that the memory of animals in the control group worsened: the animals in this group spent less time in the place where the platform used to be located than intact animals (Table 4). The administration of ULD anti-S100 didn't lead to improvement of the studied parameters (Tables 2, 3, 4). The administration of ULD anti-Si 00 + anti-eNOS led to some improvement in learning which resulted in a shortening of the latent time of the platform search time (Table 2) and covered distance (Table 3) within 4 days of training and an improvement of memory as reflected in increase of the time spent in a place where the platform used to be located (Table 4).

Table 2.

Latent period of the platform search, sec

Group	Training			
	1 st day	2 nd day	3 rd day	4 th day
Intact, n=12	54.7±6.2	30.8±2.8	26.9±5.1	20.5±3.6

Control, n=12	100.1 \pm 6.8 ***	92.4 \pm 9.3 ***	81.4 \pm 10.7 ***	77.7 \pm 9.4 ***
ULD anti-S100, n=12	106.8 \pm 7.0	99.3 \pm 7.8	95.6 \pm 9.0	80.4 \pm 11.1
ULD anti-S100 + anti-eNOS, n=12	94.4 \pm 7.2	90.7 \pm 8.2	78.3 \pm 8.6	60.1 \pm 10.2

*** - difference from intact is significant, p<0.05

Table 3.

Distance overcome to search the platform, cm

Group	Training			
	1 st day	2 nd day	3 rd day	4 th day
Intact, n=12	1055.7 \pm 94.6	659.5 \pm 62.2	564.8 \pm 119.3	406.1 \pm 61.2
Control, n=12	2587.1 \pm 217.2 ***	2559.6 \pm 250.5 ***	2397.9 \pm 312.6	2366.1 \pm 293.8 ***
ULD anti-S100, n=12	2797.2 \pm 208.9	2865.2 \pm 255.1	2857.0 \pm 300.8	2457.4 \pm 344.4
ULD anti-S100 + anti-eNOS, n=12	2434.3 \pm 222.8	2529.9 \pm 282.7	2344.2 \pm 283.0	1905.1 \pm 343.7

*** - difference from intact is significant, p<0.05

Table 4.

Time spent in a place where the platform used to be located, sec.

Group	Test		
	0-30 sec.	30-60 sec.	0-60 sec.
Intact, n=12	40.8 \pm 4.1	36.8 \pm 3.6	38.5 \pm 2.6

Control, n=12	18.4+2.8***	18.8±1.9***	18.811.7***
ULD anti-S100, n=12	13.3±2.1	21.5±2.6	17.611.3
ULD anti-S100 + anti-eNOS, n=12	19.1±4.8	23.8±2.2	21.2±2.5

*** - difference from intact is significant, p<0.05

Thus, in model of Alzheimer's disease, the administration of the complex ULD anti-S100 + anti-eNOS was more effective in comparison with administration of ULD anti-S100 and vehicle.

Example 3.

The preclinical research studied the ultra low doses (ULD) of activated - potentiated forms of polyclonal affinity purified rabbit antibodies to brain-specific protein S-100 (anti-S100) purified on antigen, and endothelial NO-synthase (anti-eNOS), obtained by super-dilution of initial matrix solution (concentration: 2,5 mg/ml) (100¹², 100³⁰, 100²⁰⁰ times), equivalent to a mixture of centesimal homeopathic dilutions C12, C30, C200 (ratio: 1:1) (ULD anti-S100+anti-eNOS) in treating ischemic stroke caused by prefrontal cerebrocortical photothrombosis in rats.

Acute cerebrovascular disease (brain stroke) ranks third among lethality causes in developed countries and one of the main causes of disability in humans (*Gusev E.I., 2003; Janardhan V., Qureshi A.I., 2004*).

The photo-induced thrombosis model meets almost all requirements to the experimental model of focal cerebral ischemia. The method developed by Watson (*Watson B. et al, 1985*) is based on the effect of light with wavelength 560 nm on photosensitive pigment Bengal rose introduced into the blood flow. Active oxygen forms are created and caused increase in adhesiveness of endothelium cells and platelets, and formation of clots closing vascular lumens. The method of ischemic brain lesion

induction by using photo-induced thrombosis is technically simple and to close to clinical forms of ischemic brain stroke. A great advantage of this model is that it is non-invasive, i.e. does not require craniotomy and, therefore, more accurately reproduces clinical picture of cerebral thrombosis.

Thirty seven male Wistar rats (weight: 150-180 g; age: 2-3 months) were included in the study of the activity of ULD anti-S100+anti-eNOS in rats with ischemic stroke caused by prefrontal cerebrocortical photothrombosis. Bilateral focal ischemic injury in prefrontal cerebral cortex in rats was induced using the photochemical thrombosis method by Watson (*Watson B. D. et al, 1985*) as modified by I.V. Viktorov (*Romanova G.A. et al, 1998*). Bengal rose (3% solution) was injected in the jugular vein of anesthetized rats (n=37) (anesthesia: chloral hydrate 300 mg/kg, intraperitoneally). Using a fiber optic bundle (3 cm in diameter) the light beam from halogen lamp (24 V, 250 W) was delivered to the skull surface above the frontal cortex of the left and right cerebral hemispheres to induce photothrombosis. Sham-operated rats (n=6) were subject to the same procedure except administration of bengal rose and exposure to halogen lamp light. The intact group included 6 rats.

Five days before and 9 days after stoke induction the following preparations were administered to rats with photothrombosis: distilled water (control-photothrombosis, 5 ml/kg daily, n=12), ULD anti-S100 (5 ml/kg daily, n=7) or ULD anti-S100+anti-eNOS (5 ml/kg daily, n=6). On Day 8 after the operation (or sham operation) conditioned passive avoidance reflex (CPAR) test was performed to assess learning capability and memory in rats. Rats were placed in a unit consisting of illuminated site and connected dark chamber, where animals were exposed to electric foot-shock of 0,45 mA due to which usually preferred dark chamber became dangerous. Development of conditioned passive avoidance reflex was tested on the next day. At that, rats were placed in the illuminated chamber. Latent period of the first entry in the dark chamber was recorded. If a rat avoided the dark chamber for a long time, a conclusion was made that it remembered the danger (electric shock). The longer the latent period of entry in the dark chamber, the better the memory.

Volume of the stroke lesion was morphologically assessed in a proportion of rats of experimental groups on Day 9.

In control rats photothrombosis caused formation of a large stroke area and, therefore, leaded to memory impairment: CPAR reproduction worsened by 9.6% compared to intact rats and by 22.9% compared to sham-operated (Table 5). Administration of ULD anti-S100 reduced the stroke volume by 42.2% and improved memory by 14.0% compared to control-photothrombosis group. Administration of ULD anti-S100+anti-eNOS was more effective: the stroke volume reduced by 44.0% , and conditioned reflex reproduction - by 33.4% compared to control-photothrombosis group.

Therefore, administration of the complex preparation of ULD anti-S100 + anti-eNOS was more efficient than monocomponent preparation of ULD anti-S100.

Table 5.

	Volume of focal stroke (mm ³); the number of animals	Latent period of CPAR (seconds), the number of animals
Intact	-	135.8 ± 28.8; n=6
Sham-operated	-	159.3 ± 18.7; n=6
Control-photothrombosis	3.41 ± 0.5; n=9	122.8 ± 20.9; n=12
Photothrombosis + ULD anti-S100	1.97 ± 0.6; n=4	140.0 ± 26.5; n=7
Photothrombosis + ULD anti-S100+anti-eNOS	1.91 ± 0.5; n=4	163.8 ± 16.2; n=6

Example 4.

Study of the combination of "activated" potentiated forms of antibodies to a C-terminal fragment of the angiotensin II AT1-receptor, in a mixture of homeopathic

dilutions of C12, C30, C200, with the activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in SHR rats in a model of hypertension.

The combination of the "activated" potentiated form of antibodies to a C-terminal fragment of the angiotensin II AT1-receptor, in a mixture of homeopathic dilutions of C12, C30, C200, and the activated potentiated form of antibodies to endothelial NO-synthase in a mixture of homeopathic dilutions of C12, C30, C200, was studied, in solution form, in the SHR rat hypertension model. Investigations were conducted on 40 SHR line male rats from (weight 350+50 g, age 4.5 - 5 months) with hypertension, which were divided into 4 groups of 10 animals each.

For 28 days, the animals were treated as follows. Group 1 - 2.5 ml/kg of the potentiated activated form of antibodies to a C-terminal fragment AT1 of human angiotensin II receptor (a mixture of aqueous dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water, Group 2 - 2.5 ml/kg of the potentiated activated form of antibodies to endothelial NO-synthase (a mixture of aqueous dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water, Group 3 - 5 ml/kg of the combination pharmaceutical composition (a mixture of aqueous dilutions C12, C30, C200 for each component), and Group 4 - 5 ml/kg of distilled water.

Systolic blood pressure (SBP) of awake rats was measured with the aid of an indirect method in a tail artery (using a cuff) once a week and 9 hours after the last administration of medicines.

All tested compositions demonstrated hypotensive effect ($p<0.05$): by 28th day, systolic blood pressure (SBD) decreased in comparison with the initial level in Group 1 by - 20.6%; in Group 2 by 14.4%; in Group 3 by 27.6%. In the control Group 4, SBD changes were 1.6% in comparison with the initial values. The results demonstrate a clear synergistic hypotensive effect of the combination pharmaceutical composition.

Example 5.

Study of the combination of the activated potentiated forms of antibodies to a C-terminal fragment of angiotensin II AT1-receptor, in a mixture of homeopathic dilutions of

C12, C30, C200, with the activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in NISAG rats in a model of hypertension.

The combination of the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1-receptor, in a mixture of homeopathic dilutions of C12, C30, C200, and the activated potentiated form of antibodies to endothelial NO-synthase in a mixture of homeopathic dilutions of C12, C30, C200, was studied, in solution form, in the NISAG rat hypertension model. Investigations were conducted on 50 NISAG line male rats (weight 300 g, age 4 months) with hereditary stipulated stress-sensitive arterial hypertension, which were divided into 5 groups by 10 animals each.

The animals were given per orally, once a day and for 28 days, the following medications: Group 1 - 2.5 ml/kg of the activated potentiated form of antibodies to a C-terminal fragment AT1 of human angiotensin II receptor (a mixture of dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water; Group 2 - 2.5 ml/kg of the activated potentiated form of antibodies to endothelial NO-synthase (a mixture of dilutions C12, C30, C200) in combination with 2.5 ml/kg of distilled water; Group 3 - 5 ml/kg of the combination pharmaceutical composition (a mixture of homeopathic aqueous dilutions C12, C30, C200 of each component); Group 4 - 5 ml/kg (10 ml/kg dose) of the comparison drug (losartan); and Group 5 - 5 ml/kg of distilled water.

Two times a week, 2 to 6 hours after administration of VSD antibodies and losartan, systolic blood pressure (SBP) was measured by an indirect method in a tail artery (using a cuff). The Table 6 shows the dynamics of changes in systolic blood pressure in NISAG line rats, measured by indirect method.

Table 6.

Indicator	Initial SBP in mmHg	SBP after 28 days of medicine administration in mmHg	Δ in comparison with the initial level, in mmHg	% of the initial level
VSD antibodies to C-terminal fragment AT1 of	176	150	-26	-14.7%

human angiotensin II receptor				
VSD antibodies to endothelial NO-synthase	175	164.5	-10.5	-6%
Combination medicine on the basis of VSD antibodies to C-tailed fragment AT1 of angiotensin II receptor and to endothelial NO-synthase	179.5	140	-39.5	-22%
Losartan	173.5	140.5	-33	-19%
Control (distilled water)	181	178	-3	-1.6%

Example 6.

The experimental study investigated the effects of antibodies to the C-terminal fragment to the insulin receptor β -subunit affinity purified on antigen, in ultra-low dose, obtained by super dilution of the initial matrix solution 100^{12} , 100^{30} , 100^{200} times (ULD anti-IR), antibodies to endothelial NO-synthase affinity purified on antigen, in ultra-low dose, obtained by hyper-dilution of the initial matrix solution 100^{12} , 100^{30} , 100^{200} (ULD anti-ULD anti-eNOS), as well as the combination of ultra-low doses of antibodies to the C-terminal fragment to the insulin receptor β -subunit and ultra-low dose of antibodies to endothelial NO-synthase (ULD anti-IR + ULD anti-eNOS).

In the study, 150 male Wistar were used (weight at beginning of study 250-300 g, age 3.5-4 months). 10 rats were intact. The rest were intravenously injected with streptozotocin at the dose of 50 mg/kg (experimental model of diabetes mellitus). 72 hours after injection of streptozotocin, rats with blood plasma glucose level not less than 12 mmol/l were selected, divided into 7 groups (20 rats in each), which over 21 days were given distilled water (5 ml/kg/day, once daily intragastrically), insulin® (8 units/kg/day, subcutaneously), Rosiglitazone® (8 mg/kg/day, twice daily intragastrically), ULD anti-IR (2.5 ml/kg/day in a volume of 5 ml/kg/day, once daily intragastrically), ULD anti-IR + ULD anti-eNOS (5 ml/kg/day, once daily intragastrically), and also Rosiglitazone® and insulin® together or ULD anti-IR + ULD anti-eNOS and insulin®,

according to regimes corresponding to each preparation (as described above). Intact rats received distilled water in the same volume. On days 7, 14 and 21 of injection of preparations in rats, fasting blood plasma glucose level measured with enzymatic method (glucose oxidase method) with utilization of "glucose FKD" kits (Russia).

Oral glucose tolerance test (OGTT) was performed on day 14 of the study (day 14 of administration of preparation) according to standard method (Du Vigneaud and Karr, 1925). The rats were starving at water for 18 hours. 60 min before the test they were last given test substances. Intact rats received distilled water in the same volume. Glucose was administered per os 50% w/w water glucose solution (1 g/kg of rat weight). Serum glucose of blood sample from tail vein was measured by using "Glucose FKD" kit (OOO "Pharamaceutical and clinical diagnostics, Russia, www.fkd.ru) at 0, 30, 60, 90, 120 min. Mean area under the curve (AUC) concentration of blood glucose over time was calculated.

Injection of streptozotocin led to a substantial increase in blood plasma glucose of rats in comparison with intact animals (18 mmol/l versus 3.5 mmol/l, p<0.05). In the ULD anti-IR group, on day 7, 14 and 21 of injection of preparation, glucose level was lower than in the control group by 22-28% on average; however, differences did not reach a statistically significant level. The combination of ULD anti-IR and anti-eNOS was more efficacious; the decrease in glucose level on days 14 and 21 of the experiment were 47% and 42%, respectively (p<0.05 versus control). The reference preparation, Rosiglitazone, also lowered glucose level by day 14 and 21 of the experiment; at that, the effect reached statistical significance on day 14 of the experiment only (36%, p<0.05 versus control).

Insulin, injected at $\frac{1}{2}$ of the effective dose (selected in the preliminary study) most effectively lowered glucose level in all observation periods (down to the level of the intact control). (Figure 1). It should be taken into account that short-acting insulin was used in the study and blood plasma glucose was measured 1 hour after its injection, which also influenced the effect of the $\frac{1}{2}$ insulin dose on blood glucose level. Against this background it was not possible to fully determine what the effect of the combined use of insulin and rosiglitazone or insulin and complex ULD anti-IR + anti-eNOS is.

Glucose tolerance disturbance (reduction in glucose utilization by the body) is one of the most important indicators in diagnostic and treatment of diabetes mellitus. In intact animals, in the oral glucose tolerance test (day 14 of injection of preparations), complex preparation ULD anti-IR + ULD anti-eNOS and insulin most effectively increased glucose tolerance when administered alone. Rosiglitazone also reduced the area under concentration over time curve (increased glucose tolerance); however, its efficacy was not statistically significant versus the control group (Figure 2).

Example 7.

The experimental study investigated the effects of antibodies to the C-terminal fragment to the insulin receptor β -subunit affinity purified on antigen, in ultra-low dose, obtained by super dilution of the initial matrix solution 100¹², 100³⁰, 100²⁰⁰ times (ULD anti-IR), antibodies to endothelial NO-synthase affinity purified on antigen, in ultra-low dose, obtained by hyper-dilution of the initial matrix solution 100¹², 100³⁰, 100²⁰⁰ (ULD anti-ULD anti-eNOS), as well as the combination of ultra-low doses of antibodies to the C-terminal fragment to the insulin receptor β -subunit and ultra-low dose of antibodies to endothelial NO-synthase (ULD anti-IR + ULD anti-eNOS).

In the study, 36 male Goto-Kakizaki rats were used (weight at beginning of study 250-280 g, age 10-12 weeks). Rats of this line are characterized by spontaneous development of non-insulin-dependent diabetes. The animals were divided into 3 groups (12 rats in each) and received either distilled water (5 ml/kg, once daily intragastrically), or ULD anti-IR (2.5 ml/kg once daily intragastrically), or ULD anti-IR + ULD anti-eNOS (5 ml/kg, once daily intragastrically) for 28 days. Blood plasma glucose level was measured with the help of a glucose analyzer (Beckman, Fullerton, California, USA) before beginning injection of preparations and on day 4, 8, 12, 16, 20, 24, 28 of injection of preparations. On day 28, a glucose tolerance test was carried out (glucose p.o., 1 g/kg).

Injection of ULD anti-IR led to a significant ($p<0.05$) drop in blood plasma glucose level of rats; however, the use of complex ULD anti-IR + ULD anti-eNOS was more efficacious ($p<0.001$ versus control) (Figure 3).

The results were confirmed by glucose tolerance test data carried out on day 28 of injection of preparations (Figure 4). Injection of ULD anti-IR led to an increase in glucose tolerance (statistically insignificant drop by 44% AUC versus control). At the same time, the reduction in this parameter (AUC) caused by injection of complex ULD anti-IR + ULD anti-eNOS was 62% and it was statistically significant versus control ($p<0.05$).

Example 8.

The following preparation were used: 300 mg tablets impregnated with aqueous alcoholic solution (3 mg / tab.) activated - potentiated form of polyclonal rabbit brain-specific proteins antibodies S-100, purified on an antigen, in ultra low dose (ULD anti-Si 00) received by super dilution of initial solution (with concentration of 2.5 mg / ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200; 300 mg tablets impregnated with pharmaceutical composition contained aqueous-alcohol solutions of (6 mg/tab) activated - potentiated forms of polyclonal affinity purified rabbit antibodies to brain-specific protein S-100 (anti S-100) and to eNOS (anti-eNOS) in ultra low dose (ULD), received by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200; (ULD anti-S100+anti-eNOS); 300 mg tablets impregnated with aqueous-alcohol solution (3 mg/tab.) of activated - potentiated form of polyclonal rabbit eNOS antibodies purified on antigen in ultra low dose (ULD anti-eNOS), received by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200; and as placebo 300 mg tablets containing excipients: lactose (lactose monohydrate) - 267 mg, microcrystal cellulose - 30 mg, magnesium stearate - 3 mg.

The effectiveness of the studied drugs in the treatment of dizziness (vertigo) and other symptoms of motion sickness was evaluated on kinetosis model or motion diseases/motion sicknesses which occurs by various vestibular vegetative disorders.

Dizziness is the typical sign of lesion of the vestibular analyzer of various genesis including dysfunction of the vestibular nerve and cochlear system, circulatory embarrassment in vertebral basilar system, pathology of the central nervous system (CNS), etc. Dizziness as a manifestation of kinetosis accompanied with other vestibular-vegetative disorders which include three types of reactions: the vestibular-motor (nystagmus and the reaction of deviation), vestibular-sensory (in addition to dizziness, nystagmus is (or reaction of post rotation), defensive movements) and vegetative (nausea, vomiting, sweating, palpitation, heat feeling, pulse and blood pressure fluctuations).

Double blind placebo controlled comparative study were conducted in parallel groups consisting of 15 somatically healthy subjects - males and females aged from 15 to 60 years (mean age 33.3 ± 0.75 years) with low (n=5; 33%) or mean (n=10; 67%) degree of motion sickness resistance in order to test anti motion sickness properties of various compositions. Group 1 was given ULD anti-S100+anti-eNOS, Group 2 was given ULD anti-S100 and Group 3 was given anti-eNOS.

To simulate the condition of motion sickness and evaluate the effectiveness of studied drugs the most appropriate and recognized kinetosis models - test with a continuous cumulative effect of accelerations by Coriolis (CCEAC) was used. Initial tolerance of CCEAC test in all study subjects was not more than 5 minutes. Vestibular-vegetative disorders provoked by kinetic effect (CCEAC) were registered with usage of complex of diagnostic methods including subject's examination, quantitative evaluation of disorders of vestibular-vegetative sensitivity (Halle scale), analysis of heart rate variability (HRV), and self-esteem of functional condition (WBAM - well-being, activity, and mood). As the criteria of efficiency of conducted therapy the dynamics of tolerance and extent of recovery period at kinetic influence were assessed as well as alteration of indexes' evidence of sensory-motor reactions (nystagmus), HRV indexes (with usage of Biocom Wellness Scan system, developed by AWS, LLC in accordance with International Standard of European Cardiologists Association and North American Electrophysiology Association) and WBAM data. The safety criteria were character, evidence and terms of emergence of probable adverse events (AE) in the treatment

period connected with medication intake; influence of studied drugs for indexes which characterize the function of central nervous system (CNS) (reaction on moving object (RMO)), the time of simple motor reaction (TSMR); the dynamics of physical and functional factors (heart rate (HR), systolic and diastolic blood pressure (SBP, DBP), Stange's test; exercise tolerance (index of Harvard step-test). Safety was assessed after single dose administration and after 7-day course administration of the combination ULD anti-S-100 and ULD anti-eNOS.

All the subjects during 1 month before being involved into the study had not taken any drugs. After screening the subjects were randomized into 4 groups (Group 1 - ULD anti-S100+anti-eNOS, Group 2 - ULD anti-S100, Group 3 - ULD anti-eNOS, and Group 4 - placebo).

On the first day of the study (Visit 1) the initial functional and psycho-physiological state of the subjects was registered, the subjects were then given 5 tablets of the respective ULD antibodies, followed by administration of the CCEAC test. The duration of the test was registered; vegetative-vestibular disorders and AEs related to motion sickness were detected with the help of a complex diagnostic examination. In the next 2-6 days the subject were given 1 tablet three times a day of the prescribed drug. At the 7th day (Visit 2) the subjects were given the same dosage as on the first day (Visit 1). The complex of diagnostic studies was conducted before and after the CCEAC test. The study was organized in such way as study crew would work only with one subject. The study was parallel and conducted in the first half of a day with participation of, as a rule, 4 persons in a day, one person for drug or placebo. The next three weeks were washout period, at the end of which the new drug or placebo was prescribed to subjects of each group; the cycle of study was being repeated (Visit 1, the course intake of a drug; Visit 2). Thus, during the study each subject took part in four cycles of study. That is, each subject participated in each group with a three-week washout period between each cycle. This allowed the researcher to level the influence of individual peculiarities of a test person on the treatment effect. The analysis of drug efficiency was conducted on the data of all the test subjects who has completed the full course of studied drug intake according to study protocol (n=15).

The evidence factors of symptoms of motion sickness (vertigo, nausea, inactivity, skin pallor, sweatiness, etc.) after kinetic influence (CCEAC) against the background of single-day intake of studied drugs evidenced that all the study subjects have gained roughly the same state of motion sickness as far as the evidence of assessed symptoms of vegetative dysfunction on Halle's scale by physician-researcher was not differed significantly in all groups (Table 7, Visit 1). However, while the kinetic affect which cause similar symptoms of motion sickness was different in four groups and was dependent on the drug which was taken by the subjects of the study (Table 8, Visit 1). One-day intake of ULD anti-S100 + anti-eNOS preparation led to most clearly anti motion sickness effect which manifested itself not only in significantly more time of tolerance of CCEAC test (104.10 ± 13.14 sec. vs. 68.50 ± 6.57 sec. - in the group of ULD anti-S100; 75.00 ± 6.79 sec. - in the group of ULD anti-eNOS and 61.30 ± 3.15 sec. - in the placebo group) but also in the least time of nystagmus (9.90 ± 1.20 sec. vs. 13.50 ± 1.51 ; 16.10 ± 1.68 and 13.30 ± 1.12 sec., respectively) and in maximal rapid recovery (96.90 ± 13.54 sec. vs. 194.20 ± 18.45 ; 202.50 ± 21.72 and 241.70 ± 38.41 sec., respectively).

Roughly similar indexes were registered at Visit 2 after receiving a course of drugs. To achieve the similar symptoms of motion sickness (Table 7, Visit 2) the longest time of kinetic impact was applied to the subjects who has been receiving the composition of ULD anti-S100 + anti-eNOS (Table 8, Visit 2) for 7 days. The most pronounced anti motion sickness effect of the composition of ULD anti-S100 + anti-eNOS was expressed in significantly less time of nystagmus (9.50 ± 1.38 sec, $p < 0.01$) and duration of the recovery period (117.90 ± 15.65 sec; $p < 0.01$). The monocomponent preparation ULD anti-S100 had anti motion sickness action as better indexes of tolerance of CCEAC test, recovery time of nystagmus and recovery than in the placebo group evidenced (Table 8, Visits 1 and 2), but the efficacy of ULD anti-S100 was inferior to composition of ULD anti-S100 + anti-eNOS. The monocomponent preparation ULD anti-eNOS did not show anti motion sickness effect since the results of CCEAC tests and subsequent recovery period had no significant difference from the placebo group (Table 8, Visits 1 and 2). Comparative analysis of indexes of CCEAC test in the groups

of ULD anti-S100 + anti-eNOS and ULD anti-S100 in one-day intake of the drugs has shown that the addition of ULD anti-eNOS increased the tolerance of the kinetic effect on the 52%, reduced the nystagmus time on 27% and contributed to the reduction the recovery period after the end of the kinetic effect on 50% including the duration of dizziness - on **49%**. However, the greatest contribution of the component of ULD anti-eNOS introduced the effectiveness of combined preparation (compositions of ULD anti-Si 00 + anti-eNOS) in course intake of a drug which was expressed in excess of 30% of the result achieved in the group of ULD anti-S100 by factors of tolerance of kinetic effect and nystagmus duration (in each of the parameters). In addition, the growth of the effect on Visit 2 by indexes of tolerance of CCEAC test and duration of the nystagmus in relation to data of Visit 1 when taking the composition ULD anti-S100 + anti-eNOS in comparison to monocomponent preparation ULD anti-S100 was expressed in a greater degree as confirmed by alteration of these indexes on 30% and **4%** (versus 21% and 0% in the ULD anti-S100 group). In assessing the effectiveness of anti motion sickness properties of drugs the special attention was paid to the possible impact of drugs on the stability of autonomic nervous system (ANS) in particular, shifting of the balance between its sympathetic and parasympathetic divisions. For this purpose, at each visit HRV parameters were analyzed at the rest condition and when performing the functional tests (breathing and orthostatic tests).

Table 7

Indexes of Halle's scale depending on applied preparation after the performance of CCEAC test

Preparation	Halle's scale (points)	
	Visit 1	Visit 2
	(one-day intake) (n=15; M±SE)	(course intake) (n=15; M±SE)
ULD anti-S100 + anti-eNOS	12.00±0.63	12.30±0.59

ULD anti-S100	13.30±0.65	12.30±0.46
ULD anti-eNOS	13.10±0.78	12.00±0.55
Placebo	13.40±0.77	13.30±0.45

Table 8

The dynamics of indexes of CCEAC test depending on applied preparation

Preparation	Visit 1 (one-day intake)		
	Tolerance of CCEAC test, sec. (n=15; M±SD)	Nystagmus time, sec. (n=15; M±SD)	Recovery time, sec. (n=15; M±SD)
ULD anti-S100 + anti-eNOS	104.10±13.14 **	9.90±1.20 *	96.90±13.54 ***
ULD anti-S100	68.50±6.57 ×	13.50±1.51	194.20±18.45 ***
ULD anti-eNOS	75.00±6.79	16.10±1.68	202.50±21.72 ***
Placebo	61.30±3.15	13.30±1.12	241.70±38.41
P value on Kruskal-Wallis test ¹	0.0182	0.0658	0,0001
Visit 2 (course intake)			
ULD anti-S100 + anti-eNOS	134,70±20,24 **	9,50±1,38 **	117,90±15,65 **
ULD anti-S100	82,70±10,33	13,50±1,69	167,50±14,72 ×
ULD anti-eNOS	74,30±9,49 ×	17,30±2,40 ***	209,20±21,62 **
Placebo	63,70±3,91	15,00±1,47	199,60±31,19
P value on Kruskal-Wallis test ¹	0,0341	0,0244	0,0061

Notes:

¹ for determination of significant difference between groups the Kruskal-Wallis test was used. If the

test showed a significant difference of $p < 0.05$ for comparison between groups against each other

the Mann-Whitney test was used.

- * the significant difference in comparison with placebo, $p < 0.05$;
- ** the significant difference in comparison with placebo, $p < 0.01$;
- *** the significant difference in comparison with placebo, $p < 0.001$.
- * the significant difference in comparison with ULD anti-S100 + anti-eNOS, $p < 0.05$;
- ** the significant difference in comparison with ULD anti-S100 + anti-eNOS, $p < 0.01$;
- *** the significant difference in comparison with ULD anti-S100 + anti-eNOS, $p < 0.001$.

The analysis of HRV at the rest condition (in sitting position) before and after the CCEAC test (Table 9) detected that in subjects receiving study drugs had a tendency to an increased rate of SDNN indicating an increase in heart rate variability due to parasympathetic influence on heart rhythm. In response to a kinetic effect in all treatment groups the value of RMS-SD increased which characterizes the activity of the parasympathetic component of autonomic regulation. In the groups receiving the composition ULD anti-S100 + anti-eNOS and ULD anti-S100 showed an increase in HF which also indicated a shift in autonomic balance toward parasympathetic link. Thus, after conducting CCEAC tests in all groups there was an increase of parasympathetic effects on heart rate.

Table 9

The HRV parameters of the study participants at rest
before and after the kinetic action

Parameter	Visit 1 (one-day intake)		Visit 2 (course intake)	
	After the drug intake	After the CCEAC test	After the drug intake	After the CCEAC test
ULD anti-S100 + anti-eNOS group (M±SD)				
SDNN, msec.	57.7±5.51	68.2±7.42	59.4±5.03	65.6±4.66
RMSSD, msec.	43.1±6.77	51.4±9.22	47.0±6.21	47.6±5.33
TP, msec. ²	979.0±186.06	1678.3±397.1 1#	1067.2±167.24	1381.01166.30
LF, msec. ²	437.5±709.6	709.6±178.72	391.9±75.61	588.5±87.48
HF, msec. ²	171.5±51.08	228.4±76.79	206.5±58.32	218.5±43.96
LF/HF, c.u.	4.2±0.82	4.9±0.83	3.3±0.83	4.2±0.91
ULD anti-S100 group (M±SD)				
SDNN, msec.	60.9±4.62	70.9±5.90	59.1±4.80	68.8±4.87
RMSSD, msec.	44.3±5.39	50.6±6.56	42.4±4.63	47.8±5.57
TP, msec. ²	832.2±124.93 *	1342.8±217.0 9	841.4±149.93	1288.0±163.52 #
LF, msec. ²	315.2±52.38 *	550.9±72.44#	313.6±66.71	540.7±87.57#
HF, msec. ²	151.4±41.19	247.0±69.53#	138.3±38.42	187.1±39.80
LF/HF, c.u.	3.0±0.54	4.0±0.72	2.8±0.53	4.0±0.52
ULD anti-eNOS group (M±SD)				
SDNN, msec.	67.4±7.73	78.6±6.14	65.8±8.68	69.0±5.23

RMSD, msec.	53.0±8.86	58.4±7.68	59.6±12.45	52.2±5.30
TP, msec. ²	1307.8±324.24	1841.1±359.7 9#	1232.3±292.51	1275.4±172.47
LF, msec. ²	576.5±167.07	849.9±194.2#	527.2±167.07	562.1±89.38
HF, msec. ²	313.3±139.90	285.3±65.92	218.9±74.78	216.3±63.72
LF/HF, c.u.	3.6±0.87	3.9±0.82	3.7±1.14	3.8±0.58
Placebo group (M±SD)				
SDNN, msec.	64.6±6.10	75.7±6.42	61.1±6.72	70.8±6.79
RMSD, msec.	50.9±7.74	53.1±6.62	44.6±6.63	44.3±5.31
TP, msec. ²	1062.2±150.02	1917.8±318.9 6#	898.8±169.62	1418.5±227.59 #
LF, msec. ²	440.6±77.30	832.4±181.15	334.8±75.94	611.4±113.64#
HF, msec. ²	253.9±59.95	266.7±61.94	166.0±48.14	174.1±44.96
LF/HF, c.u.	3.4±0.72	5.0±1.33	3.4±0.93	4.8±0.83

Note: * the significant difference in comparison with the placebo, $p \leq 0.05$;

the significant difference in comparison with baseline parameters, $p < 0.05$.

The analysis of HRV in transition states showed that one-day intake of composition ULD anti-S100 + anti-eNOS increased the reaction time (13.9 ± 1.14 ; $p \leq 0.05$) and the stabilization time (24.2 ± 1.28 ; $p \leq 0.05$) in comparison with the ULD anti-S100 and placebo. The same factors exceeded the value of the placebo group and after the kinetic effect which demonstrated the positive effect of the combined drug on the reactivity of the ANS (increase of tolerance to changes in body position). The smallest

difference between the maximum and minimum heart rate in the breath test confirmed a better balance of the two divisions of ANS after receiving a one-day composition ULD anti-S100 + anti-eNOS (25.1 ± 2.66 beats / min, $p \leq 0.05$). By the end of week course of therapy the stabilizing effect on the balance of ANS after the CCEAC test (with orthostatic and breath test) is also noticed in the group receiving the composition ULD anti-S100 + anti-eNOS (Tables 10 and 11).

Table 10
The HRV parameters of participants of the study
at orthostatic test before and after kinetic action

Parameter	Visit 1 (one-day intake)		Visit 2 (course intake)	
	After drug intake	After CCEAC test	After drug intake	After CCEAC test
ULD anti-S100 + anti-eNOS (M±SD) Group				
Exercise reaction, c.u.	1.30±0.06	1.40±0.04	1.30±0.06	1.40±0.06
Reaction time, sec.	13.9±1.14*×	12.7±1.24*	11.8±0.57	11.7±1.09
Stabilization time, sec.	24.2±1.28*×	21.9±1.44*	20.6±0.74	22.4±1.44*×
ULD anti-S100 (M±SD) Group				
Exercise reaction, c.u.	1.40±0.04	1.30±0.04	1.30±0.04	1.30±0.05
Reaction time, sec.	7.60±1.05	10.6±1.55	9.7±1.21	10.0±1.73
Stabilization time, sec.	15.1±1.16*	18.3±1.43	18.0±1.18	18.0±1.80
ULD anti-eNOS (M±SD) Group				
Exercise reaction, c.u.	1.30±0.04	1.30±0.04	1.50 ± 0.12	1.30±0.04
Reaction time, sec.	8.20±0.94	9.10±1.12	9.2 ± 0.77	8.3±0.70
Stabilization time,	16.5±1.02	17.1±1.33	19.0 ± 2.04	16.7±0.98

sec.				
Placebo group (M±SD)				
Exercise reaction, c.u.	1.30±0.04	1.30±0.04	1.40 ± 0.06	1.30±0.06
Reaction time, sec.	9.5±1.28	8.1±0.90	10.4 ± 1.58	8.8±1 .09
Stabilization time, sec.	18.3±0.94	16.811 .09	18.0 ± 1.37	16.5±1 .11

Note: * the significant difference in comparison with placebo, $p<0.05$;

× the significant difference in comparison with ULD anti-S100, $p<0.05$.

Table 11

**The HRV parameters of participants of the study
at breath test before and after kinetic action**

Parameter	Visit 1 (one-day intake)		Visit 2 (course intake)	
	After drug intake	After CCEAC test	After drug intake	After CCEAC test
ULD anti-S100 + anti-eNOS (M±SD) Group				
Corellation max HR / min HR, c.u.	1.5 ± 0.05*	1.5 ± 0.06	1.5 ± 0.05	1.5 ± 0.05
Difference max HR – min HR, beats/min.	25.1 ± 2.66*	26.5 ± 2.77	26.5 ± 2.37	24.9 ± 2.24*
ULD anti-S100 (M±SD) Group				
Corellation max HR / min HR, c.u.	1.5±0.06	1.6±0.05	1.5±0.04	1.6±0.06
Difference				

max HR - min HR, beats/min.	27.7±2.68	27.2±2.40	25.7±2.24	26.9±2.67
ULD anti-eNOS (M±SD) Group				
Corellation				
max HR / min HR, c.u.	1.5±0.05	1.5±0.04	1.5±0.06	1.6±0.05
Difference				
max HR - min HR, beats/min.	26.7±2.44	26.2±2.04	27.7±2.47	27.3±2.12
Placebo group (M±SD)				
Corellation				
max HR / min HR, c.u.	1.6±0.07	1.6±0.06	1.5±0.05	1.6±0.05
Difference				
max HR - min HR, beats/min.	31.2±3.06	28.2±2.50	27.7±2.37	29.2±2.44

Note: * the significant difference in comparison with placebo, $p<0,05$

The results of self-esteem of functional state (well-being, activity, mood) of the subjects which was conducted by the participants of the study after the simulation of motion sickness (CCEAC tests) at the beginning and at the end of therapy showed that the subjects of all the groups have given 'average' points for each of the parameters (Table 12). Thus, on the background of drugs intake the CCEAC tolerance was satisfactory. The highest growth rates compared with data of the placebo group by the end of the 7th day of intake (more than 10%) was observed in the group of composition of ULD anti-S100 + anti-eNOS.

Table 12

The dynamics of parameters of self-esteem of functional

condition (well-being-activity-mood) of study participants

Parameter	Visit 1 (one-day intake)	Visit 2 (course intake)
ULD anti-S100 + anti-eNOS (M±SE) group		
Well-being	4.3±0.26	4.6±0.27
Activity	4.2±0.20	4.2±0.22
Mood	5.0±0.16	5.2±0.13
ULD anti-S100 (M±SE) group		
Well-being	3.7±0.21	4.3±0.22
Activity	3.6±0.17	4.0±0.19
Mood	4.5±0.16	4.9±0.19
ULD anti-eNOS (M±SE) Group		
Well-being	3.9±0.25	4.1±0.26
Activity	3.8±0.25	3.9±0.23
Mood	4.4±0.19	4.6±0.19
Placebo group (M±SE)		
Well-being	4.0±0.24	4.0±0.24
Activity	3.8±0.20	3.7±0.26
Mood	4.3±0.20	4.7±0.24

The safety analysis included data from all the subjects who participated in the study. During the observation period a well tolerance of studied preparations was noticed. No adverse events associated with drug administration identified. All the

subjects of studied groups completed treatment in the terms established by the study protocol; there was not persons early dropped out.

According to the results of physical examination including indicators of heart rate, systolic and diastolic blood pressure and according to the Harvard step test data the subjects were not recorded as with any abnormalities during the study (Table 13). All identified changes were not beyond the normal range. In this case, subjectively all subjects reported satisfactory well-being.

Table 13

The dynamics of physical parameters and exercise tolerance of study participants before and after kinetic action

Parameter	Visit 1 (one-day intake)		Visit 2 (course intake)	
	After drug intake	After CCEAC test	After drug intake	After CCEAC test
ULD anti-S100 + anti-eNOS (M±SE) Group				
HR (beats/min)	74.6±3.36	68.4±3.67	74.1±3.10	67.7±2.62
Systolic blood pressure (mmhg.)	123.4±2.83	125.9±4.08	121.8±2.65	128.3±4.25
Diastolic blood pressure (mmhg.)	74.0±3.09	79.3±2.62	76.2±2.43	80.3±3.30
Step-test index	—	53.6±2.60	—	52.3±2.09
ULD anti-S100 (M±SE) Group				
HR (beats/min)	73.5±2.57	69.7±2.78	72.1±2.84	67.7±2.39
Systolic blood pressure (mmhg.)	127.5±2.55	133.5±4.77	127.1±2.55	129.9±5.06
Diastolic blood pressure (mmhg.)	75.5±2.65	82.6±3.31	74.9±2.41	82.3±3.19

Step-test index	—	50.6±1 .71	—	53.0±1 .63
ULD anti-eNOS (M±SE) Group				
HR (beats/min)	76.5±2.59	67.3±1 .98	77.3±2.02	70.1±3.23
Systolic blood pressure (mmhg.)	127.3±3.14	131 .5±5.16	123.5±3.06	129.3±4.13
Diastolic blood pressure (mmhg.)	75.2±2.24	80.3±2.66	73.9±2.83	81.0±3.22
Step-test index	—	51.8±2.12	—	51.2±2.21
Placebo group (M±SE)				
HR (beats/min)	74.5±2.78	68.9±3.46	73.9±3.23	72.3±3.58
Systolic blood pressure (mmhg.)	125.3±3.30	133.3±4.73	124.3±2.83	126.9±3.95
Diastolic blood pressure (mmhg.)	76.2±2.15	81.7±2.83	75.4±1 .86	79.7±3.03
Step-test index	—	50.0±2.03	—	50.1±1 .99

In addition to the hemodynamic parameters, for evaluation of the safety of studied drugs and its possible negative impact on the central nervous functions, the following physiological parameters were examined in subjects: (RMO (reaction on moving object), SMRT (simple motor reaction time), RA (range of attention), attention span (AS), and attention stability factor (ASF)). In addition, the Stange's test was conducted to assess tolerance to hypoxia.

According to received results (Table 9) neither one-day or course drug intake had a significant effect on the estimated parameters. Indexes of sensory motor coordination (SMRT, RMO) did not differ from the results of the placebo group before and after the

CCEAC test at both visits. Study data of such complicated functions like volume and stability of attention showed that the studied drugs both before and after the CCEAC test did not change the degree of concentration and shift in attention not being different from the placebo group.

The analysis of standard exercise tests with breath holding showed a tendency to increase of the tolerance of hypoxia by the subjects (Table 14). When holding the breath the duration of Stange's test grew after taking all study drugs. However, only intake of the combination composition ULD anti-S100 + anti-eNOS showed significantly longer time in the holding of the breath after the kinetic effect (68.1 ± 18.8 sec. at baseline and 91.7 ± 27.4 sec. after the CCEAC test; $p < 0.05$). The increase of tolerance of hypoxia was also noted when the Gench's test (Stange's test) (breath holding at expiration, $P > 0.05$) was used.

Table 14

The dynamics of parameters of psycho-physiological state of study participants before and after kinetic action

Parameter	Visit 1 (one-day intake)		Visit 2 (course intake)	
	After drug intake	After CCEAC test	After drug intake	After CCEAC test
ULD anti-S100 + anti-eNOS (M\pmSE) Group				
SMRT	257.5 \pm 8.67	268.9 \pm 10.18	269.6 \pm 9.75	279.9 \pm 12.24
RMO, c.u.	50.1 \pm 3.92	49.5 \pm 4.50	47.3 \pm 4.86	47.0 \pm 3.54
RMO, % of target hit	3.0 \pm 0.95	4.5 \pm 1.15	5.3 \pm 1.58	4.0 \pm 1.11
AS, sec.	5.2 \pm 0.34	5.2 \pm 0.35	5.2 \pm 0.41	5.1 \pm 0.40
Range of attention, sec.	41.7 \pm 2.36	39.9 \pm 2.38	38.1 \pm 2.17	37.5 \pm 2.04
ASF	17.4 \pm 1.66	17.2 \pm 1.51	18.0 \pm 1.71	18.8 \pm 1.72
Stange's test	68.1 \pm 4.85	91.7 \pm 7.07*	71.8 \pm 6.02	85.5 \pm 9.36

Gench's test	47.1±4.03	50.1±3.94	46.7±3.28	48.114.52
ULD anti-S100 (M±SE) Group				
SMRT	258.9±9.95	282.4±13.56	268.4±1 1.37	279.119.20
RMO, c.u.	58.1 ±6.40	57.5±6.34	55.115.06	53.815.02
RMO, % of target hit	3.7±1.50	2.0±0.82	2.3±0.83	5.011.69
AS, sec.	6.0±0.40	6.4±0.52	6.2±0.42	6.010.41
Range of attention, sec.	42.6±2.68	42.1±2.27	42.7±2.30	41.912.52
ASF	14.5±1.16	14.9±1.26	15.3±1.13	15.411.18
Stange's test	59.0±4.09	72.6±6.19	64.5±4.93	75.915.67
Gench's test	47.114.48	49.4±4.69	48.3±4.30	48.814.14
ULD anti-eNOS (M±SE) group				
SMRT	257.7±8.49	279.4±14.23	266.7±13.19	275.5111 .44
RMO, c.u.	48.3±3.67	51.9±4.39	52.5±4.79	49.614.22
RMO, % of target hit	2.3±0.83	2.0±0.82	3.3±1.26	5.711.68
AS, sec.	5.9±0.25	6.0±0.34	5.5+0.24	5.910.33
Range of attention, sec.	41.9±2.10	43.8±2.39	41.3±2.00	42.512.22
ASF	13.7±1.34	14.8±1.31	15.6±1.24	14.111.40
Stange's test	62.5±5.49	69.5±5.09	56.7±3.34	73.117.98
Gench's test	43.113.51	45.7±3.15	43.413.77	45.814.03
Placebo group (M±SE)				
SMRT	267.6±7.64	290.1±1 1.33	281.119.78	263 .316.85
RMO, c.u.	60.7±8.31	54.1±5.57	51.113.69	52.615.38
RMO, % of target hit	3.7±1.03	3.7±1.24	3.310.93	4.311.61
AS, sec.	6.1±0.71	5.7±0.36	5.510.32	5.910.71

Range of attention, sec.	41.9±2.09	42.4±2.81	41.3±2.18	39.612.26
ASF	14.5±1.64	14.511.79	15.3±1.55	15.911.58
Stange's test	63.7±4.71	67.916.90	64.815.94	83.011.2.24
Gench's test	44.7±2.52	47.113.30	43.712.71	47.813.78

Thus, the study using an experimental motion sickness demonstrated the effectiveness of the combination composition ULD anti-S100 + anti-eNOS and monocomponent preparation ULD-S100. The studied drugs increase the stability of the subjects to the kinetic effect after simulation of the clinical and physiological effects of motion sickness contributing to more mild clinical process of motion sickness and earlier recovery of the subjects after cessation of treatment. In addition, it was shown that the anti motion sickness effect of the combination composition (compositions ULD anti-S100 + anti-eNOS) increases the efficiency of individual components. The effectiveness of the combination composition ULD anti-S100 + anti-eNOS in the control of the vestibular-autonomic and sensory reactions of a body in experimental motion sickness increases at course intake. It should be noted that ULD anti-eNOS in the form of monopreparation does not have a protective effect against motion sickness but when combined with ULD anti-S100 significantly enhances the anti motion sickness effect of the last one which manifests itself as at one-day so at short course intake of the drug. The best ability to adjust the transient processes that is to influence to the reactivity of the parasympathetic and sympathetic parts of ANS as well as adaptive capabilities of ANS in a state of motion sickness (to increase the tolerance to sudden changes in a body position) was observed in the composition ULD anti-S100 + anti-eNOS which is an important component of anti motion sickness properties of the drug. Composition ULD anti-S100 + anti-eNOS and monocomponent preparation ULD anti-S100 when using them as anti motion sickness preparation including when performing an operator functions are safe and do not adversely impact on the physical and psycho- physiological parameters.

Combination composition ULD anti-S100 + anti-eNOS and ULD anti-S100 can be recommended for the prophylaxis and relief of kinesia in motion disease (including sea, air and car sicknesses) to persons with low and moderate degree of stability. The combination composition has high safety and no adverse effects on the quality of professional activity.

Example 9.

To study the properties of the combination pharmaceutical composition of the present application for the treatment of psychoorganic syndrome, tablets with weight of 300 mg were used. The tablets were impregnated with pharmaceutical composition containing water-alcohol solutions (6 mg/tab.) of activated-potentiated forms of polyclonal affinity purified rabbit brain-specific proteins antibodies S-100 (anti-S100) and to endothelial NO-synthase (anti-eNOS) in ultra low doses (ULD) obtained by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200 ("ULD anti-Si 00 + anti-eNOS").

The control group patients received 300 mg tablets impregnated with pharmaceutical composition containing water-alcohol solutions (6 mg/tablet) of activated-potentiated forms of polyclonal affinity purified rabbit brain-specific proteins antibodies S-100 (anti-S100) in ultra low doses (ULD) obtained by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times.

The study included patients diagnosed with psychoorganic syndrome of posttraumatic origin. Psychoorganic syndrome is characterized by the following triad of signs: weakness of memory, loop of intelligence, incontinence of affect (Walther Buell triad).

The study was an open-label randomized comparative parallel group clinical trial of efficacy and safety of the therapy in patients with psychoorganic syndrome of posttraumatic origin (the first group of patients took the preparation of ULD anti-S100, the second group of patients - the preparation of ULD anti-S100+anti-eNOS).

The study included 6 patients aged 35 to 90 years old (mean age 70.83 ± 21.95) diagnosed with psychoorganic syndrome.

Compliance of patients to following inclusion and exclusion criteria was checked:

Inclusion criteria:

1. Patients diagnosed with posttraumatic encephalopathy with psychoorganic syndrome or with encephalopathy of complex etiology (vascular, posttraumatic) with psychoorganic syndrome, confirmed by medical history, neurological examinations and medical records.
2. Patient without change in concomitant therapy within at least one month prior to Visit 1.
3. No need for change in concomitant therapy for the whole observation period.
4. No need for immunomodulatory drugs prescription for the next 6 months.
5. Patients with a level of education sufficient to adequately communicate with the researcher and study coordinator.
6. Patients assessed by the researcher as reliable and ready to perform all scheduled clinical visits, tests and procedures stipulated in the protocol.
7. Patients having a valid home address.

Exclusion criteria:

1. Any brain surgery in medical history.
2. Acute myocardial infarction.
3. Hemorrhagic stroke.
4. The diagnosis of psychosis, bipolar disorder or schizoaffective disorder in medical history.
5. Major depressive disorder according to criteria of depression module of international neuropsychiatric mini-interview (MINI).
6. Factors/conditions of medical or another character which in the opinion of the researcher may affect to the test results for patients in the study.

7. Answers "2A", "2B", "2C" or "3" in the section 1" of Beck Depression questionnaire (active suicidal ideation with some intent to act, without a specific plan, or active suicidal ideation with a specific plan and intent).
8. Autoimmune disease in medical history.
9. Acute damage of liver or severe cirrhosis (class C by Child-Pugh).
10. Non-corrected disorder of thyroid gland function.
11. Decompensated arterial hypertension in medical history.
12. Serious or decompensated cardiovascular disease, liver disease, kidney disease, metabolic, respiratory or hematological disease, symptomatic peripheral vascular disease or another medical or psychiatric condition which in the opinion of the researcher, may affect the patient's participation in the study or could lead to prolonged hospitalization or re-hospitalization during the study.
13. Diseases and conditions which in the opinion of researcher may prevent patient from the participation in the study.
14. The intake of the drug containing ULD anti-eNOS or the drug containing ULD anti-Si 00 before inclusion in the study.
15. The intake of antidepressants of any group including plant and homeopathic preparations.
16. The intake of anxiolytics of any group including plant and homeopathic preparations.
17. The intake of immunomodulators including plant and homeopathic preparations.
18. The treatment with systemic steroids within 1 month before Visit 0.
19. The participation in the study of the drug containing ULD anti-eNOS or the drug containing ULD anti-S100 if patients took at least one dose of preparation.
20. Participation in other clinical studies within 1 month before within 1 month before being enrolled in this study.
21. Pregnancy, breast feeding, impossibility to use an adequate contraception during the study period and within 1 month after the last intake of the studied drug.
22. The presence of allergy/intolerance of any component of drugs including lactose intolerance.

23. Patients taking narcotic drugs and neuroleptics, alcoholic dependence, psychiatric diseases in patients.
24. Patients are the staff of the center which directly related to the conducted study and/or are family members of the research center staff's which directly associated with the ongoing study. The "family members" are a husband (wife), parents, children, brothers (sisters).
25. Participation in the trial or presumable receiving of compensation or participation in the judicial process in the opinion of a researcher.

After the determination of patient conformity to inclusion and exclusion criteria the patients were randomized into two study groups: a group of patients receiving ULD anti-S100 (3 patients, women - 33.33%, men - 66.66%, mean age - 71.33 ± 16.25 years old), a group of patients receiving ULD anti-S100 + anti-eNOS (3 patients, women - 66.66 % men - 33.33 %, mean age - 70.33 ± 30.66 years old).

During this study the five visits were carried out. Treatment phase lasted from Visit 1 to Visit 4 for 84 ± 5 days on average. Visit 4 (Day 84 ± 5) was the first endpoint of the study followed by a follow-up observation. Follow-up phase continued from Visit 4 to Visit 5 (Day 168 ± 5 on average).

In the safety analysis the data of all patients participating in the study ($n = 6$) was included. During the study good tolerance of the drug was recorded. No adverse events were registered. All patients of studied groups have completed the treatment according to the protocol; no early dropouts.

The effect of ULD anti-S100 + anti-eNOS preparation on the main clinical signs and symptoms of psychoorganic syndrome (NPI neuropsychiatric inventory, Intensity section), on the intensity of concomitant distress of the person attending to the patient (NPI Neuropsychiatric Inventory, Distress section) as well as the on patient's cognitive functions (The Mini Mental State Examination, MMSE) were assessed. An improvement was found in the key symptoms of psychoorganic syndrome such as statistically

significant reduction of the intensity section of NPI neuropsychiatnc inventory (from 91.0 ± 15.13 to 69.0 ± 6.24 , $p < 0.05$), decrease of distress section score of NPI neuropsychiatnc inventory (from 44.33 ± 17.78 to 36.33 ± 3.21 , $p < 0.05$) at Visit 4 (Table 15).

In the group of patients receiving ULD anti-S100 alone no clinical improvement was recorded.

At that, a difference between the groups of patients in the total score of the Intensity section of NPI neuropsychiatnc inventory at the end of therapy was statistically significant at $p < 0.05$.

Table 15.

	NPI (intensity)	NPI (distress)	ADS-ADL	MMSE
ULD anti-S100+anti-eNOS before treatment	91.0 ± 15.13	44.33 ± 17.78	42.66 ± 4.93	22.33 ± 3.21
ULD anti-S100+anti-eNOS after treatment	69.0 ± 6.24 *#	36.33 ± 3.21 *	52.0 ± 5.57	22.66 ± 2.08
ULD anti-S100 before treatment	114.0 ± 25.53	45.66 ± 14.47	33.0 ± 13.89	22.33 ± 4.16
ULD anti-S100 after treatment	99.66 ± 18.0	49.0 ± 17.05	31.66 ± 10.69	23.0 ± 4.36

* - p from baseline < 0.05 ; # - p from control < 0.05

Thus, in the conducted clinical study a positive effect of combined pharmaceutical composition ULD anti-S100 + anti-eNOS on the main clinical signs and symptoms of

psychoorganic syndrome and tendency to effect cognitive functions with psychoorganic syndrome. In addition, good drug tolerability was confirmed. No drug-related adverse events were registered.

Example 10.

To study the properties of the combination pharmaceutical composition of the present application for the treatment of Alzheimer's disease, tablets with weight of 300 mg were used. The tablets were impregnated with pharmaceutical composition containing water-alcohol solutions (6 mg/tablet.) of activated-potentiated forms of polyclonal affinity purified rabbit brain-specific proteins antibodies S-100 (anti-S100) and to endothelial NO-synthase (anti-eNOS) in ultra low doses (ULD) obtained by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200 (ratio: 1:1) ("ULD anti-S100 + anti-eNOS").

The control group patients received 300 mg tablets impregnated with pharmaceutical composition containing water-alcohol solutions (3 mg/tablet) of activated-potentiated forms of polyclonal affinity purified rabbit brain-specific proteins antibodies S-100 (anti-S100) in ultra low doses (ULD) obtained by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} times, of equivalent mixture of centesimal homeopathic dilutions C12, C30, C200

The study included patients diagnosed with Alzheimer's disease. Alzheimer's disease is characterized by dementia (acquired dementia, stable impairment of cognitive activity with certain loss of previously acquired knowledge and practical skills, difficulties or impossibility to gain new knowledge).

The study was an open-label randomized comparative clinical trial of efficiency and safety of the therapy in two parallel groups (preparations of ULD anti-S100 and ULD anti-S100+anti-eNOS) in the treatment of patients with mild to moderate Alzheimer's disease.

The study included 6 patients aged 55 - 64 years old (mean age 59.0 ± 3.58) diagnosed with mild to moderate Alzheimer's disease.

Compliance of patients to following inclusion and exclusion criteria was checked:

Inclusion criteria are as follows:

1. Patients with mild to moderate Alzheimer's disease, confirmed by medical history, neurological examinations and medical records.
2. Patient without change in concomitant therapy within at least one month prior to Visit 1.
3. No need for change in concomitant therapy for the whole observation period.
4. No need for immunomodulatory drugs prescription for the next 6 months.
5. Patients with a level of education sufficient to adequately communicate with the researcher and study coordinator.
6. Patients assessed by the researcher as reliable and ready to perform all scheduled clinical visits, tests and procedures stipulated in the protocol.
7. Patients having a valid home address.

Exclusion criteria are as follows:

1. Any brain surgery in medical history.
2. Acute myocardial infarction.
3. Hemorrhagic stroke.
4. The diagnosis of psychosis, bipolar disorder or schizoaffective disorder in medical history.
5. Major depressive disorder according to criteria of depression module of international neuropsychiatric mini-interview (MINI).
6. Factors/conditions of medical or another character which in the opinion of the researcher may affect to the test results for patients in the study.

7. Answers "2A", "2B", "2C" or "3" in the section "I" of Beck Depression questionnaire (active suicidal ideation with some intent to act, without a specific plan, or active suicidal ideation with a specific plan and intent).
8. Autoimmune disease in medical history.
9. Acute damage of liver or severe cirrhosis (class C by Child-Pugh).
10. Non-corrected disorder of thyroid gland function.
11. Decompensated arterial hypertension in medical history.
12. Serious or decompensated cardiovascular disease, liver disease, kidney disease, metabolic, respiratory or hematological disease, symptomatic peripheral vascular disease or another medical or psychiatric condition which in the opinion of the researcher, may affect the patient's participation in the study or could lead to prolonged hospitalization or re-hospitalization during the study.
13. Diseases and conditions which in the opinion of researcher may prevent patient from the participation in the study.
14. The intake of the drug containing ULD anti-eNOS or the drug containing ULD anti-Si 00 before inclusion in the study.
15. The intake of antidepressants of any group including plant and homeopathic preparations.
16. The intake of anxiolytics of any group including plant and homeopathic preparations.
17. The intake of immunomodulators including plant and homeopathic preparations.
18. The treatment with systemic steroids within 1 month before Visit 0.
19. The participation in the study of the drug containing ULD anti-eNOS or the drug containing ULD anti-S100 if patients took at least one dose of preparation.
20. Participation in other clinical studies within 1 month before being enrolled in this study.
21. Pregnancy, breast feeding, impossibility to use an adequate contraception during the study period and within 1 month after the last intake of the studied drug.
22. The presence of allergy/intolerance of any component of drugs including lactose intolerance.

23. Patients taking narcotic drugs and neuroleptics, alcoholic dependence, psychiatric diseases in patients.
24. Patients are the staff of the center which directly related to the conducted study and/or are family members of the research center staff's which directly associated with the ongoing study. The "family members" are a husband (wife), parents, children, brothers (sisters).
25. Participation in the trial or presumed receiving of compensation or participation in the judicial process in the opinion of a researcher.

After the determination of patient conformity to inclusion and exclusion criteria the patients were randomized into two study groups: a group of patients receiving ULD anti-Si 00 (3 patients, women - 100%, men - 0%, mean age - 59.0 ± 3.6 years old), a group of patients receiving ULD anti-S100 + anti-eNOS (3 patients, women - 66.66 % men - 33.33 %, mean age - 59.0 ± 4.36 years old).

During this study the five visits were carried out. Treatment phase lasted from Visit 1 to Visit 4 for 84 ± 5 days on average. Visit 4 (Day 84 ± 5) was the first endpoint of the study followed by a follow-up observation. Follow-up phase continued from Visit 4 to Visit 5 (Day 168 ± 5 on average).

In the safety analysis the data of all patients participating in the study ($n = 6$) was included. During the study good tolerance of the drug was recorded. No adverse events were registered. All patients of studied groups have completed the treatment according to the protocol; no early dropouts.

The effect of ULD anti-S100 + anti-eNOS preparation on the main clinical signs and symptoms of Alzheimer's disease (NPI neuropsychiatric inventory, Intensity section), on the intensity of concomitant distress of the person attending to the patient (NPI Neuropsychiatric Inventory, Distress section) as well as the on patient's cognitive functions (The Mini Mental State Examination, MMSE) were assessed. An improvement was found in the key symptoms of Alzheimer's disease such as statistically significant

reduction of the intensity section of NPI neuropsychiatric inventory (from 24.33 ± 4.73 to 12.0 ± 3.46 , $p < 0.05$) at Visit 4 (Table 16).

A tendency for reduction of distress of the person attending to the patient was also found as well as for the reduction in activity of the patient's everyday life at the end of therapy (however, without any statistically significant difference, possibly due to the small number of patients included in the study).

Besides, a tendency for improvement of cognitive functions was found, manifested by increase of MMSE score from 23.66 ± 3.21 to 26.66 ± 1.53 points, however, the difference also failed to reach statistically significant values at the end of therapy, which may also be related to the small sample size.

The same endpoints in the group of patients receiving ULD anti-S100, showed no trend for improvement, except a statistically insignificant improvement of MMSE score from 22.66 ± 0.58 to 23.33 ± 0.58 points.

At that, a difference between the groups of patients in the total MMSE score at the end of therapy was statistically significant at $p < 0.05$.

Table 16.

	NPI (intensity)	NPI (distress)	ADCS-ADL	MMSE
ULD anti-Si 00+anti-eNOS before treatment	24.33 ± 4.73	9.66 ± 1.53	71.0 ± 6.56	23.66 ± 3.21
ULD anti-Si 00+anti-eNOS after treatment	$12.0 \pm 3.46^*$	5.0 ± 3.61	74.33 ± 2.51	$26.66 \pm 1.53\#$
ULD anti-S100 before treatment	35.66 ± 15.50	22.33 ± 5.50	61.66 ± 5.13	22.66 ± 0.58
ULD anti-S100 after treatment	38.33 ± 8.5	23.0 ± 5.0	61.33 ± 5.86	23.33 ± 0.58

* - p from baseline < 0.05 ; # - p from control < 0.05

Thus, in the conducted clinical study a positive effect of combined pharmaceutical composition ULD anti-S100 + anti-eNOS on the main clinical signs and symptoms of Alzheimer's disease and tendency to effect cognitive functions with Alzheimer's disease. In addition, good drug tolerability was confirmed. No drug-related adverse events were registered.

Example 11.

Group 1 - the active drug group was given 300 mg tablets impregnated with an aqueous-alcohol solutions (6 mg/tab) of activated-potentiated form of polyclonal rabbit antibodies to brain specific S-100 protein (anti-S-100), and to endothelial NO-synthase (anti-eNOS) in ultra low dose (ULD anti-S-100 + ULD anti-eNOS), purified on antigen, obtained by super dilution of initial solution (with concentration of 2.5 mg/ml) in 100^{12} , 100^{30} , 100^{200} time, equivalent to mixture of centesimal homeopathic dilutions C12, C30, C200;

Group 2 - the comparison group was given 300 mg tablets impregnated with an aqueous-alcohol solution (3 mg/tab) of activated-potentiated forms of polyclonal rabbit antibodies to brain-specific S-100 protein purified on antigen in ultra low dose (ULD anti-S100) obtained by super dilution of initial solution in 100^{12} , 100^{30} , 100^{50} times, of equivalent mixture homeopathic dilutions C12, C30, C50.

Group 3 - the control group (placebo) was given of 300 mg tablets having excipients (lactose monohydrate - 267 mg, microcrystal cellulose - 30 mg, magnesium stearate - 3 mg).

The effectiveness of the active drug ULD anti-S100 + anti-eNOS in the treatment of patients with syndrome of attention deficit and hyperactivity disorder (ADHD) was conducted in comparative double blind placebo-controlled study in 146 children from 6 to 12 years old (mean age 9.3 ± 0.24 years old) who were randomized into three groups depending on prescribed therapy. Within 12 weeks the patients of group No.1 (n = 46)

received the composition ULD anti-S100 + anti-eNOS, 2 tablets twice a day; the comparison group 2 members (n = 50) received ULD anti-S100, 2 tablets twice a day; the control group 3 members (n = 50) received 2 tablets twice a day. All the patients included in the study had clinically marked presentations of ADHD which was confirmed by high points on ADHD symptoms assessing scale (ADHDRS-IV-Home Version): 33.8 ± 0.92 in group 1; 32.5 ± 1.14 in group 2 and 33.6 ± 0.91 in group 3. Most of the children were characterized by a moderate degree of severity of ADHD according to the CGI-ADHD-Severity questionnaire. The total score on this scale was 4.0 ± 0.02 points in the group 1, 4.0 ± 0.03 points in the group 2, and 4.0 ± 0.00 points in the group 3. Thus, initially the patients of the three groups had comparable indicators of the severity of ADHD. According to the results of neurological, clinical - laboratory and instrumental examination at the time of enrollment to the study no abnormalities in any patient was detected. Over the 12 weeks of treatment, patients were seen six times by a doctor. During which the physician-researcher recorded the dynamics of intensity of clinical presentations of ADHD (total score on a scale ADHDRS-IV-Home Version) and disease severity (on the CGI-ADHD-Severity), supervised the prescriptions and administration of treatment and evaluated the safety of the treatment.

The analysis of the effectiveness of 12 weeks of therapy in the three groups showed a decrease of more than 25% from the initial total score on a scale ADHDRS-IV-Home Version in 75% (n = 36) of children treated with the composition ULD anti-S100 + anti-eNOS; in 66% (n = 33) of patients treated with ULD anti-S100 and in 56% (n = 28) of children receiving placebo. Differences of efficiency between the groups showing a more detailed assessment, taking into account the three-level grading of improvement of condition (reduction of total score on a scale ADHDRS-IV for <25%, 25-49.9% or $\geq 50\%$ from the baseline), are presented in Table 17. Significant improvement with a reduction in total score on 50% or more from the baseline was noted in 52% of children in group 9 who were taking ULD anti-S100 + anti-eNOS, and in 34% of children in group 2 who were taking ULD anti-S100 (vs. 8% of patients in group 3 with placebo).

Significant reduction ($p <0.001$) of clinical implications of ADHD in comparison with the initial state is already occurred after 2 weeks of therapy in all three groups of

observation. Positive dynamics was more significant in patients of groups 9 and 2 as the significant differences were identified in them between total scores ADHRS-IV-Home Version, not only in relation to the screening visit but when compared with the indexes of the group 3 with placebo. In subsequent weeks of treatment the efficacy of treatment with composition ULD anti-S100 + anti-eNOS and monocomponent preparation ULD-S100 started to grow, the most significantly in the active drug group ($p < 0.05$). The resulting decrease in total score on a scale ADHRS-IV-Home Version in children of the group 9 with ULD anti-S100 + anti-eNOS was 16.5 points, in patients of the group 2 with ULD anti-S100 - 12.4 points (compared to 6.3 points in the group 3 with placebo). As a result of 12-week of treatment the intensity of clinical implications of ADHD in children treated with the composition ULD anti-S100 + anti-eNOS decreased by almost in half (-48.8%) and in patients treated with ULD anti-S100 more than in one-third (-38.2%) compared with the baseline.

The intake of composition ULD anti-S100 + anti-eNOS or ULD anti-S100 influenced on both clusters of symptoms of ADHD which was confirmed by dynamics of assessments by two sections of the scale with ADHRS-IV-Home Version. Moreover, the treatment with the composition ULD anti-S100 + anti-eNOS was significantly higher than the effectiveness of therapy with monopreparation ULD anti-S100 in the degree of influence on the intensity of implications and attention deficit and hyperactivity/impulsivity.

The positive therapeutic effect of the active drug ULD anti-S100 + anti-eNOS and drug of comparison ULD-S100 was shown in evaluating of patients' treatment results on a scale of ADHD severity assessment (CGI-ADHR-Severity) (Table 17). Almost the fourth part of the patients in ULD anti-S100 + anti-eNOS group the severity of disease was decreased from moderate to mild and even to minimal as confirmed by a decrease in mean value on a scale CGI-ADHR-Severity on 15% after 3 months of therapy (from 4.0 ± 0.02 to 3.4 ± 0.06 ; $p < 0.001$). The effect of therapy with monopreparation ULD anti-S100 was slightly lower and indicated -10% on a scale CGI-ADHR-Severity over 3 months (vs. 5% in the placebo group). The safety analysis included data of all the patients participating in the study. During the whole period of monitoring there was both,

well comparable to placebo, the tolerance of active drug ULD anti-S100 + anti-eNOS and preparation of comparison ULD-S100. Adverse events were reported in one patient of the group with ULD anti-S100 (subside during the fourth week of the study headaches) and in one patient of the placebo group (sleepwalking during the second month of observation). These adverse events were not connected with the therapy. In addition, during the treatment the single cases of acute respiratory disease were observed which also are not associated with the therapy. All the patients of studied groups completed the treatment to schedule established by the study protocol; no early dropouts. The absence of pathological changes according to physical examination of the patients and in the course of repeated analysis of laboratory parameters confirmed the safety of studied therapy.

According to the results of physical examination (heart rate, SBP, DBP, body temperature) in patients any pathological alterations during treatment were not registered. Differences in analyzing rates according to visits and in the compared groups did not reach the statistical significance and do not exceed the limits of physiologically-allowable deviations. High rates of adherence to therapy additionally evidenced as about effectiveness so as about the safety of studied preparations. By the end of the third month of treatment the adherence was $99.8 \pm 1.15\%$ and $98.8 \pm 2.25\%$ in the group 9 with ULD anti-S100 + anti-eNOS and in the group 2 with ULD anti-S100 respectively (versus $74.6 \pm 2.54\%$ in the group 3 with placebo).

Thus, the study demonstrated the efficacy and safety of the compositions ULD anti-S100 + anti-eNOS and of monocomponent preparation ULD-S100 in the treatment of children with ADHD. The most pronounced therapeutic effect in the 12-week course was observed in complex drug (ULD anti-S100 + anti-eNOS) which was manifested by positive dynamics of clinical symptoms in the majority (75%) of children. The composition ULD anti-S100 + anti-eNOS had correcting influence to both of the clusters of symptoms of ADHD and as a result, the significant reduction of attention disorders and hyperactivity in patients with ADHD was noted.

Table 17. The dynamics of total score by the scale ADHRS-IV-Home

Version by the end of 12 weeks of therapy

Groups of patients	The proportion of patients with decrease of total score by the scale ADHRS-IV-Home Version		
	Less than 25.0% from baseline	on 25.0 – 49.9% from baseline	on 50.0% and more from baseline
ULD anti-S100 + anti-eNOS, n=48	12 (25%)	11 (23%)	25 (52%) ##
ULD anti - S100, n=50	17 (34%)	16 (32%)	17 (34%) ##
Placebo, n=50	22 (44%)	24 (48%)	4 (8%)

The difference is significant in comparison with the placebo group:

** p<0.01.

Table 18. The dynamics of evidence of clinical implications of ADHD by the scale ADHRS-IV-Home Version

Treatment stage	ULD anti-S100 + anti-eNOS, n=48		ULD anti-S100, n=50		Placebo, n=50	
	Value (M±SE)	Δ from baseline	Value (M±SE)	Δ from baseline	Value (M±SE)	Δ from baseline
Total score						
Screening	33.8 ±0.96		32.5 ± 1.14		33.6 ± 0.91	
2 weeks	24.1 ±0.97 *** #	-28.7%	25.1 ± 1.03	-22.8 %	28.8 ± 1.26	-14.3 %

			*** #		***	
4 weeks	22.6 ±0.98 *** ##	-33.1%	22.7 ± 1.23 *** ##	-30.2 %	29.9 ± 1.06 ***	-11.0 %
6 weeks	19.4 ±0.95 " * ##	-42.6%	20.8 ± 1.06 *** ##	-36.0 %	29.0 ± 1.25 ***	-13.7 %
8 weeks	18.9 ±0.94 *** um	-44.1 %	20.9 ± 1.30 *** ##	-35.7 %	27.6 ± 1.35 ***	-17.9 %
12 weeks	17.3 ±0.96 *** ## &	-48.8%	20.1 ± 1.21 *** ##	-38.2 %	27.3 ± 1.48 ***	-18.8 %

Attention disorders

Screening	18.4 ±0.55		17.4 ± 0.57		18.4 ± 0.43	
2 weeks	12.8 ±0.57 *** #	-30.4%	13.7 ± 0.68 *** #	-21.3 %	16.1 ± 0.66 ***	-12.5 %
4 weeks	11.6 ±0.56 *** ##	-37.0%	12.9 ± 0.79 *** ##	-25.9 %	16.4 ± 0.57 ***	-10.9 %
6 weeks	10.7 ±0.54 *** ##	-41.8%	11.9 ± 0.64 *** ###	-31.6 %	16.0 ± 0.70 ***	-13.0 %
8 weeks	10.3 ±0.53 *** ##	-44.0%	11.5 ± 0.70	-33.9 %	15.1 ± 0.76	-17.9 %

			*** ###		***	
12 weeks	9.7 ±0.55 *** ##&	-47.3%	11.4 ± 0.68 *** ##	-34.5 %	14.9 ± 0.78 ***	-19.0 %
Hyperactivity /impulsion						
Screening	15.4 ±0.61		15.1 ± 0.77		15.2 ± 0.62	
2 weeks	11.3 ±0.63 ***	-26.6%	11.4 ± 0.61 ***	-24.5 %	12.7 ± 0.74 ***	-16.4 %
4 weeks	11.0 ±0.62 *** ##	-28.6%	9.8 ± 0.64 *** ##	-35.1 %	13.5 ± 0.67 **	-11.2 %
6 weeks	8.7 ±0.59 *** ##	-43.5%	8.9 ± 0.64 *** ##	-41.1 %	12.9 ± 0.73 **	-15.1 %
8 weeks	8.6 ±0.60 *** ##	-44.2%	9.5 ± 0.76 *** ##	-37.1 %	12.5 ± 0.81 ***	-17.8 %
12 weeks	7.6 ±0.57 *** ## &	-50.6%	8.7 ± 0.70 *** ##	-42.4 %	12.5 ± 0.82 ***	-17.8 %

Note. The difference is significant in comparison with baseline parameter:

* p<0.05, ** p<0.01, *** p<0.001 .

The difference is significant in comparison with placebo group:

p<0.05, ## p<0.01, ### p<0.001 .

The difference is significant in comparison with the group of

ULD anti-S100: & p<0.05.

Table 19. The dynamics of severity level of ADHD by the scale

CGI-ADHD-Severity

Parameter		
	ADHD Severity	
	M±SE	Δ from baseline
ULD anti-S100 + anti-eNOS, n=48		
Screening	4.0±0.02	
4 Weeks	3.6±0.02**	-10%
12 Weeks	3.4±0.06***	-15%
ULD anti-S100, n=50		
Screening	4.0±0.03	
4 Weeks	3.8±0.06**	-5%
12 Weeks	3.6±0.08***	-10%
Placebo, n=50		
Screening	4.0±0.01	
4 Weeks	3.9±0.05	-2.5%
12 Weeks	3.8±0.06***	-2.5%

The difference is significant in comparison with the baseline parameter: ** p<0.01 , *** p<0.001 .

Example 12.

Double blind, placebo-controlled clinical study of a combination of activated potentiated forms of antibodies to the C-terminal fragment of the angiotensin II AT1-receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic

dilutions of C12, C30, C200, in human patients with chronic heart failure to evaluate key parameters of the CHF pathology.

80 patients (CHF of II-IV functional class (FC), left ventricular ejection fraction (LVEF) less than 40%) were divided in 4 equal treatment and control groups for a 6 months study. The background therapy was not discontinued (bisoprolol β -blocker, ACE inhibitor enalapril, aspirin (unless contraindicated); administration of diuretics, nitrates, digoxin was also admitted). Group 1 received the activated potentiated form of antibodies to a C-terminal fragment of the angiotensin II AT1-receptor (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 2 received the activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 3 received the combination pharmaceutical composition comprising both activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1-receptor (mixture of homeopathic dilutions C12, C30, C200) and activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, \ C200) (3 tablets/day, n=20). Group 4 received placebo (3 tablets/day, n=20). The groups were comparable in the initial study parameters: in age and sex, and severity (class of CHF and LVEF) and duration of the disease.

Before and after treatment, the patients were evaluated for the effect of the administered medications on vascular remodeling and endothelium dysfunction that is important for the CHF process and progression. The effects of the medications on the processes of vascular remodeling were evaluated by pulse wave velocity (PWV) ("Colson" system) in the carotid-femoral (CF) (elastic type) and carotid-radial (CR) (muscle type) segments of arteries.

Table 20 shows the dynamics in the rates of pulse wave velocity in the carotid-femoral (CF) (elastic type) and carotid-radial (CR) (muscle type) segments of arteries.

Table 20

Groups/ Parameters	ULDs ¹ of Abs ² to C-terminal fragment of AT1 receptor of	ULD of Abs to endothelial NO- synthase	Combination of ULDs of Abs to C-end fragment of AT1 receptor	Placebo

	angiotensin II						of angiotensine II and ULD of Abs to endothelial NO-synthase					
	^	&	Δ %	^	&	Δ %	^	&	Δ %	^	&	Δ %
CF, m/c	9, 7± 0. 5	8± 0.6	-14.8*	10 .1 ±0 .5	9.8± 0.4	- 2.9	10. 8± 7	8.6 ±0. 6	- 20.3*	8. 2± 0. 4	8.2±0. 5	0.1
CR, m/c	8. 6± 0. 2	8.9 ±0. 3	2.9	8. 8± 0. 1	8.3± 0.3	- 5.7	8.9 ±0. 5	7.6 ±0. 7	- 15.6* #\$	9. 1± 0. 3	9.7±0. 3	6.4*

(^) denotes initial value

(&) denotes 6 month after beginning of administration

(*) denotes difference from initial value is verifiable with p value < 0.05.

(#) denotes difference from the group receiving ULDs of Abs to C-terminal fragment AT1 receptor angiotensin II with verifiable difference in p value of < 0.05.

(\\$) denotes difference from the group receiving ULDs of Abs to endothelial NO-synthase with verifiable difference in p value of < 0.05.

(1) ULD denotes ultra-low doses.

(2) Abs denotes antibodies.

After 6 months of treatment, only group 3 showed a proven effect of the claimed pharmaceutical composition on the stiffness of muscular type arteries. Group 1 which received ULD of antibodies to a C-terminal fragment of angiotensin II AT1-receptor, and group 3 which received the combination pharmaceutical composition of the invention showed a proven increase in the stiffness of elastic type arteries.

Example 13.

Double blind, placebo-controlled clinical study of a combination of activated potentiated forms of antibodies to the C-terminal fragment of angiotensin II AT1-receptor, in a mixture of homeopathic dilutions of C12, C30, C200, with activated potentiated form of antibodies to endothelial NO-synthase, in a mixture of homeopathic dilutions of C12, C30, C200, in human patients with chronic heart failure to evaluate key measurement of quality of life.

80 patients (CHF of II-IV functional class (FC), left ventricular ejection fraction (LVEF) less than 40%) were divided in 4 equal treatment and control groups for a 6 months study. The background therapy was not discontinued (bisoprolol β -blocker, ACE inhibitor enalapril, aspirin (unless contraindicated); administration of diuretics, nitrates, digoxin was also admitted). Group 1 received the activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1-receptor (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 2 received the activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 3 received the combination pharmaceutical composition comprising both activated potentiated form of antibodies to a C-terminal fragment of angiotensin II AT1-receptor (mixture of homeopathic dilutions C12, C30, C200) and activated potentiated form of antibodies to endothelial NO-synthase (mixture of homeopathic dilutions C12, C30, C200) (3 tablets/day, n=20). Group 4 received placebo (3 tablets/day, n=20). The groups were comparable in the initial study parameters: in age and sex, and severity (class of CHF and LVEF) and duration of the disease. Before and after treatment, the patients were evaluated for the life quality (Minnesota and Kansas questionnaires), morphological parameters of the heart, and tolerance to physical exercise.

Table 3 shows the results of the study in the form of dynamics in the basic parameters of the treatment efficacy.

After 6 months of treatment, the patients in group 1 treated with ULD of antibodies to a C-terminal fragment of angiotensin II AT1-receptor showed a significant improvement of the life quality, improvement of the left ventricular systolic function, and an increased tolerance to physical exercise. Group 2 showed a proven decrease in the anxiety and depression levels and in the life quality, which were evaluated using the Kansas questionnaire. The study confirmed that the maximum therapeutic effect was achieved with the combination pharmaceutical composition of the invention in combination with the standard CHF therapy, which was administered to patients from group 3 that showed a proven positive dynamics in all parameters under study.

The combination of activated (potentiated) forms of antibodies to a C-terminal fragment of angiotensin II AT1-receptor and to endothelial nitric oxide synthase (NO-synthase) in the pharmaceutical composition of the invention (combination drug) provides an unexpected synergistic therapeutic effect implying an enhanced influence on vascular remodeling and endothelium dysfunction that is critical for the CHF process and progression, as also on the improvement of the patients' life quality, on morphological parameters of the heart and tolerance to physical exercise, which is confirmed by clinical trials.

The results are set forth in Table 21.

Table 21

Groups/ Parameters	ULD ¹ of Abs ² to C-terminal fragment of AT1 receptor of angiotensin II			ULD of Abs to endothelial NO-synthase			Combination of ULD of Abs to C-terminal fragment of AT1 receptor of angiotensin II and ULD of Abs to endothelial NO-synthase			Placebo		
	[^]	^{&}	$\Delta\%$	[^]	^{&}	$\Delta\%$	[^]	^{&}	$\Delta\%$	[^]	^{&}	$\Delta\%$
Minnesota ³	47.5 ± 2.8	39.1 ±3.8 **	- 17.6	48. 1± 3.7	40. 8± 3.8	- 15.2	43. 9± 2.8	32.0± 4.9 ***\$	- 27.1	48.3 ± 3.7	42.4 ±2.9 **	- 12.2
Kansas ⁴	82.1 ± 2.3	70.1 ±5.5 ***	- 14.6	81. 5±2 .5	72. 0±8 .2 *	- 11.7	87, 7± 2.3	65.7±7. 3 ***\$	- 25.1	83.8 ± 3.5	60.3 ±6.8	- 7.2

HADS ^a	15.3 ± 1.0	12.5 ±0.9 **	- 18.5	16. 2±1 .7	11. 34± 2.1 ***	- 30.3	16. 2± 1.3	8.4±0.9 *** # \$\$	- 48.1	17.3 ± 1.1	15.9 ±1.1	- 8.1
FC CHF ^b	2.7± 0.1	2.2± 0.1* **	- 17.3	2.9± 0.1	2.7± 0.2	- 7.3	3.0 ± 0.2	1.9±0.1 *** # \$	- 36.6	2.7± 0.1	2.5±0 .1	- 6.2
FF LV ^c	27.1 ± 0.9	33. 6±1 .5**	24.0	28.2 ±1.5	25. 3±1 .7	10.3	25. 3± 1.1	34.6±1 .9 *** # \$	36.7	26.4± 1.1	28.0 ±1.4	6.3
6-minute walk test	378. 7±1 2.4	419. 6±1 3.7* **	10.8	383. 1± 15.3	416. 8±1 7.2	8.8	37 8.7 ±1 2.4	450.1± 17.7** # \$	18.9	390.5 ± 11.9	409.1 ±11.5	4.8

*, **, *** - p values < 0.05, 0.01 and 0.001, respectively

#- difference from group receiving ULDs of Abs to C-terminal fragment ATI of angiotensin receptor II with verifiable with p value < 0.05

\$, \$\$ - difference from the group receiving ULDs of Abs to endothelial NO-synthase is verifiable at p values of 0.05 and 0.01, respectively.

(1)-ULD means ultra low doses

(2) Abs means antibodies

(3) "Minnesota" denotes Minnesota Questionnaire

(4) "Kansas" denotes Kansas Questionnaire

(5) HADS denotes HADS total score

(6) FC CHF denotes patients with chronic heart failure, functional class

(7) FF LV denotes fraction of functioning of left ventricle.

Example 14.

To study properties of the proposed pharmaceutical composition in the treatment of patients with a benign prostatic hyperplasia, 300 mg pills were used, saturated with the pharmaceutical composition containing water-alcohol solutions (6 mg/pill) of activated - potentiated rabbit polyclonal affinity purified antibodies to prostate specific antigen (anti-PSA) and endothelial NO synthase (anti-eNOS) in ultra low doses (ULD), produced by ultra dilution of the initial matrix solution 100^{12} , 100^{30} , 100^{200} times, equivalent to the mixture of centesimal homeopathic dilutions C12, C30, C200 (ULD anti-PSA +anti-eNOS), and 300 mg pills, saturated with the pharmaceutical composition containing water-alcohol solutions (3 mg/pill) of activated potentiated rabbit polyclonal affinity purified antibodies to prostate specific antigen in ultra low doses (ULD), obtained by an ultra dilution of the initial matrix solution 100^{12} , 100^{30} , 100^{200} times, equivalent to the mixture of centesimal homeopathic dilutions C12, C30, C200 (ULD anti-PSA).

Benign prostatic hyperplasia (BPH) is one of the most frequently occurring disorders in males (Bruskewitz R.C., 2003; Rosen R., 2003): on the one hand, epidemiological studies, carried out in Russia, point to a gradual increase in frequency of BPH from 11.3% in 40-49 year olds to 81.4% in 80 year olds (Gorilovskiy, L.M., 1999); on the other hand, demographic studies conducted by WHO confirm a significant increase in the population over 60 years old, surpassing any other age group growth.

The main symptoms of benign prostatic hyperplasia are lower urinary tract symptoms, which can cause significant discomfort and decrease quality of life (Bruskewitz R.C., 2003; Lepor H., 2004; O'Leary M.P., 2005). In severe cases, the disease can be accompanied by complications, such as acute urinary retention, urinary tract infection, erythrorhithia, kidney failure (Stepanov, V.N., 1999; Jacobsen S.J., 1997;

Lepor H., 2004). BPH is also associated with development of erectile dysfunction in patients (Bruskewitz R.C., 2003; Daly MP, 2005).

An open-label comparative parallel group study of efficacy and safety the of pharmaceutical compositions containing ULD anti-PSA + ULD anti-eNOS and ULD anti-PSA in ameliorating urinary disturbances to caused by benign prostatic hyperplasia (BPH), included 40 patients selected in accordance with inclusions/exclusions criteria. Patients were randomized in 2 groups, one group received 1 pill 3 times per day during 12 weeks (n=21) of a ULD anti-PSA +anti-eNOS, and another one 1 pill 3 times per day during 12 weeks (n=19) of a ULD anti-PSA. The groups were comparable in age, severity of BPH symptoms, urination parameters and prostate volume.

The study included patients over 45 years old with a history of BPH with corresponding symptoms of lower urinary tract for no less than 6 months, IPSS \geq 13, prostate volume according to transrectal ultrasonography >30 cm³, with maximum urinary flow speed of \geq 4 ml/s and \leq 15 ml/sc and minimum residual urine volume equal to 125 ml, with PSA level <4 ng/ml. A necessary inclusion criterion was absence of intake of the following medications in the medical records: finasteride, dutasteride, or other experimental drug 6 months prior to inclusion in the study, a1-adrenoreceptor blockers and herbal medications 4 weeks prior to the inclusion into the study, any inhibitors of phosphodiesterase type 5 and other erectile dysfunction treatments 4 weeks prior to the inclusion into the study.

The study did not include patients undergone invasive methods of treatment of BPH, including transurethral prostatic resection, thermotherapy, transurethral needle ablation, stent angioplasty and other; with malignant oncological disease, acute urination delay, bladder stones, urethral stricture, Marion's disease, genitourinary system infections in the phase of active inflammation and others.

Clinical efficacy of pharmaceutical compositions was assessed by the improvement of clinical symptoms of lower urinary tract, evaluated using IPSS questionnaire (International Prostate Symptom Score), urination parameters (maximum and average urinary flow speed, urination volume, volume of residual urine) and prostate volume

based on the data of transurethral ultrasound (TU), and also erectile function was evaluated based on the data obtained from IIEF questionnaire (International Index of Erectile Function). Results of the study are shown in tables 22 and 23.

Table 22.

	ULD anti-PSA				ULD anti-PSA + ULD anti-eNOS			
	n/N(%) ¹	In., aver.	12 weeks., aver.	Δ, cp	n/N (%) ¹	In., aver.	12 weeks., aver.	Δ, cp
IPSS, score	19/19 (100.0)	17.8	11.9	-5.9	20/21 (95.2)	16.0	10.5	-5.6
QoL/, score (quality of life)	19/19 (100.0)	3.4	2.4	-1.0	20/21 (95.2)	3.4	2.3	-1.1
IIEF, score	2/19 (10.5)	17.8	18.6	0.8	4/21 (19.0)	17.5	18.9	1.4
Qmax, ml/s (maximum urine rate)	16/19 (84.2)	10.8	13.1	2.2	15/21 (71.4)	11.7	13.7	2.0
Qave, ml/s (average urine rate)	15/19 (78.9)	5.8	7.1	1.3	18/21 (85.7)	5.8	7.1	1.3
V, ml (volume of urination)	10/19 (52.6)	218.6	206.8	-11.8	15/21 (71.4)	203.7	252.0	48.3
RV, ml (residual volume of urine)	15-19 (78.9)	23.6	19.4	-4.3	14/21 (66.6)	19.1	14.1	-5.0
PV, cm ³ (prostate volume)	18/19 (94.7)	55.9	48.9	-7.0	15/21 (71.4)	57.0	52.4	-4.6

¹ – the numerator is a number of patients (n) showing improvement, denominator is total number of patients in the study (N).

Table 23.

Dynamics of subscales of obstructive and irritative symptoms, and question 7 of IPSS questionnaire

	ULD anti-PSA		ULD anti-PSA +anti-eNOS	
	M±SD Visit 1	M±SD Visit 2	M±SD Visit 1	M±SD Visit 2
Obstructive	10.0±3.02#	6.5±2.81***	8.2±2.96	6.0±3.39**
Irrit.	7.5±2.21&	5.3±1.90***	7.8±2.16&	4.5±2.34***
7 th question	2.1±0.78	1.9±0.75	2.3±0.90	1.4±0.98***
Obstr., % ²		-33.4±26.85		-25.2±34.50
Irrit., % ²		-28.2±1730		-40.3±30.35
7 th question, % ²		- 2.0±49.61##		-37.7±39.23

* - p<0.05 vs baseline; ** - p<0.01 vs vaseline; *** - p<0.001 vs baseline

- p<0,01 vs ULD anti-PSA

² - shows decrease compared to the baseline in %, average group value

The given data confirm that both ULD anti-PSA, and ULD anti-PSA + ULD anti-eNOS were used to effectively treat symptoms of lower urinary tract, increase average and maximum urinary flow speed, improve quality of life of patients (Table 22). The course of the was not long (12 weeks), therefore, a decrease in prostate volume was not observed in any study group. ULD anti-PSA did not effect the volume of urination, which increased only in 52.6% patients, on average the group showed some statistically insignificant decrease of urination volume by 11.8 ml (5.4%) compared to the baseline values. At the same time, patients, treated with ULD anti-PSA + ULD anti-eNOS, showed an increase in urination volume in 71.4%, and on average, an increase in volume was 48.3 ml (23.7%) compared to the baseline.

An analysis of dynamics of obstructive and irritative symptoms according to IPSS subscales as well as nocturia evidence (question 7 of IPSS) showed that both pharmaceutical compositions contributed to a decrease of obstruction and irritative symptoms, and also a decrease of nocturia symptoms. At the same time, a ULD anti-

PSA +anti-eNOS was more effective compared to a ULD anti-PSA in decreasing irritative symptoms of lower urinary tract (28.2% vs. 40.3%, p<0.05) and nighttime urination urges (2.0% vs. 37.7%,).

It should be noted, that ULD anti-PSA + ULD anti-eNOS is also more effective compared to ULD anti-PSA in improving erectile function in patients. In ULD anti-PSA + ULD anti-eNOS group, the total IIEF (International Index of Erectile Dysfunction) score increased by 19% in patients (in ULD anti-PSA group by 10.5%), an average increase of IIEF score in ULD anti-PSA + ULD anti-eNOS group was 8% vs 4.5% in a ULD anti-PSA group.

The pharmaceutical compositions showed excellent safety profile, no adverse effects related to the administered medications were observed in the course of study.

Therefore, ULD anti-PSA + ULD anti-eNOS showed better efficacy compared to that of ULD anti-PSA in treating urination problems caused by benign prostatic hyperplasia. In addition, a greater positive effect of ULD anti-PSA + ULD anti-eNOS on erectile function of patients compared to ULD anti-PSA was revealed.

What is claimed is:

1. A method of increasing the effect of an activated-potentiated form of an antibody to an endogenous biological molecule, said method comprising combining said endogenous biological molecule with an activated-potentiated form of an antibody to endothelial NO-synthase.
2. The method of claim 1, further comprising administering said combination to a patient in need of treatment with said activated-potentiated form of an antibody.
3. The method of claim 2, wherein said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to S-100 protein.
4. The method of claim 2, wherein said activated-potentiated form of an antibody to an endogenous biological molecule is an antibody to prostate specific antigen.
5. The method of claim 2, wherein said activated-potentiated form an antibody to an endogenous biological molecule is an antibody to insulin receptor.
6. The method of claim 2, wherein said activated-potentiated form an antibody to an endogenous biological molecule is an antibody to angiotensin receptor II.
7. A pharmaceutical composition comprising a) an activated-potentiated form of an antibody to an endogenous biological molecule, and b) an activated-potentiated form of an antibody to NO synthase.
8. The pharmaceutical composition of claim 7, further comprising pharmaceutically acceptable solid carrier.
9. The pharmaceutical composition of claim 8, wherein said activated-potentiated form of an antibody to endothelial NO-synthase is in the form of a mixture of C12, C30, and C200 homeopathic dilutions impregnated onto said solid carrier.
10. The pharmaceutical composition of claim 7, wherein said activated-potentiated form of an antibody to an endogenous biological molecule is a monoclonal, monoclonal, or natural antibody.


11. The pharmaceutical composition of claim 10, wherein said antibody to an endogenous biological molecule is a polyclonal antibody.

12. The pharmaceutical composition of claim 7, wherein said activated-potentiated form of an antibody to an endogenous biological molecule is prepared by successive centesimal dilutions coupled with shaking of every dilution.


13. The pharmaceutical composition of claims 7, wherein said antibody to endothelial NO-synthase is monoclonal, polyclonal or natural antibody.

14. The pharmaceutical composition of claim 8, wherein said antibody to endothelial NO-synthase is a polyclonal antibody.

15. The pharmaceutical composition of claim 9, the activated-potentiated form of an antibody to endothelial NO-synthase is prepared by successive centesimal dilutions coupled with shaking of every dilution.



* differences are statistically significant relative to control (distilled water), $p < 0.05$



* differences rare statistically significant in reference to control (distilled water), p<0.05

