





(86) Date de dépôt PCT/PCT Filing Date: 2013/06/28
(87) Date publication PCT/PCT Publication Date: 2014/01/03
(85) Entrée phase nationale/National Entry: 2014/12/12
(86) N° demande PCT/PCT Application No.: US 2013/048697
(87) N° publication PCT/PCT Publication No.: 2014/005080
(30) Priorités/Priorities: 2012/06/29 (US61/666,535);
2013/03/14 (US61/782,894)

(51) Cl.Int./Int.Cl. *A61K 31/343* (2006.01),
A61K 31/4178 (2006.01), *A61K 31/4525* (2006.01),
A61P 39/00 (2006.01), *A61P 9/10* (2006.01),
C07D 493/08 (2006.01)
(71) Demandeur/Applicant:
LIXTE BIOTECHNOLOGY, INC., US
(72) Inventeur/Inventor:
KOVACH, JOHN S., US
(74) Agent: BENOIT & COTE INC.

(54) Titre : OXABICYCLOHEPTANES ET OXABICYCLOHEPTENES POUR LE TRAITEMENT D'UNE LESION DE
REPERFUSION

(54) Title: OXABICYCLOHEPTANES AND OXABICYCLOHEPTENES FOR THE TREATMENT OF REPERFUSION
INJURY



(57) Abrégé/Abstract:

A method of reducing reperfusion injury in mammalian tissue comprising contacting the tissue with a protein phosphatase 2A (PP2A) inhibitor having the structure.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
3 January 2014 (03.01.2014)



(10) International Publication Number
WO 2014/005080 A1

(51) International Patent Classification:
C07D 493/08 (2006.01)

(21) International Application Number:
PCT/US2013/048697

(22) International Filing Date:
28 June 2013 (28.06.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/666,535 29 June 2012 (29.06.2012) US
61/782,894 14 March 2013 (14.03.2013) US

(71) Applicant (for all designated States except US): **LIXTE BIOTECHNOLOGY, INC.** [US/US]; 248 Route 25a No. 2, East Setauket, NY 11733 (US).

(72) Inventor; and

(71) Applicant (for US only): **KOVACH, John, S.** [US/US]; 6 Tinker Lane, East Setauket, NY 11733 (US).

(74) Agent: **WHITE, John, P.**; Cooper & Dunham LLP, 30 Rockefeller Plaza, New York, NY 10112 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— with international search report (Art. 21(3))


[Continued on next page]

(54) Title: OXABICYCLOHEPTANES AND OXABICYCLOHEPTENES FOR THE TREATMENT OF REPERFUSION INJURY

A



(57) Abstract: A method of reducing reperfusion injury in mammalian tissue comprising contacting the tissue with a protein phosphatase 2A (PP2A) inhibitor having the structure.



WO 2014/005080 A1



-
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

OXABICYCLOHEPTANES AND OXABICYCLOHEPTENES FOR
THE TREATMENT OF REPERFUSION INJURY

5

Throughout this application various publications are referenced. The disclosures of these documents in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention
10 pertains.

BACKGROUND OF THE INVENTION

Reperfusion is a re-establishment of blood flow and re-oxygenation of an affected area following an ischemic event and is critical to
15 limit irreversible damage. However, the absence of oxygen and nutrients from the blood creates a condition in which reperfusion injury may occur. The restoration of blood flow after an ischemic event results in inflammation and oxidative damage. Upon restoration of blood flow, white blood cells release inflammatory
20 factors such as interleukins as well as free radicals. The restored blood flow reintroduces oxygen within cells that damages cellular proteins, DNA, and the plasma membrane.

As acute myocardial infarction (MI) remains the leading cause of
25 death worldwide, the possibility that a pharmacologic intervention applied promptly but after the onset of the heart attack would minimize damage to heart tissue caused by reperfusion and therefore be expected to save many lives and reduce the number of individuals with heart failure following excessive cardiac muscle damage after
30 an MI (Yellon and Hausenloy, 2005; Longacre et al, 2011). It has been suggested that a lack of commercial interest in a developing a drug that would likely be used only once in an individual has limited progress in this field (Cohen and Downey, 2011).

35 At present, the only established intervention that consistently reduces the size of myocardial infarcts in humans is by improving coronary artery flow as soon as possible after an MI either by drugs, which dissolve fresh clots and/or cardiac catheterization

with balloon angioplasty with or without placement of a fixed conduit, a stent. These methods of improving coronary artery blood flow (reperfusion) have improved patient care and decreased hospital mortality. However, delay in initiating reperfusion because of travel time to a cardiac center is a serious limitation to applying these treatments to patients with acute cardiac injury. It has also been discovered the reperfusion treatment in and of itself may cause myocardial cell death, a phenomenon called reperfusion injury. Reducing injury caused by reperfusion by pharmacologic means should improve the success of current interventions for acute heart attacks. A drug minimizing tissue damage that could be administered at the time of a MI by emergency personnel prior to arrival at a cardiac center could be a major advance in the care of heart attack victims. Acute injury due to oxygen deprivation leading to myocardial damage is also a significant problem in heart surgery. The incidence of infarction after coronary artery bypass graft surgery has been estimated to be as high as 19% with attendant cardiac morbidity (Longacre et al, 2011).

20

25


30

35

SUMMARY OF THE INVENTION

A method of reducing reperfusion injury in mammalian tissue comprising contacting the tissue with a protein phosphatase 2A (PP2A) inhibitor having the structure:

5



wherein


bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,

10 where R_9 is H, alkyl, alkenyl, alkynyl or aryl,
or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,
 $O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,

15



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12}

alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,

20 substituted aryl where the substituent is other
than chloro when R_1 and R_2 are $=O$,



4

$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,
 wherein each R_{11} is independently alkyl,
 alkenyl or alkynyl, or H;

5 R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken
 together are =O;
 R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or
 SR_{12} ,

where R_{12} is H, alkyl, alkenyl, alkynyl or aryl; and

10


each occurrence of alkyl, alkenyl, or alkynyl is branched or
 unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

15

A method of reducing tissue damage associated with reperfusion
 injury in the heart of a subject following a myocardial infarction
 comprising administering to the subject a therapeutically effective
 amount of a protein phosphatase 2A (PP2A) inhibitor having the

20 structure:



wherein

bond α is present or absent;

25 R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,


or R_1 and R_2 together are =O;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,

$\text{O}(\text{CH}_2)_{1-6}\text{R}_9$, SH, S^- , SR_9 ,

30

5



where X is O, S, NR₁₀, or N⁺R₁₀R₁₀,

where each R₁₀ is independently H, alkyl, C₂-C₁₂ alkyl, alkenyl, C₄-C₁₂ alkenyl, alkynyl, aryl, substituted aryl where the substituent is other than chloro when R₁ and R₂ are =O,



-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,

wherein each R₁₁ is independently alkyl, alkenyl or alkynyl, or H;

R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;

R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,


where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

A method of reducing vascular leakage associated with reperfusion injury in a subject suffering from sepsis comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:

6



wherein


bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,

5 where R_9 is H, alkyl, alkenyl, alkynyl or aryl,
or R_1 and R_2 together are =O;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,
 $O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,

10



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12}
alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,
15 substituted aryl where the substituent is other
than chloro when R_1 and R_2 are =O,

15



$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

wherein each R_{11} is independently alkyl,
20 alkenyl or alkynyl, or H;

20

7

R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;

R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,

5 where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

10 or a salt, enantiomer or zwitterion of the compound.

15

Brief Description of the Figures

Figure 1. Photographs of in-bred "control" rats 1-10 implanted with a pump containing sodium chloride prior to raising a skin flap.

5 Photographs were taken 7 days after creation of the skin flap. (A) control rat 1; (B) control rat 2; (C) control rat 3; (D) control rat 4; (E) control rat 5; (F) control rat 6; (G) control rat 7; (H) control rat 8; (I) control rat 9; and (J) control rat 10.

10 **Figure 2.** Photographs of in-bred "treatment" rats 1-10 implanted with a pump containing 0.55 mg of LB-100 prior to raising a skin flap. The pump was set to deliver 0.5 μ l/hour $\pm$ 10% so as to administer about 12 μ l/day for 8 days, four days prior to raising the graft and for the first four days after creation of the graft.


15 Photographs were taken 7 days after creation of the skin flap. (A) treatment rat 1; (B) treatment rat 2; (C) treatment rat 3; (D) treatment rat 4; (E) treatment rat 5; (F) treatment rat 6; (G) treatment rat 7; (H) treatment rat 8; (I) treatment rat 9; and (J) treatment rat 10.

20

DETAILED DESCRIPTION OF THE INVENTION

A method of reducing reperfusion injury in mammalian tissue comprising contacting the tissue with a protein phosphatase 2A (PP2A) inhibitor having the structure:

5



wherein


bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,

10 where R_9 is H, alkyl, alkenyl, alkynyl or aryl,
or R_1 and R_2 together are =O;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,
 $O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,

15



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12}
alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,
20 substituted aryl where the substituent is other
than chloro when R_1 and R_2 are =O,



10

-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,
wherein each R₁₁ is independently alkyl,
alkenyl or alkynyl, or H;

5 R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken
together are =O;
R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or
SR₁₂,
where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and

10

each occurrence of alkyl, alkenyl, or alkynyl is branched or
unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

15

In some embodiments, the method wherein the reduction of
reperfusion injury comprises increased phosphorylation of Akt in
the mammalian tissue that has suffered an ischemia.

20 In some embodiments, the method wherein the reduction of
reperfusion injury comprises increased activation of Akt in the
mammalian tissue that has suffered an ischemia.

In some embodiments, the method wherein the reduction of
25 reperfusion injury comprises increased phosphorylation of BAD,
mdm2, eNOS and/or GSK-3 β in the mammalian tissue that has suffered
an ischemia.

In some embodiments, the method wherein the ischemia is caused by a
30 myocardial infarction, stroke or sepsis.


In some embodiments, the method wherein the tissue is myocardial
tissue, brain tissue or endothelial tissue.

35 In some embodiments, the method wherein endothelial dysfunction is
reduced.

11

In some embodiments, the method wherein the tissue is myocardial tissue, brain tissue or endothelial tissue.

A method of reducing tissue damage associated with reperfusion injury in the heart of a subject following a myocardial infarction comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:



wherein

bond α is present or absent;


R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,

$O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12}

alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,

substituted aryl where the substituent is other

than chloro when R_1 and R_2 are $=O$,

12



-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,

wherein each R₁₁ is independently alkyl, alkenyl or alkynyl, or H;

5

R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;

R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,

10

where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and


each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

15

or a salt, enantiomer or zwitterion of the compound.

A method of reducing vascular leakage associated with reperfusion injury in a subject suffering from sepsis comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:

20



wherein

bond α is present or absent;

25

R₁ and R₂ is each independently H, O⁻ or OR₉,


where R₉ is H, alkyl, alkenyl, alkynyl or aryl,

or R₁ and R₂ together are =O;

R₃ and R₄ are each different, and each is OH, O⁻, OR₉, OR₁₀,


O(CH₂)₁₋₆R₉, SH, S⁻, SR₉,

13



where X is O, S, NR₁₀, or N⁺R₁₀R₁₀,

- 5 where each R₁₀ is independently H, alkyl, C₂-C₁₂ alkyl, alkenyl, C₄-C₁₂ alkenyl, alkynyl, aryl, substituted aryl where the substituent is other than chloro when R₁ and R₂ are =O,



- 10 -CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,
 wherein each R₁₁ is independently alkyl, alkenyl or alkynyl, or H;

- 15 R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;
 R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,
 where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and


- 20 each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

- 25 In some embodiments, wherein the reperfusion injury is caused by ischemia that is caused by septic shock.

In one embodiment, the protein phosphatase 2A inhibitor has the structure

14



wherein



bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,


5 where R_9 is H, alkyl, alkenyl, alkynyl or aryl,
or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , SH, S^- ,
 SR_9 ,

10



or



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

15

where each R_{10} is independently H, alkyl, C_2-C_{12}
alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,
substituted aryl where the substituent is other
than chloro when R_1 and R_2 are $=O$,



20

$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

15


where each R_{11} is independently alkyl,
alkenyl, alkynyl, or H;

5 R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken
together are =O; and


R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or
 SR_{12} , where R_{12} is H, aryl, alkyl, alkenyl or alkynyl,

10 or a salt, enantiomer or zwitterion of the compound.


In one embodiment, the protein phosphatase 2A inhibitor has the
structure



15 In one embodiment, the protein phosphatase 2A inhibitor has the
structure



In one embodiment, the protein phosphatase 2A inhibitor has the
20 structure




16

In one embodiment, bond α is present. In another embodiment, bond α is absent.

In one embodiment, R_1 and R_2 together are =O;

5

R_3 is O^- or OR_9 , where R_9 is H, methyl, ethyl or phenyl;



10

where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

15



$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

where R_{11} is alkyl, alkenyl or alkynyl, each of which is substituted or unsubstituted, or H;

20

R_5 and R_6 taken together are =O; and

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,

25


where R_{12} is a substituted or unsubstituted alkyl, alkenyl or alkynyl.

In one embodiment, R_3 is O^- .

30


In another embodiment, R_4 is

17




where X is O, NR_{10} , $\text{N}^+\text{R}_{10}\text{R}_{10}$

- 5 where each R_{10} is independently H, alkyl, substituted $\text{C}_2\text{-C}_{12}$ alkyl, alkenyl, substituted $\text{C}_4\text{-C}_{12}$ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro when R_1 and R_2 are =O,




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,
where R_{11} is H or alkyl.


- 15 In one embodiment, the protein phosphatase inhibitor 2A has the structure



18




In one embodiment, R₄ is



where R₁₀ is H, alkyl, substituted C₂-C₁₂ alkyl, alkenyl, substituted C₄-C₁₂ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro when R₁ and R₂ are =O,


10




-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂, where R₁₁ is H or alkyl.

15

In one embodiment, R₄ is




20 In one embodiment, R₄ is




19


where R_{10} is




5 In one embodiment, R_4 is




where R_{10} is



10 In one embodiment, R_4 is




In one embodiment, R_4 is



15

In one embodiment, R_5 and R_6 together are =O. In another embodiment, R_7 and R_8 are each H.

In one embodiment,




20

wherein bond α is present or absent;

20

R_9 is present or absent and when present is H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl or phenyl; and X is O, S, NR_{10} or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the
 5 substituent is other than chloro,




10 $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-CH_2CN$, or $-CH_2CH_2R_{16}$, where R_{11} is H or alkyl, and where R_{16} is any substituent that is a precursor to an aziridinyl intermediate,

or a salt, zwitterion or enantiomer of the compound.

15

In one embodiment, the protein phosphatase 2A inhibitor has the structure



20


wherein,

bond α is present or absent;

X is O, S, NR_{10} or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the
 25 substituent is other than chloro,

21




5 $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{CH}_2\text{CN}$, or $-\text{CH}_2\text{CH}_2\text{R}_{16}$, where R_{11} is H or alkyl, and where R_{16} is any substituent that is a aziridinyl intermediate,

or a salt, zwitterion or enantiomer of a compound.

In one embodiment,

10 X is O or NH^+R_{10} ,

where R_{10} is H, alkyl, substituted $\text{C}_2\text{-C}_{12}$ alkyl, alkenyl, substituted $\text{C}_4\text{-C}_{12}$ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,



15

In one embodiment, X is $-\text{CH}_2\text{CH}_2\text{R}_{16}$, where R_{16} is any substituent that is a precursor to an aziridinyl intermediate.


In one embodiment, X is O.

20


In another embodiment, X is NH^+R_{10} ,

where R_{10} H, alkyl, substituted $\text{C}_2\text{-C}_{12}$ alkyl, alkenyl, substituted $\text{C}_4\text{-C}_{12}$ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

25




In one embodiment, R_{10} is methyl. In another embodiment, R_{10} is



22


In one embodiment, R_{10} is



In one embodiment, R_{10} is ethyl. In another embodiment, R_{10} is absent.

5

In one embodiment, the protein phosphatase 2A inhibitor has the structure



10 wherein


bond α is present or absent;

R_9 is present or absent and when present is H, alkyl, alkenyl, alkynyl or phenyl; and

15 X is O, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2-C_{12} alkyl, alkenyl, substituted C_4-C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

20




$-CH_2CN$, $-CH_2CO_2R_{12}$, or $-CH_2COR_{12}$,
where R_{12} is H or alkyl,

25 or a salt, zwitterion, or enantiomer of the compound.

23

In one embodiment, the protein phosphatase 2A inhibitor has the structure




wherein

5 bond α is present or absent;

X is O or NH^+R_{10} ,

where R_{10} is H, alkyl, substituted $\text{C}_2\text{-C}_{12}$ alkyl, alkenyl, substituted $\text{C}_4\text{-C}_{12}$ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

10




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{12}$, or $-\text{CH}_2\text{COR}_{12}$, where R_{12} is H or alkyl.

15






In one embodiment, bond α is present. In another embodiment, bond α is absent.

In one embodiment, the protein phosphatase 2A inhibitor has the structure

20




24






, or

25




In one embodiment, the protein phosphatase 2A inhibitor has the structure


5



10




26



5

In one embodiment, the protein phosphatase 2A inhibitor has the structure




10 wherein

bond α is present or absent;

X is NH^+R_{10} ,

where R_{10} is present or absent and when present R_{10} is alkyl, substituted C2-C12 alkyl, alkenyl, substituted C4-C12 alkenyl,

15




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{12}$, or $-\text{CH}_2\text{COR}_{12}$, where R_{12} is H or alkyl.


In one embodiment of the method, the protein phosphatase 2A inhibitor has the structure

20

27



5 In one embodiment of the method, the protein phosphatase 2A inhibitor has the structure



wherein


bond α is present or absent;

10 R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, substituted alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is $O(CH_2)_{1-6}R_9$ or OR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl,

hydroxyalkyl, C_2-C_{12} alkyl, alkenyl, C_4-C_{12} alkenyl,

alkynyl, aryl, substituted aryl where the

20 substituent is other than chloro when R_1 and R_2 are $=O$,


28



$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,

where each R_{11} is independently alkyl, alkenyl or alkynyl, each of which is substituted or unsubstituted, or H;

5 or R_3 and R_4 are each different and each is OH or



R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken together are =O;

10

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,

where R_{12} is H, aryl or a substituted or unsubstituted alkyl, alkenyl or alkynyl; and

15


each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

20

In one embodiment of the method, the protein phosphatase 2A inhibitor has the structure

29



In one embodiment of the method, the bond α is present.

In one embodiment of the method, the bond α is absent.


5 In one embodiment of the method,

R_3 is OR_9 or $O(CH_2)_{1-6}R_9$,

where R_9 is aryl, substituted ethyl or substituted phenyl,

wherein the substituent is in the para position of the phenyl;

R_4 is



where X is O , S , NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H , alkyl, hydroxyalkyl, substituted C_2-C_{12} alkyl, alkenyl, substituted C_4-C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

15




$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

where R_{11} is alkyl, alkenyl or alkynyl, each of which is substituted or unsubstituted, or H ;

20


30

or where R_3 is OH and R_4 is




5 In one embodiment of the method,

R_4 is



where R_{10} is alkyl or hydroxylalkyl



or R_4 is

when R_3 is OH.

10

In one embodiment of the method,


R_1 and R_2 together are =O;

R_3 is OR_9 or $O(CH_2)_{1-2}R_9$,

15


where R_9 is aryl, substituted ethyl, or substituted phenyl, wherein the substituent is in the para position of the phenyl;

or R_3 is OH and R_4 is



20

R_4 is



where R_{10} is alkyl or hydroxyl alkyl;

31

R_5 and R_6 together are =O; and


R_7 and R_8 are each independently H.

5 In one embodiment of the method,

R_1 and R_2 together are =O;



R_3 is OH, $O(CH_2)R_9$, or OR_9 ,

where R_9 is phenyl or CH_2CCl_3 ,



10

R_4 is



15

where R_{10} is CH_3 or CH_3CH_2OH ;

R_5 and R_6 together are =O; and

R_7 and R_8 are each independently H.

20 In one embodiment of the method,

R_3 is OR_9 ,

where R_9 is $(CH_2)_{1-6}(CHNHBoc)CO_2H$, $(CH_2)_{1-6}(CHNH_2)CO_2H$, or $(CH_2)_{1-6}CCl_3$.

25 In one embodiment of the method,

R_9 is $CH_2(CHNHBoc)CO_2H$, $CH_2(CHNH_2)CO_2H$, or CH_2CCl_3 .

In one embodiment of the method,

32


R_3 is $O(CH_2)_{1-6}R_9$ or $O(CH_2)R_9$,
where R_9 is phenyl.

In one embodiment of the method,


5 R_3 is $O(CH_2)R_9$,
where R_9 is phenyl.

In one embodiment of the method,

10 R_3 is OH and R_4 is




In one embodiment of the method,




15 R_4 is
wherein R_{10} is alkyl or hydroxyalkyl.

In one embodiment of the method, R_{11} is $-CH_2CH_2OH$ or $-CH_3$.




In one embodiment of the method, the protein phosphatase 2A inhibitor has the structure



20






33




5

In one embodiment of the method, the protein phosphatase 2A inhibitor has the structure



10

34



5 For the foregoing embodiments, each embodiment disclosed herein is contemplated as being applicable to each of the other disclosed embodiments. Thus, all combinations of the various elements described herein are within the scope of the invention.

10 Definitions

As used herein, and unless otherwise stated, each of the following terms shall have the definition set forth below.

15 In particular, the invention is directed to the treatment or prevention of reperfusion injury.

As used herein, "reperfusion injury" is tissue damage, tissue death, cell damage, cell death, vascular leakage or endothelial dysfunction caused when blood supply returns to tissue, cells or blood vessels after a period of ischemia or lack of oxygen.

25 As used herein, "myocardial infarction" (MI), also known as a heart attack, is an infarction of the heart, causing cardiac tissue damage. This is most commonly due to occlusion (blockage) of a coronary artery following the rupture of a vulnerable atherosclerotic plaque, which is an unstable collection of lipids (fatty acids) and white blood cells (especially macrophages) in the wall of an artery. The resulting ischemia (restriction in blood

supply) and oxygen shortage, if left untreated for a sufficient period of time, can cause damage or death of heart muscle tissue (myocardium) due to reperfusion injury.

5 Examples of conditions caused by ischemia and that result in reperfusion injury include, but are not limited to, myocardial infarction; cerebral infarction (stroke) due to a disturbance in the blood vessels supplying blood to the brain; pulmonary infarction or lung infarction; Splenic infarction occurs when the
10 splenic artery or one of its branches are occluded, for example by a blood clot; Limb infarction caused by arterial embolisms; skeletal muscle infarction caused by diabetes mellitus; bone infarction; testicle infarction; and sepsis.

15 As used herein, a "symptom" associated with reperfusion injury includes any clinical or laboratory manifestation associated with reperfusion injury and is not limited to what the subject can feel or observe.

20 As used herein, "treatment of the diseases" or "treating", e.g. of reperfusion injury, encompasses inducing prevention, inhibition, regression, or stasis of the disease or a symptom or condition associated with the disease.

25 As used herein, "inhibition" of disease progression or disease complication in a subject means preventing or reducing the disease progression and/or disease complication in the subject.

As used herein, "alkyl" is intended to include both branched and
30 straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. Thus, C_1-C_n as in " C_1-C_n alkyl" is defined to include groups having 1, 2, ..., $n-1$ or n carbons in a linear or branched arrangement, and specifically includes methyl, ethyl, propyl, butyl, pentyl, hexyl, and so on. An embodiment can
35 be C_1-C_{12} alkyl. "Alkoxy" represents an alkyl group as described above attached through an oxygen bridge.

The term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight or branched, containing at least 1 carbon to carbon double bond, and up to the maximum possible number of non-aromatic carbon-carbon double bonds may be present. Thus, C_2-C_n alkenyl is defined to include groups having 1, 2, ..., $n-1$ or n carbons. For example, "C₂-C₆ alkenyl" means an alkenyl radical having 2, 3, 4, 5, or 6 carbon atoms, and at least 1 carbon-carbon double bond, and up to, for example, 3 carbon-carbon double bonds in the case of a C₆ alkenyl, respectively. Alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated. An embodiment can be C₂-C₁₂ alkenyl.

The term "alkynyl" refers to a hydrocarbon radical straight or branched, containing at least 1 carbon to carbon triple bond, and up to the maximum possible number of non-aromatic carbon-carbon triple bonds may be present. Thus, C_2-C_n alkynyl is defined to include groups having 1, 2, ..., $n-1$ or n carbons. For example, "C₂-C₆ alkynyl" means an alkynyl radical having 2 or 3 carbon atoms, and 1 carbon-carbon triple bond, or having 4 or 5 carbon atoms, and up to 2 carbon-carbon triple bonds, or having 6 carbon atoms, and up to 3 carbon-carbon triple bonds. Alkynyl groups include ethynyl, propynyl and butynyl. As described above with respect to alkyl, the straight or branched portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated. An embodiment can be a C₂-C_n alkynyl.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 10 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydro-naphthyl, indanyl, biphenyl, phenanthryl, anthryl or acenaphthyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring. The substituted aryls included in this invention include substitution at any suitable position with amines, substituted amines, alkylamines, hydroxys and alkylhydroxys, wherein the "alkyl"

portion of the alkylamines and alkylhydroxys is a C₂-C_n alkyl as defined hereinabove. The substituted amines may be substituted with alkyl, alkenyl, alkynyl, or aryl groups as hereinabove defined.

5 The alkyl, alkenyl, alkynyl, and aryl substituents may be unsubstituted or unsubstituted, unless specifically defined otherwise. For example, a (C₁-C₆) alkyl may be substituted with one or more substituents selected from OH, oxo, halogen, alkoxy, dialkylamino, or heterocyclyl, such as morpholinyl, piperidinyl,
10 and so on.

In the compounds of the present invention, alkyl, alkenyl, and alkynyl groups can be further substituted by replacing one or more hydrogen atoms by non-hydrogen groups described herein to the
15 extent possible. These include, but are not limited to, halo, hydroxy, mercapto, amino, carboxy, cyano and carbamoyl.

The term "substituted" as used herein means that a given structure has a substituent which can be an alkyl, alkenyl, or aryl group as
20 defined above. The term shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently
25 substituted, it is meant that the (two or more) substituents can be the same or different.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of
30 ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one group, it is understood that these
35 multiple groups may be on the same carbon or on different carbons, so long as a stable structure results.

As used herein, a "compound" is a small molecule that does not include proteins, peptides or amino acids.

As used herein, an "isolated" compound is a compound isolated from
5 a crude reaction mixture or from a natural source following an affirmative act of isolation. The act of isolation necessarily involves separating the compound from the other components of the mixture or natural source, with some impurities, unknown side products and residual amounts of the other components permitted to
10 remain. Purification is an example of an affirmative act of isolation.

As used herein, "administering" an agent may be performed using any of the various methods or delivery systems well known to those
15 skilled in the art. The administering can be performed, for example, orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery, subcutaneously,
20 intraadiposally, intraarticularly, intrathecally, into a cerebral ventricle, intraventricularly, intratumorally, into cerebral parenchyma or intraparenchymally.

The following delivery systems, which employ a number of routinely
25 used pharmaceutical carriers, may be used but are only representative of the many possible systems envisioned for administering compositions in accordance with the invention.

Injectable drug delivery systems include solutions, suspensions,
30 gels, microspheres and polymeric injectables, and can comprise excipients such as solubility-altering agents (e.g., ethanol, propylene glycol and sucrose) and polymers (e.g., polycaprylactones and PLGA's).

35 Other injectable drug delivery systems include solutions, suspensions, gels. Oral delivery systems include tablets and capsules. These can contain excipients such as binders (e.g., hydroxypropylmethylcellulose, polyvinyl pyrrolidone, other

cellulosic materials and starch), diluents (e.g., lactose and other sugars, starch, dicalcium phosphate and cellulosic materials), disintegrating agents (e.g., starch polymers and cellulosic materials) and lubricating agents (e.g., stearates and talc).

5

Implantable systems include rods and discs, and can contain excipients such as PLGA and polycaprylactone.

Oral delivery systems include tablets and capsules. These can
10 contain excipients such as binders (e.g., hydroxypropylmethylcellulose, polyvinyl pyrrolidone, other cellulosic materials and starch), diluents (e.g., lactose and other sugars, starch, dicalcium phosphate and cellulosic materials), disintegrating agents (e.g., starch polymers and cellulosic
15 materials) and lubricating agents (e.g., stearates and talc).

Transmucosal delivery systems include patches, tablets, suppositories, pessaries, gels and creams, and can contain excipients such as solubilizers and enhancers (e.g., propylene
20 glycol, bile salts and amino acids), and other vehicles (e.g., polyethylene glycol, fatty acid esters and derivatives, and hydrophilic polymers such as hydroxypropylmethylcellulose and hyaluronic acid).

25 Dermal delivery systems include, for example, aqueous and nonaqueous gels, creams, multiple emulsions, microemulsions, liposomes, ointments, aqueous and nonaqueous solutions, lotions, aerosols, hydrocarbon bases and powders, and can contain excipients such as solubilizers, permeation enhancers (e.g., fatty acids,
30 fatty acid esters, fatty alcohols and amino acids), and hydrophilic polymers (e.g., polycarbophil and polyvinylpyrrolidone). In one embodiment, the pharmaceutically acceptable carrier is a liposome or a transdermal enhancer.

35 Solutions, suspensions and powders for reconstitutable delivery systems include vehicles such as suspending agents (e.g., gums, xanthans, cellulose and sugars), humectants (e.g., sorbitol), solubilizers (e.g., ethanol, water, PEG and propylene glycol),

surfactants (e.g., sodium lauryl sulfate, Spans, Tweens, and cetyl pyridine), preservatives and antioxidants (e.g., parabens, vitamins E and C, and ascorbic acid), anti-caking agents, coating agents, and chelating agents (e.g., EDTA).

5

As used herein, "pharmaceutically acceptable carrier" refers to a carrier or excipient that is suitable for use with humans and/or animals without undue adverse side effects (such as toxicity, irritation, and allergic response) commensurate with a reasonable benefit/risk ratio. It can be a pharmaceutically acceptable solvent, suspending agent or vehicle, for delivering the instant compounds to the subject.

The compounds used in the method of the present invention may be in a salt form. As used herein, a "salt" is a salt of the instant compounds which has been modified by making acid or base salts of the compounds. In the case of compounds used to treat an infection or disease, the salt is pharmaceutically acceptable. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as phenols. The salts can be made using an organic or inorganic acid. Such acid salts are chlorides, bromides, sulfates, nitrates, phosphates, sulfonates, formates, tartrates, maleates, malates, citrates, benzoates, salicylates, ascorbates, and the like. Phenolate salts are the alkaline earth metal salts, sodium, potassium or lithium. The term "pharmaceutically acceptable salt" in this respect, refers to the relatively non-toxic, inorganic and organic acid or base addition salts of compounds of the present invention. These salts can be prepared in situ during the final isolation and purification of the compounds of the invention, or by separately reacting a purified compound of the invention in its free base or free acid form with a suitable organic or inorganic acid or base, and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate,

lactobionate, and laurylsulphonate salts and the like. (See, e.g., Berge et al. (1977) "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19).

5 As used herein, an "amount" or "dose" of an agent measured in milligrams refers to the milligrams of agent present in a drug product, regardless of the form of the drug product.

As used herein, the term "therapeutically effective amount" or
10 "effective amount" refers to the quantity of a component that is sufficient to yield a desired therapeutic response without undue adverse side effects (such as toxicity, irritation, or allergic response) commensurate with a reasonable benefit/risk ratio when used in the manner of this invention. The specific effective amount
15 will vary with such factors as the particular condition being treated, the physical condition of the patient, the type of mammal being treated, the duration of the treatment, the nature of concurrent therapy (if any), and the specific formulations employed and the structure of the compounds or its derivatives.

20

Where a range is given in the specification it is understood that the range includes all integers and 0.1 units within that range, and any sub-range thereof. For example, a range of 77 to 90% is a disclosure of 77, 78, 79, 80, and 81% etc.

25

As used herein, "about" with regard to a stated number encompasses a range of +one percent to -one percent of the stated value. By way of example, about 100 mg/kg therefore includes 99, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9, 100, 100.1, 100.2, 100.3,
30 100.4, 100.5, 100.6, 100.7, 100.8, 100.9 and 101 mg/kg. Accordingly, about 100 mg/kg includes, in an embodiment, 100 mg/kg. It is understood that where a parameter range is provided, all integers within that range, and tenths thereof, are also provided by the invention. For example, "0.2-5 mg/kg/day" is a disclosure of 0.2
35 mg/kg/day, 0.3 mg/kg/day, 0.4 mg/kg/day, 0.5 mg/kg/day, 0.6 mg/kg/day etc. up to 5.0 mg/kg/day.

All combinations of the various elements described herein are within the scope of the invention.

This invention will be better understood by reference to the
5 Experimental Details which follow, but those skilled in the art
will readily appreciate that the specific experiments detailed are
only illustrative of the invention as described more fully in the
claims which follow thereafter.

10

15

20

25


30

35

Experimental Details**Example 1. Synthesis of LB-107**


LB-107 (**5**) was prepared by reacting acid **3** with N-methylpiperazine (**4**) in the presence of EDC. In order to prepare **5** in better yields three different methods were attempted. In the first method, one pot reaction on LB-100 using thionyl chloride in methanol was attempted but no product was observed. In the second method, acid chloride of LB-100 was allowed to react with methanol in presence of triethylamine/DMAP to give the desired methyl ester. The methyl ester thus obtained was in low yields and the separation of triethylamine from the product was also tedious. Hence a two-step procedure was used. In this third method, endotal (**1**) when heated under reflux in methanol gave the desired monomethylester **3** in 95% yields. Compound **3** when treated with N-methylpiperazine (**4**) in presence of EDC and a catalytic amount of N-hydroxybenzotriazole gave the required methyl ester **5** in 39% yields after purification with column chromatography.

7-Oxa-bicyclo [2,2,1]heptane-2,3-dicarboxylic acid monomethyl ester (3**):**



The mixture of exo-7-Oxabicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride (**1**, 10.0 g, 59.5 mmole) and dry methanol (**2**, 50 mL) was heated at reflux temperature for 3.5 h. The reaction mixture became homogeneous during the reflux. The reaction mixture was then cooled down to room temperature and concentrated to give **3** (11.3 g, 95%) as crystalline white material. The crude ^1H NMR was clean enough with no extra peaks. Hence this material was utilized in the next step without further purification. ^1H NMR (DMSO- d_6) δ 1.52 (m, 4H), 2.98 (s, 2H), 3.49 (s, 3H), 4.66 (d, 2H), 12.17 (s, 1H).

3-(4-Methylpiperazine-1-carbonyl)-7-oxa-bicyclo[2,2,1]heptane -2-carboxylic acid methyl ester (5):



5

To a mixture of acid **3** (2.00 g, 10.0 mmole) in 50 mL of methylene chloride containing N-hydroxybenzotriazole (98.0 mg, 0.725 mmol) and EDC (2.09 g, 13.5 mmole) was added N-methylpiperazine (**4**, 1.45 g, 14.5 mmole) and the reaction mixture was stirred at room temperature overnight. The reaction mixture was evaporated to dryness and the product purified by column using 5% methanol in methylene chloride to give the required ester **5** as a semi solid (1.89 g, 67%). This was further purified by triturating with isopropyl ether followed by re-crystallization with a mixture of ethyl acetate /Hexane to give a white crystalline material of **5** (LB-107) (1.10 g, yield: 39%, mp 108-109°C). The mother liquor was concentrated and saved for future recrystallization. ^1H NMR ($CDCl_3$) δ 1.50 (m, 2H), 1.83 (m, 2H), 2.30-2.44 (m, 7H), 2.94 (d, $J=9.6$ Hz, 1H), 3.10 (d, $J=9.6$ Hz, 1H), 3.50 (m, 3H), 3.71 (m, 4H), 4.90 (m, 2H), ESMS: 282.

20

Example 2. Protein Phosphatase 2A Inhibitors


The compounds used in the method of the present invention are protein phosphatase 2A (PP2A) inhibitors (Lu et al., 2009; US 7,998,957 B2). Compounds LB-100 and LB-102 are inhibitors of PP2A in vitro in human cancer cells and in xenografts of human tumor cells in mice when given parenterally in mice. These compounds inhibit the growth of cancer cells in mouse model systems. It has also been shown that another structural homolog of these compounds, LB-107, is active when given orally to mice.

30


45

LB100, LB102 or LB107 are tested in an animal model of cardiac ischemia-reperfusion injury.


The structure of LB100 is:



The structure of LB102 is:



5 The structure of LB107 is:



Example 3. Increase Phosphorylation and Activation of Akt

Compounds LB-100, LB-102, LB-107, and other homologs of LB-100 disclosed herein increase phosphorylation of Akt in mammalian
 10 cells, including, but not limited to, cardiac cells, brain cells and endothelial cells. Compounds LB-100, LB-102 and LB-107 and other homologs of LB-100 disclosed herein reduce dephosphorylation and inactivation of Akt by protein phosphatase 2A (PP2A) in mammalian cells, including, but not limited to, cardiac cells,
 15 brain cells and endothelial cells. Compounds LB-100, LB-102 and LB-107 and other homologs of LB-100 disclosed herein increase activation of Akt by protein phosphatase 2A (PP2A) in mammalian cells, including, but not limited to, cardiac cells, brain cells and endothelial cells

20

Example 4. Reperfusion Injury

Compounds LB-100, LB-102, LB-107 and other homologs of LB-100 disclosed herein reduce reperfusion injury in mammalian tissue that has suffered from an ischemia. The mammalian tissue includes, but

is not limited to, cardiac tissue, brain tissue and endothelial tissue.

Example 5. Myocardial Infarction

5 Compounds LB-100, LB-102, LB-107 and other homologs of LB-100 disclosed herein reduce tissue damage associated with reperfusion injury in the heart of a subject following a myocardial infarction. Compounds LB-100, LB-102, LB-107 and other homologs of LB-100 disclosed herein prevent tissue damage associated with reperfusion
10 injury in the heart of a subject following a myocardial infarction.

Example 5. Sepsis

Compounds LB-100, LB-102, LB-107 and other homologs of LB-100 disclosed herein reduce vascular leakage associated with
15 reperfusion injury in a subject suffering from sepsis. Compounds LB-100, LB-102, LB-107 and other homologs of LB-100 disclosed herein reduce endothelial dysfunction associated with reperfusion injury in a subject suffering from sepsis.

20 **Example 6. Study of LB-100 to improve vascular integrity and reduce tissue damage following acute trauma to tissue**

The ability of LB-100 to improve vascular integrity and to reduce tissue damage following acute trauma to tissue was tested by
25 analyzing the survival of tissue at the distal end of a flap graft in in-bred rats infused with LB-100 via an implanted pump prior to raising a skin flap.

An ALZET pump (model 1007D, reservoir volume 100ul) was implanted
30 subcutaneously under anesthesia subcutaneously four days before creating a skin flap again under anesthesia. The pump in control animals contained sodium chloride and the pump in treated animals contained 0.55 mg of LB-100. The pump delivers 0.5 ul/hour +/- 10% so each animal received about 12 ul/day for 8 days, 4 days prior to
35 raising the graft and for the first 4 days after creation of the graft. There were 10 animals in each group.

The animals were sacrificed on day 7 following creation of the

flap, the amount of necrosis at the end of the flap was measured by planimetry. The area of necrosis at the end of the flap was divided by total surface area of the entire flap and expressed as a percentage. Table 1 shows the area (%) of necrosis in control rats
 5 1-10 vs rats receiving LB-100 (treatment rats).

Table 1. Area of necrosis in control rats vs rats receiving LB-100.

Control	Total	Necrosis	%	Treatment	Total	Necrosis	%
1	245751	139686	56.8404605	1	180926	55473	30.6606016
2	222271	94119	42.3442554	2	188880	63777	33.7658831
3	273475	142373	52.0607002	3	254862	66144	25.9528686
4	293026	87832	29.974132	4	154749	37181	24.0266496
5	265486	124567	46.9203649	5	227395	77089	33.9009213
6	227431	73396	32.2717659	6	200725	64147	31.9576535
7	269259	118707	44.0865486	7	231616	66886	28.8779704
8	301765	124054	41.1094726	8	251903	77327	30.6971334
9	223558	78105	34.9372422	9	231841	91515	39.4731734
10	248844	74858	30.0823006	10	277689	112275	40.431922
Average			41.0627243	Average			31.9744777

Analysis of the planimetry results revealed that the mean decrease
 10 in extent of necrosis in treated animals was 22%.

The visual appearance of the flaps was also inspected. Figures 1a-j show control rats 1-10 on day 7 and Figures 2a-j show treatment rats 1-10 on day 7. The treatment rats had received LB-100.

15

As shown in the photographs of the flaps on day 7 (Figures 1a-j and 2a-j), there was less extensive necrosis at the distal ends of the flap in those animals receiving LB-100.

20 Discussion

It has been known since the early 1970s (Maroko et al, 1971) that brief episodes of oxygen deprivation (ischemia) and relief of the ischemia (reperfusion) protects the heart against damage from a subsequent episode. In 1986, Murray et al showed that several short
 25 periods of cardiac ischemia reduced the size of tissue damage (infarct) following prolonged ischemia in a canine model. In 2003, Zhao et al. demonstrated that several rounds of ischemia and the return of blood flow (reperfusion) following the induction of an experimental heart attack, also reduced the size of the infarct,

compared to simple reperfusion after the attack. The mechanism(s) responsible for cardiac protection by pre- or post- brief cycles of ischemia has been the subject of more than 2500 papers and, although the phenomenon is demonstrable in several animal models, remains unclear. Despite a lack of understanding of the molecular basis of pharmacologic cardioprotection, however, recently there have been increasing calls for finding a way to reduce the amount of cardiac tissue damage following a heart attack (Cohen and Downey, 2011).

10

In 1998, Weinbrenner et al. reported that fostriecin (FOS), an inhibitor of protein phosphatase 2A, minimizes the size of infarction of cardiac tissue (myocardium) before and even when given after the onset of oxygen deprivation. The following year, Barancik et al. (1999) showed that another known inhibitor of serine/threonine phosphatase, okadaic acid, protected pig myocardium against ischemia (reduced the size of infarcted tissue) in an *in vivo* model and also showed that the activity of phosphatases, especially that of PP2A, were decreased in the heart tissue infused with the inhibitor.

20

FOS, a well studied inhibitor of PP2A, reduces injury even when given after the onset of coronary artery restriction and confirmation of this phenomenon by Barancik et al. (1999) using another PP2A inhibitor, okadaic acid in a pig heart model. Weinbrenner et al. (1998) studied a rabbit and pointed out that death of tissue in the rabbit heart progresses about 5 times faster than in the primate heart, leading them to speculate that administration of a PP2A inhibitor such as FOS as late as 50 minutes after the start of heart attack signs in humans might be expected to offer protection. In earlier studies, Armstrong et al (Armstrong et al. 1997; Armstrong et al. 1998) demonstrated that protein phosphatase inhibitor calyculin A, a PP1/2A inhibitor and FOS both afforded protection from injury to rabbit and pig cells in vitro. Subsequently, Fenton et al. (2005) in rats showed that okadaic acid reduced cellular death following induced MI in both young and aged animals. Fan et al. (2010) studied the effects of okadaic acid at concentrations inhibiting only PP2A and the effects

35

of cantharadin, a naturally occurring inhibitor of both PP1 and PP2A, in isolated perfused functioning rat hearts and concluded that the use of phosphatase inhibitors may provide an approach to reducing reperfusion-induced cell death.

5

The mechanism by which PP2A confers protection of myocardial damage due to oxygen deprivation is believed to be via activation of a major cell signaling pathway, the PI3K-Akt pathway (Matsui et al. 2001). Recently, Kunuthur (Kunuthur et al. 2011) showed that of
10 three Akt isoforms, Akt1 is essential for cardioprotection against ischemic-reperfusion injury.

Inhibition of PP2A by the novel inhibitors LB-100 and LB-102 and other structural homologs of these compounds have been shown to
15 result in increased phosphorylation of Akt (Lu et al. 2009; US Patent No. 8,085,268). Phosphorylation of Akt leads to its activation, which in turn increases the phosphorylation of several proteins affecting mitochondrial function and mediating cell death (Tsang et al. 2005).

20

Without wishing to be bound by any scientific theory, Akt mediates protection by phosphorylation of a number of target proteins, including GSK-3 β , endothelial nitric oxide synthase (eNOS), the proapoptotic Bcl-2 family member BAD, caspase 9, the ubiquitin
25 ligase murine double minute 2 (mdm2), and others (Fayard, E. et al, 2005). Phosphorylation of BAD suppresses apoptosis and promotes cell survival (Datta, S.R. et al. 1997). Overexpression of Akt blocks hypoxia-mediated activation of caspase 9, also blocking their proapoptotic roles (Uchiyama, T. et al. 2004). Akt
30 phosphorylates and activates the ubiquitin ligase, mdm2, which has been shown to play a role in reducing hypoxia-reoxygenation cell death in myocytes (Toth, A. 2006). Akt phosphorylates and activates eNOS, resulting in an increase in nitric oxide (NO) production, which may activate a number of pathways resulting in
35 cardioprotection (Tong, H. et al.). Akt also phosphorylates and inactivates GSK-3 β , which also provides an anti-apoptotic effect (Tong et al. 2000).

LB-100, LB-102, and LB-107 and other structural homologs are effective inhibitors of PP2A. LB-102, for example, inhibits PP2A with IC50 of about 0.4uM and to much lesser extent PP1 with an IC50 of about 8.0 uM (Lu et al. 2009(a)). LB-100 is currently entering
5 a Phase I clinical trial in which the compound is anticipated to enhance the cytotoxicity of DNA damaging agents. Although the maximum tolerated dose in humans has not yet been determined, toxicokinetic studies in rats and dogs indicate that the compound can be given with acceptable and reversible toxicity at doses known
10 to inhibit PP2A in tissue of rodents. Thus, LB-100, LB-102 and/or structural homologs can be safely administered to human beings to minimize the extent of myocardial damage following MI. Administration of PP2A is also expected to reduce the extent of tissue damage caused by ischemia in other disease states, such as
15 stroke and acute tissue injury due to trauma, either accidental or surgical injury that compromises blood supply to tissue acutely.

Tissue ischemia also results from hypotension secondary to acute bacterial infections. PP2A is activated due to the inflammatory
20 processes, particularly sepsis, and treatment with LB-100 and structural homologs may be highly beneficial, indeed lifesaving. Sepsis leads to activation of NADPH oxidase and uncoupling of endothelial nitric oxide synthase to produce superoxide, increased NO production and neuronal NOS activity, with increased 3-
25 nitrotyrosine formation and increase PP2A activity in the hind limbs of animals (Zhou et al. 2012). Zhou demonstrated that rapid injection of ascorbate protected against these effects including reduction of PP2A activation. This same mechanism was found by Ladurner (Ladurner et al. 2012). Ladurner showed that in addition
30 to ascorbate, the protein phosphatase inhibitor okadaic acid had similar effects supporting the hypothesis that endothelial damage in these model systems is mediated by PP2A. Han (Han et al. 2010) had previously demonstrated that okadaic acid at a concentration that inhibits PP2A activity decreases endothelial barrier
35 disruption caused by septic insult. Wu and Wilson (Wu et al. 2009) studied mouse skeletal muscle endothelial cells and showed that PP2A inhibition by okadaic acid preserved endothelial barrier function. Thus, LB-100 is expected to be useful in the treatment

of conditions such as septic shock in which PP2A mediates vascular leakage.

One mechanism by which fostriecin and the LB-100 compounds reduce
5 tissue damage following acute trauma to tissue may be by inhibition
of PP2A in the vasculature. The ability of LB-100 to improve
vascular integrity and to reduce tissue damage following acute
trauma to tissue was tested by analyzing survival of tissue at the
distal end of a flap graft as described in Example 6. The data from
10 Example 6 show that the mean decrease in extent of necrosis in
treated animals was 22%. Such an improvement in graft survival
would be valuable clinically where necrosis of the distal end of a
flap graft is a major limitation to this surgical intervention to
correct tissue defects.

References


- Armstrong SC et al (1997) Comparison of in vitro pre-conditioning responses of isolated pig and rabbit cardiomyocytes: effects of a protein phosphatase inhibitor fostriecin. J Mol Cell Cardiol 29:3009-3024.
- Armstrong SC et al (1998) Protein phosphatase inhibitors calyculin A and fostriecin protect rabbit cardiomyocytes in late ischemia. J Mol Cell Cardiol 30:61-73.
- Barancik M et al (1999) Okadaic acid and anisomycin are protective and stimulate the SAPK/JNK pathway. Journal of Cardiovascular Pharmacology 34(2):182-190.
- Cohen MV and Downey JM (2011) Is it time to translate ischemic preconditioning's mechanism of cardioprotection into clinical practice? J of Cardiovascular Pharmacology and Therapeutics 16(3-4):273-280.
- Datta, S.R. (1997) Akt Phosphorylation of BAD Couples Survival Signals to the Cell-Intrinsic Death Machinery. Cell, 91, 231-241.
- Fan WJ (2010) Kinases and phosphatases in ischaemic preconditioning: a re-evaluation. Basic Res Cardiol 105:495-511.
- Fenton et al (2005). Inhibition of phosphatase activity enhances pre-conditioning and limits cell death in the ischemic/reperfused aged rat heart. Life Sciences 77:3375-3388.
- Fayard E, Tintignac LA, Baudry A, Hemmings BA. (2005) Protein kinase B/Akt at a glance. J Cell Sci 118: 5675-5678.
- Kunuthur SP et al. (2011) The Akt1 isoform is an essential mediator of ischemic preconditioning. Journal of Cellular and Molecular Medicine. "Accepted Article"; doi:10.1111/j.1582-4934.2011.01491.x

- Ladurner et al. (2012) Ascorbate stimulates endothelial nitric oxide synthase enzyme activity by rapid modulation of its phosphorylation status. Free Radical Biology and Medicine, In Press.
- 5 Longacre LS et al (2011) New horizons in cardio protection. Circulation 124:1172-1179.
- Lu J et al (2009) The effect of a PP2A inhibitor on the nuclear
10 receptor corepressor pathway in glioma. Journal of Neurosurgery
DOI:10.3171/2009.11 JNS091272.
- Lu J et al (2009)(a) Inhibition of serine/threonine phosphatase
PP2A enhances cancer chemotherapy by blocking DNA damage induced
15 defense mechanisms. PNAS 106(28):11697-11702.
- Maroko PR et al (1971) Factors influencing infarct size following
experimental coronary artery occlusions. Circulation 43:67-82.
- 20 Matsui T et al (2001). Akt activation preserves cardiac function
and prevents injury after transient cardiac ischemia in vivo.
Circulation 104:330-335.
- USPTO 8,058,268 B2. Neuroprotective agents for the prevention and
25 treatment of neurodegenerative diseases.
- Tsang A et al. (2005) Myocardial postconditioning: reperfusion
injury revisited. American Journal of Physiology Heart Circulation
Physiology 289: H2-H7.
- 30 Tong H, Chen W, Steenbergen C, Murphy E. (2000) Ischemic
preconditioning activates phosphatidylinositol-3-kinase upstream of
protein kinase C. Circ Res 87: 309-315.
- 35 Toth A, Nickson P, Qin LL, Erhardt P. (2006) Differential
regulation of cardiomyocyte survival and hypertrophy by MDM2, an E3
ubiquitin ligase. J Biol Chem 281: 3679-3689.

- Uchiyama T, Engelman RM, Maulik N, Das DK. (2004) Role of Akt signaling in mitochondrial survival pathway triggered by hypoxic preconditioning. *Circulation* 109: 3042-3049.
- 5 Weinbrenner C et al (1998) Fostriecin, an inhibitor of phosphatase 2A, limits myocardial infarct size even when administered after onset of ischemia. *Circulation* 98:899-905.
- Wu, et al. (2007) Activation of Protein Phosphatase 2A by Palmitate
10 Inhibits AMP-activated Protein Kinase. *The Journal of Biological Chemistry*. 282, 13, 9777-9788.
- Yellon DM and Hausenloy DJ (2005) Realizing the clinical potential of ischemic preconditioning and postconditioning. *Nature Clinical*
15 *Practice Cardiovascular Medicine* 2(11):568-575.
- Zhou, G. et al. (2012) Ascorbate protects against vascular leakage in cecal ligation and puncture-induced septic peritonitis. *American J Physiol Regul Integr Comp Physiol* 302:409-416.
20
- Zhao Z-Q et al (2003) Inhibition of myocardial injury by ischemic postconditioning during reperfusion: comparison with preconditioning. *American Journal of Physiology* 285(2):H579-H588.

What is claimed is:

1. A method of reducing reperfusion injury in mammalian tissue comprising contacting the tissue with a protein phosphatase 2A (PP2A) inhibitor having the structure:



wherein

bond α is present or absent;


R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 ,

$O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2 - C_{12} alkyl,

alkenyl, C_4 - C_{12} alkenyl, alkynyl, aryl, substituted aryl

where the substituent is other than chloro when R_1 and R_2 are $=O$,



$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,

wherein each R_{11} is independently alkyl, alkenyl or alkynyl, or H;

R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken together are $=\text{O}$;

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,

where R_{12} is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

2. The method of claim 1, wherein the reduction of reperfusion injury comprises increased phosphorylation of Akt in the mammalian tissue that has suffered an ischemia.

3. The method of claim 2, wherein the reduction of reperfusion injury comprises increased activation of Akt in the mammalian tissue that has suffered an ischemia.


4. The method of claims 2 or 3, wherein the reduction of reperfusion injury comprises increased phosphorylation of BAD, mdm2, eNOS and/or GSK-3 β in the mammalian tissue that has suffered an ischemia.

5. The method of any of claims 2-4, wherein the ischemia is caused by a myocardial infarction, stroke or sepsis.

6. The method of any of claims 1-5, wherein the tissue is myocardial tissue, brain tissue or endothelial tissue.

7. The method of any of claims 1-6, wherein the protein phosphatase 2A inhibitor has the structure

57



wherein


bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , SH, S^- , SR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl,

C_2 - C_{12} alkyl, alkenyl, C_4 - C_{12} alkenyl, alkynyl, aryl,

substituted aryl where the substituent is other

than chloro when R_1 and R_2 are $=O$,



$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

where each R_{11} is independently alkyl, alkenyl, alkynyl, or H;

R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken together are $=O$;

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,


58

where R_{12} is H, aryl or a substituted or unsubstituted alkyl, alkenyl or alkynyl; and


each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.


8. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure



9. The method of claim 8, wherein the protein phosphatase 2A inhibitor has the structure



10. The method of claim 8, wherein the protein phosphatase 2A inhibitor has the structure



11. The method of claim 7 or 8, wherein bond α is present.

12. The method of claim 7 or 8, wherein bond α is absent.


13. The method of claim 11 or 12,
wherein

R_1 and R_2 together are =O;

R_3 is O^- or OR_9 ,

where R_9 is H, methyl, ethyl or phenyl;

R_4 is



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl,
substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12}
alkenyl, alkynyl, substituted alkynyl, aryl,
substituted aryl where the substituent is other
than chloro,



$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,

where R_{11} is alkyl, alkenyl or alkynyl, each
of which is substituted or unsubstituted, or
H;

R_5 and R_6 taken together are =O; and

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or
 SR_{12} ,


where R_{12} is a substituted or unsubstituted alkyl, alkenyl
or alkynyl.

14. The method of any of claims 7-13, wherein R_3 is O^- .

15. The method of any of claims 7-14,


60

wherein R_4 is




where X is O , NR_{10} , $N^+R_{10}R_{10}$

where each R_{10} is independently H , alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro when R_1 and R_2 are $=O$,




$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,
where R_{11} is H or alkyl.


16. The method of claim 15, wherein the protein phosphatase inhibitor 2A has the structure




61



17. The method of any of claims 7-15,




where R_{10} is R_{10} H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro when R_1 and R_2 are =O,




$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,
where R_{11} is H or alkyl.

18. The method of any of claims 7-15 or 17,


62




19. The method of any of claims 7-15 or 17,




20. The method of any of claims 7-15 or 17,



21. The method of any of claims 7-15 or 17,



22. The method of any of claims 7-15 or 17,




23. The method of any one of claims 7-15 or 17-22,
wherein R_5 and R_6 together are =O.

24. The method of any one of claims 7-15 or 17-23,
wherein R_7 and R_8 are each H.

25. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure


63



wherein bond α is present or absent;

R_9 is present or absent and when present is H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl or phenyl; and X is O, S, NR_{10} or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,




$-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-CH_2CN$, or $-CH_2CH_2R_{16}$,

where R_{11} is H or alkyl, and where R_{16} is any substituent that is a precursor to an aziridinyl intermediate,

or a salt, zwitterion or enantiomer of the compound.

26. The method of claim 25, wherein the protein phosphatase 2A inhibitor has the structure




wherein

bond α is present or absent;

X is O, S, NR_{10} or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,



$-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-CH_2CN$, or $-CH_2CH_2R_{16}$,


where R_{11} is H or alkyl, and where R_{16} is any substituent that is a aziridinyll intermediate,

or a salt, zwitterion or enantiomer of a compound.

27. The method of claim 26, wherein

X is O or NH^+R_{10} ,

where R_{10} is H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,



28. The method of claim 26,

wherein X is $-CH_2CH_2R_{16}$,

where R_{16} is any substituent that is a precursor to an aziridinyll intermediate.


29. The method of claim 26 or 27, wherein X is O.

30. The method of claim 26 or 27,

wherein X is NH^+R_{10} ,


where R_{10} H, alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,

65




31. The method of claim 26, 27 or 30, wherein R_{10} is methyl.

32. The method of claim 26, 27 or 30, wherein R_{10} is




33. The method of claim 26, 27 or 30, wherein R_{10} is



34. The method of claim 26, 27 or 30, wherein R_{10} is ethyl.

35. The method of claim 26, 27 or 30, wherein R_{10} is absent.

36. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure



wherein

bond α is present or absent;


R_9 is present or absent and when present is H, alkyl, alkenyl, alkynyl or phenyl; and

X is O, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, substituted C_2-C_{12} alkyl, alkenyl, substituted C_4-C_{12} alkenyl, alkynyl,

66

substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{12}$, or $-\text{CH}_2\text{COR}_{12}$,

where R_{12} is H or alkyl,

or a salt, zwitterion, or enantiomer of the compound.

37. The method of claim 36, where in the protein phosphatase 2A inhibitor has the structure




wherein

bond α is present or absent;

X is O or NH^+R_{10} ,

where R_{10} is H, alkyl, substituted $\text{C}_2\text{-C}_{12}$ alkyl, alkenyl, substituted $\text{C}_4\text{-C}_{12}$ alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{12}$, or $-\text{CH}_2\text{COR}_{12}$,

where R_{12} is H or alkyl.



38. The method of claim 36, wherein bond α is present.

39. The method of claim 36, wherein bond α is absent.



40. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure




68




41. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure



69



42. The method of claim 35, wherein the protein phosphatase 2A inhibitor has the structure




wherein

bond α is present or absent; X is NH^+R_{10} ,

70


where R_{10} is present or absent and when present R_{10} is is alkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl,




$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2R_{12}$, or $-\text{CH}_2\text{COR}_{12}$,

where R_{12} is H or alkyl.

43. The method of claim 7, wherein the protein phosphatase 2A inhibitor has the structure



44. The method of any of claims 7, wherein the protein phosphatase 2A inhibitor has the structure



wherein

bond α is present or absent;


R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is $O(\text{CH}_2)_{1-6}R_9$ or OR_9 , or

71



where X is O, S, NR₁₀, or N⁺R₁₀R₁₀,


where each R₁₀ is independently H, alkyl, hydroxyalkyl, C₂-C₁₂ alkyl, alkenyl, C₄-C₁₂ alkenyl, alkynyl, aryl, where the substituent is other than chloro when R₁ and R₂ are =O,



-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,

where each R₁₁ is independently H, alkyl, alkenyl, or alkynyl;

or R₃ and R₄ are each different and each is OH or



R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;

R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,


where R₁₂ is H, aryl or a substituted or unsubstituted alkyl, alkenyl or alkynyl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.


72

45. The method of claim 44, wherein the protein phosphatase 2A inhibitor has the structure




46. The method of claim 44 or 45, wherein bond α is present.
47. The method of claim 44 or 45, wherein bond α is absent.
48. The method of any of claims 44-47,
 R_3 is OR_9 or $O(CH_2)_{1-6}R_9$,
 where R_9 is aryl, substituted ethyl or substituted phenyl, wherein the substituent is in the para position of the phenyl;

R_4 is



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,


where each R_{10} is independently H, alkyl, hydroxyalkyl, substituted C_2 - C_{12} alkyl, alkenyl, substituted C_4 - C_{12} alkenyl, alkynyl, substituted alkynyl, aryl, substituted aryl where the substituent is other than chloro,



$-CH_2CN$, $-CH_2CO_2R_{11}$, $-CH_2COR_{11}$, $-NHR_{11}$ or $-NH^+(R_{11})_2$,


73

where R_{11} is alkyl, alkenyl or alkynyl, each of which is substituted or unsubstituted, or H;




or where R_3 is OH and R_4 is

49. The method of any of claims 44-48, wherein R_4 is



where R_{10} is alkyl or hydroxylalkyl



or R_4 is

when R_3 is OH.

50. The method of any of claims 44-49,


wherein

R_1 and R_2 together are =O;


R_3 is OR_9 or $O(CH_2)_{1-2}R_9$,

where R_9 is aryl, substituted ethyl, or substituted phenyl, wherein the substituent is in the para position of the phenyl;

or R_3 is OH and R_4 is



R_4 is



74

where R_{10} is alkyl or hydroxyl alkyl;
 R_5 and R_6 together are =O; and
 R_7 and R_8 are each independently H.


51. The method of any of claims 44-50,

wherein


R_1 and R_2 together are =O;

R_3 is OH, $O(CH_2)R_9$, or OR_9 ,

where R_9 is phenyl or CH_2CCl_3 ,



R_4 is



where R_{10} is CH_3 or CH_3CH_2OH ;

R_5 and R_6 together are =O; and
 R_7 and R_8 are each independently H.

52. The method of any of claims 44-51,

wherein R_3 is OR_9

where R_9 is $(CH_2)_{1-6}(CHNHBOC)CO_2H$, $(CH_2)_{1-6}(CHNH_2)CO_2H$, or
 $(CH_2)_{1-6}CCl_3$.

53. The method of claim 52,


wherein R_9 is $CH_2(CHNHBOC)CO_2H$, $CH_2(CHNH_2)CO_2H$, or CH_2CCl_3 .

75


54. The method of any of claims 44-52,
wherein R_3 is $O(CH_2)_{1-6}R_9$ or $O(CH_2)R_9$,
where R_9 is phenyl.

55. The method of any of claims 44-52, wherein R_3 is $O(CH_2)R_9$
where R_9 is phenyl.

56. The method of any of claims 44-52, wherein R_3 is OH and R_4 is




57. The method of any of claims 44-65, wherein R_4 is





wherein R_{10} is alkyl or hydroxyalkyl.

58. The method of claim 59, wherein R_{11} is $-CH_2CH_2OH$ or $-CH_3$.


59. The method of claim 44, wherein the protein phosphatase 2A inhibitor has the structure






76




, or




60. The method of claim 44, wherein the protein phosphatase 2A inhibitor has the structure



77



61. A method of reducing tissue damage associated with reperfusion injury in the heart of a subject following a myocardial infarction comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:



wherein

bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,


where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,

$O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,

78



where X is O, S, NR_{10} , or $\text{N}^+\text{R}_{10}\text{R}_{10}$,

where each R_{10} is independently H, alkyl, $\text{C}_2\text{--C}_{12}$ alkyl, alkenyl, $\text{C}_4\text{--C}_{12}$ alkenyl, alkynyl, aryl, substituted aryl where the substituent is other than chloro when R_1 and R_2 are =O,



$\text{---CH}_2\text{CN}$, $\text{---CH}_2\text{CO}_2\text{R}_{11}$, $\text{---CH}_2\text{COR}_{11}$, ---NHR_{11} or $\text{---NH}^+(\text{R}_{11})_2$,

wherein each R_{11} is independently alkyl, alkenyl or alkynyl, or H;

R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken together are =O;

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,


where R_{12} is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

79

62. A method of reducing vascular leakage associated with reperfusion injury in a subject suffering from sepsis comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:



wherein

bond α is present or absent;


R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are $=O$;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , OR_{10} ,

$O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12}

alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl,

substituted aryl where the substituent is other

than chloro when R_1 and R_2 are $=O$,



80

$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,
 wherein each R_{11} is independently alkyl,
 alkenyl or alkynyl, or H;

R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken
 together are =O;


R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or
 SR_{12} ,

where R_{12} is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or
 unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound.

63. A method of reducing tissue damage due to an acute trauma in a
 subject, comprising administering to the subject a therapeutically
 effective amount of a protein phosphatase 2A (PP2A) inhibitor
 having the structure:



wherein

bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,


where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are =O;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 ,

$\text{O}(\text{CH}_2)_{1-6}\text{R}_9$, SH, S^- , SR_9 ,

81



where X is O, S, NR₁₀, or N⁺R₁₀R₁₀,

where each R₁₀ is independently H, alkyl, C₂-C₁₂ alkyl, alkenyl, C₄-C₁₂ alkenyl, alkynyl, aryl, substituted aryl where the substituent is other than chloro when R₁ and R₂ are =O,



-CH₂CN, -CH₂CO₂R₁₁, -CH₂COR₁₁, -NHR₁₁ or -NH⁺(R₁₁)₂,

wherein each R₁₁ is independently alkyl, alkenyl or alkynyl, or H;

R₅ and R₆ is each independently H, OH, or R₅ and R₆ taken together are =O;

R₇ and R₈ is each independently H, F, Cl, Br, SO₂Ph, CO₂CH₃, or SR₁₂,

where R₁₂ is H, alkyl, alkenyl, alkynyl or aryl; and


each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound,

so as to thereby reduce tissue damage due to the acute trauma in the subject.

82

64. A method of reducing vascular leakage due to an acute trauma in a subject, comprising administering to the subject a therapeutically effective amount of a protein phosphatase 2A (PP2A) inhibitor having the structure:



wherein


bond α is present or absent;

R_1 and R_2 is each independently H, O^- or OR_9 ,

where R_9 is H, alkyl, alkenyl, alkynyl or aryl,

or R_1 and R_2 together are =O;

R_3 and R_4 are each different, and each is OH, O^- , OR_9 , $O(CH_2)_{1-6}R_9$, SH, S^- , SR_9 ,



where X is O, S, NR_{10} , or $N^+R_{10}R_{10}$,

where each R_{10} is independently H, alkyl, C_2-C_{12} alkyl, alkenyl, C_4-C_{12} alkenyl, alkynyl, aryl, substituted aryl where the substituent is other than chloro when R_1 and R_2 are =O,



$-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{CO}_2\text{R}_{11}$, $-\text{CH}_2\text{COR}_{11}$, $-\text{NHR}_{11}$ or $-\text{NH}^+(\text{R}_{11})_2$,

wherein each R_{11} is independently alkyl, alkenyl or alkynyl, or H;

R_5 and R_6 is each independently H, OH, or R_5 and R_6 taken together are =O;

R_7 and R_8 is each independently H, F, Cl, Br, SO_2Ph , CO_2CH_3 , or SR_{12} ,

where R_{12} is H, alkyl, alkenyl, alkynyl or aryl; and

each occurrence of alkyl, alkenyl, or alkynyl is branched or unbranched, unsubstituted or substituted,

or a salt, enantiomer or zwitterion of the compound,

so as to thereby reduce vascular leakage due to the acute trauma in the subject.


65. The method according to claim 63 or 64, wherein the acute trauma causes ischemia.

66. The method according to any one of claims 63-65, wherein the acute trauma is due to surgical injury.

67. The method according to claim 66, wherein the surgical injury is due to flap graft surgery.

Figure 1A-B.

A



B





Figure 1C-D.

C



D





Figure 1E-F.

E



F





Figure 1G-H.

G



H




Figure 1I-J.

I




J



Figure 2A-B.

A



B





Figure 2C-D.

C



D




Figure 2E-F.



E**F**

Figure 2G-H.

G




H



Figure 2I-J.

I



J



A

