



(12) **United States Patent**
Har-Noy

(10) **Patent No.:** **US 10,385,315 B2**
(45) **Date of Patent:** **Aug. 20, 2019**

(54) **CELLS EXPRESSING TH1 CHARACTERISTICS AND CYTOLYTIC PROPERTIES**

(71) Applicant: **Michael Har-Noy**, Modi'in (IL)

(72) Inventor: **Michael Har-Noy**, Modi'in (IL)

(73) Assignee: **IMMUNOVATIVE THERAPIES, LTD.**, Jerusalem (IL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/800,193**

(22) Filed: **Mar. 13, 2013**

(65) **Prior Publication Data**

US 2013/0195825 A1 Aug. 1, 2013

Related U.S. Application Data

(62) Division of application No. 13/214,534, filed on Aug. 22, 2011.

(60) Provisional application No. 61/401,881, filed on Aug. 20, 2010, provisional application No. 61/495,055, filed on Jun. 9, 2011.

(51) **Int. Cl.**

A61K 35/17 (2015.01)
C12N 5/0783 (2010.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 5/0636** (2013.01); **A61K 35/17** (2013.01); **C12N 5/0638** (2013.01); **A61K 2039/5158** (2013.01); **A61K 2039/57** (2013.01); **C12N 2501/51** (2013.01); **C12N 2501/515** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,402,431 B2 7/2008 Har-Noy
7,435,592 B2 10/2008 Har-Noy
7,722,863 B2 5/2010 Multhoff
2003/0175242 A1 9/2003 Gruenberg
2003/0194395 A1 10/2003 Gruenberg et al.

FOREIGN PATENT DOCUMENTS

JP 2007-500217 A 1/2007
JP 2013-536232 A 9/2013
WO 2003/034820 A1 5/2003

OTHER PUBLICATIONS

Janeway et al. "Summary to Chapter 9" in Immunobiology: The Immune System in Health and Disease. 5th edition. New York: Garland Science; 2001.*

Grossman et al. "Differential expression of granzymes A and B in human cytotoxic lymphocyte subsets and T regulatory cells", Blood 104(9): 2840-2848, 2004.*

Koeffler et al. "Human myeloid leukemia cell lines: a review", Blood 56(3): 344-350, 1980.*

BD Pharmingen "Purified NA/LE mouse anti-human CD3", technical data sheet 555336 Rev. 16, 2008.*

BD Pharmingen "Purified NA/LE hamster anti-mouse CD28", technical data sheet 553294 Rev. 17, 2011.*

ATCC "A20 Product Sheet", company's technical data sheet, copyright 2013.*

Paludan et al. "Epstein-Barr nuclear antigen 1-specific CD4+ Th1 cells kill Burkitt's lymphoma cells.", The Journal of Immunology 169 (3): 1593-1603, 2002 (Year: 2002).*

Ozaki et al. "Humanized anti-HM1. 24 antibody mediates myeloma cell cytotoxicity that is enhanced by cytokine stimulation of effector cells", Blood 93(11): 3922-3930, 1999 (Year: 1999).*

Barrett, J. et al. (2000). "The benefits of an alloresponse: graft-versus-tumor." Journal of Hematotherapy Stem Cell Research, vol. 9, No. 3, pp. 347-354.

Bodey, B. et al. (2000). "Failure of cancer vaccines: the significant limitations of this approach to immunotherapy." Anticancer Research, vol. 20, No. 4, pp. 2665-2676.

Childs, R. W. (2000). "Nonmyeloablative allogeneic peripheral blood stem-cell transplantation as immunotherapy for malignant diseases." Cancer Journal, vol. 6, No. 3, pp. 179-187.

Har-Noy, M. et al. (2008). "Completely mismatched allogeneic CD3/CD28 cross-linked Th1 memory cells elicit anti-leukemia effects in unconditioned hosts without GVHD toxicity." Leukemia Research, vol. 32, No. 12, pp. 1903-1913.

Har-Noy, M. et al. (2009). "Allogeneic CD3/CD28 cross-linked Th1 memory cells provide potent adjuvant effects for active immunotherapy of leukemia/lymphoma." Leukemia Research, vol. 33, No. 4, pp. 525-538.

Har-Noy, M. et al. (2008). "The anti-tumor effect of allogeneic bone marrow/stem cell transplant without graft vs. host disease toxicity and without a matched donor requirement?" Medical Hypotheses, vol. 70, No. 6, pp. 1186-1192.

Kantoff, P. W., C. S. Higano, et al. "Sipuleucel-T immunotherapy for castration-resistant prostate cancer." The New England Journal of Medicine, vol. 363, No. 5, pp. 411-422.

Lee, S. J. et al. (2002). "Severity of chronic graft-versus-host disease: association with treatment-related mortality and relapse." Blood Journal, vol. 100, No. 2, pp. 406-414.

Hombach et al. (2006) "Human CD4+T Cells lyse target cells via granzyme/perforin upon circumvention of MHC Class II restriction by an antibody-like immunoreceptor" J. Immunol 15;177(8): 5668-5675.

(Continued)

Primary Examiner — Emily A Cordas

(74) *Attorney, Agent, or Firm* — Westman, Champlin & Koehler, P.A.; Z. Peter Sawicki; Visala Goswitz

(57)

ABSTRACT

A novel cell type has been generated that has both Th1 characteristics and cytolytic activity. These Th1/killer cells are CD4+ cells purified from peripheral blood and manipulated to have Th1 characteristics such as production of IFN-gamma combined with cytolytic activity similar to cytotoxic T-cells (CTL). The CTL activity is targeted toward diseased cells, not normal cells. The cytolytic activity of the Th1/killer cells is mediated by Granzyme B-Perforin mechanism and results in apoptotic death of diseased cells. Methods of producing and using these Th1/killer cells include isolating CD4+ cells from peripheral blood, activating the CD4+ T-cells to form Th1/killer cells and administering these Th1/killer cells with the cytolytic activity to a patient wherein the Th1/killer cells are allogeneic to the patient.

(56)

References Cited

OTHER PUBLICATIONS

Brown et al. (2009) "IL-2 and antigen dose differentially regulate perforin- and FasL-mediated cytolytic activity in antigen specific CD4+ T cells" *Cellular Immunology* 257:69-79.

Chinese Office Action dated Feb. 13, 2014; for CN201180040232.4. Extended European Search Report dated Feb. 19, 2014; EP 11818884.6.

Jursik et al. (2009) "Large-Scale Production and Characterization of Novel CD4+ Cytotoxic T Cells with Broad Tumor Specificity for Immunotherapy" *Mol Cancer Res* 7: 339-353.

Echchakir et al. (2000) "Cutaneous T Cell Lymphoma Reactive CD4+ Cytotoxic T Lymphocyte Clones Display a Th1 Cytokine Profile and Use a Fas-Independent Pathway for Specific Tumor Cell Lysis" *Journal of Investigative Dermatology* 115(1): 74-80.

Soghoian et al. (2010) "Cytolytic CD4+ T cells in viral immunity" *Expert Rev Vaccines* 9(12): 1453-1463.

Quezada et al. (2010) "Tumor-reactive CD4+ cells develop cytotoxic activity and eradicate large established melanoma after transfer into lymphopenic hosts" *J Exp Med* 207(3): 637-650.

Search Report & Written Opinion for Singapore Application No. 201301106-9, dated Jun. 25, 2014.

Brown D.M., Cytolytic CD4 Cells: Direct Mediators in Infections Disease and Malignancy. *Cell Immunol.*, Feb. 24, 2010, vol. 262, No. 2, pp. 89-95.

Yasukawa et al. "Granule exocytosis, and not the Fas/Fas ligand system, is the main pathway of cytotoxicity mediated by alloantigen-specific CD4+ as well as CD8+ cytotoxic T-lymphocytes in humans", *Blood* 95(7): 2352-5, 2000.

Traidl et al. "Disparate cytotoxic activity of nickel-specific CD8+ and CD4+ T cell subsets against keratinocytes" *Journal of Immunology* 165: 3058-3064, 2000.

ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). Mar. 12, 2009. Identifier—NCT00861107, In-Situ Therapeutic Cancer Vaccine for Metastatic Cancer Combining AlloStim With Tumor Cryoblation. [Nov. 26, 2015] Available from: <https://clinicaltrials.gov/ct2/show/NCT00861107>.

Li, J. M., C. R. Giver, et al. (2009). "Separating graft-versus-leukemia from graft-versus-host disease in allogeneic hematopoietic stem cell transplantation." *Immunotherapy*, vol. 1, No. 4, pp. 599-621.

Resnick, I. B., M. Y. Shapira, et al. (2005). "Nonmyeloablative stem cell transplantation and cell therapy for malignant and non-malignant diseases." *Transplant Immunology*, vol. 14, Nos. 3-4, pp. 207-219.

Office Action issued in corresponding Japanese Patent Application No. 2013-526060, dated Aug. 2, 2016.

Office Action issued for related Korean patent application No. 10-2013-7004186, dated Oct. 20, 2017.

Cai, S.F. et al., "Differential Expression of Granzyme B and C in murine Cytotoxic Lymphocytes" *The Journal of Immunology*, 2009, 182: 6287-6297.

Office Action issued for related Japanese patent application No. 2016-235386, dated Jul. 24, 2018.

Kemper et al., "Activation of human CD4+ cells with CD3 and CD46 induces a T-regulatory cell 1 phenotype" (2003) *Nature*, vol. 421, pp. 388-392.

Sharma et al. "Granzyme B, a new player in activation-induced cell death, is down regulated by vasoactive intestinal peptide in Th2 but not Th1 effectors" *Journal of Immunology* 176: 97-110, 2006 (Year:2006).

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4A

FIG. 4B

FIG. 4C




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7

1

CELLS EXPRESSING TH1 CHARACTERISTICS AND CYTOLYTIC PROPERTIES

CROSS-REFERENCE TO RELATED APPLICATION

The present application is a divisional of U.S. application Ser. No. 13/214,534, filed Aug. 22, 2011 which is based on and claims the benefit of U.S. provisional patent application Ser. No. 61/401,881, filed on Aug. 20, 2010 and U.S. provisional patent application Ser. No. 61/495,055, filed on Jun. 9, 2011; the contents of all which are hereby incorporated by reference in their entirety.

BACKGROUND

Therapeutic cancer vaccination is a type of immunotherapy. Immunotherapy is a new treatment modality that is emerging to join chemotherapy, radiation and surgery as a class of drugs for treatment of cancer. Immunotherapy methods seek to harness the power of the immune system in order to treat diseases. Therapeutic vaccination is a type of immunotherapy that is potentially curative therapy against existing tumors, viruses and bacterial pathogens. Therapeutic vaccines generally consist of an antigen source derived from the target disease and an adjuvant designed to enhance the desired immune response. In some therapeutic vaccine protocols, living immune cells, either derived from the host (autologous) or from a donor (allogeneic), are components of therapeutic vaccines. Autologous and allogeneic living immune cells, such as dendritic cells, NK cells and T-cells, are also used in immunotherapy protocols for the treatment of cancer.

Lymphocytes are part of the vertebrate immune system and include large granular lymphocytes and small lymphocytes. Large granular lymphocytes include natural killer cells (NK cells). Small lymphocytes consist of T-cells and B cells. NK cells are part of the innate immune system and play a major role in defending an animal from tumors and virally infected cells. NK cells can distinguish infected and tumor cells from uninfected cells through the multihisto-compatibility complex (MHC) class I surface molecules. NK cells are activated in response to cytokines and upon activation release cytotoxic granules such as perforin and granzyme B that destroy the infected cells or tumor cells. NK cells are characterized by the cell surface markers CD16+ and CD56+, but do not express CD4.

T-cells and B-cells are part of the adaptive immune response and recognize non-self antigens. B-cells respond to non-self antigen by antibody production. Different types of T-cells respond to the non-self antigens in different ways. Cytotoxic T-cells (CTL) are characterized by the markers CD3+, CD8+ and TCR $\alpha\beta$, and do not express CD4. Tumor-specific CD8+ CTLs are mainly responsible for tumor elimination. Moreover, CTLs can specifically recognize tumor cells and do not attack normal cells of the same tissue. CTLs produce toxic granules that contain powerful enzymes including granzyme B and perforin that induce death of diseased or cancerous cells.

T helper cells are characterized by the cell surface markers CD3+, CD4+ and TCR $\alpha\beta$. T-helper cells produce cytokines or other molecules that direct the immune response through other cells or molecules. T-helper cells are not known to directly mediate killing of cells and thus do not normally contain cytotoxic granules such as granzyme B or perforin. T-helper cells are sub-categorized into two types:

2

T-helper type 1 (Th1) and the T-helper type 2 (Th2). Th1 cells mediate cellular immunity and are critical for immune-mediated tumor eradication, whereas Th2 cells mediate a humoral or antibody immunity. Th1 and Th2 cells are counter-regulatory, increased Th1 cells down regulate Th2 cells and vice versa.

A variety of immunotherapy methods and compositions have been developed in order to enhance or suppress the immune response in patients. Cell therapy methods often involve ex-vivo manipulations such as proliferation, differentiation and/or activation of cells and the transfer of these cells to a patient as therapy.

Adoptive immunotherapy involves removing lymphocytes from the patient, boosting their anti-cancer activity ex-vivo, growing the cells to large clinically relevant numbers, and then returning the cells to the patient:

Initial experiments in adoptive immunotherapy involved removing lymphocytes from the blood of a patient and growing them in the presence of the lymphokine interleukin-2 (IL-2), an immune stimulator. The cells were then returned to the patient. These lymphocytes were called lymphokine-activated killer (LAK) cells.

A stronger response against tumor cells was obtained using lymphocytes isolated from the tumor itself. These tumor-infiltrating lymphocytes (TILs) are grown in the presence of IL-2 and returned to the body to attack the tumor.

Adoptive immunotherapy is a form of immunotherapy where ex vivo processed cells are introduced into the body. Pre-clinical studies suggested anti-tumor activity could be enhanced by using interleukin-2 together with ex vivo activated and expanded autologous lymphocytes. Also, the first objective responses with high-dose bolus interleukin-2 therapy were noted in patients receiving interleukin-2 together with LAK cells prepared through in vitro activation of autologous peripheral blood lymphocytes that were harvested by lymphopheresis, and initially, it appeared that the combination of interleukin-2/LAK was more active than interleukin-2 alone. IL-2 is a cytokine produced by Th1 helper cells. Interest in the use of adoptive immunotherapy declined after clinical studies failed to demonstrate that the addition of activated LAK or TIL cells with IL-2 is any more effective than IL-2 alone.

CD4+T (Th) cells are crucial for the activation and regulation of most aspects of the host defense against infections and for adequate function of cytotoxic CD8+ lymphocytes (CTL). Much of the research in tumor immunology has been focused on naturally occurring autologous tumor-specific T cells that are predominantly CD8+ and can be isolated from melanoma and some other tumors. These cells can be expanded in vitro and reinfused. This type of adoptive immunotherapy has resulted in objective tumor responses and long-term survival in some patients whose disease is refractory to other interventions. While there is an extensive clinical experience using CD8+ cytotoxic T cells as adoptive immunotherapy, poor clinical responses underscore the need to improve these therapies in order to achieve tumor rejection in higher percentages of patients.

CD8+ lymphocytes can fail to maintain functionality in vivo in large part because of the absence of CD4+ T cell help. In vitro, CD8+ cells release large quantities of interferon- γ (IFN- γ) upon exposure to MHC-compatible cell lines and lyse autologous antigen-positive and MHC class I-positive tumors. However, the genetic instability of tumor cells frequently leads to losses in the ability to process and present endogenous antigens rendering tumors inherently unreliable targets for CD8+ cytolytic T cells. Furthermore,

CD8+ T cells appear to lack the intrinsic ability to orchestrate a broad antitumor response that seems inherent in some CD4+ T cell subsets.

While immunotherapies with adoptive transfer of CTL have demonstrated promise in many animal models, the translation of these results to humans has proven to be difficult and elusive. CD4+ cells, especially of the Th1 subtype, would be an attractive addition to the available immunotherapy armamentarium to provide natural “help” to adoptively transferred CD8+ killer cells. However, there are significant barriers in working with naturally occurring tumor-specific CD4+ cells. The genetic diversity of the class II HLA associated with CD4+ helper cells in any given population of patients is much more complex than in the case of class I HLA found in CD8+ cytolytic (killer) T-cells, making identification of epitopes and TCRs more problematic. Moreover, CD4+ T-cells expand *in vitro* less well than CD8+ cells and the culture conditions significantly affect their characteristics. Finally, there is a paucity of realistic animal models based on tumor-specific CD4+ cells. These factors have limited translational research using CD4+ T cells and limited their clinical use.

This invention discloses a method for producing clinically-relevant numbers of CD4+ cells that also contain the cytolytic granules granzyme B and perforin and function both as Th1 cells (produce IFN-gamma and not IL-4) and have NK-like activity (recognize and kill tumor cells and not normal cells) for use in therapeutic cancer vaccine and adoptive immune cell therapy protocols.

SUMMARY

Cell-based therapeutic cancer vaccines (cellular immunotherapy) hold promise as a new minimally toxic approach for treatment of cancer. The promise of this approach has been demonstrated in animal models, but it has been difficult to translate this promise to the clinic (Bodey, Bodey et al. 2000). The recent FDA approval of the first cellular immunotherapy drug, PROVENGE® (sipuleucel-T), an autologous dendritic cell immunotherapy for the treatment of asymptomatic or minimally symptomatic metastatic castrate resistant prostate cancer (Kantoff, Higano et al.), has renewed enthusiasm to develop improved cell therapy products for clinical application.

Since it has proven difficult to successfully translate the immune-mediated anti-tumor mechanisms of cellular immunotherapy strategies developed in animals, we instead concentrated on translating an anti-tumor immune mechanism already known to elicit significant clinical anti-tumor effects. Arguably, the most powerful and most effective immune-mediated anti-tumor mechanism ever described in humans is the graft vs. tumor (GVT) effect that occurs after non-myeloablative conditioning and transplant of HLA-matched stem cells (Barrett and Childs 2000; Childs 2000; Resnick, Shapira et al. 2005). Unfortunately, these GVT effects are often accompanied by chronic graft-versus-host-disease (GVHD) toxicity. Chronic GVHD is associated with significant morbidity and is the leading cause for late mortality after allogeneic hematopoietic stem cell transplantation (Lee, Klein et al. 2002), significantly limiting the clinical application of the GVT mechanism.

Separation of the beneficial GVT effects from the detrimental GVHD effects is difficult because the effects are interrelated and proportional, in that manipulations that inhibit GVHD also decrease the GVT effect (Li, Giver et al. 2009). By analyzing the known immune-mediated mechanisms of the inter-related GVT and GVHD effects, it was

hypothesized that it might be possible to develop an allogeneic cellular immunotherapy that instead of separating the GVT/GVHD effects would maintain the interrelationship of the immune mechanisms by reversing the direction of the effects from graft-to-host to host-to-graft (Har-Noy and Slavin 2008). Successful reversal of the direction of the effects would result in a host vs. tumor (HVT) effect coupled to a non-toxic host vs. graft (HVG) rejection effect.

It was hypothesized that intentionally mis-matched, activated, allogeneic Th1 cells that expressed high density CD40L and produced high amounts of interferon (IFN)- γ would be capable of eliciting coupled HVT/HVG effects. This hypothesis was tested in an animal model of chemotherapy-refractory B-cell leukemia/lymphoma (Har-Noy, Zeira et al. 2008). These studies demonstrated that significant HVT/HVG activity could be elicited by ex-vivo differentiated allogeneic Th1 cells only when the Th1 cells were activated and administered attached to anti-CD3/anti-CD28-coated microbeads. Additional studies indicated that these allogeneic CD3/CD28-crosslinked Th1 cells had both immunomodulatory effects, capable of switching systemic dominant Th2 immunity to dominant Th1 immunity; and, adjuvant effects for promoting development of Th1-specific immunity and dysregulation of tumor immunoavoidance mechanisms (Har-Noy, Zeira et al. 2009). In view of these data, human version of Th1 allografts were investigated for the possibility of translating this experimental approach to the clinic.

There were significant regulatory and logistical concerns associated with the ex vivo differentiation and expansion of allogeneic Th1 cells that were necessary to be overcome in order to use these cells for human clinical investigations. One such issue was that the mouse version of allogeneic Th1 cells required the use of exogenous cytokines, IL-2, IL-12 and anti-IL-4 mAb to cause the ex-vivo differentiation of mouse Th1 cells. The current invention discloses the successful development of a GMP-compliant process for the production of human allogeneic Th1 cells without the use of exogenous cytokines resulting in consistent batch-to-batch identity and functional characteristics for classification as a human biological drug. Upon characterization of the resulting allograft cells, the human production process resulted in an unexpected finding. The allogeneic Th1 cells generated by the methods described herein not only developed Th1 characteristics, but also developed NK-like characteristics. An allograft with a combination of Th1 and NK-like activity is unique and is not described in normal blood circulation. The method of the current invention involves the ex-vivo culture of purified, resting (CD25⁻), CD4+ T-cells with mixed memory and naïve phenotype (CD45RA and CD45RO, respectively). The cells are activated multiple times with immobilized anti-CD3/anti-CD28 mAbs in the absence of exogenous cytokines. The cells expand 25-100-fold after 9-days and differentiate into CD4+, CD45RO+, CD25+, CD62^{hi} cells. This intermediate cell can be stored in liquid nitrogen for years. When needed for patient administration, the precursor cells are incubated for 4 h with CD3/CD28 mAb coated microbeads. During this final 4 h incubation, the precursor cells further differentiate to decrease expression of CD62L and increase expression of CD40L and CD25. The final CD4+, CD45RO+, CD40L^{hi}, CD62L^{lo}, CD25+ cells produce >1000 pg/10⁶ cells of IFN-gamma and <50 pg/10⁶ cells of IL-4 (Th1 phenotype) and express cytolytic granules of granzyme B and perforin and exhibit NK-like activity with the ability to lyse tumor cells and not normal cells.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A and FIG. 1B are flow cytometry plots from immunostaining of CD4 surface antigen demonstrating the cell phenotype before and after CD4+ selection, respectively.

FIG. 2A is a series of flow cytometry plots demonstrating the phenotype of source cells having as resting CD4 naïve (CD4, CD45RA) cells.

FIG. 2B is a series of flow cytometry plots demonstrating the change in phenotype after culture to activated CD4 memory cells (CD4, CD45RO and CD25).

FIG. 3 is a flow cytometry plot demonstrating the shift in CD62L expression from high to low in CAC before activation and CFB after activation.

FIGS. 4A-4C are plots of the CD40L expression pattern in CD4+ cells, CAC and CFB, respectively demonstrating increased expression of CD40L in CFB.

FIG. 5 is a graph of the direct killing assay and shows CFB can kill cells from the tumor cell line, ARH77.

FIG. 6A is a plot of expression of Granzyme B by CAC.

FIG. 6B is a plot of expression of Granzyme B by CFB.

FIG. 6C is a picture of a western blot showing detection of Perforin in CFB.

FIG. 7 is a graph of perforin-Granzyme B activity with and without EGTA.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

This disclosure relates to compositions that include cells with unique characteristics. The composition includes cells of a cell type having both Th1 helper cell properties and cytolytic granules and NK activity. CD4+ T-cells with the combination of Th1 characteristics and NK characteristics can provide substantial therapeutic capabilities for patients having a variety of diseases. Th1 characteristics can include, for example, production of IFN-gamma without production of IL-4. NK characteristics can include killing of tumor cells but not normal cells utilizing expression of Granzyme B and perforin.

The present invention includes methods to cause the differentiation and expansion of this novel cell type to clinically-relevant numbers for adoptive immunotherapy. Methods for deriving the novel cells can include isolating T-cells, specifically CD4+ T-cells from the peripheral blood or other sources of T-cells. These CD4+ T-cells can be differentiated and expanded according to the methods described herein to clinically-relevant numbers and used for immunotherapy. Surprisingly, the CD4+ T-cells prepared according to these methods can exhibit cytolytic activity similar to the cytolytic activity generally found in NK/CTL cells and also exhibit Th1 characteristics and activity.

Generally, T-helper cells do not directly exhibit cytolytic activity but produce cytokines and other factors that then elicit other cells such as NK cells and CTLs to inactivate or kill the diseased cells. The CD4+ T-cells in the present compositions can directly kill the diseased cells as well as produce cytokines to stimulate other cells such as NK cells that then carry out the cytolytic activity. The cytolytic activity of these CD4+ T-cells is mediated through Granzyme B and perforin resulting in the killing of the tumor or infected cells.

Cytolytic activity as referred to herein relates to direct killing by the effector cells of target tumor cells or cell lines, when they interact with the target cells. Direct killing can

occur by the presence of only the effector and the target cells and without the presence of any other cell types.

The novel cell type described in the present invention will be referred to herein as Th1/killer cells. The Th1/killer cells can exhibit a combination of Th1 characteristics, killing of tumor cells and not normal cells and expression of NK activity through a mechanism of Granzyme B and perforin expression.

Clinically-relevant number of cells as referred to herein relates to sufficient number of the Th1/killer cells to elicit an anti-tumor effect either directly or indirectly. The number of Th1/killer cells administered generally is at least 1×10^6 cells, preferably at least 1×10^7 number of cells, and more preferable at least about 1×10^8 number of cells in a single dose. Doses with greater amount of cells are also within the scope of the invention. Preferably sufficient amount of cells are produced from a single blood donor to produce enough doses for multiple patients. Clinically relevant numbers include enough cells for at least one patient, more preferably for up to 10 patients and most preferably for over 100 patients.

The compositions of the present invention can be administered to a patient having a variety of diseased cells. The patient can be animal, human or other mammals. The diseased cells can be cancerous cells or cells from infected tissue. Diseased cells may be infected with viruses and/or bacterial pathogens. The compositions may be administered to patients with solid tumors such as breast cancer, lung cancer, colon cancer, stomach cancer, pancreatic cancer and the like. The compositions may also be administered to patients with hematological malignancies (e.g., Chronic Lymphocytic Leukemia, Multiple Myeloma, and non-Hodgkin's lymphomas) or viral infectious diseases (e.g., hepatitis B or C, herpes, HIV) and other disorders.

The composition described herein is live cells. By live cells, it is meant that >70% of the cells are viable as determined by appropriate assay techniques such as trypan blue extrusion, MTT or bioluminescent detection of the ATP levels such that the cells are capable of ex vivo manipulations such as expansion, differentiation, and/or activation under appropriate conditions. The compositions, however, may include some inactivated cells, radiated cells and/or non-viable cells and non-living components.

The composition generally includes CD4+ cells first derived, for example, from blood of a donor. The donor can be without disease or normal. The CD4+ cells may be processed in a manner as described herein and then formulated for infusion into the same donor or into a related or unrelated recipient.

The CD4+ source cells are preferably purified from the peripheral blood. The CD4 cells can be purified in a variety of ways. Generally, the CD4+ cells are purified from the buffy coat of peripheral blood by positive selection. Positive selection of the CD4+ from buffy coat can be performed, for example, by using magnetic beads and columns obtained from Miltenyi Biotec (Auburn, Calif.). This can result in a cell composition that is greater than about 90 percent, preferably greater than about 95 percent, and more preferably greater than about 98 percent pure CD4+ cells.

The CD4+ cells isolated from the buffy coat and used as source material for production of Th1/killer cells can be characterized by expression of a number of cell surface markers. The population of CD4+ cells purified from the buffy coat of the peripheral blood can have a mixed population of CD45RA/CD45RO markers. The CD4+ T-cells are generally predominantly CD25-. Preferably the source CD4+ cells are mostly naïve (express CD45RA and not

CD45RO). Source CD4+ T-cells can also be evaluated for CD62L expression and CD40L expression. Generally, the source CD4+ T-cells express very low amounts of CD40L. Expression of CD62L can be a bimodal (high and low) mean fluorescent intensity (MFI) peak. Source CD4+ cells preferably have a high CD62L expression.

The source CD4+ T-cells can be processed to differentiate into Th1/killer cells by multiple activation steps. Preferably the CD4+ cells are activated every three days for 9 days. These CD4+ cells are preferably activated through cross-linking of CD3 and CD28 cell surface molecules, Cross-linking of CD3 and CD28 can be accomplished by adding immobilized anti-CD3 and CD28 monoclonal antibodies. Generally, the Th1/killer cells are obtained by differentiation and expansion in cell cultures. The multiply activated CD4+ cells are harvested from the cell culture and stored frozen in liquid nitrogen. Prior to administration to a patient, the cells are activated a final time. The activating agent is left associated with the cells. The cells with the activating agent attached is formulated in a device, such as a syringe, suitable for infusion or injection. "Activation" as used herein refers to both binding of the cell surface molecules by agents such as monoclonal antibodies and cross-linking of these agents by cross-linking agents. This type of activation is, for example, described in U.S. Pat. Nos. 7,435,592 and 7,402,431, both of which are incorporated herein by reference.

The CD4+ T-cells may be activated in a variety of ways including by the use of immobilized monoclonal antibodies specific for T-cell surface molecules. Suitable activated T-cells are, for example, described in U.S. Pat. No. 7,435,592. The cells preferably have cell surface moieties that are bound by monoclonal antibodies or other binding agents that are then cross-linked. These monoclonal antibodies and/or binding agents are preferably cross-linked by an agent, for example, immobilized on a solid surface in order to activate the T-cells. These are referred to herein as cells activated in culture (CAC). These ex vivo prepared CAC can be frozen for future use or formulated for infusion. The CAC, for example, may be aliquoted and frozen in the gas phase of Liquid Nitrogen tanks. The CAC are capable of consistently meeting pre-defined identity and functional release criteria as described below. In process and final quality control tests are established to assure sterility and endotoxin-free status.

In one exemplary embodiment, the CD4+ T-cells are cultured for 9 days in the presence of CD3/CD28 CLINEXVIVO™ DYNAL BEADS® obtained from INVITROGEN®, Carlsbad, Calif. The cells are grown in Life Cell Flasks (Baxter Scientific, Deerfield, Ill.) at 37° C. and 5% CO₂ and re-stimulated with the beads at days 3 and 6 of culture. After 9 days in culture, the beads can be removed and the cells can be referred to as CAC. Other methods of culturing cells for activation are also within the scope of the invention.

In preferred embodiments, the ex vivo prepared CAC are stored frozen until needed for patient administration or other uses. Prior to administration to the patient, the CAC are thawed, if necessary, washed and reactivated in nutrient media by cross-linking of the cell surface binding moieties such as CD3 and CD28 as described, for example in U.S. Pat. No. 7,402,431. The CAC, together with the activation and cross-linking agents, can then be washed and transferred to a non-nutritive buffer such as a formulation buffer. The reactivated cells in formulation buffer are referred to herein as cells in formulation buffer (CFB). CFB may also be referred to herein as Th1/killer cells. CFB and Th1/killer cells may be used interchangeably herein.

The Th1/killer cells can be packed in a syringe or other suitable containers and transferred to a point of care site, if needed. The Th1/killer cells can then be administered to the patient for therapeutic purposes. The Th1/killer cells may be administered intravenously, intraperitoneally, intradermally or by other suitable methods. Th1/killer cells, once transferred to non-nutritive buffer have a limited shelf life. Cells can be formulated at a density of at least about 10⁶ cells per ml, preferably at about 10⁷ cells per ml or higher. In some embodiments, the cells may be formulated at a density at about 10⁸ cells per ml or higher. The specific concentration of the cells may be determined by the specific use of the cells and the therapy protocol.

The composition may also include a number of other components in addition to the Th1/killer cells. These components can include, for example, agents that maintain the cells in the desired activation state. In one exemplary embodiment, the composition can include agents that maintain the T-cells in an activated state such as DYNABEADS® CLINEXVIVO™ described below in the Examples.

Generally, the cells are transferred to non-nutritive buffer that is appropriate for infusion into a patient. The cells can be in a variety of non-nutritive buffers. Non-nutritive buffer, as referred to herein, is any type of media, buffer or other liquid that lacks the appropriate components to support cellular proliferation and/or expansion. The non-nutritive buffers generally are isotonic, USP sterile, pyrogen-free and contain the appropriate components and/or buffering system to maintain live cells intact and are licensed for human parenteral use. In an exemplary embodiment, the non-nutritive buffer is a formulation buffer that is PLASMA-LYTE® A (Baxter Scientific, Deerfield, Ill.) with 1% human serum albumin. (McKesson, San Francisco, Calif.)

In embodiments with Th1/killer cells, the activation signals for the cells are maintained even when the cells are transferred to the non-nutritive buffer. For example, in embodiments where the cells are activated by cross-linking the cell surface binding moieties, the cross-linking is preferably maintained in the non-nutritive buffer. The maintenance of the cross-linking during storage may be critical to restoring the functional characteristics of the composition after removal from storage.

The Th1/killer cells in the composition can exhibit a novel combination of functional characteristics. These functions include Th1 characteristics and cytolytic functions that are similar to NK functions. The Th1/killer cells generally produce IFN-gamma and have substantially diminished or no IL-4 production. Preferably, the IL-4 production is below 50 pg/10⁶ cells and IFN-gamma production greater than 1000 pg/10⁶ cells.

Additional functional characteristics of the Th1/killer cells can include for example, expression of functional molecules such as CD40L, FasL, perforin and granzyme B, co-stimulatory molecules 4-1BBL, CD28, CTLA4, and TNF-related activation-induced cytokine (TRANCE), TWEAK, PD-1, B7 family, adhesion molecules such as the integrins, the cadherins, and the selectins and secretion of a variety of cytokines and chemokines and expression of receptors for these cytokines and chemokines. The Th1/killer cells can produce and secrete large amounts of cytokines. The cytokines produced can be, for example, IFN α , GM-CSF and TNF- α , with only residual IL4 secretion

The Th1/killer cells described herein can also exhibit cytolytic activity. The cytolytic activity is similar to the characteristics of CTL/NK cells. Generally, the cytolytic activity of CTL/NK cells, as well as the Th1/killer cells, is specific for tumor or diseased cells but not normal cells. In

other words, the Th1/killer cells can kill diseased cells but not normal cells. The cytolytic activity can occur through a variety of mechanisms. The cytolytic activity can occur, for example, through the Granzyme B-Perforin mechanism. The cytolytic activity may lead to destruction of the cells by cytolysis or apoptosis. The cytolytic activity of the Th1/killer cells can be blocked by EGTA. Concentration of EGTA at about 1 mM, for example, can decrease the cytolytic activity of the activated Th1 cells.

Th1/killer cells of the present invention can express the molecules Granzyme B and Perforin. Granzyme B and Perforin are generally not expressed in CD4+ T-cells, whether naïve or after maturation by differentiation, activation and the like. The amount of Granzyme B and Perforin in the activated Th1 cells can vary. Generally, the amount of Granzyme B and Perforin expressed is sufficient to destroy diseased cells.

In preferred embodiments, the present invention includes compositions that include Th1/killer cells. These Th1/killer cells can inactivate a variety of tumor cells including the myeloma tumor cell lines such as ARH77 cells. The inactivation of the diseased cells can be through a mechanism sensitive to EGTA. The effector cell (Th1/killer cells) to target cell (diseased cell) ratio can vary. The effector to target cell ratio can be at least 1:9, preferably at least 1:5 and more preferably at least 1:1.

The present invention includes a method for destroying diseased cells. The diseased cells can be in a patient. The patient, for example, may have cancer or contain infected tissue due to infection by a virus. The patient may have a solid tumor or hematological malignancy. The method can include administering to the patient a composition that includes the Th1/killer cells described herein. The Th1/killer cells are preferably allogeneic to the patient. The allogeneic cells may be from one normal allogeneic donor. Alternatively, the allogeneic cells may be derived from multiple donors and combined into a therapeutic composition. Preferably, the histocompatibility mismatch between the patient and allogeneic cells is maximized. Some histocompatibility between the patient and the allogeneic cells can be tolerated and can still result in the desired effect.

The Th1/killer cells can be administered to the patient by a variety of routes including intradermally, intravenously, intraperitoneally and the like. The patient may be administered one dose or multiple doses. The number of cells administered to the patient can vary and may depend on the specific disease, the patient, the method of activation and other factors. Generally, the patient is administered at least 10^7 of activated Th1 cells, preferably at least 10^8 and more preferably at least about 10^9 cells.

The present invention also includes a method of treating a patient by administering a composition that includes Th1/killer cells. The patient may receive one or more doses of the Th1/killer cells. The amount and the frequency of the doses can vary and can depend on the specific disease. In preferred embodiments, the Th1/killer cells are allogeneic to the patient and administration of the Th1/killer cells to the patient induces a host versus tumor effect along with a host versus graft effect without graft versus host disease.

EXAMPLES

Materials:

PE-conjugated CD40L was purchased from Beckman Coulter, Brea, Calif. 7-Amino-Actinomycin D (7-AAD) (1000 \times) was purchased from Cayman Chemical Co., Ann Arbor, Mich. PLASMALYTE® A was purchased from Bax-

ter Scientific, Deerfield, Ill. Human serum albumin (HSA) was purchased from McKesson, San Francisco, Calif. FcR Binding Inhibitor was purchased from eBioscience, San Diego, Calif. DYNABEADS® CLINEXVIVO™ was purchased from INVITROGEN®, Carlsbad, Calif.

Preparation of Cells Activated in Culture (CAC)—

CD4+ T-cells were cultured for 9 days in the presence of CD3/CD28 CLINEXVIVO™ DYNAL BEADS® obtained from INVITROGEN®, (Carlsbad, Calif.). The cells were grown in Life Cell Flasks (Baxter) at 37° C. and 5% CO₂ and re-stimulated with the beads at days 3 and 6 of culture. After 9 days in culture, the beads can be removed and these cells are referred to as CAC.

Preparation of Cells in Formulation Buffer (CFB)—

CAC were placed into cRPMI media for washing. Time was recorded to indicate the beginning of the formulation protocol. The cells in cRPMI media were centrifuged, the supernatant removed and the cells resuspended in cRPMI buffer. Cell viability was determined by using Trypan Blue assays. The total cell number and the concentration of live cells were used to determine the percentage of viable cells. If the sample had greater than 80 percent cell viability, then the procedure was continued for reactivation and formulation of cells.

The CAC cells were resuspended at a concentration of 1×10^7 cells/ml. Reactivation was done at a live cell concentration of 1×10^7 cells/ml. Reactivation was done in a 24 well plate, 6 well plate or a 75 cm³ flask depending on the volume. DYNABEADS® CLINEXVIVO™ CD3/CD28 were added to reactivate the cells and incubated at 36-38° C. and 5% CO₂ for 4 hours.

After incubation for about 4 hours, then the cells were removed and transferred to a 50 ml. tube with final formulation buffer (FFB). FFB is PLASMALYTE® A with 1% HSA. The reactivated cells were centrifuged, supernatant removed and resuspended in FFB. These are referred to as cells in formulation buffer (CFB).

CFB were resuspended in FFB at a concentration 10^7 cells per ml. The CFB were resuspended for ID, IT or IV administration. 1 ml of the cell suspension was added to a 3 ml syringe as an ID formulation. IT and IV formulation were 3 ml and 5 ml, respectively. The syringes with the appropriate formulations were stored in refrigeration with an average temperature of about 4° C.

Harvesting of Samples after Storage—

The cells and supernatant were collected at different time points. The time points were as follows: 0 (initial); 2 hours at Room Temp (RT); 48 hours at 4° C.; and 48 hours at 4° C. followed by 2 hours RT.

At each time point, 100 ul cell suspensions were collected and the cells were spun at 400 g for 5 min at 4° C. The supernatant was then transferred to another tube for IFN- γ detection later using ELISA. The cells were resuspended in 150 ul staining buffer for flow cytometry. In some experiments, the cells were resuspended in 100 ul cRPMI medium and cultured in the incubator at 37° C. for 24 h with 5% CO₂. The supernatant was taken after 24 h incubation and the IFN- γ was detected by ELISA.

Flow Cytometry (CD40L and 7-AAD)—

50 ul cell suspension were transferred from above (150 ul) into 3 eppendorf tubes, labeled as unstained, CD40L and 7-AAD, respectively. The unstained tube was incubated on ice for 20 min. For the CD40L tube, the cells were pre-incubated with FcR Binding inhibitor according to the instructions of the manufacturer for 20 minutes on ice. Then 40 ul staining buffer (PBS+1% FBS) and 10 ul PE-CD40L

11

antibody was added into the cell suspension and incubated for additional 20 min on ice in the dark.

Cell viability was tested by flow cytometry of 7-AAD. 7-AAD intercalates into DNA of dead or damaged cells, thus determination of 7-AAD positive cells is an indicator of cell viability. For 7-AAD tubes, the tubes were centrifuged at 400 g for 5 min at 6 C. After removing the supernatant, the cell pellets were resuspended in 100 ul 1x7-AAD solution. The tube was incubated on ice for 15 min in the dark. 1 ml of staining buffer was added to the CD40L tube and then the 3 tubes were centrifuged together. After discarding the supernatant, the cell pellets were resuspended in 0.4 ml staining buffer and FACS was run.

Example 1

Purification and cell phenotyping of CD4+ cells. Normal donor peripheral blood was obtained and the CD4+ cells from the buffy coat were purified using Miltenyi magnetic beads and column obtained from Miltenyi Biotec (Auburn, Calif.). FIG. 1 is a flow cytometry plot of side scatter (measures size and density of cells) vs. CD4. FIG. 1 shows that CD4+ cells are about 46% of the total population (N=22). After positive selection for CD4+ cells, the result is 98.69% pure CD4+ cells (N=22). As can be seen from FIG. 1B, the population of CD4+ after the column contains both lymphocytes and monocytes.

Since every batch is produced from a different donor, antigen markers were tested at the beginning and the end of production. Cell phenotyping of CD4+ cells (FIG. 2A) versus Cells activated in culture (CAC) (FIG. 2B) were tested. The results show a very consistent shift from a naïve CD4+ phenotype (CD45RA+) to a memory phenotype (CD45RO+). FIG. 2A shows the phenotypes of the CD4+ cells for CD45RA, CD45RO and CD25, respectively. FIG. 2B shows the phenotypes of the CAC after 9 days of culturing and analysis for surface expression of CD45RA, CD45RO and CD25, respectively. From a mixed CD45RA/CD45RO population, a CD45RA negative CD45RO positive population is obtained. CD25+ cells are approximately 10% at the beginning of the expansion process. At day 9, CD25+ was approximately 85%. This is in accordance with the activation state of the cells. FIG. 2A shows CD4+ cells after column show a naïve phenotype of CD45RA+ cells (avg=44.76% n=29), CD45RO+(avg=53.44% n=29) and CD25+(avg=10.83% n=29). FIG. 2B shows that after 9 days in culture with continued stimulation, the phenotype changes into Th memory like cells. CD45RA goes down drastically (avg=4.22% n=29), CD45RO+ goes up to avg. of 97.8 (n=29), and CD25+ also goes up to average of 90.03% (n=29).

CD62L Expression pattern—FIG. 3 shows that at Day 0, CD4+ seed cells express a bimodal CD62L mean fluorescent intensity (MFI) with a maximum peak at approximately 10³ (the average arithmetic mean is 541 n=17). The MFI increases after 9 days in culture to an average arithmetic mean of 1541 (CAC) and then becomes significantly lower (average arithmetic mean of 190.3) after the final 4 hours activation (CFB).

CD40L Expression Pattern—The density of CD40L expression on CAC was consistently increased on the surface of CAC after a 4 hour activation with CD3/CD28 beads. FIG. 4 shows the change in the arithmetic mean and percentage of CD40L expression of a representative batch comparing CD4+ at day 0 (FIG. 4A), day 9 harvested CAC before final activation (FIG. 4B) and after 4 hours activation, CFB (FIG. 4C). As can be seen, the difference between the

12

un-activated and activated cells is not so much in the percent of CD40L+ cells, but rather in the arithmetic mean (the average AM for CAC is 20 (n=29) while the average for CFB is 112 (n=29))

Cytokine secretion during final activation—Samples of 9 day harvested CAC cells from each production batch were activated with CD3/CD28 beads for 4 hours and the supernatants were collected and tested for cytokine production (IFN γ , IL-4, IL-8, TNF α and GM-CSF) using ELISA assay. Table 1 shows the results of these experiments.

TABLE 1

Batch Code	IFN γ	IL4	GM-CSF	TNF α
068C	2,178.3	0.00	647.9	1,338.3
069C	2,500.0	0.00	809.9	1,245.8
070C	5,469.1	0.00	1,595.9	1,994.9
071C	6,995.4	0.00	1,739.3	2,799.0
072C	806.6	0.00	251.1	117.7
073C	2,931.5	0.00	596.6	358.3
074C	462.9	0.00	0.0	105.3
075C	2,722.5	0.00	927.0	1,189.7
076C	4,118.0	0.30	752.0	576.7
078C	1,072.0	0.00	294.2	393.1
080C	3,846.7	0.90	1,878.4	1,828.4
081C	9,702.8	0.35	9,030.4	1,522.9
082C	6,662.4	0.76	3,984.9	869.0
083C	910.8	0.00	344.4	365.5
087C	2,257.6	0.00	764.6	736.8
088C	4,342.1	0.00	2,539.4	3,961.5
090C	1,021.4	1.00	402.0	810.5
094C	1,420.2	1.03	733.6	757.2
097C	703.2	1.16	327.7	989.4
Average (n = 15)	3,816.0	0.29	1,779.7	1,358.8

Only IFN γ and IL-4 values were used for functional release criteria. The results for 19 consecutive batches are shown Table 1 (values are expressed as pg/10⁶ cells/4 hours). The reported average includes only passing batches.

Example 2

Direct Killing by activated Th1/killer cells. To test if CFB may have direct effect on tumor cells, the ability of the cells to kill the ARH77 cell line was tested. This cell line is an established myeloma cell line. The ARH77 cells were stained in CFSE to differentiate them from the CFB. CFB were mixed with the stained ARH77 cells at different Effector (CFB) to Target (ARH77) for 18 hours. After that time, 7AAD was added and cells were acquired using the FC500MPL flow cytometer. This experiment was repeated with different CAC batches and with fresh and thawed CFB. ARH77 alone, with DYNABEADS®, CD4+ cells or with CAC had no significant death (less than 10%) data not shown. The results shown in FIG. 5 indicate that CFB, but not CAC, CD4+ cells or beads, have direct killing effect on the ARH77 cell line.

The nature of the direct killing effect was examined to see if the effect was due to the Perforin-Granzyme pathway. Both internal staining and western blot techniques were used to stain for Perforin and Granzyme B. FIG. 6A shows that CAC express very little Granzyme B, as can be seen. FIG. 6B shows that CFB express significant amount of Granzyme B. FIG. 6C shows that Perforin can be detected only by the use of western blot. Lane 1 is the extract of activated PBMCs, as a positive control. Lane 2&4 are of CAC from different batches, and lanes 3&5 are of the CFB from the same batches.

Inhibition assays were also performed to elucidate the nature of the direct killing mechanism. CFB were mixed with ARH77 cells as described above and different inhibitory agents were added as shown below Table 2. Antibodies against CD40L and/or FAS-L were added to the CFB: ARH77 mix. Table 2 shows the even the highest concentration did not prevent the killing profile.

TABLE 2

	Anti C40L	no inhibition	0.1 μ g/ml	1 μ g/ml	20 μ g/ml
Anti Fas-L no inhibition		33.85	35.48	35.92	35.5
1 ng/ml	36.55		34.55		
10 ng/ml	38.86				
200 ng/ml	38.97				30.0

The effect of EGTA on the direct killing mechanism was examined. EGTA blocks the polymerization of Perforin and acts as an inhibitor of the Perforin/Granzyme B mechanism. 1 mM EGTA was added to the direct killing assay described above. As shown in FIG. 7, 1 mM EGTA reduced the killing by an average of 50% (n=7) Paired t test P value 0.0002. The direct killing effect that CFB show comes from the Perforin-Granzyme pathway.

Although the present invention has been described with reference to preferred embodiments, workers skilled in the art will recognize that changes may be made in form and detail without departing from the spirit and scope of the invention.

What is claimed is:

1. A method for killing tumor cells comprising: contacting the tumor cells from a tumor cell line with a composition comprising ex-vivo activated CD4+ Th1 cells without the presence of any other cell types, wherein the activated CD4+ Th1 cells have cytolytic activity mediated by Granzyme B and perforin, wherein the CD4+ Th1 cells are frozen after harvesting from cell culture, thawed reactivated by both binding of the CD3/CD28 cell surface molecules by agents and cross-linking of these agents in the composition by the presence of the cross-linking agents in the composition, washed and transferred to a non-nutritive buffer, wherein the CD4+ Th1 cells express IFN-gamma at greater than 1000 pg/10⁶ cells and wherein the interaction of the CD4+ Th1 cells with the tumor cells leads to the killing of the tumor cells.
2. The method of claim 1 wherein the CD4+ Th1 cells lack expression of IL-4.
3. The method of claim 1 wherein the CD4+ Th1 cells are activated by cross-linked anti-CD3/anti-CD28 monoclonal antibodies.
4. The method of claim 1 wherein the tumor cell line is ARH77.
5. The method of claim 1 wherein the ratio of the CD4+ Th1 cells to cells of the tumor cell line is at least 1:9.
6. The method of claim 1 wherein the composition comprises at least about 10⁷ cells/ml.

* * * * *