



US 20100267685A1

(19) **United States**

(12) **Patent Application Publication**
Straub et al.

(10) **Pub. No.: US 2010/0267685 A1**

(43) **Pub. Date: Oct. 21, 2010**

(54) **METHODS FOR THE PROPHYLAXIS
AND/OR TREATMENT OF
THROMBOEMBOLIC DISORDERS BY
COMBINATION THERAPY WITH
SUBSTITUTED OXAZOLIDINONES**

(75) Inventors: **Alexander Straub**, Wuppertal
(DE); **Thomas Lampe**, Dusseldorf
(DE); **Josef Pernerstorfer**,
Wuppertal (DE); **Elisabeth
Perzborn**, Wuppertal (DE); **Jens
Pohlmann**, Wuppertal (DE);
Susanne Röhrig, Essen (DE);
Karl-Heinz Schlemmer, Wuppertal
(DE)

Correspondence Address:
CONNOLLY BOVE LODGE & HUTZ, LLP
P O BOX 2207
WILMINGTON, DE 19899 (US)

(73) Assignee: **Bayer Schering Pharma
Aktiengesellschaft**, Berlin (DE)

(21) Appl. No.: **12/766,365**

(22) Filed: **Apr. 23, 2010**

Related U.S. Application Data

(62) Division of application No. 10/481,297, filed on Jun. 28, 2004, now Pat. No. 7,767,702, filed as application No. PCT/EP2002/006237 on Jun. 7, 2002.

(30) **Foreign Application Priority Data**

Jun. 20, 2001 (DE) DE 101 29 725.4

Publication Classification

(51) **Int. Cl.**

<i>A61K 31/397</i>	(2006.01)
<i>A61K 31/5377</i>	(2006.01)
<i>A61K 31/541</i>	(2006.01)
<i>A61K 31/4365</i>	(2006.01)
<i>A61K 31/428</i>	(2006.01)
<i>A61K 31/496</i>	(2006.01)
<i>A61K 31/444</i>	(2006.01)
<i>A61K 31/454</i>	(2006.01)
<i>A61K 31/422</i>	(2006.01)
<i>A61K 31/423</i>	(2006.01)
<i>A61K 31/538</i>	(2006.01)
<i>A61K 31/4439</i>	(2006.01)
<i>A61K 31/5355</i>	(2006.01)
<i>A61K 31/427</i>	(2006.01)
<i>A61P 9/08</i>	(2006.01)
<i>A61P 7/02</i>	(2006.01)
<i>A61P 9/10</i>	(2006.01)
<i>A61P 3/06</i>	(2006.01)
<i>A61P 9/00</i>	(2006.01)
<i>A61P 9/04</i>	(2006.01)
<i>A61K 31/5365</i>	(2006.01)

(52) **U.S. Cl.** **514/210.02**; 514/236.8; 514/227.8;
514/301; 514/367; 514/253.1; 514/254.02;
514/333; 514/326; 514/376; 514/375; 514/230.5;
514/343; 514/228.8; 514/230.8; 514/210.2;
514/372; 514/230.2

(57) **ABSTRACT**

The invention relates to combinations of A) oxazolidinones of formula (I) and B) other active ingredients, to a method for producing said combinations and to the use thereof as medicaments, in particular for the treatment and/or prophylaxis of thrombo-embolic diseases.

**METHODS FOR THE PROPHYLAXIS
AND/OR TREATMENT OF
THROMBOEMBOLIC DISORDERS BY
COMBINATION THERAPY WITH
SUBSTITUTED OXAZOLIDINONES**

[0001] The present invention relates to combinations of A) oxazolidinones of the formula (I) with B) other active ingredients, to a process for producing these combinations and to the use thereof as medicaments, in particular for the prophylaxis and/or treatment of thromboembolic disorders.

[0002] Oxazolidinones of the formula (I) act in particular as selective inhibitors of coagulation factor Xa and as anticoagulants.

[0003] It has been possible to demonstrate an antithrombotic effect of factor Xa inhibitors in numerous animal models (cf. WO 99/37304; WO 99/06371; J. Hauptmann, J. Stürzebecher, Thrombosis Research 1999, 93, 203; F. Al-Obeidi, J. A. Ostrem, Factor Xa inhibitors, Exp. Opin. Ther. Patents 1999, 9, 931; B.-Y. Zhu, R. M. Scarborough, Curr. Opin. Card. Pulm. Ren. Inv. Drugs 1999, 1 (1), 63; M. Samama, J. M. Walenga, B. Kaiser, J. Fareed, Specific Factor Xa Inhibitors, Cardiovascular Thrombosis: Thrombocardiology and Thromboneurology, Second Edition, edited by M. Verstraete, V. Fuster, E. J. Topol, Lippincott-Raven Publishers, Philadelphia 1998) and clinical studies on patients (The Ephesus Study, blood, Vol 96, 490a, 2000; The Penthifra Study, blood, Vol 96, 490a, 2000; The Pentamaks Study, blood, Vol 96, 490a-491a, 2000; The Pentathlon 2000 Study, blood, Vol 96, 491a, 2000). Factor Xa inhibitors can therefore be employed preferably in medicaments for the prophylaxis and/or treatment of thrombo embolic disorders.

[0004] Thromboembolic vascular disorders are the commonest cause of morbidity and mortality in industrialized countries (Thiemes Innere Medizin, Georg Thieme Verlag Stuttgart, New York; American Heart Association, 2000 heart and stroke statistical update, Dallas, Tex.: American Heart Association, 2000). Anticoagulant therapy has proved effective in the treatment of vascular disorders in order to prevent thrombotic vascular occlusions and to reopen thrombotically occluded vessels, and has great importance in the prophylaxis and treatment of coronary, peripheral and cerebral vascular disorders, and in the prophylaxis and/or treatment of venous thromboses and pulmonary embolisms.

[0005] Thromboembolic complications may be caused by atherosclerotic lesions of the vessel wall, especially disturbance of endothelial function, which may lead to acute thrombotic occlusions. Atherosclerosis is a multifactorial disorder which depends on a large number of cardiovascular risk factors. Clinical studies have shown that prophylaxis with anticoagulants does not definitively influence the course of the arterial vascular disorder. Targeted treatment of the risk factors in conjunction with an antithrombotic therapy is therefore advantageous.

[0006] Risk factors for coronary, peripheral and cerebral vascular disorders are, for example: elevated serum cholesterol levels, arterial hypertension, cigarette smoking, diabetes mellitus (Allgemeine und spetargetle Pharmacologie und Toxikologie, W. Forth, D. Henschler, W. Rummel, K. Starke; Spektrum Akademischer Verlag Heidelberg Berlin Oxford; Thiemes Inhere Medizin, Georg Thieme Verlag Stuttgart, New York). The principles of preventive medicine are based on elimination of these risk factors. Besides a change in

lifestyle, also included are pharmacological measures such as, for example, antihypertensive therapy, lipid-lowering medicaments or thrombosis prophylaxis. In addition, combination with coronary therapeutic agents is suitable for the treatment where there is pre-existent coronary heart disease.

[0007] It has now been found, surprisingly, that combinations of oxazolidinones of the formula (I) with certain other active ingredients have interesting properties and are more suitable for the prophylaxis and/or treatment of various diseases than the individual active ingredients alone.


[0008] The invention therefore relates to combinations of

[0009] A) oxazolidinones of the formula (I) with

[0010] B) other active ingredients, in particular with platelet aggregation inhibitors, anticoagulants, fibrinolytics, lipid-lowering agents, coronary therapeutic agents and/or vasodilators.

[0011] "Combinations" mean for the purposes of the invention not only dosage forms which comprise all the components (so-called fixed combinations), and combination packs which comprise the components separate from one another, but also components administered simultaneously or sequentially as long as they are employed for the prophylaxis and/or treatment of the same disease. It is likewise possible to combine two or more active ingredients together, and thus the combinations in this connection are in each case double or multiple.

[0012] Suitable oxazolidinones of the combination of the invention include, for example, compounds of the formula (I)



(I)

in which:

[0013] R¹ is optionally benzo-fused thiophene (thienyl) which may optionally be substituted one or more times;

[0014] R² is any organic radical;

[0015] R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are identical or different and are hydrogen or (C₁-C₆)-alkyl, and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.

[0016] Preference is given in this connection to compounds of the formula (I)

in which

[0017] R¹ is optionally benzo-fused thiophene (thienyl) which may optionally be substituted one or more times by a radical from the group of halogen; cyano; nitro; amino; aminomethyl; (C₁-C₈)-alkyl which may in turn be optionally substituted one or more times by halogen; (C₃-C₇)-cycloalkyl; (C₁-C₃)-alkoxy; imidazolyl; —C(=NH)NH₂; carbamoyl; and mono- and di-(C₁-C₄)-alkylaminocarbonyl,

[0018] R² is one of the following groups:

[0019] A-,


[0020] A-M-,

[0021] D-M-A-,

- [0022] B-M-A-,
- [0023] B—,
- [0024] B-M-,
- [0025] B-M-B—,
- [0026] D-M-B—,
- [0027] where:
- [0028] the radical "A" is (C₆-C₁₄)-aryl, preferably (C₆-C₁₀)-aryl, in particular phenyl or naphthyl, very particularly preferably phenyl;
- [0029] the radical "B" is a 5- or 6-membered aromatic heterocycle which comprises up to 3 heteroatoms and/or hetero chain members, in particular up to 2 heteroatoms and/or hetero chain members, from the series S, N, NO (N-oxide) and O;
- [0030] the radical "D" is a saturated or partially unsaturated, mono- or bicyclic, optionally benzo-fused 4- to 9-membered heterocycle which comprises up to three heteroatoms and/or hetero chain members from the series S, SO, SO₂, N, NO(N-oxide) and O;
- [0031] the radical "M" is —NH—, —CH₂—, —CH₂CH₂—, —O—, —NH—CH₂—, —CH₂—NH—, —OCH₂—, —CH₂O—, —CONH—, —NHCO—, —COO—, —OOC—, —S—, —SO₂— or a covalent bond;
- [0032] where
- [0033] the groups "A", "B" and "D" defined above may in each case optionally be substituted one or more times by a radical from the group of halogen; trifluoromethyl; oxo; cyano; nitro; carbamoyl; pyridyl; (C₁-C₆)-alkanoyl; (C₃-C₇)-cycloalkanoyl; (C₆-C₁₄)-arylcarbonyl; (C₅-C₁₀)-heteroarylcarbonyl; (C₁-C₆)-alkanoyloxymethoxy; (C₁-C₄)-hydroxyalkylcarbonyl; —COOR²⁷; —SO₂R²⁷; —C(NR²⁷R²⁸)=NR²⁹; —CONR²⁸R²⁹; SO₂NR²⁸R²⁹; —OR³⁰; —NR³⁰R³¹; (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl,
- [0034] where (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl in turn may optionally be substituted by a radical from the group of cyano; —OR²⁷; —NR²⁸R²⁹; —CO(NH), (NR²⁷R²⁸) and —C(NR²⁷R²⁸)=NR²⁹,
- [0035] where:
- [0036] v is either 0 or 1 and
- [0037] R²⁷; R²⁸ and R²⁹ are identical or different and are, independently of one another, hydrogen, (C₁-C₄)-alkyl, (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkanoyl, carbamoyl, trifluoromethyl, phenyl or pyridyl,
- [0038] and/or
- [0039] R²⁷ and R²⁸, or R²⁷ and R²⁹, form together with a nitrogen atom to which they are bonded a saturated or partially unsaturated 5- to 7-membered heterocycle having up to three, preferably up to two, identical or different heteroatoms from the group of N, O and S, and
- [0040] R³⁰ and R³¹ are identical or different and are, independently of one another, hydrogen, (C₁-C₄)-alkyl, (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-hydroxyalkyl, (C₁-C₄)-aminoalkyl, di-(C₁-C₄)-alkylamino-(C₁-C₄)-alkyl, —CH₂C(NR²⁷R²⁸)=NR²⁹ or —COR³³,
- [0041] where
- [0042] R³³ is (C₁-C₆)-alkoxy, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, (C₁-C₄)-alkoxycarbonyl-(C₁-C₄)-alkyl, (C₁-C₄)-aminoalkyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkanoyl-(C₁-C₄)-alkyl, (C₃-C₇)-cycloalkyl, (C₂-C₆)-alkenyl, (C₁-C₈)-alkyl which may optionally be substituted by phenyl or acetyl, or is (C₆-C₁₄)-aryl, (C₅-C₁₀)-heteroaryl, trifluoromethyl, tetrahydrofuranyl or butyrolactone,
- [0043] R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are identical or different and are hydrogen or (C₁-C₆)-alkyl, and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.
- [0044] Preference is likewise given in this connection to compounds of the general formula (I) in which
- [0045] R¹ is thiophene (thienyl), in particular 2-thiophene, which may optionally be substituted one or more times by halogen, preferably chlorine or bromine, amino, aminomethyl or (C₁-C₈)-alkyl, preferably methyl, where the (C₁-C₈)-alkyl radical may optionally in turn be substituted one or more times by halogen, preferably fluorine,
- [0046] R² is one of the following groups:
- [0047] A-,
- [0048] A-M-,
- [0049] D-M-A-,
- [0050] B-M-A-,
- [0051] B—,
- [0052] B-M-,
- [0053] B-M-B—,
- [0054] D-M-B—,
- [0055] where:
- [0056] the radical "A" is (C₆-C₁₄)-aryl, preferably (C₆-C₁₀)-aryl, in particular phenyl or naphthyl, very particularly preferably phenyl;
- [0057] the radical "B" is a 5- or 6-membered aromatic heterocycle which comprises up to 3 heteroatoms and/or hetero chain members, in particular up to 2 heteroatoms and/or hetero chain members, from the series S, N, NO(N-oxide) and O;
- [0058] the radical "D" is a saturated or partially unsaturated 4- to 7-membered heterocycle which comprises up to three heteroatoms and/or hetero chain members from the series S, SO, SO₂, N, NO(N-oxide) and O;
- [0059] the radical "M" is —NH—, —CH₂—, —CH₂CH₂—, —O—, —NH—CH₂—, —CH₂—NH—, —OCH₂—, —CH₂O—, —CONH—, —NHCO—, —COO—, —OOC—, —S— or a covalent bond;
- [0060] where
- [0061] the groups "A", "B" and "D" defined above may in each case optionally be substituted one or more times by a radical from the group of halogen; trifluoromethyl; oxo; cyano; nitro; carbamoyl; pyridyl; (C₁-C₆)-alkanoyl; (C₃-C₇)-cycloalkanoyl; (C₆-C₁₄)-arylcarbonyl; (C₅-C₁₀)-heteroarylcarbonyl; (C₁-C₆)-alkanoyloxymethoxy; —COOR²⁷; —SO₂R²⁷; —C(NR²⁷R²⁸)=NR²⁹; —CONR²⁸R²⁹; —SO₂NR²⁸R²⁹; —OR³⁰; —NR³⁰R³¹, (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl,
- [0062] where (C₁-C₆)-alkyl and (C₃-C₇)-cycloalkyl may in turn optionally be substituted by a radical from the group of cyano; —OR²⁷; —NR²⁸R²⁹; —CO(NH), (NR²⁷R²⁸) and —C(NR²⁷R²⁸)=NR²⁹;
- [0063] where:
- [0064] v is either 0 or 1, and
- [0065] R²⁷, R²⁸ and R²⁹ are identical or different and are, independently of one another, hydrogen, (C₁-C₄)-alkyl or (C₃-C₇)-cycloalkyl,
- [0066] and/or
- [0067] R²⁷ and R²⁸, or R²⁷ and R²⁹, form together with the nitrogen atom to which they are bonded a saturated

- or partially unsaturated 5- to 7-membered heterocycle having up to three, preferably up to two, identical or different heteroatoms from the group of N, O and S, and
- [0068] R^{30} and R^{31} are identical or different and are, independently of one another, hydrogen, (C_1-C_4) -alkyl, (C_3-C_7) -cycloalkyl, (C_1-C_4) -alkylsulfonyl, (C_1-C_4) -hydroxyalkyl, (C_1-C_4) -aminoalkyl, di- (C_1-C_4) -alkylamino- (C_1-C_4) -alkyl, (C_1-C_4) -alkanoyl, (C_6-C_{14}) -arylcarbonyl, (C_5-C_{10}) -heteroarylcarbonyl, (C_1-C_4) -alkylaminocarbonyl or $-\text{CH}_2\text{C}(\text{NR}^{27}\text{R}^{28})=\text{NR}^{29}$,
- [0069] R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are identical or different and are hydrogen or (C_1-C_6) -alkyl,
- and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.
- [0070] Particular preference is given in this connection to compounds of the general formula (I) in which
- [0071] R^1 is thiophene (thienyl), in particular 2-thiophene, which may optionally be substituted one or more times by halogen, preferably chlorine or bromine, or (C_1-C_8) -alkyl, preferably methyl, where the (C_1-C_8) -alkyl radical may in turn optionally be substituted one or more times by halogen, preferably fluorine,
- [0072] R^2 is one of the following groups:
- [0073] A-,
- [0074] A-M-,
- [0075] D-M-A-,
- [0076] B-M-A-,
- [0077] B-,
- [0078] B-M-,
- [0079] B-M-B-,
- [0080] D-M-B-,
- [0081] where:
- [0082] the radical "A" is phenyl or naphthyl, in particular phenyl;
- [0083] the radical "B" is a 5- or 6-membered aromatic heterocycle which comprises up to 2 heteroatoms from the series S, N, NO(N-oxide) and O;
- [0084] the radical "D" is a saturated or partially unsaturated 5- or 6-membered heterocycle which comprises up to two heteroatoms and/or hetero chain members from the series S, SO, SO_2 , N, NO(N-oxide) and O;
- [0085] the radical "M" is $-\text{NH}-$, $-\text{O}-$, $-\text{NH}-\text{CH}_2-$, $-\text{CH}_2-\text{NH}-$, $-\text{OCH}_2-$, $-\text{CH}_2\text{O}-$, $-\text{CONH}-$, $-\text{NHCO}-$ or a covalent bond;
- [0086] where
- [0087] the groups "A", "B" and "D" defined above may in each case optionally be substituted one or more times by a radical from the group of halogen; trifluoromethyl; oxo; cyano; pyridyl; (C_1-C_3) -alkanoyl; (C_6-C_{10}) -arylcarbonyl; (C_5-C_6) -heteroarylcarbonyl; (C_1-C_3) -alkanoyloxymethoxy; $-\text{C}(\text{NR}^{27}\text{R}^{28})=\text{NR}^{29}$; $-\text{CONR}^{28}\text{R}^{29}$; $-\text{SO}_2\text{NR}^{28}\text{R}^{29}$; $-\text{OH}$; $-\text{NR}^{30}\text{R}^{31}$; (C_1-C_4) -alkyl; and cyclopropyl, cyclopentyl or cyclohexyl,
- [0088] where (C_1-C_4) -alkyl and cyclopropyl, cyclopentyl or cyclohexyl may in turn optionally be substituted by a radical from the group of cyano; $-\text{OH}$; $-\text{OCH}_3$; $-\text{NR}^{28}\text{R}^{29}$; $-\text{CO}(\text{NH})_v(\text{NR}^{27}\text{R}^{28})$ and $-\text{C}(\text{NR}^{27}\text{R}^{28})=\text{NR}^{29}$,
- [0089] where:
- [0090] v is either 0 or 1, preferably 0, and
- [0091] R^{27} , R^{28} and R^{29} are identical or different and are, independently of one another, hydrogen, (C_1-C_4) -alkyl or else cyclopropyl, cyclopentyl or cyclohexyl,
- [0092] and/or
- [0093] R^{27} and R^{28} , or R^{27} and R^{29} , may form together with the nitrogen atom to which they are bonded a saturated or partially unsaturated 5- to 7-membered heterocycle having up to two identical or different heteroatoms from the group of N, O and S, and
- [0094] R^{30} and R^{31} are identical or different and are, independently of one another, hydrogen, (C_1-C_4) -alkyl, cyclopropyl, cyclopentyl, cyclohexyl, (C_1-C_4) -alkylsulfonyl, (C_1-C_4) -hydroxyalkyl, (C_1-C_4) -aminoalkyl, di- (C_1-C_4) -alkylamino- (C_1-C_4) -alkyl, (C_1-C_3) -alkanoyl or phenylcarbonyl,
- [0095] R^3 , R^4 , R^5 , R^6 , R^7 and R^8 are identical or different and are hydrogen or (C_1-C_6) -alkyl,
- and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.
- [0096] Especial preference is given in this connection to compounds of the general formula (I) in which
- [0097] R^1 is 2-thiophene which may optionally be substituted in position 5 by a radical from the group chlorine, bromine, methyl or trifluoromethyl,
- [0098] R^2 is one of the following groups:
- [0099] A-,
- [0100] A-M-,
- [0101] D-M-A-,
- [0102] B-M-A-,
- [0103] B-,
- [0104] B-M-,
- [0105] B-M-B-,
- [0106] D-M-B-,
- [0107] where:
- [0108] the radical "A" is phenyl or naphthyl, in particular phenyl;
- [0109] the radical "B" is a 5- or 6-membered aromatic heterocycle which comprises up to 2 heteroatoms from the series S, N, NO(N-oxide) and O;
- [0110] the radical "D" is a saturated or partially unsaturated 5- or 6-membered heterocycle which comprises a nitrogen atom and optionally a further heteroatom and/or hetero chain member from the series S, SO, SO_2 and O;
- [0111] the radical "M" is $-\text{NH}-$, $-\text{O}-$, $-\text{NH}-\text{CH}_2-$, $-\text{CH}_2-\text{NH}-$, $-\text{OCH}_2-$, $-\text{CH}_2\text{O}-$, $-\text{CONH}-$, $-\text{NHCO}-$ or a covalent bond;
- [0112] where
- [0113] the groups "A", "B" and "D" defined above may in each case optionally be substituted one or more times by a radical from the group of halogen; trifluoromethyl; oxo; cyano; pyridyl; (C_1-C_3) -alkanoyl; (C_6-C_{10}) -arylcarbonyl; (C_5-C_6) -heteroarylcarbonyl; (C_1-C_3) -alkanoyloxymethoxy; $-\text{CONR}^{28}\text{R}^{29}$; $-\text{SO}_2\text{NR}^{28}\text{R}^{29}$; $-\text{OH}$; $-\text{NR}^{30}\text{R}^{31}$; (C_1-C_4) -alkyl; and cyclopropyl, cyclopentyl or cyclohexyl,
- [0114] where (C_1-C_4) -alkyl and cyclopropyl, cyclopentyl or cyclohexyl may in turn optionally be substituted by a radical from the group of cyano; $-\text{OH}$; $-\text{OCH}_3$; $-\text{NR}^{28}\text{R}^{29}$; $-\text{CO}(\text{NH})_v(\text{NR}^{27}\text{R}^{28})$ and $-\text{C}(\text{NR}^{27}\text{R}^{28})=\text{NR}^{29}$,

- [0115] where:
 [0116] v is either 0 or 1, preferably 0, and
 [0117] R²⁷, R²⁸ and R²⁹ are identical or different and are, independently of one another, hydrogen, (C₁-C₄)-alkyl or else cyclopropyl, cyclopentyl or cyclohexyl,
 [0118] and/or
 [0119] R²⁷ and R²⁸, or R²⁷ and R²⁹, may form together with the nitrogen atom to which they are bonded a saturated or partially unsaturated 5- to 7-membered heterocycle having up to two identical or different heteroatoms from the group of N, O and S, and
 [0120] R³⁰ and R³¹ are identical or different and are, independently of one another, hydrogen, (C₁-C₄)-alkyl, cyclopropyl, cyclopentyl, cyclohexyl, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-hydroxyalkyl, (C₁-C₄)-aminoalkyl, di-(C₁-C₄)-alkylamino-(C₁-C₄)-alkyl, (C₁-C₃)-alkanoyl or phenylcarbonyl,
 [0121] R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are identical or different and are hydrogen or (C₁-C₄)-alkyl, and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.
 [0122] Very particular preference is given in this connection to compounds of the general formula (I) in which
 [0123] R¹ is 2-thiophene which is substituted in position 5 by a radical from the group of chlorine, bromine, methyl or trifluoromethyl,
 [0124] R² is D-A-:
 [0125] where:
 [0126] the radical "A" is phenylene;
 [0127] the radical "D" is a saturated 5- or 6-membered heterocycle which
 [0128] is linked via a nitrogen atom to "A",
 [0129] which has a carbonyl group in direct vicinity to the linking nitrogen atom, and in which a ring carbon member may be replaced by a heteroatom from the series S, N and O;
 [0130] where
 [0131] the group "A" defined above may optionally be substituted once or twice in the meta position relative to the linkage to the oxazolidinone by a radical from the group of fluorine, chlorine, nitro, amino, trifluoromethyl, methyl or cyano,
 [0132] R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are hydrogen, and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.
 [0133] Very particular preference is likewise given in this connection to the compound having the following formula



and the pharmaceutically acceptable salts, hydrates and prodrugs thereof.

[0134] To date, oxazolidinones have been described essentially only as antibiotics, and in a few cases also as MAO inhibitors and fibrinogen antagonists (Review: Riedl, B., Endermann, R., Exp. Opin. Ther. Patents 1999, 9 (5), 625), and a small 5-[acylaminoethyl] group (preferably 5-[acetylaminoethyl]) appears to be essential for the antibacterial effect.

[0135] Substituted aryl- and heteroarylphenyloxazolidinones in which a monosubstituted or polysubstituted phenyl radical may be bonded to the N atom of the oxazolidinone ring and which may have in position 5 of the oxazolidinone ring an unsubstituted N-methyl-2-thiophenecarboxamide residue, and their use as substances with antibacterial activity are disclosed in the U.S. Pat. No. 5,929,248, U.S. Pat. No. 5,801,246, U.S. Pat. No. 5,756,732, U.S. Pat. No. 5,654,435, U.S. Pat. No. 5,654,428 and U.S. Pat. No. 5,565,571.

[0136] In addition, benzamidine-containing oxazolidinones are known as synthetic intermediates in the synthesis of factor Xa inhibitors or fibrinogen antagonists (WO-A-99/31092, EP-A-623615).

[0137] The compounds of the formula (I) may, depending on the substitution pattern, exist in stereoisomeric forms which either are related as image and mirror image (enantiomers) or are not related as image and mirror image (diastereomers). Both the enantiomers or diastereomers and respective mixtures thereof are included. The racemic forms can, just like the diastereomers, be separated in a known manner into the stereoisomerically pure constituents.

[0138] Certain compounds of the formula (I) may also exist in tautomeric forms. This is known to the skilled worker, and such compounds are likewise included.

[0139] Physiologically acceptable, i.e. pharmaceutically acceptable, salts may be salts of the compounds of the invention with inorganic or organic acids. Preferred salts are those with inorganic acids such as, for example, hydrochloric acid, hydrobromic acid, phosphoric acid or sulfuric acid, or salts with organic carboxylic or sulfonic acids such as, for example, acetic acid, trifluoroacetic acid, propionic acid, maleic acid, fumaric acid, malic acid, citric acid, tartaric acid, lactic acid, benzoic acid, or methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, toluenesulfonic acid or naphthalenedisulfonic acid.

[0140] Pharmaceutically acceptable salts which may also be mentioned are salts with conventional bases, such as, for example, alkali metal salts (e.g. sodium or potassium salts), alkaline earth metal salts (e.g. calcium or magnesium salts) or ammonium salts derived from ammonia or organic amines such as, for example, diethylamine, triethylamine, ethyldiisopropylamine, procaine, dibenzylamine, N-methylmorpholine, dihydroabietylamine or methylpiperidine.

[0141] "Hydrates" refers to those forms of the compounds of the above formula (I) which form a molecular compound (solvate) in the solid or liquid state through hydration with water. In the hydrates, the water molecules are attached through secondary valencies by intermolecular forces, in particular hydrogen bonds. Solid hydrates contain water as so-called water of crystallization in stoichiometric ratios, and the water molecules do not have to be equivalent in terms of their binding state. Examples of hydrates are sesquihydrates, monohydrates, dihydrates or trihydrates. Equally suitable are also the hydrates of salts of the compounds of the invention.

[0142] "Prodrugs" refers to those forms of the compounds of the above formula (I) which may themselves be biologically active or inactive but can be converted into the corre-

sponding biologically active form (for example metabolically, solvolytically or in another way).

[0143] Halogen is fluorine, chlorine, bromine and iodine. Chlorine or fluorine are preferred.

[0144] (C_1-C_8) -Alkyl is a straight-chain or branched alkyl radical having 1 to 8 carbon atoms. Examples which may be mentioned are: methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl and n-hexyl. The corresponding alkyl groups with fewer carbon atoms are derived analogously from this definition, such as, for example, (C_1-C_6) -alkyl and (C_1-C_4) -alkyl. It is generally true that (C_1-C_4) -alkyl is preferred.

[0145] The meaning of the corresponding constituent of other more complex substituents is also derived from this definition, such as, for example, in the case of alkylsulfonyl, hydroxyalkyl, hydroxyalkylcarbonyl, alkoxy-alkyl, alkoxy-carbonyl-alkyl, alkanoylalkyl, aminoalkyl or alkylaminoalkyl.

[0146] (C_3-C_7) -Cycloalkyl is a cyclic alkyl radical having 3 to 7 carbon atoms. Examples which may be mentioned are: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl. The corresponding cycloalkyl groups with fewer carbon atoms are derived analogously from this definition, such as, for example, (C_3-C_5) -cycloalkyl. Cyclopropyl, cyclopentyl and cyclohexyl are preferred.

[0147] The meaning of the corresponding constituent of other more complex substituents such as, for example, cycloalkanoyl is also derived from this definition.

[0148] (C_2-C_6) -Alkenyl is a straight-chain or branched enyl radical having 2 to 6 carbon atoms. A straight-chain or branched alkenyl radical having 2 to 4 carbon atoms is preferred. Examples which may be mentioned are: vinyl, allyl, isopropenyl and n-but-2-en-1-yl.

[0149] (C_1-C_8) -Alkoxy is a straight-chain or branched alkoxy radical having 1 to 8 carbon atoms. Examples which may be mentioned are: methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, tert-butoxy, n-pentoxy, n-hexoxy, n-heptoxy and n-octoxy. The corresponding alkoxy groups with fewer carbon atoms are derived analogously from this definition, such as, for example, (C_1-C_6) -alkoxy and (C_1-C_4) -alkoxy. It is generally true that (C_1-C_4) -alkoxy is preferred.

[0150] The meaning of the corresponding constituent of other more complex constituents such as, for example, alkoxy-alkyl, alkoxy-carbonyl-alkyl and alkoxy-carbonyl is also derived from this definition.

[0151] Mono- or di- (C_1-C_4) -alkylaminocarbonyl is an amino group which is linked via a carbonyl group and which has a straight-chain or branched or two identical or different straight-chain or branched alkyl substituents each having 1 to 4 carbon atoms. Examples which may be mentioned are: methylamino, ethylamino, n-propylamino, isopropylamino, t-butylamino, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N-methyl-N-n-propylamino, N-isopropyl-N-n-propylamino and N-t-butyl-N-methylamino.

[0152] (C_1-C_6) -Alkanoyl is a straight-chain or branched alkyl radical having 1 to 6 carbon atoms which has a double bonded oxygen atom in position 1 and is linked via position 1. Examples which may be mentioned are: formyl, acetyl, propionyl, n-butyryl, butyryl, pivaloyl, n-hexanoyl. The corresponding alkanoyl groups with fewer carbon atoms are derived analogously from this definition, such as, for

example, (C_1-C_5) -alkanoyl, (C_1-C_4) -alkanoyl and (C_1-C_3) -alkanoyl. It is generally true that (C_1-C_3) -alkanoyl is preferred.

[0153] The meaning of the corresponding constituent of other more complex constituents such as, for example, cycloalkanoyl and alkanoylalkyl is also derived from this definition.

[0154] (C_3-C_7) -Cycloalkanoyl is a cycloalkyl radical as defined above which has 3 to 7 carbon atoms and which is linked via a carbonyl group.

[0155] (C_1-C_6) -Alkanoyloxymethoxy is a straight-chain or branched alkanoyloxymethoxy radical having 1 to 6 carbon atoms. Examples which may be mentioned are: acetoxymethoxy, propionoxymethoxy, n-butyroxymethoxy, i-butyroxymethoxy, pivaloyloxymethoxy, n-hexanoyloxymethoxy. The corresponding alkanoyloxymethoxy groups with fewer carbon atoms, such as, for example, (C_1-C_3) -alkanoyloxymethoxy, are derived analogously from this definition. It is generally true that (C_1-C_3) -alkanoyloxymethoxy is preferred.

[0156] (C_6-C_{14}) -Aryl is an aromatic radical having 6 to 14 carbon atoms. Examples which may be mentioned are: phenyl, naphthyl, phenanthrenyl and anthracenyl. The corresponding aryl groups with fewer carbon atoms, such as, for example, (C_6-C_{10}) -aryl, are derived analogously from this definition. It is generally true that (C_6-C_{10}) -aryl is preferred.

[0157] The meaning of the corresponding constituent of other more complex constituents such as, for example, aryl-carbonyl is also derived from this definition.

[0158] (C_5-C_{10}) -Heteroaryl or a 5- to 10-membered aromatic heterocycle having up to 3 heteroatoms and/or hetero chain members from the series S, O, N and/or NO(N-oxide) is a mono- or bicyclic heteroaromatic system which is linked via a ring carbon atom of the heteroaromatic system, optionally also via a ring nitrogen atom of the heteroaromatic system. Examples which may be mentioned are: pyridyl, pyridyl N-oxide, pyrimidyl, pyridazinyl, pyrazinyl, thienyl, furyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl or isoxazolyl, indolizinyll, indolyl, benzo[b]thienyl, benzo[b]furyl, indazolyl, quinolyl, isoquinolyl, naphthyridinyl, quinazolinyl. The corresponding heterocycles with a smaller ring size such as, for example, 5- or 6-membered aromatic heterocycles are derived analogously from this definition. It is generally true that 5- or 6-membered aromatic heterocycles such as, for example, pyridyl, pyridyl N-oxide, pyrimidyl, pyridazinyl, furyl and thienyl are preferred.


[0159] The meaning of the corresponding constituent of other more complex constituents such as, for example, (C_5-C_{10}) -heteroarylcarbonyl is also derived from this definition.

[0160] A 3- to 9-membered saturated or partially unsaturated, mono- or bicyclic, optionally benzo-fused heterocycle having up to 3 heteroatoms and/or hetero chain members from the series S, SO, SO₂, N, NO(N-oxide) and/or O is a heterocycle which may comprise one or more double bonds, which may be mono- or bicyclic, in which a benzene ring may be fused to two adjacent ring carbon atoms, and which is linked via a ring carbon atom or a ring nitrogen atom. Examples which may be mentioned are: tetrahydrofuryl, pyrrolidinyl, pyrrolinyl, piperidinyl, 1,2-dihydropyridinyl, 1,4-dihydropyridinyl, piperazinyl, morpholinyl, morpholinyl N-oxide, thiomorpholinyl, azepinyl, 1,4-diazepinyl and cyclohexyl. Piperidinyl, morpholinyl and pyrrolidinyl are preferred.

[0161] The corresponding cyclic systems with a small ring size, such as, for example, 5- to 7-membered cyclic systems are derived analogously from this definition.

[0162] The compounds of the formula (I) can be prepared by either, in a process alternative,

[A] reacting compounds of the general formula (II)




(II)

[0163] in which

[0164] the radicals R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0165] with carboxylic acids of the general formula (III)



(III)


[0166] in which

[0167] the radical R¹ has the meaning indicated above,

[0168] or else with the corresponding carbonyl halides, preferably carbonyl chlorides,

[0169] or else with the corresponding symmetrical or mixed carboxylic anhydrides of the carboxylic acids of the general formula (III) defined above

[0170] in inert solvents, where appropriate in the presence of an activating or coupling reagent and/or of a base, to give compounds of the general formula (I)




(I)

[0171] in which

[0172] the radicals R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0173] or else in a process alternative

[B] converting compounds of the general formula (IV)




(IV)

[0174] in which

[0175] the radicals R¹, R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0176] with a suitable selective oxidizing agent in an inert solvent into the corresponding epoxide of the general formula (V)



(V)

[0177] in which

[0178] the radicals R¹, R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0179] and are reacted in an inert solvent, where appropriate in the presence of a catalyst, with an amine of the general formula (VI)




(VI)

[0180] in which

[0181] the radical R² has the meaning indicated above,

[0182] initially preparing the compounds of the general formula (VII)




(VII)

[0183] in which

[0184] the radicals R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0185] and

[0186] subsequently cyclizing in an inert solvent in the presence of phosgene or phosgene equivalents such as, for example, carbonyldiimidazole (CDI) to the compounds of the general formula (I)



(I)

[0187] in which

[0188] the radicals R¹, R², R³, R⁴, R⁵, R⁶, R⁷ and R⁸ have the meanings indicated above,

[0189] where, both for process alternative [A] and for process alternative [B] in the case where R² is a 3- to 7-membered saturated or partially unsaturated cyclic hydrocarbon radical having one or more identical or different heteroatoms from the group of N and S, it is

possible for an oxidation with a selective oxidizing agent to the corresponding sulfone, sulfoxide or N-oxide to follow,

[0190] and/or

[0191] where, both for process alternative [A] and for process alternative [B] in the case where the compound prepared in this way has a cyano group in the molecule, it is possible for an amidation of this cyano group by conventional methods to follow,

[0192] and/or

[0193] where, both for process alternative [A] and for process alternative [B] in the case where the compound prepared in this way has a BOC amino protective group in the molecule, it is possible for an elimination of this BOC amino protective group by conventional methods to follow,

[0194] and/or


[0195] where, both for process alternative [A] and for process alternative [B] in the case where the compound prepared in this way has an aniline or benzylamine residue in the molecule, it is possible for a reaction of this amino group with various reagents such as carboxylic acids, carboxylic anhydrides, carbonyl chlorides, isocyanates, sulfonyl chlorides or alkyl halides to give the corresponding derivatives to follow,

[0196] and/or



[0197] where, both for process alternative [A] and for process alternative [B] in the case where the compound prepared in this way has a phenyl ring in the molecule, it is possible for a reaction with chlorosulfonic acid and subsequent reaction with amines to give the corresponding sulfonamides to follow.

[0198] The processes can be illustrated by way of example by the following formula diagrams:


[A]



[A]



[0199] The oxidation step described above, which takes place where appropriate, can be illustrated by way of example by the following formula diagrams:



[0200] Solvents suitable for the processes described above are in these cases organic solvents which are inert under the reaction conditions. These include halohydrocarbons such as dichloromethane, trichloromethane, tetrachloromethane, 1,2-dichloroethane, trichloroethane, tetrachloroethane, 1,2-dichloroethylene or trichloroethylene, ethers such as diethyl ether, dioxane, tetrahydrofuran, glycol dimethyl ether or diethylene glycol dimethyl ether, alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol or tert-butanol, hydrocarbons such as benzene, xylene, toluene, hexane or cyclohexane, dimethylformamide, dimethyl sulfoxide, acetonitrile, pyridine, hexamethylphosphoric triamide or water.

[0201] It is likewise possible to employ solvent mixtures composed of the aforementioned solvents.

[0202] Activating or coupling reagents suitable for the processes described above are in these cases the reagents normally used for these purposes, for example N'-(3-dimethylaminopropyl)-N-ethylcarbodiimide.HCl, N,N'-dicyclohexylcarbodiimide, 1-hydroxy-1H-benzotriazole, H₂O and the like.

[0203] Suitable bases are the usual inorganic or organic bases. These preferably include alkali metal hydroxides such as, for example, sodium or potassium hydroxide or alkali metal carbonates such as sodium or potassium carbonate or sodium or potassium methanolate or sodium or potassium ethanolate or potassium tert-butoxide or amides such as sodamide, lithium bis-(trimethylsilyl)amide or lithium diisopropylamide or amines such as triethylamine, diisopropylethylamine, diisopropylamine, 4-N,N-dimethylaminopyridine or pyridine.

[0204] The base can be employed in these cases in an amount of from 1 to 5 mol, preferably from 1 to 2 mol, based on 1 mol of the compounds of the general formula (II).

[0205] The reactions generally take place in a temperature range from -78° C. to the reflux temperature, preferably in the range from 0° C. to the reflux temperature.

[0206] The reactions can be carried out under atmospheric, elevated or reduced pressure (e.g. in the range from 0.5 to 5 bar), generally under atmospheric pressure.

[0207] Suitable selective oxidizing agents both the preparing epoxides and for the oxidation which is optionally carried out to the sulfone, sulfoxide or N-oxide are, for example,

m-chloroperbenzoic acid (MCPBA), sodium metaperiodate, N-methylmorpholine N-oxide (NMO), monoperoxyphthalic acid or osmium tetroxide.

[0208] The conditions used for preparing the epoxides are those customary for these preparations.

[0209] For detailed conditions for the process of oxidation, which is carried out where appropriate, to the sulfone, sulfoxide or N-oxide, reference may be made to the following literature: M. R. Barbachyn et al., J. Med. Chem. 1996, 39, 680 and WO-A-97/10223.

[0210] Reference is further made to Examples 14 to 16 detailed in the experimental part.

[0211] The amidation which is carried out where appropriate takes place under the usual conditions. For further details, reference may be made to Examples 31 to 35 and 140 to 147.

[0212] The compounds of the formulae (II), (III), (IV) and (VI) are known per se to the skilled worker or can be prepared by conventional methods. For oxazolidinones, in particular the 5-(aminomethyl)-2-oxoxazolidinones required, cf. WO-A-98/01446; WO-A-93/23384; WO-A-97/03072; J. A. Tucker et al., J. Med. Chem. 1998, 41, 3727; S. J. Brickner et al., J. Med. Chem. 1996, 39, 673; W. A. Gregory et al., J. Med. Chem. 1989, 32, 1673.

[0213] A preferred compound A) of the formula (I) for use in combinations is 5-chloro-N-({(5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide, the compound of Example 44.

[0214] The combinations of the invention are suitable in particular for the prophylaxis and/or treatment of arterial thromboses and embolisms associated with coronary heart disease, impairments of cerebrovascular blood flow and impairments of peripheral arterial blood flow. Combinations of oxazolidinones of the formula (I) with platelet aggregation inhibitors, anticoagulants and/or fibrinolytics are additionally suitable in particular for the prophylaxis and/or treatment of venous thromboses and pulmonary embolisms.

[0215] The individual active ingredients of the combinations are known from the literature and for the most part commercially available. They may, where appropriate, just like the oxazolidinones of the formula (I), be employed in subtherapeutically effective doses.

[0216] Suitable for the prophylaxis and/or treatment of arterial vascular disorders is a combination therapy of oxazolidinones of the formula (I) with lipid-lowering agents, in particular with HMG-CoA (3-hydroxy-3-methylglutaryl-coenzyme A) reductase inhibitors such as, for example, cerivastatin (Rivastatin, Baycol; U.S. Pat. No. 5,177,080), lovastatin (Mevacor; U.S. Pat. No. 4,231,938), simvastatin (Zocor; U.S. Pat. No. 4,444,784), pravastatin (Pravachol; U.S. Pat. No. 4,346,227), fluvastatin (Lescol; U.S. Pat. No. 5,354,772), atorvastatin (Lipitor; U.S. Pat. No. 5,273,995), or with coronary therapeutic agents/vasodilators, in particular ACE (angiotensin converting enzyme) inhibitors, such as, for example, captopril, lisinopril, enalapril, ramipril, cilazapril, benazepril, fosinopril, quinapril, perindopril; AII (angiotensin II) receptor antagonists such as, for example, embusartan (U.S. Pat. No. 5,863,930), losartan, valsartan, irbesartan, candesartan, eprosartan, temisartan; β -adrenoceptor antagonists such as, for example, carvedilol, alprenolol, bisoprolol, acebutolol, atenolol, betaxolol, carteolol, metoprolol, nadolol, penbutolol, pindolol, propranolol, timolol; alpha-1-adrenoceptor antagonists such as, for example, prazosin, bunazosin, doxazosin, terazosin; diuretics such as, for example, hydrochlorothiazide, furosemide, bumetanide, piretanide, torasemide, amiloride; dihydralazine; calcium channel blockers such as, for example, verapamil, diltiazem or dihydropyridine derivatives such as, for example, nifedipine (Adalat) or nitrendipine (Bayotensin); substances which bring about an increase in cyclic guanosine monophosphate (cGMP), such as, for example, stimulators of soluble guanylate cyclase (WO 98/16223, WO 98/16507, WO 98/23619, WO 00/06567, WO 00/06568, WO 00/06569, WO 00/21954, WO 00/66582, WO 01/17998, WO 01/19776, WO 01/19355, WO 01/19780, WO 01/19778).

[0217] The pharmacotherapeutic aim of treatment of a pre-existing coronary heart disease is to eliminate the imbalance between oxygen supply and oxygen demand in the areas of myocardium affected by the ischemic. Suitable for the treatment of a pre-existing coronary heart disease is therefore in particular combination therapy of an oxazolidinone of the formula (I) with coronary therapeutic agents, in particular with β -adrenoceptor antagonists; ACE (angiotensin converting enzyme) inhibitors; A-II (angiotensin II) receptor antagonists; nitrates such as, for example, isosorbide 5-mononitrate, isosorbide dinitrate, glycerol trinitrate; substances which bring about an increase in cyclic guanosine monophosphate (cGMP); calcium channel blockers. Most of these compounds are also employed for therapy of hypertension.

[0218] Thrombolytic therapy with plasminogen aktivators (thrombolytic/fibrinolytic agents) such as, for example, tissue plasminogen activator (t-PA), streptokinase, reteplase or urokinase has proved effective for reopening thrombotically occluded vessels. However, administration of plasminogen activators on their own does not prevent further growth of the thrombus. High doses of plasminogen activators may additionally mean an increased risk of bleeding. Combined administration of a thrombolytic agent with an oxazolidinone of the formula (I) for opening thrombotically occluded vessels associated with coronary heart disease, transient ischemic attacks, stroke, peripheral arterial occlusive diseases and pulmonary embolisms prevents further growth of the thrombus through inhibition of thrombin formation and thus reduces the risk of reocclusion. In addition, on combination therapy with a thrombolytic agent and an oxazolidinone of the formula (I) it is possible to reduce the dose of

thrombolytic agent necessary for therapy, which leads to a reduction in the bleeding complications and thus represents a considerable advantage over monotherapy.

[0219] Oxazolidinones of the formula (I) can also be given in combination with other substances with anticoagulant activity (anticoagulants) for the prophylaxis and/or treatment of arterial, intracardiac and venous thromboembolic disorders.

[0220] Combination therapy of oxazolidinones of the formula (I) in particular with heparin (UFH), lower molecular weight heparins (LMWH) such as, for example, tinzaparin, certoparin, parnaparin, nadroparin, ardeparin, enoxaparin, reviparin, dalteparin or direct thrombin inhibitors such as, for example, hirudin leads to an enhanced antithrombotic effect.

[0221] Oxazolidinones of the formula (I) can additionally also be given in combination with platelet aggregation-inhibiting substances (platelet aggregation inhibitors) for the prophylaxis and/or treatment of arterial, intracardiac and venous thromboembolic disorders. An endothelial lesion is associated with adhesion to the wall and activation of blood platelets and simultaneous stimulation of coagulation. This leads to the formation of platelet- and fibrin-containing thrombi, and the platelets contribute to stabilizing the fibrin matrix (J. Hirsh, E. W. Salzman, V. J. Marder, R. W. Colman, Overview of the Thrombotic Process and its Therapy, pages 1151-1163 in Hemostasis and Thrombosis: Basic Principles and Clinical Practice, Third Edition, edited by R. W. Colman, J. Hirsh, V. J. Marder, E. W. Salzman. J. B. Lippincott Company, Philadelphia, 1994). Simultaneous inhibition of coagulation and of platelet aggregation therefore leads to an enhanced anti-thrombotic effect. Particularly suitable for combination therapy are combinations of an oxazolidinone of the formula (I) with platelet aggregation inhibitors such as, for example, aspirin, ticlopidin (Ticlid), clopidogrel (Plavix); fibrinogen receptor antagonists; (glycoprotein antagonists) such as, for example, abciximab, eptifibatide, tirofiban, lamifiban, lefradafiban.

[0222] All usual administration forms are suitable for administering the combinations of the invention. Administration preferably takes place orally, lingually, sublingually, buccally, rectally, topically or parenterally (i.e. avoiding the intestinal tract, i.e. intravenous, intraarterial, intracardiac, intracutaneous, subcutaneous, transdermal, intraperitoneal or intramuscular).

[0223] The present invention includes pharmaceutical preparations which, besides non-toxic, inert pharmaceutically suitable excipients and/or carriers, comprise one or more combinations of the invention or which consist of a combination of the invention, and processes for producing these preparations.

[0224] The combinations of the invention are intended to be present in the above-mentioned pharmaceutical preparations in a concentration of about 0.1 to 99.5, preferably about 0.5 to 95, % by weight of the complete mixture.

[0225] The abovementioned pharmaceutical preparations may, besides the combinations of the invention, also comprise further active pharmaceutical ingredients.

[0226] The abovementioned pharmaceutical preparations can be produced in a conventional way by known methods, e.g. by mixing the active ingredient or active ingredients with the carrier(s).

[0227] It has generally proved advantageous to administer the combinations of the invention in total amounts of about 0.001 to 100 mg/kg, preferably about 0.01 to 100 mg/kg, in

particular about 0.1 to 10 mg/kg, of body weight every 24 hours, where appropriate in the form of a plurality of single doses, to achieve the desired results.

[0228] It may nevertheless be necessary where appropriate to depart from the aforementioned amounts, in particular depending on the body weight, on the nature of the administration route, the type and severity of the disorder, on the individual behavior toward the medicament, on the nature of the formulation and on the time or interval over which administration takes place. Thus, it may be sufficient in some cases to make do with less than the aforementioned minimum amount, whereas in other cases the upper limit mentioned must be exceeded. It may be advisable, for example when relatively large amounts are administered, to distribute these over the day, in particular either in a plurality of single doses or as continuous infusion.

[0229] The invention therefore further relates to the combinations defined above for the prophylaxis and/or treatment of disorders.

[0230] The invention further relates to medicaments comprising at least one of the combinations defined above and, where appropriate, further active pharmaceutical ingredients.

[0231] The invention further relates to the use of the combinations defined above for producing medicaments for the prophylaxis and/or treatment of the disorders described above, preferably thromboembolic disorders, in particular myocardial infarction, angina pectoris (including unstable angina), sudden heart death, reocclusions and restenoses after an angioplasty or aortocoronary bypass, stroke, transient ischemic attacks, peripheral arterial occlusive diseases, pulmonary embolisms or deep venous thromboses.

[0232] The percentage data in the following examples are based in each case on weight; parts are parts by weight.

EXAMPLES

A Assessment of the Physiological Activity

1. Physiological Activity of Compounds of the Formula (I)

[0233] The compounds of the formula (I) act in particular as selective inhibitors of coagulation factor Xa and do not inhibit, or also inhibit only at distinctly higher concentrations, other serine proteases such as thrombin, plasmin or trypsin.

[0234] Inhibitors of coagulation factor Xa are referred to as "selective" when their IC_{50} values for factor Xa inhibition are 100-fold, preferably 500-fold, in particular 1000-fold, smaller than the IC_{50} values for the inhibition of other serine proteases, in particular thrombin, plasmin and trypsin, reference being made concerning the test methods for the selectivity to the test methods of Examples A-1) a.1) and a.2) described below.

[0235] The particularly advantageous biological properties of the compounds of the formula (I) can be ascertained by the following methods.

a) Test Description (In Vitro)

a.1) Measurement of Factor Xa Inhibition

[0236] The enzymatic activity of human factor Xa (FXa) was measured via the conversion of an FXa-specific chromogenic substrate. In this case, factor Xa eliminates p-nitroaniline from the chromogenic substrate. The determinations were carried out in microtiter plates as follows.

[0237] The test substances were dissolved in various concentrations in DMSO and incubated with human FXa (0.5

nmol/l dissolved in 50 mmol/l tris buffer [C,C,C-tris(hydroxymethyl)-aminomethane], 150 mmol/l NaCl, 0.1% BSA (bovine serum albumine), pH=8.3) at 25° C. for 10 minutes. Pure DMSO serves as control. The chromogenic substrate (150 μ mol/l Pefachrome® FXa from Pentapharm) was then added. After incubation at 25° C. for 20 minutes, the extinction at 405 nm was determined. The extinctions of the test mixtures with test substance were compared with the control mixtures without test substance, and the IC_{50} values were calculated therefrom.

a.2) Selectivity Determination

[0238] Selective FXa inhibition was demonstrated by investigating the inhibition by the test substances of other human serine proteases such as thrombin, trypsin, plasmin. The enzymatic activity of thrombin (75 mU/ml), trypsin (500 mU/ml) and plasmin (3.2 nmol/l) was determined by dissolving these enzymes in tris buffer (100 mmol/l, 20 mmol/l $CaCl_2$, pH=8.0) and incubating with test substance or solvent for 10 minutes. The enzymatic reaction was then started by adding the appropriate specific chromogenic substrates (Chromozym Thrombin® from Boehringer Mannheim, Chromozym Trypsin® from Boehringer Mannheim, Chromozym Plasmin® from Boehringer Mannheim), and the extinction was determined at 405 nm after 20 minutes. All determinations were carried out at 37° C. The extinctions of the test mixtures with test substance were compared with the control samples without test substance, and the IC_{50} values were calculated therefrom.

a.3) Determination of the Anticoagulant Effect

[0239] The anticoagulant effect of the test substances was determined in vitro in human plasma. For this purpose, human blood was collected in a 0.11 molar sodium citrate solution in the sodium citrate/blood mixing ratio of 1/9. The blood was thoroughly mixed after collection and centrifuged at about 2000 g for 10 minutes. The supernatant was removed by pipette. The prothrombin time (PT, synonym: Quick's test) was determined in the presence of varying concentrations of test substance or the appropriate solvent using a commercially available test kit (Neoplastin® from Boehringer Mannheim). The test compounds were incubated with the plasma at 37° C. for 10 minutes. Coagulation was then induced by adding thromboplastin, and the time of onset of coagulation was determined. The concentration of test substance which brings about a doubling of the prothrombin time was found.

b) Determination of the Antithrombotic Effect (In Vivo)

b.1) Arteriovenous Shunt Model (Rat)

[0240] Fasting male rats (strain: HSD CPB:WU) weighing 200-250 g were anesthetized with a Rompun/Ketavet solution (12 mg/kg/50 mg/kg). Thrombus formation was induced in an arteriovenous shunt by a method based on that described by Christopher N. Berry et al., Br. J. Pharmacol. (1994), 113, 1209-1214. For this purpose, the left jugular vein and the right carotid artery were exposed. An extracorporeal shunt was formed between the two vessels using a 10 cm-long polyethylene tube (PE 60). This polyethylene tube was secured in the middle by tying in a further 3 cm-long polyethylene tube (PE 160) which contained a roughened nylon thread forming a loop to produce a thrombogenic surface. The extracorporeal circulation was maintained for 15 minutes. The shunt was

then removed and the nylon thread with the thrombus was immediately weighed. The blank weight of the nylon thread had been found before the start of the experiment. The test substances were administered either intravenously through the tail vein or orally by gavage to conscious animals before setting up the extracorporeal circulation. The results are shown in Table 1:

TABLE 1

Antithrombotic effect in the arteriovenous shunt model (rat) after oral or intravenous administration		
Example	ED ₅₀ [mg/kg] p.o.	ED ₅₀ [mg/kg] i.v.
1		10
17		6
44	3	
95		3
114		3
115		3
123	3	
162		3

b.2) Arterial Thrombosis Model (Rat)

[0241] Fasting male rats (strain: HSD CPB: WU) were anesthetized as described above. The rats had an average weight of about 200 g. The left carotid artery was exposed (about 2 cm). Formation of an arterial thrombus was induced by mechanical injury to the vessel by a method based on that described by K. Meng et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.* (1977), 301, 115-119. For this purpose, the exposed carotid artery was clamped off from the blood flow, cooled to -12° C. in a metal channel for 2 minutes and, to standardize the thrombus size, simultaneously compressed with a weight of 200 g. The blood flow was then additionally reduced by a clip placed around the carotid artery distal from the injured section of vessel. The proximal clamp was removed, and the wound was closed and reopened after 4 hours in order to remove the injured section of vessel. The section of vessel was opened longitudinally and the thrombus was removed from the injured section of vessel. The wet weight of the thrombi was measured immediately. The test substances were administered either intravenously via the tail vein or orally by gavage to conscious animals at the start of the experiment.

b.3) Venous Thrombosis Model (Rat)

[0242] Fasting male rats (strain: HSD CPB: WU) were anesthetized as described above. The rats had an average weight of about 200 g. The left jugular vein was exposed (about 2 cm). Formation of a venous thrombus was induced by mechanical injury to the vessel by a method based on that described by K. Meng et al., *Naunyn-Schmiedeberg's Arch. Pharmacol.* (1977), 301, 115-119. For this purpose, the exposed jugular vein was clamped off from the blood flow, cooled to -12° C. in a metal channel for 2 minutes and, to standardize the thrombus size, simultaneously compressed with a weight of 200 g. The blood flow was reopened and the wound was closed. After 4 hours, the wound was reopened in order to remove the thrombi from the injured sections of vessels. The wet weight of the thrombi was measured immediately. The test substances were administered either intrave-

nously via the tail vein or orally by gavage to conscious animals at the start of the experiment.

2. Physiological Activity of the Combinations of Compounds of the Formula (I)

a) In Vivo Investigations in a Rat Thrombosis Model

[0243] The carotid artery of rats (HSD CPB:WU, Harlan Winkelmann) was exposed under anesthesia. A piece of filter paper impregnated with an aqueous 10% strength FeCl_3 solution (dissolved in 1N aqueous hydrochloric acid) was cautiously pushed underneath the exposed vessel in accordance with the method described by Kurz et al. (Rat Model of Arterial Thrombosis Induced by Ferric Chloride, *Thrombosis Research* 60, 269-280, 1990). After 3 minutes, the piece of filter paper was removed. The carotid artery was removed after 15 minutes, and the thrombus was detached and immediately weighed. The animals (10 rats per group) had been pretreated with 1 mg/kg of each of the individual active ingredients (oxazolidinone of the formula (I) and combination active ingredient) or with the combination of 1 mg/kg oxazolidinone of the formula (I) and 1 mg/kg combination active ingredient. The animals in the control group had been treated with the corresponding solvent. The statistical significance was calculated using Student's t-test. Values with $p < 0.05$ are regarded as a statistically significant effect (Medical Statistics, M.J Campbell, D. Machin, Second Edition, John Wiley & Sons). The results are shown in Table 2:

TABLE 2

Synergistic antithrombotic effect of the combination of an oxazolidinone of the formula (I) with a platelet aggregation inhibitor Reduction in thrombus weight after oral treatment with		
Compound of Example 44 [1 mg/kg]	Clopidogrel [1 mg/kg]	Combination of the compound of Example 44 [1 mg/kg] with clopidogrel [1 mg/kg]
22%	28%	39%
no effect ($p > 0.05$)	no effect ($p > 0.05$)	effect ($p < 0.05$)

[0244] As shown in Table 2, a synergistic effect is achieved with the combination of an oxazolidinone of the formula (I) such as the compound of Example 44 with a platelet aggregation inhibitor such as clopidogrel, i.e. the two components mutually enhance their effect. In single dosage, both compounds were inactive at the dose investigated. Combination of the two compounds by contrast led to a significant reduction in the thrombus weight. Combination of oxazolidinones of the formula (I) with a platelet aggregation-inhibiting substance is therefore able to improve antithrombotic therapy considerably.

B Preparation examples

Starting Compounds


[0245] The preparation of 3-morpholinone is described in U.S. Pat. No. 5,349,045.

[0246] The preparation of N-(2,3-epoxypropyl)phthalimide is described in J.-W. Chern et al. *Tetrahedron Lett.* 1998, 39,8483.

[0247] The substituted anilines can be obtained by reacting, for example, 4-fluoronitrobenzene, 2,4-difluoronitrobenzene

or 4-chloronitrobenzene with the appropriate amines or amides in the presence of a base. This can also take place with use of Pd catalysts such as Pd(OAc)₂/DPPF/NaOt-Bu (Tetrahedron Lett. 1999, 40,2035) or copper (Renger, Synthesis 1985,856; Aebischer et al., Heterocycles 1998, 48,2225). Haloaromatic compounds without a nitro group can initially be converted into the corresponding amides in exactly the same way in order to be subsequently nitrated in position 4 (U.S. Pat. No. 3,279,880).

I. 4-(4-Morpholin-3-onyl)nitrobenzene



2 mol (202 g) of morpholin-3-one (E. Pfeil, U. Harder, Angew. Chem. 79, 1967, 188) are dissolved in 21 of N-methylpyrrolidone (NMP). 88 g (2.2 mol) of sodium hydride (60% in paraffin) are then added in portions over a period of 2 h. After hydrogen evolution ceases, 282 g (2 mol) of 4-fluoronitrobenzene are added dropwise while cooling at room temperature over the course of 1 h, and the reaction mixture is then stirred overnight. Subsequently, 1.7 l of the liquid volume are distilled out at 12 mbar and 76° C., the residue is poured into 21 of water, and this mixture is extracted twice with 1 l of ethyl acetate each time. The combined organic phases are washed with water and then dried over sodium sulfate, and the solvent is distilled off in vacuo. Purification takes place by chromatography on silica gel with hexane/ethyl acetate (1:1) and subsequent crystallization from ethyl acetate. The product is obtained as 78 g of a colorless to brownish solid in 17.6% of theory.


[0248] ¹H-NMR (300 MHz, CDCl₃): 3.86 (m, 2H, CH₂CH₂), 4.08 (m, 2H, CH₂CH₂), 4.49 (s, 2H, CH₂CO), 7.61 (d, 2H, ³J=8.95 Hz, CHCH), 8.28 (d, 2H, ³J=8.95 Hz, CHCH)

[0249] MS (r.l. %)=222 (74, M⁺), 193 (100), 164 (28), 150 (21), 136 (61), 117 (22), 106 (24), 90 (37), 76 (38), 63 (32), 50 (25)

[0250] The following compounds were synthesized analogously:

- [0251]** 3-fluoro-4-(4-morpholin-3-onyl)nitrobenzene
- [0252]** 4-(N-piperidonyl)nitrobenzene
- [0253]** 3-fluoro-4-(N-piperidonyl)nitrobenzene
- [0254]** 4-(N-pyrrolidonyl)nitrobenzene
- [0255]** 3-fluoro-4-(N-pyrrolidonyl)nitrobenzene

II. 4-(4-Morpholin-3-onyl)aniline



[0256] 63 g (0.275 mol) of 4-(4-morpholin-3-onyl)nitrobenzene are dissolved in 200 ml of tetrahydrofuran in an autoclave, 3.1 g of Pd/C (5%) are added, and the mixture is hydrogenated under a hydrogen pressure of 50 bar at 70° C. for 8 h. After filtration of the catalyst, the solvent is distilled out in vacuo and the product is purified by crystallization from ethyl acetate. The product is obtained as 20 g of a colorless to blueish solid in 37.6% of theory.

[0257] Purification can also take place by chromatography on silica gel with hexane/ethyl acetate.

[0258] ¹H-NMR (300 MHz, CDCl₃): 3.67 (m, 2H, CH₂CH₂), 3.99 (m, 2H, CH₂CH₂), 4.27 (s, 2H, CH₂CO), 6.68 (d, 2H, ³J=8.71 Hz, CHCH), 7.03 (d, 2H, ³J=8.71 Hz, CHCH)

[0259] MS (r.l. %)=192 (100, M⁺), 163 (48), 133 (26), 119 (76), 106 (49), 92 (38), 67 (27), 65 (45), 52(22), 28 (22)

[0260] The following compounds were synthesized analogously:


[0261] 3-fluoro-4-(4-morpholin-3-onyl)aniline

[0262] 4-(N-piperidonyl)aniline

[0263] 3-fluoro-4-(N-piperidonyl)aniline

[0264] 4-(N-pyrrolidonyl)aniline 3-fluoro-4-(N-pyrrolidonyl)aniline

General Method For Preparing 4-Substituted Anilines by Reacting 1-Fluoro-4-Nitrobenzenes and 1-Chloro-4-Nitrobenzenes with Primary or Secondary Amines And Subsequent Reduction



[0265] Equimolar amounts of the fluoronitrobenzene or chloronitrobenzene and of the amine are dissolved in dim-

ethyl sulfoxide or acetonitrile (0.1 M to 1 M solution) and stirred at 100° C. overnight. After cooling to RT, the reaction mixture is diluted with ether and washed with water. The organic phase is dried over MgSO₄, filtered and concentrated. If a precipitate is obtained in the reaction mixture, it is filtered off and washed with ether or acetonitrile. If product is also to be found in the mother liquor, this is worked up with ether and water as described. The crude products can be purified by chromatography on silica gel (dichloromethane/cyclohexane and dichloromethane/ethanol mixtures).

[0266] For the subsequent reduction, the nitro compound is dissolved in methanol, ethanol or ethanol/dichloromethane mixtures (0.01 M to 0.5 M solution), mixed with palladium on carbon (10%) and stirred under hydrogen of atmospheric pressure overnight. This is followed by filtration and concentration. The crude product can be purified by chromatography on silica gel (dichloromethane/ethanol mixtures) or preparative reversed phase HPLC (acetonitrile/water mixtures).

[0267] Alternatively, iron powder can also be used as reducing agent. For this purpose, the nitro compound is dissolved in acetic acid (0.1 M to 0.5 M solution) and, at 90° C., six equivalents of iron powder and water (0.3 to 0.5 times the volume of acetic acid) are added in portions over the course of 10-15 min. After a further 30 min at 90° C., the mixture is filtered and the filtrate is concentrated. The residue is worked up by extraction with ethyl acetate and 2N sodium hydroxide solution. The organic phase is dried over magnesium sulfate, filtered and concentrated. The crude product can be purified by chromatography on silica gel (dichloromethane/ethanol mixtures) or preparative reversed phase HPLC (acetonitrile/water mixtures).

[0268] The following starting compounds were prepared in an analogous manner:

III-1. Tert-butyl 1-(4-aminophenyl)-L-prolinate

[0269] MS (ESI): m/z (%)=304 (M+H+MeCN, 100), 263 (M+H, 20);

[0270] HPLC (method 4): rt=2.79 min.

III-2. 1-(4-Aminophenyl)-3-piperidinecarboxamide

[0271] MS (ESI): m/z (%)=220 (M+H, 100);

[0272] HPLC (method 4): rt=0.59 min.

III-3. 1-(4-Aminophenyl)-4-piperidinecarboxamide

[0273] MS (ESI): m/z (%)=220 (M+H, 100);

[0274] HPLC (method 4): rt=0.57 min.

III-4. 1-(4-Aminophenyl)-4-piperidinone

[0275] MS (ESI): m/z (%)=191 (M+H, 100);

[0276] HPLC (method 4): rt=0.64 min.

III-5. 1-(4-Aminophenyl)-L-prolinamide

[0277] MS (ESI): (%)=206 (M+H, 100);

[0278] HPLC (method 4): rt=0.72 min.

III-6. [1-(4-Aminophenyl)-3-piperidinyl]methanol

[0279] MS (ESI): m/z (%)=207 (M+H, 100);

[0280] HPLC (method 4): rt=0.60 min.

III-7. [1-(4-Aminophenyl)-2-piperidinyl]methanol

[0281] MS (ESI): m/z (%)=207 (M+H, 100);

[0282] HPLC (method 4): rt=0.59 min.

III-8. Ethyl 1-(4-aminophenyl)-2-piperidinecarboxylate

[0283] MS (ESI): m/z (%)=249 (M+H, 35), 175 (100);

[0284] HPLC (method 4): rt=2.43 min

III-9. [1-(4-Aminophenyl)-2-pyrrolidinyl]methanol

[0285] MS (ESI): m/z (%)=193 (M+H, 45);

[0286] HPLC (method 4): rt=0.79 min.

III-10. 4-(2-Methylhexahydro-5H-pyrrolo[3,4-d]isoxazol-5-yl)phenylamine

[0287] starting from 2-methylhexahydro-2H-pyrrolo[3,4-d]isoxazole (Ziegler, Carl B., et al.; J. Heterocycl. Chem., 25; 2; 1988; 719-723)

[0288] MS (ESI): m/z (%)=220 (M+H, 50), 171 (100);

[0289] HPLC (method 4): rt=0.54 min.

III-11. 4-(1-Pyrrolidinyl)-3-(trifluoromethyl) aniline

[0290] MS (ESI): m/z (%)=231 (M+H, 100);

[0291] HPLC (method 7): rt=3.40 min.

III-12. 3-Chloro-4-(1-pyrrolidinyl)aniline

[0292] MS (ESI): m/z (%)=197 (M+H, 100);

[0293] HPLC (method 4): rt=0.78 min

III-13. 5-Amino-2-(4-morpholinyl)benzamide

[0294] MS (ESI): m/z (%)=222 (M+H, 100);

[0295] HPLC (method 4): rt=0.77 min.

III-14. 3-Methoxy-4-(4-morpholinyl)aniline

[0296] MS (ESI): m/z (%)=209 (M+H, 100);


[0297] HPLC (method 4): rt=0.67 min.

III-15. 1-[5-Amino-2-(4-morpholinyl)phenyl]ethanone

[0298] MS (ESI): m/z (%)=221 (M+H, 100);

[0299] HPLC (method 4): rt=0.77 min.

General Method for Preparing 4-Substituted Anilines by Reacting 1-Fluoro-4-Nitrobenzenes with Amides and Subsequent Reduction



[0300] The Amide is Dissolved in DMF, and 1.5 Equivalents of Potassium Tert-Butoxide Are added. The mixture is stirred at RT for 1 h, and then 1.2 equivalents of the 1-fluoro-4-nitrobenzene are added in portions. The reaction mixture is stirred at RT overnight, diluted with ether or ethyl acetate and washed with saturated aqueous sodium bicarbonate solution. The organic phase is dried over magnesium sulfate, filtered and concentrated. The crude product can be purified by chromatography on silica gel (dichloromethane/ethanol mixtures).

[0301] For the subsequent reduction, the nitro compound is dissolved in ethanol (0.01 M to 0.5 M solution), mixed with palladium on carbon (10%) and stirred under hydrogen of atmospheric pressure overnight. This is followed by filtration and concentration. The crude product can be purified by chro-

matography on silica gel (dichloromethane/ethanol mixtures) or preparative reversed phase HPLC (acetonitrile/water mixtures).

[0302] Alternatively, iron powder can also be used as reducing agent. For this purpose, the nitro compound is dissolved in acetic acid (0.1 M to 0.5 M solution) and, at 90° C., six equivalents of iron powder and water (0.3 to 0.5 times the volume of acetic acid) are added in portions over the course of 10-15 min. After a further 30 min at 90° C., the mixture is filtered and the filtrate is concentrated. The residue is worked up by extraction with ethyl acetate and 2N sodium hydroxide solution. The organic phase is dried over magnesium sulfate, filtered and concentrated. The crude product can be purified by chromatography on silica gel (dichloromethane/ethanol mixtures) or preparative reversed phase HPLC (acetonitrile/water mixtures).

[0303] The following starting compounds were prepared in an analogous manner:

IV-1. 1-[4-Amino-2-(trifluoromethyl)phenyl]-2-pyrrolidinone

[0304] MS (ESI): m/z (%)=245 (M+H, 100);

[0305] HPLC (method 4): rt =2.98 min

IV-2. 4-(4-Amino-2-(trifluoromethyl)phenyl)-3-morpholinone

[0306] MS (ESI): m/z (%)=261 (M+H, 100);

[0307] HPLC (method 4): rt =2.54 min.

IV-3. 4-(4-Amino-2-chlorophenyl)-3-morpholinone

[0308] MS (ESI): m/z (%)=227 (M+H, 100);

[0309] HPLC (method 4): rt =1.96 min.

IV-4. 4-(4-Amino-2-methylphenyl)-3-morpholinone

[0310] MS (ESI): m/z (%)=207 (M+H, 100);

[0311] HPLC (method 4): rt =0.71 min.

IV-5. 5-Amino-2-(3-oxo-4-morpholinyl)benzonitrile

[0312] MS (ESI): m/z (%)=218 (M+H, 100);

[0313] HPLC (method 4): rt =1.85 min.

IV-6. 1-(4-Amino-2-chlorophenyl)-2-pyrrolidinone

[0314] MS (ESI): m/z (%)=211 (M+H, 100);

[0315] HPLC (method 4): rt =2.27 min.

IV-7. 4-(4-Amino-2,6-dimethylphenyl)-3-morpholinone

[0316] starting from 2-fluoro-1,3-dimethyl-5-nitrobenzene (Bartoli et al., J. Org. Chem. 1975, 40, 872):

[0317] MS (ESI): m/z (%)=221 (M+H, 100);

[0318] HPLC (method 4): rt =0.77 min.

IV-8. 4-(2,4-Diaminophenyl)-3-morpholinone

[0319] starting from 1-fluoro-2,4-dinitrobenzene:

[0320] MS (ESI): m/z (%)=208 (M+H, 100);

[0321] HPLC (method 4): rt =0.60 min.

IV-9. 4-(4-Amino-2-chlorophenyl)-2-methyl-3-morpholinone

[0322] starting from 2-methyl-3-morpholinone (Pfeil, E.; Harder, U.; Angew. Chem. 1967, 79, 188):

[0323] MS (ESI): m/z (%)=241 (M+H, 100);

[0324] HPLC (method 4): rt =2.27 min

IV-10. 4-(4-Amino-2-chlorophenyl)-6-methyl-3-morpholinone

[0325] starting from 6-methyl-3-morpholinone (EP 350 002):

[0326] MS (ESI): m/z (%)=241 (M+H, 100);

[0327] HPLC (method 4): rt =2.43 min.


Synthesis Examples

[0328] The following Examples 1 to 13, 17 to 19 and 36 to 57 relate to process variant [A].

Example 1

Preparation of 5-chloro-N-[[[(5S)-3-(3-fluoro-4-morpholinophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]-2-thiophenecarboxamide

[0329]



[0330] (5S)-5-(Aminomethyl)-3-(3-fluoro-4-morpholinophenyl)-1,3-oxazolidin-2-one (for preparation, see S. J. Brickner et al., J. Med. Chem. 1996, 39, 673) (0.45 g, 1.52 mmol), 5-chlorothiophene-2-carboxylic acid (0.25 g, 1.52 mmol) and 1-hydroxy-1H-benzotriazole hydrate (HOBT) (0.3 g, 1.3 equivalents) are dissolved in 9.9 ml of DMF. 0.31 g (1.98 mmol, 1.3 equivalents) of N¹-(3-dimethylaminopropyl)-N-ethylcarbodiimide (EDCI) is added and, at room temperature, 0.39 g (0.53 ml, 3.05 mmol, 2 equivalents) of diisopropylethylamine (DIEA) is added dropwise. The mixture is stirred at room temperature overnight. 2 g of silica gel are added and the mixture is evaporated to dryness in vacuo. The residue is chromatographed on silica gel with a toluene/ethyl acetate gradient. 0.412 g (61.5% of theory) of the target compound is obtained with a melting point (m.p.) of 197° C.

[0331] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.29 (precursor=0.0);


[0332] MS (DCI) 440.2 (M+H), Cl pattern;

[0333] ¹H-NMR (d₆-DMSO, 300 MHz) 2.95 (m, 4H), 3.6 (t, 2H), 3.72 (m, 4H), 3.8 (dd, 1H), 4.12 (t, 1H), 4.75-4.85 (m, 1H), 7.05 (t, 1H), 7.15-7.2 (m, 3H), 7.45 (dd, 1H), 7.68 (d, 1H), 8.95 (t, 1H).

Example 2

5-Chloro-N-[[[(5S)-3-(4-morpholinophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]-2-thiophenecarboxamide

[0334]



is obtained analogously from benzyl 4-morpholinophenyl-carbamate via the stage of (5S)-5-(aminomethyl)-3-(3-fluoro-4-morpholinophenyl)-1,3-oxazolidin-2-one (see Example 1).

[0335] M.p.: 198° C.;


[0336] IC₅₀=43 nM;

[0337] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.24.

Example 3

5-Chloro-N-({(5S)-3-[3-fluoro-4-(1,4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0338]



is obtained analogously from (5S)-5-(aminomethyl)-3-[3-fluoro-4-(1,4-thiazinan-4-yl)phenyl]-1,3-oxazolidin-2-one (for preparation, see M. R. Barbachyn et al., J. Med. Chem. 1996, 39, 680).

[0339] M.p.: 193° C.;


[0340] Yield: 82%;

[0341] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.47 (precursor=0.0).

Example 4

5-Bromo-N-({(5S)-3-[3-fluoro-4-(1,4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0342]




is obtained analogously from 5-bromothiophene-2-carboxylic acid.

[0343] M.p.: 200° C.

Example 5

N-({(5S)-3-[3-Fluoro-4-(1,4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-5-methyl-2-thiophenecarboxamide

[0344]




is obtained analogously from 5-methylthiophene-2-carboxylic acid.

[0345] M.p.: 167° C.

Example 6

5-Chloro-N-({(5S)-3-(6-methylthieno[2,3-b]pyridin-2-yl)-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0346]




is obtained analogously from (5S)-5-(aminomethyl)-3-(6-methylthieno[2,3-b]pyridin-2-yl)-1,3-oxazolidin-2-one (for preparation, see EP-A-785 200).

[0347] M.p.: 247° C.

Example 7

5-Chloro-N-({(5S)-3-(3-methyl-2-oxo-2,3-dihydro-1,3-benzothiazol-6-yl)-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0348]




is obtained analogously from 6-[(5S)-5-(aminomethyl)-2-oxo-1,3-oxazolidin-3-yl]-3-methyl-1,3-benzothiazol-2(3H)-one (for preparation, see EP-A-738 726).

[0349] M.p.: 217° C.

Example 8

5-Chloro-N-({(5S)-3-[3-fluoro-4-[4-(4-pyridinyl)piperazino]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0350]




is obtained analogously from (5S)-5-(aminomethyl)-3-{3-fluoro-4-[4-(4-pyridinyl)piperazino]phenyl}-1,3-oxazolidin-2-one (preparation in analogy to J. A. Tucker et al., J. Med. Chem. 1998, 41, 3727).

[0351] MS (ESI) 516 (M+H), Cl pattern.

Example 9

5-Chloro-N-({(5S)-3-[3-fluoro-4-(4-methylpiperazino)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0352]




is obtained analogously from (5S)-5-(aminomethyl)-3-{3-fluoro-4-(4-methylpiperazino)phenyl}-1,3-oxazolidin-2-one.

Example 10

5-Chloro-N-({(5S)-3-[3-fluoro-4-(4-tert-butoxycarbonylpiperazin-1-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0353]



is obtained analogously from (5S)-5-(aminomethyl)-3-{3-fluoro-4-(4-tert-butoxycarbonylpiperazin-1-yl)phenyl}-1,3-oxazolidin-2-one (for preparation, see WO-A-93/23384 which has already been cited).


[0354] M.p.: 184° C.;

[0355] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.42.

Example 11

5-Chloro-N-({(5S)-3-[3-fluoro-4-(piperazin-1-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0356]



is obtained by reacting Example 12 with trifluoroacetic acid in methylene chloride.


[0357] IC₅₀=140 nM;

[0358] ¹H-NMR [d₆-DMSO]: 3.01-3.25 (m, 8H), 3.5-3.65 (m, 2H), 3.7-3.9 (m, 1H), 4.05-4.2 (m, 1H), 4.75-4.9 (m, 1H), 7.05-7.25 (m, 3H), 7.5 (dd, 1H), 7.7 (d, 1H), 8.4 (broad s, 1H), 9.0 (t, 1H).

Example 12

5-Chloro-N-[(5S)-3-(2,4-bipyridinyl-5-yl)-2-oxo-1,3-oxazolidin-5-yl]methyl]-2-thiophenecarboxamide

[0359]



is obtained analogously from (5S)-5-(aminomethyl)-3-(2,4'-bipyridinyl-5-yl)-2-oxo-1,3-oxazolidin-2-one (for preparation, see EP-A-789 026).


[0360] R_f (SiO₂, ethyl acetate/ethanol 1:2)=0.6;

[0361] MS (ESI) 515 (M+H), Cl pattern.

Example 13

5-Chloro-N-[(5S)-2-oxo-3-(4-piperidinophenyl)-1,3-oxazolidin-5-yl]methyl]-2-thiophenecarboxamide

[0362]




is obtained from 5-(hydroxymethyl)-3-(4-piperidinophenyl)-1,3-oxazolidin-2-one (for preparation, see DE 2708236) after mesylation, reaction with potassium phthalimide, hydrazinolysis and reaction with 5-chlorothiophene-2-carboxylic acid. $R_f(\text{SiO}_2, \text{ethyl acetate/toluene } 1:1)=0.31$;

[0363] M.p. 205° C.

Example 17

5-Chloro-N-((5S)-2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0364]



[0365] 5-Chloro-N-((5S)-2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide is obtained from 1-(4-aminophenyl)pyrrolidin-2-one (for preparation, see Reppe et al., Justus Liebigs Ann. Chem.: 596; 1955; 209) in analogy to the known synthesis scheme (see S. J. Brickner et al., J. Med. Chem. 1996, 39, 673) after reaction with benzyloxycarbonyl chloride, subsequent reaction with R-glycidyl butyrate, mesylation, reaction with potassium phthalimide, hydrazinolysis in methanol and finally reaction with 5-chlorothiophene-2-carboxylic acid. The 5-chloro-N-((5S)-2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide obtained in this way has an IC_{50} of 4 nM (test method for the IC_{50} according to Example A-1. a.1) "Measurement of factor Xa inhibition" described above).

[0366] M.p.: 229° C.;

[0367] $R_f(\text{SiO}_2, \text{toluene/ethyl acetate } 1:1)=0.05$ (precursor: =0.0);

[0368] MS (ESI): 442.0 (21%, M+Na, Cl pattern), 420.0 (72%; M+H, Cl pattern), 302.3 (12%), 215 (52%), 145 (100%);

[0369] $^1\text{H-NMR}$ (d_6 -DMSO, 300 MHz): 2.05 (m, 2H), 2.45 (m, 2H), 3.6 (t, 2H), 3.77-3.85 (m, 3H), 4.15 (t, 1H), 4.75-4.85 (m, 1H), 7.2 (d, 1H), 7.5 (d, 2H), 7.65 (d, 2H), 7.69 (d, 1H), 8.96 (t, 1H).

[0370] The individual stages in the synthesis of Example 17 described above, with the respective precursors, are as follows:

[0371] 4.27 g (25.03 mmol) of benzyl chloroformate are slowly added to 4 g (22.7 mmol) of 1-(4-aminophenyl)pyrrolidin-2-one and 3.6 ml (28.4 mmol) of N,N-dimethylaniline in 107 ml of tetrahydrofuran at -20° C. The mixture is stirred at -20° C. for 30 minutes and then allowed to reach room temperature. 0.5 l of ethyl acetate is added, and the organic phase is washed with 0.5 l of saturated NaCl solution. The removed organic phase is dried with MgSO_4 , and the solvent is evaporated in vacuo. The residue is triturated with diethyl ether and filtered off with suction. 5.2 g (73.8% of theory) of benzyl 4-(2-oxo-1-pyrrolidinyl)phenylcarbamate are obtained as pale beige crystals with a melting point of 174° C.

[0372] 7.27 ml of a 2.5 M solution of n-butyllithium (BuLi) in hexane are added dropwise to 1.47 g (16.66 mmol) of isoamyl alcohol in 200 ml of tetrahydrofuran under argon at -10° C., a further 8 ml of the BuLi solution being necessary until the color of the added N-benzylidenebenzylamine indicator changed. The mixture is stirred at -10° C. for 10 minutes and cooled to -78° C., and a solution of 4.7 g (15.14 mmol) of benzyl 4-(2-oxo-1-pyrrolidinyl)phenylcarbamate is slowly added. Then a further 4 ml of n-BuLi solution are added until the color of the indicator changes to pink. The mixture is stirred at -78° C. for 10 minutes and 2.62 g (18.17 mmol) of R-glycidyl butyrate are added, and the mixture is stirred at -78° C. for 30 minutes.

[0373] The mixture is allowed to reach room temperature overnight, 200 ml of water are added to the mixture, and the THF content is evaporated in vacuo. The aqueous residue is extracted with ethyl acetate, and the organic phase is dried with MgSO_4 and concentrated in vacuo. The residue is triturated with 500 ml of diethyl ether, and the crystals which have separated out are filtered off with suction in vacuo.

[0374] 3.76 g (90% of theory) of (5R)-5-(hydroxymethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one are obtained with a melting point of 148° C. and an $R_f(\text{SiO}_2, \text{toluene/ethyl acetate } 1:1)=0.04$ (precursor=0.3).

[0375] 3.6 g (13.03 mmol) of (5R)-5-(hydroxymethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one and 2.9 g (28.67 mmol) of triethylamine are introduced into 160 ml of dichloromethane at 0° C. with stirring. 1.79 g (15.64 mmol) of methanesulfonyl chloride are added with stirring, and the mixture is stirred at 0° C. for 1.5 hours and at room temperature for 3 h.

[0376] The reaction mixture is washed with water and the aqueous phase is extracted once more with methylene chloride. The combined organic extracts are dried with MgSO_4 and evaporated. The residue (1.67 g) is then dissolved in 70 ml of acetonitrile, 2.62 g (14.16 mmol) of potassium phthalimide are added, and the mixture is stirred in a closed vessel at 180° C. in a microwave oven for 45 minutes.

[0377] The mixture is filtered off from the insoluble residue, the filtrate is concentrated in vacuo, the residue (1.9 g) is dissolved in methanol, and 0.47 g (9.37 mmol) of hydrazine hydrate is added. The mixture is boiled for 2 hours and cooled, saturated sodium bicarbonate solution is added, and the mixture is extracted six times with a total of 2 l of methylene chloride. The combined organic extracts of the crude (5S)-5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one are dried with MgSO_4 and concentrated in vacuo.

[0378] The final stage, 5-chloro-N-((5S)-2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide, is prepared by dissolving 0.32 g (1.16 mmol) of the (5S)-5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one prepared above, 5-chlorothiophene-2-carboxylic acid (0.19 g; 1.16 mmol) and 1-hydroxy-1H-benzotriazole hydrate (HOBT) (0.23 g, 1.51 mmol) in 7.6 ml of DMF. 0.29 g (1.51 mmol) of N'-(3-dimethylaminopropyl)-N-ethylcarbodiimide (EDCI) is added and at room temperature, 0.3 g (0.4 ml; 2.32 mmol, 2 equivalents) of diisopropylethylamine (DIEA) is added dropwise. The mixture is stirred at room temperature overnight.

[0379] The mixture is evaporated to dryness in vacuo, and the residue is dissolved in 3 ml of DMSO and chromatographed on an RP-MPLC with acetonitrile/water/0.5% TFA gradients. The acetonitrile content is evaporated off from the appropriate fractions, and the precipitated compound is filtered off with suction. 0.19 g (39% of theory) of the target compound is obtained.

[0380] The following were prepared in an analogous manner:

Example 18

5-Chloro-N-((5S)-2-oxo-3-[4-(1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0381] The compound 5-chloro-N-((5S)-2-oxo-3-[4-(1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide is obtained from 4-pyrrolidin-1-ylaniline (Reppe et al., Justus Liebigs Ann. Chem., 596; 1955; 151) in analogy to Example 17.

[0382] IC₅₀=40 nM;

[0383] M.p.: 216° C.;

[0384] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.31 [precursor: =0.0].

Example 19

5-Chloro-N-((5S)-2-oxo-3-[4-(diethylamino)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0385] The compound 5-chloro-N-((5S)-2-oxo-3-[4-(diethylamino)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide is obtained analogously from N,N-diethylphenyl-1,4-diamine (U.S. Pat. No. 2,811,555; 1955).

[0386] IC₅₀=270 nM;

[0387] M.p.: 181° C.;

[0388] R_f (SiO₂, toluene/ethyl acetate 1:1)=0.25 [precursor: =0.0].

Example 36

5-Chloro-N-((5S)-3-[2-methyl-4-(4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0389] starting from 2-methyl-4-(4-morpholinyl)aniline (J. E. LuValle et al. J. Am. Chem. Soc. 1948, 70, 2223);

[0390] MS (ESI): m/z (%)=436 ([M+H]⁺, 100), Cl pattern;

[0391] HPLC (method 1): rt (%)=3.77 (98).

[0392] IC₅₀: 1.26 μM

Example 37

5-Chloro-N-((5S)-3-(3-chloro-4-morpholinophenyl)-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0393] starting from 3-chloro-4-(4-morpholinyl)aniline (H. R. Snyder et al. J. Pharm. Sci. 1977, 66, 1204);

[0394] MS (ESI): m/z (%)=456 ([M+H]⁺, 100), Cl₂ pattern;

[0395] HPLC (method 2): rt (%)=4.31 (100).

[0396] IC₅₀: 33 nM

Example 38

5-Chloro-N-((5S)-3-[4-(4-morpholinylsulfonyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0397] starting from 4-(4-morpholinylsulfonyl)aniline (Adams et al. J. Am. Chem. Soc. 1939, 61, 2342);

[0398] MS (ESI): m/z (%)=486 ([M+H]⁺, 100), Cl pattern;

[0399] HPLC (method 3): rt (%)=4.07 (100).

[0400] IC₅₀: 2 μM

Example 39

5-Chloro-N-((5S)-3-[4-(1-azetidinylsulfonyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0401] starting from 4-(1-azetidinylsulfonyl)aniline;

[0402] MS (DCI, NH₃): m/z (%)=473 ([M+NH₄]⁺, 100), Cl pattern;

[0403] HPLC (method 3): rt (%)=4.10 (100).

[0404] IC₅₀: 0.84 μM

Example 40

5-Chloro-N-(((5S)-3-[4-[(dimethylamino)sulfonyl]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0405] starting from 4-amino-N,N-dimethylbenzenesulfonamide (I. K. Khanna et al. J. Med. Chem. 1997, 40, 1619);


[0406] MS (ESI): m/z (%)=444 ([M+H]⁺, 100), Cl pattern;

[0407] HPLC (method 3): rt (%)=4.22 (100).

[0408] IC₅₀: 90 nM

General method for the acylation of 5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one with carbonyl chlorides

[0409]



[0410] An approx. 0.1 molar solution of 5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one (from Example 45) (1.0 eq.) and absolute pyridine (approx. 6 eq) in absolute dichloromethane is added dropwise to the appropriate acid chloride (2.5 eq.) under argon at room temperature. The mixture is stirred at room temperature for about 4 h before approx. 5.5 eq of PS-trisamine (Argonaut Technologies) are added. The suspension is stirred gently for 2 h and, after dilution with dichloromethane/DMF (3:1), filtered (the resin is washed with dichloromethane/DMF) and the filtrate is concentrated. The resulting product is purified by preparative RP-HPLC where appropriate.

[0411] The following was prepared in an analogous manner:

Example 41

N-({2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide


[0412] LC-MS (method 6): m/z (%)=386 (M+H, 100);

[0413] LC-MS: rt (%)=3.04 (100).

[0414] IC₅₀: 1.3 μM

General method for preparing acyl derivatives starting from 5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one and carboxylic acids

[0415]



[0416] The appropriate carboxylic acid (approx. 2 eq) and a mixture of absolute dichloromethane/DMF (approx. 9:1) are added to 2.9 eq. of resin-bound carbodiimide (PS-Carbodiimide, Argonaut Technologies). After shaking gently at room temperature for about 15 min, 5-(aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one (from Example 45) (1.0 eq.) is added, and the mixture is shaken overnight before the resin is filtered off (washing with dichloromethane) and the filtrate is concentrated. The resulting product is purified by preparative RP-HPLC where appropriate.

[0417] The following were prepared in an analogous manner

Example 42

5-Methyl-N-({2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0418] LC-MS: m/z (%)=400 (M+H, 100);

[0419] LC-MS (method 6): rt (%)=3.23 (100).

[0420] IC₅₀: 0.16 μM

Example 43

5-Bromo-N-({2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0421] LC-MS: m/z (%)=466 (M+H, 100);


[0422] LC-MS (method 5): rt (%)=3.48 (78).

[0423] IC₅₀: 0.014 μM

Example 44

5-Chloro-N-({(5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0424]



a) 2-((2R)-2-Hydroxy-3-[[4-(3-oxo-4-morpholinyl)phenyl]amino]propyl)-1H-isoindole-1,3(2H)-dione

[0425] A suspension of 2-[(2S)-2-oxiranylmethyl]-1H-isoindole-1,3(2H)-dione (A. Gutcait et al. *Tetrahedron Asym.* 1996, 7, 1641) (5.68 g, 27.9 mmol) and 4-(4-aminophenyl)-3-morpholinone (5.37 g, 27.9 mmol) in ethanol/water (9:1, 140 ml) is refluxed for 14 h (the precipitate dissolves and,

after some time, there is renewed formation of a precipitate). The precipitate (desired product) is filtered off, washed three times with diethyl ether and dried. The combined mother liquers are concentrated in vacuo and, after addition of a second portion of 2-[(2S)-2-oxiranylmethyl]-1H-isoindole-1,3(2H)-dione (2.84 g, 14.0 mmol), are suspended in ethanol/water (9:1, 70 ml) and refluxed for 13 h (the precipitate dissolves and, after some time, there is renewed formation of a precipitate). The precipitate (desired product) is filtered off, washed three times with diethyl ether and dried. Overall yield: 10.14 g, 92% of theory.

[0426] MS (ESI): m/z (%)=418 ([M+Na]⁺, 84), 396 ([M+H]⁺, 93);

[0427] HPLC (method 3): rt (%)=3.34 (100).

b) 2-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-1H-isoindole-1,3(2H)-dione

[0428] N,N'-Carbonyldiimidazole (2.94 g, 18.1 mmol) and dimethylaminopyridine (catalytic amount) are added to a suspension of the amino alcohol (3.58 g, 9.05 mmol) in tetrahydrofuran (90 ml) under argon at room temperature. The reaction suspension is stirred at 60° C. for 12 h (the precipitate dissolves and, after some time, there is renewed formation of a precipitate), a second portion of N,N'-carbonyldiimidazole (2.94 g, 18.1 mmol) is added, and the mixture is stirred at 60° C. for a further 12 h. The precipitate (desired product) is filtered off, washed with tetrahydrofuran and dried. The filtrate is concentrated in vacuo and further product is purified by flash chromatography (dichloromethane/methanol mixtures). Overall yield: 3.32 g, 87% of theory.

[0429] MS (ESI): m/z (%)=422 ([M+H]⁺, 100);

[0430] HPLC (method 4): rt (%)=3.37 (100).

c) 5-Chloro-N-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)-methyl)-2-thiophenecarboxamide

[0431] Methylamine (40% strength in water, 10.2 ml, 0.142 mol) is added dropwise to a suspension of the oxazolidinone (4.45 g, 10.6 mmol) in ethanol (102 ml) at room temperature. The reaction mixture is refluxed for 1 h and concentrated in vacuo. The crude product is employed without further purification in the next reaction.

[0432] 5-Chlorothiophene-2-carbonyl chloride (2.29 g, 12.7 mmol) is added dropwise to a solution of the amine in pyridine (90 ml) under argon at 0° C. The ice cooling is removed and the reaction mixture is stirred at room temperature for 1 h, and water is added. Addition of dichloromethane and phase separation are followed by extraction of the aqueous phase with dichloromethane. The combined organic phases are dried (sodium sulfate), filtered and concentrated in vacuo. The desired product is purified by flash chromatography (dichloromethane/methanol mixtures). Overall yield: 3.92 g, 86% of theory.

[0433] M.p.: 232-233° C.;

[0434] ¹H NMR (DMSO-d₆, 200 MHz): 9.05-8.90 (t, J=5.8 Hz, 1H), 7.70 (d, J=4.1 Hz, 1H), 7.56 (d, J=9.0 Hz, 2H), 7.41 (d, J=9.0 Hz, 2H), 7.20 (d, J=4.1 Hz, 1H), 4.93-4.75 (m, 1H), 4.27-4.12 (m, 3H), 4.02-3.91 (m, 2H), 3.91-3.79 (dd, J=6.1 Hz, 9.2 Hz, 1H), 3.76-3.66 (m, 2H), 3.66-3.54 (m, 2H);

[0435] MS (ESI): m/z (%)=436 ([M+H]⁺, 100, Cl pattern);

[0436] HPLC (method 2): rt (%)=3.60 (100);

[0437] [α]_D²¹ = -38° (c 0.2985, DMSO); ee: 99%.

[0438] IC₅₀: 0.7 nM

[0439] The following were prepared in an analogous manner:

Example 45

5-Methyl-N-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0440] MS (ESI): m/z (%)=831 ([2M+H]⁺, 100), 416 ([M+H]⁺, 66);

[0441] HPLC (method 3): rt (%)=3.65 (100).

[0442] IC₅₀: 4.2 nM

Example 46

5-Bromo-N-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0443] MS (ESI): m/z (%)=480 ([M+H], 100, Br pattern);


[0444] HPLC (method 3): rt (%)=3.87 (100).

[0445] IC₅₀: 0.3 nM

Example 47

5-Chloro-N-((5S)-3-(3-isopropyl-2-oxo-2,3-dihydro-1,3-benzoxazol-6-yl)-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0446]







[0447] 200 mg (0.61 mmol) of 6-[(5S)-5-(aminomethyl)-2-oxo-1,3-oxazolidin-3-yl]-3-isopropyl-1,3-benzoxazol-2(3H)-one hydrochloride (EP 738726) are suspended in 5 ml of tetrahydrofuran, and 0.26 ml (1.83 mmol) of triethylamine and 132 mg (0.73 mmol) of 5-chlorothiophene-2-carbonyl chloride are added. The reaction mixture is stirred at room temperature overnight and then concentrated. The product is isolated by column chromatography (silica gel, methylene chloride/ethanol=50/1 to 20/1). 115 mg (43% of theory) of the desired compound are obtained.

[0448] MS (ESI): m/z (%)=436 (M+H, 100);

[0449] HPLC (method 4): rt=3.78 min.

[0450] The following compounds were prepared in an analogous manner:

Example No.	Structure	M.p. [° C.]	IC ₅₀ [μM]
48	 <p>Chiral</p>	210	0.12
49	 <p>Chiral</p>	234	0.074
50	 <p>Chiral</p>	195	1.15
51	 <p>Chiral</p>	212	1.19

-continued


Example No.	Structure	M.p. [° C.]	IC ₅₀ [μM]
57		226	0.58
58		228-230	

[0451] Examples 20 to 30 and 58 to 139 which follow relate to process variant [B], with Examples 20 and 21 describing the preparation of precursors.

Example 20

Preparation of N-allyl-5-chloro-2-thiophenecarboxamide

[0452]



[0453] 5-Chlorothiophene-2-carbonyl chloride (7.61 g, 42 mmol) is added dropwise to an ice-cooled solution of 2.63 ml (35 mmol) of allylamine in 14.2 ml of absolute pyridine and 14.2 ml of absolute THF. The ice cooling is removed and the mixture is stirred at room temperature for 3 h before being concentrated in vacuo. Water is added to the residue, and the solid is filtered off. The crude product is purified by flash chromatography on silica gel (dichloromethane).

[0454] Yield: 7.20 g (99% of theory);


[0455] MS (DCI, NH₄): m/z (%)=219 (M+NH₄, 100), 202 (M+H, 32);

[0456] HPLC (method 1): rt (%)=3.96 min (98.9).


Example 21

Preparation of 5-chloro-N-(2-oxiranylmethyl)-2-thiophenecarboxamide

[0457]



-continued




[0458] meta-Chloroperbenzoic acid (3.83 g, approx. 60% pure) is added to an ice-cooled solution of 2.0 g (9.92 mmol) of N-allyl-5-chloro-2-thiophenecarboxamide in 10 ml of dichloromethane. The mixture is stirred overnight while warming to room temperature, and then washed with 10% sodium bisulfate solution (three times). The organic phase is washed with saturated sodium bicarbonate solution (twice) and with saturated sodium chloride solution, dried over magnesium sulfate and concentrated. The product is purified by chromatography on silica gel (cyclohexane/ethyl acetate 1:1).

[0459] Yield: 837 mg (39% of theory);

[0460] MS (DCI, NH₄): m/z (%)=253 (M+NH₄, 100), 218 (M+H, 80);

[0461] HPLC (method 1): rt (%)=3.69 min (approx. 80). General method for preparing substituted N-(3-amino-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide derivatives starting from 5-chloro-N-(2-oxiranylmethyl)-2-thiophenecarboxamide



[0462] 5-Chloro-N-(2-oxiranylmethyl)-2-thiophenecarboxamide (1.0 eq.) is added in portions to a solution of primary amine or aniline derivative (1.5 to 2.5 eq.) in 1,4-dioxane, 1,4-dioxane/water mixtures or ethanol, ethanol/water

mixtures (approx. 0.3 to 1.0 mol/l) at room temperature or at temperatures up to 80° C. The mixture is stirred for 2 to 6 hours before being concentrated. The product can be isolated from the reaction mixture by chromatography on silica gel (cyclohexane/ethyl acetate mixtures, dichloromethane/methanol mixtures or dichloromethane/methanol/triethylamine mixtures).

[0463] The following were prepared in an analogous manner:

Example 22

N-[3-(Benzylamino)-2-hydroxypropyl]-5-chloro-2-thiophenecarboxamide

[0464] MS (ESI): m/z (%)=325 (M+H, 100);

[0465] HPLC (method 1): rt (%)=3.87 min (97.9).

Example 23

5-Chloro-N-[3-(3-cyanoanilino)-2-hydroxypropyl]-2-thiophenecarboxamide

[0466] MS (ESI): m/z (%)=336 (M+H, 100);

[0467] HPLC (method 2): rt (%)=4.04 min (100).

Example 24

5-Chloro-N-[3-(4-cyanoanilino)-2-hydroxypropyl]-2-thiophenecarboxamide

[0468] MS (ESI): m/z (%)=336 (M+H, 100);

[0469] HPLC (method 1): rt (%)=4.12 min (100).

Example 25

5-Chloro-N-[3-[4-(cyanomethyl)anilino]-2-hydroxypropyl]-2-thiophenecarboxamide

[0470] MS (ESI): m/z (%)=350 (M+H, 100);

[0471] HPLC (method 4): rt (%)=3.60 min (95.4).

Example 26

5-Chloro-N-[3-[3-(cyanomethyl)anilino]-2-hydroxypropyl]-2-thiophenecarboxamide

[0472] MS (ESI): m/z (%)=350 (M+H, 100);

[0473] HPLC (method 4): rt (%)=3.76 min (94.2).

Example 58

tert-Butyl 4-[(3-[(5-chloro-2-thienyl)carbonyl]amino)-2-hydroxypropyl]amino-benzylcarbamate

[0474] Starting from tert-butyl 4-aminobenzylcarbamate (*Bioorg. Med. Chem. Lett.*: 1997; 1921-1926);

[0475] MS (ES-pos): m/z (%)=440 (M+H, 100), (ES-neg): m/z (%)=438 (M-H, 100);

[0476] HPLC (method 1): rt (%)=4.08 (100).

Example 59

tert-Butyl 4-[(3-[(5-chloro-2-thienyl)carbonyl]amino)-2-hydroxypropyl]amino-phenylcarbamate

[0477] Starting from N-tert-butylloxycarbonyl-1,4-phenylenediamine:

[0478] MS (ESI): m/z (%)=426 (M+H, 45), 370 (100);

[0479] HPLC (method 1): rt (%)=4.06 (100).

Example 60

tert-Butyl 2-hydroxy-3-[[4-(2-oxo-1-pyrrolidinyl)phenyl]amino]propyl-carbamate

[0480] Starting from 1-(4-aminophenyl)-2-pyrrolidinone (*Justus Liebigs Ann. Chem.*: 1955; 596; 204);

[0481] MS (DCI, NH₃): m/z (%)=350 (M+H, 100);

[0482] HPLC (method 1): rt (%)=3.57 (97).

Example 61

5-Chloro-N-(3-[[3-fluoro-4-(3-oxo-4-morpholinyl)phenyl]amino]-2-hydroxypropyl)-2-thiophenecarboxamide

[0483] 800 mg (3.8 mmol) of 4-(4-amino-2-fluorophenyl)-3-morpholinone and 700 mg (3.22 mmol) of 5-chloro-N-(2-oxiranylmethyl)-2-thiophenecarboxamide are heated in 15 ml of ethanol and 1 ml of water under reflux for 6 hours. The mixture is evaporated in vacuo, the crystals which have separated out after treatment with ethyl acetate are filtered off with suction, and chromatography of the mother liquor results in 276 mg (17% of theory) of the target compound.

[0484] R_f(ethyl acetate): 0.25.

Example 62

(N-(3-Anilino-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide

[0485] starting from aniline:

[0486] MS (DCI, NH₃): m/z (%)=311 ([M+H]⁺, 100), Cl pattern;

[0487] HPLC (method 3): rt (%)=179 (100).

Example 63

5-Chloro-N-(2-hydroxy-3-[[4-(3-oxo-4-morpholinyl)phenyl]amino]propyl)-2-thiophenecarboxamide

[0488] starting from 4-(4-aminophenyl)-3-morpholinone:

[0489] MS (ESI): m/z (%)=410 ([M+H]⁺, 50), Cl pattern;

[0490] HPLC (method 3): rt (%)=3.58 (100).

Example 64

N-[3-([4-[Acetyl(cyclopropyl)amino]phenyl]amino)-2-hydroxypropyl]-5-chloro-2-thiophenecarboxamide

[0491] starting from N-(4-aminophenyl)-N-cyclopropylacetamide:

[0492] MS (ESI): m/z (%)=408 ([M+H]⁺, 100), Cl pattern;

[0493] HPLC (method 3): rt (%)=3.77 (100).

Example 65

N-[3-([4-[Acetyl(methyl)amino]phenyl]amino)-2-hydroxypropyl]-5-chloro-2-thiophenecarboxamide

[0494] starting from N-(4-aminophenyl)-N-methylacetamide:

[0495] MS (ESI): m/z (%)=382 (M+H, 100);

[0496] HPLC (method 4): rt=3.31 min.

Example 66

5-Chloro-N-(2-hydroxy-3-{{4-(1H-1,2,3-triazol-1-yl)phenyl}amino}propyl)-2-thiophenecarboxamide

- [0497] starting from 4-(1H-1,2,3-triazol-1-yl)aniline (Bouchet et al.; J. Chem. Soc. Perkin Trans. 2; 1974; 449);
 [0498] MS (ESI): m/z (%)=378 (M+H, 100);
 [0499] HPLC (method 4): rt=3.55 min.

Example 67

Tert-butyl 1-{{4-{{3-{{(5-chloro-2-thienyl)carbonyl}amino}-2-hydroxypropyl)-amino}phenyl}-L-prolinate

- [0500] MS (ESI): m/z (%)=480 (M+H, 100);
 [0501] HPLC (method 4): rt=3.40 min.

Example 68

1-{{4-{{3-{{(5-Chloro-2-thienyl)carbonyl}amino}-2-hydroxypropyl)amino}phenyl}-4-piperidinecarboxamide

- [0502] MS (ESI): m/z (%)=437 (M+H, 100);
 [0503] HPLC (method 4): rt=2.39 min.

Example 69

1-{{4-{{3-{{(5-Chloro-2-thienyl)carbonyl}amino)-2-hydroxypropyl)-amino}phenyl}-3-piperidinecarboxamide

- [0504] MS (ESI): m/z (%)=437 (M+H, 100);
 [0505] HPLC (method 4): rt=2.43 min.

Example 70

5-Chloro-N-(2-hydroxy-3-{{4-(4-oxo-1-piperidinyl)phenyl}amino}propyl)-2-thiophenecarboxamide

- [0506] MS (ESI): m/z (%)=408 (M+H, 100);
 [0507] HPLC (method 4): rt=2.43 min.

Example 71

1-{{4-{{3-{{(5-Chloro-2-thienyl)carbonyl}amino}-2-hydroxypropyl)amino}phenyl}-L-prolinamide

- [0508] MS (ESI): m/z (%)=423 (M+H, 100);
 [0509] HPLC (method 4): rt=2.51 min.

Example 72

5-Chloro-N-[[2-hydroxy-3-{{4-[[3-(hydroxymethyl)-1-piperidinyl]phenyl]-amino}propyl]-2-thiophenecarboxamide

- [0510] MS (ESI): m/z (%)=424 (M+H, 100);
 [0511] HPLC (method 4): rt=2.43 min.

Example 73

5-Chloro-N-[[2-hydroxy-3-{{4-[[2-(hydroxymethyl)-1-piperidinyl]phenyl]-amino}propyl]-2-thiophenecarboxamide

- [0512] MS (ESI): m/z (%)=424 (M+H, 100);
 [0513] HPLC (method 4): rt=2.49 min.

Example 74

Ethyl 1-{{4-{{3-{{(5-chloro-2-thienyl)carbonyl}amino}-2-hydroxypropyl)-amino}phenyl}-2-piperidinecarboxylate

- [0514] MS (ESI): m/z (%)=466 (M+H, 100);
 [0515] HPLC (method 4): rt=3.02 min.

Example 75

5-Chloro-N-[[2-hydroxy-3-{{4-[[2-(hydroxymethyl)-1-pyrrolidinyl]phenyl]amino}-propyl]-2-thiophenecarboxamide

- [0516] MS (ESI): m/z (%)=410 (M+H, 100);
 [0517] HPLC (method 4): rt=2.48 min.

Example 76

5-Chloro-N-(2-hydroxy-3-{{4-(2-methylhexahydro-5H-pyrrolo[3,4-d]isoxazol-5-yl)phenyl}amino}propyl)-2-thiophenecarboxamide

- [0518] MS (ESI): m/z (%)=437 (M+H, 100);
 [0519] HPLC (method 5): rt=1.74 min.

Example 77

5-Chloro-N-(2-hydroxy-3-{{4-(1-pyrrolidinyl)-3-(trifluoromethyl)phenyl}-amino}propyl)-2-thiophenecarboxamide

- [0520] MS (ESI): m/z (%)=448 (M+H, 100);
 [0521] HPLC (method 4): rt=3.30 min.

Example 78

5-Chloro-N-(2-hydroxy-3-{{4-(2-oxo-1-pyrrolidinyl)-3-(trifluoromethyl)phenyl]-amino}propyl)-2-thiophenecarboxamide

- [0522] MS (ESI): m/z (%)=462 (M+H, 100);
 [0523] HPLC (method 4): rt=3.50 min.

Example 79

5-Chloro-N-(3-{{3-chloro-4-(3-oxo-4-morpholinyl)phenyl}amino}-2-hydroxypropyl)-2-thiophenecarboxamide

- [0524] MS (ESI): m/z (%)=444 (M+H, 100);
 [0525] HPLC (method 4): rt=3.26 min.

Example 80

5-Chloro-N-(2-hydroxy-3-{{4-(3-oxo-4-morpholinyl)-3-(trifluoromethyl)phenyl]-amino}propyl)-2-thiophenecarboxamide

- [0526] MS (ESI): m/z (%)=478 (M+H, 100);
 HPLC (method 4): rt=3.37 min.

Example 81

5-Chloro-N-(2-hydroxy-3-{{3-methyl-4-(3-oxo-4-morpholinyl)phenyl}amino}-propyl)-2-thiophenecarboxamide

- [0527] MS (ESI): m/z (%)=424 (M+H, 100);
 [0528] HPLC (method 4): rt=2.86 min.

Example 82

5-Chloro-N-(3-([3-cyano-4-(3-oxo-4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0529] MS (ESI): m/z (%)=435 (M+H, 100);

[0530] HPLC (method 4): rt=3.10 min.

Example 83

5-Chloro-N-(3-([3-chloro-4-(1-pyrrolidinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0531] MS (ESI): m/z (%)=414 (M+H, 100);

[0532] HPLC (method 4): rt=2.49 min.

Example 84

5-Chloro-N-(3-([3-chloro-4-(2-oxo-1-pyrrolidinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0533] MS (ESI): m/z (%)=428 (M+H, 100);

[0534] HPLC (method 4): rt=3.39 min.

Example 85

5-Chloro-N-(3-([3,5-dimethyl-4-(3-oxo-4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0535] MS (ESI): m/z (%)=438 (M+H, 100);

[0536] HPLC (method 4): rt=2.84 min.

Example 86

N-(3-([3-(Aminocarbonyl)-4-(4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide

[0537] MS (ESI): m/z (%)=439 (M+H, 100);

[0538] HPLC (method 4): rt=2.32 min.

Example 87

5-Chloro-N-(2-hydroxy-3-([3-methoxy-4-(4-morpholinyl)phenyl]amino)propyl)-2-thiophenecarboxamide

[0539] MS (ESI): m/z (%)=426 (M+H, 100);

[0540] HPLC (method 4): rt=2.32 min.

Example 88

N-(3-([3-Acetyl-4-(4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide

[0541] MS (ESI): m/z (%)=438 (M+H, 100);

[0542] HPLC (method 4): rt=2.46 min.

Example 89

N-(3-([3-Amino-4-(3-oxo-4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide

[0543] MS (ESI): m/z (%)=425 (M+H, 100);

[0544] HPLC (method 4): rt=2.45 min.

Example 90

5-Chloro-N-(3-([3-chloro-4-(2-methyl-3-oxo-4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0545] MS (ESI): m/z (%)=458 (M+H, 100);

[0546] HPLC (method 4): rt=3.44 min.

Example 91

5-Chloro-N-(3-([3-chloro-4-(2-methyl-5-oxo-4-morpholinyl)phenyl]amino)-2-hydroxypropyl)-2-thiophenecarboxamide

[0547] MS (ESI): m/z (%)=458 (M+H, 100);

[0548] HPLC (method 4): rt=3.48 min.

Example 91a


5-Chloro-N-[2-hydroxy-3-({4-[(3-oxo-4-morpholinyl)methyl]phenyl}amino)-propyl]-2-thiophenecarboxamide

[0549] Starting from 4-(4-aminobenzyl)-3-morpholinone (Surrey et al.; J. Amer. Chem. Soc.; 77; 1955; 633):

[0550] MS (ESI): m/z (%)=424 (M+H, 100);

[0551] HPLC (method 4): rt=2.66 min

General method for preparing 3-substituted 5-chloro-N-[(2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide derivatives starting from substituted N-(3-amino-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide derivatives



[0552] Carbodiimidazole (1.2 to 1.8 eq.) or a comparable phosgene equivalent is added to a solution of substituted N-(3-amino-2-hydroxypropyl)-5-chloro-2-thiophenecarboxamide derivative (1.0 eq.) in absolute THF (approx. 0.1 mol/l) at room temperature. The mixture is stirred at room temperature or, where appropriate, at elevated temperature (up to 70° C.) for 2 to 18 h before being concentrated in vacuo. The product can be purified by chromatography on silica gel (dichloromethane/methanol mixtures or cyclohexane/ethyl acetate mixtures).

[0553] The following were prepared in an analogous manner:

Example 27

N-[(3-Benzyl-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide

[0554] MS (DCI, NH₄): m/z (%)=372 (M+Na, 100), 351 (M+H, 45);

[0555] HPLC (method 1): rt (%)=4.33 min (100).

Example 28

5-Chloro-N-[[3-(3-cyanophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl]-2-thiophenecarboxamide

[0556] MS (DCI, NH₄): m/z (%)=362 (M+H, 42), 145 (100);

[0557] HPLC (method 2): rt (%)=4.13 min (100).

Example 29

5-Chloro-N-([3-[4-(cyanomethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl)-2-thiophenecarboxamide

[0558] MS (ESI): m/z (%)=376 (M+H, 100);

[0559] HPLC (method 4): rt=4.12 min

Example 30

5-Chloro-N-([3-[3-(cyanomethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl)-2-thiophenecarboxamide

[0560] MS (ESI): m/z (%)=376 (M+H, 100);

HPLC (method 4): rt=4.17 min

Example 92

tert-Butyl 4-[5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]benzylcarbamate

[0561] starting from Example 58:

[0562] MS (ESI): m/z (%)=488 (M+Na, 23), 349 (100);

[0563] HPLC (method 1): rt (%)=4.51 (98.5).

Example 93

tert-Butyl 4-[5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenylcarbamate

[0564] starting from Example 59:

[0565] MS (ESI): m/z (%)=493 (M+Na, 70), 452 (M+H, 10), 395 (100);

[0566] HPLC (method 1): rt (%)=4.41 (100).

Example 94

[0567] tert-Butyl 2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl]methylcarbamate

[0568] starting from Example 60:


[0569] MS (DCI, NH₃): m/z (%)=393 (M+NH₄, 100);

[0570] HPLC (method 3): rt (%)=3.97 (100).

Example 95

5-Chloro-N-([3-[3-fluoro-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl)-2-thiophenecarboxamide

[0571]



[0572] 260 mg (0.608 mmol) of 5-chloro-N-(3-{[3-fluoro-4-(3-oxo-4-morpholinyl)phenyl]-amino}-2-hydroxypropyl)-2-thiophenecarboxamide (from Example 61), 197 mg (1.22 mmol) of carbonylimidazole and 7 mg of dimethylaminopyridine are boiled in 20 ml of dioxane under reflux for 5 hours. 20 ml of acetonitrile are then added, and the mixture is stirred in a closed container in a microwave oven at 180° C. for 30 minutes. The solution is concentrated in a rotary evaporator and chromatographed on an RP-HPLC column. 53 mg (19% of theory) of the target compound are obtained.

[0573] NMR (300 MHz, d₆-DMSO): δ=3.6-3.7 (m, 4H), 3.85 (dd, 1H), 3.95 (m, 2H), 4.2 (m, 1H), 4.21 (s, 2H), 4.85 (m, 1H), 4.18 (s, 2H), 7.19 (d, 1H, thiophene), 7.35 (dd, 1H), 7.45 (t, 1H), 7.55 (dd, 1H), 7.67 (d, 1H, thiophene), 8.95 (t, 1H, CONH).

Example 96

5-Chloro-N-[(2-oxo-3-phenyl-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0574] starting from Example 62:

[0575] MS (EST): m/z (%)=359 ([M+Na]⁺, 71), 337 ([M+H]⁺, 100), Cl pattern;

[0576] HPLC (method 3): rt (%)=4.39 (100).

[0577] IC₅₀: 2 μM

Example 97

5-Chloro-N-([2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl]-methyl)-2-thiophenecarboxamide

[0578] starting from Example 63:

[0579] MS (ESI): m/z (%)=458 ([M+Na]⁺, 66), 436 ([M+H]⁺, 100), Cl pattern;

HPLC (method 3): rt (%)=3.89 (100).

[0580] IC₅₀: 1.4 nM

Example 98

N-([3-[4-[Acetyl(cyclopropyl)amino]phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl)-5-chloro-2-thiophenecarboxamide

[0581] starting from Example 64:

[0582] MS (ESI): m/z (%)=456 ([M+Na]⁺, 55), 434 ([M+H]⁺, 100), Cl pattern;

[0583] HPLC (method 3): rt (%)=4.05 (100)

[0584] IC₅₀: 50 nM .

Example 99

N-[(3-{4-[Acetyl(methyl)amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide

[0585] MS (ESI): m/z (%)=408 (M+H, 30), 449 (M+H+MeCN, 100);

[0586] HPLC (method 4): rt=3.66 min.

Example 100

5-Chloro-N-({2-oxo-3-[4-(1H-1,2,3-triazol-1-yl)phenyl]-1,3-oxazolidin-5-yl}-methyl)-2-thiophenecarboxamide

[0587] MS (ESI): m/z (%)=404 (M+H, 45), 445 (M+H+MeCN, 100);

[0588] HPLC (method 4): rt=3.77 min.

Example 101

Tert-butyl 1-{4-[5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-L-prolinate

[0589] MS (ESI): m/z (%)=450 (M+H-56, 25), 506 (M+H, 100);

[0590] HPLC (method 4): rt=5.13 min.

Example 102

1-{4-[5-({[(5-Chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-4-piperidinecarboxamide

[0591] MS (ESI): m/z (%)=463 (M+H, 100);

[0592] HPLC (method 4): rt=2.51 min.

Example 103

1-{4-[5-({[(5-Chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-3-piperidinecarboxamide

[0593] MS (ESI): m/z (%)=463 (M+H, 100);

[0594] HPLC (method 4): rt=2.67 min.

Example 104

5-Chloro-N-({2-oxo-3-[4-(4-oxo-1-piperidinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0595] MS (ESI): m/z (%)=434 (M+H, 40), 452 (M+H+H₂O, 100), 475 (M+H+MeCN, 60);

[0596] HPLC (method 4): rt=3.44 min.

Example 105

1-{4-[5-({[(5-Chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-L-prolinamide

[0597] MS (ESI): m/z (%)=449 (M+H, 100);

[0598] HPLC (method 4): rt=3.54 min

Example 106

5-Chloro-N-[(3-{4-[3-(hydroxymethyl)-1-piperidinyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0599] MS (ESI): m/z (%)=450 (M+H, 100);

[0600] HPLC (method 5): rt=2.53 min

Example 107

5-Chloro-N-[(3-{4-[2-(hydroxymethyl)-1-piperidinyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0601] MS (ESI): m/z (%)=450 (M+H, 100);

[0602] HPLC (method 5): rt=2.32 min.

Example 108

Ethyl 1-{4-[5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-2-piperidinecarboxylate

[0603] MS (ESI): m/z (%)=492 (M+H, 100);

[0604] HPLC (method 5): rt=4.35 min.

Example 109

5-Chloro-N-[(3-{4-[2-(hydroxymethyl)-1-pyrrolidinyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0605] MS (ESI): m/z (%)=436 (M+H, 100);

[0606] HPLC (method 4): rt=2.98 min

Example 110

5-Chloro-N-({2-oxo-3-[4-(1-pyrrolidinyl)-3-(trifluoromethyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0607] MS (ESI): m/z (%)=474 (M+H, 100);

[0608] HPLC (method 4): rt=4.63 min.

Example 111

5-Chloro-N-({3-[4-(2-methylhexahydro-5,3-pyrrolo[3,4-d]isoxazol-5-yl)phenyl]2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0609] MS (ESI): m/z (%)=463 (M+H, 100);

[0610] HPLC (method 4): rt=2.56 min.

Example 112

5-Chloro-N-({2-oxo-3-[4-(2-oxo-1-pyrrolidinyl)-3-(trifluoromethyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0611] MS (ESI): m/z (%)=488 (M+H, 100);

[0612] HPLC (method 4): rt=3.64 min.

Example 113

5-Chloro-N-({3-[3-chloro-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0613] MS (ESI): m/z (%)=470 (M+H, 100);

[0614] HPLC (method 4): rt=3.41 min.

Example 114

5-Chloro-N-({2-oxo-3-[4-(3-oxo-4-morpholinyl)-3-(trifluoromethyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0615] MS (ESI): m/z (%)=504 (M+H, 100);

[0616] HPLC (method 4): rt=3.55 min.

Example 115

5-Chloro-N-({3-[3-methyl-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0617] MS (ESI): m/z (%) 450 (M+H, 100);

[0618] HPLC (method 4): rt=3.23 min.

Example 116

5-Chloro-N-({3-[3-cyano-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0619] MS (ESI): m/z (%)=461 (M+H, 100);

[0620] HPLC (method 4): rt=3.27 min.

Example 117

5-Chloro-N-({3-[3-chloro-4-(1-pyrrolidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0621] MS (ESI): m/z (%)=440 (M+H, 100);

[0622] HPLC (method 4): rt=3.72 min.

Example 118

5-Chloro-N-({3-[3-chloro-4-(2-oxo-1-pyrrolidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0623] MS (ESI): m/z (%)=454 (M+H, 100);

[0624] HPLC (method 4): rt=3.49 min.

Example 119

5-Chloro-N-({3-[3,5-dimethyl-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0625] MS (ESI): m/z (%)=464 (M+H, 100);

[0626] HPLC (method 4): rt=3.39 min.

Example 120

N-({3-[3-(Aminocarbonyl)-4-(4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0627] MS (ESI): m/z (%)=465 (M+H, 100);

[0628] HPLC (method 4): rt=3.07 min.

Example 121

5-Chloro-N-({3-[3-methoxy-4-(4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0629] MS (ESI): m/z (%)=452 (M+H, 100);

[0630] HPLC (method 4): rt=2.86 min.

Example 122

N-({3-[3-Acetyl-4-(4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0631] MS (ESI): m/z (%)=464 (M+H, 100);

[0632] HPLC (method 4): rt=3.52 min

Example 123

N-({3-[3-Amino-4-(3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}-methyl)-5-chloro-2-thiophenecarboxamide

[0633] MS (ESI): m/z (%)=451 (M+H, 100);

[0634] HPLC (method 6): rt=3.16 min.

Example 124

5-Chloro-N-({3-[3-chloro-4-(2-methyl-3-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0635] MS (ESI): m/z (%)=484 (M+H, 100);

[0636] HPLC (method 4): rt=3.59 min.

Example 125

5-Chloro-N-({3-[3-chloro-4-(2-methyl-5-oxo-4-morpholinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0637] MS (ESI): m/z (%)=484 (M+H, 100);

[0638] HPLC (method 4): rt=3.63 min.









Example 125a

5-Chloro-N-[(2-oxo-3-{4-[(3-oxo-4-morpholinyl)methyl]phenyl}-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide






[0639] MS (ESI): m/z (%)=450 (M+H, 100);

[0640] HPLC (method 4): rt=3.25 min.

[0641] In addition, the following compounds were prepared by the route of epoxide opening with an amine and subsequent cyclization to give the corresponding oxazolidinone:

Example No.	Structure	M.p. [° C.]	IC ₅₀ [μM]
126	 	229 decomp.	0.013
127		159	0.0007
128		198	0.002
129		196	0.001
130		206	0.0033
130a		194	
131		195	0.85

-continued

Example No.	Structure	M.p. [° C.]	IC ₅₀ [μM]
132		206	0.12
133		217	0.062
134	 <p>from 1-(4-Amino-phenyl)-piperidin-3-ol (Tong, L.K.J. et al; <i>J. Amer. Chem Soc</i> 1960; 82, 1998).</p>	207	0.48
135		202	1.1
136		239	1.2

-continued

Example No.	Structure	M.p. [° C.]	IC ₅₀ [μM]
137		219	0.044
138		95	0.42
139		217	1.7

[0642] Examples 14 to 16 which follow are exemplary embodiments of the optional oxidation process step, i.e. one which takes place where appropriate.

Example 14

5-Chloro-N-({(5S)-3-[3-fluoro-4-(1-oxo-1[lambda]⁴, 4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

50 ml of water are then added to the mixture, and the insoluble product is filtered off with suction. Washing with water and drying result in 60 mg (58% of theory) of crystals.


[0645] M.p.: 257° C.;

[0646] R_f (silica gel, toluene/ethyl acetate 1:1)=0.54 (precursor=0.46);

[0647] IC₅₀=1.1 μM;

[0648] MS (DCI) 489 (M+NH₄), Cl pattern.

[0643]




[0644] 5-Chloro-N-({(5S)-3-[3-fluoro-4-(1,4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide (0.1 g, 0.22 mmol) from Example 3 in methanol (0.77 ml) is added at 0° C. to a solution of sodium periodate (0.05 g, 0.23 mmol) in water (0.54 ml) and stirred at 0° C. for 3 h. Then 1 ml of DMF is added, and the mixture is stirred at RT for 8 h. Addition of a further 50 mg of sodium periodate is followed by stirring at RT once again overnight.

Example 15

Preparation of 5-chloro-N-({(5S)-3-[4-(1,1-dioxo-1[lambda]⁶, 4-thiazinan-4-yl)-3-fluorophenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0649]



[0650] 80 mg (0.66 mmol) of N-methylmorpholine N-oxide (NMO) and 0.1 ml of a 2.5% strength solution of osmium

tetroxide in 2-methyl-2-propanol are added to 5-chloro-N-((5S)-3-[3-fluoro-4-(1,4-thiazinan-4-yl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide from Example 3 (0.1 g, 0.22 mmol) in 3.32 ml of a mixture of 1 part of water and 3 parts of acetone. The mixture is stirred at room temperature overnight and a further 40 mg of NMO are added. After being stirred for a further night, the mixture is added to 50 ml of water and extracted three times with ethyl acetate. Drying and evaporation of the organic phase result in 23 mg, and filtration with suction of the insoluble solid from the aqueous phase results in 19 mg of the target compound (total 39% of theory).

[0651] M.p.: 238° C.;

[0652] R_f (toluene/ethyl acetate 1:1)=0.14 (precursor=0.46);

[0653] IC_{50} =210 nM;

[0654] MS (DCI): 505 (M+NH₄), Cl pattern.

Example 16

5-Chloro-N-((5S)-3-(3-fluoro-4-morpholinophenyl)-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide N-oxide

[0655] is obtained by treating 5-chloro-N-((5S)-3-(3-fluoro-4-morpholinophenyl)-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide from Example 1 with monoperoxyphthalic acid magnesium salt.

[0656] MS (ESI): 456 (M+H, 21%, Cl pattern), 439 (100%).

[0657] Examples 31 to 35 and 140 to 147 which follow relate to the optional amidination process step, i.e. one which takes place where appropriate.

General method for preparing amidines and amidine derivatives starting from cyanomethylphenyl-substituted 5-chloro-N-[(2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide derivatives

[0658] The particular cyanomethylphenyl-substituted 5-chloro-N-[(2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide derivative (1.0 eq.) is stirred together with triethylamine (8.0 eq.) in a saturated solution of hydrogen sulfide in pyridine (approx. 0.05-0.1 mol/l) at RT for one to two days. The reaction mixture is diluted with ethyl acetate (EtOAc) and washed with 2 N hydrochloric acid. The organic phase is dried with MgSO₄, filtered and evaporated in vacuo.

[0659] The crude product is dissolved in acetone (0.01-0.1 mol/l), and methyl iodide (40 eq.) is added. The reaction mixture is stirred at room temperature (RT) for 2 to 5 h and then concentrated in vacuo.

[0660] The residue is dissolved in methanol (0.01-0.1 mol/l) and, to prepare the unsubstituted amidines, ammonium acetate (3 eq.) and ammonium chloride (2 eq.) are added. The substituted amidine derivatives are prepared by adding primary or secondary amines (1.5 eq.) and acetic acid (2 eq.) to the methanolic solution. After 5-30 h, the solvent is removed in vacuo and the residue is purified by chromatography on an RP8 silica gel column (water/acetonitrile 9/1-1/1+0.1% trifluoroacetic acid).

[0661] The following were prepared in an analogous manner:

Example 31

N-((3-[4-(2-Amino-2-iminoethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-5-chloro-2-thiophenecarboxamide

[0662] MS (ESI): m/z (%)=393 (M+H, 100);

[0663] HPLC (method 4): rt=2.63 min

Example 32

5-Chloro-N-((3-[3-(4,5-dihydro-1H-imidazol-2-ylmethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0664] MS (ESI): m/z (%)=419 (M+H, 100);

[0665] HPLC (method 4): rt=2.61 min

Example 33

5-Chloro-N-[(3-[3-[2-imino-2-(4-morpholinyl)ethyl]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0666] MS (ESI): m/z (%)=463 (M+H, 100);

[0667] HPLC (method 4): rt=2.70 min

Example 34

5-Chloro-N-[(3-[3-[2-imino-2-(1-pyrrolidinyl)ethyl]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0668] MS (ESI): m/z (%)=447 (M+H, 100);

[0669] HPLC (method 4): rt=2.82 min

Example 35

N-((3-[3-(2-Amino-2-iminoethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-5-chloro-2-thiophenecarboxamide

[0670] MS (ESI): m/z (%)=393 (M+H, 100);

[0671] HPLC (method 4): rt=2.60 min

Example 140

5-Chloro-N-((3-[4-(4,5-dihydro-1H-imidazol-2-ylmethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0672] MS (ESI): m/z (%)=419 (M+H, 100);

[0673] HPLC (method 4): rt=2.65 min

Example 141

5-Chloro-N-[(3-[4-[2-imino-2-(4-morpholinyl)ethyl]phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0674] MS (ESI): m/z (%)=463 (M+H, 100);

[0675] HPLC (method 4): rt=2.65 min

Example 142

5-Chloro-N-[(3-{4-[2-imino-2-(1-piperidinyl)ethyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0676] MS (ESI): m/z (%)=461 (M+H, 100);

[0677] HPLC (method 4): rt=2.83 min

Example 143

5-Chloro-N-[(3-{4-[2-imino-2-(1-pyrrolidinyl)ethyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0678] MS (ESI): m/z (%)=447 (M+H, 100);

[0679] HPLC (method 4): rt=2.76 min

Example 144

5-Chloro-N-[(3-{4-[2-(cyclopentylamino)-2-iminoethyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0680] MS (ESI): m/z (%)=461 (M+H, 100);

[0681] HPLC (method 4): rt=2.89 min

Example 145

5-Chloro-N-[(3-{4-[2-imino-2-[(2,2,2-trifluoroethyl)amino]ethyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0682] MS (ESI): m/z (%)=475 (M+H, 100);

[0683] HPLC (method 4): rt=2.79 min

Example 146

N-({3-[4-(2-Anilino-2-iminoethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-5-chloro-2-thiophenecarboxamide

[0684] MS (ESI): m/z (%)=469 (M+H, 100);

[0685] HPLC (method 4): rt=2.83 min

Example 147

5-Chloro-N-[(3-{4-[2-imino-2-(2-pyridinylamino)ethyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide


[0686] MS (ESI): m/z (%)=470 (M+H, 100);

[0687] HPLC (method 4): rt=2.84 min

[0688] Examples 148 to 151 which follow relate to the elimination of BOC amino protective groups:

General Method for Eliminating Boc Protective Groups (Tert-Butyloxycarbonyl):

[0689]



[0690] Aqueous trifluoroacetic acid (TFA, approx. 90%) is added dropwise to an ice-cooled solution of a tert-butyloxycarbonyl-(Boc)-protected compound in chloroform or dichloromethane (approx. 0.1 to 0.3 mol/l). After about 15 min, the ice cooling is removed and the mixture is stirred at room temperature for about 2-3 h before the solution is con-

centrated and dried under high vacuum. The residue is taken up in dichloromethane or dichloromethane/methanol and washed with saturated sodium bicarbonate or 1N sodium hydroxide solution. The organic phase is washed with saturated sodium chloride solution, dried over a little magnesium sulfate and concentrated. Purification takes place where appropriate by crystallization from ether or ether/dichloromethane mixtures.

[0691] The following were prepared in an analogous manner from the appropriate Boc-protected precursors:

Example 148

N-({3-[4-(Aminomethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-5-chloro-2-thiophenecarboxamide

[0692] starting from Example 92:

[0693] MS (ESI): m/z (%)=349 (M—NH₂, 25), 305 (100);

[0694] HPLC (method 1): rt (%)=3.68 (98).

[0695] IC₅₀: 2.2 μM

Example 149


N-({3-[4-(Aminophenyl)-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide

[0696] starting from Example 93:

[0697] MS (ESI): m/z (%)=352 (M+H, 25); HPLC (method 1): rt (%)=3.50 (100).

[0698] IC₅₀: 2 μM

[0699] An enantiopure alternative synthesis of this compound is depicted in the following scheme (cf. also Delalande S. A., DE 2836305, 1979; Chem. Abstr. 90, 186926):



Example 150

5-Chloro-N-({3-[4-(glycylamino)phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0700] starting from Example 152:

[0701] MS (ES-pos): m/z (%)=408 (100); HPLC (method 3): rt (%)=3.56 (97).

IC₅₀: 2 μM

Example 151

5-(Aminomethyl)-3-[4-(2-oxo-1-pyrrolidinyl)phenyl]-1,3-oxazolidin-2-one

[0702] starting from Example 60:

[0703] MS (ESI): m/z (%)=276 (M+H, 100);

[0704] HPLC (method 3): rt (%)=2.99 (100).


[0705] IC₅₀: 2 μM

[0706] Examples 152 to 166 which follow relate to the amino group-derivatization of aniline- or benzylamine-substituted oxazolidinones with various reagents:

Example 152

5-Chloro-N-({3-[4-(N-tert-butyloxycarbonyl-glycylamino)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0707]



[0708] 754 mg (2.1 mmol) of N-{{3-[4-(aminophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide (from Example 149) are added to a solution of 751 mg (4.3 mmol) of Boc-glycine, 870 mg (6.4 mmol) of HOBt (1-hydroxy-1H-benzotriazol \times H₂O), 1790 mg (4.7 mmol) of HBTU [O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate] and 1.41 ml (12.9 mmol) of N-methylmorpholine in 15 ml of DMF/CH₂Cl₂ (1:1) at 0° C. The mixture is stirred at room temperature overnight before being diluted with water. The precipitated solid is filtered off and dried. Yield: 894 mg (79.7% of theory);


[0709] MS (DCI, NH₃): m/z (%)=526 (M+NH₄, 100);

[0710] HPLC (method 3): rt (%)=4.17 (97).

Example 153

N-{{3-[4-((Acetylamino)methyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide

[0711]



[0712] Acetic anhydride (0.015 ml, 0.164 mmol) is added to a mixture of 30 mg (0.082 mmol) of N-{{3-[4-(aminomethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide (from Example 148) in 1.5 ml of absolute THF and 1.0 ml of absolute dichloromethane, 0.02 ml of absolute pyridine at 0° C. The mixture is stirred at room

temperature overnight. The product is obtained after addition of ether and crystallization. Yield: 30 mg (87% of theory),

[0713] MS (ESI): m/z (%)=408 (M+H, 18), 305 (85);


[0714] HPLC (method 1): rt (%)=3.78 (97).

[0715] IC₅₀: 0.6 μM

Example 154

N-{{3-[4-{{[(Aminocarbonyl)amino]methyl}phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide

[0716]




[0717] 0.19 ml (0.82 mmol) of trimethylsilyl isocyanate is added dropwise to a mixture of 30 mg (0.082 mmol) of N-{{3-[4-(aminomethyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide (from Example 148) in 1.0 ml of dichloromethane at room temperature. The mixture is stirred overnight before, after addition of ether, the product is obtained by filtration. Yield: 21.1 mg (52% of theory),

[0718] MS (ESI): m/z (%)=409 (M+H, 5), 305 (72);

[0719] HPLC (method 1): rt (%)=3.67 (83).

[0720] IC₅₀: 1.3 μM

General method for acylating N-{{3-[4-(aminophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide with carbonyl chlorides:



[0721] An approx. 0.1 molar solution of N-{{3-[4-(aminophenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl}-5-chloro-2-thiophenecarboxamide (from Example 149) (1.0 eq.) in absolute dichloromethane/pyridine (19:1) is added dropwise under argon to the appropriate acid chloride (2.5 eq.). The mixture is stirred overnight before addition of approx. 5 eq of

PS-trisamine (Argonaut Technologies) and 2 ml of absolute dichloromethane. Gentle stirring for 1 h is followed by filtration and concentration of the filtrate. The products are purified where appropriate by preparative RP-HPLC.

[0722] The following were prepared in an analogous manner:

Example 155

N-({3-[4-(Acetylamino)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0723] LC-MS: m/z (%)=394 (M+H, 100);

[0724] LC-MS (method 6): rt (%)=3.25 (100).

[0725] IC₅₀: 1.2 μM

Example 156

5-Chloro-N-[(2-oxo-3-{4-[(2-thienylcarbonyl)amino]phenyl}-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0726] LC-MS: m/z (%)=462 (M+H, 100);

[0727] LC-MS (method 6): rt (%)=3.87 (100).

[0728] IC₅₀: 1.3 μM

Example 157

5-Chloro-N-[(3-{4-[(methoxyacetyl)amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)-methyl]-2-thiophenecarboxamide

[0729] LC-MS: m/z (%)=424 (M+H, 100);

[0730] LC-MS (method 6): rt (%)=3.39 (100).

[0731] IC₅₀: 0.73 μM

Example 158

N-{4-[5-({[(5-Chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-3,5-dimethyl-4-isoxazolecarboxamide


[0732] LC-MS: m/z (%)=475 (M+11, 100).

[0733] IC₅₀: 0.46 μM

Example 159

5-Chloro-N-({3-[4-[(3-chloropropyl)sulfonyl]amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl}-2-thiophenecarboxamide

[0734]



[0735] 35 mg (0.1 mmol) of N-({3-[4-(aminophenyl)-2-oxo-1,3-oxazolidin-5-yl]-methyl}-5-chloro-2-thiophenecarboxamide (from Example 149) are added to an ice-cooled solution of 26.4 mg (0.15 mmol) of 3-chloro-1-propanesulfonyl chloride and 0.03 ml (0.2 mmol) of triethylamine in 3.5 ml of absolute dichloromethane. After 30 min, the ice cooling is

removed and the mixture is stirred at room temperature overnight before adding 150 mg (approx. 5.5 eq) of PS-trisamine (Argonaut Technologies) and 0.5 ml of dichloromethane. The suspension is stirred gently for 2 h and filtered (the resin is washed with dichloromethane/methanol), and the filtrate is concentrated. The product is purified by preparative RP-HPLC. Yield: 19.6 mg (40% of theory).

[0736] LC-MS: m/z (%)=492 (M+H, 100);


[0737] LC-MS (method 5): rt (%)=3.82 (91).

[0738] IC₅₀: 1.7 μM

Example 160

5-Chloro-N-({3-[4-(1,1-dioxido-2-isothiazolidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide

[0739]



[0740] A mixture of 13.5 mg (0.027 mmol) of 5-chloro-N-({3-[4-({[(3-chloropropyl)sulfonyl]amino}phenyl)-2-oxo-1,3-oxazolidin-5-yl]methyl}-2-thiophenecarboxamide (from Example 159) and 7.6 mg (0.055 mmol) of potassium carbonate in 0.2 ml of DMF is heated at 100° C. for 2 h. Cooling is followed by dilution with dichloromethane and washing with water. The organic phase is dried and concentrated. The residue is purified by preparative thin-layer chromatography (silica gel, dichloromethane/methanol, 95:5). Yield: 1.8 mg (14.4% of theory).

[0741] MS (ESI): m/z (%)=456 (M+H, 15), 412 (100);


[0742] LC-MS (method 4): rt (%)=3.81 (90).

[0743] IC₅₀: 0.14 μM

Example 161

5-Chloro-N-(((5S)-3-{4-[(5-chloropentanoyl)amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide

[0744]



[0745] 0.5 g (1.29 mmol) of N-({(5S)-3-(4-aminophenyl)-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide (from Example 149) is dissolved in 27 ml of tetrahydrofuran, and 0.2 g (1.29 mmol) of 5-chlorovaleryl chloride and 0.395 ml (2.83 mmol) of triethylamine are added. The mixture is evaporated in vacuo and chromato-


graphed on silica gel with a toluene/ethyl acetate=1:1->ethyl acetate gradient. 315 mg (52% of theory) of a solid are obtained.

[0746] M.p.: 211° C.

Example 162

5-Chloro-N-({(5S)-2-oxo-3-[4-(2-oxo-1-piperidiny)phenyl]-1,3-oxazolidin-5-yl]-methyl)-2-thiophenecarboxamide

[0747]



[0748] 30 mg of 60 percent NaH in liquid paraffin are added under inert conditions to 5 ml of DMSO, and the mixture is heated at 75° C. for 30 min until gas evolution ceases. Then a solution of 290 mg (0.617 mmol) of 5-chloro-N-[(5S)-3-{4-[5-chloropentanoyl]amino}phenyl]-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide (from Example 161) in 5 ml of methylene chloride is added dropwise, and the mixture is stirred at room temperature overnight. The reaction is stopped and the mixture is added to 100 ml of water and extracted with ethyl acetate. The evaporated organic phase is chromatographed on an RP-8 column and eluted with acetonitrile/water. 20 mg (7.5% of theory) of the target compound are obtained.

[0749] M.p.: 205° C.;


[0750] NMR (300 MHz, d_6 -DMSO): δ =1.85 (m, 4H), 2.35 (m, 2H), 3.58 (m, 4H), 3.85 (m, 1H), 4.2 (t, 1H), 4.82 (m, 1H), 7.18 (d, 1H, thiophene), 7.26 (d, 2H), 7.5 (d, 2H), 2.68 (d, 1H, thiophene), 9.0 (t, 1H, CONH).

[0751] IC_{50} : 2.8 nM

Example 163

5-Chloro-N-[(5S)-3-{4-[(3-bromopropionyl)amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0752]




is obtained in an analogous manner from Example 149.

Example 164

5-Chloro-N-({(5S)-2-oxo-3-[4-(2-oxo-1-azetidiny)phenyl]-1,3-oxazolidin-5-yl]-methyl)-2-thiophenecarboxamide

[0753]



is obtained in an analogous manner by cyclization of the open-chain bromopropionyl compound from Example 163 using NaH/DMSO.


[0754] MS (ESI): m/z (%)=406 ($[M+11]^+$, 100), Cl pattern.

[0755] IC_{50} : 380 nM

Example 165

tert-Butyl 4-{4-[5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-3,5-dioxo-1-piperazinecarboxylate

[0756]



[0757] 300 mg (0.85 mmol) of N-([3-(4-aminophenyl)-2-oxo-1,3-oxazolidin-5-yl]-methyl)-5-chloro-2-thiophenecarboxamide in 6 ml of a mixture of DMF and dichloromethane (1:1) are added to a solution of 199 mg (0.85 mmol) of Boc-iminodiacetic acid, 300 mg (2.2 mmol) of HOBt, 0.66 ml (6 mmol) of N-methylmorpholine and 647 mg (1.7 mmol) of HBTU. The mixture is stirred overnight before, after dilution with dichloromethane, being washed with water, saturated ammonium chloride solution, saturated sodium bicarbonate solution, water and saturated sodium chloride solution. The organic phase is dried over magnesium sulfate and concentrated. The crude product is purified by chromatography on silica gel (dichloromethane/methanol 98:2). Yield: 134 mg (29% of theory);

[0758] MS (ESI): m/z (%)=571 (M+Na, 82), 493 (100);


[0759] HPLC (method 3): rt (%)=4.39 (90).

[0760] IC_{50} : 2 μ M

Example 166

N-[(5S)-3-{4-[(3R)-3-Amino-2-oxo-1-pyrrolidinyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide trifluoroacetate

[0761]



N2-(tert-Butoxycarbonyl)-N1-{4-[(5S)-5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-D-methioninamide

tert-Butyl (3R)-1-{4-[(5S)-5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-2-oxo-3-pyrrolidinylcarbamate

[0762] 429 mg (1.72 mmol) of N-BOC-D-methionine, 605 mg (1.72 mmol) of N-[(5S)-3-(4-aminophenyl)-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide, and 527 mg (3.44 mmol) of HOBT hydrate are dissolved in 35 ml of DMF, and 660 mg (3.441 mmol) of EDCI hydrochloride and then, dropwise, 689 mg (5.334 mmol) of N-ethyl-diisopropylamine are added. The mixture is stirred at room temperature for two days. The resulting suspension is filtered with suction and the residue is washed with DMF. The combined filtrates are mixed with a little silica gel, evaporated in vacuo and chromatographed on silica gel with a toluene->T10EA7 gradient. 170 mg (17% of theory) of the target compound are obtained with a melting point of 183° C.

[0763] R_f (SiO₂, toluene/ethyl acetate=1:1):0.2.

[0764] ¹H-NMR (300 MHz, d₆-DMSO): δ=1.4 (s, 1H, BOC), 1.88-1.95 (m, 2H), 2.08 (s, 3H, SMe), 2.4-2.5 (m, 2H, partly covered by DMSO), 3.6 (m, 2H), 3.8 (m, 1H), 4.15 (m, 2H), 4.8 (m, 1H), 7.2 (1H, thiophene), 7.42 (d, part of an AB system, 2H), 7.6 (d, part of an AB system, 2H), 7.7 (d, 1H, thiophene), 8.95 (t, 1H, CH₂NHCO), 9.93 (bs, 1H, NH).

[0765] 170 mg (0.292 mmol) of N2-(tert-butoxycarbonyl)-N1-{4-[(5S)-5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-D-methioninamide are dissolved in 2 ml of DMSO, and 178.5 mg (0.875 mmol) of trimethylsulfonium iodide and 60.4 mg (0.437 mmol) of potassium carbonate are added, and the mixture is stirred at 80° C. for 3.5 hours. It is then evaporated under high vacuum, and the residue is washed with ethanol. 99 mg of the target compound remain.

[0766] ¹H-NMR (300 MHz, d₆-DMSO): δ=1.4 (s, 1H, BOC), 1.88-2.05 (m, 1H), 2.3-2.4 (m, 1H), 3.7-3.8 (m, 3H), 3.8-3.9 (m, 1H), 4.1-4.25 (m, 1H), 4.25-4.45 (m, 1H), 4.75-4.95 (m, 1H), 7.15 (1H, thiophene), 7.25 (d, 1H), 7.52 (d, part of an AB system, 2H), 7.65 (d, part of an AB system, 2H), 7.65 (d, 1H, thiophene), 9.0 (broad s, 1H).

N-[(5S)-3-{4-[(3R)-3-Amino-2-oxo-1-pyrrolidinyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide trifluoroacetate

[0767] 97 mg (0.181 mmol) of tert-butyl (3R)-1-{4-[(5S)-5-({[(5-chloro-2-thienyl)carbonyl]amino}methyl)-2-oxo-1,3-oxazolidin-3-yl]phenyl}-2-oxo-3-pyrrolidinylcarbamate

are suspended in 4 ml of methylene chloride and, after addition of 1.5 ml of trifluoroacetic acid, stirred at room temperature for 1 hour. The mixture is then evaporated in vacuo and purified on an RP-HPLC (acetonitrile/water/0.1% TFA gradient). Evaporation of the relevant fraction results in 29 mg (37% of theory) of the target compound with a melting point of 241° C. (decomposition).


[07768] R_f (SiO₂, EtOH/TEA=17:1) 0.19.

[07769] ¹H-NMR (300 MHz, d₆-DMSO): δ=1.92-2.2 (m, 1H), 2.4-2.55 (m, 1H, partially covered by DMSO peak), 3.55-3.65 (m, 2H), 3.75-3.95 (m, 3H), 4.1-4.3 (m, 2H), 4.75-4.9 (m, 1H), 7.2 (1H, thiophene), 7.58 (d, part of an AB system, 2H), 7.7 (d, part of an AB system, 2H), 7.68 (d, 3H, thiophene), 8.4 (broad s, 3H, NH₃), 8.9 (t, 1H, NHCO).

[07770] Examples 167 to 170 which follow relate to the introduction of sulfonamide groups into phenyl-substituted oxazolidinones:

General method for preparing substituted sulfonamides starting from 5-chloro-N-[(2-oxo-3-phenyl-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[07771]



[07772] 5-Chloro-N-[(2-oxo-3-phenyl-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide (from Example 96) is added to chlorosulfonic acid (12 eq.) under argon at 5° C. The reaction mixture is stirred at room temperature for 2 h and then added to ice-water. The precipitate which separates out is filtered, washed with water and dried.

[07773] It is then dissolved in tetrahydrofuran (0.1 mol/l) under argon at room temperature, and the appropriate amine (3 eq.), triethylamine (1.1 eq.) and dimethylaminopyridine (0.1 eq.) are added. The reaction mixture is stirred for 1-2 h and then concentrated in vacuo. The desired product is purified by flash chromatography (dichloromethane/methanol mixtures).

[07774] The following were prepared in an analogous manner:

Example 167

5-Chloro-N-[(2-oxo-3-[4-(1-pyrrolidinyl)sulfonyl]phenyl]-1,3-oxazolidin-5-yl)-methyl]-2-thiophenecarboxamide

[07775] MS (ESI): m/z (%)=492 ([M+Na]⁺, 100), 470 ([M+H]⁺, 68), Cl pattern;

[07776] HPLC (method 3): rt (%)=4.34 (100).

[07777] IC₅₀: 0.5 μM

Example 168

5-Chloro-N-[(3-{4-[(4-methyl-1-piperazinyl)sulfonyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[07778] MS (ESI): m/z (%)=499 ([M+H]⁺, 100), Cl pattern;

[07779] HPLC (method 2): rt (%)=3.3 (100).

Example 169

5-Chloro-N-[(2-oxo-3-[4-(1-piperidinyl)sulfonyl]phenyl]-1,3-oxazolidin-5-yl)-methyl]-2-thiophenecarboxamide

[07780] MS (ESI): m/z (%)=484 ([M+H]⁺, 100), Cl pattern;

[07781] HPLC (method 2): rt (%)=4.4 (100).

Example 170

5-Chloro-N-[(3-{4-[(4-hydroxy-1-piperidinyl)sulfonyl]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide


[07782] MS (ESI): m/z (%)=500 ([M+H]⁺, 100), Cl pattern;

[07783] HPLC (method 3): rt (%)=3.9 (100).

Example 171

5-Chloro-N-[(2-oxo-3-[4-(1-pyrrolidinyl)phenyl]-1,3-oxazolidin-5-yl)methyl]-2-thiophenecarboxamide

[0784]



[0785] 780 mg (1.54 mmol) of tert-butyl 1-{4-[5-[(5-chloro-2-thienyl)carbonyl]amino]-methyl}-2-oxo-1,3-oxazolidin-3-yl]phenyl]prolinate are dissolved in 6 ml of dichloromethane and 9 ml of trifluoroacetic acid, and the mixture is stirred at 40° C. for two days. The reaction mixture is then concentrated and stirred with ether and 2 N sodium

hydroxide-solution. The aqueous phase is concentrated and stirred with ether and 2 N hydrochloric acid. The organic phase from this extraction is dried over MgSO_4 , filtered and concentrated. The crude product is chromatographed on silica gel ($\text{CH}_2\text{Cl}_2/\text{EtOH}/\text{conc. aq. NH}_3$ solution=100/1/0.1 to 20/1/0.1).

[0786] 280 mg (40% of theory) of the product are obtained.

[0787] MS (ESI): m/z (%)=406 (M+H, 100);

[0788] HPLC (method 4): rt=3.81 min

HPLC Parameters and LC-MS Parameters for the HPLC and LC-MS Data Stated in the Preceding Examples (the Unit of Retention Time (Rt) is Minutes):

[0789] [1] Column: Kromasil C18, L-R temperature: 30° C., flow rate=0.75 mlmin⁻¹, eluent: A=0.01 M HClO_4 , B= CH_3CN , gradient:-> 0.5 min 98% A-> 4.5 min 10% A-> 6.5 min 10% A

[2] Column: Kromasil C18 60*2, L-R temperature: 30° C., flow rate=0.75 mlmin⁻¹, eluent: A=0.01 M H_3PO_4 , B= CH_3CN , gradient:-> 0.5 min 90% A-> 4.5 min 10% A-> 6.5 min 10% A

[3] Column: Kromasil C18 60*2, L-R temperature: 30° C., flow rate=0.75 mlmin⁻¹, eluent: A=0.005 M HClO_4 , B= CH_3CN , gradient:-> 0.5 min 98% A-> 4.5 min 10% A-> 6.5 min 10% A

[4] Column: Symmetry C18 2.1x150 mm, column oven: 50° C., flow rate=0.6 mlmin⁻¹, eluent: A=0.6 g of 30% $\text{HCl}/1$ of water, B= CH_3CN , gradient: 0.0 min 90% A-> 4.0 min 10% A-> 9 min 10% A

[5] MHZ-2Q, Instrument Micromass Quattro LCZ

[0790] Column Symmetry C18, 50 mmx2.1 mm, 3.5 μm , temperature: 40° C., flow rate=0.5 ml min⁻¹, eluent A $\text{CH}_3\text{CN}+0.1\%$ formic acid, eluent B=water+0.1% formic acid, gradient: 0.0 min 10% A-> 4 min 90% A-> 6 min 90% A

[6] MHZ-2P, Instrument Micromass Platform LCZ

[0791] Column Symmetry C18, 50 mmx2.1 mm, 3.5 μm , temperature: 40° C., flow rate=0.5 mlmin⁻¹, eluent A= $\text{CH}_3\text{CN}+0.1\%$ formic acid, eluent B=water+0.1% formic acid, gradient: 0.0 min 10% A-> 4 min 90% A-> 6 min 90% A

[7] MHZ-7Q, Instrument Micromass Quattro LCZ


[0792] Column Symmetry C18, 50 mmx2.1 mm, 3.5 μm , temperature: 40° C., flow rate=0.5 mlmin⁻¹, eluent A= $\text{CH}_3\text{CN}+0.1\%$ formic acid, eluent B=water+0.1% formic acid, gradient: 0.0 min 5% A-> 1 min 5% A-> 5 min 90% A-> 6 min 90% A


General Method for Preparing Oxazolidinones of the General Formula B by Solid Phase-Assisted Synthesis

[0793] Reactions with various resin-bound products took place in a set of separate reaction vessels.

[0794] 5-(Bromomethyl)-3-(4-fluoro-3-nitrophenyl)-1,3-oxazolidin-2-one A (prepared from epibromohydrin and 4-fluoro-3-nitrophenyl isocyanate with $\text{LiBr}/\text{Bu}_3\text{PO}$ in xylene in analogy to U.S. Pat. No. 4,128,654, Ex. 2) (1.20 g, 3.75 mmol) and ethyldiisopropylamine (DIEA, 1.91 ml, 4.13 mmol) were dissolved in DMSO (70 ml), mixed with a secondary amine (1.1 eq, amine component 1) and reacted at 55°

C. for 5 h. TentaGel SAM resin (5.00 g, 0.25 mmol/g) was added to this solution and reacted at 75° C. for 48 h. The resin was filtered and repeatedly washed with methanol (MeOH), dimethylformamide (DMF), MeOH, dichloromethane (DCM) and diethyl ether and dried. The resin (5.00 g) was suspended in dichloromethane (80 ml), mixed with DIEA (10 eq) and 5-chlorothiophene-2-carbonyl chloride [prepared by reacting 5-chlorothiophene-2-carboxylic acid (5 eq) and 1-chloro-1-dimethylamino-2-methylpropene (5 eq) in DCM (20 ml) at room temperature for 15 minutes] and reacted at room temperature for 5 h. The resulting resin was filtered and washed repeatedly with MeOH, DCM and diethyl ether and dried. The resin was then suspended in DMF/water (v/v 9:2, 80 ml), mixed with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (5 eq) and reacted at room temperature for 18 h. The resin was again washed repeatedly with MeOH, DMF, water, MeOH, DCM and diethyl ether and dried. This resin was suspended in DCM, mixed with DIEA (10 eq) and, at 0° C., with an acid chloride (5 eq of acid derivative 1) and reacted at room temperature overnight. Before the reaction, carboxylic acids were converted into the corresponding acid chlorides by reacting with 1-dimethylamino-1-chloro-2-methylpropene (1 eq, based on the carboxylic acid) in DCM at room temperature for 15 min. The resin was washed repeatedly with DMF, water, DMF, MeOH, DCM and diethyl ether and dried. Where Fmoc-protected amino acids were used as acid derivative 1, the Fmoc-protective group was eliminated in the last reaction step by reacting with piperidine/DMF (v/v, 1/4) at room temperature for 15 minutes, and the resin was washed with DMF, MeOH, DCM and diethyl ether and dried. The products were then cleaved off the solid phase with trifluoroacetic acid (TFA)/DCM (v/v, 1/1), the resin was filtered off, and the reaction solutions were evaporated. The crude products were filtered through silica gel (DCM/MeOH, 9:1) and evaporated in order to obtain a set of products B.






Compounds prepared by solid phase-assisted synthesis:

Example 172

N-({3-[3-Amino-4-(1-pyrrolidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0795]




[0796] 5 g (1.25 mmol) of TentaGel SAM resin were reacted with pyrrolidine as amine derivative 1 in analogy to the general procedure for preparing the derivatives B. The aniline obtained after reduction with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ was eliminated from the solid phase, without a further acylation step, and evaporated. The crude product was partitioned between ethyl acetate and NaHCO_3 solution, and the organic phase was salted out with NaCl , decanted and evaporated to dryness. This crude product was purified by vacuum flash chromatography on silica gel (dichloromethane/ethyl acetate, 3:1-1:2).

[0797] $^1\text{H-NMR}$ (300 MHz, CDCl_3): 1.95-2.08, br, 4H, 3.15-3.30, br, 4H, 3.65-3.81, m, 2H, 3.89, ddd, 1H, 4.05, dd, 1H, 4.81, dddd, 1H, 6.46, dd, 1H, 6.72, dd, 1H, 6.90, dd, 1H, 6.99, dd, 1H, 7.03, dd, 1H, 7.29, d, 1H.

Example 173

N-[(3-{3-(β -Alanyl amino)-4-[(3-hydroxypropyl) amino]phenyl}-2-oxo-1,3-oxazolidin-5-yl)methyl]-5-chloro-2-thiophenecarboxamide

[0798]




[0799] 5 g (1.25 mmol) of TentaGel SAM resin were reacted with azetidine as amine derivative 1 and Fmoc- β -alanine as acid derivative 1 in analogy to the general procedure for preparing the derivatives B. The crude product obtained after elimination was stirred in methanol at room temperature for 48 h and evaporated to dryness. This crude product was purified by reversed phase HPLC with a water/TFA/acetonitrile gradient.

[0800] $^1\text{H-NMR}$ (400 MHz, CD_3OD): 2.31, tt, 2H, 3.36, t, 2H, 3.54, t, 2H, 3.62, t, 2H, 3.72, dd, 1H, 3.79, dd, 1H, 4.01, dd, 1H, 4.29, dd, 2H, 4.43, t, 2H, 4.85-4.95, m, 1H, 7.01, d, 1H, 4.48-7.55, m, 2H, 7.61, d, 1H, 7.84, d, 1H.

Example 174

N-({4-[4-(3-Amino-1-pyrrolidinyl)-3-nitrophenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0801]




[0802] 130 mg (32.5 mmol) of TentaGel SAM resin were reacted with tert-butyl 3-pyrrolidinylcarbamate as amine derivative 1 in analogy to the general procedure for preparing the derivatives B. The nitrobenzene derivative obtained after acylation with 5-chlorothiophenecarboxylic acid was eliminated from the solid phase and evaporated. This crude product was purified by reversed phase HPLC with a water/TFA/acetonitrile gradient.

[0803] $^1\text{H-NMR}$ (400 MHz, CD_3OH): 2.07-2.17, m, 1H, 2.39-2.49, m, 1H, 3.21-3.40, m, 2H, 3.45, dd, 1H, 3.50-3.60, m, 1H, 3.67, dd, 1H, 3.76, dd, 1H, 3.88-4.00, m, 2H, 4.14-4.21, t, 1H, 4.85-4.95, m, 1H, 7.01, d, 1H, 7.11, d, 1H, 7.52, d, 1H, 7.66, dd, 1H; 7.93, d, 1H.

Example 175

N-({3-[3-amino-4-(1-piperidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0804]




[0805] 130 mg (32.5 mmol) of TentaGel SAM resin were reacted with piperidine as amine derivative 1 in analogy to the general procedure for preparing the derivatives B. The aniline obtained after reduction was eliminated, without a further acylation step, from the solid phase and evaporated. This crude product was purified by reversed phase HPLC with a water/TFA/acetonitrile gradient.

[0806] ¹H-NMR (400 MHz, CD₃OH): 1.65-1.75, m, 2H, 1.84-1.95, m, 4H, 3.20-3.28, m, 4H, 3.68, dd, 1H, 3.73, dd, 1H, 3.90, dd, 1H, 4.17, dd, 1H, 4.80-4.90, m, 1H, 7.00, d, 1H, 7.05, dd, 1H, 7.30-7.38, m, 2H, 7.50, d, 1H.

Example 176

N-({3-[3-(Acetylamino)-4-(1-pyrrolidinyl)phenyl]-2-oxo-1,3-oxazolidin-5-yl}methyl)-5-chloro-2-thiophenecarboxamide

[0807]








[0808] 130 mg (32.5 mmol) of TentaGel SAM resin were reacted with pyrrolidine as amine derivative 1 and acetyl chloride as acid derivative 1 in analogy to the general procedure for preparing the derivatives B. The crude product was partitioned between ethyl acetate and NaHCO₃ solution, and the organic phase was salted out with NaCl, decanted and evaporated to dryness. This crude product was purified by vacuum flash chromatography on silica gel (dichloromethane/ethyl acetate, 1:1-0:1).

[0809] ¹H-NMR (400 MHz, CD₃OH): 1.93-2.03, br, 4H, 2.16, s, 3H, 3.20-3.30, br, 4H; 3.70, d, 2H, 3.86, dd, 1H, 4.10, dd, 1H, 4.14, dd, 1H, 4.80-4.90, m, 1H, 7.00, d, 1H, 7.07, d, 1H, 7.31, dd, 1H, 7.51, d, 1H, 7.60, d, 1H.






[0810] The following compounds were prepared in analogy to the general procedure.

Example	Structure	Ret. time	HPLC [%]
177		2.62	79.7
178		2.49	33.7





-continued

Example	Structure	Ret. time	HPLC [%]
179		4.63	46.7
180		3.37	44.8
181		2.16	83
182		2.31	93.3
183		2.7	100





-continued

Example	Structure	Ret. time	HPLC [%]
184		3.91	51
185		2.72	75.2
186		3.17	46
187		4.61	50.2
188		3.89	56.6





-continued

Example	Structure	Ret. time	HPLC [%]
189		3.37	52.9
190		3.6	63.9
191		2.52	70.1
192		3.52	46.6






-continued

Example	Structure	Ret. time	HPLC [%]
193		2.87	50.1
194		3.25	71.1
195		2.66	67
196		2.4	52.1





-continued

Example	Structure	Ret. time	HPLC [%]
197		3.13	48.9
198		2.67	75.5
199		2.72	65.7
200		2.71	57.3






-continued

Example	Structure	Ret. time	HPLC [%]
201		2.22	100
202		3.89	75.7
203		3.19	49.6
204		2.55	88.2
205		2.44	68.6





-continued

Example	Structure	Ret. time	HPLC [%]
206		2.86	71.8
207		2.8	63.6
208		2.41	77
209		2.56	67.9





-continued

Example	Structure	Ret. time	HPLC [%]
210		3.67	78.4
211		2.54	69.8
212		3.84	59.2
213		2.41	67.8
214		2.41	75.4






-continued

Example	Structure	Ret. time	HPLC [%]
215		4.01	81.3
216		3.46	49.5
217		4.4	60.2
218		3.79	70.9






-continued

Example	Structure	Ret. time	HPLC [%]
219		4.57	51.5
220		2.68	100
221		4.53	63.5
222		2.66	89.2





-continued

Example	Structure	Ret. time	HPLC [%]
223		4.76	69.3
224		3.45	77.4
225		3.97	63.2
226		3.94	61.4
227		4.15	66.3






-continued

Example	Structure	Ret. time	HPLC [%]
228		4.41	55.1
229		2.83	41.1
230		2.7	83
231		4.39	64.2
232		4.85	74.9





-continued

Example	Structure	Ret. time	HPLC [%]
233		4.17	41
234		4.21	61.8
235		2.75	100
236		3.94	50





-continued

Example	Structure	Ret. time	HPLC [%]
237		4.65	75.8
238		4.4	75.3
239		4.24	62.2
240		4.76	75.1
241		4.17	72.5






-continued

Example	Structure	Ret. time	HPLC [%]
242		4.6	74.8
243		4.12	51.6
244		4.71	66.2
245		4.86	62

-continued

Example	Structure	Ret. time	HPLC [%]
246		5.23	58.3
247		4.17	72.4
248		3.35	59.6
249		2.41	60.3


-continued

Example	Structure	Ret. time	HPLC [%]
250		3.31	65.2
251		2.86	36.5
252		2.69	89.8
253		2.81	67.4
254		2.19	75.4


[0811] All products of the solid phase-assisted synthesis were characterized by LC-MS. The following separation system was routinely used for this: HP 1100 with 1N detector (208-400 nm), 40° C. oven temperature, Waters Symmetry C18 column (50 mm×2.1 mm, 3.5 μm), mobile phase A: 99.9% acetonitrile/0.1% formic acid, mobile phase B: 99.9% water/0.1% formic acid; gradient:

Time	A: %	B: %	Flow rate
0.00	10.0	90.0	0.50
4.00	90.0	10.0	0.50
6.00	90.0	10.0	0.50
6.10	10.0	90.0	1.00
7.50	10.0	90.0	0.50

[0812] The substances were detected by means of a Micro-mass Quattro LCZ MS, ionization: ESI positive/negative. The radical(s)



or —O present in the structures detailed above always mean a




or —OH function.

1-9. (canceled)

10. A method for the prophylaxis and/or treatment of thromboembolic disorders comprising administering to a subject in need thereof an effective amount of a combination comprising

A) at least one compound of the formula (I)



in which

R¹ is 2-thiophene which is substituted in position 5 by a radical selected from chlorine, bromine, methyl and trifluoromethyl,

R² is D-A-:

where:

the radical "A" is phenylene;

the radical "D" is a saturated 5- or 6-membered heterocycle which is linked via a nitrogen atom to "A", and

which has a carbonyl group directly adjacent to the linking nitrogen atom, and

in which a ring carbon member may be replaced by a heteroatom selected from the series S, N and O;

where

the group "A" defined above may optionally be substituted once or twice in the meta position relative to the linkage to the oxazolidinone by a radical selected from fluorine, chlorine, nitro, amino, trifluoromethyl, methyl and cyano,


R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are hydrogen,

or a pharmaceutically acceptable salt or hydrate thereof, or their mixture

and

B) at least one further active pharmaceutical ingredient.

11. The method of claim 10, wherein the compound of the formula (I) is 5-chloro-N-({(5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl}methyl)-2-thiophenecarboxamide of the formula




or a pharmaceutically acceptable salt or hydrate thereof, or their mixture.

12. The method of claim 10, wherein the further active pharmaceutical ingredient B) is a platelet aggregation inhibitor, anticoagulant, fibrinolytic, lipid-lowering agent, coronary therapeutic agent and/or vasodilator.

13. The method of claim 11, wherein the further active pharmaceutical ingredient B) is a platelet aggregation inhibitor, anticoagulant, fibrinolytic, lipid-lowering agent, coronary therapeutic agent and/or vasodilator.

14. A method for the prophylaxis and/or treatment of myocardial infarction, angina pectoris, sudden heart death, reocclusions and restenoses after angioplasty or aortocoronary bypass, stroke, transient ischemic attacks, peripheral arterial occlusive diseases, pulmonary embolisms or deep venous thromboses comprising administering to a subject in need thereof an effective amount of a combination comprising

A) at least one compound of the formula (I)



(I)

in which

R¹ is 2-thiophene which is substituted in position 5 by a radical selected from chlorine, bromine, methyl and trifluoromethyl,

R² is D-A-:

where:

the radical "A" is phenylene;

the radical "D" is a saturated 5- or 6-membered heterocycle which is linked via a nitrogen atom to "A", and

which has a carbonyl group directly adjacent to the linking nitrogen atom, and

in which a ring carbon member may be replaced by a heteroatom selected from the series S, N and O;

where

the group "A" defined above may optionally be substituted once or twice in the meta position relative to the linkage to the oxazolidinone by a radical selected from fluorine, chlorine, nitro, amino, trifluoromethyl, methyl and cyano,


R³, R⁴, R⁵, R⁶, R⁷ and R⁸ are hydrogen,

or a pharmaceutically acceptable salt or hydrate thereof, or their mixture

and

B) at least one further active pharmaceutical ingredient.

15. The method of claim 14, wherein the compound of formula (I) is 5-chloro-N-((5S)-2-oxo-3-[4-(3-oxo-4-morpholinyl)phenyl]-1,3-oxazolidin-5-yl)methyl)-2-thiophenecarboxamide of the formula



or a pharmaceutically acceptable salt or hydrate thereof, or their mixture.

16. The method of claim 14, wherein the further active pharmaceutical ingredient B) is a platelet aggregation inhibitor, anticoagulant, fibrinolytic, lipid-lowering agent, coronary therapeutic agent and/or vasodilator.

17. The method of claim 15, wherein the further active pharmaceutical ingredient B) is a platelet aggregation inhibitor, anticoagulant, fibrinolytic, lipid-lowering agent, coronary therapeutic agent and/or vasodilator.

18. The method of claim 14, wherein the method is for the prophylaxis and/or treatment of angina pectoris that is unstable angina.

19. The method of claim 15, wherein the method is for the prophylaxis and/or treatment of angina pectoris that is unstable angina.

* * * * *