

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2014364520 B2**

(54) Title  
**Methods and assays relating to circulating tumor cells**

(51) International Patent Classification(s)  
**A61K 38/00** (2006.01)      **G01N 33/00** (2006.01)  
**C12Q 1/68** (2006.01)

(21) Application No: **2014364520**      (22) Date of Filing: **2014.12.18**

(87) WIPO No: **WO15/095527**

(30) Priority Data

(31) Number <b>61/918,816</b>	(32) Date <b>2013.12.20</b>	(33) Country <b>US</b>	
	<b>61/937,883</b>	<b>2014.02.10</b>	<b>US</b>

(43) Publication Date: **2015.06.25**  
(44) Accepted Journal Date: **2020.01.02**

(71) Applicant(s)  
**The General Hospital Corporation**

(72) Inventor(s)  
**Ting, David T.;Haber, Daniel A.;Maheswaran, Shyamala**

(74) Agent / Attorney  
**Pizzeys Patent and Trade Mark Attorneys Pty Ltd, GPO Box 1374, BRISBANE, QLD, 4001, AU**

(56) Related Art  
**WO 2011112903 A2**  
**WO 2009051734 A1**  
**US 20120009582 A1**  
**WO 2013134649 A1**  
**ZALATNAI, Attila et al., Molecular Background of Chemoresistance in Pancreatic Cancer. *in vivo*, 1 January 2007, Retrieved from the Internet: URL:<http://iv.iiarjournals.org/content/21/2/339.full.pdf>**  
**WO 2013109944 A1**  
**SEUNG BAE RHO et al. Insulin-like growth factor-binding protein-5 (IGFBP-5) acts as a tumor suppressor by inhibiting angiogenesis. CARCINOGENESIS, 2008, vol. 29, No. 11, pages 2106-2111**  
**WO 2003006029 A1**  
**WO 2006089091 A2**  
**SIEUWERTS, Anieta M et al., mRNA and microRNA Expression Profiles in Circulating Tumor Cells and Primary Tumors of Metastatic Breast Cancer Patients. Clinical Cancer Research, 2011 v17(11) pages 3600-3618**  
**WO 2013117705 A1**

(43) International Publication Date  
25 June 2015 (25.06.2015)(51) International Patent Classification:  
*A61K 38/00* (2006.01)   *G01N 33/00* (2006.01)  
*C12Q 1/68* (2006.01)Road, Newton, Massachusetts 02467 (US). **MAHESWARAN, Shyamala**; 24 Eastern Ave., Lexington, Massachusetts 02421 (US).(21) International Application Number:  
PCT/US2014/071169(74) Agents: **RESNICK, David S.** et al.; Nixon Peabody LLP, 100 Summer Street, Boston, Massachusetts 02110 (US).(22) International Filing Date:  
18 December 2014 (18.12.2014)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(25) Filing Language: English  
(26) Publication Language: English  
(30) Priority Data:  
61/918,816 20 December 2013 (20.12.2013) US  
61/937,883 10 February 2014 (10.02.2014) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,

(71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit St., Boston, Massachusetts 02114 (US).

(72) Inventors: **TING, David T.**; 1 Bryant Ln, Dover, Massachusetts 02030 (US). **HABER, David**; 34 Monadnock*[Continued on next page]*

(54) Title: METHODS AND ASSAYS RELATING TO CIRCULATING TUMOR CELLS



(57) Abstract: The technology described herein relates to methods of detecting circulating tumor cells (CTCs), e.g. by detecting changes in the expression of certain CTC marker genes. Aberrant expression of CTC marker genes, e.g. changes in expression indicative of CTCs can also be targeted in order to treat cancer.

**FIG. 14D**



TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

— *with international search report (Art. 21(3))*

## METHODS AND ASSAYS RELATING TO CIRCULATING TUMOR CELLS

### CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application Nos. 61/918,816 filed December 20, 2013 and 61/937,883 filed February 10, 2014, the contents of which are incorporated herein by reference in their entirety.

### GOVERNMENT SUPPORT

[0002] This invention was made with federal funding under Grant Nos. 2R01CA129933 awarded by the National Institutes of Health. The U.S. government has certain rights in the invention.

### TECHNICAL FIELD

[0003] The technology described herein relates to the diagnosis and treatment of cancer.

### BACKGROUND

[0004] Circulating Tumor Cells (CTCs) are shed from primary tumors into the bloodstream, mediating the spread of cancer to distant organs (metastasis). Thus, the presence of circulating tumor cells (CTCs) in the bloodstream ultimately leads to spread of cancer to distant organs. However, CTCs are rare, estimated at one to ten tumor cells among ten billion normal blood cells in a milliliter of blood. As such, their isolation and molecular analysis has posed a significant technological challenge (Pantel et al., Nat Rev Cancer 2008 8:329-340; Yu et al., J Cell Biol 2011 192:373-382).

### SUMMARY

[0005] As described herein, the inventors have identified a number of genes, the expression of which is characteristic of CTCs. In particular, the expression of these genes differentiates CTCs from primary tumor cells. Accordingly, provided herein are methods and assays relating to the detection of CTCs, including diagnostic and prognostic methods and assays. Further, provided herein are treatments for cancer that target these markers of CTCs, e.g., to inhibit metastasis.

[0006] In one aspect, described herein is a method of detecting circulating tumor cells (CTCs) in a sample, the method comprising: measuring the level of a PC-CTC marker gene expression product in the sample; and determining that PC-CTCs are present if the detected level of the marker gene expression product is greater than a reference level. In some embodiments, the CTCs are pancreatic cancer CTCs. In some embodiments, the method further comprises a first step of isolating the CTCs from the sample. In some embodiments, the expression product is a nucleic acid. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization. In some embodiments, the expression product is a polypeptide. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of: Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunoassay (RIA); sandwich assay;

fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunometric assay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay. In some embodiments, the CTC marker gene is selected from Table 7 or Table 8. In some embodiments, the CTC marker gene is selected from the group consisting of: ABI3BP; ADAMTS5; ADAMTS1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

**[0007]** In one aspect, described herein is a method of treating cancer in a subject, the method comprising administering a therapeutically effective amount of a CTC marker gene-targeted therapy to the subject. In some embodiments, the cancer is pancreatic cancer. In some embodiments, the CTC marker gene-targeted therapy comprises an inhibitor of a CTC marker gene. In some embodiments, the inhibitor is an antibody reagent. In some embodiments, the inhibitor is an inhibitory nucleic acid reagent. In some embodiments, the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent and a chemotherapeutic agent. In some embodiments, the subject is a subject determined to have an elevated level of CTCs and/or an elevated level of a CTC marker gene present in the blood and/or stroma of the cancer.

**[0008]** In one aspect, described herein is a method of determining if a subject is likely to respond to treatment with a CTC marker gene-targeted therapy, the method comprising measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and determining that the subject is likely to respond to the treatment if the level of the expression product is increased relative to a reference level. In some embodiments, the method further comprises a first step of isolating the CTCs from the sample. In some embodiments, the cancer is pancreatic cancer. In some embodiments, the expression product is a nucleic acid. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization. In some embodiments, the expression product is a polypeptide. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of: Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunometric assay;

immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay. In some embodiments, the PC-CTC marker gene is selected from Table 7 or Table 8. In some embodiments, the CTC marker gene is selected from the group consisting of: ABI3BP; ADAMTS5; ADAMTS1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

**[0009]** In one aspect, described herein is a method of monitoring the treatment of a subject, the method comprising: administering a cancer therapy to a subject in need thereof; measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and determining that the subject is responding if the level of the CTC marker gene expression product is decreased relative to the reference level and determining that the subject is not responding to the treatment if the CTC marker gene expression product is not decreased relative to the reference level. In some embodiments, the cancer is pancreatic cancer. In some embodiments, the reference level is the level of the gene expression product in the patient prior to the administering step. In some embodiments, the method further comprises a first step of isolating the CTCs from the sample. In some embodiments, the expression product is a nucleic acid. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization. In some embodiments, the expression product is a polypeptide. In some embodiments, the level of the expression product is determined using a method selected from the group consisting of: Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunometric assay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay. In some embodiments, the PC-CTC marker gene is selected from Table 7 or Table 8. In some embodiments, the CTC marker gene is selected from the group consisting of: ABI3BP; ADAMTS5; ADAMTS1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2;

SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2. In some embodiments, the CTC marker gene is selected from the group consisting of: ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0010] Figs. 1A-1C demonstrate the isolation and characterization of CTCs. Fig. 1A depicts a schematic of CTC-iChip negative IFD system. Fig. 1B depicts a graph of mouse WBC depletion consistency between normal and cancer mouse models. WBC depletion shown in log10. Fig. 1C depicts a graph of CTC enumeration by immunofluorescent staining (CK+/CD45-/DAPI+) from normal and KPC mice.

[0011] Fig. 2 depicts schematics of principal component analysis of single cell samples.

[0012] Figs. 3A-3B demonstrate that epithelial, mesenchymal, and stem cell genes are differentially expressed in CTC-c cells vs Tumors. Depicted are boxplot of genes that are A) downregulated (Fig. 3A) and upregulated (Fig. 3B) in CTC-c cells vs Tumors. Bar = median, box plot = quartiles, scale in log10(rpm).

[0013] Figs. 4A-4C demonstrate CTC-iChip characterization. Fig. 4A depicts a graph of the percent of WBC deflected (y-axis) as a function of the number of anti-CD45 beads per WBC (x-axis). Fig. 4B depicts a graph of the recovery of mouse PDAC cell line NB508 spiked into normal mouse blood (4 independent experiments shown). Fig. 4C depicts a graph of the captured CTCs/mL of blood from syngeneic orthotopic PDAC tumors using NB508 cell line.

[0014] Fig. 5A depicts a table of KPC mouse genotype and characteristics. Fig. 5B depicts graphs of quality metrics of single cell sequencing with % of reads aligned and total unique alignments for cell lines (NB508, MEF), CTCs, WBC, and diluted bulk RNA from matched primary tumors. Fig. 5C depicts graphs of single cell heterogeneity using mean intra-cluster correlation coefficient for each cluster (rights) and between single cell primary tumor (TuGMP3), cancer cell line (NB508), and all CTCs (Cluster 1, 3, 4, 5, 9). Circle = mean, Range = 95% CI.

[0015] Fig. 6 depicts boxplot graphs of ECM protein gene enriched in CTC-c compared to bulk primary tumors and single cell primary tumors. Bar = median, boxplot – quartiles, scale in log 10 (rpm).

[0016] Fig. 7 depicts a heatmap expression profile of human pancreatic CTCs from 3 patients. Epithelial genes used to define CTCs and enriched extracellular proteins shown. Expression shown in log10 scale.

[0017] Fig. 8 depicts a graph of quantitative RT-PCR of SPARC expression in human pancreatic cancer cell lines.

**[0018]** Fig. 9 depicts invasion assays. Decreases in invasion through Matrigel of PDAC2 and PDAC 3 cell lines with shRNA against SPARC (ShF1 and ShF3) were observed. shNT = Non-target shRNA

**[0019]** Fig. 10 depicts a graph of the number of mice with detectable metastases by in vivo luciferase imaging in non-target shRNA (NT) and SPARC shRNA (SHF1).

**[0020]** Fig. 11 depicts a schematic of the process of determining CTC heterogeneity.

**[0021]** Figs. 12A-12C demonstrate that CTC-Enriched Genes are Found in Epithelial and Stromal Components of Primary Tumors. Depicted are expression boxplots of (Fig. 12A) Aldh1a2 stem cell and CTC highly enriched genes (Fig. 12B) Klf4 and (Fig. 12C) Igfbp5 genes. Bar = median, box plot = quartiles, scale in log10(rpm).

**[0022]** Fig. 13 demonstrates that human and mouse CTCs across different epithelial cancer express high levels of ECM protein genes. Depicted are expression boxplot of highly expressed ecm genes in human pdac, breast (br), and prostate (pr) ctc. bar, median; boxplot, quartiles; scale in log10(rpm). holm-adjusted p value < 0.05 (\*), 0.01 (\*\*), 0.001 (\*\*\*)�

**[0023]** Figs. 14A-14E demonstrate that SPARC expression in human PDAC enhances invasion and metastasis. Fig. 14A depicts a graph of proliferation of PDAC3 cell lines determined by MTT. Fig. 14B depicts a graph of tumor spheres in PDAC3 shNT versus shSPARC counted per 43 field (error bars represent SD). Fig. 14C depicts a graph of invasion of shSPARC and shNT cell lines quantitated by number of nuclei/203 field. p value < 0.01 (\*\*), 0.001 (\*\*\*)�, 0.0001 (\*\*\*\*). Error bars represent SD. Fig. 14D depicts a graph of Percentage of detectable lung metastases by in vivo luciferase imaging after 3 weeks after tail vein inoculation of PDAC3 cell lines. Fisher's exact test p value is shown. Fig. 14E depicts a graph of normalized metastasis burden in mice with orthotopic pancreatic tumors from PDAC3 cell lines. Error bars represent SD (\*p < 0.05).

**[0024]** Fig. 15 depicts a Summary Model of the Role of Pancreatic CTCs in the Metastatic Cascade. Shown are the heterogeneous subsets of pancreatic CTCs with a focus on the most prominent classical CTC group, which are enriched for coexpression of epithelial (keratin) and stromal (Sparc) genes.

**[0025]** Fig. 16A depicts a graph of PDAC2 shRNA cell lines by qRT-PCR. Average shown with max and min RQ (error bars). Fig. 16B depicts a graph of proliferation rates by MTT assay similar in PDAC2 cell line between shNT and shSPARC stable lines. Fig. 16C depicts a graph of tumor sphere invasion assay (error bars =STD) formation at 2 weeks similar between shNT and shSPARC cell lines. Quantiation done per 4x magnification field (Error bars = SD). Migratory behavior reduced by shSPARC\_1 & 3 as determined by (Fig. 16D) invasion assay at 48 hours.

## DETAILED DESCRIPTION

**[0026]** As described herein, the inventors have discovered that circulating tumor cells (CTCs) are characterized by the expression of certain genes, i.e. CTC marker genes. The discovery of these CTC marker genes permit methods and assays for the detection and/or measurement of CTC levels, e.g. CTC levels in a sample from a subject. These methods and assays can provide improved speed and accuracy in the measurement of CTC levels. Furthermore, because the expression of these marker genes distinguishes CTCs from other cells, e.g., other circulating cells and/or normal tumor cells, therapies can be targeted against CTCs by binding to and/or inhibiting these marker gene expression products to reduce the level and/or metastatic potential of CTCs.

**[0027]** As used herein, “circulating tumor cell” or “CTC” refers to tumor cells which are shed from a tumor and present in the blood, i.e. in circulation. Cell markers (e.g. marker genes) that can be used to identify and/or isolate CTCs from other components of the blood are described below herein. In some embodiments, a CTC can be a pancreatic cancer CTC.

**[0028]** In one aspect, described herein is a method of detecting circulating tumor cells (CTCs) in a sample, the method comprising measuring the level of a CTC marker gene expression product in the sample; and determining that CTCs are present if the detected level of the marker gene expression product is greater than a reference level.

**[0029]** As described herein, the inventors have discovered that a number of genes are differentially regulated in CTCs, e.g. as compared to non-circulating tumor cells. Accordingly, there are provided herein methods and assays relating to the measurement of CTC levels. Elevated CTC levels can indicate a poor prognosis, e.g. an increased risk of metastasis. Accordingly, provided herein are methods and assays related to the prognosis, risk assessment, and treatment of subjects having cancer. In certain embodiments, the assays and methods are directed to determination and/or measurement of the expression level of a gene product (e.g. protein and/or gene transcript such as mRNA) in a biological sample of a subject. In certain embodiments the assays and methods are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, i.e. at least two genes, at least three genes, at least four genes, at least five genes, at least six genes, at least seven genes, at least eight genes, at least nine genes, at least 10 genes...at least 15 genes,...at least 25 genes,...at least 30 genes, or more genes, or any number of genes selected from Table 7, Table 8, and/or Table 14 as described herein.

**[0030]** In some embodiments, the marker gene(s) is selected from the group consisting of ABI3BP; ADAMTS5; ADAMTSL1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4. In some embodiments, the assays, methods,

and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or at least four genes, or, e.g. all of the following genes: ABI3BP; ADAMTS5; ADAMTS1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4.

**[0031]** In some embodiments, the marker gene(s) is selected from the group consisting of ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2. In some embodiments, the assays, methods, and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or at least four genes, or, e.g. all of the following genes: ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2.

**[0032]** In some embodiments, the marker gene(s) is selected from the group consisting of ALDH1A2; IGFBP5; KLF4; DCN; and SPARC. In some embodiments, the assays, methods, and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or at least four genes, or, e.g. all of the following genes: ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

**[0033]** In some embodiments, the marker gene(s) is selected from the group consisting of ALDH1A2; IGFBP5; KLF4; and DCN. In some embodiments, the assays, methods, and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or e.g. all of the following genes: ALDH1A2; IGFBP5; KLF4; and DCN.

**[0034]** In some embodiments, the marker gene(s) is selected from the group consisting of TPT1; HMGB1; SPON 2; SPARC; and ARSA. In some embodiments, the assays, methods, and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or at least four genes, or, e.g. all of the following genes: TPT1; HMGB1; SPON 2; SPARC; and ARSA.

**[0035]** In some embodiments, the marker gene(s) is selected from the group consisting of IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A. In some embodiments, the assays, methods, and systems described herein are directed to determination of the expression level of a gene product of at least two genes in a biological sample of a subject, e.g. at least two genes, or at least three genes, or at least four genes, or at least five genes, or at least six genes, or

at least seven genes, or at least eight genes or, e.g. all of the following genes: IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A. In some embodiments, the level of polypeptide expression products are determined for the marker gene(s) is selected from the group consisting of IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A, e.g. because, as described herein, RNA levels of cell surface proteins are lower than polypeptide levels.

[0036] **Table 7: Exemplary mouse marker genes**

MOUSE GENE SYMBOL	Gene Name
Abcb1b	ATP-binding cassette, sub-family B (MDR/TAP), member 1B
Abi3bp	ABI gene family, member 3 (NESH) binding protein
Ablim3	actin binding LIM protein family, member 3
Acad9	acyl-Coenzyme A dehydrogenase family, member 9
Acbd3	acyl-Coenzyme A binding domain containing 3
Acin1	apoptotic chromatin condensation inducer 1
Actb	actin, beta
Actg1	predicted gene 8543; actin-like 8; predicted gene 7505; predicted gene 12715; predicted gene 12003; predicted gene 8399; predicted gene 6375; actin, gamma, cytoplasmic 1; similar to gamma-actin; predicted gene 4667; similar to cytoplasmic beta-actin; predicted gene 16385
Adamts5	similar to a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2); a disintegrin-like and metallopeptidase (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2)
Adamtsl1	ADAMTS-like 1
Add3	adducin 3 (gamma)
Aebp1	AE binding protein 1
Agap1	ArfGAP with GTPase domain, ankyrin repeat and PH domain 1
Akap13	A kinase (PRKA) anchor protein 13
Akap2	A kinase (PRKA) anchor protein 2; paralemmin 2
Akr1b3	aldo-keto reductase family 1, member B3 (aldose reductase)
Akt2	similar to RAC-beta serine/threonine-protein kinase (RAC-PK-beta) (Protein kinase Akt-2) (Protein kinase B, beta) (PKB beta); thymoma viral proto-oncogene 2; similar to serine/threonine kinase
Aldh1a1	aldehyde dehydrogenase family 1, subfamily A1
Aldh1a2	aldehyde dehydrogenase family 1, subfamily A2
Alox12	arachidonate 12-lipoxygenase
Amfr	autocrine motility factor receptor
Amhr2	anti-Mullerian hormone type 2 receptor
Ang	angiogenin, ribonuclease, RNase A family, 5
Ankrd11	ankyrin repeat domain 11

Ankrd12	ankyrin repeat domain 12; similar to Ankrd12 protein
Ankrd17	ankyrin repeat domain 17
Ano6	anoctamin 6
Anp32a	acidic (leucine-rich) nuclear phosphoprotein 32 family, member A
Anxa7	annexin A7
Ap1s3	predicted gene 8532; similar to adaptor-related protein complex AP-1, sigma 3; adaptor-related protein complex AP-1, sigma 3
Ap3s1	predicted gene 7603; adaptor-related protein complex 3, sigma 1 subunit; predicted gene 5610
Ap4e1	adaptor-related protein complex AP-4, epsilon 1
Aplp1	amyloid beta (A4) precursor-like protein 1
Apol9a	apolipoprotein L 9b; apolipoprotein L 9a
App	amyloid beta (A4) precursor protein
Aqp1	aquaporin 1
Arap2	predicted gene 336; ArfGAP with RhoGAP domain, ankyrin repeat and PH domain 2
Arf2	ADP-ribosylation factor 2
Arf3	ADP-ribosylation factor 3
Arf5	similar to ADP-ribosylation factor; ADP-ribosylation factor 5
Arhgap28	Rho GTPase activating protein 28
Arhgap29	Rho GTPase activating protein 29
Arhgap5	Rho GTPase activating protein 5
Arhgef12	predicted gene 7281; predicted gene 5831; similar to SP140 nuclear body protein (predicted); Rho guanine nucleotide exchange factor (GEF) 12
Arid1a	similar to AT rich interactive domain 1A isoform a; AT rich interactive domain 1A (SWI-like)
Arid4a	AT rich interactive domain 4A (RBP1-like)
Arid4b	AT rich interactive domain 4B (RBP1-like)
Arid5b	similar to modulator recognition factor 2; AT rich interactive domain 5B (MRF1-like)
Arl3	ADP-ribosylation factor-like 3
Arl4d	ADP-ribosylation factor-like 4D; hypothetical protein LOC100044157
Arl6ip5	ADP-ribosylation factor-like 6 interacting protein 5
Armcx3	armadillo repeat containing, X-linked 3; hypothetical protein LOC100044266; predicted gene 9299
Arpc2	predicted gene 5492; actin related protein 2/3 complex, subunit 2
Arsa	arylsulfatase A
Arsb	arylsulfatase B
Ascc3	activating signal cointegrator 1 complex subunit 3
Atf3	activating transcription factor 3
Atg3	autophagy-related 3 (yeast)

Atp1a1	ATPase, Na+/K+ transporting, alpha 1 polypeptide
Atp1b1	ATPase, Na+/K+ transporting, beta 1 polypeptide
Atp2b1	ATPase, Ca++ transporting, plasma membrane 1
Atp6v1a	ATPase, H+ transporting, lysosomal V1 subunit A
Atxn2	ataxin 2
B230120H23Rik	RIKEN cDNA B230120H23 gene
B2m	beta-2 microglobulin
BC003331	similar to odorant response abnormal 4; cDNA sequence BC003331
BC005537	cDNA sequence BC005537
BC005561	THO complex 2; cDNA sequence BC005561
BC013529	cDNA sequence BC013529
Baz2a	bromodomain adjacent to zinc finger domain, 2A
Bbs4	Bardet-Biedl syndrome 4 (human)
Bbx	bobby sox homolog (Drosophila)
Bcam	basal cell adhesion molecule
Bcl10	B-cell leukemia/lymphoma 10; predicted gene 6141
Bdp1	B double prime 1, subunit of RNA polymerase III transcription initiation factor IIIB
Bicc1	bicaudal C homolog 1 (Drosophila)
Bicd1	bicaudal D homolog 1 (Drosophila)
Birc6	baculoviral IAP repeat-containing 6
Blvrb	biliverdin reductase B (flavin reductase (NADPH))
Bnc1	basonuclin 1
Bnc2	basonuclin 2
Bod1l	biorientation of chromosomes in cell division 1-like
Bptf	bromodomain PHD finger transcription factor
Braf	Braf transforming gene
Brd2	similar to mKIAA4005 protein; bromodomain containing 2
Brd4	bromodomain containing 4
Brp44l	similar to brain protein 44-like protein; brain protein 44-like; predicted gene 3452; predicted gene 8219
Bst2	bone marrow stromal cell antigen 2
Btbd2	BTB (POZ) domain containing 2
Btbd7	BTB (POZ) domain containing 7
Btf3	predicted gene 9308; basic transcription factor 3; predicted gene 3531; predicted gene 7973
Btg2	B-cell translocation gene 2, anti-proliferative
Bzw1	predicted gene 11652; predicted gene 5191; basic leucine zipper and W2 domains 1
C1d	C1D nuclear receptor co-repressor
C1ra	complement component 1, r subcomponent; predicted gene 8551
C1rl	complement component 1, r subcomponent-like

C1s	similar to Complement component 1, s subcomponent; complement component 1, s subcomponent
C2	complement component 2 (within H-2S)
C3	complement component 3; similar to complement component C3 prepropeptide, last
C4a	similar to Complement C4 precursor; complement component 4A (Rodgers blood group); similar to complement C4; complement component 4B (Chido blood group)
C4b	similar to Complement C4 precursor; complement component 4A (Rodgers blood group); similar to complement C4; complement component 4B (Chido blood group)
Calm1	predicted gene 7743; calmodulin 3; calmodulin 2; calmodulin 1; predicted gene 7308
Calm2	predicted gene 7743; calmodulin 3; calmodulin 2; calmodulin 1; predicted gene 7308
Cap1	CAP, adenylate cyclase-associated protein 1 (yeast)
Cast	calpastatin
Cav1	caveolin 1, caveolae protein
Ccdc109b	coiled-coil domain containing 109B
Ccdc34	coiled-coil domain containing 34
Ccdc80	coiled-coil domain containing 80
Ccdc88a	coiled coil domain containing 88A
Ccdc90a	coiled-coil domain containing 90A
Ccnl1	cyclin L1
Cd109	CD109 antigen
Cd200	CD200 antigen; similar to MRC OX-2 antigen homolog
Cd248	CD248 antigen, endosialin
Cd34	CD34 antigen
Cd55	CD55 antigen
Cd81	CD81 antigen
Cd82	CD82 antigen
Cd9	CD9 antigen
Cdc42ep3	CDC42 effector protein (Rho GTPase binding) 3
Cdh11	cadherin 11
Cdh3	cadherin 3
Cdk13	cell division cycle 2-like 5 (cholinesterase-related cell division controller)
Cdon	cell adhesion molecule-related/down-regulated by oncogenes
Celf2	CUG triplet repeat, RNA binding protein 2
Cep164	centrosomal protein 164
Cep57	centrosomal protein 57

Cfh	complement component factor h; similar to complement component factor H
Cfl1	cofilin 1, non-muscle; similar to Cofilin-1 (Cofilin, non-muscle isoform); predicted gene 6180
Cfl2	cofilin 2, muscle
Chd1	chromodomain helicase DNA binding protein 1
Chd2	chromodomain helicase DNA binding protein 2
Chi3l1	chitinase 3-like 1
Chst4	carbohydrate (chondroitin 6/keratan) sulfotransferase 4
Cish	cytokine inducible SH2-containing protein
Clcn3	chloride channel 3
Cldn15	claudin 15
Cldn25	predicted gene 16492
Clec1b	C-type lectin domain family 1, member b
Clec3b	C-type lectin domain family 3, member b
Clic4	chloride intracellular channel 4 (mitochondrial)
Clip1	CAP-GLY domain containing linker protein 1
Clip3	CAP-GLY domain containing linker protein 3
Cln8	ceroid-lipofuscinosis, neuronal 8
Cmah	cytidine monophospho-N-acetylneuraminc acid hydroxylase
Cmtm3	CKLF-like MARVEL transmembrane domain containing 3
Cmtm7	CKLF-like MARVEL transmembrane domain containing 7
Cnot6l	CCR4-NOT transcription complex, subunit 6-like
Cobl	cordon-bleu
Cobl1	Cobl-like 1
Col14a1	collagen, type XIV, alpha 1
Col1a2	collagen, type I, alpha 2
Col3a1	collagen, type III, alpha 1
Col4a6	collagen, type IV, alpha 6
Colec12	collectin sub-family member 12
Coq10b	hypothetical protein LOC675736; coenzyme Q10 homolog B ( <i>S. cerevisiae</i> ); predicted gene 4899
Creb3l1	cAMP responsive element binding protein 3-like 1
Creb5	RIKEN cDNA 9430076C15 gene; cAMP responsive element binding protein 5
Crebbp	CREB binding protein
Creg1	cellular repressor of E1A-stimulated genes 1
Crim1	cysteine rich transmembrane BMP regulator 1 (chordin like)
Crlns1	cardiolipin synthase 1
Cryab	crystallin, alpha B
Cryl1	crystallin, lambda 1
Crym	crystallin, mu

Csda	cold shock domain protein A
Csf1	colony stimulating factor 1 (macrophage)
Csnk1a1	casein kinase 1, alpha 1
Csrnp1	cysteine-serine-rich nuclear protein 1
Csrp1	cysteine and glycine-rich protein 1
Cuedc1	CUE domain containing 1
Cyb5	cytochrome b-5
Cybrd1	cytochrome b reductase 1
Cyp2d22	cytochrome P450, family 2, subfamily d, polypeptide 22
Cyp2s1	cytochrome P450, family 2, subfamily s, polypeptide 1
Cyr61	cysteine rich protein 61
Dab2	disabled homolog 2 (Drosophila)
Dag1	dystroglycan 1
Dagl <sup>b</sup>	diacylglycerol lipase, beta
Dapk1	death associated protein kinase 1
Dcn	decorin
Ddr1	discoidin domain receptor family, member 1
Ddr2	discoidin domain receptor family, member 2
Ddx3x	DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 3, X-linked
Ddx5	DEAD (Asp-Glu-Ala-Asp) box polypeptide 5; predicted gene 12183
Dennd5a	DENN/MADD domain containing 5A; similar to Rab6 interacting protein 1
Dhx15	DEAH (Asp-Glu-Ala-His) box polypeptide 15
Diap1	diaphanous homolog 1 (Drosophila)
Dlgap4	discs, large homolog-associated protein 4 (Drosophila)
Dmkn	dermokine
Dnaja2	Dnaj (Hsp40) homolog, subfamily A, member 2
Dnajb9	predicted gene 6568; Dnaj (Hsp40) homolog, subfamily B, member 9
Dnajc1	Dnaj (Hsp40) homolog, subfamily C, member 1
Dnmt1	DNA methyltransferase (cytosine-5) 1
Dpp4	dipeptidylpeptidase 4
Dpysl2	dihydropyrimidinase-like 2
Dpysl3	dihydropyrimidinase-like 3
Dst	dystonin; hypothetical protein LOC100047109
Dtx2	deltex 2 homolog (Drosophila)
Dusp1	dual specificity phosphatase 1
Dusp14	dual specificity phosphatase 14
Dusp3	dual specificity phosphatase 3 (vaccinia virus phosphatase VH1-related)
Dync1i2	dynein cytoplasmic 1 intermediate chain 2
Ecd	ecdysoneless homolog (Drosophila)

Eea1	early endosome antigen 1
Eef1a1	predicted gene 5869; predicted gene 7161; predicted gene 7105; predicted gene 5822; similar to eukaryotic translation elongation factor 1 alpha 1; predicted gene 6192; predicted gene 6392; predicted gene 6767; predicted gene 6170; predicted gene 6548; predicted gene 6789; eukaryotic translation elongation factor 1 alpha 1
Efemp1	epidermal growth factor-containing fibulin-like extracellular matrix protein 1
Efhd2	similar to EF hand domain containing 2; EF hand domain containing 2
Efna5	ephrin A5
Egr1	early growth response 1
Ehd2	EH-domain containing 2
Eif2s3x	eukaryotic translation initiation factor 2, subunit 3, structural gene X-linked; similar to translation initiation factor eIF-2 gamma subunit; predicted gene 2223
Eif3a	eukaryotic translation initiation factor 3, subunit A
Elf1	E74-like factor 1
Elov16	predicted gene 11295; ELOVL family member 6, elongation of long chain fatty acids (yeast)
Emp2	epithelial membrane protein 2
Enpp2	ectonucleotide pyrophosphatase/phosphodiesterase 2
Enpp4	ectonucleotide pyrophosphatase/phosphodiesterase 4
Esam	endothelial cell-specific adhesion molecule
Esf1	ESF1, nucleolar pre-rRNA processing protein, homolog (S. cerevisiae)
Espn	espin
Esyt3	family with sequence similarity 62 (C2 domain containing), member C
Etfa	predicted gene 2893; electron transferring flavoprotein, alpha polypeptide
Evpl	envoplakin
Exoc4	exocyst complex component 4
F11r	F11 receptor
Faim2	Fas apoptotic inhibitory molecule 2
Fam117a	family with sequence similarity 117, memberA
Fam134b	family with sequence similarity 134, member B
Fam53b	family with sequence similarity 53, member B
Fam63b	RIKEN cDNA B230380D07 gene
Fam76a	predicted gene 7527; family with sequence similarity 76, member A
Fam84b	RIKEN cDNA D330050I23 gene

Fas	Fas (TNF receptor superfamily member 6)
Fbln1	fibulin 1
Fermt2	fermitin family homolog 2 (Drosophila)
Fgf1	fibroblast growth factor 1
Fhl1	four and a half LIM domains 1
Filip1l	filamin A interacting protein 1-like
Fkbp5	FK506 binding protein 5
Flii	flightless I homolog (Drosophila); similar to cytoskeletal actin-modulating protein
Flnc	filamin C, gamma
Flrt2	fibronectin leucine rich transmembrane protein 2
Fmo2	flavin containing monooxygenase 2
Fmod	fibromodulin
Fndc1	fibronectin type III domain containing 1; similar to fibronectin type III domain containing 1
Fos	FBJ osteosarcoma oncogene
Foxn3	forkhead box N3
Frmd4b	FERM domain containing 4B
Fth1	ferritin heavy chain 1
Fxyd1	FXYD domain-containing ion transport regulator 1
G3bp1	Ras-GTPase-activating protein SH3-domain binding protein 1
Gabarapl1	gamma-aminobutyric acid (GABA) A receptor-associated protein-like 1
Gadd45b	growth arrest and DNA-damage-inducible 45 beta
Ganab	alpha glucosidase 2 alpha neutral subunit
Gas1	growth arrest specific 1
Gas6	growth arrest specific 6
Gata6	GATA binding protein 6
Gbp2	guanylate binding protein 2
Gbp3	guanylate binding protein 3
Gcap14	granule cell antiserum positive 14
Gcsh	predicted gene 3672; similar to Glycine cleavage system H protein, mitochondrial precursor; glycine cleavage system protein H (aminomethyl carrier)
Gda	guanine deaminase
Gem	GTP binding protein (gene overexpressed in skeletal muscle)
Gfm2	G elongation factor, mitochondrial 2
Gfpt2	glutamine fructose-6-phosphate transaminase 2
Gja1	gap junction protein, alpha 1
Gjb5	gap junction protein, beta 5
Gm10052	predicted gene 10052

Gm13251	predicted gene 13251; predicted gene, OTTMUSG00000010657; RIKEN cDNA 1700029I01 gene
Gm3893	similar to 4933409K07Rik protein; predicted gene, 665845; predicted gene 2490; predicted gene 10601; predicted gene 2163; predicted gene 3892; RIKEN cDNA 4933409K07 gene; predicted gene 3893
Gm6548	predicted gene 5869; predicted gene 7161; predicted gene 7105; predicted gene 5822; similar to eukaryotic translation elongation factor 1 alpha 1; predicted gene 6192; predicted gene 6392; predicted gene 6767; predicted gene 6170; predicted gene 6548; predicted gene 6789; eukaryotic translation elongation factor 1 alpha 1
Gm6578	predicted gene 6578
Gm6644	predicted gene 6644
Gm9199	predicted gene 9199
Gnb2	guanine nucleotide binding protein (G protein), beta 2
Golga4	golgi autoantigen, golgin subfamily a, 4
Golgb1	golgi autoantigen, golgin subfamily b, macrogolgin 1
Gpc3	glypican 3
Gpc4	glypican 4; similar to Glypican 4
Gpcpd1	preimplantation protein 4
Gpm6a	glycoprotein m6a
Gpr116	G protein-coupled receptor 116
Gpr133	G protein-coupled receptor 133
Gpr64	G protein-coupled receptor 64
Gprc5b	G protein-coupled receptor, family C, group 5, member B
Gpx8	glutathione peroxidase 8 (putative)
Gsr	similar to Glutathione reductase, mitochondrial precursor (GR) (GRase); glutathione reductase
Gsta3	glutathione S-transferase, alpha 3
Gstm1	similar to Glutathione S-transferase Mu 1 (GST class-mu 1) (Glutathione S-transferase GT8.7) (pmGT10) (GST 1-1); predicted gene 5562; glutathione S-transferase, mu 1
Gstm4	glutathione S-transferase, mu 4
Gucy1a3	guanylate cyclase 1, soluble, alpha 3
H2-D1	histocompatibility 2, D region; histocompatibility 2, D region locus 1
H2-K1	histocompatibility 2, K1, K region; similar to H-2K(d) antigen

H2-Q6	histocompatibility 2, Q region locus 1; histocompatibility 2, Q region locus 9; similar to H-2 class I histocompatibility antigen, L-D alpha chain precursor; histocompatibility 2, Q region locus 8; histocompatibility 2, Q region locus 2; similar to MHC class Ib antigen; histocompatibility 2, Q region locus 7; histocompatibility 2, Q region locus 6; hypothetical protein LOC100044307; similar to H-2 class I histocompatibility antigen, Q7 alpha chain precursor (QA-2 antigen); RIKEN cDNA 0610037M15 gene
H3f3a	predicted gene 14383; predicted gene 3835; predicted gene 14384; predicted gene 12950; predicted gene, 670915; H3 histone, family 3A; predicted gene 12657; predicted gene 6132; predicted gene 10257; predicted gene 7227; H3 histone, family 3B; predicted gene 6128; similar to histone; predicted gene 1986; predicted gene 6186; hypothetical protein LOC676337; predicted gene 6421; predicted gene 2198; predicted gene 6817; predicted gene 8095; predicted gene 12271; predicted gene 13529; predicted gene 8029; predicted gene 4938; predicted gene 7100; predicted gene 9014; similar to Histone H3.4 (Embryonic); predicted gene 7179; similar to H3 histone, family 3B; predicted gene 7900; predicted gene 2099; similar to H3 histone, family 3A; predicted gene 6749; predicted gene 6485; predicted gene 4028; predicted gene 7194
Hdac3	histone deacetylase 3
Hdac5	histone deacetylase 5
Heg1	HEG homolog 1 (zebrafish)
Herpud2	HERPUD family member 2
Hes1	hairy and enhancer of split 1 (Drosophila)
Hexb	hexosaminidase B
Hist1h1c	histone cluster 1, H1c

	<p>predicted gene 13121; predicted gene 3160; high-mobility group (nonhistone chromosomal) protein 1-like 1; predicted gene 6090; predicted gene 3851; predicted gene 8967; predicted gene 7782; predicted gene 4587; predicted gene 4689; predicted gene 3307; predicted gene 13932; predicted gene 15059; predicted gene 3565; predicted gene 15447; predicted gene 12587; predicted gene 9012; predicted gene 6115; predicted gene 9480; high mobility group box 1; predicted gene 8423; predicted gene 5853; predicted gene 8288; predicted gene 7888; predicted gene 8594; predicted gene 15387; predicted gene 5473; predicted gene 8807; similar to high mobility group box 1; similar to 2810416G20Rik protein; predicted gene 8390; predicted gene, OTTMUSG00000005439; predicted gene 5842; predicted gene 5527; predicted gene 8563; predicted gene 2710; predicted gene 12331; predicted gene 5937; predicted gene 5504; similar to high-mobility group box 1; predicted gene 10361; predicted gene 2607; predicted gene 7422; predicted gene 10075; predicted gene 12568; predicted gene 6589; predicted gene 4383; predicted gene 8031; similar to High mobility group protein 1 (HMG-1) (High mobility group protein B1) (Amphoterin) (Heparin-binding protein p30); predicted gene 7468; predicted gene 8554</p>
Hnrnph1	heterogeneous nuclear ribonucleoprotein H1
Hnrnph2	heterogeneous nuclear ribonucleoprotein H2
Hnrnpl	heterogeneous nuclear ribonucleoprotein L
Hnrnpm	heterogeneous nuclear ribonucleoprotein M
Hnrnpr	predicted gene 6159; heterogeneous nuclear ribonucleoprotein R
Hook3	hook homolog 3 (Drosophila)
Hoxa5	homeo box A5
Hp1bp3	heterochromatin protein 1, binding protein 3
Hsp90aa1	predicted gene 5511; heat shock protein 90, alpha (cytosolic), class A member 1
Hsp90ab1	heat shock protein 90 alpha (cytosolic), class B member 1
Hsp90b1	heat shock protein 90, beta (Grp94), member 1
Hspa12a	heat shock protein 12A
Hspa2	heat shock protein 2
Hspb1	heat shock protein 1
Hspb8	heat shock protein 8
Id1	inhibitor of DNA binding 1
Id2	inhibitor of DNA binding 2
Ier2	immediate early response 2
Ifi204	interferon activated gene 204
Ifi205	interferon activated gene 205
Ifi27l2a	interferon, alpha-inducible protein 27 like 2A

Ifi35	interferon-induced protein 35
Ifit3	interferon-induced protein with tetratricopeptide repeats 3
Ifitm3	interferon induced transmembrane protein 3
Ifnar2	interferon (alpha and beta) receptor 2
Ifngr1	interferon gamma receptor 1
Ifrd1	interferon-related developmental regulator 1
Ift74	intraflagellar transport 74 homolog (Chlamydomonas)
Igf1r	insulin-like growth factor I receptor
Igfbp5	insulin-like growth factor binding protein 5
Igfbp6	insulin-like growth factor binding protein 6
Il16	interleukin 16
Il17re	interleukin 17 receptor E
Il6ra	interleukin 6 receptor, alpha
Il6st	interleukin 6 signal transducer
Illdr2	immunoglobulin-like domain containing receptor 2
Ilf3	interleukin enhancer binding factor 3
Impad1	inositol monophosphatase domain containing 1
Ints10	integrator complex subunit 10; similar to integrator complex subunit 10
Iqsec1	IQ motif and Sec7 domain 1
Irak4	interleukin-1 receptor-associated kinase 4
Irf2bp2	interferon regulatory factor 2 binding protein 2
Irf7	interferon regulatory factor 7
Irs2	insulin receptor substrate 2
Itch	itchy, E3 ubiquitin protein ligase
Itga6	integrin alpha 6
Itpr2	inositol 1,4,5-triphosphate receptor 2
Jmjd1c	jumonji domain containing 1C
Jun	Jun oncogene
Junb	Jun-B oncogene
Jund	Jun proto-oncogene related gene d
Jup	junction plakoglobin
Kank1	KN motif and ankyrin repeat domains 1
Kcnab1	potassium voltage-gated channel, shaker-related subfamily, beta member 1
Kdelr1	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 1
Kdm5a	lysine (K)-specific demethylase 5A
Kdm6b	KDM1 lysine (K)-specific demethylase 6B
Kdr	kinase insert domain protein receptor
Keap1	kelch-like ECH-associated protein 1
Kif1b	kinesin family member 1B

Kif5b	kinesin family member 5B
Klf10	Kruppel-like factor 10
Klf2	Kruppel-like factor 2 (lung)
Klf4	Kruppel-like factor 4 (gut)
Klf6	Kruppel-like factor 6
Klf7	Kruppel-like factor 7 (ubiquitous)
Klf9	Kruppel-like factor 9
Kpna1	karyopherin (importin) alpha 1
Kpna3	karyopherin (importin) alpha 3
Krcc1	lysine-rich coiled-coil 1
Krt14	keratin 14
Ktn1	kinectin 1
Lama4	laminin, alpha 4
Lamp2	lysosomal-associated membrane protein 2
Lars2	leucyl-tRNA synthetase, mitochondrial
Lass2	LAG1 homolog, ceramide synthase 2
Lass4	LAG1 homolog, ceramide synthase 4
Lgals7	lectin, galactose binding, soluble 7
Limch1	LIM and calponin homology domains 1
Lims2	LIM and senescent cell antigen like domains 2
Lman1	lectin, mannose-binding, 1
Lpar2	lysophosphatidic acid receptor 2
Lrrc20	leucine rich repeat containing 20
Lrrc58	leucine rich repeat containing 58; predicted gene, OTTMUSG00000025724
Lrrc61	leucine rich repeat containing 61
Lrrn4	leucine rich repeat neuronal 4
Lrrn4cl	LRRN4 C-terminal like
Ltbp4	latent transforming growth factor beta binding protein 4
Luc7l3	RIKEN cDNA 3300001P08 gene
Maf	similar to c-Maf long form; avian musculoaponeurotic fibrosarcoma (v-maf) AS42 oncogene homolog
Maged1	melanoma antigen, family D, 1
Magt1	magnesium transporter 1
Malat1	metastasis associated lung adenocarcinoma transcript 1 (non-coding RNA)
Man1a	mannosidase 1, alpha
Manf	mesencephalic astrocyte-derived neurotrophic factor
Maoa	monoamine oxidase A
Map3k3	mitogen-activated protein kinase kinase kinase 3
Mapk1	mitogen-activated protein kinase 1
Mapkapk3	mitogen-activated protein kinase-activated protein kinase 3

Mapre2	microtubule-associated protein, RP/EB family, member 2
Marcks1	MARCKS-like 1; predicted gene 9106
Mat2a	methionine adenosyltransferase II, alpha
Mat2b	methionine adenosyltransferase II, beta
Matr3	matrin 3; similar to Matrin 3
Med13l	mediator complex subunit 13-like
Med21	mediator complex subunit 21
Mef2c	myocyte enhancer factor 2C
Meis2	Meis homeobox 2
Mesdc1	mesoderm development candidate 1
Metap2	methionine aminopeptidase 2
Mettl2	methyltransferase like 2
Mettl7a1	methyltransferase like 7A1
Mfap1a	similar to microfibrillar-associated protein 1A; microfibrillar-associated protein 1A; microfibrillar-associated protein 1B
Mfhas1	malignant fibrous histiocytoma amplified sequence 1
Mgll	monoglyceride lipase
Mgst1	microsomal glutathione S-transferase 1
Mll1	myeloid/lymphoid or mixed-lineage leukemia 1
Mll3	myeloid/lymphoid or mixed-lineage leukemia 3
Morf4l2	predicted gene 5521; similar to mortality factor 4 like 2; mortality factor 4 like 2
Mpdz	multiple PDZ domain protein
Mphosph8	M-phase phosphoprotein 8
Mras	muscle and microspikes RAS
Mrgprf	MAS-related GPR, member F
Msn	moesin
Mtap1a	microtubule-associated protein 1 A
Mtdh	metadherin
Mtmr6	myotubularin related protein 6
Mut	methylmalonyl-Coenzyme A mutase
Mxd4	Max dimerization protein 4
Myh10	myosin, heavy polypeptide 10, non-muscle
Myl7	myosin, light polypeptide 7, regulatory
Mylip	myosin regulatory light chain interacting protein
Myst4	MYST histone acetyltransferase monocytic leukemia 4
Naa25	RIKEN cDNA C330023M02 gene
Naga	N-acetyl galactosaminidase, alpha
Nckap1	NCK-associated protein 1
Ncoa1	similar to Nuclear receptor coactivator 1 (NCoA-1) (Steroid receptor coactivator 1) (SRC-1) (Nuclear receptor coactivator protein 1) (mNRC-1); nuclear receptor coactivator 1

Ncoa4	predicted gene 6768; nuclear receptor coactivator 4
Ncor1	nuclear receptor co-repressor 1
Ndn	necdin
Ndst1	N-deacetylase/N-sulfotransferase (heparan glucosaminyl) 1
Ndufa4	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 4
Nedd4	neural precursor cell expressed, developmentally down-regulated 4
Nf1	neurofibromatosis 1
Nfe2l1	nuclear factor, erythroid derived 2,-like 1
Nfia	nuclear factor I/A
Nfic	nuclear factor I/C
Nfix	nuclear factor I/X
Nfkb2	nuclear factor of kappa light polypeptide gene enhancer in B-cells 2, p49/p100
Nfkbia	nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha
Nfkbiz	nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta
Nfyc	nuclear transcription factor-Y gamma
Nid2	nidogen 2
Ninl	ninein-like
Nipal3	NIPA-like domain containing 3; similar to NIPA-like domain containing 3
Nipbl	Nipped-B homolog (Drosophila)
Nkain4	Na+/K+ transporting ATPase interacting 4
Nkd1	naked cuticle 1 homolog (Drosophila); similar to naked cuticle 1 homolog
Nnmt	nicotinamide N-methyltransferase
Nod1	nucleotide-binding oligomerization domain containing 1
Npr1	natriuretic peptide receptor 1
Nr1d1	nuclear receptor subfamily 1, group D, member 1
Nr3c1	nuclear receptor subfamily 3, group C, member 1
Nr4a1	nuclear receptor subfamily 4, group A, member 1
Nrgn	neurogranin
Nucks1	nuclear casein kinase and cyclin-dependent kinase substrate 1
Oasl2	2'-5' oligoadenylate synthetase-like 2
Oat	ornithine aminotransferase
Ogdh	oxoglutarate dehydrogenase (lipoamide)
Ogn	osteoglycin
Olfr1033	olfactory receptor 1033
Olfr613	olfactory receptor 614; hypothetical protein LOC100044261; olfactory receptor 613
Opa3	optic atrophy 3 (human)

Orai3	ORAI calcium release-activated calcium modulator 3
Osr1	odd-skipped related 1 (Drosophila)
Oxct1	3-oxoacid CoA transferase 1
Oxnad1	oxidoreductase NAD-binding domain containing 1
Pard3b	par-3 partitioning defective 3 homolog B (C. elegans)
Parp14	poly (ADP-ribose) polymerase family, member 14
Parp4	poly (ADP-ribose) polymerase family, member 4
Parvb	parvin, beta; similar to parvin, beta
Pbx1	pre B-cell leukemia transcription factor 1; region containing RIKEN cDNA 2310056B04 gene; pre B-cell leukemia transcription factor 1
Pcdh15	protocadherin 15
Pcdhgb5	protocadherin gamma subfamily B, 5
Pcm1	pericentriolar material 1
Pdap1	PDGFA associated protein 1
Pcd6ip	programmed cell death 6 interacting protein
Pde4dip	phosphodiesterase 4D interacting protein (myomegalin)
Pdia3	protein disulfide isomerase associated 3
Pdia4	protein disulfide isomerase associated 4
Pdpn	podoplanin
Pef1	penta-EF hand domain containing 1
Peli1	pellino 1
Per1	period homolog 1 (Drosophila)
Pf4	platelet factor 4
Pfn1	profilin 1
Pgcp	plasma glutamate carboxypeptidase
Pgrmc1	progesterone receptor membrane component 1
Phf21a	PHD finger protein 21A
Phf3	PHD finger protein 3
Phip	pleckstrin homology domain interacting protein
Pigt	phosphatidylinositol glycan anchor biosynthesis, class T; similar to GPI transamidase component PIG-T precursor (Phosphatidylinositol-glycan biosynthesis class T protein) (Neuronal development-associated protein 7)
Pik3c2a	phosphatidylinositol 3-kinase, C2 domain containing, alpha polypeptide
Pim1	proviral integration site 1
Pitpnm2	phosphatidylinositol transfer protein, membrane-associated 2
Pkhd1l1	polycystic kidney and hepatic disease 1-like 1
Pknox1	Pbx/knotted 1 homeobox
Pla2g4a	phospholipase A2, group IVA (cytosolic, calcium-dependent)
Plat	plasminogen activator, tissue
Plce1	phospholipase C, epsilon 1

Plk1s1	non-protein coding RNA 153
Plk2	polo-like kinase 2 (Drosophila)
Plod2	procollagen lysine, 2-oxoglutarate 5-dioxygenase 2
Plxdc1	plexin domain containing 1
Plxdc2	plexin domain containing 2
Plxna4	plexin A4
Pmp22	peripheral myelin protein 22
Pnrc1	proline-rich nuclear receptor coactivator 1
Podn	podocan
Ppap2a	phosphatidic acid phosphatase type 2A
Ppbp	pro-platelet basic protein
Ppfibp2	protein tyrosine phosphatase, receptor-type, F interacting protein, binding protein 2
Ppig	peptidyl-prolyl isomerase G (cyclophilin G)
Ppl	periplakin
Ppp1cb	protein phosphatase 1, catalytic subunit, beta isoform
Ppp1r12a	protein phosphatase 1, regulatory (inhibitor) subunit 12A
Ppp1r15a	protein phosphatase 1, regulatory (inhibitor) subunit 15A; myeloid differentiation primary response gene 116
Ppp3ca	protein phosphatase 3, catalytic subunit, alpha isoform
Pppde1	PPPDE peptidase domain containing 1
Pqlc3	PQ loop repeat containing
Prelp	proline arginine-rich end leucine-rich repeat
Prg4	proteoglycan 4 (megakaryocyte stimulating factor, articular superficial zone protein)
Prkar2a	protein kinase, cAMP dependent regulatory, type II alpha
Prpf40a	PRP40 pre-mRNA processing factor 40 homolog A (yeast)
Prr13	proline rich 13
Prss23	protease, serine, 23
Psd	pleckstrin and Sec7 domain containing
Psip1	PC4 and SFRS1 interacting protein 1
Psmb2	proteasome (prosome, macropain) subunit, beta type 2
Psmd11	predicted gene 14048; proteasome (prosome, macropain) 26S subunit, non-ATPase, 11
Psmd7	proteasome (prosome, macropain) 26S subunit, non-ATPase, 7
Ptges3	predicted gene 9769; prostaglandin E synthase 3 (cytosolic); similar to Sid3177p; predicted gene 11893
Ptgis	prostaglandin I2 (prostacyclin) synthase
Ptgs1	prostaglandin-endoperoxide synthase 1
Ptma	predicted gene 12504; predicted gene 9800; predicted gene 4617; predicted gene 6625; predicted gene 7614; similar to prothymosin alpha; prothymosin alpha; predicted gene 9009

Ptp4a2	predicted gene 13422; protein tyrosine phosphatase 4a2
Ptplad2	protein tyrosine phosphatase-like A domain containing 2
Ptprd	protein tyrosine phosphatase, receptor type, D
Ptprf	protein tyrosine phosphatase, receptor type, F
Ptrf	polymerase I and transcript release factor
Qrich1	glutamine-rich 1
Qser1	glutamine and serine rich 1
R74862	expressed sequence R74862
Rab11fip1	RAB11 family interacting protein 1 (class I)
Rab1b	RAB1B, member RAS oncogene family
Rab5c	RAB5C, member RAS oncogene family
Rab6b	RAB6B, member RAS oncogene family
Rab7	RAB7, member RAS oncogene family
Rabgap1l	RAB GTPase activating protein 1-like
Ralbp1	ralA binding protein 1
Raly	RIKEN cDNA C130057N11 gene; hnRNP-associated with lethal yellow
Rarres2	retinoic acid receptor responder (tazarotene induced) 2
Rb1cc1	RB1-inducible coiled-coil 1
Rbbp6	retinoblastoma binding protein 6
Rbbp8	retinoblastoma binding protein 8
Rbm25	RNA binding motif protein 25
Rbm27	RNA binding motif protein 27
Rbm3	predicted gene 15453; RNA binding motif protein 3
Rbpms	RNA binding protein gene with multiple splicing
Rdx	radixin
Rest	RE1-silencing transcription factor
Rgma	RGM domain family, member A
Rgs10	regulator of G-protein signalling 10
Rhob	ras homolog gene family, member B
Rhoj	ras homolog gene family, member J
Rhou	ras homolog gene family, member U
Rnase4	ribonuclease, RNase A family 4
Rnd3	Rho family GTPase 3
Rnf167	ring finger protein 167
Rnf20	ring finger protein 20
Rock1	Rho-associated coiled-coil containing protein kinase 1
Rock2	Rho-associated coiled-coil containing protein kinase 2
Rpp25	ribonuclease P 25 subunit (human)
Rras2	related RAS viral (r-ras) oncogene homolog 2
Rspo1	R-spondin homolog (Xenopus laevis)

Rtf1	Rtf1, Paf1/RNA polymerase II complex component, homolog ( <i>S. cerevisiae</i> )
Rtn1	reticulon 1
Ryk	receptor-like tyrosine kinase
Sarnp	predicted gene 6563; SAP domain containing ribonucleoprotein
Sat1	similar to spermidine/spermine N1-acetyltransferase; predicted gene 5552; spermidine/spermine N1-acetyl transferase 1
Sbsn	suprabasin
Scd1	stearoyl-Coenzyme A desaturase 1
Sdc4	syndecan 4
Sdpr	serum deprivation response
Sec62	SEC62 homolog ( <i>S. cerevisiae</i> )
Secisbp2	SECIS binding protein 2
Sema5a	sema domain, seven thrombospondin repeats (type 1 and type 1-like), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 5A
Senp6	similar to Sentrin-specific protease 6 (Sentrin/SUMO-specific protease SENP6) (SUMO-1-specific protease 1); SUMO/sentrin specific peptidase 6
Sep15	selenoprotein
Sept9	septin 9
Serinc5	serine incorporator 5
Serpincb6b	serine (or cysteine) peptidase inhibitor, clade B, member 6b
Serpinc1	serine (or cysteine) peptidase inhibitor, clade G, member 1
Serpinc1h	serine (or cysteine) peptidase inhibitor, clade H, member 1
Sesn1	sestrin 1
Setd2	SET domain containing 2
Sf3b1	splicing factor 3b, subunit 1
Sf3b4	predicted gene 7935; splicing factor 3b, subunit 4
Sfrs18	splicing factor, arginine/serine-rich 18
Shc1	predicted gene 5500; src homology 2 domain-containing transforming protein C1
Shfm1	split hand/foot malformation (ectrodactyly) type 1
Siae	sialic acid acetyl esterase
Siah1a	seven in absentia 1A
Sirt2	sirtuin 2 (silent mating type information regulation 2, homolog) 2 ( <i>S. cerevisiae</i> )
Slc10a3	solute carrier family 10 (sodium/bile acid cotransporter family), member 3
Slc16a1	solute carrier family 16 (monocarboxylic acid transporters), member 1
Slc1a5	solute carrier family 1 (neutral amino acid transporter), member 5

Slc26a3	solute carrier family 26, member 3
Slc27a3	solute carrier family 27 (fatty acid transporter), member 3
Slc38a1	solute carrier family 38, member 1
Slc39a8	solute carrier family 39 (metal ion transporter), member 8
Slc43a3	solute carrier family 43, member 3
Slc4a4	solute carrier family 4 (anion exchanger), member 4
Slc6a4	solute carrier family 6 (neurotransmitter transporter, serotonin), member 4
Slc6a6	solute carrier family 6 (neurotransmitter transporter, taurine), member 6
Slc8a1	solute carrier family 8 (sodium/calcium exchanger), member 1
Slc9a3r1	solute carrier family 9 (sodium/hydrogen exchanger), member 3 regulator 1
Slpi	secretory leukocyte peptidase inhibitor
Sltm	SAFB-like, transcription modulator
Slu7	SLU7 splicing factor homolog ( <i>S. cerevisiae</i> )
Slurp1	secreted Ly6/Plaur domain containing 1
Smad4	similar to MAD homolog 4 ( <i>Drosophila</i> ); MAD homolog 4 ( <i>Drosophila</i> )
Smarca2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2
Smarca5	predicted gene 13034; SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 5
Smc2	structural maintenance of chromosomes 2
Smc3	predicted gene 8892; structural maintenance of chromosomes 3
Smc4	structural maintenance of chromosomes 4
Smc6	structural maintenance of chromosomes 6
Smchd1	SMC hinge domain containing 1
Smpd3	sphingomyelin phosphodiesterase 3, neutral
Snrnp70	small nuclear ribonucleoprotein 70 (U1)
Sntb2	similar to beta-2-syntrophin; syntrophin, basic 2
Soat1	sterol O-acyltransferase 1
Socs3	suppressor of cytokine signaling 3
Sod3	superoxide dismutase 3, extracellular
Sorbs1	sorbin and SH3 domain containing 1
Sorbs3	sorbin and SH3 domain containing 3
Sox6	SRY-box containing gene 6
Sp100	nuclear antigen Sp100
Spag9	sperm associated antigen 9
Sparc	secreted acidic cysteine rich glycoprotein; similar to Secreted acidic cysteine rich glycoprotein
Spen	SPEN homolog, transcriptional regulator ( <i>Drosophila</i> )

Spint2	serine protease inhibitor, Kunitz type 2
Spnb2	spectrin beta 2
Spock2	sparc/osteonectin, cwcv and kazal-like domains proteoglycan 2
Spon2	spondin 2, extracellular matrix protein
Spop	speckle-type POZ protein
Src	Rous sarcoma oncogene
Srrm1	serine/arginine repetitive matrix 1
Ssh2	slingshot homolog 2 (Drosophila)
Ssr3	signal sequence receptor, gamma
St3gal1	ST3 beta-galactoside alpha-2,3-sialyltransferase 1
Stag1	stromal antigen 1
Star	steroidogenic acute regulatory protein
Stard5	StAR-related lipid transfer (START) domain containing 5
Stat3	similar to Stat3B; signal transducer and activator of transcription 3
Stim1	similar to Stromal interaction molecule 1; stromal interaction molecule 1
Stk10	serine/threonine kinase 10
Stk40	serine/threonine kinase 40
Stmn2	stathmin-like 2
Stra6	stimulated by retinoic acid gene 6
Strn3	striatin, calmodulin binding protein 3
Sulf1	sulfatase 1
Sulf2	sulfatase 2
Supt16h	suppressor of Ty 16 homolog ( <i>S. cerevisiae</i> )
Sv2a	synaptic vesicle glycoprotein 2 a
Syne1	synaptic nuclear envelope 1
Syne2	synaptic nuclear envelope 2
Syt11	synaptotagmin XI; similar to synaptotagmin XI
Syt1l1	synaptotagmin-like 1; similar to synaptotagmin-like 1
Taf3	TAF3 RNA polymerase II, TATA box binding protein (TBP)-associated factor
Taf7	TAF7 RNA polymerase II, TATA box binding protein (TBP)-associated factor
Tapbp	TAP binding protein
Tbc1d15	TBC1 domain family, member 15
Tbcel	tubulin folding cofactor E-like
Tbl1x	transducin (beta)-like 1 X-linked
Tbx18	T-box18
Tceal8	transcription elongation factor A (SII)-like 8; similar to transcription elongation factor A (SII)-like 8
Tcf7l1	transcription factor 3
Tfdp2	transcription factor Dp 2

Tgfb1i1	transforming growth factor beta 1 induced transcript 1
Tgfb2	transforming growth factor, beta 2
Tgfbr2	transforming growth factor, beta receptor II
Tgm2	transglutaminase 2, C polypeptide
Thbd	thrombomodulin
Thbs1	thrombospondin 1; similar to thrombospondin 1
Thoc2	THO complex 2; cDNA sequence BC005561
Thrap3	thyroid hormone receptor associated protein 3; predicted gene 5898
Thsd4	thrombospondin, type I, domain containing 4
Timp2	tissue inhibitor of metalloproteinase 2
Tirap	toll-interleukin 1 receptor (TIR) domain-containing adaptor protein
Tlr2	toll-like receptor 2
Tm4sf1	transmembrane 4 superfamily member 1
Tm4sf5	transmembrane 4 superfamily member 5
Tmcc3	transmembrane and coiled coil domains 3
Tmco1	transmembrane and coiled-coil domains 1
Tmco7	transmembrane and coiled-coil domains 7
Tmed2	transmembrane emp24 domain trafficking protein 2; predicted gene 10698; predicted gene 7318
Tmem119	transmembrane protein 119
Tmem140	transmembrane protein 140
Tmem151a	transmembrane protein 151A
Tmem221	transmembrane protein 221
Tmem50a	transmembrane protein 50A
Tmem98	transmembrane protein 98
Tmod3	tropomodulin 3
Tmpo	thymopoietin
Tmsb4x	thymosin, beta 4, X chromosome; similar to thymosin beta-4
Tnxb	tenascin XB
Tob2	transducer of ERBB2, 2
Topors	topoisomerase I binding, arginine/serine-rich
Tpm3	predicted gene 7848; predicted gene 7839; predicted gene 4157; similar to tropomyosin 3, gamma; tropomyosin 3, gamma; predicted gene 4903
Tppp3	tubulin polymerization-promoting protein family member 3
Tpt1	predicted gene 1974; tumor protein, translationally-controlled 1 pseudogene; tumor protein, translationally-controlled 1; predicted gene 14456
Trafd1	TRAF type zinc finger domain containing 1
Trib1	tribbles homolog 1 (Drosophila)
Trim8	tripartite motif protein 8
Trpm7	transient receptor potential cation channel, subfamily M, member 7

Tsc22d3	TSC22 domain family, member 3
Tshz1	teashirt zinc finger family member 1
Tsix	X (inactive)-specific transcript, antisense
Tspan31	tetraspanin 31
Tspan5	tetraspanin 5
Ttc28	tetratricopeptide repeat domain 28
Ttc38	tetratricopeptide repeat domain 38
Tuba1a	predicted gene 7172; similar to tubulin, alpha 1; tubulin, alpha 1A
Tubb2a	tubulin, beta 2A
Twsg1	twisted gastrulation homolog 1 (Drosophila)
Txndc5	thioredoxin domain containing 5
Txnrd1	thioredoxin reductase 1
Uap1	UDP-N-acetylglucosamine pyrophosphorylase 1
Uba7	ubiquitin-activating enzyme E1-like; RIKEN cDNA D330022A01 gene
Ube2d1	ubiquitin-conjugating enzyme E2D 1, UBC4/5 homolog (yeast)
Ube2l6	ubiquitin-conjugating enzyme E2L 6
Ube2n	ubiquitin-conjugating enzyme E2N; similar to ubiquitin-conjugating enzyme E2 UbCH-ben; similar to ubiquitin-conjugating enzyme E2N; predicted gene 5943
Ube2v1	ubiquitin-conjugating enzyme E2 variant 1; predicted gene 7181; predicted gene 12502; similar to ubiquitin-conjugating enzyme E2 variant 1
Ubqln2	ubiquilin 2
Ubxn2a	UBX domain protein 2A; predicted gene 6245
Ubxn4	UBX domain protein 4
Ugdh	UDP-glucose dehydrogenase
Upk1b	uroplakin 1B
Upk3b	uroplakin 3B
Usp16	ubiquitin specific peptidase 16
Usp2	ubiquitin specific peptidase 2
Usp25	ubiquitin specific peptidase 25
Usp54	ubiquitin specific peptidase 54
Usp8	ubiquitin specific peptidase 8
Utp20	UTP20, small subunit (SSU) processome component, homolog (yeast)
Vat1	vesicle amine transport protein 1 homolog (T californica)
Vim	vimentin
Vps13a	vacuolar protein sorting 13A (yeast)
Vwa5a	von Willebrand factor A domain containing 5A
Wac	similar to WW domain-containing adapter protein with coiled-coil; WW domain containing adaptor with coiled-coil
Wasf2	WAS protein family, member 2

Wdr26	WD repeat domain 26; similar to myocardial ischemic preconditioning upregulated protein 2
Wdr92	WD repeat domain 92
Wfdc1	WAP four-disulfide core domain 1
Wls	G protein-coupled receptor 177
Wnt4	wingless-related MMTV integration site 4
Wrnip1	Werner helicase interacting protein 1
Wt1	similar to Wilms tumor homolog; Wilms tumor 1 homolog
Wwc2	WW, C2 and coiled-coil domain containing 2
Xdh	xanthine dehydrogenase
Xist	inactive X specific transcripts
Yipf5	Yip1 domain family, member 5; predicted gene 5738
Ywhaz	tyrosine 3-monoxygenase/tryptophan 5-monoxygenase activation protein, zeta polypeptide; predicted gene 4202
Zbed6	similar to Zinc finger BED domain containing protein 4
Zbtb16	zinc finger and BTB domain containing 16
Zbtb20	zinc finger and BTB domain containing 20
Zbtb4	zinc finger and BTB domain containing 4
Zbtb7c	zinc finger and BTB domain containing 7C
Zc3h13	zinc finger CCCH type containing 13
Zc3h18	predicted gene 5939; zinc finger CCCH-type containing 18
Zcchc11	zinc finger, CCHC domain containing 11
Zcchc3	zinc finger, CCHC domain containing 3
Zfand6	zinc finger, AN1-type domain 6
Zfhx4	zinc finger homeodomain 4
Zfp148	zinc finger protein 148
Zfp277	zinc finger protein 277
Zfp281	zinc finger protein 281
Zfp318	zinc finger protein 318
Zfp353	zinc finger protein 353
Zfp36	zinc finger protein 36
Zfp385a	zinc finger protein 385A
Zfp488	zinc finger protein 488
Zfp672	zinc finger protein 672
Zfp704	zinc finger protein 704
Zmat1	zinc finger, matrin type 1
Zrsr1	zinc finger (CCCH type), RNA binding motif and serine/arginine rich 1
Zzef1	zinc finger, ZZ-type with EF hand domain 1
1110002B05Rik	RIKEN cDNA 1110002B05 gene
1110003E01Rik	RIKEN cDNA 1110003E01 gene

1110004F10Rik	predicted gene 9169; RIKEN cDNA 1110004F10 gene; similar to small acidic protein
1500003O03Rik	RIKEN cDNA 1500003O03 gene; similar to EF-hand Ca2+ binding protein p22
1600029D21Rik	RIKEN cDNA 1600029D21 gene
1810014B01Rik	RIKEN cDNA 1810014B01 gene
1810041L15Rik	RIKEN cDNA 1810041L15 gene
1810074P20Rik	RIKEN cDNA 1810074P20 gene
2010107G12Rik	RIKEN cDNA 2010107G12 gene
2210403K04Rik	hypothetical protein LOC100042498
2310030G06Rik	RIKEN cDNA 2310030G06 gene
2510002D24Rik	RIKEN cDNA 2510002D24 gene
2610034B18Rik	RIKEN cDNA 2610034B18 gene
2610101N10Rik	RIKEN cDNA 2610101N10 gene
2810474O19Rik	RIKEN cDNA 2810474O19 gene
2900002K06Rik	RIKEN cDNA 2900002K06 gene
3110062M04Rik	RIKEN cDNA 3110062M04 gene
4930402H24Rik	RIKEN cDNA 4930402H24 gene
4930523C07Rik	RIKEN cDNA 4930523C07 gene
5430435G22Rik	RIKEN cDNA 5430435G22 gene
6330406I15Rik	RIKEN cDNA 6330406I15 gene
A130040M12Rik	RIKEN cDNA A130040M12 gene
AI848100	expressed sequence AI848100
Gm16897	
kg:uc009lxf.1	
Prrc2c	
kg:uc007won.1	
kg:uc009ogv.1	
kg:uc009iln.1	
kg:uc007qca.1	
Atxn7l3b	
kg:uc008ewj.2	
kg:uc008wkn.1	
kg:uc007bgn.1	
Ces2g	
kg:uc009cvm.1	
kg:uc008ehr.1	
Tmem234	
kg:uc012hdk.1	
kg:uc008ajk.1	
eg:245190:chr7:m	
kg:uc007qse.1	

kg:uc007bvx.1	
Mob3c	
kg:uc008dzh.1	
kg:uc009okn.1	
kg:uc007zts.1	
kg:uc008jup.1	
kg:uc008tkz.1	
kg:uc007zwh.1	
kg:uc008znh.1	
Mau2	
kg:uc009mng.1	
kg:uc007ded.1	
kg:uc007ctp.1	
kg:uc007zak.1	
eg:497210:chr14:m	
kg:uc007vsr.1	
Mir3064	
kg:uc009ize.1	
Kansl1	
eg:320169:chr9:p	
kg:uc009vev.1	
kg:uc009acs.1	
kg:uc009tuw.1	
kg:uc007pff.1	
kg:uc007vnc.1	
kg:uc009igb.1	
kg:uc008oki.1	
kg:uc008tky.1	

[0037] **Table 8: Exemplary human marker genes**

HUMAN GENE SYMBOL	Gene Name
ABI3BP	ABI family, member 3 (NESH) binding protein
ABLIM3	actin binding LIM protein family, member 3
ACAD9	acyl-Coenzyme A dehydrogenase family, member 9
ACBD3	acyl-Coenzyme A binding domain containing 3
ACIN1	apoptotic chromatin condensation inducer 1
ACTB	actin, beta
ACTG1	actin, gamma 1
ADAMTS5	ADAM metallopeptidase with thrombospondin type 1 motif, 5

ADAMTSL1	ADAMTS-like 1
ADD3	adducin 3 (gamma)
AEBP1	AE binding protein 1
AGAP1	ArfGAP with GTPase domain, ankyrin repeat and PH domain 1
AKAP13	A kinase (PRKA) anchor protein 13
AKAP2	A kinase (PRKA) anchor protein 2; paralemmin 2; PALM2-AKAP2 readthrough transcript
AKT2	v-akt murine thymoma viral oncogene homolog 2
ALDH1A1	aldehyde dehydrogenase 1 family, member A1
ALDH1A2	aldehyde dehydrogenase 1 family, member A2
ALOX12	arachidonate 12-lipoxygenase
AMFR	autocrine motility factor receptor
AMHR2	anti-Mullerian hormone receptor, type II
ANG	angiogenin, ribonuclease, RNase A family, 5
ANKRD11	ankyrin repeat domain 11; hypothetical protein LOC100128265
ANKRD12	ankyrin repeat domain 12
ANKRD17	ankyrin repeat domain 17
ANO6	anoctamin 6
ANP32A	hepatopoietin PCn127; acidic (leucine-rich) nuclear phosphoprotein 32 family, member A
ANXA7	annexin A7
AP1S3	adaptor-related protein complex 1, sigma 3 subunit
AP3S1	adaptor-related protein complex 3, sigma 1 subunit
AP4E1	adaptor-related protein complex 4, epsilon 1 subunit
APLP1	amyloid beta (A4) precursor-like protein 1
APP	amyloid beta (A4) precursor protein
AQP1	aquaporin 1 (Colton blood group)
ARAP2	ArfGAP with RhoGAP domain, ankyrin repeat and PH domain 2
ARF3	ADP-ribosylation factor 3
ARF5	ADP-ribosylation factor 5
ARHGAP28	Rho GTPase activating protein 28
ARHGAP29	Rho GTPase activating protein 29
ARHGAP5	Rho GTPase activating protein 5
ARHGEF12	Rho guanine nucleotide exchange factor (GEF) 12
ARID1A	AT rich interactive domain 1A (SWI-like)
ARID4A	AT rich interactive domain 4A (RBP1-like)
ARID4B	AT rich interactive domain 4B (RBP1-like)
ARID5B	AT rich interactive domain 5B (MRF1-like)
ARL3	ADP-ribosylation factor-like 3
ARL4D	ADP-ribosylation factor-like 4D
ARL6IP5	ADP-ribosylation-like factor 6 interacting protein 5
ARMCX3	armadillo repeat containing, X-linked 3

ARPC2	actin related protein 2/3 complex, subunit 2, 34kDa
ARSA	arylsulfatase A
ARSB	arylsulfatase B
ASCC3	activating signal cointegrator 1 complex subunit 3
ATF3	activating transcription factor 3
ATG3	ATG3 autophagy related 3 homolog (S. cerevisiae)
ATP1A1	ATPase, Na+/K+ transporting, alpha 1 polypeptide
ATP1B1	ATPase, Na+/K+ transporting, beta 1 polypeptide
ATP2B1	ATPase, Ca++ transporting, plasma membrane 1
ATP6V1A	ATPase, H+ transporting, lysosomal 70kDa, V1 subunit A
ATXN2	ataxin 2
B2M	beta-2-microglobulin
BAZ2A	bromodomain adjacent to zinc finger domain, 2A
BBS4	Bardet-Biedl syndrome 4
BBX	bobby sox homolog (Drosophila)
BCAM	basal cell adhesion molecule (Lutheran blood group)
BCL10	B-cell CLL/lymphoma 10; hypothetical LOC646626
BDP1	B double prime 1, subunit of RNA polymerase III transcription initiation factor IIIB
BICC1	bicaudal C homolog 1 (Drosophila)
BICD1	bicaudal D homolog 1 (Drosophila)
BIRC6	baculoviral IAP repeat-containing 6
BLVRB	biliverdin reductase B (flavin reductase (NADPH))
BNC1	basonuclin 1
BNC2	basonuclin 2
BOD1L	biorientation of chromosomes in cell division 1-like
BPTF	bromodomain PHD finger transcription factor
BRAF	v-raf murine sarcoma viral oncogene homolog B1
BRD2	bromodomain containing 2
BRD4	bromodomain containing 4
BRP44L	brain protein 44-like
BST2	NPC-A-7; bone marrow stromal cell antigen 2
BTBD2	BTB (POZ) domain containing 2
BTBD7	BTB (POZ) domain containing 7
BTF3	basic transcription factor 3; basic transcription factor 3, like 1 pseudogene
BTG2	BTG family, member 2
BZW1	basic leucine zipper and W2 domains 1 pseudogene 1; basic leucine zipper and W2 domains 1 like 1; basic leucine zipper and W2 domains 1
C1D	C1D nuclear receptor co-repressor; similar to nuclear DNA-binding protein; similar to hCG1791993

C1RL	complement component 1, r subcomponent-like
C1S	complement component 1, s subcomponent
C2	complement component 2
C3	similar to Complement C3 precursor; complement component 3; hypothetical protein LOC100133511
C4A	complement component 4A (Rodgers blood group)
C4B	complement component 4B (Chido blood group)
CALM1	calmodulin 3 (phosphorylase kinase, delta); calmodulin 2 (phosphorylase kinase, delta); calmodulin 1 (phosphorylase kinase, delta)
CALM2	calmodulin 3 (phosphorylase kinase, delta); calmodulin 2 (phosphorylase kinase, delta); calmodulin 1 (phosphorylase kinase, delta)
CAP1	CAP, adenylate cyclase-associated protein 1 (yeast)
CAST	calpastatin
CAV1	caveolin 1, caveolae protein, 22kDa
CCDC109B	coiled-coil domain containing 109B
CCDC34	coiled-coil domain containing 34
CCDC80	coiled-coil domain containing 80
CCDC88A	coiled-coil domain containing 88A
CCDC90A	coiled-coil domain containing 90A
CCNL1	cyclin L1
CD109	CD109 molecule
CD200	CD200 molecule
CD248	CD248 molecule, endosialin
CD34	CD34 molecule
CD55	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
CD81	CD81 molecule
CD82	CD82 molecule
CD9	CD9 molecule
CDC42EP3	CDC42 effector protein (Rho GTPase binding) 3
CDH11	cadherin 11, type 2, OB-cadherin (osteoblast)
CDH3	cadherin 3, type 1, P-cadherin (placental)
CDK13	cell division cycle 2-like 5 (cholinesterase-related cell division controller)
CDON	Cdon homolog (mouse)
CELF2	CUG triplet repeat, RNA binding protein 2
CEP164	centrosomal protein 164kDa
CEP57	centrosomal protein 57kDa
CFH	complement factor H
CFL1	cofilin 1 (non-muscle)

CFL2	cofilin 2 (muscle)
CHD1	chromodomain helicase DNA binding protein 1
CHD2	chromodomain helicase DNA binding protein 2
CHI3L1	chitinase 3-like 1 (cartilage glycoprotein-39)
CHST4	carbohydrate (N-acetylglucosamine 6-O) sulfotransferase 4
CISH	cytokine inducible SH2-containing protein
CLCN3	chloride channel 3
CLDN10	claudin 10
CLDN15	claudin 15
CLDN25	claudin-like
CLEC1B	C-type lectin domain family 1, member B
CLEC3B	C-type lectin domain family 3, member B
CLIC4	chloride intracellular channel 4
CLIP1	CAP-GLY domain containing linker protein 1
CLIP3	CAP-GLY domain containing linker protein 3
CLN8	ceroid-lipofuscinosis, neuronal 8 (epilepsy, progressive with mental retardation)
CMAH	cytidine monophosphate-N-acetylneurameric acid hydroxylase (CMP-N-acetylneuraminate monooxygenase) pseudogene
CMTM3	CKLF-like MARVEL transmembrane domain containing 3
CMTM7	CKLF-like MARVEL transmembrane domain containing 7
CNOT6L	CCR4-NOT transcription complex, subunit 6-like
COBL	cordon-bleu homolog (mouse)
COBL1	COBL-like 1
COL14A1	collagen, type XIV, alpha 1
COL1A2	collagen, type I, alpha 2
COL3A1	collagen, type III, alpha 1
COL4A6	collagen, type IV, alpha 6
COLEC12	collectin sub-family member 12
COQ10B	coenzyme Q10 homolog B ( <i>S. cerevisiae</i> )
CREB3L1	cAMP responsive element binding protein 3-like 1
CREB5	cAMP responsive element binding protein 5
CREBBP	CREB binding protein
CREG1	cellular repressor of E1A-stimulated genes 1
CRIM1	cysteine rich transmembrane BMP regulator 1 (chordin-like)
CRLS1	cardiolipin synthase 1
CRYAB	crystallin, alpha B
CRYL1	crystallin, lambda 1
CRYM	crystallin, mu
CSDA	cold shock domain protein A; cold shock domain protein A pseudogene 1
CSF1	colony stimulating factor 1 (macrophage)

CSNK1A1	casein kinase 1, alpha 1
CSRNP1	cysteine-serine-rich nuclear protein 1
CSRP1	cysteine and glycine-rich protein 1
CUEDC1	CUE domain containing 1
CYBRD1	cytochrome b reductase 1
CYP2S1	cytochrome P450, family 2, subfamily S, polypeptide 1
CYR61	cysteine-rich, angiogenic inducer, 61
DAB2	disabled homolog 2, mitogen-responsive phosphoprotein ( <i>Drosophila</i> )
DAG1	dystroglycan 1 (dystrophin-associated glycoprotein 1)
DAGLB	diacylglycerol lipase, beta
DAPK1	death-associated protein kinase 1
DCN	decorin
DDR1	discoidin domain receptor tyrosine kinase 1
DDR2	discoidin domain receptor tyrosine kinase 2
DDX3X	DEAD (Asp-Glu-Ala-Asp) box polypeptide 3, X-linked
DDX5	DEAD (Asp-Glu-Ala-Asp) box polypeptide 5
DENND5A	DENN/MADD domain containing 5A
DHX15	DEAH (Asp-Glu-Ala-His) box polypeptide 15
DLGAP4	discs, large ( <i>Drosophila</i> ) homolog-associated protein 4
DMKN	dermokine
DNAJA2	DnaJ (Hsp40) homolog, subfamily A, member 2
DNAJB9	DnaJ (Hsp40) homolog, subfamily B, member 9
DNAJC1	DnaJ (Hsp40) homolog, subfamily C, member 1
DNMT1	DNA (cytosine-5-)-methyltransferase 1
DPP4	dipeptidyl-peptidase 4
DPYSL2	dihydropyrimidinase-like 2
DPYSL3	dihydropyrimidinase-like 3
DST	dystonin
DTX2	deltex homolog 2 ( <i>Drosophila</i> )
DUSP1	dual specificity phosphatase 1
DUSP14	dual specificity phosphatase 14
DUSP3	dual specificity phosphatase 3
DYNC1I2	similar to dynein cytoplasmic 1 intermediate chain 2; dynein, cytoplasmic 1, intermediate chain 2
ECD	ecdysoneless homolog ( <i>Drosophila</i> )
EEA1	early endosome antigen 1
EEF1A1	eukaryotic translation elongation factor 1 alpha-like 7; eukaryotic translation elongation factor 1 alpha-like 3; similar to eukaryotic translation elongation factor 1 alpha 1; eukaryotic translation elongation factor 1 alpha 1
EFEMP1	EGF-containing fibulin-like extracellular matrix protein 1

EFHD2	EF-hand domain family, member D2
EFNA5	ephrin-A5
EGR1	early growth response 1
EHD2	EH-domain containing 2
EIF3A	eukaryotic translation initiation factor 3, subunit A
ELF1	E74-like factor 1 (ets domain transcription factor)
ELOVL6	ELOVL family member 6, elongation of long chain fatty acids (FEN1/Elo2, SUR4/Elo3-like, yeast)
EMP2	epithelial membrane protein 2
ENPP2	ectonucleotide pyrophosphatase/phosphodiesterase 2
ENPP4	ectonucleotide pyrophosphatase/phosphodiesterase 4 (putative function)
ESAM	endothelial cell adhesion molecule
ESF1	similar to ABT1-associated protein; ESF1, nucleolar pre-rRNA processing protein, homolog (S. cerevisiae)
ESPN	espin
ESYT3	family with sequence similarity 62 (C2 domain containing), member C
ETFA	electron-transfer-flavoprotein, alpha polypeptide
EVPL	envoplakin
EXOC4	exocyst complex component 4
F11R	F11 receptor
FAIM2	Fas apoptotic inhibitory molecule 2
FAM117A	family with sequence similarity 117, member A
FAM134B	family with sequence similarity 134, member B
FAM53B	family with sequence similarity 53, member B
FAM63B	family with sequence similarity 63, member B
FAM76A	family with sequence similarity 76, member A
FAM84B	family with sequence similarity 84, member B
FAS	Fas (TNF receptor superfamily, member 6)
FBLN1	fibulin 1
FERMT2	fermitin family homolog 2 (Drosophila)
FGF1	fibroblast growth factor 1 (acidic)
FHL1	four and a half LIM domains 1
FILIP1L	filamin A interacting protein 1-like
FKBP5	FK506 binding protein 5
FLII	flightless I homolog (Drosophila)
FLNC	filamin C, gamma (actin binding protein 280)
FLRT2	fibronectin leucine rich transmembrane protein 2
FMO2	flavin containing monooxygenase 2 (non-functional)
FMOD	fibromodulin
FNDC1	fibronectin type III domain containing 1

FOS	v-fos FBJ murine osteosarcoma viral oncogene homolog
FOXN3	forkhead box N3
FRMD4B	FERM domain containing 4B
FTH1	ferritin, heavy polypeptide 1; ferritin, heavy polypeptide-like 16; similar to ferritin, heavy polypeptide 1; ferritin, heavy polypeptide-like 3 pseudogene
FXYD1	FXYD domain containing ion transport regulator 1
G3BP1	GTPase activating protein (SH3 domain) binding protein 1
GABARAPL1	GABA(A) receptors associated protein like 3 (pseudogene); GABA(A) receptor-associated protein like 1
GADD45B	growth arrest and DNA-damage-inducible, beta
GANAB	glucosidase, alpha; neutral AB
GAS1	growth arrest-specific 1
GAS6	similar to growth arrest-specific 6; growth arrest-specific 6
GATA6	GATA binding protein 6
GBP2	guanylate binding protein 2, interferon-inducible
GBP3	guanylate binding protein 3
GBP7	guanylate binding protein 7
GCSH	similar to Glycine cleavage system H protein, mitochondrial precursor; glycine cleavage system protein H (aminomethyl carrier); similar to Glycine cleavage system H protein, mitochondrial
GDA	guanine deaminase
GEM	GTP binding protein overexpressed in skeletal muscle
GFM2	G elongation factor, mitochondrial 2
GFPT2	glutamine-fructose-6-phosphate transaminase 2
GJA1	gap junction protein, alpha 1, 43kDa
GJB5	gap junction protein, beta 5, 31.1kDa
GNB2	guanine nucleotide binding protein (G protein), beta polypeptide 2
GOLGA4	golgi autoantigen, golgin subfamily a, 4
GOLGB1	golgin B1, golgi integral membrane protein
GPC3	glypican 3
GPC4	glypican 4
GPCPD1	hypothetical protein KIAA1434
GPM6A	glycoprotein M6A
GPR116	G protein-coupled receptor 116
GPR133	G protein-coupled receptor 133
GPR64	G protein-coupled receptor 64
GPRC5B	G protein-coupled receptor, family C, group 5, member B
GPX8	glutathione peroxidase 8 (putative)
GSR	glutathione reductase

GSTA3	glutathione S-transferase alpha 3
GSTM1	glutathione S-transferase mu 1
GSTM4	glutathione S-transferase mu 4
GUCY1A3	guanylate cyclase 1, soluble, alpha 3
H3F3A	H3 histone, family 3B (H3.3B); H3 histone, family 3A pseudogene; H3 histone, family 3A; similar to H3 histone, family 3B; similar to histone H3.3B
HDAC3	histone deacetylase 3
HDAC5	histone deacetylase 5
HEG1	HEG homolog 1 (zebrafish)
HERPUD2	HERPUD family member 2
HES1	hairy and enhancer of split 1, (Drosophila)
HEXB	hexosaminidase B (beta polypeptide)
HIST1H1C	histone cluster 1, H1c
HMGB1	high-mobility group box 1; high-mobility group box 1-like 10
HNRNPH1	heterogeneous nuclear ribonucleoprotein H1 (H)
HNRNPH2	ribosomal protein L36a pseudogene 51; ribosomal protein L36a pseudogene 37; ribosomal protein L36a pseudogene 49; heterogeneous nuclear ribonucleoprotein H2 (H'); ribosomal protein L36a
HNRNPL	similar to heterogeneous nuclear ribonucleoprotein L-like; heterogeneous nuclear ribonucleoprotein L
HNRNPM	heterogeneous nuclear ribonucleoprotein M
HNRNPR	heterogeneous nuclear ribonucleoprotein R
HOOK3	hook homolog 3 (Drosophila)
HOXA5	homeobox A5
HP1BP3	heterochromatin protein 1, binding protein 3
HSP90AA1	heat shock protein 90kDa alpha (cytosolic), class A member 2; heat shock protein 90kDa alpha (cytosolic), class A member 1
HSP90AB1	heat shock protein 90kDa alpha (cytosolic), class B member 1
HSP90B1	heat shock protein 90kDa beta (Grp94), member 1
HSPA12A	heat shock 70kDa protein 12A
HSPA2	heat shock 70kDa protein 2
HSPB1	heat shock 27kDa protein-like 2 pseudogene; heat shock 27kDa protein 1
HSPB8	heat shock 22kDa protein 8
ID1	inhibitor of DNA binding 1, dominant negative helix-loop-helix protein
ID2	inhibitor of DNA binding 2, dominant negative helix-loop-helix protein
IER2	immediate early response 2
IFI35	interferon-induced protein 35

IFIT3	interferon-induced protein with tetratricopeptide repeats 3
IFITM3	interferon induced transmembrane protein 3 (1-8U)
IFNAR2	interferon (alpha, beta and omega) receptor 2
IFNGR1	interferon gamma receptor 1
IFRD1	interferon-related developmental regulator 1
IFT74	intraflagellar transport 74 homolog (Chlamydomonas)
IGF1R	insulin-like growth factor 1 receptor
IGFBP5	insulin-like growth factor binding protein 5
IGFBP6	insulin-like growth factor binding protein 6
IL16	interleukin 16 (lymphocyte chemoattractant factor)
IL17RE	interleukin 17 receptor E
IL6ST	interleukin 6 signal transducer (gp130, oncostatin M receptor)
ILDR2	immunoglobulin-like domain containing receptor 2
ILF3	interleukin enhancer binding factor 3, 90kDa
IMPAD1	inositol monophosphatase domain containing 1
INTS10	integrator complex subunit 10
IQSEC1	IQ motif and Sec7 domain 1
IRAK4	interleukin-1 receptor-associated kinase 4
IRF2BP2	interferon regulatory factor 2 binding protein 2
IRF7	interferon regulatory factor 7
IRS2	insulin receptor substrate 2
ITCH	itchy E3 ubiquitin protein ligase homolog (mouse)
ITGA6	integrin, alpha 6
ITPR2	inositol 1,4,5-triphosphate receptor, type 2
JMJD1C	jumonji domain containing 1C
JUN	jun oncogene
JUNB	jun B proto-oncogene
JUND	jun D proto-oncogene
JUP	junction plakoglobin
KANK1	KN motif and ankyrin repeat domains 1; similar to ankyrin repeat domain protein 15 isoform b
KCNAB1	potassium voltage-gated channel, shaker-related subfamily, beta member 1
KDELR1	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 1
KDM5A	lysine (K)-specific demethylase 5A
KDM6B	lysine (K)-specific demethylase 6B
KDR	kinase insert domain receptor (a type III receptor tyrosine kinase)
KEAP1	kelch-like ECH-associated protein 1
KIF1B	kinesin family member 1B
KIF5B	kinesin family member 5B
KLF10	Kruppel-like factor 10

KLF2	Kruppel-like factor 2 (lung)
KLF4	Kruppel-like factor 4 (gut)
KLF6	Kruppel-like factor 6
KLF7	Kruppel-like factor 7 (ubiquitous)
KLF9	Kruppel-like factor 9
KPNA1	karyopherin alpha 1 (importin alpha 5)
KPNA3	karyopherin alpha 3 (importin alpha 4)
KRCC1	lysine-rich coiled-coil 1
KRT14	keratin 14
KTN1	kinectin 1 (kinesin receptor)
LAMA4	laminin, alpha 4
LAMP2	lysosomal-associated membrane protein 2
LARS2	leucyl-tRNA synthetase 2, mitochondrial
LASS2	LAG1 homolog, ceramide synthase 2
LASS4	LAG1 homolog, ceramide synthase 4
LGALS7	lectin, galactoside-binding, soluble, 7; lectin, galactoside-binding, soluble, 7B
LIMCH1	LIM and calponin homology domains 1
LIMS2	LIM and senescent cell antigen-like domains 2
LMAN1	lectin, mannose-binding, 1
LPAR2	lysophosphatidic acid receptor 2
LRRC20	leucine rich repeat containing 20
LRRC58	leucine rich repeat containing 58
LRRC61	leucine rich repeat containing 61
LRRN4	leucine rich repeat neuronal 4
LRRN4CL	LRRN4 C-terminal like
LTBP4	latent transforming growth factor beta binding protein 4
LUC7L3	cisplatin resistance-associated overexpressed protein
MAF	v-maf musculoaponeurotic fibrosarcoma oncogene homolog (avian)
MAGED1	melanoma antigen family D, 1
MAGT1	magnesium transporter 1
MALAT1	metastasis associated lung adenocarcinoma transcript 1 (non-protein coding)
MANF	mesencephalic astrocyte-derived neurotrophic factor
MAOA	monoamine oxidase A
MAP3K3	mitogen-activated protein kinase kinase kinase 3
MAPK1	mitogen-activated protein kinase 1
MAPKAPK3	mitogen-activated protein kinase-activated protein kinase 3
MAPRE2	microtubule-associated protein, RP/EB family, member 2
MARCKSL1	MARCKS-like 1
MAT2A	methionine adenosyltransferase II, alpha

MAT2B	methionine adenosyltransferase II, beta
MATR3	matrin 3
MED13L	mediator complex subunit 13-like
MED21	mediator complex subunit 21
MEF2C	myocyte enhancer factor 2C
MEIS2	Meis homeobox 2
MESDC1	mesoderm development candidate 1
METAP2	methionyl aminopeptidase 2
MFHAS1	malignant fibrous histiocytoma amplified sequence 1
MGLL	monoglyceride lipase
MGST1	microsomal glutathione S-transferase 1
MLL3	myeloid/lymphoid or mixed-lineage leukemia 3
MORF4L2	mortality factor 4 like 2
MPDZ	multiple PDZ domain protein
MPHOSPH8	M-phase phosphoprotein 8
MRAS	muscle RAS oncogene homolog
MRGPRF	MAS-related GPR, member F
MSN	moesin
MTDH	metadherin
MTMR6	myotubularin related protein 6
MUT	methylmalonyl Coenzyme A mutase
MXD4	MAX dimerization protein 4
MYH10	myosin, heavy chain 10, non-muscle
MYL12A	myosin, light chain 12A, regulatory, non-sarcomeric
MYL7	myosin, light chain 7, regulatory
MYLIP	myosin regulatory light chain interacting protein
MYST4	MYST histone acetyltransferase (monocytic leukemia) 4
NAA25	chromosome 12 open reading frame 30
NAGA	N-acetylgalactosaminidase, alpha-
NCKAP1	NCK-associated protein 1
NCOA1	nuclear receptor coactivator 1
NCOA4	nuclear receptor coactivator 4
NCOR1	nuclear receptor co-repressor 1
NDN	necdin homolog (mouse)
NDST1	N-deacetylase/N-sulfotransferase (heparan glucosaminyl) 1
NDUFA4	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 4, 9kDa
NEDD4	neural precursor cell expressed, developmentally down-regulated 4
NF1	neurofibromin 1
NFE2L1	nuclear factor (erythroid-derived 2)-like 1
NFIA	nuclear factor I/A
NFIC	nuclear factor I/C (CCAAT-binding transcription factor)

NFIX	nuclear factor I/X (CCAAT-binding transcription factor)
NFKB2	nuclear factor of kappa light polypeptide gene enhancer in B-cells 2 (p49/p100)
NFKBIA	nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha
NFKBIZ	nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta
NFYC	nuclear transcription factor Y, gamma
NID2	nidogen 2 (osteonidogen)
NINL	ninein-like
NIPAL3	NIPA-like domain containing 3
NIPBL	Nipped-B homolog (Drosophila)
NKAIN4	Na+/K+ transporting ATPase interacting 4
NKD1	naked cuticle homolog 1 (Drosophila)
NNMT	nicotinamide N-methyltransferase
NOD1	nucleotide-binding oligomerization domain containing 1
NPR1	natriuretic peptide receptor A/guanylate cyclase A (atrionatriuretic peptide receptor A)
NR1D1	nuclear receptor subfamily 1, group D, member 1
NR3C1	nuclear receptor subfamily 3, group C, member 1 (glucocorticoid receptor)
NR4A1	nuclear receptor subfamily 4, group A, member 1
NRGN	neurogranin (protein kinase C substrate, RC3)
NUCKS1	nuclear casein kinase and cyclin-dependent kinase substrate 1
OAT	ornithine aminotransferase (gyrate atrophy)
OGDH	oxoglutarate (alpha-ketoglutarate) dehydrogenase (lipoamide)
OGN	osteoglycin
OPA3	optic atrophy 3 (autosomal recessive, with chorea and spastic paraplegia)
ORAI3	ORAI calcium release-activated calcium modulator 3
OSR1	odd-skipped related 1 (Drosophila)
OXCT1	3-oxoacid CoA transferase 1
OXNAD1	oxidoreductase NAD-binding domain containing 1
PARD3B	par-3 partitioning defective 3 homolog B (C. elegans)
PARP14	poly (ADP-ribose) polymerase family, member 14
PARP4	poly (ADP-ribose) polymerase family, member 4
PARVB	parvin, beta
PBX1	pre-B-cell leukemia homeobox 1
PCDH15	protocadherin 15
PCDHGB5	protocadherin gamma subfamily B, 5
PCM1	pericentriolar material 1

PDAP1	PDGFA associated protein 1; similar to PDGFA associated protein 1
PCD6IP	programmed cell death 6 interacting protein
PDE4DIP	hypothetical protein LOC100134230; similar to KIAA0454 protein; similar to phosphodiesterase 4D interacting protein isoform 2; phosphodiesterase 4D interacting protein
PDIA3	protein disulfide isomerase family A, member 3
PDIA4	protein disulfide isomerase family A, member 4
PDPN	podoplanin
PEF1	penta-EF-hand domain containing 1
PELI1	pellino homolog 1 (Drosophila)
PER1	period homolog 1 (Drosophila)
PF4	platelet factor 4
PFN1	profilin 1
PGCP	plasma glutamate carboxypeptidase
PGRMC1	progesterone receptor membrane component 1
PHF21A	PHD finger protein 21A
PHF3	PHD finger protein 3
PHIP	pleckstrin homology domain interacting protein
PIGT	phosphatidylinositol glycan anchor biosynthesis, class T
PIK3C2A	phosphoinositide-3-kinase, class 2, alpha polypeptide
PIM1	pim-1 oncogene
PITPNM2	phosphatidylinositol transfer protein, membrane-associated 2
PKHD1L1	polycystic kidney and hepatic disease 1 (autosomal recessive)-like 1
PKNOX1	PBX/knotted 1 homeobox 1
PLA2G4A	phospholipase A2, group IVA (cytosolic, calcium-dependent)
PLAT	plasminogen activator, tissue
PLCE1	phospholipase C, epsilon 1
PLK1S1	non-protein coding RNA 153
PLK2	polo-like kinase 2 (Drosophila)
PLOD2	procollagen-lysine, 2-oxoglutarate 5-dioxygenase 2
PLXDC1	plexin domain containing 1
PLXDC2	plexin domain containing 2
PLXNA4	plexin A4
PMP22	peripheral myelin protein 22
PNRC1	proline-rich nuclear receptor coactivator 1
PODN	podocan
PPAP2A	phosphatidic acid phosphatase type 2A
PPBP	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)
PPFIBP2	PTPRF interacting protein, binding protein 2 (liprin beta 2)
PPIG	peptidylprolyl isomerase G (cyclophilin G)

PPL	periplakin
PPP1CB	protein phosphatase 1, catalytic subunit, beta isoform; speedy homolog A ( <i>Xenopus laevis</i> )
PPP1R12A	protein phosphatase 1, regulatory (inhibitor) subunit 12A
PPP1R15A	protein phosphatase 1, regulatory (inhibitor) subunit 15A
PPP3CA	protein phosphatase 3 (formerly 2B), catalytic subunit, alpha isoform
PPPDE1	PPPDE peptidase domain containing 1
PQLC3	PQ loop repeat containing 3
PRELP	proline/arginine-rich end leucine-rich repeat protein
PRG4	proteoglycan 4
PRKAR2A	protein kinase, cAMP-dependent, regulatory, type II, alpha
PRPF40A	PRP40 pre-mRNA processing factor 40 homolog A ( <i>S. cerevisiae</i> )
PRR13	proline rich 13
PRSS23	protease, serine, 23
PSD	pleckstrin and Sec7 domain containing
PSIP1	PC4 and SFRS1 interacting protein 1
PSMB2	proteasome (prosome, macropain) subunit, beta type, 2
PSMD11	proteasome (prosome, macropain) 26S subunit, non-ATPase, 11
PSMD7	proteasome (prosome, macropain) 26S subunit, non-ATPase, 7
PTGES3	prostaglandin E synthase 3 (cytosolic)
PTGIS	prostaglandin I2 (prostacyclin) synthase
PTGS1	prostaglandin-endoperoxide synthase 1 (prostaglandin G/H synthase and cyclooxygenase)
PTMA	hypothetical LOC728026; prothymosin, alpha; hypothetical gene supported by BC013859; prothymosin, alpha pseudogene 4 (gene sequence 112)
PTP4A2	protein tyrosine phosphatase type IVA, member 2
PTPLAD2	protein tyrosine phosphatase-like A domain containing 2
PTPRD	protein tyrosine phosphatase, receptor type, D
PTPRF	protein tyrosine phosphatase, receptor type, F
PTRF	polymerase I and transcript release factor
QRICH1	glutamine-rich 1
QSER1	glutamine and serine rich 1
RAB11FIP1	RAB11 family interacting protein 1 (class I)
RAB1B	RAB1B, member RAS oncogene family
RAB5C	RAB5C, member RAS oncogene family
RAB6B	RAB6B, member RAS oncogene family
RABGAP1L	RAB GTPase activating protein 1-like
RALBP1	hypothetical LOC100129773; ralA binding protein 1
RALY	RNA binding protein, autoantigenic (hnRNP-associated with lethal yellow homolog (mouse))

RARRES2	retinoic acid receptor responder (tazarotene induced) 2
RB1CC1	RB1-inducible coiled-coil 1
RBBP6	retinoblastoma binding protein 6
RBBP8	retinoblastoma binding protein 8
RBM25	RNA binding motif protein 25
RBM27	RNA binding motif protein 27
RBM3	RNA binding motif (RNP1, RRM) protein 3
RBPMS	RNA binding protein with multiple splicing
RDX	radixin
REST	RE1-silencing transcription factor
RGMA	RGM domain family, member A
RGS10	regulator of G-protein signaling 10
RHOB	ras homolog gene family, member B
RHOJ	ras homolog gene family, member J
RHOU	ras homolog gene family, member U
RNASE4	ribonuclease, RNase A family, 4
RND3	Rho family GTPase 3
RNF167	ring finger protein 167
RNF20	ring finger protein 20
ROCK1	similar to Rho-associated, coiled-coil containing protein kinase 1; Rho-associated, coiled-coil containing protein kinase 1
ROCK2	Rho-associated, coiled-coil containing protein kinase 2
RPP25	ribonuclease P/MRP 25kDa subunit
RRAS2	related RAS viral (r-ras) oncogene homolog 2; similar to related RAS viral (r-ras) oncogene homolog 2
RSPO1	R-spondin homolog (Xenopus laevis)
RTF1	Rtf1, Paf1/RNA polymerase II complex component, homolog (S. cerevisiae)
RTN1	reticulon 1
RYK	RYK receptor-like tyrosine kinase
SARNP	SAP domain containing ribonucleoprotein
SAT1	spermidine/spermine N1-acetyltransferase 1
SBSN	suprabasin
SDC4	syndecan 4
SDPR	serum deprivation response (phosphatidylserine binding protein)
SEC62	SEC62 homolog (S. cerevisiae)
SECISBP2	SECIS binding protein 2
SEMA5A	sema domain, seven thrombospondin repeats (type 1 and type 1-like), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 5A
SENP6	SUMO1/sentrin specific peptidase 6
SEP15	15 kDa selenoprotein

SEPT9	septin 9
SERINC5	serine incorporator 5
SERPING1	serpin peptidase inhibitor, clade G (C1 inhibitor), member 1
SERPINH1	serpin peptidase inhibitor, clade H (heat shock protein 47), member 1, (collagen binding protein 1)
SESN1	sestrin 1
SETD2	SET domain containing 2
SF3B1	splicing factor 3b, subunit 1, 155kDa
SF3B4	splicing factor 3b, subunit 4, 49kDa
SFRS18	splicing factor, arginine/serine-rich 18
SHC1	SHC (Src homology 2 domain containing) transforming protein 1
SHFM1	split hand/foot malformation (ectrodactyly) type 1
SIAE	sialic acid acetylesterase
SIRT2	sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae)
SLC10A3	solute carrier family 10 (sodium/bile acid cotransporter family), member 3
SLC16A1	solute carrier family 16, member 1 (monocarboxylic acid transporter 1)
SLC1A5	solute carrier family 1 (neutral amino acid transporter), member 5
SLC26A3	solute carrier family 26, member 3
SLC27A3	solute carrier family 27 (fatty acid transporter), member 3
SLC38A1	solute carrier family 38, member 1
SLC39A8	solute carrier family 39 (zinc transporter), member 8
SLC43A3	solute carrier family 43, member 3
SLC4A4	solute carrier family 4, sodium bicarbonate cotransporter, member 4
SLC6A4	solute carrier family 6 (neurotransmitter transporter, serotonin), member 4
SLC6A6	solute carrier family 6 (neurotransmitter transporter, taurine), member 6
SLC8A1	solute carrier family 8 (sodium/calcium exchanger), member 1
SLC9A3R1	solute carrier family 9 (sodium/hydrogen exchanger), member 3 regulator 1
SLPI	secretory leukocyte peptidase inhibitor
SLTM	SAFB-like, transcription modulator
SLU7	SLU7 splicing factor homolog (S. cerevisiae)
SLURP1	secreted LY6/PLAUR domain containing 1
SMAD4	SMAD family member 4
SMARCA2	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2

SMARCA5	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 5
SMC2	structural maintenance of chromosomes 2
SMC3	structural maintenance of chromosomes 3
SMC4	structural maintenance of chromosomes 4
SMC6	structural maintenance of chromosomes 6
SMCHD1	structural maintenance of chromosomes flexible hinge domain containing 1
SMPD3	sphingomyelin phosphodiesterase 3, neutral membrane (neutral sphingomyelinase II)
SNRNP70	small nuclear ribonucleoprotein 70kDa (U1)
SNTB2	syntrophin, beta 2 (dystrophin-associated protein A1, 59kDa, basic component 2)
SOAT1	sterol O-acyltransferase 1
SOCS3	suppressor of cytokine signaling 3
SOD3	superoxide dismutase 3, extracellular
SORBS1	sorbin and SH3 domain containing 1
SORBS3	sorbin and SH3 domain containing 3
SOX6	SRY (sex determining region Y)-box 6
SP100	SP100 nuclear antigen
SPAG9	sperm associated antigen 9
SPARC	secreted protein, acidic, cysteine-rich (osteonectin)
SPEN	spen homolog, transcriptional regulator (Drosophila)
SPINT2	serine peptidase inhibitor, Kunitz type, 2
SPOCK2	sparc/osteonectin, cwcv and kazal-like domains proteoglycan (testican) 2
SPON2	spondin 2, extracellular matrix protein
SPOP	speckle-type POZ protein
SRC	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)
SRRM1	serine/arginine repetitive matrix 1
SSH2	slingshot homolog 2 (Drosophila)
SSR3	signal sequence receptor, gamma (translocon-associated protein gamma)
ST3GAL1	ST3 beta-galactoside alpha-2,3-sialyltransferase 1
STAG1	stromal antigen 1
STAR	steroidogenic acute regulatory protein
STARD5	StAR-related lipid transfer (START) domain containing 5
STAT3	signal transducer and activator of transcription 3 (acute-phase response factor)
STIM1	stromal interaction molecule 1
STK10	serine/threonine kinase 10

STK40	serine/threonine kinase 40
STMN2	stathmin-like 2
STRA6	stimulated by retinoic acid gene 6 homolog (mouse)
STRN3	striatin, calmodulin binding protein 3
SULF1	sulfatase 1
SULF2	sulfatase 2
SUPT16H	suppressor of Ty 16 homolog (S. cerevisiae); suppressor of Ty 16 homolog (S. cerevisiae) pseudogene
SV2A	synaptic vesicle glycoprotein 2A
SYNE1	spectrin repeat containing, nuclear envelope 1
SYNE2	spectrin repeat containing, nuclear envelope 2
SYT11	synaptotagmin XI
SYTL1	synaptotagmin-like 1
TAF3	TAF3 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 140kDa
TAF7	TAF7 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 55kDa
TAPBP	TAP binding protein (tapasin)
TBC1D15	TBC1 domain family, member 15
TBCEL	tubulin folding cofactor E-like
TBL1X	transducin (beta)-like 1X-linked
TBX18	T-box 18
TCEAL8	transcription elongation factor A (SII)-like 8
TCF7L1	transcription factor 7-like 1 (T-cell specific, HMG-box)
TFDP2	transcription factor Dp-2 (E2F dimerization partner 2)
TGFB1I1	transforming growth factor beta 1 induced transcript 1
TGFB2	transforming growth factor, beta 2
TGFBR2	transforming growth factor, beta receptor II (70/80kDa)
TGM2	transglutaminase 2 (C polypeptide, protein-glutamine-gamma-glutamyltransferase)
THBD	thrombomodulin
THBS1	thrombospondin 1
THOC2	THO complex 2
THRAP3	thyroid hormone receptor associated protein 3
THSD4	thrombospondin, type I, domain containing 4
TIMP2	TIMP metallopeptidase inhibitor 2
TIRAP	toll-interleukin 1 receptor (TIR) domain containing adaptor protein
TLR2	toll-like receptor 2
TM4SF1	transmembrane 4 L six family member 1
TM4SF5	transmembrane 4 L six family member 5
TMCC3	transmembrane and coiled-coil domain family 3

TMCO1	transmembrane and coiled-coil domains 1
TMCO7	transmembrane and coiled-coil domains 7
TMED2	transmembrane emp24 domain trafficking protein 2
TMEM119	transmembrane protein 119
TMEM140	transmembrane protein 140
TMEM151A	transmembrane protein 151A
TMEM221	transmembrane protein 221
TMEM50A	transmembrane protein 50A
TMEM98	similar to transmembrane protein 98; transmembrane protein 98
TMOD3	tropomodulin 3 (ubiquitous)
TMPO	thymoprotein
TMSB4X	thymosin-like 2 (pseudogene); thymosin-like 1 (pseudogene); thymosin beta 4, X-linked
TNXB	tenascin XB; tenascin XA pseudogene
TOB2	transducer of ERBB2, 2
TOPORS	topoisomerase I binding, arginine/serine-rich
TPM3	tropomyosin 3
TPPP3	tubulin polymerization-promoting protein family member 3
TPT1	similar to tumor protein, translationally-controlled 1; tumor protein, translationally-controlled 1
TRAFD1	TRAF-type zinc finger domain containing 1
TRIB1	tribbles homolog 1 (Drosophila)
TRIM8	tripartite motif-containing 8
TRPM7	transient receptor potential cation channel, subfamily M, member 7
TSC22D3	TSC22 domain family, member 3; GRAM domain containing 4
TSHZ1	teashirt zinc finger homeobox 1
TSIX	XIST antisense RNA (non-protein coding)
TSPAN31	tetraspanin 31
TSPAN5	tetraspanin 5
TTC28	chromosome 6 open reading frame 35; hCG1820764; tetra-tricopeptide repeat domain 28
TTC38	tetra-tricopeptide repeat domain 38
TUBA1A	tubulin, alpha 1a
TUBB2A	tubulin, beta 2A
TWSG1	twisted gastrulation homolog 1 (Drosophila)
TXNDC5	thioredoxin domain containing 5 (endoplasmic reticulum); muted homolog (mouse)
TXNRD1	thioredoxin reductase 1; hypothetical LOC100130902
UAP1	UDP-N-acetylglucosamine pyrophosphorylase 1
UBA7	ubiquitin-like modifier activating enzyme 7
UBE2D1	ubiquitin-conjugating enzyme E2D 1 (UBC4/5 homolog, yeast)

UBE2L6	ubiquitin-conjugating enzyme E2L 6
UBE2N	ubiquitin-conjugating enzyme E2N (UBC13 homolog, yeast)
UBE2V1	ubiquitin-conjugating enzyme E2 variant 1; ubiquitin-conjugating enzyme E2 variant 1 pseudogene 2; transmembrane protein 189; TMEM189-UBE2V1 readthrough transcript
UBQLN2	ubiquilin 2
UBXN2A	UBX domain protein 2A
UBXN4	UBX domain protein 4
UGDH	UDP-glucose dehydrogenase
UPK1B	uroplakin 1B
UPK3B	uroplakin 3B
USP16	ubiquitin specific peptidase 16
USP2	ubiquitin specific peptidase 2
USP25	ubiquitin specific peptidase 25
USP54	ubiquitin specific peptidase 54
USP8	ubiquitin specific peptidase 8
UTP20	similar to Down-regulated in metastasis protein (Key-1A6 protein) (Novel nucleolar protein 73) (NNP73); UTP20, small subunit (SSU) processome component, homolog (yeast)
VAT1	vesicle amine transport protein 1 homolog (T. californica)
VIM	vimentin
VPS13A	vacuolar protein sorting 13 homolog A (S. cerevisiae)
VWA5A	von Willebrand factor A domain containing 5A
WAC	WW domain containing adaptor with coiled-coil
WASF2	WAS protein family, member 2
WDR26	WD repeat domain 26
WDR92	WD repeat domain 92
WFDC1	WAP four-disulfide core domain 1
WLS	G protein-coupled receptor 177
WNT4	wingless-type MMTV integration site family, member 4
WRNIP1	Werner helicase interacting protein 1
WT1	Wilms tumor 1
WWC2	WW and C2 domain containing 2
XDH	xanthine dehydrogenase
XIST	X (inactive)-specific transcript (non-protein coding)
YIPF5	Yip1 domain family, member 5
YWHAZ	tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein, zeta polypeptide
ZBTB16	zinc finger and BTB domain containing 16
ZBTB20	zinc finger and BTB domain containing 20
ZBTB4	zinc finger and BTB domain containing 4
ZBTB7C	zinc finger and BTB domain containing 7C

ZC3H13	zinc finger CCCH-type containing 13
ZC3H18	zinc finger CCCH-type containing 18
ZCCHC11	zinc finger, CCHC domain containing 11
ZCCHC3	zinc finger, CCHC domain containing 3
ZFAND6	zinc finger, AN1-type domain 6
ZFHX4	zinc finger homeobox 4
ZFP36	zinc finger protein 36, C3H type, homolog (mouse)
ZMAT1	zinc finger, matrin type 1
ZRSR1	zinc finger (CCCH type), RNA-binding motif and serine/arginine rich 1
ZZEF1	zinc finger, ZZ-type with EF-hand domain 1

**[0038]** The gene names listed in Table 7 and Table 8 are common names. NCBI Gene ID numbers for each of the genes listed in Table 7 or Table 8 can be obtained by searching the “Gene” Database of the NCBI (available on the World Wide Web at <http://www.ncbi.nlm.nih.gov/>) using the common name as the query and selecting the first returned *Homo sapiens* (for the genes in Table 8) or *Mus musculus* gene (for the genes in Table 7). Other genes may be obtained using the UCSC genome browser (available on the World Wide Web at <http://genome.ucsc.edu>) using the Gene Sorter function. Human homologs of mouse genes can be readily identified, e.g. the identified homologs in the NCBI database, or by querying databases such as BLAST. In certain embodiments, the marker gene(s) are selected from the genes listed in Table 7, Table 8, or Table 14.

**[0039]** In a CTC, the marker genes listed in Table 7, Table 8, or Table 14 can be upregulated, e.g. for marker genes listed in Table 7, Table 8, or Table 14, if the measured marker gene expression in a cell or sample is higher as compared to a reference level of that marker gene’s expression, then the cell is identified as a CTC and/or the sample is identified as comprising CTCs. Preferably, once looks at a statistically significant change. However, even if a few genes in a group do not differ from normal, a sample can be identified as comprising CTCs if the overall change of the group shows a significant change, preferably a statistically significant change. All possible combinations of 2 or more of the indicated markers are contemplated herein.

**[0040]** The level of a gene expression product of a marker gene in Table 7, Table 8, or Table 14 which is higher than a reference level of that marker gene by at least about 10% than the reference amount, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 80%, at least about 100%, at least about 200%, at least about 300%, at least about 500% or at least about 1000% or more, is indicative of the presence of a CTC.

**[0041]** In some embodiments, the reference can be a level of expression of the marker gene product in a cell or population of cells which are not CTCs, e.g. the average level in non-circulating tumor cells and/or circulating cells which are not cancer cells. In some embodiments, the reference

can also be a level of expression of the marker gene product in a control sample, a pooled sample of control individuals or a numeric value or range of values based on the same.

**[0042]** In some embodiments, the methods and assays described herein include (a) transforming the gene expression product into a detectable gene target; (b) measuring the amount of the detectable gene target; and (c) comparing the amount of the detectable gene target to an amount of a reference, wherein if the amount of the detectable gene target is statistically significantly different than the amount of the reference level, the presence and/or level of CTCs is determined. In some embodiments, if the amount of the detectable gene target is not statistically significantly different than the amount of the reference level, the sample is identified as not comprising CTCs.

**[0043]** As used herein, the term “transforming” or “transformation” refers to changing an object or a substance, e.g., biological sample, nucleic acid or protein, into another substance. The transformation can be physical, biological or chemical. Exemplary physical transformation includes, but not limited to, pre-treatment of a biological sample, e.g., from whole blood to blood serum by differential centrifugation. A biological/chemical transformation can involve at least one enzyme and/or a chemical reagent in a reaction. For example, a DNA sample can be digested into fragments by one or more restriction enzyme, or an exogenous molecule can be attached to a fragmented DNA sample with a ligase. In some embodiments, a DNA sample can undergo enzymatic replication, e.g., by polymerase chain reaction (PCR).

**[0044]** Methods to measure gene expression products associated with the marker genes described herein are well known to a skilled artisan. Such methods to measure gene expression products, e.g., protein level, include ELISA (enzyme linked immunosorbent assay), western blot, FACS, radioimmunological assay; (RIA); sandwich assay; fluorescent in situ hybridization ( FISH); immunohistological staining; immunoelectrophoresis; immunoprecipitation, and immunofluorescence using detection reagents such as an antibody or protein binding agents. Alternatively, a peptide can be detected in a subject by introducing into a subject a labeled anti-peptide antibody and other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in the subject is detected by standard imaging techniques.

**[0045]** For example, antibodies for the polypeptide expression products of the marker genes described herein are commercially available and can be used for the purposes of the invention to measure protein expression levels, e.g. anti-IGFBP5 (Cat. No. 4255; Abcam; Cambridge, MA). Alternatively, since the amino acid sequences for the marker genes described herein are known and publically available at NCBI website, one of skill in the art can raise their own antibodies against these proteins of interest for the purpose of the invention. The amino acid sequences of the marker genes described herein have been assigned NCBI accession numbers for different species such as human, mouse and rat.

**[0046]** In some embodiments, immunohistochemistry (“IHC”) and immunocytochemistry (“ICC”) techniques can be used. IHC is the application of immunochemistry to tissue sections, whereas ICC is the application of immunochemistry to cells or tissue imprints after they have undergone specific cytological preparations such as, for example, liquid-based preparations. Immunochemistry is a family of techniques based on the use of an antibody, wherein the antibodies are used to specifically target molecules inside or on the surface of cells. The antibody typically contains a marker that will undergo a biochemical reaction, and thereby experience a change color, upon encountering the targeted molecules. In some instances, signal amplification can be integrated into the particular protocol, wherein a secondary antibody, that includes the marker stain or marker signal, follows the application of a primary specific antibody.

**[0047]** In some embodiments, the assay can be a Western blot analysis. Alternatively, proteins can be separated by two-dimensional gel electrophoresis systems. Two-dimensional gel electrophoresis is well known in the art and typically involves iso-electric focusing along a first dimension followed by SDS-PAGE electrophoresis along a second dimension. These methods also require a considerable amount of cellular material. The analysis of 2D SDS-PAGE gels can be performed by determining the intensity of protein spots on the gel, or can be performed using immune detection. In other embodiments, protein samples are analyzed by mass spectroscopy.

**[0048]** Immunological tests can be used with the methods and assays described herein and include, for example, competitive and non-competitive assay systems using techniques such as Western blots, radioimmunoassay (RIA), ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, immunodiffusion assays, agglutination assays, e.g. latex agglutination, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, e.g. FIA (fluorescence-linked immunoassay), chemiluminescence immunoassays (CLIA), electrochemiluminescence immunoassay (ECLIA, counting immunoassay (CIA), lateral flow tests or immunoassay (LFIA), magnetic immunoassay (MIA), and protein A immunoassays. Methods for performing such assays are known in the art, provided an appropriate antibody reagent is available. In some embodiment, the immunoassay can be a quantitative or a semi-quantitative immunoassay.

**[0049]** An immunoassay is a biochemical test that measures the concentration of a substance in a biological sample, typically a fluid sample such as serum, using the interaction of an antibody or antibodies to its antigen. The assay takes advantage of the highly specific binding of an antibody with its antigen. For the methods and assays described herein, specific binding of the target polypeptides with respective proteins or protein fragments, or an isolated peptide, or a fusion protein described herein occurs in the immunoassay to form a target protein/peptide complex. The complex is then detected by a variety of methods known in the art. An immunoassay also often involves the use of a detection antibody.

**[0050]** Enzyme-linked immunosorbent assay, also called ELISA, enzyme immunoassay or EIA, is a biochemical technique used mainly in immunology to detect the presence of an antibody or an antigen in a sample. The ELISA has been used as a diagnostic tool in medicine and plant pathology, as well as a quality control check in various industries.

**[0051]** In one embodiment, an ELISA involving at least one antibody with specificity for the particular desired antigen (i.e. a marker gene polypeptide as described herein) can also be performed. A known amount of sample and/or antigen is immobilized on a solid support (usually a polystyrene micro titer plate). Immobilization can be either non-specific (e.g., by adsorption to the surface) or specific (e.g. where another antibody immobilized on the surface is used to capture antigen or a primary antibody). After the antigen is immobilized, the detection antibody is added, forming a complex with the antigen. The detection antibody can be covalently linked to an enzyme, or can itself be detected by a secondary antibody which is linked to an enzyme through bio-conjugation. Between each step the plate is typically washed with a mild detergent solution to remove any proteins or antibodies that are not specifically bound. After the final wash step the plate is developed by adding an enzymatic substrate to produce a visible signal, which indicates the quantity of antigen in the sample. Older ELISAs utilize chromogenic substrates, though newer assays employ fluorogenic substrates with much higher sensitivity.

**[0052]** In another embodiment, a competitive ELISA is used. Purified antibodies that are directed against a target polypeptide or fragment thereof are coated on the solid phase of multi-well plate, i.e., conjugated to a solid surface. A second batch of purified antibodies that are not conjugated on any solid support is also needed. These non-conjugated purified antibodies are labeled for detection purposes, for example, labeled with horseradish peroxidase to produce a detectable signal. A sample (e.g., tumor, blood, serum or urine) from a subject is mixed with a known amount of desired antigen (e.g., a known volume or concentration of a sample comprising a target polypeptide) together with the horseradish peroxidase labeled antibodies and the mixture is then added to coated wells to form competitive combination. After incubation, if the polypeptide level is high in the sample, a complex of labeled antibody reagent-antigen will form. This complex is free in solution and can be washed away. Washing the wells will remove the complex. Then the wells are incubated with TMB (3, 3', 5, 5'-tetramethylbenzidine) color development substrate for localization of horseradish peroxidase-conjugated antibodies in the wells. There will be no color change or little color change if the target polypeptide level is high in the sample. If there is little or no target polypeptide present in the sample, a different complex is formed, the complex of solid support bound antibody reagents-target polypeptide. This complex is immobilized on the plate and is not washed away in the wash step. Subsequent incubation with TMB will produce much color change. Such a competitive ELSA test is specific, sensitive, reproducible and easy to operate.

**[0053]** There are other different forms of ELISA, which are well known to those skilled in the art. The standard techniques known in the art for ELISA are described in "Methods in Immunodiagnosis", 2nd Edition, Rose and Bigazzi, eds. John Wiley & Sons, 1980; and Oellerich, M. 1984, *J. Clin. Chem. Clin. Biochem.* 22:895-904. These references are hereby incorporated by reference in their entirety.

**[0054]** In one embodiment, the levels of a polypeptide in a sample can be detected by a lateral flow immunoassay test (LFIA), also known as the immunochromatographic assay, or strip test. LFIA are a simple device intended to detect the presence (or absence) of antigen, e.g. a polypeptide, in a fluid sample. There are currently many LFIA tests are used for medical diagnostics either for home testing, point of care testing, or laboratory use. LFIA tests are a form of immunoassay in which the test sample flows along a solid substrate via capillary action. After the sample is applied to the test strip it encounters a colored reagent (generally comprising antibody specific for the test target antigen) bound to microparticles which mixes with the sample and transits the substrate encountering lines or zones which have been pretreated with another antibody or antigen. Depending upon the level of target polypeptides present in the sample the colored reagent can be captured and become bound at the test line or zone. LFIA are essentially immunoassays adapted to operate along a single axis to suit the test strip format or a dipstick format. Strip tests are extremely versatile and can be easily modified by one skilled in the art for detecting an enormous range of antigens from fluid samples such as urine, blood, water, and/or homogenized tumor samples etc. Strip tests are also known as dip stick test, the name bearing from the literal action of "dipping" the test strip into a fluid sample to be tested. LFIA strip tests are easy to use, require minimum training and can easily be included as components of point-of-care test (POCT) diagnostics to be use on site in the field. LFIA tests can be operated as either competitive or sandwich assays. Sandwich LFIA are similar to sandwich ELISA. The sample first encounters colored particles which are labeled with antibodies raised to the target antigen. The test line will also contain antibodies to the same target, although it may bind to a different epitope on the antigen. The test line will show as a colored band in positive samples. In some embodiments, the lateral flow immunoassay can be a double antibody sandwich assay, a competitive assay, a quantitative assay or variations thereof. Competitive LFIA are similar to competitive ELISA. The sample first encounters colored particles which are labeled with the target antigen or an analogue. The test line contains antibodies to the target/its analogue. Unlabelled antigen in the sample will block the binding sites on the antibodies preventing uptake of the colored particles. The test line will show as a colored band in negative samples. There are a number of variations on lateral flow technology. It is also possible to apply multiple capture zones to create a multiplex test.

**[0055]** The use of "dip sticks" or LFIA test strips and other solid supports have been described in the art in the context of an immunoassay for a number of antigen biomarkers. U.S. Pat. Nos. 4,943,522; 6,485,982; 6,187,598; 5,770,460; 5,622,871; 6,565,808, U. S. patent applications Ser. No.

10/278,676; U.S. Ser. No. 09/579,673 and U.S. Ser. No. 10/717,082, which are incorporated herein by reference in their entirety, are non-limiting examples of such lateral flow test devices. Examples of patents that describe the use of "dip stick" technology to detect soluble antigens via immunochemical assays include, but are not limited to US Patent Nos. 4,444,880; 4,305,924; and 4,135,884; which are incorporated by reference herein in their entireties. The apparatuses and methods of these three patents broadly describe a first component fixed to a solid surface on a "dip stick" which is exposed to a solution containing a soluble antigen that binds to the component fixed upon the "dip stick," prior to detection of the component-antigen complex upon the stick. It is within the skill of one in the art to modify the teachings of this "dip stick" technology for the detection of polypeptides using antibody reagents as described herein.

**[0056]** Other techniques can be used to detect the level of a polypeptide in a sample. One such technique is the dot blot, an adaptation of Western blotting (Towbin et al., Proc. Nat. Acad. Sci. 76:4350 (1979)). In a Western blot, the polypeptide or fragment thereof can be dissociated with detergents and heat, and separated on an SDS-PAGE gel before being transferred to a solid support, such as a nitrocellulose or PVDF membrane. The membrane is incubated with an antibody reagent specific for the target polypeptide or a fragment thereof. The membrane is then washed to remove unbound proteins and proteins with non-specific binding. Detectably labeled enzyme-linked secondary or detection antibodies can then be used to detect and assess the amount of polypeptide in the sample tested. The intensity of the signal from the detectable label corresponds to the amount of enzyme present, and therefore the amount of polypeptide. Levels can be quantified, for example by densitometry.

**[0057]** Flow cytometry is a well-known technique for analyzing and sorting cells (or other small particles) suspended in a fluid stream. This technique allows simultaneous analysis of the physical and/or chemical characteristics of single cells flowing through an optical, electronic, or magnetic detection apparatus. As applied to FACS, the flow cytometer consists of a flow cell which carries the cells in a fluid stream in single file through a light source with excites the fluorescently labeled detection marker(s) (for example, antibody reagents) and measures the fluorescent character of the cell. The fluid stream is then ejected through a nozzle and a charging ring, under pressure, which breaks the fluid into droplets. The flow cell device and fluid stream is calibrated such that there is a relatively large distance between individual cells or bound groups of cells, resulting in a low probability that any droplet contains more than a single cell or bound group of cells. The charging ring charges the droplets based on the fluorescence characteristic of the cell which is contained therein. The charged droplets are then deflected by an electrostatically-charged deflection system which diverts the droplets into various containers based upon their charge (related to the fluorescence intensity of the cell). A FACS system (e.g. the FACSARIA™ flow cytometer (BD Biosciences) and FLOWJO™ Version 7.6.4 (TreeStar)) can detect and record the number of total cells as well as the

number of cells which display one or more fluorescent characteristics, e.g. the total number of cells bound by one or more antibody reagents specific for a CTC marker gene.

**[0058]** In certain embodiments, the gene expression products as described herein can be instead determined by determining the level of messenger RNA (mRNA) expression of genes associated with the marker genes described herein. Such molecules can be isolated, derived, or amplified from a biological sample, such as a tumor biopsy. Detection of mRNA expression is known by persons skilled in the art, and comprise, for example but not limited to, PCR procedures, RT-PCR, quantitative PCR or RT-PCR, Northern blot analysis, differential gene expression, RNA protection assay, microarray analysis, hybridization methods, next-generation sequencing etc. Non-limiting examples of next-generation sequencing technologies can include Ion Torrent, Illumina, SOLiD, 454; Massively Parallel Signature Sequencing solid-phase, reversible dye-terminator sequencing; and DNA nanoball sequencing.

**[0059]** In general, the PCR procedure describes a method of gene amplification which is comprised of (i) sequence-specific hybridization of primers to specific genes or sequences within a nucleic acid sample or library, (ii) subsequent amplification involving multiple rounds of annealing, elongation, and denaturation using a thermostable DNA polymerase, and (iii) screening the PCR products for a band of the correct size. The primers used are oligonucleotides of sufficient length and appropriate sequence to provide initiation of polymerization, i.e. each primer is specifically designed to be complementary to a strand of the genomic locus to be amplified. In an alternative embodiment, mRNA level of gene expression products described herein can be determined by reverse-transcription (RT) PCR and by quantitative RT-PCR (QRT-PCR) or real-time PCR methods. Methods of RT-PCR and QRT-PCR are well known in the art. The nucleic acid sequences of the marker genes described herein have been assigned NCBI accession numbers for different species such as human, mouse and rat. Accordingly, a skilled artisan can design an appropriate primer based on the known sequence for determining the mRNA level of the respective gene.

**[0060]** Nucleic acid and ribonucleic acid (RNA) molecules can be isolated from a particular biological sample using any of a number of procedures, which are well-known in the art, the particular isolation procedure chosen being appropriate for the particular biological sample. For example, freeze-thaw and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from solid materials; heat and alkaline lysis procedures can be useful for obtaining nucleic acid molecules from urine; and proteinase K extraction can be used to obtain nucleic acid from blood (Roiff, A et al. PCR: Clinical Diagnostics and Research, Springer (1994)).

**[0061]** In general, the PCR procedure describes a method of gene amplification which is comprised of (i) sequence-specific hybridization of primers to specific genes within a nucleic acid sample or library, (ii) subsequent amplification involving multiple rounds of annealing, elongation, and denaturation using a DNA polymerase, and (iii) screening the PCR products for a band of the

correct size. The primers used are oligonucleotides of sufficient length and appropriate sequence to provide initiation of polymerization, i.e. each primer is specifically designed to be complementary to each strand of the nucleic acid molecule to be amplified.

**[0062]** In an alternative embodiment, mRNA level of gene expression products described herein can be determined by reverse-transcription (RT) PCR and by quantitative RT-PCR (QRT-PCR) or real-time PCR methods. Methods of RT-PCR and QRT-PCR are well known in the art.

**[0063]** In some embodiments, one or more of the reagents (e.g. an antibody reagent and/or nucleic acid probe) described herein can comprise a detectable label and/or comprise the ability to generate a detectable signal (e.g. by catalyzing reaction converting a compound to a detectable product). Detectable labels can comprise, for example, a light-absorbing dye, a fluorescent dye, or a radioactive label. Detectable labels, methods of detecting them, and methods of incorporating them into reagents (e.g. antibodies and nucleic acid probes) are well known in the art.

**[0064]** In some embodiments, detectable labels can include labels that can be detected by spectroscopic, photochemical, biochemical, immunochemical, electromagnetic, radiochemical, or chemical means, such as fluorescence, chemiluminescence, or chemiluminescence, or any other appropriate means. The detectable labels used in the methods described herein can be primary labels (where the label comprises a moiety that is directly detectable or that produces a directly detectable moiety) or secondary labels (where the detectable label binds to another moiety to produce a detectable signal, e.g., as is common in immunological labeling using secondary and tertiary antibodies). The detectable label can be linked by covalent or non-covalent means to the reagent. Alternatively, a detectable label can be linked such as by directly labeling a molecule that achieves binding to the reagent via a ligand-receptor binding pair arrangement or other such specific recognition molecules. Detectable labels can include, but are not limited to radioisotopes, bioluminescent compounds, chromophores, antibodies, chemiluminescent compounds, fluorescent compounds, metal chelates, and enzymes.

**[0065]** In other embodiments, the detection reagent is label with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wavelength, its presence can then be detected due to fluorescence. In some embodiments, a detectable label can be a fluorescent dye molecule, or fluorophore including, but not limited to fluorescein, phycoerythrin, phycocyanin, o-phthaldehyde, fluorescamine, Cy3<sup>TM</sup>, Cy5<sup>TM</sup>, allophycocyanine, Texas Red, peridinin chlorophyll, cyanine, tandem conjugates such as phycoerythrin-Cy5<sup>TM</sup>, green fluorescent protein, rhodamine, fluorescein isothiocyanate (FITC) and Oregon Green<sup>TM</sup>, rhodamine and derivatives (e.g., Texas red and tetrarhodamine isothiocyanate (TRITC)), biotin, phycoerythrin, AMCA, CyDyes<sup>TM</sup>, 6-carboxyfluorescein (commonly known by the abbreviations FAM and F), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 6-carboxy-4',5'-dichloro-2',7'-dimethoxyfluorescein (JOE or J), N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA or T), 6-carboxy-X-rhodamine (ROX or R), 5-

carboxyrhodamine-6G (R6G5 or G5), 6-carboxyrhodamine-6G (R6G6 or G6), and rhodamine 110; cyanine dyes, e.g. Cy3, Cy5 and Cy7 dyes; coumarins, e.g. umbelliferone; benzimide dyes, e.g. Hoechst 33258; phenanthridine dyes, e.g. Texas Red; ethidium dyes; acridine dyes; carbazole dyes; phenoxazine dyes; porphyrin dyes; polymethine dyes, e.g. cyanine dyes such as Cy3, Cy5, etc; BODIPY dyes and quinoline dyes. In some embodiments, a detectable label can be a radiolabel including, but not limited to  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ , and  $^{33}\text{P}$ . In some embodiments, a detectable label can be an enzyme including, but not limited to horseradish peroxidase and alkaline phosphatase. An enzymatic label can produce, for example, a chemiluminescent signal, a color signal, or a fluorescent signal. Enzymes contemplated for use to detectably label an antibody reagent include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-V-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-VI-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. In some embodiments, a detectable label is a chemiluminescent label, including, but not limited to lucigenin, luminol, luciferin, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester. In some embodiments, a detectable label can be a spectral colorimetric label including, but not limited to colloidal gold or colored glass or plastic (e.g., polystyrene, polypropylene, and latex) beads.

**[0066]** In some embodiments, detection reagents can also be labeled with a detectable tag, such as c-Myc, HA, VSV-G, HSV, FLAG, V5, HIS, or biotin. Other detection systems can also be used, for example, a biotin-streptavidin system. In this system, the antibodies immunoreactive (i. e. specific for) with the biomarker of interest is biotinylated. Quantity of biotinylated antibody bound to the biomarker is determined using a streptavidin-peroxidase conjugate and a chromogenic substrate. Such streptavidin peroxidase detection kits are commercially available, e. g. from DAKO; Carpinteria, CA. A reagent can also be detectably labeled using fluorescence emitting metals such as  $^{152}\text{Eu}$ , or others of the lanthanide series. These metals can be attached to the reagent using such metal chelating groups as diethylenetriaminepentaacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

**[0067]** In some embodiments of any of the aspects described herein, the level of expression products of more than one gene can be determined simultaneously (e.g. a multiplex assay) or in parallel. In some embodiments, the level of expression products of no more than 200 other genes is determined. In some embodiments, the level of expression products of no more than 100 other genes is determined. In some embodiments, the level of expression products of no more than 20 other genes is determined. In some embodiments, the level of expression products of no more than 10 other genes is determined.

**[0068]** The term “sample” or “test sample” as used herein denotes a sample taken or isolated from a biological organism, e.g., a tumor sample from a subject. Exemplary biological samples

include, but are not limited to, a biofluid sample; serum; plasma; urine; saliva; a tumor sample; a tumor biopsy and/or tissue sample etc. The term also includes a mixture of the above-mentioned samples. The term “test sample” also includes untreated or pretreated (or pre-processed) biological samples. In some embodiments, a test sample can comprise cells from subject. In some embodiments, a test sample can be a tumor cell test sample, e.g. the sample can comprise cancerous cells, cells from a tumor, and/or a tumor biopsy. In some embodiments, the test sample can be a blood sample.

**[0069]** The test sample can be obtained by removing a sample of cells from a subject, but can also be accomplished by using previously isolated cells (e.g. isolated at a prior timepoint and isolated by the same or another person). In addition, the test sample can be freshly collected or a previously collected sample.

**[0070]** In some embodiments, the test sample can be an untreated test sample. As used herein, the phrase “untreated test sample” refers to a test sample that has not had any prior sample pre-treatment except for dilution and/or suspension in a solution. Exemplary methods for treating a test sample include, but are not limited to, centrifugation, filtration, sonication, homogenization, heating, freezing and thawing, and combinations thereof. In some embodiments, the test sample can be a frozen test sample, e.g., a frozen tissue. The frozen sample can be thawed before employing methods, assays and systems described herein. After thawing, a frozen sample can be centrifuged before being subjected to methods, assays and systems described herein. In some embodiments, the test sample is a clarified test sample, for example, by centrifugation and collection of a supernatant comprising the clarified test sample. In some embodiments, a test sample can be a pre-processed test sample, for example, supernatant or filtrate resulting from a treatment selected from the group consisting of centrifugation, filtration, thawing, purification, and any combinations thereof. In some embodiments, the test sample can be treated with a chemical and/or biological reagent. Chemical and/or biological reagents can be employed to protect and/or maintain the stability of the sample, including biomolecules (e.g., nucleic acid and protein) therein, during processing. One exemplary reagent is a protease inhibitor, which is generally used to protect or maintain the stability of protein during processing. The skilled artisan is well aware of methods and processes appropriate for pre-processing of biological samples required for determination of the level of an expression product as described herein.

**[0071]** In some embodiments, the methods, assays, and systems described herein can further comprise a step of obtaining a test sample from a subject. In some embodiments, the subject can be a human subject.

**[0072]** In some embodiments, the methods and assays described herein can further comprise a step of isolating CTCs or potential CTCs from a sample prior to measuring the level the expression product of one or more of the marker genes described herein. By way of non-limitng example, CTCs

can be isolated from, e.g. a blood sample by hydrodynamic size-based separation and/or immunodepletion of other cell types present in blood samples. The CTC-iChip, described in the Examples herein combines these two approaches to isolate CTCs.

**[0073]** Subjects with high, or at least detectable, levels of CTCs are most likely to benefit from treatment with therapies that specifically target CTCs. Accordingly, provided herein is a method of determining if a subject is likely to respond to treatment with a CTC marker gene-targeted therapy, the method comprising: measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and determining that the subject is likely to respond to the treatment if the level of the expression product is increased relative to a reference level. CTC marker gene-targeted therapies are discussed below herein.

**[0074]** Decreased levels of CTCs after administration of a therapy can be indicative of an improvement in the condition of the subject, e.g. the cancer is reduced in size, growth, and/or metastatic potential. Accordingly, provided herein is a method of monitoring the treatment of a subject, the method comprising administering a cancer therapy to a subject in need thereof; measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and determining that the subject is responding if the level of the CTC marker gene expression product is decreased relative to the reference level and determining that the subject is not responding to the treatment if the CTC marker gene expression product is not decreased relative to the reference level. In some embodiments the therapy is a chemotherapy, surgical therapy, and/or radiation therapy. In some embodiments, the therapy is a CTC marker gene-targeted therapy. In some embodiments, the reference level is the level of the gene expression product in the patient prior to the administering step.

**[0075]** The CTC marker genes described herein can be targeted directly and/or used to physically target a chemotherapeutic agent to reduce the levels and/or pathogenic activity of CTCs (e.g. metastatic activity). Accordingly, described herein is a method of treating cancer in a subject, the method comprising administering a therapeutically effective amount of a CTC marker gene-targeted therapy to the subject. In some embodiments, the subject is a subject determined to have an elevated level of CTCs and/or an elevated level of a CTC marker gene present in the blood and/or stroma of the cancer.

**[0076]** In some embodiments, the CTC marker gene-targeted therapy can comprise an inhibitor of a CTC marker gene, e.g. the CTC marker gene-targeted therapy can inhibit the level and/or activity of a CTC marker gene. As used herein, the term “inhibitor” refers to an agent which can decrease the expression and/or activity of the targeted expression product (e.g. mRNA encoding the target or a target polypeptide), e.g. by at least 10% or more, e.g. by 10% or more, 50% or more, 70% or more, 80% or more, 90% or more, 95% or more, or 98 % or more. The efficacy of an inhibitor of a CTC marker gene, e.g. its ability to decrease the level and/or activity of the CTC marker gene can be

determined, e.g. by measuring the level of an expression product and/or the activity of the CTC marker gene. Methods for measuring the level of a given mRNA and/or polypeptide are known to one of skill in the art, e.g. RTPCR with primers can be used to determine the level of RNA and Western blotting with an antibody can be used to determine the level of a polypeptide. The activity of, e.g. a CTC marker gene can be determined, e.g. by measuring the levels and/or survival of CTCs using methods known in the art and described elsewhere herein. In some embodiments, the inhibitor of a CTC marker gene can be an inhibitory nucleic acid; an aptamer; an antibody reagent; an antibody; or a small molecule.

**[0077]** In some embodiments, the inhibitor of a CTC marker gene can be an antibody reagent. As used herein an "antibody" refers to IgG, IgM, IgA, IgD or IgE molecules or antigen-specific antibody fragments thereof (including, but not limited to, a Fab, F(ab')<sub>2</sub>, Fv, disulphide linked Fv, scFv, single domain antibody, closed conformation multispecific antibody, disulphide-linked scfv, diabody), whether derived from any species that naturally produces an antibody, or created by recombinant DNA technology; whether isolated from serum, B-cells, hybridomas, transfectomas, yeast or bacteria.

**[0078]** As described herein, an "antigen" is a molecule that is bound by a binding site on an antibody agent. Typically, antigens are bound by antibody ligands and are capable of raising an antibody response *in vivo*. An antigen can be a polypeptide, protein, nucleic acid or other molecule or portion thereof. The term "antigenic determinant" refers to an epitope on the antigen recognized by an antigen-binding molecule, and more particularly, by the antigen-binding site of said molecule.

**[0079]** As used herein, the term "antibody reagent" refers to a polypeptide that includes at least one immunoglobulin variable domain or immunoglobulin variable domain sequence and which specifically binds a given antigen. An antibody reagent can comprise an antibody or a polypeptide comprising an antigen-binding domain of an antibody. In some embodiments, an antibody reagent can comprise a monoclonal antibody or a polypeptide comprising an antigen-binding domain of a monoclonal antibody. For example, an antibody can include a heavy (H) chain variable region (abbreviated herein as VH), and a light (L) chain variable region (abbreviated herein as VL). In another example, an antibody includes two heavy (H) chain variable regions and two light (L) chain variable regions. The term "antibody reagent" encompasses antigen-binding fragments of antibodies (e.g., single chain antibodies, Fab and sFab fragments, F(ab')<sub>2</sub>, Fd fragments, Fv fragments, scFv, and domain antibodies (dAb) fragments (see, e.g. de Wildt et al., Eur J. Immunol. 1996; 26(3):629-39; which is incorporated by reference herein in its entirety)) as well as complete antibodies. An antibody can have the structural features of IgA, IgG, IgE, IgD, IgM (as well as subtypes and combinations thereof). Antibodies can be from any source, including mouse, rabbit, pig, rat, and primate (human and non-human primate) and primatized antibodies. Antibodies also include midibodies, humanized antibodies, chimeric antibodies, and the like.

**[0080]** The VH and VL regions can be further subdivided into regions of hypervariability,

termed "complementarity determining regions" ("CDR"), interspersed with regions that are more conserved, termed "framework regions" ("FR"). The extent of the framework region and CDRs has been precisely defined (see, Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242, and Chothia, C. et al. (1987) J. Mol. Biol. 196:901-917; which are incorporated by reference herein in their entireties). Each VH and VL is typically composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

**[0081]** The terms "antigen-binding fragment" or "antigen-binding domain", which are used interchangeably herein are used to refer to one or more fragments of a full length antibody that retain the ability to specifically bind to a target of interest. Examples of binding fragments encompassed within the term "antigen-binding fragment" of a full length antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')2 fragment, a bivalent fragment including two Fab fragments linked by a disulfide bridge at the hinge region; (iii) an Fd fragment consisting of the VH and CH1 domains; (iv) an Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) Nature 341:544-546; which is incorporated by reference herein in its entirety), which consists of a VH or VL domain; and (vi) an isolated complementarity determining region (CDR) that retains specific antigen-binding functionality. As used herein, the term "specific binding" refers to a chemical interaction between two molecules, compounds, cells and/or particles wherein the first entity binds to the second, target entity with greater specificity and affinity than it binds to a third entity which is a non-target. In some embodiments, specific binding can refer to an affinity of the first entity for the second target entity which is at least 10 times, at least 50 times, at least 100 times, at least 500 times, at least 1000 times or greater than the affinity for the third nontarget entity.

**[0082]** Additionally, and as described herein, a recombinant humanized antibody can be further optimized to decrease potential immunogenicity, while maintaining functional activity, for therapy in humans. In this regard, functional activity means a polypeptide capable of displaying one or more known functional activities associated with a recombinant antibody or antibody reagent thereof as described herein. Such functional activities include, e.g. the ability to bind to a given CTC marker gene.

**[0083]** In some embodiments, the inhibitor of a CTC marker gene can be an inhibitory nucleic acid reagent. In some embodiments, the inhibitory nucleic acid is an inhibitory RNA (iRNA). Double-stranded RNA molecules (dsRNA) have been shown to block gene expression in a highly conserved regulatory mechanism known as RNA interference (RNAi). The inhibitory nucleic acids described herein can include an RNA strand (the antisense strand) having a region which is 30 nucleotides or less in length, i.e., 15-30 nucleotides in length, generally 19-24 nucleotides in length,

which region is substantially complementary to at least part the targeted mRNA transcript. The use of these iRNAs enables the targeted degradation of mRNA transcripts, resulting in decreased expression and/or activity of the target.

**[0084]** As used herein, the term “iRNA” refers to an agent that contains RNA as that term is defined herein, and which mediates the targeted cleavage of an RNA transcript via an RNA-induced silencing complex (RISC) pathway. In one embodiment, an iRNA as described herein effects inhibition of the expression and/or activity of the target mRNA. In certain embodiments, contacting a cell with the inhibitor (e.g. an iRNA) results in a decrease in the target mRNA level in a cell by at least about 5%, about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, about 99%, up to and including 100% of the target mRNA level found in the cell without the presence of the iRNA.

**[0085]** In some embodiments, the iRNA can be a dsRNA. A dsRNA includes two RNA strands that are sufficiently complementary to hybridize to form a duplex structure under conditions in which the dsRNA will be used. One strand of a dsRNA (the antisense strand) includes a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence. The target sequence can be derived from the sequence of an mRNA formed during the expression of the target. The other strand (the sense strand) includes a region that is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. Generally, the duplex structure is between 15 and 30 inclusive, more generally between 18 and 25 inclusive, yet more generally between 19 and 24 inclusive, and most generally between 19 and 21 base pairs in length, inclusive. Similarly, the region of complementarity to the target sequence is between 15 and 30 inclusive, more generally between 18 and 25 inclusive, yet more generally between 19 and 24 inclusive, and most generally between 19 and 21 nucleotides in length, inclusive. In some embodiments, the dsRNA is between 15 and 20 nucleotides in length, inclusive, and in other embodiments, the dsRNA is between 25 and 30 nucleotides in length, inclusive. As the ordinarily skilled person will recognize, the targeted region of an RNA targeted for cleavage will most often be part of a larger RNA molecule, often an mRNA molecule. Where relevant, a “part” of an mRNA target is a contiguous sequence of an mRNA target of sufficient length to be a substrate for RNAi-directed cleavage (i.e., cleavage through a RISC pathway). dsRNAs having duplexes as short as 9 base pairs can, under some circumstances, mediate RNAi-directed RNA cleavage. Most often a target will be at least 15 nucleotides in length, preferably 15-30 nucleotides in length.

**[0086]** In yet another embodiment, the RNA of an iRNA, e.g., a dsRNA, is chemically modified to enhance stability or other beneficial characteristics. The nucleic acids featured in the invention may be synthesized and/or modified by methods well established in the art, such as those described in “Current protocols in nucleic acid chemistry,” Beaucage, S.L. *et al.* (Edrs.), John Wiley & Sons, Inc.,

New York, NY, USA, which is hereby incorporated herein by reference. Modifications include, for example, (a) end modifications, e.g., 5' end modifications (phosphorylation, conjugation, inverted linkages, etc.) 3' end modifications (conjugation, DNA nucleotides, inverted linkages, etc.), (b) base modifications, e.g., replacement with stabilizing bases, destabilizing bases, or bases that base pair with an expanded repertoire of partners, removal of bases (abasic nucleotides), or conjugated bases, (c) sugar modifications (e.g., at the 2' position or 4' position) or replacement of the sugar, as well as (d) backbone modifications, including modification or replacement of the phosphodiester linkages. Specific examples of RNA compounds useful in the embodiments described herein include, but are not limited to RNAs containing modified backbones or no natural internucleoside linkages. RNAs having modified backbones include, among others, those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified RNAs that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides. In particular embodiments, the modified RNA will have a phosphorus atom in its internucleoside backbone.

**[0087]** Modified RNA backbones can include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those) having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included. Representative U.S. patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Pat. Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; 6,028,188; 6,124,445; 6,160,109; 6,169,170; 6,172,209; 6,239,265; 6,277,603; 6,326,199; 6,346,614; 6,444,423; 6,531,590; 6,534,639; 6,608,035; 6,683,167; 6,835,826; 6,858,715; 6,867,289; 6,867,294; 6,878,805; 7,015,315; 7,041,816; 7,273,933; 7,321,029; 7,834,171; 7,919,612; 7,960,360; 7,989,603; 8,309,707; 6,524,681; and US Pat RE39464, each of which is herein incorporated by reference

**[0088]** Modified RNA backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones;

alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts. Representative U.S. patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Pat. Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and, 5,677,439, each of which is herein incorporated by reference.

**[0089]** In other RNA mimetics suitable or contemplated for use in iRNAs, both the sugar and the internucleoside linkage, *i.e.*, the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an RNA mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar backbone of an RNA is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found, for example, in Nielsen *et al.*, *Science*, 1991, 254, 1497-1500.

**[0090]** Some embodiments featured in the invention include RNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular --CH<sub>2</sub>--NH--CH<sub>2</sub>--, --CH<sub>2</sub>--N(CH<sub>3</sub>)--O--CH<sub>2</sub>--[known as a methylene (methylimino) or MMI backbone], --CH<sub>2</sub>--O--N(CH<sub>3</sub>)--CH<sub>2</sub>--, --CH<sub>2</sub>--N(CH<sub>3</sub>)--N(CH<sub>3</sub>)--CH<sub>2</sub>-- and --N(CH<sub>3</sub>)--CH<sub>2</sub>--CH<sub>2</sub>--[wherein the native phosphodiester backbone is represented as --O--P--O--CH<sub>2</sub>--] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. In some embodiments, the RNAs featured herein have morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

**[0091]** Modified RNAs can also contain one or more substituted sugar moieties. The iRNAs, e.g., dsRNAs, featured herein can include one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Exemplary suitable modifications include O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. In other embodiments, dsRNAs include one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an

intercalator, a group for improving the pharmacokinetic properties of an iRNA, or a group for improving the pharmacodynamic properties of an iRNA, and other substituents having similar properties. In some embodiments, the modification includes a 2'-methoxyethoxy (2'-O--CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78:486-504) *i.e.*, an alkoxy-alkoxy group. Another exemplary modification is 2'-dimethylaminoxyethoxy, *i.e.*, a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), *i.e.*, 2'-O--CH<sub>2</sub>--O--CH<sub>2</sub>--N(CH<sub>2</sub>)<sub>2</sub>, also described in examples herein below.

**[0092]** Other modifications include 2'-methoxy (2'-OCH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications can also be made at other positions on the RNA of an iRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. iRNAs may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,700,920; 8,084,600; 8,124,745; 8,377,644 each of which is herein incorporated by reference.

**[0093]** An iRNA can also include nucleobase (often referred to in the art simply as “base”) modifications or substitutions. As used herein, “unmodified” or “natural” nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl anal other 8-substituted adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in Modified Nucleosides in Biochemistry, Biotechnology and Medicine, Herdewijn, P. ed. Wiley-VCH, 2008; those disclosed in The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. L, ed. John Wiley & Sons, 1990, these disclosed by Englisch *et al.*, Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y S., Chapter 15, dsRNA Research and Applications, pages 289-302,

Crooke, S. T. and Lebleu, B., Ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds featured in the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and 0-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., dsRNA Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are exemplary base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

**[0094]** Representative U.S. patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Pat. No. 3,687,808, as well as U.S. Pat. Nos. 4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; 5,681,941; 6,015,886; 6,147,200; 6,166,197; 6,222,025; 6,235,887; 6,380,368; 6,528,640; 6,639,062; 6,617,438; 7,045,610; 7,427,672; and 7,495,088, each of which is herein incorporated by reference, and U.S. Pat. No. 5,750,692, also herein incorporated by reference.

**[0095]** The RNA of an iRNA can also be modified to include one or more locked nucleic acids (LNA). A locked nucleic acid is a nucleotide having a modified ribose moiety in which the ribose moiety comprises an extra bridge connecting the 2' and 4' carbons. This structure effectively "locks" the ribose in the 3'-endo structural conformation. The addition of locked nucleic acids to siRNAs has been shown to increase siRNA stability in serum, and to reduce off-target effects (Elmen, J. et al., (2005) *Nucleic Acids Research* 33(1):439-447; Mook, OR. et al., (2007) *Mol Canc Ther* 6(3):833-843; Grunweller, A. et al., (2003) *Nucleic Acids Research* 31(12):3185-3193). Representative U.S. Patents that teach the preparation of locked nucleic acid nucleotides include, but are not limited to, the following: U.S. Pat. Nos. 6,268,490; 6,670,461; 6,794,499; 6,998,484; 7,053,207; 7,084,125; and 7,399,845, each of which is herein incorporated by reference in its entirety.

**[0096]** Another modification of the RNA of an iRNA featured in the invention involves chemically linking to the RNA one or more ligands, moieties or conjugates that enhance the activity, cellular distribution, pharmacokinetic properties, or cellular uptake of the iRNA. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger *et al.*, Proc. Natl. Acid. Sci. USA, 1989, 86: 6553-6556), cholic acid (Manoharan *et al.*, Biorg. Med. Chem. Let., 1994, 4:1053-1060), a thioether, *e.g.*, beryl-S-tritylthiol (Manoharan *et al.*, Ann. N.Y. Acad. Sci., 1992, 660:306-309; Manoharan *et al.*, Biorg. Med. Chem. Let., 1993, 3:2765-2770), a thiocholesterol (Oberhauser *et al.*, Nucl. Acids Res., 1992, 20:533-538), an aliphatic chain, *e.g.*, dodecandiol or undecyl residues (Saison-Behmoaras *et al.*, EMBO J, 1991, 10:1111-1118; Kabanov *et al.*, FEBS Lett., 1990, 259:327-330; Svinarchuk *et al.*, Biochimie, 1993, 75:49-54), a phospholipid, *e.g.*, di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-phosphonate

(Manoharan *et al.*, Tetrahedron Lett., 1995, 36:3651-3654; Shea *et al.*, Nucl. Acids Res., 1990, 18:3777-3783), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, Nucleosides & Nucleotides, 1995, 14:969-973), or adamantan acetic acid (Manoharan *et al.*, Tetrahedron Lett., 1995, 36:3651-3654), a palmityl moiety (Mishra *et al.*, Biochim. Biophys. Acta, 1995, 1264:229-237), or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke *et al.*, J. Pharmacol. Exp. Ther., 1996, 277:923-937).

**[0097]** In some embodiments the CTC marker gene-targeted therapy can comprise an agent that binds to the CTC marker gene expression product and an agent that is chemotherapeutic. In some embodiments, the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent and a chemotherapeutic agent. A CTC marker gene-binding antibody reagent can be an antibody reagent that binds, e.g. a CTC marker gene polypeptide. The binding antibody reagent can be an inhibitor or can exhibit no inhibitory effect on its own. By binding to the CTC marker gene, and thereby a CTC, it concentrates and localizes the chemotherapeutic agent at CTC cells in the circulation and/or stroma of the tumor – increasing efficacy and reducing side effects.

**[0098]** In some embodiments, the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent that binds a marker gene selected from Table 14. In some embodiments, the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent that binds a marker gene selected from the group consisting of: IL6ST, SULF2, and SV2A.

**[0099]** As used herein the term "chemotherapeutic agent" refers to any chemical or biological agent with therapeutic usefulness in the treatment of diseases characterized by abnormal cell growth. Such diseases include tumors, neoplasms and cancer as well as diseases characterized by hyperplastic growth. These agents can function to inhibit a cellular activity upon which the cancer cell depends for continued proliferation. In some aspect of all the embodiments, a chemotherapeutic agent is a cell cycle inhibitor or a cell division inhibitor. Categories of chemotherapeutic agents that are useful in the methods of the invention include alkylating/alkaloid agents, antimetabolites, hormones or hormone analogs, and miscellaneous antineoplastic drugs. Most of these agents are directly or indirectly toxic to cancer cells. In one embodiment, a chemotherapeutic agent is a radioactive molecule. One of skill in the art can readily identify a chemotherapeutic agent of use ( e.g. see Slapak and Kufe, Principles of Cancer Therapy, Chapter 86 in Harrison's Principles of Internal Medicine, 14th edition; Perry *et al.* , Chemotherapy, Ch. 17 in Abeloff, Clinical Oncology 2nd ed. 2000 Churchill Livingstone, Inc; Baltzer L, Berkery R (eds): Oncology Pocket Guide to Chemotherapy, 2nd ed. St. Louis, Mosby-Year Book, 1995; Fischer D S, Knobf M F, Durivage H J (eds): The Cancer Chemotherapy Handbook, 4th ed. St. Louis, Mosby-Year Book, 1993). In some embodiments, the chemotherapeutic agent can be a cytotoxic chemotherapeutic. The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes ( e.g. At211, I131, I125, Y90, Re186,

Re188, Sm153, Bi212, P32 and radioactive isotopes of Lu), chemotherapeutic agents, and toxins, such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

**[00100]** Non-limiting examples of chemotherapeutic agents can include gemcitabine, cisplatin, paclitaxel, carboplatin, bortezomib, AMG479, vorinostat, rituximab, temozolomide, rapamycin, ABT-737, PI-103; alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylololomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; calystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, chlophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (*e.g.*, calicheamicin, especially calicheamicin gammaI and calicheamicin omegaI (see, *e.g.*, Agnew, Chem. Int. Ed. Engl., 33: 183-186 (1994)); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabicin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprime, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as folinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestramycin; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone;

etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; niraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAZANE® Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumberg, Ill.), and TAXOTERE® doxetaxel (Rhone-Poulenc Rorer, Antony, France); chloranbucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin, oxaliplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE.™vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (Camptosar, CPT-11) (including the treatment regimen of irinotecan with 5-FU and leucovorin); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; combretastatin; leucovorin (LV); oxaliplatin, including the oxaliplatin treatment regimen (FOLFOX); lapatinib (Tykerb™.); inhibitors of PKC-alpha, Raf, H-Ras, EGFR (e.g., erlotinib (Tarceva®)) and VEGF-A that reduce cell proliferation and pharmaceutically acceptable salts, acids or derivatives of any of the above.

**[00101]** In some embodiments, the binding antibody reagent and the chemotherapeutic agent can be directly conjugated and/or bound to each other, e.g. an antibody-drug conjugate. In some embodiments, binding can be non-covalent, e.g., by hydrogen, electrostatic, or van der waals interactions, however, binding may also be covalent. By "conjugated" is meant the covalent linkage of at least two molecules. In some embodiments, the composition can be an antibody-drug conjugate.

**[00102]** In some embodiments, the binding antibody reagent can be bound to and/or conjugated to multiple chemotherapeutic molecules. In some embodiments, the ratio of a given chemotherapeutic molecule to the binding antibody reagent molecule can be from about 1:1 to about 1,000:1, e.g. a single antibody binding reagent molecule can be linked to, conjugated to, etc. from about 1 to about 1,000 individual chemotherapeutic molecules.

**[00103]** In some embodiments, the binding antibody reagent and the chemotherapeutic agent can be present in a scaffold material. Scaffold materials suitable for use in therapeutic compositions are known in the art and can include, but are not limited to, a nanoparticle; a matrix; a hydrogel; and a biomaterial, biocompatible, and/or biodegradable scaffold material. As used herein, the term "nanoparticle" refers to particles that are on the order of about  $10^{-9}$  or one billionth of a meter. The

term "nanoparticle" includes nanospheres; nanorods; nanoshells; and nanoprisms; and these nanoparticles may be part of a nanonetwork.

**[00104]** The term "nanoparticles" also encompasses liposomes and lipid particles having the size of a nanoparticle. As used herein, the term "matrix" refers to a 3-dimensional structure comprising the components of a composition described herein (e.g. a binding reagent, kinase inhibitor, and/or EGFR inhibitor). Non-limiting examples of matrix structures include foams; hydrogels; electrospun fibers; gels; fiber mats; sponges; 3-dimensional scaffolds; non-woven mats; woven materials; knit materials; fiber bundles; and fibers and other material formats (See, e.g. Rockwood et al. *Nature Protocols* 2011 6:1612-1631 and US Patent Publications 2011/0167602; 2011/0009960; 2012/0296352; and U.S. Patent No. 8,172,901; each of which is incorporated by reference herein in its entirety). The structure of the matrix can be selected by one of skill in the art depending upon the intended application of the composition, e.g. electrospun matrices can have greater surface area than foams.

**[00105]** In some embodiments, the scaffold is a hydrogel. As used herein, the term "hydrogel" refers to a three-dimensional polymeric structure that is insoluble in water but which is capable of absorbing and retaining large quantities of water to form a stable, often soft and pliable, structure. In some embodiments, water can penetrate in between the polymer chains of the polymer network, subsequently causing swelling and the formation of a hydrogel. In general, hydrogels are superabsorbent. Hydrogels have many desirable properties for biomedical applications. For example, they can be made nontoxic and compatible with tissue, and they are highly permeable to water, ions, and small molecules. Hydrogels are super-absorbent (they can contain over 99% water) and can be comprised of natural (e.g., silk) or synthetic polymers, e.g., PEG.

**[00106]** As used herein, "biomaterial" refers to a material that is biocompatible and biodegradable. As used herein, the term "biocompatible" refers to substances that are not toxic to cells. In some embodiments, a substance is considered to be "biocompatible" if its addition to cells in vitro results in less than or equal to approximately 20% cell death. In some embodiments, a substance is considered to be "biocompatible" if its addition to cells in vivo does not induce inflammation and/or other adverse effects in vivo. As used herein, the term "biodegradable" refers to substances that are degraded under physiological conditions. In some embodiments, a biodegradable substance is a substance that is broken down by cellular machinery. In some embodiments, a biodegradable substance is a substance that is broken down by chemical processes.

**[00107]** In some embodiments, the methods described herein relate to treating a subject having or diagnosed as having cancer with a CTC marker-gene targeted therapy. In some embodiments, the cancer can be pancreatic cancer. Subjects having cancer can be identified by a physician using current methods of diagnosing cancer. Symptoms and/or complications of cancer, e.g. pancreatic cancer, which characterize these conditions and aid in diagnosis are well known in the art and include but are not limited to, pain in the upper abdomen, heartburn, nausea, vomiting, diarrhea, cachexia,

jaundice, pulmonary embolism, Trouseau syndrome, and diabetes mellitus. Tests that may aid in a diagnosis of, e.g. pancreatic cancer include, but are not limited to, liver function tests, CA19-9 tests, CT and endoscopic ultrasound. A family history of pancreatic cancer or exposure to risk factors for pancreatic cancer (e.g. smoking or drinking) can also aid in determining if a subject is likely to have cancer or in making a diagnosis of cancer.

**[00108]** The compositions and methods described herein can be administered to a subject having or diagnosed as having cancer, e.g. pancreatic cancer. In some embodiments, the methods described herein comprise administering an effective amount of compositions described herein, e.g. a CTC marker-gene targeted therapy to a subject in order to alleviate a symptom of a cancer. As used herein, "alleviating a symptom of a cancer" is ameliorating any condition or symptom associated with the cancer. As compared with an equivalent untreated control, such reduction is by at least 5%, 10%, 20%, 40%, 50%, 60%, 80%, 90%, 95%, 99% or more as measured by any standard technique. A variety of means for administering the compositions described herein to subjects are known to those of skill in the art. Such methods can include, but are not limited to oral, parenteral, intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), pulmonary, cutaneous, topical, injection, or intratumoral administration. Administration can be local or systemic.

**[00109]** The term "effective amount" as used herein refers to the amount of a CTC marker-gene targeted therapy needed to alleviate at least one or more symptom of the disease or disorder, and relates to a sufficient amount of pharmacological composition to provide the desired effect. The term "therapeutically effective amount" therefore refers to an amount of CTC marker-gene targeted therapy that is sufficient to provide a particular anti-cancer effect when administered to a typical subject. An effective amount as used herein, in various contexts, would also include an amount sufficient to delay the development of a symptom of the disease, alter the course of a symptom disease (for example but not limited to, slowing the progression of a symptom of the disease), or reverse a symptom of the disease. Thus, it is not generally practicable to specify an exact "effective amount". However, for any given case, an appropriate "effective amount" can be determined by one of ordinary skill in the art using only routine experimentation.

**[00110]** Effective amounts, toxicity, and therapeutic efficacy can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dosage can vary depending upon the dosage form employed and the route of administration utilized. The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio LD50/ED50. Compositions and methods that exhibit large therapeutic indices are preferred. A therapeutically effective dose can be estimated initially from cell culture assays. Also, a dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of a CTC marker-gene targeted

therapy, which achieves a half-maximal inhibition of symptoms) as determined in cell culture, or in an appropriate animal model. Levels in plasma can be measured, for example, by high performance liquid chromatography. The effects of any particular dosage can be monitored by a suitable bioassay, e.g., assay for CTC levels, among others. The dosage can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

**[00111]** In some embodiments, the technology described herein relates to a pharmaceutical composition comprising a CTC marker-gene targeted therapy as described herein, and optionally a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers and diluents include saline, aqueous buffer solutions, solvents and/or dispersion media. The use of such carriers and diluents is well known in the art. Some non-limiting examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, methylcellulose, ethyl cellulose, microcrystalline cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) lubricating agents, such as magnesium stearate, sodium lauryl sulfate and talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol (PEG); (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; (22) bulking agents, such as polypeptides and amino acids (23) serum component, such as serum albumin, HDL and LDL; (22) C<sub>2</sub>-C<sub>12</sub> alcohols, such as ethanol; and (23) other non-toxic compatible substances employed in pharmaceutical formulations. Wetting agents, coloring agents, release agents, coating agents, sweetening agents, flavoring agents, perfuming agents, preservative and antioxidants can also be present in the formulation. The terms such as "excipient", "carrier", "pharmaceutically acceptable carrier" or the like are used interchangeably herein. In some embodiments, the carrier inhibits the degradation of the active agent, e.g. a CTC marker-gene targeted therapy as described herein.

**[00112]** In some embodiments, the pharmaceutical composition comprising a CTC marker-gene targeted therapy as described herein can be a parenteral dose form. Since administration of parenteral dosage forms typically bypasses the patient's natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions. In addition, controlled-release parenteral dosage

forms can be prepared for administration of a patient, including, but not limited to, DUROS®-type dosage forms and dose-dumping.

**[00113]** Suitable vehicles that can be used to provide parenteral dosage forms of a CTC marker-gene targeted therapy as disclosed within are well known to those skilled in the art. Examples include, without limitation: sterile water; water for injection USP; saline solution; glucose solution; aqueous vehicles such as but not limited to, sodium chloride injection, Ringer's injection, dextrose Injection, dextrose and sodium chloride injection, and lactated Ringer's injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and propylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate. Compounds that alter or modify the solubility of a pharmaceutically acceptable salt of a CTC marker-gene targeted therapy as disclosed herein can also be incorporated into the parenteral dosage forms of the disclosure, including conventional and controlled-release parenteral dosage forms.

**[00114]** Pharmaceutical compositions comprising a CTC marker-gene targeted therapy can also be formulated to be suitable for oral administration, for example as discrete dosage forms, such as, but not limited to, tablets (including without limitation scored or coated tablets), pills, caplets, capsules, chewable tablets, powder packets, cachets, troches, wafers, aerosol sprays, or liquids, such as but not limited to, syrups, elixirs, solutions or suspensions in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion, or a water-in-oil emulsion. Such compositions contain a predetermined amount of the pharmaceutically acceptable salt of the disclosed compounds, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, Remington: The Science and Practice of Pharmacy, 21st Ed., Lippincott, Williams, and Wilkins, Philadelphia PA. (2005).

**[00115]** Conventional dosage forms generally provide rapid or immediate drug release from the formulation. Depending on the pharmacology and pharmacokinetics of the drug, use of conventional dosage forms can lead to wide fluctuations in the concentrations of the drug in a patient's blood and other tissues. These fluctuations can impact a number of parameters, such as dose frequency, onset of action, duration of efficacy, maintenance of therapeutic blood levels, toxicity, side effects, and the like. Advantageously, controlled-release formulations can be used to control a drug's onset of action, duration of action, plasma levels within the therapeutic window, and peak blood levels. In particular, controlled- or extended-release dosage forms or formulations can be used to ensure that the maximum effectiveness of a drug is achieved while minimizing potential adverse effects and safety concerns, which can occur both from under-dosing a drug (i.e., going below the minimum therapeutic levels) as well as exceeding the toxicity level for the drug. In some embodiments, the CTC marker-gene targeted therapy can be administered in a sustained release formulation.

**[00116]** Controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled release counterparts. Ideally, the use of an optimally

designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include: 1) extended activity of the drug; 2) reduced dosage frequency; 3) increased patient compliance; 4) usage of less total drug; 5) reduction in local or systemic side effects; 6) minimization of drug accumulation; 7) reduction in blood level fluctuations; 8) improvement in efficacy of treatment; 9) reduction of potentiation or loss of drug activity; and 10) improvement in speed of control of diseases or conditions. Kim, Cherrng-ju, Controlled Release Dosage Form Design, 2 (Technomic Publishing, Lancaster, Pa.: 2000).

**[00117]** Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, ionic strength, osmotic pressure, temperature, enzymes, water, and other physiological conditions or compounds.

**[00118]** A variety of known controlled- or extended-release dosage forms, formulations, and devices can be adapted for use with the salts and compositions of the disclosure. Examples include, but are not limited to, those described in U.S. Pat. Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; 4,008,719; 5674,533; 5,059,595; 5,591,767; 5,120,548; 5,073,543; 5,639,476; 5,354,556; 5,733,566; and 6,365,185 B1 ; each of which is incorporated herein by reference. These dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydroxypropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems (such as OROS® (Alza Corporation, Mountain View, Calif. USA)), or a combination thereof to provide the desired release profile in varying proportions.

**[00119]** The methods described herein can further comprise administering a second agent and/or treatment to the subject, e.g. as part of a combinatorial therapy. Non-limiting examples of a second agent and/or treatment can include radiation therapy, surgery, and chemotherapeutic agents as described above herein.

**[00120]** In certain embodiments, an effective dose of a composition comprising a CTC marker gene-targeted therapy as described herein can be administered to a patient once. In certain embodiments, an effective dose of a composition comprising a CTC marker gene-targeted therapy can be administered to a patient repeatedly. For systemic administration, subjects can be administered a therapeutic amount of a composition comprising a CTC marker gene-targeted therapy, such as, e.g. 0.1 mg/kg, 0.5 mg/kg, 1.0 mg/kg, 2.0 mg/kg, 2.5 mg/kg, 5 mg/kg, 10 mg/kg, 15 mg/kg, 20 mg/kg, 25 mg/kg, 30 mg/kg, 40 mg/kg, 50 mg/kg, or more.

**[00121]** In some embodiments, after an initial treatment regimen, the treatments can be administered on a less frequent basis. For example, after treatment biweekly for three months, treatment can be repeated once per month, for six months or a year or longer. Treatment according to the methods described herein can reduce levels of a marker or symptom of a condition, e.g. CTC levels by at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80 % or at least 90% or more.

**[00122]** The dosage of a composition as described herein can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment. With respect to duration and frequency of treatment, it is typical for skilled clinicians to monitor subjects in order to determine when the treatment is providing therapeutic benefit, and to determine whether to increase or decrease dosage, increase or decrease administration frequency, discontinue treatment, resume treatment, or make other alterations to the treatment regimen. The dosing schedule can vary from once a week to daily depending on a number of clinical factors, such as the subject's sensitivity to the CTC marker gene-targeted therapy. The desired dose or amount of activation can be administered at one time or divided into subdoses, e.g., 2-4 subdoses and administered over a period of time, e.g., at appropriate intervals through the day or other appropriate schedule. In some embodiments, administration can be chronic, e.g., one or more doses and/or treatments daily over a period of weeks or months. Examples of dosing and/or treatment schedules are administration daily, twice daily, three times daily or four or more times daily over a period of 1 week, 2 weeks, 3 weeks, 4 weeks, 1 month, 2 months, 3 months, 4 months, 5 months, or 6 months, or more. A composition comprising a CTC marker gene-targeted therapy can be administered over a period of time, such as over a 5 minute, 10 minute, 15 minute, 20 minute, or 25 minute period.

**[00123]** The dosage ranges for the administration of a CTC marker gene-targeted therapy, according to the methods described herein depend upon, for example, the form of the CTC marker gene-targeted therapy, its potency, and the extent to which symptoms, markers, or indicators of a condition described herein are desired to be reduced, for example the percentage reduction desired for CTC levels. The dosage should not be so large as to cause adverse side effects. Generally, the dosage will vary with the age, condition, and sex of the patient and can be determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

**[00124]** The efficacy of a CTC marker gene-targeted therapy in, e.g. the treatment of a condition described herein, or to induce a response as described herein (e.g. reduction of CTC levels) can be determined by the skilled clinician. However, a treatment is considered "effective treatment," as the term is used herein, if one or more of the signs or symptoms of a condition described herein are altered in a beneficial manner, other clinically accepted symptoms are improved, or even ameliorated, or a desired response is induced e.g., by at least 10% following treatment according to the methods described herein. Efficacy can be assessed, for example, by measuring a marker, indicator, symptom,

and/or the incidence of a condition treated according to the methods described herein or any other measurable parameter appropriate, e.g. tumor size and/or growth. Efficacy can also be measured by a failure of an individual to worsen as assessed by hospitalization, or need for medical interventions (i.e., progression of the disease is halted). Methods of measuring these indicators are known to those of skill in the art and/or are described herein. Treatment includes any treatment of a disease in an individual or an animal (some non-limiting examples include a human or an animal) and includes: (1) inhibiting the disease, e.g., preventing a worsening of symptoms (e.g. pain or inflammation); or (2) relieving the severity of the disease, e.g., causing regression of symptoms. An effective amount for the treatment of a disease means that amount which, when administered to a subject in need thereof, is sufficient to result in effective treatment as that term is defined herein, for that disease. Efficacy of an agent can be determined by assessing physical indicators of a condition or desired response, (e.g. CTC levels). It is well within the ability of one skilled in the art to monitor efficacy of administration and/or treatment by measuring any one of such parameters, or any combination of parameters. Efficacy can be assessed in animal models of a condition described herein, for example treatment of cancer, e.g. pancreatic cancer. When using an experimental animal model, efficacy of treatment is evidenced when a statistically significant change in a marker is observed, e.g. a change in CTC levels.

**[00125]** For convenience, the meaning of some terms and phrases used in the specification, examples, and appended claims, are provided below. Unless stated otherwise, or implicit from context, the following terms and phrases include the meanings provided below. The definitions are provided to aid in describing particular embodiments, and are not intended to limit the claimed invention, because the scope of the invention is limited only by the claims. Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. If there is an apparent discrepancy between the usage of a term in the art and its definition provided herein, the definition provided within the specification shall prevail.

**[00126]** For convenience, certain terms employed herein, in the specification, examples and appended claims are collected here.

**[00127]** The terms “decrease”, “reduced”, “reduction”, or “inhibit” are all used herein to mean a decrease by a statistically significant amount. In some embodiments, “reduce,” “reduction” or “decrease” or “inhibit” typically means a decrease by at least 10% as compared to a reference level (e.g. the absence of a given treatment) and can include, for example, a decrease by at least about 10%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 98%, at least about 99%, or more. As used herein, “reduction” or “inhibition” does not encompass a complete inhibition or reduction as compared to a reference level.

“Complete inhibition” is a 100% inhibition as compared to a reference level. A decrease can be preferably down to a level accepted as within the range of normal for an individual without a given disorder.

**[00128]** The terms “increased”, “increase”, “enhance”, or “activate” are all used herein to mean an increase by a statically significant amount. In some embodiments, the terms “increased”, “increase”, “enhance”, or “activate” can mean an increase of at least 10% as compared to a reference level, for example an increase of at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, or any increase between 2-fold and 10-fold or greater as compared to a reference level. In the context of a marker or symptom, a “increase” is a statistically significant increase in such level.

**[00129]** As used herein, a "subject" means a human or animal. Usually the animal is a vertebrate such as a primate, rodent, domestic animal or game animal. Primates include chimpanzees, cynomolgous monkeys, spider monkeys, and macaques, e.g., Rhesus. Rodents include mice, rats, woodchucks, ferrets, rabbits and hamsters. Domestic and game animals include cows, horses, pigs, deer, bison, buffalo, feline species, e.g., domestic cat, canine species, e.g., dog, fox, wolf, avian species, e.g., chicken, emu, ostrich, and fish, e.g., trout, catfish and salmon. In some embodiments, the subject is a mammal, e.g., a primate, e.g., a human. The terms, “individual,” “patient” and “subject” are used interchangeably herein.

**[00130]** Preferably, the subject is a mammal. The mammal can be a human, non-human primate, mouse, rat, dog, cat, horse, or cow, but is not limited to these examples. Mammals other than humans can be advantageously used as subjects that represent animal models of cancer. A subject can be male or female.

**[00131]** A subject can be one who has been previously diagnosed with or identified as suffering from or having a condition in need of treatment (e.g. cancer) or one or more complications related to such a condition, and optionally, have already undergone treatment for cancer or the one or more complications related to cancer. Alternatively, a subject can also be one who has not been previously diagnosed as having cancer or one or more complications related to cancer. For example, a subject can be one who exhibits one or more risk factors for cancer or one or more complications related to cancer or a subject who does not exhibit risk factors.

**[00132]** A “subject in need” of treatment for a particular condition can be a subject having that condition, diagnosed as having that condition, or at risk of developing that condition.

**[00133]** As used herein, the term “cancer” or “tumor” refers to an uncontrolled growth of cells which interferes with the normal functioning of the bodily organs and systems. A subject who has a

cancer or a tumor is a subject having objectively measurable cancer cells present in the subject's body. Included in this definition are benign and malignant cancers, as well as dormant tumors or micrometastases. Cancers which migrate from their original location and seed vital organs can eventually lead to the death of the subject through the functional deterioration of the affected organs.

**[00134]** The term "agent" refers generally to any entity which is normally not present or not present at the levels being administered to a cell, tissue or subject. An agent can be selected from a group including but not limited to: polynucleotides; polypeptides; small molecules; and antibodies or antigen-binding fragments thereof. A polynucleotide can be RNA or DNA, and can be single or double stranded, and can be selected from a group including, for example, nucleic acids and nucleic acid analogues that encode a polypeptide. A polypeptide can be, but is not limited to, a naturally-occurring polypeptide, a mutated polypeptide or a fragment thereof that retains the function of interest. Further examples of agents include, but are not limited to a nucleic acid aptamer, peptide-nucleic acid (PNA), locked nucleic acid (LNA), small organic or inorganic molecules; saccharide; oligosaccharides; polysaccharides; biological macromolecules, peptidomimetics; nucleic acid analogs and derivatives; extracts made from biological materials such as bacteria, plants, fungi, or mammalian cells or tissues and naturally occurring or synthetic compositions. An agent can be applied to the media, where it contacts the cell and induces its effects. Alternatively, an agent can be intracellular as a result of introduction of a nucleic acid sequence encoding the agent into the cell and its transcription resulting in the production of the nucleic acid and/or protein environmental stimuli within the cell. In some embodiments, the agent is any chemical, entity or moiety, including without limitation synthetic and naturally-occurring non-proteinaceous entities. In certain embodiments the agent is a small molecule having a chemical moiety selected, for example, from unsubstituted or substituted alkyl, aromatic, or heterocyclyl moieties including macrolides, leptomycins and related natural products or analogues thereof. Agents can be known to have a desired activity and/or property, or can be selected from a library of diverse compounds. As used herein, the term "small molecule" can refer to compounds that are "natural product-like," however, the term "small molecule" is not limited to "natural product-like" compounds. Rather, a small molecule is typically characterized in that it contains several carbon—carbon bonds, and has a molecular weight more than about 50, but less than about 5000 Daltons (5 kD). Preferably the small molecule has a molecular weight of less than 3 kD, still more preferably less than 2 kD, and most preferably less than 1 kD. In some cases it is preferred that a small molecule have a molecular mass equal to or less than 700 Daltons.

**[00135]** Aptamers are short synthetic single-stranded oligonucleotides that specifically bind to various molecular targets such as small molecules, proteins, nucleic acids, and even cells and tissues. These small nucleic acid molecules can form secondary and tertiary structures capable of specifically binding proteins or other cellular targets, and are essentially a chemical equivalent of antibodies. Aptamers are highly specific, relatively small in size, and non-immunogenic. Aptamers are generally

selected from a biopanning method known as SELEX (Systematic Evolution of Ligands by Exponential enrichment) (Ellington et al. *Nature*. 1990;346(6287):818–822; Tuerk et al., *Science*. 1990;249(4968):505–510; Ni et al., *Curr Med Chem*. 2011;18(27):4206-14; which are incorporated by reference herein in their entireties). Methods of generating an aptamer for any given target are well known in the art. Preclinical studies using, e.g. aptamer-siRNA chimeras and aptamer targeted nanoparticle therapeutics have been very successful in mouse models of cancer and HIV (Ni et al., *Curr Med Chem*. 2011;18(27):4206-14).

**[00136]** As used herein, the terms “protein” and “polypeptide” are used interchangeably herein to designate a series of amino acid residues, connected to each other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues. The terms “protein”, and “polypeptide” refer to a polymer of amino acids, including modified amino acids (e.g., phosphorylated, glycated, glycosylated, etc.) and amino acid analogs, regardless of its size or function. “Protein” and “polypeptide” are often used in reference to relatively large polypeptides, whereas the term “peptide” is often used in reference to small polypeptides, but usage of these terms in the art overlaps. The terms “protein” and “polypeptide” are used interchangeably herein when referring to a gene product and fragments thereof. Thus, exemplary polypeptides or proteins include gene products, naturally occurring proteins, homologs, orthologs, paralogs, fragments and other equivalents, variants, fragments, and analogs of the foregoing.

**[00137]** As used herein, the term “nucleic acid” or “nucleic acid sequence” refers to any molecule, preferably a polymeric molecule, incorporating units of ribonucleic acid, deoxyribonucleic acid or an analog thereof. The nucleic acid can be either single-stranded or double-stranded. A single-stranded nucleic acid can be one nucleic acid strand of a denatured double- stranded DNA. Alternatively, it can be a single-stranded nucleic acid not derived from any double-stranded DNA. In one aspect, the nucleic acid can be DNA. In another aspect, the nucleic acid can be RNA. Suitable nucleic acid molecules are DNA, including genomic DNA or cDNA. Other suitable nucleic acid molecules are RNA, including mRNA.

**[00138]** As used herein, the terms “treat,” “treatment,” “treating,” or “amelioration” refer to therapeutic treatments, wherein the object is to reverse, alleviate, ameliorate, inhibit, slow down or stop the progression or severity of a condition associated with a disease or disorder, e.g. cancer. The term “treating” includes reducing or alleviating at least one adverse effect or symptom of a condition, disease or disorder associated with a cancer. Treatment is generally “effective” if one or more symptoms or clinical markers are reduced. Alternatively, treatment is “effective” if the progression of a disease is reduced or halted. That is, “treatment” includes not just the improvement of symptoms or markers, but also a cessation of, or at least slowing of, progress or worsening of symptoms compared to what would be expected in the absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease,

stabilized (*i.e.*, not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, remission (whether partial or total), and/or decreased mortality, whether detectable or undetectable. The term "treatment" of a disease also includes providing relief from the symptoms or side-effects of the disease (including palliative treatment).

**[00139]** As used herein, the term "pharmaceutical composition" refers to the active agent in combination with a pharmaceutically acceptable carrier *e.g.* a carrier commonly used in the pharmaceutical industry. The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

**[00140]** As used herein, the term "administering," refers to the placement of a compound as disclosed herein into a subject by a method or route which results in at least partial delivery of the agent at a desired site. Pharmaceutical compositions comprising the compounds disclosed herein can be administered by any appropriate route which results in an effective treatment in the subject.

**[00141]** The term "statistically significant" or "significantly" refers to statistical significance and generally means a two standard deviation (2SD) or greater difference.

**[00142]** Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients or reaction conditions used herein should be understood as modified in all instances by the term "about." The term "about" when used in connection with percentages can mean  $\pm 1\%$ .

**[00143]** As used herein the term "comprising" or "comprises" is used in reference to compositions, methods, and respective component(s) thereof, that are essential to the method or composition, yet open to the inclusion of unspecified elements, whether essential or not.

**[00144]** The term "consisting of" refers to compositions, methods, and respective components thereof as described herein, which are exclusive of any element not recited in that description of the embodiment.

**[00145]** As used herein the term "consisting essentially of" refers to those elements required for a given embodiment. The term permits the presence of elements that do not materially affect the basic and novel or functional characteristic(s) of that embodiment.

**[00146]** The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The abbreviation, "e.g." is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation "e.g." is synonymous with the term "for example."

**[00147]** Definitions of common terms in cell biology and molecular biology can be found in “The Merck Manual of Diagnosis and Therapy”, 19th Edition, published by Merck Research Laboratories, 2006 (ISBN 0-911910-19-0); Robert S. Porter et al. (eds.), The Encyclopedia of Molecular Biology, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); Benjamin Lewin, Genes X, published by Jones & Bartlett Publishing, 2009 (ISBN-10: 0763766321); Kendrew et al. (eds.), Molecular Biology and Biotechnology: a Comprehensive Desk Reference, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8) and Current Protocols in Protein Sciences 2009, Wiley Intersciences, Coligan et al., eds.

**[00148]** Unless otherwise stated, the present invention was performed using standard procedures, as described, for example in Sambrook et al., Molecular Cloning: A Laboratory Manual (4 ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (2012); Davis et al., Basic Methods in Molecular Biology, Elsevier Science Publishing, Inc., New York, USA (1995); or Methods in Enzymology: Guide to Molecular Cloning Techniques Vol.152, S. L. Berger and A. R. Kimmel Eds., Academic Press Inc., San Diego, USA (1987); Current Protocols in Protein Science (CPPS) (John E. Coligan, et. al., ed., John Wiley and Sons, Inc.), Current Protocols in Cell Biology (CPCB) (Juan S. Bonifacino et. al. ed., John Wiley and Sons, Inc.), and Culture of Animal Cells: A Manual of Basic Technique by R. Ian Freshney, Publisher: Wiley-Liss; 5th edition (2005), Animal Cell Culture Methods (Methods in Cell Biology, Vol. 57, Jennie P. Mather and David Barnes editors, Academic Press, 1st edition, 1998) which are all incorporated by reference herein in their entireties.

**[00149]** Other terms are defined herein within the description of the various aspects of the invention.

**[00150]** All patents and other publications; including literature references, issued patents, published patent applications, and co-pending patent applications; cited throughout this application are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the technology described herein. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.

**[00151]** The description of embodiments of the disclosure is not intended to be exhaustive or to limit the disclosure to the precise form disclosed. While specific embodiments of, and examples for, the disclosure are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the disclosure, as those skilled in the relevant art will recognize. For example, while method steps or functions are presented in a given order, alternative embodiments

may perform functions in a different order, or functions may be performed substantially concurrently. The teachings of the disclosure provided herein can be applied to other procedures or methods as appropriate. The various embodiments described herein can be combined to provide further embodiments. Aspects of the disclosure can be modified, if necessary, to employ the compositions, functions and concepts of the above references and application to provide yet further embodiments of the disclosure. Moreover, due to biological functional equivalency considerations, some changes can be made in protein structure without affecting the biological or chemical action in kind or amount. These and other changes can be made to the disclosure in light of the detailed description. All such modifications are intended to be included within the scope of the appended claims.

**[00152]** Specific elements of any of the foregoing embodiments can be combined or substituted for elements in other embodiments. Furthermore, while advantages associated with certain embodiments of the disclosure have been described in the context of these embodiments, other embodiments may also exhibit such advantages, and not all embodiments need necessarily exhibit such advantages to fall within the scope of the disclosure.

**[00153]** The technology described herein is further illustrated by the following examples which in no way should be construed as being further limiting.

**[00154]** Some embodiments of the technology described herein can be defined according to any of the following numbered paragraphs:

1. A method of detecting circulating tumor cells (CTCs) in a sample, the method comprising:  
measuring the level of a PC-CTC marker gene expression product in the sample; and  
determining that PC-CTCs are present if the detected level of the marker gene  
expression product is greater than a reference level.
2. The method of paragraph 1, wherein the CTCs are pancreatic cancer CTCs.
3. The method of any of paragraphs 1-2, wherein the method further comprises a first step of  
isolating the CTCs from the sample.
4. The method of any of paragraphs 1-3, wherein the expression product is a nucleic acid.
5. The method of paragraph 4, wherein the level of the expression product is determined using a  
method selected from the group consisting of:  
RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis;  
next-generation sequencing; and RNA in situ hybridization.
6. The method of any of paragraphs 1-3, wherein the expression product is a polypeptide.
7. The method of paragraph 6, wherein the level of the expression product is determined using a  
method selected from the group consisting of:  
Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA);  
radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization

(FISH); immunohistological staining; radioimmunometric assay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay.

8. The method of any of paragraphs 1-7, wherein the CTC marker gene is selected from Table 7; Table 8; or Table 14.

9. The method of any of paragraphs 1-8, wherein the CTC marker gene is selected from the group consisting of:

ABI3BP; ADAMTS5; ADAMTSL1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4.

10. The method of any of paragraphs 1-8, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2.

11. The method of any of paragraphs 1-9, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

12. The method of any of paragraphs 1-9, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A2; IGFBP5; KLF4; and DCN.

13. The method of any of paragraphs 1-9, wherein the CTC marker gene is selected from the group consisting of:

TPT1; HMGB1; SPON 2; SPARC; and ARSA.

14. The method of any of paragraphs 1-9, wherein the CTC marker gene is selected from the group consisting of:

IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A.

15. A method of treating cancer in a subject, the method comprising administering a therapeutically effective amount of a CTC marker gene-targeted therapy to the subject.

16. The method of paragraph 15, wherein the cancer is pancreatic cancer.

17. The method of any of paragraphs 15-16, wherein the CTC marker gene-targeted therapy comprises an inhibitor of a CTC marker gene.

18. The method of paragraph 17, wherein the inhibitor is an antibody reagent.

19. The method of paragraph 17, wherein the inhibitor is an inhibitory nucleic acid reagent.
20. The method of any of paragraphs 15-19, wherein the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent and a chemotherapeutic agent.
21. The method of any of paragraphs 15-20, wherein the subject is a subject determined to have an elevated level of CTCs and/or an elevated level of a CTC marker gene present in the blood and/or stroma of the cancer.
22. The method of any of paragraphs 15-21, wherein the CTC marker gene-targeted therapy comprises a CTC marker gene-binding antibody reagent that binds a marker gene selected from the group consisting of:

IL6ST, SULF2, and SV2A.
23. A method of determining if a subject is likely to respond to treatment with a CTC marker gene-targeted therapy, the method comprising:

measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and

determining that the subject is likely to respond to the treatment if the level of the expression product is increased relative to a reference level.
24. The method of paragraph 23, wherein the method further comprises a first step of isolating the CTCs from the sample.
25. The method of any of paragraphs 23-24, wherein the cancer is pancreatic cancer.
26. The method of any of paragraphs 23-25, wherein the expression product is a nucleic acid.
27. The method of paragraph 26, wherein the level of the expression product is determined using a method selected from the group consisting of:

RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization.
28. The method of any of paragraphs 23-26, wherein the expression product is a polypeptide.
29. The method of paragraph 28, wherein the level of the expression product is determined using a method selected from the group consisting of:

Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunometric assay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay.
30. The method of any of paragraphs 23-29, wherein the PC-CTC marker gene is selected from Table 7; Table 8; or Table 14.
31. The method of any of paragraphs 23-30, wherein the CTC marker gene is selected from the group consisting of:

ABI3BP; ADAMTS5; ADAMTSL1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4.

32. The method of any of paragraphs 23-31, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2.

33. The method of any of paragraphs 23-31, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.

34. The method of any of paragraphs 23-31, wherein the CTC marker gene is selected from the group consisting of:

ALDH1A2; IGFBP5; KLF4; and DCN.

35. The method of any of paragraphs 23-31, wherein the CTC marker gene is selected from the group consisting of:

TPT1; HMGB1; SPON 2; SPARC; and ARSA.

36. The method of any of paragraphs 23-31, wherein the CTC marker gene is selected from the group consisting of:

IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A.

37. A method of monitoring the treatment of a subject, the method comprising:

administering a cancer therapy to a subject in need thereof;

measuring the level of a CTC marker gene expression product present in the blood and/or stroma of a cancer; and

determining that the subject is responding if the level of the CTC marker gene expression product is decreased relative to the reference level and determining that the subject is not responding to the treatment if the CTC marker gene expression product is not decreased relative to the reference level.

38. The method of paragraph 37, wherein the cancer is pancreatic cancer.

39. The method of any of paragraphs 37-38, wherein the reference level is the level of the gene expression product in the patient prior to the administering step.

40. The method of any of paragraphs 37-39, wherein the method further comprises a first step of isolating the CTCs from the sample.

41. The method of any of paragraphs 37-40, wherein the expression product is a nucleic acid.
42. The method of paragraph 41, wherein the level of the expression product is determined using a method selected from the group consisting of:  
RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization.
43. The method of any of paragraphs 37-40, wherein the expression product is a polypeptide.
44. The method of paragraph 43, wherein the level of the expression product is determined using a method selected from the group consisting of:  
Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunoassay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay.
45. The method of any of paragraphs 37-44, wherein the PC-CTC marker gene is selected from Table 7; Table 8; or Table 14.
46. The method of any of paragraphs 37-45, wherein the CTC marker gene is selected from the group consisting of:  
ABI3BP; ADAMTS5; ADAMTSL1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1; FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4.
47. The method of any of paragraphs 37-46, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2.
48. The method of any of paragraphs 37-46, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.
49. The method of any of paragraphs 37-46, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A2; IGFBP5; KLF4; and DCN.
50. The method of any of paragraphs 37-46, wherein the CTC marker gene is selected from the group consisting of:  
TPT1; HMGB1; SPON 2; SPARC; and ARSA.

51. The method of any of paragraphs 37-46, wherein the CTC marker gene is selected from the group consisting of:

IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A.

## EXAMPLES

**[00155] EXAMPLE 1: Single Cell RNA-Sequencing of Mouse Pancreatic Circulating Tumor Cells Reveals their Expression of ECM Proteins**

**[00156]** Circulating Tumor Cells (CTCs) are shed from primary tumors into the bloodstream, mediating the hematogenous spread of cancer to distant organs. Using a pancreatic cancer mouse model, a microfluidic device was applied to isolate CTCs independently of tumor epitopes, subjecting these to single cell RNA-sequencing. CTCs clustered into multiple subsets, distinct from primary tumors and cancer cell lines. While proliferative signatures were generally low, CTCs were enriched for MAPK, as well as WNT, TGF- $\beta$ , Neurotrophin, Toll-like receptor, and B-cell receptor signaling pathways. CTCs were highly enriched for expression of the stem-cell associated gene *Aldh1a2*. Their virtually universal expression of *Igfbp5* and *Klf4* was correlated with a subset of primary tumor cells localized to the epithelial/stromal boundary, consistent with the presence of both epithelial and mesenchymal markers in CTCs. The very high CTC expression of stromal-derived extracellular matrix proteins, including *Dcn* and *Sparc*, indicates microenvironmental contributions to metastasis and identifies unexpected therapeutic targets.

**[00157] Introduction**

**[00158]** Pancreatic ductal adenocarcinoma (PDAC) is the fourth leading cause of cancer deaths in the US, with a 6% overall survival at 5 years (Society, 2013). The high mortality of this cancer stems from the rapid dissemination of tumor cells leading to widespread metastasis. While local tissue and lymphatic invasion are evident even in early PDAC, the presence of circulating tumor cells (CTCs) in the bloodstream ultimately leads to spread of cancer to distant organs. CTCs are rare, estimated at one to ten tumor cells among ten billion normal blood cells in a milliliter of blood. As such, their isolation and molecular analysis has posed a significant technological challenge (Pantel et al., 2008; Yu et al., 2011). Given their role in blood-borne metastasis, CTC populations are likely to be enriched for metastatic precursors, and their analysis may identify potential therapeutic targets, as well as providing opportunities for early detection of pancreatic cancer.

**[00159]** Genetically engineered mouse pancreatic cancer models have provided important insight into the progression of this disease. Specifically, the genetically engineered *LSL-Kras*<sup>G12D</sup>, *Trp53*<sup>fl/fl</sup> *or*<sup>+</sup>, *Pdx1-Cre* (KPC) mouse model recapitulates the histological progression from preneoplastic pancreatic intraepithelial neoplasia (PanIN) lesions to invasive carcinoma (Bardeesy et al., 2006). Recent studies have suggested that epithelial-to-mesenchymal transition (EMT) occurs early in this model potentially enhancing tumor invasiveness (Rhim et al., 2012). In an initial molecular

characterization of mouse pancreatic CTCs, RNA sequencing of CTC-enriched populations was performed, thereby identifying activation of non-canonical WNT signaling as a recurrent event, potentially contributing to the anoikis resistance of circulating epithelial cells (Yu et al., 2012). In that study, analysis of purified CTC populations was accomplished using single molecule RNA sequencing, combined with digital subtraction of matched leukocyte RNA reads, so as to derive a CTC-enriched expression signature. However, transcriptomic analysis of such partially purified cell populations is limited by depth of coverage to the most highly differentially expressed genes, and such studies of bulk CTC populations cannot resolve the degree of heterogeneity across these poorly understood cell populations.

**[00160]** To achieve a deep RNA sequencing profile of CTCs at the single cell level, a novel inertial focusing-enhanced device, the CTC-iChip, which allows high efficiency negative depletion of normal blood cells, leaving unattached CTCs in solution where they can be selected and analyzed as single cells (Ozkumur et al., 2013) was used. By avoiding tumor epitope-specific capture, such as targeting the epithelial marker EpCAM, the CTC-iChip is unbiased in isolating cancer cells with both epithelial and mesenchymal characteristics. Further, the high quality of RNA purified from viable, untagged CTCs is particularly well suited for detailed transcriptomic analysis. Finally, the use of a mouse model of pancreatic cancer allows for simultaneous analysis of primary tumor and CTCs, while the shared driver mutations across different animals facilitates the identification of CTC-specific heterogeneity. Described herein is a comprehensive transcriptome analysis of CTCs at the single cell level, pointing to distinct cell subsets within CTC populations, signaling pathways that are enriched in CTCs, and identifying unique CTC markers and therapeutic targets.

### **[00161] Results**

**[00162] *Isolation of Mouse Pancreatic CTCs***. The CTC-iChip, an integrated microfluidic cell separation platform applied directly to whole blood specimens for isolation of CTCs (Ozkumur et al., 2013) was used in the experiments described herein. It combines initial hydrodynamic size-based separation of all nucleated cells (leukocytes (WBC) and CTCs) away from red blood cells, platelets and plasma, with subsequent inertial focusing of the nucleated cells within a single streamline to achieve high efficiency in-line magnetic sorting. While tumor epitopes are highly variable, WBC cell surface markers are well established; applying magnetic-conjugated anti-WBC antibodies to this very high throughput microfluidic cell separation device can thus exclude the vast majority of WBCs to reveal a small number of untagged CTCs (Fig. 1A). The CTC-iChip was adapted for depletion of murine hematopoietic cells and applied to the KPC pancreatic cancer mouse model. This PDAC model generates significant numbers of CTCs (Rhim et al., 2012; Yu et al., 2012). Whole blood labeling using 100 anti-CD45 beads per WBC achieved  $> 10^3$  depletion in normal mice, mice bearing orthotopic tumors, and the genetically engineered KPC mice (Figs. 1B and 4A-4C).

**[00163]** CTC recovery was measured as a mean of 95% (+/- 3% std), using GFP-tagged NB508 mouse pancreatic cancer cells spiked into whole mouse blood and processed through the CTC-iChip (Figs. 4A-4C). NB508 cells were previously generated from a pancreatic tumor arising in the same *Kras/Trp53*-driven KPC mouse model (Bardeesy et al., 2006). In comparison, only 35% recovery of the same cells was achieved using an alternative microfluidic platform based on anti-EpCAM capture of mouse CTCs (Yu et al., 2012). Applying the CTC-iChip to orthotopic tumors derived from pancreatic inoculation of GFP-tagged NB508 cells generated > 1000 CTCs/mL in all three mice tested (Figs. 4A-4C). Finally, testing the CTC-iChip with the genetically engineered KPC model, followed by dual immunofluorescence staining of isolated cells for the epithelial marker pan-cytokeratin (CK) versus the leukocyte marker CD45, revealed a median 118 CTCs/mL (mean 429 CTCs/mL; range 0-1694) (Fig. 1C). No CK positive cells were isolated from 7 healthy control mice. The vast majority of CD45 positive cells that failed to be deflected in the microfluidic device retained some immunomagnetic beads on their surface. Thus, CTCs were readily distinguished from WBCs in the CTC-iChip product, enabling single cell manipulation without requiring staining for epithelial-specific cell surface epitopes, such as EpCAM.

**[00164]** *Single CTC RNA-sequencing.* Five tumor-bearing KPC mice generated a total of 168 single CTCs that were subjected to a modified initial cDNA amplification and library protocol (Tang et al., 2010), and screened for RNA quality (*Gapdh*, *Actb*), presence of pancreatic markers (*Krt8*, *Krt18*, *Krt19*, *Pdx1*), and absence of WBC markers (*Cd45/Ptprc*) (Figs. 5A-5C). Of these, 75 (45%) were of sufficient quality to proceed to further amplification and library construction for next generation sequencing. It is noteworthy that a majority of candidate CTCs (55%) appeared morphologically intact but had degraded RNA. These cells likely represent tumor cells that have lost viability in the bloodstream. Given the rapid processing of blood samples from mouse models, the minimal shear condition in the microfluidic device, and the preserved RNA quality of control cells processed identically, it is unlikely that cells underwent such damage during in vitro purification. For comparison with pancreatic CTCs, single cell RNA-sequencing was also performed on 12 WBCs from a control mouse, 12 mouse embryonic fibroblasts (MEFs), and 16 single cells from the mouse NB508 pancreatic cancer cell line. Over 90% of single cells from NB508 and MEF cultures met criteria for sequencing quality, highlighting the high frequency of CTCs with compromised RNA templates under the same conditions. To compare CTC profiles to that of matched parental tumors harvested at the time of CTC isolation, bulk RNA from each primary tumor was diluted to 1 or 10 cell equivalents (10 or 100 pg RNA) and subjected to the same amplification and RNA-sequencing protocol (n = 34; min 8 replicates from 4 matched tumors).

**[00165]** Single cell RNA sequencing performance was comparable for all samples analyzed, with a mean 4.4-8.5 million reads, of which a mean 46-61% were uniquely aligned to the genome (Figs. 5A-5C). Genome aligned reads were annotated and counted using UCSC Known Gene transcriptome

reference and normalized in reads per million (RPM). Normalized reads were then analyzed by unsupervised hierarchical clustering (data not shown). Single cell transcriptomes from MEFs, the NB508 pancreatic cancer cell line and normal WBCs were tightly clustered, supporting the analytic reliability of the RNA sequencing strategy. Five distinct clusters of candidate CTCs were identified, all of which were distinct from matched primary tumor sequences, as well as from cancer-derived cell lines. Principal component analysis demonstrates the clustering and inter-relationships of these different groups (Fig. 2).

**[00166]** The uniform genetic drivers of PDAC in the KPC mouse model made it possible to quantify measures of cellular heterogeneity in CTCs derived from individual mice and across different mice. Single cell heterogeneity within each CTC cluster was assessed by calculating the intra-cluster correlation coefficients, where lower correlation coefficients reflect higher heterogeneity (Figs. 5A-5C). As expected, CTC clusters showed considerably more heterogeneity (mean 0.42, 95% CI 0.36-0.47) than single cells derived from the NB508 cancer cell line (mean 0.86, 95% CI 0.80-0.91, p-value  $1.2 \times 10^{-15}$ ). To assess heterogeneity of cells within a primary PDAC, a conditional Tomato/EGFP (mT/mG) expression marker (Muzumdar et al., 2007) was crossed with the KPC mouse to generate a lineage-tagged mouse tumor (KPC-mT/mG), which could be used to isolate individual EGFP positive primary tumor cells away from contaminating stromal cells. A primary tumor (TuGMP3) was disaggregated into single cell suspension and 20 EGFP positive cells were subjected to RNA sequencing. The single primary tumor cells clustered well within the previously analyzed bulk tumor material (data not shown), with a heterogeneity score (mean 0.38, 95% CI 0.28-0.47) similar to that of CTCs (p-value 0.49).

**[00167]** In summary, described herein is the single cell RNA-sequencing of mouse pancreatic CTCs isolated without positive selection bias, along with parental tumors, an established genotype-matched cancer cell line, MEFs and WBCs. CTCs clustered separately from the primary tumor (both bulk tumor and isolated single cells) and from the tumor-derived cell line, with comparable degrees of intercellular heterogeneity between CTCs and primary tumor cells.

**[00168] *Defining Subsets of Pancreatic CTCs.*** To identify and classify candidate CTCs, gene sets for known epithelial, hematopoietic, and endothelial markers were applied across all clustered samples. As expected, epithelial markers (*Krt7*, *Krt8*, *Krt18*, *Krt19*, *Epcam*, *Egfr*, *Cdh1*) were highly expressed in primary pancreatic tumors and in the cancer cell line NB508, and nearly absent in the non-epithelial MEFs and in normal WBCs (data not shown). In contrast, hematopoietic markers (*Ptprc/Cd45*, *Csf3r/Cd114*, *Cd14*, *Fcgr3/Cd16*, *Itga2b/Cd41*, *Itgb3/Cd61*) were present in normal WBCs, and absent in NB508 and MEFs. Some expression of hematopoietic markers was detectable in the bulk primary tumor samples, consistent with varying degrees of leukocytic infiltrates. No specific cluster of endothelial cells was identified, based on expression of characteristic markers

(*Cdh5/Cd144*, *Vwf*, *Thbd/Cd141*, *Pecam1/Cd31*, *Mcam/Cd146*, *Sele/E-selectin*, *Cd34*) and absence of epithelial and hematopoietic markers.

**[00169]** Interrogation of single cells isolated by CD45-depletion from tumor-bearing mice, using the epithelial, hematopoietic and endothelial markers, revealed five major candidate CTC groupings (Clusters 1, 3, 4, 5 and 9; data not shown). Clusters 3, 4, and 5 were all part of a larger grouping, showing strong expression of epithelial markers, consistent with “classical” CTCs (denoted CTC-c). A subset of these cells expressed *Cd34*, an endothelial progenitor marker that is also found in mesenchymal cells including MEFs (data not shown) and stromal cells (Krause et al., 1994), but other characteristic endothelial lineage markers were absent. Clusters 1 and 9 were more complex, with the former noteworthy for enrichment of platelet markers CD41 (*Itga2b*) and CD61 (*Itgb3*) (hence denoted CTC-plt), and the latter having a prominent cellular proliferation signature (CTC-pro).

**[00170]** To better define the characteristics of each candidate CTC cluster, a non-parametric differential gene expression analysis including a rank product (RP) methodology adapted to variations in absolute transcript levels and differences in transcriptome representation from cell to cell was used (Breitling et al., 2004). Setting very stringent parameters (FDR  $\leq 0.01$ ), the control comparison of primary tumors versus WBCs identified 927 genes relatively overexpressed in tumors and 293 genes high in WBCs, including the expected differential expression of epithelial tumor markers keratin 7, 8, 18, and 19, versus the leukocyte specific CD45 (data not shown). Comparing the “classical” CTC-c cluster to WBCs also showed enrichment for cytokeratin 18 and 19 in CTCs versus CD45 in WBCs, validating the RP methodology to identify relevant differentially expressed genes between single cell populations.

**[00171]** The most abundant CTC cluster, CTC-c, comprised 41 of 75 cells (55%) meeting established criteria for epithelial tumor cells (versus CTC-plt: 32%; CTC-pro: 13%). Of note, the only mouse with multiple gross metastases (MP7) had large numbers of CTCs within this class. Compared with matched primary tumors CTC-c cells had 878 transcripts increased in expression and 774 genes with reduced expression (Table 2). Gene Ontology (GO) analysis of CTC-c enriched genes (Table 3) indicated enrichment for signatures associated with cellular interactions with environmental signals (GO:0045785 – positive regulation of cell adhesion; GO:0048584 – positive regulation of response to stimulus), cell shape and structure (GO: 0030036 – actin cytoskeleton organization; GO:0060429 – epithelium development), and transcriptional states (GO:0045449 – regulation of transcription; GO:0051276 – chromosome organization). To evaluate the contribution of signaling pathways activated by external stimuli in CTC-c cells, the enriched genes were annotated using the KEGG database (Table 1). Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathway analysis similarly showed enrichment for focal adhesion (odds ratio [OR] 2.7, q-value 6.7 3 10.4) and regulation of actin cytoskeleton (OR 2.4, q-value 0.005). Notably, of the KEGG signaling pathways annotated, the mitogen-activated protein

kinase (MAPK) pathway was most highly enriched. Most highly represented was the MAPK pathway (OR 2.2, q-value 0.006); MAPK signaling is already activated in the *Kras*<sup>G12D</sup> driven primary tumor. However, while MSigDB Kras dependency signatures were enriched in primary tumors compared with CTCs, the latter had increased expression of *Braf*, *Mras* and *Rras2*, pointing to alternative paths to further activate MAPK in CTCs. This finding is consistent with another study that identified the MAPK pathway as being the most highly enriched in pancreatic CTCs using microarray based methodologies (Sergeant et al., 2012).

**[00172]** CTC enriched genes also had representation of well established signaling pathways involved with metastasis, including TGF-β (Ikushima and Miyazono, 2010; Siegel and Massague, 2003), WNT (Anastas and Moon, 2013; Clevers and Nusse, 2012; Katoh and Katoh, 2007), and VEGF (Carmeliet and Jain, 2011; Folkman, 1995). In this cohort of pancreatic cancer CTCs, *Wnt4* and *Tgfb2* were most highly enriched in CTCs relative to primary tumor, implicating autocrine signaling involving these major pathways. In addition to these well defined contributors to metastasis, CTC expression analyses also revealed activation of unexpected signaling pathways, including the neurotrophin, toll-like receptor, and B-cell receptor pathways. Neurotrophin pathway activation has been reported in pancreatic cancer, particularly in association with increased perineural invasion (Miknyoczki et al., 1996; Miknyoczki et al., 1999; Ohta et al., 1997; Wang et al., 2009; Zhang et al., 2005). Toll-like receptor and B-cell receptor pathways had less representation among CTC reads, but they suggest aberrant activation of immunomodulatory signaling components. Ultimately, the establishment of CTC-derived cultures will be required to test the functional significance of these activated signaling pathways.

**[00173]** While single cells within the CTC-c cluster fulfilled characteristic criteria for tumor cells, defining the identity of the non-classical CTC clusters, CTC-plt and CTC-pro, required additional analyses. Compared with CTC-c, single cells within the CTC-plt cluster had a high enrichment for wound healing and hemostasis signatures, as well as MSigDB platelet and megakaryocyte expression profiles (Table 4). This indicates that these cells are either circulating megakaryocytes/giant platelets or CTCs covered with adherent platelets. Tumor cell specific lineage tagging supports the identification of CTC-plt cells being of tumor origin. Eighteen EGFP lineage-tagged single CTCs from two KPC-mT/mG mice were subjected to single cell RNA sequencing: a total of 9 CTCs from the two mice (7/7 CTCs from mouse GMP1 and 2/11 from mouse GMP2) were included within CTC-plt, using unsupervised hierarchical clustering (data not shown). Thus, the CTC-plt cluster includes CTCs that exhibit strong platelet markers, most likely derived from transcripts encoded by adherent platelets. Interestingly, CTC-plt cells maintained their distinct segregation from CTC-c even after digital removal of all annotated platelet transcripts (data not shown). It is therefore possible that the adherence of abundant platelets may modulate the intrinsic CTC expression profile, as recently suggested by *in vitro* modeling experiments (Labelle et al., 2011).

**[00174]** The CTC-pro cluster was most similar to both the NB508 pancreatic cancer cell line and MEFs, and it was enriched for the cellular proliferation marker *Mki67* when compared to CTC-c. Multiple lineages are likely to have contributed to this complex grouping: CTCs from KPC mice with tumor-restricted, lineage-tagged EGFP expression clustered with CTC-pro (data not shown), noteworthy for abundant expression of *Mki67* and an annotated cell cycle signature in MSigDB (Whitfield et al., 2002) (data not shown). One single cell within the CTC-pro cluster was derived from the pancreatic cancer cell line NB508, while another (MP3-2) had high keratin/high E-cadherin expression characteristic of classical CTCs (data not shown). Nonetheless, another sub-cluster contained immune and dendritic cells, identified by their expression of antigen processing and presentation genes (GO:0019886 - antigen processing and presentation of exogenous peptide antigen via MHC class II; Table 5). Taken together, the CTC-pro cluster appears to represent a grouping of highly proliferative cells, of which a subset are tumor-derived.

**[00175]** Together, unbiased isolation and RNA sequencing evaluation of single pancreatic CTCs indicate that over half of these are nonviable with RNA at various stages of degradation. Among the remaining viable CTCs, three major classes are distinguishable by unsupervised clustering: the classical subset (CTC-c) accounts for 55%, with a second platelet adherent group (CTC-plt; 32%) and a third heterogeneous cluster marked by proliferative signatures (CTC-pro; 13%). Given their most clearly defined tumor-derived characteristics, we selected the CTC-c cluster for detailed analysis of metastasis-associated pathways.

**[00176]** *Pancreatic CTCs Co-express Epithelial, Mesenchymal, and Stem Cell Markers.* The relevance of EMT to early metastasis in pancreatic cancer has been supported by lineage tracing studies in the KPC mouse model (Rhim et al., 2012). In human breast cancer CTCs, a distribution of epithelial and mesenchymal markers within individual CTCs was recently reported by the inventors, reflecting both tumor histology and response or resistance to diverse therapies (Yu et al., 2013). To directly test for EMT in the mouse pancreatic CTCs, established epithelial (E) and mesenchymal (M) markers (Kalluri and Weinberg, 2009) were used to evaluate each cell within the CTC-c cluster (data not shown). Compared with the primary tumor, CTC-c cells demonstrated clear loss of the epithelial markers E-cadherin (*Cdh1*) and *Muc1*, whereas mesenchymal transcripts were mixed, with some showing increased expression (*Cdh11*, *Vim*) and others with reduced levels (*S100a4*, *Itga5*, *Sdc1*) (Figs. 3A and 3B). Notably, even the mesenchymal genes that were upregulated in CTCs showed a high degree of heterogeneous expression across single cells (data not shown). In contrast, loss of epithelial marks, including E-cadherin (*Cdh1*) was nearly universal across all classical CTCs.

**[00177]** CTCs are also thought to be enriched for metastatic precursors, capable of initiating metastatic tumor deposits. The relationship between such precursor cells and postulated cancer stem cells is uncertain, as is the relevance of established stem cell markers in identifying these cells. Proposed pancreatic cancer stem cell genes (Rasheed and Matsui, 2012; Rasheed et al., 2010) were

evaluated in the single cell RNA sequencing reads (Fig. 3B). Among all candidate markers tested (*Aldh1a1*, *Aldh1a2*, *Prom1/Cd133*, *Cd44*, *Met*, *EpCAM*), only *Aldh1a1* and *Aldh1a2* were enriched in CTCs. Classical CTCs expressed predominantly the *Aldh1a2* isoform, while CTC-plt cells were enriched for *Aldh1a1*, but these isoforms were also co-expressed within some single CTCs. MEFs, NB508 pancreatic cancer cells and normal WBCs also expressed *Aldh1a1*, but not *Aldh1a2* (data not shown). Within single CTCs, there was no correlation between expression of Aldh1 isoforms and enrichment for the mesenchymal genes *Cdh11* or *Vim*, suggesting that these two biomarkers are not intrinsically linked.

**[00178]** Given the identification of *Aldh1a2* as a potential stem-like marker expressed by CTCs, its expression within matched primary tumors was tested using RNA *in situ* hybridization (RNA-ISH). Expression patterns within tumors were heterogeneous: *Aldh1a2* expressing cells were primarily localized within the “stromal” or non-epithelial (i.e. keratin low) compartment of the tumor (data not shown). The origin of these non-epithelial cells, which are particularly abundant in pancreatic cancer, is likely to be mixed. Both histological evaluation and negative KRAS mutational analysis (Biankin et al., 2012; Ogino et al., 2005) in human pancreatic cancer have indicated that most of these cells represent reactive fibroblasts or stroma, rather than being of tumor origin. However, lineage tracing in KPC mice has recently shown that a small fraction of these supposedly stromal cells are in fact tumor-derived, presumably having undergone EMT to appear fibroblastic (Rhim et al., 2012). Interestingly, the mouse with the most metastases and the highest number of *Aldh1a2* positive CTCs, MP7, also had the primary tumor with the highest levels of *Aldh1a2*. In that case, *Aldh1a2*-positive cells were present diffusely in the stromal compartment, as well as comprising a small subpopulation of the epithelial (keratin high) component (data not shown). Thus, classical CTCs, which are keratin-high, express the stem cell-associated gene *Aldh1a2*, whose expression in primary tumors is restricted to the stromal (keratin low) compartment and only a small subpopulation of epithelial cells.

**[00179] *Classical CTCs Share Expression of Stromal Enriched Genes.*** Beside the evident diversity of CTCs, shared transcripts were sought that might provide further insight into their cell of origin within the primary tumor, the mechanisms by which they invade and survive within the bloodstream, and ultimately identify potential CTC-specific therapeutic targets. Rigorous criteria were selected to identify the most highly enriched CTC transcripts (RP score < 300), expressed at very high levels (>100 RPM) in ≥ 90% of all classical CTCs. Three genes met these criteria: Decorin (*Dcn*), a extracellular matrix proteoglycan expressed in tumor stroma across a variety of different cancers (Adany et al., 1990; Bostrom et al., 2013; Henke et al., 2012; Hunzelmann et al., 1995; Iozzo and Cohen, 1994; Mu et al., 2013; Nash et al., 2002); Insulin-like growth factor binding protein 5 (*Igfbp5*), an extracellular growth factor binding protein expressed in human PDAC reported to have both pro and anti-proliferative properties (Johnson et al., 2006; Johnson and Haun, 2009); and

Kruppel-like factor 4 (*Klf4*), one of the key stem cell (iPS) reprogramming factors (Takahashi and Yamanaka, 2006), which has been implicated in pancreatic cancer development (Brembeck and Rustgi, 2000; Prasad et al., 2005; Wei et al., 2010). By RNA-ISH, *Dcn* was expressed diffusely in the stromal elements of the tumor (Fig. 6). Remarkably, both *Igfbp5* and *Klf4* were expressed focally, predominantly within stromal-appearing cells that border the epithelial compartments of the tumor (data not shown). RNA-ISH of EGFP lineage restricted primary tumors confirmed that the *Igfbp5* positive cells at the epithelial/stromal interface are of tumor origin (data not shown). In addition to this transitional region, analysis of *Klf4* in this EGFP-tagged tumor also found expression in a subset of epithelial ducts (data not shown). Of note, while they are expressed in only a small subset of primary tumor cells, both *Igfbp5* and *Klf4* are highly co-expressed in 85% of all classical CTCs. Together with the mixed epithelial/mesenchymal markers evident in CTCs, these observations raise the possibility that many CTCs are derived from foci at the epithelial/stromal interface, that may be defined by *Igfbp5* and *Klf4* expression.

**[00180]** In addition to the three most highly expressed transcripts, CTCs were noteworthy for high level expression of genes implicated in stromal cell matrix. Gene ontology analysis of all CTC-enriched genes (Table 3) identified 60 extracellular proteins (GO:0044421, OR 1.7, q-value 6.4 x 10<sup>-3</sup>), of which 32 are found in proteinaceous extracellular matrix (ECM) (GO:0005578, OR 2.4, q-value 4.8 x 10<sup>-3</sup>). Recent studies have highlighted the importance of the reactive stroma to pancreatic cancer pathogenesis and metastasis (Feig et al., 2012; Neesse et al., 2013; Neesse et al., 2011; Olive et al., 2009; Provenzano et al., 2012), however, the expression of these stroma-associated ECM genes within tumor cells in circulation was unexpected. To identify the predominant stromal enriched genes in the mouse pancreatic tumor model, we performed RP differential expression analysis between the bulk tumor samples representing tumor cells mixed with reactive stromal cells versus purified EGFP-tagged single cells from the primary tumor (TuGMP3). A total of 51 proteinaceous ECM genes were enriched in bulk tumors versus single primary tumor cells (GO:0005578, OR 4.8, q-value 3.4 x 10<sup>-18</sup>). Of these, 6 genes (*Ccdc80*, *Col1a2*, *Col3a1*, *Dcn*, *Sparc*, *Timp2*) were shared with the previously identified CTC-enriched gene set (data not shown). Decorin (*Dcn*), as noted above, was identified as the most highly enriched (median 10,686 rpm) in CTCs with high level expression (>100 rpm) in 98% of CTCs. The second most abundant gene was *Sparc* (median 3,913 rpm) with high expression in 88% of CTCs. These two genes were co-expressed at high levels in 88% of classical CTCs. RNA-ISH of primary tumors for both *Dcn* (Fig. 6) and *Sparc* (data not shown) confirmed that these genes are expressed throughout the reactive stroma and are not present in the epithelial keratin-rich regions of primary tumors.

**[00181]** The expression of stromal-derived ECM genes is a common feature of all classical CTCs, yet a mouse-specific bias in distribution among these genes was evident, despite their identical *Kras/p53* genetic drivers. This mouse-specific clustering was evident in the unsupervised analysis (p-

value  $< 2.2 \times 10^{-16}$ ). For instance, sub-cluster 3 was over-represented with single CTCs from mouse MP6, while sub-cluster 4 was enriched for mouse MP7, and sub-cluster 5 for mouse MP2. Of 68 transcripts differentially expressed between the CTCs of mice MP2 and MP7 by RP analysis, gene ontology indicated significant enrichment for 11 extracellular proteins (GO:0044421, OR 3.8, q-value 0.06), 7 of which are found in proteinaceous ECM (GO:0005578, OR 6.3, q-value 0.05) (data not shown). Together, these data indicate that most CTCs derived from a mouse pancreatic cancer model express at high levels a set of ECM genes normally found in the stromal, rather than the epithelial compartment of the primary tumor. This may reflect the origin of many CTCs at the epithelial/stromal interface, consistent with their expression of uniquely restricted markers such as *Igfbp5* and *Klf4*. The fact that individual genetically matched mouse tumors generate CTCs with both shared and unique patterns of ECM gene expression suggests tumor-specific invasion pathways that are superimposed upon fundamental characteristics of CTCs. The high levels of extracellular proteins expressed by CTCs provide unexpected opportunities for targeting these metastatic precursors.

**[00182] *Human Pancreatic CTCs Express the ECM Protein SPARC.*** To determine the relevance of ECM protein expression to human disease, CTCs were isolated from the blood of metastatic PDAC patients and subjected to single cell RNA-sequencing. Analysis of 7 pancreatic CTCs from 3 patients revealed that the majority expressed keratins defining their epithelial origins and a total of 13 of 60 extracellular protein genes enriched in mouse CTCs were expressed at high levels ( $>100$  rpm) in at least one human pancreatic CTC (Fig. 7). Human *SPARC* was the only gene found at high levels in all human pancreatic CTCs. Analysis of human prostate and breast CTCs also show significant expression of extracellular proteins including SPARC highlighting that these targets are commonly shared in metastatic epithelial cancer cells (data not shown). RNA-ISH of *Sparc/SPARC* in both mouse and human PDAC found expression confined primarily to the stromal compartment of tumors (data not shown). *SPARC* expression was found in 196/198 (99%) human primary PDAC tumors and 36% of positive tumors had some detectable *SPARC* in epithelial tumor cells albeit the minority of the overall signal. The presence of SPARC as an extracellular protein permits antibody directed therapies that target SPARC. Together these data indicate that findings in mouse pancreatic CTCs can be found in human disease and offer both novel biomarkers and therapeutic targets.

**[00183] **Discussion****

**[00184]** Described herein is a detailed analysis of CTC composition and diversity, using single cell RNA sequencing. In total, high quality transcriptomes were achieved in 93 single mouse pancreatic CTCs, which were compared with 20 single cells from matched primary tumors, as well as bulk tumor preparations, and with 16 cells from an immortalized cell line established from the same mouse pancreatic tumor model. The use of a mouse model, which closely matches human PDAC, made it possible to compare primary tumor specimens isolated simultaneously with the CTCs. Given

the shared *Kras/Trp53* genetic drivers in the KPC mouse model, it was also possible to examine CTC heterogeneity within individual mice and across different animals. Finally, the use of the CTC-iChip technology enabled the selection of untagged CTCs, irrespective of their cell surface epitopes, thus avoiding any bias associated with tumor marker-specific cell purification. Together, these observations include the following: 1. CTCs cluster into multiple subsets, including a major “classical CTC” group, and others that are marked by platelet-derived markers or proliferative signatures; 2. While individual mouse tumors may produce CTCs that fit into each of these clusters, there are unique patterns to CTCs derived from individual mice, despite their shared genetic drivers; 3. Common markers shared by virtually all classical CTCs include both epithelial and mesenchymal markers, the *Aldh1a2* stem cell marker, and two highly expressed transcripts (*Igfbp5* and *Klf4*) that identify foci localized to the epithelial/stromal boundary of primary tumors; and 4. The most highly enriched CTC-specific transcripts shared by almost all classical CTCs encode extracellular matrix proteins associated with the tumor stromal compartment.

**[00185]** Compared with previous RNA sequencing of partially purified, bulk CTC populations, the single cell analysis reported here provides considerably more depth of tumor cell-specific reads. As such, the detailed analysis of classical CTCs from the mouse pancreatic cancer model is unprecedented. It is demonstrated herein that pancreatic cancer CTCs uniformly lose expression of the epithelial marker E-cadherin (*Cdh1*), a key feature of epithelial-to-mesenchymal transition. However, the cells do not lose expression of other epithelial markers, such as cytokeratins, nor is there a consistent increase in classical EMT mesenchymal markers such as vimentin. As such, most classical CTCs appear arrested in a biphenotypic state. Despite their expression of cytokeratins (present in the epithelial components of the primary tumor), most other highly expressed markers in CTCs were shared with the non-epithelial or “stromal” component of the primary tumor. Among these stromal genes expressed in classical CTCs is *Aldh1a2*, a putative pancreatic cancer stem cell marker (Rasheed and Matsui, 2012; Rasheed et al., 2010). Whether *Aldh1a2* is a functionally significant marker of cellular plasticity in metastatic precursors remains to be determined.

**[00186]** A provocative observation relating to the shared epithelial and mesenchymal state of classical CTCs is their virtually uniform (>85%) high level co-expression of *Igfbp5* and *Klf4*, two genes that are only expressed in a small subpopulation of cells at the epithelial/stromal interface within primary tumors. This raises the intriguing possibility that this critical location within the tumor generates a disproportionate fraction of viable CTCs. Indeed, tumor cells that are actively undergoing EMT are presumably enriched at the epithelial-stromal function, contributing to the mixed lineage of the tumor stroma, with both tumor-derived and non-malignant reactive cell types. The potential roles of both IGF signaling and Klf4 transcriptional regulation in embryonic development and pancreatic malignancy make their unique expression pattern in both tumors and CTCs particularly noteworthy.

**[00187]** Finally, the most unexpected observation from this single CTC RNA sequencing study is the very high level abundance of ECM proteins on the vast majority of classical CTCs. Notably, prior evaluation of matched primary and metastatic breast tumors identified the most prevalent gene expression difference as enrichment for ECM molecules in the metastases, comprising some 18% of differentially expressed genes (Weigelt et al., 2005). While this has been interpreted as reflecting differences in the local environment of the metastatic site, the present data indicate that ECM proteins are highly expressed by CTCs themselves. By analogy with the classical “seed versus soil” debate (Fidler, 2003), CTCs may in fact be seeds carrying some of their own soil.

**[00188]** The ultimate goal of detailed molecular analysis of CTCs is to understand the process by which they are generated and their therapeutic vulnerabilities. In this regard, an important observation derived from the present single CTC RNA sequencing analysis is the unexpected expression of extracellular proteins with a preponderance of proteins found in ECM. Two of the most abundant and commonly shared ECM proteins in CTCs are *Dcn* and *Sparc*, both of which are established tumor stromal genes. Notably, *Sparc* expressing stroma appears to bind albumin-conjugated chemotherapy-containing nanoparticles (nab-paclitaxel) allowing for increased cytotoxicity and efficacy in human PDAC (Neuzillet et al., 2013; Von Hoff et al., 2011; Yardley, 2013). Indeed, considerable effort has been directed to targeting pancreatic cancer stroma as a means of improving delivery of chemotherapeutics and stripping tumor cells of their supportive microenvironment (Neesse et al., 2011; Olive et al., 2009; Provenzano et al., 2012; Rasheed et al., 2012). The finding that these gene products are also expressed by CTCs indicates that antibody-directed therapies can be used not only against primary tumor stroma, but also to target tumor cells as they transit in the blood.

**[00189]** As described herein, the present CTC analyses to extend from matching them to known tumor-defining markers to interrogating them for unique properties that distinguish them from most primary tumor cells and may underlie their ability to survive in the bloodstream and generate distant metastases. Such insights into the cellular process of human cancer metastasis are critical to the goal of ultimately preventing the spread of a primary tumor to distant organs.

#### **[00190] Experimental Procedures**

**[00191]** *Mice and cell lines.* Mice with pancreatic cancer used in these experiments express Cre driven by *Pdx1*, *LSL-Kras*<sup>G12D</sup>, and *Trp53*<sup>lox/+</sup> or *Trp53*<sup>lox/lox</sup> as previously described (Bardeesy et al., 2006). EGFP pancreatic lineage tagged KPC mice were generated by breeding the mT/mG mouse (Jackson Laboratory - Gt(ROSA)26Sortm4(ACTB-tdTomato,-EGFP)Luo/J) into the breeder pairs used for KPC mouse generation. Normal FVB mice were purchased from Jackson Laboratory. All mice care and procedures were done under MGH SRAC approved protocols.

**[00192]** *Adaptation of CTC Enrichment Technology.* Given the desire for an unbiased enrichment system, the previously presented negative depletion technology was selected for this application

(Ozkumur et al., 2013). All processing protocols were identical to those previously identified, except a rat anti-mouse CD45 antibody (BAM114, R&D Systems, USA) was conjugated to MyOne beads.

**[00193]** *Single cell Micromanipulation, Amplification, and Sequencing.* After whole blood anti-CD45 negative depletion, the product containing enriched cells was collected in a 35mm petri dish and viewed using a Nikon Eclipse Ti<sup>TM</sup> inverted fluorescent microscope. Cells of interest were identified based on intact cellular morphology and lack of labeling with anti-CD45 magnetic beads. These target cells were individually micromanipulated with a 10  $\mu$ m transfer tip on an Eppendorf TransferMan<sup>®</sup> NK 2 micromanipulator and ejected into PCR tubes containing RNA protective lysis buffer and immediately flash frozen in liquid nitrogen. Single cells were amplified with a modified protocol (Tang et al., 2010) and sequenced on the ABI 5500XL<sup>TM</sup> system.

**[00194]** *RNA in situ Hybridization (RNA-ISH).* RNA-ISH was performed according to the Affymetrix QuantiGene ViewRNA ISH Tissue-2 Plex Assay<sup>TM</sup>.

**[00195] References**

Adany, R., Heimer, R., Caterson, B., Sorrell, J.M., and Iozzo, R.V. (1990). Altered expression of chondroitin sulfate proteoglycan in the stroma of human colon carcinoma. Hypomethylation of PG-40 gene correlates with increased PG-40 content and mRNA levels. *The Journal of biological chemistry* 265, 11389-11396.

Anastas, J.N., and Moon, R.T. (2013). WNT signalling pathways as therapeutic targets in cancer. *Nat Rev Cancer* 13, 11-26.

Bardeesy, N., Aguirre, A.J., Chu, G.C., Cheng, K.H., Lopez, L.V., Hezel, A.F., Feng, B., Brennan, C., Weissleder, R., Mahmood, U., et al. (2006). Both p16(INK4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* 103, 5947-5952.

Biankin, A.V., Waddell, N., Kassahn, K.S., Gingras, M.C., Muthuswamy, L.B., Johns, A.L., Miller, D.K., Wilson, P.J., Patch, A.M., Wu, J., et al. (2012). Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. *Nature* 491, 399-405.

Bostrom, P., Sainio, A., Kakko, T., Savontaus, M., Soderstrom, M., and Jarvelainen, H. (2013). Localization of decorin gene expression in normal human breast tissue and in benign and malignant tumors of the human breast. *Histochemistry and cell biology* 139, 161-171.

Breitling, R., Armengaud, P., Amtmann, A., and Herzyk, P. (2004). Rank products: a simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS letters* 573, 83-92.

Brembeck, F.H., and Rustgi, A.K. (2000). The tissue-dependent keratin 19 gene transcription is regulated by GKL/KLF4 and Sp1. *The Journal of biological chemistry* 275, 28230-28239.

Cann, G.M., Gulzar, Z.G., Cooper, S., Li, R., Luo, S., Tat, M., Stuart, S., Schroth, G., Srinivas, S., Ronaghi, M., et al. (2012). mRNA-Seq of single prostate cancer circulating tumor cells reveals recapitulation of gene expression and pathways found in prostate cancer. *PLoS One* 7, e49144.

Carmeliet, P., and Jain, R.K. (2011). Molecular mechanisms and clinical applications of angiogenesis. *Nature* *473*, 298-307.

Chen, C.L., Mahalingam, D., Osmulski, P., Jadhav, R.R., Wang, C.M., Leach, R.J., Chang, T.C., Weitman, S.D., Kumar, A.P., Sun, L., *et al.* (2013). Single-cell analysis of circulating tumor cells identifies cumulative expression patterns of EMT-related genes in metastatic prostate cancer. *The Prostate* *73*, 813-826.

Clevers, H., and Nusse, R. (2012). Wnt/beta-catenin signaling and disease. *Cell* *149*, 1192-1205.

Feig, C., Gopinathan, A., Neesse, A., Chan, D.S., Cook, N., and Tuveson, D.A. (2012). The pancreas cancer microenvironment. *Clin Cancer Res* *18*, 4266-4276.

Fidler, I.J. (2003). The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat Rev Cancer* *3*, 453-458.

Folkman, J. (1995). Angiogenesis in cancer, vascular, rheumatoid and other disease. *Nature medicine* *1*, 27-31.

Henke, A., Grace, O.C., Ashley, G.R., Stewart, G.D., Riddick, A.C., Yeun, H., O'Donnell, M., Anderson, R.A., and Thomson, A.A. (2012). Stromal expression of decorin, Semaphorin6D, SPARC, Sprouty1 and Tsukushi in developing prostate and decreased levels of decorin in prostate cancer. *PLoS One* *7*, e42516.

Hunzelmann, N., Schonherr, E., Bonnekoh, B., Hartmann, C., Kresse, H., and Krieg, T. (1995). Altered immunohistochemical expression of small proteoglycans in the tumor tissue and stroma of basal cell carcinoma. *The Journal of investigative dermatology* *104*, 509-513.

Ikushima, H., and Miyazono, K. (2010). TGFbeta signalling: a complex web in cancer progression. *Nat Rev Cancer* *10*, 415-424.

Iozzo, R.V., and Cohen, I. (1994). Altered proteoglycan gene expression and the tumor stroma. *Exs* *70*, 199-214.

Johnson, S.K., Dennis, R.A., Barone, G.W., Lamps, L.W., and Haun, R.S. (2006). Differential expression of insulin-like growth factor binding protein-5 in pancreatic adenocarcinomas: identification using DNA microarray. *Molecular carcinogenesis* *45*, 814-827.

Johnson, S.K., and Haun, R.S. (2009). Insulin-like growth factor binding protein-5 influences pancreatic cancer cell growth. *World journal of gastroenterology : WJG* *15*, 3355-3366.

Kalluri, R., and Weinberg, R.A. (2009). The basics of epithelial-mesenchymal transition. *J Clin Invest* *119*, 1420-1428.

Katoh, M., and Katoh, M. (2007). WNT signaling pathway and stem cell signaling network. *Clin Cancer Res* *13*, 4042-4045.

Krause, D.S., Ito, T., Fackler, M.J., Smith, O.M., Collector, M.I., Sharkis, S.J., and May, W.S. (1994). Characterization of murine CD34, a marker for hematopoietic progenitor and stem cells. *Blood* *84*, 691-701.

Labelle, M., Begum, S., and Hynes, R.O. (2011). Direct signaling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis. *Cancer Cell* 20, 576-590.

Miknyoczki, S.J., Klein-Szanto, A.J., and Ruggeri, B.A. (1996). Neurotrophin-Trk receptor interactions in neoplasia: a possible role in interstitial and perineural invasion in ductal pancreatic cancer. *Critical reviews in oncogenesis* 7, 89-100.

Miknyoczki, S.J., Lang, D., Huang, L., Klein-Szanto, A.J., Dionne, C.A., and Ruggeri, B.A. (1999). Neurotrophins and Trk receptors in human pancreatic ductal adenocarcinoma: expression patterns and effects on in vitro invasive behavior. *International journal of cancer Journal international du cancer* 81, 417-427.

Mu, Y., Chen, Y., Zhang, G., Zhan, X., Li, Y., Liu, T., Li, G., Li, M., Xiao, Z., Gong, X., *et al.* (2013). Identification of stromal differentially expressed proteins in the colon carcinoma by quantitative proteomics. *Electrophoresis* 34, 1679-1692.

Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., and Luo, L. (2007). A global double-fluorescent Cre reporter mouse. *Genesis* 45, 593-605.

Nash, M.A., Deavers, M.T., and Freedman, R.S. (2002). The expression of decorin in human ovarian tumors. *Clin Cancer Res* 8, 1754-1760.

Neesse, A., Frese, K.K., Bapiro, T.E., Nakagawa, T., Sternlicht, M.D., Seeley, T.W., Pilarsky, C., Jodrell, D.I., Spong, S.M., and Tuveson, D.A. (2013). CTGF antagonism with mAb FG-3019 enhances chemotherapy response without increasing drug delivery in murine ductal pancreas cancer. *Proc Natl Acad Sci U S A* 110, 12325-12330.

Neesse, A., Michl, P., Frese, K.K., Feig, C., Cook, N., Jacobetz, M.A., Lolkema, M.P., Buchholz, M., Olive, K.P., Gress, T.M., *et al.* (2011). Stromal biology and therapy in pancreatic cancer. *Gut* 60, 861-868.

Neuzillet, C., Tijeras-Raballand, A., Cros, J., Faivre, S., Hammel, P., and Raymond, E. (2013). Stromal expression of SPARC in pancreatic adenocarcinoma. *Cancer metastasis reviews*.

Ogino, S., Kawasaki, T., Brahmandam, M., Yan, L., Cantor, M., Namgyal, C., Mino-Kenudson, M., Lauwers, G.Y., Loda, M., and Fuchs, C.S. (2005). Sensitive sequencing method for KRAS mutation detection by Pyrosequencing. *The Journal of molecular diagnostics : JMD* 7, 413-421.

Ohta, T., Numata, M., Tsukioka, Y., Futagami, F., Kayahara, M., Kitagawa, H., Nagakawa, T., Yamamoto, M., Wakayama, T., Kitamura, Y., *et al.* (1997). Neurotrophin-3 expression in human pancreatic cancers. *The Journal of pathology* 181, 405-412.

Olive, K.P., Jacobetz, M.A., Davidson, C.J., Gopinathan, A., McIntyre, D., Honess, D., Madhu, B., Goldgraben, M.A., Caldwell, M.E., Allard, D., *et al.* (2009). Inhibition of Hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. *Science* 324, 1457-1461.

Ozkumur, E., Shah, A.M., Ciciliano, J.C., Emmink, B.L., Miyamoto, D.T., Brachtel, E., Yu, M., Chen, P.I., Morgan, B., Trautwein, J., *et al.* (2013). Inertial focusing for tumor antigen-dependent and -independent sorting of rare circulating tumor cells. *Sci Transl Med* 5, 179ra147.

Pantel, K., Brakenhoff, R.H., and Brandt, B. (2008). Detection, clinical relevance and specific biological properties of disseminating tumour cells. *Nat Rev Cancer* 8, 329-340.

Powell, A.A., Talasaz, A.H., Zhang, H., Coram, M.A., Reddy, A., Deng, G., Telli, M.L., Advani, R.H., Carlson, R.W., Mollick, J.A., *et al.* (2012). Single cell profiling of circulating tumor cells: transcriptional heterogeneity and diversity from breast cancer cell lines. *PLoS One* 7, e33788.

Prasad, N.B., Biankin, A.V., Fukushima, N., Maitra, A., Dhara, S., Elkahloun, A.G., Hruban, R.H., Goggins, M., and Leach, S.D. (2005). Gene expression profiles in pancreatic intraepithelial neoplasia reflect the effects of Hedgehog signaling on pancreatic ductal epithelial cells. *Cancer Res* 65, 1619-1626.

Provenzano, P.P., Cuevas, C., Chang, A.E., Goel, V.K., Von Hoff, D.D., and Hingorani, S.R. (2012). Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma. *Cancer Cell* 21, 418-429.

Ramskold, D., Luo, S., Wang, Y.C., Li, R., Deng, Q., Faridani, O.R., Daniels, G.A., Khrebtukova, I., Loring, J.F., Laurent, L.C., *et al.* (2012). Full-length mRNA-Seq from single-cell levels of RNA and individual circulating tumor cells. *Nat Biotechnol* 30, 777-782.

Rasheed, Z.A., and Matsui, W. (2012). Biological and clinical relevance of stem cells in pancreatic adenocarcinoma. *Journal of gastroenterology and hepatology* 27 Suppl 2, 15-18.

Rasheed, Z.A., Matsui, W., and Maitra, A. (2012). Pathology of pancreatic stroma in PDAC. In *Pancreatic Cancer and Tumor Microenvironment*, P.J. Grippo, and H.G. Munshi, eds. (Trivandrum (India)).

Rasheed, Z.A., Yang, J., Wang, Q., Kowalski, J., Freed, I., Murter, C., Hong, S.M., Koorstra, J.B., Rajeshkumar, N.V., He, X., *et al.* (2010). Prognostic significance of tumorigenic cells with mesenchymal features in pancreatic adenocarcinoma. *J Natl Cancer Inst* 102, 340-351.

Rhim, A.D., Mirek, E.T., Aiello, N.M., Maitra, A., Bailey, J.M., McAllister, F., Reichert, M., Beatty, G.L., Rustgi, A.K., Vonderheide, R.H., *et al.* (2012). EMT and dissemination precede pancreatic tumor formation. *Cell* 148, 349-361.

Sergeant, G., van Eijnsden, R., Roskams, T., Van Duppen, V., and Topal, B. (2012). Pancreatic cancer circulating tumour cells express a cell motility gene signature that predicts survival after surgery. *BMC cancer* 12, 527.

Siegel, P.M., and Massague, J. (2003). Cytostatic and apoptotic actions of TGF-beta in homeostasis and cancer. *Nat Rev Cancer* 3, 807-821.

Society, A.C. (2013). *Cancer Facts & Figures 2013* (Atlanta: American Cancer Society).

Takahashi, K., and Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126, 663-676.

Tang, F., Barbacioru, C., Nordman, E., Li, B., Xu, N., Bashkirov, V.I., Lao, K., and Surani, M.A. (2010). RNA-Seq analysis to capture the transcriptome landscape of a single cell. *Nat Protoc* 5, 516-535.

Von Hoff, D.D., Ramanathan, R.K., Borad, M.J., Laheru, D.A., Smith, L.S., Wood, T.E., Korn, R.L., Desai, N., Trieu, V., Iglesias, J.L., *et al.* (2011). Gemcitabine plus nab-paclitaxel is an active regimen in patients with advanced pancreatic cancer: a phase I/II trial. *J Clin Oncol* 29, 4548-4554.

Wang, W., Zhao, H., Zhang, S., Kang, E., Chen, Y., Ni, C., Zhang, S., and Zhu, M. (2009). Patterns of expression and function of the p75(NGFR) protein in pancreatic cancer cells and tumours. *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology* 35, 826-832.

Wei, D., Wang, L., Kanai, M., Jia, Z., Le, X., Li, Q., Wang, H., and Xie, K. (2010). KLF4alpha up-regulation promotes cell cycle progression and reduces survival time of patients with pancreatic cancer. *Gastroenterology* 139, 2135-2145.

Weigelt, B., Wessels, L.F., Bosma, A.J., Glas, A.M., Nuyten, D.S., He, Y.D., Dai, H., Peterse, J.L., and van't Veer, L.J. (2005). No common denominator for breast cancer lymph node metastasis. *Br J Cancer* 93, 924-932.

Welty, C.J., Coleman, I., Coleman, R., Lakely, B., Xia, J., Chen, S., Gulati, R., Larson, S.R., Lange, P.H., Montgomery, B., *et al.* (2013). Single cell transcriptomic analysis of prostate cancer cells. *BMC molecular biology* 14, 6.

Whitfield, M.L., Sherlock, G., Saldanha, A.J., Murray, J.I., Ball, C.A., Alexander, K.E., Matese, J.C., Perou, C.M., Hurt, M.M., Brown, P.O., *et al.* (2002). Identification of genes periodically expressed in the human cell cycle and their expression in tumors. *Molecular biology of the cell* 13, 1977-2000.

Yardley, D.A. (2013). nab-Paclitaxel mechanisms of action and delivery. *Journal of controlled release : official journal of the Controlled Release Society* 170, 365-372.

Yu, M., Stott, S., Toner, M., Maheswaran, S., and Haber, D.A. (2011). Circulating tumor cells: approaches to isolation and characterization. *J Cell Biol* 192, 373-382.

Yu, M., Ting, D.T., Stott, S.L., Wittner, B.S., Ozsolak, F., Paul, S., Ciciliano, J.C., Smas, M.E., Winokur, D., Gilman, A.J., *et al.* (2012). RNA sequencing of pancreatic circulating tumour cells implicates WNT signalling in metastasis. *Nature* 487, 510-513.

Zhang, Y., Dang, C., Ma, Q., and Shimahara, Y. (2005). Expression of nerve growth factor receptors and their prognostic value in human pancreatic cancer. *Oncology reports* 14, 161-171.

**[00196]** **Table 1:** Annotation of CTC enriched genes in KEGG defined signaling pathways. \* indicates gene found in multiple pathway gene sets.

MAPK Pathway	WNT Pathway	Neurotropin Pathway	TGF-beta Pathway	Toll-Like Receptor Pathway	VEGF Pathway
1500003o03rik*	Jund	1500003o03rik*	Akt2*	Amhr2	Akt2*
Akt2*	Map3k3*	Crebbp*	Braf*	Crebbp*	Fos*
B230120h23rik	Mapk1*	Csnk1a1	Calm1	Dcn	Ifnar2
Braf*	Mapkapk3*	Jun*	Calm2	Id1	Irak4*
Dusp1	Mef2c	Nkd1	Irak4*	Id2	Irf7
Dusp14	Mras	Ppp3ca*	Irs2	Mapk1*	Jun*
Dusp3	Nf1	Rock1*	Jun*	Rock1*	Mapk1*
Fas	Nfkb2	Rock2*	Maged1	Rock2*	Nfkbia*
Fgf1	Nr4a1	Siah1a	Map3k3*	Smad4*	Tirap
Flnc	Pla2g4a*	Smad4*	Mapk1*	Tgfb2*	Tlr2
Fos*	Ppp3ca*	Tbl1x	Nfkbia*	Tgfbr2*	
Gadd45b	Rras2	Tcf7l1	Shc1	Thbs1	
Hspa2	Tgfb2*	Wnt4	Ywhaz		
Hspb1*	Tgfbr2*				
Jun*					

[00197] **Table 2:** Significantly Expressed Genes by Rank Product (FDR < 0.01)

Count	CTC-c vs Primary Tumor Enriched Gene	Primary Tumor vs CTC-c Enriched Gene	CTC-plt vs CTC-c	CTC-pro vs CTC-c
1	Upk3b	Tff2	Clec1b	kg:uc007pge.1
2	Ier2	Wfdc2	AU023871	kg:uc007pgd.1
3	Egr1	Lamb3	Alox12	kg:uc007pgf.1
4	Nkain4	Lad1	Itga2b	kg:uc007pgg.1
5	Igfbp5	Dmbt1	Ppbp	Igj
6	Slc6a4	Npy	Gng11	kg:uc012enb.1
7	Klf4	Pmepa1	Vwf	2010001M09Rik
8	Tmem221	Kcnn4	Pf4	kg:uc009cfw.1
9	Arl4d	Serinc2	Fcer1g	kg:uc007pgi.1
10	Lrrn4	5730559C18Rik	Tmem40	kg:uc007pgh.1
11	Cldn15	Muc1	Hba-a2	kg:uc007yos.1
12	Gpm6a	Chi3l3	Stom	Coro1a
13	Atf3	Pglyrp1	Beta-s	Pou2af1
14	Ptma	Arl4c	Plek	kg:uc011yvj.1
15	Slc9a3r1	Spp1	Srgn	Glipr1
16	Fos	Col15a1	Myl9	Cd52
17	Tmem119	C1qb	Cd84	Cd79b
18	Ptgis	Tnnt2	F5	Sec11c
19	Dcn	Gkn3	Trem1	Tnfrsf17
20	Gbp2	Onecut2	Hbb-b1	Krr1
21	Dmkn	Mmp7	Itgb3	Gmfg

22	Sdc4	Cd74	Gp9	Ccr9
23	Ildr2	Ctss	Mpl	Pycard
24	Akap2	Lame2	Ctla2a	Derl3
25	Gfpt2	Olfml3	Tubb1	Rac2
26	Klf6	Lgals4	Mylk	Srgn
27	Btg2	Lcn2	F13a1	Cytip
28	Myl7	Ly6a	Slamf1	Edem2
29	Igfbp6	Pak1	Rgs10	Itgb7
30	Gpr133	Capn5	Mkrn1	Lsp1
31	Oasl2	Ptprn	Laptm5	Lcp1
32	Pfn1	Reg3b	1810058I24Rik	Cyfip2
33	Cap1	Fmn13	Itgb2	Nans
34	Nfkbia	Sdc1	Slc2a3	Slamf7
35	Malat1	Prom1	Pcmt1	Ell2
36	Rarres2	Ankrd50	Gp5	H2-Eb1
37	Rspo1	Ccl6	Ube2o	Creld2
38	Espn	Slc4a11	5430417L22Rik	Cd74
39	Klf9	Oraov1	Ptpn18	Blnk
40	Zbtb7c	Aldh1l1	Lat	Fmn11
41	Brd2	Slc20a1	Fermt3	Snrnp70
42	Olfr1033	Cldn7	Nrgn	Sec61b
43	Wt1	Acsbg1	Mrv1	Edem1
44	Esam	Las11	Lyz2	Tspan13
45	kg:uc009igb.1	C1qc	Epb4.1	Psmb8
46	Tmem151a	Lama5	Rasgrp2	Pim1
47	Mgll	Mgat4a	Trem12	Sept1
48	Csrnp1	Cldn2	Hist1h4i	Cd48
49	Cd9	Mept2	March2	Sub1
50	Gjb5	Fxyd3	Ltbp1	Lims1
51	Lrrc61	Il4ra	Nptn	Ncoa2
52	Wasf2	Itga5	Abtb1	Ctnnb1
53	Pdpn	Porcn	Ctla2b	Fdps
54	kg:uc009ogv.1	Mast3	Prkab2	Ube2j1
55	Sdpr	Scara3	Arhgdb	Mettl1
56	Gpr64	Atox1	Alas2	Lax1
57	Flnc	Arrdc1	Odc1	Rilpl2
58	Add3	Mmp2	Ptpn11	Ctse
59	Gata6	Saa3	Dhcr24	Glrx
60	Wfdc1	Serpinf1	Mfsd2b	Fut8
61	A130040M12Rik	Sox11	Gp1bb	AI662270
62	Ankrd12	Prpsap1	Rbpms2	Gramd3
63	Adamts11	Mcpt1	Fyb	Il2rg

64	C2	Mfge8	Smox	Rasgrp3
65	Prss23	Col18a1	P2rx1	Impdh1
66	Ube2v1	Lyz2	Otud7b	Plek
67	Cryab	C1qa	kg:uc007ttx.1	Ints5
68	Pkhd1l1	Acp5	Samd14	Blmh
69	Rtn1	Angptl4	Clca1	Dnmt1
70	Birc6	Ccnd1	kg:uc007tty.1	Galk1
71	Xdh	Asl	Gpr56	kg:uc007hxv.1
72	Cd34	Ctxn1	Sh3bgrl2	Cedc88b
73	Rab6b	Pgs1	Pttg1ip	Selplg
74	Dusp1	Anapc2	Nomo1	Sar1b
75	Clic4	Cp	Gnaz	Lat2
76	C3	Gpx3	Mmrn1	Slc16a6
77	Rhob	Lama3	Gp1ba	Mki67
78	Mir3064	Rbp1	Sh3bgrl3	Dnajc3
79	Thbd	Cotl1	Slc24a3	H2-Ab1
80	Dpysl2	Nek6	Sord	Ndufs6
81	Cobl	Cpxm1	Nfe2	Actr3
82	Npr1	Sfrp1	Tuba4a	Etnk1
83	Dnajb9	Ttr	Zyx	Herpud1
84	Arhgap29	Gsto1	Cnn2	Ptpn7
85	Cav1	Npepl1	Itgb5	Ctss
86	Gbp7	Usmg5	Gata1	Cs
87	Hes1	Polr2l	Hist1h1c	Fbxw7
88	Gm16897	Sphk1	Tbxas1	Ppp2r5c
89	Ppp1r12a	Asxl1	Ptplad2	Znrd1
90	Sv2a	Ctsh	Bpgm	Rfc2
91	Ang	Egfl7	Pdlim7	Preb
92	Aldh1a2	C1qtnf6	Mmd	Fcer1g
93	Cry11	Rras	G6b	Dnajb11
94	Kank1	Lgi4	kg:uc009duo.1	Slc35b1
95	2210403K04Rik	Hmga2	Lyz1	Sin3b
96	kg:uc009okn.1	Cep250	Tacc1	Nktr
97	Osr1	B4galt3	Dap	
98	kg:uc008ewj.2	Tmem223	Mast2	
99	kg:uc009tuw.1	Ltbp2	Atp2a3	
100	Gadd45b	Tnfrsf23	Snca	
101	Ablim3	Col7a1	Stx11	
102	Clec3b	Ggct	C030046I01Rik	
103	Usp25	Rab25	Trpt1	
104	Sntb2	Nedd8	Tsc22d1	
105	Rock2	9430023L20Rik	Prkar2b	

106	Col14a1	Arl2	Cd9	
107	Cd200	Wbp1	Pgm2l1	
108	kg:uc008ehr.1	H2-Ab1	Gp6	
109	Atp2b1	Preb	Pde5a	
110	Exoc4	Sgsm3	Itga6	
111	Abcb1b	Sfn	Itgal	
112	Nrgn	Prrx2	Edem1	
113	kg:uc009cvm.1	Ptpk	Isg20	
114	Ncoa4	Reg1	Cdc42ep5	
115	Ndufa4	Sdcbp2	Nipal3	
116	Upk1b	Pcbd1	Ccdc92	
117	Jun	Slc25a1	Sort1	
118	Syne2	Vamp5	Ly6g6c	
119	kg:uc007bvx.1	Crlf1	Ubash3b	
120	Ap4e1	Avil	Inf2	
121	Spock2	2700094K13Rik	Asap1	
122	Efemp1	Ctse	Sec11c	
123	Prpf40a	Penk	Gas211	
124	Tspan5	Tmc4	Parvb	
125	Lgals7	Dhrs3	Tmsb4x	
126	Kif5b	Ap1s1	kg:uc007xrw.1	
127	Psip1	Arl6ip4	Nudt3	
128	kg:uc008oki.1	9430008C03Rik	Bcl2l1	
129	1810014B01Rik	Fcer1g	B230312A22Rik	
130	Ptges3	Uqcr11	Cnp	
131	Limch1	Nhp2	Plp1	
132	Bicd1	Plbd2	Cnst	
133	Rdx	Capg	Rgs18	
134	Pcdh15	Pnpla6	Lsm12	
135	Foxn3	Ppdpf	Alox5ap	
136	Morf4l2	Hgfac	Ppif	
137	Ppp1r15a	Apoe	Spnb1	
138	Cdc42ep3	Fam40a	Ormdl3	
139	Pard3b	Lyz1	Hpse	
140	Bicc1	2200002D01Rik	Srxn1	
141	Amhr2	Laptm5	2010002N04Rik	
142	Gucy1a3	Qars	Hist1h2bc	
143	Psmb2	Tmx2	Cyba	
144	Mapkapk3	Fkbp4	Chst12	
145	Ube2l6	Plin2	kg:uc009sps.1	
146	kg:uc007pff.1	Fcgr3	Max	
147	kg:uc007ctp.1	Gkn1	Was	

148	Nedd4	Snhg1	Isca1	
149	Plxna4	Lsp1	Pdzk1ip1	
150	2010107G12Rik	Gm20605	Lyn	
151	Ifngr1	Ly6c1	Mob3a	
152	Bcam	Aim1	H2-T24	
153	Ccnl1	2310007B03Rik	Slc44a1	
154	Hoxa5	Tgfb1	Derl1	
155	Fhl1	Tsta3	Gclm	
156	1810041L15Rik	Pafah1b3	Fech	
157	2900002K06Rik	Chid1	Ywhah	
158	Hspb1	Smox	Igtp	
159	Podn	1500012F01Rik	Myl6	
160	Fam63b	Tspan4	Thbs1	
161	Hsp90b1	Agrn	Tln1	
162	Dpp4	Cfp	kg:uc009apq.1	
163	Gas1	Cdh1	Bcap31	
164	kg:uc007zak.1	Rasgrf1	Ilk	
165	Zc3h13	Nxfl	Epha1	
166	Sox6	Pdrg1	2810453I06Rik	
167	Arid4a	Polr2j	Rnf19b	
168	Tnxb	Suds3	Gsn	
169	Tsix	D0H4S114	Flna	
170	Scd1	Ccl9	Arrb1	
171	Jund	Neat1	kg:uc007pum.1	
172	Crls1	Ccdc12	Mbnl1	
173	1110003E01Rik	Prr24	Ccnd3	
174	Rnase4	Impdh1	Pdlim1	
175	Arhgef12	Card10	Ctse	
176	Irf7	Cpsf1	Tspan17	
177	Bbx	Sema4g	Gpx4	
178	Sema5a	Hes6	Bnip3l	
179	Mau2	C130074G19Rik	P2ry12	
180	Abi3bp	Ctrb1	kg:uc009vev.1	
181	Dag1	Rnaseh2a	Prkab1	
182	Cyp2s1	Golm1	F2rl2	
183	Sfrs18	Ctsz	Stk4	
184	Hspb8	Cyb561	Fhl1	
185	Cnot6l	Ndufs8	Rnf10	
186	Twsg1	Atp6ap1	Rasa3	
187	Gpc3	Srd5a1	Taldo1	
188	Lrrn4cl	Carkd	Bysl	
189	Cdh3	Cd24a	Esd	

190	Cyr61	Eng	Aldh2	
191	Cyp2d22	Tcirlg1	Rhog	
192	Hist1h1c	Slc9a3r2	kg:uc009ecr.1	
193	Aplp1	0910001L09Rik	Cald1	
194	Tbl1x	Cox5b	Wbp2	
195	Pcm1	Adipor2	Ptpn1	
196	Ifi204	Scarf2	Tpm4	
197	Nfix	Myo7a	Mxi1	
198	Flrt2	Ppap2c	Ly6g6f	
199	Heg1	Pea15a	Sla	
200	Il6ra	Sh3pxd2b	Slpi	
201	Ralbp1	H19	Bicd2	
202	Rhoj	Tpd52	Clu	
203	Ktn1	2610203C20Rik	Mtmr14	
204	Arl6ip5	Naa10	Abca7	
205	Crebbp	Fermt1	Ppp1r18	
206	Ppig	Sap301	Kif2a	
207	Akap13	Bgn	Prdx6	
208	Rab7	Timm13	kg:uc009ize.1	
209	Plxdc2	Krt20	Calm3	
210	Aldh1a1	Itga3	Dhrs1	
211	Bnc2	Pfkl	Cfl1	
212	Slc4a4	Agpat6	Glipr2	
213	Tbx18	Mrpl11	Slc25a37	
214	Zbtb16	Ramp1	Atox1	
215	Arid4b	Hmga1	BC057079	
216	Enpp2	Gpx2	Pla2g16	
217	Ptplad2	0610012G03Rik	Rnf144b	
218	Akr1b3	9130017N09Rik	Stk16	
219	Gm6644	Cygb	Rsd2	
220	Arf5	Tmprss4	Paip2	
221	Chi3l1	Paox	Capzb	
222	Gpr116	Endod1	Ppp1r12c	
223	Cd82	Cndp2	4930412F15Rik	
224	Srrm1	Suv39h1	Ninj1	
225	Fmo2	Cog4	2510009E07Rik	
226	Tgfb1i1	Trim27	kg:uc007vsr.1	
227	Qrich1	Cyhr1	Pygb	
228	Nfia	Trmt1	Tlk1	
229	Pmp22	Zfyve19	Myct1	
230	Cdh11	Esrp1	Rnasek	
231	Arid5b	kg:uc008oow.1	Ctsd	

232	Rbm3	Dync1h1	0610010K14Rik	
233	Prelp	Tab1	Bcas3	
234	kg:uc007qse.1	Pla2g6	Atpif1	
235	Ddx3x	Timp1	Serf2	
236	Sulf1	Eif3f	Becn1	
237	Spnb2	Abhd11	Tspan9	
238	Tspan31	Pmm2	Acer2	
239	Prr13	Tyrobp	Vdac3	
240	Ppp1cb	Farsb	kg:uc008kbg.1	
241	Fbln1	Plod3	Oaz2	
242	Gm6548	Abtb1	Serpine2	
243	Uap1	Brf1	Ccdc90a	
244	Mpdz	Tnk2	Ndufa1	
245	Sat1	Rfc2	Tssc1	
246	Stim1	Stxbp2	Mboat7	
247	Mll3	Pdlim7	Cd44	
248	Slurp1	A430105I19Rik	Cxx1c	
249	Cd81	Vill	Ecm1	
250	Emp2	Bmp1	Mff	
251	Trpm7	Mpzl1	Ptpn12	
252	Crym	Thy1	Mgmt	
253	Enpp4	Stab1	Cox4i1	
254	Raly	Aldh16a1	Tollip	
255	Celf2	Eif4ebp3	Cds2	
256	Ap3s1	Itpripl2	Ybx1	
257	C1s	Mrpl52	Gypc	
258	Frmd4b	2310002L13Rik	Dgkd	
259	Nr4a1	Mcm6	Pecam1	
260	Acin1	Kcnk1	Ftl2	
261	Plod2	Pmf1	Nt5c3	
262	Id1	Cuta	1700037H04Rik	
263	Creg1	Nt5dc2	Cd151	
264	Zfp318	Rmnd5b	Lpin2	
265	Tmem140	Araf	6430548M08Rik	
266	Mras	Wwp2	Pon2	
267	Vwa5a	Lamb1	Ndufa3	
268	Esyt3	Kcne3	6330578E17Rik	
269	Hexb	Uqcrq	Mfap3l	
270	Nckap1	Gps1	Mink1	
271	Nipal3	Rexo4	Ston2	
272	Ubxn4	Coro1c	Rac2	
273	Zfp36	Hras1	Fyn	

274	Hnrnpl	Spint1	Serinc3	
275	C1ra	Cblc	Maged2	
276	Nnmt	Fhod1	Ap2m1	
277	Mut	Atp13a1	Pacsin2	
278	kg:uc008jup.1	Man2c1	Ftl1	
279	Pnrc1	Vsig2	Adipor1	
280	Usp8	Bpgm	kg:uc009qdo.1	
281	Pgcp	Bap1	Snap23	
282	Junb	Smpd2	Tagln2	
283	C1rl	Ubqln4	Cox6c	
284	Slc6a6	Sirt7	Creg1	
285	kg:uc008znh.1	Krt23	Bsg	
286	Aqp1	D8Ertd738e	Cmtm6	
287	Myh10	Mapk13	Cntd1	
288	Slc43a3	kg:uc008bcq.1	Plekho2	
289	Spint2	Polr2g	Arrb2	
290	Hnrnph1	Ndufs2	Pard3b	
291	Arhgap28	Dad1	Mlec	
292	Cfh	Wnt7b	Taf10	
293	Brd4	Fam20c	Gabarapl2	
294	Fndc1	Cxxc5	Bag1	
295	Star	Polr2f	Galnt2	
296	Nfkbiz	Ltf	Hk1	
297	Arsb	2210407C18Rik	Fbxo9	
298	Rnd3	Cdip1	kg:uc009izd.1	
299	Stard5	Glrx5	Pnpo	
300	Thbs1	Gemin7	Fam46c	
301	kg:uc008wkn.1	Man1b1	Pkm	
302	Slc26a3	Heatr7a	Ap1b1	
303	Phip	Arid5a	Rap1b	
304	Usp2	Sumo3	Itgb1	
305	Golgb1	Srm	St7	
306	Rock1	Plscr3	Smap1	
307	Rgma	2210010C17Rik	Rabgap11	
308	Actg1	Fam102a	Tmbim4	
309	BC013529	Dlst	H3f3a	
310	kg:uc007zwh.1	Vps37c	Frmd8	
311	3110062M04Rik	Ngfrap1	Nlrx1	
312	Cast	Pold4	Oaz1	
313	Mob3c	Grc10	Fam125b	
314	Slc16a1	Wnt7a	Hexa	
315	Fam117a	2010111I01Rik	Tspo	

316	Pdia3	Pxdn	Dcaf12	
317	Trim8	Coasy	Nav1	
318	kg:uc009mng.1	Dctn1	Cd24a	
319	eg:245190:chr7:m	Ncor2	Uqcr11	
320	Sbsn	Postn	Wipf1	
321	Serpinb6b	Col4a2	F10	
322	Daglb	Cib1	Erlec1	
323	Smarca2	Tbc1d13	Map2k3	
324	Mef2c	Ccnl2	Stk24	
325	Prrc2c	Dcakd	Ldlrap1	
326	BC005537	Cdc34	Ehd4	
327	Hsp90ab1	Atp6v0b	Atp6v1f	
328	Snrnp70	Abhd12	Gnas	
329	Ppl	Flot2	Arhgap18	
330	Serpinh1	Sla2	Arhgap10	
331	Sorbs3	Rhbdf1	Pitpnm1	
332	Golga4	Cdh17	S100a1	
333	Acbd3	Psmb5	Bin1	
334	Hook3	Serf1	Ttyh3	
335	Map3k3	Slc15a3	Selp	
336	Rhou	Sftp1	Trappc9	
337	Smc2	Pop5	Aes	
338	C1d	Nudc	Taok3	
339	kg:uc008dzh.1	Sh2d5	Zfand3	
340	Psmd7	kg:uc007fwp.1	Stim1	
341	Dab2	Mrpl37	Rnf114	
342	Cep164	Rin1	Sep15	
343	Crim1	Podxl	kg:uc012hdk.1	
344	Rtf1	Paqr5	Lgals9	
345	Fxyd1	Sepx1	Cox6b1	
346	H2-D1	Agr2	Riok3	
347	Zfp704	Bax	Slc38a10	
348	Mtap1a	Rxrb	Rtn3	
349	Ascc3	Tes	B3gat2	
350	Med131	Hdac6	Ccndbp1	
351	Jup	1110008F13Rik	Rsu1	
352	Nid2	Mpnd	kg:uc007upr.1	
353	Kdr	Gmppa	Itm2b	
354	Ifnar2	Gramd1a	St3gal1	
355	5430435G22Rik	Wars	Sec61g	
356	Col4a6	Mtap	Ptpn1	
357	Il17re	C1qtnf5	kg:uc012bhf.1	

358	Gbp3	Mrpl28	B2m	
359	Slc39a8	Mfrp	Rasgrp3	
360	Cfl2	Kars	Memo1	
361	Slc38a1	Lbp	Slc39a4	
362	Cuedc1	Plxnb1	Sdcbp	
363	Fgf1	2700081O15Rik	Tspan14	
364	Gas6	Mrps24	Ubl7	
365	Cldn25	Klc4	Nras	
366	Sorbs1	Dctn3	Ssx2ip	
367	Hspa12a	Kcnq1	kg:uc007zbz.1	
368	kg:uc007zts.1	Smurf1	Wbp1	
369	Slc1a5	Fam162a	1110003E01Rik	
370	Nr3c1	Hip1r	Clip2	
371	Adamts5	kg:uc007hyr.2	Gapdh	
372	Gpcpd1	Gys1	Gm6578	
373	Dpysl3	Sac3d1	Actn1	
374	Colec12	Ndufs6	St3gal2	
375	Pdcd6ip	Rgl2	3110001D03Rik	
376	Dst	Atp5g1	Ctsz	
377	Ifit3	Itgb4	kg:uc007vdl.1	
378	Chst4	Sars	Fam73a	
379	Xist	2310003F16Rik	Vcl	
380	Ifi27l2a	Nhp2l1	Lims1	
381	Fkbp5	D19Wsu162e	Lars2	
382	Agap1	Cd320	Birc2	
383	Ankrd11	Pigq	Lamp2	
384	kg:uc007qca.1	Chd3	Ras110a	
385	Syt11	Zdhhc4	Mif	
386	Ptrf	Eif3l	Rab10	
387	Krc1	St8sia3	Pabpc1	
388	Zfp488	Rcan3	Wwp2	
389	Lama4	Meg3	Nqo2	
390	Aebp1	Nudt4	kg:uc007fte.1	
391	Fam134b	Gss	Plxna4	
392	Tppp3	Pih1d1	Gm1821	
393	Maf	Limd2	Gadd45a	
394	Peli1	Ap1s2	Slc25a39	
395	Zfp353	BC056474	kg:uc009pet.1	
396	Cdon	Mms19	Ubb	
397	Sarnp	Clip2	Ppp1r2	
398	Atxn7l3b	2310016M24Rik	Rab27b	
399	Pef1	Itpa	Cap1	

400	App	Slc25a10	Jarid2	
401	Mtdh	Fibp	Rnf11	
402	Lrrc20	Higd2a	Tmem50b	
403	Btbd2	Snrpd2	Myh9	
404	Gnb2	Eri3	Tmem128	
405	Pigt	Nbeal2	Stradb	
406	Efna5	Trim28	Cela1	
407	Tm4sf1	S100a4	Ndrg2	
408	Coq10b	Ivns1abp	Dhrs3	
409	Eif2s3x	Ppp1r18	Hipk1	
410	Cmah	Efemp2	Atg9a	
411	Sf3b1	Med22		
412	Eea1	Nelf		
413	Slpi	2810428I15Rik		
414	Tmod3	D2Wsu81e		
415	Ppp3ca	Trappc6a		
416	Tceal8	Trappc21		
417	Anp32a	Antxr2		
418	Actb	Rab11fip5		
419	Ddx5	Ldhd		
420	Cobll1	Npnt		
421	Cish	Acrbp		
422	Nod1	Pafah1b2		
423	Psd	Angptl2		
424	Gm10052	Fzr1		
425	Lims2	Aaas		
426	Stra6	Eif2b2		
427	kg:uc007bgn.1	1190003J15Rik		
428	Plxdc1	5730403B10Rik		
429	Nfe211	Adamts13		
430	Smpd3	Eif3b		
431	Bcl10	Znrf1		
432	Ilf3	Pkp3		
433	Fam76a	Lemd2		
434	Cybrd1	Rab34		
435	Gm3893	Mpv17l2		
436	Siae	Cdkn2b		
437	Ssh2	Snrpe		
438	Nfic	Gm14005		
439	Btf3	Prdx4		
440	Sp100	Xab2		
441	Ndn	Dpp3		

442	Matr3	Tyms		
443	Gm13251	Leprotl1		
444	Arhgap5	Uqcr10		
445	Zbtb4	Cdk5rap3		
446	Pgrmc1	Gorasp2		
447	4930402H24Rik	Wbp7		
448	Bptf	Sort1		
449	Dusp3	Ddx41		
450	Pla2g4a	Cct3		
451	Brp441	Mrps33		
452	Oxct1	Frmd8		
453	Stk40	1110049F12Rik		
454	Ddr1	Fscn1		
455	Ifi205	Ndufa2		
456	Col3a1	Dpcd		
457	Nipbl	Unc13a		
458	Plk1s1	Eif1ad		
459	Bdp1	Sgta		
460	Smc3	Chaf1a		
461	Ifitm3	Plxna1		
462	Ndst1	Hspa9		
463	Zbed6	1110014N23Rik		
464	Rest	Cd99l2		
465	kg:uc007vnc.1	Snrpa		
466	Ccdc88a	Mcm7		
467	Stat3	Tars2		
468	Arf2	Gon41		
469	Trib1	Stk38		
470	Gcap14	C1qtnf1		
471	Tbc1d15	Tbrg4		
472	Igf1r	Tmem132a		
473	Ppbp	Cox6c		
474	kg:uc008tky.1	Alcam		
475	Rab1b	Phka2		
476	Krt14	Trim3		
477	Med21	Ppp1r14b		
478	Gja1	Gpaa1		
479	Klf10	Ctps2		
480	Id2	Ptpn23		
481	Mfap1a	Endog		
482	Ogn	Mrto4		
483	Gpc4	Mrps6		

484	Bst2	Pvr		
485	Dtx2	Phgdh		
486	Wac	Itpr3		
487	Kpna3	Polr2e		
488	Kcnab1	Sec16a		
489	Orai3	Mdp1		
490	Gcsh	Fbf1		
491	Wdr92	Mcpt8		
492	Olf613	Rps6ka4		
493	Tcf7l1	Mical1		
494	Tgfb2	Mrpl34		
495	Il16	Agpat3		
496	Manf	2310044H10Rik		
497	Mgst1	Myo9b		
498	kg:uc008tkz.1	Ndufb10		
499	Creb3l1	Apex1		
500	Txndc5	Elk3		
501	Klf2	Cpsf31		
502	Slu7	Tnk1		
503	Ttc28	Pmvk		
504	1110002B05Rik	Ppp1r16a		
505	Zcchc11	Arhgef5		
506	Ptp4a2	Lonp1		
507	Pbx1	Pla2g7		
508	Clcn3	Pip5k1c		
509	Tmco7	Inf2		
510	Lrrc58	Pgk1		
511	Eif3a	Parp6		
512	Cldn10	Urm1		
513	H2-Q6	Mad2l2		
514	Ccdc80	Ing4		
515	kg:uc009iln.1	Rbck1		
516	Rab5c	Cant1		
517	Tsc22d3	Sgpl1		
518	Tm4sf5	Ehbp111		
519	Hmgb1	Runx1		
520	Sec62	Slc27a4		
521	Maoa	Ndufa7		
522	Clec1b	Mcm3ap		
523	Mphosph8	1110008P14Rik		
524	Oat	Rassf7		
525	Ncor1	Ptpmt1		

526	Cyb5	Arfgap1		
527	Trafld1	Sec61a1		
528	Rpp25	Rps6ka1		
529	kg:uc007ded.1	Ints1		
530	2610101N10Rik	Tpcn1		
531	Il6st	Iffo2		
532	Evp1	Trim44		
533	Psmd11	kg:uc012ctw.1		
534	Dync1i2	Golga2		
535	Lars2	Msto1		
536	Pdia4	Ppp6r3		
537	Cd55	Trmt2a		
538	Amfr	Appl2		
539	Zcchc3	Sparcl1		
540	Herpud2	Rapgef1		
541	Txnrd1	Zfp11		
542	Vat1	Psmc4		
543	Diap1	Mosc2		
544	Tmed2	Fam101b		
545	Arf3	1500010J02Rik		
546	Arap2	Ccdc124		
547	St3gal1	Ptges		
548	Man1a	Fam189b		
549	Rgs10	Th11		
550	Tmsb4x	Kctd2		
551	Uba7	Olfr1372-ps1		
552	C4b	Hexa		
553	Tmem98	Anapc5		
554	Lpar2	Serpina3n		
555	Gabarap11	1810046J19Rik		
556	Cmtm7	Tmem167		
557	Spon2	Gm11428		
558	Smarca5	Gcn111		
559	Mxd4	Kansl3		
560	Smc4	Fasn		
561	Thsd4	Slc50a1		
562	Gsr	Smad3		
563	Ptprd	Trip6		
564	Clip1	Atp6v1e1		
565	Cln8	Chchd5		
566	Rbm27	Adss11		
567	Zmat1	Nes		

568	Smc6	Ap1b1		
569	B2m	Fcgrt		
570	Irf2bp2	Ltbp3		
571	Ppap2a	Csf2rb		
572	Zfhx4	Ssna1		
573	Tob2	Mrps16		
574	Rabgap11	Cyba		
575	Nfkb2	Cyth2		
576	Nfyc	Igf2		
577	Ube2d1	Pisd-ps1		
578	Creb5	Atp13a2		
579	Opa3	Mlph		
580	Csnk1a1	Cyp4f16		
581	Fam84b	2010107E04Rik		
582	Ddr2	Gas5		
583	Usp54	Eif3k		
584	Akt2	Fam149a		
585	Strn3	Mif		
586	Hnrnpm	B230312A22Rik		
587	eg:497210:chr14:m	Ppp1r12c		
588	Tpt1	Tfip11		
589	Naa25	Tex10		
590	Eef1a1	Slc16a3		
591	Parp4	Stk16		
592	Msn	Epn1		
593	Zbtb20	Noc4l		
594	Fermt2	Rcc2		
595	Bod11	Rgs12		
596	Sltm	Shkbp1		
597	Dapk1	Got2		
598	Hnrnpr	Plek2		
599	Baz2a	Lilrb3		
600	Rnf167	Ndufb5		
601	Mapk1	Tesk1		
602	eg:320169:chr9:p	Rab24		
603	4930523C07Rik	Atp5j2		
604	Nfl	Commd9		
605	Fam53b	Rtkn		
606	Faim2	Prpf19		
607	Tgm2	6720401G13Rik		
608	Calm2	Ppa1		
609	AI848100	Pgp		

610	Slc10a3	Hps1		
611	Ogdh	Puf60		
612	Arl3	Mdm2		
613	Timp2	kg:uc012cgd.1		
614	Atxn2	kg:uc009uim.1		
615	Mll1	Pyy		
616	Ces2g	Zfp358		
617	Mat2a	Timm8b		
618	Esf1	Ddx39		
619	Hsp90aa1	Pgm2		
620	Zfp385a	kg:uc008gbp.1		
621	Zfp672	Sipa1		
622	Csda	Mgat1		
623	Pf4	Tmem208		
624	Arsa	Ruvbl2		
625	F11r	8430410A17Rik		
626	C4a	Bad		
627	Kpna1	Pfdn5		
628	Rbbp8	Eme1		
629	Oxnad1	kg:uc009mzj.1		
630	Rb1cc1	Igf1		
631	Setd2	Prkag1		
632	Kif1b	kg:uc009sua.1		
633	2510002D24Rik	Uap111		
634	Cep57	Trappc4		
635	Chd2	Bola2		
636	Serinc5	Usp5		
637	Marcks1	Ear2		
638	Shfm1	Cars		
639	Bbs4	1810027O10Rik		
640	Impad1	Amdhd2		
641	Tbcel	Phb		
642	Kdelr1	Kcmf1		
643	Ninl	Lsmd1		
644	Syt11	Sec11c		
645	Tpm3	Pcbp4		
646	Rbbp6	Mepce		
647	Lman1	Tpd52l2		
648	Ankrd17	Trf		
649	Naga	Hsd17b11		
650	Rbpms	Pilra		
651	Magt1	Atn1		

652	Tfdp2	Pgf		
653	Gem	Nxn		
654	Pde4dip	Inpp5k		
655	Mrgprf	Actrla		
656	kg:uc008ajk.1	Cd68		
657	Itch	Eef1g		
658	Elf1	Fbn1		
659	Meis2	Hint1		
660	Arid1a	March5		
661	Serpine1	Usp48		
662	Slc27a3	Hnflb		
663	Thoc2	Gga3		
664	Gsta3	Drosha		
665	Hnrnph2	Ubp1		
666	Soes3	Pkn3		
667	Armcx3	Tmem192		
668	Siah1a	Prpf31		
669	kg:uc009ize.1	Hspd1		
670	Irs2	Otub1		
671	Mettl7a1	Mrpl20		
672	Ppfibp2	Tead2		
673	Blvrb	Phpt1		
674	Yipf5	Neu1		
675	Plat	Pygo2		
676	Gm6578	Myeov2		
677	Mat2b	Cdk5		
678	Tmpo	Ndor1		
679	Metap2	Rbp4		
680	Zfp277	Psat1		
681	Wls	Mrpl41		
682	Mesdc1	Snrpg		
683	kg:uc009acs.1	Acot7		
684	Colla2	Vars		
685	Csf1	Nono		
686	Sulf2	Gtf2i		
687	Ifrd1	Traf3		
688	Wrnip1	Ppp2r4		
689	Flii	Actg2		
690	2810474O19Rik	Pi4k2a		
691	Sep15	Slc35b2		
692	2310030G06Rik	Ubqln1		
693	Cmtm3	Ppox		

694	Mylip	Bud31		
695	Slc8a1	Man2b1		
696	Btbd7	Nat15		
697	Hdac5	Spon1		
698	Zfand6	Cyc1		
699	Tapbp	Mpeg1		
700	Keap1	Nsun2		
701	Ube2n	Rab4a		
702	Ssr3	Mtmr11		
703	H3f3a	BC004004		
704	Myst4	B4galnt1		
705	G3bp1	Atp5k		
706	Ugdh	Lin37		
707	Lamp2	D330041H03Rik		
708	Zrsr1	Tbc1d17		
709	Pim1	March6		
710	Gm9199	2410015M20Rik		
711	Supt16h	1810013D10Rik		
712	Ano6	Eif2s1		
713	Soat1	Traf7		
714	Eci1	Rpl36al		
715	Plce1	Psenen		
716	Atg3	Aip		
717	Bnc1	Cmas		
718	Pik3c2a	Rpia		
719	Pqlc3	Ncbp1		
720	Thrap3	Mea1		
721	Irak4	Timm50		
722	Kdm6b	Ear12		
723	Apol9a	Fkbp1a		
724	Wnt4	Commd4		
725	1500003O03Rik	Col5a3		
726	Phf3	Fblim1		
727	1110004F10Rik	Cwh43		
728	Kans11	Arl2bp		
729	Fth1	Mrpl46		
730	Tmem50a	Tcn2		
731	Utp20	Add2		
732	Smad4	Specc11		
733	Stmn2	Ppcs		
734	Gstm1	Vrk3		
735	Senp6	Trim25		

736	Gda	Nfate1		
737	Nucks1	Rap1gap		
738	Ints10	Hsd17b12		
739	Syne1	Epas1		
740	Itga6	Ddx1		
741	Acad9	Prdx6		
742	Maged1	Mmp24		
743	Spen	Ndufb9		
744	Chd1	Phf23		
745	Taf3	Rpa2		
746	Ptgs1	5031439G07Rik		
747	Sparc	Rrp7a		
748	R74862	Arfip2		
749	B230120H23Rik	Efna1		
750	Tmem234	Agps		
751	Ryk	Seph8		
752	Dlgap4	Apoc2		
753	Atp1b1	Mrps27		
754	Parp14	Snn		
755	Tgfb2	Serinc3		
756	Ccdc90a	Pdcd5		
757	Ncoa1	AA986860		
758	Pppde1	Pitpna		
759	Luc713	Vac14		
760	Prg4	2810025M15Rik		
761	Rab11fip1	Def8		
762	Plk2	Hilpda		
763	Ifi35	Eif6		
764	Pdap1	Brd7		
765	Cd248	Fes		
766	Sesn1	Sbf1		
767	Ecd	Ak2		
768	Ap1s3	1810035L17Rik		
769	H2-K1	Lime1		
770	Spag9	Hspe1		
771	Tshz1	Csrp2bp		
772	Dennd5a	Uba5		
773	Stag1	Gsta4		
774	Gpx8	2900092E17Rik		
775	Sod3			
776	BC005561			
777	kg:uc009vev.1			

778	Ywhaz			
779	Ganab			
780	Rras2			
781	Dusp14			
782	kg:uc012hdk.1			
783	Nrl1d1			
784	Wwc2			
785	Ubxn2a			
786	Iqsec1			
787	kg:uc007vsr.1			
788	Cfl1			
789	Csrp1			
790	Smchd1			
791	Myl12a			
792	Ubqln2			
793	Tmcc3			
794	Kdm5a			
795	Rbm25			
796	Wdr26			
797	Vim			
798	Arpc2			
799	Calm1			
800	Dnaja2			
801	She1			
802	Vps13a			
803	Klf7			
804	1810074P20Rik			
805	BC003331			
806	Itpr2			
807	Jmjd1c			
808	Pcdhgb5			
809	Tubb2a			
810	Ehd2			
811	Ift74			
812	Per1			
813	Pitpnm2			
814	Gstm4			
815	Dnmt1			
816	Tmco1			
817	Lass4			
818	Ptpf			
819	Sirt2			

820	Gfm2			
821	Taf7			
822	Spop			
823	Zzef1			
824	Ccdc34			
825	Zfp281			
826	Tuba1a			
827	Ccdc109b			
828	Cdk13			
829	Dhx15			
830	Src			
831	Braf			
832	Mapre2			
833	Anxa7			
834	Sept9			
835	Alox12			
836	Pknox1			
837	2610034B18Rik			
838	Topors			
839	Phf21a			
840	Qser1			
841	Tirap			
842	Fas			
843	Lass2			
844	6330406I15Rik			
845	Parvb			
846	Atp1a1			
847	Mtmr6			
848	Cd109			
849	Dnajc1			
850	Hp1bp3			
851	1600029D21Rik			
852	Ttc38			
853	Mfhas1			
854	Filip11			
855	Zfp148			
856	Nkd1			
857	Usp16			
858	Tlr2			
859	Zc3h18			
860	Stk10			
861	Ltbp4			

862	Hdac3			
863	Efhd2			
864	Prkar2a			
865	Atp6v1a			
866	Sf3b4			
867	Gprc5b			
868	Clip3			
869	Mettl2			
870	Secisbp2			
871	Fmod			
872	kg:uc009lxf.1			
873	Elovl6			
874	Bzw1			
875	Etfa			
876	Hspa2			
877	kg:uc007won.1			
878	Rnf20			

**[00198]** **Table 3:** Most Significant Gene Ontology Terms in CTC-c enriched genes using BP\_FAT and CC\_FAT Datasets

q-value < 0.01	Term	Count	Odds Ratio	Benjamini (q-value)
Source	Term	Count	Odds Ratio	Benjamini (q-value)
GOTERM_BP_FAT	GO:0060429~epithelium development	35	2.92	8.72E-05
GOTERM_BP_FAT	GO:0030029~actin filament-based process	27	3.47	6.85E-05
GOTERM_BP_FAT	GO:0030036~actin cytoskeleton organization	26	3.57	4.95E-05
GOTERM_BP_FAT	GO:0007010~cytoskeleton organization	36	2.50	6.27E-04
GOTERM_BP_FAT	GO:0051173~positive regulation of nitrogen compound metabolic process	49	2.11	6.62E-04
GOTERM_BP_FAT	GO:0035295~tube development	31	2.66	7.80E-04
GOTERM_BP_FAT	GO:0010604~positive regulation of macromolecule metabolic process	54	1.93	0.001727
GOTERM_BP_FAT	GO:0031328~positive regulation of cellular biosynthetic process	49	2.01	0.0015751
GOTERM_BP_FAT	GO:0051789~response to protein stimulus	16	4.16	0.0014484
GOTERM_BP_FAT	GO:0035239~tube morphogenesis	23	3.05	0.0015064
GOTERM_BP_FAT	GO:0045449~regulation of transcription	140	1.42	0.0014097
GOTERM_BP_FAT	GO:0048729~tissue morphogenesis	28	2.66	0.0013058
GOTERM_BP_FAT	GO:0009891~positive regulation of biosynthetic process	49	1.99	0.0012408
GOTERM_BP_FAT	GO:0045935~positive regulation of nucleobase, nucleoside, nucleotide and nucleic acid metabolic process	46	2.04	0.0012061

GOTERM_BP_FAT	GO:0002009~morphogenesis of an epithelium	23	3.01	0.0012149
GOTERM_BP_FAT	GO:0048584~positive regulation of response to stimulus	24	2.92	0.0011396
GOTERM_BP_FAT	GO:0051276~chromosome organization	39	2.19	0.0012619
GOTERM_BP_FAT	GO:0045637~regulation of myeloid cell differentiation	12	5.33	0.0014358
GOTERM_BP_FAT	GO:0045785~positive regulation of cell adhesion	11	5.79	0.0016889
GOTERM_BP_FAT	GO:0045941~positive regulation of transcription	43	2.05	0.0016795
GOTERM_BP_FAT	GO:0045893~positive regulation of transcription, DNA-dependent	39	2.12	0.0019852
GOTERM_BP_FAT	GO:0051254~positive regulation of RNA metabolic process	39	2.11	0.0022107
GOTERM_BP_FAT	GO:0006357~regulation of transcription from RNA polymerase II promoter	51	1.87	0.0022801
GOTERM_BP_FAT	GO:0006325~chromatin organization	32	2.30	0.0025187
GOTERM_BP_FAT	GO:0010628~positive regulation of gene expression	43	2.00	0.0025252
GOTERM_BP_FAT	GO:0060562~epithelial tube morphogenesis	17	3.47	0.0025847
GOTERM_BP_FAT	GO:0042127~regulation of cell proliferation	45	1.89	0.0051485
GOTERM_BP_FAT	GO:0010557~positive regulation of macromolecule biosynthetic process	44	1.88	0.0071937
GOTERM_BP_FAT	GO:0002253~activation of immune response	14	3.69	0.0078441
GOTERM_BP_FAT	GO:0050778~positive regulation of immune response	18	3.00	0.0080458
GOTERM_BP_FAT	GO:0002684~positive regulation of immune system process	23	2.53	0.0088166
GOTERM_BP_FAT	GO:0045944~positive regulation of transcription from RNA polymerase II promoter	33	2.09	0.0090124
GOTERM_CC_FAT	GO:0005578~proteinaceous extracellular matrix	32	2.38	0.0047511
GOTERM_CC_FAT	GO:0031012~extracellular matrix	32	2.28	0.0051923
GOTERM_CC_FAT	GO:0044421~extracellular region part	60	1.71	0.0064365
GOTERM_CC_FAT	GO:0031981~nuclear lumen	65	1.62	0.0102413
GOTERM_CC_FAT	GO:0043233~organelle lumen	79	1.53	0.0085938
GOTERM_CC_FAT	GO:0005829~cytosol	45	1.81	0.0100772
GOTERM_CC_FAT	GO:0070013~intracellular organelle lumen	78	1.52	0.0093866
GOTERM_CC_FAT	GO:0031982~vesicle	43	1.83	0.0087123
GOTERM_CC_FAT	GO:0031974~membrane-enclosed lumen	80	1.50	0.0082696

[00199] **Table 4:** Most Significant Gene Sets Enriched in CTC-plt vs CTC-c

q-value < 0.01					
Source		Count	Odds Ratio	Benjamini (q-value)	
GOTERM_BP_FAT	GO:0042060~wound healing	18	7.8	1.86E-07	
GOTERM_BP_FAT	GO:0007596~blood coagulation	15	10.4	9.31E-08	
GOTERM_BP_FAT	GO:0050817~coagulation	15	10.4	9.31E-08	
GOTERM_BP_FAT	GO:0007599~hemostasis	15	10.3	7.59E-08	
GOTERM_BP_FAT	GO:0050878~regulation of body fluid levels	15	8.2	1.30E-06	
GOTERM_BP_FAT	GO:0030029~actin filament-based process	20	5.5	1.14E-06	
GOTERM_BP_FAT	GO:0007010~cytoskeleton organization	26	3.9	3.95E-06	
GOTERM_BP_FAT	GO:0030036~actin cytoskeleton organization	18	5.3	1.11E-05	
GOTERM_BP_FAT	GO:0009611~response to wounding	26	3.6	1.02E-05	
GOTERM_BP_FAT	GO:0007155~cell adhesion	33	2.9	2.86E-05	
GOTERM_BP_FAT	GO:0022610~biological adhesion	33	2.8	2.70E-05	
GOTERM_BP_FAT	GO:0001775~cell activation	19	3.7	4.70E-04	
GOTERM_BP_FAT	GO:0030168~platelet activation	6	18.2	1.68E-03	
GOTERM_BP_FAT	GO:0007229~integrin-mediated signaling pathway	10	6.4	2.95E-03	
GOTERM_BP_FAT	GO:0016192~vesicle-mediated transport	25	2.6	3.81E-03	
MSigDBv3.1 CGP	GNATENKO PLATELET SIGNATURE	20	55.1	3.91E-24	
MSigDBv3.1 CGP	TENEDINI MEGAKARYOCYTE MARKERS	14	15.3	1.35E-11	
MSigDBv3.1 CP:REACTOME	REACTOME FACTORS INVOLVED IN MEGAKARYOCYTE DEVELOPMENT AND PLATELET PRODUCTION	6	2.9	2.25E-02	

[00200] **Table 5:** Most Significant Gene Sets Enriched in CTC-pro vs CTC-c

q-value &lt; 0.01

Source	Term	Count	Odds Ratio	Benjamini (q-value)
GOTERM_BP_FAT	GO:0002495~antigen processing and presentation of peptide antigen via MHC class II	5	59.81	6.97E-04
GOTERM_BP_FAT	GO:0019886~antigen processing and presentation of exogenous peptide antigen via MHC class II	5	59.81	6.97E-04
GOTERM_BP_FAT	GO:0002504~antigen processing and presentation of peptide or polysaccharide antigen via MHC class II	5	50.36	7.34E-04
GOTERM_BP_FAT	GO:0002478~antigen processing and presentation of exogenous peptide antigen	5	41.60	1.10E-03
GOTERM_BP_FAT	GO:0019884~antigen processing and presentation of exogenous antigen	5	34.18	1.87E-03
GOTERM_BP_FAT	GO:0048002~antigen processing and presentation of peptide antigen	5	27.34	3.72E-03
GOTERM_BP_FAT	GO:0001775~cell activation	9	7.00	3.82E-03

GOTERM_BP_FAT	GO:0019882~antigen processing and presentation	6	13.20	7.40E-03
---------------	--	---	-------	----------

## EXAMPLE 2: SUPPLEMENTAL METHODS

**[00201]** *Mice and cell lines.* Mice with pancreatic cancer used in these experiments express Cre driven by *Pdx1*, *LSL-Kras*<sup>G12D</sup>, and *Trp53*<sup>lox/+</sup> or *Trp53*<sup>lox/lox</sup> (otherwise referred to as KPC) as previously described (Bardeesy et al., 2006). EGFP pancreatic lineage tagged KPC mice were generated by breeding the mT/mG mouse (Purchased from the Jackson Laboratory - Gt(ROSA)26Sortm4(ACTB-tdTomato,-EGFP)Luo/J) into the breeder pairs used for KPC mouse generation. Normal FVB mice were purchased from Jackson Laboratory. All mice care and procedures were done under MGH SRAC approved protocols.

**[00202]** For cardiotocentesis, animals were sedated with isofluorane, the chest wall was sterilized with ethanol and a skin incision was made above the rib cage to expose the thoracic cavity and eliminate normal skin epithelial cell contamination. A 23-gauge needle was used to draw approximately 1 mL of blood into a 1 mL syringe primed with 100  $\mu$ L of PBS-10mM EDTA pH 7.4 (Gibco). Blood EDTA concentration was raised to 5mM by either the addition of a concentrated bolus of 500mM EDTA or 1:1 dilution with 10 mM EDTA. Animals were then euthanized per animal protocol guidelines.

**[00203]** A mouse pancreatic cell line NB508 (*Pdx1-Cre/Kras*<sup>G12D</sup>/*Trp53*<sup>lox/+</sup>) previously generated from primary tumors developed in this endogenous model was GFP transfected by lentivirus (NB508-GFP). This cell line was used for spiked cell experiments and orthotopic tumor formation.

**[00204]** NB508-GFP Cell lines were maintained in standard culture conditions using RPMI-1640 medium + 10% FBS + 1% Pen/Strep (Gibco/Invitrogen).

**[00205]** For orthotopic experiments, NB508-GFP cells were orthotopically injected into the pancreas of healthy syngeneic (FVB background) mice. Briefly, mice were anesthetized with isofluorane and the left abdominal wall was treated with Nair® hair removal product, and sterilized with 70% ethanol. A small incision was made on the upper left lateral abdominal wall and the pancreas was mobilized. Approximately 1 million NB508-GFP cells in PBS in a total volume of 0.1 mL was injected into the pancreas. The peritoneum and abdominal wall was closed by sterile surgical staples. The tumors were allowed to grow for 2 weeks, at which time blood was obtained by cardiotocentesis for CTC-iChip processing.

**[00206]** *Adaptation of CTC Enrichment Technology.* Given the desire for an unbiased enrichment system, the negative depletion technology was selected for this application. All processing protocols were identical to those previously identified, except a rat anti-mouse CD45 antibody (BAM114, R&D Systems, USA) was conjugated to MyOne beads.

**[00207]** Spiked cell experiments were conducted to validate the system by spiking ~ 1000 GFP expressing NB508 cells into 1 mL of healthy mouse blood and processing to determine recovery efficiency. Orthotopic models were used to validate recovery efficiency as well as initially determine expected depletion efficiency from tumor-bearing mice. In these experiments, enriched samples were evaluated for the number of GFP+ cells observed in the product.

**[00208]** *Immunostaining of CTCs Isolated from the endogenous model.* Isolated CTCs were spun onto glass slides and immunostained using a primary-secondary approach. Primary antibodies were rabbit anti-wide spectrum cytokeratin (1:50, Abcam ab9377), and goat anti-mouse CD45 (1:500, R&D systems AF114). Secondary immunofluorescent-tagged antibodies were used for signal amplification. These were donkey anti-rabbit Alexa Fluor 594 (1:500, Invitrogen A-21207), and donkey anti-goat Alexa Fluor 488 (1:500, Invitrogen A-11055). Nuclei were then counterstained with DAPI and the slides were rinsed with PBS, cover slipped and stored at 4°C. They were imaged under 10x magnification using the BioView™ Ltd. automated imaging system (Billerica, MA) as well as an automated upright fluorescence microscope (Eclipse 90i™, Nikon, Melville, NY). Positive staining for CK, without CD45 staining, was required for scoring potential CTCs, which were then manually reviewed. Threshold and baseline signals were established using specimens from non-tumor bearing mice.

**[00209]** *Single cell micromanipulation.* After whole blood anti-CD45 negative depletion, the product containing enriched cells was collected in a 35mm petri dish and viewed using a Nikon Eclipse Ti™ inverted fluorescent microscope. Cells of interest were identified based on intact cellular morphology and lack of labeling with anti-CD45 magnetic beads. These target cells were individually micromanipulated with a 10  $\mu$ m transfer tip on an Eppendorf TransferMan® NK 2 micromanipulator and ejected into PCR tubes containing RNA protective lysis buffer (10X PCR Buffer II, 25mM MgCl<sub>2</sub>, 10% NP40, 0.1 M DTT, SUPERase-In, RNase Inhibitor, 0.5 uM UP1 Primer, 10mM dNTP and Nuclease-free water) and immediately flash frozen in liquid nitrogen.

**[00210]** *Single Cell Amplification and Sequencing.* Single cell amplification and sequencing were done as previously described (Tang et al., 2010) with slight modifications underlined below. RNA samples from extracted single circulating tumor cells were thawed on ice and incubated at 70°C for 90 seconds. To generate cDNA, samples were treated with reverse transcription master mix (0.05 uL RNase inhibitor, 0.07uL T4 gene 32 protein, and 0.33uL SuperScript™ III Reverse Transcriptase per 1X volume) and incubated on thermocycler at 50°C for 30 minutes and 70°C for 15 minutes. To remove free primer, 1.0uL of EXOSAP mix was added to each sample, which was incubated at 37°C for 30 minutes and inactivated at 80°C for 25 minutes. Next, a 3'-poly-A tail was added to the cDNA in each sample by incubating in master mix (0.6uL 10X PCR Buffer II, 0.36uL 25mM MgCl<sub>2</sub>, 0.18uL 100mM dATP, 0.3uL Terminal Transferase, 0.3uL RNase H, and 4.26uL H<sub>2</sub>O per 1X volume) at 37°C for 15 minutes and inactivated at 70°C for 10 minutes. A second strand cDNA was synthesis by

dividing each sample into 4 and incubating in master mix (2.2uL 10X High Fidelity PCR Buffer, 1.76uL 2.5mM each dNTP, 0.066uL UP2 Primer at 100uM, 0.88uL 50mM MgSO<sub>4</sub>, 0.44uL Platinum Taq DNA Polymerase, and 13.654uL H<sub>2</sub>O per 1X volume) at 95°C for 3 minutes, 50°C for 2 minutes, and 72°C for 10 minutes.

**[00211]** PCR amplification (95°C for 3 minutes, 20 cycles of 95°C for 30 seconds, 67°C for 1 minute, and 72°C for 6 minutes 6 seconds) was performed with master mix (4.1uL 10X High Fidelity PCR Buffer, 1.64uL 50mM MgSO<sub>4</sub>, 4.1uL 2.5mM each dNTP, 0.82uL AUP1 Primer at 100uM, 0.82uL AUP2 Primer at 100uM, 0.82uL Platinum Taq DNA Polymerase, and 6.7uL H<sub>2</sub>O per 1X volume). The 4 reactions of each sample were pooled and purified using the QIAGEN PCR Purification Kit (Cat. No 28106) and eluted in 50uL EB buffer. Samples were selected by testing for genes Gapdh, ActB, Ptprc (CD45), Krt8, Krt18, Krt19, and Pdx1 using qPCR. Each sample was again divided in 4 and a second round of PCR amplification (9 cycles of 98°C for 3 minutes, 67°C for 1 minute, and 72°C for 6 minutes 6 seconds) was performed with master mix (9uL 10X High Fidelity PCR Buffer, 3.6uL 50mM MgSO<sub>4</sub>, 13.5uL 2.5mM each dNTP, 0.9uL AUP1 Primer at 100uM, 0.9uL AUP2 Primer at 100uM, 1.8uL Platinum Taq DNA Polymerase, and 59.1uL H<sub>2</sub>O per 1X volume). Samples were pooled and purified using Agencourt AMPure XP beads and eluted in 40uL 1X low TE buffer.

**[00212]** *Sequencing Library Construction.* To shear the DNA using the Covaris S2<sup>TM</sup> System, 1X low TE buffer and 1.2uL shear buffer were added to each sample. Conditions of the shearing program include: 6 cycles, 5°C bath temperature, 15°C bath temperature limit, 10% duty cycle, intensity of 5, 100 cycles/burst, and 60 seconds. Then, samples were end-polished at room temperature for 30 minutes with master mix (40uL 5X Reaction Buffer, 8uL 10mM dNTP, 8uL End Polish Enzyme1, 10uL End Polish Enzyme2, and 14uL H<sub>2</sub>O per 1X volume). DNA fragments larger than 500bp were removed with 0.5X volumes of Agencourt AMPure XP<sup>TM</sup> beads. Supernatant was transferred to separate tubes. To size-select 200-500bp DNA products, 0.3X volumes of beads were added and samples were washed 2X with 70% EtOH. The products were eluted in 36uL low TE buffer. A dA-tail was added to each size-selected DNA by treating with master mix (10uL 5X Reaction Buffer, 1uL 10mM dATP, and 5uL A-Tailing Enzyme I per 1X volume) and incubated at 68°C for 30 minutes and cooled to room temperature. To label and distinguish each DNA sample for sequencing, barcode adaptors (5500 SOLiD 4464405) were ligated to DNA using the 5500 SOLiD Fragment Library Enzyme Module<sup>TM</sup> (4464413). Following barcoding, samples were purified twice using the Agencourt AMPure XP<sup>TM</sup> beads and eluted in 22uL low TE buffer. Following a round of PCR Amplification (95°C for 5 minutes, 12 cycles of 95°C for 15 seconds, 62°C for 15 seconds, and 70°C for 1 minute, and 70°C for 5 minutes), the libraries were purified with AMPure XP beads. Finally, to quantify the amount of ligated DNA, SOLiD Library TaqMan Quantitation Kit<sup>TM</sup> was used to perform qPCR.

Completed barcoded libraries were then subjected to emulsion PCR with template beads preparation and sequenced on the ABI 5500XL™.

**[00213]** *RNA in situ Hybridization (RNA-ISH).* Paraffin-embedded tissue blocks were freshly cut and frozen at -80°C. Upon removal from the freezer, slides were baked for 1 hr at 60°C and fixed in %10 formaldehyde for 1 hr at room temperature (RT). Paraffin was removed using Histo-Clear™ and RNA-ISH™ was performed according to the Affymetrix QuantiGene ViewRNA ISH Tissue-2 Plex Assay™. Tissue sections were permeabilized by pretreating in buffer solution for 10 min at 95°C and digested with protease for 10 min, before being fixed at RT in 5% formaldehyde. Target probe sets were applied and hybridized to the tissue by incubating for 2 hr at 40°C. Type 1 probes were used at a dilution of 1:50 and included Aldh1a2 (VB1-14197), Dcn (VB1-14962), Klf4 (VB1-14988), Igfbp5 (VB1-14987), and Sparc (VB1-14196). Type 6 probes included EGFP (VF6-13336) at 1:50 and pooled Krt8 (VB6-11060) and Krt18 (VB6-11059) at 1:100 each. Signal was amplified through the sequential hybridization of PreAmplifier and Amplifier QT mixes to the target probe set. Target mRNA molecules were detected by applying Type 6 Label Probe with Fast Blue substrate and Type 1 Label Probe with Fast Red substrate. Tissue was counterstained with Gill's Hematoxylin for 10 sec at RT. DAPI (Invitrogen, D3571; 3.0 µg/ml) staining was performed for 1 min. Fluorescence microscopy using a Nikon 90i was used to visualize target mRNAs. Type 1 probes were detected in the Cy3 channel and Type 6 probes in the Cy5 channel. Merged images were generated using NIS-Elements™ software.

**[00214]** *Determination of reads-per-million (rpm)* Color space reads were aligned using tophat™ version 2.0.4 (Trapnell et al., 2009) and bowtie1™ version 0.12.7 with the no-novel-juncs argument set with mouse genome version mm9 and transcriptome defined by the mm9 knownGene table from genome.ucsc.edu. Reads that did not align or aligned to multiple locations in the genome were discarded. The mm9 table knownToLocusLink from genome.ucsc.edu was used to map, if possible, each aligned read to the gene whose exons the read had aligned to. The reads count for each gene was the number of reads that were so mapped to that gene. This count was divided by the total number of reads that were mapped to any gene and multiplied by one million to form the reads-per-million (rpm) count. Rpm rather than rpkm was used because a 3' bias was noted in the alignments.

**[00215]** *Unsupervised hierarchical clustering and principal components analysis.* The minimum of 1 and the smallest positive value of the rpm matrix was added to the rpm matrix to eliminate zeros. The result was then log10 transformed, yielding what is termed the log10(rpm) matrix. The rows (corresponding to genes) of the log10(rpm) matrix with the top 2000 standard deviations were retained and the rest of the rows discarded. The result was then median polished. The result was clustered using agglomerative hierarchical clustering with average linkage with distance metric equal to 1 minus the Pearson correlation coefficient. The principal components of the log10(rpm) matrix

were computed and the coordinates of the samples with respect to the first three principal components were plotted.

**[00216]** *Measures of cellular heterogeneity.* For a collection of clusters of samples, a statistic, M, was defined as the mean over the clusters of the mean over all the pairs of samples in the cluster of the atanh of the correlation coefficient between the two columns of the rpm matrix corresponding to the pair. The “mean intra-cluster correlation coefficient” was defined as tanh(M). The jackknife estimator was used with respect to the samples to estimate a standard deviation, s, of the statistic. The 95% CI was defined as  $\tanh(M \pm s\phi^{-1}(0.975))$ , where  $\phi$  is the cumulative distribution function of the standard normal distribution. To compute a p-value for the null hypothesis that the mean of the distribution of the M statistic for a cluster is the same as the mean of the distribution of the M statistic for a collection of clusters, we let  $p=2(1-\phi(|M1-M2| / \sqrt(s_1^2 + s_2^2)))$ . Of note, bootstrap was performed on the same data as an alternative to jackknife and similar results obtained (data not shown).

**[00217]** *Supervised differential gene expression using rank product.* To find differentially expressed genes between two sets of samples, analysis was begin with the log10(rpm) matrix defined above. Columns corresponding to samples not in either set of samples were removed. Then removed rows for which the 90<sup>th</sup> percentile of the values was less than log10(10) were removed. The RP function of the Bioconductor (Gentleman et al., 2004) RankProd™ package (version 2.28.0) was used to get FDR estimates for both up and down differential expression. Genes were considered to be differentially expressed if their FDR estimate was less than 0.01, but discarded if they were both up and down differentially expressed, if there were any.

**[00218]** *Gene set enrichment.* Enrichment was considered in four gene set collections: (1) all of KEGG™, as found in DAVID™ 6.7 (Huang da et al., 2009), (2) Gene Ontology (GO) using GO\_BP as found in DAVID 6.7, and (3) GO\_CC as found in DAVID 6.7. Sets of genes found to be differentially expressed were tested for enrichment in the gene set collections using a hypergeometric test for each gene set in the collection. The resulting p-values for each collection were converted to FDR estimates using the Benjamini-Hochberg method (Benjamini and Hochberg, 1995).

**[00219]** *Digital removal of all annotated platelet transcripts* The 446 genes whose expression in the log10(rpm) matrix had an absolute value of correlation coefficient greater than 0.6 with the expression of any of the genes in the gene sets named GNATENKO\_PLATELET\_SIGNATURE and TENEDINI\_MEGAKARYOCYTE\_MARKERS in MSigDB v3.1 were removed from the log10(rpm) matrix (defined above). Clustering was then performed as described above.

**[00220] Supplemental Methods References**

Bardeesy, N., Aguirre, A.J., Chu, G.C., Cheng, K.H., Lopez, L.V., Hezel, A.F., Feng, B., Brennan, C., Weissleder, R., Mahmood, U., *et al.* (2006). Both p16(INK4a) and the p19(Arf)-p53 pathway constrain progression of pancreatic adenocarcinoma in the mouse. *Proc Natl Acad Sci U S A* **103**, 5947-5952.

Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B (Methodological)* 57, 289-300.

Gentleman, R.C., Carey, V.J., Bates, D.M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., *et al.* (2004). Bioconductor: open software development for computational biology and bioinformatics. *Genome biology* 5, R80.

Huang da, W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4, 44-57.

Tang, F., Barbacioru, C., Nordman, E., Li, B., Xu, N., Bashkirov, V.I., Lao, K., and Surani, M.A. (2010). RNA-Seq analysis to capture the transcriptome landscape of a single cell. *Nat Protoc* 5, 516-535.

Trapnell, C., Pachter, L., and Salzberg, S.L. (2009). TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25, 1105-1111.

**[00221] EXAMPLE 3**

**[00222]** A comparative analysis of mouse pancreatic CTCs indicated an enrichment of 60 extracellular proteins (Table 6). Evaluation of these particular biomarkers and therapeutic targets was undertaken in human pancreatic circulating tumor cells and the most abundant targets in human pancreatic CTCs are shown (Fig. 7). These not only represent potential biomarkers, but given their nature as proteins on the external surface of tumor cells, they are therapeutic targets. The extracellular proteins of Table 6 can be targeted, e.g. by antibody-based therapeutics (e.g. as in the cases of trastuzumab for HER2, cetuximab for EGFR, and bevacizumab for VEGF) to treat cancer.

**[00223] Table 6:** List of Pancreatic CTC enriched Extracellular Proteins.

OFFICIAL GENE SYMBOL	Gene Name
Abi3bp	ABI gene family, member 3 (NESH) binding protein
Adamts5	similar to a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2); a disintegrin-like and metallopeptidase (reprolysin type) with thrombospondin type 1 motif, 5 (aggrecanase-2)
Adamtsl1	ADAMTS-like 1
Ang	angiogenin, ribonuclease, RNase A family, 5
Arsa	arylsulfatase A
C1rl	complement component 1, r subcomponent-like
C3	complement component 3; similar to complement component C3 prepropeptide, last
C4a	similar to Complement C4 precursor; complement component 4A (Rodgers blood group); similar to complement C4; complement component 4B (Childs blood group)

C4b	similar to Complement C4 precursor; complement component 4A (Rodgers blood group); similar to complement C4; complement component 4B (Chido blood group)
Ccdc80	coiled-coil domain containing 80
Cd109	CD109 antigen
Chi3l1	chitinase 3-like 1
Clec3b	C-type lectin domain family 3, member b
Cmtm3	CKLF-like MARVEL transmembrane domain containing 3
Cmtm7	CKLF-like MARVEL transmembrane domain containing 7
Col14a1	collagen, type XIV, alpha 1
Col1a2	collagen, type I, alpha 2
Col3a1	collagen, type III, alpha 1
Col4a6	collagen, type IV, alpha 6
Csf1	colony stimulating factor 1 (macrophage)
Dag1	dystroglycan 1
Dcn	decorin
Dmkn	dermokine
Fbln1	fibulin 1
Fgf1	fibroblast growth factor 1
Fmod	fibromodulin
Gpc3	glypican 3
Gpc4	glypican 4; similar to Glypican 4
Hmgb1	high mobility group box 1
Ifnar2	interferon (alpha and beta) receptor 2
Igfbp5	insulin-like growth factor binding protein 5
Il16	interleukin 16
Lama4	laminin, alpha 4
Ltbp4	latent transforming growth factor beta binding protein 4
Mfap1a	similar to microfibrillar-associated protein 1A; microfibrillar-associated protein 1A; microfibrillar-associated protein 1B
Nid2	nidogen 2
Ogn	osteoglycin
Pdap1	PDGFA associated protein 1
Pf4	platelet factor 4
Plat	plasminogen activator, tissue
Podn	podocan
Prelp	proline arginine-rich end leucine-rich repeat
Rspo1	R-spondin homolog (Xenopus laevis)
Serpinc1	serine (or cysteine) peptidase inhibitor, clade G, member 1
Slurp1	secreted Ly6/Plaur domain containing 1
Sod3	superoxide dismutase 3, extracellular
Sparc	secreted acidic cysteine rich glycoprotein; similar to Secreted acidic cysteine rich glycoprotein
Spock2	sparc/osteonectin, cwcv and kazal-like domains proteoglycan 2
Spon2	spondin 2, extracellular matrix protein

Sulf1	sulfatase 1
Sulf2	sulfatase 2
Tgfb2	transforming growth factor, beta 2
Tgm2	transglutaminase 2, C polypeptide
Thbd	thrombomodulin
Thbs1	thrombospondin 1; similar to thrombospondin 1
Thsd4	thrombospondin, type I, domain containing 4
Timp2	tissue inhibitor of metalloproteinase 2
Tnxb	tenascin XB
Tpt1	predicted gene 1974; tumor protein, translationally-controlled 1 pseudogene; tumor protein, translationally-controlled 1; predicted gene 14456
Twsg1	twisted gastrulation homolog 1 (Drosophila)
Wnt4	wingless-related MMTV integration site 4

[00224] Extending these CTC enriched genes to human pancreatic, breast, and prostate single cell CTC data identified 5 candidate genes shown in Table 9.

Table 9: Percent of human single CTCs with high expression by RNA-seq

Cancer Type	Percent of Single CTCs > 50 RPM of Expression			
	Pancreas (N=7)	Breast (N=29)	Prostate (N=77)	ALL (N=113)
TPT1	86%	90%	90%	89%
HMGB1	43%	62%	44%	49%
SPON2	43%	7%	45%	35%
SPARC	100%	41%	9%	23%
ARSA	71%	17%	5%	12%

[00225] Focusing on pancreatic cancer, SPARC was selected as an initial gene to evaluate. SPARC RNA-ISH in mouse and human primary tumors (data not shown) demonstrated significant expression in the stromal cells of the tumor that provides essential microenvironmental signals to tumors. Much effort in the field focuses on targeting the stroma of PDAC for therapeutic efficacy [1-4] making SPARC a CTC therapeutic target as well as a stromal directed target. A total of 196/198 (99%) of human pancreatic tumors were positive for SPARC and 36% with clear epithelial tumor cell expression.

[00226] Evaluation of human pancreatic cancer cell lines identified 3 of 5 cell lines with elevated SPARC expression which correlates to increased migratory behavior, a surrogate in vitro assay that correlates with metastatic behavior (Fig. 8).

[00227] Evaluation of SPARC function in human pancreatic cancer was done using short hairpin RNA interferences (shRNA) on the two cell lines with highest SPARC expression (PDAC2 and PDAC3). Multiple in vitro assays were done including proliferation, migration, invasion, scratch, and soft agar. The most profound effects of suppressing SPARC expression was on migratory behavior (Fig. 9 and data not shown), indicating SPARC is not only present in many CTCs, but has functional consequences when inhibited in cell line models.

[00228] Given these data, in vivo tail vein inoculation was performed using PDAC-3 to determine if SPARC knockdown affected metastasis. Initial data at 2 weeks post tail vein injection indicates there is reduced metastatic potential when SPARC is inhibited by shRNA with 83% of control mice with metastases compared to 40% in cell lines with shRNA against SPARC (Fig. 10).

[00229] **Surface Protein Targets**

[00230] Most of the targets identified in Table 9 are secreted factors and analysis of genes annotated as cell surface proteins are summarized in Table 14.

[00231] Table 14: Percent of human single CTCs with high expression of surface protein genes

Cancer type	Percent of Single CTCs > 50 RPM of Expression			
	Pancreas (N=7)	Breast (N=29)	Prostate (N=77)	ALL (N=113)
IL6ST	0%	38%	8%	15%
ARSA	71%	17%	5%	12%
TIMP2	0%	21%	4%	8%
CD55	0%	17%	4%	7%
SULF2	0%	24%	0%	6%
ITGA6	0%	14%	3%	5%
SDC4	0%	14%	3%	5%
CDON	0%	7%	5%	5%
SV2A	14%	3%	1%	3%

[00232] It is contemplated herein that these genes are targets given they would be integrated into the plasma membrane of CTCs. In general, RNA expression of cell surface markers tend to be lower than actual protein levels on cells.

[00233] Contemplated herein are antibodies to **IL6ST, SULF2, and SV2A** for therapeutic utility.

1. **IL6ST** – signal transducer for IL6, LIF, CNTF, and oncostatin M.
  - a. Important for STAT3 activation downstream
  - b. Antibodies against IL6 receptor and IL6 have been developed for human disease including cancer
2. **SULF2** – sulfatase modifies heparin sulfate by removing 6-O-sulfate groups

- a. Expression enriched in cancer progression and metastasis
- b. Drugs have been developed against sulfatase activity and tested with activity in liver cancer models

3. *SV2A* – synaptic vesicle glycoprotein elevated in neuroendocrine cells

- a. A marker of neuroendocrine cells, which appear at the epithelial stromal border of human pancreatic cancer
- b. Neuroendocrine differentiation common feature in cancers and portends to more aggressive disease

[00234] REFERENCES

1. Olive, K.P., M.A. Jacobetz, C.J. Davidson, A. Gopinathan, D. McIntyre, D. Honess, B. Madhu, M.A. Goldgraben, M.E. Caldwell, D. Allard, K.K. Frese, G. Denicola, C. Feig, C. Combs, S.P. Winter, H. Ireland-Zecchini, S. Reichelt, W.J. Howat, A. Chang, M. Dhara, L. Wang, F. Ruckert, R. Grutzmann, C. Pilarsky, K. Izeradjene, S.R. Hingorani, P. Huang, S.E. Davies, W. Plunkett, M. Egorin, R.H. Hruban, N. Whitebread, K. McGovern, J. Adams, C. Iacobuzio-Donahue, J. Griffiths, and D.A. Tuveson, *Inhibition of Hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer*. Science, 2009. 324(5933): p. 1457-61.
2. Neesse, A., P. Michl, K.K. Frese, C. Feig, N. Cook, M.A. Jacobetz, M.P. Lolkema, M. Buchholz, K.P. Olive, T.M. Gress, and D.A. Tuveson, *Stromal biology and therapy in pancreatic cancer*. Gut, 2011. 60(6): p. 861-8.
3. Rasheed, Z.A., W. Matsui, and A. Maitra, *Pathology of pancreatic stroma in PDAC*, in *Pancreatic Cancer and Tumor Microenvironment*, P.J. Grippo and H.G. Munshi, Editors. 2012: Trivandrum (India).
4. Provenzano, P.P., C. Cuevas, A.E. Chang, V.K. Goel, D.D. Von Hoff, and S.R. Hingorani, *Enzymatic targeting of the stroma ablates physical barriers to treatment of pancreatic ductal adenocarcinoma*. Cancer Cell, 2012. 21(3): p. 418-29.
5. Nagasaki, T., M. Hara, H. Nakanishi, H. Takahashi, M. Sato, and H. Takeyama, *Interleukin-6 released by colon cancer-associated fibroblasts is critical for tumour angiogenesis: anti-interleukin-6 receptor antibody suppressed angiogenesis and inhibited tumour-stroma interaction*. Br J Cancer, 2014. 110(2): p. 469-78.
6. Zarogoulidis, P., L. Yarmus, and K. Zarogoulidis, *New insights for IL-6 targeted therapy as an adjuvant treatment for non-small-cell lung cancer*. Ther Deliv, 2013. 4(10): p. 1221-3.
7. Voorhees, P.M., R.F. Manges, P. Sonneveld, S. Jagannath, G. Somlo, A. Krishnan, S. Lentzsch, R.C. Frank, S. Zweegman, P.W. Wijermans, R.Z. Orlowski, B. Kranenburg, B. Hall, T. Casneuf, X. Qin, H. van de Velde, H. Xie, and S.K. Thomas, *A phase 2 multicentre study of*

*siltuximab, an anti-interleukin-6 monoclonal antibody, in patients with relapsed or refractory multiple myeloma.* Br J Haematol, 2013. 161(3): p. 357-66.

8. Betts, B.C., E.T. St Angelo, M. Kennedy, and J.W. Young, *Anti-IL6-receptor-alpha (tocilizumab) does not inhibit human monocyte-derived dendritic cell maturation or alloreactive T-cell responses.* Blood, 2011. 118(19): p. 5340-3.

9. Bayliss, T.J., J.T. Smith, M. Schuster, K.H. Dragnev, and J.R. Rigas, *A humanized anti-IL-6 antibody (ALD518) in non-small cell lung cancer.* Expert Opin Biol Ther, 2011. 11(12): p. 1663-8.

10. Khurana, A., D. Jung-Beom, X. He, S.H. Kim, R.C. Busby, L. Lorenzon, M. Villa, A. Baldi, J. Molina, M.P. Goetz, and V. Shridhar, *Matrix detachment and proteasomal inhibitors diminish Sulf-2 expression in breast cancer cell lines and mouse xenografts.* Clin Exp Metastasis, 2013. 30(4): p. 407-15.

11. Lui, N.S., A. van Zante, S.D. Rosen, D.M. Jablons, and H. Lemjabbar-Alaoui, *SULF2 expression by immunohistochemistry and overall survival in oesophageal cancer: a cohort study.* BMJ Open, 2012. 2(6).

12. Hur, K., T.S. Han, E.J. Jung, J. Yu, H.J. Lee, W.H. Kim, A. Goel, and H.K. Yang, *Up-regulated expression of sulfatases (SULF1 and SULF2) as prognostic and metastasis predictive markers in human gastric cancer.* J Pathol, 2012. 228(1): p. 88-98.

13. Phillips, J.J., E. Huillard, A.E. Robinson, A. Ward, D.H. Lum, M.Y. Polley, S.D. Rosen, D.H. Rowitch, and Z. Werb, *Heparan sulfate sulfatase SULF2 regulates PDGFRalpha signaling and growth in human and mouse malignant glioma.* J Clin Invest, 2012. 122(3): p. 911-22.

14. Zheng, X., X. Gai, S. Han, C.D. Moser, C. Hu, A.M. Shire, R.A. Floyd, and L.R. Roberts, *The human sulfatase 2 inhibitor 2,4-disulfonylphenyl-tert-butylnitrone (OKN-007) has an antitumor effect in hepatocellular carcinoma mediated via suppression of TGFB1/SMAD2 and Hedgehog/GLI1 signaling.* Genes Chromosomes Cancer, 2013. 52(3): p. 225-36.

15. Bandala, C., A. Miliar-Garcia, C.M. Mejia-Barradas, M. Anaya-Ruiz, J.P. Luna-Arias, C.I. Bazan-Mendez, M. Gomez-Lopez, S. Juarez-Mendez, and E. Lara-Padilla, *Synaptic vesicle protein 2 (SV2) isoforms.* Asian Pac J Cancer Prev, 2012. 13(10): p. 5063-7.

16. Ting, D.T., D. Lipson, S. Paul, B.W. Brannigan, S. Akhavanfard, E.J. Coffman, G. Contino, V. Deshpande, A.J. Iafrate, S. Letovsky, M.N. Rivera, N. Bardeesy, S. Maheswaran, and D.A. Haber, *Aberrant overexpression of satellite repeats in pancreatic and other epithelial cancers.* Science, 2011. 331(6017): p. 593-6.

17. Zeng, Y.J., W. Lai, L. Liu, H. Wu, X.X. Luo, J. Wang, and Z.H. Chu, *Prognostic Significance of Neuroendocrine Differentiation in Colorectal Adenocarcinoma After Radical Operation: a Meta-analysis.* J Gastrointest Surg, 2014.

18. Wang, J., B. Wei, C.T. Albarracin, J. Hu, S.C. Abraham, and Y. Wu, *Invasive neuroendocrine carcinoma of the breast: a population-based Study from the Surveillance, Epidemiology and End Results (SEER) database*. BMC Cancer, 2014. 14(1): p. 147.
19. Lipianskaya, J., A. Cohen, C.J. Chen, E. Hsia, J. Squires, Z. Li, Y. Zhang, W. Li, X. Chen, H. Xu, and J. Huang, *Androgen-deprivation therapy-induced aggressive prostate cancer with neuroendocrine differentiation*. Asian J Androl, 2014.
20. Chen, Y., I. Nowak, J. Huang, P.C. Keng, H. Sun, H. Xu, G. Wei, and S.O. Lee, *Erk/MAP kinase signaling pathway and neuroendocrine differentiation of non-small-cell lung cancer*. J Thorac Oncol, 2014. 9(1): p. 50-8.
21. Chang, P.C., T.Y. Wang, Y.T. Chang, C.Y. Chu, C.L. Lee, H.W. Hsu, T.A. Zhou, Z. Wu, R.H. Kim, S.J. Desai, S. Liu, and H.J. Kung, *Autophagy Pathway Is Required for IL-6 Induced Neuroendocrine Differentiation and Chemoresistance of Prostate Cancer LNCaP Cells*. PLoS One, 2014. 9(2): p. e88556.
22. Debes, J.D. and D.J. Tindall, *Mechanisms of androgen-refractory prostate cancer*. N Engl J Med, 2004. 351(15): p. 1488-90.

**[00235] EXAMPLE 5**

**[00236]** Circulating tumor cells (CTCs) are shed from primary tumors into the bloodstream, mediating the hematogenous spread of cancer to distant organs. To define their composition, genomewide expression profiles of CTCs were compared with matched primary tumors in a mouse model of pancreatic cancer, isolating individual CTCs using epitope-independent microfluidic capture, followed by single-cell RNA sequencing. CTCs clustered separately from primary tumors and tumor-derived cell lines, showing lowproliferative signatures, enrichment for Aldh1a2, biphenotypic expression of epithelial and mesenchymal markers, and expression of Igfbp5, a gene transcript enriched at the epithelial-stromal interface. Mouse as well as human pancreatic CTCs exhibit a very high expression of stromal-derived extracellular matrix (ECM) proteins, including SPARC, whose knockdown in cancer cells suppresses cell migration and invasiveness. The aberrant expression by CTCs of stromal ECM genes points to their contribution of microenvironmental signals for the spread of cancer to distant organs.

**[00237]** Classical CTCs expressed predominantly the Aldh1a2 isoform, while Aldh1a1 was expressed in a variety of cell types (data not shown). Within single CTCs, there was no correlation between expression of Aldh1 isoforms and either enrichment for the mesenchymal genes (Cdh11, Vim) or loss of epithelial genes (Cdh1, Muc1), indicating that stem cell and EMT markers are not intrinsically linked in CTCs. Analysis of primary pancreatic tumors for Aldh1a2 using RNA in situ hybridization (RNA-ISH) identified rare epithelial tumor cells expressing this stem cell marker, but

the majority of expression was present within the cancer associated stromal cells (Fig. 12A), consistent with immunohistochemistry for ALDH protein in human PDAC (Rasheed et al., 2010).

**[00238]** Besides the evident diversity of CTCs, shared transcripts were searched for that could provide further insight into their cell of origin within the primary tumor and the mechanisms by which they invade and survive within the bloodstream and ultimately identify potential CTC-specific therapeutic targets. Rigorous criteria were selected to identify the most highly enriched CTC-c transcripts (RP score < 300), expressed at very high levels (>100 rpm) in R90% of all classical CTCs. Three genes met these criteria: Kruppel-like factor 4 (Klf4), one of the key stem cell (iPS) reprogramming factors (Takahashi and Yamanaka, 2006), insulin-like growth factor binding protein 5 (Igfbp5), an extracellular growth factor binding protein and decorin (Dcn). RNA-ISH was utilized in primary tumor specimens to identify the potential colocalization of these three highly enriched CTC genes. In contrast to Aldh1a2, Klf4 is expressed in epithelial components of the primary tumor (Fig. 12B). Igfbp5 is of particular interest, in that it is expressed focally at the tumor epithelial-stromal interface (Fig. 12C). It is contemplated herein that this geographic area is enriched for cancer cells undergoing EMT, contributing to the mixed epithelial/stromal transcriptional programs evident by RNA-seq of single CTCs.

**[00239]** In addition to highly expressing Dcn, CTCs consistently had high levels of multiple ECM gene transcripts. GO analysis of all CTC-enriched genes (Table 3) identified 32 proteinaceous ECM genes (GO:0005578, OR 2.4, q-value 4.8 3 10.3). These genes are normally expressed in reactive stromal cells, rather than in epithelial cancer cells, and while recent studies have highlighted the importance of the stroma in supporting pancreatic cancer pathogenesis and metastasis (Feig et al., 2012; Neesse et al., 2011, 2013; Olive et al., 2009; Provenzano et al., 2012), the expression of these stroma-associated ECM genes within tumor cells in circulation was unexpected. Using RP differential expression analysis, CTCs were compared with purified EGFP-tagged primary tumor single cells (TuGMP3) and bulk tumor samples (tumor cells admixed with reactive stromal cells). Six proteinaceous ECM genes were highly expressed by CTCs and by stromal component, but not by epithelial cells within primary tumors: Dcn, Sparc, Ccdc80, Col1a2, Col3a1, and Timp2 (data not shown). RNA-ISH analysis of both Dcn and Sparc confirmed diffuse expression in stromal elements of mouse primary tumors, with rare areas where these transcripts are colocalized with keratin-expressing cells at the epithelial-stromal border (data not shown).

**[00240]** SPARC is a ECM protein gene. RNA-ISH analysis of 198 primary human PDACs demonstrates abundant stromal cell expression of SPARC transcripts in 99% of cases, with up to a third of tumors with rare epithelial cells expressing this ECM gene product (data not shown). Consistent with these observations, RNA-seq of EGFP-tagged single primary tumor cells (data not shown) identified only 1 of 20 cells (5%) with coexpression of high levels (>100 rpm) of Sparc and Krt19.

**[00241]** In summary, abundant expression of ECM genes is a common feature of all keratin-rich classical CTCs. This is in marked contrast to the primary tumor, where these gene products are secreted by supporting stromal cells and not by the epithelial cancer cells. However, rare cells at the epithelial-stromal interface of primary tumors do appear to express both keratins and ECM genes, consistent with the pattern observed in CTCs themselves.

**[00242]** To confirm the expression of proteinaceous ECM genes by human cancer cells circulating in the bloodstream, single CTCs were isolated from patients with pancreatic (n = 7), breast (n = 29), and prostate (n = 77) cancers and subjected these to single- cell RNA-seq. Six ECM protein genes were highly expressed in human CTCs (>100 rpm in >15% of all CTC samples) (Fig. 13; Table 13). Notably, three genes (SPARC, MGP, SPON2) are ECM glycoproteins, defined as part of the core matrisome (Naba et al., 2012). The core matrisome protein SPARC was particularly enriched in pancreatic CTCs being expressed at high levels (>100 rpm) in 100% of pancreatic CTCs compared to 31% of breast and 9% of prostate CTCs. The notable differences in ECM protein gene expression across human epithelial CTCs suggest microenvironment tissue specificity as well as probable redundancies in ECM protein signaling. Together, the consistent expression of ECM gene family members in human CTCs indicates that their upregulation contributes either to the generation of CTCs from primary tumors or to the survival of cancer cells deprived of microenvironmental signals as they circulate in the bloodstream.

**[00243]** In order to define the functional consequences of SPARC expression in pancreatic cancer cells, a panel of patient-derived, low-passage PDAC cell lines was screened for expression. Two human PDAC cell lines with relatively high SPARC expression were identified (PDAC2 and PDAC3), making it possible to test the consequences of small hairpin RNA (shRNA)-mediated knockdown (Fig. 8, 9, Figs. 16A-16D). Suppression of endogenous SPARC expression in both PDAC2 and PDAC3 cell lines using two independent shRNA constructs did not affect proliferation in 2D cultures or anchorage-independent tumor sphere formation (Figs. 14A-14B, Figs. 16A-16D). However, SPARC knockdown by both shRNAs significantly reduced pancreatic cancer cell migration in wound scratch assays and their invasive properties, as measured by in vitro Boyden assays (data not shown).

**[00244]** Tail vein injection of SPARC-suppressed PDAC3 cells using both shRNA constructs generated significantly fewer lung metastases than cells expressing nontargeting hairpin (shNT) controls (Fig. 14D). Metastases generated from orthotopic pancreatic xenografts were also significantly reduced for SPARC-suppressed PDAC3 cells, as measured by luciferase imaging and normalized for primary tumor size (Fig. 14E). Thus, SPARC expression by pancreatic cancer cells appears to selectively enhance their invasive and migratory properties to augment metastatic virulence. The high levels of SPARC expression evident in virtually all pancreatic CTCs thus raises the possibility that it contributes significantly to the metastatic spread of pancreatic cancer.

**[00245] DISCUSSION**

**[00246]** Described herein is the detailed analysis of CTC composition and diversity in pancreatic cancer, using single-cell RNA-seq. High-quality transcriptomes were achieved in 93 single mouse pancreatic CTCs, which were compared with bulk and single-cell preparations from matched primary tumors and from an immortalized cell line established from the same mouse pancreatic tumor model. The use of the KPC mouse model made it possible to compare simultaneously isolated primary tumor specimens and CTCs, and it allowed measurements of CTC heterogeneity across multiple mice sharing the same Kras/Trp53 genetic drivers. The large number of isolated CTCs and the high quality of the isolated RNA from these cells reflect the application of the CTC-iChip technology, which effectively depletes normal blood components, enriching for CTCs that are untagged and accessible for single-cell manipulation. Finally, the purification of CTCs irrespective of their cell-surface epitopes avoids any bias associated with their purification based on expression of common epithelial markers such as EpCAM.

**[00247]** Together, the observations made herein include the following. (1) CTC expression profiles cluster into three classes, including a major “classical CTC” group, and others that are defined by platelet-derived markers or proliferative signatures. (2) Common features shared by virtually all classical CTCs include expression of both epithelial and mesenchymal markers, the stem cell-associated gene Aldh1a2, and three highly expressed transcripts, Klf4, Igfbp5, and Dcn. The specific localization of Igfbp5-expressing cells at the epithelial-stromal boundary within primary tumors may point to a region that contributes significantly to CTC generation. (3) The most highly enriched CTC-specific transcripts shared by almost all classical CTCs encode extracellular matrix proteins, such as Sparc. (4) Aberrant expression in CTCs of this ECM gene product, which is normally abundant in the tumor stromal compartment, is observed in both mouse and human pancreatic CTCs, and its knockdown attenuates cancer cell migration and invasion in reconstituted systems. (Fig. 15) Compared with RNA-seq of partially purified, bulk CTC populations, which required digital subtraction of leukocyte-derived reads (Yu et al., 2012, 2013), the single-cell analysis reported here provides considerably more depth of tumor cell-specific transcript reads, and it allows measurements of CTC heterogeneity.

**[00248]** It is contemplated herein that in addition to the initiating mutations, somatically acquired genetic and epigenetic changes may distinguish CTCs derived from different tumors. Multiple mouse tumors contributed to each of the three distinct clusters of CTCs. Despite their atypical expression pattern, the identification of platelet-associated and proliferative CTC subsets as being tumor-derived is established by their inclusion of lineage-tagged tumor cells. The more characteristic expression pattern exhibited by the classical CTC cluster enabled detailed comparison with primary tumor cells, thereby providing further insight into the origin and properties of CTCs.

**[00249]** Mouse pancreatic classical CTCs uniformly lose expression of the epithelial marker E-cadherin (Cdh1), a key feature of epithelial-to-mesenchymal transition. However, the cells do not lose expression of other epithelial markers, such as cytokeratins, nor is there a consistent increase in classical mesenchymal markers such as vimentin. As such, most classical CTCs appear arrested in a biphenotypic state. Despite their expression of cytokeratins, which are present in the epithelial components of the primary tumor, most other highly expressed markers in CTCs are shared with the stromal component of the primary tumor. Among these stromal genes is Aldh1a2 (Rasheed and Matsui, 2012; Rasheed et al., 2010). A provocative observation relating to the shared epithelial and mesenchymal state of classical CTCs is their virtually universal (93%) expression of Igfbp5, which is uniquely expressed in a small subpopulation of cells at the epithelial/stromal interface within primary tumors. This raises the possibility that this critical location within the primary tumor generates a disproportionate fraction of viable CTCs.

**[00250]** The most unexpected observation from the single-CTC RNAseq study is the high abundance of ECM transcripts in the vast majority of classical CTCs. The coexpression of pancreaticcancer-enriched cytokeratins (Krt7 and Krt19) in single cells expressing these ECM gene products excludes the possibility that these represent circulating tumor-derived fibroblasts.

**[00251]** Consistent with the aberrant expression of SPARC in some pancreatic cancer cells, a subset of patient-derived tumor cell lines also coexpress it along with epithelial cytokeratins. The reduction in cell migration and metastatic potential exhibited by these pancreatic cell lines following SPARC knockdown indicates that it contributes to CTC-mediated metastasis. It is contemplated herein that Sparc expression contributes to metastasis, but inherent redundancies in ECM protein expression may mitigate this effect in some embodiments.

**[00252]** Considerable effort has been directed to targeting the pancreatic cancer stroma as a means of improving delivery of chemotherapeutics as well as stripping tumor cells of their supportive microenvironment (Neesse et al., 2011; Olive et al., 2009; Provenzano et al., 2012; Rasheed et al., 2012). The findings described herein, e.g., that these gene products are also expressed by CTCs themselves suggests a remarkable level of cellular plasticity. To the extent that invasive properties of CTCs are mediated in part by expression of such ECM proteins, it also raises the possibility of targeting cancer cells in the blood.

**[00253]** **Table 13: Human CTC ECM Gene Expression**

		Percent of Samples > 100 RPM			
Count	ECM Gene Symbol	All CTCs	PDAC CTCs	Breast CTCs	Prostate CTCs
1	ANXA2	36.3%	0.0%	51.7%	33.8%
2	SPON2	29.2%	0.0%	3.4%	41.6%
3	LGALS3	22.1%	42.9%	37.9%	14.3%
4	SPARC	21.2%	100.0%	31.0%	10.4%

5	LGALS3BP	16.8%	0.0%	34.5%	11.7%
6	MGP	15.9%	57.1%	44.8%	1.3%
7	LAMC1	15.0%	0.0%	6.9%	19.5%
8	SMC3	15.0%	42.9%	17.2%	11.7%
9	CALR	14.2%	0.0%	6.9%	18.2%
10	TIMP1	13.3%	14.3%	27.6%	7.8%
11	MMP24	11.5%	0.0%	10.3%	13.0%
12	DAG1	10.6%	0.0%	20.7%	7.8%
13	ERBB2IP	10.6%	14.3%	20.7%	6.5%
14	MMP19	10.6%	0.0%	10.3%	11.7%
15	AGRN	8.8%	0.0%	6.9%	10.4%
16	CRTAP	8.8%	0.0%	6.9%	10.4%
17	COL24A1	8.0%	57.1%	17.2%	0.0%
18	ANG	7.1%	0.0%	0.0%	10.4%
19	MFAP1	7.1%	0.0%	6.9%	7.8%
20	VWF	7.1%	14.3%	17.2%	2.6%
21	VWA1	7.1%	0.0%	3.4%	9.1%
22	TIMP2	6.2%	0.0%	13.8%	3.9%
23	ECM1	6.2%	0.0%	24.1%	0.0%
24	LTBP1	6.2%	28.6%	10.3%	2.6%
25	LGALS1	6.2%	0.0%	10.3%	5.2%
26	SERPINA1	6.2%	0.0%	20.7%	1.3%
27	SPOCK1	6.2%	14.3%	0.0%	7.8%
28	TFF3	6.2%	0.0%	17.2%	2.6%
29	NPNT	5.3%	0.0%	3.4%	6.5%
30	TFIP11	5.3%	14.3%	6.9%	3.9%
31	COL9A2	4.4%	0.0%	0.0%	6.5%
32	COL6A1	4.4%	0.0%	0.0%	6.5%
33	FN1	4.4%	14.3%	10.3%	1.3%
34	LAD1	4.4%	0.0%	10.3%	2.6%
35	LAMA1	4.4%	14.3%	3.4%	3.9%
36	LAMB2	4.4%	0.0%	10.3%	2.6%
37	MATN2	4.4%	14.3%	3.4%	3.9%
38	ZP3	4.4%	0.0%	0.0%	6.5%
39	ADAMTSL3	3.5%	28.6%	3.4%	1.3%
40	FRAS1	3.5%	14.3%	0.0%	3.9%
41	TIMP3	3.5%	0.0%	3.4%	3.9%
42	DST	3.5%	0.0%	6.9%	2.6%
43	GFOD2	3.5%	14.3%	0.0%	3.9%
44	LAMA3	3.5%	14.3%	0.0%	3.9%
45	LAMB1	3.5%	14.3%	0.0%	3.9%
46	MMP7	3.5%	0.0%	0.0%	5.2%
47	ANGPTL4	2.7%	0.0%	0.0%	3.9%
48	BMP4	2.7%	0.0%	0.0%	3.9%
49	LTBP2	2.7%	28.6%	3.4%	0.0%
50	LEPRE1	2.7%	0.0%	0.0%	3.9%
51	LUM	2.7%	0.0%	0.0%	3.9%
52	NID2	2.7%	14.3%	6.9%	0.0%
53	SLC1A3	2.7%	28.6%	0.0%	1.3%
54	TECTA	2.7%	14.3%	3.4%	1.3%
55	THSD4	2.7%	0.0%	6.9%	1.3%

56	ADAMTS15	1.8%	0.0%	6.9%	0.0%
57	USH2A	1.8%	14.3%	3.4%	0.0%
58	APLP1	1.8%	0.0%	0.0%	2.6%
59	COL4A3	1.8%	14.3%	3.4%	0.0%
60	COL7A1	1.8%	0.0%	3.4%	1.3%
61	COL11A1	1.8%	0.0%	6.9%	0.0%
62	COL11A2	1.8%	0.0%	0.0%	2.6%
63	COL15A1	1.8%	28.6%	0.0%	0.0%
64	CTGF	1.8%	0.0%	0.0%	2.6%
65	CRISP3	1.8%	0.0%	0.0%	2.6%
66	DCN	1.8%	0.0%	0.0%	2.6%
67	ENTPD2	1.8%	0.0%	0.0%	2.6%
68	FMOD	1.8%	0.0%	3.4%	1.3%
69	GPC1	1.8%	0.0%	0.0%	2.6%
70	HSPG2	1.8%	0.0%	0.0%	2.6%
71	LAMA5	1.8%	0.0%	3.4%	1.3%
72	LAMC2	1.8%	14.3%	0.0%	1.3%
73	MMP10	1.8%	0.0%	3.4%	1.3%
74	MMP12	1.8%	0.0%	0.0%	2.6%
75	NTN4	1.8%	0.0%	6.9%	0.0%
76	NAV2	1.8%	0.0%	6.9%	0.0%
77	PAPLN	1.8%	0.0%	3.4%	1.3%
78	SFTPA2	1.8%	0.0%	0.0%	2.6%
79	VCAN	1.8%	14.3%	0.0%	1.3%
80	ADAMTS13	0.9%	0.0%	3.4%	0.0%
81	ADAMTS3	0.9%	14.3%	0.0%	0.0%
82	ADAMTS5	0.9%	14.3%	0.0%	0.0%
83	ADAMTSL4	0.9%	0.0%	0.0%	1.3%
84	EFEMP1	0.9%	0.0%	3.4%	0.0%
85	EFEMP2	0.9%	0.0%	3.4%	0.0%
86	EGFLAM	0.9%	14.3%	0.0%	0.0%
87	KAL1	0.9%	0.0%	0.0%	1.3%
88	KAZALD1	0.9%	0.0%	0.0%	1.3%
89	MAMDC2	0.9%	14.3%	0.0%	0.0%
90	SMOC1	0.9%	0.0%	0.0%	1.3%
91	SMOC2	0.9%	0.0%	0.0%	1.3%
92	ACHE	0.9%	0.0%	0.0%	1.3%
93	AMTN	0.9%	0.0%	3.4%	0.0%
94	ANXA2P2	0.9%	0.0%	3.4%	0.0%
95	CPZ	0.9%	0.0%	3.4%	0.0%
96	CHADL	0.9%	0.0%	0.0%	1.3%
97	COCH	0.9%	0.0%	0.0%	1.3%
98	COL6A6	0.9%	14.3%	0.0%	0.0%
99	COL1A2	0.9%	0.0%	3.4%	0.0%
100	COL2A1	0.9%	0.0%	0.0%	1.3%
101	COL4A1	0.9%	14.3%	0.0%	0.0%
102	COL4A2	0.9%	0.0%	0.0%	1.3%
103	COL4A6	0.9%	0.0%	0.0%	1.3%
104	COL5A1	0.9%	14.3%	0.0%	0.0%
105	COL6A2	0.9%	0.0%	0.0%	1.3%
106	COL8A1	0.9%	14.3%	0.0%	0.0%

107	COL12A1	0.9%	14.3%	0.0%	0.0%
108	COL14A1	0.9%	14.3%	0.0%	0.0%
109	COL19A1	0.9%	14.3%	0.0%	0.0%
110	COL17A1	0.9%	14.3%	0.0%	0.0%
111	COL22A1	0.9%	14.3%	0.0%	0.0%
112	ENTPD1	0.9%	14.3%	0.0%	0.0%
113	FBN2	0.9%	0.0%	0.0%	1.3%
114	FBN3	0.9%	0.0%	3.4%	0.0%
115	FBLN1	0.9%	14.3%	0.0%	0.0%
116	FBLN7	0.9%	0.0%	0.0%	1.3%
117	GPC4	0.9%	0.0%	3.4%	0.0%
118	HMCN1	0.9%	14.3%	0.0%	0.0%
119	IMPG1	0.9%	14.3%	0.0%	0.0%
120	IMPG2	0.9%	0.0%	3.4%	0.0%
121	LAMA2	0.9%	0.0%	3.4%	0.0%
122	LAMB3	0.9%	14.3%	0.0%	0.0%
123	MEPE	0.9%	0.0%	3.4%	0.0%
124	MMP1	0.9%	14.3%	0.0%	0.0%
125	MMP2	0.9%	0.0%	3.4%	0.0%
126	MMP25	0.9%	0.0%	0.0%	1.3%
127	MMP3	0.9%	0.0%	3.4%	0.0%
128	MMP9	0.9%	14.3%	0.0%	0.0%
129	OGN	0.9%	14.3%	0.0%	0.0%
130	PI3	0.9%	0.0%	0.0%	1.3%
131	PRELP	0.9%	14.3%	0.0%	0.0%
132	PTPRZ1	0.9%	14.3%	0.0%	0.0%
133	RELN	0.9%	0.0%	3.4%	0.0%
134	ADAMTSL2	0.9%	0.0%	0.0%	1.3%
135	TGFBI	0.9%	0.0%	3.4%	0.0%
136	UCMA	0.9%	0.0%	3.4%	0.0%
137	VIT	0.9%	0.0%	3.4%	0.0%
138	WNT10A	0.9%	14.3%	0.0%	0.0%
139	WNT10B	0.9%	0.0%	0.0%	1.3%
140	WNT11	0.9%	0.0%	3.4%	0.0%
141	WNT4	0.9%	0.0%	0.0%	1.3%
142	ZP2	0.9%	14.3%	0.0%	0.0%
143	ADAMTS1	0.0%	0.0%	0.0%	0.0%
144	ADAMTS10	0.0%	0.0%	0.0%	0.0%
145	ADAMTS12	0.0%	0.0%	0.0%	0.0%
146	ADAMTS14	0.0%	0.0%	0.0%	0.0%
147	ADAMTS16	0.0%	0.0%	0.0%	0.0%
148	ADAMTS17	0.0%	0.0%	0.0%	0.0%
149	ADAMTS18	0.0%	0.0%	0.0%	0.0%
150	ADAMTS19	0.0%	0.0%	0.0%	0.0%
151	ADAMTS2	0.0%	0.0%	0.0%	0.0%
152	ADAMTS20	0.0%	0.0%	0.0%	0.0%
153	ADAMTS4	0.0%	0.0%	0.0%	0.0%
154	ADAMTS6	0.0%	0.0%	0.0%	0.0%
155	ADAMTS8	0.0%	0.0%	0.0%	0.0%
156	ADAMTS9	0.0%	0.0%	0.0%	0.0%
157	ADAMTS1	0.0%	0.0%	0.0%	0.0%

158	ADAMTS5	0.0%	0.0%	0.0%	0.0%
159	CD248	0.0%	0.0%	0.0%	0.0%
160	DGCR6	0.0%	0.0%	0.0%	0.0%
161	EGFL6	0.0%	0.0%	0.0%	0.0%
162	EMID1	0.0%	0.0%	0.0%	0.0%
163	FREM1	0.0%	0.0%	0.0%	0.0%
164	FREM2	0.0%	0.0%	0.0%	0.0%
165	RELL2	0.0%	0.0%	0.0%	0.0%
166	SPARCL1	0.0%	0.0%	0.0%	0.0%
167	ACAN	0.0%	0.0%	0.0%	0.0%
168	AMBN	0.0%	0.0%	0.0%	0.0%
169	AMELX	0.0%	0.0%	0.0%	0.0%
170	AMELY	0.0%	0.0%	0.0%	0.0%
171	ASPN	0.0%	0.0%	0.0%	0.0%
172	BGN	0.0%	0.0%	0.0%	0.0%
173	BCAN	0.0%	0.0%	0.0%	0.0%
174	CRTAC1	0.0%	0.0%	0.0%	0.0%
175	CILP2	0.0%	0.0%	0.0%	0.0%
176	CILP	0.0%	0.0%	0.0%	0.0%
177	COMP	0.0%	0.0%	0.0%	0.0%
178	CHL1	0.0%	0.0%	0.0%	0.0%
179	CHI3L1	0.0%	0.0%	0.0%	0.0%
180	CHAD	0.0%	0.0%	0.0%	0.0%
181	C6orf15	0.0%	0.0%	0.0%	0.0%
182	CCDC80	0.0%	0.0%	0.0%	0.0%
183	CTHRC1	0.0%	0.0%	0.0%	0.0%
184	COL1A1	0.0%	0.0%	0.0%	0.0%
185	COL3A1	0.0%	0.0%	0.0%	0.0%
186	COL4A4	0.0%	0.0%	0.0%	0.0%
187	COL4A5	0.0%	0.0%	0.0%	0.0%
188	COL9A1	0.0%	0.0%	0.0%	0.0%
189	COL9A3	0.0%	0.0%	0.0%	0.0%
190	COL5A2	0.0%	0.0%	0.0%	0.0%
191	COL5A3	0.0%	0.0%	0.0%	0.0%
192	COL6A3	0.0%	0.0%	0.0%	0.0%
193	COL8A2	0.0%	0.0%	0.0%	0.0%
194	COL10A1	0.0%	0.0%	0.0%	0.0%
195	COL16A1	0.0%	0.0%	0.0%	0.0%
196	COL18A1	0.0%	0.0%	0.0%	0.0%
197	COL21A1	0.0%	0.0%	0.0%	0.0%
198	COL27A1	0.0%	0.0%	0.0%	0.0%
199	COL28A1	0.0%	0.0%	0.0%	0.0%
200	COLQ	0.0%	0.0%	0.0%	0.0%
201	DMP1	0.0%	0.0%	0.0%	0.0%
202	DSPP	0.0%	0.0%	0.0%	0.0%
203	DPT	0.0%	0.0%	0.0%	0.0%
204	ELN	0.0%	0.0%	0.0%	0.0%
205	EMILIN1	0.0%	0.0%	0.0%	0.0%
206	EMILIN2	0.0%	0.0%	0.0%	0.0%
207	EMILIN3	0.0%	0.0%	0.0%	0.0%
208	ENAM	0.0%	0.0%	0.0%	0.0%

209	EPYC	0.0%	0.0%	0.0%	0.0%
210	ECM2	0.0%	0.0%	0.0%	0.0%
211	FBN1	0.0%	0.0%	0.0%	0.0%
212	FGF1	0.0%	0.0%	0.0%	0.0%
213	FGF9	0.0%	0.0%	0.0%	0.0%
214	FLRT1	0.0%	0.0%	0.0%	0.0%
215	FLRT2	0.0%	0.0%	0.0%	0.0%
216	FLRT3	0.0%	0.0%	0.0%	0.0%
217	FBLN2	0.0%	0.0%	0.0%	0.0%
218	FBLN5	0.0%	0.0%	0.0%	0.0%
219	GPLD1	0.0%	0.0%	0.0%	0.0%
220	GPC2	0.0%	0.0%	0.0%	0.0%
221	GPC3	0.0%	0.0%	0.0%	0.0%
222	GPC5	0.0%	0.0%	0.0%	0.0%
223	GPC6	0.0%	0.0%	0.0%	0.0%
224	HAPLN1	0.0%	0.0%	0.0%	0.0%
225	HAPLN2	0.0%	0.0%	0.0%	0.0%
226	HAPLN3	0.0%	0.0%	0.0%	0.0%
227	HAPLN4	0.0%	0.0%	0.0%	0.0%
228	KERA	0.0%	0.0%	0.0%	0.0%
229	LAMA4	0.0%	0.0%	0.0%	0.0%
230	LAMB4	0.0%	0.0%	0.0%	0.0%
231	LAMC3	0.0%	0.0%	0.0%	0.0%
232	LTBP4	0.0%	0.0%	0.0%	0.0%
233	LOX	0.0%	0.0%	0.0%	0.0%
234	LOXL1	0.0%	0.0%	0.0%	0.0%
235	MATN1	0.0%	0.0%	0.0%	0.0%
236	MATN3	0.0%	0.0%	0.0%	0.0%
237	MMP11	0.0%	0.0%	0.0%	0.0%
238	MMP13	0.0%	0.0%	0.0%	0.0%
239	MMP16	0.0%	0.0%	0.0%	0.0%
240	MMP17	0.0%	0.0%	0.0%	0.0%
241	MMP20	0.0%	0.0%	0.0%	0.0%
242	MMP23A	0.0%	0.0%	0.0%	0.0%
243	MMP26	0.0%	0.0%	0.0%	0.0%
244	MMP27	0.0%	0.0%	0.0%	0.0%
245	MMP28	0.0%	0.0%	0.0%	0.0%
246	MMP8	0.0%	0.0%	0.0%	0.0%
247	MFAP5	0.0%	0.0%	0.0%	0.0%
248	MFAP2	0.0%	0.0%	0.0%	0.0%
249	MFAP4	0.0%	0.0%	0.0%	0.0%
250	MUC4	0.0%	0.0%	0.0%	0.0%
251	MMRN2	0.0%	0.0%	0.0%	0.0%
252	NTN1	0.0%	0.0%	0.0%	0.0%
253	NTN3	0.0%	0.0%	0.0%	0.0%
254	NID1	0.0%	0.0%	0.0%	0.0%
255	NYX	0.0%	0.0%	0.0%	0.0%
256	ODAM	0.0%	0.0%	0.0%	0.0%
257	OPTC	0.0%	0.0%	0.0%	0.0%
258	OMD	0.0%	0.0%	0.0%	0.0%
259	OTOA	0.0%	0.0%	0.0%	0.0%

260	POSTN	0.0%	0.0%	0.0%	0.0%
261	PODN	0.0%	0.0%	0.0%	0.0%
262	PODNL1	0.0%	0.0%	0.0%	0.0%
263	PRSS36	0.0%	0.0%	0.0%	0.0%
264	RPTN	0.0%	0.0%	0.0%	0.0%
265	RBP3	0.0%	0.0%	0.0%	0.0%
266	SPN	0.0%	0.0%	0.0%	0.0%
267	ADAMTS7	0.0%	0.0%	0.0%	0.0%
268	SPOCK2	0.0%	0.0%	0.0%	0.0%
269	SPOCK3	0.0%	0.0%	0.0%	0.0%
270	SPON1	0.0%	0.0%	0.0%	0.0%
271	SFTPA1	0.0%	0.0%	0.0%	0.0%
272	SFTPД	0.0%	0.0%	0.0%	0.0%
273	TECTB	0.0%	0.0%	0.0%	0.0%
274	TNC	0.0%	0.0%	0.0%	0.0%
275	TNN	0.0%	0.0%	0.0%	0.0%
276	TNR	0.0%	0.0%	0.0%	0.0%
277	TNXB	0.0%	0.0%	0.0%	0.0%
278	THBS4	0.0%	0.0%	0.0%	0.0%
279	TFPI2	0.0%	0.0%	0.0%	0.0%
280	TGFB1	0.0%	0.0%	0.0%	0.0%
281	TINAG	0.0%	0.0%	0.0%	0.0%
282	TNFRSF11B	0.0%	0.0%	0.0%	0.0%
283	VEGFA	0.0%	0.0%	0.0%	0.0%
284	VTN	0.0%	0.0%	0.0%	0.0%
285	VWC2	0.0%	0.0%	0.0%	0.0%
286	WNT2	0.0%	0.0%	0.0%	0.0%
287	WNT1	0.0%	0.0%	0.0%	0.0%
288	WNT16	0.0%	0.0%	0.0%	0.0%
289	WNT2B	0.0%	0.0%	0.0%	0.0%
290	WNT3	0.0%	0.0%	0.0%	0.0%
291	WNT3A	0.0%	0.0%	0.0%	0.0%
292	WNT5A	0.0%	0.0%	0.0%	0.0%
293	WNT5B	0.0%	0.0%	0.0%	0.0%
294	WNT6	0.0%	0.0%	0.0%	0.0%
295	WNT7A	0.0%	0.0%	0.0%	0.0%
296	WNT7B	0.0%	0.0%	0.0%	0.0%
297	WNT8A	0.0%	0.0%	0.0%	0.0%
298	WNT8B	0.0%	0.0%	0.0%	0.0%
299	WNT9A	0.0%	0.0%	0.0%	0.0%
300	WNT9B	0.0%	0.0%	0.0%	0.0%
301	ZP1	0.0%	0.0%	0.0%	0.0%
302	ZP4	0.0%	0.0%	0.0%	0.0%

[00254] Table 10: Most significant Gene Sets Enriched in CTC-pro vs. CTC-c

q-value &lt; 0.01

Source	Term	Count	Odds Ratio	Benjamini (q-value)

GOTERM_BP_FAT	GO:0002495~antigen processing and presentation of peptide antigen via MHC class II	5	59.81	6.97E-04
GOTERM_BP_FAT	GO:0019886~antigen processing and presentation of exogenous peptide antigen via MHC class II	5	59.81	6.97E-04
GOTERM_BP_FAT	GO:0002504~antigen processing and presentation of peptide or polysaccharide antigen via MHC class II	5	50.36	7.34E-04
GOTERM_BP_FAT	GO:0002478~antigen processing and presentation of exogenous peptide antigen	5	41.60	1.10E-03
GOTERM_BP_FAT	GO:0019884~antigen processing and presentation of exogenous antigen	5	34.18	1.87E-03
GOTERM_BP_FAT	GO:0048002~antigen processing and presentation of peptide antigen	5	27.34	3.72E-03
GOTERM_BP_FAT	GO:0001775~cell activation	9	7.00	3.82E-03
GOTERM_BP_FAT	GO:0019882~antigen processing and presentation	6	13.20	7.40E-03

[00255] Table 11: Most significant Gene Sets Enriched in CTC-plt vs. CTC-c

q-value &lt; 0.01

Source	Term	Count	Odds Ratio	Benjamini (q-value)
GOTERM_BP_FAT	GO:0042060~wound healing	18	7.8	1.86E-07
GOTERM_BP_FAT	GO:0007596~blood coagulation	15	10.4	9.31E-08
GOTERM_BP_FAT	GO:0050817~coagulation	15	10.4	9.31E-08
GOTERM_BP_FAT	GO:0007599~hemostasis	15	10.3	7.59E-08
GOTERM_BP_FAT	GO:0050878~regulation of body fluid levels	15	8.2	1.30E-06
GOTERM_BP_FAT	GO:0030029~actin filament-based process	20	5.5	1.14E-06
GOTERM_BP_FAT	GO:0007010~cytoskeleton organization	26	3.9	3.95E-06
GOTERM_BP_FAT	GO:0030036~actin cytoskeleton organization	18	5.3	1.11E-05
GOTERM_BP_FAT	GO:0009611~response to wounding	26	3.6	1.02E-05
GOTERM_BP_FAT	GO:0007155~cell adhesion	33	2.9	2.86E-05
GOTERM_BP_FAT	GO:0022610~biological adhesion	33	2.8	2.70E-05
GOTERM_BP_FAT	GO:0001775~cell activation	19	3.7	4.70E-04
GOTERM_BP_FAT	GO:0030168~platelet activation	6	18.2	1.68E-03
GOTERM_BP_FAT	GO:0007229~integrin-mediated signaling pathway	10	6.4	2.95E-03
GOTERM_BP_FAT	GO:0016192~vesicle-mediated transport	25	2.6	3.81E-03
MSigDBv3.1 CGP	GNATENKO PLATELET SIGNATURE	20	55.1	3.91E-24
MSigDBv3.1 CGP	TENEDINI MEGAKARYOCYTE MARKERS	14	15.3	1.35E-11
MSigDBv3.1 CP:REACTOME	REACTOME FACTORS INVOLVED IN MEGAKARYOCYTE DEVELOPMENT	6	2.9	2.25E-02

	AND PLATELET PRODUCTION			
--	-------------------------	--	--	--

[00256] Table 12: Significantly Expressed Genes by Rank Product (FDR &lt; 0.01)

Count	CTC-c vs Primary Tumor Enriched Gene	Primary Tumor vs CTC-c Enriched Gene	CTC-plt vs CTC-c	CTC-pro vs CTC-c	Primary Tumor vs WBC	WBC vs Primary Tumor	CTC-c vs WBC	WBC vs CTC
1	Upk3b	Tff2	Clec1b	kg:uc007pge.1	Wfdc2	Ppbp	Olfr1033	Beta-s
2	Ier2	Wfdc2	AU023871	kg:uc007pgd.1	Spp1	Alas2	Crip1	Alas2
3	Egr1	Lamb3	Alox12	kg:uc007pgf.1	Cct3	Nrgn	Ppp1r12a	Hbb-b1
4	Nkain4	Lad1	Itga2b	kg:uc007pgg.1	Itga3	Cd9	Vcp	Il1b
5	Igfbp5	Dmbt1	Ppbp	Igj	Gsto1	Csf3r	Klf9	Ppbp
6	Slc6a4	Npy	Gng11	kg:uc012enb.1	Mmp2	Il1b	Mprip	Hba-a2
7	Klf4	Pmepa1	Vwf	2010001M09Rik	Mfge8	Gdpd3	Sdc4	kg:uc007pgs.1
8	Tmem221	Kcnn4	Pf4	kg:uc009cfw.1	Capg	Ms4a1	Gprc5a	kg:uc011yvj.1
9	Arl4d	Serinc2	Fcer1g	kg:uc007pgi.1	Cd63	Hbb-b1	Vat1	Coro1a
10	Lrrn4	5730559C18Rik	Tmem40	kg:uc007pgh.1	Stub1	Beta-s	Wdr92	Cd74
11	Cldn15	Muc1	Hba-a2	kg:uc007yos.1	Lad1	kg:uc007pgs.1	S100a11	Gdpd3
12	Gpm6a	Chi3l3	Stom	Coro1a	Myo1h	kg:uc011yvj.1	Clic4	Ccndbp1
13	Atf3	Pglyrp1	Beta-s	Pou2af1	Igfbp7	Rprl1	Dync1i2	kg:uc009cfw.1
14	Ptma	Arl4c	Plek	kg:uc011yvj.1	Kcnn4	Pfn1	Nfkbia	kg:uc012enb.1
15	Slc9a3r1	Spp1	Srgn	Glipr1	D8Ertd738e	Clec1b	Cyp2s1	Ptprc
16	Fos	Col15a1	Myl9	Cd52	Lamb3	Ptprc	Esam	Csf3r
17	Tmem119	C1qb	Cd84	Cd79b	Chi3l3	Stim1	Surf4	Rac2
18	Ptgis	Tnnt2	F5	Sec11c	Arl4c	Ccndbp1	Krt19	Rprl1
19	Dcn	Gkn3	Trem1	Tnfrsf17	Col18a1	Cap1	Bsg	H2-Ab1
20	Gbp2	Onecut2	Hbb-b1	Krr1	Atox1	Cd79b	Tm4sf1	Epb4.1
21	Dmkn	Mmp7	Itgb3	Gmfg	Ly6a	Alox12	Lgals3	Lyz2
22	Sdc4	Cd74	Gp9	Ccr9	Dmbt1	Hba-a2	Clic1	Ctla2b
23	Il1dr2	Ctss	Mpl	Pycard	Dync1h1	Ube2l6	Capns1	Pld4
24	Akap2	Lamc2	Ctla2a	Derl3	Adipor2	Cat	Igfbp6	kg:uc007pgt.1

25	Gfpt2	Olfml3	Tubb1	Rac2	Rpl37	Faim3	Rrbp1	Gng11
26	Klf6	Lgals4	Mylk	Srgn	Kctd10	Dusp1	Calr	Mepce
27	Btg2	Lcn2	F13a1	Cytip	Col15a1	kg:uc007 pgt.1	Rtf1	Tyrobp
28	Myl7	Ly6a	Slamf1	Edem2	Surf4	E2f2	Ildr2	Isca1
29	Igfbp6	Pak1	Rgs10	Itgb7	Dad1	Phospho 1	Mark 2	281045 3I06Rik
30	Gpr133	Capn5	Mkrn1	Lsp1	Col4a1	Abi3	Mt1	Slc30a9
31	Oasl2	Ptprn	Laptm5	Lcp1	Ap2s1	Sorl1	Akr1b 3	Treml2
32	Pfn1	Reg3b	1810058I2 4Rik	Cyfip2	Sdc1	Treml2	Gm66 44	Srgn
33	Cap1	Fmnl3	Itgb2	Nans	Rpl35	Cytip	Nkain 4	Dcaf12
34	Nfkbia	Sdc1	Slc2a3	Slamf7	Sec61a 1	B2m	Ppp2c a	Plek
35	Malat1	Prom1	Pcmt1	Eli2	Rras	Fyb	Akap 2	Cat
36	Rarres2	Ankrd50	Gp5	H2-Eb1	Oraov1	Peli1	Hspb 1	Alox12
37	Rspo1	Ccl6	Ube2o	Creld2	Ndufa2	Plek	Ptgis	Fech
38	Espn	Slc4a11	5430417L 22Rik	Cd74	Anapc2	N4bp3	Msln	Rbm5
39	Klf9	Oraov1	Ptpn18	Blnk	Pitpna	Fam117a	Emp2	Cd97
40	Zbtb7c	Aldh1l1	Lat	Fmnl1	Psap	Srgn	Capn 2	March8
41	Brd2	Slc20a1	Fermt3	Snrnp70	Atp5j2	Sept9	Rhoc	Pnpo
42	Olfr1033	Cldn7	Nrgn	Sec61b	Onecut 2	kg:uc012 hdk.1	Ptprf	Phosph o1
43	Wt1	Acsbg1	Mrvi1	Edem1	Hmga1	kg:uc009 vev.1	Bcam	Isg20
44	Esam	Las1l	Lyz2	Tspan13	Pmepa1	Ptprcap	Ogdh	March2
45	kg:uc009igb.1	C1qc	Epb4.1	Psmb8	S100a1 1	kg:uc007 pgq.1	Sparc	Lsp1
46	Tmem151a	Lama5	Rasgrp2	Pim1	Rbp1	kg:uc007 pgr.1	Ahna k	181005 8I24Rik
47	Mgll	Mgat4a	Treml2	Sept1	Rpl36al	kg:uc007 vdl.1	Oasl2	Clec1b
48	Csrnp1	Cldn2	Hist1h4i	Cd48	S100a4	Ctla2b	Wt1	Btg1
49	Cd9	Mcpt2	March2	Sub1	Atp6ap 1	Myl9	Klf4	Laptm5
50	Gjb5	Fxyd3	Ltbp1	Lims1	Ndufs2	Itpr2	Cdkn 1a	Nrgn
51	Lrrc61	Il4ra	Nptn	Ncoa2	Anapc5	kg:uc012 enb.1	Myl7	H2-Aa
52	Wasf2	Itga5	Abtb1	Ctnnbl1	Cox6b1	Isg20	Col1a 2	Fyb
53	Pdpn	Porcn	Ctla2b	Fdps	Krtcap2	Rasal3	Eif4a1	Cd24a
54	kg:uc009ogv.1	Mast3	Prkab2	Ube2j1	Atn1	Gng11	Rbpm s	Fnbp4
55	Sdpr	Scara3	Arhgdib	Mettl1	573055 9C18Rik	kg:uc009 cfw.1	Emp3	Ehbp1l1

56	Gpr64	Atox1	Alas2	Lax1	Pea15a	Tmsb4x	Scaf1 1	Ctla2a
57	Flnc	Arrdc1	Odc1	Rilpl2	Grcc10	Trem1	Col14 a1	Sgk1
58	Add3	Mmp2	Ptpn11	Ctse	Lama5	Fech	Ptrf	Glyr1
59	Gata6	Saa3	Dhcr24	Glrx	Krt18	Epb4.1	Crip2	Myl9
60	Wfdc1	Serpinf1	Mfsd2b	Fut8	Ccnd1	Sgk1	Ubxn 4	Il2rg
61	A130040M12R ik	Sox11	Gp1bb	Al662270	Arhgef5	Dgkq	Eif2s2	Mrps17
62	Ankrd12	Prpsap1	Rbpms2	Gramd3	Golm1	Snap23	S100a 6	Cdr2
63	Adamtsl1	Mcpt1	Fyb	Il2rg	Tff2	Usp25	Hectd 1	Mkrn1
64	C2	Mfge8	Smox	Rasgrp3	Plin2	Kif21b	Zc3h1 5	Gart
65	Prss23	Col18a1	P2rx1	Impdh1	H13	Irs2	Ube2 d3	Lyz1
66	Ube2v1	Lyz2	Otud7b	Plek	Rpl29	Pxk	A130 040M 12Rik	Vwf
67	Cryab	C1qa	kg:uc007t tx.1	Ints5	111003 4A24Rik	Cyp4f18	Cd34	Gadd45 a
68	Pkhd1l1	Acp5	Samd14	Blmh	Trim28	Map4k1	Igfbp 5	Mpp1
69	Rtn1	Angptl4	Clca1	Dnmt1	Ltbp3	Isca1	C1s	Stim1
70	Birc6	Ccnd1	kg:uc007t ty.1	Galk1	Fkbp1a	Itga4	Upk3 b	Psme3
71	Xdh	Asl	Gpr56	kg:uc007hxv.1	Erp29	Dock2	Gpr13 3	Ets1
72	Cd34	Ctxn1	Sh3bgrl2	Ccdc88b	Muc1	Spib	Dab2	Snap23
73	Rab6b	Pgs1	Pttg1ip	Selplg	Lamc2	2810453I 06Rik	Serpi nh1	Arhgdb
74	Dusp1	Anapc2	Nomo1	Sar1b	Plscr3	Cdr2	Upk1 b	Hmha1
75	Clic4	Cp	Gnaz	Lat2	Agrn	Naa16	Sdf4	Itpr2
76	C3	Gpx3	Mmrn1	Slc16a6	Park7	Arhgdb	Ctbp2	Ubl7
77	Rhob	Lama3	Gp1ba	Mki67	Ctnnb1	Cd79a	Psap	Ddx58
78	Mir3064	Rbp1	Sh3bgrl3	Dnajc3	Atp5g1	Rbm27	Arhge f12	Nfkbie
79	Thbd	Cotl1	Slc24a3	H2-Ab1	Eef1g	Lmnb1	Copb 2	Setd7
80	Dpysl2	Nek6	Sord	Ndufs6	Nhp2	Slc25a37	Ctsl	Stk24
81	Cobl	Cpxm1	Nfe2	Actr3	Rrbp1	Klf6	Aldh1 a2	Hvcn1
82	Npr1	Sfrp1	Tuba4a	Etnk1	Sumo3	Hist1h1c	Dcn	Plekha2
83	Dnajb9	Ttr	Zyx	Herpud1	Scyl1	Phip	Timp 3	Psme4
84	Arhgap29	Gsto1	Cnn2	Ptpn7	Cox6a1	Qrfp	Xdh	Ankrd4 4
85	Cav1	Npepl1	Itgb5	Ctss	Krt8	Fermt3	Irf7	B4galt5

	Gbp7	Usmg5	Gata1	Cs	Gsta4	Ptma	Tme m151 a	Phf20
86								
87	Hes1	Polr2l	Hist1h1c	Fbxw7	Ppp1r1 4b	Etv3	Aebp 1	Zc3hav1
88	Gm16897	Sphk1	Tbxas1	Ppp2r5c	Tnk1	Apobr	C2	Rnf11
89	Ppp1r12a	Asxl1	Ptplad2	Znrd1	D19Ws u162e	kg:uc008 ewj.2	Spen	Plk3
90	Sv2a	Ctsh	Bpgm	Rfc2	Ctsl	Malat1	kg:uc 007pf e.1	Fbxw5
91	Ang	Egfl7	Pdlim7	Preb	Timp1	March8	Krt18	Emb
92	Aldh1a2	C1qtnf6	Mmd	Fcer1g	S100a6	Coro1a	Arf4	kg:uc00 7vdl.1
93	Cryl1	Rras	G6b	Dnajb11	Rps15	Rac2	Rab1 4	Taok2
94	Kank1	Lgi4	kg:uc009d uo.1	Slc35b1	Polr2j	Glyr1	Tme m98	Dhrs11
95	2210403K04Ri k	Hmga2	Lyz1	Sin3b	Hspe1	Btg2	Prss2 3	Slc25a3 9
96	kg:uc009okn.1	Cep250	Tacc1	Nktr	Lgals4	Mtf2	Egr1	Csk
97	Osr1	B4galt3	Dap		Edf1	Nfkbie	Perp	Bcl2
98	kg:uc008ewj.2	Tmem223	Mast2		Mtch1	Cd84	Csrp1	kg:uc00 9vev.1
99	kg:uc009tuw.1	Ltbp2	Atp2a3		Rnf187	AW5498 77	Pdpn	Wipf1
100	Gadd45b	Tnfrsf23	Snca		Npy	March2	Pdcd6 ip	Sept9
101	Ablim3	Col7a1	Stx11		Cox5b	Add3	Rpl37	Rnf10
102	Clec3b	Ggct	C030046I 01Rik		Pak1	Ddx50	F11r	Pml
103	Usp25	Rab25	Trpt1		Mmp7	Prkcb	Gpm6 a	Cd9
104	Sntb2	Nedd8	Tsc22d1		Fxyd3	Klf2	Tuba 1a	D4Wsu 53e
105	Rock2	9430023L20R ik	Prkar2b		Cuta	Dcaf12	Ctnna 1	Traf7
106	Col14a1	Arl2	Cd9		Ndufb8	Il2rg	Anxa 8	Pitpnc1
107	Cd200	Wbp1	Pgm2l1		Gps1	Selplg	Tpm1	Mms19
108	kg:uc008ehr.1	H2-Ab1	Gp6		Bud31	Cd37	S100a 16	Naa16
109	Atp2b1	Preb	Pde5a		Ppap2c	Fastkd2	Chmp 4b	Sharpin
110	Exoc4	Sgsm3	Itga6		Dap	Rsad2	Tbrg1	Capza1
111	Abcb1b	Sfn	Itgal		Slc25a1	Msn	C3	Rsad2
112	Nrgn	Prrx2	Edem1		Chaf1a	2010321 M09Rik	Ptgs1	kg:uc01 2hdk.1
113	kg:uc009cvm.1	Ptprk	Isg20		Asxl1	Kif2a	Rhou	Ghitm
114	Ncoa4	Reg1	Cdc42ep5		Jmjd8	Cd97	Cdc42	Csnk1g1
115	Ndufa4	Sdcbp2	Nipal3		Tecr	Hvcn1	Gpx4	Dgkz

116	Upk1b	Pcbd1	Ccdc92		Mgp	Nipsnap3b	Ppib	B2m
117	Jun	Slc25a1	Sort1		Uqcrh	Uba7	Stub1	Irs2
118	Syne2	Vamp5	Ly6g6c		Wdr38	1810058I 24Rik	Dmkn	Emg1
119	kg:uc007bvx.1	Crlf1	Ubash3b		Col4a2	Nfrkb	Rnh1	Impact
120	Ap4e1	Avil	Inf2		Tnnt2	Pabpc1	Pdgfa	Mylip
121	Spock2	2700094K13R ik	Asap1		Ndufs8	Usp16	Rpl37a	Psmb8
122	Efemp1	Ctse	Sec11c		Tspan4	Pde1b	Rabac1	Rfk
123	Prpf40a	Penk	Gas2l1		Agpat6	Ncoa4	Timp2	Map3k5
124	Tspan5	Tmc4	Parvb		Timp3	Irf8	Serping1	Odc1
125	Lgals7	Dhrs3	Tmsb4x		Ankrd50	Ppp1cb	Rbm39	Slc11a2
126	Kif5b	Ap1s1	kg:uc007x rw.1		Ube2d3	Rgs2	Tgoln1	Eif2b1
127	Psip1	Arl6ip4	Nudt3		Sf1	Smyd4	Nfix	kg:uc00 8wjd.1
128	kg:uc008oki.1	9430008C03R ik	Bcl2l1		Csnk1d	Arid3b	Brd4	Rexo1
129	1810014B01Rik	Fcer1g	B230312A 22Rik		Reg3b	Sh3bgrl2	Tmem234	Ddx50
130	Ptges3	Uqcr11	Cnp		Flot2	Lyl1	Wbp5	Nipsnap3b
131	Limch1	Nhp2	Plp1		Lmna	Prr13	Ppig	Sp100
132	Bicd1	Plbd2	Cnst		231004 4H10Rik	Plagl2	Cd63	Uggt1
133	Rdx	Capg	Rgs18		H19	Nfkbia	Col1a1	kg:uc00 7czl.1
134	Pcdh15	Pnpla6	Lsm12		Slc20a1	Eef1a1	Mt2	Arpc5
135	Foxn3	Ppdpf	Alox5ap		672045 6B07Rik	Brd2	Zbtb7c	Nfrkb
136	Morf4l2	Hgfac	Ppif		Mdh2	Egr1	Npr1	Nap1l4
137	Ppp1r15a	Apoe	Spnb1		Eif6	Mkrn1	Tmem119	Fam117a
138	Cdc42ep3	Fam40a	Ormdl3		Phf5a	Pld4	Atf3	Sipa1l1
139	Pard3b	Lyz1	Hpse		Vps28	Aldh1a1	Ankh d1	Ttc1
140	Bicc1	2200002D01 Rik	Srxn1		Bag1	Dnajb9	Tmed10	kg:uc00 9vew.1
141	Amhr2	Laptm5	2010002N 04Rik		Cyc1	Gjb5	Slc6a4	
142	Gucy1a3	Qars	Hist1h2bc		Angptl4	Mtif2	Atxn7 l3b	
143	Psmb2	Tmx2	Cyba		Lgals3	H2- DMb2	Rpl29	
144	Mapkapk3	Fkbp4	Chst12		Farsb	Sdpr	Ccar1	
145	Ube2l6	Plin2	kg:uc009s		Mbd3	4932438	Ltbp4	

			ps.1			A13Rik		
146	kg:uc007pff.1	Fcgr3	Max		Timm13	Treml1	Scyl1	
147	kg:uc007ctp.1	Gkn1	Was		Tpd52l2	Nup153	Ap3d 1	
148	Nedd4	Snhg1	Isca1		Ptpn	Mpp1	Iqgap 1	
149	Plxna4	Lsp1	Pdzk1ip1		Crip2	Dhrs11	Cldn1 5	
150	2010107G12Rik	Gm20605	Lyn		Raver1	Lrmp	Spnb 2	
151	Ifngr1	Ly6c1	Mob3a		Eif2b2	Manf	Ano1	
152	Bcam	Aim1	H2-T24		Psma7	Mli3	Lrrn4	
153	Ccnl1	2310007B03Rik	Slc44a1		Rps6ka 4	Fam116b	Id3	
154	Hoxa5	Tgfb1	Derl1		Mgat4a	B4galt5	Eif3a	
155	Fhl1	Tsta3	Gclm		Ifitm2	kg:uc009 vew.1	Prkcd bp	
156	1810041L15Rik	Pafah1b3	Fech		Wars	Ly6d	Atp1a 1	
157	2900002K06Rik	Chid1	Ywhah		Capn5	Dguok	Dnaja 2	
158	Hspb1	Smox	Igtp		Bsg	Pnpo	Tubb 4b	
159	Podn	1500012F01Rik	Myl6		Sec16a	Tmem17 5	Hnrn pab	
160	Fam63b	Tspan4	Thbs1		Cldn7	Gm6548	Mmp 14	
161	Hsp90b1	Agrn	Tln1		Cox7a2	Rsrc2	Atp1b 1	
162	Dpp4	Cfp	kg:uc009a pq.1		Nek6	Ccdc88b	Psip1	
163	Gas1	Cdh1	Bcap31		Rpl39	Akna	Mgll	
164	kg:uc007zak.1	Rasgrf1	Ilk		Itpr3	Tsc22d3	Rnase 4	
165	Zc3h13	Nxf1	Epha1		Ctnna1	Txndc5	Ywha b	
166	Sox6	Pdrg1	2810453I0 6Rik		Tpd52	Tubb4a	Clip1	
167	Arid4a	Polr2j	Rnf19b		Mlf2	Stx11	Syn3	
168	Tnxb	Suds3	Gsn		Crip1	D4Wsu5 3e	Myl1 2a	
169	Tsix	D0H4S114	Flna		Fkbp4	Amfr	Rbm2 5	
170	Scd1	Ccl9	Arrb1		Gprc5a	Tti1	Arf2	
171	Jund	Neat1	kg:uc007p um.1		Slc4a11	Fam175b	Cav1	
172	Crls1	Ccdc12	Mbnl1		Syn3	Zfp36	Hnrn pc	
173	1110003E01Rik	Prr24	Ccnd3		Npc2	Ddx5	Syne2	
174	Rnase4	Impdh1	Pdlim1		Rpl32	Tlr7	Dst	
175	Arhgef12	Card10	Ctse		Inf2	Rfk		

176	Irf7	Cpsf1	Tspan17		Rps10	kg:uc007 ded.1		
177	Bbx	Sema4g	Gpx4		Rps26	Gnb2		
178	Sema5a	Hes6	Bnip3l		Rpl37a	Tmed5		
179	Mau2	C130074G19 Rik	P2ry12		Ctxn1	Thbs1		
180	Abi3bp	Ctrb1	kg:uc009v ev.1		Lrrc59	eg:32016 9:chr9:p		
181	Dag1	Rnaseh2a	Prkab1		Dctn1	Zfp335		
182	Cyp2s1	Golm1	F2rl2		Mtap4	Emg1		
183	Sfrs18	Ctsz	Stk4		Uqcr10	Trmt61a		
184	Hspb8	Cyb561	Fhl1		Suds3	Adipor1		
185	Cnot6l	Ndufs8	Rnf10		Ap1s1	Vwf		
186	Twsg1	Atp6ap1	Rasa3		S100a1	Aatf		
187	Gpc3	Srd5a1	Taldo1		Atp5j	Trib1		
188	Lrrn4cl	Carkd	Bysl		Aim1	Pcyt1a		
189	Cdh3	Cd24a	Esd		Plec	Stx18		
190	Cyr61	Eng	Aldh2		Prom1	Trp53bp 2		
191	Cyp2d22	Tcirc1	Rhog		Rhoc	Stk40		
192	Hist1h1c	Slc9a3r2	kg:uc009e cr.1		Mast3	Il18		
193	Aplp1	0910001L09R ik	Cald1		Olfml3	1810014 B01Rik		
194	Tbl1x	Cox5b	Wbp2		Uqcr11	Lcp2		
195	Pcm1	Adipor2	Ptpkj		Plp2	Gimap4		
196	Ifi204	Scarf2	Tpm4		Spna2	Rabl2		
197	Nfix	Myo7a	Mxi1		170001 7B05Rik	Ncf2		
198	Flrt2	Ppap2c	Ly6g6f		Anxa4	eg:49721 0:chr14: m		
199	Heg1	Pea15a	Sla		Nudc	Tpt1		
200	Il6ra	Sh3pxd2b	Slpi		Asl	Mli5		
201	Ralbp1	H19	Bicd2		Prkcsb	H3f3a		
202	Rhoj	Tpd52	Clu		Plod3	Tspan13		
203	Ktn1	2610203C20R ik	Mtmr14		Ndufa9	Il10ra		
204	Arl6ip5	Naa10	Abca7		Impdh2	Mdc1		
205	Crebbp	Fermt1	Ppp1r18		Ccnl2	Stk24		
206	Ppig	Sap30l	Kif2a		Nedd8	Myst4		
207	Akap13	Bgn	Prdx6		Atp6v1f	Zdhhc20		
208	Rab7	Timm13	kg:uc009iz e.1		Mt1	Eif2b1		
209	Plxdc2	Krt20	Calm3		Il4ra	Exoc4		
210	Aldh1a1	Itga3	Dhrs1		Cndp2	Wipf1		
211	Bnc2	Pfkl	Cfl1		Appt	Impa1		

212	Slc4a4	Agpat6	Glipr2		Preb	Tmem119		
213	Tbx18	Mrpl11	Slc25a37		Ap3d1	Pml		
214	Zbtb16	Ramp1	Atox1		Mcm6	Ubb		
215	Arid4b	Hmga1	BC057079		Ubr4	Zmat3		
216	Enpp2	Gpx2	Pla2g16		Pvrl2	Slc30a9		
217	Ptplad2	0610012G03 Rik	Rnf144b		Snrpg	Lat		
218	Akr1b3	9130017N09 Rik	Stk16		Cycts	Tgfb2		
219	Gm6644	Cygb	Rsad2		Efemp2	Ube2o		
220	Arf5	Tmprss4	Paip2		Cct4	Igfbp5		
221	Chi3l1	Paox	Capzb		Gm20605	Tspan5		
222	Gpr116	Endod1	Ppp1r12c		Smad3	Fmn1		
223	Cd82	Cndp2	4930412F 15Rik		Card10	Fnbp4		
224	Srrm1	Suv39h1	Ninj1		Krt7	Extl3		
225	Fmo2	Cog4	2510009E 07Rik		Cct2	Adcy7		
226	Tgfb1i1	Trim27	kg:uc007v sr.1		Coro1c	Enpp4		
227	Qrich1	Cyhr1	Pygb		Ltbr	Sep15		
228	Nfia	Trmt1	Tlk1		Ric8	H2-Ab1		
229	Pmp22	Zfyve19	Myct1		Ndufs6	Bnip3l		
230	Cdh11	Esrp1	Rnasek		Fibp	Slc11a2		
231	Arid5b	kg:uc008oow .1	Ctsd		Pold4	Stom		
232	Rbm3	Dync1h1	0610010K 14Rik		Rpl34	Mfhas1		
233	Prelp	Tab1	Bcas3		Rpl34- ps1	Mettl1		
234	kg:uc007qse.1	Pla2g6	Atpif1		Clic1	Rnf10		
235	Ddx3x	Timp1	Serf2		Eri3	kg:uc009 cf.1		
236	Sulf1	Eif3f	Becn1		Ets2	Klf4		
237	Spnb2	Abhd11	Tspan9		Unc13a	Psme4		
238	Tspan31	Pmm2	Acer2		Usmg5	Sema4a		
239	Prr13	Tyrobp	Vdac3		Sh3pxd 2b	Ftl2		
240	Ppp1cb	Farsb	kg:uc008k bg.1		Wdr6	Atad1		
241	Fbln1	Plod3	Oaz2		Las1l	Tspan31		
242	Gm6548	Abtb1	Serpine2		Polr2f	Srrm2		
243	Uap1	Brf1	Ccdc90a		Vamp5	Rab5c		
244	Mpdz	Tnk2	Ndufa1		Endod1	Capza1		
245	Sat1	Rfc2	Tssc1		Snrpd2	H2-Aa		
246	Stim1	Stxbp2	Mboat7		Tpi1	Fhl1		

247	Mll3	Pdlim7	Cd44		Wwp2	Cryab		
248	Slurp1	A430105I19Rik	Cxx1c		Dalrd3	Arid4b		
249	Cd81	Vill	Ecm1		Iqgap1	Gart		
250	Emp2	Bmp1	Mff		Ahsa1	1110004 F10Rik		
251	Trpm7	Mpzl1	Ptpn12		Trim27	Rnf11		
252	Crym	Thy1	Mgmt		Serpinf1	Zc3hav1		
253	Enpp4	Stab1	Cox4i1		D33004 1H03Rik	kg:uc008 bt1.1		
254	Raly	Aldh16a1	Tollip		Ppp2r5d	Rnf34		
255	Celf2	Eif4ebp3	Cds2		Minos1	Dmkn		
256	Ap3s1	Itpripl2	Ybx1		Tsta3	Btg1		
257	C1s	Mrpl52	Gypc		Prpsap1	Syt11		
258	Frmd4b	2310002L13Rik	Dgkd		Sphk1	Mtdh		
259	Nr4a1	Mcm6	Pecam1		Ldha	Med21		
260	Acin1	Kcnk1	Ftl2		Abca3	Rnf2		
261	Plod2	Pmf1	Nt5c3		B4galt3	Tcf12		
262	Id1	Cuta	1700037H04Rik		Porcn	Tacstd2		
263	Creg1	Nt5dc2	Cd151		Tmc4	Madd		
264	Zfp318	Rmnd5b	Lpin2		Serinc2	D16Ert4 72e		
265	Tmem140	Araf	6430548 M08Rik		Akr1b8	Pias1		
266	Mras	Wwp2	Pon2		Nudt4	Taok2		
267	Vwa5a	Lamb1	Ndufa3		Atp5l	Pold1		
268	Esyt3	Kcne3	6330578E 17Rik		Psmc3	Cep110		
269	Hexb	Uqcrq	Mfap3l		Hint1	A930013 F10Rik		
270	Nckap1	Gps1	Mink1		Rpl41	Tcof1		
271	Nipal3	Rexo4	Ston2		Xpnpep1	kg:uc009 bpd.1		
272	Ubxn4	Coro1c	Rac2		Nav1	kg:uc009 bpr.2		
273	Zfp36	Hras1	Fyn		Parva	Capza2		
274	Hnrnpl	Sprint1	Serinc3		Immt	Ptp4a2		
275	C1ra	Cblc	Maged2		Pafah1b3	Fth1		
276	Nnmt	Fhod1	Ap2m1		Chid1	Mepce		
277	Mut	Atp13a1	Pacsin2		Aldh1l1	Rexo1		
278	kg:uc008jup.1	Man2c1	Ftl1		Rpl31	Prg4		
279	Pnrc1	Vsig2	Adipor1		Wbp1	Ctla2a		
280	Usp8	Bpgm	kg:uc009q		Zfp622	Smarc5		

			do.1				
281	Pgcp	Bap1	Snap23		270006 OE02Rik	Icam2	
282	Junb	Smpd2	Tagln2		Hspa9	Pbx1	
283	C1rl	Ubqln4	Cox6c		Tceb2	Gnl3l	
284	Slc6a6	Sirt7	Creg1		Rpl36a	Slc2a3	
285	kg:uc008znh.1	Krt23	Bsg		Pgs1	Nnmt	
286	Aqp1	D8Ertd738e	Cmtm6		Mpnd	Rb1cc1	
287	Myh10	Mapk13	Cntd1		Cdc42	Bpgm	
288	Slc43a3	kg:uc008bcq. 1	Plekho2		Dhrs3	Lcp1	
289	Spint2	Polr2g	Arrb2		Hexa	Sipa1l1	
290	Hnrnph1	Ndufs2	Pard3b		Cpsf1	Lilrb4	
291	Arhgap28	Dad1	Mlec		Mea1	Ankrd44	
292	Cfh	Wnt7b	Taf10		Polr2e	Specc1	
293	Brd4	Fam20c	Gabarapl2		Ddb1	Rif1	
294	Fndc1	Cxxc5	Bag1		Ptcd1		
295	Star	Polr2f	Galnt2		Atp5f1		
296	Nfkbiz	Ltf	Hk1		Sec61b		
297	Arsb	2210407C18R ik	Fbxo9		Psmc5		
298	Rnd3	Cdip1	kg:uc009iz d.1		Fam89b		
299	Stard5	Glrx5	Pnpo		Lama3		
300	Thbs1	Gemin7	Fam46c		Tomm6		
301	kg:uc008wkn.1	Man1b1	Pkm		Mrpl28		
302	Slc26a3	Heatr7a	Ap1b1		Syngr2		
303	Phip	Arid5a	Rap1b		Ngfrap1		
304	Usp2	Sumo3	Itgb1		Kcmf1		
305	Golgb1	Srm	St7		Tubb4b		
306	Rock1	Plscr3	Smap1		Anapc1 1		
307	Rgma	2210010C17R ik	Rabgap1l		Vcp		
308	Actg1	Fam102a	Tmbim4		Arpp19		
309	BC013529	Dlst	H3f3a		Pglyrp1		
310	kg:uc007zwh.1	Vps37c	Frmd8		Rrp1		
311	3110062M04Ri k	Ngfrap1	Nlrx1		Gkn3		
312	Cast	Pold4	Oaz1		Atpif1		
313	Mob3c	Grcc10	Fam125b		Prickle3		
314	Slc16a1	Wnt7a	Hexa		Map4k4		
315	Fam117a	2010111l01Ri k	Tspo		Arrdc1		
316	Pdia3	Pxdn	Dcaf12		C1qtnf6		
317	Trim8	Coasy	Nav1		Hras1		

318	kg:uc009mng.1	Dctn1	Cd24a		Lamb1			
319	eg:245190:chr7:m	Ncor2	Uqcr11		Eif3d			
320	Sbsn	Postn	Wipf1		Snrpa			
321	Serpinb6b	Col4a2	F10		Tbrg1			
322	Daglb	Cib1	Erlec1		Nxf1			
323	Smarca2	Tbc1d13	Map2k3		Pdlim7			
324	Mef2c	Ccnl2	Stk24		Add1			
325	Prcc2c	Dcakd	Ldlrap1		Pfdn5			
326	BC005537	Cdc34	Ehd4		Stk16			
327	Hsp90ab1	Atp6v0b	Atp6v1f		Gm17821			
328	Snrnp70	Abhd12	Gnas		Csnk1e			
329	Ppl	Flot2	Arhgap18		Rrp7a			
330	Serpinh1	Sla2	Arhgap10		Psmb6			
331	Sorbs3	Rhbdf1	Pitpnm1		Snhg1			
332	Golga4	Cdh17	S100a1		Ssr4			
333	Acbd3	Psmb5	Bin1		Ergic3			
334	Hook3	Serf1	Ttyh3		Rnaseh2a			
335	Map3k3	Slc15a3	Selp		kg:uc009cut.1			
336	Rhou	Sftpd	Trappc9		Bgn			
337	Smc2	Pop5	Aes		Gm5506			
338	C1d	Nudc	Taok3		Uqcrq			
339	kg:uc008dzh.1	Sh2d5	Zfand3		Tmem167			
340	Psmd7	kg:uc007fwp.1	Stim1		Nasp			
341	Dab2	Mrpl37	Rnf114		Mif			
342	Cep164	Rin1	Sep15		Acaa2			
343	Crim1	Podxl	kg:uc012hdk.1		Fam162a			
344	Rtf1	Paqr5	Lgals9		Eif4ebp3			
345	Fxyd1	Sepx1	Cox6b1		Nhp2l1			
346	H2-D1	Agr2	Riok3		Prelid1			
347	Zfp704	Bax	Slc38a10		Gss			
348	Mtap1a	Rxrb	Rtn3		Lonp1			
349	Ascc3	Tes	B3gat2		Srsf2			
350	Med13l	Hdac6	Ccndbp1		Igsf8			
351	Jup	1110008F13Rik	Rsu1		Ndufa7			
352	Nid2	Mpnd	kg:uc007upr.1		Neat1			
353	Kdr	Gmppa	Itm2b		S100a1			

					3		
354	Ifnar2	Gramd1a	St3gal1		Apoa1bp		
355	5430435G22Rik	Wars	Sec61g		Fam40a		
356	Col4a6	Mtap	Ptpn1		Rps25		
357	Il17re	C1qtnf5	kg:uc012bhf.1		Eno1		
358	Gbp3	Mrpl28	B2m		Cldn2		
359	Slc39a8	Mfrp	Rasgrp3		Capn2		
360	Cfl2	Kars	Memo1		Glo1		
361	Slc38a1	Lbp	Slc39a4		Atp5c1		
362	Cuedc1	Plxnb1	Sdcbp		Rab2a		
363	Fgf1	2700081O15Rik	Tspan14		Rab25		
364	Gas6	Mrps24	Ubl7		Ncor2		
365	Cldn25	Klc4	Nras		Lgi4		
366	Sorbs1	Dctn3	Ssx2ip		Ier3		
367	Hspa12a	Kcnq1	kg:uc007zbz.1		Tmem223		
368	kg:uc007zts.1	Smurf1	Wbp1		Slc9a3r2		
369	Slc1a5	Fam162a	1110003E01Rik		Atp13a1		
370	Nr3c1	Hip1r	Clip2		Rpn2		
371	Adamts5	kg:uc007hyr.2	Gapdh		Acp5		
372	Gpcpd1	Gys1	Gm6578		Cct5		
373	Dpysl3	Sac3d1	Actn1		Sdf4		
374	Colec12	Ndufs6	St3gal2		Mppip		
375	Pcd6ip	Rgl2	3110001D03Rik		Pmm2		
376	Dst	Atp5g1	Ctsz		Snx22		
377	Ifit3	Itgb4	kg:uc007vdl.1		Arl2		
378	Chst4	Sars	Fam73a		1110008F13Rik		
379	Xist	2310003F16Rik	Vcl		Polr1d		
380	Ifi27l2a	Nhp2l1	Lims1		Dpm2		
381	Fkbp5	D19Wsu162e	Lars2		Cela1		
382	Agap1	Cd320	Birc2		2310016M24Rik		
383	Ankrd11	Pigg	Lamp2		Cep250		
384	kg:uc007qca.1	Chd3	Rasl10a		Mybbp1a		
385	Syt11	Zdhhc4	Mif		Polr2g		
386	Ptrf	Eif3l	Rab10		Bag6		

387	Krcc1	St8sia3	Pabpc1		Cpxm1			
388	Zfp488	Rcan3	Wwp2		Eif3m			
389	Lama4	Meg3	Nqo2		Prr24			
390	Aebp1	Nudt4	kg:uc007fe.1		Sra1			
391	Fam134b	Gss	Plxna4		Scara3			
392	Tppp3	Pih1d1	Gm1821		Reg1			
393	Maf	Limd2	Gadd45a		Gas5			
394	Peli1	Ap1s2	Slc25a39		Hnrnpab			
395	Zfp353	BC056474	kg:uc009pet.1		Mcpt2			
396	Cdon	Mms19	Ubb		Tgfb1			
397	Sarnp	Clip2	Ppp1r2		Capns1			
398	Atxn7l3b	2310016M24Rik	Rab27b		Fdx1l			
399	Pef1	Itpa	Cap1		S100a16			
400	App	Slc25a10	Jarid2		Nap1l1			
401	Mtdh	Fibp	Rnf11		Swi5			
402	Lrrc20	Higd2a	Tmem50b		Rpl38			
403	Btbd2	Snrpd2	Myh9		Dctn2			
404	Gnb2	Eri3	Tmem128		Pdlim1			
405	Pigt	Nbeal2	Stradb		Gemin7			
406	Efna5	Trim28	Cela1		Pnpla6			
407	Tm4sf1	S100a4	Ndrg2		Nono			
408	Coq10b	Ivns1abp	Dhrs3		Sla2			
409	Eif2s3x	Ppp1r18	Hipk1		Idh3b			
410	Cmah	Efemp2	Atg9a		Ppp2r4			
411	Sf3b1	Med22			Map2k2			
412	Eea1	Nelf			Ndufb10			
413	Slpi	2810428I15Rik			Atp5d			
414	Tmod3	D2Wsu81e			Arfgap1			
415	Ppp3ca	Trappc6a			Tmbim1			
416	Tceal8	Trappc2l			Ergic1			
417	Anp32a	Antxr2			Pdgfa			
418	Actb	Rab11fip5			Ppp2ca			
419	Ddx5	Ldhd			Hk1			
420	Cobll1	Npnt			Ltbp2			
421	Cish	Acrbp			Trim35			
422	Nod1	Pafah1b2			Gtf2i			
423	Psd	Angptl2			C1qb			
424	Gm10052	Fzr1			Ankhd1			

425	Lims2	Aaas			Podxl			
426	Stra6	Eif2b2			Rps21			
427	kg:uc007bgn.1	1190003J15Rik			Huwe1			
428	Plxdc1	5730403B10Rik			Pomp			
429	Nfe2l1	Adamts13			Dpp3			
430	Smpd3	Eif3b			Fkbp8			
431	Bcl10	Znrf1			Itga5			
432	Ilf3	Pkp3			Hes6			
433	Fam76a	Lemd2			Mrpl11			
434	Cybrd1	Rab34			Poldip3			
435	Gm3893	Mpv17l2			Scd2			
436	Siae	Cdkn2b			Tmem55b			
437	Ssh2	Snrpe			Ndufa13			
438	Nfic	Gm14005			Dcakd			
439	Btf3	Prdx4			Ubqln1			
440	Sp100	Xab2			Gpx4			
441	Ndn	Dpp3			Cyb561			
442	Matr3	Tyms			Gmppa			
443	Gm13251	Leprotl1			Ncaph2			
444	Arhgap5	Uqcr10			Pdha1			
445	Zbtb4	Cdk5rap3			Ndufs4			
446	Pgrmc1	Gorasp2			Fcer1g			
447	4930402H24Rik	Wbp7			Myof			
448	Bptf	Sort1			Ppib			
449	Dusp3	Ddx41			Mrpl52			
450	Pla2g4a	Cct3			Tes			
451	Brp44l	Mrps33			Emp3			
452	Oxct1	Frmd8			Ndufa11			
453	Stk40	1110049F12Rik			Tor1aip2			
454	Ddr1	Fscn1			Anp32b			
455	Ifi205	Ndufa2			Tnk2			
456	Col3a1	Dpcd			Mcpt1			
457	Nipbl	Unc13a			Ssr2			
458	Plk1s1	Eif1ad			Psmb3			
459	Bdp1	Sgta			2700081015Rik			
460	Smc3	Chaf1a			Pcbd1			
461	Ifitm3	Plxna1			Eif1ax			

462	Ndst1	Hspa9		Pmm1			
463	Zbed6	1110014N23 Rik		Ptprk			
464	Rest	Cd99l2		Hadha			
465	kg:uc007vnc.1	Snrpa		Calu			
466	Ccdc88a	Mcm7		Fam73a			
467	Stat3	Tars2		Atp5e			
468	Arf2	Gon4l		Hsd17b 10			
469	Trib1	Stk38		Rbm39			
470	Gcap14	C1qtnf1		Egfl7			
471	Tbc1d15	Tbrg4		Psmc1			
472	Igf1r	Tmem132a		Perp			
473	Ppbp	Cox6c		Lman2			
474	kg:uc008tky.1	Alcam		Galnt1			
475	Rab1b	Phka2		Rbx1			
476	Krt14	Trim3		Lemd2			
477	Med21	Ppp1r14b		Zglp1			
478	Gja1	Gpaa1		Ing4			
479	Klf10	Ctps2		kg:uc00 8ooow.1			
480	Id2	Ptpn23		150001 2F01Rik			
481	Mfap1a	Endog		Cox4i1			
482	Ogn	Mrto4		kg:uc00 8bcq.1			
483	Gpc4	Mrps6		Ubap2l			
484	Bst2	Pvr		Pafah1b 2			
485	Dtx2	Phgdh		Mrpl13			
486	Wac	Itpr3		Nucb1			
487	Kpna3	Polr2e		Fbn1			
488	Kcnab1	Sec16a		Adrm1			
489	Orai3	Mdp1		Itgb4			
490	Gcsh	Fbf1		Ctss			
491	Wdr92	Mcpt8		Plbd2			
492	Olfr613	Rps6ka4		Ptpmt1			
493	Tcf7l1	Mical1		Sap30l			
494	Tgfb2	Mrpl34		Ppp1r1 2c			
495	Il16	Agpat3		Sgta			
496	Manf	2310044H10 Rik		Acrbp			
497	Mgst1	Myo9b		Higd2a			
498	kg:uc008tkz.1	Ndufb10		Higd1a			
499	Creb3l1	Apex1		Tmem2			

					08		
500	Txndc5	Elk3			Cdh1		
501	Klf2	Cpsf3l			Ube2d2a		
502	Slu7	Tnk1			Suv39h1		
503	Ttc28	Pmvk			Rabac1		
504	1110002B05Rik	Ppp1r16a			Anxa5		
505	Zcchc11	Arhgef5			Ubxn6		
506	Ptp4a2	Lonp1			Tpm1		
507	Pbx1	Pla2g7			Hmga2		
508	Clcn3	Pip5k1c			Cnbp		
509	Tmco7	Inf2			Rpl21		
510	Lrrc58	Pgk1			Ndufb5		
511	Eif3a	Parp6			Sec31a		
512	Cldn10	Urm1			Znhit1		
513	H2-Q6	Mad2l2			Cyb5b		
514	Ccdc80	Ing4			Sfn		
515	kg:uc009iln.1	Rbck1			Ccdc12		
516	Rab5c	Cant1			Elov1		
517	Tsc22d3	Sgpl1			Psmb5		
518	Tm4sf5	Ehbp1l1			Slc25a11		
519	Hmgb1	Runx1			Psmd2		
520	Sec62	Slc27a4			Nsun2		
521	Maoa	Ndufa7			Slc50a1		
522	Clec1b	Mcm3ap			Eme1		
523	Mphosph8	1110008P14Rik			Bnip2		
524	Oat	Rassf7			Pxdn		
525	Ncor1	Ptpmt1			Mad2l2		
526	Cyb5	Arfgap1			Pcd6		
527	Traf1	Sec61a1			2010107E04Rik		
528	Rpp25	Rps6ka1			Abhd11		
529	kg:uc007ded.1	Ints1			Carkd		
530	2610101N10Rik	Tpcn1			Polr2l		
531	Il6st	Iflo2			Ppdpf		
532	Evpl	Trim44			Cib1		
533	Psmd11	kg:uc012ctw.1			Dgcr2		
534	Dync1i2	Golga2			Timm50		
535	Lars2	Msto1			Mrps24		
536	Pdia4	Ppp6r3			Abhd12		

537	Cd55	Trmt2a			Brf1			
538	Amfr	Appl2			Man1b 1			
539	Zcchc3	Sparcl1			kg:uc01 2cgd.1			
540	Herpud2	Rapgef1			Gpaa1			
541	Txnrnd1	Zfp11			Fmn13			
542	Vat1	Psmc4			Mapk3			
543	Diap1	Mosc2			C1qc			
544	Tmed2	Fam101b			Pgls			
545	Arf3	1500010J02Rik			Cp			
546	Arap2	Ccdc124			Serhl			
547	St3gal1	Ptges			261020 3C20Rik			
548	Man1a	Fam189b			Hsbp1			
549	Rgs10	Th1l			Tmem2 14			
550	Tmsb4x	Kctd2			Akt1			
551	Uba7	Olfr1372-ps1			kg:uc00 7pfe.1			
552	C4b	Hexa			Tmed10			
553	Tmem98	Anapc5			Ttl3			
554	Lpar2	Serpina3n			220000 2D01Rik			
555	Gabarapl1	1810046J19Rik			Tnfrsf2 3			
556	Cmtm7	Tmem167			Sgsm3			
557	Spon2	Gm11428			Atp9a			
558	Smarca5	Gcn1l1			Lcn2			
559	Mxd4	Kansl3			Pdrg1			
560	Smc4	Fasn			Tspan9			
561	Thsd4	Slc50a1			Nrd1			
562	Gsr	Smad3			Rin1			
563	Ptprd	Trip6			Ndufv1			
564	Clip1	Atp6v1e1			Naa10			
565	Cln8	Chchd5			Wnk1			
566	Rbm27	Adssl1			Heatr7a			
567	Zmat1	Nes			Slc4a2			
568	Smc6	Ap1b1			Ggct			
569	B2m	Fcgtr			573040 3B10Rik			
570	Irf2bp2	Ltbp3			Sh3glb2			
571	Ppap2a	Csf2rb			Pfkl			
572	Zfhx4	Ssna1			Tspan3			

573	Tob2	Mrps16			Gns			
574	Rabgap1l	Cyba			Sdcbp2			
575	Nfkb2	Cyth2			C13007 4G19Rik			
576	Nfyc	Igf2			Cotl1			
577	Ube2d1	Pisd-ps1			Tubb5			
578	Creb5	Atp13a2			Sec11c			
579	Opa3	Mlph			Pigg			
580	Csnk1a1	Cyp4f16			Zc3h15			
581	Fam84b	2010107E04Rik			Lsmd1			
582	Ddr2	Gas5			Ppa1			
583	Usp54	Eif3k			Chmp4b			
584	Akt2	Fam149a			Sepn1			
585	Strn3	Mif			Angptl2			
586	Hnrnpm	B230312A22Rik			Itpripl2			
587	eg:497210:chr14:m	Ppp1r12c			Ddx1			
588	Tpt1	Tfip11			Hbxip			
589	Naa25	Tex10			Cdk2ap1			
590	Eef1a1	Slc16a3			Clta			
591	Parp4	Stk16			Cpsf3l			
592	Msn	Epn1			Apoe			
593	Zbtb20	Noc4l			Ift46			
594	Fermt2	Rcc2			Sae1			
595	Bod1l	Rgs12			Gpi1			
596	Sltm	Shkbp1			Gorasp2			
597	Dapk1	Got2			150003 2L24Rik			
598	Hnrnpr	Plek2			Nsmce4a			
599	Baz2a	Lilrb3			Dlst			
600	Rnf167	Ndufb5			Bap1			
601	Mapk1	Tesk1			Pitpnb			
602	eg:320169:chr9:p	Rab24			Meg3			
603	4930523C07Rik	Atp5j2			Cyth2			
604	Nf1	Commd9			Atp5o			
605	Fam53b	Rtkn			Gon4l			
606	Faim2	Prpf19			Sox11			
607	Tgm2	6720401G13Rik			Cxxc5			

608	Calm2	Ppa1			Avil			
609	Al848100	Pgp			Alcam			
610	Slc10a3	Hps1			Eif3f			
611	Ogdh	Puf60			Cygb			
612	Arl3	Mdm2			Eif1ad			
613	Timp2	kg:uc012cgd.1			Polr3h			
614	Atxn2	kg:uc009uim.1			Araf			
615	Mll1	Pyy			Gkn1			
616	Ces2g	Zfp358			Rhog			
617	Mat2a	Timm8b			Mtap			
618	Esf1	Ddx39			Eif4ebp1			
619	Hsp90aa1	Pgm2			Akr1a1			
620	Zfp385a	kg:uc008gbp.1			Trip6			
621	Zfp672	Sipa1			Prdx6			
622	Csda	Mgat1			241001 5M20Rik			
623	Pf4	Tmem208			Rps6			
624	Arsa	Ruvbl2			Rps23			
625	F11r	8430410A17 Rik			Stxbp2			
626	C4a	Bad			Rps19			
627	Kpna1	Pfdn5			Ykt6			
628	Rbbp8	Eme1			Atp5g2			
629	Oxnad1	kg:uc009mzj.1			Serpinb1a			
630	Rb1cc1	Igf1			Col7a1			
631	Setd2	Prkag1			Mrps6			
632	Kif1b	kg:uc009sua.1			Lgals9			
633	2510002D24Rik	Uap1l1			Rcn3			
634	Cep57	Trappc4			Trim44			
635	Chd2	Bola2			Surf2			
636	Serinc5	Usp5			Rps29			
637	Marcks1	Ear2			Cdpt			
638	Shfm1	Cars			Lmf2			
639	Bbs4	1810027O10 Rik			Psenen			
640	Impad1	Amdhd2			Ltf			
641	Tbcel	Phb			Mpzl1			
642	Kdelr1	Kcmf1			Psmd6			
643	Ninl	Lsmd1			Cttn			

644	Syt1	Sec11c			Tmc6			
645	Tpm3	Pcbp4			250000 3M10Rik			
646	Rbbp6	Mepce			Atp6v0 a1			
647	Lman1	Tpd52l2			Med8			
648	Ankrd17	Trf			Prrx2			
649	Naga	Hsd17b11			Atp5b			
650	Rbpms	Pilra			Smurf1			
651	Magt1	Atn1			Carhsp1			
652	Tfdp2	Pgf			Tpcn1			
653	Gem	Nxn			Ndufb9			
654	Pde4dip	Inpp5k			Pih1d1			
655	Mrgprf	Actr1a			Hhrnpa 0			
656	kg:uc008ajk.1	Cd68			Fn1			
657	Itch	Eef1g			281042 8I15Rik			
658	Elf1	Fbn1			061001 2G03Rik			
659	Meis2	Hint1			Ube2i			
660	Arid1a	March5			Anxa3			
661	Serpinc1	Usp48			Msto1			
662	Slc27a3	Hnf1b			Eng			
663	Thoc2	Gga3			091000 1L09Rik			
664	Gsta3	Drosha			Rpl10			
665	Hnrnph2	Ubp1			kg:uc00 7xxx.1			
666	Socs3	Pkn3			Mosc2			
667	Armcx3	Tmem192			Vps37c			
668	Siah1a	Prpf31			Sgpl1			
669	kg:uc009ize.1	Hspd1			Fam166 a			
670	Irs2	Otub1			Polr2b			
671	Mettl7a1	Mrpl20			Fam101 b			
672	Ppfibp2	Tead2			Nupr1			
673	Blvrb	Phpt1			Lsm4			
674	Yipf5	Neu1			Rpl36			
675	Plat	Pygo2			061000 7C21Rik			
676	Gm6578	Myeov2			Psmc2			
677	Mat2b	Cdk5			Supt6h			
678	Tmpo	Ndor1			Rps13			

679	Metap2	Rbp4		543043 7P03Rik			
680	Zfp277	Psat1		Dsp			
681	Wls	Mrpl41		Ddx56			
682	Mesdc1	Snrpg		Tsc2			
683	kg:uc009acs.1	Acot7		Trmt2a			
684	Col1a2	Vars		Vdac2			
685	Csf1	Nono		Cant1			
686	Sulf2	Gtf2i		Eif4h			
687	Ifrd1	Traf3		Puf60			
688	Wrnip1	Ppp2r4		A43010 5I19Rik			
689	Flii	Actg2		Cacnb3			
690	2810474O19Rik	Pi4k2a		Prdx4			
691	Sep15	Slc35b2		March5			
692	2310030G06Rik	Ubqln1		Ccar1			
693	Cmtm3	Ppox		Npepl1			
694	Mylip	Bud31		Fermt1			
695	Slc8a1	Man2b1		Use1			
696	Btbd7	Nat15		Axl			
697	Hdac5	Spon1		Slc39a4			
698	Zfand6	Cyc1		111000 8P14Rik			
699	Tapbp	Mpeg1		Sema4g			
700	Keap1	Nsun2		Timm8b			
701	Ube2n	Rab4a		Krt23			
702	Ssr3	Mtmr11		Rpl28			
703	H3f3a	BC004004		Lgals3bp			
704	Myst4	B4galnt1		Hdgf			
705	G3bp1	Atp5k		111000 5A03Rik			
706	Ugdh	Lin37		Impdh1			
707	Lamp2	D330041H03Rik		Mtmr11			
708	Zrsr1	Tbc1d17		Msln			
709	Pim1	March6		Zdhhc3			
710	Gm9199	2410015M20Rik		Znrf1			
711	Supt16h	1810013D10Rik		Aldh16a1			
712	Ano6	Eif2s1		Bloc1s1			
713	Soat1	Traf7		Prkag1			
714	Eci1	Rpl36al		Plxnb1			

715	Plce1	Psenen		Crat			
716	Atg3	Aip		Phpt1			
717	Bnc1	Cmas		593043 4B04Rik			
718	Pik3c2a	Rpia		Kpnb1			
719	Pqlc3	Ncbp1		Nme2			
720	Thrap3	Mea1		E43002 5E21Rik			
721	Irak4	Timm50		Smyd2			
722	Kdm6b	Ear12		Cyhr1			
723	Apol9a	Fkbp1a		Mvp			
724	Wnt4	Commd4		Rps27l			
725	1500003O03Rik	Col5a3		Rbp4			
726	Phf3	Fblim1		Cars			
727	1110004F10Rik	Cwh43		kg:uc01 2ctw.1			
728	Kansl1	Arl2bp		Ssr1			
729	Fth1	Mrpl46		Ssu72			
730	Tmem50a	Tcn2		Usp48			
731	Utp20	Add2		Atp5k			
732	Smad4	Specc1l		Lrrk1			
733	Stmn2	Ppcs		BC0564 74			
734	Gstm1	Vrk3		Epn1			
735	Senp6	Trim25		Trappc1			
736	Gda	Nfatc1		Clk2			
737	Nucks1	Rap1gap		Sugt1			
738	Ints10	Hsd17b12		Nenf			
739	Syne1	Epas1		kg:uc00 9cuu.1			
740	Itga6	Ddx1		Ubap2			
741	Acad9	Prdx6		Rps20			
742	Maged1	Mmp24		Atp5h			
743	Spen	Ndufb9		943000 8C03Rik			
744	Chd1	Phf23		Kars			
745	Taf3	Rpa2		Mrpl37			
746	Ptgs1	5031439G07 Rik		Aimp1			
747	Sparc	Rrp7a		Trmt1			
748	R74862	Arfip2		Hspa4			
749	B230120H23Rik	Efna1		Cd164			
750	Tmem234	Agps		943002 3L20Rik			
751	Ryk	Sephs1		Rnf4			

752	Dlgap4	Apoc2		H1f0			
753	Atp1b1	Mrps27		C1qtnf1			
754	Parp14	Snn		Srd5a1			
755	Tgfbr2	Serinc3		150001 OJ02Rik			
756	Ccdc90a	Pdcd5		Rpl35a			
757	Ncoa1	AA986860		Cand2			
758	Pppde1	Pitpna		C63000 4H02Rik			
759	Luc7l3	Vac14		Acsbg1			
760	Prg4	2810025M15 Rik		Derl1			
761	Rab11fip1	Def8		Cbx5			
762	Plk2	Hilpda		Tmem6 3a			
763	Ifi35	Eif6		Hgfac			
764	Pdap1	Brd7		Stx5a			
765	Cd248	Fes		Bri3			
766	Sesn1	Sbf1		Tomm2 0			
767	Ecd	Ak2		Fam20c			
768	Ap1s3	1810035L17R ik		Cox6c			
769	H2-K1	Lime1		Tm2d2			
770	Spag9	Hspe1		Plekhb2			
771	Tshz1	Csrp2bp		Ramp1			
772	Dennd5a	Uba5		241000 1C21Rik			
773	Stag1	Gsta4		Tardbp			
774	Gpx8	2900092E17R ik		Pebp1			
775	Sod3			kg:uc00 8gbp.1			
776	BC005561			Eif3b			
777	kg:uc009vev.1			Ccna2			
778	Ywhaz			Ptges			
779	Ganab			kg:uc00 7hyr.2			
780	Rras2			Wbp5			
781	Dusp14			Chchd2			
782	kg:uc012hdk.1			Fdft1			
783	Nr1d1			Srm			
784	Wwc2			Gtf3a			
785	Ubxn2a			D0H4S1 14			
786	Iqsec1			181000 9A15Rik			

787	kg:uc007vsr.1				Rps27			
788	Cfl1				Tmem1 76b			
789	Csrp1				Ndufc1			
790	Smchd1				Lasp1			
791	Myl12a				Fam108 a			
792	Ubqln2				Mapk8i p3			
793	Tmcc3				Copa			
794	Kdm5a				Serpina 3n			
795	Rbm25				Rps17			
796	Wdr26				Dnpep			
797	Vim				Lbp			
798	Arpc2				Krt19			
799	Calm1				Ei24			
800	Dnaja2				Ap1b1			
801	Shc1				Mogs			
802	Vps13a				Uba1			
803	Klf7				Postn			
804	1810074P20Rik				Phf23			
805	BC003331				Paox			
806	Itpr2				Nploc4			
807	Jmjd1c				Ndufv2			
808	Pcdhgb5				Actr1a			
809	Tubb2a				Mxd3			
810	Ehd2				Pfdn1			
811	Ift74				Ide			
812	Per1				Foxp4			
813	Pitpnm2				181001 3D10Rik			
814	Gstm4				231000 7B03Rik			
815	Dnmt1				Xab2			
816	Tmco1				Agr2			
817	Lass4				Dctn3			
818	Ptprf				Urm1			
819	Sirt2				H2-Ke2			
820	Gfm2				Spint1			
821	Taf7				Slc38a2			
822	Spop				Ube2z			
823	Zzef1				Ctrb1			

824	Ccdc34				Fam195b			
825	Zfp281				Suclg1			
826	Tuba1a				Ube2l3			
827	Ccdc109b				Rpn1			
828	Cdk13				Mrps7			
829	Dhx15				Tsg101			
830	Src				Drosha			
831	Braf				Arfip2			
832	Mapre2				Mrto4			
833	Anxa7				Grif1			
834	Sept9				Sort1			
835	Alox12				Oaf			
836	Pknox1				Ints1			
837	2610034B18Rik				Slc44a2			
838	Topors				Dph3			
839	Phf21a				Gramd1a			
840	Qser1				Fkbp9			
841	Tirap				Fam149a			
842	Fas				1810035L17Rik			
843	Lass2				kg:uc007fte.1			
844	6330406I15Rik				Eif2s1			
845	Parvb				Smpd1			
846	Atp1a1				Eef1b2			
847	Mtmmr6				Actr10			
848	Cd109				Rab11fip5			
849	Dnajc1				Ypel3			
850	Hp1bp3				Flnb			
851	1600029D21Rik				Tcn2			
852	Ttc38				Crlf1			
853	Mfhas1				Map3k15			
854	Filip1l				Cul7			
855	Zfp148				Atp6v1g1			
856	Nkd1				Ncbp1			
857	Usp16				Atp1b3			
858	Tlr2				Mtif3			
859	Zc3h18				Aldoa			
860	Stk10				Htra1			

861	Ltbp4			Rab14			
862	Hdac3			Ppm1a			
863	Efhd2			Ndufb1 1			
864	Prkar2a			Kansl3			
865	Atp6v1a			Rab24			
866	Sf3b4			Bcl2l1			
867	Gprc5b			Lgals1			
868	Clip3			Samm5 0			
869	Mettl2			Mrps33			
870	Secisbp2			Anxa1			
871	Fmod			Chchd1			
872	kg:uc009lxf.1			Mapre1			
873	Elovl6			Ctbp2			
874	Bzw1			Rnps1			
875	Etfa			Spg7			
876	Hspa2			Tnfrsf1 2a			
877	kg:uc007won.1			H6pd			
878	Rnf20			Myo7a			
879				Mcm7			
880				Psmd13			
881				Mrpl54			
882				Atp6v0 b			
883				Prdx1			
884				Elof1			
885				Rexo4			
886				Mrps18 a			
887				Dpcd			
888				D2Wsu 81e			
889				Cd99l2			
890				Synpo			
891				Atp2a2			
892				Cdc5l			
893				Stard7			
894				Atp13a 2			
895				Sdha			
896				Hdac6			
897				Krt20			
898				Ppp6r3			

899				170003 7H04Rik			
900				Napa			
901				Pgp			
902				Cnih			
903				Atg4b			
904				Cox8a			
905				Srp68			
906				St13			
907				Gng12			
908				Cfdp1			
909				Rcc2			
910				Pisd-ps1			
911				Ivns1abp			
912				Mpv17l2			
913				Ssna1			
914				Gnl1			
915				Tmem111			
916				Hbs1l			
917				Agpat3			
918				Col6a2			
919				March6			
920				Usp39			
921				Rps11			
922				Ahnak			
923				Lcmt1			
924				Ddx41			
925				H2afv			
926				Fau			
927				Tuba1c			

[00257] The gene names listed in Table 13 and 12 are common names. NCBI Gene ID numbers for each of the genes listed in Table 13 and 12 can be obtained by searching the “Gene” Database of the NCBI (available on the World Wide Web at <http://www.ncbi.nlm.nih.gov/>) using the common name as the query and selecting the first returned *Homo sapiens* gene. Other genes may be obtained using the UCSC genome browser (available on the World Wide Web at <http://genome.ucsc.edu>) using the Gene Sorter function. In certain embodiments, the marker gene(s) are selected from the genes listed in Table 13 and/or 12.

**[00258]** In some embodiments, the marker gene(s) is selected from a marker gene indicated to be upregulated in at least one type of CTC in Table 13, e.g. marker genes 1-142. In some embodiments, the marker gene(s) is selected from a marker gene indicated to be upregulated in at least one type of CTC in Table 12, e.g. marker genes listed in the columns labeled “CTC-c vs. Primary Tumor Enriched Gene” or “CTC-c vs. WBC”.

**[00259]** In a CTC, the marker genes listed in Table 13 or 12 can be upregulated, e.g. for marker genes listed in Table 13 and/or 12, if the measured marker gene expression in a cell or sample is higher as compared to a reference level of that marker gene’s expression, then the cell is identified as a CTC and/or the sample is identified as comprising CTCs. Preferably, once looks at a statistically significant change. However, even if a few genes in a group do not differ from normal, a sample can be identified as comprising CTCs if the overall change of the group shows a significant change, preferably a statistically significant change. All possible combinations of 2 or more of the indicated markers are contemplated herein.

What is claimed herein is:

1. A method of detecting classical circulating tumor cells (CTC-cs) in a sample, the method comprising:
  - isolating CTCs from the sample;
  - measuring the level of an IGFBP5 expression product and a DCN expression product in the CTCs; and
  - determining that CTC-cs are present if the detected level of the expression products is greater than a reference level.
2. The method of claim 1, wherein the CTCs are pancreatic cancer CTCs.
3. A method of treating cancer in a subject determined to have an elevated level of CTC-cs and/or IGFBP5 and DCN present in the blood and/or stoma of the cancer, the method comprising administering a therapeutically effective amount of a DCN and IGFBP5-targeted therapy to the subject.
4. The method of claim 3, wherein the targeted therapy comprises an inhibitor of DCN and/or IGFBP5 gene.
5. The method of claim 4, wherein the inhibitor is an antibody reagent or inhibitory nucleic acid reagent.
6. The method of any of claims 3-5, wherein the targeted therapy comprises a DCN and/or IGFBP5-binding antibody reagent and a chemotherapeutic agent.
7. A method of determining if a subject is likely to respond to treatment with a CTC marker gene-targeted therapy, the method comprising:

isolating CTCs from the sample wherein the sample is blood or the stroma of a cancer; measuring the level of a DCN and IGFBP5 expression product present in the the CTCs; and


determining that the subject is likely to respond to the treatment if the level of the expression product is increased relative to a reference level.
8. The method of any of claims 1-2 or 7, wherein the CTC marker gene is selected from Table 7; Table 8; or Table 14.
9. The method of any of claims 1-2 or 7-8, wherein the CTC marker gene is selected from the group consisting of:

ABI3BP; ADAMTS5; ADAMTSL1; ANG; ARSA; C1RL; C3; C4A; C4B; CCDC80; CD109; CHI3L1; CLEC3B; CMTM3; CMTM7; COL14A1; COL1A2; COL3A1; COL4A6; CSF1; DAG1; DCN; DMKN; FBLN1; FGF1;



FMOD; GPC3; GPC4; HMGB1; IFNAR2; IGFBP5; IL16; LAMA4; LTBP4; MFAP1A; NID2; OGN; PDAP1; PF4; PLAT; PODN; PRELP; RSPO1; SERPING1; SLURP1; SOD3; SPARC; SPOCK2; SPON2; SULF1; SULF2; TGFB2; TGM2; THBD; THBS1; THSD4; TIMP2; TNXB; TPT1; TWSG1 and WNT4.

10. The method of any of claims 1-2 or 7-9, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A1; ALDH1A2; IGFBP5; KLF4; DCN; SPARC; WNT; TGFB2; VEGF; COL1A2; COL3A1; and TIMP2.
11. The method of any of claims 1-2 or 7-9, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A2; IGFBP5; KLF4; DCN; and SPARC.
12. The method of any of claims 1-2 or 7-9, wherein the CTC marker gene is selected from the group consisting of:  
ALDH1A2; IGFBP5; KLF4; and DCN.
13. The method of any of claims 1-2 or 7-9, wherein the CTC marker gene is selected from the group consisting of:  
TPT1; HMGB1; SPON 2; SPARC; and ARSA.
14. The method of any of claims 1-2 or 7-9, wherein the CTC marker gene is selected from the group consisting of:  
IL6ST; ARSA; TIMP2; CD55; SULF2; ITGA6; SDC4; CDON; and SV2A.
15. A method of monitoring the treatment of a subject, the method comprising:  
administering a cancer therapy to a subject in need thereof;  
isolating CTCs from a sample obtained from the subject wherein the sample is blood and/or the stroma of a cancer;  
measuring the level of a DCN and IFGBP5 expression product present in the CTCs; and  
determining that the subject is responding if the level of the gene expression products are decreased relative to the reference level and determining that the subject is not responding to the treatment if the gene expression products are not decreased relative to the reference level.
16. The method of claim 3, 7, or 15, wherein the cancer is pancreatic cancer.
17. The method of claim 15, wherein the reference level is the level of the gene expression product in the patient prior to the administering step.
18. The method of any of claims 1-2, or 7-17, wherein the expression product is a nucleic acid.


19. The method of claim 18, wherein the level of the expression product is determined using a method selected from the group consisting of:  
RT-PCR; quantitative RT-PCR; Northern blot; microarray based expression analysis; next-generation sequencing; and RNA in situ hybridization.
20. The method of any of claims 1-2, 7-17, wherein the expression product is a polypeptide.
21. The method of claim 20, wherein the level of the expression product is determined using a method selected from the group consisting of:  
Western blot; immunoprecipitation; enzyme-linked immunosorbent assay (ELISA); radioimmunological assay (RIA); sandwich assay; fluorescence in situ hybridization (FISH); immunohistological staining; radioimmunometric assay; immunofluorescence assay; mass spectroscopy; FACS; and immunoelectrophoresis assay.




2/24

**FIG. 1B****FIG. 1C**


3/24





4/24




5/24

**FIG. 3B**

6/24

**FIG. 4A****FIG. 4B**

7/24




***FIG. 4C***

8/24

MOUSE ID	Kras	Cre	Trp53	GENDER	AGE (WEEKS)	GROSS METASTASES	IF CK+ CTC/mL BLOOD	CELLS PICKED	CELLS SEQUENCED
MP1	G12D	Pdx1	VL	M	5.86	NO	118	3	N/A
MP2	G12D	Pdx1	VL	M	5.00	NO	1684	36	16
MP3	G12D	Pdx1	VL	M	6.14	NO	0	24	8
MP4	G12D	Pdx1	VL	F	8.00	YES	28	42	15
MP5	G12D	Pdx1	VL	M	6.00	NO	240	3	N/A
MP6	G12D	Pdx1	VL	F	6.43	NO	861	24	16
MP7	G12D	Pdx1	VL+	F	16.71	YES	63	42	22

**FIG. 5A**

9/24



10/24






FIG. 5C


11/24

**FIG. 6**


12/24



13/24


**FIG. 7**

14/24




**FIG. 8**

15/24




16/24






***FIG. 10***


17/24






18/24



**FIG. 12A****FIG. 12B****FIG. 12C**

19/24




**FIG. 14A****FIG. 14B****FIG. 14C**



21/24

**FIG. 14D****FIG. 14E**


22/24




23/24

**FIG. 16A****FIG. 16B**

24/24



**FIG. 16C**



**FIG. 16D**