



US005789551A

United States Patent [19]

[11] Patent Number: 5,789,551

Pestka

[45] Date of Patent: Aug. 4, 1998

- [54] **HUMAN LEUKOCYTE INTERFERON HU-IFN- α 001**
- [75] Inventor: **Sidney Pestka**, North Caldwell, N.J.
- [73] Assignee: **Pestka Biomedical Laboratories, Inc.**, West Caldwell, N.J.
- [21] Appl. No.: **257,784**
- [22] Filed: **Jun. 10, 1994**

Related U.S. Application Data

- [63] Continuation-in-part of Ser. No. 076,231, Jun. 11, 1993, abandoned.
- [51] Int. Cl.⁶ **C07K 14/56; C12N 15/21**
- [52] U.S. Cl. **530/351; 424/85.4; 435/69.51; 435/71.1; 435/252.3; 435/320.1; 514/2; 514/12**
- [58] **Field of Search** **530/351, 395; 435/69.51, 172.3, 240.2, 252.3, 320.1, 71.1, 71.2; 935/4, 11, 22, 52, 70, 72; 514/2, 8, 12; 424/85.4**

References Cited

U.S. PATENT DOCUMENTS

3,699,222	10/1972	Isaacs et al.	424/85
4,683,195	7/1987	Mullis et al.	435/6
4,695,623	9/1987	Stabinsky	530/351
4,801,685	1/1989	Goeddel et al.	530/351
5,008,182	4/1991	Sninsky et al.	435/5
5,175,383	12/1992	Leder et al.	800/2
5,175,384	12/1992	Krimpenfort et al.	800/2
5,175,385	12/1992	Wagner et al.	800/2
5,372,808	12/1994	Blatt et al.	424/85.4

OTHER PUBLICATIONS

- Mizoguchi et al., (1985) DNA, vol. 4, No. 3, pp. 221-232.
- Wang et al. (1994) J. Interferon Res. 14:41-6.
- Emanuel et al. (1993) J. Biol. Chem. 268:12565-9.
- Grander et al. (1992) Blood 79:2076-83.
- Zoon et al. (1992) J. Biol. Chem. 267, No.21:15210-6.
- Desai et al. (1992) J. Interferon Res. 12:s138.

- Hosoi et al. (1992) International Society for Study of the Liver (Brighton, UK):113.
- Antonelli et al. (1991) J. Infectious Dis. 163:882-5.
- Colamonici et al. (1991) J. Interferon Res. 11:s54.
- Diaz et al. (1991) J. Interferon Res. 11:s85.
- Adolf et al. (1991) Biochem. 276:511-8.
- Moormeier et al. (1989) Leukemia and Lymphoma 1:43-5.
- Freund et al. (1989) British J. Haematology 72:350-6.
- Steis et al. (1988) New Eng. J. Med. 318:1409-13.
- Hotta et al. (1988) J. Interferon Res. 8:51-60.
- Itri et al. (1987) Cancer 59:668-74.
- Von Wussow et al. (1987) J. Interferon Res. 7:680.
- Pestka et al. (1987) Ann. Rev. Biochem. 56:727-77.
- Pestka, S. (1986) Methods in Enzymology 119:3-23.
- Hotta et al. (1986) Methods in Enzymology 119:481-5.
- Foon et al. (1986) American J. Med. 80:351-6.
- Huber et al. (1985) Oncology 42, suppl. 1:7-9.
- Henco et al. (1985) J. Mol. Biol. 185:227-60.
- Quesada et al. (1985) J. Clinical Oncology 3:1522-8.
- Langer et al. (1984) J. Investigative Dermatology 83:128s-36s.
- Pestka, S. (1983) Arch. Biochem. Biophys. 221:1-37.
- Rehberg et al. (1982) J. Biol. Chem. 257:11497-502.
- Hobbs et al. (1982) J. Biol. Chem. 257:4071-6.
- Pestka et al. (1981) Methods in Enzymology (S. Pestka, ed.) 78:3-14.
- Staehelin et al. (1981) J. Biol. Chem. 256:9750-4.
- Rubinstein et al. (1981) Arch. Biochem. Biophys. 210:307-18.
- Streuli et al. (1980) Science 209:1343-7.
- Goeddel et al. (1980) Nature 287:411-6.
- Koeffler et al. (1978) Science 200:1153-4.

Primary Examiner—John Ulm

Assistant Examiner—Prema Mertz

Attorney, Agent, or Firm—Klauber & Jackson

[57]

ABSTRACT

A new class of polypeptides is disclosed, along with a method for identifying and producing such polypeptides, having the characteristic of being unique to diseased states, particularly tumors and blood-borne malignancies. These new polypeptides are active and will be useful for therapeutic purposes.

7 Claims, 5 Drawing Sheets

		ATGGCCTTG MetAlaLeu -23	9
10	TCTTTTCTTTACTGATGCTGCTGCTGCTACTCAGCTACAAATCCATCTGCTCTCTGGGC SerPheSerLeuLeuMetValValLeuValLeuSerTyrLysSerIleCysSerLeuGly -20	-10	69
70	TGTGATCTGCTCAGACCCACAGCCTGCTAATAGAGGGGCTTGAATACTCTCGCACAA CysAspLeuProGlnThrIleSerLeuArgAsnArgAlaLeuIleLeuLeuAlaGln -10	-20	129
130	ATGGGAGAAATCTCTCTTCTCTCTGCTGCTGAGGACAGACATGAATTCAGATTCCAGAG MetGlyArgIleSerProPheSerCysLeuLysAspArgHisGluPheArgPheProGlu -30	-40	189
190	GAGGAGTTGATGGCACCAGTCCAGAAAGCTCAACCATCTCTGCTCCTCATGAGATG GluGluPheAspGlyHisGlnPheGlnLysThrGlnAlaIleSerValLeuAlaGluMet -40	-50	249
250	ATCCAGCAGACCTTCAATCTCTTCCAGCAGAGGAGGACTCTCTGCTGCTGGGACAGAGC IleGlnGlnThrPheAsnLeuPheSerThrGluAspSerSerAlaAlaTrpGluGlnSer -70	-80	309
310	CTCTAGAAAAATTTTCCACTGAACCTTACAGCACTGAATGACCTGGAGCATGTGTG LeuLeuGluLysPheSerThrGluLeuTyrGlnGlnLeuAsnAspLeuGluAlaCysVal -80	-90	369
370	ATACAGAGAGTTGGGTTGAGAGAGACTCCCTCTGTAATGAGGACTCCATCTGCTGCTG IleGlnGluValGlyValGluGlnThrProLeuMetAsnGluAspSerIleLeuAlaVal -110	-120	429
430	AGCAATACTTCCAGAAATCACTCTTATCTAAGCAGAGAAATACAGCCCTGTGCTG ArgLysTyrPheGlnArgIleThrLeuTyrIleuThrGluLysLysTyrSerProCysAla -130	-140	489
490	TGGGAGTTGTGAGAGCAGAAATCATGACATCCCTCTGCTTTTCAACAACTTGCAGAAA TrpGluValValArgAlaGluIleMetArgSerLeuSerPheSerThrAsnLeuGlnLys -150	-160	549
550	AGATTAGGAGGAGCATGTA ArgLeuArgArgLysAspEnd	570 168	

9	ATGGCCTTG	
	MetAlaLeu	
	-23	
69		
10	TCCTTTTCTTACTGATGGTCGTGCTGGTACTCAGCTACAAATCCATCTGCTCTCTGGGC	
	SerPheSerLeuLeuMetValValLeuValLeuSerTyrLysSerIleCysSerLeuGly	
	-20	-1
70	TGTGATCTGCCCTCAGACCCACAGCCTGCCGTAATAGGAGGGCCTTGATACTCCTGGCACAA	
	CysAspLeuProGlnThrHisSerLeuArgAsnArgArgAlaLeuIleLeuLeuAlaGln	
	1	20
130	ATGGGAAGAAATCTCTCCTTCTCCTGCTTGAAGGACAGACATGAATTCAGATTCCCAGAG	
	MetGlyArgIleSerProPheSerCysLeuLysAspArgHisGluPheArgPheProGlu	
	30	40
190	GAGGAGTTTGATGGCCACCACTTCCAGAAAGACTCAAGCCATCTCTGTCTCCATGAGATG	
	GluGluPheAspGlyHisGlnPheGlnLysThrGlnAlaIleSerValLeuHisGluMet	
	50	60
250	ATCCAGCAGACCTTCAATCTCTTTCAGCACAGAGACTCATCTGCTGCTTGGGAACAGAGC	
	IleGlnGlnThrPheAsnLeuPheSerThrGluAspSerSerAlaAlaTrpGluGlnSer	
	70	80
309		

FIG. 1A

310	CTCCTAGAAAATTTTCCACTGAACTTTAC <u>CAGCAACTGAATG</u> ACCTGGAAAGCATGTGTG	369
	LeuLeuGluLysPheSerThrGluLeuTyrGlnGlnLeuAsnAspLeuGluAlaCysVal	100
	90	
370	ATACAGGAGGTTGGGGTGGAGAGACTCCCTGATGAATGAGGACTCCATCCTGGCTGTG	429
	IleGlnGluValGlyValGluGluThrProLeuMetAsnGluAspSerIleLeuAlaVal	120
	110	
430	AGGAAATACTTCCAAAGAATCACTCTTTATCTAACAGAGAGAAATACAGCCCTTGTGCC	489
	ArgLysTyrPheGlnArgIleThrLeuTyrLeuThrGluLysLysTyrSerProCysAla	140
	130	
490	TGGGAGGTTGTCAGAGCAGAAATCATGAGATCCCCTCTCGTTTTCACACAAACTTGCAAAA	549
	TrpGluValValArgAlaGluIleMetArgSerLeuSerPheSerThrAsnLeuGlnLys	160
	150	
550	AGATTAGGAGGAAGGATTGA	570
	ArgLeuArgArgLysAspEnd	166

FIG. 1B

1	MALSFSL	MVVL	SYKSI	CSLG	CDLP	QTHSL	RNRRA	LILLA	QMGRI	SPF	50
1	MARFSL	LMVVL	SYKSI	CSLG	CDLP	QTHSL	RNRRA	LILLA	QMGRI	SPF	50
51	SCLKDR	HEFR	FPEEE	FDGH	QFQK	TQAIS	VLHEMI	QQTFN	LFSTED	SSAAW	100
51	SCLKDR	HEFR	FPEEE	FDGH	QFQK	TQAIS	VLHEMI	QQTFN	LFSTED	SSAAW	100
101	EQSLLE	KFST	ELYQ	QLND	LEAC	VIQEV	GVVEET	PLMNED	SILAVR	KYFQRI	150
101	EQSLLE	KFST	ELYQ	QLND	LEAC	VIQEV	GVVEET	PLMNED	FILAVR	KYFQRI	150
151	TLYLTE	KKYSP	CAWE	VVRAE	IMRSL	SFSTN	LQKRL	RRKD			189
151	TLYLME	KKYSP	CAWE	VVRAE	IMRSL	SFSTN	LKKGL	RRKD			189

FIG. 2




FIG. 3




FIG. 4

HUMAN LEUKOCYTE INTERFERON HU- IFN- α 001

CROSS REFERENCE TO RELATED APPLICATION

This application is a continuation in part of U.S. application Ser. No. 08/076,231, filed Jun. 11, 1993 now abandoned.

FIELD OF THE INVENTION

This invention relates to the field of biotechnology and more particularly to polypeptides having antitumor, antiviral, immunomodulatory and other activities, DNA that codes for such polypeptides, recombinant vectors that include such DNA, host organisms transformed with the recombinant vector that produces the polypeptide and therapeutic application of the polypeptide.

The publications and other materials hereof used to illuminate the background of the invention, and in particular cases, to provide additional details respecting its practice are hereby incorporated by reference, and for convenience are numerically referenced by the following text and respectively grouped in the appended bibliography.

BACKGROUND OF THE INVENTION

Human leukocyte interferon was first discovered and prepared in the form of very crude fractions by Isaacs and Lindenmann (1, 2). Efforts to purify and characterize the material have led to the preparation of relatively homogeneous leukocyte interferons derived from normal or leukemic (chronic myelogenous leukemia or "CML") donors' leukocytes. These interferons are a family of proteins characterized by a potent ability to confer a virus-resistant state in their target cells. In addition, interferon can inhibit cell proliferation, modulate immune responses and alter expression of proteins. These properties have prompted the clinical use of leukocyte interferon as a therapeutic agent for the treatment of viral infections and malignancies.

With the advent of recombinant DNA technology, the controlled microbial production of an enormous variety of useful polypeptides has become possible. The workhorse of recombinant DNA technology is the plasmid, a non-chromosomal circle of double-stranded DNA found in bacteria and other microbes, oftentimes in multiple copies per cell. Included in the information encoded in the plasmid DNA is that required to reproduce the plasmid (i.e., an origin of replication) and ordinarily, one or more selection characteristics such as, in the case of bacteria, resistance to antibiotics which permit clones of the host cell containing the plasmid of interest to be recognized and preferentially grown in selective media. The utility of plasmids lies in the fact that they can be specifically cleaved by one or another restriction endonuclease or "restriction enzyme," each of which recognizes a specific site in the DNA. Heterologous genes or gene fragments may be inserted into the plasmid at the cleavage site. To construct vectors with specific sequences inserted, DNA recombination is performed outside the cell, but the resulting "recombinant" plasmid can be introduced into cells by a process known as transformation and large quantities of the heterologous gene-containing recombinant plasmid obtained by growing the transformant. Moreover, where a promoter which governs the transcription of the encoded DNA message, is properly placed upstream (5') of a coding sequence or a gene, the resulting expression vector can be used to produce the polypeptide sequence for

which the inserted sequence or gene codes, a process referred to as expression.

Expression is initiated in a region known as the promoter which is recognized by and bound by RNA polymerase. In many cases promoter regions are overlapped by "control" regions such as the bacterial operators. Operators are DNA sequences which are recognized by so-called repressor proteins which serve to regulate the frequency of transcription initiation at a particular promoter. The polymerase travels along the DNA, transcribing the information contained in the coding strand from its 5' to 3' end into messenger RNA (mRNA) which is in turn translated into a polypeptide having the amino acid sequence for which the DNA codes. Each amino acid is encoded by a nucleotide triplet or "codon" within the coding sequence, i.e., that part which encodes the amino acid sequence of the expressed product. In bacteria (e.g. *Escherichia coli*) the mRNA contains a ribosome binding site, a translation initiation or "start" signal (ordinarily ATG in the DNA, which in the resulting mRNA becomes AUG), the nucleotide codons within the coding sequence itself, one or more stop codons, and an additional sequence of messenger RNA, the 3' untranslated region. Ribosomes bind to the binding site provided on the messenger RNA, in bacteria ordinarily as the mRNA is formed, and produce the encoded polypeptide, beginning at the translation start signal and ending at the stop signal. The desired product is produced if the sequences encoding the ribosome binding site are positioned properly with respect to the AUG initiator codon and if all remaining codons follow the initiator codon in phase. The resulting product may be obtained from the host cell and recovered by appropriate purification. In other systems, proteins may be secreted from the host cells. A wide variety of expression vectors and host systems exist so that RNA and proteins may be expressed in prokaryotic and eukaryotic cells as well as in intact animals and plants.

During the past several decades a large number of human and animal interferons have been produced, identified, purified and cloned (see ref. 1-72). Several of the interferon preparations have been prepared for clinical trial in both crude form, for some of the original interferon preparations, as well as in purified form. Several individual recombinant interferon species have been cloned and expressed. The proteins have then been purified by various procedures and formulated for clinical use in a variety of formulations (73). Most of the interferons in clinical use that have been approved by various regulatory agencies throughout the world are mixtures or individual species of human interferon (Hu-IFN- α). In some countries Hu-IFN- β and γ have also been approved for clinical trial and in some cases approved for therapeutic use (56,74). The major thesis underlying clinical use of these interferons was that they were natural molecules produced by normal individuals. Indeed, the specific thesis was that all the interferons prepared for clinical use, be they natural- or recombinant-generated products, represented interferons that were produced naturally by normal people. This is true for a large number of interferons as well as specific growth factors, lymphokines, cytokines, hormones, clotting factors and other proteins that have been produced (17, 21, 22, 25-27, 29-34, 39, 40, 45-51, 53-57, 62-64, 68-72).

Reports have suggested that Hu-IFN- α A (also designated Hu-IFN- α 2a and by the trade name Roferon A) was not represented in interferons produced by a normal population of individuals (75-79). Believing that certain interferons (or, more generically, certain polypeptides) are uniquely found in diseased cells, the inventor of the present invention

undertook to identify interferons which are so uniquely characterized. For convenience the inventor began by screening known interferons, in particular, the sources of the several variants of Hu-IFN- α 2 that have been described. As discussed more fully below, it was found that the source of two of the variants of Hu-IFN- α 2, Hu-IFN- α 2a and Hu-IFN- α 2c, are not present in normal individuals. Only Hu-IFN- α 2b is found in normal individuals (79).

DESCRIPTION OF THE DRAWINGS

FIG. 1. Nucleotide and Amino Acid Sequence of Hu-IFN- α 001 (SEQ ID NO:12). The location of the AlwNI site is underlined. The signal peptide is shown as the 23 amino acids labeled -1 to -23.

FIG. 2. Comparison of the Protein Sequence of Hu-IFN- α 001 with that of Hu-IFN- α J. The signal peptide represents the first 23 amino acids at the amino terminus.

FIG. 3. Expression vector for Hu-IFN- α 001. The structure of the plasmid pHu-IFN- α 001 is shown. The NsiI site represents nucleotide position #1. The P_R promoter drives expression of Hu-IFN- α 001.

FIG. 4. SDS-Polyacrylamide Gel Electrophoresis of the Purified Hu-IFN- α 001. Hu-IFN- α 001 was placed in lanes 1, 2 and 3 in amounts of 3 μ g, 1.5 μ g and 0.75 μ g, respectively. The columns labeled M represent the molecular weight markers with the values in kilodaltons given to the left of each respective molecular weight marker.

SUMMARY OF INVENTION

An extensive analysis of normal individuals from various ethnic and racial backgrounds as well as two tumor cell lines has shown that certain interferons originated from two cell lines that were obtained from patients with disease, in particular, malignancies of the hemopoietic system (79). These results lead the inventor to conclude that there is a new class of interferon molecules which are present in diseased states, specifically in tumors and blood borne malignancies. This discovery of a new class of interferons provides a wide variety of potentially new interferons for clinical and therapeutic use. These interferons include not only Hu-IFN- α species, but also Hu-IFN- β , Hu-IFN- γ and Hu-IFN- ω , as well as other newly described interferons in other animals and species. The observations suggest that growth factors, cytokines, lymphokines, clotting factors, peptide and polypeptide hormones, adhesion factors and many other molecules are also modified in disease processes. Therefore, modified forms of all these cytokines, lymphokines, growth factors, adhesion molecules, enzymes, clotting factors, peptide and polypeptide hormones, etc. will also occur in tumors and other diseases. Based on two presently identified members of this class (not previously recognized as such), these interferons are active, are as active as the standard molecules, and in fact have been used effectively for therapeutic purposes. A paper co-authored by the inventor and listed hereafter as Reference No. 79 is particularly related to the invention and is incorporated herein by reference insofar as may be needed for a full understanding of the invention. That paper is more fully described, and is furnished as an attachment to a contemporaneously filed Information Disclosure Statement.

DETAILED DESCRIPTION OF THE INVENTION

Four distinct classes of interferons (IFNs) are known to exist in humans. The IFN- α family represents the predomi-

nant class of IFNs and are produced by stimulated peripheral blood leukocytes (10-15, 17-27, 29, 50, 51, 57-59, 61, 63, 64, 68, 70), and lymphoblastoid and myeloblastoid cell lines (28, 30, 60). Cloning of the IFN- α genes from these cells has revealed that IFN- α is encoded by a multigene family consisting of about 15 functional genes and four pseudogenes (17, 26, 27, 29, 31, 50, 51, 53, 54, 57, 61, 63, 64, 65). It has been uncertain whether or not some of the cloned human IFN- α genes and cDNAs with few nucleotide differences, such as the Hu-IFN- α A, Hu-IFN- α 2 and Hu-IFN- α 2(Arg) genes, are allelic variants or represent distinct genes.

To determine if these sequences do indeed represent separate genes or are instead polymorphic variants of a single gene, sequences representing only the Hu-IFN- α A, Hu-IFN- α 2 and Hu-IFN- α 2(Arg) genes were amplified by nested polymerase chain reaction (PCR) from human genomic DNAs of healthy consenting individuals. These sequences were then subcloned and examined by sequencing of individual clones. In addition, the DNAs were examined from KG-1 (80) and Namalwa (81) cell lines from which the Hu-IFN- α A and Hu-IFN- α 2 (Arg) cDNAs, respectively, were cloned.

MODES FOR CARRYING OUT INVENTION

Three oligodeoxynucleotides were prepared by the phosphoramidite method (82, 83) and purified (84). Primer I (5'-TGGGCTGTGATCTGCCTC-3') (SEQ ID NO: 1) complementary to nucleotides 125 to 142 at the 5' end was used with Primer II (5'-CATGATTCTCTGCTGACAACC-3') (SEQ ID NO:2) complementary to nucleotides 552 to 573 at the 3' end to amplify the desired nucleotide sequences. The DNA, as amplified by the polymerase chain reaction (PCR) with this primer pair, was expected to represent sequences from most of the IFN- α gene family (79). This conserved PCR product was then used as template in a second amplification reaction with the same 3' oligonucleotide but with a 5' oligonucleotide specific for the human IFN- α A, IFN- α 2 and IFN- α 2(Arg) genes only (79). The second reaction produced a product of 430 bp when Primer III (5'-AACCACAGCCTGGGTAG-3') (SEQ ID NO:3) complementary to nucleotides 144 to 161 was substituted for the Primer I. The 430 bp DNA was purified and cloned into the SmaI site of pBluescript-SK⁺ (Stratagene, LaJolla, Calif.) as described (79, 85, 86).

DNA of the plasmids was prepared by the alkaline lysis miniprep procedure (86, 87) from 1 ml cultures grown overnight in LB medium containing 100 μ g/ml ampicillin. The resultant DNA pellet was sequenced by the dideoxy chain termination procedure (79, 88, 89). The reactions were run on 6% polyacrylamide gels which were then dried and exposed to X-ray film overnight at room temperature with an intensifying screen.

Reverse transcriptase PCR (RT-PCR) was used to analyze the expression of the IFN- α subtypes α A, α 2 and α 2(Arg) in the KG-1 and Namalwa cell lines (90). RNA was isolated at 6 hours after induction from Sendai virus-induced KG-1 cells (60) and at 8 hours post induction from NDV-stimulated Namalwa cells (91, 92).

DNA was extracted from the human myeloblastoid cell line KG-1 and from the lymphoblastoid Namalwa cell line by a modification of the method of Pellicer et al. (93). After obtaining informed consent, human genomic DNA was prepared from whole blood samples collected from normal, healthy individuals by ammonium acetate precipitation as described (79, 94).

METHODOLOGICAL BASIS FOR INVENTION

The DNA from 11 normal individuals was amplified by nested PCR then cloned and sequenced as described above. The number of sequences corresponding to the various human IFN- α species is shown in Table 1. It can be seen that neither the sequence for the αA gene nor the $\alpha 2(\text{Arg})$ gene was detected in any of the normal individuals examined in this study. As shown in Table 2, however, the αA sequence was detected in the DNA from the KG-1 cell line, but not in Namalwa cells; and the $\alpha 2(\text{Arg})$ sequence was detected in the DNA from the Namalwa cell line, but not in KG-1 cells.

TABLE 1

Frequency of Hu-IFN- αA , - $\alpha 2$ and $\alpha 2(\text{Arg})$ Clones From Normal Individuals

Interferon Variant	Number of Clones
IFN- $\alpha 2$	165
IFN- αA	0
IFN- $\alpha 2(\text{Arg})$	0
Other ¹	36
Total	201

¹Other refers to sequences which contained one or more mutations in an area unrelated to the αA and $\alpha 2(\text{Arg})$ specific differences. It should be noted that the frequency of mutations detected is in the range or slightly lower than that predicted from the combined error rates of Taq DNA polymerase and Sequenase DNA polymerase (95, 96). Previous analysis of IFN- $\alpha 2$ genes have been reported (97, 98), but did not discern any differences in their representation in the DNA from normal individuals. Descriptions and abbreviations relevant to interferons are described in detail in several references (10-12, 61, 99, 100).

TABLE 2

Frequency of IFN- α Clones From KG-1 and Namalwa Cell Lines

Cells	IFN- $\alpha 2$	IFN- αA	IFN- $\alpha 2(\text{Arg})$	Other	Total
KG-1 cells	15	10	0	16	41
Namalwa cells	22	0	13	2	37

Restriction endonuclease analysis to detect the IFN- αA gene was also performed on DNA from five of the individuals from whom clones had been sequenced and on DNA from seven additional people that were not examined by DNA sequencing. It was found that the restriction endonuclease analysis of the amplified DNA from all of these individuals showed no IFN- αA gene present (See Ref. 79, FIG. 2).

PREFERRED EMBODIMENT OF INVENTION

From the foregoing analysis, it can be concluded that in human DNA from a wide variety of humans only Hu-IFN- $\alpha 2$ is present. The species Hu-IFN- αA and Hu-IFN- $\alpha 2(\text{Arg})$, not present in the DNA of 11 normal individuals, apparently arose during the development of the disease and/or the establishment of the cell lines in culture. It is noteworthy that the expression of these alleles of Hu-IFN- $\alpha 2$ yields IFN- α species with high activity in a wide variety of assays (63, 68, 69, 101-115). The specific activities of all three of these IFN- α species are comparable. Furthermore, it has been reported that patients treated with Hu-IFN- αA produced a higher level of anti-interferon antibodies than patients treated with Hu-IFN- $\alpha 2$ or Hu-IFN- αn (Welferol: a preparation of mixed Hu-IFN- α species produced by induced Namalwa cells) (116-124). Some of the new interferons produced by the described invention may be able to by-pass neutralization by the antibodies produced in patients treated with IFN- α preparations in current use. Such new

IFN- α species should be able to be used to treat patients who have relapsed because of neutralization of the administered IFN- α species.

While the inventor has, for convenience, used Hu-IFN- $\alpha 2$ and its known variants for establishing his hypothesis of the existence of a class of super or tumor interferons, it will be apparent to those skilled in the art that the results extend to an entire class of such interferons, as well as other polypeptides. Illustrative of this conclusion is the extraordinary high percentage of variant forms of the IFN- $\alpha 2$ and αA genes in KG-1 cells—i.e., 39% (19/41), much more than could be explained by experimental error, as shown in the column labeled "Other" of Table 2.

It will also be apparent that the method of the invention, as illustrated above for Hu-IFN- $\alpha 2$, can be applied to any protein. In the general case, a primer pair is chosen to encompass part or all of the nucleotide coding sequence with the use of DNA from tumor cells or from cultured cells as templates for the PCR. The PCR product is then cloned and sequenced. The amino acid sequence predicted by the nucleotide sequence so obtained is compared to the sequence of the protein in normal individuals. Proteins with amino acid sequences different from those proteins in normal individuals are then cloned in appropriate expression vectors (11, 12, 14, 17, 45, 53, 54, 57, 63, 69, 86, 103), produced, purified and characterized. Those with desirable activities are then developed for therapeutic use.

The origin of the tumor interferons or super interferons is unknown. Yet, it is clear that they are developed during the pathological process. It is believed that the cells producing these interferons have been selected during development and progress of the disease.

The presence of allelic forms of IFN- $\alpha 2$ in the KG-1 and Namalwa cells is most noteworthy. DNA from leukocytes from normal individuals did not contain these variants. Because both the KG-1 and Namalwa cells originated from patients with leukemia or lymphoma, it is believed that this alteration is an early change in progression of these diseases. Indeed, it has been reported that there are significant gross changes in restriction endonuclease digestion patterns for the IFN- α genes in acute leukemias (125, 126).

The disease mechanisms involved in developing malignant cells and selection of those cells produce a wide variety of genetic changes in the resultant tumor cells. In order for cancer cells to grow unfettered, to escape the normal controls and to metastasize, the usual regulatory network of the immune system, involving growth inhibitors as the interferons and growth factors and hormones, may be modified. The control of cell growth and nonmalignant behavior is a delicate balance of many regulatory factors, a few of which gone astray can alter the normal growth patterns. Although it has been reported that changes in the DNA of cancer cells occurs, the changes have been focussed on oncogenes and tumor suppressor genes that lead to the malignant phenotype. The inventor has provided data that the changes are more pervasive than expected, not merely those changes focussed on oncogenes and tumor suppressors. Furthermore, by genetic changes (mutations in DNA) and selection of tumor cells for aggressiveness, many alterations will be embodied in the final tumor cell population. The new proteins produced will have lower, the same or higher activities than the normal proteins. By identifying those modified proteins associated with changes in activity, it will be possible to identify those proteins with new and/or enhanced activities.

From an analysis of initial clones obtained from the KG1 cell line (53, 54, 63), it was shown that several abnormal interferons exist in this cell line (also see ref. 61 for list of IFN- α species). This is especially evident in that αB (not previously recognized as an abnormal interferon) has an

insertion and compensating deletion making an abnormal protein that differs from Hu-IFN- α 8, the normal counterpart. The presence of an insertion and a compensating deletion producing a normal sized molecule suggests some enormous selective pressures to produce these interferons. The fact that an insertion and a deletion would be incorporated into a molecule simply by random stochastic processes without external pressures is highly unlikely. Thus, these modified interferons are seen to represent an entire new family of molecules that have been developed under the pressure of enormous external forces to provide for the selection of these species.

With respect to the interferons produced by mechanisms to enhance or combat the disease process, some may indeed have unusual properties and may be more active than some of the interferons produced by normal cells. For example, since interferon is a growth inhibitory molecule, production of a new interferon that could down regulate the receptors for interferon in cells and help select for cells without interferon receptors or low levels of interferon receptors may enhance the disease process. Such interferons could also help select cells with an altered signal transduction mechanism, but normal receptor number. Thus, a cell producing spontaneously some interferon, could be expected to have initially a low level of receptors due to down regulation and its growth would likely be reduced. Nevertheless, during the multiplication of such cells, cells would be selected that would have low levels of receptors so that they could escape the inhibition of the endogenous interferon. The same would hold for a wide variety of other molecules such as cytokines, lymphokines, tumor suppressors, growth factors, anti-growth factors, matrix molecules, hormones, angiogenic factors, clotting factors, etc., all molecules that can control growth and/or metastases in one manner or another.

The altered proteins described herein are found in tumor cells or cultured cells obtained from tumors. Furthermore, selection of cell-lines in culture can also produce some of the alterations as selection in vivo.

CLONING OF Hu-IFN- α 001

A new, interferon was amplified from the genomic DNA of KG-1 cells (ATCC CCL 246) based on the strategies outlined hereinbefore and by the procedures described herein and elsewhere (79). The primers used to amplify the genes are shown in Table 3. The 5' primer contains an Apal site, and the 3' primer contains an XbaI site for cloning. The PCR reactions were carried out in 50 μ l with 100 ng KG-1 template DNA, 100 ng of each primer (6431 and 6432), 0.2

and XbaI sites of plasmid pBluescript II KS+(Stratagene) and then transformed into competent *E. coli* strain DH5 α cells. Competent cells were prepared in 12% PEG and 36% glycerol in Luria-Bertani medium (L-broth medium, 10 g tryptone, 5 g yeast extract, 10 g NaCl, pH 7.3) from Digene (Silver Spring, Md. 20904, Cat. No. 3500-1002) as described (127). Plasmid DNA was isolated from 2.0 ml of overnight cultures grown at 37° C. by a modified alkaline lysis procedure as reported (128). The size of the inserts was determined by digestion with restriction endonucleases KpnI and SacI that flanked the cloning sites in the vector pBluescript. A total of 10 independent colonies were identified that contained a 700 base pair insert.

TABLE 3

Primers for PCR Amplification

Primer	Sequence	Length	Primer No.
5'	GCGGGCCCCAATGGCCYTGYCCTTT (SEQ ID NO:4)	25	6431
3'	GCTCTAGAAATCATGAAAGYGTGA (SEQ ID NO:5)	24	6432

The sequence of the primers are given in the 5' to 3' direction. The "Y" represents a pyrimidine (T or C).

The DNA from one of the clones (plasmid pBS001) was sequenced in both directions. Automated DNA sequencing was performed on a Genesis 2000 Automated DNA Sequencer (DuPont, Wilmington, Del.) with the primers shown in Table 4 by methods previously reported (86,88, 89). All sequences were performed on both strands. Automated sequencing was carried out and the results were compiled to create a consensus sequence. The sequence determined from the T3 primer represents the 5' end of the insert; the T7-derived sequence represents the 3' end.

The sequence so determined is designated Hu-IFN- α 001 and is shown in FIG. 1. The location of the AlwNI restriction endonuclease recognition site (5' CAGNNNCTG 3') that was used for the splicing of the Hu-IFN- α 001 insert into the expression vector TGATG (129) is indicated in the figure by underlining. The signal peptide is shown as the 23 amino acids labeled -1 to -23. As seen in FIG. 1, the mature protein contains 166 amino acids.

TABLE 4

Primers used for Sequencing

Designation	Sequence	Primer No.	Position in Hu-IFN- α 001	Direction
IFN-A1	CTTGAAGGACAGACATG (SEQ ID NO:6)	6942	157-172	F
IFN-A2	CTGTCTTCATGAGATG (SEQ ID NO:7)	6941	233-249	F
IFN-A3	GGTCATTCAGCTGCTGG (SEQ ID NO:8)	6940	339-355	R
IFN-A4	TCCTCCTTCATCAGGGG (SEQ ID NO:9)	6939	397-413	R
T3	ATTAAACCCTCACTAAAG (SEQ ID NO:10)	T3	Vector	F
T7	TAATACGACTCACTATA (SEQ ID NO:11)	T7	Vector	R

All primers are shown from 5' to 3' orientation. The column designated "Direction" represents the direction of sequencing with respect to the sequence of the Hu-IFN- α ; "F" represents forward; "R" - reverse. Oligodeoxynucleotides were synthesized on an Applied Biosystem DNA synthesizer model 380B by the phosphoramidite method (83,130).

mM of each dNTP, and 2.5 units of Taq DNA Polymerase for 30 cycles of 94° C. 30 seconds, 50° C. 30 seconds, 72° C. 30 seconds in the Perkin Elmer model 9600 thermocycler. Products of the PCR amplification were cloned into the Apal

A comparison of the protein sequence with other human interferon alpha species (Hu-IFN- α) demonstrates that Hu-IFN- α 001 is most closely related to Hu-IFN- α J. That comparison is graphically depicted in FIG. 2. A summary of

the known Hu-IFN- α sequences has been previously reported (61). There are a total of six amino acid changes compared to Hu-IFN- α J. The data clearly demonstrate that this tumor derived Hu-IFN- α species is different from any other known Hu-IFN- α species previously reported. Furthermore, it would not have been possible to predict this specific sequence as the number of possible proteins with alterations in these six positions is 20^6 or 64,000,000. One of the amino acid changes is in the signal peptide sequence; the remaining five alterations are in the mature protein. It is also to be emphasized that the derived Hu-IFN- α species presented here is a natural interferon derived from tumor cells. It is not a synthetic construct prepared by simply mutating six positions.

Expression of the Hu-IFN- α 001 gene was accomplished in two steps. The plasmid pBS001 was digested with restriction endonuclease KpnI (5' end of Hu-IFN- α 001 sequence). The KpnI ends were made blunt by incubation with T4 DNA polymerase in the following reaction mixture: 1 μ g of DNA; 33 mM Tris acetate, pH 7.9; 66 mM potassium acetate; 10 mM magnesium acetate; 0.5 mM dithiothreitol; 100 μ g/ml BSA (bovine serum albumin); 2 mM of each of the four dNTPs; 5 units of T4 DNA polymerase (United States Biochemical Corp.); total volume of 18 μ l. Incubation was performed for 5 minutes at 37° C. to prepare the blunt ends. The plasmid DNA was then digested with XbaI (3' end of Hu-IFN- α 001 sequence) to release the insert containing the Hu-IFN- α 001 sequence. The DNA fragments were then purified as described (86). The TGATG vector was prepared by digestion with restriction endonuclease SacI, followed by preparing blunt ends with T4 DNA polymerase as described above, and then digested with restriction endonuclease XbaI. The fragment containing the Hu-IFN- α 001 insert was then ligated to the pTGATG expression vector (129). After ligation the DNA was transformed into competent *E. coli* DH5 α cells. Colonies were analyzed by growing the cells as described above to isolate plasmid DNA. The plasmids were then digested with restriction endonucleases EcoRI and XbaI to determine the size of the insert. An expression vector for Hu-IFN- α J was prepared as previously described for the expression plasmids for Hu-IFN- α B2 and Hu-IFN- α A/D (131).

The nucleotide sequences encoding Hu-IFN- α J and Hu-IFN- α 001 contain an AlwNI site in identical positions of the sequence (FIG. 3); and, as illustrated in FIG. 3, which shows the structure of the plasmid pHu-IFN- α 001 containing the expression vector for Hu-IFN- α 001, there is a second AlwNI site in the vector itself.

In addition, because the AlwNI recognition sites (CAGNNN^{*}CTG) have three unspecified nucleotides (NNN) in the 3' overhang, the religations are specific and asymmetric. Accordingly, pTGATG vectors (129) encoding Hu-IFN- α J and Hu-IFN- α 001 were digested with restriction endonuclease AlwNI to isolate the large vector and Hu-IFN- α 001 (3' end) fragments, respectively. The Hu-IFN- α 001 fragment was then ligated into the vector fragment from plasmid pHu-IFN- α J to yield the *E. coli* expression vector pHu-IFN- α 001, as shown in FIG. 3, which was transfected into competent *E. coli* (DH5 α) cells (86).

Plasmid pHu-IFN- α 001 is deposited with the American Type Culture Collection (ATCC) at 12301 Parklawn Drive, Rockville, Md. 20852; Accession number: 69640; Deposit date Jun. 7, 1994; and designated as plasmid pHu-IFN- α 001 (*E. coli* DH5 α /pHu-IFN- α 001 as the host vector system).

The *E. coli* (DH5 α) cells containing the expression vector pHu-IFN- α 001 were grown in 875 ml of Medium A over-

night at 30° C. in one 2 liter flask with rotary shaking. Medium A consists of KH₂PO₄ (4.5 g/L), Na₂HPO₄·7H₂O (18.9 g/L), NH₄Cl (1.5 g/L), NaCl (0.75 g/L), glucose (15 g/L), casamino acids (7.5 g/L), MgSO₄·7H₂O (0.369 g/L), thiamine hydrochloride (0.0015 g/L), leucine (0.04 g/L), proline (0.04 g/L) and ampicillin (0.05 g/L) adjusted to pH 7.4. The overnight culture was used to inoculate 22.5 liters of Medium A in a fermentor. The *E. coli* containing the expression vector were grown at 30° C. until the A₅₅₀ reached 7.0 at which time the temperature was raised to 42° C. The cells were harvested 3 hrs after temperature induction at 42° C. by centrifugation and cell pellets divided into 50 g portions prior to freezing at -80° C. The cells were stored at -80° C. until used for preparation of interferon.

PURIFICATION OF Hu-IFN- α 001

For purification of Hu-IFN- α 001, frozen *E. coli* cell paste was thawed by suspension in 10 volumes of Buffer A (50 mM Tris-HCl, pH 8.0, 50 mM NaCl, 10 mM EDTA, 0.1 mM PMSF, phenylmethylsulfonyl fluoride). After the addition of egg white lysozyme (0.2 mg/ml) the suspension was sonicated four times with 30 second bursts while kept in an ice bath, then incubated at 23° C. overnight while stirring vigorously to eliminate viscosity contributed by DNA. The suspension was centrifuged for 20 minutes at 12,000 rpm at 4° C. The pellet was resuspended again in 10 volumes of Buffer A with 1% Triton X-100, 50 mM EDTA and 0.5M NaCl and incubated for at least 2 hours (2-16 hrs) at room temperature with shaking and then centrifuged for 20 min at 12,000 rpm at 4° C. Once again, the pellet was resuspended in 5 volumes of Buffer A with 0.5M NaCl and incubated for 60 min at room temperature with shaking and then centrifuged for 20 min at 12,000 rpm at 4° C; the supernatant was discarded. The pellet was dispersed in 2 volumes of Buffer A in the presence of a mixture of oxidized/reduced forms of glutathione (0.2 mM/2.0 mM) and solid guanidine.HCl (2.5 times bacterial weight) was added and the solution was stirred at room temperature for 7 hours. After this, the mixture was diluted tenfold with Buffer A and allowed to stand overnight. Renaturation of the interferon was carried out by very slow addition of 7M guanidine.HCl to 0.7M. The refolding of Hu-IFN- α 001 in solution takes more than 15 hours. Since Hu-IFN- α 001 contains two disulfide bonds, this step involves slow oxidation of the protein during dilution from guanidine-containing solution. Then suspension was then centrifuged to remove debris. Solid (NH₄)₂SO₄ was added to the supernatant to a final concentration 1M, and the solution, after clarification by centrifugation, was loaded at 5 ml/min onto a column (Pharmacia XK 26/20 #18-1000-72) packed with 100 ml of the sorbent Phenyl-Toyopearl 650 S (20-50 μ m) (Supelco, #8-14477; 100 g), previously equilibrated with 3-4 column volumes of Buffer B (50 mM Tris.HCl, pH 7.4, 0.5 M guanidine-HCl and 1M (NH₄)₂SO₄). The column effluent was monitored at 280 nm. After loading, the column was washed with Buffer B until the A₂₈₀ of the effluent returned to near baseline level and then was eluted sequentially with 2-3 column volumes of Buffer C (50 mM Tris.HCl, 0.5M guanidine.HCl, 0.6M (NH₄)₂SO₄) with which the Hu-IFN- α 001 was eluted. Peak fractions showing maximum bands of Hu-IFN- α 001 on SDS-polyacrylamide gel electrophoresis were pooled. The Phenyl-Toyopearl column was regenerated in situ with 100 ml 0.5M NaOH and 1M NaCl solution; and was stored in 0.01% sodium azide. Fractions with Hu-IFN-001 as measured by antiviral activity and/or gel electrophoresis were pooled and concentrated 10-fold with an Amicon Centriprep 10 concentrator. The solution was then diluted 3-fold with

Buffer D (20 mM Tris.HCl, pH 8.0, 5% glycerol) and was loaded onto a FPLC monoQ HR 10/10 ion exchange column (Pharmacia #17-0556-01) equilibrated with Buffer D. The column was washed with about 10 ml of Buffer D until the A₂₈₀ reached baseline. Elution of Hu-IFN- α 001 was accomplished with a linear gradient of Buffer D and Buffer E (Buffer D plus 1M NaCl) at a flow rate of 1.5 ml/min from 0 to 100% Buffer E over 3 hours. The Hu-IFN- α 001 was eluted at 0.15M NaCl in a single peak. The fractions were pooled, analyzed by sodium dodecylsulfate (SDS) polyacrylamide gel electrophoresis and assayed for antiviral activity. From 6 g of bacterial pellet (wet weight), about 8–10 mg of purified Hu-IFN- α 001 was obtained.

The purified protein was mixed with 15 μ l of SDS sample buffer (0.5M Tris.HCl, pH 6.8, 1% (v/v) β -mercaptoethanol, 1% (w/v) sodium dodecylsulfate (SDS), 12% (v/v) glycerol, 2 mM ethylenediaminetetraacetic acid (EDTA), bromphenol blue) in a total volume of 35 μ l. The solution was boiled for two minutes after which 25 μ l was loaded onto a 12.5% polyacrylamide gel with a 4% polyacrylamide stacking gel. The separating gel was buffered in 0.3M Tris.HCl, 0.08% SDS, 2 mM EDTA, pH 8.8. The stacking gel was in 0.065M Tris.HCl, pH 6.8, and 0.05% SDS. The chamber buffer was 25 mM Tris.HCl, 0.1% SDS, 0.2M glycine. Electrophoresis was carried out for 1 hour at 150 V, 20 mA in the BioRad miniprotein II apparatus (132). The gel was stained with Coomassie Blue R-250 (2.4%, w/v, Coomassie Blue in 45% methanol, 9%, v/v, acetic acid) for 1 hour at room temperature; and destained in 8% acetic acid. From SDS-polyacrylamide gel electrophoresis it was apparent that the purified Hu-IFN- α 001 migrated with a M_r of 20,000 as shown in FIG. 4. As indicated in that figure, Hu-IFN- α 001 was placed in lanes 1, 2 and 3 in amounts of 3 μ g, 1.5 μ g and 0.75 μ g, respectively. The columns labeled M represent the molecular weight markers with the values in kilodaltons given to the left of each respective molecular weight marker. As can be seen, the Hu-IFN- α 001 exhibited a slightly slower mobility than Hu-IFN- α J on SDS-polyacrylamide gel electrophoresis (SDS PAGE, ref. 132).

Antiviral activity of Hu-IFN- α 001 was assayed on bovine MDBK and human FS7 cells with vesicular stomatitis virus (VSV) (Table 5) as described previously (133). The antiviral units were determined with respect to the human IFN- α A international standard Gxa01-901-535. There was approximately equal antiviral activity on human and bovine cells (Table 5) as is seen with many Hu-IFN- α species (17,27, 30,100,103,134).

TABLE 5

Antiviral Assay of Interferon			
Sample	Interferon Titer (units/ml)		Ratio (FS-7/MDBK)
	FS-7 Cells	MDBK Cells	
α 001	1×10^8	1×10^8	1.0

The interferon titer is given in units/mg as described (10–12, 99, 100, 133, 135) with respect to the international standard for human interferon alpha A Gxa01-901-535 from the National Institutes of Health. Vesicular stomatitis virus (VSV) was used as the challenge virus with human FS-7 and bovine MDBK cells. The ratio of the antiviral activity of the interferon on FS-7 to that on MDBK cells is given in the last column. The samples of Hu-IFN- α 001 were prepared as described in the text. Protein was determined by the method of Bradford (136).

Herein has been described an entire new class of molecules designated as super proteins, proteins not present in normal cells, but present in the cells in various diseased states and a method for identifying, producing and express-

ing such molecules. Although the present embodiment of the invention has been described in detail, it should be understood that various changes, alterations and substitutions can be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

BIBLIOGRAPHY

1. Isaacs, A., and Lindenmann, J., Production of Viral Interfering Substances, U.S. Pat. No. 3,699,222 (application Ser. No. 757,188) filed Aug. 5, 1968, issued Oct. 17, 1972.
2. Isaacs, A., and Lindenmann, J. (1957) "Virus Interference. I. The interferon," *Proc. Royal Society, London Ser. B* 147, 258–267.
3. Nagano, Y., and Kojima, Y. (1958) "Inhibition de l'infection vaccinale par un facteur liquide dans le tissu infecté par le virus homologue," *C. R. Seances Soc. Biol. Ses Fil.* 152, 1627–1629.
4. Wheelock, E. F. (1965) "Interferon-like virus-inhibitor induced human leukocytes by phytohemagglutinin," *Science* 149, 310–311.
5. Wheelock, E. F. (1966) "Virus replication and high-titered interferon production in human leukocyte cultures inoculated with Newcastle disease virus," *J. Bact.* 92, 1415–1421.
6. Youngner, J. S., and Stinebring, W. R. (1965) "Interferon appearance by endotoxin, bacteria, or viruses in mice pretreated with *Escherichia coli*, endotoxin, or infected with *Mycobacterium tuberculosis*," *Nature* 208, 456–458.
7. Youngner, J. S., and Salvin, S. B. (1973) "Production and properties of migration inhibitory factor and interferon in the circulation of mice with delayed hypersensitivity," *J. Immunol.* 111, 1914–1922.
8. Cantell, K., and Tovell, D. R. (1971) "Substitution of milk for serum in the production of human leukocyte interferon," *Appl. Microbiol.* 22, 625–628.
9. Stewart, W. E., II (1979) "The Interferon System." Springer-Verlag, New York, pp. 421.
10. Pestka, S. (1981) "Interferons, Part A," *Methods in Enzymology*, (S. Pestka, ed.) Academic Press, New York, Vol. 78, pp 632.
11. Pestka, S. (1981) "Interferons, Part B," *Methods in Enzymology*, (S. Pestka, ed.) Academic Press, New York, Vol. 79, pp 677.
12. Pestka, S. (1986) "Interferons, Part C," *Methods in Enzymology*, (S. Pestka, ed.) Academic Press, New York, Vol. 119, pp 845.
13. Baron, S., and Dianzani, F. (1977) "The Interferon System: A Current Review to 1978," *Texas Reports on Biology and Medicine* Vol. 35, University of Texas Medical Branch, Galveston, pp. 573.
14. Baron, S., Dianzani, F., and Stanton, G. J. (1982) "The Interferon System: A Review to 1982—Part I," *Texas Reports on Biology and Medicine* Vol. 41, University of Texas Medical Branch, Galveston, pp. 410.
15. Baron, S., Dianzani, F., and Stanton, G. J. (1982) "The Interferon System: A Review to 1982—Part II," *Texas Reports on Biology and Medicine* Vol. 41, University of Texas Medical Branch, Galveston, pp. 411–715.
16. Pestka, S., McInnes, J., Havell, E., and Vilcek, J. (1975) "Cell-free Synthesis of Human Interferon," *Proc. Natl. Acad. Sci. U.S.A.* 72, 3898–3901.
17. Pestka, S. (1983) "The Human Interferons—From Protein Purification and Sequence to Cloning and Expression in Bacteria: Before, Between, and Beyond," *Arch. Biochem. Biophys.* 221, 1–37.
18. Bose, S., Gurari-Rotman, D., Th. Ruegg, U., Corley, K., and C. B. Anfinsen (1976) "Apparent dispensability of the

- carbohydrate moiety of human interferon for antiviral activity," *J. Biol. Chem.* 251, 1659-1662.
19. Bose, S., and Hickman, J. (1977) "Role of the carbohydrate moiety in determining the survival of interferon in the circulation," *J. Biol. Chem.* 252, 8336-8337.
 20. Bridgen, P. J., Anfinsen, C. B., Corley, L., Bose, S., Zoon, K. C., Th. Ruegg, U., and Buckler, C. E. (1977) "Human lymphoblastoid interferon. Large scale production and partial purification," *J. Biol. Chem.* 252, 6585-6587.
 21. Rubinstein, M., Rubinstein, S., Familletti, P. C., Gross, M. S., Miller, R. S., Waldman, A. A., and Pestka, S. (1978) "Human Leukocyte Interferon Purified to Homogeneity," *Science* 202, 1289-1290.
 22. Rubinstein, M., Rubinstein, S., Familletti, P. C., Miller, R. S., Waldman, A. A., and Pestka, S. (1979) "Human Leukocyte Interferon: Production, Purification to Homogeneity, and Initial Characterization," *Proc. Natl. Acad. Sci. U.S.A.* 76, 640-644.
 23. Rubinstein, M., Rubinstein, S., Familletti, P. C., Brink, L. D., Hershberg, R. D., Gutterman, J., Hester, J., and Pestka, S. (1979) "Human Leukocyte Interferon Production and Purification to Homogeneity by HPLC," in *Peptides: Structure and Biological Function* (E. Gross and J. Meienhofer, eds.), Pierce Chemical Company, Rockford, Ill., 99-103.
 24. Rubinstein, M., Rubinstein, S., Familletti, P., Gutterman, J., Hester, J., and Pestka, S. (1980) "An Alternative Source of Human Leukocyte Interferon," in *Interferon: Properties and Clinical Uses* (A. Khan, N. O. Hill, and G. L. Dorn, eds.), Leland Fikes Foundation Press, Dallas, Tex., 45-55.
 25. Zoon, K. C., Smith, M. E., Bridgen, P. J., zur Nedden, D., and Anfinsen, C. B. (1979) "Purification and partial characterization of human lymphoblastoid interferon," *Proc. Natl. Acad. Sci. U.S.A.* 76, 5601-5605.
 26. Allen, G., and Fantes, F. H. (1980) "A family of structural genes for human lymphoblastoid (leukocyte-type) interferon," *Nature* (London) 287, 408-411.
 27. Rubinstein, M., Levy, W. P., Moschera, J. A., Lai, C.-Y., Hershberg, R. D., Bartlett, R. T., and Pestka, S. (1981) "Human Leukocyte Interferon: Isolation and Characterization of Several Molecular Forms," *Arch. Biochem. Biophys.* 210, 307-318.
 28. Zoon, K. C. (1981) "Purification and Characterization of Human Interferon from Lymphoblastoid (Namalwa) Cultures," *Methods in Enzymology* 78, 457-464.
 29. Berg, K., and Heron, I. (1981) "Antibody Affinity Chromatography of Human Leukocyte Interferon," *Methods in Enzymology* 78, 487-499.
 30. Hobbs, D. S., and Pestka, S. (1982) "Purification and Characterization of Interferons from a Continuous Myeloblastic Cell Line," *J. Biol. Chem.* 257, 4071-4076.
 31. Berg, K. (1982) "Purification and characterization of murine and human interferons. A review of the literature of the 1970s," *Acta Path. Microbiol. Immunol. Scand., Section C, Suppl.* 279, 1-136.
 32. Pestka, S. (1989) "The Interferons," p.433-480, in *Journal of Chromatography Library—Vol. 43, "Natural Products Isolation"* (Edited by G. H. Wagman and R. Cooper) Elsevier, N.Y., pp 619.
 33. Knight, E., Jr. (1976) "Interferon: purification and initial characterization from human diploid cells," *Proc. Natl. Acad. Sci. U.S.A.* 73, 520-523.
 34. Berthold, W., Tan, C., and Tan, Y. H. (1978) "Purification and in vitro labeling of interferon from a human fibroblastoid cell line," *J. Biol. Chem.* 253, 5206-5212.

35. Iwakura, Y., Yonehara, S., and Kawade, Y. (1978) "Purification of mouse L cell interferon. Essentially pure preparations with associated cell growth inhibitory activity," *J. Biol. Chem.* 253, 5074-5079.
36. Kawakita, M., Cabrer, B., Taira, H., Rebello, M., Slattery, E., Weideli, H., and Lengyel, P. (1978) "Purification of interferon from mouse Ehrlich ascites tumor cells," *J. Biol. Chem.* 253, 598-602.
37. De Maeyer-Guignard, J., Tovey, M. G., Gresser, I., and De Maeyer, E. (1978) "Purification of mouse interferon by sequential affinity chromatography on poly(U)—and antibody—agarose columns," *Nature*, London 271, 622-625.
38. Cabrer, B., Taira, H., Broeze, R. J., Kempe, T. D., Williams, K., Slattery, W. H., Konigsberg, W. H., and Lengyel, P. (1979) "Structural characteristics of interferons from mouse Ehrlich ascites tumor cells," *J. Biol. Chem.* 254, 3681-3684.
39. Stein, S., Kenny, C., Friesen, H.-J., Shively, J., Del Valle, U., and Pestka, S. (1980) "NH₂-Terminal Amino Acid Sequence of Human Fibroblast Interferon," *Proc. Natl. Acad. Sci. U.S.A.* 77, 5716-5719.
40. Friesen, H.-J., Stein, S., Evinger, M., Familletti, P. C., Moschera, J., Meienhofer, J., Shively, J., and Pestka, S. (1981) "Purification and Molecular Characterization of Human Fibroblast Interferon," *Arch. Biochem. Biophys.* 206, 432-450.
41. De Maeyer-Guignard, J. (1981) "Purification of Mouse C-243 Cell Interferon by Affinity Chromatography and Polyacrylamide Gel Electrophoresis," *Methods in Enzymology* 78, 513-522.
42. Okamura, H., Berthold, W., Hood, L., Hunakpiller, M., Inoue, M., Smith-Johannsen, H., and Tan, Y. H. (1981) "Human fibroblastoid interferon: immunosorbent column chromatography and N-terminal amino acid sequence," *Biochemistry* 19, 3831-3835.
43. Kawade, Y., Fujisawa, J., Yonehara, S., Iwakura, Y., and Yamamoto, Y. (1981) "Purification of L Cell Interferon," *Methods in Enzymology* 78, 522-535.
44. Knight, E., Jr., and Fahey, D. (1982) "Human Interferon-Beta: Effects of Deglycosylation," *J. Interferon Res.* 2, 421-429.
45. Goeddel, D. V., Shepard, H. M., Yelverton, E., Leung, D., Crea, R., Sloma, A., and Pestka, S. (1980) "Synthesis of Human Fibroblast Interferon by *E. coli*," *Nucleic Acids Res.* 8, 4057-4074.
46. Taniguchi, T., Ohno, S., Fujii-Kuriyama, Y., and Muramatsu, M. (1980) "The nucleotide sequence of human fibroblast interferon cDNA," *Gene*, 10, 11-15.
47. Derynck, R., Content, J., De Clercq, E., Volckaert, G., Tavernier, J., Devos, R., and Fiers, W. (1980) "Isolation and structure of a human fibroblast interferon gene," *Nature* (London) 285, 542-547.
48. Houghton, M., Steward, A. G., Doel, S. W., Emtage, J. S., Eaton, M. A. W., Smith, J. S., Patel, T. P., Lewis, H. M., Porter, A. G., Birch, J. R., Cartwright, T., and Carey, N. H. (1980) "The amino-terminal sequence of human fibroblast interferon as deduced from reverse transcripts obtained using synthetic oligonucleotide primers," *Nucleic Acids Res.* 8, 1913-1931.
49. Friesen, H.-J., and Pestka, S., Preparation of Homogeneous Human Fibroblast Interferon, U.S. Pat. No. 4,289,689 (application Ser. No. 160,889) filed Jun. 19, 1980, issued Sep. 15, 1981.
50. Pestka, S., and Rubinstein, M., Protein Purification Process and Product, U.S. Pat. No. 4,289,690 (application Ser. No. 167,165) filed Jul. 9, 1980, issued Sep. 15, 1981.

51. Pestka, S., and Rubinstein, M., Protein Purification Process and Product, U.S. Pat. No. 4,503,035 (application Ser. No.465,979) filed Feb. 14, 1983, issued Mar. 5, 1985.
52. Kung, H.-F., Miller, D. L., and Pestka, S., Crystalline Human Leukocyte Interferon, U.S. Pat. No. 4,672,108 (application Ser. No. 751,753) filed Jul. 3, 1985, issued Jun. 9, 1987.
53. Goeddel, D. V., and Pestka, S., Microbial Production of Mature Human Leukocyte Interferon K and L, U.S. Pat. No. 4,801,685 (application Ser. No.56,623), filed Jun. 1, 1987, issued Jan. 31, 1989; European Patent Application 82107337.6.
54. Goeddel, D. V., and Pestka, S., Microbial Production of Mature Human Leukocyte Interferon K and L, U.S. Pat. No. 4,810,645 (application Ser. No. 822,984) filed Jan. 27, 1986, issued Mar. 7, 1989.
55. Friesen, H.-J., and Pestka, S., Preparation of Homogeneous Human Fibroblast Interferon, U.S. Pat. No. 5,015,730 (application Ser. No. 386,088), filed Jul. 14, 1989, issued May 14, 1991.
56. Baron, S., Copenhagen, D. H., Dianzani, F., Fleischmann, W. R., Jr., Hughes, T. K., Jr., Klimpel, G. R., Niesel, D. W., Stanton, G. J., and Tying, S. K., editors (1992) "Interferon: Principles and Medical Applications," The University of Texas Medical Branch at Galveston, Galveston, pp. 624.
57. Pestka, S., Langer, J. A., Zoon, K. C., and Samuel, C. E. (1987) "Interferons and Their Actions," *Annu. Rev. Biochem.* 56, 727-777.
58. Havell, E. A., Berman, B., Ogburn, C. A., Berg, K., Paucker, K., and Vilcek, J. (1975) "Two Antigenically Distinct Species of Human Interferon," *Proc. Natl. Acad. Sci. USA* 72, 2185-2187.
59. Cavalieri, R. L., Havell, E. A., Vilcek, J., and Pestka, S. (1977) "Synthesis of Human Interferon by *Xenopus laevis* Oocytes: Two Structural Genes for Interferons in Human Cells," *Proc. Natl. Acad. Sci. U.S.A.* 74, 3287-3291.
60. Familletti, P. C., McCandliss, R., and Pestka, S. (1981) "Production of High Levels of Human Leukocyte Interferon from a Continuous Human Myeloblastoid Cell Culture," *Antimicrob. Agents Chemother.* 20, 5-9.
61. Pestka, S. (1986) "Interferon from 1981 to 1986," *Methods in Enzymology* (S. Pestka, ed.), Academic Press, New York 119, 3-14.
62. Maeda, S., McCandliss, R., Gross, M., Sloma, P. C., Familletti, J. M., Tabor, M., Evinger, M., Levy, W. P., and Pestka, S. (1981) "Construction and identification of bacterial plasmids containing nucleotide sequence for human leukocyte interferon," *Proc. Natl. Acad. Sci. USA* 77, 7010-7013 (1980); 78, 4648.
63. Goeddel, D. V., and Pestka, S. (1982) Polypeptides, process for their microbial production, intermediates therefor and compositions containing them [Microbial Production of Mature Human Leukocyte Interferons] European Patent Application 81105067.3
64. Henco, K., Brosius, J., Fujisawa, A., Fujisawa, J.-I., Haynes, J. R., Hochstadt, J., Kovacic, T., Pasek, M., Schamboek, A., Schmid, J., Todokoro, K., Wälichli, M., Nagata, S., and Weissmann, C. (1985) "Structural Relationship of Human Interferon Alpha Genes and Pseudogenes," *J. Mol. Biol.* 185, 227-260.
65. Diaz, M. O., Pomykala, H., Bohlander, S., Maltepe, E., Olopade, O. (1991) "A Complete Physical Map of the Type-I Interferon Gene Cluster," *J. Interferon Res.*, 11, S85.
66. Owerback, D., Rutter, W. J., Shows, T. B., Gray, P., Goeddel, D. V., and Lawn, R. M. (1981) "Leukocyte and Fibroblast Interferon Genes are Located on Chromosome 9," *Proc. Natl. Acad. Sci. USA* 78, 3123-3127.
67. Trent, J. M., Olson, S., and Lawn, R. M. (1982) "Chromosomal Localization of Human Leukocyte, Fibro-

- blast and Immune Interferon Genes by Means of in situ Hybridization," *Proc. Natl. Acad. Sci. USA* 79, 7809-7813.
68. Langer, J. A., and Pestka, S. (1984) "Purification, Bacterial Expression and Biological Activities of the Human Interferons," *J. Invest. Dermatol.* 83, 128-136.
69. Goeddel, D. V., Yelverton, E., Ullrich, A., Heyneker, H. L., Miozzari, G., Holmes, W., Seeburg, P. H., Dull, T., May, L., Stebbing, N., Crea, R., Maeda, S., McCandliss, A., Sloma, J. M., Tabor, J. M., Gross, M., Familletti, P. C., and Pestka, S. (1981) "Human leukocyte interferon produced in *E. coli* is biologically active," *Nature* 287, 411-416.
70. Streuli, M., Nagata, S., and Weissmann, C. (1980) "At Least Three Human Type α Interferons: Structure of $\alpha 2$," *Science* 209, 1343-1347.
71. Lawn, R. M., Gross, M., Houk, C. M., Franke, A. E., Gray, P. V., and Goeddel, D. V. (1981) "DNA Sequence of a Major Human Leukocyte Interferon Gene," *Proc. Natl. Acad. Sci. USA* 78, 5435-5439.
72. Dworkin-Rastl, E., Dworkin, M. B., and Swetly, P. (1982) "Molecular cloning of human alpha and beta interferon genes from Namalwa cells," *J. Interferon Research* 2, 575-585.
73. Physicians' Desk Reference, PDR, 47th Edition, 1993: pages 1078-1079; 1879-1881; 2006-2008; 2194-2201.
74. Dianzani, F. (1992) Biological Basis for Therapy and for Side Effects in "Interferon: Principles and Medical Applications," (Baron, S., Copenhagen, D. H., Dianzani, F., Fleischmann, W. R., Jr., Hughes, T. K., Jr., Klimpel, G. R., Niesel, D. W., Stanton, G. J., and Tying, S. K., editors) The University of Texas Medical Branch at Galveston, Galveston, 409-416.
75. Hosoi, H., Miyaki, K., and Yamanaka, M. (1992) "The Interferon $\alpha 2$ Gene in Japanese Patients with Chronic Viral Hepatitis Who Developed Antibodies After Recombinant Interferon $\alpha 2A$ Treatment," *International Society for Study of the Liver*, Brighton, UK, 3-6th Jun., 1992; p 113, Abstract.
76. Desai, M., Hussain, M., Lee, N., Ni, D., Liao, M.-J., and Testa, D. (1992) "Identification of IFN- $\alpha 2$ Transcripts in Sendai Virus Induced Human Leukocytes by PCR," *J. Interferon Res.* 12, S138.
77. Adolf, G. R., Kalsner, L., Ahorn, H., Maurer-Fogy, I., and Cantell, K. (1991) "Natural Human Interferon Alpha-2 is O-glycosylated," *Biochem. J.* 276, 511-518.
78. Zoon, K. C., Miller, D., Bekisz, J., zur Nedden, D., Ny, J. C., Nguyen, N. Y., and Hu, R. (1992) "Purification and Characterization of Multiple Components of Human Lymphoblastoid Interferon- α ," *J. Biol. Chem.* 267, 15210-15216.
79. Emanuel, S. L., and Pestka, S. (1993) "Human interferon- αA , - $\alpha 2$ and - $\alpha 2(Arg)$ Genes in Genomic DNA," *J. Biol. Chem.* 268, 12565-12569.
80. Koeffler, H. P., and Golde, D. W. (1978) "Acute Myelogenous Leukemia: A Human Cell Line Responsive to Colony-Stimulating Activity," *Science* 200, 1153-1154.
81. Nadkarni, J. S., Nadkarni, J. J., Clifford, P., Manolov, G., Fenyo, E. M., and Klein, E. (1969) "Characteristics of new cell lines derived from Burkitts lymphomas," *Cancer* 23, 64-79.
82. Beaucage, S. L., and Carothers, M. H. (1981) "Deoxynucleoside Phosphoramidites-A New Class of Key Intermediates for Deoxypolynucleotide Synthesis," *Tetrahedron Lett.* 22, 1859-1862.
83. Chow, R., Kempe, T., and Palm, G. (1981) "Synthesis of Oligodeoxyribonucleotides on Silica Gel Support," *Nucleic Acids Res.* 9, 2807-2817.
84. Johnson, B. A., Mc Clain, S. G., and Doran, E. R. (1990) "Rapid Purification of Synthetic Oligonucleotides: A Convenient Alternative to High-Performance Liquid

- Chromatography and Polyacrylamide Gel Electrophoresis," *Biotechniques* 8, 424-429.
85. Altling-Mees, M. A., and Short, J. M. (1989) "pBluescript II: gene mapping vectors," *Nucleic Acids Res.* 17, 9494.
 86. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*. Three Volume Set, Cold Spring Harbor Laboratory, New York.
 87. Birnboim, H. C., and Doly, J. (1979) "A Rapid Alkaline Extraction Procedure for Screening Recombinant Plasmid DNA," *Nucleic Acids Res.* 7, 1513-1523.
 88. Sanger, F., Nicklen, S., and Coulson, A. R. (1977) "DNA Sequencing with Chain Terminating Inhibitors," *Proc. Natl. Acad. Sci. USA* 74, 5463-5467.
 89. Tabor, S., and Richardson, C. C. (1987) "DNA Sequence Analysis with a Modified Bacteriophage T7 DNA Polymerase," *Proc. Natl. Acad. Sci. USA* 84, 4767-4771.
 90. Kawasaki, E. S., and Wang, A. M. (1989) "Detection of Gene Expression," In: *PCR Technology: Principles and Applications of DNA Amplification*. (Erlich, H. A. ed.) Stockton Press, Inc., New York, N.Y., pp 89-97.
 91. McCandliss, R., Sloma, A., and Pestka, S. (1981) "Isolation and Cell-free Translation of Human Interferon mRNA from Fibroblasts and Leukocytes," in *Methods in Enzymology*, Vol. 79 (S. Pestka, ed.), Academic Press, New York, 51-59.
 92. Chomczynski, P., and Sacchi, N. (1987) "Single Step Method of RNA Isolation by Acid Guanidinium Thiocyanate-Phenol-Chloroform Extraction," *Anal. Biochem.* 162, 156-159.
 93. Pellicer, A., Wigler, M., and Axel, R. (1978) "The Transfer and Stable Integration of the HSV Thymidine Kinase Gene into Mouse Cells," *Cell* 14, 133-14.
 94. Gross-Bellard, M., Oudet, P., and Chambon, P. (1973) "Isolation of High-Molecular-Weight DNA from Mammalian Cells," *Eur. J. Biochem.* 36, 32-38.
 95. Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A., and Arnheim, N. (1985) "Enzymatic Amplification of Beta-Globin Genomic Sequences and Restriction Site Analysis for Diagnosis of Sickle Cell Anemia," *Science* 230, 1350-1354.
 96. Keohavong, P., and Thilly, W. G. (1989) "Fidelity of DNA Polymerases in DNA Amplification," *Proc. Natl. Acad. Sci. USA* 86, 9253-9257.
 97. Hotta, K., Collier, K. J., and Pestka, S. (1986) "Detection of a Single Base Substitution Between Human Leukocyte Interferon α A and α 2 Genes with Octadecyl Deoxyoligonucleotide Probes," in *Methods in Enzymology* (S. Pestka, ed.), Academic Press, New York 119, 481-485.
 98. Hotta, K., Monahan, J., Collier, K. J., and Pestka, S. (1988) "Detection of Human Leukocyte Interferon α A and α 2 Genes in Genomic DNAs by the Use of Deoxyoctadecyloligonucleotide Probes," *J. Interferon Res.* 8, 51-60.
 99. Pestka, S., and Baron, S. (1981) "Definition and Classification of the Interferons," in *Methods in Enzymology* (S. Pestka, ed.), Academic Press, New York, 78, 3-14.
 100. Pestka, S. (1986) "Interferon Standards and General Abbreviations," in *Methods in Enzymology* (S. Pestka, ed.), Academic Press, New York, 119, 14-23.
 101. Evinger, M., Maeda, S., and Pestka, S. (1981) "Recombinant Human Leukocyte Interferon Produced in Bacteria Has Antiproliferative Activity," *J. Biol. Chem.* 256, 2113-2114.
 102. Herberman, R. B., Ortaldo, J. R., Mantovani, A., Hobbs, D. S., Kung, H.-F., and Pestka, S. (1982) "Effect of Human Recombinant Interferon on Cytotoxic Activity of Natural Killer (NK) Cells and Monocytes," *Cell Immunol.* 67, 160-167.
 103. Rehberg, E., Kelder, B., Hoal, E. G., and Pestka, S. (1982) "Specific Molecular Activities of Recombinant and Hybrid Leukocyte Interferons," *J. Biol. Chem.* 257, 11497-11502.

104. Jones, C. M., Varesio, L., Herberman, R. B., and Pestka, S. (1982) "Interferon Activates Macrophages to Produce Plasminogen Activator," *J. Interferon Res.* 2, 377-386.
105. Grant, S., Bhalla, K., Weinstein, I. B., Pestka, S., and Fisher, P. B. (1982) "Differential Effect of Recombinant Human Leukocyte Interferon on Human Leukemic and Normal Myeloid Progenitor Cells," *Biochem. Biophys. Res. Commun.* 108, 1048-1055.
106. Ortaldo, J. R., Herberman, R. B., and Pestka, S. (1982) "Augmentation of Human Natural Killer Cells with Human Leukocyte and Human Recombinant Leukocyte Interferon," in *NK Cells and Other Natural Effector Cells* (R. B. Herberman, ed.), Academic Press, New York, 1279-1283.
107. Ortaldo, J. R., Mason, A., Rehberg, E., Moschera, J., Kelder, B., Pestka, S., and Herberman, R. B. (1983) "Effects of Recombinant and Hybrid Recombinant Human Leukocyte Interferons on Cytotoxic Activity of Natural Killer Cells," *J. Biol. Chem.* 258, 15011-15015.
108. Fisher, P., Miranda, A. F., Babiss, L. E., Pestka, S., and Weinstein, I. B. (1983) "Opposing Effects of Interferon Produced in Bacteria and of Tumor Promoters on Myogenesis in Human Myoblast Cultures," *Proc. Natl. Acad. Sci. U.S.A.* 80, 2961-2965.
109. Sen, G. C., Herz, R. E., Davatellis, V., and Pestka, S. (1984) "Antiviral and Protein-Inducing Activities of Recombinant Human Leukocyte Interferons and Their Hybrids," *J. Virol.* 50, 445-450.
110. Giacomini, P., Aguzzi, A., Pestka, S., Fisher, P. B., and Ferrone, S. (1984) "Modulation by Recombinant DNA Leukocyte (α) and Fibroblast (β) Interferons of the Expression and Shedding of HLA and Tumor Associated Antigens by Melanoma Cells," *J. Immunol.* 133, 1649-1655.
111. Greiner, J. W., Hand, P. H., Noguchi, P., Fisher, P. B., Pestka, S., and Schlom, J. (1984) "Enhanced Expression of Surface Tumor-Associated Antigens on Human Breast and Colon Tumor Cells After Recombinant Human Leukocyte α -Interferon Treatment," *Cancer Res.* 44, 3208-3214.
112. Fisher, P. B., Prignoli, D. R., Hermon, H., Jr., Weinstein, I. B., and Pestka, S. (1985) "Effects of Combined Treatment with Interferon and Mezerin on Melanogenesis and Growth in Human Melanoma Cells," *J. Interferon Res.* 5, 11-22.
113. Grant, S., Bhalla, K., Weinstein, I. B., Pestka, S., Mileno, M. D., and Fisher, P. B. (1985) "Recombinant Human Interferon Sensitizes Resistant Myeloid Leukemic Cells to Induction of Terminal Differentiation," *Biochem. Biophys. Res. Commun.* 130, 379-388.
114. Guadagni, F., Schlom, J., Johnston, W. W., Szpak, C. Z., Goldstein, D., Smalley, R., Simpson, J. F., Borden, E. C., Pestka, S., and Greiner, J. W., (1989) "Selective Interferon-Induced Enhancement of Tumor-Associated Antigens on a Spectrum of Freshly Isolated Human Adenocarcinoma Cells," *J. Natl. Cancer Inst.* 81, 502-512.
115. Sperber, S. J., Gocke, D. J., Habertzettl, C., Kuk, R., Schwartz, B., and Pestka, S. (1992) "Anti-HIV-1 Activity of Recombinant and Hybrid Species of Interferon Alpha," *J. Interferon Res.* 12, 363-368.
116. Huber, C., Flener, R., and Gastl, G. (1985) "Interferon-Alpha-2c in the Treatment of Advanced Hairy Cell Leukemia," *Oncology* 42, Suppl 1, 7-9.
117. Foon, K. A., Maluish, A. E., Abrams, P. G., Wrightington, S., Stevenson, H. C., Alarif, A., Fer, M. F., Overton, W. R., and Poole, M. (1986) "Recombinant Leukocyte A Interferon Therapy for Advanced Hairy Cell Leukemia," *Am. J. Med.* 80, 351-356.
118. Steis, R. G., Smith, J. W. II, Urba, W. J., Clark, J. W., Itri, L. M., Evans, L. M., Schoenberger, C., and Longo, D.

- L. (1988) "Resistance to Recombinant Interferon Alfa-2 in Hairy Cell Leukemia Associated with Neutralizing Anti-interferon Antibodies," *New Engl. J. Med.* 318, 1409-1413.
119. Von Wussow, P., Freund, M., Hartmann, F., Diedrich, H., Poliwoda, H., and Deicher, H. (1987) "Anti Interferon Antibodies: Pharmacokinetics and Clinical Significance," *J. Interferon Res.* 7, 680.
120. Freund, M., Von Wussow, P., Diedrich, H., Eisert, R., Link, H., Wilke, H., Buchholz, F., LeBlanc, S., Fonatsch, C., Deicher, H., and Poliwoda, H. (1989) "Recombinant Human Interferon (IFN) Alpha-2b in Chronic Myelogenous Leukemia: Dose Dependency of Response and Frequency of Neutralizing Anti-Interferon Antibodies," *Br. J. Haematol.* 72, 350-356.
121. Itri, L. M., Campion, M., Dennin, R. A., Palleroni, A. V., Gutterman, J. O., Groopman, J. E., and Trown, P. W. (1987) "Incidence and Clinical Significance of Neutralizing Antibodies in Patients Receiving Recombinant Interferon Alpha-1A by Intramuscular Injection," *Cancer* 59, 668-674.
122. Moormeier, J. A., Westbrook, C. A., Ratain, M. J., and Golomb, H. M. (1989) "Interferon Alfa-2b Antibodies and Clinical Resistance in a Patient with Hairy Cell Leukemia," *Leuk. Lymphoma* 1, 43-45.
123. Quesada, J. R., Rios, A., Swanson, D. A., Trown, P., and Gutterman, J. U. (1985) "Antitumor Activity of Recombinant-derived Interferon Alpha in Metastatic Renal Cell Carcinoma," *J. Clin. Oncol.* 3, 1522-1528.
124. Antonelli, G., Currenti, M., Turriziani, O., and Dianzani, F. (1991) "Neutralizing Antibodies to Interferon α : Relative Frequency in Patients Treated with Different Interferon Preparations," *J. Infect. Dis.* 163, 882-885.
125. Colamonici, O., Porterfield, B., and Diaz, M. O. (1991) "Interferon Sensitivity of Human Leukemia Cell Lines With and Without Deletion of the Interferon Genes," *J. Interferon Res.* 11, S54.
126. Grandé, D., Heyman, M., Brøndum-Nielsen, K., Liu, Y., Lundgren, E., Söderhäll, S., and Einhorn, S. (1992) "Interferon System in Primary Acute Lymphocytic Leukemia Cells With or Without Deletions of the α/β -Interferon Genes," *Blood* 79, 2076-2083.

127. Nishimura, A., Morita, M., Nishimura, Y. and Sugino, Y. (1990) "A Rapid and Highly Efficient Method for Preparation of Competent *Escherichia coli* Cells," *Nucleic Acids Research* 18, 6169.
128. Lee, S. and Rashid, S. (1990) "A Simple Procedure for Maximum Yield of High-quality Plasmid DNA," *Bio-Techniques* 9, 676-679.
129. Mashko, S. V., Veiko, V. P., Lapidus, A. L., Lebedeva, M. I., Mochulsky, A. V., Shechter, I. I., Trukhan, M. E., Ratmanova, K. I., Rebentish, B. A., Kaluzhsky, V. E. and Debabov, V. G. (1990) TGATG vector: a new expression system for cloned foreign genes in *Escherichia coli* cells. *Gene* 88, 121-126.
130. Caruthers, M. H., Barone, A. D., Beaucage, S. L., Dodds, D. R., Fisher, E. F., McBride, L. J., Matteucci, M., Stabinsky, Z., Tang, J. Y. (1987) "Chemical Synthesis of Deoxyoligonucleotides by the Phosphoramidite Method," *Methods in Enzymology* 154, 287-313.
131. Wang, P., Izotova, L., Mariano, T. M., Donnelly, R. J. and Pestka, S. (1994) "Construction and Activity of Phosphorylatable Human Interferon- α B2 and Interferon- α A/D," *J. Interferon Res.* 14, 41-46.
132. Laemmli, U. K. (1990) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* 227: 680-685.
133. Familletti, P. C., Rubinstein, S., and Pestka, S. (1981) "A Convenient and Rapid Cytopathic Effect Inhibition Assay for Interferon," in *Methods in Enzymology*, Vol. 78 (S. Pestka, ed.), Academic Press, New York, 387-394.
134. Staehelin, T., Hobbs, D. S., Kung, H.-F., Lai, C.-Y., and Pestka, S. (1981) "Purification and Characterization of Recombinant Human Leukocyte Interferon (IFLrA) with Monoclonal Antibodies," *J. Biol. Chem.* 256, 9750-9754.
135. Pestka, S. (1981) "Standard Media and General Abbreviations," in *Methods in Enzymology*, Vol. 78 (S. Pestka, ed.), Academic Press, New York, 22-25.
136. Bradford, M. M. (1976) "A Rapid and Sensitive Method for the quantitation of Microgram quantities of Protein Utilizing the Principle of Protein-dye Binding," *Anal. Biochem.* 72, 248-254.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i i i) NUMBER OF SEQUENCES: 12

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TGGGCTGTGA TCTGCCTC

1 8

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single

-continued

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 2:

CATGATTTCT GCTCTGACAA CC 2 2

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 3:

AACCCACAGC CTGGGTAG 1 8

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 4:

GCGGGCCCCA ATGGCCYTGY CCTTT 2 5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 5:

GCTCTAGAAY TCATGAAAGY GTGA 2 4

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 6:

CTTGAAGGAC AGACATG 1 7

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(i i) MOLECULE TYPE: Genomic DNA

(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 7:

CTGTCCTCCA TGAGATG 17

(2) INFORMATION FOR SEQ ID NO: 8:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(i i) MOLECULE TYPE: Genomic DNA
(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 8:

GGTCATTTCAG CTGCTGG 17

(2) INFORMATION FOR SEQ ID NO: 9:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(i i) MOLECULE TYPE: Genomic DNA
(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 9:

TCCTCCTTCA TCAGGGG 17

(2) INFORMATION FOR SEQ ID NO: 10:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(i i) MOLECULE TYPE: Genomic DNA
(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 10:

ATTAACCCTC ACTAAAG 17

(2) INFORMATION FOR SEQ ID NO: 11:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(i i) MOLECULE TYPE: Genomic DNA
(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 11:

TAATACGACT CACTATA 17

(2) INFORMATION FOR SEQ ID NO: 12:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 570 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(i i) MOLECULE TYPE: Genomic DNA
(x i) SEQUENCE DESCRIPTION: SEQ ID. NO: 12:

ATG GCC TTG 9
Met Ala Leu
- 23

-continued

TCC	TTT	TCT	TTA	CTG	ATG	GTC	GTG	CTG	GTA	CTC	AGC	TAC	AAA	TCC	54
Ser	Phe	Ser	Leu	Leu	Met	Val	Val	Leu	Val	Leu	Ser	Tyr	Lys	Ser	
				- 20					- 15					- 10	
ATC	TGC	TCT	CTG	GGC	TGT	GAT	CTG	CCT	CAG	ACC	CAC	AGC	CTG	CGT	99
Ile	Cys	Ser	Leu	Gly	Cys	Asp	Leu	Pro	Gln	Thr	His	Ser	Leu	Arg	
- 5					1				5					10	
AAT	AGG	AGG	GCC	TTG	ATA	CTC	CTG	GCA	CAA	ATG	GGA	AGA	ATC	TCT	144
Asn	Arg	Arg	Ala	Leu	Ile	Leu	Leu	Ala	Gln	Met	Gly	Arg	Ile	Ser	
				15					20					25	
CCT	TTC	TCC	TGC	TTG	AAG	GAC	AGA	CAT	GAA	TTC	AGA	TTC	CCA	GAG	189
Pro	Phe	Ser	Cys	Leu	Lys	Asp	Arg	His	Glu	Phe	Arg	Phe	Pro	Glu	
				30					35					40	
GAG	GAG	TTT	GAT	GGC	CAC	CAG	TTC	CAG	AAG	ACT	CAA	GCC	ATC	TCT	234
Glu	Glu	Phe	Asp	Gly	His	Gln	Phe	Gln	Lys	Thr	Gln	Ala	Ile	Ser	
				45					50					55	
GTC	CTC	CAT	GAG	ATG	ATC	CAG	CAG	ACC	TTC	AAT	CTC	TTC	AGC	ACA	279
Val	Leu	His	Glu	Met	Ile	Gln	Gln	Thr	Phe	Asn	Leu	Phe	Ser	Thr	
				60					65					70	
GAG	GAC	TCA	TCT	GCT	GCT	TGG	GAA	CAG	AGC	CTC	CTA	GAA	AAA	TTT	324
Glu	Asp	Ser	Ser	Ala	Ala	Trp	Glu	Gln	Ser	Leu	Leu	Glu	Lys	Phe	
				75					80					85	
TCC	ACT	GAA	CTT	TAC	CAG	CAA	CTG	AAT	GAC	CTG	GAA	GCA	TGT	GTG	369
Ser	Thr	Glu	Leu	Tyr	Gln	Gln	Leu	Asn	Asp	Leu	Glu	Ala	Cys	Val	
				90					95					100	
ATA	CAG	GAG	GTT	GGG	GTG	GAA	GAG	ACT	CCC	CTG	ATG	AAT	GAG	GAC	414
Ile	Gln	Glu	Val	Gly	Val	Glu	Glu	Thr	Pro	Leu	Met	Asn	Glu	Asp	
				105					110					115	
TCC	ATC	CTG	GCT	GTG	AGG	AAA	TAC	TTC	CAA	AGA	ATC	ACT	CTT	TAT	459
Ser	Ile	Leu	Ala	Val	Arg	Lys	Tyr	Phe	Gln	Arg	Ile	Thr	Leu	Tyr	
				120					125					130	
CTA	ACA	GAG	AAG	AAA	TAC	AGC	CCT	TGT	GCC	TGG	GAG	GTT	GTC	AGA	504
Leu	Thr	Glu	Lys	Lys	Tyr	Ser	Pro	Cys	Ala	Trp	Glu	Val	Val	Arg	
				135					140					145	
GCA	GAA	ATC	ATG	AGA	TCC	CTC	TCG	TTT	TCA	ACA	AAC	TTG	CAA	AAA	549
Ala	Glu	Ile	Met	Arg	Ser	Leu	Ser	Phe	Ser	Thr	Asn	Leu	Gln	Lys	
				150					155					160	
AGA	TTA	AGG	AGG	AAG	GAT	TGA									570
Arg	Leu	Arg	Arg	Lys	Asp	End									
				165											

I claim the following:

1. Isolated human leukocyte interferon wherein the interferon has the amino acid sequence from position 1 to position 166 as depicted for Hu-IFN- α 001 in FIG. 1 (SEQ ID NO:12).

2. A recombinantly produced polypeptide having interferon activity and comprising the amino acid sequence of Hu-IFN- α 001 (SEQ ID NO:12):

CysAspLeuProGlnThrHisSerLeuArgAsnArgArgAlaLeulleuLeuAlaGln
1 10 20
MetGlyArgIleSerProPheSerCysLeuLysAspArgHisGluPheArgPheProGlu
30 40
GluGluPheAspGlyHisGlnPheGlnLysThrGlnAlaIleSerValLeuHisGluMet
50 60
IleGlnGlnThrPheAsnLeuPheSerThrGluAspSerSerAlaAlaTrpGluGlnSer
70 80
LeuLeuGluLysPheSerThrGluLeuTyrGlnGlnLeuAsnAspLeuGluAlaCysVal
90 100
IleGlnGluValGlyValGluGluThrProLeuMetAsnGluAspSerIleLeuAlaVal
110 120
ArgLysTyrPheGlnArgIleThrLeuTyrLeuThrGluLysLysTyrSerProCysAla
130 140

45

-continued

TrpGluValValArgAlaGluIleMetArgSerLeuSerPheSerThrAsnLeuGlnLys
150 160
ArgLeuArgArgLysAspEnd
166

50

55

60

65

3. A pharmaceutical composition for providing interferon therapy to a human comprising an effective amount of the polypeptide of claim 2 admixed with a pharmaceutically acceptable vehicle or carrier.

4. The polypeptide of claim 2 wherein said polypeptide is derived from diseased cells.

5. The recombinantly produced polypeptide of claim 4 wherein said diseased cells are derived from hemopoietic cells.

6. The recombinantly produced polypeptide of claim 4 wherein said diseased cells are derived from leukemic leukocytes.

7. The recombinantly produced polypeptide of claim 4 wherein said diseased cells are derived from human malignancies.

* * * * *