



(86) Date de dépôt PCT/PCT Filing Date: 2007/01/12
(87) Date publication PCT/PCT Publication Date: 2007/07/19
(45) Date de délivrance/Issue Date: 2013/07/30
(85) Entrée phase nationale/National Entry: 2008/04/15
(86) N° demande PCT/PCT Application No.: US 2007/000994
(87) N° publication PCT/PCT Publication No.: 2007/082098
(30) Priorité/Priority: 2006/01/13 (US60/758,701)


(51) Cl.Int./Int.Cl. *C07D 213/79* (2006.01),
A01N 43/40 (2006.01), *C07C 25/13* (2006.01),
C07C 39/26 (2006.01), *C07C 39/27* (2006.01),
C07C 39/28 (2006.01), *C07C 47/55* (2006.01),
C07C 47/575 (2006.01), *C07C 49/80* (2006.01),
C07D 405/04 (2006.01), *C07F 5/02* (2006.01)

(72) Inventeurs/Inventors:
BALKO, TERRY WILLIAM, US;
SCHMITZER, PAUL RICHARD, US;
DAEUBLE, JOHN FILSON, US;
SIDDALL, THOMAS LYMAN, US;
YERKES, CARLA NANETTE, US

(73) Propriétaire/Owner:

(54) Titre : 6-(POLY-ARYL SUBSTITUTE)-4-AMINOPICOLINATES ET UTILISATIONS DE CEUX-CI COMME
HERBICIDES

(54) Title: 6-(POLY-SUBSTITUTED ARYL)-4-AMINOPICOLINATES AND THEIR USE AS HERBICIDES



(57) Abrégé/Abstract:

4-Aminopicolinic acids having tri- and tetra-substituted aryl substituents in the 6-position, and their amine and acid derivatives of the formula I are potent herbicides demonstrating a broad spectrum of weed control.



(73) Propriétaires(suite)/Owners(continued):DOW AGROSCIENCES LLC, US

(74) Agent: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2007 (19.07.2007)

PCT

(10) International Publication Number
WO 2007/082098 A3

(51) International Patent Classification:

C07D 213/79 (2006.01) *C07C 47/575* (2006.01)
C07C 39/27 (2006.01) *C07C 49/80* (2006.01)
C07C 39/28 (2006.01) *A01N 43/40* (2006.01)
C07F 5/02 (2006.01) *C07D 405/04* (2006.01)
C07C 25/13 (2006.01) *C07C 39/26* (2006.01)
C07C 47/55 (2006.01)

(74) Agent: MIXAN, Craig, E.; Dow AgroSciences LLC,
9330 Zionsville Road, Indianapolis, IN 46268 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2007/000994

(22) International Filing Date: 12 January 2007 (12.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/758,701 13 January 2006 (13.01.2006) US

(71) Applicant (*for all designated States except US*): DOW AGROSCIENCES LLC [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268-1054 (US).

(72) Inventors: BALKO, Terry, William; 1410 Bowman Drive, Greenfield, IN 46140 (US). SCHMITZER, Paul, Richard; 7960 North Whittier Place, Indianapolis, IN 46250 (US). DAEUBLE, John, Filson; 1829 Horseguard Close, Carmel, IN 46032 (US). YERKES, Carla, Nanette; 4571 South State Road 47, Crawfordsville, IN 47933 (US). SIDDALL, Thomas, Lyman; 1925 South 900 East, Zionsville, IN 46077 (US). YERKES, Carla, Nanette; 4571 South State Road 47, Crawfordsville, IN 47933 (US).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
8 November 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 6-(POLY-SUBSTITUTED ARYL)-4-AMINOPICOLINATES AND THEIR USE AS HERBICIDES



(57) Abstract: 4-Aminopicolinic acids having tri- and tetra-substituted aryl substituents in the 6-position, and their amine and acid derivatives of the formula I are potent herbicides demonstrating a broad spectrum of weed control.

WO 2007/082098 A3

73776-249

6-(POLY-SUBSTITUTED ARYL)-4-AMINOPICOLINATES AND THEIR USE
AS HERBICIDES


5

This invention relates to certain novel 6-(poly-substituted aryl)-4-aminopicolinate and their derivatives and to the use of these compounds as herbicides.

A number of picolinic acids and their pesticidal properties have been described in the art. U.S. Patent 6,784,137 B2 discloses a genus of 6-aryl-4-aminopicolinic acids and their derivatives and their use as herbicides. It has now been discovered that certain particular subclasses of the genus disclosed in '137 have greatly improved herbicidal activity and selectivity.

It has now been found that certain 6-(poly-substituted aryl)-4-aminopicolinic acids and their derivatives are superior herbicides with a broad spectrum of weed control against woody plants, grasses and sedges as well as broadleaves and with excellent crop selectivity. The compounds further possess excellent toxicological or environmental profiles.

The invention includes compounds of Formula I:



I

20

wherein


Q_1 represents H or F;

Q_2 represents a halogen with the proviso that when Q_1 is H then Q_2 is Cl or Br;

5 R_1 and R_2 independently represent H, C_1 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, hydroxy, C_1 - C_6 alkoxy, amino, C_1 - C_6 acyl, C_1 - C_6 carboalkoxy, C_1 - C_6 alkylcarbamyl, C_1 - C_6 alkylsulfonyl, C_1 - C_6 trialkylsilyl or C_1 - C_6 dialkyl phosphonyl or R_1 and R_2 taken together with N represent a 5- or 6-membered saturated ring; and

10 Ar represents a polysubstituted aryl group selected from the group consisting of

a)



15 wherein

W_1 represents halogen;


X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, -CN, 20 - NR_3R_4 or fluorinated acetyl or propionyl;

Y_1 represents C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen or $-CN$, or, when X_1 and Y_1 are taken together, represents $-O(CH_2)_nO-$ wherein $n = 1$ or 2 ; and

R_3 and R_4 independently represent H or C_1 - C_4 alkyl;

5

b)



wherein


W_2 represents F or Cl;

X_2 represents F, Cl, $-CN$, $-NO_2$, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, $-NR_3R_4$ or fluorinated acetyl or propionyl;

Y_2 represents halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl or $-CN$, or, when W_2 represents F, X_2 and Y_2 taken together represent $-O(CH_2)_nO-$ wherein $n = 1$ or 2 ; and

R_3 and R_4 independently represent H or C_1 - C_6 alkyl; and

c)



wherein

73776-249

Y_3 represents halogen, $-\text{CN}$ or $-\text{CF}_3$;

Z_3 represents F, Cl, $-\text{CN}$, $-\text{NO}_2$, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ alkylthio, $\text{C}_1\text{-C}_4$ alkylsulfinyl, $\text{C}_1\text{-C}_4$ alkylsulfonyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_1\text{-C}_4$ alkoxy-substituted $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy-substituted $\text{C}_1\text{-C}_4$ alkoxy, $-\text{NR}_3\text{R}_4$ or fluorinated acetyl or propionyl; and

R_3 and R_4 independently represent H, or $\text{C}_1\text{-C}_6$ alkyl;

and agriculturally acceptable derivatives of the carboxylic acid group.

Compounds of Formula I wherein Y_1 , Y_2 or Y_3 represents Cl, Br or CF_3 , wherein W_1 or W_2 represents Cl or F, wherein X_1 or X_2 represents $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ thioalkoxy, $\text{C}_1\text{-C}_4$ haloalkyl or NR_3R_4 , and wherein Z represents $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_1\text{-C}_4$ thioalkoxy, $\text{C}_1\text{-C}_4$ haloalkyl or NR_3R_4 are independently preferred.

The invention includes herbicidal compositions comprising a herbicidally effective amount of a compound of Formula I and agriculturally acceptable derivatives of the carboxylic acid group in admixture with an agriculturally acceptable adjuvant or carrier. The invention also includes a method of use of the compounds and compositions of the present invention to kill or control undesirable vegetation by application of an herbicidal amount of the compound to the vegetation or to the locus of the vegetation as well as to the soil prior to emergence of the vegetation. The invention further includes intermediates for the preparation of the compounds.

73776-249


The invention includes a compound of formula

ArMetal

wherein

Ar represents a polysubstituted aryl group selected from the group
5 consisting of

a)



wherein


W₁ represents halogen;

10 X₁ represents F, Cl, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -CN, -NR₃R₄ or fluorinated acetyl or propionyl;

15 Y₁ represents C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen or -CN, or, when X₁ and Y₁ are taken together, represents -O(CH₂)_nO- wherein n=1 or 2; and

R₃ and R₄ independently represent H or C₁-C₄ alkyl;

b)



73776-249(S)

wherein

W_2 represents F or Cl;


X_2 represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl;

Y_2 represents halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or -CN, or, when W_2 represents F, X_2 and Y_2 taken together represent -O(CH₂)_nO- wherein n=1 or 2; and

R_3 and R_4 independently represent H or C₁-C₆ alkyl; and

10

c)



wherein

Y_3 represents halogen, -CN or -CF₃;

Z_3 represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl; and

R_3 and R_4 independently represent H, or C₁-C₆ alkyl; and

Metal represents Mg-halide, Zn-halide, tri-(C₁-C₄ alkyl)tin, lithium, copper, or B(OR₈)(OR₉), where R₈ and R₉ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group.

73776-249(S)


According to an embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which R_1 and R_2 represent H.

5 According to another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q_1 represents H.

According to still another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q_2 represents Cl.

10 According to yet another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q_1 represents H and Q_2 represents Cl.

According to a further embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative
15 of the carboxylic acid group, in which Ar represents



wherein

W_1 represents halogen;

X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio,
20 C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy,

73776-249(S)

C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -CN, -NR₃R₄ or fluorinated acetyl or propionyl;

Y₁ represents C₁-C₄ alkyl, C₁-C₄ haloalkyl, halogen or -CN, or, when X₁ and Y₁ are taken together, represents -O(CH₂)_nO- wherein n=1 or 2; and


5 R₃ and R₄ independently represent H or C₁-C₄ alkyl.

According to yet a further embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which W₁ represents Cl or F.

10 According to still a further embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which X₁ represents C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkyl or -NR₃R₄.

15 According to another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Y₁ represents Cl, Br or -CF₃.

According to yet another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Ar represents



20 wherein

W₂ represents F or Cl;

73776-249(S)

X_2 represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl;


5 Y_2 represents halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or -CN, or, when W₂ represents F, X₂ and Y₂ taken together represent -O(CH₂)_nO- wherein n=1 or 2; and

R₃ and R₄ independently represent H or C₁-C₆ alkyl.

10 According to another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which X₂ represents C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkyl or -NR₃R₄.

According to still another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Y₂ represents Cl, Br or -CF₃.

15 According to yet another embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Ar represents



wherein

20 Y₃ represents halogen, -CN or -CF₃;

Z₃ represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl,

73776-249(S)


C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl; and

R₃ and R₄ independently represent H, or C₁-C₆ alkyl.

5 According to a further embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Y₃ represents Cl, Br or -CF₃.

10 According to yet a further embodiment of the present invention, there is provided a compound as described herein, or an agriculturally acceptable derivative of the carboxylic acid group, in which Z₃ represents C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkyl or -NR₃R₄.

The herbicidal compounds of the present invention are derivatives of 4-aminopicolinic acids of Formula II:




wherein

Q_1 represents H or F;

Q_2 represents a halogen with the proviso that when Q_1 is H then Q_2 is Cl or Br; and

Ar represents a polysubstituted aryl group selected from the group consisting of

a)



wherein


W_1 represents halogen;

X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, -CN, fluorinated acetyl or propionyl, or $-NR_3R_4$;

Y^1 represents C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen or $-CN$ or, when X_1 and Y_1 are taken together, represents $-O(CH_2)_nO-$ wherein $n = 1$ or 2 ; and

R_3 and R_4 independently represent H, or C_1 - C_4 alkyl;

5 b)



wherein


W_2 represents F or Cl;

X_2 represents , F, Cl, $-CN$, $-NO_2$, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, fluorinated acetyl or propionyl, or $-NR_3R_4$;

Y_2 represents halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl or $-CN$, or, when W_2 represents F, X_2 and Y_2 taken together represent $-O(CH_2)_nO-$ wherein $n = 1$ or 2 ; and

R_3 and R_4 independently represent H, or C_1 - C_6 alkyl; and

c)



wherein

Y₃ represents halogen, -CN or -CF₃;

Z₃ represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy fluorinated acetyl or propionyl, or -NR₃R₄; and

R₃ and R₄ independently represent H, or C₁-C₆ alkyl.

These compounds are characterized by possessing a Cl or F in the 3-position, H or F in the 5-position and a tri- or tetra-substituted aryl group in the 6-position of the pyridine ring. Compounds in which Cl is in the 3-position and H is in the 5-position are generally preferred. Preferred substituted aryl groups include 2,3,4-trisubstituted, 2-fluoro-4,5-trisubstituted and 2-fluoro-4,5,6-tetrasubstituted phenyl groups. Particularly preferred substituted aryl groups include those wherein Y₁, Y₂ or Y₃ represents Cl, Br or CF₃, wherein W₁ or W₂ represents Cl or F, wherein X₁ or X₂ represents C₁-C₄ alkoxy, C₁-C₄ thioalkoxy, C₁-C₄ haloalkyl or NR₃R₄, and wherein Z represents C₁-C₄ alkoxy, C₁-C₄ thioalkoxy, C₁-C₄ haloalkyl or NR₃R₄.

The amino group at the 4-position of the pyridine ring can be unsubstituted or substituted with one or more C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₆ alkynyl, hydroxy, C₁-C₆ alkoxy or amino substituents. The amino group can be further derivatized as an amide, a carbamate, a urea, a sulfonamide, a silylamine or a phosphoramidate. Such derivatives are capable of breaking down into the amine. An unsubstituted amino group is preferred.

The carboxylic acids of Formula I are believed to be the compounds that actually kill or control undesirable vegetation and are typically preferred. Analogs of these compounds in which the acid group of the picolinic acid is derivatized to form a related substituent that can be transformed within

plants or the environment to a acid group possess essentially the same herbicidal effect and are within the scope of the invention. Therefore, an “agriculturally acceptable derivative”, when used to describe the carboxylic acid functionality at the 2-position, is defined as any salt, ester, acylhydrazide, imidate, thioimidate, 5 amidine, amide, orthoester, acylcyanide, acyl halide, thioester, thionoester, dithiolester, nitrile or any other acid derivative well known in the art which (a) does not substantially affect the herbicidal activity of the active ingredient, i.e., the 6-aryl-4-aminopicolinic acid, and (b) is or can be hydrolyzed, oxidized or metabolized in plants or soil to the picolinic acid of Formula I that, depending 10 upon the pH, is in the dissociated or the undissociated form. The preferred agriculturally acceptable derivatives of the carboxylic acid are agriculturally acceptable salts, esters and amides. Likewise, an “agriculturally acceptable derivative”, when used to describe the amine functionality at the 4-position, is defined as any salt, silylamine, phosphorylamine, phosphinimine, 15 phosphoramidate, sulfonamide, sulfilimine, sulfoximine, amina, hemiaminal, amide, thioamide, carbamate, thiocarbamate, amidine, urea, imine, nitro, nitroso, azide, or any other nitrogen containing derivative well known in the art which (a) does not substantially affect the herbicidal activity of the active ingredient, i.e., the 6-aryl-4-aminopicolinic acid, and (b) is or can be hydrolyzed in plants or soil 20 to a free amine of Formula II. *N*-Oxides, which are also capable of breaking into the parent picoline of Formula II, are also covered by the scope of this invention.

Suitable salts include those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Preferred cations include sodium, potassium, magnesium, and aminium cations of the formula:



wherein R_5 , R_6 , and R_7 each, independently represents hydrogen or C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl or C_3 - C_{12} alkynyl, each of which is optionally substituted by one or more hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio or phenyl groups, provided that R_5 , R_6 , and R_7 are sterically compatible. Additionally, any two of R_5 , R_6 , and R_7

together may represent an aliphatic difunctional moiety containing 1 to 12 carbon atoms and up to two oxygen or sulfur atoms. Salts of the compounds of Formula I can be prepared by treatment of compounds of Formula I with a metal hydroxide, such as sodium hydroxide, or an amine, such as ammonia, trimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, morpholine, cyclododecylamine, or benzylamine. Amine salts are often preferred forms of the compounds of Formula I because they are water-soluble and lend themselves to the preparation of desirable aqueous based herbicidal compositions.

- 10 Suitable esters include those derived from C₁-C₁₂ alkyl, C₃-C₁₂ alkenyl or C₃-C₁₂ alkynyl alcohols, such as methanol, *iso*-propanol, butanol, 2-ethylhexanol, butoxyethanol, methoxypropanol, allyl alcohol, propargyl alcohol or cyclohexanol. Esters can be prepared by coupling of the picolinic acid with the alcohol using any number of suitable activating agents such as those used for
- 15 peptide couplings such as dicyclohexylcarbodiimide (DCC) or carbonyl diimidazole (CDI), by reacting the corresponding acid chloride of a picolinic acid of Formula I with an appropriate alcohol or by reacting the corresponding picolinic acid of Formula I with an appropriate alcohol in the presence of an acid catalyst. Suitable amides include those derived from ammonia or from C₁-C₁₂
- 20 alkyl, C₃-C₁₂ alkenyl or C₃-C₁₂ alkynyl mono- or di-substituted amines, such as but not limited to dimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, cyclododecylamine, benzylamine or cyclic or aromatic amines with or without additional heteroatoms such as but not limited to aziridine, azetidine, pyrrolidine, pyrrole, imidazole, tetrazole or morpholine.
- 25 Amides can be prepared by reacting the corresponding picolinic acid chloride, mixed anhydride, or carboxylic ester of Formula I with ammonia or an appropriate amine.


The terms “alkyl”, “alkenyl” and “alkynyl”, as well as derivative terms such as “alkoxy”, “acyl”, “alkylthio” and “alkylsulfonyl”, as used herein, include within their scope straight chain, branched chain and cyclic moieties. The terms “alkenyl” and “alkynyl” are intended to include one or more unsaturated
5 bonds.

The term “aryl”, as well as derivative terms such as “aryloxy”, refers to a phenyl.

Unless specifically limited otherwise, the term “halogen” including derivative terms such as “halo” refers to fluorine, chlorine, bromine, and iodine.
10 The terms “haloalkyl” and “haloalkoxy” refer to alkyl and alkoxy groups substituted with from 1 to the maximum possible number of halogen atoms.

The compounds of Formula I can be made using well-known chemical procedures. The required starting materials are commercially available or readily synthesized utilizing standard procedures.


15 The 6-substituted arylpicolines of Formula I can be prepared from a number of ways, which are well known in the art, e.g., by reaction of an appropriately substituted pyridine with a facile leaving group in the 6-position (III) with an organometallic compound of the type (IV) in an inert solvent in the presence of a transition metal catalyst.



20 In this case “L” can be chlorine, bromine, iodo or trifluoromethanesulfonate, “Metal” can be Mg-halide, Zn-halide, tri-(C₁-C₄ alkyl)tin, lithium, copper, or B(OR₈)(OR₉), where R₈ and R₉ are, independent of one another, hydrogen, C₁-C₄

alkyl, or when taken together form an ethylene or propylene group, and
 “Catalyst” is a transition metal catalyst, in particular a palladium catalyst such as
 palladium diacetate, bis(triphenylphosphine)palladium(II)dichloride, or a nickel
 catalyst such as nickel(II)acetylacetonate, bis(triphenylphosphine)nickel(II)
 5 chloride.

Alternatively, compounds of Formula I can be prepared by
 reaction of an appropriately substituted 6-metal substituted pyridine (V) with an
 aryl compound of the type (VI) in an inert solvent in the presence of a transition
 metal catalyst.



10 In this case “L” can be chlorine, bromine, iodo or trifluoromethanesulfonate and
 “Metal” can be Mg-halide, Zn-halide, tri-(C₁-C₄ alkyl)tin, lithium, copper, or
 B(OR₈)(OR₉), where R₈ and R₉ are independent of one another, hydrogen, C₁-
 C₄ alkyl, or when taken together form an ethylene or propylene group, and
 “Catalyst” can be a transition metal catalyst, in particular a palladium catalyst
 15 such as palladium diacetate, bis(triphenylphosphine)palladium(II)dichloride, or a
 nickel catalyst such as nickel(II)acetylacetonate, bis(triphenylphosphine)nickel(II)
 chloride.

Reactions with boronic acids or esters are well known as
 exemplified by the following references:

- 20 (1) W.J. Thompson and J. Gaudino, J. Org. Chem., 49, 5223 (1984);
 (2) S. Gronowitz and K. Lawitz, Chem. Scr., 24, 5 (1984);
 (3) S. Gronowitz et al., Chem. Scr., 26, 305 (1986);

- (4) J. Stavenuiter et al., Heterocycles, 26, 2711 (1987);
- (5) V. Snieckus et al., Tetrahedron Letters, 28, 5093 (1987);
- (6) V. Snieckus et al., Tetrahedron Letters, 29, 2135 (1988);
- (7) M.B. Mitchell et al., Tetrahedron Letters, 32, 2273 (1991); Tetrahedron,
5 48, 8117 (1992);
- (8) JP-A 93/301870.

Reactions with Grignard compounds (metal = Mg-Hal):

- (9) L.N. Pridgen, J. Heterocyclic Chem., 12, 443 (1975);
- (10) M. Kumada et al., Tetrahedron Letters, 21, 845 (1980);
- 10 (11) A. Minato et al., J. Chem. Soc. Chem. Commun., 5319 (1984).

Reaction with organozinc compounds (metal = Zn-Hal):

- (12) A.S. Bell et al., Synthesis, 843 (1987);
- (13) A.S. Bell et al., Tetrahedron Letters, 29, 5013 (1988);
- (14) J.W. Tilley and S. Zawoiski, J. Org. Chem., 53, 386 (1988); see also
15 ref.(9).

Reactions with organotin compounds (metal = Sn(C₁-C₄(alkyl)₃):

- (15) T.R. Bailey et al., Tetrahedron Letters, 27, 4407 (1986);
- (16) Y. Yamamoto et al., Synthesis, 564 (1986); see also ref.(6)

The coupling of III+IV, or V+VI may, where appropriate, be followed by reactions on either ring to obtain further derivatives of the compounds of Formula I.

Appropriately substituted pyridines of Formula III where L is
5 chloro, bromo, iodo or trifluoromethanesulfonate can be easily obtain by well-known methods; see WO 0151468. For example, 6-bromo analogs can be prepared by the reduction of several key intermediates, e.g., the corresponding 6-bromo-4-azido, 6-bromo-4-nitro, and 6-bromo-4-nitro pyridine *N*-oxide analogs. These intermediates, in turn, can be prepared either by nucleophilic displacement
10 of 6-bromo-4-halo analogs with NaN_3 or by electrophilic nitration of the corresponding 6-bromopyridine-*N*-oxides. Alternatively, such analogs can be prepared by direct amination of the corresponding 4,6-dibromo analogs.

4-*N*-Amide, carbamate, urea, sulfonamide, silylamine and phosphoramidate amino derivatives can be prepared by the reaction of the free
15 amino compound with, for example, a suitable acid halide, chloroformate, carbamyl chloride, sulfonyl chloride, silyl chloride or chlorophosphate.

Substituted 4-amino analogs can be prepared by reacting the corresponding 4-halopyridine-2-carboxylate or any other displaceable 4-substituent with the substituted amine.

20 The compounds of Formula I, obtained by any of these processes, can be recovered by conventional means. Typically, the reaction mixture is acidified with an aqueous acid, such as hydrochloric acid, and extracted with an organic solvent, such as ethyl acetate or methylene chloride. The organic solvent and other volatiles can be removed by distillation or evaporation to obtain the
25 desired compound of Formula I, which can be purified by standard procedures, such as by recrystallization or chromatography.

The compounds of Formula I have been found to be useful as pre-emergence and post-emergence herbicides. They can be employed at non-selective (higher) rates of application to control a broad spectrum of the vegetation in an area or at lower rates of application for the selective control of undesirable vegetation. Areas of application include pasture and rangelands, roadsides and rights of way, power lines and any industrial areas where control of undesirable vegetation is desirable. Another use is the control of unwanted vegetation in crops such as corn, rice and cereals. They can also be used to control undesirable vegetation in tree crops such as citrus, apple, rubber, oil palm, forestry and others. It is usually preferred to employ the compounds post-emergence. It is further usually preferred to use the compounds to control a wide spectrum of woody plants, broadleaf and grass weeds, and sedges. Use of the compounds to control undesirable vegetation in established crops is especially indicated. While each of the 6-aryl-4-aminopicolinate compounds encompassed by Formula I is within the scope of the invention, the degree of herbicidal activity, the crop selectivity, and the spectrum of weed control obtained varies depending upon the substituents present. An appropriate compound for any specific herbicidal utility can be identified by using the information presented herein and routine testing.

The term herbicide is used herein to mean an active ingredient that kills, controls or otherwise adversely modifies the growth of plants. An herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect and includes deviations from natural development, killing, regulation, desiccation, retardation, and the like. The terms plants and vegetation include germinant seeds, emerging seedlings and established vegetation.

Herbicidal activity is exhibited by the compounds of the present invention when they are applied directly to the plant or to the locus of the plant at

any stage of growth or before planting or emergence. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific
5 compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical applied. These and other factors can be adjusted as is known in the art to promote non-selective or selective herbicidal action. Generally, it is preferred to apply the compounds of Formula I postemergence to relatively immature undesirable vegetation to achieve the
10 maximum control of weeds.

Application rates of 0.1 to 1,000 g/Ha are generally employed in postemergence operations; for preemergence applications, rates of 1 to 2,000 g/Ha are generally employed. The higher rates designated generally give non-selective control of a broad variety of undesirable vegetation. The lower rates
15 typically give selective control and can be employed in the locus of crops.

The herbicidal compounds of the present invention are often applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction with other herbicides, the presently claimed compounds can be formulated with the other
20 herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the compounds of the present invention include: *amide herbicides* such as allidochlor, beflubutamid, benzadox, benzipram, bromobutide, cafenstrole, CDEA, chlorthiamid, cyprazole,
25 dimethenamid, dimethenamid-P, diphenamid, epronaz, etnipromid, fentrazamide, flupoxam, fomesafen, halosafen, isocarbamid, isoxaben, napropamide, naptalam, pethoxamid, propyzamide, quinonamid and tebutam; *anilide herbicides* such as chloranocryl, cisanilide, clomeprop, cypromid, diflufenican, etobenzanid,

fenasulam, flufenacet, flufenican, mefenacet, mefluidide, metamifop, monalide, naproanilide, pentanochlor, picolinafen and propanil; *arylalanine herbicides* such as benzoylprop, flamprop and flamprop-M; *chloroacetanilide herbicides* such as acetochlor, alachlor, butachlor, butenachlor, delachlor, diethatyl, dimethachlor, metazachlor, metolachlor, S-metolachlor, pretilachlor, propachlor, propisochlor, prynachlor, terbuchlor, thenylchlor and xylachlor; *sulfonanilide herbicides* such as benzofluor, perfluidone, pyrimisulfan and profluazol; *sulfonamide herbicides* such as asulam, carbasulam, fenasulam and oryzalin; *antibiotic herbicides* such as bilanafos; *benzoic acid herbicides* such as chloramben, dicamba, 2,3,6-TBA and tricamba; *pyrimidinyloxybenzoic acid herbicides* such as bispyribac and pyriminobac; *pyrimidinylthiobenzoic acid herbicides* such as pyriithiobac; *phthalic acid herbicides* such as chlorthal; *picolinic acid herbicides* such as aminopyralid, clopyralid and picloram; *quinolinecarboxylic acid herbicides* such as quinclorac and quinmerac; *arsenical herbicides* such as cacodylic acid, CMA, DSMA, hexaflurate, MAA, MAMA, MSMA, potassium arsenite and sodium arsenite; *benzoylcyclohexanedione herbicides* such as mesotrione, sulcotrione, tefuryltrione and tembotrione; *benzofuranyl alkylsulfonate herbicides* such as benfuresate and ethofumesate; *carbamate herbicides* such as asulam, carboxazole chlorprocarb, dichlormate, fenasulam, karbutilate and terbucarb; *carbanilate herbicides* such as barban, BCPC, carbasulam, carbetamide, CEPC, chlorbufam, chlorpropham, CPPC, desmedipham, phenisopham, phenmedipham, phenmedipham-ethyl, propham and swep; *cyclohexene oxime herbicides* such as alloxydim, butroxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, tepraloxym and tralkoxydim; *cyclopropylisoxazole herbicides* such as isoxachlortole and isoxaflutole; *dicarboximide herbicides* such as benzfendazole, cinidon-ethyl, flumezin, flumiclorac, flumioxazin and flumipropyn; *dinitroaniline herbicides* such as benfluralin, butralin, dinitramine, ethalfluralin, fluchloralin, isopropalin, methalpropalin, nitralin, oryzalin, pendimethalin, prodiamine, profluralin and trifluralin; *dinitrophenol herbicides*

such as dinofenate, dinoprop, dinosam, dinoseb, dinoterb, DNOC, etinofen and medinoterb; *diphenyl ether herbicides* such as ethoxyfen; *nitrophenyl ether herbicides* such as acifluorfen, aclonifen, bifenox, chlomethoxyfen, chlornitrofen, etnipromid, fluorodifen, fluoroglycofen, fluoronitrofen, fomesafen, furyloxyfen, halosafen, lactofen, nitrofen, nitrofluorfen and oxyfluorfen; *dithiocarbamate herbicides* such as dazomet and metam; *halogenated aliphatic herbicides* such as alorac, chloropon, dalapon, flupropanate, hexachloroacetone, iodomethane, methyl bromide, monochloroacetic acid, SMA and TCA; *imidazolinone herbicides* such as imazamethabenz, imazamox, imazapic, imazapyr, imazaquin and imazethapyr; *inorganic herbicides* such as ammonium sulfamate, borax, calcium chlorate, copper sulfate, ferrous sulfate, potassium azide, potassium cyanate, sodium azide, sodium chlorate and sulfuric acid; *nitrile herbicides* such as bromobonil, bromoxynil, chloroxynil, dichlobenil, iodobonil, ioxynil and pyraclonil; *organophosphorus herbicides* such as amiprofos-methyl, anilofos, bensulide, bilanafos, butamifos, 2,4-DEP, DMPA, EBEP, fosamine, glufosinate, glyphosate and piperophos; *phenoxy herbicides* such as bromofenoxim, clomeprop, 2,4-DEB, 2,4-DEP, difenopenten, disul, erbon, etnipromid, fenteracol and trifopsime; *phenoxyacetic herbicides* such as 4-CPA, 2,4-D, 3,4-DA, MCPA, MCPA-thioethyl and 2,4,5-T; *phenoxybutyric herbicides* such as 4-CPB, 2,4-DB, 3,4-DB, MCPB and 2,4,5-TB; *phenoxypropionic herbicides* such as cloprop, 4-CPP, dichlorprop, dichlorprop-P, 3,4-DP, fenoprop, mecoprop and mecoprop-P; *aryloxyphenoxypropionic herbicides* such as chlorazifop, clodinafop, clofop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-P, fenthiaprop, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, isoxapyrifop, metamifop, propaquizafop, quizalofop, quizalofop-P and trifop; *phenylenediamine herbicides* such as dinitramine and prodiamine; *pyrazolyl herbicides* such as benzofenap, pyrazolynate, pyrasulfotole, pyrazoxyfen, pyroxasulfone and topramezone; *pyrazolylphenyl herbicides* such as fluazolate and pyraflufen; *pyridazine herbicides* such as credazine, pyridafol and pyridate;

pyridazinone herbicides such as brompyrazon, chloridazon, dimidazon, flufenpyr, metflurazon, norflurazon, oxapyrazon and pydanon; *pyridine herbicides* such as aminopyralid, clidinate, clopyralid, dithiopyr, fluroxypyr, haloxydine, picloram, picolinafen, pyriclor, thiazopyr and triclopyr; *pyrimidinediamine herbicides* such as iprymidam and tioclorim; *quaternary ammonium herbicides* such as cyperquat, diethamquat, difenzoquat, diquat, morfamquat and paraquat; *thiocarbamate herbicides* such as butylate, cycloate, di-allate, EPTC, esprocarb, ethiolate, isopolinate, methiobencarb, molinate, orbencarb, pebulate, prosulfocarb, pyributicarb, sulfallate, thiobencarb, tiocarbazil, tri-allate and vernolate; *thiocarbonate herbicides* such as dimexano, EXD and proxan; *thiourea herbicides* such as methiuron; *triazine herbicides* such as dipropetryn, triaziflam and trihydroxytriazine; *chlorotriazine herbicides* such as atrazine, chlorazine, cyanazine, cyprazine, eglinazine, ipazine, mesoprazine, procyazine, proglinazine, propazine, sebuthylazine, simazine, terbuthylazine and trietazine; *methoxytriazine herbicides* such as atraton, methometon, prometon, secbumeton, simeton and terbumeton; *methylthiotriazine herbicides* such as ametryn, aziprotryne, cyanatryn, desmetryn, dimethametryn, methoprotryne, prometryn, simetryn and terbutryn; *triazinone herbicides* such as ametridione, amibuzin, hexazinone, isomethiozin, metamitron and metribuzin; *triazole herbicides* such as amitrole, cafenstrole, epronaz and flupoxam; *triazolone herbicides* such as amicarbazone, bencarbazone, carfentrazone, flucarbazone, propoxycarbazone, sulfentrazone and thienicarbazone-methyl; *triazolopyrimidine herbicides* such as cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam and pyroxsulam; *uracil herbicides* such as butafenacil, bromacil, flupropacil, isocil, lenacil and terbacil; *3-phenyluracils*; *urea herbicides* such as benzthiazuron, cumyluron, cycluron, dichloralurea, diflufenzopyr, isonoruron, isouron, methabenzthiazuron, monisouron and noruron; *phenylurea herbicides* such as anisuron, buturon, chlorbromuron, chloreturon, chlorotoluron, chloroxuron, daimuron, difenoxuron, dimefuron, diuron, fenuron, fluometuron, fluothiuron, isoproturon, linuron,

methiuron, methyldymron, metobenzuron, metobromuron, metoxuron, monolinuron, monuron, neburon, parafluron, phenobenzuron, siduron, tetrafluron and thidiazuron; *pyrimidinylsulfonylurea herbicides* such as amidosulfuron, azimsulfuron, bensulfuron, chlorimuron, cyclosulfamuron, ethoxysulfuron, 5 flazasulfuron, flucetosulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, mesosulfuron, nicosulfuron, orthosulfamuron, oxasulfuron, primisulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron and trifloxysulfuron; *triazinylsulfonylurea herbicides* such as chlorsulfuron, cinosulfuron, ethametsulfuron, iodosulfuron, metsulfuron, prosulfuron, 10 thifensulfuron, triasulfuron, tribenuron, triflusulfuron and tritosulfuron; *thiadiazolylurea herbicides* such as buthiuron, ethidimuron, tebuthiuron, thiazafluron and thidiazuron; and *unclassified herbicides* such as acrolein, allyl alcohol, azafenidin, benazolin, bentazone, benzobicyclon, buthidazole, calcium cyanamide, cambendichlor, chlorfenac, chlorfenprop, chlorflurazole, 15 chlorflurenol, cinmethylin, clomazone, CPMF, cresol, ortho-dichlorobenzene, dimepiperate, endothal, fluoromidine, fluridone, flurochloridone, flurtamone, fluthiacet, indanofan, methazole, methyl isothiocyanate, nipyraclufen, OCH, oxadiargyl, oxadiazon, oxaziclomefone, pentachlorophenol, pentoxazone, phenylmercury acetate, pinoxaden, prosulfalin, pyribenzoxim, pyriftalid, 20 quinoclamine, rhodethanil, sulglycapin, thidiazimin, tridiphane, trimeturon, tripropindan and tritac. The herbicidal compounds of the present invention can, further, be used in conjunction with glyphosate, glufosinate or 2,4-D on glyphosate-tolerant, glufosinate-tolerant or 2,4-D-tolerant crops. It is generally preferred to use the compounds of the invention in combination with herbicides 25 that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate employed. It is further generally preferred to apply the compounds of the invention and other complementary herbicides at the same time, either as a combination formulation or as a tank mix.

The compounds of the present invention can generally be employed in combination with known herbicide safeners, such as benoxacor, benthocarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlormid, dicyclonon, dimepiperate, disulfoton, fenchlorazole-ethyl, fencloirim, 5 flurazole, fluxofenim, furilazole, isoxadifen-ethyl, mefenpyr-diethyl, MG 191, MON 4660, naphthalic anhydride (NA), oxabetrinil, R29148 and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity. They can additionally be employed to control undesirable vegetation in many crops that have been made tolerant to or resistant to them or to other herbicides by genetic manipulation or 10 by mutation and selection. For example, corn, wheat, rice, soybean, sugarbeet, cotton, canola, and other crops that have been made tolerant or resistant to compounds that are acetolactate synthase inhibitors in sensitive plants can be treated. Many glyphosate and glufosinate tolerant crops can be treated as well, alone or in combination with these herbicides. Some crops (e.g. cotton) have 15 been made tolerant to auxinic herbicides such as 2,4-dichlorophenoxyacetic acid. These herbicides may be used to treat such resistant crops or other auxin tolerant crops.

While it is possible to utilize the 6-aryl-4-amino-picolinate compounds of Formula I directly as herbicides, it is preferable to use them in 20 mixtures containing a herbicidally effective amount of the compound along with at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with the compounds of 25 Formula I or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water dispersible

granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions.

Suitable agricultural adjuvants and carriers that are useful in preparing the herbicidal mixtures of the invention are well known to those skilled
5 in the art.

Liquid carriers that can be employed include water, toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol
10 monomethyl ether, methanol, ethanol, isopropanol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, and the like. Water is generally the carrier of choice for the dilution of concentrates.

Suitable solid carriers include talc, pyrophyllite clay, silica, attapulgius clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium
15 carbonate, bentonite clay, Fuller's earth, cotton seed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, and the like.

It is usually desirable to incorporate one or more surface-active agents into the compositions of the present invention. Such surface-active agents are advantageously employed in both solid and liquid compositions, especially
20 those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. Typical surface-active agents include salts of alkyl sulfates, such as diethanol-ammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecyl-
25 benzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as

tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkylnaphthalene-sulfonate salts, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethyl-
5 ammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono and dialkyl phosphate esters.

Other adjuvants commonly used in agricultural compositions include compatibilizing agents, antifoam agents, sequestering agents, neutralizing
10 agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids, sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid
15 fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

The concentration of the active ingredients in the herbicidal compositions of this invention is generally from 0.001 to 98 percent by weight. Concentrations from 0.01 to 90 percent by weight are often employed. In
20 compositions designed to be employed as concentrates, the active ingredient is generally present in a concentration from 5 to 98 weight percent, preferably 10 to 90 weight percent. Such compositions are typically diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds generally contain about 0.0001 to 1 weight
25 percent active ingredient and preferably contain 0.001 to 0.05 weight percent.

The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators,

by addition to irrigation water, and by other conventional means known to those skilled in the art.

Many of the starting materials useful for the preparation of the compounds of the present invention, e.g., 4-amino-3,6-dichloropyridine-2-carboxylic acid, 4-amino-3,5,6-trifluoro-2-cyanopyridine, methyl 4-amino-6-bromo-3,5-difluoropyridine-2-carboxylate and methyl 4-amino-6-bromo-3-chloropyridine-2-carboxylate, are described in US Patent 6,297,197 B1.

Examples:

1. Preparation of 3-bromo-6-chloro-2-fluorophenol

A solution of 1-bromo-4-chloro-2-fluorobenzene (20.4 g, 0.100 mol) in tetrahydrofuran (THF; 50 mL) was slowly added to lithium diisopropyl amide (LDA; 0.125 mol) in THF (600 mL) at -50° C. After addition the solution was warmed to -20° C and then cooled to -50° C and a solution of trimethyl borate (13.5 g, 0.130 mol) in tetrahydrofuran (20 mL) was added slowly and the temperature was warmed to -20° C. The mixture was then cooled to -70° C and solution of peracetic acid (32 percent in acetic acid, 0.150 mol) was slowly added and the mixture was warmed to ambient temperature. Water (250 mL) was added and solution extracted with ethyl acetate (2 x 200 mL). The combined organic phases were dried and concentrated. The black oil was purified by column chromatography (20 percent ethyl acetate in hexanes) to give 3-bromo-6-chloro-2-fluorophenol (14.1g, 0.063 mol): ¹H NMR (CDCl₃): δ 7.05 (m, 2H), 5.5 (bs, 1H).

The following compounds were prepared according to the procedure of Example 1.

3-Bromo-2,6-dichlorophenol: mp 69-70° C.

3-Bromo-2-fluoro-6-trifluoromethylphenol: ^1H NMR (CDCl_3): δ 7.20 (m, 2H), 5.85 (bs, 1H).

3-Bromo-2-chloro-6-fluorophenol: LC/MS ($m/z=225$).

2. Preparation of 1-bromo-4-chloro-2-fluoro-3-methoxybenzene

5 A heterogeneous mixture of 3-bromo-6-chloro-2-fluorophenol (14.4 g, 0.064 mol), methyl iodide (13.5 g, 0.096 mol) and potassium carbonate (8.8 g, 0.064 mol) in acetonitrile (100 mL) was heated under reflux for 2 hours. The mixture was cooled, diluted with water (100 mL) and extracted with ethyl ether (2 x 150 mL). The combined extracts were dried and concentrated. The
10 dark oil was purified by chromatography (5 percent ethyl acetate in hexanes) to give 1-bromo-4-chloro-2-fluoro-3-methoxybenzene (14.8 g, 0.062 mol): ^1H NMR (CDCl_3): δ 7.20 (m, 1H), 7.10 (dd, 1 H), 4.0 (s, 3H).

The following compounds were prepared according to the procedure of Example 2.

15 1-Bromo-4-chloro-3-ethoxy-2-fluorobenzene: ^1H NMR (CDCl_3): δ 7.20 (m, 1H), 7.10 (dd, 1H), 4.20 (q, 2H), 1.50 (t, 3H).

1-Bromo-4-chloro-2-fluoro-3-isopropoxybenzene: ^1H NMR (CDCl_3): δ 7.20 (m, 1H), 7.10 (dd, 1H), 4.5 (m, 1H), 1.40 (d, 6H).

1-Bromo-4-chloro-2-fluoro-3-(2-methoxyethoxy) benzene: ^1H NMR (CDCl_3): δ
20 7.25 (m, 1H), 7.15 (dd, 1H), 4.25 (t, 2H), 3.75 (t, 2H), 3.5 (s, 3H).

1-Bromo-2-fluoro-3-methoxy-4-trifluoromethylbenzene: ^1H NMR (CDCl_3): δ 7.39 (d, 1H), 7.21 (d, 1H), 6.18 (tt, 1H), 4.24 (td, 2H).

73776-249(S)

1-Bromo-2,4-dichloro-3-ethoxybenzene: ^1H NMR (CDCl_3): δ 7.32 (d, 1H), 7.17 (d, 1H), 4.10 (q, 2H), 1.47 (t, 3H).

1-Bromo-2,4-dichloro-3-methoxybenzene: ^1H NMR (CDCl_3): δ 7.35 (d, 1H), 7.15 (d, 1H), 3.95 (s, 3H).

5 1-Chloro-3,5-difluoro-2-methoxybenzene: GC-MS ($m/z=178$).

1-Chloro-3,5-difluoro-2-ethoxybenzene: GC-MS ($m/z=192$) bp 80-85°/ 30 mm.

1,3-Dichloro-5-fluoro-2-methoxybenzene: GC-MS ($m/z=194$).

1-Bromo-3-butoxy-4-chloro-2-fluorobenzene: GC-PCI ($m/z=180$).

1-Bromo-4-chloro-2-fluoro-3-methoxymethoxybenzene: GC-MS ($m/z=269$).

10 1-Bromo-2-chloro-4-fluoro-3-methoxybenzene: GC-MS ($m/z=239$).

3-Chloro-5-fluoro-2-methoxybenzaldehyde: GC-MS ($m/z=188$).

1,3-Difluoro-3-ethoxybenzene: GC-MS ($m/z=158$).

3. Preparation of 1-bromo-4-chloro-2-fluoro-5-methoxybenzene

A solution of 4-chloro-2-fluoro-5-methoxyaniline (25.0 g, 0.143 mol) in 10 percent HBr (250 mL) was cooled to 0° C and a solution of sodium nitrite (15.0 g, 0.218 mol) in water (20 mL) was slowly added. Methylene chloride (50 mL) and cupric bromide (30.0 g, 0.244 mol) were added slowly and then the mixture was warmed to ambient temperature and stirred for 1 hour. The reaction mixture was filtered through a bed of celite and extracted with methylene chloride (2 x 100 mL) and the combined organic phases were dried (sodium sulfate) and concentrated. Chromatography of the dark oil (5 percent ethyl

15

20

acetate in hexanes) gave 1-bromo-4-chloro-2-fluoro-5-methoxybenzene (16.6 g, 0.070 mol): ^1H NMR (CDCl_3): δ 7.20 (m, 1H) 7.05 (dd, 1H), 4.00 (s, 3H).

4. Preparation of 1-bromo-4-chloro-2-fluoro-3-difluoromethoxybenzene

To a solution of 3-bromo-6-chloro-2-fluorophenol (1.00 g, 4.44 mmol) and sodium chlorodifluoroacetate in dimethylformamide (DMF; 9 mL) was added potassium carbonate (1.22 g, 5.32 mmol) and water (1.77 mL) and the resulting mixture heated to 100° C for 4 hours. The solution was cooled to ambient temperature and concentrated hydrochloric acid (2.5 mL) and water (4 mL) were added and stirred at ambient temperature overnight. The solution was cooled in an ice bath and neutralized with 2N sodium hydroxide and then extracted with ethyl ether (2 x 25 mL). The combined extracts were washed with brine, dried (sodium sulfate), and concentrated to give 1-bromo-4-chloro-2-fluoro-3-difluoromethoxybenzene (1.20 g, 4.2 mmol). This material was used without further purification.

5. Preparation of 1-bromo-4-chloro-3-(2,2-difluoroethoxy)-2-fluorobenzene

A solution of 3-bromo-6-chloro-2-fluorophenol (15.4 g, 0.068 mol) in DMF (25 mL) was slowly added to a suspension of sodium hydride (60 percent dispersion in mineral oil) (4.0 g, 0.10 mol) in DMF (100 mL) and the mixture was stirred 1 hour. A solution of methanesulfonic acid 2,2 difluoroethyl ester (17.5 g, 0.109 mol) in DMF (10 mL) was slowly added. The resulting solution was heated to 70° C for 18 hours. The solution was diluted with water (200 mL) and extracted with ethyl ether. The combined organic phases were dried (sodium sulfate) and concentrated. The residual oil was purified by chromatography (in hexanes) to give 1-bromo-4-chloro-3-(2,2-difluoroethoxy)-2-fluorobenzene (9.0 g, 0.031 mol): ^1H NMR (CDCl_3): δ 7.3 (m, 1H), 7.10 (dd, 1H), 6.15 (dt, 1H), 4.35 (m, 2H).

The following compound was prepared according to the procedure of Example 5.

1-Bromo-2,4-dichloro-3-(2,2-difluoroethoxy)benzene: ^1H NMR (CDCl_3): δ 7.35 (m, 1H), 7.15 (dd, 1H), 6.15 (tt, 1H), 4.35 (dt, 2H).

5 6. Preparation of 1-bromo-4-chloro-2-fluoro-3-(methylthio)benzene

A solution of 1-bromo-4-chloro-2-fluorobenzene (20.4 g, 0.100 mol) in THF (50 mL) was slowly added to LDA (0.125 mol) in THF (600 mL) at -50°C . After addition, the solution was warmed to -20°C and then cooled to -50°C and a solution of dimethyldisulfide (18.8 g, 0.20 mol) in THF (50 mL) was
10 slowly added and the mixture was warmed to ambient temperature. The reaction was quenched with water (200 mL) and extracted with ethyl acetate (2 x 150 mL) and the combined organic phases dried (sodium sulfate) and concentrated. The residual red oil was purified by chromatography (5 percent ethyl acetate in hexanes) to give 1-bromo-4-chloro-2-fluoro-3-(methylthio)benzene (23.9 g,
15 0.094mol): ^1H NMR (CDCl_3): δ 7.40 (m, 1H), 7.15 (dd, 1H), 2.50 (s, 3H).

The following compound was prepared according to the procedure of Example 6.

1-Bromo-2,4-dichloro-3-(methylthio)benzene: ^1H NMR (CDCl_3): δ 7.52 (d, 1H), 7.25 (d, 1H), 2.46 (s, 3H).

20 7. Preparation of 3-bromo-6-chloro-2-fluorobenzaldehyde

A solution of 1-bromo-4-chloro-2-fluorobenzene (20.4 g, 0.100 mol) in THF (50 mL) was slowly added to LDA (0.125 mol) in THF (600 mL) at -50°C . After addition the solution was warmed to -20°C and then cooled to -50°C and a solution of DMF (14.6 g, 0.20 mol) in THF (50 mL) was slowly added
25 and the reaction mixture was warmed to ambient temperature. The reaction was

quenched with water (250 mL) and extracted with ethyl acetate (2 x 150 mL) and the combined organic phases were dried (sodium sulfate) and concentrated. The solid residue was recrystallized from hexane to give 3-bromo-6-chloro-2-fluorobenzaldehyde (40.0 g, 0.169 mol): mp 92-93° C.

5 The following compounds were prepared according the procedure of Example 7 by quenching with appropriate electrophile, e.g., an aldehyde, ester or ketone.

1-(3-Bromo-6-chloro-2-fluorophenyl)-1-ethanol: ¹H NMR (CDCl₃): δ 7.40 (m, 1H), 7.15 (m, 1H), 5.40 (m, 1H), 2.50 (m, 1H), 1.85 (d, 3H).

10 3-Bromo-2,4-dichlorobenzaldehyde: mp 96-97° C.

1-(3-Bromo-2,6-dichlorophenyl)-1-ethanol: ¹H NMR (CDCl₃): δ 7.60 (d, 1H), 7.15 (d, 1H), 5.45 (m, 1H), 3.0 (d, 1H), 1.85 (d, 3H).

1-Bromo-4-chloro-3-ethyl-2-fluorobenzene: GC/PCI (m/z=236).

15 1-(3-Bromo-6-chloro-2-fluorophenyl)-2,2,2-trifluoroethanone: GC/PCI (m/z=304).

1-(3-Bromo-6-chloro-2-fluorophenyl)-1-propanol: ¹H NMR (CDCl₃): δ 7.40 (m, 1H), 7.15 (m, 1H), 5.40 (m, 1H), 2.50 (m, 1H), 1.90 (m, 4H), 1.00 (t, 3H).

2-(3-Bromo-6-chloro-2-fluorophenyl)-2-propanol: ¹H NMR (CDCl₃): δ 7.40 (m, 1H), 7.15 (m, 1H), 3.45 (s, 1H), 1.80 (m, 6H).

20 8. Preparation of 3-chloro-5-fluoro-2-hydroxybenzaldehyde

A solution of 2-chloro-4-fluorophenol (15.0 g, 102 mmol) in 150 g of 50 percent sodium hydroxide, water (30 mL) and chloroform (45 mL) were heated under reflux for 8 hours, additional chloroform (45 mL) was added every 2

hours. The mixture was then cooled and allowed to stand for 18 hours and the precipitated sodium salt was collected. The solid was slurried in water (200 mL) and the pH adjusted to 1.5 with 6N hydrochloric acid and then extracted with ethyl acetate (2 x 100 mL). The combined extracts were washed with brine, dried
5 and concentrated. The residue was purified by chromatography (1:1 methylene chloride:hexanes) to give 3-chloro-5-fluoro-2-hydroxybenzaldehyde (5.0 g, 29 mmol): LC/MS (m/z=174).

9. Preparation of 1-bromo-4-chloro-2-fluoro-3-difluoromethylbenzene

Diethylamino sulfur trifluoride (15.3 g, 0.096 mol) was added
10 slowly to a solution of 3-bromo-6-chloro-2-fluorobenzaldehyde (7.50 g, 0.032 mol) in methylene chloride at 0° C and the mixture stirred for 1 hour after allowing to warm to ambient temperature. The reaction was carefully quenched with a saturated solution of sodium bicarbonate in water (100 mL) and extracted with methylene chloride (2 x 75 mL) and the combined extracts dried (sodium
15 sulfate) and concentrated to give 1-bromo-4-chloro-2-fluoro-3-difluoromethylbenzene (7.20 g, 0.028 mol): ¹H NMR (CDCl₃): δ 7.60 (m, 1H), 7.05 (m, 1H), 7.00 (d, 1H).

The following compounds were prepared according to the procedure in Example 9.

20 1-Bromo-2,4-dichloro-3-difluoromethylbenzene: LC/MS ESI (m/z=271).

1-Difluoromethyl-3-fluoro-4-iodo-2-methoxybenzene: LC/MS (m/z=302).

2-Chloro-4-difluoromethyl-3-methoxy-1-nitrobenzene: LC/MS (m/z=237) (used without complete purification).

3-Chloro-1-difluoromethyl-2-methoxybenzene: LC/MS (m/z=210).

10. Preparation of 2-chloro-4-difluoromethyl-3-methoxyaniline

A solution of crude 2-chloro-4-difluoromethyl-3-methoxy-1-nitrobenzene in ethanol (50 mL) containing 5 percent palladium on charcoal (250 mg) was hydrogenated (50 psi) for 5 hours. The solution was filtered and
5 concentrated to give 2-chloro-4-difluoromethyl-3-methoxyaniline (1.3 g, 48 mmol): LC/MS ($m/z=207$).

11. Preparation of *N*-pivaloyl-2-bromo-4-chloro-3-methoxyaniline

To solution of *N*-pivaloyl-4-chloro-3-methoxyaniline (5.0 g, 21 mmol) in tetrahydrofuran (60 mL) at -60°C was added *n*-butyl lithium (2.5M, 44
10 mmol). The solution was allowed to warm to 0°C and stir for 3 hours. Ethylene dibromide (9.9 g, 53 mmol) was added and the solution stirred for 18 hours before being quenched with ammonium chloride (saturated solution, 20 mL) and extracted with ethyl ether (2 x 50 mL). The combined extracts were washed with brine, dried (sodium sulfate) and concentrated. The residue was purified by
15 chromatography (10 percent ethyl acetate/hexanes) to give *N*-pivaloyl-2-bromo-4-chloro-3-methoxyaniline (3.7 g, 1.2 mmol): LC/MS ($m/z=320$).

12. Preparation of 2-bromo-4-chloro-1-iodo-3-methoxybenzene

A solution of *N*-pivaloyl-2-bromo-4-chloro-3-methoxyaniline (3.7 g, 1.2 mmol) in dioxane (35 mL) was treated with concentrated hydrochloric acid
20 (35 mL) and the solution heated under reflux for 2 hours. After the solution was cooled the pH was adjusted to 10 by addition of sodium hydroxide (50 percent solution) and extracted with ethyl ether (2 x 50 mL). The combined extracts were washed with brine, dried (sodium sulfate) and concentrated. The crude aniline was dissolved in 2N hydrochloric acid and sodium nitrite (750 mg, 11 mmol) was
25 added and the solution stirred for 20 minutes. This solution was poured into a vigorously stirred solution of sodium iodide (3.3 g, 22 mmol) in water (20 mL)

and methylene chloride (30 mL) and stirred for 20 minutes. The solution was then extracted with methylene chloride (2 x 25mL) and the combined extracts washed with brine, dried (sodium sulfate) and concentrated. The residue was purified by chromatography (hexanes) to give 2-bromo-4-chloro-1-iodo-3-methoxybenzene (1.5 g, 0.6 mmol): LC/MS (m/z=267).

The following compounds were prepared according to the procedure of Example 12.

4-Chloro-2,6-difluoro-5-methyl-1-iodobenzene: bp 60-70° C/ 0.75 mm.

2-Chloro-4-difluoromethyl-3-methoxy-1-iodobenzene: LC/MS (m/z=318).

10 1-chloro-2-difluoromethyl-3,5-difluoro-4-iodobenzene: LC/MS (m/z=324).

13. Preparation of 3-chloro-2-methoxy-4-nitrotoluene

A solution of 4-nitro-2-hydroxytoluene (5.0 g, 33 mmol) in chloroform (40 mL) was heated under reflux and treated with a stream of chlorine for 1 hour. Upon cooling, the solution was washed with sodium bicarbonate (saturated solution), brine, dried (sodium sulfate) and concentrated. The residue was dissolved in acetonitrile (75 mL), treated with potassium carbonate (9.1 g, 66 mmol) and iodomethane (2.5 mL, 5.7 g, 40 mmol) and heated under reflux for 2 hours. Upon cooling, the solution was concentrated and the residue was taken up in ethyl ether (75 mL) and washed with water (50 mL), brine, dried (sodium sulfate) and concentrated. The crude product was purified by chromatography on silica with 1-5 percent ethyl acetate/hexane to 3-chloro-2-methoxy-4-nitrotoluene (3.0 g, 15 mmol): LC/MS (m/z=201).

14. Preparation of 3-chloro-2-methoxy-4-nitrobenzaldehyde

A solution of 3-chloro-2-methoxy-4-nitrotoluene (1.0 g, 5.0 mmol) in acetic acid (10 mL) and acetic anhydride (10 mL) was cooled to 5° C and

treated with 1 mL conc. H₂SO₄. Chromium trioxide (1.4 g, 14 mmol) was added in portions over 10 minutes while the reaction temperature was maintained at 0-5° C. After 30 minutes at this temperature, the mixture was poured onto ice-water (20 g) and the mixture was extracted with ethyl acetate (2 x 25 mL). The combined
5 organic phases were washed with brine, dried (sodium sulfate) and concentrated. The crude bis-acetate was combined with water (10 mL), ethanol (10 mL) and concentrated sulfuric acid (1 mL) and heated to reflux for 40 minutes. Upon cooling, the mixture was diluted with water (20 mL) and extracted with ethyl acetate (2 x 50 mL). The combined extracts were washed with brine, dried
10 (sodium sulfate) and concentrated. The crude mixture was chromatographed (10 percent ethyl acetate/hexanes) to give 3-chloro-2-methoxy-4-nitrobenzaldehyde (400 mg, 1.9 mmol): LC/MS (m/z=215).

15. Preparation of (3-bromo-6-chloro-2-fluorophenyl) methanol

To a solution of 3-bromo-6-chloro-2-fluorobenzaldehyde (20.0 g, 0.084 mol) in ethanol (250 mL) was added sodium borohydride (6.4 g, 0.168 mol) and the reaction mixture was stirred at ambient temperature for 1 hour. The mixture was diluted with water (300 mL) and extracted with ethyl acetate (2 x 200 mL), dried (sodium sulfate) and concentrated. The crude solid was recrystallized from hexanes to give (3-bromo-6-chloro-2-fluorophenyl) methanol
20 (10.4 g, 0.043 mol): ¹H NMR (CDCl₃): δ 7.50 (m, 1H), 7.15 (dt, 1H), 4.90 (s, 2H), 2.10 (bs, 1H).

The following compounds were prepared according to the procedure of Example 15.

1-(3-Bromo-6-chloro-2-fluorophenyl)-2,2,2-trifluoroethanol: GC/PCI (m/z=306).

25 3-Fluoro-4-iodo-2-methoxyphenyl)methanol: LC/MS (m/z=282).

16. Preparation of 1-bromo-4-chloro-3-fluoromethyl-2-fluorobenzene

A solution of diethylamino sulfur trifluoride (7.6 g, 0.047 mol) in methylene chloride (25 mL) was slowly added to (3-bromo-6-chloro-2-fluorophenyl) methanol (10.3 g, 0.043 mol) in methylene chloride (150 mL) at 5 0° C. The solution was stirred at ambient temperature for 1 hour. A saturated solution of sodium bicarbonate (100 mL) was cautiously added and the mixture extracted with methylene chloride (2 x 100 mL). The combined organic phases were dried (sodium sulfate) and concentrated. The dark residual oil was purified by chromatography (5 percent ethyl acetate in hexanes) to give 1-bromo-4-chloro-10 3-fluoromethyl-2-fluorobenzene (6.20 g, 0.024 mol): ¹H NMR (CDCl₃): δ 7.55(m, 1H) 7.18 (dd, 1H) 5.6 (d, 2H).

The following compounds were prepared according to the procedure in Example 16.

1-Bromo-4-chloro-2-fluoro-3-(1-fluoro-1-methylethyl)benzene: ¹H NMR 15 (CDCl₃): δ 7.42 (m, 1H), 7.10 (m, 1H), 6.05 (dq, 1H) 1.95 (dd, 6H).

1-Bromo-2,4-dichloro-3-(1-fluoroethyl)benzene: GC/PCI (m/z=270).

1-Bromo-4-chloro-2-fluoro-3-(1-fluoroethyl)benzene: GC/PCI (m/z=255).

1-Bromo-4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)benzene: GC/EI (m/z=308).

20 1-Bromo-4-chloro-2-fluoro-3-(1-fluoropropyl)benzene: GC/EI (m/z=267).

2-Fluoro-4-fluoromethyl-1-iodo-3-methoxybenzene: LC/MS (m/z=284).

17. Preparation of 1-bromo-4-chloro-2-fluoro-3-methoxymethylbenzene

A solution of 1-bromo-4-chloro-2-fluorobenzene (20.4 g, 0.100 mol) in THF (50 mL) was slowly added to LDA (0.125 mol) in THF (600 mL) at -50° C. After addition the solution was warmed to -20° C and then cooled to -50° C a solution of bromomethoxymethane (25 g, 0.200 mol) in THF (25 mL) was slowly added and the reaction mixture was warmed to ambient temperature. The reaction was quenched with water (400 mL) and extracted with diethyl ether (2 x 150 mL). The combined organic phases were dried (sodium sulfate) and concentrated. The residual oil was distilled (70°-75° C/ 0.5 mm) to give 1-bromo-4-chloro-2-fluoro-3-methoxymethylbenzene (18.0 g, 0.071 mol): ¹H NMR (CDCl₃): δ 7.50 (m, 1H), 7.15 (dd, 1H), 4.65 (s, 1H), 3.40 (s, 1H).

The following compound was prepared according to the procedure of Example 17.

1-Bromo-4-chloro-2-fluoro-3-methylbenzene: ¹H NMR (CDCl₃): δ 7.30 (m, 1H), 7.05 (dd, 1H), 2.35 (s, 3H).

18. Preparation of 2-(4-chloro-2-fluoro-3-methoxyphenyl)-[1,3,2]-dioxaborinane

To a solution of 1-bromo-4-chloro-2-fluoro-3-methoxybenzene (10.4 g, 0.043 mol) in diethyl ether (150 mL) at -78° C was slowly added *n*-butyl lithium (2.5M, 19.0 mL, 0.0475 mol) and the solution was stirred for 30 minutes. A solution of triisopropyl borate (12.0 g, 0.064 ml) in THF (25 mL) was slowly added and the solution was warmed to a 0° C. Acetyl chloride (10.0 g, 0.13 mol) was added. After stirring for 1 hour, the solution was concentrated and the solid residue taken into ethyl acetate (150 mL) and a 1N solution of sodium hydroxide (50 mL). Ice was added to the aqueous phase which was then acidified with sufficient concentrated hydrochloric acid to obtain a pH of 2. The heterogeneous

mixture was extracted with ethyl acetate (2 x 150 mL) and the combined organic phases dried (sodium sulfate) and concentrated. The resulting solid was slurried in toluene and propane-1,3-diol (6.6 g, 0.09 mol) was added and the mixture heated under reflux with water being removed with a Dean-Stark trap. After 2
5 hours the mixture was cooled and concentrated. The resulting oil was taken into methylene chloride (50 mL) and washed with water (25 mL), then dried (sodium sulfate) and concentrated to give 2-(4-chloro-2-fluoro-3-methoxyphenyl)-[1,3,2]-dioxaborinane (6.4 g, 0.062 mol): ¹H NMR (CDCl₃): δ 7.15 (m, 1H), 6.95 (dd, 1H), 4.05 (t, 4H), 3.8 (s, 3H), 1.95 (t, 2H).

10 The following compounds were prepared according to the procedure of Example 18.

2-(4-Chloro-2-fluoro-5-methoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃): δ 7.25 (d, 1H), 7.05 (d, 1H), 4.20 (t, 4H), 4.15 (s, 3H), 2.10 (t, 2H).

2-(4-Chloro-2-fluoro-3-ethoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃):
15 δ 7.30 (m, 1H), 7.05 (dd, 1H), 4.20 (m, 7H), 2.05 (t, 3H), 1.50 (t, 3H).

2-(4-Chloro-2-fluoro-3-(methylthio)phenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃): δ 7.50 (m, 1H), 7.18 (dd, 1H), 4.20 (t, 4H), 2.50 (s, 3H), 2.05 (t, 2H).

2-(4-Chloro-2-fluoro-3-methoxymethylphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃): δ 7.6 (dt, 1H), 7.25 (dd, 1H), 4.76 (s, 2H), 4.20 (t, 4H), 3.40 (s, 3H),
20 2.05 (t, 2H).

2-(4-Chloro-2-fluoro-3-isopropoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃): δ 7.25 (m, 1H), 7.15 (dd, 1H), 4.5 (q, 1H), 4.20 (t, 4H), 2.05 (t, 2H), 1.50 (d, 6H).

2-(4-Chloro-2-fluoro-3-difluoromethylphenyl)-[1,3,2]-dioxaborinane ¹H NMR (CDCl₃): δ 7.75 (m, 1H), 7.15 (dd, 1H), 6.90-7.15 (t, 1H) 4.20 (t, 4H), 2.05 (t, 2H).

2-(4-Chloro-2-fluoro-3-fluoromethylphenyl)-[1,3,2]-dioxaborinane ¹H NMR
5 (CDCl₃): δ 7.70 (m, 1H), 7.25 (dd, 1H), 5.8 (d, 2H), 4.20 (t, 4H), 2.05 (t, 2H).

4-[1,3,2]-Dioxaborinan-2-yl-3-fluoro-2-methoxybenzonitrile: ¹H NMR (CDCl₃):
δ 7.40 (m, 1H), 7.3 (dd, 1H), 4.25 (t, 4H), 4.15 (s, 3H), 2.10 (t, 3H).

2-(4-Chloro-2-fluoro-3-methoxyethoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR
(CDCl₃): δ 7.35 (m, 1H), 7.15 (dd, 1H), 4.25 (m, 6H), 3.75 (d, 2H), 3.48 (s, 3H),
10 2.15 (t, 3H).

2-(2, 4-Dichloro-3-ethoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (DMSO-*d*₆): δ
7.40 (d, 1H), 7.29 (d, 1H), 4.08 (m, 4H), 4.00 (q, 2H), 1.99 (m, 2H), 1.34 (t, 3H).

2-[4-Chloro-2-fluoro-3-(2,2-difluoroethoxyphenyl)]-[1,3,2]-dioxaborinane: ¹H
NMR (CDCl₃): δ 7.45 (m, 1H), 7.15 (dd, 1H), 6.15 (tt, 1H), 4.38 (t, 4H), 4.20 (t,
15 2H), 2.10 (t, 2H).

2-(4-Chloro-2-fluoro-5-ethoxyphenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃):
δ 7.25 (d, 1H), 7.05 (d, 1H), 4.20 (t, 4H), 4.15 (t, 2H), 2.10 (t, 2H), 1.45 (t, 3H).

2-(2,4-Dichloro-3-(methylthio)phenyl)-[1,3,2]-dioxaborinane: ¹H NMR (CDCl₃):
δ 7.45-7.28 (m, 2H), 3.86 (m, 4H), 2.42 (s, 3H), 1.80 (m, 2H).

20 6-Chloro-3-[1,3,2]-dioxaborinan-2-yl-2-fluorobenzonitrile: ¹H NMR (CDCl₃): δ
7.85 (m, 1H), 6.25 (m, 1 H), 4.20 (m, 4H), 2.10 (m, 2H).

19. Preparation of 1-fluoro-2,3-methylenedioxybenzene

Alliquat 336 (methyltrioctylammonium chloride, 0.63 g, 1.6 mmol), dibromomethane (40.7 g, 234.2 mmol) and water (31 mL) were placed in a 500 mL 3-necked flask equipped with an addition funnel, condenser and a stir bar. The addition funnel was charged with a solution of 3-fluorocatechol (20.0 g, 160 mmol) in 5M sodium hydroxide (80 mL). The mixture in the flask was heated to reflux and the solution of the catechol was added dropwise with good stirring over 2 hours and the resulting dark mixture heated an additional 2 hours at reflux. After cooling to room temperature, the reaction was diluted with methylene chloride and the layers separated. The aqueous layer was extracted with methylene chloride and the combined organic layers dried (Na₂SO₄ with charcoal). Filtration and concentration to a constant weight on the rota-vap gave 1-fluoro-2,3-methylenedioxybenzene (14.6 g, 104.2 mmol) as a dark yellow oil: ¹H NMR (CDCl₃): 6.80 (m, 1H), 6.68 (m, 2H), 6.04 (s, 2H).

20. Preparation of 2-fluoro-3,4-methylenedioxyphenylboronic acid

1-Fluoro-2,3-methylenedioxybenzene (5.0 g, 35.7 mmol) was dissolved in THF (70 mL) and the solution cooled to -65° C in a dry ice acetone bath. *n*-Butyl lithium (2.5 g, 15.7 mL, 39.3 mmol) was added to the solution via syringe with stirring. The reaction was allowed to warm to -35° C over 1 hour then re-cooled to -65° C and treated with trimethylborate (4.1 g, 39.3 mmol) via syringe. The reaction was allowed to warm slowly to room temperature then quenched with 1N HCl (50 mL), stirred for 15 minutes then diluted with ether and the layers separated. The organic layer was extracted with 1N sodium hydroxide and this extract separated from the ether and acidified with 1N hydrochloric acid. The acidic aqueous layer was extracted with two portions of ether and these combined ether extracts were dried (sodium sulfate), filtered and concentrated to a oily solid which was triturated with methylene dichloride. The resulting solid

was collected by filtration, washed with methylene dichloride and dried to give 1-fluoro-2,3-methylenedioxyphenylboronic acid (1.4 g, 7.6 mmol) as a tan solid: ^1H NMR ($\text{DMSO}-d_6$): δ 8.05 (bs, 2H), 7.08 (dd, 1H), 6.76 (d, 1H), 6.08 (s, 2H).

5 The following compounds were prepared according to the procedure in Example 20.

2-Fluoro-3-methoxy-4-methylphenylboronic acid: ^1H NMR (300 MHz, d_6 -DMSO + H_2O): δ 7.32 (bs, 2H), 7.11 (dd, 1H, $J=7.2, 5.7$ Hz), 6.92 (d, 1H, $J=7.2$ Hz), 3.74 (s, 3H), 2.19 (s, 3H).

10

4-Bromo-2-fluoro-3-methoxyphenylboronic acid: ^1H NMR (300 MHz, $\text{DMSO } d_6 + \text{D}_2\text{O}$): δ 8.36 (bs, 2H), 7.38 (dd, 1H, $J=8.4$), 7.17 (dd, 1H), 3.82 (d, 3H).

15 21. Preparation of 2-(2-fluoro-3,4-methylenedioxyphenyl)-[1,3,2]-dioxaborinane

2-Fluoro-3,4-methylenedioxyphenylboronic acid (1.4 g, 7.6 mmol) was slurried in toluene and propane-1,3-diol (0.6 g, 7.9 mmol) and the mixture heated under reflux for 0.5 hour. The mixture was cooled, filtered through glass wool and concentrated to give 2-(2-fluoro-3,4-methylenedioxyphenyl)-[1,3,2]-dioxaborinane (1.6 g, 7.1 mmol): ^1H NMR ($\text{DMSO}-d_6$): δ 7.12 (dd, 1H, $J=7.8, 5.7$ Hz), 6.75 (d, 1H), 6.09 (s, 2H), 4.06 (t, 4H), 1.98 (m, 2H).

20

22. Preparation of 3-bromo-6-chloro-2-fluorobenzonitrile

25 A suspension of 3-bromo-6-chloro-2-fluorobenzaldehyde (9.0 g, 0.04 mol) and *O*-sulfinic acid hydroxyamine (7.50 g, 0.07 mole) in water (300 mL) was warmed to 50° C for 18 hours. The suspension was cooled and the solid collected to give 3-bromo-6-chloro-2-fluorobenzonitrile (8.8 g, 0.04 mol): ^1H NMR (CDCl_3): δ 7.75 (m, 1H), 7.25 (m, 1H).

23. Preparation of 3-bromo-2-fluoro-6-chlorobenzamide

Concentrated sulfuric acid (15 mL) was placed in a 100 mL 3-neck flask equipped with an internal thermometer and then heated to 55° C. 3-Bromo-2-fluoro-6-chlorobenzonitrile (11.0 g, 47 mmol) was added portion-wise to the acid with stirring maintaining the temperature above 50° C. The dark solution was heated at 65° C for 24 hours then cooled to room temperature and poured over ice and cautiously neutralized with concentrated ammonium hydroxide. The mixture was extracted with two portions of ethyl acetate and the combined organic layers dried over Na₂SO₄ (with charcoal). Filtration and concentration gave 3-bromo-2-fluoro-6-chlorobenzamide (11.5 g, 45.5 mmol) as a light orange solid: mp 157-158° C, ¹H NMR (CDCl₃): δ 7.54 (t, 1H), 7.14 (dd, 1H), 6.03 (bs, 1H) 5.81 (bs, 1H).

24. Preparation of 3-bromo-2,6-dichlorobenzamide

3-Bromo-2,6-dichlorobenzoic acid (7.2 g, 26.7 mmol) was treated with thionyl chloride (30 mL, 400.0 mmol) in a 250 mL round bottom flask and the mixture was heated to reflux for 1 hour then cooled to room temperature. Approximately one third of the light yellow solution was removed and concentrated *in vacuo*. The residue was dissolved in THF (16 mL) and treated with concentrated ammonium hydroxide (6 mL) with good stirring. The mixture was partitioned between ethyl acetate and saturated NaHCO₃ and the layers separated and the organics dried over Na₂SO₄. Filtration and concentration gave 3-Bromo-2,6-dichlorobenzamide (2.19 g) as a foam: mp 117-119 °C, ¹H NMR (300 MHz, d₆-DMSO): δ 8.10 (bs, 1H), 7.88 (bs, 1H), 7.79 (d, 1H, J=8.), 7.44 (d, 1H).

25. Preparation of 3-bromo-6-chloro-2-fluoroaniline

Sodium hydroxide (4.0 g, 100.0 mmol) was dissolved in water (70 mL) in a 250 mL round bottomed flask and the resulting solution cooled in an ice

bath and treated with bromine (4.7 g, 29.7 mmol) to give a yellow solution. 3-Bromo-2-fluoro-6-chlorobenzenecarboxamide (5.0 g, 19.9 mmol) was added as a solid, slowly with good stirring and the orange mixture was heated to reflux for 2 hours. The cooled reaction mixture was diluted with methylene dichloride and the layers separated and the methylene dichloride dried (Na₂SO₄). Crystallization of the concentrated filtrate from cold hexanes gave 3-bromo-6-chloro-2-fluoroaniline (2.8 g, 12.6 mmol) as a off white solid: mp 61-62° C, ¹H NMR (CDCl₃): δ 6.94 (dd, 1H), 6.83 (dd, 1H), 4.16 (bs, 2H).

The following compound was prepared according to the procedure in Example 25.

3-Bromo-2,6-dichloroaniline: mp 71-72° C.

26. Preparation of *N*-(3-bromo-6-chloro-2-fluorophenyl)-*N,N*-dimethylamine

3-Bromo-6-chloro-2-fluoroaniline (2.5 g, 11.1 mmol) was dissolved in THF (25 mL) in a nitrogen flushed 250 mL round bottomed flask and treated with 37 percent formaldehyde (0.84 g, 2.1 mL, 27.8 mmol), dibutyl tin dichloride (0.07 g, 0.22 mmol) and phenyl silane (1.33 g, 12.3 mmol) and allowed to stir at room temperature under nitrogen for 48 hours. The reaction mixture was concentrated *in vacuo* and purified by flash silica gel chromatography (hexanes) to give *N*-(3-bromo-6-chloro-2-fluorophenyl)-*N,N*-dimethylamine (2.0 g 7.9 mmol) as a oil: ¹H NMR (CDCl₃): δ 7.19 (dd, 1H), 7.04 (dd, 1H), 2.88 (s, 3H), 2.87 (s, 3H).

The following compounds were prepared according to the procedure of Example 26.

N-(3-Bromo-6-chloro-2-fluorophenyl)-*N,N*-diethylamine: ¹H NMR (300 MHz, CDCl₃): δ 7.24 (dd, 1H), 7.07 (dd, 1H), 3.16 (dd, 4H), 1.01 (t, 6H).

N-(3-Bromo-2,6-dichlorophenyl)- *N,N* -dimethylamine: ¹H NMR (300 MHz, CDCl₃): δ 7.33 (d, 1H), 7.13 (d, 1H), 2.88 (s, 6H).

N-(2-Chloro-4,6-difluorophenyl)- *N,N* -dimethylamine: Colorless oil. ¹H NMR (300 MHz, CDCl₃): δ 6.93 (ddd, 1H), 6.73 (m, 1H), 2.82 (d, 6H).

27. Preparation of *N*-(3-bromo-2-fluorophenyl), *N*-methyl amine

3-Bromo-6-chloro-2-fluoroaniline (1.2 g, 5.2 mmol) was dissolved in THF (12 mL) in a 100 mL round bottomed flask and treated with tetra-*N*-butyl ammonium bromide (0.015 g, 0.05 mmol) and powdered sodium hydroxide (0.83 g, 20.9 mmol) with stirring. After 5 minutes, the dark mixture was treated with dimethyl sulfate (1.3 g, 10.4 mmol) and the mixture heated at 60° C for 2 hours. Analysis by TLC (hexanes: ethyl acetate / 20:1) indicated some starting material remained. Added additional dimethyl sulfate (0.66 g, 5.2 mmol) and continued heating for 1.5 hour then stirred at room temperature overnight. The reaction was partitioned between diethyl ether and water and the layers separated and the aqueous phase extracted with ether. The combined ether layers were washed with water and dried (MgSO₄). Filtration, concentration and purification by flash silica gel chromatography (hexanes) gave *N*-(3-bromo-6-chloro-2-fluorophenyl)-*N*-methylamine (0.81 g) as a colorless oil: ¹H NMR (300 MHz, CDCl₃): δ 6.93 (dd, 1H, J=8.4), 6.83 (dd, 1H), 4.00(bs, 1H), 3.09 (d, 3H).

28. Preparation of 1,3-difluoro-2-iodo-4-methoxy-5-methylbenzene

A solution of 1,5-difluoro-2-methoxy-3-methylbenzene (0.65 g, 4.1 mmol) in THF (1 mL) was added to LDA (4.3 mmol) in THF (10 mL) at -55° C and the solution stirred for 45 minutes. Solid iodine (1.15 g, 4.5 mmol) was added and the solution allowed to warm to ambient temperature. The solution was diluted with ethyl acetate and washed with sodium bicarbonate (saturated solution 15 mL) and sodium thiosulfate (saturated solution, 15 mL), dried (sodium sulfate) and concentrated. The residue was purified by

chromatography (hexanes) to give 1,3-difluoro-2-iodo-4-methoxy-5-methylbenzene: ^1H NMR (300 MHz, CDCl_3): δ 6.72 (dd, 1H), 3.84 (d, 3H), 2.26 (s, 3H).

The following compounds were prepared according to the procedure of
5 Example 28.

3,5-Difluoro-4-iodo-2-methoxybenzonitrile: mp 106-110° C.

N-(6-Chloro-2,4-difluoro-3-iodophenyl)- *N,N* -dimethylamine: ^1H NMR (300 MHz, CDCl_3): δ 6.98 (dd, 1H), 2.82 (d, 6H).

1,3-Dichloro-5-fluoro-4-iodo-2-methoxybenzene: GC-MS ($m/z=320$).

10 1,3-Difluoro-2-iodo-4-methoxy-5-methylbenzene: ^1H NMR (CDCl_3): δ 6.72 (dd, 1H), 3.84 (d, 3H), 2.26 (s, 3H).

2,3-Difluoro-4-iodo-benzonitrile: LC/MS ($m/z=265$).

3-Chloro-1-difluoromethyl-5-fluoro-4-iodo-2-methoxybenzene: LC/MS ($m/z=336$).

15 2-Ethoxy-1,3-difluoro-4-iodobenzene: GC-MS ($m/z=284$).

1,3-Difluoro-2-iodo-4-methoxy-5-methylbenzene: ^1H NMR (CDCl_3): δ 6.72 (dd, 1H), 3.84 (d, 3H), 2.26 (s, 3H).

3,5 Difluoro-4-iodo-2-methoxybenzonitrile: mp. 106-110° C.

(6-Chloro-2,4-difluoro-3-iodophenyl) dimethyl amine: ^1H NMR (CDCl_3): δ 6.98
20 (dd, 1H), 2.82 (d, 6H).

2,4,6-Trifluoro-3-iodobenzonitrile: LC/MS ($m/z=283$).

1-Bromo-3,5-difluoro-4-iodo-2-methoxybenzene: LC/MS ($m/z=349$).

1,3-Dichloro-5-fluoro-4-iodo-2-methoxybenzene: GC-MS ($m/z=320$).

29. Preparation of 1-chloro-3,5-difluoro-4-iodo-2-methoxybenzene.

To a solution of 1-chloro-3,5-difluoro-2-methoxybenzene (2.0 g, 0.01 mol) in THF at -75°C was added *n*-butyl lithium (2.5M in hexanes 6.7mL) and the resulting solution was stirred at -75°C for 1 hour. A solution of iodine (5.1 g, 0.02 mol) in THF (10 mL) was added and the reaction mixture was allowed to warm to ambient temperature. The reaction mixture was diluted with ethyl ether (50 mL) and aqueous sodium thiosulfate (10 percent, 50 mL) was added and stirred for 1 hour. After separating the phases, the organic phase was dried (Na_2SO_4) and concentrated to give 1-chloro-3,5-difluoro-4-iodo-2-methoxybenzene as a white solid: mp $62-64^{\circ}\text{C}$.

The following compound was prepared according to the procedure in Example 29.

1-Chloro-2-ethoxy-3,5-difluoro-4-iodobenzene: GC-MS ($m/z=178$).

30. Preparation of 3-fluoro-4-iodo-2-methoxybenzonitrile.

Sodium hydride (60 mg, 2.5 mmol) was slurried in DMF (15 mL) and treated with dry methanol (120 μl , 96 mg, 3.0 mmol). After stirring for 10 minutes at 25°C , the solution was cooled to -25°C and treated with 2,3-difluoro-4-iodobenzonitrile (500 mg, 1.9 mmol). After 25 minutes, the mixture was quenched by addition of 10 percent citric acid solution (5 mL) and warmed to 25°C . The mixture was extracted with ethyl ether (2 x 15 mL), the combined ether phases washed with water (5 mL), brine, dried (sodium sulfate) and evaporated to give 3-fluoro-4-iodo-2-methoxybenzonitrile (500 mg, 1.8 mmol): LC/MS ($m/z=277$).

31. Preparation of 3-fluoro-4-iodo-2-methoxybenzaldehyde

3-Fluoro-4-iodo-2-methoxybenzonitrile (1.0 g, 3.6 mmol) was dissolved in toluene (7 mL), cooled to 0° C and treated in portions with 25 wt. percent diisobutyl aluminum hydride (DIBAL) in toluene (3.1 ml, 4.7 mmol)
5 After 30 minutes, the reaction was quenched by addition of methanol (5 mL), poured into 1M sulfuric acid (15 mL) and extracted with ethyl acetate (2 x 15 mL). The combined extracts were washed with water (5 mL), brine, dried (sodium sulfate) and evaporated to give 3-fluoro-4-iodo-2-methoxybenzaldehyde (450 mg, 1.6 mmol): LC/MS (m/z=260).

10 32. Preparation of 3-chloro-4-iodo-2-methoxyaniline

3-Chloro-2-methoxyaniline (1 g, 6.4 mmol) was dissolved in methylene chloride (10 mL) in a 250 mL round bottomed flask and treated with water (10 mL), NaHCO₃ (1.1 g, 12.7 mmol) and iodine (1.61 g, 6.4 mmol) and the resulting mixture heated to reflux for 2 hours by which time TLC
15 (hexanes:ethyl acetate / 4:1) indicated nearly complete conversion. The reaction mixture was diluted with ethyl acetate and washed with saturated Na₂S₂O₃. The organic layer was separated and dried (Na₂SO₄). Filtration, concentration and purification by flash silica gel chromatography (hexanes:ethyl acetate / 8:1) gave 0.71 g of a red oil which was identified as a 2.5:1 mixture of the 3-chloro-4-iodo-
20 2-methoxyaniline and starting material: ¹H NMR (300 MHz, CDCl₃) δ 7.36 (d, 1H, J=8.4 Hz), 6.45 (d, 1H, J=8.4 Hz), 4.10 (bs, 2H), 3.82 (s, 3H). Other runs with excess iodine produced similar results. The mixture was combined with a previous run and used as is without further purification.

33. Preparation of 4-chloro-2,6-difluoro-3-methylaniline

25 A solution of 2,6-difluoro-3-methylaniline (15 g, 105 mmol) in acetic acid (80 mL) was heated to 70° C and a solution of sulfuryl chloride (16.0 g, 115 mmol) in acetic acid (40 mL) was added dropwise over 20 minutes. The

temperature was maintained for 3 hours, then cooled and concentrated. The residue taken into water (100 mL) and the pH was adjusted to 9 with 2N sodium hydroxide and then extracted with methylene chloride (2 x 75 mL) and the combined extracts washed with brine, dried (sodium sulfate) and concentrated.

5 The residue was distilled to give 4-chloro-2,6-difluoro-3-methylaniline (12.2 g, 69 mmol): bp 55°-65° C/55 mm.

34. Preparation of 1-bromo-3-chloro-4-iodo-2-methoxybenzene

A mixture of 3-chloro-2-methoxy aniline and 3-chloro-4-iodo-2-methoxyaniline (2.8 g, 2.5:1) was dissolved in dioxane (20 mL) and the resulting solution cooled on an ice bath while being treated with 48 percent HBr (20 mL). The dark purple mixture was then treated with NaNO₂ (1.4 g, 19.5 mmol) in water (6 mL) with good stirring. After 10 minutes, CuBr (5.1 g, 35.5 mmol) was added and the mixture was removed from the ice bath and allowed to warm to room temperature and stir for 0.5 hour. The dark purple mixture was diluted with ethyl acetate and water then filtered through celite with ethyl acetate. The layers were separated and the organic layer was washed with saturated sodium sulfite and saturated ammonium chloride and dried (Na₂SO₄). Filtration, concentration and purification by flash silica gel chromatography (hexanes) gave 2.95 g of a colorless oil which was determined to be a 2.3:1 mixture of 1-bromo-3-chloro-2-methoxybenzene and 1-bromo-3-chloro-4-iodo-2-methoxybenzene which was used as is without further purification: ¹H NMR (300 MHz, CDCl₃): δ 7.49 (d, 1H), 7.19 (d, 1H), 3.90 (s, 3H).

10
15
20

35. Preparation of 1-chloro-3,5-difluoro-4-trimethylsilyl-benzene

To a solution of 3,5-difluoro-1-chlorobenzene (10.0 g, 67 mmol) and tetramethylethylenediamine (TMEDA; 7.8 g, 67 mmol) in THF (75 mL) at -75° C, was added *n*-butyl lithium (2.5M, 68 mmol) and the resulting solution stirred for 30 minutes. A solution of trimethylsilyl chloride (7.6 g, 70 mmol) in THF (15 mL) was added and the solution allowed to warm to ambient

25

temperature and stir for 25 hours. Ammonium chloride (saturated solution 15 mL) was added and the mixture extracted with ethyl ether (2 x 100 mL) and the combined extracts washed with brine, dried (sodium sulfate) and concentrated. The residual oil was distilled to give 1-chloro-3,5-difluoro-4-

5 trimethylsilylbenzene (13.0 g, 60 mmol): bp 70-75° C.

36. Preparation of 6-chloro-2,4-difluoro-3-trimethylsilylbenzaldehyde

A solution of 1-chloro-3,5-difluoro-4-trimethylsilylbenzene (1.5 g, 6.8 mmol) in THF (10 mL) was added to a solution of lithium tetramethylpiperdine (14 mmol) in THF (15 mL) at -75° C. The solution was
10 stirred for 2 hours then DMF (1.5 g, 20 mmol) was added and the solution allowed to warm to ambient temperature. The reaction was quenched with a saturated solution of ammonium chloride (25 mL) and extracted with ethyl ether (2 x 25 mL) and the combined extracts dried (sodium sulfate) and concentrated. The residue was purified by chromatography (hexanes) to give 6-chloro-2,4-
15 difluoro-3-trimethylsilylbenzaldehyde (1.4 g, 5.6 mmol): LC/MS (m/z=248).

37. Preparation of 6-chloro-2,4-difluoro-3-iodobenzaldehyde

A solution of 6-chloro-2,4-difluoro-3-trimethylsilylbenzaldehyde (600 mg, 2.4 mmol) in methylene chloride (7 mL) was cooled to 0° C and treated with solution of iodochloride (780 mg, 4.8 mmol) in methylene chloride (10 mL)
20 and stirred at ambient temperature for 2 hours. Additional iodochloride (900 mg) was added and the solution heated under reflux for 40 minutes. The solution was allowed to cool and was washed with sodium bisulfite (15 ml, 5 percent solution), water (10 mL), and a saturated solution of sodium bicarbonate (10 mL). The organic phase was dried (sodium sulfate) and concentrated to give 6-chloro-2,4-
25 difluoro-3-iodobenzaldehyde (620 mg, 3.9 mmol): LC/MS (m/z=302).

38. Preparation of 1-bromo-2, 4-difluoro-5-methoxybenzene.

To a solution of 1,3 dibromo-4,6-difluorobenzene (10.0 g, 37 mmol) in THF (50 mL) at -20° C was added isopropyl magnesium chloride (2.0M in THF, 42 mmol) and the resulting solution warm to 0° C and stirred for 30 minutes. Trimethylborate (4.7 g, 45 mmol) was added and the mixture stirred at ambient temperature for 1 hour. The solution was recooled to -20° C and peracetic acid (32 percent, 50 mmol) was added and the solution was stirred for 30 minutes at ambient temperature. The solution was then quenched with sodium bisulfite (5 percent solution, 75 mL) and then acidified with 6 N hydrochloric acid and extracted with ethyl ether (2 x 75 mL). Combined extracts were dried (sodium sulfate) and concentrated. The crude phenol was dissolved in acetonitrile (40 mL) and potassium carbonate (10 g, 74 mmol) and iodomethane (5.7 g, 40 mmol) were added and stirred at ambient temperature for 20 hours. The solution was concentrated and the residue taken into water (50 mL) and extracted with ethyl ether (2 x 75 mL) and the combined extracts were washed with brine, dried (sodium sulfate), and concentrated. The residue was purified by chromatography (hexanes) to give 1-bromo-2,4-difluoro-5-methoxybenzene (2.0 g, 8.6 mmol): LC/MS (m/z=223).

39. Preparation of 2-chloro-4,6-difluoro-1-iodo-3-methoxybenzene

A solution of 4-chloro-2,6-difluoro-1-iodo-3-methoxybenzene (2.0 g, 6.6 mmol) in THF (7 mL) was added to LDA (8.0 mmol) in THF(15 mL) at -55° C and the reaction stirred for 2 hours. The solution was diluted with ethyl ether (50 mL) and washed with ammonium chloride (saturated solution 25 mL), brine and then dried (sodium sulfate) and concentrated. The residual oil was purified by reverse phase chromatography (70 percent acetonitrile in water) to give 2-chloro-4,6-difluoro-1-iodo-3-methoxybenzene (500 mg. 1.7 mmol): LC/MS (m/z=304).

40. Preparation 4-chloro-3-(dimethylamino)-2-fluorophenylboronic acid

N-(3-Bromo-6-chloro-2-fluorophenyl)-*N,N*-dimethylaniline (0.88 g 3.5 mmol) was dissolved in ether (10 mL) in an oven dried, nitrogen flushed, 50 mL 3-necked flask equipped with a thermometer. The solution was cooled to -60° C under nitrogen. *n*-Butyl lithium (0.23 g, 3.6 mmol, 1.45 mL of a 2.5M solution) was added dropwise via syringe keeping the temperature under -55° C. After 0.5 hours, trimethylborate (0.40 g, 0.38 mmol) was added via syringe and the reaction was allowed to warm to room temperature. 1N HCl (3.5 mL) was added and the mixture was stirred for 0.5 hours. The mixture was diluted with water and ether and the ether layer was separated and dried over Na₂SO₄. Filtration and concentration left 0.753 g of a foam which was triturated with hexanes. The resulting solids were collected by filtration and dried to give 4-chloro-3-(dimethylamino)-2-fluorophenylboronic acid (0.5 g, 2.3 mmol) as an off white solid: ¹H NMR (DMSO-*d*₆) revealed the solid to be a mixture of what appears to be the boronic acid and anhydrides. The solid was used as is without further purification or characterization.

The following compounds were prepared according to the procedure in Example 40.

4-Chloro-3-(diethylamino)-2-fluorophenylboronic acid: ¹H NMR (300 MHz, DMSO *d*₆ + D₂O): δ 8.26 (bs, 2H), 7.26 (m, 2H), 3.05 (q, 4H), 0.91 (t, 6H)

2,4-Dichloro-3-dimethylaminophenylboronic acid. This material was isolated as an oily mixture of boronic acid and anhydride and used without further purification or characterization.

2-Chloro-4-fluoro-3-methoxyphenylboronic acid (used without purification)

73776-249(S)

41. Preparation of 6-chloro-2-fluoro-3-(4,4,5,5-tetramethyl-
{1,2,3}dioxaborolan-2-yl)-phenyl methyl amine.

N-(3-bromo-6-chloro-2-fluorophenyl)-*N*-methylaniline (0.84 g, 3.5
5 mmol) was dissolved in DMF (10 mL) in a nitrogen flushed 50 mL 3-neck flask
and the resulting solution was purged with nitrogen for 0.5 hour.
Bis(pinacolato)diboron (0.98 g, 3.9 mmol), potassium acetate (1.04 g, 10.6 mmol)
and [1,1'-bis(diphenylphosphino) ferrocene]dichloropalladium(II) · CH₂Cl₂
complex (0.14 g, 0.18 mmol) were added all at once and the mixture heated to
10 110° C with a nitrogen purge of the head space through 60° C and then under an
open nitrogen atmosphere for 1.5 hours. The reaction mixture was cooled to
room temperature and partitioned between ethyl acetate and brine and the layers
separated. The aqueous phase was extracted with ethyl acetate and the combined
organics washed with brine and dried (Na₂SO₄). Filtration, concentration and
15 purification by flash silica gel chromatography (hexanes:ethyl acetate / 5:1) gave
[6-chloro-2-fluoro-3-(4,4,5,5-tetramethyl-{1,3,2}dioxaborolan-2-yl)-phenyl
methyl amine (0.85 g): light green solid: ¹H NMR (300 MHz, CDCl₃): δ 7.05 (m,
2H), 3.96 (bs, 1H), 3.06 (d, 3H), 1.35 (s, 12 H).

42. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-thiomethyl)-
20 phenyl)pyridine-2-carboxylic acid methyl ester (Compound 1).

A solution of 4-amino-3,6-dichloropyridine-2-carboxylic acid
methyl ester (2.6 g, 0.012 mol), 2-(4-chloro-2-fluoro-3-thiomethylphenyl)-[1,3,2]-
dioxaborinane (4.1 g, 0.016 mol), 1,4-bis(diphenylphosphino)butane (0.610 g,
0.0015 mol) and cesium fluoride (6.7 g, 0.045 mol) in acetonitrile (75 mL) were
25 degassed with nitrogen for 15 minutes. Palladium acetate (0.30 g, 0.0015 mol)
was added and the mixture heated to reflux for 4 hours. The reaction mixture was
quenched with water and extracted with ethyl acetate (2 x 100 mL) and the
combined extracts were dried (sodium sulfate) and concentrated. The residual

solid was purified by chromatography (25 percent-40 percent ethyl acetate in hexanes) to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-thiomethylphenyl)pyridine-2-carboxylic acid methyl ester (2.0g, 0.005mol): ¹H NMR (DMSO-*d*₆): δ 7.78 (t, 1H), 7.50 (d, 1H), 7.25 (s, 1H), 3.95 (s, 3H), 2.50 (s, 3H).

The following compounds were prepared according to the procedure of Example 42 (in some cases the crude boronic acid was used instead of the ester).

4-Amino-3-chloro-6-(4-chloro-3-cyano-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester (Compound 2): mp 189-190° C.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 3): mp 116-117° C.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 4): mp 145-146° C.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 5): ¹H NMR (CDCl₃): δ 8.05 (dt, 1H), 7.25 (d, 1H), 7.15 (s, 1H), 5.65 (dd, 2H), 4.95 (bs, 2H), 4.00 (s, 3H).

4-Amino-3-chloro-6-(4-cyano-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 6): ¹H NMR (CDCl₃): δ 7.70 (m, 1H), 7.20 (dd, 1H), 7.10 (d, 1H), 4.95 (bs, 2H), 4.00 (s, 3H).

4-Amino-3-chloro-6-(2,4-dichloro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 7): mp 132-133° C.

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl]pyridine-2-carboxylic acid methyl ester (Compound 8): mp 134-136° C.

- 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 9): mp 115-116° C.
- 4-Amino-3-chloro-6-[2,4-dichloro-3-(2,2-difluoroethoxy)phenyl]pyridine-2-carboxylic acid methyl ester (Compound 10): mp 113-115° C.
- 5 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methoxyethoxy)phenyl]pyridine-2-carboxylic acid methyl ester (Compound 11): mp 89-90° C.
- 4-Amino-3-chloro-5-fluoro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 12): mp 151-152° C.
- 10 4-Amino-3-chloro-6-(2,4-dichloro-3-methylthiophenyl)pyridine-2-carboxylic acid methyl ester (Compound 13): ¹H NMR (CDCl₃): δ 7.45 (d, 1H), 7.40 (d, 1H), 6.98 (s, 1H), 4.97 (bs, 2H), 3.98 (s, 3H), 2.44 (s, 3H).
- 4-Amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 14): mp 172-173° C.
- 15 4-Amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 15): mp 168-169° C.
- 4-Amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 16): mp 104-106 °C.
- 4-Amino-3-chloro-6-[4-chloro-3-(diethylamino)-2-fluorophenyl]pyridine-2-carboxylic acid, methyl ester (Compound 17): mp 120-121 °C.
- 20 43. Preparation of 4-acetylamino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 18).

A solution of 4-acetylamino-3,6-dichloropyridine carboxylic acid methyl ester (4.3 g, 0.016 mol), 2-(4-chloro-2-fluoro-5-methoxyphenyl)-[1,3,2]-

73776-249(S)

dioxaborinane (4.5 g, 0.018 mol), cesium fluoride (3.5 g, 0.025 mol), and 1,4-bis(diphenylphosphino)butane (0.360 g, .0016 mol) in acetonitrile (100 mL) was degassed with nitrogen for 15 minutes before palladium acetate (0.180 g, 0.0016 mol) was added and the solution heated under reflux for 18 hours. Water (150
 5 mL) was added and the resulting solid collected and dried to give 4-acetylamino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (4.5 g, 0.012 mol): mp 180-182° C.

The following compounds were prepared according to the procedure of Example 43.

10 4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 19): mp 151-152° C.

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 20): mp 159-160° C.

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-
 15 carboxylic acid methyl ester (Compound 21): ¹NMR (CDCl₃): δ 9.0 (bs, 1H) 8.00 (m, 1H), 7.60 (m, 1H), 7.25 (s, 1H), 4.25 (q, 2H), 4.00 (s, 3H), 2.35 (s, 3H), 1.50 (t, 3H).

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 22): ¹NMR (CDCl₃): δ 8.90 (s, 1H), 8.05 (s, 1H),
 20 7.40 (d, 1H), 7.30 (d, 1H), 4.05 (s, 3H), 3.95 (s, 3H), 2.35 (s, 3H).

4-Acetylamino-3-chloro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester: mp 132-134° C.

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester: ¹H NMR (CDCl₃): δ 8.86 (s, 1H), 8.00 (bs, 1H),
 25 7.37 (d, 1H), 7.26 (d, 1H), 4.11 (q, 2H), 4.00 (s, 3H), 2.32 (s, 3H), 1.47 (t, 3H).

4-Acetylamino-3-chloro-6-[2,4-dichloro-3-(2,2-difluoroethoxy)phenyl]pyridine-2-carboxylic acid methyl ester: ^1H NMR (CDCl_3): δ 8.81 (s, 1H), 8.10 (bs, 1H), 7.32 (d, 1H), 7.29 (d, 1H), 6.16 (tt, 1H), 4.22 (td, 2H), 3.95 (s, 3H), 2.28 (s, 3H).

44. Preparation of 4-acetylamino-3-chloro-6-(2,4-difluoro-3-methylphenyl)-pyridine-2-carboxylic acid methyl ester (Compound 23).

A solution of 1-bromo-2,4-difluoro-3-methylbenzene (2.10 g, 0.09 mol), 4-acetylamino-3-chloro-6-trimethylstannylpyridine-2-carboxylic acid methyl ester (4.00 g, 0.01 mol), 1,4-bis(diphenylphosphino)butane (0.140 g, 0.0003 mol) and cesium fluoride (4.5 g, 0.03 mol) in acetonitrile (100 mL) was degassed with nitrogen. Palladium acetate (0.70 g, 0.0003 mol) was added the solution heated to reflux for 2 hours. The reaction mixture was filtered through celite and the filtrate diluted with water (200 mL) and then extracted with ethyl acetate (2 x 150 mL). The combined organic phases were dried and concentrated and the residual solid was purified by chromatography (50 percent ethyl acetate in hexanes) to give 4-acetylamino-3-chloro-6-(2,4-difluoro-3-methylphenyl)-pyridine-2-carboxylic acid methyl ester (0.800 g, 0.002 mol): mp 152-153° C.

The following compound was prepared according to the procedure of Example 44.

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 24): mp 172-174° C.

45. Preparation of (2-fluoro-3-methoxy-4-trifluoromethylphenyl)trimethylstannane.

Dioxane (28 mL) was added to an oven dried, nitrogen swept 100 mL 3-neck flask equipped with a condenser and a magnetic stirbar and was de-aerated with a purge of nitrogen for 15 minutes. Hexamethylditin (5.0 g, 15.3 mmol), and 1-bromo-2-fluoro-3-methoxy-4-(trifluoromethyl)benzene (3.8 g, 13.9

73776-249(S)

- mmol) were added and the purging continued for 5 minutes. 1,4-bis(diphenylphosphino)butane (0.6 g, 1.4 mmol) and palladium acetate (0.3 g, 1.4 mmol) were added and the head space of the reaction vessel purged with nitrogen while heating to reflux with stirring. The reaction was then stirred at reflux under an open nitrogen atmosphere for 6 hours. TLC (hexanes) indicated complete consumption of starting material. The cooled reaction mixture was concentrated and purified by flash silica gel chromatography (hexanes) to give (2-fluoro-3-methoxy-4-trifluoromethylphenyl)trimethyl stannane (3.93 g, 11.0 mmol) as an oil: ¹H NMR (CDCl₃): δ 7.31 (d, 1H), 7.11 (dd, 1H), 3.98 (d, 3H), 0.38 (s, 9H). All signals had appropriate tin satellites.

The following compounds were prepared according to the procedure in Example 45.

15. (3-Butoxy-4-chloro-2-fluorophenyl)-trimethyl stannane (used without purification).
- [4-Chloro-2-fluoro-3-(1-fluoro-1-methyl-ethyl)phenyl]-trimethyl stannane (used without purification).
- 20 46. Preparation of 4-acetylamino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid methyl ester.

- In an oven dried, nitrogen flushed 100 mL 3 neck flask equipped with a condenser and a magnetic stirbar was dissolved methyl-4-acetamido-3,6-dichloropyridine-2-carboxylate (1.2 g, 4.6 mmol) and (2-fluoro-3-methoxy-4-trifluoromethylphenyl)trimethyl stannane (1.7 g, 4.6 mmol) in acetonitrile (15 mL) and the resulting solution was de-aerated with a purge of nitrogen for 15 minutes. Cesium fluoride (2.1 g, 13.9 mmol), 1,4-bis(diphenylphosphino)butane (0.2 g, 0.46 mmol) and palladium acetate (0.1 g, 0.46 mmol) were added and the

head space of the reaction was purged with nitrogen as the reaction was heated to reflux with stirring. The reaction was then stirred under an open nitrogen atmosphere for 5 hours by which time TLC (hexane:ethyl acetate/1:2) indicated complete consumption of starting material. The cooled reaction was filtered
5 through celite with ethyl acetate and the filtrate concentrated and purified by flash silica gel chromatography (hexanes:ethyl acetate/2:1) to give 4-acetylamino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid methyl ester (0.19 g, 0.46 mmol) as an oil: ¹H NMR (CDCl₃): δ 9.03 (d, 1H), 8.02 (bs, 1H), 7.68 (bt, 1H), 7.42 (bd, 1H), 4.03 (d, 3H), 4.01 (s, 3H), 2.33 (s,
10 3H).

The following compounds were prepared according to the procedure in Example 46.

4-Acetylamino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester: LC/MS ESI (m/z=428).

15 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoro-1-methylethyl)phenyl]-pyridine-2-carboxylic acid, methyl ester (Compound 25): mp 122-123° C.

47. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-5-nitrophenyl)-pyridine-2-carboxylic acid methyl ester.

Methyl 4-amino-3-chloro-6-(4-chloro-2-fluorophenyl)pyridine-2-
20 carboxylate (1.0 g, 3.2 mmol) was added as a solid to well stirred concentrated sulfuric acid (16 mL) at 0 °C. Sodium nitrate (0.29 g, 3.5 mmol) was added to the mixture and the resulting yellow solution was stirred at 0° C for 1.5 hours then diluted with a large quantity of ice and allowed to warm to room temperature. The solution was filtered to remove some solids then the filtrate extracted with 3
25 portions of ethyl acetate and the combined ethyl acetate layers washed with saturated aqueous NaHCO₃ and dried (Na₂SO₄). Filtration and concentration gave

4-amino-3-chloro-6-(4-chloro-2-fluoro-5-nitrophenyl)pyridine-2-carboxylic acid methyl ester (0.96 g, 2.7 mmol) as a yellow solid: mp 161-163° C: ¹H NMR (CDCl₃): δ 8.73 (d, 1H), 7.36 (d, 1H), 7.24 (d, 1H), 4.93 (bs, 2H), 4.02 (s, 3H).

48. Preparation of 4-amino-3-chloro-6-(5-amino-4-chloro-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester.

Iron powder (0.78 g, 13.9 mmol) was added to a slurry of methyl 4-amino-3-chloro-6-(4-chloro-2-fluoro-5-nitrophenyl)pyridine-2-carboxylate (0.5 g, 1.39 mmol) in acetic acid (20 mL) in a 100 mL round bottomed flask. The mixture was heated to 85° C for 15 minutes. then cooled to room temperature and filtered through celite with ethanol and the filtrate concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated NaHCO₃, the layers separated, the aqueous phase extracted with ethyl acetate and the combined organics dried (Na₂SO₄). Filtration and concentration gave 4-amino-3-chloro-6-(5-amino-4-chloro-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester (0.44 g, 1.33 mmol) as a light brown solid: mp 99-101° C.

49. Preparation of 4-amino-3-chloro-6-(4-chloro-5-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester (Compound 26).

Methyl 4-amino-3-chloro-6-(5-amino-4-chloro-2-fluorophenyl)pyridine-2-carboxylate (0.38 g, 1.13 mmol)) was dissolved in THF (10 mL) in a nitrogen flushed 250 mL round bottomed flask and treated with 37 percent formaldehyde (0.068 g, 0.17 mL, 2.27 mmol), dibutyl tin dichloride (0.007 g, 0.023 mmol) and phenyl silane (0.135 g, 1.25 mmol) and allowed to stir at room temperature for 72 hours. The reaction mixture was concentrated *in vacuo* and purified by flash silica gel chromatography (hexanes:ethyl acetate/2:1) to give 4-amino-3-chloro-6-(4-chloro-5-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic methyl ester (0.283 g 0.8 mmol) as a soft white solid: mp 113-115° C,

¹H NMR (CDCl₃): δ 7.66 (d, 1H), 7.14 (d, 1H), 7.14 (d, 1H). 4.91 (bs, 2H), 3.99 (s, 3H), 2.79 (s, 6H) .

50. Preparation of 4-acetylamino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester.

5 To a solution of 4-acetylamino-3-chloro-6-trimethylstannanylp¹pyridine-2-carboxylic acid, methyl ester (21.0g, 54mmol) in methylene chloride (200mL) was added iodine (13.7g, 108mmol) and the solution was stirred 1 hour at ambient temperature. The solution was filtered through a bed of celite and then washed with sodium thiosulfate (50mL, 10 percent solution), dried
10 (sodium sulfate) and concentrated. The residue was purified by chromatography (10 percent ethyl acetate in hexanes) to give 4-acetylamino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester (5.6g, 18mmol): GC-MS (m/z=354).

15 51. Preparation of 4-amino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester.

To solution of 4-acetylamino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester (5.0g, 14mmol) in methanol (25 mL) was added acetyl chloride (1 mL) and the solution heated under reflux for 1 hour. The
20 solution was cooled and water (25mL) added and the precipitate was collected. Recrystallization from methanol gave 4-amino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester (3.5 g, 11mmol) mp 152-153° C.

52. Preparation of 4-amino-6-(4-bromo-2-fluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid methyl ester (Compound 27).

25 A solution of 4-amino-3-chloro-6-iodopyridine-2-carboxylic acid, methyl ester (1.1 g, 3.0 mmol), 4-bromo-2-fluoro-3-methoxyphenyl boronic acid (1.3 g, 4.5 mmol) and cesium fluoride (0.60 g, 4.0 mmole) in dimethoxyethane (2 mL) and water (2 mL) was de-aerated with a stream of nitrogen for 15 minuets before adding dichlorobis(triphenylphosphine) palladium (0.25 g, 0.4 mmol) and

then heated to 85° C for 2 hours. The reaction mixture was diluted with water (15 mL) and extracted with ethyl acetate (2 x 15 mL). The combined extracts were washed with brine, dried (sodium sulfate) and concentrated. The residue was purified by chromatography (33 percent ethyl acetate in hexanes) to give 4-acetylamino-6-(4-bromo-2-fluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (1.2 g, 2.8 mmol): mp 164-165° C.

The following compounds were prepared according to the procedure in Example 52.

4-Acetylamino-3-chloro-6-(2,4,6-trifluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without further purification).

4-Acetylamino-3-chloro-(2,4-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 28): mp 114-115° C.

4-Acetylamino-3-chloro-(2-chloro-4-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 29): mp 153-154.5° C.

53. Preparation of 4-acetylamino-3-chloro-6-[2,4-dichloro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid methyl ester.

A solution of 2,4 dichloro-3-(1-fluoroethyl)-bromobenzene (3.4 g, 12.5 mmol), 4-acetylamino-3-chloro-6-trimethylstannyl-pyridine-2-carboxylic acid, methyl ester (4.7 g, 12.5 mmol), copper iodide (0.4 g, 2.1 mmol), cesium fluoride (3.6 g, 25 mmol) and dichlorobis(triphenylphosphine)palladium (0.14 g, 2 mmol) in DMF (50 mL) was heated to 70° C for 2 hours. Water (100 mL) was added the mixture extracted with ethyl acetate (2 x 100 mL). The combined extracts were washed with brine, dried (sodium sulfate) and concentrated. The residue was purified by chromatography (15 percent-30 percent ethyl acetate in hexanes) to give 4-acetylamino-3-chloro-6-[2,4 dichloro-3-(1-

73776-249(S)

fluoroethyl)phenyl]pyridine-2-carboxylic acid, methyl ester which was used without further purification.

The following compounds were made according to the procedure of Example 53.

- 5 4-Acetylamino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester: mp 134-136°C.
- 4-Acetylamino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester: mp 146-147°C.
- 4-Acetylamino-3-chloro-6-(4-chloro-3-dimethylamino-2,6-difluorophenyl)pyridine-2-carboxylic acid, methyl ester (used without further purification).
- 10 4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without further purification).
- 4-Acetylamino-3-chloro-6-(2,4-dichloro-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without further purification).
- 15 4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester: (used without further purification).
- 4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, methyl ester (used without further purification).
- 4-Acetylamino-3-chloro-6-[2,4-dichloro-3-(difluoromethyl)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 30): mp 152-153°C.
- 20 4-Acetylamino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester: GC/PCI (m/z=236).

4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)-phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 31): LC/MS ESI ($m/z=456$).

5 4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)-phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 32): LC/MS ESI ($m/z=416$).

4-Acetylamino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 33): mp 169-171° C.

10 4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxy-phenyl)pyridine-2-carboxylic acid, methyl ester (Compound 34) mp 122-123° C.

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethoxyphenyl)pyridine, carboxylic acid, methyl ester (Compound 35): mp 146° C.

4-Acetylamino-3-chloro-6-(4-difluoromethyl-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

15 4-Acetylamino-3-chloro-6-(2-fluoro-4-fluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methylphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

20 4-Acetylamino-6-(2-bromo-4-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-6-(4-bromo-2,6-difluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-(3-difluoromethyl-2,4,6-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-(2-chloro-4-difluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

- 5 4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-(2-chloro-4,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

- 10 4-Acetylamino-3-chloro-6-(2,4-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-(4-chloro-3-cyclopropyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester (used without purification).

4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2,2-trifluoroacetyl)phenyl]pyridine-2-carboxylic acid, methyl ester (used without purification).

- 15 54. Preparation of 4-amino-3-chloro-6-(2-fluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 36).

- 20 A solution of 4-amino-3,6-dichloropyridine-2-carboxylic acid, methyl ester (0.24 g, 1.1 mmol), 2-fluoro-3-methoxy-4-methylphenyl boronic acid (0.30 g, 1.63 mmol) and cesium fluoride (3.0 g, 3.26 mmole) in dimethoxyethane (2 mL) and water (2 mL) was purged with a stream of nitrogen for 15 minutes before adding dichlorobis(triphenylphosphine) palladium (0.07 g, 0.1 mmol) and then heated to 85° C for 2 hours. The reaction mixture was diluted with water (15 mL) and extracted with ethyl acetate (2 x 15 mL). The combined extracts were washed with brine, dried (sodium sulfate) and concentrated. The residue

was purified by chromatography (33 percent ethyl acetate in hexanes) to give 4-amino-3-chloro-6-(2-fluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester (0.08 g, 0.2 mmol): mp 95-96° C.

The following compounds were prepared according the procedure
5 in Example 52.

4-Amino-3-chloro-6-[2,4-dichloro-3-(dimethylamino)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 37): ¹H NMR (300 MHz, CDCl₃): δ 7.30 (d, 1H), 7.19 (d, 1H), 6.93 (s, 1H) 4.88 (bs, 2H), 3.96 (s, 3H), 2.88 (s, 6H).

10 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methylamino)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 38): mp 148-149° C.

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methoxycarbonyl) phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 39): ¹H NMR (300 MHz, CDCl₃): δ 7.53
15 (d, 1H), 7.37 (d, 1H), 6.94 (s, 1H), 4.97 (bs, 2H), 3.97 (s, 3H), 3.96 (s, 3H).

4-Acetylamino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester: mp 146-147° C.

20 4-Acetylamino-3-chloro-6-[4-chloro-3-(dimethylamino)-2,6-difluoro-phenyl]pyridine-2-carboxylic acid, methyl ester: ¹H NMR (300 MHz, CDCl₃): δ 8.70 (s, 1H), 8.03 (bs, 1H), 7.03 (dd, 1H), 4.00 (s, 3H), 2.83 (d, 6H), 2.32 (s, 3H).

25 4-Acetylamino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester: mp 134-136° C.

4-Acetylamino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester: mp 129-130° C.

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 40): mp 144-144.5° C.

55. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)-
5 pyridine-2-carboxylic acid methyl ester (Compound 41).

To a solution of 4-acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester (0.5 g, 0.0013 mol) in methanol (10 mL) was added acetyl chloride (1 mL) and the solution heated under reflux for 1 hour. Water (2 mL) was added and resulting solid collected
10 and dried to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester: mp 154-156° C.

The following compounds were prepared according to the procedure of Example 55.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic
15 acid methyl ester (Compound 42): mp 141-142° C.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 43): ¹NMR (CDCl₃): δ 7.60 (d, 1H), 7.25 (s, 1H), 7.20 (m, 1H), 4.95 (bs, 2H), 4.05 (s, 3H), 3.95 (s, 3H).

4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid
20 methyl ester (Compound 44): mp 168-170° C.

4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 45): mp 144-146° C.

4-Amino-3-chloro-6-(2-fluoro-3,4 methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester (Compound 46): mp 139-141° C.

4-Amino-3-chloro-6-(2,4-difluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 47): mp 129-130° C.

4-Amino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 48): ¹H NMR (CDCl₃): 7.30 (d, 1H), 7.25 (d, 1H), 6.95 (s, 1H).

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 49): mp 116-118° C.

4-Amino-3-chloro-6-[4-chloro-3-(dimethylamino)-2,6-difluorophenyl]pyridine-2-carboxylic acid, methyl ester (Compound 50): ¹H NMR (300 MHz, CDCl₃): δ 7.00 (dd, 1H), 6.81 (s, 1H) 4.93 (bs, 2H), 3.96 (s, 3H), 2.81 (d, 6H).

4-Amino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 51): Off white foam, ¹H NMR (300 MHz, CDCl₃): δ 6.83 (s, 1H), 6.74 (dd, 1H), 4.88 (bs, 2H), 3.97 (s, 3H), 3.84 (d, 3H), 2.28 (s, 3H).

4-Amino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (Compound 52): mp 154-155 °C.

4-Amino-3-chloro-(2-chloro-4-difluoromethyl-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 53): LC/MS (m/z=380).

4-Amino-3-chloro-6-(4-chloro-3-difluoromethyl-2,6-difluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 54): LC/MS (m/z=383).

- 4-Amino-3-chloro-(2-chloro-4,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 55): LC/MS ($m/z=348$).
- 4-Amino-3-chloro-6-(2,4,6-trifluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 56): HPLC-MS ($m/z=346$).
- 5 4-Amino-3-chloro-6-(2,4-dichloro-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 57): HPLC-MS ($m/z=378$).
- 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 58): HPLC-MS ($m/z=362$).
- 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 59): HPLC-MS ($m/z=376$).
- 10 4-Amino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 60): LC/MS ESI ($m/z=386$).
- 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 61): LC/MS ESI ($m/z=360$).
- 15 4-Amino-3-chloro-6-(2,4-dichloro-3-difluoromethylphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 62): LC/MS ESI ($m/z=380$).
- 4-Amino-3-chloro-6-[2,4-dichloro-3-(1-fluoro-1-ethylphenyl)]pyridine-2-carboxylic acid, methyl ester (Compound 63): LC/MS ESI ($m/z=362$).
- 4-Amino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 64): mp 120-122° C.
- 20 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)phenyl]-pyridine-2-carboxylic acid, methyl ester (Compound 65): LC/MS ESI ($m/z=414$).

- 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 66): ^1H NMR (300 MHz, CDCl_3): δ 7.90 (m, 1H), 7.25 (m, 1H), 7.10 (s, 1H), 5.96 (dq, 1H), 4.90 (bs, 2H), 4.05 (s, 3H), 2.30 (m, 1H), 2.05 (m, 1H), 1.05 (t, 3H).
- 5 4-Amino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 67): mp 163-164° C.
- 10 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 68): mp 99° C.
- 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 69): mp 164° C
- 15 4-Amino-3-chloro-6-(2,4-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 70): LC/MS (m/z = 328).
- 4-Amino-3-chloro-6-(4-difluoromethyl-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 71): mp 145-147° C.
- 20 4-Amino-3-chloro-6-(2-fluoro-4-fluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 72): LC/MS (m/z =342).
- 4-Amino-3-chloro-6-(2,4-difluoro-5-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 73): mp 128-130° C.
- 25 4-Amino-3-chloro-6-(2-chloro-4-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 74): mp 145-146° C.

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methylphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 75): LC/MS ($m/z=346$).

5 4-Amino-6-(2-bromo-4-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (Compound 76): LC/MS ($m/z=405$).

4-Amino-6-(4-bromo-2,6-difluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester (Compound 77): LC/MS ($m/z=407$).

10 4-Amino-3-chloro-6-(3-difluoromethyl-2,4,6-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 78): LC/MS ($m/z=366$).

4-Amino-3-chloro-6-(2-chloro-4-difluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 79): mp 134-136° C.

15 4-Amino-3-chloro-6-(2,4-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 80): mp 158-159° C.

20 4-Amino-3-chloro-6-(4-chloro-3-cyclopropyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester (Compound 81): mp 135-136.5° C.

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2,2-trifluoroacetyl)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 82): LC/MS ($m/z=410$).

56. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfinylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 83) and 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfonylphenyl)pyridine-2-carboxylic acid methyl ester (Compound 84).

5 To a solution of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methylthiophenyl)pyridine-2-carboxylic acid methyl ester (0.800 g, 2 mmol) in methylene chloride (20 mL) was slowly added peracetic acid (32 percent in acetic acid, 0.004 mol). The reaction mixture was quenched with sodium bisulfite (10 percent in water, 20 mL) and extracted with methylene chloride (20 mL). The
10 organic phase was dried (sodium sulfate) and concentrated. The solid residue was purified by chromatography (15 percent-50 percent ethyl acetate in hexanes). The first fraction was 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfonylphenyl)pyridine-2-carboxylic acid methyl ester (0.250 g, 0.7 mmole): mp 108-110° C. The second fraction was 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfinylphenyl)pyridine-2-carboxylic acid methyl ester (0.100
15 g, 0.03 mmole): mp 122-124° C.

57. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethylthiophenyl)pyridine-2-carboxylic acid methyl ester (Compound 85).

To a solution of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methylthiophenyl)pyridine-2-carboxylic acid, methyl ester (1.50 g, 4.1 mmol) in
20 acetonitrile (40 mL) was added 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane-bis(tetrafluoroborate) (F-TEDA; SELECTFLUORTM fluorinating agent; 1.48 g, 8.2 mmol) and the solution was stirred at ambient temperature for 2 hours. A saturated solution of sodium
25 bicarbonate (25 mL) was added the solution extracted with methylene chloride (2 x 50 mL). The combined extracts were dried (sodium sulfate) and concentrated and the residue purified by chromatography (20 percent-50 percent ethyl acetate

in hexanes) to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethyl-thiophenyl)pyridine-2-carboxylic acid methyl ester (0.09 g, 0.2 mmol): mp 104-105° C.

58. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-pyridine-2-carboxylic acid, ethyl ester (Compound 86).

To a solution of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (200 mg, 0.9 mmol) in ethanol (5 mL) was added a catalytic amount of titanium tetrakisopropoxide and the solution heated under reflux for 2 hours. The solution was cooled and partitioned between ethyl acetate (10 mL) and water (10 mL) and the organic phase was dried (sodium sulfate) and concentrated. The residue was passed over a plug of silica gel (1:1 ethyl acetate: hexanes) to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, ethyl ester (150 mg, 0.6 mmol): mp 74-75° C.

The following compounds were prepared according to the procedure in Example 58.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, 2-butoxy-ethyl ester (Compound 87): LC/MS (m/z=430).

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester (Compound 88): LC/MS (m/z=404).

4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester (Compound 89): LC/MS ESI (m/z=402).

4-Amino-3-chloro-6-(4-bromo-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester (Compound 90): LC/MS ESI (m/z=430).

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, butyl ester (Compound 91): LC/MS ESI ($m/z=402$).

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, 2-butoxy-ethyl ester (Compound 92): mp 72-74° C.

59. Preparation of 4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid.

A slurry of 4-acetylamino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine (0.13 g, 0.33 mmol) in methanol (5 mL) was treated with 1N sodium hydroxide (1.7 mL) and stirred at ambient temperature for 1 hour. The solution was diluted with water (15 mL) and acidified with 1N hydrochloric acid (1.7 mL) and the solid collected (0.095 g). Analysis by ¹H NMR showed the 4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid and an impurity. This material was used without further purification.

The following compounds were prepared according to the procedure of example 59.

4-Amino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid (Compound 93): mp 167-168° C.

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-dimethylaminophenyl)pyridine-2-carboxylic acid (Compound 94): mp 171-172° C.

60. Preparation of 4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (Compound 95).

A slurry of 4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid in tetrahydrofuran (10 mL) and

methanol (3 mL) was treated with trimethylsilyldiazomethane (0.64 g, 0.28 mL of a 2M solution in hexanes) at ambient temperature. After 1 hour the reaction mixture was quenched with acetic acid (2 mL) and water (10 mL). The mixture was extracted with ethyl acetate (2 x 20 mL) and the combined extracts dried (Na₂SO₄), and concentrated. The residue was purified by chromatography (30-100 percent ethyl acetate in hexanes) to give 4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester (0.083 g, 0.24 mmol): ¹H NMR (CDCl₃) δ 7.15 (dd, 1H), 6.85 (s, 1H), 4.98 (bs, 2H), 4.08 (d, 3h), 3.98 (s, 3H).

10 61. Preparation of 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl)pyridine-2-carboxylic acid (Compound 96).

A solution of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl)pyridine-2-carboxylic acid methyl ester (0.300 g, 0.0008 mol) in methanol (5 mL) and sodium hydroxide (1N, 2 mL) was heated to reflux 1 hour and then acidified to pH 3 (concentrated hydrochloric acid) and allowed to cool. The resulting solid was collected and dried to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl)pyridine-2-carboxylic acid (0.270 g, 0.0007 mol): mp 183-184° C dec.

The following compounds were prepared according to the procedure of Example 61.

4-Amino-3-chloro-(2,4-difluoro-3-methylphenyl)pyridine-2-carboxylic acid (Compound 97): mp 189-190° C dec.

4-Amino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid (Compound 98): mp 170-172° C dec.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid (Compound 99): mp 179-181° C dec.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid (Compound 100): mp 174-175° C dec.

- 5 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid (Compound 101): ¹NMR (DMSO-*d*₆): δ 13.80 (bs, 1H) 7.55 (m, 2H), 7.25 (s, 1H), 3.95 (s, 3H).

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid (Compound 102): mp 177-178° C dec.

- 10 4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid (Compound 103): mp 179-180° C dec.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylthiophenyl)pyridine-2-carboxylic acid (Compound 104): ¹NMR (DMSO-*d*₆) δ 13.80 (bs, 1H), 7.90 (t, 1H), 7.55 (d, 1H), 7.10 (s, 1H), 6.95 (bs, 2H), 2.50 (s, 3H).

- 15 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethylphenyl)pyridine-2-carboxylic acid (Compound 105): mp 187-188° C dec.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid (Compound 106): mp 182-183° C dec.

- 20 4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid (Compound 107): mp 177-178° C dec.

4-Amino-3-chloro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid (Compound 108): mp 202-203° C dec.

- 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-ethoxyphenyl)pyridine-2-carboxylic acid (Compound 109): mp 178-179° C dec.
- 4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid (Compound 110): mp 187-188° C dec.
- 5 4-Amino-3-chloro-6-(4-chloro-5-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid (Compound 111): mp 180-181° C dec.
- 4-Amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid (Compound 112): mp 185-186° C dec.
- 10 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyethoxyphenyl)pyridine-2-carboxylic acid (Compound 113): mp 169-170° C dec.
- 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethylphenyl)pyridine-2-carboxylic acid (Compound 114): ¹H NMR (DMSO-*d*₆): 8.00 (dt, 1H) 7.50 (dd, 1H), 6.90 (bs, 2H), 5.60 (dd, 1H).
- 15 4-Amino-3-chloro-5-fluoro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid (Compound 115): mp 179-180° C.
- 4-Amino-3-chloro-6-[4-chloro-3-(diethylamino)-2-fluorophenyl]pyridine-2-carboxylic acid (Compound 116): mp 174-175° C
- 4-Amino-3-chloro-6-[2,4-dichloro-3-(dimethylamino)phenyl]pyridine-2-carboxylic acid (Compound 117): mp 177-178° C.
- 20 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methylamino)phenyl]pyridine-2-carboxylic acid (Compound 118): mp 185-186° C.
- 4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid (Compound 119): mp 178-179° C.

73776-249(S)

4-Amino-6-(4-bromo-2-fluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid (Compound 120): mp 176-177° C.

4-Amino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid (Compound 121): mp 192-193° C.

5 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid (Compound 122): mp 180-181° C.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid (Compound 123): mp 178-179° C.

10 4-Amino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid (Compound 124): mp 157-158° C dec.

4-Amino-3-chloro-6-[2,4 dichloro-3-(1-fluorothyl)phenyl]pyridine-2-carboxylic acid (Compound 125): mp 169-170°C dec.

4-Amino-3-chloro-6-(2,4-dichloro-3-difluoromethylphenyl)pyridine-2-carboxylic acid (Compound 126): mp 182-183° C dec.

15 4-Amino-3-chloro-6-[2,4-dichloro-3-(1-fluoro-1-methylethylphenyl)pyridine-2-carboxylic acid (Compound 127): mp >250 °C .

4-Amino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid (Compound 128): mp 192-193° C dec.

20 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)-phenyl]pyridine-2-carboxylic acid (Compound 129): mp 175-176° C dec.

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)phenyl]pyridine-2-carboxylic acid, methyl ester (Compound 130): mp 165-167° C dec.

4-Amino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid (Compound 131): mp. 178-179° C dec.

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxyphenyl)pyridine-2-carboxylic acid (Compound 132): mp. 159° C dec.

5 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid (Compound 133): LC/MS (m/z=293).

4-Amino-3-chloro-6-(4-chloro-3-difluoromethoxy-2-fluorophenyl)pyridine-2-carboxylic acid (Compound 134): mp 179° C dec.

62. Preparation of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-
10 pyridine-2-carboxylic acid, triethyl amine salt (Compound 135).

To solution of 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-pyridine-2-carboxylic acid (1.0 g, 3.1 mmol) in methanol (30 mL) was added triethylamine (10 mL) and the solution stirred at ambient temperature for 1 hour and was then concentrated to give 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-
15 methoxyphenyl)pyridine-2-carboxylic acid, triethylamine salt (1.1 g, 2.6 mmol): ¹NMR (DMSO-*d*₆): δ 7.60 (m, 1H), 7.45 (m, 1H), 7.00 (s, 1H), 4.00 (s, 3H), 3.00 (t, 6H), 1.15 (q, 9H).

The following compound was prepared according to the procedure of Example 62.

20 4-Amino-3-chloro-6-(2,4 dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid, triethyl amine salt (Compound 136): ¹NMR (DMSO-*d*₆) δ 7.55 (d, 1H), 7.435 (d, 1H), 7.00 (s, 1H), 4.00 (s, 3H), 3.00 (t, 6H), 1.15 (q, 9H).

63. Preparation of Herbicidal Compositions

In the following illustrative compositions, parts and percentages
25 are by weight.

EMULSIFIABLE CONCENTRATES

Formulation A

	<u>WT%</u>
Compound 1	26.2
Polyglycol 26-3	5.2
Nonionic emulsifier-(di-sec-butyl)-phenyl-poly(oxypropylene)block polymer with (oxyethylene). The polyoxyethelene content is 12 moles.	
Witconate P12-20 (Anionic emulsifier-calcium dodecylbenzene sulfonate-60 wt. percent active)	5.2
Aromatic 100 (Xylene range aromatic solvent)	63.4

Formulation B

	<u>WT%</u>
Compound 12	3.5
Sunspray 11N (paraffin oil)	40.0
Polyglycol 26-3	19.0
Oleic acid	1.0
Xylene range aromatic solvent	36.5

Formulation C

	<u>WT%</u>
Compound 14	13.2
Stepon C-65	25.7
Ethomeen T/25	7.7
Ethomeen T/15	18.0
Xylene range aromatic solvent	35.4

Formulation D

	<u>WT%</u>
Compound 18	30.0
Agrimer Al-10LC (emulsifier)	3.0
N-methyl-2-pyrrolidone	67.0

5

Formulation E

	<u>WT%</u>
Compound 41	10.0
Agrimul 70-A (dispersant)	2.0
Amsul DMAP 60 (thickener)	2.0
Emulsogen M (emulsifier)	8.0
Attagel 50 (suspension aid)	2.0
Crop oil	76.0

These concentrates can be diluted with water to give emulsions of suitable concentrations for controlling weeds.

WETTABLE POWDERS

Formulation FWT%

Compound 95	26.0
Polyglycol 26-3	2.0
Polyfon H	4.0
Zeosyl 100 (Precipitated hydrated SiO ₂)	17.0
Barden clay + inerts	51.0

5

Formulation GWT%

Compound 101	62.4
Polyfon H (sodium salt of lignin sulfonate)	6.0
Sellogen HR (sodium naphthalene sulfonate)	4.0
Zeosyl 100	27.6

Formulation HWT%

Compound 109	1.4
Kunigel V1 (carrier)	30.0
Stepanol ME Dry (wetter)	2.0
Tosnanon GR 31A (binder)	2.0
Kaolin NK-300 Clay (filler)	64.6

The active ingredient is applied to the corresponding carriers and then these are mixed and ground to yield wettable powders of excellent wettability and suspension power. By diluting these wettable powders with water it is possible to obtain suspensions of suitable concentrations for controlling weeds.

WATER DISPERSIBLE GRANULES

Formulation I

	<u>WT%</u>
Compound 63	26.0
Sellogen HR	4.0
Polyfon H	5.0
Zeosyl 100	17.0
Kaolinite clay	48.0

The active ingredient is added to the hydrated silica, which is then mixed with the other ingredients and ground to a powder. The powder is agglomerated with water and sieved to provide granules in the range of -10 to +60 mesh. By dispersing these granules in water it is possible to obtain suspensions of suitable concentrations for controlling weeds.

GRANULES

Formulation J

	<u>WT%</u>
Compound 114	5.0
Celetom MP-88	95.0

The active ingredient is applied in a polar solvent such as *N*-methyl pyrrolidinone, cyclohexanone, *gamma*-butyrolactone, etc. to the Celetom

MP 88 carrier or to other suitable carriers. The resulting granules can be applied by hand, granule applicator, airplane, etc. in order to control weeds.

Formulation K

	<u>WT%</u>
Compound 117	1.0
Polyfon H	8.0
Nekal BA 77	2.0
Zinc Stearate	2.0
Barden Clay	87.0

5 All materials are blended and ground to a powder then water is added and the clay mixture is stirred until a paste is formed. The mixture is extruded through a die to provide granules of proper size.

10 Water Soluble Liquids

Formulation L

	<u>WT%</u>
Compound 136	3.67
Monoethanolamine pH buffer	0.5
Water	95.83

The active ingredient is dissolved in an appropriate amount of water and the additional monoethanolamine is added as a buffer. A water-soluble surfactant may be added. Other aids may be incorporated to improve physical, chemical and/or formulation properties.

15

64. Evaluation of Postemergence Herbicidal Activity

Seeds or nutlets of the desired test plant species were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of 30 percent, in plastic pots with a surface area of 64 square centimeters. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 7-21 days in a greenhouse with an approximate 15 hour photoperiod which was maintained at 23-29° C during the day and 22-28° C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the first or second true leaf stage.

A weighed amount, determined by the highest rate to be tested, of each test compound was placed in a 25 mL glass vial and was dissolved in 4 mL of a 97:3 v/v (volume/volume) mixture of acetone and dimethyl sulfoxide (DMSO) to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions obtained were diluted with 20 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Atplus 411F crop oil concentrate, and Triton[®] X-155 surfactant in a 48.5:39:10:1.5:1.0:0.02 v/v ratio to obtain spray solutions containing the highest application rates. Additional application rates were obtained by serial dilution of 12 mL of the high rate solution into a solution containing 2 mL of 97:3 (v/v) mixture of acetone and dimethyl sulfoxide (DMSO) and 10 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Atplus 411F crop oil concentrate, and Triton X-155 surfactant in a 48.5:39:10:1.5:1.0:0.02 (v/v) ratio to obtain 1/2X, 1/4X, 1/8X and 1/16X rates of the high rate. Compound requirements are based

upon a 12 mL application volume at a rate of 187 L/ha. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters at a spray height of 18 inches (43 cm) above the
 5 average plant canopy height. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 14 days, the condition of the test plants as compared with that
 10 of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

By applying the well-accepted probit analysis as described by J. Berkson in *Journal of the American Statistical Society*, 48, 565 (1953) and by D. Finney in "*Probit Analysis*" Cambridge University Press (1952), the above data
 15 can be used to calculate GR₅₀ and GR₈₀ values, which are defined as growth reduction factors that correspond to the effective dose of herbicide required to kill or control 50 percent or 80 percent, respectively, of a target plant.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 1.

20

Table 1. Post-emergent control of weeds.

Compound #	Rate (g ai/ha)	Percent Control		
		ABUTH	AMARE	CHEAL
1	140	85	100	100
2	140	85	90	80
3	140	100	100	80
4	140	100	100	85
5	140	100	35	90
6	140	90	100	95
7	140	100	90	95

8	140	80	100	95
9	140	85	100	100
10	140	90	100	100
11	140	90	95	80
12	140	100	100	100
13	140	100	100	90
14	140	100	100	100
15	140	100	100	100
16	140	100	100	100
17	140	100	100	100
25	140	100	100	100
26	140	100	100	100
27	140	100	90	100
28	140	100	100	95
29	140	95	40	98
30	140	100	90	95
32	140	60	80	50
33	140	95	95	90
36	140	100	100	100
37	140	100	100	100
38	140	100	100	100
39	140	90	75	95
40	140	95	95	100
41	140	95	85	98
42	140	100	100	100
43	140	100	100	100
44	140	100	100	100
45	140	100	80	95
46	140	100	100	95
47	140	100	100	100
48	101	98	80	85
49	140	100	80	90
50	140	100	100	100
51	140	100	100	100
52	140	100	100	95
53	140	70	80	95
54	140	100	100	100
55	140	100	100	90
56	140	100	100	100
57	140	100	100	100
58	140	100	100	100
59	140	100	90	90
60	140	55	90	90
61	140	100	100	100
62	140	100	100	100
63	140	100	100	100

64	140	95	100	100
65	140	95	100	90
66	140	90	100	95
67	140	100	100	100
68	140	95	100	100
69	140	100	100	100
70	140	100	100	100
71	140	100	80	85
72	140	100	20	80
73	140	90	100	100
74	140	100	90	100
75	140	100	95	95
76	140	100	100	100
77	140	100	100	100
78	140	90	100	100
79	140	95	90	100
80	140	100	100	95
81	140	100	100	95
82	140	100	100	95
83	140	90	85	85
84	140	80	75	70
85	140	100	95	100
86	140	100	100	100
87	140	100	100	100
90	140	100	85	100
91	140	95	100	100
92	140	100	100	100
93	140	100	100	100
94	140	100	100	100
95	140	100	85	90
96	140	50	100	90
97	140	65	85	80
98	140	90	70	80
99	140	80	90	90
100	140	100	100	100
101	140	100	100	100
102	140	80	85	95
103	140	95	100	100
104	140	85	80	80
105	140	95	98	85
106	140	90	100	90
107	140	100	80	90
108	140	100	100	95
110	140	80	90	85
112	140	100	100	100
114	140	95	85	95

115	140	100	100	100
116	140	80	90	100
117	140	100	100	100
118	140	100	100	100
119	140	100	100	95
120	140	100	90	95
121	140	100	95	90
123	140	100	100	100
126	140	90	100	100
127	140	90	100	100
129	140	85	95	90
130	140	80	95	95
131	140	100	100	100
134	140	95	100	90

ABUTH = velvetleaf (*Abutilon theophrasti*)

AMARE = redroot pigweed (*Amaranthus retroflexus*)

CHEAL = lambsquarter (*Chenopodium album*)

5

65. Evaluation of Preemergence Herbicidal Activity

Seeds of the desired test plant species were planted in a soil matrix prepared by mixing a loam soil (43 percent silt, 19 percent clay, and 38 percent sand, with a pH of 8.1 and an organic matter content of 1.5 percent) and sand in a 70 to 30 ratio. The soil matrix was contained in plastic pots with a surface area of 113 square centimeters. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied.

A weighed amount, determined by the highest rate to be tested, of each test compound was placed in a 25 mL glass vial and was dissolved in 6 mL of a 97:3 v/v (volume/volume) mixture of acetone and DMSO to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The stock solutions obtained were diluted with 18 mL of a 0.1 percent v/v aqueous solution of Tween[®] 20 surfactant to obtain spray solutions containing the highest application rate. Additional

application rates were obtained by serial dilution of 12 mL of the high rate solution into a solution containing 3 mL of 97:3 v/v (volume/volume) mixture of acetone and DMSO and 9 mL of the 0.1 percent v/v aqueous solution of Tween[®] 20 surfactant to obtain 1/2X, 1/4X, 1/8X and 1/16X rates of the high rate.

- 5 Compound requirements are based upon a 12 mL application volume at a rate of 187 L/ha. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters at a spray height of 18 inches (43 cm) above the soil surface. Control plants were sprayed in the
- 10 same manner with the solvent blank.

The treated pots and control pots were placed in a greenhouse maintained with an approximate 15 hour photoperiod and temperatures of 23-29 °C during the day and 22-28° C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal

15 halide 1000-Watt lamps as necessary. The water was added by top-irrigation. After 20-22 days, the condition of the test plants that germinated and grew as compared with that of the untreated plants that emerged and grew was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill or no emergence.

- 20 Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 2.

Table 2. Pre-emergent control of weeds.

Compound #	Rate (g ai/ha)	Percent Control		
		ABUTH	AMARE	CHEAL
1	140	80	100	90
3	140	98	100	95

10	280	40	50	80
12	140	100	100	100
14	140	100	100	100
15	140	60	100	100
16	140	90	100	100
26	140	10	10	90
27	140	100	100	90
41	140	80	50	90
42	140	100	100	100
43	140	98	100	100
44	140	95	30	95
49	140	100	100	99
50	140	100	100	100
52	140	100	100	90
58	140	100	100	100
59	140	90	100	60
62	140	100	100	100
63	140	100	100	100
64	140	80	100	100
70	140	100	100	100
74	140	100	95	95
77	140	100	100	100
82	140	90	100	100
93	140	100	100	100
94	140	100	100	100
95	140	90	90	90
96	140	20	70	40
99	140	100	100	95
100	140	100	100	100
101	140	100	100	100
102	140	85	95	0
103	123	90	100	100
104	140	100	100	80
105	140	100	100	95
108	140	90	100	90
110	140	100	40	40
111	140	35	100	100

112	140	100	100	100
113	140	0	0	55
115	140	100	100	100
116	140	100	100	75
121	140	100	100	100
122	140	100	100	100
123	140	100	100	100
125	140	100	100	100
131	140	90	100	100

ABUTH = velvetleaf (*Abutilon theophrasti*)

AMARE = redroot pigweed (*Amaranthus retroflexus*)

CHEAL = lambsquarter (*Chenopodium album*)

5

66. Evaluation of Herbicidal Activity in Transplanted Paddy Rice

Weed seeds or nutlets of the desired test plant species were planted in puddled soil (mud) prepared by mixing a non-sterilized mineral soil (28 percent silt, 18 percent clay, and 54 percent sand, with a pH of 7.3 to 7.8 and an organic matter content of 1.0 percent) and water at a ratio of 100 kg of soil to 19 L of water. The prepared mud was dispensed in 250 mL aliquots into 480 mL non-perforated plastic pots with a surface area of 91.6 square centimeters leaving a headspace of 3 centimeters in each pot. Rice seeds were planted in Sun Gro MetroMix® 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of 30 percent, in plastic plug trays. Seedlings at the second or third leaf stage of growth were transplanted into 650 mL of mud contained in 960 mL non-perforated plastic pots with a surface area of 91.6 square centimeters 4 days prior to herbicide application. The paddy was created by filling the 3 centimeter headspace of the pots with water. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 4-14 days in a greenhouse with an approximate 14 hour photoperiod which was maintained at 29° C during the day and 26° C during the night. Nutrients were added as

10

15

20

Osmocote (17:6:10, N:P:K + minor nutrients) at 2 g per pot. Water was added on a regular basis to maintain the paddy flood, and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the second or third true leaf stage.

5 A weighed amount, determined by the highest rate to be tested, of each test compound was placed in a 120 mL glass vial and was dissolved in 20 mL of acetone to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions obtained were diluted with 20 mL of an aqueous mixture
10 containing 0.01 percent Tween 20 (v/v). Application rates of 1/2X, 1/4X, 1/8X and 1/16X of the high rate were obtained by injecting an appropriate amount of the stock solution into the aqueous layer of the paddy. Control plants were treated in the same manner with the solvent blank.

 The treated plants and control plants were placed in a greenhouse
15 as described above and water was added as needed to maintain a paddy flood. After 3 weeks the condition of the test plants as compared with that of the control plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

 Some of the compounds tested, application rates employed, plant
20 species tested, and results are given in Table 3.

Table 3. Herbicidal Activity in Transplant Rice.

Compound #	Rate (g ai/ha)	Percent Control				
		ORYSA	ECHCG	MOOVA	CYPDI	SCPJU
4	17.5	0	0	100	95	70
5	70	0	20	100	100	90
7	70	10	100	100	100	85
17	35	0	70	100	100	90
27	17.5	0	45	100	100	99
38	70	0	50	100	100	95
41	17.5	10	50	100	100	90
42	17.5	5	40	100	100	90
44	70	0	50	90	100	95
46	35	0	20	100	100	100
52	70	0	60	99	100	95
58	17.5	0	30	100	100	50
61	35	0	80	100	100	100
70	70	0	45	100	95	95
77	35	0	70	100	100	99
86	70	10	25	100	100	95
88	70	10	100	100	100	100
89	70	0	85	100	100	90
90	17.5	0	30	100	100	90
91	35	0	99	100	100	99
92	17.5	0	40	100	99	90
100	17.5	10	30	100	100	80
103	70	10	40	100	100	95
120	35	0	40	100	100	95
125	70	5	100	100	100	100
135	17.5	10	30	100	100	85
136	35	0	15	100	100	80

ORYSA = rice (*Oryza sativa* var. Japonica)ECHCG = *Echinochloa crus-galli*SCPJU = *Scirpus juncooides*CYPDI = *Cyperus difformis*MOOVA = *Monochoria vaginalis*

67. Evaluation of Postemergence Herbicidal Activity in Cereal Crops

Seeds of the desired test plant species were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of 30 percent, in plastic pots with a surface area of 103.2 square centimeters. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 7-36 days in a greenhouse with an approximate 14 hour photoperiod which was maintained at 18° C during the day and 17° C during the night. Nutrients and water were added on a regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the second or third true leaf stage.

A weighed amount, determined by the highest rate to be tested, of each test compound was placed in a 25 mL glass vial and was dissolved in 8 mL of a 97:3 v/v mixture of acetone and DMSO to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions obtained were diluted with 16 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton[®] X-77 surfactant in a 64.7:26.0:6.7:2.0:0.7:0.01 v/v ratio to obtain spray solutions containing the highest application rates. Additional application rates were obtained by serial dilution of 12 mL of the high rate solution into a solution containing 4 mL of 97:3 v/v mixture of acetone and DMSO and 8 mL of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton X-77 surfactant in a 48.5:39.0:10.0:1.5:1.0:0.02 v/v ratio to obtain 1/2X, 1/4X, 1/8X and 1/16X rates of the high rate. Compound requirements are based upon a 12 mL application volume at a rate of 187 L/ha. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer

equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters at a spray height of 18 inches (43 cm) above average plant canopy height. Control plants were sprayed in the same manner with the blank.

5 The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 20-22 days, the condition of the test plants as compared with that of the control plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete
10 kill.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 4 .

Table 4. Post-emergent Control of Several Key Weeds in Cereal Crops

Compound #	Rate (g ai/ha)	Percent Control					
		TRZAS	HORVS	GALAP	LAMPU	PAPRH	VERPE
1	35	5	10	85	80	80	50
14	35	0	0	90	95	95	65
16	17.5	10	10	95	100	95	95
17	17.5	0	0	80	100	100	99
26	70	0	10	80	95	95	40
27	17.5	5	5	99	100	100	50
36	70	5	0	95	99	99	50
38	17.5	5	0	85	95	100	99
42	17.5	5	5	99	100	99	60
43	70	0	5	65	99	85	30
44	35	10	5	65	99	90	60
45	17.5	0	0	90	100	90	80
50	70	0	0	95	100	95	60

51	35	0	0	95	70	100	30
56	70	0	10	100	99	100	95
58	17.5	0	0	100	100	99	60
61	17.5	10	0	85	99	90	75
65	35	15	5	85	95	90	40
70	17.5	0	0	99	100	99	60
71	17.5	0	0	95	95	95	50
73	70	0	0	95	99	95	60
77	17.5	0	0	99	100	99	65
87	17.5	0	0	100	100	100	50
90	17.5	0	10	99	100	95	25
91	17.5	20	10	90	100	60	45
92	17.5	0	0	99	100	95	30
100	17.5	10	5	90	100	95	65
101	70	10	5	60	99	100	60
103	35	0	0	60	95	100	75
120	17.5	0	0	80	95	95	50
133	17.5	0	0	100	99	100	70
135	17.5	0	0	90	100	100	60

TRZAS = wheat (*Triticum aestivum*)HORVS = barley (*Hordeum vulgare*)GALAP = *Galium aparine*LAMPU = *Lamium purpureum*PAPRH = *Papaver rhoeas*VERPE = *Veronica persica*

5 68. Evaluation of Postemergence Herbicidal Safening in Cereal Crops

Seeds of the desired test plant species were planted in Sun Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of 30 percent, in plastic pots with a surface area of 103.2 square centimeters. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 7-36 days in a greenhouse with an approximate 14 hour photoperiod which was maintained at 18 ° C during the day and 17 ° C during the night. Nutrients and water were added on a regular basis

and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the second or third true leaf stage.

Treatments consisted of compound 42 and cloquintocet-mexyl
5 alone and in combination. Weighed amounts were placed in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to obtain 9 mg ai/mL stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions were diluted to 3 mg ai/mL with the addition of 2 volumes of an aqueous mixture containing acetone, water,
10 isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton[®] X-77 surfactant in a 64.7:26.0:6.7:2.0:0.7:0.01 v/v ratio. A dilution solution was prepared by mixing 1 volume of 97:3 v/v acetone/DMSO and 2 volumes of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton[®] X-77 surfactant in a 64.7:26.0:6.7:2.0:0.7:0.01
15 v/v ratio. Compound requirements are based upon a 12 mL application volume at a rate of 187 L/ha. Spray solutions of cloquintocet-mexyl and compound 42 mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients at 1:1, 1:2, and 1:4 (herbicide:safener) ratios. Formulated compounds were applied to
20 the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters at a spray height of 18 inches (43 cm) above average plant canopy height. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse
25 as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 3 weeks the condition of the test plants as compared with that

of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 5.

Table 5. Safening of herbicidal injury on cereal crops

Compound #42	<u>Application</u> Rate (g ai/ha)		<u>Percent Control</u>							
	Cloquintocet- mexyl		TRZAS		HORVW		GALAP		PAPRH	
			Ob	Ex	Ob	Ex	Ob	Ex	Ob	Ex
35	0		35	--	0	--	99	--	100	--
70	0		40	--	15	--	100	--	100	--
0	35		0	--	0	--	0	--	0	--
0	70		0	--	0	--	0	--	0	--
0	140		0	--	0	--	0	--	0	--
0	280		0	--	0	--	0	--	0	--
35	35		0	35	0	0	100	99	100	100
35	70		0	35	0	0	100	99	100	100
35	140		0	35	0	0	100	99	100	100
70	70		0	40	0	15	100	100	100	100
70	140		0	40	0	15	100	100	100	100
70	280		0	40	0	15	100	100	100	100

TRZAS = *Triticum aestivum* (spring wheat)HORVW = *Hordeum vulgare* (winter barley)GALAP = *Galium aparine*PAPRH = *Papaver rhoeas*

Ob = observed values

Ex = expected, calculated values

69. Evaluation of Postemergence Herbicidal Activity of Mixtures in Cereal Crops

5 Seeds of the desired test plant species were planted and reared as described in "Evaluation of Postemergence Herbicidal Safening in Cereal Crops" (section 37).

10 Treatments consisted of compound (as listed in Tables 6 and 7), fluroxypyr methylheptyl ester (MHE), and clopyralid monoethanolamine (MEA) salt alone and in combination. Weighed amounts were placed in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to obtain 4.5 mg ai/mL stock solutions. If the experimental compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions were

diluted to 1.5 mg ai/mL with the addition of 2 volumes of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton[®] X-77 surfactant in a 64.7:26.0:6.7:2.0:0.7:0.01 v/v ratio. A dilution solution was prepared by mixing 1 volume of 97:3 v/v acetone/DMSO and 2 volumes of an aqueous mixture containing acetone, water, isopropyl alcohol, DMSO, Agri-dex crop oil concentrate, and Triton[®] X-77 surfactant in a 64.7:26.0:6.7:2.0:0.7:0.01 v/v ratio. Compound requirements are based upon a 12 mL application volume at a rate of 187 L/ha. Spray solutions of fluroxypyr MHE, clopyralid MEA, and experimental compound mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients in two- and three-way combinations. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 square meters at a spray height of 18 inches (43 cm) above average plant canopy. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 3 weeks the condition of the test plants as compared with that of the control plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.).

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

5 The following equation was used to calculate the expected activity of mixtures containing three active ingredients, A, B, and C:

$$\text{Expected} = A + B + C - (A \times B + B \times C + A \times C)/100 + (A \times B \times C)/10000$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

10 B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

C = observed efficacy of active ingredient C at the same concentration as used in the mixture.

15 Some of the compounds tested, application rates employed, plant species tested, and results are given in Table 6 and Table 7.

Table 6. Safening and synergistic activity of herbicidal combinations.

Application Rate (g ai/ha)					TRZAS		<u>Percent Control</u> HORVW		MATCH	
Compound #93	Compound #74	Compound #58	Fluroxypyr MHE	Clopyralid MEA	Ob	Ex	Ob	Ex	Ob	Ex
0	0	0	0	35	0	--	0	--	20	--
0	0	0	35	0	0	--	0	--	0	--
0	0	0	35	35	0	0	0	0	25	20
35	-	-	0	0	15	--	10	--	45	--
35	-	-	35	35	0	15	0	10	85	56
-	35	-	0	0	25	--	0	--	10	--
-	35	-	35	35	10	25	0	0	80	28
-	-	35	0	0	35	--	15	--	0	--
-	-	35	35	35	30	35	10	15	70	20

TRZAS = *Triticum aestivum* (spring wheat)
 HORVW = *Hordeum vulgare* (winter barley)
 MATCH = *Matricaria chamomilla*

Ob = observed values
 Ex = expected, calculated values

MHE = methyl heptyl ester
 MEA = monoethanolamine salt

Table 7. Safening and synergistic activity of herbicidal combinations.

Compound #42	<u>Application</u> Rate (g ai/ha)		<u>Percent Control</u>					
	Clopyralid MEA	Fluroxypyr MHE	TRZAS		HORVW		MATCH	
			Ob	Ex	Ob	Ex	Ob	Ex
35	0	0	30	--	25	--	50	--
70	0	0	45	--	30	--	60	--
0	35	0	0	--	0	--	5	--
0	70	0	0	--	0	--	40	--
0	0	35	0	--	0	--	0	--
0	0	70	0	--	0	--	0	--
0	35	35	0	0	0	0	20	5
0	70	70	0	0	0	0	20	40
35	35	0	25	30	15	30	99	52
70	70	0	40	45	25	30	100	76
35	0	35	30	30	15	25	0	50
70	0	70	35	45	20	30	90	60
35	35	35	30	30	15	25	70	52
70	70	70	30	45	25	30	99	76
35	70	35	20	30	15	25	99	70
35	35	70	10	30	15	25	99	52
70	35	35	30	45	25	30	70	62

TRZAS = *Triticum aestivum* (spring wheat)
HORVW = *Hordeum vulgare* (winter barley)
MATCH = *Matricaria chamomilla*


Ob = observed values
Ex = expected, calculated values

MHE = methyl heptyl ester
MEA = monoethanolamine salt

73776-249(S)

CLAIMS:

1. A compound of formula I



I)

wherein


- 5 Q_1 represents H or F;

Q_2 represents a halogen with the proviso that when Q_1 is H then Q_2 is Cl or Br;

R_1 and R_2 independently represent H, C_1 - C_6 alkyl, C_3 - C_6 alkenyl, C_3 - C_6 alkynyl, hydroxy, C_1 - C_6 alkoxy, amino, C_1 - C_6 acyl, C_1 - C_6 carboalkoxy, C_1 - C_6 alkylcarbonyl, C_1 - C_6 alkylsulfonyl, C_1 - C_6 trialkylsilyl or C_1 - C_6 dialkyl phosphonyl or

- 10 R_1 and R_2 taken together with N represent a 5- or 6-membered saturated ring; and

Ar represents a polysubstituted aryl group selected from the group consisting of



a)


wherein W_1 represents halogen;

73776-249(S)

X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, -CN, - NR_3R_4 or fluorinated acetyl or propionyl;

- 5 Y_1 represents C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen or -CN, or, when X_1 and Y_1 are taken together, represents - $O(CH_2)_nO$ - wherein $n=1$ or 2; and

R_3 and R_4 independently represent H or C_1 - C_4 alkyl;



b)


wherein

- 10 W_2 represents F or Cl;

X_2 represents F, Cl, -CN, - NO_2 , C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, - NR_3R_4 or fluorinated acetyl or propionyl;

- 15 Y_2 represents halogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl or -CN, or, when W_2 represents F, X_2 and Y_2 taken together represent - $O(CH_2)_nO$ - wherein $n=1$ or 2; and

R_3 and R_4 independently represent H or C_1 - C_6 alkyl; and



c)

73776-249(S)

wherein

Y_3 represents halogen, -CN or -CF₃;

Z_3 represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl; and

R₃ and R₄ independently represent H, or C₁-C₆ alkyl;

or an agriculturally acceptable derivative of the carboxylic acid group,

wherein the alkyl, alkenyl and alkynyl groups are straight chain moieties, or branched chain or cyclic moieties to the extent allowed by the number of carbon atoms present in the groups.

2. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which R₁ and R₂ represent H.


3. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q₁ represents H.

4. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q₂ represents Cl.

5. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Q₁ represents H and Q₂ represents Cl.

6. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Ar represents

73776-249(S)



wherein

W_1 represents halogen;

X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio,

- 5 C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, -CN, - NR_3R_4 or fluorinated acetyl or propionyl;

Y_1 represents C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen or -CN, or, when X_1 and Y_1 are taken together, represents $-O(CH_2)_nO-$ wherein $n=1$ or 2 ; and

- 10 R_3 and R_4 independently represent H or C_1 - C_4 alkyl.


7. A compound of claim 6, or an agriculturally acceptable derivative of the carboxylic acid group, in which W_1 represents Cl or F.

8. A compound of claim 6, or an agriculturally acceptable derivative of the carboxylic acid group, in which X_1 represents C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 haloalkyl
15 or - NR_3R_4 .

9. A compound of claim 6, or an agriculturally acceptable derivative of the carboxylic acid group, in which Y_1 represents Cl, Br or - CF_3 .

10. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Ar represents

73776-249(S)



wherein

W_2 represents F or Cl;

X_2 represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy,

- 5 C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl;

Y_2 represents halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or -CN, or, when W_2 represents F, X_2 and Y_2 taken together represent -O(CH₂)_nO- wherein n=1 or 2; and


- 10 R_3 and R_4 independently represent H or C₁-C₆ alkyl.

11. A compound of claim 10, or an agriculturally acceptable derivative of the carboxylic acid group, in which X_2 represents C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ haloalkyl or -NR₃R₄.

12. A compound of claim 10, or an agriculturally acceptable derivative of the
15 carboxylic acid group, in which Y_2 represents Cl, Br or -CF₃.

13. A compound of claim 1, or an agriculturally acceptable derivative of the carboxylic acid group, in which Ar represents

73776-249(S)



wherein

Y_3 represents halogen, $-CN$ or $-CF_3$;

Z_3 represents F , Cl , $-CN$, $-NO_2$, C_1-C_4 alkyl, C_1-C_4 alkoxy,

- 5 C_1-C_4 alkylthio, C_1-C_4 alkylsulfinyl, C_1-C_4 alkylsulfonyl, C_1-C_4 haloalkyl, C_1-C_4 haloalkoxy, C_1-C_4 alkoxy-substituted C_1-C_4 alkyl, C_1-C_4 alkoxy-substituted C_1-C_4 alkoxy, $-NR_3R_4$ or fluorinated acetyl or propionyl; and

R_3 and R_4 independently represent H , or C_1-C_6 alkyl.

14. A compound of claim 13, or an agriculturally acceptable derivative of the
10 carboxylic acid group, in which Y_3 represents Cl , Br or $-CF_3$.

15. A compound of claim 13, or an agriculturally acceptable derivative of the carboxylic acid group, in which Z_3 represents C_1-C_4 alkoxy, C_1-C_4 alkylthio, C_1-C_4 haloalkyl or $-NR_3R_4$.

16. A herbicidal composition comprising an herbicidally effective amount of a
15 compound according to any one of claims 1 to 15, or an agriculturally acceptable derivative of the carboxylic acid group, in admixture with an agriculturally acceptable adjuvant or carrier.

17. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil to prevent the emergence of
20 vegetation an herbicidally effective amount of a compound according to any one of claims 1 to 15, or an agriculturally acceptable derivative of the carboxylic acid group.

73776-249(S)

18. A compound of formula


ArMetal

wherein

Ar represents a polysubstituted aryl group selected from the group consisting

5 of

a)



wherein

 W_1 represents halogen;

10 X_1 represents F, Cl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkylsulfinyl, C_1 - C_4 alkylsulfonyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkyl, C_1 - C_4 alkoxy-substituted C_1 - C_4 alkoxy, -CN, -NR₃R₄ or fluorinated acetyl or propionyl;


 Y_1 represents C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, halogen or -CN, or, when X_1 and

15 Y_1 are taken together, represents -O(CH₂)_nO- wherein n=1 or 2; and

 R_3 and R_4 independently represent H or C_1 - C_4 alkyl;

73776-249(S)

b)



wherein


W₂ represents F or Cl;

- 5 X₂ represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl;

- 10 Y₂ represents halogen, C₁-C₄ alkyl, C₁-C₄ haloalkyl or -CN, or, when W₂ represents F, X₂ and Y₂ taken together represent -O(CH₂)_nO- wherein n=1 or 2; and

R₃ and R₄ independently represent H or C₁-C₆ alkyl; and

c)



wherein

- 15 Y₃ represents halogen, -CN or -CF₃;

73776-249(S)

Z₃ represents F, Cl, -CN, -NO₂, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄ alkylthio, C₁-C₄ alkylsulfinyl, C₁-C₄ alkylsulfonyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ alkoxy-substituted C₁-C₄ alkyl, C₁-C₄ alkoxy-substituted C₁-C₄ alkoxy, -NR₃R₄ or fluorinated acetyl or propionyl; and

5 R₃ and R₄ independently represent H, or C₁-C₆ alkyl; and

Metal represents Mg-halide, Zn-halide, tri-(C₁-C₄ alkyl)tin, lithium, copper, or B(OR₈)(OR₉), where R₈ and R₉ are independent of one another, hydrogen, C₁-C₄ alkyl, or when taken together form an ethylene or propylene group,

10 wherein the alkyl, alkenyl and alkynyl groups are straight chain moieties, or branched chain or cyclic moieties to the extent allowed by the number of carbon atoms present in the groups.

19. A compound according to claim 18, which is:

2-(4-chloro-2-fluoro-3-methoxyphenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-5-methoxyphenyl)-[1,3,2]-dioxaborinane;

15 2-(4-chloro-2-fluoro-3-ethoxyphenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-3-(methylthio)phenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-3-methoxymethylphenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-3-isopropoxyphenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-3-difluoromethylphenyl)-[1,3,2]-dioxaborinane;

20 2-(4-chloro-2-fluoro-3-fluoromethylphenyl)-[1,3,2]-dioxaborinane;

4-[1,3,2]-dioxaborinan-2-yl-3-fluoro-2-methoxybenzonitrile;

2-(4-chloro-2-fluoro-3-methoxyethoxyphenyl)-[1,3,2]-dioxaborinane;

73776-249(S)

2-(2,4-dichloro-3-ethoxyphenyl)-[1,3,2]-dioxaborinane;

2-[4-chloro-2-fluoro-3-(2,2-difluoroethoxyphenyl)-[1,3,2]-dioxaborinane;

2-(4-chloro-2-fluoro-5-ethoxyphenyl)-[1,3,2]-dioxaborinane;

2-(2,4-dichloro-3-(methylthio)phenyl)-[1,3,2]-dioxaborinane;

5 2-fluoro-3,4-methylenedioxyphenylboronic acid;

2-fluoro-3-methoxy-4-methylphenylboronic acid;

4-bromo-2-fluoro-3-methoxyphenylboronic acid;

2-(2-fluoro-3,4-methylenedioxyphenyl)-[1,3,2]-dioxaborinane;

4-chloro-3-(diethylamino)-2-fluorophenylboronic acid;

10 2,4-dichloro-3-dimethylaminophenylboronic acid;

2-chloro-4-fluoro-3-methoxyphenylboronic acid;

6-chloro-2-fluoro-3-(4,4,5,5-tetramethyl-{1,3,2}-dioxaborolan-2-yl)-phenyl
methyl amine;

(2-fluoro-3-methoxy-4-trifluoromethylphenyl)trimethyl stannane;

15 (3-butoxy-4-chloro-2-fluorophenyl)trimethyl stannane; or

[4-chloro-2-fluoro-3-(1-fluoro-1-methylethyl)phenyl]trimethyl stannane.

20. A compound according to claim 1, which is:

4-amino-3-chloro-6-(4-chloro-2-fluoro-3-thiomethylphenyl)pyridine-2-
carboxylic acid methyl ester;

73776-249(S)

4-Amino-3-chloro-6-(4-chloro-3-cyano-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethylphenyl)pyridine-2-carboxylic acid methyl ester;

5 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethylphenyl)pyridine-2-carboxylic acid methyl ester;

10 4-Amino-3-chloro-6-(4-cyano-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl]pyridine-2-carboxylic acid methyl ester;

15 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-[2,4-dichloro-3-(2,2-difluoroethoxy)phenyl]pyridine-2-carboxylic acid methyl ester;

20 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methoxyethoxy)phenyl]pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-5-fluoro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-3-methylthiophenyl)pyridine-2-carboxylic acid methyl ester;

73776-249(S)

4-Amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

5 4-Amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-[4-chloro-3-(diethylamino)-2-fluorophenyl]pyridine-2-carboxylic acid, methyl ester;

10 4-acetylamino-3-chloro-6-(4-chloro-2-fluoro-5-methoxy-phenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

15 4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

20 4-Acetylamino-3-chloro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-ethoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-[2,4-dichloro-3-(2,2-difluoroethoxy)phenyl]-2-pyridine-2-carboxylic acid methyl ester;

73776-249(S)

4-acetylamino-3-chloro-6-(2,4-difluoro-3-methylphenyl)-pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester;

5 4-acetylamino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoro-1-methylethyl)phenyl]-pyridine-2-carboxylic acid, methyl ester;

4-amino-3-chloro-6-(4-chloro-2-fluoro-5-nitrophenyl)-pyridine-2-carboxylic acid methyl ester;

4-amino-3-chloro-6-(5-amino-4-chloro-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester;

15 4-amino-3-chloro-6-(4-chloro-5-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid methyl ester;

4-amino-6-(4-bromo-2-fluoro-3-methoxyphenyl)-3-chloro-pyridine-2-carboxylic acid methyl ester;

20 4-Acetylamino-3-chloro-6-(2,4,6-trifluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-(2,4-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-(2-chloro-4-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-acetylamino-3-chloro-6-[2,4-dichloro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid methyl ester;

5 4-Acetylamino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-3-dimethylamino-2,6-difluorophenyl)-pyridine-2-carboxylic acid, methyl ester;

10 4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(2,4-dichloro-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

15 4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-[2,4-dichloro-3-(difluoromethyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

20 4-Acetylamino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)-phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)-phenyl]pyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-Acetylamino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

5 4-Acetylamino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethoxyphenyl)pyridine, carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-difluoromethyl-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Acetylamino-3-chloro-6-(2-fluoro-4-fluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methylphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-6-(2-bromo-4-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

15 4-Acetylamino-6-(4-bromo-2,6-difluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(3-difluoromethyl-2,4,6-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester;

20 4-Acetylamino-3-chloro-6-(2-chloro-4-difluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(2-chloro-4,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-Acetylamino-3-chloro-6-(2,4-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-3-cyclopropyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

5 4-Acetylamino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2,2-trifluoroacetyl)phenyl]-pyridine-2-carboxylic acid, methyl ester;

4-amino-3-chloro-6-(2-fluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Amino-3-chloro-6-[2,4-dichloro-3-(dimethylamino)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methylamino)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

15 4-Acetylamino-3-chloro-6-[4-chloro-3-(dimethylamino)-2,6-difluorophenyl]pyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester;

20 4-Acetylamino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

4-Acetylamino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-amino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)-pyridine-2-carboxylic acid methyl ester;

73776-249(S)

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

5 4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid methyl ester;

10 4-Amino-3-chloro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(2,4-difluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester;

15 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid methyl ester;

4-Amino-3-chloro-6-[4-chloro-3-(dimethylamino)-2,6-difluorophenyl]pyridine-2-carboxylic acid, methyl ester;

20 4-Amino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-(2-chloro-4-difluoromethyl-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-Amino-3-chloro-6-(4-chloro-3-difluoromethyl-2,6-difluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-(2-chloro-4,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

5 4-Amino-3-chloro-6-(2,4,6-trifluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-6-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

15 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-3-difluoromethylphenyl)pyridine-2-carboxylic acid, methyl ester;

20 4-Amino-3-chloro-6-[2,4-dichloro-3-(1-fluoro-1-ethylphenyl)]pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)phenyl]-pyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester;

5 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Amino-3-chloro-6-(2,4-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-difluoromethyl-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2-fluoro-4-fluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

15 4-Amino-3-chloro-6-(2,4-difluoro-5-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2-chloro-4-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

20 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methylphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-6-(2-bromo-4-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

4-Amino-6-(4-bromo-2,6-difluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid, methyl ester;

73776-249(S)

4-Amino-3-chloro-6-(3-difluoromethyl-2,4,6-trifluorophenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2-chloro-4-difluoromethyl-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

5 4-Amino-3-chloro-6-(2,4-difluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(4-chloro-3-cyclopropyl-2-fluorophenyl)pyridine-2-carboxylic acid, methyl ester;

10 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(2,2,2-trifluoroacetyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfinyl-phenyl)pyridine-2-carboxylic acid methyl ester;

4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methanesulfonylphenyl)pyridine-2-carboxylic acid methyl ester;

15 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-pyridine-2-carboxylic acid, ethyl ester;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, 2-butoxy-ethyl ester;

20 4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester;

4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester;

73776-249(S)

4-Amino-3-chloro-6-(4-bromo-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, butyl ester;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid, butyl ester;

5 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, 2-butoxy-ethyl ester;

4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid;

10 4-Amino-3-chloro-6-(2,6-difluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-dimethylaminophenyl)pyridine-2-carboxylic acid;

4-amino-3-chloro-6-(4-cyano-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid, methyl ester;

15 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-(2,2-difluoroethoxy)phenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-(2,4-difluoro-3-methylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(2,4-dichloro-3-methylphenyl)pyridine-2-carboxylic acid;

20 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid;

73776-249(S)

4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-ethoxyphenyl)pyridine-2-carboxylic acid;

5 4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methylthiophenyl)pyridine-2-carboxylic acid;

10 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-difluoromethylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-trifluoromethylphenyl)pyridine-2-carboxylic acid;

15 4-Amino-3-chloro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-ethoxyphenyl)pyridine-2-carboxylic acid;

20 4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-5-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluorophenyl)pyridine-2-carboxylic acid;

73776-249(S)

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyethoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-fluoromethylphenyl)pyridine-2-carboxylic acid;

5 4-Amino-3-chloro-5-fluoro-6-(2-fluoro-3,4-methylenedioxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-[4-chloro-3-(diethylamino)-2-fluorophenyl]pyridine-2-carboxylic acid;

10 4-Amino-3-chloro-6-[2,4-dichloro-3-(dimethylamino)phenyl]pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(methylamino)phenyl]pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(2-fluoro-3-methoxy-4-methylphenyl)pyridine-2-carboxylic acid;

15 4-Amino-6-(4-bromo-2-fluoro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid;

4-Amino-6-(4-bromo-2-chloro-3-methoxyphenyl)-3-chloropyridine-2-carboxylic acid;

20 4-Amino-3-chloro-6-(4-chloro-2-fluoro-5-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-3-butoxy-2-fluorophenyl)pyridine-2-carboxylic acid;

73776-249(S)

4-Amino-3-chloro-6-[2,4-dichloro-3-(1-fluoroethyl)phenyl]pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(2,4-dichloro-3-difluoromethylphenyl)pyridine-2-carboxylic acid;

5 4-Amino-3-chloro-6-[2,4-dichloro-3-(1-fluoro-1-methylethylphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-3-ethyl-2-fluorophenyl)pyridine-2-carboxylic acid;

10 4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1,2,2,2-tetrafluoroethyl)-phenyl]pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-[4-chloro-2-fluoro-3-(1-fluoropropyl)phenyl]pyridine-2-carboxylic acid, methyl ester;

4-Amino-3-chloro-6-(2,3,4-trifluorophenyl)pyridine-2-carboxylic acid;


15 4-Amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxymethoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-2,6-difluoro-3-methoxyphenyl)pyridine-2-carboxylic acid;

4-Amino-3-chloro-6-(4-chloro-3-difluoromethoxy-2-fluorophenyl)pyridine-2-carboxylic acid;

20 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-pyridine-2-carboxylic acid, triethyl amine salt; or

4-Amino-3-chloro-6-(2,4-dichloro-3-methoxyphenyl)pyridine-2-carboxylic acid, triethyl amine salt.



D)