



(51) International Patent Classification:

C07K 16/40 (2006.01) *A61P 35/00* (2006.01)
A61K 39/395 (2006.01)

(21) International Application Number:

PCT/IL2014/050678

(22) International Filing Date:

24 July 2014 (24.07.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicant: **YEDA RESEARCH AND DEVELOPMENT CO. LTD.** [IL/IL]; at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot (IL).

(72) Inventors: **SAGI, Irit**; 12 Neve Metz, The Weizmann Institute of Science, 7610002 Rehovot (IL). **GROSSMAN, Moran**; c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot (IL). **BEN-CHETRIT, Nir**; c/o Yeda Research and Development Co. Ltd., at the Weizmann Institute of Science, P.O. Box 95, 7610002 Rehovot (IL).

(74) Agents: **EHRlich, Gal** et al.; G. E. EHRlich (1995) LTD., The Rogovin-Tidhar Tower, 15th Floor, 11 Mena-chem Begin Road, 5268104 Ramat Gan (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))


Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTIBODIES TARGETED AGAINST LOXL-2 FOR THE TREATMENT OF COLLAGEN-ASSOCIATED PATHOLOGIES

FIG. 5B



(57) Abstract: An article of manufacture is disclosed which comprises at least two active agents, wherein a first of the two active agents comprises an anti-cancer agent or an antifibrotic agent and a second of the at least two active agents downregulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of altering the structure of the extracellular matrix.

ANTIBODIES TARGETED AGAINST LOXL-2 FOR THE TREATMENT OF COLLAGEN-ASSOCIATED PATHOLOGIES

5 FIELD AND BACKGROUND OF THE INVENTION

The present invention, in some embodiments thereof, relates to antibodies targeted against lysyl-oxidase like protein-2 (LOXL-2) which are capable of altering the architecture of pathologic collagen assembly.

The extracellular matrix (ECM) is an essential mediator of tissue function that
10 provides both chemical and mechanical stimuli to influence cellular behavior in both health and disease. The ECM consists of a mesh of fibrillar and non-fibrillar collagens, elastic fibers, glycoproteins and proteoglycans that assemble into complex networks in a tissue specific manner which determine the biophysical properties of tissues such as stiffness, compliance and resilience. Remodeling of the ECM by a variety of tissue
15 remodeling enzymes that cross-link, glycosylate and/or degrade the ECM, can create space for cells to migrate, regulate tissue architecture, and activate, deactivate, or alter the activity of signaling molecules. Tightly controlled ECM homeostasis is therefore essential for life, and dysregulated ECM remodeling can result with pathologic outcomes such as cancer and fibrosis. Indeed, over-expression of tissue remodeling enzymes is one
20 of the hallmarks of cancer and examination of ECM derived from solid tumors reveals massive deposition of collagen, altered ECM morphology and increased tissue stiffness. In addition, proteomic analyses have demonstrated that the composition of the tumor extracellular matrix changes with tumor metastatic potential. Thus, controlling ECM morphology and composition is considered as potent therapeutic strategy for cancer
25 patients (Siddikuzzaman, Grace, V.M. & Guruvayoorappan, C. Inflammopharmacology 19, 117-29 (2011)). However, despite the growing amount of data correlating between ECM biophysical properties and tumor malignancy, how to clinically interfere with the altered biophysical ECM during the onset and progression of cancer is unknown.

The lysyl oxidases (LOX) comprise a family of five copper-dependent amine
30 oxidases homologous enzymes (LOX and LOXL1-4) that initiate the process of covalent intra- and intermolecular cross-linking of collagens and elastin. Among the LOX family, LOX and LOXL2 are highly expressed in many human cancers, partly and adversely in correlation to clinical outcome. Activation of LOX family members during cancer has been demonstrated to increase the stiffness of the tissue due to extensive crosslinking of

collagen fibrils and to effect downstream signals by modification of snail and integrins, repression of E-cadherin leading to induction of epithelial to mesenchymal (EMT) transition. Interestingly, LOXL2 is highly over-expressed mainly in invasive metastatic tumors more than noninvasive tumors, and is linked to promoting tumor cell invasion, angiogenesis and remodeling of the tumor microenvironment. LOXL2 has been proposed to function both extracellularly and intracellularly to activate oncogenic signaling pathways leading to epithelial–mesenchymal transition (EMT) by and invasion of breast cancer cells. In the tumor stroma, LOXL2 mediates fibroblast activation through integrin engagement and FAK signaling. It has been recently shown that LOXL2 and the transcription factor, E47, contribute to early steps of metastatic colonization by cell and noncell autonomous functions regulating the recruitment of bone marrow progenitor cells to the lungs and by direct transcriptional regulation of fibronectin and cytokines TNF α , ANG-1 and GM-CSF. Genetic, chemical or antibody inhibition of LOXL2 significantly reduced lungs, liver and bone metastasis through its effect on matrix remodeling and cell invasion (Barker, H.E. et al. Cancer Res 71, 1561-72 (2011); Barry-Hamilton, V. et al. Nat Med 16, 1009-17 (2010)).

An antibody developed to specifically target the fourth SRCR domain of LOXL-2 is disclosed in Barry-Hamilton V. et al., Nat Med. 2010 Sep;16(9):1009-17.

An antibody targeted to the catalytic site of LOXL-2 is disclosed in U.S. Patent No. 8,168,180.

SUMMARY OF THE INVENTION

According to an aspect of some embodiments of the present invention there is provided an article of manufacture comprising at least two active agents, wherein a first of the two active agents comprises an anti-cancer agent or an antifibrotic agent and second of the at least two active agents downregulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of altering the structure of the extracellular matrix.

According to an aspect of some embodiments of the present invention there is provided an isolated antibody comprising an antigen recognition region which specifically binds to lysyl-oxidase like protein-2 (LOXL-2) and being capable of down-regulating crosslinking of type I collagen in vitro.

According to an aspect of some embodiments of the present invention there is provided an isolated antibody comprising an antigen recognition region which comprises CDR amino acid sequences as set forth in SEQ ID NOs: 3, 4, 5, 6, 7 and 8.

According to an aspect of some embodiments of the present invention there is
5 provided a method of treating a disease associated with aberrant collagen deposition in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the antibody disclosed herein, thereby treating the disease.

According to an aspect of some embodiments of the present invention there is provided an article of manufacture comprising an isolated antibody comprising an
10 antigen recognition region which comprises CDR amino acid sequences set forth in SEQ ID NO: 3, 4, 5, 6, 7 and 8 and an anti-cancer agent or an antifibrotic agent.

According to an aspect of some embodiments of the present invention there is provided a method of treating a disease associated with aberrant collagen deposition in a subject in need thereof, comprising administering to the subject a therapeutically
15 effective amount of an agent which downregulates the activity and/or expression of the LOXL-2 and an additional agent selected from the group consisting of an anti-cancer agent and an antifibrotic agent, thereby treating the disease.

According to an aspect of some embodiments of the present invention there is provided a combination of an agent which downregulates the activity and/or expression
20 of the LOXL-2 and an additional agent selected from the group consisting of an anti-cancer agent and an antifibrotic agent for use in treating a disease associated with aberrant collagen deposition.

According to an aspect of some embodiments of the present invention there is provided a pharmaceutical composition comprising the antibody described herein and a
25 pharmaceutically acceptable carrier.

According to an aspect of some embodiments of the present invention there is provided a pharmaceutical composition comprising an anti-cancer agent or an antifibrotic agent as a first active agent and an agent which downregulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of
30 altering the structure of the extracellular matrix as a second active agent, and a pharmaceutically acceptable carrier.

According to an aspect of some embodiments of the present invention there is provided a bispecific antibody comprising a first antigen-binding domain that specifically binds to LOXL-2 and a second antigen-binding domain that specifically binds to an epitope of a protein that is associated with cancer.

5 According to some embodiments of the invention, the second agent is an antibody which comprises an antigen recognition region which specifically binds to the LOXL-2.

According to some embodiments of the invention, the antibody specifically binds the catalytic site of the LOXL-2.

10 According to some embodiments of the invention, the antibody does not bind to the fourth scavenger receptor-cysteine-rich (SRCR) domain of the LOXL-2.

According to some embodiments of the invention, the second agent is a polynucleotide agent which downregulates the expression of the LOXL-2.

15 According to some embodiments of the invention, the first agent and the second agent are packaged in separate packaging.

According to some embodiments of the invention, the first agent and the second agent are formulated in a single pharmaceutical composition.

According to some embodiments of the invention, the second agent is capable of down-regulating crosslinking of type I collagen in vitro.

20 According to some embodiments of the invention, the antibody is capable of specifically binding to the catalytic site of the LOXL-2.

According to some embodiments of the invention, the antibody does not bind to the fourth scavenger receptor-cysteine-rich (SRCR) domain of the LOXL-2.

25 According to some embodiments of the invention, the antibody is for use in treating a disease associated with aberrant collagen deposition.

According to some embodiments of the invention, the agent which downregulates the activity and/or expression of the LOXL-2 does not affect angiogenesis in vivo.

30 According to some embodiments of the invention, the antigen recognition region comprises CDR amino acid sequences as set forth in SEQ ID NOs: 3, 4, 5, 6, 7 and 8.

According to some embodiments of the invention, the amino acid sequence of the V_H of the antibody is as set forth in SEQ ID NO: 1.

According to some embodiments of the invention, the amino acid sequence of the V_L of the antibody is as set forth in SEQ ID NO: 2.

According to some embodiments of the invention, the anti-cancer agent comprises a chemotherapeutic agent.

5 According to some embodiments of the invention, the chemotherapeutic agent is cisplatin.

According to some embodiments of the invention, the anti-cancer agent is an antibody.

10 According to some embodiments of the invention, the antibody binds specifically to MMP-9.

According to some embodiments of the invention, the disease is cancer.

According to some embodiments of the invention, the disease is pulmonary alveolar proteinosis (PAP).

According to some embodiments of the invention, the cancer is breast cancer.

15 According to some embodiments of the invention, the cancer is colon cancer.

According to some embodiments of the invention, the breast cancer is triple negative breast cancer.

According to some embodiments of the invention, the cancer has metastasized.

20 According to some embodiments of the invention, the administering of the anti-cancer agent or antifibrotic agent is effected following administering of the agent which downregulates the activity and/or expression of the LOXL-2.

According to some embodiments of the invention, the disease is a fibrotic disease.

25 According to some embodiments of the invention, the agent which downregulates the activity and/or expression of the LOXL-2 is an antibody which comprises an antigen recognition region which specifically binds to the LOXL-2, the antibody being capable of down-regulating crosslinking of type I collagen in vitro.

30 According to some embodiments of the invention, the agent which downregulates the activity and/or expression of the LOXL-2 does not affect angiogenesis in vivo.

According to some embodiments of the invention, the antigen recognition region comprises CDR amino acid sequences as set forth in SEQ ID NO: 3, 4, 5, 6, 7 and 8.

According to some embodiments of the invention, the amino acid sequence of the V_H of the antibody is as set forth in SEQ ID NO: 1.

5 According to some embodiments of the invention, the amino acid sequence of the V_L of the antibody is as set forth in SEQ ID NO: 2.

Unless otherwise defined, all technical and/or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although methods and materials similar or equivalent to those
10 described herein can be used in the practice or testing of embodiments of the invention, exemplary methods and/or materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be necessarily limiting.

BRIEF DESCRIPTION OF THE DRAWINGS

15 Some embodiments of the invention are herein described, by way of example only, with reference to the accompanying images. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of embodiments of the invention. In this regard, the description taken with the drawings makes apparent to those skilled in the art how
20 embodiments of the invention may be practiced.

In the drawings:

FIGs. 1A-C Characterization of ECM maturation by human dermal fibroblasts. A) representative two-photon SHG images of collagen at various time points, depicting collagen synthesis (day 5), assembly into linear fibrils (day 7) and maturation into cross-
25 linked collagen fibers (4 weeks). Scale bar 50µm. B) LOXL2 therapeutic window. A scheme illustrating over-expression of LOXL2 at later stages of collagen assembly based on qPCR analysis of representative collagen crosslinking enzymes (Lox and Loxl2), and major collagens found in basement membrane (Col I and Col IV) (see Figures 2A-D for qPCR full analysis). C) Domain organization of LOXL2: pre domain (pink), four SRCR
30 domains (blue), and Cu²⁺-dependent catalytic domain (residues 545-775). This domain was cloned into recombinant expression system, and used for immunizing mice in order to develop specific antibodies against the catalytic activity of LOXL2.

FIGs. 2A-D are graphs illustrating qPCR analysis of LOX (Figure 2A), LOXL2 (Figure 2B), Collagen I (Figure 2C) and Collagen IV (Figure 2D).

FIGs. 3A-C GS341 is a nanomolar binder of LOXL2 and inhibits LOXL2 enzymatic activity. A) ELISA assay for GS341 binding to the catalytic domains of LOXL2. B) Binding analysis in solution by microscale thermophoresis. Values on the Y-axis represent the percentage of the maximal thermophoretic response observed, or in the case of no detectable binding values on the Y-axis represent the thermophoretic shift of the labeled protein. All binding curves were determined in triplicate by MST and are represented as the mean $\pm$ SD. Error bars represent standard deviation of representative experiments done in triplicate. C) Inhibition of collagen cross-linking by GS341.

FIGs. 4A-F Inhibition of LOXL2 by GS341 changes the morphology of ECM secreted by fibroblast cells. A) 3D spheroid invasion assay of MDA-231 cells through Matrigel in the presence of vehicle or GS341 (10uM) demonstrated reduction in invasion by 70% upon treatment with GS341. B) Representative SHG images of control and GS341 treated fibroblasts demonstrate wavy-like organization of collagen fibers in the control samples, while treatment Scale bar 50uM. C) Raman (scale bar 10uM) analyses of ECM derived from the fibroblast grown in the presence of PBS or GS341 (100ngr) for four weeks, demonstrate dramatic differences in collagen morphology. D) MS analysis of ECM derived from the control and GS341 treated cells did not detect statistically significant differences in ECM composition indicating that GS341 affects only ECM morphology (The scale represent relative fold change within each protein compared with the other group). E) SEM analysis of nanoscale changes in collagen fiber morphology upon treatment with GS341 of same sample as in B. F) Fibril orientation analysis of the fibrils done by Fiji. Inhibition of collagen cross-linking by GS341 led to multi-oriented fibril organization.

FIG. 5A Representative two-photon SHG images of collagen secreted by human dermal fibroblast treated with PBS at various time points, depicting collagen synthesis (day 5), assembly into linear fibrils (day 7) and maturation into cross-linked collagen fibers (4 weeks). Scale bar 50 μ m.

FIG. 5B SHG images of cells treated with 100ngr GS341 did not mature into crosslinked collagen fibers as in Figure 5A (4 weeks), but remained linear,

FIG. 6 is a Raman spectra for a non-treated and GS341 treated decellularized ECM sample, demonstrating differences in collagen bond chemistry.

FIGs. 7A-H De-remodeling of the tumor ECM reduces metastasis seeding to the lungs. A) Treatment with GS341 (30mg/kg) reduce tumor size in experimental metastasis model in which MDA-231 cells are implanted into the mammary fatpad of mice after 8 weeks (n=4 for each group), $P^* < 0.05$. B) Quantification of lung metastases formed after orthotopic injection of MDA-MB-231 cells. Mice were either untreated or treated with GS341. n=5 mice per group. C) Inhibition by GS341 reduced tumor cell proliferation in vivo demonstrated by immunofluorescence staining for phosphohistone h3 (ph3) of paraffin tumor sections $P^* < 0.05$. D) Mass spectrometry analysis of ECM from tumors reveal changes in ECM composition (n=2 per group). E) Representative SHG imaging of collagen of the tumors demonstrate a shift from linear collagen organization of the vehicle treated tumors into random orientation upon treatment with GS341. F) Analysis of fiber orientation. G) SEM analysis of nanoscale changes in collagen fiber morphology upon treatment with GS341 are similar to that observed for the fibroblast cells (2E). H) Quantification of fibril diameter by ImageJ depicts a shift to thinner fibrils upon treatment with GS341 (n=100) $P^{***} < 0.0001$. * Difference from a previous time point is statistically significant with $P < 0.0001$. Error bars represent standard error of the mean of 20 images. n=2 animals for each time point. Scale bar represents 25 μm .

FIGs. 8A-B is a schematic illustration of the sequence of the VH (Figure 8A – SEQ ID NO: 1) and VL (Figure 8B - SEQ ID NO: 2).

FIGs. 9A-C is a tube formation assay demonstrating that treatment with GS341 reduces the number of nodes formed by endothelial cells during formation of tubes.

FIGs. 9D-F is a 2D invasion assay of MDA-MB-231 cells through matrigel in the presence of control antibody and GS341. Treatment with GS341 reduce cell invasion by 50%.

FIGs. 9G-H is a 2D invasion assay of MDA-MB-231 cells through matrigel in the presence of control antibody and a combination of GS341 and anti-MMP-9 antibody.

FIGs. 10A-E CCL4 mice model of liver metastasis. CCL4 is injected into mice. Fibrosis picks up after 24 hours. Thick branched collagen forms in the liver. TIMP is

elevated after 24 hours and inhibits collagenases (Figure 10E). The tissue is stained with Sirius red collagen stain. From 72 hours, the tissue regenerates-loss of red stain (Figures 10A-D).

FIGs. 11A-C Treatment of CCL4 mice with GS341 significantly attenuates
5 fibrosis (loss of red stain) resulting in a speeding up of tissue regeneration.

FIGs. 12A-B Treatment with Fab GS341 provides access of collagenases (possibly from immune cells) to the collagen fibrotic scar. Green-tissue collagen zymography indicative of collagenase activity in the tissue red-staining of TIMPs (endogenous MMP inhibitors) and blue-DAPI. Note the TIMP signal is not localized on
10 the scarred fibers in the treated tissue which means no interference with collagenase activity during treatment with GS341 (possibly secreted from different cells).

DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

The present invention, in some embodiments thereof, relates to antibodies
15 targeted against lysyl-oxidase like protein-2 (LOXL-2) which are capable of altering the architecture of pathological collagen assembly.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not necessarily limited in its application to the details set forth in the following description or exemplified by the Examples. The invention is
20 capable of other embodiments or of being practiced or carried out in various ways.

The extracellular medium (ECM) is an extremely versatile biomaterial regulating intricately cell-cell connections, interactions and communications. Tissue stiffness via remodeling of collagen, a major component of the ECM, is often linked with pathological scenarios including fibrosis and poor prognosis in several solid tumors.

25 The present inventors characterized ECM synthesis, assembly and maturation by human dermal fibroblast which served as a model system for stromal ECM. By characterizing LOX/LOXL2 expression levels profiles, the present inventors found that LOXL2 is involved in the later stages of covalent collagen assembly (Figures 1A-B). The present inventors targeted the enzymatic activity of LOXL-2 at the ECM by
30 developing an inhibitory monoclonal antibody of LOXL-2. They screened for antibodies that demonstrates a direct effect on ECM architecture. This effect was quantified by an integrated spectroscopic approach involving two photon second

harmonic generation, raman microspectroscopy and scanning electron microscopy (Figures 4A-F, 5A, 6 and 7A-H). The ECM interfering antibody demonstrated therapeutic potential in mice models of breast cancer metastasis (Figure 5B), by inducing morphological changes in collagen fibers organization in the primary tumor
5 crucial for cancer cell invasion and metastasis (Figures 9A-H, 10A-E and 11A-C).

The results demonstrate shifts of cancerous collagen linearization to random orientation distribution which led to significant decrease of primary tumor progression and attenuation of lung metastasis via collagen disassembly mechanisms. Altogether the present results highlight LOXL-2 antibodies as effective inhibitors targeting pathological
10 collagen assembly. These inhibitors may be used as primary inhibitors, as well as in adjuvant therapy for cancer, fibrosis and other collagen based pathologies.

Thus, according to one aspect of the present invention there is provided an agent which down-regulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of altering the structure of the extracellular matrix.

The term "lysyl-oxidase like protein-2 (LOXL-2)" refers to human LOXL-2 having the EC 1.4.3.13, which has a collagen crosslinking activity and has an amino acid sequence at least 90 % homologous, at least 91 % homologous, at least 92 % homologous, at least 93 % homologous, at least 94 % homologous, at least 95 % homologous, at least 96 % homologous, at least 97 % homologous, at least 98 %
20 homologous, at least 99 % homologous, and more preferably 100 % homologous to the amino acid sequence as set forth in SEQ ID NO: 9.

The agents of this aspect of the present invention are capable of specifically binding LOXL-2 or at least a part of the LOXL-2 protein or DNA or mRNA encoding same. According to a particular embodiment, the agents bind to a region spanning the
25 catalytic site of LOXL-2 (or DNA or mRNA encoding same) and inhibiting its activity when introduced into the mammalian tissue.

According to a particular embodiment, the agents are capable of down-regulating crosslinking of type I collagen in an *in vitro* assay system as further described in Example 1 herein below. Further, the agents are capable of down-regulating
30 crosslinking of pathological (or aberrant) type I collagen *in vivo* following administration. Thus, for example, the present inventors contemplate antibodies which down-regulate the crosslinking activity of type I collagen with an IC_{50} between 1nM-

10 μ M. More preferably between 1nM-1 μ M. According to a specific embodiment, the antibody down-regulates the crosslinking activity of type I collagen with an IC₅₀ with an IC₅₀ of about 1 μ M.

Since the agents of this aspect of the present invention are capable of interfering
5 with type I collagen crosslinking, they alter the structure of the extracellular matrix (ECM), relaxing the collagen fibers, changing them from an ordered orientation to a more random orientation. Thus, the agents of this aspect of the present invention alter the fibrillation of collagen (and accordingly strength of the collagen) without affecting the amount of collagen.

10 Methods of analyzing collagen architecture are known in the art and include for example second harmonic generation (SHG) microscopy, electron microscopy imaging and Raman microscopy.

According to another embodiment, the agents of this aspect of the present invention do not affect angiogenesis *in vivo*.

15 For example, the Cultrex[®] Directed *In Vivo* Angiogenesis Assay (DIVAA[™]) may be performed. During the course of the assay, implant grade silicone cylinders closed at one end, called angioreactors, are filled with basement membrane extract (BME) premixed with or without angiogenesis modulating factors. These angioreactors are then implanted subcutaneously in the dorsal flanks of nude mice. If filled with
20 angiogenic factors, vascular endothelial cells migrate into, and proliferate in the BME to form vessels in the angioreactor. As early as nine days post-implantation, there are enough cells to determine an effective dose response to angiogenic factors.

One example, of an agent capable of downregulating LOXL-2 is an antibody or antibody fragment capable of specifically binding LOXL-2. Preferably, the antibody
25 specifically binds at least one epitope of LOXL-2 (e.g. the catalytic domain). Thus, for example the antibody may bind to residues 545-775 of LOXL-2.

As used herein, the term "binding" or "specifically binding" refers to the binding of the antibody to an epitope of the antigen (either VEGF or ANG2) in an in-vitro assay, e.g. in an ELISA assay or in a surface plasmon resonance assay (BIAcore, GE-
30 Healthcare Uppsala, Sweden or BIO-RAD ProteOn XPR36) with purified wild-type antigen. The affinity of the binding is defined by the terms K_a (rate constant for the

association of the antibody from the antibody/antigen complex), k_D (dissociation constant), and K_D (k_D/k_a).

Preferably, the antibody binds with an affinity less than 10×10^{-9} mol/l, more preferably less than 9×10^{-9} mol/l, more preferably less than 8×10^{-9} mol/l, more preferably less than 7×10^{-9} mol/l, more preferably less than 6×10^{-9} mol/l, more preferably less than 5×10^{-9} mol/l, more preferably less than 4×10^{-9} mol/l, more preferably less than 3×10^{-9} mol/l, more preferably less than 2×10^{-9} mol/l, and more preferably less than 1×10^{-9} mol/l. Preferably the antibody binds with an affinity between preferably 10^{-9} M to 10^{-13} mol/l.

As used herein, the term "epitope" refers to any antigenic determinant on an antigen to which the paratope of an antibody binds.

According to a particular embodiment, the antibody binds the catalytic site of the LOXL-2 protein (situated between residues 545-775) with at least 2 fold higher, 5 fold higher, 10 fold higher, or even 20 fold higher affinity than to the fourth scavenger receptor-cysteine-rich (SRCR) domain of LOXL-2 (situated between residues 435-544).

According to still another embodiment, the antibody doesn't bind to the fourth scavenger receptor-cysteine-rich (SRCR) domain of LOXL-2 (situated between residues 435-544).

Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or carbohydrate side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

The term "antibody" as used in this invention includes intact molecules as well as functional fragments thereof, such as Fab, $F(ab')_2$, and Fv that are capable of binding to macrophages. These functional antibody fragments are defined as follows: (1) Fab, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule, can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) Fab' , the fragment of an antibody molecule that can be obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule; (3) $(Fab')_2$, the fragment of the antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; $F(ab')_2$ is a dimer of two Fab' fragments held together by

two disulfide bonds; (4) Fv, defined as a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (5) Single chain antibody ("SCA"), a genetically engineered molecule containing the variable region of the light chain and the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule.

Methods of producing polyclonal and monoclonal antibodies as well as fragments thereof are well known in the art (See for example, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, 1988, incorporated herein by reference).

Antibody fragments according to some embodiments of the invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* or mammalian cells (e.g. Chinese hamster ovary cell culture or other protein expression systems) of DNA encoding the fragment. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted F(ab')₂. This fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce 3.5S Fab' monovalent fragments. Alternatively, an enzymatic cleavage using pepsin produces two monovalent Fab' fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, U.S. Pat. Nos. 4,036,945 and 4,331,647, and references contained therein, which patents are hereby incorporated by reference in their entirety. See also Porter, R. R. [Biochem. J. 73: 119-126 (1959)]. Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

Fv fragments comprise an association of VH and VL chains. This association may be noncovalent, as described in Inbar et al. [Proc. Nat'l Acad. Sci. USA 69:2659-62 (1972)]. Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. Preferably, the Fv fragments

comprise VH and VL chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a structural gene comprising DNA sequences encoding the VH and VL domains connected by an oligonucleotide. The structural gene is inserted into an expression vector, which is subsequently introduced into a host cell such as *E. coli*. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing sFvs are described, for example, by [Whitlow and Filpula, *Methods* 2: 97-105 (1991); Bird et al., *Science* 242:423-426 (1988); Pack et al., *Bio/Technology* 11:1271-77 (1993); and U.S. Pat. No. 4,946,778, which is hereby incorporated by reference in its entirety.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells. See, for example, Larrick and Fry [*Methods*, 2: 106-10 (1991)].

As used herein "recombinant antibody" refers to intact antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., mouse) that is transgenic for immunoglobulin genes (e.g., human immunoglobulin genes) or hybridoma prepared therefrom; (b) antibodies isolated from a host cell (e.g. prokaryotic cells) transformed to express the antibody; (c) antibodies isolated from a recombinant antibody library; and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of immunoglobulin gene sequences to other DNA sequences. In certain embodiments immunoglobulin of the present invention may have variable and constant regions derived from human germline immunoglobulin sequences. In other embodiments, such recombinant human antibodies can be subjected to in vitro mutagenesis and thus the amino acid sequences of the V_H and V_L regions of the recombinant antibodies comprise sequences that while derived from and related to human germline V_H and V_L sequences, may not naturally exist within the human antibody germline repertoire in vivo.

The following exemplary embodiments of antibodies (both monospecific and bispecific) are encompassed by the scope of the invention.

As used herein "human antibody" refers to intact antibodies having variable regions in which both the framework and CDR regions are derived from human immunoglobulin sequences as described, for example, by Kabat et al. (see Kabat 1991, Sequences of proteins of immunological Interest, 5th Ed. NIH Publication No. 91-3242).

5 The constant region of the human antibody is also described from human immunoglobulin sequences. The human antibodies may include amino residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site directed mutagenesis in vitro or somatic mutation in vivo). However, the term "human antibody", as used herein, is not intended to include antibodies in which
10 CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

As used herein, a "chimeric antibody" refers to an intact antibody in which the variable regions derive from a first species and the constant regions are derived from a second species. Chimeric immunoglobulins can be constructed by genetic engineering
15 from immunoglobulin gene segments belonging to different species (e.g., VH and VL domains from a mouse antibody with constant domains of human origin).

As used herein "humanized immunoglobulin" refers to an intact antibody in which the minimum mouse part from a non-human (e.g., murine) antibody is transplanted onto a human antibody; generally humanized antibodies are 5-10 % mouse
20 and 90-95 % human.

In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The
25 humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

Humanized forms of non-human (e.g., murine) antibodies are chimeric
30 molecules of immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab').sub.2 or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized

antibodies include human immunoglobulins (recipient antibody) in which residues form a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as import residues, which are typically taken from an import variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature* 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such humanized antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and

Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introduction of human immunoglobulin loci into transgenic animals, e.g., mice
5 in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following
10 scientific publications: Marks et al., *Bio/Technology* 10,: 779-783 (1992); Lonberg et al., *Nature* 368: 856-859 (1994); Morrison, *Nature* 368 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14: 826 (1996); and Lonberg and Huszar, *Intern. Rev. Immunol.* 13, 65-93 (1995).

According to specific embodiments, the antibody of this aspect of the present
15 invention comprises CDR sequences as set forth in SEQ ID NOs: 3-8.

According to other embodiments, the amino acid sequence of the V_H of the antibody is at least 90 % identical to SEQ ID NO: 1, more preferably at least 91 % identical to SEQ ID NO: 1, more preferably at least 92 % identical to SEQ ID NO: 1, more preferably at least 93 % identical to SEQ ID NO: 1, more preferably at least 94 %
20 identical to SEQ ID NO: 1, more preferably at least 95 % identical to SEQ ID NO: 1, more preferably at least 96 % identical to SEQ ID NO: 1, more preferably at least 97 % identical to SEQ ID NO: 1, more preferably at least 98 % identical to SEQ ID NO: 1, more preferably at least 99 % identical to SEQ ID NO: 1, more preferably 100 % identical as determined using the Standard protein-protein BLAST [blastp] software of
25 the NCBI to the sequence as set forth in SEQ ID NO: 1, wherein the amino acids of the CDR regions are not replaced by other amino acids.

According to other embodiments, the amino acid sequence of the V_L of the antibody is at least 90 % homologous to SEQ ID NO: 2, more preferably at least 91 % homologous to SEQ ID NO: 2, more preferably at least 92 % homologous to SEQ ID
30 NO: 2, more preferably at least 93 % homologous to SEQ ID NO: 2, more preferably at least 94 % homologous to SEQ ID NO: 2, more preferably at least 95 % homologous to SEQ ID NO: 2, more preferably at least 96 % homologous to SEQ ID NO: 2, more

preferably at least 97 % homologous to SEQ ID NO: 2, more preferably at least 98 % homologous to SEQ ID NO: 2, more preferably at least 99 % homologous to SEQ ID NO: 2, more preferably 100 % homologous as determined using the Standard protein-protein BLAST [blastp] software of the NCBI to the sequence as set forth in SEQ ID NO: 2, wherein the amino acids of the CDR regions are not replaced by other amino acids.

The antibodies of the present invention may be conjugated to a functional moiety such as a detectable or a therapeutic moiety.

Various types of detectable or reporter moieties may be conjugated to the antibody of the invention. These include, but not are limited to, a radioactive isotope (such as ^[125]iodine), a phosphorescent chemical, a chemiluminescent chemical, a fluorescent chemical (fluorophore), an enzyme, a fluorescent polypeptide, an affinity tag, and molecules (contrast agents) detectable by Positron Emission Tomography (PET) or Magnetic Resonance Imaging (MRI).

Table 1 provides non-limiting examples of identifiable moieties which can be conjugated to the antibody of the invention.

Table 1

<i>Nucleic Acid sequence (GenBank Accession No.)</i>	<i>Amino Acid sequence (GenBank Accession No.)</i>	<i>Identifiable Moiety</i>
AF435427	AAL33912	Green Fluorescent protein
AY042185	AAK73766	Alkaline phosphatase
A00740	CAA00083	Peroxidase
Nucleotides 790-807 of GenBank Accession No. AF329457	Amino acids 264-269 of GenBank Accession No. AAK09208	Histidine tag
Nucleotides 817-849 of GenBank Accession No. AF329457	Amino acids 273-283 of GenBank Accession No. AAK09208	Myc tag
		Biotin lygase tag
AF435432	AAL33917	orange fluorescent protein
EU626139	ACH42114	Beta galactosidase
AF283893	AAM49066	Streptavidin

As mentioned, the antibody may be conjugated to a therapeutic moiety. The therapeutic moiety can be, for example, a cytotoxic moiety, a toxic moiety, a cytokine moiety and a second antibody moiety comprising a different specificity to the antibodies of the invention.

Non-limiting examples of therapeutic moieties which can be conjugated to the antibody of the invention are provided in Table 2, hereinbelow.

Table 2

<i>Nucleic acid sequence (GenBank Accession No.)</i>	<i>Amino acid sequence (GenBank Accession No.)</i>	<i>Therapeutic moiety</i>
EU090068	ABU63124	Pseudomonas exotoxin
AY820132.1	AAV70486	Diphtheria toxin
A02159	CAA00227	interleukin 2
X03884	P07766	CD3
NM_000569.6	NP_000560.5	CD16
NM_000589.2	NP_000580.1	interleukin 4
K02883	P01892	HLA-A2
M57627	P22301	interleukin 10
EQ975183	EEF27734	Ricin toxin

The functional moiety may be conjugated to the V_H or the V_L sequence at either the N- or C-terminus or be inserted into other protein sequences in a suitable position.

Downregulation of LOXL-2 can be also achieved by RNA silencing. As used herein, the phrase "RNA silencing" refers to a group of regulatory mechanisms [e.g. RNA interference (RNAi), transcriptional gene silencing (TGS), post-transcriptional gene silencing (PTGS), quelling, co-suppression, and translational repression] mediated by RNA molecules which result in the inhibition or "silencing" of the expression of a corresponding protein-coding gene. RNA silencing has been observed in many types of organisms, including plants, animals, and fungi.

As used herein, the term "RNA silencing agent" refers to an RNA which is capable of specifically inhibiting or "silencing" the expression of a target gene. In

certain embodiments, the RNA silencing agent is capable of preventing complete processing (e.g. the full translation and/or expression) of an mRNA molecule through a post-transcriptional silencing mechanism. RNA silencing agents include noncoding RNA molecules, for example RNA duplexes comprising paired strands, as well as precursor RNAs from which such small non-coding RNAs can be generated. Exemplary RNA silencing agents include dsRNAs such as siRNAs, miRNAs and shRNAs. In one embodiment, the RNA silencing agent is capable of inducing RNA interference. In another embodiment, the RNA silencing agent is capable of mediating translational repression.

According to an embodiment of the invention, the RNA silencing agent is specific to the target RNA (e.g., SEQ ID NO: 9) and does not cross inhibit or silence a gene or a splice variant which exhibits 99% or less global homology to the target gene, e.g., less than 98%, 97%, 96%, 95%, 94%, 93%, 92%, 91%, 90%, 89%, 88%, 87%, 86%, 85%, 84%, 83%, 82%, 81% global homology to the target gene.

RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs). The corresponding process in plants is commonly referred to as post-transcriptional gene silencing or RNA silencing and is also referred to as quelling in fungi. The process of post-transcriptional gene silencing is thought to be an evolutionarily-conserved cellular defense mechanism used to prevent the expression of foreign genes and is commonly shared by diverse flora and phyla. Such protection from foreign gene expression may have evolved in response to the production of double-stranded RNAs (dsRNAs) derived from viral infection or from the random integration of transposon elements into a host genome via a cellular response that specifically destroys homologous single-stranded RNA or viral genomic RNA.

The presence of long dsRNAs in cells stimulates the activity of a ribonuclease III enzyme referred to as dicer. Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs). Short interfering RNAs derived from dicer activity are typically about 21 to about 23 nucleotides in length and comprise about 19 base pair duplexes. The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence

complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex.

Accordingly, some embodiments of the invention contemplate use of dsRNA to
5 downregulate protein expression from mRNA.

According to one embodiment, the dsRNA is greater than 30 bp. The use of long dsRNAs (i.e. dsRNA greater than 30 bp) has been very limited owing to the belief that these longer regions of double stranded RNA will result in the induction of the interferon and PKR response. However, the use of long dsRNAs can provide numerous
10 advantages in that the cell can select the optimal silencing sequence alleviating the need to test numerous siRNAs; long dsRNAs will allow for silencing libraries to have less complexity than would be necessary for siRNAs; and, perhaps most importantly, long dsRNA could prevent viral escape mutations when used as therapeutics.

Various studies demonstrate that long dsRNAs can be used to silence gene
15 expression without inducing the stress response or causing significant off-target effects - see for example [Strat et al., *Nucleic Acids Research*, 2006, Vol. 34, No. 13 3803–3810; Bhargava A et al. *Brain Res. Protoc.* 2004;13:115–125; Diallo M., et al., *Oligonucleotides*. 2003;13:381–392; Paddison P.J., et al., *Proc. Natl Acad. Sci. USA*. 2002;99:1443–1448; Tran N., et al., *FEBS Lett.* 2004;573:127–134].

In particular, the invention according to some embodiments thereof
20 contemplates introduction of long dsRNA (over 30 base transcripts) for gene silencing in cells where the interferon pathway is not activated (e.g. embryonic cells and oocytes) see for example Billy et al., *PNAS* 2001, Vol 98, pages 14428-14433. and Diallo et al., *Oligonucleotides*, October 1, 2003, 13(5): 381-392. doi:10.1089/154545703322617069.

The invention according to some embodiments thereof also contemplates
25 introduction of long dsRNA specifically designed not to induce the interferon and PKR pathways for down-regulating gene expression. For example, Shinagwa and Ishii [*Genes & Dev.* 17 (11): 1340-1345, 2003] have developed a vector, named pDECAP, to express long double-strand RNA from an RNA polymerase II (Pol II) promoter.
30 Because the transcripts from pDECAP lack both the 5'-cap structure and the 3'-poly(A) tail that facilitate ds-RNA export to the cytoplasm, long ds-RNA from pDECAP does not induce the interferon response.

Another method of evading the interferon and PKR pathways in mammalian systems is by introduction of small inhibitory RNAs (siRNAs) either via transfection or endogenous expression.

The term "siRNA" refers to small inhibitory RNA duplexes (generally between 18-30 basepairs) that induce the RNA interference (RNAi) pathway. Typically, siRNAs are chemically synthesized as 21mers with a central 19 bp duplex region and symmetric 2-base 3'-overhangs on the termini, although it has been recently described that chemically synthesized RNA duplexes of 25-30 base length can have as much as a 100-fold increase in potency compared with 21mers at the same location. The observed increased potency obtained using longer RNAs in triggering RNAi is theorized to result from providing Dicer with a substrate (27mer) instead of a product (21mer) and that this improves the rate or efficiency of entry of the siRNA duplex into RISC.

It has been found that position of the 3'-overhang influences potency of an siRNA and asymmetric duplexes having a 3'-overhang on the antisense strand are generally more potent than those with the 3'-overhang on the sense strand (Rose et al., 2005). This can be attributed to asymmetrical strand loading into RISC, as the opposite efficacy patterns are observed when targeting the antisense transcript.

The strands of a double-stranded interfering RNA (e.g., an siRNA) may be connected to form a hairpin or stem-loop structure (e.g., an shRNA). Thus, as mentioned the RNA silencing agent of some embodiments of the invention may also be a short hairpin RNA (shRNA).

The term "shRNA", as used herein, refers to an RNA agent having a stem-loop structure, comprising a first and second region of complementary sequence, the degree of complementarity and orientation of the regions being sufficient such that base pairing occurs between the regions, the first and second regions being joined by a loop region, the loop resulting from a lack of base pairing between nucleotides (or nucleotide analogs) within the loop region. The number of nucleotides in the loop is a number between and including 3 to 23, or 5 to 15, or 7 to 13, or 4 to 9, or 9 to 11. Some of the nucleotides in the loop can be involved in base-pair interactions with other nucleotides in the loop. Examples of oligonucleotide sequences that can be used to form the loop include 5'-UUCAAGAGA-3' (Brummelkamp, T. R. et al. (2002) Science 296: 550) and 5'-UUUGUGUAG-3' (Castanotto, D. et al. (2002) RNA 8:1454). It will be recognized

by one of skill in the art that the resulting single chain oligonucleotide forms a stem-loop or hairpin structure comprising a double-stranded region capable of interacting with the RNAi machinery.

Synthesis of RNA silencing agents suitable for use with some embodiments of the invention can be effected as follows. First, the LOXL-2 mRNA sequence is scanned downstream of the AUG start codon for AA dinucleotide sequences. Occurrence of each AA and the 3' adjacent 19 nucleotides is recorded as potential siRNA target sites. Preferably, siRNA target sites are selected from the open reading frame, as untranslated regions (UTRs) are richer in regulatory protein binding sites. UTR-binding proteins and/or translation initiation complexes may interfere with binding of the siRNA endonuclease complex [Tuschl ChemBiochem. 2:239-245]. It will be appreciated though, that siRNAs directed at untranslated regions may also be effective, as demonstrated for GAPDH wherein siRNA directed at the 5' UTR mediated about 90 % decrease in cellular GAPDH mRNA and completely abolished protein level.

Second, potential target sites are compared to an appropriate genomic database (e.g., human, mouse, rat etc.) using any sequence alignment software, such as the BLAST software available from the NCBI server. Putative target sites which exhibit significant homology to other coding sequences are filtered out.

Qualifying target sequences are selected as template for siRNA synthesis. Preferred sequences are those including low G/C content as these have proven to be more effective in mediating gene silencing as compared to those with G/C content higher than 55 %. Several target sites are preferably selected along the length of the target gene for evaluation. For better evaluation of the selected siRNAs, a negative control is preferably used in conjunction. Negative control siRNA preferably include the same nucleotide composition as the siRNAs but lack significant homology to the genome. Thus, a scrambled nucleotide sequence of the siRNA is preferably used, provided it does not display any significant homology to any other gene.

For example, a suitable LOXL-2 siRNA can be the siRNA disclosed in US Patent Application No. 20130022617, incorporated herein by reference.

It will be appreciated that the RNA silencing agent of some embodiments of the invention need not be limited to those molecules containing only RNA, but further encompasses chemically-modified nucleotides and non-nucleotides.

In some embodiments, the RNA silencing agent provided herein can be functionally associated with a cell-penetrating peptide." As used herein, a "cell-penetrating peptide" is a peptide that comprises a short (about 12-30 residues) amino acid sequence or functional motif that confers the energy-independent (i.e., non-
5 endocytotic) translocation properties associated with transport of the membrane-permeable complex across the plasma and/or nuclear membranes of a cell. The cell-penetrating peptide used in the membrane-permeable complex of some embodiments of the invention preferably comprises at least one non-functional cysteine residue, which is either free or derivatized to form a disulfide link with a double-stranded ribonucleic acid
10 that has been modified for such linkage. Representative amino acid motifs conferring such properties are listed in U.S. Pat. No. 6,348,185, the contents of which are expressly incorporated herein by reference. The cell-penetrating peptides of some embodiments of the invention preferably include, but are not limited to, penetratin, transportan, pIsl, TAT(48-60), pVEC, MTS, and MAP.

15 mRNAs to be targeted using RNA silencing agents include, but are not limited to, those whose expression is correlated with an undesired phenotypic trait. Exemplary mRNAs that may be targeted are those that encode truncated proteins i.e. comprise deletions. Accordingly the RNA silencing agent of some embodiments of the invention may be targeted to a bridging region on either side of the deletion. Introduction of such
20 RNA silencing agents into a cell would cause a down-regulation of the mutated protein while leaving the non-mutated protein unaffected.

According to another embodiment the RNA silencing agent may be a miRNA or miRNA mimic.

The term "microRNA", "miRNA", and "miR" are synonymous and refer to a
25 collection of non-coding single-stranded RNA molecules of about 19-28 nucleotides in length, which regulate gene expression. miRNAs are found in a wide range of organisms and have been shown to play a role in development, homeostasis, and disease etiology.

The term "microRNA mimic" refers to synthetic non-coding RNAs that are
30 capable of entering the RNAi pathway and regulating gene expression. miRNA mimics imitate the function of endogenous microRNAs (miRNAs) and can be designed as mature, double stranded molecules or mimic precursors (e.g., or pre-miRNAs). miRNA

mimics can be comprised of modified or unmodified RNA, DNA, RNA-DNA hybrids, or alternative nucleic acid chemistries (e.g., LNAs or 2'-O,4'-C-ethylene-bridged nucleic acids (ENA)). For mature, double stranded miRNA mimics, the length of the duplex region can vary between 13-33, 18-24 or 21-23 nucleotides. The miRNA may also
5 comprise a total of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 nucleotides. The sequence of the miRNA may be the first 13-33 nucleotides of the pre-miRNA. The sequence of the miRNA may also be the last 13-33 nucleotides of the pre-miRNA.

Another agent capable of downregulating LOXL-2 is a DNAzyme molecule
10 capable of specifically cleaving an mRNA transcript or DNA sequence of the LOXL-2. DNAzymes are single-stranded polynucleotides which are capable of cleaving both single and double stranded target sequences (Breaker, R.R. and Joyce, G. Chemistry and Biology 1995;2:655; Santoro, S.W. & Joyce, G.F. Proc. Natl. Acad. Sci. USA 1997;94:4262). A general model (the "10-23" model) for the DNAzyme has been
15 proposed. "10-23" DNAzymes have a catalytic domain of 15 deoxyribonucleotides, flanked by two substrate-recognition domains of seven to nine deoxyribonucleotides each. This type of DNAzyme can effectively cleave its substrate RNA at purine:pyrimidine junctions (Santoro, S.W. & Joyce, G.F. Proc. Natl. Acad. Sci. USA 199; for rev of DNAzymes see Khachigian, LM [Curr Opin Mol Ther 4:119-21 (2002)]).

20 Examples of construction and amplification of synthetic, engineered DNAzymes recognizing single and double-stranded target cleavage sites have been disclosed in U.S. Pat. No. 6,326,174 to Joyce et al. DNAzymes of similar design directed against the human Urokinase receptor were recently observed to inhibit Urokinase receptor expression, and successfully inhibit colon cancer cell metastasis in vivo (Itoh et al, 25 20002, Abstract 409, Ann Meeting Am Soc Gen Ther www.asgt.org). In another application, DNAzymes complementary to bcr-abl oncogenes were successful in inhibiting the oncogenes expression in leukemia cells, and lessening relapse rates in autologous bone marrow transplant in cases of CML and ALL.

Downregulation of LOXL-2 can also be effected by using an antisense
30 polynucleotide capable of specifically hybridizing with an mRNA transcript encoding the LOXL-2.

Design of antisense molecules which can be used to efficiently downregulate a LOXL-2 must be effected while considering two aspects important to the antisense approach. The first aspect is delivery of the oligonucleotide into the cytoplasm of the appropriate cells, while the second aspect is design of an oligonucleotide which specifically binds the designated mRNA within cells in a way which inhibits translation thereof.

The prior art teaches of a number of delivery strategies which can be used to efficiently deliver oligonucleotides into a wide variety of cell types [see, for example, Luft J Mol Med 76: 75-6 (1998); Kronenwett et al. Blood 91: 852-62 (1998); Rajur et al. Bioconjug Chem 8: 935-40 (1997); Lavigne et al. Biochem Biophys Res Commun 237: 566-71 (1997) and Aoki et al. (1997) Biochem Biophys Res Commun 231: 540-5 (1997)].

In addition, algorithms for identifying those sequences with the highest predicted binding affinity for their target mRNA based on a thermodynamic cycle that accounts for the energetics of structural alterations in both the target mRNA and the oligonucleotide are also available [see, for example, Walton et al. Biotechnol Bioeng 65: 1-9 (1999)].

Such algorithms have been successfully used to implement an antisense approach in cells. For example, the algorithm developed by Walton et al. enabled scientists to successfully design antisense oligonucleotides for rabbit beta-globin (RBG) and mouse tumor necrosis factor-alpha (TNF alpha) transcripts. The same research group has more recently reported that the antisense activity of rationally selected oligonucleotides against three model target mRNAs (human lactate dehydrogenase A and B and rat gp130) in cell culture as evaluated by a kinetic PCR technique proved effective in almost all cases, including tests against three different targets in two cell types with phosphodiester and phosphorothioate oligonucleotide chemistries.

In addition, several approaches for designing and predicting efficiency of specific oligonucleotides using an in vitro system were also published (Matveeva et al., Nature Biotechnology 16: 1374 - 1375 (1998)).

Another agent capable of downregulating LOXL-2 is a ribozyme molecule capable of specifically cleaving an mRNA transcript encoding LOXL-2. Ribozymes are being increasingly used for the sequence-specific inhibition of gene expression by the

cleavage of mRNAs encoding proteins of interest [Welch et al., Curr Opin Biotechnol. 9:486-96 (1998)]. The possibility of designing ribozymes to cleave any specific target RNA has rendered them valuable tools in both basic research and therapeutic applications. In the therapeutics area, ribozymes have been exploited to target viral
 5 RNAs in infectious diseases, dominant oncogenes in cancers and specific somatic mutations in genetic disorders [Welch et al., Clin Diagn Virol. 10:163-71 (1998)].

An additional method of regulating the expression of a LOXL-2 gene in cells is via triplex forming oligonucleotides (TFOs). Recent studies have shown that TFOs can be designed which can recognize and bind to polypurine/polypyrimidine regions in
 10 double-stranded helical DNA in a sequence-specific manner. These recognition rules are outlined by Maher III, L. J., et al., Science, 1989; 245:725-730; Moser, H. E., et al., Science, 1987; 238:645-630; Beal, P. A., et al., Science, 1992; 251:1360-1363; Cooney, M., et al., Science, 1988; 241:456-459; and Hogan, M. E., et al., EP Publication 375408. Modification of the oligonucleotides, such as the introduction of intercalators and
 15 backbone substitutions, and optimization of binding conditions (pH and cation concentration) have aided in overcoming inherent obstacles to TFO activity such as charge repulsion and instability, and it was recently shown that synthetic oligonucleotides can be targeted to specific sequences (for a recent review see Seidman and Glazer, J Clin Invest 2003; 112:487-94).

20 In general, the triplex-forming oligonucleotide has the sequence correspondence:

oligo	3'--A	G	G	T
duplex	5'--A	G	C	T
duplex	3'--T	C	G	A

However, it has been shown that the A-AT and G-GC triplets have the greatest
 25 triple helical stability (Reither and Jeltsch, BMC Biochem, 2002, Sept12, Epub). The same authors have demonstrated that TFOs designed according to the A-AT and G-GC rule do not form non-specific triplexes, indicating that the triplex formation is indeed sequence specific.

Thus for any given sequence in the LOXL-2 regulatory region a triplex forming
 30 sequence may be devised. Triplex-forming oligonucleotides preferably are at least 15, more preferably 25, still more preferably 30 or more nucleotides in length, up to 50 or 100 bp.

Transfection of cells (for example, via cationic liposomes) with TFOs, and formation of the triple helical structure with the target DNA induces steric and functional changes, blocking transcription initiation and elongation, allowing the introduction of desired sequence changes in the endogenous DNA and resulting in the specific downregulation of gene expression. Examples of such suppression of gene expression in cells treated with TFOs include knockout of episomal supFG1 and endogenous HPRT genes in mammalian cells (Vasquez et al., Nucl Acids Res. 1999;27:1176-81, and Puri, et al, J Biol Chem, 2001;276:28991-98), and the sequence- and target specific downregulation of expression of the Ets2 transcription factor, important in prostate cancer etiology (Carbone, et al, Nucl Acid Res. 2003;31:833-43), and the pro-inflammatory ICAM-1 gene (Besch et al, J Biol Chem, 2002;277:32473-79). In addition, Vuyisich and Beal have recently shown that sequence specific TFOs can bind to dsRNA, inhibiting activity of dsRNA-dependent enzymes such as RNA-dependent kinases (Vuyisich and Beal, Nuc. Acids Res 2000;28:2369-74).

Additionally, TFOs designed according to the abovementioned principles can induce directed mutagenesis capable of effecting DNA repair, thus providing both downregulation and upregulation of expression of endogenous genes (Seidman and Glazer, J Clin Invest 2003;112:487-94). Detailed description of the design, synthesis and administration of effective TFOs can be found in U.S. Patent Application Nos. 2003017068 and 20030096980 to Froehler et al, and 20020128218 and 20020123476 to Emanuele et al, and U.S. Pat. No. 5,721,138 to Lawn.

It will be appreciated that since the agent which targets LOXL-2 serves to relax the collagen fibers of the ECM, the present inventors contemplate administering the agent together with an additional agent, whereby the LOXL-2 targeting agent allows the additional agent to gain better access to the site of the disease (e.g. tumor or fibrosis). In this way, it is anticipated that the dose of the additional agent may be lower than the commonly administered dose (i.e. is lower than its effective clinical concentration when administered alone) and accordingly the LOXL-2 targeting agent may serve to lower the side-effects of the additional agent.

In one preferred embodiment, the amount provided of the additional agent of the invention is below the minimum dose required for therapeutic, prophylactic and/or pain palliative effectiveness when used as a single therapy (e.g. 10-99%, preferably 25 to

75% of that minimum dose). This allows for reduction of the side effects caused by the additional agent but the therapy is rendered effective because in combination with the LOXL-2 targeting agent, the additional agents of the invention are effective overall. In a further preferred embodiment the anti-LOXL-2 agents is used at a level below the minimum normal therapeutic dose that brings about angiogenesis, for example 10-99% of the normal minimum therapeutic dose, preferably 25 to 75% of their normal therapeutic dose. This again serves to reduce the danger of side effects and allows a higher level of total effectiveness without exposing the subject to unacceptable side effects.

In one preferred aspect of the present invention, the anti-LOXL-2 agent and the at least one other additional agent of the invention are synergistic with respect to their dosages. That is to say that the effect provided by the anti-LOXL-1 agent of the present invention is greater than would be anticipated from the additive effects of the incorporated doses of anti-LOXL-2 and at least one other additional agent when used separately. In an alternative but equally preferred embodiment, the anti-LOXL-2 agent and the at least one additional agent are synergistic with respect to their side effects. That is to say that the side-effects caused by agents of the present invention are less than would be anticipated when the equivalent therapeutic effect is provided by either anti-LOXL-2 agent or by the additional agent when used separately.

According to embodiments of the present invention, the additional agent is an anti-cancer agent.

According to one embodiment, the anti-cancer agent comprises a chemotherapeutic agent.

Examples of chemotherapeutic agents include but are not limited to abarelix, aldesleukin, aldesleukin, alemtuzumab, alitretinoin, allopurinol, altretamine, amifostine, anastrozole, arsenic trioxide, asparaginase, azacitidine, bevacuzimab, bexarotene, bleomycin, bortezomib, busulfan, calusterone, capecitabine, carboplatin, carmustine, celecoxib, cetuximab, cisplatin, cladribine, clofarabine, cyclophosphamide, cytarabine, dacarbazine, dactinomycin, actinomycin D, Darbepoetin alfa, Darbepoetin alfa, daunorubicin liposomal, daunorubicin, decitabine, Denileukin diftitox, dexrazoxane, dexrazoxane, docetaxel, doxorubicin, dromostanolone propionate, Elliott's B Solution, epirubicin, Epoetin alfa, erlotinib, estramustine, etoposide, exemestane,

Filgrastim, floxuridine, fludarabine, fluorouracil 5-FU, fulvestrant, gefitinib, gemcitabine, gemtuzumab ozogamicin, goserelin acetate, histrelin acetate, hydroxyurea, Ibritumomab Tiuxetan, idarubicin, ifosfamide, imatinib mesylate, interferon alfa 2a, Interferon alfa-2b, irinotecan, lenalidomide, letrozole, leucovorin, Leuprolide Acetate, levamisole, lomustine, CCNU, mecllorethamine, nitrogen mustard, megestrol acetate, melphalan, L-PAM, mercaptopurine 6-MP, mesna, methotrexate, mitomycin C, mitotane, mitoxantrone, nandrolone phenpropionate, nelarabine, Nofetumomab, Oprelvekin, Oprelvekin, oxaliplatin, paclitaxel, palifermin, pamidronate, pegademase, pegaspargase, Pegfilgrastim, pemetrexed disodium, pentostatin, pipobroman, plicamycin mithramycin, porfimer sodium, procarbazine, quinacrine, Rasburicase, Rituximab, sargramostim, sorafenib, streptozocin, sunitinib maleate, tamoxifen, temozolomide, teniposide VM-26, testolactone, thioguanine 6-TG, thiotepa, thiotepa, topotecan, toremifene, Tositumomab, Trastuzumab, tretinoin ATRA, Uracil Mustard, valrubicin, vinblastine, vinorelbine, zoledronate and zoledronic acid.

According to a particular embodiment, the anti-cancer agent is cisplatin.

According to another embodiment, the anti-cancer agent is an antibody.

Examples of antibodies that may be used in the context of the present invention include those listed in Table 3 herein below. It will be appreciated that the present invention contemplates additional antibodies which bind to the targets listed herein below.

Table 3

Name	Target	Use
3F8	GD2	neuroblastoma
8H9	B7-H3	neuroblastoma, sarcoma, metastatic brain cancers
Abagovomab	CA-125 (imitation)	ovarian cancer
Abciximab	CD41 (integrin alpha-IIb)	platelet aggregation inhibitor
Actoxumab	<i>Clostridium difficile</i>	<i>Clostridium difficile</i> infection
Adalimumab	TNF- α	Rheumatoid arthritis, Crohn's Disease, Plaque Psoriasis, Psoriatic Arthritis, Ankylosing Spondylitis, Juvenile Idiopathic Arthritis, Hemolytic disease of the newborn

Name	Target	Use
Adecatumumab	EpCAM	prostate and breast cancer
Aducanumab	beta-amyloid	Alzheimer's disease
Afelimomab	TNF- α	sepsis
Afutuzumab	CD20	lymphoma
Alacizumab pegol	VEGFR2	cancer
ALD518	IL-6	rheumatoid arthritis
Alemtuzumab	CD52	Multiple sclerosis
Alirocumab	NARP-1	hypercholesterolemia
Altumomab pentetate	CEA	colorectal cancer (diagnosis)
Amatuximab	mesothelin	cancer
Anatumomab mafenatox	TAG-72	non-small cell lung carcinoma
Anifrolumab	interferon α/β receptor	systemic lupus erythematosus
Anrukinzumab (= IMA-638)	IL-13	
Apolizumab	HLA-DR	hematological cancers
Arcitumomab	CEA	gastrointestinal cancers (diagnosis)
Aselizumab	L-selectin (CD62L)	severely injured patients
Atinumab	RTN4	
Atlizumab (= tocilizumab)	IL-6 receptor	rheumatoid arthritis
Atorolimumab	Rhesus factor	hemolytic disease of the newborn
Bapineuzumab	beta amyloid	Alzheimer's disease
Basiliximab	CD25 (α chain of IL-2 receptor)	prevention of organ transplant rejections
Bavituximab	phosphatidylserine	cancer, viral infections
Bectumomab	CD22	non-Hodgkin's lymphoma (detection)
Belimumab	BAFF	non-Hodgkin lymphoma etc.
Benralizumab	CD125	asthma
Bertilimumab	CCL11 (eotaxin-1)	severe allergic disorders
Besilesomab	CEA-related antigen	inflammatory lesions and metastases (detection)
Bevacizumab	VEGF-A	metastatic cancer, retinopathy of prematurity
Bezlotoxumab	<i>Clostridium difficile</i>	<i>Clostridium difficile</i> infection
Biciromab	fibrin II, beta chain	thromboembolism (diagnosis)

Name	Target	Use
Bimagrumab	ACVR2B	myostatin inhibitor
Bivatuzumab mertansine	CD44 v6	squamous cell carcinoma
Blinatumomab	CD19	cancer
Blosozumab	SOST	osteoporosis
Brentuximab vedotin	CD30 (TNFRSF8)	hematologic cancers
Briakinumab	IL-12, IL-23	psoriasis, rheumatoid arthritis, inflammatory bowel diseases, multiple sclerosis
Brodalumab	IL-17	inflammatory diseases
Canakinumab	IL-1	rheumatoid arthritis
Cantuzumab mertansine	mucin CanAg	colorectal cancer
Cantuzumab ravtansine	MUC1	cancers
Caplacizumab	VWF	
Capromab pendetide	prostatic carcinoma cells	prostate cancer (detection)
Carlumab	MCP-1	oncology/immune indications
Catumaxomab	EpCAM, CD3	ovarian cancer, malignant ascites, gastric cancer
CC49	TAG-72	tumor detection
cBR96-doxorubicin immunoconjugate	Lewis-Y antigen	cancer
Cedelizumab	CD4	prevention of organ transplant rejections, treatment of autoimmune diseases
Certolizumab pegol	TNF- α	Crohn's disease
Cetuximab	EGFR	metastatic colorectal cancer and head and neck cancer
Ch.14.18		neuroblastoma
Citatumumab bogatox	EpCAM	ovarian cancer and other solid tumors
Cixutumumab	IGF-1 receptor	solid tumors
Clazakizumab	<i>Oryctolagus cuniculus</i>	rheumatoid arthritis
Clenoliximab	CD4	rheumatoid arthritis
Clivatuzumab tetraxetan	MUC1	pancreatic cancer

Name	Target	Use
Conatumumab	TRAIL-R2	cancer
Concizumab	TFPI	bleeding
Crenezumab	1-40- β -amyloid	Alzheimer's disease
CR6261	Influenza A hemagglutinin	infectious disease/influenza A
Dacetuzumab	CD40	hematologic cancers
Daclizumab	CD25 (α chain of IL-2 receptor)	prevention of organ transplant rejections
Dalotuzumab	insulin-like growth factor I receptor	cancer etc.
Daratumumab	CD38 (cyclic ADP ribose hydrolase)	
Demcizumab	DLL4	cancer
Denosumab	RANKL	osteoporosis, bone metastases etc.
Detumomab	B-lymphoma cell	lymphoma
Dorlimomab aritox		
Drozitumab	DR5	cancer
Duligotumab	HER3	
Dupilumab	IL4	atopic diseases
Dusigitumab	ILGF2	cancer
Ecromeximab	GD3 ganglioside	malignant melanoma
Eculizumab	C5	paroxysmal nocturnal hemoglobinuria
Edobacomab	endotoxin	sepsis caused by Gram-negative bacteria
Edrecolomab	EpCAM	colorectal carcinoma
Efalizumab	LFA-1 (CD11a)	psoriasis (blocks T-cell migration)
Efungumab	Hsp90	invasive <i>Candida</i> infection
Eldelumab	interferon gamma-induced protein	Crohn's disease, ulcerative colitis
Elotuzumab	SLAMF7	multiple myeloma
Elsilimomab	IL-6	
Enavatuzumab	TWEAK receptor	cancer
Enlimomab pegol	ICAM-1 (CD54)	
Enokizumab	IL9	asthma
Enoticumab	DLL4	
Ensituximab	5AC	cancer
Epitumomab cituxetan	episialin	

Name	Target	Use
Epratuzumab	CD22	cancer, SLE
Erlizumab	ITGB2 (CD18)	heart attack, stroke, traumatic shock
Ertumaxomab	HER2/neu, CD3	breast cancer etc.
Etaracizumab	integrin $\alpha_v\beta_3$	melanoma, prostate cancer, ovarian cancer etc.
Etrolizumab	integrin $\alpha_7\beta_7$	inflammatory bowel disease
Evolocumab	PCSK9	hypcholesterolemia
Exbivirumab	hepatitis B surface antigen	hepatitis B
Fanolesomab	CD15	appendicitis (diagnosis)
Faralimomab	interferon receptor	
Farletuzumab	folate receptor 1	ovarian cancer
Fasinumab	HNGF	
FBTA05	CD20	chronic lymphocytic leukaemia
Felvizumab	respiratory syncytial virus	respiratory syncytial virus infection
Fezakinumab	IL-22	rheumatoid arthritis, psoriasis
Ficlatuzumab	HGF	cancer etc.
Figitumumab	IGF-1 receptor	adrenocortical carcinoma, non-small cell lung carcinoma etc.
Flanvotumab	TYRP1(glycoprotein 75)	melanoma
Fontolizumab	IFN- γ	Crohn's disease etc.
Foralumab	CD3 epsilon	
Foravirumab	rabies virus glycoprotein	rabies (prophylaxis)
Fresolimumab	TGF- β	idiopathic pulmonary fibrosis, focal segmental glomerulosclerosis, cancer
Fulranumab	NGF	pain
Futuximab	EGFR	
Galiximab	CD80	B-cell lymphoma
Ganitumab	IGF-I	cancer
Gantenerumab	beta amyloid	Alzheimer's disease
Gavilimomab	CD147 (basigin)	graft versus host disease
Gemtuzumab ozogamicin	CD33	acute myelogenous leukemia
Gevokizumab	IL-1 β	diabetes.

Name	Target	Use
Girentuximab	carbonic anhydrase 9 (CA-IX)	clear cell renal cell carcinoma ^[64]
Glembatumumab vedotin	GPNMB	melanoma, breast cancer
Golimumab	TNF- α	rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis
Gomiliximab	CD23 (IgE receptor)	allergic asthma
Guselkumab	IL13	psoriasis
Ibalizumab	CD4	HIV infection
Ibritumomab tiuxetan	CD20	non-Hodgkin's lymphoma
Icrucumab	VEGFR-1	cancer etc.
Igovomab	CA-125	ovarian cancer (diagnosis)
IMAB362	CLDN18.2	gastrointestinal adenocarcinomas and pancreatic tumor
Imciromab	cardiac myosin	cardiac imaging
Imgatuzumab	EGFR	cancer
Inclacumab	selectin P	
Indatuximab ravtansine	SDC1	cancer
Infliximab	TNF- α	rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis, psoriasis, Crohn's disease, ulcerative colitis
Intetumumab	CD51	solid tumors (prostate cancer, melanoma)
Inolimomab	CD25 (α chain of IL-2 receptor)	graft versus host disease
Inotuzumab ozogamicin	CD22	cancer
Ipilimumab	CD152	melanoma
Iratumumab	CD30 (TNFRSF8)	Hodgkin's lymphoma
Itolizumab	CD6	
Ixekizumab	IL-17A	autoimmune diseases
Keliximab	CD4	chronic asthma
Labetuzumab	CEA	colorectal cancer
Lambrolizumab	PDCD1	antineoplastic agent
Lampalizumab	CFD	
Lebrikizumab	IL-13	asthma
Lemalesomab	NCA-90 (granulocyte	diagnostic agent

Name	Target	Use
	antigen)	
Lerdelimumab	TGF beta 2	reduction of scarring after glaucoma surgery
Lexatumumab	TRAIL-R2	cancer
Libivirumab	hepatitis B surface antigen	hepatitis B
Ligelizumab	IGHE	
Lintuzumab	CD33	cancer
Lirilumab	KIR2D	
Lodelcizumab	PCSK9	hypercholesterolemia
Lorvotuzumab mertansine	CD56	cancer
Lucatumumab	CD40	multiple myeloma, non-Hodgkin's lymphoma, Hodgkin's lymphoma
Lumiliximab	CD23 (IgE receptor)	chronic lymphocytic leukemia
Mapatumumab	TRAIL-R1	cancer
Margetuximab	ch4D5	cancer
Maslimomab	T-cell receptor	
Mavrilimumab	GMCSF receptor α -chain	rheumatoid arthritis
Matuzumab	EGFR	colorectal, lung and stomach cancer
Mepolizumab	IL-5	asthma and white blood cell diseases
Metelimumab	TGF beta 1	systemic scleroderma
Milatuzumab	CD74	multiple myeloma and other hematological malignancies
Minretumomab	TAG-72	
Mitumomab	GD3 ganglioside	small cell lung carcinoma
Mogamulizumab	CCR4	cancer
Morolimumab	Rhesus factor	
Motavizumab	respiratory syncytial virus	respiratory syncytial virus (prevention)
Moxetumomab pasudotox	CD22	cancer
Muromonab-CD3	CD3	prevention of organ transplant rejections
Nacolumab tafenoatox	C242 antigen	colorectal cancer
Namilumab	CSF2	

Name	Target	Use
Naptumomab estafenatox	5T4	non-small cell lung carcinoma, renal cell carcinoma
Narnatumab	RON	cancer
Natalizumab	integrin α_4	multiple sclerosis, Crohn's disease
Nebacumab	endotoxin	sepsis
Necitumumab	EGFR	non-small cell lung carcinoma
Nerelimomab	TNF- α	
Nesvacumab	angiopoietin 2	cancer
Nimotuzumab	EGFR	squamous cell carcinoma, head and neck cancer, nasopharyngeal cancer, glioma
Nivolumab	IgG4	cancer
Nofetumomab merpentan		cancer (diagnosis)
Ocaratuzumab	CD20	cancer
Ocrelizumab	CD20	rheumatoid arthritis, lupus erythematosus etc.
Odulimomab	LFA-1 (CD11a)	prevention of organ transplant rejections, immunological diseases
Ofatumumab	CD20	chronic lymphocytic leukemia etc.
Olaratumab	PDGF-R α	cancer
Olokizumab	IL6	
Omalizumab	IgE Fc region	allergic asthma
Onartuzumab	human scatter factor receptor kinase	cancer
Ontuxizumab	TEM1	cancer
Oportuzumab monatox	EpCAM	cancer
Oregovomab	CA-125	ovarian cancer
Orticumab	oxLDL	
Otelixizumab	CD3	diabetes mellitus type 1
Otlertuzumab	CD37	cancer
Oxelumab	OX-40	asthma
Ozanezumab	NOGO-A	ALS and multiple sclerosis
Ozoralizumab	TNF- α	inflammation
Pagibaximab	lipoteichoic acid	sepsis (<i>Staphylococcus</i>)
Palivizumab	F protein of respiratory syncytial virus	respiratory syncytial virus (prevention)

Name	Target	Use
Panitumumab	EGFR	colorectal cancer
Pankomab	tumor specific glycosylation of MUC1	ovarian cancer
Panobacumab	<i>Pseudomonas aeruginosa</i>	<i>Pseudomonas aeruginosa</i> infection
Parsatuzumab	EGFL7	cancer
Pascolizumab	IL-4	asthma
Pateclizumab	LTA	TNF
Patritumab	HER3	cancer
Pemtumomab	MUC1	cancer
Perakizumab	IL17A	arthritis
Pertuzumab	HER2/neu	cancer
Pexelizumab	C5	reduction of side effects of cardiac surgery
Pidilizumab	PD-1	cancer and infectious diseases
Pinatuzumab vedotin	CD22	cancer
Pintumomab	adenocarcinoma antigen	adenocarcinoma (imaging)
Placulumab	human TNF	
Polatuzumab vedotin	CD79B	
Ponezumab	human beta-amyloid	Alzheimer's disease
Priliximab	CD4	Crohn's disease, multiple sclerosis
Pritoxaximab	<i>E. coli</i> shiga toxin type-1	
Pritumumab	vimentin	brain cancer
PRO 140	CCR5	HIV infection
Quilizumab	IGHE	
Racotumomab	<i>N</i> -glycolylneuraminic acid	cancer
Radretumab	fibronectin extra domain-B	cancer
Rafivirumab	rabies virus glycoprotein	rabies (prophylaxis)
Ramucirumab	VEGFR2	solid tumors
Ranibizumab	VEGF-A	macular degeneration (wet form)

Name	Target	Use
Raxibacumab	anthrax toxin, protective antigen	anthrax (prophylaxis and treatment)
Regavirumab	cytomegalovirus glycoprotein B	cytomegalovirus infection
Reslizumab	IL-5	inflammations of the airways, skin and gastrointestinal tract
Rilotumumab	HGF	solid tumors
Rituximab	CD20	lymphomas, leukemias, some autoimmune disorders
Robatumumab	IGF-1 receptor	cancer
Roledumab	RHD	
Romosozumab	sclerostin	osteoporosis
Rontalizumab	IFN- α	systemic lupus erythematosus
Rovelizumab	CD11, CD18	haemorrhagic shock etc.
Ruplizumab	CD154 (CD40L)	rheumatic diseases
Samalizumab	CD200	cancer
Sarilumab	IL6	rheumatoid arthritis, ankylosing spondylitis
Satumomab pendetide	TAG-72	cancer (diagnosis)
Secukinumab	IL-17A	uveitis, rheumatoid arthritis psoriasis
Seribantumab	ERBB3	cancer
Setoxaximab	<i>E. coli</i> shiga toxin type-1	
Sevirumab	cytomegalovirus	cytomegalovirus infection
Sibrotuzumab	FAP	cancer
SGN-CD19A	CD19	acute lymphoblastic leukemia and B-cell non-Hodgkin lymphoma
SGN-CD33A	CD33	Acute myeloid leukemia
Sifalimumab	IFN- α	SLE, dermatomyositis, polymyositis
Siltuximab	IL-6	cancer
Simtuzumab	LOXL2	fibrosis
Siplizumab	CD2	psoriasis, graft-versus-host disease (prevention)
Sirukumab	IL-6	rheumatoid arthritis
Solanezumab	beta amyloid	Alzheimer's disease
Solitomab	EpCAM	

Name	Target	Use
Sonepcizumab	sphingosine-1-phosphate	choroidal and retinal neovascularization
Sontuzumab	episialin	
Stamulumab	myostatin	muscular dystrophy
Sulesomab	NCA-90 (granulocyte antigen)	osteomyelitis (imaging)
Suvizumab	HIV-1	viral infections
Tabalumab	BAFF	B-cell cancers
Tacatuzumab tetraxetan	alpha-fetoprotein	cancer
Tadocizumab	integrin $\alpha_{IIb}\beta_3$	percutaneous coronary intervention
Talizumab	IgE	allergic reaction
Tanezumab	NGF	pain
Taplitumomab paptox	CD19	cancer
Tefibazumab	clumping factor A	<i>Staphylococcus aureus</i> infection
Telimomab aritox		
Tenatumomab	tenascin C	cancer
Teneliximab	CD40	
Teplizumab	CD3	diabetes mellitus type 1
Teprotumumab	CD221	hematologic tumors
TGN1412	CD28	chronic lymphocytic leukemia, rheumatoid arthritis
Ticilimumab (= tremelimumab)	CTLA-4	cancer
Tildrakizumab	IL23	immunologically mediated inflammatory disorders
Tigatuzumab	TRAIL-R2	cancer
TNX-650	IL-13	Hodgkin's lymphoma
Tocilizumab (= atlizumab)	IL-6 receptor	rheumatoid arthritis
Toralizumab	CD154 (CD40L)	rheumatoid arthritis, lupus nephritis etc.
Tositumomab	CD20	follicular lymphoma
Tovetumab	CD140a	cancer
Tralokinumab	IL-13	asthma etc.
Trastuzumab	HER2/neu	breast cancer
TRBS07	GD2	melanoma

Name	Target	Use
Tregalizumab	CD4	
Tremelimumab	CTLA-4	cancer
Tucotuzumab celmoleukin	EpCAM	cancer
Tuvirumab	hepatitis B virus	chronic hepatitis B
Ublituximab	MS4A1	cancer
Urelumab	4-1BB	cancer etc.
Urtioxazumab	<i>Escherichia coli</i>	diarrhoea caused by <i>E. coli</i>
Ustekinumab	IL-12, IL-23	multiple sclerosis, psoriasis, psoriatic arthritis
Vantictumab	Frizzled receptor	cancer
Vapaliximab	AOC3 (VAP-1)	
Vatelizumab	ITGA2	
Vedolizumab	integrin $\alpha_4\beta_7$	Crohn's disease, ulcerative colitis
Veltuzumab	CD20	non-Hodgkin's lymphoma
Vepalimomab	AOC3 (VAP-1)	inflammation
Vesencumab	NRP1	
Visilizumab	CD3	Crohn's disease, ulcerative colitis
Volociximab	integrin $\alpha_5\beta_1$	solid tumors
Vorsetuzumab mafodotin	CD70	cancer
Votumumab	tumor antigen CTAA16.88	colorectal tumors
Zalutumumab	EGFR	squamous cell carcinoma of the head and neck
Zanolimumab	CD4	rheumatoid arthritis, psoriasis, T-cell lymphoma
Zatuximab	HER1	cancer
Ziralimumab	CD147 (basigin)	
Zolimomab aritox	CD5	

According to a particular embodiment, the anti-cancer agent is an antibody that targets a matrix metalloproteinase. Methods of generating such antibodies are disclosed in US Patent No. 8,697,078, International Patent Application WO2004/087042 and

5 WO2008/102359, the contents of each are incorporated herein by reference. According

to a particular embodiment, the matrix metalloproteinase is Matrix metalloproteinase 9 (MMP-9).

When the anti-LOXL-2 agent is an antibody and the additional anti-cancer agent is an antibody, the present invention further contemplates combining the two antibodies to produce a bisppecific antibody.

As used herein, the term "bispecific antibody" refers to an antibody which comprises two antigen binding sites, each binding to a different epitope of a different antigen. The bispecific antibodies of this aspect of the present invention do not share common light chains or common heavy chains.

Bispecific antibodies can be produced by many methods known in the art including for example chemical cross-linkage, genetic engineering, or somatic hybridization, as further described herein below.

Methods of generating bispecific antibodies are disclosed in U.S. Patent No. 8,268,314, U.S. Patent Application No. 20140017244 and WO 2012/123949, the entire contents of which are incorporated herein by reference.

The bispecific antibodies according to the invention may be "bivalent" – denoting the presence of two binding sites.

According to another embodiment, the antibodies of this aspect of the present invention are trivalent or tetravalent.

The agents of the present invention (or combination of agents) may be tested in vitro for a therapeutic effect. Thus, for example, the agents of the present invention can be tested for their ability to prevent or retard metastasis of tumor cells.

Testing a metastatic phenotype of transformed tumor cells can be performed in vitro since nearly all steps of the metastatic process, including attachment, matrix degradation and migration, can be modeled experimentally in vitro by measuring invasion of a reconstituted basement membrane (RBM). Metastatic invasiveness of tumor cell can be modeled by migration of tumor cells into reconstituted basement membrane (RBM) in the presence and absence of a chemoattractant, such as fibroblast conditioned medium (FCM). The assay determines cells that have attached to the RBM, degraded the RBM enzymatically and, finally, cells that have penetrated the FCM side of the membrane.

Since in vitro metastasis events correspond to steps observed in the metastatic spread of tumor cells through the basement membrane in vivo, in vitro invasiveness of cells can be assayed by the methods described in Albini et al., 1987 Cancer Research 47: 3239-3245, which is incorporated herein by reference in its entirety. Invasiveness assays and other methods for assessing metastatic affects, are described in Leyton et al., 5 1994 Cancer Research 54: 3696-3699, which is incorporated by reference herein in its entirety. Reconstituted basement membrane preparations for use in accordance with the hereinabove described assays are readily available from numerous commercial suppliers. One suitable example membrane in this regard is "MATRIGEL" available 10 from Collaborative Biomedical Products of Bedford, Mass.

In vitro evaluation of tumor cell metastatic phenotype can also be effected by determining level and pattern of expression of one or more metastasis associated markers such protease markers, which are considered to be an integral part of tumor metastasis (see U.S. Pat. No. 6,303,318). One example is the arachidonic acid, the 15 release of which in cells can serve to indicate metastatic potential of a tumor (U.S. Pat. No. 6,316,416). In this regard, determining phospholipase A-2 (PLA₂) activity, and the activity or abundance of factors that affect the activity of PLA₂, such as uteroglobin protein (U.S. Pat. No. 6,316,416) can serve as an indication of metastatic potential.

According to another embodiment, the additional agent is an antifibrotic agent.

20 The term "antifibrotic agent" refers to one or more chemical or biological compounds that have antifibrotic activity in mammals. These compounds may have different mechanisms of action, some reducing the formation of collagen or another protein, and others enhancing the metabolism or removal of collagen in the affected area of the body. All such compounds having activity in the reduction of the presence of 25 fibrous tissue are included herein, without regard to the particular mechanism of action by which each such drug functions.

An antifibrotic agent, includes, but is not limited to, an agent that degrades or causes the dissolution or shrinkage of fibrotic tissue or a portion thereof, such as for example, the fibrotic tissue in or surrounding the tendon sheaths that pass through the 30 carpal tunnel. In some embodiments, the antifibrotic agent will stretch or relax fibrous tissue (e.g., normal or abnormal fibrous tissue). Abnormal fibrous tissue includes tissue that has an excessive amount or abnormal amount of fibrosis tissue. Normal fibrous

tissue includes tissue that has a normal amount of fibrous tissue that is not pathological or indicative of disease. For example, in carpal tunnel syndrome, in some embodiments, the flexor retinaculum generally has normal amounts of fibrous tissue. However, it is beneficial to degrade this normal fibrous tissue so the flexor retinaculum can stretch more easily to accommodate the excess tissue within the carpal tunnel. In some 5 embodiments, the antifibrotic agent can degrade, stretch, and/or relax the normal fibrous tissue, which can be beneficial in treatment of conditions such as scoliosis, tarsal tunnel syndrome, carpal tunnel syndrome and other conditions, where degrading, stretching and/or relaxing healthy or normal fibrosis tissue would be beneficial.

10 In some embodiments, the antifibrotic agent will enzymatically degrade or shrink the fibrotic tissue. For example, in some embodiments when the antifibrotic agent is an enzyme, it will reduce pressure by degrading proteoglycans (PG) so that PGs are not available to hold water. In some embodiments, the antifibrotic agent can be a protease or glycanase, which is not proteolytic. In some embodiments, the antifibrotic 15 agent, instead of an enzyme, can be an agent that dehydrates the fibrotic tissue, such as for example, a polycationic polymer. In some embodiments, the anti-fibrotic agent may be a hormone, such as for example, relaxin, which inhibits collagen production and stimulates collagen degradation. In some embodiments, the anti-fibrotic agent may be a cytokine, drug, cell, or nucleic-acid-based material that influences the function, viability, 20 or proliferation of fibroblasts or other cells in the fibrotic tissue. In some embodiments, the antifibrotic agent may be cells that inhibit collagen production and/or stimulates collagen degradation.

Antifibrotic agents include, one or more of pancreatic elastase, elastase-2a, elastase-2b, neutrophil elastase, proteinase-3, endogenous vascular elastase, cathepsin G, 25 mast cell chymase, mast cell tryptase, plasmin, thrombin, granzyme B, cathepsin S, cathepsin K, cathepsin L, cathepsin B, cathepsin C, cathepsin H, cathepsin F, cathepsin G, cathepsin O, cathepsin R, cathepsin V (cathepsin 12), cathepsin W, calpin 1, calpin 2, chondroitinase ABC, chondroitinase AC, hyaluronidase, chymopapain, chymotrypsin, legumain, cathepsin Z (cathepsin X), cathepsin D, cathepsin E, collagenase, matrix 30 metalloproteinases, such as for example, MMP-1 (collagenase-1), MMP-9, MMP-7 (matrilysin), MMP-8 (collagenase-2), MMP-13 (collagenase-3), MMP-18 (collagenase-4), MMP-2 (gelatinase a), MMP-9 (gelatinase b), MMP-3 (stromelysin-1), MMP-10

(stromelysin-2), MMP-11 (stromelysin-3), MMP-7 (matrilysin), MMP-26 (matrilysin), MMP-12 (metalloelastase), MMP-14 (MT1-MMP), MMP-15 (MT2-MMP), MMP-16 (MT3-MMP), MMP-17 (MT4-MMP), MMP-24 (MT5-MMP) transmembrane, MMP-25 (MT6-MMP), gpl anchor, MMP-19, MMP-20 (enamelysin), MMP-x, MMP-23, MMP-27, MMP-28 (epilysin), ADAMTS-1, ADAMTS-2, ADAMTS-3, ADAMTS-4 (aggrecanase-1), ADAMTS-5(aggrecanase-2), ADAMTS-14, papain, subtilisin, subtilisin A, heparanase. tyrosine kinase inhibitors: imatinib mesylate, dasatinib, nilotinib, inhibitors of PKC-delta and other kinases, TGF-beta receptor inhibitors, HMG-CoA inhibitors, angiotensin inhibitors: angiotensin-converting enzyme inhibitors, angiotensin-II receptor antagonist, pirfenidone, rosiglitazone, cannabinoid receptor, trabedersen, lerdelimumab, metelimumab, mycophenolate mofetil, interferon, or a combination thereof. In some embodiments, the antifibrotic factors include, but are not limited to, interleukins, interferons, cytokines, chemokines, chemotactic molecules, macrophages, lymphocytes, tumor necrosis factor alpha (TNF-alpha), T cells, interferon gamma (IFN-gamma), relaxin, hormones (e.g., progesterone, estrogen, testosterone, growth hormone, thyroid hormone, parathyroid hormone, etc.) or a combination thereof.

The agents of the present invention may be used in diagnostic and therapeutic applications and as such may be included in therapeutic or diagnostic kits.

Thus, compositions of the present invention may, if desired, be presented in a pack or dispenser device, such as an FDA approved kit, which may contain one or more unit dosage forms containing the active ingredient e.g., antibody. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accommodated by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising a preparation of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition, as is further detailed above.

As mentioned, the agents of the present invention are useful for treating diseases associated with aberrant collagen deposition.

Examples of diseases associated with aberrant collagen deposition include cancer (e.g. metastasized cancer), tumor fibrosis, fibrotic diseases.

5 The term "fibrosis" refers to the formation of fibrous tissue or "fibrotic tissue", usually as a reparative or a reactive process. As used herein, "fibrosis" or "fibroproliferative disease" includes those disorders or disease states that are caused by or accompanied by the abnormal deposition of scar tissue, or by excessive accumulation of collagenous connective tissue. In other embodiments, "fibrosis" includes formation of
10 fibrous tissue that may be a normal part of an organ or tissue. Thus, the antifibrotic agent may be administered to normal or healthy fibrous tissue to provide benefit to the patient (e.g., stretching, relaxing tissue to reduce or prevent pain and/or inflammation). Fibrosis may occur in any organ including, for example, kidney, lung, liver, skin, central nervous system, bone, bone marrow, cardiovascular system, an endocrine organ or the
15 gastrointestinal system. By "fibrosis-associated condition" is meant any condition that is related to fibrosis. Thus, fibrosis-associated conditions may be caused by, be concomitant with, or cause fibrosis.

 Examples of pathologic and excessive fibrotic accumulations include, but are not limited to, pulmonary fibrosis, asthma, adult respiratory distress syndrome (ARDS),
20 chronic obstructive pulmonary disease (COPD), idiopathic pulmonary fibrosis (IPF), acute lung injury (ALI), pulmonary fibrosis due to infectious or toxic agents, such as radiation therapy or chemotherapy, pulmonary fibrosis due to particle inhalation, post-transplant pulmonary fibrosis, perirenal fascitis, glomerulonephritis (GN), diabetic nephropathy, renal interstitial fibrosis, renal fibrosis resulting from complications of drug
25 exposure, HIV-associated nephropathy, transplant necropathy, retroperitoneal fibrosis, perivascular fibrosis in Systemic Lupus Erythematosus (SLE), obstruction-induced fibrosis in kidneys or spleen, benign prostate hypertrophy, fibrocystic breast disease, uterine fibroids, ovarian cysts, endometriosis, coronary infarcts, myocardial fibrosis, cerebral infarcts, congestive heart failure, dilated cardiomyopathy, myocarditis,
30 myelofibrosis, vascular stenosis, progressive systemic sclerosis, polymyositis, scleroderma (which affects the skin and the lungs), dermatomyositis, Raynaud's syndrome, retroperitoneal fibrosis, progressive massive fibrosis, nephrogenic systemic

fibrosis, Crohn's disease, stenosing tenosynovitis (trigger finger), Dupuytren's disease (palmar fibromatosis), Ledderhose's disease (plantar fibromatosis), Peyronie's disease, fibromatosis colli, keloids, mediastinal fibrosis, rheumatoid arthritis, musculoskeletal fibrosis, post-surgical adhesions, liver fibrosis, autoimmune hepatitis, cirrhosis including
5 primary biliary cirrhosis, viral hepatitis including HIV- or Hepatitis C-induced hepatitis, renal fibrotic disease, fibrotic vascular disease, e.g., atherosclerosis, varix, or varicose veins, scleroderma, Alzheimer's disease, diabetic retinopathy, glaucoma, proliferative vitreoretinopathy, fibrosis associated with ocular surgery, chronic transplant rejection, graft vs. host disease, radiation-induced fibrosis, and excessive or hypertrophic scar
10 and/or keloid formation in the dermis occurring during wound healing resulting from trauma or surgical wounds.

Fibrosis also includes healthy or normal production of fibrous tissue. In some embodiments, the antifibrotic agent will stretch or relax fibrous tissue (e.g., healthy or normal fibrous tissue, such as muscles, cartilage, tendons, ligaments, etc.) anywhere in the
15 body. For example, in carpal tunnel syndrome, in some embodiments, the flexor retinaculum generally has normal amounts of fibrous tissue. However, it is beneficial to degrade this normal fibrous tissue so the flexor retinaculum can stretch more easily to accommodate the excess tissue within the carpal tunnel. In some embodiments, the antifibrotic agent can degrade, stretch, and/or relax the normal fibrous tissue, which can
20 be beneficial in treatment of conditions such as scoliosis, tarsal tunnel syndrome, carpal tunnel syndrome and other conditions, where degrading, stretching and/or relaxing healthy or normal fibrosis tissue would be beneficial.

In one embodiment, the disorder to be treated is a disorder that results in fibrosis or sclerosis, including but not limited to groups of disorders selected from skeletal muscle
25 fibrosis, irradiation-induced fibrosis, autoimmune-related fibrosis, cardiovascular fibrosis, arteriosclerotic disorders, pulmonary fibrosis, adult respiratory distress syndrome, inflammatory disorders, scleroderma, cirrhosis, keloids, adhesions, hypertrophic scars; skeletal muscle fibrosis associated with a condition, such as muscular dystrophy, denervation atrophy induced by neuromuscular disease, or traumatic injury-induced
30 denervation atrophy; cardiovascular fibrosis selected from left ventricular hypertrophy secondary to hypertension, fibrosis associated with myocardial infarction, fibrosis associated with ischemiareperfusion injury, or fibrosis associated with myocarditis;

dermal fibrosis, keloid formation, hypertrophic scar formation, or adhesion formation; pulmonary fibrosis, pulmonary fibrosis due to adult respiratory distress syndrome and irradiation induced fibrosis, or a combination thereof.

In some embodiments, the fibrosis is a localized type of fibrosis, such as for example, stenosing tenosynovitis (trigger finger), Dupuytren's disease (palmar fibromatosis), Ledderhose's disease (plantar fibromatosis), Peyronie's disease, fibromatosis colli, keloids, mediastinal fibrosis, carpal tunnel syndrome, tarsal tunnel syndrome or a combination thereof. Examples of cancer include but are not limited to carcinoma, lymphoma, blastoma, sarcoma, and leukemia. Particular examples of cancerous diseases but are not limited to: Myeloid leukemia such as Chronic myelogenous leukemia. Acute myelogenous leukemia with maturation. Acute promyelocytic leukemia, Acute nonlymphocytic leukemia with increased basophils, Acute monocytic leukemia. Acute myelomonocytic leukemia with eosinophilia; Malignant lymphoma, such as Birkitt's Non-Hodgkin's; Lymphocytic leukemia, such as Acute lymphoblastic leukemia. Chronic lymphocytic leukemia; Myeloproliferative diseases, such as Solid tumors Benign Meningioma, Mixed tumors of salivary gland, Colonic adenomas; Adenocarcinomas, such as Small cell lung cancer, Kidney, Uterus, Prostate, Bladder, Ovary, Colon, Sarcomas, Liposarcoma, myxoid, Synovial sarcoma, Rhabdomyosarcoma (alveolar), Extraskelitel myxoid chondrosarcoma, Ewing's tumor; other include Testicular and ovarian dysgerminoma, Retinoblastoma, Wilms' tumor, Neuroblastoma, Malignant melanoma, Mesothelioma, breast, skin, prostate, and ovarian.

According to a particular embodiment, the cancer is colorectal cancer, lung cancer, breast cancer (e.g. triple negative breast cancer), renal cancer, ovarian cancer, gastric cancer, bladder cancer, liver cancer, ovarian cancer, fallopian cancer, glioblastoma, leukemia, including acute myeloid leukemia and lymphoma.

According to a particular embodiment, the cancer is a solid tumor.

Treatment of diseases may be effected by administering the agents per se, or together with a carrier as a pharmaceutical composition.

According to one embodiment, the anti-cancer agent or anti-fibrotic agent and the anti-LOXL-2 agent are formulated together in a single composition (e.g. pharmaceutical composition).

According to another embodiment, the anti-cancer agent (or anti-fibrotic agent) and the anti-LOXL-2 agent are formulated in two separate compositions (e.g. two individual pharmaceutical agents).

When administered as separate compositions, the present invention contemplates
5 administering the anti-LOXL-2 prior to administration of the anti-cancer agent (or anti-fibrotic agent), concomitant with administration of the anti-cancer agent (or anti-fibrotic agent) or following administration of the anti-cancer agent (or anti-fibrotic agent).

As used herein, "pharmaceutical carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption
10 delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g. by injection or infusion).

A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or
15 mode of administration will vary depending upon the desired results. To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent.

20 Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutical carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

25 The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intra-arterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid,
30 intraspinal, epidural and intrasternal injection and infusion.

The present invention also contemplates ex vivo cell therapy using the antibodies described herein, whereby cells and/or tissues are genetically modified to

express and secrete the antibodies. Following genetic modification, the cells are then introduced back into the body. The cells may be derived from the patient himself or may be obtained from a donor. These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of
5 presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions.

10 In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

Regardless of the route of administration selected, the agents of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable
15 dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the
20 patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions
25 employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical art.

The composition must be sterile and fluid to the extent that the composition is deliverable by syringe. In addition to water, the carrier preferably is an isotonic buffered saline solution.

30 Proper fluidity can be maintained, for example, by use of coating such as lecithin, by maintenance of required particle size in the case of dispersion and by use of surfactants. In many cases, it is preferable to include isotonic agents, for example,

sugars, polyalcohols such as mannitol or sorbitol, and sodium chloride in the composition.

Other contemplated uses of the antibodies of the present invention include purification of analytes; in immunohistochemistry and enzyme immunoassays; for
5 radioimaging and radioimmunotherapy and for drug delivery.

Other contemplated uses are set forth in Cao Y, Suresh MR. Bispecific antibodies as novel bioconjugates. *Bioconjug Chem.* 1998 Nov-Dec;9(6):635-44, incorporated herein by reference.

It is expected that during the life of a patent maturing from this application many
10 relevant anti-cancer and antifibrotic agents will be developed and the scope of these terms is intended to include all such new technologies *a priori*.

As used herein the term "about" refers to $\pm 10\%$.

The terms "comprises", "comprising", "includes", "including", "having" and their conjugates mean "including but not limited to".

15 The term "consisting of" means "including and limited to".

The term "consisting essentially of" means that the composition, method or structure may include additional ingredients, steps and/or parts, but only if the additional ingredients, steps and/or parts do not materially alter the basic and novel characteristics of the claimed composition, method or structure.

20 Throughout this application, various embodiments of this invention may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as
25 individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

30 As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known

manners, means, techniques and procedures by practitioners of the chemical, pharmacological, biological, biochemical and medical arts.

As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition or substantially preventing the appearance of clinical or aesthetical symptoms of a condition.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

Various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below find experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions illustrate some embodiments of the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659

and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Culture of Animal Cells - A Manual of Basic Technique" by Freshney, Wiley-Liss, N. Y. (1994), Third Edition; "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

GENERAL MATERIALS AND METHODS

Cell lines and culture: The human dermal fibroblast cells were cultured in high-glucose DMEM (Invitrogen) supplemented with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS, Invitrogen), and 100 U/ml penicillin and 100 g/ml streptomycin. Medium was exchanged every 2–3 days and passaged after reaching 80–90% confluence with 0.05% trypsin EDTA (Biological industries). All cell lines were maintained at 37°C in a humidified atmosphere containing 5% CO₂. Fibroblasts were grown on glass coverslips in 24-well dishes until reaching contact inhibition. Then, the medium was replaced with same medium containing 5ng/ml EGF, 5µg/ml insulin and

150µg/ml L-ascorbic acid phosphate magnesium salt n-hydrate to induce ECM secretion, in the presence of PBS or A341 (100ng/ul). Medium was exchanged every 3 days.

LOXL2 expression and purification: The catalytic domain of human LOXL2 (545-775) was cloned into the pET28 expression vector. The expression vector was transformed into competent Escherichia coli BL21, and colonies were selected for kanamycin resistance. Bacteria were grown in LB medium containing 100 µg/ml kanamycin in a shaker flask at 37°C. Protein expression was induced with 0.5mM IPTG at an OD600 = 0.6, and cell growth was continued for further 4 hours. Following expression, the enzyme was accumulated in the fraction of inclusion bodies. Cells were harvested by centrifugation and resuspended in a buffer containing 25% sucrose, 50mM Tris-HCl pH 8, 0.1M NaCl, 0.2mM EDTA, 1mM DTT, ~10mg lysozyme and 1 pill of complete protease inhibitor, and stirred for 10~20min in cold room. The suspension was sonicated (6 cycles, 30-seconds cycles) and centrifuged at 22000 g for 30 min. at 4C. The pellet was further suspended in buffer containing 2M urea, 2% Triton, 50mM Tris-HCl pH 8, 0.1M NaCl, 0.2mM EDTA, 1mM DTT and centrifuged at 9,000 rpm for 1 hour, sonicated and centrifuged. The pellet was solubilized in 8 M urea, 50 mM MOPS pH 7.5. Then the protein was purified by anion exchange using 6M Urea, 25 mM MOPS pH 7.5 and a gradient of NaCl concentration. The collected protein-containing fractions were diluted <50ug/1ml in 6M urea, 25 mM MOPS pH 7.5, 150mM BME and refolded by using a multi-step dialysis against solution containing 0.4M arginine (non-neutralized), 25 mM MOPS pH 7.5, 200mM NaCl, 40uM CuCl₂, 1mM glutathion reduced, 0.1mM glutathion oxidized and decreasing concentration of urea (from 2M to 0M). Finally, the enzyme was purified by size exclusion chromatography using a High Load™ 16/60 Superdex™ 75 (Amersham Biosciences) and eluted with 50 mM MOPS pH 7.5, 150 mM NaCl. The eluted fractions were checked for purity on 15% gel by SDS-PAGE, and those containing the LOXL2 protein were pooled and concentrated.

Antibody generation: Female BALB/c mice were immunized on day 1 with complete Freund's adjuvant (Disco) and 50 µg of the catalytic domain of LOXL2 and boosted every 2 weeks with incomplete Freund's adjuvant by subcutaneous injection. To generate antibodies, spleens were collected, and B cells were fused with NSO murine myeloma cells³⁴. Hybridomas were screened with ELISA for immunoreactivity against

the catalytic domain of LOXL2 (as described below). Selected hybridomas positive for LOXL2 were subcloned twice and then expanded in tissue culture.

Antibody purification: Hybridoma cells of GS341 were grown in DCCM. Cells were collected by centrifugation at 1200rpm and the supernatant was collected. Supernatant was dialyzed against 20mM phosphate buffer pH 8. 5ml HiTrap Protein A HP column was equilibrated with 100mM phosphate buffer pH 8 and the supernatant was loaded at 5ml/min. Antibody was eluted with 100mM citrate buffer pH 6 and dialyzed against PBS or TBS buffers.

Antibody digestion with papain: Papain was activated in 0.5M Tris-HCl pH 8, 10mM EDTA and 5mM DTT for 15mins at 37°C. Active papain was added to a solution of intact ab341 at a ratio of 1:1000, the digestion process was carried for 3hrs at 37°C. Digestion reaction was terminated with the addition of 20mM iodoacetamide in the dark at room temperature for 30 mins. The fab fragment was isolated from the Fc by Protein A column, the fab fragment was collected from the flow through and dialyzed against PBS. The purity of the Fab fragment was estimated by 12% SDS-PAGE gel.

Antibody sequencing: Total RNA was extracted from the hybridoma cells and cDNA was prepared using SuperScript kit (Invitrogen) as recommended by the manufacturer. The cDNA was used as a template for the PCR reaction using Reddy mix (Termo). For VH and VL amplification the primer mix was used according to Zhou et al. (Zhou et al., 1994). PCR products were purified by gel extraction and used for direct T-A cloning into the pGEM-T plasmid. Ligation products were transformed into DH5 α bacteria and grown on LB plates supplemented with Ampicilin, X-gal and IPTG. Positive clones were sent for sequencing, and sequences were analyzed using IMGT-V-QUEST tool.

LOXL2 ELISA binding assay: A 96 well plate (Nunc) was coated with LOXL2 or BSA at 5 μ g/ml. After the plate was coated, it was incubated with the antibody for 1 h at 25°C. The bound antibody was detected by peroxidase-conjugated antibody goat anti mouse according to standard procedures. The EC₅₀ was calculated from a four-parametric sigmoidal-curve fitting analysis.

Microscale thermophoresis: LOXL2 was labeled with 100 nM Alexa647-labeled and incubated with increasing concentration of antibody from 0.1 to 1000 μ M in binding buffer containing 50 mM Tris pH 8, 150 mM NaCl, 10 mM MgCl₂, 1% BSA

and 0.05% Tween. After 15 min incubation, samples were measured using a Monolith NT.115 (Nanotemper, Germany) with 50% Laser-power, Laser on time 35 s, and 40% LED-power.

MST competition assay: Labeled LOXL2 was preincubated with 100 nM bAPN at room temperature for 15 min. Next, the LOXL2:bAPN samples were incubated with increasing concentration of antibody from 0.1 to 1000 μ M in binding buffer. After 15 min incubation samples were measured at using a Monolith NT.115 (Nanotemper, Germany) with 50% Laser-power, Laser on time 35s, and 40% LED-power.

Enzyme inhibition assay: The enzymatic activity of LOXL2 was measured by coupling horse radish peroxidase (HRP) activity to LOXL2 and using the conversion of Amplex Red (Invitrogen) to resofurin as the readout, as described previously, expect collagen type I was used as a substrate (Barry Hamilton et al., Nat Med 16, 1009-17 (2010)). Briefly, collagen type I from rat tail (AdvanceBioMatrix) was naturalized by addition of PBS and NaOH according to manufacture procedure. Prior to the enzymatic assay, LOXL2 was incubated with increasing concentration of antibody for 1 hour in room temperature. The enzymatic reaction was started by adding substrate mixture (50 mM sodium borate, pH 8.0, 100 μ M Amplex Red reagent, 0.25mg/ml Collagen) to the enzyme mixture (50 mM sodium borate, pH 8.0, 2 U/mL HRP, 25 nM LOXL2), and fluorescence was measured using SynergyHT plate reader in kinetics mode with the excitation wavelength at 540 nm and the emission wavelength set to 590 nm. Measurements were made at 30 second intervals for 1 hour at 37 °C. As a background, measurements of solution mixture without LOXL2 were conducted, and the signal obtained was subtracted from kinetic measurements. The IC₅₀ (the concentration of inhibitor that results in a 50% decrease in activity relative to no inhibitor) was determined by dividing the reaction initial velocities at different antibody concentration with the activity at zero concentration, and fitting the data to the following equation:

ECM isolation from tissue culture: ECM from individual cultures was decellularized by incubating the cells with 0.5% triton and 20mM EDTA for 3 days in room temperature, followed by gentle washing with PBS.

Two-photon microscopy and second harmonics generation: Samples were imaged using a two-photon microscope in the *in-vivo* imaging (2PM:Zeiss LSM 510 META NLO; equipped with a broadband Mai Tai-HP-femtosecond single box tunable

Ti-sapphire oscillator, with automated broadband wavelength tuning 700–1020 nm from Spectraphysics, for two-photon excitation). For collagen second harmonic imaging a wavelength of 800 nm was used (detection at 400nm), using the X20 objective.

Raman microscopy: Raman spectra were acquired using a WITec (Ulm, Germany) Alpha300 R/A/S confocal Raman microscope as described previously^{19,20} using excitation wavelength of 532 nm. Imaging was performed by scanning of the laser beam over the sample using an image size is in the range of 150µm X 150µm, with 300 points per line, and integration time of 0.5sec. Data analysis was performed using the WITec Project Plus software. Raman images were constructed by integrating the raw data over the vibrational bands of interest. Integration over selected spectral regions was performed by summing up the CCD counts between two frequencies. For example, the C–H stretching band between 2850 cm⁻¹ and 3000 cm⁻¹ was used to image collagen morphology.

Scanning electron microscopy: The ECM was washed with PBS and fixated in 4% paraformaldehyde in PBS for 20 min, followed by 3 washes in PBS. The samples were further incubated with SST for 40 minutes in room temperature followed by 3 washes in water. Ethanol dehydration of samples was then carried out by consecutive washing in increasing concentrations of ethanol (from 30% to 100%), followed by ethanol exchange with liquid CO₂ and critical point drying. Finally, samples were coated with a thin layer of gold and imaged in an Ultra microscope (Zeiss).

Metastasis tests in animals: Female CB-17 SCID mice (Harlan Laboratories, Haslett, MI; 5 per group) were implanted in the fat pad with RFP-MDA-MB-231 cells (1.4 x10⁶ cells/mouse). Three weeks later, mice were treated twice a week with PBS or 30mg/kg GS341 for another four weeks. For lung metastasis, lungs were removed and washed, and images were acquired using a fluorescent binocular. In the second experiment, mice were pretreated for four week with PBS or 30mg/kg Fab GS341 prior to injection of RFP-MDA-MB-231 cells (1.4 x10⁶ cells/mouse) into the mice tail vein, and mice were continued to be treated with PBS or 30mg/kg Fab GS341 for another four weeks. Metastasis to the lungs was visualized as stated above. Tumors and lungs tissues were preserved in OCT and frozen in -80 °C until use.

ECM isolation from tissue: Tumors were sectioned to 150 µm sections and washed three times with PBS and water. Then, the sample underwent six cycles of

freezing on dry ice and thawing in water followed by replacement of water. Final cells removal was accomplished by treatment with NH_4OH for 20 minutes, followed by three wash with water.

Immunofluorescence analysis: Murine tumors were fixed in PFA 4%,
5 embedded in paraffin and sectioned. Tumor sections underwent deparafinization and antigen retrieval according to the primary antibody manufacturer's instructions. Sections were permeabilized with 0.5% Triton X-100 in PBS for 5 minutes and blocked with 5% bovine serum albumin (in PBS containing 0.1% Triton) for 1 hour at room temperature. For immunostaining, the cells were incubated overnight at 40 °C with
10 phosphorylated histone H3 (pH3) antibody (Santa Cruz) which were used to analyze cell proliferation. After three washes with PBS, the cells were stained for 1 hour at room temperature with secondary antibodies (abcam) followed by 10 minutes of DAPI (4',6'-diamidino-2-phenylindole dihydrochloride) staining. The cells were viewed under a Nikon eclipse 90i fluorescence microscope. Pictures were taken with a 1310 digital
15 camera (DVC, Austin, TX).

Statistical analyses: All analyses were performed in triplicate for $n=3$. Tumor growth measurements used the Exact-sig (2x1-tailed) Mann-Whitney test. Statistical analysis was performed with Microsoft Excel using student t-test. Significance levels were set at $P \leq 0.05$, $P \leq 0.01$, and $P \leq 0.001$. Unless otherwise indicated, all graphical data
20 are reported as $\pm \text{SE}$.

EXAMPLE 1

LOXL2 is critical for maturation of collagen in the ECM

ECM secreted by human dermal fibroblast served as a model system to study
25 ECM buildup by stromal cells. Two photon second harmonic generation (SHG) microscopy, which allows label free imaging of collagen fibers on native tissue and cells was used to follow the process of collagen synthesis and assembly at different time points (Figure 1A). Only after 5 days, a premature form of collagen fibers was visualized. These were still not organized. At day 7 the assembly of linear collagen
30 fibers was observed, which later matured to the characteristic collagen wavy-like higher organization. The expression levels of enzymes belonging to the LOX/LOXL family as well as representative ECM proteins were monitored along a period of four weeks by

real-time PCR (Figures 2A-D). Interestingly, a gradual increase in LOX expression levels until day 7 was observed followed by total decrease of LOX expression afterward, while expression of LOXL enzymes, and among them LOXL2, was only elevated at later stages (day 14) (Figure 1B). Therefore, it can be concluded that crosslinking by LOXL enzymes are required for maturation of collagen fibers in the ECM at later stages of assembly of the ECM while LOX is required at the earlier stages. Thus the overexpression of LOXL2 at later stages of collagen assembly, can serve as a therapeutic window to interfere with ECM synthesis during the progression of a disease.

EXAMPLE 2

GS341 inhibits collagen type I crosslinking and maturation of collagen fibers

In order to specifically interfere with LOXL2 catalytic activity at the extracellular matrix, the present inventors generated a recombinant catalytic domain of the enzyme composing of residues 545-775 (Figure 1C). Antibodies were generated using the hybridoma technique by immunizing mice with the generated catalytic fragment, and screening for monoclonal antibodies clones that bind the catalytic domain of LOXL2. Several purified antibodies from positive clones displayed binding affinities to the catalytic domain of LOXL2 as well as to the full length enzyme at the sub-nano molar range using an ELISA binding assay as well as thermoscale thermophoresis measurements of the binding dissociation constant in solution (Figures 3A-C). Next, the present inventors screened for an antibody that has an inhibitory activity on collagen crosslinking by LOXL2 by performing an enzymatic activity assay in the presence of monomeric collagen type I as the substrate.

A decrease in the catalytic activity of the enzyme correlated with increasing concentrations GS341 antibody, with an IC_{50} of $1\mu M$. GS341 bound both human and mouse LOXL2 and was found to be a specific inhibitor of LOXL2. Since invasion is the first process during metastasis, the present inventors further evaluated the inhibitory activity of GS341 in 3D invasion cell-based assay using human metastatic breast cancer cell line MDA-MB-231, that overexpresses almost only LOXL2. The antibody reduced the ability of the cells to invade into the basement membrane by 70 % (Figure 4A). Thus, targeting the enzymatic activity of LOXL2 at the ECM by GS341 has a beneficial effect on a key biological process associated with cancer metastasis.

In order to evaluate the direct effect of GS341 on ECM formation and composition, the present inventors looked into collagen deposition by human dermal fibroblast cells in the presence or absence of the antibody in their growth medium. SHG analysis of control and antibody treated samples did not detect differences in morphology or amount of collagen during the first week of ECM assembly (Figure 5A). Only after four weeks of growth, it was observed that the control samples displayed a wave-like fibril organization of collagen, while the collagen fibers in the GS341 treated samples remained linear (Figure 4B). This observation was also confirmed by Raman microscopy of decellularized ECM samples, demonstrating similar effect on ECM organization upon inhibition of LOXL2 (Figure 4C). In addition, comparison of the averaged Raman spectra for the control and GS341 treated samples further demonstrated shift in vibration peak associated with collagen indicating changes in the chemical properties of the samples (Figure 6). Mass spectrometry analysis of ECM derived from the control and GS341 treated cells did not detect statistically significant differences in ECM composition indicating that GS341 affects only ECM morphology (Figure 4D). These results confirm that the enzymatic activity of LOXL2 is required at later stages of ECM deposition during basement membrane formation, and that inhibition of LOXL2 by GS341 prevents maturation of collagen fibers. High resolution scanning electron microscopy imaging of these samples further demonstrated nanoscale changes in fibril directionality within the collagen fibers of the samples. While collagen fibrils of the control samples were assembled in one dominant direction, the collagen fibrils in the antibody treated sample were dispersed and orientated in many directions (Figure 4E and 4F). This indicates that crosslinking of collagen fibrils by LOXL2 is required also for proper alignment in the extra-cellular space.

25

EXAMPLE 3

GS341 affect breast cancer tumor ECM morphology leading to attenuation of lung metastasis

Overexpression of LOXL2 is associated with poor clinical prognosis for patients and breast cancer tumor are characterized with increasing ECM stiffening with the progression of tumor malignancy. To determine whether inhibition of collagen maturation by LOXL2 has a therapeutic potential on an ECM associated disease, the

30

present inventors quantified the effect of GS341 on breast cancer metastasis. Specifically, triple negative breast cancer MDA-MB-231 cells implanted into the mammary fat pads of immunocompromised nude mice were used. Previously, genetic knock down of LOXL2 in a similar breast cancer model using MB-MDA-231 cells indicated that LOXL2 is not required for primary tumor growth. Therefore the present inventors started treatment with GS341 three weeks after implantation (30 mg per kg body weight twice weekly). Four weeks later both tumor size (Figure 7A) and metastases in the lungs (Figure 7B) were measured. Treatment with GS341 not only led to significant decrease in final tumor volume ($P=0.02$) but also to a significant reduction in metastasis to the lungs. This effect was similar to that obtained by genetic knock-down of LOXL2, indicating that targeting extracellular LOXL2 by an antibody can solely reduce metastasis. In addition, treatment of GS341 significantly reduced cell proliferation in the primary tumors, as shown by immunostaining analysis for markers of cell cycle activity (Figure 7C).

In order to link between metastasis reduction and the direct effect of LOXL2 on the ECM, changes in the ECM of the primary tumor associated antibody treatment were quantified. Mass spectrometry analysis of tissue samples from the primary tumor detected changes in ECM composition between control and GS341 treated tumors (Figure 7D). SHG imaging of representative cryo-sections of control tumors depicted massive and linear collagen fibers orientated in a preferred directionality, characteristic of tumors. In sharp contrast, collagen fibers in the GS341 treated tumors, were highly branched and disorientated (Figure 7E and 7F). SEM analysis of decellularized tumor sections further depicts that collagen fibrils in the treated tumors have a difference diameter distribution and are thinner compared with the control tumors (Figure 7G and 7H). Thus, inhibition of cross-linking by LOXL2 led to the buildup of a different ECM architecture at the tumor microenvironment. Importantly, this effect on collagen architecture is a mirror-image of that observed in the fibroblast experiment *in vitro*. Thus, inhibition of collagen crosslinking by LOXL2 has a direct effect on the ECM of the primary tumor, leading to attenuation of metastatic seeding in the lungs. Since tumor cells are thought to migrate along linear collagen fibers during the process of tumor invasion and metastasis, it is proposed that GS341 inhibits metastasis by blocking the formation of tumor collagen “high-ways”.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope
5 of the appended claims.

All publications, patents and patent applications mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference. In addition, citation or
10 identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention. To the extent that section headings are used, they should not be construed as necessarily limiting.

WHAT IS CLAIMED IS:

1. An article of manufacture comprising at least two active agents, wherein a first of said two active agents comprises an anti-cancer agent or an antifibrotic agent and a second of said two active agents downregulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of altering the structure of the extracellular matrix.
2. The article of manufacture of claim 1, wherein said second agent is an antibody which comprises an antigen recognition region which specifically binds to said LOXL-2.
3. The article of manufacture of claim 2, wherein said antibody specifically binds the catalytic site of said LOXL-2.
4. The article of manufacture of claim 2, wherein said antibody does not bind to the fourth scavenger receptor-cysteine-rich (SRCR) domain of said LOXL-2.
5. The article of manufacture of claim 1, wherein said second agent is a polynucleotide agent which downregulates the expression of said LOXL-2.
6. The article of manufacture of claim 1, wherein said first agent and said second agent are packaged in separate packaging.
7. The article of manufacture of claim 1, wherein said first agent and said second agent are formulated in a single pharmaceutical composition.
8. The article of manufacture of any one of claims 1-5, wherein said second agent is capable of down-regulating crosslinking of type I collagen in vitro.

9. An isolated antibody comprising an antigen recognition region which specifically binds to lysyl-oxidase like protein-2 (LOXL-2) and being capable of down-regulating crosslinking of type I collagen in vitro.

10. The antibody of claim 9, specifically binding to the catalytic site of said LOXL-2.

11. The antibody of claim 9, wherein said antibody does not bind to the fourth scavenger receptor-cysteine-rich (SRCR) domain of said LOXL-2.

12. An isolated antibody comprising an antigen recognition region which comprises CDR amino acid sequences as set forth in SEQ ID NOs: 3, 4, 5, 6, 7 and 8.

13. A method of treating a disease associated with aberrant collagen deposition in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of the antibody of claims 9 or 12, thereby treating the disease.

14. The antibody of claim 9 or 12 for use in treating a disease associated with aberrant collagen deposition.

15. The article of manufacture or isolated antibody of claims 1, 9 or 12, wherein said agent which downregulates the activity and/or expression of said LOXL-2 does not affect angiogenesis in vivo.

16. The article of manufacture or isolated antibody of claims 2 or 9, wherein said antigen recognition region comprises CDR amino acid sequences as set forth in SEQ ID NOs: 3, 4, 5, 6, 7 and 8.

17. The article of manufacture or isolated antibody of any one of claims 2, 9 or 12 wherein the amino acid sequence of the V_H of the antibody is as set forth in SEQ ID NO: 1.

18. The article of manufacture or isolated antibody of any one of claims 2, 9 or 12, wherein the amino acid sequence of the V_L of the antibody is as set forth in SEQ ID NO: 2.

19. An article of manufacture comprising an isolated antibody comprising an antigen recognition region which comprises CDR amino acid sequences set forth in SEQ ID NO: 3, 4, 5, 6, 7 and 8 and an anti-cancer agent or an antifibrotic agent.

20. The article of manufacture of claims 1 or 19, wherein said anti-cancer agent comprises a chemotherapeutic agent.

21. The article of manufacture of claim 20, wherein said chemotherapeutic agent is cisplatin.

22. The article of manufacture of claims 1 or 19, wherein said anti-cancer agent is an antibody.

23. The article of manufacture of claim 22, wherein said antibody binds specifically to MMP-9.

24. A method of treating a disease associated with aberrant collagen deposition in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of an agent which downregulates the activity and/or expression of said LOXL-2 and an additional agent selected from the group consisting of an anti-cancer agent and an antifibrotic agent, thereby treating the disease.

25. A combination of an agent which downregulates the activity and/or expression of said LOXL-2 and an additional agent selected from the group consisting of an anti-cancer agent and an antifibrotic agent for use in treating a disease associated with aberrant collagen deposition.

26. The method, antibody or combination of any one of claims 13, 14, 24 or 25, wherein said disease is cancer.

27. The method or combination of any one of claims 13, 14, 24 or 25, wherein said disease is pulmonary alveolar proteinosis (PAP).

28. The method, antibody or combination of claim 26, wherein said cancer is breast cancer.

29. The method, antibody or combination of claim 26, wherein said cancer is colon cancer.

30. The method, antibody or combination of claim 28, wherein said breast cancer is triple negative breast cancer.

31. The method, antibody or combination of claim 26, wherein said cancer has metastasized.

32. The method of claim 24, wherein said administering of the anti-cancer agent or antifibrotic agent is effected following administering of the agent which downregulates the activity and/or expression of said LOXL-2.

33. The method, antibody or combination of any one of claims 13, 14, 24 or 25, wherein said disease is a fibrotic disease.

34. The method or combination of claims 24 or 25, wherein the agent which downregulates the activity and/or expression of said LOXL-2 is an antibody which comprises an antigen recognition region which specifically binds to said LOXL-2, the antibody being capable of down-regulating crosslinking of type I collagen in vitro.

35. The method or combination of claims 24 or 25, wherein the agent which downregulates the activity and/or expression of said LOXL-2 does not affect angiogenesis in vivo.

36. The method or combination of claim 34, wherein said antigen recognition region comprises CDR amino acid sequences as set forth in SEQ ID NO: 3, 4, 5, 6, 7 and 8.

37. The method, antibody or combination of any one of claims 13, 14 or 34, wherein the amino acid sequence of the V_H of the antibody is as set forth in SEQ ID NO: 1.

38. The method, antibody or combination of any one of claims 13, 14 or 34, wherein the amino acid sequence of the V_L of the antibody is as set forth in SEQ ID NO: 2.

39. A pharmaceutical composition comprising the antibody of any one of claims 9 or 12 and a pharmaceutically acceptable carrier.

40. A pharmaceutical composition comprising an anti-cancer agent or an antifibrotic agent as a first active agent and an agent which downregulates the activity and/or expression of lysyl-oxidase like protein-2 (LOXL-2) and which is capable of altering the structure of the extracellular matrix as a second active agent, and a pharmaceutically acceptable carrier.

41. A bispecific antibody comprising a first antigen-binding domain that specifically binds to LOXL-2 and a second antigen-binding domain that specifically binds to an epitope of a protein that is associated with cancer.




FIG. 1B




FIG. 1C




FIG. 2B




FIG. 2D




FIG. 2A








FIG. 2C











FIG. 4F






FIG. 5B





FIG. 6
















FIG. 7A

FIG. 7E

FIG. 7G

FIG. 8A




FIG. 8B






FIG. 9A



1µM Control IgG

FIG. 9B



1µM anti-LOXL2-341

FIG. 9C






FIG. 9D



1µM Control IgG

FIG. 9E



1µM anti-LOXL2-341

FIG. 9F






FIG. 9G



1µM Control IgG

FIG. 9H



1µM anti-LOXL2 + anti-MMP-9





FIG. 10E

FIG. 11A




FIG. 11B

Fab
GS341




FIG. 11C

PBS




FIG. 12B

PBS





FIG. 12A

Fab GS341



INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2014/050678

A. CLASSIFICATION OF SUBJECT MATTER IPC (2014.01) C07K 16/40, A61K 39/395, A61P 35/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC (2014.01) C07K, A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Databases consulted: NCBI, Sequence Manipulation Suite, Esp@cenet, Google Patents, CAPLUS, MEDLINE, Google Scholar Search terms used: Loxl2, extracellular matrix, collagen, LOXL2 catalytic site, fourth scavenger receptor cysteine rich, SRCR, angiogenesis.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2012139045 A1 GILEAD BIOLOGICS, INC. ; MARSHALL, DEREK. ; SPANGLER, RHYANNON. ; SMITH, VICTORIA 11 Oct 2012 (2012/10/11) The whole document	1-8,15,20-22,40
X	BARRY-HAMILTON, Vivian, et al. Allosteric inhibition of lysyl oxidase-like-2 impedes the development of a pathologic microenvironment. Nature medicine, 2010, 16(9): 1009-1017. <DOI:10.1038/nm.2208> 05 Sep 2010 (2010/09/05) The whole document	1-40
X	EP 2537529 A1 GILEAD BIOLOGICS INC 26 Dec 2012 (2012/12/26) paragraphs [0166]-[0174]	41
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 19 Oct 2014		Date of mailing of the international search report 21 Oct 2014
Name and mailing address of the ISA: Israel Patent Office Technology Park, Bldg.5, Malcha, Jerusalem, 9695101, Israel Facsimile No. 972-2-5651616		Authorized officer VILSKER Olga Telephone No. 972-2-5651780

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing filed or furnished:

a. (means)

☐

on paper

☒

in electronic form

b. (time)

☐

in the international application as filed

☒

together with the international application in electronic form

☐

subsequently to this Authority for the purposes of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/IL2014/050678

Patent document cited search report	Publication date	Patent family member(s)	Publication Date
WO 2012139045 A1	11 Oct 2012	WO 2012139045 A1	11 Oct 2012
		EP 2694554 A1	12 Feb 2014
		US 2014186340 A1	03 Jul 2014
EP 2537529 A1	26 Dec 2012	EP 2537529 A1	26 Dec 2012
		AU 2008282739 A1	05 Feb 2009
		AU 2008282739 B2	19 Dec 2013
		AU 2008299784 A1	19 Mar 2009
		AU 2008299784 B2	24 Apr 2014
		AU 2014204536 A1	07 Aug 2014
		CA 2693208 A1	05 Feb 2009
		CA 2693310 A1	19 Mar 2009
		CN 101835490 A	15 Sep 2010
		CN 101842114 A	22 Sep 2010
		DK 2182981 T3	02 Apr 2013
		EP 2182981 A2	12 May 2010
		EP 2182981 B1	09 Jan 2013
		EP 2185198 A1	19 May 2010
		EP 2543389 A2	09 Jan 2013
		EP 2543389 A3	24 Apr 2013
		ES 2402334 T3	30 Apr 2013
		HK 1143736 A1	05 Jul 2013
		HR P20130287 T1	31 May 2013
		JP 2010535205 A	18 Nov 2010
		JP 5312459 B2	09 Oct 2013
		JP 2010535219 A	18 Nov 2010
		JP 2013231035 A	14 Nov 2013
		PT 2182981 E	18 Apr 2013
		SI 2182981 T1	31 May 2013
		US 2009053224 A1	26 Feb 2009
		US 8461303 B2	11 Jun 2013

Information on patent family members

PCT/IL2014/050678