



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2356404 C 2010/03/30

(11)(21) **2 356 404**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1999/12/22
(87) Date publication PCT/PCT Publication Date: 2000/06/29
(45) Date de délivrance/Issue Date: 2010/03/30
(85) Entrée phase nationale/National Entry: 2001/06/19
(86) N° demande PCT/PCT Application No.: GB 1999/004381
(87) N° publication PCT/PCT Publication No.: 2000/037099
(30) Priorités/Priorities: 1998/12/22 (US09/218,913);
1999/11/17 (US09/441,966)

(51) Cl.Int./Int.Cl. *A61K 38/57* (2006.01),
A61K 9/72 (2006.01), *A61P 11/12* (2006.01)
(72) Inventeurs/Inventors:
HALL, RODERICK L., GB;
POLL, CHRISTOPHER T., GB;
NEWTON, BENJAMIN B., GB;
TAYLOR, WILLIAM J. A., GB
(73) Propriétaire/Owner:
BAYER AKTIENGESELLSCHAFT, DE
(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : TECHNIQUE PERMETTANT D'ACCELERER LE DEBIT DE CLAIRANCE MUCOCILIAIRE
(54) Title: METHOD FOR ACCELERATING THE RATE OF MUCOCILIARY CLEARANCE

(57) **Abrégé/Abstract:**

The instant invention provides for a composition and method for using Kunitz-type serine protease inhibitors, e.g., aprotinin or bikunin, for stimulating the rate of mucociliary clearance of mucus and sputum in lung airways of subjects afflicted with mucociliary dysfunctions such as cystic fibrosis.



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 38/57, A61P 11/12	A3	(11) International Publication Number: WO 00/37099 (43) International Publication Date: 29 June 2000 (29.06.00)
(21) International Application Number: PCT/GB99/04381 (22) International Filing Date: 22 December 1999 (22.12.99) (30) Priority Data: 09/218,913 22 December 1998 (22.12.98) US 09/441,966 17 November 1999 (17.11.99) US (71) Applicant (for all designated States except US): BAYER AKTIENGESSELLSCHAFT [DE/DE]; D-51368 Leverkusen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): HALL, Roderick [GB/GB]; 10 Acacia Close, Woodham, Surrey KT15 3SJ (GB). POLL, Christopher, T. [GB/GB]; 11 Scarborough Way, Windsor Meadows, Slough SL1 9JY (GB). NEW- TON, Benjamin, B. [GB/GB]; 35 St. Bartholomew's Road, Reading, Berkshire RG1 3QA (GB). TAYLOR, William, J., A. [GB/GB]; 'Wyndin' the Willows, Maidenhead Road, Windsor SL4 5UB (GB). (74) Agent: PETT, C., P.; Frank B. Dehn & Co., 179 Queen Victoria Street, London EC4V 4EL (GB).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 26 October 2000 (26.10.00)
(54) Title: METHOD FOR ACCELERATING THE RATE OF MUCOCILIARY CLEARANCE BY USING A KUNITZ-TYPE SERINE PROTEASE INHIBITOR (57) Abstract The instant invention provides for a composition and method for using Kunitz-type serine protease inhibitors, e.g., aprotinin or bikunin, for stimulating the rate of mucociliary clearance of mucus and sputum in lung airways of subjects afflicted with mucociliary dysfunctions such as cystic fibrosis.		

WO 00/37099

PCT/GB99/04381

Method for Accelerating the Rate of Mucociliary Clearance

5

Field of the Invention

10 The present invention relates to compositions comprising serine protease inhibitor proteins which stimulate the rate of mucociliary clearance of mucus and sputum in lung airways. The present invention also relates to methods for stimulating the rate of mucociliary clearance in mammals.

15 Background of the Invention

Problem Addressed

Mucociliary dysfunction, characterized by the inability of ciliated epithelium to clear mucus and sputum in lung airways, is a serious complication
20 of chronic obstructive lung diseases such as Chronic Bronchitis (CB), Bronchiectasis (BE), asthma and, especially, Cystic Fibrosis (CF). Patients suffering from mucociliary dysfunction are particularly vulnerable to secondary bacterial infections. Treatment and maintenance modalities for CF and other respiratory diseases associated with mucociliary dysfunction and the need for
25 improved treatments have been described. See, for instance, Braga "Drugs in Bronchial Mucology, Raven Press, New York, 1989; Lethem et al, Am Rev. Respir. Dis. 142:1053-1058, 1990; U.S. Patent No. 5,830,436..

Cystic Fibrosis

30 Cystic fibrosis (CF) is an autosomal recessive disease that causes abnormalities in fluid and electrolyte transport in exocrine epithelia. Mutations within the DNA coding for a protein termed the cystic fibrosis transmembrane conductance regulator (CFTR) have been found in virtually all CF patients. Cells of the lung are particularly affected. Di Santagrese et al, Am J. Med. 66: 121-132
35 (1979).

In CF, the luminal border of the airway mucosal cell is unresponsive to cAMP-dependant protein kinase activation of membrane chloride ion channels. The cell permeability to Cl^- is impaired and Na^+ absorption across the cell

membrane is accelerated. Both of these electrolyte imbalances tend to reduce the level of hydration of the airway mucus thus contributing to the viscous lung secretions characteristic of CF. Knowles, Clin. Chest. Med. 11: 75 (1986).

Adventitious bacteria and mycoplasmas enter the lung airways and establish colonies within the mucus. The thick mucus associated with CF isolates these pathogens from the immune system. Since mucociliary clearance is reduced in CF patients, bacterial clearance is also reduced. Lung congestion and infection are thus common. The prolonged presence of these pathogenic agents invariably initiates inflammatory reactions that compromise lung function. Bedrossian et al., Human Pathol. 7:195-204, 1976.

Mucus viscosity in CF lungs is in part due to the decreased hydration of the mucus as related to Cl⁻ channel malfunction and modification of sodium (Na⁺) ion concentration in the airway surface liquid (ASL) that change the rate of airway mucociliary clearance (MCC). The mechanisms involved in mucus transport have been studied in vitro and in vivo. CB, CF, and BE sputa are transported slowly by the mammalian ciliated epithelium of the mucus depleted bovine trachea (MDBT) (Wills et al, J. Clin. Invest. 97(1): 9-13, 1995). Slow transportability of diseased sputum on the MDBT may be linked to its low electrolyte/osmolyte content (Wills et al, J. Resp. Crit. Care Med. 151(4): 1255-1258, 1997). Indeed, diseased sputum is known to have low electrolyte content relative to plasma (Matthews et al, Am. Rev. Resp. Dis. 88: 199-204, 1963 ; Potter et al, Am. Rev. Resp. Dis. 67(1): 83-87, 1967 ; Tomkiewicz et al, Am. Rev. Resp. Dis. 148(4, Pt. 1): 1002-1007, 1993).

Further studies on the MDBT have shown that transportability of diseased mucus is markedly improved following treatment with sodium chloride (Wills et al 1995). Furthermore, clinical studies have shown that inhalation of hypertonic saline, or of the epithelial sodium channel (ENaC) blocker amiloride can significantly increase MCC in diseased patients (Robinson et al, Thorax 52(10): 900-903, 1997; App et al, Am. Rev. Resp. Dis. 141, 605-612, 1990). Recently, the relationship between mucus clearance and its ionic composition in vivo in the guinea-pig model of tracheal mucus velocity (TMV) has been elucidated. In vivo studies showed that a 5 minute aerosol of hypertonic saline transiently increased TMV. An increase in TMV was observed 1 min after hypertonic saline (14.4%) aerosol. TMV was 5.1 ± 1.0 mm.min⁻¹ (n=9) in 0.9% saline-exposed animals compared to 11.3 ± 1.3 mm.min⁻¹ in hypertonic saline exposed animals (n=9; p≤0.001)(Newton & Hall, 1997). Inhaled amiloride also caused an increase in TMV. A significant increase in TMV was observed 15 minutes after a 20

minute aerosol of amiloride (10mM). TMV was 3.2 ± 2.5 mm.min⁻¹ (n=9) in water-exposed animals compared to 8.1 ± 0.3 mm.min⁻¹ in amiloride-exposed animals (n=8; p $\leq$ 0.05) Newton et al, Ped. Pulm. S17, Abs. 364, 1998). These agents would appear to act by increasing the ionic content of airway surface liquid (ASL).

Clinical genetic evidence from subjects with systemic pseudohypoaldosteronism (SPHA) also supports the view that down-regulation of the activity of airway epithelial sodium channels will increase mucociliary clearance in the lung. SPHA patients with loss of function mutations in the genes for the epithelial sodium channel subunits had no sodium absorption from airways surfaces, they had increased sodium ion concentration in nasal surface liquid compared to normal subjects, and they had 4-fold increased lung mucociliary clearance rate compared to normal subjects (Kerem et al, New England J. Med. 341, 156-162, 1999).

Recently, a serine protease termed channel activating protease-1 (CAP-1) has been found in the apical membrane of amphibian *Xenopus* kidney epithelial cells (A6 cells) (Vallet et al, Nature 389(6651): 607-610, 1997). CAP-1 appears to modulate Na⁺ channel activity in these cells. Exposure of the apical membrane to the prototypical bovine Kunitz inhibitor, aprotinin, reduced transepithelial Na⁺ transport (Vallet et al 1997 : Chraïbi et al, J. Gen. Physio. 111(1): 127-138, 1998). The effect of Bukinin (1-170), a two Kunitz domain human homologue of bovine aprotinin (Delaria et al, J. Biol. Chem. 272(18): 12209-12214, 1997; Marlor et al, J. Biol. Chem. 272(18): 12202-12208, 1997), was evaluated using normal cultured human bronchial epithelial cell (HBE) short circuit current (Isc) in vitro (McAulay et al, Ped. Pulm. S17, Abs. 141, 1998). Bikunin (1.5ug.ml⁻¹ : 70nM) significantly inhibited 54% Na⁺ Isc in normal HBE cells (n=5-8 ; p $\leq$ 0.05). Overall, Bikunin (70nM) inhibited 58% of the baseline Isc in 90 minutes. In a further study, Bikunin (5ug.ml⁻¹) significantly inhibited 84% Na⁺ Isc in normal HBE cells (n=6; p $\leq$ 0.01) whilst the serpin-family serine protease inhibitor alpha(1)-protease inhibitor (α_1 -PI)(50 ug.ml⁻¹) was without a significant effect. In cultured human cystic fibrosis airway epithelial cells in vitro, Isc was inhibited by bikunin (1-170) (1 ug/mL), and was inhibited by aprotinin (20 ug/mL).

Two recent studies by a single research group have demonstrated a protease inhibitor induced effect on TMV. α_1 -PI (10mg) given either 30 min before antigen challenge, or 1 h after challenge, attenuated antigen-induced reduction in TMV in allergic sheep, 6h after challenge (O'Riordan et al, Am. J. Resp. Crit. Care Med. 97(5): 1522-1528, 1997). In Fig 1 in the O'Riordan et al 1997

WO 00/37099

PCT/GB99/04381

paper, the authors showed that α_1 -PI administered on its own (no antigen challenge) to the airways of allergic sheep, had no effect on baseline TMV over a 6h period. In the second study, α_1 -PI was given 6 h after antigen challenge and caused only a significant reversal of the antigen-induced fall in TMV at 24 h after challenge (O'Riordan et al, J. App. Physio. 85(3): 1086-1091, 1998). The authors argue that the mechanism for the effect of α_1 -PI is associated with its anti-neutrophil elastase property, where neutrophil elastase is believed to be the enzyme responsible for the reduced rate of mucociliary clearance in their model. They reasoned that α_1 -PI could be used to treat mucociliary dysfunction brought about by allergy-induced neutrophil elastase release in asthma (O'Riordan et al 1998); they did not speculate on a potential role in other respiratory diseases.

Brief Summary of the Invention

The instant invention is directed to the use of Kunitz-family serine protease inhibitors that stimulate the rate of mucociliary clearance (MCC) of mucus and sputum in the airways of the lung. Kunitz-serine protease inhibitors can be used to treat lung diseases such as Cystic Fibrosis (CF), Chronic Bronchitis (CB) and Bronchiectasis (BE) where the retention and accumulation of mucus is a major clinical problem. Until now, prior art has not associated protease inhibitors with the ability to increase the rate of MCC above baseline rate. Kunitz-type serine protease inhibitors can also be used to treat chronic sinusitis and glue ear where the retention and accumulation of mucus is a clinical problem.

The instant invention contemplates the use of serine protease inhibitor proteins which include Kunitz domains or Kunitz-like domains for use in a method for stimulating MCC. In one embodiment of the invention, bovine serine protease inhibitor proteins such as aprotinin and variants and fragments thereof such as the ones described in EP 821007, published January 28, 1998 may be used in practicing the invention.

In another embodiment of the invention, human serine protease inhibitors are contemplated for use in the method for stimulating the rate of MCC. Representative examples of human serine protease inhibitors include Bikunin and variants and fragments thereof such as the ones described in WO 97/33996, published September 18, 1997 (Bayer Corp.), and U.S. Patent No. 5,407,915, issued April 18, 1995 (Bayer AG).

Description of the Drawings

The invention will be better understood from a consideration of the following detailed description and claims, taken in conjunction with the drawings, in which:

Figure 1 depicts the nucleotide sequence of EST R35464 (SEQ ID NO:12) and the translation of this DNA sequence (labeled "ORF") which yielded an open reading frame with some sequence similarity to aprotinin. Amino acids 1-110 of the translation correspond to SEQ ID NO:13; amino acids 112-130 correspond to SEQ ID NO:72. The translation product contains 5 of the 6 cysteines in the correct spacing that is characteristic for Kunitz-like inhibitor domains (indicated in bold). The position normally occupied by the remaining cysteine (at codon 38) contained instead a phenylalanine (indicated by an asterisk).

Figure 2 depicts the nucleotide sequence of EST R74593 (SEQ ID NO:14), and the translation of this DNA sequence (labeled "ORF") which yielded an open reading frame with homology to the Kunitz class of serine protease inhibitor domains. Amino acids 3-22 of the translation correspond to SEQ ID NO:15; amino acids 24-131 correspond to SEQ ID NO:73; amino acids 136-166 correspond to SEQ ID NO:74. The translation product contained 6 cysteines in the correct spacing that is characteristic for Kunitz-like inhibitor domains (indicated in bold). However, the reading frame sequence includes stop codons at codon 3 and 23.

Figure 3 depicts a deduced nucleic acid sequence of human placental bikunin (SEQ ID NO:9) labeled "consensus" and matched with the translated protein amino acid sequence labeled "translated". Amino acids —18-179 of the translation correspond to SEQ ID NO:10; amino acids 184-214 correspond to SEQ ID NO:76. Also as comparison are shown the nucleic acid sequence for ESTs H94519 (SEQ ID NO:16), N39798 (SEQ ID NO:17), R74593 (corrected by the insertion of G at position 114) (SEQ ID NO:75) and R35464 (SEQ ID NO:12). The underlined nucleotides in the consensus sequence correspond to the site of PCR primers described in the Examples. Underlined amino acids in the translated consensus sequence are residues whose identity have been confirmed by amino acid sequencing of purified native human placental bikunin. Nucleotide and amino acid code are standard single letter code, "N" in the nucleic acid code indicates an unassigned nucleic acid, and "*" indicates a stop codon in the amino acid sequence.

Figure 4A depicts the original overlay of a series of ESTs with some nucleic acid sequence homology to ESTs encoding human placental bikunin, or portions thereof. Shown for reference are the relative positions of bikunin (7-64) and bikunin (102-159), labeled KID1 and KID2 respectively.

Figure 4B depicts a subsequent more comprehensive EST overlay incorporating additional ESTs. Numbers on the upper X-axis refer to length in base pairs, starting at the first base from the most 5' EST sequence. The length of each bar is in proportion to the length in base pairs of the individual ESTs including gaps. The EST accession numbers are indicated to the right of their respective EST bars.

Figure 4C depicts the corresponding alignment of the oligonucleotide sequences of each of the overlapping ESTs shown schematically in Figure 4B. The upper sequence (SEQ ID NO:51) labeled bikunin represents the consensus oligonucleotide sequence derived from the overlapping nucleotides at each position. The numbers refer to base-pair position within the EST map. The oligonucleotides in EST R74593 (SEQ ID NO:89) that are bold underlined (at map positions 994 and 1005) are base insertions observed in R74593 that were consistently absent in each of the other overlapping ESTs. In Figure 4C, N40851 corresponds to SEQ ID NO:77; N39876 corresponds to SEQ ID NO:78; R87894 corresponds to SEQ ID NO:79; H16866 corresponds to SEQ ID NO:80; R34808 corresponds to SEQ ID NO:81; T66058 corresponds to SEQ ID NO:82; N57450 corresponds to SEQ ID NO:83; N57374 corresponds to SEQ ID NO:84; R35464 corresponds to SEQ ID NO:85; H94519 corresponds to SEQ ID NO:86; N39798 corresponds to SEQ ID NO:87; H87300 corresponds to SEQ ID NO:88; R74593 corresponds to SEQ ID NO:89; R31730 corresponds to SEQ ID NO:90; R34701 corresponds to SEQ ID NO:91; H02982 corresponds to SEQ ID NO:92; R32676 corresponds to SEQ ID NO:93; T47439 corresponds to SEQ ID NO:94; R73968 corresponds to SEQ ID NO:95; H39840 corresponds to SEQ ID NO:96; H95233 corresponds to SEQ ID NO:97; H39841 corresponds to SEQ ID NO:98; N30199 corresponds to SEQ ID NO:99; T52966 corresponds to SEQ ID NO:100; N29508 corresponds to SEQ ID NO:101; N26919 corresponds to SEQ ID NO:102; N26910 corresponds to SEQ ID NO:103;

H16757 corresponds to SEQ ID NO:104; and N27732 corresponds to SEQ ID NO:105.

Figure 4D depicts the amino acid translation of the consensus oligonucleotide sequence for bikunin depicted in Figure 4C (SEQ ID NO.: 45).

Figure 4E depicts the nucleotide sequence (SEQ ID NO.: 46) and corresponding amino acid translation (SEQ ID NO.: 47) of a placental bikunin encoding sequence that was derived from a human placental cDNA library by PCR-based amplification.

Figure 4F depicts the nucleotide sequence (SEQ ID NO.: 48) and corresponding amino acid translation (SEQ ID NO.: 49) of a native human placental bikunin encoding clone that was isolated from a human placental lambda cDNA library by colony hybridization.

Figure 4G compares the alignment of the amino acid translated oligonucleotide sequences for placental bikunin obtained by EST overlay (SEQ ID NO.: 45), PCR based cloning (SEQ ID NO.: 47), and conventional lambda colony hybridization (SEQ ID NO.: 49).

Figure 5 shows a graph of purification of human placental bikunin from placental tissue after Superdex 75 Gel-Filtration. The plot is an overlay of the protein elution profile as measured by OD 280 nm (solid line), activity of eluted protein in a trypsin inhibition assay (% inhibition shown by circles), and activity of eluted protein in a kallikrein inhibition assay (% inhibition shown by squares).

Figure 6 shows a graph which plots the purification of human placental bikunin from placental tissue using C18 Reverse-Phase Chromatography. The plot is an overlay of the protein elution profile as measured by OD 215 nm (solid line), activity of eluted protein in a trypsin inhibition assay (% inhibition shown by circles), and activity of eluted protein in a kallikrein inhibition assay (% inhibition shown by squares).

Figure 7 depicts a silver stained SDS-PAGE gel of highly purified placental bikunin (lane 2), and a series of molecular size marker proteins (lane 1) of the indicated sizes in kilodaltons. Migration was from top to bottom.

Figure 8 shows the amount of trypsin inhibitory activity present in the cell-free fermentation broth from the growth of yeast strains SC101 (panel 8A) or WHL341 (panel 8B) that were stably transformed with a plasmid (pS604) that directs the expression of placental bikunin (102-159).

5 Figure 9 shows both a silver stained SDS-PAGE (Figure 9A) and a Western blot with anti-placental bikunin (102-159) pAb (Figure 9B) of cell-free fermentation broth from the growth of yeast strain SC101 (recombinants 2.4 and 2.5) that was stably transformed with a plasmid directing the expression of either bovine aprotinin, or placental bikunin (102-159). Migration was from top to
10 bottom.

Figure 10 is a photograph which shows a silver stained SDS-PAGE of highly purified placental bikunin (102-159) (lane 2) and a series of molecular size marker proteins (lane 1) of the indicated sizes in Kilodaltons. Migration was from top to bottom.

15 Figure 11 is a photograph which shows the results of Northern blots of mRNA from various human tissues that was hybridized to a ^{32}P labeled cDNA probe encoding either placental bikunin (102-159) (Figure 11A) or encoding placental bikunin (1-213) (Figure 11B). Migration was from top to bottom. The numbers to the right of each blot refer to the size in kilobases of the adjacent
20 RNA markers. The organs from which mRNA was derived is described under each lane of the blot.

Figure 12 depicts an immunoblot of placental derived placental bikunin with rabbit antiserum raised against either synthetic reduced placental bikunin (7-64) (Figure 12A) or 102-159 (Figure 12b). For each panel, contents were: molecular size
25 markers (lanes 1); native placental bikunin isolated from human placenta (lanes 2); synthetic placental bikunin (7-64) (lanes 3) and synthetic placental bikunin (102-159) (lanes 4). Tricine 10-20% SDS-PAGE gels were blotted and developed with protein A-purified primary polyclonal antibody (8 ug IgG in 20 ml 0.1% BSA/Tris-buffered saline (pH 7.5), followed by alkaline phosphatase-conjugated goat anti-rabbit
30 secondary antibody. Migration was from top to bottom.

Figure 13 depicts a Coomassie Blue stained 10-20% Tricine SDS-PAGE gel of 3 micrograms of highly purified placental bikunin (1-170) derived from a baculovirus / Sf9 expression system (lane 2). Lane 1 contains molecular size markers. Migration was from top to bottom.

35 Figure 14 depicts a comparison of the effect of increasing concentrations of either Sf9-derived human placental bikunin (1-170) (filled circles), synthetic placental bikunin (102-159) (open circles), or aprotinin (open squares) on the activated partial

thromboplastin time of human plasma. Clotting was initiated with CaCl_2 . The concentration of proteins are plotted versus the -fold prolongation in clotting time. The uninhibited clotting time was 30.8 seconds.

Figure 15 illustrates the effect of Bikunin at dosage levels of 2 μM and 0.2 μM relative to amiloride (100 μM) and Hank's Balanced salt solution (HBSS) vehicle (control) on potential differences in guinea pig trachea 3 hours post-treatment.

Figure 16 illustrates (a) the positioning of the instillment syringe and beta probe relative to the guinea pig trachea; (b) a representative graph for measurement of trachea mean velocity (TMV) using ^{32}P -labelled *S.cerevisiae*; and (c) the sustained increase in TMV in vivo in guinea pig in response to Bikunin (5 μg) relative to HBSS vehicle control at 1.5, 1.75, 2.0, 2.25 and 2.5 hours following tracheal instillment.

Figure 17 illustrates that Bikunin (70 nM) decreases sodium current in cultured human bronchial epithelial cells in vitro relative to amiloride (10 μM).

Figure 18 illustrates the effect of a 5 min aerosol of hypertonic saline (14.4%) on increasing TMV, following aerosol treatment in guinea pig trachea.

Figure 19 illustrates the effect of a 20 minute aerosol of amiloride (10 mM) on TMV, following aerosol treatment in guinea pig trachea.

Figure 20 illustrates that Bikunin (5 $\mu\text{g}/\text{mL}$), aprotinin (5 $\mu\text{g}/\text{mL}$), and aprotinin double mutein (0.5 $\mu\text{g}/\text{mL}$, 1.5 $\mu\text{g}/\text{mL}$ and 5 $\mu\text{g}/\text{mL}$) decrease sodium short circuit current in cultured human bronchial cells in vitro.

Figure 21 illustrates that Aprotinin (1 mg/mL) inhibited I_{sc} in vitro in human CF bronchial epithelial cells.

Figure 22 illustrates that Bikunin aerosol (3 mL of 3 mg/mL) significantly increased TMV in sheep relative to PBS control.

Figure 23 illustrates that Bikunin (50 $\mu\text{g}/\text{mL}$) inhibited sodium current in vitro in guinea pig tracheal epithelial cells over a 30 minute period.

Figure 24 illustrates that Bikunin (100 $\mu\text{g}/\text{mL}$) significantly inhibited sodium current in vitro in ovine tracheal epithelial cells over a 90 minute period.

Figure 25 illustrates that (a) exposure to LPS caused a significant PMN influx and that (b) Bikunin significantly inhibited potential difference in guinea pigs pre-exposed to LPS.

Figure 26 illustrates that Aprotinin double mutein (10 μg) increased TMV in vivo in the guinea pig relative to HBSS over a sustained period of 1.5 to 2.5 hours following administration.

Figure 27 represents a plasmid map of pBC-BK (CMV-IE = cytomegalovirus immediate early; DHFR = dihydrofolate reductase; AMP-r = ampicillin resistance)

Figure 28 illustrates that (a) CHO expressed Bikunin (1-170) (10 ug/mL) decreases sodium current in vitro in human CF bronchial epithelial cells over a 90 minute period and (b) CHO Bikunin (1-170) at 1, 5, and 10 ug/mL and Aprotinin at 20 ug/mL decreases sodium current at 90 minutes after apical application to human CF bronchial epithelial cells in vitro.

Figure 29 illustrates the process steps for purifying Bikunin (1-170) from a CHO cell expression system.

Figure 30A shows a graph of the migration of the isoforms of purified CHO bikunin (1-170) using C18 Reverse-Phase Chromatography. The plot is an overlay of the protein elution profile as measured by Absorbance at 280 nm (solid line) and the percentage of acetonitrile in 0.1% Trifluoroacetic acid used to elute the protein (diamonds).

Figure 30B is a photograph which shows a silver stained SDS-PAGE of purified bikunin (1-170) glycosylated isoforms (lanes 45-55) expressed from a CHO cell expression system and a series of molecular size marker proteins (between lanes 54 and 55) of the indicated sizes in Kilodaltons. Migration was from top to bottom.

Figure 31 is a photograph which shows a silver stained SDS-PAGE of N-Glycosidase F treated (lanes 2 and 4) and untreated (lanes 1 and 3) of CHO purified bikunin (1-170) isoforms and a series of molecular size marker proteins of the indicated sizes in Kilodaltons. Migration was from top to bottom.

Detailed Description of the Invention

The present invention relates to compositions comprising Kunitz-type serine protease inhibitor proteins and fragments thereof which stimulate the rate of mucociliary clearance of mucus and sputum in lung airways. The compositions also encompass a newly identified human protein herein called human placental bikunin that contains two serine protease inhibitor domains of the Kunitz class.

The present invention also provides methods for stimulating the rate of mucociliary clearance in patients with mucociliary dysfunction, wherein an effective amount of the disclosed serine protease inhibitors of the present invention, in a biologically compatible vehicle, is administered to the patient.

A preferred application for placental bikunin, isolated domains, and other variants is for stimulating mucociliary clearance in CF patients as part of disease therapy and management. These methods and compositions reduce or eliminate mucus and sputum buildup in lung airways in patients with chronic obstructive

lung disease, thereby reducing the risk of secondary lung infections and other adverse side effects, as well as avoiding or delaying the need for lung transplant surgery in CF patients.

The method of the present invention contemplates the use of aprotinin to stimulating MCC. Aprotinin has been shown to reduce transepithelial Na⁺ transport in the apical membrane of amphibian *Xenopus* kidney epithelial cells (A6 cells) (Vallet et al 1997 : Chraïbi et al 1998). The mechanism of aprotinin action has been proposed to involve inhibition of CAP-1, a protease involved in modulating Na⁺ channel activity in A6 cells. Bikunin (1-170), a two Kunitz domain human homologue of bovine aprotinin (Delaria et al 1997 : Marlor et al 1997), was also shown to significantly inhibit normal cultured human bronchial epithelial cell (HBE) short circuit current (Isc) in vitro (McAulay et al 1998). Bikunin (1.5ug.ml⁻¹ : 70nM) significantly inhibited 54% Na⁺ Isc in normal HBE cells (n=5-8 ; p≤0.05). Overall, Bikunin (70nM) inhibited 58% of the baseline Isc in 90 minutes. In a further study, Bikunin (5ug.ml⁻¹) significantly inhibited 84% Na⁺ Isc in normal HBE cells (n=6; p≤0.01) whilst the serpin-family serine protease inhibitor alpha(1)-protease inhibitor (α₁-PI)(50ug.mL⁻¹) was without a significant effect. In cultured human cystic fibrosis airway epithelial cells in vitro, Isc was inhibited by bikunin (1-170) (1 ug/mL), and was inhibited by aprotinin (20 ug/mL).

In light of these observations, Kunitz-type serine inhibitors such as aprotinin, placental bikunin and fragments thereof are contemplated as therapeutics for treating mucociliary dysfunction, including cystic fibrosis.

By "Kunitz inhibitor" is meant an inhibitor of proteases; structurally, a "Kunitz inhibitor" or "Kunitz domain" is a protein, or protein domain, typically of about 60 amino acids in length and containing three disulfide bonds. (See Laskowske & Kato, *Ann. Rev. Biochem.* 49, 593-626, 1980).

A significant advantage of the Kunitz domains of the serine protease inhibitor Bikunin and fragments and analogs thereof of the present invention is that they are human proteins, and also less positively charged than Trasylol® (Example 1), thereby reducing the risk of kidney damage on administration of large doses of the proteins. Being of human origin, the protein of the instant invention can thus be administered to human patients with significantly reduced risk of undesired immunological reactions as compared to administration of similar doses of Trasylol®. Furthermore, it was found that bikunin(102-159), bikunin(7-64), and bikunin(1-170) are significantly more potent inhibitors of plasma kallikrein than Trasylol® *in vitro* (Example 3, 4 and 10). Thus bikunin and

fragments thereof are expected to be more effective *in vivo* relative to aprotinin.

The amount of the pharmaceutical composition to be employed will depend on the recipient and the condition being treated. The requisite amount may be determined without undue experimentation by protocols known to those skilled in the art. Alternatively, the requisite amount may be calculated, based on a determination of the amount of target protease such as plasmin, kallikrein or prostatic acid phosphatase which must be inhibited in order to treat the condition. As the active materials contemplated in this invention are deemed to be nontoxic, treatment preferably involves administration of an excess of the optimally required amount of active agent.

For stimulating the rates of mucociliary clearance in patients with chronic obstructive lung disease, the proteins of the instant invention may be used like aprotinin Trasylol® while taking into account the differences in potency. The use of Trasylol® is outlined in the Physicians Desk Reference, 1995, listing for Trasylol® supplement A. Briefly, with the patient in a supine position, the loading dose of placental bikunin, isolated domain or other variant is given by infusion slowly over about 20 to 30 minutes. In general, a total dose of between about 2×10^6 KIU (kallikrein inhibitory units) and 8×10^6 KIU will be used, depending on such factors as patient weight and condition. Preferred loading doses are those that contain a total of 1 to 2 million kallikrein inhibitory units (KIU).

The proteins of the instant invention are employed in pharmaceutical compositions formulated in the manner known to the art. Such compositions contain active ingredient(s) plus one or more pharmaceutically acceptable carriers, diluents, fillers, binders, and other excipients, depending on the administration mode and dosage form contemplated. Examples of therapeutically inert inorganic or organic carriers known to those skilled in the art include, but are not limited to, lactose, corn starch or derivatives thereof, talc, vegetable oils, waxes, fats, polyols such as polyethylene glycol, water, saccharose, alcohols, glycerin and the like. Various preservatives, emulsifiers, dispersants, flavorants, wetting agents, antioxidants, sweeteners, colorants, stabilizers, salts, buffers and the like can also be added, as required to assist in the stabilization of the formulation or to assist in increasing bioavailability of the active ingredient(s) or to yield a formulation of acceptable flavor or odor in the case of oral, nasal or pulmonary dosing. The inhibitor employed in such compositions may be in the form of the original compound itself, or optionally, in the form of a pharmaceutically acceptable salt. The compositions so

formulated are selected as needed for administration of the inhibitor by any suitable mode known to those skilled in the art.

Parenteral administration modes include intravenous (*i.v.*), subcutaneous (*s.c.*), intraperitoneal (*i.p.*), and intramuscular (*i.m.*) routes. Intravenous
5 administration can be used to obtain acute regulation of peak plasma concentrations of the drug as might be needed. Alternatively, the drug can be administered at a desired rate continuously by *i.v.* catheter. Suitable vehicles include sterile, non-pyrogenic aqueous diluents, such as sterile water for injection, sterile-buffered solutions or sterile saline. The resulting composition is
10 administered to the patient prior to and/or during surgery by intravenous injection or infusion.

Improved half life and targeting of the drug to phagosomes such as neutrophils and macrophage involved in inflammation may be aided by entrapment of the drug in liposomes. It should be possible to improve the
15 selectivity of liposomal targeting by incorporating into the outside of the liposomes ligands that bind to macromolecules specific to target organs/tissues such as the GI tract and lungs. Alternatively, *i.m.* or *s.c.* deposit injection with or without encapsulation of the drug into degradable microspheres (e.g., comprising poly-DL-lactide-co-glycolide) or protective formulations containing
20 collagen can be used to obtain prolonged sustained drug release. For improved convenience of the dosage form it is possible to use an *i.p.* implanted reservoir and septum such as the percuseal system. Improved convenience and patient compliance may also be achieved by use of either injector pens (e.g., the Novo Pin or Q-pen) or needle-free jet injectors (e.g., from Bioject, Mediject or Becton
25 Dickinson). Precisely controlled release can also be achieved using implantable pumps with delivery to the desired site via a cannula. Examples include the subcutaneously implanted osmotic pumps available from ALZA such as the ALZET osmotic pump.

Oral delivery may be achieved by incorporating the drug into tablets,
30 coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions, suspensions or enteric coated capsules designed to release the drug into the colon where digestive protease activity is low. Examples of the latter include the OROS-CT/Osmet™ system of ALZA, and the PULSINCAP™ system of Scherer Drug Delivery Systems. Other systems use azo-crosslinked polymers that are
35 degraded by colon-specific bacterial azoreductases, or pH sensitive polyacrylate polymers that are activated by the rise in pH in the colon. The above systems may be used in conjunction with a wide range of available absorption enhancers.

Rectal delivery may be achieved by incorporating the drug into suppositories.

Nasal delivery may be achieved by incorporating the drug into bioadhesive particulate carriers (<200 mm) such as those comprising cellulose, polyacrylate or polycarbophil, in conjunction with suitable absorption enhancers such as phospholipids or acylcarnitines. Commercially available systems include those developed by Dan Biosys and Scios Nova.

For stimulating the rate of mucociliary clearance, the preferred mode of administration of the placental bikunin variants of the present invention is pulmonary delivery. The Kunitz-type serine protease inhibitors disclosed herein may be administered to the lungs of a subject by any suitable means, but are preferably administered by administering an aerosol suspension of respirable particles comprised of the active compound, which the subject inhales. The respirable particles may be liquid or solid. Micron-sized dry powders containing the medicament in a suitable carrier such as mannitol, sucrose or lactose may be delivered to the lung airway surface using dry powder inhalers such as those of Inhale™, Dura™, Fisons (Spinhaler™), and Glaxo (Rotahaler™), or Astra (Turbohaler™) propellant based metered dose inhalers. Solution formulations with or without liposomes may be delivered using nebulizers.

Aerosols of liquid particles comprising the proteins may be produced by any suitable means, such as with a pressure-driven aerosol nebulizer or an ultrasonic nebulizer. See, e.g., U.S. Pat. No. 4,501,729. Nebulizers are commercially available devices which transform solutions or suspensions of the active ingredient into a therapeutic aerosol mist either by means of acceleration of compressed gas, typically air or oxygen, through a narrow venturi orifice or by means of ultrasonic agitation. Suitable formulations for use in nebulizers consist of the active ingredient in a liquid carrier. The carrier is typically water (and most preferably sterile, pyrogen-free water) or a dilute aqueous alcoholic solution, preferably made isotonic with body fluids by the addition of, for example, sodium chloride. Optional additives include preservatives if the formulation is not made sterile, for example, methyl hydroxybenzoate, antioxidants, flavoring agents, volatile oils, buffering agents and surfactants.

Aerosols of solid particles comprising the protein may likewise be produced with any solid particulate medicament aerosol generator. Aerosol generators for administering solid particulate medicaments to a subject produce particles which are respirable, as explained above, and generate a volume of aerosol containing a predetermined metered dose of a medicament at a rate suitable for human administration. One illustrative type of solid particulate

aerosol generator is an insufflator. Suitable formulations for administration by insufflation include finely comminuted powders which may be delivered by means of an insufflator or taken into the nasal cavity in the manner of a snuff. In the insufflator, the powder (e.g., a metered dose thereof effective to carry out the treatments described herein) is contained in capsules or cartridges, typically made of gelatin or plastic, which are either pierced or opened in situ and the powder delivered by air drawn through the device upon inhalation or by means of a manually-operated pump. The powder employed in the insufflator consists either solely of the protein or of a powder blend comprising the protein, a suitable powder diluent, such as lactose, and an optional surfactant. A second type of illustrative aerosol generator comprises a metered dose inhaler. Metered dose inhalers are pressurized aerosol dispensers, typically containing a suspension or solution formulation of the active ingredient in a liquified propellant. During use these devices discharge the formulation through a valve adapted to deliver a metered volume, typically from 10 to 200 uL, to produce a fine particle spray containing the protein. Suitable propellants include certain chlorofluorocarbon compounds, for example, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane and mixtures thereof. The formulation may additionally contain one or more co-solvents, for example, ethanol, surfactants, such as oleic acid or sorbitan trioleate, antioxidants and suitable flavoring agents.

For metered dose inhaler or dry powder inhaler devices, the aerosol, whether formed from solid or liquid particles, may be produced by the aerosol generator at a rate of from about 5 to 150 liters per minute, more preferably from about 10 to 100 liters per minute, and most preferably for metered dose inhalers from about 10 to 50 liters per minute, and most preferably for dry powder inhalers about 60 liters per minute. Aerosols generated by nebulizer, jet or ultrasonic, may be produced by the aerosol generator at a rate of from about 1 to 100 liters per minute, more preferably from about 4 to 10 liters per minute. Aerosols containing greater amounts of protein may be administered more rapidly.

The dosage of the protease inhibitor will vary depending on the condition being treated and the state of the subject. The daily dose may be divided among one or several unit dose administrations. The daily dose by weight may range from about 0.01 to 20 milligrams of respirable particles for a human subject, depending upon the age and condition of the subject.

Solid or liquid particulate pharmaceutical formulations containing

protease inhibitors of the present invention should include particles of respirable size: that is, particles of a size sufficiently small to pass through the mouth and larynx upon inhalation and into the bronchi and alveoli of the lungs. In general, particles ranging from about 1 to 8 microns in size (more particularly, less than about 6 microns in size) are respirable. Particles of non-respirable size which are included in the aerosol tend to be deposited in the throat and swallowed, and the quantity of non-respirable particles in the aerosol is preferably minimized. For nasal administration, a particle size in the range of 10-500 microns is preferred to ensure retention in the nasal cavity.

10 In the manufacture of a formulation according to the invention, the protease inhibitor is typically admixed with, inter alia, an acceptable carrier. The carrier must, of course, be acceptable in the sense of being compatible with any other ingredients in the formulation and must not be deleterious to the patient. The carrier may be a solid or a liquid, or both, and is preferably formulated with the compound as a unit-dose formulation, for example, a capsule, which may contain from 0.5% to 99% by weight of the active compound. One or more active compounds may be incorporated in the formulations of the invention, which formulations may be prepared by any of the well-known techniques of pharmacy consisting essentially of admixing the components.

20 Compositions containing respirable dry particles of protease inhibitor may be prepared by grinding the inhibitor with a mortar and pestle, and then passing the micronized composition through a 400 mesh screen to break up or separate out large agglomerates.

25 The pharmaceutical composition may optionally contain a dispersant which serves to facilitate the formation of an aerosol. A suitable dispersant is lactose, which may be blended with the active agent in any suitable ratio (e.g., a 1 to 1 ratio by weight).

30 If desired, general ex vivo and in vivo gene therapy strategies may employed to deliver nucleic acid constructs encoding Kunitz-type serine protease inhibitor proteins such as Bikunin, Aprotinin or fragments and variants thereof such as the ones described in WO 97/33996 (Bayer Corp.) and U.S. Patent No. 5,407,915.(Bayer AG). Gene therapy techniques that are primarily virus-based have been used to transform pulmonary cells as a means for treating the manifestations of CF in the lung and associated extrapulmonary tissues. See 35 WO 93/03709, published March 3, 1993 which describes the use of retroviral and non-retroviral vectors (e.g., adenoviruses and adeno-associated viruses) for the stable expression of the CFTR gene in CF patients. Alternatively, non-viral

methods for delivery of exogenous nucleic acids are also known and are contemplated for use in the instant invention. See WO 93/12240, published June 24, 1993 and references cited therein, describing a transcription or expression cassettes including the coding sequence for a CFTR molecule operably joined to
5 regulatory sequences functional in a mammal. The nucleic acid constructs are then supplied to the airways and alveoli of the lung in a number of ways including aerosolized delivery alone or in combination with lipid-based complexes, e.g., Lipofectin.TM WO 95/26356, published October 5, 1995 describes representative examples of lipids useful for transfection. It is therefore
10 contemplated in the instant invention that nucleic acid molecules encoding Kunitz-type serine protease inhibitors such as Bikunin, Aprotinin or variants and fragments thereof may be similarly administered to lung airways by any suitable gene therapy method as a means for stimulating the rate of mucociliary clearance of mucus and sputum in a subject in need of such treatment.

15

Searching Human Sequence Data

The existence of a distinct human protein homologous in function to aprotinin, was deduced following a unique analysis of sequence entries to the expressed-sequence-tag data-base (hereafter termed dbEST) at the NCBI
20 (National Center for Biological Information, Maryland). Using the TBlastN algorithm (BLAST, or Basic Local Alignment Search Tool uses the method of Altschul et al., (1990) J. Mol Biol 215, 00 403-410, to search for similarities between a query sequence and all the sequences in a data-base, protein or nucleic acid in any combination), the data-base was examined for nucleotide sequences bearing
25 homology to the sequence of bovine pre-pro-aprotinin, Trasylol[®]. This search of numerous clones was selectively narrowed to two particular clones which could possibly encode for a deduced amino acid sequence that would correspond to a human protein homologous in function to aprotinin. The selected nucleic acid sequences were R35464 (SEQ ID NO: 12) and R74593 (SEQ ID NO: 14) that were
30 generated from a human placental nucleic acid library. The translated protein sequence in the longest open reading frame for R35464 (SEQ ID NO: 13) was missing one of the 6 cysteines that are critical for formation of the Kunitz-domain covalent structure, meaning that the nucleic acid sequence of R35464 could not yield a functional inhibitor. Similarly, the longest translated open reading frame
35 from clone R74593 (SEQ ID NO: 15) contained a stop codon 5' to the region encoding the Kunitz like sequence, meaning that this sequence, could not be translated to yield a functional secreted Kunitz domain. The significance of these

sequences alone was unclear. It was possible that they represented a) the products of pseudogenes, b) regions of untranslated mRNA, or c) the products of viable mRNA which had been sequenced incorrectly.

5 Discovery of Human Bikunin

To specifically isolate and determine the actual human sequence, cDNA primers were designed to be capable of hybridizing to sequences located 5' and 3' to the segment of cDNA encoding our proposed Kunitz like sequences found within R35464 and R74593. The primers used to amplify a fragment encoding the Kunitz like sequence of R74593 were CGAAGCTTCATCTCCGAAGCTCCAGACG (the 3' primer with a HindIII site; SEQ ID NO.:33) and AGGATCTAGACAATAATTACCTGACCAAGGA (the 5' primer with an XbaI site; SEQ ID NO.:34).

These primers were used to amplify by PCR (30 cycles) a 500 base pair product from a human placental cDNA library from Clontech (MATCHMAKER, Cat #HL4003AB, Clontech Laboratories, Palo Alto, CA), which was subcloned into Bluescript-SK+ and sequenced with the T3 primer with a Sequenase™ kit version 2.0. Surprisingly, the sequence of the fragment obtained using our primers was different from the sequence listed in the dbEST data base for clone R74593. In particular, our new sequence contained an additional guanosine base inserted 3' to the putative stop codon, but 5' to the segment encoding the Kunitz-like sequence (Figure 3). The insertion of an additional G shifted the stop codon out of the reading frame for the Kunitz-like domain (G at base pair 114 of the corrected sequence for R74593; Figure 3).

Subsequent query of the dbEST for sequences homologous to the Kunitz-like peptide sequence of R74593 yielded H94519 derived from human retina library and N39798. These sequences contained a Kunitz-like sequence that was almost identical to the Kunitz-like domain encoded in R35464 except that it contained all six of the characteristic cysteines. Overlay of each of the nucleotide sequences with that of R74593 (corrected by the insertion of G at b,p, 114) and R35464 was used to obtain a consensus nucleotide sequence for a partial human placental bikunin (SEQ ID NO.: 9; Figure 3). The translated consensus sequence yielded an open reading frame extending from residue -18 to +179 (Figure 3; full translation SEQ ID NO.: 10) that contained two complete Kunitz-like domain sequences, within the region of amino acid residues 17-64 and 102-159 respectively.

Further efforts attempted to obtain additional 5' sequence by querying

dbEST with the sequence of R35464. Possible matches from such searches, that possessed additional 5' sequence were then in turn used to re-query the dbEST. In such an iterative fashion, a series of overlapping 5' sequences were identified which included clones H16866, T66058, R34808, R87894, N40851 and N39876 (Figure 4). Alignment of some of these sequences suggested the presence of a 5' ATG which might serve as a start site for synthesis of the consensus translated protein sequence. From this selected information, it was now possible to selectively screen for, and determine the nucleic acid and polypeptide sequences of a human protein with homologous function to aprotinin.

Re-interrogation of the dbEST revealed a number of new EST entries shown schematically in Figure 4B. Overlap with these additional ESTs allowed us to construct a much longer consensus oligonucleotide sequence (Figure 4C) that extended both 5' and 3' beyond the original oligonucleotide sequence depicted in Figure 3. In fact, the new sequence of total length 1.6 kilobases extended all the way to the 3' poly-A tail. The increased number of overlapping ESTs at each base-pair position along the sequence improved the level of confidence in certain regions such as the sequence overlapping with the 3' end of EST R74593 (Figure 3). Several overlapping ESTs in this region corroborated two critical base deletions relative to R74593 (located as bold underlined in Figure 4C, map positions 994 and 1005). Translation of the new consensus sequence (Figure 4D) in the bikunin encoding frame yielded a form of placental bikunin that was larger (248 amino acids) than the mature sequence (179 amino acids) encoded from the original consensus (SEQ ID NO.: 1), and was terminated by an in-frame stop codon within the oligonucleotide consensus. The size increase was due to a frame shift in the 3' coding region resulting from removal of the two base insertions unique to EST R74593. The frame shift moved the stop codon of the original consensus (Figure 3) out of frame enabling read through into a new frame encoding the additional amino acid sequence. The new translation product (Figure 4D) was identical to the original protein consensus sequence (SEQ ID NO.: 1) between residues +1 to +175 (encoding the Kunitz domains), but contained a new C-terminal extension exhibiting a putative 24 residue long transmembrane domain (underlined in Figure 4D) followed by a short 31 residue cytoplasmic domain. The precise sequence around the initiator methionine and signal peptide was somewhat tentative due to considerable heterogeneity amongst the overlapping ESTs in this region.

Analysis of the protein sequence by Geneworks™, highlighted asparagine residues at positions 30 and 67 as consensus sites for putative N-linked

glycosylation. Asparagine 30 was not observed during N-terminal sequencing of the full length protein isolated from human placenta, consistent with it being glycosylated.

5 Cloning of Human Bikunin

The existence of a human mRNA corresponding to the putative human bikunin nucleotide sequence inferred from the analysis of Figure 3, was confirmed as follows. The nucleic acid primer hybridizing 5' to the Kunitz-encoding cDNA sequence of R35464 (b.p. 3-27 of consensus nucleotide sequence in Figure 3): GGTCTAGAGGCCGGGTCGTTTCTCGCCTGGCTGGGA (a 5' primer derived from R35464 sequence with an XbaI site; SEQ ID NO.: 35), and the nucleic acid primer hybridizing 3' to the Kunitz encoding sequence of R74593 (b.p. 680-700 of consensus nucleotide sequence in Figure 3), was used to PCR amplify, from a Clontech human placental library, a fragment of the size (ca. 670 b.p.) expected from a cDNA consensus nucleotide sequence encoding the placental bikunin sequence of Figure 3 (Shown schematically in Figure 4A).

Using a 5' primer hybridizing to a sequence in R87894 that is 126 b.p 5' to the putative ATG start site discussed above, (shown schematically in Figure 4A at b.p. 110) plus the same 3' primer to R74593 as used above, it was possible to amplify a fragment from a Clontech human placental library of the expected size (approximately 872 b.p) predicted by EST overlay (Shown schematically in Figure 4).

Sequencing of the 872 b.p. fragment showed it to contain nucleotide segment corresponding to b.p. 110 to 218 of EST R87894 at its 5' end and b.p. 310 to 542 of the consensus sequence for placental bikunin inferred from the EST overlay analysis (of Figure 3), at its 3' end. This 3' nucleotide sequence contained all of the Kunitz-like domain encoded by placental bikunin (102-159).

To obtain a cDNA encoding the entire extracellular region of the protein, the following 5' PCR primer: CACCTGATCGCGAGACCCC (SEQ ID NO.: 36) designed to hybridize to a sequence within EST R34808 was used with the same 3' primer to EST 74593 to amplify (30 cycles) an approximately 780 base-pair cDNA product from the human placental cDNA library. This product was gel purified, and cloned into the TA vector (Invitrogen) for DNA sequencing by the dideoxy method (Sanger F., et al., (1977) Proc. Natl. Acad. Sci (USA), 74, pp 5463-5467) with the following primers:

Vector Specific: GATTTAGGTGACACTATAG (SP6) (SEQ ID NO.: 37)

TAATACGACTCACTATAGGG (T7) (SEQ ID NO.: 38)

Gene Specific: TTACCTGACCAAGGAGGAGTGC (SEQ ID NO.: 39)
 AATCCGCTGCATTCCTGCTGGTG (SEQ ID NO.: 40)
 5 CAGTCACTGGGCCTTGCCGT (SEQ ID NO.: 41)

The resulting cDNA sequence is depicted in Figure 4E together with its translation product. At the nucleotide level, the sequence exhibited only minor differences from the consensus EST sequence (Figure 4D). Translation of the
 10 sequence yielded a coding sequence containing an in-frame initiator ATG site, signal peptide and mature placental bikunin sequence and transmembrane domain. The translated sequence of the PCR product was missing the last 12 amino acid residues from the cytoplasmic domain as a consequence of the choice of selection of the 3' primer for PCR amplification. This choice of 3' PCR primer
 15 (designed based on the sequence of R74593) was also responsible for the introduction of an artifactual S to F mutation at amino acid position 211 of the translated PCR-derived sequence. The signal peptide deduced from translation of the PCR fragment was somewhat different to that of the EST consensus.

To obtain a full length placental bikunin cDNA, the PCR derived product
 20 (Figure 4E) was gel purified and used to isolate a non-PCR based full length clone representing the bikunin sequence. The PCR derived cDNA sequence was labeled with ³²P-CTP by High Prime (Boehringer Mannheim) and used to probe a placental cDNA Library (Stratagene, Unizap™ λ library) using colony hybridization techniques. Approximately 2 X 10⁶ phage plaques underwent 3
 25 rounds of screening and plaque purification. Two clones were deemed full length (~1.5 kilobases) as determined by restriction enzyme analysis and based on comparison with the size of the EST consensus sequence (see above). Sequencing of one of these clone by the dideoxy method yielded the oligonucleotide sequence depicted in Figure 4F. The translation product from
 30 this sequence yielded a protein with inframe initiator methionine, signal peptide and mature placental bikunin sequence. The mature placental bikunin sequence was identical to the sequence of the mature protein derived by translation of the EST consensus although the signal peptide sequence lengths and sequences differed. Unlike the PCR derived product, the cDNA derived by colony
 35 hybridization contained the entire ectodomain, transmembrane domain, cytoplasmic domain and in-frame stop codon. In fact, the clone extended all the way to the poly-A tail. The initiator methionine was followed by a hydrophobic

signal peptide which was identical to the signal peptide encoded in the PCR derived clone. Subsequently, we expressed and purified soluble fragments of placental bikunin, Bikunin (1-170), from Sf9 cells (Example 9) and CHO cells (Example 17), and found them to be functional protease inhibitors (Examples 10 and 18). Furthermore, we isolated from human placenta a soluble fragment of placental bikunin which was also an active protease inhibitor (Example 7).

Based on the above observations, it seems that full length placental bikunin has the capacity to exist as a transmembrane protein on the surface of cells as well as a soluble protein. Other transmembrane proteins that contain Kunitz domains are known to undergo proteolytic processing to yield mixtures of soluble and membrane associated forms. These include two forms of the Amyloid Precursor Protein termed APP751 (Esch F., et al., (1990) Science, 248, pp 1122-1124) and APP 770 (Wang R., et al., (1991), J. Biol Chem, 266, pp16960-16964).

Contact activation is a process which is activated by exposure of damaged vascular surfaces to components of the coagulation cascade. Angiogenesis is a process that involves local activation of plasmin at endothelial surfaces. The specificity of placental bikunin and its putative capacity to anchor to cell surfaces, suggest that the physiologic functions of transmembranous placental bikunin may include regulation of contact activation and angiogenesis.

The amino acid sequences for placental bikunin (7-64), bikunin (102-159), and full length placental bikunin (Figure 4F) were searched against the PIR (Vers. 46.0) and PatchX (Vers. 46.0) protein databases as well as the GeneSeq (Vers. 20.0) protein database of patented sequences using the Genetics Computer Group program FastA. Using the Genetics Computer Group program TFASTA (Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85:2444-2448), these same protein sequences were searched versus the six-frame translations of the GenBank (Vers. 92.0 with updates to 1/26/96) and EMBL (modified Vers. 45.0) nucleotide databases as well as the GeneSeq (Vers. 20.0) nucleotide database of patented sequences. The EST and STS subsets of GenBank and EMBL were not included in this set of searches. The best matches resulting from these searches contained sequences which were only about 50% identical over their full length to the 58-amino acid protein sequence derived from our analysis of clones R74593 and R35464.

35

Isolation of Human Bikunin

As mentioned above, synthetic peptides corresponding to bikunin (7-64)

5

10

15.

20

25

30

35

-AEICLLPLDY GPCRALLRY YYRYRTQSCR QFLYGGCEGN ANNFYTWEC DDACWRI

22

WO 00/37099

PCT/GB99/04381

-PSFCYSPKDE GLCSANVTRY YFNPRYRTCD AFTYTGCGGN DNNFVSREDC KRACAKA

Amyloid precursor protein homologue (SEQ ID NO: 23)

-KAVCSQEAMT GPCRAVMPRT TFDLSKGKCV RFITGGCGGN RNNFESEDYC MAVCKAM

5

Aprotinin (SEQ ID NO: 24)

RPDFCLEPPYT GPCKARIIRY FYNAGAGLCQ TFVYGGCRAK RNNFKSAEDC MRTCGGA

Inter- α -trypsin inhibitor precursor (SEQ ID NOs: 25)

10 ----CQLGYSA GPCMGMTSRY FYNGTSMACE TFQYGGCMGN GNNFVTEKEC LQTC

Inter- α -trypsin inhibitor precursor (SEQ ID NOs: 26) VAACNLPIVR GPCRAFIQLW

AFDAVKGKCV LFPYGGCQGN GNKFYSEKEC REYCGVP

15 Amyloid precursor protein (SEQ ID NO: 27)

-EVCCSEQAET GPCRAMISRW YFDVTEGKCA PFFYGGCGGN RNNFDTEEYC MAVCGSA

Collagen α -3(VI) precursor (SEQ ID NO: 28)

----CKLPKDE GTCRDFILKW YYDPNTKSCA RFWYGGCGGN ENKFQSQKEC EKVC

20

HKL-B9 (SEQ ID NO: 29)

-PNVCAFPMEK GPCOTYMTRW FFNFETGECE LFAYGGCGGN SNNFLRKEKC EKFCCKFT

25 The placental bikunin, isolated domains or other variants of the present invention may be produced by standard solid phase peptide synthesis using either t-Boc chemistry as described by Merrifield R.B. and Barany G., in: The peptides, Analysis, Synthesis, Biology, 2, Gross E. et al., Eds. Academic Press (1980) Chapter 1; or using F-moc chemistry as described by Carpino L.A., and Han G.Y., (1970) J. Amer Chem Soc., 92, 5748-5749, and illustrated in Example 2.

30 Alternatively, expression of a DNA encoding the placental bikunin variant may be used to produce recombinant placental bikunin variants.

The instant invention provides for the use of a purified human serine protease inhibitor which can specifically inhibit kallikrein, that has been isolated from human placental tissue via affinity chromatography. The human serine

35 protein inhibitor, herein called human placental bikunin, contains two serine protease inhibitor domains of the Kunitz class. In one particular embodiment, the instant invention embodies a protein having the amino acid sequence:

WO 00/37099

PCT/GB99/04381

5 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179

10 (SEQ ID NO.: 1)

In a preferred embodiment the instant invention provides for a bikunin protein (Bikunin (1-170)) having the amino acid sequence:

15 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 20 ACMLRCFRQQ ENPPLPLGSK 170

(SEQ ID NO.: 52)

25 In one aspect, the biological activity of the protein useful in practicing the
 instant invention is that it can bind to and substantially inhibit the biological
 activity of trypsin, human plasma and tissue kallikreins, human plasmin and
 Factor XIIa. In a preferred embodiment, the present invention provides for a
 native human placental bikunin protein, in glycosylated form. In a further
 embodiment the instant invention encompasses native human bikunin protein
 30 which has been formed such that it contains at least one cysteine-cysteine
 disulfide bond. In a preferred embodiment, the protein contains at least one
 intra-chain cysteine-cysteine disulfide bond formed between a pair of cysteines
 selected from the group consisting of CYS11-CYS61, CYS20-CYS44, CYS36-
 CYS57, CYS106-CYS156, CYS115-CYS139, and CYS131-CYS152, wherein the
 35 cysteines are numbered according to the amino acid sequence of native human
 placental bikunin. One of ordinary skill will recognize that the protein of the
 instant invention may fold into the proper three-dimensional conformation, such
 that the biological activity of native human bikunin is maintained, where none,
 one or more, or all of the native intra-chain cysteine-cysteine disulfide bonds are
 40 present. In a most preferred embodiment, the protein of the instant invention is
 properly folded and is formed with all of the proper native cysteine-cysteine
 disulfide bonds.

45 Active protein for use in the instant invention can be obtained by
 purification from human tissue, such as placenta, or via synthetic protein
 chemistry techniques, as illustrated by the Examples below. It is also understood

that the protein for use in the instant invention may be obtained using molecular biology techniques, where self-replicating vectors are capable of expressing the protein of the instant invention from transformed cells. Such protein can be made as non-secreted, or secreted forms from transformed cells. In order to
5 facilitate secretion from transformed cells, to enhance the functional stability of the translated protein, or to aid folding of the bikunin protein, certain signal peptide sequences may be added to the NH2-terminal portion of the native human bikunin protein.

In one embodiment, the instant invention thus provides for the native
10 human bikunin protein with at least a portion of the native signal peptide sequence intact. Thus one embodiment of the invention provides for native human bikunin with at least part of the signal peptide, having the amino acid sequence:

15	A G S F L A W L G S L L L S G V L A	-1
	A D R E R S I H D F C L V S K V V G R C R A S M P	25
	R W W Y N V T D G S C Q L F V Y G G C D G N S N N	50
	Y L T K E E C L K K C A T V T E N A T G D L A T S	75
	R N A A D S S V P S A P R R Q D S E D H S S D M F	100
20	N Y E E Y C T A N A V T G P C R A S F P R W Y F D	125
	V E R N S C N N F I Y G G C R G N K N S Y R S E E	150
	A C M L R C F R Q Q E N P P L P L G S K V V V L A	175
	G A V S	179

25 (SEQ ID NO.: 2)

In a preferred embodiment, the instant invention provides for the use of a native human placental bikunin protein with part of the leader sequence intact, having the amino acid sequence of SEQ ID NO.: 52 with an intact leader segment having
30 the amino acid sequence:

MAQLCGL RRSRAFLALL GSLLLSGVLA -1

(SEQ ID NO.: 53)

35 In another embodiment, the instant invention provides for the use of bikunin protein with part of the leader sequence intact, having the amino acid sequence of SEQ ID NO.: 52 with the intact leader segment having the amino acid sequence:

MLR AEADGVSRLL GSLLLSGVLA -1

40

(SEQ ID NO.: 54)

In a preferred numbering system used herein the amino acid numbered +1 is assigned to the NH2-terminus of the amino acid sequence for native human

placental bikunin. One will readily recognize that functional protein fragments can be derived from native human placental bikunin, which will maintain at least part of the biological activity of native human placental bikunin, and act as serine protease inhibitors.

5 In one embodiment, the protein for use in the method of the instant invention comprises a fragment of native human placental bikunin, which contains at least one functional Kunitz-like domain, having the amino acid sequence of native human placental bikunin amino acids 7-159, hereinafter called "bikunin (7-159)". Thus the instant invention embodies a method that employs a
10 protein having the amino acid sequence:

15
R W W Y N V T D G S C Q L F V Y G G C D G N S N N 25
Y L T K E E C L K K C A T V T E N A T G D L A T S 50
R N A A D S S V P S A P R R Q D S E D H S S D M F 75
N Y E E Y C T A N A V T G P C R A S F P R W Y F D 100
V E R N S C N N F I Y G G C R G N K N S Y R S E E 125
A C M L R C F R Q 150
159

20 (SEQ ID NO.: 3)

where the amino acid numbering corresponds to that of the amino acid sequence of native human placental bikunin. Another functional variant of this embodiment can be the fragment of native human placental bikunin, which
25 contains at least one functional Kunitz-like domain, having the amino acid sequence of native human placental bikunin amino acids 11-156, bikunin (11-156)

30
R W W Y N V T D G S C Q L F V Y G G C D G N S N N 25
Y L T K E E C L K K C A T V T E N A T G D L A T S 50
R N A A D S S V P S A P R R Q D S E D H S S D M F 75
N Y E E Y C T A N A V T G P C R A S F P R W Y F D 100
V E R N S C N N F I Y G G C R G N K N S Y R S E E 125
A C M L R C 150
35 156

(SEQ ID NO.: 50).

One can recognize that the individual Kunitz-like domains are also fragments of the native placental bikunin. In particular, the instant invention
40 contemplates the use of a protein having the amino acid sequence of a first Kunitz-like domain consisting of the amino acid sequence of native human placental bikunin amino acids 7-64, hereinafter called "bikunin (7-64)". Thus in one embodiment the instant invention encompasses a protein which contains at least one Kunitz-like domain having the amino acid sequence:

45
I H D F C L V S K V V G R C R A S M P 25

WO 00/37099

PCT/GB99/04381

R W W Y N	V T D G S	C Q L F V	Y G G C D	G N S N N	50
Y L T K E	E C L K K	C A T V			64

(SEQ ID NO.: 4)

5

where the amino acid numbering corresponds to that of the amino acid sequence of native human placental bikunin. Another form of the protein of the instant invention can be a first Kunitz-like domain consisting of the amino acid sequence of native human placental bikunin amino acids 11-61, "bikunin (11-61)" having the amino acid sequence:

10

				C L V S K	V V G R C	R A S M P	25
R W W Y N	V T D G S	C Q L F V	Y G G C D	G N S N N			50
Y L T K E	E C L K K	C					61

15

(SEQ ID NO.: 5)

The instant invention also provides for a protein having the amino acid sequence of a Kunitz-like domain consisting of the amino acid sequence of native human placental bikunin amino acids 102-159, hereinafter called "bikunin (102-159)". Thus one embodiment the instant invention encompasses a protein which contains at least one Kunitz-like domain having the amino acid sequence:

20

25	Y E E Y	C T A N A	V T G P C	R A S F P	R W Y F D	125
	V E R N S	C N N F I	Y G G C R	G N K N S	Y R S E E	150
	A C M L R	C F R Q				159

(SEQ ID NO.: 6)

30

where the amino acid numbering corresponds to that of the amino acid sequence of native human placental bikunin. Another form of this domain can be a Kunitz-like domain consisting of the amino acid sequence of native human placental bikunin amino acids 106-156, "bikunin (106-156)" having the amino acid sequence:

35

		C T A N A	V T G P C	R A S F P	R W Y F D	125
V E R N S	C N N F I	Y G G C R	G N K N S	Y R S E E		150
A C M L R	C					156

40

(SEQ ID NO.: 7)

Thus one of ordinary skill will recognize that fragments of the native human bikunin protein can be made which will retain at least some of the native

protein biological activity. Such fragments can also be combined in different orientations or multiple combinations to provide for alternative proteins which retain some of, the same, or more biological activity of the native human bikunin protein.

5 One will readily recognize that biologically active protein employed in the method of the instant invention may comprise one or more of the instant Kunitz-like domains in combination with additional Kunitz-like domains from other sources. Biologically active protein of the method of the instant invention may comprise one or more of the instant Kunitz-like domains in combination with
10 additional protein domains from other sources with a variety of biological activities. The biological activity of the protein useful in practicing the instant invention can be combined with that of other known protein or proteins to provide for multifunctional fusion proteins having predictable biological activity. Thus, in one embodiment, the method of instant invention encompasses
15 the use of a protein which contains at least one amino acid sequence segment the same as, or functionally equivalent to the amino acid sequence of either SEQ ID NO.: 5 or SEQ ID NO.: 7.

An open reading frame which terminates at an early stop codon can still code for a functional protein. The instant invention encompasses such alternative
20 termination, and in one embodiment provides for the use of a protein of the amino acid sequence:

25	A D R E R	S I H D F	C L V S K	V V G R C	R A S M P	25
	R W W Y N	V T D G S	C Q L F V	Y G G C D	G N S N N	50
	Y L T K E	E C L K K	C A T V T	E N A T G	D L A T S	75
	R N A A D	S S V P S	A P R R Q	D S		92

(SEQ ID NO.: 8)

30 In one embodiment, the instant invention provides for the use of substantially purified, or recombinantly produced native human bikunin protein with an intact segment of the leader sequence, and at least a portion of the native transmembrane region intact. Thus one embodiment of the invention provides for the use of native human bikunin, with an intact leader sequence, and with at
35 least part of the transmembrane domain (underlined), having an amino acid sequence selected from:

40	EST		MLR	AEADGVSRL	GSLLLSGVLA	-1
	PCR		MAQLCGL	RRSRAFLAL	GSLLLSGVLA	-1
	λ cDNA		MAQLCGL	RRSRAFLAL	GSLLLSGVLA	-1
	EST	ADRERSIHDF	CLVSKVVGRC	RASMPRWWYN	VTGSCQLFV	YGGCDGNSNN 50
	PCR	ADRERSIHDF	CLVSKVVGRC	RASMPRWWYN	VTGSCQLFV	YGGCDGNSNN 50
	λ cDNA	ADRERSIHDF	CLVSKVVGRC	RASMPRWWYN	VTGSCQLFV	YGGCDGNSNN 50

EST YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
PCR YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
λcDNA YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100

5 EST NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
PCR NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
λcDNA NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150

10 EST ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
PCR ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
λcDNA ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200

EST QERALRTVWS SGDDKEQLVK NTYVL 225
PCR QERALRTVWS FGD 213
15 λcDNA QERALRTVWS SGDDKEQLVK NTYVL 225

where EST is EST derived consensus SEQ ID NO.: 45, PCR is PCR clone SEQ ID NO.:47, and λcDNA is lambda cDNA clone SEQ ID NO.:49. In a preferred embodiment a protein of the method of the instant invention comprises one of the amino acid sequence of SEQ ID NO.: 45, 47 or 49 wherein the protein has been cleaved in the region between the end of the last Kunitz domain and the transmembrane region (underlined).

The instant invention also embodies the use of the protein wherein the signal peptide is deleted. Thus the method of the instant invention provides for a protein having the amino acid sequence of SEQ ID NO.: 52 continuous with a transmembrane amino acid sequence:

EST VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
EST QERALRTVWS SGDDKEQLVK NTYVL 225

30 (SEQ ID NO.: 69)

a transmembrane amino acid sequence:

35 PCR VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
PCR QERALRTVWS FGD 213

(SEQ ID NO.: 68)

40 or a transmembrane amino acid sequence:

λcDNA VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
λcDNA QERALRTVWS SGDDKEQLVK NTYVL 225

45 (SEQ ID NO.: 67).

The invention also embodies the use the following proteins:

ADRERSIHDF CLVSKVVGRC RASMPRWYN VTDGSCQLFV YGGCDGNSNN 50
50 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150

12-01-2001

GB 009904381

ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 70),

5 or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 10 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 71).

15 The protein amino acid sequences for use in the instant invention clearly
 teach one skilled in the art the appropriate nucleic acid sequences which can be
 used in molecular biology techniques to produce the proteins for use in the
 instant invention. Thus, one embodiment of the instant invention provides for
 use of a nucleic acid sequence which encodes for a human bikunin having the

consensus DNA sequence of Figure 3 (SEQ ID NO.: 9), which translates into the amino acid sequence for native human placental bikunin sequence of Figure 3 (SEQ ID NO.: 10). In another embodiment, the instant invention provides for a consensus nucleic acid sequence of Figure 4C (SEQ ID NO.: 51) which encodes for an amino acid sequence of Figure 4D (SEQ ID NO.: 45).

In a preferred embodiment, the instant invention provides for the use of a nucleic acid sequence which encodes for native human placental bikunin having the DNA sequence of Figure 4F (SEQ ID NO.: 48) which encodes for the protein sequence of SEQ ID NO.: 49. In another embodiment, the instant invention provides for a nucleic acid sequence of Figure 4E (SEQ ID NO.: 46) which encodes for a protein sequence of SEQ ID NO.: 47.

One can easily recognize that certain allelic mutations, and conservative substitutions made in the nucleic acid sequence can be made which will still result in a protein amino acid sequence encompassed by the method of the instant invention. One of skill in the art can recognize that certain natural allelic mutations of the protein of the instant invention, and conservative substitutions of amino acids in the protein of the instant invention will not significantly alter the biological activity of the protein, and are encompassed by the instant invention.

The instant invention also provides for pharmaceutical compositions containing human placental bikunin and fragments thereof that are useful for stimulating MCC in patients impaired by mucociliary dysfunction.

The present invention also provides methods for stimulating MCC in a patient suffering from mucociliary dysfunction, wherein an effective amount of the disclosed human serine protease inhibitors of the present invention in a biologically compatible vehicle is administered to the patient.

The present invention also provides for a method for stimulating MCC that employs variants of placental bikunin, and the specific Kunitz domains described above, that contain amino acid substitutions that alter the protease specificity. Preferred sites of substitution are indicated below as positions Xaa¹ through Xaa³² in the amino acid sequence for native placental bikunin. Substitutions at Xaa¹ through Xaa¹⁶ are also preferred for variants of bikunin (7-64), while substitutions at Xaa¹⁷ through Xaa³² are preferred for variants of bikunin (102-159).

Thus the method of the present invention embodies the use of a protein having an amino acid sequence:

Ala Asp Arg Glu Arg Ser Ile Xaa¹ Asp Phe

10

WO 00/37099

PCT/GB99/04381

	Cys Leu Val Ser Lys Val Xaa ² Gly Xaa ³ Cys	20
	Xaa ⁴ Xaa ⁵ Xaa ⁶ Xaa ⁷ Xaa ⁸ Xaa ⁹ Trp Trp Tyr Asn	30
	Val Thr Asp Gly Ser Cys Gln Leu Phe Xaa ¹⁰	40
	Tyr Xaa ¹¹ Gly Cys Xaa ¹² Xaa ¹³ Xaa ¹⁴ Ser Asn Asn	50
5	Tyr Xaa ¹⁵ Thr Lys Glu Glu Cys Leu Lys Lys	60
	Cys Ala Thr Xaa ¹⁶ Thr Glu Asn Ala Thr Gly	70
	Asp Leu Ser Thr Ser Arg Asn Ala Ala Asp	80
	Ser Ser Val Pro Ser Ala Pro Arg Arg Gln	90
	Asp Ser Glu His Asp Ser Ser Asp Met Phe	100
10	Asn Tyr Xaa ¹⁷ Glu Tyr Cys Thr Ala Asn Ala	110
	Val Xaa ¹⁸ Gly Xaa ¹⁹ Cys Xaa ²⁰ Xaa ²¹ Xaa ²² Xaa ²³ Xaa ²⁴	120
	Xaa ²⁵ Trp Tyr Phe Asp Val Glu Arg Asn Ser	130
	Cys Asn Asn Phe Xaa ²⁶ Tyr Xaa ²⁷ Gly Cys Xaa ²⁸	140
	Xaa ²⁹ Xaa ³⁰ Lys Asn Ser Tyr Xaa ³¹ Ser Glu Glu	150
15	Ala Cys Met Leu Arg Cys Phe Arg Xaa ³² Gln	160
	Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys	170
	Val Val Val Leu Ala Gly Ala Val Ser	179
	(SEQ ID NO: 11).	

20 where Xaa¹ - Xaa³² each independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one of the amino acid residues Xaa¹-Xaa³² is different from the corresponding amino acid residue of the native sequence.

25 In the present context, the term "naturally occurring amino acid residue" is intended to indicate any one of the 20 commonly occurring amino acids, i.e., Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val.

30 By substituting one or more amino acids in one or more of the positions indicated above, it may be possible to change the inhibitor specificity profile of native placental bikunin or that of the individual Kunitz-like domains, bikunin(7-64) or bikunin (102-159) so that it preferentially inhibits other serine proteases such as, but not limited to, the enzymes of the complement cascade, TF/FVIIa, FXa, prostatic, thrombin, neutrophil elastase, cathepsin G or proteinase-3.

35 Examples of preferred variants of placental bikunin include those wherein Xaa¹ is an amino acid residue selected from the group consisting of His, Glu, Pro, Ala, Val or Lys, in particular wherein Xaa¹ is His or Pro; or wherein Xaa² is

an amino acid residue selected from the group consisting of Val, Thr, Asp, Pro, Arg, Tyr, Glu, Ala, Lys, in particular wherein Xaa² is Val or Thr; or wherein Xaa³ is an amino acid residue selected from the group consisting of Arg, Pro, Ile, Leu, Thr, in particular wherein Xaa³ is Arg or Pro; or wherein Xaa⁴ is an amino acid residue selected from the group consisting of Arg, Lys and Ser, Gln, in particular wherein Xaa⁴ is Arg or Lys; or wherein Xaa⁵ is an amino acid residue selected from the group consisting of Ala, Gly, Asp, Thr, in particular wherein Xaa⁵ is Ala; or wherein Xaa⁶ is an amino acid residue selected from the group consisting of Ser, Ile, Tyr, Asn, Leu, Val, Arg, Phe, in particular wherein Xaa⁶ is Ser or Arg; or wherein Xaa⁷ is an amino acid residue selected from the group consisting of Met, Phe, Ile, Glu, Leu, Thr and Val, in particular wherein Xaa⁷ is Met or Ile; or wherein Xaa⁸ is an amino acid residue selected from the group consisting of Pro, Lys, Thr, Gln, Asn, Leu, Ser or Ile, in particular wherein Xaa⁸ is Pro or Ile; or wherein Xaa⁹ is an amino acid residue selected from the group consisting of Arg, Lys or Leu, in particular wherein Xaa⁹ is Arg; or wherein Xaa¹⁰ is an amino acid residue selected from the group consisting of Val, Ile, Lys, Ala, Pro, Phe, Trp, Gln, Leu and Thr, in particular wherein Xaa¹⁰ is Val; or wherein Xaa¹¹ is an amino acid residue selected from the group consisting of Gly, Ser and Thr, in particular wherein Xaa¹¹ is Gly; or wherein Xaa¹² is an amino acid residue selected from the group consisting of Asp, Arg, Glu, Leu, Gln, Gly, in particular wherein Xaa¹² is Asp or Arg; or wherein Xaa¹³ is an amino acid residue selected from the group consisting of Gly and Ala; or wherein Xaa¹⁴ is an amino acid residue selected from the group consisting of Asn or Lys; or wherein Xaa¹⁵ is an amino acid residue selected from the group consisting of Gly, Asp, Leu, Arg, Glu, Thr, Tyr, Val, and Lys, in particular wherein Xaa¹⁵ is Leu or Lys; or wherein Xaa¹⁶ is an amino acid residue selected from the group consisting of Val, Gln, Asp, Gly, Ile, Ala, Met, and Val, in particular wherein Xaa¹⁶ is Val or Ala; or wherein Xaa¹⁷ is an amino acid residue selected from the group consisting of His, Glu, Pro, Ala, Lys and Val, in particular wherein Xaa¹⁷ is Glu or Pro; or wherein Xaa¹⁸ is an amino acid residue selected from the group consisting of Val, Thr, Asp, Pro, Arg, Tyr, Glu, Ala or Lys, in particular wherein Xaa¹⁸ is Thr; or wherein Xaa¹⁹ is an amino acid residue selected from the group consisting of Arg, Pro, Ile, Leu or Thr, in particular wherein Xaa¹⁹ is Pro; or wherein Xaa²⁰ is an amino acid residue selected from the group consisting of Arg, Lys, Gln and Ser, in particular wherein Xaa²⁰ is Arg or Lys; or wherein Xaa²¹ is an amino acid residue selected from the group consisting of Ala, Asp, Thr or Gly; in particular wherein Xaa²¹ is Ala; or wherein Xaa²² is an amino acid residue selected from

the group consisting of Ser, Ile, Tyr, Asn, Leu, Val, Arg or Phe, in particular wherein Xaa²² is Ser or Arg ; or wherein Xaa²³ is an amino acid residue selected from the group consisting of Met, Phe, Ile, Glu, Leu, Thr and Val, in particular wherein Xaa²³ is Phe or Ile; or wherein Xaa²⁴ is an amino acid residue selected from the group consisting of Pro, Lys, Thr, Asn, Leu, Gln, Ser or Ile, in particular wherein Xaa²⁴ is Pro or Ile; or wherein Xaa²⁵ is an amino acid residue selected from the group consisting of Arg, Lys or Leu, in particular wherein Xaa²⁵ is Arg; or wherein Xaa²⁶ is an amino acid residue selected from the group consisting of Val, Ile, Lys, Leu, Ala, Pro, Phe, Gln, Trp and Thr, in particular wherein Xaa²⁶ is Val or Ile; or wherein Xaa²⁷ is an amino acid residue selected from the group consisting of Gly, Ser and Thr, in particular wherein Xaa²⁷ is Gly; or wherein Xaa²⁸ is an amino acid residue selected from the group consisting of Asp, Arg, Glu, Leu, Gly or Gln, in particular wherein Xaa²⁸ is Arg; or wherein Xaa²⁹ is an amino acid residue selected from the group consisting of Gly and Ala; or wherein Xaa³⁰ is an amino acid residue selected from the group consisting of Asn or Lys; or wherein Xaa³¹ is an amino acid residue selected from the group consisting of Gly, Asp, Leu, Arg, Glu, Thr, Tyr, Val, and Lys, in particular wherein Xaa³¹ is Arg or Lys; or wherein Xaa³² is an amino acid residue selected from the group consisting of Val, Gln, Asp, Gly, Ile, Ala, Met, and Thr, in particular wherein Xaa³² is Gln or Ala.

The invention also relates to DNA constructs that encode the Placental bikunin protein variants of the present invention. These constructs may be prepared by synthetic methods such as those described in Beaucage S.L. and Caruthers M.H., (1981) Tetrahedron Lett, 22, pp1859-1862; Matteucci M.D and Caruthers M.H., (1981), J. Am. Chem. Soc. 103, p 3185; or from genomic or cDNA which may have been obtained by screening genomic or cDNA libraries with cDNA probes designed to hybridize with placental bikunin encoding DNA sequence. Genomic or cDNA sequence can be modified at one or more sites to obtain cDNA encoding any of the amino acid substitutions or deletions described in this disclosure.

The instant invention also relates to expression vectors containing the DNA constructs encoding the placental bikunin, isolated domains or other variants of the present invention that can be used for the production of recombinant placental bikunin variants. The cDNA should be connected to a suitable promoter sequence which shows transcriptional activity in the host cell of choice, possess a suitable terminator and a poly-adenylation signal. The cDNA encoding the placental bikunin variant can be fused to a 5' signal peptide

that will result in the protein encoded by the cDNA to undergo secretion. The signal peptide can be one that is recognized by the host organism. In the case of a mammalian host cell, the signal peptide can also be the natural signal peptide present in full length placental bikunin. The procedures used to prepare such
5 vectors for expression of placental bikunin variants are well known in the art and are for example described in Sambrook et al., Molecular Cloning: A laboratory Manual, Cold Spring Harbor, New York, (1989).

The instant invention also relates to transformed cells containing the DNA constructs encoding the placental bikunin, isolated domains or other variants of
10 the present invention that can be used for the production of recombinant placental bikunin variants. A variety of combinations of expression vector and host organism exist which can be used for the production of the placental bikunin variants. Suitable host cells include baculovirus infected Sf9 insect cells, mammalian cells such as BHK, CHO, Hela and C-127, bacteria such as E. coli,
15 and yeasts such as Saccharomyces cerevisiae. Methods for the use of mammalian, insect and microbial expressions systems needed to achieve expression of placental bikunin are well known in the art and are described, for example, in Ausubel F.M et al., Current Protocols in Molecular Biology, John Wiley & Sons (1995), Chapter 16. For fragments of placental bikunin containing a single Kunitz
20 inhibitor domain such as bikunin (7-64) and (102-159), yeast and E. coli expression systems are preferable, with yeast systems being most preferred. Typically, yeast expression would be carried out as described in US patent 5,164,482 for aprotinin variants and adapted in Example 5 of the present specification for placental bikunin (102-159). E.coli expression could be carried
25 out using the methods described in US patent 5,032,573. Use of mammalian and yeast systems are most preferred for the expression of larger placental bikunin variants containing both inhibitor domains such as the variant bikunin(7-159).

DNA encoding variants of placental bikunin that possess amino acid substitution of the natural amino sequence can be prepared for expression of
30 recombinant protein using the methods of Kunkel T.A., (1985) Proc. Natl. Acad. Sci USA 82, pp 488-492. Briefly, the DNA to be mutagenized is cloned into a single stranded bacteriophage vector such as M13. An oligonucleotide spanning the region to be changed and encoding the substitution is hybridized to the single stranded DNA and made double stranded by standard molecular biology
35 techniques. This DNA is then transformed into an appropriate bacterial host and verified by dideoxynucleotide sequencing. The correct DNA is then cloned into the expression plasmid. Alternatively, the target DNA may be mutagenized by

standard PCR techniques, sequenced, and inserted into the appropriate expression plasmid.

The following particular examples are offered by way of illustration, and not limitation, of certain aspects and preferred embodiments of the instant invention. All patents, patent applications and literature references cited in this application are incorporated by reference in their entirety.

Example 1

Preparation of synthetic placental bikunin (102-159)

Materials and methods/Reagents used. The fluorogenic substrate Tos-Gly-Pro-Lys-AMC was purchased from Bachem BioScience Inc (King of Prussia, PA). PNGB, Pro-Phe-Arg-AMC, Ala-Ala-Pro-Met-AMC, bovine trypsin (type III), human plasma kallikrein, and human plasmin were from Sigma (St. Louis, MO).

Recombinant aprotinin (Trasylol®) was from Bayer AG (Wuppertal, Germany). Pre-loaded Gln Wang resin was from Novabiochem (La Jolla, CA). Thioanisole, ethanedithiol and t-butyl methyl ether was from Aldrich (Milwaukee, WI).

Quantification of functional placental bikunin (7-64) (SEQ ID NO: 4) and (102-159)

The amount of trypsin inhibitory activity present in the refolded sample at various stages of purification was measured using GPK-AMC as a substrate. Bovine trypsin (200 pmoles) was incubated for 5 min at 37°C with bikunin (7-64) or (102-159), from various stages of purification, in buffer A (50 mM Hepes, pH 7.5, 0.1 M NaCl, 2 mM CaCl₂ and 0.01% triton X-100). GPK-AMC was added (20 µM final) and the amount of coumarin produced was determined by measuring the fluorescence (ex = 370 nm, em = 432 nm) on a Perkin-Elmer LS-50B fluorimeter over a 2 min. period. For samples being tested the % inhibition for each was calculated according to equation 1; where R₀ is the rate of fluorescence increase in the presence of inhibitor and R₁ is the rate determined in the absence of added sample. One unit of activity for the inhibitor is defined as the amount needed to achieve 50% inhibition in the assay using the conditions as described.

$$\% \text{ inhibition} = 100 \times [1 - R_0/R_1] \quad (1)$$

Synthesis. Placental bikunin (102-159)(SEQ ID NO: 6) was synthesized on

WO 00/37099

PCT/GB99/04381

an Applied Biosystems model 420A peptide synthesizer using NMP-HBTU Fmoc chemistry. The peptide was synthesized on pre loaded Gln resin with an 8-fold excess of amino acid for each coupling. Cleavage and deprotection was performed in 84.6% trifluoroacetic acid (TFA), 4.4% thioanisole, 2.2% ethanedithiol, 4.4% liquified phenol, and 4.4% H₂O for 2 hours at room temperature. The crude peptide was precipitated, centrifuged and washed twice in t-butyl methyl ether. The peptide was purified on a Dynamax 60A C18 reverse-phase HPLC column using a TFA/acetonitrile gradient. The final preparation (61.0 mg) yielded the correct amino acid composition and molecular mass by Electrospray mass spectroscopy (MH⁺ = 6836.1; calcd = 6835.5) for the predicted sequence:

YEEYCTANAV TGPCRASFPF WYFDVERNCS NNFIYGGCRG NKNSYRSEEA
CMLRCFRQ (SEQ ID NO.: 6)

15

Purification. Refolding of placental bikunin (102-159) was performed according to the method of Tam et al., (J. Am. Chem. Soc. 1991, 113; 6657-62). A portion of the purified peptide (15.2 mg) was dissolved in 4.0 ml of 0.1 M Tris, pH 6.0, and 8 M urea. Oxidation of the disulfides was accomplished by dropwise addition of a solution containing 23% DMSO, and 0.1 M Tris, pH 6.0 to obtain a final concentration of 0.5 mg/ml peptide in 20% DMSO, 0.1 M Tris, pH 6.0, and 1 M urea. The solution was allowed to stir for 24 hr at 25°C after which it was diluted 1:10 in buffer containing 50 mM Tris, pH 8.0, and 0.1 M NaCl. The material was purified using a kallikrein affinity column made by covalently attaching 30 mg of bovine pancreatic kallikrein (Bayer AG) to 3.5 mls of CNBr activated Sepharose* (Pharmacia) according to the manufacturers instructions. The refolded material was loaded onto the affinity column at a flow rate of 1 ml/min and washed with 50 mM Tris, pH 8.0, and 0.1 M NaCl until absorbance at 280 nm of the wash could no longer be detected. The column was eluted with 3 volumes each of 0.2 M acetic acid, pH 4.0 and 1.7. Active fractions were pooled (see below) and the pH of the solution adjusted to 2.5. The material was directly applied to a Vydac* C18 reverse-phase column (5 micron, 0.46 x 25 cm) which had been equilibrated in 22.5% acetonitrile in 0.1% TFA. Separation was achieved using a linear gradient of 22.5 to 40% acetonitrile in 0.1% TFA at 1.0 ml/min over 40 min. Active fractions were pooled, lyophilized, redissolved in 0.1% TFA, and stored at -20°C until needed.

35

Results. Synthetic placental bikunin (102-159) was refolded using

*Trademark

WO 00/37099

PCT/GB99/04381

20% DMSO as the oxidizing agent as described above, and purified by a 2-step purification protocol as shown below, to yield an active trypsin inhibitor (Table 1 below).

5 Table 1 Purification table for the isolation of synthetic placental bikunin (102-159)

TABLE 1						
Purification Step	Vol (ml)	mg/ml	mg	Units ^c (U)	SpA (U/mg)	Yield
8.0 M Urea	4.0	3.75 ^a	15.0	0	0	-
20% DMSO	32.0	0.47 ^a	15.0	16,162	1,078	100
Kallikrein affinity	9.8	0.009 ^b	0.09	15,700	170,000	97
C18	3.0	0.013 ^{ab}	0.04	11,964	300,000	74

^aProtein determined by AAA.

10 ^bProtein determined by OD280 nm using the extinction coefficient determined for the purified protein ($1.7 \times 10^4 \text{ Lmol}^{-1} \text{ cm}^{-1}$).

^cOne Unit is defined as the amount of material required to inhibit 50% of trypsin activity in a standard assay.

15 Chromatography of the crude refolded material over an immobilized bovine pancreatic kallikrein column selectively isolated 6.0% of the protein and 97% of the trypsin inhibitory activity present. Subsequent chromatography using C18 reverse-phase yielded a further purification of 2-fold, with an overall recovery of 74%. On RPHPLC, the reduced and refolded placental bikunin (102-159), exhibited elution times of 26.3 and 20.1 minutes, respectively. Mass spectroscopy analysis of the purified material revealed a molecular mass of 6829.8; a loss of 6 mass units from the starting material. This demonstrates the complete formation of the 3 disulfides predicted from the peptide sequence.

20 The isoelectric points of the purified, refolded synthetic placental bikunin (102-159) was determined using a Multiphor^{*} II Electrophoresis System (Pharmacia) run according to the manufacturers suggestions, together with pI standards, using a precast Ampholine[®] PAGplate (pH 3.5 to 9.5) and focused for 1.5 hrs. After staining, the migration distance from the cathodic edge of the gel to the different protein bands was measured. The pI of each unknown was determined by using a standard curve generated by a plot of the migration distance of standards versus the corresponding pI's. With this technique, the pI of placental bikunin (102-159) was determined to be 8.3, in agreement with the value predicted from the amino acid sequence. This is lower than the value of

*Trademark

WO 00/37099

PCT/GB99/04381

10.5 established for the pI of aprotinin. (Tenstad et al., 1994, Acta Physiol. Scand. 152:33-50).

Example 2

5 Preparation of synthetic placental bikunin (7-64)

Placental bikunin (7-64) (SEQ ID NO: 4) was synthesized, refolded and purified essentially as described for placental bikunin (102-159) (SEQ ID NO:6) in Example 1 but with the following modifications: during refolding, the synthetic peptide was stirred for 30 hr as a solution in 20% DMSO at 25°C; purification by
10 C18 RP-HPLC was achieved with a linear gradient of 25 to 45% acetonitrile in 0.1% TFA over 40 min (1ml/min). Active fractions from the first C18 run were reapplied to the column and fractionated with a linear gradient (60 min, 1 ml/min) of 20 to 40% acetonitrile in 0.1% TFA.

15 **Results.** The final purified reduced peptide exhibited an MH⁺ = 6563, consistent with the sequence:

IHDFCLVSKV VGRCRASMPR WWYNVTDGSC QLFVYGGCDG NSNNYLTKEE
CLKKCATV (SEQ ID NO.: 4)

20

The refolding and purification yielded a functional Kunitz domain that was active as an inhibitor of trypsin (Table 2 below).

25 **Table 2A Purification table for the isolation of synthetic placental bikunin (7-64)**

TABLE 2A						
Purification Step	Vol (ml)	mg/ml	mg	Units (U)	SpA (U/mg)	Yield
8.0 M Urea	8.0	2.5	20.0	0	0	-
20% DMSO	64.0	0.31	20.0	68,699	3,435	100
Kall affinity pH 4.0	11.7	0.10	1.16	43,333	36,110	62
Kall affinity pH 1.7	9.0	0.64	5.8	4972	857	7.2
C18-1	4.6	0.14	0.06	21,905	350,143	31.9
C18-2	1.0	0.08	0.02	7,937	466,882	11.5

The purified refolded protein exhibited an $MH^+ = 6558$, i.e. 5 ± 1 mass units less than for the reduced peptide. This demonstrates that refolding caused the formation of at least one appropriate disulfide bond.

5 The pI of placental bikunin (7-64) was determined using the methods employed to determine the pI of placental bikunin (102-159). Placental bikunin (7-64) exhibited a pI that was much higher than the predicted value ($pI = 7.9$). Refolded placental bikunin (7-64) migrated to the cathodic edge of the gel (pH 9.5) and an accurate pI could not be determined under these conditions.

10 *Continued Preparation of synthetic placental bikunin (7-64)*

Because the synthetic placental bikunin (7-64) may not have undergone complete deprotection prior to purification and refolding, refolding was repeated using protein which was certain to be completely deprotected. Placental bikunin (7-64) was synthesized, refolded and purified essentially as described for placental
15 bikunin (102-159) but with the following modifications: during refolding, the synthetic peptide (0.27 mg/ml) was stirred for 30 hr as a solution in 20% DMSO at 25 C; purification by C18 RP-HPLC was achieved with a linear gradient of 22.5 to 50% acetonitrile in 0.1% TFA over 40 min (1 ml/min).

20 **Results.** The final purified reduced peptide exhibited an $MH^+ = 6567.5$, consistent with the sequence:

IHDFCLVSKV VGRCRASMPRW WYNVTDGSC QLFVYGGCDG NSNNYLTKKE
CLKKCATV (SEQ ID NO. : 4)

25 The refolding and purification yielded a functional Kunitz domain that was as active as an inhibitor of trypsin (Table 2B below).

Table 2B Purification table for the isolation of synthetic placental bikunin (7-64)

TABLE 2B						
Purification Step	Vol (ml)	mg/ml	mg	Units (U)	SpA (U/mg)	Yield
8.0 M Urea	4.9	2.1	10.5	0	0	-
20% DMSO	39.0	0.27	10.5	236,000	22,500	100
Kallikrein Affinity (pH 2)	14.5	0.3	0.43	120,000	279,070	50.9
C18 Reverse-Phase	0.2	1.2	0.24	70,676	294,483	30.0

The purified refolded protein exhibited an $MH^+ = 6561.2$, i.e. 6.3 mass units less than for the reduced peptide. This demonstrates that refolding caused the formation of the expected three disulfide bonds.

The pI of refolded placental bikunin (7-64) was determined using the methods employed to determine the pI of placental bikunin (102-159). Refolded placental bikunin (7-64) exhibited a pI of 8.85, slightly higher than the predicted value (pI = 7.9).

Example 3

In vitro specificity of functional placental bikunin fragment (102-159)

Proteases. Bovine trypsin, human plasmin, and bovine pancreatic kallikrein quantitation was carried out by active site titration using p-nitrophenyl p'-guanidinobenzoate HCl as previously described (Chase, T., and Shaw, E., (1970) *Methods Enzymol.* 19, 20-27). Human kallikrein was quantitated by active site titration using bovine aprotinin as a standard and PFR-AMC as a substrate assuming a 1:1 complex formation. The K_m for GPK-AMC with trypsin and plasmin under the conditions used for each enzyme was 29 μM and 726 μM , respectively; the K_m for PFR-AMC with human plasma kallikrein and bovine pancreatic kallikrein was 457 μM and 81.5 μM , respectively; the K_m for AAPR-AMC with elastase was 1600 μM . Human tissue kallikrein (Bayer, Germany) quantification was carried out by active site titration using p-nitrophenyl p'-guanidinobenzoate HCl as previously described (Chase, T., and Shaw, E., (1970) *Methods Enzymol.* 19, 20-27).

Inhibition Kinetics: The inhibition of trypsin by placental bikunin (102-159) (described in Example 1) or aprotinin was measured by the incubation of 50 pM trypsin with placental bikunin (102-159) (0-2 nM) or aprotinin (0-3 nM) in buffer A in a total volume of 1.0 ml. After 5 min. at 37°C, 15 μl of 2 mM GPK-AMC was added and the change in fluorescence (as above) was monitored. The

WO 00/37099

PCT/GB99/04381

inhibition of human plasmin by placental bikunin (102-159) and aprotinin was determined with plasmin (50 pM) and placental bikunin (102-159) (0-10 nM) or aprotinin (0-4 nM) in buffer containing 50 mM Tris-HCl (pH 7.5), 0.1 M NaCl, and 0.02% triton x-100. After 5 min. incubation at 37°C, 25 µl of 20 mM GPK-AMC was added and the change in fluorescence monitored. The inhibition of human plasma kallikrein by placental bikunin (102-159) or aprotinin was determined using kallikrein (2.5 nM) and placental bikunin (102-159) (0-3 nM) or aprotinin (0-45 nM) in 50 mM Tris-HCl (pH 8.0), 50 mM NaCl, and 0.02% triton x-100. After 5 min. at 37°C 15 µl of 20 mM PFR-AMC was added and the change in fluorescence monitored. The inhibition of bovine pancreatic kallikrein by placental bikunin (102-159) and aprotinin was determined in a similar manner with kallikrein (92 pM), placental bikunin (102-159) (0-1.6 nM) and aprotinin (0-14 pM) and a final substrate concentration of 100 µM. The apparent inhibition constant K_i^* was determined using the nonlinear regression data analysis program Enzfitter software (Biosoft, Cambridge, UK): The kinetic data from each experiment were analyzed in terms of the equation for a tight binding inhibitor:

$$V_i/V_o = 1 - (E_o + I_o + K_i^* - [(E_o + I_o + K_i^*)^2 - 4 E_o I_o]^{1/2}) / 2 E_o \quad (2)$$

where V_i/V_o is the fractional enzyme activity (inhibited vs. uninhibited rate), and E_o and I_o are the total concentrations of enzyme and inhibitor, respectively. K_i values were obtained by correcting for the effect of substrate according to the equation:

$$K_i = K_i^* / (1 + [S_o]/K_m) \quad (3)$$

(Boudier, C., and Bieth, J. G., (1989) *Biochim Biophys Acta*. 995: 36-41)

For the inhibition of human neutrophil elastase by placental bikunin (102-159) and aprotinin, elastase (19 nM) was incubated with placental bikunin (102-159) (150 nM) or aprotinin (0-7.5 µM) in buffer containing 0.1 M Tris-HCl (pH 8.0), and 0.05% triton X-100. After 5 min at 37°C, AAPM-AMC (500 µM or 1000 µM) was added and the fluorescence measured over a two-minute period. K_i values were determined from Dixon plots of the form $1/V$ versus $[I]$ performed at two different substrate concentrations (Dixon et al., 1979).

The inhibition of human tissue kallikrein by aprotinin, placental bikunin fragment (7-64) or placental bikunin fragment (102-159) was measured by the

incubation of 0.35 nM human tissue kallikrein with placental bikunin (7-64) (0-40 nM) or placental bikunin (102-159) (0-2.5 nM), or aprotinin (0-0.5 nM) in a 1 ml reaction volume containing 50 mM Tris-HCl buffer pH 9.0, 50 mM NaCl, and 0.1% triton x-100. After 5 min. at 37°C, 5 ul of 2 mM PFR-AMC was added achieving 10 uM final and the change in fluorescence monitored. The K_m for PFR-AMC with human tissue kallikrein under the conditions employed was 5.7 uM. The inhibition of human factor Xa (American Diagnostica, Inc, Greenwich, CT) by synthetic placental bikunin (102-159), recombinant placental bikunin, and aprotinin was measured by the incubation of 0.87 nM human factor Xa with increasing amounts of inhibitor in buffer containing 20 mM Tris (pH 7.5), 0.1 M NaCl, and 0.1% BSA. After 5 min. at 37°C, 30 ul of 20 mM LGR-AMC (Sigma) was added and the change in fluorescence monitored. The inhibition of human urokinase (Sigma) by Kunitz inhibitors was measured by the incubation of urokinase (2.7 ng) with inhibitor in a total volume of 1 ml buffer containing 50 mM Tris-HCl (pH 8.0), 50 mM NaCl, and 0.1% Triton x-100. After 5 min. at 37°C, 35 ul of 20 mM GGR-AMC (Sigma) was added and the change in fluorescence monitored. The inhibition of Factor XIa (from Enzyme Research Labs, Southbend, IN) was measured by incubating FXIa (0.1 nM) with either 0 to 800 nM placental bikunin (7-64), 0 to 140 nM placental bikunin (102-159) or 0 to 40 uM aprotinin in buffer containing 50 mM Hepes pH 7.5, 100 mM NaCl, 2 mM $CaCl_2$, 0.01% triton x-100, and 1% BSA in a total volume of 1 ml. After 5 min at 37 C, 10 ul of 40 mM Boc-Glu(OBzl)-Ala-Arg-AMC (Bachem Biosciences, King of Prussia, PA) was added and the change in fluorescence monitored.

Results: A direct comparison of the inhibition profiles of placental bikunin (102-159) and aprotinin was made by measuring their inhibition constants with various proteases under identical conditions. The K_i values are listed in Table 3 below.

Table 3 Ki values for the inhibition of various proteases by bikunin (102-159)

TABLE 3				
Protease (concentration)	Bikunin (102-159) Ki (nM)	Aprotinin Ki (nM)	Substrate (concentration)	Km (mM)
Trypsin (48.5 pM)	0.4	0.8	GPK-AMC (0.03 mM)	0.022
Chymotrypsin (5 nM)	0.24	0.86	AAPF-pNA (0.08 mM)	0.027
Bovine Pancreatic Kallikrein (92.0 pM)	0.4	0.02	PFR-AMC (0.1 mM)	0.08
Human Plasma Kallikrein (2.5 nM)	0.3	19.0	PFR-AMC (0.3 mM)	0.46
Human Plasmin (50 pM)	1.8	1.3	GPK-AMC (0.5 mM)	0.73
Human Neutrophil Elastase (19 nM)	323.0	8500.0	AAPM-AMC (1.0 μ M)	1.6
Factor XIIa	>300.0	12,000.0	PFR-AMC (0.2 μ M)	0.35
Human Tissue Kallikrein (0.35 nM)	0.13	0.004	PFR-AMC (10 μ M)	0.0057
factor Xa (0.87 nM)	274	N.I. at 3 μ M	LGR-AMC (0.6 mM)	N.D.
Urokinase	11000	4500	GGR-AMC (0.7 mM)	N.D.
factor Xia (0.1 nM)	15	288	E(OBz)AR-AMC (0.4 mM)	0.46

Placental bikunin (102-159) and aprotinin inhibit bovine trypsin and human plasmin to a comparable extent under the conditions employed. Aprotinin inhibited elastase with a K_i of 8.5 μ M. Placental bikunin (102-159) inhibited elastase with a K_i of 323 nM. The K_i value for the placental bikunin (102-159) inhibition of bovine pancreatic kallikrein was 20-fold higher than that of aprotinin inhibition. In contrast, placental bikunin (102-159) is a more potent inhibitor of human plasma kallikrein than aprotinin and binds with a 56-fold higher affinity.

Because placental bikunin (102-159) is greater than 50 times more potent than Trasylol[®] as an inhibitor of kallikrein, smaller amounts of human placental bikunin, or fragments thereof (i.e. placental bikunin (102-159)) are needed than Trasylol[®] in order to maintain the effective patient doses of inhibitor in KIU. This reduces the cost per dose of the drug and reduces the likelihood of adverse nephrotoxic effects upon re-exposure of the medicament to patients. Furthermore, the protein is human derived, and thus much less immunogenic in man than aprotinin which is derived from cows. This results in significant reductions in the risk of incurring adverse immunologic events upon re-exposure of the medicament to patients.

Example 4**In vitro specificity of functional placental bikunin fragment (7-64)**

In vitro specificity of the functional human placental bikunin (7-64) described in Example 2 was determined using the materials and methods as described in the Examples above.

Results: The table below shows the efficacy of placental bikunin (7-64) as an inhibitor of various serine proteases *in vitro*. Data is shown compared against data obtained for screening inhibition using either placental bikunin (102-159), or aprotinin (Trasylol®).

Table 4 A Ki values for the inhibition of various proteases by bikunin(7-64)

TABLE 4A			
Protease (concentration)	bikunin(7-64) Ki (nM)	Aprotinin Ki (nM)	Bikunin (102-159) Ki (nM)
Trypsin (48.5 pM)	0.17	0.8	0.4
Bovine Pancreatic Kallikrein (92.0 pM)	0.4	0.02	0.4
Human Plasma Kallikrein (2.5 nM)	2.4	19.0	0.3
Human Plasmin (50 pM)	3.1	1.3	1.8
Bovine chymotrypsin (5 nM)	0.6	0.9	0.2
Factor XIIa	>300	12000	>300
elastase	>100	8500	323

The results show that the amino acid sequence encoding placental bikunin (7-64) can be refolded to obtain an active serine protease inhibitor that is effective against at least four trypsin-like serine proteases.

Table 4B below also shows the efficacy of refolded placental bikunin (7-64) as an inhibitor of various serine proteases *in vitro*. Refolded placental bikunin (7-64) was prepared from protein that was certain to be completely deprotected prior to purification and refolding. Data is shown compared against data obtained for screening inhibition using either placental bikunin (102-159), or aprotinin (Trasylol®).

Table 4B Ki values for the inhibition of various proteases by refolded bikunin (7-64)

TABLE 4B			
Protease (concentration)	bikunin (7-64) Ki (nM)	Aprotinin Ki (nM)	bikunin (102-159) Ki (nM)
Trypsin (50 pM)	0.2	0.8	0.3
Human Plasma Kallikrein (0.2 nM)	0.7	19.0	0.7
Human Plasmin (50 pM)	3.7	1.3	1.8
Factor XIIa	not done	12,000	4,500
Factor XIa (0.1 nM)	200	288	15
Human Tissue Kallikrein	2.3	0.004	0.13

5 Suprisingly, placental bikunin (7-64) was more potent than aprotinin at inhibiting human plasma kallikrein, and at least similar in efficacy as a plasmin inhibitor. These data show that placental bikunin (7-64) is at least as effective as aprotinin, using *in vitro* assays, and that one would expect better or similar potency *in vivo*.

10 Example 5

Expression of placental bikunin variant (102-159) in yeast

15 The DNA sequence encoding placental bikunin 102-159 (SEQ ID NO.: 6) was generated using synthetic oligonucleotides. The final DNA product consisted (5' to 3') of 15 nucleotides from the yeast α -mating factor propeptide sequence fused to the in-frame cDNA sequence encoding placental bikunin (102-159), followed by an in-frame stop codon. Upon cloning into a yeast expression vector pS604, the cDNA would direct the expression of a fusion protein comprising an N-terminal yeast α -mating factor propeptide fused to the 58 amino acid sequence of placental bikunin (102-159). Processing of this fusion protein at a KEX-2 cleavage site at the junction between the α -mating factor and Kunitz domain was designed to liberate the Kunitz domain at its native N-terminus.

25 A 5' sense oligonucleotide of the following sequence and containing a HindIII site for cloning was synthesized:

WO 00/37099

PCT/GB99/04381

GAA GGG GTA AGC TTG GAT AAA AGA TAT GAA GAA TAC TGC ACC GCC
 AAC GCA GTC ACT GGG CCT TGC CGT GCA TCC TTC CCA CGC TGG TAC
 TTT GAC GTG GAG AGG (SEQ ID NO. : 42)

- 5 A 3' antisense oligonucleotide of the following sequence and containing both a BamHI site for cloning and a stop codon was synthesized:

CGC GGA TCC CTA CTG GCG GAA GCA GCG GAG CAT GCA GGC CTC CTC
 10 AGA GCG GTA GCT GTT CTT ATT GCC CCG GCA GCC TCC ATA GAT GAA
 GTT ATT GCA GGA GTT CCT CTC CAC GTC AAA GTA CCA GCG
 (SEQ ID NO. : 43)

- The oligonucleotides were dissolved in 10 mM Tris buffer pH 8.0 containing 1 mM EDTA, and 12 ug of each oligo were added combined and brought to 0.25M NaCl. To hybridize, the oligonucleotides were denatured by boiling for 5 minutes and allowed to cool from 65°C to room temp over 2 hrs. Overlaps were extended using the Klenow fragment and digested with HindIII and BamHI. The resulting digested double stranded fragment was cloned into pUC19 and sequence confirmed. A clone containing the fragment of the correct sequence was digested with BamHI/HindIII to liberate the bikunin containing fragment with the following + strand sequence:

GAA GGG GTA AGC TTG GAT AAA AGA TAT GAA GAA TAC TGC ACC GCC
 AAC GCA GTC ACT GGG CCT TGC CGT GCA TCC TTC CCA CGC TGG TAC
 25 TTT GAC GTG GAG AGG AAC TCC TGC AAT AAC TTC ATC TAT GGA GGC
 TGC CGG GGC AAT AAG AAC AGC TAC CGC TCT GAG GAG GCC TGC ATG
 CTC CGC TGC TTC CGC CAG TAG GGA TCC (SEQ ID. : 44)

- which was then gel purified and ligated into BamHI/HindIII cut pS604. The ligation mixture was extracted into phenol/chloroform and purified over a S-200 minispin column. The ligation product was directed transformed into yeast strains SC101 and WHL341 and plated on ura selection plates. Twelve colonies from each strain were re-streaked on ura drop out plates. A single colony was inoculated into 2 ml of ura DO media and grown over night at 30°C. Cells were pelleted for 2 minutes at 14000x g and the supernatants evaluated for their content of placental bikunin (102-159).

- Detection of expression of placental bikunin (102-159) in transformed yeast*
 40 Firstly, the supernatants (50 ul per assay) were evaluated for their

WO 00/37099

PCT/GB99/04381

capacity to inhibit the *in vitro* activity of trypsin using the assay methods as described in Example 1 (1 ml assay volume). An un-used media only sample as well as a yeast clone expressing an inactive variant of aprotinin served as negative controls. A yeast clone expressing natural aprotinin served as a positive control and is shown for comparison.

The second method to quantify placental bikunin (102-159) expression exploited use of polyclonal antibodies (pAbs) against the synthetic peptide to monitor the accumulation of the recombinant peptide using Western blots. These studies were performed only with recombinants derived from strain SC101, since these produced greater inhibitory activity than recombinants derived from strain WHL341.

To produce the pAb, two 6-8 week old New Zealand White female rabbits (Hazelton Research Labs, Denver, Pa) were immunized on day zero with 250 ug of purified reduced synthetic placental bikunin (102-159), in Complete Freund's adjuvant, followed by boosts on days 14, 35 and 56 and 77 each with 125 ug of the same antigen in Incomplete Freund's adjuvant. Antiserum used in the present studies was collected after the third boost by established procedures. Polyclonal antibodies were purified from the antiserum over protein A.

Colonies 2.4 and 2.5 from transformation of yeast SC101 (Figure 8) as well as an aprotinin control were grown overnight in 50 ml of ura DO media at 30°C. Cells were pelleted and the supernatant concentrated 100-fold using a Centriprep 3 (Amicon, Beverly, MA) concentrator. Samples of each (30 µl) were subjected to SDS-PAGE on 10-20% tricine buffered gels (Novex, San Diego, CA) using the manufacturers procedures. Duplicate gels were either developed with a silver stain kit (Integrated Separation Systems, Nantick, MA) or transferred to nitrocellulose and developed with the purified polyclonal antibody elicited to synthetic bikunin (102-159). Alkaline-phosphatase conjugated goat anti-rabbit antibody was used as the secondary antibody according to the manufacturer's directions (Kirkegaard and Perry, Gaithersburg, MD).

Purification of placental bikunin (102-159) from a transformed strain of SC101

Fermentation broth from a 1L culture of SC101 strain 2.4 was harvested by centrifugation (4,000 g x 30 min.) then applied to a 1.0 ml column of anhydrochymotrypsin-sepharose (Takara Biochemical Inc., CA), that was previously equilibrated with 50 mM Hepes buffer pH 7.5 containing 0.1M NaCl, 2 mM CaCl₂ and 0.01% (v/v) triton X-100. The column was washed with the same buffer but containing 1.0 M NaCl until the A280nm declined to zero,

whereupon the column was eluted with 0.1M formic acid pH 2.5. Eluted fractions were pooled and applied to a C18 column (Vydac, 5um, 4.6 x 250 mm) previously equilibrated with 0.1% TFA, and eluted with a 50 min. linear gradient of 20 to 80% acetonitrile in 0.1% TFA. Fractions containing placental bikunin (102-159) were pooled and re-chromatographed on C18 employing elution with a linear 22.5 to 50% acetonitrile gradient in 0.1% TFA.

Results. Figure 8 shows the percent trypsin activity inhibited by twelve colonies derived from the transformation of each of strains SC101 and WHL341. The results show that all twelve colonies of yeast strain SC101 transformed with the trypsin inhibitor placental bikunin (102-159) had the ability to produce a substantial amount of trypsin inhibitory activity compared to the negative controls both of which showed no ability to inhibit trypsin. The activity is therefore related to the expression of a specific inhibitor in the placental bikunin variant (102-159) transformed cells. The yeast WHL341 samples contained minimal trypsin inhibitory activity. This may be correlated to the slow growth observed with this strain under the conditions employed.

Figure 9 shows the SDS-PAGE and western analysis of the yeast SC101 supernatants. Silver stained SDS-PAGE of supernatants derived from recombinant yeasts 2.4 and 2.5 expressing placental bikunin (102-159) as well as from the yeast expressing aprotinin yielded a protein band running at approximated 6 kDa, corresponding to the size expected for each recombinant Kunitz inhibitor domain. Western analysis showed that the 6 kDa bands expressed by stains 2.4. and 2.5 reacted with the pAb elicited to placental bikunin (102-159). The same 6 kDa band in the aprotinin control did not react with the same antibody, demonstrating the specificity of the antibody for the placental bikunin variant (102-159).

The final preparation of placental bikunin C-terminal domain was highly pure by silver-stained SDS-PAGE (Figure 10). The overall recovery of broth-derived trypsin inhibitory activity in the final preparation was 31%. N-terminal sequencing of the purified inhibitor indicated that 40% of the protein is correctly processed to yield the correct N-terminus for placental bikunin (102-159) while about 60 % of the material contained a portion of the yeast α -mating factor. The purified material comprised an active serine protease inhibitor exhibiting an apparent K_i of 0.35 nM for the *in vitro* inhibition of plasma kallikrein.

In conclusion, the accumulation both of a protease inhibitor activity and a protein immunochemically related to synthetic bikunin (102-159) in fermentation

WO 00/37099

PCT/GB99/04381

broth as well as the isolation of placental bikunin (102-159) from one of the transformed lines provided proof of expression of placental bikunin in the recombinant yeast strains described herein, showing for the first time the utility of yeasts for the production of placental bikunin fragments.

5 Additional constructs were prepared in an effort to augment the expression level of the Kunitz domain contained within placental bikunin 102-159, as well as to increase the yield of protein with the correct N-terminus. We hypothesized that the N-terminal residues of placental bikunin 102-159 (YEEY--) may have presented a cleavage site that is only poorly recognized by the yeast KEX-2 protease that
10 enzymically removes the yeast a-factor pro-region. Therefore, we prepared yeast expression constructs for the production of placental bikunin 103-159 (N-terminus of EEY...), 101-159 (N-terminus of NYEEY...) and 98-159 (DMFNYEEY..) in order to modify the P' subsites surrounding the KEX-2 cleavage site. To attempt to augment the levels of recombinant protein expression, we also used the yeast preferred codons
15 rather than mammalian preferred codons in preparing some of the constructs described below. The constructs were essentially prepared as described above for placental bikunin 102-159 (defined as construct #1) but with the following modifications:

20 Construct #2 placental bikunin 103-159, yeast codon usage
A 5' sense oligonucleotide

GAAGGGGTAA GCTTGGATAA AAGAGAAGAA TACTGTACTG CTAATGCTGT
TACTGGTCCA TGTAGAGCTT CTTTCCAAG ATGGTACTTT GATGTTGAAA GA
25 (SEQ ID NO.: 55)

and 3' antisense oligonucleotide

ACTGGATCCT CATTGGCGAA AACATCTCAA CATAAGGCT TCTTCAGATC
30 TGTAAGAATT TTTATTACCT CTACAACCAC CGTAAATAAA ATTATTACAA
GAATTTCTTT CAACATCAAA GTACCATCT (SEQ ID NO.: 56)

were manipulated as described for the production of an expression construct (construct #1 above) for the expression of placental bikunin 102-159
35

Construct #3 placental bikunin 101-159, yeast codon usage
A 5' sense oligonucleotide

WO 00/37099

PCT/GB99/04381

GAAGGGGTAA GCTTGGATAA AAGAAATTAC GAAGAATACT GTACTGCTAA
TGCTGTTACT GGTCATGTA GAGCTTCTTT TCCAAGATGG TACTTTGATG
TTGAAAGA (SEQ ID NO.: 57)

5

and the same 3' antisense oligonucleotide as used for construct #2, were manipulated as described for the production of an expression construct (construct #1 above) for the expression of placental bikunin 102-159.

10 Construct #4 placental bikunin 98-159, yeast codon usage
A 5' sense oligonucleotide

GAAGGGGTAA GCTTGGATAA AAGAGATATG TTTAATTACG AAGAATACTG
TACTGCTAAT GCTGTTACTG GTCCATGTAG AGCTTCTTTT CCAAGATGGT
15 ACTTTGATGT TGAAAGA (SEQ ID NO.: 58)

and the same 3' antisense oligonucleotide as used for construct #2, were manipulated as described for the production of an expression construct (construct #1 above).

20 Yeast strain SC101 (MAT α , ura 3-52, suc 2) was transformed with the plasmids containing each of the above cDNAs, and proteins were expressed using the methods that were described above for the production of placental bikunin 102-159 with human codon usage. Approximately 250 ml of each yeast culture was harvested, and the supernatant from centrifugation (15 min x 3000 RPM) separately subjected to
25 purification over 1 ml columns of kallikrein-sepharose as described above. The relative amount of trypsin inhibitory activity in the aplysate, the amount of purified protein recovered and the N-terminal sequence of the purified protein were determined and are listed below in Table 7.

30 Table 7 Relative production levels of different proteins containing the C-terminal Kunitz domain of placental bikunin

TABLE 7					
Construct	Relative conc. of inhibitor in aplysate	N-terminal sequencing Amount (pmol)	sequence	Comments	
#2 103-159	none detected	none	none	no expression	
#3 101-159	25 % inhibition	none	none	low expression	

40

WO 00/37099

PCT/GB99/04381

5	#4 98-159 product	93 % inhibition	910	DMFNYE-	good expression correct
10	#1 102-159	82 % inhibition	480	AKEEGV-	expression of active incorrectly processed protein

The results show that placental bikunin fragments of different lengths that contain the C-terminal Kunitz domain show wide variation in capacity to express functional secreted protein. Constructs expressing fragments 101-159 and 103-159 yielded little or low enzymic activity in the supernatants prior to purification, and N-terminal sequencing of 0.05 ml aliquots of each purified fraction yielded undetectable amounts of inhibitor. On the other hand expression either of placental bikunin 102-159 or 98-159 yielded significant amounts of protease activity prior to purification. N-terminal sequencing however showed that the purified protein recovered from expression of 102-159 was once again largely incorrectly processed, exhibiting an N-terminus consistent with processing of the majority of the pre-protein at a site within the yeast α -mating factor pro-sequence. The purified protein recovered from expression of placental bikunin 98-159 however was processed entirely at the correct site to yield the correct N-terminus. Furthermore, nearly twice as much protein was recovered as compared to the recovery of placental bikunin 102-159. Placental bikunin 98-159 thus represents a preferred fragment length for the production of the C-terminal Kunitz domain of placental bikunin by the α -mating factor pre-pro sequence/KEX-2 processing system of *S. cerevisiae*,

Example 6

Alternative procedure for yeast expression

The 58 amino acid peptide derived from the R74593 translation product can also be PCR amplified from either the R87894-R74593 PCR product cloned into the TA vectorTM (Invitrogen, San Diego, CA) after DNA sequencing or from human placental cDNA. The amplified DNA product will consist of 19 nucleotides from the yeast α -mating factor leader sequence mated to the R74593 sequence which codes for the YEEY--CFRQ (58 residues) so as to make the translation product in frame, constructing an α -mating factor/Kunitz domain fusion protein. The protein sequence also contains a kex 2 cleavage which will

WO 00/37099

PCT/GB99/04381

liberate the Kunitz domain at its native N-terminus.

The 5' sense oligonucleotide which contains a HindIII site for cloning will contain the following sequence:

5

GCCAAGCTTG GATAAAAGAT ATGAAGAAT ACTGCACCGC CAACGCA
(SEQ ID NO.: 30)

The 3' antisense oligonucleotide contains a BamHI site for cloning as well as a stop codon and is of the following sequence:

10

GGGGATCCTC ACTGCTGGCG GAAGCAGCGG AGCAT (SEQ ID NO.: 31)

The full 206 nucleotide cDNA sequence to be cloned into the yeast expression vector is of the following sequence:

15

CCAAGCTTGG ATAAAAGATA TGAAGAATAC TGCACCGCCA ACGCAGTCAC
TGGGCCTTGC CGTGCATCCT TCCCACGCTG GTACTTTGAC GTGGAGAGGA
ACTCCTGCAA TAACTTCATC TATGGAGGCT GCCGGGGCAA TAAGAACAGC
20 TACCGCTCTG AGGAGGCCTG CATGCTCCGC TGCTTCCGCC AGCAGTGAGG
ATCCCC (SEQ ID NO.: 32)

After PCR amplification, this DNA will be digested with HindIII, BamHI and cloned into the yeast expression vector pMT15 (see US patent 5,164,482, incorporated by reference in the entirety) also digested with HindIII and BamHI. The resulting plasmid vector is used to transform yeast strain SC 106 using the methods described in US patent 5,164,482. The URA 3+ yeast transformants are isolated and cultivated under inducing conditions. The yield of recombinant Placental bikunin variants is determined according to the amount of trypsin inhibitory activity that accumulated in the culture supernatants over time using the *in vitro* assay method described above. Fermentation broths are centrifuged at 9000 rpm for 30 minutes. The supernatant is then filtered through a 0.4 then a 0.2 µm filter, diluted to a conductivity of 7.5 ms, and adjusted to pH 3 with citric acid. The sample is then batch absorbed onto 200 ml of S-sepharose fast flow (Pharmacia) in 50 mM sodium citrate pH 3 and stirred for 60 min. The gel is subsequently washed sequentially with 2 L of each of: 50 mM sodium citrate pH 3.0; 50 mM Tris-HCL pH 9.0; 20 mM HEPES pH 6.0. The washed gel is transferred into a suitable column and eluted with a linear gradient of 0 to 1 M

30

35

sodium chloride in 20 mM HEPES pH 6.0. Eluted fractions containing *in vitro* trypsin inhibitory activity are then pooled and further purified either by a) chromatography over a column of immobilized anhydrotrypsin (essentially as described in Example 2); b) by chromatography over a column of immobilized bovine kallikrein; or c) a combination of conventional chromatographic steps including gel filtration and /or anion-exchange chromatography.

Example 7

Isolation and characterization of native human placental bikunin from placenta

Bikunin protein was purified to apparent homogeneity from whole frozen placenta (Analytical Biological Services, Inc, Wilmington, DE). The placenta (740 gm) was thawed to room temperature and cut into 0.5 to 1.0 cm pieces, placed on ice and washed with 600 ml PBS buffer. The wash was decanted and 240 ml of placenta pieces placed into a Waring blender. After adding 300 ml of buffer consisting of 0.1 M Tris (pH 8.0), and 0.1 M NaCl, the mixture was blended on high speed for 2 min, decanted into 750.0 ml centrifuge tubes, and placed on ice. This procedure was repeated until all material was processed. The combined slurry was centrifuged at 4500 x g for 60 minutes at 4°C. The supernatant was filtered through cheese cloth and the placental bikunin purified using a kallikrein affinity column made by covalently attaching 70 mg of bovine pancreatic kallikrein (Bayer AG) to 5.0 mls of CNBr activated Sepharose (Pharmacia) according to manufacturers instruction. The material was loaded onto the affinity column at a flow rate of 2.0 ml/min and washed with 0.1 M Tris (pH 8.0), 0.1 M NaCl until absorbance at 280 nm of the wash could no longer be detected. The column was further washed with 0.1 M Tris (pH 8.0), 0.5 M NaCl and then eluted with 3 volumes of 0.2 M acetic acid, pH 4.0. Fractions containing kallikrein and trypsin inhibitory (see below) activity were pooled, frozen, and lyophilized. Placental bikunin was further purified by gel-filtration chromatography using a Superdex 75 10/30 (Pharmacia) column attached to a Beckman System Gold HPLC system. Briefly, the column was equilibrated in 0.1 M Tris, 0.15 M NaCl, and 0.1% Triton X-100 at a flow rate of 0.5 ml/min. The lyophilized sample was reconstituted in 1.0 ml of 0.1 M Tris, pH 8.0 and injected onto the gel-filtration column in 200 µl aliquots. Fractions were collected (0.5 ml) and assayed for trypsin and kallikrein inhibitory activity. Active fractions were pooled, and the pH of the solution adjusted to 2.5 by addition of TFA. The material was directly applied to a Vydac C18 reverse-phase column (5 micron, 0.46 x 25 cm) which had been equilibrated in 20% acetonitrile in 0.1 %TFA.

Separation was achieved using a linear gradient of 20 to 80% acetonitrile in 0.1% TFA at 1.0 ml/min over 50 minutes after an initial 20 minute wash at 20% acetonitrile in 0.1% TFA. Fractions (1ml) were collected and assayed for trypsin and kallikrein inhibitory activity. Fractions containing inhibitory activity were
 5 concentrated using a speed-vac concentrator (Savant) and subjected to N-terminal sequence analysis.

Functional assays for Placental Bikunin:

Identification of functional placental bikunin was achieved by measuring
 10 its ability to inhibit bovine trypsin and human plasma kallikrein. Trypsin inhibitory activity was performed in assay buffer (50 mM Hepes, pH 7.5, 0.1 M NaCl, 2.0 mM CaCl₂, 0.1% Triton x-100) at room temperature in a 96-well microtiter plate (Perkin Elmer) using Gly-Pro-Lys-Aminomethylcoumarin as a substrate. The amount of coumarin produced by trypsin was determined by
 15 measuring the fluorescence (ex = 370 nm, em = 432 nm) on a Perkin-Elmer LS-50B fluorimeter equipped with a plate reader. Trypsin (23 µg in 100 µl buffer) was mixed with 20 µl of the sample to be tested and incubated for 10 minutes at 25°C. The reaction was started by the addition of 50 µl of the substrate GPK-AMC (33 µM final) in assay buffer. The fluorescence intensity was measured and
 20 the % inhibition for each fraction was determined by:

$$\% \text{ inhibition} = 100 \times [1 - F_0/F_1]$$

where F_0 is the fluorescence of the unknown and F_1 is the fluorescence of the
 25 trypsin only control. Kallikrein inhibitory activity of the fractions was similarly measured using 7.0 nM kallikrein in assay buffer (50 mM Tris, pH 8.0, 50 mM NaCl, 0.1% triton x-100) and 66.0 µM Pro-Phe-Arg-AMC as a substrate.

Determination of the in vitro specificity of placental bikunin

30 The *In vitro* specificity of native human placental bikunin was determined using the materials and methods as described in the preceding examples above. Placental bikunin was quantified by active site titration against a known concentration of trypsin using GPK-AMC as a substrate to monitor the fraction of unbound trypsin.

35

Protein Sequencing

The 1 ml fraction (C18-29 Delaria) was reduced to 300 µl in volume, on a

WO 00/37099

PCT/GB99/04381

Speed Vac, to reduce the amount of organic solvent. The sample was then loaded onto a Hewlett-Packard miniature biphasic reaction column, and washed with 1 ml of 2% trifluoroacetic acid. The sample was sequenced on a Hewlett-Packard Model G1005A protein sequencing system using Edman degradation. Version 3.0 sequencing methods and all reagents were supplied by Hewlett-Packard. Sequence was confirmed for 50 cycles.

Results. Placental Bikunin was purified to apparent homogeneity by sequential kallikrein affinity, gel-filtration, and reverse-phase chromatography (see purification table below):

Table 5 Purification table for native Placental Bikunin (1-179)

TABLE 5					
Step	Vol (ml)	OD 280 (/ml)	OD 280	Units ^a (U)	Units/OD 280
Placenta Supernatant	1800.0	41.7	75,060	3,000,000	40.0
Kallikrein Affinity pH 4.0	20.0	0.17	3.36	16,000	4,880
Kallikrein Affinity pH 1.7	10.2	0.45	4.56	12,000	2,630
Superdex 75	15.0	0.0085	0.13	3,191	24,546

^aOne Unit is defined as that amount which inhibits 50% of trypsin activity in a standard assay.

The majority of the kallikrein and trypsin inhibitory activity eluted from the kallikrein affinity column in the pH 4.0 elution. Subsequent gel-filtration chromatography (Figure 5) yielded a peak of kallikrein and trypsin inhibitory activity with a molecular weight range of 10 to 40 kDa as judged by a standard curve generated by running molecular weight standards under identical conditions. Reverse-phase C18 chromatography (Figure 6) yielded 4 peaks of inhibitory activity with the most potent eluting at approximately 30 % acetonitrile. The activity associated with the first peak to elute from C18 (fraction 29) exhibited an amino acid sequence starting with amino acid 1 of the predicted amino acid sequence of placental bikunin (ADRER...; SEQ ID NO.: 1), and was identical to the predicted sequence for 50 cycles of sequencing (underlined amino acids in Figure 3). Cysteine residues within this sequence stretch were silent as expected for sequencing of oxidized protein. The cysteine residues at amino acid positions 11 and 20 of mature placental bikunin were later identified from sequencing of the S-pyridylethylated protein whereupon PTH-pyridylethyl-

cysteine was recovered at cycles 11 and 20.

Interestingly, the asparagine at amino acid residue number 30 of the sequence (Figure 3) was silent showing that this site is likely to be glycosylated. Fraction 29 yielded one major sequence corresponding to that of placental bikunin starting at residue #1 (27 pmol at cycle 1) plus a minor sequence (2 pmol) also derived from placental bikunin starting at residue 6 (SIHD...). This shows that the final preparation sequenced in fraction 29 is highly pure, and most likely responsible for the protease inhibitory activity associated with this fraction (Figure 6).

Accordingly, the final preparation of placental bikunin from C18 chromatography was highly pure based on a silver-stained SDS-PAGE analysis (Figure 7), where the protein migrated with an apparent Mr of 24 kDa on a 10 to 20 % acrylamide tricine gel (Novex, San Diego, CA) calibrated with the following molecular weight markers: insulin (2.9 kDa); bovine trypsin inhibitor (5.8 kDa); lysozyme (14.7 kDa); β -lactaglobulin (18.4 kDa); carbonic anhydrase (29 kDa); and ovalbumin (43 kDa). The above size of placental bikunin on SDS-PAGE is consistent with that predicted from the full length coding sequence (Figure 4F).

As expected based on the N-terminal sequencing results described above, the purified protein reacted with an antibody elicited to placental bikunin (7-64) to yield a band with the same Mr (Figure 12A) as observed for the purified preparation detected on gels by silver stain (Figure 7). However, when the same preparation was reacted with an antibody elicited to synthetic placental bikunin (102-159), a band corresponding to the full length protein was not observed. Rather, a fragment that co-migrated with synthetic bikunin (102-159) of approximately 6 kDa was observed. The simplest interpretation of these results is that the purified preparation had undergone degradation subsequent to purification to yield an N-terminal fragment comprising the N-terminal domain and a C-terminal fragment comprising the C-terminal domain. Assuming that the fragment reactive against antiserum to placental bikunin (7-64) is devoid of the C-terminal end of the full length protein, the size (24 kDa) would suggest a high state of glycosylation.

Table 6. below shows the potency of *in vitro* inhibition of various serine proteases by placental bikunin. Data are compared with that obtained with aprotinin (Trasyol®).

Table 6 Ki values for the inhibition of various proteases by placental bikunin

TABLE 6

WO 00/37099

PCT/GB99/04381

Protease (concentration)	Placental Bikunin Ki (nM)	Aprotinin Ki (nM)
Trypsin (48.5 pM)	0.13	0.8
Human Plasmin (50 pM)	1.9	1.3

The results show that placental bikunin isolated from a natural source (human placenta) is a potent inhibitor of trypsin-like serine proteases.

5 Example 8

Expression pattern of placental bikunin amongst different human organs and tissues

A multiple tissue northern was purchased from Clontech which contained 2 µg of polyA⁺ RNA from human heart, brain, placenta, lung, liver, skeletal muscle, kidney, and pancreas. Two different cDNA probes were used: 1) a gel purified cDNA encoding placental bikunin (102-159); 2) the 780 base pair PCR-derived cDNA (Figure 4E) liberated from a TA clone by digestion with EcoRI and gel purified. Each probe was labeled using ³²P-dCTP and a random priming labeling kit from Boehringer Mannheim Biochemicals (Indiana), then used to hybridize to the multiple tissue northern according to the manufacturers specifications. Autoradiographs were generated using Biomax^{*} film with an 18 hr exposure time, and developed using a Umax^{*} Scanner and scanned using Adobe Photoshop^{*}.

Results. The pattern of tissue expression observed using a placental bikunin (102-159) probe (Figure 11A) or a larger probe containing both Kunitz domains of placental bikunin (Figure 11B) was essentially the same as might be expected. The placental bikunin mRNA was most abundant in pancreas and placenta. Significant levels were also observed in lung, brain and kidney, while lower levels were observed in heart and liver, and the mRNA was undetectable in skeletal muscle. The transcript size was 1.95 kilobases in all cases, in close agreement with the predicted size of placental bikunin deduced both from EST overlay and cloning of full length cDNA described in preceding sections.

The broad tissue distribution of the mRNA shows that placental bikunin is broadly expressed. Since the protein also contains a leader sequence it would have ample exposure to the human immune system, requiring that it become recognized as a self protein. Additional evidence for a broad tissue distribution of placental bikunin mRNA expression was derived from the fact that some of

WO 00/37099

PCT/GB99/04381

the EST entries with homology to placental bikunin (Figure 4B) were derived from human adult and infant brain, and human retina, breast, ovary, olfactory epithelium, and placenta. It is concluded therefore that administration of the native human protein to human patients would be unlikely to elicit an immune response.

Interestingly, the expression pattern of placental bikunin is somewhat reminiscent of that for bovine aprotinin which is found in high levels in bovine lung and pancreas. To further elucidate the expression pattern of placental bikunin, RT-PCR of total RNA from the following human cells was determined: un-stimulated human umbilical vein endothelial cells (HUVECs), HK-2 (line derived from kidney proximal tubule), TF-1 (erythroleukemia line) and phorbol ester (PMA)-stimulated human peripheral blood leukocytes. The probes used:

CACCTGATCGCGAGACCCC (sense; SEQ ID NO.: 59);
CTGGCGGAAGCAGCGGAGCATGC (antisense; SEQ ID NO.: 60),

were designed to amplify a 600 b.p placental bikunin encoding cDNA fragment. Comparisons were normalized by inclusion of actin primers to amplify an 800 b.p. actin fragment. Whereas the 800 b.p fragment identified on agarose gels with ethidium bromide was of equal intensity in all lanes, the 600 b.p. placental bikunin fragment was absent from the HUVECs but present in significant amounts in each of the other cell lines. We conclude that placental bikunin is not expressed in at least some endothelial cells but is expressed in some leukocyte populations.

Example 9

Purification and properties of Placental Bikunin (1-170) highly purified from a Baculovirus / Sf9 expression system

A large fragment of Placental bikunin containing both Kunitz domains (Bikunin (1-170) (SEQ ID NO:52) was expressed in Sf9 cells as follows. Placental bikunin cDNA obtained by PCR (Figure 4E) and contained within a TA vector (see previous Examples) was liberated by digestion with HindIII and XbaI yielding a fragment flanked by a 5' XbaI site and 3' HindIII site. This fragment was gel purified and then cloned into the M13mp19 vector (New England Biolabs, Beverly, MA). In vitro mutagenesis (Kunkel T.A., (1985) Proc. Natl. Acad. Sci. USA, 82: 488-492) was used to generate a PstI site 3' to the XbaI site at the 5' end, but 5' to the sequence

WO 00/37099

PCT/GB99/04381

encoding the ATG start site, natural placental bikunin signal peptide and mature placental bikunin coding sequence. The oligonucleotide used for the mutagenesis had the sequence:

5 5' CGC GTC TCG GCT GAC CTG GCC CTG CAG ATG GCG CAC GTG TGC GGG 3'
(SEQ ID NO.: 61)

A stop codon (TAG) and BglII / XmaI site was similarly engineered at the 3' end of the cDNA using the oligonucleotide:

10

5' CTG CCC CTT GGC TCA AAG TAG GAA GAT CTT CCC CCC GGG GGG GTG
GTT CTG GCG GGG CTG 3' (SEQ ID NO.: 62).

15

The stop codon was in frame with the sequence encoding placental bikunin and caused termination immediately following the Lysine at amino acid residue 170, thus encoding a truncated placental bikunin fragment devoid of the putative transmembrane domain. The product from digestion with PstI and BglII was isolated and cloned into the BacPac8 vector for expression of Placental bikunin fragment (1-170) which contains both Kunitz domains but which is truncated immediately N-terminal to the putative transmembrane segment.

20

25

30

The expression of Bikunin by Sf-9 insect cells was optimal at a multiplicity of infection of 1 to 1 when the medium was harvested at 72 h post infection. After harvesting, the baculovirus cell culture supernatant (2L) was adjusted to pH 8.0 by the addition of Tris-HCl. Bikunin was purified by chromatography using a 5 ml bovine pancreatic kallikrein affinity column as previously described in Example 7 for the purification of native placental bikunin from placenta. Eluted material was adjusted to pH 2.5 with TFA and subjected to chromatography on a C18 reverse-phase column (1.0 x 25 cm) equilibrated in 10% acetonitrile in 0.1% TFA at a flow rate of 1 ml/min. The bikunin was eluted with a linear gradient of 10 to 80% acetonitrile in 0.1% TFA over 40 min. Active fractions were pooled, lyophilized, redissolved in 50 mM Hepes (pH 7.5), 0.1 M NaCl, 2 mM CaCl₂, and 0.1% triton x-100, and stored at -20°C until needed. The concentration of recombinant bikunin was determined by amino acid analysis.

35

Results. Recombinant bikunin was purified from baculovirus cell culture supernatant using a 2-step purification protocol as shown below, to yield an active trypsin inhibitor (Table 8 below).

WO 00/37099

PCT/GB99/04381

Table 8 Purification of recombinant bikunin from transformed culture supernatant

TABLE 8

Purification Step	Vol (ml)	OD 280/ml	OD 280 total	Units (U)	Specific activity (U/OD)
Supernatant	2300.0	9.0	20,700	6,150,000	297
Kallikrein affinity	23.0	0.12	2.76	40,700	14,746
C18 reverse-phase	0.4	3.84	1.54	11,111	72,150

Chromatography of the crude material over an immobilized bovine pancreatic kallikrein affinity column selectively isolated 0.013 % of the protein and 0.67 % of the trypsin inhibitory activity present. The majority of the trypsin inhibitory activity present in the starting supernatant did not bind to the immobilized kallikrein and is not related to bikunin (results not shown). Subsequent chromatography using C18 reverse-phase yielded a further purification of 5-fold, with a recovery of 0.2%. The final preparation was highly pure by SDS-PAGE (Figure 13), exhibiting an Mr of 21.3 kDa, and reacted on immunoblots to rabbit anti-placental bikunin 102-159 (not shown). N-terminal sequencing (26 cycles) yielded the expected sequence for mature placental bikunin (Figure 4F) starting at residue +1(ADRER....), showing that the signal peptide was correctly processed in Sf9 cells.

Purified placental bikunin from Sf9 cells (100 pmol) was pyridylethyl-alkylated, CNBr digested and then sequenced without resolution of the resulting fragments. Sequencing for 20 cycles yielded the following N-terminii:

Sequence	Amount	Placental bikunin residue #
LRCFrQQENPP-PLG-----	21 pmol	154 - 168 (SEQ ID NO.: 63)
ADRERSIHDFCLVSKVVGRC	20 pmol	1 - 20 (SEQ ID NO.: 64)
FNYeEYCTANAVTGPCRASf	16 pmol	100 - 119 (SEQ ID NO.: 65)
Pr--Y-V-dGS-Q-F-Y-G	6 pmol	25 - 43 (SEQ ID NO.: 66)

Thus N-terminii corresponding to each of the expected four fragments were recovered. This confirms that the Sf9 expressed protein contained the entire ectodomain sequence of placental bikunin (1-170).

Example 10

Inhibition specificity of purified placental bikunin (1-170) derived from Sf9 cells.

The *in vitro* specificity of recombinant bikunin was determined using the

WO 00/37099

PCT/GB99/04381

materials and methods as described in Examples 3, 4 and 7. In addition, the inhibition of human tissue kallikrein by bikunin was measured by the incubation of 0.35 nM human tissue kallikrein recombinant bikunin in buffer containing 50 mM Tris (pH 9.0), 50 mM NaCl, and 0.01% triton x-100. After 5 min. at 37°C, 5 µl of 2 mM PFR-
5 AMC was added and the change in fluorescence monitored.

Inhibition of tissue plasminogen activator (tPA) was also determined as follows: tPA (single chain form from human melanoma cell culture from Sigma Chemical Co, St Louis, MO) was pre-incubated with inhibitor for 2 hr at room temperature in 20 mM Tris buffer pH 7.2 containing 150 mM NaCl, and 0.02% sodium
10 azide. Reactions were subsequently initiated by transfer to a reaction system comprising the following initial component concentrations: tPA (7.5 nM), inhibitor 0 to 6.6 µM, Dlle-Lpro-Larg-pNitroaniline (1mM) in 28 mM Tris buffer pH 8.5 containing 0.004 % (v/v) triton x-100 and 0.005% (v/v) sodium azide. Formation of p-Nitroaniline was determined from the A405nm measured following incubation at 37
15 C for 2hr.

The table below show the efficacy of recombinant bikunin as an inhibitor of various serine proteases *in vitro*. Data is shown compared against data obtained for screening inhibition using either recombinant bikunin, or aprotinin.

5 Table 9 Comparisons of Ki values for the inhibition of various proteases by recombinant placental bikunin (1-170) or aprotinin

TABLE 9		
Protease (concentration)	Recombinant Bikunin Ki (nM)	Aprotinin Ki (nM)
Trypsin (48.5 pM)	0.064	0.8
Human Plasma Kallikrein (2.5 nM)	0.18	19.0
Human Tissue Kallikrein (0.35 nM)	0.04	0.004
Bovine Pancreatic Kallikrein (100 pM)	0.12	0.02
Human Plasmin (50 pM)	0.23	1.3
factor Xa (0.87 nM)	180	5% Inhibition at 31 μ M
factor XIa (0.1 nM)	3.0	288
tissue plasminogen activator (7.5 nM)	< 60	no inhibition at 6.6 μ M
Tissue Factor VIIa	800	no inhibition at 1 μ M

10 The results show that recombinant bikunin can be expressed in insect cells to yield an active protease inhibitor that is effective against at least five different serine protease inhibitors. Recombinant bikunin was more potent than aprotinin against human plasma kallikrein, trypsin and plasmin. Surprisingly, the recombinant bikunin was more potent than the synthetically derived bikunin fragments (7-64) and (102-159) against all enzymes tested. These data show that recombinant bikunin is more
15 effective than aprotinin, using *in vitro* assays, and that one would expect better *in vivo* potency.

Besides measuring the potencies against specific proteases, the capacity of placental bikunin (1-170) to prolong the activated partial thromboplastin time (APTT) was evaluated and compared with the activity associated with aprotinin. Inhibitor
20 was diluted in 20 mM Tris buffer pH 7.2 containing 150 mM NaCl and 0.02% sodium azide and added (0.1 ml) to a cuvette contained within an MLA Electra^R 800 Automatic Coagulation Timer coagulometer (Medical Laboratory Automation, Inc., Pleasantville, N.Y.). The instrument was set to APTT mode with a 300 sec. activation time and the duplicate mode. Following addition of 0.1 ml of plasma (Specialty
25 Assayed Reference Plasma lot 1-6-5185, Helena Laboratories, Beaumont, TX), the APTT reagent (Automated APTT-lot 102345, from Organon Teknika Corp., Durhan,

NC) and 25 mM CaCl₂ were automatically dispensed to initiate clotting, and the clotting time was monitored automatically. The results (Figure 14) showed that a doubling of the clotting time required approximately 2 μ M final aprotinin, but only 0.3 μ M Sf9 derived placental bikunin. These data show that placental bikunin is an effective anticoagulant, and usefull as a medicament for diseases involving pathologic activation of the intrinsic pathway of coagulation.

Example 11

Measurement of Tracheal Potential Difference in the Guinea-pig

The aim of this study was to investigate the effect of the Kunitz serine protease inhibitor Bikunin, and the sodium channel blocker amiloride on guinea-pig tracheal potential difference 3 hours post treatment. These agents were delivered into the cephalad trachea by topical instillation. TPD was monitored 2 hours later for 60 minutes. The procedure used in this Example is described in Newton et al. in "Cilia, Mucus and Mucociliary Interactions," Ed., Baum, G.L. et al., Marcel Dekker, New York, 1998; Newton et al., Ped. Pulm. S17, Abs. 364, 1998).

Materials and methods/Reagents used

Aqueous formulations of Bikunin (1-170)(5 and 50 μ g/mL (SEQ ID NO: 52)) (as described in Example 17 below) and amiloride (obtained from Sigma Chemicals, St. Louis, MO, USA)(100 μ M) were prepared, sterile filtered and endotoxin tested prior to use. These formulations were prepared in Hank's Balanced salt solution (HBSS) and contained 137 mM NaCl, 3 mM KCl, 3 mM KH₂PO₄, 8 mM Na₂HPO₄, 0.2% Tween-80, pH 7.1) was prepared, sterile filtered and endotoxin tested for use in this example. HBSS was used as a control solution. Hypnorm® (Fentanyl citrate 0.315 mg/mL and Fluanisone 10 mg/mL) was obtained from Janssen Animal Health and Hypnovel® (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea pigs (550-750 g) were supplied by David Hall, UK. Thermistor probes were obtained from Kane-May Ltd, UK.

Induction of anaesthesia and administration of Bikunin into tracheal airway

Animals were anaesthetised using halothane. Once a satisfactory level of anaesthesia was induced a small incision was made below the lower jaw. The trachea was exposed and 100 μ l volume of vehicle, bikunin (0.5 μ g or 5 μ g) or amiloride (100 μ M) was instilled onto the tracheal surface using a needle and syringe. Once injected, the skin incision was sealed using Vetbond® (cyanocacrylate tissue glue). The animals were then allowed to recover.

Preparation of guinea-pig for measurement of tracheal potential difference

Two hours following agent treatment, guinea-pigs were anaesthetised for a second time with Hypnom® and Hypnovel® and immobilised in a supine position.

5 Rectal temperature, measured with a thermistor probe was maintained at 37° C by manual adjustment of a heat lamp. A ventral midline incision was made from the lower jaw to the clavicles. Using blunt dissection a length of trachea was exposed and bisected at the upper edge of the sternum. The external jugular vein was exposed and cannulated. The caudal part of the trachea was then cannulated to allow the animal

10 to spontaneously breath room air. The animal was then placed supine and its body temperature maintained using the heat lamp. 20 min. following induction of i.m. anaesthesia the tracheal agar electrode was inserted into the cephalad trachea and tracheal potential difference was measured for 60 minutes. The reference electrode was placed under cephalad trachea in contact with the trachea cartilage. The wound

15 site was covered to prevent drying.

Results

As shown in Figure 15, Bikunin (5 ug) inhibited the potential difference in guinea pig trachea in vivo following three hours of treatment relative to vehicle. The

20 effect of Amiloride (100 uM) and Bikunin (0.5 ug) is shown for comparison.

Example 12**The Effect of Bikunin on Tracheal Mucus Velocity in the Guinea-pig**

The aim of this study was to investigate the effect of the Kunitz family serine protease inhibitor Bikunin on guinea-pig tracheal mucus velocity 1.5 hours post

25 treatment. This agent was delivered into the cephalad trachea by topical instillation. TMV was monitored 1.5 hours later for 60 mins. The procedure used in this Example is described in Newton et al. in "Cilia, Mucus and Mucociliary Interactions," Ed., Baum, G.L. et al., Marcel Dekker, New York, 1998; Newton et al., Ped. Pulm. S17, Abs.

30 364, 1998).

Materials and methods/Reagents used:

A Bikunin (1-170) formulation (50 ug/mL Bikunin (SEQ ID NO: 52) (as described in Example 17 below) was prepared in HBBS containing 137 mM NaCl, 3

35 mM KCl, 3 mM KH₂PO₄, 8 mM Na₂HPO₄, 0.2% Tween-80, pH 7.1). The formulation was sterile filtered and endotoxin tested prior to use in this example. HBSS was used as a control solution. Hypnorm® (Fentanyl citrate 0.315 mg/mL and Fluanisone 10

WO 00/37099

PCT/GB99/04381

mg/mL) was obtained from Janssen Animal Health and Hypnovel® (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea pigs (550-750 g) were supplied by David Hall, UK. Thermistor probes were obtained from Kane-May Ltd, UK.

5 *Induction of anaesthesia and administration of Bikunin into tracheal airway*

Animals were anaesthetized using halothane. Once a satisfactory level of anaesthesia was induced, a small incision was made below the lower jaw. The trachea was exposed and 100 ul volume of vehicle or bikunin (5 ug) was instilled onto the tracheal surface using a needle and syringe. Once instilled, the skin incision was
10 sealed using Vetbond® (cyanoacrylate tissue glue). The animals were then allowed to recover.

Measurement of tracheal mucus velocity (TMV)

TMV was monitored using a lead collimated miniature Beta particle detector
15 probe arranged to detect the radioactivity emitted from an injected aliquot of ³²P-labelled *Saccharomyces cerevisiae* as it was transported on the tracheal mucociliary layer of an anaesthetized guinea pig (Newton and Hall 1998) Figure 16(a) illustrates the arrangement of the syringe and beta probe. Figure 16(b) illustrates the counts detected by the probe as the ³²P-labelled *S.cerevisiae* is transported along the tracheal
20 mucociliary layer.

70 minutes following instillation of bikunin, each animal was anaesthetized for a second time using Hyponorm® and Hypnovel® and immobilized in a supine position. The first TMV measurement was made 20 minutes afterwards. Subsequent measurements were taken every 15 minutes. The procedure for TMV measurements
25 is described, in detail, in Newton et al., "Cilia. Mucus and Mucociliary Interactions." Ed. Baum, G.L., Preil, Z., Roth, Y., Liron., Ostfield, E., Marcel Dekker. New York, 1990 and Newton et al. in *Pediatric Pulmonology* S17, Abs 364, 1998.

Results

30 As shown in Figure 16(c), Bikunin (5 ug) increased TMV in vivo in guinea pig, relative to saline, over a sustained period of 1.5 to 2.5 hours following administration.

Example 13

**Bikunin decreases sodium current in cultured human bronchial epithelial (HBE)
35 cell short circuit current (I_{sc}) in vitro.**

Tertiary HBE cell monolayers grown to confluence were mounted in modified Ussing chambers, immersed in Krebs buffer (KBR) solution and bubbled with

WO 00/37099

PCT/GB99/04381

95%O₂/5%CO₂ warmed to 37C.

Cells were left to equilibrate for 20 minutes before calibrating for background noise and fluid resistance. Transepithelial potential difference was then clamped to 0 mV using a WPI EVC 4000 voltage clamp. Ag/AgCl electrodes were used to monitor
 5 Isc. Once a stable baseline was achieved (typically 10-20 min), cells were treated with amiloride (10 uM). Once a response to amiloride was seen, it was washed out with KBR solution. After return to baseline and equilibration, Bikunin (1-170) (as described in Example 17 below) (0.5-50 ug/mL in PBS) or PBS control was added. 90 minutes following agent treatment, amiloride (10 uM) was added. Once the current
 10 was stable forskolin (10 uM) and then bumetanide (100 uM) was added.

Results

As shown in Figure 17, Bukinin (70 nM) inhibited sodium current in vitro in human bronchial epithelial cells over a 90 minute period. Forskolin induced cAMP-
 15 mediated chloride secretion and monolayer resistance was unaffected.

Example 14

The effect of hypertonic saline (14.4%) on TMV in the guinea pig

The aim of this comparative study was to investigate the effect of hypertonic saline (14.4% x 5 min) on guinea-pig tracheal mucus velocity. This agent was
 20 delivered into the cephalad trachea by aerosol. TMV was monitored immediately and every 15 minutes for 30 minutes. The procedure used in this Example is described in Newton et al. in "Cilia, Mucus and Mucociliary Interactions," Ed., Baum, G.L. et al., Marcel Dekker, New York, 1998; Newton et al., Ped. Pulm. S17, Abs. 364, 1998).

25

Materials and methods/Reagents used

Hypnorm® (Fentanyl citrate 0.315 mg/mL and Fluanisone 10 mg/mL) was obtained from Janssen Animal Health and Hypnovel® (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea pigs (550-750 g) were supplied by
 30 Harlan UK Ltd. Thermistor probes were obtained from Kane-May Ltd, UK.

Measurement of tracheal mucus velocity:

Animals were anaesthetized using Hypnorm® and Hypnovel®. TMV was monitored using a lead collimated miniature Beta particle detector probe arranged to
 35 detect the radioactivity emitted from an injected aliquot of ³²P-labelled *Saccharomyces cerevisiae* as it was transported on the tracheal mucociliary layer of an anaesthetized guinea pig (Newton and Hall 1998)

The first TMV measurement (run 1) was made 20 minutes after administration. Subsequent measurements were taken every 15 minutes. At a time point 6 minutes before the second run, a 5 minute aerosol of saline (0.9%) or hypertonic saline (14.4%) was administered. The radiolabelled tracer particles were given via the 0.5 um hole made in the trachae. An aerosol of ether saline (0.9%) or hypertonic saline (14.4%) was generated by a Pari pressure nebulizer. The aerosol was switched off one minute before the second run. The procedure for TMV measurements is described, in detail, in Newton et al., "Cilia, Mucus and Mucociliary Interactions." Ed. Baum, G.L., Preil, Z., Roth, Y., Liron., Ostfield, E., Marcel Dekker. New York, 1990 and Newton et al. in *Pediatric Pulmonology* S17, Abs 364, 1998.

Results

As shown in Figure 18, hypertonic saline (14.4% x 5 mins) caused a transient increase in TMV immediately after aerosol.

Example 15

Effect of Amiloride on TMV in the Guinea-pig

The aim of this study was to investigate the effect of amiloride (10mM x 20 min.) on guinea-pig tracheal mucus in the anaesthetized spontaneously breathing guinea pig. This agent was delivered into the cephalad trachea by aerosolization as described in Example 14. The TMV measurement procedure used in this Example is described in Newton et al. in "Cilia, Mucus and Mucociliary Interactions," Ed., Baum, G.L. et al., Marcel Dekker, New York, 1998; Newton et al., *Ped. Pulm.* S17, Abs. 364, 1998).

Materials and methods/Reagents used

An amiloride formulation (10 mM) in water was prepared for this example. Hypnorm® (Fentanyl citrate 0.315 mg/mL and Fluanisone 10 mg/mL) was obtained from Janssen Animal Health and Hypnovel® (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea pigs (550-750 g) were supplied by Harlan UK Ltd. Thermistor probes were obtained from Kane-May Ltd, UK.

Measurement of tracheal mucus velocity

Animals were anaesthetized using Hypnorm® and Hypnovel®. TMV was monitored using a lead collimated miniature Beta particle detector probe arranged to detect the radioactivity emitted from an injected aliquot of ³²P-labelled *Saccharomyces cerevisiae* as it was transported on the tracheal mucociliary layer of an anaesthetized

WO 00/37099

PCT/GB99/04381

guinea pig. Guinea pigs were anaesthetized with Hypnorm® and Hyponovel at time 0. Amiloride (10 mM x 20 min) was administered by aerosol. The first TMV measurement was made immediately afterwards and subsequent measurements were taken every 15 minutes.

5

Results

As shown in Figure 19, amiloride (10 mM x 20 mins) caused a statistically significant increase in TMV 15 minutes after aerosol.

10 **Example 16**

Aprotinin double mutein decreases sodium current in cultured human bronchial epithelial (HBE) cell short circuit current (Isc)

The aim of this study was to investigate the effect of the Kunitz family serine protease inhibitor Aprotinin double mutein on Isc in vitro. Tertiary HBE
15 cell monolayers grown to confluence were mounted in modified Ussing chambers, immersed in Krebs buffer (KBR) solution and bubbled with 95% O₂/5%CO₂ warmed to 37°C. Aprotinin double mutein is Des Pro2-Ser10-Arg15-Asp24-Thr26-Glu31-Asn41-Glu53-Aprotinin which is described in Example 1 of EP 821 007, published January 28, 1998.

20 Cells were left to equilibrate for 20 minutes before calibrating for background noise and fluid resistance. Transepithelial potential difference was then clamped to 0 mV using a WPI EVC 4000 voltage clamp. Ag/AgCl electrodes were used to monitor Isc. Once a stable baseline was achieved (typically 10-20 mins), cells were treated with a amiloride (10uM). Once a
25 response to amiloride was seen, it was washed out with KBR solution. After return to baseline and equilibration, Bikunin (5 ug/mL), Aprotinin double mutein (0.5 to 5 ug/mL), Aprotinin (1.5 to 5 ug/mL) or PBS was added. 90 minutes following agent treatment, amiloride (10uM) was added.

30 **Results**

As shown in Figure 20, Aprotinin double mutein (0.5 to 5 ug/mL) dose dependently inhibited sodium current in vitro in human bronchial epithelial cells over a 90 minute period.

35 **Example 17**

Expression, purification and comparative protease inhibitory activity of placental Bikunin (1-170) expressed in Chinese Hamster Ovary (CHO) cells

(a) Development of stable, high-producing CHO cell lines that express Bikunin

Stable production cell lines that secrete high quantities of bikunin were developed by transfecting CHO (dhfr-) cells with the expression vector shown in Figure 27. The vector was constructed using standard recombinant DNA techniques. A description of the construction of the expression vector and CHO cell expression system can be found in U.S.S.N. 09/441,654, filed November 12, 1999, entitled "Method of Producing Glycosylated Bikunin," by Inventor Sam Chan. Briefly, the expression vector pBC-BK was constructed by cloning bikunin cDNA immediately downstream of the cytomegalovirus immediate early promoter and upstream of the polyadenylation signal sequence. The expression vector pBC-BK consists of a transcriptional unit for bikunin, dihydrofolate reductase, and ampicillin resistance. Bikunin cDNA was released from the cloning vector by restriction enzymes, blunt-ended, and ligated to linearized pBC. The linearization of pBC was done by a single restriction enzyme digestion. The orientation of bikunin cDNA was confirmed by sequencing.

About 1×10^6 CHO (Chinese hamster ovary) cells were transfected with 10 μ g of pBC-BK using Lipofectin reagents (Life Technology, Bethesda, Maryland) according to manufacturer's instructions. The cells were then selected in the presence of 50 nM methotrexate and grown in DME/F12 media deficient in thymidine and hypoxanthine plus 5% dialyzed fetal bovine serum. Cell populations were screened for bikunin production with a chromogenic assay. Briefly, bikunin standards or culture fluid was serially diluted and incubated with an equal volume of kallikrein at 37° C for 30 minutes after which a chromogenic substrate, N-benzoyl-Pro-Phe-Arg-pNA, was added. The reaction was incubated for 15 minutes before the addition of 50% acetic acid. The amount of p-nitroanilide released was measured at 405 nM. The high producing populations were further selected in media containing increasing concentrations of methotrexate (100 to 400 nM methotrexate) and screened for the production of bikunin. Limiting dilution cloning was then applied to derive clones with high and stable productivity. The cloning was done in the absence of methotrexate using standard tissue culture techniques by depositing 1 cell/well in 96-well plates. A clone designated FD3-1 was chosen for productivity evaluation in a bioreactor and was deposited on November 12, 1999 with the American Type Culture Collection (ATCC), Rockville, MD, and was assigned Patent Deposit

WO 00/37099

PCT/GB99/04381

Designation PTA-940.

(b) Serum-free production of bikunin in a perfusion bioreactor

Continuous production of bikunin was done by continuous perfusion fermentation. A 1.5 liter Wheaton fermenter was inoculated with a stable CHO cell line at 2×10^6 cells/ml and perfused at a medium exchange rate of 0.5 liters/day. The production medium was a DME/F12-based medium supplemented with insulin (10 μ g/ml) and $\text{FeSO}_4 \cdot \text{EDTA}$ (50 μ M). The cell density was maintained at 4×10^6 cells/ml. The average daily yield of the fermenter was ~20 mg/day. The production of bikunin was stably maintained for 21 days.

(c) Purification of bikunin (1-170) produced from a CHO cell expression system

Bikunin produced from CHO cells was purified using standard chromatography techniques involving ion exchange, metal chelate, and size exclusion chromatography as outlined in Figure 29.

The SP column (18 x 10 cm, 2.5 L) was prepared with SP-Sepharose Fast Flow (Pharmacia), and equilibrated. Cold filtered CHO cell harvest (TCF) was diluted 1:2.5 with cold sterile water, and the pH was adjusted to 5.0. Chromatography was performed at ambient temperature with cold buffers. The cold starting material was loaded on the column at 800 mL/min (189 cm/hr). The amount of bikunin loaded onto the column ranged from 0.888 - 1.938 g (approximately 14 mg/L). After loading, the column was washed with equilibration buffer and the bikunin eluted with elution buffer. The eluate was collected at 2 - 8 °C (in an ice bath) and immediately adjusted to pH 7 with 6 N NaOH. The column was washed, then sanitized with cold (2 - 8 °C) 1 N NaOH, and stored at 2 - 8 °C in 20% ethanol until its next use. The equilibration and wash buffer contained 50 mM NaCl, 30 mM NaH_2PO_4 , pH 5.0; the elution buffer contained 350 mM NaCl, 30 mM NaH_2PO_4 , pH 5.0; and the pH adjusting buffer was 1 M citric acid, 1 M NaH_2PO_4 , pH 2.4.

Thawed SP-Sepharose eluate was concentrated by ultrafiltration (UF) approximately 10-fold to reduce the volume before a 5 to 7-fold diafiltration (DF) was performed in preparation for anion exchange chromatography. All operations were performed at ambient temperature in a horizontal flow hood. UF/DF utilized a Pellicon 2 "mini" filter system from Millipore* (Bedford, MA) and two 10 kDa regenerated cellulose cartridges (P2C010C01). Flux rates were

approximately 130 ± 20 mL/min for the two-cartridge system and were maintained by regulating the inlet and outlet pressures between 22 to 26 psi and 12 to 16 psi respectively. Circulation was with a peristaltic pump; recirculation was set to 500 to 600 mL per minute before transmembrane pressure adjustment.

5 Diafiltration was performed with cold 10 mM NaH_2PO_4 buffer, pH 8.1.

The Q-Sepharose column chromatography was performed as follows. A 13 x 9 cm, 1.2 L column of Q-Sepharose Fast Flow (Pharmacia) was washed with 5 column volumes (CV) of sterile water and equilibrated with approximately 10 CV's equilibration buffer. Diafiltered SP eluate was adjusted to pH 8.1 and

10 applied on the Q-Sepharose column at 100 mL/min (45 cm/hr). The amount of bikunin loaded onto the column ranged from 1121-2607 mg (approximately 15 mg/mL). After loading, the column was washed with equilibration buffer until the UV absorbance at A280 reached baseline; then the bikunin was eluted. The eluate was collected and used as feed material for Zn-IMAC, zinc immobilized

15 metal ion adsorption chromatography. The column was cleaned with 1 M NaOH, rinsed with sterile water, and stored in 20% ethanol. All operations were performed at 2 - 8 °C. Equilibration and washing buffer for Q-Sepharose column contained 10 mM NaH_2PO_4 , pH 8.1. The elution buffer contained 10 mM NaH_2PO_4 , 100 mM NaCl, pH 8.1.

20 A Zn-IMAC column (5 x 10 cm) containing approximately 200 mL bed volume of Chelating Sepharose Fast Flow (Pharmacia) was charged with 3 volumes of ZnSO_4 solution (see below); washed with 2 volumes of sterile water, and equilibrated with 6 volumes of buffer as described below. The Q-Sepharose eluate was adjusted to pH 7.4 and 300 mM NaCl (by addition of NaCl solid),

25 applied to Zn-IMAC at 30 mL/min (92 cm/hr linear rate), and then the column was washed with equilibration buffer until the UV absorbance reached baseline. The amount of bikunin loaded onto the column ranged from 0.097 - 1.681 g (approximately 0.63 mg/mL). The flowthrough and wash were collected for UF. The column was stripped, and sanitized with 0.5 M NaOH. All operations of this

30 step were performed at 2 - 8 °C. Equilibration buffer for Zn-IMAC contained 10 mM NaH_2PO_4 , 300 mM NaCl, pH 7.4; the strip buffer contained 50 mM EDTA, 10 mM NaH_2PO_4 , 300 mM NaCl, pH 7.4; charging solution, 2 mg/mL $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$.

The Zn-IMAC flow-through was concentrated by ultrafiltration 5-fold

35 with the Pellicon 2 system (Millipore) for Sephacryl S-200 chromatography. Permeate flux rates were approximately 60 to 70 mL/min and were maintained as described earlier. Recirculation was at 400 to 500 mL/min.

WO 00/37099

PCT/GB99/04381

A column (10 x 58.5 cm) containing 4.59 L of Sephacryl S-200 High Resolution (Pharmacia) was equilibrated with 137 mM NaCl, 2.7 mM KCl, 2.9 mM KH₂PO₄, 10 mM NaH₂PO₄, 8 mM Na₂HPO₄, 2 mg/L Tween^{*}80, pH 7.2. Flow rate was 30 mL/min (23 cm/hr). In a typical run, 1475 mg of bikunin in a volume of 95 mL was applied to the column.

The post-S-200 pool was treated with Etox resin in batch mode for removal of any potential pyrogen. ActiClean Etox^{*}, 2705 resin (Sterogene Bioseparations, Inc.) was rinsed with 1 M NaOH and incubated for 5 hours in 1 M NaOH at room temperature with agitation. The resin was washed and equilibrated with PBS, pH 7.2. Eighty mL of the filtered and equilibrated resin was added to 1063 mL of bikunin S200 pool (5460.11 mg), and agitated overnight at 2 - 8°C. The ETOX resin was then removed by filtration with 0.2 micron Nalgene flask filter.

Results. Table 10 shows the average yield afforded by each step.

Table 10

Purification Step	Average Yield (%)
SP-Sepharose	88.1
UF/DF plus filtration	81
Q-Sepharose	59 ± 14
Zn-IMAC	99.5
Sephacryl S-200	81
ETOX Resin	93

Overall yield of bikunin was about 30% with a purity of 95%. Mass spectroscopy data also suggested that in addition to full length Bikunin (1-170) molecules, species lacking three (G-S-K) and four (L-G-S-K) amino acids from the carboxy end of Bikunin (1-170) were present in the pure protein pool. The material produced was shown to be stable to degradation when exposed to 72-hours incubation at ambient temperature or at 37°C, neutral pH. N-terminal sequencing, gel electrophoresis, immunoblotting, and amino acid analysis indicated that the bikunin was substantially pure (no other sequences were detected). An additional reverse phase chromatography step revealed that the CHO-derived purified bikunin was still able to be fractionated into several species (Figure 30A). CHO bikunin (8.5 mg) was adjusted to pH 2.5 with trifluoroacetic acid (TFA, 0.1% final concentration) and subjected to

chromatography on a C18 reverse-phase column (Vydac, 2.5 x 25 cm) equilibrated in 17.5% acetonitrile and 0.1% TFA at a flow rate of 2 ml/min. CHO bikunin was eluted with a linear gradient of 17.5-40% acetonitrile in 0.1% TFA over 60 min. Figure 30B shows the silver stained SDS-PAGE profile of these fractions (lane between 54 and 55 represents molecular size markers).

Preliminary carbohydrate analysis was performed on the glycosylated isoforms of CHO bikunin having a MW ranging from about 21 kDa to about 38 kDa. The total sugar content was found to be 7.5%. Both N-linked sites (Asn-30 and Asn-67) were found to be occupied with carbohydrate structures. Chromatographic and mass spectrometric analysis confirmed the presence of very heterogeneous and highly branched oligosaccharide structures contributing to the size heterogeneity observed for the purified bikunin. About 90% of the oligosaccharides were sialylated and the remaining structures were neutral. When treated with N-Glycosidase F, the glycosylated isoforms of CHO bikunin (Figure 30B) were converted to a single 18 kDa isoform (See Figure 31).

The sialic acid content of bikunin was analyzed by incubation with sialidase in 50 mM sodium acetate buffer, pH 5.0, for 18 hours in a capped microfuge tube. Sialic acids were separated on a Carbo Pac PA1 anion-exchange column using a buffer gradient of 20-250 mM sodium acetate in 100 mM NaOH for 50 minutes at a flow rate of 1 ml/min. Detection was done with a pulsed electrochemical detector and quantitated by comparing retention times and peak areas of samples to standard sialic acids (N-acetylneuraminic acid and N-glutarylneuraminic acid). The results are shown in Table 11.

Table 11. Sialic acid composition of bikunin

Sialic acid	Contents (g/100g of bikunin)
N-Acetylneuraminic acid	5.4
N-Glutarylneuraminic acid	0

Example 18

Comparative Protease Inhibitor Activity of Placental Bikunin (1-170) expressed in Chinese Hamster Ovary (CHO) cells

General. The *in vitro* specificity of recombinant bikunin was determined using the materials and methods as described in Examples 3, 4, 7, and 10. Table 12 below shows the efficacy of recombinant bikunin as an inhibitor of various serine proteases *in vitro*. Data is shown using either recombinant bikunin or aprotinin.

Proteases. Human plasmin and human plasma kallikrein quantitation was carried out by active site titration using p-nitrophenyl p'-guanidinobenzoate HCl as previously described (Chase, T., and Shaw, E., (1970) *Methods Enzymol.* 19, 20-27). Human tissue kallikrein (Bayer, Germany) was quantitated by active site titration using bovine aprotinin as a standard and PFR-AMC as a substrate assuming a 1:1 complex formation. The K_m for GPK-AMC with plasmin under the conditions used was 726 μ M; the K_m for PFR-AMC with human plasma kallikrein was 457 μ M; the K_m for PFR-AMC with human tissue kallikrein was 5.7 μ M.

Inhibition Kinetics: The inhibition of human plasmin by CHO expressed placental bikunin (1-170) and aprotinin was determined with plasmin (50 pM) and CHO expressed placental bikunin (1-170) (0-2 nM) or aprotinin (0-4 nM) in buffer containing 50 mM Tris-HCl (pH 7.5), 0.1 M NaCl, and 0.02% triton x-100. After 30 min. incubation at 37°C, 25 μ l of 20 mM GPK-AMC was added and the change in fluorescence monitored. The inhibition of human plasma kallikrein by CHO expressed placental bikunin (1-170) or aprotinin was determined using kallikrein (0.2 nM) and CHO expressed placental bikunin (1-170) (0-4 nM) or aprotinin (0-45 nM) in 50 mM Tris-HCl (pH 8.0), 50 mM NaCl, and 0.02% triton x-100. After 30 min. at 37°C, 5 μ l of 20 mM PFR-AMC was added and the change in fluorescence monitored.

The inhibition of human tissue kallikrein by aprotinin or CHO expressed placental bikunin (1-170) was measured by the incubation of 0.35 nM human tissue kallikrein with CHO expressed placental bikunin (1-170) (0-10 nM) or aprotinin (0-0.5 nM) in a 1 ml reaction volume containing 50 mM Tris-HCl buffer pH 9.0, 50 mM NaCl, and 0.1% triton x-100. After 5 min. at 37°C, 5 μ l of 2 mM PFR-AMC was added achieving 10 μ M final concentration and the change in fluorescence monitored.

The inhibition of Factor XIa (from Enzyme Research Labs, South Bend, IN) was measured by incubating FXIa (0.1 nM) with either 0 to 40 nM CHO expressed placental bikunin (1-170) or 0 to 4 μ M aprotinin in buffer containing 50 mM Hepes pH 7.5, 100 mM NaCl, 2 mM CaCl_2 , 0.01% triton x-100, and 1% BSA in a total volume of 1 ml. After 30 min at 37°C, 10 μ l of 40 mM Boc-Glu(OBzl)-Ala-Arg-AMC (Bachem Biosciences, King of Prussia, PA) was added and the change in fluorescence monitored.

The apparent inhibition constant K_i^* was determined using the nonlinear

regression data analysis program Enzfitter software (Biosoft, Cambridge, UK):
The kinetic data from each experiment were analyzed in terms of the equation
for a tight binding inhibitor:

$$V_i/V_o = 1 - (E_o + I_o + K_i^* - [(E_o + I_o + K_i^*)^2 - 4 E_o I_o]^{1/2}) / 2E_o \quad (2)$$

where V_i/V_o is the fractional enzyme activity (inhibited vs. uninhibited rate),
and E_o and I_o are the total concentrations of enzyme and inhibitor, respectively.
Ki values were obtained by correcting for the effect of substrate according to the
equation:

$$K_i = K_i^* / (1 + [S_o]/K_m) \quad (3)$$

(Boudier, C., and Bieth, J. G., (1989) *Biochim Biophys Acta*. 995: 36-41).

Results: The K_i values are listed in Table 12 below.

**Table 12. Comparison of K_i values for the inhibition of various proteases by
CHO Bikunin (1-170) or aprotinin**

Protease	CHO bikunin (1-170) K_i (nM)	Aprotinin K_i (nM)
Human plasma kallikrein	0.5	23.0
Human tissue kallikrein	0.3	0.004
Human plasmin	0.2	0.2
Human FXIa	1.9	270.0

The results show that recombinant bikunin can be expressed in CHO cells
to yield an active protease inhibitor that is effective against at least three
different serine proteases. Recombinant bikunin was more potent than aprotinin
against human plasma kallikrein, and human FXIa. It was equipotent with
aprotinin at inhibiting human plasmin.

Example 19

**Aprotinin decreases sodium current in cultured human cystic fibrosis
bronchial epithelial cell short circuit current (I_{sc}) in vitro**

HBE cells were isolated from CF patient lung transplant tissue and grown
in collagen coated flasks for one week. The cells were then passaged and seeded

WO 00/37099

PCT/GB99/04381

onto collagen coated Costar Transwell filters (0.33 cm²) and grown in DMEM/F12 media supplemented with 2% Ultros^{*}er G. Cells were grown at an air liquid interface and used 2 to 4 weeks after seeding:

Cells on the Transwell filters were mounted in modified Costar Ussing chambers and studied under Isc conditions. Baseline Isc values were recorded (0 to 10 minutes), and then aprotinin (1mg/mL in PBS) was added to the apical side. Isc was recorded for 100 minutes and then the apical bath fluid was exchanged with fresh buffer. Trypsin (100 BAEE units/mL) was added to the apical side and Isc recorded for 10 minutes. Amiloride (10 μ M) was then added to the apical side and Isc was recorded for a further 10 minutes.

Results

As shown in Figure 21, aprotinin (1mg/mL in PBS) inhibited Isc in vitro in human CF bronchial epithelial cells over a 100 minute period. After washout, Isc was increased by treatment of the apical surface with the serine protease, trypsin. Finally, addition of amiloride (10 μ M) demonstrated that the changes in Isc were the result of changes in sodium dependent current.

Example 20

Assessment of activity of Bikunin (1-170) following nebulization

The aim of this study was to assess the anti-protease activity of Bikunin (1-170) described in Example 17 following nebulization. All studies were performed with Bikunin formulated in phosphate buffered saline, pH 7.4 (137 mM NaCl, 3 mM KCl, 3mM KH₂PO₄, 8 mM Na₂HPO₄, 0.02 g/L Tween 80).

Methods of nebulization

Raindrop® Medication Nebulizer: Bikunin was aerosolized at concentrations of 1 and 3 mg/mL. The nebulizer cup was filled with 2.5 mL of Bikunin solution. Aerosolisation was performed at 7.35 L/min or 35 psi.

Collison Nebulizer: Bikunin was aerosolised at a concentration of 4.7 mg/mL. The nebulizer cup was filled with 2.5 mL of Bikunin solution. Aerosolisation was performed at 35 psi.

Collection of nebulised samples

Aerosolised Bikunin was collected using a twin impinger (6.4 μ m mean aerodynamic particle cut-off size at 60 L/min through the system). The impinger works on the principle of liquid impingement and divided the aerosol into a non-respirable fraction (>6.4 μ m collected in Stage 1) and a respirable fraction (<6.4 μ m collected in Stage 2).

Measurement of anti-protease activity

Bikunin activity was measured in vitro by its inhibition of human plasma kallikrein.

Results

5 Raindrop® Medication Nebulizer: Activity (K_i) values for the pre- and post-nebulization samples were as follows: 1 mg/mL Bikunin: K_i values were 0.47(± 0.02) and 0.76 (± 0.04) respectively; and for 3 mg/mL Bikunin(1-170) the K_i values were 0.52 (± 0.03) and 0.62 (± 0.03) respectively.

10 Collison Nebulizer: Activity (K_i) values for the pre- and post-nebulization samples were 0.27 (± 0.03) and 0.45 (± 0.03) respectively.

Conclusion:

15 Pre- and post-nebulized Bikunin samples from both the Raindrop and Collison nebulizers exhibited similar activities (similar K_i values within assay variability), indicating that Bikunin (1-170) was stable to aerosolization and retained its anti-protease activity following nebulization.

Example 21**The effect of Bikunin on Tracheal Mucus Velocity in the Sheep**

20 The aim of this study was to investigate the effect of the Kunitz family serine protease inhibitor Bikunin (1-170) described in Example 17 on sheep tracheal mucus velocity over 8 hours after treatment. This agent was delivered by nebulised aerosol administration to the airways. The procedure used in this example is described in O'Riordan et al., J. Applied Physiol. 85(3), 1086-1091, 1998.

25 *Measurement of TMV*

30 Adult ewes were restrained in an upright position, with their heads immobilized, in a specialized body harness. They were nasally intubated with an endotracheal tube, with the cuff placed just below the vocal cords. The inspired air was warmed and humidified. To minimize possible impairment of TMV caused by inflated cuffs, the endotracheal tube cuff remained deflated throughout the study except for the period of aerosol administration.

35 To measure TMV 5 to 10 radiopaque Teflon particles (approximately 1 mm in diameter, 0.8 mm thick, and weighing 1.5 to 2 mg) were insufflated into the trachea via a catheter placed within the endotracheal tube. The movement of the Teflon particles was then measured over a 1 minute period. The procedure used in this example is described in Russi et al., J. Applied Physiol. 59(5), 1416-1422, 1985. A collar containing radiopaque markers of known length was

applied to the exterior of the animals and used as a standard to convert distance traversed by the particles on the video screen to actual distance traveled. TMV was calculated from the average distance in a cephalad direction traveled per minute for 5 to 10 Teflon particles. Baseline TMV was measured immediately prior to administration of aerosol.

Test substances; PBS, 1 mg/mL Bikunin in PBS, or 3 mg/mL Bikunin in PBS; were delivered to the sheep airways as an aerosol (3mL) generated using a Raindrop jet nebuliser operated at a flow rate that produced droplets of mass median aerodynamic diameter of 3.6 μ m. TMV was measured immediately after administration of test substance (0 hours), then again at 0.5, 1, 2, 3, 4, 5, 6, 7, and at 8 hours.

Results

As shown in Figure 22, 9 mg Bikunin aerosol (3 mL of 3 mg/mL) delivered to sheep airways significantly increased TMV at 0, 0.5, 3, 4, 5, 6, 7, and 8 hours compared to the same time points for a group of animals receiving PBS vehicle aerosol. At 24 hours TMV had returned to baseline rates in both the Bikunin treatment and in the PBS vehicle groups.

At a lower dose (3mg Bikunin aerosol (3mL of 1 mg/mL)), no significant differences were observed in TMV between treatment and vehicle groups at any time point studied.

Example 22

Bikunin (50 μ g/mL) decreases sodium current in cultured guinea pig tracheal epithelial (GPTE) cell short circuit current (Isc)

The aim of this study was to investigate the effect of Bikunin (1-170) (See Example 17) on Isc on GPTE cells in vitro. GPTE cells were seeded onto 1.2 cm diameter, 0.4 μ m pore size Snapwell™ inserts (Costar UK). Cells were grown to confluence and mounted in modified Ussing chambers 2-4 days after placement on air-liquid interface. Inserts were immersed in Krebs buffer (KBR) solution and bubbled with 95% O₂/5%CO₂ warmed to 37° C.

After a 20 minute equilibration period, transepithelial potential difference was then clamped to 0 mV using a WPI EVC 4000 voltage clamp. Ag/AgCl electrodes were used to monitor Isc. Once a stable baseline was achieved (typically 20-30 mins), cells were treated with amiloride (30 μ M). Once a response to amiloride was obtained, it was washed out with KBR solution. After return to baseline and equilibration, Bikunin (1-170) described in Example 17 (10 to 50 μ g/mL) or PBS was added. 30 minutes following agent treatment

WO 00/37099

PCT/GB99/04381

amiloride (30 uM) was added.

Results

As shown in Figure 23, Bikunin (1-170) (50 ug/mL) inhibited sodium
5 current in vitro in guinea pig tracheal epithelial cells over a 30 minute period.

Example 23

**Bikunin (1-170) (100 ug/mL) decreases sodium current in cultured ovine
tracheal epithelial (OTE) cell short circuit current (Isc)**

10 The aim of this study was to investigate the effect of Bikunin (1-170) (See
Example 17), on Isc in OTE cells *in vitro*. OTE cells were seeded onto 1.2 cm
diameter, 0.4 um pore size Snapwell™ inserts (Costar UK). Cells were grown to
confluence and mounted in modified Ussing chambers 3-5 days after placement
on air-liquid interface. Inserts were immersed in Krebs buffer (KBR) solution
15 and bubbled with 95%O₂/5%CO₂ warmed to 37° C.

After a 20 minute equilibration period, transepithelial potential difference
was then clamped to 0 mV using a WPI EVC 4000 voltage clamp. Ag/ AgCl
electrodes were used to monitor Isc. Once a stable baseline was achieved
(typically 30 mins), cells were treated with amiloride (10 uM). Once a response
20 to amiloride was obtained, it was washed out with KBR solution. After return to
baseline, Bikunin (1-170) described in Example 17 (25, 50 or 100 ug/mL) or PBS
was added. 90 minutes following agent treatment amiloride (10uM) was again
added.

Results

25 As shown in Figure 24, Bikunin (1-170) (100 ug/mL) significantly inhibited
sodium current in vitro in ovine tracheal epithelial cells over a 90 minute period.

Example 24

**The effect of Bikunin (1-170) on tracheal potential difference in guinea pigs
30 pretreated with LPS**

Polymorphonuclear (PMN) leukocyte (neutrophil) dominated airway
inflammation is often a feature of CF lung disease. In the guinea pig, exposure to
an aerosol of E. coli lipopolysaccharide (LPS) induces a marked PMN influx in
the bronchoalveolar lavage fluid 24 hours post challenge. The aim of this study
35 was to investigate the effect of Bikunin (1-170) described in Example 17 on
tracheal potential difference in guinea pigs pre-exposed to an aerosol of LPS.
Agents were delivered into the cephalad trachea by topical instillation. TPD was

WO 00/37099

PCT/GB99/04381

monitored for 60 minutes, 23 hours after exposure to LPS.

Materials/Reagents

Bikunin (1-170) was formulated in Hank's Balanced salt solution (HBSS). Amiloride was obtained from Sigma Chemicals and formulated in HBSS.

- 5 Vehicle control was HBSS. Hypnorm (Fentanyl citrate 0.315 mg/mL and Fluanisone^{*} 10 mg/mL) was obtained from Janssen Animal Health and Hypnovel (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea pigs (600-700 g) were supplied by Harlan UK. Thermistor probes were obtained from Kane-May Ltd, UK.

10 *Induction of PMN Influx*

Individual animals were exposed to an aerosol of 0.03 mg/mL LPS or PBS for 10 minutes.

Preparation of guinea pig for measurement to tracheal potential difference

- 23.4 hours following LPS treatment, guinea pigs were anaesthetised with
- 15 Hypnorm and Hypnovel and immobilised in a supine position. A ventral midline incision was made from the lower jaw to the clavicles. Using blunt dissection a length of trachea was exposed and bisected at the upper edge of the sternum. The external jugular vein was exposed and cannulated. The caudal part of the trachea was then cannulated to allow the animal to spontaneously
- 20 breath room air. The animal was then placed supine and body temperature was maintained at 37°C by manual adjustment of a heat lamp. Rectal temperature was monitored with a thermistor probe.

- 20 minutes following induction of anaesthesia, Bikunin (50 µg/mL) or amiloride (100 µM) was topically instilled into the cephalad trachea. The
- 25 tracheal agar electrode was then inserted into the cephalad trachea and tracheal potential difference was measured for 60 minutes. The reference electrode was placed under the cephalad trachea in contact with the trachea cartilage. The wound site was covered to prevent drying.

Results

- 30 As shown in Figure 25(a), exposure to LPS caused a significant PMN influx. Bikunin significantly inhibited potential difference in guinea pigs pre-exposed to LPS, as shown in Figure 25(b).

Example 25

- 35 **The effect of Aprotinin double mutein on tracheal mucus velocity in the guinea pig**

The aim of this study was to investigate the effect of Aprotinin double

mutein described in Example 16 on guinea pig tracheal mucus velocity 1.5 hours post treatment. This agent was delivered into the cephalad trachea by topical instillation. TMV was monitored 1.5 hours later for 60 minutes.

Materials/Reagents

5 Aprotinin double mutein (see Example 16) was obtained from Biotechnologie, Bayer AG, Germany USA and formulated in Hank's Balanced salt solution (HBSS). Hypnorm (Fentanyl citrate 0.315 mg/mL and Fluanisone 10 mg/mL) was obtained from Janssen Animal Health and Hypnovel (Midazolam 5 mg/mL) was obtained from Roche. Male Dunkin-Hartley guinea
10 pigs (600-750 g) were supplied by Harlan UK. Thermistor probes were obtained from Kane-May Ltd., UK.

Induction of anaesthesia:

Animals were anaesthetised using halothane. Once a satisfactory level of anaesthesia was induced a small incision was made below the lower jaw. The
15 trachea was exposed and 100 ul vehicle or Aprotinin double mutein (10 ug) was injected into the airway lumen using a needle and syringe. Once injected the skin overlying the tracheal injection site was repaired. The animals were then allowed to recover.

Measurement of tracheal mucus velocity

20 Tracheal mucus velocity (TMV) was monitored using a lead collimated miniature Beta-particle detector probe arranged to detect the radioactivity emitted from an injected aliquot of phosphorus-32-labelled *Saccharomyces cerevisiae* as it was transported on the tracheal mucociliary layer of an anaesthetised guinea pig (Newton and Hall 1998). 70 minutes following
25 instillation of test agent, guinea pigs were anaesthetised for a second time with hypnorm and hypnovel and immobilised in a supine position. The first TMV measurement was made 20 minutes afterwards.

Results:

30 As shown in Figure 26, Aprotinin double mutein (10 ug) increased TMV in vivo in the guinea pig, relative to HBSS, over a sustained period of 1.5 to 2.5 hours following administration.

Example 26

**Bikunin (1-170) decreases sodium current in cultured human cystic fibrosis
35 bronchial epithelial cell short circuit current (I_{sc}) in vitro**

HBE cells were isolated from CF patient lung transplant tissue and grown in collagen coated flasks for one week. The cells were then passaged and seeded

onto collagen coated Costar Transwell filters (0.33 cm²) and grown in DMEM/F12 media supplemented with 2% Ultrosor G. Cells were grown at an air liquid interface and used 2 to 4 weeks after seeding.

Cells on the Transwell filters were mounted in modified Costar Ussing chambers and studied under Isc conditions. Baseline Isc values were recorded (0 to 20 minutes), and then bikunin (1-170) (see Example 17) (10 ug/mL in PBS) was added to the apical side. Isc was recorded for 90 minutes and then amiloride (10uM) was added and Isc was recorded for a further 10 minutes. The apical bath fluid was then exchanged with fresh buffer, and Isc was recorded for a further 15 minutes. Trypsin (1 uM) was added to the apical side and Isc recorded for 15 minutes. Amiloride (10 uM) was then added to the apical side and Isc was recorded for a further 10 minutes.

Results

As shown in Figure 28(a), bikunin (1-170)(10 ug/mL in PBS) inhibited Isc in vitro in human CF bronchial epithelial cells over a 90 minute period. Isc was further reduced by addition of amiloride (10 uM). On washout, Isc was increased back to the current achieved before the addition of amiloride. After a further 15 minutes, the apical surface was treated with trypsin (1 uM) and this further increased Isc to the baseline current level (i.e. that observed at 20 minutes). Finally, addition of amiloride (10 uM) inhibited the majority of the current, and demonstrated that the changes in Isc were the result of changes in sodium dependent current.

Figure 28(b) shows that bikunin (1-170) at 1, 5, and 10 ug/mL, and aprotinin at 20 ug/mL inhibited Isc at 90 minutes after apical application to human CF bronchial epithelial cells in vitro.

Although certain embodiments of the invention have been described in detail for the purpose of illustration, it will be readily apparent to those skilled in the art that the methods and formulations described herein may be modified without departing from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

WO 00/37099

PCT/GB99/04381

SEQUENCE LISTING FREE TEXT

- <210> 9
<223> /note= "n at positions 622, 679, 707 is any nucleic acid"
- 5 <210> 10
<223> /note= "Xaa at positions 201, 226, and 233 are nonsense or stop codons"
- 10 <210> 11
<223> /note= "Xaa at positions 8, 17, 21-26, 40, 42, 45-47, 52, 64, 103, 112, 114, 116-121, 135, 137, 140-142, 147, and 159 is any amino acid residue"
- 15 <210> 12
<223> /note= "residue 361 is any nucleic acid"
<223> /note= "residue 367 is any nucleic acid"
<223> /note= "residue 384 is any nucleic acid"
<223> /note= "residue 390 is any nucleic acid"
- 20 <210> 13
<223> /label= signal peptide
<223> /note= "Xaa at positions 111, 120, 122, 128, and 130 represents a nonsense or stop codon"
- 25 <210> 14
<223> /note= "n at positions 425, 482, and 510 is any nucleic acid"
- 30 <210> 15
<223> /note= "Xaa at positions 2, 23, 132, 160, and 167 represent a nonsense or stop codon"
- 35 <210> 16
<223> /note= "n at positions 3, 11, 12, 17, 51 and 429 represent any nucleic acid"
- <210> 17
<223> /note= "n at positions 6, 310 and 408 represent any nucleic acid"

WO 00/37099

PCT/GB99/04381

- <210> 18
<223> /note= "Tissue factor pathway inhibitor precursor 1"
- <210> 19
5 <223> /note= "Tissue factor pathway inhibitor precursor 1"
- <210> 20
<223> /note= "Tissue factor pathway inhibitor precursor"
- 10 <210> 21
<223> /note= "Tissue factor pathway inhibitor precursor 2"
- <210> 22
<223> /note= "Tissue factor pathway inhibitor precursor 2"
- 15 <210> 23
<223> /note= "Amyloid Precursor Protein homologue"
- <210> 24
20 <223> /note= "Aprotinin"
- <210> 25
<223> /note= "Inter alpha-trypsin inhibitor precursor"
- 25 <210> 26
<223> /note= "Inter alpha-trypsin inhibitor precursor"
- <210> 27
<223> /note= "Amyloid precursor protein"
- 30 <210> 28
<223> /note= "Collagen alpha-3 (VI) precursor"
- <210> 29
35 <223> /note= "HKI-B9"
- <210> 32
<223> /note= "cDNA of human Bikunin protein fragment"

WO 00/37099

PCT/GB99/04381

<210> 45
 <223> /label= signal peptide

5 <210> 47
 <223> /label= signal peptide

<210> 49
 <223> /label= signal peptide

10 <210> 50
 <223> /note= "Human Bikunin protein fragment"

<210> 51
 <223> /note= "Human Bikunin protein fragment"

15 <210> 52
 <223> /note= "Human Bikunin protein fragment"

<210> 53
 20 <223> /note= "Signal peptide of Human Bikunin protein"

<210> 54
 <223> Human Bikunin protein fragment

25 <210> 55
 <223> /note= "Oligomer for preparing expression construct"

<210> 56
 <223> Oligomer for preparing expression construct

30 <210> 57
 <223> /note= "Oligomer for preparing expression construct"

<210> 58
 35 <223> /note= "Oligomer for preparing expression construct"

<210> 61
 <223> /note= "Oligomer for preparing Bikunin expression construct"

WO 00/37099

PCT/GB99/04381

<210> 62

<223> /note= "Oligomer for preparing Bikunin construct"

<210> 63

5 <223> /note= "Human Bikunin protein fragment"

<210> 64

<223> /note= "Human Bikunin protein fragment"

10 <210> 65

<223> /note= "Human Bikunin protein fragment"

<210> 66

15 <223> /note= "Human Bikunin protein fragment"

<210> 67

<223> /note= "Human Bikunin protein fragment"

<210> 68

20 <223> /note= "Human Bikunin protein fragment"

<210> 69

<223> /note= "Human Bikunin protein fragment"

25 <210> 70

<223> /note= "Human Bikunin protein fragment"

<210> 71

30 <223> /note= "Human Bikunin protein fragment"

SEQUENCE LISTING

<110> Bayer Aktiengesellschaft
 <120> Method For Accelerating The Rate Of Mucociliary Clearance
 <130> 08891832CA
 <140> 2,356,404
 <141> 1998-12-22
 <150> PCT/GB99/04381
 <151> 1999-12-22
 <150> US 09/441,966
 <151> 1999-11-17
 <150> US 09/218,913
 <151> 1998-12-22
 <160> 105
 <170> PatentIn version 3.1
 <210> 1
 <211> 179
 <212> PRT
 <213> Homo sapiens
 <400> 1

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val
 1 5 10 15

Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr
 20 25 30

Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser
 35 40 45

Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val
 50 55 60

Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp
 65 70 75 80

Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser
 85 90 95

Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr
 100 105 110

Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg
 115 120 125

Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn
 130 135 140

Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln
 145 150 155 160

Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly
 165 170 175

Ala Val Ser

<210> 2
 <211> 197
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SIGNAL
 <222> (1)..(18)
 <223>

<400> 2

Ala Gly Ser Phe Leu Ala Trp Leu Gly Ser Leu Leu Leu Ser Gly Val
 1 5 10 15

Leu Ala Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser
 20 25 30

Lys Val Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn
 35 40 45

Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly
 50 55 60

Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala
 65 70 75 80

Thr Val Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala
 85 90 95

Ala Asp Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp
 100 105 110

His Ser Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala
 115 120 125

Val Thr Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val
 130 135 140

Glu Arg Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn
 145 150 155 160

Lys Asn Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg
 165 170 175

Gln Gln Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu
 180 185 190

Ala Gly Ala Val Ser
 195

<210> 3
 <211> 153
 <212> PRT
 <213> Homo sapiens

<400> 3

Ile His Asp Phe Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg Ala
 1 5 10 15

Ser Met Pro Arg Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln Leu
 20 25 30

Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr Lys
 35 40 45

Glu Glu Cys Leu Lys Lys Cys Ala Thr Val Thr Glu Asn Ala Thr Gly
 50 55 60

Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp Ser Ser Val Pro Ser Ala
 65 70 75 80

Pro Arg Arg Gln Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn Tyr
85 90 95

Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala Ser
100 105 110

Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn Phe
115 120 125

Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu Glu
130 135 140

Ala Cys Met Leu Arg Cys Phe Arg Gln
145 150

<210> 4
<211> 58
<212> PRT
<213> Homo sapiens

<400> 4

Ile His Asp Phe Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg Ala
1 5 10 15

Ser Met Pro Arg Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln Leu
20 25 30

Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr Lys
35 40 45

Glu Glu Cys Leu Lys Lys Cys Ala Thr Val
50 55

<210> 5
<211> 51
<212> PRT
<213> Homo sapiens

<400> 5

Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg Ala Ser Met Pro Arg
1 5 10 15

Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly
20 25 30

Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu
 35 40 45

Lys Lys Cys
 50

<210> 6
 <211> 58
 <212> PRT
 <213> Homo sapiens

<400> 6

Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala
 1 5 10 15

Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn
 20 25 30

Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu
 35 40 45

Glu Ala Cys Met Leu Arg Cys Phe Arg Gln
 50 55

<210> 7
 <211> 51
 <212> PRT
 <213> Homo sapiens

<400> 7

Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala Ser Phe Pro Arg
 1 5 10 15

Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn Phe Ile Tyr Gly
 20 25 30

Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu Glu Ala Cys Met
 35 40 45

Leu Arg Cys
 50

<210> 8
 <211> 92

<212> PRT
 <213> Homo sapiens

<400> 8

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val
 1 5 10 15

Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr
 20 25 30

Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser
 35 40 45

Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val
 50 55 60

Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp
 65 70 75 80

Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser
 85 90

<210> 9
 <211> 708
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Consensus DNA sequence of human Bikunin (Fig. 3).

<220>
 <221> misc_feature
 <222> (622)..(622)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (679)..(679)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (707)..(707)
 <223> "n" is any nucleotide.

<400> 9
 ggccgggtcg tttctcgccct ggctgggata gctgctcttc tctgggggtcc tggcggccga 60

```

ccgagaacgc agcatccacg acttctgcct ggtgtcgaag gtgggtgggca gatgccgggc 120
ctccatgcct aggtggtggt acaatgtcac tgacggatcc tgccagctgt ttgtgtatgg 180
gggctgtgac ggaaacagca ataattacct gaccaaggag gagtgcctca agaaatgtgc 240
cactgtcaca gagaatgcca cgggtgacct ggccaccagc aggaatgcag cggattcctc 300
tgtcccaagt gctcccagaa ggcaggattc tgaagaccac tccagcgata tgttcaacta 360
tgaagaatac tgcaccgcca acgcagtcac tgggccttgc cgtgcatact tcccacgctg 420
gtactttgac gtggagagga actcctgcaa taacttcac tatggaggct gccggggcaa 480
taagaacagc taccgctctg aggaggcctg catgctcgc tgcttccgcc agcaggagaa 540
tcctcccctg ccccttggct caaagggtgt ggttctggcc ggggctgttt cgtgatggtg 600
ttgatecttt tcctggggag cntccatggt cttactgatt ccgggtggca aggaggaacc 660
aggagcgtgc cctgcgganc gtctggagct tcggagatga caagggnt 708

```

<210> 10

<211> 197

<212> PRT

<213> Artificial Sequence

<220>

<223> Amino acids -18 to 179 of the translation of the consensus DNA sequence in Fig. 3.

<400> 10

```

Ala Gly Ser Phe Leu Ala Trp Leu Gly Ser Leu Leu Leu Ser Gly Val
1           5           10           15

```

```

Leu Ala Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser
          20           25           30

```

```

Lys Val Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn
          35           40           45

```

```

Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly
          50           55           60

```

```

Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala
65           70           75           80

```

```

Thr Val Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala
          85           90           95

```

Ala Asp Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp
 100 105 110

His Ser Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala
 115 120 125

Val Thr Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val
 130 135 140

Glu Arg Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn
 145 150 155 160

Lys Asn Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg
 165 170 175

Gln Gln Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu
 180 185 190

Ala Gly Ala Val Ser
 195

<210> 11
 <211> 179
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Variants of human Bikunin.

<220>
 <221> MISC_FEATURE
 <222> (8)..(8)
 <223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE
<222> (19)..(19)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (21)..(26)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (40)..(40)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (42)..(42)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (45)..(47)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
<221> MISC_FEATURE
<222> (64)..(64)
<223> Each "Xaa" independently represents a naturally occurring amino

acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (103)..(103)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (112)..(112)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (114)..(114)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (116)..(121)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (135)..(135)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>

<221> MISC_FEATURE

<222> (137)..(137)

<223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
 <221> MISC_FEATURE
 <222> (140)..(142)
 <223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
 <221> MISC_FEATURE
 <222> (147)..(147)
 <223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<220>
 <221> MISC_FEATURE
 <222> (159)..(159)
 <223> Each "Xaa" independently represents a naturally occurring amino acid residue except Cys, with the proviso that at least one "Xaa" in SEQ ID NO:11 is different from the corresponding amino acid residue of the native sequence.

<400> 11

Ala Asp Arg Glu Arg Ser Ile Xaa Asp Phe Cys Leu Val Ser Lys Val
 1 5 10 15

Xaa Gly Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Trp Trp Tyr Asn Val Thr
 20 25 30

Asp Gly Ser Cys Gln Leu Phe Xaa Tyr Xaa Gly Cys Xaa Xaa Xaa Ser
 35 40 45

Asn Asn Tyr Xaa Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Xaa
 50 55 60

Thr Glu Asn Ala Thr Gly Asp Leu Ser Thr Ser Arg Asn Ala Ala Asp
 65 70 75 80

Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu His Asp Ser
 85 90 95

Ser Asp Met Phe Asn Tyr Xaa Glu Tyr Cys Thr Ala Asn Ala Val Xaa
 100 105 110

Gly Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Trp Tyr Phe Asp Val Glu Arg
 115 120 125

Asn Ser Cys Asn Asn Phe Xaa Tyr Xaa Gly Cys Xaa Xaa Xaa Lys Asn
 130 135 140

Ser Tyr Xaa Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg Xaa Gln
 145 150 155 160

Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly
 165 170 175

Ala Val Ser

<210> 12
 <211> 393
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (361)..(361)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (367)..(367)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (384)..(384)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (390)..(390)
 <223> "n" is any nucleotide.

<400> 12
 ggccggggtcg tttctcgcct ggctgggatac gctgctcttc tctgggggtcc tggccgggccc 60
 accgagaacg cagcatccac gacttctgcc tgggtgtcgaa ggtgggtgggc agattccggg 120
 cctccatgcc taggtggtgg tacaatgtca ctgacggatac ctgccagctg tttgtgtatg 180

```

ggggctgtga cggaaacagc aataattacc tgaccaagga ggagtgcctc aagaaatgtg      240
ccactgtcac agagaatgcc acgggtgacc tggccaccag caggaatgca gcggattcct      300
ctgtcccaag tgctcccaga aggcaggatt cttgaagacc acttcagcga tatgtttcaa      360
ntattgnaag aataattgca ccgncaacgn att                                     393

```

```

<210> 13
<211> 110
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> SIGNAL
<222> (1)..(18)
<223>

```

```

<400> 13

```

```

Pro Gly Arg Phe Ser Pro Gly Trp Asp Arg Cys Ser Ser Leu Gly Ser
1              5              10              15

```

```

Trp Pro Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser
          20              25              30

```

```

Lys Val Val Gly Arg Phe Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn
          35              40              45

```

```

Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly
50              55              60

```

```

Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala
65              70              75              80

```

```

Thr Val Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala
          85              90              95

```

```

Ala Asp Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser
          100              105              110

```

```

<210> 14
<211> 510
<212> DNA
<213> Homo sapiens

```

```

<220>

```

<221> misc_feature
 <222> (424)..(424)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (481)..(481)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (509)..(509)
 <223> "n" is any nucleotide.

<400> 14
 gcaataatta cctgaccaag gaggagtgcc tcaagaaatg tgccactgtc acagagaatg 60
 ccacgggtga cctggccacc agcaggaatg cagcggattc ctctgtccca agtctcccag 120
 aaggcaggat tctgaagacc actccagcga tatgttcaac tatgaagaat actgcaccgc 180
 caacgcagtc actgggcctt gccgtgcac cttcccacgc tggactttg acgtggagag 240
 gaactcctgc aataacttca tctatggagg ctgccggggc aataagaaca gctaccgctc 300
 tgaggaggcc tgcattgctcc gctgcttccg ccagcaggag aatcctcccc tgccccttgg 360
 ctcaaaggtg gtggttcttg ccggggctgt ttcgtgatgg tggtgatcct tttcctgggg 420
 agcntccatg gtcttactga ttccgggtgg caaggaggaa ccaggagcgt gccctgcgga 480
 ncgtctggag cttcggagat gacaaggnt 510

<210> 15
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 15

Leu Pro Asp Gln Gly Gly Val Pro Gln Glu Met Cys His Cys His Arg
 1 5 10 15

Glu Cys His Gly
 20

<210> 16
 <211> 427
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (3)..(3)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (11)..(12)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (17)..(17)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (48)..(48)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (425)..(425)
 <223> "n" is any nucleotide.

<400> 16
 gcngcgcggt nntcgcntgc tgggatcgct gcacctctct ggggtcgngg cggccgaccg 60
 agaacgcagc atccacgact tctgcctggt gtcgaagggt gtgggcagat gccgggcctc 120
 catgcctagg tggtggtaca atgtcactga cggatcctgc cagctgtttg tgtatggggg 180
 ctgtgacgga aacagcaata attacctgac caaggaggag tgcctcaaga aatgtgccac 240
 tgtcacagag aatgccacgg gtgacctggc caccagcagg aatgcagcgg attcctctgt 300
 cccaagtgct ccagaaggc aggattctga agaccactcc agcgatatgt tcaactatga 360
 agaatactgg caccgccaac gcattcactg ggcctgcgtg catccttccc acgctgggtac 420
 tttgncg 427

<210> 17
 <211> 423
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (6)..(6)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (401)..(401)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (407)..(407)
 <223> "n" is any nucleotide.

<400> 17
 tgggantcgc tgctcctctc tggggtcctg gcggccgacc gagaacgcag catccacgac 60
 ttctgcctgg tgtcgaaggt ggtgggcaga tgccgggcct ccatgcctag gtggtggtac 120
 aatgtcactg acggatcctg ccagctgttt gtgtatgggg gctgtgacgg aaacagcaat 180
 aattacctga ccaaggagga gtgcctcaag aaatgtgcca ctgtcacaga gaatgccacg 240
 ggtgacctgg ccaccagcag gaatgcagcg gattcctctg tccaagtgc tcccagaagg 300
 caggattctg aagaccactc cagcgatatg ttcaactatg aagaatactg caccgccaac 360
 gcagtcactg ggccttgctg ggaatccttt cccacgctgg naatttngac gttgagaagg 420
 aac 423

<210> 18
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of tissue factor pathway inhibitor precursor 1.

<400> 18

His Ser Phe Cys Ala Phe Lys Ala Asp Asp Gly Pro Cys Lys Ala Ile
 1 5 10 15

Met Lys Arg Phe Phe Phe Asn Ile Phe Thr Arg Gln Cys Glu Glu Phe
 20 25 30

Ile Tyr Gly Gly Cys Glu Gly Asn Gln Asn Arg Phe Glu Ser Leu Glu
 35 40 45

Glu Cys Lys Lys Met Cys Thr Arg Asp
 50 55

<210> 19
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of tissue factor pathway inhibitor precursor 1.
 <400> 19

Pro Asp Phe Cys Phe Leu Glu Glu Asp Pro Gly Ile Cys Arg Gly Tyr
 1 5 10 15

Ile Thr Arg Tyr Phe Tyr Asn Asn Gln Thr Lys Gln Cys Glu Arg Phe
 20 25 30

Lys Tyr Gly Gly Cys Leu Gly Asn Met Asn Asn Phe Glu Thr Leu Glu
 35 40 45

Glu Cys Lys Asn Ile Cys Glu Asp Gly
 50 55

<210> 20
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of tissue factor pathway inhibitor precursor.
 <400> 20

Pro Ser Trp Cys Leu Thr Pro Ala Asp Arg Gly Leu Cys Arg Ala Asn
 1 5 10 15

Glu Asn Arg Phe Tyr Tyr Asn Ser Val Ile Gly Lys Cys Arg Pro Phe
 20 25 30

Lys Tyr Ser Gly Cys Gly Gly Asn Glu Asn Asn Phe Thr Ser Lys Gln
 35 40 45

Glu Cys Leu Arg Ala Cys Lys Lys Gly
 50 55

<210> 21
 <211> 57
 <212> PRT
 <213> Unknown

<220>

<223> Kunitz-like domain of tissue factor pathway inhibitor precursor 2.

<400> 21

Ala Glu Ile Cys Leu Leu Pro Leu Asp Tyr Gly Pro Cys Arg Ala Leu
 1 5 10 15

Leu Leu Arg Tyr Tyr Tyr Arg Tyr Arg Thr Gln Ser Cys Arg Gln Phe
 20 25 30

Leu Tyr Gly Gly Cys Glu Gly Asn Ala Asn Asn Phe Tyr Thr Trp Glu
 35 40 45

Ala Cys Asp Asp Ala Cys Trp Arg Ile
 50 55

<210> 22

<211> 57

<212> PRT

<213> Unknown

<220>

<223> Kunitz-like domain of tissue factor pathway inhibitor precursor 2.

<400> 22

Pro Ser Phe Cys Tyr Ser Pro Lys Asp Glu Gly Leu Cys Ser Ala Asn
 1 5 10 15

Val Thr Arg Tyr Tyr Phe Asn Pro Arg Tyr Arg Thr Cys Asp Ala Phe
 20 25 30

Thr Tyr Thr Gly Cys Gly Gly Asn Asp Asn Asn Phe Val Ser Arg Glu
 35 40 45

Asp Cys Lys Arg Ala Cys Ala Lys Ala
 50 55

<210> 23

<211> 57

<212> PRT

<213> Unknown

<220>

<223> Kunitz-like domain of amyloid precursor protein homologue.

<400> 23

Lys Ala Val Cys Ser Gln Glu Ala Met Thr Gly Pro Cys Arg Ala Val
 1 5 10 15

Met Pro Arg Thr Thr Phe Asp Leu Ser Lys Gly Lys Cys Val Arg Phe
 20 25 30

Ile Thr Gly Gly Cys Gly Gly Asn Arg Asn Asn Phe Glu Ser Glu Asp
 35 40 45

Tyr Cys Met Ala Val Cys Lys Ala Met
 50 55

<210> 24
 <211> 58
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of aprotinin.

<400> 24

Arg Pro Asp Phe Cys Leu Glu Pro Pro Tyr Thr Gly Pro Cys Lys Ala
 1 5 10 15

Arg Ile Ile Arg Tyr Phe Tyr Asn Ala Lys Ala Gly Leu Cys Gln Thr
 20 25 30

Phe Val Tyr Gly Gly Cys Arg Ala Lys Arg Asn Asn Phe Lys Ser Ala
 35 40 45

Glu Asp Cys Met Arg Thr Cys Gly Gly Ala
 50 55

<210> 25
 <211> 51
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of inter-alpha-trypsin inhibitor precursor.

<400> 25

Cys Gln Leu Gly Tyr Ser Ala Gly Pro Cys Met Gly Met Thr Ser Arg
 1 5 10 15

Tyr Phe Tyr Asn Gly Thr Ser Met Ala Cys Glu Thr Phe Gln Tyr Gly
 20 25 30

Gly Cys Met Gly Asn Gly Asn Asn Phe Val Thr Glu Lys Glu Cys Leu
 35 40 45

Gln Thr Cys
 50

<210> 26
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of inter-alpha-trypsin inhibitor precursor.

<400> 26

Val Ala Ala Cys Asn Leu Pro Ile Val Arg Gly Pro Cys Arg Ala Phe
 1 5 10 15

Ile Gln Leu Trp Ala Phe Asp Ala Val Lys Gly Lys Cys Val Leu Phe
 20 25 30

Pro Tyr Gly Gly Cys Gln Gly Asn Gly Asn Lys Phe Tyr Ser Glu Lys
 35 40 45

Glu Cys Arg Glu Tyr Cys Gly Val Pro
 50 55

<210> 27
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of amyloid precursor protein.

<400> 27

Glu Val Cys Cys Ser Glu Gln Ala Glu Thr Gly Pro Cys Arg Ala Met
 1 5 10 15

Ile Ser Arg Trp Tyr Phe Asp Val Thr Glu Gly Lys Cys Ala Pro Phe
 20 25 30

Phe Tyr Gly Gly Cys Gly Gly Asn Arg Asn Asn Phe Asp Thr Glu Glu

35

40

45

Tyr Cys Met Ala Val Cys Gly Ser Ala
 50 55

<210> 28
 <211> 51
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of collagen alpha-3(VI) precursor.

<400> 28

Cys Lys Leu Pro Lys Asp Glu Gly Thr Cys Arg Asp Phe Ile Leu Lys
 1 5 10 15

Trp Tyr Tyr Asp Pro Asn Thr Lys Ser Cys Ala Arg Phe Trp Tyr Gly
 20 25 30

Gly Cys Gly Gly Asn Glu Asn Lys Phe Gly Ser Gln Lys Glu Cys Glu
 35 40 45

Lys Val Cys
 50

<210> 29
 <211> 57
 <212> PRT
 <213> Unknown

<220>
 <223> Kunitz-like domain of HKI-B9.

<400> 29

Pro Asn Val Cys Ala Phe Pro Met Glu Lys Gly Pro Cys Gln Thr Tyr
 1 5 10 15

Met Thr Arg Trp Phe Phe Asn Phe Glu Thr Gly Glu Cys Glu Leu Phe
 20 25 30

Ala Tyr Gly Gly Cys Gly Gly Asn Ser Asn Asn Phe Leu Arg Lys Glu
 35 40 45

Lys Cys Glu Lys Phe Cys Lys Phe Thr
 50 55

<210> 30
 <211> 46
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 5' sense oligonucleotide used in Example 6.

<400> 30
 gccaaagcttg gataaaagat atgaagaata ctgcaccgcc aacgca 46

<210> 31
 <211> 35
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 3' antisense oligonucleotide used in Example 6.

<400> 31
 ggggatcctc actgctggcg gaagcagcgg agcat 35

<210> 32
 <211> 206
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Cloned bikunin cDNA fragment in Example 6.

<400> 32
 ccaagcttgg ataaaagata tgaagaatac tgcaccgccca acgcagtcac tgggccttgc 60
 cgtgcatacct tcccacgctg gtactttgac gtggagagga actcctgcaa taacttcac 120
 tatggaggct gccggggcaa taagaacagc taccgctctg aggaggcctg catgctccgc 180
 tgcttccgcc agcagtgagg atcccc 206

<210> 33
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 3' PCR primer used to amplify EST R74593.

<400> 33
 cgaagcttca tctccgaagc tccagacg 28

<210> 34

<211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 5' PCR primer used to amplify EST R74593.

<400> 34
 aggatctaga caataattac ctgaccaagg a

31

<210> 35
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 5' PCR primer used to amplify EST R35464.

<400> 35
 ggtctagagg ccgggtcggtt tctcgccctgg ctggga

36

<210> 36
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 5' PCR primer used to amplify EST R34808.

<400> 36
 cacctgatcg cgagacccc

19

<210> 37
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Vector specific DNA sequencing primer (SP6).

<400> 37
 gatttaggtg acactatag

19

<210> 38
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Vector specific DNA sequencing primer (T7).

<400> 38
 taatacgact cactataggg

20

<210> 39
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Gene specific DNA sequencing primer.

<400> 39
 ttacctgacc aaggaggagt gc 22

<210> 40
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Gene specific DNA sequencing primer.

<400> 40
 aatccgctgc attcctgctg gtg 23

<210> 41
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Gene specific DNA sequencing primer.

<400> 41
 cagtcactgg gccttgccgt 20

<210> 42
 <211> 105
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> 5' sense oligonucleotide used in Example 5.

<400> 42
 gaaggggtaa gcttggataa aagatatgaa gaatactgca ccgccaacgc agtcactggg 60
 ccttgccgtg catccttccc acgctggtac tttgacgtgg agagg 105

<210> 43
 <211> 129
 <212> DNA
 <213> Artificial Sequence

<220>

<223> 3' antisense oligonucleotide used in Example 5.

<400> 43

cgcggatccc tactggcgga agcagcggag catgcaggcc tcctcagagc ggtagctggt 60

cttattgccc cggcagcctc catagatgaa gttattgcag gagttcctct ccacgtcaaa 120

gtaccagcg 129

<210> 44

<211> 207

<212> DNA

<213> Artificial Sequence

<220>

<223> Cloned bikunin fragment in Example 5.

<400> 44

gaaggggtaa gcttggataa aagatatgaa gaatactgca ccgccaacgc agtcactggg 60

ccttgccgtg catccttccc acgctggtac tttgacgtgg agaggaactc ctgcaataac 120

ttcatctatg gaggctgccg gggcaataag aacagctacc gctctgagga ggccctgcatg 180

ctccgctgct tccgccagta gggatcc 207

<210> 45

<211> 248

<212> PRT

<213> Artificial Sequence

<220>

<223> EST derived consensus sequence of human Bikunin (Figs. 4D and 4G).

<220>

<221> SIGNAL

<222> (1)..(23)

<223>

<400> 45

Met Leu Arg Ala Glu Ala Asp Gly Val Ser Arg Leu Leu Gly Ser Leu
1 5 10 15Leu Leu Ser Gly Val Leu Ala Ala Asp Arg Glu Arg Ser Ile His Asp
20 25 30Phe Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg Ala Ser Met Pro
35 40 45

Arg Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr
50 55 60

Gly Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys
65 70 75 80

Leu Lys Lys Cys Ala Thr Val Thr Glu Asn Ala Thr Gly Asp Leu Ala
85 90 95

Thr Ser Arg Asn Ala Ala Asp Ser Ser Val Pro Ser Ala Pro Arg Arg
100 105 110

Gln Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn Tyr Glu Glu Tyr
115 120 125

Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala Ser Phe Pro Arg
130 135 140

Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn Phe Ile Tyr Gly
145 150 155 160

Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu Glu Ala Cys Met
165 170 175

Leu Arg Cys Phe Arg Gln Gln Glu Asn Pro Pro Leu Pro Leu Gly Ser
180 185 190

Lys Val Val Val Leu Ala Gly Leu Phe Val Met Val Leu Ile Leu Phe
195 200 205

Leu Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala Arg Arg Asn Gln
210 215 220

Glu Arg Ala Leu Arg Thr Val Trp Ser Ser Gly Asp Asp Lys Glu Gln
225 230 235 240

Leu Val Lys Asn Thr Tyr Val Leu
245

<210> 46

<211> 782

<212> DNA

<213> Homo sapiens

```

<400> 46
acctgatcgc gagaccccaa cggctggtgg cgtcgctgc gcgtctcggc tgagctggcc 60
atggcgcagc tgtgcgggct gaggcggagc cgggcgtttc tcgccctgct gggatcgctg 120
ctcctctctg gggtcctggc ggccgaccga gaacgcagca tccacgactt ctgcctggtg 180
tcgaaggtgg tgggcagatg cggggcctcc atgcctaggt ggtggtacaa tgtcactgac 240
ggatcctgcc agctgtttgt gtatgggggc tgtgacggaa acagcaataa ttacctgacc 300
aaggaggagt gcctcaagaa atgtgccact gtcacagaga atgccacggg tgacctggcc 360
accagcagga atgcagcgga ttctctctgt ccaagtgtc ccagaaggca ggattctgaa 420
gaccactcca gcgatatgtt caactatgaa gaatactgca ccgccaacgc agtcactggg 480
ccttgccgtg catccttccc acgctggtac tttgacgtgg agaggaactc ctgcaataac 540
ttcatctatg gaggctgccg gggcaataag aacagctacc gctctgagga ggctgcatg 600
ctccgctgct tccgccagca ggagaatcct cccctgcccc ttggtcaaa ggtggtgggt 660
ctggcggggc tgttcgtgat ggtgttgatc ctcttcctgg gagcctccat ggtctacctg 720
atccgggtgg cacggaggaa ccaggagcgt gccctgcgca ccgtctggag cttcggagat 780
ga 782

```

```

<210> 47
<211> 240
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> SIGNAL
<222> (1)..(27)
<223>

```

```

<400> 47

```

```

Met Ala Gln Leu Cys Gly Leu Arg Arg Ser Arg Ala Phe Leu Ala Leu
1          5          10          15

```

```

Leu Gly Ser Leu Leu Leu Ser Gly Val Leu Ala Ala Asp Arg Glu Arg
20          25          30

```

```

Ser Ile His Asp Phe Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg
35          40          45

```

```

Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln
50          55          60

```

Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr
 65 70 75 80

Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val Thr Glu Asn Ala Thr
 85 90 95

Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp Ser Ser Val Pro Ser
 100 105 110

Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn
 115 120 125

Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala
 130 135 140

Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn
 145 150 155 160

Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu
 165 170 175

Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln Glu Asn Pro Pro Leu
 180 185 190

Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly Leu Phe Val Met Val
 195 200 205

Leu Ile Leu Phe Leu Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala
 210 215 220

Arg Arg Asn Gln Glu Arg Ala Leu Arg Thr Val Trp Ser Phe Gly Asp
 225 230 235 240

<210> 48
 <211> 1544
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1358)..(1358)
 <223> "n" is any nucleotide.

<400> 48

```

gcacgagttg ggaggtgtag cgcggctctg aacgcgctga gggccggtga gtgtcgcagg      60
cggcgagggc gcgagtgagg agcagaccca ggcacgcgc gccgagaagg ccgggcgtcc      120
ccacactgaa ggtccggaaa ggcgacttcc gggggctttg gcacctggcg gaccctcccg      180
gagcgtcggc acctgaacgc gaggcgctcc attgcgcgtg cgcgttgagg ggcttcccgc      240
acctgatcgc gagaccccaa cggctgggtg cgtcgcctgc gcgtctcggc tgagctggcc      300
atggcgcagc tgtgcgggct gaggcggagc cgggcgtttc tcgccttgcg gggatcgcgtg      360
ctcctctctg gggtcctggc ggccgaccga gaacgcagca tccacgactt ctgcctgggtg      420
tcgaagggtg tgggcagatg ccgggcctcc atgcctaggt ggtggtacaa tgtcactgac      480
ggatcctgcc agctgtttgt gtatgggggc tgtgacggaa acagcaataa ttacctgacc      540
aaggaggagt gcctcaagaa atgtgccact gtcacagaga atgccacggg tgacctggcc      600
accagcagga atgcagcggg ttctctctgt ccaagtgtc ccagaaggca ggattctgaa      660
gaccactcca gcgatatgtt caactatgaa gaatactgca ccgccaacgc agtcactggg      720
ccttgccgtg catccttccc acgctggtac tttgacgtgg agaggaaact ctgcaataac      780
ttcatctatg gaggtgccc gggcaataag aacagctacc gctctgagga ggcctgcatg      840
ctccgctgct tcgccagca ggagaatcct ccctgcccc ttggctcaaa ggtggtggtt      900
ctggcggggc tgttcgtgat ggtgttgatc ctcttcttgg gagcctccat ggtctacctg      960
atccgggttg cacggaggaa ccaggagcgt gccttgcgca ccgtctggag ctccggagat     1020
gacaaggagc agctggtgaa gaacacatat gtctgtgac cgccctgtcg ccaagaggac     1080
tggggaaggg aggggagact atgtgtgagc tttttttaa tagagggatt gactcggatt     1140
tgagtgatca ttagggctga ggtctgtttc tctgggaggt aggacggctg cttcctggtc     1200
tggcagggat gggtttgctt tggaaatcct ctaggaggct cctcctcgca tggcctgcag     1260
tctggcagca gccccgagtt gtttctcgc tgatcgatct ctttctcca ggtagagttt     1320
tctttgctta tggtgaattc cattgcctcc ttttctcnat cacagaagtg atgttggaat     1380
cgtttctttt gtttgtctga tttatggttt ttttaagtat aaacaaaagt tttttattag     1440
cattctgaaa gaaggaaagt aaaatgtaca agtttaataa aaaggggcct tcccctttag     1500
aataaatttc cagcatgttg ctttcaaaaa aaaaaaaaaa aaaa                        1544

```

<210> 49

<211> 252

<212> . PRT

<213> Homo sapiens

<220>

<221> SIGNAL

<222> (1)..(27)

<223>

<400> 49

Met Ala Gln Leu Cys Gly Leu Arg Arg Ser Arg Ala Phe Leu Ala Leu
1 5 10 15

Leu Gly Ser Leu Leu Leu Ser Gly Val Leu Ala Ala Asp Arg Glu Arg
20 25 30

Ser Ile His Asp Phe Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg
35 40 45

Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln
50 55 60

Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr
65 70 75 80

Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val Thr Glu Asn Ala Thr
85 90 95

Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp Ser Ser Val Pro Ser
100 105 110

Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn
115 120 125

Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala
130 135 140

Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn
145 150 155 160

Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu
165 170 175

Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln Glu Asn Pro Pro Leu
180 185 190

Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly Leu Phe Val Met Val
 195 200 205

Leu Ile Leu Phe Leu Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala
 210 215 220

Arg Arg Asn Gln Glu Arg Ala Leu Arg Thr Val Trp Ser Ser Gly Asp
 225 230 235 240

Asp Lys Glu Gln Leu Val Lys Asn Thr Tyr Val Leu
 245 250

<210> 50
 <211> 146
 <212> PRT
 <213> Homo sapiens

<400> 50

Cys Leu Val Ser Lys Val Val Gly Arg Cys Arg Ala Ser Met Pro Arg
 1 5 10 15

Trp Trp Tyr Asn Val Thr Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly
 20 25 30

Gly Cys Asp Gly Asn Ser Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu
 35 40 45

Lys Lys Cys Ala Thr Val Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr
 50 55 60

Ser Arg Asn Ala Ala Asp Ser Ser Val Pro Ser Ala Pro Arg Arg Gln
 65 70 75 80

Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys
 85 90 95

Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp
 100 105 110

Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly
 115 120 125

Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu

130

135

140

Arg Cys
145

<210> 51
<211> 1530
<212> DNA
<213> Artificial Sequence

<220>
<223> Consensus bikunin sequence of Fig. 4C.

<220>
<221> misc_feature
<222> (46)..(46)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (117)..(117)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (313)..(313)
<223> "n" is any nucleotide.

<400> 51
gcgacctccg cgcgttggga ggtgtagcgc ggctctgaac gcgtgnaggg ccgttgagtg 60
tcgcaggcgg cgagggcgcg agtgaggagc agaccaggc atcgcgcgcc gagaagncgg 120
gcgtcccccac actgaaggtc cggaaaggcg acttccgggg gctttggcac ctggcggacc 180
ctcccggagc gtcggcacct gaacgcgagg cgctccattg cgcgtgcgtt tgaggggctt 240
cccgcacctg atcgcgagac cccaacggct ggtggcgctc ctgcgcgtct cggctgagct 300
ggccatggcg cantgttgcg ggctgaggcg gacggcgttt ctgcctgct gggatcgctg 360
ctcctctctg gggtcctggc ggccgaccga gaacgcagca tccacgactt ctgcctggtg 420
tcgaaggtgg tgggcagatg ccgggcctcc atgcctaggt ggtggtacaa tgtcactgac 480
ggatcctgcc agctgtttgt gtatgggggc tgtgacggaa acagcaataa ttacctgacc 540
aaggaggagt gcctcaagaa atgtgccact gtcacagaga atgccacggg tgacctggcc 600
accagcagga atgcagcgga ttctctgtc ccaagtgtc ccagaaggca ggattctgaa 660
gaccactcca gcgatatgtt caactatgaa gaatactgca ccgccaacgc agtcactggg 720

```

ccttgccgtg catccttccc acgctggtac tttgacgtgg agaggaactc ctgcaataac 780
ttcatctatg gaggctgccg gggcaataag aacagctacc gctctgagga ggcctgcatg 840
ctccgctgct tccgccagca ggagaatcct cccctgcccc ttggctcaaa ggtggtggtt 900
ctggcgggggc tgttcgtgat ggtgttgatc ctcttcctgg gagcctccat ggtctacctg 960
atccgggtgg cacggaggaa ccaggagcgt gccctgcgca ccgtctggag ctccggagat 1020
gacaaggagc agctggtgaa gaacacatat gtcctgtgac cgccctgtcg ccaagaggac 1080
tggggaaggg aggggagact atgtgtgagc tttttttaa tagagggatt gactcggatt 1140
tgagtgatca ttagggctga ggtctgtttc tctgggaggt aggacggctg cttcctggtc 1200
tggcagggat gggtttgctt tggaaatcct ctaggaggct cctcctcgca tggcctgcag 1260
tctggcagca gccccgagtt gtttcctcgc tgatcgattt ctttcctcca ggtagagttt 1320
tctttgctta tggtgaattc cattgcctct tttctcatca cagaagtgat gttggaatcg 1380
tttcttttgt ttgtctgatt tatggttttt ttaagtataa aaaaagtgtt tttattagca 1440
ttctgaaaga aggaaagtaa aatgtacaag ttaataaaa aggggccttc ccctttagaa 1500
taaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1530

```

```

<210> 52
<211> 170
<212> PRT
<213> Homo sapiens

```

```

<400> 52

```

```

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val
1          5          10          15

```

```

Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr
          20          25          30

```

```

Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser
          35          40          45

```

```

Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val
          50          55          60

```

```

Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp
65          70          75          80

```

Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser
85 90 95

Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr
100 105 110

Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg
115 120 125

Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn
130 135 140

Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln
145 150 155 160

Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys
165 170

<210> 53
<211> 27
<212> PRT
<213> Homo sapiens

<400> 53

Met Ala Gln Leu Cys Gly Leu Arg Arg Ser Arg Ala Phe Leu Ala Leu
1 5 10 15

Leu Gly Ser Leu Leu Leu Ser Gly Val Leu Ala
20 25

<210> 54
<211> 23
<212> PRT
<213> Homo sapiens

<400> 54

Met Leu Arg Ala Glu Ala Asp Gly Val Ser Arg Leu Leu Gly Ser Leu
1 5 10 15

Leu Leu Ser Gly Val Leu Ala
20

<210> 55
<211> 102
<212> DNA

<213> Artificial Sequence

<220>

<223> 5' sense oligonucleotide used for construct #2 in Example 5.

<400> 55

gaaggggtaa gcttggataa aagagaagaa tactgtactg ctaatgctgt tactgggtcca 60

tgtagagctt cttttccaag atggtacttt gatgttgaaa ga 102

<210> 56

<211> 129

<212> DNA

<213> Artificial Sequence

<220>

<223> 3' antisense oligonucleotide used for construct #2 in Example 5.

<400> 56

actggatcct cattggcgaa aacatctcaa catacaggct tcttcagatc tgtaagaatt 60

tttattacct ctacaaccac cgtaaataaa attattacaa gaatttcttt caacatcaaa 120

gtaccatct 129

<210> 57

<211> 108

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' sense oligonucleotide used for construct #3 in Example 5.

<400> 57

gaaggggtaa gcttggataa aagaaattac gaagaatact gtactgctaa tgctgttact 60

gggccatgta gagcttcttt tccaagatgg tactttgatg ttgaaaga 108

<210> 58

<211> 117

<212> DNA

<213> Artificial Sequence

<220>

<223> 5' sense oligonucleotide used for construct #4 in Example 5.

<400> 58

gaaggggtaa gcttggataa aagagatatg tttaattacg aagaatactg tactgctaat 60

gctgttactg gtccatgtag agcttctttt ccaagatggg actttgatgt tgaaaga 117

<210> 59

<211> 19

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Sense oligonucleotide used in PCR in Example 8.

<400> 59
 cacctgatcg cgagacccc 19

<210> 60
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Antisense oligonucleotide used in PCR in Example 8.

<400> 60
 ctggcggaag cagcggagca tgc 23

<210> 61
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide used in in vitro mutagenesis in Example 9.

<400> 61
 cgcgtctcgg ctgacctggc cctgcagatg gcgcacgtgt gcggg 45

<210> 62
 <211> 60
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide used in in vitro mutagenesis in Example 9.

<400> 62
 ctgccccttg gctcaaagta ggaagatctt ccccccgggg ggggtggttct ggcgggggctg 60

<210> 63
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 63

Leu Arg Cys Phe Arg Gln Gln Glu Asn Pro Pro Pro Leu Gly
 1 5 10

<210> 64
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 64

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val
 1 5 10 15

Val Gly Arg Cys
 20

<210> 65
 <211> 20
 <212> PRT
 <213> Homo sapiens

<400> 65

Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys
 1 5 10 15

Arg Ala Ser Phe
 20

<210> 66
 <211> 11
 <212> PRT
 <213> Homo sapiens

<400> 66

Pro Arg Tyr Val Asp Gly Ser Gln Phe Tyr Gly
 1 5 10

<210> 67
 <211> 55
 <212> PRT
 <213> Homo sapiens

<400> 67

Val Val Val Leu Ala Gly Leu Phe Val Met Val Leu Ile Leu Phe Leu
 1 5 10 15

Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala Arg Arg Asn Gln Glu
 20 25 30

Arg Ala Leu Arg Thr Val Trp Ser Ser Gly Asp Asp Lys Glu Gln Leu

35

40

45

Val Lys Asn Thr Tyr Val Leu
 50 55

<210> 68
 <211> 43
 <212> PRT
 <213> Homo sapiens

<400> 68

Val Val Val Leu Ala Gly Leu Phe Val Met Val Leu Ile Leu Phe Leu
 1 5 10 15

Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala Arg Arg Asn Gln Glu
 20 25 30

Arg Ala Leu Arg Thr Val Trp Ser Phe Gly Asp
 35 40

<210> 69
 <211> 55
 <212> PRT
 <213> Homo sapiens

<400> 69

Val Val Val Leu Ala Gly Leu Phe Val Met Val Leu Ile Leu Phe Leu
 1 5 10 15

Gly Ala Ser Met Val Tyr Leu Ile Arg Val Ala Arg Arg Asn Gln Glu
 20 25 30

Arg Ala Leu Arg Thr Val Trp Ser Ser Gly Asp Asp Lys Glu Gln Leu
 35 40 45

Val Lys Asn Thr Tyr Val Leu
 50 55

<210> 70
 <211> 213
 <212> PRT
 <213> Homo sapiens

<400> 70

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val

1	5	10	15
Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr	20	25	30
Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser	35	40	45
Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val	50	55	60
Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp	65	70	75
Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser	85	90	95
Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr	100	105	110
Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg	115	120	125
Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn	130	135	140
Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln	145	150	155
Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly	165	170	175
Leu Phe Val Met Val Leu Ile Leu Phe Leu Gly Ala Ser Met Val Tyr	180	185	190
Leu Ile Arg Val Ala Arg Arg Asn Gln Glu Arg Ala Leu Arg Thr Val	195	200	205
Trp Ser Phe Gly Asp	210		

<210> 71
 <211> 225

<212> PRT

<213> Homo sapiens

<400> 71

Ala Asp Arg Glu Arg Ser Ile His Asp Phe Cys Leu Val Ser Lys Val
 1 5 10 15

Val Gly Arg Cys Arg Ala Ser Met Pro Arg Trp Trp Tyr Asn Val Thr
 20 25 30

Asp Gly Ser Cys Gln Leu Phe Val Tyr Gly Gly Cys Asp Gly Asn Ser
 35 40 45

Asn Asn Tyr Leu Thr Lys Glu Glu Cys Leu Lys Lys Cys Ala Thr Val
 50 55 60

Thr Glu Asn Ala Thr Gly Asp Leu Ala Thr Ser Arg Asn Ala Ala Asp
 65 70 75 80

Ser Ser Val Pro Ser Ala Pro Arg Arg Gln Asp Ser Glu Asp His Ser
 85 90 95

Ser Asp Met Phe Asn Tyr Glu Glu Tyr Cys Thr Ala Asn Ala Val Thr
 100 105 110

Gly Pro Cys Arg Ala Ser Phe Pro Arg Trp Tyr Phe Asp Val Glu Arg
 115 120 125

Asn Ser Cys Asn Asn Phe Ile Tyr Gly Gly Cys Arg Gly Asn Lys Asn
 130 135 140

Ser Tyr Arg Ser Glu Glu Ala Cys Met Leu Arg Cys Phe Arg Gln Gln
 145 150 155 160

Glu Asn Pro Pro Leu Pro Leu Gly Ser Lys Val Val Val Leu Ala Gly
 165 170 175

Leu Phe Val Met Val Leu Ile Leu Phe Leu Gly Ala Ser Met Val Tyr
 180 185 190

Leu Ile Arg Val Ala Arg Arg Asn Gln Glu Arg Ala Leu Arg Thr Val
 195 200 205

Trp Ser Ser Gly Asp Asp Lys Glu Gln Leu Val Lys Asn Thr Tyr Val
 210 215 220

Leu
 225

<210> 72
 <211> 19
 <212> PRT
 <213> Homo sapiens

<220>
 <221> MISC_FEATURE
 <222> (9)..(9)
 <223> "Xaa" is Ile, Thr, Asn, or Ser.

<220>
 <221> MISC_FEATURE
 <222> (11)..(11)
 <223> "Xaa" is Val, Ala, Glu, or Gly.

<220>
 <221> MISC_FEATURE
 <222> (17)..(17)
 <223> "Xaa" is Ser, Pro, Thr, or Ala.

<220>
 <221> MISC_FEATURE
 <222> (19)..(19)
 <223> "Xaa" is Tyr, His, Asn, or Asp.

<400> 72

Arg Pro Leu Gln Arg Tyr Val Ser Xaa Ile Xaa Arg Ile Ile Ala Pro
 1 5 10 15

Xaa Thr Xaa

<210> 73
 <211> 108
 <212> PRT
 <213> Homo sapiens

<400> 73

Pro Gly His Gln Gln Glu Cys Ser Gly Phe Leu Cys Pro Lys Ser Pro
 1 5 10 15

Arg Arg Gln Asp Ser Glu Asp His Ser Ser Asp Met Phe Asn Tyr Glu
20 25 30

Glu Tyr Cys Thr Ala Asn Ala Val Thr Gly Pro Cys Arg Ala Ser Phe
35 40 45

Pro Arg Trp Tyr Phe Asp Val Glu Arg Asn Ser Cys Asn Asn Phe Ile
50 55 60

Tyr Gly Gly Cys Arg Gly Asn Lys Asn Ser Tyr Arg Ser Glu Glu Ala
65 70 75 80

Cys Met Leu Arg Cys Phe Arg Gln Gln Glu Asn Pro Pro Leu Pro Leu
85 90 95

Gly Ser Lys Val Val Val Leu Ala Gly Ala Val Ser
100 105

<210> 74
<211> 31
<212> PRT
<213> Homo sapiens

<220>
<221> MISC_FEATURE
<222> (25)..(25)
<223> "Xaa" is Asp or Glu.

<400> 74

Ser Phe Ser Trp Gly Ala Ser Met Val Leu Leu Ile Pro Gly Gly Lys
1 5 10 15

Glu Glu Pro Gly Ala Cys Pro Ala Xaa Arg Leu Glu Leu Arg Arg
20 25 30

<210> 75
<211> 511
<212> DNA
<213> Artificial Sequence

<220>
<223> Corrected version of EST R74593 (Fig. 3).

<220>
<221> misc_feature
<222> (425)..(425)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (482)..(482)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (510)..(510)

<223> "n" is any nucleotide.

<400> 75

```

gcaataatta cctgaccaag gaggagtgcc tcaagaaatg tgccactgtc acagagaatg      60
ccacgggtga cctggccacc agcaggaatg cagcggattc ctctgtccca agtgctccca      120
gaaggcagga ttctgaagac cactccagcg atatgttcaa ctatgaagaa tactgcaccg      180
ccaacgcagt cactgggcct tgccgtgcat ccttcccacg ctggtacttt gacgtggaga      240
ggaactcctg caataacttc atctatggag gctgccgggg caataagaac agctaccgct      300
ctgaggagggc ctgcatgctc cgctgcttcc gccagcagga gaatcctccc ctgccccttg      360
gctcaaaggt ggtgggttctg gccgggggctg ttctgtgatg gtgttgatcc ttttcctggg      420
gagcntccat ggtcttactg attccgggtg gcaaggagga accaggagcg tgccctgcgg      480
ancgtctgga gcttcggaga tgacaagggn t                                     511

```

<210> 76

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Amino acids 184-214 of the translation of the consensus DNA sequence in Fig. 3.

<220>

<221> MISC_FEATURE

<222> (25)..(25)

<223> "Xaa" is Asp or Glu.

<400> 76

```

Ser Phe Ser Trp Gly Ala Ser Met Val Leu Leu Ile Pro Gly Gly Lys
1           5           10           15

```

```

Glu Glu Pro Gly Ala Cys Pro Ala Xaa Arg Leu Glu Leu Arg Arg

```

20

25

30

<210> 77
 <211> 312
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (45)..(45)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (49)..(49)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (118)..(118)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (231)..(231)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (305)..(305)
 <223> "n" is any nucleotide.

<400> 77
 gcgacctcgc gcggttggga ggtgtagcgc ggctctgaac gcgtngagng gccgttgagt 60
 gtcgcaggcg gcgagggcgc gagtgaggag cagacccagg catcgcgcg gcagaagncg 120
 ggcgtoccca cactgaaggt ccggaaaggc gacttccggg ggctttggca cctggcggac 180
 cctcccggag cgtcggcacc tgaacgcgag gcgctccatt gcgcgtgcgt ntgaggggct 240
 tcccgcacct gatcgcgaga ccccaacggc tggtggcgtc gcctgcgcgt ctgggctgag 300
 ctggncatgt cg 312

<210> 78
 <211> 330
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (117)..(117)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (123)..(123)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (321)..(321)
 <223> "n" is any nucleotide.

<400> 78
 gcgacctccg cgcgttggga ggtgtagcgc ggctctgaac gcgtgcaggg ccgttgagtg 60
 tcgcaggcgg cgagggcgcg agtgaggagc agaccaggc atcgcgcgcc gagaagncgg 120
 gcntccccac actgaaggtc cggaaggcg acttccgggg gctttggcac ctggcggacc 180
 ctcccggagc gtggcacctg aacgcgaggc gctccattgc gcgtgcgttt gaggggcttc 240
 ccgcacctga tcgcgagacc ccaacggctg gtggcgtcgc ctgcgcgtct cggctgagct 300
 ggccatggcg cactgtgcgg ngctgaggcg 330

<210> 79
 <211> 283
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (9)..(9)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (11)..(11)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (222)..(222)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature

<222> (231)..(231)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (262)..(262)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (267)..(267)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (274)..(274)
 <223> "n" is any nucleotide.

<400> 79
 ttgagtgtng naggcggcga gggcgcgagt gaggagcaga cccagggcatc gcgcgccgag 60
 aaggccgggc gtccccacac tgaagggtccg gaaaggcgac ttccgggggc tttggcacct 120
 ggcggaccct cccggagcgt cggcacctga acgcgaggcg ctccattgcg cgtgcgtttg 180
 aggggcttcc cgcacctgat cgcgagaccc caacggctgg tngcgtcgct ncgcgtctcg 240
 gctgagcttg gccatggcgc antgttnccg gctnaggcgg acg 283

<210> 80
 <211> 423
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (44)..(44)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (46)..(46)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (76)..(76)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (114)..(114)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (187)..(187)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (268)..(268)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (309)..(309)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (317)..(317)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (332)..(332)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (370)..(370)
 <223> "n" is any nucleotide.

<400> 80
 ggcgacctcc ggcggttggg aggtgtagcg cgctctgaac gggnggggc cgttgagtgt 60
 cgcaggcggc agggcngagt gaggagcaga cccaggcatc gcgcgccgag aagncgggcg 120
 tccccacact gaaggtccgg aaaggcgact tccggggggt ttggcacctg gcggacgtcc 180
 cggagcnggc acctgaacgc gaggcgctcc attgcgcgtg cgtttgaggg gcttcccgca 240
 cctgatcgcg agaccccaac ggctggtngc gtcgctggcg cgttctcggc tgagctggcc 300
 atggcgcant gttgcgngct gaggcggacc gncgtttttc ttcgccttgc tgggattcgc 360
 ttgcttcctn tctggggggt cctggggcggc cgaccgagaa cgcagcatcc aagaattttt 420

gcc

423

<210> 81
<211> 344
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (35)..(35)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (148)..(148)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (235)..(235)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (261)..(261)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (272)..(272)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (293)..(293)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (300)..(300)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (313)..(313)
<223> "n" is any nucleotide.

<220>

<221> misc_feature
 <222> (320)..(320)
 <223> "n" is any nucleotide.

<400> 81
 ggaggagcag acccaggcat cgcgcgccga gaagncgggc gtccccacac tgaagggtccg 60
 gaaaggcgac ttccgggggc tttggcacct ggcggaccct cccggagcgt cggcacctga 120
 acgcgaggcg ctccattgcg cgtgcgtntg gaggggcttc ccgcacctga tcgcgagacc 180
 ccaacggctg gtgggcgctg ctgcgcgtct tcggctgagc tgggccatgg cgcanttggt 240
 gcgggctgag gcggacgcgg ncgttttttc gnccttgctg ggattcggtg ttnctctctn 300
 ggggttctgg ggnggccgan cgagaacgca agcattcacg attt 344

<210> 82
 <211> 253
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (56)..(56)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (137)..(137)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (145)..(145)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (159)..(159)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (233)..(233)
 <223> "n" is any nucleotide.

<400> 82
 ggaccctccc ggagcgtcgg cacctgaacg cgaggcctcc attgcggtgc gtgtgnaggg 60

gcttcccgcga cctgatcgcg agaccccaac ggctgggtggc gtcgctgcgc gtctcggctg 120
agctggccat ggcgcantgt tgcgngctga ggcggcggnc gttttctcgc ctgctgggat 180
cgctgctcct ctctgggggtc ctggcggccg accgagaacg cagcatccac gantttcttc 240
tggtgttcga agg 253

<210> 83
<211> 419
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (26)..(26)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (95)..(95)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (292)..(292)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (313)..(315)
<223> "n" is any nucleotide.

<400> 83
ttagcgcggc tctgaacgcn agaagnggcc gttgagtgtc gcaggcggcg agggcgcgag 60
tgaggagcag acccaggcat cgcgcgccga gaagncgggc gtccccacac tgaaggtccg 120
gaaaggcgac ttccgggggc tttggcacct ggcggaccct cccggagcgt cggcacctga 180
acgcgaggcg ctccattgcg cgtgcgtttg aggggcttcc cgcacctgat cgcgagaccc 240
caacggctgg tggcgtcgcc tgcgcgtctc ggctgagctg gccatggcgc antggtgcgg 300
gcttgaggcg gannngccgt ttctcgctg ctgggategc tgctcctctc tgggggtctg 360

gcggccgacc gagaacgcag catccacgac ttctgcctgg tgtcgaaggt ggtgggcag 419

<210> 84
<211> 477
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (27)..(27)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (139)..(139)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (223)..(223)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (232)..(232)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (302)..(302)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (310)..(310)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (322)..(322)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (328)..(328)
<223> "n" is any nucleotide.

<220>

<221> misc_feature
<222> (357)..(357)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (375)..(375)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (392)..(392)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (398)..(398)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (405)..(405)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (427)..(427)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (437)..(437)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (449)..(449)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (458)..(458)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (474)..(474)
<223> "n" is any nucleotide.

<400> 84

```

agacccaggg atcgcgcgcc gagaagncgg gcgtccccac actgaaggtc cggaaaggcg      60
acttccgggg gctttggcac ctggcggacc ctcccggagc gtcggcacct gaacgcgagg      120
cctccattgc cgtgcgttng aggggcttcc cggaacttga tcgcgagacc ccaacggctg      180
gtggcgtcgc tgcgcgtcct cggctgagct ggccatggcg cantggtgcc gngctgaggg      240
cggagggccg gtttctcgcc ttgctgggat cgctgctcct ctctggggtc ctggcggccg      300
ancgaagaan gcagcaatcc angaattnct gcctggtggt cgaaagttgg tgggcanatt      360
ccggggcctt catgnctaag gttggttggt anaatgtnaa ttaangattc ttgcaactgt      420
ttgtgtnatt ggggctntta aacggaaana caataatnac ctgaccaaag aagnaatt      477

```

<210> 85

<211> 393

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (361)..(361)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (367)..(367)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (384)..(384)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (390)..(390)

<223> "n" is any nucleotide.

<400> 85

```

ggccgggtcg tttctcgctt ggctgggata gctgctcctc tctgggggtcc tggccggccg      60
accgagaacg cagcatccac gacttctgcc tggtgtcgaa ggtggtgggc agattccggg      120
cctccatgcc taggtggtgg tacaatgtca ctgacggata ctgccagctg tttgtgtatg      180
ggggctgtga cggaaacagc aataattacc tgaccaagga ggagtgcctc aagaaatgtg      240

```

ccactgtcac agagaatgcc acgggtgacc tggccaccag caggaatgca gcggattcct 300
 ctgtcccaag tgctcccaga aggcaggatt cttgaagacc acttcagcga tatgtttcaa 360
 ntattgnaag aataattgca ccgncaacgn att 393

<210> 86
 <211> 428
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (3)..(3)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (11)..(12)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (17)..(17)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (48)..(48)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (425)..(425)
 <223> "n" is any nucleotide.

<400> 86
 gcngcgcgtt nntcgcntgc tgggatcgct gcacctctct ggggtcgngg cggccgaccg 60
 agaacgcagc atccacgact tctgcctggt gtcgaagggt gtgggcagat gccgggcctc 120
 catgcctagg tggtggtaca atgtcactga cggatcctgc cagctgtttg tgtatggggg 180
 ctgtgacgga aacagcaata attacctgac caaggaggag tgcctcaaga aatgtgccac 240
 tgtcacagag aatgccacgg gtgacctggc caccagcagg aatgcagcgg attcctctgt 300
 cccaagtgct ccagaaggc aggattctga agaccactcc agcgatatgt tcaactatga 360
 agaatactgg caccgccaac gcattcactg ggcctgcgtg catccttccc acgctgggtac 420

tttgncgt

428

<210> 87
 <211> 425
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (7)..(7)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (403)..(403)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (409)..(409)
 <223> "n" is any nucleotide.

<400> 87
 ctgggantcg ctgctcctct ctgggggtcct ggcggccgac cgagaacgca gcatccacga 60
 cttctgcctg gtgtcgaagg tgggtgggcag atgccggggcc tccatgccta ggtggtggta 120
 caatgtcact gacggatcct gccagctgtt tgtgtatggg ggctgtgacg gaaacagcaa 180
 taattacctg accaaggagg agtgcctcaa gaaatgtgcc actgtcacag agaatgccac 240
 ggggtgacctg gccaccagca ggaatgcagc ggattcctct gtcccaagtg ctcccagaag 300
 gcaggattct gaagaccact ccagcgatat gttcaactat gaagaatact gcaccgccaa 360
 cgcagtcact ggggccttgc gtggaatcct ttcccacgct ggnaatttng acgttgagaa 420
 ggaac 425

<210> 88
 <211> 343
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (48)..(48)
 <223> "n" is any nucleotide.

<220>

<221> misc_feature
 <222> (62)..(62)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (211)..(211)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (232)..(232)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (245)..(245)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (309)..(309)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (318)..(318)
 <223> "n" is any nucleotide.

<400> 88
 gattcggcac aggggaaaca gcaataatta cctgaccaag gaggagtncc tcaagaaatg 60
 tnccactgtc acagagaatg ccacgggtga cctggccacc agcaggaatg cagcggattc 120
 ctctgtccca agtgctccca gaaggcagga ttctgaagac cactccagcg atatgttcaa 180
 ctatgaagaa tactgcaccg ccaacgcagt ncactgggcc ttgcgtggca tnccttccca 240
 cgctngtact ttgacgtgga gaggaactcc tggcaataac ttcatttatg gaggcttgcc 300
 ggggcaatna agaacagntt accgctcttt aggaggcctg cat 343

<210> 89
 <211> 510
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (424)..(424)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (481)..(481)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (509)..(509)

<223> "n" is any nucleotide.

<400> 89

```

gcaataatta cctgaccaag gaggagtgcc tcaagaaatg tgccactgtc acagagaatg      60
ccacgggtga cctggccacc agcaggaatg cagcggattc ctctgtccca agtctcccag      120
aaggcaggat tctgaagacc actccagcga tatgttcaac tatgaagaat actgcaccgc      180
caacgcagtc actgggcctt gccgtgcac cttcccacgc tgggtactttg acgtggagag      240
gaactcctgc aataacttca tctatggagg ctgccggggc aataagaaca gctaccgctc      300
tgaggaggcc tgcattgctc gctgcttccg ccagcaggag aatcctcccc tgccccttgg      360
ctcaaaggtg gtggttcttg cgggggctgt ttcgtgatgg tgttgatcct tttcctgggg      420
agcntccatg gtcttactga ttccgggtgg caaggaggaa ccaggagcgt gccctgcgga      480
ncgtctggag ctccggagat gacaagggnt                                     510

```

<210> 90

<211> 293

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (257)..(257)

<223> "n" is any nucleotide.

<400> 90

```

gctaccgctc tgaggaggcc tgcattgctc gctgcttccg ccagcaggag aatcctcccc      60
tgccccttgg ctcaaaggtg gtggttcttg cgggggctgt cgtgatgggt ttgatcctct      120
tcctggggag cctccatggt ctacctgac cgggtggcac ggagggaacc agggagcgtg      180
ccctgcgcac cgtctgggag ctccggagat gacaagggag cagctgggtg aagaacacat      240
atgttctctg tgaccgncct gttcgccaag aggattgggg gaaggagggg gga                                     293

```

<210> 91
 <211> 282
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (19)..(19)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (147)..(147)
 <223> "n" is any nucleotide.

<400> 91
 ttccgccaaag caggaaaant cctccccctcc cccttggctc aaaggtggtg gttcctggcg 60
 gggctgttcg tgatggtggt gatccctcct tcccgggagc ctcccatggt cctaccctga 120
 tccgggtggc acggaggaac ccaggancgt gccctgcgca ccgtctggag ctccggagat 180
 gacaaggagc agctggtgaa gaacacatat gtcctgtgac cgccctgtcg ccaagaggac 240
 tggggaaggg aggggagact atgtgtgagc tttttttaaa ta 282

<210> 92
 <211> 390
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (33)..(33)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (55)..(55)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (118)..(118)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (213)..(213)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (228)..(228)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (259)..(259)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (267)..(267)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (324)..(324)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (333)..(333)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (344)..(344)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (387)..(387)
 <223> "n" is any nucleotide.

<400> 92
 gagaggaact cctgcaataa cttcatctat gnggctgcc ggggaataag aacanctacc 60
 gctctgagga ggctgcgtg ctccgtgct tccgtgtgt gttctcttcc aggccagcag 120
 gagaatcctc ccctgcccct tggctcaaag gtggtgggtc tggcggggct gttcgtgatg 180
 gtgttgatcc tcttctggg agcctccatg gtntacctga tccgggtngc acggaggaac 240
 cagggagcgt gcctgcgna ccgtctngga gctccggaga tgacaaggag cagctggtga 300
 agaacacata tgtcctgtga ccgncctgtt cgncaagagg actnggggaa aggggagggg 360

agattatgtg ttgagttttt tttaaantag

390

<210> 93
<211> 406
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (306)..(306)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (328)..(328)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (342)..(342)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (365)..(365)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (370)..(370)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (377)..(377)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (382)..(382)
<223> "n" is any nucleotide.

<220>
<221> misc_feature
<222> (402)..(402)
<223> "n" is any nucleotide.

<400> 93

gattcgggaac gaggagccgg ggcaataaga acagctaccg ctctgaggag gcctgcatgc 60
 tccgctgctt ccgccagcag gagaatcctc ccctgccctt tggctcaaag gtggtgggtc 120
 tggcgggggt gtctgtgatg gtgttgatcc tcttcctggg agcctccatg gtctacctga 180
 tccgggtggc acggaggaac caggagcgt gccctgcgca ccgtctggga gctccggaga 240
 tgacaaggga gcagctggtg aagaacacat atgttcctgt tgaccgccct gttcgccaag 300
 agggantggg ggaaggggag ggggaganta ttgttggtga gntttttttt aaaattagga 360
 ggggnttgan ttcgggnttt tnagttgatc catttagggg gntgag 406

<210> 94
 <211> 360
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (1)..(1)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (142)..(142)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (339)..(339)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (347)..(347)
 <223> "n" is any nucleotide.

<400> 94
 nggccttgca gtgctccgct gcttccgcca gcaggagaat cctccccctgc cccttggtc 60
 aaagggtggtg gttctggcgg ggctgttcgt gatggtgttg atcctcttcc tgggagcctc 120
 catggtctac ctgatccggg tngcacggag gaaccaggag cgtgccctgc gcaccgtctg 180
 gagctccgga gatgacaagg agcagctggt gaagaacaca tatgtcctgt gaccgccctg 240
 tcgccaagag gactggggaa gggaggggag actatgtgtg agcttttttt aaatagaggg 300
 attgactcgg atttgagtga tcattagggc tgaggtctnt ttctctngga ggtaggacga 360

<210> 95
 <211> 438
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (334)..(334)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (368)..(368)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (376)..(376)
 <223> "n" is any nucleotide.

<400> 95
 cggggctggt cgtgatggtg ttgatcctct tcctgggagc ctccatggtc tacctgatcc 60
 ggggtggcacg gaggaaccag gagcgtgcc tgccgaccgt ctggagctcc ggagatgaca 120
 aggagcagct ggtgaagaac acatatgtcc tgtgaccgcc ctgtcgccaa gaggactggg 180
 gaagggaggg gagactatgt gtgagctttt tttaaataga gggattgact cggatttgag 240
 tgatcattag ggctgaggtc tgtttctctg ggaggttagga cggctgcttc ctgggtcttg 300
 gcagggatgg ggtttgcttt gggaaatcct cttnggaggc tcctccttcg catgggcctt 360
 gcagtctngg cagcancccc cgagtttttt tccttcgctg atccgatttc ttttcctcca 420
 ggtaagaatt tttctttt 438

<210> 96
 <211> 448
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (108)..(108)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (261)..(261)
 <223> "n" is any nucleotide.

<400> 96

```

gggaaccagg agcgtgccct gcgcaccggt ctggagctcc ggagatgaca aggagcagct      60
ggtgaagaac acatatgtcc tgtgaccgcc ctgtcgccaa gaggactngg gaagggaggg      120
gagactatgt gtgagctttt tttaaataga gggattgact cggatttgag tgatcattag      180
ggctgaggtc tgtttctctg ggaggtagga cggctgcttc ctggtctggc agggatgggt      240
ttgctttgga gaatcctcta ngaggctcct cctcgcatgg cctgcagtct ggcagcagcc      300
ccgagttggt tcctcgctga tcgattttct tcctccaggt agagttttct ttgcttatgt      360
tgaattccat tgcctctttt ctcatcacag aagtgatggt ggaatcgttt cttttgtttt      420
gtctgattta tgggtttttt ttaagtat                                     448

```

<210> 97

<211> 331

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (20)..(20)

<223> "n" is any nucleotide.

<220>

<221> misc_feature

<222> (30)..(30)

<223> "n" is any nucleotide.

<400> 97

```

attagggctg aggtctgttn ctctgggagn taggacggct gccttcctgg tctggcaggg      60
atgggtttgc tttggaaatc ctctaggagg ctctcctcgc catggcctgc agttctgcag      120
cagccccgag ttgtttcctc gctgatcgat ttctttcctc caggtagagt tttctttgct      180
tatgttgaat tccattgcct cttttctcat cacagaagtg atgttggaat cgtttctttt      240
gtttgtctga tttatggttt ttttaagtat aaacaaaagt tttttattag cattctgaaa      300
gaaggaaagt aaaatgtaca agtttaataa a                                     331

```

<210> 98

<211> 373

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature
 <222> (45)..(45)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (102)..(102)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (105)..(105)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (159)..(159)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (174)..(174)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (213)..(213)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (337)..(337)
 <223> "n" is any nucleotide.

<400> 98
 gattgactcg gatttgagtg atcattaggg ctgaggtctg tttcnctggg aggtaggacg 60
 gctgctcccc tgggtctggca gggatggggtt tgctttggaa anccnctagg aggctcctcc 120
 tcgcatggcc tgcagtctgg cagcagcccc gagttgttnc ctcgctgac gatntctttc 180
 ccccaggtag agttttcttt gcttatgttg aantccattg cctcttttct catcacagaa 240
 gtgatgttgg aatcgtttct tttgtttgtc tgatttatgg tttttttaag tataaacaaa 300
 agttttttat tagcattctg aaagaaggaa agtaaatgt acaagtttaa taaaaagggg 360
 ccttcccctt taa 373

<210> 99
 <211> 380
 <212> DNA
 <213> Homo sapiens

<400> 99
 gattgactcg gatttggagt gatcattagg gctgaggtct gtttctctgg gaggtaggac 60
 ggctgcttcc tggctctggca gggatggggt tgctttggaa atcctctagg aggctcctcc 120
 ttgcgatggc ctgcagtctg gcagcagccc cgagttgttt cctcgctgat cgatttcttt 180
 cctccaggta gagttttctt tgcttatgtt gaattccatt gcctcttttc tcatcacaga 240
 agtgatgttg gaatcgtttc ttttgtttgt ctgatttatg gtttttttaa gtataaacia 300
 aagtttttta ttagcattct gaaagaagga aagtaaaatg tacaagttta ataaaaaggg 360
 gccttcccct ttagaataaa 380

<210> 100
 <211> 320
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (304)..(304)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (309)..(309)
 <223> "n" is any nucleotide.

<400> 100
 tctggcaggg atgggtttgc tttggaaatc ctctaggagg ctctcctcg catggcctgc 60
 agtctggcag cagcccgagt tgtttcctcg ctgatcgatt tctttcctcc aggtagagtt 120
 ttctttgctt atgttgaatt ccattgcctc ttttctcatc acagaagtga tgttggaatc 180
 gtttcttttg tttgtctgat ttatggtttt ttttaagtata aacaaaagtt ttttattagc 240
 attctgaaag aaggaaagta aaatgtacaa gttaataaaa aaggggcctt cccctttagg 300
 aatnaaaaana aaaaagggtg 320

<210> 101
 <211> 397
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (24)..(24)
 <223> "n" is any nucleotide.

<400> 101
 gattgactcg gatttgagtg atcnattagg gctgaggtct gtttctctgg gaggtaggac 60
 ggctgcttca tgggtctggca gggatggggt tgctttggaa atcctctagg aggctcctcc 120
 tcgcatggcc tgcagtctgc agcagccccg agttgtttcc tcgctgatcg atttctttcc 180
 tccaggtaga gttttctttg cttatgttga attccattgc ctcttttctc atcacagaag 240
 tgatgttgga atcgtttctt ttgtttgtct gatttatggg ttttttaagt ataaacaaaa 300
 gttttttatt agcattctga aagaaggaaa gtaaaatgta caagtttaat aaaaaggggc 360
 cttccccttt agaataaatt tcagcatgtg ctttcaa 397

<210> 102
 <211> 289
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (61)..(61)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (74)..(74)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (122)..(122)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (184)..(184)
 <223> "n" is any nucleotide.

<400> 102
 gaggtcctc ctgcgatggc ctgcagtctt ggcagcagcc ccgagttggt tectcgctga 60
 ncgatttctt tccnccaggt agagtcttct ttgcttatgt tgaattccat tgccctctttt 120
 cncatcacag aagtgatgtt ggaatcgttt cttttgtttg tctgatttat gggttttttta 180

agtntaaaca aaagtttttt attagcattc tgaaagaagg aaagtaaaat gtacaagttt 240
 aataaaaagg ggccttcccc tttagaataa aaaaaaaaaa aaaaaaaaaa 289

<210> 103
 <211> 311
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (7)..(7)
 <223> "n" is any nucleotide.

<400> 103
 cttttgnaaa tcctctagga ggctcctcct cgcattggcct gcagtctgca gcagccccga 60
 gttgttttct cgcctgatcg atttctttcc tccaggtaga gttttctttg cttatgttga 120
 attccattgc ctcttttctc atcacagaag tgatgttgga atcgtttctt ttgtttgtct 180
 gatttatggt ttttttaagt ataaacaaaa gttttttatt agcattctga aagaaggaaa 240
 gtaaaatgta caagtttaat aaaaaggggc cttccccctt agaataaatt tcagcatgtg 300
 ctttcaaaaa a 311

<210> 104
 <211> 338
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (32)..(32)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (67)..(67)
 <223> "n" is any nucleotide.

<220>
 <221> misc_feature
 <222> (136)..(136)
 <223> "n" is any nucleotide.

<400> 104
 ggtctggcag ggatggggtt gcctttggaa anctctagg aggctcctcc tcgcatggcc 60

```

tgcagtnctg gcagcagacc ccgagttggt tctctgctga tcgattttctt tacccccagg 120
tagagttttc ctttgnctta tggtgaattc cattgcctct tttactcatc acagaagtga 180
tggtggaatc gtttcttttg tttgtctgat ttatgggtttt ttttaagtata aacaaaagtt 240
ttttattagc attctgaaag aaggaaagta aaatgtacaa gtttaataaa aaggggcctt 300
cccctttaga ataaaaaaaa aaaaaaaaaa aaaaaaaaaa 338

```

```

<210> 105
<211> 343
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (13)..(13)
<223> "n" is any nucleotide.

```

```

<220>
<221> misc_feature
<222> (19)..(19)
<223> "n" is any nucleotide.

```

```

<220>
<221> misc_feature
<222> (107)..(107)
<223> "n" is any nucleotide.

```

```

<400> 105
ccctgggtcc tgncaaggna tggggtttgc tttggaaatc ctcttaggag gctcctcctc 60
gcatggcctg cagtctggca gcagccccga gttgtttcct cgctganoga tttctttcct 120
ccaggtagag ttttctttgc ttatgttgaa ttccattgcc tcttttctca tcacagaagt 180
gatgttggaa tcgtttcttt tgtttgtctg atttatggtt tttttaagta taaacaaaag 240
ttttttatta gcattctgaa agaaggaaag taaaatgtac aagtttaata aaaaggggcc 300
ttccccttta gaataaaaaa aaaaaaaaaa aaaaaaaaaa aaa 343

```

Claims:

1. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier therefore, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

MAQLCGL RRSRAFLALL GSLLL SGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS SGDDKEQLVK NTYVL 225
(SEQ ID NO.: 49),

```

```

AGSFLAWL GSLLL SGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGAVS 179
(SEQ ID NO.: 2),

```

```

MLR AEADGVSRLL GSLLL SGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS SGDDKEQLVK NTYVL 225
(SEQ ID NO.: 45),

```

MAQLCGL RRSRAFLALL GSLLLSGVLA -1

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 47),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 71),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 70),

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATV 64
 (SEQ ID NO.: 4),

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK C 61
 (SEQ ID NO.: 5),

YEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQ 159
 (SEQ ID NO.: 6),

CTANAVTGPC RASFPRWYFD VERNSCNNFI YGGCRGNKNS YRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 7),

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQ 159
 (SEQ ID NO.: 3),

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 50),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 25
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179
 (SEQ ID NO.: 1),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK 170
 (SEQ ID NO.: 52),

or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DS 92
 (SEQ ID NO.: 8).

2. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

MAQLCGL RRSRAFLALL GSLLLLSGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS SGDDKEQLVK NTYVL 225
(SEQ ID NO.: 49),

```

```

AGSFLAWL GSLLLLSGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGAVS 179
(SEQ ID NO.: 2),

```

```

MLR AEADGVSRLG GSLLLLSGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS SGDDKEQLVK NTYVL 225
(SEQ ID NO.: 45),

```

```

MAQLCGL RRSRAFLALL GSLLLLSGVLA -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS FGD 213
(SEQ ID NO.: 47),

```

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 71),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 70),

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATV 64
 (SEQ ID NO.: 4),

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK C 61
 (SEQ ID NO.: 5),

YEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQ 159
 (SEQ ID NO.: 6),

CTANAVTGPC RASFPRWYFD VERNSCNNFI YGGCRGNKNS YRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 7),

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQ 159
 (SEQ ID NO.: 3),

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 50),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 25
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179
 (SEQ ID NO.: 1),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK 170
 (SEQ ID NO.: 52),

or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DS 92
 (SEQ ID NO.: 8).

3. The use according to claim 1 or claim 2, wherein the Kunitz-type serine protease inhibitor is glycosylated.
4. The use according to claim 1 or claim 2 wherein the Kunitz-type serine protease inhibitor contains at least one intrachain cysteine-cysteine disulfide bond.
5. The use according to claim 1 or claim 2, wherein the Kunitz-type serine protease inhibitor contains at least one intrachain cysteine-cysteine disulfide bond selected from the cysteine-cysteine paired groups consisting of CYS11-CYS61, CYS20-CYS44, CYS36-CYS57, CYS106-CYS156, CYS115-CYS139, and CYS131-CYS152, wherein the cysteine residues are numbered according to the amino acid sequence of native human placental bikunin.
6. The use according to any one of claims 1 to 5, wherein the inhibitor is for administration in lung airways.
7. The use according to any one of claims 1 to 5, wherein said inhibitor is for administration directly by aerosolization.
8. The use according to any one of claims 1 to 5, wherein said composition is for administration directly as an aerosol suspension into a mammal's respiratory tract.
9. The use according to claim 8, wherein said aerosol suspension includes respirable particles ranging in size from about 1 to about 10 microns.
10. The use according to claim 8, wherein said aerosol suspension includes respirable particles ranging in size from about 1 to about 5 microns.

11. The use according to any one of claims 8 to 10, wherein said aerosol suspension is for administration to said subject by a pressure driven nebulizer.
12. The use according to any one of claims 8 to 10, wherein said aerosol suspension is for administration to said subject by an ultrasonic nebulizer.
13. The use according to any one of claims 8 to 10, wherein said aerosol suspension is for administration to said subject by a non-toxic propellant.
14. The use according to claim 1, wherein said physiologically acceptable carrier is a member selected from the group consisting of a physiologically buffered solution; an isotonic saline, normal saline, and combinations thereof.
15. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

                                MAQLCGL RRSRAFLALL GSLLLSGVLA   -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS SGDDKEQLVK NTYVL                                225
(SEQ ID NO.: 49) .

```

16. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

                                MAQLCGL RRSRAFLALL GSLLLLSGVLA   -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN   50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF  100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE  150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN  200
QERALRTVWS SGDDKEQLVK NTYVL                                     225
(SEQ ID NO.: 49).

```

17. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

                                AGSFLAWL GSLLLLSGVLA   -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN   50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF  100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE  150
ACMLRCFRQQ ENPPLPLGSK VVVLAGAVS                               179
(SEQ ID NO.: 2).

```

```

                                MLR AEADGVSRLI GSLLLLSGVLA   -1
ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN   50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF  100

```

NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 45),

MAQLCGL RRSRAFLALL GSLLLSGVLA -1
 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 47),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 71),

or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 70).

18. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

AGSFLAWL GSLLLLSGVLA -1
 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179
 (SEQ ID NO.: 2),

MLR AEADGVSRLI GSLLLLSGVLA -1
 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 45),

MAQLCGL RRSRAFLALL GSLLLLSGVLA -1
 ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS FGD 213
 (SEQ ID NO.: 47),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
 QERALRTVWS SGDDKEQLVK NTYVL 225
 (SEQ ID NO.: 71),

or

```

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK VVVLAGLFVM VLILFLGASM VYLIRVARRN 200
QERALRTVWS FGD 213
(SEQ ID NO.: 70) .

```

19. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATV 64
(SEQ ID NO.: 4) ,

```

```

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK C 61
(SEQ ID NO.: 5) ,

```

```

YEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQ 159
(SEQ ID NO.: 6) ,

```

CTANAVTGPC RASFPRWYFD VERNSCNNFI YGGCRGNKNS YRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 7),

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQ 159
 (SEQ ID NO.: 3),

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRC 156
 (SEQ ID NO.: 50),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 25
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179
 (SEQ ID NO.: 1),

or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK 170
 (SEQ ID NO.: 52).

20. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50
YLTKEECLKK CATV	64
(SEQ ID NO.: 4),	

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50
YLTKEECLKK C	61
(SEQ ID NO.: 5),	

YEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE	150
ACMLRCFRQ	159
(SEQ ID NO.: 6),	

CTANAVTGPC RASFPRWYFD VERNSCNNFI YGGCRGNKNS YRSEE	150
ACMLRC	156
(SEQ ID NO.: 7),	

IHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF	75
NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE	125
ACMLRCFRQ	159
(SEQ ID NO.: 3),	

CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRC 156

(SEQ ID NO.: 50),

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 25
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 75
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 125
 ACMLRCFRQQ ENPPLPLGSK VVVLGAVS 179

(SEQ ID NO.: 1),

or

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
 NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE 150
 ACMLRCFRQQ ENPPLPLGSK 170

(SEQ ID NO.: 52).

21. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DS 92
 (SEQ ID NO.: 8).

22. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
 YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DS 92
 (SEQ ID NO.: 8) .

23. The use according to any one of claims 15 to 22, wherein the Kunitz-type serine protease inhibitor is glycosylated.
24. The use according to any one of claims 15 to 22, wherein the Kunitz-type serine protease inhibitor contains at least one intra-chain cysteine-cysteine disulfide bond.
25. The use according to any one of claims 15 to 22, wherein the Kunitz-type serine protease inhibitor contains at least one intra-chain cysteine-cysteine disulfide bond selected from the cysteine-cysteine paired groups consisting of CYS11-CYS61, CYS20-CYS44, CYS36-CYS57, CYS106-CYS156, CYS115-CYS139, and CYS131-CYS152, wherein the cysteine residues are numbered according to the amino acid sequence of native human placental bikunin.
26. The use according to any one of claims 15, 17, 19 and 21, wherein said physiologically acceptable carrier is a member selected from the group consisting of a physiologically buffered solution, an isotonic saline, normal saline and a combination thereof.
27. A use of a Kunitz-type serine protease inhibitor for preparing a medicament for accelerating the rate of mucociliary clearance in a subject in need of such treatment, the medicament comprising an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor and a physiologically acceptable carrier wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS' CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK                                     170
(SEQ ID NO.: 52) .

```

28. A use of an effective mucociliary clearance stimulatory amount of a Kunitz-type serine protease inhibitor for accelerating the rate of mucociliary clearance in a subject in need of such treatment, wherein the Kunitz-type serine protease inhibitor comprises the amino acid sequence:

```

ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN 50
YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF 100
NYEEYCTANA VTGPCRASFP RWYFDVERNS' CNNFIYGGCR GNKNSYRSEE 150
ACMLRCFRQQ ENPPLPLGSK                                     170
(SEQ ID NO.: 52) .

```

29. A use of a therapeutically effective mucociliary clearance stimulatory amount of a serine protease inhibitor comprising a Kunitz-type domain comprising at least 5 to 6 cysteine residues, wherein the serine protease inhibitor comprises at least one intra-chain cysteine-cysteine disulfide bond, for treating a mucociliary clearance disorder.
30. A use of a therapeutically effective mucociliary clearance stimulatory amount of a serine protease inhibitor comprising a Kunitz-type domain comprising at least 5 to 6 cysteine residues, wherein the serine protease inhibitor comprises at least one intra-chain cysteine-cysteine disulfide bond, for preparing a medicament for treating a mucociliary clearance disorder.
31. A use of a therapeutically effective amount of the protease inhibitor defined in any one of claims 1 to 30 for treating cystic fibrosis.

WO 00/37099

PCT/GB99/04381

1/52

FIG. 1

R35464	GGCCGGGTCG TTTCTCGCCT GGCTGGGATC GCTGCTCCTC TCTGGGGTCC	50
ORF	P G R F S P G W D R C S S L G S	16
R35464	TGGCCGGCCG ACCGAGAACG CAGCATCCAC GACTTCTGCC TGGTGTGCGAA	100
ORF	W P A D R E R S I H D F C L V S K	33
R35464	GGTGGTGGGC AGATTCCGGG CCTCCATGCC TAGGTGGTGG TACAATGTCA	150
ORF	V V G R F R A S M P R W W Y N V T	50
R35464	CTGACGGATC CTGCCAGCTG TTTGTGTATG GGGGCTGTGA CGGAAACAGC	200
ORF	D G S C Q L F V Y G G C D G N S	66
R35464	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	250
ORF	N N Y L T K E E C L K K C A T V T	83
R35464	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	300
ORF	E N A T G D L A T S R N A A D S S	100
R35464	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CTTGAAGACC ACTTCAGCGA	350
ORF	V P S A P R R Q D S * R P L Q R	116
R35464	TATGTTTCAA NTATTGNAAG AATAATTGCA CCGNCAACGN ATT-----	393
ORF	Y V S * I * R I I A P * T *	130

KEY

R35464 = Nucleic acid sequence of EST R35464 (SEQ ID NO:12)

ORF = EST R35464 Open Reading Frame Translation (SEQ ID NO: 13)

WO 00/37099

PCT/GB99/04381

2/52

FIG. 2

R74593	GCAATAATTA CCTGACCAAG GAGGAGTGCC TCAAGAAATG TGCCACTGTC	50
ORF	Q * L P D Q G G V P Q E M C H C H	17
R74593	ACAGAGAATG CCACGGGTGA CCTGGCCACC AGCAGGAATG CAGCGGATTC	100
ORF	R E C H G * P G H Q Q E C S G F	33
R74593	CTCTGTCCCA AGTCTCCAG AAGGCAGGAT TCTGAAGACC ACTCCAGCGA	150
ORF	L C P K S P R R Q D S E D H S S D	50
R74593	TATGTTCAAC TATGAAGAAT ACTGCACCGC CAACGCAGTC ACTGGGCCTT	200
ORF	M F N Y E E Y C T A N A V T G P C	67
R74593	GCCGTGCATC CTTCCCACGC TGGTACTTTG ACGTGGAGAG GAACTCCTGC	250
ORF	R A S F P R W Y F D V E R N S C	83
R74593	AATAACTTCA TCTATGGAGG CTGCCGGGGC AATAAGAACA GCTACCGCTC	300
ORF	N N F I Y G G C R G N K N S Y R S	100
R74593	TGAGGAGGCC TGCATGCTCC GCTGCTTCCG CCAGCAGGAG AATCCTCCCC	350
ORF	E E A C M L R C F R Q Q E N P P L	117
R74593	TGCCCCTTGG CTCAAAGGTG GTGGTTCTGG CCGGGGCTGT TTCGTGATGG	400
ORF	P L G S K V V V L A G A V S * W	133
R74593	TGTTGATCCT TTTCTGGGG AGCNTCCATG GTCTTACTGA TTCCGGGTGG	450
ORF	C * S F S W G A S M V L L I P G G	150
R74593	CAAGGAGGAA CCAGGAGCGT GCCCTGCGGA NCGTCTGGAG CTTCCGAGAT	500
ORF	K E E P G A C P A X R L E L R R *	167
R74593	GACAAGGGNT	510
ORF	Q G	169

KEY

R74593 = Nucleic acid sequence of EST R74593 (SEQ ID NO: 14)

ORF = EST R74593 Open Reading Frame Translation (SEQ ID NO: 15)

WO 00/37099

PCT/GB99/04381

3/52

FIG. 3

R35464	GGCCGGGTCGT TTCTCGCCTG GCTGGGA-TC GCTGCTCCTC TCTGGGGTCC	50
N39798	TGGGANTC GCTGCTCCTC TCTGGGGTCC	28
H94519	GCNGCG-CGT TNNTCGCNT- GCTGGGA-TC GCTGCACCTC TCTGGGGTCC	47
R74593 corr.	-----	
Consensus	GGCCGGGTCGT TTCTCGCCTG GCTGGGA-TC GCTGCTCCTC TCTGGGGTCC	50
Translation	A G S F L A W L G S L L L S G V	-3
R35464	TGGCCGGCCG ACCGAGAACG CAGCATCCAC GACTTCTGCC TGGTGTGCGAA	100
N39798	TGG-CGGCCG ACCGAGAACG CAGCATCCAC GACTTCTGCC TGGTGTGCGAA	77
H94519	NGG-CGGCCG ACCGAGAACG CAGCATCCAC GACTTCTGCC TGGTGTGCGAA	96
R74593 corr.	-----	
Consensus	TGG-CGGCCG ACCGAGAACG CAGCATCCAC GACTTCTGCC TGGTGTGCGAA	99
Translation	L A A D R E R S I H D F C L V S K	15
R35464	GGTGGTGGGC AGATTCCGGG CCTCCATGCC TAGGTGGTGG TACAATGTCA	150
N39798	GGTGGTGGGC AGATGCCGGG CCTCCATGCC TAGGTGGTGG TACAATGTCA	127
H94519	GGTGGTGGGC AGATGCCGGG CCTCCATGCC TAGGTGGTGG TACAATGTCA	146
R74593 corr.	-----	
Consensus	GGTGGTGGGC AGATGCCGGG CCTCCATGCC TAGGTGGTGG TACAATGTCA	149
Translation	V V G R C R A S M P R W W Y N V T	32
R35464	CTGACGGATC CTGCCAGCTG TTTGTGTATG GGGGCTGTGA CGGAAACAGC	200
N39798	CTGACGGATC CTGCCAGCTG TTTGTGTATG GGGGCTGTGA CGGAAACAGC	177
H94519	CTGACGGATC CTGCCAGCTG TTTGTGTATG GGGGCTGTGA CGGAAACAGC	196
R74593 corr.	-----	
Consensus	CTGACGGATC CTGCCAGCTG TTTGTGTATG GGGGCTGTGA CGGAAACAGC	199
Translation	D G S C Q L F V Y G G C D G N S	48
R35464	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	250
N39798	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	227
H94519	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	246
R74593 corr.	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	52
Consensus	AATAATTACC TGACCAAGGA GGAGTGCCTC AAGAAATGTG CCACTGTCAC	249
Translation	N N Y L T K E E C L K K C A T V T	65
R35464	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	300
N39798	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	277
H94519	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	296
R74593 corr.	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	102
Consensus	AGAGAATGCC ACGGGTGACC TGGCCACCAG CAGGAATGCA GCGGATTCCT	299
Translation	E N A T G D L A T S R N A A D S S	82
R35464	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CTTGAAGACC ACTTCAGCGA	350
N39798	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CT-GAAGACC ACTCCAGCGA	326
H94519	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CT-GAAGACC ACTCCAGCGA	345
R74593 corr.	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CT-GAAGACC ACTCCAGCGA	151
Consensus	CTGTCCCAAG TGCTCCCAGA AGGCAGGATT CT-GAAGACC ACTCCAGCGA	348
Translation	V P S A P R R Q D S E D H S S D	98

WO 00/37099

PCT/GB99/04381

4/52

FIG. 3 (Cont.)

R35464	TATGTTTCAA	NTATTGNAAG	AATAATTGCA	CCGNCAACGN	ATT-----	393												
N39798	TATGTT-CAA	CTA-TG-AAG	AATACT-GCA	CCGCCAACGC	AGTCACTGGG	372												
H94519	TATGTT-CAA	CTA-TG-AAG	AATACTGGCA	CCGCCAACGC	ATTCACCTGGG	392												
R74593 corr.	TATGTT-CAA	CTA-TG-AAG	AATACT-GCA	CCGCCAACGC	AGTCACTGGG	197												
Consensus	TATGTT-CAA	CTA-TG-AAG	AATACT-GCA	CCGCCAACGC	AGTCACTGGG	394												
Translation	M	F	N	Y	E	E	Y	C	T	A	N	A	V	T	G	113		
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----		
N39798	CCTTGC-GTG	GAATCCTTTC	CCACGCTGGN	AATTTNGACG	TTGAGAAGGA	421												
H94519	CCT-GC-GTG	-CATCCTT-C	CCACGCTGGT	ACTTT-GNCG	-----	427												
R74593 corr.	CCTTGCCGTG	-CATCCTT-C	CCACGCTGGT	ACTTT-GACG	TGGAGA-GGA	243												
Consensus	CCTTGCCGTG	-CATCCTT-C	CCACGCTGGT	ACTTT-GACG	TGGAGA-GGA	440												
Translation	P	C	R	A	S	F	P	R	W	Y	F	D	V	E	R	N	129	
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----		
N39798	AC-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	423	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	ACTCCTGCAA	TAACTTCATC	TATGGAGGCT	GCCGGGGCAA	TAAGAACAGC	293												
Consensus	ACTCCTGCAA	TAACTTCATC	TATGGAGGCT	GCCGGGGCAA	TAAGAACAGC	490												
Translation	S	C	N	N	F	I	Y	G	G	C	R	G	N	K	N	S	145	
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
N39798	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	TACCGCTCTG	AGGAGGCCTG	CATGCTCCGC	TGCTTCCGCC	AGCAGGAGAA	343												
Consensus	TACCGCTCTG	AGGAGGCCTG	CATGCTCCGC	TGCTTCCGCC	AGCAGGAGAA	540												
Translation	Y	R	S	E	E	A	C	M	L	R	C	F	R	Q	Q	E	N	162
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
N39798	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	TCCTCCCCTG	CCCCTTGGCT	CAAAGGTGGT	GGTTCTGGCC	GGGGCTGTTT	393												
Consensus	TCCTCCCCTG	CCCCTTGGCT	CAAAGGTGGT	GGTTCTGGCC	GGGGCTGTTT	590												
Translation	P	P	L	P	L	G	S	K	V	V	V	L	A	G	A	V	S	179
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
N39798	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	CGTGATGGTG	TTGATCCTTT	TCCTGGGGAG	CNTCCATGGT	CTTACTGATT	443												
Consensus	CGTGATGGTG	TTGATCCTTT	TCCTGGGGAG	CNTCCATGGT	CTTACTGATT	640												
Translation	*	W	C	*	S	F	S	W	G	A	S	M	V	L	L	I	195	
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
N39798	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	CCGGGTGGCA	AGGAGGAACC	AGGAGCGTGC	CCTGCGGANC	GTCTGGAGCT	493												
Consensus	CCGGGTGGCA	AGGAGGAACC	AGGAGCGTGC	CCTGCGGANC	<u>GTCTGGAGCT</u>	690												
Translation	P	G	G	K	E	E	P	G	A	C	P	A	*	R	L	E	L	212
R35464	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
N39798	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
H94519	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	
R74593 corr.	TCGGAGATGA	CAAGGGNT															511	
Consensus	<u>TCGGAGATGA</u>	CAAGGGNT															708	
Translation	R	R	*	Q	G												217	

WO 00/37099

PCT/GB99/04381

5/52


FIG. 3 (Cont.)


KEY

R35464 = Nucleic acid sequence of EST R35464 (SEQ ID NO.: 12)
N39798 = Nucleic acid sequence of EST N39798 (SEQ ID NO.: 17)
H94519 = Nucleic acid sequence of EST H94519 (SEQ ID NO.: 16)
R74593 corr = Corrected version of (SEQ ID NO.: 14) G at b.p. 114
Consensus = Nucleic acid sequence for human bikunin (SEQ ID NO. 9)
Translation = Amino acid Translation of Consensus (SEQ ID NO.: 10)

6/52

FIG. 4A





8/52

FIG. 4C

	1				50
Bikunin	GCGA	CCTCCGCGCG	TTGGGAGGTG	TAGCGCGGCT	CTGAACGCGT
N40851	GCGA	CCTCCGCGCG	TTGGGAGGTG	TAGCGCGGCT	CTGAACGCGT
N39876	GCGA	CCTCCGCGCG	TTGGGAGGTG	TAGCGCGGCT	CTGAACGCGT
R87894					
H16866	GGCGA	CCTCCGCGCG	TTGGGAGGTG	TAGCGCG.CT	CTGAACGGGN
R34808					
T66058					
N57450			T	TAGCGCGGCT	CTGAACGCNA
N57374					
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

FIG. 4C (Cont.)

	51				100
Bikunin	GNA GGGCCG	TTGAGTGTCG	CAGGCGGCCGA	GGGCGCGAGT	GAGGAGCAGA
N40851	NGAGNNGGCCG	TTGAGTGTCG	CAGGCGGCCGA	GGGCGCGAGT	GAGGAGCAGA
N39876	GCA.GGGCCG	TTGAGTGTCG	CAGGCGGCCGA	GGGCGCGAGT	GAGGAGCAGA
R87894		TTGAGTGTCG	NAGGCGGCCGA	GGGCGCGAGT	GAGGAGCAGA
H16866	..ANGGGCCG	TTGAGTGTCG	CAGGCGGC.A	GGGCN.GAGT	GAGGAGCAGA
R34808				G	GAGGAGCAGA
T66058					
N57450	GAAGNNGGCCG	TTGAGTGTCG	CAGGCGGCCGA	GGGCGCGAGT	GAGGAGCAGA
N57374					AGA
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

10/52

FIG. 4C (Cont.)

	101				150
Bikunin	CCCAGGCATC	GCGCGCCGAG	AAGNC	GGGC	GTCCCCACAC TGAAGGTCCG
N40851	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	GTCCCCACAC TGAAGGTCCG
N39876	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	NTCCCCACAC TGAAGGTCCG
R87894	CCCAGGCATC	GCGCGCCGAG	AAGGCCGGGC		GTCCCCACAC TGAAGGTCCG
H16866	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	GTCCCCACAC TGAAGGTCCG
R34808	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	GTCCCCACAC TGAAGGTCCG
T66058					
N57450	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	GTCCCCACAC TGAAGGTCCG
N57374	CCCAGGCATC	GCGCGCCGAG	AAGNC	.GGGC	GTCCCCACAC TGAAGGTCCG
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

11/52

FIG. 4C (Cont.)

	151				200
Bikunin	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
N40851	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
N39876	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
R87894	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
H16866	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACG.T	CCCGGAGCN.
R34808	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
T66058				...GGACCCT	CCCGGAGCGT
N57450	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
N57374	GAAAGGCGAC	TTCCGGGGGC	TTTGGCACCT	GGCGGACCCT	CCCGGAGCGT
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

12/52

FIG. 4C (Cont.)

	201		250		
Bikunin	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTTTG	.AGGGGCTTC
N40851	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTNTG	.AGGGGCTTC
N39876	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTTTG	.AGGGGCTTC
R87894	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTTTG	.AGGGGCTTC
H16866	.GGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTTTG	.AGGGGCTTC
R34808	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTNTG	GAGGGGCTTC
T66058	CGGCACCTGA	ACGCGAGGC.	CTCCATTGCG	.GTGCGTGTG	NAGGGGCTTC
N57450	CGGCACCTGA	ACGCGAGGCG	CTCCATTGCG	CGTGCGTTTG	.AGGGGCTTC
N57374	CGGCACCTGA	ACGCGAGGC.	CTCCATTGC.	CGTGCGTTNG	.AGGGGCTTC
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

13/52

FIG. 4C (Cont.)

	251				300
Bikunin	CCGCACCT G	ATCGCGAGAC	CCCAACGGCT	GGTGG CGTC	GC TG CGCG
N40851	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTGG.CGTC	GCCTG.CGCG
N39876	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTGG.CGTC	GCCTG.CGCG
R87894	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTNG.CGTC	GC.TN.CGCG
H16866	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTNG.CGTC	GC.TGGCGCG
R34808	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTGGGCGTC	GC.TG.CGCG
T66058	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTGG.CGTC	GC.TG.CGCG
N57450	CCGCACCT.G	ATCGCGAGAC	CCCAACGGCT	GGTGG.CGTC	GCCTG.CGCG
N57374	CCGGAACCTG	ATCGCGAGAC	CCCAACGGCT	GGTGG.CGTC	GC.TG.CGCG
R35464					
H94519					
N39798					
H87300					
R74593					
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

WO 00/37099

PCT/GB99/04381

14/52

FIG. 4C (Cont.)

	301		350
Bikunin	TC TCGGCTG AGCT GGCCA TGGCGCANT	GTTGC GGGC T	GAGGC GG
N40851	TC.TCGGCTG AGCT.GGNCA TGTCG		
N39876	TC.TCGGCTG AGCT.GGCCA TGGCGCACT.	G.TGCGGNGC	T.GAGGC.G
R87894	TC.TCGGCTG AGCTTGGCCA TGGCGCANT.	GTTNC.GGGC	T.NAGGC.GG
H16866	TTCTCGGCTG AGCT.GGCCA TGGCGCANT.	GTTGC.GNGC	T.GAGGC.GG
R34808	TCTTCGGCTG AGCTGGGCCA TGGCGCANTT	GTTGC.GGGC	T.GAGGC.GG
T66058	TC.TCGGCTG AGCT.GGCCA TGGCGCANT.	GTTGC.GNGC	T.GAGGC.GG
N57450	TC.TCGGCTG AGCT.GGCCA TGGCGCANT.	GGTGC.GGGC	TTGAGGC.GG
N57374	TCCTCGGCTG AGCT.GGCCA TGGCGCANT.	GGTGCCGNGC	T.GAGGCCGG
R35464			GGCCGG
H94519			
N39798			
H87300			
R74593			
R31730			
R34701			
H02982			
R32676			
T47439			
R73968			
H39840			
H95233			
H39841			
N30199			
T52966			
N29508			
N26919			
N26910			
H16757			
N27732			

15/52

FIG. 4C (Cont.)

	351					400
Bikunin	AC GG CG	TTTCTCG	CC TGCTGGG	A TCGCT GC	T CCTCTCT	
R87894	ACG.					
H16866	AC..CGNCGT	TTTTCTTCG.	CCTTGCTGGG	ATTCGCTTGC	TTCCTNTCTG	
R34808	ACGCGGNCG.	.TTTTTTCGN	CCTTGCTGGG	ATTCG.TTG.	TTNCTCTCTN	
T66058	...CGGNCG.	.TTTCTCG.	CC.TGCTGGG	A.TCGCT.GC	T.CCTCTCT.	
N57450	ANN.NGCCG.	.TTTCTCG.	CC.TGCTGGG	A.TCGCT.GC	T.CCTCTCT.	
N57374	AG..GGCCGG	.TTTCTCG.	CCTTGCTGGG	A.TCGCT.GC	T.CCTCTCTG	
R35464	GTCG.	.TTTCTCG.	CCTGGCTGGG	A.TCGCT.GC	T.CCTCTCT.	
H94519	.GCNGCGCG.	.TTNNTCG.	CN.TGCTGGG	A.TCGCT.GC	A.CCTCTCT.	
N39798			CTGGG	ANTCGCT.GC	T.CCTCTCT.	
H87300						
R74593						
R31730						
R34701						
H02982						
R32676						
T47439						
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						

FIG. 4C (Cont.)

	401						450
Bikunin	GGGG TCCTG	G	CGGCCGA	CCGA	GAACG	CA GCA TCC	ACGACTT CT
H16866	GGGGTTCCTG	GG	.CGGCCGA	CCGA	.GAACG	CA.GCA.TCC	AAGAATTTTT
R34808	GGGGTTC.TG	GGG	NGGCCGA	NCGA	.GAACG	CAAGCA.TTC	ACGA.TTT
T66058	GGGG.TCCTG	G..	CGGCCGA	CCGA	.GAACG	CA.GCA.TCC	ACGANTT.CT
N57450	GGGG.TCCTG	G..	CGGCCGA	CCGA	.GAACG	CA.GCA.TCC	ACGACTT.CT
N57374	GGGG.TCCTG	G..	CGGCCGA	NCGA	AAGAANG	CA.GCAATCC	ANGAATTNCT
R35464	GGGG.TCCTG	G.	CCGCCGA	CCGA	.GAACG	CA.GCA.TCC	ACGACTT.CT
H94519	GGGG.TCGNG	G..	CGGCCGA	CCGA	.GAACG	CA.GCA.TCC	ACGACTT.CT
N39798	GGGG.TCCTG	G..	CGGCCGA	CCGA	.GAACG	CA.GCA.TCC	ACGACTT.CT
H87300							
R74593							
R31730							
R34701							
H02982							
R32676							
T47439							
R73968							
H39840							
H95233							
H39841							
N30199							
T52966							
N29508							
N26919							
N26910							
H16757							
N27732							

FIG. 4C (Cont.)

	451					500
Bikunin	GCCTGGTGT	CGAAGGT GG	TGGGCAGATG	CCGGG	CCTC	CATGCCTA G
H16866	GCC					
T66058	TCCTGGTGTT	CGAAGG				
N57450	GCCTGGTGT.	CGAAGGT.GG	TGGGCAG			
N57374	GCCTGGTGTT	CGAAAGTTGG	TGGGCANATT	CCGGGGCCTT	CATGNCTAAG	
R35464	GCCTGGTGT.	CGAAGGT.GG	TGGGCAGATT	CCGGG.CCTC	CATGCCTA.G	
H94519	GCCTGGTGT.	CGAAGGT.GG	TGGGCAGATG	CCGGG.CCTC	CATGCCTA.G	
N39798	GCCTGGTGT.	CGAAGGT.GG	TGGGCAGATG	CCGGG.CCTC	CATGCCTA.G	
H87300						
R74593						
R31730						
R34701						
H02982						
R32676						
T47439						
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						

18/52

FIG. 4C (Cont.)

	501					550
Bikunin	G TGGT GGT ACAATGTCAC	TGACGGATCC	TGCCAGCTGT	TTGTGT	ATG	
N57374	GTTGGTTGGT ANAATGTNAA	TTAANGATT	TTGCAACTGT	TTGTGT	NATT	
R35464	G.TGGT.GGT ACAATGTCAC	TGACGGATCC	TGCCAGCTGT	TTGTGT	.ATG	
H94519	G.TGGT.GGT ACAATGTCAC	TGACGGATCC	TGCCAGCTGT	TTGTGT	.ATG	
N39798	G.TGGT.GGT ACAATGTCAC	TGACGGATCC	TGCCAGCTGT	TTGTGT	.ATG	
H87300						
R74593						
R31730						
R34701						
H02982						
R32676						
T47439						
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						
	551					600
Bikunin	GGGGCTGTGA CGGAAACA	GCAATAATTA	CCTGACCAAG	GA	GGAGTGC	
N57374	GGGGCTNTTA AACGGAAANA	.CAATAATNA	CCTGACCAAA	GAAGNAAT.	.	
R35464	GGGGCTGTGA ..CGGAAACA	GCAATAATTA	CCTGACCAAG	GA.	GGAGTGC	
H94519	GGGGCTGTGA ..CGGAAACA	GCAATAATTA	CCTGACCAAG	GA.	GGAGTGC	
N39798	GGGGCTGTGA ..CGGAAACA	GCAATAATTA	CCTGACCAAG	GA.	GGAGTGC	
H87300	GATTCGGCAC AGGGGAAACA	GCAATAATTA	CCTGACCAAG	GA.	GGAGTNC	
R74593		GCAATAATTA	CCTGACCAAG	GA.	GGAGTGC	
R31730						
R34701						
H02982						
R32676						
T47439						
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						

19/52

FIG. 4C (Cont.)

	601				650
Bikunin	CTCAAGAAAT	GTGCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
R35464	CTCAAGAAAT	GTGCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
H94519	CTCAAGAAAT	GTGCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
N39798	CTCAAGAAAT	GTGCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
H87300	CTCAAGAAAT	GTNCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
R74593	CTCAAGAAAT	GTGCCACTGT	CACAGAGAAT	GCCACGGGTG	ACCTGGCCAC
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					
	651				700
Bikunin	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGTGCTCCC	AGAAGGCAGG
R35464	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGTGCTCCC	AGAAGGCAGG
H94519	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGTGCTCCC	AGAAGGCAGG
N39798	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGTGCTCCC	AGAAGGCAGG
H87300	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGTGCTCCC	AGAAGGCAGG
R74593	CAGCAGGAAT	GCAGCGGATT	CCTCTGTCCC	AAGT.CTCCC	AGAAGGCAGG
R31730					
R34701					
H02982					
R32676					
T47439					
R73968					
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

20/52

FIG. 4C (Cont.)

	701	750
Bikunin	ATTCT GAAG ACCACTCCAG CGATATGTT CAACTAT G AAGAATACTG	
R35464	ATTCTTGAAG ACCACTTCAG CGATATGTTT CAANTATTGN AAGAATAATT	
H94519	ATTCT.GAAG ACCACTCCAG CGATATGTT. CAACTAT..G AAGAATACTG	
N39798	ATTCT.GAAG ACCACTCCAG CGATATGTT. CAACTAT..G AAGAATACTG	
H87300	ATTCT.GAAG ACCACTCCAG CGATATGTT. CAACTAT..G AAGAATACTG	
R74593	ATTCT.GAAG ACCACTCCAG CGATATGTT. CAACTAT..G AAGAATACTG	
R31730		
R34701		
H02982		
R32676		
T47439		
R73968		
H39840		
H95233		
H39841		
N30199		
T52966		
N29508		
N26919		
N26910		
H16757		
N27732		

[illegible]

21/52

FIG. 4C (Cont.)

	801					850
Bikunin	GCTGGTACTT	T	GACGTGGA	GA	GGAACTC	CTG CAATAA CTTCATCTAT
H94519	GCTGGTACTT	T	GNCGT			
N39798	GCTGGNAATT	T	NGACGTTGA	GA	GAAGGAAC	
H87300	GCTNGTACTT	T	GACGTGGA	GA	GGAACTC	CTGGCAATAA CTTCATCTAT
R74593	GCTGGTACTT	T	GACGTGGA	GA	GGAACTC	CTG.CAATAA CTTCATCTAT
R31730						
R34701						
H02982			GA	GA	GGAACTC	CTG.CAATAA CTTCATCTAT
R32676						G ATTC..GGAA
T47439						
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						
	851					900
Bikunin	GGAGGCT	GC	CGGGGCAAT	AAGAACAG	C	TACCGCTC T GAGGAGGCCT
H87300	GGAGGCTTGC	CGGGGCAATN	AAGAACAGNT	TACCGCTCTT		TAGGAGGCCT
R74593	GGAGGCT.GC	CGGGGCAAT.	AAGAACAG.C	TACCGCTC.T		GAGGAGGCCT
R31730				G.C	TACCGCTC.T	GAGGAGGCCT
R34701						
H02982	GGNGGCT.GC	CGGGG.AAT.	AAGAACA.NC	TACCGCTC.T		GAGGAGGCCT
R32676	CGAGGA..GC	CGGGGCAAT.	AAGAACAG.C	TACCGCTC.T		GAGGAGGCCT
T47439						NGGCCT
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						

WO 00/37099

PCT/GB99/04381

22/52

FIG. 4C (Cont.)

	901					950
Bikunin	GCA TGCTC	CGCTGCTTCC	GC			CA GCAGGA
H87300	.GCA.T....	.				
R74593	.GCA.TGCTC	CGCTGCTTCC	GC.....			.CA.GCAGGA
R31730	.GCA.TGCTC	CGCTGCTTCC	GC.....			.CA.GCAGGA
R34701		TTCC	GC.....			.CAAGCAGGA
H02982	.GCG.TGCTC	CGCTGCTTCC	GCTGTGTGTT	CTCTTCCAGG	CCA.	GCAGGA
R32676	.GCA.TGCTC	CGCTGCTTCC	GC.....			.CA.GCAGGA
T47439	TGCAGTGCTC	CGCTGCTTCC	GC.....			.CA.GCAGGA
R73968						
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						
	951					1000
Bikunin	GAA TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC	TGG	CGGGGC
R74593	GAA.TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC.	TGG	CGGGGC
R31730	GAA.TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC.	TGG.	CGGGGC
R34701	AAANTCCTCC	CCTCCCCCTT	GGCTCAAAGG	TGGTGGTTCC	TGG.	CGGGGC
H02982	GAA.TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC.	TGG.	CGGGGC
R32676	GAA.TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC.	TGG.	CGGGGC
T47439	GAA.TCCTCC	CCTGCCCCCTT	GGCTCAAAGG	TGGTGGTTC.	TGG.	CGGGGC
R73968						CGGGGC
H39840						
H95233						
H39841						
N30199						
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						

23/52

FIG. 4C (Cont.)

	1001				1050
Bikunin	TGTT CGTGA TGGTGTGAT	CC T CTTCC	TGGG AGCCT	CC ATGGTC	
R74593	TGTTTCGTGA TGGTGTGAT	CCTT..TTCC	TGGGGAGCCT	CC.ATGGTCT	
R31730	TGTT.CGTGA TGGTGTGAT	CC.T.CTTCC	TGGGGAGCCT	CC.ATGGTC.	
R34701	TGTT.CGTGA TGGTGTGAT	CCCTCCTTCC	CGGG.AGCCT	CCCATGGTCC	
H02982	TGTT.CGTGA TGGTGTGAT	CC.T.CTTCC	TGGG.AGCCT	CC.ATGGTN.	
R32676	TGTT.CGTGA TGGTGTGAT	CC.T.CTTCC	TGGG.AGCCT	CC.ATGGTC.	
T47439	TGTT.CGTGA TGGTGTGAT	CC.T.CTTCC	TGGG.AGCCT	CC.ATGGTC.	
R73968	TGTT.CGTGA TGGTGTGAT	CC.T.CTTCC	TGGG.AGCCT	CC.ATGGTC.	
H39840					
H95233					
H39841					
N30199					
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					
	1051				1100
Bikunin	TACC TGAT CCGGGTGGCA	CGGAGG AAC	C AGG AGCG	TGCCCTGCGC	
R74593	TAC..TGATT CCGGGTGGCA	AGGAGG.AAC	C.AGG.AGCG	TGCCCTGCGG	
R31730	TACC.TGAT. CCGGGTGGCA	CGGAGGGAAC	C.AGGGAGCG	TGCCCTGCGC	
R34701	TACCCTGAT. CCGGGTGGCA	CGGAGG.AAC	CCAGG.ANCG	TGCCCTGCGC	
H02982	TACC.TGAT. CCGGGTNGCA	CGGAGG.AAC	C.AGGGAGCG	TGCCCTGCGN	
R32676	TACC.TGAT. CCGGGTGGCA	CGGAGG.AAC	C.AGGGAGCG	TGCCCTGCGC	
T47439	TACC.TGAT. CCGGGTNGCA	CGGAGG.AAC	C.AGG.AGCG	TGCCCTGCGC	
R73968	TACC.TGAT. CCGGGTGGCA	CGGAGG.AAC	C.AGG.AGCG	TGCCCTGCGC	
H39840		...GGG.AAC	C.AGG.AGCG	TGCCCTGCGC	
H95233					
H39841					
N30199		..GAGGAACC	C.ANG.AGCT	TCCCCTGCGC	
T52966					
N29508					
N26919					
N26910					
H16757					
N27732					

24/52

FIG. 4C (Cont.)

	1101					1150
Bikunin	ACCG TCT G	GAGCTCCGGA	GATGACAAGG	AGCAGCTGG	TGAAGAAC	
R74593	ANCG.TCT.G	GAGCTTCGGA	GATGACAAGG	GNT		
R31730	ACCG.TCTGG	GAGCTCCGGA	GATGACAAGG	GAGCAGCTGG	GTGAAGAAC.	
R34701	ACCG.TCT.G	GAGCTCCGGA	GATGACAAGG	.AGCAGCTGG	.TGAAGAAC.	
H02982	ACCG.TCTNG	GAGCTCCGGA	GATGACAAGG	.AGCAGCTGG	.TGAAGAAC.	
R32676	ACCG.TCTGG	GAGCTCCGGA	GATGACAAGG	GAGCAGCTGG	.TGAAGAAC.	
T47439	ACCG.TCT.G	GAGCTCCGGA	GATGACAAGG	.AGCAGCTGG	.TGAAGAAC.	
R73968	ACCG.TCT.G	GAGCTCCGGA	GATGACAAGG	.AGCAGCTGG	.TGAAGAAC.	
H39840	ACCGGTCT.G	GAGCTCCGGA	GATGACAAGG	.AGCAGCTGG	.TGAAGAAC.	
H95233						
H39841						
N30199	ACCG.TCT.G	GAGCTCCGGA	GATNACAANG	.AGCAGCTGN	.TGAAGAACC	
T52966						
N29508						
N26919						
N26910						
H16757						
N27732						
	1151					1200
Bikunin	ACATATGT C	CTGT GACCG	CCCTGT CGC	C AAGAGG A	CT GGGGAA	
R31730	ACATATGTTC	CTGTTGACCG	NCCTGTTCGC	C.AAGAGG.A	TTGGGGGAA.	
R34701	ACATATGT.C	CTGT.GACCG	CCCTGT.CGC	C.AAGAGG.A	CT.GGGGAA.	
H02982	ACATATGT.C	CTGT.GACCG	NCCTGTTCGN	C.AAGAGG.A	CTNGGGGAAA	
R32676	ACATATGTTC	CTGTTGACCG	CCCTGTTCGC	C.AAGAGGGA	NTGGGGGAA.	
T47439	ACATATGT.C	CTGT.GACCG	CCCTGT.CGC	C.AAGAGG.A	CT.GGGGAA.	
R73968	ACATATGT.C	CTGT.GACCG	CCCTGT.CGC	C.AAGAGG.A	CT.GGGGAA.	
H39840	ACATATGT.C	CTGT.GACCG	CCCTGT.CGC	C.AAGAGG.A	CT.NGGGAA.	
H95233						
H39841		C.	CCCTGT.CGC	CCAAAAGG.A	CT.GGGGAA.	
N30199	ACATATGT.C	CTGT.GACCG	CCCTNT.CGC	C.AAGAGG.A	CT.GGGNAAA	
T52966						
N29508		CC.	CCCTNT.CGC	C.AAGAGG.A	CT.GGG.AA.	
N26919						
N26910						
H16757						
N27732						

25/52

FIG. 4C (Cont.)

	1201					1250
Bikunin	GGGAGGGG	AGACTAT G	TGT GA GCT	TTTTTT	AA A TAGA	GG
R31730	.GGGAGGGG	A				
R34701	.GGGAGGGG.	AGACTAT.G.	TGT.GA.GCT	TTTTTT..AA	A.TA	
H02982	GGGGAGGGG.	AGATTAT.G.	TGTTGA.GTT	TTTTTT..AA	ANTAG	
R32676	GGGGAGGGGG	AGANTATTGT	TGTTGA.GNT	TTTTTTTAAA	ATTAGGAGGG	
T47439	.GGGAGGGG.	AGACTAT.G.	TGT.GA.GCT	TTTTTT..AA	A.TAGA..GG	
R73968	.GGGAGGGG.	AGACTAT.G.	TGT.GA.GCT	TTTTTT..AA	A.TAGA..GG	
H39840	.GGGAGGGG.	AGACTAT.G.	TGT.GA.GCT	TTTTTT..AA	A.TAGA..GG	
H95233						
H39841	.GGGAGGGGA	AAACNAT.G.	TGT.GAACCT	TTTTTT.AAA	A.TAGA..GG	
N30199	.GGGAGGNG.	AGACTAT.G.	TGT.AA.GCT	TTTTTT..AA	A.TAGA..GG	
T52966						
N29508	.GGGAGGGG.	AGACTA..G.	TGT.GA.GCT	TTTTTT..AA	A.TAGA..GG	
N26919						
N26910						
H16757						
N27732						
	1251					1300
Bikunin	GATTGACTC	GGATTTG A	GT GATC A	TTAGGG	CT GAGGTCTGTT	
R32676	GNTTGANTTC	GGGNTTTTNA	GTGATCCAT	TTAGGGGGNT	GAG	
T47439	GATTGACTC.	.GGATTTG.A	GT.GATC.A.	TTAGGG..CT	GAGGTCTNTT	
R73968	GATTGACTC.	.GGATTTG.A	GT.GATC.A.	TTAGGG..CT	GAGGTCTGTT	
H39840	GATTGACTC.	.GGATTTG.A	GT.GATC.A.	TTAGGG..CT	GAGGTCTGTT	
H95233			A.	TTAGGG..CT	GAGGTCTGTT	
H39841	GATTGACTC.	.GGATTTG.A	GT.GATC.A.	TTAGGG..CT	GAGGTCTGTT	
N30199	GATTGACTC.	.GGATTTGGA	GT.GATC.A.	TTAGGG..CT	GAGGTCTGTT	
T52966						
N29508	GATTGACTC.	.GGATTTG.A	GT.GATCNA.	TTAGGG..CT	GAGGTCTGTT	
N26919						
N26910						
H16757						
N27732						
	1301					1350
Bikunin	TCTCTGGGAG	GTAGGACGGC	TGCTTCC TG	G TC TGGCA	GGGATGGG	
T47439	TCTCTNGGAG	GTAGGACGA				
R73968	TCTCTGGGAG	GTAGGACGGC	TGCTTCC.TG	GGTCTTGGCA	.GGGATGGGG	
H39840	TCTCTGGGAG	GTAGGACGGC	TGCTTCC.TG	G.TC.TGGCA	.GGGATGGG.	
H95233	NCTCTGGGAG	NTAGGACGGC	TGCCTTCCTG	G.TC.TGGCA	.GGGATGGG.	
H39841	TCNCTGGGAG	GTAGGACGGC	TGCTCCCCTG	G.TC.TGGCA	.GGGATGGG.	
N30199	TCTCTGGGAG	GTAGGACGGC	TGCTTCC.TG	G.TC.TGGCA	.GGGATGGG.	
T52966				..TC.TGGCA	.GGGATGGG.	
N29508	TCTCTGGGAG	GTAGGACGGC	TGCTTCA.TG	G.TC.TGGCA	.GGGATGGG.	
N26919						
N26910						
H16757			G	G.TC.TGGCA	.GGGATGGG.	
N27732			CCCTG	GGTCCTGNCA	AGGNATGGGG	

26/52

FIG. 4C (Cont.)

	1351		1400
Bikunin	TTTG CTTTG G AAATCCTC T AGGAGGCT CCTCCT CGC ATGG CC TG		
R73968	TTTG.CTTTG GGAAATCCTC TTNGGAGGCT CCTCCTTCGC ATGGGCCTTG		
H39840	TTTG.CTTTG GAGAATCCTC T.ANGAGGCT CCTCCT.CGC ATGG.CC.TG		
H95233	TTTG.CTTTG G.AAATCCTC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
H39841	TTTG.CTTTG G.AAANCCNC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
N30199	TTTG.CTTTG G.AAATCCTC T.AGGAGGCT CCTCCTTCGC ATGG.CC.TG		
T52966	TTTG.CTTTG G.AAATCCTC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
N29508	TTTG.CTTTG G.AAATCCTC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
N26919	 GAGGCT CCTCCT.CGC ATGG.CC.TG		
N26910	CTTTT GNAAATCCTC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
H16757	TTTGCCTTTG G.AAANCCTC T.AGGAGGCT CCTCCT.CGC ATGG.CC.TG		
N27732	TTTG.CTTTG G.AAATCCTC TTAGGAGGCT CCTCCT.CGC ATGG.CC.TG		
	1401		1450
Bikunin	CAGT CT GG CAGCAG CCC CGAGTTGTTT CC TCGCTG ATC GATTTC		
R73968	CAGT.CTNGG CAGCANCCCC CGAGTTTTTT TCCTTCGCTG ATCCGATTTC		
H39840	CAGT.CT.GG CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
H95233	CAGTTCT..G CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
H39841	CAGT.CT.GG CAGCAG.CCC CGAGTTGTTN .CC.TCGCTG ATC.GATNTC		
N30199	CAGT.CT.GG CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
T52966	CAGT.CT.GG CAGCAG..CC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
N29508	CAGT.CT..G CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
N26919	CAGT.CTTGG CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ANC.GATTTC		
N26910	CAGT.CT..G CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ATCGGATTTC		
H16757	CAGTNCT.GG CAGCAGACCC CGAGTTGTTT .CC.TCGCTG ATC.GATTTC		
N27732	CAGT.CT.GG CAGCAG.CCC CGAGTTGTTT .CC.TCGCTG ANC.GATTTC		
	1451		1500
Bikunin	TTT CCTCCA GGTAG AGT TTTC TTTG CTTATGTTGA ATTCCATTGC		
R73968	TTTTCCTCCA GGTAAGAATT TTTCTTTT		
H39840	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
H95233	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
H39841	TTT.CCCCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ANTCCATTGC		
N30199	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
T52966	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
N29508	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
N26919	TTT.CCNCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
N26910	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		
H16757	TTTACCCCCA GGTAG..AGT TTTCCTTTGN CTTATGTTGA ATTCCATTGC		
N27732	TTT.CCTCCA GGTAG..AGT TTTC.TTTG. CTTATGTTGA ATTCCATTGC		

27/52

FIG. 4C (Cont.)

	1501				1550
Bikunin	CTCTTTT CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT GT
H39840	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTTTGT
H95233	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
H39841	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
N30199	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
T52966	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
N29508	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
N26919	CTCTTTT.CN	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
N26910	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
H16757	CTCTTTTACT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
N27732	CTCTTTT.CT	CATCACAGAA	GTGATGTTGG	AATCGTTTCT	TTTGTTT.GT
	1551				1600
Bikunin	CTGATTATG G	TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
H39840	CTGATTATG	GGTTTTTTT	AAGTAT		
H95233	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
H39841	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
N30199	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
T52966	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
N29508	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
N26919	CTGATTATG	G..TTTTTTT	AAGTNTAAAC	AAAAGTTTTT	TATTAGCATT
N26910	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
H16757	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
N27732	CTGATTATG	G..TTTTTTT	AAGTATAAAC	AAAAGTTTTT	TATTAGCATT
	1601				1650
Bikunin	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
H95233	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAA	
H39841	CTGAAAGAAG	GAAAGTAAAN	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
N30199	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
T52966	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
N29508	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
N26919	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
N26910	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
H16757	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
N27732	CTGAAAGAAG	GAAAGTAAAA	TGTACAAGTT	TAATAAAAAG	GGGCCTTCCC
	1651			1689	
Bikunin	CTTTAG AAT	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	
H39841	CTTTAA.				
N30199	CTTTAG.AAT	AAA			
T52966	CTTTAGGAAT	NAAAANAAAA	AAGGGTG		
N29508	CTTTAG.AAT	AAATTTTCAGC	ATGTGCTTTC	AA	
N26919	CTTTAG.AAT	AAAAAAAAAA	AAAAAAAAAA	A	
N26910	CTTTAG.AAT	AAATTTTCAGC	ATGTGCTTTC	AAAAAA	
H16757	CTTTAG.AAT	AAAAAAAAAA	AAAAAAAAAA	AAAAAA	
N27732	CTTTAG.AAT	AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	

WO 00/37099

PCT/GB99/04381

28/52

FIG. 4D

EST consensus MLRAEADGVS RLLGSLLL SG VLAADRERSI HDFCLVSKVV GRCRASMPRW 50
EST consensus WYNVTDGSCQ LFVYGGCDGN SNNYLTKEEC LKKCATVTEN ATGDLATSRN 100
EST consensus AADSSVPSAP RRQDSEDHSS DMFNYEEYCT ANAVTGPCRA SFPRWYFDVE 150
EST consensus RNSCENNFIYG GCRGNKNSYR SEEACMLRCF RQENPPLPL GSKVVVLAGL 200
EST consensus FVMVLILFLG ASMVYLIRVA RRNQERALRT VWSSGDDKEQ LVKNITYVL 248

WO 00/37099

PCT/GB99/04381

29/52

FIG. 4E

cDNA		ACC	3
translation		T	-47
cDNA	TGATCGCGAG ACCCCAACGG CTGGTGGCGT CGCCTGCGCG TCTCGGCTGA	53	
translation	. S R D P N G W W R R L R V S A E	-30	
cDNA	GCTGGCCATG GCGCAGCTGT GCGGGCTGAG GCGGAGCCGG GCGTTTCTCG	103	
translation	L A M A Q L C G L R R S R A F L A	-13	
cDNA	CCCTGCTGGG ATCGCTGCTC CTCTCTGGGG TCCTGGCGGC CGACCGAGAA	153	
translation	L L G S L L L S G V L A A D R E	4	
cDNA	CGCAGCATCC ACGACTTCTG CCTGGTGTCTG AAGGTGGTGG GCAGATGCCG	203	
translation	R S I H D F C L V S K V V G R C R	21	
cDNA	GGCCTCCATG CCTAGGTGGT GGTACAATGT CACTGACGGA TCCTGCCAGC	253	
translation	A S M P R W W Y N V T D G S C Q L	38	
cDNA	TGTTTGTGTA TGGGGGCTGT GACGGAAACA GCAATAATTA CCTGACCAAG	303	
translation	F V Y G G C D G N S N N Y L T K	54	
cDNA	GAGGAGTGCC TCAAGAAATG TGCCACTGTC ACAGAGAATG CCACGGGTGA	353	
translation	E E C L K K C A T V T E N A T G D	71	
cDNA	CCTGGCCACC AGCAGGAATG CAGCGGATTC CTCTGTCCCA AGTGCTCCCA	403	
translation	L A T S R N A A D S S V P S A P R	88	
cDNA	GAAGGCAGGA TTCTGAAGAC CACTCCAGCG ATATGTTCAA CTATGAAGAA	453	
translation	R Q D S E D H S S D M F N Y E E	104	
cDNA	TACTGCACCG CCAACGCAGT CACTGGGCCT TGCCGTGCAT CCTTCCCACG	503	
translation	Y C T A N A V T G P C R A S F P R	121	
cDNA	CTGGTACTTT GACGTGGAGA GGAATCCTG CAATAACTTC ATCTATGGAG	553	
translation	W Y F D V E R N S C N N F I Y G G	138	
cDNA	GCTGCCGGGG CAATAAGAAC AGCTACCGCT CTGAGGAGGC CTGCATGCTC	603	
translation	C R G N K N S Y R S E E A C M L	154	
cDNA	CGCTGCTTCC GCCAGCAGGA GAATCCTCCC CTGCCCCTTG GCTCAAAGGT	653	
translation	R C F R Q Q E N P P L P L G S K V	171	
cDNA	GGTGGTTCTG GCGGGGCTGT TCGTGATGGT GTTGATCCTC TTCCTGGGAG	703	
translation	<u>V V L A G L F V M V L I L F L G A</u>	188	
cDNA	CCTCCATGGT CTACCTGATC CGGGTGGCAC GGAGGAACCA GGAGCGTGCC	753	
translation	<u>S M V Y L I</u> R V A R R N Q E R A	204	
cDNA	CTGCGCACCG TCTGGAGCTT CGGAGATGA	782	
translation	L R T V W S F G D	213	

WO 00/37099

PCT/GB99/04381

30/52

FIG. 4F

cDNA	GCACGAGTTG	GGAGGTGTAG	CGCGGCTCTG	AACGCGCTGA	GGGCCGTTGA	50
cDNA	GTGTGCGCAGG	CGGCGAGGGC	GCGAGTGAGG	AGCAGACCCA	GGCATCGCGC	100
cDNA	GCCGAGAAGG	CCGGGCGTCC	CCACACTGAA	GGTCCGGAAA	GGCGACTTCC	150
cDNA	GGGGGCTTTG	GCACCTGGCG	GACCCTCCCG	GAGCGTCGGC	ACCTGAACGC	200
cDNA	GAGGCGCTCC	ATTGCGCGTG	CGCGTTGAGG	GGCTTCCCGC	ACCTGATCGC	250
cDNA	GAGACCCCAA	CGGCTGGTGG	CGTCGCCTGC	GCGTCTCGGC	TGAGCTGGCC	300
cDNA	ATGGCGCAGC	TGTGCGGGCT	GAGGCGGAGC	CGGGCGTTTC	TCGCCCTGCT	350
translation	M	A	Q	L	C	G
	L	R	R	S	R	A
	F	L	A	L	L	-11
cDNA	GGGATCGCTG	CTCCTCTCTG	GGGTCCTGGC	GGCCGACCGA	GAACGCAGCA	400
translation	G	S	L	L	L	S
	G	V	L	A	A	D
	R	E	R	S	I	7
cDNA	TCCACGACTT	CTGCCTGGTG	TCGAAGGTGG	TGGGCAGATG	CCGGGCCTCC	450
translation	H	D	F	C	L	V
	S	K	V	V	G	R
	C	R	A	S		23
cDNA	ATGCCTAGGT	GGTGGTACAA	TGTCACTGAC	GGATCCTGCC	AGCTGTTTGT	500
translation	M	P	R	W	W	Y
	N	V	T	D	G	S
	C	Q	L	F	V	40
cDNA	GTATGGGGGC	TGTGACGGAA	ACAGCAATAA	TTACCTGACC	AAGGAGGAGT	550
translation	Y	G	G	C	D	G
	N	S	N	N	Y	L
	T	K	E	E	C	57
cDNA	GCCTCAAGAA	ATGTGCCACT	GTCACAGAGA	ATGCCACGGG	TGACCTGGCC	600
translation	L	K	K	C	A	T
	V	T	E	N	A	T
	G	D	L	A		73
cDNA	ACCAGCAGGA	ATGCAGCGGA	TTCCTCTGTC	CCAAGTGCTC	CCAGAAGGCA	650
translation	T	S	R	N	A	A
	D	S	S	V	P	S
	A	P	R	R	Q	90
cDNA	GGATTCTGAA	GACCACTCCA	GCGATATGTT	CAACTATGAA	GAATACTGCA	700
translation	D	S	E	D	H	S
	S	D	M	F	N	Y
	E	E	Y	C	T	107
cDNA	CCGCCAACGC	AGTCACTGGG	CCTTGCCGTG	CATCCTTCCC	ACGCTGGTAC	750
translation	A	N	A	V	T	G
	P	C	R	A	S	F
	P	R	W	Y		123
cDNA	TTTGACGTGG	AGAGGAACTC	CTGCAATAAC	TTCATCTATG	GAGGCTGCCG	800
translation	F	D	V	E	R	N
	S	C	N	N	F	I
	Y	G	G	C	R	140
cDNA	GGGCAATAAG	AACAGCTACC	GCTCTGAGGA	GGCCTGCATG	CTCCGCTGCT	850
translation	G	N	K	N	S	Y
	R	S	E	E	A	C
	M	L	R	C	F	157
cDNA	TCCGCCAGCA	GGAGAATCCT	CCCCTGCCCC	TTGGCTCAAA	GGTGGTGGTT	900
translation	R	Q	Q	E	N	P
	P	L	P	L	G	S
	K	<u>V</u>	<u>V</u>	<u>V</u>		173
cDNA	CTGGCGGGGC	TGTTGCTGAT	GGTGTGATC	CTCTTCCTGG	GAGCCTCCAT	950
translation	<u>L</u>	<u>A</u>	<u>G</u>	<u>L</u>	<u>F</u>	<u>V</u>
	<u>M</u>	<u>V</u>	<u>L</u>	<u>I</u>	<u>L</u>	<u>F</u>
	<u>L</u>	<u>G</u>	<u>A</u>	<u>S</u>	<u>M</u>	190
cDNA	GGTCTACCTG	ATCCGGGTGG	CACGGAGGAA	CCAGGAGCGT	GCCCTGCGCA	1000
translation	<u>V</u>	<u>Y</u>	<u>L</u>	<u>I</u>	<u>R</u>	<u>V</u>
	<u>A</u>	<u>R</u>	<u>R</u>	<u>N</u>	<u>Q</u>	<u>E</u>
	<u>R</u>	<u>A</u>	<u>L</u>	<u>R</u>	<u>T</u>	207
cDNA	CCGTCTGGAG	CTCCGGAGAT	GACAAGGAGC	AGCTGGTGAA	GAACACATAT	1050
translation	V	W	S	S	G	D
	D	K	E	Q	L	V
	K	N	T	Y		223
cDNA	GTCCTGTGAC	CGCCCTGTCT	CCAAGAGGAC	TGGGGAAGGG	AGGGGAGACT	1100
translation	V	L	*			
						225

31/52

FIG. 4F (Cont.)

cDNA	ATGTGTGAGC	TTTTTTTAAA	TAGAGGGATT	GACTCGGATT	TGAGTGATCA	1150
cDNA	TTAGGGCTGA	GGTCTGTTTC	TCTGGGAGGT	AGGACGGCTG	CTTCCTGGTC	1200
cDNA	TGGCAGGGAT	GGGTTTGCTT	TGGAAATCCT	CTAGGAGGCT	CCTCCTCGCA	1250
cDNA	TGGCCTGCAG	TCTGGCAGCA	GCCCCGAGTT	GTTTCCTCGC	TGATCGATTT	1300
cDNA	CTTTCCTCCA	GGTAGAGTTT	TCTTTGCTTA	TGTTGAATTC	CATTGCCTCC	1350
cDNA	TTTTCTCNAT	CACAGAAGTG	ATGTTGGAAT	CGTTTCCTTT	GTTTGTCTGA	1400
cDNA	TTTATGGTTT	TTTTAAGTAT	AAACAAAAGT	TTTTTATTAG	CATTCTGAAA	1450
cDNA	GAAGGAAAGT	AAAATGTACA	AGTTTAATAA	AAAGGGGCCT	TCCCCTTTAG	1500
cDNA	AATAAATTC	CAGCATGTTG	CTTTCAAAAA	AAAAAAAAAA	AAAA	1550

32/52

FIG. 4G

EST consensus	MLR AEADGVSRL	GSLLLSGVLA	-1
PCR clone	MAQLCGL RRSRAFLALL	GSLLLSGVLA	-1
λcDNA clone	MAQLCGL RRSRAFLALL	GSLLLSGVLA	-1
EST consensus	ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50	
PCR clone	ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50	
λcDNA clone	ADRERSIHDF CLVSKVVGRC RASMPRWWYN VTDGSCQLFV YGGCDGNSNN	50	
EST consensus	YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF	100	
PCR clone	YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF	100	
λcDNA clone	YLTKEECLKK CATVTENATG DLATSRNAAD SSVPSAPRRQ DSEDHSSDMF	100	
EST consensus	NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE	150	
PCR clone	NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE	150	
λcDNA clone	NYEEYCTANA VTGPCRASFP RWYFDVERNS CNNFIYGGCR GNKNSYRSEE	150	
EST consensus	ACMLRCFRQQ ENPPLPLGSK <u>VVVLAGLFVM VLILFLGASM VYLIRVARRN</u>	200	
PCR clone	ACMLRCFRQQ ENPPLPLGSK <u>VVVLAGLFVM VLILFLGASM VYLIRVARRN</u>	200	
λcDNA clone	ACMLRCFRQQ ENPPLPLGSK <u>VVVLAGLFVM VLILFLGASM VYLIRVARRN</u>	200	
EST consensus	QERALRTVWS SGDDKEQLVK NTYVL	225	
PCR clone	QERALRTVWS FGD	213	
λcDNA clone	QERALRTVWS SGDDKEQLVK NTYVL	225	

33/52

FIG. 5






FIG. 6



35/52

FIG. 7



WO 00/37099

PCT/GB99/04381

FIG. 8A

36/52





FIG. 8B



37/52

FIG. 9A

SDS-PAGE






FIG. 9B

Western



38/52

FIG. 10



39/52

FIG. 11A




FIG. 11B




FIG. 12A





FIG. 12B




WO 00/37099

PCT/GB99/04381

40/52

FIG. 13



41/52

FIG. 14








FIG. 15





43/52

FIG. 17





FIG. 18



44/52

FIG. 19







FIG. 20





46/52

FIG. 23





FIG. 24



47/52

FIG. 25A





FIG. 25B



48/52

FIG. 26





FIG. 27



49/52

FIG. 28A





FIG. 28B



50/52

FLOW CHART OF CHO rhBIKUNIN PURIFICATION
PROCESS TRAIN

FIG. 29





FIG. 30A




WO 00/37099

PCT/GB99/04381

51/52

FIG. 30B




WO 00/37099

PCT/GB99/04381

52/52

FIG. 31



Application number/ Numéro de demande : 2356404

Documents of poor quality scanned
(request original documents in File Prep. Section on the 10th floor)

Documents de piètre qualité numérisés
(Pour obtenir les documents originaux, veuillez vous adresser à la Section de préparation
des dossiers, située au 10^e étage)