




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 305/14	A1	(11) International Publication Number: WO 93/06093 (43) International Publication Date: 1 April 1993 (01.04.93)
(21) International Application Number: PCT/US92/07935 (22) International Filing Date: 22 September 1992 (22.09.92) (30) Priority data: 763,805 23 September 1991 (23.09.91) US 900,408 18 June 1992 (18.06.92) US (71) Applicant: FLORIDA STATE UNIVERSITY [US/US]; 2035 East Paul Dirac Drive, 109 Herb Morgan Building, Tallahassee, FL 32310 (US). (72) Inventors: HOLTON, Robert, A. ; Florida State University, Dittmer Laboratory of Chemistry, Chemistry Department, Room 715, Tallahassee, FL 32306 (US). CHEN, Shu-Hui ; FARINA, Vittorio ; Bristol-Myers Squibb Company, #5 Research Parkway, Wallingford, CT 06492-7660 (US).		(74) Agents: HEJLEK, Edward, J. et al.; Senniger, Powers, Leavitt & Roedel, One Metropolitan Square, 16th Floor, St. Louis, MO 63102 (US). (81) Designated States: AU, CA, CS, FI, HU, JP, KP, KR, NO, PL, RU, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE). Published <i>With international search report.</i>

(54) Title: 10-DESACETOXYTAXOL DERIVATIVES



(57) Abstract

The present invention relates to 10-desacetoxytaxol and derivatives thereof, which are useful as antitumor agents. These compounds have formula (1) wherein R₂ is hydrogen, hydroxy or a protected hydroxy group; R₃ and R₄ are independently hydrogen, hydroxy, a protected hydroxy group, methyl, -SH, -NH₂, or -NR₈R₉; R₅ is R₁₀, or -OR₁₀; R₆ and R₇ are independently hydrogen, alkyl, or aryl; R₈ and R₉ are independently hydrogen, alkyl, alkenyl, alkynyl, or aryl; and R₁₀ is alkoxy, alkyl, alkenyl, alkynyl, or aryl.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain			US	United States of America

10-DESACETOXYTAXOL DERIVATIVES


REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part application of U.S. Serial No. 07/863,849, filed April 6, 1992, which in turn is a continuation-in-part application of U.S. Serial No. 07/763,805, filed September 23, 1991, now abandoned, and this application is a continuation-in-part application of U.S. Serial No. 07/900,408, filed June 18, 1992.

10 BACKGROUND OF THE INVENTION

The present invention relates to novel antitumor agents. More particularly, the present invention relates to 10-desacetoxytaxol and derivatives thereof.


Taxol is a natural product extracted from the bark of yew trees. It has been shown to have excellent antitumor activity in in vivo animal models, and recent studies have elucidated its unique mode of action, which involves abnormal polymerization of tubulin and disruption of mitosis. It is currently undergoing clinical trials against ovarian, breast and other types of cancer in the United States and France and preliminary results have confirmed it as a most promising chemotherapeutic agent. The structure of taxol and the numbering system conventionally used is shown below; this numbering system is also applicable to compounds of the present invention.



SUMMARY OF THE INVENTION

Among the objects of the present invention, therefore, is the provision of the novel taxane, 10-desacetoxytaxol, and derivatives thereof, which exhibit anti-tumor activity.

In accordance with the present invention, a compound is provided having the following structural formula (1):



(1)

wherein R₂ is hydrogen, hydroxy or a protected hydroxy group; R₃ and R₄ are independently hydrogen, hydroxy, a protected hydroxy group, methyl, -SH, -NH₂, or -NR₈R₉; R₅ is R₁₀, or -OR₁₀; R₆ and R₇ are independently hydrogen, alkyl, or aryl; R₈ and R₉ are independently hydrogen, alkyl, alkenyl, alkynyl, or aryl; and R₁₀ is alkoxy, alkyl, alkenyl, alkynyl, or aryl.

Also provided are the compounds 10-desacetoxytaxol and 10-deoxytaxotere.


Other objects and features of this invention will be in part apparent and in part pointed out hereinafter.

DETAILED DESCRIPTION OF THE INVENTION

As used herein "Ar" means aryl; "Ph" means phenyl; "Ac" means acetyl; "R" means alkyl; "11-ene" means

a double bond between carbon atoms 11 and 12 of a compound of formula (1); and "10,12-diene" means double bonds between carbon atoms 10 and 11, and 12 and 18 of a compound of formula (1). "Hydroxy protecting group" includes, but is not limited to, ethers such as methyl, t-butyl, benzyl, p-methoxybenzyl, p-nitrobenzyl, allyl, trityl, methoxymethyl, methoxyethoxymethyl, ethoxyethyl, tetrahydropyranyl, tetrahydrothiopyranyl, and trialkylsilyl ethers such as trimethylsilyl ether, triethylsilyl ether, dimethylarylsilyl ether, triisopropylsilyl ether and t-butyl dimethylsilyl ether; esters such as benzoyl, acetyl, phenylacetyl, formyl, mono-, di-, and trihaloacetyl such as chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl; and carbonates including but not limited to alkyl carbonates having from one to six carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl; isobutyl, and n-pentyl; alkyl carbonates having from one to six carbon atoms and substituted with one or more halogen atoms such as 2,2,2-trichloroethoxymethyl and 2,2,2-trichloroethyl; alkenyl carbonates having from two to six carbon atoms such as vinyl and allyl; cycloalkyl carbonates have from three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and phenyl or benzyl carbonates optionally substituted on the ring with one or more C₁₋₆alkoxy, or nitro.


The present invention provides novel 10-des-acetoxytaxol derivatives of the formula (1)



(1)


wherein R₂ is hydrogen, hydroxy or a protected hydroxy group; R₃ and R₄ are independently hydrogen, hydroxy, a
 5 protected hydroxy group, methyl, -SH, -NH₂, or -NR₈R₉; R₅
 is R₁₀, or -OR₁₀; R₆ and R₇ are independently hydrogen,
 alkyl, or aryl; R₈ and R₉ are independently hydrogen,
 alkyl, alkenyl, alkynyl, or aryl; and R₁₀ is alkoxy, alkyl,
 alkenyl, alkynyl, or aryl.

10 Exemplary compounds within the generic formula
 are depicted hereinbelow:




(2)


5




(3)




(4)




(5)




(6)



(7)



(8)




(9)

Preferably, one of R_3 and R_4 is hydrogen or methyl, while the other is hydroxy, or a protected hydroxy group. Preferably R_5 is phenyl, p-substituted phenyl or lower alkoxy, and most preferably phenyl, methoxy, ethoxy, or t-butoxy. Preferably, one of R_6 and R_7 is hydrogen or lower alkyl and the other is phenyl or naphthyl. Preferably R_8 is hydrogen or lower alkyl and R_9 is an amine protecting group.

In one preferred embodiment of compounds of formula (1), R_2 is hydroxy or a protected hydroxy group; R_3 is hydrogen; R_4 is hydroxy or a protected hydroxy group; R_5 is phenyl; R_6 is hydrogen and R_7 is phenyl. In another preferred embodiment, R_5 is t-butoxy. Where either R_3 or R_4 is a protected hydroxy group, it is preferably one of the carbonate type, most preferably it is selected from the group consisting of 2,2,2-trichloroethoxycarbonyl, allyloxycarbonyl, and benzyloxycarbonyl; or one of the trialkylsilyl type, most preferably triethylsilyl. The most preferred embodiments of compounds of formula (1) are: 1) 10-desacetoxytaxol; and 2) 10-deoxytaxotere.

A method for preparing compounds of formula (1) is provided by following the reaction sequence depicted in Scheme I.

SCHEME I



As illustrated, the starting materials used for the preparation of compounds of formula (1) using Scheme 1 include 10-deacetyltaxol, which may be obtained from taxol by treating the latter with zinc bromide in chloroform/

methanol as reported in G. Samaranayake et al., J. Org. Chem., 1991, 56:5114-5119, hereby incorporated by reference.

In Scheme I, P₁ is a hydroxy protecting group. Thus in the first reaction step, the 2'-hydroxy group of
5 10-deacetyltaxol is protected; the protecting group may be one that can be introduced and removed without unacceptably affecting the rest of the molecule, more preferably it may be one that preferentially blocks the 2'-hydroxy group over the two secondary hydroxy groups on the ring, namely, the
10 7- and 10-hydroxy groups. Protection of a hydroxy group may be accomplished by methods well known in the art; for example, reacting with a carboxylic acid or its acylating equivalent to form an ester, reacting with an alkyl halide in the presence of a base to form an ether, or reacting
15 with a chloroformate to form the carbonate.

The preferred 2'-hydroxy protecting group in this reaction is a carbonate which can be formed by treating 10-deacetyltaxol with an appropriate chloroformate, for example, allyl or benzyl chloroformate to form the
20 corresponding carbonate. The reaction is carried out in an inert organic solvent such as dichloromethane, tetrahydrofuran, acetonitrile, dimethylformamide, benzene, pyridine, p-dioxan, and preferably in the presence of an acid scavenger such as an amine base, for example, pyridine,
25 diisopropylethylamine, 4-aminopyridine, triethylamine and the like, or an inorganic base such as potassium carbonate or tetrabutylammonium hydroxide, at a suitable temperature which may be -78°C to about 50°C, depending on the particular reagents chosen. Generally from about one to
30 ten equivalents of the protecting group reagent relative to the taxol compound may be used; it is advantageous to monitor the reaction for example by thin-layer chromatography so as to achieve the desired degree of protection.

The 2'-hydroxy protected 10-deacetyltaxol thus obtained is then treated with 1,1,2-trifluoro-2-chlorotriethylamine (hereinafter TFCT) to give 2'-hydroxy protected 7-Q-chlorofluoroacetyl-10-desacetoxy-11,12-dihydrotaxol-10,12(18)-diene. TFCT can be prepared from diethylamine and trifluorochloroethylene according to the method reported in N. N. Yarovenko, J. Org. Chem. USSR (English), 1959, 29:2125-8, hereby incorporated by reference. The reaction of 2'-hydroxy protected 10-deacetyltaxol and TFCT is carried out in an inert organic solvent such as halogenated hydrocarbons, for example, dichloromethane, chloroform, and carbon tetrachloride, at a temperature of from about 0°C up to the refluxing temperature of the reaction solution, preferably the reaction is run at ambient temperature. The TFCT is used in at least 1 equivalent to the taxol reactant, but preferably it is used in excess; typically from about one to about ten equivalents of TFCT may be used relative to the taxol reactant.


The 2'-hydroxy protecting group can be removed using methods known in the art that are suitable for the particular protecting group used; for example, acid or base catalyzed hydrolysis, reduction, hydrogenolysis, and the like. Thus the allyl carbonate can be removed by tributyltin hydride and tetrakis(triphenylphosphine)-palladium; the benzyl carbonate can be removed by catalytic hydrogenolysis.

7-Q-Chlorofluoroacetyl-10-desacetoxy-11,12-dihydrotaxol-10,12(18)-diene can be converted to 10-des-acetoxy-7-Q-chlorofluoroacetyltaxol by catalytic hydrogenation in which the catalyst may be, for example, palladium, platinum, rhodium and the like. In cases where the 2'-hydroxy protecting group used in the previous steps of the reaction sequence is benzyloxycarbonyl, catalytic

hydrogenation converts the 10,12-diene into the 11-ene, and removes the 2'- protecting group in one step. It is understood that the order of deprotecting the 2'-hydroxy group and hydrogenation is not crucial, and either may be
5 effected prior to the other.

A further method for preparing compounds of the present invention provides compounds of formula (1) wherein R_2 is hydrogen, hydroxy, or protected hydroxy; and R_5 is phenyl. Compounds of this type may be prepared from
10 10-deacetyltaxol or 10-deacetyl-7-epitaxol as depicted in Scheme II.

SCHEME II



In Scheme II, R_a is hydroxy, R_b is hydrogen, R_2' is OP_2 , and R_3' is hydrogen; or R_a is hydrogen, R_b is hydroxy, R_2' is hydrogen, and R_3' is hydroxy; P_2 is a hydroxy protecting group. When 10-deacetyltaxol (i.e. R_a

is hydroxy and R_D is hydrogen) is the starting material, it is desirable to protect both the 2'- and the 7-hydroxy groups, and that the 10-hydroxy group be left free; preferably the protecting group is 2,2,2-trichloroethoxy-carbonyl which can be introduced by reacting the taxol
5 reactant with 2,2,2-trichloroethyl chloroformate in the presence of a base such as pyridine, diisopropylamine, triethylamine, dimethylaminopyridine, and the like. To control the degree of protection and to minimize blocking
10 the 10-hydroxy group, the base is usually used in about one to two equivalents relative to taxol and the chloroformate in about 0.6 to about 1.5 equivalents relative to taxol. When 10-deacetyl-7-epitaxol (i.e. R_A is hydrogen and R_D is hydroxy) is the starting material, only the 2'-hydroxy
15 group is protected since the 7-hydroxy group is substantially more inert than the 2'-hydroxy group, and is relatively insusceptible to acylation, either with a chloroformate, or with the TFCT in the subsequent step.


The 2',7-bishydroxy protected 10-deacetyltaxol or
20 2'-hydroxy protected 10-deacetyl-7-epitaxol thus obtained is then treated with TFCT to give 2',7-bishydroxy protected 10-desacetoxy-11,12-dihydrotaxol-10,12(18)-diene or 2'-protected 10-desacetoxy-11,12-dihydro-7-epitaxol-10,12(18)-diene, respectively. The reaction with TFCT is
25 carried out in an inert organic solvent such as halogenated hydrocarbons such as dichloromethane, chloroform and carbon tetrachloride, at a temperature of from about 0°C to about the refluxing temperature of the reaction solution, preferably the reaction is run at ambient temperature. The
30 TFCT is used in at least 1 equivalent to the taxol reactant, but preferably it is used in excess; typically TFCT is used from about one to about ten equivalents. The 2'- and, if present, the 7-hydroxy protecting group is then removed using a method suitable for the particular

protecting group to provide 10-desacetoxy-11,12-dihydro-taxol-10,12(18)-diene or 10-desacetoxy-11,12-dihydro-7-epitaxol-10,12(18)-diene; for example, trichloroethoxy-carbonyl group is removed by a zinc reagent, benzyloxy-carbonyl is removed by hydrogenolysis, and allyl carbonate
5 is removed by tributyltin hydride and tetrakis(triphenylphosphine)palladium. 10-Desacetoxy-11,12-dihydro-10,12(18)-diene and 10-desacetoxy-11,12-dihydro-7-epitaxol-10,12(18)-diene may then be subjected to catalytic
10 hydrogenation as previously discussed to provide 10-desacetoxytaxol and 10-desacetoxy-7-epitaxol, respectively.

2',7-Bishydroxy protected 10-desacetoxy-11,12-dihydro-10,12(18)-diene or 2'-hydroxy protected 10-desacetoxy-11,12-dihydro-7-epitaxol-10,12(18)-diene can
15 also be subjected to catalytic hydrogenation to give 2',7-bishydroxy protected 10-desacetoxytaxol or 2'-hydroxy protected 10-desacetoxy-7-epitaxol, respectively. The hydroxy protecting groups may be removed as previously described to provide 10-desacetoxytaxol or 10-desacetoxy-
20 7-epitaxol.

A further method for preparing compounds of the present invention provides compounds of formula (1) wherein R_2 is hydroxy or a protected hydroxy group, R_3 is hydrogen; R_4 is hydrogen, hydroxy, or a hydroxy protecting group; R_5
25 is t-butoxy; R_6 is hydrogen and R_7 is phenyl. Compounds of this type may be prepared from 7-protected 10-desacetoxy baccatin III or 10-desacetoxy-7-epi-baccatin III using the process shown in Scheme III.

SCHEME III



In Scheme III, R_a is hydroxy, R_b is hydrogen, R_2' is $-\text{OP}_3$, and R_3' is hydrogen; or R_a is hydrogen, R_b is hydroxy, R_2' is hydrogen, and R_3' is hydroxy; P_3 is a


hydroxy protecting group; preferably it is a trialkylsilyl group, most preferably triethylsilyl. 7-Hydroxy protected 10-desacetoxy baccatin III or 10-desacetoxy-7-epi-baccatin III is treated with a strong base such as n-butyl lithium
5 to generate the corresponding 13-alkoxide. The alkoxide is reacted with (3R,4S)-1-t-butoxycarbonyl-4-phenyl-3-protected hydroxy-2-azetidinone in an inert organic solvent such as tetrahydrofuran to provide the corresponding 2',7-bishydroxy protected 10-deoxytaxotere. The reaction
10 is carried out at a low temperature in the range of about -20°C to about 25°C, preferably at about 0°C. The hydroxy protecting groups are then removed to provide 10-deoxy-taxotere using conventional methods, for example triethylsilyl group may be removed by acid hydrolysis with hydro-
15 chloric acid.

The reaction sequence depicted in Scheme III shows the preparation of 10-deoxytaxotere or 10-deoxy-7-epitaxotere; however, it is equally applicable for preparing 10-desacetoxytaxol or 10-desacetoxy-7-epitaxol.
20 Thus, 10-desacetoxytaxol and 10-desacetoxy-7-epitaxol may be obtained in an analogous manner by replacing the β -lactam shown in Scheme III with (3R,4S)-1-benzoyl-4-phenyl-3-hydroxy protected-2-azetidinone.

Other derivatives of 10-desacetoxytaxol within
25 the scope of formula (1) may readily be prepared by selection of the proper substituents for the β -lactam which forms the β -amido ester side chain at C-13 of the taxol structure.

10-Desacetoxybaccatin III and 10-desacetoxy-
30 7-epibaccatin III may be obtained from 10-deacetyl baccatin III and 10-deacetyl-7-epibaccatin III by the process shown in Scheme IV.

SCHEME IV



In Scheme IV, R_2' is hydrogen and R_3' is hydroxy, or R_2' is protected hydroxy and R_3' is hydrogen. Preferably, the hydroxy protecting group is trialkylsilyl ether type, for example, triethylsilyl. Thus 10-deacetyl-
 10 baccatin III is first treated with a strong base such as n-butyl lithium, and alkoxide generated in situ is reacted with pentafluorophenyl chlorothionoformate to provide the corresponding 10-(pentafluorophenyl)thiocarbonate. The reaction is carried out in an inert organic solvent such as
 15 tetrahydrofuran, at a temperature in the range of about -78°C to about room temperature. The thiocarbonate thus

obtained is treated with a suitable reagent such as tributyltin hydride, triphenyltin hydride, trimethylsilane, and tributyltin chloride and sodium cyanoborohydride, in the presence of a radical initiator such as azobisisobutyronitrile (AIBN). The reaction is carried out in an organic solvent such as t-butanol, N-methylpyrrolidone, dimethylsulfoxide, dimethylformamide, toluene and benzene at a temperature from about 80°C to about 115°C to provide 7-protected hydroxy-10-desacetoxy baccatin III or 7-epi-baccatin III. Alternatively, the reaction can be carried out under photochemical conditions (triorganotin hydride, $h\nu$, benzene, 0°C - room temperature). The hydroxy protecting group may be removed using methods known in the art; for example, triethylsilyl can be removed by acid hydrolysis using hydrochloric acid.

The 10-desacetoxytaxol derivatives of the present invention are useful agents for inhibiting tumor growth in animals and human. Compounds were evaluated in in vitro cytotoxicity activity against human colon carcinoma cells HCT-116 and HCT-116/VM46. The HCT116/VM cells are cells that have been selected for teniposide resistance and express the multidrug resistance phenotype, including resistance to taxol. Cytotoxicity was assessed in HCT-116 human colon carcinoma cells by XTT (2,3-bis(2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino)carbonyl]2H-tetrazolium hydroxide) assay as reported in D.A. Scudiero, et al., "Evaluation of soluble tetrazolium/formazan assay for cell growth and drug sensitivity in culture using human and other tumor cell lines," Cancer Res. 48:4827-4833, 1988. Cells were plated at 4000 cells/well in 96 well microtiter plates and 24 hours later drugs were added and serially diluted. The cells were incubated at 37°C for 72 hours at which time the tetrazolium dye, XTT, was added. A dehydrogenase enzyme in live cells reduces the XTT to a

form that absorbs light at 450 nm which can be quantitated spectrophotometrically. The greater the absorbance, the greater the number of live cells. The results are expressed as an IC_{50} , which is the drug concentration required to inhibit cell proliferation (i.e., absorbance at 450 nm) to 50% of that of untreated control cells. The IC_{50} values for 10-desacetoxytaxol and taxol are given in Table I.

TABLE I

10 In vitro cytotoxicity for taxol analogs against human colon carcinoma cells.

		IC_{50} (μM)	
Compound		HCT-116	HCT-116/VM46
	10-Desacetoxytaxol	0.008	1.22 (153)
15	Taxol	0.004	0.440 (124)

* Value in parenthesis is fold resistance relative to HCT-116 cells.

Compounds of formula (1) of the instant invention are useful for inhibiting tumor growth in animals including humans and are preferably administered in the form of a pharmaceutical composition comprising an effective anti-tumor amount of compound of the instant invention in combination with a pharmaceutically acceptable carrier or diluent.

25 Antitumor compositions herein may be made up in any suitable form appropriate for desired use; e.g., oral, parenteral or topical administration. Examples of

parenteral administration are intramuscular, intravenous, intraperitoneal, rectal and subcutaneous administration.

The diluent or carrier ingredients should not be such as to diminish the therapeutic effects of the antitumor compounds.

Suitable dosage forms for oral use include tablets, dispersible powders, granules, capsules, suspensions, syrups, and elixirs. Inert diluents and carriers for tablets include, for example, calcium carbonate, sodium carbonate, lactose and talc. Tablets may also contain granulating and disintegrating agents such as starch and alginic acid, binding agents such as starch, gelatin and acacia, and lubricating agents such as magnesium stearate, stearic acid and talc. Tablets may be uncoated or may be coated by unknown techniques; e.g., to delay disintegration and absorption. Inert diluents and carriers which may be used in capsules include, for example, calcium carbonate, calcium phosphate and kaolin. Suspensions, syrups and elixirs may contain conventional excipients, for example, methyl cellulose, tragacanth, sodium alginate; wetting agents, such as lecithin and polyoxyethylene stearate; and preservatives, e.g., ethyl-p-hydroxybenzoate.

Dosage forms suitable for parenteral administration include solutions, suspensions, dispersions, emulsions and the like. They may also be manufactured in the form of sterile solid compositions which can be dissolved or suspended in sterile injectable medium immediately before use. They may contain suspending or dispersing agents known in the art.


One aspect of the invention herein is directed to therapeutically inhibiting tumor growth in an animal host having a tumor sensitive to the compounds of the instant invention which comprises administering to said host an

effective antitumor dose of said compound. It will be appreciated that the actual preferred amount of compound of the instant invention will vary according to the particular compound, the particular composition formulated, the mode of application and the particular situs, host and disease being treated. Many factors that modify the action will be taken into account by those skilled in the art; e.g., age, body weight, sex, diet, time of administration, route of administration, rate of excretion, condition of the host, drug combinations, reaction sensitivities and severities and severity of disease. Administration can be carried out continuously or periodically within the maximum tolerated dose. Optimal application rates for a given set of conditions can be ascertained by those skilled in the art using conventional dosage administration tests in view of the above guidelines.

The following examples are provided in order to more fully illustrate the present invention.

EXAMPLE 1

20 2',7-bis'O-(2,2,2-trichloroethoxycarbonyl)-10-deacetyl taxol (12).



(12)


10-Deacetyl taxol (140 mg, 0.173 mmol) in dry dichloro-
methane (3.5 mL) was treated at 0°C with pyridine (0.028
mL, 0.346 mmol) and trichloroethyl chloroformate (0.0724
mL, 0.260 mmol). After 1h at this temperature, the cold
5 bath was removed and the mixture was stirred at room
temperature overnight. The solvent was evaporated and the
residue chromatographed on silica gel (30-50% ethyl acetate
in hexane) to afford 12 as a foam (92.3 mg, 46%). Further
elution afforded unreacted starting material (35 mg, 25%).
10 Carbonate 12 had NMR (300 MHz, CDCl₃) δ 8.14 (d, J=8.5 Hz,
2H) 7.75 (d, J=8.5 Hz, 2H) 7.65-7.35 (m, 11H) 6.94 (exch.d,
J=9.3 Hz, 1H) 6.27 (br t, 1H) 6.04 (dd, J=9.3 Hz, J'=2.6
Hz, 1H) 5.71 (d, J=6.9 Hz, 1H) 5.54 (d, J=2.6 Hz, 1H)
5.43-5.37 (m, 2H) 4.96 (d, J=7.9 Hz, 1H) 4.85-4.67 (m, 4H)
15 4.29 (AB q, 2H) 4.04-4.01 (m, 2H) 2.69- 1.80 (m, 14H incl.
singlets at 2.58, 1.96, 1.89, 3H each) 1.20 (s, 3H) 1.09
(s, 3H).

High Res. Mass Spectrum: Calcd for C₅₁H₅₁NO₁₇Cl₆K (MK+)
1198.0925, found 1198.0949.

20

EXAMPLE 2

2',7-bis'O-(2,2,2-trichloroethoxycarbonyl)-10-desacetoxy-
11,12-dihydrotaxol-10,12(18)-diene (13).




(13)

Bis-carbonate 12 (92.3 mg, 0.079 mmol) in dry dichloromethane (2 mL) was treated at room temperature with TFCT (0.0384 mL, 0.238 mmol). The solution was stirred overnight. The solvent was evaporated and the residue
5 purified by column chromatography (25% ethyl acetate in hexane) to afford 13 as a white powder (42.8 mg, 47.3%).
NMR (300 MHz, CDCl₃) δ 8.18 (d, J=8.5 Hz, 2H) 7.69 (d, J=8.5 Hz, 2H) 7.58-7.28 (m, 11H) 6.90 (exch. d, J=9.7 Hz, 1H) 6.31 (s, 1H) 6.25 (br t, 1H) 6.04 (dd, J=9.7 Hz, J'=2.2
10 Hz, 1H) 5.80 (d, J=7.8 Hz, 1H) 5.51 (d, J=2.2 Hz, 1H) 5.38 (d, J=2.3 Hz, 1H) 5.31 (dd, J= 11.0 Hz, J'=7.3 Hz, 1H) 4.99 (d, J=2.3 Hz, 1H) 4.97 (br d, 1H) 4.80-4.65 (m, 4H) 4.30 (AB q, 2H) 3.75 (d, J=7.8 Hz, 1H) 2.60 (m, 1H) 2.55 (s, 3H) 2.16-2.09 (m, 3H) 1.83 (s, 3H) 1.77 (s, 1H) 1.13 (s, 3H)
15 1.07 (s, 3H).
High Res. Mass Spectrum: Calcd for C₅₁H₄₉NO₁₆Cl₆K (MK+) 1180.0820, found 1180.0777.

EXAMPLE 3

10-Desacetoxy-11,12-dihydrotaxol-10,12(18)-diene (14).



(14)

Dienone 13 (180 mg, 0.157 mmol) was dissolved in methanol (3 mL) and treated with acid-washed Zn dust (300 mg, 4.72 mmol). The slurry was refluxed for 20 min, filtered, and


the filtrate evaporated. Chromatography of the residue (40-60% ethyl acetate in hexane) gave 14 as a foam (18 mg, 14%), together with its 7-epi isomer (97 mg, 77.7%).

Compound 14 had NMR (300 MHz, CDCl₃) δ 8.19 (d, J=8.5 Hz, 2H) 7.69 (d, J=8.5 Hz, 2H) 7.59-7.25 (m, 11H) 7.00 (exch. d, J=10.4 Hz, 1H) 6.20 (br t, 1H) 5.96 (s, 1H) 5.83-5.77 (m, 2H) 5.18 (d, J=2.2 Hz, 1H) 4.94 (dd, J=7.2 Hz, J'=2.1 Hz, 1H) 4.79 (d, J=2.2 Hz, 1H) 4.72 (dd, J=4.3 Hz, J'=2.2 Hz, 1H) 4.30 (AB q, 2H) 4.00 (m, 1H) 3.65 (d, J=7.8 Hz, 1H) 3.51 (exch. d, J=4.3 Hz, 1H) 2.60 (m, 1H) 2.40 (s, 3H) 2.30-1.80 (m, 5H) 1.70 (s, 3H) 1.16 (s, 3H) 1.06 (s, 3H). High Res. Mass Spectrum: Calcd for C₄₅H₄₈NO₁₂ (MH⁺) 794.3177, found 794.3152.

Alternatively, dienone 13 (39 mg, 0.034 mmol) was dissolved in methanol (0.5 mL) and acetic acid (0.5 mL), and treated with acid-washed zinc dust (66.4 mg, 1.020 mmol). The slurry was heated at 40°C for 1h, filtered and the filtrate evaporated. Chromatography of the residue with 60% ethyl acetate/hexane gave 14 as a foam (22 mg, 81%). Spectral data are the same as the previous ones.

EXAMPLE 4

10-Desacetoxytaxol (15).




(15)

Dienone 14 (22 mg, 0.028 mmol) in ethyl acetate (0.7 mL) was hydrogenated at atmospheric pressure in the presence of palladium on charcoal (10%, 14.7 mg, 0.014 mmol Pd) After 5.5 h at RT, filtration (rinsing with ethyl acetate),
 5 evaporation and chromatography (60% ethyl acetate in hexane) gave 10-deoxytaxol 15 (15.0 mg, 68%) as a white foam. NMR (CDCl₃) δ 8.12(d, J=8.5 Hz, 2H) 7.61 (d, J=8.4 Hz, 2H) 7.60-7.27 (m, 11H) 7.06 (exch. d, J=8.9 Hz, 1H) 6.09 (br t, 1H) 5.78 (dd, J= 8.9 Hz, J'=2.6 Hz, 1H) 5.66
 10 (d, J=6.8 Hz, 1H) 4.91 (d, J=7.6 Hz, 1H) 4.76 (dd, J= 5.1 Hz, J'= 2.6 Hz, 1H) 4.29-4.20 (m, 3H) 4.01 (d, J=6.8 Hz, 1H) 3.75 (d, J=15.8 Hz, 1H) 3.60 (exch. d, J=5.1 Hz, 1H) 3.39 (br d, 1H) 2.60 (m, 1H) 2.34 (s, 3H) 2.34-2.22 (m, 2H) 1.90- 1.71 (m, 2H) 1.62 (s, 3H) 1.61 (s, 3H) 1.53 (exch. d,
 15 J=7.8 Hz, 1H) 1.14 (s, 3H) 1.12 (s, 3H).
 High Res. Mass Spectrum: Calcd for C₄₅H₅₀NO₁₂ (MH⁺) 796.3333, found 796.3319.

EXAMPLE 5

10-Desacetoxy-7-O-triethylsilyl baccatin III (16).



(16)

A. 10-desacetyl-7-O-triethylsilyl baccatin III (Compound A).

10-Desacetyl baccatin III (from *Taxus baccata*, 628.0 mg, 1.150 mmol) was dissolved in dry DMF (6 mL), cooled to 0°C,
 25 and treated with imidazole (312.8 mg, 4.595 mmol) and

chlorotriethylsilane (0.772 mL, 4.60 mmol). The mixture was stirred at 0°C for 4 h, then diluted with ethyl acetate (150 mL) and washed exhaustively with water and brine. The organic layer was dried and concentrated. The residue
5 was chromatographed (silica, 50% ethyl acetate in hexane), to afford compound A as a foam (586 mg, 77%). This compound is described by Greene et al. in J. Am. Chem. Soc., 1988, 110, 5917.

10 B. 10-desacetyl-7-*O*-triethylsilyl-10-*O*-(pentafluorophenylthiocarbonyl) baccatin III (Compound B).


Compound A (319 mg, 0.485 mmol) was dissolved in dry THF (5 mL), cooled to -40°C, and treated with *n*-butyllithium (1.58M in hexanes, 0.384 mL, 0.606 mmol). After 40 min at this temperature, pentafluorophenyl chlorothionoformate
15 (0.086 mL, 0.536 mmol) was added neat by syringe. The reaction mixture was stirred at -20°C for 90 min, then quenched with saturated ammonium chloride solution, and extracted with ethyl acetate. The ethyl acetate layer was dried, evaporated and the residue chromatographed (silica,
20 40% ethyl acetate in hexane) to afford Compound B as a foam (320 mg, 74%). NMR (CDCl₃) δ 8.09 (d, 2H) 7.56 (t, 1H) 7.44 (m, 2H) 6.78 (s, 1H) 5.64 (d, J=6.9 Hz, 1H) 4.96-4.89 (m, 2H) 4.49 (dd, J=10.2 Hz, J'=6.6 Hz, 1H) 4.12 (AB q, 2H) 3.80 (d, J=6.9 Hz, 1H) 2.55-0.44 (m, 43H).
25 Mass Spectrum: 884 (MH⁺).

C. 10-desacetoxy-7-*O*-triethylsilyl baccatin III.

Thionocarbonate (compound B, 119 mg, 0.135 mmol) was dissolved in dry toluene (3 mL) and treated with 2,2'-azobisisobutyronitrile (AIBN, 2 mg). The solution was
30 degassed with dry nitrogen, then tributyltin hydride (0.055 mL, 0.202 mmol) was added and the solution was heated for 1

h (90°C). Solvent evaporation and chromatography (silica, 40% ethyl acetate in hexane) gave the title product (87 mg, 99%) as a colorless foam. NMR (CDCl₃) δ 8.07 (d, J=8.2 Hz, 2H) 7.56 (br t, 1H) 7.44 (m, 2H) 5.57 (d, J=6.7 Hz, 1H) 4.92 (d, J=9.3 Hz, 1H) 4.78 (br s, 1H) 4.48 (dd, J=10.4 Hz, J'=6.6 Hz, 1H) 4.09 (AB q, 2H) 4.06 (d, J=6.7 Hz, 1H) 3.74 (d, J=14.8 Hz, 1H) 3.35 (br d, 1H) 2.44 (m, 1H) 2.25 (s, 3H) 2.22-0.45 (m, 42H).
Mass spectrum: 642 (MH⁺).

10

EXAMPLE 610-Deoxytaxotere (17).

(17)

Compound 16 (100 mg, 0.156 mmol) was placed in a flask under Argon and dissolved in dry THF (1.5 mL). Upon cooling to -40°C, n-butyllithium (1.45M in hexanes, 0.119 mL, 0.170 mmol) was added dropwise, followed by (3R,4S)-1-tert-butoxycarbonyl-4-phenyl-3-triethylsilyloxy-2-azetidinone 94.2 mg, 0.25 mmol) in THF (0.5 mL) over a period of 2 min. The mixture was immediately warmed to 0°C and stirred for 45 min before quenching with saturated ammonium chloride (3 mL). The mixture was extracted with ethyl acetate, dried, and evaporated. Silica gel chromatography (30% ethyl acetate in hexane) afforded 2',-7-bis-Q-(triethylsilyl)-10-deoxytaxotere as a foam (125

mg. 76%). This compound (100mg, 0.098 mmol) was immediately dissolved in acetonitrile (2 mL) at -5°C and treated with hydrochloric acid (0.037 mL, 36%, 12M). The mixture was stirred for 2h at -5°C, then it was quenched with aqueous bicarbonate, extracted with ethyl acetate, and dried. Evaporation was followed by silica chromatography (75% ethyl acetate in hexane) to afford the title compound as a foam (80.5mg, 80%). NMR (CDCl₃) δ 8.10 (d, J=8.2 Hz, 2H) 7.64-7.29 (m, 8H) 6.11 (br t, 1H) 5.68 (d, J=6.9 Hz, 1H) 5.43 (br d, 1H) 5.25 (br d, 1H) 4.93 (d, J=7.7 Hz, 1H) 4.60 (br s, 1H) 4.30-4.18 (m, 3H) 4.02 (d, J=7.7 Hz, 1H) 3.80 (d, J=15.8 Hz, 1H) 3.46-3.40 (m, 2H) 2.62 (m, 1H) 2.35 (s, 3H) 2.35-2.25 (m, 2H) 1.89-1.65 (m, 5H) 1.63 (s, 3H) 1.35 (s, 9H) 1.19 (s, 3H) 1.16 (s, 3H).

15

EXAMPLE 7

Preparation of (3R,4S)-1-tertbutoxycarbonyl-4-phenyl-3-triethylsilyloxy-2-azetidinone (see Scheme V).

(L)-Threonine methyl ester hydrochloride (1.26 g, 7.44 mmol) in anhydrous dichloromethane (15 mL) was stirred with imidazole (1.01 g, 14.89 mmol) and t-butoxydiphenylsilyl chloride (2.274 g, 7.816 mmol) for 16 h at room temperature. The reaction mixture was partitioned between water and dichloromethane. The organic phase was washed with 5% aqueous sodium bicarbonate and water, dried and concentrated to give 2.88 g of a crude oil, which was used directly in the next step; ¹H-NMR (CDCl₃) δ: 7.70-7.25 (m, 10H) 4.44 (m, 1H) 3.62 (s, 3H) 3.31 (d, J=3 Hz, 1H) 2.12 (bs, 2H) 1.3-1.15 (m, 12H).

This oil (548 mg, 1.414 mmol) in anhydrous dichloromethane (10 mL) was treated with benzaldehyde (0.158 mL, 1.55 mmol) at rt overnight in the presence of 4Å molecular sieves to afford compound of formula XVIIa in

situ. Upon cooling the solution containing compound XVIIa to -40°C , triethylamine (0.20 mL, 1.698 mmol) was added, followed by acetoxyacetyl chloride (XVIa) (0.182 mL, 1.698 mmol) over 10 min. The mixture was allowed to reach rt
5 over 4 h and the product was partitioned between dichloromethane and water. The organic phase was further washed with water and brine, dried and concentrated. Silica gel chromatography (being eluted with 1:4 EtOAc/hexane) gave 411 mg of compound XVIIIa as a ca. 10:1 mixture of 3R,4S :
10 3S,4R diastereomers.

This mixture of diastereomers (245.1 mg, 0.414 mmol) in dry THF (2 mL) was treated with acetic acid (0.15 mL) and tetrabutylammonium fluoride (TBAF, 1M in THF, 1.20 mL). The solution was stirred for 14 h at rt, then
15 partitioned between ethyl acetate and 5% aqueous sodium bicarbonate. The organic phase was dried and concentrated. Flash silica gel chromatography using 1:1 ethyl acetate : hexane as eluant gave 66 mg (Y: 50%) of compound XIXa as a foam in one diastereomer; $^1\text{H-NMR}$ (CDCl_3) δ : 7.42–7.25 (m, 5H) 5.90 (d, $J=4.8$ Hz, 1H) 5.09 (d, $J=4.8$ Hz, 1H) 4.28 (m, 1H) 4.01 (d, $J=4.8$ Hz, 1H) 3.70 (s, 3H) 1.73 (s, 3H) 1.19 (d, $J=6.6$ Hz, 3H).
20

Compound of formula XIXa (9.8 g, 0.0305 mol) in dry dichloromethane (100 mL) was treated at -78°C with
25 triethylamine (9.40 mL, 0.0671 mol) and methanesulfonyl chloride (MsCl , 3.50 mL, 0.0457 mol). The solution was allowed to reach rt overnight. The reaction mixture was partitioned between water and dichloromethane. The organic layer was washed with 5% aqueous sodium bicarbonate, dilute
30 aqueous HCl , water and brine, and concentrated to afford compound XXa as a crude oily residue. The crude residue (10.0 g) was dissolved in dichloromethane (250 mL) and ozonized at -78°C until the color of the solution stayed as blue. Addition of methyl sulfide (11 mL) and concentration

of the reaction mixture gave compound of formula XXIIa (crude).

The compound of formula XXIIa was dissolved in THF (150 mL) and treated at -78°C with hydrazine hydrate (10 mL). After 2 h, the mixture was poured into dilute aqueous HCl and ethyl acetate, and the two phases were separated. The organic phase was washed with more acid, water and brine and concentrated to afford a crude product, which was purified by silica gel chromatography using 1-5% methanol in methylene chloride to yield 4.40 g (Y: 71%) of compound of formula XXIIa; ¹H-NMR (CDCl₃) δ: 7.38-7.24 (m, 5H) 6.31 (bs, 1H) 5.87 (bm, 1H) 5.04 (d, J=4.8 Hz, 1H) 1.67 (s, 3H).


To a cooled (-5°C) mixture of 1M aqueous KOH (140 mL) and acetonitrile (100 mL), a solution of compound XXIIa (2.39 g, 11.22 mmol) in acetonitrile (130 mL) was added dropwise. The mixture was stirred at 0°C for 1 h and diluted with ethyl acetate (300 mL), water (50 mL) and saturated aqueous bicarbonate (50 mL). The organic phase was separated, and the aqueous layer further extracted with ethyl acetate (3x200 mL). The organic phases were combined, dried, filtered and concentrated to give compound of formula XXIIIa (crude), which was recrystallized from hexane/acetone (mp, 184-6°C); yield, 1.53 g (Y: 82%).

To azetidinone XXIIIa (580 mg, 3.55 mmol) in dry THF (5.0 mL) was added imidazole (265.5 mg, 3.90 mmol), followed by triethylsilyl chloride (0.654 mL, 3.90 mmol). The mixture was allowed to be stirred for 1 h. Ethyl acetate was added and the organic layer was washed with brine, 10% aqueous HCl and dried. Silica gel chromatography (being eluted with 25% ethyl acetate in hexane) gave 670 mg (Y: 68%) of compound XXIVa as a foam.

To a stirred solution of compound XXIVa (2.20 g, 7.92 mmol) in dry THF (25 mL) was added ethyl diisopropylamine (1.65 mL, 9.51 mmol) at 0°C under argon atmosphere. The solution was stirred for 5 min, then di-tert-butyl

carbonate (2.08 g, 9.51 mmol) and 4-dimethylaminopyridine (193.6 mg, 1.58 mmol) were added. The reaction mixture was stirred at 0°C for 60 min. The reaction was diluted with ethyl acetate (25 mL), and the mixture was washed with
5 brine, 10% aqueous sodium bicarbonate, 10% aqueous HCl, dried over magnesium sulfate, and concentrated to leave an oil. Silica gel flash chromatography (being eluted with 15% ethyl acetate in hexane) gave 2.40 g (Y: 83%) of compound XVa as a white solid; ¹H-NMR (CDCl₃) δ 7.28 (m,
10 5H) 5.03 (m, 2H) 1.38 (s, 9H) 0.76 (t, J = 7.56, 9H) 0.43 (m, 6H).

SCHEME V




In view of the above, it will be seen that the several objects of the invention are achieved.

As various changes could be made in the above compositions and processes without departing from the scope of the invention, it is intended that all matter contained in the above description be interpreted as illustrative and not in a limiting sense.

CLAIMS

We Claim:

1. A compound of the formula



- wherein R_2 is hydrogen, hydroxy or a protected hydroxy group; R_3 and R_4 are independently hydrogen, hydroxy, a
 5 protected hydroxy group, methyl, $-SH$, $-NH_2$, or $-NR_8R_9$; R_5 is R_{10} , or $-OR_{10}$; R_6 and R_7 are independently hydrogen, alkyl, or aryl; R_8 and R_9 are independently hydrogen, alkyl, alkenyl, alkynyl, or aryl; and R_{10} is alkoxy, alkyl, alkenyl, alkynyl, or aryl; provided that when R_2 is
 10 hydrogen, R_3 and R_4 are independently hydrogen, methyl, $-SH$, $-NH_2$, or $-NR_8R_9$.

2. A compound as set forth in claim 1 wherein R_3 is hydrogen and R_4 is hydroxy or a protected hydroxy group, R_5 is phenyl or t-butoxy, R_6 is hydrogen, and R_7 is phenyl.

3. A compound as set forth in claim 2 wherein R_5 is phenyl.

4. A compound as set forth in claim 2 wherein R_5 is t-butoxy.

5. A compound as set forth in claim 1 wherein R_3 is hydrogen and R_4 is $-NH_2$.

6. A compound as set forth in claim 5 wherein R_5 is phenyl or t-butoxy.

7. A compound as set forth in claim 1 wherein R_3 is methyl, R_4 is $-SH$, R_6 is aryl, and R_7 is methyl.

8. A compound as set forth in claim 7 wherein R_5 is phenyl or t-butoxy.

9. A compound as set forth in claim 1 wherein R_2 , R_3 and R_6 are hydrogen; R_4 is $-NR_8R_9$; R_5 is R_{10} and R_7 is phenyl.

10. A compound as set forth in claim 9 wherein R_8 is hydrogen, R_9 is an amine protecting group and R_{10} is t-butoxy.

11. A compound as set forth in claim 1 wherein R_2 is hydroxy.

12. A compound as set forth in claim 11 wherein R_3 is hydrogen and R_4 is hydroxy or a protected hydroxy group, R_5 is phenyl or t-butoxy, R_6 is hydrogen, and R_7 is phenyl.

13. A compound as set forth in claim 11 wherein R_3 is hydrogen and R_4 is $-NH_2$.

14. A compound as set forth in claim 13 wherein R_5 is phenyl or t-butoxy.

15. A compound as set forth in claim 11 wherein R_3 is methyl, and R_4 is -SH.

16. A compound as set forth in claim 15 wherein R_5 is phenyl or t-butoxy.

17. A compound as set forth in claim 11 wherein R_3 and R_6 are hydrogen, R_4 is $-NR_8R_9$, R_5 is phenyl or t-butoxy, and R_7 is phenyl.

18. A compound as set forth in claim 17 wherein R_8 is hydrogen and R_9 is an amine protecting group.

19. The compound 10-desacetoxytaxol.

20. The compound 10-deoxytaxotere.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US92/07935**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) :C07D 305/14

US CL :549/214, 510, 511

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 549/214, 510, 511

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS structure search.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	J. Org. Chem. 1991, Kingston et al., "Modified Taxols. Reaction Of Taxol With Electrophilic Reagents And Preparation Of A Rearranged Taxol Derivative With Tubulin Assembly Activity." Vol. 56, No. 17, pp. 5114 to 5119.	1-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be part of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 NOVEMBER 1992

Date of mailing of the international search report

07 DEC 1992


Name and mailing address of the ISA/ US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

BA TRINH

Telephone No. (703) 308-1235


 NGUYEN NGOC-HO
 INTERNATIONAL DIVISION