

(19) **DANMARK**

(10) **DK/EP 3590949 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 07 H 21/04 (2006.01)** **C 12 N 15/09 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-07-11**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2022-05-18**
- (86) Europæisk ansøgning nr.: **19177059.3**
- (86) Europæisk indleveringsdag: **2011-10-03**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-01-08**
- (30) Prioritet: **2010-10-01 US 40441310 P**
- (62) Stamansøgningsnr: **11830061.5**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **ModernaTX, Inc., 200 Technology Square, Cambridge, MA 02139, USA**
- (72) Opfinder: **SCHRUM, Jason P, 20 Watertown Street, Unit 129, Watertown, MA 02472-2580, USA**
Siddiqi, Suhaib, 37 University Avenue, Burlington, MA 01803, USA
EJEBE, Kenechi, 3 East 101 Street, Apt 6I, New York, NY 10029, USA
- (74) Fuldmægtig i Danmark: **Budde Schou A/S, Dronningens Tværgade 30, 1302 København K, Danmark**
- (54) Benævnelse: **RIBONUKLEINSYRER INDEHOLDENDE N1-METHYL-PSEUDOURACIL OG ANVENDELSE HERAF**
- (56) Fremdragne publikationer:
WO-A2-2007/024708
KARIKÓ KATALIN ET AL: "Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability", MOLECULAR THERAPY, NATURE PUBLISHING GROUP, GB, vol. 16, no. 11, 1 November 2008 (2008-11-01), pages 1833-1840, XP002598556, ISSN: 1525-0024, DOI: 10.1038/MT.2008.200 [retrieved on 2008-09-16]
KARIKÁ, KATALIN ET AL: "Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability", MOLECULAR THERAPY, NATURE PUBLISHING GROUP, GB, vol. 16, no. 11, 1 November 2008 (2008-11-01), pages 1833-1840, XP002598556, ISSN: 1525-0024, DOI: 10.1038/MT.2008.200 [retrieved on 2008-09-16]

DESCRIPTION

CLAIM OF PRIORITY

[0001] Priority is claimed from U.S. Provisional Application Serial No. 61/404,413, filed on October 1, 2010.

BACKGROUND

[0002] Naturally occurring RNAs are synthesized from four basic ribonucleotides: ATP, CTP, UTP and GTP, but may contain post-transcriptionally modified nucleotides. Further, approximately one hundred different nucleoside modifications have been identified in RNA (Rozenski, J, Crain, P, and McCloskey, J. (1999). The RNA Modification Database: 1999 update. Nucl Acids Res 27: 196-197). The role of nucleoside modifications on the immunostimulatory potential, stability, and on the translation efficiency of RNA, and the consequent benefits to this for enhancing protein expression and producing therapeutics however, is unclear.

[0003] There are multiple problems with prior methodologies of effecting protein expression. For example, heterologous deoxyribonucleic acid (DNA) introduced into a cell can be inherited by daughter cells (whether or not the heterologous DNA has integrated into the chromosome) or by offspring. Introduced DNA can integrate into host cell genomic DNA at some frequency, resulting in alterations and/or damage to the host cell genomic DNA. In addition, multiple steps must occur before a protein is made. Once inside the cell, DNA must be transported into the nucleus where it is transcribed into RNA. The RNA transcribed from DNA must then enter the cytoplasm where it is translated into protein. This need for multiple processing steps creates lag times before the generation of a protein of interest. Further, it is difficult to obtain DNA expression in cells; frequently DNA enters cells but is not expressed or not expressed at reasonable rates or concentrations. This can be a particular problem when DNA is introduced into cells such as primary cells or modified cell lines.

[0004] There is a need in the art for biological modalities to address the modulation of intracellular translation of nucleic acids.

[0005] WO 2007/024708 A2 relates to RNA comprising certain modified nucleosides, particularly pseudouridine, which was found to reduce the immunogenicity of RNA.

SUMMARY

[0006] The present invention is defined in the claims and thus provides a method of

synthesizing an mRNA, comprising the steps of:

1. a) providing a complementary deoxyribonucleic acid (cDNA) that encodes a protein of interest;
2. b) selecting a nucleotide that is known to disrupt binding of a major groove binding partner with a nucleic acid, wherein the nucleotide has decreased binding affinity to the major groove binding partner;
3. c) contacting the provided cDNA and the selected nucleotide with an RNA polymerase under conditions such that the mRNA is synthesized,

wherein the nucleotide comprises N1-methyl-pseudouridine, and


wherein 100% of nucleotides comprising uracil in the RNA are replaced with nucleotides comprising N1-methyl-pseudouridine.

[0007] The invention also provides an mRNA wherein 100% of nucleotides comprising uracil in the mRNA are replaced with nucleotides comprising N1-methyl-pseudouridine.

[0008] The present disclosure relates, *inter alia*, to modified nucleosides, modified nucleotides, and modified nucleic acids which can exhibit a reduced innate immune response when introduced into a population of cells, both *in vivo* and *ex vivo*. Further, these modified nucleosides, modified nucleotides, and modified nucleic acids described herein can disrupt binding of a major groove interacting partner with the nucleic acid. Because of the reduced immunogenicity and the decrease in major groove interactions, these modified nucleosides, modified nucleotides, and modified nucleic acids can be more efficient during protein production than, e.g., unmodified nucleic acids.

[0009] Thus, the present disclosure relates to compounds comprising nucleotides that can disrupt binding of a major groove binding partner with a nucleic acid, wherein the nucleotide has decreased binding affinity to the major groove binding partner.


[0010] The present disclosure further relates to compounds having Formula I:




I

wherein constituent variables are provided herein.

[0011] The present disclosure further relates to nucleic acid sequences of at least two nucleotides comprising a compound of Formula I-d:





I-d

wherein constituent variables are provided herein.

[0012] The present disclosure further relates to compositions comprising at least one compound of Formula I.

[0013] The present disclosure further relates to pharmaceutical compositions comprising a compound of Formula I.

[0014] The present disclosure further relates to methods of preparing nucleic acid sequences of at least two nucleotides of a compound of Formula I-d.

[0015] The present disclosure further relates to methods of amplifying nucleic acid sequences of at least two nucleotides of a compound of Formula I-d.

[0016] The present disclosure further relates to kits comprising a compound of Formula I.

[0017] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use relating to the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0018] In case of conflict with references mentioned herein, the present specification including definitions, will control.

BRIEF DESCRIPTION OF DRAWINGS

[0019]

FIGs. 1A and 1B depict images of non-denaturing agarose gels of each *in vitro*-transcribed modified RNA.

FIGs. 2A and 2B depict images of an Enzyme-linked immunosorbent assay (ELISA) for Human Granulocyte-Colony Stimulating Factor (G-CSF) of *in vitro* transfected Human Keratinocyte cells with each indicated modRNA encoding human G-CSF and the line indicates a saturating level of maximum detectable limit of secreted G-CSF in the assay.

FIGs. 3A-N depict line graphs of a series of Enzyme-linked immunosorbent assays (ELISA) for Human Granulocyte-Colony Stimulating Factor (G-CSF) secreted from *in vitro*-transfected Human Keratinocyte cells at different time points with each indicated human G-CSF-encoding modRNA at the indicated doses. The line indicates a saturating level of maximum detectable limit of secreted G-CSF in the assay.

FIGs. 4A and 4B depict bar graphs of a series of Enzyme-linked immunosorbent assays (ELISA) for endogenous cellular human Tumor Necrosis Factor- α (TNF- α) secreted from *in vitro*-transfected Human Keratinocyte cells at 24 hours with each indicated hu-G-CSF-encoding modRNA at increasing doses.

FIGs. 4C and 4D depict bar graphs of a series of Enzyme-linked immunosorbent assays (ELISA) for endogenous cellular human Interferon- β (IFN- β) secreted from *in vitro*-transfected Human Keratinocyte cells at 24 hours with each indicated hu-G-CSF-encoding modRNA at increasing doses.

FIGs. 4E and 4F depict bar graphs of a series of Enzyme-linked immunosorbent assays (ELISA) for human-G-CSF secreted from *in vitro*-transfected Human Keratinocyte cells at 24 hours with each indicated hu-G-CSF-encoding modRNA at increasing doses.

FIG. 5A is a table showing results from an Enzyme-linked immunosorbent assay (ELISA) for human-G-CSF secreted from *in vitro*-transfected Human Keratinocyte cells sampled from individual wells in a co-culture 24-well tissue culture plate 42 hours post-transfection with 750ng of each indicated hu-G-CSF-encoding modRNA.

FIG. 5B depicts an image of an agarose gel of RT-PCR hu-G-CSF modRNA products from co-culture cell extracts 42 hours post-transfection of the human keratinocyte feeder layer with hu-G-CSF modRNA and the un-transfected Kasumi-1 and KG-1 insert culture cells.

FIGs. 5C and 5D depict graphs of results from a hu-G-CSF-modRNA-induced cell proliferation assay of Kasumi-1 (FIG. 5C) and KG-1 (FIG. 5D) cells normalized to untransfected cells. Hu-G-CSF modRNA identity transfected into human keratinocyte feeder cells is indicated.

FIGs. 6A-L depict graphs of the UV absorbance spectra for exemplary modRNA molecules that incorporate the indicated modified nucleotide.

DETAILED DESCRIPTION

[0020] The present disclosure relates, *inter alia*, to modified nucleosides, modified nucleotides, and modified nucleic acids that exhibit a reduced innate immune response when introduced into a population of cells. The modified nucleosides, modified nucleotides, and modified nucleic acids can be chemically modified on the major groove face, thereby disrupting major groove binding partner interactions, which cause innate immune responses.

[0021] In general, exogenous unmodified nucleic acids, particularly viral nucleic acids, introduced into cells induce an innate immune response, resulting in cytokine and interferon (IFN) production and cell death. However, it is of great interest for therapeutics, diagnostics, reagents and for biological assays to deliver a nucleic acid, e.g., a ribonucleic acid (RNA) inside a cell, either *in vivo* or *ex vivo*, such as to cause intracellular translation of the nucleic acid and production of the encoded protein. Of particular importance is the delivery and function of a non-integrative nucleic acid, as nucleic acids characterized by integration into a target cell are generally imprecise in their expression levels, deleteriously transferable to progeny and neighbor cells, and suffer from the substantial risk of causing mutation. Described herein in part are nucleic acids encoding useful polypeptides capable of modulating a cell's function and/or activity, and methods of making and using these nucleic acids and polypeptides. As described herein, these nucleic acids are capable of reducing the innate immune activity of a population of cells into which they are introduced, thus increasing the efficiency of protein production in that cell population. Further, one or more additional advantageous activities and/or properties of the nucleic acids and proteins of the present disclosure are described.

[0022] Further, the modified nucleosides, modified nucleotides, and modified nucleic acids described herein can be modified on the major groove face. These major groove modifications can allow for alterations, e.g. a decrease, in the interaction of the modified nucleosides, modified nucleotides, and modified nucleic acids with a binding groove partner.

[0023] Accordingly, in a first aspect, the present disclosure describes compounds comprising a nucleotide that can disrupts binding of a major groove interacting, e.g. binding, partner with a nucleic acid, wherein the nucleotide has decreased binding affinity to major groove interacting, e.g. binding, partners.

[0024] In another aspect, the present disclosure describes compounds comprising a nucleotide that contains chemical modifications, wherein the nucleotide can have altered binding to major groove interacting, e.g. binding, partners.

[0025] In some cases , the chemical modifications are located on the major groove face of the nucleobase, and wherein the chemical modifications can include replacing or substituting an atom of a pyrimidine nucleobase with an amine, an SH, an alkyl (e.g., methyl or ethyl), or a halo (e.g., chloro or fluoro).

[0026] In some cases , the chemical modifications can be located on the major groove face of the nucleobase, and wherein the chemical modification can include replacing or substituting an atom of a pyrimidine nucleobase with an amine, an SH, a methyl or ethyl, or a chloro or fluoro.

[0027] In some cases , the chemical modifications can be located on the sugar moiety of the nucleotide.

[0028] In some cases , the chemical modifications can be located on the phosphate backbone of the nucleotide.

[0029] In some cases , the chemical modifications can alter the electrochemistry on the major groove face of the nucleotide.

[0030] In another aspect, the present disclosure relates to nucleotides that contain chemical modifications, wherein the nucleotide reduces the cellular innate immune response, as compared to the cellular innate immune induced by a corresponding unmodified nucleic acid.

[0031] In another aspect, the present disclosure relates to nucleic acid sequences comprising at least two nucleotides, the nucleic acid sequence comprising a nucleotide that disrupts binding of a major groove interacting partner with the nucleic acid sequence, wherein the nucleotide has decreased binding affinity to the major groove binding partner.

[0032] In another aspect, the present disclosure relates to compositions comprising a compound as described herein.

[0033] In some cases , the composition is a reaction mixture.

[0034] In some cases , the composition is a pharmaceutical composition.

[0035] In some cases , the composition is a cell culture.

[0036] In some cases , the compositions further comprise an RNA polymerase and a cDNA template.

[0037] In some cases , the compositions further comprise a nucleotide selected from the group consisting of adenosine, cytosine, guanosine, and uracil.

[0038] In a further aspect, the present disclosure relates to methods of synthesizing a pharmaceutical nucleic acid, comprising providing a complementary deoxyribonucleic acid (cDNA) that encodes a pharmaceutical protein of interest; selecting a nucleotide that is known to disrupt a binding of a major groove binding partner with a nucleic acid, wherein the nucleotide has decreased binding affinity to the major groove binding partner; and contacting the provided cDNA and the selected nucleotide with an RNA polymerase, under conditions such that the pharmaceutical nucleic acid is synthesized.

[0039] In some cases , the pharmaceutical nucleic acid is a ribonucleic acid (RNA).

[0040] In a further aspect, the present disclosure relates to methods of making a pharmaceutical formulation comprising a physiologically active secreted protein, comprising transfecting a first population of human cells with a pharmaceutical nucleic acid made by the methods described herein, wherein the secreted protein is active upon a second population of

human cells.

[0041] In some cases , the secreted protein is capable of interacting, e.g. binding, with a receptor on the surface of at least one cell present in the second population.

[0042] In some cases , the secreted protein is Granulocyte-Colony Stimulating Factor (G-CSF).

[0043] In some cases , the second population contains myeloblast cells that express the G-CSF receptor.

[0044] In a further aspect, the present disclosure relates to methods of making a pharmaceutical formulation comprising human cells comprising a physiologically active secreted protein, comprising transfecting a first population of human cells with a pharmaceutical nucleic acid made by the methods described herein, wherein the secreted protein is active upon a second population of human cells.

Definitions

[0045] At various places in the present specification, substituents of compounds of the present disclosure are disclosed in groups or in ranges. It is specifically intended that the present disclosure include each and every individual subcombination of the members of such groups and ranges. For example, the term "C₁₋₆ alkyl" is specifically intended to individually disclose methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl.

[0046] It is further intended that the compounds of the present disclosure are stable. As used herein "stable" refers to a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and preferably capable of formulation into an efficacious therapeutic agent.

[0047] It is further appreciated that certain features of the present disclosure, which are, for clarity, described separately , can also be provided in combination.

[0048] Conversely, various features of the present disclosure which are, for brevity, described in the context of a single aspect , can also be provided separately or in any suitable subcombination.

[0049] As used herein, the term "alkyl" is meant to refer to a saturated hydrocarbon group which is straight-chained or branched. Example alkyl groups include methyl (Me), ethyl (Et), propyl (e.g., n-propyl and isopropyl), butyl (e.g., n-butyl, isobutyl, t-butyl), pentyl (e.g., n-pentyl, isopentyl, neopentyl), and the like. An alkyl group can contain from 1 to about 20, from 2 to about 20, from 1 to about 12, from 1 to about 8, from 1 to about 6, from 1 to about 4, or from 1

to about 3 carbon atoms.

[0050] As used herein, "alkenyl" refers to an alkyl group having one or more double carbon-carbon bonds. Example alkenyl groups include ethenyl, propenyl, and the like.

[0051] As used herein, "alkoxy" refers to an -O-alkyl group. Example alkoxy groups include methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), t-butoxy, and the like.

[0052] As used herein, "alkynyl" refers to an alkyl group having one or more triple carbon-carbon bonds. Example alkynyl groups include ethynyl, propynyl, and the like.

[0053] As used herein, "aryl" refers to monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbons such as, for example, phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some cases, aryl groups have from 6 to about 20 carbon atoms.

[0054] As used herein, "halo" or "halogen" includes fluoro, chloro, bromo, and iodo.

[0055] As used herein, "therapeutic agent" refers to any agent that, when administered to a subject, has a therapeutic, diagnostic, and/or prophylactic effect and/or elicits a desired biological and/or pharmacological effect.

[0056] As used herein, "animal" refers to any member of the animal kingdom. In some cases, "animal" refers to humans at any stage of development. In some cases, "animal" refers to non-human animals at any stage of development. In certain cases, the non-human animal is a mammal (e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). In some cases, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, and worms. In some cases, the animal is a transgenic animal, genetically-engineered animal, or a clone.

[0057] As used herein, "approximately" or "about," as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain cases, the term "approximately" or "about" refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0058] As used herein, "associated with," "conjugated," "linked," "attached," and "tethered," when used with respect to two or more moieties, means that the moieties are physically associated or connected with one another, either directly or via one or more additional moieties that serves as a linking agent, to form a structure that is sufficiently stable so that the moieties remain physically associated under the conditions in which the structure is used, e.g., physiological conditions.

[0059] As used herein, "biologically active" refers to a characteristic of any substance that has activity in a biological system and/or organism. For instance, a substance that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active. In particular cases, where a nucleic acid is biologically active, a portion of that nucleic acid that shares at least one biological activity of the whole nucleic acid is typically referred to as a "biologically active" portion.

[0060] As used herein, "conserved" refers to nucleotides or amino acid residues of a polynucleotide sequence or amino acid sequence, respectively, that are those that occur unaltered in the same position of two or more related sequences being compared. Nucleotides or amino acids that are relatively conserved are those that are conserved amongst more related sequences than nucleotides or amino acids appearing elsewhere in the sequences. In some cases, two or more sequences are said to be "completely conserved" if they are 100% identical to one another. In some cases, two or more sequences are said to be "highly conserved" if they are at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some cases, two or more sequences are said to be "highly conserved" if they are about 70% identical, about 80% identical, about 90% identical, about 95%, about 98%, or about 99% identical to one another. In some cases, two or more sequences are said to be "conserved" if they are at least 30% identical, at least 40% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some cases, two or more sequences are said to be "conserved" if they are about 30% identical, about 40% identical, about 50% identical, about 60% identical, about 70% identical, about 80% identical, about 90% identical, about 95% identical, about 98% identical, or about 99% identical to one another.

[0061] As used herein, "expression" of a nucleic acid sequence refers to one or more of the following events: (1) production of an RNA template from a DNA sequence (e.g., by transcription); (2) processing of an RNA transcript (e.g., by splicing, editing, 5' cap formation, and/or 3' end processing); (3) translation of an RNA into a polypeptide or protein; and (4) post-translational modification of a polypeptide or protein.

[0062] As used herein, a "functional" biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

[0063] As used herein, "*in vitro*" refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, *etc.*, rather than within an organism (e.g., animal, plant, or microbe).

[0064] As used herein, "*in vivo*" refers to events that occur within an organism (e.g., animal, plant, or microbe).

[0065] As used herein, "isolated" refers to a substance or entity that has been (1) separated from at least some of the components with which it was associated when initially produced

(whether in nature or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated agents are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is "pure" if it is substantially free of other components.

[0066] As used herein, "subject" or "patient" refers to any organism to which a composition in accordance with the present disclosure may be administered, e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals (e.g., mammals such as mice, rats, rabbits, non-human primates, and humans) and/or plants.

[0067] As used herein, "substantially" refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term "substantially" is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

[0068] An individual who is "suffering from" a disease, disorder, and/or condition has been diagnosed with or displays one or more symptoms of a disease, disorder, and/or condition.

[0069] An individual who is "susceptible to" a disease, disorder, and/or condition has not been diagnosed with and/or may not exhibit symptoms of the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition (for example, cancer) may be characterized by one or more of the following: (1) a genetic mutation associated with development of the disease, disorder, and/or condition; (2) a genetic polymorphism associated with development of the disease, disorder, and/or condition; (3) increased and/or decreased expression and/or activity of a protein and/or nucleic acid associated with the disease, disorder, and/or condition; (4) habits and/or lifestyles associated with development of the disease, disorder, and/or condition; (5) a family history of the disease, disorder, and/or condition; and (6) exposure to and/or infection with a microbe associated with development of the disease, disorder, and/or condition. In some cases, an individual who is susceptible to a disease, disorder, and/or condition will develop the disease, disorder, and/or condition. In some cases, an individual who is susceptible to a disease, disorder, and/or condition will not develop the disease, disorder, and/or condition.

[0070] As used herein, "therapeutically effective amount" means an amount of an agent to be delivered (e.g., nucleic acid, drug, therapeutic agent, diagnostic agent, prophylactic agent, etc.) that is sufficient, when administered to a subject suffering from or susceptible to a disease, disorder, and/or condition, to treat, improve symptoms of, diagnose, prevent, and/or delay the onset of the disease, disorder, and/or condition.

[0071] As used herein, "transcription factor" refers to a DNA-binding protein that regulates transcription of DNA into RNA, for example, by activation or repression of transcription. Some transcription factors effect regulation of transcription alone, while others act in concert with other proteins. Some transcription factor can both activate and repress transcription under certain conditions. In general, transcription factors bind a specific target sequence or sequences highly similar to a specific consensus sequence in a regulatory region of a target gene. Transcription factors may regulate transcription of a target gene alone or in a complex with other molecules.

[0072] As used herein, "treating" refers to partially or completely alleviating, ameliorating, improving, relieving, delaying onset of, inhibiting progression of, reducing severity of, and/or reducing incidence of one or more symptoms or features of a particular disease, disorder, and/or condition. For example, "treating" cancer may refer to inhibiting survival, growth, and/or spread of a tumor. Treatment may be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a disease, disorder, and/or condition for the purpose of decreasing the risk of developing pathology associated with the disease, disorder, and/or condition. In some cases, treatment comprises delivery of a protein associated with a therapeutically active nucleic acid to a subject in need thereof.

[0073] As used herein, "unmodified" refers to a nucleic acid prior to being modified, e.g. adenosine, guanosine, cytosine, thymidine, and uracil, or a naturally occurring amino acid. The compounds described herein can be asymmetric (e.g., having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically active starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present disclosure. Cis and trans geometric isomers of the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms.

[0074] Compounds of the present disclosure also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone - enol pairs, amide - imidic acid pairs, lactam - lactim pairs, amide - imidic acid pairs, enamine - imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

[0075] Compounds of the present disclosure can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

[0076] The term "compound," as used herein, is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted.

[0077] The compounds of the present disclosure may be substantially isolated. By "substantially isolated" is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compound of the present disclosure. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound of the present disclosure, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

[0078] The compounds of the present disclosure, and salts thereof, can also be prepared in combination with solvent or water molecules to form solvates and hydrates by routine methods.

[0079] The present disclosure also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and Journal of Pharmaceutical Science, 66, 2 (1977).

[0080] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.


[0081] The present disclosure also includes prodrugs of the compounds described herein. As

used herein, "prodrugs" refer to any carriers, typically covalently bonded, which release the active parent drug when administered to a mammalian subject. Prodrugs can be prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds. Prodrugs include compounds wherein hydroxyl, amino, sulfhydryl, or carboxyl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, sulfhydryl, or carboxyl group respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of the present disclosure. Preparation and use of prodrugs is discussed in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

Modified Nucleosides and Nucleotides

[0082] The present disclosure relates to modified nucleosides and nucleotides. As described herein "nucleoside" is defined as a compound containing a five-carbon sugar molecule (a pentose or ribose) or derivative thereof, and an organic base, purine or pyrimidine, or a derivative thereof. As described herein, "nucleotide" is defined as a nucleoside consisting of a phosphate group. The nucleosides and nucleotides described herein are generally chemically modified on the major groove face. In some cases, the major groove chemical modifications can include an amino group, a thiol group, an alkyl group, or a halo group.

[0083] Table 1 below identifies the chemical faces of each canonical nucleotide. Circles identify the atoms comprising the respective chemical regions.



-

OH OH

OH OH

OH OH

[0084] Modified nucleosides include pyridin-4-one ribonucleoside, 5-aza-uridine, 2-thio-5-aza-uridine, 2-thiouridine, 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxyuridine, 3-methyluridine, 5-carboxymethyl-uridine, 1-carboxymethyl-pseudouridine, 5-propynyl-uridine, 1-propynyl-pseudouridine, 5-taurinomethyluridine, 1-taurinomethyl-pseudouridine, 5-taurinomethyl-2-thio-uridine, 1-taurinomethyl-4-thio-uridine, 5-methyl-uridine, 1-methyl-pseudouridine, 4-thio-1-methyl-pseudouridine, 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouridine, dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, and 4-methoxy-2-thio-pseudouridine.


[0085] Modified nucleosides include 5-aza-cytidine, pseudoisocytidine, 3-methyl-cytidine, N4-acetylcytidine, 5-formylcytidine, N4-methylcytidine, 5-hydroxymethylcytidine, 1-methyl-pseudoisocytidine, pyrrolo-cytidine, pyrrolo-pseudoisocytidine, 2-thio-cytidine, 2-thio-5-methyl-cytidine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytidine, 2-methoxy-5-methyl-cytidine, 4-methoxy-pseudoisocytidine, and 4-methoxy-1-methyl-pseudoisocytidine.


[0086] Modified nucleosides include 2-aminopurine, 2, 6-diaminopurine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-aminopurine, 7-deaza-8-aza-2-aminopurine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyladenosine, N6-methyladenosine, N6-isopentenyladenosine, N6-(cis-hydroxyisopentenyl)adenosine, 2-methylthio-N6-(cis-hydroxyisopentenyl) adenosine, N6-glycinylicarbamoyl-adenosine, N6-threonylicarbamoyl-adenosine, 2-methylthio-N6-threonyl carbamoyl-adenosine, N6,N6-dimethyladenosine, 7-methyladenine, 2-methylthio-adenine, and 2-methoxy-adenine.

[0087] Modified nucleosides include inosine, 1-methyl-inosine, wyosine, wybutosine, 7-deaza-guanosine, 7-deaza-8-aza-guanosine, 6-thio-guanosine, 6-thio-7-deaza-guanosine, 6-thio-7-deaza-8-aza-guanosine, 7-methyl-guanosine, 6-thio-7-methyl-guanosine, 7-methylinosine, 6-methoxy-guanosine, 1-methyl-guanosine, N2-methyl-guanosine, N2,N2-dimethyl-guanosine, 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 1-methyl-6-thio-guanosine, N2-methyl-6-thio-guanosine, and N2,N2-dimethyl-6-thio-guanosine.

[0088] The nucleotide can be modified on the major groove face and can include replacement of the hydrogen on C-5 of uracil with a methyl group or a halo group.

[0089] The nucleoside and nucleotide can be a compound of Formula I:





I

wherein:


Z is O or S;

each of Y¹ is independently selected from -OR^{a1}, -NR^{a1}R^{b1}, and -SR^{a1};each of Y² is independently selected from O, NR^a, S or a linker comprising an atom selected from the group consisting of C, O, N, and S;each of Y³ is independently selected from O and S;Y⁴ is selected from H, -OR^a, -SR^a, and -NHR^a;


n is 0, 1, 2, or 3;

m is 0, 1, 2 or 3;


B is a nucleobase;

R^a is H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl;R^{a1} and R^{b1} are each independently H or a counterion; and-Y³-R^{c1} is OH or SH at a pH of about 1 or -Y³-R^{c1} is O⁻ or S⁻ at physiological pH;or -Y³-R^{c1} is C₁₋₂₀ alkoxy, C₂₋₂₀ -O-alkenyl, or C₁₋₂₀ -O-alkynyl;wherein when B is an unmodified nucleobase selected from cytosine, guanine, uracil and adenine, then at least one of Z, Y¹ or Y² is not O or OH.**[0090]** In some cases, B is a nucleobase of Formula II-a, II-b, or II-c:

II-a



II-b



II-c

wherein:



denotes a single or double bond;

X is O or S;

U and W are each independently C or N;

V is O, S, C or N;

wherein when V is C then R^1 is H, C_{1-6} alkyl, C_{1-6} alkenyl, C_{1-6} alkynyl, halo, or $-OR^c$, wherein C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl are each optionally substituted with $-OH$, $-NR^aR^b$, $-SH$, $-C(O)R^c$, $-C(O)OR^c$, $-NHC(O)R^c$, or $-NHC(O)OR^c$;

and wherein when V is O, S, or N then R^1 is absent;

R^2 is H, $-OR^c$, $-SR^c$, $-NR^aR^b$, or halo;

or when V is C then R^1 and R^2 together with the carbon atoms to which they are attached can form a 5- or 6-membered ring optionally substituted with 1-4 substituents selected from halo, $-OH$, $-SH$, $-NR^aR^b$, C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, C_{1-20} alkoxy, or C_{1-20} thioalkyl;

R^3 is H or C_{1-20} alkyl;

R^4 is H or C_{1-20} alkyl; wherein when




denotes a double bond then R^4 is absent, or $N-R^4$, taken together, forms a positively charged N substituted with C_{1-20} alkyl;

R^a and R^b are each independently H, C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, or C_{6-20} aryl; and

R^c is H, C_{1-20} alkyl, C_{2-20} alkenyl, phenyl, benzyl, a polyethylene glycol group, or an amino-polyethylene glycol group.


[0091] In some cases, B is a nucleobase of Formula II-a1, II-a2, II-a3, II-a4, or II-a5:



[0092] In some cases, B is a nucleobase selected from the group consisting of cytosine, guanine, adenine, and uracil.


[0093] In some cases, B is a pyrimidine or derivative thereof.

[0094] In some cases the nucleotide is a compound of Formula I-a:




I-a.

[0095] In some cases the nucleotide is a compound of Formula I-b;




I-b.


[0096] Formula I-c;




I-c.

or is selected from the group consisting of:






[0097] In some cases , the nucleotide is selected from the group consisting of:




[0098] For example, the modified nucleotide can be:




[0099] In some cases , the major groove chemical modification can include replacement of the C-H group at C-5 with an -NH- group or a -NH(CH₃)- group.

[0100] For example, the modified nucleotide can be:




[0101] In another case, the major groove chemical modification can include replacement of the hydrogen at C-5 of cytosine with a halo group or a methyl group.

[0102] For example, the modified nucleotide can be:




[0103] In yet a further case, the major groove chemical modification can include a fused ring that is formed by the NH₂ at the C-4 position and the carbon atom at the C-5 position.


[0104] For example, the modified nucleotide can be:



[0105] In some cases, a modified nucleotide is 5'-O-(1-Thiophosphate)-Adenosine, 5'-O-(1-Thiophosphate)-Cytidine, 5'-O-(1-Thiophosphate)-Guanosine, 5'-O-(1-Thiophosphate)-Uridine or 5'-O-(1-Thiophosphate)-Pseudouridine.




5'-O-(1-Thiophosphate)-Adenosine







5'-O-(1-Thiophosphate)-Cytidine



5'-O-(1-Thiophosphate)-Guanosine



5'-O-(1-Thiophosphate)-Uridine



5'-O-(1-Thiophosphate)-Pseudouridine

[0106] The α -thio substituted phosphate moiety is provided to confer stability to RNA and DNA polymers through the unnatural phosphorothioate backbone linkages. Phosphorothioate DNA and RNA have increased nuclease resistance and subsequently a longer half-life in a cellular environment. Phosphorothioate linked nucleic acids are expected to also reduce the innate immune response through weaker binding/activation of cellular innate immune molecules.

[0107] Further examples of modified nucleotides and modified nucleotide combinations are provided below in Table 2.

Table 2

Modified Nucleotide	Modified Nucleotide Combination
6-aza-cytidine	α -thio-cytidine/5-iodo-uridine
2-thio-cytidine	α -thio-cytidine/N 1-methyl-pseudo-uridine
α -thio-cytidine	α -thio-cytidine/ α -thio-uridine
Pseudo-iso-cytidine	α -thio-cytidine/5-methyl-uridine
5-aminoallyl-uridine	α -thio-cytidine/pseudo-uridine
5-iodo-uridine	Pseudo-iso-cytidine/5-iodo-uridine

Modified Nucleotide	Modified Nucleotide Combination
N1-methyl-pseudouridine	Pseudo-iso-cytidine/N 1-methyl-pseudo-uridine
5,6-dihydro-uridine	Pseudo-iso-cytidine/ α -thio-uridine
α -thio-uridine	Pseudo-iso-cytidine/5-methyl-uridine
4-thio-uridine	Pseudo-iso-cytidine/Pseudo-uridine
6-aza-uridine	Pyrrolo-cytidine
5-hydroxy-uridine	Pyrrolo-cytidine/5-iodo-uridine
Deoxy-thymidine	Pyrrolo-cytidine/N 1-methyl-pseudo-uridine
Pseudo-uridine	Pyrrolo-cytidine/ α -thio-uridine
Inosine	Pyrrolo-cytidine/5-methyl-uridine
α -thio-guanosine	Pyrrolo-cytidine/Pseudo-uridine
8-oxo-guanosine	5-methyl-cytidine/5-iodo-uridine
O6-methyl-guanosine	5-methyl-cytidine/N 1-methyl-pseudo-uridine
7-deaza-guanosine	5-methyl-cytidine/ α -thio-uridine
No modification	5-methyl-cytidine/5-methyl-uridine
N 1-methyl-adenosine	5-methyl-cytidine/Pseudo-uridine
2-amino-6-Chloro-purine	5-methyl-cytidine
N6-methyl-2-aminopurine	25% Pseudo-iso-cytidine
6-Chloro-purine	25% N1-methyl-pseudo-uridine
N6-methyl-adenosine	25% N 1-Methyl-pseudo-uridine/75%-pseudo-uridine
α -thio-adenosine	5-methyl-uridine
8-azido-adenosine	5-iodo-cytidine
7-deaza-adenosine	

Synthesis of Modified Nucleotides

[0108] The modified nucleosides and nucleotides disclosed herein can be prepared from readily available starting materials using the following general methods and procedures. It is understood that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given; other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

[0109] The processes described herein can be monitored according to any suitable method


known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ^1H or ^{13}C) infrared spectroscopy, spectrophotometry (e.g., UV-visible), or mass spectrometry, or by chromatography such as high performance liquid chromatography (HPLC) or thin layer chromatography.


[0110] Preparation of modified nucleosides and nucleotides can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Greene, et al., *Protective Groups in Organic Synthesis*, 2d. Ed., Wiley & Sons, 1991.

[0111] The reactions of the processes described herein can be carried out in suitable solvents, which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially nonreactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, i.e., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected. Resolution of racemic mixtures of modified nucleosides and nucleotides can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a "chiral resolving acid" which is an optically active, salt-forming organic acid. Suitable resolving agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids. Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (e.g., dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.


[0112] Exemplary syntheses of modified nucleotides are provided below in Schemes 1 and 2.

Scheme 1





Scheme 2



[0113] Modified nucleosides and nucleotides can also be prepared according to the synthetic methods described in Ogata et al. *Journal of Organic Chemistry* 74:2585-2588, 2009; Purmal et al. *Nucleic Acids Research* 22(1): 72-78, 1994; Fukuhara et al. *Biochemistry* 1(4): 563-568, 1962; and Xu et al. *Tetrahedron* 48(9): 1729-1740, 1992.

Modified Nucleic Acids

[0114] The present disclosure relates to nucleic acids, including RNAs such as mRNAs that contain one or more modified nucleosides (termed "modified nucleic acids") or nucleotides as described herein, which have useful properties including the significant decrease or lack of a substantial induction of the innate immune response of a cell into which the mRNA is introduced, or the suppression thereof. Because these modified nucleic acids enhance the efficiency of protein production, intracellular retention of nucleic acids, and viability of contacted cells, as well as possess reduced immunogenicity, of these nucleic acids compared to

unmodified nucleic acids, having these properties are termed "enhanced nucleic acids" herein.

[0115] In addition, the present disclosure provides nucleic acids, which have decreased binding affinity to a major groove interacting, e.g. binding, partner. For example, the nucleic acids are comprised of at least one nucleotide that has been chemically modified on the major groove face as described herein.

[0116] The term "nucleic acid," in its broadest sense, includes any compound and/or substance that is or can be incorporated into an oligonucleotide chain. Exemplary nucleic acids for use in accordance with the present disclosure include, but are not limited to, one or more of DNA, RNA including messenger mRNA (mRNA), hybrids thereof, RNAi-inducing agents, RNAi agents, siRNAs, shRNAs, miRNAs, antisense RNAs, ribozymes, catalytic DNA, RNAs that induce triple helix formation, aptamers, vectors, *etc.*, described in detail herein.

[0117] Described are modified nucleic acids containing a translatable region and one, two, or more than two different nucleoside modifications. In some cases, the modified nucleic acid exhibits reduced degradation in a cell into which the nucleic acid is introduced, relative to a corresponding unmodified nucleic acid. Exemplary nucleic acids include ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), locked nucleic acids (LNAs) or a hybrid thereof. In the invention, the modified nucleic acid is messenger RNA (mRNA). As described herein, the nucleic acids of the present disclosure do not substantially induce an innate immune response of a cell into which the mRNA is introduced.

[0118] In certain cases, it is desirable to intracellularly degrade a modified nucleic acid introduced into the cell, for example if precise timing of protein production is desired. Thus, the present disclosure provides a modified nucleic acid containing a degradation domain, which is capable of being acted on in a directed manner within a cell.


[0119] Other components of nucleic acid are optional, and are beneficial in some cases. For example, a 5' untranslated region (UTR) and/or a 3'UTR are described, wherein either or both may independently contain one or more different nucleoside modifications. In such cases, nucleoside modifications may also be present in the translatable region. Also described are nucleic acids containing a Kozak sequence.

[0120] Additionally, described are nucleic acids containing one or more intronic nucleotide sequences capable of being excised from the nucleic acid.

[0121] Further, described are nucleic acids containing an internal ribosome entry site (IRES). An IRES may act as the sole ribosome binding site, or may serve as one of multiple ribosome binding sites of an mRNA. An mRNA containing more than one functional ribosome binding site may encode several peptides or polypeptides that are translated independently by the ribosomes ("multicistronic mRNA"). When nucleic acids are provided with an IRES, further optionally provided can be a second translatable region. Examples of IRES sequences that can

be used according to the present disclosure include without limitation, those from picornaviruses (e.g. FMDV), pest viruses (CFFV), polio viruses (PV), encephalomyocarditis viruses (ECMV), foot-and-mouth disease viruses (FMDV), hepatitis C viruses (HCV), classical swine fever viruses (CSFV), murine leukemia virus (MLV), simian immune deficiency viruses (SIV) or cricket paralysis viruses (CrPV).

[0122] In some cases, the nucleic acid sequences comprise a compound of Formula I-d:



I-d

wherein:

Z is O or S;

each of Y¹ is independently selected from -OR^{a1}, -NR^{a1}R^{b1}, and -SR^{a1};

each of Y² is independently selected from O, NR^a, S or a linker comprising an atom selected from the group consisting of C, O, N, and S;

each of Y³ is independently selected from O and S;

Y⁴ is selected from H, -OR^a, -SR^a, and -NHR^a;

B is a nucleobase;

R^a is H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl;


R^{a1} and R^{b1} are each independently H or a counterion; and


-Y³-R^{c1} is OH or SH at a pH of about 1 or -Y³-R^{c1} is O⁻ or S⁻ at physiological pH;

or -Y³-R^{c1} is C₁₋₂₀ alkoxy, C₂₋₂₀ -O-alkenyl, or C₁₋₂₀ -O-alkynyl;


wherein when B is an unmodified nucleobase selected from cytosine, guanine, thymidine, uracil and adenine, then at least one of Z, Y¹ or Y² is not O or OH.

[0123] In some cases, B is a nucleobase of Formula II-a, II-b, or II-c:






II-a



II-b



II-c

wherein:



denotes a single or double bond;

X is O or S;

U and W are each independently C or N;

V is O, S, C or N;

wherein when V is C then R¹ is H, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, halo, or -OR^c, wherein C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl are each optionally substituted with -OH, -NR^aR^b, -SH, -C(O)R^c, -C(O)OR^c, -NHC(O)R^c, or -NHC(O)OR^c;

and wherein when V is O, S, or N then R¹ is absent;

R² is H, -OR^c, -SR^c, -NR^aR^b, or halo;

or when V is C then R¹ and R² together with the carbon atoms to which they are attached can form a 5- or 6-membered ring optionally substituted with 1-4 substituents selected from halo, -OH, -SH, -NR^aR^b, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, C₁₋₂₀ alkoxy, or C₁₋₂₀ thioalkyl;

R³ is H or C₁₋₂₀ alkyl;

R⁴ is H or C₁₋₂₀ alkyl; wherein when









denotes a double bond then R⁴ is absent, or N-R⁴, taken together, forms a positively charged N substituted with C₁₋₂₀ alkyl;

R^a and R^b are each independently H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl; and

R^c is H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, phenyl, benzyl, a polyethylene glycol group, or an amino-polyethylene glycol group.

[0124] In some cases, B is a nucleobase of Formula II-a1, II-a2, II-a3, II-a4, or II-a5:





[0125] In some cases , at least 25% of the cytosines are replaced by a compound of Formula I-a (e.g., at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

[0126] In some cases , at least 25% of the uracils are replaced by a compound of Formula I-a (e.g., at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

[0127] In some cases , at least 25% of the cytosines and 25% of the uracils are replaced by a compound of Formula I-a (e.g., at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

Major Groove Interacting Partners

[0128] As described herein, the phrase "major groove interacting partner" refers RNA recognition receptors that detect and respond to RNA ligands through interactions, e.g. binding, with the major groove face of a nucleotide or nucleic acid. As such, RNA ligands comprising modified nucleotides or nucleic acids as described herein decrease interactions with major groove binding partners, and therefore decrease an innate immune response, or expression and secretion of pro-inflammatory cytokines, or both.

[0129] Example major groove interacting, e.g. binding, partners include, but are not limited to the following nucleases and helicases. Within membranes, TLRs (Toll-like Receptors) 3, 7, and 8 can respond to single- and double-stranded RNAs. Within the cytoplasm, members of the superfamily 2 class of DEX(D/H) helicases and ATPases can sense RNAs to initiate antiviral responses. These helicases include the RIG-I (retinoic acid-inducible gene I) and MDA5 (melanoma differentiation-associated gene 5). Other examples include laboratory of genetics and physiology 2 (LGP2), HIN-200 domain containing proteins, or Helicase-domain containing proteins.

Prevention or reduction of innate cellular immune response activation using modified nucleic acids

[0130] The term "innate immune response" includes a cellular response to exogenous nucleic acids, including single stranded nucleic acids, generally of viral or bacterial origin, which involves the induction of cytokine expression and release, particularly the interferons, and cell death. Protein synthesis is also reduced during the innate cellular immune response. While it is advantageous to eliminate the innate immune response in a cell, the present disclosure provides modified mRNAs that substantially reduce the immune response, including interferon signaling, without entirely eliminating such a response. In some embodiments, the immune response is reduced by 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99%, 99.9%, or greater than 99.9% as compared to the immune response induced by a corresponding unmodified nucleic acid. Such a reduction can be measured by expression or activity level of Type 1 interferons or the expression of interferon-regulated genes such as the toll-like receptors (e.g., TLR7 and TLR8). Reduction of innate immune response can also be measured by decreased cell death following one or more administrations of modified RNAs to a cell population; e.g., cell death is 10%, 25%, 50%, 75%, 85%, 90%, 95%, or over 95% less than the cell death frequency observed with a corresponding unmodified nucleic acid. Moreover, cell death may affect fewer than 50%, 40%, 30%, 20%, 10%, 5%, 1%, 0.1%, 0.01% or fewer than 0.01% of cells contacted with the modified nucleic acids.

[0131] Described herein is the repeated introduction (e.g., transfection) of modified nucleic acids into a target cell population, e.g., *in vitro*, *ex vivo*, or *in vivo*. The step of contacting the cell population may be repeated one or more times (such as two, three, four, five or more than five times). The step of contacting the cell population with the modified nucleic acids may be repeated a number of times sufficient such that a predetermined efficiency of protein translation in the cell population is achieved. Given the reduced cytotoxicity of the target cell population provided by the nucleic acid modifications, such repeated transfections are achievable in a diverse array of cell types.

Polypeptide variants

[0132] Described are nucleic acids that encode variant polypeptides, which have a certain identity with a reference polypeptide sequence. The term "identity" as known in the art, refers to a relationship between the sequences of two or more peptides, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between peptides, as determined by the number of matches between strings of two or more amino acid residues. "Identity" measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., "algorithms"). Identity of related peptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds.,

Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo et al., SIAM J. Applied Math. 48, 1073 (1988).

[0133] In some cases, the polypeptide variant has the same or a similar activity as the reference polypeptide. Alternatively, the variant has an altered activity (e.g., increased or decreased) relative to a reference polypeptide. Generally, variants of a particular polynucleotide or polypeptide of the present disclosure will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to that particular reference polynucleotide or polypeptide as determined by sequence alignment programs and parameters described herein and known to those skilled in the art.

[0134] Protein fragments, functional protein domains, and homologous proteins are also described, for example, any protein fragment of a reference protein (meaning a polypeptide sequence at least one amino acid residue shorter than a reference polypeptide sequence but otherwise identical) 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or greater than 100 amino acids in length. In another example, any protein that includes a stretch of about 20, about 30, about 40, about 50, or about 100 amino acids which are about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, or about 100% identical to any of the sequences described herein can be utilized in accordance with the present disclosure. In certain cases, a protein sequence to be utilized in accordance with the present disclosure includes 2, 3, 4, 5, 6, 7, 8, 9, 10, or more mutations as shown in any of the sequences provided or referenced herein.

Polypeptide libraries

[0135] Also described are polynucleotide libraries containing nucleoside modifications, wherein the polynucleotides individually contain a first nucleic acid sequence encoding a polypeptide, such as an antibody, protein binding partner, scaffold protein, and other polypeptides known in the art. Preferably, the polynucleotides are mRNA of the invention, in a form suitable for direct introduction into a target cell host, which in turn synthesizes the encoded polypeptide.

[0136] Multiple variants of a protein, each with different amino acid modification(s), may be produced and tested to determine the best variant in terms of pharmacokinetics, stability, biocompatibility, and/or biological activity, or a biophysical property such as expression level. Such a library may contain 10 , 10^2 , 10^3 , 10^4 , 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , or over 10^9 possible variants (including substitutions, deletions of one or more residues, and insertion of one or more residues).

Polypeptide-nucleic acid complexes

[0137] Proper protein translation involves the physical aggregation of a number of polypeptides and nucleic acids associated with the mRNA. Described herein are protein-nucleic acid complexes, containing a translatable mRNA having one or more nucleoside modifications (e.g., at least two different nucleoside modifications) and one or more polypeptides bound to the mRNA. Generally, the proteins are provided in an amount effective to prevent or reduce an innate immune response of a cell into which the complex is introduced.

Untranslatable Modified Nucleic Acids

[0138] Also described herein are mRNAs having sequences that are substantially not translatable. Such mRNA is effective as a vaccine when administered to a mammalian subject.

[0139] Also described are modified nucleic acids that contain one or more noncoding regions. Such modified nucleic acids are generally not translated, but are capable of binding to and sequestering one or more translational machinery component such as a ribosomal protein or a transfer RNA (tRNA), thereby effectively reducing protein expression in the cell. The modified nucleic acid may contain a small nucleolar RNA (sno-RNA), micro RNA (miRNA), small interfering RNA (siRNA) or Piwi-interacting RNA (piRNA).

Synthesis of Modified Nucleic Acids

[0140] Nucleic acids for use in accordance with the present disclosure may be prepared according to any available technique including, but not limited to chemical synthesis, enzymatic synthesis, which is generally termed *in vitro* transcription, enzymatic or chemical cleavage of a longer precursor, etc. Methods of synthesizing RNAs are known in the art (see, e.g., Gait, M.J. (cd.) Oligonucleotide synthesis: a practical approach, Oxford [Oxfordshire], Washington, DC: IRL Press, 1984; and Herdewijn, P. (ed.) Oligonucleotide synthesis: methods and applications, Methods in Molecular Biology, v. 288 (Clifton, N.J.) Totowa, N.J.: Humana Press, 2005).

[0141] The modified nucleosides and nucleotides disclosed herein can be prepared from readily available starting materials using the following general methods and procedures. It is understood that where typical or preferred process conditions (*i.e.*, reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given; other process conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.

[0142] The processes described herein can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ^1H or ^{13}C) infrared spectroscopy, spectrophotometry (e.g., UV-visible), or mass spectrometry, or by chromatography such as

high performance liquid chromatography (HPLC) or thin layer chromatography.

[0143] Preparation of modified nucleosides and nucleotides can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in Greene, et al., *Protective Groups in Organic Synthesis*, 2d. Ed., Wiley & Sons, 1991.

[0144] The reactions of the processes described herein can be carried out in suitable solvents, which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially nonreactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, i.e., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected.

[0145] Resolution of racemic mixtures of modified nucleosides and nucleotides can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a "chiral resolving acid" which is an optically active, salt-forming organic acid. Suitable resolving agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids. Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (e.g., dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art. Modified nucleic acids need not be uniformly modified along the entire length of the molecule. Different nucleotide modifications and/or backbone structures may exist at various positions in the nucleic acid. One of ordinary skill in the art will appreciate that the nucleotide analogs or other modification(s) may be located at any position(s) of a nucleic acid such that the function of the nucleic acid is not substantially decreased. A modification may also be a 5' or 3' terminal modification. Nucleic acids described herein may contain at a minimum one and at maximum 100% modified nucleotides, or any intervening percentage, such as at least 5% modified nucleotides, at least 10% modified nucleotides, at least 25% modified nucleotides, at least 50% modified nucleotides, at least 80% modified nucleotides, or at least 90% modified nucleotides. For example, nucleic acids may contain a modified pyrimidine such as uracil or cytosine. In the invention, 100% of the uracil in the nucleic acid is replaced with a modified uracil, namely by using N1-methyl-pseudouridine.

[0146] In some embodiments, at least 5%, at least 10%, at least 25%, at least 50%, at least 80%, at least 90% or 100% of the cytosine in the nucleic acid is replaced with a modified cytosine. The modified cytosine can be replaced by a compound having a single unique structure, or can be replaced by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures).


[0147] Generally, the shortest length of a modified mRNA of the present disclosure can be the length of an mRNA sequence that is sufficient to encode for a dipeptide. In another embodiment, the length of the mRNA sequence is sufficient to encode for a tripeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a tetrapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a pentapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a hexapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a heptapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for an octapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a nonapeptide. In another embodiment, the length of an mRNA sequence is sufficient to encode for a decapeptide.

[0148] Examples of dipeptides that the modified nucleic acid sequences can encode for include, but are not limited to, carnosine and anserine.

[0149] In a further embodiment, the mRNA is greater than 30 nucleotides in length. In another embodiment, the RNA molecule is greater than 35 nucleotides in length. In another embodiment, the length is at least 40 nucleotides. In another embodiment, the length is at least 45 nucleotides. In another embodiment, the length is at least 55 nucleotides. In another embodiment, the length is at least 60 nucleotides. In another embodiment, the length is at least 60 nucleotides. In another embodiment, the length is at least 80 nucleotides. In another embodiment, the length is at least 90 nucleotides. In another embodiment, the length is at least 100 nucleotides. In another embodiment, the length is at least 120 nucleotides. In another embodiment, the length is at least 140 nucleotides. In another embodiment, the length is at least 160 nucleotides. In another embodiment, the length is at least 180 nucleotides. In another embodiment, the length is at least 200 nucleotides. In another embodiment, the length is at least 250 nucleotides. In another embodiment, the length is at least 300 nucleotides. In another embodiment, the length is at least 350 nucleotides. In another embodiment, the length is at least 400 nucleotides. In another embodiment, the length is at least 450 nucleotides. In another embodiment, the length is at least 500 nucleotides. In another embodiment, the length is at least 600 nucleotides. In another embodiment, the length is at least 700 nucleotides. In another embodiment, the length is at least 800 nucleotides. In another embodiment, the length is at least 900 nucleotides. In another embodiment, the length is at least 1000 nucleotides. In another embodiment, the length is at least 1100 nucleotides. In another embodiment, the length is at least 1200 nucleotides. In another embodiment, the length is at least 1300 nucleotides. In another embodiment, the length is at least 1400 nucleotides. In another embodiment, the length is at least 1500 nucleotides. In another embodiment, the length is at least 1600 nucleotides. In another embodiment, the length is at least 1800 nucleotides. In another embodiment, the length is at least 2000 nucleotides. In another embodiment, the length is at least 2500 nucleotides. In another embodiment, the length is at least 3000 nucleotides. In another embodiment, the length is at least 4000 nucleotides. In another embodiment, the length is at least 5000 nucleotides, or greater than 5000 nucleotides.

[0150] The present disclosure relates to methods of preparing a nucleic acid sequence

comprising a nucleotide that disrupts binding of a major groove interacting partner with the nucleic acid sequence, wherein the nucleic acid sequence comprises a compound of Formula I-d:



I-d

wherein:

Z is O or S;

each of Y¹ is independently selected from -OR^{a1}, -NR^{a1}R^{b1}, and -SR^{a1};

each of Y² is independently selected from O, NR^a, S or a linker comprising an atom selected from the group consisting of C, O, N, and S;

each of Y³ is independently selected from O and S;

Y⁴ is selected from H, -OR^a, -SR^a, and -NHR^a;

B is a nucleobase;

R^a is H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl;

R^{a1} and R^{b1} are each independently H or a counterion; and


-Y³-R^{c1} is OH or SH at a pH of about 1 or -Y³-R^{c1} is O⁻ or S⁻ at physiological pH;

or -Y³-R^{c1} is C₁₋₂₀ alkoxy, C₂₋₂₀ -O-alkenyl, or C₁₋₂₀ -O-alkynyl;

wherein when B is an unmodified nucleobase selected from cytosine, guanine, uracil and adenine, then at least one of Z, Y¹ or Y² is not O or OH;

the method comprising:

reacting a compound of Formula I-c:




I-c

with an RNA polymerase, and a cDNA template.

[0151] In some embodiments, the reaction is repeated from 1 to about 7,000 times.

[0152] In some embodiments, B is a nucleobase of Formula II-a, II-b, or II-c:



wherein:



denotes a single or double bond;

X is O or S;

U and W are each independently C or N;

V is O, S, C or N;

wherein when V is C then R¹ is H, C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, halo, or -OR^c, wherein C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl are each optionally substituted with -OH, -NR^aR^b, -SH, -C(O)R^c, -C(O)OR^c, -NHC(O)R^c, or -NHC(O)OR^c;

and wherein when V is O, S, or N then R¹ is absent;

R² is H, -OR^c, -SR^c, -NR^aR^b, or halo;

or when V is C then R¹ and R² together with the carbon atoms to which they are attached can form a 5- or 6-membered ring optionally substituted with 1-4 substituents selected from halo, -OH, -SH, -NR^aR^b, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, C₁₋₂₀ alkoxy, or C₁₋₂₀ thioalkyl;

R³ is H or C₁₋₂₀ alkyl;

R⁴ is H or C₁₋₂₀ alkyl; wherein when




denotes a double bond then R⁴ is absent, or N-R⁴, taken together, forms a positively charged N substituted with C₁₋₂₀ alkyl;

R^a and R^b are each independently H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl; and

R^c is H, C_{1-20} alkyl, C_{2-20} alkenyl, phenyl, benzyl, a polyethylene glycol group, or an amino-polyethylene glycol group.

[0153] In some cases, B is a nucleobase of Formula II-a1, II-a2, II-a3, II-a4, or II-a5:




[0154] In some cases, the methods further comprise a nucleotide selected from the group consisting of adenosine, cytosine, guanosine, and uracil.

[0155] In some cases, the nucleobase is a pyrimidine or derivative thereof.

[0156] In a further aspect, the present disclosure includes methods of amplifying a nucleic acid sequence comprising a nucleotide that disrupts binding of a major groove binding partner with the nucleic acid sequence, the method comprising:

reacting a compound of Formula I-c:



Z is O or S;

each of Y^1 is independently selected from $-OR^{a1}$, $-NR^{a1}R^{b1}$, and $-SR^{a1}$;

each of Y^2 is independently selected from O, NR^a , S or a linker comprising an atom selected from the group consisting of C, O, N, and S;

each of Y^3 is independently selected from O and S;

Y^4 is selected from H, $-OR^a$, $-SR^a$, and $-NHR^a$;

B is a nucleobase;

R^a is H, C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, or C_{6-20} aryl;


R^{a1} and R^{b1} are each independently H or a counterion; and

$-Y^3-R^{c1}$ is OH or SH at a pH of about 1 or $-Y^3-R^{c1}$ is O^- or S^- at physiological pH;


or $-Y^3-R^{c1}$ is C_{1-20} alkoxy, C_{2-20} -O-alkenyl, or C_{1-20} -O-alkynyl;

wherein when B is an unmodified nucleobase selected from cytosine, guanine, uracil and adenine, then at least one of Z, Y^1 or Y^2 is not O or OH;
with a primer, a cDNA template, and an RNA polymerase.


[0157] In some cases, B is a nucleobase of Formula II-a, II-b, or II-c:



II-a



II-b



II-c

wherein:



denotes a single or double bond;

X is O or S;

U and W are each independently C or N;

V is O, S, C or N;

wherein when V is C then R^1 is H, C_{1-6} alkyl, C_{1-6} alkenyl, C_{1-6} alkynyl, halo, or $-OR^c$, wherein C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl are each optionally substituted with -OH, $-NR^aR^b$, -SH, $-C(O)R^c$, $-C(O)OR^c$, $-NHC(O)R^c$, or $-NHC(O)OR^c$;

and wherein when V is O, S, or N then R^1 is absent;

R^2 is H, $-OR^c$, $-SR^c$, $-NR^aR^b$, or halo;

or when V is C then R^1 and R^2 together with the carbon atoms to which they are attached can form a 5- or 6-membered ring optionally substituted with 1-4 substituents selected from halo, -OH, -SH, $-NR^aR^b$, C_{1-20} alkyl, C_{2-20} alkenyl, C_{2-20} alkynyl, C_{1-20} alkoxy, or C_{1-20} thioalkyl;

R^3 is H or C_{1-20} alkyl;

R^4 is H or C_{1-20} alkyl; wherein when




denotes a double bond then R^4 is absent, or N- R^4 , taken together, forms a positively charged N substituted with C_{1-20} alkyl;

R^a and R^b are each independently H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, C₂₋₂₀ alkynyl, or C₆₋₂₀ aryl; and

R^c is H, C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, phenyl, benzyl, a polyethylene glycol group, or an amino-polyethylene glycol group.

[0158] In some cases, B is a nucleobase of Formula II-a1, II-a2, II-a3, II-a4, or II-a5:



[0159] In some embodiments, the methods further comprise a nucleotide selected from the group consisting of adenosine, cytosine, guanosine, and uracil.

[0160] In some embodiments, the nucleobase is a pyrimidine or derivative thereof.

Uses of Modified Nucleic Acids

Therapeutic Agents

[0161] The modified mRNA of the invention and the proteins translated therefrom can be used as therapeutic agents. For example, a modified mRNA of the invention can be administered to a subject, wherein the modified nucleic acid is translated in vivo to produce a therapeutic peptide in the subject. Accordingly, the mRNA can be used in compositions, methods, kits, and reagents for treatment or prevention of disease or conditions in humans and other mammals. The active therapeutic agents of the present disclosure include modified mRNA, cells containing modified mRNA or polypeptides translated from the modified mRNA, polypeptides translated from modified mRNA, and cells contacted with cells containing modified mRNA or polypeptides translated from the modified mRNA.

[0162] Described are combination therapeutics containing one or more modified mRNA containing translatable regions that encode for a protein or proteins that boost a mammalian subject's immunity along with a protein that induces antibody-dependent cellular toxicity. For example, therapeutics containing one or more mRNA that encode trastuzumab and granulocyte-colony stimulating factor (G-CSF). In particular, such combination therapeutics are useful in Her2+ breast cancer patients who develop induced resistance to trastuzumab. (See, e.g., Albrecht, Immunotherapy. 2(6):795-8 (2010)). The mRNA of the invention is useful in

methods of inducing translation of a recombinant polypeptide in a cell population. Such translation can be *in vivo*, *ex vivo*, *in culture*, or *in vitro*. The cell population is contacted with an effective amount of a composition containing the mRNA encoding the recombinant polypeptide. The population is contacted under conditions such that the mRNA is localized into one or more cells of the cell population and the recombinant polypeptide is translated in the cell from the nucleic acid.

[0163] An effective amount of the composition is provided based, at least in part, on the target tissue, target cell type, means of administration, physical characteristics of the nucleic acid (e.g., size, and extent of modified nucleosides), and other determinants. In general, an effective amount of the composition provides efficient protein production in the cell, more efficient than a composition containing a corresponding unmodified nucleic acid. Increased efficiency may be demonstrated by increased cell transfection (i.e., the percentage of cells transfected with the nucleic acid), increased protein translation from the nucleic acid, decreased nucleic acid degradation (as demonstrated, e.g., by increased duration of protein translation from a modified nucleic acid), or reduced innate immune response of the host cell.

[0164] Also described are methods of inducing *in vivo* translation of a recombinant polypeptide in a mammalian subject in need thereof. Therein, an effective amount of a composition containing a nucleic acid that has at least one nucleoside modification and a translatable region encoding the recombinant polypeptide is administered to the subject using the delivery methods described herein. The nucleic acid is provided in an amount and under other conditions such that the nucleic acid is localized into a cell of the subject and the recombinant polypeptide is translated in the cell from the nucleic acid. The cell in which the nucleic acid is localized, or the tissue in which the cell is present, may be targeted with one or more than one rounds of nucleic acid administration.

[0165] Other aspects of the present disclosure relate to transplantation of cells containing modified nucleic acids to a mammalian subject. Administration of cells to mammalian subjects is known to those of ordinary skill in the art, such as local implantation (e.g., topical or subcutaneous administration), organ delivery or systemic injection (e.g., intravenous injection or inhalation), as is the formulation of cells in pharmaceutically acceptable carrier. Compositions containing modified nucleic acids are formulated for administration intramuscularly, transarterially, intraperitoneally, intravenously, intranasally, subcutaneously, endoscopically, transdermally, or intrathecally. In some cases, the composition is formulated for extended release.

[0166] The subject to whom the therapeutic agent is administered suffers from or is at risk of developing a disease, disorder, or deleterious condition.

[0167] The administered modified nucleic acid may direct production of one or more recombinant polypeptides that provide a functional activity which is substantially absent in the cell in which the recombinant polypeptide is translated. For example, the missing functional activity may be enzymatic, structural, or gene regulatory in nature.

[0168] In other cases , the administered modified nucleic acid directs production of one or more recombinant polypeptides that replace a polypeptide (or multiple polypeptides) that is substantially absent in the cell in which the recombinant polypeptide is translated. Such absence may be due to genetic mutation of the encoding gene or regulatory pathway thereof. Alternatively, the recombinant polypeptide functions to antagonize the activity of an endogenous protein present in, on the surface of, or secreted from the cell. Usually, the activity of the endogenous protein is deleterious to the subject, for example, do to mutation of the endogenous protein resulting in altered activity or localization. Additionally, the recombinant polypeptide antagonizes, directly or indirectly, the activity of a biological moiety present in, on the surface of, or secreted from the cell. Examples of antagonized biological moieties include lipids (e.g., cholesterol), a lipoprotein (e.g., low density lipoprotein), a nucleic acid, a carbohydrate, or a small molecule toxin.

[0169] The recombinant proteins described herein are engineered for localization within the cell, potentially within a specific compartment such as the nucleus, or are engineered for secretion from the cell or translocation to the plasma membrane of the cell.

[0170] As described herein, a useful feature of the modified mRNA of the present invention is the capacity to reduce the innate immune response of a cell to an exogenous nucleic acid. This allows for performing the titration, reduction or elimination of the immune response in a cell or a population of cells. In some cases , the cell is contacted with a first composition that contains a first dose of a first exogenous nucleic acid including a translatable region and at least one nucleoside modification, and the level of the innate immune response of the cell to the first exogenous nucleic acid is determined. Subsequently, the cell is contacted with a second composition, which includes a second dose of the first exogenous nucleic acid, the second dose containing a lesser amount of the first exogenous nucleic acid as compared to the first dose. Alternatively, the cell is contacted with a first dose of a second exogenous nucleic acid. The second exogenous nucleic acid may contain one or more modified nucleosides, which may be the same or different from the first exogenous nucleic acid or, alternatively, the second exogenous nucleic acid may not contain modified nucleosides. The steps of contacting the cell with the first composition and/or the second composition may be repeated one or more times. Additionally, efficiency of protein production (e.g., protein translation) in the cell is optionally determined, and the cell may be re-transfected with the first and/or second composition repeatedly until a target protein production efficiency is achieved.

Therapeutics for diseases and conditions

[0171] The mRNA of the invention as defined in the claims can be used in methods for treating or preventing a symptom of diseases characterized by missing or aberrant protein activity, by replacing the missing protein activity or overcoming the aberrant protein activity. Because of the rapid initiation of protein production following introduction of modified mRNAs, as compared to viral DNA vectors, the compounds of the present disclosure are particularly advantageous in

treating acute diseases such as sepsis, stroke, and myocardial infarction. Moreover, the lack of transcriptional regulation of the modified mRNAs of the present disclosure is advantageous in that accurate titration of protein production is achievable.

[0172] Diseases characterized by dysfunctional or aberrant protein activity include, but not limited to, cancer and proliferative diseases, genetic diseases (e.g., cystic fibrosis), autoimmune diseases, diabetes, neurodegenerative diseases, cardiovascular diseases, and metabolic diseases. The mRNA of the invention as defined in the claims can be used in a method for treating such conditions or diseases in a subject by introducing nucleic acid or cell-based therapeutics containing the modified mRNA provided herein, wherein the modified nucleic acids encode for a protein that antagonizes or otherwise overcomes the aberrant protein activity present in the cell of the subject. Specific examples of a dysfunctional protein are the missense mutation variants of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which produce a dysfunctional protein variant of CFTR protein, which causes cystic fibrosis.

[0173] Multiple diseases are characterized by missing (or substantially diminished such that proper protein function does not occur) protein activity. Such proteins may not be present, or are essentially non-functional. The mRNA of the invention as defined in the claims can be used in a method for treating such conditions or diseases in a subject by introducing the mRNA or cell-based therapeutics containing the mRNA, wherein the mRNA encodes for a protein that replaces the protein activity missing from the target cells of the subject. Specific examples of a dysfunctional protein are the nonsense mutation variants of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which produce a nonfunctional protein variant of CFTR protein, which causes cystic fibrosis.

[0174] Thus, methods of treating cystic fibrosis in a mammalian subject may include contacting a cell of the subject with a modified mRNA having a translatable region that encodes a functional CFTR polypeptide, under conditions such that an effective amount of the CFTR polypeptide is present in the cell. Preferred target cells are epithelial cells, such as the lung, and methods of administration are determined in view of the target tissue; i.e., for lung delivery, the RNA molecules are formulated for administration by inhalation.

[0175] Also described is a method for treating hyperlipidemia in a subject, by introducing into a cell population of the subject with a modified mRNA molecule encoding Sortilin, a protein recently characterized by genomic studies, thereby ameliorating the hyperlipidemia in a subject. The *SORT1* gene encodes a trans-Golgi network (TGN) transmembrane protein called Sortilin. Genetic studies have shown that one of five individuals has a single nucleotide polymorphism, rs12740374, in the *lpl3* locus of the *SORT1* gene that predisposes them to having low levels of low-density lipoprotein (LDL) and very-low-density lipoprotein (VLDL). Each copy of the minor allele, present in about 30% of people, alters LDL cholesterol by 8 mg/dL, while two copies of the minor allele, present in about 5% of the population, lowers LDL cholesterol 16 mg/dL. Carriers of the minor allele have also been shown to have a 40% decreased risk of myocardial infarction. Functional *in vivo* studies in mice describes that

overexpression of *SORT1* in mouse liver tissue led to significantly lower LDL-cholesterol levels, as much as 80% lower, and that silencing SORT1 increased LDL cholesterol approximately 200% (Musunuru K et al. From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. Nature 2010; 466: 714-721).

Methods of cellular nucleic acid delivery

[0176] Methods described herein enhance nucleic acid delivery into a cell population, *in vivo*, *ex vivo*, or *in culture*. For example, a cell culture containing a plurality of host cells (e.g., eukaryotic cells such as yeast or mammalian cells) is contacted with a composition that contains an enhanced nucleic acid having at least one nucleoside modification and, optionally, a translatable region. The composition also generally contains a transfection reagent or other compound that increases the efficiency of enhanced nucleic acid uptake into the host cells. The enhanced nucleic acid exhibits enhanced retention in the cell population, relative to a corresponding unmodified nucleic acid. The retention of the enhanced nucleic acid is greater than the retention of the unmodified nucleic acid. In some embodiments, it is at least about 50%, 75%, 90%, 95%, 100%, 150%, 200% or more than 200% greater than the retention of the unmodified nucleic acid. Such retention advantage may be achieved by one round of transfection with the enhanced nucleic acid, or may be obtained following repeated rounds of transfection.

[0177] In some cases, the enhanced nucleic acid is delivered to a target cell population with one or more additional nucleic acids. Such delivery may be at the same time, or the enhanced nucleic acid is delivered prior to delivery of the one or more additional nucleic acids. The additional one or more nucleic acids may be modified nucleic acids or unmodified nucleic acids. It is understood that the initial presence of the enhanced nucleic acids does not substantially induce an innate immune response of the cell population and, moreover, that the innate immune response will not be activated by the later presence of the unmodified nucleic acids. In this regard, the enhanced nucleic acid may not itself contain a translatable region, if the protein desired to be present in the target cell population is translated from the unmodified nucleic acids.

Targeting Moieties

[0178] In some cases, modified nucleic acids are provided to express a protein-binding partner or a receptor on the surface of the cell, which functions to target the cell to a specific tissue space or to interact with a specific moiety, either *in vivo* or *in vitro*. Suitable protein-binding partners include antibodies and functional fragments thereof, scaffold proteins, or peptides. Additionally, modified nucleic acids can be employed to direct the synthesis and extracellular localization of lipids, carbohydrates, or other biological moieties.

Permanent Gene Expression Silencing

[0179] A method for epigenetically silencing gene expression in a mammalian subject, comprising a nucleic acid where the translatable region encodes a polypeptide or polypeptides capable of directing sequence-specific histone H3 methylation to initiate heterochromatin formation and reduce gene transcription around specific genes for the purpose of silencing the gene. For example, a gain-of-function mutation in the Janus Kinase 2 gene is responsible for the family of Myeloproliferative Diseases.

Pharmaceutical Compositions

[0180] Proteins can be generated from modified mRNAs. Pharmaceutical compositions may optionally comprise one or more additional therapeutically active substances. A method of administering pharmaceutical compositions comprising one or more proteins to be delivered to a subject in need thereof is described. In some cases, compositions are administered to humans. For the purposes of the present disclosure, the phrase "active ingredient" generally refers to a protein or protein-containing complex as described herein.

[0181] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to animals of all sorts. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with merely ordinary, if any, experimentation. Subjects to which administration of the pharmaceutical compositions is contemplated include, but are not limited to, humans and/or other primates; mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, cats, dogs, mice, and/or rats; and/or birds, including commercially relevant birds such as chickens, ducks, geese, and/or turkeys.

[0182] Formulations of the pharmaceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of bringing the active ingredient into association with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, shaping and/or packaging the product into a desired single- or multi-dose unit.

[0183] A pharmaceutical composition in accordance with the present disclosure may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a "unit dose" is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be

administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[0184] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the present disclosure will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition may comprise between 0.1% and 100% (w/w) active ingredient.

[0185] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's The Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro (Lippincott, Williams & Wilkins, Baltimore, MD, 2006)

[0186] discloses various excipients used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional excipient medium is incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated in this present disclosure.

[0187] A pharmaceutically acceptable excipient may be at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% pure. In some cases, an excipient is approved for use in humans and for veterinary use. In some cases, an excipient is approved by United States Food and Drug Administration. In some cases, an excipient is pharmaceutical grade. In some cases, an excipient meets the standards of the United States Pharmacopoeia (USP), the European Pharmacopoeia (EP), the British Pharmacopoeia, and/or the International Pharmacopoeia.

[0188] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in pharmaceutical formulations. Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and/or perfuming agents can be present in the composition, according to the judgment of the formulator.

[0189] Exemplary diluents include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, *etc.*,

and/or combinations thereof.

[0190] Exemplary granulating and/or dispersing agents include, but are not limited to, potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (Veegum), sodium lauryl sulfate, quaternary ammonium compounds, *etc.*, and/or combinations thereof.

[0191] Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (e.g. bentonite [aluminum silicate] and Veegum[®] [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g. carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (e.g. carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (e.g. polyoxyethylene sorbitan monolaurate [Tween[®]20], polyoxyethylene sorbitan [Tween[®]60], polyoxyethylene sorbitan monooleate [Tween[®]80], sorbitan monopalmitate [Span[®]40], sorbitan monostearate [Span[®]60], sorbitan tristearate [Span[®]65], glyceryl monooleate, sorbitan monooleate [Span[®]80]), polyoxyethylene esters (e.g. polyoxyethylene monostearate [Myrj[®]45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and Solutol[®]), sucrose fatty acid esters, polyethylene glycol fatty acid esters (e.g. Cremophor[®]), polyoxyethylene ethers, (e.g. polyoxyethylene lauryl ether [Brij[®]30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, Pluronic[®]F 68, Poloxamer[®]188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, *etc.* and/or combinations thereof.

[0192] Exemplary binding agents include, but are not limited to, starch (e.g. cornstarch and starch paste); gelatin; sugars (e.g. sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol,); natural and synthetic gums (e.g. acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum[®]), and larch arabogalactan); alginates; polyethylene

oxide; polyethylene glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; *etc.*; and combinations thereof.

[0193] Exemplary preservatives may include, but are not limited to, antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and/or other preservatives. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and/or sodium sulfite. Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate, edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and/or trisodium edetate. Exemplary antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and/or thimerosal. Exemplary antifungal preservatives include, but are not limited to, butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and/or sorbic acid. Exemplary alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and/or phenylethyl alcohol. Exemplary acidic preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and/or phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, Glydant Plus[®], Phenonip[®], methylparaben, Germall[®]115, Germaben[®]II, Neolone[™], Kathon[™], and/or Euxyl[®].

[0194] Exemplary buffering agents include, but are not limited to, citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, d-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, *etc.*, and/or combinations thereof.

[0195] Exemplary lubricating agents include, but are not limited to, magnesium stearate,

calcium stearate, stearic acid, silica, talc, malt, glyceryl behanate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, *etc.*, and combinations thereof.

[0196] Exemplary oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and/or combinations thereof.

[0197] Liquid dosage forms for oral and parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as Cremophor[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

[0198] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

[0199] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to

use.

[0200] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[0201] Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing compositions with suitable non-irritating excipients such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

[0202] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, an active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient such as sodium citrate or dicalcium phosphate and/or fillers or extenders (e.g. starches, lactose, sucrose, glucose, mannitol, and silicic acid), binders (e.g. carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia), humectants (e.g. glycerol), disintegrating agents (e.g. agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate), solution retarding agents (e.g. paraffin), absorption accelerators (e.g. quaternary ammonium compounds), wetting agents (e.g. cetyl alcohol and glycerol monostearate), absorbents (e.g. kaolin and bentonite clay), and lubricants (e.g. talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate), and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may comprise buffering agents.

[0203] Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. Solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type may be employed as fillers in soft and hard-filled

gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0204] Dosage forms for topical and/or transdermal administration of a composition may include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants and/or patches. Generally, an active ingredient is admixed under sterile conditions with a pharmaceutically acceptable excipient and/or any needed preservatives and/or buffers as may be required. Additionally, the present disclosure contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms may be prepared, for example, by dissolving and/or dispensing the compound in the proper medium. Alternatively or additionally, rate may be controlled by either providing a rate controlling membrane and/or by dispersing the compound in a polymer matrix and/or gel.

[0205] Suitable devices for use in delivering intradermal pharmaceutical compositions described herein include short needle devices such as those described in U.S. Patents 4,886,499; 5,190,521; 5,328,483; 5,527,288; 4,270,537; 5,015,235; 5,141,496; and 5,417,662. Intradermal compositions may be administered by devices which limit the effective penetration length of a needle into the skin, such as those described in PCT publication WO 99/34850 and functional equivalents thereof. Jet injection devices which deliver liquid compositions to the dermis via a liquid jet injector and/or via a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Jet injection devices are described, for example, in U.S. Patents 5,480,381; 5,599,302; 5,334,144; 5,993,412; 5,649,912; 5,569,189; 5,704,911; 5,383,851; 5,893,397; 5,466,220; 5,339,163; 5,312,335; 5,503,627; 5,064,413; 5,520,639; 4,596,556; 4,790,824; 4,941,880; 4,940,460; and PCT publications WO 97/37705 and WO 97/13537. Ballistic powder/particle delivery devices which use compressed gas to accelerate vaccine in powder form through the outer layers of the skin to the dermis are suitable. Alternatively or additionally, conventional syringes may be used in the classical mantoux method of intradermal administration.

[0206] Formulations suitable for topical administration include, but are not limited to, liquid and/or semi liquid preparations such as liniments, lotions, oil in water and/or water in oil emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions. Topically-administrable formulations may, for example, comprise from about 1% to about 10% (w/w) active ingredient, although the concentration of active ingredient may be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration may further comprise one or more of the additional ingredients described herein.

[0207] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration via the buccal cavity. Such a formulation may comprise dry particles which comprise the active ingredient and which have a diameter in the range from about 0.5 nm to about 7 nm or from about 1 nm to about 6 nm. Such compositions are conveniently in the form of dry powders for administration using a device comprising a dry powder reservoir to which a stream of propellant may be directed to disperse the powder

and/or using a self propelling solvent/powder dispensing container such as a device comprising the active ingredient dissolved and/or suspended in a low-boiling propellant in a sealed container. Such powders comprise particles wherein at least 98% of the particles by weight have a diameter greater than 0.5 nm and at least 95% of the particles by number have a diameter less than 7 nm. Alternatively, at least 95% of the particles by weight have a diameter greater than 1 nm and at least 90% of the particles by number have a diameter less than 6 nm. Dry powder compositions may include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

[0208] Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50% to 99.9% (w/w) of the composition, and active ingredient may constitute 0.1% to 20% (w/w) of the composition. A propellant may further comprise additional ingredients such as a liquid nonionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

[0209] Pharmaceutical compositions formulated for pulmonary delivery may provide an active ingredient in the form of droplets of a solution and/or suspension. Such formulations may be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising active ingredient, and may conveniently be administered using any nebulization and/or atomization device. Such formulations may further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. Droplets provided by this route of administration may have an average diameter in the range from about 0.1 nm to about 200 nm.

[0210] Formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutical composition. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 μm to 500 μm . Such a formulation is administered in the manner in which snuff is taken, *i.e.* by rapid inhalation through the nasal passage from a container of the powder held close to the nose.

[0211] Formulations suitable for nasal administration may, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of active ingredient, and may comprise one or more of the additional ingredients described herein. A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may, for example, 0.1% to 20% (w/w) active ingredient, the balance comprising an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations suitable for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising active ingredient. Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, may have an average particle and/or droplet size in the range

from about 0.1 nm to about 200 nm, and may further comprise one or more of any additional ingredients described herein.

[0212] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for ophthalmic administration. Such formulations may, for example, be in the form of eye drops including, for example, a 0.1/1.0% (w/w) solution and/or suspension of the active ingredient in an aqueous or oily liquid excipient. Such drops may further comprise buffering agents, salts, and/or one or more other of any additional ingredients described herein. Other ophthalmically-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this present disclosure.

[0213] General considerations in the formulation and/or manufacture of pharmaceutical agents may be found, for example, in Remington: The Science and Practice of Pharmacy 21st ed., Lippincott Williams & Wilkins, 2005.

[0214] Administration proteins or complexes in accordance with the present disclosure can be used in methods comprising administering proteins or complexes in accordance with the present disclosure to a subject in need thereof. Proteins or complexes, or pharmaceutical, imaging, diagnostic, or prophylactic compositions thereof, may be administered to a subject using any amount and any route of administration effective for preventing, treating, diagnosing, or imaging a disease, disorder, and/or condition (e.g., a disease, disorder, and/or condition relating to working memory deficits). The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease, the particular composition, its mode of administration, its mode of activity, and the like. Compositions are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present disclosure will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or appropriate imaging dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[0215] Proteins to be delivered and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof may be administered to animals, such as mammals (e.g., humans, domesticated animals, cats, dogs, mice, rats, *etc.*). Pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof may be administered to humans.

[0216] Proteins to be delivered and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof may be administered by any route. Proteins and/or pharmaceutical,

prophylactic, diagnostic, or imaging compositions thereof, may be administered by one or more of a variety of routes, including oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, transdermal, interdermal, rectal, intravaginal, intraperitoneal, topical (e.g. by powders, ointments, creams, gels, lotions, and/or drops), mucosal, nasal, buccal, enteral, vitreal, intratumoral, sublingual; by intratracheal instillation, bronchial instillation, and/or inhalation; as an oral spray, nasal spray, and/or aerosol, and/or through a portal vein catheter. In some cases, proteins or complexes, and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof, are administered by systemic intravenous injection. In specific cases, proteins or complexes and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof may be administered intravenously and/or orally. In specific cases, proteins or complexes, and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof, may be administered in a way which allows the protein or complex to cross the blood-brain barrier, vascular barrier, or other epithelial barrier.

[0217] However, the present disclosure relates to the delivery of proteins or complexes, and/or pharmaceutical, prophylactic, diagnostic, or imaging compositions thereof, by any appropriate route taking into consideration likely advances in the sciences of drug delivery.

[0218] In general the most appropriate route of administration will depend upon a variety of factors including the nature of the protein or complex comprising proteins associated with at least one agent to be delivered (e.g., its stability in the environment of the gastrointestinal tract, bloodstream, etc.), the condition of the patient (e.g., whether the patient is able to tolerate particular routes of administration), etc. The present disclosure relates to the delivery of the pharmaceutical, prophylactic, diagnostic, or imaging compositions by any appropriate route taking into consideration likely advances in the sciences of drug delivery.

[0219] Compositions in accordance with the present disclosure may be administered at dosage levels sufficient to deliver from about 0.0001 mg/kg to about 100 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, from about 0.1 mg/kg to about 40 mg/kg, from about 0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, or from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic, diagnostic, prophylactic, or imaging effect. The desired dosage may be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. The desired dosage may be delivered using multiple administrations (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

[0220] Proteins or complexes may be used in combination with one or more other therapeutic, prophylactic, diagnostic, or imaging agents. By "in combination with," it is not intended to imply that the agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope of the present disclosure. Compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. In general, each agent will be administered

at a dose and/or on a time schedule determined for that agent. In some cases, the present disclosure encompasses the delivery of pharmaceutical, prophylactic, diagnostic, or imaging compositions in combination with agents that improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body.

[0221] It will further be appreciated that therapeutically, prophylactically, diagnostically, or imaging active agents utilized in combination may be administered together in a single composition or administered separately in different compositions. In general, it is expected that agents utilized in combination will be utilized at levels that do not exceed the levels at which they are utilized individually. In some cases, the levels utilized in combination will be lower than those utilized individually.

[0222] The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, a composition useful for treating cancer in accordance with the present disclosure may be administered concurrently with a chemotherapeutic agent), or they may achieve different effects (e.g., control of any adverse effects).

Kits

[0223] Also described are a variety of kits for conveniently and/or effectively carrying out methods of the present disclosure. Typically kits will comprise sufficient amounts and/or numbers of components to allow a user to perform multiple treatments of a subject(s) and/or to perform multiple experiments.

[0224] Described in one aspect are kits for protein production, comprising a first isolated nucleic acid comprising a translatable region and a nucleic acid modification, wherein the nucleic acid is capable of evading an innate immune response of a cell into which the first isolated nucleic acid is introduced, and packaging and instructions.

[0225] Described in one aspect are kits for protein production, comprising: a first isolated nucleic acid comprising a translatable region, provided in an amount effective to produce a desired amount of a protein encoded by the translatable region when introduced into a target cell; a second nucleic acid comprising an inhibitory nucleic acid, provided in an amount effective to substantially inhibit the innate immune response of the cell; and packaging and instructions.

[0226] Described in one aspect are kits for protein production, comprising a first isolated nucleic acid comprising a translatable region and a nucleoside modification, wherein the nucleic acid exhibits reduced degradation by a cellular nuclease, and packaging and instructions.

[0227] Described in one aspect are kits for protein production, comprising a first isolated nucleic acid comprising a translatable region and at least two different nucleoside modifications, wherein the nucleic acid exhibits reduced degradation by a cellular nuclease, and packaging and instructions.

[0228] Described in one aspect are kits for protein production, comprising a first isolated nucleic acid comprising a translatable region and at least one nucleoside modification, wherein the nucleic acid exhibits reduced degradation by a cellular nuclease; a second nucleic acid comprising an inhibitory nucleic acid; and packaging and instructions.

[0229] The first isolated nucleic acid may comprise messenger RNA (mRNA). The mRNA may comprise at least one nucleoside selected from the group consisting of pyridin-4-one ribonucleoside, 5-aza-uridine, 2-thio-5-aza-uridine, 2-thiouridine, 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxyuridine, 3-methyluridine, 5-carboxymethyl-uridine, 1-carboxymethyl-pseudouridine, 5-propynyl-uridine, 1-propynyl-pseudouridine, 5-taurinomethyluridine, 1-taurinomethyl-pseudouridine, 5-taurinomethyl-2-thio-uridine, 1-taurinomethyl-4-thio-uridine, 5-methyl-uridine, 1-methyl-pseudouridine, 4-thio-1-methyl-pseudouridine, 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouridine, dihydropseudouridine, 2-thio-dihydrouridine, 2-thio-dihydropseudouridine, 2-methoxyuridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, and 4-methoxy-2-thio-pseudouridine.

[0230] The mRNA may comprise at least one nucleoside selected from the group consisting of 5-aza-cytidine, pseudoisocytidine, 3-methyl-cytidine, N4-acetylcytidine, 5-formylcytidine, N4-methylcytidine, 5-hydroxymethylcytidine, 1-methyl-pseudoisocytidine, pyrrolo-cytidine, pyrrolo-pseudoisocytidine, 2-thio-cytidine, 2-thio-5-methyl-cytidine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytidine, 2-methoxy-5-methyl-cytidine, 4-methoxy-pseudoisocytidine, and 4-methoxy-1-methyl-pseudoisocytidine.

[0231] The mRNA may comprise at least one nucleoside selected from the group consisting of 2-aminopurine, 2, 6-diaminopurine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-aminopurine, 7-deaza-8-aza-2-aminopurine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyladenosine, N6-methyladenosine, N6-isopentenyladenosine, N6-(cis-hydroxyisopentenyl)adenosine, 2-methylthio-N6-(cis-hydroxyisopentenyl) adenosine, N6-glycincylcarbamoyladenosine, N6-threonylcarbamoyladenosine, 2-methylthio-N6-threonyl carbamoyladenosine, N6,N6-dimethyladenosine, 7-methyladenine, 2-methylthio-adenine, and 2-methoxy-adenine.

[0232] The mRNA may comprise at least one nucleoside selected from the group consisting of inosine, 1-methyl-inosine, wyosine, wybutosine, 7-deaza-guanosine, 7-deaza-8-aza-guanosine, 6-thio-guanosine, 6-thio-7-deaza-guanosine, 6-thio-7-deaza-8-aza-guanosine, 7-

methyl-guanosine, 6-thio-7-methyl-guanosine, 7-methylinosine, 6-methoxy-guanosine, 1-methylguanosine, N2-methylguanosine, N2,N2-dimethylguanosine, 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 1-methyl-6-thio-guanosine, N2-methyl-6-thio-guanosine, and N2,N2-dimethyl-6-thio-guanosine.

[0233] Described in another aspect are compositions for protein production, comprising a first isolated nucleic acid comprising a translatable region and a nucleoside modification, wherein the nucleic acid exhibits reduced degradation by a cellular nuclease, and a mammalian cell suitable for translation of the translatable region of the first nucleic acid. The invention provides an mRNA wherein 100% of nucleotides comprising uracil are replaced with nucleotides comprising N1-methyl-pseudouridine.

EXAMPLES

Example 1. Modified mRNA In Vitro Transcription

Materials and Methods

[0234] Modified mRNAs (modRNAs) were made using standard laboratory methods and materials for in vitro transcription with the exception that the nucleotide mix contained modified nucleotides. The open reading frame (ORF) of the gene of interest is flanked by a 5' untranslated region (UTR) containing a strong Kozak translational initiation signal and an alpha-globin 3' UTR terminating with an oligo(dT) sequence for templated addition of a polyA tail for modRNAs not incorporating Adenosine analogs. Adenosine-containing modRNAs were synthesized without an oligo (dT) sequence to allow for post-transcription poly (A) polymerase poly-(A) tailing. The modRNAs were modified by incorporating chemically modified nucleotides indicated in Table 3 (below) during the in vitro transcription with 100% replacement of the corresponding natural nucleotide or partial replacement of the corresponding natural nucleotide at the indicated percentage.

[0235] Table 3 indicates the chemical identity of each chemically-distinct modified nucleotide incorporated into a modified mRNA with the given chemistry designation number.

Table 3

Modified Nucleotide	Chemistry #	Modified Nucleotide Combination	Chemistry #
6-aza-cytidine	Chem 1	α -thio-cytidine/5-iodo-uridine	Chem 29
2-thio-cytidine	Chem 2	α -thio-cytidine/N1-methyl-pseudo-uridine	Chem 30
α -thio-cytidine	Chem 3	α -thio-cytidine/ α -thio-uridine	Chem 31

Modified Nucleotide	Chemistry #	Modified Nucleotide Combination	Chemistry #
Pseudo-iso-cytidine	Chem 4	α -thio-cytidine/5-methyl-uridine	Chem 32
5-aminoallyl-uridine	Chem 5	α -thio-cytidine/pseudo-uridine	Chem 33
5-iodo-uridine	Chem 6	Pseudo-iso-cytidine/5-iodo-uridine	Chem 34
N1-methyl-pseudouridine	Chem 7	Pseudo-iso-cytidine/N1-methyl-pseudo-uridine	Chem 35
5,6-dihydrouridine	Chem 8	Pseudo-iso-cytidine/ α -thio-uridine	Chem 36
α -thio-uridine	Chem 9	Pseudo-iso-cytidine/5-methyl-uridine	Chem 37
4-thio-uridine	Chem 10	Pseudo-iso-cytidine/Pseudo-uridine	Chem 38
6-aza-uridine	Chem 11	Pyrrolo-cytidine	Chem 39
5-hydroxy-uridine	Chem 12	Pyrrolo-cytidine/5-iodo-uridine	Chem 40
Deoxy-thymidine	Chem 13	Pyrrolo-cytidine/N1-methyl-pseudo-uridine	Chem 41
Pseudo-uridine	Chem 14	Pyrrolo-cytidine/ α -thio-uridine	Chem 42
Inosine	Chem 15	Pyrrolo-cytidine/5-methyl-uridine	Chem 43
α -thio-guanosine	Chem 16	Pyrrolo-cytidine/Pseudo-uridine	Chem 44
8-oxo-guanosine	Chem 17	5-methyl-cytidine/5-iodo-uridine	Chem 45
O6-methyl-guanosine	Chem 18	5-methyl-cytidine/N1-methyl-pseudo-uridine	Chem 46
7-deaza-guanosine	Chem 19	5-methyl-cytidine/ α -thio-uridine	Chem 47
No modification	Chem 20	5-methyl-cytidine/5-methyl-uridine	Chem 48
N1-methyl-adenosine	Chem 21	5-methyl-cytidine/Pseudo-uridine	Chem 49
2-amino-6-Chloro-purine	Chem 22	5-methyl-cytidine	Chem 50
N6-methyl-2-aminopurine	Chem 23	25% Pseudo-iso-cytidine	Chem 51
6-Chloro-purine	Chem 24	25% N1-methyl-pseudo-uridine	Chem 52
N6-methyl-adenosine	Chem 25	25% N1-Methyl-pseudo-uridine/75%-pseudo-uridine	Chem 53
α -thio-adenosine	Chem 26	5-methyl-uridine	Chem 54
8-azido-adenosine	Chem 27	5-iodo-cytidine	Chem 55
7-deaza-adenosine	Chem 28		

[0236] Agarose Gel Electrophoresis of modRNA: Individual modRNAs (200-400 ng in a 20 μ l volume) were loaded into a well on a non-denaturing 1.2% Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes according to the manufacturer protocol (FIG. 1A). Tables 4 and 5 below indicate the modified nucleotide (Table 4) or nucleic acid (Table 5) loaded in each lane. These data indicate which chemically modified nucleotides were transcribed into chemically-modified mRNAs and the quality of each individual modRNA. These data demonstrate that nucleotides with chemical modifications on the major groove and minor groove face of the nucleotide were capable of being transcribed into a modRNA.

Table 4

Lane	Modified NTP
1	α -thio-cytidine
2	Pseudo-iso-cytidine
3	5-aminoallyl-uridine
4	5-iodo-uridine
5	N1-methyl-pseudo-uridine
6	α -thio-uridine
7	4-thio-uridine
8	5-hydroxy-uridine
9	Deoxy-thymidine
10	Pseudo-uridine
11	Inosine
12	α -thio-guanosine
13	8-oxo-guanosine
14	N1-methyl-guanosine
15	O6-methyl-guanosine
16	No modification
17	N1-methyl-adenosine
18	2-amino-6-Chloro-purine
19	N6-methyl-2-amino-purine
20	6-Chloro-purine
21	α -thio-adenosine
22	8-azido-adenosine
23	7-deaza-adenosine
24	6-aza-cytidine
25	2-thio-cytidine
26	5,6-dihydro-uridine
27	6-aza-uridine
28	7-deaza-guanosine
29	N6-methyl-adenosine

Table 5

Lane	Modified NTP combination
1	α -thio-cytidine/5-iodo-uridine
2	α -thio-cytidine/N1-methyl-pseudouridine
3	α -thio-cytidine/ α -thio-uridine
4	α -thio-cytidine/5-methyl-uridine
5	α -thio-cytidine/pseudouridine
6	5-iodo-cytidine/5-iodo-uridine
7	5-iodo-cytidine/N1-methyl-pseudouridine
8	5-iodo-cytidine/ α -thio-uridine
9	5-iodo-cytidine/5-methyl-uridine
10	5-iodo-cytidine/pseudouridine

11	Pseudo-Iso-cytidine/5-iodo-uridine
12	Pyrrolo-cytidine
13	Pyrrolo-cytidine/5-iodo-uridine
14	Pyrrolo-cytidine/N1-methyl-pseudouridine
15	Pyrrolo-cytidine/ α -thio-uridine
16	Pyrrolo-cytidine/5-methyl-uridine
17	Pyrrolo-cytidine/pseudouridine
18	5-methyl-cytidine/5-iodo-uridine
19	5-methyl-cytidine/N1-methyl-uridine
20	5-methyl-cytidine/ α -thio-uridine
21	5-methyl-cytidine/5-methyl-uridine
22	5-methyl-cytidine/pseudouridine
23	Pseudo-Iso-cytidine/N1-methyl-pseudouridine
24	Pseudo-Iso-cytidine/ α -thio-uridine
25	Pseudo-Iso-cytidine/5-methyl-uridine
26	Pseudo-Iso-cytidine/pseudouridine
27	5-methyl-cytidine
28	25% pseudo-Iso-cytidine
29	25% N1-methyl-pseudouridine
30	25% N1-methyl-pseudouridine/75% pseudouridine

[0237] Agarose Gel Electrophoresis of RT-PCR products: Individual reverse transcribed-PCR products (200-400ng) were loaded into a well of a non-denaturing 1.2% Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes according to the manufacturer protocol (FIG. 1B). Table 5 below indicates the modified nucleotide loaded in each lane.

[0238] Nanodrop modRNA quantification and UV spectral data: modRNAs in TE buffer (1 μ l) were used for Nanodrop UV absorbance readings to quantitate the yield of each modRNA from an *in vitro* transcription reaction (UV absorbance traces are shown in Figures 6A-6L). These data indicate which chemically modified nucleotides were transcribed into chemically-modified mRNAs. These data also demonstrate that nucleotides with chemical modifications on the major groove and minor groove face of the nucleotide were capable of being transcribed into a modRNA. These data further demonstrate that the nucleotides of the present invention are transcription-competent and compatible with incorporation into a modRNA, which may have altered UV spectra due to the presence of a given modified nucleotide. For example, Pyrrolo-C containing modRNAs have an increase in UV absorbance at a lower wavelength due to the presence of the pyrrolo ring of the modified C nucleotide. In another example, 2-amino-adenine nucleotide-containing modRNAs have an increase in UV absorbance at a higher wavelength due to the presence of an exocyclic amine off the purine ring. Nucleotides that are not transcription-competent and cannot be incorporated into a modRNA have a scrambled UV spectrum indicating no product from the transcription reaction.

Example 2. Modified RNA Transfection

[0239] Reverse Transfection: For experiments performed in a 24-well collagen-coated tissue culture plate, Keratinocytes were seeded at a cell density of 1×10^5 . For experiments

performed in a 96-well collagen-coated tissue culture plate, Keratinocytes were seeded at a cell density of 0.5×10^5 . For each modRNA to be transfected, modRNA: RNAiMAX was prepared as described and mixed with the cells in the multi-well plate within a period of time, e.g., 6 hours, of cell seeding before cells had adhered to the tissue culture plate.

[0240] Forward Transfection: In a 24-well collagen-coated tissue culture plate, Keratinocytes were seeded at a cell density of 0.7×10^5 . For experiments performed in a 96-well collagen-coated tissue culture plate, Keratinocytes were seeded at a cell density of 0.3×10^5 . Keratinocytes were then grown to a confluency of >70% for over 24 hours. For each modRNA to be transfected, modRNA: RNAiMAX was prepared as described and transfected onto the cells in the multi-well plate over 24 hours after cell seeding and adherence to the tissue culture plate.

modRNA Translation Screen: G-CSF ELISA

[0241] FIGs. 2A and 2B show an Enzyme-linked immunosorbent assay (ELISA) for Human Granulocyte-Colony Stimulating Factor (hu-G-CSF) of *in vitro* transfected Human Keratinocyte cells. Keratinocytes were grown in EpiLife medium with Supplement S7 from Invitrogen at a confluence of >70%. FIG. 2A keratinocytes were reverse transfected with 300 ng of the indicated chemically modified mRNA complexed with RNAiMAX from Invitrogen. FIG. 2B keratinocytes were forward transfected with 300 ng modRNA complexed with RNAiMAX from Invitrogen. The RNA:RNAiMAX complex was formed by first incubating the RNA with Supplement-free EpiLife media in a 5X volumetric dilution for 10 minutes at room temperature. In a second vial, RNAiMAX reagent was incubated with Supplement-free EpiLife Media in a 10X volumetric dilution for 10 minutes at room temperature. The RNA vial was then mixed with the RNAiMAX vial and incubated for 20-30 at room temperature before being added to the cells in a drop-wise fashion. Secreted huG-CSF concentration in the culture medium was measured at 18 hours post-transfection for each of the chemically modified mRNAs in triplicate. Secretion of Human Granulocyte-Colony Stimulating Factor (G-CSF) from transfected human keratinocytes was quantified using an ELISA kit from Invitrogen or R&D Systems (Minneapolis, MN) following the manufacturers recommended instructions. These data show that huG-CSF modRNAs comprised of chemically distinct nucleotide analogs (SEQ ID NO: 2) is capable of being translated in Human Keratinocyte cells and that huG-CSF is transported out of the cells and released into the extracellular environment. These data indicate which modified nucleotides were translated into protein when incorporated into a chemically modified mRNA. These data show that modified RNA containing nucleotides with chemical modifications on the major groove face of pyrimidine analogs have the highest levels of secreted hu-G-CSF into the cell culture medium.

modRNA Dose and Duration: G-CSF ELISA

[0242] FIGs. 3A-N show Enzyme-linked immunosorbent assays (ELISA) for Human Granulocyte-Colony Stimulating Factor (G-CSF) of *in vitro* transfected Human Keratinocyte cells. Keratinocytes were grown in EpiLife medium with Supplement S7 from Invitrogen at a confluence of >70%. Keratinocytes were reverse transfected with 0ng, 46.875ng, 93.75ng, 187.5ng, 375ng, 750ng, or 1500ng modRNA complexed with RNAiMAX from Invitrogen. The modRNA:RNAiMAX complex was formed as described. Secreted huG-CSF concentration in the culture medium was measured at 0, 6, 12, 24, and 48 hours post-transfection for each concentration of each modRNA in triplicate. Secretion of Human Granulocyte-Colony Stimulating Factor (G-CSF) from transfected human keratinocytes was quantified using an ELISA kit from Invitrogen or R&D Systems following the manufacturers recommended instructions. These data show that huG-CSF modRNAs comprised of chemically distinct nucleotide analogs (SEQ ID NO: X and Table 6) secreted hu-G-CSF protein in a modRNA dose-dependent manner from Human Keratinocyte cells and that huG-CSF is transported out of the cells and released into the extracellular environment. These data indicate which modified RNAs containing modified nucleotide analogs sustain hu-G-CSF expression for the longest and at the highest levels. These data show that modified RNA containing modified nucleotides with chemical modifications on the major groove face of pyrimidine analogs have the highest levels of secreted hu-G-CSF into the cell culture medium and that 750ng of modRNA elicits the highest level of secreted hu-G-CSF.

Example 3. Cellular Innate Immune Response to modRNA

IFN- β ELISA and TNF- α ELISA:

[0243] FIGs. 4A-F show an Enzyme-linked immunosorbent assay (ELISA) for Human Tumor Necrosis Factor- α (TNF- α) (FIGs. 4A and 4B); Human Interferon- β (IFN- β) (FIGs. 4C and 4D); and Human Granulocyte-Colony Stimulating Factor (G-CSF) (FIGs. 4E and 4F) secreted from *in vitro*-transfected Human Keratinocyte cells. Keratinocytes were grown in EpiLife medium with Human Keratinocyte Growth Supplement in the absence of hydrocortisone from Invitrogen at a confluence of >70%. In FIGs. 4A and 4B, keratinocytes were reverse transfected with 0ng, 93.75ng, 187.5ng, 375ng, 750ng, 1500ng or 3000ng of the indicated chemically modified mRNA complexed with RNAiMAX from Invitrogen as described in triplicate. Secreted TNF- α in the culture medium was measured 24 hours post-transfection for each of the chemically modified mRNAs using an ELISA kit from Invitrogen according to the manufacturer protocols.

[0244] In FIGs. 4C and 4D, secreted IFN- β in the same culture medium was measured 24 hours post-transfection for each of the chemically modified mRNAs using an ELISA kit from Invitrogen according to the manufacturer protocols. In FIGs. 4E and 4F, secreted hu-G-CSF concentration in the same culture medium was measured at 24 hours post-transfection for each of the chemically modified mRNAs. Secretion of Human Granulocyte-Colony Stimulating Factor (G-CSF) from transfected human keratinocytes was quantified using an ELISA kit from

Invitrogen or R&D Systems (Minneapolis, MN) following the manufacturers recommended instructions. These data indicate which modified RNAs containing modified nucleotides were capable of eliciting a reduced cellular innate immune response in comparison to natural and other chemically modified nucleotides by measuring exemplary type I cytokines TNF- α and IFN- β . These data show that modified RNAs containing modified nucleotides with chemical modifications on the major groove face of pyrimidine analogs have the lowest levels of secreted TNF- α and IFN- β into the cell culture medium while maintaining high levels of modRNA-encoding hu-G-CSF secretion into the cell culture medium.

Example 4. Human Granulocyte-Colony Stimulating Factor-modified RNA-induced Cell Proliferation Assay

[0245] FIGs. 5A-D show modRNA-encoding hu-G-CSF produced by a human keratinocyte feeder cell layer induced the proliferation of both human myeloblast cells KG-1 and Kasumi-1 that express the G-CSF-receptor where the cell populations are separated by a semi-permeable membrane.

[0246] Human keratinocytes were grown in EpiLife medium with Supplement S7 from Invitrogen at a confluence of >70% in a 24-well collagen-coated Transwell® (Corning, Lowell, MA) co-culture tissue culture plate. Keratinocytes were reverse transfected with 750ng of the indicated chemically modified mRNA complexed with RNAiMAX from Invitrogen as described in triplicate. The modRNA:RNAiMAX complex was formed as described. Keratinocyte media was exchanged 6-8 hours post-transfection. 42-hours post-transfection, the 24-well Transwell® plate insert with a 0.4 μ m-pore semi-permeable polyester membrane was placed into the hu-G-CSF modRNA-transfected keratinocyte containing culture plate. FIG. 5A is a table showing the results from an Enzyme-linked immunosorbent assay (ELISA) for human-G-CSF secreted from *in vitro*-transfected Human Keratinocyte cells sampled from individual wells in a co-culture 24-well tissue culture plate 42 hours post-transfection with 750ng of each indicated hu-G-CSF-encoding modRNA.

[0247] Human myeloblast cells, Kasumi-1 cells (FIG. 5C) or KG-1 (FIG. 5D) (0.2×10^5 cells), were seeded into the insert well and cell proliferation was quantified 42 hours post-co-culture initiation using the CyQuant Direct Cell Proliferation Assay (Invitrogen) in a 100-120 μ l volume in a 96-well plate. modRNA-encoding hu-G-CSF-induced myeloblast cell proliferation was expressed as a percent cell proliferation normalized to untransfected keratinocyte/myeloblast co-culture control wells. Secreted hu-G-CSF concentration in both the keratinocyte and myeloblast insert co-culture wells was measured at 42 hours post-co-culture initiation for each modRNA in duplicate. Secretion of Human Granulocyte-Colony Stimulating Factor (G-CSF) was quantified using an ELISA kit from Invitrogen following the manufacturers recommended instructions.

[0248] Transfected hu-G-CSF modRNA in human keratinocyte feeder cells and untransfected

human myeloblast cells were detected by RT-PCR. Total RNA from sample cells was extracted and lysed using RNeasy kit (Qiagen, Valencia, CA) according to the manufacturer instructions. Extracted total RNA was submitted to RT-PCR for specific amplification of modRNA-G-CSF using ProtoScript® M-MuLV *Taq* RT-PCR kit (New England BioLabs, Ipswich, MA) according to the manufacturer instructions with hu-G-CSF-specific primers (see below). RT-PCR products were visualized by 1.2% agarose gel electrophoresis (FIG. 5B). Table 6 below shows which modRNAs were run on the agarose gel.

Table 6

Lane	Cell type	RT-PCR hu-G-CSF modRNA Target
1	Keratinocyte KG-1 Feeder	Vehicle
2	Keratinocyte KG-1 Feeder	Scramble RNA
3	Keratinocyte KG-1 Feeder	No Modification
4	Keratinocyte KG-1 Feeder	Chem 7
5	Keratinocyte KG-1 Feeder	Chem 6
6	Keratinocyte KG-1 Feeder	Chem 37
7	Keratinocyte Kasumi-1 Feeder	Vehicle
8	Keratinocyte Kasumi-1 Feeder	Scramble RNA
9	Keratinocyte Kasumi-1 Feeder	No Modification
10	Keratinocyte Kasumi-1 Feeder	Chem 7
11	Keratinocyte Kasumi-1 Feeder	Chem 6
12	Keratinocyte Kasumi-1 Feeder	Chem 37
13	Keratinocyte KG-1 Feeder	Chem 46
14	Keratinocyte KG-1 Feeder	Chem 48
15	Keratinocyte KG-1 Feeder	Chem 49
16	Keratinocyte KG-1 Feeder	Chem 53
17	Keratinocyte Kasumi-1 Feeder	Chem 46
18	Keratinocyte Kasumi-1 Feeder	Chem 48
19	Keratinocyte Kasumi-1 Feeder	Chem 49
20	Keratinocyte Kasumi-1 Feeder	Chem 53
21	Kasumi-1	Vehicle
22	KG-1	Vehicle
23	Kasumi-1	Vehicle
24	Kasumi-1	Scramble RNA
25	Kasumi-1	No Modification
26	Kasumi-1	Chem 7
27	Kasumi-1	Chem 6
28	Kasumi-1	Chem 37
29	Kasumi-1	Chem 46

30	Kasumi-1	Chem 48
31	Kasumi-1	Chem 49
32	Kasumi-1	Chem 53
33	KG-1	Vehicle
34	KG-1	Scramble RNA
35	KG-1	No Modification
36	KG-1	Chem 7
37	Empty	Empty
38	Empty	Empty
39	Empty	Empty
40	Empty	Empty
41	Empty	Empty
42	Empty	Empty
43	Empty	Empty
44	Empty	Empty

[0249] These data show that human keratinocyte cells containing hu-G-CSF modRNAs comprised of chemically distinct nucleotide analogs secreted hu-G-CSF protein and that the secreted hu-G-CSF was physiologically-active in inducing the proliferation of human myeloblast cells expressing the G-CSF receptor. These data also show the secreted hu-G-CSF protein was permeable across a semi-permeable membrane and acted on a different non-G-CSF-producing cell population. Additionally, these data show that hu-G-CSF modRNA-transfected into human keratinocyte cells in a co-culture environment was present in only the transfected keratinocyte cells and not the un-transfected myeloblast cells. Further, these data show that the modified nucleotide chemical composition of hu-G-CSF modRNA did not affect resultant protein activity.

Example 5. The Effect of modRNA on Cellular Viability

Cytotoxicity and Apoptosis:

[0250] This experiment demonstrates cellular viability, cytotoxicity and apoptosis for distinct modRNA-*in vitro* transfected Human Keratinocyte cells. Keratinocytes are grown in EpiLife medium with Human Keratinocyte Growth Supplement in the absence of hydrocortisone from Invitrogen at a confluence of >70%. Keratinocytes are reverse transfected with 0ng, 46.875ng, 93.75ng, 187.5ng, 375ng, 750ng, 1500ng, 3000ng, or 6000ng of modRNA complexed with RNAiMAX from Invitrogen. The modRNA:RNAiMAX complex is formed. Secreted huG-CSF concentration in the culture medium is measured at 0, 6, 12, 24, and 48 hours post-transfection for each concentration of each modRNA in triplicate. Secretion of Human Granulocyte-Colony Stimulating Factor (G-CSF) from transfected human keratinocytes is quantified using an ELISA kit from Invitrogen or R&D Systems following the manufacturers recommended instructions. Cellular viability, cytotoxicity and apoptosis is measured at 0, 12, 48, 96, and 192 hours post-transfection using the ApoToxGlo kit from Promega (Madison, WI) according to manufacturer instructions.

Example 6. Co-culture

[0251] The modified mRNA comprised of chemically-distinct modified nucleotides encoding human Granulocyte-Colony Stimulating Factor (G-CSF) may stimulate the cellular proliferation of a transfection incompetent cell in co-culture environment. The co-culture includes a highly transfectable cell type such as a human keratinocyte and a transfection incompetent cell type such as a white blood cell (WBC). The modified mRNA encoding G-CSF may be transfected into the highly transfectable cell allowing for the production and secretion of G-CSF protein into the extracellular environment where G-CSF acts in a paracrine-like manner to stimulate the white blood cell expressing the G-CSF receptor to proliferate. The expanded WBC population

may be used to treat immune-compromised patients or partially reconstitute the WBC population of an immunosuppressed patient and thus reduce the risk of opportunistic infections.

[0252] Another example, a highly transfectable cell such as a fibroblast may be transfected with certain growth factors to support and simulate the growth, maintenance, or differentiation of poorly transfectable embryonic stem cells or induced pluripotent stem cells.

Example 7. 5'-Guanosine Capping on Modified Nucleic Acids (modRNAs)

[0253] The cloning, gene synthesis and vector sequencing was performed by DNA2.0 Inc. (Menlo Park, CA). Sequence and insert sequence are set forth herein. The ORF was restriction digested using XbaI and used for cDNA synthesis using tailed-or tail-less-PCR. The tailed-PCR cDNA product was used as the template for the modified mRNA synthesis reaction using 25mM mixture each modified nucleotide (all modified nucleotides were custom synthesized or purchased from TriLink Biotech, San Diego, CA except pyrrolo-C triphosphate purchased from Glen Research, Sterling VA; unmodified nucleotides were purchased from Epicenter Biotechnologies, Madison, WI) and CellScript MegaScript™ (Epicenter Biotechnologies, Madison, WI) complete mRNA synthesis kit. The *in vitro* transcription reaction was run for 4 hours at 37°C. modRNAs incorporating adenosine analogs were poly (A) tailed using yeast Poly (A) Polymerase (Affymetrix, Santa Clara, CA). PCR reaction used HiFi PCR 2X Master Mix™ (Kapa Biosystems, Woburn, MA). modRNAs were post-transcriptionally capped using recombinant Vaccinia Virus Capping Enzyme (New England BioLabs, Ipswich, MA) and a recombinant 2'-O-methyltransferase (Epicenter Biotechnologies, Madison, WI) to generate the 5'-guanosine Cap1 structure. Cap 2 structure and Cap 3 structures may be generated using additional 2'-O-methyltransferases. The *in vitro* transcribed mRNA product was run on an agarose gel and visualized. modRNA was purified with Ambion/Applied Biosystems (Austin, TX) MEGAClear RNA™ purification kit. PCR used PureLink™ PCR purification kit (Invitrogen, Carlsbad, CA). The product was quantified on Nanodrop™ UV Absorbance (ThermoFisher, Waltham, MA). Quality, UV absorbance quality and visualization of the product was performed on an 1.2% agarose gel. The product was resuspended in TE buffer.

5' Capping Modified Nucleic Acid (mRNA) Structure:

[0254] 5'-modRNA capping may be completed concomitantly during the *in vitro*-transcription reaction using the following chemical RNA cap analogs to generate the 5'-guanosine cap structure according to manufacturer protocols: 3'-O-Me-m⁷G(5')ppp(5')G; G(5')ppp(5')A; G(5')ppp(5')G; m⁷G(5')ppp(5')A; m⁷G(5')ppp(5')G (New England BioLabs, Ipswich, MA). 5'-modRNA capping may be completed post-transcriptionally using a Vaccinia Virus Capping Enzyme to generate the "Cap 0" structure: m⁷G(5')ppp(5')G (New England BioLabs, Ipswich,

MA). Cap 1 structure may be generated using both Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m7G(5')ppp(5')G-2'-O-methyl. Cap 2 structure may be generated from the Cap 1 structure followed by the 2'-O-methylation of the 5'-antepenultimate nucleotide using a 2'-O methyl-transferase. Cap 3 structure may be generated from the Cap 2 structure followed by the 2'-O-methylation of the 5'-preantepenultimate nucleotide using a 2'-O methyl-transferase. Enzymes are preferably derived from a recombinant source.

Sequences:

[0255]

G-CSF cDNA:

agcttttgagccctcgtagagaagctaatacactcactatagggaaataagagagaaaagaagagtaagaagaaatataagag
ccaccatggccgggtcccgcgacccaaagcccatgaaacttatggccctgcagttgctgctttggcactcggccctctggacagtccaaga
agcgactcctctcgacctgcctcatcgttgcgcagtcattcctttgaaagtgtctggagcaggtgcgaaagattcagggcgatggagccg
cactccaagagaagctctgcgcgacatacaaaactttgccatcccgaggagctcgtactgctcgggcacagcttggggattccctgggctcc
tctctcgtcctgtccgtcgcaggtttgcagttggcaggggtgcctttccagctccactccggtttgtttgtatcagggactgctgcaagccc
ttgagggaaatctcgccagaattgggcccgcagctggacacgttgacgtcgcagctggcggttgcgaacaaccatctggcagcagatgg
aggaactggggatggcaccgcgctgcagcccacgcagggggcaatgccggcctttgcgtccgcgtttcagcgcagggcggtggagt
cctcgtagcagccacctcaatcttttgaagtctcgtaccgggtgctgagacatcttgcgcagccgtgaagcgctgccttctcgggggc
ttgccttctggccatgcccttctctccttgcacctgtacctcttggtctttgaataaagcctgagtaggaaggcgccgctcgagcatgca
tctagagggcccaattcgccctattcgaaagtcg (SEQ ID NO: 1)

G-CSF mRNA:

agcuuuuggaccucguacagaagcuauuacgacucacuauagggaauaagagagaaaagaagaguaagaagaaau
auaagagccaccauaggccggucccgacccaaagcccaugaaacuauaggccugcaguugcugcuuuggcacucggcccu
cuggacaguccaagaagcgacuccucucggaccugccuauugccgcagucuuuccuuuugaagugucugggagcagggug
cgaaagauucagggcgauaggcgcacuccaagagaagcucugcgcgacauacaaacuugccaucccgaggagcucguacu
gcucgggcacagcuuuggggaauuccugggcuccucucucguccuguccgucgcagggcuuugcaguuggcagggugccuuuc
ccaguccacuccggguuuguuuguaucaaggagcugcgaagcccuugagggaauucgccaagaauugggcccgcgcug
gacacguugcagcucgacgugcggaauucgcaacaaccaucuggcagcagauggaggaacuggggauggcaccgcgcugc
agcccacgcagggggcaaugccggccuuugcguccgcguuucagcgcagggcggguggaguccuagcguagcagccaccuuca
aucauuuuuggaagucucguaccgggugcugagacauuugcgcagccgugaagcgcugccuucugcggggcuugccuuc
ggccaugcccuucucuccuugcaccuguaaccuucugguucuuugaauaaagccugaguagggaaggcgccgcgcgagca
ugcaucuagagggcccaauucgcccuaucgaagucg (SEQ ID NO: 2)

G-CSF protein:

MAGPATQSPMKLMALQLLLWHSALWTVQEATPLGPASSLPQSFLKCLEQVRKI
QGDGAALQEKLVSSEATYKLCHPEELVLLGHS LGIPWAPLSSCPSQALQLAGCLS QLHS
GLFLYQGLLQALEGISPELGPTLDTLQLDVADFATTIWQQMEELGMAPALQPTQGAMPA
FASAFQRRAGGVLVASHLQSFLEVS YRVLRLHLAQP (SEQ ID NO: 3)

cDNA synthesis primers:

Forward Primer: 5'- TTG GAC CCT CGT ACA GAA GCT AAT ACG (SEQ ID NO: 4)

Reverse Primer for template Poly (A) tailing: 5'- T₍₁₂₀₎CT TCC TAC TCA GGC TTT ATT CAA AGA CCA (SEQ ID NO: 5)

Reverse Primer for post-transcriptional Poly (A) Polymerase tailing: 5'- CTT CCT ACT CAG GCT TTA TTC AAA GAC CA (SEQ ID NO: 6)

G-CSF modRNA RT-PCR primers:

Forward Primer: 5'- TGG CCG GTC CCG CGA CCC AA (SEQ ID NO: 7)

Reverse Primer: 5'- GCT TCA CGG CTG CGC AAG AT (SEQ ID NO: 8)

SEQUENCE LISTING

[0256]

<110> SCHRUM, JASON

<120> MODIFIED NUCLEOSIDES, NUCLOETIDES, AND NUCLEIC ACIDS, AND USES THEREOF

<130> M2107-7008WO

<140> PCT/US2011/054617

<141> 2011-10-03

<150> 61/404,413

<151> 2010-10-01

<160> 8

<170> PatentIn version 3.5

<210> 1

<211> 852

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 1

acgtttttga ccttctaga gaagtaata cactcacta taggaata agagagaaa

```

agccccgga cccccgaca gaagccaca cgaccaca cagggauca agagagauaa    60
gaagagtaag aagaaatata agagccacca tggccggtcc cgcgacccaa agcccatga    120
aacttatggc cctgcagttg ctgctttggc actcggccct ctggacagtc caagaagcga    180
ctcctctcgg acctgcctca tcgttgccgc agtcattcct ttggaagtgt ctggagcagg    240
tgcgaaagat tcagggcgat ggagccgcac tccaagagaa gctctgcgcg acatacaaac    300
tttgccatcc cgaggagctc gtactgctcg ggcacagctt ggggattccc tgggctcctc    360
tctcgtcctg tccgtcgcag gctttgcagt tggcagggtg cctttcccag ctccactccg    420
gtttgttctt gtatcagggg ctgctgcaag cccttgaggg aatctcgcca gaattgggcc    480
cgacgctgga cacgttcgag ctgcacgtgg cggatttcgc aacaaccatc tggcagcaga    540
tggaggaact ggggatggca cccgcgctgc agcccacgca gggggcaatg ccggcctttg    600
cgtccgcggt tcagcgcagg gcgggtggag tcctcgtagc gagccacctt caatcatttt    660
tggaaagtct gtaccgggtg ctgagacatc ttgcgcagcc gtgaagcgtc gccttctgcg    720
gggcttgctt tctggccatg cccttcttct ctcccttgca cctgtacctc ttggtctttg    780
aataaagcct gagtaggaag gcggccgctc gagcatgcat ctagagggcc caattcgccc    840
tattcgaagt cg                                         852

```

<210> 2

<211> 852

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 2

```

agcuuuugga cccucguaca gaagcuaaua cgacucacua uagggaaaua agagagaaaa    60
gaagaguaag aagaaauaua agagccacca uggccggucc cgcgacccaa agcccgaug    120
aacuuauagg ccugcaguug cugcuuuggc acucggcccu cuggacaguc caagaagcga    180
cuccucucgg accugccuca ucguugccgc agucauuccu uuugaagugu cuggagcagg    240
ugcgaaagau ucagggcgau ggagccgcac uccaagagaa gcucugcgcg acauacaaac    300
uuugccaucc cgaggagcuc guacugcucg ggcacagcuu ggggauuccc ugggcuccuc    360
ucucguccug uccgucgcag gcuuugcagu uggcagggug ccuuucccag cuccacuccg    420
guuuguucuu guaucagggg cugcugcaag ccuugagggg aaucucgcca gaauugggcc    480
cgacgcugga cacguugcag cucgacgug cggauuucgc aacaaccauc uggcagcaga    540
uggaggaacu ggggauggca cccgcgcugc agcccacgca gggggcaaug ccggccuuug    600
cguccgcguu ucagcgcagg gcggguggag uccucguagc gagccaccuu caaucuuuu    660
uggaagucuc guaccgggug cugagacauc uugcgcagcc gugaagcgcu gccuucugcg    720
gggcuugccu ucuggccaug cccuucuuuc cucccuugca ccuguaccuc uuggucuuug    780
aauaaagccu gaguaggaag gcggccgcuc gagcaugcau cuagagggcc caauucgccc    840
uauucgaagu cg                                         852

```

<210> 3

<211> 207

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 3

Met	Ala	Gly	Pro	Ala	Thr	Gln	Ser	Pro	Met	Lys	Leu	Met	Ala	Leu	Gln
1				5					10					15	

Leu	Leu	Leu	Trp	His	Ser	Ala	Leu	Trp	Thr	Val	Gln	Glu	Ala	Thr	Pro
			20					25					30		

Leu	Gly	Pro	Ala	Ser	Ser	Leu	Pro	Gln	Ser	Phe	Leu	Leu	Lys	Cys	Leu
		35					40					45			

Glu	Gln	Val	Arg	Lys	Ile	Gln	Gly	Asp	Gly	Ala	Ala	Leu	Gln	Glu	Lys
	50					55					60				

Leu	Val	Ser	Glu	Cys	Ala	Thr	Tyr	Lys	Leu	Cys	His	Pro	Glu	Glu	Leu
65					70					75					80

Val	Leu	Leu	Gly	His	Ser	Leu	Gly	Ile	Pro	Trp	Ala	Pro	Leu	Ser	Ser
				85					90					95	

Cys	Pro	Ser	Gln	Ala	Leu	Gln	Leu	Ala	Gly	Cys	Leu	Ser	Gln	Leu	His
			100					105					110		

Ser	Gly	Leu	Phe	Leu	Tyr	Gln	Gly	Leu	Leu	Gln	Ala	Leu	Glu	Gly	Ile
		115					120					125			

Ser	Pro	Glu	Leu	Gly	Pro	Thr	Leu	Asp	Thr	Leu	Gln	Leu	Asp	Val	Ala
	130					135					140				

Asp	Phe	Ala	Thr	Thr	Ile	Trp	Gln	Gln	Met	Glu	Glu	Leu	Gly	Met	Ala
145					150					155					160

Pro	Ala	Leu	Gln	Pro	Thr	Gln	Gly	Ala	Met	Pro	Ala	Phe	Ala	Ser	Ala
				165					170					175	

Phe	Gln	Arg	Arg	Ala	Gly	Gly	Val	Leu	Val	Ala	Ser	His	Leu	Gln	Ser
		180						185					190		

Phe	Leu	Glu	Val	Ser	Tyr	Arg	Val	Leu	Arg	His	Leu	Ala	Gln	Pro	
	195						200					205			

<210> 4

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 4

ttggaccctc gtacagaagc taatacg 27

<210> 5

<211> 149

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 5

tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt 60

tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt tttttttttt 120

cttcctactc aggctttatt caaagacca 149

<210> 6

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 6

cttcctactc aggctttatt caaagacca 29

<210> 7

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 7

tggccggtcc cgcgacccaa 20

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 8

gcttcacggc tgcgcaagat 20

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61404413 \[0001\]](#)
- [WO2007024708A2 \[0005\]](#)
- [US4886499A \[0205\]](#)
- [US5190521A \[0205\]](#)
- [US5328483A \[0205\]](#)
- [US5527288A \[0205\]](#)
- [US4270537A \[0205\]](#)
- [US5015235A \[0205\]](#)
- [US5141496A \[0205\]](#)
- [US5417662A \[0205\]](#)
- [WO9934850A \[0205\]](#)
- [US5480381A \[0205\]](#)
- [US5599302A \[0205\]](#)
- [US5334144A \[0205\]](#)
- [US5993412A \[0205\]](#)
- [US5649912A \[0205\]](#)
- [US5569189A \[0205\]](#)
- [US5704911A \[0205\]](#)
- [US5383851A \[0205\]](#)
- [US5893397A \[0205\]](#)
- [US5466220A \[0205\]](#)
- [US5339163A \[0205\]](#)
- [US5312335A \[0205\]](#)

- [US5503627A](#) [0205]
- [US5064413A](#) [0205]
- [US5520639A](#) [0205]
- [US4596556A](#) [0205]
- [US4790824A](#) [0205]
- [US4941880A](#) [0205]
- [US4940460A](#) [0205]
- [WO9737705A](#) [0205]
- [WO9713537A](#) [0205]
- [US2011054617W](#) [0256]
- [US61404413B](#) [0256]

Non-patent literature cited in the description

- **ROZENSKI, JCRAIN, PMCCLOSKEY, J**The RNA Modification Database: 1999 updateNucl Acids Res, 1999, vol. 27, 196-197 [0002]
- Remington's Pharmaceutical SciencesMack Publishing Company198500001418- [0079]
- Journal of Pharmaceutical Science, 1977, vol. 66, 2- [0079]
- **T. HIGUCHI**Pro-drugs as Novel Delivery SystemsA.C.S. Symposium Series, vol. 14, [0081]
- Bioreversible Carriers in Drug DesignAmerican Pharmaceutical Association and Pergamon Press19870000 [0081]
- **GREENE et al.**Protective Groups in Organic SynthesisWiley & Sons19910000 [0110] [0143]
- **OGATA et al.**Journal of Organic Chemistry, 2009, vol. 74, 2585-2588 [0113]
- **PURMAL et al.**Nucleic Acids Research, 1994, vol. 22, 172-78 [0113]
- **FUKUHARA et al.**Biochemistry, 1962, vol. 1, 4563-568 [0113]
- **XU et al.**Tetrahedron, 1992, vol. 48, 91729-1740 [0113]
- Computational Molecular BiologyOxford University Press19880000 [0132]
- Biocomputing: Informatics and Genome ProjectsAcademic Press19930000 [0132]
- Computer Analysis of Sequence Data, Part 1Humana Press19940000 [0132]
- **VON HEINJE, G.**Sequence Analysis in Molecular BiologyAcademic Press19870000 [0132]
- Sequence Analysis PrimerM. Stockton Press19910000 [0132]
- **CARILLO et al.**SIAM J. Applied Math., 1988, vol. 48, 1073- [0132]
- Oligonucleotide synthesis: a practical approachIRL Press19840000 [0140]
- Oligonucleotide synthesis: methods and applications**CLIFTON, N.J.**Methods in Molecular BiologyHumana Press20050000vol. 288, [0140]
- **ALBRECHT**Immunotherapy, 2010, vol. 2, 6795-8 [0162]
- **MUSUNURU K et al.**From noncoding variant to phenotype via SORT1 at the 1p13

cholesterol locusNature, 2010, vol. 466, 714-721 [\[0175\]](#)

- Remington's The Science and Practice of PharmacyLippincott, Williams & Wilkins20060000 [\[0185\]](#)
- Remington: The Science and Practice of PharmacyLippincott Williams & Wilkins20050000 [\[0213\]](#)

Patentkrav**1.** Fremgangsmåde til syntetisering af et mRNA, omfattende trinnene:

- a) tilvejebringelse af en komplementær deoxyribonukleinsyre (cDNA), der koder for et protein af interesse;
- b) udvælgelse af et nukleotid, der er kendt for at forstyrre binding af en hoveddrillebindingspartner med en nukleinsyre, hvor nukleotidet har nedsat bindingsaffinitet for hoveddrillebindingspartneren;
- c) at bringe det tilvejebragte cDNA og det valgte nukleotid i kontakt med en RNA-polymerase under betingelser, således at mRNA'et syntetiseres, hvor nukleotidet omfatter N1-methyl-pseudouridin, og hvor 100 % af nukleotiderne omfattende uracil i RNA'et erstattes med nukleotider, der omfatter N1-methyl-pseudouridin.

2. Fremgangsmåde ifølge krav 1, hvor hoveddrillebindingspartneren er valgt fra gruppen bestående af toll-lignende receptor (TLR) 3, TLR7, TLR8, retinsyre-inducerbart gen I (RIG-I), melanomdifferentieringsassocieret gen 5 (MDA5) og laboratorium for genetik og fysiologi 2 (LGP2).

3. mRNA, hvori 100 % af nukleotiderne omfattende uracil i mRNA'et erstattes med nukleotider omfattende N1-methyl-pseudouridin.

4. mRNA ifølge krav 3, hvor mRNA'et er mindst 300 nukleotider langt.

5. mRNA ifølge et hvilket som helst af kravene 3 eller 4 omfattende en polyA-hale.

6. mRNA ifølge et hvilket som helst af kravene 3-5, hvor hoveddrillebindingspartneren er RIG-I, MDA5 eller LGP2.

7. mRNA ifølge et hvilket som helst af kravene 3-5, hvor hoveddrillebindingspartneren er TLR3, TLR7 eller TLR8.

DRAWINGS




Fig. 1A




Fig. 1B




Fig. 2A




Fig. 2B




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 3D




Fig. 3E




Fig. 3F




Fig. 3G




Fig. 3H




Fig. 3I




Fig. 3J




Fig. 3K




Fig. 3L




Fig. 3M





Fig. 3N








Fig. 4C




Fig. 4D




Fig. 4E




Fig. 4F

Hu-G-CSF modRNA	Keratinocyte well hu-G-CSF (ng/ml)	KG-1 well hu-G-CSF (ng/ml)
Vehicle	0.0177	0.0189
Scramble	0.6217	0.0086
Chem 7	≥5	≥5
Chem 20	≥5	≥5
Chem 46	≥5	≥5
Chem 53	≥5	≥5

Hu-G-CSF modRNA	Keratinocyte well hu-G-CSF (ng/ml)	Kasumi-1 well hu-G-CSF (ng/ml)
Vehicle	0.0434	0.0256
Scramble	0.0655	0.3317
Chem 6	≥5	≥5
Chem 7	≥5	≥5
Chem 20	≥5	≥5
Chem 37	≥5	≥5
Chem 46	≥5	≥5
Chem 48	≥5	≥5
Chem 49	≥5	≥5
Chem 53	≥5	≥5

Fig. 5A






Fig. 5B



#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
21	UNMODIFIED NTP's	lab	8/15/2011 11:45:55 AM	1599.6	ng/ μ l	39.990	17.841	2.24	2.41	RNA	40.00




Fig. 6A

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
2	N1-methyl-A	lab	8/16/2011 11:35:49 AM	20.1	ng/ μ l	0.503	0.288	1.75	0.92	RNA	40.00




Fig. 6B

Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
N6-methyl-2-amino-Purine	lab	8/16/2011 11:38:29 AM	1112.4	ng/ μ l	27.811	17.607	1.58	1.40	RNA	40.00




Fig. 6C

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
2	2-thio-CTP	lab	8/15/2011 11:26:59 AM	259.2	ng/ μ l	6.479	4.719	1.37	1.30	RNA	40.00




Fig. 6D

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
4	pseudo-iso-CTP	lab	8/15/2011 11:29:10 AM	1332.5	ng/ μ l	33.311	13.674	2.44	1.79	RNA	40.00




Fig. 6E

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
6	5-iodo-UTP	lab	8/15/2011 11:31:06 AM	1179.4	ng/ μ l	29.486	16.726	1.76	1.10	RNA	40.00




Fig. 6F

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
7	N1 methyl-pseudoUTP	lab	8/15/2011 11:32:15 AM	1408.1	ng/ μ l	35.202	18.565	1.90	2.17	RNA	40.00




Fig. 6G

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
15	ITP	lab	8/15/2011 11:40:05 AM	225.1	ng/ μ l	5.628	5.781	0.97	1.05	RNA	40.00




Fig. 6H

#	Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
18	N1-methyl-GTP	lab	8/15/2011 11:43:10 AM	161.5	ng/ μ l	4.036	2.917	1.38	0.67	RNA	40.00




Fig. 6l

Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
pseudo-iso-CTP/N1-methyl-pseudo-UTP	lab	9/2/2011 3:48:37 PM	1268.3	ng/ μ l	31.707	17.208	1.84	1.64	RNA	40.00




Fig. 6J

Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
Pyr-CTP	lab	9/12/2011 2:03:00 PM	1103.7	ng/ μ l	27.592	11.101	2.49	1.14	RNA	40.00




Fig. 6K

Sample ID	User name	Date and Time	Nucleic Acid Conc.	Unit	A260	A280	260/280	260/230	Sample Type	Factor
5-Me-CTP/N1-Me-pseudo-UTP	lab	9/12/2011 3:47:12 PM	1094.3	ng/ μ l	27.358	16.135	1.70	2.11	RNA	40.00




Fig. 6L