



US 20160312199A1

(19) **United States**

(12) **Patent Application Publication**
Joung et al.

(10) **Pub. No.: US 2016/0312199 A1**

(43) **Pub. Date: Oct. 27, 2016**

(54) **ENGINEERED CRISPR-CAS9 NUCLEASES
WITH ALTERED PAM SPECIFICITY**

on May 22, 2015, provisional application No. 62/239,737, filed on Oct. 9, 2015, provisional application No. 62/258,402, filed on Nov. 20, 2015.

(71) Applicant: **The General Hospital Corporation,**
Boston, MA (US)

Publication Classification

(72) Inventors: **J. Keith Joung**, Winchester, MA (US);
Benjamin Kleinstiver, Medford, MA (US)

(51) **Int. Cl.**
C12N 9/22 (2006.01)
C12N 15/10 (2006.01)

(21) Appl. No.: **15/208,461**

(52) **U.S. Cl.**
CPC **C12N 9/22** (2013.01); **C12N 15/102**
(2013.01); **C12Y 301/00** (2013.01); **C12N**
2800/80 (2013.01)

(22) Filed: **Jul. 12, 2016**

Related U.S. Application Data

(63) Continuation of application No. 15/060,448, filed on Mar. 3, 2016.

(60) Provisional application No. 62/127,634, filed on Mar. 3, 2015, provisional application No. 62/165,517, filed

(57) **ABSTRACT**

Engineered CRISPR-Cas9 nucleases with altered and improved PAM specificities and their use in genomic engineering, epigenomic engineering, and genome targeting.




FIG. 1A




FIG. 1C




FIG. 1B




FIG. 1D




FIG. 1E









FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 3C

FIG. 3E




FIG. 3F




FIG. 3G




FIG. 4A




FIG. 4B




FIG. 4C





FIG. 4F




FIG. 4G

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F




FIG. 5G




FIG. 5H




FIG. 5I





FIG. 5J






FIG. 6




FIG. 7




FIG. 8A






FIG. 8B







FIG. 11



expression plasmid		selection plasmid		survival %	
Cas9	sgRNA	spacer	PAM	mean	S.E.M.
wild-type	spacer 1	spacer 1	NGG	68.70	6.90
		spacer 2	NGG	n.d. (<0.20%)	-
		none	none	n.d. (<0.20%)	-
wild-type	spacer 2	spacer 2	NGG	26.28	2.75
		spacer 1	NGG	n.d. (<0.11%)	-
wild-type	spacer 1	spacer 1	NAG	1.32	0.39
			NGA	n.d. (<0.07%)	-
			NAA	n.d. (<0.21%)	-
D10A/H840A	spacer 1	spacer 1	NGG	n.d. (<0.08%)	-
R1333Q	spacer 1	spacer 1	NAG	n.d. (<0.17%)	-
R1335Q			NGA	n.d. (<0.08%)	-
R1333Q/R1335Q			NAA	n.d. (<0.17%)	-
spacer 1 = GGGCAGCGGCAGCTTGCCCG				n.d. - not detectable at indicated %	
spacer 2 = GTGGCCCTCGAACTTCACCT					

FIG. 12A




FIG. 12B





FIG. 13A





FIG. 13B



FIGS. 13C-D





FIG. 14

th1	-	Mutations in 15/17 sequences	SEQ ID NO:
CGTAAGGAGCGCGAGGCGCGGCCGCGCGGCGGAGGCTGCAGGACTGAGCGAGCAGATCGTGTGAGG	<u>Wild-type</u>		107
CGTAAGGAGCGG	-----AGCAGATCGTGTGAGG	-41	108
CGTAAGGAGCGGAGG	-----CGAGCAGATCGTGTGAGG	-34	109
CGTAAGGAGCGCGAGGC	-----TGAGCGAGCAGATCGTGTGAGG	-29	110
CGTAAGGAGCGCGAGCGCGCGG	-----GAGCAGATCGTGTGAGG	-28	111
CGTAAGGAGCGCGAGCGCGCGCGCagatc	-----GAGCGAGCAGATCGTGTGAGG	-16 (-21, +5)	112
CGTAAGGAGCGCGAGCGCGCGCGCGCGT	-----AGCGAGCAGATCGTGTGAGG	-15 (-16, +1)	113
CGTAAGGAGCGCGAGCGCGCGCGCGCGCGG	-----GCGAGCAGATCGTGTGAGG	-15	114
CGTAAGGAGCGCGAGCGCGCGCGCGCGG	-----TGAGCGAGCAGATCGTGTGAGG	-14	115
CGTAAGGAGCGCGAGCGCGCGCGCGCGG	-----GCGAGCAGATCGTGTGAGG	-13 (-32, +19)	116
CGTAAGGAGCGCGAGCGCGCGCGCGCGGgag	-----GCGAGCAGATCGTGTGAGG	-13 (-37, +24)	117
CGTAAGGAGCGCGAGCGCGCGCGCGCGG	-----CTGAGCGAGCAGATCGTGTGAGG	-8	118
CGTAAGGAGCGCGAGCGCGCGCGCGCGCGG	-----ACTGAGCGAGCAGATCGTGTGAGG	-1 [2x]	119
CGTAAGGAGCGCGAGCGCGCGCGCGCGG	-----GagcagACTGAGCGAGCAGATCGTGTGAGG	+2 (-3, +5)	120
CGTAAGGAGCGCGAGCGCGCGCGCGCGG	-----GcagcagagcagatcGTGTGAGG	+2 (-17, +19)	121
th1/1	-	Mutations in 17/27 sequences	SEQ ID NO:
TGTCGGGAACCTCTCCAGGGATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	<u>Wild-type</u>		122
TGTCGGGAl	-----GTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-12 (-15, +3)	123
TGTCGGGAACCTCT	-----GTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-8 [x4]	124
TGTCGGGAACCTCTCC	-----TGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-5	125
TGTCGGGAACCTCTCCA	-----TGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-4 [x3]	126
TGTCGGGAACCTCTCC	-----GGGATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-1	127
TGTCGGGAACCTCTCCAl	-----GATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	-1 (-2, +1)	128
TGTCGGGAACCTCTCCAGGAGc	-----TTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	0 (-1, +1)	129
TGTCGGGAACCTCTgaaccG	-----GATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCAGATC	+1 (-4, +5)	130
TGTCGGGAACCTCTCgttacgag	-----GATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTCA	+4 (-4, +8)	131
TGTCGGGAACCTCTCCAlgttac	-----GGATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTC	+5 (-1, +6)	132
TGTCGGGAACCTCTCCAlgtgtGG	-----ATGTTACGGAGGCGCCCTCATCTCTCAAGTGTCTCTC	+5 (-1, +6)	133
TGTCGGGAACCTCTCCAlgtgagagg	-----aattattalaaataataaTTACGGAGGCGCCCT	+20 (-11, +31)	134
th	-	Mutations in 6/20 sequences	SEQ ID NO:
CATGGCGACCGGGGCGGAACCTACTGCTCTCCAACCGAGGCGGAGAAATCGGGGGCGGACG	<u>wild-type</u>		135
CATGGCGACCGGGGCGGGAACCTACTG	-----ACCAGAGCGGAGAAATCGGGGGCGGACG	-6	136
CATGGCGACCGGGGCGGGAACCTACTGCT	-----CAGAGCGGAGAAATCGGGGGCGGACG	-5	137
CATGGCGACCGGGGCGGGAACCTACTGCT	-----CCAGAGCGGAGAAATCGGGGGCGGACG	-4	138
CATGGCGACCGGGGCGGGAACCTACTGCTg	-----CCAGAGCGGAGAAATCGGGGGCGGACG	-4 (-5, +1)	139
CATGGCGACCGGGGCGGGAACCTACTGCTCTCagagg	-----CCAGAGCGGAGAAATCGGGGGCGG +3 (-3, +6)		140
CATGGCGACCGGGGCGGAGCTACTGCTCTClactgtctatc	-----ACCAGAGCGGAGAAATCGG +10 (-2, +12)		141

FIG. 15



FIGS. 16A-B







FIG. 17A







FIGS. 18A-C



FIGS. 19A-F



FIGS. 20A-B






FIG. 21D






FIG. 21E



FIGs. 21F






FIG. 23A




FIG. 23B




FIG. 23D




FIG. 23C




FIG. 23E




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28






FIG. 29








FIG. 31A




FIG. 31B




FIG. 32






FIG. 33



d




FIG. 34D




FIG. 34E




FIG. 35B




FIG. 35C

FIG. 35D




FIG. 36B




FIG. 36A




FIG. 37A




FIG. 37B

ENGINEERED CRISPR-CAS9 NUCLEASES WITH ALTERED PAM SPECIFICITY

CLAIM OF PRIORITY

[0001] This application is a continuation application of U.S. patent application Ser. No. 15/060,448, filed Mar. 3, 2016, which claims the benefit of U.S. Provisional Patent Application Ser. No. 62/127,634, filed on Mar. 3, 2015; 62/165,517, filed on May 22, 2015; 62/239,737, filed on Oct. 9, 2015; and 62/258,402, filed on Nov. 20, 2015. The entire contents of the foregoing are hereby incorporated by reference.

FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0002] This invention was made with Government support under Grant Nos. DP1 GM105378, NIH R01 GM107427, and R01 GM088040 awarded by the National Institutes of Health. The Government has certain rights in the invention.

TECHNICAL FIELD

[0003] The invention relates, at least in part, to engineered Clustered Regularly Interspaced Short Palindromic Repeats (CRISPRs)/CRISPR-associated protein 9 (Cas9) nucleases with altered and improved Protospacer Adjacent Motif (PAM) specificities and their use in genomic engineering, epigenomic engineering, and genome targeting.

SEQUENCE LISTING

[0004] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Jul. 12, 2016, is named 29539-0163003_SL.txt and is 925,050 bytes in size.

BACKGROUND

[0005] CRISPR-Cas9 nucleases enable efficient, customizable genome editing in a wide variety of organisms and cell types (Sander & Joung, Nat Biotechnol 32, 347-355 (2014); Hsu et al., Cell 157, 1262-1278 (2014); Doudna & Charpentier, Science 346, 1258096 (2014); Barrangou & May, Expert Opin Biol Ther 15, 311-314 (2015)). Target site recognition by Cas9 is directed by two short RNAs known as the crRNA and tracrRNA (Deltcheva et al., Nature 471, 602-607 (2011); Jinek et al., Science 337, 816-821 (2012)), which can be fused into a chimeric single guide RNA (sgRNA) (Jinek et al., Science 337, 816-821 (2012); Jinek et al., Elife 2, e00471 (2013); Mali et al., Science 339, 823-826 (2013); Cong et al., Science 339, 819-823 (2013)). The 5' end of the sgRNA (derived from the crRNA) can base pair with the target DNA site, thereby permitting straightforward re-programming of site-specific cleavage by the Cas9/sgRNA complex (Jinek et al., Science 337, 816-821 (2012)). However, Cas9 must also recognize a specific protospacer adjacent motif (PAM) that lies proximal to the DNA that base pairs with the sgRNA (Mojica et al., Microbiology 155, 733-740 (2009); Shah et al., RNA Biol 10, 891-899 (2013); Jinek et al., Science 337, 816-821 (2012); Sapranaukas et al., Nucleic Acids Res 39, 9275-9282 (2011); Horvath et al., J Bacteriol 190, 1401-1412 (2008)), a requirement that is needed to initiate sequence-specific recognition (Sternberg et al., Nature 507, 62-67 (2014)) but that can also constrain

the targeting range of these nucleases for genome editing. The broadly used *Streptococcus pyogenes* Cas9 (SpCas9) recognizes a short NGG PAM (Jinek et al., Science 337, 816-821 (2012); Jiang et al., Nat Biotechnol 31, 233-239 (2013)), which occurs once in every 8 bps of random DNA sequence. By contrast, other Cas9 orthologues characterized to date can recognize longer PAMs (Horvath et al., J Bacteriol 190, 1401-1412 (2008); Fonfara et al., Nucleic Acids Res 42, 2577-2590 (2014); Esvelt et al., Nat Methods 10, 1116-1121 (2013); Ran et al., Nature 520, 186-191 (2015); Zhang et al., Mol Cell 50, 488-503 (2013)). For example, *Staphylococcus aureus* Cas9 (SaCas9), one of several smaller Cas9 orthologues that are better suited for viral delivery (Horvath et al., J Bacteriol 190, 1401-1412 (2008); Ran et al., Nature 520, 186-191 (2015); Zhang et al., Mol Cell 50, 488-503 (2013)), recognizes a longer NNGRRT (SEQ ID NO:46) PAM that is expected to occur once in every 32 bps of random DNA. Broadening the targeting range of Cas9 orthologues is important for various applications including the modification of small genetic elements (e.g., transcription factor binding sites (Canver et al., Nature, 527(7577):192-7 (2015); Vierstra et al., Nat Methods. 12(10):927-30 (2015)) or performing allele-specific alterations by positioning sequence differences within the PAM (Courtney, D. G. et al. Gene Ther. 23(1):108-12 (2015)).

SUMMARY

[0006] As described herein, the commonly used *Streptococcus pyogenes* Cas9 (SpCas9) as well as the *Staphylococcus aureus* Cas9 (SaCas9) were engineered to recognize novel PAM sequences using structural information, bacterial selection-based directed evolution, and combinatorial design. These altered PAM specificity variants enable robust editing of endogenous gene sites in zebrafish and human cells that cannot be efficiently targeted by wild-type SpCas9 or SaCas9. In addition, we identified and characterized another SpCas9 variant that exhibits improved PAM specificity in human cells, possessing reduced activity on sites with non-canonical NAG and NGA PAMs. Furthermore, we found that two smaller-size Cas9 orthologues with completely different PAM specificities, *Streptococcus thermophilus* Cas9 (St1Cas9) and *Staphylococcus aureus* Cas9 (SaCas9), function efficiently in our bacterial selection system and in human cells, suggesting that our engineering strategies could be extended to Cas9s from other species. Our findings provide broadly useful SpCas9 and SaCas9 variants, referred to collectively herein as "variants" or "the variants".

[0007] In a first aspect, the invention provides isolated *Streptococcus pyogenes* Cas9 (SpCas9) proteins with mutations at one or more of the following positions: G1104, S1109, L1111, D1135, S1136, G1218, N1317, R1335, T1337, e.g., comprising a sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO:1. In some embodiments, the variant SpCas9 proteins comprise one or more of the following mutations: G1104K; S1109T; L1111H; D1135V; D1135E; D1135N; D1135Y; S1136N; G1218R; N1317K; R1335E; R1335Q; and T1337R. In some embodiments, the variant SpCas9 proteins comprise the following mutations: D1135; D1135V/R1335Q/T1337R (VQR variant); D1135E/R1335Q/T1337R (EQR variant); D1135V/G1218/R1335Q/T1337R (VRQR variant); D1135N/G1218R/R1335Q/T1337R (NRQR variant);

D1135Y/G1218R/R1335Q/T1337R (YRQR variant); G1104K/D1135V/G1218R/R1335Q/T1337R (KVRQR variant); S1109T/D1135V/G1218R/R1335Q/T1337R (TVRQR variant); L1111H/D1135V/G1218R/R1335Q/T1337R (HVRQR variant); D1135V/51136N/G1218R/R1335Q/T1337R (VNRQR variant); D1135V/G1218R/N1317K/R1335Q/T1337R (VRKQR variant); or D1135V/G1218R/R1335E/T1337R (VRER variant).

[0008] In some embodiments, the variant SpCas9 proteins comprise one or more mutations that decrease nuclease activity selected from the group consisting of mutations at D10, E762, D839, H983, or D986; and at H840 or N863.

[0009] In some embodiments, the mutations are: (i) D10A or D10N, and (ii) H840A, H840N, or H840Y.

[0010] Also provided herein are isolated *Staphylococcus aureus* Cas9 (SaCas9) proteins with mutations at one or more of the following positions: E782, N968, and/or R1015, e.g., comprising a sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO:2. Also provided herein are isolated *Staphylococcus aureus* Cas9 (SaCas9) proteins with mutations at one, two or more of the following positions: E735, E782, K929, N968, A1021, K1044 and/or R1015. In some embodiments, the variant SaCas9 proteins comprise one or more of the following mutations: R1015Q, R1015H, E782K, N968K, E735K, K929R, A1021T, K1044N. In some embodiments, the variant SaCas9 proteins comprise one or more mutations that decrease nuclease activity selected from the group consisting of mutations at D10, D556, H557, and/or N580. In some embodiments, the variant SaCas9 proteins comprise mutations at D10A, D556A, H557A, N580A, e.g., D10A/H557A and/or D10A/D556A/H557A/N580A.

[0011] SpCas9 variants described herein can include the amino acid sequence of SEQ ID NO:1, with mutations at one or more of the following positions: D1135, G1218, R1335, T1337. In some embodiments, the SpCas9 variants can include one or more of the following mutations: D1135V; D1135E; G1218R; R1335E; R1335Q; and T1337R. In some embodiments, the SpCas9 variants can include one of the following sets of mutations: D1135V/R1335Q/T1337R (VQR variant); D1135V/G1218R/R1335Q/T1337R (VRQR variant); D1135E/R1335Q/T1337R (EQR variant); or D1135V/G1218R/R1335E/T1337R (VRER variant).

[0012] SaCas9 variants described herein can include the amino acid sequence of SEQ ID NO:2, with mutations at one or more of the following positions: E735, E782, K929, N968, R1015, A1021, and/or K1044. In some embodiments, the SaCas9 variants can include one or more of the following mutations: R1015Q, R1015H, E782K, N968K, E735K, K929R, A1021T, K1044N. In some embodiments, the SaCas9 variants can include one of the following sets of mutations: E782K/N968K/R1015H (KKH variant); E782K/K929R/R1015H (KRH variant); or E782K/K929R/N968K/R1015H (KRRH variant).

[0013] Also provided herein are fusion protein comprising the isolated variant SaCas9 or SpCas9 proteins described herein fused to a heterologous functional domain, with an optional intervening linker, wherein the linker does not interfere with activity of the fusion protein. In some embodiments, the heterologous functional domain is a transcriptional activation domain. In some embodiments, the transcriptional activation domain is from VP64 or NF- κ B p65. In some embodiments, the heterologous functional domain is a transcriptional silencer or transcriptional repression

domain. In some embodiments, the transcriptional repression domain is a Krueppel-associated box (KRAB) domain, ERF repressor domain (ERD), or mSin3A interaction domain (SID). In some embodiments, the transcriptional silencer is Heterochromatin Protein 1 (HP1), e.g., HP1 α or HP1 β . In some embodiments, the heterologous functional domain is an enzyme that modifies the methylation state of DNA. In some embodiments, the enzyme that modifies the methylation state of DNA is a DNA methyltransferase (DNMT) or a TET protein. In some embodiments, the TET protein is TET1. In some embodiments, the heterologous functional domain is an enzyme that modifies a histone subunit. In some embodiments, the enzyme that modifies a histone subunit is a histone acetyltransferase (HAT), histone deacetylase (HDAC), histone methyltransferase (HMT), or histone demethylase. In some embodiments, the heterologous functional domain is a biological tether. In some embodiments, the biological tether is MS2, Csy4 or lambda N protein. In some embodiments, the heterologous functional domain is FokI.

[0014] Also provided herein are isolated nucleic acids encoding the variant SaCas9 or SpCas9 proteins described herein, as well as vectors comprising the isolated nucleic acids, optionally operably linked to one or more regulatory domains for expressing the variant SaCas9 or SpCas9 proteins described herein. Also provided herein are host cells, e.g., mammalian host cells, comprising the nucleic acids described herein, and optionally expressing the variant SaCas9 or SpCas9 proteins described herein.

[0015] Also provided herein are methods of altering the genome of a cell, by expressing in the cell an isolated variant SaCas9 or SpCas9 protein described herein, and a guide RNA having a region complementary to a selected portion of the genome of the cell.

[0016] Also provided herein are methods for altering, e.g., selectively altering, the genome of a cell by expressing in the cell the variant proteins, and a guide RNA having a region complementary to a selected portion of the genome of the cell.

[0017] Also provided are methods for altering, e.g., selectively altering, the genome of a cell by contacting the cell with a protein variant described herein, and a guide RNA having a region complementary to a selected portion of the genome of the cell.

[0018] In some embodiments, the isolated protein or fusion protein comprises one or more of a nuclear localization sequence, cell penetrating peptide sequence, and/or affinity tag.

[0019] In some embodiments of the methods described herein, the cell is a stem cell, e.g., an embryonic stem cell, mesenchymal stem cell, or induced pluripotent stem cell; is in a living animal; or is in an embryo, e.g., a mammalian, insect, or fish (e.g., zebrafish) embryo or embryonic cell.

[0020] Further, provided herein are methods, e.g., in vitro methods, for altering a double stranded DNA (dsDNA) molecule. The methods include contacting the dsDNA molecule with one or more of the variant proteins described herein, and a guide RNA having a region complementary to a selected portion of the dsDNA molecule.

[0021] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable

methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

[0022] Other features and advantages of the invention will be apparent from the following detailed description and FIGs, and from the claims.

DESCRIPTION OF DRAWINGS

[0023] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[0024] FIGS. 1A-J|Evolution and characterization of SpCas9 variants with altered PAM specificities. A, Rational mutation of the SpCas9 residues that make base-specific contacts to the PAM bases is insufficient to alter PAM specificity in the U2OS human cell-based Enhanced Green Fluorescent Protein (EGFP) disruption assay. Disruption frequencies were quantified by flow cytometry; the mean level of disruption observed with the background control is represented by the dashed red line for this and subsequent panels (C, G, H, and J); error bars represent s.e.m., n=3. B, Schematic of the two-plasmid positive selection assay used to alter the PAM specificity of SpCas9. Cleavage of a target site within the positive selection plasmid by a functional Cas9/sgRNA complex is necessary for survival when bacteria are plated on selective media (see also FIGS. 12A-B). C, Combinatorial assembly and testing of mutations obtained from the positive selection for SpCas9 variants that can cleave a target site containing an NGA PAM. SpCas9 Variants were paired with sgRNAs that target sites containing either an NGG or an NGA PAM and activity was assessed using the EGFP disruption assay. Error bars represent s.e.m., n=3. D, Schematic of the negative selection assay, in which cleavage of the selection plasmid results in cell death when bacteria are plated on selective media. This system was adapted to profile the PAM specificity of Cas9 by generating a library of plasmids that contain a randomized sequence adjacent to the 3' end of the protospacer (see also FIG. 13B). E, Scatterplot of the post-selection PAM depletion values (PPDVs) of wild-type SpCas9 with two randomized PAM libraries (each with a different protospacer). PAMs were grouped and plotted by their 2nd/3rd/4th positions. The red dashed line indicates the cutoff for statistically significant depletion (obtained from a dCas9 control experiment, see FIG. 13C), and the gray dashed line represents five-fold depletion (PPDV of 0.2). F, PPDV scatterplots for the VQR and EQR SpCas9 variants that recognize PAMs distinct from those recognized by wild-type SpCas9. G, EGFP disruption frequencies for wild-type, VQR, and EQR SpCas9 on sites with NGAN and NGNG PAMs. Error bars represent s.e.m., n=3. H, Combinatorial assembly and testing of mutations obtained from the positive selection for SpCas9 variants that can cleave a target site containing an NGCG PAM. sgRNAs that target sites containing either an NGGG or an NGCG PAM were assessed for Cas9 targeting using the EGFP disruption assay. Error bars represent s.e.m., n=3. I, PPDV scatterplot for the VRER variant. J, EGFP disruption frequencies for wild-type and

VRER SpCas9 on sites with NGCN and NGNG PAMs. Error bars represent s.e.m., n=3.

[0025] FIGS. 2A-F|SpCas9 variants with evolved PAM specificities robustly modify endogenous sites in zebrafish embryos and human cells. A, Quantification of mutagenesis frequencies in zebrafish embryos induced by wild-type or VQR SpCas9 on endogenous gene sites bearing NGAG PAMs. Mutation frequencies were determined using the T7E1 assay; error bars represent s.e.m., n=5 to 9 individual embryos. B, Mutation frequencies of the VQR variant quantified by T7E1 assay at 16 target sites in four endogenous human genes with sgRNAs targeted to sites containing NGAG, NGAT, and NGAA PAMs. Error bars represent s.e.m., n=3. C, Mutation frequencies of wild-type SpCas9 on endogenous human gene target sites with NGA PAMs. For ease of comparison, the mutation frequencies for the VQR variant using the same sgRNAs are re-presented here (same data shown in panel B). Error bars represent s.e.m., n=3; n.d., not detectable by T7E1. D, Mutation frequencies of wild-type, VRER, and VQR SpCas9 at nine target sites containing NGCG PAMs in three endogenous human genes quantified by T7E1 assay. sgRNA complementarity lengths of 19 and 20 nt were used; error bars represent s.e.m., n=3. E, Representation of the number sites in the human genome with 20 nt spacers targetable by wild-type, VQR, and VRER SpCas9. F, Number of off-target cleavage sites identified by GUIDE-seq for the VQR and VRER SpCas9 variants using sgRNAs from panels B and D.

[0026] FIGS. 3A-G|A D1135E mutation improves the PAM recognition and spacer specificity of SpCas9. A, PPDV scatterplots for wild-type and D1135E SpCas9 (left and right panels, respectively) for the two randomized PAM libraries. PAMs were grouped and plotted by their 2nd/3rd/4th positions. The data shown for wild-type SpCas9 is the same as the plot from FIG. 1D and is re-presented here for ease of comparison. The red dashed line indicates PAMs that are statistically significantly depleted (see FIG. 13C), and the gray dashed line indicates a five-fold depletion cutoff (PPDV of 0.2). B, EGFP disruption activities of wild-type and D1135E SpCas9 on sites that contain NGG, NAG, and NGA PAMs in human cells. Disruption frequencies were quantified by flow cytometry; the mean level of disruption observed with the background control is represented by the dashed red line for this panel and (D); error bars represent s.e.m., n=3; mean fold change in activity is shown. C, Mutagenesis frequencies detected by T7E1 for wild-type and D1135E SpCas9 at six endogenous sites in human cells. Error bars represent s.e.m., n=3; mean fold change in activity is shown. D, Titration of the amount of wild-type or D1135E SpCas9-encoding plasmid transfected for EGFP disruption experiments in human cells. The amount of sgRNA plasmid used for all of these experiments was fixed at 250 ng. Two sgRNAs targeting different EGFP sites were used; error bars represent s.e.m., n=3. E, Targeted deep-sequencing of on- and off-target sites for 3 sgRNAs using wild-type and D1135E SpCas9 (SEQ ID NOS 660-687, respectively, in order of appearance). The on-target site is shown at the top, with off-target sites listed below highlighting mismatches to the on-target. Fold decreases in activity with D1135E relative to wild-type SpCas9 at off-target sites greater than the change in activity at the on-target site are highlighted in green; control indel levels for each amplicon are reported. F, Summary of the targeted deep-sequencing data, plotted as the fold-decrease in activity at on- and

off-target sites using D1135E relative to the indel frequency observed with wild-type SpCas9. G, Summary of GUIDE-seq detected changes in specificity between wild-type and D1135E at off-target sites, plotted as the normalized fold-change in specificity using D1135E versus the read counts at that off-target site using wild-type SpCas9 (see also FIG. 18C). Estimated fold-gain in specificity at sites without read-counts for D1135E are not plotted (see FIG. 18C).

[0027] FIGS. 4A-G|Characterization of St1Cas9 and SaCas9 orthologues in bacteria and human cells. A, PPDV scatterplots for St1Cas9 using the two randomized PAM libraries. PAMs were grouped and plotted by their 3rd/4th/5th/6th positions. sgRNA complementarity lengths of 20 and 21 nucleotides were used to program St1Cas9 for both libraries (left and right panels, respectively). The red dashed line indicates PAMs that are statistically significantly depleted (see FIG. 13C), and the gray dashed line represents five-fold depletion (PPDV of 0.2); α , PAM previously predicted by a bioinformatic approach²⁷; β , PAMs previously identified under stringent experimental conditions²⁰; *, novel PAMs discovered in this study; γ , PAMs previously identified under moderate experimental conditions²⁰ (PAM 1 disclosed as SEQ ID NO: 3). B, PPDV scatterplots for SaCas9 using the two randomized PAM libraries. PAMs were grouped and plotted by their 3rd/4th/5th/6th positions. sgRNA complementarity lengths of 21 and 23 nucleotides were used to program SaCas9 for both libraries (left and right panels, respectively). PAMs identified for SaCas9 are shown, with PAMs 1-3 consistently depleted across all

combinations of spacer and spacer length used in these experiments (PAMs 1-3 disclosed as SEQ ID NOS 4, 6 and 5, respectively, in order of appearance). C, Survival percentages of St1Cas9 and SaCas9 in the bacterial positive selection when challenged with selection plasmids that harbor different target sites and PAMs indicated on the x-axis. Highly depleted PAMs from panels (A) and (B) for St1Cas9 and SaCas9 were used for the target sites in the positive selection plasmids (SEQ ID NOS 3, 3, 5, 5, 5 and 3, respectively, in order of appearance). D, E, EGFP disruption activities of St1Cas9 (panel D) or SaCas9 (panel E) on sites in EGFP that contain NNAGAA (SEQ ID NO:3) or NNGGGT (SEQ ID NO:4)/NNGAGT (SEQ ID NO:5) PAMs, respectively. Matched sgRNAs of different lengths for the same site are indicated; disruption frequencies were quantified by flow cytometry; the mean frequency of EGFP disruption obtained with a negative control is represented by the dashed red line; error bars represent s.e.m., n=3. F, G, Mutation frequencies of St1Cas9 (panel F) and SaCas9 (panel G) quantified by T7E1 assay at sites in four endogenous human genes that contain NNAGAA (SEQ ID NO:3) or NNGGGT (SEQ ID NO:4)/NNGAGT (SEQ ID NO:5)/NNGAAT (SEQ ID NO:6) PAMs, respectively. Error bars represent s.e.m., n=3; n.d., not detectable by T7E1.

FIGS. 5A-J. Sequences and Maps—plasmids used in this study

[0028]

FIG	Name	SEQ ID NO	Description
5A	BPK764	7	T7-humanSpCas9-NLS-3xFLAG-T7-BsaIcassette-SpgRNA T7 promoters: nts 1-17 and 4360-4376; human codon optimized <i>S. pyogenes</i> Cas9 88-4224; Nuclear Localization Signal (NLS) (CCCAAGAAGAAGAGGAAAGTC (SEQ ID NO: 650)) at nts 4198-4218, 3xFLAG tag 4225-4290, BsaI sites 4379-4384 and 4427-4432, gRNA (GTTTGTAGAGCTAGAAATAGCAAGTTAAATAAGGCTAGTCCG TTATCAACTTGAAAAAGTGGCACCAGTCGGTGC (SEQ ID NO: 651)) 4434-4509, T7 terminator 4252-4572 of SEQ ID NO:7
	MSP712	8	T7-humanSpdCas9 (D10A/H840A) -T7-BsaIcassette-SpgRNA T7 promoters at nts 1-17 and 4360-4376, human codon optimized <i>S. pyogenes</i> Cas9 88-4293, modified codons iat 115-117 and 2605-2607, bold and underlined, NLS (CCCAAGAAGAAGAGGAAAGTC (SEQ ID NO: 650)) at nts 4198-4218, 3xFLAG tag 4225-4290, BsaI sites 4379-4384 and 4427-4432, gRNA (GTTTGTAGAGCTAGAAATAGCAAGTTAAATAAGGCTAGT CCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGC (SEQ ID NO: 651)) at nts 4434-4509, T7 terminator 4252-4572 of SEQ ID NO: 8
5B	BPK2169	9	T7-humanSt1Cas9-NLS-T7-BspMICassette-St1gRNA T7 promoters at 1-17 and 3555-3571, human codon optimized <i>S. thermophilus</i> Cas9 at 88-3489, NLS at 3454 to 3486; BspMI sites at 3577-3582 and 3625-3630, gRNA at 3635-3763, T7 terminator 3778-3825 of SEQ ID NO: 9.
5C	BPK2101	10	T7-humanSaCas9-NLS-3xFLAG-T7-BsaIcassette-SagRNA T7 promoters at 1-17 and 3418-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3352, NLS at 3256-3276, 3xFLAG tag at 3283-3348, BsaI sites at 3437-3442 and 3485-3490, gRNA at 3492-3616, T7 terminator at 3627-2674 of SEQ ID NO: 10.
5D	p11-lacY-wtx1 ¹⁷	—	BAD-ccDB-Amp ^R -AraC-lacY(A177C)

-continued

FIG	Name	SEQ ID NO	Description
5E	JDS246	11	CMV-T7-humanSpCas9-NLS-3xFLAG ADDGENE ID: 43861 Human codon optimized <i>S. pyogenes</i> Cas9 1-4206, NLS at 4111-4131, 3xFLAG tag at 4138-4203 of SEQ ID NO: 11.
	MSP469	12	CMV-T7-humanSpCas9 (D1135V/R1335Q/T1337R)-NLS-3xFLAG (VQR variant) Human codon optimized <i>S. pyogenes</i> Cas9 1-4206, modified codons at 3403-3405, 4003-4005, and 4009-4011, NLS at 411-4131, 3xFLAG tag 4138-4203 of SEQ ID NO: 12.
	MSP680	13	CMV-T7-humanSpCas9 (D1135E/R1335Q/T1337R)-NLS-3xFLAG (EQR variant) Human codon optimized <i>S. pyogenes</i> Cas9 1-4206, modified codons at 3403-3405, 3652-3654, 4003-4005, and 4009-4011, NLS at 411-4131, 3xFLAG tag 4138-4203 of SEQ ID NO: 13.
	MSP1101	14	CMV-T7-humanSpCas9 (D1135V/G1218R/R1335E/T1337R)-NLS-3xFLAG (VRER variant) Human codon optimized <i>S. pyogenes</i> Cas9 1-4206, modified codons at 3403-3405, 4003-4005, and 4009-4011, NLS at 411-4131, 3xFLAG tag 4138-4203 of SEQ ID NO: 14
	MSP977	15	CMV-T7-humanSpCas9 (D1135E)-NLS-3xFLAG Human codon optimized <i>S. pyogenes</i> Cas9 1-4206, modified codons at 3403-3405, NLS at 411-4131, 3xFLAG tag 4138-4203 of SEQ ID NO: 15.
5F	MSP1393	16	CAG-humanStlCas9-NLS Human codon optimized <i>S. thermophilus</i> 1 Cas9 1-3402, NLS at 3367-3399 of SEQ ID NO: 16.
5G	BPK2139	17	CAG-humanSaCas9-NLS-3xFLAG Human codon optimized <i>S. aureus</i> Cas9 1-3195, NLS 3169-3189, 3xFLAG tag 3196-3261 of SEQ ID NO: 17.
5H	BPK1520	18	U6-BsmBICassette-SpgRNA U6 promoter at 1-318, BsmBI sites at 320-325 and 333-338, <i>S. pyogenes</i> gRNA 339-422, U6 terminator 416-422 of SEQ ID NO: 18.
5I	BPK2301	19	U6-BsmBICassette-StlgrRNA U6 promoter 1-318, BsmBI sites at 320-325 and 333-338, <i>S. thermophilus</i> 1 gRNA 340-471, U6 terminator 464-471 of SEQ ID NO: 19.
5J	VVT1	20	U6-BsmBICassette-SagRNA U6 promoter 1-318, BsmBI sites at 320-325 and 333-338, <i>S. aureus</i> gRNA 340-466, U6 terminator 459-466 of SEQ ID NO: 20.

[0029] FIG. 6|Alignment of Cas9 orthologues to predict PAM-interacting residues of SaCas9. The PAM-interacting domains of SpCas9, SaCas9, and 11 other Cas9 orthologues were aligned to identify PAM contacting residues in SaCas9, based on what is known for SpCas9. Top, Top, *S. Pyogenes*, amino acids 1229-1368 of SEQ ID NO:1, then SEQ ID NOs:29-40, respectively.

[0030] FIG. 7|Substitutions in SaCas9 assessed for activity against different PAMs in the bacterial screen. Based on the alignment from FIG. 6, single amino acid substitutions were tested in the bacterial positive selection to screen for effects on activity against a canonical NNGAGT (SEQ ID NO:5) and non-canonical NNAAGT (SEQ ID NO:41) and NNAGGT (SEQ ID NO:42) PAMs. Bacterial colonies on the selective media suggest that the SaCas9 variant has activity against a site containing the indicated PAM.

[0031] FIGS. 8A-B|Summary of amino acid substitutions that enable SaCas9 variants to target NNARRT (SEQ ID

NO:43) PAMs. Amino acid sequences of the PAM-interacting domain of 52 selected mutant SaCas9 clones that enabled survival in bacteria against sites containing an NNARRT (SEQ ID NO:43) PAM; the sequences presented are partial sequences of SEQ ID NOs:53-104 shown in Table 6 (SEQ ID NOS 967-1019, respectively, in order of appearance). Figure discloses "IIKKG" as SEQ ID NO: 966.

[0032] FIG. 9|Human cell activity of wild-type and engineered SaCas9 variants. Activity of wild-type, KKQ, and KKH SaCas9 was assessed in the human cell EGFP reporter assay against sites containing NNRRRT (SEQ ID NO:45) PAMs.

[0033] FIG. 10. SaCas9 activity against non-canonical PAMs in bacteria, and how directed mutations at R1015 impact activity against the same non-canonical PAMs (SEQ ID NOS 5, 511-514, 5, 511-512, 5, 513-514, 5 and 513-514, respectively, in order of appearance).

[0034] FIG. 11. Engineered variants can recognize PAMs of the form NNNRRT (SEQ ID NOS 5, 41-42, and 511-514, respectively, in order of appearance, on both the left and right side of the figure)

[0035] FIGS. 12A-B|Bacterial-based positive selection used to engineer altered PAM specificity variants of SpCas9. A, Expanded schematic of the positive selection from FIG. 1B (left panel), and validation that SpCas9 behaves as expected in the positive selection (right panel). Spacer 1, SEQ ID NO:105; Spacer 2, SEQ ID NO:106. B, Schematic of how the positive selection was adapted to select for SpCas9 variants that have altered PAM recognition specificities. A library of SpCas9 clones with randomized PAM-interacting (PI) domains (residues 1097-1368) is challenged by a selection plasmid that harbors an altered PAM (target site disclosed as SEQ ID NO: 688). SpCas9 variants that survive the selection by cleaving the positive selection plasmid are sequenced to determine the mutations that enable altered PAM specificity.

[0036] FIGS. 13A-D|Bacterial cell-based site-depletion assay for profiling the global PAM specificities of Cas9 nucleases. A, Expanded schematic illustrating the negative selection from FIG. 1D (left panel), and validation that wild-type SpCas9 behaves as expected in a screen of sites with functional (NGG) and non-functional (NGA) PAMs (right panel). B, Schematic of how the negative selection was used as a site-depletion assay to screen for functional PAMs by constructing negative selection plasmid libraries containing 6 randomized base pairs in place of the PAM. Selection plasmids that contain PAMs cleaved by a Cas9/sgRNA of interest are depleted while PAMs that are not cleaved (or poorly cleaved) are retained. The frequencies of the PAMs following selection are compared to their pre-selection frequencies in the starting libraries to calculate the post-selection PAM depletion value (PPDV). Spacer 1, SEQ ID NO: 689; Spacer 2, SEQ ID NO: 690. C, D, A cutoff for statistically significant PPDVs was established by plotting the PPDV of PAMs for catalytically inactive SpCas9 (dCas9) (grouped and plotted by their 2nd/3rd/4th positions) for the two randomized PAM libraries (C). A threshold of 3.36 standard deviations from the mean PPDV for the two libraries was calculated (red lines in (D)), establishing that any PPDV deviation below 0.85 is statistically significant compared to dCas9 treatment (red dashed line in (C)). The gray dashed line in (C) indicates a five-fold depletion in the assay (PPDV of 0.2).

[0037] FIG. 14|Concordance between the site-depletion assay and EGFP disruption activity. Data points represent the average EGFP disruption of the two NGAN and NGNG PAM sites for the VQR and EQR SpCas9 variants (FIG. 1G) plotted against the mean PPDV observed for library 1 and 2 (FIG. 1F) for the corresponding PAM. The red dashed line indicates PAMs that are statistically significantly depleted (PPDV of 0.85, see FIG. 13C), and the gray dashed line represents five-fold depletion (PPDV of 0.2). Mean values are plotted with the 95% confidence interval.

[0038] FIG. 15|Insertion or deletion mutations induced by the VQR SpCas9 variant at endogenous zebrafish sites containing NGAG PAMs. For each target locus, the wild-type sequence is shown at the top with the protospacer highlighted in yellow (highlighted in green if present on the complementary strand) and the PAM is marked as red underlined text. Deletions are shown as red dashes highlighted in gray and insertions as lower case letters high-

lighted in blue. The net change in length caused by each indel mutation is shown on the right (+, insertion; -, deletion). Note that some alterations have both insertions and deletions of sequence and in these instances the alterations are enumerated in parentheses. The number of times each mutant allele was recovered (if more than once) is shown in brackets.

[0039] FIGS. 16A-B|Endogenous genes targeted by wild-type and evolved variants of SpCas9. A, Sequences targeted by wild-type, VQR, and VRER SpCas9 are shown in blue, red, and green, respectively. Sequences of sgRNAs and primers used to amplify these loci for T7E1 are provided in Tables 1 and 2, below. B, Mean mutagenesis frequencies detected by T7E1 for wild-type SpCas9 at eight target sites bearing NGG PAMs in the four different endogenous human genes (corresponding to the annotations in the top panel). Error bars represent s.e.m., n=3.

[0040] FIGS. 17A-B|Specificity profiles of the VQR and VRER SpCas9 variants determined using GUIDE-seq. The intended on-target site is marked with a black square, and mismatched positions within off-target sites are highlighted. A, The specificity of the VQR variant was assessed in human cells by targeting endogenous sites containing NGA PAMs: EMX1 site 4 (SEQ ID NO:142 and variants disclosed as SEQ ID NOS 691-692, respectively, in order of appearance), FANCF site 1 (SEQ ID NO:143 and variants disclosed as SEQ ID NOS 693-699, respectively, in order of appearance), FANCF site 3 (SEQ ID NO:144 and variants disclosed as SEQ ID NOS 700-702, respectively, in order of appearance), FANCF site 4 (SEQ ID NO:145 and variant disclosed as SEQ ID NO: 703), RUNX1 site 1 (SEQ ID NO:146 and variants disclosed as SEQ ID NOS 704-714, respectively, in order of appearance), RUNX1 site 3 (SEQ ID NO:147 and variants disclosed as SEQ ID NOS 715-771, respectively, in order of appearance), VEGFA site 1 (SEQ ID NO:148 and variant disclosed as SEQ ID NO: 772), and ZSCAN2 (SEQ ID NO:149 and variants disclosed as SEQ ID NOS 773-794, respectively, in order of appearance). B, The specificity of the VRER variant was assessed in human cells by targeting endogenous sites containing NGCG PAMs: FANCF site 3 (SEQ ID NO:150 and variants disclosed as SEQ ID NOS 795-796, respectively, in order of appearance), FANCF site 4 (SEQ ID NO:151 and variants disclosed as SEQ ID NOS 797-798, respectively, in order of appearance), RUNX1 site 1 (SEQ ID NO:152 and variant disclosed as SEQ ID NO: 799), VEGFA site 1 (SEQ ID NO:153 and variants disclosed as SEQ ID NOS 800-804, respectively, in order of appearance), and VEGFA site 2 (SEQ ID NO:154 and variant disclosed as SEQ ID NO: 805).

[0041] FIGS. 18A-C|Activity differences between D1135E and wild-type SpCas9 at off-target sites detected by GUIDE-seq. A, Mean frequency of oligo tag integration at the on-target sites, estimated by restriction fragment length polymorphism analysis. Error bars represent s.e.m., n=4. B, Mean mutagenesis frequencies at the on-target sites detected by T7E1. Error bars represent s.e.m., n=4. C, GUIDE-seq read-count differences between wild-type SpCas9 and D1135E at 3 endogenous human cell sites (EMX1 site 3 (SEQ ID NO:155 and variants disclosed as SEQ ID NOS 806-812, respectively, in order of appearance); ZNF629 site (SEQ ID NO:156 and variants disclosed as SEQ ID NOS 813-826, respectively, in order of appearance), VEGFA site 3 (SEQ ID NO:157 and variants disclosed as SEQ ID NOS 827-873, respectively, in order of appearance). The on-target

site is shown at the top and off-target sites are listed below with mismatches highlighted. In the table, a ratio of off-target activity to on-target activity is compared between wild-type and D1135E to calculate the normalized fold-changes in specificity (with gains in specificity highlighted in green). For sites without detectable GUIDE-seq reads, a value of 1 has been assigned to calculate an estimated change in specificity (indicated in orange). Off-target sites analyzed by deep-sequencing in FIG. 3E are numbered to the left of the EMX1 site 3 and VEGFA site 3 off-target sites.

[0042] FIGS. 19A-F | Additional PAMs for St1Cas9 and SaCas9 and activities based on spacer lengths in human cells. A, PPDV scatterplots for St1Cas9 comparing the sgRNA complementarity lengths of 20 and 21 nucleotides obtained with a randomized PAM library for spacer 1 (top panel) or spacer 2 (bottom panel). PAMs were grouped and plotted by their 3rd/4th/5th/6th positions ("nnAGAA" disclosed as SEQ ID NO: 3). The red dashed line indicates PAMs that are statistically significantly depleted (see FIG. 13C) and the gray dashed line represents five-fold depletion (PPDV of 0.2). B, Table of PAMs with PPDVs of less than 0.2 for St1Cas9 under each of the four conditions tested. PAM numbering shown on the left is the same as in FIG. 4A (PAM 1 disclosed as SEQ ID NO: 3). C, PPDV scatterplots for SaCas9 comparing the sgRNA complementarity lengths of 21 and 23 nucleotides obtained with a randomized PAM library for spacer 1 (top panel) or spacer 2 (bottom panel). PAMs were grouped and plotted by their 3rd/4th/5th/6th positions. The red and gray dashed lines are the same as in (A). D, Table of PAMs with PPDVs of less than 0.2 for SaCas9 under each of the four conditions tested. PAM numbering is the same as in FIG. 4B ("nnGGGT" disclosed as SEQ ID NO: 4, "nnGAAT" disclosed as SEQ ID NO: 6, "nnGAGT" disclosed as SEQ ID NO: 5, "nnAAGT" disclosed as SEQ ID NO 41, "nnAGGT" disclosed as SEQ ID NO: 42 and "nnCAGT" disclosed as SEQ ID NO: 511). E, F, Human cell activity of St1Cas9 and SaCas9 across various spacer lengths via EGFP disruption (panel E, data from FIGS. 4D, 4E) and endogenous gene mutagenesis detected by T7E1 (panel F, data from FIGS. 4F, 4G). Activity for all replicates shown (n=3 or 4); bars illustrate mean and 95% confidence interval; number of sites per spacer length indicated.

[0043] FIGS. 20A-B | Structural and functional roles of D1135, G1218, and T1337 in PAM recognition by SpCas9. A, Structural representations of the six residues implicated in PAM recognition. The left panel illustrates the proximity of D1135 to 51136, a residue that makes a water-mediated, minor groove contact to the 3rd base position of the PAM15. The right panel illustrates the proximity of G1218, E1219, and T1337 to R1335, a residue that makes a direct, base-specific major groove contact to the 3rd base position of the PAM15. Angstrom distances indicated by yellow dashed lines; non-target strand guanine bases dG2 and dG3 of the PAM are shown in blue; other DNA bases shown in orange; water molecules shown in red; images generated using PyMOL from PDB:4UN3. B, Mutational analysis of six residues in SpCas9 that are implicated in PAM recognition. Clones containing one of three types of mutations at each position were tested for EGFP disruption with two sgRNAs targeted to sites harboring NGG PAMs. For each position, we created an alanine substitution and two non-conservative mutations. S1136 and R1335 were previously reported to mediate contacts to the 3rd guanine of the PAM15, and

D1135, G1218, E1219, and T1337 are reported in this study. EGFP disruption activities are quantified by flow cytometry; background control represented by the dashed red line; error bars represent s.e.m., n=3.

[0044] FIGS. 21A-F | Selection and assembly of SaCas9 variants with altered PAM specificities (A) Phylogenetic tree of Cas9 orthologues with SpCas9 and SaCas9 highlighted. (B) Activity of SaCas9 variants with single amino acid substitutions assessed in the bacterial positive selection assay (see also FIG. 31B). Error bars represent s.e.m., n=3; NS=no survival. (C) Human cell activity of wild-type and R1015H SaCas9 (SEQ ID NOS 41, 511, 5 and 513, respectively, in order of appearance). EGFP disruption activity quantified by flow cytometry; error bars represent s.e.m., n=3, mean level of background EGFP loss represented by dashed red line (for this and panel E). (D) Total number of substitutions observed at each amino acid position when selecting for SaCas9 variants with altered PAM specificities (SEQ ID NOS 43 and 47, respectively, in order of appearance). Starter mutations at R1015 are not counted. (E) Human cell EGFP disruption activity of variants containing mutations observed when selecting for altered PAM specificities (SEQ ID NOS 41, 5, 42, 4 and 511, 513, 512 and 514, respectively, in order of appearance). (F) Mean post-selection PAM depletion value (PPDV) scatterplot of wild-type SaCas9 versus the KKH variant (n=2, see also FIG. 34C) (SEQ ID NOS 43, 47, 46, 48, 514, 513, 6, 4, 42, 512, 5, 41 and 511, respectively, in order of appearance). Two libraries with different protospacers and 8 randomized basepairs in place of the PAM were used to determine which PAMs are targetable by each Cas9. Statistically significant depletion indicated by the red dashed line (relative to a dCas9 control, see FIGS. 34A and 34B), and 5-fold depletion by the grey dashed line.

[0045] FIGS. 22A-F. Activity of the SaCas9 KKH variant targeted to endogenous sites in human cells (A) Mutagenesis frequencies across 55 different sites bearing NNNRRT PAMs (SEQ ID NOS 43, 47, 46 and 48, respectively, in order of appearance) induced by KKH SaCas9, determined by T7E1 assay. Error bars represent s.e.m., n=3, ND, not detectable by T7E1 assay. (B) KKH variant preference for the third position of the PAM. Mean activities from data in panel A are shown for this and panels B and C. (C) KKH variant preference for the fourth and fifth positions of the PAM. (D) Spacer length preference of the KKH SaCas9 variant. (E) Comparison of the human cell EGFP disruption activity of wild-type and KKH SaCas9 targeted to various sites containing NNNRRT PAMs (SEQ ID NOS 43, 47, 46 and 48, respectively, in order of appearance). EGFP disruption quantified by flow cytometry; error bars represent s.e.m., n=3, mean level of background EGFP loss represented by dashed red line. (F) Mutagenesis frequencies of wild-type SaCas9 against one site for each of the 16 possible NNNRRT sites from panel A (sites with the highest KKH activity were selected). Error bars represent s.e.m., n=3, ND, not detectable by T7E1 assay.

[0046] FIGS. 23A-E | Genome-wide specificity profiles of wild-type and KKH SaCas9 (A) and (B) Direct comparison of wild-type and KKH SaCas9 targeted to sites containing NNGRRT (SEQ ID NO:46) PAMs, represented by total number of off-targets (panel A) and mismatches observed at each off-target site (panel B) at EMX site 6 (SEQ ID NO:158 and variants disclosed as SEQ ID NOS 874-876, respectively, in order of appearance) and VEGF site 8 (SEQ ID

NO:159 and variants disclosed as SEQ ID NOS 891-925, respectively, in order of appearance). FANCF site 10 wild-type and variants disclosed as SEQ ID NOS 877 and 878; FANCF site 13 wildtype and variants disclosed as SEQ ID NOS 879-883; RUNX1 site 13 wildtype and variants disclosed as SEQ ID NOS 884-887; and RUNX1 site 14 variants disclosed as SEQ ID NOS 888-890, all respectively, in order of appearance). For panels B and E, GUIDE-seq read counts at each site are indicated; on-target sequences are marked with a black box; mismatched positions within off-target sites are highlighted; sequences have been corrected for cell-type specific SNPs; sites with potential sgRNA or DNA bulge nucleotides are indicated by a small red-bordered base or a dash, respectively. (C) Venn diagram highlighting the differences in off-target site cleavage by wild-type and KKH SaCas9 at VEGFA site 8. (D) and (E) Specificity profile of the KKH variant targeted to sites containing NNHRRT (SEQ ID NO:44) PAMs, EMX site 1 (SEQ ID NO:160 and variants disclosed as SEQ ID NOS 926-941, respectively, in order of appearance), EMX site 4 (SEQ ID NO:161 and variants disclosed as SEQ ID NOS 942-951, respectively, in order of appearance), EMX site 10 (SEQ ID NO:162 and variants disclosed as SEQ ID NOS 952-957, respectively, in order of appearance), FANCF site 9 (SEQ ID NO:163 and variants disclosed as SEQ ID NOS 958-962, respectively, in order of appearance), and FANCF site 16 (SEQ ID NO:164 and variant disclosed as SEQ ID NO: 963), represented by total number of off-targets (panel D) and mismatches observed at each off-target site (panel E).

[0047] FIG. 24: Activity of VQR-derivative clones in the bacterial 2-plasmid screen. Testing of 24 different VQR derivative variants against sites in bacteria that contain NGAN PAMs. Survival on the selective plate, relative to the non-selective plate, is indicative of activity against the indicated PAM.

[0048] FIG. 25: Human cell EGFP disruption activity of SpCas9-VQR derivatives. EGFP disruption activity of the SpCas9 variants is a measure of activity against sites that contain the indicated PAM.

[0049] FIG. 26: Human cell EGFP disruption activity of SpCas9-VQR and -VRQR variants. EGFP disruption activity of the SpCas9 variants is a measure of activity against sites that contain the indicated PAM.

[0050] FIG. 27: Activity of SpCas9-VRQR derivative variants in the bacterial 2-plasmid screen. Testing of 12 different VQR derivative variants against sites in bacteria that contain NGAN PAMs, compared to the VQR and VRQR variants. Survival on the selective plate, relative to the non-selective plate, is indicative of activity against the indicated PAM.

[0051] FIG. 28: Human cell EGFP disruption activity of SpCas9-VRQR variants. EGFP disruption activity of the SpCas9 variants is a measure of activity against sites that contain the indicated PAM.

[0052] FIG. 29 Protein domain alignment of Cas9 orthologues (from FIG. 21A). The domain structure of SpCas9 is shown at the top (based on PDB:4UN3; Anders et al., 2014); the PAM contacting residues of SpCas9 are highlighted; the region of SaCas9 mutagenized to select for altered PAM specificity variants is shown.

[0053] FIGS. 30A-B Primary sequence alignment of Cas9 orthologues for identification of PAM-interacting residues; SEQ ID NOS:165-176, respectively. SpCas9 residues previously identified (Anders et al., 2014; Examples 1-2) to be important for contacting the PAM are highlighted in blue,

residues capable of modulating SaCas9 PAM specificity (identified in this study) are highlighted in orange, and positively charged residues adjacent to R1015 are highlighted in yellow. The structurally predicted PAM-interacting domain of SpCas9 is highlighted with a blue dashed line (based on PDB:4UN3; Anders et al., 2014), and the conservative estimate of the SaCas9 PAM-interacting domain used as a boundary for PCR mutagenesis is indicated with an orange dashed line.

[0054] FIGS. 31A-B Schematic of the bacterial positive selection assay (A) The selection plasmids can be modified to screen for Cas9 variants that are able to recognize alternative PAM sequences (SEQ ID NO: 1020). (B) Schematic of the positive selection plasmids (left panel) and expected outcomes (right panel) when screening functional or non-functional Cas9/sgRNA pairs in the positive selection (SEQ ID NO: 1021).

[0055] FIG. 32 Addition of the K929R mutation to the KNH and KKH variants (SEQ ID NOS 41-42, 511-512, 5, 4 and 513-514, respectively, in order of appearance). EGFP disruption activity quantified by flow cytometry; error bars represent s.e.m, n=3, mean level of background EGFP loss represented by the dashed red line.

[0056] FIG. 33 Schematic of the bacterial site-depletion assay. Site-depletion plasmids with 8 randomized nucleotides in place of the PAM that are refractory to cleavage by wild-type or KKH SaCas9 are sequenced. Library 1 Spacer sequence, SEQ ID NO:964; library 2 spacer sequence, SEQ ID NO: 965. Targetable PAMs are inferred by their depletion relative to the input library, calculated as the post-selection PAM depletion value (PPDV).

[0057] FIGS. 34A-E Site-depletion assay results for wild-type and KKH SaCas9 (A) PPDV values for dCas9 control experiments on both libraries. The red dashed line indicates statistical significance (PPDV=0.794, see panel B); grey dashed line indicates 5-fold depletion; PPDVs for a window comprising the 3rd/4th/5th/6th positions of the PAM are plotted (for this and panel C). (B) Statistically significant post-selection PAM depletion values (PPDVs) were determined from the dCas9 control experiments in panel A. Statistical significance was determined by setting the threshold at 3.36 times the standard deviation. (C) Comparison of the PPDVs for wild-type and KKH SaCas9 for each of the two libraries containing 8 randomized nucleotides in place of the PAM. Library 1 discloses SEQ ID NOS 514, 513, 6, 4, 42, 512, 5, 511 and 41, and Library 2 discloses SEQ ID NOS 513-514, 5-6, 4, 42, 41 and 511-512, respectively, in order of appearance. (D) and (E) PAMs and corresponding PPDV values for all PAMs depleted greater than 5-fold for wild-type and KKH SaCas9, respectively. Sequence motifs are shown for PAMs in two categories: 1) greater than 10 fold or 2) 5- to 10-fold depleted.

[0058] FIGS. 35A-D Additional characteristics of endogenous sites targeted by KKH SaCas9 (A) Activity for each of the 55 endogenous site sgRNAs, binned based on the 16 possible NRR motifs of an NNNRRT PAM. Mean activities from FIG. 2A are shown for this and panels B and C. (B) and (C) Relationship between endogenous gene disruption activity and GC content of the spacer and PAM, respectively. (D) Sequence logos for the spacer and PAM of target sites binned based on activity. Sites were grouped based on mean mutation frequency (from FIG. 2A) into low (0-10%, 17 sites), medium (10-30%, 17 sites), or high (>30%, 21 sites) activity.

[0059] FIGS. 36A-B On-target tag integration and mutagenesis frequencies for GUIDE-seq experiments (A) Restriction fragment length polymorphism (RFLP) analysis to determine the mean GUIDE-seq tag integration frequencies. Error bars represent s.e.m., n=3 (for this and panel B). (B) Mean mutagenesis detected by T7E1 assay.

[0060] FIGS. 37A-B A truncated repeat: anti-repeat sgRNA outperforms the full length sgRNA, similar to previous results (Ran et al., 2015) (A) Human cell EGFP disruption activity for wild-type SaCas9 against 4 sites that contain NNGRRT (SEQ ID NO:46) PAMs (SEQ ID NOS 4, 5 and 4, respectively, in order of appearance). EGFP disruption activity quantified by flow cytometry; error bars represent s.e.m, n=3, mean level of background EGFP loss represented by dashed red line (for this and panel B). (B) Human cell EGFP disruption activity for KKH SaCas9 against 8 sites that contain NNNRRT PAMs (SEQ ID NOS 41-42, 511-512, 5, 4 and 513-514, respectively, in order of appearance).

DETAILED DESCRIPTION

[0061] Although CRISPR-Cas9 nucleases are widely used for genome editing¹⁻⁴, the range of sequences that Cas9 can cleave is constrained by the need for a specific protospacer adjacent motif (PAM) in the target site^{5,6}. For example, SpCas9, the most robust and widely used Cas9 to date, primarily recognizes NGG PAMs. As a result, it can often be difficult to target double-stranded breaks (DSBs) with the precision that is necessary for various genome editing applications. In addition, imperfect PAM recognition by Cas9 can lead to the creation of unwanted off-target mutations^{7,8}. The ability to evolve Cas9 derivatives with purposefully altered or improved PAM specificities would address these limitations but, to the present inventors' knowledge, no such Cas9 variants have been described.

[0062] A potential strategy for improving the targeting range of orthogonal Cas9s that recognize extended PAMs is to alter their PAM recognition specificities. As described herein, PAM recognition specificity of SpCas9 can be altered using a combination of structure-guided design and directed evolution performed with a bacterial cell-based selection system; see Examples 1 and 2. Also described herein are variants that have been evolved to have relaxed or partially relaxed specificities for certain positions within the PAM; see Example 3. These variants expand the utility of Cas9 orthologues that specify longer PAM sequences.

[0063] Engineered Cas9 Variants with Altered PAM Specificity

[0064] The SpCas9 variants engineered in this study greatly increase the sites accessible by wild-type SpCas9, further enhancing the opportunities to use the CRISPR-Cas9 platform to practice efficient HDR, to target NHEJ-mediated indels to small genetic elements, and to exploit the requirement for a PAM to distinguish between two different alleles in the same cell. The altered PAM specificity SpCas9 variants can efficiently disrupt endogenous gene sites that are not currently targetable by SpCas9 in both zebrafish embryos and human cells, suggesting that they will work in a variety of different cell types and organisms. Importantly, GUIDE-seq experiments show that the global profiles of the VQR and VRER SpCas9 variants are similar to or better than those observed with wild-type SpCas9. In addition, the improved specificity D1135E variant that we identified and characterized provides a superior alternative to the widely

used wild-type SpCas9. D1135E has similar activity to wild-type SpCas9 on sites with canonical NGG PAMs but reduces genome-wide cleavage of off-target sites bearing mismatched spacer sequences and either canonical or non-canonical PAMs.

[0065] All of the SpCas9 and SaCas9 variants described herein can be rapidly incorporated into existing and widely used vectors, e.g., by simple site-directed mutagenesis, and because they require only a small number of mutations contained within the PAM-interacting domain, the variants should also work with other previously described improvements to the SpCas9 platform (e.g., truncated sgRNAs (Tsai et al., Nat Biotechnol 33, 187-197 (2015); Fu et al., Nat Biotechnol 32, 279-284 (2014)), nickase mutations (Mali et al., Nat Biotechnol 31, 833-838 (2013); Ran et al., Cell 154, 1380-1389 (2013)), dimeric FokI-dCas9 fusions (Guilinger et al., Nat Biotechnol 32, 577-582 (2014); Tsai et al., Nat Biotechnol 32, 569-576 (2014)).

[0066] Beyond the mutations to R1335 that presumably contact the 3rd PAM base position, the SpCas9 variants evolved in this study bear amino acid substitutions at D1135, G1218, and T1337, all of which are located near or adjacent to residues that make direct or indirect contacts to the 3rd PAM position in the SpCas9-PAM structure but do not themselves mediate contacts with the PAM bases (Anders et al., Nature 513, 569-573 (2014)) (FIG. 20A). Consistent with this, we found that various mutations at these positions do not appear to affect SpCas9-mediated cleavage of sites bearing an NGG PAM (FIG. 20B). These results, together with the nature of the amino acid substitutions at G1218 and T1337 in the VQR and VRER SpCas9 variants, suggest that alterations at these two positions may be gain-of-function mutations. For example, it is possible that the T1337R mutation is forming backbone or base-specific contacts near or to the 4th position of the PAM, particularly in the case of the VRER variant. The mechanistic role of mutations at D1135 remain less clear but they may perhaps influence the activity of the adjacent S1136 residue, which has been implicated in making a water-mediated contact through the minor groove to the guanine in the third position of the PAM (Anders et al., Nature 513, 569-573 (2014)). The D1135E mutation might improve specificity by disrupting this network, perhaps reducing the overall interaction energy of the SpCas9/gRNA complex with the target site, a mechanism we have previously proposed might reduce off-target effects by making cleavage of these unwanted sequences less energetically favorable (Fu et al., Nat Biotechnol 32, 279-284 (2014)).

[0067] The present results clearly establish the feasibility of engineering Cas9 nucleases with altered PAM specificities. Characterization of additional Cas9 orthologues (Esvelt et al., Nat Methods 10, 1116-1121 (2013); Fonfara et al., Nucleic Acids Res 42, 2577-2590 (2014)) or generation of domain-swapped Cas9 chimeras (Nishimasu et al., Cell. 156(5):935-49 (2014)) as previously described also provide potential avenues for targeting different PAMs. The engineering strategy delineated herein can also be performed with such orthologues or synthetic hybrid Cas9s to further diversify the range of targetable PAMs. St1Cas9 and SaCas9 make particularly attractive frameworks for future engineering efforts given their smaller sizes relative to SpCas9 and our demonstration of their robust genome editing activities in our bacterial selection systems and in human cells.

[0068] Our results strongly suggested that R1015 in wild-type SaCas9 contacts the G in the third PAM position. Without wishing to be bound by theory, the R1015H substitution may remove this contact and relax specificity at the third position; however, loss of the R1015 to G contact could also conceivably reduce the energy associated with target site binding, which may explain why the R1015H mutation alone is not sufficient for robust activity at NNNRRT sites in human cells. Because the E782K and N968K substitutions both add positive charge, it is possible that they may make non-specific interactions with the DNA phosphate backbone to compensate energetically for the loss of the R1015 to guanine contact.

[0069] The genetic approach described here does not require structural information and therefore should be applicable to many other Cas9 orthologues. The only requirement

to evolve Cas9 nucleases with broadened PAM specificities is that they function in a bacterial-based selection. While previous studies demonstrated that PAM recognition can be altered by swapping the PAM-interacting domains of highly related Cas9 orthologues (Nishimasu et al., Cell (2014)), it remains to be determined whether this strategy is generalizable or effective when using more divergent orthologues. By contrast, the evolution strategies we have described herein can be used engineer PAM recognition specificities beyond those encoded within naturally occurring Cas9 orthologues. This overall strategy can be employed to expand the targeting range and extend the utility of the numerous Cas9 orthologues that exist in nature.

[0070] SpCas9 Variants with Altered Specificity

[0071] Thus, provided herein are spCas9 variants. The SpCas9 wild type sequence is as follows:

(SEQ ID NO: 1)

10	20	30	40	50	60
MDKKYSIGLD	IGTNSVGWAV	ITDEYKVPSK	KFKVLGNTDR	HSIKKNLIGA	LLFDSGETAE
70	80	90	100	110	120
ATRLKRTARR	RYTRKRNRC	YLQEIFSNE	AKVDDSFHR	LEESFLVEED	KKHERHPIFG
130	140	150	160	170	180
NIVDEVAYHE	KYPTIYHLRK	KLVDSTDKAD	LRLIYLALAH	MIKFRGHFLI	EGDLNPDNSD
190	200	210	220	230	240
VDKLFQILVQ	TYNQLFEENP	INASGVDAKA	ILSARLSKSR	RLENLIAQLP	GEKKNGLFGN
250	260	270	280	290	300
LIALSLGLTP	NFKSNFDLAE	DAKLQLSKDT	YDDDLNLLA	QIGDQYADLF	LAAKNLSDAI
310	320	330	340	350	360
LLSDILRVNT	EITKAPLSAS	MIKRYDEHHQ	DLTLLKALVR	QQLPEKYKEI	FFDQSKNGYA
370	380	390	400	410	420
GYIDGGASQE	EFYKFIKPIL	EKMDGTEELL	VKLNREDLLR	KQRTFDNGSI	PHQIHLGELH
430	440	450	460	470	480
AILRRQEDFY	PFLKDNREKI	EKILTFRIPY	YVGPLARGNS	RFAWMTRKSE	ETITPWNFEE
490	500	510	520	530	540
VVDKGASAQ	FIERMTNFDK	NLPNEKVLPK	HSLLYEYFTV	YNELTKVKYV	TEGMRKPAFL
550	560	570	580	590	600
SGEQKKAIVD	LLFKTNRKVT	VKQLKEDYFK	KIECFDSVEI	SGVEDRFNAS	LGTYHDLLKI
610	620	630	640	650	660
IKDKDFLDNE	ENEDILEDIV	LTLTLFEDRE	MIEERLKTYA	HLPDDKVMQK	LKRRRYTGWG
670	680	690	700	710	720
RLSRKLINGI	RDQSGKTIL	DFLKSDGFAN	RNFMQLIHDD	SLTFKEDIQK	AQVSGQGDSL
730	740	750	760	770	780
HEHIANLAGS	PAIKKGILQT	VKVVDLVKV	MGRHKPENIV	IEMARENQTT	QKGQKNSRER
790	800	810	820	830	840
MKRIEBGIKE	LGSQILKEHP	VENTQLQNEK	LYLYYLQNGR	DMYVDQELDI	NRLSDYDVDH
850	860	870	880	890	900
IVPQSFLKDD	SIDNKVLTRS	DKNRGKSDNV	PSEEVVKKMK	NYWRQLLNAK	LITQRKFDNL
910	920	930	940	950	960
TKAERGGLSE	LDKAGFIKRQ	LVETRQITKH	VAQILDSRMN	TKYDENDKLI	REVKVITLKS
970	980	990	1000	1010	1020
KLVSDFRKDF	QFYKVREINN	YHHAHDAYLN	AVVGTALIKK	YPKLESEFVY	GDYKVIDVRK
1030	1040	1050	1060	1070	1080
MIAKSEQEIG	KATAKYFFYS	NIMNFFKTEI	TLANGEIRKR	PLIETNGETG	EIVWDKGRDF
1090	1100	1110	1120	1130	1140
ATVRKVLSP	QVNIIVKTEV	QTGGFSKESI	LPKRNSDKLI	ARKKDWDPKK	YGGFDSPTVA

-continued

1150	1160	1170	1180	1190	1200
YSVLVVAKVE	KGKSKKLKSV	KELLGITIME	RSSFENPID	FLEAKGYKEV	KKDLIIKLPK
1210	1220	1230	1240	1250	1260
YSLFELENGR	KRMLASAGEL	QKGNELALPS	KYVNFLYLAS	HYEKLKGSPE	DNEQKQLFVE
1270	1280	1290	1300	1310	1320
QHKHYLDEII	EQISEFSKRV	ILADANLDKV	LSAYNKHRDK	PIREQAENII	HLFTLTNLGA
1330	1340	1350	1360		
PAAFKYFDTT	IDRKRYTSTK	EVLDTLTIHQ	SITGLYETRI	DLSQLGGD	

[0072] The SpCas9 variants described herein can include mutations at one or more of the following positions: D1135, G1218, R1335, T1337 (or at positions analogous thereto). In some embodiments, the SpCas9 variants include one or more of the following mutations: D1135V; D1135E; G1218R; R1335E; R1335Q; and T1337R. In some embodiments, the SpCas9 variants are at least 80%, e.g., at least 85%, 90%, or 95% identical to the amino acid sequence of SEQ ID NO:1, e.g., have differences at up to 5%, 10%, 15%, or 20% of the residues of SEQ ID NO:1 replaced, e.g., with conservative mutations. In preferred embodiments, the variant retains desired activity of the parent, e.g., the nuclease activity (except where the parent is a nickase or a dead Cas9), and/or the ability to interact with a guide RNA and target DNA).

[0073] To determine the percent identity of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). The length of a reference sequence aligned for comparison purposes is at least 80% of the length of the reference sequence, and in some embodiments is at least 90% or 100%. The nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein nucleic acid “identity” is equivalent to nucleic acid “homology”). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. Percent identity between two polypeptides or nucleic acid sequences is determined in various ways that are within the skill in the art, for instance, using publicly available computer software such as Smith Waterman Alignment (Smith, T. F. and M. S. Waterman (1981) *J Mol Biol* 147:195-7); “BestFit” (Smith and Waterman, *Advances in Applied Mathematics*, 482-489 (1981)) as incorporated into GeneMatcher Plus™, Schwarz and Dayhoff (1979) *Atlas of Protein Sequence and Structure*, Dayhoff, M. O., Ed, pp 353-358; BLAST program (Basic Local Alignment Search Tool; (Altschul, S. F., W. Gish, et al.

(1990) *J Mol Biol* 215: 403-10), BLAST-2, BLAST-P, BLAST-N, BLAST-X, WU-BLAST-2, ALIGN, ALIGN-2, CLUSTAL, or Megalign (DNASTAR) software. In addition, those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the length of the sequences being compared. In general, for proteins or nucleic acids, the length of comparison can be any length, up to and including full length (e.g., 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%). For purposes of the present compositions and methods, at least 80% of the full length of the sequence is aligned using the BLAST algorithm and the default parameters.

[0074] For purposes of the present invention, the comparison of sequences and determination of percent identity between two sequences can be accomplished using a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

[0075] Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenyl-alanine, tyrosine.

[0076] In some embodiments, the SpCas9 variants include one of the following sets of mutations: D1135V/R1335Q/T1337R (VQR variant); D1135V/G1218R/R1335Q/T1337R (VRQR variant); D1135E/R1335Q/T1337R (EQR variant); or D1135V/G1218R/R1335E/T1337R (VRER variant).

[0077] In some embodiments, the SpCas9 variants also include one of the following mutations, which reduce or destroy the nuclease activity of the Cas9: D10, E762, D839, H983, or D986 and H840 or N863, e.g., D10A/D10N and H840A/H840N/H840Y, to render the nuclease portion of the protein catalytically inactive; substitutions at these positions could be alanine (as they are in Nishimasu et al., *Cell* 156, 935-949 (2014)), or other residues, e.g., glutamine, asparagine, tyrosine, serine, or aspartate, e.g., E762Q, H983N, H983Y, D986N, N863D, N863S, or N863H (see WO 2014/152432). In some embodiments, the variant includes mutations at D10A or H840A (which creates a single-strand nickase), or mutations at D10A and H840A (which abrogates nuclease activity; this mutant is known as dead Cas9 or dCas9).

[0078] Also provided herein are SaCas9 variants. The SaCas9 wild type sequence is as follows:

(SEQ ID NO: 2)

10	20	30	40	50
MKRNYILGLD	IGITSVGYGI	IDYETRDVID	AGVRLFKKAN	VENNEGRRSK
60	70	80	90	100
RGARRLKRRL	RHRIQRVKKL	LFDYNLLTDH	SELSGINPYE	ARVKGLSQKL

-continued

110	120	130	140	150
SEEEFSAALL	HLAKRRGVHN	VNEVEEDTGN	ELSTKEQISR	NSKALEEKYV
160	170	180	190	200
AELQLERLKK	DGEVRGSINR	FKTSDYVKEA	KQLLKVQKAY	HQLDQSFIDT
210	220	230	240	250
YIDLLETRRT	YYEGPGEKSP	FGWKDIKEWY	EMLMGHCTYF	PEELRSVKYA
260	270	280	290	300
YNADLYNALN	DLNNLVITRD	ENEKLEYEYK	FQIIENVFKQ	KKKPTLKQIA
310	320	330	340	350
KEILVNEEDI	KGYRVSTGK	PEFTNLKVIH	DIKDITARKE	IIENAELLDQ
360	370	380	390	400
IAKILTIYQS	SEDIQEELTN	LNSELTQEEL	EQISNLKGYT	GTHNLSLKAI
410	420	430	440	450
NLILDELWHT	NDNQIAIFNR	LKLVPKKVDL	SQQKEIPTTL	VDDFILSPVV
460	470	480	490	500
KRSFIQSIKV	INAIKKYGL	PNDIIELAR	EKNSKDAQKM	INEMQKRNRQ
510	520	530	540	550
TNERIEEIR	TTGKENAKYL	IEKIKLHDMQ	EGKCLYSLEA	IPLEDLLNNP
560	570	580	590	600
FNYEVDHIIP	RSVSFDNSFN	NKVLVKQEN	SKKGNRTPFQ	YLSSSDSKIS
610	620	630	640	650
YETFKKHILN	LAKGKGRISK	TKKEYLLEER	DINRFSVQKD	FINRNLDVTR
660	670	680	690	700
YATRGLMNL	RSYFRVNNLD	VKVKSSINGG	TSFLRRKWK	KKERNKGYKH
710	720	730	740	750
HAEDALIIN	ADFIFKEWKK	LDKAKVMEN	QMFEKQAES	MPEIETEY
760	770	780	790	800
KEIFITPHQI	KHIKDFKDYK	YSHRVDKPN	RELINDTLYS	TRKDDKGNTL
810	820	830	840	850
IVNNLNGLYD	KDNDKLKLI	NKSPEKLLMY	HHDPQTYQKL	KLIMEQYGDE
860	870	880	890	900
KNPLYKYEE	TGNYLTKYSK	KDNGPVIKKI	KYYGNKLNH	LDITDDYPNS
910	920	930	940	950
RNKVVKLSLK	PYRFDVYLDN	GVYKFVTVKN	LDVIKENY	EVNSKCYEEA
960	970	980	990	1000
KKLKKISQA	EFIASFYNN	LIKINGELYR	VIGVNNDDL	RIEVNMDIT
1010	1020	1030	1040	1050
YREYLENMND	KRPPIIKTI	ASKTQSIKKY	STDILGNLYE	VKSKKHPQII

KKG

[0079] The SaCas9 variants described herein include mutations at one or more of the following positions: E782, N968, and/or R1015 (or at positions analogous thereto). In some embodiments, the variants include one or more of the following mutations: R1015Q, R1015H, E782K, N968K, E735K, K929R, A1021T, K1044N. In some embodiments, the SaCas9 variants include mutations E782K, K929R, N968K, and R1015X, wherein X is any amino acid other than R. In some embodiments, the SaCas9 variants are at least 80%, e.g., at least 85%, 90%, or 95% identical to the amino acid sequence of SEQ ID NO:2, e.g., have differences at up to 5%, 10%, 15%, or 20% of the residues of SEQ ID NO:2 replaced, e.g., with conservative mutations. In preferred embodiments, the variant retains desired activity of the parent, e.g., the nuclease activity (except where the

parent is a nickase or a dead Cas9), and/or the ability to interact with a guide RNA and target DNA).

[0080] In some embodiments, the SaCas9 variants also include one of the following mutations, which may reduce or destroy the nuclease activity of the SaCas9: D10A, D556A, H557A, N580A, e.g., D10A/H557A and/or D10A/D556A/H557A/N580A, to render the nuclease portion of the protein catalytically inactive; substitutions at these positions could be alanine (as they are in Nishimasu et al., Cell 156, 935-949 (2014)), or other residues, e.g., glutamine, asparagine, tyrosine, serine, or aspartate. In some embodiments, the variant includes mutations at D10A, D556A, H557A, or N580A (which may create a single-strand nickase), or mutations at D10A/H557A and/or D10A/D556A/H557A/N580A

may (which may abrogate nuclease activity by analogy to SpCas9; these are referred to as dead Cas9 or dCas9).

[0081] Also provided herein are isolated nucleic acids encoding the SpCas9 and/or SaCas9 variants, vectors comprising the isolated nucleic acids, optionally operably linked to one or more regulatory domains for expressing the variant proteins, and host cells, e.g., mammalian host cells, comprising the nucleic acids, and optionally expressing the variant proteins.

[0082] The variants described herein can be used for altering the genome of a cell; the methods generally include expressing the variant proteins in the cells, along with a guide RNA having a region complementary to a selected portion of the genome of the cell. Methods for selectively altering the genome of a cell are known in the art, see, e.g., U.S. Pat. No. 8,697,359; US2010/0076057; US2011/0189776; US2011/0223638; US2013/0130248; WO/2008/108989; WO/2010/054108; WO/2012/164565; WO/2013/098244; WO/2013/176772; US20150050699; US20150045546; US20150031134; US20150024500; US20140377868; US20140357530; US20140349400; US20140335620; US20140335063; US20140315985; US20140310830; US20140310828; US20140309487; US20140304853; US20140298547; US20140295556; US20140294773; US20140287938; US20140273234; US20140273232; US20140273231; US20140273230; US20140271987; US20140256046; US20140248702; US20140242702; US20140242700; US20140242699; US20140242664; US20140234972; US20140227787; US20140212869; US20140201857; US20140199767; US20140189896; US20140186958; US20140186919; US20140186843; US20140179770; US20140179006; US20140170753; Makarova et al., "Evolution and classification of the CRISPR-Cas systems" 9(6) Nature Reviews Microbiology 467-477 (1-23) (June 2011); Wiedenheft et al., "RNA-guided genetic silencing systems in bacteria and archaea" 482 Nature 331-338 (Feb. 16, 2012); Gasiunas et al., "Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria" 109(39) Proceedings of the National Academy of Sciences USA E2579-E2586 (Sep. 4, 2012); Jinek et al., "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity" 337 Science 816-821 (Aug. 17, 2012); Carroll, "A CRISPR Approach to Gene Targeting" 20(9) Molecular Therapy 1658-1660 (September 2012); U.S. Appl. No. 61/652,086, filed May 25, 2012; Al-Attar et al., "Clustered Regularly Interspaced Short Palindromic Repeats (CRISPRs): The Hallmark of an Ingenious Antiviral Defense Mechanism in Prokaryotes, Biol Chem. (2011) vol. 392, Issue 4, pp. 277-289; Hale et al., "Essential Features and Rational Design of CRISPR RNAs That Function With the Cas RAMP Module Complex to Cleave RNAs, Molecular Cell, (2012) vol. 45, Issue 3, 292-302.

[0083] The variant proteins described herein can be used in place of the SpCas9 proteins described in the foregoing references with guide RNAs that target sequences that have PAM sequences according to the following Table 4.

TABLE 4

Variant protein	Stronger PAM	Weaker PAM
SpCas9-D1135E	NGG	NAG, NGA, and NNGG
SpCas9-VQR	NGAN and NGCG	NGGG, NGTG, and NAAG
SpCas9-VRQR	NGAN	

TABLE 4-continued

Variant protein	Stronger PAM	Weaker PAM
SpCas9-EQR	NGAG	NGAT, NGAA, and NGCG
SpCas9-VRER	NGCG	NGCA, NGCC, and NGCT
SaCas9-KKH	NNNRRT	
SaCas9-KKQ	NNNRRT (SEQ ID NO: 45)	NNNRRT
SaCas9-KKE	NNCRRT (SEQ ID NO: 47)	NNNRRT
SaCas9- (KKL or KKM)	NNTRRT (SEQ ID NO: 48)	NNNRRT

[0084] In addition, the variants described herein can be used in fusion proteins in place of the wild-type Cas9 or other Cas9 mutations (such as the dCas9 or Cas9 nickase described above) as known in the art, e.g., a fusion protein with a heterologous functional domains as described in WO 2014/124284. For example, the variants, preferably comprising one or more nuclease-reducing or killing mutation, can be fused on the N or C terminus of the Cas9 to a transcriptional activation domain or other heterologous functional domains (e.g., transcriptional repressors (e.g., KRAB, ERD, SID, and others, e.g., amino acids 473-530 of the ets2 repressor factor (ERF) repressor domain (ERD), amino acids 1-97 of the KRAB domain of KOX1, or amino acids 1-36 of the Mad mSIN3 interaction domain (SID); see Beerli et al., PNAS USA 95:14628-14633 (1998)) or silencers such as Heterochromatin Protein 1 (HP1, also known as swi6), e.g., HP1 α or HP1 β ; proteins or peptides that could recruit long non-coding RNAs (lncRNAs) fused to a fixed RNA binding sequence such as those bound by the MS2 coat protein, endoribonuclease Csy4, or the lambda N protein; enzymes that modify the methylation state of DNA (e.g., DNA methyltransferase (DNMT) or TET proteins); or enzymes that modify histone subunits (e.g., histone acetyltransferases (HAT), histone deacetylases (HDAC), histone methyltransferases (e.g., for methylation of lysine or arginine residues) or histone demethylases (e.g., for demethylation of lysine or arginine residues)) as are known in the art can also be used. A number of sequences for such domains are known in the art, e.g., a domain that catalyzes hydroxylation of methylated cytosines in DNA. Exemplary proteins include the Ten-Eleven-Translocation (TET)1-3 family, enzymes that converts 5-methylcytosine (5-mC) to 5-hydroxymethylcytosine (5-hmC) in DNA.

[0085] Sequences for human TET1-3 are known in the art and are shown in the following table:

GenBank Accession Nos.		
Gene	Amino Acid	Nucleic Acid
TET1	NP_085128.2	NM_030625.2
TET2*	NP_001120680.1 (var 1) NP_060098.3 (var 2)	NM_001127208.2 NM_017628.4
TET3	NP_659430.1	NM_144993.1

*Variant (1) represents the longer transcript and encodes the longer isoform (a). Variant (2) differs in the 5' UTR and in the 3' UTR and coding sequence compared to variant 1. The resulting isoform (b) is shorter and has a distinct C-terminus compared to isoform a.

[0086] In some embodiments, all or part of the full-length sequence of the catalytic domain can be included, e.g., a catalytic module comprising the cysteine-rich extension and

the 2OGFeDO domain encoded by 7 highly conserved exons, e.g., the Tet catalytic domain comprising amino acids 1580-2052, Tet2 comprising amino acids 1290-1905 and Tet3 comprising amino acids 966-1678. See, e.g., FIG. 1 of Iyer et al., *Cell Cycle*, 2009 Jun. 1; 8(11):1698-710. Epub 2009 Jun. 27, for an alignment illustrating the key catalytic residues in all three Tet proteins, and the supplementary materials thereof for full length sequences (see, e.g., seq 2c); in some embodiments, the sequence includes amino acids 1418-2136 of Tet1 or the corresponding region in Tet2/3.

[0087] Other catalytic modules can be from the proteins identified in Iyer et al., 2009.

[0088] In some embodiments, the heterologous functional domain is a biological tether, and comprises all or part of (e.g., DNA binding domain from) the MS2 coat protein, endoribonuclease Csy4, or the lambda N protein. These proteins can be used to recruit RNA molecules containing a specific stem-loop structure to a locale specified by the dCas9 gRNA targeting sequences. For example, a dCas9 variant fused to MS2 coat protein, endoribonuclease Csy4, or lambda N can be used to recruit a long non-coding RNA (lncRNA) such as XIST or HOTAIR; see, e.g., Keryer-Bibens et al., *Biol. Cell* 100:125-138 (2008), that is linked to the Csy4, MS2 or lambda N binding sequence. Alternatively, the Csy4, MS2 or lambda N protein binding sequence can be linked to another protein, e.g., as described in Keryer-Bibens et al., *supra*, and the protein can be targeted to the dCas9 variant binding site using the methods and compositions described herein. In some embodiments, the Csy4 is catalytically inactive. In some embodiments, the Cas9 variant, preferably a dCas9 variant, is fused to FokI as described in WO 2014/204578.

[0089] In some embodiments, the fusion proteins include a linker between the dCas9 variant and the heterologous functional domains. Linkers that can be used in these fusion proteins (or between fusion proteins in a concatenated structure) can include any sequence that does not interfere with the function of the fusion proteins. In preferred embodiments, the linkers are short, e.g., 2-20 amino acids, and are typically flexible (i.e., comprising amino acids with a high degree of freedom such as glycine, alanine, and serine). In some embodiments, the linker comprises one or more units consisting of GGGGS (SEQ ID NO:188) or GGGGS (SEQ ID NO:189), e.g., two, three, four, or more repeats of the GGGGS (SEQ ID NO:188) or GGGGS (SEQ ID NO:189) unit. Other linker sequences can also be used.

[0090] Expression Systems

[0091] To use the Cas9 variants described herein, it may be desirable to express them from a nucleic acid that encodes them. This can be performed in a variety of ways. For example, the nucleic acid encoding the Cas9 variant can be cloned into an intermediate vector for transformation into prokaryotic or eukaryotic cells for replication and/or expression. Intermediate vectors are typically prokaryote vectors, e.g., plasmids, or shuttle vectors, or insect vectors, for storage or manipulation of the nucleic acid encoding the Cas9 variant for production of the Cas9 variant. The nucleic acid encoding the Cas9 variant can also be cloned into an expression vector, for administration to a plant cell, animal cell, preferably a mammalian cell or a human cell, fungal cell, bacterial cell, or protozoan cell.

[0092] To obtain expression, a sequence encoding a Cas9 variant is typically subcloned into an expression vector that

contains a promoter to direct transcription. Suitable bacterial and eukaryotic promoters are well known in the art and described, e.g., in Sambrook et al., *Molecular Cloning, A Laboratory Manual* (3d ed. 2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel et al., eds., 2010). Bacterial expression systems for expressing the engineered protein are available in, e.g., *E. coli*, *Bacillus* sp., and *Salmonella* (Palva et al., 1983, *Gene* 22:229-235). Kits for such expression systems are commercially available. Eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known in the art and are also commercially available.

[0093] The promoter used to direct expression of a nucleic acid depends on the particular application. For example, a strong constitutive promoter is typically used for expression and purification of fusion proteins. In contrast, when the Cas9 variant is to be administered in vivo for gene regulation, either a constitutive or an inducible promoter can be used, depending on the particular use of the Cas9 variant. In addition, a preferred promoter for administration of the Cas9 variant can be a weak promoter, such as HSV TK or a promoter having similar activity. The promoter can also include elements that are responsive to transactivation, e.g., hypoxia response elements, Gal4 response elements, lac repressor response element, and small molecule control systems such as tetracycline-regulated systems and the RU-486 system (see, e.g., Gossen & Bujard, 1992, *Proc. Natl. Acad. Sci. USA*, 89:5547; Oligino et al., 1998, *Gene Ther.*, 5:491-496; Wang et al., 1997, *Gene Ther.*, 4:432-441; Neering et al., 1996, *Blood*, 88:1147-55; and Rendahl et al., 1998, *Nat. Biotechnol.*, 16:757-761).

[0094] In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the nucleic acid in host cells, either prokaryotic or eukaryotic. A typical expression cassette thus contains a promoter operably linked, e.g., to the nucleic acid sequence encoding the Cas9 variant, and any signals required, e.g., for efficient polyadenylation of the transcript, transcriptional termination, ribosome binding sites, or translation termination. Additional elements of the cassette may include, e.g., enhancers, and heterologous spliced intronic signals.

[0095] The particular expression vector used to transport the genetic information into the cell is selected with regard to the intended use of the Cas9 variant, e.g., expression in plants, animals, bacteria, fungus, protozoa, etc. Standard bacterial expression vectors include plasmids such as pBR322 based plasmids, pSKF, pET23D, and commercially available tag-fusion expression systems such as GST and LacZ.

[0096] Expression vectors containing regulatory elements from eukaryotic viruses are often used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 late promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[0097] The vectors for expressing the Cas9 variants can include RNA Pol III promoters to drive expression of the guide RNAs, e.g., the H1, U6 or 7SK promoters. These human promoters allow for expression of Cas9 variants in mammalian cells following plasmid transfection.

[0098] Some expression systems have markers for selection of stably transfected cell lines such as thymidine kinase, hygromycin B phosphotransferase, and dihydrofolate reductase. High yield expression systems are also suitable, such as using a baculovirus vector in insect cells, with the gRNA encoding sequence under the direction of the polyhedrin promoter or other strong baculovirus promoters.

[0099] The elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of recombinant sequences.

[0100] Standard transfection methods are used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of protein, which are then purified using standard techniques (see, e.g., Colley et al., 1989, J. Biol. Chem., 264:17619-22; Guide to Protein Purification, in Methods in Enzymology, vol. 182 (Deutscher, ed., 1990)). Transformation of eukaryotic and prokaryotic cells are performed according to standard techniques (see, e.g., Morrison, 1977, J. Bacteriol. 132:349-351; Clark-Curtiss & Curtiss, Methods in Enzymology 101:347-362 (Wu et al., eds, 1983)).

[0101] Any of the known procedures for introducing foreign nucleotide sequences into host cells may be used. These

include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, nucleofection, liposomes, microinjection, naked DNA, plasmid vectors, viral vectors, both episomal and integrative, and any of the other well-known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see, e.g., Sambrook et al., supra). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing the Cas9 variant.

[0102] The present invention includes the vectors and cells comprising the vectors.

EXAMPLES

[0103] The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

[0104] Methods

[0105] The following materials and methods were used in Examples 1 and 2.

[0106] Plasmids and Oligonucleotides

[0107] Schematic maps and DNA sequences for parent constructs used in this study can be found in FIGS. 5A-J and SEQ ID NOs:7-20. Sequences of oligonucleotides used to generate the positive selection plasmids, negative selection plasmids, and site-depletion libraries are available in Table 1. Sequences of all gRNA targets in this study are available in Table 2. Point mutations in Cas9 were generated by PCR.

TABLE 1

		SEQ ID NO:
Oligos used to generate positive and negative selection plasmids		
sequence	description	
ctagaGGGCACGGGCAGCTTGCCGGTGGgcatg	top oligo to clone site 1 into the positive selection vector (XbaI/SphI cut p11-lacY-wtx1)	190
cccACCGGCAAGCTGCCCCGTGCCct	bottom oligo to clone site 1 into the positive selection vector	191
ctagaGGTGCCTTCGAACCTTACCTCGGgcatg	top oligo to clone site 2 into the positive selection vector (XbaI/SphI cut p11-lacY-wtx1)	192
cccGAGGTGAAGTTCGAGGGGACct	bottom oligo to clone site 2 into the positive selection vector	193
aattcGGGCACGGGCAGCTTGCCGGTGGgcatg	top oligo to clone site 1 into the negative selection vector (EcoRI/SphI cut p11-lacY-wtx1)	194
cccACCGGCAAGCTGCCCCGTGCCg	bottom oligo to clone site 1 into the negative selection vector	195
aattcGGTGCCTTCGAACCTTACCTCGGgcatg	top oligo to clone site 2 into the negative selection vector (EcoRI/SphI cut p11-lacY-wtx1)	196
cccGAGGTGAAGTTCGAGGGGACc	bottom oligo to clone site 2 into the negative selection vector	197
Oligos used to generate libraries for site-depletion experiments		
sequence	description	
GCAGgaattcGGGCACGGGCAGCTTGCCGGN	top strand oligo for site 1 PAM library, cut with EcoRI once filled in	198
NNNNNCTNNNGCGCAGGTACGAGGCATG	top strand oligo for site 2 PAM library, cut with EcoRI once filled in	199
GCAGgaattcGTGCGCTCGAACCTTACCTN	top strand oligo for site 2 PAM library, cut with EcoRI once filled in	200
NNNNNCTNNNGCGCAGGTACGAGGCATG	reverse primer to fill in library oligos	
/5Phos/ccTcGTGAcTgGcGc		

TABLE 1-continued

		SEQ ID NO:
Primers used to amplify site-depletion libraries for sequencing		
sequence	description	
GATACCGCTCGCCGCAGC	forward primer	201
CTGCGTTCTGATTAACTGTATCAGGC	reverse primer	202
Primers used for T7E1 experiments		
sequence	description	
GGAGATGTAAATCACCTCCATCTGA	forward primer targeted to th1 in zebrafish	203
ATGTTAGCCTACCTCGAAAACCTTC	reverse primer targeted to th1 in zebrafish	204
CCTGTGCTCTCCTGTTTTAGGTAT	forward primer targeted to tialL in zebrafish	205
AACATGGTAAGAAGCGTGAGTGT	reverse primer targeted to tialL in zebrafish	206
CAGGCTGTTGAACCGTAGATTAGT	forward primer targeted to fh in zebrafish	207
TCCACATGTTTTGAGTTTGAGAGTC	reverse primer targeted to fh in zebrafish	208
GGACAGCTGGTCAGAGGGG	forward primer targeted to EMX1 in U2OS human cells	209
CCATAGGGAAGGGGACACTGG	reverse primer targeted to EMX1 in U2OS human cells	210
GGCCCGGAAAGAGTTGCTG	forward primer tar- geted to FANCF in U2OS human cells	211
GCCCTACATCTGCTCTCCCTCC	reverse primer tar- geted to FANCF in U2OS human cells	212
CCAGCACAACCTACTCGCACTTGAC	forward primer tar- geted to RUNX1 in U2OS human cells	213
CATCACCAACCCACAGCCAAGG	reverse primer tar- geted to RUNX1 in U2OS human cells	214
GATGAGGGCTCCAGATGGCAC	forward primer tar- geted to VEGFA in U2OS human cells	215
GAGGAGGGAGCAGGAAAGTGAGG	reverse primer tar- geted to VEGFA in U2OS human cells	216

TABLE 2

<i>S. pyogenes</i> gRNAs						
EGFP						
NXX gRNAs						
Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
FYF1320	NGG 1-20	20	GGGCACGGGCAGCTTGCCGG	217	GGGCACGGGCAGCTTGCCGGTGGT	218
BPK1345	NGG 2-20	20	GTCGCCCTCGAACTTCACCT	219	GTCGCCCTCGAACTTCACCTCGGC	220
MSP792	NGG 3-20	20	GGTCGCCACCATGGTGAGCA	221	GGTCGCCACCATGGTGAGCAAGGG	222
MSP795	NGG 4-20	20	GGTCAGGGTGGTCACGAGGG	223	GGTCAGGGTGGTCACGAGGGTGGG	224
FYF1328	NGG 5-20	20	GGTGGTGCAGATGAACCTCA	225	GGTGGTGCAGATGAACCTCAGGGT	226
MSP160	NAG 1-20	20	GGGTGGTGCCCATCTGGTC	227	GGGTGGTGCCCATCTGGTCGAGC	228
MSP161	NAG 2-20	20	GACGTAAACGGCCACAAGTT	229	GACGTAAACGGCCACAAGTTCAGC	230
MSP162	NAG 3-20	20	GTGCAGATGAACCTCAGGGT	231	GTGCAGATGAACCTCAGGGTCAGC	232
MSP163	NAG 4-20	20	GGGTGGTCACGAGGGTGGGC	233	GGGTGGTCACGAGGGTGGGCCAGG	234
MSP164	NAA 1-20	20	GGTCGAGCTGGACGGCGACG	235	GGTCGAGCTGGACGGCGACGTAAA	236
MSP165	NAA 2-20	20	GTCGAGCTGGACGGCGACGT	237	GTCGAGCTGGACGGCGACGTAAAC	238
MSP168	NGA 1-20	20	GGGGTGGTGCCCATCTGGT	239	GGGGTGGTGCCCATCTGGTGCAG	240
MSP366	NGA 2-20	20	GCCACCATGGTGAGCAAGGG	241	GCCACCATGGTGAGCAAGGGCGAG	242
MSP171	NGA 3-20	20	GTCGCCCTCCAGCTCGACCA	243	GTCGCCCTCCAGCTCGACCAAGGAT	244
BPK1466	NGA 4-20	20	GCATCGCCCTCGCCCTCGCC	245	GCATCGCCCTCGCCCTCGCCGGAC	246
BPK1468	NGA 5-20	20	GTTTCGAGGGCGACACCTGG	247	GTTTCGAGGGCGACACCTGGTGAA	248
NGXX gRNAs						
Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
BPK1468	NGAA 1-20	20	GTTTCGAGGGCGACACCTGG	249	GTTTCGAGGGCGACACCTGGTGAA	250
MSP807	NGAA 2-20	20	GTTTCACAGGGTGTGCGCCCT	251	GTTTCACAGGGTGTGCGCCCTCGAA	252
BPK1469	NGAA 3-20	20	GCAGAAGAACGGCATCAAGG	253	GCAGAAGAACGGCATCAAGGTGAA	254
MSP787	NGAA 3-17	17	GAAGAACGGCATCAAGG	255	GAAGAACGGCATCAAGGTGAA	256
MSP170	NGAC 1-20	20	GCCACCCCTCGTGACCAACC	257	GCCACCCCTCGTGACCAACCCTGAC	258
MSP790	NGAC 2-20	20	GCCCTTGCTCACCATGGTGG	259	GCCCTTGCTCACCATGGTGCGAC	260

TABLE 2-continued

MSP171	NGAT 1-20	20	GTCGCCGTCAGCTCGACCA	261.	GTCGCCGTCAGCTCGACCA	262.
BPK1979	NGAT 1-17	17	GCCGTCCAGCTCGACCA	263.	GCCGTCCAGCTCGACCA	264.
MSP169	NGAT 2-20	20	GTGTCCGGCGAGGGCGAGGG	265.	GTGTCCGGCGAGGGCGAGGG	266.
BPK1464	NGAT 3-20	20	GGGCAGCTTGCCGGTGGTGC	267.	GGGCAGCTTGCCGGTGGTGC	268.
MSP788	NGAT 3-19	19	GGCAGCTTGCCGGTGGTGC	269.	GGCAGCTTGCCGGTGGTGC	270.
MSP789	NGAT 3-18	18	GCAGCTTGCCGGTGGTGC	271.	GCAGCTTGCCGGTGGTGC	272.
MSP168	NGAG 1-20	20	GGGGTGGTGCCCATCCTGGT	273.	GGGGTGGTGCCCATCCTGGT	274.
MSP783	NGAG 1-19	19	GGGTGGTGCCCATCCTGGT	275.	GGGTGGTGCCCATCCTGGT	276.
MSP784	NGAG 1-18	18	GGTGGTGCCCATCCTGGT	277.	GGTGGTGCCCATCCTGGT	278.
MSP785	NGAG 1-17	17	GTGGTGCCCATCCTGGT	279.	GTGGTGCCCATCCTGGT	280.
MSP366	NGAG 2-20	20	GCCACCATGGTGAGCAAGGG	281.	GCCACCATGGTGAGCAAGGG	282.
MSP368	NGAG 3-20	20	GCCGTAGGTGAGGGTGGTCA	283.	GCCGTAGGTGAGGGTGGTCA	284.
BPK1974	NGAG 3-17	17	GTAGGTGAGGGTGGTCA	285.	GTAGGTGAGGGTGGTCA	286.
MSP376	NGAG 4-20	20	GCTGCCCGACAACCACTACC	287.	GCTGCCCGACAACCACTACC	288.
BPK1978	NGAG 4-17	17	GCCCCGACAACCACTACC	289.	GCCCCGACAACCACTACC	290.
MSP1028	NGCA 1-20	20	GCGAGGGCGATGCCACCTAC	291.	GCGAGGGCGATGCCACCTAC	292.
MSP1030	NGCA 2-20	20	GTGGTCGGGGTAGCGGCTGA	293.	GTGGTCGGGGTAGCGGCTGA	294.
MSP1032	NGCC 1-20	20	GGAGCTGTTACCCGGGGTGG	295.	GGAGCTGTTACCCGGGGTGG	296.
MSP1033	NGCC 2-20	20	GAACCTGTGGCCGTTACGT	297.	GAACCTGTGGCCGTTACGT	298.
MSP1036	NGCT 1-20	20	GGTGAAACAGCTCCTCGCCCT	299.	GGTGAAACAGCTCCTCGCCCT	300.
MSP1037	NGCT 2-20	20	GGTGGTGCCCATCCTGGTCCG	301.	GGTGGTGCCCATCCTGGTCCG	302.
MSP800	NGCG 1-20	20	GCCACAAGTTGAGCGTGTCC	303.	GCCACAAGTTGAGCGTGTCC	304.
MSP801	NGCG 2-20	20	GCGTGTCCGGCGAGGGCGAG	305.	GCGTGTCCGGCGAGGGCGAG	306.
MSP1360	NGCG 2-18	18	GTGTCCGGCGAGGGCGAG	307.	GTGTCCGGCGAGGGCGAG	308.
MSP802	NGCG 3-20	20	GCCCCAAGGCTACGTCCAGG	309.	GCCCCAAGGCTACGTCCAGG	310.
MSP803	NGCG 4-20	20	GTCGTCTTGAAGAAGATGG	311.	GTCGTCTTGAAGAAGATGG	312.
MSP1366	NGCG 4-17	17	GTCCTTGAAGAAGATGG	313.	GTCCTTGAAGAAGATGG	314.
MSP792	NGGG 1-20	20	GGTCGCCACCATGGTGAGCA	315.	GGTCGCCACCATGGTGAGCA	316.
MSP794	NGGG 2-20	20	GGTGGTCACGAGGGTGGGCC	317.	GGTGGTCACGAGGGTGGGCC	318.
MSP796	NGTG 1-20	20	GATCCACCGGTGCCACCAT	319.	GATCCACCGGTGCCACCAT	320.
MSP799	NGTG 2-20	20	GTAACCGGCCACAAGTTGAG	321.	GTAACCGGCCACAAGTTGAG	322.

Endogenous genes
EMX1

Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
FYF1548	NGG 1-20	20	GAGTCCGAGCAGAAGAAGAA	323.	GAGTCCGAGCAGAAGAAGGGC	324.
MSP809	NGG 2-20	20	GTCACCTCCAATGACTAGGG	325.	GTCACCTCCAATGACTAGGGTGGG	326.
MSP811	NGA 1-20	20	GAGGAGGAAGGGCCTGAGTC	327.	GAGGAGGAAGGGCCTGAGTCCGAG	328.
MSP812	NGA 2-20	20	GGTTGCCACCCCTAGTCATT	329.	GGTTGCCACCCCTAGTCATTGGAG	330.
MSP813	NGA 3-20	20	GCTGAGCTGAGAGCCTGATG	331.	GCTGAGCTGAGAGCCTGATGGGAA	332.
MSP814	NGA 4-20	20	GCCACGAAGCAGGCCAATGG	333.	GCCACGAAGCAGGCCAATGGGGAG	334.

FANCF

Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
DR348	NGG 1-20	20	GGAATCCCTTCTGCAGCACC	335.	GGAATCCCTTCTGCAGCACC	336.
MSP815	NGG 2-20	20	GCTGCAGAAGGGATTCCATG	337.	GCTGCAGAAGGGATTCCATGAGGT	338.
MSP818	NGA 1-20	20	GAATCCCTTCTGCAGCACCT	339.	GAATCCCTTCTGCAGCACCTGGAT	340.
MSP819	NGA 2-20	20	GTGCTGCAGAAGGGATTCCA	341.	GTGCTGCAGAAGGGATTCCATGAG	342.
MSP820	NGA 3-20	20	GCGGCGGCTGCACAACCACT	343.	GCGGCGGCTGCACAACCACTGGAG	344.
MSP885	NGA 4-20	20	GGTTGTGCAGCCGCGCTCC	345.	GGTTGTGCAGCCGCGCTCCAGAG	346.
MSP1060	NGCG 1-20	20	GAGGCAAGAGGGCGGCTTTG	347.	GAGGCAAGAGGGCGGCTTTGGGCG	348.
MSP1061	NGCG 2-19	19	GGGGTCCAGTTCCGGGATT	349.	GGGGTCCAGTTCCGGGATTAGCG	350.
MSP1062	NGCG 3-20	20	GCAGAAGGGATTCCATGAGG	351.	GCAGAAGGGATTCCATGAGGTGCG	352.
MSP1063	NGCG 4-19	19	GAAGGGATTCCATGAGGTG	353.	GAAGGGATTCCATGAGGTGCGCG	354.

RUNX1

Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
MSP823	NGG 1-20	20	GCTGAAACAGTGACCTGTCT	355.	GCTGAAACAGTGACCTGTCTGGT	356.
MSP824	NGG 2-20	20	GATGTAGGGCTAGAGGGGTG	357.	GATGTAGGGCTAGAGGGGTGAGGC	358.
MSP826	NGA 1-20	20	GGTGCAATTTTCAGGAGGAAG	359.	GGTGCAATTTTCAGGAGGAAGCGAT	360.
MSP827	NGA 2-20	20	GTTTTCGCTCCGAAGGTAAAG	361.	GTTTTCGCTCCGAAGGTAAAGAA	362.
MSP828	NGA 3-20	20	GAGATGTAGGGCTAGAGGGG	363.	GAGATGTAGGGCTAGAGGGGTGAG	364.
MSP829	NGA 4-20	20	GCAGAGGGGAGAAGAAAGAG	365.	GCAGAGGGGAGAAGAAAGAGAT	366.
MSP1068	NGCG 1-19	19	GGGTGCATTTTCAGGAGGA	367.	GGGTGCATTTTCAGGAGGAAGCG	368.

TABLE 2-continued

VEGFA						
Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
VC228	NGG 1-20	20	GGTGAGTGAGTGTGTCGCTG	369.	GGTGAGTGAGTGTGTCGCTGTGGG	370.
MSP830	NGG 2-20	20	GTTGGAGCGGGGAGAAGGCC	371.	GTTGGAGCGGGGAGAAGGCCAGGG	372.
BPK1846	NGA 1-20	20	GCGAGCAGCGCTCTTCGAGAG	373.	GCGAGCAGCGCTCTTCGAGAGTGAG	374.
BPK1848	NGA 2-20	20	GACGTGTGTGTCTGTGTGGG	375.	GACGTGTGTGTCTGTGTGGGTGAG	376.
BPK1850	NGA 3-20	20	GGTTGAGGGCGTTGGAGCGG	377.	GGTTGAGGGCGTTGGAGCGGGGAG	378.
MSP831	NGA 4-20	20	GCTTTGGAAGGGGGTGGGG	379.	GCTTTGGAAGGGGGTGGGGGGAG	380.
MSP1074	NGCG 1-20	20	GCAGACGGCAGTCACTAGGG	381.	GCAGACGGCAGTCACTAGGGGGCG	382.
MSP1075	NGCG 2-20	20	GCTGGGTGAATGGAGCGAGC	383.	GCTGGGTGAATGGAGCGAGCAGCG	384.
MSP1076	NGCG 3-19	19	GTGTGGGTGAGTGAGTGTG	385.	GTGTGGGTGAGTGAGTGTGTGCG	386.
MSP1077	NGCG 4-19	19	GTGTGCGTGTGGGGTTGAG	387.	GTGTGCGTGTGGGGTTGAGGGCG	388.
<i>S. aureus</i> gRNAs EGFP						
Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
MSP1395	Site 1-20	20	GTCGTGCTGCTTCATGTGGT	389.	GTCGTGCTGCTTCATGTGGTCGGGGT	390.
MSP1405	Site 1-23	23	GAAGTCGTGCTGCTTCATGTGGT	391.	GAAGTCGTGCTGCTTCATGTGGTCGGGGT	392.
MSP1396	Site 2-21	21	GCCGGTGGTGCGATGAACTT	393.	GCCGGTGGTGCGATGAACTTCAGGGT	394.
MSP1397	Site 3-21	21	GCCGTAGGTCAGGGTGGTCAC	395.	GCCGTAGGTCAGGGTGGTCACAGGGT	396.
MSP1400	Site 4-21	21	GCAACATCCTGGGGCACAAGC	397.	GCAACATCCTGGGGCACAAGCTGGAGT	398.
MSP1404	Site 4-22	22	GGCAACATCCTGGGGCACAAGC	399.	GGCAACATCCTGGGGCACAAGCTGGAGT	400.
MSP1398	Site 5-21	21	GAAGCACTGCACGCCGTAGGT	401.	GAAGCACTGCACGCCGTAGGTCAAGGT	402.
MSP1408	Site 5-24	24	GCTGAAGCACTGCACGCCGTAGGT	403.	GCTGAAGCACTGCACGCCGTAGGTCAAGGT	404.
MSP1428	Site 6-21	21	GCCCTCGAACTTCACCTCGGC	405.	GCCCTCGAACTTCACCTCGGC CGGGT	406.
MSP1409	Site 6-24	24	GTCGCCCTCGAACTTCACCTCGGC	407.	GTCGCCCTCGAACTTCACCTCGGC CGGGT	408.
MSP1403	Site 7-22	22	GCAAGGGCGAGGAGCTGTTAC	409.	GCAAGGGCGAGGAGCTGTTACCCGGGT	410.
MSP1406	Site 7-24	24	GAGCAAGGGCGAGGAGCTGTTAC	411.	GAGCAAGGGCGAGGAGCTGTTACCCGGGT	412.
MSP1410	Site 8-24	24	GCCCTTCAGCTCGATGCGGTTTAC	413.	GCCCTTCAGCTCGATGCGGTTTACCCAGGT	414.
<i>S. thermophilus</i> gRNAs EGFP						
Prep Name	Name	Spacer length (nt)	Sequence	SEQ ID NO:	Sequence with extended PAM	SEQ ID NO:
MSP1412	Site 1-20	20	GTCTATATCATGGCCGACAA	415.	GTCTATATCATGGCCGACAAGCAGAA	416.
MSP1414	Site 2-21	21	GCAGCTCGCCGACCACTACCA	417.	GCAGCTCGCCGACCACTACCAGCAGAA	418.
MSP1417	Site 2-23	23	GTGCAGCTCGCCGACCACTACCA	419.	GTGCAGCTCGCCGACCACTACCAGCAGAA	420.
MSP1413	Site 3-21	21	GCCTTCGGGCATGGCGGACTT	421.	GCCTTCGGGCATGGCGGACTTGAAGAA	422.
MSP1418	Site 3-24	24	GTAGCCTTCGGGCATGGCGGACTT	423.	GTAGCCTTCGGGCATGGCGGACTTGAAGAA	424.
MSP1416	Site 4-23	23	GTCTATATCATGGCCGACAA	425.	GTCTATATCATGGCCGACAAGCAGAAGAA	426.
MSP1415	Site 5-23	23	GTCTTGTAGTTGCCGTCGTCTT	427.	GTCTTGTAGTTGCCGTCGTCTTGAAGAA	428.
MSP1419	Site 5-24	24	GGTCTTGTAGTTGCCGTCGTCTT	429.	GGTCTTGTAGTTGCCGTCGTCTTGAAGAA	430.

[0108] Bacterial Cas9/sgRNA expression plasmids were constructed with two T7 promoters to separately express Cas9 and the sgRNA. These plasmids encode human codon optimized versions of Cas9 for *S. pyogenes* (BPK764, SpCas9 sequence subcloned from JDS246¹⁷), *S. thermophilus* Cas9 from CRISPR locus 1 (MSP1673, St1Cas9 sequence modified from previous published description²⁰), and *S. aureus* (BPK2101, SaCas9 sequence codon optimized from Uniprot J7RUA5). Previously described sgRNA sequences were utilized for SpCas9^{34, 35} and St1Cas9²⁰, while the SaCas9 sgRNA sequence was determined by searching the European Nucleotide Archive sequence HE980450 for crRNA repeats using CRISPRfinder and identifying the tracrRNA using a bioinformatic approach similar to one previously described³⁶. Annealed oligos to complete the spacer complementarity region of the sgRNA were ligated into BsaI cut BPK764 and BPK2101, or BspMI cut MSP1673 (append 5'-ATAG to the spacer to generate the

top oligo and append 5'-AAAC to the reverse complement of the spacer sequence to generate the bottom oligo).

[0109] Residues 1097-1368 of SpCas9 were randomly mutagenized using Mutazyme II (Agilent Technologies) at a rate of ~5.2 substitutions/kilobase to generate mutagenized PAM-interacting (PI) domain libraries. The theoretical complexity of each PI domain library was estimated to be greater than 10⁷ clones based on the number of transformants obtained. Positive and negative selection plasmids were generated by ligating annealed target site oligos into XbaI-SphI or EcoRI/SphI cut p11-lacY-wtx1¹⁷, respectively.

[0110] Two randomized PAM libraries (each with a different protospacer sequence) were constructed using Klenow(-exo) to fill-in the bottom strand of oligos that contained six randomized nucleotides directly adjacent to the 3' end of the protospacer (see Table 1). The double-stranded product was cut with EcoRI to leave EcoRI/SphI ends for ligation into cut p11-lacY-wtx1. The theoretical complexity

of each randomized PAM library was estimated to be greater than 10^6 based on the number of transformants obtained.

[0111] SpCas9 and SpCas9 variants were expressed in human cells from vectors derived from JDS246¹⁶. For St1Cas9 and SaCas9, the Cas9 ORFs from MSP1673 and BPK2101 were subcloned into a CAG promoter vector to generate MSP1594 and BPK2139, respectively. Plasmids for U6 expression of sgRNAs (into which desired spacer oligos can be cloned) were generated using the sgRNA sequences described above for the SpCas9 sgRNA (BPK1520), the St1Cas9 sgRNA (BPK2301), and the SaCas9 gRNA (VVT1). Annealed oligos to complete the spacer complementarity region of the sgRNA were ligated into the BsmBI overhangs of these vectors (append 5'-CACC to the spacer to generate the top oligo and append 5'-AAAC to the reverse complement of the spacer sequence to generate the bottom oligo).

[0112] Bacterial-based positive selection assay for evolving SpCas9 variants Competent *E. coli* BW25141(λ DE3)²³ containing a positive selection plasmid (with embedded target site) were transformed with Cas9/sgRNA-encoding plasmids. Following a 60 minute recovery in SOB media, transformations were plated on LB plates containing either chloramphenicol (non-selective) or chloramphenicol+10 mM arabinose (selective). Cleavage of the positive selection plasmid was estimated by calculating the survival frequency: colonies on selective plates/colonies on non-selective plates (see also FIG. 12).

[0113] To select for SpCas9 variants that can cleave novel PAMs, PI-domain mutagenized Cas9/sgRNA plasmid libraries were electroporated into *E. coli* BW25141(λ DE3) cells containing a positive selection plasmid that encodes a target site+PAM of interest. Generally 50,000 clones were screened to obtain between 50-100 survivors. The PI domains of surviving clones were subcloned into fresh backbone plasmid and re-tested in the positive selection. Clones that had greater than 10% survival in this secondary screen for activity were sequenced. Mutations observed in the sequenced clones were chosen for further assessment based on their frequency in surviving clones, type of substitution, proximity to the PAM bases in the SpCas9/sgRNA crystal structure (PDB:4UN3)¹⁴, and (in some cases) activities in a human cell-based EGFP disruption assay.

[0114] Bacterial-Based Site-Depletion Assay for Profiling Cas9 PAM Specificities

[0115] Competent *E. coli* BW25141(λ DE3) containing a Cas9/sgRNA expression plasmid were transformed with negative selection plasmids harboring cleavable or non-cleavable target sites. Following a 60 minute recovery in SOB media, transformations were plated on LB plates containing chloramphenicol+carbenicillin. Cleavage of the negative selection plasmid was estimated by calculating the colony forming units per μ g of DNA transformed (see also FIG. 13).

[0116] The negative selection was adapted to determine PAM specificity profiles of Cas9 nucleases by electroporating each randomized PAM library into *E. coli* BW25141(λ DE3) cells that already harbored an appropriate Cas9/sgRNA plasmid. Between 80,000-100,000 colonies were plated at a low density spread on LB+chloramphenicol+carbenicillin plates. Surviving colonies containing negative selection plasmids refractory to cleavage by Cas9 were harvested and plasmid DNA isolated by maxi-prep (Qiagen). The resulting plasmid library was amplified by PCR using

Phusion Hot-start Flex DNA Polymerase (New England BioLabs) followed by an Agencourt Ampure XP cleanup step (Beckman Coulter Genomics). Dual-indexed Tru-Seq Illumina deep-sequencing libraries were prepared using the KAPA HTP library preparation kit (KAPA BioSystems) from ~500 ng of clean PCR product for each site-depletion experiment. The Dana-Farber Cancer Institute Molecular Biology Core performed 150-bp paired-end sequencing on an Illumina MiSeq Sequencer.

[0117] The raw FASTQ files outputted for each MiSeq run were analyzed with a Python program to determine relative PAM depletion. The program (see Methods) operates as follows: First, a file dialog is presented to the user from which all FASTQ read files for a given experiment can be selected. For these files, each FASTQ entry is scanned for the fixed spacer region on both strands. If the spacer region is found, then the six variable nucleotides flanking the spacer region are captured and added to a counter. From this set of detected variable regions, the count and frequency of each window of length 2-6 nt at each possible position was tabulated. The site-depletion data for both randomized PAM libraries was analyzed by calculating the post-selection PAM depletion value (PPDV): the post-selection frequency of a PAM in the selected population divided by the pre-selection library frequency of that PAM. PPDV analyses were performed for each experiment across all possible 2-6 length windows in the 6 bp randomized region. The windows we used to visualize PAM preferences were: the 3 nt window representing the 2nd, 3rd, and 4th PAM positions for wild-type and variant SpCas9 experiments, and the 4 nt window representing the 3rd, 4th, 5th, 6th PAM positions for St1Cas9 and SaCas9.

[0118] Two significance thresholds for the PPDVs were determined based on: 1) a statistical significance threshold based on the distribution of dCas9 versus pre-selection library log read count ratios (see FIGS. 13C & 13D), and 2) a biological activity threshold based on an empirical correlation between depletion values and activity in human cells. The statistical threshold was set at 3.36 standard deviations from the mean PPDV for dCas9 (equivalent to a relative PPDV of 0.85), corresponding to a normal distribution two-sided p-value of 0.05 after adjusting for multiple comparisons (i.e. $p=0.05/64$). The biological activity threshold was set at 5-fold depletion (equivalent to a PPDV of 0.2) because this level of depletion serves as a reasonable predictor of activity in human cells (see also FIG. 14). The 95% confidence intervals in FIG. 14 were calculated by dividing the standard deviation of the mean, by the square root of the sample size multiplied by 1.96.

[0119] Human Cell Culture and Transfection

[0120] U2OS.EGFP cells harboring a single integrated copy of a constitutively expressed EGFP-PEST reporter gene¹⁵ were cultured in Advanced DMEM media (Life Technologies) supplemented with 10% FBS, 2 mM Gluta-Max (Life Technologies), penicillin/streptomycin, and 400 μ g/ml of G418 at 37° C. with 5% CO₂. Cells were co-transfected with 750 ng of Cas9 plasmid and 250 ng of sgRNA plasmid (unless otherwise noted) using the DN-100 program of a Lonza 4D-nucleofactor according to the manufacturer's protocols. Cas9 plasmid transfected together with an empty U6 promoter plasmid was used as a negative control for all human cell experiments. Target sites for

endogenous gene experiments were selected within 200 bp of NGG sites cleavable by wild-type SpCas9 (see FIG. 16A and Table 2).

[0121] Zebrafish Care and Injections

[0122] Zebrafish care and use was approved by the Massachusetts General Hospital Subcommittee on Research Animal Care. Cas9 mRNA was transcribed with PmeI-digested JDS246 (wild-type SpCas9) or MSP469 (VQR variant) using the mMESSAGE mMACHINE T7 ULTRA Kit (Life Technologies) as previously described²¹. All sgRNAs in this study were prepared according to the cloning-independent sgRNA generation method²⁴. sgRNAs were transcribed by the MEGAscript SP6 Transcription Kit (Life Technologies), purified by RNA Clean & Concentrator-5 (Zymo Research), and eluted with RNase-free water.

[0123] sgRNA- and Cas9-encoding mRNA were co-injected into one-cell stage zebrafish embryos. Each embryo was injected with ~2-4.5 nL of solution containing 30 ng/4 gRNA and 300 ng/μL Cas9 mRNA. The next day, injected embryos were inspected under a stereoscope for normal morphological development, and genomic DNA was extracted from 5 to 9 embryos.

[0124] Human Cell EGFP Disruption Assay

[0125] EGFP disruption experiments were performed as previously described¹⁶. Transfected cells were analyzed for EGFP expression ~52 hours post-transfection using a Fortessa flow cytometer (BD Biosciences). Background EGFP loss was gated at approximately 2.5% for all experiments (graphically represented as a dashed red line).

[0126] T7E1 Assay, Targeted Deep-Sequencing, and GUIDE-Seq to Quantify Nuclease-Induced Mutation Rates

[0127] T7E1 assays were performed as previously described for human cells¹⁵ and zebrafish²¹. For U2OS. EGFP human cells, genomic DNA was extracted from transfected cells ~72 hours post-transfection using the Agencourt DNAdvance Genomic DNA Isolation Kit (Beckman Coulter Genomics). Target loci from zebrafish or human cell genomic DNA were amplified using the primers listed in Table 1. Roughly 200 ng of purified PCR product was denatured, annealed, and digested with T7E1 (New England BioLabs). Mutagenesis frequencies were quantified using a Qiaxcel capillary electrophoresis instrument (Qiagen), as previously described for human cells¹⁵ and zebrafish²¹.

[0128] For targeted deep-sequencing, previously characterized on- and off-target sites (Tsai et al., Nat Biotechnol 33, 187-197 (2015); Fu et al., Nat Biotechnol 31, 822-826 (2013); Fu et al., Nat Biotechnol 32, 279-284 (2014)) were amplified using Phusion Hot-start Flex with the primers listed in Table 1. Genomic loci were amplified for a control

condition (empty sgRNA), wild-type, and D1135E SpCas9. An Agencourt Ampure XP cleanup step (Beckman Coulter Genomics) was performed prior to pooling ~500 ng of DNA from each condition for library preparation. Dual-indexed Tru-Seq Illumina deep-sequencing libraries were generated using the KAPA HTP library preparation kit (KAPA Bio-Systems). The Dana-Farber Cancer Institute Molecular Biology Core performed 150-bp paired-end sequencing on an Illumina MiSeq Sequencer. Mutation analysis of targeted deep-sequencing data was performed as previously described (Tsai et al., Nat Biotechnol 32, 569-576 (2014)). Briefly, Illumina MiSeq paired end read data was mapped to human genome reference GRCh37 using bwa (Li et al., Bioinformatics 25, 1754-1760 (2009)). High-quality reads (quality score >=30) were assessed for indel mutations that overlapped the target or off-target sites. 1-bp indel mutations were excluded from the analysis unless they occurred within 1-bp of the predicted breakpoint. Changes in activity at on- and off-target sites comparing D1135E versus wild-type SpCas9 were calculated by comparing the indel frequencies from both conditions (for rates above background control amplicon indel levels).

[0129] GUIDE-seq experiments were performed as previously described (Tsai et al., Nat Biotechnol 33, 187-197 (2015)). Briefly, phosphorylated, phosphorothioate-modified double-stranded oligodeoxynucleotides (dsODNs) were transfected into U2OS cells with Cas9 nuclease along with Cas9 and sgRNA expression plasmids, as described above. dsODN-specific amplification, high-throughput sequencing, and mapping were performed to identify genomic intervals containing DSB activity. For wild-type versus D1135E experiments, off-target read counts were normalized to the on-target read counts to correct for sequencing depth differences between samples. The normalized ratios for wild-type and D1135E SpCas9 were then compared to calculate the fold-change in activity at off-target sites. To determine whether wild-type and D1135E samples for GUIDE-seq had similar oligo tag integration rates at the intended target site, restriction fragment length polymorphism (RFLP) assays were performed by amplifying the intended target loci with Phusion Hot-Start Flex from 100 ng of genomic DNA (isolated as described above) using primers listed in Table 1. Roughly 150 ng of PCR product was digested with 20 U of NdeI (New England BioLabs) for 3 hours at 37° C. prior to clean-up using the Agencourt Ampure XP kit. RFLP results were quantified using a Qiaxcel capillary electrophoresis instrument (Qiagen) to approximate oligo tag integration rates. T7E1 assays were performed for a similar purpose, as described above.

Software - for analyzing PAM depletion MiSeq data

Run in the command prompt (in the directory containing the file) using the command "python PAM_depletion.py"

```
-----
import numpy as np
import pandas as pd
import glob
import fnmatch
import os
from collections import Counter
from Bio.Seq import Seq
from Bio import SeqIO
import itertools
import re
```

-continued

Software - for analyzing PAM depletion MiSeq data

```

from pandas import ExcelWriter
import Tkinter, tkFileDialog
__author__ = "Ved V. Topkar"
__version__ = "1.0"
"""
IUPAC_notation_regex describes a mapping between certain base characters and the
relevant regex string
(Useful for parsing out ambiguous base strings)
"""
IUPAC_notation_regex = {
    'N' : '[ATCG]',
    'Y' : '[CT]',
    'R' : '[AG]',
    'W' : '[AT]',
    'S' : '[CG]',
    'A' : 'A',
    'T' : 'T',
    'C' : 'C',
    'G' : 'G'
}
def ambiguous_PAMs(length):
    """
    Given an inputted length, return a list of strings describing all possible PAM
sequences
    NOTE: Returned strings include ambiguous base characters
    """
    permutations = itertools.product(['N', 'A', 'T', 'C', 'G'], repeat=length)
    PAMs = [ ]
    for item in permutations:
        PAMs.append(''.join(item))
    return PAMs
def unambiguous_PAMs(length):
    permutations = itertools.product(['A', 'T', 'C', 'G'], repeat=length)
    PAMs = [ ]
    for item in permutations:
        PAMs.append(''.join(item))
    return PAMs
def regex_from_seq(seq):
    """
    Given a sequence with ambiguous base characters, returns a regex that matches
for
the explicit (unambiguous) base characters
    """
    regex = ''
    for c in seq:
        regex += IUPAC_notation_regex[c]
    return regex
def regex_match_count(regex, list_of_counts):
    """
    Given a list of strings and a regex, return the number of strings in the list
that the regex matches.
    """
    c = 0
    for item in list_of_counts:
        if re.search(regex, item):
            c += 1
    return c
def tabulate_substring_frequencies(pams, indices):
    """
    Given a list of raw pams and substring indices, tabulates the frequency of
tabulate_substring_frequencies
    RETURNS a Pandas Series
    """
    base_PAMs = unambiguous_PAMs(indices[1] - indices[0])
    tmp_PAMs = Counter([pam[indices[0]:indices[1]] for pam in pams])
    c = Counter( )
    for base_PAM in base_PAMs:
        c[base_PAM] = tmp_PAMs[base_PAM]
    PAMs = pd.Series(c)
    PAMs.sort(ascending=False)
    excel_PAMs = pd.DataFrame( )
    excel_PAMs[ 'PAM' ] = PAMs.index
    excel_PAMs[ 'Count' ] = PAMs.values
    excel_PAMs[ 'Frequency' ] = PAMs.values.astype(float)/sum(PAMs.values)

```


-continued

Software - for analyzing PAM depletion MiSeq data

```

return excel_PAMs
def generate_raw_PAM_counts(filepaths, targetsites, PAM_length):
    """
    Here, we get all of our relevant PAM sequences from the inputted files
    by searching for the targetsites and looking at the flanking region
    """
    reverse_target_sequences = {targetsite:
    str(Seq(targetsites[targetsite]).reverse_complement()) for targetsite in
    targetsites}
    all_pams = {targetsite: [] for targetsite in targetsites}
    # Iterate through each file and collect the PAMs of each sequence
    # Checks both forward and reverse reads
    for filepath in filepaths:
        print 'Scanning file: ' + os.path.basename(filepath)
        pams = []
        records = SeqIO.parse(filepath, filepath.split('.')[-1])
        for record in records:
            seq = str(record.seq)
            for targetsite in targetsites:
                target_seq = targetsites[targetsite]
                target = seq.find(targetsites[targetsite])
                if target > -1:
                    index = target + len(target_seq)
                    all_pams[targetsite].append(seq[index:index + PAM_length])
                else:
                    target = seq.find(reverse_target_sequences[targetsite])
                    if target > -1:
                        index = target
                        all_pams[targetsite].append(str(Seq(seq[index -
PAM_length:index]).reverse_complement()))
            return all_pams
def analyze_PAM_depletion_data(filepaths, targetsites, PAM_length=3):
    """
    Given a directory that contains a given file extension and a target sequence,
    do the entire PAM depletion analysis
    """
    # Make sure that dirnames and target sequences are inputted
    if filepaths is None:
        raise Exception('Please specify a directory name')
    if targetsites is None:
        raise Exception('Please specify a target sequence')
    if PAM_length is None or PAM_length < 3:
        raise Exception('Please enter a valid PAM length')
    all_pams = generate_raw_PAM_counts(filepaths, targetsites, PAM_length)
    letters = ['A', 'T', 'C', 'G']
    all_counters = {targetsite: Counter(all_pams[targetsite]) for targetsite in
    targetsites}
    for targetsite in targetsites:
        pams = all_pams[targetsite]
        base_counters = [Counter() for x in range(PAM_length)]
        for pam in pams:
            for i, c in enumerate(pam):
                base_counters[i][c] += 1
        raw_PAM_counts = pd.Series(all_counters[targetsite])
        raw_PAM_counts.sort(ascending=False)
        raw_counts_df = pd.DataFrame()
        raw_counts_df['PAM'] = raw_PAM_counts.index
        raw_counts_df['Count'] = raw_PAM_counts.values
        single_base_counts = pd.DataFrame(base_counters)
        single_base_frequencies =
single_base_counts.divide(single_base_counts.sum(axis=1).ix[0])
    # Prepare substring counts and frequencies
    writer = ExcelWriter('out/' + os.path.basename(filepath).split('.')[-1][0] +
    '_' + targetsite + '.xlsx')
    single_base_counts.to_excel(writer, 'Single Base Counts')
    single_base_frequencies.to_excel(writer, 'Single Base Frequencies')
    raw_counts_df.to_excel(writer, 'Raw PAM Counts')
    # Designate which windows should be analyzed and name them
    settings = {
        'XXXXNNN': [0,3],
        'NXXXXNN': [1,4],
        'NNXXXXN': [2,5],
        'NNNXXXX': [3,6],
        'XXXXNNN': [0,4],
    }

```

-continued

```

Software - for analyzing PAM depletion MiSeq data


---


'NXXXXXN': [1,5],
'NNXXXX': [2,6],
'XXNNNN': [0,2],
'NXNNNN': [1,3],
'NNXXNN': [2,4],
'NNNXXN': [3,5],
'NNNNXX': [4,6],
'XXXXXXN': [0,5],
'NXXXXX': [1,6],
'XXXXXX': [0,6],
}
for item in settings:
    df = tabulate_substring_frequencies(pams, settings[item])
    df.to_excel(writer, item)
writer.save()
print 'Saved excel output for ' + targetsite
if __name__ == "__main__":
    # Display the filepicker, accepting only FASTQ files
    root = Tkinter.Tk()
    root.withdraw()
    file_paths = tkFileDialog.askopenfilenames(parent=root, title='Choose FASTQ
files', filetypes=[("FastQ files", "*.fastq")])
    # Describe the targetsite(s) to search for
    targetsites = {'EGFP site 1': 'GTCGCCCTCGAACTTCACCT'}
    # Run the analysis on the inputted filepaths and targetsite for a given
variable nucleotide region length
    analyze_PAM_depletion_data(file_paths, targetsites, PAM_length=6)

```

Example 1

[0130] One potential solution to address targeting range limitations would be to engineer Cas9 variants with novel PAM specificities. A previous attempt to alter PAM specificity utilized structural information about base-specific SpCas9-PAM interactions to mutate arginine residues (R1333 and R1335) that contact guanine nucleotides at the second and third PAM positions, respectively (Anders et al., *Nature* 513, 569-573 (2014)). Substitution of both arginines with glutamines (whose side-chains might be expected to interact with adenines) failed to yield SpCas9 variants that could cleave targets harboring the expected NAA PAM in vitro (Anders et al., *Nature* 513, 569-573 (2014)). Using a human cell-based U2OS EGFP reporter gene disruption assay in which nuclease-induced indels lead to loss of fluorescence (Reyon et al., *Nat Biotechnol* 30, 460-465 (2012); Fu et al., *Nat Biotechnol* 31, 822-826 (2013)), we confirmed that an R1333Q/R1335Q SpCas9 variant failed to efficiently cleave target sites with NAA PAMs (FIG. 1A). Additionally, we found that single R1333Q and R1335Q SpCas9 variants each failed to efficiently cleave target sites with their expected NAG and NGA sites, respectively (FIG. 1A). We therefore reasoned that re-engineering PAM specificity might require additional mutations at positions other than R1333 and R1335. For example, available structural information shows that K1107 and S1136 make direct and indirect minor groove contacts to the second and third bases in the PAM, respectively (Anders et al., *Nature* 513, 569-573 (2014)). Therefore, it is plausible that additional alterations at or near these positions might be needed to alter PAM specificity.

[0131] To identify additional positions that might be critical for modifying PAM specificity, we adapted a bacterial selection system previously used to study properties of homing endonucleases (hereafter referred to as the positive

selection) (Chen & Zhao, *Nucleic Acids Res* 33, e154 (2005); Doyon et al., *J Am Chem Soc* 128, 2477-2484 (2006)). In our adaptation of this system, Cas9-mediated cleavage of a positive selection plasmid encoding an inducible toxic gene enables cell survival, due to subsequent degradation and loss of the linearized plasmid (FIG. 1B and FIG. 12A). After establishing that SpCas9 can function in the positive selection system, we tested both wild-type and the R1335Q variant for their ability to cleave a selection plasmid harboring a target site with an NGA PAM and failed to observe survival, as expected (FIG. 12A). To screen for gain-of-function mutations, we generated libraries of wild-type and R1335Q SpCas9 bearing randomly mutagenized PAM-interacting domains (amino acid positions 1097-1368) with a mean rate of 5.2 mutations per kilobase (FIG. 12B and Methods). These libraries were introduced into bacteria with a positive selection plasmid containing a target site with an NGA PAM and plated on selective medium. Sequences of surviving clones from the R1335Q-based library revealed that the most frequent substitutions in addition to the pre-existing R1335Q mutation were D1135V/Y/N/E and T1337R (Table 3). We obtained fewer survivors with the wild-type SpCas9-based library selection but the sequences of these clones also included D1135V/Y/N and R1335Q mutations. We next assembled and tested SpCas9s bearing all possible single, double, and triple combinations of the D1135V/Y/N/E, R1335Q, and T1337R mutations using the human cell-based EGFP disruption assay. This analysis showed that SpCas9 variants with substitutions at all three positions displayed the highest activities on an NGA PAM, but also the lowest activities on an NGG PAM (FIG. 1C). We chose two SpCas9 variants, D1135V/R1335Q/T1337R and D1135E/R1335Q/T1337R (hereafter referred to as the VQR and EQR SpCas9 variants, respectively), because they possessed the greatest discrimination between NGA and NGG PAMs (FIG. 1C), for further characterization.

[0132] To assess the global PAM specificity profiles of our novel SpCas9 variants, we used a bacterial-based negative selection system (FIG. 1D and FIG. 13A). Previous studies have used similar types of selection systems to identify the cleavage site preferences of Cas9 nucleases (Jiang et al., *Nat Biotechnol* 31, 233-239 (2013); Esvelt et al., *Nat Methods* 10, 1116-1121 (2013)). In our version of this assay (which we refer to as the site-depletion assay), a library of plasmids bearing randomized 6 bp sequences placed adjacent to a protospacer is tested for cleavage by a Cas9/sgRNA complex in *E. coli* (FIG. 13B). Plasmids with protospacer-adjacent sequences resistant to cleavage by a Cas9/sgRNA complex enable cell survival due to the presence of an antibiotic resistance gene, whereas plasmids bearing cleavable sequences are degraded and therefore depleted from the library (FIG. 13B). High-throughput sequencing of 100,000 non-targetable sequences enabled us to calculate a post-selection PAM depletion value (PPDV) for any given PAM. The PPDV of a PAM (or group of PAMs) is defined as the frequency of that PAM in the post-selection population divided by its frequency in the pre-selection library. This quantitative value provides an estimate of Cas9 activity on that PAM. Profiles obtained with catalytically inactive Cas9 (dCas9) on two randomized PAM libraries (each with a different protospacer) enabled us to define what represents a statistically significant change in PPDV for any given PAM or group of PAMs (FIG. 13C). We then validated our site-depletion assay by demonstrating that the PPDVs for wild-type SpCas9 obtained with the two randomized PAM libraries recapitulated its previously described profile of targetable PAMs (Jiang et al., *Nat Biotechnol* 31, 233-239 (2013)) (FIG. 1E).

[0133] Using the site-depletion assay, we obtained PAM specificity profiles for the VQR and EQR SpCas9 variants using the two randomized PAM libraries. The VQR variant strongly depleted sites bearing NGAN and NGCG PAMs, and more weakly NGGG, NGTG, and NAAG PAMs (FIG. 1F). In contrast, the EQR variant strongly depleted NGAG PAMs and more weakly NGAT, NGAA, and NGCG PAMs (FIG. 1F), demonstrating a potentially more limited targeting range relative to the VQR variant. To test whether PAMs identified by the site-depletion assay could also be recognized in human cells, we assessed cleavage by the VQR and EQR SpCas9 variants on target sites using the EGFP disruption assay. The VQR variant robustly cleaved sites in EGFP bearing NGAN PAMs (with relative efficiencies NGAG>NGAT=NGAA>NGAC), and also sites bearing NGCG, NGGG, and NGTG PAMs with generally lower efficiencies (FIG. 1G). The EQR variant also recapitulated its preference for NGAG and NGNG PAMs over the other NGAN PAMs in human cells, again all at lower activities than with the VQR variant (FIG. 1G). Collectively, these results in human cells strongly mirror what was observed with the bacterial site-depletion assay (FIG. 14) and suggested that PPDVs of 0.2 (representing a five-fold depletion) in the bacterial assay provide a reasonable predictive threshold for activity in human cells (FIG. 14).

[0134] We next sought to extend the generalizability of our engineering strategy by attempting to identify SpCas9 variants capable of recognizing an NGC PAM. We first designed Cas9 mutants bearing amino acid substitutions of R1335 that might be expected to interact with a cytosine (D, E, S, or T) and found no activity on an NGC PAM site using the positive selection system. We then randomly mutagen-

ized the PAM-interacting domain of each of these singly substituted SpCas9 variants but still failed to obtain surviving colonies in positive selections. Because the T1337R mutation had increased the activities of our VQR and EQR SpCas9 variants (FIG. 1C), we combined this mutation with R1335 substitutions of A, D, E, S, T, or V, and again randomly mutagenized their PAM-interacting domains. Selections using two of these six mutagenized libraries (bearing pre-existing R1335E/T1337R and R1335T/T1337R substitutions) yielded surviving colonies harboring a variety of additional mutations (Table 3). Characterization of various selected clones using both bacterial and human cell-based assays suggested that substitutions at four positions in particular (D1135V, G1218R, R1335E, and T1337R) appeared to be important for cleavage of NGC PAMs. Assembly and testing of all potential single, double, triple, and quadruple combinations of these mutations using the EGFP disruption assay established that the quadruple VRER variant displayed the highest activity on an NGCG PAM and minimal activity on an NGGG PAM (FIG. 1H). Analysis of the VRER variant using the site-depletion assay revealed it to be highly specific for NGCG PAMs (FIG. 1I). Consistent with this result, EGFP disruption assays performed in human cells with the VRER variant revealed efficient cleavage of sites with NGCG PAMs, greatly decreased and inconsistent cleavage of sites with NGCA, NGCC, and NGCT PAMs, and essentially no activity on sites with NGAG, NGTG, and NGGG PAMs (FIG. 1J).

[0135] To demonstrate directly that our VQR and VRER SpCas9 variants can enable targeting of sites not currently modifiable by wild-type SpCas9, we tested their activities on endogenous genes in zebrafish embryos and human cells. In single cell zebrafish embryos, we found that the VQR variant could efficiently modify endogenous gene sites bearing NGAG PAMs with mean mutagenesis frequencies of 20 to 43% (FIG. 2A) and that the indels originated at the predicted cleavage sites (FIG. 15). In human cells, we found that the VQR variant robustly modified 16 sites across four different endogenous genes that harbored NGAG, NGAT, and NGAA PAMs (range of 6 to 53%, mean of 33%; FIG. 2B and FIG. 16A). Importantly, we verified that wild-type SpCas9 was unable to efficiently alter most of the same sites with NGAG and NGAT PAMs in zebrafish and human cells (FIGS. 2A and 2C), yet was able to efficiently modify nearby sites bearing NGG PAMs (FIG. 16B). Similarly, when examining VRER variant activity at nine sites with NGCG PAMs across three endogenous human genes, we also observed robust mean disruption frequencies (range of 5 to 36%, mean of 21%; FIG. 2D). Consistent with our site-depletion data (FIGS. 1E & 1F), the VQR variant altered NGCG PAM sites efficiencies similar to that observed with the VRER variant, while wild-type SpCas9 was unable to do so (FIG. 2D). Computational analysis of the reference human genome sequence shows that the addition of our VQR and VRER SpCas9 variants doubles the range of potential target sites compared with what was previously possible with only wild-type SpCas9 (FIG. 2E). Taken together, these results demonstrate that our engineered SpCas9 variants expand the targeting range of SpCas9 by enabling modification of previously inaccessible endogenous sites in zebrafish embryos and human cells.

[0136] To determine the genomewide specificity of our VQR and VRER SpCas9 nucleases, we used the recently described GUIDE-seq (Genome-wide Unbiased Identifica-

tion of Double-stranded breaks Enabled by sequencing) method¹⁰ to profile off-target cleavage events of these SpCas9 variants in human cells. We profiled the genome-wide activities of the VQR and VRER SpCas9 variants using a total of 13 different sgRNAs (eight for VQR and five for VRER from FIGS. 2B and 2D, respectively), which we had shown could induce high efficiencies of modification at their intended on-target sites. These GUIDE-seq experiments yielded a number of important observations: The numbers of off-target DSBs induced by our SpCas9 variants in human cells are comparable to (or, in the case of the VRER variant, perhaps even better than) what has been previously observed with wild-type SpCas9 (FIG. 2F). We note that the high genome-wide specificities observed with VRER might result both from its restricted specificity for NGCG PAMs and perhaps from the relative depletion of sites with NGCG PAMs in the human genome (FIG. 2E)²¹. Additionally, the off-target sites observed generally possess the expected PAM sequences predicted by our site-depletion experiments, including some tolerance for PAMs “shifted” 3' by one base (compare PAMs from FIGS. 1F and 1I with those in the sites of FIG. 17). Finally, the position and numbers of mismatches found in the off-target sites of our VQR and VRER SpCas9 variants (FIG. 17) are similar in their distributions to what we previously observed with wild-type SpCas9 for sgRNAs targeted to non-repetitive sequences¹⁰.

[0137] Previous studies have shown that imperfect PAM recognition by SpCas9 can lead to recognition of unwanted sites that contain non-canonical NAG, NGA, and other PAMs in human cells (Hsu et al., Nat Biotechnol 31, 827-832 (2013); Tsai et al., Nat Biotechnol 33, 187-197 (2015); Jiang et al., Nat Biotechnol 31, 233-239 (2013); Mali et al., Nat Biotechnol 31, 833-838 (2013); Zhang et al., Sci Rep 4, 5405 (2014)). Therefore, we were interested in exploring if mutations at or near residues that mediate PAM-interaction might improve SpCas9 PAM specificity. While engineering the VQR variant we had noticed that a D1135E SpCas9 mutant appeared to better discriminate between a canonical NGG PAM and a non-canonical NGA PAM compared to wild-type SpCas9 (FIG. 1C). Given this observation, we comprehensively assessed the PAM recognition profile of this D1135E variant using our site-depletion assay. This experiment revealed a decrease in depletion of non-canonical NAG, NGA, and NNGG PAMs with D1135E SpCas9 relative to wild-type SpCas9 (FIG. 3A). Interestingly, this effect was more prominent for one of the two protospacers we used, suggesting that the impact of the D1135E substitution on non-canonical PAM recognition may vary to some degree in a protospacer-dependent manner. Importantly, we did not observe the emergence of any new non-canonical PAM specificities.

[0138] We next tested whether the improved PAM specificity of D1135E SpCas9 also could be observed in human cells. In direct comparisons of wild-type and D1135E SpCas9 on eight target sites with non-canonical NAG or NGA PAMs, we observed that these sites were consistently less efficiently cleaved by D1135E than by wild-type SpCas9 in the EGFP disruption assay (FIG. 3B, mean fold-decrease in activity of 1.94). Importantly, wild-type and D1135E SpCas9 both showed comparable activities on four EGFP reporter gene sites and six endogenous human gene sites with canonical NGG PAMs (FIGS. 3B and 3C, respectively), demonstrating that the D1135E variant does not appreciably affect cleavage of on-target sites with NGG

PAMs (mean fold-decrease in activity of 1.04 across all ten sites). Titration experiments in which we decreased the concentration of Cas9-encoding plasmid transfected into human cells revealed no substantial differences in the activities of wild-type and D1135E SpCas9 when they were targeted to the same sites (FIG. 3D), implying that the increased specificity observed with the D1135E variant is not simply the result of protein destabilization.

[0139] To more directly assess whether the introduction of D1135E could reduce off-target cleavage effects of SpCas9, we used deep-sequencing to compare mutation rates induced by wild-type and D1135E SpCas9 on 25 previously known off-target sites of three different sgRNAs (Hsu et al., Nat Biotechnol 31, 827-832 (2013); Tsai et al., Nat Biotechnol 33, 187-197 (2015); Fu et al., Nat Biotechnol 31, 822-826 (2013)). These 25 sites included off-target sites with various mismatches in the spacer sequence and both canonical NGG and non-canonical PAMs (FIG. 3E). The results of these deep-sequencing experiments revealed that the D1135E variant showed reduced mutation frequencies at 19 of the 22 off-target sites with activity above background indel rates, relative to the mutation frequency observed at the three on-target sites (FIGS. 3E & 3F). Interestingly, these reduced off-target mutation frequencies were observed at many sites with a canonical PAM, suggesting that the gain in specificity with D1135E is not restricted only to sites with non-canonical PAMs. To assess the improvements in specificity associated with D1135E on a genome-wide scale, we performed GUIDE-seq experiments using wild-type and D1135E SpCas9 with three different sgRNAs (two of which were previously known to have off-target sites with canonical and non-canonical PAMs (Hsu et al., Nat Biotechnol 31, 827-832 (2013); Tsai et al., Nat Biotechnol 33, 187-197 (2015); Fu et al., Nat Biotechnol 31, 822-826 (2013)). We observed a generalized improvement in genome-wide specificity when using the D1135E SpCas9 variant compared with wild-type SpCas9 (FIG. 3G). For all three sgRNAs we tested, these improvements in specificity were observed at off-target sites that contained mismatched spacers with canonical or non-canonical PAMs (FIG. 18). Importantly, these GUIDE-seq experiments demonstrated that the introduction of the D1135E mutation does not increase the number of off-target effects induced by SpCas9. Collectively, these results show that the D1135E substitution can increase the global specificity of SpCas9.

[0140] Although all of the experiments described above were performed with SpCas9, there are many Cas9 orthologues from other bacteria that could make attractive candidates for characterizing and engineering Cas9s with novel PAM specificities (Fonfara et al., Nucleic Acids Res 42, 2577-2590 (2014); Ran et al., Nature 520, 186-191 (2015)). To explore the feasibility of doing this, we determined whether two smaller-size orthologues, *Streptococcus thermophilus* Cas9 from the CRISPR1 locus (St1Cas9) (Deveau et al., J Bacteriol 190, 1390-1400 (2008); Horvath et al., J Bacteriol 190, 1401-1412 (2008)) and *Staphylococcus aureus* (SaCas9) (Hsu et al., Cell 157, 1262-1278 (2014); Ran et al., Nature 520, 186-191 (2015)), might also function in our bacterial selection assays. While the PAM of St1Cas9 has previously been characterized as NNAGAA (SEQ ID NO:3) (Esvelt et al., Nat Methods 10, 1116-1121 (2013); Fonfara et al., Nucleic Acids Res 42, 2577-2590 (2014); Deveau et al., J Bacteriol 190, 1390-1400 (2008); Horvath et al., J Bacteriol 190, 1401-1412 (2008)), our attempts to

bioinformatically derive the SaCas9 PAM using a previously described approach (Fonfara et al., *Nucleic Acids Res* 42, 2577-2590 (2014)) failed to yield a consensus sequence (data not shown). Therefore, we used our site-depletion assay to determine the PAM for SaCas9 and, as a positive control, for St1Cas9. These experiments were performed using the two different protospacers and sgRNAs with two different complementarity lengths for each protospacer, resulting in four selections for each Cas9. For St1Cas9, we identified two novel PAMs in addition to the six PAMs that had been previously described (Esvelt et al., *Nat Methods* 10, 1116-1121 (2013); Fonfara et al., *Nucleic Acids Res* 42, 2577-2590 (2014); Horvath et al., *J Bacteriol* 190, 1401-1412 (2008)) (FIG. 4A and FIGS. 19C and 19D, consistent with a recent definition of SaCas9 PAM specificity (Ran et al., *Nature* 520, 186-191 (2015))). For SaCas9, there was PPDV variability among the four selections mainly due to the restricted PAM preferences observed with one protospacer. As a result, only three PAMs were depleted greater than 5-fold in all four experiments: NNGGGT (SEQ ID NO:4), NNGAAT (SEQ ID NO:6), NNGAGT (SEQ ID NO:5) (FIG. 4B). We did, however, identify many more targetable PAMs with the second protospacer library, implying that SaCas9 might recognize numerous additional PAMs (FIGS. 18C and 18D). Using PAMs identified in our site-depletion experiments (NNAGAA (SEQ ID NO:3) for St1Cas9 and NNGAGT (SEQ ID NO:5) for SaCas9), we found that both St1Cas9 and SaCas9 can function efficiently in the bacterial positive selection system (FIG. 4C), suggesting that their PAM specificities could be modified by mutagenesis and selection.

[0141] Because not all Cas9 orthologues function efficiently outside of their native context (Esvelt et al., *Nat Methods* 10, 1116-1121 (2013)), we tested whether St1Cas9 and SaCas9 can robustly cleave target sites in human cells. St1Cas9 has been previously shown to function as a nuclease in human cells but on only a few sites (Esvelt et al., 2013; Cong et al., *Science* 339, 819-823 (2013)). We assessed St1Cas9 activity on sites harboring NNAGAA (SEQ ID NO:3) PAMs using sgRNAs with variable-length complementarity regions and found high activity at three of the five target sites (FIG. 4D). For SaCas9, we observed efficient activity at eight sites harboring NNGGGT (SEQ ID NO:4) or NNGAGT (SEQ ID NO:5) PAMs (FIG. 4E). For both St1Cas9 and SaCas9 no obvious correlation between activity and length of spacer complementarity was observed (FIG. 19E). We next determined whether St1Cas9 and SaCas9 could efficiently modify endogenous loci in human cells. For St1Cas9, 7 out of 11 sites across 4 genes were disrupted efficiently as judged by T7E1 assay (1 to 25%, mean of 13%; FIG. 4F), while SaCas9 displayed somewhat more robust activities at 16 sites tested across 4 genes (1% to 37%, mean of 19%; FIG. 4G). Once again, no distinct trend was observed when considering sgRNA spacer length for St1Cas9 and SaCas9 (FIG. 19F). Collectively, our results show that St1Cas9 and SaCas9 function robustly both in our bacterial-based selection and in human cells, making them attractive candidates for engineering additional SpCas9 variants with novel PAM specificities.

TABLE 3

	SEQ ID NO:
Wild-type SpCas9? sequence from K1097-D1368? of SEQ ID NO: 1	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	aa
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	1097-
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	1368
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	of SEQ ID NO: 1
Selected mutant clones for VQR and EQR variant, sequence from K1097-D1368	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	431.
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	432.
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	433.
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	434.
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI	435.
MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLELENGRKRMLASAGELQKGNELALPSKYVNFYLYLAS	
HYEKLKGPSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF	
TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	436.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	437.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	438.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	439.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLEATLIHQSI TGLYETRIDLSQLGGD	440.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP MD FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLA SHYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	441.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	442.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFESPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPSAFKYFDTTIDRKQYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	443.
KTEVQTGGFSKESIFPKRNSDKLIARKKDWDPKKYGGLYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEIKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLASH YEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLFT LTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	444.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDKEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	445.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	446.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	447.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASARELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDATIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	448.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGS LEDNEQKQLFVEQHRHYLDEII EQISEFSKRVI LADANLDKVL SAYNKYRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	449.
KTEVQTGGFSKESILPKRNSDKLIARKKVWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERS SF EKNP ID FLEAKGYKEVKKDLI IKLPKYSLFEL ENGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPS PEDNEQKQLFVEQHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIREQAENI IHLF TLTNLGAPAAFKYFDTTIDRKQYTS TKEVLDATLIHQSI TGLYETRIDLSQLGGD	450.

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	451.
KTEVQTGRFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	452.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHL FTLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	453.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	454.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHL FTLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLVGD	455.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	456.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	457.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	458.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	459.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	460.
ETEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAFSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	461.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSILVVAKEKGSKKLKSVKDLLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	462.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSILVVAKEKGSKKLKSVKELLGIT NMERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAEELQKGNELALPSKYVNFLYLA SHYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHL FTLTNLGAPAAFKYFDATIDRKQYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	463.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSILVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIIYRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	464.

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVANVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKQYTSKEVLDATLIHQSI TGLYETRIDLSQLGGD	465.
Selected mutant clones for VRER variant, sequence from K1097-D1368	
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASARELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQFFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHL FTLTNLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	466.
KTEMQTGGFSKESVLPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKVEKGSKRLKSVKELLGI TIMERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGVLQKGNELALPSKYVNFLYL SHYEKLKGPSDDNEQKRLFVEQHKHYLDEII EQISEFSKRVILADANRDKVL SAYNKHRDKPIREQAENIIH LFTLTNLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	467.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSKDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	468.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGVLQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	469.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKQYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	470.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASARELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	471.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKDEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTKLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	472.
KTEVQTGGFSKESILPKRNSDKV IARKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	473.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGVLQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	474.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEIKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLASH YEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKTIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	475.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHHSITGLYETRIDLSQLGGD	476.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADGNLDKVL SAYNKHRDKPIREQAENIIHL FTLTNLGAPAAFKYFDDTTIDRKEYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	477.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKVEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYNLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTDLGAPAAFKYFDDTTIDRKQYRSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	478.

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	479.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPKVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKQYRSTKEVLDTLIHQSI TGLYETRIDMSQLGGD	480.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	481.
KTEBQTGGFSKESIHPKRNSDKLIARKKDWDPKKYGGFHSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNMHRDKPIREQAENIIHL FTLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLEGD	482.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHL FTLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	483.
KTEVQTGRFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	484.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKVKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGVLQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	485.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HFEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	486.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQPKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	487.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFVSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	488.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	489.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGVLQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	490.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	491.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFYSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDDKVL SAYNKHDKPIREQAENIIHLF TLTNLGAPAAFKYFDTTIDRKEYRSTKEVLDTLIHQSI TGLYETRIDLSQLGGD	492.

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADPNLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	493.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	494.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	495.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVIELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	496.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	497.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKTKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVIKDFIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLASH YEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLFT LTNLGAPAAFKYFDDTTIDRKQYRSPKEVL DATLIHQSI TGLYETRIDLSQLGGD	498.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	499.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVPVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELSGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKEYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	500.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKNKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	501.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHL FTLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	502.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLMIKLPKYSLFELKNGRKRMLASAGELQKGNELALPSKYVNFLYLA SHYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHL FTLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHHSI TGLYETRIDLSQLGGD	503.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASARELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	504.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKRDIIKLPKYSLFELKNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	505.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEKGSKKLKSVKELLGITI MERSSFEKNPIDFLEAKGYKEVKKDLIIKLPKYSLFELNGRKRMLASAGELQKGNELALPSKYVNFLYLAS HYEKLKGPSPEDNEQKQLFVEQHKHYLDEII EQISEFSKRVILADANLDKVL SAYNKHRDKPIREQAENIIHLF TLTNLGAPAAFKYFDDTTIDRKTYRSTKEVL DATLIHQSI TGLYETRIDLSQLGGD	506.

TABLE 3-continued

	SEQ ID NO:
KTEVQTGGFSKESILPKGNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKEGKSKKLKSVKELLGITI MERSSEKPNIDFLEAKGYKEVKKDLIIKLPKYSLEFLENGRKRMLASAGELQKGNELALPSKYVNFLYLASH HYEKLKGGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLF LTNLGAPAAPKYFDTTIDRKYRSTKEVLDTLIHQSIITGLYETRIDLSQLGGD	507.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKDLLGITI MERSSEKPNIDFLEAKGYKEVKKDLIIKLPKYSLEFLENGRKRMLASARELQKGNELALPSKYVNFLYLASH SHYEKLKGGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIVH LFTLTNLGAPAAPKYFDTTIDRKYRSTKEVLDTLIHQSIITGLYETRIDLSQLGGD	508.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFNSPTVAYSVLVVAKEGKSKKLKSVKELLGITI MERSSEKPNIDFLEAKGYKEIKKDLIIKLPKYSLEFLENGRKRMLASAGELQKGNELALPSKYVNFLYLASH YEKLKGGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKHRDKPIREQAENIIHLFT LTNLGAPAAPKYFDTTIDRKYRSTKEVLDTLIHQSIITGLYETRIDLSQLGGD	509.
KTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVAYSVLVVAKEGKSKKLKSVKELLGITI MERSSEKPNIDFLEAKGYKEVKKDLIIKLPKYSLEFLENGRKRMLASARELQKGNELALPSKYVNFLYLASH HYEKLKGGSPEDNEQKQLFVEQHKHYLDEIIIEQISEFSKRVIADANLDKVL SAYNKDRDKPIREQAENIIHLFT LTNLGAPAAPKYFDTTIDRKYRSTKEVLDTLIHQSIITGLYETRIDLSQLGGD	510.

Example 2

Engineering the PAM Specificity of *Staphylococcus aureus* Cas9

[0142] Because we knew what residues of *Streptococcus pyogenes* Cas9 (SpCas9) were important for PAM recognition (R1333 and R1335), we generated an alignment of Cas9 orthologues to look for homologous residues in the PAM-interacting domain (PI domain) of *Staphylococcus aureus* Cas9 (SaCas9) (see FIG. 6). We and others have previously shown that the PAM of SaCas9 is NNGRRT (SEQ ID NO:46) (where N is any nucleotide, and R is an A or G). The preference for a G at the 3rd position of the PAM appeared to be the most strict requirement based on our data, so we hypothesized that positively charged residues like lysine (K) or arginine (R) might be mediating that interaction. As shown in FIG. 6, there are a number of candidate residues in SaCas9 in the homologous region to R1333 and R1335 of SpCas9, including K1101, R1012, R1015, K1018, and K1023.

[0143] We generated alanine (A) and glutamine (Q) substitutions at these five positions to determine if the mutant clones could still cleave a site containing the canonical NNGRRT PAM (SEQ ID NO:46), or possibly cleave the previously non-targetable PAM of NNARRT (SEQ ID NO:43) (FIG. 7). We utilized our bacterial assay (described in the previous patent application), where activity of Cas9 can be visualized by survival of bacterial colonies when plated under a selective condition. The relative activity of Cas9 can be quantified by calculating the ratio of bacterial colonies growing on the selective versus the non-selective media. In FIG. 7, we show that only the R1015A and R1015Q mutations affect the ability of SaCas9 to recognize a canonical NNGAGT (SEQ ID NO:5) PAM, while no mutations enable targeting of NNARRT (SEQ ID NO:43) PAMs (NNAAGT (SEQ ID NO:41) or NNAGGT (SEQ ID NO:42)). These results suggested to us that R1015 plays a role in PAM recognition by SaCas9.

[0144] We then selected randomly mutagenized either wild-type SaCas9, or the R1015Q variant and selected for altered PAM specificity clones against sites containing NNAAGT (SEQ ID NO:41) or NNAGGT (SEQ ID NO:42) PAMs (as previously described for SpCas9). We identified, re-screened, and sequenced a number of mutant clones that

could target these PAMs, with their amino acid sequences shown in FIG. 8 (and Table 6). In summary of these sequences, a number of changes appear to be very important for altering SaCas9 specificity (R1015Q, R1015H, E782K), while many other mutations may also contribute (N968K, E735K, K929R, A1021T, K1044N).

[0145] After identifying the positions and mutations essential for altering the PAM specificity of SaCas9 to NNARRT (SEQ ID NO:43), we assessed the contributions of the most abundant mutations to the specificity change by making single, double, and triple mutants combinations (Table 5). When testing these mutations against various PAMs in our positive selection (as previously described), we observed that a number of mutations allowed activity on both a canonical NNGAGT (SEQ ID NO:5) and non-canonical NNAAGT (SEQ ID NO:41) or NNAGGT (SEQ ID NO:42) PAMs, whereas the wild-type SaCas9 enzyme had very low activity on the non-canonical PAMs. Specifically, it appeared as though the triple mutations enabled a relaxed specificity at the third position of the PAM (KKQ, KKH, GKQ, GKH—named based on mutations to positions E782/N968/R1015), leading to a consensus PAM motif of NNRRRT (SEQ ID NO:45) versus the canonical NNGRRT (SEQ ID NO:46). This relaxation of the PAM requirement theoretically doubles the targeting range of SpCas9. Henceforth, variants will be named based on their identities at positions 782, 968, and 1015. For example, E782K/N968K/R1015H would be named the SaCas9 KKH variant.

TABLE 5

SaCas9? mutant activity in the bacterial screen			
mutation(s)	NNGAGT (SEQ ID NO: 5)	NNAAGT (SEQ ID NO: 41)	NNAGGT (SEQ ID NO: 42)
E782 N? 968 R1015	% activity	% activity	% activity
	100.0	21.4	15.7
Q	0.0	4.3	0.0
H	100.0	100.0	57.1

TABLE 5-continued				
SaCas9? mutant activity in the bacterial screen				
mutation(s)	NNGAGT (SEQ ID NO: 5)	NNAAGT (SEQ ID NO: 41)	NNAGGT (SEQ ID NO: 42)	
E782 N? 968R1015	% activity	% activity	% activity	
K	85.7	61.4	57.1	
G	85.7	57.1	57.1	
K	100.0	57.1	57.1	
K	Q	85.7	92.9	85.7
K	H	100.0	100.0	85.7
G	Q	71.4	85.7	71.4
G	H	100.0	85.7	85.7
K	Q	85.7	85.7	85.7
K	H	85.7	92.9	92.9
K	K	71.4	71.4	71.4
G	K	85.7	71.4	71.4
K	K	Q	100.0	100.0
K	K	H	92.9	100.0
G	K	Q	92.9	100.0
G	K	H	100.0	100.0

[0146] We next assessed two of the triple mutants in the human cell EGFP disruption assay (as previously described) to determine whether the engineered variants could target non-canonical PAMs in a human cell context (FIG. 9). Variants capable of targeting sites within the EGFP gene containing non-canonical PAMs will disrupt the EGFP coding frame, leading to loss of signal. The results revealed that both the KKQ and KKH mutants retained similar activity to wild-type SaCas9 on canonical NNGRRT (SEQ ID NO:46) PAMs, but had much higher activity on NNARRT (SEQ ID NO:43) PAMs.

[0147] Overall, we've identified mutations in SaCas9 (KKQ or KKH variants) that appear to relax the preference of the wild-type enzyme at the third position of the PAM

from a G to an R (A or G). This effectively relaxes the targeting of SaCas9 from an NNGRRT (SEQ ID NO:46) PAM constraint to an NNRRRT (SEQ ID NO:45) PAM.

[0148] Because we had successfully derived variants that could target NNARRT (SEQ ID NO:43) PAMs in human cells, we next asked the question of whether we could engineer variants with specificity for NNCRRRT (SEQ ID NO:47) or NNTRRT (SEQ ID NO:48). To do so, we first mutated R1015 to E (in the case of specifying a C at the 3rd position of the PAM) and to L or M (in the case of specifying a T at the 3rd position of the PAM), and tested these against their expected PAMs in our bacterial positive selection assay (previously described) (FIG. 10). We observed that wild-type SaCas9 could inefficiently cleave a site containing an NNCAGT (SEQ ID NO:511) PAM, that an R1015E variant had slightly better activity on the same site, and that wild-type or any of the other directed mutations did not convey activity against other PAMs (FIG. 10). This suggested that as we saw with R1015Q, other mutations would be necessary to engineer SaCas9 variants that could target NNCRRRT (SEQ ID NO:47) and NNTRRT (SEQ ID NO:48) PAMs.

[0149] For the SaCas9 evolved variants against NNARRT (SEQ ID NO:43) PAMs, the E782K and N968K mutations were necessary and essential along with the R1015(H/Q). To test whether these mutations would increase the activity of the R1015(E/L/M) variants against their expected PAM, we generated the KKE, KKL, and KKM variants. As shown in FIG. 11, the KKE, KKL, and KKM all had robust activity against their expected PAMs.

[0150] We were also curious as to whether the KKQ, KKH, KKE, KKL, or KKM variants had relaxed specificity against any nucleotide at the 3rd position of the PAM, so we interrogated a number of sites in our bacterial positive selection assay containing NNNRRT PAMs. As shown in FIG. 11, with a few exceptions nearly all of these variants can cleave all sites tested that contain NNNRRT PAMs. This indicated that they had a relaxed specificity at the 3rd position of the PAM as they can efficiently target NNNRRT sites. This is in contrast to the wild-type protein (ENR) that can only efficiently target the NNGAGT (SEQ ID NO:5) site, with very low activity on a few NNNRRT sites. In summary, the KKH (and other similar derivatives shown in FIG. 11) variant can target sites containing NNNRRT PAMs in bacteria, effectively quadrupling the targeting range of SaCas9.

[0151] Thus, the KKH variant (and some of the other variants in FIG. 6) can target NNNRRT PAMs in bacteria, effectively quadrupling the targeting range of SaCas9.

TABLE 6	
residues A652-G1053 of SaCas9	SEQ ID NO:
Wild Type SaCas9	
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKPKKERNKGYKXHAEDALIIAN	Aa
ADFIFKEWKKLDKAKVMENQMFEKQAESMPETETEQEYKEIFITPHQIKHIKDFKDYK	652-
YSHRVDKKPNRELINDTLYSTRKDDKGNLTIVNNLNGLYDKDNDKLKLINKSPEKLLMY	1053 of
HHDPQTYQKLKLIMEQYQDEKNPLYKYEETGNLYTKYSKKDNGPVIKKIKYGNKLNA	SEQ ID
HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKENYEVNSKCYEE	NO: 2.
AKKLKKSINQAEIFASFPYNNDLIKINGELRVIGVNNDLLNRIEVNMIDITYREYLENMND	
KRPRIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
Sequences of selected clones of SaCas9 variants	
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	53.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFMASFYKNDLIKFN GEL YRVIGVNNDLNRIEVNMDITYREYLENMND DKRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	54.
ATRGLMNLRSYFRVNNLDIKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IANA DFIKKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMYH HDPTQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN NITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEAK KKLKISNQAEFISSFYNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMNDKRP PHIIKRIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	55.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMNDKRP PHIIKRIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	56.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIRINGEL YRVIGVNNDLNRIEVNMDITYREYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	57.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDFK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYRGYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	58.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	59.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	60.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWFKKERNKGKXHAEDALI IAN ADFIFKEWKRLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKDNGPVIKKIKYGNKLN HLDITDDYPNSRNKVVKLSLKP YRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKISNQAEFIASFYNNDLIKINGEL YRVIGVNNDLNRIEVNMDITYREYLENMND KRPPHI IKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	61.

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVRNLDVIKKENHYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	62.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMYH HDPPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A DITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNTKCYEEAK KKKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIVKKG	63.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	64.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A LDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEA KKKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	65.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNMDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	66.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVRNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	67.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKR LDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A LDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVRNLDVIKKENYEVNSKCYEEA KKKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIK*G	68.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	69.
ATRGLMNLRLRNYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFITPHQIKHIKDFKDYK YSHRVDKKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVKKIKYYGNKLN A HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQII*KG	70.

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN AH LDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVIVKNLDVI KKENY YEVNSKCYEEA KKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMNDK RPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIVKKG	71.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMND KRPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	72.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMND KRPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	73.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDY KYSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLM YHHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN AHLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYE EAKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMN DKRPPIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	74.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN AH LDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVIVKNLDVI KKENY YEVNSKCYEEA KKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMNYK RPPQIIKT IASKTQSIKKYSSDILGNLYEVKSKKHP*IIKKG	75. and 966
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMND KRPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	76.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMND KRPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	77.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVI KKENY YEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENVNDK RPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	78.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F ITPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I KKI KYGNKLN A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGIYKFVTVKNLDVI KKENY YEVNSKCYEKA KKLKKISNQAEFIASFYNNDLIKINGELYRVI GVNNDLLNRIEVMIDITYREYLENMNDK RPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	79.

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	80.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPHIIKKG	81.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMYH HDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKIRYGNKLNAHL DITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEAK KKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDKR PPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	82.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYFENMNV KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	83.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	84.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHIMDFKDY KYSHRVDKKPNRELINDTLYSTRKDEKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLM YHHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVRNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	85.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAVSMPEIETE QEYKEIFINPHQIKHI KDFKDY KYSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKYNDKLLKL INKSPKLLM YHHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSRKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYRENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	86.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KDFKDYK YSHRVDKKPNRELINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENINGK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	87.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNNGYKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEIFI TPHQIKHI KEFKDYK YSHRVDKKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPTTYQKLKLIMEQYGDEKNPLYKYEEETGIYLT KYSKKDN GPVIKKI KYGNKLNAH LDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYGEA KKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLEIMNDKR PPQIIKTIASKTQSIKKYSTDILGNLYEVKSNKHPQIIKKG	88.

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A H LDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEA KKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	89.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	90.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNTIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKPKKHPQIIKKG	91.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIIITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHP*IIKKG	92. and 966
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P Y Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	93.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDNKKVMENQMFEEKRAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D F K YSHMVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLIY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	94.
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWIRLDDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHP*IIKKG	95. and 966
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALIVAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVINLNGLYDKDNDKLLKL INKSPKLLMYH HDPPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A H DITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEEAK KKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDKR PPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHP*IIKKG	96. and 966
ATRGLMNLRLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKXHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAESMPEIETE QEYKEI F I T P H Q I K H I K D F K D Y K YSHRVDKPKPNRKLINDTLYSTRKDDKGNTLIVNNLNGLYDKDNDKLLKL INKSPKLLMY HHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDN GPV I K K I K Y Y G N K L N A HLDITDDYPNSRNKVVKLSLKPYPFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMNDK RPPQIIKTIASKTQSIKKYSTDILGNLYEVKSKKHP*IIKKG	97. and 966

TABLE 6-continued

residues A652-G1053 of SaCas9	SEQ ID NO:
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y K E I F I T P H Q I K H I K D F K D Y KYSHRVDDKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLM YHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPDSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPHI IKT IASKTQSIKKYSTDILGNLYEVKSQKHPQIIKKG	98. and 966
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQAGSMPEI E T E Q E Y K E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLESMNDK RPPQIIKT IASKTQTIKKYSTDILGNLYEVKSKKHPQIIKKG	99.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y K E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRELINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLM YHDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA YLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPHI IKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	100.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y K E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKT IATKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	101.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y K E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIK*G	102.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y N E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLYVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	103.
ATRGLMNLRSYFRVNNLDVKVKSINGGFTSFLRRKWKFKKERNKGKHHHAEDALI IAN ADFIFKEWKKLDKAKKVMENQMFEEKQASMP E I E T E Q E Y K E I F I T P H Q I K H I K D F K D Y YSHRVDDKPNRKLINDTL YSTRKDDKGNTLIVNNLNGLYDKDNDKLLKLINKSPEKLLMY HDPQTYQKLKLIMEQYGDEKNPLYKYEEETGNYLT KYSKKDNGPVIKKI KYGNKLNA HLDITDDYPNSRNKVVKLSLKPYRFDVYLDNGVYKFVTVKNLDVIKKENYEVNSKCYEE AKLKKISNQAEFIASFYNNDLIKINGELYRVIGVNNDDLNR IEVNMIDITYREYLENMND KRPPQIIKT IASKTQSIKKYSTDILGNLYEVKSKKHPQIIKKG	104.

Methods for Example 3

[0152] The following materials and methods were used in Example 3.

Plasmids and Oligonucleotides

[0153] Oligonucleotides are listed in Table 11, sgRNA target sites are listed in Table 12, and plasmids used in this study are listed in Table 10.

[0154] Bacterial Cas9/sgRNA expression plasmids were used to express both a human codon optimized version of SaCas9 and the sgRNA, each expressed under a separate T7 promoter. Bacterial expression plasmids used in the selec-

tions were derived from BPK2101 (see Examples 1-2) while those used in the site-depletion assay were modified to express a sgRNA with a shortened repeat:anti-repeat sequence (see below). All sgRNAs in these bacterial expression plasmids included two guanines at the 5' end of the spacer sequence for proper expression from the T7 promoter.

[0155] To generate libraries of SaCas9 variants, amino acids M657-G1053 of SaCas9 were randomly mutagenized using Mutazyme II (Agilent Technologies) at a frequency of ~5.5 mutations/kilobase. Both wild-type and R1015Q SaCas9 were used as starting template for mutagenesis, resulting in two libraries with estimated complexities of greater than 6×10^6 clones.

[0156] Positive selection plasmids were assembled by ligating oligonucleotide duplexes encoding target sites into XbaI/SphI-digested p11-lacY-wtx1 (Chen, Z. & Zhao, H. A highly sensitive selection method for directed evolution of homing endonucleases. *Nucleic Acids Res* 33, e154 (2005)). For the site-depletion experiments, two separate libraries containing different spacer sequences were generated. For each library, an oligonucleotide containing 8 randomized nucleotides adjacent to the spacer sequence (in place of the PAM) was complexed with a bottom strand primer and filled in using Klenow(-exo) (refer to Table 11). The resulting product was digested with EcoRI and ligated into EcoRI/SphI-digested p11-lacY-wtx1. Estimated complexities of the two site-depletion libraries were greater than 4×10^6 clones.

[0157] For human cell experiments, human codon-optimized wild-type and variant SaCas9s were expressed from a plasmid containing a CAG promoter (Table 12). sgRNA expression plasmids (containing a U6 promoter) were generated by ligating oligonucleotide duplexes encoding the spacer sequence into BsmBI digested VVT1 (See Examples 1-2 or BPK2660 (containing the full length 120 nt crRNA: tracrRNA sgRNA or a 84 nt shortened repeat:anti-repeat version, respectively). All sgRNAs used in this study for human expression included one guanine at the 5' end of the spacer to ensure proper expression from the U6 promoter, and also used a shortened sgRNA (FIG. 37A-B) similar to that previously described (Ran, F. A. et al. In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 520, 186-191 (2015)).

Bioinformatic Analysis of Cas9 Orthologue Sequences

[0158] Similar to alignments performed in previous studies (Fonfara, I. et al. Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems. *Nucleic Acids Res* 42, 2577-2590 (2014); Ran, F. A. et al. In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 520, 186-191 (2015); Anders, C., Niewoehner, O., Duerst, A. & Jinek, M. Structural basis of PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature* 513, 569-573 (2014)), Cas9 orthologues similar to both SpCas9 and SaCas9 were aligned using ClustalW2 (ebi.ac.uk/Tools/msa/clustalw2/). The resulting phylogenetic tree and protein alignment were visualized using Geneious version 8.1.6 and ESPript (esprict.ibcp.fr/ESPript/ESPript/).

Bacterial-Based Positive Selection Assay

[0159] The bacterial positive selection assays were performed as previously described (See Examples 1-2). Briefly, Cas9/sgRNA plasmids were transformed into *E. coli* BW25141(λ DE3) (Kleinstiver et al., *Nucleic Acids Res* 38, 2411-2427 (2010)) containing a positive selection plasmid. Transformations were plated on both non-selective (chloramphenicol) and selective (chloramphenicol+10 mM arabinose) conditions. Cas9 cleavage of the selection plasmid was estimated by calculating the percent survival: (# of colonies on selective plates/# of colonies on non-selective plates) $\times 100$. To select for SaCas9 variants capable of recognizing alternative PAMs, the wild-type and R1015Q libraries with mutagenized PI domains were transformed into competent *E. coli* BW25141(λ DE3) containing positive selection plasmids with NNAAGT (SEQ ID NO:41), NNAGGT (SEQ ID NO:42), NNCAGT (SEQ ID NO:511),

NNCGGT (SEQ ID NO:512), NNTAGT (SEQ ID NO:513), or NNTGGT (SEQ ID NO:514) PAMs. Approximately 1×10^5 clones were screened by plating on selective conditions, and surviving colonies containing SaCas9 variants presumed to cleave the selection plasmid were mini-prepped (MGH DNA Core). All variants were re-screened individually in the positive selection assay, and those with greater than ~20% survival were sequenced to determine the mutations required for recognition of the alternate PAM.

Bacterial-Based Site-Depletion Assay

[0160] The site-depletion experiments were performed as previously described (See Examples 1-2). Briefly, the randomized PAM libraries were electroporated into competent *E. coli* BW25141(λ DE3) containing either wild-type, catalytically inactive (D10A/H557A), or KKH variant SaCas9/sgRNA plasmids. Greater than 1×10^5 colonies were plated on chloramphenicol/carbenicillin plates, and selection plasmids with PAMs resistant to Cas9 targeting contained within the surviving colonies were isolated by maxiprep (Qiagen). The region of the plasmid containing the spacer sequence and PAM was PCR-amplified using the primers listed in Table 11. The KAPA HTP library preparation kit (KAPA BioSystems) was used to generate a dual-indexed Tru-seq Illumina sequencing library using ~500 ng purified PCR product from each site-depletion condition prior to an Illumina MiSeq high-throughput sequencing run at the Dana-Farber Cancer Institute Molecular Biology Core. The data from the site-depletion experiments was analyzed as previously described (See Examples 1-2), with the exception that the script was modified to analyze 8 randomized nucleotides. Cas9 ability to recognize PAMs was determined by calculating the post-selection PAM depletion value (PPDV) of any given PAM: the ratio of the post-selection frequency of that PAM to the pre-selection library frequency. A control experiment using catalytically inactive SaCas9 was used to establish that a PPDV of 0.794 represents statistically significant depletion relative to the input library.

Human Cell Culture and Transfection

[0161] U2OS cells obtained from our collaborator T. Cathomen (Freiburg) and U2OS.EGFP cells harboring a single integrated copy of an EGFP-PEST reporter gene (Reyon, D. et al. FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol* 30, 460-465 (2012)) were cultured in Advanced DMEM medium (Life Technologies) with 10% FBS, penicillin/streptomycin, and 2 mM GlutaMAX (Life Technologies) at 37° C. with 5% CO₂. Cell line identities were validated by STR profiling (ATCC) and deep sequencing, and cells were tested bi-weekly for *mycoplasma* contamination. U2OS.EGFP culture medium was additionally supplemented with 400 μ g/mL G418. Cells were co-transfected with 750 ng Cas9 plasmid and 250 ng sgRNA plasmid using the DN-100 program of a Lonza 4D-nucleofactor following the manufacturer's instructions.

Human Cell EGFP Disruption Assay

[0162] EGFP disruption experiments were performed as previously described (Fu, Y. et al. High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. *Nat Biotechnol* 31, 822-826 (2013); Reyon, D. et al. FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol* 30, 460-465 (2012)). Approximately

52 hours post-transfection, a Fortessa flow cytometer (BD Biosciences) was used to measure EGFP fluorescence in transfected U2OS.EGFP cells. Negative control transfections of Cas9 and empty U6 promoter plasmids were used to establish background EGFP loss at ~2.5% for all experiments (represented as a red dashed lined in FIGs).

T7E1 Assay

[0163] T7E1 assays were performed as previously described (Reyon, D. et al. FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol* 30, 460-465 (2012)) to quantify Cas9-induced mutagenesis at endogenous loci in human cells. Approximately 72 hours post-transfection, genomic DNA was isolated using the Agencourt DNAdvance Genomic DNA Isolation Kit (Beckman Coulter Genomics). Target loci were PCR-amplified from ~100 ng of genomic DNA using the primers listed in Table 11. Following an Agencourt Ampure XP clean-up step (Beckman Coulter Genomics), ~200 ng purified PCR product was denatured and hybridized prior to digestion with T7E1 (New England Biolabs). Following a second clean-up step, mutagenesis frequencies were quantified using a Qiaxcel capillary electrophoresis instrument (Qiagen).

GUIDE-Seq Experiments

[0164] GUIDE-seq experiments were performed and analyzed as previously described (Tsai, S. Q. et al. GUIDE-seq

enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015)). Briefly, U2OS cells were transfected as described above with Cas9 and sgRNA plasmids, as well as 100 pmol of a phosphorylated, phosphorothioate-modified double-stranded oligodeoxynucleotide (dsODN) with an embedded NdeI site. Restriction fragment length polymorphism (RFLP) analyses were performed to determine frequency of dsODN-tag integration frequencies ((See Examples 1-2; Tsai, S. Q. et al. GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015)), and T7E1 assays were performed to quantify on-target Cas9 mutagenesis frequencies. dsODN tag-specific amplification and library preparation (Tsai, S. Q. et al. GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015)) was performed prior to high-throughput sequencing using an Illumina MiSeq Sequencer. When mapping potential off-target sites, the cut-off for alignment to the on-target spacer sequence was set at 8 mismatches for 21 nucleotide spacers, 9 mismatches for 22 nucleotide spacers, and 10 mismatches for 23 nucleotide spacers. Off-target sites with potential DNA- or RNA-bulges (Lin, Y. et al. CRISPR/Cas9 systems have off-target activity with insertions or deletions between target DNA and guide RNA sequences. *Nucleic Acids Res* 42, 7473-7485 (2014)) were identified by manual alignment.

TABLE 10

Plasmids used in Example 3		
Name	SEQ ID NO:	Description
BPK2101	10	T7-humanSaCas9-NLS-3xFLAG-T7-BsaIcassette-Sa-sgRNA(120) Addgene ID: 65770 T7 promoters at 1-17 and 3418-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3352, NLS at 3256-3276, 3xFLAG tag at 3283-3348, BsaI sites at 3437-3442 and 3485-3490, gRNA at 3492-3616, T7 terminator at 3627-2674 of SEQ ID NO: 10.
MSP2283	21	T7-humanSaCas9-NLS-3xFLAG-T7-site1-Sa-sgRNA(84) T7 promoters at nts 1-17 and 3418-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, NLS at 3256-3276, 3xFLAG tag at 3243-33348, site 1 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 21
MSP2262	22	T7-humanSadCas9(D10A, H557A)-NLS-3xFLAG-T7-site1-Sa-sgRNA(84) T7 promoters at nts 1-17 and 3418-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, modified codons at 118-120 and 1759-1761, NLS at 3256-3276, 3xFLAG tag at 3243-33348, site 1 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 22
MSP2253	23	T7-humanSaCas9(E782K, N968K, R1015H)-NLS-3xFLAG-T7-site1-Sa-sgRNA(84) T7 promoters at nts 1-17 and 3418-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, modified codons at 2434-2436, 2992-2994, and 3133-3135, NLS at 3256-3276, 3xFLAG tag at 3243-33348, site 1 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 23
MSP2266	24	T7-humanSaCas9-NLS-3xFLAG-T7-site2-Sa-sgRNA(84) T7 promoters at 1-17 and 3419-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, NLS at 3256-3276, 3xFLAG tag at 3283-3348, site 2 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 24
MSP2279	25	T7-humanSadCas9(D10A, H557A)-NLS-3xFLAG-T7-site2-Sa-sgRNA(84) T7 promoters at 1-17 and 3419-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, modified codons at 118-120 and 1759-1761, NLS at 3256-3276, 3xFLAG tag at 3283-3348, site 2 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 25
MSP2292	26	T7-humanSaCas9(E782K, N968K, R1015H)-NLS-3xFLAG-T7-site2-Sa-sgRNA(84) T7 promoters at 1-17 and 3419-3434, human codon optimized <i>S. aureus</i> Cas9 at 88-3351, modified codons at 2434-2436, 2992-2994, and 3133-3135, NLS at 3256-3276, 3xFLAG tag at 3283-3348, site 2 spacer at 3435-3455, sgRNA(84) at 3456-3539, T7 terminator at 3562-3609 of SEQ ID NO: 26

TABLE 10-continued

Plasmids used in Example 3		
Name	SEQ ID NO:	Description
p11-lacY-wtx1	—	BAD-ccDB-Amp ^R -AraC-lacY(A177C) (Chen et al, 2005)
BPK2139	17	CAG-humanSaCas9-NLS-3xFLAG Addgene ID: 65776 Human codon optimized <i>S. aureus</i> Cas9 1-3195, NLS 3169-3189, 3xFLAG tag 3196-3261 of SEQ ID NO: 17.
MSP1830	27	CAG-humanSaCas9(E782K, N968K, R1015H)-NLS-3xFLAG (KKH variant) Human codon optimized <i>S. aureus</i> Cas9 1-3264, NLS 3169-3189, modified codons at 2347-2349, 2905-2907, and 3046-3048, 3xFLAG tag 3196-3261 of SEQ ID NO: 27
VVT1	20	U6-BsmBIcassette-Sa-sgRNA(120) Addgene ID: 65779 U6 promoter 1-318, BsmBI sites at 320-325 and 333-338, <i>S. aureus</i> gRNA 340-466, U6 terminator 459-466 of SEQ ID NO: 20.
BPK2660	28	U6-BsmBIcassette-Sa-sgRNA(84) U6 promoter 1-318, BsmBI sites at 320-325 and 333-338, <i>S. aureus</i> gRNA 340-423, U6 terminator 424-430 of SEQ ID NO: 28.

TABLE 11

Oligonucleotides used in Example 3		
Sequence	Description	SEQ ID NO:
Oligos used to generate positive selection plasmids		
ctagaGGGtGGGcGGGaGGGTCGCCCTCGAACTT CACCTtgGAGTgcatg	top oligo to clone site 2 with an NNGAGT (SEQ ID NO: 5) PAM into the positive selection vector (XbaI/SphI cut p11-lacY-wtx1)	515
cACTCcaAGGTGAAGTTCGAGGGCGACCCtCCCG CCCACCCt	bottom oligo to clone site 2 into the positive selection vector	516
Oligos used to generate libraries for site-depletion experiments		
GCAGgaattcGGGAGGGGCACGGGCAGCTTGCCG GNNNNNNNNCTNNNGCGCAGGTCACGAGGCATG	top strand oligo for site 1 PAM library, cut with EcoRI once filled in	517
GCAGgaattcGGAGGGTCGCCCTCGAACTTCACC TNNNNNNNNCTNNNGCGCAGGTCACGAGGCATG	top strand oligo for site 2 PAM library, cut with EcoRI once filled in	518
/5Phos/CCTCGTGACCTGCGC	reverse primer to fill in library oligos	200
Primers used to amplify site-depletion libraries for sequencing		
GATACCGCTCGCCGAGC	forward primer	201
CTGCGTTCTGATTTAATCTGTATCAGGC	reverse primer	202
Primers used for T7E1 and RFLP experiments		
GGAGCAGCTGGTCAGAGGGG	forward primer targeted to EMX1 in U2OS human cells	209
CCATAGGGAAGGGGACACTGG	reverse primer targeted to EMX1 in U2OS human cells	210
GGGCCGGGAAGAGTTGCTG	forward primer targeted to FANCF in U2OS human cells	211
GCCCTACATCTGCTCTCCCTCC	reverse primer targeted to FANCF in U2OS human cells	212
CCAGCACAACCTACTCGCACTTGAC	forward primer targeted to RUNX1 in U2OS human cells	213
CATCACCAACCCACAGCCAAGG	reverse primer targeted to RUNX1 in U2OS human cells	214
TCCAGATGGCACATTGTGACG	forward primer targeted to VEGFA in U2OS human cells	652
AGGGAGCAGGAAAGTGAGGT	reverse primer targeted to VEGFA in U2OS human cells	653

TABLE 12

sgRNA target sites for Example 3						
In VVT1 (120) EGFP						
Prep Name	Name	Spacer length (nt)	Spacer Sequence	SEQ ID NO.: Sequence with PAM	SEQ ID NO.:	
MSP1428	NNGRRT 1 (SEQ ID NO: 46)	21	GCCCTCGAACTTCACCTCGGC	405 GCCCTCGAACTTCACCTCGGC	406	
MSP1400	NNGRRT 2 (SEQ ID NO: 46)	21	GCAACATCCTGGGGCACAAGC	397 GCAACATCCTGGGGCACAAGCTGGAGT	398	
MSP1401	NNGRRT 3 (SEQ ID NO: 46)	21	GTTGTACTCCAGCTTGTGCCC	519 GTTGTACTCCAGCTTGTGCCCCAGGAT	520	
MSP1403	NNGRRT 4 (SEQ ID NO: 46)	22	GCAAGGGCGAGGAGCTGTTTAC	409 GCAAGGGCGAGGAGCTGTTTACCGGGGT	410	
MSP1748	NNARRT 1 (SEQ ID NO: 43)	20	GGACGGCGACGTAAACGGCC	521 GGACGGCGACGTAAACGGCCACAAGT	522	
MSP1754	NNARRT 5 (SEQ ID NO: 43)	21	GAACCTCAGGGTCAGCTTGCC	523 GAACCTCAGGGTCAGCTTGCCGTAGGT	524	
MSP2030	NNCRRT 2 (SEQ ID NO: 47)	20	GTCGATGCCCTTCAGCTCGA	525 GTCGATGCCCTTCAGCTCGATCGGT	526	
MSP2034	NNCRRT 4 (SEQ ID NO: 47)	22	GTGACCACCCTGACCTACGGCG	527 GTGACCACCCTGACCTACGGCGTGCAGT	528	
MSP2040	NNTRRT 1 (SEQ ID NO: 48)	20	GATATAGACGTTGTGGCTGT	529 GATATAGACGTTGTGGCTGTTGTAGT	530	
MSP2045	NNTRRT 3 (SEQ ID NO: 48)	21	GGTGAAGTTCGAGGGCGACAC	531 GGTGAAGTTCGAGGGCGACACCTGGT	532	
In BPK2660 (84) EGFP						
Prep Name	Name	Spacer length (nt)	Spacer Sequence	Sequence with PAM		
MSP2149*	NNARRT 1 (SEQ ID NO: 43)	20	GGACGGCGACGTAAACGGCC	521 GGACGGCGACGTAAACGGCCACAAGT	522	
MSP2152	NNARRT 2 (SEQ ID NO: 43)	21	GTAGTTGCCGTCGTCCTTGAA	654 GTAGTTGCCGTCGTCCTTGAAGAAGAT	655	
MSP2153	NNARRT 3 (SEQ ID NO: 43)	22	GCCACCTACGGCAAGCTGACCC	656 GCCACCTACGGCAAGCTGACCTGAAGT	657	
MSP2154	NNARRT 4 (SEQ ID NO: 43)	23	GACGGCAACTACAAGACCCGCGC	658 GACGGCAACTACAAGACCCGCGCCGAGGT	659	
MSP2150*	NNARRT 5 (SEQ ID NO: 43)	21	GAACCTCAGGGTCAGCTTGCC	523 GAACCTCAGGGTCAGCTTGCCGTAGGT	524	
MSP2155	NNCRRT 1 (SEQ ID NO: 47)	20	GCGTGTCCGGCGAGGGCGAG	305 GCGTGTCCGGCGAGGGCGAGGGCGAT	533	
MSP2156*	NNCRRT 2 (SEQ ID NO: 47)	20	GTCGATGCCCTTCAGCTCGA	525 GTCGATGCCCTTCAGCTCGATCGGT	526	
MSP2158	NNCRRT 3 (SEQ ID NO: 47)	22	GCTCGACCAGGATGGGCACCAC	534 GCTCGACCAGGATGGGCACCACCCCGGT	535	
MSP2159*	NNCRRT 4 (SEQ ID NO: 47)	22	GTGACCACCCTGACCTACGGCG	527 GTGACCACCCTGACCTACGGCGTGCAGT	528	

TABLE 12-continued

sgRNA target sites for Example 3						
MSP2145*	NNGRRT 1 (SEQ ID NO: 46)	21	GCCCTCGAACTTCACCTCGGC	405	GCCCTCGAACTTCACCTCGGC	406
MSP2146*	NNGRRT 2 (SEQ ID NO: 46)	21	GCAACATCCTGGGGCACAAGC	397	GCAACATCCTGGGGCACAAGCTGGAGT	398
MSP2147	NNGRRT 3 (SEQ ID NO: 46)	21	GTTGTACTCCAGCTTGTGCCC	519	GTTGTACTCCAGCTTGTGCCCCAGGAT	520
MSP2148	NNGRRT 4 (SEQ ID NO: 46)	22	GCAAGGGCGAGGAGCTGTTTAC	409	GCAAGGGCGAGGAGCTGTTTACCGGGGT	410
MSP2161*	NNTRRT 1 (SEQ ID NO: 48)	20	GATATAGACGTTGTGGCTGT	529	GATATAGACGTTGTGGCTGTTGTAGT	530
MSP2162	NNTRRT 2 (SEQ ID NO: 48)	21	GGGCGAGGAGCTGTTTACCGG	536	GGGCGAGGAGCTGTTTACCGGGGTGGT	537
MSP2164*	NNTRRT 3 (SEQ ID NO: 48)	21	GGTGAAGTTCGAGGGCGACAC	531	GGTGAAGTTCGAGGGCGACACCCTGGT	532
MSP2163	NNTRRT 4 (SEQ ID NO: 48)	21	GCACTGCACGCCGTAGGTCAG	538	GCACTGCACGCCGTAGGTCAGGGTGGT	539
Endogenous genes EMX1						
Prep Name	Name	Spacer length (nt)	Spacer Sequence	Sequence with PAM		
MSP2184**	EMX1 1	22	GTGTGGTTCAGAACCGGAGGA	540	GTGTGGTTCAGAACCGGAGGACAAAGT	541
MSP2185	EMX1 2	21	GCAGGCTCTCCGAGGAGAAGG	542	GCAGGCTCTCCGAGGAGAAGGCAAGT	543
MSP2183	EMX1 3	23	GCCCTCCCTCCCTGGCCAGGT	544	GCCCTCCCTCCCTGGCCAGGTGAAGGT	545
MSP2199**	EMX1 4	21	GCTCAGCCTGAGTGTTGAGGC	546	GCTCAGCCTGAGTGTTGAGGCCCCAGT	547
MSP2202	EMX1 5	21	GCCTGCTTCGTGGCAATGCGCC	548	GCCTGCTTCGTGGCAATGCGCCACCGGT	549
MSP2168**	EMX1 6	21	GCAACCACAAACCCACGAGGG	550	GCAACCACAAACCCACGAGGGCAGAGT	551
MSP2169	EMX1 7	21	GGCCTCCCCAAAGCCTGGCCA	552	GGCCTCCCCAAAGCCTGGCCAGGGAGT	553
MSP2170	EMX1 8	23	GCAGAAGCTGGAGGAGGAAGGGC	554	GCAGAAGCTGGAGGAGGAAGGGCCTGAGT	555
MSP2201	EMX1 9	21	GCTTCGTGGCAATGCGCCACCG	556	GCTTCGTGGCAATGCGCCACCGGTTGAT	557
MSP2200**	EMX1 10	22	GGCTCTCCGAGGAGAAGGCCA	558	GGCTCTCCGAGGAGAAGGCCAAGTGGT	559
FANCF						
Prep Name	Name	Spacer length (nt)	Spacer Sequence	Sequence with PAM		
MSP2189	FANCF 1	22	GCCTCTCTGCAATGCTATTGGT	560	GCCTCTCTGCAATGCTATTGGTCGAAAT	561
MSP2190	FANCF 2	21	GCGTACTGATTGGAACATCCG	562	GCGTACTGATTGGAACATCCGCGAAAT	563
MSP2186	FANCF 3	23	GACGTCACAGTGACCGAGGGCCT	564	GACGTCACAGTGACCGAGGGCCTGGAAGT	565
MSP2187	FANCF 4	23	GCCCGGCGCACGGTGGCGGGGTC	566	GCCCGGCGCACGGTGGCGGGGTCCAGGT	567
MSP2188	FANCF 5	21	GGCGGGGTCCCAGGTGCTGAC	568	GGCGGGGTCCCAGGTGCTGACGTAGGT	569
MSP2205	FANCF 6	21	GGCGTATCATTTTCGCGGATGT	570	GGCGTATCATTTTCGCGGATGTTCCAAT	571
MSP2208	FANCF 7	22	GAGACCGCCAGAACTCGGAAA	572	GAGACCGCCAGAACTCGGAAAAGCGAT	573
MSP2204	FANCF 8	21	GGATCGCTTTTCCGAGCTTCT	574	GGATCGCTTTTCCGAGCTTCTGGCGGT	575
MSP2207**	FANCF 9	22	GCGCCCACTGCAAGGCCCGCG	576	GCGCCCACTGCAAGGCCCGCGCACGGT	577
MSP2172**	FANCF 10	21	GTAGGGCCTTCGCGCACCTCA	578	GTAGGGCCTTCGCGCACCTCATGGAAT	579
MSP2174	FANCF 11	22	GCAGCCGCGCTCCAGAGCCGT	580	GCAGCCGCGCTCCAGAGCCGTGCGAAT	581
MSP2332	FANCF 12	22	GGCCATGCCGACCAAAGCGCCG	582	GGCCATGCCGACCAAAGCGCCGATGGAT	583
MSP2171**	FANCF 13	21	GCAAGGCCCGCGCACGGTGG	584	GCAAGGCCCGCGCACGGTGGCGGGGT	585
MSP2173	FANCF 14	22	GAGGCAAGAGGGCGGCTTTGGG	586	GAGGCAAGAGGGCGGCTTTGGGCGGGGT	587
MSP2206	FANCF 15	22	GTGACCGAGGGCCTGGAAGTTC	588	GTGACCGAGGGCCTGGAAGTTCGCTAAT	589
MSP2203**	FANCF 16	21	GGGGTCCCAGGTGCTGACGTA	590	GGGGTCCCAGGTGCTGACGTAGGTAGT	591
MSP2209	FANCF 17	22	GTACTGATTGGAACATCCGCGA	592	GTACTGATTGGAACATCCGCGAAATGAT	593

TABLE 12-continued

sgRNA target sites for Example 3					
RUNX1					
Prep Name	Name	Spacer length (nt)	Spacer Sequence	Sequence with PAM	
MSP2192	RUNX1 1	23	GTCTGAAGCCATCGCTTCCTCCT	594. GTCTGAAGCCATCGCTTCCTCCTGAAAAT	595.
MSP2193	RUNX1 2	21	GGTTTTTCGCTCCGAAGGTAAA	596. GGTTTTTCGCTCCGAAGGTAAAAGAAAT	597.
MSP2195	RUNX1 3	21	GGGACTCCCCAAGCCCTATTA	598. GGGACTCCCCAAGCCCTATTAAGAAAT	599.
MSP2235	RUNX1 4	22	GCAGCTTGTTTCACCTCGGTGC	600. GCAGCTTGTTTCACCTCGGTGCAGAGAT	601.
MSP2194	RUNX1 5	22	GACCTGTCTTGGTTTTCGCTCC	602. GACCTGTCTTGGTTTTCGCTCCGAAGGT	603.
MSP2216	RUNX1 6	23	GCTTCCATCTGATTAGTAAGTAA	604. GCTTCCATCTGATTAGTAAGTAATCCAAT	605.
MSP2214	RUNX1 7	22	GTGCAGAGATGCCTCGGTGCCT	606. GTGCAGAGATGCCTCGGTGCCTGCCAGT	607.
MSP2211	RUNX1 8	21	GAGGGTGCATTTTCAGGAGGA	608. GAGGGTGCATTTTCAGGAGGAAGCGAT	609.
MSP2217	RUNX1 9	23	GTTTCACCTCGGTGCAGAGATGC	610. GTTTCACCTCGGTGCAGAGATGCCTCGGT	611.
MSP2176	RUNX1 10	22	GCGATGGCTTCAGACAGCATAT	612. GCGATGGCTTCAGACAGCATATTTGAGT	613.
MSP2177	RUNX1 11	22	GCTCCGAAGGTAAAAGAAATCA	614. GCTCCGAAGGTAAAAGAAATCATTGAGT	615.
MSP2334	RUNX1 12	22	GAGGCATATGATTACAAGTCTA	616. GAGGCATATGATTACAAGTCTATTGGAT	617.
MSP2175**	RUNX1 13	21	GAAAGAGAGATGTAGGGCTAG	618. GAAAGAGAGATGTAGGGCTAGAGGGGT	619.
MSP2178**	RUNX1 14	23	GTACTCACCTCTCATGAAGCACT	620. GTACTCACCTCTCATGAAGCACTGTGGGT	621.
MSP2210	RUNX1 15	21	GAGGTGAGTACATGCTGGTCT	622. GAGGTGAGTACATGCTGGTCTTGTAAT	623.
MSP2213	RUNX1 16	22	GAGAGGAATTCAAAGTGGGCA	624. GAGAGGAATTCAAAGTGGGCATATGAT	625.
MSP2212	RUNX1 17	21	GAGGCTGAAACAGTGACCTGT	626. GAGGCTGAAACAGTGACCTGTCTTGGT	627.
VEGFA					
Prep Name	Name	Spacer length (nt)	Spacer Sequence	Sequence with PAM	
MSP2196	VEGFA 1	21	GTACATGAAGCAACTCCAGTC	628. GTACATGAAGCAACTCCAGTCCCAAAT	629.
MSP2198	VEGFA 2	21	GACGGGTGGGGAGAGGGACAC	630. GACGGGTGGGGAGAGGGACACAGAT	631.
MSP2197	VEGFA 3	22	GTCCCAAATATGTAGCTGTTTG	632. GTCCCAAATATGTAGCTGTTTGGGAGGT	633.
MSP2219	VEGFA 4	21	GGCCAGGGGTCACTCCAGGAT	634. GGCCAGGGGTCACTCCAGGATTCCAAT	635.
MSP2220	VEGFA 5	22	GCCAGAGCCGGGTGTGCAGAC	636. GCCAGAGCCGGGTGTGCAGACGGCAGT	637.
MSP2181	VEGFA 6	22	GAGGACGTGTGTGTCTGTGTGG	638. GAGGACGTGTGTGTCTGTGTGGGTGAGT	639.
MSP2336	VEGFA 7	22	GGGAGAAGGCCAGGGGTCACTC	640. GGGAGAAGGCCAGGGGTCACTCAGGAT	641.
MSP2179**	VEGFA 8	21	GGGTGAGTGAGTGTGTGCGTG	642. GGGTGAGTGAGTGTGTGCGTGTGGGT	643.
MSP2180	VEGFA 9	22	GAGTGAGGACGTGTGTGTCTGT	644. GAGTGAGGACGTGTGTGTCTGTGTGGGT	645.
MSP2182	VEGFA 10	22	GCGTTGGAGCGGGGAGAAGGCC	646. GCGTTGGAGCGGGGAGAAGGCCAGGGGT	647.
MSP2218	VEGFA 11	21	GCTCCATTACCCAGCTTCCC	648. GCTCCATTACCCAGCTTCCCTGTGGT	649.

*Used in FIGS. 1C and 1E, FIG. 32

**Used for GUIDE-seq experiments in FIG. 3, FIGS. 36A-B

Example 3

Engineering the PAM Specificity of *Staphylococcus aureus* Cas9

[0165] Site-specific DNA cleavage by CRISPR-Cas9 nucleases is primarily guided by RNA-DNA interactions, but also requires Cas9-mediated recognition of a protospacer adjacent motif (PAM). Although the commonly used *Streptococcus pyogenes* Cas9 specifies only two nucleotides within its NGG PAM, other Cas9 orthologues with desirable properties recognize longer PAMs. While potentially advantageous from the perspective of specificity, extended PAM sequences can limit the targeting range of Cas9 orthologues for genome editing applications. One possible strategy to broaden the range of sequences targetable by such Cas9 orthologues might be to evolve variants with relaxed speci-

ficity for certain positions within the PAM. Here we used molecular evolution to modify the NNGRRT (SEQ ID NO:46) PAM specificity of *Staphylococcus aureus* Cas9 (SaCas9), a smaller size orthologue that is useful for applications requiring viral delivery. One variant we identified, referred to as KKH SaCas9, shows robust genome editing activities at endogenous human target sites with NNNRRT PAMs. Importantly, using the GUIDE-seq method, we showed that both wild-type and KKH SaCas9 induce comparable numbers of off-target effects in human cells. KKH SaCas9 increased the targeting range of SaCas9 by nearly two- to four-fold, enabling targeting of sequences that cannot be altered with the wild-type nuclease. More generally, these results demonstrate the feasibility of relaxing PAM specificity to broaden the targeting range of Cas9 orthologues. Our molecular evolution strategy does not

require structural information or a priori knowledge of specific residues that contact the PAM, and therefore should be applicable to a wide range of Cas9 orthologues.

Results

[0166] We devised an unbiased genetic approach for engineering Cas9 variants with relaxed PAM recognition specificities that does not require structural information. We tested this strategy using SaCas9, for which no structural data was available at the time we initiated these studies. In an initial step, we sought to conservatively estimate the PAM-interacting domain for SaCas9 by sequence comparisons with the structurally well-characterized SpCas9 (Jiang et al., *Science* 348, 1477-1481 (2015); Anders et al., *Nature* 513, 569-573 (2014); Jinek et al., *Science* (2014); Nishimasu et al., *Cell* (2014)). Although SpCas9 and SaCas9 differ substantially at the primary sequence level (FIG. 21A, FIG. 29), alignment of both with 10 additional orthologues enabled us to conservatively define a predicted PAM-interacting domain for SaCas9 (See Methods for Example 3; FIGS. 29 and 30).

[0167] Because the guanine at the third position in the SaCas9 PAM is the most strictly specified base (Ran et al., *Nature* 520, 186-191 (2015)), we randomly mutagenized the predicted PI domain and used our previously described bacterial cell-based method (see Examples 1-2) to attempt to select for mutants capable of cleaving sites with each of the three other possible nucleotides at the 3rd PAM position (i.e., NN[A/C/T]RRT PAMs (NNHRRT (SEQ ID NO:44)); FIG. 31A). All but one of the surviving variants from the selections against sites containing NNARRT (SEQ ID NO:43) and NNCRRRT (SEQ ID NO:47) PAMs harbored an R1015H mutation, whereas we did not obtain any variants from the selections with NNTRRT (SEQ ID NO:48) PAMs. These results strongly suggested that R1015 might participate in recognition of the guanine at the third position of the SpCas9 PAM. Indeed, in our alignments we found that R1015 of SaCas9 is in the vicinity of SpCas9 R1335 (FIG. 30), a residue previously implicated in recognition of the third base position of the PAM ((See Examples 1-2; Anders, C., Niewoehner, O., Duerst, A. & Jinek, M. Structural basis of PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature* 513, 569-573 (2014)). Consistent with this, we found that mutation of R1015 to an alanine or glutamine substantially decreased SaCas9 activity on a target site containing an NNGRRRT (SEQ ID NO:46) PAM (FIG. 21B) when tested in our bacterial selection system (FIG. 31B). Alanine or glutamine substitutions of other positively charged residues in the vicinity of R1015 did not have as strong of an effect on SaCas9 activity (FIG. 21B, FIG. 30).

[0168] Our bacterial-based selection results also suggested that the R1015H mutation might at least partially relax the specificity of SaCas9 at the third position of the PAM. However, we found that the R1015H single mutant had suboptimal activity in our previously described human cell-based EGFP disruption assay (Fu et al., *Nat Biotechnol* 31, 822-826 (2013); Reyon et al., *Nat Biotechnol* 30, 460-465 (2012)) when tested against sites with any nucleotide at the 3rd position of NNNRRT PAMs (FIG. 21C). Because this suggested that additional mutations might be required to increase or optimize the activity of the R1015H mutant in human cells, we randomly mutagenized a region encompassing the predicted PI domain of an SaCas9 that also

harbored a R1015Q mutation. We then selected for variants from this library that could cleave target sites with each of the three different NNHRRT (SEQ ID NO:44) PAMs using our bacterial selection system. We used R1015Q because, unlike R1015H, this mutant did not show activity in bacteria (FIG. 21B). Although no surviving clones were again observed when selecting against NNTRRT (SEQ ID NO:48) PAMs, selections with the R1015Q variant against NNARRT (SEQ ID NO:43) or NNCRRRT (SEQ ID NO:47) yielded mutations at E782, K929, N968, and, surprisingly, mutation of the Q at 1015 to H.

[0169] Combined with the selection results from wild-type SaCas9, the most frequent missense mutations identified across all selections were E782K, K929R, N968K, and R1015H (FIG. 21D), suggesting that a combination of these mutations might permit efficient cleavage of sites that contain an A or C at the third position of the SaCas9 PAM. We therefore tested SaCas9 variants containing different combinations of these mutations using the human cell-based EGFP disruption assay with sgRNAs targeted to sites harboring each of the 4 bases at the third position of the PAM (i.e., on NNNRRT PAMs) (FIG. 21E, FIG. 32). We found that the variants with the triple mutant combinations E782K/N968K/R1015H and E782K/K929R/R1015H were highly active at sites with NNNRRT PAMs (FIG. 21E, FIG. 32), whereas the quadruple mutant variant containing all four mutations (E782K/K929R/N968K/R1015H) had generally lower activities on these sites (FIG. 32). We chose the E782K/N968K/R1015H (hereafter referred to as the KKH variant) for further characterization, and verified using our human cell-based EGFP disruption assay that all three substitutions comprising the KKH variant were required for activity (FIG. 21E).

[0170] To more comprehensively define the PAM specificities of KKH and wild-type SaCas9, we used our previously described bacterial cell-based site-depletion assay (See Examples 1-2) (FIG. 33). This method yields Cas9 PAM specificity profiles by identifying the relative cleavage (and therefore depletion in bacterial cells) of DNA plasmids bearing randomized PAM sequences, quantified as a post-selection PAM depletion value (PPDV). We performed site-depletion experiments with both wild-type and KKH SaCas9 using libraries with two different spacer sequences each with 8 randomized bases in place of the PAM (FIG. 33). Control experiments using catalytically inactive SaCas9 showed little depletion of any PAM sequence (FIG. 34A), enabling us to establish a threshold for statistically significant depletion as a PPDV of 0.794 (FIG. 34B). Previous experiments have shown that PAMs with PPDVs of <0.2 in our bacterial site-depletion assay can be efficiently cleaved in our human cell-based EGFP disruption assay (See Examples 1-2). With wild-type SaCas9, the most depleted PAMs (based on mean PPDVs obtained from the two libraries) were, as expected, the four NNGRRRT (SEQ ID NO:46) (PAMs (FIG. 21F and FIG. 34C). Interestingly, other PAMs with mean PPDVs <0.1 included those of the form NNGRRN (SEQ ID NO:49) (FIG. 34), suggesting that for some spacer sequences the last position of the PAM may not be fully specified as a T in our bacterial-based assay (although a previous report demonstrated by an in vitro PAM depletion assay, ChIP-seq, and targeting of endogenous human sites that a thymine at the sixth position of the PAM was highly preferred (Ran, F. A. et al. In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 520,

186-191 (2015))). By contrast, with the KKH variant, PAMs with mean PPDVs of <0.2 included not only the NNGRRT (SEQ ID NO:46) PAMs but also all four NNARRT (SEQ ID NO:43), all four NNCRRT (SEQ ID NO:47), and three of the four NNTRRT (SEQ ID NO:48) PAMs (FIG. 21F, FIGS. 34C and 34E). These results suggested that KKH SaCas9 appears to have a broadened PAM targeting range relative to its wild-type counterpart.

[0171] To assess the robustness of the KKH SaCas9 variant in human cells, we tested its activity on 55 different endogenous gene target sites containing a variety of NNNRRT PAMs (FIG. 22A). The KKH variant showed efficient activity with a mean mutagenesis frequency of 24.7% across all sites, with 80% of sites (44 of 55 sites) showing greater than 5% disruption. Analysis of KKH SaCas9 activity across all 55 sites revealed ordered preferences for the 3rd position of the PAM (NN[G>A=C>T]RRT; FIG. 22B) as well as the 4th/5th positions of the PAM (NNN[AG>GG>GA>AA]T; FIG. 22C). Consistent with this, we observed differences among the 16 possible combinations of the 3rd/4th/5th positions of an NNNRRT PAM (FIG. 35A). KKH SaCas9 functioned efficiently on spacer lengths ranging from 21-23 nucleotides (FIG. 22D), spacer sequences with variable GC content (FIG. 35B), and PAMs with variable GC content (FIG. 35C). Sequence logos derived from sites cleaved with low, medium, and high efficiencies (0-10%, 10-30%, and >30% mean mutagenesis frequencies, respectively) revealed little sequence preference across the entire target site other than at the 4th and 5th positions of the NNNRRT PAM, and perhaps a slight preference for guanine at the 2nd PAM position on sites cleaved with high efficiencies (FIG. 35D).

[0172] To demonstrate that the KKH variant enables modification of PAMs that cannot be targeted by wild-type SaCas9, we performed direct comparisons of these nucleases in human cells on sites bearing various NNNRRT PAMs. Assessment of 16 sites using our EGFP disruption assay and 16 endogenous human gene targets (FIGS. 22E and 22F, respectively) showed that KKH SaCas9 robustly modified target sites bearing NNNRRT PAMs whereas wild-type SaCas9 efficiently targeted only sites with NNGRRT (SEQ ID NO:46) PAMs. For all 24 sites with NNHRRT (SEQ ID NO:44) PAMs, the KKH variant induced substantially higher rates of mutagenesis than wild-type SaCas9; on the eight sites with NNGRRT (SEQ ID NO:46) PAMs, KKH SaCas9 induced comparable or slightly lower levels of mutagenesis compared with wild-type (FIGS. 22E and 22F). These results collectively demonstrate that the KKH variant can cleave sites with NNNRRT PAMs, thereby enabling targeting of sites with NNHRRT (SEQ ID NO:44) PAMs that currently cannot be efficiently altered by wild-type SaCas9 in human cells.

[0173] To assess the impact of the KKH mutations on the genome-wide specificity of SaCas9, we used the GUIDE-seq (Genome-wide Unbiased Identification of DSBs Enabled by sequencing) method (Tsai, S. Q. et al. GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015)) to directly compare the off-target profiles of wild-type and KKH SaCas9 with the same sgRNAs. When tested with sgRNAs targeted to six endogenous human gene sites containing NNGRRT (SEQ ID NO:46) PAMs, we observed that wild-type and KKH SaCas9 induced nearly identical GUIDE-seq tag integration rates and on-target cleavage

frequencies for all six sites (FIGS. 36A and 36B, respectively). Furthermore, wild-type and KKH SaCas9 induce mutations at similar numbers of off-target sites with each of the six sgRNAs (FIGS. 23A and 23B). Off-target sites for the KKH variant generally adhered to the NNNRRT PAM motif, and off-target sites for wild-type SaCas9 adhered to an NNGRR|T>G| motif (FIG. 22B). With one of the sgRNAs, which induced the highest number of off-target sites among the six sgRNAs tested, we observed a similar number of off-target sites with wild-type and KKH SaCas9. However, the off-target sites were only partially overlapping between wild-type and KKH SaCas9, as might be expected given their different PAM specificities (FIGS. 23B and 23C). Although we would not advocate the use of the KKH variant for targeting sites with NNGRRT (SEQ ID NO:46) PAMs (because wild-type SaCas9 can show higher on-target activities than KKH for these sites), these results suggest that KKH SaCas9 only cleaves off-target sites with the expected PAMs and generally induces numbers of off-target sites comparable to those observed with wild-type SaCas9.

[0174] To further examine the genome-wide specificity of KKH SaCas9, we tested five additional sgRNAs targeted to sites containing NNHRRT (SEQ ID NO:44) PAMs (FIGS. 23D and 23E). Off-target sites detected by GUIDE-seq were generally low in number (comparable to the numbers observed with wild-type SpCas9 and SpCas9 variants in previously published experiments (See Examples 1-2 (Tsai, S. Q. et al. GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015))), displayed potential DNA- and RNA-bulged off-targets (Lin, Y. et al. CRISPR/Cas9 systems have off-target activity with insertions or deletions between target DNA and guide RNA sequences. *Nucleic Acids Res* 42, 7473-7485 (2014))), and contained expected PAM sequences. Taken together, our experiments demonstrate that the genome-wide specificities of wild-type and KKH SaCas9 are similar and generally show low numbers of off-target mutations in human cells as judged by GUIDE-seq.

[0175] Although wild-type SaCas9 remains the most optimal choice for targeting NNGRRT (SEQ ID NO:46) PAMs, the KKH SaCas9 variant we describe here can robustly target sites with NNARRT (SEQ ID NO:43) and NNCRRT (SEQ ID NO:47) PAMs and has a reasonable success rate for sites with NNTRRT (SEQ ID NO:48) PAMs. Thus, we conservatively estimate that the KKH variant increases the targeting range of SaCas9 by nearly two- to four-fold in random DNA sequence, thereby improving the prospects for more broadly utilizing SaCas9 in a variety of different applications that require highly precise targeting. Using GUIDE-seq, we demonstrated that KKH SaCas9 induces similar numbers of off-target mutations as wild-type SaCas9 when targeted to the same sites that contain NNGRRT (SEQ ID NO:46) PAMs. Also, KKH SaCas9 induces only a small number of off-target mutations when targeted to sites bearing NNHRRT (SEQ ID NO:44) PAMs. Although KKH SaCas9 recognizes a modified PAM sequence relative to wild-type SaCas9, our findings are not entirely surprising given that the total combined length of the protospacer and PAM is still long enough with the KKH variant (24 to 26 bps) to be reasonably orthogonal to the human genome. Furthermore, it is possible that modifying PAM recognition can improve specificity by altering the energetics of Cas9/sgRNA interaction with its target site (similar to the previ-

ously proposed mechanisms for improved specificities of truncated sgRNAs (Fu, Y., Sander, J. D., Reyon, D., Cascio, V. M. & Joung, J. K. Improving CRISPR-Cas nuclease specificity using truncated guide RNAs. *Nat Biotechnol* 32, 279-284 (2014)) or the D1135E SpCas9 mutant (See Examples 1-2)).

Example 4

Improving the Activity of the SpCas9-VQR Variant

[0176] Because the SpCas9-VQR variant has a preference for NGAN PAMs of: NGAG>NGAA=NGAT>NGAC, we sought to select for derivative variants that had improved activity against NGAH PAMs (where H=A, C, or T). Selections with the R1335Q library (with PI domain randomly mutagenized) against cells that contain target sites with either an NGAA, NGAT, or NGAC PAM enabled us to sequence additional clones that contained mutations that convey an altered PAM specificity. The sequences of these clones revealed additional mutations that might be important for altering PAM specificity towards NGAA, NGAT, or NGAC PAMs.

[0177] Based on the results of these selections, the VQR variant and 24 other derivative variants were tested against NGAG, NGAA, NGAT, and NGAC PAM sites in bacteria. A number of these derivative variants survived better than the VQR variant on NGAH PAM sites, most of which contained the G1218R mutation (Table 7 and FIG. 24).

TABLE 7

Table of variants and their corresponding amino acid changes.						
variant	D1135	G1218	E1219	R1335	T1337	
A1 VRQ	V	R	—	Q	—	
A2 NRQ	N	R	—	Q	—	
A3 YRQ	Y	R	—	Q	—	
A4 VRQL	V	R	—	Q	L	
A5 VRQM	V	R	—	Q	M	
A6 VRQR	V	R	—	Q	R	
A7 VRQE	V	R	—	Q	E	
A8 VRQQ	V	R	—	Q	Q	
A9 NRQL	N	R	—	Q	L	
A10 NRQM	N	R	—	Q	M	
A11 NRQR	N	R	—	Q	R	
A12 NRQE	N	R	—	Q	E	
B1 NRQQ	N	R	—	Q	Q	
B2 YRQL	Y	R	—	Q	L	
B3 YRQM	Y	R	—	Q	M	
B4 YRQR	Y	R	—	Q	R	
B5 YRQE	Y	R	—	Q	E	
B6 YRQQ	Y	R	—	Q	Q	

TABLE 7-continued

Table of variants and their corresponding amino acid changes.						
variant	D1135	G1218	E1219	R1335	T1337	
B7 VRVQE	V	R	V	Q	E	
B8 NRVQE	N	R	V	Q	E	
B9 YRVQE	Y	R	V	Q	E	
B10 VVQE	V	—	V	Q	E	
B11 NVQE	N	—	V	Q	E	
B12 YVQE	Y	—	V	Q	E	
C1 VQR	V	—	—	Q	R	

[0178] Given that the results from the bacterial screen demonstrated that some of these additional mutations improved activity against NGAH PAM sites, we tested some of the best candidates in human cells in the EGFP disruption assay. What we observed is that a number of these variants outperformed the VQR variant at targeting NGAH sites, including the VRQR, NRQR, and YRQR variants (Table 8 and FIG. 25). The main difference between these clones and the VQR variant is that they include a G1218R mutation.

TABLE 8

Table of SpCas9-VQR derivatives and their corresponding amino acid changes				
variant	D1135	G1218	R1335	T1337
VQR	V	—	Q	R
YRQ	Y	R	Q	—
VRQR	V	R	Q	R
VRQQ	V	R	Q	Q
NRQR	N	R	Q	R
NRQQ	N	R	Q	Q
YRQR	Y	R	Q	R
YRQQ	Y	R	Q	Q

[0179] Because the VRQR variant appeared to be the most robust of those tested, we compared its activity to that of the VQR against 9 different endogenous sites in human cells (2 sites for each NGAA, NGAC, NGAT, and NGAG PAMs, and 1 site for an NGCG PAM). This data reveals that the VRQR variant outperforms the VQR variant at all sites tested in human cells (FIG. 26).

[0180] After demonstrating that VRQR variant has improved activity relative to the VQR variant, we sought to determine whether adding additional substitutions could further improve activity. Because we observed additional mutations in the selections that were in close proximity to the PAM interacting pocket of SpCas9, a subset of these mutations were added to the VQR and VRQR variants and screened in bacteria against sites containing NGAG, NGAA, NGAT, and NGAC PAMs (Table 9 and FIG. 27). A number

of derivative variants appears to have higher activity against NGAT and NGAC PAM sites, so we proceeded to test these variants in human cells. We tested in the human cell EGFP disruption assay additional variants that contained added mutations to either the VQR or VRQR background. These experiments again revealed that the VRQR has more robust activity against NGAH PAMs than the VQR variant, and that additional mutations to the VRQR backbone are beneficial.

TABLE 9

Table of variants and their corresponding amino acid changes		
variant	mutations	
1 VQR + L1111H	L1111H/D1135V/R1335Q/T1337R	
2 VRQR + L1111H	L1111H/D1135V/G1218R/R1335Q/T1337R	
3 VQR + E1219K	D1135V/E1219K/R1335Q/T1337R	
4 VQR + E1219V	D1135V/E1219V/R1335Q/T1337R	
5 VQR + N1317K	D1135V/N1317K/R1335Q/T1337R	
6 VRQR + N1317K	D1135V/G1218R/N1317K/R1335Q/T1337R	
7 VQR + G1104K	G1104R/D1135V/R1335Q/T1337R	
8 VRQR + G1104K	G1104R/D1135V/G1218R/R1335Q/T1337R	
9 VQR + S1109T	S1109T/D1135V/R1335Q/T1337R	
10 VRQR + S1109T	S1109T/D1135V/G1218R/R1335Q/T1337R	
11 NQR + S1136N	D1135N/S1136N/R1335Q/T1337R	
12 NRQR + S1136N	D1135N/S1136N/G1218R/R1335Q/T1337R	
13 VQR	D1135V/R1335Q/T1337R	
14 VRQR	D1135V/G1218R/R1335Q/T1337R	

[0181] Taken together, these results suggest that including additional mutations in the SpCas9-VQR variant can improve activity against sites that contain NGAN PAMs, specifically sites that contain NGAH PAMs.

REFERENCES

[0182] 1. Sander, J. D. & Joung, J. K. CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol* 32, 347-355 (2014).

[0183] 2. Hsu, P. D., Lander, E. S. & Zhang, F. Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157, 1262-1278 (2014).

[0184] 3. Doudna, J. A. & Charpentier, E. Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* 346, 1258096 (2014).

[0185] 4. Barrangou, R. & May, A. P. Unraveling the potential of CRISPR-Cas9 for gene therapy. *Expert Opin Biol Ther* 15, 311-314 (2015).

[0186] 5. Jinek, M. et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816-821 (2012).

[0187] 6. Sternberg, S. H., Redding, S., Jinek, M., Greene, E. C. & Doudna, J. A. DNA interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature* 507, 62-67 (2014).

[0188] 7. Hsu, P. D. et al. DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol* 31, 827-832 (2013).

[0189] 8. Tsai, S. Q. et al. GUIDE-seq enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases. *Nat Biotechnol* 33, 187-197 (2015).

[0190] 9. Hou, Z. et al. Efficient genome engineering in human pluripotent stem cells using Cas9 from *Neisseria meningitidis*. *Proc Natl Acad Sci USA* (2013).

[0191] 10. Fonfara, I. et al. Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems. *Nucleic Acids Res* 42, 2577-2590 (2014).

[0192] 11. Esvelt, K. M. et al. Orthogonal Cas9 proteins for RNA-guided gene regulation and editing. *Nat Methods* 10, 1116-1121 (2013).

[0193] 12. Cong, L. et al. Multiplex genome engineering using CRISPR/Cas systems. *Science* 339, 819-823 (2013).

[0194] 13. Horvath, P. et al. Diversity, activity, and evolution of CRISPR loci in *Streptococcus thermophilus*. *J Bacteriol* 190, 1401-1412 (2008).

[0195] 14. Anders, C., Niewoehner, O., Duerst, A. & Jinek, M. Structural basis of PAM-dependent target DNA recognition by the Cas9 endonuclease. *Nature* 513, 569-573 (2014).

[0196] 15. Reyon, D. et al. FLASH assembly of TALENs for high-throughput genome editing. *Nat Biotechnol* 30, 460-465 (2012).

[0197] 16. Fu, Y. et al. High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells. *Nat Biotechnol* 31, 822-826 (2013).

[0198] 17. Chen, Z. & Zhao, H. A highly sensitive selection method for directed evolution of homing endonucleases. *Nucleic Acids Res* 33, e154 (2005).

[0199] 18. Doyon, J. B., Pattanayak, V., Meyer, C. B. & Liu, D. R. Directed evolution and substrate specificity profile of homing endonuclease I-SceI. *J Am Chem Soc* 128, 2477-2484 (2006).

[0200] 19. Jiang, W., Bikard, D., Cox, D., Zhang, F. & Marraffini, L. A. RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat Biotechnol* 31, 233-239 (2013).

[0201] 20. Mali, P. et al. RNA-guided human genome engineering via Cas9. *Science* 339, 823-826 (2013).

[0202] 21. Hwang, W. Y. et al. Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol* 31, 227-229 (2013).

[0203] 22. Chylinski, K., Le Rhun, A. & Charpentier, E. The tracrRNA and Cas9 families of type II CRISPR-Cas immunity systems. *RNA Biol* 10, 726-737 (2013).

[0204] 23. Kleinstiver, B. P., Fernandes, A. D., Gloor, G. B. & Edgell, D. R. A unified genetic, computational and experimental framework identifies functionally relevant residues of the homing endonuclease I-BmoI. *Nucleic Acids Res* 38, 2411-2427 (2010).

[0205] 24. Gagnon, J. A. et al. Efficient mutagenesis by Cas9 protein-mediated oligonucleotide insertion and large-scale assessment of single-guide RNAs. *PLoS One* 9, e98186 (2014).

OTHER EMBODIMENTS

[0206] It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

 SEQUENCE LISTING

The patent application contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<http://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US20160312199A1>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

What is claimed is:

1. An isolated *Staphylococcus aureus* Cas9 (SaCas9) protein with mutations at one or more of the following positions: E782, N968, and/or R1015.

2. The isolated protein of claim 1, comprising a sequence that is at least 80% identical to the amino acid sequence of SEQ ID NO:2.

3. The isolated protein of claim 1, comprising one or more of the following mutations: R1015Q, R1015H, E782K, N968K, E735K, K929R, A1021T, K1044N, E782K/N968K/R1015H (KKH variant); E782K/K929R/R1015H (KRH variant); or E782K/K929R/N968K/R1015H (KRKH variant).

4. The isolated protein of claim 1, further comprising one or more mutations that decrease nuclease activity selected from the group consisting of mutations at one or more of D10, D556, H557, and/or N580.

5. The isolated protein of claim 4, wherein the mutations are one or more of D10A, D556A, H557A, or N580A.

6. The isolated protein of claim 4, wherein the mutations are:

- (i) D10A/H557A, or
- (ii) D10A/D556A/H557A/N580A.

7. A fusion protein comprising the isolated protein of claim 1, fused to a heterologous functional domain, with an optional intervening linker, wherein the linker does not interfere with activity of the fusion protein.

8. The fusion protein of claim 7, wherein the heterologous functional domain is a transcriptional activation domain.

9. The fusion protein of claim 8, wherein the transcriptional activation domain is from VP64 or NF- κ B p65.

10. The fusion protein of claim 7, wherein the heterologous functional domain is a transcriptional silencer or transcriptional repression domain.

11. The fusion protein of claim 10, wherein the transcriptional repression domain is a Krueppel-associated box (KRAB) domain, ERF repressor domain (ERD), or mSin3A interaction domain (SID).

12. The fusion protein of claim 10, wherein the transcriptional silencer is Heterochromatin Protein 1 (HP1).

13. The fusion protein of claim 7, wherein the heterologous functional domain is an enzyme that modifies the methylation state of DNA.

14. The fusion protein of claim 13, wherein the enzyme that modifies the methylation state of DNA is a DNA methyltransferase (DNMT) or a TET protein.

15. The fusion protein of claim 14, wherein the TET protein is TET1.

16. The fusion protein of claim 7, wherein the heterologous functional domain is an enzyme that modifies a histone subunit.

17. The fusion protein of claim 7, wherein the enzyme that modifies a histone subunit is a histone acetyltransferase (HAT), histone deacetylase (HDAC), histone methyltransferase (HMT), or histone demethylase.

18. The fusion protein of claim 7, wherein the heterologous functional domain is a biological tether.

19. The fusion protein of claim 18, wherein the biological tether is MS2, Csy4 or lambda N protein.

20. The fusion protein of claim 7, wherein the heterologous functional domain is FokI.

21. An isolated nucleic acid encoding the protein of claim 1.

22. A vector comprising the isolated nucleic acid of claim 21.

23. The vector of claim 22, wherein the isolated nucleic acid of claim 21 is operably linked to one or more regulatory domains for expressing an isolated *Staphylococcus aureus* Cas9 (SaCas9) protein with mutations at one or more of the following positions: E782, N968, and/or R1015.

24. A host cell comprising the nucleic acid of claim 23.

25. A method of altering the genome of a cell, the method comprising expressing in the cell, or contacting the cell with, the isolated protein or fusion protein of claim 1, and a guide RNA having a region complementary to a selected portion of the genome of the cell.

26. The method of claim 25, wherein the isolated protein or fusion protein comprises one or more of a nuclear localization sequence, cell penetrating peptide sequence, and/or affinity tag.

27. The method of claim 25, wherein the cell is a stem cell.

28. The method of claim 27, wherein the cell is an embryonic stem cell, mesenchymal stem cell, or induced pluripotent stem cell; is in a living animal; or is in an embryo.

29. A method of altering a double stranded DNA (dsDNA) molecule, the method comprising contacting the dsDNA molecule with the isolated protein or fusion protein of claim 1, and a guide RNA having a region complementary to a selected portion of the dsDNA molecule.

30. The method of claim 29, wherein the dsDNA molecule is in vitro.

* * * * *