

(19) AUSTRALIAN PATENT OFFICE

- (54) Title
Pharmaceutical composition comprising a bispecific antibody for EpCAM
- (51)⁶ International Patent Classification(s)
C07K 16/30 (2006.01) ^{8BMEP} **C07K**
C07K 16/28 (2006.01) 16/28
C07K 16/30 20060101ALI2005100
20060101AFI2005100 ^{8BMEP}
PCT/EP2004/005687
- (21) Application No: 2004242847 (22) Application Date: 2004 .05 .26
- (87) WIPO No: W004/106383
- (30) Priority Data
- | (31) Number | (32) Date | (33) Country |
|--------------|--------------|--------------|
| 03 01 2133.9 | 2003 .05 .31 | EP |
| 03 01 2134.7 | 2003 .05 .31 | EP |
- (43) Publication Date : 2004 .12 .09
- (71) Applicant(s)
Micromet AG
- (72) Inventor(s)
Wolf, Andreas; Bauerle, Patrick; Lenkkeri-Schutz, Ulla; Kufer, Peter; Offner, Sonja ; Raum, Tobias; Berry, Meera; Kohleisen, Birgit; Brischwein, Klaus
- (74) Agent/Attorney
Griffith Hack, Level 3 509 St Kilda Road, Melbourne, VIC, 3004
- (56) Related Art
WO 1999/025818 A1
Flieger, D. et al. 2000 Can. Immunol. Immunother. Vol. 49. p. 441-448
WO 2000/006605 A2
WO 2000/003016 A1
WO 2001/071005 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
9 December 2004 (09.12.2004)

PCT

(10) International Publication Number
WO 2004/106383 A1

- (51) International Patent Classification⁷: **C07K 16/30**,
16/28, A61K 39/395, 16/46
- (21) International Application Number:
PCT/EP2004/005687
- (22) International Filing Date: 26 May 2004 (26.05.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
03 01 2134.7 31 May 2003 (31.05.2003) EP
03 01 2133.9 31 May 2003 (31.05.2003) EP
- (71) Applicant (for all designated States except US): **MI-
CROMET AG** [DE/DE]; Staffelseestrasse 2, 81477
München (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **KUFER, Peter**
[DE/DE]; Am Kapellenacker 13, 85368 Moosburg (DE).
BERRY, Meera [IN/DE]; Steingasse 10, 89073 Ulm (DE).
OFFNER, Sonja [DE/DE]; Zamboniinstr. 23, 80638
München (DE). **BRISCHWEIN, Klaus** [DE/DE]; Licht-
entfellerstrasse 18, 81243 Munich (DE). **WOLF, Andreas**
[DE/DE]; Einsteinstr. 10, 82152 Planegg (DE). **RAUM,**
Tobias [DE/DE]; Zentnerstrasse 20, 80789 München (DE).
KOHLSEISEN, Birgit [DE/DE]; Staffelseestr. 2, 81477
München (DE). **LENKKERI-SCHÜTZ, Ulla** [FI/DE];
Nelkenstrasse 23, 85386 Eching (DE). **BÄUERLE,**
- (74) Agent: **VOSSIUS & PARTNER**; Siebertstrasse 4, 81675
München (DE).
- (81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GI, GM, GR, GU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PI, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.
- (84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 2004/106383 A1

(54) Title: PHARMACEUTICAL COMPOSITION COMPRISING A BISPECIFIC ANTIBODY FOR EPCAM

(57) Abstract: The present invention provides a pharmaceutical composition comprising a bispecific single chain antibody construct. Said bispecific single chain antibody construct is characterized to comprise or consist of at least two domains, whereby one of said at least two domains specifically binds to human EPCAM and comprises at least one CDR-H3 region comprising the amino acid sequence NXID antigen; and a second domain binds to human CD3 antigen. The invention further provides a process for the production of the pharmaceutical composition of the invention, a method for the prevention, treatment or amelioration of a tumorous disease and the use of the disclosed bispecific single chain antibody construct and corresponding means in the prevention, treatment or amelioration of a tumorous disease.

**PHARMACEUTICAL COMPOSITION COMPRISING
A BISPECIFIC ANTIBODY SPECIFIC FOR EPCAM**

5

The invention relates to a pharmaceutical composition comprising a bispecific single chain antibody construct. Said bispecific single chain antibody construct is characterized to comprise or consist of at least two domains, whereby one of said at least two domains specifically binds to human EpCAM antigen and comprises at least one CDR-H3 region comprising the amino acid sequence NXD and a second domain binds to human CD3 antigen. The invention further provides a process for the production of the pharmaceutical composition of the invention, a method for the prevention, treatment or amelioration of a tumorous disease and the use of the disclosed bispecific single chain antibody construct and corresponding means in the prevention, treatment or amelioration of a tumorous disease.

A variety of documents is cited throughout this specification. The disclosure content of said documents is herewith incorporated by reference.

Epithelial cell adhesion molecule (EpCAM, also called 17-1A antigen, KSA, EGP40, GA733-2, ks1-4 or esa) is a 40-kDa membrane-integrated glycoprotein of 314 amino acids with specific expression in certain epithelia and on many human carcinomas (reviewed in Balzar, J. Mol. Med. 1999, 77, 699-712). EpCAM was discovered and subsequently cloned through its recognition by the murine monoclonal antibody 17-1A/edrecolomab (Goettlinger, Int J Cancer. 1986; 38, 47-53 and Simon, Proc. Natl. Acad. Sci. USA. 1990; 87, 2755-2759). Monoclonal antibody 17-1A was generated by immunization of mice with human colon carcinoma cells (Koprowski, Somatic Cell Genet. 1979, 5, 957-971).

The EGF-like repeats of EpCAM were shown to mediate lateral and reciprocal interactions in homophilic cell adhesion (Balzar, Mol. Cell. Biol. 2001, 21, 2570-2580) and, for that reason, is predominantly located between epithelial cells (Litvinov, J Cell Biol. 1997, 139, 1337-1348, Balzar, J Mol Med. 1999, 77, 699-712 and Trebak, J Biol. Chem. 2001, 276, 2299-2309). EpCAM serves to adhere epithelial cells in an oriented and highly ordered fashion (Litvinov, J Cell Biol. 1997, 139, 1337-1348). Data from experiments with transgenic mice and rats expressing human EpCAM on their epithelia suggest that EpCAM on normal tissue may however not be accessible to systemically administered antibody (McLaughlin, Cancer Immunol. Immunother., 1999, 48, 303-311). Upon malignant transformation of epithelial cells the rapidly growing tumor cells are abandoning the high cellular order of epithelia. Consequently, the surface distribution of EpCAM becomes less restricted and the molecule better exposed on tumor cells. Due to their epithelial cell origin, tumor cells from most carcinomas still express EpCAM on their surface.

15

In vivo, expression of EpCAM is related to increased epithelial proliferation and negatively correlates with cell differentiation (for review see Balzar, 1999, J. Mol. Med. 77, 699-712). Expression of EpCAM, as detected by immunohistochemistry using anti-EpCAM monoclonal antibodies, is essentially seen with all major carcinomas (reviewed in Balzar, J Mol Med. 1999, 77, 699-712). Best EpCAM expression was observed with non-small cell lung cancer (De Bree, Nucl Med Commun. 1994, 15, 613-27) and prostate cancer (Zhang, Clin Cancer Res. 1998, 4, 295-302) where 100% of tumor patient samples showed positive EpCAM staining. In these studies, EpCAM is also reported to homogeneously stained tumor tissues indicating that the antigen is expressed on a large proportion of cells of a given tumor. Because of its widespread expression, EpCAM is referred to as a "pan-carcinoma" antigen.

EpCAM has been shown in various studies to be beneficial in diagnosis and therapy of various carcinomas. Furthermore, in many cases, tumor cells were

30

observed to express EpCAM to a much higher degree than their parental epithelium or less aggressive forms of said cancers. For example, EpCAM expression was shown to be significantly higher on neoplastic tissue and in adenocarcinoma than on normal prostate epithelium (n=76; p<0.0001), suggesting
5 that increased EpCAM expression represents an early event in the development of prostate cancer (Poczatek, J Urol., 1999, 162, 1462-1644). In addition, in the majority of both squamous and adenocarcinomas of the cervix a strong EpCAM expression correlates with an increased proliferation and the disappearance of markers for terminal differentiation (Litvinov, Am. J. Pathol. 1996, 148, 865-75).
10 One example is breast cancer where overexpression of EpCAM on tumor cells is a predictor of survival (Gastl, Lancet. 2000, 356, 1981-1982). Furthermore, EpCAM has been described as a marker for the detection of disseminated tumor cells in patients suffering from squamous cell carcinoma of the head, neck and lung (Chaubal, Anticancer Res 1999, 19, 2237-2242, Piyathilake, Hum Pathol. 2000, 31,
15 482-487). Normal squamous epithelium, as found in epidermis, oral cavity, epiglottis, pharynx, larynx and esophagus did not significantly express EpCAM (Quak, Hybridoma, 1990, 9, 377-387).

In addition to the above-mentioned carcinomas, EpCAM has been shown to be
20 expressed on the majority of primary, metastatic, and disseminated NSCLC (non small cell lung cancer cells) (Passlick, Int J Cancer, 2000, 87, 548-552), on gastric and gastro-oesophageal junction adenocarcinomas (Martin, J Clin Pathol 1999, 52, 701-4) and in cell lines derived from colorectal, pancreatic carcinomas and breast carcinomas (Szala, Proc Natl Acad Sci U S A 1990, 87, 3542-6, Packeisen,
25 Hybridoma, 1999, 18, 37-40).

Clinical trials have shown that the use of antibodies directed against 17-1A (EpCAM) for treatment of patients with surgically completely resected colorectal carcinoma leads to a significant benefit concerning the overall survival and the frequency of distant metastasis (Riethmüller, Lancet, 1994, 343, 1177-1183).
30 Murine monoclonal antibody against EpCAM was found to reduce the 5-year

mortality (Riethmüller, Lancet, 1994, 343, 1177-1183) and also the 7-year mortality (Riethmüller, Proceedings of the American Society of Clinical Oncology, 1996, 15, 444) of patients with minimal residual disease. Example of murine monoclonal antibody recognizing EpCAM is Edrecolomab (Panorex) (Koprowski, Somatic Cell
5 Genet. 1979, 5, 957-971 and Herlyn, Cancer Res., 1980, 40, 717-721). However, the first administration of Panorex during adjuvant immunotherapy of colon cancer led to the development and exacerbation of Wegener's granulomatosis suggesting that mAb 17-1A should be applied cautiously in a patient with autoimmune disease (Franz, Onkologie, 2000, 23, 472-474). The limitations of Panorex are the rapid
10 formation of human anti-mouse antibodies (HAMA), the limited ability to interact by its murine IgG2a Fc-portion with human immune effector mechanisms and the short half-life in circulation (Frodin, Cancer Res., 1990, 50, 4866-4871). Furthermore, the murine antibody caused immediate-type allergic reactions and anaphylaxis upon repeated injection in patients (Riethmüller, Lancet. 1994, 343,
15 1177-1183, Riethmüller, J Clin Oncol., 1998, 16, 1788-1794 and Mellstedt, Annals New York Academy of Sciences. 2000, 910, 254-261).

Humanized anti-EpCAM antibody called 3622W94 resulted in pancreatitis and increased serum levels of amylase, as being indicative for damage of pancreas
20 epithelium, which were a dose-limiting toxicity of this high-affinity anti-EpCAM monoclonal antibody (LoBuglio, Proceedings of the American Society of Clinical Oncology (Abstract). 1997, 1562 and Khor, Proceedings of the American Society of Clinical Oncology (Abstract), 1997, 847).

25 Bispecific antibodies comprising a region directed against EpCAM and a region directed against CD3 have also been described. The authors of Möller & Reisfeld 1991 Cancer Immunol. Immunother. 33:210-216 describe the construction of two different bispecific antibodies by fusing a hybridoma producing monoclonal antibody against EpCAM with either of the two hybridomas OKT3 and 9.3.
30 Furthermore, Kroesen, Cancer Research, 1995, 55:4409-4415 describe a

quadroma bispecific monoclonal antibodies against CD3 (BIS-1) and EpCAM.

Other examples of bispecific antibodies against EpCAM comprise the bispecific antibody, BiUII, (anti-CD3 (rat IgG2b) x anti-EpCAM (mouse IgG2a)) a complete Ig molecule which also binds and activates Fc-receptor positive accessory cells (like monocytes/macrophages, NK cells and dendritic cells) through its Fc-region (Zeidler, J. Immunol., 1999, 163:1247-1252) and an anti-EpCAMxanti-CD3 bispecific antibody in the arrangement $V_{L17-1A}-V_{H17-1A}-V_{Hanti-CD3}-V_{Lanti-CD3}$ (Mack, Proc. Natl. Acad. Sci., 1995, 92:7021-7025).

10

In addition, other formats of antibody constructs comprising EpCAM have been described; e.g. a bispecific diabody having the structure $V_{H anti-CD3}-V_{L anti-EpCAM}-V_{H anti-EpCAM}-V_{L anti-CD3}$ (Helfrich, Int. J. Cancer, 1998, 76:232-239) and a trispecific antibody having two different tumour antigen specificities (two antigen binding regions which bind two different antigens on a tumour cell) and which may have a further specificity for an antigen localized on an effector cell (DE 195 31 348).

15

There exist various descriptions in the prior art of using phage display technology to identify antibodies or fragments thereof, which specifically bind to the human EpCAM antigen (De Kruif JMB, 1995, 248:97-105, WO 99/25818). However, it has been extremely difficult to identify antibodies against EpCAM, which show cytotoxic activity sufficient for therapeutic applications in a bispecific format.

20

It is therefore an aim of the present invention to provide a bispecific single chain molecule with a binding domain specific for EpCAM with strong cytotoxic activity mediated by target specific activation of T cells.

25

Thus, the technical problem underlying the present invention was to provide means

and methods for the generation of well tolerated and convenient medicaments for the treatment and or amelioration of tumorous diseases.

The solution to said technical problem is achieved by providing the embodiments characterized in the claims.

It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

SUMMARY

A first aspect provides a pharmaceutical composition comprising a bispecific single chain antibody construct, wherein said construct comprises or consists of at least two domains, wherein one of said domains binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region comprising the amino acid sequence NXD preferably in position 102 to 104 of SEQ ID NOs: 80, 88 and 96, or preferably in position 106 to 108 of SEQ ID NOs: 84 and 92, wherein X is an aromatic amino acid.

A second aspect provides a pharmaceutical composition comprising a bispecific single chain antibody construct, wherein said construct comprises or consists of at least two domains, wherein one of said at least two domains specifically binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region of at least 9 amino acid residues and wherein said binding domain specific for EpCAM has a K_D value of more than 5×10^{-9} M.

A third aspect provides a pharmaceutical composition comprising a nucleic acid sequence encoding a bispecific single chain antibody construct as defined in the first

6a

or second aspect.

A fourth aspect provides a pharmaceutical composition comprising a vector which comprises a nucleic acid sequence as defined in the third aspect.

5

A fifth aspect provides a pharmaceutical composition comprising a host transformed or transfected with a vector as defined in the fourth aspect.

10

A sixth aspect provides a process for the production of a pharmaceutical composition according to the first, second, third, fourth, or fifth aspect, said process comprising culturing a host defined in the fifth aspect under conditions allowing the expression of the bispecific single chain antibody construct as defined in the first or second aspect and recovering the produced bispecific single chain antibody construct from the culture.

15

A seventh aspect provides use of a bispecific single chain antibody construct as defined in the first or second aspect, a nucleic acid sequence as defined in the third aspect, a vector as defined in the fourth aspect, a host as defined in the fifth aspect and/or produced by a process according to the sixth aspect for the preparation of a pharmaceutical composition for the prevention, treatment or amelioration of a tumorous disease.

20

An eighth provides a method for the prevention, treatment or amelioration of a tumorous disease, comprising the step of administering to a subject in need of such a prevention, treatment or amelioration a pharmaceutical composition of the first, second, third, fourth, or fifth aspect.

25

A ninth aspect provides a kit comprising a bispecific single chain antibody construct as defined in the first or second aspect, a nucleic acid sequence as defined in the third aspect, a vector as defined in the fourth aspect, a host as defined in the fifth aspect

30

6b

and/or produced by a process according to the sixth aspect.

Accordingly, the present invention relates to a composition, preferably a pharmaceutical composition, comprising a bispecific single chain antibody construct, whereby said construct comprises or consists of at least two binding domains, whereby one of said domains binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region comprising the amino acid sequence NXD preferably in position 102 to 104 of SEQ ID NOs: 80, 88 and 96, or preferably in position 106 to 108 of SEQ ID NOs: 84 and 92, wherein X is an aromatic amino acid.

Preferably or alternatively, the present invention relates to a composition, preferably a pharmaceutical composition, comprising a bispecific single chain antibody construct, whereby said construct comprises or consists of at least two domains, whereby one of said at least two domains specifically binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region of least 9 amino acid residues and wherein said binding domain specific for EpCAM has a K_D value of more than 5×10^{-9} M.

DESCRIPTION

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

In accordance with this invention, the term "pharmaceutical composition" relates to a composition for administration to a patient, preferably a human patient. In a preferred embodiment, the pharmaceutical composition comprises a composition for parenteral,

2029358_1 (CHMediCare) P58001 AU 11-May-11

2004242847 11 May 2011

6c

transdermal, intraluminal, intra-arterial, intrathecal or intravenous administration or for direct injection into the tumor. It is in particular envisaged that said pharmaceutical composition is administered to a patient via infusion or injection. Administration of the suitable compositions may be effected by different

ways, e.g., by intravenous, subcutaneous, intraperitoneal, intramuscular, topical or intradermal administration. The pharmaceutical composition of the present invention may further comprise a pharmaceutically acceptable carrier. Examples of suitable pharmaceutical carriers are well known in the art and include phosphate
5 buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions, etc. Compositions comprising such carriers can be formulated by well known conventional methods. These pharmaceutical compositions can be administered to the subject at a suitable dose. The dosage regimen will be determined by the attending physician and clinical
10 factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A preferred dosage for administration might be in the range of 0.24 μ g to 48 mg, preferably
15 0.24 μ g to 24 mg, more preferably 0.24 μ g to 2.4 mg, even more preferably 0.24 μ g to 1.2 mg and most preferably 0.24 μ g to 240 μ g units per kilogram of body weight per day. Particularly preferred dosages are recited herein below. Progress can be monitored by periodic assessment. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10^6 to 10^{12} copies of the
20 DNA molecule. The compositions of the invention may be administered locally or systematically. Administration will generally be parenteral, e.g., intravenous; DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target site or by catheter to a site in an artery. In an preferred embodiment, the pharmaceutical composition is administered subcutaneously and
25 in an even more preferred embodiment intravenously. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or
30 suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated

Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishes, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. In addition, the pharmaceutical composition of the present invention might comprise proteinaceous carriers, like, e.g., serum albumine or immunoglobuline, preferably of human origin. It is envisaged that the pharmaceutical composition of the invention might comprise, in addition to the proteinaceous bispecific single chain antibody constructs or nucleic acid molecules or vectors encoding the same (as described in this invention), further biologically active agents, depending on the intended use of the pharmaceutical composition. Such agents might be drugs acting on the gastro-intestinal system, drugs acting as cytostatica, drugs preventing hyperurikemia, agents such as T-cell co-stimulatory molecules or cytokines, drugs inhibiting immune reactions (e.g. corticosteroids) and/or drugs acting on the circulatory system, e.g. on the blood pressure, known in the art. Possible indications for administration of the composition(s) of the invention are tumorous diseases especially epithelial cancers/carcinomas such as breast cancer, colon cancer, prostate cancer, head and neck cancer, skin cancer, cancers of the genito-urinary tract, e.g. ovarian cancer, endometrial cancer, cervix cancer and kidney cancer, lung cancer, gastric cancer, cancer of the small intestine, liver cancer, pancreas cancer, gall bladder cancer, cancers of the bile duct, esophagus cancer, cancer of the salivatory glands and cancer of the thyroid gland. The administration of the composition(s) of the invention is especially indicated for minimal residual disease preferably early solid tumor, advanced solid tumor or metastatic solid tumor, which is characterized by the local and non-local reoccurrence of the tumor caused by the survival of single cells. The invention further envisages the co-administration protocols with other compounds, e.g. bispecific antibody constructs, targeted toxins or other compounds, which act via T cells. The clinical regimen for co-administration of the inventive compound(s) may encompass co-administration at the same time, before or after the administration of the other component.

- A possible approach to demonstrate the efficacy/activity of the inventive constructs is an in vivo model like mouse. Suitable models may be transgenic and chimeric mouse models. Mouse models expressing human CD3 and human EpCAM, a chimeric mouse model expressing murine CD3 and into which tumour cells expressing human EpCAM can be transfected and chimeric mouse models comprising nude mice into which human tumours expressing EpCAM can be transplanted or tumour cells expressing human EpCAM can be injected and, additionally, human PBMCs are injected. The term "bispecific single chain antibody construct" relates to a construct comprising two antibody derived binding domains.
- 5 One of said binding domains consists of variable regions (or parts thereof) of an antibody, antibody fragment or derivative thereof, capable of specifically binding to/interacting with human EpCAM antigen (target molecule 1). The second binding domain consists of variable regions (or parts thereof) of an antibody, antibody fragment or derivative thereof, capable of specifically binding to/interacting with
- 10 human CD3 antigen (target molecule 2). As will be detailed below, a part of a variable region may be at least one CDR ("Complementary determining region"), most preferably at least the CDR3 region. Said two domains/regions in the single chain antibody construct are preferably covalently connected to one another as a single chain. This connection can be effected either directly (domain 1 [specific for the CD3 antigen] – domain 2 [specific for the EpCAM antigen] or domain 1 [specific for the EpCAM antigen] – domain 2 [specific for the CD3 antigen]) or through an additional polypeptide linker sequence (domain1 – linker sequence – domain2). In the event that a linker is used, this linker is preferably of a length and sequence sufficient to ensure that each of the first and second domains can, independently
- 15 from one another, retain their differential binding specificities. Most preferably and as documented in the appended examples, the "bispecific single chain antibody construct" to be employed in the pharmaceutical composition of the invention is a bispecific single chain Fv (scFv). Bispecific single chain molecules are known in the art and are described in WO 99/54440, Mack, J. Immunol. (1997), 158, 3965-3970, Mack, PNAS, (1995), 92, 7021-7025, Kufer, Cancer Immunol. Immunother., (1997), 45, 193-197, Löffler, Blood, (2000), 95, 6, 2098-2103 and Brühl, J.
- 20
- 25
- 30

Immunol., (2001), 166, 2420-2426 . A particularly preferred molecular format of the invention provides a polypeptide construct wherein the antibody-derived region comprises one V_H and one V_L region. The intramolecular orientation of the V_H-domain and the V_L-domain, which are linked to each other by a linker-domain, in
5 the scFv format is not decisive for the recited bispecific single chain constructs. Thus, scFvs with both possible arrangements (V_H-domain – linker domain – V_L-domain; V_L-domain – linker domain – V_H-domain) are particular embodiments of the recited bispecific single chain construct.
The antibody construct may also comprise additional domains, e.g. for the isolation
10 and/or preparation of recombinantly produced constructs.

A corresponding format for a bispecific single chain antibody construct is described in the appended example 1.

The term "single-chain" as used in accordance with the present invention means
15 that said first and second domain of the bispecific single chain construct are covalently linked, preferably in the form of a co-linear amino acid sequence encodable by a single nucleic acid molecule.

The term "binding to/interacting with" as used in the context with the present invention defines a binding/interaction of at least two "antigen-interaction-sites" with
20 each other. The term "antigen-interaction-site" defines, in accordance with the present invention, a motif of a polypeptide which shows the capacity of specific interaction with a specific antigen or a specific group of antigens. Said binding/interaction is also understood to define a "specific recognition". The term "specifically recognizing" means in accordance with this invention that the antibody
25 molecule is capable of specifically interacting with and/or binding to at least two amino acids of each of the human target molecule as defined herein. Said term relates to the specificity of the antibody molecule, i.e. to its ability to discriminate between the specific regions of the human target molecule as defined herein. The specific interaction of the antigen-interaction-site with its specific antigen may result
30 in an initiation of a signal, e.g. due to the induction of a change of the conformation of the antigen, an oligomerization of the antigen, etc. Further, said binding may be

exemplified by the specificity of a "key-lock-principle". Thus, specific motifs in the amino acid sequence of the antigen-interaction-site and the antigen bind to each other as a result of their primary, secondary or tertiary structure as well as the result of secondary modifications of said structure. The specific interaction of the antigen-interaction-site with its specific antigen may result as well in a simple binding of said site to the antigen.

The term "specific interaction" as used in accordance with the present invention means that the bispecific single chain construct does not or essentially does not cross-react with (poly)peptides of similar structures. Cross-reactivity of a panel of bispecific single chain construct under investigation may be tested, for example, by assessing binding of said panel of bispecific single chain construct under conventional conditions (see, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1988 and *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1999) to the (poly)peptide of interest as well as to a number of more or less (structurally and/or functionally) closely related (poly)peptides. Only those antibodies that bind to the (poly)peptide/protein of interest but do not or do not essentially bind to any of the other (poly)peptides are considered specific for the (poly)peptide/protein of interest. Examples for the specific interaction of an antigen-interaction-site with a specific antigen comprise the specificity of a ligand for its receptor. Said definition particularly comprises the interaction of ligands which induce a signal upon binding to its specific receptor. Examples for corresponding ligands comprise cytokines which interact/bind with/to its specific cytokine-receptors. Also particularly comprised by said definition is the binding of an antigen-interaction-site to antigens like antigens of the selectin family, integrins and of the family of growth factors like EGF. An other example for said interaction, which is also particularly comprised by said definition, is the interaction of an antigenic determinant (epitope) with the antigenic binding site of an antibody.

The term "binding to/interacting with" may also relate to a conformational epitope, a structural epitope or a discontinuous epitope consisting of two regions of the human target molecules or parts thereof. In context of this invention, a

conformational epitope is defined by two or more discrete amino acid sequences separated in the primary sequence which come together on the surface of the molecule when the polypeptide folds to the native protein (Sela, (1969) Science 166, 1365 and Laver, (1990) Cell 61, 553-6).

5 The term "discontinuous epitope" means in context of the invention non-linear epitopes that are assembled from residues from distant portions of the polypeptide chain. These residues come together on the surface of the molecule when the polypeptide chain folds into a three-dimensional structure to constitute a conformational/structural epitope.

10 The constructs of the present invention are also envisaged to specifically bind to/interact with a conformational/structural epitope(s) composed of and/or comprising the two regions of the human CD3 complex described herein or parts thereof as disclosed herein below.

Accordingly, specificity can be determined experimentally by methods known in the art and methods as disclosed and described herein. Such methods comprise, but
15 are not limited to Western blots, ELISA-, RIA-, ECL-, IRMA-, EIA-tests and peptide scans.

The term "antibody fragment or derivative thereof" relates to single chain antibodies, or fragments thereof, synthetic antibodies, antibody fragments, such as
20 Fab, a F(ab₂)', Fv or scFv fragments etc., or a chemically modified derivative of any of these. Antibodies to be employed in accordance with the invention or their corresponding immunoglobulin chain(s) can be further modified using conventional techniques known in the art, for example, by using amino acid deletion(s), insertion(s), substitution(s), addition(s), and/or recombination(s) and/or any other
25 modification(s) (e.g. posttranslational and chemical modifications, such as glycosylation and phosphorylation) known in the art either alone or in combination. Methods for introducing such modifications in the DNA sequence underlying the amino acid sequence of an immunoglobulin chain are well known to the person skilled in the art; see, e.g., Sambrook (1989), loc. cit.

30 The term "(poly)peptide" as used herein describes a group of molecules which comprise the group of peptides, as well as the group of polypeptides. The group of

peptides is consisting of molecules with up to 30 amino acids, the group of polypeptides is consisting of molecules with more than 30 amino acids.

The term "antibody fragment or derivative thereof" particularly relates to (poly)peptide constructs comprising at least one CDR.

- 5 Fragments or derivatives of the recited antibody molecules define (poly)peptides which are parts of the above antibody molecules and/or which are modified by chemical/biochemical or molecular biological methods. Corresponding methods are known in the art and described inter alia in laboratory manuals (see Sambrook et al.; Molecular Cloning: A Laboratory Manual; Cold Spring Harbor Laboratory Press, 10 2nd edition 1989 and 3rd edition 2001; Gerhardt et al.; Methods for General and Molecular Bacteriology; ASM Press, 1994; Lefkovits; Immunology Methods Manual: The Comprehensive Sourcebook of Techniques; Academic Press, 1997; Golemis; Protein-Protein Interactions: A Molecular Cloning Manual; Cold Spring Harbor Laboratory Press, 2002).

- 15 Bispecific antibodies that specifically recognize the EpCAM antigen and the CD3 antigen are described in the prior art, e.g., in Mack (Proc. Natl. Acad. Sci., 1995, 92:7021-7025).

- 20 As mentioned above, the said variable domains comprised in the herein described bispecific single chain constructs are connected by additional linker sequences. The term "peptide linker" defines in accordance with the present invention an amino acid sequence by which the amino acid sequences of the first domain and the second domain of the defined construct are linked with each other. An essential 25 technical feature of such peptide linker is that said peptide linker does not comprise any polymerization activity. A particularly preferred peptide linker is characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. (Gly)₄Ser, or polymers thereof, i.e. ((Gly)₄Ser)_x. The characteristics of said peptide linker, which comprise the absence of the promotion of secondary structures are known in the art and 30 described e.g. in Dall'Acqua et al. (Biochem. (1998) 37, 9266-9273), Cheadle et al. (Mol Immunol (1992) 29, 21-30) and Raag and Whitlow (FASEB (1995) 9(1), 73-

- 80). Also particularly preferred are peptide linkers which comprise less amino acid residues. An envisaged peptide linker with less than 5 amino acids can comprise 4, 3, 2 or one amino acids. A particularly preferred "single" amino acid in context of said "peptide linker" is Gly. Accordingly, said peptide linker may consist of the single amino acid Gly. Furthermore, peptide linkers which also do not promote any secondary structures are preferred. The linkage of said domains to each other can be provided by, e.g. genetic engineering, as described in the examples. Methods for preparing fused and operatively linked bispecific single chain constructs and expressing them in mammalian cells or bacteria are well-known in the art (e.g. WO 99/54440, Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. 1989 and 1994 or Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).
- 15 The bispecific single chain antibody constructs described herein above and below may be humanized or deimmunized antibody constructs. Methods for the humanization and/or deimmunization of (poly)peptides and, in particular, antibody constructs are known to the person skilled in the art.
- 20 Here it was surprisingly found that domains with specificity for the EpCAM antigen, comprising at least one CDR-H3 region comprising the amino acid sequence NXD (asparagine-X-aspartic acid) preferably in position 102 to 104 of SEQ ID NOs: 80, 88 and 96, or in position 106 to 108 of SEQ ID NOs: 84 and 92, wherein X is an aromatic amino acid are particularly useful in the specific format of a bispecific single chain antibody construct. These bispecific single chain antibody constructs are particularly useful as pharmaceutical compositions since these constructs are advantageous over constructs which do not comprise said amino acids.
- 25 Furthermore, it was surprisingly found that domains with specificity for the EpCAM antigen, comprising at least one CDR-H3 region of at least 9 amino acid residues and having a K_D value of more than 5×10^{-9} M are particularly useful in the specific format of a bispecific single chain antibody construct. These bispecific single chain
- 30

antibody construct are particularly useful as pharmaceutical compositions since these constructs are advantageous over constructs of less than 9 amino acid residues and wherein said binding domain specific for EpCAM has a K_D value of less than 5×10^{-9} M.

5

The prior art constructs are characterized by less advantageous EC_{50} values and/or less efficient or complete purifications as shown in the appended examples. It was in particular surprising that the domain of the single chain constructs with specificity for the CD3 antigen to be employed in accordance with the invention are

10 highly bioactive in N- as well as C-terminal position, wherein in particular arrangements in $V_{H(\text{anti-CD3})}$ - $V_{L(\text{anti-CD3})}$ are preferred. The constructs to be employed in the pharmaceutical composition of the invention are characterized by advantageous production and purification properties as well as by their high bioactivity, i.e. their desired cytotoxic activity. In particular, when the cytotoxic

15 activity of the constructs of the invention were compared with cytotoxic activity of conventional M79xanti-CD3 and HD70xanti-CD3 constructs, the constructs of the invention showed clearly higher bioactivity (Figure 11B). The corresponding high bioactivity is reflected by low to very low EC_{50} values as determined in cytotoxicity tests. The lower the EC_{50} value of the molecule is, the higher cytotoxicity, i.e. the

20 effectivity in the cell lysis, of the construct is higher. On the other hand, the higher the EC_{50} value, the less effective the molecule is in inducing cell lysis. The term " EC_{50} " corresponds, in context of this invention, to EC_{50} values as determined according to the methods known in the art and as illustrated in the appended examples: A standard dose-response curve is defined by four parameters: the

25 baseline response (Bottom), the maximum response (Top), the slope, and the drug concentration that provokes a response halfway between baseline and maximum (EC_{50}). EC_{50} is defined as the concentration of a drug or molecule that provokes a response half way between the baseline (Bottom) and maximum response (Top). A lower K_D value of the constructs of the invention depicts higher binding affinity. E.g.

30 a low K_D of 10^{-9} M shows high binding affinity of the binding construct. On the other hand a high K_D value of e.g. 10^{-6} M relates to lower binding affinity of the binding

domain of the construct.

The percentage of cell lysis (i.e. cytotoxic activity) may be determined by, inter alia, release assays disclosed herein above, for example, ^{51}Cr release assays, LDH-release assays and the like. Most preferably, in context of this invention
 5 fluorochrome release assays is employed as illustrated in the appended examples. Here, strong cytotoxic activity against EpCAM-positive cells (see CHO-EpCAM cells in appended example 3) of the bispecific single chain constructs described herein relates to a molecule comprising EC_{50} values preferably ≤ 500 pg/ml, more preferably ≤ 400 pg/ml, even more preferably ≤ 300 pg/ml, even more preferably $\leq$
 10 250 pg/ml, most preferably ≤ 200 pg/ml ≤ 100 pg/ml, ≤ 50 pg/ml.

The bispecific constructs comprised in the pharmaceutical compositions of the present invention show a surprisingly high cytotoxic activity (preferably in the range of about 10 pg/ml to 170 pg/ml) compared to the prior art M79xanti-CD3 construct
 15 (VL_{17-1A}- VH_{17-1A}- VH_{CD3}- VL_{CD3}; 8628 pg/ml). A skilled person is aware that EC_{50} values may vary depending to the bioactivity assay. Factors affecting EC_{50} value may comprise type of effector cells, activity of effector cells, type of target cells, E:T ratio, incubation time, incubation temperature and other external circumstances. Different EC_{50} values of same constructs in different experiments may be
 20 compared with the EC_{50} values of controls. A construct having high cytotoxic activity according to the invention has at least 2.5 time lower EC_{50} value than the control (at least 2.5 times higher cytotoxicity than the control), preferably at least three times lower EC_{50} value and more preferably at least five times lower EC_{50} value.

25 Furthermore, the constructs of the invention bind EpCAM with a surprisingly high affinity measured by surface plasmon resonance (BIAcore®). The prior art EpCAM and CD3 binding construct M79xanti-CD3 has a K_D of 4×10^{-6} M and the constructs of the invention a K_D of $2,3 \times 10^{-7} - 2,5 \times 10^{-7}$ M.

30 Preferably, the X in said NXD motif is W (tryptophan) or Y (tyrosine).

It is further envisaged that the pharmaceutical composition of the invention comprises a bispecific single chain antibody construct, wherein the CDR-H3 of the EpCAM specific domain comprises at least 9 amino acid residues, preferably at least 14 amino acids. Preferably the CDR-H3 comprises less than 18 amino acids, more preferably less than 15 amino acids. Thus, preferably the CDR-H3 comprises 9 to 17 amino acids, more preferably 9 to 15 amino acids and most preferably 10 or 14 amino acids.

Bispecific single chain antibody construct comprising a corresponding EpCAM specific domain have been surprisingly found to be advantageous in the format of the above described construct over other EpCAM specific domain known in the art. Such effect is demonstrated in appended examples 3, 4 and 5. The prior art EpCAM binding antibody M79 comprises eight amino acids in its CDR-H3 region and does not comprise the sequence NXD (Figure 11A).

The pharmaceutical composition according to the invention may also comprise constructs, wherein said binding domain specific for EpCAM has a K_D value of more than 5×10^{-9} M.

The pharmaceutical composition may additionally be characterized by the feature that said binding domain specific for the CD3 antigen has a K_D value of more than 10^{-7} M.

The K_D value is a physical value defining the tendency of a complex to dissociate. For the binding equilibrium $A+B \leftrightarrow AB$, the dissociation constant is given as the ratio of the two kinetic rate constants k_{off} and k_{on} : $[A][B] (k_{on})/[AB] (k_{off})$. The smaller the dissociation constant the tighter A and B bind to each other. In biological systems a good, specific binder has a dissociation constant in the range of 10^{-9} - 10^{-7} M. K_D can be measured with a number of methods known to the person skilled in the art, e.g. surface plasmon resonance (SPR, e.g. with BIAcore®), analytical ultracentrifugation, isothermal titration calorimetry, fluorescence anisotropy, fluorescence spectroscopy or by radiolabeled ligand binding assays.

The K_D s of the constructs of the invention have been measured using the surface

plasmon resonance (SPR) spectroscopy. The ligand is injected over the immobilized antigen chip surface and the change in optical density on the chip surface upon binding is measured. The change in optical density, monitored by a change in reflection angle, correlates directly to the amount of ligand binding to the chip surface - the biophysical phenomenon used is called surface plasmon resonance.

One of the interaction partners has to be immobilized on the surface of the sensor chip of the apparatus based on surface plasmon resonance (e.g. BIAcore®). The kinetics of association and dissociation of ligand with the immobilized antigen on the chip surface are observed in real time. The binding curves are fitted for kinetic rate constants k_{on} and k_{off} , resulting in an apparent equilibrium dissociation constant (K_D).

It is particularly preferred, that said binding domain specific for EpCAM has a K_D value in a range between 1×10^{-7} and 5×10^{-9} M and said binding domain specific for CD3 has a K_D value in a range between 1×10^{-6} and 5×10^{-9} M.

In a particularly preferred embodiment, the pharmaceutical composition may additionally be characterized by the feature that said binding domain specific for the CD3 antigen has a K_D value of $>$ (more than) 1×10^{-7} M.

The constructs of the invention have the advantage that they may be used a number of times for killing tumour cells since the EpCAM binding part has an affinity with a K_D value of more than 5×10^{-9} M. If the affinity of a bispecific construct for binding an EpCAM-expressing tumour cell is too high, the construct binds one EpCAM expressing tumour cell and remains on its surface even when it has been killed and cannot continue to another tumour cell to be killed. A further advantage of the construct of the invention is that the binding domain specific for EpCAM binds with a high affinity (corresponds to lower K_D value), thus leading the circulating T-cells to the tumour cells marked with the bispecific construct. Therefore, the K_D of the binding domain specific for EpCAM of the bispecific construct is preferably in the range of 10^{-7} - 5×10^{-9} M and the K_D of the binding

domain specific for CD3 is preferably in the range of 10^{-6} - 5×10^{-9} M. In a preferred embodiment, the KD value of the EpCAM binding domain is lower than the KD value of the CD3 binding domain corresponding to a higher affinity of the EpCAM binding domain compared to the CD3 binding domain.

5

Further it is envisaged that the pharmaceutical composition of the invention comprises a bispecific single chain antibody construct, wherein the CDR-H3 of the EpCAM specific domain comprises at least 9 amino acids, preferably at least 14 amino acids. Preferably the CDR-H3 comprises less than 18 amino acids, more preferably less than 15 amino acids. Thus, preferably the CDR-H3 comprises 9 to 17 amino acids, more preferably 9 to 15 amino acids and most preferably 10 or 14 amino acids.

10

In a preferred embodiment of the pharmaceutical composition of the invention the V_H chain of the domain specific for human EpCAM antigen is selected from the group consisting of:

15

(a) an amino acid sequence as shown in any of SEQ ID NO: 80, SEQ ID NO: 84, SEQ ID NO: 88, SEQ ID NO: 92 and SEQ ID NO: 96;

(b) an amino acid sequence encoded by a nucleic acid sequence as shown in SEQ ID NO: 79, SEQ ID NO: 83, SEQ ID NO: 87, SEQ ID NO: 91 and SEQ ID NO: 95;

20

(c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b) under stringent hybridization conditions;

25

(d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).

The term "hybridizing" as used herein refers to polynucleotides/nucleic acid sequences which are capable of hybridizing to the polynucleotides encoding bispecific single chain constructs as defined herein or parts thereof. Therefore, said polynucleotides may be useful as probes in Northern or Southern Blot analysis of

30

RNA or DNA preparations, respectively, or can be used as oligonucleotide primers in PCR analysis dependent on their respective size. Preferably, said hybridizing polynucleotides comprise at least 10, more preferably at least 15 nucleotides in length while a hybridizing polynucleotide of the present invention to be used as a probe preferably comprises at least 100, more preferably at least 200, or most preferably at least 500 nucleotides in length.

It is well known in the art how to perform hybridization experiments with nucleic acid molecules, i.e. the person skilled in the art knows what hybridization conditions s/he has to use in accordance with the present invention. Such hybridization conditions are referred to in standard text books such as Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (2001) N.Y. Preferred in accordance with the present inventions are polynucleotides which are capable of hybridizing to the polynucleotides of the invention or parts thereof, under stringent hybridization conditions.

- 15 "Stringent hybridization conditions" refer, e.g. to an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 x SSC at about 65°C. Also contemplated are nucleic acid
- 20 molecules that hybridize to the polynucleotides of the invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an
- 25 overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1 X SSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X
- 30 SSC). It is of note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to

- suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above,
- 5 due to problems with compatibility.
- The recited nucleic acid molecules may be, e.g., DNA, cDNA, RNA or synthetically produced DNA or RNA or a recombinantly produced chimeric nucleic acid molecule comprising any of those polynucleotides either alone or in combination.
- 10 Preferably said pharmaceutical composition of the invention may comprise a bispecific single chain construct, wherein the V_L chain domains specific for human EpCAM antigen is selected from the group consisting of:
- (a) an amino acid sequence as shown in any of SEQ ID NO: 82, SEQ ID NO: 86, SEQ ID NO: 90, SEQ ID NO: 94 and SEQ ID NO: 98;
 - 15 (b) an amino acid sequence encoded by a nucleic acid sequence as shown in SEQ ID NO: 81, SEQ ID NO: 85, SEQ ID NO: 89, SEQ ID NO: 93 and SEQ ID NO: 97 ;
 - (c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b)
 - 20 under stringent hybridization conditions;
 - (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).
- 25 In a preferred embodiment of the pharmaceutical composition of this invention, the V_H and V_L regions of said human CD3 specific domain are derived from an CD3 specific antibody selected from the group consisting of X35-3, VIT3, BMA030 (BW264/56), CLB-T3/3, CRIS7, YTH12.5, F111-409, CLB-T3.4.2, TR-66, WT32, SPv-T3b, 11D8, XIII-141, XIII-46, XIII-87, 12F6, T3/RW2-8C8, T3/RW2-4B6,
- 30 OKT3D, M-T301, SMC2, WT31 and F101.01. These CD3-specific antibodies are well known in the art and, inter alia, described in Tunncliffe (1989), Int. Immunol.

1, 546-550. In a more preferred embodiment, said V_H and V_L regions of said CD3 specific domain are derived from OKT-3 (as defined and described above). Even more preferred (and as illustrated in the appended examples) said V_H and V_L regions are or are derived from an antibody/antibody derivative with specificity for the CD3 molecule described by Traunecker (1991), EMBO J. 10, 3655-3659. In accordance with this invention, said V_H and V_L regions are derived from antibodies/antibody derivatives and the like which are capable of specifically recognizing the human CD3- ϵ chain in the context of other TCR subunits, e.g. in mouse cells transgenic for human CD3- ϵ chain. These transgenic mouse cells express human CD3- ϵ chain in a native or near native conformation. Accordingly, the V_H and V_L regions derived from an CD3- ϵ chain specific antibody is most preferred in accordance with this invention and said (parental) antibodies should be capable of specifically binding epitopes reflecting the native or near native structure or a conformational epitope of human CD3 presented in context of the TCR complex. Such antibodies have been classified by Tunnacliffe (1989) as "group II" antibodies. Further classifications in Tunnacliffe (1989) comprise the definition of "group I" and "group III" antibodies directed against CD3. "Group I" antibodies, like UCHT1, recognize CD3- ϵ chain expressed as recombinant protein and as part of the TCR on the cell surface. Therefore, "group I" antibodies are highly specific for CD3- ϵ chain. In contrast, the herein preferred "group II antibodies" recognize CD3- ϵ chain only in the native TCR complex in association with other TCR subunits. Without being bound by theory, it is speculated in context of this invention that in "group II" antibodies, the TCR context is required for recognition of CD3- ϵ chain. CD3- γ chain and δ chain, being associated with ϵ chain, are also involved in binding of "group II antibodies". All three subunits express immunoreceptor-tyrosine activation motifs (ITAMs) which can be tyrosine phosphorylated by protein tyrosine-based kinases. For this reason group II antibodies induce T cell signaling via CD3- ϵ chain, γ chain and δ chain, leading to a stronger signal compared to group I antibodies selectively inducing T cell signaling via CD3- ϵ chain. Yet, since for therapeutic applications induction of a strong T cell signaling is desired, the V_H (anti-CD3) / V_L (anti-CD3)- regions (or parts thereof) to be employed in the bispecific single

chain constructs comprised in the inventive pharmaceutical composition, are preferably derived from antibodies directed against human CD3 and classified in "group II" by Tunnacliffe (1989), loc.cit.

- 5 In one embodiment the present invention relates to a pharmaceutical composition wherein said bispecific single chain antibody construct comprises an amino acid sequence selected from the group of:
 - (a) an amino acid sequence as shown in any of SEQ ID NOs: 2, 4, 8, 10, 12, 14, 16, 18, 20, 30, 36, 39, 42, 44, 46, 48, 50, 52, 54, 56, 58 and 60;
 - 10 (b) an amino acid sequence encoded by a nucleic acid sequence as shown in any of SEQ ID NOs: 1, 3, 7, 9, 11, 13, 15, 17, 19, 29, 35, 38, 41, 43, 45, 47, 49, 51, 53, 55, 57 and 59;
 - (c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b) under stringent hybridization conditions;
 - 15 (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).
- 20 The present invention also provides for a pharmaceutical composition comprising a nucleic acid sequence encoding a bispecific single chain antibody construct as defined above.

Said nucleic acid molecule may be a natural nucleic acid molecule as well as a recombinant nucleic acid molecule. The nucleic acid molecule may, therefore, be

- 25 of natural origin, synthetic or semi-synthetic. It may comprise DNA, RNA as well as PNA (peptide nucleic acid) and it may be a hybrid thereof.

Thus, the present invention relates to a pharmaceutical composition comprising a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- 30 (a) a nucleotide sequence encoding the mature form of a protein comprising the amino acid sequence of the bispecific single chain antibody constructs

defined herein, preferably as given in SEQ ID Nos: 2, 4, 8, 10, 12, 14, 16, 18, 20, 30, 36, 39, 42, 44, 46, 48, 50, 52, 54, 56, 58 and 60;

- (b) a nucleotide sequence comprising or consisting of the DNA sequence as given in SEQ ID Nos: 1, 3, 7, 9, 11, 13, 15, 17, 19, 29, 35, 38, 41, 43, 45, 47, 49, 51, 53, 55, 57 and 59;
- (c) a nucleotide sequence hybridizing with the complementary strand of a nucleotide sequence as defined in (b) under stringent hybridization conditions;
- (d) a nucleotide sequence encoding a protein derived from the protein encoded by a nucleotide sequence of (a) or (b) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence encoded by the nucleotide sequence of (a) or (b);
- (e) a nucleotide sequence encoding a protein having an amino acid sequence at least 60 % identical to the amino acid sequence encoded by the nucleotide sequence of (a) or (b);
- (f) a nucleotide sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (a) to (e);

The term "mature form of the protein" defines in context with the present invention a protein translated from its corresponding mRNA and optional subsequently modified.

The term "hybridizing" has been defined in the context of the present invention herein above.

It is evident to the person skilled in the art that regulatory sequences may be added to the nucleic acid molecule comprised in the pharmaceutical composition of the invention. For example, promoters, transcriptional enhancers and/or sequences which allow for induced expression of the polynucleotide of the invention may be employed. A suitable inducible system is for example tetracycline-regulated gene expression as described, e.g., by Gossen and Bujard (Proc. Natl. Acad. Sci. USA 89 (1992), 5547-5551) and Gossen et al. (Trends Biotech. 12 (1994), 58-62), or a dexamethasone-inducible gene expression system as described, e.g. by Crook (1989) EMBO J. 8, 513-519 .

Furthermore, it is envisaged for further purposes that nucleic acid molecules may contain, for example, thioester bonds and/or nucleotide analogues. Said modifications may be useful for the stabilization of the nucleic acid molecule
5 against endo- and/or exonucleases in the cell. Said nucleic acid molecules may be transcribed by an appropriate vector containing a chimeric gene which allows for the transcription of said nucleic acid molecule in the cell. In this respect, it is also to be understood that such polynucleotide can be used for "gene targeting" or "gene therapeutic" approaches. In another embodiment said nucleic acid molecules are
10 labeled. Methods for the detection of nucleic acids are well known in the art, e.g., Southern and Northern blotting, PCR or primer extension. This embodiment may be useful for screening methods for verifying successful introduction of the nucleic acid molecules described above during gene therapy approaches.

Said nucleic acid molecule(s) may be a recombinantly produced chimeric nucleic
15 acid molecule comprising any of the aforementioned nucleic acid molecules either alone or in combination. Preferably, the nucleic acid molecule is part of a vector.

The present invention therefore also relates to a pharmaceutical composition comprising a vector comprising the nucleic acid molecule described in the present
20 invention.

Many suitable vectors are known to those skilled in molecular biology, the choice of which would depend on the function desired and include plasmids, cosmids, viruses, bacteriophages and other vectors used conventionally in genetic engineering. Methods which are well known to those skilled in the art can be used
25 to construct various plasmids and vectors; see, for example, the techniques described in Sambrook et al. (loc cit.) and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1989), (1994). Alternatively, the polynucleotides and vectors of the invention can be reconstituted into liposomes for delivery to target cells. As discussed in further details below, a
30 cloning vector was used to isolate individual sequences of DNA. Relevant sequences can be transferred into expression vectors where expression of a

particular polypeptide is required. Typical cloning vectors include pBluescript SK, pGEM, pUC9, pBR322 and pGBT9. Typical expression vectors include pTRE, pCAL-n-EK, pESP-1, pOP13CAT.

Preferably said vector comprises a nucleic acid sequence which is a regulatory
5 sequence operably linked to said nucleic acid sequence encoding a bispecific single chain antibody constructs defined herein.

Such regulatory sequences (control elements) are known to the artisan and may include a promoter, a splice cassette, translation initiation codon, translation and insertion site for introducing an insert into the vector. Preferably, said nucleic acid
10 molecule is operatively linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells.

It is envisaged that said vector is an expression vector comprising the nucleic acid molecule encoding a bispecific single chain antibody constructs defined herein.

The term "regulatory sequence" refers to DNA sequences which are necessary to
15 effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoters, ribosomal binding sites, and terminators. In eukaryotes generally control sequences include promoters, terminators and, in some instances, enhancers, transactivators or transcription
20 factors. The term "control sequence" is intended to include, at a minimum, all components the presence of which are necessary for expression, and may also include additional advantageous components.

The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner.
25 A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is preferably used.

Thus, the recited vector is preferably an expression vector. An "expression vector"
30 is a construct that can be used to transform a selected host and provides for expression of a coding sequence in the selected host. Expression vectors can for

instance be cloning vectors, binary vectors or integrating vectors. Expression comprises transcription of the nucleic acid molecule preferably into a translatable mRNA. Regulatory elements ensuring expression in prokaryotes and/or eukaryotic cells are well known to those skilled in the art. In the case of eukaryotic cells they

5 comprise normally promoters ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the P_L , *lac*, *trp* or *tac* promoter in *E. coli*, and examples of regulatory elements permitting expression in eukaryotic host cells are the *AOX1* or

10 *GAL1* promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells.

Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as

15 the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Furthermore, depending on the expression system used leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the recited nucleic acid sequence and are well known in the art; see also, e.g., the appended examples.

20 The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an N-terminal identification peptide imparting desired

25 characteristics, e.g., stabilization or simplified purification of expressed recombinant product; see supra. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pEF-Neo, pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogene), pEF-DHFR and pEF-ADA, (Raum et al. Cancer Immunol Immunother (2001) 50(3), 141-

30 150) or pSPORT1 (GIBCO BRL).

Preferably, the expression control sequences will be eukaryotic promoter systems

in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and as desired, the collection and purification of the polypeptide of the invention may follow; see, e.g., the appended examples.

An alternative expression system which could be used to express a cell cycle interacting protein is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The coding sequence of a recited nucleic acid molecule may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of said coding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which the protein of the invention is expressed (Smith, J. Virol. 46 (1983), 584; Engelhard, Proc. Nat. Acad. Sci. USA 91 (1994), 3224-3227).

Additional regulatory elements may include transcriptional as well as translational enhancers. Advantageously, the above-described vectors of the invention comprises a selectable and/or scorable marker.

Selectable marker genes useful for the selection of transformed cells and, e.g., plant tissue and plants are well known to those skilled in the art and comprise, for example, antimetabolite resistance as the basis of selection for dhfr, which confers resistance to methotrexate (Reiss, Plant Physiol. (Life Sci. Adv.) 13 (1994), 143-149); npt, which confers resistance to the aminoglycosides neomycin, kanamycin and paromycin (Herrera-Estrella, EMBO J. 2 (1983), 987-995) and hyg, which confers resistance to hygromycin (Marsh, Gene 32 (1984), 481-485). Additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 (1988), 8047); mannose-6-phosphate isomerase which allows cells to utilize mannose (WO 94/20627) and

- ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed.) or deaminase from *Aspergillus terreus* which confers resistance to
- 5 Blasticidin S (Tamura, Biosci. Biotechnol. Biochem. 59 (1995), 2336-2338).
- Useful scorable markers are also known to those skilled in the art and are commercially available. Advantageously, said marker is a gene encoding luciferase (Giacomin, Pl. Sci. 116 (1996), 59-72; Scikantha, J. Bact. 178 (1996), 121), green fluorescent protein (Gerdes, FEBS Lett. 389 (1996), 44-47) or β -glucuronidase
- 10 (Jefferson, EMBO J. 6 (1987), 3901-3907). This embodiment is particularly useful for simple and rapid screening of cells, tissues and organisms containing a recited vector.
- As described above, the recited nucleic acid molecule can be used alone or as part of a vector to express the encoded polypeptide in cells, for, e.g., gene therapy. The
- 15 nucleic acid molecules or vectors containing the DNA sequence(s) encoding any one of the above described bispecific single chain antibody constructs is introduced into the cells which in turn produce the polypeptide of interest. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable
- 20 vectors, methods or gene-delivery systems for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813; Verma, Nature 389 (1994), 239; Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086;
- 25 Onodera, Blood 91 (1998), 30-36; Verma, Gene Ther. 5 (1998), 692-699; Nabel, Ann. N.Y. Acad. Sci. 811 (1997), 289-292; Verzeletti, Hum. Gene Ther. 9 (1998), 2243-51; Wang, Nature Medicine 2 (1996), 714-716; WO 94/29469; WO 97/00957, US 5,580,859; US 5,589,466; or Schaper, Current Opinion in Biotechnology 7 (1996), 635-640. The recited nucleic acid molecules and vectors may be designed
- 30 for direct introduction or for introduction via liposomes, or viral vectors (e.g., adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell,

embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell. An example for an embryonic stem cell can be, inter alia, a stem cell as described in, Nagy, Proc. Natl. Acad. Sci. USA 90 (1993), 8424-8428.

In accordance with the above, the present invention relates to methods to derive
5 vectors, particularly plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering that comprise a nucleic acid molecule encoding the polypeptide sequence of a bispecific single chain antibody constructs defined herein. Preferably, said vector is an expression vector and/or a gene transfer or targeting vector. Expression vectors derived from viruses such as
10 retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the recited polynucleotides or vector into targeted cell populations. Methods which are well known to those skilled in the art can be used to construct recombinant vectors; see, for example, the techniques described in Sambrook et al. (loc cit.), Ausubel (1989, loc cit.) or other standard
15 text books. Alternatively, the recited nucleic acid molecules and vectors can be reconstituted into liposomes for delivery to target cells. The vectors containing the nucleic acid molecules of the invention can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas
20 calcium phosphate treatment or electroporation may be used for other cellular hosts; see Sambrook, supra.

The recited vector may be the pEF-DHFR, pEF-ADA or pEF-neo.

The vectors pEF-DHFR and pEF-ADA have been described in the art, e.g. in Mack et al. (PNAS (1995) 92, 7021-7025) and Raum et al. (Cancer Immunol Immunother
25 (2001) 50(3), 141-150).

It is further envisaged that the pharmaceutical composition of the invention comprises a host transformed or transfected with a vector defined herein above.

Said host may be produced by introducing said at least one of the above described
30 vector or at least one of the above described nucleic acid molecules into the host. The presence of said at least one vector or at least one nucleic acid molecule in

the host may mediate the expression of a gene encoding the above described bespecific single chain antibody constructs.

The described nucleic acid molecule or vector which is introduced in the host may either integrate into the genome of the host or it may be maintained
5 extrachromosomally.

The host can be any prokaryote or eukaryotic cell.

The term "prokaryote" is meant to include all bacteria which can be transformed or transfected with a DNA or RNA molecules for the expression of a protein of the invention. Prokaryotic hosts may include gram negative as well as gram positive
10 bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. The term "eukaryotic" is meant to include yeast, higher plant, insect and preferably mammalian cells. Depending upon the host employed in a recombinant production procedure, the protein encoded by the polynucleotide of the present invention may be glycosylated or may be non-glycosylated. Especially
15 preferred is the use of a plasmid or a virus containing the coding sequence of the polypeptide of the invention and genetically fused thereto an N-terminal FLAG-tag and/or C-terminal His-tag. Preferably, the length of said FLAG-tag is about 4 to 8 amino acids, most preferably 8 amino acids. An above described polynucleotide can be used to transform or transfect the host using any of the techniques
20 commonly known to those of ordinary skill in the art. Furthermore, methods for preparing fused, operably linked genes and expressing them in, e.g., mammalian cells and bacteria are well-known in the art (Sambrook, loc cit.).

Preferably, said the host is a bacteria, an insect, fungal, plant or animal cell.

It is particularly envisaged that the recited host may be a mammalian cell, more
25 preferably a human cell or human cell line.

Particularly preferred host cells comprise CHO cells, COS cells, myeloma cell lines like SP2/0 or NS/0.

The pharmaceutical composition of the invention may also comprise a
30 proteinaceous compound capable of providing an activation signal for immune effector cells useful for cell proliferation or cell stimulation.

The proteinaceous compound is not understood as an additional domain of the above defined bispecific single chain antibody construct, but at least one additional component of the pharmaceutical composition of the invention.

In the light of the present invention, said "proteinaceous compounds" providing an
5 activation signal for immune effector cells" may be, e.g. a further activation signal for T cells (e.g. a further costimulatory molecule: molecules of the B7-family, Ox40 L, 4.1 BBL), or a further cytokine: interleukin (e.g. IL-2), or an NKG-2D engaging compound. Preferred formats of proteinaceous compounds comprise additional bispecific antibodies and fragments or derivatives thereof, e.g. bispecific scFv.
10 Proteinaceous compounds can comprise, but are not limited to scFv fragments specific for the T cell receptor or superantigens. Superantigens directly bind to certain subfamilies of T cell receptor variable regions in an MHC-independent manner thus mediating the primary T cell activation signal. The proteinaceous compound may also provide an activation signal for immune effector cell which is a
15 non-T cell. Examples for immune effector cells which are non-T cells comprise, inter alia, NK cells.

An additional technical feature of the pharmaceutical composition of the invention is that said pharmaceutical composition is thermostable at $\geq 37^{\circ}\text{C}$.

20

An alternative embodiment of the invention relates to a process for the production of a pharmaceutical composition of the invention, said process comprising culturing a host defined herein above under conditions allowing the expression of the construct and recovering the produced bispecific single chain antibody construct
25 from the culture.

The transformed hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The polypeptide of the invention can then be isolated from the growth medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the, e.g., microbially
30 expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological

separations such as those involving the use of monoclonal or polyclonal antibodies directed, e.g., against a tag of the polypeptide of the invention or as described in the appended examples.

The conditions for the culturing of a host which allow the expression are known in the art and discussed herein above. The same holds true for procedures for the purification/recovery of said constructs.

A further alternative embodiment of the invention relates to the use of a bispecific single chain antibody construct as defined above, a nucleic acid sequence as defined above, a vector as defined above, a host as defined above and/or produced by a process as defined above for the preparation of a pharmaceutical composition for the prevention, treatment or amelioration of a tumorous disease.

In particular, the pharmaceutical composition of the present invention may be particularly useful in preventing, ameliorating and/or treating cancer.

Preferably said tumorous disease is epithelial cancer or a minimal residual cancer.

It is envisaged by the present invention that the above defined bispecific single chain antibody construct, nucleic acid molecules and vectors are administered either alone or in any combination using standard vectors and/or gene delivery systems, and optionally together with a pharmaceutically acceptable carrier or excipient. Subsequent to administration, said nucleic acid molecules or vectors may be stably integrated into the genome of the subject.

On the other hand, viral vectors may be used which are specific for certain cells or tissues and persist in said cells. Suitable pharmaceutical carriers and excipients are well known in the art. The pharmaceutical compositions prepared according to the invention can be used for the prevention or treatment or delaying the above identified diseases.

Furthermore, it is possible to use a pharmaceutical composition of the invention which comprises described nucleic acid molecules or vectors in gene therapy. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses,

adenoviruses, and adeno-associated viruses, among others. Delivery of nucleic acids to a specific site in the body for gene therapy may also be accomplished using a biolistic delivery system, such as that described by Williams (Proc. Natl. Acad. Sci. USA 88 (1991), 2726-2729). Further methods for the delivery of nucleic acids comprise particle-mediated gene transfer as, e.g., described in Verma, Gene Ther. 15 (1998), 692-699.

Furthermore the invention relates to a method for the prevention, treatment or amelioration of a tumorous disease comprising the step of administering to a subject in the need thereof an effective amount a bispecific single chain antibody construct as defined above, a nucleic acid sequence as defined above, a vector as defined as defined above, a host as defined above and/or produced in by a process as defined above.

Preferably said subject is a human.

- 15 The method for the prevention, treatment or amelioration of the invention may comprise the co-administration of an above defined proteinaceous compound capable of an activation signal for immune effector cells to the subject. The co-administration may be a simultaneous co-administration or a non-simultaneous co-administration.
- 20 It is particularly preferred for the use and the method of the invention that said tumorous disease is epithelial cancer, preferably adenocarcinomas, or a minimal residual cancer, preferably early solid tumor, advanced solid tumor or metastatic solid tumor.
- 25 Finally, the present invention relates to a kit comprising a bispecific single chain antibody construct as defined above, a nucleic acid sequence as defined above, a vector as defined above and/or a host as defined above. It is also envisaged that the kit of this invention comprises a pharmaceutical composition as described herein above, either alone or in combination with further medicaments to be
- 30 administered to a patient in need of medical treatment or intervention.

The Figures show:

Figure 1:

DNA and amino acid sequence of the anti-CD3-anti-EpCAM constructs **A)** anti-CD3 VHVL stL x 3-1 VHVL (SEQ ID NO.:11,12), **B)** anti-CD3 VHVL aL x 4-7 VHVL (SEQ ID NO.:1,2), **C)** anti-CD3 VHVL aL Ser x 4-7 VHVL (SEQ ID NO.:7, 8), **D)** anti-CD3 VHVL stL x 4-7 VHVL (SEQ ID NO.:13,14), **E)** anti-CD3 VHVL stL x 4-7 VLVH (SEQ ID NO.:15,16), **F)** anti-CD3 VHVL aL x 5-10 VHVL (SEQ ID NO.:3,4), **G)** anti-CD3 VHVL aL Ser x 5-10 VHVL (SEQ ID NO.:9, 10), **H)** anti-CD3 VHVL stL x 5-10 VHVL (SEQ ID NO.:17,18), **I)** anti-CD3 VHVL stL x 5-10 VLVH (SEQ ID NO.:19,20), **J)** anti-CD3 VHVL aL x 3-1 VHVL (SEQ ID NO.:45, 46), **K)** anti-CD3 VHVL aL Ser x 3-1 VHVL (SEQ ID NO.:47,48), **L)** anti-CD3 VHVL aL x 3-5 VHVL (SEQ ID NO.:49,50), **M)** anti-CD3 VHVL aL Ser x 3-5 VHVL (SEQ ID NO.:51,52), **N)** anti-CD3 VHVL stL x 3-5 VHVL (SEQ ID NO.:53,54), **O)** anti-CD3 VHVL aL x 4-1 VHVL (SEQ ID NO.:55,56), **P)** anti-CD3 VHVL aL Ser x 4-1 VHVL (SEQ ID NO.:57,58) and **Q)** anti-CD3 VHVL stL x 4-1 VHVL (SEQ ID NO.:59,60).

Figure 2:

FACS analysis of the constructs **A)** anti-CD3 VHVL stL x 5-10 VHVL (SEQ ID NO.:18), **B)** anti-CD3 VHVL stL x 4-7 VHVL (SEQ ID NO.:14), **C)** anti-CD3 VHVL aL x 5-10 VHVL (SEQ ID NO.:4), **D)** anti-CD3 VHVL aL x 4-7 VHVL (SEQ ID NO.:2), **E)** anti-CD3 VHVL aL Ser x 5-10 VHVL (SEQ ID NO.:10), **F)** anti-CD3 VHVL aL Ser x 4-7 VHVL (SEQ ID NO.:8), **G)** anti-CD3 VHVL stL x 3-1 VHVL (SEQ ID NO.:12), **H)** anti-CD3 VHVL stL x 5-10 VLVH (SEQ ID NO.:20) and **I)** anti-CD3 VHVL stL x 4-7 VLVH (SEQ ID NO.:16) in CD3 positive Jurkat and EpCAM-positive Kato III cells. A shift to the right shows binding. In Jurkat and KatoIII cells the dotted line indicates the shift of the negative control (only secondary antibody), dashed line shows the binding of an anti-EpCAM-anti-CD3 control antibody and the bold line shows the bispecific construct of interest.

Figure 3:

DNA and amino acid sequence of the anti-EpCAM-anti-CD3- constructs **A)** 4-7 VLVHx anti-CD3 VHVL (SEQ ID NO.:41,42), **B)** 3-5 VLVHx anti-CD3 VHVL (SEQ ID NO.:29,30), **C)** 3-1 VLVHx anti-CD3 VHVL (SEQ ID NO.:35,36), **D)** 4-1 VLVHx anti-CD3 VHVL (SEQ ID NO.:38,39) and **E)** 5-10 VLVHx anti-CD3 VHVL (SEQ ID NO.:43,44).

Figure 4: FACS analysis of the constructs **A)** 4-7 VLVHx anti-CD3 VHVL (SEQ ID NO.:42), **B)** 3-5 VLVHx anti-CD3 VHVL (SEQ ID NO.:30), **C)** 3-1 VLVHx anti-CD3 VHVL (SEQ ID NO.:36), **D)** 4-1 VLVHx anti-CD3 VHVL (SEQ ID NO.:39) and **E)** 5-10 VLVHx anti-CD3 VHVL (SEQ ID NO.: 44) constructs in CD3 positive Jurkat and EpCAM-positive Kato III cells. A shift to the right shows binding.

Figure 5:

A representative elution pattern of an EpCAM bispecific antibody containing protein fractions from a Zn-Chelating Fractogel® column at 280 nm. High adsorption at 280 nm from 50-450 ml retention time was due to non-bound protein in the column flow through. The arrow at the peak at 530.09 ml indicates the EpCAM bispecific construct containing protein fraction that was used for further purification.

Figure 6:

A representative protein elution pattern from a Sephadex® S200 gel filtration column at 280 nm. The protein peak at 82.66 ml containing bispecific antibodies against CD3 and EpCAM corresponds to a molecular weight of ca. 52 kD. Fractions were collected from 40-140 ml retention time.

Figure 7

A) Cation exchange chromatogram of 3-1 x anti-CD3 (SEQ ID NO.:36) shows the overall charge isoforms of the protein. Cation exchange chromatography was performed on a MiniS® (Amersham) column. After washing with 20 mM MES buffer pH 5.5, the protein was eluted with a gradient of elution buffer containing 1 M NaCl: 0-30% in 60 column volumes. The bispecific construct was eluted at 23,58 ml. Unspecific protein was eluted with 1 M NaCl starting at 50 ml.

- B) Cation exchange chromatogram of 5-10 x anti-CD3 (SEQ ID NO.:44) shows the overall charge isoforms of the protein. Cation exchange chromatography was performed as in Figure 7A. The bispecific construct was eluted at a shoulder at 35,77 ml. Unspecific protein was eluted with 1 M NaCl starting at 50 ml.

Figure 8:

- A) Representative SDS-PAGE analysis of EpCAM bispecific single chain antibody protein fractions. Lane **M**: Molecular weight marker Lane **1**: cell culture supernatant; lane **2**: IMAC flow through; lane **3**: IMAC wash; lane **4**: IMAC eluate; lane **5**: purified antibody against EpCAM and CD3 obtained from gel filtration.
- B) Representative Western blot analysis of purified EpCAM bispecific single chain antibody protein fractions Lane **1**: cell culture supernatant; lane **2**: IMAC flow through; lane **3**: IMAC wash; lane **4**: IMAC eluate; lane **5**: purified antibody against EpCAM and CD3 obtained from gel filtration.

Figure 9:

- Cytotoxicity assay of C-terminal EpCAM binders anti-CD3x3-1 (SEQ ID NO.:46), anti-CD3 x-5-10 (SEQ ID NO.:4), and anti-CD3 x4-7 (SEQ ID NO.:2). CB15 T cell clone and CHO-EpCAM cells were used in an E:T ratio of 5:1. CHO-EpCAM cells were stained with PKH26 dye and the cells were counted after bispecific single chain antibody incubation with FACS analysis.

Figure 10:

- Cytotoxicity assay of N-terminal EpCAM binders 3-1xanti-CD3 (SEQ ID NO.:36), and 5-10xanti-CD3 (SEQ ID NO.:44) . CB15 T cell clone and CHO-EpCAM cells were used in an E:T ration of 5:1. CHO-EpCAM cells were stained with PKH26 dye and the cells were counted after bispecific single chain antibody incubation with FACS analysis.

Figure 11:

- A) Sequence alignment of the CDR3 of the VH chains of EpCAM 3-1 (SEQ ID NO.:80), EpCAM 4-1 (SEQ ID NO.: 88), EpCAM 5-10 (SEQ ID NO.: 96), EpCAM 3-5 (SEQ ID NO.: 84), EpCAM 4-7 (SEQ ID NO.:92), compared with CDR3 of the VH chain of EpCAM M79, HD70 and 3B10. The NXD motif is depicted as bold.
- B) Comparison of the cytotoxic activity of 3-1xanti-CD3 (SEQ ID NO.: 36), 5-10xanti-CD3 (SEQ ID NO.:44), anti-CD3x4-7 (SEQ ID NO.:2) and anti-CD3x5-10 (SEQ ID NO.:18) with M79Xanti-CD3 and HD70xanti-CD3 controls. PBMC cells and Kato III cells were used in a E:T ratio of 10:1. KatoIII cells were stained with propidium iodide and the cells were counted after bispecific single chain antibody incubation with FACS analysis.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of scope of the present invention.

Example 1: Cloning and expression of the EpCAM constructs

A number of constructs comprising anti-CD3 and anti-EpCAM in various structures and domain arrangements were generated. Anti-EpCAM VH and VL variable domains of the antibodies 3-1 are shown in SEQ ID NO.:79, 80, 81, 82, 3-5 in SEQ ID NO.:83, 84, 85, 86, 4-1 in SEQ ID NO.:87, 88, 89, 90, 4-7 SEQ ID NO.:91, 92, 93, 94 and 5-10 in SEQ ID NO.:95, 96, 97, 98. The constructs are summarized in Table 1.

Table 1. anti-CD3-anti-EpCAM and anti-EpCAM-anti-CD3 constructs

SEQ ID NO.: Construct No.	Construct	Domain arrangement	Distinctive feature
anti-CD3xanti-EpCAM constructs			
SEQ ID NO.: 1,2	anti-CD3x4-7	VH-VLXVH-VL	
SEQ ID NO.: 3, 4	anti-CD3x5-10	VH-VLXVH-VL	
SEQ ID NO.: 45,46	anti-CD3x3-1	VH-VLXVH-VL	
SEQ ID NO.: 49,50	anti-CD3x3-5	VH-VLXVH-VL	
SEQ ID NO.: 55,56	anti-CD3x4-1	VH-VLXVH-VL	
SEQ ID NO.: 7,8	anti-CD3x4-7Cys-Ser	VH-VLXVH-VL	Cys-Ser mutation
SEQ ID NO.: 9,10	anti-CD3x5-10Cys-Ser	VH-VLXVH-VL	Cys-Ser mutation
SEQ ID NO.: 47,48	anti-CD3x3-1	VH-VLXVH-VL	Cys-Ser mutation
SEQ ID NO.: 51,52	anti-CD3x3-5	VH-VLXVH-VL	Cys-Ser mutation
SEQ ID NO.: 57,58	anti-CD3x4-1	VH-VLXVH-VL	Cys-Ser mutation
SEQ ID NO.: 11,12	anti-CD3x3-1	VH-VLXVH-VL	(G ₄ S) ₃ -linker
SEQ ID NO.: 13,14	anti-CD3x4-7	VH-VLXVH-VL	(G ₄ S) ₃ -linker
SEQ ID NO.: 15,16	anti-CD3x4-7	VH-VLXL-VH	(G ₄ S) ₃ -linker
SEQ ID NO.: 17,18	anti-CD3x5-10	VH-VLXVH-VL	(G ₄ S) ₃ -linker
SEQ ID NO.: 19,20	anti-CD3x5-10	VH-VLXL-VH	(G ₄ S) ₃ -linker
SEQ ID NO.: 53,54	anti-CD3x3-5	VH-VLXVH-VL	(G ₄ S) ₃ -linker
SEQ ID NO.: 59,60	anti-CD3x4-1	VH-VLXVH-VL	(G ₄ S) ₃ -linker
anti-EpCAM- anti-CD3 constructs			
SEQ ID NO.: 29,30	3-5xanti-CD3	VL-VHxVH-VL	
SEQ ID NO.: 35,36	3-1xanti-CD3	VL-VHxVH-VL	
SEQ ID NO.: 38,39	4-1xanti-CD3	VL-VHxVH-VL	
SEQ ID NO.: 41,42	4-7xanti-CD3	VL-VHxVH-VL	
SEQ ID NO.: 43,44	5-10xanti-CD3	VL-VHxVH-VL	

1.1 Cloning of C-terminal EpCAM-binders

1.1.1 Preparation of anti-CD3 PCR products

a) Anti-CD3 constructs with original 18 amino acid linker (SEQ ID NOs.:1, 2, 3 and 4)

- 5 The N-terminal original anti-CD3 containing the 18 amino acid linker (SEQ ID NO.:70) was obtained by PCR using the CD19xCD3 construct (Löffler A et al., Blood 2000 95:2098-103) as template and the following primers (CD3 VH *BsrGI*: AGGTGTACACTCCGATATCAAAGTGCAGCAG (SEQ ID NO.:5), CD3 VL *BspEI*: AATCCGGATTTCAGCTCCAGCTTGG (SEQ ID NO.:6)).

0 b) Anti-CD3 constructs with original 18 amino acid linker and Cys to Ser mutation in CDRH3 (SEQ ID Nos. 7,8, 9 and 10)

- The N-terminal original anti-CD3 containing the 18 amino acid linker (Seq ID NO.:70) and the Cys to Ser mutation was obtained by PCR using a CD19xanti-CD3 (C→S mutation) construct as template and the primers CD3 VH *BsrGI* and
 5 CD3 VL *BspEI* (Seq ID Nos. 5 and 6). The CDRH3 sequence with the Cys-Ser mutation is shown in SEQ ID NO.:78.

c) Anti-CD3-anti-EpCAM constructs with (G₄S)₃ linker (Seq ID Nos. 11, 12, 13, 14, 15, 16, 17, 18, 19 and 20)

- The N-terminal anti-CD3 containing the 15 amino acid standard (G₄S)₃ linker (SEQ ID NO.:99) was obtained by PCR using the CD19xCD3 (Löffler A et al., Blood 2000
 20 95:2098-103) as template. The anti-CD3 VH region and the anti-CD3 VL region were separately amplified by the following primers (CD3 VH: CD3 VH *BsrGI*: AGGTGTACACTCCGATATCAAAGTGCAGCAG (SEQ ID NO.:5), 3'CD3 VH GS15 GGAGCCGCCGCCGAGAACCAACCACCTGAGGAGACTGTGA
 25 GAGTGGTGCCTTG (SEQ ID NO.:21); CD3 VL: 5'CD3 VL GS15 GGCGGCCGCCGCTCCGGTGGTGGTCTGACATTGAGC
 TGACCCAGTCTCC (SEQ ID NO.:22), CD3 VL *BspEI*: AATCCGGATTTCAGCTCCAGCTTGG (SEQ ID NO.:6)). Overlapping complementary sequences introduced into the PCR products were used to form the
 30 coding sequence of a 15-amino acid (G₄S)₃ (single-letter amino acid code) (SEQ ID

NO.:99) linker during the subsequent fusion PCR. This amplification step was performed with the primer pair CD3 VH *BsrGI* (SEQ ID NO.:5) and CD3 VL *BspEI* (SEQ ID NO.:6).

1.1.2 Cloning of the anti-CD3xanti EpCAM constructs in VH_{anti-CD3}-VL_{anti-CD3} x

- 5 VH_{anti-EpCAM}-VL_{anti-EpCAM} orientation (SEQ ID NO.:1,2, SEQ ID NO.:3,4, SEQ ID NO.:7,8, SEQ ID NO.:9,10, SEQ ID NO.:11,12, SEQ ID NO.:13,14 and SEQ ID NO.:17,18)

The N-terminal original anti-CD3 containing the 18 amino acid linker (SEQ ID NO.:70) or the N-terminal original anti-CD3 containing the 15 amino acid standard (G₄S)₃ linker (SEQ ID NO.:99) was cleaved with the restriction enzymes *BsrGI* and *BspEI* and subsequently cloned into the bluescript KS vector (Stratagene, La Jolla, CA), containing the amino acid sequence of an eukaryotic secretory signal (leader peptide) as a *EcoRI/BsrGI*-Fragment. After cleavage of this construct with *EcoRI* and *BspEI* the resulting DNA fragment comprising the respective anti-CD3 scFv
15 with the leader peptide was cloned into a *EcoRI/BspEI* cleaved plasmid containing the c-terminal EpCAM binders 3-1 (SEQ ID NO.:79-82), 4-7 (SEQ ID NO.:91-94), or 5-10 (SEQ ID NO.:95-98) in pEFDHFR. pEFDHFR was described in Mack et al. Proc. Natl. Acad. Sci. USA 92 (1995) 7021-7025).

1.1.3. Cloning of the anti-CD3xanti EpCAM constructs in VH_{anti-CD3}-VL_{anti-CD3} x

- 20 VL_{anti-EpCAM}-VH_{anti-EpCAM} orientation (SEQ ID Nos.: 15, 16, 19 and 20)

The C-terminal anti-EpCAM antibody 4-7 (SEQ ID NO.:91-94) in VLVH orientation containing the 15 amino acid standard linker (SEQ ID NO.:99) was obtained by PCR. The 4-7 VH region and the 4-7 VL region were separately amplified by the following primers (4-7 VL: 4-7 VL *BspEI* FOR
15 CTGAAATCCGGAGGTGGTGGATCCGAGCTCGTGATGACCCAGACTCC (SEQ ID NO.:100), 4-7 VL GS15 REV GGAGCCGCCGCCGAGACCA
CCACCTTTGATCTCAAGCTTGGTCCCC (SEQ ID NO.:101); 4-7 VH: 4-7 VH GS15
FOR GGCGGCGCGGCTCCGGTGGTGGTGGTCTGAGGTGCAGCTGCTCGAGCA
10 G (SEQ ID NO.:23), 4-7 VH *SaI* REV TTTTAAGTCGACCTAATGATGATGAT-

GATGATGTGAGGAGACGGTGACCGTGG (SEQ ID NO.:24)). Overlapping complementary sequences introduced into the PCR products were used to form the coding sequence of a 15-amino acid (G₄S)₃ (single-letter amino acid code) linker (SEQ ID NO.:99) during the subsequent fusion PCR. This amplification step was
 5 performed with the primer pair 4-7 VL *BspEI* FOR and 4-7 VH *SaI* REV (SEQ ID NO.100, SEQ ID NO.:24).

The C-terminal anti-EpCAM antibody 5-10 (SEQ ID NO.:95-98) in VLVH orientation containing the 15 amino acid standard linker (SEQ ID NO.:99) was obtained by PCR. The 5-10 VH region and the 5-10 VL region were separately amplified by the
 10 following primers (5-10 VL: 5-10 VL *BspEI* FOR
 CTGAAATCCGGAGGTGGTGGATCCGAGCTCGTGATGACACAGTCTCCAT
 (SEQ ID NO.:25), 5-10 VL GS15 REV
 GGAGCCGCCGCCGAGAACCAACCAACCTTTGATCTCAAGCTTGGTCCCA
 G (SEQ ID NO.: 26); 5-10 VH: 5-10 VH GS15 FOR
 15 GGCGGCGGCGGCTCCGGTGGTGGTTCTGAGGTGCAGCTGCTCGAGC
 (SEQ ID NO.:27), 5-10 VH *SaI* REV
 TTTTAAGTCGACCTAATGATGATGATGATGATGTGAGGAGACGGTGACCGTG
 G (SEQ ID NO.:28)). Overlapping complementary sequences introduced into the PCR products were used to form the coding sequence of a 15-amino acid (G₄S)₃
 20 linker (SEQ ID NO.:99) during the subsequent fusion PCR. This amplification step was performed with the primer pair 5-10 VL *BspEI* FOR and 5-10 VH *SaI* REV (SEQ ID NO.:25, SEQ ID NO.:28).

These PCR products (5-10 VLVH and 4-7 VLVH) were cleaved with *BspEI* and *SaI* and ligated in the *BspEI/SaI* cleaved anti-CD3 VHVL stLx5-10 VHVL (SEQ ID
 25 NO.:17,18) or anti-CD3 VHVL stL x 4-7 (SEQ ID NO.:13, 14) VHVL in pEFDHFR replacing the 5-10 VHVL DNA fragment.

1.1.4. Expression and binding of the anti-CD3-EpCAM constructs

After confirmation of the sequence coding for the bispecific single chain by sequencing the plasmid was transfected into DHFR deficient CHO cells for
 30 eukaryotic expression. Eukaryotic protein expression in DHFR deficient CHO cells

was performed as described in Kaufmann R.J. (1990) Methods Enzymol. 185, 537-566). The transfected cells were then expanded and 1 litre of supernatant produced. Expression and binding of the bispecific single chain molecules were confirmed by FACS analyses. For that purpose the EpCAM positive human gastric cancer cell line Kato III (obtained from American Type Culture Collection (ATCC) Manassas, VA 20108 USA, ATCC number: HTB-103) was used. Binding of the anti-CD3 part was demonstrated on Jurkat cells (ATCC TIB 152).

Cells were cultured according to the recommendations of the supplier and ca. 200000 cells were incubated with 10µg/ml of the construct in 50µl PBS with 2%FCS. The binding of the construct was detected with an anti-His antibody (Penta-His Antibody, BSA free, obtained from Qiagen GmbH, Hilden, FRG) at 2µg/ml in 50µl PBS with 2%FCS. As a second step reagent a R-Phycoerythrin-conjugated affinity purified F(ab')₂ fragment, goat anti-mouse IgG, Fc-gamma fragment specific antibody, diluted 1:100 in 50µl PBS with 2% FCS (obtained from Dianova, Hamburg, FRG) was used. The samples were measured on a FACSScan (BD biosciences, Heidelberg, FRG). All the constructs comprising anti-CD3 and anti-EpCAM showed stronger binding affinity to CD3 and to EpCAM than the prior art anti-EpCAM (M79)xanti-CD3 bispecific antibody (Figure 2).

1.2 N-terminal EpCAM binders

1.2.1 Cloning of the anti-EpCAMxanti-CD3 constructs

Cloning of the construct 3-5xanti-CD3 (SEQ ID NOs.29, 30):

The C-terminal 3-5 in VH-VL orientation was obtained by PCR for the construction of 3-5 xanti-CD3 (SEQ ID NO.:29) molecule. Fragments I and II were amplified by PCR using primer pairs me 81 (SEQ ID NO.:31) /me 90 (SEQ ID NO.:34) and me 83 (SEQ ID NO.:32) /me 84 (SEQ ID NO.:33), respectively. Hot Start PCR was done using the Expand High Fidelity System of Roche Diagnostics. 20 cycles (94°C/30 sec; 60°C/1min;72°C/1min) were used for amplification followed by one cycle of 3 min at 72°C.

PCR fragments I and II were subjected to electrophoresis on a 1.5% agarose gel.

Fragments were mixed (1 ng of each) and used as a template for the next PCR

reaction performed with primer pair me 81 (SEQ ID NO.:31) and me 84 (SEQ ID NO.:33) for amplification of fragment III. PCR was performed as described above. Fragment III was purified on an agarose gel and digested with BssHII and BspEI (Biolabs), purified and subsequently cloned into the corresponding sites of the

5 pEF-dHFR-signal peptide (77/78)- anti-CD3 cloning vector, which facilitates cloning of anti-target variable regions in front of the anti-CD3 region. The vector has a unique BssHII site just after the signal peptide followed by BspEI site, linker (G₄S) and anti-CD3 region. The cloned region was verified by restriction digests and by DNA-sequencing.

10 Sequences of the Primers used:

Me 81: 5'- GGA TGC GCG CGA GCT CGT GAT GAC CCA GAC TCCA CTC TCC -3' (SEQ ID NO.:31)

Me 83: 5'- GGT TCT GGC GGC GGC GGC TCC GGT GGT GGT TCT GAG GTG CAG CTG CTC GA CAG TCT G -3' (SEQ ID NO.:32)

15 Me 84: 5'- GTG CTC CGG AGG AGA CGG TGA CCG TGG TCC CTT GGC CCC AG -3' (SEQ ID NO.:33)

Me 90: 5'- CCG GAG CCG CCG CCG CCA GAA CCA CCA CCT TTG ATC TCA AGC TTG GTC CC -3' (SEQ ID NO.:34)

Cloning of the construct 3-1xanti-CD3 (SEQ ID NO.:35, 36):

20 The C-terminal 3-1 in VH-VL orientation was obtained by PCR for the construction of 3-1 xanti-CD3 (SEQ ID NO.:35) molecule. Fragments I and II were amplified by PCR using primer pairs me 91a (SEQ ID NO.:37) /me 90 (SEQ ID NO.:34) and me 83 (SEQ ID NO.:32) /me 84 (SEQ ID NO.:33) , respectively. PCR was performed as above.

25 Agarose gel fragments comprising PCR fragments I and II were used as a template for the next PCR reaction performed with primer pair me 91a (SEQ ID NO.:37) and me 84 (SEQ ID NO.:33) for amplification of fragment III. PCR was performed as described above except, annealing was performed at 68°C instead of at 60°C. Fragment III was purified on an agarose gel and digested with BsrGI and BspEI

(Biolabs), purified and subsequently cloned into the corresponding sites of the pEF-dHFR-M79 X anti-CD3 cloning vector. The cloned region was verified by restriction digests and by DNA-sequencing.

Me 91a: 5'- GGA TTG TAC A CTCC GA GCT CGT CAT GAC CCA GTC TCC ATC

5 TTA TCT TGC TGC -3' (SEQ ID NO.:37)

Cloning of the construct 4-1xanti-CD3 (SEQ ID NO.:38, 39) :

The C-terminal 4-1 in VH-VL orientation was obtained by PCR for the construction of 4-1 xanti-CD3 (SEQ ID NO.:38, 39) molecule. Fragments I and II were amplified by PCR using primer pairs me 92a (SEQ ID NO.:40) /me 90 (SEQ ID NO.:34) and
10 me 83 (SEQ ID NO.:32) /me 84 (SEQ ID NO.:33), respectively. PCR was performed as above in annealing temperature of 60°C.

Agarose gel fragments comprising PCR fragments I and II were used as a template for the next PCR reaction performed with primer pair me 92a (SEQ ID NO.:40) and me 84 (SEQ ID NO.:33) for amplification of fragment III. PCR was performed as
15 described above except, annealing was performed at 68°C instead of at 60°C. Fragment III was purified on an agarose gel and digested with BsrGI and BspEI (Biolabs), purified and subsequently cloned into the corresponding sites of the pEF-dHFR-M79 X anti-CD3 cloning vector. The cloned region was verified by restriction digests and by DNA-sequencing.

20 Me 92a: 5'- GGA TTG TAC A CTCC GA GCT CGT GAT GAC ACA GTCTCC ATC CTC C -3' (SEQ ID NO.:40)

Cloning of the construct 4-7xanti-CD3 (SEQ ID NO.:41,42)

The C-terminal 4-7 in VH-VL orientation was obtained by PCR for the construction of 4-7 xanti-CD3 (SEQ ID NO.:41, 42) molecule. Fragments I and II were amplified
25 by PCR using primer pairs me 81 (SEQ ID NO.:31) /me 90 (SEQ ID NO.:34) and me 83 (SEQ ID NO.:32) /me 84 (SEQ ID NO.:33), respectively. PCR was performed as above with an annealing temperature of 60°C.

Agarose gel fragments comprising PCR fragments I and II were used as a template for the next PCR reaction performed with primer pair me 81 (SEQ ID NO.:31) and

me 84 (SEQ ID NO.:33) for amplification of fragment III. PCR was performed as described above. Fragment III was purified on an agarose gel and digested with BssHII and BspEI (Biolabs), purified and subsequently cloned into the corresponding sites of the pEF-dhfr-signal peptide (77/78)- anti-CD3 cloning vector. The cloned region was verified by restriction digests and by DNA-sequencing.

Cloning of the construct 5-10xanti-CD3 (SEQ ID NO.:43, 44) :

The C-terminal 5-10 in VH-VL orientation was obtained by PCR for the construction of 5-10xanti-CD3 (SEQ ID NO.:43, 44) molecule. Fragments I and II were amplified by PCR using primer pairs me 92a (SEQ ID NO.:40) /me 90 (SEQ ID NO.:34) and me 83 (SEQ ID NO.:32) /me 84 (SEQ ID NO.:33), respectively. PCR was performed as above with an annealing temperature of 60°C.

Agarose gel fragments comprising PCR fragments I and II were used as a template for PCR with primer pair me 92a (SEQ ID NO.:40) and me 84 (SEQ ID NO.:33) for amplification of fragment III. PCR was performed as described above except, annealing was performed at 68°C instead of at 60°C. Fragment III was purified on an agarose gel and digested with BsrGI and BspEI (Biolabs), purified and subsequently cloned into the corresponding sites of the pEF-dhfr-M79 X anti-CD3 cloning vector. The cloned region was verified by restriction digests and by DNA-sequencing.

1.2.2 Expression of anti-EpCAMxanti-CD3 bispecific molecules

CHO-cells lacking DHFR gene were maintained in alpha MEM medium (Life Technologies, cat.no: 32561) supplemented with 10% fetal Calf Serum (Life Technologies, heat inactivated at 65°C for 30 minutes) and with HT (Hypoxanthin and Thymidine; Life Technologies, cat. no: 41065-012). The cells were transfected with pEF-dHFR-3-1xanti-CD3 (SEQ ID NO.:35, 36), pEF-dHFR-3-5xanti-CD3 (SEQ ID NO.:29, 30), pEF-dHFR-4-1xanti-CD3 (SEQ ID NO.:38, 39), pEF-dHFR-4-7xanti-CD3 (SEQ ID NO.:41, 42) and pEF-dHFR-5-10xanti-CD3 (SEQ ID NO.:43, 44) using Lipofectamine 2000 kit (Invitrogen; cat. no:11668-019) according to the instructions provided by the Manufacturer. After 48 hrs, the cells were subjected to

selection by transferring the transfected cells into the selection medium (alpha MEM medium (cat. no:32561) containing heat inactivated 10% dialysed fetal Calf Serum (Life Technologies). After 2-3 weeks of selection, the cells were grown for 8 to 9 days (in 500 ml of selection medium) for production of bispecific molecules in 2
5 litre Tissue culture Roller Bottles (Falcon (cat. no: 353068;Becton Dickinson Labware). The tissue culture medium was centrifuged at 4°C for 10 minutes at 300g (1300rpm) to remove the cells and cell debris. The supernatant containing the secreted bispecific molecules was stored at -20°C until further analysis.

1.2.3 Binding assays of bispecific anti EpCAMxanti CD3 variants

10 In order to analyze the binding strength of the bispecific anti-EpCAMxanti-CD3 single chain constructs of the invention, the following binding assay was carried out.

250000 Jurkat cells (for CD3 binding) and Kato cells (for EpCAM binding) were independently incubated with crude supernatants (50µl) containing bispecific
15 construct for 45 min. at 4°C. Thereafter, the cells were washed twice in FACS buffer (phosphate-buffered saline containing 1% fetal calf serum (FCS) and 0.05% sodium azide) and incubated with mouse anti-His antibody (Dianova, DIA910) for 60 min. at 4°C. Washing steps were performed as above.

The cells were finally incubated either with goat anti-mouse-FITC-conjugated
20 antibody (BD 550003) or with anti-mouse-PE conjugated antibody (IgG) (Sigma, P8547). After washing steps, 10,000 events were analysed using FACS Calibur (B&D). All the EpCAM constructs showed strong binding (Figure 4).

Example 2. Purification of the EpCAM constructs

25 In order to purify the bispecific single chain constructs comprising anti-EpCAM and anti-CD3 the CHO-EpCAM cells were grown in roller bottles with HiClone® CHO modified DMEM medium (HiQ) for 7 days before harvest. The cells were removed by centrifugation and the supernatant containing the expressed protein was stored at -20°C.

Åkta FPLC System® (Pharmacia) and Unicorn Software® were used for chromatography. All chemicals were of research grade and purchased from Sigma (Deisenhofen) or Merck (Darmstadt).

IMAC was performed, using a Fractogel® column (Pharmacia) that was loaded
5 with ZnCl_2 according to the manufacturers protocol. The column was equilibrated
with buffer A2 (20 mM NaPP pH 7.5, 0.4 M NaCl) and the cell culture supernatant
(500ml) was applied to the column (10 ml) with a flow rate of 3 ml/min. The column
was washed with buffer A2 to remove unbound sample. Bound protein was eluted
using a 2-step gradient of buffer B2 (20 mM NaPP pH 7.5, 0.4 M NaCl, 0.5 M
10 Imidazol). In Step 1 20% buffer B2 in 10 column volumes was used and in Step2
100% buffer B2 in 10 column volumes was used. Eluted protein fractions from the
100% step were pooled for further purification.

Gel filtration chromatography was performed on a Sephadex S200 HiPrep®
column (Pharmacia) equilibrated with PBS (Gibco). Eluted protein samples (flow
15 rate 1ml/min) were subjected to SDS-Page and Western Blot for detection.

The column was previously calibrated for molecular weight determination
(molecular weight marker kit, Sigma MW GF-200).

Protein concentrations were determined using protein assay dye (MicroBCA,
Pierce) and IgG (Biorad) as standard protein. The yields of the protein are shown
20 in Table 2. All constructs were producible.

Table 2. Yields of the single-chain bispecific constructs comprising anti-EpCAM and anti-CD3

Construct	Yield [µg purified protein per liter culture]
4-1 x anti-CD3 (SEQ ID NO.:39)	172.5
3-5 x anti-CD3 (SEQ ID NO.:30)	265
4-7 x anti-CD3 (SEQ ID NO.:42)	37
anti-CD3 x 4-7. (SEQ ID NO.:2)	112.5
anti-CD3 Cys-Ser x 4-7 (SEQ ID NO.:8)	140
3-1 x anti-CD3 (SEQ ID NO.:36)	265
5-10 x anti-CD3 (SEQ ID NO.:44)	400
anti-CD3 x 5-10 (SEQ ID NO.:4)	195

- A further high resolution cation exchange chromatography was performed on a
- 5 MiniS® column (Amersham), equilibrated with 20mM MES buffer pH 5.5. The sample was diluted 1:3 with the same buffer before loading to the column. Bound protein was eluted with a gradient of equilibration buffer containing 1M NaCl: 0-30% in 60 column volumes. Remaining protein was eluted in 3 column volumes of 1M NaCl (**Figure 7**).
- 10 The EpCAM bispecific single chain construct proteins were isolated in a two-step purification process including immobilized metal affinity chromatography (IMAC) (**Figure 5**) and gel filtration (**Figure 6**). The main product had a molecular weight of 52 kDa under native conditions as determined by gel filtration in PBS.
- Purified bispecific protein was analyzed in SDS PAGE under reducing conditions
- 15 performed with precast 4-12% Bis Tris gels (Invitrogen). Sample preparation and application were according to the manufacturers protocol. The molecular weight was determined with MultiMark® protein standard (Invitrogen). The gel was stained with colloidal Coomassie (Invitrogen protocol). The purity of the isolated protein was shown to be >95% (**Figure 8A**). Western Blot was performed with an Optitran
- 20 BA-S83® membrane and the Invitrogen Blot Module according to the manufacturers protocol. The antibodies used were Penta His (Qiagen) and Goat-anti-Mouse-Ig labeled with alkaline phosphatase (AP) (Sigma), the chromogenic

substrate solution was BCIP/NBT liquid (Sigma). The EpCAM bispecific protein could be specifically detected by Western Blot (**Figure 8B**). The main signal corresponds to the main band in the SDS PAGE at 52 kD corresponding to the purified bispecific molecule.

5

Example 3. Cytotoxicity assays of the constructs comprising anti-CD3 and anti-EpCAM

In order to test the bioactivity of the constructs comprising anti-EpCAM and anti-CD3 a FACS based cytotoxicity test was performed.

- 10 For the cytotoxicity test, CHO cells from the American Type Cell Culture Collection (ATCC, Manassas, USA) were transfected with epithelial cell adhesion molecule (EpCAM). A cell clone derived from this transfection, referred to as CHO-EpCAM cells, was used for the experiments. CHO-EpCAM (1.5×10^7) cells were washed free of serum two times with PBS and incubated with PKH26 dye (Sigma-Aldrich
- 15 Co.) according to the manufacturers instructions. After staining cells were washed two times with RPMI/10% FCS.

- Cells were counted and mixed with CB15 effector cells. The CD4-positive T cell clone CB15 was provided by Dr. Fickenscher, University of Erlangen/Nuernberg, Germany. Cells were cultured as recommended by the suppliers. The resulting cell
- 20 suspension contained 400.000 target and 2×10^6 effector cells per ml. 50 μ l of the mixture was used per well in a 96 well round bottom plate.

- Antibodies were diluted in RPMI/10% FCS to the required concentration and 50 μ l of this solution was added to the cell suspension. A standard reaction was incubated for 16 h at 37°C / 5% CO₂. Propidium iodide was added to a final
- 25 concentration of 1 μ g/ml. After 10 min of incubation at room temperature cells were analysed by FACS. PKH26 fluorescence was used for positive identification of target cells. Cytotoxicity was measured as ratio of PI positive over all target cells. Sigmoidal dose response curves typically had R² values >0.97 as determined by Prism Software (GraphPad Software Inc., San Diego, USA) (**Figure 9 and 10**).

EC₅₀ values calculated by the analysis program were used for comparison of bioactivity. All the constructs of the invention show at least 50 times better cytotoxicity (maximum EC₅₀-value 169 pg/ml) than the prior art construct M79xanti-CD3 (8628 pg/ml).

5

Example 4. Determination of the binding affinity by BIAcore™ 2000 of the constructs comprising anti-EpCAM and anti-CD3 to EpCAM

In order to show the superior binding affinity of the constructs of the invention, the KD values of the constructs and of the prior art anti-EpCAM construct (M79)xanti-CD3 were determined.

10

Kinetic binding experiments were performed using surface plasmon resonance on the BIAcore™ 2000, Biacore AB (Uppsala, Sweden) with a flow rate of 5 µL/min and HBS-EP (0.01 M HEPES, pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% surfactant P20) as running buffer at 25 °C. The extracellular domain of the EpCAM antigen (residues 17-265) was immobilized onto flow cells 2-4 on a CM5 sensor chip. The chip surface was activated injecting 80 µL of 0.1 M sodium-hydroxysuccinimid, 0.4 M N-ethyl-N'(3—dimethylaminepropyl)-carbodiimid (NHS/EDC). The antigen was coupled by manual injection of 60 µg/mL EpCAM in 0.01 M sodium-acetate, pH 4.7. Different densities of antigen were immobilized on flow cells 2-4 adjusting the amount of manual injection times. Flow cell 1 was left empty while flow cell 2 was coated with the highest density of EpCAM (4100 RU). Flow cell 3 was coated with ¼ of the amount of antigen immobilized on flow cell 2 (974 RU) and flow cell 4 was coated with lowest density of Ep-CAM antigen (265 RU). The activated surface of the sensor chip was blocked injecting 85 µL of 1 M ethanolamine and the chip was left to equilibrate over night at a constant flow of 5 µL/min of HBS-EP.

15

20

25

Binding kinetics of the bispecific constructs were measured injecting 10 µL of protein solution at concentrations ranging from 4 µM-0.07 µM and monitoring the dissociation for 100 sec. Protein was buffered in HBS-EP. The data were fitted using BIAevaluation™ software determining the rate constant for dissociation and

30

association kinetics with a 1:1 Langmuir binding equation (1, 2). Where A is the concentration of injected analyte and B[0] is Rmax.

$$dB/dt = -(ka*[A]*[B] - kd*[AB]) \quad (1)$$

$$dAB/dt = -(ka*[A]*[B] - kd*[AB]) \quad (2)$$

- 5 Kinetic binding curves were determined in four concentrations of each bispecific construct analysed. The independent fitting of the raw data resulted in dissociation and association rate constants that were used to calculate the equilibrium dissociation constant (KD). The calculated KD values were unbiased for concentration indicating reliable data analysis. The average of the independently
10 determined dissociation constants as well as the standard deviation are summarized in table 3.

The analysed bispecific constructs bind to the Ep-CAM antigen immobilized on the chip surface within a well defined affinity range. The standard deviation for the calculated average dissociation constant is as expected.

15

Table 3: Dissociation constants for the bispecific constructs binding to EpCAM.

	KD (M)
M79 x anti-CD3 (control)	$4,0 \times 10^{-6}$
4-1 x anti-CD3 (SEQ ID NO.:39)	$2,5 \times 10^{-7}$
3-5 x anti-CD3 (SEQ ID NO.:30)	$2,3 \times 10^{-7}$

- The prior art anti-EpCAM x anti-CD3 construct M79xCD3 had a KD of $4,0 \times 10^{-6}$ M
20 while surprisingly the constructs of the invention have a KD in the range of $2,3 \times 10^{-7}$ - $2,5 \times 10^{-7}$ M. Thus, the constructs of the invention have more than 15 times stronger binding affinity than the prior art construct.

Example 5. Comparison of the cytotoxic activity of the constructs of the invention with prior art constructs

In order to compare the bioactivity of constructs having the NXD motif with conventional M79xCD3 and HD70xCD3 constructs the following cytotoxic assay was carried out.

KatIII cells (ATCC HTB-103) were used as target cells and grown in RPMI supplemented with 10% fetal calf serum at 37°C in a 5% CO₂ humidified incubator. Subconfluent cultures were treated with 0.25% trypsin, counted in a Neubauer chamber slide and checked for vitality by trypan-blue exclusion. Only cultures with >95% vitality were used for cytotoxicity assays. Target cells were stained with PKH26 fluorescent membrane dye according to the manufacturers manual (Sigma-Aldrich GmbH, Germany, PKH26-GL). Cell number was adjusted to 8×10^5 cells/ml in RPMI complete medium.

Human peripheral blood mononuclear cells (PBMCs) were used as effector cells and isolated from healthy donors using ficoll density gradient centrifugation with subsequent 100 x g centrifugation to remove thrombocytes. The pellet was resuspended in 10 vol. erythrocyte lysing buffer and incubated at room temperature for 10 min. Lysing reaction was stopped by addition PBS. PBMCs were resuspended in RPMI 1640 complete medium and cell number adjusted to 8×10^6 cells/ml.

Equal volumes of target and effector cell suspension were mixed and 50 µl of this suspension transferred to each well of a 96 well round bottom plate, 50 µl of EpCAM bispecific antibody serial dilution or RPMI complete medium as a negative control was added. Plates were incubated for 16 to 20 hrs at 37°C, 5% CO₂ in a humidified incubator. 50 µl propidium iodide was added to a final concentration of 1 µg/ml and incubated 15 min at room temperature. Samples were analysed by flow cytometry (FACSCalibur, Becton Dickinson). 2×10^4 events were counted.

Target cells were identified by their PKH26 fluorescent label and cytotoxicity within this population was subsequently determined. Viable cells were separated from dead cells by propidium iodide staining and the percentage of dead target cells was used as a measure for cytotoxicity. Mean values were plotted against the concentration of the bispecific antibody on a logarithmic scale, resulting in a dose response curve (Figure 11B). The corresponding EC₅₀ values were obtained after nonlinear fitting of data with the GraphPad Prism software.

The cytotoxic activity of constructs having the NXD motif (SEQ ID NO.: 36, 44, 2 and 18) was compared with conventional constructs M79xanti-CD3 and HD70-xanti-CD3 (Fig. 11B). A sequence alignment of the CDR3 regions of the VH chains of 3-1, 5-10, 4-7, 3-5 and 4-1 with M79, HD70 and 3B10 is shown in Figure 11A. Only 3-1, 5-10, 4-7, 3-5 and 4-1 have the NXD motif and furthermore, the lengths of the CDR3 regions differ. As can be seen from Figure 11A, 3-1, 4-1 and 5-10 have a CDR-H3 region of 10 amino acids, 3-5 and 4-7 have 14 amino acids whereas the prior art M79 has 8 amino acids, 3B10 has 6 amino acids and HD70 has 18 amino acids.

SEQ ID NO.: 36, 44, 2 and 18 showed a clearly better bioactivity compared to the conventional M79 and HD70 constructs (2250 pg/ml and less compared to 71460 and 11327 pg/ml of the prior art constructs, respectively) demonstrating the advantageous effects of the constructs of the invention.

Claims

1. A pharmaceutical composition comprising a bispecific single chain antibody construct, wherein said construct comprises or consists of at least two domains, wherein one of said domains binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region comprising the amino acid sequence NXD preferably in position 102 to 104 of SEQ ID NOs: 80, 88 and 96, or preferably in position 106 to 108 of SEQ ID NOs: 84 and 92, wherein X is an aromatic amino acid.
2. The pharmaceutical composition of claim 1, wherein X is W or Y.
3. The pharmaceutical composition of claim 1 or claim 2, wherein the CDR-H3 comprises at least 9 amino acid residues.
4. The pharmaceutical composition of any one of claims 1 to 3, wherein said binding domain specific for EpCAM has a K_D value of more than 5×10^{-9} M.
5. The pharmaceutical composition of any one of claims 1 to 4, wherein said binding domain specific for EpCAM has a K_D value in a range between 1×10^{-7} and 5×10^{-9} M and said binding domain specific for CD3 has a K_D value in a range between 1×10^{-6} and 5×10^{-9} M.
6. A pharmaceutical composition comprising a bispecific single chain antibody construct, wherein said construct comprises or consists of at least two domains, wherein one of said at least two domains specifically binds to human EpCAM antigen and a second domain binds to human CD3 antigen, wherein said binding domain specific for EpCAM comprises at least one CDR-H3 region of at least 9 amino acid residues and wherein said binding domain specific for

56

EpCAM has a K_D value of more than 5×10^{-9} M.

5

7. The pharmaceutical composition of any one of claims 1 to 6, wherein said binding domain specific for CD3 has a K_D value of more than 10^{-7} M.

10

8. The pharmaceutical composition of any one of claims 1 to 7, wherein the CDR-H3 region comprises at least 14 amino acids.

15

9. The pharmaceutical composition of any one of claims 1 to 8, wherein the V_H chain domains specific for human EpCAM antigen is selected from the group consisting of:

- (a) an amino acid sequence as shown in any one of SEQ ID NO: 80, SEQ ID NO: 84, SEQ ID NO: 88, SEQ ID NO: 92 and SEQ ID NO: 96;
- (b) an amino acid sequence encoded by a nucleic acid sequence as shown in any one of SEQ ID NO: 79, SEQ ID NO: 83, SEQ ID NO: 87, SEQ ID NO: 91 and SEQ ID NO: 95;
- (c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b) under stringent hybridization conditions; and
- (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).

25

10. The pharmaceutical composition of any one of claims 1 to 9, wherein the V_L chain domains specific for human EpCAM antigen is selected from the group consisting of:

- (a) an amino acid sequence as shown in any one of SEQ ID NO: 82, SEQ ID NO: 86, SEQ ID NO: 90, SEQ ID NO: 94 and SEQ ID NO: 98;
- (b) an amino acid sequence encoded by a nucleic acid sequence as shown in any one of SEQ ID NO: 81, SEQ ID NO: 85, SEQ ID NO: 89, SEQ ID NO: 93 and SEQ ID NO: 97;

30

2026305_1 (GPMaSeam) P58861 AU 4-Apr-11

- (c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b) under stringent hybridization conditions; and
 - (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).
 11. The pharmaceutical composition of any one of claims 1 to 10, wherein the binding domains specific for the CD3 antigen is derived from an antibody selected from the group consisting of: X35-3, VIT3, BMA030 (BW264/56), CLB-T3/3, CRIS7, YTH12.5, F111-409, CLB-T3.4.2, TR-66, WT32, SPv-T3b, 11D8, XIII-141, XIII-46, XIII-87, 12F6, T3/RW2-8C8, T3/RW2-4B6, OKT3D, M-T301, SMC2, WT31 and F101.01.
 12. The pharmaceutical composition of any one of claims 1 to 11, wherein said bispecific single chain antibody construct comprises an amino acid sequence selected from the group of:
 - (a) an amino acid sequence as shown in any one of SEQ ID NOs: 2, 4, 8, 10, 12, 14, 16, 18, 20, 30, 36, 39, 42, 44, 46, 48, 50, 52, 54, 56, 58 and 60;
 - (b) an amino acid sequence encoded by a nucleic acid sequence as shown in any one of SEQ ID NOs: 1, 3, 7, 9, 11, 13, 15, 17, 19, 29, 35, 38, 41, 43, 45, 47, 49, 51, 53, 55, 57 and 59;
 - (c) an amino acid sequence encoded by a nucleic acid sequence hybridizing with the complementary strand of a nucleic acid sequence as defined in (b) under stringent hybridization conditions; and
 - (d) an amino acid sequence encoded by a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence of any one of (b) and (c).

11 May 2011

2004242847

58

13. A pharmaceutical composition comprising a nucleic acid sequence encoding a bispecific single chain antibody construct as defined in any one of claims 1 to 12.
- 5 14. A pharmaceutical composition comprising a vector which comprises a nucleic acid sequence as defined in claim 13.
- 10 15. The pharmaceutical composition of claim 14, wherein said vector further comprises a regulatory sequence which is operably linked to said nucleic acid sequence defined in claim 13.
16. The pharmaceutical composition of claim 14 or claim 15, wherein said vector is an expression vector.
- 15 17. A pharmaceutical composition comprising a host transformed or transfected with a vector as defined in any one of claims 14 to 16.
18. The pharmaceutical composition of any one of claims 1 to 17, further comprising a proteinaceous compound capable of providing an activation signal for immune effector cells.
- 20 19. The pharmaceutical composition of any one of claims 1 to 18, wherein the pharmaceutical composition is thermostable at $\geq 37^{\circ}\text{C}$.
- 25 20. A process for the production of a pharmaceutical composition according to any one of claims 1 to 19, said process comprising culturing a host defined in claim 17 under conditions allowing the expression of the bispecific single chain antibody construct as defined in any one of claims 1 to 12 and recovering the produced bispecific single chain antibody construct from the culture.
- 30

2029250_1 (G+MarSer) P58001 AU 4-Apr-11

21. Use of a bispecific single chain antibody construct as defined in any one of claims 1 to 12, a nucleic acid sequence as defined in claim 13, a vector as defined in any of claims 14 to 16, a host as defined in claim 17 and/or produced by a process according to claim 20 for the preparation of a pharmaceutical composition for the prevention, treatment or amelioration of a tumorous disease.
22. A method for the prevention, treatment or amelioration of a tumorous disease, comprising the step of administering to a subject in need of such a prevention, treatment or amelioration a pharmaceutical composition of any one of claims 1 to 19.
23. The method of claim 22, wherein said subject is a human.
24. The use of claim 21 or the method of claim 22 or claim 23, wherein said tumorous disease is epithelial cancer or a minimal residual cancer.
25. A kit comprising a bispecific single chain antibody construct as defined in any one of claims 1 to 12, a nucleic acid sequence as defined in claim 13, a vector as defined in any one of claims 14 to 16, a host as defined in claim 17 and/or produced by a process according to claim 20.
26. A pharmaceutical composition according to any one of claims 1, 6, 13, 14, or 17, a process according to claim 20, use according to claim 21, a method according to claim 22, or a kit according to claim 25, substantially as hereinbefore described with reference to any one of the examples or figures.

Figure 1

A) anti-CD3 VHVL stL x 3-1 VHVL (SEQ ID NO: 11)

GATATCAAACGTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACAGAGGCCCTGGACAGGGTCTTGGAATGGATTGGAT
 ACAATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAGTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTAATTAATGCTTGAATGCTGGGCCAAGGCACCACTCTCACAGTCTCCTCAGGTGGTGGTGT
 CTGGCGGGCGGGCTCCGGTGGTGGTGGTTCTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGAGAAGGTCACCATGACCTGCAGAGCCAGTTCAAGTGTAACTTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCAAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTACAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTATTACTGCCAA
 CAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGAGGTGGTGGATCCGA
 GGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTGAACCTGGGGCCTCAGTGAAGATATCCTGCAAGGCTT
 CTGGATACGCCCTTCACTAACTACTGGCTAGGTGGGTAAAGCAGAGGCCCTGGACATGGACTTGAGTGGATTGGA
 GATCTTTCCCTGGAAGTGGTAATACTCACTACAATGAGAGGTTCAAGGGCAAGCCACACTGACTGCAGACAA
 ATCCTCGAGCACAGCCTTTATGCAGCTCAGTAGCCTGACATCTCAGGACTCTGCTGTCTATTCTGTGCAAGAT
 TGAGGAACCTGGGACGAGGCTATGGACTACTGGGCCAAGGGACCAAGTCCCGTCTCCTCAGGTGGTGGTGGT
 TCTGGCGGGCGGGCTCCGGTGGTGGTGGTTCTGAGCTCGTCAAGCCAGTCTCCATCTTATCTTGTGCTGCATC
 TCCTGGAGAAACCATTAATTAATGCAGGGCAAGTAAGAGCATTAGCAAAATATTAGCCTGGTATCAAGAGA
 AACCTGGGAAACATAAAGCTTCTTATCTACTCTGGATCCACTTTGCAATCTGGAATTCATCAAGGTTTCAGT
 GGCAGTGGATCTGGTACAGATTTCACTCTCACCATCAGTAGCCTGGAGCCTGAAG

Figure 1 A) continued

ATTTTGCAATGTATTACTGTCAACAGCATAATGAATATCCGTACACGTTCCGAGGGGGACCAAGCTTGAGATC
AAACATCATCACCATCATCATTAG

(SEQ ID NO: 12)

DIKLOQSGAELARPGASVKMSCKTSGYTFTRRYTMHWVKQRPQGLIEWIGYINPSRGYTNYNQKFKDKATLTIDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGGGGGGGGGSDIQLTQSPAIMSAS
PGEKVMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPRFSGSGSGTSYSLTISMEAEADAATYYCQ
QWSSNPLTFGAGTKLELKSGGGSEVQLLEQSGAELVKPGASVKISCKASGYAFTNHWLGVVKQRPCHGLEWIG
DLFPGSGNTHYNERFRGKATLTADKSSSTAFMQLSSLTSEDSAVYFCARLRNWDAMDYWGQGTTLTVSSGGGG
SGGGSGGGGSELVMTQSPSYLAASPGETITINCRASKSISKYLAWYQEKPGKTNKLLIYSGSTLQSGCIPSRFS
GSGSGTDFTLTISSELPEDFAMYYCQQHNEYPTFGGGTKLEIKHHHHH

Figure 1

B) anti-CD3 VHVL aL x 4-7 VHVL (SEQ ID NO:1)

GATATCAAACTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCTCAGTGAAGATGTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATCTAATTAACAATCAGAAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCACTCTATTACTGTGCAAGATA
 TTATGATGATCAATTACTGCCTTGACTACTCTGGGGCCAAAGGCACCACTCTCACAGTCTCCTCAGTCGAAGGTGGAA
 GTGGAGGTTCTGGTGGAAAGTGGAGGTTTCAGGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAAGTCAACCATGACCTGCAGAGCCAGTTCAAAGTGAAGTTACATGAAGTGGTACCA
 GCAGAAGTCAGGCACCTCCCCCAAAGATGGATTATGACACATCCAAAGTGGCTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGTCTGGACCTCATACTCTCTCACAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAAACAGTGGAGTAGTAACCCGCTCACGTTCCGTTGCTGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGCGAGGCTGGGCTTCAGTGAAGCTGTCTCT
 GCAAGGCTTCTGGCTACACCTTCACAAACTATGGTTTAAGCTGGGTGAAGCAGAGGCCCTGGACAGGTCTCTTGAG
 TGGATTGGAGAGGTTTATCCTAGAAATGGTAATGCTTACTACAATGAGAAAGTTCAAGGGCAAGGCCACACTGAC
 TGCAGACAAAATCTCCAGCACAGGTCCTCATGGAGCTCCGAGCCTGACCTCTGAGGACTCTGGGTCTATTCT
 GTGCAAGACCGGGATCTACGATACTAATACGACTGCTACTCTGATGCTCTGGGGCCAAAGGACCCACGGTCACC
 GTCTCCTCAGGTGGTGGTCTGGCGGGCGGGCTCCGGTGGTCTGGTCTGAGCTCGTGTGATGACCCAGAC
 TCCACTCTCCCTGCTCAGTCTTGGAGATCAAGCCTCCATCTCTGCAGATCTAGTCAGAGCCTTGTACACA
 GTAAATGGAAACACCTATTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAG

Figure 1 B) continued

CTCCTGATCTACAAAAGTTTCCAAACCGATTTCTGGGGTCCAGACAGGTTCA GTGGCAGTGGATCAGGGACAGA
 TTTTCACTCAAGATCAGCAGAGTGAGGCTGAGGATCTGGAGTTATTCTGCTCTCAAAGTACACATGTTTC
 CGTACACGTTGGAGGGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 2)

DIKLQSGAEIARPGASVKMSCKTSGYTFTRYTMHWVKQRPCQGLEWIGYINPSRGYTNYNQKFKDKATLTDDK
 SSTAYMQLSSLTSEDSAVYVCARYDDHYCLDYWGQGTTLTVSSVEGGSGSGSGGGVDDIQLTQSPAIM
 SASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYSLTSSMEAEADAATY
 YCQQWSSNPLTFGAGTKLELKS GGGSEVQLLEQSGAEIARPGASVKLSCKASGYTFTNYGLSWVKQRPQGVLE
 WIGEVYPRIGNAYNEKFKCKATLTADKSSSTASMEI RSLTSEDSAVYFCARRGSYDTNIDWYFDVWGQGTIVT
 VSSGGSGSGSGGGGSELVMTQTPLSLPVSLGDOASISCRSSQSLVHSNGNTYLVHWYLOKPGQSPKLLIYKV
 SNRFGVPPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSEQSTHVPYTFGGGTKLEIKHHHHHH

Figure 1

C) anti-CD3 VHVL aL Ser x 4-7 VHVL (SEQ ID NO: 7)

GATATCAAACTGCAGCAGTCAGGGGCTGAAC'TGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCAC'TGGTAAACACAGAGGCCCTGGACAGGGTCTGGAATGGATGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTACAATCAGAAGTTCAGGACAAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCATACATGCAACTAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCA'TACTCCCTTGACTACTGGGGCCCAAGGCCACTCTCACAGTCTCCTCAGTCGAAGGTGGAA
 GTGGAGGTTC'TGGTGAAGTGGAGGTTCAGGTGGAGTCCAGCGACATTCAGTGAACCCAGTCTCCAGCAATCATG
 TCTGCA'TCTCCAGGGGAGAAGTCAACATGACCTGCAGAGCCAGTTCAGTGTAAAGTTACATGAAC'TGGTACCA
 GCAGAACTCAGGCACCTCCCCAAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGGTCTGGGACCTCATACTCTCTCACAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAACAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGCGAGGCCCTGGGGCTTCAGTGAAGCTGTCCCT
 GCAAGGCTTCTGGCTACACCTTCACAAACTATGGTTTAAAGCTGGGTGAAGCAGAGGCCCTGGACAGGTCTCTTGAG
 TGGATTGGAGAGGTTTA'CCTAGAA'TGGTAATGCTTACTACAATGAGAAGTTCAGGGCAAGGCCACACTGAC
 TGCAGACAAATCCTCCAGCACAGCGTCCATGGAGCTCCGCAGCCTGACCTCTGAGGACTCTGCGGTCTATTCT
 GTGCAAGACGGGGATCCTACGATACTAACTACGACTGGTACTTCGATGTCTGGGGCCAGGGGACCAACGGTCACC
 GTCTCCTCAGGTGGTGGTCTTGCGGGCGGGGCTCCGGTGGTGGTCTGAGTCTGAGTCTCGTGTGATGACCCAGAC
 TCCACTCTCCCTGCCTGT'CAGTCTTGAGATCAAGCCTCCATCTCTTGACAGATCTAGTCAGAGCCTTGTACACA
 GTAATGGAACACACCTATTATACAT'TGGTACCTGCAGAGCCAGGCCAGTCT'CCAAAGCTCCTGATCTACAAAGTT
 TCCAACCGATTTC'TGGGGTCCCAGACAGGTTCA'GTGGCAGTGGATCAGGGACAG

Figure 1 C) continued

ATTTCACACTCAAGATCAGCAGAGTGGAGGCTGAGGATCTGGAGTTTATTCTGCTCTCAAAGTACACATGTT
CCGTACACGTTTCGGAGGGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 8)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQLEWIGYINPSRGYNNYNQKFKDKATLTIDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYSLDYWGQGTTLTVSSVEGGSGGSGGVDDIQLTQSPAIM
SASPGEKVMTTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPRFSGSGTSYSLTSSMEAEADAATY
YCOQWSSNPLTFGAGTKLELKSQGGSEVQLLEQSGAELARPGASVKLSCKASGYFTFTNYGLSWVKQRPQGVLE
WIGEVYPRIGNAYYNEKFKGKATLTADKSSSTASMEIRSLTSEDSAVYFCARRGSYDNTYDWFYDVGQGTVT
VSSGGGSGGGGGGGSEIVMTQTPELSPVSLGDQASISCRSSQSLVHSNGNTYLHWYLRKPGQSPKLLIYKV
SNRFSGVPRDRFSGSGGTDFTLKISRVEAEDLGVYFCSQSTHVPYTFGGGTKEIKHHHHH

Figure 1

D) anti-CD3 VHVL stL x 4-7 VHVL (SEQ ID NO: 13)

GATATCAAACCTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCTCAGTGAAGATGTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACAGAGGCCCTGGACAGGCTCTGGAATGGATTTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTAATTACTGTGCAAGATA
 TTATGATGATCATTAATGCTTACTGCTTACTGCTTGGGGCCCAAGCCACTCTCACAGTCTCCTCAGGTGGTGGTT
 CTGGGGGGGGGCTCCGGTGGTGGTGGTTCTGACATTCAGCTGACCCAGTCTCCAGCAATCATGCTGTCATCT
 CCAAGGGAGAAGGTCAACCATGACCTGCAGAGCCAGTTCAAGTGAAGTTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCCCAAGATGGATTTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTCAAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTATTACTGCCAA
 CAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGGTGGATCCGA
 GGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGCGAGGCCCTGGGGCTTCAGTGAAGCTGTCTGCAAGGCTT
 CTGGCTACACCTTCACAACTATGGTTAAAGCTGGGTGAAGCAGAGGCCCTGGACAGGTCCCTTGAGTGGATTGGA
 GAGGTTTATCCTAGAATTGGTAATGCTTACTACAATGAGAAGTTCAAGGCAAGGCCACACTGACTGCAGACAA
 ATCCTCCAGCACAGGCTCCATGGAGCTCCGAGCCTGACCTCTGAGGACTCTGCGGTCTAATTCTGTGCAAGAC
 GGGATCCTACGATACTAATACGACTGGTACTTCGATGTCTGGGGCCAAAGGGACCAAGTCCCGTCTCCTCA
 GGTGGTGGTGGTTCTGGCGGGCGGCTCCGGTGGTGGTGGTTCTGAGCTCGTGATGCCAGACTCCACTCTC
 CTTGCCCTGTGAGTCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCCTTGTACACAGTAATGGAA
 ACACCTATTATACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACAAAGTTTCCAAACCGA
 TTTTCTGGGGTCCCAGACAGGTTTCAGTGGCAGTGGATCAGGGACAGATTTACAC

Figure 1 D) continued

TCAAAGATCAGCAGAGTGAGGCTGAGGATCTGGGAGTTTATTCTGCTCTCAAAGTACACATGTTCCGTACACG
 TTCGGAGGGGGACCAAGCTTGAGATCAAACATCATCACCATCATCATTAG

(SEQ ID NO: 14)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGGLWIGYINPSRGYTNYNQKFKDKATLTIDK
 SSTAXMQLSLSLTSEDSAVYYCARYDDHYCLDYWGQGITTLTVSSGGGGGGGGSDIQLTQSPAIMSAS
 PGEKVMTCRASSSVYMNWYQOKSGTSPKRWIYDTSKVASGVPYRFSGSGGTSYSLTSSMEAEADAATYYCQ
 QWSSNPLTFGAGTKLELKSGGGSEVQLLEQSGAELARPGASVKLSCKASGYTFTNYGLSWVKQRPQGVLWIG
 EVYPRIGNAYYNEKFKGKATLTADKSSSTASMEILRLSLTSEDSAVYFCARRGSYDTNYDWYFDVWGQGTITVSS
 GGGSGGGGGGGSEIVMTQTPLSLPVSLGDAQASISCRSSQSLVHSNGNTYLNHWYLNQKPGQSPKLLIYKVSNR
 FSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSTHVPYTFGGGTKLEIKHHHHH

Figure 1

E) anti-CD3 VHVL stL x 4-7 VLVH (SEQ ID NO: 15)

GATATCAAACCTGCAGCAGTCAGGGGGCTGAACCTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACACAGAGGCCCTGCACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATACAAATCAGAAGTTCAAGGACAAGGCCACATTTGACTACAGACAAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTTACTTGCCTTGACTACTGGGGCCCAAGGCCACTCTCACAGTCTCCCTCAGGTGGTGGTGT
 CTGGCGGGCGGGCTCCGGTGGTGGTGGTCTTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGAGAAAGTCAACATGACCTGCAGAGCCAGTTCAGTGTAAAGTTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCCCAAAAGATGGATTTATGACACATCCAAAGTGGCTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTCAAAATCAGCAGCATGGAGGTGAAGATGCTGCCACTTATTACTGCCAA
 CAGTGGAGTAGTAACCGCTCACGTTCCGTTGGTGGACCAAGCTGGAGCTGAAATCCGGAGGTGGTGGATCCGA
 GCTCGTGATGACCCAGACTCCACTCTCCCTGCCCTGTAGTCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTA
 GTCAGAGCCTTGTACACAGTAATGGAACACCTATTATTTACATTTGGTACCTGCAGAAAGCCAGGCCAGTCTCCAAAG
 CTCCTGATCTACAAAAGTTTCCAAACCGATTTCTTGGGGTCCCAGACAGGTTCAGTGGCAGTGGATCAGGGACAGA
 TTTACACTCAAGATCAGCAGAGTGGAGCTGAGGATCTGGGAGTTTATTCTGCTCTCAAAGTACACATGTTC
 CGTACACGTTCCGAGGGGGACCAAGCTTCAGATCAAAGTGGTGGTGTCTTGGCGGGCGGGCTCCGGTGGT
 GGTGGTCTGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGCGAGGCCCTGGGGCTTCAGTGAAGCTGTC
 CTGCAAGGCTTCTGGCTACACCTTCACAAACTATGGTTTAAAGTGGTGAAGCAGAGGCCCTGGACAGGTCTCTTG
 AGTGGATTGGAGAGGTTTATCCTAGAATTGGTAATGCTTACTACAATGAGAAGTTCAAGGGCAAGGCCACACTG
 ACTGCAGACAAATCCTCCAGCACAGCGTCCATGGAGCTCCGACGCTGACCTCTG

Figure 1 E) continued

AGGACTCTGCGGTCATATTCTGTGCAAGACGGGATCCTACGATACTAACTACGACTGGTACTTCGATGCTGTGG
GGCCAAGGACCACCGGTACCGTCTCCTCACATCATCACCATCATCATTAG

(SEQ ID NO: 16)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGGLWIGYINPSRGYTNYNQKFKDKATLTIDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGSGGGSGGGSDIQLTQSPAIMSAS
PGEKVTMTCRASSVSVMNWYQOKSGTSPKRWIYDTSKVASGVPRFSGSGTSYSLTSSMEAEADAATYYCQ
QWSSNPLTFGAGTKLELKSGGGSELVMTQTPLSLPVSLGDOASISCRSSQSLVHSNGNTYLLHWYLQKPGQSPK
LLTYKVSNRFGVPDRFSGSGGTDFTLKISRVEAEDLGVIYFCSTHVPYTFGGGTKLEIKGGGGSGGGSGG
GGSEVQLLEQSGAELARPGASVKLSCKASGYTFFTNYGLSWVKQRPQGVLEWIGEVYPRIGNAYYNEKFKGKATL
TADKSSSTASMEIARSLTSEDSAVYFCARRGSDTNDYFDVWGQGTITVTVSSHHHHH

Figure 1

F) anti-CD3 VHVL aL x 5-10 VHVL (SEQ ID NO:3)

GATATCAAAACTGCAGCAGTCAGGGGCTGAACTGGCAAGACCTGGGGCCTCAGTGAAGATGTCCTGCAAAGACTTCTGGCTACACCTTTACTAGGTACAGATGCACTGGGTAAAAACAGAGGCTGGACAGGTCCTGGAATGGATTGGATACATTAATCTAGCCGTGTTATCTAATTAATCAGAACTTCAAGGACAAGGCCACATTGACTACAGACAAATCCTCCAGCACAGCTACATGCAACTGAGCAGCTGACATCTGAGGACTCTGAGTCTATTAATGTCGAAGATA TTAATGATGATCATTAATGCTCCCTTGACTACTGGGGCCAAAGCCACCACTCTCACAGTCTCCTCAGTCGAAGGTGGAA GTGGAGGTCTTGTTGGAAGTGGAGGTTTCAGGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG TCTGCATCTCCAGGGGAGAAGTCAACATGACCTGCAGAGCCAGTTCAAGTGAAGTTACATGAACCTGGTACCA GCAGAAGTCAGGCACTCCCCCAAAGATGGATTAATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATTCGCT TCAGTGGCAGTGGGTCTGGGACCTCATACTCTCTCAAAACAGACGATGGAGGCTGAAGATGCTGCCACTTAT TACTGCCAAACAGTGGAGTAGTAACCGCTCACGCTTCAGGTCCTGGGACCAAGCTGGAGCTGAATCCGGAGGTGG TGGATCCGAGGTGAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCTTGGACTTCAGTGAACATATCCT GCAAGGCTCTGTGAATAAGCCTTCACTAACATCTAGGCTAGCTTGGTAAAGCAGAGGCTGGACATGGACTTGAG TGGATTGGAGATATTTCTCCCTGGAAGTGTAAATCCACTACAATGAGAACTCAAGGGCAAAGCCACACTGAC TGCAGACAAATCTTCGAGCACAGCTTAATGATCGAGCTCAGTAGCTGACATTTGAGGACTCTGCTGTCTATTTCT TGCAGAAACTGAGGAACCTGGGACGAGCTATGGACTACTGGGGCCAAAGGACCAAGGTCACCGTCTCCTCAGGT GTTGGTGGTCTTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGAGCTCGTGATGACACAGTCTCCATCCTCCCT GACTGTGACAGCAGGAGAGAAGTCACTATGAGCTGCAAGTCCAGTCAGAGTCTGTTAAACAGTGGAAATCAAA AGAACTACTTGACCTGGTACGACAGAAACGAGGCGAGCTCTTAAACTGTTGATCTACTGGGCATCCACTAGG GAATCTGGGGTCCCTGATCGCTTCCAGGCAAGTGGATCTGGAAACAGATTTCACTC

Figure 1 F) continued

TCACCATCAGCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGT CAGAAATGATTATAGTTATCCGCTCACG
 TTCGGTGTGGACCAAGCTTGAGATCAAACATCATCACCATCATCATTAG

(SEQ ID NO: 4)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKORPGQGLEWIGYINPSRGYTNYNQKFKDKATLTIDK
 SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSVEGGSGSGSGGVDDIQLTQSPAIM
 SASPGEKVTMTCRASSVSVMNWYQQKSGTSPKRWIYDTSKVASGVPIRFSGSGTSYSLTSSMEAEADAATY
 YCQWSSNPLTTFGAGTKLELKSGGGSEVQLLEQSGAELVRPGTSVKISCKASGYAFINYLWLGWVKORPGHGLE
 WIGDIFPGSGNIHYNEKFKGKATLTADKSSSTAYMQLSSLTFEDSAVYFCARLRNWDPEPMDYWGQGTITVTVSSG
 GGGSGGGSGGGSELVMTQSPSSITVTAGEKVTMSCKSSQSLNSGNQNYLTWYQQKPGQPPKLLIYWASTR
 ESGVPDRFTGSGSGTDFITLTISVQAEDLAVYYCQNDYSYPLTTFGAGTKLEIKHHHHHH

Figure 1

G) anti-CD3 VHVL aL Ser x 5-10 VHVL (SEQ ID NO: 9)

GATATCAAACTGCAGCAGTCAGGGGCTGAACTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACTGGGTAAACACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAGTTCAGGACCAAGGCCACATGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCACTACTCCCTTGACTACTGGGGCCAAAGGCCACACTCTCACAGTCTGCTCAGTCGAAGGTGGAA
 GTGGAGGTTCTGGTGGAGTGGAGGTTCAAGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAGGTCACCATGACCTCCAGAGCCAGTCAAGTGAAGTTACATGAATGGTACCA
 GCAGAGTCAGGCACC"CCCCCAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGGTCTGGGACCTCATACTCTCACAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAACAGTGGAGTAGTAACCCGCTCACGTTGGTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCCCTGGAGCTTCACTGAAGATATCCT
 GCAAGGCTTCTGGATACGCCCTTCACTAACTACTGGCTAGGTTGGTTAAGCAGAGGCCCTGGACATGGACTTGAG
 TGGATTGGAGATAATTTCCCTGGAAGTGGTAATA"CCACTACAATGAGAAGTTCGAAGGCCAAAGCCACACTGAC
 TGCAGACAAATCTTCGAGCACAGCCTATATGCAGTCACTAGCTTGACATTTGAGGACTCTGCTGTCTATTCT
 GTGCAAGACTGAGGAACCTGGGACGAGCCTATGGACTACTGGGGCCAAAGGCCACCGTCAACCGTCTCCTCAGGT
 GGTGGTGGTTCTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGAGCTCGTGTGATGACACAGTCTCCATCCTCCCT
 GACTGTGACAGCAGGAGAGAAGTCACTATGAGCTGCAAGTCCAGTCCAGAGTCTGTTAAACAGTGGAAATCAAA
 AGAACTACTTGAACCTGGTACCAGCAGAAACCCAGGGCAGCCTCCTAAACTGTTGATC

Figure 1 G) continued

TACTGGGCATCCACTAGGGAATCTGGGTCCTGATCGCTTACAGGCAGTGGATCTGGAACAGATTTCACCTCT
 CACCATCAGCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTCAGAATGATTATAGTTATCCGCTCACGT
 TCGGTGCTGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 10)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYIMHWVKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTIDK
 SSSTAYMQLSSLTSEDSAVYYCARYDDHYSLDYWGQGTTLTVSSVEGGSGSGGSGVDDIQLTQSPAIM
 SASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGGTSYSLTSSMEAEADAATY
 YCQOWSSNPLTFGAGTKLELKS GGGSEVQLLEQSGAELVRPGTSVKISCKASGYAFNYYLWVWKQRPGHGLE
 WIGDIFPGSGNIHYNEKFKGKATLTADKSSSTAYMQLSSLTFEDSAVYFCARLRNWDEPMDYWGQGTITVYSSG
 GGGSGGGGGGGSELVMTQSPSSLTVTAGEKVTMSCKSSQSLNSGNQKNYLTWYQQKPGQPPKLLIYWASTR
 ESGVPRFTGSGSGTDFTLTITSSVQAEDLAVYYCQNDYSYPLTFGAGTKLEIKHHHHH

Figure 1

H) anti-CD3 VHV1-stL x 5-10 VHV1 (SEQ ID NO: 17)

GATATCAAACTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCTCAGTGAAGATGTCCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACAGATGCACCTGGGTAAAACAGAGGCTGGACAGGCTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAGTTCAAGGACAAGGCCACATTTACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTTACTGCCTTGACTACTGGGGCCAGGCCACCTCTCACAGTCTCTCAGGTGGTGGTGGTT
 CTGGCGGGCGGGCTCCGGTGGTGGTGGTTCTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGGAGAAGGTCACCATGACCTGCAGAGCCAGTTCAAGTTAAGTTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCCAAAAGATGGATTTATGACACATCCAAAGTGGCTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGCTCTGGGACCTCATACTCTCTCACAAATCAGCAGCATGGAGCTGAAGATGCTGCCACTTATTACTGCCAA
 CAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGGTGGATCCGA
 GGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCTGGGACTTCAGTGAAGATATCCCTGCAAGGCTT
 CTGGATACGCCTTCACTAACTACTGGCTAGGTTGGGTAAAGCAGAGGCTGGACATGGACTTGAGTGGATTGGA
 GATATTTCCCTGGAAGTGGTAATATCCACTACAATGAGAAGTTCAAGGGCAAGCCACACTGACTGCAGACAA
 ATCTTCGAGCACAGCCTATATGCAGCTCAGTAGCCTGACATTTGAGGACTCTGCTGTCTATTCTGTGCAAGAC
 TGAGGAACTGGGACGAGCCTATGGACTACTGGGGCCAAAGGACCAAGCTCACCGTCTCTCAGGTGGTGGTGGT
 TCTGGCGGGCGGGCTCCGGTGGTGGTGGTTCTGAGCTCGTGTGATGACACAGTCTCCATCTCCCTGACTGTGAC
 AGCAGAGAGAAGGTCATATGAGCTGCAAGTCCAGTCCAGTCTGTAAACAGTGGAAATCAAAAGAACTACT
 TGACCTGGTACCAGCAGAAACCAAGGCAGCCTCTAAACTGTTGATCTACTGGGCATCCACTAGGGAATCTGGG
 GTCCCTGATCGCTTCACAGGCAGTGGATCTCCAACACATTTCACTCTCACCATCA

Figure 1 H) continued

GCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTGAGAATGATTATAGTTATCCGCTCACGTTCCGGTGCT
GGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 18)

DIKLQQSGAELARPGASVFMSCKTSGYTFTTRYTMHWVKQRPQGQGLEWIGYINPSRGYTNYNQFKDKATLTTDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGGGGGGGSDIQLTQSPAIMSAS
PGEKVTMTCRASSSVSYMNYQQKSGTSPKRWIYDTSKVASGVPIRFSGSGSTSYSLTSSMEAEADAATYYCQ
QWSSNPLTTFGAGTKLELKSGGGGSEVQLLEQSGAELVRPGTSVKISCKASGYAFTNYWLVKQRPQGHGLEWIG
DIFPGSGNIHYNEKFKGKATLTADKSSSTAYMQLSSLTFEDSAVYFCARLRNWDPEMDYWGQGTTVTVSSGGGG
SGGGGGGGGSELVMTQSPSSLTVTAGKVTMSCKSSQSLNSGNQKNYLTWYQQKPGQPPKLLIYWASTRESG
VPDRFTGSGSGTDFLTITISSVQAEDLAVYYCQNDYSYPLTFGAGTKLEIKHHHHH

Figure 1

D) anti-CD3 VHL stL x 5-10 VL VH (SEQ ID NO: 19)

GATATCAAACCTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAAACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATAC'TAATTACAAATCAGAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTTACTGTGCAAGATA
 TTATGATGATCATTAATGCTTGCCTTGACTACTTGGGGCCAAAGGCACCACTCTCACAGTCTCCTCAGGTGGTGGTGT
 CTGGCGGGCGGCTCCGGTGGTGGTGTCTTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGGAGAAGGTCACCATGACCTGCAGAGCCAGTTCAAGTGTAAATTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCCAAAAGATGGATTTATGACACATCCAAAAGTGGCTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTCACAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTATTTACTGCCAA
 CAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGGTGGATCCGA
 GCTCGTGTGATGACACAGTCTCCATCCTCCCTGACTGTGACAGCAGGAGAGAAGGTCACTATGAGCTGCAAGTCCA
 GTCAGAGTCTGTAAACAGTGGAAATCAAAAGAACTACTTGACCTGGTACCCAGTCCAGCAGAAAACAGGGCAGCCTCCT
 AAACCTTTGATCTACTGGGCATCCACTAGGGAATCTGGGTCCCTGATCGCTTCACAGGCAGTGGATCTGGAAC
 AGATTTCACTCTCACCATCAGCAGTGTGCAGGCTGAAGACCCTGGCAGTTTATTTACTGT'CAGAAATGATTAAGTT
 ATCCGCTCACGTTCCGTTCTGGGACCAAGCTTGACATCAAAGTGGTGGTGGTCTGGCGCGGGGCTCCGGT
 GGTGGTGGTCTGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCCTGGGACTTCAGTGAAGAT
 ATCCTGCAAGGCTTTCTGGATACGCCTTCACTAACTACTGGCTAGGTGGTAAAGCAGAGGCCCTGGACATGGAC
 TTGAGTGGATGGAGATATTTCCCTGGAAAGTGGTAATATCCACTACAATGAGAAGTTCAAGGGCAAGCCACA
 CTGACTGCAGACAAATCTTCGAGCACAGCCTATATGCAGCTCAGTAGCCTGACAT

Figure 1 D) continued

TTGAGGACTCTGCTGTCTATTCTGTGCAAGACTGAGGAACCTGGGACGAGCCTATGGACTACTGGGGCCAAGGG
 ACCACGGTCACCGTCTCCTCACATCATCACCATCATCATTAG

(SEQ ID NO: 20)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTIDK
 SSSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGGGGGGGGGSDIQLTQSPAIMSAS
 PGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPIRFSGSGTSYSLTISSEAEADAATYYCQ
 QWSSNPLTFGAGTKLELKSGGGGSELVMTQSPSSLTVTAGEKVTMSCSSQSLINSGNQKNYLTWYQQKPGQPP
 KLLIYWASTRESGVPRFTGSGSGTDFTLTISVQAEDLAVYYCQNDYSYPLTFAGATKLEIKGGGGSGGGSSG
 GGGSEVQLLEQSGAELVRPGTSVKISCKASGYAFITNYWLGWVKQRPQGHGLEWIGDIFPGSGNIHYNEKFKGKAT
 LTADKSSSTAYMQLSSLTFEDSAVYFCARLRNWDEPMDYWGQGTITVTVSSHHHHHH

Figure 1

D) anti-CD3 VHVL aL x 3-1 VHVL (SEQ ID NO: 45)

GATATCAAACTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGTGGTAAACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTACAATCAGAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTTACTGCCCTTGACTACTGGGGCCCAAGGCCACCACTCTCACAGTCTCCTCAGTCGAAGGTGGAA
 GTGGAGGTTCTGGTGAAGTGGAGGTTCAAGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAGGTCACCATGACCTGCAGAGGCCAGTCAAGTGTAAAGTTACATGAACCTGGTACCA
 GCAGAAGTCAGGCACCTCCCCAAAAGATGGATTTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGGTCTGGGACCTCATACTCTCTCACAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAACAGTGGAGTAGTAACCCGCTCACGTTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTGAACCTGGGGCCCTCAGTGAAGATATCCT
 GCAAGGCTTCTGGATACGCCCTTCACTAACTACTGGCTAGGTTGGGTAAGCAGAGGCCCTGGACATGGACTTGAG
 TGGATTGGAGATCTTTCCCTGGAAGTGGTAAATACTCACTACAATGAGAGGTTCAGGGGCAAGCCACACTGAC
 TGCAGACAAATCCTCGAGCACAGCCTTTATGGAGCTCAGTAGCCTGACATCTGAGGACTCTGCTGTCTATTTCT
 GTGCAAGATTGAGGAACCTGGGACGAGGCTATGGACTACTGGGGCCCAAGGACCCACGGTCACTCTCCCTCAGGT
 GGTGGTGGTTCTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGAGCTCGTCATGACCCAGTCTCCATCTTATCT
 TGCTGCATCTCCTGGAGAAACCATTAATTAATTCAGGGCAAGTAAGAGCATTAGCAAAATATTTAGCCTGGT
 ATCAAGAGAAACCTGGGAAAACATAAAGCTTCTTATCTACTCTGGATCCACTTGGCAATCTGGAATTCATCA
 AGGTTCAAGTGGCAGTGGATCTGGTACAGATTTCACTCTCACCATCAGTAGCCTGG

Figure 1 J) continued

AGCCTGAAGATTTTGCAATGTATTACTGTCAACAGCATAATGAATATCCGTACACGTTTCGGAGGGGGGACCAAG
CTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 46)

DIKLQQSGAELARPCASVKMSCKTSGYTFTRYTMHWVKQRPQGLEWIGYINPSRGYTNYNQKFKDKATLTIDK
SSSTAYMQLSSLTSEDSAVYICARYYDDHYCLDYWGQGTTLTVSSVEGGSGGSGGVDIQLTQSPAIM
SASPGKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVYRFSGSGGTSYSLTSSMEAEADAATY
YCQWSSNPLTFGAGTKLELKSGGGGSEVQLLEQSGAELVKPGASVKISCKASGYAFNINYLWLVKQRPFGHGLE
WIGDLFPGSGNTHYNERFRGKATLTADKSSSTAFMQLSSLTSEDSAVYFCARLRNWDAMDYWGQGTTVTVSSG
GGSGGGGGGGSELVMTQSPSYLAASPGETITINCRASKSISKYLAWYQEKPGKTNKLLIYSGSTLQSGIPS
RFSGSGSGTDFTLTISLLEPEDEFAMYCCQQHNEYPTYTFGGGTKLEIKHHHHH

Figure 1

K) anti-CD3 VHVL aL Ser x 3-1 VHVL (SEQ ID NO: 47)

GATAACAACTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCTCAGTGAAGATGTCCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACACAGAGGCCCTGGACAGGCTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAGTTCAGGACAAGGACCAAGGTCACATTTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTTACTGTGCAAGATA
 TTATGATGATCATTTACTCCCTTGACTACTGGGGCCCAAGGCCACTCTCACAGTCTCTCAGTCGAAGGTGGAA
 GTGAGGTTCTGGTGGAAAGTGGAGTTTCAGGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAGTCAACATGACCTGCAGAGCCAGTTCAAAGTGAAGTTACATGAACCTGGTACCA
 GCAGAGTCAGGCACCTCCCCCAAAAGATGGATTTATGACACATCCAAAGTGGCTCTGGAGTCCCCTTATCGCT
 TCAGTGGCAGTGGGTCTGGGACCTCATACTCTCTCAACAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAACAGTGGAGTAGTAACCCGCTCACGTTGGTGGTGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTGAACCTGGGCCCTCAGTGAAGATATCCT
 GCAAGGCTTCTGGATAAGCCTTCACTAACTACTGGCTAGGTTGGGTAAAGCAGAGGCCCTGGACATGGACTTGAG
 TGGATTGGAGATCTTTCCCTGGAAGTGGTAATACTCACTACAATGAGAGGTTCAGGGGCAAAAGCCACACTGAC
 TGCAGACAAATCCTCGAGCACAGCCTTTATGCAGCTCAGTAGCCTGACATCTGAGGACTCTGCTGTCTATTTCT
 GTGCAAGATTGAGGAACCTGGGACGAGGCTATGGACTACTGGGGCCCAAGGACCAAGGTCACCGTCTCCCTCAGGT
 GGTGGTGGTTCTGGCGGGCGCGGCTCCGGTGGTGGTGGTCTGAGCTCGTCATGACCCAGTCTCCATCTTATCT
 TGCTGCATCTCCTGGAGAAACCATTAATAATGAGGGCAAGTAAGAGCATTAGCAAAATATTAGCCTGGT
 ATCAAGACAAACCTGGGAAACCTAATAAGCTTCTTATCTACTCTGATCCACTTTG

Figure 1 K) continued

CAATCTGGAATTCATCAAGGTTCAAGTGGCAGTGGATCTGGTACAGATTTCACCTCACCATCAGTAGCCTGGA
 GCCTGAAGATTTTGCAATGTATTACTGTCAACAGCATAAATGAATATCCGTACACGTTCCGAGGGGGACCAAGC
 TTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 48)

DIKLQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTDDK
 SSSTAYMQLSSLTSEDSAVYCYARYDDHYSLDYWGQGTTLTVSSVEGGSGGSGGVDIQLTQSPAIM
 SASPGEKVTMTCRASSVSVMNWYQQKSGTSPKRWIYDTSKVASGVPRFSGSGSTSYSLTISSEAEADAATY
 YCQWSSNPLTFGAGTKLELKS GGGGSEVQLLEQSGAELVKPGASVKISCKASGYAFINYLWVGWVKQRPQGHGLE
 WIGDLFPGSGNTHYNERFRGKATLTADKSSSTAFMQLSSLTSEDSAVYFCARLRNWDEAMDYWGQGTTLTVSSG
 GGGSGGGSGGSELVMTQSPSYLAASPGETITINCRASKSISKYLAWYQEKPGKTNKLLIYSGSTLQSGIPS
 RFSGSGSGTDFTLTISLLEPEDFAMYYCQQHNEYPPYTFGGGKLEIKHHHHH

Figure 1

L) anti-CD3 VHVL aL x 3-5 VHVL (SEQ ID NO: 49)

GATATCAAACTGCAGCAGTCAGGGGCTGAACTGGCAAGACCCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTTACTAGGTACACGATGCACTGGGTAAACACAGAGGCCCTGGACAGGGTCTGGATGGATGGAT
 ACATTAAATCCCTAGCCGTGGTTATACTAATTAACAATCAGAACTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTTACTGCTTGACTACTGGGGCCAAAGGCCACCACTCTCACAGTCTCTCAGTCGAAGGTGGAA
 GTGGAGGTTCTGGTGGAGTGGAGGTTCAAGTGGAGTCCAGCACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAGGTCAACATGACCTGCAGAGCCAGTTCAAAGTGAAGTTACATGAACCTGGTACCA
 GCAGAGTCAGGCACCTCCCCCAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGTCTGGGACCTCATCTCTCACAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAAACAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCCCTGGGACTTCAGTGAAGCTGTCT
 GCAAGGCTTCTGGCTACACCTTCACAAGCTATGGTTAAAGCTGGTGAAGCAGAGAACTGGACAGGGCCTTGAG
 TGGATTGGAGAGGTTTATCCTAGAAATGGTAATGCTTACTACAATGAGAACTTCAAGGGCAAGGCCACACTGAC
 TGCAGACAAAATCCTCCAGCACAGCGTCCATGGAGTCCGAGCTTGACATCTGAGGACTCTGCGGTCYATTTCT
 GTGCAAGACAGGGGATCCTACGGTAGTAACAGACTCCGAGCTTCGATGCTCTGGGGCCAAAGGCCACCGGTCAAC
 GTCTCTCAGTGGTGGTGGTCTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGAGCTCGTATGATACCCAGAC
 TCCACTCTCCCTGCCCTGTCAGTCTTGGAGATCAAGCCCTCCATCTCTTGCAGATCTAGTCAGAGCCTTGTACACA
 GTAATGGAAACACCTATTTACATTTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACAAAGTT
 TCCAAACCGATTTCTCTGGGTCCCAAGACAGGTTTCAAGTGGCAGTGGATCAGGGACAG

Figure 1 L) continued

ATTTCACACTCAAGATCAGCAGAGTGGAGGCTGAGGATCTGGAGTTTATTCTGCTCTCAAAGTACACATGTT
CCGTACACGTTCCGAGGGGGGACCAAGCTTGAGATCAAAACATCATCCATCATCATTAG

(SEQ ID NO: 50)

DIKLQOSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQGLEWIGYINPSRGYTNYNQKFKDKATLTTDK
SSSTAYMQLSSLTSEDSAVYICARYDDHYCLDYWGQGTTLTVSSVEGGSGSGGSGGVDIQLTQSPAIM
SASPGEKVMTCRASSVSVMNWYQQKSGTSPKRWIYDTSKVASGVPIRFRFSGSGGTSYSLTSSMEAEAAATY
YCQWSSNPLTFGAGTKLELKSGGGGSEVQLLEQSGAELVRPGTSVKLSCKASGYTFTSYGLSWVKQRTGGGLE
WIGEVYPRIGNAYNEKFKGKATLTADKSSSTASMEILRSLTSEDSAVYFCARRGSGSYSNYDWYFDVWVGQGTIVT
VSSGGGSGGGSGGSELVMTQTPLSLPVSGLGDAQASISCRSSQSLVHSNGNTYIHWYIQKPGQSPKLLIYKY
SNRFSGVDPDRFSGSGGTDETLKISRVEAEDLGVYFCSTHVPYTFGGGKLEIKHHHHHH

Figure 1

M) anti-CD3 VHVLaL Ser x 3-5 VHVLaL (SEQ ID NO: 51)

GATATCAAACTGCAGCAGTCAGGGGCTGAACTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCATGGGTAAACAGAGGCCCTGGACAGGGTCTGGAAATGGATTGGAT
 ACATTAAATCCTAGCGTGGTTATATACTAATACAATCAGAAAGTTCAAGGACAAGGCCACATTCAGTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTAATCTCCCTTGACTACTGGGGCCAAAGGCACCTCTCACAGTCTCTCAGTCGAAGGTGGAA
 GTGGAGGTTCTGGTGGAAAGTGGAGGTTCAAGTGGAGTCCGACCATTCAGTGAACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAGGTCACCATGACCTGCAGAGCCAGTTCAAAGTGAAGTTACATGAATGGTACCA
 GCAGAAAGTCAGGCACCTCCCCCAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCT
 TCAGTGGCAGTGGTCTGGGACCTCATACTCTCTCAAAATCAGCAGCATGGAGGCTGAAGATGCTGGCCACTTAT
 TACTGCCAAACAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGTTAAGGCTGGGACTTCAGTGAAGCTGTCTCT
 GCAAGGCTTCTGGCTACACCTTCACAAGCTATGTTAAGCTGGTGAAGCAGAGAACTGGACAGGGGCTTGAG
 TGGATTGGAGAGGTTTATCCTAGAATTGGTAATGCTTACTACAATCAGAAAGTTCAAGGGCAAGGCCACACTGAC
 TGCAGACAAAATCCTCCAGCACAGCCTCCATGGAGCTCCGAGCTGACATCTGAGGACTCTGGGGTCTATTTCT
 GTGCAAGACAGGGGATCCTACGGTAGTAACACTAGACTGGTACTTCGATGTCTGGGGCCCAAGGACCCACGGTCAAC
 GTCTCCTCAGGTGGTGGTGGTCTGGCGGGGGGGCTCCGGTGGTGGTGGTCTGAGCTCGTGAATGACCCAGAC
 TCCACTCTCCCTGCCCTGTCAGTCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCCTTGTACACA
 GTAATGGAAAACACCTATTTACATTTGTAACCTGCAGAAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACAAAGTT
 TCCAACCGATTTCCTGGGCTCCAGACAGGTTTCAGTGGCAGTGGATCAGGGACAG

Figure 1 M) continued

ATTTCACACTCAAGATCAGCAGAGTGCAGGCTGAGGATCTGGGAGTTTATTCTGCTCTCAAAGTACACATGTT
CCGTACACAGTTCGGAGGGGGACCAAGCTTGAGATCAAACATCATCACCATCATCATTAG

(SEQ ID NO: 52)

DIKLQOSGAELARPGASVKMSCKTSGYTFTRYTMHWVKORPGQGLEWIGYINPSRGYTNYNQKFKDKATLTDDR
SSSTAYMQLSSLTSEDSAVYYCARYDDHYSLDYWGQGTTLTVSSVEGGSGSGSGGVDDIQLTQSPAIM
SASPGEKVTMTCRASSVSVMNWYQQKSGTSPKRWIYDTSKVASGVPIRFSGSGSGTSYSLTISMEAEADAATY
YCQQWSSNPLTFGAGTKLELKSGGGGSEVQLLEQSGAELVRPGTSVKLSCKASGYTFTSYGLSWVKQRTGQGLE
WIGEVYPRIGNAYYNEKFKGKATLTADKSSSTASMEELRSLTSEDSAVYFCARRGSYGSNYDWFYFDVWGQGTIVT
VSSGGGSGGGGGGSELVMTQTPLSLPVSLGDAQASISCRSSQSLVHSNGNTYLHWYLQKPGQSPKLLIYKV
SNRFGVPDRFRFGSGSGTDFTLKISRVEAEDLGVYFCSSQSTHVPYTFGGGTGLEIKHHHHHH

Figure 1

N) anti-CD3 VHVL stL x 3-5 VHVL (SEQ ID NO: 53)

GATATCAAACTGCAGCAGTCAGGGGCTGAACCTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAACTCCTAGCCGTGGTTATACATAATTACAATCAGAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCACTCTATTACTGTGCAAGATA
 TTATGATGATCATTAATGCTTGAATGCTGGGGCCCAAGGCACCACTCTCACAGTCTCCTCAGGTGGTGGTGGTT
 CTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGGAGAAGGTCAACCATGACCTGACAGCCAGTTCAAGTGAAGTTACATGAACCTGGTACCAGCAGAAGTC
 AGCACCTCCCCCAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTACAATCAGCAGCATGGAGCTGAAGATGCTGCCACTTATTAATGACCA
 CAGTGGAGTAGTAACCCGCTCAGCTCGGTGCTGGGACCAAGCTGGAGCTGAAGTGAAGCTGTCTGCAAGGCTT
 GGTCAGCTGCTCGAGCAGTCTGGAGCTGAGCTGGTAAAGGCTGGGACTTCAGTGAAGCTGTCTGCAAGGCTT
 CTGGCTACACCTTCACAAGCTATGGTTAAAGCTGGGTGAAGCAGAGAACTGGACAGGGCCTTGAGTGGATTGGA
 GAGGTTTATCCTAGAAATGGTAATGCTTACTACATGAGAAGTTCAAGGGCAAGGCCACACTGACTGCAGACAA
 ATCCTCCAGCACAGCGTCCATGGAGCTCCGAGCTGACATCTGAGGACTCTGCGGCTATTTCTGTGCAAGAC
 GGGATCCTACGGTAGTAACGACTGGTACTTCGATGTCCTGGGGCCCAAGGGACCACGGTCACCGTCTCTCTCA
 GGTGGTGGTGGTTCTGGCGGGCGGGCTCCGGTGGTGGTGGTCTGAGCTCGTGATGACCCAGACTCCACTCTC
 CTTGCCCTGTCACTCTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCCTTGTACACACAGTAATGGAA
 ACACCTAATTTACATTGGTACCTGCAGAAGCCAGGCCAGTCTCCAAAGCTCCTGATCTACAAAGTTTCCCAACCGA
 TTTTCTGGGGTCCACAGACAGGTTCAAGTGGCAGTGGATCAGGGACAGATTTCACAC

Figure 1 N) continued

TCAAGATCAGCAGAGTGGAGGCTGAGGATCTGGGAGTTATTTCTGCTCTCAAAGTACACATGTTCCGTACACG
TTCGGAGGGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATAG

(SEQ ID NO: 54)

DIKIQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTITDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGGGGGGGGGGGSDIQLTQSPAIMSAS
PGEKVTMTCRASSSVSYMNWYQOKSGTSPKRWIYDTSKVASGVPIRFSGSGSTSYSLTSSMEAEADAATYYCQ
QWSSNPLTFAGTKLELKSGGGGSEVQLLEQSGAELVRPGTSVKLSCKASGYTFTSYGLSWVKQRTGQGLEWIG
EVYPRIGNAYYNEKFKGKATLTADKSSSTASMEILRSLTSEDSAVYFCARRGSYGSNYDWFVDVWGQGTITVTVSS
GGGSGGGSGGGGSELVMTQTPLSLPVSLGDAQASISCRSSQSLVHSNGNTYLNHWYLNQKPGQSPKLLIYKVSNR
FSGVPDRFSGSGSGTDFTLKISRVEAEDLGVYFCQSQSTHVPYTFGGGTGLEIKHHHHHH

Figure 1

O) anti-CD3 VHVL aL x 4-1 VHVL (SEQ ID NO: 55)

GATATCAAACTGCAGCAGTCAGGGGCTGAACTGGCAAGACCTGGGGCCCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACTGGGTAAACAGAGGCCCTGGACAGGGTCTGGAAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATCAAAATCAGAAAGTTCAAGGACAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTAATGCTTGTGCTTACTTGGGGCCAAAGGCACCACTCTCACAGTCTCTCAGTCGAAGGTGGAA
 GTGAGGTTCTGGTGAAGTGGAGGTTCAAGTGGAGTCGACGACATTCAGCTGACCCAGTCTCCAGCAATCATG
 TCTGCATCTCCAGGGGAGAAGGTCAACATGACCTGCAAGGCCAGTTCAAAGTGAAGTTACATGAACTGGTACCA
 GCAGAACTCAGGCACCTCCCCCAAAGATGGATTATGACACATCCAAAGTGGCTTCTGGAGTCCCTTTATCGCT
 TCAGTGGCAGTGGGTCTGGACCTCATACTCTCTACAAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTAT
 TACTGCCAAACAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGG
 TGGATCCGAGGTGCAGCTGCTCGAGCAGTCTGAGCTGGTAAAGCCCTGGGACTTCAGTGAAGATATCCT
 GCAAGGCTTCTGGATACGCCCTTCACTAACTACTGGCTAGGTTGGTTAAGCAGAGGCCCTGGACATGGACTTGAA
 TGGCTTGGAGATATTTCCCTGGAAGTGGTAATGCTCACTACAATGAGAAGTTCAAGGGCAAGGCCACACTGAC
 TGCAGACAAAGTCCCTCGTACACAGCCCTATATGCAGCTCAGTAGCCCTGACATCTGAGGACTCTGGTCTATTCT
 GTGCAAGATTCGGGAATCGGACGAGGCTATGGACTACTGGGGCCAAAGGACCAAGGTCACCGTCTCCCTCAGGT
 GGTGGTGGTCTGGCGGGGGGGGCTCCGGTGGTGGTCTGAGCTCGTGATGACACAGTCTCCATCCCTCCCT
 GAGTGTGTACGAGGAGAGAAGGTCACTATGAGCTGCAAGTCCAGTCCAGTCTGTAAACAGTGGAAATCAAA
 AGAACTACTTGGCCTGGTACCAGCAGAAACAGGGCAGCCCTCCTAACTGTTGATCTACGGGGCATCCCACTAGG
 GAATCTGGGGTCCCTGATCGCTTCACAGGCAGTGCATCTGGAACAGATTTCACTC

Figure 1 O) continued

TCACCATCAGCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTCAGAATGATTATAGTTATCCGTACACG
 TTCGGAGGGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 56)

DIKLQQSGAELARPGASVKMSCKTSGYTFTRYTMHWKQRPQGQGLEWIGYINPSRGYTNYNQKFKDKATLTIDK
 SSSTAYMQLSSLTSEDSAVYICARYDDHYCLDYWGQGTTLTVSSVEGGSGSGGVDIQLTQSPAIM
 SASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVPYRFSGSGTSYSLTSSMEADAATY
 YCQOWSSNPLTFGAGTKLELKS GGGSEVQLLEQSGAELVRPGTSVKISCKASGYAFNWLGWVKQRPQGHGLE
 WVGDIFFPGSGNAHYNEKFKGKATLTADKSSYTAYMQLSSLTSEDSAVYFCARLRNWDAMDYWGQGTTVTVSSG
 GGGSGGGSGGGGSELVMTQSPSSISVSAGEKVTMSCKSSQSLNSGNQKNYLAWYQQKPGQPPKLLIYGASTR
 ESGVPDRFTTSGSGGTDFTLTISSVQAEDLAVYYCQNDYSYPYTFGGGTKLEIKHHHHHH

P) anti-CD3 VHVL aL Ser x 4-1 VHVL (SEQ ID NO: 57)

[illegible]

Figure 1 P) continued

TCACCATCAGCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTGAGAATGATTATAGTTATCCGTACACG
 TTCGAGGGGGGACCAAGCTTGAGATCAAAACATCATCACCATCATCTAG

(SEQ ID NO: 58)

DIKLQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGQLEWIGYINPSRGYTNYNQKFKDKATLTDDK
 SSSTAYMQLSLTSSEDSAVYYCARYDDHYSLDYWGQGTTLTVSSVEGGSGSGSGGVDIQLTQSPAIM
 SASPGEKVTMTCRASSSVSYMNWYQQKSGTSPKRWIYDTSKVASGVYPYRFSGSGSGTSYSLTISSEAEADAATY
 YCQWSSNPLTFGAGTKLELKS GGGSEVQLLEQSGAELVRPCTSVKISCKASGYAFITNYWLGWVKQRPQGHGLE
 WVGDIFFPGSGNAHYNEKFKGKATLTADKSSYTAYMQLSLTSSEDSAVYFCARLRNWDEAMDYWGQGTITVTVSSG
 GGGSGGGSGGGSELVMTQSPSSLSVSAGEKVTMSCKSSQSLNSGNQKNYLAWYQQKPGQPPKLLIYGASTR
 ESGVPDRFTGSGSGTDFTLTITSSVQAEDLAVYYCQNDYSYPYTFGGGGTKLEIKHHHHH

Figure 1

Q) anti-CD3 VHVL stL x 4-1 VHVL (SEQ ID NO: 59)

GATATCAAACTGCAGCAGTCAGGGGCTCAACTGGCAACACCTGGGGCTCAGTGAAGATGTCTCTGCAAGACTTC
 TGGCTACACCTTTACTAGGTACACGATGCACCTGGGTAAACACAGAGGCCCTGGACAGGGTCTGGAATGGATTGGAT
 ACATTAATCCTAGCCGTGGTTATACTAATTAACAATCAGAAAGTTCAGGACAAAGGCCACATTGACTACAGACAAA
 TCCTCCAGCACAGCCTACATGCAACTGAGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATA
 TTATGATGATCATTAATGCTTGAATGAGTGGGGCCAAAGCAACCACTCTCACAGTCTCTCAGGTGGTGGTGGTT
 CTGGCGCGGGCTCCGGTGGTGGTGGTCTGACATTCAGCTGACCCAGTCTCCAGCAATCATGTCTGCATCT
 CCAGGGAGAAAGTCAACATGACCTGCAGAGCCAGTTCAAGTGAAGTTACATGAACCTGGTACCAGCAGAAGTC
 AGGCACCTCCCCAAAGATGGATTTATGACACATCCAAAGTGGCTTCGGAGTCCCTTATCGCTTCAGTGGCA
 GTGGTCTGGGACCTCATACTCTCTCACAATCAGCAGCATGGAGGCTGAAGATGCTGCCACTTATTAAGTCCAA
 CAGTGGAGTAGTAACCCGCTCACGTTCCGTGCTGGGACCAAGCTGGAGCTGAAATCCGGAGGTGGTGGATCCGA
 GGTGCAGTCTCGAGCAGTCTGGAGCTGAGCTGGTAAGGCTGGGACTTCAGTGAAGATATCCTGCAAGGCTT
 CTGGATACGCCCTTCACTAACTACTGGCTAGGTGGTTAAGCAGAGGCCCTGGACATGGACTGAATGGGTGGGA
 GATATTTCCCTGGAGTGGTAATGCTCACTACAATGAGAAAGTTCAAGGGCAAGCCACACTGACTGCAGACAA
 GTCCCTGTACACAGCCTATATGCAGCTCAGTAGCCTGACATCTGAGGACTCTGCTGTCTATTTCTGTGCAAGAT
 TGGGAACTGGGACGAGGCTATGGACTACTGGGGCCAAAGGACCAGGTCACCGTCTCCTCAGGTGGTGGTGGT
 TCTGGCGCGCGGCTCCGGTGGTGGTGGTCTGAGCTCGTATGACACAGTCTCCATCTCCCTGAGTGTCTC
 AGCAGGAGAGAAAGTCACTATGAGCTGCAAGTCCAGTCAAGTCTGTAAACAGTGGAAATCAAAAGAACTACT
 TGGCCTGGTACCAGCAGAAACCAAGGCAGCCTCCTAAACTGTTGATCTACGGGGCATCCACTAGGGAATCTGGG
 GTCCCTGATCGCTTCACAGGCAGTGGATCTGGAAACAGATTTCACTCTCACCATCA

Figure 1 Q) continued

GCAGTGTGCAGGCTGAAGACCTGGCAGTTTATTACTGTCAGAATGATTATAGTTATCCGTACACGTTCCGAGGG
GGGACCAAGCTTGAGATCAAAACATCATCACCATCATCATTAG

(SEQ ID NO: 60)

DIKLQSGAELARPGASVKMSCKTSGYTFTRYTMHWVKQRPQGLEWIGYINPSRGYTNYNQKFKDKATLTDDK
SSSTAYMQLSSLTSEDSAVYYCARYDDHYCLDYWGQGTTLTVSSGGGGGGGGGGSDIQLTQSPALMSAS
PGEKVTMTCRASSSVSYMNWYQOKSGTSPKRWIYDTSKVASGVPYRFSGSGSGTSYSLTSSMEADAATYYCQ
QWSSNPLTFGAGTKLELKSGGGGSEVQLLEQSGAELVRPGTSVKISCKASGYAF"NNYWLGVVKQRPQGHLEWVG
DIFPGSGNAHYNEKFKGKATLTADKSSYTA"YMQQLSSLTSEDSAVYFCARLRNWDEAMDYWGQGT"TVTVSSGGGG
SGGGGGGGSELVMTQSPSSLSVSAAGEKVTMSCKSSQSLINSGNQKNYLAWYQQKPGQPPKLLIYGASTRESG
VPDRFTGSGSGTDFTLTITSSVQAEDLAVYYCQNDYSYPYTFGGGTKLEIKHHHHH


Figure 2 A

Figure 2 B





Figure 2C



Figure 2D**anti-CD3 VHVL aL x 4-7 (SEQ ID NO: 2)**


Figure 2E**anti-CD3VHVL aL Ser x 5-10 (SEQ ID NO: 10)**

Figure 2F

anti-CD3 VHVL aL Ser x 4-7 (SEQ ID NO: 8)






Figure 2G



Figure 2H**anti-CD3 VHVL stL x 5-10 VLVH (SEQ ID NO: 20)**


Figure 2I**anti-CD3 VHVL stL x 4-7 VLVH (SEQ ID NO: 16)**

Figure 3A

4-7 (vLvH) x anti-CD3 (SEQ ID NO: 42)

```

1  MGWSCIILFL VATATGVHSA RELVMTQTPL SLPVSLGDQA SISCRSSQSL
51  VHSNGNTYLH WYLQKPGQSP KLLIYKVSNR FSGVPDRFSG SSGTDFTLK
101 ISRVEAEDLG VFCSQSTHV PYTFGGGTKL EIKGGGGSGG GSGGGGSEV
151 QLLEQSGAEL ARPGASVKLS CKASGYTFN YGLSWVKQRP GQVLEWIGEV
201 YPRIGNAYYN EKFKGKATLT ADKSSSTASM ELRSLTSEDS AVYFCARRGS
251 YDTNYDWYFD VWGQGTTVTV SSGGGGSDIK LQQSGAELAR PGASVKMSCK
301 TSGYTFTRYT MHWVKQRPQG GLEWIGYINP SRGYTNYNQK FKDKATLTID
351 KSSSTAYMQL SSLTSEDSAV YYCARYYDDH YCLDYWGQGT TLTVSSVEGG
401 SGGSGGSGGS GGVDIQLTQ SPAIMASPG EKVTMTCRAS SSVSYMNWYQ
451 QKSGTSPKRW IYDTSKVASG VPYRFGSGGS GTSYSLTISS MEAEDAATYY
501 CQQWSSNPLT FGAGTKLELK HHHHHH*

```

Figure 3A (continued)

SEQ ID NO: 41

```

1  ATGGGATGGA GCTGTATCAT CCTCTTCTTG GTAGCAACAG CTACAGGTGT
51  ACACTCCGCG CGCGAGCTCG TGATGACCCA GACTCCACTC TCCCTGCCCTG
101 TCAGTCTTGG AGATCAAGCC TCCATCTCTT GCAGATCTAG TCAGAGCCCTT
151 GTACACAGTA ATGGAACAC CTATTTACAT TGGTACCTGC AGAAGCCAGG
201 CCAGTCTCCA AAGCTCCTGA TCTACAAAGT TTCCAACCGA TTTTCTGGG
251 TCCAGACAGG GTTCAGTGGC AGTGGATCAG GGACAGATTT CACACTCAAG
301 ATCAGCAGAG TGGAGGCTGA GGATCTGGGA GTTTATTCT GCTCTCAAG
351 TACACATGTT CCGTACACGT TCGGAGGGGG GACCAAGCTT GAGATCAAA
401 GTGGTGGTGG TTCTGGCGGC GGCGGCTCCG GTGGTGGTGG TTCTGAGGTG
451 CAGCTGCTCG AGCAGTCTGG AGCTGAGCTG GCGAGGCCCTG GGGCTTCAGT
501 GAAGCTGTCC TGCAAGGCTT CTGGCTACAC CTTCACAAA TATGGTTTAA
551 GCTGGGTGAA GCAGAGGCCCT GGACAGGTCC TTGAGTGGAT TGGAGAGGTT
601 TATCCTAGAA TTGGTAATGC TTAATAAAT GAGAAATTCA AGGCAAGGC
651 CACACTGACT GCAGACAAAT CCTCCAGCAC AGCTCCATG GAGCTCCGCA
701 GCCTGACCTC TGAGGACTCT GCGGTCTATT TCTGTGCAAG ACGGGGATCC
751 TACGATACTA ACTACGACTG GTACTTCGAT GTCTGGGGCC AAGGGACCA
801 GGTCAACCGTC TCCTCCGGAG GTGGTGGATC CGATATCAA CTGCAGCAGT
851 CAGGGGCTGA ACTGGCAAGA CCTGGGGCCT CAGTGAAGAT GTCCTGCAAG

```

Figure 3A (continued)

```

901 ACTTCTGGCT ACACCTTTAC TAGGTACACG ATGCACCTGGG TAAACACAGAG
951 GCCTGGACAG GGTCTGGAAT GGATTGGATA CATTAAATCCT AGCCGTGGTT
1001 ATACTAATTA CAATCAGAA G TCAAGGACA AGGCCACATT GACTACAGAC
1051 AAATCCTCCA GCACAGCCTA CATGCAACTG AGCAGCCTGA CATCTGAGGA
1101 CTCTGCAGTC TATTACTGTG CAAGATATTA TGATGATCAT TACTGCCCTTG
1151 ACTACTGGGG CCAAGGCACC ACTCTCACAG TCTCCTCAGT CGAAGGTGGA
1201 AGTGGAGGTT CTGGTGGAAG TGGAGGTTCA GGTGGAGTGG ACGACATTCA
1251 GCTGACCCAG TCTCCAGCAA TCATGTCTGC ATCTCCAGGG GAGAAGGTCA
1301 CCATGACCTG CAGAGCCAGT TCAAGTGTA GTTACATGAA CTGGTACCAG
1351 CAGAAAGTCAG GCACCTCCCC CAAAAGATGG ATTTATGACA CATCCAAAAGT
1401 GGCTTCTGGA GTCCCTTATC GCTTCAGTGG CAGTGGGTCT GGGACCTCAT
1451 ACTCTCTCAC AATCAGCAGC ATGGAGGCTG AAGATGCTGC CACTTATTAC
1501 TGCCAAACAGT GGAGTAGTAA CCCGCTCAGG TTCGGTGCTG GGACCAAGCT
1551 GGAGCTGAAA CATCATCACC ATCATCATTA G

```

Figure 3B

3-5 (VLVH) x anti-CD3 (SEQ ID NO: 30)

```

1  MGWSCIIILFL VATAAGVHSA RELVMTQTPL SLPVSLGDQA SISCRSSQSL
51  VHSNGNTYLH WYIQKPGQSP KLLIYKVSNR FSGVPDRFSG SGSGTDFTLK
101 ISRVEAEDLG VFYCSQSTHV PYTFGGGTKL EIKGGGGSGG GSGGGGGSEV
151 QLEQSGAEL VRPGTSVKLS CKASGYTFTS YGLSWVKQRT GQGLEWIGEV
201 YPRIGNAYYN EKFKGKATLT ADKSSSTASM ELRLTSEDS AVYFCARRGS
251 YGSNYDWYFD VWGQGTTVTV SSGGGGSDIK LQQSGAELAR PGASVKMSCK
301 TSGYTFTRYT MHWVKQRPQ GLEWIGYINP SRGYTNYNQK FKDKATLTID
351 KSSSTAYMQL SSLTSEDSAV YYCARYYDDH YCLDYWGQGT TLTVSSVEGG
401 SSGSGGSGGS GGVDIIQLTQ SPAIMSASPG EKVTMTCRAS SSVSYMNWYQ
451 QKSGTSPKRW IYDTSKVASG VPYRFSGSGS GTSYSLTSS MEAEADAATY
501 CQQWSSNPLT FGAGTKLELK HHHHHH*

```

Figure 3B (continued)

SEQ ID NO:29:

1 ATGGGATGGA GCTGTATCAT CCTCTTCTTG GTAGCAACAG CTACAGGTGT
 51 ACACCTCCGG CGCGAGCTCG TGATGACCCA GACTCCACTC TCCCTGCCCTG
 101 TCAGTCTTGG AGATCAAGCC TCCATCTCTT GCAGATCTAG TCAGAGCCCTT
 151 GTACACAGTA ATGGAACAC CTATTACAT TGGTACCTGC AGAAGCCAGG
 201 CCAGTCTCCA AAGTCCCTGA TCTACAAAGT TTCCAACCGA TTTTCTGGGG
 251 TCCCAGACAG GTTCAGTGGC AGTGGATCAG GGACAGATTT CACACTCAAG
 301 ATCAGCAGAG TGGAGGCTGA GGATCTGGGA GTTTATTCT GCTCTCAAAG
 351 TACACATGTT CCGTACACGT TCGGAGGGGG GACCAAGCTT GAGATCAAAG
 401 GTGGTGGTGG TTCTGGCGGC GGCGGCTCCG GTGGTGGTGG TTCTGAGGTG
 451 CAGCTGCTCG AGCAGTCTGG AGCTGAGCTG GTAAGGCCCTG GGACTTCAGT
 501 GAAGCTGTCC TGCAAGGCTT CTGGCTACAC CTTCAACAAG TATGGTTTAA
 551 GCTGGGTGAA GCAGAGAACT GGACAGGGCC TTGAGTGGAT TGGAGAGGTT
 601 TATCCTAGAA TTGGTAATGC TTAATAAAT GAGAAATTCA AGGCAAGGC
 651 CACACTGACT GCAGACAAAT CCTCCAGCAC AGCGTCCATG GAGTCCGCA
 701 GCCTGACATC TGAGGACTCT GCGGTCTATT TCTGTGCAAG ACGGGATCC
 751 TACGGTAGTA ACTACGACTG GTACTTCGAT GTCTGGGGCC AAGGACCCAC
 801 GGTCAACCGTC TCCTCCGGAG GTGGTGGATC CGATATCAAA CTGCAGCAGT
 851 CAGGGGCTGA ACTGGCAAGA CCTGGGGCCT CAGTGAAGAT GTCCTGCAAG
 901 ACTTCTGGCT ACACCTTTAC TAGGTACACG ATGCACCTGGG TAAACACAG

Figure 3B (continued)

```

951 GCCTGGACAG GGTCTGGAAT GGATTGGATA CATTAAATCCT AGCCGTGGTT
1001 ATACTAATTA CAATCAGAAG TTCAAGGACA AGGCCACATT GACTACAGAC
1051 AAATCCTCCA GCACAGCCTA CATGCAACTG AGCAGCCTGA CATCTGAGGA
1101 CTCTGCAGTC TATTAAGTGT CAAAGGACCC ACTCTCACAG TCTCCTCAGT CGAAGGTGGA
1151 ACTACTGGG CCAAGGCACC CTGGTGGAA TGGAGGTTCA GGTGGAGTCG ACGACATTCA
1201 AGTGGAGGTT CTGGTGGAA TCTCCAGCAA TCAATGTCCTG ATCTCCAGGG GAGAAGGTCA
1251 GCTGACCCAG TCTCCAGCAA TCAATGTCCTG ATCTCCAGGG GAGAAGGTCA
1301 CCATGACCTG CAGAGCCAGT TCAAGTGTA GTTACATGAA CTGGTACCAG
1351 CAGAAGTCAG GCACCTCCCC GCTTCAGTGG CAGTGGTCTT GGGACCTCAT
1401 GGCTTCTGGA GTCCCTTATC GCTTCAGTGG CAGTGGTCTT GGGACCTCAT
1451 ACTCTCTCAC AATCAGCAGC ATGGAGGCTG AAGATGCTGC CACTTATTAC
1501 TGCCCAACAGT GGAGTAGTAA CCCGCTCACG TTCGGTGTCTG GGACCAAGCT
1551 GGAGCTGAAA CATCATCACC ATCACATTA G

```

Figure 3C

3-1 (vLvH) x anti-CD3 (SEQ ID NO: 36)

```

1  MGWSCIIILFL VATAATGVHSE LVMTQSPSYL AASPGETITI NCRASKSISK
51  YLAWYQEKPG KTNKLLIYSG STLQSGIPSR FSGSGSGTDF TLTISSLEPE
101 DFAMYQCQOH NEYPYTFGGG TKLEIKGGG SGGGSGGGG SEVQLLEQSG
151 AELVKPGASV KISCKASGYA FTNYWLGWVK QRPBGHLEWI GDLEFPGSGNT
201 HYNFRFRGKA TLTADKSSST AFMQLSSLTS EDSAVYFCAR LRNWDEAMDY
251 WGQGTTVTVS SGGGSDIKL QQSGAELARP GASVKMSCKT SGYTFTRYTM
301 HWVKQRPQGQ LEWIGYINPS RGYTNYNQKF KDKATLTIDK SSSTAYMQLS
351 SLTSEDSAVY YCARYYDDHY CLDYWGQGT LTVSSVEGGS GSGGSGGSGG
401 GVDDIQLTQS PAIMSASPGE KVTMTCRASS SVSYMNWYQQ KSGTSPKRWI
451 YDTSKVASGV PYRFSGSGSG TSYSLTISSM EAEDAATYYC QQWSSNPPLTF
501 GAGTKLELKH HHHH*

```

Figure 3C (continued)

SEQ ID NO: 35

```

1  ATGGGATGGA  GCTGTATCAT  CCTCTTCTTG  GTAGCAACAG  CTACAGGTGT
51  AACTCCGAG  CTCGTATGA  CCCAGTCTCC  ATCTTATCTT  GCTGCATCTC
101 CTGGAGAAAC  CATTACTATT  AATTGCAGGG  CAAGTAAGAG  CATTAGCAAA
151 TATTAGCCT  GGTATCAAGA  GAAACCTGGG  AAAACTAATA  AGCTTCTTAT
201 CTACTCTGA  TCCACTTTGC  AATCTGGAAT  TCCATCAAGG  TTCAGTGGCA
251 GTGGATCTGG  TACAGATTTC  ACTCTACCA  TCAGTAGCCT  GGAGCCTGAA
301 GATTTTGCA  TGTATTACTG  TCAACAGCAT  AATGAATATC  CGTACACGTT
351 CGGAGGGGG  ACCAAGCTTG  AGATCAAAGG  TGGTGGTGGT  TCTGGCGGCG
401 GCGGCTCCG  TGGTGGTGGT  TCTGAGGTGC  AGCTGCTCGA  GCAGTCTGGA
451 GCTGAGCTG  TGAACCTGG  GGCCTCAGTG  AAGATATCCT  GCAAGGCTTC
501 TGGATACGC  TTCACTAACT  ACTGGCTAGG  TTGGGTAAAG  CAGAGGCTG
551 GACATGGACT  TGAGTGGATT  GGAGATCTTT  TCCCTGGAAG  TGCTAATACT
601 CACTACAATG  AGAGGTTTCA  GGGCAAAGCC  AACTGACTG  CAGACAAATC
651 CTCGAGCACA  GCCTTTATGC  AGCTCAGTAG  CCTGACATCT  GAGGACTCTG
701 CTGTCATTT  CTGTGCAAGA  TTGAGGAACT  GGGACGAGGC  TATGGACTAC
751 TGGGGCCAA  GGACCACGGT  CACCGTCTCC  TCCGGAGGTG  GTGGATCCGA
801 TATCAAAC TG  CAGCAGTCAG  GGGCTGAAC  TGGGCTCAG
851 TGAAGATGTC  CTGCAAGACT  TCTGGCTACA  CCTTTACTAG  GTACACGATG
901 CACTGGGTAA  AACAGAGGCC  TGGACAGGGT  CTGGAATGGA  TTGGATACAT

```

Figure 3C (continued)

```

951 TAATCCTAGC CGTGGTTATA CTAATTACAA TCAGAAAGTTC AAGGACAAGG
1001 CCACATTGAC TACAGACAAA TCCTCCAGCA CAGCCTACAT GCAACTGAGC
1051 AGCCTGACAT CTGAGGACTC TGCAGTCTAT TACTGTGCAA GATATTATGA
1101 TGATCAATTAC TGCCTTGACT ACTGGGGCCA AGGCACCACT CTCACAGTCT
1151 CCTCAGTCGA AGGTGGAAGT GGAGGTTCTG GTGGAAGTGG AGGTTCAGGT
1201 GGAGTCGACG ACAATTCAGT GACCCAGTCT CCAGCAATCA TGTCTGCCATC
1251 TCCAGGGGAG AAGTCAACCA TGACCTGCAG AGCCAATTCA AGTGTAAGTT
1301 ACATGAACTG GTACCAGCAG AAGTCAGGCA CCTCCCCCAA AAGATGGATT
1351 TATGACACAT CCAAAGTGGC TTCTGGAGTC CATTATCGCT TCAGTGGCAG
1401 TGGGTCTGGG ACCTCATACT CTCTCACAAT CAGCAGCATG GAGGCTGAAG
1451 ATGCTGCCAC TTATTACTGC CAACAGTGGG GTAGTAACCC GCTCACGTTT
1501 GGTGCTGGGA CCAAGCTGGA GCTGAAACAT CATCACCATC ATCATTTAG

```

Figure 3D

4-1 (vLvH) x anti-CD3 (SEQ ID NO: 39)

```

1  MGWSCIILFL VATAIGVHSE LVMTQSPSSL SVSAGEKVTM SCKSSQSLIN
51  SGNQKNYLAW YQOKPGQPPK LLTYGASTRE SGVDRFTGS GSGTDFTLTI
101 SSVQAEDLAV YYCQNDYSYP YTFGGGTKLE IKGGGGSGGG GSGGGGSEVQ
151 LLEQSGAELV RPTSVKISC KASGYAFTNY WLGWVKQRPQ HGLEWVGDI
201 PGSGNAHYNE KFKGKATLTA DKSSYTAYMQ LSLTSEDSA VYFCARLRNW
251 DEAMDYWGQG TTVTVSSGGG GSDIKLQOSG AELARPGASV KMSCKTSGYT
301 FTRYTMHWVK QRPQGLEWI GYINPSRGYT NYNQKFKDKA TLTDDKSSST
351 AYMQLSSLTS EDSAVYYCAR YYDDHYCLDY WQGTTLTVS SVEGGSGGSG
401 GSGSGGVDD IQLTQSPAIM SASPGEKVTM TCRASSSVSY MNWYQQKSGT
451 SPKRWIYDTS KVASGVPRF SGSGSGTSYS LTSSMEAEED AATYYCQQWS
501 SNPLTFGAGT KLELKHNNHH H*

```

Figure 3D (continued)

SEQ ID NO: 38:

```

1  ATGGGATGGA  GCTGTATCAT  CCTCTTCTTG  GTAGCAACAG  CTACAGGTGT
51  ACACTCCGAG  CTCGTGATGA  CACAGTCTCC  AT'CTTCCCTG  AGTGTGT'CA
101  CAGGAGAGAA  GGTCACTATG  AGCTGCAAGT  CCAGTCAGAG  TCT'CTTAAAC
151  AGTGGAAATC  AAAAGAACTA  CTTGGCCTGG  TACCAGCAGA  AACCAAGGCA
201  GCTTCCATAA  CTGTTGATCT  ACGGGGCATC  CACTAGGGAA  TCTGGGGTCC
251  CTGATCGCTT  CACAGGCAGT  GGATCTGGAA  CAGATTTTAC  TCTCACCATC
301  AGCAGTGTGC  AGGCTGAAGA  CCTGGCAGTT  TATTACTGTC  AGAATGATTA
351  TAGTTATCCG  TACACGTTTC  GAGGGGGGAC  CAAGCTTGAG  ATCAAAGGTG
401  GTGCTGCTTC  TGGCGCGGCG  GGCTCCGGTG  GTGGTGGTTC  TGAGGTGCAG
451  CTGCTCGAGC  AGTCTGGAGC  TGAGCTGGTA  AGGCCTGGGA  CTT'CAGT'GAA
501  GATATCCTGC  AAGGCTTCTG  GATACGCCCT  CACTAACTAC  TGGCTAGGTT
551  GGGTTAAGCA  GAGGCCCTGA  CATGGACTTG  AATGGGTTGG  AGATA'TTTC
601  CCTGGAAGTG  GTAATGCTCA  CTACAATGAG  AAGTTCAAGG  GCAAAGCCAC
651  ACTGACTGCA  GACAAGT'CT  CGTACACAGC  CTATATGCAG  CTCAGTAGCC
701  TGACATCTGA  GGACTCTGCT  GTCTATTTCT  GTGCAAGATT  GCGGAAC'TGG
751  GACGAGGCTA  TGGACTACTG  GGGCCAAAGG  ACCACGGTCA  CCGTCTCCTC
801  CGGAGGTGGT  GGATCCGATA  TCAAACTGCA  GCAGTCAGGG  GCTGAAC'TGG
851  CAAGACCTGG  GGCCTCAGTG  AAGATGTCCT  GCAAGACTTC  TGGCTACACC
901  TTTACTAGGT  ACACGATGCA  CTGGGTAAAA  CAGAGGCCCTG  GACAGGGTCT

```

Figure 3D (continued)

```

951 GGAATGGATT GGATACATTA ATCCTAGCCG TGGTTATACT AATTACAATC
1001 AGAAGTTCAA GGACAAGGCC ACATTGACTA CAGACAAATC CTCACAGCACA
1051 GCCTACATGC AACTGAGCAG CCTGACATCT GAGGACTCTG CAGTCTATTA
1101 CTGTGCAAGA TATPATGATG ATCATTACTG CCTTGACTAC TGGGGCCAAG
1151 GCACCACTCT CACAGTCTCC TCAGTCGAAG GTGGAAGTGG AGGTTCTGGT
1201 GGAAGTGGAG GTTCAGGTGG AGTCGACGAC ATTCAGCTGA CCCAGTCTCC
1251 AGCAATCATG TCTGCCATCTC CAGGGGAGAA GGTCAACCATG ACCTGCAGAG
1301 CCAGTTC AAG TGTAAAGTTAC ATGAAGTGGT ACCAGCAGAA GTCAGGCACC
1351 TCCCCCAAAA GATGGATTTA TGACACATCC AAAGTGGCTT CTGGAGTCCC
1401 TTATCGCTTC AGTGGCAGTG GGTCTGGGAC CTCATACTCT CTCACAAATCA
1451 GCAGCATGGA GGCTGAAGAT GCTGCCACTT ATTACTGCCA ACAGTGGAGT
1501 AGTAACCCGC TCACGTTCCG TGCTGGGACC AAGCTGGAGC TGAAACATCA
1551 TCACCATCAT CATTAG

```

Figure 3E

5-10 (vLvH) x anti-CD3 (SEQ ID NO: 44)

```

1  MGWSCIIILFL VATAATGVHSE LVMTQSPSSL TVTAGEKVTM SCKSSQSLLN
51  SGNQKNYLTW YQKPGQPPK LLIYWASTRE SGVPDRFTGS GSGTDFTLTI
101 SSVQAEIDLAV YYQNDYSYP LTFGAGTKLE IKGGGGSGGG GSGGGGSEVQ
151 LLEQSGAELV RPTSVKISC KASGYAFTNY WLGWVKQRPQ HGLEWIGDIF
201 PGSGNIHYNE KFKGKATLTA DKSSSTAYMQ LSSLTFEDSA VYFCARLRNW
251 DEPMDYWGQG TTVTVSSGGG GSDIKLQQSG AELARPGASV KMSCKTSGYT
301 FTRYTMHWK QRPQGGLWI GYINPSRGYT NYNQKFKDKA TLTDDKSSST
351 AYMQLSSLTS EDSAVYYCAR YYDDHYCLDY WQQTTLTVS SVEGGSGGSG
401 GSGGGGVDD IQLTQSPAIM SASPGEKVTM TCRASSSVSY MNWYQQKSGT
451 SPKRWIYDTS KVASGVPIRF SSGSGTSYS LTSSMEAEED AATYYCQQWS
501 SNPLTFGAGT KLELKHSHHHH*

```


Figure 3E (continued)**SEQ ID NO: 43**

```

1  ATGGGATGGA GCTGTATCAT CCTCTTCTTG GTAGCAACAG CTACAGGTGT
51  ACACTCCGAG CTCGTGATGA CACAGTCTCC ATCCTCCCTG ACTGTGACAG
101 CAGGAGAGAA GGTCACTATG AGCTGCAAGT CCAGTCAGAG TCTGTAAAC
151 AGTGGAAATC AAAAGAACTA CTTGACCTGG TACCAGCAGA AACCAGGGCA
201 GCCTCCTAAA CTGTTGATCT ACTGGGCATC CACTAGGGAA TCTGGGGTCC
251 CTGATCGCTT CACAGGCAGT GGATCTGGAA CAGATTTCAC TCTCACCATC
301 AGCAGTGTGC AGGCTGAAGA CCTGGCAGTT TATTACTGTC AGAATGATTA
351 TAGTTATCCG CTCACGTTTC GTGCTGGGAC CAAGCTTGAG ATCAAAGGTG
401 GTGGTGGTTC TGGCGGCGGC GGCTCCGGTG GTGGTGGTTC TGAGGTGCAG
451 CTGCTCGAGC AGTCTGGAGC TGAGCTGGTA AGCCTGGGA CTTCACTGAA
501 GATATCCTGC AAGGCTTCTG GATACGCCCT CACTAACTAC TGGCTAGGTT
551 GGTAAAGCA GAGGCCTGGA CATGGACTTG AGTGGATTGG AGATATTTC
601 CCTGGAAGTG GTAATATCCA CTACAATGAG AAGTTCAAGG GCAAAGCCAC
651 ACTGACTGCA GACAAATCTT CGAGCACAGC CTATATGCAG CTCAGTAGCC
701 TGACATTGGA GGA CTCTGCT GTCTATTTCT GTGCAAGACT GAGGAAC TGG
751 GACGAGCCTA TGGACTACTG GGGCCAAGG ACCACGGTCA CCGTCTCTC
801 CGGAGGTGGT GGATCCGATA TCAAAC TGA GCACTGCA GCTGAAC TGG
851 CAAGACCTGG GGCCTCAGTG AAGATGTCCT GCAAGACTTC TGGCTACACC
901 TTACTAGGT ACACGATGCA CTGGGTAAAA CAGAGGCCCTG GACAGGGTCT

```


Figure 3E (continued)

```

951 GGAATGGATT GGATACATTG ATCCTAGCCG TGGTTATACT AATTACAATC
1001 AGAAGTTCAA GGACAAGGCC ACATTGACTA CAGACAAATC CTCCAGCACA
1051 GCCTACATGC AACTGAGCAG CCTGACATCT GAGGACTCTG CAGTCTATTA
1101 CTGTGCAAGA TATTATGATG ATCATTACTG CCTTGACTAC TGGGGCCAAAG
1151 GCACCACTCT CACAGTCTCC TCAGTCGAAG GTGGAAGTGG AGGTTCTGGT
1201 GGAAGTGGAG GTTCAGGTGG AGTCGACGAC ATTCAGCTGA CCCAGTCTCC
1251 AGCAATCATG TCTGCATCTC CAGGGAGAA GGTACCCATG ACCTGCAGAG
1301 CCAGTTCAAG TGTAAAGTTAC ATGAACTGGT ACCAGCAGAA GTCAGGCACC
1351 TCCCCCAAAA GATGGATTGA TGACACATCC AAAGTGGCTT CTGGAGTCCC
1401 TTATCGCTTC AGTGGCAGTG GGTCTGGGAC CTCATACTCT CTCACAATCA
1451 GCAGCATGGA GGCTGAAGAT GCTGCCACTT ATTACTGCCA ACAGTGGAGT
1501 AGTAACCCGC TCACGTTCCG TGCTGGGACC AAGCTGGAGC TGAACACATCA
1551 TCACCATCAT CATTAG

```

Figure 4A




4-7xanti-CD3
(SEQ ID NO:42)

Figure 4B

EpCAM Binding (Kato cells)

CD3 Binding (Jurkat cells)




3-5xanti-CD3

(SEQ ID NO: 30)

Figure 4C

CD3 Binding (Jurkat cells)

EpCAM Binding (Kato cells)



**3-1xanti-CD3
(SEQ ID NO:36)**

Figure 4D





Figure 4E

Figure 5





Figure 6






Figure 7B





Figure 8


Figure 9

Figure 10





Figure 11A

3-1	LR NW DEAMDY
4-1	LR NW DEAMDY
5-10	LR NW DEPMDY
3-5	RGSYGS NYD WYFDV
4-7	RGSYDT NYD WYFDV
M79	MENWSFAY
HD70	DMGWGSGWRPYYYYGMDV
3B10	FTSPDY

Figure 11B



1
SEQUENCE LISTING

<110> Micromet AG

<120> Pharmaceutical composition comprising a construct specific for EpCAM

<130> H1656 PCT

<150> EP 03012133.9

<151> 2003-05-31

<150> EP 03012134.7

<151> 2003-05-31

<160> 101

<170> PatentIn version 3.1

<210> 1

<211> 1521

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL aL x 4-7 VHVL

<400> 1

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagtc	360
gaagggtgaa gtggagggtc tgggtgaagt ggagggtcag gtggagtcga cgacattcag	420
ctgaccagat ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc	480
agagccagtt caagtgaag ttacatgaac tgggtaccagc agaagtcagg cacctcccc	540

2

```

aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg cttcagtggc 600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc 660
acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg 720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag 780
ctggcgaggc ctggggcttc agtgaagctg tcctgcaagg cttctggcta caccttcaca 840
aactatggtt taagctgggt gaagcagagg cctggacagg tccttgagtg gattggagag 900
gtttatccta gaattggtaa tgcttactac aatgagaagt tcaagggcaa ggccacactg 960
actgcagaca aatcctccag cacagcgtcc atggagctcc gcagcctgac ctctgaggac 1020
tctgcggtct atttctgtgc aagacgggga tcctacgata ctaactacga ctggtacttc 1080
gatgtctggg gccaaaggac caggtcacc gtctcctcag gtggtggtgg ttctggcggc 1140
ggcggctccg gtggtggtgg ttctgagctc gtgatgaccc agactccact ctccctgcct 1200
gtcagctctg gagatcaagc ctccatctct tgcagatcta gtcagagcct tgtacacagt 1260
aatgaaaca cctattaca ttggtacctg cagaagccag gccagctctc aaagctcctg 1320
atctacaaag ttccaaccg attttctggg gtcccagaca ggttcagtgg cagtggatca 1380
gggacagatt tcacactcaa gatcagcaga gtggaggctg aggatctggg agtttatttc 1440
tgcttcaaa gtacacatgt tccgtacacg ttcggagggg ggaccaagct tgagatcaaa 1500
catcatcacc atcatcatta g 1521

```

<210> 2

<211> 506

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL x 4-7 VHVL

<400> 2

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1      5      10      15

```

```

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20      25      30

```

```

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35      40      45

```

```

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50      55      60

```

```

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65      70      75      80

```

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala Ser Val Lys Leu Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Gly Leu Ser Trp Val Lys
 275 280 285
 Gln Arg Pro Gly Gln Val Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg
 290 295 300
 Ile Gly Asn Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr
 340 345 350

4

Asp Thr Asn Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr
 355 360 365
 Val Thr Val Ser Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly
 370 375 380
 Gly Gly Gly Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro
 385 390 395 400
 Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser
 405 410 415
 Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys
 420 425 430
 Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe
 435 440 445
 Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 450 455 460
 Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe
 465 470 475 480
 Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys
 485 490 495
 Leu Glu Ile Lys His His His His His His
 500 505

<210> 3

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL aL X 5-10 VHVL

<400> 3

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcttgcaaga cttctggcta cacccttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagtc	360
gaaggtggaa gtggaggttc tggtggaagt ggaggttcag gtggagtcga cgacattcag	420

5

```

ctgacccagt ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc 480
agagccagtt caagtgttaag ttacatgaac tggtagcagc agaagtcagg cacctcccc 540
aaaagatgga ttatgacac atccaaagtg gcttctggag tcccttatcg cticagtggc 600
agtggtgctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc 660
acttattact gccaacagtg gagtagtaac ccgtcacgt tgggtgctgg gaccaagctg 720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag 780
ctggtaaagg ctgggacttc agtgaagata tcttgaagg ctcttgata cgccttcact 840
aactactggc taggttgggt aaagcagagg cctggacatg gacttgagtg gattggagat 900
attttccctg gaagtggtaa tatccactac aatgagaagt tcaagggcaa agccacactg 960
actgcagaca aatcttcgag cacagcctat atgcagctca gtagcctgac atttgaggac 1020
tctgtgtctt atttctgtc aagactgagg aactgggacg agcctatgga ctactggggc 1080
caagggacca cggtcaccgt ctctcaggt ggtggtggtt ctggcggcgg cggctccggt 1140
ggtggtggtt ctgagctcgt gatgacacag tctccatcct ccctgactgt gacagcagga 1200
gagaagggtca ctatgagctg caagtccagt cagagtctgt taaacagtgg aatcaaaag 1260
aactacttga cctggtacca gcagaaacca gggcagcctc ctaaactgtt gatctactgg 1320
gcatccacta gggaatctgg ggtccctgat cgcttcacag gcagtggatc tggaacagat 1380
ttcactctca ccatcagcag tgtgcaggct gaagacctgg cagtttatta ctgtcagaat 1440
gattatagtt atccgctcac gttcgggtct gggaccaagc ttgagatcaa acatcatcac 1500
catcatcatt ag 1512

```

<210> 4

<211> 503

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL AL x 5-10 VHVL

<400> 4

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe

50 55 6 60
 Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Ile Gly Asp Ile Phe Pro Gly
 290 295 300
 Ser Gly Asn Ile His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu
 325 330 335

7

Thr Phe Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350

Asp Glu Pro Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365

Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 370 375 380

Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Thr Val Thr Ala Gly
 385 390 395 400

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
 405 410 415

Gly Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln
 420 425 430

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 435 440 445

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 450 455 460

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
 465 470 475 480

Asp Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Ile
 485 490 495

Lys His His His His His His
 500

<210> 5

<211> 31

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VH BsrGI primer

<400> 5

aggtgtacac tccgatatca aactgcagca g

31

<210> 6

<211> 25

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VL BspEI primer

<400> 6

aatccggatt tcagctccag cttgg

25

<210> 7

<211> 1521

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 4-7 VHVL

<400> 7

gatataaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctcagtc	360
gaagggtgaa gtggagggtc tgggtggaagt ggagggtcag gtggagtcga cgacattcag	420
ctgaccaggt ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc	480
agagccagtt caagtgttaag ttacatgaac tggtagcagc agaagtcagg cacctcccc	540
aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg cttcagtggc	600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc	660
acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg	720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag	780
ctggcgaggc ctggggcttc agtgaagctg tcctgcaagg cttctggcta caccttcaca	840
aactatggtt taagctgggt gaagcagagg cctggacagg tccttgagtg gattggagag	900
gtttatccta gaattggtaa tgcttactac aatgagaagt tcaagggcaa ggccacactg	960
actgcagaca aatcctccag cacagcgtcc atggagctcc gcagcctgac ctctgaggac	1020
tctgcggtct atttctgtgc aagacgggga tcctacgata ctaactacga ctgggtacttc	1080
gatgtctggg gccaaaggac cagggtcacc gtctcctcag gtggtggtgg ttctggcggc	1140
ggcggctccg gtggtggtgg ttctgagctc gtgatgacct agactccact ctccctgcct	1200
gtcagtcctg gagatcaagc ctccatctct tgcagatcta gtcagagcct tgtacacagt	1260

9

aatggaaaca cctatttaca ttggtacctg cagaagccag gccagtcctc aaagctcctg 1320
 atctacaaag ttccaaccg atttctggg gtcccagaca gggtcagtg cagtggtatca 1380
 gggacagatt tcacactcaa gatcagcaga gtggaggctg aggatctggg agtttatttc 1440
 tgctctcaaa gtacacatgt tccgtacacg ttcggagggg ggaccaagct tgagatcaaa 1500
 catcatcacc atcatcatta g 1521

<210> 8

<211> 506

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 4-7 VHVL

<400> 8

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160

Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser

165 170¹⁰ 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala Ser Val Lys Leu Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr Gly Leu Ser Trp Val Lys
 275 280 285
 Gln Arg Pro Gly Gln Val Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg
 290 295 300
 Ile Gly Asn Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr
 340 345 350
 Asp Thr Asn Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr
 355 360 365
 Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Gly
 370 375 380
 Gly Gly Gly Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro
 385 390 395 400
 Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser
 405 410 415
 Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys
 420 425 430
 Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe
 435 440 445

11

Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
450 455 460

Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe
465 470 475 480

Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys
485 490 495

Leu Glu Ile Lys His His His His His His
500 505

<210> 9

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL AL Ser x 5-10 VHVL

<400> 9

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga cttctggcta cacccttact aggtacacga tgcactgggt aaaacagagg	120
cctggcaggg gtctggaatg gattggatac attaatccta gccgtggtta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctcagtc	360
gaagggtgaa gtggagggtc tgggtggaagt ggagggttcag gtggagtcga cgacattcag	420
ctgacccagt ctccagcaat catgtctgca tctccagggg agaagggtcac catgacctgc	480
agagccagtt caagtgttaag ttacatgaac tggtagcagc agaagtcagg cactcccc	540
aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg cttcagtggc	600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc	660
acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg	720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag	780
ctggttaaggc ctgggacttc agtgaagata tcttgcaagg cttctggata cgccttcact	840
aactactggc taggttgggt aaagcagagg cctggacatg gacttgagtg gattggagat	900
attttccctg gaagtggtaa tatccactac aatgagaagt tcaagggcaa agccacactg	960
actgcagaca aatcttcgag cacagcctat atgcagctca gtagcctgac atttgaggac	1020
tctgctgtct atttctgtgc aagactgagg aactgggacg agcctatgga ctactggggc	1080

12

caagggacca cggtcaccgt ctctcaggt ggtggtggtt ctggcggcgg cggctccggt	1140
ggtggtggtt ctgagctcgt gatgacacag tctccatcct cctgactgt gacagcagga	1200
gagaaggta ctatgagctg caagtccagt cagagtctgt taaacagtgg aaatcaaaag	1260
aactacttga cctggtacca gcagaaacca gggcagcctc ctaaactgtt gatctactgg	1320
gcattcacta ggaatctgg ggtccctgat cgttcacag gcagtggatc tggaacagat	1380
ttcactctca ccatcagcag tgtgcaggct gaagacctgg cagtttatta ctgtcagaat	1440
gattatagtt atccgctcac gttcgggtgct gggaccaagc ttgagatcaa acatcatcac	1500
catcatcatt ag	1512

<210> 10

<211> 503

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 5-10 VHVL

<400> 10

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu¹³ Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Ile Gly Asp Ile Phe Pro Gly
 290 295 300
 Ser Gly Asn Ile His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu
 325 330 335
 Thr Phe Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350
 Asp Glu Pro Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 370 375 380
 Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Thr Val Thr Ala Gly
 385 390 395 400
 Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
 405 410 415
 Gly Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln

14

420 425 430

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 435 440 445

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 450 455 460

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
 465 470 475 480

Asp Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Ile
 485 490 495

Lys His His His His His
 500

<210> 11

<211> 1485

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 3-1 VHVL

<400> 11

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtggtta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcaggt	360
ggtggtggtt ctggcggcgg cggtccggt ggtggtggtt ctgacattca gctgaccag	420
tctccagcaa tcatgtctgc atctccaggg gagaaggta ccatgacctg cagagccagt	480
tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaagatgg	540
atttatgaca catccaaagt ggcttctgga gtcccttata gcttcagtgg cagtgggtct	600
gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac	660
tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa	720
tccggagggt gtggatccga ggtgcagctg ctcgagcagt ctggagctga gctggtgaaa	780
cctggggcct cagtgaagat atcctgcaag gcttctggat acgccttcac taactactgg	840
ctaggttggg taaagcagag gcctggacat ggacttgagt ggattggaga tcttttcct	900
ggaagtggta atactcacta caatgagagg ttacggggca aagccacact gactgcagac	960

15

```

aaatcctcga gcacagcctt tatgcagctc agtagcctga catctgagga ctctgctgtc 1020
tatttctgtg caagattgag gaactgggac gaggttatgg actactgggg ccaagggacc 1080
acggtcaccg tctcctcagg tgggtggtgt tctggcggcg gcggctccgg tgggtggtgt 1140
tctgagctcg tcatgaccca gtctccatct tatcttctgt catctcctgg agaaaccatt 1200
actattaatt gcagggcaag taagagcatt agcaaatatt tagcctggta tcaagagaaa 1260
cctgggaaaa ctaataagct tcttatctac tctggatcca ctttgcaatc tggaattcca 1320
tcaaggttca gtggcagtgg atctgttaca gatttcactc tcaccatcag tagcctggag 1380
cctgaagatt ttgcaatgta ttactgtcaa cagcataatg aatatccgta cacgttcgga 1440
ggggggacca agcttgagat caaacatcat caccatcatc attag 1485

```

<210> 12

<211> 494

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 3-1 VHVL

<400> 12

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 115 120 125

Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile

130 135 16 140
 Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160
 Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175
 Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190
 Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205
 Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220
 Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240
 Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 245 250 255
 Glu Leu Val Lys Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser
 260 265 270
 Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro
 275 280 285
 Gly His Gly Leu Glu Trp Ile Gly Asp Leu Phe Pro Gly Ser Gly Asn
 290 295 300
 Thr His Tyr Asn Glu Arg Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp
 305 310 315 320
 Lys Ser Ser Ser Thr Ala Phe Met Gln Leu Ser Ser Leu Thr Ser Glu
 325 330 335
 Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Ala
 340 345 350
 Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly
 355 360 365
 Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Leu Val
 370 375 380
 Met Thr Gln Ser Pro Ser Tyr Leu Ala Ala Ser Pro Gly Glu Thr Ile
 385 390 395 400
 Thr Ile Asn Cys Arg Ala Ser Lys Ser Ile Ser Lys Tyr Leu Ala Trp
 405 410 415

17

Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys Leu Leu Ile Tyr Ser Gly
 420 425 430

Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly Ser Gly Ser
 435 440 445

Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu Asp Phe
 450 455 460

Ala Met Tyr Tyr Cys Gln Gln His Asn Glu Tyr Pro Tyr Thr Phe Gly
 465 470 475 480

Gly Gly Thr Lys Leu Glu Ile Lys His His His His His His
 485 490

<210> 13

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 4-7 VHVL

<400> 13
 gatatacaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcttgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attacigtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcaggt 360
 ggtggtggtt ctggcggcgg cggctccggt ggtggtggtt ctgacattca gctgaccag 420
 tctccagcaa tcatgtctgc atctccaggg gagaaggta ccatgacctg cagagccagt 480
 tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaaagatgg 540
 atttatgaca catccaaagt ggcttctgga gtcccttata gcttcagtgg cagtgggtct 600
 gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 660
 tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa 720
 tccggagggtg gtggatccga ggtgcagctg ctgagcagct ctggagctga gctggcgagg 780
 cctggggctt cagtgaagct gtcttgaag gcttctggt acaccttcac aaactatggt 840
 ttaagctggg tgaagcagag gcctggacag gtccttgagt ggattggaga ggtttatcct 900
 agaattggtg atgcttacta caatgagaag ttcaagggca aggccacact gactgcagag 960

18

```

aaatcctcca gcacagcgtc catggagctc cgcagcctga cctctgagga ctctgcggtc 1020
tatttctgtg caagacgggg atcctacgat actaactacg actggtactt cgatgtctgg 1080
ggccaaggga ccacgggtcac cgtctcctca ggtggtggtg gttctggcgg cggcggctcc 1140
ggtggtggtg gttctgagct cgtgatgacc cagactccac tctccctgcc tgtcagtcctt 1200
ggagatcaag cctccatctc ttgcagatct agtcagagcc ttgtacacag taatggaaac 1260
acctatttac attggtacct gcagaagcca ggccagtctc caaagctcct gatctacaaa 1320
gtttccaacc gattttctgg ggtcccagac aggttcagtg gcagtggtac agggacagat 1380
ttcacactca agatcagcag agtggaggct gaggatctgg gattttatct ctgctctcaa 1440
agtacacatg ttccgtacac gttcggaggg gggaccaagc ttgagatcaa acatcatcac 1500
catcatcatt ag 1512

```

<210> 14
 <211> 503
 <212> PRT
 <213> artificial sequence

<220>
 <223> CD3 VHVL stL x 4-7 VHVL
 <400> 14

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1      5      10      15
Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20     25     30
Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35     40     45
Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50     55     60
Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65     70     75     80
Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85     90     95
Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
100    105    110
Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
115    120    125

```

Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile
 130 135 140
 Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160
 Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175
 Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190
 Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205
 Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220
 Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240
 Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 245 250 255
 Glu Leu Ala Arg Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser
 260 265 270
 Gly Tyr Thr Phe Thr Asn Tyr Gly Leu Ser Trp Val Lys Gln Arg Pro
 275 280 285
 Gly Gln Val Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn
 290 295 300
 Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp
 305 310 315 320
 Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu Thr Ser Glu
 325 330 335
 Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr Asp Thr Asn
 340 345 350
 Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val
 355 360 365
 Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 370 375 380
 Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu
 385 390 395 400
 Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His

<400> 15	gataatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
	tctctgaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg	120
	cctggacagg gtctggaatg gattggatac attaatccta gccgtggtta tactaattac	180
	aatcagaagt tcaaggacaa ggcacattg actacagaca aatcctccag cacagcctac	240
	atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
	gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcaggt	360
	ggtggtggtt ctggcggcgg cggctccggt ggtggtggtt ctgacattca gctgaccacg	420
	tctccagcaa tcatgtctgc atctccaggg gagaaggcca ccatgacctg cagagccagt	480
	tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaagatgg	540
	atttatgaca catccaaagt ggcttctgga gtccctttat gttctcagtg cagtgggtct	600
	gggacctcat actctctcac aatcagcagc atggaggctg aagatgtctc cacttattac	660
	tgccaacagt ggagtagtaa cccgctcacg ttccggtctg ggaccaagct ggagctgaaa	720
	tccggagggt gtggatccga gctcgtgatg acccagactc cactctccct gcctgtcagt	780
	cttgagatc aagcctccat ctcttcgaga tctagtcaga gccttgata cagtaatgga	840

21

```

aacacctatt tacattgga cctgcagaag ccaggccagt ctccaaagct cctgatctac    900
aaagtttcca accgattttc tgggttccca gacaggttca gtggcagtg atcaggga    960
gatttcacac tcaagatcag cagagtggag gctgaggatc tgggagtta tttctgctct    1020
caaagtacac atgttccgta cacgttcgga ggggggacca agcttgagat caaagggtgt    1080
ggtggttctg gcggcggcgg ctccggtggt ggtggttctg aggtgcagct gctcgagcag    1140
tctggagctg agctggcgag gcctggggct tcagtgaagc tgcctgcaa ggccttctggc    1200
tacaccttca caaactatgg tttaagctgg gtgaagcaga ggcctggaca ggccttgag    1260
tggattggag aggtttatcc tagaattggt aatgcttact acaatgagaa gttcaagggc    1320
aaggccacac tgactgcaga caaatcctcc agcacagcgt ccattggagct ccgcagcctg    1380
acctctgagg actctgcggt ctatttctgt gcaagacggg gatcctacga tactaactac    1440
gactggtact tcgatgtctg gggccaaggg accacggtca ccgtctctc acatcatcac    1500
catcatcatt ag    1512

```

<210> 16

<211> 503

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 4-7 VLVH

<400> 16

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1           5           10           15

```

```

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20           25           30

```

```

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35           40           45

```

```

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50           55           60

```

```

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65           70           75           80

```

```

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85           90           95

```

```

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
100          105          110

```

22

Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 115 120 125
 Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile
 130 135 140
 Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160
 Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175
 Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190
 Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205
 Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220
 Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240
 Ser Gly Gly Gly Gly Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser
 245 250 255
 Leu Pro Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser
 260 265 270
 Gln Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu
 275 280 285
 Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn
 290 295 300
 Arg Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr
 305 310 315 320
 Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val
 325 330 335
 Tyr Phe Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly
 340 345 350
 Thr Lys Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly Gly Gly Ser
 355 360 365
 Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu
 370 375 380

Leu Ala Arg Pro Gly Ala Ser Val Lys Leu ²³ Ser Cys Lys Ala Ser Gly
 385 390 395 400
 Tyr Thr Phe Thr Asn Tyr Gly Leu Ser Trp Val Lys Gln Arg Pro Gly
 405 410 415
 Gln Val Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn Ala
 420 425 430
 Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys
 435 440 445
 Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu Thr Ser Glu Asp
 450 455 460
 Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr Asp Thr Asn Tyr
 465 470 475 480
 Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 485 490 495
 Ser His His His His His His
 500

<210> 17

<211> 1503

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 5-10 VHVL

<400> 17
 gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagggt 360
 ggtggtggtt ctggcggcgg cggtccgggt ggtggtggtt ctgacattca gctgaccag 420
 tctccagcaa tcatgtctgc atctccaggg gagaaggta ccatgacctg cagagccagt 480
 tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaagatgg 540
 atttatgaca catccaaagt ggcttctgga gtcccttacc gcttcagtgg cagtgggtct 600
 gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 660

24

tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa	720
tccggagggtg gtggatccga ggtgcagctg ctcgagcagt ctggagctga gctggtaagg	780
cctgggactt cagtgaagat atcctgcaag gcttctggat acgccttcac taactactgg	840
ctaggttggg taaagcagag gcctggacat ggacttgagt ggattggaga tattttccct	900
ggaagtggta atatecacta caatgagaag ttcaagggca aagccacact gactgcagac	960
aaatcttcga gcacagccta tatgcagctc agtagcctga catttgagga ctctgctgtc	1020
tattttctgt caagactgag gaactgggac gagcctatgg actactgggg ccaagggacc	1080
acggtcaccg tctcctcagg tgggtggtgt tctggcggcg gcggctccgg tgggtggtgt	1140
tctgagctcg tgatgacaca gtctccatcc tccctgactg tgacagcagg agagaaggtc	1200
actatgagct gcaagtccag tcagagtctg ttaaacagtg gaaatcaaaa gaactacttg	1260
acctggatcc agcagaaacc agggcagcct cctaaactgt tgatctactg ggcattccact	1320
agggaatctg gggctcctga tcgcttcaca ggcagtggtat ctggaacaga tttcactctc	1380
accatcagca gtgtgcaggc tgaagacctg gcagtttatt actgtcagaa tgattatagt	1440
tatccgctca cgttcggtgc tgggaccaag cttgagatca aacatcatca ccacatcat	1500
tag	1503

<210> 18

<211> 500

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 5-10 VHVL

<400> 18

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

25

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 115 120 125
 Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile
 130 135 140
 Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160
 Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175
 Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190
 Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205
 Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220
 Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240
 Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 245 250 255
 Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys Lys Ala Ser
 260 265 270
 Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro
 275 280 285
 Gly His Gly Leu Glu Trp Ile Gly Asp Ile Phe Pro Gly Ser Gly Asn
 290 295 300
 Ile His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp
 305 310 315 320
 Lys Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Phe Glu
 325 330 335
 Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Pro
 340 345 350
 Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly
 355 360 365

26

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Leu Val
 370 375 380
 Met Thr Gln Ser Pro Ser Ser Leu Thr Val Thr Ala Gly Glu Lys Val
 385 390 395 400
 Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser Gly Asn Gln
 405 410 415
 Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys
 420 425 430
 Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val Pro Asp Arg
 435 440 445
 Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 450 455 460
 Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn Asp Tyr Ser
 465 470 475 480
 Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Ile Lys His His
 485 490 495
 His His His His
 500

<210> 19

<211> 1503

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 5-10 VLVH

<400> 19
 gatatacaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcttgcaga cttctggcta cactttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcaggt 360
 ggtggtggtt ctggcggcgg cggtccggt ggtggtggtt ctgacattca gctgaccacg 420
 tctccagcaa tcatgtctgc atctccaggg gagaaggcca ccatgacctg cagagccagt 480
 tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaagatgg 540

27

```

atttatgaca catccaaagt ggcttctgga gtcccttata gcttcagtgg cagtgggtct 600
gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 660
tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa 720
tccggagggtg gtggatccga gctcgtgatg acacagtctc catcctccct gactgtgaca 780
gcaggagaga aggtcactat gagctgcaag tccagtcaga gtctgttaaa cagtggaaat 840
caaaagaact acttgacctg gtaccagcag aaaccagggc agcctcctaa actgttgatc 900
tactgggcat ccactaggga atctggggtc cctgatcgct tcacaggcag tggatctgga 960
acagatttca ctctcaccat cagcagtgtg caggctgaag acctggcagt ttattactgt 1020
cagaatgatt atagttatcc gctcacgttc ggtgctggga ccaagcttga gatcaaaggt 1080
ggtggtggtt ctggcgccgg cggctccggt ggtggtggtt ctgagggtga gctgctcgag 1140
cagtctggag ctgagctggt aaggcctggg acttcagtga agatatctg caaggcttct 1200
ggatacgctt tcactaacta ctggctaggt tgggtaaagc agaggcctgg acatggactt 1260
gagtggtatt gagatatttt ccctggaagt ggtaatatcc actacaatga gaagttcaag 1320
ggcaaagcca cactgactgc agacaaatct tcgagcacag cctatatgca gctcagtagc 1380
ctgacatttg aggactctgc tgtctatttc tgtgcaagac tgaggaactg ggacgagcct 1440
atggactact ggggccaagg gaccacggtc accgtctcct cacatcatca ccatcatcat 1500
tag 1503

```

<210> 20

<211> 500

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 5-10 VL VH

<400> 20

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1           5           10           15

```

```

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20           25           30

```

```

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35           40           45

```

```

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50           55           60

```

```

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr

```

-164-

29

Gly Thr Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Gly Gly Gly Gly
 355 360 365

Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 370 375 380

Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys Lys Ala Ser
 385 390 395 400

Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro
 405 410 415

Gly His Gly Leu Glu Trp Ile Gly Asp Ile Phe Pro Gly Ser Gly Asn
 420 425 430

Ile His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp
 435 440 445

Lys Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Phe Glu
 450 455 460

Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Pro
 465 470 475 480

Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser His His
 485 490 495

His His His His
 500

<210> 21

<211> 57

<212> DNA

<213> artificial sequence

<220>

<223> 3' CD3 VH GS15 primer

<400> 21
 ggagccgccc ccgccagaac caccaccacc tgaggagact gtgagagtgg tgccttg

57

<210> 22

<211> 53

<212> DNA

<213> artificial sequence

30

<220>

<223> 5' CD3 VL GS15 primer

<400> 22

ggcggcggcg gctccggtgg tgggtggttct gacattcagc tgaccagtc tcc 53

<210> 23

<211> 51

<212> DNA

<213> artificial sequence

<220>

<223> 4-7 VH GS15 FOR

<400> 23

ggcggcggcg gctccggtgg tgggtggttct gaggtgcagc tgctcgagca g 51

<210> 24

<211> 53

<212> DNA

<213> artificial sequence

<220>

<223> 4-7 VH SalI REV primer

<400> 24

ttttaagtcg acctaatgat gatgatgatg atgtgaggag acggtgaccg tgg 53

<210> 25

<211> 49

<212> DNA

<213> artificial sequence

<220>

<223> 5-10 VLBspEI38 primer

<400> 25

ctgaaatccg gaggtgggtgg atccgagctc gtgatgacac agtctccat 49

<210> 26

<211> 53

<212> DNA

31

<213> artificial sequence

<220>

<223> 5-10 VLGS15REV primer

<400> 26
 ggagccgccc cgcgcagaac caccaccacc ttgatctca agcttgggtcc cag 53

<210> 27

<211> 49

<212> DNA

<213> artificial sequence

<220>

<223> 5-10 VH GS15 FOR primer

<400> 27
 ggcgcgcgcg gctccggtgg tgggtgttct gaggtgcagc tgctcgagc 49

<210> 28

<211> 53

<212> DNA

<213> artificial sequence

<220>

<223> 5-10 VHSa1REV primer

<400> 28
 ttttaagtcg acctaatgat gatgatgatg atgtgaggag acggtgaccg tgg 53

<210> 29

<211> 1581

<212> DNA

<213> artificial sequence

<220>

<223> 3-5(VL-VH)xanti-CD3

<400> 29
 atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacaggtgt acactccgcg 60
 cgcgagctcg tgatgaccca gactccactc tccctgcctg tcagtcttgg agatcaagcc 120
 tccatctctt gcagatctag tcagagcctt gtacacagta atggaaacac ctattttacat 180

32

tggtagctgc	agaagccagg	ccagctctcca	aagctcctga	tctacaaagt	ttccaaccga	240
ttttctgggg	tcccagacag	gttcagtggc	agtggatcag	ggacagattt	cacactcaag	300
atcagcagag	tggaggctga	ggatctggga	gtttatttct	gctctcaaag	tacacatgtt	360
ccgtacacgt	tcggaggggg	gaccaagctt	gagatcaaag	gtggtggtgg	ttctggcggc	420
ggcggctccg	gtggtggtgg	ttctgaggtg	cagctgctcg	agcagtcctg	agctgagctg	480
gtaaggcctg	ggacttcagt	gaagctgtcc	tgcaaggctt	ctggctacac	cttcacaagc	540
tatgggttaa	gctgggtgaa	gcagagaact	ggacagggcc	ttgagtggat	tggagaggtt	600
tatcctagaa	ttggtaatgc	ttactacaat	gagaagtcca	agggcaaggc	cacactgact	660
gcagacaaat	cctccagcac	agcgtccatg	gagctccgca	gcctgacatc	tgaggactct	720
gcgggtctatt	tctgtgcaag	acggggatcc	tacggtagta	actacgactg	gtacttcgat	780
gtctggggcc	aagggaccac	ggtcaccgtc	tcctccggag	gtggtggatc	cgatatcaaa	840
ctgcagcagt	caggggctga	actggcaaga	cctggggcct	cagtgaagat	gtcctgcaag	900
acttctggct	acacctttac	taggtacacg	atgcactggg	taaaacagag	gcctggacag	960
ggtctggaat	ggatttgata	cattaatcct	agccgtgggt	atactaatta	caatcagaag	1020
ttcaaggaca	agccacatt	gactacagac	aaatcctcca	gcacagccta	catgcaactg	1080
agcagcctga	catctgagga	ctctgcagtc	tattactgtg	caagatatta	tgatgatcat	1140
tactgccttg	actactgggg	ccaaggcacc	actctcacag	tctctcagt	cgaagggtga	1200
agtggaggtt	ctggtggaag	tggaggttca	ggtggagtcg	acgacattca	gctgaccacg	1260
tctccagcaa	tcatgtctgc	atctccaggg	gagaaggtca	ccatgacctg	cagagccagt	1320
tcaagtgtaa	gttacatgaa	ctggtaccag	cagaagtcag	gcacctcccc	caaaagatgg	1380
atttatgaca	catccaaagt	ggcttctgga	gtcccttata	gcttcagtgg	cagtgggtct	1440
gggacctcat	actctctcac	aatcagcagc	atggaggctg	aagatgctgc	cacttattac	1500
tgccaacagt	ggagtagtaa	cccgtcacg	ttcgggtcctg	ggaccaagct	ggagctgaaa	1560
catcatcacc	atcatcatta	g				1581

<210> 30

<211> 526

<212> PRT

<213> artificial sequence

<220>

<223> 3-5(VL-VH)xanti-CD3

<400> 30

Met	Gly	Trp	Ser	Cys	Ile	Ile	Leu	Phe	Leu	Val	Ala	Thr	Ala	Thr	Gly
1				5					10					15	

Val His Ser Ala Arg Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu
 20 25 30 33
 Pro Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln
 35 40 45
 Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln
 50 55 60
 Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg
 65 70 75 80
 Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp
 85 90 95
 Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr
 100 105 110
 Phe Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr
 115 120 125
 Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Gly Gly Gly Ser Gly
 130 135 140
 Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu
 145 150 155 160
 Val Arg Pro Gly Thr Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr
 165 170 175
 Thr Phe Thr Ser Tyr Gly Leu Ser Trp Val Lys Gln Arg Thr Gly Gln
 180 185 190
 Gly Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn Ala Tyr
 195 200 205
 Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser
 210 215 220
 Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser
 225 230 235 240
 Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr Gly Ser Asn Tyr Asp
 245 250 255
 Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 260 265 270
 Gly Gly Gly Gly Ser Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu
 275 280 285
 Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr

290 295 34 300
 Thr Phe Thr Arg Tyr Thr Met His Trp Val Lys Gln Arg Pro Gly Gln
 305 310 315 320
 Gly Leu Glu Trp Ile Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn
 325 330 335
 Tyr Asn Gln Lys Phe Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser
 340 345 350
 Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser
 355 360 365
 Ala Val Tyr Tyr Cys Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp
 370 375 380
 Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly
 385 390 395 400
 Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile
 405 410 415
 Gln Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys
 420 425 430
 Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp
 435 440 445
 Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr
 450 455 460
 Ser Lys Val Ala Ser Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser
 465 470 475 480
 Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala
 485 490 495
 Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly
 500 505 510
 Ala Gly Thr Lys Leu Glu Leu Lys His His His His His His
 515 520 525

<210> 31

<211> 40

<212> DNA

<213> artificial sequence

35

<220>

<223> Me81 primer

<400> 31

ggatgcgcgc gagctcgtga tgaccagac tccactctcc

40

<210> 32

<211> 60

<212> DNA

<213> artificial sequence

<220>

<223> Me83 primer

<400> 32

ggttctggcg gcggcgctc cgggtggtgt ggttctgagg tgcagctgct cgacagtctg

60

<210> 33

<211> 41

<212> DNA

<213> artificial sequence

<220>

<223> Me84 primer

<400> 33

gtgctccgga ggagacggtg accgtggtcc cttggcccca g

41

<210> 34

<211> 53

<212> DNA

<213> artificial sequence

<220>

<223> Me90 primer

<400> 34

ccggagccgc cgccgccaga accaccacca ctttgatct caagcttggt ccc

53

<210> 35

<211> 1548

<212> DNA

<213> artificial sequence

<220>

<223> 3-1(VLVH)xanti-cd3

<400> 35

```

atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacaggtgt acactccgag    60
ctcgtcatga cccagtctcc atcttatctt gctgcatctc ctggagaaac cattactatt    120
aattgcaggg caagtaagag cattagcaaa tatitagcct ggtatcaaga gaaacctggg    180
aaaactaata agcttcttat ctactctgga tccactttgc aatctggaat tccatcaagg    240
ttcagtgga gtagatctgg tacagatttc actctcacca tcagtagcct ggagcctgaa    300
gattttgcaa tgtattactg tcaacagcat aatgaatata cgtacacgtt cggagggggg    360
accaagcttg agatcaaagg tgggtggtgt tctggcggcg gcggctccgg tgggtggtgt    420
tctgaggtgc agctgctcga gcagtctgga gctgagctgg tgaaacctgg ggcctcagtg    480
aagatactct gcaaggcttc tggatacgcc ttcactaact actggctagg ttgggtaaaag    540
cagaggcctg gacatggact tgagtggatt ggagatcttt tccctggaag tggtataact    600
cactacaatg agaggttcag gggcaaagcc aactgactg cagacaaatc ctcgagcaca    660
gcctttatgc agctcagtag cctgacatct gaggactctg ctgtctatct ctgtgcaaga    720
ttgaggaact gggacgaggc tatggactac tggggccaag ggaccacggt caccgtctcc    780
tccggagggtg gtggatccga tatcaaactg cagcagtcag gggctgaact ggcaagacct    840
ggggcctcag tgaagatgtc ctgcaagact tctggctaca cctttactag gtacacgatg    900
cactgggtraa aacagaggcc tggacagggt ctggaatgga ttggatacat taatcctagc    960
cgtggttata ctaattacaa tcagaagtgc aaggacaagg ccacattgac tacagacaaa   1020
tcctccagca cagcctacat gcaactgagc agcctgacat ctgaggactc tgcagtcctat   1080
tactgtgcaa gatattatga tgatcattac tgacctgact actggggcca aggcaccact   1140
ctcacagtct cctcagtcga aggtggaagt ggaggttctg gtggaagtgg aggttcaggt   1200
ggagtgcagc acattcagct gaccagctct ccagcaatca tgtctgcata tccaggggag   1260
aaggtcacca tgacctgcag agccagttca agtgtaagtt acatgaactg gtaccagcag   1320
aagtcaggca cctcccccaa aagatggatt tatgacacat ccaaagtggc ttctggagtc   1380
ccttatcgct tcagtggcag tgggtctggg acctcatact ctctcacaat cagcagcatg   1440
gaggctgaag atgctgccac ttattactgc caacagtgga gtagtaacct gctcacgttc   1500
ggtgctggga ccaagctgga gctgaaacat catcaccatc atcattag    1548

```

<210> 36

<211> 515

<212> PRT

<213> artificial sequence

<220>

<223> 3-1(VLVH)xanti-CD3

<400> 36

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
 1 5 10 15
 Val His Ser Glu Leu Val Met Thr Gln Ser Pro Ser Tyr Leu Ala Ala
 20 25 30
 Ser Pro Gly Glu Thr Ile Thr Ile Asn Cys Arg Ala Ser Lys Ser Ile
 35 40 45
 Ser Lys Tyr Leu Ala Trp Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys
 50 55 60
 Leu Leu Ile Tyr Ser Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg
 65 70 75 80
 Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 85 90 95
 Leu Glu Pro Glu Asp Phe Ala Met Tyr Tyr Cys Gln Gln His Asn Glu
 100 105 110
 Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Gly Gly
 115 120 125
 Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Glu Val Gln
 130 135 140
 Leu Leu Glu Gln Ser Gly Ala Glu Leu Val Lys Pro Gly Ala Ser Val
 145 150 155 160
 Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu
 165 170 175
 Gly Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp Ile Gly Asp
 180 185 190
 Leu Phe Pro Gly Ser Gly Asn Thr His Tyr Asn Glu Arg Phe Arg Gly
 195 200 205
 Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Phe Met Gln
 210 215 220
 Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg
 225 230 235 240
 Leu Arg Asn Trp Asp Glu Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr

245 250 38 255
 Val Thr Val Ser Ser Gly Gly Gly Gly Ser Asp Ile Lys Leu Gln Gln
 260 265 270
 Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys
 275 280 285
 Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr Thr Met His Trp Val Lys
 290 295 300
 Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly Tyr Ile Asn Pro Ser
 305 310 315 320
 Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe Lys Asp Lys Ala Thr Leu
 325 330 335
 Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu
 340 345 350
 Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala Arg Tyr Tyr Asp Asp
 355 360 365
 His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val Ser
 370 375 380
 Ser Val Glu Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly
 385 390 395 400
 Gly Val Asp Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile Met Ser Ala
 405 410 415
 Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val
 420 425 430
 Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg
 435 440 445
 Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro Tyr Arg Phe
 450 455 460
 Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met
 465 470 475 480
 Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn
 485 490 495
 Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys His His His
 500 505 510
 His His His
 515

<210> 37
 <211> 52
 <212> DNA
 <213> artificial sequence

<220>

<223> Me91a primer

<400> 37
 ggattgtaca ctccgagctc gtcattgaccc agtctccatc ttatcttgct gc 52

<210> 38
 <211> 1566
 <212> DNA
 <213> artificial sequence

<220>

<223> 4-1(VLVH)xanti-CD3

<400> 38
 atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacaggtgt acactccgag 60
 ctctgtgatga cacagtctcc atcctccctg agtgtgtcag caggagagaa ggtcactatg 120
 agctgcaagt ccagtcagag tctgttaaac agtggaaatc aaaagaacta cttggccttg 180
 taccagcaga aaccagggca gcctcctaaa ctgttgatct acggggcatc cactagggaa 240
 tctgggtgcc ctgacgctt cacaggcagt ggatctggaa cagatttcac tctcaccatc 300
 agcagtgctc aggctgaaga cctggcagtt tattactgtc agaattgatta tagttatccg 360
 tacacgttcg gaggggggac caagcttgag atcaaagggtg gtggtggttc tggcggcggc 420
 ggctccggtg gtggtggttc tgaggtgcag ctgctcgagc agtctggagc tgagctggta 480
 aggcctggga cttcagtga gatattctgc aaggcttctg gatacgctt cactaactac 540
 tggctaggtt ggggttaagca gaggcctgga catggacttg aatgggttg agatatttc 600
 cctggaagtg gtaatgctca ctacaatgag aagttcaagg gcaaagccac actgactgca 660
 gacaagtcct cgtacacagc ctatatgcag ctcatgagc tgacatctga ggactctgct 720
 gtctatttct gtgcaagatt gcggaactgg gacgaggcta tggactactg gggccaaggg 780
 accacggcta ccgtctcctc cggagggtgt ggatccgata tcaaactgca gcagtcaggg 840
 gctgaactgg caagacctgg ggcctcagt aagatgtcct gcaagacttc tggctacacc 900
 ttactaggt acacgatgca ctgggtaaaa cagaggcctg gacagggtct ggaatggatt 960
 ggatacatta atcctagccg tgggttatact aattacaatc agaagttcaa ggacaaggcc 1020

40

```

acattgacta cagacaaatc ctccagcaca gcctacatgc aactgagcag cctgacatct 1080
gaggactctg cagtctatta ctgtgcaaga tattratgatg atcattactg ccttgactac 1140
tggggccaag gcaccactct cacagtctcc tcagtcgaag gtggaagtgg aggttctggt 1200
ggaagtggag gttcaggtgg agtcgacgac attcagctga ccagtcctcc agcaatcatg 1260
tctgcatctc caggggagaa ggtcaccatg acctgcagag ccagttcaag tgtaagttac 1320
atgaactggt accagcagaa gtcaggcacc tcccccaaaa gatggattta tgacacatcc 1380
aaagtggctt ctggagtccc ttatcgcttc agtggcagtg ggtctgggac ctcatactct 1440
ctcacaatca gcagcatgga ggctgaagat gctgccactt attactgcca acagtggagt 1500
agtaaccctg tcacgttcgg tgctgggacc aagctggagc tgaaacatca tcaccatcat 1560
cattag 1566

```

<210> 39
 <211> 521
 <212> PRT
 <213> artificial sequence

<220>
 <223> 4-1(VLVH)xanti-CD3
 <400> 39

```

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
1           5           10           15
Val His Ser Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Ser Val
20           25           30
Ser Ala Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu
35           40           45
Leu Asn Ser Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys
50           55           60
Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu
65           70           75           80
Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe
85           90           95
Thr Leu Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr
100          105          110
Cys Gln Asn Asp Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys
115          120          125

```


Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly
 130 135 140 41
 Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val
 145 150 155 160
 Arg Pro Gly Thr Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala
 165 170 175
 Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro Gly His Gly
 180 185 190
 Leu Glu Trp Val Gly Asp Ile Phe Pro Gly Ser Gly Asn Ala His Tyr
 195 200 205
 Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser
 210 215 220
 Tyr Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala
 225 230 235 240
 Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Ala Met Asp Tyr
 245 250 255
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser
 260 265 270
 Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 275 280 285
 Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 290 295 300
 Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 305 310 315 320
 Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 325 330 335
 Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 340 345 350
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 355 360 365
 Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 370 375 380
 Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 385 390 395 400
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser

```

<220>
<223> 4-7(VL-VH)xanti CD3
<400> 41
atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacagggtg tactctccgc 60
cgcgagctcg tgatgaccca gactccaact tccttgctcg tcagtcttgg agatcaagcc 120
tccatctctt gcagatctag tcagagcctt gtacacagta atggaacac ctatttcat 180

```

43

```

tggtacctgc agaagccagg ccagtcctca aagctcctga tctacaaagt ttccaaccga 240
ttttctgggg tcccagacag gttcagtggc agtggatcag ggacagattt cacactcaag 300
atcagcagag tggaggctga ggatctggga gtttatttct gctctcaaag tacacatggt 360
ccgtacacgt tcggaggggg gaccaagctt gagatcaaag gtgggtgggg ttctggcggc 420
ggcggctccg gtgggtgggg ttctgaggtg cagctgctcg agcagctctg agctgagctg 480
gcgaggcctg gggcttcagt gaagctgtcc tgcaaggctt ctggctacac cttcacaac 540
tatggtttaa gctgggtgaa gcagaggcct ggacaggctc ttgagtggat tggagagggt 600
tatcctagaa ttggtaatgc ttactacaat gagaagttca agggcaaggc cacactgact 660
gcagacaaat cctccagcac agcgtccatg gagctccgca gcctgacctc tgaggactct 720
gcggtctatt tctgtgcaag acggggatcc tacgatacta actacgactg gtacttcgat 780
gtctggggcc aagggaccac ggtcaccgtc tcctccggag gtgggtggat cgatatcaaa 840
ctgcagcagt caggggctga actggcaaga cctggggcct cagtgaagat gtctgcaag 900
acttctggtt acacctttac taggtacacg atgcactggg taaaacagag gcctggacag 960
ggtctggaat ggattggata cattaatctt agcgtgggtt atactaatta caatcagaag 1020
ttcaaggaca aggccacatt gactacagac aaatcctcca gcacagccta catgcaactg 1080
agcagcctga catctgagga ctctgcagtc tattactgtg caagatatta tgatgatcat 1140
tactgccttg actactgggg ccaaggcacc actctcacag tctcctcagt cgaagggtga 1200
agtggagggt ctggtggaag tggaggttca ggtggagtcg acgacattca gctgaccag 1260
tctccagcaa tcatgtctgc atctccaggg gagaagggtc ccatgacctg cagagccagt 1320
tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaagatgg 1380
atttatgaca catccaaagt ggcttctgga gtcccttatt gcttcagtgg cagtgggtct 1440
gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 1500
tgccaacagt ggagtagtaa ccgctcacg ttcgggtgctg ggaccaagct ggagctgaaa 1560
catcatcacc atcatcatta g 1581

```

<210> 42

<211> 526

<212> PRT

<213> artificial sequence

<220>

<223> 4-7(VL-VH)xanti CD3

<400> 42

```

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
1           5           10           15

```

44

Val His Ser Ala Arg Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu
 20 25 30
 Pro Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln
 35 40 45
 Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln
 50 55 60
 Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg
 65 70 75 80
 Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp
 85 90 95
 Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr
 100 105 110
 Phe Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr
 115 120 125
 Lys Leu Glu Ile Lys Gly Gly Gly Gly Ser Gly Gly Gly Ser Gly
 130 135 140
 Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu
 145 150 155 160
 Ala Arg Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr
 165 170 175
 Thr Phe Thr Asn Tyr Gly Leu Ser Trp Val Lys Gln Arg Pro Gly Gln
 180 185 190
 Val Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn Ala Tyr
 195 200 205
 Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser
 210 215 220
 Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser
 225 230 235 240
 Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr Asp Thr Asn Tyr Asp
 245 250 255
 Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 260 265 270
 Gly Gly Gly Gly Ser Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu
 275 280 285

Ala Arg Pro Gly Ala Ser Val Lys Met Ser Cys⁴⁵ Lys Thr Ser Gly Tyr
 290 295 300
 Thr Phe Thr Arg Tyr Thr Met His Trp Val Lys Gln Arg Pro Gly Gln
 305 310 315 320
 Gly Leu Glu Trp Ile Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn
 325 330 335
 Tyr Asn Gln Lys Phe Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser
 340 345 350
 Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser
 355 360 365
 Ala Val Tyr Tyr Cys Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp
 370 375 380
 Tyr Trp Gly Gln Gly Thr Leu Thr Val Ser Ser Val Glu Gly Gly
 385 390 395 400
 Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile
 405 410 415
 Gln Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys
 420 425 430
 Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp
 435 440 445
 Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr
 450 455 460
 Ser Lys Val Ala Ser Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser
 465 470 475 480
 Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala
 485 490 495
 Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly
 500 505 510
 Ala Gly Thr Lys Leu Glu Leu Lys His His His His His His
 515 520 525

<210> 43

<211> 1566

<212> DNA

<213> artificial sequence

<220>

<223> 5-10(VLVH)xanti-CD3

<400> 43

atgggatgga gctgtatcat cctcttcttg gtagcaacag ctacaggtgt acactccgag	60
ctcgtgatga cacagtctcc atcctccctg actgtgacag caggagagaa ggtcactatg	120
agctgcaagt ccagtcagag tctgtttaa acgtggaatc aaaagaacta cttgacctgg	180
taccagcaga aaccaggga gcctcctaaa ctgttgatct actgggcatc cactagggaa	240
tctggggctc ctgactgctt cacaggcagt ggatctggaa cagatttcac tctcaccatc	300
agcagtgtgc aggtcgaaga cctggcagtt tattactgtc agaatgatta tagttatccg	360
ctcacgttcg gtgctgggac caagcttgag atcaaagggtg gtgggtggtc tggcggcggc	420
ggctccgggtg gtgggtggtc tgagggtcag ctgctcgagc agtctggagc tgagctggta	480
aggcctggga cttcagtga gatattctgc aaggcttctg gatacgctt cactaactac	540
tggctagggt gggtaagca gaggcctgga catggacttg agtgattgg agatattttc	600
cctggaagtg gtaatatcca ctacaatgag aagttcaagg gcaaagccac actgactgca	660
gacaaatctt cgagcacagc ctatatgcag ctcatagacc tgacatttga ggactctgct	720
gtctattttt gtgcaagact gaggaactgg gacgagccta tggactactg gggccaaggg	780
accacgggtc ccgtctcttc cggagggtgt ggatccgata tcaaactgca gcagtcaggg	840
gctgaactgg caagacctgg ggcctcagtg aagatgtcct gcaagacttc tggctacacc	900
tttactaggt acacgatgca ctgggtaaaa cagaggcctg gacagggtct ggaatggatt	960
ggatacatta atcctagccg tgggtatact aattacaatc agaagttcaa ggacaaggcc	1020
acattgacta cagacaaatc ctccagcaca gcctacatgc aactgagcag cctgacatct	1080
gaggactctg cagtctatta ctgtgaaga tattatgatg atcattactg ccttgactac	1140
tggggccaag gcaccactct cacagtctcc tcagtcgaag gtggaagtgg aggttctggt	1200
ggaagtggag gttcagggtg agtcgacgac attcagctga cccagtctcc agcaatcatg	1260
tctgcatctc caggggagaa ggtcaccatg acctgcagag ccagttcaag tgtaagttac	1320
atgaactggg accagcagaa gtcaggcacc tccccaaaa gatggattta tgacacatcc	1380
aaagtggctt ctggagtccc ttatcgcttc agtggcagtg ggtctgggac ctcatactct	1440
ctcacaatca gcagcatgga ggctgaagat gctgccattt attactgcca acagtggagt	1500
agtaaccgcg tcacgttcgg tgctgggacc aagctggagc tgaaacatca tcaccatcat	1560
cattag	1566

<210> 44

<211> 521

<212> PRT

<213> artificial sequence

<220>

<223> 5-10(VLVH)xanti-CD3

<400> 44

Met Gly Trp Ser Cys Ile Ile Leu Phe Leu Val Ala Thr Ala Thr Gly
 1 5 10 15
 Val His Ser Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Thr Val
 20 25 30
 Thr Ala Gly Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu
 35 40 45
 Leu Asn Ser Gly Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys
 50 55 60
 Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu
 65 70 75 80
 Ser Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe
 85 90 95
 Thr Leu Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr
 100 105 110
 Cys Gln Asn Asp Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys
 115 120 125
 Leu Glu Ile Lys Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly
 130 135 140
 Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val
 145 150 155 160
 Arg Pro Gly Thr Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala
 165 170 175
 Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro Gly His Gly
 180 185 190
 Leu Glu Trp Ile Gly Asp Ile Phe Pro Gly Ser Gly Asn Ile His Tyr
 195 200 205
 Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser
 210 215 220
 Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Phe Glu Asp Ser Ala
 225 230 235 240
 Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Pro Met Asp Tyr

245 250 48 255
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser
 260 265 270
 Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 275 280 285
 Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 290 295 300
 Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 305 310 315 320
 Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 325 330 335
 Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 340 345 350
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 355 360 365
 Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 370 375 380
 Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 385 390 395 400
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 405 410 415
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 420 425 430
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 435 440 445
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 450 455 460
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 465 470 475 480
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 485 490 495
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 500 505 510
 Glu Leu Lys His His His His His His
 515 520

<210> 45
 <211> 1494
 <212> DNA
 <213> artificial sequence

<220>

<223> CD3 VHVL aL x 3-1 VHVL

<400> 45
 gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcctgcaaga cttctggcta caccctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagtc 360
 gaaggtgaa gtggagggtc tgggtgaagt ggagggtcag gtggagtcga cgacattcag 420
 ctgacccagt ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc 480
 agagccagtt caagtgtaa ttacatgaac tggtagcagc agaagtcagg caccctcccc 540
 aaaagatgga ttatgacac atccaaagt gcttctggag tcccttatcg ctctcagtggtc 600
 agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc 660
 acttattact gccaacagt gagtagtaac ccgtcacgt tgggtgctgg gaccaagctg 720
 gagctgaaat ccggaggtgg tggatccgag gtgcagctgc tgcagcagtc tggagctgag 780
 ctgggtgaaac ctggggcctc agtgaagata tctgcaagg ctcttgata cgccttcact 840
 aactactggc taggttgggt aaagcagagg cctggacatg gacttgagt gattggagat 900
 cttttccctg gaagtggtaa tactactac aatgagaggt tcaggggcaa agccacactg 960
 actgcagaca aatcctcgag cacagccttt atgcagctca gtagcctgac atctgaggac 1020
 tctgtgtct atttctgtgc aagattgagg aactgggacg aggtatgga ctactggggc 1080
 caagggacca cggtcaccgt ctctcaggt ggtggtggtt ctggcggcgg cggctccggt 1140
 ggtggtggtt ctgagctcgt catgaccag tctccatctt atcttgctgc atctcctgga 1200
 gaaaccatta ctattaattg cagggaagt aagagcatta gcaaatattt agcctggtat 1260
 caagagaaac ctgggaaaac taataagctt cttatctact ctggatccac tttgcaatct 1320
 ggaattccat caaggttcag tggcagtga tctggtacag atttactct caccatcagt 1380
 agcctggagc ctgaagattt tgcaatgtat tactgtcaac agcataatga atatccgtac 1440
 acgttcggag gggggaccaa gcttgagatc aaacatcatc accatcatca ttag 1494

<210> 46

50

<211> 497

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL x 3-1 VHVL

<400> 46

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160

Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175

Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190

Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205

Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220

51

Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Lys Pro Gly Ala Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Ile Gly Asp Leu Phe Pro Gly
 290 295 300
 Ser Gly Asn Thr His Tyr Asn Glu Arg Phe Arg Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Phe Met Gln Leu Ser Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350
 Asp Glu Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 370 375 380
 Glu Leu Val Met Thr Gln Ser Pro Ser Tyr Leu Ala Ala Ser Pro Gly
 385 390 395 400
 Glu Thr Ile Thr Ile Asn Cys Arg Ala Ser Lys Ser Ile Ser Lys Tyr
 405 410 415
 Leu Ala Trp Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys Leu Leu Ile
 420 425 430
 Tyr Ser Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly
 435 440 445
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 450 455 460
 Glu Asp Phe Ala Met Tyr Tyr Cys Gln Gln His Asn Glu Tyr Pro Tyr
 465 470 475 480
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys His His His His His
 485 490 495

His

<210> 47

<211> 1494

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 3-1 VHVL

<400> 47

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggtta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctcagtc	360
gaagggtgaa gtggagggtc tgggtgaagt ggagggtcag gtggagtcga cgacattcag	420
ctgaccagtc ctccagcaat catgtctgca tctccagggg agaagggtcac catgacctgc	480
agagccagtt caagtgtaa ttacatgaac tggtagcagc agaagtcagg caccctcccc	540
aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg ctccagtggc	600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc	660
acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg	720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag	780
ctggtgaaac ctggggcctc agtgaagata tcctgcaagg cttctggata cgccttccact	840
aactactggc taggttgggt aaagcagagg cctggacatg gacttgagtg gattggagat	900
cttttccctg gaagtggtaa tactcactac aatgagaggt tcagggggcaa agccacactg	960
actgcagaca aatcctcagc cacagccttt atgcagctca gtagcctgac atctgaggac	1020
tctgctgtct atttctgtgc aagattgagg aactgggacg aggctatgga ctactggggc	1080
caagggacca cggtcaccgt ctctcaggt ggtggtggtt ctggcgccgg cggctccggt	1140
ggtggtggtt ctgagctcgt catgaccag tctccatctt atcttctgtc atctcctgga	1200
gaaaccatta ctattaattg cagggcaagt aagagcatta gcaaataatt agcctgggtat	1260
caagagaaac ctgggaaaac taataagctt cttatctact ctggatccac ttgcaatct	1320
ggaattccat caagggtcag tggcagtggg tctggtacag atttcaactc caccatcagt	1380
agcctggagc ctgaagattt tgcaatgtat tactgtcaac agcataatga atatccgtac	1440
acgttcggag gggggaccaa gcttgagatc aaacatcatc accatcatca ttag	1494

<210> 48

<211> 497

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 3-1 VHVL

<400> 48

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160

Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175

Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190

Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205

54

Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Lys Pro Gly Ala Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Ile Gly Asp Leu Phe Pro Gly
 290 295 300
 Ser Gly Asn Thr His Tyr Asn Glu Arg Phe Arg Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Phe Met Gln Leu Ser Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350
 Asp Glu Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 370 375 380
 Glu Leu Val Met Thr Gln Ser Pro Ser Tyr Leu Ala Ala Ser Pro Gly
 385 390 395 400
 Glu Thr Ile Thr Ile Asn Cys Arg Ala Ser Lys Ser Ile Ser Lys Tyr
 405 410 415
 Leu Ala Trp Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys Leu Leu Ile
 420 425 430
 Tyr Ser Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly
 435 440 445
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 450 455 460
 Glu Asp Phe Ala Met Tyr Tyr Cys Gln Gln His Asn Glu Tyr Pro Tyr
 465 470 475 480

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile⁵⁵ Lys His His His His His
 485 490 495

His

<210> 49

<211> 1521

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL AL x 3-5 VHVL

<400> 49
 gatatacaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggtta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagtc 360
 gaagggtgaa gtggagggtc tgggtgaagt ggagggtcag gtggagtcga cgacattcag 420
 ctgaccaggt ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc 480
 agagccagtt caagtgtaa ttacatgaac tggtagcagc agaagtcagg cacctcccc 540
 aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg ctctcagtggc 600
 agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgtgccc 660
 acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg 720
 gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag 780
 ctggttaaggc ctgggacttc agtgaagctg tcctgcaagg cttctggcta caccttcaca 840
 agctatggtt taagctgggt gaagcagaga actggacagg gccttgagtg gattggagag 900
 gtttatccca gaattggtaa tgcttactac aatgagaagt tcaagggcaa ggccacactg 960
 actgcagaca aatcctccag cacagcgtcc atggagctcc gcagcctgac atctgaggac 1020
 tctgcggctc atttctgtgc aagacgggga tcctacggtg gtaactacga ctggtacttc 1080
 gatgtctggg gccaaaggac cacggtcacc gtctcctcag gtgggtgggg ttctggcggc 1140
 ggcggctccg gtggtggtgg ttctgagctc gtgatgacct agactccact ctccctgcct 1200
 gtcagctctg gagatcaagc ctccatctct tcagatcta gtcagagcct tgtacacagt 1260
 aatggaaaca cctatttaca ttggtacctg cagaagccag gccagtctcc aaagctcctg 1320
 atctacaaag tttccaaccg attttctggg gtcccagaca gggtcagtggt cagtggatca 1380

56
 gggacagatt tcacactcaa gatcagcaga gtggaggctg aggatctggg agttttatttc 1440
 tgctctcaaa gtacacatgt tccgtacacg ttcggagggg ggaccaagct tgagatcaaa 1500
 catcatcacc atcatcatta g 1521

<210> 50

<211> 506

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL x 3-5 VHVL

<400> 50

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140

Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160

Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175

Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190

Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Leu Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Gly Leu Ser Trp Val Lys
 275 280 285
 Gln Arg Thr Gly Gln Gly Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg
 290 295 300
 Ile Gly Asn Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr
 340 345 350
 Gly Ser Asn Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr
 355 360 365
 Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser Gly
 370 375 380
 Gly Gly Gly Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro
 385 390 395 400
 Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser
 405 410 415
 Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys
 420 425 430
 Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe
 435 440 445
 Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 450 455 460

58

Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe
 465 470 475 480

Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys
 485 490 495

Leu Glu Ile Lys His His His His His His
 500 505

<210> 51

<211> 1521

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL AL Ser x 3-5 VHVL

<400> 51

```

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg    60
tcttgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg    120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac    180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac    240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat    300
gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctcagtc    360
gaagggtgaa gtggagggtc tgggtggaagt ggagggttcag gtggagtcga cgacattcag    420
ctgaccagat ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc    480
agagccagtt caagtgtaa ttacatgaac tgggtaccagc agaagtcagg cacctccccc    540
aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg cttcagtggc    600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc    660
acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg    720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag    780
ctggtaaggg ctgggacttc agtgaagctg tcttgcaagg cttctggcta caccttcaca    840
agctatggtt taagctgggt gaagcagaga actggacagg gccttgagtg gattggagag    900
gtttatccta gaattggtaa tgcttactac aatgagaagt tcaagggcaa ggccacactg    960
actgcagaca aatcctccag cacagcgtcc atggagctcc gcagcctgac atctgaggac   1020
tctgcggtct atttctgtgc aagacgggga tcttacggtg gtaactacga ctggtacttc   1080
gatgtctggg gccaaaggac cacggtcacc gtctcctcag gtggtggtgg ttctggcggc   1140
ggcggctccg gtggtggtgg ttctgagctc gtgatgaccc agactccact ctccctgcct   1200
gtcagtcctg gagatcaagc ctccatctct tgcagatcta gtcagagcct tgtacacagt   1260

```

59

aatggaaca cctatttaca ttggtacctg cagaagccag gccagtcctc aaagtcctc 1320
 atctacaaag tttccaaccg attttctggg gtcccagaca ggttcagtgg cagtggatca 1380
 gggacagatt tcacactcaa gatcagcaga gtggaggctg aggatctggg agtttatctc 1440
 tgctctcaaa gtacacatgt tccgtacacg ttcggagggg ggaccaagct tgagatcaaa 1500
 catcatcacc atcatcatta g 1521

<210> 52

<211> 506

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL at Ser x 3-5 VHVL

<400> 52

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15
 Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30
 Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser

165 170 60 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Leu Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr Gly Leu Ser Trp Val Lys
 275 280 285
 Gln Arg Thr Gly Gln Gly Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg
 290 295 300
 Ile Gly Asn Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr
 340 345 350
 Gly Ser Asn Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr
 355 360 365
 Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
 370 375 380
 Gly Gly Gly Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro
 385 390 395 400
 Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser
 405 410 415
 Leu Val His Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys
 420 425 430
 Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe
 435 440 445

61

Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe
 450 455 460

Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe
 465 470 475 480

Cys Ser Gln Ser Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys
 485 490 495

Leu Glu Ile Lys His His His His His His
 500 505

<210> 53

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 3-5 VHVL

<400> 53
 gatatacaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcttcgaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagggt 360
 ggtggtggtt ctggcgccgg cggctccggt ggtggtggtt ctgacattca gctgaccag 420
 tctccagcaa tcatgtctgc atctccaggg gagaaggta ccatgacctg cagagccagt 480
 tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctccc caaagatgg 540
 atttatgaca catccaaagt ggcttctgga gtcccttata gcttcagtgg cagtgggtct 600
 gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 660
 tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa 720
 tccggagggt gtggatccga ggtgcagctg ctcgagcagt ctggagctga gctggttaagg 780
 cctgggactt cagtgaagct gtcttgcaag gcttctggct acaccttcac aagctatggt 840
 ttaagctggg tgaagcagag aactggacag ggccttgagt ggattggaga ggtttatcct 900
 agaattggtg atgcttacta caatgagaag ttcaagggca aggccacact gactgcagac 960
 aaatcctcca gcacagcgtc catggagctc cgcagcctga catctgagga ctctgcggtc 1020
 tatttctgtg caagacgggg atcctacggt agtaactacg actggtactt cgaagtctgg 1080

62

```

ggccaaggga ccacggtcac cgtctcctca ggtgggtggtg gttctggcgg cggcggtccc 1140
ggtgggtggtg gttctgagct cgtgatgacc cagactccac tctccctgcc tgtcagtcct 1200
ggagatcaag cctccatctc ttgcagatct agtcagagcc ttgtacacag taatggaac 1260
acctatttac attggtacct gcagaagcca ggccagtctc caaagctcct gatctacaaa 1320
gtttccaacc gattttctgg ggtcccagac aggttcagtg gcagtggatc agggacagat 1380
ttcacactca agatcagcag agtggaggct gaggatctgg gagtttattt ctgctctcaa 1440
agtacacatg ttccgtacac gttcggaggg gggaccaagc ttgagatcaa acatcatcac 1500
catcatcatt ag 1512

```

<210> 54
 <211> 503
 <212> PRT
 <213> artificial sequence

<220>

<223> CD3 VHVL stL x 3-5 VHVL

<400> 54

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1      5      10      15
Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20     25     30
Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35     40     45
Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50     55     60
Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65     70     75     80
Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85     90     95
Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
100    105    110
Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
115    120    125
Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile
130    135    140

```

Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160

Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175

Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190

Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205

Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220

Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240

Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 245 250 255

Glu Leu Val Arg Pro Gly Thr Ser Val Lys Leu Ser Cys Lys Ala Ser
 260 265 270

Gly Tyr Thr Phe Thr Ser Tyr Gly Leu Ser Trp Val Lys Gln Arg Thr
 275 280 285

Gly Gln Gly Leu Glu Trp Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn
 290 295 300

Ala Tyr Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp
 305 310 315 320

Lys Ser Ser Ser Thr Ala Ser Met Glu Leu Arg Ser Leu Thr Ser Glu
 325 330 335

Asp Ser Ala Val Tyr Phe Cys Ala Arg Arg Gly Ser Tyr Gly Ser Asn
 340 345 350

Tyr Asp Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val
 355 360 365

Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 370 375 380

Ser Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu
 385 390 395 400

Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His
 405 410 415

Ser Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln

64

420 425 430

Ser Pro Lys₄₃₅ Leu Leu Ile Tyr Lys₄₄₀ Val Ser Asn Arg Phe₄₄₅ Ser Gly Val

Pro Asp Arg Phe Ser Gly Ser₄₅₅ Gly Ser Gly Thr Asp Phe₄₆₀ Thr Leu Lys

Ile Ser Arg Val Glu Ala₄₇₀ Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln₄₈₀
465

Ser Thr His Val₄₈₅ Pro Tyr Thr Phe Gly Gly₄₉₀ Gly Thr Lys Leu Glu Ile₄₉₅

Lys His His₅₀₀ His His His

<210> 55

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL AL X 4-1 VHVL

<400> 55
gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
tcttgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
cctggacagg gtctggaatg gattggatac attaatccta gccgtggtta tactaattac 180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagtc 360
gaagggtgaa gtggagggtc tgggtggaagt ggaggttcag gtggagtcga cgacattcag 420
ctgacccagt ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc 480
agagccagtt caagtgaag ttacatgaac tggtagcagc agaagtcagg cacctcccc 540
aaaagatgga ttatgacac atccaaagtg gcttctggag tcccttatcg ctctcagtcg 600
agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc 660
acttattact gccaacagtg gagtagtaac ccgctcacgt tgggtgctgg gaccaagctg 720
gagctgaaat ccggagggtg tggatccgag gtgcagctgc tcgagcagtc tggagctgag 780
ctggtgaagg ctgggacttc agtgaagata tcttgcaagg cttctggata cgccttcact 840
aactactggc taggttgggt taagcagagg cctggacatg gacttgaatg ggttggagat 900
attttccctg gaagtggtaa tgctcactac aatgagaagt tcaagggcaa agccacactg 960

65

actgcagaca agtcctcgta cacagcctat atgcagctca gtagcctgac atctgaggac 1020
 tctgctgtct atttctgtgc aagattgagg aactgggacg aggcctatgga ctactggggc 1080
 caagggacca cggtcaccgt ctctcaggt ggtggtggtt ctggcggcgg cggctccggt 1140
 ggtggtggtt ctgagctcgt gatgacacag tctccatcct ccctgagtggt gtcagcagga 1200
 gagaaggta ctatgagctg caagtccagt cagagtctgt taaacagtgg aaatcaaaag 1260
 aactacttgg cctggtacca gcagaaacca gggcagcctc ctactggtt gatctacggg 1320
 gcatccacta gggaatctgg ggtccctgat cgcttcacag gcagtggatc tggacacgat 1380
 ttcactctca ccatcagcag tgtgcaggct gaagacctgg cagtttatta ctgtcagaat 1440
 gattatagtt atccgtacac gttcggaggg gggaccaagc ttgagatcaa acatcatcac 1500
 catcatcatt ag 1512

<210> 56

<211> 503

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL AL X 4-1 VHVL

<400> 56

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125

66

Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Val Gly Asp Ile Phe Pro Gly
 290 295 300
 Ser Gly Asn Ala His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Tyr Thr Ala Tyr Met Gln Leu Ser Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350
 Asp Glu Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser
 370 375 380
 Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Ser Val Ser Ala Gly
 385 390 395 400

Glu Lys Val Thr Met Ser Cys Lys Ser Ser⁶⁷ Gln Ser Leu Leu Asn Ser
 405 410 415
 Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 420 425 430
 Pro Pro Lys Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu Ser Gly Val
 435 440 445
 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 450 455 460
 Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
 465 470 475 480
 Asp Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 485 490 495
 Lys His His His His His
 500

<210> 57

<211> 1512

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL AL Ser x 4-1 VHVL

<400> 57
 gatatacaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcctgcaaga cttctggcta cacctttact aggtacacga tgcacitgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctcagtc 360
 gaaggtggaa gtggagggtc tgggtggaagt ggaggttcag gtggagtcga cgacattcag 420
 ctgaccagtc ctccagcaat catgtctgca tctccagggg agaaggtcac catgacctgc 480
 agagccagtt caagtgaag ttacatgaac tgggtaccagc agaagtcagg cacctcccc 540
 aaaagatgga tttatgacac atccaaagtg gcttctggag tcccttatcg cttcagtggc 600
 agtgggtctg ggacctcata ctctctcaca atcagcagca tggaggctga agatgctgcc 660
 acttattact gccaacagtg gagtagtaac ccgctcacgt tcggtgctgg gaccaagctg 720
 gagctgaaat ccggaggtgg tggatccgag gtgcagctgc tcgagcagtc tggagctgag 780

68

```

ctggttaaggc ctgggacttc agtgaagata tcctgcaagg cttctggata cgccttcact 840
aactactggc taggttgggt taagcagagg cctggacatg gacttgaatg gggtggagat 900
attttccctg gaagtggtaa tgctcactac aatgagaagt tcaagggcaa agccacactg 960
actgcagaca agtcctcgta cacagcctat atgcagctca gtagcctgac atctgaggac 1020
tctgctgtct atttctgtgc aagattgcgg aactgggacg aggcctatgga ctactggggc 1080
caagggacca cggtcaccgt ctctcagggt ggtggtggtt ctggcggcgg cggctccggt 1140
ggtggtggtt ctgagctcgt gatgacacag tctccatcct cctgagtggt gtcagcagga 1200
gagaaggcca ctatgagctg caagtcaggt cagagctctgt taaacagtgg aaatcaaaag 1260
aactacttgg cctggtacca gcagaaacca gggcagcctc ctaaactggt gatctacggg 1320
gcaccacta ggaatctgg ggtccctgat cgcttcacag gcagtggatc tggaacagat 1380
ttcactctca ccatcagcag tgtgcaggct gaagacctgg cagtttatta ctgtcagaat 1440
gattatagtt atccgtacac gttcggaggg gggaccaagc ttgagatcaa acatcatcac 1500
catcatcatt ag 1512

```

<210> 58

<211> 503

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL aL Ser x 4-1 VHVL

<400> 58

```

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
1      5      10      15
Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
20     25     30
Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35     40     45
Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
50     55     60
Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
65     70     75     80
Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85     90     95
Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly
100    105    110

```

Thr Thr Leu Thr Val Ser Ser Val Glu Gly Gly Ser Gly Gly Ser Gly
 115 120 125
 Gly Ser Gly Gly Ser Gly Gly Val Asp Asp Ile Gln Leu Thr Gln Ser
 130 135 140
 Pro Ala Ile Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys
 145 150 155 160
 Arg Ala Ser Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser
 165 170 175
 Gly Thr Ser Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser
 180 185 190
 Gly Val Pro Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
 195 200 205
 Leu Thr Ile Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
 210 215 220
 Gln Gln Trp Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 225 230 235 240
 Glu Leu Lys Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln
 245 250 255
 Ser Gly Ala Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys
 260 265 270
 Lys Ala Ser Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys
 275 280 285
 Gln Arg Pro Gly His Gly Leu Glu Trp Val Gly Asp Ile Phe Pro Gly
 290 295 300
 Ser Gly Asn Ala His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu
 305 310 315 320
 Thr Ala Asp Lys Ser Ser Tyr Thr Ala Tyr Met Gln Leu Ser Ser Leu
 325 330 335
 Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp
 340 345 350
 Asp Glu Ala Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 355 360 365
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 370 375 380

70

Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Ser Val Ser Ala Gly
385 390 395 400

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
405 410 415

Gly Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
420 425 430

Pro Pro Lys Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu Ser Gly Val
435 440 445

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
450 455 460

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
465 470 475 480

Asp Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
485 490 495

Lys His His His His His His
500

<210> 59

<211> 1503

<212> DNA

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 4-1 VHVL

<400> 59
gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
tcctgcaaga cttctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctcagggt 360
ggtggtggtt ctggcggcgg cggctccggt ggtggtggtt ctgacattca gctgaccag 420
tctccagcaa tcattgtctgc atctccaggg gagaaggta ccatgacctg cagagccagt 480
tcaagtgtaa gttacatgaa ctggtaccag cagaagtcag gcacctcccc caaaagatgg 540
atttatgaca catccaaagt ggcttcggga gtcccttata gcttcagtgg cagtgggtct 600
gggacctcat actctctcac aatcagcagc atggaggctg aagatgctgc cacttattac 660

71

tgccaacagt ggagtagtaa cccgctcacg ttcggtgctg ggaccaagct ggagctgaaa 720
 tccggagggtg gtggatccga ggtgcagctg ctcgagcagt ctggagctga gctggtaagg 780
 cctgggacctt cagtgaagat atcctgcaag gcttctggat acgccttcac taactactgg 840
 ctaggttggg ttaagcagag gcctggacat ggacttgaat gggttggaga tattttccct 900
 ggaagtggta atgtcacta caatgagaag ttcaagggca aagccacact gactgcagac 960
 aagtccctgt acacagccta tatgcagctc agtagcctga catctgagga ctctgctgtc 1020
 tattttctgtg caagattgctg gaactgggac gaggctatgg actactgggg ccaaggggacc 1080
 acggtcaccg tctcctcagg tgggtggtgt tctggcgcg gcggctcgg tgggtggtgt 1140
 tctgagctcg tgatgacaca gtctccatcc tccctgagtg tgcagcagg agagaaggtc 1200
 actatgagct gcaagtccag tcagagtctg ttaaacagtg gaaatcaaaa gaactacttg 1260
 gcctgggtacc agcagaaacc agggcagcct cctaaactgt tgatctacgg ggcattccact 1320
 agggaatctg gggtccttga tcgcttcaca ggcagtggat ctggaacaga ttctactctc 1380
 accatcagca gtgtgcaggc tgaagacctg gcagtttatt actgtcagaa tgattatagt 1440
 tatccgtaca cgttcgagg ggggaccaag cttgagatca aacatcatca ccatcatcat 1500
 tag 1503

<210> 60

<211> 500

<212> PRT

<213> artificial sequence

<220>

<223> CD3 VHVL stL x 4-1 VHVL

<400> 60

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys

85 90 72 95
 Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Leu Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
 115 120 125
 Ser Gly Gly Gly Gly Ser Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile
 130 135 140
 Met Ser Ala Ser Pro Gly Glu Lys Val Thr Met Thr Cys Arg Ala Ser
 145 150 155 160
 Ser Ser Val Ser Tyr Met Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser
 165 170 175
 Pro Lys Arg Trp Ile Tyr Asp Thr Ser Lys Val Ala Ser Gly Val Pro
 180 185 190
 Tyr Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile
 195 200 205
 Ser Ser Met Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp
 210 215 220
 Ser Ser Asn Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 225 230 235 240
 Ser Gly Gly Gly Gly Ser Glu Val Gln Leu Leu Glu Gln Ser Gly Ala
 245 250 255
 Glu Leu Val Arg Pro Gly Thr Ser Val Lys Ile Ser Cys Lys Ala Ser
 260 265 270
 Gly Tyr Ala Phe Thr Asn Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro
 275 280 285
 Gly His Gly Leu Glu Trp Val Gly Asp Ile Phe Pro Gly Ser Gly Asn
 290 295 300
 Ala His Tyr Asn Glu Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp
 305 310 315 320
 Lys Ser Ser Tyr Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu
 325 330 335
 Asp Ser Ala Val Tyr Phe Cys Ala Arg Leu Arg Asn Trp Asp Glu Ala
 340 345 350
 Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly
 355 360 365

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Glu Leu Val
 370 375 380

Met Thr Gln Ser Pro Ser Ser Leu Ser Val Ser Ala Gly Glu Lys Val
 385 390 395 400

Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser Gly Asn Gln
 405 410 415

Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys
 420 425 430

Leu Leu Ile Tyr Gly Ala Ser Thr Arg Glu Ser Gly Val Pro Asp Arg
 435 440 445

Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 450 455 460

Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn Asp Tyr Ser
 465 470 475 480

Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys His His
 485 490 495

His His His His
 500

<210> 61

<211> 10

<212> PRT

<213> artificial sequence

<220>

<223> CDRH3 M1 mutant

<400> 61

His Tyr Asp Asp His Tyr Cys Leu Asp Tyr
 1 5 10

<210> 62

<211> 10

<212> PRT

<213> artificial sequence

<220>

74

<223> CDRH3 M4 mutant

<400> 62

Tyr Ser Asp Asp His Tyr Cys Leu Asp Tyr
1 5 10

<210> 63

<211> 10

<212> PRT

<213> artificial sequence

<220>

<223> CDRH3 M7 mutant

<400> 63

Tyr Tyr Asp Ala His Tyr Cys Leu Asp Tyr
1 5 10

<210> 64

<211> 10

<212> PRT

<213> artificial sequence

<220>

<223> CDRH3 M9 mutant

<400> 64

Tyr Tyr Asp Asp Gln Tyr Cys Leu Asp Tyr
1 5 10

<210> 65

<211> 10

<212> PRT

<213> artificial sequence

<220>

<223> CDRH3 M10 mutant

<400> 65

Tyr Tyr Asp Asp Pro Tyr Cys Leu Asp Tyr
1 5 10

<210> 66
<211> 10
<212> PRT
<213> artificial sequence

<220>
<223> CDRH3 M11 mutant
<400> 66
Tyr Phe Asn Asp His Tyr Cys Leu Asp Tyr
1 5 10

<210> 67
<211> 10
<212> PRT
<213> artificial sequence

<220>
<223> CDRH3 M13 mutant
<400> 67
Tyr Tyr Asn Asp Gln Tyr Cys Leu Asp Tyr
1 5 10

<210> 68
<211> 10
<212> PRT
<213> artificial sequence

<220>
<223> CDRH3 M20 mutant
<400> 68
Tyr His Asp Asp Pro Tyr Cys Leu Asp Tyr
1 5 10

<210> 69
<211> 10
<212> PRT
<213> artificial sequence

<220>

<223> CDRH3 M76 mutant

<400> 69

Tyr Tyr Asp Asp Asn Tyr Cys Leu Asp Tyr
 1 5 10

<210> 70

<211> 18

<212> PRT

<213> artificial sequence

<220>

<223> original linker

<400> 70

Val Glu Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly
 1 5 10 15

Val Asp

<210> 71

<211> 357

<212> DNA

<213> artificial sequence

<220>

<223> anti-CD3 VH

<400> 71
 gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg 60
 tcctgcaaga ctctggcta cacctttact aggtacacga tgcactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatccta gccgtgggta tactaattac 180
 aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac 240
 atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat 300
 gatgatcatt actgccttga ctactggggc caaggcacca ctctcacagt ctctca 357

<210> 72

<211> 119

<212> PRT

77

<213> artificial sequence

<220>

<223> anti-CD3VH

<400> 72

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr
 20 25 30

Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Tyr Tyr Asp Asp His Tyr Cys Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Leu Thr Val Ser Ser
 115

<210> 73

<211> 318

<212> DNA

<213> artificial sequence

<220>

<223> anti-CD3 VL

<400> 73

gacattcagc tgaccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc	60
atgacctgca gagccagttc aagtgttaagt tacatgaact ggtaccagca gaagtcaggc	120
acctccccca aaagatggat ttatgacaca tccaaagtgg cttctggagt cccttatcgc	180
ttcagtgcca gtgggtctgg gacctcatac tctctcaca tcagcagcat ggaggctgaa	240
gatgctgcca cttattactg ccaacagtgg agtagtaacc cgctcacgtt cgggtgctggg	300
accaagctgg agctgaaa	318

78

<210> 74
 <211> 106
 <212> PRT
 <213> artificial sequence

<220>
 <223> anti-CD3 VL
 <400> 74
 Asp Ile Gln Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15
 Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ser Tyr Met
 20 25 30
 Asn Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
 35 40 45
 Asp Thr Ser Lys Val Ala Ser Gly Val Pro Tyr Arg Phe Ser Gly Ser
 50 55 60
 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
 65 70 75 80
 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Leu Thr
 85 90 95
 Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

<210> 75
 <211> 10
 <212> PRT
 <213> artificial sequence

<220>
 <223> VH CDR1 anti-CD3
 <400> 75
 Gly Tyr Thr Phe Thr Arg Tyr Thr Met His
 1 5 10
 <210> 76
 <211> 357

<212> . DNA

<213> artificial sequence

<220>

<223> vH anti-CD3 cys->ser

<400> 76

gatatcaaac tgcagcagtc aggggctgaa ctggcaagac ctggggcctc agtgaagatg	60
tcctgcaaga ctcttggtta cacctttact aggtacacga tgcactgggt aaaacagagg	120
cctggacagg gtctggaatg gattggatac attaatccta gccgtgggtta tactaattac	180
aatcagaagt tcaaggacaa ggccacattg actacagaca aatcctccag cacagcctac	240
atgcaactga gcagcctgac atctgaggac tctgcagtct attactgtgc aagatattat	300
gatgatcatt actcccttga ctactggggc caaggcacca ctctcacagt ctctca	357

<210> 77

<211> 119

<212> PRT

<213> artificial sequence

<220>

<223> vH anti-CD3 cys->ser

<400> 77

Asp Ile Lys Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly Ala	1	5	10	15
Ser Val Lys Met Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Arg Tyr	20	25	30	
Thr Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile	35	40	45	
Gly Tyr Ile Asn Pro Ser Arg Gly Tyr Thr Asn Tyr Asn Gln Lys Phe	50	55	60	
Lys Asp Lys Ala Thr Leu Thr Thr Asp Lys Ser Ser Ser Thr Ala Tyr	65	70	75	80
Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys	85	90	95	
Ala Arg Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr Trp Gly Gln Gly	100	105	110	
Thr Thr Leu Thr Val Ser Ser				

80

115

<210> 78
 <211> 10
 <212> PRT
 <213> artificial sequence

<220>
 <223> VH CDR3 anti-CD3 cys->ser
 <400> 78

Tyr Tyr Asp Asp His Tyr Ser Leu Asp Tyr
 1 5 10

<210> 79
 <211> 360
 <212> DNA
 <213> artificial sequence

<220>
 <223> EpCAM 3-1 VH

<400> 79
 gaggtgcagc tgctcgagca gtctggagct gagctggtga aacctggggc ctcaagtgaag 60
 atatcctgca aggccttctgg atacgccttc actaactact ggctagggttg ggtaaagcag 120
 aggcctggac atggacttga gtggattgga gatcttttcc ctggaagtgg taataactcac 180
 tacaatgaga gggtcagggg caaagccaca ctgactgcag acaaatcctc gagcacagcc 240
 tttatgcagc tcagtagcct gacatctgag gactctgctg tctatttctg tgcaagattg 300
 aggaactggg acgaggctat ggactactgg ggccaaggga ccacgggtcac cgtctcctca 360

<210> 80
 <211> 120
 <212> PRT
 <213> artificial sequence

<220>
 <223> EpCAM 3-1 VH

<400> 80
 Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val Lys Pro Gly
 1 5 10 15

81

Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala Phe Thr Asn
20 25 30

Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp
35 40 45

Ile Gly Asp Leu Phe Pro Gly Ser Gly Asn Thr His Tyr Asn Glu Arg
50 55 60

Phe Arg Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala
65 70 75 80

Phe Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe
85 90 95

Cys Ala Arg Leu Arg Asn Trp Asp Glu Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 81

<211> 321

<212> DNA

<213> artificial sequence

<220>

<223> EpCAM 3-1 VL

<400> 81
gagctcgta tgaccagtc tccatcttat ctgctgcat ctctggaga aaccattact 60
attaattgca gggcaagtaa gagcattagc aaatatttag cctggatatca agagaaacct 120
gggaaaacta ataagcttct tatctactct ggatccactt tgcaatctgg aattccatca 180
agggtcagtg gcagtggatc tggtagatg ttactcttca ccatcagtag cctggagcct 240
gaagattttg caatgtatta ctgtcaacag cataatgaat atccgtacac gttcggaggg 300
gggaccaagc ttgagatcaa a 321

<210> 82

<211> 107

<212> PRT

<213> artificial sequence

<220>

82

<223> EpCAM 3-1 VL

<400> 82

Glu Leu Val Met Thr Gln Ser Pro Ser Tyr Leu Ala Ala Ser Pro Gly
 1 5 10 15

Glu Thr Ile Thr Ile Asn Cys Arg Ala Ser Lys Ser Ile Ser Lys Tyr
 20 25 30

Leu Ala Trp Tyr Gln Glu Lys Pro Gly Lys Thr Asn Lys Leu Leu Ile
 35 40 45

Tyr Ser Gly Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro
 65 70 75 80

Glu Asp Phe Ala Met Tyr Tyr Cys Gln Gln His Asn Glu Tyr Pro Tyr
 85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 83

<211> 372

<212> DNA

<213> artificial sequence

<220>

<223> EpCAM 3-5 VH

<400> 83

gaggtgcagc tgctcgagca gtctggagct gagctggtaa ggcctgggac ttcagtgaag 60
 ctgtcctgca aggtctctgg ctacaccttc acaagctatg gtttaagctg ggtgaagcag 120
 agaactggac agggccttga gtggattgga gaggtttatc ctagaattgg taatgcttac 180
 tacaatgaga agttcaaggg caaggccaca ctgactgcag acaaatcctc cagcacagcg 240
 tccatggagc tccgcagcct gacatctgag gactctgcgg tctatttctg tgcaagacgg 300
 ggatcctacg gtagtaacta cgactggtac ttcgatgtct ggggccaagg gaccacggtc 360
 accgtctcct ca 372

<210> 84

<211> 124

<212> PRT

83

<213> artificial sequence

<220>

<223> EpCAM 3-5 VH

<400> 84

Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val Arg Pro Gly
1 5 10 15

Thr Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser
20 25 30

Tyr Gly Leu Ser Trp Val Lys Gln Arg Thr Gly Gln Gly Leu Glu Trp
35 40 45

Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn Ala Tyr Tyr Asn Glu Lys
50 55 60

Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala
65 70 75 80

Ser Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe
85 90 95

Cys Ala Arg Arg Gly Ser Tyr Gly Ser Asn Tyr Asp Trp Tyr Phe Asp
100 105 110

Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 85

<211> 336

<212> DNA

<213> artificial sequence

<220>

<223> EpCAM 3-5 VL

<400> 85

gagctcgtga tgaccagac tccactctcc ctgcctgtca gtcttgagaga tcaagcctcc	60
atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg	120
tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt	180
tctgggtgcc cagacagggtt cagtggcagt ggatcaggga cagatttcac actcaagatc	240
agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttccg	300
tacacgttcg gaggggggac caagcttgag atcaaa	336

84

<210> 86
 <211> 112
 <212> PRT
 <213> artificial sequence

<220>

<223> EpCAM 3-5 VL

<400> 86

Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
 85 90 95

Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 87
 <211> 360
 <212> DNA
 <213> artificial sequence

<220>

<223> EpCAM 4-1 VH

<400> 87

gaggtgcagc tgctcgagca gtctggagct gagctggtaa ggcctgggac ttcagtgaag 60

atatcctgca aggccttctg atacgccttc actaactact ggctagggtg ggttaagcag 120

aggcctggac atggacttga atgggttgga gatattttcc ctggaagtgg taatgctcac 180

tacaatgaga agttcaaggg caaagccaca ctgactgcag acaagtcctc gtacacagcc 240

tatatgcagc tcagtagcct gacatctgag gactctgctg tctatttctg tgcaagattg 300

85

cggaactggg acgaggctat ggactactgg ggccaaggga ccacgggtcac cgtctcctca 360

<210> 88

<211> 120

<212> PRT

<213> artificial sequence

<220>

<223> EpCAM 4-1 VH

<400> 88

Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val Arg Pro Gly
1 5 10 15

Thr Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala Phe Thr Asn
20 25 30

Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp
35 40 45

Val Gly Asp Ile Phe Pro Gly Ser Gly Asn Ala His Tyr Asn Glu Lys
50 55 60

Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Tyr Thr Ala
65 70 75 80

Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe
85 90 95

Cys Ala Arg Leu Arg Asn Trp Asp Glu Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 89

<211> 339

<212> DNA

<213> artificial sequence

<220>

<223> EpCAM 4-1 VL

<400> 89

gagctcgtga tgacacagtc tccatcctcc ctgagtgtgt cagcaggaga gaaggctcact 60

86

atgagctgca agtcacgtca gagtctgtta aacagtggaa atcaaaagaa ctacttgccc	120
tgggtaccagc agaaaccagg gcagcctcct aaactgttga tctacggggc atccactagg	180
gaatctgggg tccctgatcg cttcacaggc agtggatctg gaacagattt cactctcacc	240
atcagcagtg tgcaggctga agacctggca gtttattact gtcagaatga ttatagttat	300
ccgtacacgt tcggaggggg gaccaagctt gagatcaaa	339

<210> 90

<211> 113

<212> PRT

<213> artificial sequence

<220>

<223> EpCAM 4-1 VL

<400> 90

Glu	Leu	Val	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Val	Ser	Ala	Gly
1			5					10					15		

Glu	Lys	Val	Thr	Met	Ser	Cys	Lys	Ser	Ser	Gln	Ser	Leu	Leu	Asn	Ser
		20					25						30		

Gly	Asn	Gln	Lys	Asn	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln
	35					40					45				

Pro	Pro	Lys	Leu	Leu	Ile	Tyr	Gly	Ala	Ser	Thr	Arg	Glu	Ser	Gly	Val
	50				55						60				

Pro	Asp	Arg	Phe	Thr	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr
65				70					75					80	

Ile	Ser	Ser	Val	Gln	Ala	Glu	Asp	Leu	Ala	Val	Tyr	Tyr	Cys	Gln	Asn
			85					90						95	

Asp	Tyr	Ser	Tyr	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile
		100						105					110		

Lys

<210> 91

<211> 372

<212> DNA

<213> artificial sequence

87

<220>

<223> EpCAM 4-7 VH

<400> 91
 gaggtgcagc tgctcgagca gtctggagct gagctggcga ggcctggggc ttcagtgaag 60
 ctgtcctgca aggccttctg ctacaccttc acaaaactatg gttaagctg ggtgaagcag 120
 aggcctggac aggtccttga gtggattgga gaggtttatc ctagaattgg taatgcttac 180
 tacaatgaga agttcaaggc caaggccaca ctgactgcag acaaatcctc cagcacagcg 240
 tccatggagc tccgcagcct gacctctgag gactctgcgg tctatttctg tgcaagacgg 300
 ggatcctacg atactaacta cgactggtac ttcgatgtct ggggccaagg gaccacggtc 360
 accgtctcct ca 372

<210> 92

<211> 124

<212> PRT

<213> artificial sequence

<220>

<223> EpCAM 4-7 VH

<400> 92
 Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Ala Arg Pro Gly
 1 5 10 15
 Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn
 20 25 30
 Tyr Gly Leu Ser Trp Val Lys Gln Arg Pro Gly Gln Val Leu Glu Trp
 35 40 45
 Ile Gly Glu Val Tyr Pro Arg Ile Gly Asn Ala Tyr Tyr Asn Glu Lys
 50 55 60
 Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala
 65 70 75 80
 Ser Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe
 85 90 95
 Cys Ala Arg Arg Gly Ser Tyr Asp Thr Asn Tyr Asp Trp Tyr Phe Asp
 100 105 110
 Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> 93

<211> 336
 <212> DNA
 <213> artificial sequence

<220>

<223> EPCAM 4-7 VL

<400> 93
 gagctcgtga tgaccagac tccactctcc ctgcctgtca gtcttgaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180
 tctggggccc cagacagggt cagtggcagt ggatcagga cagatttcac actcaagatc 240
 agcagagtgg aggtgagga tctgggagti tatttctgct ctcaaagtac acatgttccg 300
 tacacgttcg gaggggggac caagcttgag atcaaa 336

<210> 94
 <211> 112
 <212> PRT
 <213> artificial sequence

<220>

<223> EPCAM 4-7 VL

<400> 94
 Glu Leu Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
 85 90 95
 Thr His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 95
 <211> 360
 <212> DNA
 <213> artificial sequence

<220>

<223> EpCAM 5-10 VH

<400> 95
 gaggtgcagc tgctcgagca gtctggagct gagctggtaa ggcctgggac ttcagtgaag 60
 atatcctgca aggcctctgg atacgccttc actaactact ggctaggttg ggtaaagcag 120
 aggcctggac atggacttga gtggattgga gatattttcc ctggaagtgg taatatccac 180
 tacaatgaga agttcaaggc caaagccaca ctgactgcag acaaatcttc gagcacagcc 240
 tatatgcagc tcagtagcct gacatttgag gactctgctg tctattttctg tgcaagactg 300
 aggaactggg acgagcctat ggactactgg ggccaaggga ccacggtcac cgtctcctca 360

<210> 96
 <211> 120
 <212> PRT
 <213> artificial sequence

<220>

<223> EpCAM 5-10 VH

<400> 96

Glu Val Gln Leu Leu Glu Gln Ser Gly Ala Glu Leu Val Arg Pro Gly
 1 5 10 15
 Thr Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala Phe Thr Asn
 20 25 30
 Tyr Trp Leu Gly Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp
 35 40 45
 Ile Gly Asp Ile Phe Pro Gly Ser Gly Asn Ile His Tyr Asn Glu Lys
 50 55 60
 Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala
 65 70 75 80
 Tyr Met Gln Leu Ser Ser Leu Thr Phe Glu Asp Ser Ala Val Tyr Phe
 85 90 95

90

Cys Ala Arg Leu Arg Asn Trp Asp Glu Pro Met Asp Tyr Trp Gly Gln
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> 97

<211> 339

<212> DNA

<213> artificial sequence

<220>

<223> EpCAM 5-10 VL

<400> 97

gagctcgtga tgacacagtc tccatcctcc ctgactgtga cagcaggaga gaaggtcact 60
 atgagctgca agtccagtca gagtctgtta aacagtggaa atcaaaagaa ctacttgacc 120
 tggatcaccg agaaaccagg gcagcctcct aaactgttga tctactgggc atccactagg 180
 gaatctgggg tccctgatcg cttcacaggc agtggatctg gaacagattt cactctcacc 240
 atcagcagtg tgcaggctga agacctggca gtttattact gtcagaatga ttatagttat 300
 ccgctcacgt tcggtgctgg gaccaagctt gagatcaaa 339

<210> 98

<211> 113

<212> PRT

<213> artificial sequence

<220>

<223> EpCAM 5-10 VL

<400> 98

Glu Leu Val Met Thr Gln Ser Pro Ser Ser Leu Thr Val Thr Ala Gly
 1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Asn Ser
 20 25 30

Gly Asn Gln Lys Asn Tyr Leu Thr Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Asn
 85 90 95

Asp Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Ile
 100 105 110

Lys

<210> 99

<211> 15

<212> PRT

<213> artificial sequence

<220>

<223> standard linker

<400> 99

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
 1 5 10 15

<210> 100

<211> 47

<212> DNA

<213> artificial sequence

<220>

<223> 4-7 VL BspEI FOR

<400> 100
 ctgaaatccg gaggtggtgg atccgagctc gtgatgaccc agactcc

47

<210> 101

<211> 52

<212> DNA

<213> artificial sequence

<220>

<223> 4-7 VL GS15 REV5148

<400> 101