




(86) Date de dépôt PCT/PCT Filing Date: 2000/08/11
(87) Date publication PCT/PCT Publication Date: 2001/02/22
(45) Date de délivrance/Issue Date: 2011/01/25
(85) Entrée phase nationale/National Entry: 2002/02/12
(86) N° demande PCT/PCT Application No.: US 2000/022445
(87) N° publication PCT/PCT Publication No.: 2001/012621
(30) Priorités/Priorities: 1999/08/13 (US60/148,795);
1999/11/22 (US60/166,922); 2000/06/14 (US60/211,517)

(51) Cl.Int./Int.Cl. *C07D 413/04* (2006.01),
A61K 31/341 (2006.01), *A61K 31/4155* (2006.01),
A61K 31/4192 (2006.01), *A61K 31/42* (2006.01),
C07D 401/04 (2006.01), *C07D 403/04* (2006.01),
C07D 405/04 (2006.01), *C07D 413/14* (2006.01)

(72) Inventeurs/Inventors:
GREEN, JEREMY, US;
BEMIS, GUY, US;
GRILLOT, ANNE-LAURE, US;
LEDEBOER, MARK, US;
SALITURO, FRANCIS, US;
...

(73) Propriétaire/Owner:

(54) Titre : INHIBITEURS DE c-JUN N-TERMINAL KINASES (JNK) ET D'AUTRES PROTEINES-KINASES
(54) Title: INHIBITORS OF c-JUN N-TERMINAL KINASES (JNK) AND OTHER PROTEIN KINASES



(57) **Abrégé/Abstract:**

The present invention provides compounds of formula (I) where R¹ is H, CONH₂, T_(n)-R, or T_(n)-Ar², n may be zero or one, and G, XYZ, and Q are as described below. These compounds are inhibitors of protein kinase, particularly inhibitors of JNK, a mammalian protein kinase involved cell proliferation, cell death and response to extracellular stimuli. The invention also relates to methods for producing these inhibitors. The invention also provides pharmaceutical compositions comprising the inhibitors of the invention and methods of utilizing those compositions in the treatment and prevention of various disorders.

(72) **Inventeurs(suite)/Inventors(continued)**: HARRINGTON, EDMUND, US; GAO, HUI, US; BAKER, CHRISTOPHER, US; CAO, JINGRONG, US; HALE, MICHAEL, US

(73) **Propriétaires(suite)/Owners(continued)**: VERTEX PHARMACEUTICALS INCORPORATED, US

(74) **Agent**: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12621 A1(51) International Patent Classification⁷: **C07D 401/04**,
405/04, 403/04, 413/04, A61K 31/341, 31/4155, 31/4192,
31/42Lanes End, Natick, MA 07160 (US). **BAKER, Christo-
pher** [US/US]; 33 Curve Street, Bedford, MA 01730 (US).
CAO, Jingrong [CN/US]; 45 Madison Avenue, Newton,
MA 02460 (US). **HALE, Michael** [US/US]; 42 Sunset
Road, Bedford, MA 01730 (US).

(21) International Application Number: PCT/US00/22445

(22) International Filing Date: 11 August 2000 (11.08.2000)

(74) Agents: **SILVERMAN, Ian**; Vertex Pharmaceuticals Inc.,
130 Waverly Street, Cambridge, MA 02139-4242 et al.
(US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/148,795 12 August 1999 (12.08.1999) US
60/166,922 22 November 1999 (22.11.1999) US
60/211,517 14 June 2000 (14.06.2000) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(71) Applicant (*for all designated States except US*): **VER-
TEX PHARMACEUTICALS INCORPORATED**
[US/US]; Legal Dept., 130 Waverly Street, Cambridge,
MA 02139-4242 (US).


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GREEN, Jeremy**
[US/US]; 21 Greystone Court, Burlington, MA 01803
(US). **BEMIS, Guy** [US/US]; 256 Appleton Street,
Arlington, MA 02476 (US). **GRILLOT, Anne-Laure**
[FR/US]; 31 Regent Street #3R, Cambridge, MA 02140
(US). **LEDEBOER, Mark** [US/US]; 36 Faulkner Hill
Road, Acton, MA 01720 (US). **SALITURO, Francis**
[US/US]; 25 Baker Drive, Marlboro, MA 01752 (US).
HARRINGTON, Edmund [IE/US]; 460 E. 9th Street,
South Boston, MA 02127 (US). **GAO, Huai** [CN/US]; 26(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF c-JUN N-TERMINAL KINASES (JNK) AND OTHER PROTEIN KINASES



(I)

(57) **Abstract:** The present invention provides compounds of formula (I) where R¹ is H, CONH₂, T_(n)-R, or T_(n)-Ar², n may be zero or one, and G, XYZ, and Q are as described below. These compounds are inhibitors of protein kinase, particularly inhibitors of JNK, a mammalian protein kinase involved cell proliferation, cell death and response to extracellular stimuli. The invention also relates to methods for producing these inhibitors. The invention also provides pharmaceutical compositions comprising the inhibitors of the invention and methods of utilizing those compositions in the treatment and prevention of various disorders.



WO 01/12621 A1

79580-10

-1-

INHIBITORS OF c-JUN N-TERMINAL KINASES (JNK) AND
OTHER PROTEIN KINASES

TECHNICAL FIELD OF INVENTION

5 The present invention relates to
inhibitors of protein kinase, especially c-Jun N-
terminal kinases (JNK), which are members of the
mitogen-activated protein (MAP) kinase family. There
are a number of different genes and isoforms which
encode JNKs. Members of the JNK family regulate
signal transduction in response to environmental
stress and proinflammatory cytokines and have been
10 implicated to have a role in mediating a number of
different disorders. The invention also relates to
methods for producing these inhibitors. The
invention also provides pharmaceutical compositions
comprising the inhibitors of the invention and
15 methods of utilizing those compositions in the
treatment and prevention of various disorders.

-2-

BACKGROUND OF THE INVENTION

Mammalian cells respond to extracellular stimuli by activating signaling cascades that are mediated by members of the mitogen-activated protein (MAP) kinase family, which include the extracellular signal regulated kinases (ERKs), the p38 MAP kinases and the c-Jun N-terminal kinases (JNKs). MAP kinases (MAPKs) are activated by a variety of signals including growth factors, cytokines, UV radiation, and stress-inducing agents. MAPKs are serine/threonine kinases and their activation occur by dual phosphorylation of threonine and tyrosine at the Thr-X-Tyr segment in the activation loop. MAPKs phosphorylate various substrates including transcription factors, which in turn regulate the expression of specific sets of genes and thus mediate a specific response to the stimulus.

One particularly interesting kinase family are the c-Jun NH₂-terminal protein kinases, also known as JNKs. Three distinct genes, JNK1, JNK2, JNK3 have been identified and at least ten different splicing isoforms of JNKs exist in mammalian cells [Gupta et al., EMBO J., 15:2760-70 (1996)]. Members of the JNK family are activated by proinflammatory cytokines, such as tumor necrosis factor- α (TNF α) and interleukin-1 β (IL-1 β), as well as by environmental stress, including anisomycin, UV irradiation, hypoxia, and osmotic shock [Minden et al., Biochemica et Biophysica Acta, 1333:F85-F104 (1997)].

-3-

The down-stream substrates of JNKs include transcription factors c-Jun, ATF-2, Elk1, p53 and a cell death domain protein (DENN) [Zhang et al. Proc. Natl. Acad. Sci. USA, 95:2586-91 (1998)]. Each JNK

5 isoform binds to these substrates with different affinities, suggesting a regulation of signaling pathways by substrate specificity of different JNKs *in vivo* (Gupta et al., *supra*).

JNKs, along with other MAPKs, have been

10 implicated in having a role in mediating cellular response to cancer, thrombin-induced platelet aggregation, immunodeficiency disorders, autoimmune diseases, cell death, allergies, osteoporosis and heart disease. The therapeutic targets related to

15 activation of the JNK pathway include chronic myelogenous leukemia (CML), rheumatoid arthritis, asthma, osteoarthritis, ischemia, cancer and neurodegenerative diseases.

Several reports have detailed the

20 importance of JNK activation associated with liver disease or episodes of hepatic ischemia [Nat. Genet. 21:326-9 (1999); FEBS Lett. 420:201-4 (1997); J. Clin. Invest. 102:1942-50 (1998); Hepatology 28:1022-30 (1998)]. Therefore, inhibitors of JNK

25 may be useful to treat various hepatic disorders.

A role for JNK in cardiovascular disease such as myocardial infarction or congestive heart failure has also been reported as it has been shown JNK mediates hypertrophic responses to various forms

30 of cardiac stress [Circ. Res. 83:167-78 (1998); Circulation 97:1731-7 (1998); J. Biol. Chem. 272:28050-6 (1997); Circ. Res. 79:162-73 (1996);

-4-

Circ. Res. 78:947-53 (1996); J. Clin. Invest. 97:508-14 (1996)].

It has been demonstrated that the JNK cascade also plays a role in T-cell activation, including activation of the IL-2 promoter. Thus, inhibitors of JNK may have therapeutic value in altering pathologic immune responses [J. Immunol. 162:3176-87 (1999); Eur. J. Immunol. 28:3867-77 (1998); J. Exp. Med. 186:941-53 (1997); Eur. J. Immunol. 26:989-94 (1996)].

A role for JNK activation in various cancers has also been established, suggesting the potential use of JNK inhibitors in cancer. For example, constitutively activated JNK is associated with HTLV-1 mediated tumorigenesis [Oncogene 13:135-42 (1996)]. JNK may play a role in Kaposi's sarcoma (KS) because it is thought that the proliferative effects of bFGF and OSM on KS cells are mediated by their activation of the JNK signaling pathway [J. Clin. Invest. 99:1798-804 (1997)]. Other proliferative effects of other cytokines implicated in KS proliferation, such as vascular endothelial growth factor (VEGF), IL-6 and TNF α , may also be mediated by JNK. In addition, regulation of the c-jun gene in p210 BCR-ABL transformed cells corresponds with activity of JNK, suggesting a role for JNK inhibitors in the treatment for chronic myelogenous leukemia (CML) [Blood 92:2450-60 (1998)].

JNK1 and JNK2 are widely expressed in a variety of tissues. In contrast, JNK3, is selectively expressed in the brain and to a lesser

-5-

extent in the heart and testis [Gupta et al., *supra*;
Mohit et al., Neuron 14:67-78 (1995); Martin et al.,
Brain Res. Mol. Brain Res. 35:47-57 (1996)]. JNK3
has been linked to neuronal apoptosis induced by
5 kainic acid, indicating a role of JNK in the
pathogenesis of glutamate neurotoxicity. In the
adult human brain, JNK3 expression is localized to a
subpopulation of pyramidal neurons in the CA1, CA4
and subiculum regions of the hippocampus and layers
10 3 and 5 of the neocortex [Mohit et al., *supra*]. The
CA1 neurons of patients with acute hypoxia showed
strong nuclear JNK3-immunoreactivity compared to
minimal, diffuse cytoplasmic staining of the
hippocampal neurons from brain tissues of normal
15 patients [Zhang et al., *supra*]. Thus, JNK3 appears
to be involved involved in hypoxic and ischemic
damage of CA1 neurons in the hippocampus.

In addition, JNK3 co-localizes
immunochemically with neurons vulnerable in
20 Alzheimer's disease [Mohit et al., *supra*].
Disruption of the JNK3 gene caused resistance of
mice to the excitotoxic glutamate receptor agonist
kainic acid, including the effects on seizure
activity, AP-1 transcriptional activity and
25 apoptosis of hippocampal neurons, indicating that
the JNK3 signaling pathway is a critical component
in the pathogenesis of glutamate neurotoxicity (Yang
et al., Nature, 389:865-870 (1997)).

Based on these findings, JNK signalling,
30 especially that of JNK3, has been implicated in the
areas of apoptosis-driven neurodegenerative diseases
such as Alzheimer's Disease, Parkinson's Disease,

ALS (Amyotrophic Lateral Sclerosis), epilepsy and seizures, Huntington's Disease, traumatic brain injuries, as well as ischemic and hemorrhaging stroke.

5 There is a high unmet medical need to develop JNK specific inhibitors that are useful in treating the various conditions associated with JNK activation, especially considering the currently available, relatively inadequate treatment options
10 for the majority of these conditions.

 Recently, we have described crystallizable complexes of JNK protein and adenosine monophosphate, including complexes comprising JNK3, in WO 99/57253. Such information has been extremely
15 useful in identifying and designing potential inhibitors of various members of the JNK family, which, in turn, have the described above therapeutic utility.


 Much work has been done to identify and
20 develop drugs that inhibit MAPKs, such as p38 inhibitors. See, e.g., WO 98/27098 and WO 95/31451. However, to our knowledge, no MAPK inhibitors have been shown to be specifically selective for JNKs versus other related MAPKs.

25 Accordingly, there is still a great need to develop potent inhibitors of JNKs, including JNK3 inhibitors, that are useful in treating various conditions associated with JNK activation.

-7-

SUMMARY OF THE INVENTION

It has now been found that compounds of this invention and pharmaceutical compositions thereof are effective as inhibitors of c-Jun N-terminal kinases (JNK). These compounds have the
 5 general formula I:



where R^1 is H, CONH_2 , $T_{(n)}\text{-R}$, or $T_{(n)}\text{-Ar}^2$, n may be zero or one, and G, XYZ, and Q are as
 10 described below. Preferred compounds are those where the XYZ-containing ring is an isoxazole. Preferred G groups are optionally substituted phenyls and preferred Q are pyrimidine, pyridine or pyrazole rings.


15 These compounds and pharmaceutical compositions thereof are useful for treating or preventing a variety of disorders, such as heart disease, immunodeficiency disorders, inflammatory diseases, allergic diseases, autoimmune diseases,
 20 destructive bone disorders such as osteoporosis, proliferative disorders, infectious diseases and viral diseases. The compositions are also useful in methods for preventing cell death and hyperplasia and therefore may be used to treat or prevent
 25 reperfusion/ischemia in stroke, heart attacks, and organ hypoxia. The compositions are also useful in methods for preventing thrombin-induced platelet aggregation. The compositions are especially useful

79580-10


-8-

for disorders such as chronic myelogenous leukemia (CML),
 rheumatoid arthritis, asthma, osteoarthritis, ischemia,
 cancer, liver disease including hepatic ischemia, heart
 disease such as myocardial infarction and congestive heart
 5 failure, pathologic immune conditions involving T cell
 activation and neurodegenerative disorders.

According to one aspect of the present invention,
 there is provided a compound having the formula:



wherein: X-Y-Z is selected from one of the following:




R^1 is H, CONH_2 , $\text{T}_{(n)}\text{-R}$, or $\text{T}_{(n)}\text{-Ar}^2$; R is an aliphatic or
 15 substituted aliphatic group; n is zero or one; T is $\text{C}(=\text{O})$,
 CO_2 , CONH , $\text{S}(\text{O})_2$, $\text{S}(\text{O})_2\text{NH}$, COCH_2 or CH_2 ; each R^2 is
 independently selected from hydrogen, -R, $-\text{CH}_2\text{OR}$, $-\text{CH}_2\text{OH}$,
 $-\text{CH}=\text{O}$, $-\text{CH}_2\text{SR}$, $-\text{CH}_2\text{S}(\text{O})_2\text{R}$, $-\text{CH}_2(\text{C}=\text{O})\text{R}$, $-\text{CH}_2\text{CO}_2\text{R}$, $-\text{CH}_2\text{CO}_2\text{H}$,
 $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{NHR}$, $-\text{CH}_2\text{N}(\text{R})_2$, $-\text{CH}=\text{N}-\text{OR}$, $-\text{CH}=\text{NNHR}$, $-\text{CH}=\text{NN}(\text{R})_2$,
 20 $-\text{CH}=\text{NNHCOR}$, $-\text{CH}=\text{NNHCO}_2\text{R}$, $-\text{CH}=\text{NNHSO}_2\text{R}$, -aryl, -substituted
 aryl, $-\text{CH}_2(\text{aryl})$, $-\text{CH}_2(\text{substituted aryl})$, $-\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{NHCOR}$,
 $-\text{CH}_2\text{NHCONHR}$, $-\text{CH}_2\text{NHCON}(\text{R})_2$, $-\text{OCH}_2\text{NRCOR}$, $-\text{CH}_2\text{NHCO}_2\text{R}$, $-\text{CH}_2\text{CONHR}$,
 $-\text{CH}_2\text{CON}(\text{R})_2$, $-\text{CH}_2\text{SO}_2\text{NH}_2$, $-\text{CH}_2(\text{heterocyclyl})$, $-\text{CH}_2(\text{substituted}$
 heterocyclyl), $-(\text{heterocyclyl})$, or $-(\text{substituted}$
 25 heterocyclyl); each R^3 is independently selected from
 hydrogen, R, COR, CO_2R or $\text{S}(\text{O})_2\text{R}$; G is R or Ar^1 ; Ar^1 is aryl,
 substituted aryl, aralkyl, substituted aralkyl,
 heterocyclyl, or substituted heterocyclyl, wherein Ar^1 is
 optionally fused to a partially unsaturated or fully


79580-10

-8a-

unsaturated five to seven membered ring containing zero to three heteroatoms; Q-NH is



5 wherein the H of Q-NH is optionally replaced by R^3 ; A is N or CR^3 ; U is CR^3 , O, S, or NR^3 ; Ar^2 is aryl, substituted aryl, heterocyclyl or substituted heterocyclyl, wherein Ar^2 is optionally fused to a partially unsaturated or fully
 10 unsaturated five to seven membered ring containing zero to three heteroatoms; wherein each substitutable carbon atom in Ar^2 , including the fused ring when present, is optionally and independently substituted by halo, R, OR, SR, OH, NO_2 , CN, NH_2 , NHR, $N(R)_2$, $NHCOR$, $NHCONHR$, $NHCON(R)_2$, $NRCOR$, $NHCO_2R$, CO_2R , CO_2H , COR, CONHR, $CON(R)_2$, $S(O)_2R$, $SONH_2$, $S(O)R$, SO_2NHR ,
 15 or $NHS(O)_2R$, and wherein each saturated carbon in the fused ring is further optionally and independently substituted by $=O$, $=S$, $=NNHR$, $=NNR_2$, $=N-OR$, $=NNHCOR$, $=NNHCO_2R$, $=NNHSO_2R$ or $=NR$; and wherein each substitutable nitrogen atom in Ar^2 is optionally substituted by R, COR, $S(O)_2R$, or CO_2R ; provided
 20 that: (i) when compound of formula I is:




25

then R^1 is not phenyl, 3- OCH_3 -phenyl, 3- CH_3 -phenyl, 4- $N(CH_3)_2$ -phenyl, 4- $N(CH_2CH_3)_2$ -phenyl, 4-Et-phenyl, 3,5- $(CH_3)_2$ -phenyl, or 3- $N(CH_3)_2$ -phenyl; (ii) when a compound of formula I is:


79580-10

-8b-



5

then: when R^1 is allyl or phenyl, then R^2 is not D-arabinose; (iii) when a compound of formula I is:




10

then: G and R^1 are not simultaneously phenyl; or when G is methyl, then R^1 is not hydrogen, 4-chlorophenyl, 4-OCH₃-phenyl, or α -naphthyl.

15

According to another aspect of the present invention, there is provided a compound having the formula:




20 wherein A is N or CH; PG is hydrogen or a nitrogen protecting group; R^1 is H, $T_{(n)}-R$, or $T_{(n)}-Ar^2$; R is an aliphatic or substituted aliphatic group; n is zero or one; T is C(=O), CO₂, CONH, S(O)₂, S(O)₂NH, COCH₂ or CH₂; and each R^2 is independently selected from hydrogen, -R, -CH₂OR,
 25 -CH₂OH, -CH=O, -CH₂SR, -CH₂S(O)₂R, -CH₂(C=O)R, -CH₂CO₂R,


79580-10

-8c-

- CH₂CO₂H, -CH₂CN, -CH₂NHR, -CH₂N(R)₂, -CH=N-OR, -CH=NNHR,
 -CH=NN(R)₂, -CH=NNHCOR, -CH=NNHCO₂R, -CH=NNHSO₂R, -aryl,
 -substituted aryl, -CH₂(aryl), -CH₂(substituted aryl),
 -CH₂NH₂, -CH₂NHCOR, -CH₂NHCONHR, -CH₂NHCON(R)₂, -CH₂NRCOR,
 5 -CH₂NHCO₂R, -CH₂CONHR, -CH₂CON(R)₂, -CH₂SO₂NH₂,
 -CH₂(heterocyclyl), -CH₂(substituted heterocyclyl),
 -(heterocyclyl), or -(substituted heterocyclyl); provided
 that said compound is not



- 10 According to still another aspect of the present invention, there is provided a compound having the formula:




- 15 wherein: X-Y is N-O or O-N providing an isoxazole or reverse isoxazole ring; A is N or CH; G is R, aryl or substituted aryl; R is aliphatic or substituted aliphatic, R² is selected from hydrogen, -R, -CH₂OR, -CH₂OH, -CH=O, -CH₂SR, -CH₂S(O)₂R, -CH₂(C=O)R, -CH₂CO₂R, -CH₂CO₂H, -CH₂CN, -CH₂NHR,
 20 -CH₂N(R)₂, -CH=N-OR, -CH=NNHR, -CH=NN(R)₂, -CH=NNHCOR, -CH=NNHCO₂R, -CH=NNHSO₂R, -aryl, -substituted aryl, -CH₂(aryl), -CH₂(substituted aryl), -CH₂NH₂, -CH₂NHCOR, -CH₂NHCONHR, -CH₂NHCON(R)₂, -CH₂NRCOR, -CH₂NHCO₂R, -CH₂CONHR, -CH₂CON(R)₂, -CH₂SO₂NH₂, -CH₂(heterocyclyl), -CH₂(substituted

79580-10


-8d-

heterocyclyl), -(heterocyclyl), or -(substituted heterocyclyl); and R^1 is selected from halogen, NH_2 , SR , or SO_2R ; provided that: when A is CH , X is O , Y is N , and G is 4-fluorophenyl, then R^1 is not F , Br or Cl .

5 According to yet another aspect of the present invention, there is provided a use of a compound of formula I:




10 wherein: $X-Y-Z$ is selected from one of the following:




R^1 is H , $CONH_2$, $T_{(n)}-R$, or $T_{(n)}-Ar^2$; R is an aliphatic or substituted aliphatic group; n is zero or one; T is $C(=O)$, CO_2 , $CONH$, $S(O)_2$, $S(O)_2NH$, $COCH_2$ or CH_2 ; each R^2 is independently selected from hydrogen, $-R$, $-CH_2OR$, $-CH_2OH$, $-CH=O$, $-CH_2SR$, $-CH_2S(O)_2R$, $-CH_2(C=O)R$, $-CH_2CO_2R$, $-CH_2CO_2H$, $-CH_2CN$, $-CH_2NHR$, $-CH_2N(R)_2$, $-CH=N-OR$, $-CH=NNHR$, $-CH=NN(R)_2$, $-CH=NNHCOR$, $-CH=NNHCO_2R$, $-CH=NNHSO_2R$, $-aryl$, $-substituted$
 20 $aryl$, $-CH_2(aryl)$, $-CH_2(substituted\ aryl)$, $-CH_2NH_2$, $-CH_2NHCOR$, $-CH_2NHCONHR$, $-CH_2NHCON(R)_2$, $-CH_2NRCOR$, $-CH_2NHCO_2R$, $-CH_2CONHR$, $-CH_2CON(R)_2$, $-CH_2SO_2NH_2$, $-CH_2(heterocyclyl)$, $-CH_2(substituted\ heterocyclyl)$, $-(heterocyclyl)$, or $-(substituted\ heterocyclyl)$, each R^3 is independently selected from
 25 hydrogen, R , COR , CO_2R or $S(O)_2R$; G is R or Ar^1 ; Ar^1 is $aryl$, $substituted\ aryl$, $aralkyl$, $substituted\ aralkyl$, $heterocyclyl$, or $substituted\ heterocyclyl$, wherein Ar^1 is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring containing zero to
 30 three heteroatoms; $Q-NH$ is

79580-10



wherein the H of Q-NH is optionally replaced by R^3 ; A is N or CR^3 ; U is CR^3 , O, S, or NR^3 ; Ar^2 is aryl, substituted aryl, heterocyclyl or substituted heterocyclyl, wherein Ar^2 is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring containing zero to three heteroatoms; wherein each substitutable carbon atom in Ar^2 , including the fused ring when present, is optionally and independently substituted by halo, R, OR, SR, OH, NO_2 , CN, NH_2 , NHR, $N(R)_2$, NHCOR, NHCONHR, $NHCON(R)_2$, NRCOR, $NHCO_2R$, CO_2R , CO_2H , COR, CONHR, $CON(R)_2$, $S(O)_2R$, $SONH_2$, $S(O)R$, SO_2NHR , or $NHS(O)_2R$, and wherein each saturated carbon in the fused ring is further optionally and independently substituted by $=O$, $=S$, $=NNHR$, $=NNR_2$, $=N-OR$, $=NNHCOR$, $=NNHCO_2R$, $=NNHSO_2R$ or $=NR$; and wherein each substitutable nitrogen atom in Ar^2 is optionally substituted by R, COR, $S(O)_2R$, or CO_2R , for treating a disease state or condition in a mammal said disease state or condition characterized by being alleviated by treatment with a protein kinase inhibitor, and in the preparation of a medicament for treating such a disease state or condition.

According to one aspect of the present invention, there is provided a compound having the formula:




wherein:

79580-10

-8f-

X-Y-Z is selected from one of the following:



R^1 is H, CONH_2 , $\text{T}_{(n)}-\text{R}$, or $\text{T}_{(n)}-\text{Ar}^2$;

R is an optionally substituted C_{1-12} aliphatic;

5 n is zero or one;

T is $\text{C}(=\text{O})$, CO_2 , CONH , $\text{S}(\text{O})_2$, $\text{S}(\text{O})_2\text{NH}$, COCH_2 or CH_2 ;

each R^2 is independently selected from hydrogen, $-\text{R}$, $-\text{CH}_2\text{OR}$, $-\text{CH}_2\text{OH}$, $-\text{CH}=\text{O}$, $-\text{CH}_2\text{SR}$, $-\text{CH}_2\text{S}(\text{O})_2\text{R}$, $-\text{CH}_2(\text{C}=\text{O})\text{R}$, $-\text{CH}_2\text{CO}_2\text{R}$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{NHR}$, $-\text{CH}_2\text{N}(\text{R})_2$, $-\text{CH}=\text{N}-\text{OR}$, $-\text{CH}=\text{NNHR}$,
 10 $-\text{CH}=\text{NN}(\text{R})_2$, $-\text{CH}=\text{NNHCOR}$, $-\text{CH}=\text{NNHCO}_2\text{R}$, $-\text{CH}=\text{NNHSO}_2\text{R}$, optionally substituted 6 to 14-membered aryl, optionally substituted 6 to 14-membered $\text{CH}_2(\text{aryl})$, $-\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{NHCOR}$, $-\text{CH}_2\text{NHCONHR}$, $-\text{CH}_2\text{NHCON}(\text{R})_2$, $-\text{CH}_2\text{NRCOR}$, $-\text{CH}_2\text{NHCO}_2\text{R}$, $-\text{CH}_2\text{CONHR}$, $-\text{CH}_2\text{CON}(\text{R})_2$, $-\text{CH}_2\text{SO}_2\text{NH}_2$, optionally substituted 5 to 8-membered

15 $-\text{CH}_2(\text{heterocyclyl})$ comprising one or more heteroatoms selected from O, N and S, and optionally substituted 5 to 8-membered $-(\text{heterocyclyl})$ comprising one or more ring heteroatoms selected from O, N and S;

each R^3 is independently selected from hydrogen, R, COR, CO_2R
 20 and $\text{S}(\text{O})_2\text{R}$;


G is R or Ar^1 ;

Ar^1 is an optionally substituted 6 to 14-membered aryl or aralkyl, or an optionally substituted 5 to 8-membered heterocyclyl containing one or more ring heteroatoms
 25 selected from N, O and S, wherein Ar^1 is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring comprising zero to three heteroatoms selected from O, N and S;

79580-10

-8g-

Q-NH is



wherein the H of Q-NH is optionally replaced by R^3 ;

A is N or CR^3 ;

5 Ar^2 is an optionally mono- or poly-substituted 6 to 14-membered aryl or a 5 to 8-membered heterocyclyl comprising one or more heteroatoms selected from O, N and S, wherein Ar^2 optionally comprises an optionally mono- or poly-substituted, partially unsaturated or fully unsaturated
10 five to seven membered ring comprising zero to three heteroatoms selected from O, N and S fused to the aryl or the heterocyclyl;

wherein the one or more substituents on carbon atoms in Ar^2 are independently selected from halo, R, OR, SR, OH, NO_2 , CN, NH_2 , NHR, $N(R)_2$, NHCOR, NHCONHR, $NHCON(R)_2$, NRCOR, $NHCO_2R$, CO_2R , CO_2H , COR, CONHR, $CON(R)_2$, $S(O)_2R$, $SONH_2$, $S(O)R$, SO_2NHR , and $NHS(O)_2R$, and wherein the one or more substituents on saturated carbon atoms in the optional, fused ring in Ar^2 are further independently selected from =O, =S, =NNHR, =NNR₂,
20 =N-OR, =NNHCOR, =NNHCO₂R, =NNHSO₂R, and =NR; and

wherein the one or more substituents on nitrogen atoms in Ar^2 are independently selected from R, COR, $S(O)_2R$, and CO_2R .


DETAILED DESCRIPTION OF THE INVENTION

This invention provides novel compounds, and
25 pharmaceutically acceptable derivatives thereof, that are useful as JNK inhibitors. These compounds have the general

79580-10


-8h-

formula I:



wherein:

5 X-Y-Z is selected from one of the following:

R¹ is H, CONH₂, T_(n)-R, or T_(n)-Ar²;

R is an aliphatic or substituted aliphatic group;

10 n is zero or one;

T is C(=O), CO₂, CONH, S(O)₂, S(O)₂NH, COCH₂ or CH₂;each R² is independently selected from hydrogen, -R, -CH₂OR,-CH₂OH, -CH=O, -CH₂SR, -CH₂S(O)₂R, -CH₂(C=O)R, -CH₂CO₂R,-CH₂CO₂H, -CH₂CN, -CH₂NHR, -CH₂N(R)₂, -CH=N-OR, -CH=NNHR,15 -CH=NN(R)₂, -CH=NNHCOR, -CH=NNHCO₂R, -CH=NNHSO₂R, -aryl,

-9-


-substituted aryl, -CH₂(aryl), -CH₂(substituted aryl), -CH₂NH₂, -CH₂NHCOR, -CH₂NHCONHR, -CH₂NHCON(R)₂, -CH₂NRCOR, -CH₂NHCO₂R, -CH₂CONHR, -CH₂CON(R)₂, -CH₂SO₂NH₂, -CH₂(heterocyclyl),
 5 -CH₂(substituted heterocyclyl), -(heterocyclyl), or -(substituted heterocyclyl);

each R³ is independently selected from hydrogen, R, COR, CO₂R or S(O)₂R;

G is R or Ar¹;

10 Ar¹ is aryl, substituted aryl, aralkyl, substituted aralkyl, heterocyclyl, or substituted heterocyclyl, wherein Ar¹ is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring containing zero to three
 15 heteroatoms;

Q-NH is



wherein the H of Q-NH is optionally replaced by R³;
 A is N or CR³;

20 U is CR³, O, S, or NR³;

Ar² is aryl, substituted aryl, heterocyclyl or substituted heterocyclyl, wherein Ar² is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring
 25 containing zero to three heteroatoms; and

wherein each substitutable carbon atom in Ar², including the fused ring when present, is optionally and independently substituted by halo, R, OR, SR, OH, NO₂, CN, NH₂, NHR, N(R)₂, NHCOR, NHCONHR, NHCON(R)₂, NRCOR, NHCO₂R, CO₂R, CO₂H, COR,
 30 CONHR, CON(R)₂, S(O)₂R, SONH₂, S(O)R, SO₂NHR, or

-10-

NHS(O)₂R, and wherein each saturated carbon in the fused ring is further optionally and independently substituted by =O, =S, =NNHR, =NNR₂, =N-OR, =NNHCOR, =NNHCO₂R, =NNHSO₂R, or =NR;

5 wherein each substitutable nitrogen atom in Ar² is optionally substituted by R, COR, S(O)₂R, or CO₂R.

As used herein, the following definitions shall apply unless otherwise indicated. The term "aliphatic" as used herein means straight chained,
10 branched or cyclic C₁-C₁₂ hydrocarbons, preferably one to six carbons, which are completely saturated or which contain one or more units of unsaturation. For example, suitable aliphatic groups include substituted or unsubstituted linear, branched or
15 cyclic alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl. The term "alkyl" and "alkoxy" used alone or as part of a larger moiety refers to both straight and branched
20 chains containing one to twelve carbon atoms. The terms "alkenyl" and "alkynyl" used alone or as part of a larger moiety shall include both straight and branched chains containing two to twelve carbon atoms. The terms "haloalkyl", "haloalkenyl" and
25 "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I. The term "heteroatom" means N, O or S and shall include any oxidized form of nitrogen and sulfur,
30 and the quaternized form of any basic nitrogen.

The term "aryl", used alone or as part of a larger moiety as in "aralkyl", refers to aromatic

-11-

ring groups having five to fourteen members, such as phenyl, benzyl, 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl, and heterocyclic aromatic groups or heteroaryl groups such as 2-furanyl, 3-furanyl, N-
5 imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxadiazolyl, 5-oxadiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-
10 pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 3-pyridazinyl, 3-pyridazinyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 5-tetrazolyl, 2-triazolyl, 5-triazolyl, 2-thienyl, or 3-thienyl.

Aryl groups also include fused polycyclic
15 aromatic ring systems in which a carbocyclic aromatic ring or heteroaryl ring is fused to one or more other rings. Examples include tetrahydronaphthyl, benzimidazolyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, benzodiazepinyl,
20 benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinolinyl, isoindolyl, acridinyl, benzoisoxazolyl, and the like. Also included within the scope of the term "aryl", as it is used herein, is a group in which one or more carbocyclic aromatic
25 rings and/or heteroaryl rings are fused to a cycloalkyl or non-aromatic heterocyclyl, for example, indanyl or tetrahydrobenzopyranyl.

The term "heterocyclic ring" or "heterocyclyl" refers to a non-aromatic ring which
30 includes one or more heteroatoms such as nitrogen, oxygen or sulfur in the ring. The ring can be five, six, seven or eight-membered and/or fused to another

-12-

ring, such as a cycloalkyl or aromatic ring.
Examples include 3-1H-benzimidazol-2-one, 3-1-alkyl-
benzimidazol-2-one, 2-tetrahydrofuranyl, 3-
tetrahydrofuranyl, 2-tetrahydropyranyl, 3-
5 tetrahydropyranyl, 4-tetrahydropyranyl, 2-
tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-
morpholino, 3-morpholino, 4-morpholino, 2-
thiomorpholino, 3-thiomorpholino, 4-thiomorpholino,
1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-
10 piperazinyl, 2-piperazinyl, 1-piperidinyl, 2-
piperidinyl, 3-piperidinyl, 4-piperidinyl, 4-
thiazolidinyl, diazolonyl, N-substituted diazolonyl,
1-phthalimidinyl, benzoxane, benzotriazol-1-yl,
benzopyrrolidine, benzopiperidine, benzoxolane,
15 benzothiolane, and benzothiane.

A compound of this invention may contain a
ring that is fused to a partially saturated or fully
unsaturated five to seven membered ring containing
zero to three heteroatoms. Such a fused ring may be
20 an aromatic or non-aromatic monocyclic ring,
examples of which include the aryl and heterocyclic
rings described above.

An aryl group (carbocyclic and
heterocyclic) or an aralkyl group, such as benzyl or
25 phenethyl, may contain one or more substituents.
Examples of suitable substituents on the unsaturated
carbon atom of an aryl group include a halogen, -R,
-OR, -OH, -SH, -SR, protected OH (such as acyloxy),
phenyl (Ph), substituted Ph, -OPh, substituted
30 -OPh, -NO₂, -CN, -NH₂, -NHR, -N(R)₂, -NHCOR,
-NHCONHR, -NHCON(R)₂, -NRCOR, -NHCO₂R, -CO₂R, -CO₂H,
-COR, -CONHR, -CON(R)₂, -S(O)₂R, -SONH₂, -S(O)R,

-13-

-SO₂NHR, or -NHS(O)₂R, where R is an aliphatic group or a substituted aliphatic group.

An aliphatic group or a non-aromatic heterocyclic ring may contain one or more substituents. Examples of suitable substituents on the saturated carbon of an aliphatic group or of a non-aromatic heterocyclic ring include those listed above for the unsaturated carbon, such as in an aromatic ring, as well as the following: =O, =S, =NNHR, =NNR₂, =N-OR, =NNHCOR, =NNHCO₂R, =NNHSO₂R, or =NR.

A substitutable nitrogen on an aromatic or non-aromatic heterocyclic ring may be optionally substituted. Suitable substituents on the nitrogen include R, COR, S(O)₂R, and CO₂R, where R is an aliphatic group or a substituted aliphatic group.

Compounds derived by making isosteric or bioisosteric replacements of carboxylic acid or ester moieties of compounds described herein are within the scope of this invention. Isosteres, which result from the exchange of an atom or group of atoms to create a new compound with similar biological properties to the parent carboxylic acid or ester, are known in the art. The bioisosteric replacement may be physicochemically or topologically based. An example of an isosteric replacement for a carboxylic acid is CONHSO₂(alkyl) such as CONHSO₂Me.


It will be apparent to one skilled in the art that certain compounds of this invention may exist in tautomeric forms or hydrated forms, all such forms of the compounds being within the scope

-14-

of the invention. Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center.


5 Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds
10 which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -
15 enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

One embodiment of this invention relates to compounds of formula **I** where the XYZ-containing
20 ring is an isoxazole, as shown by the general formula **IA** below:



where R^2 is preferably alkyl, such as methyl, or CH_2 (heterocyclyl), such as CH_2 (N-morpholinyl); G is
25 preferably Ar^1 ; and R^1 is preferably $\text{T}_{(n)}\text{-Ar}^2$ or $\text{T}_{(n)}\text{-R}$, wherein n is most preferably zero. Most preferred are those compounds where G, R^1 , and R^2 are as just described, and Q-NH is an aminopyridine or aminopyrimidine where the NH is at the 2 position of
30 the ring:

-15-



or Q-NH is an amino pyrazole:









Table 1 below shows representative
 5 examples of **IA** compounds where Q is a pyrimidine,
 pyridine or pyrazole and R^1 is Ar^2 , represented by
 formula **IIA**.



10

IIA**Q1****Q2****Q3**

15 Table 1. Examples of Compounds of Formula **IIA**



No.	G	Q	R^2	R^3	R^4	R^5	R^6	R^7
IIA-1	Ph	Q1	Me	H	H	H	H	H
IIA-2	Ph	Q1	Me	H	H	OMe	H	H
IIA-3	Ph	Q1	Me	H	OMe	OMe	H	H
IIA-4	Ph	Q1	Me	Me	H	H	H	H





-16-


No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-5	Ph	Q1	Me	Me	H	CONH ₂	H	H
IIA-6	Ph	Q1	Me	Me	H	CN	H	H
IIA-7	Ph	Q1	Me	H	CN	H	H	H
IIA-8	Ph	Q1	Me	Me	F	H	H	H
IIA-9	Ph	Q1	Me	Me	H	F	H	H
IIA-10	Ph	Q1	Me	CF ₃	H	H	H	H
IIA-11	4-F-Ph	Q1	Me	H	H	H	H	H
IIA-12	2,3-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-13	2,4-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-14	2-Cl-Ph	Q1	Me	H	H	H	H	H
IIA-15	3,4-Cl ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-16	Ph	Q2	Et	H	CN	H	H	H
IIA-17	Ph	Q2	Et	H	CO ₂ H	H	H	H
IIA-18	Ph	Q2	Me	H	F	H	H	H
IIA-19	Ph	Q2	Me	H	H	F	H	H
IIA-20	Ph	Q2	Me	H	H	COMe	H	H
IIA-21	Ph	Q2	Me	H	H	COPh	H	H
IIA-22	Et	Q1	Me	H	H	H	H	H
IIA-23	PhCH ₂ OCH ₂ -	Q1	Me	H	H	H	H	H
IIA-24	Ph	Q2	Me	H	H	CONH ₂	H	H
IIA-25	3-F-Ph	Q1	Me	H	CN	H	H	H
IIA-26	3-F-Ph	Q1	Me	H	H	CN	H	H
IIA-27	3-F-Ph	Q1	Me	H	F	H	H	H
IIA-28	3-F-Ph	Q1	Me	H	H	F	H	H
IIA-29	3-F-Ph	Q1	Me	H	Me	CN	H	H
IIA-30	3-F-Ph	Q1	Me	H	F	CN	H	H
IIA-31	3-F-Ph	Q1	Me	H	H	SMe	H	H
IIA-32	Ph	Q1	Me	H	F	CN	H	H
IIA-33	Ph	Q1	Me	H	F	H	H	H
IIA-34	Ph	Q1	Me	H	H	CN	H	H
IIA-35	Ph	Q1	Me	H	H	COMe	H	H
IIA-36	Ph	Q1	Me	H	CH=C H	H	H	H
IIA-37	Ph	Q1	Me	H	SMe	H	H	H
IIA-38	Ph	Q1	Me	H	Me	CN	H	H

SUBSTITUTE SHEET (RULE 26)

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-39	Ph	Q1	Me	H	COMe	H	H	H
IIA-40	Ph	Q2	Et	H	H	H	H	H
IIA-41	Ph	Q1	Me	OMe	H	H	H	H
IIA-42	Ph	Q1	Me	H	H	F	H	H
IIA-43	Ph	Q2	Me	H	CO ₂ H	H	H	H
IIA-44	Ph	Q1	Me	H	H	Ph	H	H
IIA-45	Ph	Q1	Me	H	Me	H	Me	H
IIA-46	Ph	Q1	Me	H	H	SMe	H	H
IIA-47	Ph	Q2	Me	H	H	OMe	H	H
IIA-48	Ph	Q2	Me	H	OMe	H	H	H
	Ph	Q1	Me	OMe	H	H	CN	H
IIA-50	Ph	Q2	Me	H	CO ₂ Me	H	H	H
IIA-51	Ph	Q1	Me	F	H	H	CN	H
IIA-52	Ph	Q2	Me	H	H	H	H	H
IIA-53	Ph	Q2	Me	H	H	CO ₂ H	H	H
IIA-54	Ph	Q1	Me	Me	H	CN	H	H
IIA-55	2-F-Ph	Q1	Me	H	H	H	H	H
IIA-56	Ph	Q1	Me	F	H	F	H	H
IIA-57	Ph	Q1	Me	Me	H	CONH ₂	H	H
IIA-58	Ph	Q1	Me	Me	Cl	H	H	H
IIA-59	Ph	Q1	Me	F	H	H	H	H
IIA-60	2,6-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-61	Ph	Q1	Me	Me	H	OMe	H	H
IIA-62	Ph	Q1	Me	OMe	H	H	H	H
IIA-63	Ph	Q1	Me	H	H	SO ₂ Me	H	H
IIA-64	Ph	Q2	Me	H	H	CO ₂ Me	H	H
IIA-65	Ph	Q1	Me	NO ₂	H	H	H	H
IIA-66	3-F-Ph	Q1	Me	H	H	H	H	H
IIA-67	Ph	Q2	Me	H	CN	H	H	H
IIA-68	Ph	Q2	Me	H	H	CN	H	H
IIA-69	Ph	Q1	Me	CH:C H	H	H	H	H
IIA-70	Ph	Q1	Me	Me	F	H	H	H
IIA-71	Ph	Q1	Me	Cl	H	H	OMe	H
IIA-72	Ph	Q1	Me	H	Me	OMe	H	H

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-73	Ph	Q1	Me	OMe	H	H	OMe	H
IIA-74	2,5-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-75	2-Cl-6-F-Ph	Q1	Me	H	H	H	H	H
IIA-76	2-Cl-Ph	Q1	Me	H	H	H	H	H
IIA-77	3,4-Cl ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-78	Ph	Q1	Me	Me	H	F	H	H
IIA-79	2-Br-Ph	Q1	Me	H	H	H	H	H
IIA-80	2,3-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-81	Ph	Q1	Me	SMe	H	H	H	H
IIA-82	3-CF ₃ -Ph	Q1	Me	H	H	H	H	H
IIA-83	3,5-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-84	2,6-Cl ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-85	2,3-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-86	Me	Q1	Me	H	H	H	H	H
IIA-87	cyclopropyl	Q1	Me	H	H	H	H	H
IIA-88	cyclohexyl	Q1	Me	H	H	H	H	H
IIA-89	2,4-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-90	t-butyl	Q1	Me	H	H	H	H	H
IIA-91	2,6-F ₂ -Ph	Q1	Me	H	H	COMe	H	H
IIA-92	2,6-F ₂ -Ph	Q1	Me	H	CN	H	H	H
IIA-93	2,6-F ₂ -Ph	Q1	Me	H	H	CN	H	H
IIA-94	2,6-F ₂ -Ph	Q1	Me	H	F	H	H	H
IIA-95	2,6-F ₂ -Ph	Q1	Me	H	H	F	H	H
IIA-96	2,6-F ₂ -Ph	Q1	Me	H	CN	F	H	H
IIA-97	2,6-F ₂ -Ph	Q1	Me	H	H	SMe	H	H
IIA-98	Ph	Q2	Me	H	H	NMe ₂	H	H
IIA-99	Ph	Q2	Me	H	NO ₂	H	H	H
IIA-100	Ph	Q2	Me	H	NHAc	H	H	H
IIA-101	Ph	Q2	Me	H	NH ₂	H	H	H
IIA-102	Ph	Q1	Me	H	Me	H	H	H
IIA-103	Ph	Q1	Me	H	H	Me	H	H
IIA-104	2-Me-Ph	Q1	Me	H	H	H	H	H
IIA-105	2-Me-Ph	Q1	Me	H	F	CN	H	H
IIA-106	2-Me-Ph	Q1	Me	H	F	H	H	H
IIA-107	2-Me-Ph	Q1	Me	H	H	CN	H	H


No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-108	2-Me-Ph	Q1	Me	H	Me	H	H	H
IIA-109	2-Me-Ph	Q1	Me	H	CN	H	H	H
IIA-110	2-CF ₃ -Ph	Q1	Me	H	F	CN	H	H
IIA-111	2-CF ₃ -Ph	Q1	Me	H	CN	H	H	H
IIA-112	2-CF ₃ -Ph	Q1	Me	H	H	H	H	H
IIA-113	3,4-(OCH ₂ O)-Ph	Q1	Me	H	F	CN	H	H
IIA-114	3,4-(OCH ₂ O)-Ph	Q1	Me	H	CN	H	H	H
IIA-115	3,4-(OCH ₂ O)-Ph	Q1	Me	H	H	H	H	H
IIA-116	 3,4-(OCH ₂ O)	Q1	Me	 bis-N,N'-4-cyanophenyl				
IIA-117	3-OBn-Ph	Q1	Me	H	F	CN	H	H
IIA-118	3-OBn-Ph	Q1	Me	H	CN	H	H	H
IIA-119	3-OBn-Ph	Q1	Me	H	H	H	H	H
IIA-120	3-NO ₂ -Ph	Q1	Me	H	F	CN	H	H
IIA-121	3-NO ₂ -Ph	Q1	Me	bis-N,N'-4-cyanophenyl				
IIA-122	3-NO ₂ -Ph	Q1	Me	H	CN	H	H	H
IIA-123	3-NO ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-124	3-CN-Ph	Q1	Me	H	F	CN	H	H
IIA-125	3-CN-Ph	Q1	Me	H	H	CN	H	H
IIA-126	3-CN-Ph	Q1	Me	H	CN	H	H	H
IIA-127	3-CN-Ph	Q1	Me	H	H	H	H	H
IIA-128	3-NO ₂ -Ph	Q1	Me	H	H	CO ₂ Et	H	H
IIA-129	3-CN-Ph	Q1	Me	H	CO ₂ Me	H	H	H
IIA-130	Ph	Q1	Me	H	CO ₂ Et	H	H	H
IIA-131	Ph	Q1	Me	N	H	NO ₂	H	H
IIA-132	Ph	Q2	Me					
IIA-133	Ph	Q2	Me					
IIA-134	Ph	Q2	Me	H	CH ₂	H	H	H

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
					OH			
IIA-135	Ph	Q2	Me					
IIA-136	Ph	Q3	Me	H	CN	H	H	H
IIA-137	Ph	Q3	Me	H	H	CN	H	H
IIA-138	Ph	Q3	Me	H	COM _e	H	H	H


For compounds of Formula **IIA** where R¹ is phenyl, preferred phenyl substituents are selected from hydrogen and one or more halo, aliphatic, substituted aliphatic (preferably haloalkyl), alkoxy, CN, CO₂H, CO₂(alkyl), S(alkyl), CONH₂, CO(alkyl), SO₂(alkyl), CO(phenyl), or NO₂. Preferred G groups are phenyl rings optionally substituted with one or more groups independently selected from alkyl, alkoxy or halogen.
















Examples of compounds of Formula **IIA** where R¹ is other than phenyl are shown below in Table 2.

Table 2. Examples of Compounds of Formula **IIA** (R¹ is other than phenyl)




























IIA (R¹ is other than phenyl)

No.	G	A	R ¹
IIAA-1	Ph	CH	

No.	G	A	R ¹
IIAA-2	Ph	CH	
IIAA-3	Ph	N	
IIAA-4	Ph	N	
IIAA-5	Ph	N	
IIAA-6	Ph	N	
IIAA-7	Ph	N	
IIAA-8	Ph	N	
IIAA-9	3-F-Ph	N	
IIAA-10	Ph	N	
IIAA-11	Ph	N	
IIAA-12	Ph	N	
IIAA-13	Ph	N	
IIAA-14	2,6-F ₂ -Ph	N	
IIAA-15	Ph	N	
IIAA-16	Ph	N	

-22-

No.	G	A	R ¹
IIAA-17	Ph	N	
IIAA-18	Ph	N	
IIAA-19	2-Me-Ph	N	
IIAA-20	2-Me-Ph	N	
IIAA-21		N	
IIAA-22	3-NO ₂ -Ph	N	
IIAA-23	3-CN-Ph	N	
IIAA-24	Ph	N	
IIAA-25	Ph	N	
IIAA-26	Ph	N	
IIAA-27	Ph	N	
IIAA-28	Ph	N	
IIAA-29	Ph	N	
IIAA-30	Ph	N	

No.	G	A	R ¹
IIAA-31	Ph	N	
IIAA-32	Ph	N	
IIAA-33	Ph	N	
IIAA-34	Ph	N	
IIAA-35	Ph	N	
IIAA-36	Ph	N	
IIAA-37	Ph	N	
IIAA-38	Ph	N	
IIAA-39	Ph	CH	
IIAA-40	Ph	CH	

Preferred **IIA** compounds are those where Ar¹ is an unsubstituted phenyl or a phenyl substituted with one or more halo, alkyl or alkoxy. More preferred


5 **IIA** compounds are those where Ar¹ is as just described, and Ar² is a naphthyl or phenyl optionally substituted with one or more halo, alkyl, alkoxy, haloalkyl, carboxyl, alkoxycarbonyl, cyano, or CONH₂, or an indanone (as in compound IIAA-11). Also

10 preferred are **IIA** compounds where R¹ is an optionally substituted alkyl or optionally substituted cycloalkyl, more preferably alkoxyalkyl,


-24-

alkoxycarbonylalkyl, hydroxyalkyl, pyridinylalkyl, alkoxyalkyl, alkoxyalkyl, alkoxyalkyl, or hydroxyalkyl. Examples of these preferred compounds include IIAA-24, IIAA-33 through IIAA-36, 5 IIAA-38 and IIAA-40.

One embodiment of this invention relates to compounds of formula **IA** where Q is a pyrimidine ring and R¹ is T-Ar² where T is selected from CO, CO₂, CONH, S(O)₂, S(O)₂NH, COCH₂ and CH₂. When R¹ is 10 T-Ar², preferred compounds are those where T is C(=O), represented by formula **IIIA**. Table 3 below shows representative examples of **IIIA** compounds.



15

Table 3. Examples of **IIIA** Compounds

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-1	phenyl	Q1	H	H	H	H	H
IIIA-2	phenyl	Q1	Br	H	H	H	H
IIIA-3	phenyl	Q1	F	H	H	H	H
IIIA-4	phenyl	Q1	Cl	H	H	H	H
IIIA-5	phenyl	Q1	CH ₃	H	H	H	H
IIIA-6	phenyl	Q1	H	CH ₃	H	H	H
IIIA-7	phenyl	Q1	H	H	OCH ₃	H	H
IIIA-8	phenyl	Q1	H	OCH ₃	OCH ₃	H	H

-25-

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-9	phenyl	Q1	OC H ₃	H	OCH ₃	H	H
IIIA-10	phenyl	Q1	OC H ₃	H	H	H	OCH 3
IIIA-11	phenyl	Q1	H	H	CN	H	H
IIIA-12	5-fluorophenyl	Q1	H	H	OCH ₃	H	H
IIIA-13	phenyl	Q1	H	OCH ₃	OCH ₃	OC H ₃	H
IIIA-14	phenyl	Q1	H	H	F	H	H
IIIA-15	phenyl	Q1	Ar ² is 2-thienyl				
IIIA-16	phenyl	Q1	Ar ² is 1-oxo-indan-5-yl				
IIIA-17	phenyl	Q1	Ar ² is 4-pyridyl				
IIIA-18	2-CH ₃ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-19	2-CH ₃ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-20	2-CH ₃ -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-21	2-CH ₃ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-22	2-CF ₃ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-23	2-CF ₃ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-24	2-CF ₃ -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-25	2-CF ₃ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-26	benzo[3,5]dioxole	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-27	benzo[3,5]dioxole	Q1	H	OCH ₃	H	H	H
IIIA-28	benzo[3,5]dioxole	Q1	H	H	OCH ₃	H	H
IIIA-29	benzo[3,5]dioxole	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-30	3-benzyloxy- phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-31	3-benzyloxy- phenyl	Q1	H	OCH ₃	H	H	H
IIIA-32	3-benzyloxy -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-33	3-benzyloxy-phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-34	3-nitrophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-35	3-nitrophenyl	Q1	H	OCH ₃	H	H	H
IIIA-36	3-nitrophenyl	Q1	H	H	OCH ₃	H	H
IIIA-37	3-nitrophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-38	3-cyanophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-39	3-cyanophenyl	Q1	H	OCH ₃	H	H	H
IIIA-40	3-cyanophenyl	Q1	H	H	OCH ₃	H	H
IIIA-41	3-cyanophenyl	Q1	H	OCH ₃	H	OCH ₃	H

-26-

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-42	phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-43	phenyl	Q1	H	CN	H	H	H
IIIA-44	phenyl	Q1	H	H	CO ₂ Me	H	H
IIIA-45	3-fluorophenyl	Q1	H	Cl	H	H	H
IIIA-46	3-fluorophenyl	Q1	H	OCH ₃	H	H	H
IIIA-47	3-fluorophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-48	3-fluorophenyl	Q1	H	Me	H	H	H
IIIA-49	3-fluorophenyl	Q1	H	H	F	H	H
IIIA-50	3-fluorophenyl	Q1	H	H	Me	H	H
IIIA-51	3-fluorophenyl	Q1	H	CN	H	H	H
IIIA-52	3-fluorophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-53	3-fluorophenyl	Q1	Ar ² is 2-naphthyl				
IIIA-54	2-fluorophenyl	Q1	H	Cl	H	H	H
IIIA-55	2-fluorophenyl	Q1	H	OCH ₃	H	H	H
IIIA-56	2-fluorophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-57	2-fluorophenyl	Q1	H	Me	H	H	H
IIIA-58	2-fluorophenyl	Q1	H	H	OCH ₃	H	H
IIIA-59	2-fluorophenyl	Q1	H	H	F	H	H
IIIA-60	2-fluorophenyl	Q1	H	H	Me	H	H
IIIA-61	2-fluorophenyl	Q1	H	CN	H	H	H
IIIA-62	2-fluorophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-63	2-fluorophenyl	Q1	Ar ² is 2-naphthyl				
IIIA-64	2,6-F ₂ -phenyl	Q1	H	Cl	H	H	H
IIIA-65	2,6-F ₂ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-66	2,6-F ₂ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-67	2,6-F ₂ -phenyl	Q1	H	Me	H	H	H
IIIA-68	2,6-F ₂ -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-69	2,6-F ₂ -phenyl	Q1	H	H	F	H	H
IIIA-70	2,6-F ₂ -phenyl	Q1	H	H	Me	H	H
IIIA-71	2,6-F ₂ -phenyl	Q1	H	CN	H	H	H
IIIA-72	2,6-F ₂ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-73	2,6-F ₂ -phenyl	Q1	Ar ² is 2-naphthyl				
IIIA-74	phenyl	Q1	H	NO ₂	H	H	H
IIIA-75	phenyl	Q1	H	NHAc	H	H	H

SUBSTITUTE SHEET (RULE 26)







-27-

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-76	phenyl	Q1	H	COMe	H	H	H
IIIA-77	phenyl	Q2	H	COMe	H	H	H
IIIA-78	phenyl	Q2	H	CN	H	H	H
IIIA-79	phenyl	Q3	H	H	H	H	H
IIIA-80	phenyl	Q3	H	OCH ₃	H	H	H
IIIA-81	phenyl	Q3	H	H	OCH ₃	H	H
IIIA-82	phenyl	Q3	H	CN	H	H	H
IIIA-83	phenyl	Q3	H	OCH ₃	H	OCH ₃	H
IIIA-84	phenyl	Q3	H	H	F	H	H
IIIA-85	phenyl	Q3	H	COMe	H	H	H
IIIA-86	phenyl	Q3	H	H	COMe	H	H
IIIA-87	phenyl	Q3	OC H ₃	H	H	H	H
IIIA-88	phenyl	Q3	2-thienyl				
IIIA-89	phenyl	Q3	2-furanyl				
IIIA-90	3-OMe-phenyl	Q3	H	OCH ₃	H	H	H
IIIA-91	Cyclohexyl	Q3	H	OCH ₃	H	H	H
IIIA-92	4-Cl-phenyl	Q3	H	OCH ₃	H	H	H
IIIA-93	3-Cl-phenyl	Q3	H	OCH ₃	H	H	H
IIIA-94	4-F-phenyl	Q3	H	OCH ₃	H	H	H
IIIA-95	3-F-phenyl	Q3	H	OCH ₃	H	H	H
IIIA-96	4-pyridyl	Q3	H	OCH ₃	H	H	H
IIIA-97	3-pyridyl	Q3	H	OCH ₃	H	H	H

Preferred **IIIA** compounds are those compounds where Ar¹ is an unsubstituted phenyl or a phenyl substituted with one or more substituents independently selected from halogen. More preferred **IIIA** compounds are those where Ar¹ is just described, and Ar² is a thienyl, an unsubstituted phenyl or a phenyl substituted with one or more substituents independently selected from halogen, alkyl, alkoxy, CO₂H or CO₂R.


-28-

Examples of other compounds where R^1 is T- Ar^1 are shown below where A is N or CH, and T is one of the following: CH_2 (exemplified by **IVA-1**), $S(O)_2$ (**VA-1**), CONH (**VIA-1**), $COCH_2$ (**VIIA-1**), CO_2 , (**VIIIA-1**),
 5 and $S(O)_2NH$ (**IXA-1**). In other examples of these embodiments the phenyl rings may be optionally substituted as described above.

**IVA-1** ($R^2=CH_3$)**VA-1****VIA-1** ($R^{13} = H$)**VIA-2** ($R^{13} =$ **VIA-3** (thiourea**VIIA-1****VIIIA-1****IXA-1**


Another embodiment of this invention relates to compounds of formula **IA** where R^1 is T-R, R is a C_3-C_6 cycloalkyl ring or a C_1-C_6 straight chain or branched alkyl or alkenyl group optionally substituted by halogen and T is as described above.
 20 When R^1 is T-R, preferred compounds are those where T is $C(=O)$ as represented by formula **XA**. Table 4 below shows representative examples of **XA** compounds.

-29-

Table 4. Examples of **XA** Compounds (R^2 is CH_3)

No.	Ar^1	R
XA-1	phenyl	CH_3
XA-2	4-F-phenyl	CH_3
XA-3	phenyl	Cyclopentyl
XA-4	phenyl	isobutyl
XA-5	phenyl	propyl


5 Preferred R^2 groups of formula **I** include
 $-CH_2OR$, $-CH_2OH$, $-CH_2$ (heterocyclyl), $-CH_2$ (substituted
heterocyclyl), $-CH_2N(R)_2$, and an R group such as
methyl. Representative examples of compounds wherein
 R^2 is other than methyl (formula **IXA**) are shown in
10 Table 5 below.

**IXA** (R^2 is other than CH_3)Table 5. Examples of Compound **IXA**


No.	Ar^1	A	R^1	R^2
XIA-1	phenyl	CH	phenyl	CH_2 (morpholin-4-yl)
XIA-2	phenyl	CH	phenyl	$CH_2N(CH_3)_2$
XIA-3	phenyl	CH	phenyl	CH_2NEt_2
XIA-4	phenyl	CH	phenyl	$CH_2N(CH_3)CH_2Ph$
XIA-5	phenyl	CH	phenyl	CH_2 (1-t-butoxycarbonylpiperazin-4-yl)

No.	Ar ¹	A	R ¹	R ²
XIA-6	phenyl	CH	benzyl	CH ₂ (morpholin-4-yl)
XIA-7	phenyl	CH	cyclohexyl	CH ₂ (morpholin-4-yl)
XIA-8	phenyl	CH	4-[1,2-(OMe) ₂ -phenyl]	CH ₂ (morpholin-4-yl)
XIA-9	phenyl	CH	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-10	phenyl	CH	phenyl	CH ₂ N(CH ₃)CH ₂ CH ₂ N(CH ₃) ₂
XIA-11	phenyl	CH	phenyl	CH ₂ N(CH ₃)CH ₂ CO ₂ CH ₃
XIA-12	phenyl	CH	phenyl	CH ₂ (piperazin-1-yl)
XIA-13	phenyl	N	2-thienoyl	CH ₂ Br
XIA-14	phenyl	N	2-thienoyl	CH ₂ (morpholin-4-yl)
XIA-15	4-F-phenyl	CH	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-16	4-F-phenyl	CH	3-cyanophenyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-17	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-18	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-19	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-20	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-21	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-22	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-23	4-F-phenyl	CH	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-24	4-F-phenyl	CH	cyclohexyl	CH ₂ (morpholin-4-yl)
XIA-25	4-F-phenyl	CH	3-cyanophenyl	CH ₂ (morpholin-4-yl)
XIA-26	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ (morpholin-4-yl)
XIA-27	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ (morpholin-4-yl)
XIA-28	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₃
XIA-29	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₃
XIA-30	4-F-phenyl	CH	3-cyanophenyl	CH ₂ OCH ₃
XIA-31	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₃
XIA-32	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ OCH ₃
XIA-33	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₃
XIA-34	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₃
XIA-35	4-F-phenyl	CH	3-cyanophenyl	CH ₂ OCH ₃

-31-

No.	Ar ¹	A	R ¹	R ²
XIA-36	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₃
XIA-37	4-F-phenyl	CH	4-cyclohexanol	CH ₂ O(tetrahydrofuran-3-yl)
XIA-38	4-F-phenyl	CH	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-39	phenyl	N	2-thienoyl	CH ₂ (piperidin-1-yl)
XIA-40	phenyl	N	2-thienoyl	CH ₂ (piperazin-1-yl)
XIA-41	4-F-phenyl	CH	4-methoxybenzyl	CH ₂ OCH ₃
XIA-42	4-F-phenyl	N	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-43	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₃
XIA-44	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ (phenyl)
XIA-45	4-F-phenyl	N	cyclohexyl	CH ₂ OH
XIA-46	4-F-phenyl	N	CH ₂ CH ₂ (pyridin-2-yl)	CH ₂ OH
XIA-47	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₃
XIA-48	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₃
XIA-49	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-50	4-F-phenyl	N	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-51	4-F-phenyl	N	cyclohexyl	
XIA-52	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ (phenyl)
XIA-53	4-F-phenyl	N	CH ₂ CH ₂ (pyridin-2-yl)	CH ₂ OCH ₂ (phenyl)




The XYZ-containing ring of formula **I** may be an isoxazole ring as shown above or it may an isomeric isoxazole or "reverse" isoxazole (**IB**). In this embodiment Q is preferably a pyrimidine or pyridine ring where A is N or CH, or Q is a pyrazole ring, and R² is aliphatic or substituted aliphatic.

**IB**





Examples of **IB** compounds are shown in Table 6 below.

SUBSTITUTE SHEET (RULE 26)




-32-

Table 6**IB-1****IB-2****IB-3**













5.

**IB-4****IB-5****IB-6****IB-7**

10



**IB-8****IB-9****IB-10**

-33-


**IB-11****IB-12****IB-13****IB-14****IB-15****IB-16****IB-17****IB-18****IB-19****IB-20****IB-21****IB-22**

10

-34-

**IB-23****IB-24**




In another embodiment of this invention,
 5 the XYZ-containing ring is a pyrazole ring of
 formula **IC**:

**IC**

10



For compounds of formula **IC**, G is preferably an
 optionally substituted aryl. Specific examples of
IC compounds are shown in Table 7 below.

15



**Q1****Q2****Q3**Table 7. Examples of **IC** Compounds

No.	G	Q	R ¹	R ²
IC-1	4-F-phenyl	Q2	Phenyl	H
IC-2	4-F-phenyl	Q2	Cyclohexyl	H

-35-

No.	G	Q	R ¹	R ²
IC-4	4-F-phenyl	Q2	6-MeO-naphthalen-2-yl	H
IC-5	4-F-phenyl	Q2	4-cyclohexanol	H
IC-6	4-F-phenyl	Q1	Phenyl	H
IC-7	4-F-phenyl	Q1	Cyclohexyl	H
IC-8	4-F-phenyl	Q1	4-cyclohexanol	H
IC-9	4-F-phenyl	Q2	Cyclohexyl	CH ₃
IC-10	4-F-phenyl	Q2	Cyclohexyl	CH ₂ -N 
IC-11	Phenyl	Q2	Cyclohexyl	CH ₂ -N 

Other embodiments of this invention relate to compounds where the XYZ-containing ring is a furan (**ID**) or a triazole (**IE**). These embodiments are exemplified below where R¹ is phenyl, R²' is hydrogen, and A is N or CH.

**ID-1** (R² = H)**ID-2** (R² = CH₃)**IE-1** (R³ = H)**IE-1** (R³ = CH₃)

10

For compounds of formula **IB-IE**, the phenyl rings of Ar¹ and Ar² may be optionally substituted as shown above for the isoxazoles of formula **IA**.


The compounds of this invention may be prepared in general by methods known to those skilled in the art for analogous compounds, as

15

-36-

illustrated by the general schemes below and by the preparative examples that follow.

Scheme I




(a) NCS, cat. py, CHCl_3 ; (b) $(\text{CH}_3\text{CO})_2\text{CH}_2$, Et_3N , EtOH ; (c) DMA-DMF, reflux; (d) guanidine hydrochloride, NaOMe, MeOH, reflux; (e) PhBr, Pd_2dba_3 , BINAP, NaOtBu, toluene; (f) RCOCl , py, benzene, reflux


Scheme I above shows a route for making isoxazoles where Q is a pyrimidine ring. The starting benzaldehyde oxime **1** may be converted to the α -chlorobenzaldehyde oxime **2** using N-chlorosuccinimide and a catalytic amount of pyridine. Condensation of **2** with 2,4-pentanedione provides the isoxazole **3** which may be treated with dimethylformamide dimethylacetal to obtain the enamine **4**. After an aqueous work-up and without purification, **4** may be cyclized with guanidine hydrochloride to the aminopyrimidine **5**. Compounds of formula **IIA** may be obtained from **5** according to step (e) using the appropriate arylbromide in the presence of tris(dibenzylideneacetone) dipalladium.

-37-

Alternatively, **5** may be treated with the appropriate acid chloride in a pyridine/benzene solvent according to step (f) to give compounds of formula **IVA**. If the acid chloride is a Ar^2COCl , compounds of **5** formula **IIIA** may be obtained in a similar manner.

Scheme II


-38-



Reagents: (a) i. LDA, ii. 2-benzyloxy-N-methoxy-N-methyl-acetamide, -78°C to rt; (b) Et_3N , EtOH, rt to reflux; (c) oxone; (d) iodotrimethylsilane; (e) PPh_3 , CBr_4 ; (f) morpholine, Et_3N ; (g) 4-aminocyclohexanol, DMSO, 80°C ; (h) NaOEt , EtOH; (i) cyclohexylamine, DMSO, 80°C ; (j) 3:1 trifluoroacetic acid/ H_2O ; 100°C .

Scheme II above shows a route for making isoxazoles of this invention where Q is a pyrimidine ring and R^2 is modified by various groups.

Scheme III



Reagents: (a) i. LDA, ii. N-methoxy-N-methyl-acetamide, -78°C to rt; (b) Et_3N , EtOH, rt to reflux; (c) N-bromosuccinimide, AIBN, CCl_4 , reflux; (d)


-39-

morpholine, K_2CO_3 , DMF; (e) NaOMe, MeOH; (f) aniline, $Pd_2(dba)_3$, BINAP, NaOtBu, toluene, $80^\circ C$; (g) cyclohexylamine, $Pd_2(dba)_3$, BINAP, NaOtBu, toluene, $80^\circ C$.

5

Scheme III above shows a synthetic route for making isoxazoles of this invention where Q is a pyridine and R^2 is modified by various groups. In Scheme II and Scheme III, the isoxazole ring is first constructed and then the 2-position of the pyrimidine or pyridine ring is elaborated with the appropriate NHR^1 substitution. It will be apparent to one skilled in the art that position 2 of the pyrimidine or pyridine ring can be elaborated with the appropriate NHR^1 substitution before the isoxazole ring is constructed. Accordingly, isoxazoles of this invention may be obtained by performing step (b) using an appropriate intermediate having the formula **XII**:

20


**XII**

where A is N or CH; R^1 and R^2 are as described above; and PG is hydrogen or a nitrogen protecting group. Nitrogen protecting groups are well-known and include groups such as benzyl or CO_2R , where R is preferably alkyl, allyl or benzyl.

25

-40-

Scheme IV




- Reagents: (a) Et_3N , EtOH ; (b) DIBAL, toluene, 0°C ; (c) CH_3MgBr , THF; (d) $(\text{COCl})_2$, DMSO, Et_3N , CH_2Cl_2 ; (e) DMF·DMA, toluene, reflux; (f) i. thiourea, MeONa, MeOH, ii. pyridine, chloroform, CH_3I (g) m-CPBA, CH_2Cl_2 ; (h) R^1NH_2 , DMSO

Scheme IV above shows a synthetic route for making reverse isoxazoles of this invention where Q is a pyrimidine ring.

-41-


Scheme V




Reagents: (a) i. LDA, ii. N-methoxy-N-methylbenzamide; (b) Cl-C(=N-OH)CO₂Et, EtOH, Et₃N, 80° C; (c) diisobutylaluminum hydride, CH₂Cl₂, room temperature; (d) PPh₃, CBr₄, CH₂Cl₂; (e) piperidine, K₂CO₃, DMF; (f) BINAP, Pd₂(dba)₃, NaOtBu, cyclohexylamine, toluene, 80° C.

Scheme V above shows a synthetic route for making reverse isoxazoles of this invention where Q is a pyridine ring.

Scheme VI




-42-



Reagents: a) $\text{NH}_2\text{OH}/\text{HCl}$, $\text{H}_2\text{O}/\text{EtOH}$; Na_2CO_3 ; (b) NCS ,
 5 cat. pyridine, CHCl_3 ; (c) $\text{CH}_3\text{COCH}_2\text{CO}_2\text{CH}_3$, Et_3N , EtOH ;
 (d) i. NaOH , MeOH , H_2O ; then, ii. SOCl_2 , heat; (e)
 $\text{HO}_2\text{CCH}_2\text{CN}$, $n\text{-BuLi}$, -78 to 0°C ; (f) H_2NNH_2 , EtOH ; (g)
 R-X , dioxane.

10 Scheme VI above shows a general route for
 preparing compounds of this invention wherein Q is a
 pyrazole ring.

Scheme VII




15

Reagents: (a) LDA , THF ; (b) $4\text{-F-C}_6\text{H}_4\text{CO}_2\text{Et}$; (c)
 $\text{DMF}\cdot\text{DMA}$, toluene, reflux; (d) $\text{H}_2\text{NNH}_2\cdot\text{H}_2\text{O}$, EtOH ,
 20 reflux; (e) R^1NH_2 , sealed tube, 140°C .

-43-

Scheme VII above shows a general route for preparing compounds of this invention wherein the XYZ ring is a pyrazole ring.

Certain of the intermediates that are
 5 useful for making the kinase inhibitors of this invention are believed to be novel. Accordingly, one embodiment of this invention relates to compounds **XII** above and compounds represented by formula **XIII**:

**XIII**

wherein:

X-Y is N-O or O-N providing an isoxazole or reverse isoxazole ring;

15 A is N or CH;

G is R, aryl or substituted aryl;

R is aliphatic or substituted aliphatic

R² is selected from hydrogen, -R, -CH₂OR, -CH₂OH, -CH=O, -CH₂SR, -CH₂S(O)₂R, -CH₂(C=O)R, -CH₂CO₂R,

20 -CH₂CO₂H, -CH₂CN, -CH₂NHR, -CH₂N(R)₂, -CH=N-OR, -CH=NNHR, -CH=NN(R)₂, -CH=NNHCOR, -CH=NNHCO₂R, -CH=NNHSO₂R, -aryl, -substituted aryl, -CH₂(aryl), -CH₂(substituted aryl), -CH₂NH₂, -CH₂NHCOR, -CH₂NHCONHR, -CH₂NHCON(R)₂, -CH₂NRCOR, -CH₂NHCO₂R, -CH₂CONHR, -CH₂CON(R)₂, -CH₂SO₂NH₂,
 25 -CH₂(heterocyclyl), -CH₂(substituted heterocyclyl), -(heterocyclyl), or -(substituted heterocyclyl); and

R¹ is selected from halogen, NH₂, SR, or SO₂R.

-44-

The activity of the JNK inhibitors of this invention may be assayed *in vitro*, *in vivo* or in a cell line. *In vitro* assays include assays that determine inhibition of either the kinase activity
5 or ATPase activity of activated JNK. For example, see the testing examples described below. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to JNK and may be measured either by radiolabelling the inhibitor prior to binding,
10 isolating the inhibitor/JNK complex and determining the amount of radiolabel bound, or by running a competition experiment where new inhibitors are incubated with JNK bound to known radioligands. One may use any type or isoform of JNK, depending upon
15 which JNK type or isoform is to be inhibited.

The JNK inhibitors or pharmaceutical salts thereof may be formulated into pharmaceutical compositions for administration to animals or humans. These pharmaceutical compositions, which
20 comprise an amount of JNK inhibitor effective to treat or prevent a JNK-mediated condition and a pharmaceutically acceptable carrier, are another embodiment of the present invention.

The term "JNK-mediated condition", as used
25 herein means any disease or other deleterious condition in which JNK is known to play a role. Such conditions include, without limitation, inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders,
30 cancer, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia,

-45-

vascular hyperplasia, cardiac hypertrophy, thrombin-induced platelet aggregation, and conditions associated with prostaglandin endoperoxidase synthase-2.

5 Inflammatory diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, acute pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome.

10 Autoimmune diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, chronic thyroiditis, Graves' disease,
15 autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease,
20 psoriasis, or graft vs. host disease.

 Destructive bone disorders which may be treated or prevented by the compounds of this invention include, but are not limited to, osteoporosis, osteoarthritis and multiple myeloma-
25 related bone disorder.

 Proliferative diseases which may be treated or prevented by the compounds of this invention include, but are not limited to, acute myelogenous leukemia, chronic myelogenous leukemia,
30 metastatic melanoma, Kaposi's sarcoma, multiple myeloma and HTLV-1 mediated tumorigenesis.

-46-

Angiogenic disorders which may be treated or prevented by the compounds of this invention include solid tumors, ocular neovascularization, infantile haemangiomas. Infectious diseases which
5 may be treated or prevented by the compounds of this invention include, but are not limited to, sepsis, septic shock, and Shigellosis.

Viral diseases which may be treated or prevented by the compounds of this invention
10 include, but are not limited to, acute hepatitis infection (including hepatitis A, hepatitis B and hepatitis C), HIV infection and CMV retinitis.

Neurodegenerative diseases which may be treated or prevented by the compounds of this
15 invention include, but are not limited to, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis (ALS), epilepsy, seizures, Huntington's disease, traumatic brain injury, ischemic and hemorrhaging stroke, cerebral
20 ischemias or neurodegenerative disease, including apoptosis-driven neurodegenerative disease, caused by traumatic injury, acute hypoxia, ischemia or glutamate neurotoxicity.

"JNK-mediated conditions" also include
25 ischemia/reperfusion in stroke, heart attacks, myocardial ischemia, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, hepatic ischemia, liver disease, congestive heart failure, pathologic immune responses such as that caused by T cell
30 activation and thrombin-induced platelet aggregation.

-47-

In addition, JNK inhibitors of the instant invention may be capable of inhibiting the expression of inducible pro-inflammatory proteins. Therefore, other "JNK-mediated conditions" which may
5 be treated by the compounds of this invention include edema, analgesia, fever and pain, such as neuromuscular pain, headache, cancer pain, dental pain and arthritis pain.

The compounds of this invention are also
10 useful as inhibitors of Src-family kinases, especially Src and Lck. For a general review of these kinases see Thomas and Brugge, *Annu. Rev. Cell Dev. Biol.* (1997) **13**, 513; Lawrence and Niu, *Pharmacol. Ther.* (1998) **77**, 81; Tatosyan and
15 Mizenina, *Biochemistry (Moscow)* (2000) **65**, 49. Accordingly, these compounds are useful for treating diseases or conditions that are known to be affected by the activity of one or more Src-family kinases. Such diseases or conditions include hypercalcemia,
20 restenosis, hypercalcemia, osteoporosis, osteoarthritis, symptomatic treatment of bone metastasis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, T-cell mediated hypersensitivity
25 disease, Hashimoto's thyroiditis, Guillain-Barre syndrome, chronic obstructive pulmonary disorder, contact dermatitis, cancer, Paget's disease, asthma, ischemic or reperfusion injury, allergic disease, atopic dermatitis, and allergic rhinitis. Diseases
30 that are affected by Src activity, in particular, include hypercalcemia, osteoporosis, osteoarthritis, cancer, symptomatic treatment of bone metastasis,

-48-

and Paget's disease. Diseases that are affected by Lck activity, in particular, include autoimmune diseases, allergies, rheumatoid arthritis, and leukemia. Compounds of formula II-A and I-B wherein
5 Ar² is aryl are especially useful for treating diseases associated with the Src-family kinases, particularly Src or Lck.

In addition to the compounds of this invention, pharmaceutically acceptable derivatives
10 or prodrugs of the compounds of this invention may also be employed in compositions to treat or prevent the above-identified disorders.

A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable
15 salt, ester, salt of an ester or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active
20 metabolite or residue thereof. Particularly favored derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal (e.g., by allowing an orally administered compound
25 to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

Pharmaceutically acceptable prodrugs of
30 the compounds of this invention include, without limitation, esters, amino acid esters, phosphate esters, metal salts and sulfonate esters.

-49-

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

-50-

Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum
5 proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium
10 hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes,
15 polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally,
20 buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and
25 intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

Sterile injectable forms of the compositions of this invention may be aqueous or
30 oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and

-51-

suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or

-52-

solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral
5 administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening,
10 flavoring or coloring agents may also be added.

Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the
15 agent with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

20 The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the
25 skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract can be effected in a rectal
30 suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

-53-

For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers.

5 Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively,
10 the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral
15 oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

For ophthalmic use, the pharmaceutical compositions may be formulated as micronized
20 suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the
25 pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared
30 according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or

-54-

other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

5 The amount of JNK inhibitor that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so
10 that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

 It should also be understood that a specific dosage and treatment regimen for any
15 particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the
20 treating physician and the severity of the particular disease being treated. The amount of inhibitor will also depend upon the particular compound in the composition.

 According to another embodiment, the
25 invention provides methods for treating or preventing a JNK-mediated condition comprising the step of administering to a patient one of the above-described pharmaceutical compositions. The term "patient", as used herein, means an animal,
30 preferably a human.

 Preferably, that method is used to treat or prevent a condition selected from inflammatory

-55-

diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases, degenerative diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, and thrombin-induced platelet aggregation, or any specific disease or disorder described above.

Depending upon the particular JNK-mediated condition to be treated or prevented, additional drugs, which are normally administered to treat or prevent that condition, may be administered together with the inhibitors of this invention. For example, chemotherapeutic agents or other anti-proliferative agents may be combined with the JNK inhibitors of this invention to treat proliferative diseases.

Those additional agents may be administered separately, as part of a multiple dosage regimen, from the JNK inhibitor-containing composition. Alternatively, those agents may be part of a single dosage form, mixed together with the JNK inhibitor in a single composition.

In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

30 Example 1

Benzaldehyde oxime. To benzaldehyde (10.0 g, 94 mmol) in ethanol (50 mL) was added hydroxylamine

-56-

hydrochloride (6.5 g, 94 mmol in H₂O (50 mL) followed by Na₂CO₃ in H₂O (50 mL). Reaction solution was stirred for 2 hr. Poured into brine and extracted twice with diethyl ether. Combined extracts were
5 dried over MgSO₄. Evaporation afforded benzaldehyde oxime (11.0 g, 96.5% yield) as a colorless oil. ¹H NMR (CDCl₃) δ 7.40-7.50 (m, 3H), 7.60-7.70 (m, 2H), 8.22 (s, 1H), 9.1 (bs, 1H).

10

Example 2α-Chlorobenzaldehyde oxime (Benzoyl chloride oxime).

To benzaldehyde oxime (12.2 g, 0.1 mol) in chloroform was added catalytic amount of pyridine, followed by N-chlorosuccinimide (13.35 g, 0.1 mol)
15 at room temperature. The reaction mixture was stirred for 1.5 h, then saturated aqueous NaCl was added. The organic phase was washed with saturated aqueous NaCl (twice) and dried with MgSO₄. The solvent was removed under reduced pressure. 13.85g
20 α-chlorobenzaldehyde oxime was obtained. The yield was 87%.

Example 31-(5-Methyl-3-phenyl-isoxazol-4-yl)-ethanone

(Compound 3). To a solution of pentane-2,4-dione
25 (13.23 g, 0.132 mol) and triethylamine (13.35 g, 0.132 mol) in ethanol was added α-chlorobenzaldehyde oxime (13.70 g, 0.088 mol) at room temperature. The reaction mixture was stirred overnight at room temperature. To the reaction was added ethyl acetate
30 and saturated aqueous NaCl. The organic phase was washed with saturated aqueous NaCl (twice) and dried

-57-

with MgSO_4 , and the organic solvent was removed under reduced pressure to provide 17.7g of the title compound. The yield was 100%.

Example 4

5 4-(5-methyl-3-phenyl-isoxazol-4-yl)-pyrimidin-2-ylamine (Compound 5). The above Compound 3 (17.7 g, 0.088 mol) and dimethylformamide dimethyl acetal (DMF·DMA) (160 g, 0.132mol) were refluxed overnight. To the reaction mixture was added ethyl acetate and
10 saturated aqueous NaCl. The organic phase was washed with saturated aqueous NaCl (twice) and dried (MgSO_4). The organic solvent was removed under reduced pressure, and the crude product material was dissolved in 200 mL methanol. To the solution was
15 added guanidine hydrochloride (10.5 g, 0.110 mol) in 100 mL methanol, followed by sodium methoxide (6.17 g, 0.114 mol) in 100 mL methanol. The reaction mixture was refluxed overnight and then was cooled to room temperature. The reaction solvent was
20 concentrated to approximately 100mL total volume, and the precipitated product was filtered. The filtration cake afforded the title compound (9.3 g). The overall yield for two steps was 46%.

25

Example 5

[4-(5-Methyl-3-phenyl-isoxazol-4-yl)-pyrimidin-2-yl]-phenyl-amine (Compound IIA). To a solution of 50 mg (0.2 mmol) of 4-(5-methyl-3-phenyl-isoxazole-4-yl)-pyrimidin-2-ylamine in 1 mL of toluene was
30 added successively 63 μL (0.6 mmol) of bromobenzene, 10 mg of tris(dibenzylideneacetone) dipalladium, 10

-58-

mg of BINAP and 39 mg (0.4 mmol) of sodium *tert*-butoxide. The mixture was heated at reflux for 16 h, diluted with ethyl acetate, filtered, washed successively with saturated aqueous sodium bicarbonate and brine, dried (MgSO₄) and concentrated *in vacuo*. The residue was purified by column chromatography over silica gel eluted with ethyl acetate-hexanes 1:3, to afford 24 mg (36%) of the title compound as a yellow oil.

10

Example 6

5-Methyl-3-phenyl-isoxazole-4-carboxylic acid methyl ester. An ethanol solution of freshly prepared benzoyl chloride oxime (14.0 g, 90 mmol) (100 mL) was added dropwise, at 5 °C to methyl acetoacetate (11.18 g, 96 mmol) and triethyl amine (13 mL, 103 mmol) in ethanol (50 mL). After stirring for 12 hr at ambient temperature, the solution was diluted with CH₂Cl₂, washed with 1N HCl, saturated NaHCO₃, brine, dried over MgSO₄ and evaporated to give amber oil. Flash chromatography (silica) with 10% ethyl acetate in hexanes afforded the title compound (7.56 g, 39% yield) as a white solid: MS *m/z* MH⁺218 (100); ¹H NMR (CDCl₃) δ 2.78 (s, 3H), 3.81 (s, 3H), 7.45-7.55 (m, 3H), 7.65-7.69 (m, 2H).

25

Example 7

5-Methyl-3-phenyl-isoxazole-4-carboxylic acid. To 5-Methyl-3-phenyl-isoxazole-4-carboxylic acid methyl ester (0.853 g, 3.69 mmol) in methanol (12 mL) was added 2N NaOH (8 mL) the reaction solution was stirred at ambient temperature for 60 hr. The

30

-59-

solution was dilute with water and extracted twice with ethyl acetate. The combined extract was washed with brine and dried over MgSO_4 and concentrated. Recrystallization (hexanes / ethyl acetate) afforded
5 a white solid (0.540 g, 72% yield).

Example 8

5-Methyl-3-phenyl-isoxazole-4-carbonyl chloride. 5-Methyl-3-phenyl-isoxazole-4-carboxylic acid (0.54g,
10 2.56 mmol) was treated with SOCl_2 (2 mL) at 70 °C for 1 hr. Concentration in vacuum gave a yellow oil which was used without purification.

Example 9

15 3-(5-Methyl-3-phenyl-isoxazol-4-yl)-3-oxo-propionitrile. To cyanoacetic acid (0.43 g, 5.12 mmol) in THF at -78 °C, containing one crystal of 1,1'-bipyridyl was added n-butyl lithium (6.4 mL, 10.24 mmol). The temperature was allowed to warm to
20 0 °C resulting in a pink colored solution. After cooling to -78 °C, 5-Methyl-3-phenyl-isoxazole-4-carbonyl chloride (0.567 g, 2.56 mmol) in THF (5 mL) was added dropwise. The mixture was stirred at -78 °C for 1 hr. and at ambient temperature for an
25 addition 1 hr. The reaction was quenched with 1N HCl (13 mL) and extracted twice with CH_2Cl_2 . Combined extracts were washed with saturated NaHCO_3 , brine, dried over MgSO_4 to give the title compound (0.391 g, 67 % yield).

30

Example 10

-60-

N-[5-(5-Methyl-3-phenyl-isoxazol-4-yl)-2H-pyrazol-3-yl]-benzamide. 3-(5-Methyl-3-phenyl-isoxazol-4-yl)-3-oxo-propionitrile (0.391 g, 1.73 mmol) in Ethanol (3 mL) was treated with hydrazine (0.168 mL, 3.46 mmol) and heated to reflux. Evaporation in vacuum gave 5-(5-Methyl-3-phenyl-isoxazol-4-yl)-2H-pyrazol-3-ylamine used without purification. To the resulting amine (0.039 g, 0.16 mmol) in dioxane was added triethyl amine followed by benzyl chloride (0.019 mL, 0.16 mmol). The reaction was stirred at 10 °C for 1 hr and 2 hr at ambient temperature. The solution was diluted with water extracted with ethyl acetate, washed with saturated NaHCO₃, Brine, dried over MgSO₄ and concentrated in vacuum. HPLC purification afforded 1.4 mg of title compound.

Example 11

1-Benzyloxy-3-(2-methylsulfanylpirimidin-4-yl)-propan-2-one (Compound 7). To a stirred solution of 4-methyl-2-methylsulfanylpirimidine (9.60g, 68.5 mmol) in THF (150 mL) at -78 °C was added LDA (2.0 M THF/Hex, 41.1 mL, 82.2 mmol) dropwise over 10 min. The solution was stirred at -78 °C for 15 minutes, warmed to 0 °C for 10 minutes and recooled to -78 °C for 15 minutes. Then, a solution of 3-benzyloxy-N-methyl-N-methoxyacetamide (17.2 g, 82.2 mmol) in THF (30 mL) was added dropwise over 45 minutes. After 15 min. at -78 °C, the solution was warmed to 0 °C and stirred for 30 min. The reaction was quenched with HCl (1M, 85 mL) and stirred for 1 h. The solution was poured into saturated NaHCO₃ (300 mL), extracted with Et₂O (3 × 200 mL), dried (MgSO₄),

-61-

filtered and concentrated. Flash chromatography (SiO₂, 20% EtOAc-hexanes) provided the title compound (13.75 g, 47.7 mmol, 69% yield).

5 Example 12

4-[5-Benzyloxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-2-methylsulfanyl-pyrimidine (Compound 8). To a stirred solution of the above compound 7 (13.75 g, 47.7 mmol) and Et₃N (14.6 mL, 105 mmol) in EtOH (200 mL), was added a solution of 4-fluoro-benzoylchloride oxime (56 mmol) in EtOH (50 mL) over 30 min. The solution was stirred at 25 °C for 15 min. Then, the solution was heated to reflux for 90 min. The solution was cooled to 25 °C. Additional Et₃N (7.3 mL, 52 mmol) was added followed by dropwise addition of a solution of 4-fluoro-benzoylchloride oxime (38.5 mmol) in EtOH (50 mL) over 1 h. to drive the reaction to completion. The solution was refluxed for 1 h. until TLC indicated that all of the starting isoxazole was consumed. The solution was cooled to 25 °C and concentrated. The crude material was picked up in CH₂Cl₂ (50mL) and poured into saturated aqueous NaHCO₃ (150 mL), extracted with CH₂Cl₂ (3 × 150 mL), dried (MgSO₄), filtered and concentrated. Flash chromatography (SiO₂, 20% EtOAc-hexanes) provided the title compound (14.2 g, 34.8 mmol, 60%) in sufficient purity (> 85%) for use in the next reaction.

30 Example 13

4-[5-Benzyloxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-2-methanesulfonyl-pyrimidine (Compound 9). To a

-62-

stirred solution of the above compound 8 (2.00 g, 4.91 mmol) in MeOH (50 mL) at 25 °C was added dropwise a solution of oxone (7.07 g, 11.5 mmol) in H₂O (50 mL) over 10 min. After 20 h., the solution
5 was poured into H₂O (75 mL), extracted with CH₂Cl₂ (3 × 75 mL), dried (MgSO₄), filtered and concentrated. Flash chromatography (SiO₂, 45% EtOAc-hexanes) provided the title compound (1.60 g, 3.64 mmol, 74%).

10

Example 14

[3-(4-Fluoro-phenyl)-4-(2-methanesulfonyl-pyrimidin-4-yl)-isoxazol-5-yl]-methanol (Compound 10). To a stirred solution of the above compound 9 (750 mg, 1.70 mmol) in CHCl₃ (8.5 mL) at 0 °C was added
15 trimethylsilyl iodide (0.73 mL, 5.1 mmol). The reaction was stirred at 0 °C for 30 min. Then, additional trimethylsilyl iodide (0.48 mL, 3.4 mmol) was added. After 40 min. the solution was warmed to 25 °C and stirring was continued for 22 h. The
20 solution was quenched with H₂O-MeOH (2 mL) and stirred for 1 h. The solution was poured into saturated aqueous NaHCO₃ (30 mL), extracted with EtOAc (3 × 30 mL), and concentrated. Flash chromatography (SiO₂, 80% EtOAc-hexanes) provided the
25 title compound (530 mg, 1.52 mmol, 89%).

Example 15

4-[5-(Bromomethyl)-3-(4-fluoro-phenyl)-isoxazol-4-yl]-2-methanesulfonyl-pyrimidine (Compound 11). To a stirred solution of the above compound 10 (250 mg, 0.716 mmol) and CBr₄ (473 mg, 1.43 mmol) in CH₂Cl₂
30 (14 mL) at 25 °C was added PPh₃ (244 mg, 0.93 mmol).

-63-

After 10 min., additional PPh₃ (50 mg, 0.19 mmol) was added to drive the reaction to completion. After 15 min., the solution was concentrated. Flash chromatography (SiO₂, 50% EtOAc-hexanes) provided the
5 title compound (265 mg, 0.643 mmol, 90%).

Example 16

4-[3-(4-Fluoro-phenyl)-4-(2-methanesulfonyl-pyrimidin-4-yl)-isoxazol-5-ylmethyl]-morpholine

10 (Compound 12). To a stirred solution of the above compound 11 (41 mg, 0.099 mmol) and Et₃N (20 μL, 0.15 mmol) in CH₃CN (0.5 mL) at 25 °C was added morpholine (9.6 μL, 0.11 mmol). After 15 min. the solution was concentrated. Preparative thin layer chromatography
15 (SiO₂, EtOAc) provided the title compound (29 mg, 0.069 mmol, 70%).

Example 17

4-{4-[3-(4-Fluoro-phenyl)-5-(morpholin-4-ylmethyl)-isoxazol-4-yl]pyrimidin-2-ylamino}cyclohexanol

20 (Compound XIA-42). A stirred solution of Compound 13 (29 mg, 0.069 mmol) and *trans*-4-aminocyclohexanol (24 mg, 0.21 mmol) in DMSO (0.21 mL) was heated to 80 °C for 4 h. The solution was poured into half-saturated aqueous NaHCO₃ (5 mL), extracted with EtOAc
25 (5 × 5 mL), dried (MgSO₄), filtered and concentrated. Flash chromatography (SiO₂, 10% MeOH- CH₂Cl₂) provided material which was further purified by ion exchange chromatography (SCX resin, eluent: 0.25M NH₃
30 in 50% MeOH- CH₂Cl₂) to give the title compound (27 mg, 0.057mmol, 83%).

-64-

Example 18

4-[5-Ethoxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-2-methylsulfanyl-pyrimidine (Compound 13). To
5 a stirred solution of the above compound 8 (103 mg, 0.27 mmol) in EtOH (2.0 mL) at 25 °C was added NaOEt (21% w/v EtOH, 0.40 mL, 1.23 mmol). After 2 h. the reaction was quenched with saturated aqueous NH₄Cl (3 mL), CH₂Cl₂ (3 × 5 mL), dried (MgSO₄), filtered and
10 concentrated. Flash chromatography (SiO₂, 25% EtOAc-hexanes) provided the title compound (58 mg, 0.17 mmol, 62%).

Example 19

15 4-[5-Ethoxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-2-methanesulfonyl-pyrimidine (Compound 14). This compound was prepared in a manner similar to that described above in Example 13, except starting from the above compound 13 (58 mg, 0.17 mmol) to
20 provide the title compound (64 mg, 0.17 mmol, 100%) which was used directly in the next reaction without purification or characterization.

Example 20

25 Cyclohexyl-{4-[5-ethoxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-pyrimidin-2-yl}amine (Compound XIA-43) This compound was prepared in a manner similar to that described above in Example 17, starting from
30 the above compound 14 (64 mg, 0.17 mmol) and cyclohexylamine (58 µL, 0.51 mmol) to provide the title compound as crude product. After HPLC

-65-

purification (C-18, gradient elution, 10-90% H₂O-CH₃CN) and extraction into EtOAc, the crude product was converted to the HCl salt with HCl-Et₂O (1M, 1 mL). The solvents were removed in vacuo the give
5 the title compound as the HCl salt (55 mg, 0.13 mmol, 76% over two steps from compound 13).

Example 21

Cyclohexyl-{4-[5-benzyloxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-pyrimidin-2-yl}amine
10 (Compound XIA-44) This compound was prepared in a manner similar to that described above in Example 17 starting from the above compound 9 (500 mg, 1.14 mmol) and cyclohexylamine (340 μ L, 3.42 mmol).
15 Flash chromatography (SiO₂, 30% EtOAc-hexanes) provided the title compound (488 mg, 1.06 mmol, 93%).

Example 22

20 [4-(2-Cyclohexylamino-pyrimidin-4-yl)-3-(4-fluoro-phenyl)-isoxazol-5-yl]methanol (Compound XIA-45) A stirred solution of the above compound XIA-44 (461 mg, 1.01 mmol) in TFA-H₂O (3:1, 8 mL) was heated to 80 °C for 20 h. The solution was concentrated, and
25 the crude mixture was taken up in CH₂Cl₂ (25 mL), poured into saturated aqueous NaHCO₃ (30 mL), extracted with CH₂Cl₂ (3 $\times$ 25 mL), dried (MgSO₄), filtered and concentrated. TLC (50% EtOAc-hexanes) indicated about 50% consumption of starting
30 compound XIA-44. The crude material was dissolved in TFA-H₂O (3:1, 8 mL) and the resulting solution was heated to 100 °C for 22 h. The solution was

-66-

concentrated, and the crude mixture was taken up in CH_2Cl_2 (25 mL), poured into saturated aqueous NaHCO_3 (30 mL), extracted with CH_2Cl_2 (3 $\times$ 25 mL), dried (MgSO_4), filtered and concentrated. Flash chromatography (SiO₂, 40% EtOAc-hexanes) provided the title compound (313 mg, 0.85 mmol, 84%).

Example 23

1-(2-Bromo-pyridin-4-yl)-propan-2-one (Compound 16).

To a stirred solution of 2-bromo-4-methylpyridine (Compound 15) (20.20g, 117.4 mmol) in THF (250 mL) at -78 °C was added LDA (2.0 M THF/Hex, 70.5 mL, 141 mmol) dropwise over 10 min. The solution was stirred at -78 °C for 35 min. Then a solution of *N*-methoxy-*N*-methyl acetamide (14.5 g, 141 mmol) in THF (30 mL) was added dropwise over 10 min. After 15 min. at -78 °C, the solution was warmed to 0 °C and stirred for 1 h. The solution was poured into H₂O (250 mL), extracted with Et₂O (3 $\times$ 250 mL), dried (MgSO_4), filtered and concentrated. Flash chromatography (SiO₂, 20% EtOAc-hexanes) provided the title compound (16.75 g, 78.2 mmol, 67%).

Example 24

2-Bromo-4-(5-methyl-3-phenyl-isoxazol-4-yl)-pyridine (Compound 17a). To a stirred solution of Compound 16 (1.71 g, 8.0 mmol) and Et₃N (2.23 mL, 16 mmol) in EtOH (16 mL) was added a solution of benzoylchloride oxime (1.62 g, 10.4 mmol) in EtOH (16 mL) over 90 min. The solution was stirred at 25 °C for 90 min. Then, the solution was heated to reflux for 24 h.

-67-

The solution was cooled to 25 °C and concentrated. The crude material was taken up in CH₂Cl₂ (50mL) and poured into saturated aqueous NaHCO₃ (50 mL), extracted with CH₂Cl₂ (3 × 50 mL), dried (Na₂SO₄), and
5 filtered. Flash chromatography (SiO₂, 20% EtOAc-hexanes) provided the title compound (1.32 g, 4.19 mmol, 52%).

2-Bromo-4-[3-(4-fluoro-phenyl)-5-methyl-isoxazol-4-yl]-pyridine (Compound 17b) was similarly prepared
10 starting with 4-fluorobenzoylchloride oxime.

Example 25

2-Bromo-4-(5-bromomethyl-3-phenyl-isoxazol-4-yl)-pyridine (Compound 18a). A stirred solution of the
15 above Compound 17a (404 mg, 1.28 mmol), N-bromosuccinimide (239 mg, 1.35 mmol) and AIBN (11 mg, 0.064 mmol) in CCl₄ (3 mL) was heated to reflux and placed under a 300 W lamp for 18 h. The solution was diluted with CH₂Cl₂ (15 mL), extracted
20 with H₂O (3 × 10 mL), brine (40 mL), dried (MgSO₄), filtered and concentrated. Flash chromatography (SiO₂, 15-20% EtOAc-hexanes) provided the title compound (287 mg, 0.728 mmol, 57%).

2-Bromo-4-[5-bromomethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl]-pyridine (Compound 18b) was similarly
25 prepared starting with Compound 17b.

Example 26

2-Bromo-4-(5-methoxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl)-pyridine (Compound 19b). To the
30 above Compound 18b (200 mg, 0.485 mmol) was added NaOMe (0.5 M in MeOH, 2.0 mL, 1.0 mmol). The

-68-

solution was stirred at 25 °C for 90 min. Then, the solution was poured into brine, extracted with EtOAc (4 × 15 mL), dried (MgSO₄), filtered through a silica plug. Evaporation of the solvent provided the title compound (175 mg; 0.482 mmol, 99%).

Example 27

4-(4-(2-Bromo-pyridin-4-yl)-3-phenyl-isoxazol-5-ylmethyl)-morpholine (Compound 20a). A stirred solution of the above Compound 18a (484 mg, 1.22 mmol), morpholine (0.45 mL, 5.1 mmol) and K₂CO₃ (340 mg, 2.45 mmol) in anhydrous DMF (2 mL) was warmed to 40 °C for 18 h. The solution was poured into brine (10 mL), extracted with CH₂Cl₂ (3 × 15 mL), dried (MgSO₄), and filtered. Flash chromatography (SiO₂, 50% EtOAc-hexanes) provided the title compound (461 mg, 1.15 mmol, 94%).

Example 28

[4-(5-Methyl-3-phenyl-isoxazol-4-yl)-pyridin-2-yl]phenyl-amine (Compound IIA-52). To a stirred solution of the above Compound 17a (20 mg, 0.063 mmol), aniline (7.0 µL, 0.076 mmol) and BINAP (5.6 mg, 0.009 mmol) in toluene (0.6 mL) at 25 °C was added Pd₂(dba)₃ (2.7 mg, 0.003 mmol) followed by NaOtBu (9.1 mg, 0.095 mmol). The solution was heated to 80 °C for 2 h. The solution was cooled, filtered and concentrated. Preparative thin layer chromatography (SiO₂, 5% EtOAc/CH₂Cl₂) provided the title compound (12.6 mg, 0.0385 mmol, 61%).

-69-

Example 29

Cyclohexyl-[4-(5-methoxymethyl-3-(4-fluoro-phenyl)-isoxazol-4-yl)-pyridin-2-yl]-amine (Compound XIA-29). To a stirred solution of the above Compound 19b (20 mg, 0.050 mmol), cyclohexylamine (11 μ L, 0.13 mmol), and BINAP (4.7 mg, 0.0075 mmol) in toluene (0.4 mL) at 25 °C was added $\text{Pd}_2(\text{dba})_3$ (2.3 mg, 0.0025 mmol) followed by NaOtBu (12 mg, 0.13 mmol). The solution was heated to 80 °C for 15 h. The solution was cooled, poured into H_2O (5 mL), extracted with EtOAc (4 $\times$ 5 mL), dried (MgSO_4), filtered and concentrated. HPLC (gradient elution, 90-10% H_2O - CH_3CN) provided the title compound (9.1 mg, 0.022 mmol, 44%).

15

Example 30

3-Methyl-5-phenyl-isoxazole-4-carbonitrile (Compound 24). To an ethyl alcohol solution of benzoylacetonitrile was added 1.5 eq of triethyl amine, followed by 1.5 eq of acetylchloride oxime, the reaction mixture was stirred at r.t. for 4 hours. To the reaction mixture was added ethyl acetate and brine. The organic phase was dried with magnesium sulfate and the solvent was removed under reduced pressure. After chromatographic purification the title compound was obtained in 72% yield.

25

Example 31

3-Methyl-5-phenyl-isoxazole-4-carbaldehyde (Compound 25). To a toluene solution of the above compound 24 was added 1.2 eq of DIBAL-H/HAX at 0°C. The reaction

30

-70-

was stirred at 0°C for 3 hours, allowed to warm to room temperature and was stirred at r.t. overnight. The reaction mixture was transferred to 1N HCl slowly and then extracted with ethyl acetate. The organic
5 phase was dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by chromatograph providing the title compound in 57% yield.

10

Example 321-(3-Methyl-5-phenyl-isoxazol-4-yl)-ethanol

(Compound 26). To the THF solution of the above Compound 25 was slowly added 1.4 eq of methylmagnesium bromide at room temperature. The
15 reaction mixture was stirred at r.t. for 1 h. To the reaction mixture was added ethyl acetate and 1N HCl. The organic phase was washed with brine and dried over magnesium sulfate. The solvent was removed under reduced pressure, and the crude
20 product, obtained in 96% yield, was used directly for the next step without purification.

Example 331-(3-Methyl-5-phenyl-isoxazol-4-yl)-ethanone

(Compound 27). To a dichloromethane solution of
25 oxalyl chloride was added DMSO at -78 °C, the mixture was stirred at -78 °C for 15 min and followed by addition of a dichloromethane solution of compound the above Compound 26. The reaction mixture was stirred for 30 min at -78 °C, then
30 triethylamine was added, after which the reaction mixture was allowed to warm to room temperature gradually. To the reaction mixture was added ethyl

-71-

acetate and brine. The organic phase was dried over magnesium sulfate, and the solvent was removed under reduced pressure. The crude product, obtained in 94% yield, was used directly for the next step without purification.

Example 34

3-Dimethylamino-1-(3-methyl-5-phenyl-isoxazol-4-yl)-propenone (Compound 28). A toluene solution of the above Compound 27 and excess DMF-DMA was refluxed for 20 hours. To the reaction mixture was added ethyl acetate and brine, the organic phase was dried over magnesium sulfate, and the solvent was then removed under reduced pressure. The crude product was used for the next step without purification.

Example 35

4-(3-Methyl-5-phenyl-isoxazol-4-yl)-2-methylsulfanyl-pyrimidine (Compound 29). A methanol suspension of the above Compound 28, 2 equivalents of thiourea and 1.5 equivalents of sodium methoxide was refluxed for 2 days. To the reaction mixture was added ethyl acetate and 1N HCl, the organic phase was washed with brine and dried over magnesium sulfate, and the solvent was then removed under reduced pressure. The crude product was dissolved in chloroform, to it was added 1.5 eq of iodomethane and 1.5 eq of pyridine. The reaction mixture was stirred at r.t. for 2 hours. To the reaction mixture was added dichloromethane and 1N HCl, the organic phase was washed with brine and dried with

-72-

magnesium sulfate. The solvent was removed under reduced pressure, and the crude product was purified by chromatography to provide the title compound. The yield was 32%.

5

Example 36

4-(3-Methyl-5-phenyl-isoxazol-4-yl)-2-methanesulfonyl-pyrimidine (Compound 30). To a dichloromethane solution of the above Compound 29
10 was added 2 eq of m-CPBA, and the reaction was stirred at r.t. for overnight. The reaction mixture was washed with 1N NaOH twice and brine twice and dried with magnesium sulfate. The solvent was removed under reduced pressure and the crude product
15 was purified by chromatograph to provide the title compound in 79% yield.

Example 37

Compounds IB. A DMSO solution of the above Compound
20 30 and 3 equivalents of desired amine was heated at 80 °C for 4 hours. After analytical HPLC indicated the reaction was completed, the crude product was purified by reversed HPLC to provide the desired Compound IB. The yield is generally greater than
25 80%.

The following examples demonstrate how the compounds of this invention may be tested as protein kinase inhibitors, especially inhibitors of c-Jun-N-terminal kinases.

30

Example 38

Cloning, Expression and Purification of JNK3 Protein

-73-

A BLAST search of the EST database using the published JNK3 α 1 cDNA as a query identified an EST clone (#632588) that contained the entire coding sequence for human JNK3 α 1. Polymerase chain

5 reactions (PCR) using *pfu* polymerase (Stratagene) were used to introduce restriction sites into the cDNA for cloning into the pET-15B expression vector at the NcoI and BamHI sites. The protein was expressed in *E. coli*. Due to the poor solubility of

10 the expressed full-length protein (Met 1-Gln 422), an N-terminally truncated protein starting at Ser residue at position 40 (Ser 40) was produced. This truncation corresponds to Ser 2 of JNK1 and JNK2 proteins, and is preceded by a methionine

15 (initiation) and a glycine residue. The glycine residue was added in order to introduce an NcoI site for cloning into the expression vector. In addition, systematic C-terminal truncations were performed by PCR to identify a construct that give rise to

20 diffraction-quality crystals. One such construct encodes amino acid residues Ser40-Glu402 of JNK3 α 1 and is preceded by Met and Gly residues.

The construct was prepared by PCR using deoxyoligonucleotides:

25 5' GCTCTAGAGCTCCATGGGCAGCAAAAGCAAAGTTGACAA 3'
(forward primer with initiation codon underlined) (SEQ ID NO:1) and

5' TAGCGGATCCTCATTCTGAATTCATTACTTCCTTGTA 3' (reverse primer with stop codon underlined) (SEQ ID NO:2) as

30 primers and was confirmed by DNA sequencing.

Control experiments indicated that the truncated JNK3 protein had an equivalent kinase activity

-74-

towards myelin basic protein when activated with an upstream kinase MKK7 *in vitro*.

E. coli strain BL21 (DE3) (Novagen) was transformed with the JNK3 expression construct and
5 grown at 30°C in LB supplemented with 100 µg/ml carbenicillin in shaker flasks until the cells were in log phase (OD₆₀₀ ~ 0.8). Isopropylthio-β-D-galactosidase (IPTG) was added to a final concentration of 0.8 mM and the cells were harvested
10 2 hours later by centrifugation.

E. coli cell paste containing JNK3 was resuspended in 10 volumes/g lysis buffer (50 mM HEPES, pH 7.2, containing 10% glycerol (v/v), 100 mM NaCl, 2 mM DTT, 0.1 mM PMSF, 2 µg/ml Pepstatin,
15 1µg/ml each of E-64 and Leupeptin). Cells were lysed on ice using a microfluidizer and centrifuged at 100,000 x g for 30 min at 4 °C. The 100,000 x g supernatant was diluted 1:5 with Buffer A (20 mM HEPES, pH 7.0, 10% glycerol (v/v), 2 mM DTT) and
20 purified by SP-Sepharose (Pharmacia) cation-exchange chromatography (column dimensions: 2.6 x 20 cm) at 4 °C. The resin was washed with 5 column volumes of Buffer A, followed by 5 column volumes of Buffer A containing 50 mM NaCl. Bound JNK3 was eluted with a
25 7.5 column volume linear gradient of 50-300 mM NaCl. JNK3 eluted between 150-200 mM NaCl.

Example 39

Activation of JNK3

30 5 mg of JNK3 was diluted to 0.5 mg/ml in 50 mM HEPES buffer, pH 7.5, containing 100 mM NaCl, 5 mM DTT, 20 mM MgCl₂ and 1 mM ATP. GST-MKK7(DD)

-75-

was added at a molar ratio of 1:2.5 GST-MKK7:JNK3. After incubation for 30 minutes at 25°C, the reaction mixture was concentrated 5-fold by ultrafiltration in a Centriprep-30 (Amicon, Beverly, MA), diluted to 10 ml and an additional 1 mM ATP added. This procedure was repeated three times to remove ADP and replenish ATP. The final addition of ATP was 5 mM and the mixture incubated overnight at 4°C.

The activated JNK3/GST-MKK7(DD) reaction mixture was exchanged into 50 mM HEPES buffer, pH 7.5, containing 5 mM DTT and 5% glycerol (w/v) by dialysis or ultrafiltration. The reaction mixture was adjusted to 1.1 M potassium phosphate, pH 7.5, and purified by hydrophobic interaction chromatography (at 25 °C) using a Rainin Hydropore column. GST-MKK7 and unactivated JNK3 do not bind under these conditions such that when a 1.1 to 0.05 M potassium phosphate gradient is developed over 60 minutes at a flow rate of 1 ml/minute, doubly phosphorylated JNK3 is separated from singly phosphorylated JNK. Activated JNK3 (i.e. doubly phosphorylated JNK3) was stored at -70°C at 0.25-1 mg/ml.

25

Example 40

JNK Inhibition Assays

Compounds were assayed for the inhibition of JNK3 by a spectrophotometric coupled-enzyme assay. In this assay, a fixed concentration of activated JNK3 (10 nM) was incubated with various concentrations of a potential inhibitor dissolved in DMSO for 10 minutes at 30°C in a buffer containing

-76-

0.1 M HEPES buffer, pH 7.5, containing 10 mM MgCl₂, 2.5 mM phosphoenolpyruvate, 200 μ M NADH, 150 μ g/mL pyruvate kinase, 50 μ g/mL lactate dehydrogenase, and 200 μ M EGF receptor peptide. The EGF receptor peptide has the sequence KRELVEPLTPSGEAPNQALLR, and is a phosphoryl acceptor in the JNK3-catalyzed kinase reaction. The reaction was initiated by the addition of 10 μ M ATP and the assay plate is inserted into the spectrophotometer's assay plate compartment that was maintained at 30°C. The decrease of absorbance at 340 nm was monitored as a function of time. The rate data as a function of inhibitor concentration was fitted to competitive inhibition kinetic model to determine the K_i.

For selected compounds of this invention, activity in the JNK inhibition assay is shown in Table 8. Compounds having a K_i less than 0.1 micromolar (μ M) are rated "A", compounds having a K_i between 0.1 and 1 μ M are rated "B" and compounds having a K_i greater than 1 μ M are rated "C".

Table 8. Activity in the JNK3 Inhibition Assay.

No.	Activity	No.	Activity	No.	Activity
IIA-1	A	IIA-2	-	IIA-3	A
IIA-4	-	IIA-5	A	IIA-6	A
IIA-7	A	IIA-8	A/B	IIA-9	B
IIA-10	B	IIA-11	A	IIA-12	B/C
IIA-13	C	IIA-14	B	IIA-15	B
IIA-16	-	IIA-17	-	IIA-18	-
IIA-19	-	IIA-20	-	IIA-21	-
IIA-22	-	IIA-23	-	IIA-24	-
IIA-25	-	IIA-26	-	IIA-27	-
IIA-28	-	IIA-29	-	IIA-30	-
IIA-31	-	IIA-32	A	IIA-33	A
IIA-34	A	IIA-35	A	IIA-36	A
IIA-37	A	IIA-38	A	IIA-39	A

-77-

No.	Activity	No.	Activity	No.	Activity
IIA-40	A	IIA-41	A	IIA-42	A
IIA-43	A	IIA-44	A	IIA-45	A
IIA-46	A	IIA-47	A	IIA-48	A
IIA-49	A	IIA-50	A	IIA-51	A
IIA-52	A	IIA-53	A	IIA-54	A
IIA-55	A	IIA-56	A	IIA-57	A
IIA-58	A	IIA-59	A	IIA-60	A
IIA-61	A	IIA-62	A	IIA-63	A
IIA-64	A	IIA-65	A	IIA-66	A
IIA-67	A	IIA-68	A	IIA-69	A
IIA-70	A/B	IIA-71	A/B	IIA-72	A/B
IIA-73	B	IIA-74	B	IIA-75	B
IIA-76	B	IIA-77	B	IIA-78	B
IIA-79	B	IIA-80	B	IIA-81	B
IIA-82	B	IIA-83	B	IIA-84	B
IIA-85	C	IIA-86	C	IIA-87	C
IIA-88	-	IIA-89	-	IIA-90	A
IIA-91	A	IIA-92	A	IIA-93	A
IIA-94	A	IIA-95	A	IIA-96	A
IIA-97	A	IIA-98	A	IIA-99	A
IIA-100	A	IIA-101	A	IIA-102	A
IIA-103	A	IIA-104	A	IIA-105	A
IIA-106	B	IIA-107	C	IIA-108	A
IIA-109	A	IIA-110	C	IIA-111	C
IIA-112	C	IIA-113	B	IIA-114	B
IIA-115	B	IIA-116	C	IIA-117	B
IIA-118	B	IIA-119	B	IIA-120	B
IIA-121	C	IIA-122	B	IIA-123	B
IIA-124	B	IIA-125	B	IIA-126	B
IIA-127	B	IIA-128	B	IIA-129	B
IIA-130	A	IIA-131	A	IIA-132	A
IIA-133	A	IIA-134	A	IIA-135	B
IIA-136	-	IIA-137	-	IIA-138	-
IIAA-1	-	IIAA-2	-	IIAA-3	-
IIAA-4	B	IIAA-5	-	IIAA-6	-
IIAA-7	-	IIAA-8	-	IIAA-9	-
IIAA-10	A	IIAA-11	A	IIAA-12	A
IIAA-13	A	IIAA-14	A	IIAA-15	B
IIAA-16	A	IIAA-17	C	IIAA-18	B
IIAA-19	A	IIAA-20	B	IIAA-21	B
IIAA-22	B	IIAA-23	B	IIAA-24	A
IIAA-25	A	IIAA-26	C	IIAA-27	B
IIAA-28	C	IIAA-29	B	IIAA-30	C
IIAA-31	A	IIAA-32	B	IIAA-33	A
IIAA-34	A	IIAA-35	A	IIAA-36	A

-78-

No.	Activity	No.	Activity	No.	Activity
IIAA-37	A	IIAA-38	A	IIAA-39	B
IIIA-1	B	IIIA-2	C	IIIA-3	B
IIIA-4	C	IIIA-5	C	IIIA-6	B
IIIA-7	B	IIIA-8	B	IIIA-9	C
IIIA-10	C	IIIA-11	B	IIIA-12	B
IIIA-13	-	IIIA-14	B	IIIA-15	A
IIIA-16	-	IIIA-17	-	IIIA-18	B
IIIA-19	B	IIIA-20	B	IIIA-21	B
IIIA-22	C	IIIA-23	C	IIIA-24	C
IIIA-25	C	IIIA-26	C	IIIA-27	C
IIIA-28	C	IIIA-29	C	IIIA-30	B
IIIA-31	B	IIIA-32	B	IIIA-33	B
IIIA-34	C	IIIA-35	C	IIIA-36	C
IIIA-37	C	IIIA-38	C	IIIA-39	C
IIIA-40	C	IIIA-41	C	IIIA-42	B
IIIA-43	A	IIIA-44	B	IIIA-45	B
IIIA-46	B	IIIA-47	B	IIIA-48	B
IIIA-49	B	IIIA-50	B	IIIA-51	B
IIIA-52	B	IIIA-53	B	IIIA-54	B
IIIA-55	B	IIIA-56	B	IIIA-57	B
IIIA-58	B	IIIA-59	B	IIIA-60	B
IIIA-61	B	IIIA-62	B	IIIA-63	B
IIIA-64	B	IIIA-65	B	IIIA-66	B
IIIA-67	B	IIIA-68	B	IIIA-69	B
IIIA-70	B	IIIA-71	B	IIIA-72	B
IIIA-73	B	IIIA-74	A	IIIA-75	B
IIIA-76	-	IIIA-77	-	IIIA-78	-
IIIA-79	-	IIIA-80	-	IIIA-81	-
IIIA-82	-	IIIA-83	-	IIIA-84	-
IIIA-85	-	IIIA-86	-	IIIA-87	-
IIIA-88	-	IIIA-89	-	IIIA-90	-
IIIA-91	-	IIIA-92	-	IIIA-93	-
IIIA-94	-	IIIA-95	-	IIIA-96	-
IIIA-97	-				
XA-1	B	XA-2	C	XA-3	B
XA-4	B	XA-5	B	XA-6	-
XIA-1	-	XIA-2	-	XIA-3	-
XIA-4	-	XIA-5	-	XIA-6	-
XIA-7	-	XIA-8	-	XIA-9	-
XIA-10	-	XIA-11	-	XIA-12	-
XIA-13	-	XIA-14	-	XIA-15	-
XIA-16	-	XIA-17	-	XIA-18	-
XIA-19	-	XIA-20	-	XIA-21	-
XIA-22	-	XIA-23	-	XIA-24	-
XIA-25	-	XIA-26	-	XIA-27	-

-79-

No.	Activity	No.	Activity	No.	Activity
XIA-28	-	XIA-29	-	XIA-30	-
XIA-31	-	XIA-32	-	XIA-33	-
XIA-34	-	XIA-35	-	XIA-36	-
XIA-37	-	XIA-38	-	XIA-39	-
XIA-40	-	XIA-41	-	XIA-42	-
XIA-43	-	XIA-44	-	XIA-45	A
XIA-46	A	XIA-47	A	XIA-48	A
XIA-49	A	XIA-50	A	XIA-51	A
XIA-52	A	XIA-53	A		

Example 41Src Inhibition Assays

5 The compounds were assayed as inhibitors
 of full length recombinant human Src kinase (from
 Upstate Biotechnology, cat. no. 14-117) expressed
 and purified from baculo viral cells. Src kinase
 activity was monitored by following the
 10 incorporation of ^{33}P from ATP into the tyrosine of a
 random poly Glu-Tyr polymer substrate of
 composition, Glu:Tyr = 4:1 (Sigma, cat. no. P-0275).
 The following were the final concentrations of the
 assay components: 0.05 M HEPES, pH 7.6, 10 mM MgCl_2 ,
 15 2 mM DTT, 0.25 mg/ml BSA, 10 μM ATP (1-2 μCi ^{33}P -ATP
 per reaction), 5 mg/ml poly Glu-Tyr, and 1-2 units
 of recombinant human Src kinase. In a typical
 assay, all the reaction components with the
 exception of ATP were pre-mixed and aliquoted into
 20 assay plate wells. Inhibitors dissolved in DMSO
 were added to the wells to give a final DMSO
 concentration of 2.5%. The assay plate was
 incubated at 30 °C for 10 min before initiating the
 reaction with ^{33}P -ATP. After 20 min of reaction, the
 25 reactions were quenched with 150 μl of 10%
 trichloroacetic acid (TCA) containing 20 mM Na_3PO_4 .

-80-

The quenched samples were then transferred to a 96-well filter plate (Whatman, UNI-Filter GF/F Glass Fiber Filter, cat no. 7700-3310) installed on a filter plate vacuum manifold. Filter plates were washed four times with 10% TCA containing 20 mM Na_3PO_4 and then 4 times with methanol. 200 μl of scintillation fluid was then added to each well. The plates were sealed and the amount of radioactivity associated with the filters was quantified on a TopCount scintillation counter.

The most active compounds in the Src assay were found to be those compounds of formula **I** where G is an optionally substituted aryl and R^1 is Ar^2 .

15

Example 42Lck Inhibition Assays

The compounds were assayed as inhibitors of lck kinase purified from bovine thymus (from Upstate Biotechnology, cat. no. 14-106). Lck kinase activity was monitored by following the incorporation of ^{33}P from ATP into the tyrosine of a random poly Glu-Tyr polymer substrate of composition, Glu:Tyr = 4:1 (Sigma, cat. no. P-0275). The following were the final concentrations of the assay components: 0.05 M HEPES, pH 7.6, 10 mM MgCl_2 , 2 mM DTT, 0.25 mg/ml BSA, 10 μM ATP (1-2 μCi ^{33}P -ATP per reaction), 5 mg/ml poly Glu-Tyr, and 1-2 units of lck kinase. In a typical assay, all the reaction components with the exception of ATP were pre-mixed and aliquoted into assay plate wells. Inhibitors dissolved in DMSO were added to the wells to give a final DMSO concentration of 2.5%. The assay plate

-81-

was incubated at 30 °C for 10 min before initiating the reaction with ^{33}P -ATP. After 20 min of reaction, the reactions were quenched with 150 μl of 10% trichloroacetic acid (TCA) containing 20 mM Na_3PO_4 .

5 The quenched samples were then transferred to a 96-well filter plate (Whatman, UNI-Filter GF/F Glass Fiber Filter, cat no. 7700-3310) installed on a filter plate vacuum manifold. Filter plates were washed four times with 10% TCA containing 20 mM
10 Na_3PO_4 and then 4 times with methanol. 200 μl of scintillation fluid was then added to each well. The plates were sealed and the amount of radioactivity associated with the filters was quantified on a TopCount scintillation counter.

15 The most active compounds in the Lck assay were found to be those compounds of formula **I** where G is an optionally substituted aryl and R^1 is Ar^2 .


While we have described a number of
embodiments of this invention, it is apparent that
20 our basic examples may be altered to provide other embodiments which utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the
25 specific embodiments which have been represented by way of example.

79580-10

- 82 -

CLAIMS:

1. A compound having the formula:



5 wherein:

X-Y-Z is selected from one of the following:



R^1 is H, CONH_2 , $\text{T}_{(n)}-\text{R}$, or $\text{T}_{(n)}-\text{Ar}^2$;

R is an optionally substituted C_{1-12} aliphatic;

10 n is zero or one;

T is $\text{C}(=\text{O})$, CO_2 , CONH , $\text{S}(\text{O})_2$, $\text{S}(\text{O})_2\text{NH}$, COCH_2 or CH_2 ;

each R^2 is independently selected from hydrogen, -R, $-\text{CH}_2\text{OR}$, $-\text{CH}_2\text{OH}$, $-\text{CH}=\text{O}$, $-\text{CH}_2\text{SR}$, $-\text{CH}_2\text{S}(\text{O})_2\text{R}$, $-\text{CH}_2\text{C}(=\text{O})\text{R}$, $-\text{CH}_2\text{CO}_2\text{R}$, $-\text{CH}_2\text{CO}_2\text{H}$, $-\text{CH}_2\text{CN}$, $-\text{CH}_2\text{NHR}$, $-\text{CH}_2\text{N}(\text{R})_2$, $-\text{CH}=\text{N}-\text{OR}$, $-\text{CH}=\text{NNHR}$,
 15 $-\text{CH}=\text{NN}(\text{R})_2$, $-\text{CH}=\text{NNHCOR}$, $-\text{CH}=\text{NNHCO}_2\text{R}$, $-\text{CH}=\text{NNHSO}_2\text{R}$, optionally substituted 6 to 14-membered aryl, optionally substituted 6 to 14-membered $\text{CH}_2(\text{aryl})$, $-\text{CH}_2\text{NH}_2$, $-\text{CH}_2\text{NHCOR}$, $-\text{CH}_2\text{NHCONHR}$, $-\text{CH}_2\text{NHCON}(\text{R})_2$, $-\text{CH}_2\text{NRCOR}$, $-\text{CH}_2\text{NHCO}_2\text{R}$, $-\text{CH}_2\text{CONHR}$, $-\text{CH}_2\text{CON}(\text{R})_2$, $-\text{CH}_2\text{SO}_2\text{NH}_2$, optionally substituted 5 to 8-membered

20 $-\text{CH}_2(\text{heterocyclyl})$ comprising one or more heteroatoms selected from O, N and S, and optionally substituted 5 to 8-membered $-(\text{heterocyclyl})$ comprising one or more ring heteroatoms selected from O, N and S;

each R^3 is independently selected from hydrogen, R, COR, CO_2R
 25 and $\text{S}(\text{O})_2\text{R}$;

79580-10

- 83 -


G is R or Ar¹;

Ar¹ is an optionally substituted 6 to 14-membered aryl or aralkyl, or an optionally substituted 5 to 8-membered heterocyclyl containing one or more ring heteroatoms

5 selected from N, O and S, wherein Ar¹ is optionally fused to a partially unsaturated or fully unsaturated five to seven membered ring comprising zero to three heteroatoms selected from O, N and S;

Q-NH is

10



wherein the H of Q-NH is optionally replaced by R³;

A is N or CR³;

Ar² is an optionally mono- or poly-substituted 6 to 14-membered aryl or a 5 to 8-membered heterocyclyl

15 comprising one or more heteroatoms selected from O, N and S, wherein Ar² optionally comprises an optionally mono- or poly-substituted, partially unsaturated or fully unsaturated five to seven membered ring comprising zero to three heteroatoms selected from O, N and S fused to the aryl or
20 the heterocyclyl;

wherein the one or more substituents on carbon atoms in Ar² are independently selected from halo, R, OR, SR, OH, NO₂, CN, NH₂, NHR, N(R)₂, NHCOR, NHCONHR, NHCON(R)₂, NRCOR, NHCO₂R, CO₂R, CO₂H, COR, CONHR, CON(R)₂, S(O)₂R, SONH₂, S(O)R, SO₂NHR, 25 and NHS(O)₂R, and wherein the one or more substituents on saturated carbon atoms in the optional, fused ring in Ar² are further independently selected from =O, =S, =NNHR, =NNR₂, =N-OR, =NNHCOR, =NNHCO₂R, =NNHSO₂R, and =NR; and

79580-10

- 84 -

wherein the one or more substituents on nitrogen atoms in Ar^2 are independently selected from R, COR, $\text{S}(\text{O})_2\text{R}$, and CO_2R .

2. The compound of claim 1 where G is Ar^1 .

3. The compound of claim 2 where Q-NH is selected
5 from:




4. The compound of claim 3 where R^1 is $-(\text{C}_{1-12} \text{ alkyl})-$, $\text{O}-(\text{C}_{1-12} \text{ alkyl})-$, $-(\text{C}_{1-12} \text{ alkyl})-(\text{CO})-\text{O}-(\text{C}_{1-12} \text{ alkyl})-$, $-(\text{C}_{1-12} \text{ alkyl})(\text{OH})$, $(\text{C}_{1-12} \text{ alkyl})\text{pyridinyl}$, $-(\text{C}_{1-12} \text{ cycloalkyl})-\text{O}-(\text{C}_{1-12} \text{ alkyl})$, $\text{C}_{1-12} \text{ cycloalkyl}$, $-(\text{C}_{1-12} \text{ cycloalkyl})-(\text{CO})-\text{O}-(\text{C}_{1-12} \text{ alkyl})$, $(\text{C}_{1-12} \text{ cycloalkyl})(\text{OH})$, Ar^2 or $\text{T}-\text{Ar}^2$ where T is $\text{C}(=\text{O})$.
10

5. The compound of claim 4 where R^1 is cyclohexyl, cyclohexanol-4-yl, cyclohexanon-4-yl, 2-propan-1-ol, 2-methoxy-1-methylethyl, 3-butyryl alkyl ester, 2-pyridinyl-2-ethyl, or an optionally substituted phenyl, naphthyl, pyridyl, quinolinyl, thienyl or indanyl.
15

6. The compound of claim 5 where R^2 is an optionally substituted C_{1-12} alkyl.

7. A compound according to claim 1 selected from one
20 of the following compounds:


a) compounds of formula IIA:




IIA

where Q is:

79580-10



and Ar² is  ; wherein compounds are further defined as:

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-1	Ph	Q1	Me	H	H	H	H	H
IIA-2	Ph	Q1	Me	H	H	OMe	H	H
IIA-3	Ph	Q1	Me	H	OMe	OMe	H	H
IIA-4	Ph	Q1	Me	Me	H	H	H	H
IIA-5	Ph	Q1	Me	Me	H	CONH ₂	H	H
IIA-6	Ph	Q1	Me	Me	H	CN	H	H
IIA-7	Ph	Q1	Me	H	CN	H	H	H
IIA-8	Ph	Q1	Me	Me	F	H	H	H
IIA-9	Ph	Q1	Me	Me	H	F	H	H
IIA-10	Ph	Q1	Me	CF ₃	H	H	H	H
IIA-11	4-F-Ph	Q1	Me	H	H	H	H	H
IIA-12	2,3-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-13	2,4-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-14	2-Cl-Ph	Q1	Me	H	H	H	H	H
IIA-15	3,4-Cl ₂ -Ph	Q1	Me	H	H	H	H	H

79580-10

- 86 -

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-16	Ph	Q2	Et	H	CN	H	H	H
IIA-17	Ph	Q2	Et	H	CO ₂ H	H	H	H
IIA-18	Ph	Q2	Me	H	F	H	H	H
IIA-19	Ph	Q2	Me	H	H	F	H	H
IIA-20	Ph	Q2	Me	H	H	COMe	H	H
IIA-21	Ph	Q2	Me	H	H	COPh	H	H
IIA-22	Et	Q1	Me	H	H	H	H	H
IIA-23	PhCH ₂ OC H ₂ -	Q1	Me	H	H	H	H	H
IIA-24	Ph	Q2	Me	H	H	CONH ₂	H	H
IIA-25	3-F-Ph	Q1	Me	H	CN	H	H	H
IIA-26	3-F-Ph	Q1	Me	H	H	CN	H	H
IIA-27	3-F-Ph	Q1	Me	H	F	H	H	H
IIA-28	3-F-Ph	Q1	Me	H	H	F	H	H
IIA-29	3-F-Ph	Q1	Me	H	Me	CN	H	H
IIA-30	3-F-Ph	Q1	Me	H	F	CN	H	H
IIA-31	3-F-Ph	Q1	Me	H	H	SMe	H	H
IIA-32	Ph	Q1	Me	H	F	CN	H	H
IIA-33	Ph	Q1	Me	H	F	H	H	H
IIA-34	Ph	Q1	Me	H	H	CN	H	H
IIA-35	Ph	Q1	Me	H	H	COMe	H	H
IIA-36	Ph	Q1	Me	H	CH=CH	H	H	H
IIA-37	Ph	Q1	Me	H	SMe	H	H	H
IIA-38	Ph	Q1	Me	H	Me	CN	H	H

79580-10

- 87 -

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-39	Ph	Q1	Me	H	COMe	H	H	H
IIA-40	Ph	Q2	Et	H	H	H	H	H
IIA-41	Ph	Q1	Me	OMe	H	H	H	H
IIA-42	Ph	Q1	Me	H	H	F	H	H
IIA-43	Ph	Q2	Me	H	CO ₂ H	H	H	H
IIA-44	Ph	Q1	Me	H	H	Ph	H	H
IIA-45	Ph	Q1	Me	H	Me	H	Me	H
IIA-46	Ph	Q1	Me	H	H	SMe	H	H
IIA-47	Ph	Q2	Me	H	H	OMe	H	H
IIA-48	Ph	Q2	Me	H	OMe	H	H	H
IIA-49	Ph	Q1	Me	OMe	H	H	CN	H
IIA-50	Ph	Q2	Me	H	CO ₂ Me	H	H	H
IIA-51	Ph	Q1	Me	F	H	H	CN	H
IIA-52	Ph	Q2	Me	H	H	H	H	H
IIA-53	Ph	Q2	Me	H	H	CO ₂ H	H	H
IIA-54	Ph	Q1	Me	Me	H	CN	H	H
IIA-55	2-F-Ph	Q1	Me	H	H	H	H	H
IIA-56	Ph	Q1	Me	F	H	F	H	H
IIA-57	Ph	Q1	Me	Me	H	CONH ₂	H	H
IIA-58	Ph	Q1	Me	Me	Cl	H	H	H
IIA-59	Ph	Q1	Me	F	H	H	H	H
IIA-60	2,6-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-61	Ph	Q1	Me	Me	H	OMe	H	H

79580-10

- 88 -

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-62	Ph	Q1	Me	OMe	H	H	H	H
IIA-63	Ph	Q1	Me	H	H	SO ₂ Me	H	H
IIA-64	Ph	Q2	Me	H	H	CO ₂ Me	H	H
IIA-65	Ph	Q1	Me	NO ₂	H	H	H	H
IIA-66	3-F-Ph	Q1	Me	H	H	H	H	H
IIA-67	Ph	Q2	Me	H	CN	H	H	H
IIA-68	Ph	Q2	Me	H	H	CN	H	H
IIA-69	Ph	Q1	Me	CH:CH	H	H	H	H
IIA-70	Ph	Q1	Me	Me	F	H	H	H
IIA-71	Ph	Q1	Me	Cl	H	H	OMe	H
IIA-72	Ph	Q1	Me	H	Me	OMe	H	H
IIA-73	Ph	Q1	Me	OMe	H	H	OMe	H
IIA-74	2,5-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-75	2-Cl-6-F-Ph	Q1	Me	H	H	H	H	H
IIA-76	2-Cl-Ph	Q1	Me	H	H	H	H	H
IIA-77	3,4-Cl ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-78	Ph	Q1	Me	Me	H	F	H	H
IIA-79	2-Br-Ph	Q1	Me	H	H	H	H	H
IIA-80	2,3-F ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-81	Ph	Q1	Me	SMe	H	H	H	H
IIA-82	3-CF ₃ -Ph	Q1	Me	H	H	H	H	H
IIA-83	3,5-F ₂ -Ph	Q1	Me	H	H	H	H	H



79580-10

- 89 -

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-84	2,6-Cl ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-85	2,3-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-86	Me	Q1	Me	H	H	H	H	H
IIA-87	cyclopropyl	Q1	Me	H	H	H	H	H
IIA-88	cyclohexyl	Q1	Me	H	H	H	H	H
IIA-89	2,4-(MeO) ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-90	t-butyl	Q1	Me	H	H	H	H	H
IIA-91	2,6-F ₂ -Ph	Q1	Me	H	H	COMe	H	H
IIA-92	2,6-F ₂ -Ph	Q1	Me	H	CN	H	H	H
IIA-93	2,6-F ₂ -Ph	Q1	Me	H	H	CN	H	H
IIA-94	2,6-F ₂ -Ph	Q1	Me	H	F	H	H	H
IIA-95	2,6-F ₂ -Ph	Q1	Me	H	H	F	H	H
IIA-96	2,6-F ₂ -Ph	Q1	Me	H	CN	F	H	H
IIA-97	2,6-F ₂ -Ph	Q1	Me	H	H	SMe	H	H
IIA-98	Ph	Q2	Me	H	H	NMe ₂	H	H
IIA-99	Ph	Q2	Me	H	NO ₂	H	H	H
IIA-100	Ph	Q2	Me	H	NHAc	H	H	H
IIA-101	Ph	Q2	Me	H	NH ₂	H	H	H
IIA-102	Ph	Q1	Me	H	Me	H	H	H
IIA-103	Ph	Q1	Me	H	H	Me	H	H
IIA-104	2-Me-Ph	Q1	Me	H	H	H	H	H
IIA-105	2-Me-Ph	Q1	Me	H	F	CN	H	H




79580-10

- 90 -


No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-106	2-Me-Ph	Q1	Me	H	F	H	H	H
IIA-107	2-Me-Ph	Q1	Me	H	H	CN	H	H
IIA-108	2-Me-Ph	Q1	Me	H	Me	H	H	H
IIA-109	2-Me-Ph	Q1	Me	H	CN	H	H	H
IIA-110	2-CF ₃ -Ph	Q1	Me	H	F	CN	H	H
IIA-111	2-CF ₃ -Ph	Q1	Me	H	CN	H	H	H
IIA-112	2-CF ₃ -Ph	Q1	Me	H	H	H	H	H
IIA-113	3,4- (OCH ₂ O)-Ph	Q1	Me	H	F	CN	H	H
IIA-114	3,4- (OCH ₂ O)-Ph	Q1	Me	H	CN	H	H	H
IIA-115	3,4- (OCH ₂ O)-Ph	Q1	Me	H	H	H	H	H
IIA-116	 3,4-(OCH ₂ O)	Q1	Me	 bis-N,N'-4-cyanophenyl				
IIA-117	3-OBn-Ph	Q1	Me	H	F	CN	H	H
IIA-118	3-OBn-Ph	Q1	Me	H	CN	H	H	H
IIA-119	3-OBn-Ph	Q1	Me	H	H	H	H	H
IIA-120	3-NO ₂ -Ph	Q1	Me	H	F	CN	H	H
IIA-121	3-NO ₂ -Ph	Q1	Me	bis-N,N'-4-cyanophenyl				
IIA-122	3-NO ₂ -Ph	Q1	Me	H	CN	H	H	H
IIA-123	3-NO ₂ -Ph	Q1	Me	H	H	H	H	H
IIA-124	3-CN-Ph	Q1	Me	H	F	CN	H	H
IIA-125	3-CN-Ph	Q1	Me	H	H	CN	H	H

79580-10


- 91 -

No.	G	Q	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
IIA-126	3-CN-Ph	Q1	Me	H	CN	H	H	H
IIA-127	3-CN-Ph	Q1	Me	H	H	H	H	H
IIA-128	3-NO ₂ -Ph	Q1	Me	H	H	CO ₂ Et	H	H
IIA-129	3-CN-Ph	Q1	Me	H	CO ₂ Me	H	H	H
IIA-130	Ph	Q1	Me	H	CO ₂ Et	H	H	H
IIA-131	Ph	Q1	Me	N	H	NO ₂	H	H
IIA-132	Ph	Q2	Me					
IIA-133	Ph	Q2	Me					
IIA-134	Ph	Q2	Me	H	CH ₂ OH	H	H	H
IIA-135	Ph	Q2	Me					

b) compounds having the formula:













**IIA**

5 wherein compounds are further defined as:

No.	G	A	R ¹
IIAA-1	Ph	CH	













79580-10

- 92 -

No.	G	A	R ¹
IIAA-2	Ph	CH	
IIAA-3	Ph	N	
IIAA-4	Ph	N	
IIAA-5	Ph	N	
IIAA-6	Ph	N	
IIAA-7	Ph	N	
IIAA-8	Ph	N	
IIAA-9	3-F-Ph	N	
IIAA-10	Ph	N	
IIAA-11	Ph	N	
IIAA-12	Ph	N	
IIAA-13	Ph	N	












79580-10

- 93 -

No.	G	A	R ¹
IIAA-14	2,6-F ₂ -Ph	N	
IIAA-15	Ph	N	
IIAA-16	Ph	N	
IIAA-17	Ph	N	
IIAA-18	Ph	N	
IIAA-19	2-Me-Ph	N	
IIAA-20	2-Me-Ph	N	
IIAA-21		N	
IIAA-22	3-NO ₂ -Ph	N	
IIAA-23	3-CN-Ph	N	
IIAA-24	Ph	N	






79580-10

- 94 -


No .	G	A	R ¹
IIAA-25	Ph	N	
IIAA-26	Ph	N	
IIAA-27	Ph	N	
IIAA-28	Ph	N	
IIAA-29	Ph	N	
IIAA-30	Ph	N	
IIAA-31	Ph	N	
IIAA-32	Ph	N	
IIAA-33	Ph	N	
IIAA-34	Ph	N	
IIAA-35	Ph	N	

79580-10


- 95 -

No.	G	A	R ¹
IIAA-36	Ph	N	
IIAA-37	Ph	N	
IIAA-38	Ph	N	
IIAA-39	Ph	CH	
IIAA-40	Ph	CH	


c) compounds having one of the formulas:



wherein Q is one of the following groups:



Q1



Q2;

79580-10

- 96 -

wherein compounds are further defined as:

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-1	phenyl	Q1	H	H	H	H	H
IIIA-2	phenyl	Q1	Br	H	H	H	H
IIIA-3	phenyl	Q1	F	H	H	H	H
IIIA-4	phenyl	Q1	Cl	H	H	H	H
IIIA-5	phenyl	Q1	CH ₃	H	H	H	H
IIIA-6	phenyl	Q1	H	CH ₃	H	H	H
IIIA-7	phenyl	Q1	H	H	OCH ₃	H	H
IIIA-8	phenyl	Q1	H	OCH ₃	OCH ₃	H	H
IIIA-9	phenyl	Q1	OCH ₃	H	OCH ₃	H	H
IIIA-10	phenyl	Q1	OCH ₃	H	H	H	OCH ₃
IIIA-11	phenyl	Q1	H	H	CN	H	H
IIIA-12	5-fluorophenyl	Q1	H	H	OCH ₃	H	H
IIIA-13	phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-14	phenyl	Q1	H	H	F	H	H
IIIA-15	phenyl	Q1	Ar ² is 2-thienyl				
IIIA-16	phenyl	Q1	Ar ² is 1-oxo-indan-5-yl				
IIIA-17	phenyl	Q1	Ar ² is 4-pyridyl				
IIIA-18	2-CH ₃ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-19	2-CH ₃ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-20	2-CH ₃ -phenyl	Q1	H	H	OCH ₃	H	H

79580-10

- 97 -

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-21	2-CH ₃ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-22	2-CF ₃ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-23	2-CF ₃ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-24	2-CF ₃ -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-25	2-CF ₃ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-26	benzo[3,5]dioxole	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-27	benzo[3,5]dioxole	Q1	H	OCH ₃	H	H	H
IIIA-28	benzo[3,5]dioxole	Q1	H	H	OCH ₃	H	H
IIIA-29	benzo[3,5]dioxole	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-30	3-benzyloxy-phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-31	3-benzyloxy-phenyl	Q1	H	OCH ₃	H	H	H
IIIA-32	3-benzyloxy-phenyl	Q1	H	H	OCH ₃	H	H
IIIA-33	3-benzyloxy-phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-34	3-nitrophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-35	3-nitrophenyl	Q1	H	OCH ₃	H	H	H
IIIA-36	3-nitrophenyl	Q1	H	H	OCH ₃	H	H
IIIA-37	3-nitrophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-38	3-cyanophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-39	3-cyanophenyl	Q1	H	OCH ₃	H	H	H
IIIA-40	3-cyanophenyl	Q1	H	H	OCH ₃	H	H
IIIA-41	3-cyanophenyl	Q1	H	OCH ₃	H	OCH ₃	H

79580-10

- 98 -


No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-42	phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-43	phenyl	Q1	H	CN	H	H	H
IIIA-44	phenyl	Q1	H	H	CO ₂ Me	H	H
IIIA-45	3-fluorophenyl	Q1	H	Cl	H	H	H
IIIA-46	3-fluorophenyl	Q1	H	OCH ₃	H	H	H
IIIA-47	3-fluorophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-48	3-fluorophenyl	Q1	H	Me	H	H	H
IIIA-49	3-fluorophenyl	Q1	H	H	F	H	H
IIIA-50	3-fluorophenyl	Q1	H	H	Me	H	H
IIIA-51	3-fluorophenyl	Q1	H	CN	H	H	H
IIIA-52	3-fluorophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-53	3-fluorophenyl	Q1	Ar ² is 2-naphthyl				
IIIA-54	2-fluorophenyl	Q1	H	Cl	H	H	H
IIIA-55	2-fluorophenyl	Q1	H	OCH ₃	H	H	H
IIIA-56	2-fluorophenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-57	2-fluorophenyl	Q1	H	Me	H	H	H
IIIA-58	2-fluorophenyl	Q1	H	H	OCH ₃	H	H
IIIA-59	2-fluorophenyl	Q1	H	H	F	H	H
IIIA-60	2-fluorophenyl	Q1	H	H	Me	H	H
IIIA-61	2-fluorophenyl	Q1	H	CN	H	H	H
IIIA-62	2-fluorophenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H

79580-10

- 99 -

No.	Ar ¹	Q	Ar ²				
			R ²	R ³	R ⁴	R ⁵	R ⁶
IIIA-63	2-fluorophenyl	Q1	Ar ² is 2-naphthyl				
IIIA-64	2,6-F ₂ -phenyl	Q1	H	Cl	H	H	H
IIIA-65	2,6-F ₂ -phenyl	Q1	H	OCH ₃	H	H	H
IIIA-66	2,6-F ₂ -phenyl	Q1	H	OCH ₃	H	OCH ₃	H
IIIA-67	2,6-F ₂ -phenyl	Q1	H	Me	H	H	H
IIIA-68	2,6-F ₂ -phenyl	Q1	H	H	OCH ₃	H	H
IIIA-69	2,6-F ₂ -phenyl	Q1	H	H	F	H	H
IIIA-70	2,6-F ₂ -phenyl	Q1	H	H	Me	H	H
IIIA-71	2,6-F ₂ -phenyl	Q1	H	CN	H	H	H
IIIA-72	2,6-F ₂ -phenyl	Q1	H	OCH ₃	OCH ₃	OCH ₃	H
IIIA-73	2,6-F ₂ -phenyl	Q1	Ar ² is 2-naphthyl				
IIIA-74	phenyl	Q1	H	NO ₂	H	H	H
IIIA-75	phenyl	Q1	H	NHAc	H	H	H
IIIA-76	phenyl	Q1	H	COMe	H	H	H
IIIA-77	phenyl	Q2	H	COMe	H	H	H
IIIA-78	phenyl	Q2	H	CN	H	H	H

d) compounds having the formula:



XA


79580-10

- 100 -

wherein R^2 is CH_3 , and compounds are further defined as:

No.	Ar^1	R
XA-1	phenyl	CH_3
XA-2	4-F-phenyl	CH_3
XA-3	phenyl	cyclopentyl
XA-4	phenyl	isobutyl
XA-5	phenyl	propyl

e) compounds having the formula:



5

IXA

wherein compounds are defined as:

No.	Ar^1	A	R^1	R^2
XIA-1	phenyl	CH	phenyl	CH_2 (morpholin-4-yl)
XIA-2	phenyl	CH	phenyl	$CH_2N(CH_3)_2$
XIA-3	phenyl	CH	phenyl	CH_2NEt_2
XIA-4	phenyl	CH	phenyl	$CH_2N(CH_3)CH_2Ph$
XIA-5	phenyl	CH	phenyl	CH_2 (1-t-butoxycarbonylpiperazin-4-yl)
XIA-6	phenyl	CH	benzyl	CH_2 (morpholin-4-yl)
XIA-7	phenyl	CH	cyclohexyl	CH_2 (morpholin-4-yl)

79580-10

- 101 -

No.	Ar ¹	A	R ¹	R ²
XIA-8	phenyl	CH	4-[1,2-(OMe) ₂ -phenyl]	CH ₂ (morpholin-4-yl)
XIA-9	phenyl	CH	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-10	phenyl	CH	phenyl	CH ₂ N(CH ₃)CH ₂ CH ₂ N(CH ₃) ₂
XIA-11	phenyl	CH	phenyl	CH ₂ N(CH ₃)CH ₂ CO ₂ CH ₃
XIA-12	phenyl	CH	phenyl	CH ₂ (piperazin-1-yl)
XIA-13	phenyl	N	2-thienoyl	CH ₂ Br
XIA-14	phenyl	N	2-thienoyl	CH ₂ (morpholin-4-yl)
XIA-15	4-F-phenyl	CH	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-16	4-F-phenyl	CH	3-cyanophenyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-17	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-18	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-19	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-20	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-21	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-22	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ OCH ₂ CH ₂ OCH ₃
XIA-23	4-F-phenyl	CH	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-24	4-F-phenyl	CH	cyclohexyl	CH ₂ (morpholin-4-yl)
XIA-25	4-F-phenyl	CH	3-cyanophenyl	CH ₂ (morpholin-4-yl)
XIA-26	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ (morpholin-4-yl)
XIA-27	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ (morpholin-4-yl)


79580-10

- 102 -

No.	Ar ¹	A	R ¹	R ²
XIA-28	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₃
XIA-29	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₃
XIA-30	4-F-phenyl	CH	3-cyanophenyl	CH ₂ OCH ₃
XIA-31	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₃
XIA-32	4-F-phenyl	CH	1-benzyl-piperidin-4-yl	CH ₂ OCH ₃
XIA-33	4-F-phenyl	CH	4-cyclohexanol	CH ₂ OCH ₃
XIA-34	4-F-phenyl	CH	cyclohexyl	CH ₂ OCH ₃
XIA-35	4-F-phenyl	CH	3-cyanophenyl	CH ₂ OCH ₃
XIA-36	4-F-phenyl	CH	2-(2-pyridinyl)ethyl	CH ₂ OCH ₃
XIA-37	4-F-phenyl	CH	4-cyclohexanol	CH ₂ O(tetrahydrofuran-3-yl)
XIA-38	4-F-phenyl	CH	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-39	phenyl	N	2-thienoyl	CH ₂ (piperidin-1-yl)
XIA-40	phenyl	N	2-thienoyl	CH ₂ (piperazin-1-yl)
XIA-41	4-F-phenyl	CH	4-methoxybenzyl	CH ₂ OCH ₃
XIA-42	4-F-phenyl	N	4-cyclohexanol	CH ₂ (morpholin-4-yl)
XIA-43	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₃
XIA-44	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ (phenyl)
XIA-45	4-F-phenyl	N	cyclohexyl	CH ₂ OH
XIA-46	4-F-phenyl	N	CH ₂ CH ₂ (pyridin-2-yl)	CH ₂ OH
XIA-47	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₃
XIA-48	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₃
XIA-49	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ CH ₂ OCH ₃


79580-10

- 103 -


No.	Ar ¹	A	R ¹	R ²
XIA-50	4-F-phenyl	N	cyclohexyl	CH ₂ O(tetrahydrofuran-3-yl)
XIA-51	4-F-phenyl	N	cyclohexyl	CH ₂ O- 
XIA-52	4-F-phenyl	N	cyclohexyl	CH ₂ OCH ₂ (phenyl)
XIA-53	4-F-phenyl	N	CH ₂ CH ₂ (pyridin-2-yl)	CH ₂ OCH ₂ (phenyl)

and


f) compounds having one of the structures:



IB-1




IB-2




IB-3


5



IB-4



IB-5




IB-6


10

79580-10


- 104 -




IB-8




IB-9




IB-10




IB-11




IB-12




IB-13



IB-14







IB-15







IB-16

79580-10

- 105 -


**IB-17****IB-18****IB-19**

5

IB-20**IB-21****IB-22****IB-23****IB-24.**

and

8. A compound having the formula:

**XIII**

79580-10

- 106 -

wherein:

X-Y is N-O or O-N providing an isoxazole or reverse isoxazole ring;

A is N or CH;

5 G is R, an optionally substituted 6 to 14-membered aryl group;

R is an optionally substituted C₁₋₁₂ aliphatic;

R² is selected from hydrogen, -R, -CH₂OR, -CH₂OH, -CH=O, -CH₂SR, -CH₂S(O)₂R, -CH₂(C=O)R, -CH₂CO₂R, -CH₂CO₂H, -CH₂CN, 10 -CH₂NHR, -CH₂N(R)₂, -CH=N-OR, -CH=NNHR, -CH=NN(R)₂, -CH=NNHCOR, -CH=NNHCO₂R, -CH=NNHSO₂R, optionally substituted 6 to 14-membered aryl, optionally substituted 6 to 14-membered CH₂(aryl), -CH₂NH₂, -CH₂NHCOR, -CH₂NHCONHR, -CH₂NHCON(R)₂, -CH₂NRCOR, -CH₂NHCO₂R, -CH₂CONHR, -CH₂CON(R)₂, 15 -CH₂SO₂NH₂, optionally substituted 5 to 8-membered -CH₂(heterocyclyl) comprising one or more heteroatoms selected from O, N and S, and optionally substituted 5 to 8-membered-(heterocyclyl) comprising one or more ring heteroatoms selected from O, N and S; and 20 R¹ is selected from halogen, NH₂, SR, and SO₂R;

provided that:

when A is CH, X is O, Y is N, and G is 4-fluorophenyl, then R¹ is not F, Br or Cl.

9. A pharmaceutical composition comprising a compound 25 according to any one of claims 1-8, and a pharmaceutically acceptable carrier.

10. The pharmaceutical composition of claim 9 for treating a disease state or condition in a mammal that is

79580-10

- 107 -

alleviated by treatment with an inhibitor of c-Jun N-terminal kinases.

11. The pharmaceutical composition of claim 10 wherein the disease state or condition is selected from inflammatory
5 diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy,
10 thrombin-induced platelet aggregation and conditions associated with proinflammatory cytokines.

12. The pharmaceutical composition of claim 9 for treating an inflammatory disease selected from acute pancreatitis, chronic pancreatitis, asthma, allergies, and
15 adult respiratory distress syndrome.

13. The pharmaceutical composition of claim 9 for treating an autoimmune disease selected from glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, chronic thyroiditis, Graves'
20 disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, and graft
25 vs. host disease.

14. The pharmaceutical composition of claim 9 for treating a destructive bone disorder selected from osteoarthritis, osteoporosis and multiple myeloma-related bone disorder.

30 15. The pharmaceutical composition of claim 9 for treating a proliferative disease selected from acute

79580-10

- 108 -

myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma.

16. The pharmaceutical composition of claim 9 for treating a neurodegenerative disease selected from
5 Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity and hypoxia.

17. The pharmaceutical composition of claim 9 for
10 treating ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia or thrombin-induced platelet aggregation.

18. The pharmaceutical composition of claim 9 for treating a condition associated with T-cell activation or
15 pathologic immune responses.

19. The pharmaceutical composition of claim 9 for treating an angiogenic disorder selected from solid tumors, ocular neovascularization, and infantile haemangiomas.

20. The pharmaceutical composition of claim 10 wherein
20 the disease state or condition is osteoporosis, osteoarthritis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, cancer, asthma, ischemic or reperfusion injury, allergic disease, atopic dermatitis, or allergic
25 rhinitis.

21. The pharmaceutical composition of claim 9 for treating a disease selected from inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases,
30 neurodegenerative diseases, allergies, reperfusion/ischemia

79580-10

- 109 -

in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, and thrombin-induced platelet aggregation or for treating a condition associated with proinflammatory cytokines.

5 22. The pharmaceutical composition of claim 9 for treating:

an inflammatory disease selected from acute pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome;

10 an autoimmune disease selected from glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, chronic thyroiditis, Graves' disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia,
15 atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, and graft vs. host disease;

a destructive bone disorder selected from
20 osteoarthritis, osteoporosis and multiple myeloma-related bone disorder;

a proliferative disease selected from acute myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma;

25 a neurodegenerative disease selected from Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity and hypoxia; or

79580-10

- 110 -

ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia, thrombin-induced platelet aggregation, conditions associated with T-cell activation or pathologic immune responses, or an
5 angiogenic disorder selected from solid tumors, ocular neovascularization, and infantile haemangiomas.

23. The pharmaceutical composition of claim 9 for treating a disease selected from: hypercalcemia, restenosis, osteoporosis, osteoarthritis, symptomatic treatment of bone
10 metastasis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, T-cell mediated hypersensitivity disease, Hashimoto's thyroiditis, Guillain-Barre syndrome, chronic obstructive pulmonary disorder, contact dermatitis, cancer,
15 Paget's disease, asthma, ischemic injury, reperfusion injury, allergic disease, atopic dermatitis, and allergic rhinitis.

24. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition
20 for treating a disease state or condition in a mammal that is alleviated by treatment with an inhibitor of c-Jun N-terminal kinases.

25. Use of claim 24 wherein the disease state or condition is selected from inflammatory diseases, autoimmune
25 diseases, destructive bone disorders, proliferative disorders, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin-induced platelet aggregation
30 and conditions associated with proinflammatory cytokines.

79580-10

- 111 -

26. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating an inflammatory disease selected from acute pancreatitis, chronic pancreatitis, asthma, allergies, and
5 adult respiratory distress syndrome.

27. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating an autoimmune disease selected from glomerulonephritis, rheumatoid arthritis, systemic lupus
10 erythematosus, scleroderma, chronic thyroiditis, Graves' disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease,
15 ulcerative colitis, Crohn's disease, psoriasis, and graft vs. host disease.

28. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating a destructive bone disorder selected from
20 osteoarthritis, osteoporosis and multiple myeloma-related bone disorder.

29. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating a proliferative disease selected from acute
25 myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma.

30. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating a neurodegenerative disease selected from
30 Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia

79580-10

- 112 -

or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity and hypoxia.

31. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition
5 for treating ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia or thrombin-induced platelet aggregation.

32. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition
10 for treating a condition associated with T-cell activation or pathologic immune responses.

33. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition
for treating an angiogenic disorder selected from solid
15 tumors, ocular neovascularization, and infantile haemangiomas.

34. Use of claim 24 wherein the disease state or condition is osteoporosis, osteoarthritis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, cancer, asthma,
20 ischemic or reperfusion injury, allergic disease, atopic dermatitis, or allergic rhinitis.

35. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition
for treating a disease selected from inflammatory diseases,
25 autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, and
30 thrombin-induced platelet aggregation or for treating a condition associated with proinflammatory cytokines.

79580-10

- 113 -

36. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating:

an inflammatory disease selected from acute
5 pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome;

an autoimmune disease selected from
glomerulonephritis, rheumatoid arthritis, systemic lupus
erythematosus, scleroderma, chronic thyroiditis, Graves'
10 disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, and graft
15 vs. host disease;

a destructive bone disorder selected from osteoarthritis, osteoporosis and multiple myeloma-related bone disorder;

a proliferative disease selected from acute
20 myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma;

a neurodegenerative disease selected from Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia
25 or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity and hypoxia; or

ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia, thrombin-induced platelet aggregation, conditions associated
30 with T-cell activation or pathologic immune responses, or an

79580-10

- 114 -

angiogenic disorder selected from solid tumors, ocular neovascularization, and infantile haemangiomas.

37. Use of a compound as defined in any one of claims 1 to 8 in preparation of a pharmaceutical composition for treating a disease selected from: osteoporosis, osteoarthritis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, cancer, asthma, ischemic injury, reperfusion injury, allergic disease, atopic dermatitis, and allergic rhinitis.

38. Use of a compound as defined in any one of claims 1 to 8 for treating a disease state or condition in a mammal that is alleviated by treatment with an inhibitor of c-Jun N-terminal kinases.

39. Use of claim 38 wherein the disease state or condition is selected from inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin-induced platelet aggregation and conditions associated with proinflammatory cytokines.

40. Use of a compound as defined in any one of claims 1 to 8 for treating an inflammatory disease selected from acute pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome.

41. Use of a compound as defined in any one of claims 1 to 8 for treating an autoimmune disease selected from glomerulonephritis, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, chronic thyroiditis, Graves' disease, autoimmune gastritis, diabetes, autoimmune

79580-10

- 115 -

hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease, ulcerative colitis, Crohn's disease, psoriasis, and graft
5 vs. host disease.

42. Use of a compound as defined in any one of claims 1 to 8 for treating a destructive bone disorder selected from osteoarthritis, osteoporosis and multiple myeloma-related bone disorder.

10 43. Use of a compound as defined in any one of claims 1 to 8 for treating a proliferative disease selected from acute myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma.

15 44. Use of a compound as defined in any one of claims 1 to 8 for treating a neurodegenerative disease selected from Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia or neurodegenerative disease caused by
20 traumatic injury, glutamate neurotoxicity and hypoxia.

45. Use of a compound as defined in any one of claims 1 to 8 for treating ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia or thrombin-induced platelet aggregation.

25 46. Use of a compound as defined in any one of claims 1 to 8 for treating a condition associated with T-cell activation or pathologic immune responses.

47. Use of a compound as defined in any one of claims 1 to 8 for treating an angiogenic disorder selected

79580-10

- 116 -

from solid tumors, ocular neovascularization, and infantile haemangiomas.

48. Use of claim 38 wherein the disease state or condition is osteoporosis, osteoarthritis, rheumatoid
5 arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, cancer, asthma, ischemic or reperfusion injury, allergic disease, atopic dermatitis, or allergic rhinitis.

49. Use of a compound as defined in any one of
10 claims 1 to 8 for treating a disease selected from inflammatory diseases, autoimmune diseases, destructive bone disorders, proliferative disorders, infectious diseases, neurodegenerative diseases, allergies, reperfusion/ischemia in stroke, heart attacks, angiogenic disorders, organ
15 hypoxia, vascular hyperplasia, cardiac hypertrophy, and thrombin-induced platelet aggregation or for treating a condition associated with proinflammatory cytokines.

50. Use of a compound as defined in any one of claims 1 to 8 for treating:

20 an inflammatory disease selected from acute pancreatitis, chronic pancreatitis, asthma, allergies, and adult respiratory distress syndrome;

an autoimmune disease selected from glomerulonephritis, rheumatoid arthritis, systemic lupus
25 erythematosus, scleroderma, chronic thyroiditis, Graves' disease, autoimmune gastritis, diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, inflammatory bowel disease,
30 ulcerative colitis, Crohn's disease, psoriasis, and graft vs. host disease;

79580-10

- 117 -

a destructive bone disorder selected from osteoarthritis, osteoporosis and multiple myeloma-related bone disorder;

a proliferative disease selected from acute
5 myelogenous leukemia, chronic myelogenous leukemia, metastatic melanoma, Kaposi's sarcoma, and multiple myeloma;


a neurodegenerative disease selected from Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, cerebral ischemia
10 or neurodegenerative disease caused by traumatic injury, glutamate neurotoxicity and hypoxia; or

ischemia/reperfusion in stroke or myocardial ischemia, renal ischemia, heart attacks, organ hypoxia, thrombin-induced platelet aggregation, conditions associated
15 with T-cell activation or pathologic immune responses, or an angiogenic disorder selected from solid tumors, ocular neovascularization, and infantile haemangiomas.

51. Use of a compound as defined in any one of claims 1 to 8 for treating a disease selected from:
20 osteoporosis, osteoarthritis, rheumatoid arthritis, inflammatory bowel disease, multiple sclerosis, psoriasis, lupus, graft vs. host disease, cancer, asthma, ischemic injury, reperfusion injury, allergic disease, atopic dermatitis, and allergic rhinitis.

SMART & BIGGAR
OTTAWA, CANADA

PATENT AGENTS



I