



(51) International Patent Classification:

C07D 413/04 (2006.01) **A61K 31/42** (2006.01)
C07D 261/02 (2006.01) **A61P 29/00** (2006.01)

(21) International Application Number:

PCT/US20 16/0648 18

(22) International Filing Date:

2 December 2016 (02.12.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/263,481 4 December 2015 (04.12.2015) US
62/300,740 26 February 2016 (26.02.2016) US
62/364,814 20 July 2016 (20.07.2016) US

(71) **Applicant: DENALI THERAPEUTICS INC.** [US/US];
151 Oyster Point Blvd., 2nd Floor, South San Francisco,
California 94080 (US).

(72) **Inventors: BONANOMI, Giorgio;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **ESTRADA, Anthony A.;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **FENG, Jianwen A.;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **FOX, Brian;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **LESLIE, Colin Philip;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **LYSSIKATOS, Joseph P.;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **NAPOLITANO, Carmela;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **POZZAN, Alfonso;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **SUDHAKAR, Anantha;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **SWEENEY, Zachary K.;**

c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **TONELLI, Federica;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US). **DE VICENTE FD3ALGO, Javier;** c/o Denali Therapeutics Inc., 151 Oyster Blvd., 2nd Floor, South San Francisco, California 94080 (US).

(74) **Agents: TANNER, Lorna L.** et al; Sheppard Mullin Richter & Hampton LLP, 379 Lytton Avenue, Palo Alto, California 94301-1479 (US).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(in))

Published:

— with international search report (Art. 21(3))

(54) **Title:** ISOXAZOLIDINE DERIVED INHIBITORS OF RECEPTOR INTERACTING PROTEIN KINASE 1 (RIPK 1)

(57) **Abstract:** The present disclosure relates generally to methods and compositions for preventing or arresting cell death and/or inflammation.



ISOXAZOLIDINE DERIVED INHIBITORS OF RECEPTOR INTERACTING PROTEIN
KINASE 1 (RIPK 1)

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit under 35 U.S.C. 119(e) to U.S. Provisional Application Ser.
Nos. 62/263,481, filed December 4, 2015, 62/300,740, filed February 26, 2016, and 62/364,814, filed
July 20, 2016, where the contents of each is incorporated herein by reference in its entirety.

FIELD

The present disclosure relates generally to methods and compositions for preventing or arresting
cell death and/or inflammation.

BACKGROUND

Programmed necrotic cell death, also called necroptosis, is a form of cell death in which various
stimuli such as TNF α , certain toll-like receptor (TLR) agonists and ischemia can induce cellular necrosis.
Necroptosis is a highly inflammatory form of cell death and is thought to be an important contributor to
pathology in multiple degenerative and inflammatory diseases. These diseases include
neurodegenerative diseases, stroke, coronary heart disease and myocardial infarction, retinal degenerative
diseases, inflammatory bowel disease, kidney disease, liver disease and others.

Necrosis is characterized by cell membrane and organelle disruption, cell swelling and
mitochondrial impairment, followed by cell lysis. Also, cell lyses typically are accompanied by an
inflammatory response. Some of the underlying biochemical events in this process are now understood
and the activity of receptor interacting protein kinase 1 has been shown to be important for cells to
undergo necroptosis. Furthermore, this activity is also known to promote the release of inflammatory
mediators such as TNF alpha from cells which can induce inflammation and also promote further
necroptosis. Therefore, identifying and preparing low molecular weight molecules that prevent necrotic
cell death and/or inflammation by inhibiting this or by other mechanisms can provide useful compounds
for therapeutic intervention in diseases characterized by necrotic cell death and/or inflammation.

While progress has been made, there remains a need in the art for improved compounds for
preventing and treating diseases involving cell death and/or inflammation. The present disclosure
provides this and related benefits.

SUMMARY

Provided herein is a method for inhibiting receptor interacting protein kinase 1 in a cell,
comprising contacting the cell with a receptor interacting protein kinase 1 inhibitor. Also provided are
compounds that are useful as inhibitors of receptor interacting protein kinase 1. The disclosure also
provides compositions, including pharmaceutical compositions, kits that include the compounds and
methods of using and making the compounds. The disclosure further provides compounds or
compositions thereof for use in a method of treating a disease, disorder or condition that is mediated by
receptor interacting protein kinase 1. Moreover, the disclosure provides uses of the compounds or
compositions thereof in the manufacture of a medicament for the treatment of a disease, disorder or
condition that is mediated by (or mediated, at least in part, by) receptor interacting protein kinase 1.

These and other aspects of the disclosure will be apparent upon reference to the following detailed description. To this end, various references are set forth herein which describe in more detail certain background information, procedures, compounds and/or compositions and are each hereby incorporated by reference in their entirety.

DETAILED DESCRIPTION

The following description sets forth exemplary embodiments of the present technology. It should be recognized, however, that such description is not intended as a limitation on the scope of the present disclosure but is instead provided as a description of exemplary embodiments.

1. Definitions

As used in the present specification, the following words, phrases and symbols are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise.

A dash ("-") that is not between two letters or symbols is used to indicate a point of attachment for a substituent. For example, $-C(=O)NH_2$ is attached through the carbon atom. A dash at the front or end of a chemical group is a matter of convenience; chemical groups may be depicted with or without one or more dashes without losing their ordinary meaning. A wavy line or a dashed line drawn through a line in a structure indicates a specified point of attachment of a group. Unless chemically or structurally required, no directionality or stereochemistry is indicated or implied by the order in which a chemical group is written or named.

The prefix " C_{u-v} " indicates that the following group has from u to v carbon atoms. For example, " C_{1-6} alkyl" indicates that the alkyl group has from 1 to 6 carbon atoms.

Reference to "about" a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se. In certain embodiments, the term "about" includes the indicated amount $\pm 10\%$. In certain embodiments, the term "about" includes the indicated amount $\pm 5\%$. In certain embodiments, the term "about" includes the indicated amount $\pm 1\%$. Also, the term "about X" includes description of "X". Also, the singular forms "a" and "the" include plural references unless the context clearly dictates otherwise. Thus, e.g., reference to "the compound" includes a plurality of such compounds and reference to "the assay" includes reference to one or more assays and equivalents thereof known to those skilled in the art.

"Alkyl" refers to an unbranched or branched saturated hydrocarbon chain. As used herein, alkyl has 1 to 20 carbon atoms (i.e., C_{1-20} alkyl), 1 to 12 carbon atoms (i.e., C_{1-12} alkyl), 1 to 8 carbon atoms (i.e., C_{1-8} alkyl), 1 to 6 carbon atoms (i.e., C_{1-6} alkyl) or 1 to 4 carbon atoms (i.e., C_{1-4} alkyl). Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, pentyl, 2-pentyl, isopentyl, neopentyl, hexyl, 2-hexyl, 3-hexyl and 3-methylpentyl. When an alkyl residue having a specific number of carbons is named by chemical name or identified by molecular formula, all positional isomers having that number of carbons may be encompassed; thus, for example, "butyl" includes n-butyl (i.e., $-(CH_2)_3CH_3$), sec-butyl (i.e., $-CH(CH_3)CH_2CH_3$), isobutyl (i.e., $-CH_2CH(CH_3)_2$) and tert-butyl (i.e., $-C(CH_3)_3$); and "propyl" includes n-propyl (i.e., $-(CH_2)_2CH_3$) and isopropyl (i.e., $-CH(CH_3)_2$).

Certain commonly used alternative chemical names may be used. For example, a divalent group such as a divalent "alkyl" group, a divalent "aryl" group, etc., may also be referred to as an "**alkylene**" group or an "**alkylenyl**" group, an "**arylene**" group or an "**arylenyl**" group, respectively. Also, unless indicated explicitly **otherwise**, where combinations of groups are **referred** to herein as one moiety, e.g. **arylalkyl** or **aralkyl**, the last mentioned group contains the atom by which the moiety is attached to the rest of the molecule.

"**Aikenyl**" refers to an alkyl group containing at least one carbon-carbon double bond and having from 2 to 20 carbon atoms (i.e., C_{2-20} aikenyl), 2 to 8 carbon atoms (i.e., C_{2-8} aikenyl), 2 to 6 carbon atoms (i.e., C_{2-6} aikenyl) or 2 to 4 carbon atoms (i.e., C_{2-4} aikenyl). Examples of aikenyl groups include **ethenyl**, **propenyl**, **butadienyl** (including **1,2-butadienyl** and **1,3-butadienyl**).

"**Alkynyl**" refers to an alkyl group containing at least one carbon-carbon triple bond and having from 2 to 20 carbon atoms (i.e., C_{2-20} alkynyl), 2 to 8 carbon atoms (i.e., C_{2-8} alkynyl), 2 to 6 carbon atoms (i.e., C_{2-6} alkynyl) or 2 to 4 carbon atoms (i.e., C_{2-4} alkynyl). The term "alkynyl" also includes those groups having one triple bond and one double bond.

"**Alkoxy**" refers to the group "alkyl-Q-". Examples of **alkoxy** groups include methoxy, ethoxy, **n-propoxy**, **iso-propoxy**, **n-butoxy**, **tert-butoxy**, **sec-butoxy**, **n-pentoxy**, **n-hexoxy** and **1,2-dimethylbutoxy**.

"**Alkoxyalkyl**" refers to the group "alkyl -O-alkyl".

"**Alkylthio**" refers to the group "alkyl -S-".

"**Acyl**" refers to a group -C(=O)R, wherein R is hydrogen, alkyl, aikenyl, alkynyl, **cycloalkyl**, heterocyclyl, **aryl**, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein. Examples of acyl include **formyl**, **acetyl**, **cyclohexylcarbonyl**, **cyclohexylmethyl-carbonyl** and **benzoyl**.

"**Amido**" refers to both a "C-amido" group which refers to the group -C(Q)NR^yR^z and an "N-amido" group which refers to the group -NR^yC(=O)R^z, wherein R^y and R^z are independently hydrogen, alkyl, aikenyl, alkynyl, cycloalkyl, heterocyclyl, **aryl**, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

"**Amino**" refers to the group -NR^yR^z wherein R^y and R^z are independently hydrogen, alkyl, aikenyl, alkynyl, cycloalkyl, heterocyclyl, **aryl**, heteroalkyl or **heteroaryl**; each of which may be optionally substituted, as defined herein.

"**Amidino**" refers to -C(NR)(NR₂), wherein each R is independently hydrogen, alkyl, aikenyl, alkynyl, cycloalkyl, heterocyclyl, **aryl**, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

"**Aryl**" refers to an aromatic **carbocyclic** group having a single ring (e.g. monocyclic) or multiple rings (e.g. **bicyclic** or **tricyclic**) including fused systems. As used herein, **aryl** has 6 to 20 ring carbon atoms (i.e., C_{6-20} **aryl**), 6 to 18 carbon ring atoms (i.e., C_{6-18} **aryl**), 6 to 12 carbon ring atoms (i.e., C_{6-12} **aryl**) or 6 to 10 carbon ring atoms (i.e., C_{6-10} **aryl**). Examples of aryl groups include **phenyl**, **naphthyl**, **fluorenyl** and **anthryl**. Aryl, however, does not encompass or overlap in any way with heteroaryl defined

below. If **one** or more **aryl** groups are fused with a heteroaryl, the resulting ring system is heteroaryl. If one or more aryl groups are fused with a heterocyclyl, the resulting ring system is heterocyclyl.

"Arylalkyl" or "**Aralkyl**" refers to **the** group "aryl-alkyl-".

"Carbamoyl" refers to both an "**O**-carbamoyl" group which refers to the group $\sim\text{O}\sim\text{C}(\text{O})\text{NR}^y\text{R}^z$ and an "**N**-carbamoyl" group which refers to the group $\sim\text{NR}^y\text{C}(\text{O})\text{OR}^z$, wherein R^y and R^z are independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

"Carboxyl ester" or "ester" **refer to** both **-OC(O)R** and **-C(O)OR**, wherein R is alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

"Cycloalkyl" refers to a saturated or partially unsaturated cyclic alkyl group having a single ring or multiple rings including fused, bridged and spiro ring systems. The term "cycloalkyl" includes **cycloalkenyl** groups (i.e., the cyclic group having at least one double bond). As used herein, cycloalkyl has from 3 to 20 ring carbon atoms (i.e., C_{3-20} cycloalkyl), 3 to 15 ring carbon atoms (i.e., C_{3-15} cycloalkyl), 3 to 12 ring carbon atoms (i.e., C_{3-12} cycloalkyl), 3 to 10 ring carbon atoms (i.e., C_{3-10} cycloalkyl), 3 to 8 ring carbon atoms (i.e., C_{3-8} cycloalkyl) or 3 to 6 ring carbon atoms (i.e., C_{3-6} cycloalkyl). Cycloalkyl also includes "spiro cycloalkyl" when there are two positions for substitution on **the** same carbon atom. Monocyclic radicals include, for example, **cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl** and **cyclooctyl**. Polycyclic radicals include, for example, **adamantyl, norbornyl, decalinyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl** and the like. Further, the term cycloalkyl is intended to encompass any non-aromatic ring which may be fused to an aryl ring, regardless of the attachment to the remainder of the molecule.

"Cycloalkylalkyl" refers to the group "**cycloalkyl-alkyl**".

"Guanidino" refers to **-NRC(NR)(NR)**, wherein each R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

"Imino" refers to a group **-C(NR)R**, wherein each R is independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or **heteroaryl**; each of which may be optionally substituted, as defined herein.

"Imido" refers to a group **-C(O)NRC(O)R**, wherein each R is independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each **of which** may be optionally substituted, as defined herein.

"Halogen" or "halo" includes **fluoro, chloro, bromo** and **iodo**.

"Haloalkyl" refers to an **unbranched** or branched alkyl group as defined above, wherein one or more hydrogen atoms are replaced by a halogen. For example, where a residue is substituted **with** more than one halogen, it may be referred to by using a prefix corresponding to the number of halogen moieties attached. Dihaloalkyl and **trihaloalkyl** refer to alkyl substituted with two ("di") or three ("tri") halo groups, which may be, but are **not** necessarily, the same halogen. Examples of haloalkyl include

trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2-difluoroethyl, 3-bromo-2-fluoropropyl, 1,2-dibromoethyl and the like.

"Haloalkoxy" refers to an alkoxy group as defined above, wherein one or more hydrogen atoms are replaced by a halogen.

5 "Hydroxyalkyl" refers to an alkyl group as defined above, wherein one or more hydrogen atoms are replaced by a hydroxy group.

"Heteroalkyl" refers to an alkyl group in which one or more of the carbon atoms (and any associated hydrogen atoms) are each independently replaced with the same or different heteroatomic group, provided the point of attachment to the remainder of the molecule is through a carbon atom. The
 10 term "heteroalkyl" includes unbranched or branched saturated chain having carbon and heteroatoms. By way of example, 1, 2 or 3 carbon atoms may be independently replaced with the same or different heteroatomic group. Heteroatomic groups include, but are not limited to, -NR-, -O-, -S-, $\sim S(0)\sim$, -S(0)₂-, and the like, where R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl, or heteroaryl; each of which may be optionally substituted, as defined herein. Examples of heteroalkyl
 15 groups include -CH₂OCH₃, -CH₂SCH₃, and -CH₂NRCH₃, where R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl, or heteroaryl; each of which may be optionally substituted, as defined herein. As used herein, heteroalkyl includes 1 to 10 carbon atoms, 1 to 8 carbon atoms, or 1 to 4 carbon atoms; and 1 to 3 heteroatoms, 1 to 2 heteroatoms, or 1 heteroatom.

"Heteroaryl" refers to an aromatic group (e.g., a 5-14 membered ring system) having a single
 20 ring, multiple rings or multiple fused rings, with one or more ring heteroatoms independently selected from nitrogen, oxygen and sulfur. As used herein, heteroaryl includes 1 to 20 ring carbon atoms (i.e., C₁₋₂₀ heteroaryl), 1 to 13 ring carbon atoms (i.e., C₃₋₁₂ heteroaryl), 3 to 12 ring carbon atoms (i.e., C₃₋₁₂ heteroaryl) or 3 to 8 carbon ring atoms (i.e., C₃₋₈ heteroaryl); and 1 to 6 heteroatoms, 1 to 5 heteroatoms, 1 to 4 heteroatoms, 1 to 3 ring heteroatoms, 1 to 2 ring heteroatoms or 1 ring heteroatom independently
 25 selected from nitrogen, oxygen and sulfur. Examples of heteroaryl groups include azepinyl, acridinyl, benzimidazolyl, benzothiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxany, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl,
 30 carbazolyl, cinnolinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isomdoil, indolyl, isoindolyl, isoquinolyl, indolizyl, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl,
 35 quinazolinyl, quinoxalinyl, quinolinyl, quinuclidinyl, isoquinolinyl, tetrahydroquinolinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazmyl and thiophenyl (i.e., thienyl). Examples of the fused-heteroaryl rings include, but are not limited to, benzo[d]thiazolyl, quinolinyl, isoquinolinyl, benzo[b]thiophenyl, indazolyl, benzo[d]imidazolyl, pyrazolo[1,5-a]pyridinyl and imidazo[1,5-a]pyridinyl, where the

heteroaryl can be bound via either ring of the fused system. Any aromatic ring, having a single or multiple fused rings, containing at least one heteroatom, is considered a heteroaryl regardless of the attachment to the remainder of the molecule (i.e., through any one of the fused rings). Heteroaryl does not encompass or overlap with aryl as defined above.

5 "Heteroarylalkyl" refers to the group "heteroaryl-alkyl-".

"Heterocyclyl" refers to a saturated or unsaturated cyclic alkyl group, with one or more ring heteroatoms independently selected from nitrogen, oxygen and sulfur. The term "heterocyclyl" includes heterocycloalkenyl groups (i.e., the heterocyclyl group having at least one double bond), bridged-heterocyclyl groups, fused-heterocyclyl groups and spiro-heterocyclyl groups. A heterocyclyl may be a
 10 single ring or multiple rings wherein the multiple rings may be fused, bridged or spiro, and may comprise one or more oxo (C=O) or N-oxide (N-O-) moieties. Any non-aromatic ring containing at least one heteroatom is considered a heterocyclyl, regardless of the attachment (i.e., can be bound through a carbon atom or a heteroatom). Further, the term heterocyclyl is intended to encompass any non-aromatic ring containing at least one heteroatom, which ring may be fused to an aryl or heteroaryl ring, regardless of
 15 the attachment to the remainder of the molecule. As used herein, heterocyclyl has 2 to 20 ring carbon atoms (i.e., C₂₋₂₀ heterocyclyl), 2 to 12 ring carbon atoms (i.e., C₂₋₁₂ heterocyclyl), 2 to 10 ring carbon atoms (i.e., C₂₋₁₀ heterocyclyl), 2 to 8 ring carbon atoms (i.e., C₂₋₈ heterocyclyl), 3 to 12 ring carbon atoms (i.e., C₃₋₁₂ heterocyclyl), 3 to 8 ring carbon atoms (i.e., C₃₋₈ heterocyclyl) or 3 to 6 ring carbon atoms (i.e., C₃₋₆ heterocyclyl); having 1 to 5 ring heteroatoms, 1 to 4 ring heteroatoms, 1 to 3 ring
 20 heteroatoms, 1 to 2 ring heteroatoms or 1 ring heteroatom independently selected from nitrogen, sulfur or oxygen. Examples of heterocyclyl groups include dioxolanyl, thienyl [1,3]dithianyl, decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholinyl, octahydroindolyl, octahydroisindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidinyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholinyl, thiomorpholinyl, 1-oxo-thiomorpholinyl and 1,1-dioxo-thiomorpholinyl. Examples of the spiro-heterocyclyl rings include bicyclic and tricyclic ring systems, such as 2-oxa-7-azaspiro[3.5]nonanyl, 2-oxa-6-azaspiro[3.4]octanyl and 6-oxa-1-azaspiro[3.3]heptanyl. Examples of the fused-heterocyclyl rings include, but are not limited to, 1,2,3,4-tetrahydroisoquinolyl, 4,5,6,7-tetrahydrothieno[2,3-c]pyridinyl, indolyl and isoindolyl,
 30 where the heterocyclyl can be bound via either ring of the fused system.

"Heterocyclylalkyl" refers to the group "heterocyclyl-alkyl-".

"Oxime" refers to the group -CR(=NOH) wherein R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein.

35 "Sulfonyl" refers to the group -S(O)₂R, where R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroaryl; each of which may be optionally substituted, as defined herein. Examples of sulfonyl are methylsulfonyl, ethylsulfonyl, phenyl sulfonyl and toluenesulfonyl.

"Sulfinyl" refers to the group $-S(O)R$, where R is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroarvl; each of which may be optionally substituted, as defined herein. Examples of sulfinyl are methylsulfinyl, ethylsulimyl, phenylsulfinyl and toluenesulfinyl.

"Sulfonamide" refers to the groups $-SO_2NRR$ and $-NRSO_2R$, where each R is independently hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, heteroalkyl or heteroarvl; each of which may be optionally substituted, as defined herein.

"Alkylsulfonyl" refers to the group $-S(O)_2R$, where R is alkyl.

"Alkylsulfonyl" refers to the group $-S(O)R$, where R is alkyl.

The terms "optional" or "optionally" means that the subsequently described event or circumstance may or may not occur and that the description includes instances where said event or circumstance occurs and instances in which it does not. Also, the term "optionally substituted" refers to any one or more hydrogen atoms on the designated atom or group may or may not be replaced by a moiety other than hydrogen.

The term "substituted" used herein means any of the above groups (i.e., alkyl, alkenyl, alkynyl, aikylene, alkoxy, haloalkyl, haloalkoxy, cycloalkyl, aryl, heterocyclyl, heteroarvl, and/or heteroalkyl) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atom such as, but not limited to alkyl, alkenyl, alkynyl, alkoxy, alkylthio, acyl, amido, amino, amidino, aryl, aralkyl, azido, carbamoyl, carboxyl, carboxyl ester, cyano, cycloalkyl, cycloalkylalkyl, guanadino, halo, haloalkyl, haloalkoxy, hydroxyalkyl, heteroalkyl, heteroarvl, heteroarylalkyl, heterocyclyl, heterocyclylalkyl, hydrazine, hydrazone, imino, imido, hydroxy, oxo, oxime, nitro, sulfonyl, sulfinyl, alkylsulfonyl, alkylsulfonyl, thiocyanate, sulfinic acid, sulfonic acid, sulfonamide, thiol, thioxo, N-oxide, or $-Si(R^{100})_3$ wherein each R^{100} is independently hydrogen, alkyl, alkenyl, alkynyl, heteroalkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl.

The term "substituted" used herein means any of the above groups (i.e., alkyl, aikylene, alkoxy, haloalkoxy, aryl, cycloalkyl, haloalkyl, heterocyclyl, heteroaryl, hydroxyalkyl and/or alkoxyalkyl) wherein at least one hydrogen atom is replaced by a bond to a non-hydrogen atom such as, but not limited to: an alkyl group, a haloalkyl group, a halogen atom such as F, Cl, Br, and I; an alkenyl, a haloalkenyl group, an alkynyl group, a haloalkynyl group, a cyclic group such as an aryl, heteroaryl, cycloalkyl, or heterocyclyl group, an oxygen atom in groups such as hydroxyl groups, alkoxy groups, and ester groups; a sulfur atom in groups such as thiol groups, thioalkyl groups, thiohaloalkyl groups, sulfone groups, sulfonyl groups, and sulfoxide groups; a nitrogen atom in groups such as amines, amides, aikylamines, diaikylamines, arylamines, aikylaryl amines, diarylamines, N-oxides, imides, and enamines; a silicon atom in groups such as trialkylsilyl groups, dialkylarylsilyl groups, alkyl diarylsilyl groups, and triarylsilyl groups; and other heteroatoms in various other groups. "Substituted" also means any of the above groups in which one or more hydrogen atoms are replaced by a higher-order bond (e.g., a double- or triple-bond) to a heteroatom such as oxygen in oxo, carbonyl, formyl, carboxyl, carbonate, and ester groups; and nitrogen in groups such as imines, oximes, hydrazones, and nitriles.

In certain embodiments, "substituted" includes any of the above groups in which one or more hydrogen atoms are replaced **with** $-\text{NR}^g\text{R}^h$, $-\text{NR}^g\text{C}(=\text{O})\text{R}^h$, $-\text{R}^g\text{C}(\text{O})\text{NR}^g\text{R}^h$, $-\text{NR}^g\text{C}(=\text{O})\text{GR}^h$, $-\text{NR}^g\text{SO}_2\text{R}^h$, $-\text{OC}(=\text{O})\text{NR}^g\text{R}^h$, $-\text{OR}^g$, $-\text{SR}^g$, $-\text{SOR}^g$, $-\text{SO}_2\text{R}^g$, $-\text{OSO}_2\text{R}^g$, $-\text{SO}_2\text{OR}^g$, $=\text{NSO}_2\text{R}^g$, and $-\text{SO}_2\text{NR}^g\text{R}^h$.

"Substituted" also means any of the above groups in which one or more hydrogen atoms are replaced

5 with $-(\text{O})\text{R}^g$, $-\text{C}(=\text{O})\text{R}^g$, $-(\text{O})\text{NR}^g\text{R}^h$, $-\text{CH}_2\text{S}(\text{O})_2\text{R}^g$, $-\text{Cl}\text{SO}_2\text{NR}^g\text{R}^h$. "Substituted" further means any of the above groups in which one or more hydrogen atoms are replaced by $-\text{NR}^g\text{S}(\text{O})_{1-2}\text{NR}^g\text{R}^h$, $-\text{CH}_2\text{S}(\text{O})\text{R}^g$, $-\text{CH}_2\text{S}(\text{O})\text{NR}^{3/4-}$, $-\text{QC}(=\text{O})\text{OR}^g$, $-\text{SF}_5$, $-\text{SCF}_3$ or $-\text{OCF}_3$. "Substituted" further means any of the above groups in which one or more hydrogen atoms are replaced by a bond to an amino, cyano, **hydroxyl**, imino, nitro, oxo, thiooxo, halo, alkyl, aikoxy, **alkylamino**, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, heterocyclyl, N-heterocyclyl, heterocyclalkyl, heteroaryl, and/or

10 heteroarylalkyl group. In the foregoing, R^g and R^h and R^i are the same or different and independently hydrogen, halo, alkyl, alkenyl, alkynyl, aikoxy, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, and/or heteroarylalkyl. In an embodiment, each of said hydrogen, alkyl, alkenyl, **alkynyl**, aikoxy, thioalkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl,

15 haloalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, and/or heteroarylalkyl are independently optionally substituted with one or more oxo, alkyl, halo, amino, hydroxyl or **alkoxyl**. In addition, each of the foregoing substituents may also be optionally substituted with one or more of the above substituents.

Polymers or similar indefinite structures arrived at by defining substituents with further substituents appended ad infinitum (e.g., a substituted aryl having a substituted alkyl which is itself substituted with a substituted aryl group, which is further substituted by a substituted heteroalkyl group, etc.) are not intended for inclusion herein. Unless otherwise noted, the maximum number of serial substitutions in compounds described herein is three. For example, serial substitutions of substituted aryl groups with two other substituted aryl groups are limited to ((substituted aryl)substituted **aryl**) substituted **aryl**. Similarly, the above definitions are not intended to include impermissible substitution patterns (e.g.,

25 methyl substituted with 5 fluorines or **heteroaryl** groups having two adjacent oxygen ring atoms). Such impermissible substitution patterns are well known to the skilled artisan. **When** used to modify a chemical group, **the** term "substituted" may describe other chemical groups defined herein. Unless specified **otherwise**, where a group is described as optionally substituted, any substituents of **the** group are themselves unsubstituted. For example, **in** certain embodiments, the term "substituted alkyl" refers to

30 an alkyl group having one or more substituents including hydroxy, halo, aikoxy, **acyl**, oxo, amino, cycloalkyl, heterocyclyl, aryl and heteroaryl. **In** certain embodiments, the one or more substituents may be **further** substituted **with** halo, alkyl, haloalkyl, hydroxy, aikoxy, cycloalkyl, heterocyclyl, aryl or heteroaryl, each of which is substituted. **In** certain embodiments, the substituents may be **further** substituted **with** halo, alkyl, haloalkyl, aikoxy, hydroxy, cycloalkyl, heterocyclyl, aryl or heteroaryl, each

35 of which is unsubstituted.

Any compound or structure given herein, is also intended to represent **unlabeled** forms as well as **isotopically** labeled forms of the compounds. **Isotopically** labeled compounds have structures depicted by the **formulas** given herein except that one or more atoms are replaced by an atom having a selected

atomic mass or mass number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine and iodine, such as ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I and ^{125}I , respectively. Various isotopically labeled compounds of the present disclosure, for example those into which radioactive isotopes such as ^3H , ^{13}C and ^{14}C are incorporated. Such isotopically labelled compounds may be useful in metabolic studies, reaction kinetic studies, detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays or in radioactive treatment of patients.

The disclosure also includes "deuterated analogs" of compounds described herein in which from 1 to n hydrogens attached to a carbon atom is/are replaced by deuterium, in which n is the number of hydrogens in the molecule. Such compounds exhibit increased resistance to metabolism and are thus useful for increasing the half-life of any compound when administered to a mammal, particularly a human. See, for example, Foster, "Deuterium Isotope Effects in Studies of Drug Metabolism," Trends Pharmacol. Sci. 5(12):524-527 (1984). Such compounds are synthesized by means well known in the art, for example by employing starting materials in which one or more hydrogens have been replaced by deuterium.

Deuterium labelled or substituted therapeutic compounds of the disclosure may have improved DMPK (drug metabolism and pharmacokinetics) properties, relating to distribution, metabolism and excretion (ADME). Substitution with heavier isotopes such as deuterium may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life, reduced dosage requirements and/or an improvement in therapeutic index. An ^{18}F , ^3H , ^{11}C labeled compound may be useful for PET or SPECT or other imaging studies. Isotopically labeled compounds of this disclosure and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent. It is understood that deuterium in this context is regarded as a substituent in a compound described herein.

The concentration of such a heavier isotope, specifically deuterium, may be defined by an isotopic enrichment factor. In the compounds of this disclosure any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a position is designated specifically as " ^1H " or "hydrogen", the position is understood to have hydrogen at its natural abundance isotopic composition. Accordingly, in the compounds of this disclosure any atom specifically designated as a deuterium (D) is meant to represent deuterium.

In many cases, the compounds of this disclosure are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto.

Provided are also pharmaceutically acceptable salts, hydrates, solvates, tautomeric forms, stereoisomers and prodrugs of the compounds described herein. "Pharmaceutically acceptable" or "physiologically acceptable" refer to compounds, salts, compositions, dosage forms and other materials

which are useful in preparing a pharmaceutical composition that is suitable for veterinary or human pharmaceutical use.

The term "pharmaceutically acceptable salt" of a given compound refers to salts that retain the biological effectiveness and properties of the given compound and which are not biologically or otherwise undesirable. "Pharmaceutically acceptable salts" or "physiologically acceptable salts" include, for example, salts with inorganic acids and salts with an organic acid. In addition, if the compounds described herein are obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds. Those skilled in the art will recognize various synthetic methodologies that may be used to prepare nontoxic pharmaceutically acceptable addition salts. Pharmaceutically acceptable acid addition salts may be prepared from inorganic and organic acids. Salts derived from inorganic acids include hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like. Salts derived from organic acids include acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluene-sulfonic acid, salicylic acid and the like. Likewise, pharmaceutically acceptable base addition salts can be prepared from inorganic and organic bases. Salts derived from inorganic bases include, by way of example only, sodium, potassium, lithium, ammonium, calcium and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines, such as alkyl amines (i.e., $\text{NR}_3(\text{alkyl})$), dialkyl amines (i.e., $\text{HN}(\text{alkyl})_2$), trialkyl amines (i.e., $\text{N}(\text{alkyl})_3$), substituted alkyl amines (i.e., $\text{NH}_2(\text{substituted alkyl})$), di(substituted alkyl) amines (i.e., $\text{HN}(\text{substituted alkyl})_2$), tri(substituted alkyl) amines (i.e., $\text{N}(\text{substituted alkyl})_3$), alkenyl amines (i.e., $\text{NH}_2(\text{alkenyl})$), dialkenyl amines (i.e., $\text{HN}(\text{alkenyl})_2$), trialkenyl amines (i.e., $\text{N}(\text{alkenyl})_3$), substituted alkenyl amines (i.e., $\text{NH}_2(\text{substituted alkenyl})$), di(substituted alkenyl) amines (i.e., $\text{HN}(\text{substituted alkenyl})_2$), tri(substituted alkenyl) amines (i.e., $\text{N}(\text{substituted alkenyl})_3$), mono-, di- or tri- cycloalkyl amines (i.e., $\text{NH}_2(\text{cycloalkyl})$, $\text{HN}(\text{cycloalkyl})_2$, $\text{N}(\text{cycloalkyl})_3$), mono-, di- or tri- arylamines (i.e., $\text{NH}_2(\text{aryl})$, $\text{HN}(\text{aryl})_2$, $\text{N}(\text{aryl})_3$) or mixed amines, etc. Specific examples of suitable amines include, by way of example only, isopropylamine, trimethyl amine, diethyl amine, tri(iso-propyl) amine, tri(n-propyl) amine, ethanamine, 2-dimethylaminoethanol, piperazine, piperidine, morpholine, N-ethylpiperidine and the like.

The term "hydrate" refers to the complex formed by the combining of a compound described herein and water.

A "solvate" refers to an association or complex of one or more solvent molecules and a compound of the disclosure. Examples of solvents that form solvates include, but are not limited to, water, isopropanol, ethanol, methanol, dimethylsulfoxide, ethylacetate, acetic acid and ethanamine.

Some of the compounds exist as tautomers. Tautomers are in equilibrium with one another. For example, amide containing compounds may exist **in** equilibrium with imidic **acid** tautomers. Regardless of which **tautomer** is shown and regardless of the nature of the equilibrium among tautomers, the compounds are understood by one of ordinary- skill **in** the art **to** comprise both amide and imidic acid
5 tautomers. **Thus**, the amide containing compounds are understood to include their imidic acid tautomers. Likewise, the imidic acid containing compounds are understood **to** include their amide tautomers.

The compounds of the invention, **or their pharmaceutically** acceptable salts include an asymmetric center and may thus give rise to enantiomers, diastereomers, and other stereoisomeric **forms** **that** may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or as (**D**)- or (**L**)- for amino
10 acids. **The** present invention is meant to include all such possible isomers, as well as their **racemic** and optically pure forms. Optically active (+) and (-), (*R*)- and (*S*)-, or (**D**)- and (**L**)- isomers may be prepared using **chiral synthons** or **chiral** reagents, or resolved using conventional techniques, **for** example, chromatography and fractional crystallization. Conventional techniques for **the** preparation/isolation of individual enantiomers include chiral synthesis from a suitable optically pure precursor or resolution of
15 the **racemate** (or the **racemate** of a salt or derivative) using, for example, chiral high pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centres of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include **both** E and Z geometric isomers. Likewise, all tautomeric **forms** are also intended to be included.

A "stereoisomer" refers to a compound made up **of the** same atoms bonded by the same bonds
20 but having different three-dimensional **structures**, which are not interchangeable. The present invention contemplates various stereoisomers and mixtures thereof and includes "enantiomers", which refers **to two** stereoisomers whose molecules are **nonsuperimposable** mirror images **of one another**.

"Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other.

"Prodrugs" means any compound which releases an active parent drug according to a structure
25 described herein *in vivo* when such prodrug is administered **to** a mammalian subject. Prodrugs of a compound described herein are prepared by modifying functional groups present in the compound described herein in such a way that the modifications may be cleaved *in vivo* to release the parent compound. **Prodrugs** may be prepared by modifying **functional** groups present in the compounds in such
30 a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds. Prodrugs include compounds described herein **wherein** a hydroxy, amino, carboxy or sulphydryl group in a compound described herein is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxy, amino or sulphydryl group, respectively. Examples **of prodrugs** include, but are **not** limited to esters (e.g., acetate, formate and **benzoate** derivatives), amides, guanidines, carbamates
35 (e.g., N,N-dimethylaminocarbonyl) of hydroxy functional groups in compounds described herein and the like. Preparation, selection and use of prodrugs is discussed **in** T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the **A.C.S.** Symposium Series; "Design **of Prodrugs**," ed. H. Bundgaard, Elsevier, 1985; and in Bioreversible Carriers in **Drug** Design, ed. Edward B. Roche,

American Pharmaceutical Association and Pergamon Press, 1987, each of which are hereby incorporated by reference in their entirety.


As used herein, "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" or "excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The term "necrotic cell disease" refers to diseases associated with or caused by cellular necrosis. Exemplary necrotic cell diseases include, but are not limited to, acute diseases such as trauma, ischemia, stroke, cardiac infarction, anthrax lethal toxin induced septic shock, sepsis, cell death induced by LPS and HIV induced T-cell death leading to immunodeficiency. The term "necrotic cell disease" also includes but is not limited to chronic neurodegenerative diseases, such as Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, Alzheimer's disease, infectious encephalopathies, dementia such as HIV associated dementia. The term "necrotic cell disease" also includes but is not limited to diseases such as inflammatory bowel disease and acute and chronic kidney disease which are characterized by inflammation and cell death.

The chemical names used herein are generated using the MarvinSketch Version 6.1.6 (ChemAxon) or the ChemDraw Ultra Version 13.0 software naming programs.

2. Compounds

Provided are compounds having the following structure (X):



or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof, wherein:

A is -C(=O)-, -S(=O)-, -S(=O)₂- or -S(=O)(=NH)-;

R¹ is -NR^{1'}R^{1''}, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R² is aryl or heteroaryl;

R^{2'} is hydrogen or R² and R^{2'} together form a C₃₋₁₀ cycloalkyl or heterocyclyl;

R³ is hydrogen, deuterium, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R^{3'} is hydrogen, deuterium, or R³ and R^{3'} together form a C₃₋₁₀ cycloalkyl or heterocyclyl;

R⁴ is hydrogen, halo, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R^{4'} is hydrogen or R⁴ and R^{4'} together form a C₃₋₁₀ cycloalkyl or heterocyclyl;

or R^{2'} and R^{4'} or R^{3'} and R^{4'} together form a C₃₋₁₀ cycloalkyl or heterocyclyl;

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each **Ci-6** alkyl, **C3-6** alkoxy, **C3-10** cycloalkyl, heterocyclyl, heteroaryl and aryl group of R^1 , R^2 , $R^{2'}$, R^3 , R^3 , R^4 , R^4 , R^5 and R^6 is optionally substituted with one, two, three or four R^{10} ;

R^{10} in each instance is independently halo, hydroxy, cyano, nitro, **C1-6** alkyl, **C1-6** alkoxy, **C2-6** aikenyl, **C2-6** alkynyl, **C3-10** cycloalkyl, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$,
 5 $-C(=O)NR^{11}$, $-Si(=O)(R^{11})_2$, $-S(=O)_2R^{11}$, $-NR^{11}R^{11}$, $-NR^{11}C(=O)R^{11}$ or $-NR^{11}C(=O)OR^{11}$;

wherein the **C-6** alkyl, **C1-6** alkoxy, **C2-6** aikenyl, **C2-6** aikynyl, **C3-10** cycloalkyl, heterocyclyl, aryl and heteroaryl of R^{10} are optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, **C1-6** alkyl, **C1-6** alkoxy, **C1-6** haloalkoxy, **C2-6** aikenyl, **C2-6** alkynyl, **C3-10** cycloalkyl, heterocyclyl, aryl or heteroaryl; and

10 R^{11} in each instance independently is hydrogen, **C1-6** alkyl, **C3-10** cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the **C1-6** alkyl of R^{11} is optionally substituted with halo or oxo and wherein the **C3-10** cycloalkyl, heterocyclyl, heteroaryl or aryl of R^{11} is optionally substituted with one, two or three **C1-6** alkyl;

15 provided the compound is not ethyl 3-phenylisoxazolidine-2-carboxylate, benzyl 3-(4-amino-2-oxopyrimidin-1(2H)-yl)isoxazolidine-2-carboxylate, 2-(methanesulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate, 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one.
 20

Also provided are compounds having structure (X), or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof, wherein:

A is $-C(=O)-$, $-S(=O)-$, $-S(=O)_2-$ or $-Si(=O)(=NR^{11})-$;

R^1 is $-NR^5R^6$, **Ci-6** alkyl, **C3-6** alkoxy, **C3-10** cycloalkyl, heterocyclyl or heteroaryl;

25 R^2 is aryl or heteroaryl;

$R^{2'}$ is hydrogen or R^2 and R^2 together form a **C3-10** cycloalkyl or heterocyclyl;

R^3 is hydrogen, deuterium, **C-6** alkyl or **C3-10** cycloalkyl;

$R^{3'}$ is hydrogen, deuterium, or R^3 and $R^{3'}$ together form a **C3-10** cycloalkyl or heterocyclyl;

R^4 is hydrogen, halo, **C1-6** alkyl or **C3-10** cycloalkyl;

30 $R^{4'}$ is hydrogen or R^4 and $R^{4'}$ together form a **C3-10** cycloalkyl or heterocyclyl;

or $R^{2'}$ and $R^{4'}$ or $R^{3'}$ and $R^{4'}$ together form a **C3-10** cycloalkyl or heterocyclyl;

R^5 and R^6 are each independently H, **C1-6** alkyl, **C3-10** cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each **C1-6** alkyl, **C3-6** alkoxy, **C3-10** cycloalkyl, heterocyclyl, heteroaryl and aryl group of R^1 , R^2 , $R^{2'}$, R^3 , R^3 , R^4 , R^4 , R^5 and R^6 is optionally substituted with one, two or three R^{10} ;

35 R^{10} in each instance is independently halo, hydroxy, cyano, nitro, **C1-6** alkyl, **C1-6** alkoxy, **C2-6** aikenyl, **C2-6** alkynyl, **C3-10** cycloalkyl, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}$, $-S(=O)R^{11}$, $-S(=O)_2R^{11}$, $-NR^{11}R^{11}$, $-NR^{11}C(=O)R^{11}$ or $-NR^{11}C(=O)OR^{11}$;

wherein the C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl and heteroaryl of R^{10} are optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R^{11} in each instance independently is hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl of R^{11} is optionally substituted with one, two or three C_{1-6} alkyl;

provided the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate. Further the compound is not ethyl 3-phenylisoxazolidine-2-carboxylate, benzyl 3-(4-amino-2-oxopyrimidin-1(2H)-yl)isoxazolidine-2-carboxylate, 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one.

Also provided are compounds having structure (X), or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof, wherein:

A is $\sim C(=O)\sim$, $\sim S(=O)\sim$, $\sim S(=O)_2\sim$ or $\sim S(=O)(=NH)\sim$;

R^1 is $\sim NR^5R^6$, C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl, heterocyclyl or heteroaryl;

R^2 is aryl or heteroaryl;

$R^{2'}$ is hydrogen or R^2 and $R^{2'}$ together form a C_{3-10} cycloalkyl or heterocyclyl;

R^3 is hydrogen, C_{1-6} alkyl or C_{3-10} cycloalkyl;

$R^{3'}$ is hydrogen or R^3 and $R^{3'}$ together form a C_{3-10} cycloalkyl or heterocyclyl;

R^4 is hydrogen, halo, C_{1-6} alkyl or C_{3-10} cycloalkyl;

$R^{4'}$ is hydrogen or R^4 and $R^{4'}$ together form a C_{3-10} cycloalkyl or heterocyclyl;

or $R^{2'}$ and $R^{4'}$ or $R^{3'}$ and $R^{4'}$ together form a C_{3-10} cycloalkyl or heterocyclyl;

R^5 and R^6 are each independently H, C_{1-6} alkyl, C_{3-10} cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl and aryl group of R^1 , R^2 , $R^{2'}$, R^3 , $R^{3'}$, R^4 , R^4 , R^5 and R^6 is optionally substituted with one, two or three R^{10} ;


R^{10} in each instance is independently halo, hydroxy, cyano, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl, heteroaryl, $\sim C(=O)R^{11}$, $\sim C(=O)OR^{11}$, $\sim C(=O)NR^{11}R^{11}$, $\sim Si(=O)R^{11}$, $\sim S(=O)_2R^{11}$, $\sim NR^{11}R^{11}$, $\sim NR^{11}C(=O)R^{11}$ or $\sim NR^{11}C(=O)QR^{11}$;

wherein the C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl and heteroaryl of R^{10} are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R^{11} in each instance independently is hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl;

provided the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate. Further the compound is not ethyl 3-phenylisoxazolidine-2-carboxylate, benzyl 3-(4-amino-2-oxypyrimidin-1(2H)-yl)isoxazolidine-2-carboxylate, 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one.

Also provided herein are compounds having the following structure (Xa):



or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers

thereof, wherein:

A is -C(=O)-, -Si(=O)-, -S(=O)- or -Si(=O)(=NH)-;

R¹ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R² is aryl or heteroaryl;

R³ is hydrogen, deuterium, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R^{3'} is hydrogen or deuterium;

R⁴ is hydrogen, halo, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl and aryl group of R¹, R², R³, R^{3'}, R⁴, R⁵ and R⁶ is optionally substituted with one, two or three R¹⁰;

R¹⁰ in each instance is independently deuterium, halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₂₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl, heteroaryl, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹¹, -S(=O)R¹¹, -Si(=O)R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl,

heterocyclyl, aryl and heteroaryl of R¹⁰ are optionally substituted with one, two or three

substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl; and


R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl of R¹¹ is optionally substituted with one, two or three C₁₋₆ alkyl;

provided the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or 2,2-

dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one.

Also provided herein are compounds having the following structure (I):



or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof, wherein:

A is -C(=O)-, -S(=O)-, -S(=O)₂- or ~S(=O)(=NH)-;

R¹ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocycli or heteroiyl;

R² is aryl or heteroiyl;

R³ is hydrogen, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R⁴ is hydrogen, halo, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroiyl or heterocycli, wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocycli, heteroiyl and aryl group of R¹, R², R³, R⁴, R⁵ and R⁶ is optionally substituted with one, two or three R¹⁰;

R¹⁰ in each instance is independently halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycli, aryl, heteroiyl, ~C(=O)R¹¹, -C(=O)OR¹¹, ~C(=O)NR¹¹R¹¹, -S(=O)R¹¹, -S(=O)₂R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycli, aryl and heteroiyl of R¹⁰ are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycli, aryl or heteroiyl; and

R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocycli, heteroiyl or aryl;

provided the compound is not 2-{methylsulfonyl}-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or tert-butyl 5-(1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate.

In certain embodiments, the present disclosure is directed to a compound of structure (I), or a pharmaceutically acceptable salt, prodrug or stereoisomer thereof, wherein:

A is -C(=O)-, -S(=O)-, -S(=O)₂- or -S(=O)(=NH)-;

R¹ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocycli or heteroiyl;

R² is aryl or heteroiyl; and

R³ is hydrogen, C₁₋₅ alkyl or C₃₋₁₀ cycloalkyl;

R⁴ is hydrogen, halo, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R⁵ and R⁶ are each independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroiyl or


heterocycli,

wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl and aryl group is optionally substituted with one or more substituent;

provided herein, the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate.

Or alternatively, in certain embodiments, any C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl group may be optionally substituted with one or more substituent as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (II):



or a pharmaceutically acceptable salt, prodrug or stereoisomer thereof, wherein:

A is -C(=O)-, -S(=O)-, -S(=O)-, or -Si(=O)-; or -NH-;

R¹ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R² is aryl or heteroaryl;

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl and aryl group of R¹, R², R⁵ and R⁶ is optionally substituted with one, two or three R¹⁰;

R¹⁰ in each instance is independently halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl, heteroaryl, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹¹, -Si(=O)R¹¹, -S(=O)₂R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl and heteroaryl of R¹⁰ are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl;

provided the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine.

In certain embodiments, the present disclosure is directed to a compound of structure (II), or a pharmaceutically acceptable salt, prodrug or stereoisomer thereof, wherein:

A is -C(=O)-, -S(=O)-, -S(=O)-, or -Si(=O)-; or -NH-;

R¹ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R² is aryl or heteroaryl; and

R⁵ and R⁶ are each independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or

heterocyclyl,

wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl and aryl group is optionally substituted with one or more substituent;

provided the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine.

In certain embodiments, the present disclosure is directed to a compound of structure (II), or a pharmaceutically acceptable salt, prodrug or stereoisomer thereof, wherein:

A is $-\text{C}(=\text{O})\text{K}^+$, $-\text{S}(=\text{O})_2-$, $-\text{S}(=\text{O})_2-$ or $-\text{S}(=\text{O})(=\text{M})-$;


R¹ is $-\text{NR}^5\text{R}^6$, C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R² is aryl or heteroaryl; and


R⁵ and R⁶ are each independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each C₁₋₆ alkyl, C₃₋₆ alkoxy, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl and aryl group is optionally substituted with one or more substituent.

Compounds of structure (I) in enantiomerically enriched or enantiomerically pure form are also provided in various embodiments. Accordingly, in certain embodiments the compound has one of the following structures (I') and (II'):



where A, R¹, R², R³ and R⁴ are as defined herein. In certain embodiments the compound has one of the following structures (I'') and (II''):



where A, R¹, R², R³ and R⁴ are as defined herein.


In certain embodiments, the compound has structure (I'). In certain embodiments, the compound has structure (I''). In certain embodiments, the compound has structure (II'). In certain embodiments, the compound has structure (II'').

In certain embodiments, the compound is not 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(naphthalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate.

In certain embodiments, the compound is not ethyl 3-phenylisoxazolidine-2-carboxylate, benzyl 3-(4-amino-2-oxopyrimidin-1(2H)-yl)isoxazolidine-2-carboxylate, 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one. In certain embodiments, the compound is not 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one. In certain embodiments, the compound is not 2,2-dimethyl-1-


(3-phenylisoxazolidin-2-yl)butan-1-one. In certain embodiments, when R^1 is not pyrimidinyl substituted with one or more oxo and/or one or more hydroxy.

In certain embodiments, A is $-C(=O)-$ and the compound has the following structure (1a):




5 where R^1 and R^2 are as defined herein.

In certain embodiments, the compound has structure (1a'):




where R^1 and R^2 are as defined herein. In certain embodiments, the compound has structure (1a''):




10 where R^1 and R^2 are as defined herein.

In certain embodiments, A is $-S(=O)-$ and the compound has the following structure (1b):




where R^1 and R^2 are as defined herein.

15 In certain embodiments, the compound has structure (1b'):




where R^1 and R^2 are as defined herein. In certain embodiments, the compound has structure (1b''):




20 where R^1 and R^2 are as defined herein.

In certain embodiments, A is $-S(=O)_2-$ and the compound has the following structure (Ic):




where R^1 and R^2 are as defined herein.

In certain embodiments, the compound has structure (Ic'):




where R^1 and R^2 are as defined herein. In certain embodiments, the compound has structure (Ic''):




where R^1 and R^2 are as defined herein.

In certain embodiments, A is $-S(=O)(=NH)-$ and the compound has the following structure (Id):




where R^1 and R^2 are as defined herein.

In certain embodiments, the compound has structure (Id'):



where R^1 and R^2 are as defined herein. In certain embodiments, the compound has structure (Id''):



where R^1 and R^2 are as defined herein.

The R^1 and R^2 moieties in any of the structures described herein, are optionally substituted with one or more substituents. In certain embodiments, the R^1 and R^2 moieties in any of the structures described herein, are optionally substituted with one, two, three or four substituents. In certain embodiments, R^1 and R^2 are both substituted. In certain embodiments, R^1 and R^2 are both unsubstituted. In certain embodiments, R^1 is substituted and R^2 is unsubstituted. In certain embodiments, R^1 is

unsubstituted and R^2 is substituted. In certain embodiments, the optional substituents for R^1 and R^2 are selected from any of the optional substituents listed above **with** respect to the term "substituted." In certain embodiments, the optional substituents for R^1 and R^2 are selected from amino, hydroxy, **cyano**, halo (e.g., F, Cl or Br), optionally substituted C_{1-6} **alkyl**, optionally substituted C_{1-6} alkoxy, optionally substituted C_{1-6} haloalkyl and optionally substituted C_{1-6} **haloalkoxy**.


In certain embodiments, R^1 is substituted with at least one substituent selected from amino, hydroxy, cyano, halo (e.g., F, Cl or Br), optionally substituted C_{1-6} alkyl, optionally substituted C_{1-6} alkoxy, optionally substituted C_{1-6} haloalkyl and optionally substituted C_{1-6} haloalkoxy.

In certain embodiments, R^2 is substituted with at least one substituent selected from amino, hydroxy, cyano, halo (e.g., F, Cl or Br), optionally substituted C_{1-6} alkyl, optionally substituted C_{1-6} alkoxy, optionally substituted C_{1-6} haloalkyl and optionally substituted C_{1-6} haloalkoxy.

In certain embodiments, R^1 is C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-6} **alkylaminyl**, C_{3-10} **cycloalkyl**, heterocycli, heteroiyl or arylaminyl. In certain embodiments, R^1 is optionally substituted C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl or heterocycli. In certain embodiments, R^1 is optionally substituted C_{1-6} alkyl. In certain embodiments, R^1 is optionally substituted C_{10-20} alkyl.

In certain embodiments, R^1 is substituted with at least one substituent selected from deuterium, hydroxy, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano, C_{3-10} cycloalkyl, heterocycli and heteroiyl, wherein the heterocycli or heteroiyl is optionally further substituted with one, two or three C_{1-6} alkyl. In certain embodiments, R^1 is substituted with heterocycli or **heteroaryl**, wherein the heterocycli or **heteroaryl** is optionally further substituted with one, two or three substituents selected from deuterium, hydroxy, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano, C_{3-10} cycloalkyl, heterocycli and heteroaryl. In certain embodiments, R^1 is substituted with at least one substituent selected from hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy and cyano. In certain embodiments, R^1 is substituted with one, two, three or four substituents independently selected from hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy and cyano. In certain embodiments, R^1 is substituted with at least one substituent selected from hydroxy, alkoxy and cyano.

In certain embodiments, R^1 is optionally substituted branched C_{1-6} **alkyl**. In certain embodiments, R^1 is of the formula:



wherein each R^{2n} is independently hydrogen, halo, hydroxy, cyano, nitro, C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocycli, aryl, heteroaryl, $-C(=O)R^{26}$, $-C(=O)OR^{26}$, $-C(=O)NR^{26}R^{26}$, $-S(=O)R^{26}$, $-S(=O)_2R^{26}$, $-NR^{26}R^{26}$, $-NR^{26}C(=O)R^{26}$ or $-NR^{26}C(=O)OR^{26}$;

wherein the C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocycli, aryl and heteroiyl of R^{2n} are optionally substituted **with** one, two or three substituents independently selected from deuterium, halo,

hydroxy, **cyano**, amino, **nitro**, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ **haloalkoxy**, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ **cycloalkyl**, heterocyclyl, **aryl** or heteroaryl;

each R²⁵ is independently hydrogen, halo, C₁₋₆ alkyl, C₃₋₁₀ **cycloalkyl**, **heterocyclyl**, heteroaryl or aryl; or **two** R²⁵ together with the atom to which they are attached form a C₃₋₁₀ cycloalkyl or heterocyclyl;


5 wherein the C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl of R²⁵ is optionally-substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ **haloalkoxy**, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or **heteroaryl**; and

10 R²⁶ in each instance independently is hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the C₁₋₆ alkyl of R²⁶ is optionally substituted **with** halo or **oxo** and wherein the C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl of R²⁶ is optionally substituted **with** one, **two** or three C₁₋₆ alkyl;

provided that **at** least one of the following occurs: a) two R²⁵ together with the atom **to** which 15 they are attached form a C₃₋₁₀ cycloalkyl or heterocyclyl, b) at least one R²⁴ is other than hydrogen, or c) both R²⁵ are optionally substituted C₁₋₆ alkyl.

In certain embodiments, **two** R²⁵ together **with** the atom **to** which they are attached form a C₃₋₁₀ cycloalkyl or heterocyclyl. In certain embodiments, at least one R²⁴ is other than hydrogen. In certain 20 embodiments, both R²⁵ are optionally substituted C₁₋₆ alkyl. In certain embodiments, at least one R²⁵ is methyl. In certain embodiments, R¹ is of **the formula**:



wherein each R²⁴ is independently hydrogen, halo, hydroxy, cyano, nitro, C₁₋₆ alkoxy, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, **aryl**, **heteroaryl**, -C(=O)R²⁶, -C(=O)OR²⁶, -C(=O)NR²⁶R²⁶, -S(=O)R²⁶, -S(=O)₂R²⁶, -NR²⁶R²⁶, -NR²⁶C(=O)R²⁶ or -NR²⁶C(=O)OR²⁶;

25 wherein the C₁₋₆ alkoxy, C₂₋₆ alkynyl, C₃₋₁₀ **cycloalkyl**, heterocyclyl, aryl and heteroaryl of R²⁴ are optionally substituted **with** one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ **haloalkoxy**, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl; and

30 R²⁶ in each instance independently is hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the C₁₋₆ alkyl of R²⁶ is optionally substituted **with** halo or **oxo** and wherein the C₃₋₁₀ cycloalkyl, heterocyclyl, **heteroaryl** or aryl of R²⁶ is optionally substituted **with** one, **two** or three C₁₋₆ alkyl;


provided that at least one R²⁴ is other than hydrogen.

35 In certain embodiments, each R²⁴ is independently selected from hydrogen, deuterium, hydroxy, halo, C₁₋₆ **haloalkyl**, C₁₋₆ alkoxy, C₁₋₆ **haloalkoxy**, cyano, C₃₋₁₀ cycloalkyl, heterocyclyl and heteroaryl,


wherein the heterocyclyl or heteroaryl is optionally further substituted with one, two or three halo, cyano, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl. In certain embodiments, each R²⁴ is independently selected from hydrogen, hydroxy, halo, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy and cyano. In certain embodiments, each R²⁴ is independently hydrogen, halo or cyano. In certain embodiments, one R²⁴ is hydrogen and the other two

5 R^{2*} are halo.


In certain embodiments, R^1 may have one of the following structures:



In certain embodiments, R¹ may have one of the following structures:



In certain embodiments, R^1 has one of the following structures:




In certain embodiments, R^1 is optionally substituted C_{3-6} alkoxy. For example, in certain embodiments R^1 is t-butoxy.


- 5 In certain embodiments, R^1 is $-NR^3R^4$. In certain embodiments, R^1 is $-NR^3R^4$ and at least one of R^3 or R^4 is optionally substituted C_{5-6} alkyl or optionally substituted aryl. For example, in certain embodiments, R^1 is dimethylamino, t-butylamino or **phenylamino**.

- 10 In certain embodiments, R^1 is optionally substituted C_{3-10} **cycloalkyl**. For example, in some of these embodiments the C_{3-10} cycloalkyl is substituted with at least one substituent selected from hydroxy, halo, **cyano**, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{1-6} alkyl, C_{1-6} **haloalkyl**, C_{1-6} hydroxyalkyl and C_{1-6} alkoxyalkyl. In certain embodiments the C_{3-10} **cycloalkyl** is substituted **with** at least one substituent selected from hydroxy, halo, cyano, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{1-6} alkyl, C_{1-6} **haloalkyl**, C_{1-6} hydroxyalkyl and C_{1-6} alkoxyalkyl.


- 15 In certain embodiments, R^1 is C_{3-10} **cycloalkyl** and C_{3-10} **cycloalkyl** is **cyclopropyl**, **cyclobutyl**, bicyclo[1.1.1]pentyl. In certain embodiments, R^1 is C_{3-10} cycloalkyl and cycloalkyl is cyclopropyl or cyclobutyl. For example, in certain embodiments R^1 has one **of the** following structures:




In certain embodiments R^1 has one of the following structures:



In certain embodiments, R^1 has one of the following structures:




In some different embodiments, $\mathbb{R}^1$ has one of the following structures:



In certain embodiments, R¹ is optionally substituted heterocyclyl or optionally substituted heteroaryl. In certain embodiments, R¹ is oxetanyl, pyrrolidinyl, tetrahydropyranyl or azetidiny. In certain embodiments R¹ is oxetanyl or pyrrolidinyl. In certain embodiments, R¹ is pyrrolyl,


pyrazolopyridinyl or **benzoxoxazolyl**. In certain embodiments, R¹ is **heterocyclyl** or optionally substituted heteroaryl and R¹ is substituted **with** a substituent selected from **C₁₋₆ alkyl** and cyano. In certain embodiments, R¹ is heterocyclyl or optionally substituted **heteroaryl** and R¹ is substituted with a **C₁₋₆ alkyl**, cyano or both.

5 In certain embodiments, R^1 has one of the following structures:




a stereoisomer or mixture thereof.


In certain embodiments, R^1 has one of the following structures:




In certain embodiments, R^1 has one of the following structures:




In certain embodiments, R² is aryl, such as phenyl. In certain embodiments, R² is **heteroaryl**, such as **pyridinyl**. In certain embodiments, R² is unsubstituted. In certain embodiments R² is substituted, for example **with** one or more halo, cyano, C₁₋₆ alkyl, C₁₋₆ **haloalkyl**, C₁₋₆ alkoxy, -NH₂ or (34.10) **cycloalkyl**. In certain embodiments R² is substituted, for example with one or more halo atoms. In certain embodiments, R² has one of the following structures:







In certain embodiments, R^2 has one of the following structures:



In certain embodiments, R^2 has one of the following structures:



In certain embodiments, the present disclosure is directed to a compound having the following structure (III):



wherein:

each X^1 , X^2 , X^3 , X^4 and X^5 is either N or CR²¹;

R^3 is hydrogen, deuterium, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

$R^{3'}$ is hydrogen or deuterium;

R^{20} is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocycyl or heteroaryl;

wherein the C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocycyl or heteroaryl of R^{20} are optionally substituted with one, two, three or four substituents independently selected from deuterium, halo, hydroxy, cyano, amino, -S(=O)₂R¹¹, nitro, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycyl, aryl or heteroaryl optionally further substituted with C₁₋₆ alkyl; and

R^5 and R^6 are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocycyl;

each R^{21} is independently selected from hydrogen, halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycyl, aryl, heteroaryl, -C(=Q)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹¹, -S(=O)R¹¹, -S(=O)₂R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycyl, aryl and heteroaryl of R^{21} are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocycyl, aryl or heteroaryl; and

R^{11} in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocycyl, heteroaryl or aryl.

In certain embodiments, the present disclosure is directed to a compound of structure (III), wherein:

each X^1 , X^2 , X^3 , X^4 and X^5 is either N or CR²¹;

R³ is hydrogen, deuterium, C₁₋₆ alkyl or C₃₋₁₀ cycloalkyl;

R^{3'} is hydrogen or deuterium;

R²⁰ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

wherein the C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl of R²⁰ are optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, -S(=O)₂R¹¹, nitro, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl optionally further substituted with C₁₋₆ alkyl; and

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl;

each R²¹ is independently selected from hydrogen, halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl, heteroaryl, -C(=O)R¹¹,

-C(=O)OR¹¹, -C(=O)NR¹¹R¹¹, -S(=O)₂R¹¹, -S(=O)₂R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;


wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl,

heterocyclyl, aryl and heteroaryl of R²¹ are optionally substituted with one, two or three

substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IIIa):



wherein:

each X^1 , X^2 , X^3 , X^4 and X^5 is either N or CR²¹;

R²⁰ is -NR⁵R⁶, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl or heteroaryl;

R⁵ and R⁶ are each independently H, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, aryl, heteroaryl or heterocyclyl;

each R²¹ is independently selected from hydrogen, halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl, heteroaryl, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹¹, -S(=O)₂R¹¹, -S(=O)₂R¹¹, -NR¹¹R¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl and heteroaryl of R²¹ are optionally substituted with one, two or three substituents independently selected


from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, heterocyclyl, heteroaryl or aryl.

5 In certain embodiments, each of X¹, X², X³, X⁴ and X⁵ is CR²¹. In certain embodiments, at least one of X¹, X², X³, X⁴ or X⁵ is N. In certain embodiments, X¹ is N. In certain embodiments, X² is N. In certain embodiments, X³ is N. In certain embodiments, X⁴ or X⁵ is N. In certain embodiments, X⁵ is N. In certain embodiments, at least two of X¹, X², X³, X⁴ or X⁵ is N.

10 In certain embodiments, each R²¹ is independently hydrogen, halo, hydroxy, cyano, C₁₋₆ alkyl or C₁₋₆ alkoxy. In certain embodiments, at least one of X¹, X², X³, X⁴ and X⁵ is CR²¹ and R²¹ is halo.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IIIb) or (IIIc):



wherein R³, R^{3'}, R²⁰ and R²² are as defined herein;


15 **m** is 1 or 2; and

each R²² is independently selected from halo, hydroxy, cyano, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, **aryl**, **heteroaryl**, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹, -S(=O)₂R¹¹, -Si(=O)₂R¹¹, -NR¹¹, -NR¹¹C(=O)R¹¹ or -NR¹¹C(=O)OR¹¹;

20 wherein the C₁₋₆ alkyl, C₁₋₆ alkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ cycloalkyl, heterocyclyl, aryl and **heteroaryl** of R²² are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkoxy, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₁₀ **cycloalkyl**, heterocyclyl, aryl or heteroaryl; and


25 R¹¹ in each instance is independently hydrogen, C₁₋₆ alkyl, C₃₋₁₀ **cycloalkyl**, heterocyclyl, heteroaryl or aryl.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IIIb') or (IIIc'):




wherein **m**, R³, R^{3'}, R²⁰ and R²² are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IIIb'') or (IIIc''):




wherein m, R², R³, R²⁰ and R²² are as defined herein.

5 In certain embodiments, the present disclosure is directed to a compound having the following structure (IVa), (IVb) or (IVc):




wherein R²⁰ and R²² are as defined herein.

10 In certain embodiments, the present disclosure is directed to a compound having the following structure (IVa'), (IVb') or (IVc'):




wherein R²⁰ and R²² are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IVa''), (IVb'') or (IVc''):




wherein R²⁰ and R²² are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IVd), (IVe), (IVf), (IVg) or (IVh):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 5 structure (IVd'), (IVe'), (IVf'), (IVg') or (IVh'):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 10 structure (IVd''), (IVe''), (IVf''), (IVg'') or (IVh''):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 5 structure (IVi), (IVj), (IVk) or (IVl):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 structure (IVi'), (IVj'), (IVk'), or (IVl'):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 structure (IVi''), (IVj''), (IVk''), or (IVl''):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IVm), (IVn), (IVo), or (IVp):




5 wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound **having** the following structure (IVm'), (IVn'), (IVo'), or (IVp'):




10 wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (IVm''), (IVn''), (IVo''), or (IVp''):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 5 structure (V):




wherein m is 1 or 2; and R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 structure (V'):




wherein m , R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 structure (V''):




wherein m , R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (Va):




wherein R²⁰ and R²² are as defined herein.

5 In certain embodiments, the present disclosure is directed to a compound having the following structure (Va'):




wherein R²⁰ and R²² are as defined herein.

10 In certain embodiments, the present disclosure is directed to a compound having the following structure (Va''):




wherein R²⁰ and R²² are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (Vb):




15 wherein R²⁰ and R²² are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (Vb'):




wherein R^{20} and R^{22} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (Vb''):



5 wherein R^{20} and R^{22} are as defined herein.


In certain embodiments, the present disclosure is directed to a compound having the following structure (VI):



wherein m is 1 or 2; R^{20} and R^{22} are as defined herein; and


10 R^{23} is hydrogen, deuterium, C_{1-12} alkyl, $-C(=O)OC_{1-12}$ alkyl, or C_{3-10} cycloalkyl, wherein said alkyl and cycloalkyl are each independently optionally substituted with one or more oxo, halo, hydroxy, C_{1-6} alkoxy, or amino.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VI'):




15 wherein m , R^{20} , R^{22} and R^{23} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VI''):



20 wherein m , R^{20} , R^{22} and R^{23} are as defined herein.


In certain embodiments, the present disclosure is directed to a compound having the following structure (VII):



wherein m is 1 or 2; R^{20} and R^{22} are as defined herein; and


5 R^{23} is hydrogen, deuterium, C_{1-12} alkyl, $-C(=O)OC_{1-12}$ alkyl, or C_{3-10} cycloalkyl, wherein said alkyl and cycloalkyl are each independently optionally substituted with one or more oxo, halo, hydroxy, C_{1-6} alkoxy, or amino.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VII'):




10 wherein m , R^{20} , R^{22} and R^{23} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VII''):



15 wherein m , R^{20} , R^{22} and R^{23} are as defined herein.


In certain embodiments, the present disclosure is directed to a compound having the following structure (VIII):



wherein m is 1 or 2; R^{20} and R^{22} are as defined herein; and


R^{23} is hydrogen, deuterium, C_{1-12} alkyl, $-C(=O)OC_{1-12}$ alkyl, or C_{3-10} cycloalkyl, wherein said alkyl and cycloalkyl are each independently optionally substituted with one or more **oxo**, halo, hydroxy, C_{1-6} **alkoxy**, or amino.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VIII):



wherein m , R^{20} , R^{22} and R^{23} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following structure (VIII''):



wherein m , R^{20} , R^{22} and R^{23} are as defined herein.

In certain embodiments, each R^{22} is independently selected from halo, hydroxy, cyano, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-10} cycloalkyl, or heterocyclyl. In certain embodiments, each R^{22} is independently selected from halo. In certain embodiments, each R^{22} is independently selected from hydroxy. In certain embodiments, each R^{22} is independently selected from cyano. In certain embodiments, each R^{22} is independently selected from C_{1-6} alkyl. In certain embodiments, each R^{22} is independently selected from C_{3-10} cycloalkyl. In certain embodiments, each R^{22} is independently selected from heterocyclyl.


In certain embodiments of the compounds of any **structure** above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), each R^{22} is independently fluoro, methyl, or cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (HI)-(Vni) or a sub-structure thereof), one R^{22} is cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a **sub-structure** thereof), one R^{22} is fluoro. In certain embodiments, one R^{22} is methyl.

In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is optionally substituted C_{1-6} alkyl. In certain embodiments of the compounds of any **structure** above (e.g., any of structures (III)-(VIII) or a **sub-structure** thereof), R^{20} is C_{1-6} **cyanoalkyl** or C_{1-6} haloalkyl. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is C_{1-6} haloalkyl.

In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is C_{1-6} haloalkyl and, each R^{22} is independently fluoro, methyl, or cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is C_{1-6} haloalkyl and one R^{22} is fluoro. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is C_{1-6} haloalkyl and one R^{22} is cyano.

In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(Vni) or a sub-structure thereof), R^{20} is optionally substituted C_{3-10} cycloalkyl, or optionally substituted branched C_{1-6} alkyl.

In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), R^{20} is optionally substituted branched C_{1-6} alkyl. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VII) or a sub-structure thereof), R^{20} is of the formula:



wherein each R^{24} is independently hydrogen, halo, hydroxy, cyano, nitro, C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(C\equiv C)R^{24}$, $-C(=O)OR^{26}$, $-C(=O)NR^{26}R^{26}$, $-Si(=C)R^{26}$, $-S(=O)_2R^{26}$, $-NR^{26}R^{26}$, $-NR^{26}C(=O)R^{26}$ or $-NR^{26}C(=O)OR^{26}$;

wherein the C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl and heteroaryl of R^{24} are optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or heteroaryl;

each R^{25} is independently hydrogen, halo, C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl; or two R^{25} together with the atom to which they are attached form a C_{3-10} cycloalkyl or heterocyclyl;

wherein the C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl of R^{25} is optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R^{26} in each instance independently is hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl;


wherein the C_{1-6} alkyl of R^{26} is optionally substituted with halo or oxo and wherein the C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl of R^{26} is optionally substituted with one, two or three C_{1-6} alkyl;

provided that at least one of the following occurs: a) two R^{25} together with the atom to which they are attached form a C_{3-10} cycloalkyl or heterocyclyl, b) at least one R^{24} is other than hydrogen, or c) both R^{25} are optionally substituted C_{1-6} alkyl.

In certain embodiments of the compounds of any **structure** above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), two R^{25} together with the atom to which they are attached form a C_{3-10} cycioalkyl or **heterocyclyl**. In certain embodiments of the compounds of any structure above (e.g., any of **structures (IIT)-(VIII)** or a sub-structure thereof), at least one R^{34} is other than hydrogen. In certain

5 embodiments of the compounds of any structure above (e.g., any of stmcures (III)-(VIII) or a sub-**structure** thereof), both R^{25} are optionally substituted C_{1-6} **alkyl**.

In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(VIII) or a sub-structure thereof), **at** least one R^{25} is methyl. In certain embodiments of the compounds of any **structure** above (e.g., any of structures (III)-(VIII) or a **sub-structure thereof**), R^{20} is of the **formula**:



wherein each R^{34} is independently hydrogen, halo, hydroxy, cyano, **nitro**, C_{1-6} **alkoxy**, C_{2-6} alkynyl, C_{3-10} cycioalkyl, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{26}$, $-C(=O)OR^{26}$, $-C(=O)NR^{26}R^{26}$, $-S(=O)_2R^{26}$, $-S(=O)_2R^{26}$, $-NR^{26}R^{26}$, $-NR^{26}C(=O)R^{26}$ or $-NR^{26}C(=O)OR^{26}$;

wherein the C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} cycioalkyl, heterocyclyl, **aryl** and heteroaryl of R^{24} are

15 optionally substituted **with** one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} **haloalkoxy**, C_{2-6} **alkenyl**, C_{2-6} **alkynyl**, C_{3-10} cycioalkyl, heterocyclyl, aryl or **heteroaryl**; and

R^{26} in each instance independently is hydrogen, C_{1-6} alkyl, C_{3-10} cycioalkyl, heterocyclyl, heteroaryl or aryl;


wherein **the** C_{1-6} alkyl of R^{26} is optionally substituted with halo or oxo and wherein the C_{3-10} cycioalkyl, heterocyclyl, heteroaryl or aryl of R^{26} is optionally substituted with one, two or **three** C_{1-6} alkyl;

provided that at least one R^{24} is other than hydrogen.

In certain embodiments, each R^{24} is independently selected from hydrogen, deuterium, hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano, C_{3-10} cycioalkyl, heterocyclyl and heteroaryl, wherein the heterocyclyl or heteroaryl is optionally further substituted with one, two or three halo, cyano C_{1-6} alkyl or C_{3-10} cycioalkyl. In certain embodiments, each R^{24} is independently selected from hydrogen, hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy and cyano. In certain embodiments, each R^{24} is independently hydrogen, halo or cyano. In certain embodiments, one R^{24} is hydrogen and the other two

30 R^{24} are halo.

In certain embodiments, the present disclosure is directed to a compound having the following structure (TXa), (IXb), (IXc) or (IXd):



wherein:

R^{22} is halo, hydroxy, **cyano**, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} **alkenyl**, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{26}$, $-C(=G)QR^{26}$, $-C(=O)NR^{26}R^{26}$, $-S(=O)R^{26}$, $-S(=O)_2R^{26}$, $-NR^{26}R^{26}$, $-NR^{26}C(=O)R^{26}$ or $-NR^{26}C(=O)OR^{26}$;

wherein the C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} **cycloalkyl**, heterocyclyl, aryl and **heteroaryl** of R^{22} are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or **heteroaryl**;

wherein each R^{24} is independently hydrogen, halo, hydroxy, cyano, **nitro**, C_{1-6} alkoxy, C_{2-6} alkynyl, C_{3-10} **cycloalkyl**, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{26}$, $-C(=O)OR^{26}$, $-C(=O)NR^{26}R^{26}$, $-S(=O)R^{26}$, $-S(=G)_2R^{26}$, $-NR^{26}R^{26}$, $-NR^{26}C(=O)R^{26}$ or $-NR^{26}C(=O)OR^{26}$;

wherein the C_{1-6} alkoxy, C_{2-6} **alkynyl**, C_{3-10} cycloalkyl, heterocyclyl, aryl and **heteroaryl** of R^{24} are optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or **heteroaryl**;

each R^{25} is independently hydrogen, halo, C_{1-6} alkyl, C_{3-10} **cycloalkyl**, heterocyclyl, heteroaryl or **aryl**; or two R^{25} together with the atom to which they are attached form a C_{3-10} cycloalkyl or heterocyclyl;


wherein the C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, **heteroaryl** or aryl of R^{25} is optionally substituted with one, two or three substituents independently selected from deuterium, halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} **cycloalkyl**, heterocyclyl, **aryl** or heteroaryl; and

R^{26} in each instance independently is hydrogen, C_{1-6} alkyl, C_{3-10} **cycloalkyl**, heterocyclyl, heteroaryl or aryl;

wherein the C_{1-6} alkyl of R^{26} is optionally substituted with halo or oxo and wherein the C_{3-10} cycloalkyl, heterocyclyl, **heteroaryl** or **aryl** of R^{26} is optionally substituted with one, two or three C_{1-6} alkyl;


provided that at least one of the following occurs: a) two R^{25} together with the atom to which they are attached form a C_{3-10} cycloalkyl or heterocyclyl, b) at least one R^{24} is other than hydrogen, or c) both R^{25} are optionally substituted C_{1-6} alkyl.

In certain embodiments, the present disclosure is directed to a compound having the following
 5 structure (IXa'), (IXb'), (IXc') or (IXd'):



wherein R^{22} , R^{24} and R^{25} are as defined herein.

In certain embodiments, the present disclosure is directed to a compound having the following
 10 structure (IXa''), (IXb''), (IXc'') or (IXd''):



wherein R^{22} , R^{24} and R^{25} are as defined herein.

In certain embodiments of the compounds of structure (IX) or a sub-structure thereof, two R^{25}
 15 together with the atom to which they are attached form a C_{3-10} cycloalkyl or heterocyclyl. In certain
 embodiments of the compounds of structure (IX) or a sub-structure thereof, at least one R^{24} is other than
 hydrogen. In certain embodiments of the compounds of structure (IX) or a sub-structure thereof, both R^{25}
 are optionally substituted C_{1-6} alkyl.


In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), R^{22} is halo or cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), R^{22} is halo. In certain embodiments, R^{22} is **fluoro**. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), R^{22} is cyano.




In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), each R^{24} is independently selected from hydrogen, deuterium, hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano, C_{3-10} cycloalkyl, heterocyclyl and heteroaryl, wherein the heterocyclyl or heteroaryl is optionally further substituted with one, **two** or three C_{1-6} **alkyl**.




In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), each R^{24} is independently selected from hydrogen, hydroxy, halo, C_{1-6} **haloalkyl**, C_{1-6} alkoxy, C_{1-6} haloalkoxy and cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), each R^{24} is independently hydrogen, halo or cyano. In certain embodiments of the compounds of any structure above (e.g., any of structures (III)-(IX) or a **sub-structure** thereof), one R^{24} is hydrogen and the other two R^{24} are halo.




In certain embodiments, the compound is a compound selected from Table 1A, 1B or 1C. Also included within the disclosure are stereoisomers and mixtures of stereoisomers thereof.

Table 1A
























Table 1B




98




99




100




101




102




103




104




105




106




107




108




109



110



111



















112 Diastereoisomer 1


113 Diastereoisomer 2

114 Diastereoisomer 1


115 Diastereoisomer 2




134




140




146



135



147




141 Diastereoisomer 1


142 Diastereoisomer 2

136 Diastereoisomer 1


137 Diastereoisomer 2




143




148




138




144




149




139




145




150




151




156




161




152



157




162




163

Diastereoisomer 2
Diastereoisomer 1


153




158




164




154




159




165




155




160




166




167




172




177




168




173




178




169




174



170



175




179

Diastereoisomer 2


180

Diastereoisomer 1


171




176








181




182







190




191




192




193




194




195




196




197




198



199



200












































Table 1C








In certain embodiments, the compound of any structure or formula provided herein is provided in the form of a pharmaceutically acceptable salt. Exemplary salts for this purpose are described herein.

In certain embodiments, the disclosure provides a prodrug which converts to a compound of any structure provided herein *in vivo*. Exemplary prodrugs for this purpose are known in the art and described herein.

Enantiomerically enriched compositions of the compounds disclosed herein are also provided. In certain embodiments, the enantiomeric ratio of the composition is greater than 50:50. In certain embodiments, the enantiomeric ratio is calculated only with respect to a single stereocenter (e.g., R²) without regard to other stereocenters which may be present on the molecule. In certain embodiments, the composition comprises a single enantiomer of the compound and is substantially free (i.e., having less than or about 40%, 30%, 25%, 20%, 15%, 10%, 5%, 1%, 0.05%, or 0.01%) of the other enantiomer (or diastereomers).

3. General Synthesis

In certain embodiments, provided is a method of preparing a compound of structure (X), comprising coupling a compound of formula (A-1):




with a compound of formula (B-1):



under conditions to provide the compound of structure (X), wherein A, R¹, R², R³, R^{3'}, and R⁴ are as defined herein and LG is a leaving group.

5 In certain embodiments, provided is a method of preparing a compound of structure (Xa), comprising coupling a compound of formula (A-2):



with a compound of formula (B-2):




10 under conditions to provide the compound of structure (Xa), wherein A, R¹, R², R³, R^{3'}, and R⁴ are as defined herein and LG is a leaving group.

The following General Reaction Scheme I further illustrates a method of making compounds of formula (I). Other methods for preparing compounds of structure (I) and related intermediates are provided in the Examples and/or known in the art. It is understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. In general, starting components and reagents may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (see, for example, Advanced Organic Chemistry: Reactions, Mechanisms and Structure, 5th edition (Wiley, December 2000)) or prepared as described in this disclosure.

15
20

Scheme I



Referring to General Reaction Scheme 1, compounds of structure (I), wherein R¹ and R² are as defined herein and A is -(C=O)~, may be prepared from compound A or B. Appropriate compounds of structure A or B can be prepared according to the more specific methods described in the Examples which follow or by methods known to one of skill in the art. Treatment of compound A with an activating agent, such as methanesulfonyl chloride, followed by removal of the Boc protecting group under acidic conditions, results in oxazolidine C. Alternatively, compound C may be obtained by treatment of B with an appropriately substituted alkylene (e.g., ethylene gas) and an appropriate Lewis acid (e.g., BF₃). Treatment of C with an appropriate carbonyl compound (i.e., compound D, wherein X is

25
30

halo or OH) under amide coupling conditions yields the desired product. It should be noted that although General Reaction Scheme 1 depicts an embodiment wherein A is -C(=O)-, other compounds of structure (I) wherein A is -S(=O)-, -S(=O)₂- or -S(=O)(=NH)- can be made via analogous procedures by substituting compound D with an appropriate sulfur-based analogue.

When enantiomerically pure or enriched compounds are desired, dual chromatography and/or enantiomerically pure or enriched starting materials (e.g., compound A) may be used as described in the Examples. It will also be appreciated by those skilled in the art that in the process described herein the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (for example, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl and the like. Suitable protecting groups for amino, amido and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl and the like. Suitable protecting groups for mercapto include -C(0)-R" (where R" is alkyl, aryl or arylalkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. Protecting groups may be added or removed in accordance with standard techniques, which are known to one skilled in the art and as described herein. The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wutz, *Protective Groups in Organic Synthesis* (1999), 3rd Ed., Wiley. As one of skill in the art would appreciate, the protecting group may also be a polymer resin such as a Wang resin, Rink resin or a 2-chlorotrityl-chloride resin.

It will also be appreciated by those skilled in the art, although such protected derivatives of compounds of this disclosure may not possess pharmacological activity as such, they may be administered to a mammal and thereafter metabolized in the body to form compounds of the disclosure which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds of this disclosure are included within the scope of the disclosure. Further, the above general scheme can be applied to any other structure or compound as described herein.

Furthermore, all compounds of structure (I) or any other structure or compound as described herein which exist in free base or acid form can be converted to their pharmaceutically acceptable salts by treatment with the appropriate inorganic or organic base or acid by methods known to one skilled in the art. Salts of the compounds of the disclosure can be converted to their free base or acid form by standard techniques.

4. Methods of Treatment

"Treatment" or "treating" is an approach for obtaining beneficial or desired results including clinical results. Beneficial or desired clinical results may include one or more of the following: a) inhibiting the disease or condition (e.g., decreasing one or more symptoms resulting from the disease or condition and/or diminishing the extent of the disease or condition); b) slowing or arresting the development of one or more clinical symptoms associated with the disease or condition (e.g., stabilizing the disease or condition, preventing or delaying the worsening or progression of the disease or condition and/or preventing or delaying the spread (e.g., metastasis) of the disease or condition); and/or c) relieving

the disease, that is, causing the regression of clinical symptoms (e.g., ameliorating the disease state, providing partial or total remission of the disease or condition, enhancing effect of another medication, delaying the progression of the disease, increasing the quality of life and/or prolonging **survival**).

"Prevention" or "preventing" means any treatment of a disease or condition that causes the clinical **symptoms** of the disease or condition not to develop. Compounds may, in certain embodiments, be administered to a subject (including a human) who **is at risk** or has a family **history** of the disease or condition.

"Subject" refers to an animal, such as a mammal (including a human), that has been or will be the object of treatment, observation or experiment. The methods described herein may be useful in human therapy and/or veterinary applications. In certain embodiments, the subject is a mammal. In certain embodiments, the subject is a human.

The term "therapeutically effective amount" or "effective amount" of a compound described herein or a pharmaceutically acceptable **salt, tautomer**, stereoisomer, mixture of stereoisomers, prodrug or **deuterated** analog thereof means an amount sufficient to effect treatment when administered to a subject, to provide a therapeutic benefit such as amelioration of symptoms or slowing of disease progression. For example, a therapeutically effective amount may be an amount sufficient to decrease a symptom of a disease or condition of as described herein. The therapeutically effective amount may vary depending on the subject and disease or condition being treated, the weight and age of the subject, the severity of the disease or condition and the manner of administering, which can readily be determined by one of ordinary skill in the art.

The term "trauma" as used herein refers to any physical damage to the body caused by violence, accident, fracture **etc.** The term "ischemia" refers to a cardiovascular disorder characterized by a low oxygen state usually due to the **obstruction** of the arterial blood supply or inadequate blood flow leading to hypoxia in the tissue. The term "stroke" refers to cardiovascular disorders caused by a blood clot or bleeding in the brain, most commonly caused by an interruption in the flow of blood in the brain as from clot blocking a blood vessel and in certain embodiments of the disclosure the term stroke refers to ischemic stroke or hemorrhagic stroke. The term "myocardial infarction" refers to a cardiovascular disorder characterized by localized necrosis resulting from **obstruction** of the blood supply.

The methods described herein may be applied to cell populations *in vivo* or *ex vivo*. "*In vivo*" means within a living individual, as within an animal or human. In this context, the methods described herein may be used therapeutically in an individual. "*Ex vivo*" means outside of a living individual. Examples of *ex vivo* cell populations include *in vitro* cell cultures and biological samples including fluid or tissue samples obtained from individuals. Such samples may be obtained by methods well known in the art. Exemplary biological fluid samples include blood, cerebrospinal fluid, urine and saliva. In this context, the compounds and compositions described herein may be used for a variety of purposes, including therapeutic and experimental purposes. For example, the compounds and compositions described herein may be used *ex vivo* to determine the optimal schedule and/or dosing of administration of a compound of the present disclosure for a given indication, cell type, individual and other parameters.

information gleaned from such use may be used for experimental purposes or in the clinic to set protocols for *in vivo* treatment. Other *ex vivo* uses for which the compounds and compositions described herein may be suited are described below or will become apparent to those skilled in the art. The selected compounds may be further characterized to examine the safety or tolerance dosage in human or non-human subjects. Such properties may be examined using commonly known methods to those skilled in the art.

Experiments with knockout animal models and Necrostatin 1, a receptor interacting protein kinase 1 inhibitor, have demonstrated the effectiveness of receptor interacting protein kinase 1 inhibition in protecting tissues from inflammatory bowel diseases (e.g., ulcerative colitis and Crohn's disease), psoriasis, retinal-detachment-induced photoreceptor necrosis, retinitis pigmentosa, cerulein-induced acute pancreatitis and sepsis/systemic inflammatory response syndrome (SIRS) and alleviating ischemic brain injury, retinal ischemia/reperfusion injury, Huntington's disease, renal ischemia reperfusion injury, cisplatin induced kidney injury, traumatic brain injury, hematological and solid organ malignancies, bacterial infections and viral infections (e.g., tuberculosis and influenza) and lysosomal storage diseases.

The receptor interacting protein kinase 1 inhibitors of the present disclosure are therefore useful for treating diseases and conditions mediated by receptor interacting protein kinase 1, including but not limited to inflammatory diseases or disorders, necrotic cell diseases, neurodegenerative diseases, central nerve system (CNS) diseases, ocular diseases, infections and malignancies. In certain embodiments, the receptor interacting protein kinase 1 inhibitors described herein can inhibit inflammation, protect tissue or cell from damage or undesired cell death (e.g., necrosis or apoptosis), ameliorate symptoms and improve immune response in a patient suffering from any of the prescribed diseases or conditions. Moreover, the compounds may be suitable for treatment of immune-mediated disease, such as but not limited to, allergic diseases, autoimmune diseases and prevention of transplant rejection.

Necrotic Cell Diseases

The compounds described herein may be used for the treatment of diseases/disorders caused or otherwise associated with cellular necrosis. In particular, the disclosure provides methods for preventing or treating a disorder associated with cellular necrosis in a mammal, comprising the step of administering to said mammal a therapeutically effective amount of a compound or composition described herein. The term "necrotic cell disease" refers to diseases associated with or caused by cellular necrosis, for example trauma, ischemia, stroke, cardiac infarction, infection, Gaucher's disease, Krabbe disease, sepsis, Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease, HIV-associated dementia, retinal degenerative disease, glaucoma, age-related macular degeneration, rheumatoid arthritis, psoriasis, psoriatic arthritis or inflammatory bowel disease.

The necrotic cell diseases can be acute diseases such as trauma, ischemia, stroke, cardiac infarction, anthrax lethal toxin induced septic shock, sepsis, cell death induced by LPS and HIV induced T-cell death leading to immunodeficiency. In certain embodiments the disorder is an ischemic disease of organs including but not limited to brain, heart, kidney and liver.

The necrotic cell diseases also include chronic neurodegenerative diseases, such as Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, Alzheimer's disease, infectious encephalopathies, dementia such as HIV associated dementia.

In some different embodiments, the disorder is an ocular disorder such as retinal degenerative disease, glaucoma or age-related macular degeneration. In some different embodiments, the disorder is a central nervous system (CNS) disorder.

Inflammatory Diseases or Disorders

The receptor interacting protein kinase 1 inhibitors described herein may be used to treat inflammatory diseases and disorders. Inflammatory diseases and disorders typically exhibit high levels of inflammation in the connective tissues or degeneration of these tissues.

Non-limiting examples of inflammatory diseases and disorders include Alzheimer's disease, ankylosing spondylitis, arthritis including osteoarthritis, rheumatoid arthritis (RA), psoriasis, asthma, atherosclerosis, Crohn's disease, colitis, dermatitis, diverticulitis, fibromyalgia, hepatitis, irritable bowel syndrome (IBS), inflammatory bowel disease (IBD), systemic lupus erythematosus (SLE), nephritis, Parkinson's disease and ulcerative colitis. In certain embodiments, the compounds and compositions of the present disclosure are useful for treating an autoimmune disorder, such as rheumatoid arthritis, psoriasis, psoriatic arthritis, encephalitis, allograft rejection, autoimmune thyroid diseases (such as Graves' disease and Hashimoto's thyroiditis), autoimmune uveoretinitis, giant cell arteritis, inflammatory bowel diseases (including Crohn's disease, ulcerative colitis, regional enteritis, granulomatous enteritis, distal ileitis, regional ileitis, and terminal ileitis), insulin-dependent diabetes mellitus, multiple sclerosis, pernicious anemia, sarcoidosis, scleroderma, and systemic lupus erythematosus. In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein are useful for treating autoimmune encephalitis.

In certain embodiments, the compounds and compositions are useful for treating rheumatoid arthritis (RA). In certain embodiments, the compounds and compositions are useful for treating ulcerative colitis. In certain embodiments, the compounds and compositions are useful for treating psoriasis.

In certain embodiments, the disorder is an inflammatory disease of the intestines such as Crohn's disease or ulcerative colitis (both generally known together as inflammatory bowel disease). In certain embodiments, the mammal is a primate, canine or feline subject. In certain embodiments, the mammal is a human subject. While not wishing to be bound by theory-, it is believed that inhibition of receptor interacting protein kinase 1 by the presently disclosed compounds is responsible, at least in part, for their anti-inflammatory activity. Accordingly, embodiments of the disclosure also include methods for inhibiting receptor interacting protein kinase 1, either in vitro or in a subject in need thereof, the method comprises contacting a receptor interacting protein kinase 1 with a compound disclosed herein. In some of these embodiments, inhibiting receptor interacting protein kinase 1 is effective to block (partially or fully) the release of inflammatory mediators such as TNF and/or IL6.

Ocular Conditions

The receptor interacting protein kinase 1 inhibitors described herein can also be used to treat ocular conditions, for example to reduce or prevent the loss of photoreceptor and/or retinal pigment epithelial cell viability.

5 In one aspect, the disclosure provides a method of preserving the visual function of an eye of a subject with an ocular condition, wherein a symptom of the ocular condition is the loss of photoreceptor cell viability in the retina of the eye with the condition. The method comprises administering to the eye of the subject an effective amount of a compound or composition described herein, thereby preserving the viability of the photoreceptor cells disposed within the retina of the eye. After administration, the visual
10 function (e.g., visual acuity) of the eye may be preserved or improved relative to the visual function of the eye prior to administration.

The ocular condition may be a condition selected from the group consisting of age-related macular degeneration (AMD), retmosis pigmentosa (RP), macular edema, diabetic retinopathy, central areolar choroidal dystrophy, BEST disease, adult vitelliform disease, pattern dystrophy, myopic
15 degeneration, central serous retinopathy, Stargardt's disease, Cone-Rod dystrophy, North Carolina dystrophy, infectious retinitis, inflammatory retinitis, uveitis, toxic retinitis and light-induced toxicity. AMD may be the neovascular or the dry form of AMD. Retinal detachment may be a rhegmatogenous, a serous or a tractional retinal detachment.

In another aspect, the disclosure provides a method of preserving the viability of retinal pigment
20 epithelial (RPE) cells within the retina of a subject with an ocular condition, wherein a symptom of the ocular condition is the loss of retinal pigment epithelial cells in the retina of the eye with the condition. The method comprises administering to the eye of the subject an effective amount of a compound or composition described herein, thereby preserving the viability of the retinal pigment epithelial cells. The ocular condition may be selected from the group consisting of AMD, BEST disease, myopic
25 degeneration, Stargardt's disease, uveitis, adult foveomacular dystrophy, fundus flavimacuiatus, multiple evanescent white dot syndrome, serpiginous choroidopathy, acute multifocal posterior placoid epitheliopathy (AMPPE) and other uveitis disorders.

The ocular condition may be a condition selected from the group consisting of age-related macular degeneration (AMD), retinosis pigmentosa (RP), macular edema, diabetic retinopathy, central
30 areolar choroidal dystrophy, BEST disease, adult vitelliform disease, pattern dystrophy, myopic degeneration, central serous retinopathy, Stargardt's disease, Cone-Rod dystrophy, North Carolina dystrophy, infectious retinitis, inflammatory retinitis, uveitis, toxic retinitis and light-induced toxicity. Therefore, in certain embodiments, the method comprises administering to the eye an effective amount of a compound or composition described herein, thereby preseving the viability of the photoreceptor cells
35 disposed within the retina of the subject with a condition.

In another aspect, the disclosure provides a method of preserving the viability of photoreceptor cells disposed within a retina of a mammalian eye following retinal detachment. The method comprises administering a compound or composition described herein to the eye in which a region of the retina has

been detached in amounts sufficient to preserve the viability of photoreceptor cells disposed within the region of the detached retina.

In certain embodiments, the retinal detachment may be a rhegmatogenous retinal detachment, tractional retinal detachment or serous retinal detachment. In certain embodiments, the retinal detachment
5 may occur as a result of a retinal tear, retinoblastoma, melanoma or other cancers, diabetic retinopathy, uveitis, choroidal neovascularization, retinal ischemia, pathologic myopia or trauma.

In another aspect, the disclosure provides a method of preserving visual function of an eye of a subject with an ocular condition selected from the group consisting of AMD, RP, macular edema, central areolar choroidal dystrophy, retinal detachment, diabetic retinopathy, BEST disease, adult vitelliform
10 disease, pattern dystrophy, myopic degeneration, central serous retinopathy, Stargardt's disease, Cone-Rod dystrophy, North Carolina dystrophy, infectious retinitis, inflammatory retinitis, uveitis, toxic retinitis and light-induced toxicity, wherein a symptom of the ocular condition is the loss of photoreceptor cells viability in the retina of the eye, wherein the method comprises treating the subject with a compound or composition described herein to the subject.

In another aspect, the disclosure provides a method of preserving the visual function of an eye of
15 a subject with an ocular condition, wherein a symptom of the ocular condition is the loss of photoreceptor cell viability and/or RPE viability in the retina of the eye wherein the method comprises treating the subject with a compound or composition described herein to the subject.

In certain embodiments is provided a method of preserving the visual function of an eye of a
20 subject with ocular conditions, wherein a symptom of the ocular condition is the loss of retinal ganglion cell viability in the retina of the eye with the conditions. The method comprises administering to the eye of the subject an effective amount of a compound or composition, thereby preserving the viability of the retinal ganglion cells disposed within the retina of the eye. After administration of the compound or composition, the visual function of the eye may be preserved or improved relative to the visual function
25 of the eye prior to administration. Further, after the administration, the preserved retinal ganglion cell is capable of supporting axonal regeneration.

In each of the foregoing methods, the ocular condition, wherein a symptom of the condition is the loss of retinal ganglion cell viability in the retina of the eye, includes but is not limited to glaucoma, optic nerve injury, optic neuritis, optic neuropathies, diabetic retinopathy, central retinal artery occlusion
30 and central retinal vein occlusion. It is contemplated that the foregoing methods may be used for the treatment of optic neuropathies such as ischemic optic neuropathy (e.g., arteritic or non-arteritic anterior ischemic neuropathy and posterior ischemic optic neuropathy), compressive optic neuropathy, infiltrative optic neuropathy, traumatic optic neuropathy, mitochondrial optic neuropathy (e.g., Leber's optic neuropathy), nutritional optic neuropathy, toxic optic neuropathy and hereditary optic neuropathy (e.g.,
35 Leber's optic neuropathy, Dominant Optic Atrophy, Behr's syndrome).

Also disclosed is a method of preserving the visual function of an eye of a subject with an ocular condition selected from the group consisting of glaucoma, optic nerve injury, optic neuropathies, diabetic retinopathy, central retinal artery occlusion and central retinal vein occlusion. The method comprises

administering to the eye of the subject an effective amount of a compound or composition described herein, thereby preserving the viability of the retinal ganglion cells disposed within the retina of the eye and the visual function of the eye.

In another aspect, disclosed herein is a method of preserving the viability of retinal ganglion cells disposed within a retina of a mammalian eye affected by, for example, glaucoma, optic nerve injury, optic neuritis, optic neuropathies, diabetic retinopathy, central retinal artery occlusion and central retinal vein occlusion. The method comprises administering a compound or composition described herein to the eye in which a region of the retina has been affected in amounts sufficient to preserve the viability of retinal ganglion cells disposed within the region of the affected retina. The preserved retinal ganglion cell is capable of supporting axonal regeneration.

Also disclosed is a method for promoting axon regeneration in an eye of a subject with an ocular condition, wherein a symptom of the ocular condition is the loss of retinal ganglion cell viability in the retina of the eye with the condition. The method comprises administering to the eye of the subject an effective amount of a compound or composition described herein, thereby promoting axon regeneration of the retinal ganglion cell within the retina of the eye.

In each of the foregoing embodiments, it is understood that the methods and compositions described herein can be used to preserve the viability and/or promote axon regeneration of retinal ganglion cells during treatment of the underlying conditions including, but not limited to, glaucoma, optic nerve injury, optic neuritis, optic neuropathies, diabetic retinopathy, central retinal artery occlusion and central retinal vein occlusion.

Neurodegenerative and CNS Diseases

The receptor interacting protein kinase 1 inhibitors described herein may also be used to treat neurodegenerative diseases. Neurodegenerative diseases can affect many of the body's activities, such as balance, movement, talking, breathing and heart function. Neurodegenerative diseases can be genetic or caused by medical conditions such as alcoholism, tumors, strokes, toxins, chemicals and viruses.

Non-limiting examples of neurodegenerative diseases and CNS diseases include Niemann-Pick disease, type CI (NPCI), Alzheimer's disease, amyotrophic lateral sclerosis (ALS), Friedreich's ataxia, Huntington's disease, Lewy body disease, Parkinson's disease and spinal muscular atrophy.

In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein may be used to treat NPCI via inhibiting necroptosis that causes neuronal loss. In certain embodiments, the compounds and compositions of the present disclosure are useful for treating Alzheimer's disease. In certain embodiments, the compounds and compositions of the present disclosure are useful for treating Parkinson's disease. In certain embodiments, the compounds and compositions of the present disclosure are useful for treating amyotrophic lateral sclerosis (ALS).

More generally, the receptor interacting protein kinase 1 inhibitors described herein can be used to preserve neuron viability and promote axon growth and nerve functions within the central nervous system (CNS). Accordingly, the compounds may be used to reduce or even reverse the loss of cognitive,

motor and sensor}' functions associated with a CNS disease or disorder, by **preserving** neuron **viability** and/or promoting axon regeneration and/or **nerve functions**.

The receptor interacting protein kinase 1 inhibitors described herein can be used in a method for promoting axon regeneration in a CNS neuron, such as a CNS **sensory** neuron, a motor neuron, a cortical neuron, a cerebellar neuron, a **hippocampal** neuron and a midbrain neuron. The receptor interacting protein kinase I inhibitors described herein can be used in a method for promoting nerve function or **preserving** the viability following injur}' to a CNS neuron. In certain embodiments, these compounds can be used to promote regeneration of an axon in a CNS neuron **that** is degenerated in the CNS disease or disorder. The receptor interacting protein kinase 1 inhibitors may be administered **by** any conventional means, such as locally **to** the neuron or applied *ex vivo* before **re-implantation**.

Accordingly, in one aspect, the disclosure provides a method of treating a CNS disorder in a subject in need thereof, wherein a symptom of the CNS disorder is axon degeneration or injury within a CNS neuron. The method comprises administering to **the** subject an effective amount of a compound or composition disclosed herein thereby to promote regeneration of an axon in a CNS neuron affected **by** the CNS disorder. Following administration, neural functions may be measured, for example, as an indication of axon regeneration. It is also contemplated that, following administration of the compound or composition, the neuron function of the CNS neuron is preserved or improved relative **to** the neuron function **prior to** administration.

The CNS disorder includes, but is not limited **to**, brain injur}', spinal cord injury, dementia, stroke, Alzheimer's disease, amyotrophic lateral sclerosis (ALS/Lou Gehrig's Disease), Parkinson's disease, Huntington's disease, multiple sclerosis, diabetic neuropathy, polyglutamine (**polyQ**) diseases, stroke, **Fahr** disease, **Menke's** disease, Wilson's disease, cerebral ischemia and a prion disorder. In exemplary embodiments, the CNS disorder is brain **injury** or spinal cord **injury**.

Also provided herein are methods for promoting neuron survival and axon regeneration in the CNS. CNS disorders characterized by impaired or failing axon growth or axon degeneration may arise from CNS neuron **injury** (e.g., **trauma**, **surgery**, **nerve compression**, **nerve** contusion, nerve **transection**, neurotoxicity or other physical injur}'- **to** the brain or spinal cord) or neurodegenerative CNS disease, wherein a symptom of the disorder is axon degeneration (e.g., Alzheimer's disease, amyotrophic lateral sclerosis (ALS/Lou Gehrig's Disease), Parkinson's disease, multiple sclerosis, diabetic neuropathy, polyglutamine (polyQ) diseases, stroke, Fahr disease, Menke's disease, Wilson's disease, cerebral ischemia, prion disorder (e.g., **Creutzfeldt-Jakob** disease). In certain embodiments, the CNS disorder is brain injur}' (e.g., traumatic brain injury) or spinal cord injury (e.g., chronic, acute or traumatic spinal cord injury). In certain embodiments, the CNS disorder affects a subject's basic **vital** life functions such as breathing, heart beat and blood pressure, e.g., an **injury** to or aneurysm in the **brain** stem.

In certain embodiments, the CNS disorder affects a subject's cognitive ability, such as, brain injury to the cerebral cortex or a neurodegenerative CNS disorder, such as, Alzheimer's disease, **frontotemporal** dementia, dementia with **Lewy** bodies, corticobasal degeneration, progressive supranuclear palsy and prion disorders.

In certain embodiments, the CNS disorder affects a subject's movement and/or strength, such as injury to the brain or spinal cord or a neurodegenerative CNS disorder such as Parkinson's disease, frontotemporal dementia, dementia with Lewy bodies, corticobasal degeneration, progress supranuclear palsy, Huntington's disease, multiple system atrophy, amyotrophic lateral sclerosis and hereditary spastic paresis.

In certain embodiments, the CNS disorder affects a subject's coordination, such as brain injury to the cerebellum or a neurodegenerative CNS disorder such as spinocerebellar atrophies, Friedreich's ataxia and prion disorders.

In each of the foregoing methods, the CNS disorder includes, but is not limited to, brain injury, spinal cord injury, Alzheimer's disease, amyotrophic lateral sclerosis (ALS/Lou Gehrig's Disease), Parkinson's disease, multiple sclerosis, diabetic neuropathy, polyglutamine (polyQ) diseases, stroke, Falir disease, Menke's disease, Wilson's disease, cerebral ischemia, a prion disorder (e.g., Creutzfeldt-Jakob disease), dementia (e.g., frontotemporal dementia, dementia with Lewy bodies), corticobasal degeneration, progressive supranuclear palsy, multiple system atrophy, hereditary spastic paraparesis and spinocerebellar atrophies.

Tissue Injuries or Damages

The ability of the compounds described herein to inhibit inflammation and cell death makes them suitable for ameliorating tissue injuries or damages. The tissue injuries or damages may be a result of any of the diseases or conditions described above. For example, the compounds may be used for amelioration of brain tissue injury or damage following ischemic brain injury or traumatic brain injury or for amelioration of heart tissue injury or damage following myocardial infarction or for amelioration of brain tissue injury or damage associated with Huntington's disease, Alzheimer's disease or Parkinson's disease or for amelioration of liver tissue injury or damage associated with non-alcohol steatohepatitis, alcohol steatohepatitis, autoimmune hepatitis autoimmune hepatobiliary diseases or primary sclerosing cholangitis or for the amelioration of liver tissue injury or damage associated with overdose of acetaminophen or for amelioration of kidney tissue injury or damage following renal transplant or the administration of nephrotoxic drugs or substances.

Non-limiting examples of brain injury or damage include stroke (e.g., hemorrhagic and non-hemorrhagic), traumatic brain injury (TBI), cerebral hemorrhage, subarachnoid hemorrhage, intracranial hemorrhage secondary to cerebral arterial malformation, cerebral infarction, perinatal brain injury, non-traumatic brain injury, Alzheimer's disease, Parkinson's disease, Huntington's disease, multiple sclerosis, amyotrophic lateral sclerosis, brain hemorrhage, brain infections, brain tumor, subclinical brain injury, spinal cord injury, anoxic-ischemic brain injury, focal cerebral ischemia, global cerebral ischemia, and hypoxic hypoxia.

In an embodiment, the compounds and compositions of the present disclosure may be used to treat peritoneal tissue injury. Non-limiting examples of peritoneal tissue injury include peritoneal deterioration, peritoneal sclerosis, and peritoneal cancer. For example, the receptor interacting protein

kinase 1 inhibitors described herein may be used to treat peritoneal damage caused by peritoneal dialysis fluid (PDF) and **PD-related** side effects.

Liver Injury and Diseases

In an embodiment, the compounds and compositions of the present disclosure may be used to treat liver **injury** and diseases. Non-limiting examples of liver **injury** or damage include not only degeneration or necrosis **of liver** parenchyma cells which results from **injury** caused by a certain factor, but also undesirable phenomena caused by biological reactions **to the injury**, such as mobilization, infiltration, activation of Kupffer cells, leukocytes and the like, fibrosis **of the** liver tissue, etc., which reactions occur alone or **in** combination. In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein may be used **to** treat steatohepatitis and hepatocellular carcinoma via inhibiting receptor interacting protein kinase 1 activity-dependent apoptosis **of hepatocytes** and **hepatocarcinogenesis**. In an embodiment, the receptor interacting protein kinase 1 **inhibitors** described herein may be used to treat acute cholestasis and liver injury.

Kidney Injury and Diseases

In an embodiment, the compounds and compositions of the present disclosure may be used to treat kidney injury and diseases. Non-limiting examples of kidney diseases include chronic kidney disease (**CKD**) (e.g., glomerular diseases, tubulointerstitial diseases, **obstruction**, polycystic kidney-disease), acute kidney **injury (AKI)**, diabetic nephropathy, **glomerulonephritis**, focal glomerulosclerosis, immune complex nephropathy or lupus nephritis. Kidney disease may be caused by drug-induced renal injury or kidney graft rejection. Kidney disease may be characterized as nephrotic syndrome or renal insufficiency. In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein may be used to treat kidney diseases (e.g., AKI) via inhibiting cell death pathway in kidney diseases. In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein may be used **to** treat patient with kidney stones and **to** prevent **crystal-induced** cytotoxicity and acute kidney injury via inhibiting receptor interacting protein kinase 3-**MLKL-mediated necroptosis**.

Malignancies

In an embodiment, the compounds and compositions of the present disclosure are useful for treating malignancies/cancers such as carcinoma, sarcoma, melanoma, lymphoma **or** leukemia. **Non-limiting** examples of malignancies suitably treated **by** the receptor interacting protein kinase 1 inhibitors described herein include lung cancer (e.g. non-small cell lung cancer, small-cell lung cancer), hepatocellular cancer, melanoma, pancreatic cancer, urological cancer, bladder cancer, colorectal cancer, colon cancer, breast cancer, prostate cancer, renal cancer, thyroid cancer, gall bladder cancer, peritoneal cancer, ovarian cancer, cervical cancer, gastric cancer, endometrial cancer, esophageal cancer, head and neck cancer, neuroendocrine cancer, CNS cancer, brain tumors (e.g., glioma, anaplastic oligodendroglioma, adult glioblastoma multiforme, and adult anaplastic astrocytoma), bone cancer, soft tissue sarcoma, retinoblastomas, neuroblastomas, peritoneal effusions, malignant pleural effusions, mesotheliomas, Wilms tumors, trophoblastic neoplasms, hemangiopericytomas, Kaposi's sarcomas, myxoid carcinoma, round cell carcinoma, squamous cell carcinomas, esophageal squamous cell

carcinomas, oral carcinomas, vulval cancer, cancers of the adrenal cortex, ACTH producing tumors, lymphoma, and leukemia.

Infectious Diseases

In an embodiment, the compounds and compositions of the present disclosure are useful for treating infectious diseases resulting from the presence of pathogenic agents, including pathogenic viruses, pathogenic bacteria, fungi, protozoa, multicellular parasites and aberrant proteins known as prions. Non-limiting examples of infectious diseases suitably treated by the receptor interacting protein kinase 1 inhibitors described herein include virus infectious diseases and bacterial infectious diseases. The viais infectious disease is not particularly limited and includes, for example, infectious diseases with respiratory infectious viruses (e.g., infectious diseases due to **respiratory** infectious **viruses** such as influenza vims, rhino virus, corona virus, parainfluenza vims, RS virus, adeno vims, reo vims and the like), herpes zoster caused by **herpes** virus, diarrhea caused by **rotavirus**, viral hepatitis, AIDS and **the** like. The bacterial infectious disease is not particularly limited and includes, for example, infectious diseases caused by *Bacillus cereus*, *Vibrioparaahaemolyticus*, *Enterohemorrhagic Escherichia coli*, *Staphylococcus aureus*, MRSA, *Salmonella*, *Botulinus*, *Candida* and the like.

Bone Diseases

In an embodiment, the compounds and compositions of the present disclosure are useful for treating bone diseases that may result from a bone remodeling disorder whereby the balance between bone formation and bone resorption is shifted. **Non-limiting** examples of bone remodeling disorders include osteoporosis, Paget's disease, osteoarthritis, rheumatoid arthritis, achondroplasia, **osteochodrytis**, hypeø arathyroidism, osteogenesis imperfecta, congenital hypophosphatasia, fribromatous lesions, fibrous displasia, multiple myeloma, abnormal bone turnover, osteolytic bone disease and periodontal disease. Additional examples of bone diseases suitably treated by the receptor interacting protein kinase 1 inhibitors described herein include bone fracture, bone trauma, or a bone deficit condition associated with post-traumatic bone **surgery**, post-prosthetic joint **surgery**, post-piastic bone **surgery**, post-dentai **surgery**, bone chemotherapy treatment or bone radiotherapy treatment. Additional examples of diseases affecting bone or bone joints suitably treated by the receptor interacting protein kinase 1 inhibitors described herein include metastatic bone cancer, rheumatic diseases such as rheumatoid arthritis, **osteoarthritis** and other **inflammatory arthropathies**. In an embodiment, the receptor interacting protein kinase 1 inhibitors described herein may be used to treat postmenopausal osteoporosis via inhibiting **osteocyte necroptosis** and trabecular deterioration.

Cardiovascular Diseases

In an embodiment, the compounds and compositions **of** the present disclosure are useful **for** treating cardiovascular diseases that may be relate to the cardiovascular disorders of fragile plaque disorder, occlusive disorder and stenosis. Non-limiting cardiovascular diseases include **coronary artery** disorders and peripheral arterial disorders, including, among others, atherosclerosis, arterial occlusion, aneurysm **formation**, thrombosis, post-traumatic aneurysm **formation**, restenosis, and post-operative graft occlusion. It is believed that atherosclerosis results from maladaptive inflammation driven primarily **by**

macrophages. Thus, the compounds and compositions of the present disclosure may be used to treat atherosclerosis via inhibiting macrophage necroptosis.

Transplantation

In an embodiment, the compounds and compositions of the present disclosure are useful for treating transplant patients. Non-limiting examples of transplant patient suitably treated by the receptor interacting protein kinase 1 inhibitors described herein include patients with solid and non-solid organ and tissue transplantations and transplants, such as liver, heart, kidney, and heterologous and autologous bone marrow transplantations/transplants. Typically, immunosuppressive therapy is used to avoid graft rejection in recipients of solid organ transplants. Recipients of bone marrow transplants are usually subjected to extensive irradiation and chemotherapy prior to transplantation. It is believed that receptor interacting protein kinase 1 and NF- κ B signaling in dying cells determines cross-priming of CD8⁺ T cells. Thus, the receptor interacting protein kinase 1 inhibitors described herein may be used to treat transplant patient and avoid graft rejection by modulating cross-priming of CD8⁺ T cells.

Other Diseases and Conditions

Additional examples of diseases and disorders suitably treated by the receptor interacting protein kinase 1 inhibitors described herein include Gaucher disease, organ failure, pancreatitis, atopic dermatitis, spondylarthritis, gout, systemic onset juvenile idiopathic arthritis (SoJIA), systemic lupus erythematosus (SLE), Sjogren's syndrome, systemic scleroderma, anti-phospholipid syndrome (APS), vasculitis, primary sclerosing cholangitis (PSC), acetaminophen toxicity, kidney damage/injury (nephritis, renal transplant, surgery, administration of nephrotoxic drugs e.g. cisplatin, acute kidney injury(AKI)), Celiac disease, autoimmune idiopathic thrombocytopenic purpura (autoimmune ITP), cerebrovascular accident (CVA, stroke), myocardial infarction (MI), allergic diseases (including asthma), diabetes, Wegener's granulomatosis, pulmonary sarcoidosis, Behcet's disease, interleukin-1 converting enzyme (ICE/caspase-1) associated fever syndrome, chronic obstructive pulmonary disease (COPD), tumor necrosis factor receptor-associated periodic syndrome (TRAPS), periodontitis, NEMO-deficiency syndrome (F-kappa-B essential modulator gene (also known as IKK gamma or IKKG) deficiency syndrome), HOIL-1 deficiency ((also known as RBCK1) heme-oxidized IRP2 ubiquitin 3igase-1 deficiency), linear ubiquitin chain assembly complex (LUBAC) deficiency syndrome, hematological and solid organ malignancies, bacterial infections and viral infections (e.g., tuberculosis and influenza) and lysosomal storage diseases.

Non-limiting examples of lysosomal storage diseases include Gaucher disease, GM2 Gangliosidosis, alpha-mannosidosis, aspartylglucosaminuria, cholesteryl ester storage disease, chronic hexosaminidase A deficiency, cystinosis, Danon disease, Fabry disease, Farber disease, fucosidosis, galactosialidosis, GM1 gangliosidosis, mucopolidosis, infantile free sialic acid storage disease, juvenile hexosaminidase A deficiency, Krabbe disease, lysosomal acid lipase deficiency, metachromatic leukodystrophy, mucopolysaccharidoses disorders, multiple sulfatase deficiency, Niemann-Pick disease, neuronal ceroid lipofuscinoses, Pompe disease, pycnodysostosis, Sandhoff disease, Schindler disease, sialic acid storage disease, Tay-Sachs and Wolman disease.

In certain embodiments, provided are compounds and compositions for use in medicine. In certain embodiments, **the** compounds and compositions are for use in the treatment of a receptor interacting protein kinase 1-mediated disease or disorder. Also provided is a method of treating a receptor interacting protein kinase 1-mediated disease or disorder comprising administering a therapeutically effective amount of a compound or pharmaceutical composition disclosed herein to a subject in need thereof.

5. Compositions

For the purposes of administration, the **compounds** of the present disclosure may be administered as a raw chemical or may be **formulated** as pharmaceutical compositions. **Pharmaceutical** compositions of the present disclosure comprise a compound of **structure (I)** and a pharmaceutically acceptable carrier, diluent or excipient. Accordingly, different embodiments are directed to pharmaceutical compositions comprising any one or more of the foregoing compounds of structure **(I)** or a pharmaceutically acceptable salt, **prodrug** or stereoisomer thereof and a **pharmaceutically** acceptable carrier, diluent or excipient are also provided in various embodiments.

The pharmaceutical compositions of the present disclosure may be specially formulated for administration **in** solid or liquid form, including those adapted for the following: oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual and systemic **absorption**, boluses, powders, granules, pastes for application to **the** tongue; parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension or sustained-release formulation; topical application, for example, as a cream, ointment or a **controlled-release** patch or spray applied to the skin; **intravaginally** or intrarectally, for example, as a **pessary**, cream or foam; **sublingually**; ocularly; **transdermally**; or nasally, **pulmonary** and to other mucosal surfaces.

The phrase "pharmaceutically acceptable" is employed herein **to** refer to those compounds, materials, compositions and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "**pharmaceutically**-acceptable carrier" as used herein means a pharmaceutically - acceptable material, composition **or** vehicle, such **as** a liquid or solid filler, diluent, excipient or solvent encapsulating material, involved in **carrying** or transporting the subject compound from one organ or portion **of the** body, to another organ or portion **of the** body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients **of** the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically -acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose and its derivatives, such as sodium **carboxymethyl** cellulose, ethyl cellulose and cellulose acetate; surfactants, such as **polysorbate** 80 (i.e., Tween 80); powdered tragacanth; malt; gelatin; talc; **excipients**, such as cocoa butter and suppository waxes; oils, such as peanut **oil**, cottonseed **oil**, safflower **oil**, sesame **oil**,

olive oil, com oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; pH buffered solutions; polyesters, polycarbonates and/or

5 polyanhydrides; and other non-toxic compatible substances employed in pharmaceutical formulations.

Examples of such formulations include, but are not limited to DMSO, 10 mM DMSO, 8%

hydroxypropyl-beta-cyclodextrin in PBS, propylene glycol, etc. For example, in a certain embodiment the compounds of the disclosure can be used as 4 mM solution in 8% hydroxypropyl-beta-cyclodextrin in PBS for parenteral administration. In another certain embodiments, the compounds of the disclosure can
10 be used as a suspension in 0.5% aqueous CMC containing 0.1% TWEEN 80.

As set out herein, certain embodiments of the present compounds may contain a basic functional group, such as amino or methylamino (NCH_3) and are, thus, capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable acids. The term "pharmaceutically-acceptable salts" in this respect refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of
15 the present disclosure. These salts can be prepared in situ in the administration vehicle or the dosage form manufacturing process or by separately reacting a purified compound of the disclosure in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed during subsequent purification. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate,
20 maleate, fumarate, succinate, tartrate, naphylate, mesylate, glucoheptonate, lactobionate and laurylsulphonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", J. Pharm. Sci. 66:1-19.)

The pharmaceutically acceptable salts of the subject compounds include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from non-toxic organic or inorganic
25 acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluene sulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic and the
30 like.

In other cases, the compounds of the present disclosure may contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically-acceptable salts with pharmaceutically-acceptable bases. The term "pharmaceutically-acceptable salts" in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present
35 disclosure. These salts can likewise be prepared in situ in the administration vehicle or the dosage form manufacturing process or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation, with ammonia or with a pharmaceutically-acceptable organic primary, secondary or tertiary

amine. Representative alkali or alkaline earth salts include the lithium, sodium, potassium, calcium, magnesium and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include **ethylamine**, **diethylamine**, ethylenediamine, ethanolamine, diethanolamine, **piperazine** and the like.

5 Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and **perfuming** agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically -acceptable antioxidants include: water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the
10 like; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol and the like; and metal chelating agents, such as citric acid, **ethylenediamine** tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid and the like.

Formulations of the present disclosure include those suitable for oral, nasal, topical (including
15 buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will **vary** depending upon the host being treated, the **particular** mode of administration. **The** amount of active ingredient that can be combined with a carrier material to produce a
20 single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, this amount **will** range from about 1% to about 99% of active ingredient, preferably from about 5% to about 70%, most preferably from about 10% **to** about 30%.

In certain embodiments, a formulation of the present disclosure comprises an **excipient** selected from **the** group consisting of cyclodextrins, liposomes, micelle forming agents, e.g., bile acids and
25 polymeric carriers, e.g., polyesters and **polyanhydrides**; and a compound of the present disclosure. **In** certain embodiments, an aforementioned formulation renders orally bioavailable a compound of the present disclosure.

Methods of preparing these formulations or compositions include the step of bringing into association a compound of the present disclosure with **the** carrier and, optionally, one or more accessory
30 ingredients. In general, the formulations are prepared by uniformly and intimately bringing **into** association a compound of the present disclosure with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product-

Formulations of the disclosure suitable **for** oral administration may be **in** the form **of** capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or **tragacanth**),
35 powders, granules or as a solution **or** a suspension in an aqueous or **non**-aqueous liquid or as an oil-in-water or **water-in-oil** liquid emulsion or as an elixir or syrup or as pastilles (using an inert base, such as gelatin and glycerin or sucrose and acacia) and/or as mouth washes and the like, each containing a

predetermined amount of a compound of the present disclosure as an active ingredient. A compound of the present disclosure may also be administered as a bolus, electuary or paste.

In solid dosage forms of the disclosure for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically -
5 acceptable carriers, such as sodium citrate or dicalcium phosphate and/or any of the following: fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol and/or silicic acid; binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; humectants, such as glycerol; disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; solution retarding agents, such as
10 paraffin; absorption accelerators, such as quaternary ammonium compounds; wetting agents, such as, for example, cetyl alcohol, glycerol monostearate and non-ionic surfactants; absorbents, such as kaolin and bentonite clay; lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate and mixtures thereof; and coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar
15 type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory-ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium
20 starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made in a suitable machine in which a mixture of the powdered compound is moistened with an inert liquid diluent.

The tablets and other solid dosage forms of the pharmaceutical compositions of the present disclosure, such as dragees, capsules, pills and granules, may optionally be scored or prepared with
25 coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical -formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through
30 a bacteria-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved in sterile water or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include
35 polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

Liquid dosage forms for oral administration of the compounds of the disclosure include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In

addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and
5 sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for
10 example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth and mixtures thereof.

Formulations of the pharmaceutical compositions of the disclosure for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the disclosure with one or more suitable nonirritating excipients or carriers comprising, for
15 example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present disclosure which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are
20 known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of a compound of this disclosure include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier and with any preservatives, buffers or propellants which may be required.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this disclosure, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide or mixtures thereof.

Powders and sprays can contain, in addition to a compound of this disclosure, excipients such as
30 lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the present disclosure to the body. Dissolving or dispersing the compound in the proper medium can
35 make such dosage forms. Absorption enhancers can also be used to increase the flux of the compound across the skin. Either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel can control the rate of such flux.

Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this disclosure.

Pharmaceutical compositions of this disclosure suitable for parenteral administration comprise one or more compounds of the disclosure in combination with one or more pharmaceutically -acceptable
5 sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain sugars, alcohols, antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers, which may be employed in the
10 pharmaceutical compositions of the disclosure include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like) and suitable mixtures thereof, vegetable oils, such as olive oil and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

15 These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms upon the subject compounds may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenyl sorbic acid and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride and the like into the compositions. In addition, prolonged
20 absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of
25 the drug then depends upon its rate of dissolution, which in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer
30 and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions, which are compatible with body tissue.

6. Dosing

35 The phrases "parenteral administration" and "administered parenteral!}" as used herein means modes of administration other than enteral and topical administration, usually by injection and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital,

intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

These compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray, rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

Regardless of the route of administration selected, the compounds of the present disclosure, which may be used in a suitable hydrated form and/or the pharmaceutical compositions of the present disclosure, are formulated into pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this disclosure may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present disclosure employed or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion or metabolism of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated and like factors well known in the medical arts. A daily, weekly or monthly dosage (or other time interval) can be used.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the disclosure employed in the pharmaceutical composition at levels lower than that required to achieve the desired therapeutic effect and then gradually increasing the dosage until the desired effect is achieved.

In general, a suitable daily dose of a compound of the disclosure will be that amount of the compound that is the lowest dose effective to produce a therapeutic effect (e.g., inhibit necrosis). Such an effective dose will generally depend upon the factors described above. Generally doses of the compounds of this disclosure for a patient, when used for the indicated effects, will range from about 0.0001 to about 100 mg per kg of body weight per day. Preferably the daily dosage will range from 0.001 to 50 mg of compound per kg of body weight and even more preferably from 0.01 to 10 mg of compound per kg of body weight.

If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms.

In certain embodiments, the present disclosure relates to compounds for inhibiting cell death, wherein the compounds are represented by structures (I). In certain embodiments, the compounds of the present disclosure are inhibitors of cell death. In any event, the compounds of the present disclosure preferably exert their effect on inhibiting cell death at a concentration less than about 50 micromolar, more preferably at a concentration less than about 10 micromolar and most preferably at a concentration less than 1 micromolar.

The compounds of the disclosure can be tested in standard animal models of stroke and standard protocols such as described by Hara, *et al.* Proc Natl Acad Sci U S A, 1997. 94(5): 2007-12.

When the compounds of the present disclosure are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1% to 99.5% (more preferably, 0.5% to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

The compounds of the present application or the compositions thereof may be administered once, twice, three or four times daily, using any suitable mode described above. Also, administration or treatment with the compounds may be continued for a number of days: for example, commonly treatment would continue for at least 7 days, 14 days or 28 days, for one cycle of treatment. Treatment cycles are well known and are frequently alternated with resting periods of about 1 to 28 days, commonly about 7 days or about 14 days, between cycles. The treatment cycles, in certain embodiments, may also be continuous.

When administered orally, the total daily dosage for a human subject may be between 1 mg and 1,000 mg, between about 1,000-2,000 mg/day, between about 10-500 mg/day, between about 50-300 mg/day, between about 75-200 mg/day or between about 100-150 mg/day.

The daily dosage may also be described as a total amount of a compound described herein administered per dose or per day. Daily dosage of a compound may be between about 1 mg and 4,000 mg, between about 2,000 to 4,000 mg/day, between about 1 to 2,000 mg/day, between about 1 to 1,000 mg/day, between about 10 to 500 mg/day, between about 20 to 500 mg/day, between about 50 to 300 mg/day, between about 75 to 200 mg/day or between about 15 to 150 mg/day.

In certain embodiments, the method comprises administering to the subject an initial daily dose of about 1 to 800 mg of a compound described herein and increasing the dose by increments until clinical efficacy is achieved. Increments of about 5, 10, 25, 50 or 100 mg can be used to increase the dose. The dosage can be increased daily, every other day, twice per week or once per week.

In certain embodiments, a compound or pharmaceutical preparation is administered orally. In certain embodiments, the compound or pharmaceutical preparation is administered intravenously. Alternative routes of administration include sublingual, intramuscular and transdermal administrations.

The preparations of the present disclosure may be given orally, parenteral¹⁾, topically or rectally. They are of course given in forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by
5 suppositories. In certain embodiments, the administration is oral.

7. Combinations

In another aspect of the disclosure the compounds can be administered in combination with other agents, including (but not limited to) compounds that are apoptosis inhibitors; PARP poly(ADP-ribose) polymerase inhibitors; Src inhibitors; agents for the treatment of cardiovascular disorders; anti-
10 inflammatory agents, anti-thrombotic agents; fibrinolytic agents; anti-platelet agents, lipid reducing agents, direct thrombin inhibitors; glycoprotein IIb/IIIa receptor inhibitors; calcium channel blockers; beta-adrenergic receptor blocking agents; cyclooxygenase (*e.g.*, COX-1 and COX-2) inhibitors; angiotensin system inhibitor (*e.g.*, angiotensin-converting enzyme (ACE) inhibitors); renin inhibitors; and/or agents that bind to cellular adhesion molecules and inhibit the ability of white blood cells to attach
15 to such molecules (*e.g.*, polypeptides, polyclonal and monoclonal antibodies).

Embodiments of the disclosure also provide combinations of two or more compounds that inhibit cellular necrosis (*e.g.*, a compound as disclosed herein and an additional agent for inhibiting necrosis). The disclosure also provides combinations of one or more compounds that inhibit cellular necrosis combined with one or more additional agents or compounds (*e.g.*, other therapeutic compounds for
20 treating a disease, condition or infection such as an apoptosis inhibitor).

8. Kits

Provided herein are also kits that include a compound of the disclosure, combinations or a pharmaceutically acceptable salt, tautomer, stereoisomer, mixture of stereoisomers, prodrug or deuterated analog thereof and suitable packaging. In certain embodiments, a kit further includes instructions for use.
25 In one aspect, a kit includes a compound of the disclosure or a pharmaceutically acceptable salt, tautomer, stereoisomer, mixture of stereoisomers, prodrug or deuterated analog thereof and a label and/or instructions for use of the compounds in the treatment of the indications, including the diseases or conditions, described herein.

Provided herein are also articles of manufacture that include a compound described herein or a
30 pharmaceutically acceptable salt, tautomer, stereoisomer, mixture of stereoisomers, prodrug or deuterated analog thereof in a suitable container. The container may be a vial, jar, ampoule, preloaded syringe and intravenous bag.

The kit can also contain instructions for using the compounds according to the disclosure. The kit can be compartmentalized to receive the containers in close confinement. As used herein, a kit such as a
35 compartmentalized kit includes any kit in which compounds or agents are contained in separate containers. Illustrative examples of such containers include, but are not limited to, small glass containers, plastic containers or strips of plastic or paper. Particularly preferred types of containers allow the skilled worker to efficiently transfer reagents from one compartment to another compartment such that the

samples and reagents are not cross-contaminated and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers include, but are not limited to, a container that will accept a compound or combination of compounds and/or other agents of the disclosure. One or more compounds or agents can be provided as a powder (e.g. lyophilized powder) or precipitate. Such compound(s) can be resuspended prior to administration in a solution that may be provided as part of the kit or separately available. A kit can contain compounds or agents in other forms such as liquids, gels, solids, as described herein. Different compounds and/or agents may be provided in different forms in a single kit.

EXAMPLES

The examples and preparations provided below further illustrate and exemplify the compounds of the present disclosure and methods for testing such compounds. It is to be understood that the scope of the present disclosure is not limited in any way by the scope of the following examples. In the following examples and throughout the specification and claims, molecules with a chiral center, unless otherwise noted, exist as a racemic mixture. Single enantiomers may be obtained by methods known to those skilled in the art and described herein. Compounds were named by using either ChemBioDraw Ultra 13.0 or ChemAxon.

General Procedures

All solvents used were commercially available and were used without further purification. Reactions were typically run using anhydrous solvents under an inert atmosphere of nitrogen.

Analytical Methods

¹H Nuclear magnetic resonance (NMR) spectroscopy was carried out using one of the following instruments: a Bruker Avance 400 Ultrashield™ instrument equipped with probe DUAL 400MHz S1, a Bruker Avance 400 Ultrashield™ instrument equipped with probe 6 S1 400 MHz 5mm ¹H-¹³C ID, a Bruker Avance III 400 Ultrashield™ instrument with nanobay equipped with probe Broadband BBFO 5 mm direct, a 400 MHz Agilent Direct Drive instrument with ID AUTO-X PFG probe, a Bruker Mercury Plus 400 instrument equipped with a Bruker 400 BBO probe, a Bruker Avance 400 Ultrashield™ instrument equipped with a BBO probe, a Bruker Avance 400 Ultrashield™ instrument equipped with a BBFO probe, all operating at 400 MHz, or a Bruker Avance III 300 Ultrashield™ instrument equipped with a BBFO probe operating at 300 MHz. All deuterated solvents contained typically 0.03% to 0.05% v/v tetramethylsilane, which was used as the reference signal (set at δ 0.00 for both ¹H and ¹³C). The spectra were acquired in the stated solvent at around room temperature unless otherwise stated. In all cases, NMR data were consistent with the proposed structures. Characteristic chemical shifts (δ) are given in parts-per-million using conventional abbreviations for designation of major peaks: e.g. s, singlet; d, doublet; t, triplet; q, quartet; dd, doublet of doublets; dt, doublet of triplets; br, broad.

Where thin layer chromatography (TLC) has been used it refers to silica gel TLC using silica gel F254 (Merck) plates. Alternatively, thin layer chromatography (TLC) was performed on Alugram* (Silica gel 60 F254) from Manchem-Nagel. R_f is the distance travelled by the compound divided by the

distance travelled by the solvent on a TLC plate. UV was typically used to visualize the spots.

Additional visualization methods were also employed in some cases. In these cases the TLC plate was developed with iodine (generated by adding approximately 1 g of I_2 to 10 g silica gel and thoroughly mixing), ninhydrin (available commercially from Aldrich), or Magic Stain (generated by thoroughly mixing 25 g $(NH_4)_2MoO_4 \cdot 4H_2O$, 5 g $(NH_4)_2Ce(IV)(NO_3)_6$ in 450 mL water and 50 ml, concentrated H_2SO_4) to visualize the compound.

Column chromatography was performed using an automatic flash chromatography (Biotage SP1 or Isolera) system over Biotage silica gel cartridges (KP-Sil or KP-NH) or in the case of reverse phase chromatography over Biotage C18 cartridges (KP-C 18). Alternatively, column chromatography was performed using 40-63 μm (230-400 mesh) silica gel from Silicycle following analogous techniques to those disclosed in Still, W.C.; Kahn, M.; and Mitra, M. Journal of Organic Chemistry, 1978, 43, 2923. Typical solvents used for column chromatography or thin layer chromatography were mixtures of chloroform/methanol, dichloromethane/methanol, ethyl acetate/methanol, hexanes/ethyl acetate, cyclohexane/ethyl acetate.

HPLC analyses were performed on a SHIMADZU UFLC with two LC20 AD pump and a SPD-M20A Photodiode Array Detector. The column used was an XBridge C18, 3.5 μm , 4.60 $\times$ 100 mm. A linear gradient was applied, starting at 90 % A (A: 0.05% TFA in water) and ending at 95% B (B: 0.05% TFA in MeCN) over 10 min with a total run time of 15 min. The column temperature was at 40 °C with the flow rate of 1.5 mL/min. The Diode Array Detector was scanned from 200-400 nm.

Unless otherwise stated, LCMS analyses were performed on LCMS analyses were performed on a SHIMADZU LCMS consisting of an UFLC 20-AD and LCMS 2020 MS detector. The Diode Array Detector was scanned from 190-400 nm. The mass spectrometer was equipped with an electrospray ion source (ESI) operated in a positive or negative mode. The mass spectrometer was scanned between m/z 90-900 with a scan time from 0.5 to 1.0 s.

Liquid Chromatography-Mass Spectrometry Method A: Total ion current (TIC) and DAD UV chromatographic traces together with MS and UV spectra associated with the peaks were taken on a UPLC/MS Acquity™ system equipped with PDA detector and coupled to a Waters single quadrupole mass spectrometer operating in alternated positive and negative electrospray ionization mode. [LC/MS-ES (+/-): analyses performed using an Acquity UPLC™ CSH, C18 column (50 $\times$ 2.1mm, 1.7 μm particle size), column temperature 40 °C, mobile phase: A-water + 0.1% HCOOH/ B- CH_3CN + 0.1% FICOOH, flow rate: 1.0 mL/min, runtime = 2.0 min, gradient: t=0 min 3%B, t= 1.5 min 99.9% B, t = 1.9 min 99.9% B, t= 2.0 min 3% B, stop time 2.0 min. Positive ES 100-1000, Negative ES 100-1000, UV detection DAD 210-350 nm.

Liquid Chromatography-Mass Spectrometry Method B: Total ion current (TIC) and DAD UV chromatographic traces together with MS and UV spectra associated with the peaks were taken on a UPLC/MS Acquity™ system equipped with PDA detector and coupled to a Waters single quadrupole mass spectrometer operating in alternated positive and negative electrospray ionization mode. [LC/MS-ES (+/-): analyses performed using an Acquity UPLC™ BEH, C18 column (50 $\times$ 2.1mm, 1.7 μm particle

size), column temperature 40 °C, mobile phase: A - 0.1% v/v aqueous ammonia solution pH 10/ B- CH₃CN, flow rate: 1.0 mL/min, runtime = 2.0 min, gradient: t=0 min 3%B, t= 1.5 min 99.9% B, t = 1.9 min 99.9% B, t= 2.0 min 3% B, stop time 2.0 min. Positive ES 100-1000, Negative ES 100-1000, UV detection DAD 210-350 nm.

Liquid Chromatography-Mass Spectrometry Method C: The column used was an Shim-pack XR-ODS, 2.2 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 1.8 min with a total run time of 2.1 min. The column temperature was at 45 °C with the flow rate of 1.2 mL/min.

Liquid Chromatography-Mass Spectrometry Method D: The column used was an Shim-pack XR-ODS, 2.2 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 3.2 min with a total run time of 3.6 min. The column temperature was at 40 °C with the flow rate of 1.0 mL/min.

Liquid Chromatography-Mass Spectrometry Method E: The column used was an Shim-pack XR-ODS, 2.2 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 4.7 min with a total run time of 5.2 min. The column temperature was at 40 °C with the flow rate of 1.0 mL/min.

Liquid Chromatography-Mass Spectrometry Method F: The column used was an Ascentis Express C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 2.8 min with a total run time of 3.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method G: The column used was an Poroshell HPH-C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 90% A (A: 0.05%, NH₄HCO₃ in water) and ending at 95% B (B: 0.05% NH₄HCO₃ in MeCN) over 1.8 min with a total run time of 2 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method H: The column used was an Poroshell HPH-C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 90% A (A: 0.05% NH₄HCO₃ in water) and ending at 95% B (B: 0.05% NH₄HCO₃ in MeCN) over 2.7 min with a total run time of 3.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method I: The column used was an Poroshell HPH-C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 90% A (A: 0.05% NH₄HCO₃ in water) and ending at 95% B (B: 0.05% NH₄HCO₃ in MeCN) over 4.5 min with a total run time of 5.2 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method J: The column used was an Agilent Poroshell HPH-C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 90% A (A: 0.05% NH₄HCO₃ in water) and ending at 95% B (B: 0.05% NH₄HCO₃ in MeCN) over 2.7 min with a total run time of 3.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method K: The column used was an Ascentis Express C18, 2.7 µm, 3.0 x 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in

water) and ending at 100% B (B: 0.05% TFA in MeCN) over 1.8 min with a total run time of 2.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method L: The column used was an Ascentis Express C18, 2.7 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 95% B (B: 0.05% TFA in MeCN) over 2.7 min with a total run time of 3.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method M: The column used was an Ascentis Express C18, 2.7 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 1.8 min with a total run time of 2.0 min.

The column temperature was at 40 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method N: The column used was an Ascentis Express C18, 2.7 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 2.8 min with a total run time of 3.0 min.

The column temperature was at 40 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method O: The column used was an Shim-pack XR-ODS, 2.2 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 3.2 min with a total run time of 3.6 min.

The column temperature was at 40 °C with the flow rate of 1.0 mL/min.

Liquid Chromatography-Mass Spectrometry Method P: The column used was an Kinetex 2.6 μm XB-C18, 2.6 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 2.7 min with a total run time of 3.0 min.

The column temperature was at 40 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method Q: The column used was a Kinetex XB-C18100A, 2.6 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 90% A (A: 0.1% FA in water)

and ending at 95% B (B: 0.1% formic acid in MeCN) over 2.7 min with a total run time of 3.0 min. The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method Li: The column used was an Ascentis Express C18, 2.7 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 95% A (A: 0.05% TFA in water) and ending at 100% B (B: 0.05% TFA in MeCN) over 4.6 min with a total run time of 5.2 min.

The column temperature was at 45 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method S: The column used was an Kinetex EVO C18, 2.6 μm , 3.0 $\times$ 50 mm. A linear gradient was applied, starting at 90% A (A: 5 mM NH_4HCO_3 in water) and ending at 95% B (MeCN) over 4.7 min with a total run time of 5.0 min. The column temperature was at 40 °C with the flow rate of 1.5 mL/min.

Liquid Chromatography-Mass Spectrometry Method T: Total ion current (TIC) and DAD UV chromatographic traces together with MS and UV spectra associated with the peaks were taken on LCMS Agilent Technologies system equipped with DAD detector and coupled to an Agilent single quadrupole mass spectrometer operating in alternated positive and negative electrospray ionization

mode. [LC/MS-ES (+): analyses performed using an Agilent Technologies 1200 Series, Atlantis dC18 column (50 x 4.6 mm, 5µm particle size), column temperature 25 °C, mobile phase: A-water + 0.1% TFA/ B-CH₃CN + 0.1% TFA, flow rate 1.5 mL/min, total runtime = 6.0 min, gradient: t = 0 min, 10% B, t = 2.5 min 95% B, t = 4.5 min 95% B, t = 4.6 min 10% B, t = 6.0 min 10% B, stop time 6.0 min. Positive ES 100-900, UV detection DAD 220-400 nm].

Compound Preparation

Where the preparation of starting materials is not described, these are commercially available, known in the literature or readily obtainable by those skilled in the art using standard procedures. Where it is stated that compounds were prepared analogously to earlier examples or intermediates, it will be appreciated by the skilled person that the reaction time, number of equivalents of reagents and temperature can be modified for each specific reaction and that it may be necessary or desirable to employ different work-up or purification techniques. Where reactions are carried out using microwave irradiation, the microwave used is a Biotage Initiator. The actual power supplied varies during the course of the reaction in order to maintain a constant temperature.

General Procedure A: Amide coupling conditions

N,N-Diisopropylethylamine (158 µL, 1.34 mmol), 1-hydroxybenzotriazole (70 mg, 0.52 mmol) and EDC-HCl (99 mg, 0.52 mmol) were added to a stirred solution of carboxylic acid (0.26 mmol) and (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) in CH₂Cl₂ (1.5 mL) and the resulting mixture was stirred at room temperature for 18 h. The reaction mixture was diluted with CH₂Cl₂ and washed twice with sat. NaHCO₃ solution. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography to afford the title compound.

Exemplary compounds of the disclosure and intermediates made in the following examples are summarized herein, including inhibitor activity for receptor interacting protein kinase 1 (IC₅₀, micromolar) for representative compounds together with LCMS method (M), LC retention time (RT) in minutes and Mass Spec m/z values (molecular weight).

INTERMEDIATE 1

PREPARATION OF TERT-BUTYL N-[(3R)-3-HYDROXY-3-PHENYLPROPOXY]CARBAMATE

NaH (180 mg, 4.51 mmol) was added to a solution of tert-butyl N-hydroxy carbamate (542 mg, 4.10 mmol) in DMF (10 mL) which had been pre-cooled to 0 °C. After 20 min, a solution of (1R)-3-chloro-1-phenylpropan-1-ol (350 mg, 2.05 mmol) in DMF (3 mL) was added dropwise. The reaction mixture was stirred at 0 °C for 10 min, warmed up to room temperature, stirred for 72 h and quenched with sat. NH₄Cl solution. The aqueous portion was extracted with EtOAc. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound (450 mg, 82%) as a colorless wax. ¹H NMR (400 MHz, DMSO-d₆) δ 10.02–9.90 (br s, 1H), 7.37–7.28 (m, 4H), 7.26–7.19 (m, 1H), 5.22 (d, J=4.5 Hz, 1H), 4.72–4.64 (m, 1H), 3.82 (dt, J=9.8, 7.0 Hz, 1H), 3.69 (dt,

, 1=91, 6.3 Hz, 1H), 1.87-1.74 (m, 2H), 1.40 (s, 9H). LC-MS (Method A): m/z = 268.3 [M+H]⁺, 0.91 min.

INTERMEDIATE 2

PREPARATION OF 3-PHENYL-1,2-OXAZOLIDINE

5 Methane sulfonyl chloride (153 μ L, 1.98 mmol) was added to a solution of tert-butyl N-[(3R)-3-hydroxy-3-phenyl]propoxy]carbamate (459 mg, 1.72 mmol) and Et₃N (478 μ L, 3.44 mmol) in CH₂Cl₂ (40 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 1.5 h. The reaction mixture was diluted with CH₂Cl₂ and washed with water. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford crude tert-butyl 3-phenyl-1,2-oxazolidine-2-carboxylate, which was directly progressed to the next step. LC-MS (Method A): m/z = 250.3 [M+Hf, 194.2 [M+H-tBu]⁺, 1.03 min.

TFA (5 mL) was added to a solution of the crude tert-butyl 3-phenyl-1,2-oxazolidine-2-carboxylate in CH₂Cl₂ (50 mL). The reaction mixture was stirred at room temperature for 30 min and concentrated under reduced pressure. The residue was dissolved in CH₂Cl₂ and the organic portion was washed with sat. NaHCO₃, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water-CH₃CN, 100:0 to 50:50) to afford the title compound (150 mg, 58%) as a colorless oil. ¹H NMR (400 MHz, DMSO-d₆) δ 7.42-7.16 (m, 5H), 6.52-6.12 (br s, 1H), 4.47-4.23 (br s, 1H), 3.91 (td, J =8.0, 5.3 Hz, 1H), 3.85-3.66 (br s, 1H), 2.64-2.53 (m, 1H), 2.09 (dddd, J =12.1, 8.7, 6.7, 5.5 Hz, 1H). LC-MS (Method A): m/z = 150.1 [M+Hf, 0.40 min.

Examples 1 and 2, which were prepared from this intermediate, were formed as -85:15 mixtures of enantiomers. It is therefore assumed that this batch of 3-phenyl-1,2-oxazolidine has -70% enantiomeric excess (e.e.).

INTERMEDIATE 3

PREPARATION OF (3S)-3-PHENYL-1,2-OXAZOLIDINE

25 TFA (5 mL) was added to a solution of tert-butyl (3S)-3-phenyl-1,2-oxazolidine-2-carboxylate in CH₂Cl₂ (50 mL). The reaction mixture was stirred at room temperature for 30 min and concentrated under reduced pressure. The residue was dissolved in CH₂Cl₂ and the organic portion was washed with sat. NaHCO₃, dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (1.76 g, 99%). ¹H NMR (400MHz, DMSO-d₆) δ 7.41-7.35 (m, 2H), 7.35-7.28 (m, 2H), 7.27-7.21 (m, 1H), 6.67-6.33 (br. s, 1H), 4.40-4.29 (m, 1H), 3.92 (dt, J =5.7, 8.0 Hz, 1H), 3.84-3.70 (m, 1H), 2.59 (did. J =5.4, 8.5, 11.8 Hz, 1H), 2.54-2.54 (m, 1H), 2.10 (dddd, J =5.8, 6.6, 8.8, 12.2 Hz, 1H). LC-MS (Method A): m/z = 150.1 [M+H]⁺, 0.41 min.

35 A batch of Example 1 prepared from this intermediate was formed as a -98:2 mixture of enantiomers. It is therefore assumed that this batch of (3S)-3-phenyl-1,2-oxazolidine has 96% enantiomeric excess (e.e.).

INTERMEDIATE 4

PREPARATION OF (E)-N-[(2-CHLORO-6-FLUOROPHENYL)METHYLIDENE] HYDROXYLAMINE

An aqueous solution of NaOH (1.13 g, 28.5 mmol) in water (1 mL) was added to a stirred mixture of 2-chloro-6-fluorobenzaldehyde (1.5 g, 9.5 mmol), EtOH (3 mL), ice and water (9 mL) and hydroxylamine hydrochloride (849 mg, 12.3 mmol). The mixture was stirred at room temperature for 1.5 h and extracted with Et₂O to remove impurities. The aqueous layer was neutralized with 2M aq HCl solution and extracted three times with Et₂O. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude title compound (1.3 g), which was directly progressed to the next step. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.85 (s, 1H), 8.23 (s, 1H), 7.50-7.37 (m, 2H), 7.35-7.27 (m, 1H). LC-MS (Method A): m/z = 174.1 [M+H]⁺, 0.83 min.

INTERMEDIATE 5

PREPARATION OF (E)-[(TERT-BUTYLDIMETHYLSILYL)OXY] [(2-CHLORO-6-FLUOROPHENYL)METHYLIDENE]AMINE

TBSCl (520 mg, 3.5 mmol) and imidazole (472 mg, 6.9 mmol) were added to a solution of (E)-N-[(2-chloro-6-fluorophenyl)methylidene]hydroxy]amine (400 mg, 2.3 mmol) in DMF (20 mL). The reaction was stirred at room temperature for 50 h, diluted with EtOAc and washed twice with water. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 95:5) to afford the title compound (400 mg, 60%) as a colorless oil. LC-MS (Method A): m/z = 288.2 [M+H]⁺, 1.61 min.

INTERMEDIATE 6

PREPARATION OF 3-(2-CHLORO-6-FLUOROPHENYL)-1,2-OXAZOLIDINE

Ethylene (purity >99.95% GC, 00489 Sigma-Aldrich) was bubbled for 30 min through a solution of (E)-[(tert-butyldimethylsilyl)oxy][(2-chloro-6-fluorophenyl)methylidene]amine (100 mg, 0.35 mmol) in CH₂Cl₂ (7 mL) which had been pre-cooled to -78 °C. The reaction was warmed to room temperature and BF₃·Et₂O (98 μL, 0.77 mmol) was added. The solution was stirred at 60 °C for 16 h under an ethylene atmosphere, cooled to room temperature, diluted with CH₂Cl₂ and washed with sat. NaHCO₃ solution. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude reaction mixture (147 mg) was joined together with the crude reaction mixture (132 mg) from a second reaction identical to that described above in all premises but with a longer heating time of 48h. The resulting mixture was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 70:30) to afford the title compound (7 mg). ¹H NMR (400 MHz, CDCl₃) δ 7.26-7.21 (m, 2H), 7.06-6.99 (m, 1H), 4.97 (t, J=8.5 Hz, 1H), 4.25 (dt, J=5.4, 8.1 Hz, 1H), 4.08 (q, J=8.0 Hz, 1H), 2.67-2.55 (m, 1H), 2.50-2.38 (m, 1H). LC-MS (Method A): m/z = 202.1 [M+H]⁺, 0.69 min.

INTERMEDIATE 7

PREPARATION OF 1-(3-FLUOROPHENYL)PROP-2-EN-1-OL

Vinylmagnesium bromide (1M solution in THF, 12.0 mL, 12.0 mmol) was added dropwise to a solution of 3-fluorobenzaldehyde (1.0 g, 8.0 mmol) in THF (10 mL), which had been pre-cooled to -78 °C. The reaction was stirred at -78 °C for 2 h and quenched with sat. NH₄Cl solution. The aqueous

portion was extracted twice with EtOAc. The combined organic portions were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 80:20) to afford the title compound (800 mg, 66%) as a yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 7.34 (td, $J=7.9, 5.8$ Hz, 1H), 7.18-7.08 (m, 2H), 6.98 (tdd, $J=8.4, 2.5, 1.0$ Hz, 1H), 6.11-5.94 (m, 1H), 5.37 (dt, $J=17.3, 1.3$ Hz, 1H), 5.27-5.18 (m, 2H), 1.96 (d, $J=3.8$ Hz, 1H). LC-MS (Method A): m/z = 135.0 $[\text{M}-\text{H}_2\text{O}+\text{H}]^+$, 0.81 min.

INTERMEDIATE 8

PREPARATION OF 1-(3-FLUOROPHENYL)PROP-2-EN-1-ONE

Dess-Martin periodinane (3.35 g, 7.9 mmol) was added to a solution of 1-(3-fluorophenyl)prop-2-en-1-ol (800 mg, 5.3 mmol) in CH_2Cl_2 (20 mL). The reaction mixture was stirred at room temperature for 1.5 h, diluted with CH_2Cl_2 and quenched with a 1:1 sat. NaHCO_3 solution / 2M $\text{Na}_2\text{S}_2\text{O}_3$ solution (30 mL). The mixture was vigorously stirred at room temperature for 30 min. The layers were separated and the organic portion was washed with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford crude title compound (779 mg), which was directly progressed to the next step. ^1H NMR (400 MHz, CDCl_3) δ 7.75 (dq, $J=7.8, 0.8$ Hz, 1H), 7.66 (ddd, $J=9.3, 2.4, 1.5$ Hz, 1H), 7.49 (td, $J=8.0, 5.5$ Hz, 1H), 7.35-7.28 (m, 1H), 7.14 (dd, $J=7.2, 10.7$ Hz, 1H), 6.52-6.45 (m, 1H), 6.00 (dd, $J=10.5, 1.5$ Hz, 1H). LC-MS (Method A): m/z = 151.0 $[\text{M}+\text{H}]^+$, 0.92 min.

INTERMEDIATE 9

PREPARATION OF 3-CHLORO-1-(3-FLUOROPHENYL)PROPAN-1-ONE

Hydrochloric acid (4M solution in dioxane, 30 mL) was added to a solution of 1-(3-fluorophenyl)prop-2-en-1-one (779 mg) in CH_2Cl_2 (60 mL), which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 1 h and concentrated under reduced pressure to afford the title compound as a brown oil, which was directly progressed to the next step. ^1H NMR (400 MHz, CDCl_3) δ 7.80-7.74 (m, 1H), 7.67 (dt, $J=9.3, 2.0$ Hz, 1H), 7.50 (td, $J=8.0, 5.5$ Hz, 1H), 7.32 (td, $J=8.2, 2.5$ Hz, 1H), 3.95 (t, $J=6.7$ Hz, 2H), 3.47 (t, $J=6.8$ Hz, 2H).

INTERMEDIATE 10

PREPARATION OF 3-CHLORO-1-(3-FLUOROPHENYL)PROPAN-1-OL

Sodium borohydride (201 mg, 5.3 mmol) was added to a solution of 3-chloro-1-(3-fluorophenyl)propan-1-one in MeOH (40 mL). The reaction was stirred at room temperature for 30 min, concentrated to half of its initial volume under reduced pressure and diluted with EtOAc. The organic portion was washed with sat. NH_4Cl solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (778 mg, 78% over three steps) as an orange oil. ^1H NMR (400 MHz, CDCl_3) δ 7.35 (td, $J=7.9, 5.8$ Hz, 1H), 7.20-7.08 (m, 2H), 7.00 (tdd, $J=8.4, 2.6, 0.9$ Hz, 1H), 4.99 (dt, $J=8.4, 4.1$ Hz, 1H), 3.77 (ddd, $J=11.0, 8.3, 5.3$ Hz, 1H), 3.63-3.54 (m, 1H), 2.28-2.17 (m, 1H), 2.02-2.16 (m, 2H).

INTERMEDIATE 11

PREPARATION OF TERT-BUTYL N-[3-(3-FLUOROPHENYL)-3-HYDROXYPROPOXY] CARBAMATE

Sodium hydride (60% dispersion in mineral oil, 413 mg, 10.3 mmol) was added in two portions to a stirred solution of tert-butyl N-hydroxy carbamate (1.10 g, 8.27 mmol) in DMF (18 mL), which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 30 min. A solution of 3-chloro-1-(3-fluorophenyl)propan-1-ol (778 mg, 4.14 mmol) in DMF (5 mL) was added dropwise. The reaction was gradually allowed to reach room temperature, stirred for 16 h and quenched with sat. NH_4Cl solution. The aqueous portion was extracted twice with EtOAc. The combined organic portions were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound (1.10 g, 93%) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.98 (br. s, 1H), 7.36 (td, $J=7.9, 6.0$ Hz, 1H), 7.22–7.12 (m, 2H), 7.09–6.98 (m, 1H), 5.37 (d, $J=4.8$ Hz, 1H), 4.73 (dt, $J=8.7, 4.5$ Hz, 1H), 3.84 (dt, $J=8.8, 6.9$ Hz, 1H), 3.68 (ddd, $J=9.9, 6.7, 5.3$ Hz, 1H), 1.91–1.68 (m, 1H), 1.41 (s, 9H). LC-MS (Method A): m/z = 286.1 $[\text{M}+\text{H}]^+$, 0.97 min.

INTERMEDIATE 12

PREPARATION OF 3-(3-FLUOROPHENYL)-1,2-OXAZOLIDINE

Methane sulfonyl chloride (358 μL , 4.63 mmol) was added to a solution of tert-butyl N-[3-(3-fluorophenyl)-3-hydroxypropoxy]carbamate (1.10 g, 3.86 mmol) and Et_3N (1.1 mL, 7.71 mmol) in CH_2Cl_2 (40 mL), which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 1 h, diluted with CH_2Cl_2 and washed with sat. NaHCO_3 solution and sat. NH_4Cl solution. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the title compound in a mixture with tert-butyl 3-(3-fluorophenyl)-1,2-oxazolidine-2-carboxylate (1.22 g). This mixture was dissolved in CH_2Cl_2 (40 mL) and treated with trifluoroacetic acid (5 mL). The reaction mixture was stirred at room temperature for 1 h and concentrated under reduced pressure. The crude product was dissolved in CH_2Cl_2 and the organic portion was washed twice with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3CN , 100:0 to 20:80) to afford the title compound (253 mg, 39% over two steps) as a pale yellow wax. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.45–7.29 (m, 1H), 7.27–7.15 (m, 2H), 7.12–6.97 (m, 1H), 6.59–6.29 (m, 1H), 4.54–4.30 (m, 1H), 3.91 (td, $J=8.0, 5.1$ Hz, 1H), 3.82–3.57 (m, 1H), 2.68–2.55 (m, 1H), 2.08 (dddd, $J=12.0, 8.5, 7.0, 5.1$ Hz, 1H). LC-MS (Method A): m/z = 168.2 $[\text{M}+\text{H}]^+$, 0.57 min.

INTERMEDIATE 13

PREPARATION OF 3-CHLORO-1-(2-FLUOROPHENYL)PROPAN-1-ONE

To a solution of 1-bromo-2-fluorobenzene (5.0 g, 28.5 mmol) in dry THF (7.0 mL) was added dropwise at 0 °C isopropylmagnesium chloride lithium chloride complex (1.3M solution in THF, 24.1 mL, 31.3 mmol). The mixture was stirred at this temperature for 4 h, then diluted with THF (50 mL) and the resulting solution was added dropwise over 20 min to a solution of 3-chloropropionyl chloride (3.8 g, 30 mmol) in dry THF (25 mL) at 0 °C. The mixture was then stirred at 0 °C for 1 h 40 min then a sat.

aqueous NH_4Cl solution was added and the mixture was extracted twice with EtOAc. The combined organic phases were washed with sat. aqueous NaHCO_3 solution-water (1:1) and then with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 80:20) to afford the title compound (313 mg) contaminated with unknown impurities (estimated purity ~70% by ^1H NMR) as a pale yellow oil. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.93 (dt, $J=8.0, 2.0$ Hz, 1H), 7.61-7.54 (m, 1H), 7.31-7.25 (m, 1H), 7.21-7.15 (m, 1H), 3.93 (t, $J=8.0$ Hz, 2H), 3.50 (dt, $J=8.0, 4.0$ Hz, 2H). LC-MS (Method A): $m/z = 187$ $[\text{M}+\text{H}]^+$, 1.00 min.

INTERMEDIATE 14

PREPARATION OF 3-CHLORO-1-(2-FLUOROPHENYL)PROPAN-1-OL

Sodium borohydride (20 mg, 0.53 mmol) was added at 0 °C to a solution of 3-chloro-1-(2-fluorophenyl)propan-1-one (100 mg, 0.53 mmol) in dry THF (1.5 mL). The mixture was then stirred at this temperature for 1 h 30 min. Sat. aqueous NH_4Cl solution was added and the mixture was stirred at room temperature for 20 min, then extracted twice with EtOAc. The combined organic phases were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give crude title compound as a yellow oil (72 mg).

In a second experiment, sodium borohydride (39 mg, 1.03 mmol) was added at 0 °C to a solution of 3-chloro-1-(2-fluorophenyl)propan-1-one (193 mg, 1.03 mmol) in dry MeOH (3.5 mL). The mixture was then stirred at this temperature for 30 min. Sat. aqueous NH_4Cl solution was added and the mixture was stirred at 0 °C for 20 min, then extracted three times with ethyl acetate. The combined organic phases were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give crude title compound as a yellow oil (188 mg).

The two erodes obtained above were combined and purified by column chromatography (cyclohexane-EtOAc, 95:5 to 60:40) to afford the title compound (154 mg) contaminated with unknown impurities (estimated purity ~70% by ^1H NMR) as a colorless oil. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.49 (dt, $J=8.0, 2.0$ Hz, 1H), 7.34-7.27 (m, 1H), 7.22-7.16 (m, 1H), 7.10-7.02 (m, 1H), 5.30-5.23 (m, 1H), 3.83-3.75 (m, 1H), 3.70-3.62 (m, 1H), 2.33-2.18 (m, 2H), 2.12 (d, $J=4.7$ Hz, 1H).

INTERMEDIATE 15

PREPARATION OF TERT-BUTYL N-[3-(2-FLUOROPHENYL)-3-HYDROXYPROPOXY]CARBAMATE

NaH (60% dispersion in mineral oil, 72 mg, 1.80 mmol) was added to a solution of tert-butyl N-hydroxycarbamate (218 mg, 1.63 mmol) in dry DMF (2.6 mL) which had been pre-cooled to 0 °C. After 20 min, a solution of 3-chloro-1-(2-fluorophenyl)propan-1-ol (154 mg, 0.82 mmol) in dry DMF (1.5 mL) was added dropwise. The reaction was stirred at 0 °C for 45 min, then warmed to room temperature. After 4 h 30 min the mixture was quenched with sat. aqueous NH_4Cl solution and extracted three times with EtOAc. The combined organic phases were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product (340 mg) was purified by column chromatography (cyclohexane-EtOAc, 93:7 to 40:60) to afford the title compound (185 mg, 2% over 3 steps) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.62 (dt, $J=8.0, 1.7$ Hz, 1H), 7.32 (br. s, 1H), 7.28-7.21 (m, 1H), 7.20-7.15 (m,

1H), 7.05- 6.99 (m, 1H), 5.35-5.29 (m, 1H), 4.16-4.05 (m, 2H), 2.17-2.09 (m, 1H), 2.01-1.91 (m, 1H), 1.52 (s, 9H). LC-MS (Method A): m/z = 286.1 [M+H]⁺, 0.94 min.

INTERMEDIATE 16

PREPARATION OF 3-(2-FLUOROPHENYL)-1,2-OXAZOLIDINE

5 Trifluoroacetic acid (0.166 mL, 2.17 mmol) was added at 0 °C to a solution of tert-butyl 3-(2-fluorophenyl)-1,2-oxazolidine-2-carboxylate (58 mg, 0.217 mmol) in dry CH₂Cl₂ (2 mL). The mixture was then stirred at room temperature for 3 h, then concentrated under reduced pressure. The residue was dissolved in CH₂Cl₂ and the organic portion was washed twice with sat. aqueous NaHCO₃ solution, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give the title compound as a colorless
10 oil (31 mg, 85%). This was used in the following experiment without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.53 (dt, J=8.0, 1.0 Hz, 1H), 7.31-7.23 (m, 1H), 7.15 (dt, J=8.0, 1.0 Hz, 1H), 7.06 (ddd, J=10.6, 8.2, 1.0 Hz, 1H), 4.80-4.71 (m, 1H), 4.15-4.06 (m, 1H), 4.02-3.92 (m, 1H), 2.78-2.67 (m, 1H), 2.35-2.24 (m, 1H). LC-MS (Method A): m/z = 168.2 [M+H]⁺, 0.57 min.

INTERMEDIATE 17

PREPARATION OF 3-CHLORO-1-(4-FLUOROPHENYL)PROPAN-1-OL

15 NaBH₄ (200 mg, 5.37 mmol) was added portionwise to a solution of 3-chloro-1-(4-fluorophenyl)propan-1-one (1g, 5.37 mmol) in MeOH (10 mL) at 0 °C. The reaction mixture was allowed to reach room temperature and stirred at this temperature for 1 h. A sat. NH₄Cl solution was added and the mixture was stirred at 0 °C for 20 min, then extracted twice with EtOAc. The combined
20 organic phases were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford title compound (830 mg, 82% yield) as colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.40-7.31 (m, 2H), 7.13-7.01 (m, 2H), 4.97 (dt, J=8.4, 4.1 Hz, 1H), 3.76 (ddd, J=10.9, 8.3, 5.5 Hz, 1H), 3.57 (dt, J=11.0, 5.9 Hz, 1H), 2.24 (ddt, J=14.4, 8.5, 5.7 Hz, 1H), 2.14-2.02 (m, 2H). LC-MS (Method A): m/z = 171.0 [M+H]⁺, 0.97 min.

INTERMEDIATE 18

PREPARATION OF TERT-BUTYL N-[3-(4-FLUOROPHENYL)-3-HYDROXYPROPOXY]CARBAMATE

30 NaH (60% dispersion in mineral oil, 186 mg, 4.66 mmol) was added to a solution of tert-butyl N-hydroxycarbamate (566 mg, 4.25 mmol) in DMF (5 mL) which had been pre-cooled to 0 °C. After 20 min, a solution of 3-chloro-1-(4-fluorophenyl)propan-1-ol (400 mg, 2.12 mmol) in DMF (2 mL) was added dropwise. The reaction mixture was stirred at 0 °C for 10 min, warmed to room temperature, stirred for 5 h and quenched with sat. NH₄Cl solution. The aqueous portion was extracted with EtOAc. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude
35 product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound (430 mg, 71% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.44-7.34 (m, 2H), 7.29 (s, 1H), 7.10-6.99 (m, 2H), 5.02 (dt, J=8.4, 3.9 Hz, 1H), 4.19-3.99 (m, 2H), 2.10-1.88 (m, 2H), 1.52 (s, 9H). LC-MS (Method A): m/z = 308.1 [M+Na]⁺, 1.03 min.

INTERMEDIATE 19

PREPARATION OF 3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINE

TFA (0.66 mL) was added to a solution of tert-butyl 3-(4-fluorophenyl)-1,2-oxazolidine-2-carboxylate (232 mg, 0.87 mmol) in CH_2Cl_2 (7 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at room temperature for 1.5 h and concentrated under reduced pressure. The residue was dissolved in CH_2Cl_2 and the organic portion was washed with sat. NaHCO_3 , dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3CN , 100:0 to 50:50) to afford the title compound (90 mg, 62% yield) as a colorless oil. ^1H NMR (400 MHz, $\text{O}(\text{X})$) δ 7.45-7.33 (m, 1H), 7.15-6.96 (m, 2H), 4.58-4.42 (m, 1H), 4.18-3.94 (m, 2H), 2.80-2.56 (m, 1H), 2.40-2.16 (m, 1H). LC-MS (Method A): m/z = 168.2 $[\text{M} + \text{H}]^+$, 0.57 min.

INTERMEDIATE 20

PREPARATION OF 1-(6-FLUOROPYRIMIDIN-3-YL)PROP-2-EN-1-OL

Vinylmagnesium bromide (1M solution in THF, 24.8 mL, 24.8 mmol) was added dropwise to a solution of 6-fluoropyridine-3-carbaldehyde (1.00 g, 8.00 mmol) in THF (10 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 2 h and quenched with sat. NH_4Cl solution. The aqueous portion was extracted twice with EtOAc. The combined organic portions were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude title compound (1.09 g), which was directly progressed to the next step. ^1H NMR (400 MHz, CDCl_3) δ 8.23 (d, $J=2.3$ Hz, 1H), 7.84 (td, $J=8.2, 2.5$ Hz, 1H), 6.95 (dd, $J=8.4, 2.9$ Hz, 1H), 6.11-5.97 (m, 1H), 5.47-5.37 (m, 1H), 5.34-5.26 (m, 2H), 2.08 (br s, 1H). LC-MS (Method A): m/z = 154.0 $[\text{M} + \text{H}]^+$, 0.57 min.

INTERMEDIATE 21

PREPARATION OF 1-(6-FLUOROPYRIDIN-3-YL)PROP-2-EN-1-ONE

Dess-Martin periodinane (3.35 g, 7.90 mmol) was added to a solution of 1-(6-fluoropyridin-3-yl)prop-2-en-1-ol (1.09 g) in CH_2Cl_2 (70 mL). The reaction was stirred at room temperature for 16 h, and quenched with a 1:1 mixture of sat. NaHCO_3 solution/2M $\text{Na}_2\text{S}_2\text{O}_3$ solution. After being stirred vigorously for 30 min, the layers were separated. The organic portion was washed with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude title compound, which was directly progressed to the next step. LC-MS (Method A): m/z = 152.0 $[\text{M} + \text{H}]^+$, 0.67 min.

INTERMEDIATE 22

PREPARATION OF 3-CHLORO-1-(6-FLUOROPYRIDIN-3-YL)PROPAN-1-ONE

A solution of hydrochloric acid (4M in dioxane, 30 mL) was added to a solution of 1-(6-fluoropyridin-3-yl)prop-2-en-1-one in CH_2Cl_2 (60 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 40 min and concentrated under reduced pressure to give the crude title compound, which was directly progressed to the next step. LC-MS (Method A): m/z = 188.0 $[\text{M} + \text{H}]^+$, 0.80 min.

INTERMEDIATE 23

PREPARATION OF 3-CHLORO-1-(6-FLUOROPYRIDIN-3-YL)PROPAN-1-OL

Sodium borohydride (273 mg, 7.18 mmol) was added to a solution of 3-chloro-1-(6-fluoropyridin-3-yl)propan-1-one in MeOH (50 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 20 min. Volatiles were removed under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 20:80) to give the title compound (590 mg, 26% over four steps) as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 8.24 (d, *J*=2.5 Hz, 1H), 7.85 (td, *J*=8.0, 2.4 Hz, 1H), 6.97 (dd, *J*=8.4, 2.9 Hz, 1H), 5.08 (dt, *J*=8.8, 4.1 Hz, 1H), 3.84-3.77 (m, 1H), 3.61 (dt, *J*=11.1, 5.6 Hz, 1H), 2.31-2.21 (m, 1H), 2.16 (d, *J*=3.8 Hz, 1H), 2.14-2.03 (m, 1H). LC-MS (Method A): *m/z* = 190.0 [M+H]⁺, 0.74 min.

INTERMEDIATE 24

PREPARATION OF *tert*-BUTYL N-[3-(6-FLUOROPYRIDIN-3-YL)-3-HYDROXYPROPOXY]CARBAMATE

Sodium hydride (60% dispersion in mineral oil, 237 mg, 5.93 mmol) was added to a stirred solution of *tert*-butyl N-hydroxycarbamate (391 mg, 2.96 mmol) in DMF (8 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 10 min. A solution of 3-chloro-1-(6-fluoropyridin-3-yl)propan-1-ol (590 mg, 3.12 mmol) in DMF (5 mL) was then added dropwise. The reaction mixture was gradually allowed to reach room temperature, stirred for 5 h and quenched with sat. NH₄Cl solution. The aqueous portion was extracted twice with EtOAc. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 20:80) to afford the title compound (366 mg, 41%) as a yellow oil. LC-MS (Method A): *m/z* = 287.2 [M+H]⁺, 0.78 min.

INTERMEDIATE 25

PREPARATION OF *tert*-BUTYL 3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

Methanesulfonyl chloride (148 μL, 1.90 mmol) was added to a solution of *tert*-butyl N-[3-(6-fluoropyridin-3-yl)-3-hydroxypropoxy]carbamate (366 mg, 1.28 mmol) and Et₃N (355 μL, 2.55 mmol) in CH₂Cl₂ (7 mL) which had been pre-cooled to 0 °C. The reaction mixture was gradually allowed to reach room temperature and stirred for 1 h. The reaction mixture was diluted with CH₂Cl₂ and washed with sat. NH₄Cl solution. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 20:80) to afford the title compound (161 mg, 47%) as a mixture of enantiomers as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 8.24-8.18 (m, 1H), 7.84 (td, *J*=8.0, 2.5 Hz, 1H), 6.94 (dd, *J*=8.1, 2.9 Hz, 1H), 5.26 (dd, *J*=8.8, 5.8 Hz, 1H), 4.27-4.20 (m, 1H), 3.98-3.88 (m, 1H), 2.91-2.77 (m, 1H), 2.34-2.24 (m, 1H), 1.50 (s, 9H). LC-MS (Method A): *m/z* = 269.2 [M+H]⁺, 0.90 min.

INTERMEDIATE 26

PREPARATION OF (2E)-3-(6-FLUOROPYRIDIN-3-YL)PROP-2-ENAL

6-Fluoropyridine-3-carbaldehyde (2.0 g, 16.0 mmol) was added to a solution of (triphenylphosphoranylidene)acetaldehyde (6.32 g, 20.8 mmol) in CHCl₃ (100 mL) and the resulting

mixture was stirred at room temperature for 18 h. The reaction was diluted with CH_2Cl_2 and the organic portion was washed with water, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (1.17 g, 48%) as a pale yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 9.76 (d, $J=7.3$ Hz, 1H), 8.44 (d, $J=2.5$ Hz, 1H), 8.09-7.98 (m, 1H), 7.50 (d, $J=16.1$ Hz, 1H), 7.06 (dd, $J=8.5$, 3.0 Hz, 1H), 6.74 (dd, $J=16.2$, 7.4 Hz, 1H). LC-MS (Method A): m/z = 152.0 $[\text{M}+\text{H}]^+$, 0.61 min.

INTERMEDIATE 27

PREPARATION OF *tert*-BUTYL N-[(1S)-1-(6-FLUOROPYRIDIN-3-YL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

(2E)-3-(6-Fluoropyridin-3-yl)prop-2-enal (**1.17 g**, 7.75 mmol) and *tert*-butyl N-hydroxycarbamate (1.23 g, 9.30 mmol) were sequentially added to a solution of (S)-(-)- α,α -diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (504 mg, 1.55 mmol) in CH_2Cl_2 (30 mL). The resulting mixture was stirred at room temperature for 5 h, diluted with MeOH (30 mL) and cooled to 0 °C. Sodium borohydride (442 mg, 11.6 mmol) was added in three portions and the resulting mixture was stirred at 0 °C for 45 min before being quenched with sat. NH_4Cl solution. The layers were separated and the aqueous portion was extracted twice with CH_2Cl_2 . The combined organic portions were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to give the desired compound (745 mg, 34%) as a colorless wax. ^1H NMR (400 MHz, CDCl_3) δ 8.25 (d, $J=2.5$ Hz, 1H), 7.95 (td, $J=8.0$, 2.5 Hz, 1H), 6.94 (dd, $J=8.4$, 2.9 Hz, 1H), 5.30 (dd, $J=10.9$, 5.1 Hz, 1H), 3.92-3.77 (m, 2H), 2.44 (dddd, $J=14.5$, 10.9, 7.2, 3.5 Hz, 1H), 2.12-1.95 (m, 1H), 1.49 (s, 9H). LC-MS (Method A): m/z = 287.2 $[\text{M}+\text{H}]^+$, 0.69 min.

INTERMEDIATE 28

PREPARATION OF *tert*-SVTYL (3S)-3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

Diisopropyl azodicarboxylate (511 μL , 2.60 mmol) was added dropwise to a solution of *tert*-butyl N-[(1S)-1-(6-fluoropyridin-3-yl)-3-hydroxypropyl]-N-hydroxycarbamate (745 mg, 2.60 mmol) and triphenylphosphine (817 mg, 3.12 mmol) in THF (30 mL), which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 5 min, diluted with EtOAc and washed with brine. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3CN , 100:0 to 0:100) to give the title compound (560 mg, 80%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 8.21 (d, $J=2.3$ Hz, 1H), 7.84 (td, $J=8.1$, 2.6 Hz, 1H), 6.94 (dd, $J=8.4$, 2.9 Hz, 1H), 5.26 (dd, $J=8.7$, 5.6 Hz, 1H), 4.23 (td, $J=8.0$, 3.4 Hz, 1H), 3.92 (td, $J=8.7$, 7.2 Hz, 1H), 2.84 (dddd, $J=12.3$, 8.8, 7.0, 3.5 Hz, 1H), 2.29 (dddd, $J=12.3$, 9.0, 7.8, 5.5 Hz, 1H), 1.50 (s, 9H). LC-MS (Method A): m/z = 269.2 $[\text{M}+\text{H}]^+$, 0.90 min. e.e.=90.6% as determined on a Chiraipak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/EtOH 50/50% v/v.

INTERMEDIATE 29

PREPARATION OF 2-FLUORO-5-[(3S)-1,2-OXAZOLIDIN-3-YL]PYRIDINE

Trifluoroacetic acid (1 mL) was added to a solution of *tert*-butyl (3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidine-2-carboxylate (560 mg, 2.09 mmol) in CH₂Cl₂ (5 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 20 min, warmed to room temperature and stirred for a further 3 h. Volatiles were removed under reduced pressure. The residue was partitioned between 3/4 3/4 and 1M NaOH solution. The layers were separated and the aqueous portion was extracted twice with CH₂Cl₂. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (320 mg) as a colorless oil which was directly used in the synthesis of Compound 61 (Example 31B). ¹H NMR (400 MHz, CDCl₃) δ 8.23 (d, *J* = 5.5 Hz, 1H), 7.89 (td, *J* = 8.0, 2.5 Hz, 1H), 6.94 (dd, *J* = 8.4, 2.9 Hz, 1H), 5.77-5.15 (m, 1H), 4.74-4.47 (m, 1H), 4.20-4.11 (m, 1H), 4.06-3.82 (m, 1H), 2.81-2.70 (m, 1H), 2.39-2.21 (m, 1H). LC-MS (Method A): *m/z* = 169.0 [M+H]⁺, 0.39 min.

INTERMEDIATE 30

PREPARATION OF (2E)-3-(5-CHLOROPYRIDIN-2-YL)PROP-2-ENAL

5-Chloropyridine-2-carbaldehyde (2.0 g, 14.2 mmol) was added to a solution of 2-(triphenylphosphoranylidene)acetaldehyde (5.6 g, 18.4 mmol) in CH₂Cl₂ (100 mL). The reaction was stirred at room temperature for 18 h, diluted with CH₂Cl₂ and washed with water. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (1.25 g, 53%) as a light brown solid. ¹H NMR (400 MHz, CDCl₃) δ 9.82 (d, *J* = 7.5 Hz, 1H), 8.67 (d, *J* = 2.5 Hz, 1H), 7.76 (dd, *J* = 8.4, 2.4 Hz, 1H), 7.53-7.47 (m, 2H), 7.09 (dd, *J* = 15.8, 7.8 Hz, 1H). LC-MS (Method A): *m/z* = 168.0 [M+H]⁺, 0.77 min.

INTERMEDIATE 31

PREPARATION OF TERT-BUTYL N-[(1S)-1-(5-CHLOROPYRIDIN-2-YL)-3-OXOPROPYL]-N-HYDROXYCARBAMATE

(2E)-3-(5-chloropyridin-2-yl)prop-2-enal (1.25 g, 7.48 mmol) and *tert*-butyl N-hydroxycarbamate (1.19 g, 8.98 mmol) were sequentially added to a solution of (S)-(-)-α,α-diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (608 mg, 1.87 mmol) in CH₂Cl₂ (30 mL). The resulting solution was stirred at room temperature for 24 h, diluted with CH₂Cl₂ and washed with sat. NH₄Cl solution and brine. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford an impure fraction of the title compound (1.10 g), which was progressed to the next step without any further purification. LC-MS (Method A): *m/z* = 301.1 [M+H]⁺, 0.88 min.

INTERMEDIATE 32

PREPARATION OF TERT-BUTYL N-[(1S)-1-(5-CHLOROPYRIDIN-2-YL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

Sodium borohydride (139 mg, 3.67 mmol) was added in three portions to a solution of tert-butyl N-[(1S)-1-(5-chloropyridin-2-yl)-3-oxopropyl]-N-hydroxycarbamate (1.10 g) in MeOH (35 mL), which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 45 min and quenched with sat. NH₄Cl solution. The aqueous portion was extracted twice with Et₂O. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (850 mg, 38% over two steps) as a yellow wax. ¹H NMR (400 MHz, CDCl₃) δ 8.47 (d, J=2.0 Hz, 1H), 8.08-7.94 (m, 1H), 7.71 (dd, J=8.3, 2.5 Hz, 1H), 7.34-7.28 (m, 1H), 5.47 (dd, J=10.8, 4.3 Hz, 1H), 3.86-3.78 (m, 2H), 2.41-2.29 (m, 2H), 2.17-2.07 (m, 1H), 1.49 (s, 9H). LC-MS (Method A): m/z = 303.2 [M+H]⁺, 0.80 min.

INTERMEDIATE 33

PREPARATION OF TERT-BUTYL (3S)-3-(5-CHLOROPYRIDIN-2-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

DIAD (556 μL, 2.83 mmol) was added dropwise to a solution of tert-butyl N-[(1S)-1-(5-chloropyridin-2-yl)-3-hydroxypropyl]-N-hydroxycarbamate (850 mg, 2.83 mmol) and triphenylphosphine (890 mg, 3.40 mmol) in THF (30 mL), which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 5 min, diluted with EtOAc and washed with brine. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound (700 mg, 87%) as a 5.5:94.5 mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 50/50% v/v and a flow rate of 1.8 mL/min to afford the title compound (599 mg, 74%) as a pale yellow wax. ¹H NMR (400 MHz, CDCl₃) δ 8.52 (dd, J=2.5, 0.5 Hz, 1H), 7.66 (dd, J=8.5, 2.5 Hz, 1H), 7.49 (d, J=8.5 Hz, 1H), 5.34 (dd, J=8.9, 4.9 Hz, 1H), 4.19-4.11 (m, 1H), 3.93 (q, J=8.3 Hz, 1H), 2.85-2.74 (m, 1H), 2.57 (dtd, J=12.7, 8.0, 4.9 Hz, 1H), 1.50 (s, 9H). LC-MS (Method A): m/z = 285.1 [M+H]⁺, 1.01 min. e.e. = 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate = 1.0 mL/min, retention time: 12.2 min.

INTERMEDIATE 34

PREPARATION OF 5-CHLORO-2-[(3S)-1,2-OXAZOLIDIN-3-YL]PYRIDINE

TFA (1 mL) was added to a solution of tert-butyl (3S)-3-(5-chloropyridin-2-yl)-1,2-oxazolidine-2-carboxylate (599 mg, 2.10 mmol) in CH₂Cl₂ (5 mL). The reaction was stirred at room temperature for 3h. Volatiles were removed under reduced pressure. The crude residue was dissolved in CH₂Cl₂ and the organic portion was washed with 1M NaOH solution and water, dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (348 mg) as a pale yellow wax. ¹H NMR (400 MHz, CDCl₃) δ 8.51 (d, J=2.3 Hz, 1H), 7.65 (dd, J=8.3, 2.5 Hz, 1H), 7.58-7.32 (m, 1H),

6.08–5.36 (m, 1H), 4.72–4.41 (m, 1H), 4.21–3.82 (m, 2H), 2.80–2.56 (m, 1H), 2.53–2.33 (m, 1H). LC-MS (Method A): $m/z = 185.0$ $[M+H]^+$, 0.52 min.

INTERMEDIATE 35

PREPARATION OF 3,3-DIFLUORO-2-METHYLPROP-2-ENOIC ACID

5 Sodium borohydride (250 mg, 6.57 mmol) was added to a solution of 2- (trifluoromethyl)prop-2-enoic acid (460 mg, 3.28 mmol) in THF (10 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 30 min and quenched with 1M HCl solution. The organic solvent was evaporated under reduced pressure and the aqueous portion was extracted twice with EtOAc. The combined organic portions were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford a 3:2 mixture of 3,3,3-trifluoro-2-methylpropanoic acid and the title 3,3-difluoro-2-methylprop-2-enoic acid (270 mg). This mixture was progressed to the next step without further purification. Analytical data for 3,3-difluoro-2-methylprop-2-enoic acid: LC-MS (Method A): $m/z = 120.9$ $[M-H]^-$, 0.57 min.

INTERMEDIATE 36

PREPARATION OF 3,3-DIFLUORO-2-METHYLPROPANOIC ACID

15 Pd/C (10% wt Degussa type) was added to a solution of the mixture containing 3,3-difluoro-2-methylprop-2-enoic acid (270 mg) in MeOH (7 mL) and the resulting mixture was stirred under an atmosphere of H_2 for 20 h. The mixture was filtered from the catalyst and the filtrate was concentrated under reduced pressure to afford a 2:1 mixture of 3,3,3-trifluoro-2-methylpropanoic acid and the title 3,3-difluoro-2-methylpropanoic acid (100 mg). This mixture was progressed to the next step without further purification. Analytical data for 3,3-difluoro-2-methylpropanoic acid: LC-MS (Method A): $m/z = 122.9$ $[M-H]^-$, 0.53 min.

INTERMEDIATE 37

PREPARATION OF 3-CYANO-2,2-DIMETHYLPROPANOIC ACID

25 To a stirred solution of methyl 3-cyano-2,2-dimethylpropanoate (100 mg, 0.7 mmol) in a mixture of THF/MeOH/ H_2O (2:1:1, 4 mL), LiOH (34 mg, 1.4 mmol) was added and the mixture was stirred at room temperature for 1.5 h. The mixture was concentrated under reduced pressure. The residue was diluted with water and extracted with EtOAc. The organic portion was acidified to pH ~ 5 with 1N HCl solution and then extracted twice with EtOAc. The combined organic phases were dried over Na_2SO_4 and evaporated under reduced pressure to afford the title compound (68 mg, 76%) as a white solid. ^1H NMR (400 MHz, $CDCl_3$) δ 2.65 (s, 2H), 1.44 (s, 6H). LC-MS (Method A): $m/z = 127.9$ $[M+H]^+$, 0.50 min.

INTERMEDIATE 38

PREPARATION OF TERT-BUTYL N-[(1S)-1-(4-FLUOROPHENYL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

35 To a solution of (S)-(-)- α,α -diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (1.3 g, 4 mmol) in CH_2Cl_2 (45 mL) was added 4-fluorocinnamaldehyde (3.0 g, 20 mmol) and tert-butyl N-hydroxycarbamate (3.19 g, 24 mmol) and the mixture was stirred at room temperature for 20 h. The reaction mixture was diluted with MeOH (45 mL) and cooled to 0°C, then $NaBH_4$ (1.56 g, 40 mmol) was

added portionwise. The reaction mixture was stirred for 30 min at room temperature, then cooled again to 0°C and quenched with 1N HCl solution. The reaction mixture was extracted twice with CH₂Cl₂ and the organic phase was dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 70:30) to afford the title compound (1.7 g, 30%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.43-7.36 (m, 2H), 7.06-6.98 (m, 2H), 5.21 (dd, J=10.3, 5.2 Hz, 1H), 3.88-3.73 (m, 2H), 2.46-2.39 (m, 1H), 2.08-1.99 (m, 1H), 1.45 (s, 9H).

INTERMEDIATE 39

PREPARATION OF TERT-BUTYL 3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a solution of tert-butyl N-[(1S)-1-(4-fluorophenyl)-3-hydroxypropyl]-N-hydroxycarbamate (1.7 g, 6 mmol) and triphenylphosphine (1.88 g, 7.2 mmol) in anhydrous THF (20 mL) at 0°C, DIAD (1.17 mL, 6 mmol) was added dropwise. The mixture was stirred for 15 min then concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 70:30) to afford the title compound (1.4 g) as a 18.3:81.7 mixture of enantiomers as a white solid. This mixture of enantiomers was resolved by chiral HPLC on a Chiralcel OJ-H (25 x 3.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 70/30 % v/v and a flow rate of 1.8 mL/min to afford the title compound (940 mg, 58%) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.39-7.29 (m, 2H), 7.10-6.98 (m, 2H), 5.19 (dd, J=8.7, 5.5 Hz, 1H), 4.24-4.14 (m, 1H), 3.90 (ddd, J=8.9, 8.1, 7.0 Hz, 1H), 2.78 (dddd, J=12.3, 8.7, 7.0, 3.7 Hz, 1H), 2.28 (dddd, J=12.2, 9.0, 7.7, 5.6 Hz, 1H), 1.47 (s, 9H). Chiralcel (S)-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 1 mL/min, retention time: 7.3 min.

INTERMEDIATE 40

PREPARATION OF (3S)-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINE

Trifluoroacetic acid (2.5 mL) was added at 0 °C to a solution of tert-butyl 3-(4-fluorophenyl)-1,2-oxazolidine-2-carboxylate (900 mg, 3.37 mmol) in CH₂Cl₂ (15 mL). The mixture was stirred at room temperature for 4h, then concentrated under reduced pressure. The residue was dissolved in CH₂Cl₂ and the organic portion was washed twice with sat. NaHCO₃ solution, dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (380 mg, 67%) as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 7.43-7.31 (m, 2H), 7.12-6.98 (m, 2H), 4.49 (dd, J=8.1, 6.1 Hz, 1H), 4.14-3.94 (m, 2H), 2.68 (dtd, J=12.2, 8.3, 5.6 Hz, 1H), 2.27 (dddd, J=12.5, 8.7, 6.8, 6.1 Hz, 1H).

INTERMEDIATE 41

PREPARATION OF 4-[(1E)-3-OXOPROP-1-EN-1-YL]BENZONITRILE

4-Cyanobenzaldehyde (20 g, 152 mmol) was added to a solution of (triphenylphosphoranylidene) acetaldehyde (60.3 g, 198 mmol) in CH₂Cl₂ (500 mL). The reaction was stirred at room temperature for 20 hours. The reaction was concentrated under reduced pressure. The crude product was triturated with Et₂O and the solid was filtered off washing several times with Et₂O. The filtrate and washings were evaporated under reduced pressure and the resulting crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 80:20) to afford the title compound (9.97 g, 41%) as a light yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 9.78 (d, J=7.4 Hz, 1H), 7.78-7.72 (m, 2H), 7.71-7.65

(m, 2H), 7.48 (d, $J=16.1$ Hz, 1H), **6.79** (dd, $J=16.1, 7.2$ Hz, 1H). LC-MS (Method A): $m/z = 158.0$ [M+H]⁺, 0.76 min.

INTERMEDIATE 42

PREPARATION OF TERT-BUTYL N-[1-(4-CYANOPHENYL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

To a solution of catalyst (S)-(-)- α,α -diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (3.9 g, 12 mmol) in CH_2Cl_2 (400 mL) was added 4-[(1E)-3-oxoprop-1-en-1-yl]benzonitrile (**9.97** g, 60 mmol) and tert-butyl N-hydroxycarbamate (**9.57** g, 72 mmol) and mixture was stirred at room temperature for 24 h. The reaction mixture was concentrated under reduced pressure, then diluted with MeOH (300 mL) and cooled to 0°C. NaBH_4 (4.7 g, 120 mmol) was added portionwise, then the mixture was stirred for 30 min at 0°C and at room temperature for 2h. The mixture was cooled to 0°C and quenched with 1N HCl solution. The mixture was extracted twice with CH_2Cl_2 and the combined organic phases were dried over Na_2SO_4 and evaporated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100: to 30:70) to afford the title compound (**6.97** g, 39%) as a yellow oil. ¹H NMR (400 MHz, CDCl_3) δ 7.66-7.60 (m, 2H), 7.56-7.51 (m, 2H), 5.26 (dd, $J=10.9, 4.9$ Hz, 1H), 3.92-3.74 (m, 2H), 2.40 (dddd, $J=14.6, 11.0, 7.5, 3.6$ Hz, 1H), 2.08-1.98 (m, 1H), **1.43** (s, 9H).

INTERMEDIATE 43

PREPARATION OF TERT-BUTYL (3S)-3-(4-CYANOPHENYL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a solution of tert-butyl N-[1-(4-cyanophenyl)-3-hydroxypropyl]-N-hydroxycarbamate (6.9 g, 23.8 mmol) and triphenylphosphine (7.48 g, 28.56 mmol) in anhydrous THF (150 mL) at 0°C, DIAD (4.67 mL, 23.8 mmol) was added dropwise. The mixture was stirred for 1.5 h allowing the temperature to gradually reach room temperature. The mixture was concentrated under reduced pressure, and the crude product was dissolved in the minimum quantity of CH_2Cl_2 . Cyclohexane was added until a white solid, that resulted to be triphenylphosphine oxide, precipitated. The solid was filtered off and the filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound as a **6:94** mixture of enantiomers (5.2 g, 79%). 1.7 g of this mixture of enantiomers was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1 % isopropylamine) 50/50 v/v and a flow rate of 18 mL/min to afford the title compound (1.43 g) as a colorless oil. ¹H NMR (400 MHz, CDCl_3) δ 7.68-7.62 (m, 2H), 7.52-7.47 (m, 2H), 5.27 (dd, $J=8.8, 5.7$ Hz, 1H), 4.24-4.16 (m, 1H), 3.95-3.86 (m, 1H), 2.85 (dddd, $J=12.3, 8.8, 6.9, 3.4$ Hz, 1H), 2.26 (dddd, $J=12.2, 9.1, 7.7, 5.6$ Hz, 1H), **1.48** (s, 9H).

INTERMEDIATE 44

PREPARATION OF 4-(1,2-OXAZOLIDIN-3-YL)BENZONITRILE

Trifluoroacetic acid (9.77 mL) was added at 0 °C to a solution of tert-butyl 3-(4-cyanophenyl)-1,2-oxazolidine-2-carboxylate (**6:94** mixture of enantiomers, 3.5 g, 12.77 mmol) in CH_2Cl_2 (50 mL). The mixture was stirred at room temperature for 2h. The mixture was then concentrated under reduced

pressure. The residue was dissolved in CH_2Cl_2 and washed twice with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (1.45 g, 65%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 7.68-7.60 (m, 2H), 7.57-7.49 (m, 2H), 5.47 (br.s, 1H), 4.61 (br.s, 1H), 4.13 (td, $J=8.2, 5.0$ Hz, 1H), 3.89 (br.s, 1H), 2.76 (dtd, $J=11.1, 8.5, 5.0$ Hz, 1H), 2.27 (dddd, $J=12.3, 8.5, 7.3, 4.9$ Hz, 1H). LC-MS (Method A): $m/z = 175.1$ $[\text{M}+\text{H}]^+$, 0.54 min.

INTERMEDIATE 45

PREPARATION OF METHYL 3-BROMO-2,2-DIMETHYLPROPANOATE

To a solution of 3-bromo-2,2-dimethylpropanoic acid (900 mg, 5 mmol) in toluene/methanol (4:1, 20 mL) at 0°C was added (trimethylsilyl)diazomethane (2 M in hexane, 5 mL, 10 mmol) slowly under a nitrogen atmosphere. The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture was diluted with water and extracted with EtOAc. The organic phase was dried over Na_2SO_4 and evaporated under reduced pressure to afford the title compound (450 mg, 46%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 3.73 (s, 3H), 3.51 (s, 2H), 1.33 (s, 6H).

INTERMEDIATE 46

PREPARATION OF 2,2-DIMETHYL-3-(1-H-PYRAZOL-1-YL)PROPANOIC ACID

A suspension of methyl 3-bromo-2,2-dimethylpropanoate (195 mg, 1 mmol), pyrazole (82 mg, 1.2 mmol), CS_2CO_3 (390 mg, 1.2 mmol) and tetrabutylammonium iodide (184 mg, 0.5 mmol) in DMF (2 mL) was stirred at 60°C for 4h. The mixture was filtered and the organic phase was evaporated under reduced pressure. The residue was re-dissolved in a 2:1 mixture of THF/100 (2.5 mL), LiOH (96 mg, 4 mmol) was added and the mixture was stirred at room temperature overnight. The reaction mixture was washed with EtOAc, the aqueous phase was acidified to pH ~ 5 with 1N HCl solution and then extracted with EtOAc. The organic portion was dried over Na_2SO_4 and evaporated under reduced pressure. The crude product was purified by reverse phase column chromatography (water-acetonitrile, 100:0 to 60:40) to afford the title compound (50 mg, 30%). ^1H NMR (400 MHz, CDCl_3) δ 7.55 (d, $J=2.0$ Hz, 1H), 7.46 (d, $J=2.0$ Hz, 1H), 6.26 (t, $J=2.1$ Hz, 1H), 4.35 (s, 2H), 1.23 (s, 6H).

INTERMEDIATE 47

PREPARATION OF 6-CHLORO-5-FLUOROPYRIDINE-3-CARBALDEHYDE

Butyllithium (2.5 M in hexane, 13.8 mL, 34.4 mmol) was added to a solution of 5-bromo-2-chloro-3-fluoro-pyridine (5.14 g, 24.6 mmol) in anhydrous diethyl ether (80 mL) at -78°C . After 30 minutes, anhydrous DMF (24.8 mL, 320 mmol) was added and the mixture was stirred at -78°C . After 30 min water was added and the mixture was extracted three times with EtOAc. The combined organic phases were washed with brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 95:5 to 60:40) to afford the title compound (2.19 g) as a pale yellow solid. LC-MS (Method A): $m/z = 177.9$ $[\text{M}+\text{H}+\text{H}_2\text{G}]^+$, 0.71 min.

INTERMEDIATE 48

PREPARATION OF (2E)-3-(5-FLUORO-6-METHYLPYRIDIN-3-YL)PROP-2-ENAL

6-Chloro-5-fluoropyridine-3-carbaldehyde (2.19 g) was dissolved in 1,4-dioxane (24 mL) and trimethylboroxine (2.32 mL, 16.4 mmol) was added, followed by K_2CO_3 (5.68 g, 41.1 mmol). The mixture was degassed by applying alternatively vacuum and nitrogen. $Pd(PPh_3)_4$ (791 mg, 0.68 mmol) was then added. The mixture was then heated at 100 °C for 1 h 30 min, then cooled to room temperature and filtered over a pad of Celite. The solvent was evaporated under reduced pressure to give a red solid (3.33 g). This solid was dissolved in CH_2Cl_2 (50 mL) and (triphenylphosphoranylidene)acetaldehyde (5.0 g, 16.4 mmol) was then added at 10 °C. The mixture was allowed to warm to room temperature and stirred overnight. The mixture was then concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 88:12 to 0:100) to afford the title compound (1.21 g, 30% yield over 3 steps) as a yellow solid. ^1H NMR (400 MHz, $CDCl_3$) δ 9.76 (d, $J=7.3$ Hz, 1H), 8.50 (s, 1H), 7.56-7.45 (m, 2H), 6.74 (dd, $J=16.0, 7.3$ Hz, 1H), 2.61 (d, $J=3.0$ Hz, 3H). LC-MS (Method A): m/z = 166.0 $[M+H]^+$, 0.67 min.

INTERMEDIATE 49

PREPARATION OF TERT-BUTYL N-[(1S)-1-(5-FLUORO-6-METHYLPYRIDIN-3-YL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

To a solution of (5)-(-)-a,a-diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (241 mg, 0.74 mmol) in anhydrous CH_2Cl_2 (10 mL) were added a solution of (2E)-3-(5-fluoro-6-methylpyridin-3-yl)prop-2-enal (612 mg, 3.70 mmol) in anhydrous CH_2Cl_2 (8 mL) and tert-butyl N-hydroxy carbamate (592 mg, 4.45 mmol) and the mixture was stirred at room temperature. After 27 h the reaction mixture was diluted with MeOH (13 mL) and cooled to 0 °C. $NaBH_4$ (280 mg, 7.4 mmol) was added portionwise then the mixture was stirred at 0 °C for 10 min and quenched with sat. NH_4Cl solution and stirred at room temperature for 10 min. The mixture was then extracted twice with CH_2Cl_2 and twice with EtOAc. The combined organic phases were dried over Na_2SO_4 and evaporated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 80:20 to 0:100) to afford the title compound (206 mg, 18% yield) as a brown oil. ^1H NMR (400 MHz, $CDCl_3$) δ 8.22 (s, 1H), 7.46 (dd, $J=10.0, 2.0$ Hz, 1H), 5.33 (dd, $J=10.5, 5.0$ Hz, 1H), 3.84-3.73 (m, 2H), 2.54-2.34 (m, 1H), 2.43 (d, $J=3.0$ Hz, 3H), 2.06-1.96 (m, 1H), 1.49 (s, 9H). LC-MS (Method A): m/z = 301.2 $[M+H]^+$, 0.66 min.

INTERMEDIATE 50

PREPARATION OF TERT-BUTYL (3S)-3-(5-FLUORO-6-METHYLPYRTOIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a solution of tert-butyl N-[(1S)-1-(5-fluoro-6-methylpyridin-3-yl)-3-hydroxypropyl]-N-hydroxycarbamate (206 mg, 0.68 mmol) and PPh_3 (214 mg, 0.816 mmol) in anhydrous THF (7 mL) at 0 °C, DIAD (0.135 mL, 0.68 mmol) was added dropwise. After 10 min a further addition of DIAD (0.045 mL, 0.23 mmol) was made. The mixture was then concentrated under reduced pressure and the crude product was purified twice by column chromatography (first column cyclohexane-EtOAc, 88:12 to 0:100; second column cyclohexane-EtOAc, 70:30 to 0:100) to afford the title compound as a colorless oil

(120 mg, 62% yield). ^1H NMR (400 MHz, CDCl_3) δ 8.30 (s, 1H), 7.39 (dd, $J=10.0, 1.5$ Hz, 1H), 5.25 (dd, $J=8.5, 5.5$ Hz, 1H), 4.21 (dd, $J=8.0, 3.5$ Hz, 1H), 3.95-3.88 (m, 1H), 2.87-2.78 (m, 1H), 2.53 (d, $J=3.0$ Hz, 3H), 2.34-2.24 (m, 1H), 1.50 (s, 9H). LC-MS (Method A): $m/z = 283.3$ $[\text{M}+\text{H}]^+$, 0.89 min.

INTERMEDIATE 51

PREPARATION OF 3-FLUORO-2-METHYL-5-[(3S)-1,2-OXAZOLIDIN-3-YL]PYRIDINE

Trifluoroacetic acid (0.49 mL, 6.38 mmol) was added at 0 °C to a solution of tert-butyl (3S)-3-(5-fluoro-6-methylpyridin-3-yl)-1,2-oxazolidine-2-carboxylate (120 mg, 0.42 mmol) in anhydrous CH_2Cl_2 (4 mL). The mixture was stirred at room temperature for 3h30min then concentrated under reduced pressure. The residue was dissolved in CH_2Cl_2 and washed twice with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give the title compound as a pale yellow oil (65 mg, 85% yield), which was used in the following experiment without further purification. ^1H NMR (400 MHz, CDCl_3) δ 8.30 (s, 1H), 7.46 (d, $J=10.0$, 1H), 5.43 (br.s, 1H), 4.58 (br.s, 1H), 4.18-4.10 (m, 1H), 3.93 (br.s, 1H), 2.80-2.68 (m, 1H), 2.53 (d, $J=3.0$ Hz, 3H), 2.34-2.24 (m, 1H). LC-MS (Method A): $m/z = 183.1$ $[\text{M}+\text{H}]^+$, 0.40 min.

INTERMEDIATE 52

PREPARATION OF 6-CYANO-2-OXASPIRO[3.3]HEPTANE-6-CARBOXYLIC ACID

To a stirred solution of methyl 6-cyano-2-oxaspiro[3.3]heptane-6-carboxylate (181 mg, 1 mmol) in a mixture of THF/MeOH/ H_2O 2:1:1 (4 mL) was added LiOH (48 mg, 2 mmol) and the mixture was stirred at room temperature overnight. The mixture was concentrated under reduced pressure, the residue was diluted with water and washed with EtOAc. The organic portion was acidified to pH ~ 5 with IN HCl solution and then extracted twice with EtOAc. The combined organic phases were dried over Na_2SO_4 and evaporated under reduced pressure but very little product was recovered. The aqueous phase was then evaporated under reduced pressure to afford the title compound (163 mg, 97%) as white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 4.51 (d, $J=4.4$ Hz, 4H), 2.68-2.60 (m, 2H), 2.50-2.45 (m, 2H). LC-MS (Method A): $m/z = 168.0$ $[\text{M}+\text{H}]^+$, 0.44 min.

INTERMEDIATE 53

4-[(1E)-3-HYDROXY(3,3- $^2\text{H}_2$)PROP-1-EN-1-YL]BENZONITRILE

A suspension of (2E)-3-(4-cyanophenyl)prop-2-enoic acid (750 mg, 4.33 mmol), oxalyl chloride (0.385 mL, 4.54 mmol) and 1 drop of DMF in CH_2Cl_2 (15 mL) was stirred at room temperature for 30 min to give a clear solution. After checking the complete formation of the acyl chloride, the mixture was evaporated and the residue was dissolved in THF (15 mL), then NaBD_4 (544 mg, 13 mmol) was added at room temperature. The resulting mixture was stirred at room temperature for 15 min, then CD_3OD (3.75 mL) was added dropwise over 1h. The reaction was quenched with D_2O , then with IN HCl solution, and the aqueous phase was extracted with Et_2O . The organic phase was separated and washed with brine, dried over Na_2SO_4 and filtered. The solvent was evaporated under reduced pressure to give a white solid. The crude product purified by column chromatography (cyclohexane-EtOAc, 85:15 to 65:35) to afford the title compound (350 mg, 50%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.86-7.73 (m, 2H), 7.68-7.56 (m, 2H), 6.73-6.53 (m, 2H), 4.99-4.86 (m, 1H). LC-MS (Method A): $m/z = 162.1$ $[\text{M}+\text{H}]^+$,

0.70 min.

INTERMEDIATE 54

4-[(1E)-3-oxo(3,3-²H)prop-1-en-1-yl]benzonitrile

A suspension of 4-[(1E)-3-hydroxy(3,3-²H)prop-1-en-1-yl]benzonitrile (350 mg, 2.17 mmol) and MnO₂ (943 mg, 10.8 mmol) in CH₂Cl₂ (20 mL) was stirred at room temperature for 6 h. Further MnO₂ (313 mg, 3.6 mmol) was added, and the mixture was stirred at room temperature overnight. Further MnO₂ (383 mg, 4.4 mmol) was added and the mixture was stirred at room temperature overnight. Further MnO₂ (380 mg, 4.4 mmol) was added and the mixture was stirred for 5 h. The suspension was filtered and the solvent was evaporated under reduced pressure to give the title compound (303 mg, 88%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.98-7.91 (m, 4H), 7.85-7.77 (m, 1H), 7.01 (d, J=16.1 Hz, 1H). LC-MS (Method A): m/z = 159.0 [M+Hf, 0.78 min.

INTERMEDIATE 55

TERT-BUTYL N-[(1S)-1-(4-CYANOPHENYL)-3-HYDROXY(3,3-²H)PROPYL]-N-HYDROXYCARBAMATE

tert-Butyl N-hydroxycarbamate (305 mg, 2.29 mmol) was added to a solution of 4-[(1E)-3-oxo(3-²H)prop-1-en-1-yl]benzonitrile (303 mg, 1.91 mmol) and (S)-(-)-a,a-diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (190 mg, 0.58 mmol) in CH₂Cl₂ (9 mL). The mixture was stirred at room temperature for 6 h, then CD₃OD (5 mL) was added and the mixture was cooled to 0 °C and NaBD₄ (120 mg, 2.86 mmol) was added in small portions. The resulting mixture was stirred at 0 °C for 30 min. The mixture was quenched with D₂O, then with sat. NH₄Cl solution. The layers were separated and the aqueous portion was extracted twice with CH₂Cl₂. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 80:20 to 25:75) to afford the title compound (145 mg, 42%) as a pale yellow oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.24 (s, 1H), 7.85-7.73 (m, 2H), 7.56-7.48 (m, 2H), 5.20 (dd, J=6.9, 9.0 Hz, 1H), 4.50 (s, 1H), 2.11 (dd, J=13.7, 8.9 Hz, 1H), 1.86 (dd, J=13.8, 6.0 Hz, 1H), 1.35 (s, 9H). LC-MS (Method A): m/z = 295.2 [M+Hf, 0.77 min.

INTERMEDIATE 56

TERT-BUTYL N-[(1S)-1-(4-CYANOPHENYL)-3-HYDROXY(3,3-²H)PROPYL]-N-HYDROXYCARBAMATE

A solution of D1AD (0.1 mL, 0.49 mmol) in anhydrous THF (1 mL) was added dropwise to a solution of tert-butyl N-[(1S)-1-(4-cyanophenyl)-3-hydroxy(3,3-²H)propyl]-N-hydroxy carbamate (145 mg, 0.49 mmol) and triphenylphosphine (154 mg, 0.59 mmol) in anhydrous THF (12 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 15 min, diluted with EtOAc and washed with brine. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 75:25 to 60:40) to afford the title compound in a ~1:1 mixture with N'-[(propan-2-yloxy)carbonyl](propan-2-yloxy)carbohydrazide (190 mg) as a white solid. This mixture was progressed in the following step without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.69-7.60 (m, 2H), 7.53-7.43 (m, 2H), 5.25 (dd, J=8.8, 5.5 Hz, 1H), 2.82 (dd, J=12.3, 9.0 Hz, 1H), 2.24 (dd, J=12.4, 5.6 Hz, 1H), 1.47 (s, 9H). LC-MS (Method A): m/z = 277.0 [M+Hf, 1.00 min.

INTERMEDIATE 57

4-[(3S)-(5,5-²H₂)-1,2-oxazolidin-3-yl]benzonitrile

A solution of tert-butyl (3S)-3-(4-cyanophenyl)(5,5-²H₂)-L2-oxazolidine-2-carboxylate (190 mg) in CH₂Cl₂ (15 ml) and TFA (3 mL) was stirred at 0 °C for 30 minutes, then at room temperature for 3h. The mixture was evaporated to dryness. The residue was purified by ion exchange chromatography (SCX, MeOH then 0.2M NH₃ in MeOH) to afford the title compound (60 mg, 69% over two steps) as a colorless oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.82-7.73 (m, 2H), 7.58 (d, *J* = 8.3 Hz, 2H), 6.55 (br.s, 1H), 4.52 (br.s, 1H), 2.72-2.58 (m, 1H), 2.05 (dd, *J* = 12.0, 4.5 Hz, 1H). LC-MS (Method A): *m/z* = 177.0 [M+H]⁺, 0.53 min.

INTERMEDIATE 58

TERT-BUTYL (3S)-3-(4-CYANOPHENYL)-5-HYDROXY-1,2-OXAZOLIDINE-2-CARBOXYLATE

tert-Butyl N-hydroxy carbamate (631 mg, 4.74 mmol) was added to a solution of 4-[(1E)-3-oxo(3-²H)prop-1-en-1-yl]benzonitrile (622 mg, 3.95 mmol) and (S)-(-)-α,α-diphenyl-γ-pyrrolidinemethanol trimethylsilyl ether (321 mg, 0.98 mmol) in CH₂Cl₂ (30 mL). The mixture was stirred at room temperature overnight. The solution was concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 80:20 to 65:35) to afford the title compound (689 mg, 60%) as a pale yellow solid as a mixture of isomers. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.86-7.76 (m, 2H), 7.55-7.45 (m, 2H), 6.95 (br.s, 1H), 5.62 (br.s, 1H), 5.30 (t, *J* = 8.3 Hz, 1H), 2.64 (dd, *J* = 12.4, 8.4 Hz, 1H), 2.07 (ddd, *J* = 12.5, 8.1, 4.5 Hz, 1H), 1.47-1.36 (m, 9H). LC-MS (Method A): *m/z* = 291.1 [M+H]⁺, 0.90 min.

INTERMEDIATE 59

TERT-BUTYL N-[(1S)-1-(4-CYANOPHENYL)-3-HYDROXYBUTYL]-N-HYDROXYCARBAMATE

MeMgBr (3M solution in diethyl ether, 0.67 mL, 2.01 mmol) was added dropwise to a solution of tert-butyl (3S)-3-(4-cyanophenyl)-5-hydroxy-1,2-oxazolidine-2-carboxylate (585 mg, 2.01 mmol) in anhydrous THF (25 mL) under nitrogen at -20 °C. The mixture was stirred at -20 °C for 1h, then at 0 °C for 2h. Further MeMgBr (0.67 mL, 2.01 mmol) was added and the mixture was allowed to warm to room temperature and stirred overnight. The reaction was quenched with sat. NH₄Cl solution, then the aqueous phase was extracted with EtOAc. The combined organic phases were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 70:30 to 50:50) to afford the title compound (190 mg, 31%) as a -65:35 mixture of diastereoisomers. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.22 (s, 1H), 7.85-7.76 (m, 2H), 7.58-7.46 (m, 2H), 5.32-5.12 (m, 1H), 4.61-4.45 (m, 1H), 3.76-3.34 (m, 1H), 2.15-1.83 (m, 2H), 1.40-1.32 (m, 9H), 1.13-1.04 (m, 3H). LC-MS (Method A): *m/z* = 307.2 [M+H]⁺, 0.86 min.

INTERMEDIATE 60

4-[(3S)-5-METHYL-1,2-OXAZOLIDIN-3-YL]BENZONITRILE

A solution of DIAD (0.122 mL, 0.62 mmol) in anhydrous THF (2 mL) was added dropwise to a solution of tert-butyl N-[(1S)-1-(4-cyanophenyl)-3-hydroxybutyl]-N-hydroxycarbamate (190 mg, 0.62 mmol) and triphenylphosphine (195 mg, 0.744 mmol) in anhydrous THF (13 mL) which had been pre-

cooled to 0 °C. The reaction was stirred at 0 °C for 15 min, then it was concentrated under reduced pressure to give a yellow oil. The residue was dissolved in CH₂Cl₂ (12 mL) and TFA (3 mL) was added. The resulting solution was stirred at room temperature for 1h then the solvent was evaporated under reduced pressure. The crude product was purified by SCX cartridge. The desired product was purified by ion exchange chromatography (SCX, MeOH then 0.2M NH₃ in MeOH) to afford the title compound (100 mg, 85%) as a mixture of diastereoisomers as a yellow oil. LC-MS (Method A): m/z = 189.1 [M+H]⁺, 0.62 min.

INTERMEDIATE 61

TERT-BUTYL 2-(AZETIDIN-1-YL)-2-METHYLPROPANOATE

A mixture of tert-butyl α-bromoisobutyrate (0.5 mL, 5.4 mmol), azetidine (0.43 mL, 6.4 mmol) and potassium carbonate (2.23 g, 16.2 mmol) in anhydrous DMF (10 mL) was stirred at 60 °C under nitrogen for 1h 30 min. The mixture was diluted with EtOAc and washed with water. The organic phase was separated, washed several times with brine, dried over Na₂SO₄ and filtered. The solvent was evaporated under reduced pressure to give a colorless oil which was purified by column chromatography (cyclohexane-EtOAc, 20:80 to 40:60) to afford the title compound (300 mg, 56%) a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 3.49-3.37 (in. 4H), 2.07-1.91 (m, 2H), 1.50 (s, 9H), 1.14 (s, 6H). LC-MS (Method A): m/z = 200.2 [M+H]⁺, 0.36 min.

INTERMEDIATE 62

2-(AZETH)IN-1-YL)-2-METHYLPROPANOIC ACID HYDROCHLORIDE

A solution of tert-butyl 2-(azetidin-1-yl)-2-methylpropanoate (300 mg, 1.5 mmol) in HCl (4M in dioxane, 12 mL) was stirred at 0 °C for 30 min, then at room temperature overnight. Further HCl (4M in dioxane, 5 mL) was added and the mixture was stirred at room temperature for 2h. The volatiles were evaporated under reduced pressure, and the resulting solid residue was triturated with Et₂O and filtered to afford the title compound (214 mg, 79%) as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 14.55-13.71 (br.s, 1H), 11.27 (br.s, 1H), 4.07 (br.s, 4H), 2.42-2.14 (m, 2H), 1.46 (s, 6H). LC-MS (Method A): m/z = 144.0 [M+H]⁺, 0.15 min.

INTERMEDIATE 63

PREPARATION OF (2E)-3~[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]PROP-2-ENAL

A mixture of 5-bromo-2-(difluoromethyl)pyridine (1770 mg, 8.5 mmol), tris-di(4-aceiato)bis(o-(di-o-tolyl-phosphino)benzyl)dipalladium(II) (1600 mg, 1.7 mmol) and sodium acetate (79? mg, 9.35 mmol) was placed in a microwave tube under nitrogen. The tube was dried under vacuum, then acetonitrile (15 mL) and 2-propenal (4.66 mL, 30.6 mmol) was added consecutively to the reaction mixture. The mixture heated in a Biotage Initiator microwave oven at 140°C (7 cycles of 15 min each). The solids were filtered off and washed with MeOH. The filtrate was concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 0:100) to afford unreacted 5-bromo-2-(difluoromethyl)pyridine (690 mg) and the title compound (700 mg, 26% yield) as an orange oil. LC-MS (Method A): m/z = 184.1 [M+H]⁺, 0.69 min.

INTERMEDIATE 64

PREPARATION OF TERT-BUTYL N-[(1S)-1-[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

To a solution of catalyst (5)-(-)-a,a-diphenyl-2-pyrrolidinemethanol trimethylsilyl ether (0.540 mg, 1.66 mmol) in CH_2Cl_2 (40 mL), were added 3-[6-(difluoromethyl)pyridine-3-yl]prop-2-enal (0.76 g, 4.15 mmol) and tert-butyl N-hydroxycarbamate (662 mg, 5 mmol). The mixture was stirred at room temperature for 20h. The reaction was diluted with MeOH (40 mL), and NaBH_4 (0.324 g, 8.3 mmol) was added portionwise. The mixture was then stirred for 1.5 hour at room temperature. Volatiles were removed under reduced pressure and the crude product was purified by column chromatography (CH_2Cl_2 -MeOH, 100:0 to 95:5) to afford a first batch of the title compound (97 mg, 5%) as a yellow oil. LC-MS (Method A): $m/z = 319.2$ $[\text{M}+\text{H}]^+$, 0.92 min. The column was eluted further with CH_2Cl_2 -MeOH 80:20 to afford a second batch of the title compound (310 mg, 16%) as an orange oil. UPLC LC-MS (Method A): $m/z = 319.2$ $[\text{M}+\text{H}]^+$, 0.92 min.

INTERMEDIATE 65

PREPARATION OF TERT-BUTYL (3S)-3-[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a solution of tert-butyl N-[(1S)-1-[6-(difluoromethyl)pyridine-3-yl]-3-hydroxypropyl]-N-hydroxycarbamate (407 mg, 1.27 mmol) and PPh_3 (499 mg, 1.92 mmol) in anhydrous THF (9 mL) at room temperature, was added DIAD (0.377 mL, 1.92 mmol) dropwise. The mixture was stirred at the same temperature for 1 hour, then it was concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound (154 mg, 40 mmol) as a yellow oil. ^1H NMR (400MHz, CDCl_3) δ 8.65 (d, $J=2.0$ Hz, 1H), 7.88 (dd, $J=8.2, 2.1$ Hz, 1H), 7.64 (d, $J=8.0$ Hz, 1H), 6.81-6.49 (t, 1H), 5.31 (dd, $J=8.7, 5.6$ Hz, 1H), 4.23 (dt, $J=3.4, 8.0$ Hz, 1H), 3.93 (dt, $J=7.0, 8.7$ Hz, 1H), 2.87 (dddd, $J=12.3, 8.8, 7.0, 3.5$ Hz, 1H), 2.30 (dddd, $J=12.3, 9.1, 7.7, 5.8$ Hz, 1H), 1.49 (s, 9H). LC-MS (Method A): $m/z = 301.2$ $[\text{M}+\text{H}]^+$, 0.94 min.

INTERMEDIATE 66

PREPARATION OF 2-(DIFLUOROMETHYL)-5-[(3S)-1,2-OXAZOLIDIN-3-YL]PYRIDINE

To a solution of tert-butyl (3S)-3-[6-(difluoromethyl)-pyridin-3-yl]-1,2-oxazolidine-2-carboxylate (154 mg, 0.51 mmol) in CH_2Cl_2 , trifluoroacetic acid (0.5 mL, 6.6 mmol) was added. The mixture was stirred for 2 h. Sat. NaHCO_3 solution was added and the mixture was extracted twice with CH_2Cl_2 . The combined organic phases were concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 30:70) to afford the title compound (49 mg, 45%) as a yellow oil. ^1H NMR (400MHz, CDCl_3) δ 8.66 (d, $J=1.5$ Hz, 1H), 7.94 (dd, $J=8.2, 1.9$ Hz, 1H), 7.63 (d, $J=8.0$ Hz, 1H), 6.81-6.48 (t, $J=55.2$ Hz, 1H), 5.63-5.35 (m, 1H), 4.65 (br.s, 1H), 4.22-4.10 (m, 1H), 3.92 (br.s, 1H), 2.85-2.72 (m, 1H), 2.38-2.24 (m, 1H). LC-MS (Method A): $m/z = 201.1$ $[\text{M}+\text{H}]^+$, 0.51 min.

INTERMEDIATE 67

PREPARATION OF (2E)-3-(1-METHYL-1H-PYRAZOL-4-YL)PROP-2-ENAL

To a solution of 1-methyl-1H-pyrazole-4-carbaldehyde (4 g, 36.3 mmol) in N,N-dimethylformamide (50 mL) was added (formylmethylene)triphenylphosphorane (12.17 g, 40.0 mmol).

The reaction mixture was heated to 120 °C and stirred overnight. After being cooled to room temperature, the reaction mixture was quenched by the addition of water (50 mL), and extracted with EtOAc (3 x 50 mL). The combined organic layers was washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by reverse phase column chromatography (water-acetonitrile, 5:1) to afford the title compound (1.00 g, 20%) as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 9.59 (d, *J* = 7.8 Hz, 1H), 7.74 (s, 1H), 7.64 (s, 1H), 7.36 (d, *J* = 15.9 Hz, 1H), 6.50-6.42 (m, 1H), 3.96 (s, 3H).

INTERMEDIATE 68

PREPARATION OF TERT-BUTYL N-HYDROXY-N-[(1S)-3-HYDROXY-1-(1-METHYL-1H-PYRAZOL-4-YL)PROPYL] CARBAMATE

To a stirred solution of (S)-2-(diphenyl(trimethylsilyloxy)methyl)pyrrolidine (383 mg, 1.18 mmol) in CHCl₃ (20 mL) were added (2E)-3-(1-methyl-1H-pyrazol-4-yl)prop-2-enal (800 mg, 5.90 mmol) and tert-butyl N-hydroxycarbamate (940 mg, 7.00 mmol). After being stirred at room temperature for 9.5 days, the reaction mixture was diluted with methanol (20 mL) and cooled to 0 °C. NaBH₄ (447 mg, 11.8 mmol) was added portionwise. After being stirred at room temperature for 30 min, the mixture was cooled to 0 °C, quenched with sat. NH₄Cl solution (60 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (60 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:1) to afford the title compound (200 mg, 12%) as a yellow oil. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.09 (s, 1H), 7.98 (d, *J* = 1.2 Hz, 1H), 7.61 (d, *J* = 1.2 Hz, 1H), 5.12-5.07 (m, 1H), 4.49-4.46 (m, 1H), 3.85 (s, 3H), 3.44-3.39 (m, 1H), 3.33-3.28 (m, 1H), 2.01-1.94 (m, 1H), 1.83-1.76 (m, 1H), 1.39 (s, 9H). LC-MS (Method M): *m/z* = 272.2 [M+H]⁺, 0.77 min.

INTERMEDIATE 69

PREPARATION OF TERT-BUTYL (3S)-3-(1-METHYL-1H-PYRAZOL-4-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a stirred solution of tert-butyl N-hydroxy-N-[(1S)-3-hydroxy-1-(1-methyl-1H-pyrazol-4-yl)propyl]carbamate (200 mg, 0.74 mmol) and triphenylphosphine (640 mg, 2.44 mmol) in anhydrous THF (10 mL) was added DIAD (448 mg, 2.22 mmol) dropwise. After being stirred overnight at room temperature, the reaction mixture was concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:2) to afford the title compound (100 mg, 53%) as a white solid. LC-MS (Method M): *m/z* = 254.2 [M+H]⁺, 0.79 min.

INTERMEDIATE 70

PREPARATION OF (3S)-3-(1-METHYL-1H-PYRAZOL-4-YL)-1,2-OXAZOLIDINE-2-CARBOXYLIC ACID

To a solution of tert-butyl (3S)-3-(1-methyl-1H-pyrazol-4-yl)-1,2-oxazolidine-2-carboxylate (100 mg, 0.9 mmol) in 1,4-dioxane (2 mL) was added a solution of hydrogen chloride in 1,4-dioxane (4 M, 5 mL). After being stirred overnight at room temperature, the reaction mixture was concentrated under reduced pressure to afford the title compound (90.0 mg, crude) as a yellow solid, which was used directly in the next step without further purification. LC-MS (Method M): $m/z = 154.3$ $[M+H]^+$, 0.30 min.

INTERMEDIATE 71

PREPARATION OF (6-METHYLPYRAZIN-2-YL)METHANOL

To a solution of 6-methylpyrazine-2-carboxylic acid (8.00 g, 58.0 mmol) in N,N-dimethylformamide (50 mL) were added Cs_2CO_3 (37.8 g, 116 mmol) and iodomethane (12.3 g, 87.0 mmol). After being stirred overnight at room temperature, the reaction mixture was quenched with water (100 mL) and extracted with EtOAc (5 x 150 mL). The combined organic layers were washed with brine (100 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was then diluted with water (50 mL). NaBH_4 (12.5 g, 330 mmol) was added portionwise at 0 °C. After addition, the resulting mixture was stirred at room temperature for 30 min, cooled to 0 °C, diluted with water (150 mL) and extracted with EtOAc (8 x 100 mL). The combined organic layers were washed with brine (200 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the title compound (2.9 g, crude) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.50 (s, 1H), 8.42 (s, 1H), 5.55 (t, $J = 5.8$ Hz, 1H), 4.59 (d, $J = 5.5$ Hz, 2H), 2.47 (s, 3H).

INTERMEDIATE 72

PREPARATION OF 6-METHYLPYRAZINE-2-CARBALDEHYDE

To a solution of (6-methylpyrazin-2-yl)methanol (1.50 g, 12.1 mmol) in CH_2Cl_2 (30 mL) was added Dess-Martin periodinane (7.70 g, 18.1 mmol) at -15 °C. After being stirred at -15 °C for 3 hours, the reaction mixture was quenched by the addition of water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (60 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:9) to afford the title compound (540 mg, 36%) as a yellow solid. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.02 (s, 1H), 8.91 (s, 1H), 8.82 (s, 1H), 2.61 (s, 3H).

INTERMEDIATE 73

PREPARATION OF (2E)-3-(6-METHYLPYRAZIN-2-YL)PROP-2-ENAL

To a solution of 6-methylpyrazine-2-carbaldehyde (620 mg, 5.10 mmol) in THF (20 mL) was added (formylmethylene)triphenylphosphorane (1.70 g, 5.60 mmol). After being stirred overnight at room temperature, the reaction mixture was quenched by the addition of water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:3) to afford the title compound (370 mg, 49%) as a yellow

solid. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 9.81 (d, J = 7.9 Hz, 1H), 8.85 (s, 1H), 8.61 (s, 1H), 7.83 (d, J = 15.8 Hz, 1H), 7.16-7.08 (m, 1H), 2.57 (s, 3H).

INTERMEDIATE 74

PREPARATION OF N-HYDROXY-N-[(1S)-3-HYDROXY-1-(6-METHYLPYRAZIN-2-YL)PROPYL]-2,2-DIMETHYLPROPANAMIDE

To a stirred solution of (S)-2-(diphenyl(trimethylsilyloxy)methyl)pyrrolidine (198 mg, 0.60 mmol) in CHCl_3 (30 mL) were added (2E)-3-(6-methylpyrazin-2-yl)prop-2-enal (300 mg, 2.00 mmol) and N-hydroxypivalamide (284 mg, 2.4 mmol). After being stirred overnight at room temperature, the reaction mixture was diluted with MeOH (15 mL) and cooled to 0 °C. NaBH_4 (154 mg, 4.00 mmol) was added. The reaction mixture was stirred at room temperature for 30 min, then cooled to 0 °C and quenched with sat. NH_4Cl solution (30 mL). The mixture was extracted with CH_2Cl_2 (3 x 30 mL). The combined organic layers were washed with brine (30 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 4:1) to afford the title compound (180 mg, 33%) as a yellow oil. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.28 (s, 1H), 8.40 (s, 1H), 8.32 (s, 1H), 5.71-5.67 (m, 1H), 4.60 (t, J = 4.9 Hz, 1H), 3.54-3.39 (m, 2H), 2.46 (s, 3H), 2.28-2.00 (m, 2H), 1.20 (s, 9H). LC-MS (Method K): m/z = 268.3 $[\text{M}+1]^+$, 0.75 min.

INTERMEDIATE 75

PREPARATION OF 5-FORMYLPYRIDINE-3-CARBONITRILE

A mixture of 5-bromopyridine-3-carbaldehyde (10 g, 53.8 mmol) and cuprous cyanide (7.20 g, 80.6 mmol) in N,N-dimethylformamide (40 mL) was heated to 140 °C and stirred under a nitrogen atmosphere overnight. After being cooled to room temperature, the reaction mixture was diluted with water (80 mL) and filtered. The filtrate was extracted with EtOAc (3 x 50 mL). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:4) to afford the title compound (1.86 g, 26%) as a yellow solid. ^1H NMR (400 MHz, CDCl_3) δ 10.17 (s, 1H), 9.30 (s, 1H), 9.14 (s, 1H), 8.45 (t, J = 2.0 Hz, 1H).

INTERMEDIATE 76

PREPARATION OF 5-[(1E)-3-OXOPROP-1-EN-1-YL]PYRIDINE-3-CARBONITRILE

A mixture of 5-formylpyridine-3-carbonitrile (1.82 g, 13.78 mmol) and (formylmethylene)triphenylphosphorane (4.28 g, 14.06 mmol) in DMSO (15 mL) was stirred overnight at 120 °C under a nitrogen atmosphere. After being cooled to room temperature, the reaction mixture was diluted with water (30 mL) and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:3) to afford the title compound (490 mg, 22%) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ 9.79 (d, J = 7.2 Hz, 1H), 8.99 (d, J = 2.1 Hz, 1H), 8.92 (d, J = 2.1 Hz, 1H), 8.14 (m, 1H), 7.52 (d, J = 15.6 Hz, 1H), 6.87-6.79 (m, 1H).

INTERMEDIATE 77

PREPARATION OF TERT-BUTYL N-[(1S)-1-(5-CYANOPYRIDIN-3-YL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

To a solution of (S)-2-(diphenyl(trimethylsilyloxy)methyl)pyrrolidine (280 mg, 0.86 mmol) in CHCl_3 (20 mL) were added 5-[(1E)-3-oxoprop-1-en-1-yl]pyridine-3-carbonitrile (450 mg, 2.85 mmol) and tert-butyl N-hydroxycarbamate (570 mg, 4.28 mmol). The reaction mixture was stirred overnight at room temperature. MeOH (10 mL) was added to the reaction solution, then NaBFL (220 mg, 5.82 mmol) was added at 0 °C. After being stirred at room temperature for 30 min, the reaction mixture was quenched by the addition of sat. NH_4Cl solution (30 mL) and extracted with CH_2Cl_2 (3 x 20 mL). The combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 3:1) to afford the title compound (280 mg, 30%) as a yellow solid. LC-MS (Method K): $m/z = 294.3$ $[\text{M}+\text{H}]^+$, 0.67 min.

INTERMEDIATE 78

PREPARATION OF TERT-BUTYL (3S)-3-(5-CYANOPYRIDIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a stirred solution of tert-butyl N-[(1S)-1-(5-cyanopyridin-3-yl)-3-hydroxypropyl]-N-hydroxycarbamate (280 mg, 0.95 mmol) and triphenylphosphine (300 mg, 1.14 mmol) in THF (5 mL) was added dropwise DIAD (193 mg, 0.95 mmol) at 0 °C. The reaction mixture was stirred overnight at room temperature under a nitrogen atmosphere. Upon concentration under reduced pressure, the crude product was purified by preparative thin layer chromatography (EtOAc-petroleum ether, 2:3) to afford the title compound (130 mg, 49%) as a yellow solid. LC-MS (Method K): $m/z = 276.3$ $[\text{M}+\text{H}]^+$, 0.83 min.

The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 60/40 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.82 (d, $J=2.0$ Hz, 2H), 8.02 (t, $J=2.0$ Hz, 1H), 5.34 (dd, $J=8.8, 5.5$ Hz, 1H), 4.25 (dt, $J=3.4, 8.0$ Hz, 1H), 3.93 (dt, $J=7.0, 8.8$ Hz, 1H), 2.90 (dddd, $J=12.3, 8.9, 7.2, 3.3$ Hz, 1H), 2.35-2.24 (m, 1H), 1.52 (s, 9H). LC-MS (Method A): $m/z = 276.2$ $[\text{M}+\text{H}]^+$, 0.85 min. **e.e.=100%** as determined on a Chiralpak IC (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/ethanol 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 8.9 min.

INTERMEDIATE 79

PREPARATION OF 5-[(3S)-1,2-OXAZOLIDIN-3-YL]PYRIDINE-3-CARBONITRILE TRIFLUOROACETATE

Trifluoroacetic acid (5 mL) was added to a solution of tert-butyl (3S)-3-(5-cyanopyridin-3-yl)-1,2-oxazolidine-2-carboxylate (100 mg, 0.36 mmol) in CH_2Cl_2 (10 mL). After being stirred at room temperature for 1 hour, the mixture was concentrated under reduced pressure to afford the title compound (105 mg, crude) as a yellow solid. LC-MS (Method C): $m/z = 176.1$ $[\text{M}+\text{H}]^+$, 0.76 min.

INTERMEDIATE 80

PREPARATION OF 3-FLUORO-5-[(1E)-3-OXOPROP-1-EN-1-YL]BENZONITRILE

A mixture of 3-fluoro-5-formylbenzonitrile (1 g, 6.71 mmol) and (formylmethylene)triphenylphosphorane (2.04 g, 6.70 mmol) in THF (20 mL) was stirred overnight at 75

°C under a nitrogen atmosphere. After being cooled to room temperature, the reaction mixture was diluted with water (30 mL) and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over Na⁺SO₄ and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:5) to afford the title compound (830 mg, 71%) as a yellow solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 9.69 (d, *J* = 7.6 Hz, 1H), 8.15 (s, 1H), 8.08–8.04 (m, 1H), 7.98–7.94 (m, 1H), 7.72 (d, *J* = 16.0 Hz, 1H), 7.09–7.02 (m, 1H).

INTERMEDIATE 81

PREPARATION OF N-[(1S)-1-(3-CYANO-5-FLUOROPHENYL)-3-HYDROXYPROPYL]-N-HYDROXY-2,2-DIMETHYLPROPANAMIDE

To a solution of (S)-2-(diphenyl(trimethylsilyloxy)methyl)pyrro3idine (110 mg, 0.34 mmol) in CHCl₃ (10 mL) were added 3-fluoro-5-[(1E)-3-oxoprop-1-en-1-yl]benzonitrile (200 mg, 1.14 mmol) and N-hydroxypivalamide (200 mg, 1.71 mmol). The reaction mixture was stirred overnight at room temperature. MeOH (10 mL) was added to the mixture, then NaBH₄ (90 mg, 2.38 mmol) was added at 0 °C. After being stirred at room temperature for 30 min, the reaction mixture was quenched by the addition of sat. NH₄Cl solution (20 mL) and extracted with CH₂Cl₂ (3 x 15 mL). The combined organic layers were dried over Na⁺SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc-petroleum ether, 1:2) to afford the title compound (100 mg, 29%) as a yellow solid. LC-MS (Method G): *m/z* = 294.9 [M+Hf, 0.89 min.

INTERMEDIATE 82

PREPARATION OF TERT-BUTYL (3S)-3-(5-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

The title compound was isolated by chiral HPLC on a Chiraipak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50% v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.65 (d, *J* = 1.5 Hz, 1H), 8.40 (d, *J* = 0.8 Hz, 1H), 5.36 (dd, *J* = 8.9, 5.1 Hz, 1H), 4.24–4.16 (m, 1H), 4.00–3.91 (m, 1H), 2.78 (dddd, *J* = 12.3, 8.6, 7.5, 4.1 Hz, 1H), 2.67–2.56 (m, 4H), 1.51 (s, 9H). LC-MS (Method A): *m/z* = 266.1 [M+Hf, 0.76 min. e.e. = 100% as determined on a Chiraipak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50% v/v, flow rate: 1 mL/min, retention time: 23.8 min.

INTERMEDIATE 83

PREPARATION OF (2E)-3-[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]PROP-2-ENAL

A mixture of 5-bromo-2-(difluoromethyl)pyridine (1770 mg, 8.5 mmol), trans-di(p-acetato)bis[o-(di-o-toiyi-phosphino)benzyl]dipalladium(II) (1600 mg, 1.7 mmol) and sodium acetate (797 mg, 9.35 mmol) was introduced into a microwave reaction tube under nitrogen. The tube was dried under vacuum, then acetonitrile (10 mL) and 2-propenal (4.66 mL, 30.6 mmol) were added consecutively to the reaction mixture. The mixture was heated in a microwave oven to 140°C (7 cycles of 15 min each). After cooling, the solid was filtered and washed with MeOH, and the filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane/EtOAc, 100:0 to 0:100) to afford the title compound (825 mg, 47%) as an orange oil. ¹H

NMR (400MHz, CDCl_3) δ 9.78 (d, $J=7.5$ Hz, 1H), 8.83 (d, $J=1.0$ Hz, 1H), 8.05 (dd, $J=8.2, 2.1$ Hz, 1H), 7.74 (d, $J=8.3$ Hz, 1H), 7.53 (d, $J=6.3$ Hz, 1H), 6.89-6.52 (m, 1H). LC-MS (Method A): $m/z = 184.1$ $[\text{M}+\text{H}]^+$, 0.69 min.

INTERMEDIATE 84

PREPARATION OF 2-METHYL-5-[(1E)-3-OXOPROP-1-EN-1-YL]PYRIDINE-3-CARBONITRILE

A solution of 5-bromo-2-methylnicotinonitrile (1000 mg, 5.08 mmol) and 3,3-diethoxyprop-1-ene (2.78 mL, 18.27 mmol) in anhydrous MeCN (15 mL) was degassed by applying alternatively vacuum and nitrogen for 3 cycles. Palladium(II) acetate (342 mg, 1.5 mmol), tri(o-tolyl)phosphine (541 mg, 1.78 mmol) and Et_3N (1.9 mL, 13.7 mmol) were then added and the mixture was heated in a microwave oven to 100°C (4 cycles of 15 min each). After cooling, the solid was filtered and washed with methanol, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography (cyclohexane/EtOAc, 100:0 to 0:100) to afford the intermediate 5-[(1E)-3,3-diethoxyprop-1-en-1-yl]-2-methylpyridine-3-carbonitrile (360 mg) as an orange oil. This intermediate was dissolved in THF (10 mL) and treated with hydrochloric acid solution (10 mL, 10% in water) at room temperature. After 4 h, the solution was basified with sat. NaHCO_3 solution and extracted twice with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 , and concentrated under reduced pressure to afford the title compound (320 mg, 35%) as a yellow solid. ^1H NMR (400MHz, CDCl_3) δ 9.77 (d, $J=7.3$ Hz, 1H), 8.86 (d, $J=2.3$ Hz, 1H), 8.08 (d, $J=2.3$ Hz, 1H), 7.46 (d, $J=16.3$ Hz, 1H), 6.80 (dd, $J=16.2, 7.4$ Hz, 1H), 2.85 (s, 3H). LC-MS (Method A): $m/z = 173.1$ $[\text{M}+\text{H}]^+$, 0.64 min.

INTERMEDIATE 85

PREPARATION OF 3-BROMO-5-FORMYLBENZONITRILE

Isopropylmagnesium chloride solution (2M in THF, 20.9 mL, 41.88 mmol) was added at 0°C to a stirred solution of 3,5-dibromobenzonitrile (9.036 g, 34.9 mmol) in anhydrous THF (36 mL). After 30min, anhydrous DMF (8.14 mL, 104.7 mmol) was added at 0°C and the mixture was stirred at this temperature for 1h then water (200 mL) and diethyl ether (150 mL) were added to the mixture. The mixture was stirred and acidified to pH 2-3 with 6N aqueous HCl solution. The clear biphasic mixture thus obtained was separated and the aqueous phase was extracted twice again with diethyl ether. The combined organic phases were washed twice with brine, dried over Na_2SO_4 and evaporated under reduced pressure to give a yellow solid (7.39 g). This crude product was pooled with the crude product (0.73 g) isolated from a similar reaction run on 1.0 g of 3,5-dibromobenzonitrile and together they were purified by column chromatography (cyclohexane/EtOAc, 100:0 to 30:70) to afford the title compound (6.41 g, 79%). ^1H NMR (400MHz, CDCl_3) δ 10.01 (s, 1H), 8.25 (t, $J=1.5$ Hz, 1H), 8.11 (t, $J=1.4$ Hz, 1H), 8.05 (t, $J=2.0$ Hz, 1H).

INTERMEDIATE 86

PREPARATION OF (2E)-3-(6-NITROPYRIDIN-3-YL)PROP-2-ENAL

A mixture of 5-bromo-2-nitropyridine (1.92 g, 9.53 mmol), 3,3-diethoxy-1-propenylboronic acid pinacol ester (2.93 g, 11.43 mmol) and cesium carbonate (6.21 g, 19.06 mmol) in 1,4-dioxane (28 mL) was degassed by alternatively applying vacuum and nitrogen for 3 cycles. Water (4 mL) and $[\text{I}, \text{I}'-$

bis(diphenylphosphino)ferrocene]dichloropalladium(II) (0.695 g, 0.95 mmol) were then added and the mixture was stirred at 100 °C for 4h. The mixture was filtered over a pad of Celite, diluted with EtOAc and washed twice with water. The organic phase was combined with the organic phase from a second reaction run in parallel on the same scale and using the same conditions. The pooled organic phases were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford a brown oil (7.23 g). This material was dissolved in THF (75 mL) and treated with 1M aqueous HCl solution (75 mL). The mixture was stirred at room temperature for 1h then extracted three times with EtOAc. The combined organic portions were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford a brown solid (5.2 g). The crude product was purified by column chromatography (cyclohexane/EtOAc, 80:20 to 0:100) to give the title compound (2.28 g, 67%) as an orange solid. ¹H NMR (400MHz, CDCl₃) δ 9.84 (d, J=7.0 Hz, 1H), 8.83 (d, J=2.3 Hz, 1H), 8.36 (d, J=8.5 Hz, 1H), 8.23 (dd, J=8.4, 2.1 Hz, 1H), 7.59 (d, J=16.3 Hz, 1H), 6.91 (dd, J=16.2, 7.2 Hz, 1H). LC-MS (Method A): m/z = 179.1 [M+H]⁺, 0.60 min.

INTERMEDIATE 87

PREPARATION OF TERT-BUTYL 3-(6-METHYLPYRIDIN-3-YL)-5-OXO-1,2-OXAZOLIDINE-2-CARBOXYLASE

To a stirred solution of 6-methylpyridine-3-carbaldehyde (1 g, 8.26 mmol) in EtOAc (20 mL) were added 2,2-dimethyl-1,3-dioxane-4,6-dione (1.19 g, 8.26 mmol), 1,4-diazabicyclo[2.2.2]octane (93 mg, 0.83 mmol) and tert-butyl N-hydroxy carbamate (1.10 g, 8.26 mmol). After being stirred at room temperature for 16 h, the reaction mixture was quenched with water (50 mL), and extracted with EtOAc (3 x 30 mL). The combined organic layers were washed with brine (30 mL), dried over Na₂SO₄ and concentrated under reduced pressure to afford the title compound (2.10 g, crude) as a yellow oil. LC-MS (Method G): m/z = 279.3 [M+H]⁺, 0.843 min.

INTERMEDIATE 88

PREPARATION OF TERT-BUTYL N-HYDROXY-N-[3-HYDROXY-1-(6-METHYLPYRIDIN-3-YL)PROPYL] CARBAMATE

To a stirred solution of tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate (2.10 g, 7.55 mmol) in THF (30 mL) was added lithium borohydride (493 mg, 22.6 mmol) in portions. After being stirred at room temperature for 5 min, the reaction mixture was quenched by the addition of water (50 mL), and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄ and concentrated under reduced pressure to afford the title compound (2.10 g, crude) as a yellow oil. LC-MS (Method G): m/z = 283.3 [M+H]⁺, 0.795 min.

INTERMEDIATE 89

PREPARATION OF TERT-BUTYL 3-(6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

To a stirred solution of tert-butyl N-hydroxy-N-[3-hydroxy-1-(6-methylpyridin-3-yl)propyl]carbamate (2.10 g, 7.44 mmol) and triphenylphosphine (2.93 g, 11.2 mmol) in THF (30 mL) under a nitrogen atmosphere was added dropwise DIAD (2.26 g, 11.2 mmol) at 5 °C. After being stirred at 5 °C for 10 min, the mixture was concentrated under reduced pressure to afford the title compound (7.50 g, crude) as a brown oil. LC-MS (Method G): m/z = 265.3 [M+H]⁺, 0.967 min.

INTERMEDIATE 90

PREPARATION OF 2-METHYL-5-(1,2-OXAZOLIDIN-3-YL)PYRIDINE

A mixture of tert-butyl 3-(6-methylpyridin-3-yl)-1,2-oxazolidine-2-carboxylate (7.50 g, crude) and 4 M hydrogen chloride in dioxane (50 mL) was stirred at room temperature for 30 min. The solvent was removed under reduced pressure. The residue was diluted with water (30 mL) and washed with CH₂Cl₂ (4 x 30 mL). The aqueous layer was basified with sat. Na₂CO₃ solution and extracted with CH₂Cl₂ (4 x 30 mL). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to afford the title compound (1.20 g, crude) as a yellow oil. LC-MS (Method G): m/z = 165.3 [M+H]⁺, 0.464 min.

INTERMEDIATE 91

PREPARATION OF TERT-BUTYL 3-(5-CYANTHIOPHEN-2-YL)-5-OXO-1,2-OXAZOLIDINE-2-CARBOXYLATE

This compound was synthesized by the same method as described for tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate except 5-formylthiophene-2-carbonitrile (1 g, 7.30 mmol) was used instead of 6-methylpyridine-3-carbaldehyde to afford the title compound (1.6 g, 74%). LC-MS (Method C): m/z = 295.1 [M+H]⁺, 0.793 min.

INTERMEDIATE 92

PREPARATION OF TERT-BUTYL N-[1-(5-CYANTHIOPHEN-2-YL)-3-HYDROXYPROPYL]-N-HYDROXYCARBAMATE

This compound was synthesized by the same method as described for tert-butyl N-hydroxy-N-[3-hydroxy-1-(6-methylpyridin-3-yl)pyrrolidine-2-carboxylate except tert-butyl 3-(5-cyanothiophen-2-yl)-5-oxo-1,2-oxazolidine-2-carboxylate (1.4 g, 4.76 mmol) was used instead of tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate to afford the title compound (900 mg, 63%). LC-MS (Method C): m/z = 299.1 [M+H]⁺, 0.730 min.

INTERMEDIATE 93

PREPARATION OF TERT-BUTYL 3-(5-CYANTHIOPHEN-2-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE

This compound was synthesized by the same method as described for tert-butyl 3-(6-methylpyridin-3-yl)-1,2-oxazolidine-2-carboxylate except tert-butyl N-[1-(5-cyanothiophen-2-yl)-3-hydroxypropyl]-N-hydroxycarbamate (900 mg, 3.02 mmol) was used instead of tert-butyl N-hydroxy-N-[3-hydroxy-1-(6-methylpyridin-3-yl)pyrrolidine-2-carboxylate to afford the title compound (2.00 g, crude). LC-MS (Method Q): m/z = 281.1 [M+H]⁺, 0.636 min.

INTERMEDIATE 94

PREPARATION OF 5-(1,2-OXAZOLIDIN-3-YL)THIOPHENE-2-CARBONITRILE

This compound was synthesized by the same method as described for 2-methyl-5-(1,2-oxazolidin-3-yl)pyridine except tert-butyl 3-(5-cyanothiophen-2-yl)-1,2-oxazolidine-2-carboxylate (500 mg, crude) was used instead of tert-butyl 3-(6-methylpyridin-3-yl)-1,2-oxazolidine-2-carboxylate to afford the title compound (50 mg). LC-MS (Method Q): m/z = 181.1 [M+H]⁺, 0.536 min.

INTERMEDIATE 95

PREPARATION OF TERT-BUTYL 3-(5-CYANO-1-METHYL-1H-PYRROL-3-YL)-5-OXO-1,2-
OXAZOLIDINE-2-CARBOXYLATE

This compound was synthesized by the same method as described for tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate except 4-formyl-1-methyl-1H-pyrrole-2-carbonitrile (1 g, 7.46 mmol) was used instead of 6-methylpyridine-3-carbaldehyde to afford the title compound (1.4 g, 64%). LC-MS (Method M): $m/z = 192.2$ $[\text{M}-\text{Boc}+\text{H}]^+$, 0.973 min.

INTERMEDIATE 96

PREPARATION OF TERT-BUTYL N-[1-(5-CYANO-1-METHYL-1H-PYRROL-3-YL)-3-HYDROXYPROPYL]-
N-HYDROXYCARBAMATE

This compound was synthesized by the same method as described for tert-butyl N-hydroxy-N-(3-hydroxy-1-(6-methylpyridin-3-yl)propyl)carbamate except tert-butyl 3-(5-cyano-1-methyl-1H-pyrrol-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate (1.4 g, 4.81 mmol) was used instead of tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate to afford the title compound (1.15 g, 80%). LC-MS (Method G): $m/z = 196.2$ $[\text{M}-\text{Boc}+\text{H}]^+$, 0.727 min.

INTERMEDIATE 97

PREPARATION OF TERT-BUTYL 3-(5-CYANO-1-METHYL-1H-PYRROL-3-YL)-1,2-OXAZOLIDINE-2-
CARBOXYLATE

This compound was synthesized by the same method as described for tert-butyl 3-(6-methylpyridin-3-yl)-5-oxo-1,2-oxazolidine-2-carboxylate except tert-butyl N-[1-(5-cyano-1-methyl-1H-pyrrol-3-yl)-3-hydroxypropyl]-N-hydroxycarbamate (500 mg, 1.69 mmol) was used instead of 6-methylpyridine-3-carbaldehyde to afford the title compound (620 mg, crude). LC-MS (Method G): $m/z = 178.3$ $[\text{M}-\text{Boc}+\text{H}]^+$, 0.922 min.


INTERMEDIATE 98

PREPARATION OF 1-METHYL-4-(1,2-OXAZOLIDIN-3-YL)-1H-PYRROLE-2-CARBONITRILE

This compound was synthesized by the same method as described for 2-methyl-5-(1,2-oxazolidin-3-yl)pyridine except tert-butyl 3-(5-cyano-1-methyl-1H-pyrrol-3-yl)-1,2-oxazolidine-2-carboxylate (620 mg, crude) was used instead of tert-butyl 3-(6-methylpyridin-3-yl)-1,2-oxazolidine-2-carboxylate to afford the title compound (150 mg, crude). LC-MS (method g): $m/z = 178.2$ $[\text{m}+\text{h}]^+$, 0.488 min.

EXAMPLE 1

PREPARATION OF 1-METHYL-5-[(3S)-3-PHENYL-1,2-OXAZOLIDINE-2-CARBONYL]-1H-PYRROLE-2-CARBONITRILE (1)




5 N,N-Diisopropylethylamine (233 μ L, 1.34 mmol), 1-hydroxybenzotriazole (91 mg, 0.67 mmol) and EDC-HCl (129 mg, 0.67 mmol) were added to a stirred solution of 5-cyano-1-methyl-1H-pyrrole-2-carboxylic acid (50 mg, 0.33 mmol) and 3-phenyl-1,2-oxazolidine (50 mg, 0.33 mmol) in THF (2 mL) and the resulting mixture was stirred at room temperature for 18 h. The reaction mixture was diluted with EtOAc and washed twice with sat. NaHCO_3 solution. The organic portion was dried over Na_2SO_4 ,
 10 filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water~ CH_3CN , 100:0 to 20:80) to afford an 84:16 mixture of enantiomers (80.3 mg, 86%) as a white solid. This mixture was resolved by chiral HPLC on a Whelk O-1 (R,R) (25 x 2.0 cm), 10 μ m column using a mobile phase of n-hexane/iEtOH/ $\text{CF}_3\text{CO}_2\text{H}$ 9/1 + 0.1% isopropylamine) 10/90 % v/v and a flow rate of 20 mL/min to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.41-7.29 (m, 5H), 6.98 (d, $J=4.3$ Hz, 1H), 6.77 (d, $J=4.5$ Hz, 1H), 5.60 (dd, $J=8.7, 6.2$ Hz, 1H), 4.34 (td, $J=7.8, 3.9$ Hz, 1H), 4.06-3.96 (m, 4H), 2.96-2.84 (m, 1H), 2.49-2.39 (m, 1H). LC-MS (Method A): m/z = 282.2 $[\text{M}+\text{H}]^+$, 1.03 min. e.e.=100% as determined on a Whelk O-1 (R,R) (25 x 0.46 cm), 10 μ m column using a mobile phase of n-hexane/(EtOH + 0.1% isopropylamine) 20/80 % v/v, flow rate: 1 mL/min, retention time: 6.4 min.

20

EXAMPLE 2

PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (2)




Pivaloyl chloride (53 μ L, 0.433 mmol) was added to a solution of 3-phenyl-1,2-oxazolidine (40 mg, 0.268 mmol) and pyridine (54 μ L, 0.670 mmol) in CH_2Cl_2 (2.5 mL). The reaction mixture was
 25 stirred at room temperature for 4 h. Volatiles were removed under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3CN , 100:0 to 20:80) to afford an 85:15 mixture of enantiomers (45 mg, 72%) as a colorless oil. This mixture was resolved by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/EtOH 70/30 % v/v to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.38-7.31 (m, 2H), 7.29-7.21 (m, 3H), 5.31 (dd,

$J=8.7, 6.9$ Hz, 1H), 4.28 (td, $J=7.7, 2.3$ Hz, 1H), 3.84 (ddd, $J=10.0, 8.0, 6.4$ Hz, 1H), 2.87 (dddd, $J=12.0, 9.0, 6.4, 2.8$ Hz, 1H), 2.19-2.07 (ro, 1H), 1.22 (s, 9H). LC-MS (Method A): $m/z = 234.3$ $[M+H]^+$, 1.08 min. e.e.=97.6% as determined on a Chiralpak 1C (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/EtOH 70/30 % v/v.

5

EXAMPLE 3

PREPARATION OF 5-[3-(2-CHLORO-6-FLUOROPHENYL)-1,2-OXAZOLIDINE-2-CARBONYL]-1-METHYL-1H-PYRROLE-2-CARBONITRILE (3)




N,N-Diisopropylethylamine (18 μ L, 0.10 mmol), 1-hydroxybenzotriazole (6.7 mg, 0.05 mmol) and EDC-HCl (9.6 mg, 0.05 mmol) were added to a stirred solution of 5-cyano-1-methyl-1H-pyrrole-2-carboxylic acid (3.7 mg, 0.025 mmol) and 3-(2-chloro-6-fluorophenyl)-1,2-oxazolidine (5.0 mg, 0.025 mmol) in THF (2 mL) and the resulting mixture was stirred at room temperature for 72 h. The reaction mixture was diluted with EtOAc and washed twice with sat. NaHCO_3 solution. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford an impure fraction of the title compound. This material was further purified by reverse phase chromatography (water- CH_3CN , 100:0 to 30:70) to afford the title compound as a mixture of enantiomers. ^1H NMR (400 MHz, CDCl_3) δ 7.24-7.20 (m, 2H), 7.05-6.96 (m, 2H), 6.75 (d, $J=4.3$ Hz, 1H), 6.01 (t, $J=8.7$ Hz, 1H), 4.51-4.43 (m, 1H), 3.97-4.05 (m, 4H), 2.93-2.81 (m, 1H), 2.66-2.53 (m, 1H). LC-MS (Method A): $m/z = 334.2$ $[M+H]^+$, 1.09 min.

20

EXAMPLE 4

PREPARATION OF (3S)-N,3-DIPHENYL-1,2-OXAZOLIDINE-2-CARBOXAMIDE (4)




A solution of (3S)-3-phenyl-1,2-oxazolidine (25 mg, 0.16 mmol) in CH_2Cl_2 (0.5 mL) was added to a solution of phenyl isocyanate (18 μL , 0.16 mmol) in CH_2Cl_2 (0.5 mL) which had been pre-cooled to 0 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 1 h.

Volatiles were removed under reduced pressure. The crude product was purified by column

chromatography (cyclohexane-EtOAc, 95:5 to 70:30) to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.36 (s, 1H), 7.66-7.62 (m, 2H), 7.42-7.34 (m, 4H), 7.31-7.25 (m, 3H), 7.05-7.00 (m, 1H), 5.43 (dd, $J=8.5, 5.8$ Hz, 1H), 4.24 (td, $J=8.0, 3.1$ Hz, 1H), 3.92-3.85 (m, 1H), 2.85 (dddd, $J=12.1, 8.7, 7.1, 3.3$ Hz, 1H), 2.23 (dddd, $J=12.0, 9.4, 7.9, 5.8$ Hz, 1H). LC-MS (Method A): m/z =269.2 $[\text{M}+\text{H}]^+$, 1.07 min.

EXAMPLE 5

PREPARATION OF (3S)-N-tert-BUTYL-3-PHENYL-1,2-OXAZOLIDINE-2-CARBOXAMIDE (5)




A solution of (3S)-3-phenyl-1,2-oxazolidine (25 mg, 0.16 mmol) in CH_2Cl_2 (0.5 mL) was added to a solution of tert-Butyl isocyanate (18 μL , 0.16 mmol) in CH_2Cl_2 (0.5 mL) which had been pre-cooled to 0 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred overnight.

Volatiles were removed under reduced pressure. The crude product was purified by column

chromatography (cyclohexane-EtOAc, 95:5 to 70:30) to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.35-7.29 (m, 4H), 7.28-7.21 (m, 1H), 6.49 (br. s, 1H), 5.30 (dd, $J=8.5, 5.5$ Hz, 1H), 4.14-4.08 (m, 1H), 3.77-3.70 (m, 1H), 2.78-2.67 (m, 1H), 2.14 (dddd, $J=12.0, 9.3, 7.8, 5.5$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): m/z =249.3 $[\text{M}+\text{H}]^+$, 1.05 min.

EXAMPLE 6

PREPARATION OF (3S)-N,N-DIMETHYL-3-PHENYL-1,2-OXAZOLIDINE-2-CARBOXAMIDE (6)




4-(Dimethylamino)pyridine (2 mg, 0.016 mmol) was added to a stirred solution of (3S)-3-phenyl-1,2-oxazolidine (50 mg, 0.33 mmol), dimethylcarbamyl chloride (46 μL , 0.50 mmol) and pyridine (180 μL) in CH_2Cl_2 (1.0 mL) and the resulting mixture was stirred at room temperature for 18 h. The reaction mixture was diluted with CH_2Cl_2 and washed twice with sat. NH_4Cl solution. The organic

portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 95:5 to 50:50) to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.40-7.31 (m, 4H), 7.28-7.22 (m, 1H), 5.25 (dd, $J=8.7, 4.6$ Hz, 1H), 4.05 (td, $J=7.9, 4.3$ Hz, 1H), 3.83 (q, $J=8.0$ Hz, 1H), 2.89 (s, 6H), 2.71 (dtd, $J=12.4, 8.1, 4.0$ Hz, 1H), 2.20-2.10 (m, 1H). LC-MS (Method A): $m/z = 221.2$ $[\text{M}+\text{H}]^+$, 0.85 min.

EXAMPLE 7


PREPARATION OF (3S)-3-PHENYL-2-(PYRROLIDINE-1-CARBONYL)-1,2-OXAZOLIDINE (7)



4-(Dimethylamino)pyridine (2 mg, 0.016 mmol) was added to a stirred solution of (3S)-3-phenyl-1,2-oxazolidine (50 mg, 0.33 mmol), 1-pyrrolidinecarbonyl chloride (55 μL , 0.50 mmol) and pyridine (180 μL) in CH_2Cl_2 (1.0 mL) and the resulting mixture was stirred at room temperature for 18 h. The reaction mixture was diluted with 3:1 and washed twice with sat. NH_4Cl solution. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 95:5 to 50:50) to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.40-7.30 (m, 4H), 7.27-7.21 (m, 1H), 5.38 (dd, $J=8.7, 4.9$ Hz, 1H), 4.07 (td, $J=7.8, 4.0$ Hz, 1H), 3.76 (q, $J=7.8$ Hz, 1H), 3.38 (t, $J=5.6$ Hz, 4H), 2.77-2.67 (m, 1H), 2.15 (dtd, $J=12.1, 8.3, 4.9$ Hz, 1H), 1.85-1.69 (m, 4H). LC-MS (Method A): $m/z = 247.3$ $[\text{M}+\text{H}]^+$, 0.94 min.

EXAMPLE 8


PREPARATION OF (3S)-2-(3-METHYLOXETANE-3-CARBONYL)-3-PHENYL-1,2-OXAZOLIDINE (8)



3-Methyloxetane-3-carboxylic acid (30 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.39-7.33 (m, 2H), 7.31-7.25 (m, 3H), 5.32 (dd, $J=8.5, 6.5$ Hz, 1H), 4.86 (d, $J=6.3$ Hz, 1H), 4.74 (td, $J=6.0$ Hz, 1H), 4.29-4.20 (m, 3H), 3.97-3.88 (m, 1H), 2.92 (ddt, $J=15.3, 8.9, 3.2$ Hz, 1H), 2.24-2.13 (m, 1H), 1.60 (s, 3H). LC-MS (Method A): $m/z = 248.1$ $[\text{M}+\text{H}]^+$, 0.81 min.

EXAMPLE 9


PREPARATION OF 2-HYDROXY-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]ETHAN-1-ONE (9)



Glycolic acid (19 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.38-7.33 (m, 2H), 7.32-7.25 (m, 3H), 5.32 (dd, $J=8.7, 6.1$ Hz, 1H), 4.82 (t, $J=6.3$ Hz, 1H), 4.29-4.13 (m, 3H), 3.86 (ddd, $J=9.5, 8.0, 6.7$ Hz, 1H), 2.92-2.84 (m, 1H), 2.18 (dddd, $J=12.2, 9.5, 7.5, 6.1$ Hz, 1H). LC-MS (Method A): $m/z = 208.0$ $[\text{M}+\text{H}]^+$, 0.65 min.

EXAMPLE 10


PREPARATION OF 1-[(3S)-3-PHENYL-1,2-OXAZOLIDINE-2-CARBONYL]CYCLOPROPAN-1-OL (10)



1-Hydroxycyclopropane-1-carboxylic acid (26 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.38-7.31 (m, 4H), 7.29-7.23 (m, 1H), 6.01 (s, 1H), 5.59 (dd, $J=8.8, 4.8$ Hz, 1H), 4.15 (td, $J=7.7, 3.9$ Hz, 1H), 3.86 (q, $J=7.8$ Hz, 1H), 2.86-2.78 (m, 1H), 2.28-2.19 (m, 1H), 1.09-0.97 (m, 2H), 0.84 (dd, $J=4.8, 1.8$ Hz, 1H), 0.78 (dd, $J=5.3, 2.0$ Hz, 1H). LC-MS (Method A): $m/z = 234.3$ $[\text{M}+\text{H}]^+$, 0.71 min.

EXAMPLE 11

PREPARATION OF (3S)-2-(OXETANE-3-CARBONYL)-3-PHENYL-1,2-OXAZOLIDINE (11)




Oxetane-3-carboxylic acid (26 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR

(400 MHz, DMSO- d_6) δ 7.39-7.24 (m, 5H), 5.33 (dd, $J=8.4, 6.7$ Hz, 1H), 4.76-4.63 (m, 4H), 4.19 (td, $J=7.7, 3.1$ Hz, 2H), 3.88-3.80 (m, 1H), 2.89 (dddd, $J=12.2, 8.8, 6.6, 3.3$ Hz, 1H), 2.18 (dddd, $J=12.1, 9.5, 7.5, 6.3$ Hz, 1H). LC-MS (Method A): $m/z = 234.0$ $[M+H]^+$, 0.73 min.

EXAMPLE 12


PREPARATION OF (3S)-2-CYCLOPROPANECARBONYL-3-PHENYL-1,2-OXAZOLIDINE (12)



Cyclopropanecarboxylic acid (20 μ L, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.39-7.23 (m, 5H), 5.35 (dd, $J=8.7, 6.1$ Hz, 1H), 4.28 (td, $J=7.7, 3.1$ Hz, 1H), 3.86 (ddd, $J=9.3, 8.0, 6.8$ Hz, 1H), 2.88 (dddd, $J=12.1, 8.8, 6.7, 3.3$ Hz, 1H), 2.21 (dddd, $J=12.1, 9.4, 7.6, 6.0$ Hz, 2H), 0.89-0.73 (m, 4H). LC-MS (Method A): $m/z = 218.4$ $[M+H]^+$, 0.90 min.

EXAMPLE 13


PREPARATION OF 2-METHOXY-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]ETHAN-1-ONE (13)



2-Methoxyacetic acid (19 μ L, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.39-7.24 (m, 5H), 5.31 (dd, $J=8.7, 6.4$ Hz, 1H), 4.28-4.18 (m, 3H), 3.88 (ddd, $J=9.6, 8.0, 6.5$ Hz, 1H), 3.30 (s, 3H), 2.88 (dddd, $J=12.1, 9.1, 6.5, 3.0$ Hz, 1H), 2.17 (dddd, $J=12.1, 9.6, 7.5, 6.4$ Hz, 1H). LC-MS (Method A): $m/z = 222.3$ $[M+H]^+$, 0.77 min.

EXAMPLE 14


PREPARATION OF (2R)-2-HYDROXY-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (14)



D-(-)-Lactic acid (23 mg, 0.26 mmol) was reacted with (3S)-3-phenyl- 1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.39-7.24 (m, 5H), 5.36-5.27 (m, 1H), 5.04-4.97 (m, 1H), 4.53-4.42 (m, 1H), 4.30-4.21 (m, 1H), 3.88 (ddd, *J*=9.2, 8.0, 6.7 Hz, 1H), 2.88 (dddd, *J*=12.2, 8.8, 6.7, 3.4 Hz, 1H), 2.21 (dddd, *J*=12.2, 9.2, 7.5, 6.1 Hz, 1H), 1.12-1.22 (m, 3H). LC-MS (Method A): *m/z* = 222.0 [*M*+*H*]⁺, 0.68 min.

EXAMPLE 15


PREPARATION OF (2S)-2-HYDROXY-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (15)



L-(+)-Lactic acid (23 mg, 0.26 mmol) was reacted with (3S)-3-phenyl- 1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.40-7.23 (m, 5H), 5.34 (dd, *J*=6.4, 8.9 Hz, 1H), 4.84 (d, *J*=7.0 Hz, 1H), 4.54-4.44 (m, 1H), 4.25 (dt, *J*=2.6, 7.8 Hz, 1H), 3.86 (ddd, *J*=6.8, 8.0, 9.5 Hz, 1H), 2.93-2.85 (m, 1H), 2.18 (dddd, *J*=6.3, 7.6, 9.6, 12.2 Hz, 1H), 1.30 (d, *J*=6.8 Hz, 3H). LC-MS (Method A): *m/z* = 222.2 [*M*+*H*]⁺, 0.73 min.

EXAMPLE 16

PREPARATION OF 1-[(3S)-3-PHENYL-1,2-OXAZOLIDINE-2-CARBONYL]CYCLOPROPANE-1-CARBONITRILE (16)




1-Cyanocyclopropane-1-carboxylic acid (41 mg, 0.37 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (50 mg, 0.33 mmol) using general procedure A for amide coupling with DMF as solvent to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.23-7.45 (m, 5 H), 5.43 (dd, $J=8.5, 6.0$ Hz, 1 H), 4.45 (td, $J=7.9, 3.3$ Hz, 1 H), 4.03-4.21 (m, 1 H), 2.93 (dddd, $J=12.3, 8.9, 6.9, 3.5$ Hz, 1 H), 2.47 (dddd, $J=12.4, 9.3, 7.7, 6.0$ Hz, 1 H), 1.66-1.77 (m, 1 H), 1.49-1.64 (m, 3 H). LC-MS (Method A): $m/z = 243.3$ $[\text{M}+\text{H}]^+$, 0.87 min.

EXAMPLE 17

PREPARATION OF 2,2-DIMETHYL-3-OXO-3-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]PROPANENITRILE


(17)



2-Cyano-2,2-dimethylacetic acid (42 mg, 0.37 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (50 mg, 0.33 mmol) using general procedure A for amide coupling with DMF as solvent to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.22-7.43 (m, 5 H), 5.41 (dd, $J=8.9, 6.4$ Hz, 1 H), 4.39 (td, $J=8.2, 2.8$ Hz, 1 H), 4.18 (ddd, $J=9.7, 8.2, 6.7$ Hz, 1 H), 2.87-2.99 (m, 1 H), 2.46 (dddd, $J=12.4, 9.8, 7.8, 6.4$ Hz, 1 H), 1.73 (s, 3 H), 1.61 (s, 3 H). LC-MS (Method A): $m/z = 245.1$ $[\text{M}+\text{H}]^+$, 0.95 min.

EXAMPLE 18


PREPARATION OF 3-[(3S)-3-PHENYL-1,2-OXAZOLIDINE-2-CARBONYL]-2,1-BENZOXAZOLE (18)



2,1-Benzoxazole-3-carboxylic acid (42 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.91 (td, $J=0.9, 9.0$ Hz, 1 H), 7.78 (d, $J=9.3$ Hz, 1 H), 7.51 (ddd, $J=0.9, 6.3, 9.1$ Hz, 1 H), 7.45-7.37 (m, 4 H), 7.34-7.27 (m, 2 H), 5.71 (dd, $J=6.3, 8.3$ Hz, 1 H), 4.40 (dt, $J=3.5, 7.5$ Hz, 1 H), 4.09 (ddd, $J=6.8, 8.0, 9.0$ Hz, 1 H), 3.01 (dddd, $J=3.5, 6.8, 8.7, 12.1$ Hz, 1 H), 2.34 (dddd, $J=6.0, 7.4, 9.0, 12.2$ Hz, 1 H). LC-MS (Method A): $m/z = 295.1$ $[\text{M}+\text{H}]^+$, 1.03 min.

EXAMPLE 19


PREPARATION OF 2-HYDROXY-2-METHYL-1-[(3S)-3-PHENYL-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE
(19)



2-Hydroxy-2-methylpropanoic acid (27 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.39-7.23 (m, 5H), 5.39 (dd, $J=6.5, 8.3$ Hz, 1H), 4.77 (s, 1H), 4.27 (t, $J=3.1, 7.8$ Hz, 1H), 3.92-3.84 (m, 1H), 2.94-2.84 (m, 1H), 2.24-2.11 (m, 1H), 1.39 (s, 3H), 1.37 (s, 3H). LC-MS (Method A): $m/z = 236.3$ $[\text{M}+\text{H}]^+$, 0.84 min.

EXAMPLE 20

PREPARATION OF (3S)-2-(1-METHYLCYCLOPROPANECARBONYL)-3-PHENYL-1,2-OXAZOLIDINE (20)




1-Methylcyclopropane-1-carboxylic acid (26 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.38-7.23 (m, 5H), 5.33 (dd, $J=6.3, 8.8$ Hz, 1H), 4.25 (dt, $J=3.0, 7.5$ Hz, 1H), 3.85 (ddd, $J=6.7, 7.9, 9.5$ Hz, 1H), 2.85 (dddd, $J=3.0, 6.5, 9.0, 12.0$ Hz, 1H), 2.16 (dddd, $J=7.7, 9.6, 12.1$ Hz, 1H), 1.35 (s, 3H), 1.05-0.98 (m, 1H), 0.88 (ddd, $J=4.0, 6.0, 9.9$ Hz, 1H), 0.62-0.56 (m, 1H), 0.53-0.48 (m, 1H). LC-MS (Method A): $m/z = 232.3$ $[\text{M}+\text{H}]^+$, 0.96 min.

EXAMPLE 21

PREPARATION OF (3S)-3-PHENYL-2-{PYRAZOLO[1,5-a]PYRIDINE-3-CARBONYL}-1,2-OXAZOLIDINE


(21)



5 Pyrazolo[1,5-a]pyridine-3-carboxylic acid (42 mg, 0.26 mmol) was reacted with (3S)-3-phenyl-1,2-oxazolidine (40 mg, 0.26 mmol) using general procedure A for amide coupling to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.86 (td, *J*=1.0, 7.0 Hz, 1H), 8.64 (s, 1H), 8.25 (td, *J*=1.2, 8.8 Hz, 1H), 7.55 (ddd, *J*=1.0, 6.8, 9.0 Hz, 1H), 7.43-7.36 (m, 4H), 7.31-7.26 (m, 1H), 7.15 (dt, *J*=1.3, 6.9 Hz, 1H), 5.62 (dd, *J*=6.3, 8.5 Hz, 1H), 4.37 (dt, *J*=2.9, 7.7 Hz, 1H), 3.88 (ddd, *J*=6.8, 8.0, 9.5 Hz, 1H), 2.92 (dddd, *J*=7.0, 6.5, 9.0, 12.0 Hz, 1H), 2.27 (dddd, *J*=6.3, 7.7, 9.6, 12.1 Hz, 1H). LC-MS (Method A): *m/z* = 294.3 [*M*+*H*]⁺, 0.98 min.

EXAMPLE 22


PREPARATION OF TERT-BUTYL (3S)-3-PHENYL-1,2-OXAZOLIDINE-2-CARBOXYLATE (22)



15 Methanesulfonyl chloride (1.4 mL, 18.91 mmol) was added to a solution of tert-butyl N-[(3R)-3-hydroxy-3-phenylpropoxy] carbamate (4.4 g, 16.45 mmol) and Et₃N (4.5 mL, 32.9 mmol) in CH₂Cl₂ (383 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 1.5 h. The reaction was diluted with CH₂Cl₂ and washed with water. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound. ³H NMR (400 MHz, DMSO-*d*₆) δ 7.40-7.18 (m, 5H), 5.15 (dd, *J*=5.9, 8.7 Hz, 1H), 4.14 (dt, *J*=2.9, 8.1 Hz, 1H), 3.72 (ddd, *J*=6.9, 8.0, 9.4 Hz, 1H), 2.80 (dddd, *J*=3.1, 6.8, 8.8, 12.1 Hz, 1H), 2.20-2.07 (m, 1H), 1.38 (s, 9H). LC-MS (Method A): *m/z* = 194.1 [*M*+*H*]⁺ and 250.3 [*M*+*H*]⁺, 1.07 min.

EXAMPLE 23


PREPARATION OF 1-[3-(3-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (23)



5 **Pivaloyl chloride** (53 μL , 0.430 mmol) was added to a solution 3-(3-fluorophenyl)- 1,2-
oxazolidine (60 mg, 0.359 mmol) and Et_3N (75 μL , 0.539 mmol) in CH_2Cl_2 (2.5 mL). The reaction
mixture was stirred at room temperature for 40 min, diluted with CH_2Cl_2 and washed with sat. NH_4Cl
solution. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure.
The mixture of **enantiomers** was resolved by **chiral HPLC** on a Chiralcel **OJ-H** (25 x 2.0 cm), 5 μm
10 column using a mobile phase of **n-hexane/2-propanol** 90/10 % v/v to afford the **title** compound. ^1H NMR
(400 MHz, CDCl_3) δ 7.34-7.27 (m, 1H), 7.11-7.07 (m, 1H), 7.02 (dt, $J=9.8, 2.0$ Hz, 1H), 6.95 (tdd,
 $J=8.4, 2.6, 0.9$ Hz, 1H), 5.42 (dd, $J=8.8, 6.5$ Hz, 1H), 4.31-4.24 (m, 1H), 3.90 (ddd, $J=9.8, 8.0, 6.5$ Hz,
1H), 2.83 (dddd, $J=12.1, 9.1, 6.4, 2.8$ Hz, 1H), 2.30 (dddd, $J=12.3, 9.8, 7.7, 6.4$ Hz, 1H), 1.31 (s, 9H).
LC-MS (Method A): $m/z = 252.2$ $[\text{M}+\text{H}]^+$, 1.14 min. **e.e.=100%** as determined on a Chiralcel **OJ-H** (25 x
15 0.46 cm), 5 μm column using a mobile phase of **n-hexane/2-propanol** 90/10 % v/v.

EXAMPLE 24

PREPARATION OF 1-[3-(2-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (24)




20 **3-(2-Fluorophenyl)- 1,2-oxazolidine** (31 mg, 0.18 mmol) was dissolved in dry CH_2Cl_2 (1.76 mL).
Pyridine (30 μL , 0.36 mmol) and a solution of pivaloyl chloride (27 μL , 0.216 mmol) in dry CH_2Cl_2 (0.1
mL) were then added at 0 $^\circ\text{C}$. The mixture was stirred at room temperature for 3 h, then a second addition
of pyridine (15 μL , 0.18 mmol) and a solution of pivaloyl chloride (13.5 μL , 0.108 mmol) in dry CH_2Cl_2
(0.05 mL) was made. The reaction mixture was stirred at room temperature for a further 3 h then the
25 mixture was concentrated under reduced pressure. The residue was dissolved in EtOAc and washed with
sat. aqueous NH_4Cl solution, with sat. aqueous NaHCO_3 solution and with brine. The organic phase was
dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The **crude** product (44 mg) was
purified by column chromatography (**cyclohexane-EtOAc**, 95:5 to 60:40) to afford the title compound as

a mixture of enantiomers. ^1H NMR (400 MHz, CDCl_3) δ 7.31-7.21 (m, 2H), 7.14-7.02 (m, 2H), 5.69 (dd, $J=8.8, 8.0$ Hz, 1H), 4.25 (dt, $J=8.5, 3.0$ Hz, 1H), 3.96-3.88 (m, 1H), 2.95-2.85 (m, 1H), 2.32-2.22 (m, 1H), 1.33 (s, 9H). LC-MS (Method A): $m/z = 252.3$ $[\text{M}+\text{H}]^+$, 1.14 min.

EXAMPLE 25

PREPARATION OF 1-[3-(4-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE


(25)



Pivaloyl chloride (0.1 mL, 0.8 mmol) was added to a solution of 3-(4-fluorophenyl)-1,2-oxazolidine (90 mg, 0.54 mmol) and pyridine (0.1 mL, 1.35 mmol) in CH_2Cl_2 (4 mL). The reaction mixture was stirred at room temperature for 2 h. Volatiles were removed under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3CN , 100:0 to 70:30) to afford the title compound as a mixture of enantiomers. ^1H NMR (400 MHz, CDCl_3) δ 7.34-7.22 (m, 2H), 7.08-6.96 (m, 2H), 5.40 (dd, $J=8.8, 6.5$ Hz, 1H), 4.27 (tdd, $J=7.8, 2.9, 0.8$ Hz, 1H), 3.89 (ddd, $J=9.9, 8.0, 6.5$ Hz, 1H), 2.88-2.74 (m, 1H), 2.30 (dddd, $J=12.3, 9.9, 7.7, 6.5$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 252.2$ $[\text{M}+\text{H}]^+$, 1.17 min.

EXAMPLE 26

PREPARATION OF TERT-BUTYL 3-(2-FLUOROPHENYL)-1,2-OXAZOLIDINE-2-CARBOXYLATE (26)




Methanesulfonyl chloride (57 μL , 0.746 mmol) was added to a solution of tert-butyl N-[3-(2-fluorophenyl)-3-hydroxypropoxy]carbamate (185 mg, 0.65 mmol) and Et_3N (0.182 mL, 1.3 mmol) in dry CH_2Cl_2 (15 mL) at 0 $^\circ\text{C}$. The reaction mixture was stirred at 0 $^\circ\text{C}$ for 2 h then a second addition of methanesulfonyl chloride (0.11 mL, 0.15 mmol) and Et_3N (36 μL , 0.26 mmol) was made. After 5 h the reaction was diluted with CH_2Cl_2 and washed with sat. aqueous NaHCO_3 solution and sat. aqueous NH_4Cl solution. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product (178 mg) was purified by column chromatography (cyclohexane-EtOAc, 95:5 to 60:40) to afford the title compound as a mixture of enantiomers. ^1H NMR (400 MHz, CDCl_3) δ 7.50 (dt, $J=8.0, 1.5$ Hz, 1H), 7.28-7.22 (m, 1H), 7.17-7.11 (m, 1H), 7.08-7.01 (m, 1H), 5.54-5.48 (m,

1H), 4.17 (dt, J=8.0, 3.7 Hz, 1H), 3.96-3.88 (m, 1H), 2.90-2.80 (m, 1H), 2.30-2.20 (m, 1H), 1.48 (s, 9H). LC-MS (Method A): m/z = 268.1 [M+H]⁺, 1.17 min.

EXAMPLE 27


PREPARATION OF TERT-BUTYL 3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINE-2-CARBOXYLATE (27)



Methane sulfonyl chloride (130 μ L, 1.73 mmol) was added to a solution of tert-butyl N-[3-(4-fluorophenyl)-3-hydroxypropoxy]carbamate (430 mg, 1.5 mmol) and F₃N (410 μ L, 3.0 mmol) in CH₂Cl₂ (15 mL) which had been pre-cooled to 0 °C. The reaction mixture was stirred at 0 °C for 1 h and at room temperature for 1 h. The reaction mixture was diluted with CH₂Cl₂ and washed with sat. NaHCO₃ solution and sat. NH₄Cl solution. The organic portions were combined, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 80:20) to afford the title compound as a mixture of enantiomers. ¹H NMR (400 MHz, CDCl₃) δ 7.40-7.30 (m, 2H), 7.09-6.97 (m, 2H), 5.20 (dd, J=8.7, 5.5 Hz, 1H), 4.28-4.13 (m, 1H), 3.91 (ddd, J=8.9, 8.2, 7.0 Hz, 1H), 2.79 (dddd, J=12.3, 8.7, 7.0, 3.7 Hz, 1H), 2.28 (dddd, J=12.2, 8.9, 7.6, 5.6 Hz, 1H), 1.48 (s, 9H). LC-MS (Method A): m/z = 212.2 [M-tBuF], 1.10 min.

EXAMPLES 28 AND 29

PREPARATION OF 1-[(3R)-3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE AND 1-[(3S)-3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (62 AND 61)



Trifluoroacetic acid (500 μ L) was added to a solution of tert-butyl 3-(6-fluoropyridin-3-yl)-1,2-oxazolidine-2-carboxylate (161 mg, 0.60 mmol) in CH₂Cl₂ (5 mL). The reaction was stirred at room temperature for 30 min and concentrated under reduced pressure to give crude 2-fluoro-5-[(1,2-oxazolidin-3-yl)pyridine] as its trifluoroacetic acid salt (157 mg). Et₃N (300 μ L, 2.15 mmol) and pivaloyl chloride (100 μ L, 0.851 mmol) were sequentially added to a solution of the crude 2-fluoro-5-[(1,2-oxazolidin-3-yl)pyridine] trifluoroacetic acid salt (75 mg) in CH₂Cl₂ (3 mL). The reaction was stirred at room temperature for 1 h, diluted with CH₂Cl₂ and washed twice with sat. NaHCO₃ solution. The organic


portion was dried over Na_2SO_4 filtered and concentrated under reduced pressure. The crude product was purified by reverse phase chromatography (water-CITCN, 100:0 to 20:80) to afford the title compound as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/2-propanol 80/20% v/v to afford the two separated enantiomers.

First eluting enantiomer, 1-[(3R)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one. ^1H NMR (400 MHz, CDCl_3) δ 8.20 (d, $J=2.5$ Hz, 1H), 7.74 (td, $J=8.0, 2.4$ Hz, 1H), 6.92 (dd, $J=8.4, 2.9$ Hz, 1H), 5.45 (dd, $J=8.8, 6.8$ Hz, 1H), 4.35-4.28 (m, 1H), 3.93 (ddd, $J=10.0, 8.1, 6.5$ Hz, 1H), 2.87 (dddd, $J=12.4, 8.9, 6.4, 2.6$ Hz, 1H), 2.32 (dddd, $J=12.4, 10.0, 7.7, 6.8$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 253.2$ $[\text{M}+\text{H}]^+$, 0.92 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/2-propanol 80/20% v/v.

Second eluting enantiomer, 1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one. ^1H NMR (400 MHz, CDCl_3) δ 8.22-8.18 (m, 1H), 7.74 (td, $J=7.9, 2.5$ Hz, 1H), 6.92 (dd, $J=8.4, 2.9$ Hz, 1H), 5.51-5.41 (m, 1H), 4.32 (td, $J=8.0, 2.4$ Hz, 1H), 3.98-3.85 (m, 1H), 2.94-2.80 (m, 1H), 2.39-2.25 (m, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 253.3$ $[\text{M}+\text{H}]^+$, 0.92 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/2-propanol 80/20% v/v.

EXAMPLE 30

PREPARATION OF 1-[(3S)-3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (61)




Pivaloyl chloride (256 μE , 2.09 mmol) was added to a solution of 2-fluoro-5-[(3S)-1,2-oxazolidin-3-yl]pyridine (320 mg) and Et_3N (396 μL , 2.85 mmol) in CH_2Cl_2 (10 mL) which had been pre-cooled to 0 $^\circ\text{C}$. The reaction mixture was stirred at 0 $^\circ\text{C}$ for 30 min, then diluted with CH_2Cl_2 and washed with water. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 0:100) to give the title compound (452 mg) as a 4.7:95.3 mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/2-propanol 80/20% v/v to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 8.20 (d, $J=2.5$ Hz, 1H), 7.74 (td, $J=8.0, 2.6$ Hz, 1H), 6.92 (dd, $J=8.5, 3.0$ Hz, 1H), 5.45 (dd, $J=8.8, 6.8$ Hz, 1H), 4.32 (td, $J=7.8, 2.6$ Hz, 1H), 3.93 (ddd, $J=10.0, 8.2, 6.4$ Hz, 1H), 2.92-2.83 (m, 1H), 2.32 (dddd, $J=12.4, 10.0, 7.6, 6.9$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 253.2$ $[\text{M}+\text{H}]^+$, 0.92 min. e.e.=100% as

determined on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/2-propanol 80/20% v/v.

EXAMPLE 31 AND 32

PREPARATION OF 1-[3-(4-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE

ENANTIOMER 1 AND ENANTIOMER 2 (73 AND 74)




Pivaloyl chloride (0.1 mL, 0.8 mmol) was added to a solution of 3-(4-fluorophenyl)-1,2-oxazolidine (90 mg, 0.54 mmol) and pyridine (0.1 mL, 1.35 mmol) in CH_2Cl_2 (4 mL). The reaction mixture was stirred at room temperature for 2 h. Volatiles were removed under reduced pressure. The crude product was purified by reverse phase chromatography (water- CH_3QSI , 100:0 to 70:30) to afford (S)-1-[3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one and (R)-1-[3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralcel OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(EtOH/MeOH 1/1+0.1% isopropylamine) 85/15 % v/v to afford the two title compounds as separated enantiomers.

First eluting enantiomer, Enantiomer 1. ^1H NMR (400 MHz, CDCl_3) δ 7.36-7.22 (m, 3H), 7.11-6.92 (m, 2H), 5.40 (dd, $J=8.8, 6.5$ Hz, 1H), 4.27 (dt, $J=7.9, 2.8$ Hz, 1H), 3.90 (ddd, $J=9.9, 7.9, 6.5$ Hz, 1H), 2.82 (dddd, $J=12.1, 9.1, 6.4, 2.8$ Hz, 1H), 2.30 (dddd, $J=12.3, 9.9, 7.6, 6.7$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 252.2$ $[\text{M}+\text{H}]^+$, 1.10 min. e.e.=100% as determined on a Chiralcel OJ-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(EtOH/MeOH 1/1+0.1% isopropylamine) 85/15 % v/v.

Second eluting enantiomer, Enantiomer 2. ^1H NMR (400 MHz, CDCl_3) δ 7.33-7.24 (m, 2H), 7.07-6.98 (m, 2H), 5.40 (dd, $J=8.8, 6.8$ Hz, 1H), 4.27 (dt, $J=2.8, 7.8$ Hz, 1H), 3.90 (ddd, $J=9.9, 8.0, 6.5$ Hz, 1H), 2.82 (dddd, $J=12.2, 9.1, 6.4, 2.9$ Hz, 1H), 2.30 (dddd, $J=12.3, 9.9, 7.6, 6.7$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 252.2$ $[\text{M}+\text{H}]^+$, 1.11 min. e.e.=100% as determined on a Chiralcel OJ-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(EtOH/MeOH 1/1+0.1% isopropylamine) 85/15 % v/v.

EXAMPLE 33


PREPARATION OF 4-[(3S)-2-(2,2-D(METHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL) BENZONITRILE (87)



Pivaloyl chloride (1.52 mL, 12.5 mmol) was added to a solution of 4-(1,2-oxazolidin-3-yl)benzonitrile (1.45 g, 8.33 mmol) and pyridine (1.68 mL, 20.82 mmol) in CH_2Cl_2 (20 mL). The reaction mixture was stirred at room temperature for 1 h. Volatiles were removed under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 70:30) to afford the title compound (2.06 g, 96%) as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μm column using a mobile phase of *n*-hexane/(ethanol + 0.1 % isopropylamine) 70/30 v/v and a flow rate of 18 mL/min to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.68–7.60 (m, 2H), 7.45–7.37 (m, 2H), 5.43 (dd, $J=9.0, 6.8$ Hz, 1H), 4.34–4.24 (m, 1H), 3.91 (ddd, $J=10.1, 8.1, 6.3$ Hz, 1H), 2.87 (dddd, $J=12.3, 8.9, 6.4, 2.6$ Hz, 1H), 2.27 (dddd, $J=12.3, 10.0, 7.7, 6.8$ Hz, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 259.2$ $[\text{M}+\text{H}]^+$, 1.03 min.

EXAMPLE 34

PREPARATION OF 4-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL] BENZONITRILE (88)




To a solution of 3,3-difluoro-2,2-dimethylpropanoic acid (43 mg, 0.316 mmol) in CH_2Cl_2 (2 mL) were added oxalylchloride (26 μL , 0.316 mmol) and a drop of DMF. The mixture was stirred at room temperature for 2 h, then concentrated under reduced pressure. The residue was redissolved in CH_2Cl_2 (2 mL) and this solution was then added slowly to a stirred solution of 4-(1,2-oxazolidin-3-yl)benzonitrile (50 mg, 0.28 mmol) and DIPEA (0.12 mL, 0.70 mmol) in CH_2Cl_2 (2 mL) at 0 °C. The mixture was stirred at room temperature for 2 h, then the solvent was evaporated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 1:1) to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.69–7.61 (m, 2H), 7.43–7.35 (m, 2H), 6.32 (t, $J=57.2$ Hz, 1H), 5.44 (dd, $J=8.9, 6.8$ Hz, 1H), 4.38–4.28 (m, 1H), 3.97 (ddd, $J=9.8, 8.1, 6.4$ Hz, 1H), 2.91 (dddd, $J=12.0, 9.1, 6.4,$

2.9 Hz, 1H), 2.31 (dddd, $J = 12.4, 9.9, 7.6, 6.7$ Hz, 1H), 1.42-1.36 (m, 6H). LC-MS (Method A): $m/z = 295.1$ $[M+H]^+$, 1.03 min.

EXAMPLE 35


PREPARATION OF 1-[(3S)-3-(5-CHLOROPYRIDIN-2-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (89)



Pivaloyl chloride (22 μ L, 0.179 mmol) was added to a solution of 5-chloro-2-[(3S)-1,2-oxazolidin-3-yl]pyridine (32 mg, 0.174 mmol) and triethylamine (34 μ L, 0.244 mmol) in CH_2Cl_2 (3 mL), which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 30 min, diluted with CH_2Cl_2 and washed with water. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 0:100) to afford the title compound. ^1H NMR (400 MHz, $CDCl_3$) δ 8.54-8.52 (m, 1H), 7.63 (dd, $J = 8.5, 2.5$ Hz, 1H), 7.32 (d, $J = 8.5$ Hz, 1H), 5.50 (dd, $J = 9.0, 6.3$ Hz, 1H), 4.33-4.22 (m, 1H), 3.94 (ddd, $J = 9.4, 7.9, 6.8$ Hz, 1H), 2.85-2.75 (m, 1H), 2.71-2.60 (m, 1H), 1.31 (s, 9H). LC-MS (Method A): $m/z = 269.1$ $[M+H]^+$, 1.02 min.

EXAMPLE 36

PREPARATION OF (3S)-2-[1-(2,2-DIFLUOROETHYL)-3-METHYLAZETIDINE-3-CARBONYL]-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINE (90)




Et_3N (37 μ L, 0.264 mmol) and 2,2-difluoroethyl methanesulfonate (28 mg, 0.132 mmol) were sequentially added to a solution of (3S)-3-(4-fluorophenyl)-2-(3-methylazetidine-3-carbonyl)-1,2-oxazolidine (24 mg, 0.088 mmol) in CH_2Cl_2 (5 mL) which had been pre-cooled to 0 °C. The reaction was stirred at 0 °C for 2h, diluted with CH_2Cl_2 and washed with water. The organic portion was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (KP-NH, cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound. ^1H NMR (400 MHz, $CDCl_3$) δ 7.31-7.25 (m, 2H), 7.09-7.01 (m, 2H), 5.74 (tt, $J = 56.2, 5.0$ Hz, 1H), 5.37 (dd,

,7=8.7, 6.1 Hz, **IB**), 4.24 (td, J=7.6, 3.6 Hz, 1H), 3.92 (ddd, J=9.0, 8.0, 6.8 Hz, 1H), 3.57 (d, J=7.5 Hz, 1H), 3.42-3.37 (m, 3H), 2.73-2.91 (m, 3H), 2.35 (dddd, J=12.4, 9.2, 7.5, 6.3 Hz, 1H), 1.65 (s, 3H). LC-MS (Method A): m/z = 329.2 [M+H]⁺, 0.50 min.

EXAMPLES 37 AND 38

PREPARATION OF 4-[(3S)-2-(3,3-DIFLUORO-2-METHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILES (91) AND (92)




DMF (20 μ L) and oxalyl chloride (50 μ L, 0.590 mmol) were sequentially added to a solution of the mixture containing 3,3-difluoro-2-methylpropanoic acid (100 mg) and the reaction was stirred at room temperature for 1 h. A solution of 4-[(3S)-1,2-oxazolidin-3-yl]benzonitrile (100 mg, 0.575 mmol) and DIPEA (250 μ L, 1.44 mmol) in CH₂Cl₂ (2 mL) was added. The reaction was stirred at room temperature for 1.5 h, diluted with 3:1 and washed twice with water. The organic portion was dried over Na₂SO₄, filtered and concentrated under reduced pressure to provide 4-[(3S)-2-((S)-3,3-difluoro-2-methylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile and 4-[(3S)-2-((R)-3,3-difluoro-2-methylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 0:100) to afford the first eluting diastereoisomer of the title compound and a mixture of the second eluting diastereoisomer with several byproducts. The latter was further purified by semi-preparative HPLC to afford the second eluting diastereoisomer of the title compound (2.1 mg).

First eluting diastereoisomer, Diastereoisomer **1**: ¹H NMR (400 MHz, **OMSO-d₆**) δ 7.89-7.80 (m, 2H), 7.53-7.45 (m, 2H), 6.13 (td, J=5.7, 6.3 Hz, 1H), 5.51-5.37 (m, 1H), 4.30 (td, J=7.5, 2.5 Hz, 1H), 3.99-3.85 (m, 1H), 3.52-3.36 (m, 1H), 3.06-2.84 (m, 1H), 2.29-2.10 (m, 1H), 1.24 (d, J=7.0 Hz, 3H). LC-MS (Method A): m/z = 281.2 [M+H]⁺, 0.95 min.

Second eluting diastereoisomer, Diastereoisomer **2**: ¹H NMR (400 MHz, DMSO-d₆) δ 7.91-7.80 (m, 2H), 7.51-7.44 (m, 2H), 6.19 (td, J=6.5, 9.5 Hz, 1H), 5.42 (t, J=7.7 Hz, 1H), 4.29 (td, J=7.6, 3.1 Hz, 1H), 3.97 (ddd, J=9.4, 8.0, 6.4 Hz, 1H), 3.56-3.38 (m, **IB**). 2.95 (dddd, J=12.2, 9.1, 6.2, 3.1 Hz, 1H), 2.28-2.10 (m, 1H), 1.14 (d, J=7.0 Hz, 3H). LC-MS (Method A): m/z = 281.2 [M+H]⁺, 0.94 min.

EXAMPLE 39


PREPARATION OF 2-FLUORO-5-[(3S)-2-[(1-METHYLCYCLOPROPYL)SULFONYL]-1,2-OXAZOLIDIN-3-YL]PYRIDINE (93)



5 A solution of 2-fluoro-5-[(3S)-1,2-oxazolidin-3-yl]pyridine (30 mg, 0.179 mmol), 1-methylcyclopropane-1-sulfonyl chloride (30 mg, 0.197 mmol), pyridine (40 μ L, 0.358 mmol) and DMAP (5 mg, 0.041 mmol) in CH_2Cl_2 (1.5 mL) was stirred at room temperature for 20 h. Although not complete, the reaction was diluted with CH_2Cl_2 and washed with 0.5M HCl solution. The organic portion was washed with 1M HCl solution and sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 70:30) to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 8.29-8.24 (m, 1H), 7.84 (td, $J=8.0, 2.3$ Hz, 1H), 6.95 (dd, $J=8.5, 2.8$ Hz, 1H), 5.44 (dd, $J=8.3, 5.5$ Hz, 1H), 4.43-4.34 (m, 1H), 4.24-4.16 (m, 1H), 2.98-2.87 (m, 1H), 2.35 (dd, $J=12.3, 8.2, 5.5$ Hz, 1H), 1.68-1.58 (m, 4H), 1.48 (ddd, $J=10.6, 6.5, 5.5$ Hz, 1H), 1.03 (ddd, $J=9.1, 6.8, 5.4$ Hz, 1H), 0.86 (ddd, $J=9.0, 7.0, 5.3$ Hz, 1H). LC-MS (Method A): m/z = 287.1 $[\text{M}+\text{H}]^+$, 0.90 min.

EXAMPLES 40 AND 41

PREPARATION OF 3,3,3-TRIFLUORO-1-[(3S)-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2-METHYLPROPAN-1-ONES (94) AND (95)



20 A solution of (3S)-3-(4-fluorophenyl)-1,2-oxazolidine (30 mg, 0.179 mmol), 3,3,3-trifluoro-2-methylpropanoic acid (28 mg, 0.197 mmol), T_3P (50% wt solution in EtOAc, 234 μ L, 0.394 mmol) and pyridine (40 μ L, 0.537 mmol) in EtOAc (1 mL) was stirred at room temperature for 1.5 h. The reaction was diluted with EtOAc and the organic portion was washed with 1M HCl solution and sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to provide (R)-3,3,3-trifluoro-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one and (S)-3,3,3-trifluoro-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one. The crude product was purified


by chiral HPLC on a Chiralcel OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol + 0.1% isopropylamine) 30/70% v/v and a flow rate of 1.8 mL/min to afford the two separated diastereoisomers of the title compound.

First eluting diastereoisomer, Diastereoisomer 1: ^1H NMR (400 MHz, CDCl_3) δ 7.31–7.22 (m, 2H), 7.00–7.08 (m, 2H), 5.41 (dd, $J=8.5, 5.8$ Hz, 1H), 4.31 (td, $J=7.6, 3.9$ Hz, 1H), 4.03–3.93 (m, 1H), 3.92–3.79 (m, 1H), 2.88 (dddd, $J=12.4, 8.7, 6.8, 4.0$ Hz, 1H), 2.44–2.34 (m, 1H), 1.38 (d, $J=7.0$ Hz, 3H). LC-MS (Method A): $m/z = 292.1$ $[\text{M}+\text{H}]^+$, 1.05 min.

Second eluting diastereoisomer, Diastereoisomer 2: ^1H NMR (400 MHz, CDCl_3) δ 7.29–7.21 (m, 2H), 7.06–6.98 (m, 2H), 5.37 (dd, $J=8.8, 5.3$ Hz, 1H), 4.32 (td, $J=7.7, 4.5$ Hz, 1H), 4.05 (td, $J=8.2, 7.0$ Hz, 1H), 2.96–2.85 (m, 1H), 2.42–2.31 (m, 1H), 1.97–1.85 (m, 1H), 1.48 (br. s., 3H). LC-MS (Method A): $m/z = 292.1$ $[\text{M}+\text{H}]^+$, 1.07 min.

EXAMPLE 42


PREPARATION OF 1-[(3S)-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-3-HYDROXY-2,2-DIMETHYLPROPAN-1-ONE (96)



A mixture of (3S)-3-(4-fluorophenyl)-1,2-oxazolidine (100 mg, 0.6 mmol), 3-hydroxy-2,2-dimethylpropanoic acid (85 mg, 0.72 mmol), HATU (456 mg, 1.2 mmol) and DIPEA (0.2 mL, 1.2 mmol) in CH_2Cl_2 (7 mL) was stirred at room temperature for 6h. The mixture was evaporated under reduced pressure and the crude product was purified twice by column chromatography (first column CH_2Cl_2 -MeOH, 100:0 to 90:10; second column cyclohexane-EtOAc, 100:0 to 50:50) and then by Prep-HPLC with the following conditions: column: Gemini C18 110A AXIA Column, 100x30 mm, 5 μ m; Mobile Phase A: 10 mmol/L ammonium bicarbonate aqueous solution adjusted to pH 10 with ammonia; Mobile Phase B: acetonitrile (20% to 70% over 10 min); Detector, UV 210 to 350 nm to afford the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.31–7.21 (m, 2H), 7.09–6.99 (m, 2H), 5.38 (dd, $J=8.8, 6.3$ Hz, 1H), 4.35–4.26 (m, 1H), 3.94 (ddd, $J=9.7, 8.0, 6.5$ Hz, 1H), 3.62–3.48 (m, 2H), 3.09 (t, $J=7.1$ Hz, 1H), 2.84 (dddd, $J=12.1, 8.9, 6.6, 3.1$ Hz, 1H), 2.30 (dddd, $J=12.4, 9.7, 7.6, 6.3$ Hz, 1H), 1.37 (s, 3H), 1.29 (s, 3H). LC-MS (Method B): $m/z = 267.0$ $[\text{M}+\text{H}]^+$, 0.86 min.

EXAMPLE 43


1-[(3S)-3-(5-FLUORO-6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (97)



3-Fluoro-2-methyl-5-[(3S)-1,2-oxazolidin-3-yl]pyridine (38 mg, 0.208 mmol) was dissolved in CH_2Cl_2 (1.0 mL). A solution of pyridine (0.034 mL, 0.416 mmol) in CH_2Cl_2 (0.5 mL) and a solution of pivaloyl chloride (0.033 mL, 0.27 mmol) in CH_2Cl_2 (0.5 mL) were then added at 0 °C. The mixture was then stirred at room temperature for 30 min then the mixture was concentrated under reduced pressure. The residue was dissolved in EtOAc and washed with sat. NH_4Cl solution, twice with sat. NaHCO_3 solution and with brine. The organic phase was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 88:12 to 0:100) to afford the title compound as an unequal mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1 % isopropylamine) 75/25% v/v and a flow rate of 18 mL/min.

Second eluting enantiomer, Enantiomer 2: ^1H NMR (400 MHz, CDCl_3) δ 8.29 (s, 1H), 7.27 (m, 1H), 5.43 (t, $J=8.8, 6.8$ Hz, 1H), 4.30 (td, $J=8.0, 2.5$ Hz, 1H), 3.96-3.88 (m, 1H), 2.91-2.82 (m, 1H), 2.51 (d, $J=3.0$ Hz, 3H), 2.36-2.26 (m, 1H), 1.30 (s, 9H). LC-MS (Method A): m/z = 267.2 $[\text{M} + \text{H}]^+$, 0.92 min. e.e.=98.8% as determined on a Chiralpak TC (25 x 0.46 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 75/25% v/v, flow rate: 1 mL/min, retention time: 7.4 min.

EXAMPLE 44

4-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-(5,5- $^2\text{H}_2$)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE (98)


A solution of 4-[(3S)-(5,5- $^2\text{H}_2$)-1,2-oxazolidin-3-yl]benzonitrile (30 mg, 0.17 mmol), pivaloyl chloride (0.023 mL, 0.19 mmol) and Et_3N (0.035 mL, 0.25 mmol) in CH_2Cl_2 (4 mL) was stirred at room temperature for 1 h. The mixture was diluted with CH_2Cl_2 and washed with brine. The organic phase was

separated, dried over Na_2SO_4 , filtered and evaporated under reduced pressure. The crude product was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 65/35 % v/v and a flow rate of 18 mL/min.

Second eluting enantiomer, Enantiomer 2: ^1H NMR (400 MHz, CDCl_3) δ 7.68-7.58 (m, 2 H) 7.45-7.35 (m, 2 H) 5.43 (dd, $J=9.0$, 6.8 Hz, 1 H) 2.84 (dd, $J=12.3$, 9.0 Hz, 1 H) 2.32-2.18 (m, 1 H) 1.30 (s, 9 H). LC-MS (Method A): m/z = 261.2 $[\text{M}+\text{H}]^+$, 1.03 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 65/35 % v/v, flow rate: 1 mL/min, retention time: 7.3 min.

EXAMPLES 45 AND 46

10 4-[(3S,5S)-2-(2,2-DIMETHYLPROPANOYL)-5-METHYL-1,2-OXAZOLIDIN-3-YL]BENZONITRILE (99) AND 4-[(3S,5R)-2-(2,2-DIMETHYLPROPANOYL)-5-METHYL-1,2-OXAZOLIDIN-3-YL]BENZONITRILE (100)



A solution of 4-[(3S)-5-methyl-1,2-oxazolidin-3-yl]benzonitrile (50 mg, 0.26 mmol), pivaloyl chloride (0.036 mL, 0.29 mmol) and Et_3N (0.054 mL, 0.39 mmol) in CH_2Cl_2 (4 mL) was stirred at room temperature for 1h 30 min. The mixture was diluted with CH_2Cl_2 and washed with 0.5M HCl solution and brine. The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude products were purified by column chromatography (cyclohexane-EtOAc, 90:10 to 65:35) to give a mixture of diastereoisomers as a colorless oil. This mixture was resolved by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1 % isopropylamine) 70/30 % v/v and a flow rate of 18 mL/min.


4-[(3S,5S)-2-(2,2-dimethylpropanoyl)-5-methyl-1,2-oxazolidin-3-yl]benzonitrile (First eluted diastereoisomer): ^1H NMR (400 MHz, CDCl_3) δ 7.62 (d, $J=8.4$ Hz, 2H), 7.40 (d, $J=8.0$ Hz, 2H), 5.43 (t, $J=8.41$ Hz, 1H), 4.16 (dqum, $J=11.0$, 5.6 Hz, 1H), 2.93 (ddd, $J=12.3$, 8.9, 5.1 Hz, 1H), 1.85 (ddd, $J=12.3$, 10.5, 8.0 Hz, 1H), 1.42 (d, $J=5.8$ Hz, 3H), 1.29 (s, 9H). LC-MS (Method A): m/z = 273.2 $[\text{M}+\text{H}]^+$, 1.11 min. e.e.= 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1 % isopropylamine) 70/30 % v/v, flow rate: 1 mL/min, retention time: 6.5 min.

4-[(3S,5R)-2-(2,2-dimethylpropanoyl)-5-methyl-1,2-oxazolidin-3-yl]benzonitrile (Second eluted diastereoisomer): ^1H NMR (400 MHz, CDCl_3) δ 7.63 (d, $J=8.4$ Hz, 2H), 7.38 (d, $J=8.2$ Hz, 2H), 5.39 (dd, $J=8.8$, 5.2 Hz, 1H), 4.61 (sextet, $J=6.0$ Hz, 1H), 2.44 (ddd, $J=12.3$, 8.9, 5.2 Hz, 1H), 2.29 (dt, $J=12.1$, 5.9 Hz, 1H), 1.36 (d, $J=6.2$ Hz, 3H), 1.29 (s, 9H). LC-MS (Method A): m/z = 273.2 $[\text{M}+\text{H}]^+$, 1.09 min.

e.e.= 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n~hexane/(ethanol + 0.1 % isopropylamine) 70/30 %v/v, flow rate:1 mL/min, retention time: 7.6 min.

EXAMPLE 47


3,3-DIFLUORO-1-[(3S)-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE
(101)



A solution of 3,3-difluoro-2,2-dimethylpropanoic acid (55 mg, 0.4 mmol), oxalyl chloride (0.034 mL, 0.4 mmol) and DMF (1 drop) in anhydrous CH_2Cl_2 (4 mL) was stirred at room temperature 1h 30 min, then a solution of (3S)-3-(4-fluorophenyl)-1,2-oxazolidine (60 mg, 0.36 mmol) and DIPEA (0.157 mL) 0.9 mmol) in anhydrous CH_2Cl_2 (3 mL) was added and the resulting mixture was stirred at room temperature for 1h. The mixture was diluted with CH_2Cl_2 and washed with HCl solution, sat. NaHCO_3 solution and brine. The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude product was purified by column chromatography (cyclohexane-EtOAc, 90:10 to 70:30) to give the title compound containing some residual 3,3-difluoro-2,2-dimethylpropanoic acid. This mixture was dissolved in CH_2Cl_2 and washed with 0.1N NaOH solution and brine. The organic phase was concentrated under reduced pressure to give the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.29-7.21 (m, 2H), 7.08-6.97 (m, 2H), 6.50-6.14 (m, 1H), 5.40 (dd, $J=8.8, 6.5$ Hz, 1H), 4.30 (dt, $J=3.3, 7.5$ Hz, 1H), 3.95 (ddd, $J=9.5, 8.0, 6.5$ Hz, 1H), 2.91-2.76 (m, 1H), 2.33 (dddd, $J=12.4, 9.6, 7.6, 6.5$ Hz, 1H), 1.40-1.32 (m, 6H). LC-MS (Method A): m/z = 288.2 $[\text{M}+\text{H}]^+$, 1.10 min.

EXAMPLES 48 AND 49

PREPARATION OF 1-[(3R)-3-(3,5-DIFLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE AND 1-[(3S)-3-(3,5-DIFLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (102) AND (103)




Hydrochloric acid solution (4M in **dioxane**, 3 mL, 12 mmol), was added at room temperature to a solution of tert-butyl 3-(3,5-difluorophenyl)-1,2-oxazolidine-2-carboxylate (50 mg, 0.17 mmol) in anhydrous CH_2Cl_2 (2 mL). The mixture was then stirred at room temperature for 2 h. The mixture was concentrated under reduced pressure. The residue was dissolved in CH_2Cl_2 (5 mL) and **pivaloyl** chloride (0.031 mL, 0.255 mmol) was added followed by sat. NaHCO_3 solution (5 mL). The mixture was stirred for 18 h then the organic portion was separated and washed with sat. NaHCO_3 solution, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the title compounds as a mixture of enantiomers. This mixture of enantiomers was resolved by chiral HPLC on Chiralpak AS-H (25 x 2.0 cm), 5 μm column using a mobile phase of **n-hexane/2-propanol 90/10 % v/v** and a flow rate of 18 mL/min to afford the two separated title compounds.

First eluting enantiomer, **1-[(3R)-3-(3,5-difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one**. ^1H NMR (400 MHz, CDCl_3) δ 6.83 (dd, $J=8.3, 2.0$ Hz, 2 H), 6.74-6.65 (m, 1 H), 5.38 (dd, $J=8.9, 6.7$ Hz, 1 H), 4.32-4.18 (m, 1 H), 3.95-3.82 (m, 1 H), 2.89-2.76 (m, 1 H), 2.33-2.22 (m, 1 H), 1.31 (s, 9 H). LC-MS (Method A): $m/z = 269.0$ $[\text{M}+\text{H}]^+$, 1.18 min. e.e.=100% as determined on a Chiralpak AS-H (25 x 0.46 cm) 5 μm column using a mobile phase of **n-hexane/2-propanol 90/10 % v/v**, flow rate: 18 mL/min, retention time: 4.2 min.

Second eluting enantiomer, **1-[(3S)-3-(3,5-difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one**. ^1H NMR (400 MHz, CDCl_3) δ 6.79-6.89 (m, 2 H), 6.69 (tt, $J=8.9, 2.4$ Hz, 1 H), 5.38 (dd, $J=8.8, 6.5$ Hz, 1 H), 4.26 (td, $J=7.9, 2.8$ Hz, 1 H), 3.89 (ddd, $J=9.8, 8.0, 6.5$ Hz, 1 H), 2.83 (dddd, $J=12.2, 9.2, 6.5, 2.8$ Hz, 1 H), 2.27 (dddd, $J=12.3, 9.8, 7.6, 6.7$ Hz, 1 H), 1.31 (s, 9 H). LC-MS (Method A): $m/z = 269.0$ $[\text{M}+\text{H}]^+$, 1.18 min. e.e.=99 % as determined on a Chiralpak AS-H (25 x 0.46 cm) 5 μm column using a mobile phase of **n-hexane/2-propanol 90/10 % v/v**, flow rate: 18 mL/min, retention time: 5.5 min.

EXAMPLE 50

PREPARATION OF 3,3-DIMETHYL-4-[(3S)-3-(5-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDIN-2-YL]-4-OXOBUTANENITRILE (104)




Thionyl chloride (0.029 mL, 0.39 mmol) was added to a solution of 3-cyano-2,2-dimethylpropanoic acid (45 mg, 0.35 mmol) and DMF in CH_2Cl_2 (2 mL). The reaction was stirred at room temperature for 2 h. A solution of tert-butyl (3S)-3-[4-(difluoromethyl)phenyl]-1,2-oxazolidine-2-

carboxylate (45 mg, 0.35 mmol) and Et₃N (0.13 mL, 1.08 mmol) in CH₂Cl₂ was added to the reaction and stirring was continued at room temperature for 1 h. The reaction was diluted with CH₂Cl₂ and washed with sat. NaHCO₃ solution. The organic phase was concentrated under reduced pressure and the crude product was purified by column chromatography (cyclohexane-EtOAc, 100:0 to 0:100) to afford the title compound as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiraipak IC (25 x 2.0 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1%isopropylamine) 50/50 % v/v and a flow rate of 18 mL/min to afford the title compound. ¹H NMR (400MHz, CDCl₃) δ 8.53 (d, J=1.3 Hz, 1H), 8.42 (d, J=0.8 Hz, 1H), 5.51 (dd, J=8.8, 6.0 Hz, 1H), 4.41 (dt, J=4.0, 7.5 Hz, 1H), 4.15-3.95 (m, 1H), 2.88-2.76 (in. 1H), 2.73-2.62 (m, 3H), 2.57 (s, 3H), 1.49 (s, 3H), 1.47 (s, 3H). EC-MS (Method A): m/z = 275.3 [M-H]⁺, 0.68 min. e.e.=97.3% as determined on a Chiraipak IC (25 x 0.46 cm) 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1%isopropylamine) 50/50 % v/v, flow rate: 1 mL/min, retention time: 11.2 min.

EXAMPLE 51


PREPARATION OF 1-[(3S)-3-[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (105)



To a solution of test-butyl (3S)-3-[6-(difluoromethyl)pyridine -3-yl]-1,2-oxazolidine -2-carboxylate (24 mg, 0.11 mmol) in CH₂Cl₂ (2 mL) were added F³⁴N (0.031 mL, 0.22 mmol) and pivaloylchloride (0.015 mL, 0.12 mmol). The mixture was stirred for 2 h. The volatiles were removed under reduced pressure to afford the title compound as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiraipak AD-H (25 x 2 cm) 5 μm column using a mobile phase of n-hexane/ethanol 70/30 % v/v and a flow rate of 18 mL/min to afford then purified further by column chromatography (cyclohexane-EtOAc, 100:0 to 50:50) to afford the title compound. ¹H NMR (400MHz, CDCl₃) δ 8.62 (d, J=2.0 Hz, 1H), 7.77 (dd, J=8.0, 2.3 Hz, 1H), 7.62 (d, J=8.0 Hz, 1H), 6.79-6.47 (m, 1H), 5.48 (dd, J=8.5, 7.0 Hz, 1H), 4.32 (dt, J=2.3, 7.9 Hz, 1H), 3.94 (ddd, J=10.0, 8.2, 6.4 Hz, 1H), 2.96-2.85 (m, 1H), 2.38-2.26 (m, 1H), 1.30 (s, 9H). LC-MS (Method A): m/z = 285.93 [M-H]⁺, 0.97 min. e.e.=100% as determined on a Chiraipak AD-H (25 x 0.46 cm) 5 μm column using a mobile phase of n-hexane/ethanol 60/40 % v/v, flow rate: 1 mL/min, retention time: 7.2 min.

EXAMPLE 52


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(1-METHYL-1H-PYRAZOL-4-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (106)



5 To a stirred mixture of (3S)-3-(1-methyl-1H-pyrazol-4-yl)-1,2-oxazolidine hydrochloride (90 mg, 0.47 mmol) and Et₃N (143 mg, 1.42 mmol) in CH₂Cl₂ (10 mL) was added dropwise pivaloyl chloride (57 mg, 0.47 mmol) at 0 °C. The resulting mixture was warmed to room temperature and stirred for 1 hour. The reaction mixture was quenched with water (15 mL) and extracted with CH₂Cl₂ (2×15 mL). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following conditions: column: XBndge Prep C18 Column, 150×19⁴mm, 5 μm; Mobile Phase A: water (10 mmol/L ammonium bicarbonate); Mobile Phase B: acetonitrile (25% to 75% over 7 min); Detector, UV 220 & 254 nm to afford the title compound (35 mg, 31%) as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak IC (25 x 2 cm), 5 μm column using a mobile phase of hexane/ethanol 15 70/30% v/v and a flow rate of 20 mL/min to afford the title compound. First eluting enantiomer. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.57 (s, 1H), 7.31 (s, 1H), 5.26-5.22 (m, 1H), 4.25-4.20 (m, 1H), 3.82-3.76 (m, 4H), 2.70-2.60 (m, 1H), 2.30-2.22 (m, 1H), 1.15 (s, 9H). LC-MS (Method F): m/z = 238.0 [M+H]⁺, 1.00 min. e.e.=99.9% as determined on a Chiralpak IC-3 (10 x 0.46 cm), 3 μm column using a mobile phase of (n-hexane + 0.1% trifluoroacetic acid)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 5.58 20 min.

EXAMPLE 53

PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(6-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (107)




25 To a solution of N-hydroxy-N-[(1S)-3-hydroxy-1-(6-methylpyrazin-2-yl)propyl]-2,2-dimethylpropanamide (170 mg, 0.64 mmol) and triphenylphosphine (566 mg, 2.10 mmol) in anhydrous

THF (10 mL) was added DIAD (387 mg, 1.90 mmol) dropwise at room temperature. After being stirred overnight at room temperature, the reaction mixture was concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following conditions: column: XBridge Prep C18 Column, 150x19mm, 5 μ m; Mobile Phase A: Water (10 mmol/L NH_4HCO_3); Mobile Phase B: Acetonitrile (20% to 50% over 7 min); Detector, UV 220 & 254 nm to afford the title compound as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Phenomenex Lux 5u Cellulose-4AXLA Packed (25 x 2.12 cm), 5 μ m column using a mobile phase of hexane/ethanol 95/5% v/v and a flow rate of 20 mL/min to afford the title compound. First eluted enantiomer. ^1H NMR (400 MHz, CD_3OD) δ 8.39 (s, 1H), 8.34 (s, 1H), 5.44-5.40 (m, 1H), 4.45-4.31 (m, 1H), 4.07-4.01 (m, 1H), 2.99-2.78 (m, 1H), 2.68-2.43 (m, 4H), 1.27 (s, 9H). LC-MS (Method N): m/z -250.2 $[\text{M}+\text{H}]^+$, 1.12 min. e.e.=99.9% as determined on a Lux Cellulose-4 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 95/5 % v/v, flow rate: 1.0 mL/min, retention time: 3.23 min.

EXAMPLE 54


PREPARATION OF 5-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]PYRIDINE-3-CARBONITRILE (108)



To a stirred mixture of 5-[(3S)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile trifluoroacetate (105 mg, 0.36 mmol) and Et_3N (163 mg, 1.61 mmol) in CH_2Cl_2 (10 mL) was added pivaloylchloride (44 mg, 0.36 mmol) at 0 $^\circ\text{C}$. After being stirred at room temperature for 1 hour, the reaction mixture was quenched by the addition of water (15 mL) and extracted with CH_2Cl_2 (3 x 15 mL). The combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column, 5 μ m, 150 x 19 mm; Mobile phase: Phase A: Water (10 mmol/L ammonium bicarbonate); Phase B: Acetonitrile (25% up to 55% in 7 min); Detector, UV 254 & 220 nm to afford the title compound. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.92 (d, J = 1.8 Hz, 1H), 8.73 (d, J = 2.1 Hz, 1H), 8.16-8.15 (m, 1H), 5.39 (t, J = 7.2 Hz, 1H), 4.35-4.29 (m, 1H), 3.92-3.84 (m, 1H), 2.90-2.85 (m, 1H), 2.28-2.21 (m, 1H), 1.19 (s, 9H). LC-MS (Method H): m/z = 260.0 $[\text{M}+\text{H}]^+$, 1.22 min. e.e.=95.9% as determined on a Chiralpak IA-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 1.94 min.

EXAMPLE 55


PREPARATION OF 3-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-5-
FLUOROBENZONITRILE (109)



To a stirred solution of N-[(1S)-1-(3-cyano-5-fluorophenyl)-3-hydroxypropyl]-N-hydroxy-2,2-dimethylpropanamide (100 mg, 0.34 mmol) and triphenylphosphine (213 mg, 0.81 mmol) in THF (10 mL) was added dropwise DIAD (137 mg, 0.68 mmol) at 0 °C. After being stirred overnight at room temperature under a nitrogen atmosphere, the reaction mixture was concentrated under reduced pressure. The crude product was purified by preparative thin layer chromatography (EtOAc-petroleum ether, 1:3) to afford the title compound as a mixture of enantiomers. This mixture was resolved by chiral HPLC on a Chiralpak IC (25 x 2 cm) 5 μ m column using a mobile phase of hexane/ethanol 80/20% v/v and a flow rate of 20 mL/min to afford the title compound. Second eluting enantiomer. ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.79-7.75 (m, 1H), 7.58 (s, 1H), 7.46-7.41 (m, 1H), 5.37 (t, *J*=7.2 Hz, 1H), 4.32-4.27 (m, 1H), 3.90-3.82 (m, 1H), 2.90-2.86 (m, 1H), 2.22-2.14 (m, 1H), 1.21 (s, 9H). LC-MS (Method H): *m/z* = 276.9 [M+H]⁺, 1.56 min. e.e.-99.9% as determined on a Chiralpak IC-3 (10 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 3.54 min.

EXAMPLE 56

PREPARATION OF 1-[(3S)-3-{IMIDAZO[1,2-a]PYRIDIN-7-YL}-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (110)




The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1+0.1% isopropylamine) 40/60% v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.09 (dd, *J*=7.0, 0.8 Hz, 1H), 7.62 (d, *J*=1.3 Hz,

1H), 7.57-7.53 (m, 2H), 6.77 (dd, $J=7.0$, 1.8 Hz, 1H), 5.45 (dd, $J=9.0$, 6.5 Hz, 1H), 4.33-4.26 (m, 1H), 3.93 (ddd, $J=9.6$, 8.1, 6.7 Hz, 1H), 2.85 (dddd, $J=12.1$, 9.1, 6.5, 2.9 Hz, 1H), 2.34 (dddd, $J=12.3$, 9.7, 7.7, 6.7 Hz, 1H), 1.31 (s, 9H). LC-MS (Method A): $m/z = 274.2$ $[M+H]^+$, 0.43 min. e.e.= 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1+0.1% isopropylamine) 40/60% v/v, flow rate: 1 mL/min, retention time: 5.0 min.

EXAMPLE 57


PREPARATION OF 1-[(3S)-3-{IMIDAZO[1,2-A]PYRIDIN-2-YL}-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (111)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 40/60% v/v, flow rate: 18 mL/min. ^1H NMR (400 MHz, CDCl_3) δ 8.04 (dt, $J=6.8$, 1.2 Hz, 1H), 7.61-7.54 (m, 2H), 7.15 (ddd, $J=9.2$, 6.8, 1.3 Hz, 1H), 6.76 (td, $J=6.8$, 1.0 Hz, 1H), 5.69 (dd, $J=8.9$, 5.9 Hz, 1H), 4.38-4.30 (m, 1H), 3.96 (ddd, $J=9.2$, 7.8, 6.9 Hz, 1H), 2.99-2.86 (m, 1H), 2.84-2.74 (m, 1H), 1.32 (s, 9H). LC-MS (Method A): $m/z = 274.2$ $[M+H]^+$, 0.46 min. e.e.= 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 40/60% v/v, flow rate: 1 mL/min, retention time: 5.5 min.

EXAMPLES 58 AND 59

PREPARATION OF 5-[(3S)-2-(2,2-DIFLUORO-1-METHYLCYCLOPROPANECARBONYL)-1,2-OXAZOLIDIN-3-YL]-2-FLUOROPYRIDINES (112) AND (113)



The title compounds ((S)-5-[(3S)-2-(2,2-difluoro-1-methylcyclopropanecarbonyl)-1,2-oxazolidin-3-yl]-2-fluoropyridine and (R)-5-[(3S)-2-(2,2-difluoro-1-methylcyclopropanecarbonyl)-1,2-oxazolidin-3-yl]-2-fluoropyridine) were isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30% v/v, flow rate: 18 mL/min.

First eluting diastereoisomer, Diastereoisomer i. ^1H NMR (400 MHz, CDCl_3) δ 8.18 (d, $J=2.5$

Hz, 1H), 7.70 (td, $J=8.0, 2.6$ Hz, 1H), 6.92 (dd, $J=8.5, 3.0$ Hz, 1H), 5.42 (dd, $J=8.8, 5.5$ Hz, 1H), 4.41-4.31 (m, 1H), 4.08 (td, $J=8.4, 6.8$ Hz, 1H), 2.97 (dddd, $J=12.4, 8.9, 6.8, 4.0$ Hz, 1H), 2.47-2.32 (m, 1H), 1.90 (ddd, $J=13.7, 8.6, 4.5$ Hz, 1H), 1.50 (dd, $J=2.8, 1.3$ Hz, 3H), 1.26 (ddd, $J=11.4, 8.3, 5.1$ Hz, 1H).

LC-MS (Method A): $m/z = 287.1 [M+H]^+$, 0.84 min. Chiralpak AD-H (25 x 0.46 cm), 5 μ m column


5 using a mobile phase of n-hexane/ethanol 70/30% v/v, flow rate: 1 mL/min, retention time: 10.8 min.

Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400 MHz, CDCl_3) δ 8.25-8.18 (m, 1H), 7.76 (td, $J=7.8, 2.4$ Hz, 1H), 6.94 (dd, $J=8.7, 2.9$ Hz, 1H), 5.43 (dd, $J=8.5, 6.8$ Hz, 1H), 4.34 (td, $J=7.8, 3.0$ Hz, 1H), 4.02 (ddd, $J=9.9, 8.2, 6.5$ Hz, 1H), 2.93-2.83 (m, 1H), 2.37 (ddt, $J=12.4, 9.9, 7.2$ Hz, 1H), 2.04 (ddd, $J=14.2, 8.3, 4.6$ Hz, 1H), 1.59 (dd, $J=3.0, 1.5$ Hz, 3H), 1.24 (ddd, $J=11.4, 8.3, 5.1$ Hz,

10 1H). LC-MS (Method A): $m/z = 287.1 [M+H]^+$, 0.90 min. Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30% v/v, flow rate: 1 mL/min, retention time: 12.1 min.

EXAMPLES 60 AND 61

PREPARATION OF (3S)-2-(2,2-DIFLUORO-1-METHYLCYCLOPROPANECARBONYL)-3-(4-FLUOROPHENYL)-1,2-OXAZOLIDINES (114) AND (115)




15 The title compounds ((3S)-2-((R)-2,2-difluoro-1-methylcyclopropanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine and (3S)-2-((S)-2,2-difluoro-1-methylcyclopropanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine) were isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min.

20 First eluting diastereoisomer, Diastereoisomer 1. ^1H NMR (400 MHz, CDCl_3) δ 7.34-7.26 (m, 2H), 7.00-7.09 (m, 2H), 5.38 (dd, $J=8.5, 6.5$ Hz, 1H), 4.30 (td, $J=7.8, 3.0$ Hz, 1H), 4.04-3.94 (m, 1H), 2.87-2.78 (m, 1H), 2.35 (ddt, $J=12.2, 9.6, 7.2$ Hz, 1H), 2.03 (ddd, $J=14.1, 8.3, 4.8$ Hz, 1H), 1.60 (dd, $J=3.0, 1.5$ Hz, 3H), 1.21 (ddd, $J=11.5, 8.3, 5.0$ Hz, 1H). LC-MS (Method A): $m/z = 286.1 [M+H]^+$, 1.06 min. Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min, retention time: 4.9 min.

25 Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400 MHz, CDCl_3) δ 7.29-7.21 (m, 2H), 7.06-6.98 (m, 2H), 5.37 (dd, $J=8.8, 5.3$ Hz, 1H), 4.32 (td, $J=7.7, 4.5$ Hz, 1H), 4.05 (td, $J=8.2, 7.0$ Hz, 1H), 2.96-2.85 (m, 1H), 2.42-2.31 (m, 1H), 1.97-1.85 (m, 1H), 1.48 (br. s., 3H), 1.29-1.19 (m, 1H). LC-MS (Method A): $m/z = 286.1 [M+H]^+$, 1.00 min. Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min, retention time: 6.6 min.

EXAMPLES 62 AND 63

PREPARATION OF 2-CYCLOPROPYL-1-[(3S)-3-(6-FLUOROPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONES (116) AND (117)




The title compounds ((R)-2-cyclopropyl-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]propan-1-one and (S)-2-cyclopropyl-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]propan-1-one) were isolated by chiral HPLC on a Chiraipak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 70/30% v/v, flow rate: 18 mL/min.

First eluting diastereoisomer, Diastereoisomer 1. ^1H NMR (400 MHz, CDCl₃) δ 8.23-8.20 (m, 1H), 7.76 (td, $J=8.0, 2.5$ Hz, 1H), 6.93 (dd, $J=8.4, 2.9$ Hz, 1H), 5.49 (dd, $J=8.3, 6.3$ Hz, 1H), 4.26 (td, $J=7.8, 3.5$ Hz, 1H), 3.88 (ddd, $J=9.3, 8.2, 6.8$ Hz, 1H), 2.94-2.81 (m, 1H), 2.42-2.26 (m, 2H), 1.21 (d, $J=6.8$ Hz, 3H), 1.09-0.97 (m, 1H), 0.61-0.43 (m, 2H), 0.43-0.34 (m, 1H), 0.19 (dt, $J=9.5, 4.7$ Hz, 1H). LC-MS (Method A): $m/z = 265.2$ [$M+H$]⁺, 0.90 min. Chiraipak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 70/30% v/v, flow rate: 1 mL/min, retention time: 6.5 min.


Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400 MHz, CDCl₃) δ 8.20 (d, $J=2.5$ Hz, 1H), 7.75 (td, $J=8.0, 2.5$ Hz, 1H), 6.92 (dd, $J=8.5, 2.8$ Hz, 1H), 5.49 (dd, $J=9.0, 6.3$ Hz, 1H), 4.28 (td, $J=8.0, 2.6$ Hz, 1H), 3.88 (ddd, $J=9.6, 8.1, 6.7$ Hz, 1H), 2.89 (dddd, $J=12.3, 9.2, 6.3, 3.0$ Hz, 1H), 2.35 (dddd, $J=12.3, 9.6, 7.7, 6.3$ Hz, 1H), 2.30-2.19 (m, 1H), 1.28 (d, $J=7.0$ Hz, 3H), 1.04 (dt, $J=9.8, 8.1, 5.0$ Hz, 1H), 0.61-0.47 (m, 2H), 0.25-0.11 (m, 2H). LC-MS (Method A): $m/z = 265.2$ [$M+H$]⁺, 0.89 min. Chiraipak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 70/30% v/v, flow rate: 1 mL/min, retention time: 8.0 min.

EXAMPLE 64


PREPARATION OF TERT-BUTYL (3S)-3-(5-CHLOROPYRIDIN-3-YL)-1,2-OXAZOLIDINE-2-CARBOXYLATE (118)



The title compound was isolated by chiral SFC on a Chiralpak AD-H (25 x 2.0 cm) 5 μ m column using a mobile phase of 7% isopropanol in CO₂, flow rate: 45 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.49 (d, J =2.5 Hz, 1H), 8.47 (d, J =2.0 Hz, 1H), 7.77–7.70 (m, 1H), 5.26 (dd, J =8.8, 5.5 Hz, 1H), 4.27–4.18 (m, 1H), 3.91 (ddd, J =9.0, 8.3, 7.0 Hz, 1H), 2.85 (dddd, J =12.3, 8.9, 7.0, 3.4 Hz, 1H), 2.29 (dddd, J =12.3, 9.1, 7.8, 5.6 Hz, 1H), 1.49 (s, 9H). SFC on a Chiralpak AD-H (25 x 0.46 cm) 5 μ m column using a mobile phase of 7% isopropanol in CO₂, flow rate: 2.5 mL/min, retention time: 8.4 min.

EXAMPLE 65**PREPARATION OF TERT-BUTYL (3S)-3-[2-(DIFLUOROMETHYL)PYRIDIN-4-YL]-1,2-OXAZOLIDINE-2-CARBOXYLATE (119)**

The title compound was isolated by chiral HPLC on a Chiralpak IA (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 75/25 v/v, flow rate: 18 mL/min. ¹H NMR (400MHz, CDCl₃) δ 8.63 (d, J =5.0 Hz, 1H), 7.65 (s, 1H), 7.44 (dd, J =5.0, 0.8 Hz, 1H), 6.66 (t, J =55.0 Hz, 1H), 5.29 (dd, J =8.9, 5.6 Hz, 1H), 4.21 (dt, J =3.3, 7.9 Hz, 1H), 3.92 (dt, J =7.0, 8.7 Hz, 1H), 2.88 (dddd, J =12.3, 8.9, 7.0, 3.4 Hz, 1H), 2.28 (dddd, J =12.3, 9.0, 7.7, 5.6 Hz, 1H), 1.50 (s, 9H). LC-MS (Method A): m/z = 301.1 [M+H]⁺, 0.93 min. Chiralpak IA (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 75/25 v/v, flow rate: 1 mL/min, retention time: 6.7 min.

EXAMPLE 66**PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-[6-(TRIFLUOROMETHYL)PYRIDIN-3-YL]-1,2-OXAZOLIDINE-2-YL]PROPAN-1-ONE (120)**


The title compound was isolated by chiral SFC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of 5% isopropanol in CO₂, flow rate: 45 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.70 (d, J =2.0 Hz, 1H), 7.79 (dd, J =8.3, 2.0 Hz, 1H), 7.67 (d, J =8.3 Hz, 1H), 5.50 (dd, J =9.0,

7.0 Hz, 1H), 4.33 (td, $J=8.3, 2.0$ Hz, 1H), 4.0-3.91 (m, 1H), 2.96-2.87 (m, 1H), 2.37-2.27 (m, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 303.2$ $[M+H]^+$, 1.07 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of 5% isopropanol in CO_2 , flow rate: 2.5 mL/min, retention time: 7.6 min.

5

EXAMPLE 67

PREPARATION OF 1-[(3S)-3-(5-CHLORO-3-FLUOROPYRIDIN-2-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (121)




The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 75/25 % v/v, flow rate: 18 mL/min. ^1H NMR (400 MHz, $CDCl_3$) δ 8.38 (d, $J=2.0$ Hz, 1H), 7.43 (dd, $J=9.0, 2.0$ Hz, 1H), 5.67 (dd, $J=8.8, 6.8$ Hz, 1H), 4.44 (td, $J=8.0, 2.5$ Hz, 1H), 4.04-3.97 (m, 1H), 2.80-2.70 (m, 1H), 2.62-2.52 (m, 1H), 1.26 (s, 9H). LC-MS (Method A): $m/z = 287.2$ $[M+H]^+$, 1.04 min. e.e.= 100% as determined on a Chiralpak IC (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 75/25 % v/v, flow rate: 1 mL/min, retention time: 6.4 min.

15

EXAMPLE 68

PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-[5-(TRIFLUOROMETHYL)PYRIDIN-2-YL]-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (122)




20

The title compound was isolated by chiral SFC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of 6% isopropanol in CO_2 , flow rate: 45 mL/min. ^1H NMR (400 MHz, $CDCl_3$) δ 8.84 (s, 1H), 7.91 (dd, $J=8.3, 2.0$ Hz, 1H), 7.48 (d, $J=8.3$ Hz, 1H), 5.57 (dd, $J=9.0, 6.3$ Hz, 1H), 4.31 (td, $J=7.0, 3.2$ Hz, 1H), 4.0-3.91 (m, 1H), 2.90-2.81 (m, 1H), 2.70-2.60 (m, 1H), 1.32 (s, 9H). LC-MS (Method A): $m/z = 303.2$ $[M+H]^+$, 1.08 min. e.e.=100% as determined on a Chiralpak AD-H (25 x

0.46 cm), 5 μ m column using a mobile phase of 5% isopropanol in CO₂, flow rate: 2.5 mL/min, retention time: 7.7 min.

EXAMPLE 69


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-[4-(TRIFLUOROMETHYL)-1,3-TETRAZOL-2-YL]-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (123)



The title compound was isolated by chiral HPLC on a Whelk O-1 (R,R) (25 x 2.0 cm), 10 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 80/20 % v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 7.72 (s, 1H), 5.78 (dd, J =8.8, 5.5 Hz, 1H), 4.31 (td, J =7.8, 4.0 Hz, 1H), 3.98 (q, J =7.8 Hz, 1H), 3.0-2.82 (m, 2H), 1.33 (s, 9H). LC-MS (Method A): m/z = 309.2 [M+H]⁺, 1.13 min. e.e.=100% as determined on a Whelk O-1 (R,R) (25 x 0.46 cm), 10 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 80/20 % v/v, flow rate: 1 mL/min, retention time: 5.2 min.

EXAMPLE 70


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-{PYRAZOLO[1,5-A]PYRIDIN-2-YL}-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (124)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 65/35 % v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.40 (dd, J =6.0, 1.0 Hz, 1H), 7.46 (dt, J =8.8, 1.0 Hz, 1H), 7.08 (ddd, J =7.8, 6.8, 1.0 Hz, 1H), 6.7 (td, J =7.0, 1.5 Hz, 1H), 6.49 (s, 1H), 5.78 (dd, J =7.8, 6.5 Hz, 1H), 4.36-4.30 (m, 1H), 3.97 (q, J =7.5 Hz, 1H), 2.84-2.77 (m, 2H), 1.33 (s, 9H). LC-MS (Method A): m/z = 274.2 [M+H]⁺, 0.97 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 65/35 % v/v, flow rate: 1 mL/min, retention time: 7.0 min.

EXAMPLE 71


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(2-METHYLPYRIDIN-5-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (125)



5 The title compound was isolated by chiral HPLC on a Chiraipak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60 % v/v, flow rate: 18 mL/min. ^1H NMR (400 MHz, CDCl_3) δ 8.59 (s, 2H), 5.40 (dd, $J=6.8, 8.8$ Hz, 1H), 4.32 (dt, $J=2.8, 7.9$ Hz, 1H), 3.94 (ddd, $J=6.5, 8.2, 9.9$ Hz, 1H), 2.87 (dddd, $J=2.8, 6.4, 9.2, 12.3$ Hz, 1H), 2.75 - 2.69 (m, 3H), 2.32 (tdd, $J=7.3, 9.7, 12.2$ Hz, 1H), 1.32 - 1.23 (m, 9H). LC-MS (Method A): m/z = 250.2 [M+H] $^+$, 0.75 min. e.e.=100% as determined on a Chiraipak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60 v/v, flow rate: 1 mL/min, retention time: 12.0 min.

EXAMPLE 72


PREPARATION OF 3-(2,2-DIMETHYL-1-[(3S)-3-(2-METHYLPYRIDIN-5-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (126)



20 The title compound was isolated by chiral HPLC on a Chiraipak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60 % v/v, flow rate: 18 mL/min. ^1H NMR (400 MHz, CDCl_3) δ 8.58 (s, 2H), 6.48-6.09 (m, 1H), 5.40 (dd, $J=8.8, 6.8$ Hz, 1H), 4.36 (dt, $J=2.8, 8.0$ Hz, 1H), 4.00 (ddd, $J=9.7, 8.2, 6.3$ Hz, 1H), 2.91 (dddd, $J=12.3, 9.2, 6.4, 3.0$ Hz, 1H), 2.78-2.68 (m, 3H), 2.44-2.29 (m, 1H), 1.37 (s, 3H), 1.36 (s, 3H). LC-MS (Method A): m/z = 286.2 [M+H] $^+$, 0.77 min. e.e.=100% as determined on a Chiraipak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60 v/v, flow rate: 1 mL/min, retention time: 14.7 min.

EXAMPLE 73


PREPARATION OF 1-[(3S)-3-(5-FLUOROPYRIDIN-2-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (127)



5 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 85/15 % v/v, flow rate: 18 mL/min. ^1H NMR (400 MHz, CDCl_3) δ 8.46-8.38 (m, 1H), 7.40-7.33 (m, 2H), 5.51 (dd, $J=8.9, 6.1$ Hz, 1H), 4.33-4.25 (m, 1H), 3.93 (ddd, $J=9.4, 7.9, 6.8$ Hz, 1H), 2.83-2.59 (m, 2H), 1.30 (s, 9H). LC-MS (Method A): m/z = 253.2 $[\text{M}+\text{H}]^+$, 0.91 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a
10 mobile phase of n-hexane/ethanol 85/15 % v/v, flow rate: 1 mL/min, retention time: 8.0 min.

EXAMPLE 74


PREPARATION OF 1-[(3S)-3-[4-(DIFLUOROMETHYL)PHENYL]-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (128)



15 The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 85/15 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHZ, CDCl_3) δ 7.52- 7.45 (m, 2H), 7.43-7.36 (m, 2H), 6.79-6.46 (m, 1H), 5.45 (dd, $J=8.7, 6.9$ Hz, 1H), 4.28 (dt, $J=2.6, 7.8$ Hz, 1H), 3.91 (ddd, $J=10.0, 8.0, 6.4$ Hz, 1H), 2.84 (dddd, $J=12.2, 9.1, 6.3, 2.8$ Hz, 1H), 2.30 (dddd, $J=12.2, 9.9, 7.5, 6.8$ Hz, 1H), 1.31 (s, 9H). LC-MS (Method A): m/z =
20 284.2 $[\text{M}+\text{H}]^+$, 1.13 min. e.e.=100% as determined on a Chiralpak IC (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 85/15 % v/v, flow rate: 1.0 mL/min, retention time: 7.4 min.

EXAMPLE 75

PREPARATION OF 3,3,3-TRIFLUORO-2-METHYL-1-[(3S)-3-{5-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (129)




DIASTEREOISOMER 1

5 The title compound was isolated by chiral HPLC on a Chiralcel OJ-H (25 x 2 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 75/25 % v/v, and a flow rate of 18 mL/min. First eluting diastereoisomer, Diastereoisomer 1. ^1H NMR (400MHz, CDCl_3) δ 8.55 (d, $J=1.5$ Hz, 1H), 8.43 (d, $J=1.0$ Hz, 1H), 5.54 (dd, $J=8.5, 6.3$ Hz, 1H), 4.38 (dt, $J=3.9, 7.5$ Hz, 1H), 4.11-3.99 (m, 1H), 3.89 (dd, $J=8.0, 7.3$ Hz, 1H), 2.93-2.69 (m, 2H), 2.58 (s, 3H), 1.41 (d, $J=7.0$ Hz, 3H). LC-MS (Method A): $m/z = 290.2$ $[\text{M}+\text{H}]^+$, 0.80 min. e.e.=100% as determined on a Chiralcel OJ-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 75/25 % v/v, flow rate: 1 mL/min, retention time: 6.1 min.

EXAMPLE 76

PREPARATION OF 3,3,3-TRIFLUORO-2-METHYL-1-[(3S)-3-{5-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (130)




DIASTEREOISOMER 2

20 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v and a flow rate of 18 mL/min. Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400MHz, CDCl_3) δ 8.54 (d, $J=1.3$ Hz, 1H), 8.42 (s, 1H), 5.51 (dd, $J=8.8, 5.5$ Hz, 1H), 4.38 (dt, $J=7.7, 4.8$ Hz, 1H), 4.09-4.00 (m, 1H), 3.86 (td, $J=7.5, 15.4$ Hz, 1H), 2.91-2.81 (m, 1H), 2.80-2.70 (m, 1H), 2.58 (s, 3H), 1.41 (d, $J=7.3$ Hz, 3H). LC-MS (Method A): $m/z = 290.2$ $[\text{M}+\text{H}]^+$, 0.79 min. e.e.=100% as determined on a Chiralpak AD-

H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1 mL/min, retention time: 10.9 min.

EXAMPLE 77


PREPARATION OF 4-[(3S)-3-[6-(DIFLUOROMETHYL)PYRIDIN-3-YL]-1,2-OXAZOLIDIN-2-YL]-3,3-DIMETHYL-4-OXOBUTANENITRILE (131)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60% v/v and a flow rate of 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.62 (d, $J=1.8$ Hz, 1H), 7.77 (ud, $J=8.2$, 2.1 Hz, 1H), 7.64 (d, $J=8.3$ Hz, 1H), 6.80–6.47 (m, 1H), 5.48 (dd, $J=8.5$, 7.0 Hz, 1H), 4.39 (dt $J=2.9$, 7.8 Hz, 1H), 4.02 (ddd, $J=9.7$, 8.2, 6.4 Hz, 1H), 2.95 (dddd, $J=12.3$, 9.2, 6.3, 2.9 Hz, 1H), 2.78–2.62 (m, 2H), 2.37 (tdd, $J=12.1$, 9.5, 7.3 Hz, 1H), 1.51–1.49 (m, 3H), 1.49–1.46 (m, 3H). LC-MS (Method A): m/z = 310.23 $[\text{M}+\text{H}]^+$, 0.85 min. e.e.=95% as determined on a Chiralpak AD-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 40/60% v/v, flow rate: 1 mL/min, retention time: 24.1 min.

EXAMPLE 78

PREPARATION OF 2,2-D(METHYL-1-[(3S)-3-{PYRAZOLO[5,4-B]PYRIDIN-5-YL}-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (132)




The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 50/50% v/v and a flow rate of 18 mL/min. ^1H NMR (400 MHz, CDCl_3) δ 8.43 (d, $J=7.3$ Hz, 1H), 7.93 (d, $J=2.3$ Hz, 1H),

1 H), 7.50-7.40 (m, 1 H), 6.70 (dd, $J=13$, 1.8 Hz, 1 H), 6.47 (dd, $J=23$, 0.8 Hz, 1 H), 5.42 (dd, $J=8.8$, 6.8 Hz, 1 H), 4.30 (td, $J=7.7$, 2.8 Hz, 1 H), 3.91 (ddd, $J=10.0$, 8.1, 6.5 Hz, 1 H), 2.91-2.75 (tn, 1 H), 2.34 (dddd, $J=12.3$, 9.9, 7.6, 6.8 Hz, 1 H), 1.32 (s, 9 H). LC-MS (Method A): m/z = 274.2 $[M+H]^+$, 0.93 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1 % isopropylamine) 50/50% v/v, flow rate: 1 mL/min, retention time: 12.9 min.

EXAMPLE 79


PREPARATION OF 1-[(3S)-3-(5-FLUOROPYRTOIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (133)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm) 5 μ m column using a mobile phase of n-hexane/ethanol 70/30% v/v and a flow rate of 18 mL/min. ^1H NMR (400 MHz, $CDCl_3$) δ 8.42 (t, $J=1.5$ Hz, 1 H), 8.38 (d, $J=2.8$ Hz, 1 H), 7.34 (dt, $J=9.2$, 2.1 Hz, 1 H), 5.47 (dd, $J=8.9$, 6.7 Hz, 1 H), 4.34-4.27 (m, 1 H), 3.92 (ddd, $J=10.0$, 8.1, 6.5 Hz, 1 H), 2.94-2.83 (m, 1 H) 2.32 (dddd, $J=12.4$, 9.9, 7.7, 6.7 Hz, 1 H) 1.30 (s, 9 H). LC-MS (Method B): m/z = 252.0 $[M+H]^+$, 0.90 min. e.e.=98.2% as determined on a Chiralpak AD-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/ethanol 70/30% v/v, flow rate: 1 mL/min, retention time: 7.7 min.

EXAMPLES 80 AND 81

PREPARATION OF 1-[(3R)-3-(2,4-DIFLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE AND 1-[(3S)-3-(2,4-DIFLUOROPHENYL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (134) AND (135)



The title compounds were isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/2-propanol 90/10% v/v, flow rate: 17 mL/min.


1-[(3R)-3-(2,4-difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one. ^1H NMR (400 MHz, $CDCl_3$) δ 7.24 (td, $J=8.6$, 6.4 Hz, 1 H), 6.89-6.76 (m, 2 H), 5.61 (dd, $J=8.7$, 6.7 Hz, 1 H), 4.25 (td,

$J=1.7$, 2.6 Hz, 1 H), 3.90 (ddd, $J=9.8$, 8.0, 6.5 Hz, 1 H), 2.94-2.80 (m, 1 H), 2.34-2.16 (m, 1 H), 1.31 (s, 9 H). LC-MS (Method B): $m/z = 269.0$ $[M+H]^+$, 1.16 min. e.e.=95.8% as determined on a Chiralpak IC (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/2-propanol 90/10% v/v, flow rate: 1 mL/min, retention time: 6.4 min.

5 1-[(3S)-3-(2,4-difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one. ^1H NMR (400 MHz, $CDCl_3$) δ 7.27-7.20 (m, 1 H), 6.88-6.77 (m, 2 H), 5.62 (dd, $J=8.7$, 6.7 Hz, 1 H), 4.25 (td, $J=7.7$, 2.5 Hz, 1 H), 3.91 (ddd, $J=9.8$, 8.0, 6.5 Hz, 1 H), 2.90-2.85 (m, 1 H), 2.32-2.17 (m, 1 H), 1.32 (s, 9 H). LC-MS (Method B): $m/z = 269.0$ $[M+H]^+$, 1.16 min. e.e.=100% as determined on a Chiralpak IC (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/2-propanol 90/10% v/v, flow rate: 1 mL/min, retention time: 9.6 min.

EXAMPLES 82 AND 83

PREPARATION OF (3S)-3-(4-FLUOROPHENYL)-2-((S)-2-METHYLOXOLANE-2-CARBONYL)-1,2-OXAZOLIDINES (136) AND (137)




15 The title compounds ((3S)-3-(4-fluorophenyl)-2-((S)-2-methyloxolane-2-carbonyl)-1,2-oxazolidine and (3S)-3-(4-fluorophenyl)-2-((R)-2-methyloxolane-2-carbonyl)-1,2-oxazolidine) were isolated by chiral HPLC on a Chiralpak OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1.8 mL/min.

First eluting diastereoisomer, Diastereoisomer 1. ^1H NMR (400 MHz, $CDCl_3$) δ 7.34-7.24 (m, 2 H), 7.08-6.98 (m, 2 H), 5.57-5.45 (m, 1 H), 4.25 (td, $J=7.8$, 4.0 Hz, 1 H), 3.98 (q, $J=8.0$ Hz, 1 H), 3.89-3.80 (m, 1 H), 3.79-3.68 (m, 1 H), 3.68-3.57 (m, 1 H), 2.82 (dddd, $J=12.3$, 8.7, 7.2, 4.0 Hz, 1 H), 2.66-2.55 (m, 1 H), 2.39-2.24 (m, 1 H), 1.96-1.73 (m, 3 H), 1.56 (s, 3 H). LC-MS (Method B): $m/z = 280.1$ $[M+H]^+$, 0.89 min. e.e.=100% as determined on a Chiralpak OJ-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min, retention time: 6.0 min.

25 Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400 MHz, $CDCl_3$) δ 7.32-7.26 (m, 2 H), 7.06-6.99 (m, 2 H), 5.52-5.45 (m, 1 H), 4.32-4.23 (td, $J=7.8$, 3.5 Hz, 1 H), 4.04-3.92 (m, 2 H), 3.92-3.83 (m, 1 H), 2.83 (dddd, $J=12.2$, 8.7, 6.9, 3.8 Hz, 1 H), 2.51-2.41 (m, 1 H), 2.34 (dddd, $J=12.3$, 9.1, 7.6, 5.9 Hz, 1 H), 2.02-1.78 (m, 3 H), 1.50 (s, 3 H). LC-MS (Method B): $m/z = 280.1$ $[M+H]^+$, 0.90 min. e.e.=100% as determined on a Chiralpak OJ-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min, retention time: 6.9 min.

EXAMPLE 84


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(5-METHYLPYRAZIN-2-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (138)



5 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 60/40 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.53 (d, $J=1.3$ Hz, 1H), 8.41 (d, $J=1.0$ Hz, 1H), 5.53 (dd, $J=8.8, 6.3$ Hz, 1H), 4.33 (td, $J=7.4, 3.3$ Hz, Hi), 3.97 (ddd, $J=9.4, 7.9, 6.5$ Hz, 1H), 2.86-2.74 (m, 1H), 2.64 (dddd, $J=12.3, 9.4, 7.7, 6.3$ Hz, Hi), 2.56 (s, 3H), 1.30 (s, 9H). LC-MS (Method A): m/z = 250.2 $[\text{M}+\text{H}]^+$, 0.80 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ ethanol 60/40 v/v, flow rate: 1 mL/min, retention time: 9.8 min.

EXAMPLE 85


PREPARATION OF 1-[(3S)-3-(5-FLUOROPYRazin-2-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYL-3-(TRIFLUOROMETHOXY)PROPAN-1-ONE (139)



15 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 70/30 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.43-8.38 (m, 2H), 7.36-7.30 (m, 1H), 5.49 (dd, $J=8.8, 7.0$ Hz, 1H), 4.40-4.27 (m, Hi), 4.16 (s, 2H), 3.99 (ddd, $J=9.6, 8.1, 6.4$ Hz, 1H), 2.92 (ddd, $J=12.4, 5.9, 3.3$ Hz, 1H), 2.42-2.23 (m, 1H), 1.36 (s, 6H). LC-MS (Method A): m/z = 336.0 $[\text{M}+\text{H}]^+$, 1.04 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 70/30 % v/v, flow rate: 1 mL/min, retention time: 9.0 min.

EXAMPLE 86


PREPARATION OF 2,2-D(METHYL-1-[(3S)-3-(PYRAZOLO[1,5-A]PYRIDIN-7-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (140)



5 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 40/60 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.99 (d, $J=2.3$ Hz, 1H), 7.56-7.46 (m, 1H), 7.13 (dd, $J=8.9, 6.9$ Hz, 1H), 6.74 (d, $J=7.0$ Hz, 1H), 6.58 (d, $J=2.3$ Hz, 1H), 6.16 (dd, $J=8.8, 6.0$ Hz, 1H), 4.29-4.17 (m, 1H), 4.01 (ddd, $J=9.3, 8.0, 6.8$ Hz, 1H), 3.34-3.17 (m, 1H), 2.30 (dddd, $J=12.5, 9.3, 7.8, 5.8$ Hz, 1H), 1.38 (s, 10 9H). LC-MS (Method A): $m/z = 274.2$ $[\text{M}+\text{H}]^+$, 1.05 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 40/60 % v/v, flow rate: 1 mL/min, retention time: 5.1 min.

EXAMPLES 87 AND 88

PREPARATION OF 2-FLUORO-5-[(3S)-2-(3-METHYLOXOLANE-3-CARBONYL)-1,2-OXAZOLIDIN-3-YL]PYRIDINES (141) AND (142)



The title compounds (2-fluoro-5-[(3S)-2-((S)-3-methyloxolane-3-carbonyl)-1,2-oxazolidin-3-yl]pyridine and 2-fluoro-5-[(3S)-2-((R)-3-methyloxolane-3-carbon}4)-1,2-oxazolidin-3-yl]pyridine) were isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1+0.1% isopropylamine) 10/90 % v/v, flow rate: 16 mL/min. 20


First eluting diastereoisomer, Diastereoisomer 1. ^1H NMR (400MHz, CDCl_3) δ 8.20 (d, $J=2.8$ Hz, 1H), 7.73 (td, $J=8.0, 2.6$ Hz, 1H), 6.93 (dd, $J=8.5, 3.0$ Hz, 1H), 5.44 (dd, $J=8.8, 6.5$ Hz, 1H), 4.33 (td, $J=7.9, 3.0$ Hz, 1H), 4.17 (d, $J=9.0$ Hz, 1H), 4.01-3.79 (m, 3H), 3.73 (d, $J=9.0$ Hz, 1H), 2.91 (ddt, $J=15.4, 9.0, 3.2$ Hz, 1H), 2.56-2.30 (m, 2H), 1.86 (ddd, $J=12.7, 7.3, 5.5$ Hz, 1H), 1.43 (s, 3H). LC-MS (Method 25 B): $m/z = 280.0$ $[\text{M}+\text{H}]^+$, 0.72 min. Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase

of n-hexane/(ethanol/methanol 1/1+0.1% isopropylamine) 10/90 % v/v, flow rate: 1 mL/min, retention time: 17.8 min.

Second eluting diastereoisomer, Diastereoisomer 2. ^1H NMR (400MHz, CDCl_3) δ 8.20 (d, $J=23$ Hz, 1H), 7.73 (dt, $J=2.5, 8.0$ Hz, 1H), 6.92 (dd, $J=1.5, 3.0$ Hz, 1H), 5.42 (dd, $J=8.7, 6.7$ Hz, 1H), 4.33 (td, $J=7.8, 2.9$ Hz, 1H), 4.04-3.72 (m, 5H), 2.91 (dddd, $J=12.3, 9.1, 6.2, 3.0$ Hz, 1H), 2.60 (ddd, $J=12.9, 8.1, 6.9$ Hz, 1H), 2.36 (dddd, $J=12.5, 9.7, 7.6, 6.7$ Hz, 1H), 1.81 (ddd, $J=12.9, 7.4, 5.8$ Hz, 1H), 1.46 (s, 3H). LC-MS (Method B): $m/z = 280.1$ $[\text{M}+\text{H}]^+$, 0.90 min. Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/(ethanol/methanol 1/1+0.1% isopropylamine) 10/90 % v/v, flow rate: 1 mL/min, retention time: 19.6 min.

EXAMPLE 89


PREPARATION OF 5-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]PYRIDINE-2-CARBONITRILE (143)



The title compound was isolated by chiral HPLC on a Chiralpak IA (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 50/50 % v/v, flow rate: 15 mL/min. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.66 (d, $J=1.8$ Hz, 1H), 8.02 (d, $J=7.8$ Hz, 1H), 7.88 (dd, $J=7.8, 2.1$ Hz, 1H), 5.43 (t, $J=8.1$ Hz, 1H), 4.36-4.30 (m, 1H), 3.94-3.86 (m, 1H), 2.94-2.90 (m, 1H), 2.24-2.17 (m, 1H), 1.21 (s, 9H). LC-MS (Method E): $m/z = 260.1$ $[\text{M}+\text{H}]^+$, 2.67 min. e.e.=99.9% as determined on a Chiralpak IA-3 (5 x 0.46 cm), 3 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 3.02 min.

EXAMPLE 90

PREPARATION OF 1-[(3S)-3-(2-METHOXPYRIDIN-4-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (144)




The title compound was isolated by chiral HPLC on a **Chiralpak IC** (25 x 2.0 cm), 5 μ m column using a mobile phase of **n-hexane/ethanol** 80/20 % v/v, flow rate: 20 mL/min. ^1H NMR (400 MHz, **DMSO-*d*₆**) δ 8.10 (d, $J=5.2$ Hz, 1H), 6.84 (dd, $J=5.2, 1.2$ Hz, 1H), 6.60 (s, 1H), 5.27-5.23 (m, 1H), 4.27-4.22 (m, 1H), 3.85-3.79 (m, 4H), 2.87-2.82 (m, 1H), 2.10-2.04 (m, 1H), 1.19 (s, 9H). LC-MS (Method F): m/z = 265.0 $[\text{M}+\text{H}]^+$, 1.04 min. e.e.=99.9% as determined on a Chiralpak **IC-3** (5 x 0.46 cm), 5 μ m column using a mobile phase of (**n-hexane** + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 2.59 min.

EXAMPLE 91


PREPARATION OF 1-[(3S)-3-(6-METHOXPYRIDIN-3-YL)-1[^]-OXAZOLIDIN-2-YL]-2,2-

DIMETHYLPROPAN-1-ONE (145)



The title compound was isolated by chiral SFC on a **Chiralart Amylose-SA** (25 x 2.0 cm), 5 μ m column using a mobile phase of carbon dioxide/(methanol + 2 mM ammonia) 80/20 % v/v, flow rate: 40 mL/min. ^1H NMR (400 MHz, **DMSO-*d*₆**) δ 8.03 (d, $J=2.4$ Hz, 1H), 7.55 (dd, $J=8.4, 2.4$ Hz, 1H), 6.79 (d, $J=8.8$ Hz, 1H), 5.28-5.25 (m, 1H), 4.29-4.27 (m, 1H), 3.85-3.79 (m, 4H), 2.82-2.79 (m, 1H), 2.23-2.10 (m, 1H), 1.17 (s, 9H). LC-MS (Method O): m/z = 265.1 $[\text{M}+\text{H}]^+$, 1.82 min. e.e.=99.9% as determined on a Chiralpak **AD-3** (3 x 100 mm), 3 μ m column using a mobile phase of carbon dioxide/(methanol + 0.1% diethylamine) 95/5 % v/v, flow rate: 2.0 mL/min, retention time: 1.54 min.

EXAMPLE 92


PREPARATION OF 2-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1[^],2-OXAZOLIDIN-3-YL] BENZONITRILE (146)

The title compound was isolated by chiral HPLC on a Chiralpak **IC** (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/isopropanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (400 MHz, **OMSO-*d*₆**) δ 7.83 (dd, $J=7.6, 1.2$ Hz, 1H), 7.73-7.69 (m, 1H), 7.49-7.41 (m, 2H), 5.49 (t, $J=8.0$ Hz, 1H), 4.37-4.33 (m, 1H), 3.96-3.90 (m, 1H), 2.99-2.96 (m, 1H), 2.14-2.09 (m, 1H), 1.21 (s, 9H). LC-MS

(Method H): $m/z = 258.9$ $[\text{M}+\text{H}]^+$, 1.48 min. e.e.=98.9% as determined on a Chiraipak IC-3 (5 x 0.46 cm), 3 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/isopropanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 2.59 min.

EXAMPLE 93


5 PREPARATION OF 34(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE (147)



The title compound was isolated by chiral HPLC on a Chiraipak IC (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, DMSO- d_6) δ 7.76-7.64 (m, 2H), 7.60-7.51 (m, 2H), 5.36-5.31 (m, 1H), 4.25-4.31 (m, 1H), 3.88-3.80 (m, 1H), 2.92-2.82 (m, 1H), 2.20-2.08 (m, 1H), 1.19 (s, 9H). LC-MS (Method F): $m/z = 259.0$ $[\text{M}+\text{H}]^+$, 1.31 min. e.e.=99.9% as determined on a Chiraipak IC-3 (5 x 0.46 cm), 5 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 2.32 min.

EXAMPLE 94


15 PREPARATION OF 1-[(3S)-3-(6-CYCLOPROPYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (148)



The title compound was isolated by chiral HPLC on a Chiraipak IC (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, DMSO- d_6) δ 8.26 (d, $J=2.1$ Hz, 1H), 7.47 (dd, $J=8.1, 2.1$ Hz, 1H), 7.23 (d, $J=7.8$ Hz, 1H), 5.34-5.18 (m, 1H), 4.31-4.25 (m, 1H), 3.87-3.79 (m, 1H), 2.88-2.70 (m, 1H), 2.24-1.92 (m, 2H), 1.17 (s, 9H), 0.96-0.82 (m, 4H). LC-MS (Method F): $m/z = 275.0$ $[\text{M}+\text{H}]^+$, 0.80 min. e.e.=99.9% as determined on a Chiraipak IC-3 (5 x 0.46 cm), 5 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 2.89 min.

EXAMPLE 95


PREPARATION OF 1-[(3S)-3-(2-AMINOPYRIDIN-4-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (149)



5 The title compound was isolated by chiral HPLC on a Chiralpak IA (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, $\text{DMSO-}d_6$) δ 7.83 (d, $J=5.4$ Hz, 1H), 6.37 (dd, $J=5.4$, 1.5 Hz, 1H), 6.31 (s, 1H), 6.05 (brs, 2H), 5.15-5.10 (m, 1H), 4.25-4.24 (m, 1H), 3.82-3.78 (m, 1H), 2.85-2.80 (m, 1H), 2.07-2.00 (m, 1H), 1.21 (s, 9H). LC-MS (Method J): m/z = 250.3 [M+H] $^+$, 1.01 min. e.e.=99.9% as determined on a Chiralpak IA-3 (5 x 0.46
10 cm), 3 μ m column using a mobile phase of (n-hexane + 0.2% isopropylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 2.57 min.

EXAMPLE 96


PREPARATION OF 3-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-4-FHJROBENZOMTRILE (150)



15 The title compound was isolated by chiral HPLC on a Lux 5u Cellulose-3, AXIA Packed (25 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 90/10 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, $\text{DMSO-}d_6$) δ 7.89-7.83 (m, 1H), 7.61 (dd, $J=6.9$, 1.8 Hz, 1H), 7.50-7.43 (m, 1H), 5.47-5.42 (m, 1H), 4.34-4.29 (m, 1H), 3.91-3.83 (m, 1H), 2.95-2.79 (m, 1H), 2.18-2.11 (m, 1H), 1.20 (s, 9H). LC-MS (Method H): m/z = 276.9 [M+H] $^+$, 1.51 min. e.e.=99.9% as determined on a Lux Cellulose-3 (15 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/isopropanol
20 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 3.19 min.

EXAMPLE 97


PREPARATION OF 4-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-FLUOROBENZONITRILE (151)



5 The title compound was isolated by chiral HPLC on a Chiralpak IF (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 18 mL/min. ^1H NMR (300 MHz, $\text{DMSO-}d_6$) δ 7.94-7.89 (m, 1H), 7.39-7.27 (m, 2H), 5.42-5.36 (m, 1H), 4.37-4.24 (m, 1H), 3.90-3.82 (m, 1H), 2.92-2.88 (m, 1H), 2.10-2.17 (m, 1H), 1.22 (s, 9H). LC-MS (Method D): m/z = 277.1 $[\text{M}+\text{H}]^+$, 2.07 min. e.e.=99.9% as determined on a Chiralpak IF-3 (10 x 0.46 cm), 3 μ m column using a mobile
10 phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 3.36 min.

EXAMPLE 98


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (152)



15 The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, $\text{DMSO-}d_6$) δ 8.32 (d, J =2.1 Hz, 1H), 7.50 (dd, J =8.1, 2.4 Hz, 1H), 7.20 (d, J =8.1 Hz, 1H), 5.29 (t, J =6.9 Hz, 1H), 4.32-4.26 (m, 1H), 3.88-3.80 (m, 1H), 2.86-2.82 (m, 1H), 2.42 (s, 3H), 2.21-2.11 (m, 1H), 1.18
20 (s, 9H). LC-MS (Method J): m/z = 249.3 $[\text{M}+\text{H}]^+$, 1.20 min. e.e.=99.9% as determined on a Chiralpak IC-3 (10 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 4.17 min.

EXAMPLE 99

PREPARATION OF 5-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-FLUOROBENZONITRILE (153)




5 The title compound was isolated by chiral HPLC on a Phenomenex Lux 5u Cellulose-4, AXIA Packed (25 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.79-7.77 (m, 1H), 7.66-7.62 (m, 1H), 7.54-7.49 (in. 1H), 5.34 (t, $J=8.4$ Hz, 1H), 4.32-4.28 (m, 1H), 3.88-3.82 (m, 1H), 2.88-2.85 (m, 1H), 2.19-2.14 (m, 1H), 1.20 (s, 9H). LC-MS (Method L): m/z = 277.2 $[\text{M}+\text{H}]^+$, 1.56 min. e.e.=99.9% as determined on a Lux

10 Cellulose-3 (10 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1 % diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 2.44 min.

EXAMPLE 100

PREPARATION OF 4-[(3S)-2-[3-(DIFLUOROMETHOXY)-2,2-DIMETHYLPROPANOYL]-1,2-OXAZOLIDIN-3-YL]BENZONITRILE (154)




15 The title compound was isolated by chiral HPLC on a EnantioPak A 1-5 (25 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, CD_3OD) 7.72 (d, $J=8.4$ Hz, 2H), 7.50 (d, $J=8.1$ Hz, 2H), 6.42 (t, $J=7.6$ Hz, 1H), 5.44 (t, $J=7.2$ Hz, 1H), 4.40-4.34 (m, 1H), 4.13-4.00 (m, 3H), 3.03-2.93 (m, 1H), 2.32-2.26 (m, 1H), 1.33 (s, 6H). LC-MS

20 (Method D): m/z = 325.1 $[\text{M}+\text{H}]^+$, 2.00 min. e.e.=99.9% as determined on a Chiralpak IC-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 1.53 min.

EXAMPLE 101

PREPARATION OF 3,3-DIFLUORO-1-[(3S)-3-(5-FLUORO-6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-2,2-DIMETHYLPROPAN-1-ONE (195)




5 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.28 (s, 1H), 7.24 (dd, *J*=9.8, 1.8 Hz, 1H), 6.32 (t, *J*=57.7 Hz, 1H), 5.44 (dd, *J*=8.7, 6.7 Hz, 1H), 4.35 (td, *J*=7.8, 3.0 Hz, 1H), 3.98 (ddd, *J*=9.5, 8.3, 6.5 Hz, 1H), 2.90 (dddd, *J*=12.4, 9.2, 6.3, 3.3 Hz, 1H), 2.52 (d, *J*=3.0 Hz, 3H), 2.35 (dddd, *J*=12.4, 9.6, 7.6, 6.8 Hz, 1H), 1.40-1.36 (m, 6H). LC-MS

10 (Method A): *m/z* = 302.9 [M+H]⁺, 0.93 min. e.e.= 100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 1 mL/min, retention time: 5.5 min.

EXAMPLE 102


PREPARATION OF 4-[(3S)-3-(5-FLUORO-6-METHYL PYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]-3,3-DIMETHYL-4-OXOBUTANENITRILE (196)



- 5 The title compound was isolated by chiral HPLC on a Chiralcel OJ-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 50/50% v/v, flow rate: 18 mL/min. ¹H NMR (400 MHz, CDCl₃) δ 8.30-8.27 (m, 1H), 7.26 (dd, *J*=9.8, 1.8 Hz, 1H), 5.43 (dd, *J*=8.7, 6.7 Hz, 1H), 4.41-4.32 (m, 1H), 4.01 (ddd, *J*=9.5, 8.0, 6.5 Hz, 1H), 2.91 (dddd, *J*=12.3, 9.2, 6.4, 3.0 Hz, 1H), 2.73 (d, *J*=16.8 Hz, 1H), 2.67 (d, *J*=17.1 Hz, 1H), 2.52 (d, *J*=2.8 Hz, 3H), 2.36 (dddd, *J*=12.5, 9.6, 7.5, 6.5 Hz, 1H), 1.50 (s, 3H), 1.48 (s, 3H). LC-MS (Method A): *m/z* = 291.9 [M+H]⁺, 0.79 min. e.e. = 100% as determined on a Chiralcel OJ-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/ethanol 65/35% v/v, flow rate: 1 mL/min, retention time: 6.4 min.
- 10

EXAMPLE 103


3,3-DIFLUORO-2,2-DIMETHYL-1-[(3S)-3-(5-METHYLPYRAZOL-2-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (197)



¹H NMR (400 MHz, CDCl₃) δ 8.51 (d, *J*=1.5 Hz, 1H), 8.41 (d, *J*=1.0 Hz, 1H), 6.32 (t, *J*=58.2 Hz, 1H), 5.51 (dd, *J*=8.9, 6.1 Hz, 1H), 4.39 (td, *J*=7.8, 3.9 Hz, 1H), 4.03 (ddd, *J*=9.0, 7.9, 6.7 Hz, 1H), 2.88-2.77 (m, 1H), 2.74-2.62 (in. 1H), 2.57 (s, 3H), 1.39-1.34 (m, 6H). LC-MS (Method A): *m/z* = 286.4 [M+H]⁺, 0.81 min.

EXAMPLE 104

PREPARATION OF (S)-3,3-DIFLUORO-2,2-DIMETHYL-1-[(3-(1-METHYL-1H-PYRAZOL-4-YL)ISOXAZOLIDIN-2-YL)PROPAN-1-ONE (198)




The title compound was isolated by chiral HPLC on a Chiralpak 1C (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.59 (s, 1H), 7.32 (s, 1H), 6.33 (t, *J*=56.9 Hz, 1H), 5.28-5.24 (m, 1H), 4.30-4.25 (m, 1H),

3.90–3.84 (m, 1H), 3.76 (s, 3H), 2.73–2.65 (m, 1H), 2.33–2.25 (m, 1H), 1.21 (s, 6H). LC-MS (Method F): $m/z = 273.9 [M+H]^+$, 0.98 min. e.e.=99.9% as determined on a Chiralpak IC-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 1.86 min.

5

EXAMPLE 105

PREPARATION OF (S)-3-(2-(3-CYANO-2,2-DIMETHYLPROPANOYL)ISOXAZOLIDIN-3-YL)-5-FLUOROBENZONITRILE (199)




The title compound was isolated by chiral HPLC on a Chiralpak IA (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, CD_3OD) δ 7.55–7.53 (m, 1H), 7.51–7.39 (m, 2H), 5.49–5.36 (m, 1H), 4.46–4.33 (m, 1H), 4.08–3.99 (m, 1H), 3.07–2.91 (m, 1H), 2.91–2.70 (m, 2H), 2.37–2.25 (m, 1H), 1.47 (s, 3H), 1.43 (s, 3H). LC-MS (Method Q): $m/z = 302.0 [M+H]^+$, 1.33 min. e.e.=99.9% as determined on a Chiralpak IA-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 1.99 min.

15

EXAMPLE 106

PREPARATION OF (S)-5-(2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)ISOXAZOLIDIN-3-YL)NICOTINONITRILE (200)




20

The title compound was isolated by chiral HPLC on a EnantioPak AI-5 (25 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (400

MHz, DMSO- d_6) δ 8.94 (d, $J=2.0$ Hz, 1H), 8.74 (d, $J=2.2$ Hz, 1H), 8.18–8.16 (m, 1H), 6.38 (t, $J=56.7$ Hz, 1H), 5.52–5.34 (m, 1H), 4.39–4.34 (m, 1H), 3.99–3.93 (m, 1H), 2.94–2.87 (m, 1H), 2.33–2.24 (m, 1H), 1.27 (s, 3H), 1.25 (s, 3H). LC-MS (Method F): m/z = 295.9 $[M+H]^+$, 1.11 min. e.e.=97.0% as determined on a Chiraipak AS-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 1.37 min.

EXAMPLE 107


PREPARATION OF (S)-3-(3,3-DIFLUORO-2,2-DIMETHYL-1-(3-(6-METHYLPYRAZIN-2-YL)OXAZOLIDIN-2-YL)PROPAN-1-ONE (201)



The title compound was isolated by chiral HPLC on a Chiralcel OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 20 mL/min. ^1H NMR (400 MHz, CD_3OD) δ 8.40 (s, 1H), 8.34 (s, 1H), 6.36 (t, $J=57.1$ Hz, 1H), 5.46–5.42 (m, 1H), 4.45–4.40 (m, 1H), 4.12–4.07 (m, 1H), 2.93–2.85 (m, 1H), 2.59–2.50 (m, 4H), 1.34–1.32 (m, 6H). LC-MS (Method R): m/z = 286.0 $[M+H]^+$, 2.55 min. e.e.=99.9% as determined on a Chiraipak IA-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 1.56 min.

EXAMPLE 108

PREPARATION OF 3,3-DIFLUORO-2,2-DIMETHYL-1-[(3S)-3-(6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE (202)




To a stirred solution of 3,3-difluoro-2,2-dimethylpropanoic acid (150 mg, 1.09 mmol), N,N,N',N'-tetraethyl-0-(7-azabenzotriazol-1-yl)uronium hexafluorophosphate (619 mg, 1.63 mmol) and N,N-diisopropylethylamine (421 mg, 3.26 mmol) in DMF (6 mL) was added 2-methyl-5-(1,2-oxazolidin-3-yl)pyridine (357 mg, 2.17 mmol). After being stirred at room temperature for 4 h, the reaction mixture was quenched by the addition of water (20 mL), and extracted with EtOAc (3 x 20 mL). The combined organic layers were washed with brine (20 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The residue was purified by preparative-HPLC with the following conditions: column: Xbridge Prep C18, 19 x 150 mm, 5 μ m; Mobile phase: Phase A: water (10 mmol/L

NH_4HCO_3); Phase B: acetonitrile (20% to 85% over 10 min); Detector, UV 220 & 254 nm to afford 3,3-difluoro-2,2-dimethyl-1-[3-(6-methylpyridin-3-yl)-1,2-oxazolidin-2-yl]propan-1-one as a mixture of enantiomers (180 mg, 58%) as a white solid. This mixture was resolved by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 80/20 % v/v and a flow rate of 20 mL/min to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 8.34 (d, J = 2.0 Hz, 1H), 7.52 (dd, J = 8.0, 2.4 Hz, 1H), 7.23 (d, J = 8.0 Hz, 1H), 6.39 (t, J = 56.8 Hz, 1H), 5.35-5.31 (m, 1H), 4.38-4.33 (m, 1H), 3.96-3.90 (m, 1H), 2.93-2.85 (m, 1H), 2.44 (s, 3H), 2.24-2.17 (m, 1H), 1.27 (s, 3H), 1.26 (s, 3H). LC-MS (Method J): m/z = 285.4 $[\text{M}+\text{H}]^+$, 1.156 min. e.e. = 99.9% as determined on a Chiralpak IC-3 (5 x 0.46 cm), 3 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 2.056 min.

EXAMPLES 109 AND 110

PREPARATION OF 5-[(3R)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL] THIOPHENE-2-CARBONITRILE AND 5-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL] THIOPHENE-2-CARBONITRILE (203 AND 204)



These compound was synthesized by the same method as described for 3,3-difluoro-2,2-dimethyl-1-[3-(6-methylpyridin-3-yl)-1,2-oxazolidin-2-yl]propan-1-one except 5-(1,2-oxazolidin-3-yl)thiophene-2-carbonitrile (50 mg, 0.28 mmol) was used instead of 2-methyl-5-(1,2-oxazolidin-3-yl)pyridine. The title compounds were isolated by chiral HPLC on a CHIRALART Cellulose-SB-KSB99S05-2520WX12806 (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/ethanol 85/15 % v/v, flow rate: 20 mL/min.

5-[(3R)-2-(3,3-difluoro-2,2-dimethylprop-1-yn-1-yl)-1,2-oxazolidin-3-yl]thiophene-2-carbonitrile.

^1H NMR (400 MHz, CD_3OD) δ 7.62 (d, J = 4.0 Hz, 1H), 7.13 (d, J = 3.6 Hz, 1H), 6.33 (t, J = 57.2 Hz, 1H), 5.72-5.68 (m, 1H), 4.41-4.36 (m, 1H), 4.06-4.00 (m, 1H), 2.98-2.92 (m, 1H), 2.52-2.45 (m, 1H), 1.33 (s, 6H). LC-MS (Method D): m/z = 301.1 $[\text{M}+\text{H}]^+$, 1.751 min. e.e. = 97.7% as determined on a CHIRAL Cellulose-SB (0.46 x 15 cm) 5 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 4.296 min.


5-[(3S)-2-(3,3-difluoro-2,2-dimethylprop-1-yn-1-yl)-1,2-oxazolidin-3-yl]thiophene-2-carbonitrile.

^1H NMR (400 MHz, CD_3OD) δ 7.62 (d, J = 4.0 Hz, 1H), 7.13 (d, J = 4.0 Hz, 1H), 6.33 (t, J = 57.2 Hz, 1H), 5.72-5.68 (m, 1H), 4.41-4.36 (m, 1H), 4.06-4.00 (m, 1H), 2.99-2.91 (m, 1H), 2.52-2.45 (m, 1H), 1.32 (s, 6H). LC-MS (Method D): m/z = 301.1 $[\text{M}+\text{H}]^+$, 1.751 min. e.e. = 99.7% as determined on a

CHIRAL Cellulose-SB (0/46 x 15 cm) 5 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 4.786 min.

EXAMPLE 111


PREPARATION OF 4-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-1-METHYL-1H-PYRROLE-2-CARBONITRILE (205)



The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, DMSO- d_6) δ 7.09 (d, J = 1.8 Hz, 1H), 6.83 (d, J = 1.8 Hz, 1H), 5.24-5.18 (m, 1H), 4.25-4.20 (m, 1H), 3.83-3.75 (m, 1H), 3.70 (s, 3H), 2.73-2.60 (tn, 1H), 2.27-2.17 (m, 1H), 1.16 (s, 9H). LC-MS (Method F): m/z = 262.0 $[\text{M}+\text{H}]^+$, 1.233 min. e.e.=99.9% as determined on a Lux Cellulose-4 (10 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 2.939 min.

EXAMPLE 112

PREPARATION OF (Z)-4-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-N'-HYDROXYBENZENE-1-CARBOXIMIDAMIDE (206)




To a solution of 4-[(3S)-2-(3,3-difluoro-2,2-dimethylpropionyl)-1,2-oxazolidin-3-yl]benzonitrile (1 g, 3.4 mmol) in DMF (10 mL) was added hydroxyamine (50% aq. solution) (1.12 mL, 16 mmol). The reaction mixture was allowed to stir at room temperature for 2 h. After completion of the reaction, DMF was removed under reduced pressure. The residue was diluted with EtOAc (50 mL) and washed with sat. NaHCO₃ solution (20 mL). The aqueous layer was extracted further with EtOAc (3 x 30 mL). The combined organic layers were washed with brine and dried over anhydrous Na₂SO₄ and evaporated under reduced pressure. The crude product was purified by column chromatography (Petroleum ether/EtOAc,

60:40 to 57:43) to afford the title compound (960 mg, 86%) as a white solid. ^1H NMR (400MHz, DMSO- d_6) δ 9.61 (s, 1H), 7.63 (d, $J=8.4$ Hz, 2H), 7.24 (d, $J=8.4$ Hz, 2H), 6.41 (t, $J=56.8$ Hz, 1H), 5.79 (s, 2H), 5.37-5.30 (m, 1H), 4.50-4.31 (m, 1H), 3.97-3.87 (m, 1H), 2.94-2.86 (m, 1H), 2.26-2.12 (m, 1H), 1.32-1.24 (m, 6H). LC-MS (Method T): m/z = 328.3 [$\text{M}+\text{H}$] $^+$, 2.014 min. Melting range = 67.3°-75.6°C, $[\alpha]_D^{23.7} = -132.89^\circ$ ($c = 0.152$, MeOH). e.e. = 100% as determined on a Lux C2 (25 x 0.46 cm), 5 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 9.439 min.

EXAMPLE 113


PREPARATION OF (Z)-4-[(3S)-2-(3,3-Difluoro-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]-N'-hydroxybenzene-1-carboximidamide (207)



(Z)-4-[(3S)-2-(3,3-Difluoro-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]-N'-hydroxybenzene-1-carboximidamide (3.4 g, 10.4 mmol) was suspended in ethanol (10.2 mL) and triethyl orthoformate (10.2 mL). The reaction mixture was heated to 100°C for 2 days in a sealed tube. After completion of the reaction, the reaction mixture was cooled to room temperature and ethanol was removed under reduced pressure. The crude product was purified by column chromatography (petroleum ether/EtOAc, 75:25 to 72:27) and then by preparative HPLC with the following conditions: column: Sunfire OBD, C18; 250x30 mm, 5 μm ; Mobile phase: 0.1% TFA in $\text{H}_2\text{O}/\text{ACN}$, Flow rate: 22 mL/min, to afford the title compound (950 mg, 27%) as an off-white solid. ^1H NMR (400MHz, CDCl_3) δ 8.77 (s, 1H), 8.18-8.07 (m, 2H), 7.48-7.39 (m, 2H), 6.35 (t, $J=57.0$ Hz, 1H), 5.55-5.43 (m, 1H), 4.40-4.30 (m, 1H), 4.05-3.94 (m, 1H), 2.98-2.87 (m, 1H), 2.47-2.33 (m, 1H), 1.46-1.34 (m, 6H). LC-MS (Method T): m/z = 338.3 [$\text{M}+\text{H}$] $^+$, 2.696 min. Melting range = 87.9°-88.9°C, $[\alpha]_D^{28.4} = -156.87^\circ$ ($c = 0.160$, MeOH). e.e. = 100% as determined on a Lux C2 (25 x 0.46 cm), 5 μm column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 6.708 min.

EXAMPLE 114


PREPARATION OF 1-[(3S)-3-[4-(5-CYCLOPROPYL-1,2,4-OXAZOL-3-YL)PHENYL]-1,2-OXAZOLIDIN-2-YL]-3,3-DIFLUORO-2,2-DIMETHYLPROPAN-1-ONE (208)



To a solution of cyclopropane carboxylic acid (421 mg, 4.9 mol) and (5)-4-(2-(3,3-difluoro-2,2-diethylpropanoyl)isoxazolidin-3-yl)-N-hydroxybenzimidamide (1 g, 3.1 mmol) in EtOAc (10 mL) was added Et₃N (2.11 mL, 15.1 mmol) followed by T3P (50% solution in EtOAc, 5.8 mL, 7 mmol). The reaction mixture was heated to 80°C under an inert atmosphere for 20 h. The reaction mixture was cooled to room temperature, diluted with ice water (30 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine and dried over Na₂SO₄. The crude product was purified twice by column chromatography (petroleum ether/EtOAc, 80:20 to 75:25) to afford the title compound (360 mg, 31%) as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ 8.08-7.98 (m, 2H), 7.43-7.33 (m, 2H), 6.34 (t, J=57.2 Hz, 1H), 5.52-5.41 (m, 1H), 4.38-4.28 (m, 1H), 4.03-3.92 (m, 1H), 2.95-2.82 (m, 1H), 2.44-2.30 (m, 1H), 2.30-2.20 (m, 1H), 1.45-1.18 (m, 10H). LC-MS (Method T): m/z = 378.3 [M+H]⁺, 2.924 min. Melting range = 92.1° - 95.6°C, [α]_D^{25.2} = -147.61° (c = 0.105, MeOH). e.e. = 99.3% as determined on a Chiralcel OD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-heptane/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 6.141 min.

EXAMPLE 115

PREPARATION OF 4-[(3S)-2-[(2S)-3-CYANO-2-METHYLPROPANOYL]-1,2-OXAZOLIDIN-3-YL]BENZONITRILE DIASTEREOISOMER 1 (209)




DIASTEREOISOMER 1

The title compound was isolated by chiral HPLC on a Chiralpak AS-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.69-7.64 (m, 2H), 7.44-7.39 (m, 2H), 5.45 (dd, $J=8.8, 6.3$ Hz, 1H), 4.32 (dt, $J=3.1, 7.8$ Hz, 1H), 4.00 (ddd, $J=9.5, 8.2, 6.7$ Hz, 1H), 3.36 (sextet, $J=7.2$ Hz, 1H), 3.01-2.88 (m, 1H), 2.69 (dd, $J=16.8, 7.8$ Hz, 1H), 2.45 (dd, $J=16.8, 6.8$ Hz, 1H), 2.35 (dddd, $J=12.5, 9.5, 7.6, 6.3$ Hz, 1H), 1.40 (d, $J=7.0$ Hz, 3H). LC-MS (Method A): $m/z = 270.2$ $[\text{M}+\text{H}]^+$, 0.84 min. e.e.=100% as determined on a Chiralpak AS-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 7.7 min.

EXAMPLE 116

PREPARATION OF 4-[(3S)-2-[(2S)-3-CYANO-2-METHYLPROPANOYL]-1,2-OXAZOLIDIN-3-
YL] BENZONITRILE DIASTEREOMER 2 (210)




DIASTEREOMER. 2

The title compound was isolated by chiral HPLC on a Chiralpak AS-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.70-7.63 (m, 2H), 7.47-7.40 (m, 2H), 5.41 (dd, $J=8.8, 6.3$ Hz, 1H), 4.34 (dt, $J=3.9, 7.7$ Hz, 1H), 4.06-3.95 (m, 1H), 3.37 (sextet, $J=7.0$ Hz, 1H), 3.00-2.87 (in. 1H), 2.69 (dd, $J=16.6, 6.5$ Hz, 1H), 2.54 (dd, $J=17.3, 8.0$ Hz, 1H), 2.37 (dddd, $J=12.5, 9.1, 7.4, 6.1$ Hz, 1H), 1.31 (d, $J=7.0$ Hz, 3H). LC-MS (Method A): $m/z = 270.2$ $[\text{M}+\text{H}]^+$, 0.81 min. e.e.=100% as determined on a Chiralpak AS-H (25 x 0.46 cm) 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 6.6 min.

EXAMPLE 117


PREPARATION OF (3S)-3-(3,4-DIFLUOROPHENYL)-2-(3-METHYLOXOLANE-3-CARBONYL)-1,2-
OXAZOLIDINE DIASTEREISOMER 1 (211)



5 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 60/40 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.18-7.07 (m, 2H), 7.06- 6.99 (m, 1H), 5.35 (dd, J =8.7, 6.4 Hz, 1H), 4.28 (dt, J =3.1, 7.8 Hz, 1H), 4.18 (d, J =9.0 Hz, 1H), 4.00- 3.79 (m, 3H), 3.73 (d, J =9.0 Hz, 1H), 2.86 (dddd, J =12.2, 9.2, 6.3, 3.1 Hz, 1H), 2.47 (td, J =7.8, 12.7 Hz, 1H), 2.32 (dddd, J =12.5, 9.6, 7.5, 6.5 Hz, 1H), 1.86 (ddd, J =12.8, 7.3, 5.5 Hz, 1H), 1.43 (s, 3H). LC-MS (Method A): m/z = 298.2 $[\text{M}+\text{H}]^+$, 0.96 min. e.e.=97.5% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 7.8 min.

EXAMPLE 118

15 PREPARATION OF (3S)-3-(3,4-DIFLUOROPHENYL)-2-(3-METHYLOXOLANE-3-CARBONYL)-1,2-
OXAZOLIDINE DIASTEREISOMER 2 (212)




20 The title compound was isolated by chiral HPLC on a Chiralpak AD-H (2.5 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 60/40 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.18-7.08 (m, 2H), 7.07-7.00 (m, 1H), 5.34 (dd, J =8.8, 6.5 Hz, 1H), 4.29 (dt, J =3.0, 7.9 Hz, 1H), 3.98 (d, J =9.0 Hz, 1H), 3.96-3.88 (m, 2H), 3.84 (dt, J =5.8, 8.3 Hz, 1H), 3.77 (d, J =9.0 Hz, 1H), 2.91-2.81 (m, 1H), 2.62 (ddd, J =12.8, 8.2, 6.9 Hz, 1H), 2.32 (dddd, J =12.5,

9.6, 7.5, 6.5 Hz, 1H), 1.81 (ddd, $J=13.0$, 7.3, 6.0 Hz, 1H), 1.48 (s, 3H). LC-MS (Method A): m/z = 298.2 [M+H]⁺, 0.97 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 60/40 % v/v, flow rate: 1.0 mL/min, retention time: 8.7 min.

5

EXAMPLE 119

**PREPARATION OF 4-[(3S)-2-(2-CYCLOPROPYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE
DIASTEREISOISOMER 1 (213)**




DIASTEREISOISOMER 1

The title compound was isolated by chiral HPLC on a Chiralpak OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18 mL/min. ¹H NMR (400MHz, CDCl₃) δ 7.70-7.62 (m, 2H), 7.47-7.42 (m, 2H), 5.48 (dd, $J=9.0$, 6.3 Hz, 1H), 4.24 (dt, $J=3.1$, 7.8 Hz, 1H), 3.87 (ddd, $J=9.5$, 8.1, 6.8 Hz, 1H), 2.94-2.83 (m, 1H), 2.41-2.24 (m, 2H), 1.22 (d, $J=7.0$ Hz, 3H), 1.12-0.98 (m, 1H), 0.62-0.36 (m, 3H), 0.26-0.15 (m, 1H). LC-MS (Method A): m/z = 271.2 [M+H]⁺, 1.01 min. e.e.=100% as determined on a Chiralpak OJ-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 7.0 min.

15

EXAMPLE 120

**PREPARATION OF 4-[(3S)-2-(2-CYCLOPROPYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE
DIASTEREISOISOMER 2 (214)**



DIASTEREISOISOMER 2


20

The title compound was isolated by chiral HPLC on a Chiralpak OJ-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18

mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.68-7.61 (m, 2H), 7.47-7.40 (m, 2H), 5.48 (dd, $J=8.8, 6.5$ Hz, 1H), 4.25 (dt, $J=2.6, 7.7$ Hz, 1H), 3.88 (ddd, $J=9.7, 8.0, 6.7$ Hz, 1H), 2.90 (dddd, $J=12.2, 9.2, 6.5, 3.0$ Hz, 1H), 2.39-2.19 (m, 2H), 1.29 (d, $J=7.0$ Hz, 3H), 1.04 (td, $J=8.1, 5.0, 9.7$ Hz, 1H), 0.63-0.45 (m, 2H), 0.24-0.12 (m, 2H). LC-MS (Method A): $m/z = 271.2$ $[\text{M}+\text{H}]^+$, 1.01 min. e.e.=100% as determined on a Chiralpak OJ-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 9.6 min.

EXAMPLE 121

PREPARATION OF 4-[(3S)-2-(2-TRANS-METHYLCYCLOPROPANECARBONYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE DIASTEREISOMER 1 (215)




DIASTEREISOMER 1

The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μm column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1% isopropylamine) 30/70 % v/v, flow rate: 17 mL/min. ^1H NMR (400MHz, $\text{DMSO}-d_6$) δ 7.86-7.79 (m, 2H), 7.54-7.46 (m, 2H), 5.41 (dd, $J=8.8, 6.3$ Hz, 1H), 4.28 (dt, $J=3.0, 7.9$ Hz, 1H), 3.85 (ddd, $J=9.5, 8.0, 6.8$ Hz, 1H), 2.92 (dddd, $J=12.1, 9.1, 6.5, 2.9$ Hz, 1H), 2.19 (dddd, $J=12.2, 9.5, 7.7, 6.3$ Hz, 1H), 1.97 (far. s., 1H), 1.26-1.16 (m, 1H), 1.10 (d, $J=6.0$ Hz, 3H), 1.04-0.94 (m, 1H), 0.75-0.66 (m, 1H). LC-MS (Method A): $m/z = 257.1$ $[\text{M}+\text{H}]^+$, 0.90 min.

EXAMPLE 122

PREPARATION OF 4-[(3S)-2-(2-TRANS-METHYLCYCLOPROPANECARBONYL)-1,2-OXAZOLIDIN-3-YL]BENZONITRILE DIASTEREISOMER 2 (216)




DIASTEREISOMER 2

The title compound was isolated by chiral HPLC on a ChiraSpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol/methanol 1/1 + 0.1% isopropylamine) 30/70 % v/v, flow rate: 17 mL/min. ^1H NMR (400MHz, DMSO- d_6) δ 7.87-7.78 (m, 2H), 7.52-7.46 (m, 2H), 5.43 (dd, $J=8.8, 6.3$ Hz, 1H), 4.29 (dt, $J=2.8, 7.8$ Hz, 1H), 3.86 (ddd, $J=9.5, 8.0, 6.8$ Hz, 1H), 2.93 (dddd, $J=12.2, 9.2, 6.5, 3.0$ Hz, 1H), 2.19 (dddd, $J=12.1, 9.5, 7.7, 6.0$ Hz, 1H), 1.96 (br. s, 1H), 1.24-1.15 (m, 1H), 1.11-1.04 (m, 3H), 0.99 (ddd, $J=8.5, 4.8, 3.5$ Hz, 1H), 0.77-0.68 (m, 1H). LC-MS (Method A): $m/z = 257.1$ $[\text{M}+\text{H}]^+$, 0.89 min.

EXAMPLE 123

PREPARATION OF 2-CYCLOPROPYL-1-[(3S)-3-(5-FLUORO-6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE DIASTEREOISOMER 1 (217)




DIASTEREOISOMER 1

The title compound was isolated by chiral HPLC on a ChiraSpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 17 mL/min. ^1H NMR (400MHz, CDCl₃) δ 8.31 (s, 1H), 7.30 (dd, $J=9.8, 1.8$ Hz, 1H), 5.47 (dd, $J=8.8, 6.0$ Hz, 1H), 4.25 (dt, $J=3.4, 7.7$ Hz, 1H), 3.88 (ddd, $J=9.1, 8.1, 6.9$ Hz, 1H), 2.88 (dddd, $J=12.3, 8.9, 6.7, 3.4$ Hz, 1H), 2.52 (d, $J=2.8$ Hz, 3H), 2.39-2.29 (m, 2H), 1.22 (d, $J=7.0$ Hz, 3H), 1.09-0.99 (m, 1H), 0.61-0.38 (m, 3H), 0.20 (qd, $J=4.8, 9.4$ Hz, 1H). LC-MS (Method A): $m/z = 279.3$ $[\text{M}+\text{H}]^+$, 0.91 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 9.8 min.

EXAMPLE 124

PREPARATION OF 2-CYCLOPROPYL-1-[(3S)-3-(5-FLUORO-6-METHYLPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL]PROPAN-1-ONE DIASTEREOISOMER 2 (218)




DIASTEREOISOMER 2

The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 17 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.30 (s, 1H), 7.31--7.27 (m, 1H), 5.48 (dd, $J=8.8, 6.3$ Hz, 1H), 4.27 (dt, $J=2.9, 7.7$ Hz, 1H), 3.89 (ddd, $J=9.3, 8.1, 6.7$ Hz, 1H), 2.93-2.84 (m, 1H), 2.52 (d, $J=3.0$ Hz, 3H), 2.41-2.21 (m, 2H), 1.29 (d, $J=6.8$ Hz, 3H), 1.10-1.00 (m, 1H), 0.62-0.48 (m, 2H), 0.24-0.13 (m, 2H). LC-MS (Method A): $m/z = 279.3$ $[\text{M}+\text{H}]^+$, 0.90 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 80/20 % v/v, flow rate: 1.0 mL/min, retention time: 14.8 min.

EXAMPLE 125


PREPARATION OF 5-[(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-METHYLPYRIDINE-3-CARBONITRILE (219)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.65 (d, $J=2.5$ Hz, 1H), 7.82 (d, $J=2.3$ Hz, 1H), 5.42 (dd, $J=8.8, 7.0$ Hz, 1H), 4.33 (dt, $J=2.5, 7.9$ Hz, 1H), 3.93 (ddd, $J=10.1, 8.2, 6.3$ Hz, 1H), 2.89 (dddd, $J=12.4, 8.8, 6.3, 2.5$ Hz, 1H), 2.77 (s, 3H), 2.29 (tdd, $J=7.4, 12.2, 9.9$ Hz, 1H), 1.29 (s, 9H). LC-MS (Method A): $m/z = 274.1$ $[\text{M}+\text{H}]^+$, 0.90 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 24.0 min.

EXAMPLE 126

PREPARATION OF 5-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-METHYLPYRIDINE-3-CARBONITRILE (220)




The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18

mL/min. ¹H NMR (400MHz, CDCl₃) δ 8.64 (d, *J*=2.3 Hz, 1H), 7.79 (d, *J*=2.5 Hz, 1H), 6.30 (t, *J*=57.0 Hz, 1H), 5.43 (dd, *J*=8.8, 7.0 Hz, 1H), 4.38 (dt, *J*=2.8, 7.7 Hz, 1H), 3.99 (ddd, *J*=9.9, 8.2, 6.3 Hz, 1H), 2.94 (dddd, *J*=12.5, 8.9, 6.2, 2.8 Hz, 1H), 2.78 (s, 3H), 2.34 (tdd, *J*=7.4, 12.4, 9.8 Hz, 1H), 1.39-1.35 (m, 6H). LC-MS (Method A): *m/z* = 310.3 [M+H]⁺, 0.94 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μm column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 17.9 min.

EXAMPLE 127


PREPARATION OF 1-[(3S)-3-[6-(DiFLuOROMETIm)PYRIDIN-3-YL]-1,2-oxAz oLiDiN-2-YL]-3,3-D(FLUORO-2,2-DIMETHYLPROPAN -1-ONE (221)



The title compound was isolated by chiral HPLC on a Chiralpak AD-H (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.61 (d, $J=2.3$ Hz, 1H), 7.75 (dd, $J=8.0, 2.3$ Hz, 1H), 7.64 (d, $J=8.0$ Hz, 1H), 6.64 (t, $J=55.5$ Hz, 1H), 6.31 (t, $J=57.0$ Hz, 1H), 5.49 (dd, $J=8.5, 7.0$ Hz, 1H), 4.37 (dt, $J=2.6, 7.7$ Hz, 1H), 4.00 (ddd, $J=9.8, 8.2, 6.4$ Hz, 1H), 2.94 (dddd, $J=12.2, 9.2, 6.3, 2.9$ Hz, 1H), 2.37 (dddd, $J=12.3, 9.8, 7.5, 6.8$ Hz, 1H), 1.41- 1.36 (m, 6H). LC-MS (Method A): m/z = 321.0 $[\text{M}+\text{H}]^+$, 0.96 min. e.e.=100% as determined on a Chiralpak AD-H (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 7.4 min.

EXAMPLE 128


**PREPARATION OF 3-BROMO-5- [(3S)-2-(2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-
YL]BENZONITRILE (222)**



The title compound was isolated by chiral HPLC on a Chiralpak IC (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 17 mL/min. ^1H NMR (400MHz, CDCl_3) δ 7.71-7.64 (m, 2H), 7.53 (t, $J=1.3$ Hz, 1H), 5.37 (dd, $J=8.9, 6.9$ Hz, 1H), 4.29 (dt, $J=30, 7.5$ Hz, 1H), 3.90 (ddd, $J=10.2, 8.2, 6.3$ Hz, 1H), 2.87 (dddd, $J=12.5, 8.9, 6.3, 2.5$ Hz, 1H), 2.32-2.19 (m, 1H), 1.30 (s, 9H). LC-MS (Method A): $m/z = 336.0$ $[\text{M}+\text{H}]^+$, 1.19 min. e.e.=100% as determined on a Chiralpak IC (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 8.2 min.

EXAMPLE 129


PREPARATION OF 2,2-DIMETHYL-1-[(3S)-3-(6-NITROPYRIDIN-3-YL)-1,2-OXAZOLIDIN-2-YL] PROPAN-1-ONE (223)



The title compound was isolated by chiral HPLC on a Chiralpak IA (25 x 2.0 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 18 mL/min. ^1H NMR (400MHz, CDCl_3) δ 8.61 (d, $J=2.3$ Hz, 1H), 8.24 (d, $J=8.3$ Hz, 1H), 7.96 (dd, $J=8.3, 2.3$ Hz, 1H), 5.56 (dd, $J=9.0, 7.0$ Hz, 1H), 4.36 (dt, $J=2.4, 8.0$ Hz, 1H), 3.98 (ddd, $J=10.0, 8.3, 6.3$ Hz, 1H), 3.02-2.91 (m, 1H), 2.34 (tdd, $J=7.4, 12.3, 10.0$ Hz, 1H), 1.31 (s, 9H). LC-MS (Method A): $m/z = 280.3$ $[\text{M}+\text{H}]^+$, 0.92 min. e.e.=100% as determined on a Chiralpak IA (25 x 0.46 cm), 5 μ m column using a mobile phase of n-hexane/(ethanol + 0.1% isopropylamine) 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 11.3 min.

EXAMPLE 130

5-[(3S)-2-(3,3-D(FLUORO-2,2-DIMETHYLBUTANOYL)-1,2-OXAZOLIDIN-3-YL)PYRIDINE-3-CARBONYL] FURAN-2-CARBOXYLIC ACID (224)




^1H NMR (400MHz, $\text{DMSO}-d_6$) δ 8.95 (d, $J=1.8$ Hz, 1H), 8.76 (d, $J=2.3$ Hz, 1H), 8.19 (t, $J=2.0$ Hz, 1H), 5.45 (dd, $J=8.5, 7.0$ Hz, 1H), 4.35 (dt, $J=2.4, 7.8$ Hz, 1H), 3.94 (ddd, $J=9.9, 8.0, 6.4$ Hz, 1H), 3.01-2.82 (m, 1H), 2.27 (tdd, $J=7.4, 12.2, 9.9$ Hz, 1H), 1.67 (t, $J=19.7$ Hz, 3H), 1.42 (s, 3H),

1.37 (s, 3H). LC-MS (Method A): $m/z = 310.2$ $[M+H]^+$, 0.93 min.

EXAMPLE 131

PREPARATION OF 5-[(3S)-2-(PROPANE-2-SULFONYL)-1,2-OXAZOLIDIN-3-YL]PYRIDINE-3-CARBONITRILE (225)



5


The title compound was isolated by chiral HPLC on a Chiralpak IA (15 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (400 MHz, DMSO- d_6) δ 8.97 (d, $J = 1.6$ Hz, 1H), 8.87 (d, $J = 2.0$ Hz, 1H), 8.29-8.24 (m, 1H), 5.52-5.46 (m, 1H), 4.33-4.12 (m, 2H), 3.80-3.66 (m, 1H), 2.96-2.82 (m, 1H), 2.42-2.30 (m, 1H), 1.36-1.28 (m, 6H). LC-MS (Method F): $m/z = 281.9$ $[M+H]^+$, 1.024 min. e.e. = 99.3% as determined on a Chiralpak AS-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 3.39 min.

10

EXAMPLE 132

PREPARATION OF 3-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-5-FLUOROBENZONITRILE (226)

15




The title compound was isolated by chiral HPLC on a Lux 5 μ Cellulose-4, AXIA Packed (25 x 2.12 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 70/30 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, CD_3OD) δ 7.50-7.46 (m, 2H), 7.40-7.35 (m, 1H), 6.39 (t, $J = 57.0$ Hz, 1H), 5.45-5.40 (m, 1H), 4.43-4.37 (m, 1H), 4.06-3.98 (m, 1H), 3.00-2.95 (m, 1H), 2.34-2.27 (m, 1H), 1.36 (s, 3H), 1.34 (s, 3H). LC-MS (Method D): $m/z = 313.1$ $[M+H]^+$, 1.796 min. e.e. = 99.9% as determined on a Lux Cellulose-4 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 70/30 % v/v, flow rate: 1.0 mL/min, retention time: 1.543 min.

20

EXAMPLE 133


PREPARATION OF 5-[(3S)-2-(3,3-DIFLUORO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-(DIFLUOROMETHYL)BENZONITRILE (227)



5 The title compound was isolated by chiral HPLC on a Phenomenex Lux (25 x 2.1 cm), 5 μ m column using a mobile phase of n-hexane/ethanol 75/25 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.92 (s, 1H), 7.84 (d, J = 8.1 Hz, 1H), 7.73 (d, J = 8.2 Hz, 1H), 7.26 (t, J = 54.2 Hz, 1H), 6.42 (t, J = 56.8 Hz, 1H), 5.50-5.40 (m, 1H), 4.45-4.31 (m, 1H), 4.10-3.92 (m, 1H), 3.03-2.90 (m, 1H), 2.33-2.16 (m, 1H), 1.30 (s, 3H), 1.29 (s, 3H). LC-MS (Method D): m/z = 345.1 $[\text{M}+\text{H}]^+$, 1.804 min. e.e.=99.9% as determined on a Chiralpak IF-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 3.05 min.




EXAMPLE 134

PREPARATION OF 5-[(3S)-2-(3-CYANO-2,2-DIMETHYLPROPANOYL)-1,2-OXAZOLIDIN-3-YL]-2-(DIFLUOROMETHYL)BENZONITRILE (228)









15 The title compound was isolated by chiral HPLC on a Chiralpak IA (15 x 2.1 cm), 5 μ m column using a mobile phase of n-hexane/ 2-propanol 50/50 % v/v, flow rate: 20 mL/min. ^1H NMR (300 MHz, CD_3OD) δ 7.82 (s, 1H), 7.80-7.71 (m, 2H), 7.01 (t, J = 54.6 Hz, 1H), 5.52-5.41 (m, 1H), 4.46-4.34 (m, 1H), 4.11-4.00 (m, 1H), 3.10-2.93 (m, 1H), 2.91-2.70 (m, 2H), 2.37-2.21 (m, 1H), 1.46 (s, 3H), 1.43 (s, 3H). LC-MS (Method S): m/z = 334.4 $[\text{M}+\text{H}]^+$, 2.349 min. e.e.=99.9% as determined on a Chiralpak IA-3 (5 x 0.46 cm), 3 μ m column using a mobile phase of (n-hexane + 0.1% diethylamine)/ethanol 50/50 % v/v, flow rate: 1.0 mL/min, retention time: 1.60 min.




The following compounds were prepared according to the Examples above and/or general procedures described herein.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
28	4-[(3S)-3-(4-Fluorophenyl)-1,2-oxazolidine-2-carbonyl]oxane-4-carbonitrile		¹ H NMR (500 MHz, CDCl ₃) δ 7.32–7.24 (m, 2H), 7.05 (t, J=8.7 Hz, 2H), 5.39 (dd, J=8.7, 6.7 Hz, 1H), 4.38 (td, J=8.1, 2.7 Hz, 1H), 4.17 (ddd, J=9.8, 8.3, 6.9 Hz, 1H), 4.05–3.91 (m, 2H), 3.85–3.78 (m, 2H), 2.93 (dddd, J=12.2, 9.1, 6.5, 2.9 Hz, 1H), 2.47–2.33 (m, 2H), 2.19 (ddd, J=13.9, 11.6, 4.4 Hz, 1H), 2.07 (ddd, J=13.7, 11.4, 4.3 Hz, 1H), 2.00 (dq, J=2.3, 14.0 Hz, 1H).	(Method B): m/z = 304.0 [M+H] ⁺ , 0.93 min
29	(3S)-3-(4-fluorophenyl)-2-[1-(trifluoromethyl)cyclopentanecarbonyl]-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.26 (m, 2H), 7.07–7.00 (m, 2H), 5.41 (dd, J=8.8, 6.8 Hz, 1H), 4.29 (td, J=8.0, 2.8 Hz, 1H), 4.00–3.91 (m, 1H), 2.89–2.80 (m, 1H), 2.47–2.28 (m, 3H), 2.24–2.11 (m, 2H), 1.80–1.60 (m, 4H).	(Method A): m/z = 332.2 [M+H] ⁺ , 1.24 min
30	1-[(3R)-3-(2,3-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.12–7.00 (m, 3H), 5.68 (dd, J=8.8, 6.8 Hz, 1H), 4.27 (td, J=8.0, 2.8 Hz, 1H), 3.96–3.89 (m, 1H), 2.97–2.88 (m, 1H), 2.33–2.22 (m, 1H), 1.33 (s, 9H).	(Method A): m/z = 270.3 [M+H] ⁺ , 1.16 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
31	1-[(3S)-3-(2,3-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.12–7.00 (m, 3H), 5.68 (dd, <i>J</i> =8.8, 6.8 Hz, 1H), 4.27 (td, <i>J</i> =8.0, 2.8 Hz, 1H), 3.96–3.89 (m, 1H), 2.97–2.88 (m, 1H), 2.33–2.22 (m, 1H), 1.33 (s, 9H).	(Method A): <i>m/z</i> = 270.3 [M+H] ⁺ , 1.16 min
32	(3S)-2-{Bicyclo[1.1.1]pentane-1-carbonyl}-3-(4-fluorophenyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.34–7.25 (m, 2H), 7.07–6.99 (m, 2H), 5.42 (dd, <i>J</i> =8.7, 5.9 Hz, 1H), 4.25 (td, <i>J</i> =7.8, 3.1 Hz, 1H), 3.89–3.77 (m, 1H), 2.82 (dddd, <i>J</i> =12.2, 8.9, 7.0, 3.3 Hz, 1H), 2.49 (br s, 1H), 2.33 (dddd, <i>J</i> =12.3, 9.3, 7.7, 5.8 Hz, 1H), 2.19 (br s, 6H).	LC-MS (Method A): <i>m/z</i> = 262.2 [M+H] ⁺ , 1.02 min
33	(3S)-3-(4-Fluorophenyl)-2-(3-methyloxetane-3-carbonyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.23 (m, 2H), 7.12–7.00 (m, 2H), 5.39 (dd, <i>J</i> =8.5, 6.0 Hz, 1H), 5.08 (d, <i>J</i> =6.3 Hz, 1H), 4.91 (d, <i>J</i> =6.0 Hz, 1H), 4.41 (d, <i>J</i> =6.3 Hz, 1H), 4.38 (d, <i>J</i> =6.5 Hz, 1H), 4.25 (td, <i>J</i> =7.9, 3.8 Hz, 1H), 3.93 (ddd, <i>J</i> =9.0, 8.0, 6.8 Hz, 1H), 2.92–2.82 (m, 1H), 2.42–2.31 (m, 1H), 1.71 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 266.2 [M+H] ⁺ , 0.83 min
34	(3S)-2-(1-Fluorocyclopropanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.37–7.30 (m, 2H), 7.10–7.00 (m, 2H), 5.60–5.48 (m, 1H), 4.31 (dd, <i>J</i> =15.4, 4.6 Hz, 1H), 4.10–4.02 (m, 1H), 2.95–2.86 (m, 1H), 2.40 (dtd, <i>J</i> =12.5, 7.8, 5.0 Hz, 1H), 1.50–1.39 (m, 1H), 1.36–1.11 (m, 3H).	LC-MS (Method A): <i>m/z</i> = 254.1 [M+H] ⁺ , 0.91 min





Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
35	(3S)-3-(4-Fluorophenyl)-2-[3-methyloxane-3-carbonyl]-1,2-oxazolidine diastereoisomer 2 (Second eluting diastereoisomer)	 DIASTEREOISOMER 2	¹ H NMR (400 MHz, CDCl ₃) δ 7.31-7.26 (m, 5H) 7.02 (t, J=8.8 Hz, 2H) 5.50-5.38 (m, 1H) 4.24 (s, 2H) 3.96-3.86 (m, 1H) 3.64 (d, J=5.3 Hz, 2H) 3.51 (d, J=11.5 Hz, 1H) 2.87-2.75 (m, 1H) 2.36-2.24 (m, 1H) 2.22-2.12 (m, 1H) 1.71-1.59 (m, 3H) 1.29 (s, 3H)	
36	(3S)-3-(4-Fluorophenyl)-2-[3-methyloxane-3-carbonyl]-1,2-oxazolidine diastereoisomer 1 (First eluting diastereoisomer)	 DIASTEREOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 7.29-7.25 (m, 2H), 7.04-6.98 (m, 2H), 5.43 (dd, J=8.8, 6.6 Hz, 1H), 4.27 (td, J=7.7, 2.5 Hz, 1H), 4.18 (dd, J=11.5, 1.4 Hz, 1H), 3.92 (ddd, J=9.8, 8.0, 6.6 Hz, 1H), 3.73-3.64 (m, 1H), 3.61-3.53 (m, 1H), 3.37 (d, J=1.4 Hz, 1H), 2.82 (dddd, J=12.1, 9.1, 6.4, 2.9 Hz, 1H), 2.36-2.21 (m, 2H), 1.81-1.68 (m, 1H), 1.68-1.59 (m, 2H), 1.26 (s, 3H).	
37	(3S)-3-(4-Fluorophenyl)-2-[1-(trifluoromethyl)cyclopropanecarbonyl]-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.32-7.22 (m, 2H), 7.10-7.00 (m, 2H), 5.42 (dd, J=8.8, 6.0 Hz, 1H), 4.31 (td, J=7.7, 3.3 Hz, 1H), 3.99 (ddd, J=9.2, 8.1, 6.8 Hz, 1H), 2.87 (dddd, J=12.3, 8.8, 6.9, 3.5 Hz, 1H), 2.37 (dddd, J=12.3, 9.2, 7.7, 6.0 Hz, 1H), 1.41-1.16 (m, 4H).	(Method A): m/z = 304.1 [M+H] ⁺ , 1.06 min




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
38	(3S)-2-(3-Ethyloxetane-3-carbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ ppm 7.32–7.25 (m, 2H) 7.05 (t, <i>J</i> =8.7 Hz, 2H) 5.44–5.38 (m, 1H) 5.07–5.02 (m, 1H) 4.83 (d, <i>J</i> =6.5 Hz, 1H) 4.45 (d, <i>J</i> =6.5 Hz, 1H) 4.39 (d, <i>J</i> =6.5 Hz, 1H) 4.26–4.19 (m, 1H) 3.94–3.84 (m, 1H) 2.92–2.78 (m, 1H) 2.42–2.29 (m, 1H) 2.24–2.05 (m, 2H) 0.94 (t, <i>J</i> =7.5 Hz, 3H).	(Method A): <i>m/z</i> = 280.1 [M+H] ⁺ , 0.97 min
39	(3S)-3-(4-Fluorophenyl)-2-[1-(trifluoromethyl)cyclobutanecarbonyl]-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.26 (m, 2H), 7.10–7.01 (m, 2H), 5.41 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.26 (td, <i>J</i> =7.8, 3.5 Hz, 1H), 3.98–3.90 (m, 1H), 2.90–2.73 (m, 2H), 2.67–2.49 (m, 3H), 2.40–2.30 (m, 1H), 2.17–2.03 (m, 1H), 1.92–1.80 (m, 1H).	(Method A): <i>m/z</i> = 318.2 [M+H] ⁺ , 1.15 min
40	(3S)-3-(4-Fluorophenyl)-2-(3-methyloxolane-3-carbonyl)-1,2-oxazolidine diastereoisomer 1 (First eluting diastereoisomer)	 DIASTEREOISOMER 1	¹ H NMR (500 MHz, CDCl ₃) δ 7.32–7.22 (m, 2H), 7.03 (t, <i>J</i> =8.8 Hz, 2H), 5.36 (dd, <i>J</i> =8.6, 6.6 Hz, 1H), 4.28 (td, <i>J</i> =7.8, 3.4 Hz, 1H), 3.98 (d, <i>J</i> =9.3 Hz, 1H), 3.95–3.87 (m, 2H), 3.83 (td, <i>J</i> =8.3, 5.9 Hz, 1H), 3.77 (d, <i>J</i> =8.8 Hz, 1H), 2.84 (dddd, <i>J</i> =12.2, 9.1, 6.4, 3.2 Hz, 1H), 2.61 (dt, <i>J</i> =12.7, 7.6 Hz, 1H), 2.39–2.29 (m, 1H), 1.80 (dt, <i>J</i> =12.8, 6.5 Hz, 1H), 1.46 (s, 3H).	(Method B): <i>m/z</i> = 279.0 [M+H] ⁺ , 0.92 min




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
41	(3S)-3-(4-Fluorophenyl)-2-(3-methyloxolane-3-carbonyl)-1,2-oxazolidine diastereoisomer 2 (Second eluting diastereoisomer)	 DIASTEREOISOMER 2	¹ H NMR (500 MHz, CDCl ₃) δ 7.31-7.24 (m, 2H), 7.03 (t, J=8.8 Hz, 2H), 5.38 (dd, J=8.6, 6.6 Hz, 1H), 4.28 (td, J=7.7, 3.2 Hz, 1H), 4.18 (d, J=8.8 Hz, 1H), 3.73 (d, J=8.8 Hz, 1H), 3.97-3.78 (m, 3H), 2.84 (ddt, J=15.4, 9.0, 3.2 Hz, 1H), 2.48 (dt, J=12.7, 7.6 Hz, 1H), 2.39-2.29 (m, 1H), 1.85 (ddd, J=12.8, 7.2, 5.9 Hz, 1H), 1.43 (s, 3H).	(Method B): m/z = 279.0 [M+H] ⁺ , 0.91 min
42	(3S)-2-(1,3-Dimethylazetidine-3-carbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine		¹ H NMR (400 MHz, DMSO-d ₆) δ 7.36-7.27 (m, 2H), 7.22-7.13 (m, 2H), 5.29 (dd, J=8.5, 6.8 Hz, 1H), 4.23 (td, J=7.8, 3.3 Hz, 1H), 3.86 (ddd, J=9.6, 8.0, 6.5 Hz, 1H), 3.23 (d, J=7.0 Hz, 1H), 3.10 (s, 2H), 3.06 (d, J=7.0 Hz, 1H), 2.88 (dddd, J=12.0, 9.0, 6.3, 3.0 Hz, 1H), 2.22-2.11 (m, 4H), 1.49 (s, 3H).	LC-MS (Method A): m/z = 279.1 [M+H] ⁺ , 0.47 min
43	(3S)-3-(4-Fluorophenyl)-2-(3-methylazetidine-3-carbonyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.35-7.25 (m, 2H), 7.12-6.99 (m, 2H), 5.39 (dd, J=8.7, 6.1 Hz, 1H), 4.24 (td, J=7.8, 3.6 Hz, 1H), 4.17 (d, J=8.5 Hz, 1H), 4.00 (d, J=8.3 Hz, 1H), 3.96-3.86 (m, 1H), 3.42-3.33 (m, 2H), 2.92-2.80 (m, 1H), 2.42-2.28 (m, 1H), 1.68 (s, 3H).	LC-MS (Method A): m/z = 265.2 [M+H] ⁺ , 0.43 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
44	tert-Butyl 3-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidine-2-carbonyl]-3-methylazetidine-1-carboxylate		¹ H NMR (400 MHz, CDCl ₃) δ 7.34–7.24 (m, 2H), 7.10–7.01 (m, 2H), 5.37 (dd, <i>J</i> =8.7, 6.1 Hz, 1H), 4.35 (d, <i>J</i> =8.8 Hz, 1H), 4.27 (td, <i>J</i> =7.7, 3.5 Hz, 1H), 4.16 (d, <i>J</i> =8.8 Hz, 1H), 3.96 (ddd, <i>J</i> =9.0, 8.0, 6.8 Hz, 1H), 3.71 (d, <i>J</i> =8.8 Hz, 1H), 3.67 (d, <i>J</i> =9.0 Hz, 1H), 2.92–2.82 (m, 1H), 2.42–2.31 (m, 1H), 1.62 (s, 3H), 1.46 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 365.3 [M+H] ⁺ , 1.13 min
45	1-[(3R)-3-(2-Fluoropyridin-4-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =5.3 Hz, 1H), 7.14–7.10 (m, 1H), 6.87 (s, 1H), 5.43 (dd, <i>J</i> =6.8, 8.8 Hz, 1H), 4.29 (td, <i>J</i> =7.8, 2.2 Hz, 1H), 3.96–3.89 (m, 1H), 2.93–2.84 (m, 1H), 2.33–2.22 (m, 1H), 1.32 (s, 9H).	(Method A): <i>m/z</i> = 253.2 [M+H] ⁺ , 0.93 min
46	1-[(3S)-3-(2-Fluoropyridin-4-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =5.3 Hz, 1H), 7.14–7.10 (m, 1H), 6.87 (s, 1H), 5.43 (dd, <i>J</i> =6.8, 8.8 Hz, 1H), 4.29 (td, <i>J</i> =7.8, 2.2 Hz, 1H), 3.96–3.89 (m, 1H), 2.93–2.84 (m, 1H), 2.33–2.22 (m, 1H), 1.32 (s, 9H).	(Method A): <i>m/z</i> = 253.2 [M+H] ⁺ , 0.93 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
47	1-[(3S)-3-(4-Chlorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.34–7.29 (m, 2H), 7.28–7.23 (m, 2H), 5.39 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.27 (td, <i>J</i> =7.7, 2.6 Hz, 1H), 3.90 (ddd, <i>J</i> =9.9, 8.0, 6.4 Hz, 1H), 2.82 (dddd, <i>J</i> =12.2, 9.1, 6.3, 2.8 Hz, 1H), 2.29 (dddd, <i>J</i> =12.3, 9.9, 7.7, 6.8 Hz, 1H), 1.30 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 268.2 [M+H] ⁺ , 1.11 min
48	1-[(3R)-3-(4-Chlorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.35–7.29 (m, 2H), 7.28–7.23 (m, 2H), 5.39 (dd, <i>J</i> =8.9, 6.7 Hz, 1H), 4.27 (td, <i>J</i> =7.8, 2.3 Hz, 1H), 3.90 (ddd, <i>J</i> =9.8, 8.0, 6.5 Hz, 1H), 2.82 (dddd, <i>J</i> =12.1, 9.1, 6.4, 2.8 Hz, 1H), 2.29 (dddd, <i>J</i> =12.3, 9.9, 7.7, 6.5 Hz, 1H), 1.30 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 268.2 [M+H] ⁺ , 1.12 min
49	<i>tert</i> -Butyl 3-(4-chlorophenyl)-1,2-oxazolidine-2-carboxylate		¹ H NMR (400 MHz, CDCl ₃) δ 7.36–7.28 (m, 4H), 5.19 (dd, <i>J</i> =5.5, 8.5 Hz, 1H), 4.20 (dt, <i>J</i> =3.3, 7.8 Hz, 1H), 3.91 (dt, <i>J</i> =7.0, 8.5 Hz, 1H), 2.79 (dddd, <i>J</i> =3.6, 7.0, 8.7, 12.2 Hz, 1H), 2.33–2.22 (m, 1H), 1.48 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 284.1 [M+H] ⁺ , 1.13 min
50	1-[(3S)-3-(3,4-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.18–7.07 (m, 2H), 7.06–7.00 (m, 1H), 5.37 (dd, <i>J</i> =8.7, 6.9 Hz, 1H), 4.27 (td, <i>J</i> =7.8, 2.6 Hz, 1H), 3.89 (ddd, <i>J</i> =9.8, 8.0, 6.5 Hz, 1H), 2.88–2.76 (m, 1H), 2.27 (dddd, <i>J</i> =12.3, 9.9, 7.5, 6.7 Hz, 1H), 1.30 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 270.1 [M+H] ⁺ , 1.09 min




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
51	1-[(3R)-3-(3,4-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.18–7.08 (m, 2H), 7.07–7.00 (m, 1H), 5.37 (dd, <i>J</i> =8.8, 6.8 Hz, 1H), 4.28 (td, <i>J</i> =7.8, 2.8 Hz, 1H), 3.89 (ddd, <i>J</i> =10.0, 8.1, 6.5 Hz, 1H), 2.82 (dddd, <i>J</i> =12.1, 9.1, 6.4, 2.8 Hz, 1H), 2.28 (dddd, <i>J</i> =12.3, 9.9, 7.7, 6.5 Hz, 1H), 1.30 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 270.1 [M+H] ⁺ , 1.09 min.
52	(3S)-2-(Oxetane-2-carbonyl)-3-phenyl-1,2-oxazolidine diastereoisomer 2 (Second eluting diastereoisomer)	 DIASTEREISOISOMER 2	¹ H NMR (400 MHz, CDCl ₃) δ 7.40–7.29 (m, 5H), 5.63–5.45 (m, 2H), 4.80–4.66 (m, 2H), 4.26 (dt, <i>J</i> =7.5, 4.5 Hz, 1H), 3.92 (br s, 1H), 3.02–2.76 (m, 3H), 2.47–2.35 (m, 1H).	(Method A): <i>m/z</i> = 234.1 [M+H] ⁺ , 0.73 min
53	(3S)-2-(Oxetane-2-carbonyl)-3-phenyl-1,2-oxazolidine diastereoisomer 1 (First eluting diastereoisomer)	 DIASTEREISOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 7.40–7.29 (m, 5H), 5.58–5.48 (m, 2H), 4.73 (t, <i>J</i> =7.5 Hz, 2H), 4.24 (dt, <i>J</i> =7.7, 3.6 Hz, 1H), 3.91 (q, <i>J</i> =7.9 Hz, 1H), 3.07–2.82 (m, 3H), 2.45–2.35 (m, 1H).	(Method A): <i>m/z</i> = 234.1 [M+H] ⁺ , 0.75 min
54	(3S)-3-Phenyl-2-[1-(trifluoromethyl)cyclopropanecarbonyl]-1,2-oxazolidine		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.39–7.33 (m, 2H), 7.30–7.24 (m, 3H), 5.34 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.29 (dt, <i>J</i> =3.3, 7.7 Hz, 1H), 3.94 (ddd, <i>J</i> =9.7, 7.9, 6.5 Hz, 1H), 2.91 (s, 1H), 2.24–2.12 (m, 1H), 1.29 (dd, <i>J</i> =7.2, 2.4 Hz, 4H).	(Method A): <i>m/z</i> = 286.2 [M+H] ⁺ , 1.08 min




Comp. No.	Name	Structure	^1H NMR	M+H ⁺
55	3,3,3-Trifluoro-2,2-dimethyl-1-[(3S)-3-phenyl-1,2-oxazolidin-2-yl]propan-1-one		^1H NMR (400 MHz, CDCl_3) δ 7.38 - 7.28 (m, 5H), 5.45 (dd, $J=8.8, 6.5$ Hz, 1H), 4.31 (dt, $J=8.0, 2.8$ Hz, 1H), 3.94 (ddd, $J=9.7, 8.0, 6.7$ Hz, 1H), 2.84 (dddd, $J=12.0, 9.1, 6.5, 3.0$ Hz, 1H), 2.36 (dddd, $J=12.3, 9.8, 7.8, 6.5$ Hz, 1H), 1.58 (s, 3H), 1.57 - 1.56 (m, 3H).	(Method A): m/z = 288.1 [M+H] ⁺ , 1.19 min
56	2-Fluoro-2-methyl-1-[(3S)-3-phenyl-1,2-oxazolidin-2-yl]propan-1-one		^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.39-7.34 (m, 2H), 7.30-7.25 (m, 3H), 5.37 (dd, $J=8.5, 6.5$ Hz, 1H), 4.29 (dt, $J=7.7, 3.3$ Hz, 1H), 3.89 (ddd, $J=9.5, 7.9, 6.7$ Hz, 1H), 2.91 (dddd, $J=12.2, 9.1, 6.3, 3.0$ Hz, 1H), 2.19 (dddd, $J=12.1, 9.6, 7.7, 6.5$ Hz, 1H), 1.61 (s, 3H), 1.55 (s, 3H).	(Method A): m/z = 238.2 [M+H] ⁺ , 0.95 min
57	(3S)-2-(3,3-difluoro-1-methylcyclobutanecarbonyl)-3-phenyl-1,2-oxazolidine		^1H NMR (400 MHz, CDCl_3) δ 7.41- 7.29 (m, 5H), 5.41 (dd, $J=8.9, 6.1$ Hz, 1H), 4.28 (dt, $J=7.9, 4.0$ Hz, 1H), 3.96 (ddd, $J=9.0, 8.0, 6.8$ Hz, 1H), 3.25-3.11 (m, 1H), 3.02-2.82 (m, 2H), 2.59-2.34 (m, 3H), 1.58 (s, 3H).	(Method A): m/z = 282.3 [M+H] ⁺ , 1.11 min






Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
58	2,2-Dimethyl-1-[3-(pyridin-3-yl)-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.50–8.45 (m, 2H), 7.64 (dt, <i>J</i> =8.0, 1.8 Hz, 1H), 7.41–7.34 (m, 1H), 5.36 (dd, <i>J</i> =8.8, 7.0 Hz, 1H), 4.32 (td, <i>J</i> =7.7, 2.5 Hz, 1H), 3.88 (ddd, <i>J</i> =10.0, 8.0, 6.4 Hz, 1H), 2.95–2.84 (m, 1H), 2.19 (ddt, <i>J</i> =12.0, 9.9, 7.3 Hz, 1H), 1.21 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 235.2 [M+H] ⁺ , 0.50 min
59	5-[(3R)-3-(4-fluorophenyl)-1,2-oxazolidine-2-carbonyl]-1-methyl-1H-pyrrole-2-carbonitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.41–7.32 (m, 2H), 7.07 (t, <i>J</i> =8.6 Hz, 2H), 7.00 (d, <i>J</i> =4.3 Hz, 1H), 6.78 (dd, <i>J</i> =4.3, 0.7 Hz, 1H), 5.63–5.54 (m, 1H), 4.35 (td, <i>J</i> =7.8, 3.5 Hz, 1H), 4.04–3.94 (m, 4H), 2.95–2.85 (m, 1H) 2.48–2.34 (m, 1H).	(Method A): <i>m/z</i> = 300.2 [M+H] ⁺ , 1.05 min
60	5-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidine-2-carbonyl]-1-methyl-1H-pyrrole-2-carbonitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.41–7.31 (m, 2H), 7.12–7.01 (m, 2H), 7.00 (d, <i>J</i> =4.3 Hz, 1H), 6.77 (d, <i>J</i> =4.3 Hz, 1H), 5.58 (dd, <i>J</i> =8.7, 6.1 Hz, 1H), 4.39–4.29 (m, 1H), 4.01 (s, 3H), 4.02–3.93 (m, 1H), 2.89 (dddd, <i>J</i> =12.2, 8.7, 6.7, 3.5 Hz, 1H), 2.40 (dddd, <i>J</i> =12.4, 9.3, 7.5, 6.1 Hz, 1H).	(Method A): <i>m/z</i> = 300.2 [M+H] ⁺ , 1.05 min





Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
64	1-[(3S)-3-(4-Fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.34–7.27 (m, 2H), 7.09–6.98 (m, 2H), 5.40 (dd, <i>J</i> =8.8, 5.9 Hz, 1H), 4.32–4.22 (m, 1H), 3.90 (ddd, <i>J</i> =9.2, 8.0, 6.8 Hz, 1H), 3.08–2.99 (m, 1H), 2.83 (dddd, <i>J</i> =12.3, 8.8, 6.8, 3.5 Hz, 1H), 2.35 (dddd, <i>J</i> =12.2, 9.2, 7.6, 5.9 Hz, 1H), 1.22 (d, <i>J</i> =6.9 Hz, 3H), 1.12 (d, <i>J</i> =6.8 Hz, 3H).	(Method A): <i>m/z</i> = 238.2 [M+H] ⁺ , 0.97 min.
65	1-[(3R)-3-(4-Fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.35–7.25 (m, 2H), 7.09–6.98 (m, 2H), 5.40 (dd, <i>J</i> =8.8, 5.9 Hz, 1H), 4.31–4.22 (m, 1H), 3.90 (ddd, <i>J</i> =9.2, 8.0, 6.8 Hz, 1H), 3.08–2.99 (m, 1H), 2.83 (dddd, <i>J</i> =12.3, 8.8, 6.8, 3.5 Hz, 1H), 2.34 (dddd, <i>J</i> =12.3, 9.2, 7.5, 5.9 Hz, 1H), 1.22 (d, <i>J</i> =6.9 Hz, 3H), 1.12 (d, <i>J</i> =6.9 Hz, 3H).	(Method A): <i>m/z</i> = 238.2 [M+H] ⁺ , 0.96 min.
63	1-[(3R)-3-(2-Fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.13 (dt, <i>J</i> =4.8, 1.5 Hz, 1H), 7.72–7.65 (m, 1H), 7.18 (td, <i>J</i> =5.0, 2.4 Hz, 1H), 5.59 (dd, <i>J</i> =8.7, 6.9 Hz, 1H), 4.26 (td, <i>J</i> =7.9, 2.8 Hz, 1H), 3.93 (ddd, <i>J</i> =9.8, 8.0, 6.5 Hz, 1H), 3.00–2.90 (m, 1H), 2.35–2.21 (m, 1H), 1.34 (s, 9H).	




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
66	1-[(3S)-3-(2-Fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.12 (dt, <i>J</i> =4.8, 1.5 Hz, 1H), 7.68 (ddt, <i>J</i> =9.7, 6.7, 1.3 Hz, 1H), 7.18 (ddd, <i>J</i> =7.2, 5.1, 1.8 Hz, 1H), 5.59 (dd, <i>J</i> =8.9, 6.7 Hz, 1H), 4.29–4.22 (m, 1H), 3.93 (ddd, <i>J</i> =9.8, 8.0, 6.5 Hz, 1H), 3.00–2.87 (m, 1H), 2.34–2.22 (m, 1H), 1.33 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 253.2 [M+H] ⁺ , 0.92 min
67	1-[(3R)-3-(2,6-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.26–7.16 (m, 1H), 6.92–6.83 (m, 2H), 5.68 (t, <i>J</i> =8.8, 1H), 4.40 (td, <i>J</i> =7.8, 1.8 Hz, 1H), 3.95–3.87 (m, 1H), 2.81–2.73 (m, 1H), 2.56–2.45 (m, 1H), 1.27 (s, 9H).	(Method A): <i>m/z</i> = 270.2 [M+H] ⁺ , 1.09 min
68	1-[(3S)-3-(2,6-Difluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.26–7.16 (m, 1H), 6.92–6.83 (m, 2H), 5.68 (t, <i>J</i> =8.8, 1H), 4.40 (td, <i>J</i> =7.8, 1.8 Hz, 1H), 3.95–3.87 (m, 1H), 2.81–2.73 (m, 1H), 2.56–2.45 (m, 1H), 1.27 (s, 9H).	(Method A): <i>m/z</i> = 270.2 [M+H] ⁺ , 1.09 min
69	(3S)-2-(4-methyloxane-4-carbonyl)-3-phenyl-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.41–7.23 (m, 5H) 5.48 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.27 (td, <i>J</i> =7.5, 3.3 Hz, 1H) 3.91 (ddd, <i>J</i> =9.5, 8.0, 6.5 Hz, 1H) 3.85–3.72 (m, 2H) 3.59 (dddd, <i>J</i> =11.9, 9.3, 6.3, 2.9 Hz, 2H) 2.90–2.77 (m, 1H) 2.47–2.27 (m, 2H) 2.21 (ddt, <i>J</i> =13.6, 4.7, 2.6 Hz, 1H) 1.86–1.46 (m, 2H) 1.34 (s, 3H).	LC-MS (Method B): <i>m/z</i> = 275.00 [M+H] ⁺ , 0.94 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
75	(1r,3r)-3-[(3S)-3-phenyl-1,2-oxazolidine-2-carbonyl]cyclobutan-1-ol		¹ H NMR (400 MHz, CDCl ₃) δ 7.39–7.29 (m, 5H), 5.49–5.36 (m, 1H), 4.58–4.49 (m, 1H), 4.24 (dt, <i>J</i> =7.8, 3.8 Hz, 1H), 3.91–3.82 (m, 1H), 3.49 (br s, 1H), 2.84 (dddd, <i>J</i> =12.3, 8.8, 6.8, 3.8 Hz, 1H), 2.71–2.55 (m, 2H), 2.38 (dddd, <i>J</i> =12.3, 9.0, 7.5, 5.8 Hz, 1H), 2.32–2.23 (m, 1H), 2.18 (br s, 1H), 1.78 (d, <i>J</i> =5.3 Hz, 1H).	(Method A): <i>m/z</i> = 248.2 [M+H] ⁺ , 0.71 min
76	(3S)-2-[1-(methoxymethyl)cyclopropanecarbonyl]-3-phenyl-1,2-oxazolidine		¹ H NMR (400 MHz, , DMSO- <i>d</i> ₆) δ 7.39–7.22 (m, 5H), 5.36 (dd, <i>J</i> =8.7, 6.1 Hz, 1H), 4.24 (dt, <i>J</i> =7.8, 3.0 Hz, 1H), 3.89–3.80 (m, 2H), 3.35–3.33 (m, 1H), 3.23 (s, 3H), 2.86 (tdd, <i>J</i> =15.2, 9.0, 3.2 Hz, 1H), 2.16 (dddd, <i>J</i> =12.1, 9.6, 7.7, 6.3 Hz, 1H), 1.07–1.00 (m, 1H), 0.91 (dt, <i>J</i> =6.0, 2.8 Hz, 1H), 0.75–0.66 (m, 2H).	(Method A): <i>m/z</i> = 262.3 [M+H] ⁺ , 0.90 min
77	(3S)-2-(1-methylcyclobutanecarbonyl)-3-phenyl-1,2-oxazolidine		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.39–7.21 (m, 5H), 5.29 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.21 (dt, <i>J</i> =7.8, 3.0 Hz, 1H), 3.84 (ddd, <i>J</i> =9.7, 7.9, 6.5 Hz, 1H), 2.94 – 2.82 (m, 1H), 2.48–2.41 (m, 1H), 2.37–2.28 (m, 1H), 2.15 (dddd, <i>J</i> =12.1, 9.7, 7.5, 6.5 Hz, 1H), 2.02–1.88 (m, 1H), 1.84–1.59 (m, 3H), 1.43 (s, 3H).	(Method A): <i>m/z</i> = 246.1 [M+H] ⁺ , 1.08 min




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
78	2-Methyl-1-[(3S)-3-phenyl-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.39–7.23 (m, 5H), 5.31 (dd, <i>J</i> =8.8, 6.3 Hz, 1H), 4.25 (dt, <i>J</i> =7.7, 2.9 Hz, 1H), 3.86 (ddd, <i>J</i> =9.4, 8.0, 6.7 Hz, 1H), 3.03–2.92 (m, 1H), 2.92–2.83 (m, 1H), 2.24–2.12 (m, 1H), 1.10 (d, <i>J</i> =7.0 Hz, 3H), 1.00 (d, <i>J</i> =6.8 Hz, 3H).	(Method A): <i>m/z</i> = 220.0 [M+H] ⁺ , 0.95 min
79	1-[(3S)-3-Phenyl-1,2-oxazolidine-2-carbonyl]cyclobutane-1-carbonitrile		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.43–7.34 (m, 2H), 7.33–7.25 (m, 3H), 5.36 (dd, <i>J</i> =8.5, 6.8 Hz, 1H), 4.34 (dt, <i>J</i> =7.7, 2.5 Hz, 1H), 3.99 (ddd, <i>J</i> =9.9, 7.9, 6.5 Hz, 1H), 2.97 (dddd, <i>J</i> =12.0, 9.0, 6.3, 2.8 Hz, 1H), 2.77–2.62 (m, 2H), 2.61–2.45 (m, 2H), 2.31–2.08 (m, 2H), 1.91 (ttd, <i>J</i> =11.4, 9.4, 5.6 Hz, 1H)	(Method A): <i>m/z</i> = 257.2 [M+H] ⁺ , 0.98 min
80	2-Methoxy-2-methyl-1-[(3S)-3-phenyl-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.39–7.22 (m, 5H), 5.40 (dd, <i>J</i> =8.5, 6.5 Hz, 1H), 4.26 (dt, <i>J</i> =7.7, 3.0 Hz, 1H), 3.86 (ddd, <i>J</i> =9.4, 7.9, 6.8 Hz, 1H), 3.13 (s, 3H), 2.87 (dddd, <i>J</i> =12.1, 9.1, 6.5, 3.0 Hz, 1H), 2.15 (dddd, <i>J</i> =12.1, 9.5, 7.5, 6.3 Hz, 1H), 1.37 (s, 3H), 1.36 (s, 3H).	(Method A): <i>m/z</i> = 250.1 [M+H] ⁺ , 0.87 min



Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
15	(2S)-2-Hydroxy-1-[(3S)-3-phenyl-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.40–7.23 (m, 5H), 5.34 (dd, <i>J</i> =8.9, 6.4 Hz, 1H), 4.84 (d, <i>J</i> =7.0 Hz, 1H), 4.54–4.44 (m, 1H), 4.25 (dt, <i>J</i> =7.8, 2.6 Hz, 1H), 3.86 (ddd, <i>J</i> =9.5, 8.0, 6.8 Hz, 1H), 2.93–2.85 (m, 1H), 2.18 (dddd, <i>J</i> =12.2, 9.6, 7.6, 6.3 Hz, 1H), 1.30 (d, <i>J</i> =6.8 Hz, 3H).	(Method A): <i>m/z</i> = 222.2 [M+H] ⁺ , 0.73 min.
81	1-(3-(2-fluoropyridin-3-yl)isoxazolidin-2-yl)-2,2-dimethylpropan-1-one			
82	(R)-1-(3-(3-fluorophenyl)isoxazolidin-2-yl)-2,2-dimethylpropan-1-one			
83	tert-butyl 3-phenylisoxazolidine-2-carboxylate			
84	(R)-2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)propan-1-one			



Comp. No.	Name	Structure	³ / ₄ NMR	M+H ⁺
85	(R)-1-methyl-5-(3-phenylisoxazolidine-2-carbonyl)-1H-pyrrole-2-carbonitrile			
86	2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)propan-1-one			
155	(3S)-2-(3,3-difluoro-1-methylcyclobutanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine		¹ H NMR (400 MHz, CDCl ₃) δ 7.33-7.23 (m, 2H), 7.11-6.98 (m, 2H), 5.38 (dd, J=8.7, 6.1 Hz, 1H), 4.28 (dt, J=3.6, 7.6 Hz, 1H), 3.95 (ddd, J=9.2, 8.0, 6.7 Hz, 1H), 3.25-3.07 (m, 1H), 3.02-2.79 (ro, 2H), 2.62-2.29 (m, 3H), 1.57 (s, 3H)	LC-MS (Method A): m/z = 287.2 [M+H] ⁺ , 1.18 min
156	2-(azetidin-1-yl)-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one		¹ H NMR (500 MHz, CDCl ₃) δ 7.32-7.23 (m, 2H), 7.05-6.96 (m, 2H), 5.59-5.44 (m, 1H), 4.24 (td, J=7.8, 2.7 Hz, 1H), 3.96-3.84 (m, 1H), 3.53-3.35 (m, 4H), 2.84-2.75 (in, 1H), 2.33-2.24 (m, 1H), 2.01 (quin, J=7.1 Hz, 2H), 1.36-1.28 (m, 6H)	LC-MS (Method A): m/z = 293.2 [M+H] ⁺ , 0.91 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
157	2,2-dimethyl-1-[3-(pyridazin-4-yl)-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 9.2–9.13 (m, 2H), 7.39 (dd, <i>J</i> =5.3, 2.2 Hz, 1H), 5.44 (dd, <i>J</i> =8.8, 6.8 Hz, 1H), 4.31 (td, <i>J</i> =8.0, 2.5 Hz, 1H), 4.0–3.91 (m, 1H), 2.97–2.89 (m, 1H), 2.36–2.26 (m, 1H), 1.32 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 236.2 [M+H] ⁺ , 0.69 min.
158	4-[(3S)-2-(1-cyclopropylecyclopropanecarbonyl)-1,2-oxazolidin-3-yl]benzonitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.66 (d, <i>J</i> =8.3 Hz, 2H), 7.47 (d, <i>J</i> =8.5 Hz, 2H), 5.51 (dd, <i>J</i> =8.8, 6.0 Hz, 1H), 4.29 (td, <i>J</i> =8.0, 3.3 Hz, 1H), 4.0–3.94 (m, 1H), 2.94–2.85 (m, 1H), 2.38–2.28 (m, 1H), 1.66–1.58 (m, 1H), 1.06–0.99 (m, 1H), 0.90–0.84 (m, 1H), 0.68–0.61 (m, 1H), 0.55–0.39 (m, 3H), 0.27–0.15 (m, 2H).	LC-MS (Method A): <i>m/z</i> = 283.2 [M+H] ⁺ , 1.00 min.
159	5-[(3S)-2-(1-cyclopropylecyclopropanecarbonyl)-1,2-oxazolidin-3-yl]-2-fluoropyridine		¹ H NMR (400 MHz, CDCl ₃) δ 8.23 (d, <i>J</i> =2.5 Hz, 1H), 7.78 (td, <i>J</i> =8.0, 2.5 Hz, 1H), 6.93 (dd, <i>J</i> =8.0, 2.5 Hz, 1H), 5.51 (dd, <i>J</i> =9.0, 6.0 Hz, 1H), 4.32 (td, <i>J</i> =8.0, 3.3 Hz, 1H), 4.0–3.94 (m, 1H), 2.94–2.85 (m, 1H), 2.41–2.31 (m, 1H), 1.66–1.58 (m, 1H), 1.06–0.99 (m, 1H), 0.90–0.84 (m, 1H), 0.68–0.61 (m, 1H), 0.55–0.39 (m, 3H), 0.27–0.15 (m, 2H).	LC-MS (Method A): <i>m/z</i> = 277.2 [M+H] ⁺ , 0.91 min.

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
160	2-fluoro-5-[(3S)-2-{3-oxabicyclo[3.1.0]hexan-6-carbonyl}-1,2-oxazolidin-3-yl]pyridine		¹ H NMR (400 MHz, CDCl ₃) δ 8.21 (d, <i>J</i> =2.0 Hz, 1H), 7.78 (td, <i>J</i> =8.1, 2.4 Hz, 1H), 6.92 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 5.46 (dd, <i>J</i> =8.9, 5.9 Hz, 1H), 4.35–4.29 (m, 1H), 3.99 (d, <i>J</i> =8.8 Hz, 1H), 3.98–3.89 (m, 2H), 3.81–3.77 (m, 2H), 2.94–2.83 (m, 1H), 2.43–2.25 (m, 2H), 2.21–2.17 (m, 2H).	LC-MS (Method A): <i>m/z</i> = 279.2 [M+H] ⁺ , 0.66 min.
161	1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-bis(² H ₃)methyl(² H ₃)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.20 (d, <i>J</i> =2.3 Hz, 1H), 7.74 (td, <i>J</i> =7.9, 2.5 Hz, 1H), 6.92 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 5.46 (dd, <i>J</i> =8.9, 6.9 Hz, 1H), 4.35–4.28 (m, 1H), 3.93 (ddd, <i>J</i> =9.9, 8.1, 6.4 Hz, 1H), 2.87 (dddd, <i>J</i> =12.1, 9.1, 6.4, 2.8 Hz, 1H), 2.38–2.26 (m, 1H).	LC-MS (Method A): <i>m/z</i> = 262.2 [M+H] ⁺ , 0.92 min.
162	3,3,3-trifluoro-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =2.5 Hz, 1H), 7.72 (td, <i>J</i> =8.0, 2.6 Hz, 1H), 6.93 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 5.45 (dd, <i>J</i> =8.8, 6.0 Hz, 1H), 4.34 (td, <i>J</i> =7.8, 3.6 Hz, 1H), 3.98 (td, <i>J</i> =8.7, 6.8 Hz, 1H), 3.85 (dt, <i>J</i> =15.2, 7.6 Hz, 1H), 2.98–2.89 (m, 1H), 2.40 (dddd, <i>J</i> =12.5, 9.0, 7.5, 6.1 Hz, 1H), 1.39 (d, <i>J</i> =7.0 Hz, 3H).	LC-MS (Method A): <i>m/z</i> = 293.1 [M+H] ⁺ , 0.89 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
163	3,3,3-trifluoro-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one	 DIASTEREOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 8.21 (d, <i>J</i> =2.5 Hz, 1H), 7.75 (td, <i>J</i> =7.9, 2.5 Hz, 1H), 6.94 (dd, <i>J</i> =8.5, 3.0 Hz, 1H), 5.49 (dd, <i>J</i> =8.8, 6.3 Hz, 1H), 4.34 (td, <i>J</i> =8.0, 2.9 Hz, 1H), 3.99 (ddd, <i>J</i> =9.7, 8.2, 6.8 Hz, 1H), 3.94–3.84 (m, 1H), 2.93 (dddd, <i>J</i> =12.3, 9.2, 6.5, 2.9 Hz, 1H), 2.42 (dddd, <i>J</i> =12.5, 9.7, 7.7, 6.5 Hz, 1H), 1.43 (d, <i>J</i> =7.3 Hz, 3H).	LC-MS (Method A): <i>m/z</i> = 293.1 [M+H] ⁺ , 0.91 min.
164	1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethyl-3-(trifluoromethoxy)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =2.5 Hz, 1H), 7.76–7.68 (m, 1H), 6.92 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 5.46 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.39–4.31 (m, 1H), 4.16 (s, 2H), 3.99 (ddd, <i>J</i> =9.7, 8.1, 6.5 Hz, 1H), 2.89 (dddd, <i>J</i> =12.2, 9.2, 6.3, 2.9 Hz, 1H), 2.39–2.27 (m, 1H), 1.34–1.36 (m, 6H).	LC-MS (Method A): <i>m/z</i> = 337.2 [M+H] ⁺ , 1.06 min.
165	4-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-3,3-dimethyl-4-oxobutanenitrile		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =2.5 Hz, 1H), 7.73 (td, <i>J</i> =8.0, 2.6 Hz, 1H), 6.93 (dd, <i>J</i> =8.5, 3.0 Hz, 1H), 5.45 (dd, <i>J</i> =8.7, 6.9 Hz, 1H), 4.38 (td, <i>J</i> =7.9, 2.5 Hz, 1H), 4.00 (ddd, <i>J</i> =9.7, 8.1, 6.5 Hz, 1H), 2.92 (dddd, <i>J</i> =12.3, 9.2, 6.3, 2.9 Hz, 1H), 2.73 (d, <i>J</i> =16.6 Hz, 1H), 2.66 (d, <i>J</i> =16.6 Hz, 1H), 2.36 (dddd, <i>J</i> =12.5, 9.7, 7.5, 6.7 Hz, 1H), 1.49 (s, 3H), 1.47 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 278.2 [M+H] ⁺ , 0.79 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
166	(2S)-3,3,3-trifluoro-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2-hydroxy-2-methylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.23 (d, <i>J</i> =1.8 Hz, 1H), 7.75 (ddd, <i>J</i> =8.3, 7.5, 2.8 Hz, 1H), 6.97 (dd, <i>J</i> =8.4, 3.4 Hz, 1H), 5.48 (dd, <i>J</i> =8.7, 6.9 Hz, 1H), 4.70 (s, 1H), 4.42 (td, <i>J</i> =8.2, 2.4 Hz, 1H), 4.04 (ddd, <i>J</i> =10.2, 8.4, 6.5 Hz, 1H), 2.98 (dddd, <i>J</i> =12.6, 8.9, 6.7, 2.4 Hz, 1H), 2.43 (dddd, <i>J</i> =12.7, 10.2, 7.8, 6.7 Hz, 1H), 1.76 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 309.2 [M+H] ⁺ , 0.80 min.
167	2,2-dicyclopropyl-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]ethan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.21 (d, <i>J</i> =2.5 Hz, 1H), 7.76 (td, <i>J</i> =8.1, 2.9 Hz, 1H), 6.93 (dd, <i>J</i> =8.5, 2.8 Hz, 1H), 5.52 (dd, <i>J</i> =8.7, 6.1 Hz, 1H), 4.24 (td, <i>J</i> =7.8, 2.8 Hz, 1H), 3.85 (ddd, <i>J</i> =9.5, 8.0, 6.8 Hz, 1H), 2.95–2.84 (m, 1H), 2.34 (dddd, <i>J</i> =12.3, 9.6, 7.6, 6.1 Hz, 1H), 1.68 (t, <i>J</i> =9.4 Hz, 1H), 1.26–1.07 (m, 2H), 0.68–0.55 (m, 2H), 0.54–0.38 (m, 2H), 0.36–0.24 (m, 3H), 0.17–0.04 (m, 1H).	LC-MS (Method A): <i>m/z</i> = 291.2 [M+H] ⁺ , 0.99 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
168	5-[(3S)-2-[1-(difluoromethyl)cyclobutanecarbonyl]-1,2-oxazolidin-3-yl]-2-fluoropyridine		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =2.3 Hz, 1H), 7.72 (td, <i>J</i> =8.0, 2.6 Hz, 1H), 6.93 (dd, <i>J</i> =8.5, 2.8 Hz, 1H), 6.24 (t, <i>J</i> =57.5 Hz, 1H), 5.47 (dd, <i>J</i> =8.5, 6.5 Hz, 1H), 4.29 (td, <i>J</i> =7.9, 3.3 Hz, 1H), 3.96 (ddd, <i>J</i> =9.4, 8.2, 6.5 Hz, 1H), 3.01–2.83 (m, 1H), 2.69–2.58 (m, 1H), 2.54–2.29 (m, 4H), 2.08–1.94 (m, 1H), 1.94–1.81 (m, 1H).	LC-MS (Method A): <i>m/z</i> = 301.2 [M+H] ⁺ , 0.95 min.
169	3,3-difluoro-1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.19 (d, <i>J</i> =2.5 Hz, 1H), 7.72 (td, <i>J</i> =7.9, 2.5 Hz, 1H), 6.93 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 6.32 (t, <i>J</i> =58.0 Hz, 1H), 5.46 (dd, <i>J</i> =8.5, 6.8 Hz, 1H), 4.36 (td, <i>J</i> =7.7, 2.6 Hz, 1H), 3.99 (ddd, <i>J</i> =9.7, 8.1, 6.5 Hz, 1H), 2.91 (dddd, <i>J</i> =12.3, 9.1, 6.3, 2.9 Hz, 1H), 2.36 (dddd, <i>J</i> =12.5, 9.7, 7.5, 6.8 Hz, 1H), 1.40–1.36 (m, 6H).	LC-MS (Method A): <i>m/z</i> = 289.2 [M+H] ⁺ , 0.94 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
170	5-[(3S)-2-[1-(difluoromethyl)cyclopropanecarbonyl]-1,2-oxazolidin-3-yl]-2-fluoropyridine		¹ H NMR (400 MHz, CDCl ₃) δ 8.20 (d, <i>J</i> =2.5 Hz, 1H), 7.75 (td, <i>J</i> =8.0, 2.6 Hz, 1H), 6.94 (dd, <i>J</i> =8.4, 2.9 Hz, 1H), 6.65 (dd, <i>J</i> =59.5, 58.5 Hz, 1H), 5.51 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.36 (td, <i>J</i> =7.9, 3.0 Hz, 1H), 3.98 (ddd, <i>J</i> =9.5, 8.3, 6.8 Hz, 1H), 2.99–2.83 (m, 1H), 2.38 (dddd, <i>J</i> =12.5, 9.5, 7.7, 6.5 Hz, 1H), 1.44–1.33 (m, 1H), 1.31–1.09 (m, 3H).	LC-MS (Method A): <i>m/z</i> = 287.1 [M+H] ⁺ , 0.88 min.
171	4-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-3,3-dimethyl-4-oxobutanenitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.23 (m, 2H), 7.10–6.99 (m, 2H), 5.40 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.34 (td, <i>J</i> =7.8, 3.1 Hz, 1H), 3.98 (ddd, <i>J</i> =9.6, 8.1, 6.5 Hz, 1H), 2.86 (dddd, <i>J</i> =12.2, 8.7, 6.5, 3.1 Hz, 1H), 2.79–2.63 (m, 2H), 2.34 (dddd, <i>J</i> =12.4, 9.6, 7.6, 6.5 Hz, 1H), 1.49 (s, 6H).	LC-MS (Method A): <i>m/z</i> = 277.2 [M+H] ⁺ , 0.97 min.
172	6-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidine-2-carbonyl]-2-oxaspiro[3.3]heptane-6-carbonitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.24 (m, 2H), 7.12–7.00 (m, 2H), 5.33 (dd, <i>J</i> =8.8, 6.2 Hz, 1H), 4.82 (s, 2H), 4.64–4.52 (m, 2H), 4.41–4.31 (m, 1H), 4.18–4.07 (m, 1H), 3.06–2.86 (m, 4H), 2.81 (dd, <i>J</i> =13.0, 2.5 Hz, 1H), 2.43 (dddd, <i>J</i> =12.5, 9.4, 7.6, 6.1 Hz, 1H).	LC-MS (Method A): <i>m/z</i> = 317.2 [M+H] ⁺ , 0.89 min.





Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
173	1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-bis(² H ₃)methyl(² H ₃)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.24 (m, 2H), 7.08–6.97 (m, 2H), 5.41 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.27 (tdd, <i>J</i> =7.8, 2.9, 0.7 Hz, 1H), 3.89 (ddd, <i>J</i> =9.8, 8.0, 6.5 Hz, 1H), 2.88–2.75 (m, 1H), 2.30 (dddd, <i>J</i> =12.2, 9.8, 7.7, 6.4 Hz, 1H).	LC-MS (Method A): <i>m/z</i> = 261.2 [M+H] ⁺ , 1.09 min
174	1-[(3S)-3-(5-chloropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.49 (d, <i>J</i> =2.3 Hz, 1H), 8.47 (d, <i>J</i> =2.0 Hz, 1H), 7.60 (t, <i>J</i> =2.1 Hz, 1H), 5.43 (dd, <i>J</i> =8.9, 6.7 Hz, 1H), 4.36–4.26 (m, 1H), 3.92 (ddd, <i>J</i> =10.0, 8.1, 6.4 Hz, 1H), 2.88 (dddd, <i>J</i> =12.4, 9.0, 6.4, 2.6 Hz, 1H), 2.31 (dddd, <i>J</i> =12.3, 10.0, 7.7, 6.7 Hz, 1H), 1.31 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 269.1 [M+H] ⁺ , 0.97 min
175	1-[(3S)-3-[2-(difluoromethyl)pyridin-4-yl]-1,2-oxazolidin-2-yl]-2,2-dimethylpropan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.65–8.58 (m, 1H), 7.56 (s, 1H), 7.34–7.33 (m, 1H), 6.64 (t, <i>J</i> =55.6 Hz, 1H), 5.45 (dd, <i>J</i> =9.0, 6.7 Hz, 1H), 4.35–4.25 (m, 1H), 3.93 (ddd, <i>J</i> =9.9, 8.1, 6.4 Hz, 1H), 2.96–2.84 (m, 1H), 2.35–2.21 (m, 1H), 1.32 (s, 9H).	LC-MS (Method A): <i>m/z</i> = 285.2 [M+H] ⁺ , 0.95 min
176	1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethyl-3-(trifluoromethoxy)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.31–7.22 (m, 2H), 7.08–6.97 (m, 2H), 5.41 (dd, <i>J</i> =8.8, 6.4 Hz, 1H), 4.30 (td, <i>J</i> =7.9, 3.3 Hz, 1H), 4.16 (s, 2H), 3.96 (ddd, <i>J</i> =9.5, 8.0, 6.6 Hz, 1H), 2.83 (dddd, <i>J</i> =12.2, 8.8, 6.5, 3.2 Hz, 1H), 2.31 (dddd, <i>J</i> =12.3, 9.5, 7.6, 6.4 Hz, 1H), 1.35 (s, 6H).	LC-MS (Method A): <i>m/z</i> = 336.2 [M+H] ⁺ , 1.20 min

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
177	1-[(3S)-3-(5-chloropyridin-3-yl)-1,2-oxazolidin-2-yl]-2,2-dimethyl-3-(trifluoromethoxy)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.50 (d, <i>J</i> =2.5 Hz, 1H), 8.46 (d, <i>J</i> =2.0 Hz, 1H), 7.60 (t, <i>J</i> =1.9 Hz, 1H), 5.45 (dd, <i>J</i> =8.9, 6.6 Hz, 1H), 4.35 (td, <i>J</i> =7.7, 2.8 Hz, 1H), 4.16 (s, 2H), 4.04–3.93 (m, 1H), 2.91 (dddd, <i>J</i> =12.1, 9.2, 6.4, 3.0 Hz, 1H), 2.34 (dddd, <i>J</i> =12.4, 9.7, 7.5, 6.5 Hz, 1H), 1.36 (s, 6H)	LC-MS (Method A): <i>m/z</i> = 353.2 [M+H] ⁺ , 1.09 min
178	1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethyl-3-(1H-pyrazol-1-yl)propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 7.47 (dd, <i>J</i> =1.9, 0.7 Hz, 1H), 7.39 (dd, <i>J</i> =2.3, 0.7 Hz, 1H), 7.35–7.26 (m, 2H), 7.08–6.97 (m, 2H), 6.19 (t, <i>J</i> =2.1 Hz, 1H), 5.41 (dd, <i>J</i> =8.8, 6.6 Hz, 1H), 4.58 (d, <i>J</i> =14.0 Hz, 1H), 4.38–4.25 (m, 2H), 4.15 (ddd, <i>J</i> =9.7, 7.8, 6.5 Hz, 1H), 2.88 (dddd, <i>J</i> =12.0, 8.8, 6.5, 3.1 Hz, 1H), 2.33 (dddd, <i>J</i> =12.2, 9.6, 7.6, 6.6 Hz, 1H), 1.33 (s, 3H), 1.26 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 318.2 [M+H] ⁺ , 0.94 min
179	4-[(3S)-2-(3,3,3-trifluoro-2-methylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile (Second eluting diastereoisomer)		¹ H NMR (400 MHz, CDCl ₃) δ 7.70–7.63 (m, 2H), 7.44–7.38 (m, 2H), 5.45 (dd, <i>J</i> =8.9, 6.1 Hz, 1H), 4.33 (td, <i>J</i> =7.8, 3.6 Hz, 1H), 3.99 (td, <i>J</i> =8.6, 6.6 Hz, 1H), 3.92–3.83 (m, 1H), 3.01–2.88 (m, 1H), 2.44–2.31 (m, 1H), 1.39 (d, <i>J</i> =7.1 Hz, 3H).	LC-MS (Method A): <i>m/z</i> = 299.1 [M+H] ⁺ , 0.99min




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
180	4-[(3S)-2-(3,3,3-trifluoro-2-methylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile (First eluting diastereoisomer)	 DIASTEREOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 7.70–7.62 (m, 2H), 7.46–7.38 (m, 2H), 5.48 (dd, <i>J</i> =8.9, 6.5 Hz, 1H), 4.37–4.27 (m, 1H), 4.04–3.82 (m, 2H), 3.00–2.87 (m, 1H), 2.38 (dddd, <i>J</i> =12.4, 9.8, 7.7, 6.5 Hz, 1H), 1.45 (d, <i>J</i> =7.2 Hz, 3H).	LC-MS (Method A): <i>m/z</i> = 299.1 [M+H] ⁺ , 1.00min
181	4-[(3S)-2-(3-cyano-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.70–7.61 (m, 2H), 7.45–7.37 (m, 2H), 5.44 (dd, <i>J</i> =8.9, 6.8 Hz, 1H), 4.40–4.30 (m, 1H), 3.99 (ddd, <i>J</i> =9.8, 8.1, 6.3 Hz, 1H), 2.91 (dtd, <i>J</i> =12.0, 6.3, 3.2 Hz, 1H), 2.78–2.63 (m, 2H), 2.32 (dddd, <i>J</i> =12.4, 9.8, 7.6, 6.8 Hz, 1H), 1.50 (s, 3H), 1.47 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 284.2 [M+H] ⁺ , 0.89min
182	3-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2,2-dimethyl-3-oxopropanenitrile		¹ H NMR (400 MHz, CDCl ₃) δ 7.32–7.25 (m, 2 H), 7.09–7.00 (m, 2 H), 5.37 (dd, <i>J</i> =8.8, 6.5 Hz, 1 H), 4.42–4.34 (m, 1 H), 4.16 (ddd, <i>J</i> =9.8, 8.3, 6.8 Hz, 1 H), 2.99–2.87 (m, 1 H), 2.43 (dddd, <i>J</i> =12.5, 9.9, 7.8, 6.5 Hz, 1 H), 1.71 (s, 3 H), 1.60 (s, 3 H).	LC-MS (Method A): <i>m/z</i> = 263.2 [M+H] ⁺ , 0.97 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
183	1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-(trifluoromethoxy)propan-1-one (First eluting diastereoisomer)	 DIASTEREOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 7.33–7.22 (m, 2H), 7.09–7.00 (m, 2H), 5.45–5.32 (m, 1H), 5.26–5.14 (m, 1H), 4.30 (td, J=7.6, 4.1 Hz, 1H), 3.99 (td, J=8.2, 6.9 Hz, 1H), 2.88 (dddd, J=12.5, 8.6, 6.8, 4.3 Hz, 1H), 2.49–2.30 (m, 1H), 1.54 (d, J=6.5 Hz, 3H).	LC-MS (Method A): m/z = 308.1 [M+H] ⁺ , 1.07 min.
184	1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2-(trifluoromethoxy)propan-1-one (Second eluting diastereoisomer)	 DIASTEREOISOMER 2	¹ H NMR (400MHz, CDCl ₃) δ 8.19 (d, J=2.5 Hz, 1H), 7.73 (dt, J=2.6, 8.0 Hz, 1H), 6.94 (dd, J=2.9, 8.4 Hz, 1H), 5.41 (dd, J=8.7, 6.1 Hz, 1H), 5.20 (q, J=6.8 Hz, 1H), 4.35 (dt, J=4.0, 7.7 Hz, 1H), 4.00 (dt, J=6.8, 8.5 Hz, 1H), 3.02–2.86 (m, 1H), 2.51–2.36 (m, 1H), 1.58–1.50 (m, 3H).	LC-MS (Method A): m/z = 309.1 [M+H] ⁺ , 0.91 min.
185	1-[(3S)-3-(6-fluoropyridin-3-yl)-1,2-oxazolidin-2-yl]-2-(trifluoromethoxy)propan-1-one (First eluting diastereoisomer)	 DIASTEREOISOMER 1	¹ H NMR (400MHz, CDCl ₃) δ 8.22–8.17 (m, 1H), 7.74 (dt, J=2.6, 8.0 Hz, 1H), 6.94 (dd, J=8.5, 2.8 Hz, 1H), 5.50 (dd, J=8.7, 6.4 Hz, 1H), 5.24 (q, J=6.8 Hz, 1H), 4.36 (dt, J=2.6, 7.8 Hz, 1H), 3.95 (ddd, J=9.8, 8.3, 6.8 Hz, 1H), 2.99–2.88 (m, 1H), 2.41 (dddd, J=12.5, 9.7, 7.7, 6.3 Hz, 1H), 1.65–1.57 (m, 3H)	LC-MS (Method A): m/z = 274.2 [M+H] ⁺ , 1.05 min.





Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
186	2-(dimethyl-1,2-oxazol-4-yl)-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]propan-1-one (Second eluting diastereoisomer)	 DIASTEREISOISOMER 2	¹ H NMR (400MHz, CDCl ₃) δ 7.16–7.08 (m, 2H), 7.03–6.94 (m, 2H), 5.35 (dd, J=8.5, 6.0 Hz, 1H), 4.21 (td, J=7.8, 3.8 Hz, 1H), 3.98–3.82 (m, 2H), 2.90–2.76 (m, 1H), 2.34–2.24 (m, 4H), 2.20 (s, 3H), 1.35 (d, J=7.3 Hz, 3H)	LC-MS (Method A): m/z = 319.2 [M+H] ⁺ , 0.97 min.
187	4-[(3S)-2-[1-(difluoromethyl)cyclopropanecarbonyl]-1,2-oxazolidin-3-yl]benzonitrile	 DIASTEREISOISOMER 2	¹ H NMR (400MHz, CDCl ₃) δ 7.69–7.62 (m, 2H), 7.45–7.39 (m, 2H), 6.83–6.49 (m, 1H), 5.50 (dd, J=8.9, 6.4 Hz, 1H), 4.33 (dt, J=2.8, 7.8 Hz, 1H), 3.97 (ddd, J=9.6, 8.1, 6.7 Hz, 1H), 2.91 (dddd, J=12.2, 9.2, 6.5, 3.0 Hz, 1H), 2.33 (dddd, J=12.3, 9.6, 7.7, 6.5 Hz, 1H), 1.43–1.32 (m, 1H), 1.29–1.12 (m, 3H)	LC-MS (Method A): m/z = 293.17 [M+H] ⁺ , 0.97 min
188	3,3-difluoro-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one (First eluting diastereoisomer)	 DIASTEREISOISOMER 1	¹ H NMR (400 MHz, CDCl ₃) δ 7.32–7.24 (m, 2H), 7.09–7.00 (m, 2H), 5.95 (ddd, J=56.2, 54.7, 6.8 Hz, 2H), 5.42 (dd, J=9.2, 5.9 Hz, 1H), 4.33–4.25 (m, 1H), 3.94 (ddd, J=9.4, 8.0, 6.9 Hz, 1H), 3.60–3.44 (m, 1H), 2.86 (dddd, J=12.2, 8.8, 6.8, 3.3 Hz, 1H), 2.37 (dddd, J=12.3, 9.5, 7.8, 6.0 Hz, 1H), 1.34 (d, J=7.3 Hz, 3H).	LC-MS (Method A): m/z = 296.0 [M+Na] ⁺ , 1.02 min.




Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
189	3,3-difluoro-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one (Second eluting diastereoisomer)	 DIASTEREOISOMER 2	¹ H NMR (400 MHz, CDCl ₃) δ 7.34–7.22 (m, 2H), 7.08–7.01 (m, 2H), 6.00 (td, <i>J</i> =56.7, 7.5 Hz, 1H), 5.40 (dd, <i>J</i> =8.7, 5.9 Hz, 1H), 4.38–4.22 (m, 1H), 4.04–3.91 (m, 1H), 3.58–3.35 (m, 1H), 2.97–2.78 (m, 1H), 2.48–2.30 (m, 1H), 1.27 (d, <i>J</i> =6.5 Hz, 3H).	LC-MS (Method A): <i>m/z</i> = 296.0 [M+Na] ⁺ , 1.00 min.
190	2,2-dimethyl-1-[(3S)-3-(1H-pyrazol-5-yl)-1,2-oxazolidin-2-yl]propan-1-one		¹ H NMR (400 MHz, CDCl ₃) δ 8.10 (brs, 1H), 7.67 (d, <i>J</i> =2.2 Hz, 1H), 6.32 (d, <i>J</i> =2.2 Hz, 1H), 5.58 (t, <i>J</i> =7.7 Hz, 1H), 4.35–4.31 (m, 1H), 3.99–3.85 (m, 1H), 2.85–2.79 (m, 1H), 2.72–2.65 (m, 1H), 1.27 (s, 9H).	LC-MS (Method F): <i>m/z</i> = 224.0 [M+H] ⁺ , 0.90 min.
191	(5S)-5-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidine-2-carbonyl]-5-methylpyrrolidin-2-one		¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) δ 7.80 (s, 1H), 7.31–7.26 (m, 2H), 7.18–7.12 (m, 2H), 5.29 (t, <i>J</i> =7.7 Hz, 1H), 4.28–4.23 (m, 1H), 3.95–3.87 (m, 1H), 2.89–2.85 (m, 1H), 2.49–2.10 (m, 5H), 1.46 (s, 3H).	LC-MS (Method H): <i>m/z</i> = 585.0 [2M+H] ⁺ , 1.03 min.
192	2-[(3S)-2-(2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]pyridine-4-carbonitrile		¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.80 (dd, <i>J</i> =4.5, 1.5 Hz, 1H), 7.79 (dd, <i>J</i> =4.5, 1.5 Hz, 1H), 7.66 (s, 1H), 5.42–5.38 (m, 1H), 4.35–4.30 (m, 1H), 3.95–3.89 (m, 1H), 2.85–2.80 (m, 1H), 2.41–2.36 (m, 1H), 1.22 (s, 9H).	LC-MS (Method H): <i>m/z</i> = 260.0 [M+H] ⁺ , 1.23 min.





Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
193	2-cyclopropyl-1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methylpropan-1-one		¹ H NMR (300 MHz, CD ₃ OD) δ 7.36–7.31 (m, 2H), 7.10–7.04 (m, 2H), 5.40–5.37 (m, 1H), 4.36–4.31 (m, 1H), 4.03–3.94 (m, 1H), 2.93–2.87 (m, 1H), 2.30–2.22 (m, 1H), 1.43–1.38 (m, 1H), 1.14 (s, 3H), 1.10 (s, 3H), 0.40–0.31 (m, 4H).	LC-MS (Method P): m/z = 278.2 [M+H] ⁺ , 1.66 min.
194	1-[(3S)-3-(4-fluorophenyl)-1,2-oxazolidin-2-yl]-2-methanesulfonyl-2-methylpropan-1-one		¹ H NMR (300 MHz, CD ₃ OD) δ 7.37–7.33 (m, 2H), 7.09 (d, J=8.7 Hz, 2H), 5.44–5.39 (m, 1H), 4.40–4.34 (m, 1H), 4.11–4.03 (m, 1H), 3.06 (s, 3H), 2.98–2.88 (m, 1H), 2.36–2.24 (m, 1H), 1.62 (s, 6H).	LC-MS (Method P): m/z = 316.2 [M+H] ⁺ , 1.22 min.
229	4-[(3S)-2-[1-(difluoromethyl)cyclobutanecarbonyl]-1,2-oxazolidin-3-yl]benzonitrile		¹ H NMR (400MHz, CDCl ₃) δ 7.66 (d, J=8.2 Hz, 2H), 7.40 (d, J= 8.6 Hz, 2H), 6.03–6.58 (m, 1H), 5.46 (dd, J=8.6, 6.7 Hz, 1H), 4.26 (dt, J=3.1, 7.8 Hz, 1H), 3.94 (ddd, J=9.4, 8.2, 6.7 Hz, 1H), 2.92 (dddd, J=12.3, 9.2, 6.3, 3.1 Hz, 1H), 2.73–2.58 (m, 1H), 2.52–2.26 (m, 4H), 2.11–1.94 (m, 1H), 1.93–1.80 (m, 1H).	LC-MS (Method B): m/z = 307.2 [M+H] ⁺ , 1.03 min.


Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
230	4-[(3S)-2-(4,4,4-trifluoro-2-methylbutanoyl)-1,2-oxazolidin-3-yl]benzonitrile Diastereoisomer 1	 DIASTEREISOIMER 1	¹ H NMR (400MHz, CDCl ₃) δ 7.69–7.62 (m, 2H), 7.45–7.38 (m, 2H), 5.44 (dd, J=8.8, 6.3 Hz, 1H), 4.29 (dt, J=3.8, 7.8 Hz, 1H), 3.92 (ddd, J=9.5, 8.0, 6.8 Hz, 1H), 3.47–3.35 (m, 1H), 2.95–2.86 (m, 1H), 2.77 (dq, J=15.1, 11.0, 8.8 Hz, 1H), 2.33 (dddd, J=12.4, 9.6, 7.7, 6.3 Hz, 1H), 2.10 (dq, J=15.2 10.9, 4.5 Hz, 1H), 1.33 (d, J=7.0 Hz, 3H)	LC-MS (Method B): m/z = 312.0 [M+H] ⁺ , 1.03 min.
231	4-[(3S)-2-(4,4,4-trifluoro-2-methylbutanoyl)-1,2-oxazolidin-3-yl]benzonitrile Diastereoisomer 2	 DIASTEREISOIMER 2	¹ H NMR (400MHz, CDCl ₃) δ 7.71–7.61 (m, 2H), 7.44–7.38 (m, 2H), 5.41 (dd, J=8.8, 6.3 Hz, 1H), 4.31 (dt, J=3.4, 7.7 Hz, 1H), 4.03–3.94 (m, 1H), 3.47–3.30 (m, 1H), 2.98–2.87 (m, 1H), 2.73 (dq, J=15.0, 11.1, 7.8 Hz, 1H), 2.35 (dddd, J=12.4, 9.2, 7.3, 6.3 Hz, 1H), 2.23–2.07 (m, 1H), 1.24 (d, J=7.0 Hz, 3H)	LC-MS (Method B): m/z = 312.0 [M+H] ⁺ , 1.01 min.
232	4-[(3S)-2-(4,4-difluoro-2-methylbutanoyl)-1,2-oxazolidin-3-yl]benzonitrile Diastereoisomer 1	 DIASTEREISOIMER 1	¹ H NMR (400MHz, CDCl ₃) δ 7.70–7.61 (m, 2H), 7.45–7.38 (m, 2H), 6.07–5.73 (m, 1H), 5.43 (dd, J=8.8, 6.3 Hz, 1H), 4.28 (dt, J=2.8, 7.8 Hz, 1H), 3.93 (ddd, J=9.5, 8.2, 6.7 Hz, 1H), 3.37–3.22 (m, 1H), 2.95–2.85 (m, 1H), 2.47–2.26 (m, 2H), 1.87 (tdt, J=16.9, 14.5, 4.8 Hz, 1H), 1.31 (d, J=7.3 Hz, 3H).	LC-MS (Method B): m/z = 294.0 [M+H] ⁺ , 0.98 min.

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
233	4-[(3S)-2-(4,4-difluoro-2-methylbutanoyl)-1,2-oxazolidin-3-yl]benzonitrile Diastereoisomer 2	 DIASTEREISOIMER 2	¹ H NMR (400MHz, CDCl ₃) δ 7.72–7.61 (m, 2H), 7.44–7.38 (m, 2H), 6.01–5.68 (m, 1H), 5.42 (dd, J=8.8, 6.3 Hz, 1H), 4.31 (dt, J=3.5, 7.8 Hz, 1H), 4.02–3.93 (m, 1H), 3.37–3.18 (m, 1H), 2.97–2.87 (m, 1H), 2.46–2.27 (m, 2H), 1.93 (ttd, J=15.1, 5.3, 17.3 Hz, 1H), 1.21 (d, J=7.0 Hz, 3H).	LC-MS (Method B): m/z = 294.0 [M+H] ⁺ , 0.95 min.
234	5-[(3S)-2-(4,4,4-trifluoro-2,2-dimethylbutanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, CDCl ₃) δ 8.82 (d, J=2.0 Hz, 1H), 8.79 (d, J=2.3 Hz, 1H), 7.88 (t, J=1.9 Hz, 1H), 5.49–5.39 (m, 1H), 4.37 (dt, J=2.6, 8.0 Hz, 1H), 4.04 (ddd, J=9.9, 8.1, 6.4 Hz, 1H), 2.95 (dddd, J=12.2, 9.1, 6.3, 2.8 Hz, 1H), 2.80 (qd, J=11.5, 15.4 Hz, 1H), 2.58 (qd, J=11.3, 15.5 Hz, 1H), 2.35 (tdd, J=7.3, 12.4, 9.8 Hz, 1H), 1.43 (s, 3H), 1.39 (s, 3H).	LC-MS (Method A): m/z = 328.3 [M+H] ⁺ , 0.95 min.
235	5-[(3S)-2-(2-cyclopropyl-2-methylpropanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, CDCl ₃) δ 8.80 (t, J=1.9 Hz, 2H), 7.91 (t, J=1.9 Hz, 1H), 5.51 (dd, J=8.8, 7.0 Hz, 1H), 4.34 (dt, J=2.8, 7.8 Hz, 1H), 3.99 (ddd, J=10.2, 8.2, 6.5 Hz, 1H), 3.01–2.89 (m, 1H), 2.38–2.25 (m, 1H), 1.41–1.32 (m, 1H), 1.17 (s, 3H), 1.12 (s, 3H), 0.49–0.31 (m, 4H).	LC-MS (Method A): m/z = 286.3 [M+H] ⁺ , 0.96 min.

Comp. No.	Name	Structure	^1H NMR	M+H ⁺
236	(3S)-2-(2,2-difluorocyclopropanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine Diastereoisomer 1	 DIASTEREOISOMER 1	^1H NMR (400MHz, CDCl_3) δ 7.36- 7.29 (m, 2H), 7.08-7.01 (m, 2H), 5.45 (dd, $J=8.7, 6.1$ Hz, 1H), 4.33 (dt, $J=3.3, 7.9$ Hz, 1H), 3.99-3.90 (m, 1H), 3.11-2.97 (m, 1H), 2.87 (dddd, $J=12.2, 8.9, 6.8, 3.1$ Hz, 1H), 2.38 (dddd, $J=12.4, 9.5, 7.8, 6.0$ Hz, 1H), 2.16 (dtd, $J=12.6, 8.0, 5.9$ Hz, 1H), 1.78-1.66 (m, 1H).	LC-MS (Method A): m/z = 272.0 [M+Hf, 0.96 min.
237	(3S)-2-(2,2-difluorocyclopropanecarbonyl)-3-(4-fluorophenyl)-1,2-oxazolidine Diastereoisomer 2	 DIASTEREOISOMER 2	^1H NMR (400MHz, CDCl_3) δ 7.34- 7.26 (m, 2H), 7.09-7.00 (m, 2H), 5.47- 5.36 (m, 1H), 4.31 (dt, $J=3.6, 7.7$ Hz, 1H), 3.98 (q, $J=8.1$ Hz, 1H), 3.03 (br. s., 1H), 2.93- 2.82 (m, 1H), 2.45-2.33 (m, 1H), 2.21-2.09 (m, 1H), 1.76- 1.64 (m, 1H).	LC-MS (Method A): m/z = 272.3 [M+Hf, 0.97 min.
238	4-[(3S)-2-(cubane-1-carbonyl)-1,2-oxazolidin-3-yl]benzonitrile		^1H NMR (300 MHz, CD_3OD) δ 7.69 (d, $J = 8.4$ Hz, 2H), 7.48 (d, $J = 8.4$ Hz, 2H), 5.44-5.32 (m, 1H), 4.31- 4.20 (m, 4H), 4.05- 3.83 (m, 5H), 3.04-2.91 (m, 1H), 2.32-2.17 (m, 1H).	LC-MS (Method E): m/z = 305.1 [M+H] ⁺ , 2.902 min.
239	3-[(3S)-2-(3,3-difluoro-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]benzonitrile		^1H NMR (400 MHz, CD_3OD) δ 7.64-7.58 (m, 3H), 7.58-7.52 (m, 1H), 6.38 (i. $J = 57.2$ Hz, 1H), 5.45-5.38 (m, 1H), 4.41-4.36 (m, 1H), 4.04- 3.98 (m, 1H), 2.98-2.94 (m, 1H), 2.34-2.26 (m, 1H), 1.34 (s, 3H), 1.33 (s, 3H).	LC-MS (Method E): m/z = 295.1 [M+Hf, 2.991 min.

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
240	5-[(3S)-2-[1-(trifluoromethyl)cyclopropanecarbonyl]-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, CDCl ₃) δ 8.83 (d, <i>J</i> =1.8 Hz, 1H), 8.79 (d, <i>J</i> =2.3 Hz, 1H), 7.88 (t, <i>J</i> =2.1 Hz, 1H), 5.49 (dd, <i>J</i> =8.8, 6.5 Hz, 1H), 4.38 (dt, <i>J</i> =2.9, 8.0 Hz, 1H), 4.03 (ddd, <i>J</i> =9.7, 8.4, 6.5 Hz, 1H), 2.99 (dddd, <i>J</i> =12.3, 9.2, 6.5, 2.9 Hz, 1H), 2.47–2.32 (m, 1H), 1.47–1.20 (m, 4H).	LC-MS (Method A): <i>m/z</i> = 312.2 [M+H] ⁺ , 0.85 min.
241	5-[(3S)-2-(3,3,3-trifluoro-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, CDCl ₃) δ 8.83 (d, <i>J</i> =2.0 Hz, 1H), 8.79 (d, <i>J</i> =2.3 Hz, 1H), 7.88 (t, <i>J</i> =2.1 Hz, 1H), 5.49 (dd, <i>J</i> =8.8, 7.3 Hz, 1H), 4.40 (dt, <i>J</i> =2.4, 8.0 Hz, 1H), 3.99 (ddd, <i>J</i> =10.2, 8.3, 6.3 Hz, 1H), 3.03–2.90 (m, 1H), 2.36 (tdd, <i>J</i> =7.4, 12.5, 10.2 Hz, 1H), 1.58 (s, 3H), 1.57 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 314.2 [M+H] ⁺ , 0.94 min.
242	5-[(3S)-2-(tricyclopropylacetyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, CDCl ₃) δ 8.80 (s, 2H), 7.91 (t, <i>J</i> =1.9 Hz, 1H), 5.55 (dd, <i>J</i> =8.8, 7.1 Hz, 1H), 4.31 (dt, <i>J</i> =1.8, 7.9 Hz, 1H), 3.97 (ddd, <i>J</i> =10.4, 8.1, 6.3 Hz, 1H), 2.98–2.88 (m, 1H), 2.28 (tdd, <i>J</i> =7.5, 12.0, 10.4 Hz, 1H), 0.86 (tt, <i>J</i> =8.5, 5.7 Hz, 3H), 0.69–0.62 (m, 6H), 0.46–0.36 (m, 6H).	LC-MS (Method B): <i>m/z</i> = 338.3 [M+H] ⁺ , 1.11 min.

Comp. No.	Name	Structure	¹ H NMR	M+H ⁺
243	5-[(3S)-2-(4,4-difluoro-2,2-dimethylbutanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile		¹ H NMR (400MHz, DMSO- <i>d</i> ₆) δ 8.94 (d, <i>J</i> =2.0 Hz, 1H), 8.77 (d, <i>J</i> =2.0 Hz, 1H), 8.16 (t, <i>J</i> =2.1 Hz, 1H), 6.13 (tt, <i>J</i> =56.5, 5.05 Hz, 1H), 5.43–5.34 (m, 1H), 4.36 (dt, <i>J</i> =2.6, 7.7 Hz, 1H), 3.95 (ddd, <i>J</i> =9.7, 8.2, 6.0 Hz, 1H), 2.92 (dddd, <i>J</i> =12.1, 9.0, 6.1, 2.9 Hz, 1H), 2.36–2.20 (m, 3H), 1.28 (s, 3H), 1.26 (s, 3H).	LC-MS (Method A): <i>m/z</i> = 310.2 [M+H] ⁺ , 0.90 min.
244	5-[(3S)-2-(3-cyclopropyl-3,3-difluoro-2,2-dimethylpropanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile			
245	5-[(3S)-2-(2-(difluoromethyl)-2-methylbutanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile			
246	5-[(3S)-2-(3,3,3-trifluoro-2-methylpropanoyl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile			

Comp. No.	Name	Structure	³ / ₄ NMR	M+H ⁺
247	5-[(3S)-2-(2,2-dicyclopropyl-5,5-difluoropentanovl)-1,2-oxazolidin-3-yl]pyridine-3-carbonitrile			

Further details for the separation of the compounds described herein are as follows:

No.	RT first eluting*	RT second eluting*	e.e.	Flow rate / column preparative	Flow rate / column analytical	Solvent	Comment
Int 28	4.6	14.0	90.6		1 mL/min / as described	as described	second eluting enantiomer is major enantiomer 4.7:95.3
2	6.4	13.9	97.6	18 mL/min / as described	1 mL/min / as described, column (25 x 0.46 cm)	as described	only first eluting enantiomer is described
23	6.0	7.6	100	18 mL/min / as described	1 mL/min / as described, column (25 x 0.46 cm)	as described	only first eluting enantiomer is described
61 and 62	5.0	6.9	100 and 100	16 mL/min / as described	1 mL/min / as described	as described	
61	5.0	7.1	100	18 mL/min / as described	1 mL/min / as described	as described	Only second eluting enantiomer is described
73 and 74	4.7	6.3	100 and 100	18 mL/min / as described	1 mL/min / as described	as described	

No.	RT first eluting*	RT second eluting*	e.e.	Flow rate / column preparative	Flow rate / column analytical	Solvent	Comment
30		7.7	100	18 mL/min / Whelk 0-1 (R,R) (25 x 2.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ (EtOH/MeOH 1/1) 30/70% v/v	
31	3.8		100	18 mL/min / Whelk O-1 (R,R) (25 x 2.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ (EtOH/MeOH 1/1) 30/70% v/v	
35		10.1	100	18 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 50/50% v/v	
36	5.8		100	18 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 50/50% v/v	
40	12.0		100	17 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 75/25% v/v	
41		17.0	100	17 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 75/25% v/v	
45		8.1	100	18 mL/min / Chiraipak IC (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak IC (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 50/50% v/v	

No.	RT first eluting*	RT second eluting*	e.e.	Flow rate / column preparative	Flow rate / column analytical	Solvent	Comment
46	5.0		100	18 mL/min / Chiralpak IC (25 x 2.0 cm) 5 mM	1 mL/min / Chiralpak IC (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 50/50% v/v	
47		6.1	100	18 mL/min / Chiralpak IC (25 x 2.0 cm) 5 mM	1 mL/min / Chiralpak IC (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 80/20% v/v	
48	4.7		100	18 mL/min / Chiralpak IC (25 x 2.0 cm) 5 mM	1 mL/min / Chiralpak IC (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 80/20% v/v	
50		9.8	100	45 mL/min / Chiralpak IC (25 x 2.0 cm) 5 mM	2.5 mL/min / Chiralpak IC (25 x 0.46 cm) 5 mM	2-propanol 4% SFC	
51	6.0		100	45 mL/min / Chiralpak IC (25 x 2.0 cm) 5 mM	2.5 mL/min / Chiralpak IC (25 x 0.46 cm) 5 mM	2-propanol 4% SFC	
52							Separated by column chromatography (Cyclohexane- EtOAc, 95:5 to 50:50)
53							Separated by column chromatography (Cyclohexane- EtOAc, 95:5 to 50:50)

No.	RT first eluting*	RT second eluting*	e.e.	Flow rate / column preparative	Flow rate / column analytical	Solvent	Comment
59		15.7	100	40 mL/min / Whelk O-1 (R,R) (25 x 3.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ EtOH 40/60% v/v	
60	6.7		100	40 mL/min / Whelk O-1 (R,R) (25 x 3.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ EtOH 40/60% v/v	
64		8.1	100	18 mL/min / Chiraipak OJ-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak QJ- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 70/30% v/v	
65	4.8		100	18 mL/min / Chiraipak OJ-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak OJ- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 70/30% v/v	
63		6.8	97	18 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 82/18% v/v	
66	5.8		100	18 mL/min / Chiraipak AD-H (25 x 2.0 cm) 5 mM	1 mL/min / Chiraipak AD- H (25 x 0.46 cm) 5 mM	n-hexane/ EtOH 82/18% v/v	

No,	RT first eluting*	RT second eluting*	e.e.	Flow rate / column preparative	Flow rate / column analytical	Solvent	Comment
67		8.8	100	40 mL/min / Whelk O-1 (R,R) (25 x 3.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ EtOH 65/35% v/v	
68	5.5		100	40 mL/min / Whelk O-1 (R,R) (25 x 3.0 cm) 10 mM	1 mL/min / Whelk O-1 (R,R) (25 x 0.46 cm) 10 mM	n-hexane/ EtOH 65/35% v/v	

*Enantiomer or Diastereomer

IN VITRO ASSAY 1

RECEPTOR INTERACTING PROTEIN KINASE 1 INHIBITION BY COMPOUNDS OF STRUCTURE (I)

Compound potency at receptor interacting protein kinase 1 and selectivity at receptor interacting protein kinase 2 and receptor interacting protein kinase 3 was determined in autophosphorylation mode, using an ADP-Glo luminescence assay, which measures the conversion of ATP to ADP.

GST-hRIPK1 (8-327) enzyme was generated by Proteros GmbH by Baculovirus expression system. GST-hRIPK2 (1-299) and GST-hRIPK3 (1-518) enzymes were purchased from SignaiChem.

Test compounds were diluted in DMSO and 0.1 μ L of solution was dispensed to each well of a 384-well white solid microplates. The assay buffer was 50mM HEPES pH 7.5, 50mM NaCl, 30mM MgCl₂. The buffer was supplemented with 0.02% CHAPS, 0.01% of Piuronic F127, 0.1mg/mL BSA and 1mM DTT. Only for receptor interacting protein kinase 1 5mM MnCl₂, was included in the assay buffer on the day of the experiment. The enzymatic reaction consisted of 1.5pg/mL GST-hRIPK1 (8-327) and 50 μ M ATP for receptor interacting protein kinase 1, 0.1 μ g/mL GST-hRIPK2 (1-299) and 150 μ M ATP, 4 μ g/mL GST-hRIPK3 (1-518) and 15 μ M ATP. 5 μ L of enzyme and 5 μ L of ATP were added to the plate at twice the final assay concentration and incubated at room temperature for 3 hours. Following this reaction, 10 μ L of ADP-Glo reagent (Promega) was added to each well and incubated for 40 minutes at room temperature. This stops the kinase reaction and depletes any remaining ATP. 20 μ L of ADP-Glo detection reagent was then added to each well and incubated at room temperature for at least 15 minutes. The detection reagent converts ADP to ATP and introduces luciferase and luciferin to detect ATP. The luminescence is then measured with the Envision (PerkinElmer) plate reader. Test compound inhibition was expressed as percent inhibition of internal assay controls. For concentration response curves, normalized data is fit and pIC₅₀ determined using XL-fit (IDBS) for Excel. The pIC₅₀ were averaged to determine a mean value, for a minimum of two independent experiments.

Fluorescent Polarization Binding (FP Binding) assay (Berger S.B. et al. (2015) *Cell Death Discovery*, 1: 15009; Maki J.L. et al. (2012) *Anal Biochem.*, 427(2): 164-174) was performed in polystyrene low volume 384-well black plate, at Room Temperature (RT) in a final volume of 10.1 μ L/well using 10 nM of GST-hRIPK1 (8-327) enzyme and 5 nM of fluorescent-labeled ligand (14-(2-([3-({2-([4-(cyanomethyl)phenyl)amino]-6-[(5-cyclopropyl-1H-pyrazol-3-yl)amino]-4-pyrimidinyl)amino]propylamine)-2-oxoethyl)-16,16,18,18-tetramethyl-6,7,7a,8a,9,10,16,18-octahydrobenzo[2",3"]indolizino[8",7":5',6']pyrano[3',2':3,4]pyrido[1,2-a]indol-5-yl)-2-sulfonate.

Test compounds were serially diluted in DMSO at 100 fold final concentrations in the assay (1% DMSO final). In each well of a 384-well Plate were dispensed 0.1 μ L of compound solution (or DMSO for controls) followed by 5 μ L of GST-hRTPK1 (8-327) at twice the final concentrations in assay buffer (50 mM HEPES pH 7.5, 10 mM NaCl, 50 mM MgCl₂, 0.02% CHAPS, 0.5 mM DTT and 0.01% Pluronic F127). For negative control the enzyme addition was replaced by assay buffer only.

After addition of 5 μ L of fluorescent-labeled ligand at twice the final concentrations in assay buffer, the plate was incubated at RT for 30 min. At the end, the binding was measured as FP value with the Envision (PerkinElmer) plate reader using filter for an excitation λ = 531 nm FP and an emission λ = 595 nm FP (S & P-pol).

Certain compounds were tested in the following. Compound potency at receptor interacting protein kinase 1 and selectivity at receptor interacting protein kinase 2 and receptor interacting protein kinase 3 was determined in autophosphorylation mode, using an ADP-Glo luminescence assay (Berger S.B. et al. (2015) *Cell Death Discovery*, 1: 15009), which measures the conversion of ATP to ADP.

GST-hRIPK1 (8-327) enzyme was generated by Proteros GmbH by Baculovirus expression system.

Test compounds were diluted in DMSO and 0.1 μ L of solution was dispensed to each well of a 384-well white solid microplate. The assay buffer was 50 mM HEPES pH 7.5, 50 mM NaCl, 30 mM MgCl₂. The buffer was supplemented with 0.02% CHAPS, 0.01% of Pluronic F127, 0.1 mg/mL BSA and 1 mM DTT. Only for receptor interacting protein kinase 1 5 mM MnCl₂, was included in the assay buffer on the day of the experiment. The enzymatic reaction comprised 1.5 μ g/mL GST-hRTPK1 (8-327) and 50 μ M ATP for receptor interacting protein kinase 1 and 15 μ M ATP. 5 μ L of enzyme and 5 μ L of ATP were added to the plate at twice the final assay concentration and incubated at room temperature for 3 hours. Following this reaction, 10 μ L of ADP-Glo reagent (Promega) was added to each well and incubated for 40 minutes at room temperature. This stops the kinase reaction and depletes any remaining ATP. 20 μ L of ADP-Glo detection reagent was then added to each well and incubated at room temperature for at least 15 minutes. The detection reagent converts ADP to ATP and introduces luciferase and luciferin to detect ATP. The luminescence is then measured with the Envision (PerkinElmer) plate reader. Test compound inhibition was expressed as percent inhibition of internal assay controls. For concentration response curves, normalized data is fit and IC₅₀ determined using XL-fit (IDBS) for Excel. The IC₅₀ were averaged to determine a mean value, for a minimum of two independent experiments.

Test compound inhibition was expressed as percent inhibition of internal assay controls. For concentration response curves, normalized data is fit and IC_{50} determined using XL-fit (IDBS) for Excel. IC_{50} were averaged to determine a mean value, for a minimum of two independent experiments.

Receptor interacting protein kinase 1 activity of exemplarv' compounds was determined according to the above general procedures. Results are summarized in Table 2.

Table 2

No	ADP IC_{50}	FP IC_{50}
1	+++	+++
2	+++	+++
3	+++	
4	+++	+++
5	++	
6	++	
7	++	
8	+++	+++
9	+	
10	+	
11	++	
12	+++	+++
13	++	
14	+	
15		+
16	+++	+++
17	+++	+++
18	+++	+++
19	+++	+++
20	+++	+++
21		++
22	++	
23	+++	+++
28		+++
29		+++
30		+++
31		+++
32		+++
33		+++
34		+++

No	ADP IC_{50}	FP IC_{50}
35		+++
36		+++
37		+++
38		+++
39		+++
40		+++
41		+++
42		+
43		+
44		++
45		++
46		+++
47		+++
48		+
49		+++
50		+++
51		++
52		+
53		++
54		+++
55		+++
56		+++
57		+++
58		+++
59		+
60		+++
61		+++
62		++
63		++
64	+++	+++

No	ADP IC_{50}	FP IC_{50}
65		+
66		+++
67		++
68		+++
69	+++	+++
70	+++	+++
71	+++	+++
72	+++	+++
73	+++	+++
74		+
75		++
76		++
77	+++	+++
78	+++	+++
79	+++	+++
80	+++	+++
81	+++	+++
82		++
83	++	
84	+	
85	+	
87	+++	+++
88	+++	+++
89		+++
90		+++
91		+++
92		+++
93		++
94		+++
95		+++

WO 2017/096301

No	ADP ICso	F P ICso
96		4-4-4-
97	+++	4-4-4-
98		444-
99		44-4-
100		4-4-4-
101	+ + - +	+4-4-
102		+++
103		+++
104		+++
105		+++
106	4-4-4-	+++
107		+++
108	44-4-	+++
109		+++
110		+++
111		++
112		+++
113		+++
114		+++
115		+++
116		+++
117		+++
118		N/A
119		N/A
120		+++
121		+++
122		+++
123		++
124		+++
125		+++
126		+++
127		+++
128		+++
129		+++
130		+++
131		+++

PCT/US2016/064818

No	ADP ICso	F P ICso
132		+++
133		+++
134		+++
135		+++
136		++
137		4-4-
138		+++
139		+++
140		+++
141		+++
142		+++
143		+++
144		+++
145		+++
146		+++
147		+++
148		+++
149		++
150		+++
151		+++
152		+++
153		+++
154		+++
155		+++
156		++
157		++
158		+++
159		+++
160		++
161		+++
162		+++
163		+++
164		+++
165		+++
166		++
167		+++

No	ADP ICso	F P ICso
168		+++
169		+++
170		+++
171		+++
172		+++
173		+++
174		+++
175		+++
176		+++
177		+++
178		+++
179		+++
180		+++
181		+++
182		+++
183		+++
184		+++
185		++
186		++
187		+++
188		+++
189		+++
190		+++
191		+++
192		+++
193		+++
194		+4-
195		+++
196		+++
198		+++
200		+++
202		+++
203		+++
204		+++
205		+++
206		++

No	ADP IC ₅₀	FP IC ₅₀
207		+++
208		4-4-4-
209		44-4-
210		4-4-
211		4-4-4-
212		4-4-4-
213		4-4-4-
214		4-4-4-
215		4-4-4-
216		4-4-

No	ADP IC ₅₀	FP IC ₅₀
217		+++
218		+++
219		+4-4-
220		4-4+
221		+++
222		4-4-4
223		4-4-4
225		++
226		+++
227		++++

No	ADP IC ₅₀	FP IC ₅₀
228		+++
229		+++
230		4-4-4-
231		4-4-4-
232		+++
233		4-4-4-
236		4-4-4-
237		+++
238		+++
239		4-4-4-

44-4- indicates IC₅₀ less than 1 μ M

4-4- indicates IC₅₀ from 1 μ M to 10 μ M

4 indicates IC₅₀ greater than 10 μ M

Receptor interacting protein kinase 1 activity of certain compounds was determined according to

the above general procedures. These results in nM are summarized in Table 3.

Table 3

No	FP IC ₅₀ (nM)
61	28
88	16
101	8
109	7

No	FP IC ₅₀ (nM)
171	11
193	5
195	10
196	9

No	FP IC ₅₀ (nM)
198	51
200	9
221	70
242	5


Although various embodiments of the disclosure are disclosed herein, many adaptations and modifications may be made within the scope of the disclosure in accordance with the common general knowledge of those skilled in this art. Such modifications include the substitution of known equivalents for any aspect of the disclosure in order to achieve the same result in substantially the same way.

Numeric ranges are inclusive of the numbers defining the range. The word "comprising" is used herein as an open-ended term, substantially equivalent to the phrase "including, but not limited to" and the word "comprises" has a corresponding meaning. As used herein, the singular forms "a", "an" and "the"

include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a thing" includes more than one such thing. Citation of references herein is not an admission that such references are prior art to the present disclosure. Any priority document(s) and all publications, including but not limited to patents and patent applications, cited in this specification are incorporated herein by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein and as though fully set forth herein. The disclosure includes all embodiments and variations substantially as hereinbefore described and with reference to the examples and drawings.

CLAIMS

1. A method for inhibiting receptor interacting protein kinase 1 in a cell, comprising contacting the cell with a receptor interacting protein kinase 1 inhibitor.
2. A method for screening a compound for receptor interacting protein kinase 1 inhibitory activity in a cell, comprising contacting the cell with a compound and measuring the cellular conversion of ATP to ADP.
3. The method of claim 2, which the measuring is a colorimetric assay, a fluorescence assay, or a luminometric assay.
4. A compound having the following structure (Xa):



or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof, wherein:

A is $-C(=O)-$, $-S(=O)-$, $-S(=O)_2-$ or $-Si(=O)(=NH)-$;

R^1 is $-NR^5R^6$, C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl, heterocyclyl or heteroaryl;

R^2 is aryl or heteroaryl;

R^3 is hydrogen, deuterium, C_{1-6} alkyl or C_{3-10} cycloalkyl;

$R^{3'}$ is hydrogen or deuterium;

R^4 is hydrogen, halo, C_{1-6} alkyl or C_{3-10} cycloalkyl;

R^5 and R^6 are each independently H, C_{1-6} alkyl, C_{3-10} cycloalkyl, aryl, heteroaryl or heterocyclyl,

wherein each C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl and aryl group of R^1 , R^2 , R^3 , $R^{3'}$, R^4 , R^5 and R^6 is optionally substituted with one, two or three R^{10} ;

R^{10} in each instance is independently deuterium, halo, hydroxy, cyano, nitrile, C_{1-6} alkyl, C_{2-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{11}$, $-S(=O)R^{11}$, $-S(=O)_2R^{11}$, $-NR^{11}R^{11}$, $-R^{11}C(=O)SR^{11}$ or $-NR^{11}C(=O)OR^{11}$;

wherein the C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl and heteroaryl of R^{10} are optionally substituted with one, two or three substituents independently selected from halo, hydroxy, cyano, amino, nitro, C_{1-6} alkyl, C_{2-6} alkoxy, C_{1-6} haloalkoxy, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-10} cycloalkyl, heterocyclyl, aryl or heteroaryl; and

R^{11} in each instance is independently hydrogen, C_{1-6} alkyl, C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl;

wherein the C_{3-10} cycloalkyl, heterocyclyl, heteroaryl or aryl of R^{11} is optionally substituted with one, two or three C_{1-6} alkyl;

provided the compound is not ethyl 3-phenylisoxazolidine-2-carboxylate, benzyl 3-(4-amino-2-oxopyrimidin-1(2H)-yl)isoxazolidine-2-carboxylate, 2-(methylsulfonyl)-3-phenyl-isoxazolidine, tert-

butyl 5-benzyl-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-(napb.thalen-2-ylmethyl)-3-phenylisoxazolidine-2-carboxylate, tert-butyl 5-([1,1'-biphenyl]-4-ylmethyl)-3-phenylisoxazolidine-2-carboxylate or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)butan-1-one or 2,2-dimethyl-1-(3-phenylisoxazolidin-2-yl)but-3-en-1-one.

5. The compound of claim 4, wherein R^1 is optionally substituted C_{1-6} alkyl, C_{3-6} alkoxy, C_{3-10} cycloalkyl or heterocyclyl.

6. The compound of claim 4, wherein R^1 is optionally substituted C_{1-6} alkyl.

7. The compound of claim 5 or 6, wherein R^1 is substituted with at least one substituent selected from deuterium, hydroxy, halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy, cyano, C_{3-10} cycloalkyl, heterocyclyl and heteroaryl;

wherein the heterocyclyl or heteroaryl is optionally substituted with one, two or three C_{1-6} alkyl.

8. The compound of claim 6, wherein R^1 is substituted with at least one substituent selected from hydroxy, halo, C_{1-6} haloalkyl, C_{1-6} alkoxy, C_{1-6} haloalkoxy and cyano.

9. The compound of claim 4, wherein R^1 is optionally substituted C_{3-6} alkoxy.

10. The compound of claim 9, wherein R^1 is t-butoxy.

11. The compound of claim 4, wherein R^1 is $-NR^5R^6$ and at least one of R^5 or R^6 is optionally substituted C_{1-6} alkyl or optionally substituted aryl.

12. The compound of claim 11, wherein R^1 is dimethylamino, t-butylamino or phenylamino.

13. The compound of claim 4, wherein R^1 is optionally substituted C_{3-10} cycloalkyl.

14. The compound of claim 13, wherein the C_{3-10} cycloalkyl is substituted with at least one substituent selected from hydroxy, halo, cyano, C_{1-6} alkoxy, C_{1-6} haloalkoxy, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} hydroxy alkyl and C_{1-6} alkoxyalkyl.

15. The compound of claim 13, wherein C_{3-10} cycloalkyl is cyclopropyl, cyclobutyl, or bicycle [1.1.1]pentyl.

16. The compound of claim 4, wherein R^1 is optionally substituted heterocyclyl or optionally substituted heteroaryl.

17. The compound of claim 16, wherein R^1 is oxetanyl, pyrrolidinyl, tetrahydropyranyl or azetidiny.

18. The compound of claim 16, wherein R^1 is pyrrolyl, pyrazolopyndinyl or benzoisoxazolyl.

19. The compound as in one of claims 16-18, wherein R^1 is optionally substituted with a C_{1-6} alkyl, cyano or both.

20. The compound as in one of claims 4-19, wherein R^2 is aryl.

21. The compound of claim 20, wherein R^2 is phenyl.


22. The compound as in one of claims 4-19, wherein R^2 is heteroaryl.

23. The compound of claim 22, wherein R^2 is pyridinyl.

24. The compound as in one of claims 20-23, wherein R^2 is unsubstituted.

25. The compound as in one of claims 20-23, wherein R^2 is substituted.

26. **The** compound of claim 25, wherein R^2 is substituted **with** one or more halo.
27. A compound selected from Table 1A, Table 1B, or Table 1C, or a pharmaceutically acceptable salt, prodrug, stereoisomer or a mixture of stereoisomers thereof.
28. A composition comprising a pharmaceutically acceptable carrier and a **compound** as in one of claims 4-27.
29. A method for treating a necrotic cell disease, the method comprising administering an effective amount of the composition of claim 28 to a subject **in** need thereof.
30. **The** method of claim 29, wherein the necrotic cell disease is trauma, ischemia, stroke, cardiac infarction, infection, **Gaucher's** disease, Krabbe disease, sepsis, Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis, Huntington's disease, HIV-associated dementia, retinal degenerative disease, glaucoma, age-related macular degeneration, rheumatoid arthritis, psoriasis, psoriatic arthritis or inflammatory bowel disease.
31. A method for treating an inflammatory disorder, the method comprising administering an effective amount of the composition of claim 29 to a subject in need thereof.
32. **The** method of claim 31, wherein **the** inflammatory' disorder is inflammatory' bowel disease.
33. **The** method of claim 32, wherein the **inflammatory** disorder is Crohn's disease or ulcerative colitis.
34. A compound as in one of claims 4-27 for use in therapy.
35. A compound as **in** one of claims 4-27 **for** use in **the** treatment of a necrotic cell disease or an inflammatory disease.
36. Use of a compound as **in** one of claims 4-27 for the manufacture of a medicament for treating a necrotic cell disease or an **inflammatory** disease.
37. A method of preparing a compound of structure (Xa) of claim 4, comprising coupling a compound of formula (A-2):



with a compound of formula (B-2):



under conditions to provide the compound of structure (Xa), wherein A, R^1 , R^2 , R^3 , R^3' , and R^4 are as defined in claim 4 and LG is a leaving group.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/064818

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D413/04 C07D261/02 A61K31/42 A61P29/00 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal , WPI Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PHILIP A. HARRIS ET AL: "Discovery of Small Molecule RIP1 Kinase Inhibitors for the Treatment of Pathologies Associated with Necroptosis", ACS MEDICINAL CHEMISTRY LETTERS, vol. 4, no. 12, 12 December 2013 (2013-12-12) , pages 1238-1243 , XP055123759 , ISSN: 1948-5875 , DOI: 10.1021/ml400382p	1-3
A	page 1239: paragraph connecting left and right hand column (luminescent kinase assay measuring formation of ADP) page 1242: fourth line in last paragraph . -----	4-37
X	WO 2014/125444 A1 (GLAXOSMITHKLINE IP DEV LTD [GB]) 21 August 2014 (2014-08-21)	1-3
A	page 203, line 12 - page 204, line 6 ----- -/- .	4-37
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 8 March 2017		Date of mailing of the international search report 20/03/2017
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Panday, Narendra

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/064818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2014/145022 AI (HARVARD COLLEGE [US] ; SHANGHAI INST ORGANIC CHEM [CN] ; UNIV TUFTS [US]) 18 September 2014 (2014-09-18)	1-3
A	page 57, line 6 - page 58, line 4 -----	4-37
X	wo 2014/152182 AI (HARVARD COLLEGE [US] ; SHANGHAI INST ORGANIC CHEM [CN] ; BRIGHAM & WOMEN) 25 September 2014 (2014-09-25)	1-3
A	page 28, line 26 - page 29, line 19 -----	4-37
X	wo 2013/013826 AI (UNIV FRIEDRICH ALEXANDER ER [DE] ; GUENTHER CLAUDIA [DE] ; BECKER CHRIST) 31 January 2013 (2013-01-31)	1-3
A	page 18, line 27 - page 19, line 6 -----	4-37
Y	DE 23 39 185 AI (LEPETIT SPA) 28 February 1974 (1974-02-28) 2-carbethoxy-3-phenyl isoxazolidine mentioned in last paragraph "d" ; page 11 -----	37
Y	DATABASE CA [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ROMEO, ROBERTO ET AL: "Truncated Reverse Isoxazolidinyl Nucleosides: A New Class of Allosteric HIV-1 Reverse Transcriptase Inhibitors" , XP002767829 , retrieved from STN Database accession no. 2012 :196899 abstract	37
A	& ROMEO, ROBERTO ET AL: "Truncated Reverse Isoxazolidinyl Nucleosides: A New Class of Allosteric HIV-1 Reverse Transcriptase Inhibitors" , CHEMMEDCHEM , 7(4) , 565-569 CODEN: CHEMGX; ISSN: 1860-7179 , 2012 , DOI : 10.1002/CMDC.201200022 10.1002/CMDC.201200022 -----	1-36
X	DATABASE CA [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; LEMEN, GEORGIA S. ET AL: "Influence of Hydroxyl amine Conformation on Stereocontrol in Pd-Catalyzed Isoxazolidine-Forming Reactions" , XP002767830, retrieved from STN Database accession no. 2009 :244095 abstract	4,5 ,7,9 , 10,20, 21,24
Y	----- -/--	37

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/064818

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	& LEMEN, GEORGIA S. ET AL: "Infl uence of Hydroxy' lamine Conformati on on Stereocontrol i n Pd- Catalyzed I soxazol i dine-Formi ng Reacti ons" , JOURNAL OF ORGANIC CHEMISTRY , 74(6) , 2533-2540 CODEN: JOCEAH; ISSN: 0022-3263 , 2009 , DOI : 10. 1021/J08027399 10. 1021/J08027399 ----- CLAUDIO GIOIA ET AL: "Organocatalyti c Asymmetri c Formal 3 + 2₁ Cycl oaddi ti on with i n Si tu-Generated N -Carbamoyl Ni trones " , JOURNAL OF THE AMERICAN CHEMICAL SOCI ETY, vol . 131, no. 28, 22 July 2009 (2009-07-22) , pages 9614-9615 , XP055003071 , ISSN: 0002-7863 , DOI : 10. 1021/ja902458m scheme 2: conversi on of 7 to 8 -----	37
Y	WANG WENTIAN ET AL: "The use of Mosher deri vati ves for the determi nati on of the absol ute confi gurati on of substi tuted i soxazol i di nes" , TETRAHEDRON ASYMMETRY, PERGAMON PRESS LTD, OXFORD, GB, vol . 24, no. 4, 26 February 2013 (2013-02-26) , pages 223-228, XP028986698, ISSN: 0957-4166, DOI : 10. 1016/J .TETASY.2013 .01 .016 scheme 1: conversi on of 4 to 5 -----	37

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/064818

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
wo 2014125444	AI	21-08-2014	
		AU 2014217453 AI	13-08-2015
		CA 2900695 AI	21-08-2014
		CL 2015002279 AI	08-07-2016
		CN 105121432 A	02-12-2015
		CR 20150420 A	19-10-2015
		DO P2015000196 A	15-05-2016
		EA 201591506 AI	29-01-2016
		EP 2956452 AI	23-12-2015
		HK 1213545 AI	08-07-2016
		JP 2016512488 A	28-04-2016
		KR 20150119174 A	23-10-2015
		NZ 628447 A	26-08-2016
		PE 17522015 AI	25-11-2015
		PH 12015501675 AI	19-10-2015
		SG 11201505796T A	28-08-2015
		TW 201443042 A	16-11-2014
		US 2015353533 AI	10-12-2015
		US 2017008878 AI	12-01-2017
		UY 35330 A	29-08-2014
		wo 2014125444 AI	21-08-2014
wo 2014145022	AI	18-09-2014	
		EP 2968276 AI	20-01-2016
		JP 2016514693 A	23-05-2016
		US 2014323489 AI	30-10-2014
		US 2016024098 AI	28-01-2016
		wo 2014145022 AI	18-09-2014
wo 2014152182	AI	25-09-2014	NONE
wo 2013013826	AI	31-01-2013	NONE
DE 2339185	AI	28-02-1974	
		AR 195048 AI	30-08-1973
		AT 326120 B	25-11-1975
		AU 5872373 A	06-02-1975
		BE 803722 AI	17-12-1973
		CA 995680 A	24-08-1976
		CH 575934 A5	31-05-1976
		DD 107045 A5	12-07-1974
		DE 2339185 AI	28-02-1974
		ES 417674 AI	16-02-1976
		FR 2196166 AI	15-03-1974
		GB 1381249 A	22-01-1975
		HU 166257 B	28-02-1975
		IL 42890 A	30-07-1976
		JP S4956975 A	03-06-1974
		LU 68243 AI	23-10-1973
		NL 7311234 A	19-02-1974
		RO 66744 A	30-05-1981
		SU 481157 A3	15-08-1975
		US 3853903 A	10-12-1974
		ZA 7304775 B	26-06-1974