




(86) 1996/03/05

(87) 1996/09/19

(45) 2001/07/10

(72) CHEN, Jian, US

(72) HAMMONDS, R. Glenn, US

(72) GODOWSKI, Paul J., US

(72) MARK, Melanie R., US

(72) MATHER, Jennie P., US

(72) LI, Ronghao, US

(73) GENENTECH, INC., US


(51) Int.Cl.⁶ C12N 15/12, C07K 14/475, C07K 14/47, A61K 38/18,
A61K 38/17, C12N 5/08, C12N 5/06

(30) 1995/03/10 (08/402253) US

(30) 1995/05/10 (08/438,861) US

(54) **ACTIVATION DE RECEPTEURS AU MOYEN DU GENE 6
SPECIFIQUE DE L'ARRET DE CROISSANCE (GAS6)**

(54) **RECEPTOR ACTIVATION BY GAS6**



(57) An activator of the Rse and Mer receptor protein tyrosine kinases has been identified. The activator is encoded by growth arrest-specific gene 6 (gas6). Accordingly, the invention provides a method of activating the Rse or Mer receptor using gas6 polypeptide. Furthermore, the invention provides a method of enhancing survival, proliferation or differentiation of cells having the Rse or Mer receptor incorporated in their cell membranes which involves exposing those cells to gas6 polypeptide. The types of cells which can be treated according to the method include glial cells such as Schwann cells and mononuclear cells. Kits and articles of manufacture which include gas6 polypeptide are further provided. The invention also relates to gas6 variants.

P0929P2

RECEPTOR ACTIVATION BY GAS6**ABSTRACT OF THE DISCLOSURE**

An activator of the Rse and Mer receptor protein tyrosine kinases has been identified. The activator is encoded by growth arrest-specific gene 6 (gas6). Accordingly, the invention
5 provides a method of activating the Rse or Mer receptor using gas6 polypeptide. Furthermore, the invention provides a method of enhancing survival, proliferation or differentiation of cells having the Rse or Mer receptor incorporated in their cell membranes which involves exposing those cells to gas6 polypeptide. The types of cells which can be treated according to the method include glial cells such as Schwann cells and mononuclear cells. Kits and articles of
10 manufacture which include gas6 polypeptide are further provided. The invention also relates to gas6 variants.

P0929P2

RECEPTOR ACTIVATION BY GAS6

BACKGROUND OF THE INVENTIONField of the Invention

The invention relates generally to methods of activating the Rse or Mer tyrosine kinase
 5 receptors. More particularly, the invention relates to methods of enhancing survival, proliferation and/or differentiation of cells comprising the Rse receptor (such as glial cells) or Mer receptor (*e.g.*, monocytes) using gas6. The invention also relates to gas6 variants, particularly those which are less γ -carboxylated than gas6 isolated from nature.

Description of Related Art

10 Specific signals that control the growth and differentiation of cells in developing and adult tissues often exert their effects by binding to and activating cell surface receptors containing an intrinsic tyrosine kinase activity. Mark *et al.* recently described the human and murine complementary DNA sequences of the receptor tyrosine kinase Rse that is preferentially expressed in the adult brain (Mark *et al.*, J. Biol. Chem. **269**:10720 [1994]).
 15 The extracellular domain of Rse receptor is composed of two immunoglobulin-like (Ig-L) repeats followed by two fibronectin type III repeats. Complementary DNA sequences encoding proteins identical to human (Ohashi *et al.*, Oncogene **9**:699 [1994]) and murine Rse (Lai *et al.*, Oncogene **9**:2567 [1994]) have been reported independently, and termed Sky and Tyro3, respectively. See also Fujimimoto and Yamamoto Oncogene **9**: 693 (1994) concerning
 20 the murine equivalent to Rse they call brt and Dai *et al.* Oncogene **9**: 975 (1994) with respect to the human molecule they call tif.

The expression of Rse in various tissues has been investigated. Lai *et al.*, *supra*, found that, in the adult brain, Rse mRNA is localized in neurons of the neocortex, cerebellum and hippocampus. Schulz *et al.* similarly found that Rse is expressed at high levels in the cerebral
 25 cortex, the lateral septum, the hippocampus, the olfactory bulb and in the cerebellum. The highest levels of Rse expression in the brain were found to be associated with neurons. (Schulz *et al.* Molec. Brain Res. **28**: 273-280 [1995]). In the central nervous system (CNS) of mice, the expression of Rse is detected at highest levels during late embryonic stages and post birth, coincident with the establishment and maintenance of synaptic circuitry in cortical
 30 and hippocampal neurons (Lai *et al.*, *supra* and Schneider *et al.*, Cell **54**:787-793 [1988]).

P0929P2

This process is believed to be regulated locally, by cells that are in direct contact or positioned close to one another. By Northern blot analysis, Mark *et al.*, *supra*, found that high levels of Rse mRNA were present in samples of RNA from the brain and kidney. Dai *et al.*, *supra* found that Rse was highly expressed in human ovary and testes. The expression of Rse in
 5 various human cell lines was also analyzed by Mark *et al.*, *supra*. Little, or no, Rse mRNA was detected by Northern blotting of mRNA samples from the monocyte cell line THP-1 or the lymphoblast-like RAJI cells. However, the Rse transcript was detected in a number of hematopoietic cell lines, including cells of the myeloid (*i.e.*, myelogenous leukemia line K562 and myelomonocytic U937 cells) and the megakaryocytic leukemia lines DAMI and CMK11-
 10 5, as well as the human breast carcinoma cell line MCF-7. In the cell lines examined, the highest level of expression was observed in Hep 3B cells, a human hepatocarcinoma cell line.

Rse is structurally related to Axl (also known as Ufo or Ark) and shares 43% overall amino acid sequence identity with this tyrosine kinase receptor. See O'Bryan *et al.*, Mol. Cell. Biol. 11:5016 (1991), Janssen *et al.*, Oncogene 6:2113 (1991), Rescigno *et al.* Oncogene
 15 5:1908 (1991) and Bellosta *et al.* 15: 614 (1995) concerning Axl. Rse and Axl, together with Mer (Graham *et al.*, Cell Growth Differ. 5:647 [1994]), define a class of receptor tyrosine kinases whose extracellular domains resemble neural cell recognition and adhesion molecules (reviewed by Ruitishauser, U. in Current Opin. Neurobiology 3:709 [1993] and Brummendorf and Rathjen in J. Neurochemistry 61:1207 [1993]). Like Rse, Axl is also expressed in the
 20 nervous system, but is more widely expressed than Rse in peripheral tissues.

Mer mRNA is detected in peripheral blood mononuclear cells, bone marrow mononuclear cells and in monocytes, but not in granulocytes. Despite the fact that Mer mRNA is expressed in neoplastic B and T cell lines, it is not detected in normal B or T lymphocytes (Graham *et al.*, Cell Growth Differ. 5:647 [1994]). Mer is widely expressed in human tissues,
 25 but the highest levels of Mer mRNA are detected in the testis, ovary, prostate, lung and kidney (Graham *et al.*, Cell Growth Differ. 5:647 [1994]).

Disregulated expression of Mer, Rse and Axl is associated with cellular transformation. For example, Axl was isolated from DNA of patients with chronic myelogenous leukemia (O'Bryan *et al.*, *supra*) and chronic myeloproliferative disorder
 30 (Janssen *et al.*, *supra*) using a transfection/tumorigenicity assay. Mer was cloned from a neoplastic B cell line and is expressed in numerous transformed T acute lymphocytic leukemia cell lines (Graham *et al.*, *supra*). Rse and Axl, when overexpressed in fibroblasts, induce

P0929P2

cellular transformation (O'Bryan *et al.*, *supra*; Ohashi *et al.*, Oncogene 9:669 [1994]; Taylor *et al.*, J. Biol. Chem. 270:6872-6880 [1995]; and McCloskey *et al.*, Cell Growth and Diff. 5:1105-1117 [1994]). Rse mRNA and protein are also overexpressed in mammary tumors derived from transgenic animals that overexpress either the *wnt-1* or *fgf-3* oncogenes (Taylor *et al.*, J. Biol. Chem. 270: 6872-6880 [1994]).

Putative ligands for the Rse and Axl receptors have been reported. Varnum *et al.* Nature 373:623 (1995) and Stitt *et al.* Cell 80: 661-670 (1995) recently reported that gas6 (for growth arrest-specific gene 6) is a ligand for Axl. Gas6 belongs to a set of murine genes which are highly expressed during serum starvation in NIH 3T3 cells (Schneider *et al.*, Cell 54:787-793 [1988]). These genes were designated growth arrest-specific genes, since their expression is negatively regulated during growth induction. The human homolog of murine gas6 was also cloned and sequenced by Manfioletti *et al.* in Molec. Cell Biol. 13(8):4976-4985 (1993). They concluded that gas6 is a vitamin K-dependent protein and speculated that it may play a role in the regulation of a protease cascade relevant in growth regulation. Gas6 is expressed in a variety of tissues including the brain. See also Colombo *et al.* Genome 2:130-134 (1992) and Ferrero *et al.* J. Cellular Physiol. 158:263-269 (1994) concerning gas6.

Stitt *et al.*, *supra* further reported that protein S is the ligand for Tyro3. Protein S is a vitamin K-dependent plasma protein that functions as an anticoagulant by acting as a cofactor to stimulate the proteolytic inactivation of factors Va and VIIIa by activated protein C. Reviewed in Easmon *et al.* Atheroscler. Thromb. 12:135 (1992). Accordingly, protein S is an important negative regulator of the blood-clotting cascade. See Walker *et al.*, J. Biol. Chem. 255:5521-5524 (1980), Walker *et al.*, J. Biol. Chem. 256:11128-11131 (1981), Walker *et al.*, Arch. Biochem. Biophys. 252: 322-328 (1991), Griffin *et al.* Blood 79: 3203 (1992) and Easmon, D., Atheroscler. Thromb. 12:135 (1992). The discovery that about half of the protein S in human plasma is bound to C4BP further supports the notion that protein S is involved in the complement cascade. Dahlback *et al.*, PNAS(USA) 78: 2512-2516 (1981). The role of protein S as a mitogen for smooth muscle cells has also been reported. Gasic *et al.*, PNAS(USA) 89:2317-2320 (1992).

Protein S can be divided into four domains (see Figs. 1A, 1C and 1D herein). Residues 1-52 (Region A) are rich in γ -carboxyglutamic acid (Gla) residues which mediate the Ca^{2+} dependent binding of protein S to negatively charged phospholipids (Walker, J. Biol. Chem. 259:10335 [1984]). Region B includes a thrombin-sensitive loop. Region C contains

P0929P2

four epidermal growth factor (EGF)-like repeats. Region D is homologous to the steroid hormone binding globulin (SHBG) protein (Hammond *et al.*, FEBS Lett. **215**:100 [1987]). As discussed by Joseph and Baker (FASEB J. **6**:2477 [1994]), this region is homologous to domains in the A chain of laminin (23% identity) and merosin (22% identity) and to a domain
 5 in the *Drosophila* crumbs (19%).

Murine and human gas6 cDNAs encode proteins having 43 and 44% amino acid sequence identity respectively to human protein S.

SUMMARY OF THE INVENTION

The foregoing invention relates to gas6 variants which are essentially not γ
 10 carboxylated or are substantially less γ carboxylated than gas6 derived from an endogenous source of the molecule. Examples of such variants include gas6 variants lacking one or more glutamic acid residues from the A domain of gas6 which are normally γ carboxylated, fragments of gas6 which lack the A domain as well as fragments which consist essentially of the D domain of gas6 (or a G domain fragment of gas6).

15 The invention further provides a method of activating Rse or Mer receptor by exposing a cell (preferably a human cell) comprising the Rse or Mer receptor to exogenous gas6 in an amount effective to activate the Rse or Mer receptor. The Rse or Mer receptor is normally cell-bound and the gas6 is preferably human gas6. The invention also provides a method of enhancing survival, proliferation and/or differentiation of a cell which has the Rse or Mer
 20 receptor incorporated in the cell membrane thereof by exposing the cell to gas6 in an amount effective to enhance survival, proliferation and/or differentiation of the cell. The cell may be a neuron or a glial cell, such as a Schwann cell, or a monocyte (*e.g.* a macrophage). The cell may be present in cell culture or in a mammal (*e.g.* a human) which is suffering from a neurologic disease or disorder. Often, the gas6 is provided in a physiologically acceptable
 25 carrier.

The invention also provides kits and articles of manufacture comprising gas6 polypeptide. The article of manufacture usually comprises instructions for using the gas6 in an *in vitro* cell culture or for administering the gas6 to a mammal.

30

BRIEF DESCRIPTION OF THE DRAWINGS

P0929P2

Figs. 1A-1D provide a schematic representation of the structure of protein S and gas6 (Fig. 1A) and comparison of the amino acid homology between the bovine (b) and human (h) forms of protein S (Figs. 1C and 1D, respectively) with human gas6 (Fig. 1B). For h gas6, boxes represent the Gla region (*i.e.* the A domain), the loop region (*i.e.* the B domain), the 4 EGF-like repeats (labeled C₁-C₄) which form the C domain, and the region homologous to sex-hormone binding globulin (*i.e.* the D domain), which is also related to the G domains of laminin A chain and merosin and to *Drosophila* crumbs protein. The percentage of amino acid identity shared between h gas6 and either b protein S or h protein S is indicated within the corresponding boxes. The amino acids at the boundaries of each of the regions are indicated above the boxes.

Fig. 2 shows a comparison of the amino acid sequences of murine gas6 (m gas6) [SEQ ID NO:1], h gas6 [SEQ ID NO:2] and h protein S [SEQ ID NO: 3]. Residues of the "pre" and "pro" sequences are indicated (with the arrow indicating the last residue of each precursor sequence). The A-D domains are delineated, as are the two G domains which reside in the D domain (*i.e.* G domain 1 and G domain 2).

Figs. 3A-3D are graphs depicting characterization of the Rse-L in fetal bovine serum (FBS). Fig. 3A shows binding of ¹²⁵I-Rse-IgG as a function of FBS concentration. Binding, percent of total counts added that are membrane associated (100 x B/T, *i.e.* bound/total), is plotted as a function of FBS concentration. The data were fit to a 4 parameter model which gave an EC₅₀ of 0.58% v/v. Fig 3B illustrates binding of ¹²⁵I-Rse-IgG as a function of Ca²⁺ concentration, with constant FBS concentration. Binding was performed as in Fig. 3A either in the presence of 10% diafiltered FBS or in its absence and varying the concentration of added Ca²⁺. The EC₅₀ of Ca²⁺ as judged by a 4 parameter fit to the data is 0.18 mM. Fig 3C is a Scatchard analysis of ¹²⁵I-Rse-IgG binding to CMK11-5 membranes mediated by FBS. A single concentration of ¹²⁵I-Rse-IgG, FBS and Ca²⁺ was used with increasing concentrations of unlabeled Rse-IgG, and binding plotted vs the ratio of bound and free (B/F) after correction for nonspecific binding. Experiments at both 1% (K_d = 0.82 nM) and 10% (K_d = 2.2 nM) FBS are shown. Fig 3D is a KIRA analysis of dose dependent activation of Rse phosphorylation by the Q-sepharose^{*} enriched (QSE) fraction of FBS. The inset shows Rse-L activity was specifically neutralized by incubation with Rse-IgG. Rse phosphorylation is shown in serum starved cells (-); or cells treated with QSE fraction in the absence of added IgG proteins (QSE); or with QSE incubated with Rse-IgG or CD4-IgG as indicated.

P0929P2

Fig. 4 is a flow chart /cartoon illustrating the KIRA ELISA for the Rse receptor described in Example 4.

Fig. 5 displays inhibition of binding of ^{125}I -Rse-IgG to gas6 by unlabeled Rse-IgG. Increasing amounts of unlabeled Rse-IgG were added to tubes with constant ^{125}I -Rse-IgG and gas6. A nonlinear least squares fit to the data using a single class of sites gave an estimated equilibrium dissociation constant of 0.46 ± 0.04 nM. The inset shows a Scatchard plot of bound (B) vs bound/free (B/F) after correction for nonspecific binding.

Figs. 6A-6C show Rse-L activity in astrocyte cultures. To determine if astrocytes secrete Rse ligand, serum free media that was conditioned for 3 days was concentrated 10-fold in a Centricon-10^{*} centrifugal ultrafiltration device, and added directly to assay tubes to give the final concentrations indicated. In Fig. 6A binding of ^{125}I -Rse-IgG to CMK11-5 membranes was enabled by addition of astrocyte conditioned medium (ACM), with a half maximum effect achieved at 13% v/v ACM. Fig. 6B is a KIRA analysis of phosphorylation of Rse by ACM. Fig. 6C shows that the phosphorylation of Rse by ACM was inhibited by incubation with Rse-IgG, but not CD4-IgG. Neutralization was carried out as described in Fig. 3 legend.

As shown in Fig. 7, a deletion analysis of gas6 indicated that the G domains are sufficient for binding to Rse *in vitro*. Epitope tagged (gD) gas6 or protein S, or N-terminal truncation variants of gas6 (containing the indicated residues) were constructed and transiently expressed in 293 cells essentially following the procedure described in Example 6. Proteins of the correct molecular weight could be detected in unfractionated (input) cell supernatants using an antibody directed against the epitope tag. In contrast to protein S, the gas6-derivatives were precipitated from the cell supernatants by Rse-IgG. The binding was specific to the extracellular domain of Rse because proteins were not precipitated by control human Fc. For quantification purposes, the unfractionated (input) lanes represented 20% of the material used for precipitation.

Fig. 8 shows that gas6 induces the proliferation of P45 rat Schwann cells in a dose responsive fashion. Cells were plated in 24-well plates in F12/DME medium with 10 $\mu\text{g/ml}$ insulin and transferrin and 5 $\mu\text{g/ml}$ Vitamin E with the indicated concentrations of recombinant human gas6. Cells were counted with a Coulter counter after 48 h. The mean and standard deviation for six wells for each treatment are shown.

*-trademark

P0929P2

Fig. 9 illustrates that gas6 induced proliferation of p45 rat Schwann cells is neutralized by Rse-IgG. Cells were plated as described in Fig. 8 legend. Control cells received no further additions. All other cells were treated with two different purifications of gas6 (*i.e.* lot #15 and lot #9) and 10 μ g/ml of either Rse-IgG (labeled Rse) or CD4-IgG (labeled CD4Fc).

5 Fig. 10 shows a dose response curve for activation of Rse phosphorylation in the KIRA assay as described in Example 10.

Fig. 11 illustrates ion exchange chromatography of media conditioned by cells expressing human recombinant gas6. Media (700 ml) was dialyzed against buffer A (50 mM Tris-HCl pH 7.5, 5 mM benzamidine), adjusted to 0.1 % CHAPS, and loaded on a 6 ml
10 Resource-Q*column (Pharmacia) at 10 ml/minute. The column was washed with buffer A, and eluted with a 70 ml linear gradient of 0 to 0.4 M NaCl in buffer A at a flow rate of 1.0 ml/min collecting fractions of 2.0 ml. The fractions were analyzed for their ability to bind and activate Rse using the barium chloride binding method described in Example 6 and the KIRA assay described in Example 4. The binding activity is expressed as the percent of total
15 radioactivity added which is precipitated by barium chloride. The KIRA activity is expressed in units/ml relative to a standard.

Figs. 12A-12C depict the effect of gas6 and other growth factors on human Schwann cell growth and DNA synthesis. All data presented as mean $\pm$ standard error (n=4). Fig. 12A shows dose responsive curves of human Schwann cells to gas6 in different conditions. Cell
20 numbers were counted with coulter counter at 84 hours after culture with the indicated concentrations of gas6. Fig. 12B shows that gas6 increased thymidine incorporation in Schwann cells cultured as in Fig. 12A in the presence of different concentrations of gas6. ^{3} H-[methyl]-thymidine (0.5 μ Ci/ml) was added at 48 hours of culture. Cells were harvested at 96 hours of culture and processed for measurement of the radioactivity incorporated in DNA.
25 Fig. 12C shows the influence of growth factors on Schwann cell growth in the presence of 8F. Schwann cells were plated in 8F with or without PDGF (10 ng/ml, basic FGF (20 ng/ml), IL-1 α (1 ng/ml), TGF- β 1 (1 ng/ml) and gas6 (30 ng/ml). Cell numbers were counted after 108 hours.

Fig. 13 illustrates a time course of human Schwann cell growth in culture. Human
30 Schwann cells were plated at 2×10^4 cells/well in 24 well multiplates in F12/DME (1:1) supplemented with 8F with or without gas6 or 10% dia-filtered fetal bovine serum (FBS).

P0929P2

Four wells of cultures were taken from each group for cell counting every 24 hours. Data shown are mean $\pm$ standard error (n=4).

Fig. 14 depicts neutralization of gas6 induced phosphorylation of Rse by receptor-Fc fusion proteins measured in a KIRA assay. The percentage of Rse phosphorylation observed in CHO Rse.gD cells treated with purified gas6 in the presence of the indicated concentrations of receptor fusion protein relative to that observed in cells treated with just gas6 is shown.

Figs. 15A-15D show a kinetic analysis of ligand binding to Mer-Fc. Mer-Fc was coupled to the carboxymethylated dextran layer on the surface a BIAcore™ biosensor. Purified gas6 (Figs. 15A, B and C) or Protein S (Fig. 15D) at a concentration of either 100 nM (broken line) or 140 nM (solid line) was injected over the surface of the biosensor at 160 sec. At 340 seconds, the injector loop was switched to buffer to follow dissociation. The binding of gas6 to Mer-Fc on the chip was blocked by preincubation with soluble Mer-Fc (Fig. 15B) but not CD4-Fc (Fig. 15C). No binding of Protein S to Mer-Fc was observed (Fig. 15D).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

1. Definitions

As used herein, the terms "gas6" and "gas6 polypeptide" (unless indicated otherwise) refer to a polypeptide which is able to activate the Rse receptor or Mer receptor and encompass the mature, pre-, prepro- and pro- forms of gas6 polypeptide, either purified from a natural source, chemically synthesized or recombinantly produced. The present definition specifically includes "human" gas6 polypeptide comprising the amino acid sequence published in Manfioletti *et al.*, Mol. Cell. Biol. 13(8):4976-4985 (1993) (available from EMBL/GenBank/DDBJ under accession number X59846) and other mammalian gas6 polypeptides (such as murine gas6, see Manfioletti *et al.*, *supra*). Where the gas6 polypeptide has the amino acid sequence of a gas6 polypeptide found in nature, it is referred to herein as a "native" or "native sequence" polypeptide regardless of the method by which it is produced (e.g. it can be isolated from an endogenous source of the molecule or produced by synthetic techniques).

Gas6 has various amino acid "regions" or "domains" which are delineated in Figs. 1A-B and Fig. 2. The "A domain" or "Gla region" at the amino terminus of the polypeptide has residues which are rich in γ -carboxyglutamic acid (Gla residues) which appear to mediate calcium dependent binding of gas6 to negatively charged phospholipids in cell membranes.

P0929P2

The A-domain stretches from about residue 46-86 of murine gas6 and about residue 49-89 of human gas6. The following "B domain" comprises a thrombin sensitive "loop" and extends from about residue 87-114 of murine gas6 and about residue 90-117 of human gas6. The third domain called the "C domain" herein has four epidermal growth factor (EGF)-like repeats (C₁-
 5 C₄ in Fig. 1B). This C domain extends from about residue 115-275 of murine gas6 and about residue 118-278 of human gas6. The remaining "D domain" is homologous to steroid hormone binding globulin (SHBG) protein and comprises about residues 276-673 of murine gas6 and residues 279-678 of human gas6. The D domain comprises a pair of "G domains" called "G Domain 1" (*i.e.* about residues 311-468 for murine gas6 and about residues 314-471
 10 for human gas6) and "G Domain 2" (*i.e.* about residues 500-666 for murine gas6 and about residues 503-671 for human gas6).

The terms "gas6" and "gas6 polypeptide" also encompass "variants" or "mutants" of native gas6. Such variants include fragments of the human gas6 sequence; polypeptides wherein one or more amino acid residues are added at the N- or C-terminus of, or within, the
 15 human gas6 sequence; one or more amino acid residues are deleted, and optionally substituted by one or more amino acid residues; and derivatives of the above proteins, polypeptides, or fragments thereof, wherein an amino acid residue has been covalently modified so that the resulting product is a non-naturally occurring amino acid. Gas6 variants may be made synthetically, for example, by site-directed or PCR mutagenesis, or may exist naturally, as in
 20 the case of allelic forms and other naturally occurring variants of the translated amino acid sequence set forth in Manfioletti *et al.* that may occur in human and other animal species.

A gas6 variant is included within the scope of the invention provided that it is functionally active. As used herein, "functionally active" and "functional activity" in reference to gas6 means that the gas6 is able to activate the Rse receptor and/or Mer receptor
 25 and/or promote the proliferation, survival, and/or differentiation of cells comprising the Rse receptor or Mer receptor such as neurons, glial cells or monocytic cells. A "glial cell" is derived from the central and peripheral nervous system and can be selected from oligodendroglial, astrocyte, ependymal, or microglial cells as well as satellite cells of ganglia and the neurolemmal or Schwann cells around peripheral nerve fibers. A "monocytic cell"
 30 is a mononuclear leukocyte such as a macrophage.

Often gas6 variants will share at least about 75% (preferably greater than 80% and more preferably greater than 90%) sequence identity with the translated amino acid sequence

P0929P2

encoding mature gas6 or fragments thereof after aligning the sequences to provide for maximum homology, as determined, for example, by the Fitch *et al.*, PNAS (USA) **80**:1382-1386 (1983), version of the algorithm described by Needleman *et al.*, J. Mol. Biol. **48**:443-453 (1970). In order to screen for functionally active gas6 variants, a variant can be subjected to

5 one or more of the following functional activity tests/assays:

- (a) Receptor activation assays which measure downregulation or activation of receptor tyrosine kinase activity (*e.g.* western blotting using an anti-phosphotyrosine antibody to determine whether the variant is able to activate Rse receptor or Mer receptor, see Example 3 herein).
- 10 (b) KIRA ELISA to determine Rse or Mer receptor activation-capability of the variant as described in Example 4 below.
- (c) Schwann cell proliferation assay to establish whether or not the variant is able to enhance Schwann cell proliferation in cell culture. See Example 9 herein.

Amino acid sequence variants of gas6 can be prepared by introducing appropriate

15 nucleotide changes into gas6 DNA and thereafter expressing the resulting modified DNA in a host cell, or by *in vitro* synthesis. Such variants include, for example, deletions from, or insertions or substitutions of, amino acid residues within the gas6 amino acid sequence set forth in Manfioletti *et al.* Any combination of deletion, insertion, and substitution may be made to arrive at an amino acid sequence variant of gas6, provided that such variant possesses

20 the desired characteristics described herein. Changes that are made in the amino acid sequence to arrive at an amino acid sequence variant of gas6 may also result in further modifications of gas6 upon its expression in host cells, for example, by virtue of such changes introducing or moving sites of glycosylation.

There are two principal variables in the construction of amino acid sequence variants

25 of gas6: the location of the mutation site and the nature of the mutation. These are variants from the human gas6 amino acid sequence, and may represent naturally occurring allelic forms of gas6, or predetermined mutant forms of gas6 made by mutating gas6 DNA, either to arrive at an allele or a variant not found in nature. In general, the location and nature of the mutation chosen will depend upon the gas6 characteristic to be modified.

30 For example, due to the degeneracy of nucleotide coding sequences, mutations can be made in the human gas6 nucleotide sequence without affecting the amino acid sequence of the gas6 encoded thereby. Other mutations can be made that will result in a gas6 that has an

P0929P2

amino acid sequence different from that set forth in Manfiolleti *et al.*, but which is functionally active. Such functionally active amino acid sequence variants of gas6 are selected, for example, by substituting one or more amino acid residues in the human gas6 amino acid sequence with other amino acid residues of a similar or different polarity or charge.

One useful approach is called "alanine scanning mutagenesis". Here, an amino acid residue or group of target residues is/are identified (*e.g.*, charged residues such as arg, asp, his, lys, and glu) and, by means of recombinant DNA technology, replaced by a neutral or negatively charged amino acid (most preferably alanine or polyalanine) to affect the interaction of the amino acids with the surrounding aqueous environment in or outside the cell. Cunningham *et al.*, Science 244:1081-1085 (1989). Those domains demonstrating functional sensitivity to the substitutions are then refined by introducing further or other variants at or for the sites of substitution.

Thus, while the site for introducing an amino acid sequence variation is predetermined, the nature of the mutation *per se* need not be predetermined. For example, to optimize the performance of a mutation at a given site, ala scanning or random mutagenesis is conducted at the target codon or region and the expressed gas6 variants are screened for functional activity as discussed above.

Amino acid sequence deletions generally range from about 1 to 30 residues, more preferably about 1 to 10 residues, and typically are contiguous. Deletions from regions of substantial homology with other tyrosine kinase receptor ligands, for example, are more likely to affect the functional activity of gas6. Generally, the number of consecutive deletions will be selected so as to preserve the tertiary structure of gas6 in the affected domain, *e.g.*, β -pleated sheet or α helix. Preferred deletion mutants include those which lack one or more glutamic acid residues in the A domain of gas6 (*i.e.* those E residues in the A domain of gas6 shown in Fig. 2) or lack the A domain entirely. A preferred deletion mutant of gas6 is the D domain of gas6 or one of the G domains thereof.

Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one amino acid residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions (*i.e.*, insertions made within the human gas6 amino acid sequence) may range generally from about 1 to 10 residues, more preferably 1 to 5, most preferably 1

P0929P2

to 3. Examples of terminal insertions include gas6 with an N-terminal methionyl residue (such as may result from the direct expression of gas6 in recombinant cell culture), and gas6 with a heterologous N-terminal signal sequence to improve the secretion of gas6 from recombinant host cells. Other insertions include the fusion to the N- or C-terminus of gas6 of immunogenic polypeptides (for example, bacterial polypeptides such as β -lactamase or an enzyme encoded by the *E. coli* trp locus, or yeast protein), and C-terminal fusions with proteins having a long half-life such as immunoglobulin constant regions, albumin, or ferritin, as described in PCT Pub. No. WO 89/02922 (published April 6, 1989).

The third group of variants are those in which at least one amino acid residue in the gas6 amino acid sequence, and preferably only one, has been removed and a different residue inserted in its place. The sites of greatest interest for making such substitutions are in the regions of the gas6 amino acid sequence that have the greatest homology with other tyrosine kinase receptor ligands. Those sites are likely to be important to the functional activity of the gas6. Accordingly, to retain functional activity, those sites, especially those falling within a sequence of at least three other identically conserved sites, are substituted in a relatively conservative manner. Such conservative substitutions are shown in Table 1 under the heading of "Preferred Substitution". If such substitutions do not result in a change in functional activity, then more substantial changes, denominated "Exemplary Substitutions" in Table 1, or as further described below in reference to amino acid classes, may be introduced and the resulting variant gas6 analyzed for functional activity.

TABLE 1

Original Residue	Exemplary Substitutions	Preferred Substitution
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	ala; pro	ala
His (H)	asn; gln; lys; arg	arg

P0929P2

	Original Residue	Exemplary Substitutions	Preferred Substitution
	Ile (I)	leu; val; met; ala; phe; norleucine	leu
	Leu (L)	norleucine; ile; val; met; ala; phe	ile
	Lys (K)	arg; gln; asn	arg
	Met (M)	leu; phe; ile	leu
5	Phe (F)	leu; val; ile; ala; tyr	leu
	Pro (P)	ala	ala
	Ser (S)	thr	thr
	Thr (T)	ser	ser
	Trp (W)	tyr; phe	tyr
10	Tyr (Y)	trp; phe; thr; ser	phe
	Val (V)	ile; leu; met; phe; ala; norleucine	leu

Insertional, deletional, and substitutional changes in the gas6 amino acid sequence may be made to improve the stability of gas6. For example, trypsin or other protease cleavage sites are identified by inspection of the encoded amino acid sequence for an arginyl or lysinyl residue. These are rendered inactive to protease by substituting the residue with another residue, preferably a basic residue such as glutamine or a hydrophobic residue such as serine; by deleting the residue; or by inserting a prolyl residue immediately after the residue. Also, any cysteine residues not involved in maintaining the proper conformation of gas6 for functional activity may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking.

DNA encoding amino acid sequence variants of gas6 is prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants of gas6) or preparation by site-directed (or oligonucleotide-mediated) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared DNA encoding a variant or a non-variant form of gas6.

Site-directed mutagenesis is a preferred method for preparing substitution, deletion, and insertion variants of gas6 DNA. This technique is well known in the art (see, *e.g.*, Zoller *et al.*, Meth. Enz. **100**:4668-500 [1983]; Zoller *et al.*, Meth. Enz. **154**:329-350 [1987]; Carter, Meth. Enz. **154**:382-403 [1987]; and Horwitz *et al.*, Meth. Enz. **185**:599-611 [1990]), and has been used, for example, to produce amino acid sequence variants of trypsin and T4 lysozyme,

P0929P2

which variants have certain desired functional properties. Perry *et al.*, Science 226:555-557 (1984); and Craik *et al.*, Science 228:291-297 (1985).

Briefly, in carrying out site-directed mutagenesis of gas6 DNA, the gas6 DNA is altered by first hybridizing an oligonucleotide encoding the desired mutation to a single strand
5 of such gas6 DNA. After hybridization, a DNA polymerase is used to synthesize an entire second strand, using the hybridized oligonucleotide as a primer, and using the single strand of gas6 DNA as a template. Thus, the oligonucleotide encoding the desired mutation is incorporated in the resulting double-stranded DNA.

PCR mutagenesis is also suitable for making amino acid sequence variants of gas6.
10 See Higuchi, in PCR Protocols, pp.177-183 (Academic Press, 1990); and Vallette *et al.*, Nuc. Acids Res. 17:723-733 (1989). Briefly, when small amounts of template DNA are used as starting material in a PCR, primers that differ slightly in sequence from the corresponding region in a template DNA can be used to generate relatively large quantities of a specific DNA fragment that differs from the template sequence only at the positions where the primers differ
15 from the template.

Another method for preparing variants, cassette mutagenesis, is based on the technique described by Wells *et al.*, Gene 34:315-323 (1985). The starting material is the plasmid (or other vector) comprising the gas6 DNA to be mutated. The codon(s) in the gas6 DNA to be mutated are identified. There must be a unique restriction endonuclease site on each side of
20 the identified mutation site(s). If no such restriction sites exist, they may be generated using the above-described oligonucleotide-mediated mutagenesis method to introduce them at appropriate locations in the gas6 DNA. The plasmid DNA is cut at these sites to linearize it. A double-stranded oligonucleotide encoding the sequence of the DNA between the restriction sites but containing the desired mutation(s) is synthesized using standard procedures, wherein
25 the two strands of the oligonucleotide are synthesized separately and then hybridized together using standard techniques. This double-stranded oligonucleotide is referred to as the cassette. This cassette is designed to have 5' and 3' ends that are compatible with the ends of the linearized plasmid, such that it can be directly ligated to the plasmid. This plasmid now contains the mutated gas6 DNA sequence.

30 Covalent modifications of gas6 molecules also are included within the scope of this invention. For example, covalent modifications are introduced into gas6 by reacting targeted amino acid residues of the gas6 with an organic derivatizing agent that is capable of reacting with selected amino acid side chains or the N- or C-terminal residues.

Cysteinyll residues most commonly are reacted with α -haloacetates (and corresponding
35 amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or

P0929P2

carboxyamidomethyl derivatives. Cysteinyl residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing α -amino-containing residues include imidoesters such as methyl picolinimate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK_a of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ^{125}I or ^{131}I to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ($\text{R}'\text{-N}=\text{C}=\text{N-R}'$), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking gas6 to a water-insoluble support matrix or surface for diagnostic and/or therapeutic use. Commonly used crosslinking agents include, *e.g.*, 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional

P0929P2

imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimide yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Glutaminyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains, acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group. Creighton, Proteins: Structure and Molecular Properties, pp.79-86 (W.H. Freeman & Co., 1983). Gas6 also is covalently linked to nonproteinaceous polymers, *e.g.* polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,179,337; 4,301,144; 4,496,689; 4,640,835; 4,670,417; or 4,791,192.

The preferred gas6 is one which is "non-immunogenic in a human" which means that upon contacting the polypeptide in a pharmaceutically acceptable carrier and in a therapeutically effective amount with the appropriate tissue of a human, no state of sensitivity or resistance to the polypeptide is demonstrable upon the second administration of the polypeptide after an appropriate latent period (*e.g.*, 8 to 14 days).

A preferred gas6 variant is one which is essentially not " γ carboxylated" or is less carboxylated than "native" gas6 derived from an endogenous source of the molecule (*e.g.* serum) or native gas6 made by a recombinant cell wherein the conditions for culturing such a cell facilitate γ carboxylation of the gas6 (*e.g.* Vitamin K is present in the culture media). Vitamin K is a cofactor for the carboxylase enzyme. The A domain of native gas6 has several glutamic acid residues which are normally γ carboxylated (see Manfioletti *et al.*, *supra*). Accordingly, a convenient way to make a non- γ -carboxylated variant gas6 is to generate gas6 variants which lack one or more of the E residues from the A domain of native gas6 (see Fig. 2) or other gas6 fragments which lack this domain. The extent of γ carboxylation can be measured by amino acid sequence analysis or the barium chloride assay described in Example 11.

P0929P2

"gas6 antagonist" or "antagonist" refers to a substance that opposes or interferes with a functional activity of gas6. Examples of gas6 antagonists include neutralizing antibodies, Rse-IgG, Rse extracellular domain (Rse ECD), Axl-IgG, Axl ECD, Mer-IgG and Mer ECD.

5 The term "antibody" is used in the broadest sense and specifically covers single anti-gas6 monoclonal antibodies (including agonist and antagonist antibodies) and anti-gas6 antibody compositions with polyepitopic specificity.

10 The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

15 The monoclonal antibodies herein include hybrid and recombinant antibodies produced by splicing a variable (including hypervariable) domain of an anti-gas6 antibody with a constant domain (*e.g.* "humanized" antibodies), or a light chain with a heavy chain, or a chain from one species with a chain from another species, or fusions with heterologous proteins, regardless of species of origin or immunoglobulin class or subclass designation, as well as antibody fragments (*e.g.*, Fab, F(ab')₂, and Fv), so long as they exhibit the desired biological activity. (See, *e.g.* US Pat No 4,816,567 and Mage & Lamoyi, in Monoclonal Antibody Production Techniques and Applications, pp.79-97 (Marcel Dekker, Inc., New York [1987])).

25 Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler & Milstein, Nature 256:495 (1975), or may be made by recombinant DNA methods (U.S. Patent No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage libraries generated using the techniques described in McCafferty *et al.*, Nature 348:552-554 (1990), for example.

35 "Humanized" forms of non-human (*e.g.* murine) antibodies are specific chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human

P0929P2

immunoglobulins (recipient antibody) in which residues from a complementarity determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework region (FR) residues of the human
 5 immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or
 10 substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

The term "neutralizing antibody" as used herein refers to an antibody that is capable
 15 of specifically binding to gas6, and which is capable of substantially inhibiting or eliminating the functional activity of gas6 *in vivo* and/or *in vitro*. Typically a neutralizing antibody will inhibit the functional activity of gas6 at least about 50%, and preferably greater than 80%, as determined, for example, by KIRA ELISA (see Example 4 below).

Polyclonal antibodies directed toward gas6 generally are raised in animals by multiple
 20 subcutaneous or intraperitoneal injections of gas6 and an adjuvant. It may be useful to conjugate gas6 or a peptide fragment thereof to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, or soybean trypsin inhibitor, using a bifunctional or derivatizing agent, for example, maleimidobenzoyl sulfosuccinimide ester (conjugation through cysteine residues),
 25 N-hydroxysuccinimide (conjugation through lysine residues), glutaraldehyde, succinic anhydride, SOCl_2 , or $\text{R}^1\text{N} = \text{C} = \text{NR}$, where R and R^1 are different alkyl groups.

Animals are immunized with such gas6-carrier protein conjugates combining 1 mg or 1 μg of conjugate (for rabbits or mice, respectively) with 3 volumes of Freund's complete adjuvant and injecting the solution intradermally at multiple sites. One month later the
 30 animals are boosted with 1/5th to 1/10th the original amount of conjugate in Freund's complete adjuvant by subcutaneous injection at multiple sites. Seven to 14 days later animals are bled and the serum is assayed for anti-gas6 antibody titer. Animals are boosted until the antibody titer plateaus. Preferably, the animal is boosted by injection with a conjugate of the same gas6 with a different carrier protein and/or through a different cross-linking agent.
 35 Conjugates of gas6 and a suitable carrier protein also can be made in recombinant cell culture

P0929P2

as fusion proteins. Also, aggregating agents such as alum are used to enhance the immune response.

Monoclonal antibodies directed toward gas6 are produced using any method which provides for the production of antibody molecules by continuous cell lines in culture.

5 Examples of suitable methods for preparing monoclonal antibodies include the original hybridoma method of Kohler *et al.*, Nature 256:495-497 (1975), and the human B-cell hybridoma method, Kozbor, J., Immunol. 133:3001 (1984); Brodeur *et al.*, Monoclonal Antibody Production Techniques and Applications, pp.51-63 (Marcel Dekker, Inc., New York, 1987).

10 Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be performed following methods known in the art (Jones *et al.*, Nature 321:522-525 [1986];
15 Riechmann *et al.*, Nature 332:323-327 [1988]; and Verhoeven *et al.*, Science 239:1534-1536 [1988]), by substituting rodent complementarity-determining regions (CDRs) for the corresponding regions of a human antibody.

Alternatively, it is now possible to produce transgenic animals (*e.g.*, mice) that are capable, upon immunization, of producing a full repertoire of human antibodies in the absence
20 of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the production of human antibodies upon antigen challenge. See, for example,
25 Jakobovits *et al.*, PNAS 90:2551-2555 (1993); Jakobovits *et al.*, Nature 362:255-258 (1993); and Bruggermann *et al.*, Year in Immuno. 7:33 (1993). Human antibodies can also be produced in phage-display libraries. Hoogenboom *et al.*, J. Mol. Biol. 227:381 (1991); and Marks *et al.*, J. Mol. Biol. 222:581 (1991).

The term "immunoadhesin" is used interchangeably with the expressions "gas6-
30 immunoglobulin chimera" ("gas6-Ig"), "Rse-immunoglobulin chimera" ("Rse-Ig") and "Mer-immunoglobulin chimera" ("Mer-Ig") and refers to a chimeric molecule that combines a functionally active gas6 (*e.g.* the D domain thereof), Rse or Mer (*e.g.* the ECDs thereof) with an immunoglobulin sequence. The immunoglobulin sequence preferably, but not necessarily, is an immunoglobulin constant domain. The immunoglobulin moiety in the chimeras of the

P0929P2

present invention may be obtained from IgG₁, IgG₂, IgG₃ or IgG₄ subtypes, IgA, IgE, IgD or IgM, but preferably IgG₁ or IgG₃.

Chimeras constructed from a protein sequence (*e.g.* Rse or Mer receptor ECD) linked to an appropriate immunoglobulin constant domain sequence (immunoadhesins) are known in the art. Immunoadhesins reported in the literature include fusions of the T cell receptor (Gascoigne *et al.*, PNAS (USA) **84**:2936-2940 [1987]); CD4 (Capon *et al.*, Nature **337**:525-531 [1989]; Traunecker *et al.*, Nature **339**:68-70 [1989]; Zettmeissl *et al.*, DNA Cell Biol. USA **9**:347-353 [1990]; and Byrn *et al.*, Nature **344**:667-670 [1990]); L-selectin (Watson *et al.*, J. Cell. Biol. **110**:2221-2229 [1990]; and Watson *et al.*, Nature **349**:164-167 [1991]); CD44 (Aruffo *et al.*, Cell **61**:1303-1313 [1990]); CD28 and B7 (Linsley *et al.*, J. Exp. Med. **173**:721-730 [1991]); CTLA-4 (Lisley *et al.*, J. Exp. Med. **174**:561-569 [1991]); CD22 (Stamenkovic *et al.*, Cell **66**:1133-1144 [1991]); and TNF receptor (Ashkenazi *et al.*, PNAS (USA) **88**:10535-10539 [1991]).

The simplest and most straightforward immunoadhesin design combines the functionally active region(s) of the "adhesin" protein with the hinge and Fc regions of an immunoglobulin heavy chain. Ordinarily, when preparing the gas6-, Mer-, or Rse-immunoglobulin chimeras of the present invention, nucleic acid encoding the extracellular domain of Rse or Mer receptor or encoding gas6 (or a fragment thereof) will be fused C-terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, C_H2 and C_H3 domains of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately N-terminal to the C_H1 of the heavy chain or the corresponding region of the light chain.

The precise site at which the fusion is made is not critical; particular sites are well known and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the Rse-, Mer- or gas6-immunoglobulin chimeras.

In some embodiments, the Rse-, Mer- or gas6-immunoglobulin chimeras are assembled as monomers, or hetero- or homo-multimers, and particularly as dimers or tetramers, essentially as illustrated in WO 91/08298.

In a preferred embodiment, the gas6 sequence, Rse or Mer receptor extracellular domain sequence is fused to the N-terminus of the Fc domain of immunoglobulin G₁ (IgG₁). It is possible to fuse the entire heavy chain constant region to the gas6, Mer or Rse receptor sequence. However, more preferably, a sequence beginning in the hinge region just upstream

P0929P2

of the papain cleavage site which defines IgG Fc chemically (*i.e.* residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the Rse or Mer receptor or gas6 amino acid sequence is fused to (a) the hinge region and C_H2 and C_H3 or (b) the C_H1, hinge, C_H2 and C_H3 domains, of an IgG₁, IgG₂, or IgG₃ heavy chain. The precise site at which the fusion is made is not critical, and the optimal site can be determined by routine experimentation.

In some embodiments, the Rse-, Mer- or gas6-immunoglobulin chimeras are assembled as multimers, and particularly as homo-dimers or tetramers. Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in serum. In the case of multimer, each of the four units may be the same or different.

Alternatively, the Rse, Mer or gas6 sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the Rse, Mer or gas6 sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the C_H2 domain, or between the C_H2 and C_H3 domains. Similar constructs have been reported by Hoogenboom *et al.*, Mol. Immunol. 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to a Rse, Mer or gas6-immunoglobulin heavy chain fusion polypeptide, or directly fused to the Rse or Mer receptor or gas6. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the Rse-, Mer- or gas6-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567.

In a preferred embodiment, the immunoglobulin sequences used in the construction of the immunoadhesins of the present invention are from an IgG immunoglobulin heavy chain constant domain. For human immunoadhesins, the use of human IgG₁ and IgG₃

P0929P2

immunoglobulin sequences is preferred. A major advantage of using IgG₁ is that IgG₁ immunoadhesins can be purified efficiently on immobilized protein A. In contrast, purification of IgG₃ requires protein G, a significantly less versatile medium. However, other structural and functional properties of immunoglobulins should be considered when choosing the Ig fusion partner for a particular immunoadhesin construction. For example, the IgG₃ hinge is longer and more flexible, so it can accommodate larger "adhesin" domains that may not fold or function properly when fused to IgG₁. Another consideration may be valency; IgG immunoadhesins are bivalent homodimers, whereas Ig subtypes like IgA and IgM may give rise to dimeric or pentameric structures, respectively, of the basic Ig homodimer unit. For Rse-, Mer- or gas6-immunoadhesins designed for *in vivo* application, the pharmacokinetic properties and the effector functions specified by the Fc region are important as well. Although IgG₁, IgG₂ and IgG₄ all have *in vivo* half-lives of 21 days, their relative potencies at activating the complement system are different. IgG₄ does not activate complement, and IgG₂ is significantly weaker at complement activation than IgG₁. Moreover, unlike IgG₁, IgG₂ does not bind to Fc receptors on mononuclear cells or neutrophils. While IgG₃ is optimal for complement activation, its *in vivo* half-life is approximately one third of the other IgG isotypes. Another important consideration for immunoadhesins designed to be used as human therapeutics is the number of allotypic variants of the particular isotype. In general, IgG isotypes with fewer serologically-defined allotypes are preferred. For example, IgG₁ has only four serologically-defined allotypic sites, two of which (G1m and 2) are located in the Fc region; and one of these sites, G1m1, is non-immunogenic. In contrast, there are 12 serologically-defined allotypes in IgG₃, all of which are in the Fc region; only three of these sites (G3m5, 11 and 21) have one allotype which is nonimmunogenic. Thus, the potential immunogenicity of a γ 3 immunoadhesin is greater than that of a γ 1 immunoadhesin.

Gas6, Mer and Rse immunoadhesins are most conveniently constructed by fusing the cDNA sequence encoding the gas6, Mer or Rse portion in-frame to an Ig cDNA sequence. However, fusion to genomic Ig fragments can also be used (see, *e.g.* Gascoigne *et al.*, *supra*; Aruffo *et al.*, Cell 61:1303-1313 [1990]; and Stamenkovic *et al.*, Cell 66:1133-1144 [1991]). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the Ig parts of the immunoadhesin are inserted in tandem into a plasmid vector that directs efficient expression in the chosen host cells. For expression in mammalian cells, pRK5-based vectors (Schall *et al.*, Cell 61:361-370 [1990]) and CDM8-based vectors (Seed,

P0929P2

Nature 329:840 [1989]) are useful. The exact junction can be created by removing the extra sequences between the designed junction codons using oligonucleotide-directed deletional mutagenesis (Zoller and Smith, Nucleic Acids Res. 10:6487 [1982]; and Capon *et al.*, Nature 337:525-531 [1989]). Synthetic oligonucleotides can be used, in which each half is
 5 complementary to the sequence on either side of the desired junction; ideally, these are 36- to 48-mers. Alternatively, PCR techniques can be used to join the two parts of the molecule in-frame with an appropriate vector.

The choice of host cell line for the expression of the immunoadhesin depends mainly on the expression vector. Another consideration is the amount of protein that is required.
 10 Milligram quantities often can be produced by transient transfections. For example, the adenovirus EIA-transformed 293 human embryonic kidney cell line can be transfected transiently with pRK5-based vectors by a modification of the calcium phosphate method to allow efficient immunoadhesin expression. CDM8-based vectors can be used to transfect COS cells by the DEAE-dextran method (Aruffo *et al.*, Cell 61:1303-1313 [1990]; and
 15 Zettmeissl *et al.*, DNA Cell Biol. (US) 9:347-353 [1990]). If larger amounts of protein are desired, the immunoadhesin can be expressed after stable transfection of a host cell line. For example, a pRK5-based vector can be introduced into Chinese hamster ovary (CHO) cells in the presence of an additional plasmid encoding dihydrofolate reductase (DHFR) and conferring resistance to G418. Clones resistant to G418 can be selected in culture. These
 20 clones are grown in the presence of increasing levels of DHFR inhibitor methotrexate and clones are selected in which the number of gene copies encoding the DHFR and immunoadhesin sequences is co-amplified. If the immunoadhesin contains a hydrophobic leader sequence at its N-terminus, it is likely to be processed and secreted by the transfected cells. The expression of immunoadhesins with more complex structures may require uniquely
 25 suited host cells. For example, components such as light chain or J chain may be provided by certain myeloma or hybridoma host cells (Gascoigne *et al.*, *supra*; and Martin *et al.*, J. Virol. 67:3561-3568 [1993]).

Immunoadhesins can be conveniently purified by affinity chromatography. The suitability of protein A as an affinity ligand depends on the species and isotype of the
 30 immunoglobulin Fc domain that is used in the chimera. Protein A can be used to purify immunoadhesins that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark *et al.*, J. Immunol. Meth. 62:1-13 [1983]). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss *et al.*, EMBO J. 5:1567-1575 [1986]). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable
 35 matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow

P0929P2

rates and shorter processing times than can be achieved with agarose. The conditions for binding an immunoadhesin to the protein A or G affinity column are dictated entirely by the characteristics of the Fc domain; that is, its species and isotype. Generally, when the proper ligand is chosen, efficient binding occurs directly from unconditioned culture fluid. One distinguishing feature of immunoadhesins is that, for human $\gamma 1$ molecules, the binding capacity for protein A is somewhat diminished relative to an antibody of the same Fc type. Bound immunoadhesin can be efficiently eluted either at acidic pH (at or above 3.0), or in a neutral pH buffer containing a mildly chaotropic salt. This affinity chromatography step can result in an immunoadhesin preparation that is >95% pure.

10 The expression "Rse extracellular domain" or "Rse ECD" when used herein refers to a polypeptide sequence that shares a ligand-binding function of the extracellular domain of the Rse receptor. "Ligand-binding function" refers to the ability of the polypeptide to bind a Rse ligand, such as gas6. Accordingly, it is often not necessary to include the entire extracellular domain since smaller segments are commonly found to be adequate for ligand binding. The term ECD encompasses polypeptide sequences in which the cytoplasmic domain and hydrophobic transmembrane sequence (and, optionally, 1-20 amino acids amino-terminal to the transmembrane domain) of the Rse receptor have been deleted. Generally the ECD of the Rse receptor comprises amino acid residues from about 1-428 of the mature Rse receptor sequence disclosed in Mark *et al.*, *supra*.

20 The expression "Mer extracellular domain" or "Mer ECD" when used herein refers to a polypeptide sequence that shares a ligand-binding function of the extracellular domain of the Mer receptor. "Ligand-binding function" refers to the ability of the polypeptide to bind a Mer ligand, such as gas6. Accordingly, it may be unnecessary to include the entire extracellular domain since smaller segments are commonly found to be adequate for ligand binding. The term ECD encompasses polypeptide sequences in which the cytoplasmic domain and hydrophobic transmembrane sequence (and, optionally, 1-20 amino acids amino-terminal to the transmembrane domain) of the Mer receptor have been deleted. Generally the ECD of the Mer receptor comprises amino acid residues from about 1-499 of the mature human Mer receptor sequence disclosed in the GenBank database (Accession Number U08023).

30 The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising functionally active gas6 fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody thereagainst can be made, yet is short enough such that it does not interfere with functional activity of the gas6. The tag polypeptide preferably also is fairly unique so that the antibody thereagainst does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least

35

P0929P2

6 amino acid residues and usually between about 8-50 amino acid residues (preferably between about 9-30 residues). The epitope tag is generally provided at the amino- or carboxyl-terminus of the gas6. Such epitope tagged forms of the gas6 are desirable, as the presence thereof can be detected using a labelled antibody against the tag polypeptide. Also, provision
 5 of the epitope tag enables the gas6 to be readily purified by affinity purification using the anti-tag antibody.

Tag polypeptides and their respective antibodies are well known in the art. Examples include the flu HA tag polypeptide and its antibody 12CA5, (Field *et al.*, Mol. Cell. Biol. 8:2159-2165 [1988]); the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies
 10 thereto (Evan *et al.*, Molecular and Cellular Biology 5(12):3610-3616 [1985]); and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody (Paborsky *et al.*, Protein Engineering 3(6):547-553 [1990]). Other tag polypeptides have been disclosed. Examples include the Flag-peptide (Hopp *et al.*, BioTechnology 6:1204-1210 [1988]); the KT3 epitope peptide (Martin *et al.*, Science 255:192-194 [1992]); an α -tubulin epitope peptide (Skinner *et al.*, J.
 15 Biol. Chem 266:15163-15166 [1991]); and the T7 gene 10 protein peptide tag (Lutz-Freyermuth *et al.*, Proc. Natl. Acad. Sci. USA 87:6393-6397 [1990]). Once the tag polypeptide has been selected, an antibody thereto can be generated using the techniques disclosed herein.

Gas6-tag polypeptide fusions are most conveniently constructed by fusing the cDNA
 20 sequence encoding the gas6 portion in-frame to the tag polypeptide DNA sequence and expressing the resultant DNA fusion construct in appropriate host cells. Ordinarily, when preparing the gas6-tag polypeptide chimeras of the present invention, nucleic acid encoding the gas6 (or a fragment thereof) will be fused at its 3' end to nucleic acid encoding the N-terminus of the tag polypeptide, however 5' fusions are also possible.

25 Epitope tagged gas6 can be conveniently purified by affinity chromatography using the anti-tag antibody. The matrix to which the affinity antibody is attached is most often agarose, but other matrices are available [e.g. controlled pore glass or poly(styrenedivinyl)benzene]. The epitope tagged gas6 can be eluted from the affinity column by varying the buffer pH or ionic strength or adding chaotropic agents, for example.

30 An "exogenous" compound is defined herein to mean a compound that is foreign to a cell and/or mammal to be treated with the compound, or homologous to a compound found in the cell or mammal but produced outside the cell or mammal.

"Isolated", when used to describe the various proteins disclosed herein, means protein that has been identified and separated and/or recovered from a component of its natural
 35 environment. Contaminant components of its natural environment are materials that would

P0929P2

interfere with diagnostic or therapeutic uses for the protein, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the protein will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain.

"Essentially pure" protein means a composition comprising at least about 90% by weight of the protein, based on total weight of the composition, preferably at least about 95% by weight. "Essentially homogeneous" protein means a composition comprising at least about 99% by weight of protein, based on total weight of the composition.

Mammalian "Rse receptors" or "Rse receptor protein tyrosine kinases" (*i.e.* "rPTKs") have been described by Mark *et al.* in J. Biol. Chem. **269**: 10720 (1994). When used throughout this application, the expression "Rse receptor" refers to endogenous Rse receptor present in a cell of interest as well as Rse receptor which is present in a cell by virtue of the cell having been transformed with nucleic acid encoding the Rse receptor, for example. Accordingly, the Rse receptor may be an amino acid or covalent variant of one of the native Rse receptors described by Mark *et al.*, provided it is still "functionally active" (*i.e.* is able to be activated by a Rse ligand such as gas6). The preferred Rse receptor is endogenous human Rse receptor present in the cell membrane of a human cell.

The phrase "activating Rse receptor" refers to the step of causing the intracellular kinase domain of the Rse receptor to phosphorylate tyrosine residues in a substrate polypeptide. Often, the tyrosine residues are intrinsic to the Rse receptor (*i.e.* the "substrate" comprises the intracellular domain of the Rse receptor). Therefore, the degree of activation correlates with Rse receptor "autophosphorylation". Rse receptor autophosphorylation can be detected by Western blotting using an anti-phosphotyrosine antibody (see Example 3) or by KIRA ELISA (see Example 4). However, activation of the Rse receptor may correlate with phosphorylation of a substrate other than the Rse receptor (*e.g.* a tyrosine kinase existing adjacent the Rse receptor). This can be detected by measuring tyrosine phosphorylation of the substrate (*e.g.* by Western blotting).

Mammalian "Mer receptors" have been described in Graham *et al.*, Cell Growth Differ. **5**:647 (1994) (see GenBank database Accession Number U08023 for the correct human Mer sequence) and Graham *et al.*, Oncogene **10**(12):2349-2359 (1995). When used throughout this application, the expression "Mer receptor" refers to endogenous Mer receptor present in a cell of interest as well as Mer receptor which is present in a cell by virtue of the

P0929P2

cell having been transformed with nucleic acid encoding the Mer receptor, for example. The preferred Mer receptor is endogenous human Mer receptor present in a human cell.

The phrase "activating Mer receptor" refers to the step of causing the intracellular kinase domain of the Mer receptor to phosphorylate tyrosine residues in a substrate polypeptide. Often, the tyrosine residues are intrinsic to the Mer receptor (*i.e.* the "substrate" comprises the intracellular domain of the Mer receptor). Therefore, the degree of activation correlates with Mer receptor "autophosphorylation". Mer receptor autophosphorylation can be detected by Western blotting using an anti-phosphotyrosine antibody or by KIRA ELISA (see below). However, activation of the Mer receptor may correlate with phosphorylation of a substrate other than the Mer receptor (*e.g.* a tyrosine kinase existing adjacent the Mer receptor). This can be detected by measuring tyrosine phosphorylation of the substrate (*e.g.* by Western blotting).

The expression "enhancing survival of a cell" refers to the act of increasing the period of existence of a cell, relative to an untreated cell which has not been exposed to gas6, either *in vitro* or *in vivo*.

The phrase "enhancing proliferation of a cell" encompasses the step of increasing the extent of growth and/or reproduction of the cell, relative to an untreated cell, either *in vitro* or *in vivo*. An increase in cell proliferation in cell culture can be detected by counting the number of cells before and after exposure to gas6 (see Example 9 herein). The extent of proliferation can be quantified via microscopic examination of the degree of confluency. Cell proliferation can also be quantified by measuring ³H uptake by the cells.

By "enhancing differentiation of a cell" is meant the act of increasing the extent of the acquisition or possession of one or more characteristics or functions which differ from that of the original cell (*i.e.* cell specialization). This can be detected by screening for a change in the phenotype of the cell (*e.g.* identifying morphological changes in the cell, see Example 9 below).

"Physiologically acceptable" carriers, excipients, or stabilizers are ones which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar

P0929P2

alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween*, Pluronics* or polyethylene glycol (PEG).

The terms "treating", "treatment", and "therapy" refer to curative therapy, prophylactic therapy, and preventative therapy.

- 5 The term "mammal" refers to any mammal classified as a mammal, including humans, cows, horses, dogs and cats. In a preferred embodiment of the invention, the mammal is a human.

10 The expression "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, a ribosome binding site, and possibly, other as yet poorly understood sequences. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

15 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally,
20 "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

25 2. Gas6 Production

Techniques suitable for the production of native gas6 or gas6 variants are well known in the art and include isolating gas6 from an endogenous source of this polypeptide (e.g. from serum), peptide synthesis (using a peptide synthesizer) and recombinant techniques (or any combination of these techniques). The preferred technique for production of native gas6 or
30 a gas6 variant is a recombinant technique. The preferred gas6 variants are those which are essentially not γ carboxylated. This can be achieved in a number of ways but most conveniently involves creating a molecule which lacks one or more of the glutamic acid residues in the A domain of native gas6 which are normally γ carboxylated. Optionally, the entire A domain may be removed from the native molecule by enzymatic cleavage, but

P0929P2

normally a nucleic acid molecule will be isolated which encodes the desired fragment (*e.g.* the D domain or a G domain therefrom). This nucleic acid molecule can be derived from the native gas6 nucleic acid.

Nucleic acid encoding native gas6 can be isolated from a cDNA library prepared from
 5 tissue believed to possess the polypeptide mRNA and to express it at a detectable level (*e.g.* brain tissue, see Example 6 below). Libraries are screened with probes (such as antibodies or oligonucleotides of about 20-80 bases) designed to identify the gas6 gene or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures as described in chapters 10-12 of Sambrook *et al.*,
 10 Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989).

Techniques for generating gas6 mutants via modification of the wildtype nucleic acid have been discussed above. The nucleic acid (*e.g.*, cDNA or genomic DNA) encoding the native gas6 or gas6 variant is inserted into a replicable vector for further cloning
 15 (amplification of the DNA) or for expression. Many vectors are available. The vector components generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence.

The gas6 polypeptide may be produced as a fusion polypeptide with a signal sequence
 20 or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the DNA that is inserted into the vector. The heterologous signal sequence selected preferably is one that is recognized and processed (*i.e.*, cleaved by a signal peptidase) by the host cell. For prokaryotic host cells, the signal sequence may be substituted by a
 25 prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the native signal sequence may be substituted by, *e.g.*, the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Pat. No. 5,010,182 issued 23 April 1991), or acid phosphatase leader, the *C. albicans* glucoamylase
 30 leader (EP 362,179 published 4 April 1990), or the signal described in WO 90/13646 published 15 November 1990. In mammalian cell expression the native gas6 signal sequence is satisfactory, although other mammalian signal sequences may be suitable as well as viral secretory leaders, for example, the herpes simplex gD signal. The DNA for such precursor region is ligated in reading frame to DNA encoding the native gas6/gas6 variant.

P0929P2

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Generally, in cloning vectors this sequence is one that enables the vector to replicate independently of the host chromosomal DNA, and includes origins of replication or autonomously replicating sequences. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (the SV40 origin may typically be used only because it contains the early promoter).

Expression and cloning vectors should contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, *e.g.*, ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, *e.g.*, the gene encoding D-alanine racemase for *Bacilli*. One example of a selection scheme utilizes a drug to arrest growth of a host cell. Those cells that are successfully transformed with a heterologous gene produce a protein conferring drug resistance and thus survive the selection regimen. Examples of such dominant selection use the drugs neomycin (Southern *et al.*, J. Molec. Appl. Genet. 1:327 [1982]), mycophenolic acid (Mulligan *et al.*, Science 209:1422 [1980]) or hygromycin (Sugden *et al.*, Mol. Cell. Biol. 5:410-413 [1985]). The three examples given above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid), or hygromycin, respectively.

Another example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the *gas6* nucleic acid, such as DHFR or thymidine kinase. The mammalian cell transformants are placed under selection pressure that only the transformants are uniquely adapted to survive by virtue of having taken up the marker. Selection pressure is imposed by culturing the transformants under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes *gas6* variant. Increased quantities of *gas6* are synthesized from the amplified DNA. Other examples of amplifiable genes include metallothionein-I and -II, preferably primate metallothionein genes, adenosine deaminase, ornithine decarboxylase, *etc.*

For example, cells transformed with the DHFR selection gene are first identified by culturing all of the transformants in a culture medium that contains methotrexate (Mtx), a

P0929P2

competitive antagonist of DHFR. An appropriate host cell when wild-type DHFR is employed is the Chinese hamster ovary (CHO) cell line deficient in DHFR activity, prepared and propagated as described by Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216 (1980). The transformed cells are then exposed to increased levels of methotrexate. This leads to the synthesis of multiple copies of the DHFR gene, and, concomitantly, multiple copies of other DNA comprising the expression vectors, such as the DNA encoding the gas6. This amplification technique can be used with any otherwise suitable host, e.g., ATCC No. CCL61 CHO-K1, notwithstanding the presence of endogenous DHFR if, for example, a mutant DHFR gene that is highly resistant to Mtx is employed (EP 117,060).

Alternatively, host cells (particularly wild-type hosts that contain endogenous DHFR) transformed or co-transformed with DNA sequences encoding gas6, wild-type DHFR protein, and another selectable marker such as aminoglycoside 3'-phosphotransferase (APH) can be selected by cell growth in medium containing a selection agent for the selectable marker such as an aminoglycosidic antibiotic, e.g., kanamycin, neomycin, or G418. See U.S. Patent No. 4,965,199.

A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 (Stinchcomb *et al.*, Nature 282:39 [1979]; Kingsman *et al.*, Gene 7:141 [1979]; or Tschemper *et al.*, Gene 10:157 [1980]). The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 (Jones, Genetics 85:12 [1977]). The presence of the *trp1* lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan. Similarly, *Leu2*-deficient yeast strains (ATCC 20,622 or 38,626) are complemented by known plasmids bearing the *Leu2* gene.

In addition, vectors derived from the 1.6 μ m circular plasmid pKD1 can be used for transformation of *Kluyveromyces* yeasts. Bianchi *et al.*, Curr. Genet. 12:185 (1987). More recently, an expression system for large-scale production of recombinant calf chymosin was reported for *K. lactis*. Van den Berg, Bio/Technology 8:135 (1990). Stable multi-copy expression vectors for secretion of mature recombinant human serum albumin by industrial strains of *Kluyveromyces* have also been disclosed. Fler *et al.*, Bio/Technology 9:968-975 (1991).

Expression and cloning vectors usually contain a promoter that is recognized by the host organism and is operably linked to the gas6 nucleic acid. A large number of promoters recognized by a variety of potential host cells are well known. These promoters are operably linked to gas6-encoding DNA by removing the promoter from the source DNA by restriction enzyme digestion and inserting the isolated promoter sequence into the vector.

P0929P2

Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems (Chang *et al.*, Nature **275**:615 [1978]; and Goeddel *et al.*, Nature **281**:544 [1979]), alkaline phosphatase, a tryptophan (trp) promoter system (Goeddel, Nucleic Acids Res., **8**:4057 [1980] and EP 36,776) and hybrid promoters such as the tac promoter (deBoer *et al.*, Proc. Natl. Acad. Sci. USA **80**:21-25 [1983]). However, other known bacterial promoters are suitable. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to DNA encoding the gas6 (Siebenlist *et al.*, Cell **20**:269 [1980]) using linkers or adaptors to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably
 5 linked to the DNA encoding the gas6.
 10

Promoter sequences are known for eukaryotes. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3'
 15 end of most eukaryotic genes is an AATAAA sequence that may be the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase (Hitzeman *et al.*, J. Biol. Chem. **255**:2073 [1980])
 20 or other glycolytic enzymes (Hess *et al.*, J. Adv. Enzyme Reg. **7**:149 [1968]; and Holland, Biochemistry **17**:4900 [1978]), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters
 25 for use in yeast expression are further described in Hitzeman *et al.*, EP 73,657A. Yeast enhancers also are advantageously used with yeast promoters.
 30

Gas6 transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma
 35 virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most

P0929P2

preferably Simian Virus 40 (SV40), from heterologous mammalian promoters, *e.g.*, the actin promoter or an immunoglobulin promoter or from heat-shock promoters.

The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment that also contains the SV40 viral origin of replication. Fiers *et al.*, Nature 273:113 (1978); Mulligan and Berg, Science 209:1422-1427 (1980); Pavlakis *et al.*, Proc. Natl. Acad. Sci. USA 78:7398-7402 (1981). The immediate early promoter of the human cytomegalovirus is conveniently obtained as a *Hind*III E restriction fragment. Greenaway *et al.*, Gene 18:355-360 (1982). A system for expressing DNA in mammalian hosts using the bovine papilloma virus as a vector is disclosed in U.S. Patent No. 4,419,446. A modification of this system is described in U.S. Patent No. 4,601,978. See also Gray *et al.*, Nature 295:503-508 (1982) on expressing cDNA encoding immune interferon in monkey cells; Reyes *et al.*, Nature 297:598-601 (1982) on expression of human β -interferon cDNA in mouse cells under the control of a thymidine kinase promoter from herpes simplex virus; Canaani and Berg, Proc. Natl. Acad. Sci. USA 79:5166-5170 (1982) on expression of the human interferon β 1 gene in cultured mouse and rabbit cells; and Gorman *et al.*, Proc. Natl. Acad. Sci. USA 79:6777-6781 (1982) on expression of bacterial CAT sequences in CV-1 monkey kidney cells, chicken embryo fibroblasts, Chinese hamster ovary cells, HeLa cells, and mouse NIH-3T3 cells using the Rous sarcoma virus long terminal repeat as a promoter.

Transcription of DNA encoding the gas6 by higher eukaryotes is often increased by inserting an enhancer sequence into the vector. Enhancers are relatively orientation and position independent, having been found 5' (Laimins *et al.*, Proc. Natl. Acad. Sci. USA 78:993 [1981]) and 3' (Lusky *et al.*, Mol. Cell Bio. 3:1108 [1983]) to the transcription unit, within an intron (Banerji *et al.*, Cell 33:729 [1983]), as well as within the coding sequence itself (Osborne *et al.*, Mol. Cell Bio. 4:1293 [1984]). Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. See also Yaniv, Nature 297:17-18 (1982) on enhancing elements for activation of eukaryotic promoters. The enhancer may be spliced into the vector at a position 5' or 3' to the gas6-encoding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic

P0929P2

or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding the gas6.

Construction of suitable vectors containing one or more of the above listed components employs standard ligation techniques. Isolated plasmids or DNA fragments are cleaved,
5 tailored, and religated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or
10 sequenced by the method of Messing *et al.*, Nucleic Acids Res. **9**:309 (1981) or by the method of Maxam *et al.*, Methods in Enzymology **65**:499 (1980).

Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding gas6. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host
15 cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector. Sambrook *et al.*, *supra*, pp. 16.17 - 16.22. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient positive identification of polypeptides encoded by cloned DNAs, as well as for the rapid screening of gas6 variants
20 having desired binding specificities/affinities.

Other methods, vectors, and host cells suitable for adaptation to the synthesis of the gas6 in recombinant vertebrate cell culture are described in Gething *et al.*, Nature **293**:620-625 (1981); Mantei *et al.*, Nature **281**:40-46 (1979); Levinson *et al.*; EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of gas6 is
25 pRK5 (EP 307,247) or pSVI6B (PCT pub. no. WO 91/08291 published 13 June 1991).

The choice of host cell line for the expression of gas6 depends mainly on the expression vector. Where it is desired to make a gas6 variant which is essentially not γ carboxylated, it may be desirable to select a host cell which does not have γ carboxylase enzyme, especially where the nucleic acid encoding gas6 also encodes the A domain thereof.
30 Often, a useful host for this purpose is a non-mammalian cell (*e.g.* a prokaryotic cell known to be deficient in this enzyme). Alternatively a mammalian cell line can be utilized which has been made deficient in this enzyme.

Suitable host cells for cloning or expressing the vectors herein are prokaryote, yeast, or other higher eukaryote cells described above. Suitable prokaryotes for this purpose include
35 eubacteria, such as Gram-negative or Gram-positive organisms, for example,

P0929P2

Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), *Pseudomonas* such as *P. aeruginosa*, and

5 *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* X1776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting. Strain W3110 is a particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell should secrete minimal amounts of

10 proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins, with examples of such hosts including *E. coli* W3110 strain 27C7. The complete genotype of 27C7 is *tonAΔ ptr3 phoAΔE15 Δ(argF-lac)169 ompTΔ degP41kan'*. Strain 27C7 was deposited on 30 October 1991 in the American Type Culture Collection as ATCC No. 55,244. Alternatively, the strain of *E. coli* having mutant

15 periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990 may be employed. Alternatively, methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for gas6-encoding vectors. *Saccharomyces cerevisiae*,

20 or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe* (Beach and Nurse, Nature 290:140 [1981]; EP 139,383 published May 2, 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer *et al.*, *supra*) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574;

25 Louvencourt *et al.*, J. Bacteriol., 737 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilarum* (ATCC 36,906; Van den Berg *et al.*, *supra*), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna *et al.*, J. Basic Microbiol. 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case *et al.*, Proc.

30 Natl. Acad. Sci. USA 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance *et al.*, Biochem. Biophys. Res. Commun. 112:284-289 [1983]; Tilburn *et al.*, Gene 26:205-221 [1983]; Yelton *et al.*, Proc. Natl. Acad.

35 Sci. USA 81:1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J. 4:475-479 [1985]).

P0929P2

Suitable host cells for the expression of glycosylated gas6 are derived from multicellular organisms. Such host cells are capable of complex processing and glycosylation activities. In principle, any higher eukaryotic cell culture is workable, whether from vertebrate or invertebrate culture. Examples of invertebrate cells include plant and insect
 5 cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. See, e.g., Luckow *et al.*, Bio/Technology 6:47-55 (1988); Miller *et al.*, in Genetic Engineering, Setlow *et al.*, eds., Vol. 8 (Plenum Publishing, 1986), pp. 277-279; and Maeda
 10 *et al.*, Nature 315:592-594 (1985). A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, and tobacco can
 15 be utilized as hosts. Typically, plant cells are transfected by incubation with certain strains of the bacterium *Agrobacterium tumefaciens*, which has been previously manipulated to contain the gas6 DNA. During incubation of the plant cell culture with *A. tumefaciens*, the DNA encoding the gas6 is transferred to the plant cell host such that it is transfected, and will, under appropriate conditions, express the gas6 DNA. In addition, regulatory and signal
 20 sequences compatible with plant cells are available, such as the nopaline synthase promoter and polyadenylation signal sequences. Depicker *et al.*, J. Mol. Appl. Gen. 1:561 (1982). In addition, DNA segments isolated from the upstream region of the T-DNA 780 gene are capable of activating or increasing transcription levels of plant-expressible genes in recombinant DNA-containing plant tissue. EP 321,196 published 21 June 1989.

25 Propagation of vertebrate cells in culture (tissue culture) has become a routine procedure in recent years (Tissue Culture, Academic Press, Kruse and Patterson, editors [1973]). Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham *et al.*, J. Gen Virol. 36:59 [1977]);
 30 baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA 77:4216 [1980]); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 [1980]); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34);
 35 buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL

P0929P2

75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather *et al.*, Annals N.Y. Acad. Sci. 383:44-68 [1982]); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

Host cells are transfected with the above-described expression or cloning vectors of this invention and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. Depending on the host cell used, transfection is done using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in section 1.82 of Sambrook *et al.*, *supra*, or electroporation is generally used for prokaryotes or other cells that contain substantial cell-wall barriers. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw *et al.*, Gene 23:315 (1983) and WO 89/05859 published 29 June 1989. In addition, plants may be transfected using ultrasound treatment as described in WO 91/00358 published 10 January 1991.

For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology 52:456-457 (1978) is preferred. General aspects of mammalian cell host system transformations have been described by Axel in U.S. Patent No. 4,399,216 issued 16 August 1983. Transformations into yeast are typically carried out according to the method of Van Solingen *et al.*, J. Bact. 130:946 (1977) and Hsiao *et al.*, Proc. Natl. Acad. Sci. (USA) 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, *e.g.*, polybrene, polyornithine, *etc.*, may also be used. For various techniques for transforming mammalian cells, see Keown *et al.*, Methods in Enzymology (1989), Keown *et al.*, Methods in Enzymology 185:527-537 (1990), and Mansour *et al.*, Nature 336:348-352 (1988).

Prokaryotic cells used to produce the gas6 polypeptide of this invention are cultured in suitable media as described generally in Sambrook *et al.*, *supra*.

The mammalian host cells used to produce the gas6 of this invention may be cultured in a variety of media. Commercially available media such as Ham's F10 (Sigma), Minimal Essential Medium ([MEM], Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ([DMEM], Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham and Wallace, Meth. Enz. 58:44 (1979), Barnes and Sato, Anal. Biochem. 102:255 (1980), U.S. Patent Nos. 4,767,704; 4,657,866; 4,927,762; or 4,560,655; WO 90/03430; WO 87/00195; U.S. Patent Re. 30,985; or U.S. Patent No. 5,122,469, may be used as culture media

P0929P2

for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleosides (such as adenosine and thymidine), antibiotics (such as GentamycinTM drug), trace
5 elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art.

In certain embodiments, it is desirable to culture the transformed host cells in the
10 absence of Vitamin K as this can reduce γ carboxylation of the A domain of the gas6 polypeptide. Alternatively, the transformed host cells can be cultured in the presence of a carboxylase inhibitor, such as warfarin

The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled
15 artisan. In general, principles, protocols, and practical techniques for maximizing the productivity of mammalian cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed., IL Press, 1991. The host cells referred to in this disclosure encompass cells in culture as well as cells that are within a host animal.

Gas6 preferably is recovered from the culture medium as a secreted polypeptide,
20 although it also may be recovered from host cell lysates.

When gas6 is produced in a recombinant cell other than one of human origin, it is completely free of proteins or polypeptides of human origin. However, it is necessary to purify gas6 from cell proteins or polypeptides to obtain preparations that are substantially homogeneous as to gas6. As a first step, the particulate debris, either host cells or lysed
25 fragments, is removed, for example, by centrifugation or ultrafiltration; optionally, the protein may be concentrated with a commercially available protein concentration filter, followed by separating the gas6 from other impurities by one or more steps selected from heparin Sepharose chromatography, immunoaffinity chromatography, ion-exchange column fractionation (e.g., on DEAE or matrices containing carboxymethyl or sulfopropyl groups),
30 chromatography on Blue-Sepharose, CM Blue-Sepharose, MONO-Q, MONO-S, lentil lectin-Sepharose, WGA-Sepharose, Con A-Sepharose, Ether Toyopearl,^{*} Butyl Toyopearl, Phenyl Toyopearl, or protein A Sepharose, SDS-PAGE chromatography, silica chromatography, chromatofocusing, reverse phase HPLC (e.g., silica gel with appended aliphatic groups), gel filtration using, e.g., Sephadex^{*} molecular sieve or size-exclusion chromatography,
35 chromatography on columns that selectively bind the gas6, and ethanol or ammonium sulfate

P0929P2

precipitation. A protease inhibitor may be included in any of the foregoing steps to inhibit proteolysis and antibiotics may be included to prevent the growth of adventitious contaminants. Examples of suitable protease inhibitors include phenylmethanesulfonyl fluoride (PMSF), leupeptin, pepstatin, aprotinin, 4-(2-aminoethyl)-benzenesulfonyl fluoride
 5 hydrochloride-bestatin, chymostatin, and benzamidine.

Gas6 variants in which residues have been deleted, inserted, or substituted are recovered in the same fashion as native gas6, taking account of any substantial changes in properties occasioned by the variation. For example, preparation of "epitope tagged" gas6 facilitates purification using an immunoaffinity column containing antibody to the antigen to
 10 adsorb the fusion polypeptide. Immunoaffinity columns such as a rabbit polyclonal anti-gas6 column can be employed to absorb the gas6 variant by binding it to at least one remaining immune epitope. One skilled in the art will appreciate that purification methods suitable for native gas6 may require modification to account for changes in the character of gas6 or its variants upon production in recombinant cell culture.

15 3. *In Vitro* and *In Vivo* Uses for Gas6

The present invention provides methods for activating Rse receptor or Mer receptor and/or enhancing survival, proliferation or differentiation of cells comprising the Rse or Mer receptor using gas6. The gas6 useful in the practice of the present invention can be prepared in a number of ways which have been described in the previous section (see also Example 6
 20 below).

The gas6 may be from a human or any non-human species. For instance, a mammal may be treated with gas6 from a different mammalian species (*e.g.*, mice can be treated with human gas6). There is substantial homology (about 81% amino acid identity) between murine gas6 and human gas6, and thus, it is expected that gas6 from different mammalian species can
 25 be employed. Preferably, however, the mammal is treated with homologous gas6 (*e.g.*, humans are treated with human gas6) to avoid potential immunogenicity of the gas6 in the mammal.

The present invention includes methods of activating Rse or Mer receptor and/or enhancing survival, proliferation or differentiation of cells comprising the Rse or Mer receptor
 30 *in vivo* and *in vitro*. Normally, the cells will be treated with the gas6 polypeptide. However, gene therapy approaches have been described in the art and are encompassed by the present invention. These techniques include gene delivery to a cell using adenovirus, herpes simplex I virus or adeno-associated virus as well as lipid-based delivery systems (*e.g.* liposomes). Retroviruses are useful for *ex vivo* gene therapy approaches. Accordingly, it is possible to

P0929P2

administer the nucleic acid encoding gas6, resulting in expression of the gas6 polypeptide in the patient or in tissue culture. For exemplary gene therapy techniques see WO 93/25673 and the references cited therein.

In accordance with the *in vitro* methods of the invention, cells comprising the Rse or Mer receptor are provided and placed in a cell culture medium. Examples of Rse-receptor-containing cells include neural cells, *e.g.*, brain cells (such as neurons of the neocortex, cerebellum and hippocampus); glial cells (*e.g.* Schwann cells or astrocytes); kidney or breast-derived cells; cells derived from the ovary or testes; fibroblast cells such as mouse 3T3 cells; cells from the hematopoietic system such as CMK11-5. Examples of Mer-receptor-containing cells include peripheral blood mononuclear cells, bone marrow mononuclear cells, monocytes, primary hematopoietic cells and cells derived from testis, ovary, prostate, lung, kidney, spleen, peripheral blood leukocyte, placenta, thymus, small intestine, colon or liver. Exemplary cell lines to be cultured using gas6 include T lymphocyte leukemia cell lines (*e.g.* CCRF-HSB-2, JURKAT, HPB-ALL and Peer); K-562 cell line; monocytic leukemia/lymphoma cell lines (such as U-937); megakaryoblastic leukemia cell lines (*e.g.* UT-7) and other cell lines which express Mer receptor as described in Graham *et al.*, Cell Growth Differ. 5:647 (1994).

Suitable tissue culture media are well known to persons skilled in the art and include, but are not limited to, Minimal Essential Medium (MEM), RPMI-1640, and Dulbecco's Modified Eagle's Medium (DMEM). These tissue culture medias are commercially available from Sigma Chemical Company (St. Louis, MO) and GIBCO (Grand Island, NY). The cells are then cultured in the cell culture medium under conditions sufficient for the cells to remain viable and grow in the presence of an effective amount of gas6. The cells can be cultured in a variety of ways, including culturing in a clot, agar, or liquid culture.

The cells are cultured at a physiologically acceptable temperature such as 37°C, for example, in the presence of an effective amount of gas6. The amount of gas6 may vary, but preferably is in the range of about 10 ng/ml to about 1mg/ml. The gas6 can of course be added to the culture at a dose determined empirically by those in the art without undue experimentation. The concentration of gas6 in the culture will depend on various factors, such as the conditions under which the cells and gas6 are cultured. The specific temperature and duration of incubation, as well as other culture conditions, can be varied depending on such factors as, *e.g.*, the concentration of the gas6, and the type of cells and medium. Those skilled in the art will be able to determine operative and optimal culture conditions without undue experimentation. Proliferation, differentiation and/or survival of the cells (*e.g.* neurons

P0929P2

or mononuclear cells) in the cultures can be determined by various assays known in the art such as those described above.

It is contemplated that using gas6 to enhance cell survival, growth and/or differentiation *in vitro* will be useful in a variety of ways. For instance, neural cells cultured
 5 *in vitro* in the presence of gas6 can be infused into a mammal suffering from reduced levels of the cells. In other embodiments, gas6 may be used to culture hematopoietic cells (such as monocytes/macrophages) *ex vivo* which can be administered to a patient having decreased levels of these blood cells (for example where the patient has undergone chemo- or radiation therapy). Stable *in vitro* cultures can also be used for isolating cell-specific factors and for
 10 expression of endogenous or recombinantly introduced proteins in the cell. Gas6 may also be used to enhance cell survival, proliferation and/or differentiation of cells which support the growth and/or differentiation of other cells in cell culture (*e.g.* stromal cells supporting bone marrow non-adherent cells). In this manner, Schwann cells may promote neuronal survival in cell culture.

15 Gas6 is considered to be particularly useful for growing Schwann cells *ex vivo*. It is desirable to have such populations of cells in cell culture for isolation of cell-specific factors *e.g.* P75^{NGFR} which is a Schwann cell specific marker. Such factors are useful as diagnostic tools or, in the case of P75^{NGFR}, can be used as antigens to generate antibodies for diagnostic use. It is also desirable to have stable populations of Schwann cells in cell culture to facilitate
 20 characterization of other mitogens and growth inhibitory agents for these cells.

The invention also provides *in vivo* uses for gas6. Based on the ability of gas6 to promote proliferation of glial cells (see Example 9), it is believed that this molecule will be particularly useful for treating diseases which involve demyelination, damage or loss of glial cells (*e.g.* multiple sclerosis).

25 Gas6 is also believed to be useful in promoting the development, maintenance, and/or regeneration of neurons *in vivo*, including central (brain and spinal chord), peripheral (sympathetic, parasympathetic, sensory, and enteric neurons), and motoneurons. Accordingly, gas6 may be utilized in methods for the diagnosis and/or treatment of a variety of "neurologic diseases or disorders" which effect the nervous system of a mammal, such as
 30 a human.

Such diseases or disorders may arise in a patient in whom the nervous system has been damaged by, *e.g.*, trauma, surgery, stroke, ischemia, infection, metabolic disease, nutritional deficiency, malignancy, or toxic agents. The agent is designed to promote the survival or growth of neurons. For example, gas6 can be used to promote the survival or growth of
 35 motoneurons that are damaged by trauma or surgery. Also, gas6 can be used to treat

P0929P2

motoneuron disorders, such as amyotrophic lateral sclerosis (Lou Gehrig's disease), Bell's palsy, and various conditions involving spinal muscular atrophy, or paralysis. Gas6 can be used to treat human "neurodegenerative disorders", such as Alzheimer's disease, Parkinson's disease, epilepsy, multiple sclerosis, Huntington's chorea, Down's Syndrome, nerve deafness, and Meniere's disease.

Further, gas6 can be used to treat neuropathy, and especially peripheral neuropathy. "Peripheral neuropathy" refers to a disorder affecting the peripheral nervous system, most often manifested as one or a combination of motor, sensory, sensorimotor, or autonomic neural dysfunction. The wide variety of morphologies exhibited by peripheral neuropathies can each be attributed uniquely to an equally wide number of causes. For example, peripheral neuropathies can be genetically acquired, can result from a systemic disease, or can be induced by a toxic agent. Examples include but are not limited to distal sensorimotor neuropathy, or autonomic neuropathies such as reduced motility of the gastrointestinal tract or atony of the urinary bladder. Examples of neuropathies associated with systemic disease include post-polio syndrome; examples of hereditary neuropathies include Charcot-Marie-Tooth disease, Refsum's disease, Abetalipoproteinemia, Tangier disease, Krabbe's disease, Metachromatic leukodystrophy, Fabry's disease, and Dejerine-Sottas syndrome; and examples of neuropathies caused by a toxic agent include those caused by treatment with a chemotherapeutic agent such as vincristine, cisplatin, methotrexate, or 3'-azido-3'-deoxythymidine.

Given expression of Rse receptor and Mer receptor on hematopoietic cells, gas6 may be used to enhance repopulation of mature blood cell lineages in patients having undergone chemo- or radiation therapy or bone marrow transplantation therapy. It is contemplated that gas6 will act via an enhancement of the proliferation and/or differentiation of hematopoietic cells (*e.g.* monocytes and megakaryocytes). Gas6 may similarly be useful for treating diseases characterized by a decrease in blood cells. Examples of these diseases include anemia (including macrocytic and aplastic anemia); thrombocytopenia; monocytopenia; hypoplasia; immune (autoimmune) thrombocytopenic purpura (ITP); and HIV induced ITP. Gas6 may also be used to promote growth and/or repair of tissues (*e.g.* testis, ovary, prostate, lung or kidney) which express either, or both, of these receptors. Also, gas6 may be used to improve reproductive function, given high level expression of Mer receptor in the testes and ovaries.

In that Mer is expressed on mononuclear cells, it is contemplated that gas6 may be used to treat conditions in which proliferation and/or differentiation of these cells is desired. For example, gas6 may be used to increase monocyte (*e.g.* macrophage) levels in a patient where this is required or desired.

P0929P2

In other embodiments, gas6 may be used to modulate function of cells possessing the Mer or Rse receptor. For example, gas6 may be used to activate monocytes/macrophages in situations where such activation is desired (*e.g.* to treat infections).

In still further embodiments of the invention, gas6 antagonists, and especially anti-
 5 gas6 antibodies, can be administered to patients suffering from diseases or disorders (*e.g.*, neurologic diseases or disorders) characterized by excessive production of gas6 and/or excessive Rse or Mer receptor activation by gas6. Gas6 antagonist may be used in the prevention of aberrant regeneration of sensory neurons such as may occur post-operatively, or in the selective ablation of sensory neurons, for example, in the treatment of chronic pain
 10 syndromes. Gas6 antagonists may also be used to treat monocytosis or malignancies characterized by excessive Rse or Mer receptor activation (such as lymphoid malignancies) or to treat monocyte/macrophage-mediated inflammation.

Therapeutic formulations of gas6 and gas6 antagonist are prepared by mixing gas6 or gas6 antagonist, having the desired degree of purity, with optional pharmaceutically
 15 acceptable carriers, excipients, or stabilizers which are well known. Acceptable carriers, excipients or stabilizers are nontoxic to the patient at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as
 20 polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, Pluronic or polyethylene glycol (PEG).

25 It may be desirable to adsorb gas6 onto a membrane, such as a silastic membrane, which can be implanted in proximity to damaged neural tissue, or to incorporate gas6 into liposomes. PCT Pub. No. WO 91/04014 (published April 4, 1991). In another embodiment, the gas6 used for therapeutic effect is gas6 covalently joined to another protein, such as an immunoglobulin domain (for example, to produce gas6-IgG).

30 Gas6 optionally is combined with or administered in concert with other neurotrophic factors to achieve a desired therapeutic effect. For example, gas6 may be used together with nerve growth factor (NGF), neurotrophins (NT-3), bone derived nerve factor (BDNF), neurotrophins-4 and -5 (NT-4/5), an insulin-like growth factor (*e.g.*, IGF-1 or IGF-2) or another neurotrophic factor to achieve a synergistic stimulatory effect on the growth of
 35 sensory neurons, wherein the term "synergistic" means that the effect of the combination of

P0929P2

gas6 with a second substance is greater than that achieved with either substance used individually.

For use in hematopoiesis, gas6 may be administered in concert with one or more cytokines. Included among the cytokines are growth hormone, insulin-like growth factors, human growth hormone, N-methionyl human growth hormone, bovine growth hormone, parathyroid hormone, thyroxine, insulin, proinsulin, relaxin, prorelaxin, glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH), hematopoietic growth factor, hepatic growth factor, fibroblast growth factor, prolactin, placental lactogen, tumor necrosis factor- α and - β , mullerian-inhibiting substance, mouse gonadotropin-associated peptide, inhibin, activin, vascular endothelial growth factor, integrin, thrombopoietin, nerve growth factors such as NGF- β , platelet-growth factor, transforming growth factors (TGFs) such as TGF- α and TGF- β , insulin-like growth factor-I and -II, erythropoietin (EPO), osteoinductive factors, interferons such as interferon- α , - β , and - γ , colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF), granulocyte-macrophage-CSF (GM-CSF), and granulocyte-CSF (G-CSF), interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12 and other polypeptide factors including LIF, SCF, and kit-ligand. As used herein the foregoing terms are meant to include proteins from natural sources or from recombinant cell culture. Similarly, the terms are intended to include biologically active equivalents; *e.g.*, differing in amino acid sequence by one or more amino acids or in type or extent of glycosylation.

Gas6 and gas6 antagonist to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration of a solution of gas6 or gas6 antagonist through sterile filtration membranes. Thereafter, the filtered solution may be placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle. The filtered solution also may be lyophilized to produce sterile gas6 or gas6 antagonist in a powder form.

Methods for administering gas6 and gas6 antagonists *in vivo* include injection or infusion by intravenous, intraperitoneal, intracerebral, intrathecal, intramuscular, intraocular, intraarterial, or intralesional routes, and by means of sustained-release formulations.

Sustained-release formulations generally consist of gas6 or gas6 antagonist and a matrix from which the gas6 or gas6 antagonist are released over some period of time. Suitable matrices include semipermeable polymer matrices in the form of shaped articles, for example, membranes, fibers, or microcapsules. Sustained release matrices may comprise polyesters, hydrogels, polylactides, U.S. Pat. No. 3,773,919, copolymers of L-glutamic acid

P0929P2

and γ -ethyl-L-glutamate, Sidman *et al.*, Biopolymers 22:547-556 (1983), poly (2-hydroxyethyl-methacrylate), or ethylene vinyl acetate, Langer *et al.*, J. Biomed. Mater. Res. 15:167-277 (1981); and Langer, Chem. Tech. 12:98-105 (1982).

In one embodiment of the invention, the therapeutic formulation comprises gas6 or gas6 antagonist entrapped within or complexed with liposomes. For example, gas6 covalently joined to a glycerophosphatidyl-inositol moiety may be used to form a liposome comprising gas6. In a further embodiment, the therapeutic formulation comprises cells actively producing gas6 or gas6 antagonist. Such cells may be directly introduced into the tissue of a patient, or may be encapsulated within porous membranes which are then implanted in a patient, in either case providing for the delivery of gas6 or gas6 antagonist into areas within the body of the patient in need of increased or decreased concentrations of gas6. Alternatively, an expression vector comprising gas6 DNA may be used for *in vivo* transformation of a patient's cells to accomplish the same result.

An effective amount of gas6 or gas6 antagonist to be employed therapeutically will depend, for example, upon the therapeutic objectives, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage might range from about 1 μ g/kg to up to 100 mg/kg or more, depending on the factors mentioned above. Where possible, it is desirable to determine appropriate dosage ranges first *in vitro*, for example, using assays for cell survival or growth which are known in the art, and then in suitable animal models, from which dosage ranges for human patients may be extrapolated. In a specific embodiment of the invention, a pharmaceutical composition effective in promoting the survival or growth of neurons will provide a local gas6 concentration *in vivo* of between about 0.1 and 10 ng/ml.

The invention further provides an article of manufacture and kit containing materials useful for activating the Rse or Mer receptor or enhancing survival, proliferation or differentiation of cells comprising the Rse or Mer receptor. The article of manufacture comprises a container with a label. Suitable containers include, for example, bottles, vials, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for activating the Rse or Mer receptor and/or enhancing survival, proliferation and/or differentiation of cells having the receptor of interest. The active agent in the composition is gas6. The label on the container indicates that the composition is used for activating the Rse or Mer receptor and/or enhancing survival, proliferation and/or differentiation of cells having this receptor, and may also indicate directions for either *in vivo* or *in vitro* use, such as those described above.

P0929P2

The kit of the invention comprises the container described above and a second container comprising a buffer. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use.

- 5 The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention.

Example 1

Production of Rse-IgG Fusion Protein

- 10 To identify a source of a Rse ligand (Rse-L), a fusion protein containing the extracellular domain of human Rse followed by the Fc portion of human IgG (Rse-IgG) was used as a probe to screen cells for surface bound Rse-L using flow cytometry (see Example 2 below). Rse-IgG was constructed by fusing the sequence encoding the extracellular domain (amino acids 1-428) of human Rse (Mark *et al.*, Journal of Biological Chemistry
15 269(14):10720-10728 [1994]) to amino acids 216-443 of human IgG γ_1 through a BstEII linker (adding amino acids Val and Thr). The linker was added to Rse sequences by PCR using the primers (5'- TCAAGACAATGGAACCCAGG [SEQ ID NO: 4] and 5'- CATGGAATTCGGTGACCGATGTGCGGCTGTGAGGAG [SEQ ID NO: 5]). The cDNA encoding Rse-IgG was transferred into an SV40 based expression vector and
20 introduced into DHFR⁻ CHO cells by electroporation (250 volts, 960 μ F). DHFR⁺ cells were selected and Rse-IgG expression in individual clones was determined using a human Fc-specific ELISA. Rse-IgG was purified on a protein A-Sepharose column (Pharmacia).

Example 2

Binding Analysis

- 25 Fluorescence activated cell sorting (FACS) analysis using Rse-IgG was performed as described in Goodwin *et al.*, Cell 73:447 (1993). The megakaryocytic leukemia line CMK11-5 cells (Adachi *et al.*, Exp Hematol. 19:923 [1991]) specifically bound Rse-IgG but not control fusion proteins containing the identical Fc domain such as HGFr-IgG (Mark *et al.*, J. Biol. Chem. 267:26166 [1992]) or CD4-IgG (Capon *et al.*, Nature 337:525 [1989]). Binding
30 of Rse-IgG was increased by the addition of Ca²⁺ and abolished by treatment with 2 mM EDTA.

Subsequently, an *in vitro* binding assay was established to characterize the interaction of ¹²⁵I-Rse-IgG with the putative cell surface bound Rse-L. CMK11-5 cells were suspended

P0929P2

in 10 mM TrisCl, pH 7.5 for 10 min on ice, lysed by a combination of sonication and shearing, and whole membranes collected by centrifugation and stored in 50 mM TrisCl, pH 7.5, 20% glycerol at -80°C. Membranes equivalent to 200,000 cells were combined with fetal bovine serum (FBS) or column fractions, competitors, and ¹²⁵I-Rse-IgG in a total volume of 0.1-0.12 ml. After a 30 min incubation at room temperature, 1 ml of ice cold assay buffer was added to each tube. Then, the membrane associated radioactivity was collected by centrifugation for 4 min at 15000g, separated from unbound radioactivity by aspiration of the supernatant fluid and counted in a γ counter. The assay buffer was 50 mM Tris-HCl, 0.05% Tween-20, 0.1% BSA, 5 mM CaCl₂.

Because flow cytometric analyses were performed in the presence of serum, the effect of FBS in the membrane binding assay was determined. Binding was found to be absolutely dependent on FBS concentration; no displaceable binding was seen in the absence of FBS and half maximum binding was observed with 0.58% FBS (Fig. 3A).

Binding was also Ca²⁺-dependent; half maximum binding was obtained with 0.18 mM Ca²⁺ (Fig. 3B). Although the apparent number of binding sites for Rse-IgG was dependent on the concentration of FBS, the affinity was not greatly changed [*K_d* of 0.82 nM in 1% FBS vs. 2.2 nM in 10% FBS] (Fig. 3C). Binding was specific; other recombinant IgG fusion proteins, such as CD4-IgG, did not compete for binding with ¹²⁵I-Rse-IgG.

Example 3

Epitope-Tagged Rse Receptor and Activation Thereof

Chinese Hamster Ovary (CHO) cells expressing a version of Rse receptor having a Herpes simplex virus type I (HSV-1) C-terminal glycoprotein D (gD) flag (Paborsky *et al.*, Protein Engineering 3(6):547-553 [1990]) were generated in order to further characterize a Rse-L.

Synthetic double stranded oligonucleotides were used to reconstitute the coding sequence for the C-terminal 10 amino acids (880 - 890) of human Rse and add an additional 21 amino acids containing the gD epitope for the antibody 5B6 (Paborksy *et al.*, *supra*) and a stop codon. The final sequence of the synthetic portion of the fusion gene was:

coding strand:

5'-GCAAGGGCTACTGCCACACTCGAGCTGCGCAGATGCTAGCCTCAAGATGGCTG
ATCCAAATCGATTCCGCGGCAAAGATCTTCCGGTCCTGTAGA-3' [SEQ ID NO: 6]

noncoding strand:

AGCTTCTACAGGACCGGAAGATCTTTGCCGCGGAATCGATTTGGATCAGCCAT

P0929P2

CTTG AGGCTAGCATCTGCGCAGCTCGAGTGTGGCAGTAGCCCTTGCTGCA-3' [SEQ ID NO: 7].

The synthetic DNA was ligated with the cDNA encoding amino acids 1-880 of human Rse at the PstI site beginning at nucleotide 2644 of the published human Rse cDNA sequence (Mark *et al.*, Journal of Biological Chemistry **269**(14):10720-10728 [1994]) and HindIII sites in the polylinker of the expression vector pSVI7.ID.LL (see PCT/US94/13329) derived from the vector pRK (Suva *et al.*, Science, **237**:893-896 [1987]) to create the expression plasmid pSV.ID.Rse.gD. Briefly, the expression plasmid comprises a dicistronic primary transcript which contains sequence encoding DHFR bounded by 5' splice donor and 3' splice acceptor intron splice sites, followed by sequence that encodes the Rse.gD. The full length (non-spliced) message contains DHFR as the first open reading frame and therefore generates DHFR protein to allow selection of stable transformants.

dp12.CHO cells (EP 307,247 published 15 March 1989) were electroporated with pSV.ID.Rse.gD which had been linearized at a unique NotI site in the plasmid backbone. The DNA was ethanol precipitated after phenol/chloroform extraction and was resuspended in 10 μ l 10/1 Tris/EDTA. Then, 20 μ g of DNA was incubated with 10⁷ CHO.dp12 cells in 1 ml of PBS on ice for 10 min. before electroporation at 350 volts and 330 μ f. Cells were returned to ice for 10 min. before being plated into non-selective medium. After 24 hours cells were fed nucleoside-free medium to select for stable DHFR+ clones.

To identify a cell line that expresses Rse.gD nucleic acid, candidate clones were screened by FACS analysis using the polyclonal antiserum 19B which recognizes epitopes in the extracellular domain of Rse. To confirm that clones that scored positive in the FACS assay express full-length Rse.gD nucleic acid, cell lysates were prepared (Lokker *et al.*, EMBO J, **11**:2503-2510 [1992]) and solubilized Rse.gD was immunoprecipitated with the 19B antisera. The immunoprecipitated proteins were fractionated under reducing conditions using 7% PAGE, blotted onto nitrocellulose and then probed with the anti-gD 5B6 antibody (Paborsky *et al.*, *supra*) which was detected with a horseradish peroxidase conjugated anti-mouse IgG antibody.

The ability of Rse.gD in cell clones to undergo autophosphorylation in response to 20% FBS, partially purified fractions of FBS containing the Rse receptor binding activity (*i.e.* 1:10 dilution of the QSE fraction obtained in Example 5 below) or control (*i.e.* no additions) was determined by Western blotting. Briefly, 5 x 10⁵ dp12.CHO cells transformed with Rse.gD nucleic acid as described above were seeded on a 60mm dish in the presence of serum for 6h. The cells were then washed in phosphate buffered saline (PBS) and serum-starved for 16h. The serum starved cells were then exposed to the sample for 10 min. The Rse.gD

P0929P2

protein was immunoprecipitated from CHO cell lysates using the anti-gD 5B6 monoclonal antibody. Proteins were fractionated on 7% SDS-PAGE under reducing conditions and transferred to nitrocellulose. Phosphorylation of Rse was detected with labelled anti-phosphotyrosine antibody 4G10 (obtained commercially from UBI, New York).

- 5 Addition of either 20% FBS or partially purified fractions of FBS containing the Rse-IgG binding activity to serum-starved cells expressing Rse-gD resulted in phosphorylation of the 140 kDa Rse receptor on tyrosine residues. The Rse receptor was not activated by the control.

Example 4

10

KIRA ELISA

The activity in FBS that activated Rse.gD was further characterized using an ELISA-based "KIRA" (for *Kinase Receptor Activation*) assay that allows high-throughput analysis of potential Rse-L sources. See Figure 4 for a schematic representation of this assay.

- Rse.gD transformed dp12.CHO cells produced as described in Example 3 were seeded
15 (5x10⁴ per well) in the wells of a flat-bottom-96 well culture plate in 100 µl media and cultured overnight at 37°C in 5% CO₂. The following morning the well supernatants were decanted, and the plates were lightly tamped on a paper towel. 50µl of media containing QSE fraction obtained as described in Example 5 below or control (*i.e.* media alone) was then added to each well. For neutralization experiments, potential ligand sources were treated at
20 room temperature for 30min with Rse-IgG or CD4-IgG (100µg/ml) prior to addition to the cells. The cells were stimulated at 37°C for 30 min., the well supernatants were decanted, and the plates were once again lightly tamped on a paper towel. To lyse the cells and solubilize the receptors, 100 µl of lysis buffer was added to each well. Lysis buffer consisted of 150 mM NaCl containing 50 mM HEPES (Gibco), 0.5 % Triton-X 100* (Gibco), 0.01 % thimerosal,
25 30 KIU/ml aprotinin (ICN Biochemicals, Aurora, OH), 1mM 4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride (AEBSF; ICN Biochemicals), 50 µM leupeptin (ICN Biochemicals), and 2 mM sodium orthovanadate (Na₃VO₄; Sigma Chemical Co, St. Louis, MO), pH 7.5. The plate was then agitated gently on a plate shaker (Bellco Instruments, Vineland, NJ) for 60 min. at room temperature.

- 30 While the cells were being solubilized, an ELISA microtiter plate (Nunc Maxisorp*, Inter Med, Denmark) coated overnight at 4°C with the 5B6 monoclonal anti-gD antibody (0.5 µg/ml in 50 mM carbonate buffer, pH 9.6, 100 µl/well) was decanted, tamped on a paper towel and blocked with 150 µl/well of Block Buffer* (PBS containing 0.5 % BSA [Intergen Company, Purchase, NY] and 0.01 % thimerosal) for 60 min. at room temperature with gentle

P0929P2

agitation. After 60 minutes, the anti-gD 5B6 coated plate was washed 6 times with wash buffer (PBS containing 0.05 % Tween-20 and 0.01 % thimerosal) using an automated plate washer (ScanWasher*300, Skatron Instruments, Inc, Sterling, VA).

5 The lysate containing solubilized Rse.gD from the cell-culture microtiter well was transferred (85 μ l/well) to anti-gD 5B6 coated and blocked ELISA well and was incubated for 2 h at room temperature with gentle agitation. The unbound Rse.gD was removed by washing with wash buffer and 100 μ l of biotinylated 4G10 (anti-phosphotyrosine) at 0.15 μ g/ml in buffer (PBS containing 0.5 % BSA, 0.05 % Tween-20, 5 mM EDTA, and 0.01 % thimerosal), was added to each well. After incubation for 2 h at room temperature the plate
10 was washed and 100 μ l of HRPO-conjugated streptavidin (Zymed Laboratories, S. San Francisco, CA) diluted 1:6x10⁴ in dilution buffer was added to each well. The plate was incubated for 30 minutes at room temperature with gentle agitation. The free avidin-conjugate was washed away and 100 μ l freshly prepared substrate solution (tetramethyl benzidine [TMB]; 2-component substrate kit; Kirkegard and Perry, Gaithersburg, MD) was added to
15 each well. The reaction was allowed to proceed for 10 minutes, after which the color development was stopped by the addition of 100 μ l/well 1.0 M H₃PO₄. The absorbance at 450 nm was read with a reference wavelength of 650 nm (ABS_{450/650}), using a *vmax* plate reader (Molecular Devices, Palo Alto, CA) controlled with a Macintosh Centris*650 (Apple Computers, Cupertino, CA) and DeltaSoft*software (BioMetallics, Inc, Princeton, NJ).

20 Phosphorylation of Rse.gD was stimulated in a dose dependent fashion and this activity was neutralized by Rse-IgG but not by the control CD4-IgG (Fig. 3D). These data show that a ligand capable of activating Rse is present in FBS.

Example 5

Rse Ligand Characterization

25 The Rse-L was purified from FBS by ion exchange and Rse affinity chromatography (see Table 2 below).

P0929P2

TABLE 2
Purification of a Rse Ligand from FBS

	Protein (mg)	Units	Yield (%)	Specific Activity (SA) (Units/mg)	Fold Purification
FBS	2800	196	100	0.07	1
QSE	12.8	94	48	7.4	105
5 Rse-IgG Affinity	0.183	22	11	119	1701

Fetal bovine serum (FBS) was dialyzed (molecular weight cut off 6000 Da) against 50 mM Tris HCl pH 7.5 and sterile filtered (0.22 μ cellulose nitrate, Corning) before loading onto a Q-Sepharose column equilibrated in buffer A, 10 mM Tris HCl, pH 7.5. Buffer B was
 10 buffer A with 1M NaCl. The column was eluted with a 1 column volume gradient from 0 to 18% B, then a 10 column volume gradient of 18 to 60% B. Active fractions, eluting near 0.4 M NaCl, were pooled and dialyzed against 50 mM Tris HCl pH 7.5, 5 mM benzamidine. This Q-Sepharose enriched fraction (QSE) was applied to a Rse-IgG affinity column. The column was washed with 50 mM Tris HCl, pH 7.5, 5 mM benzamidine and eluted with 4 M Urea,
 15 0.1M Tris HCl, pH 7.5, 5 mM benzamidine. The eluate was concentrated and dialyzed by centrifugal ultrafiltration (Centricon 10). Rse-IgG columns were prepared using 2 mg of Rse-IgG per ml Emphase resin according to the supplier's instructions (Pierce). The quantities tabulated are relative to 100 ml of FBS starting material. One unit of binding activity is defined as the amount present in 1 ml of a sample having an EC₅₀ of 1% v/v in the *in vitro*
 20 binding activation assay described in Example 2 above.

Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) of the affinity-purified Rse-L preparation showed a broad band centered at 60 kDa with unreduced samples which resolved into several closely spaced bands from 65 to 68kDa upon reduction. Fractions were heated for 10 min at 90°C in sample buffer, resolved on a 4-20% SDS
 25 polyacrylamide gel (Novex) and visualized by silver staining. The eluate of Rse-IgG affinity purified Rse-L was reduced with 25 mM DTT prior to electrophoresis.

The Rse-L preparation was separated by SDS-PAGE under reducing conditions, electroblotted and sequenced. Electroblotting onto Millipore Immobilon[®]PSQ membranes was carried out for 1 h at 250 mA constant current in a Biorad transblot transfer cell as described
 30 (P. Matsudaira, J. Biol. Chem. 262:10035 [1987]). The membrane was stained with 0.1%

P0929P2

Coomassie Blue R-250 in 50% methanol for 30 sec, destained with 10% acetic acid in 50% methanol for 2 to 3 min, thoroughly washed with distilled water, dried, and stored at 4° C. Automated protein sequencing was performed on models 473A and 490A Applied Biosystems Sequencers equipped with on-line PTH analyzers. Peaks were integrated with
5 Justice Innovation software using Nelson Analytical 760 interfaces. Sequence interpretation was performed as described (Henzel *et al.* J. Chromatogr. **404**:41 [1987]).

The preparation gave an amino-terminal sequence of XQVLIRRXRANTL [SEQ ID NO: 8], corresponding to that of bovine protein S. Protein S sequences were obtained from several independent preparations of Rse-L. After SDS-PAGE, some preparations were
10 characterized by the presence of a 14 kDa species having an N-terminal sequence of ANTL [SEQ ID NO: 9], as previously reported for bovine protein S, along with 60-70 kDa species with sequences corresponding to cleavage within the thrombin sensitive loop region of bovine protein S. After CnBr cleavage of the sequencing filter >99% of all identifiable residues were accounted for by a mixture of protein S CnBr fragments. Furthermore, Rse-L activity could
15 not be separated from protein S by anion exchange chromatography in the presence of Ca²⁺, cation exchange chromatography, hydrophobic interaction chromatography, Blue Sepharose chromatography, or nondenaturing gel electrophoresis. The Rse-L activity present in FBS and purified fractions could be neutralized by protein S polyclonal antisera.

Human serum or recombinant human protein S expressed in 293 cells showed low
20 activity in either the KIRA or Rse-IgG binding assays. Human serum was obtained from Pierce and from local blood banks. Human protein S (Calbiochem, Enzyme Research Labs, or Celsus labs) had an EC₅₀ of >250 nM in the membrane binding assay. In comparison, the purified bovine protein S had an EC₅₀ of 1.2 nM in this assay. In the KIRA assay, concentrations as high as 150 nM human protein S resulted in low phosphorylation of Rse.
25 Human protein S cDNAs were obtained by PCR using 1 µg of human fetal liver cDNA (Clontech) as template with Pfu DNA polymerase (Stratagene) as described in Mark *et al.* (1992), *supra*. Human protein S was expressed in 293 cells grown in the presence of 2 µg/ml Vitamin K exactly as described below for human gas6, and expression was verified by metabolic labeling of cultures and/or by western blotting with a polyclonal anti-protein S
30 antiserum. Purified human protein S bound ¹²⁵I-Rse-IgG, but with ~200 fold lower affinity than purified bovine protein S.

It was hypothesized that a homologue of protein S might be more effective. A search of the GENBANK data base revealed substantial similarity (44% amino acid identity, and a similar domain structure) between the amino acid sequence of human protein S and the
35 predicted product of human growth arrest specific gene 6 (gas6) (Manfioletti *et al.*, *supra*).

Example 6**Recombinant Production of gas6**

It was determined whether human gas6 was a ligand for Rse. Gas6 cDNA clones were obtained by polymerase chain reaction cloning from reverse transcribed human brain cDNA.

5 The full-length human gas6 clone was constructed by linking together cDNAs encoding amino acids 1-318 and 319-678. Gas6 cDNAs were obtained by PCR using 1 µg of human fetal brain cDNA (Clontech) as template with Pfu DNA polymerase as described (Mark *et al.*, J. Biol. Chem. **267**:26166 [1992]). Forward and reverse primers designed to obtain the 5' and 3' portions of hgas6 were:

10 (5'-GATATCGATCCATGGCCCCTTCGCTCTC [SEQ ID NO:10];
5'-CATGGATCCTACCGGAAGTCAAACCTCAGCTA [SEQ ID NO: 11]) and
(5'-GATATCGATGAGTGTGAAGTCCTTGAC [SEQ ID NO: 12];
5'-GTCGGATCCGACAGAGACTGAGAAGCC [SEQ ID NO: 13]), respectively.

Human fetal kidney 293 cells were transiently transfected as described in Mark *et al.*,
15 J. Biol. Chem. **267**:26166 (1992). After a 4 h incubation, the media was replaced with growth media plus antibiotics and 2 µg/ml Vitamin K. Conditions for metabolic labeling with ³⁵S-Cys and ³⁵S-Met were as described in Mark *et al.* For precipitation with IgG-fusion proteins, radiolabeled supernatants were first precleared with pansorbin (Calbiochem) for 30 min at room temperature, then incubated with 10 µg of the IgG fusion protein for 4 h at 4°C. Fusion
20 proteins were precipitated with 20 µl of pansorbin, the complexes were collected by centrifugation at 14,000 x g for 1 min, and then washed 3 times with PBS containing 0.1% Triton-X 100. Precipitates were analyzed by SDS-PAGE under reducing conditions (Capon *et al.*, Nature **337**:525 [1989]). Radioactivity in the dried gel was analyzed with a Fuji phosphoimager.

25 Conditioned media from cells metabolically labeled after transfection with a gas6 expression vector contained a 70 kDa protein that could be selectively precipitated by the Rse-IgG fusion protein but not by the control fusion protein CD4-IgG. Conditioned media from unlabeled transfections enhanced binding of ¹²⁵I-Rse-IgG to membranes, and induced phosphorylation of Rse receptor expressed in CHO cells. These data indicated that
30 recombinant human gas6 binds to and activates human Rse receptor.

Recombinant gas6 was purified from conditioned media by affinity chromatography. Human fetal kidney 293 cells were transiently transfected as described in Mark *et al.* (1992), *supra*. After a 4 h incubation, the media was replaced with serum free growth media plus antibiotics and 2 µg/ml Vitamin K. Conditioned media were collected at two and 4 days
35 following transfection. The conditioned media of the transfected cells, but not those of either

P0929P2

nontransfected or mock transfected 293 cells, activated binding of ^{125}I -Rse-IgG. A liter of pooled conditioned media was clarified by centrifugation, diluted with 1 volume of buffer A (50 mM TrisHCl, pH 7.5, 0.1% CHAPS, 5 mM benzamidine), and applied to a 6 ml Resource Q column (Pharmacia) previously equilibrated with buffer A. The column was eluted with a
 5 12 column volume gradient of 0 to 0.4 M NaCl in buffer A. The active fractions were pooled and diluted with 1 volume buffer A and applied to a Rse-IgG affinity column that was washed and developed as described (see Example 5 above).

The identity of recombinant gas6 was verified by amino terminal sequence. The sequence of the recombinant material begins with the sequence $^{49}\text{AFQVFEEA}^{56}$ [SEQ ID NO:
 10 14]. The signal from the glutamic acid residues in this sequence was weak, consistent with γ carboxylation.

A well known characteristic of Gla containing proteins is their coprecipitation with insoluble barium salts (Dahlbeck, Biochem. J. **209**:837 [1983]; Discipio and Davie, Biochemistry **18**:899 [1979]). An assay based on this property allowed us to analyze the
 15 binding of purified gas6 to ^{125}I -Rse-IgG in the absence of cell membranes. Samples containing various dilutions of Rse-L in 25 mM HEPES pH 7.2, 0.1% BSA and 0.05% Tween-20 were combined and mixed with ^{125}I -Rse-IgG diluted in the same buffer in a total volume of 100-120 ml. After a 45 min incubation at room temperature, 1 ml of a freshly prepared ice cold suspension of BaCl_2 (10 mM) in phosphate buffered saline was added to
 20 each tube and precipitable radioactivity was collected by centrifugation and aspiration of the supernatant fluid. The dissociation constant for Rse-IgG and gas6 measured in this assay was 0.46 nM (Fig. 5).

Purified gas6 stimulated phosphorylation of Rse in a dose dependent fashion. A time course experiment showed that phosphorylation of Rse was induced within two minutes after
 25 addition of purified gas6. Activation of Rse phosphorylation by gas6 was neutralized by Rse-IgG but not by CD4-IgG.

Example 7

Gas6 Expression and Characterization

Gas6 and Rse receptor expression in adult human brain tissues was investigated. A
 30 blot containing 2 μg of polyadenylated RNA from human brain tissues (Clontech) was hybridized with random-primed labeled probes corresponding to amino acids 1- 420 of Rse or to amino acids 358-605 of gas6. The tissues were amygdala, caudate nucleus, corpus callosum, hippocampus, hypothalamus, substantia nigra, subthalamic nucleus, and thalamus.

Consistent with the hypothesis that gas6 might be a ligand for Rse, it was found that gas6 and Rse mRNA are co-expressed in each of these adult human brain tissues.

Astrocytes have been reported to synthesize neurotrophic factors that support the growth and survival of neurons. Moretto *et al.*, *J. Neuropath & Exp Neuro.* **53**:78 (1994) and Lin *et al.* *Science* **260**:1130 (1993). It was determined whether cultured rat astrocytes also synthesize a ligand for Rse. A northern blot was prepared which contained 1 µg pf polyadenylated RNA from postnatal day 1 astrocytes or hippocampal neurons prepared from E18 rat embryos. Astrocytes were prepared as described (Banker and Goslin, *Culturing Nerve Cells* [MIT Press, Cambridge, 1991], pp 260-261) and then cultured in serum-free media for 1 day, 3 days, or 5 days. Hippocampal neurons were cultured in serum free defined media for 0 days, 3 days or 4 days. The blot was hybridized with a ³²P-labeled probe corresponding to amino acids 1-460 of murine gas6. The blot was stripped then hybridized with a ³²P-labeled actin probe to confirm the integrity of the RNA samples.

Gas6 mRNA was detected in cultured type 1 astrocytes prepared from postnatal day 1 rats, but could not be detected in E18 hippocampal neurons.

Expression data for gas6 and Rse obtained herein and elsewhere is summarized in the following table.

TABLE 3
Expression of Gas6 and Rse in Primary Cell Culture and Cell Lines

Cell Culture	Gas6	Rse
Neuronal System		
Astrocyte	+	+
Hippocampus neurons	-	Schulz <i>et al.</i> , <i>supra</i>
P45 Schwann's	-	+
Neuroglioma		
G28	+	-
G59	+	-
G111	+	-
Glioblastoma, astrocytorum		

P0929P2

5

U87MG	-	-
U373MG	+	-
Hematopoietic System		
JM, a CML line	n/a	+
CMK11-5	n/a	+
Jurkat	n/a	+
NIH3T3	+	+
293	+	-

10 The ability of cultured rat astrocytes to also synthesize a ligand for the Rse receptor was investigated. See figure legends for Figs. 6A-6C. Astrocyte conditioned media contained a factor which bound ^{125}I -Rse-IgG (Fig. 6A) and stimulated tyrosine phosphorylation of Rse (Fig. 6B). This activity was neutralized by Rse-IgG but not CD4-IgG (Fig. 6C).

Example 8

Gas6 Variants

15 To further characterize the interactions of gas6 with cell membranes and with Rse, a series of N-terminal deletion variants containing an epitope tag were constructed.

The coding sequences for the gD signal sequence and epitope tag (Mark *et al.*, [1992] *supra*) were fused via an XhoI site that was added by PCR to coding sequences immediately before the first amino acid of mature gas6 (gD.gas6; forward primer 5'-
 20 AGCTGCTCGAGGCGCTGTTGCCGGCGC [SEQ ID NO: 15]) or protein S (gD.protein S; forward primer 5'-AGCTGCTCGAGGCAAATTCTTTACTTGAA [SEQ ID NO: 16], or amino acids 118 (gD.gas6.118-C; forward primer 5'-AGCTGCTCGAGGACCAGTGCACGCCCAACC [SEQ ID NO: 17]) and 279 (gD.gas6.279-C; forward primers 5'- GCTGCTCGAGGACATCTTGCCGTGCGTG [SEQ ID NO: 18]) of
 25 gas6. The reverse primer for gD.gas6 and gD.gas6.118-C was 5'-CATGGATCCTACCGGAAGTCAAACCTCAGCTA [SEQ ID NO: 11]. The reverse primers for gD.gas6.279-C and gD.protein S were 5'-GTCGGATCCGACAGAGACTGAGAAGCC [SEQ ID NO: 13] and 5'-CATTCAATTATGTCAAATTCA [SEQ ID NO: 19], respectively. Gas6.gD was constructed by fusing the coding sequences of gas6 to the C-terminal gD tag
 30 used for Rse.gD through an NheI site which was added by PCR using the primers 5'-

P0929P2

ATGGAGATCAAGGTCTG [SEQ ID NO: 20] and 5'-CATCTTGAGGCTAGCGGCTGCGGCGGGCTCCAC [SEQ ID NO: 21]. The polypeptides were expressed in 293 cells using the procedure essentially as described for full length gas6 in Example 6.

- 5 gD.gas6.118-C and gD.gas6.279-C, containing the EGF repeats and tandem G domains within the D domain, or just the G domains, respectively, were precipitated by Rse-IgG (Fig. 7) from cell culture supernatants. Human protein S was not precipitated in this assay which is consistent with the above observations that human protein S binds Rse with a lower affinity than human gas6. These derivatives of gas6 that were truncated for the Gla domain
10 (*i.e.* the A domain) also fail to associate with membranes in a Ca^{2+} fashion.

This data shows that gas6 binds to Rse through the G domains, that the membrane binding and Rse-binding activities are separable, and suggests that the Gla domain is required for Ca^{2+} dependent association with cell membranes.

- The gas6 variants described in this example were functionally active. In particular,
15 gD.gas6.118-C and gD.gas6.279-C activated Rse phosphorylation in the KIRA assay described in Example 4 as effectively as full-length gD-tagged gas6 (see Fig. 7).

Example 9

Schwann Cell Proliferation Assay

- Rse mRNA, but not gas6 mRNA was also detected in the rat Schwann cell line rhESC,
20 which is derived from rat E14 dorsal root ganglia. Addition of purified gas6 to these cells resulted in a dose dependent increase in cell number (50% increase at 48 hours) with an EC_{50} of ~ 0.3 nM (Fig. 8). Gas6 treatment also altered the morphology of these cells; untreated cells were multipolar with numerous branched processes whereas gas6 treated cells became spindle-shaped with two major smooth processes and aligned themselves in a parallel array.
25 It was also demonstrated that gas6-induced proliferation was neutralized by Rse-IgG but not CD4-IgG. See Fig. 9.

Using Rse- and Axl- specific antibodies, Rse and Axl receptor tyrosine kinases were also detected in human Schwann cells. The ability of gas6 to enhance the proliferation of human Schwann cells was determined.

- 30 Peripheral nerve tissues were obtained at the University of Miami School of Medicine, with appropriate patient consent, as previously described (Levi *et al.*, J. Neuroscience 15(2):1329-1340 [1995]). Pieces of peripheral nerve fibers were placed in Belzer's UW solution and shipped to California. Upon receipt, the nerve fibers were washed with fresh F12/DME (1:1) and incubated with 1% collagenase/dispase solution (Boehringer) at 37°C for

P0929P2

30 minutes. Then, the tissue was gently washed 3x by transferring the tissue to fresh tissue culture medium. The fibers were plated in 100 mm petri dishes in serum free medium supplemented with the following formula for rat Schwann cells: F12/DME (1:1) supplemented with insulin (10 μ g/ml), transferrin (10 μ g/ml), α -tocopherol (5 μ g/ml),
 5 recombinant human heregulin- β 1₁₇₇₋₂₄₄ produced as described in Holmes *et al.*, Science, **256**: 1205-1210 (1992) (10 nmole/L), forskolin (5 μ molar), progesterone (3×10^{-8} molar), and bovine pituitary extract (BPE) (3 μ l/ml). The Schwann cells were cultured in suspension for 48 hours to allow partial demyelination. The nerve fibers were then pooled by centrifugation at 1000rpm for 5 minutes and resuspended and dispersed by gentle pipetting.

10 The dispersed Schwann cells were replated on laminin (Gibco BRL) coated tissue culture 48 well multiplates at 8×10^3 cells/well in defined medium with the addition of aprotinin (25 μ g/ml) and 50 μ L/ml chemically defined lipids (Sigma Cat# 11905-015; Gibco BRL). These cultures were designated the "primary culture". Medium was changed every 5 days. Confluent cultures of pure Schwann cells could be obtained within 2 weeks. At the first
 15 and second passage, cells were removed from the plate using collagenase/dispase (Boehringer Mannheim), washed with medium containing 3% BSA, and plated as described. The media used were "6F" medium: F12/DME (1:1) supplemented with insulin (10 μ g/ml), transferrin (10 μ g/ml), α -tocopherol (5 μ g/ml), progesterone (3×10^{-8} molar), aprotinin (25 μ g/ml) and chemically defined lipids (Sigma Cat# 11905-015). "8F" medium contains the supplements
 20 of 6F medium as well as recombinant human heregulin- β 1₁₇₇₋₂₄₄ (10nmole/L) and forskolin (5 μ M). The effect of gas6 on Schwann cell survival and proliferation was studied by adding gas6 to either or these culture mediums.

Gas6 stimulates human Schwann cell growth in a dose dependent manner (Fig. 12A) with a significant effect seen at 1 ng/ml (14pM) and maximal effect with doses over 10 ng/ml.
 25 Gas6 alone produces a significant increase in Schwann cell number compared to control medium. In the presence of the cAMP activator, forskolin, the increase in total cell number with gas6 is more pronounced. A synergistic effect is also observed between gas6 and heregulin. Gas6 increased both cell number and thymidine incorporation even in the presence of preferred concentrations of both forskolin and heregulin (Fig. 12B).

30 In the presence of the preferred concentrations of both heregulin and forskolin, other growth factors previously reported to stimulate Schwann cell growth had no effect (PDGF, FGF- β) or reduce cell number (IL-1 α and TGF- β 1) (Fig. 12C). Addition of human or bovine protein S at 10ng-5 μ g did not increase Schwann cell number after 5 days of culture. In contrast, gas6 at 30 μ g/ml maximally increased the cell number. The combination of gas6

P0929P2

with forskolin and heregulin results in maximal cell growth over a 5 day period comparable to that seen in the combination of 6F + forskolin + heregulin + 5% FBS.

Gas6 has a marked effect on cell morphology as determined by viewing phase contrast micrographs of human Schwann cells grown in 6F+heregulin; 6F+heregulin+gas6; 8F+gas6; and 8F+10% fetal bovine serum. Micrographs were taken after 96 hours of culture. The Schwann cell grown in the presence of gas6 have processes which are much longer than those seen in cells grown in the presence of heregulin or heregulin with forskolin. Mitotic figures may also clearly be seen in the 8F + gas6 cultures even in those cells with the fully developed Schwann cell spindle-shaped morphology. The addition of serum to the 8F cultures alters cell morphology causing the cells to flatten, spread and eventually become vacuolated.

Cells were stained by immunofluorescence for the Schwann cell markers GFAP and S100 protein. Briefly, passage 4 human Schwann cells grown in 8F+gas6 were cultured for 24 hours on laminin coated Chamber slides and fixed in 10% formalin in PBS. Fixed cells were blocked with 10% goat serum and incubated with rabbit anti-GFAP (ICN) and anti-S-100 protein (ICN) at dilutions recommended by the distributor. Specific binding of the primary antibody was stained with goat anti-rabbit IgG (Fab')₂-FITC conjugates. Cells were counter-stained with DNA dye propidium iodide. Negative controls were run on WI-38 cells which stained negative. Cells grown showed 100% immunofluorescent staining for the Schwann cell markers GFAP and S100 protein after 4 subcultures.

The ability of gas6 to stimulate human Schwann cell proliferation through the Axl-Rse family of tyrosine kinase receptors was investigated. Human Schwann cells were stimulated with 0, 0.01, 0.1 or 1 µg/ml of human gas6 (hgas6) for 15 min at 37°C incubator. Cell lysates were prepared and immunoprecipitated with rabbit anti-hRseFc fusion protein antibody and rabbit anti-hAxl antibody. Tyrosine phosphorylation of hRse and hAxl receptor was detected with 4G10 anti-phosphorylation antibody. 10⁶ human Schwann cells were grown to near confluence in defined media (8F+gas6) and changed to 6F 24 hours before experiment. Cells were treated with purified recombinant hgas6 for 15 min in 37°C incubator and lysed on ice with 1 ml of lysis buffer (20 mM HEPES, pH7.4, 135 mM NaCl, 50 mM NaF, 1 mM sodium vanadate and 1 mM sodium molybdate, 2 mM EDTA and 2 mM EGTA, 10% glycerol, 1%NP40, 1 µM okadaic acid, 1 mM PMSF and 1 mM AEBSF). Cell lysates were clarified by centrifuging at 14000xg 4°C for 10 min. Immunoprecipitations were performed using 1 µg of rabbit anti-hRseFc fusion protein antibody or 2 µl of rabbit anti-hAxl antiserum raised against the 10 amino acids at the COOH-terminal of hAxl at 4°C for 2 hrs. Immunocomplex were collected with 10 µl of Protein A Sepharose CL-4B beads. Proteins were separated on Novex* 4-12% gradient gel and transferred onto nitrocellulose membrane. Anti-

P0929P2

phosphotyrosine immunoblots were performed using 4G10 mouse anti-phosphotyrosine antibody (UBI), goat anti-mouse horseradish peroxidase conjugate and ECL developing kit (Amersham). Addition of human gas6 to human Schwann cells causes autophosphorylation of both Axl and Rse receptors on tyrosine residue(s). Activation of Axl and Rse could be detected at 1.4-14 μ M gas6. Such phosphorylation of Axl and Rse is not observed in cultures stimulated with heregulin. Gas6 expression in cultured rat Schwann cells was not detected by northern blot. Furthermore, gas6 activity in rat Schwann cell conditioned medium was not seen. Without being bound by any one theory, it is possible that gas6 is produced by growing axons, or by nearby fibroblast cells (from which gas6 was initially cloned). This activation of Axl-Rse receptors on Schwann cells by gas6 is highly specific, since growth factors known to act via other tyrosine kinase receptors, such as PDGF and FGF, do not increase human Schwann cell proliferation under these defined conditions. The Schwann cell growth factors, GGF/heregulin, acting independently through the *erbB* receptor family, synergize with gas6 in this study.

It is beneficial to have populations of mammalian Schwann cells (preferably human Schwann cells) for use as cellular prostheses for transplantation into areas of damaged spinal cord in an effort to influence regeneration of interrupted central axons, for assisting in the repair of peripheral nerve injuries and as alternatives to multiple autografts. See Levi *et al.*, J. Neuroscience 14(3):1309-1319 (1994). The use of cell culture techniques to obtain an abundant source of autologous graft material from a small biopsy has already met with clinical success in providing human epidermal cells to cover extensive burns (Gallico *et al.*, N. Eng J. Med., 311:338-451 [1984]). Furthermore, it has been shown that Schwann cells from human xenografts are capable of myelinating regenerating peripheral axons from mice which have been immunosuppressed (Aguayo *et al.*, Nature 268:753-755 [1977], and Aguayo *et al.*, Soc. Neurosci. Symp. 4:361-383 [1979]). Accordingly, it is expected that the above approach will meet with success in mammals, including humans.

Example 10

Gas6 Immunoadhesin

gD.gas6.279-C.IgG was constructed by fusing the coding sequences of gD.gas6.279-C (see Example 8) to amino acids 216-443 of human IgG γ 1 through a BstEII linker (adding amino acids Val and Thr). The linker was added to gD.gas6.279-C sequences by PCR using the primers 5'-ATGGAGATCAAGGTCTG [SEQ ID NO: 20] and 5'-GTCGGTGACCGCTGCTGCGGGCTCCAC [SEQ ID NO: 22].

P0929P2

The gas6 immunoadhesin thus formed was subjected to the KIRA assay described in Example 4 above. Briefly, different dilutions of conditioned media from cells transiently expressing gD.gas6.279-C.IgG were tested in the KIRA assay. The starting material had a concentration of gD.gas6.279-C.IgG of ~230 ng/ml. The EC₅₀ for activation was approximately 0.4 nM. See Fig. 10. Activity was not observed in conditioned media from transiently transfected control cell lines.

Example 11

Activation of Rse by Non- γ Carboxylated Gas6

Media (700 ml) conditioned for 3 days by 293 cells transfected with human gas6 (hgas6.17) was dialyzed against 2 x 8 L of 50 mM Tris-HCl pH 7.5, 5 mM benzimidine (buffer A). The dialyzate was adjusted to 0.1 % CHAPS, and loaded on a 6 ml Resource-Q column (Pharmacia) at 10 ml/minute. The column was washed with buffer A, and eluted with a 70 ml linear gradient of 0 to 0.4 M NaCl in buffer A, at a flow rate of 1.0 ml/min, collecting fractions of 2.0 ml.

The fractions were analyzed for their ability to bind and activate Rse using the barium chloride binding method described in Example 6 and the KIRA assay described in Example 4. The barium chloride assay can only detect binding of Gla containing Rse ligands, while the KIRA assay is sensitive to all Rse activators. Binding activity was centered at fraction 31, while KIRA showed an additional earlier eluting peak centered at fraction 24.

Aliquots (10 μ l) of fractions 20 to 44 were analyzed on 8% acrylamide (Novex) SDS gels, and proteins visualized by silver staining. In these fractions a 75 kD band comigrated with standard hgas6. Integrated intensities of the 75 kD band were measured by laser scanning (Molecular Dynamics) and image analysis (NIH Image). Peak intensities were found in fractions 24 and 31, corresponded to the 2 regions of KIRA activity. The amount of hgas6 in each fraction was estimated from the staining intensity of a known quantity (0.34 μ g) of a standard preparation of hgas6 run on the same gel, assuming a linear relationship between staining intensity and protein load.

Sequence analysis of the 75 kD bands from fractions 24 and 31 was performed after SDS-PAGE and electrophoretic transfer to PVDF membranes. The amino terminal sequence of both bands was unambiguously identified as that of hgas6 (AFQVF), but the two could be differentiated by the presence or absence of modified glutamic acid residues in later cycles. The sequence from fraction 31 lacked a signal from glutamic acid in cycles 6,7,14, and 16, consistent with a γ -carboxyl modification of these residues. The sequence from fraction 24 was consistent with unmodified glutamic acid at all these positions.

P0929P2

Both sequence analysis and binding behavior of the early eluting form of hgas6 are consistent with its identification as an unmodified form of hgas6, lacking the characteristic γ -carboxyl modification of glutamic acid. This second discovered form of recombinant hgas6 appears to be more active than the first described Gla containing form. The specific activity of the two forms was calculated from the KIRA data in Fig. 11 and from the densitometric quantitation of hgas6. The specific activity of fraction 31 (form 1) is 1170 KIRA units/mg P, while that of fraction 24 (form 2) is 3158. This indicates that form 2, lacking the Gla modification, is more potent than the Gla containing form 1 in activating Rse.

Example 12

10 Activation of Mer Receptor by gas6

A cDNA encoding human Mer was obtained by screening a human testis cDNA library

prepared in lambda DR2 (Clontech, Palo Alto CA) with oligonucleotide probes:

5'-GAAATTACAGATCCGCAGCCCCGGGATGGGGCCGGCCCCGCTGCCGCTGC

15 [SEQ ID NO: 23],

5'-CCTTGGATTCTAGCAAGCACGACTGAAGGAGCCCCATCAGTAGCACCTTT [SEQ ID NO: 24], and

5'-TCTTAAAATTAAGCTTCAGCTGCTCCTTGATATTAACCTTTGTACAGAGT-3' [SEQ ID NO: 25].

20 Positive phage were purified and insert sizes were determined. Three overlapping clones hMer.cl4, hMer.cl25, and hMer.cl6 (corresponding to nt 1-561, nt 223-2025, and nt 1902-3608 of published sequence respectively) were combined to make a cDNA encoding the entire open reading frame of human Mer.

25 Mer-Fc was constructed by fusing the sequence encoding amino acids 1-499 of human Mer to amino acids 216-443 of human IgG γ 1 through a NarI-BstEII linker (5'-GCGCCTGGCAACGCG-3' [SEQ ID NO: 26], 5'-GTGACCGCGTTGCCAG-3' [SEQ ID NO: 27]) (adding amino acids glycine and histidine). Human embryonic kidney (HEK) 293 cells expressing Mer-Fc were screened using a human Fc-specific ELISA. Mer-Fc was purified on a protein A-sepharose column (Pharmacia).

30 Epitope tagged gD.Mer was constructed by fusing the coding sequences for amino acids 1-53 of herpes simplex virus type 1 (HSV I) glycoprotein D (gD) to the sequences encoding amino acids 36-961 of human Mer. See Example 3 above. Oligos (5'-CGAATTCCTCGAGCCGGGACCTTTTCCAGGGAGC-3' [SEQ ID NO: 28], and

P0929P2

5'-CCAACTGTGTGTTTGAAGGCAAGAGGCGG-3' [SEQ ID NO: 29]) were used to add a XhoI site to the human Mer cDNA by PCR. The gD.Mer cDNA was inserted into a CMV-based expression vector and HEK 293 cells were transfected, selected and screened by western blotting and fluorescent activated cell sorting as described above in Example 3.

5 Recombinant human gas6 and Protein S were expressed as described in Godowski *et al.* Cell 82:355-358 (1995). Expression was verified by metabolic labeling of cultures with ³⁵S-Cys and ³⁵S-Met (Mark *et al.*, J. Biol. Chem. 267:26166 [1992]) and/or by western blotting with rabbit polyclonal anti-gas6 or anti-Protein S antiserum (Sigma). N-terminal tagged versions of Protein S or gas6 were constructed as described in Godowski *et al.*, (1995)
10 *supra*.

Methods to detect Rse phosphorylation using the KIRA ELISA and Western analysis using anti-phosphotyrosine antibodies have been described above in Examples 3 and 4. For neutralization experiments, potential ligand sources were treated at room temperature for 30 min with the indicated Fc fusion protein prior to addition to cells. To analyze the ability of
15 potential ligands to induce phosphorylation of gD.Mer, 500,000 cells were seeded on a 60 mm dish in the presence of serum for 6 h. The cells were then washed twice in PBS and serum-starved for 16 h. Potential ligands were added to the cells and gD.Mer was immunoprecipitated from lysates using 5B6, analyzed by SDS-PAGE under reducing conditions, and blotted with anti-phosphotyrosine antibody (4G10).

20 For binding to Rse-Fc, Axl-Fc, and Mer-Fc, conditioned media containing 5-10 nM gas6 or protein S was precleared with protein A-Sepharose (Calbiochem) for 30 min at room temperature, then incubated with 5 µg of the receptor Fc fusion protein for 4 h at 4°C. Fusion proteins were immunoprecipitated with 20 µl of protein A sepharose and the complexes were collected by centrifugation at 14,000 x g for 1 min, and then washed 3 times with PBS
25 containing 0.1% Triton X-100. Precipitates were analyzed by SDS-PAGE under reducing conditions. Western blots of the SDS-PAGE gels were probed with antibody 5B6.

Protein interaction analysis using BIAcore™ instruments were performed with Mer-Fc coupled to BIAcore CM5 sensor chips using purified gas6 and Protein S. For neutralization experiments, 5 µgs of either Mer-Fc or CD4-Fc was mixed with gas6 for thirty minutes at
30 room temperature prior to injection over the chip. Sensorgrams were analyzed with BIA evaluation 2.1 software from Pharmacia Biosensor AB. Apparent dissociation rate constants (kd) and association rate constants (ka) were obtained by evaluating the sensorgram with A+B=AB type I fitting. Equilibrium dissociation constant K_d was calculated as kd/ka.

Results

P0929P2

The ability of Mer-Fc to neutralize gas6-induced phosphorylation of Rse in the ELISA based KIRA assay was investigated. Activation of Rse by gas6 was blocked by Rse-Fc and Mer-Fc in a dose-responsive fashion (Fig. 14). Neutralization was specific to Rse-Fc and Mer-Fc, because the control protein CD4-Fc did not inhibit in this assay. Rse-Fc was somewhat
5 more potent than Mer-Fc in neutralizing gas6. These data suggest that Mer-Fc, like Rse-Fc, blocks activation of Rse by binding to gas6.

The ability of Axl-Fc, Rse-Fc and Mer-Fc to bind directly to gas6 was determined using a coprecipitation assay. The ability of these receptors to bind Protein S was also determined. To allow a more quantitative comparison of their binding properties, versions of
10 gas6 and Protein S that contained epitope tags were utilized. Conditioned media from cells expressing tagged gas6 or Protein S was incubated with each of the receptor fusion proteins. The Fc fusion proteins, and proteins bound to them, were recovered with protein A and washed extensively. Tagged proteins that bound to the Fc-fusion proteins were revealed by Western blotting and detection with an antibody directed against the epitope tag. Both Axl-Fc
15 and Rse-Fc bound tagged gas6 but not Protein S. An identical result was observed with Mer-Fc. The binding was specific in that neither gas6 nor Protein S bound the control CD4-Fc.

The kinetics of the interaction of gas6 with the extracellular domains of Mer and Rse was compared using a BIAcore instrument. Mer-Fc was immobilized on a biosensor chip, and various concentrations of gas6 were passed over the chip. A representative sensorgram from
20 such an experiment is shown in Fig. 15A. No binding was observed to Protein S and binding of gas6 to Mer-Fc immobilized on the chip was blocked by soluble Mer-Fc, but not CD4-Fc (Fig. 15). The dissociation constant (K_d) for the interaction of gas6 with Mer-Fc was 6 nM. The K_d for binding of gas6 to Rse-Fc was 4.2 nM.

HEK 293 cells were transfected with an expression vector encoding a version of Mer
25 containing an amino-terminal epitope tag. A candidate cell line (termed 293.gD.Mer) expressing gD.Mer was identified by fluorescence-activated cell sorting using 5B6, a monoclonal antibody that recognizes the epitope tag. Incubation of these cells with antibody 5B6 resulted in rapid phosphorylation of novel proteins of approximately 180 and 200 kDa that were absent in control, untransfected HEK 293 cells. Receptor phosphorylation was not
30 observed when the cells were incubated with a control antibody. These results indicate that Mer encodes a functional tyrosine kinase.

The ability of human gas6 and Protein S to activate the kinase activity of Mer was investigated. Mer did not become phosphorylated when these cells were treated with Protein S. However, receptor phosphorylation was detected in cells treated with gas6. A time course
35 experiment demonstrated that Mer was phosphorylated within minutes of addition of gas6 and

P0929P2

phosphorylation could be detected after stimulation of cells with gas6 at concentrations of 1-3nM.

5 The ability of gas6 to induce the phosphorylation of Mer expressed endogenously was determined. Mer mRNA was detected in the monocytic leukemia cell line THP-1 by reverse transcription PCR. Using a polyclonal antibody directed against the extracellular domain of Mer, expression of Mer on the surface of THP-1 cells was confirmed. Treatment of these cells with gas6 induced the rapid phosphorylation of a 180 kDa protein that was immunoprecipitated with antibodies to Mer. These observations show that gas6 is a functional ligand for Mer.

P0929P2

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Genentech, Inc.
Chen, Jian
5 Godowski, Paul J.
Hammonds, R. Glenn
Mark, Melanie
Mather, Jennie P.
Li, Ronghao
- 10 (ii) TITLE OF INVENTION: RECEPTOR ACTIVATION BY GAS6
- (iii) NUMBER OF SEQUENCES: 29
- (iv) CORRESPONDENCE ADDRESS:
15 (A) ADDRESSEE: Genentech, Inc.
(B) STREET: 460 Point San Bruno Blvd
(C) CITY: South San Francisco
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 94080
- 20 (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: 3.5 inch, 1.44 Mb floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: WinPatin (Genentech)
- 25 (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
- 30 (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/402253
(B) FILING DATE: 3/10/95
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/438861
(B) FILING DATE: 5/10/95
- 35 (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Lee, Wendy M.
(B) REGISTRATION NUMBER: 00,000
(C) REFERENCE/DOCKET NUMBER: P0929P2PCT
- 40 (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 415/225-1994
(B) TELEFAX: 415/952-9881

P0929P2

(C) TELEX: 910/371-7168

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 673 amino acids
 (B) TYPE: Amino Acid
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	Met	Pro	Pro	Pro	Pro	Gly	Pro	Ala	Ala	Ala	Leu	Gly	Thr	Ala	Leu	
	1				5					10					15	
10	Leu	Leu	Leu	Leu	Leu	Ala	Ser	Glu	Ser	Ser	His	Thr	Val	Leu	Leu	
					20					25					30	
	Arg	Ala	Arg	Glu	Ala	Ala	Gln	Phe	Leu	Arg	Pro	Arg	Gln	Arg	Arg	
					35					40					45	
15	Ala	Tyr	Gln	Val	Phe	Glu	Glu	Ala	Lys	Gln	Gly	His	Leu	Glu	Arg	
					50					55					60	
	Glu	Cys	Val	Glu	Glu	Val	Cys	Ser	Lys	Glu	Glu	Ala	Arg	Glu	Val	
					65					70					75	
	Phe	Glu	Asn	Asp	Pro	Glu	Thr	Glu	Tyr	Phe	Tyr	Pro	Arg	Tyr	Gln	
					80					85					90	
20	Glu	Cys	Met	Arg	Lys	Tyr	Gly	Arg	Pro	Glu	Glu	Lys	Asn	Pro	Asp	
					95					100					105	
	Phe	Ala	Lys	Cys	Val	Gln	Asn	Leu	Pro	Asp	Gln	Cys	Thr	Pro	Asn	
					110					115					120	
25	Pro	Cys	Asp	Lys	Lys	Gly	Thr	His	Ile	Cys	Gln	Asp	Leu	Met	Gly	
					125					130					135	
	Asn	Phe	Phe	Cys	Val	Cys	Thr	Asp	Gly	Trp	Gly	Gly	Arg	Leu	Cys	
					140					145					150	
	Asp	Lys	Asp	Val	Asn	Glu	Cys	Val	Gln	Lys	Asn	Gly	Gly	Cys	Ser	
					155					160					165	
30	Gln	Val	Cys	His	Asn	Lys	Pro	Gly	Ser	Phe	Gln	Cys	Ala	Cys	His	
					170					175					180	
	Ser	Gly	Phe	Ser	Leu	Ala	Ser	Asp	Gly	Gln	Thr	Cys	Gln	Asp	Ile	
					185					190					195	
35	Asp	Glu	Cys	Thr	Asp	Ser	Asp	Thr	Cys	Gly	Asp	Ala	Arg	Cys	Lys	
					200					205					210	

P0929P2

	Asn	Leu	Pro	Gly	Ser	Tyr	Ser	Cys	Leu	Cys	Asp	Glu	Gly	Tyr	Thr
					215					220					225
	Tyr	Ser	Ser	Lys	Glu	Lys	Thr	Cys	Gln	Asp	Val	Asp	Glu	Cys	Gln
					230					235					240
5	Gln	Asp	Arg	Cys	Glu	Gln	Thr	Cys	Val	Asn	Ser	Pro	Gly	Ser	Tyr
					245					250					255
	Thr	Cys	His	Cys	Asp	Gly	Arg	Gly	Gly	Leu	Lys	Leu	Ser	Pro	Asp
					260					265					270
10	Met	Asp	Thr	Cys	Glu	Asp	Ile	Leu	Pro	Cys	Val	Pro	Phe	Ser	Met
					275					280					285
	Ala	Lys	Ser	Val	Lys	Ser	Leu	Tyr	Leu	Gly	Arg	Met	Phe	Ser	Gly
					290					295					300
	Thr	Pro	Val	Ile	Arg	Leu	Arg	Phe	Lys	Arg	Leu	Gln	Pro	Thr	Arg
					305					310					315
15	Leu	Leu	Ala	Glu	Phe	Asp	Phe	Arg	Thr	Phe	Asp	Pro	Glu	Gly	Val
					320					325					330
	Leu	Phe	Phe	Ala	Gly	Gly	Arg	Ser	Asp	Ser	Thr	Trp	Ile	Val	Leu
					335					340					345
20	Gly	Leu	Arg	Ala	Gly	Arg	Leu	Glu	Leu	Gln	Leu	Arg	Tyr	Asn	Gly
					350					355					360
	Val	Gly	Arg	Ile	Thr	Ser	Ser	Gly	Pro	Thr	Ile	Asn	His	Gly	Met
					365					370					375
	Trp	Gln	Thr	Ile	Ser	Val	Glu	Glu	Leu	Glu	Arg	Asn	Leu	Val	Ile
					380					385					390
25	Lys	Val	Asn	Lys	Asp	Ala	Val	Met	Lys	Ile	Ala	Val	Ala	Gly	Glu
					395					400					405
	Leu	Phe	Gln	Leu	Glu	Arg	Gly	Leu	Tyr	His	Leu	Asn	Leu	Thr	Val
					410					415					420
30	Gly	Gly	Ile	Pro	Phe	Lys	Glu	Ser	Glu	Leu	Val	Gln	Pro	Ile	Asn
					425					430					435
	Pro	Arg	Leu	Asp	Gly	Cys	Met	Arg	Ser	Trp	Asn	Trp	Leu	Asn	Gly
					440					445					450
	Glu	Asp	Ser	Ala	Ile	Gln	Glu	Thr	Val	Lys	Ala	Asn	Thr	Lys	Met
					455					460					465
35	Gln	Cys	Phe	Ser	Val	Thr	Glu	Arg	Gly	Ser	Phe	Phe	Pro	Gly	Asn

P0929P2

				470						475				480				
				Gly	Phe	Ala	Thr	Tyr	Arg	Leu	Asn	Tyr	Thr	Arg	Thr	Ser	Leu	Asp
								485						490				495
5				Val	Gly	Thr	Glu	Thr	Thr	Trp	Glu	Val	Lys	Val	Val	Ala	Arg	Ile
								500					505					510
				Arg	Pro	Ala	Thr	Asp	Thr	Gly	Val	Leu	Leu	Ala	Leu	Val	Gly	Asp
								515					520					525
				Asp	Asp	Val	Val	Ile	Ser	Val	Ala	Leu	Val	Asp	Tyr	His	Ser	Thr
								530					535					540
10				Lys	Lys	Leu	Lys	Lys	Gln	Leu	Val	Val	Leu	Ala	Val	Glu	Asp	Val
								545					550					555
				Ala	Leu	Ala	Leu	Met	Glu	Ile	Lys	Val	Cys	Asp	Ser	Gln	Glu	His
								560					565					570
15				Thr	Val	Thr	Val	Ser	Leu	Arg	Glu	Gly	Glu	Ala	Thr	Leu	Glu	Val
								575					580					585
				Asp	Gly	Thr	Lys	Gly	Gln	Ser	Glu	Val	Ser	Thr	Ala	Gln	Leu	Gln
								590					595					600
				Glu	Arg	Leu	Asp	Thr	Leu	Lys	Thr	His	Leu	Gln	Gly	Ser	Val	His
								605					610					615
20				Thr	Tyr	Val	Gly	Gly	Leu	Pro	Glu	Val	Ser	Val	Ile	Ser	Ala	Pro
								620					625					630
				Val	Thr	Ala	Phe	Tyr	Arg	Gly	Cys	Met	Thr	Leu	Glu	Val	Asn	Gly
								635					640					645
25				Lys	Ile	Leu	Asp	Leu	Asp	Thr	Ala	Ser	Tyr	Lys	His	Ser	Asp	Ile
								650					655					660
				Thr	Ser	His	Ser	Cys	Pro	Pro	Val	Glu	His	Ala	Thr	Pro		
								665					670			673		

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- 30 (A) LENGTH: 678 amino acids
- (B) TYPE: Amino Acid
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

35	Met	Ala	Pro	Ser	Leu	Ser	Pro	Gly	Pro	Ala	Ala	Leu	Arg	Arg	Ala
	1				5					10					15

P0929P2

	Pro	Gln	Leu	Leu	Leu	Leu	Leu	Leu	Ala	Ala	Glu	Cys	Ala	Leu	Ala	
					20					25					30	
	Ala	Leu	Leu	Pro	Ala	Arg	Glu	Ala	Thr	Gln	Phe	Leu	Arg	Pro	Arg	
					35					40					45	
5	Gln	Arg	Arg	Ala	Phe	Gln	Val	Phe	Glu	Glu	Ala	Lys	Gln	Gly	His	
					50					55					60	
	Leu	Glu	Arg	Glu	Cys	Val	Glu	Glu	Leu	Cys	Ser	Arg	Glu	Glu	Ala	
					65					70					75	
10	Arg	Glu	Val	Phe	Glu	Asn	Asp	Pro	Glu	Thr	Asp	Tyr	Phe	Tyr	Pro	
					80					85					90	
	Arg	Tyr	Leu	Asp	Cys	Ile	Asn	Lys	Tyr	Gly	Ser	Pro	Tyr	Thr	Lys	
					95					100					105	
	Asn	Ser	Gly	Phe	Ala	Thr	Cys	Val	Gln	Asn	Leu	Pro	Asp	Gln	Cys	
					110					115					120	
15	Thr	Pro	Asn	Pro	Cys	Asp	Arg	Lys	Gly	Thr	Gln	Ala	Cys	Gln	Asp	
					125					130					135	
	Leu	Met	Gly	Asn	Phe	Phe	Cys	Leu	Cys	Lys	Ala	Gly	Trp	Gly	Gly	
					140					145					150	
20	Arg	Leu	Cys	Asp	Lys	Asp	Val	Asn	Glu	Cys	Ser	Gln	Glu	Asn	Gly	
					155					160					165	
	Gly	Cys	Leu	Gln	Ile	Cys	His	Asn	Lys	Pro	Gly	Ser	Phe	His	Cys	
					170					175					180	
	Ser	Cys	His	Ser	Gly	Phe	Glu	Leu	Ser	Ser	Asp	Gly	Arg	Thr	Cys	
					185					190					195	
25	Gln	Asp	Ile	Asp	Glu	Cys	Ala	Asp	Ser	Glu	Ala	Cys	Gly	Glu	Ala	
					200					205					210	
	Arg	Cys	Lys	Asn	Leu	Pro	Gly	Ser	Tyr	Ser	Cys	Leu	Cys	Asp	Glu	
					215					220					225	
30	Gly	Phe	Ala	Tyr	Ser	Ser	Gln	Glu	Lys	Ala	Cys	Arg	Asp	Val	Asp	
					230					235					240	
	Glu	Cys	Leu	Gln	Gly	Arg	Cys	Glu	Gln	Val	Cys	Val	Asn	Ser	Pro	
					245					250					255	
	Gly	Ser	Tyr	Thr	Cys	His	Cys	Asp	Gly	Arg	Gly	Gly	Leu	Lys	Leu	
					260					265					270	
35	Ser	Gln	Asp	Met	Asp	Thr	Cys	Glu	Asp	Ile	Leu	Pro	Cys	Val	Pro	

P0929P2

				275					280					285				
				Phe	Ser	Val	Ala	Lys	Ser	Val	Lys	Ser	Leu	Tyr	Leu	Gly	Arg	Met
				290									295					300
5				Phe	Ser	Gly	Thr	Pro	Val	Ile	Arg	Leu	Arg	Phe	Lys	Arg	Leu	Gln
				305									310					315
				Pro	Thr	Arg	Leu	Val	Ala	Glu	Phe	Asp	Phe	Arg	Thr	Phe	Asp	Pro
				320									325					330
				Glu	Gly	Ile	Leu	Leu	Phe	Ala	Gly	Gly	His	Gln	Asp	Ser	Thr	Trp
				335									340					345
10				Ile	Val	Leu	Ala	Leu	Arg	Ala	Gly	Arg	Leu	Glu	Leu	Gln	Leu	Arg
				350									355					360
				Tyr	Asn	Gly	Val	Gly	Arg	Val	Thr	Ser	Ser	Gly	Pro	Val	Ile	Asn
				365									370					375
15				His	Gly	Met	Trp	Gln	Thr	Ile	Ser	Val	Glu	Glu	Leu	Ala	Arg	Asn
				380									385					390
				Leu	Val	Ile	Lys	Val	Asn	Arg	Asp	Ala	Val	Met	Lys	Ile	Ala	Val
				395									400					405
				Ala	Gly	Asp	Leu	Phe	Gln	Pro	Glu	Arg	Gly	Leu	Tyr	His	Leu	Asn
				410									415					420
20				Leu	Thr	Val	Gly	Gly	Ile	Pro	Phe	His	Glu	Lys	Asp	Leu	Val	Gln
				425									430					435
				Pro	Ile	Asn	Pro	Arg	Leu	Asp	Gly	Cys	Met	Arg	Ser	Trp	Asn	Trp
				440									445					450
25				Leu	Asn	Gly	Glu	Asp	Thr	Thr	Ile	Gln	Glu	Thr	Val	Lys	Val	Asn
				455									460					465
				Thr	Arg	Met	Gln	Cys	Phe	Ser	Val	Thr	Glu	Arg	Gly	Ser	Phe	Tyr
				470									475					480
				Pro	Gly	Ser	Gly	Phe	Ala	Phe	Tyr	Ser	Leu	Asp	Tyr	Met	Arg	Thr
				485									490					495
30				Pro	Leu	Asp	Val	Gly	Thr	Glu	Ser	Thr	Trp	Glu	Val	Glu	Val	Val
				500									505					510
				Ala	His	Ile	Arg	Pro	Ala	Ala	Asp	Thr	Gly	Val	Leu	Phe	Ala	Leu
				515									520					525
35				Trp	Ala	Pro	Asp	Leu	Arg	Ala	Val	Pro	Leu	Ser	Val	Ala	Leu	Val
				530									535					540

P0929P2

	Asp	Tyr	His	Ser	Thr	Lys	Lys	Leu	Lys	Lys	Gln	Leu	Val	Val	Leu	
					545					550					555	
	Ala	Val	Glu	His	Thr	Ala	Leu	Ala	Leu	Met	Glu	Ile	Lys	Val	Cys	
					560					565					570	
5	Asp	Gly	Gln	Glu	His	Val	Val	Thr	Val	Ser	Leu	Arg	Asp	Gly	Glu	
					575					580					585	
	Ala	Thr	Leu	Glu	Val	Asp	Gly	Thr	Arg	Gly	Gln	Ser	Glu	Val	Ser	
					590					595					600	
10	Ala	Ala	Gln	Leu	Gln	Glu	Arg	Leu	Ala	Val	Leu	Glu	Arg	His	Leu	
					605					610					615	
	Arg	Ser	Pro	Val	Leu	Thr	Phe	Ala	Gly	Gly	Leu	Pro	Asp	Val	Pro	
					620					625					630	
	Val	Thr	Ser	Ala	Pro	Val	Thr	Ala	Phe	Tyr	Arg	Gly	Cys	Met	Thr	
					635					640					645	
15	Leu	Glu	Val	Asn	Arg	Arg	Leu	Leu	Asp	Leu	Asp	Glu	Ala	Ala	Tyr	
					650					655					660	
	Lys	His	Ser	Asp	Ile	Thr	Ala	His	Ser	Cys	Pro	Pro	Val	Glu	Pro	
					665					670					675	
20	Ala	Ala	Ala													
					678											

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 676 amino acids

(B) TYPE: Amino Acid

(D) TOPOLOGY: Linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

	Met	Arg	Val	Leu	Gly	Gly	Arg	Cys	Gly	Ala	Leu	Leu	Ala	Cys	Leu	
	1				5					10					15	
30	Leu	Leu	Val	Leu	Pro	Val	Ser	Glu	Ala	Asn	Phe	Leu	Ser	Lys	Gln	
					20					25					30	
	Gln	Ala	Ser	Gln	Val	Leu	Val	Arg	Lys	Arg	Arg	Ala	Asn	Ser	Leu	
					35					40					45	
	Leu	Glu	Glu	Thr	Lys	Gln	Gly	Asn	Leu	Glu	Arg	Glu	Cys	Ile	Glu	
					50					55					60	
35	Glu	Leu	Cys	Asn	Lys	Glu	Glu	Ala	Arg	Glu	Val	Phe	Glu	Asn	Asp	

P0929P2

					65					70					75				
					Pro	Glu	Thr	Asp	Tyr	Phe	Tyr	Pro	Lys	Tyr	Leu	Val	Cys	Leu	Arg
					80									85					90
5					Ser	Phe	Gln	Thr	Gly	Leu	Phe	Thr	Ala	Ala	Arg	Gln	Ser	Thr	Asn
					95									100					105
					Ala	Tyr	Pro	Asp	Leu	Arg	Ser	Cys	Val	Asn	Ala	Ile	Pro	Asp	Gln
					110									115					120
					Cys	Ser	Pro	Leu	Pro	Cys	Asn	Glu	Asp	Gly	Tyr	Met	Ser	Cys	Lys
					125									130					135
10					Asp	Gly	Lys	Ala	Ser	Phe	Thr	Cys	Thr	Cys	Lys	Pro	Gly	Trp	Gln
					140									145					150
					Gly	Glu	Lys	Cys	Glu	Phe	Asp	Ile	Asn	Glu	Cys	Lys	Asp	Pro	Ser
					155									160					165
15					Asn	Ile	Asn	Gly	Gly	Cys	Ser	Gln	Ile	Cys	Asp	Asn	Thr	Pro	Gly
					170									175					180
					Ser	Tyr	His	Cys	Ser	Cys	Lys	Asn	Gly	Phe	Val	Met	Leu	Ser	Asn
					185									190					195
					Lys	Lys	Asp	Cys	Lys	Asp	Val	Asp	Glu	Cys	Ser	Leu	Lys	Pro	Ser
					200									205					210
20					Ile	Cys	Gly	Thr	Ala	Val	Cys	Lys	Asn	Ile	Pro	Gly	Asp	Phe	Glu
					215									220					225
					Cys	Glu	Cys	Pro	Glu	Gly	Tyr	Arg	Tyr	Asn	Leu	Lys	Ser	Lys	Ser
					230									235					240
25					Cys	Glu	Asp	Ile	Asp	Glu	Cys	Ser	Glu	Asn	Met	Cys	Ala	Gln	Leu
					245									250					255
					Cys	Val	Asn	Tyr	Pro	Gly	Gly	Tyr	Thr	Cys	Tyr	Cys	Asp	Gly	Lys
					260									265					270
					Lys	Gly	Phe	Lys	Leu	Ala	Gln	Asp	Gln	Lys	Ser	Cys	Glu	Val	Val
					275									280					285
30					Ser	Val	Cys	Leu	Pro	Leu	Asn	Leu	Asp	Thr	Lys	Tyr	Glu	Leu	Leu
					290									295					300
					Tyr	Leu	Ala	Glu	Gln	Phe	Ala	Gly	Val	Val	Leu	Tyr	Leu	Lys	Phe
					305									310					315
35					Arg	Leu	Pro	Glu	Ile	Ser	Arg	Phe	Ser	Ala	Glu	Phe	Asp	Phe	Arg
					320									325					330

P0929P2

	Thr	Tyr	Asp	Ser	Glu	Gly	Val	Ile	Leu	Tyr	Ala	Glu	Ser	Ile	Asp	
					335					340					345	
	His	Ser	Ala	Trp	Leu	Leu	Ile	Ala	Leu	Arg	Gly	Gly	Lys	Ile	Glu	
					350					355					360	
5	Val	Gln	Leu	Lys	Asn	Glu	His	Thr	Ser	Lys	Ile	Thr	Thr	Gly	Gly	
					365					370					375	
	Asp	Val	Ile	Asn	Asn	Gly	Leu	Trp	Asn	Met	Val	Ser	Val	Glu	Glu	
					380					385					390	
10	Leu	Glu	His	Ser	Ile	Ser	Ile	Lys	Ile	Ala	Lys	Glu	Ala	Val	Met	
					395					400					405	
	Asp	Ile	Asn	Lys	Pro	Gly	Pro	Leu	Phe	Lys	Pro	Glu	Asn	Gly	Leu	
					410					415					420	
	Leu	Glu	Thr	Lys	Val	Tyr	Phe	Ala	Gly	Phe	Pro	Arg	Lys	Val	Glu	
					425					430					435	
15	Ser	Glu	Leu	Ile	Lys	Pro	Ile	Asn	Pro	Arg	Leu	Asp	Gly	Cys	Ile	
					440					445					450	
	Arg	Ser	Trp	Asn	Leu	Met	Lys	Gln	Gly	Ala	Ser	Gly	Ile	Lys	Glu	
					455					460					465	
20	Ile	Ile	Gln	Glu	Lys	Gln	Asn	Lys	His	Cys	Leu	Val	Thr	Val	Glu	
					470					475					480	
	Lys	Gly	Ser	Tyr	Tyr	Pro	Gly	Ser	Gly	Ile	Ala	Gln	Phe	His	Ile	
					485					490					495	
	Asp	Tyr	Asn	Asn	Val	Ser	Ser	Ala	Glu	Gly	Trp	His	Val	Asn	Val	
					500					505					510	
25	Thr	Leu	Asn	Ile	Arg	Pro	Ser	Thr	Gly	Thr	Gly	Val	Met	Leu	Ala	
					515					520					525	
	Leu	Val	Ser	Gly	Asn	Asn	Thr	Val	Pro	Phe	Ala	Val	Ser	Leu	Val	
					530					535					540	
30	Asp	Ser	Thr	Ser	Glu	Lys	Ser	Gln	Asp	Ile	Leu	Leu	Ser	Val	Glu	
					545					550					555	
	Asn	Thr	Val	Ile	Tyr	Arg	Ile	Gln	Ala	Leu	Ser	Leu	Cys	Ser	Asp	
					560					565					570	
	Gln	Gln	Ser	His	Leu	Glu	Phe	Arg	Val	Asn	Arg	Asn	Asn	Leu	Glu	
					575					580					585	
35	Leu	Ser	Thr	Pro	Leu	Lys	Ile	Glu	Thr	Ile	Ser	His	Glu	Asp	Leu	

P0929P2

		590		595		600									
	Gln	Arg	Gln	Leu	Ala	Val	Leu	Asp	Lys	Ala	Met	Lys	Ala	Lys	Val
					605					610					615
5	Ala	Thr	Tyr	Leu	Gly	Gly	Leu	Pro	Asp	Val	Pro	Phe	Ser	Ala	Thr
					620					625					630
	Pro	Val	Asn	Ala	Phe	Tyr	Asn	Gly	Cys	Met	Glu	Val	Asn	Ile	Asn
					635					640					645
	Gly	Val	Gln	Leu	Asp	Leu	Asp	Glu	Ala	Ile	Ser	Lys	His	Asn	Asp
					650					655					660
10	Ile	Arg	Ala	His	Ser	Cys	Pro	Ser	Val	Trp	Lys	Lys	Thr	Lys	Asn
					665					670					675
	Ser														
	676														

(2) INFORMATION FOR SEQ ID NO:4:

- 15 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCAAGACAAT GGAACCCAGG 20

(2) INFORMATION FOR SEQ ID NO:5:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 36 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CATGGAATTC GGTGACCGAT GTGCGGCTGT GAGGAG 36

- 30 (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 95 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single

P0929P2

(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCAAGGGCTA CTGCCACACT CGAGCTGCGC AGATGCTAGC CTCAAGATGG 50

CTGATCCAAA TCGATTCCGC GGCAAAGATC TTCCGGTCCT GTAGA 95

5 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 103 base pairs

(B) TYPE: Nucleic Acid

(C) STRANDEDNESS: Single

10 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGCTTCTACA GGACCGGAAG ATCTTTGCCG CGGAATCGAT TTGGATCAGC 50

CATCTTGAGG CTAGCATCTG CGCAGCTCGA GTGTGGCAGT AGCCCTTGCT 100

GCA 103

15 (2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids

(B) TYPE: Amino Acid

(D) TOPOLOGY: Linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Xaa	Gln	Val	Leu	Ile	Arg	Arg	Xaa	Arg	Ala	Asn	Thr	Leu
1				5				10				13

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

25 (A) LENGTH: 4 amino acids

(B) TYPE: Amino Acid

(D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ala	Asn	Thr	Leu
30	1		4

(2) INFORMATION FOR SEQ ID NO:10:

P0929P2

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 5 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GATATCGATC CATGGCCCCT TCGCTCTC 28

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 10 (A) LENGTH: 31 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

15 CATGGATCCT ACCGGAAGTC AAACCTCAGCT A 31

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 28 base pairs
 (B) TYPE: Nucleic Acid
 20 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GATATCGATG AGTGTGAAGT CCTTGTAC 28

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 25 (A) LENGTH: 27 base pairs
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GTCGGATCCG ACAGAGACTG AGAAGCC 27

(2) INFORMATION FOR SEQ ID NO:14:

P0929P2

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 8 amino acids
 - (B) TYPE: Amino Acid
 - (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ala Phe Gln Val Phe Glu Glu Ala
 1 5 8

(2) INFORMATION FOR SEQ ID NO:15:

- 10 (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

15 AGCTGCTCGA GGCGCTGTTG CCGGCGC 27

(2) INFORMATION FOR SEQ ID NO:16:

- 20 (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AGCTGCTCGA GGCAAATTCT TTACTTGAA 29

(2) INFORMATION FOR SEQ ID NO:17:

- 25 (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AGCTGCTCGA GGACCAGTGC ACGCCCAACC 30

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:

P0929P2

- (A) LENGTH: 28 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GCTGCTCGAG GACATCTTGC CGTGCGTG 28

(2) INFORMATION FOR SEQ ID NO:19:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CATTCATTTA TGTCAAATTC A 21

15 (2) INFORMATION FOR SEQ ID NO:20:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 17 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATGGAGATCA AGGTCTG 17

(2) INFORMATION FOR SEQ ID NO:21:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 33 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

30 CATCTTGAGG CTAGCGGCTG CGGCGGGCTC CAC 33

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:

P0929P2

- (A) LENGTH: 27 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GTCGGTGACC GCTGCTGCGG GCTCCAC 27

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 50 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GAAATTACAG ATCCGCAGCC CCGGGATGGG GCCGGCCCCG CTGCCGCTGC 50

15 (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 50 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CCTTGGATTC TAGCAAGCAC GACTGAAGGA GCCCCATCAG TAGCACCTTT 50

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 50 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

30 TCTTAAAATT AAGCTTCAGC TGCTCCTTGA TATTACCTT TGTACAGAGT 50

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:

P0929P2

- (A) LENGTH: 15 base pairs
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GCGCCTGGCA ACGCG 15

(2) INFORMATION FOR SEQ ID NO:27:

- 10 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GTGACCGCGT TGCCAG 16

15 (2) INFORMATION FOR SEQ ID NO:28:

- 20 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 34 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

CGAATTCCTC GAGCCGGGAC CTTTCCAGG GAGC 34

(2) INFORMATION FOR SEQ ID NO:29:

- 25 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 29 base pairs
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

30 CCAACTGTGT GTTTGAAGGC AAGAGGCGG 29

WHAT IS CLAIMED IS:

1. Variant gas6 polypeptide which lacks one or more glutamic acid residues from the A domain of native gas6.
2. The variant gas6 of claim 1 which lacks the A domain of native gas6.
3. The variant gas6 of claim 2 which is the D domain of gas6.
4. The variant gas6 of claim 2 which is a G domain of gas6.
5. The variant gas6 of claim 1 which is derived from human gas6.
6. A composition comprising the variant gas6 of claim 1 and a physiologically acceptable carrier.
7. Nucleic acid encoding the gas6 variant of claim 1.
8. A vector comprising the nucleic acid of claim 7.
9. A host cell comprising the nucleic acid of claim 7.
10. A method of making variant gas6 polypeptide comprising culturing the host cell of claim 9 so that the nucleic acid is expressed and recovering the gas6 variant from the cell culture.
11. A method of making variant gas6 comprising:
 - (a) culturing a host cell comprising nucleic acid encoding gas6 under conditions such that the nucleic acid is expressed and the gas6 polypeptide thus produced is essentially not γ -carboxylated, and
 - (b) recovering the gas6 variant from the cell culture.
12. The method of claim 11 wherein the culturing is carried out in the absence of Vitamin K.
13. The method of claim 11 wherein the host cell is deficient in γ -carboxylase enzyme.
14. The method of claim 13 wherein the host cell is non-mammalian.
15. The use of an effective amount of exogenous gas6 polypeptide for activating Rse receptor on a cell comprising the Rse receptor.
16. The use of claim 15 wherein the gas6 comprises human gas6.
17. The use of claim 15 wherein the gas6 comprises a variant gas6 which is essentially not γ -carboxylated.
18. The use of claim 15 wherein the gas6 comprises an immunoadhesin.
19. The use of claim 15 wherein the cell is a glial cell.
20. The use of claim 19 wherein the glial cell is a Schwann cell.

21. The use of an effective amount of an exogenous gas6 polypeptide for enhancing survival, proliferation or differentiation of a cell comprising the Rse receptor.
22. The use of claim 21 wherein the gas6 is human gas6.
23. The use of claim 22 wherein the cell is a human cell.
24. The use of claim 21 which comprises enhancing proliferation of the cell.
25. An article of manufacture, comprising: a container; a label on said container; and a composition contained within said container; wherein the composition is effective for enhancing survival, differentiation or proliferation of a cell comprising the Rse receptor, the label on said container indicates that the composition can be used for enhancing survival, differentiation or proliferation of said cell, and the composition comprises gas6 polypeptide and a physiologically acceptable carrier.
26. The use of an effective amount of an exogenous gas6 polypeptide for activating Mer receptor of a cell comprising the Mer receptor.
27. The use of claim 26 wherein the gas6 comprises human gas6.
28. The use of claim 26 wherein the cell is a mononuclear cell.
29. The use of an effective amount of an exogenous gas6 polypeptide for enhancing survival, proliferation or differentiation of a cell comprising the Mer receptor.

Gla Region Loop EGF-like Repeats SHBG-like Domain




FIG.1A

49	90	118	279	678		
A	B	C ₁	C ₂	C ₃	C ₄	D

h Gas6

FIG.1B

42	83	119	284	675
68	21	45	56	59
38				

b Protein S

FIG.1C

42	83	119	284	676		
78	21	45	59	46	59	40

h Protein S

FIG.1D




FIG.2A

C domain	mGas 6	D I D E C T - D S D T C G D A R C K N L P G S Y S C L C D E G Y T Y S S K E K T C C Q R E
	hGas 6	D I D E C A - D S E A C G D A R C K N L P G S Y S C L C D E G Y T Y S S K E K T A C C
	hProts	D V D E C S L K P S I C G D A V C K N T P G D F E C E C P E G Y F A R Y N L K S K
	mGas 6	D V D E C Q Q D R C E Q T C V N S P G S Y T C H C D G R G G L K L S P D M D T C E
	hGas 6	D V D E C L Q G R C E Q V C V N S P G S Y T C H C D G R G G L K L S P D M D T C E
	hProts	D T D E C S E N M C A Q L C V N Y P G G Y T C Y C D G K K G F K L A Q D Q K S C E
	mGas 6	276 D I L P C V P F S M A K S V K S L Y L G R M F S G T P V I
	hGas 6	279 D I L P C V P F S V A K S V K S L Y L G R M F S G T P V I
	hProts	284 V V S V C L P L N L D T K Y E L L Y L A E Q F A G V - V T L
		→ G domain 1
		R L R F K R L Q P T R L L A E F D F R T F D P E G V L F F
		R L R F K R L Q P T R L L A E F D F R T F D P E G V L F F
	mGas 6	A G G R S D S T W I V L G L R A G R L E L Q L R Y N G V G
	hGas 6	A G G H Q D S T W I V L A L R A G R L E L Q L R Y N G V G
	hProts	A E S I D H S A W L L I A L R G G K I E V Q L K N E H T S
		R I T S S G P T I N H G M W Q T I S V E E L E R N L V I K
		R V T S S G P V I N H G M W Q T I S V E E L A R N L V I K
		K I T T G G D V I N N G L W N M V S V E E L E H S I S I K
D domain	mGas 6	V N K D A V M K I A V A G E L F Q L E R G L - Y H L N L T
	hGas 6	V N R D A V M K I A V A G D L F Q L E R G L - Y H L N L T
	hProts	I A K E A V M D I N K P G P L F K P E N G L L E T K V Y F
		V G G I I F K E S E L V Q P I N P R L D G C M R S W N W L
		V G G I P F E H E S E L V Q P I N P R L D G C M R S W N W L
		A G F P R K V E S E L T K P I N P R L D G C T R S W N L M

FIG.2B




FIG.2C

5/17




FIG. 3A




FIG. 3B




FIG. 3C




FIG.3D

7/17




FIG.4




FIG. 5




FIG. 6A




FIG. 6B






FIG. 6C

**FIG.8****FIG.9**

12/17




FIG. 10

13/17






FIG. 11

14 / 17






FIG.13




FIG.12C

16/17




FIG. 14

17/17




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D

Gla Region Loop EGF-like Repeats

SHBG-like Domain

