



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2348302 C 2008/08/26

(11)(21) **2 348 302**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1999/11/17
(87) Date publication PCT/PCT Publication Date: 2000/05/25
(45) Date de délivrance/Issue Date: 2008/08/26
(85) Entrée phase nationale/National Entry: 2001/04/25
(86) N° demande PCT/PCT Application No.: JP 1999/006405
(87) N° publication PCT/PCT Publication No.: 2000/029026
(30) Priorité/Priority: 1998/11/17 (JP10/326251)

(51) Cl.Int./Int.Cl. *A61K 47/38* (2006.01),
A61K 9/02 (2006.01)
(72) Inventeurs/Inventors:
FUJIMORI, TOMOKO, JP;
SUGITA, KIMIKO, JP
(73) Propriétaire/Owner:
TAISHO PHARMACEUTICAL CO., LTD., JP
(74) Agent: TORYS LLP

(54) Titre : SUPPOSITOIRES COMPRENANT DE L'HYDROXYPROPYLMETHYL CELLULOSE HYDROPHOBE
(54) Title: SUPPOSITORIES COMPRISING HYDROPHOBIC HYDROXYPROPYL METHYLCELLULOSE

(57) **Abrégé/Abstract:**

The present invention provides a suppository comprising a hydrophobic hydroxypropyl methylcellulose. The present invention is useful as a suppository base with little interaction with a drug and has an excellent retainability.



ABSTRACT

The present invention provides a suppository comprising a hydrophobic hydroxypropyl methylcellulose.

The present invention is useful as a suppository base with little interaction with a drug and has an excellent retainability.

1

DESCRIPTION

SUPPOSITORIES COMPRISING HYDROPHOBIC HYDROXYPROPYL
METHYLCELLULOSE

TECHNICAL FIELD

The present invention relates to
suppositories, and more particularly, relates to
suppositories which is excellently retainable between
5 the anus and the lower region of the rectum.

BACKGROUND ART

It is known that a suppository, after
administered rectally, is disadvantageously spread in
the rectum so that the drug is diluted around the
10 affected site. There are reported a number of
techniques for inhibition of such spreading, for
example, a technique combining metal salts of
polyacrylic acid in JP-A-54-26325, a technique
combining a mixture of polyacrylic acid and
15 polyvinylpyrrolidone in JP-A-6-40889, a technique
combining carboxyvinyl polymers in JP-A-63-280016 and
JP-A-1-143825, a technique combining a gel-forming
agent such as alkali metal salts of polygum in JP-A-59-
55817, a technique combining polyvinyl alcohol, pectin
20 and pullulan in JP-A-61-109710, a technique combining
water-swelling clay minerals such as Laponite in JP-A-
2-15024, a technique combining water-insoluble
absorptive resins such as starch grafted acrylate in
JP-A-4-164023, etc.

However, in order to use a suppository as a pharmaceutical preparation, it is necessary to satisfy not only retainability but also many other requirements such as safety, stability or the like. For this
5 reason, acrylate polymers, especially carboxyvinyl polymers, are actually used in many cases.

Carboxyvinyl polymers, when used as a suppository base, have an excellent retainability around the affected site, but are liable to cause an
10 interaction with the main drug, therefore they have a drawback of lowering the stability of the main drug at times. For example, mixing a certain basic drug with carboxyvinyl polymers causes coloring, precipitating or lowering the release of the main drug at times.
15 Furthermore, in order to enhance the retainability of the suppository, it is necessary to elevate the viscosity of the gel by combining a base component for neutralization of the carboxyvinyl polymers, and in this case, some of the drugs are reacted with the base
20 component to cause coloring or precipitating.

Thus, use of carboxyvinyl polymers as a suppository base has a problem that only a limited number of drugs can be applied. An object of the present invention is to provide a novel suppository
25 base with little interaction with the drug, and has an excellent retainability.

DISCLOSURE OF THE INVENTION

As a result of various studies in order to solve the above problem, the present inventors have found that a suppository that can contain various kinds of drugs has a high retainability can be obtained by combining a suppository base with a hydrophobic hydroxypropyl methylcellulose in place of conventional carboxyvinyl polymers to accomplish the present invention.

That is, the present invention is directed to a suppository comprising a hydrophobic hydroxypropyl methylcellulose.

It is known that hydrophobic hydroxypropyl methylcellulose can be used as a base of external preparations for skin use or gel preparations for periodontal use as described in JP-A-4-74108, JP-A-3-223301 or JP-A-6-166614, but it has not been known that hydrophobic hydroxypropyl methylcellulose can be combined in suppositories.

The hydrophobic hydroxypropyl methylcellulose in the present invention refers to hydroxypropyl methylcellulose (HPMC) into which a small amount of a long chain alkyloxyhydroxypropoxyl group has been introduced to impart a hydrophobic property to the HPMC. Here, the long chain alkyl group as a part of the alkyloxyhydroxypropoxyl group refers to a straight or branched chain alkyl group having 6 to 26 carbon atoms, and specific examples thereof are a stearyl

group, a palmityl group, a myristyl group and a lauryl group. The most preferred long chain alkyloxyhydroxypropoxyl group in such hydrophobic hydroxypropyl methylcellulose is a stearyloxyhydroxypropoxyl group.

5 The properties of the hydrophobic hydroxypropyl methylcellulose in the present invention vary largely depending on the amount and molecular weight of the long chain alkyloxyhydroxypropoxyl group introduced. Preferred are hydrophobic hydroxypropyl
10 methylcelluloses having a viscosity of 70 - 250 mm²/s (determined with Ubbelohde viscometer, at 25°C, 0.5 W/W%, a mixture of water and isopropanol (6:4 W/W)). When the viscosity is less than 70 mm²/s, the retainability of the suppository may be insufficient,
15 and when the viscosity exceeds 250 mm²/s, it may be difficult to formulate the hydrophobic hydroxypropyl methylcellulose into a preparation.

 The hydrophobic hydroxypropyl methylcellulose comprises preferably 0.1 - 20% by mass, and especially
20 preferably 0.5 - 5% by mass of the preparation. In case the hydrophobic hydroxypropyl methylcellulose comprises less than 0.1% by mass, since sufficient viscosity cannot be obtained, the retainability is insufficient. In case the hydrophobic hydroxypropyl
25 methylcellulose comprises more than 20% by mass, it will be difficult to produce the preparation because the workability is bad due to the high viscosity.

 Introduction of a long chain alkyloxyhydroxy-

propoxyl group into HPMC can be made by a method of reacting an alkyloxyhydroxypropoxylating agent (e.g., a halide, an epoxide or an isocyanate) with the HPMC.

The thus-obtained hydrophobic hydroxypropyl
5 methylcellulose is dispersed into an ordinary suppository base to form a gelling base.

The ordinary suppository base to be used includes oleophilic bases and water-soluble bases, and they can be used in combination. The oleophilic bases
10 include cacao butter, lanolin fat and hard fats. The hard fats include, for example, Witepsol (tradename, manufactured by Huls Inc.), Suppocire (tradename, manufactured by Gattefosse Inc.), Isocacao (tradename, manufactured by Kao Corp.), Pharmasol (tradename,
15 manufactured by NOF Corp.), etc. The water-soluble bases include Macrogol.

The suppository of the present invention can be prepared by a method comprising melt-mixing an ordinary suppository base with a hydrophobic
20 hydroxypropyl methylcellulose, uniformly mixing this mixture with a drug and additives, and filling the resulting mixture into a container, a mold or the like, and cold-solidifying the filling. The method for mixing to be used includes any conventional methods.

25 BEST MODE FOR CARRYING OUT THE INVENTION

The present invention is illustrated in more detail by the following examples and experiments. The

hydrophobic hydroxypropyl methylcellulose to be used in Examples is Sangelose 60L (tradename, manufactured by Sankyo Chemicals Co.), i.e., a hydroxypropyl methylcellulose into which stearyloxyhydroxypropoxyl group has been introduced.

Example 1

	Tetrahydrozoline hydrochloride	1 g
	Lidocaine	60 g
	Hydrocortisone acetate	5 g
10	Allantoin	20 g
	Tocopherol acetate	60 g
	Light anhydrous silicic acid	20 g
	Hydrophobic hydroxypropyl methylcellulose	75 g
15	Pharmasol B115 (tradename)	1409 g

In the heat-molten (50 - 70°C) Pharmasol B115 (base) were successively dispersed the other components while stirring. After cooling to about 40°C, the mixture was filled into suppository containers and cold-molded to obtain suppositories.

Example 2

	Phenylephrine hydrochloride	2 g
	Dibucaine	3 g
	Hydrocortisone acetate	2 g
25	Zinc oxide	66 g
	Light anhydrous silicic acid	66 g
	Hydrophobic hydroxypropyl methylcellulose	70 g

Pharmasol B115 (tradename)	1441 g
----------------------------	--------

The same procedures as in Example 1 were repeated with the above formulation to obtain suppositories.

5 Example 3

	Naphazoline hydrochloride	1 g
	Lidocaine	60 g
	Prednisolone acetate	1 g
	Allantoin	20 g
10	Tocopherol acetate	60 g
	Hydrophobic hydroxypropyl methylcellulose	80 g
	Witepsol H15 (tradename: base)	1428 g

The same procedures as in Example 1 were repeated with the above formulation to obtain suppositories.

Example 4

	Naphazoline hydrochloride	1 g
	Lidocaine	60 g
20	Hydrocortisone acetate	5 g
	Diphenhydramine hydrochloride	10 g
	Allantoin	20 g
	Tocopherol acetate	50 g
	Zinc oxide	100 g
25	Hydrophobic hydroxypropyl methylcellulose	70 g
	Witepsol H15 (tradename: base)	1334 g

The same procedures as in Example 1 were

repeated with the above formulation to obtain suppositories.

Example 5

	Tetrahydrozoline hydrochloride	1 g
5	Lidocaine	60 g
	Prednisolone acetate	1 g
	Crotamiton	50 g
	Chlorhexidine hydrochloride	5 g
	Aluminum chlorohydroxy allantoinate	5 g
10	Tocopherol acetate	50 g
	Hydrophobic hydroxypropyl methylcellulose	65 g
	Witepsol W35 (tradename: base)	1413 g

The same procedures as in Example 1 were repeated with the above formulation to obtain suppositories.

Example 6

	Tetrahydrozoline hydrochloride	0.1 g
	Chlorphenylamine maleate powder	0.4 g
20	Chlorhexidine hydrochloride	0.5 g
	Hydrocortisone acetate	0.5 g
	Allantoin	2.0 g
	Lidocaine	6.0 g
	Vitamin E acetate	6.0 g
25	L-menthol	1.0 g
	Hydrophobic hydroxypropyl methylcellulose	6.5 g
	Aerosil 200 (tradename)	2.0 g

Witepsol E85 (tradename)	28 g
Witepsol W35 (tradename)	112 g

The same procedures as in Example 1 were repeated with the above formulation to obtain 100
5 suppositories.

Comparative Example 1

The same procedures as in Example 1 were repeated with the same formulation as in Example 1 except that the hydrophobic hydroxypropyl
10 methylcellulose was replaced with carboxyvinyl polymer in the same amount as that of the hydrophobic hydroxypropyl methylcellulose to obtain suppositories for comparison.

Experiment 1 In Vitro Retaining Test

15 The retainability of suppositories was determined according to the method of Setnikar-Fantelli (J. Pharm. Pharmac., 1971, 23, 490 - 494). Thus, a cellulose membrane tube was placed in the interior of a glass vessel. A suppository was inserted into the
20 lower part of the cellulose membrane tube, and then the tube was occlusively tied by thread under the suppository, after which the glass tube was filled with water at 37 °C, thereby the water pressure acted as a force to move the drug upward, the weight of the
25 suppository remained in the lower part of the cellulose membrane tube was determined to evaluate the retainability of the suppository.

The suppositories of Example 1 and

Comparative Example 1 were used as test samples. The weights of the suppositories were determined in a section of 0 - 10 cm after 10, 30 and 60 minutes, the remaining rates were calculated, and the results are shown in Table 1.

Table 1 In vitro retaining test

Time (min.)	Weight percent of Suppository	
	Example 1	Comparative Example 1
10	97.7	96.8
30	98.0	97.0
60	98.5	97.0

It is apparent from the above table that the suppository of the present invention exhibits a retainability as high as the suppository prepared from conventional carboxyvinyl polymer.

10 Experiment 2

The stability of chlorphenylamine maleate used in the suppository of Example 6 was determined.

Samples were stored at 5°C, room temperature and 35°C for one month, the residual rates of chlorphenylamine maleate in the samples were determined according to an HPLC method. As a result, there were obtained high residual rates, i.e., 100.6%, 99.8% and 99.2% at 5 °C, room temperature and 35 °C, respectively.

INDUSTRIAL APPLICABILITY

The present invention makes it possible to provide an excellent suppository which can contain various kinds of drug and can retain the preparation
5 around the affected site when administered.

In the claims:

1. A suppository comprising a hydrophobic hydroxypropyl methylcellulose having a long chain alkyloxyhydroxypropoxyl group.
- 5 2. A suppository base comprising a hydrophobic hydroxypropyl methylcellulose having a long chain alkyloxyhydroxypropoxyl group.
3. The suppository according to claim 1 wherein the hydrophobic hydroxypropyl methylcellulose comprises 0.1 to 20% by mass of the suppository.
- 10 4. The suppository according to claim 1 wherein the hydrophobic hydroxypropyl methylcellulose is a hydroxypropyl methylcellulose having a stearyloxyhydroxypropoxyl group.
5. The suppository according to claim 1 wherein the long chain alkyl of the alkyloxyhydroxypropoxyl group is a straight or branched chain alkyl group having 6 to 26 carbon atoms.
- 15 6. The suppository base according to claim 2 wherein the long chain alkyl of the alkyloxyhydroxypropoxyl group is a straight or branched chain alkyl group having 6 to 26 carbon atoms.
- 20 7. The suppository base according to claim 2 wherein the hydrophobic hydroxypropyl methylcellulose comprises 0.1 to 20% by mass of the suppository base.
8. The suppository base according to claim 2 wherein the hydrophobic hydroxypropyl methylcellulose is a hydroxypropyl methylcellulose having a stearyloxyhydroxypropoxyl group.
- 25 9. The suppository base according to claim 2 wherein the alkyl of the alkyloxyhydroxypropoxyl group is selected from the group consisting of stearyl, palmityl, myristyl and lauryl.
10. The suppository according to claim 1 wherein the alkyl of the alkyloxyhydroxypropoxyl group is selected from the group consisting of stearyl, palmityl, myristyl and lauryl.
- 30 11. The suppository according to claim 1 wherein said hydrophobic hydroxypropyl methyl cellulose has a viscosity of 70 - 250 mm²/s.
12. The suppository base according to claim 2 wherein said hydrophobic hydroxypropyl methyl cellulose has a viscosity of 70 - 250 mm²/s.
- 35 13. The suppository base according to claim 2 further comprising a base component selected from the group consisting of coco butter, lanolin fat, hard fats and Macrogol.
14. The suppository according to claim 1 further comprising a base component selected from the group consisting of coco butter, lanolin fat, hard fats and Macrogol.