

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
25 October 2007 (25.10.2007)

PCT

(10) International Publication Number
WO 2007/120980 A2

(51) International Patent Classification: Not classified

(21) International Application Number: PCT/US2007/062311

(22) International Filing Date: 16 February 2007 (16.02.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/774,761 17 February 2006 (17.02.2006) US

(71) Applicant (for all designated States except US): RIGEL PHARMACEUTICALS, INC. [US/US]; 1180 Veterans Boulevard, South San Francisco, California 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): CLOUGH, Jeffrey Wayne [US/US]; 3520 Bay Road, Redwood City, California 94063 (US). BHAMIDIPATI, Somasekhar [IN/US]; 888 Foster City Blvd., #Q2, Foster City, California 94404 (US). SINGH, Rajinder [US/US]; 1832 Hillman Avenue, Belmont, California 94002 (US). MASUDA, Esteban [US/US]; 232 Oakhurst Place, Menlo Park, California 94025 (US). ZHAO, Haoran [CN/US]; 330 Port Royal Avenue, Foster City, California 94404 (US).

(74) Agents: FLICK, Karen E. et al.; FOLEY & LARDNER LLP, 1530 Page Mill Road, Palo Alto, California 94304-1125 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 2,4-PYRIMIDINEDIAMINE COMPOUNDS FOR TREATING OR PREVENTING AUTOIMMUNE DISEASES



(57) Abstract: The present invention provides 3-hydroxyphenyl-2,4-pyrimidinediamine compounds of formula (I), as well as related compositions and methods for treating a variety of autoimmune diseases.

2,4-PYRIMIDINEDIAMINE COMPOUNDS FOR TREATING OR PREVENTING AUTOIMMUNE DISEASES

5

FIELD OF THE INVENTION

The present invention relates generally to 3-hydroxyphenyl-2,4-pyrimidinediamine compounds, pharmaceutical compositions comprising the compounds, intermediates and synthetic methods of making the compounds, and methods of using the compounds and compositions in a variety of contexts, such as in the treatment or prevention of various diseases.

10

BACKGROUND OF THE INVENTION

Crosslinking of Fc receptors, such as the high affinity receptor for IgE (Fc ϵ RI) and/or the high affinity receptor for IgG (Fc γ RI) activates a signaling cascade in mast, basophil, and other immune cells that results in the release of chemical mediators responsible for numerous adverse events. For example, such crosslinking leads to the release of preformed mediators of Type I (immediate) anaphylactic hypersensitivity reactions, such as histamine, from storage sites in granules *via* degranulation. It also leads to the synthesis and release of other mediators, including leukotrienes, prostaglandins, and platelet-activating factors (PAFs), that play important roles in inflammatory reactions. Additional mediators that are synthesized and released upon crosslinking Fc receptors include cytokines and nitric oxide.

15

The signaling cascade(s) activated by crosslinking Fc receptors such as Fc ϵ RI and/or Fc γ RI comprises an array of cellular proteins. Among the most important intracellular signal propagators are the tyrosine kinases. An important tyrosine kinase involved in the signal transduction pathways associated with crosslinking the Fc ϵ RI and/or Fc γ RI receptors, as well as other signal transduction cascades, is Syk kinase (see Valent *et al.*, 2002, *Intl. J. Hematol.* 75(4):257-362 for review).

20


Recently, various classes of 2,4-pyrimidinediamine compounds have been discovered that inhibit the Fc ϵ RI and/or Fc γ RI signaling cascades, and that have myriad

therapeutic uses. *See, e.g.*, U.S. application Serial No. 10/355,543 filed January 31, 2003 (US 2004/0029902A1), international application Serial No. PCT/US03/03022 filed January 31, 2003 (WO 03/063794), U.S. application Serial No. 10/631,029 filed July 29, 2003, international application Serial No. PCT/US03/24087 (WO 2004/014382), U.S. application Serial No. 10/903,263 filed July 30, 2004 (US2005/0234049), international application Serial No. PCT/US2004/24716, U.S. application Serial No. 10/903,870 filed July 30, 2004.

As the mediators released as a result of Fc ϵ RI and Fc γ RI receptor cross-linking are responsible for, or play important roles in, the manifestation of numerous adverse events, the availability of compounds capable of inhibiting the signaling cascade(s) responsible for their release would be highly desirable. Moreover, owing to the critical role that Syk kinase plays these and other receptor signaling cascade(s), the availability of compounds capable of inhibiting Syk kinase would also be highly desirable.

SUMMARY OF THE INVENTION

The present invention relates to a compound, stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof according to formula I:



wherein:

Y is selected from the group consisting of S, O, SO, SO₂, and C(R⁷)₂;
each R³⁵ is independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, and halo, or both R³⁵ together with the carbon to which they are attached form a carbonyl group;

W is selected from the group consisting of C=O, C=S, C=NH, C(R⁷)₂, and NR³⁷;
Z is C=O or NR³⁷, provided that Z and W are not both NR³⁷ and provided that when Z is C=O, then W is C(R⁷)₂ or NR³⁷;

X is CH or N;

each R^{31} is independently (C_1 - C_4)alkyl or both R^{31} together form a (C_1 - C_2)alkylene group optionally substituted with one to two (C_1 - C_4)alkyl groups or substituted with one (C_3 - C_7) spirocycloalkyl group;

each R^7 is independently hydrogen or (C_1 - C_4)alkyl; and

5 R^{37} is hydrogen or methyl optionally substituted with phenyl or pyridyl, wherein said phenyl or pyridyl is optionally substituted with (C_1 - C_4)alkoxy.

The present invention also relates to various related features, such as methods, compositions, and uses relating to compounds of formula I and II, that will be readily apparent from the following detailed description.

10 It will be appreciated by one of skill in the art that the implementations summarized above may be used together in any suitable combination to generate implementations not expressly recited above and that such implementations are considered to be part of the present invention.

DETAILED DESCRIPTION

15 Throughout this application, the text refers to various embodiments of the present compounds, compositions, and methods. The various embodiments described are meant to provide a variety of illustrative examples and should not be construed as descriptions of alternative species. Rather it should be noted that the descriptions of various embodiments provided herein may be of overlapping scope. The embodiments discussed herein are 20 merely illustrative and are not meant to limit the scope of the present invention.

Definitions

As used herein, the following definitions shall apply unless otherwise indicated.

“Alkyl” or “Alkanyl” by itself or as part of another substituent refers to monovalent saturated aliphatic hydrocarbyl groups having the stated number of carbon atoms (*i.e.*, C_1 - C_4 means one to four carbon atoms). This term includes, by way of example, linear and branched hydrocarbyl groups such as methyl (CH_3 -), ethyl (CH_3CH_2 -), *n*-propyl ($CH_3CH_2CH_2$ -), isopropyl ($(CH_3)_2CH$ -), *n*-butyl ($CH_3CH_2CH_2CH_2$ -), isobutyl ($(CH_3)_2CHCH_2$ -), *sec*-butyl ($(CH_3)(CH_3CH_2)CH$ -), *t*-butyl ($(CH_3)_3C$ -), *n*-pentyl ($CH_3CH_2CH_2CH_2CH_2$ -), and neopentyl ($(CH_3)_3CCH_2$ -).

“Benzyl” by itself or as part of another substituent refers to the group (C₆H₅)CH₂-.

“Carbonyl” refers to the group >C=O. “Thiocarbonyl” refers to the group >C=S.

“Cyano” by itself or as part of another substituent refers to the group –CN.

“Halogen” or “Halo” by themselves or as part of another substituent, unless
5 otherwise stated, refer to fluoro, chloro, bromo, and iodo.

“Haloalkyl” by itself or as part of another substituent refers to an alkyl group in which one or more of the hydrogen atoms is replaced with a halogen. Thus, the term “haloalkyl” is meant to include monohaloalkyls, dihaloalkyls, trihaloalkyls, etc. up to perhaloalkyls. For example, the expression “(C₁-C₂) haloalkyl” includes fluoromethyl, 10 difluoromethyl, trifluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl, 1,2-difluoroethyl, 1,1,1-trifluoroethyl, perfluoroethyl, etc.

“Nitro” by itself or as part of another substituent refers to –NO₂.

The above-defined groups may include prefixes and/or suffixes that are commonly used in the art to create additional well-recognized substituent groups. As examples, 15 “alkyloxy” or “alkoxy” refers to a group of the formula –OR", where R" is alkyl and includes alkoxy groups such as methoxy and ethoxy. As another example, “haloalkoxy” or “haloalkyloxy” refers to a group of the formula –OR", where R" is a haloalkyl. In other examples, “4-methoxybenzyl” refers to substitution of benzyl at the 4-*para* position with methoxy and “2-pyridylmethyl” refers to substitution of methyl with a 2-pyridyl group.

20 “Alkenyl” by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon double bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkene. The group may be in either the *cis* or *trans* conformation about the double bond(s). Typical alkenyl groups include, but are not limited to, ethenyl; propenyls such as prop-1-en-1-yl, 25 prop-1-en-2-yl, prop-2-en-1-yl, prop-2-en-2-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl; butenyls such as but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, etc.; and the like. As used herein, “lower alkenyl” means (C₂-C₈) alkenyl.

“Alkynyl” by itself or as part of another substituent refers to an unsaturated branched, straight-chain or cyclic alkyl having at least one carbon-carbon triple bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkyne. Typical alkynyl groups include, but are not limited to, ethynyl; propynyls such as 5 prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butynyls such as but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like. As used herein, “lower alkynyl” means (C2-C8) alkynyl.

“Alkyldiyl” by itself or as part of another substituent refers to a saturated or unsaturated, branched, straight-chain or cyclic divalent hydrocarbon group having the stated 10 number of carbon atoms (*i.e.*, C1-C6 means from one to six carbon atoms) derived by the removal of one hydrogen atom from each of two different carbon atoms of a parent alkane, alkene or alkyne, or by the removal of two hydrogen atoms from a single carbon atom of a parent alkane, alkene or alkyne. The two monovalent radical centers or each valency of the divalent radical center can form bonds with the same or different atoms. Typical alkyldiyl groups include, but are not limited to, methandiyl; ethyldiyls such as ethan-1,1-diyl, 15 ethan-1,2-diyl, ethen-1,1-diyl, ethen-1,2-diyl; propyldiyls such as propan-1,1-diyl, propan-1,2-diyl, propan-2,2-diyl, propan-1,3-diyl, cyclopropan-1,1-diyl, cyclopropan-1,2-diyl, prop-1-en-1,1-diyl, prop-1-en-1,2-diyl, prop-2-en-1,2-diyl, prop-1-en-1,3-diyl, cycloprop-1-en-1,2-diyl, cycloprop-2-en-1,2-diyl, cycloprop-2-en-1,1-diyl, prop-1-yn-1,3-diyl, etc.; butyldiyls such as, butan-1,1-diyl, 20 butan-1,2-diyl, butan-1,3-diyl, butan-1,4-diyl, butan-2,2-diyl, 2-methyl-propan-1,1-diyl, 2-methyl-propan-1,2-diyl, cyclobutan-1,1-diyl; cyclobutan-1,2-diyl, cyclobutan-1,3-diyl, but-1-en-1,1-diyl, but-1-en-1,2-diyl, but-1-en-1,3-diyl, but-1-en-1,4-diyl, 2-methyl-prop-1-en-1,1-diyl, 2-methanylidene-propan-1,1-diyl, buta-1,3-dien-1,1-diyl, buta-1,3-dien-1,2-diyl, buta-1,3-dien-1,3-diyl, buta-1,3-dien-1,4-diyl, 25 cyclobut-1-en-1,2-diyl, cyclobut-1-en-1,3-diyl, cyclobut-2-en-1,2-diyl, cyclobuta-1,3-dien-1,2-diyl, cyclobuta-1,3-dien-1,3-diyl, but-1-yn-1,3-diyl, but-1-yn-1,4-diyl, buta-1,3-diyn-1,4-diyl, etc.; and the like. Where specific levels of 30 saturation are intended, the nomenclature alkanyldiyl, alkenyldiyl and/or alkynyldiyl is used. Where it is specifically intended that the two valencies are on the same carbon atom, the nomenclature “alkylidene” is used. In some embodiments, the alkyldiyl group is (C1-C8) alkyldiyl. Specific embodiments include saturated acyclic alkanyldiyl groups in

which the radical centers are at the terminal carbons, *e.g.*, methandiyyl (methano); ethan-1,2-diyyl (ethano); propan-1,3-diyyl (propano); butan-1,4-diyyl (butano); and the like (also referred to as alkylenos, defined *infra*).

“Alkyleno” by itself or as part of another substituent refers to a straight-chain 5 saturated or unsaturated alkylidyyl group having two terminal monovalent radical centers derived by the removal of one hydrogen atom from each of the two terminal carbon atoms of straight-chain parent alkane, alkene or alkyne. The locant of a double bond or triple bond, if present, in a particular alkyleno is indicated in square brackets. Typical alkyleno groups include, but are not limited to, methano; ethylenos such as ethano, etheno, ethyno; 10 propylenos such as propano, prop[1]eno, propa[1,2]dieno, prop[1]yno, etc.; butylenos such as butano, but[1]eno, but[2]eno, buta[1,3]dieno, but[1]yno, but[2]yno, buta[1,3]diyno, etc.; and the like. Where specific levels of saturation are intended, the nomenclature alkano, alkano and/or alkyno is used. In some embodiments, the alkyleno group is (C1-C8) or 15 (C1-C3) alkyleno. Specific embodiments include straight-chain saturated alkano groups, *e.g.*, methano, ethano, propano, butano, and the like.

“Heteroalkyl,” “Heteroalkanyl,” “Heteroalkenyl,” “Heteroalkynyl,” “Heteroalkyldiyyl” and “Heteroalkylene” by themselves or as part of another substituent refer to alkyl, alkanyl, alkenyl, alkynyl, alkylidyyl and alkylene groups, respectively, in 20 which one or more of the carbon atoms are each independently replaced with the same or different heteratoms or heteroatomic groups. Typical heteroatoms and/or heteroatomic groups which can replace the carbon atoms include, but are not limited to, -O-, -S-, -S-O-, -NR'-, -PH-, -S(O)-, -S(O)₂-, -S(O) NR'-, -S(O)₂NR'-, and the like, including combinations thereof, where each R' is independently hydrogen or (C1-C8) alkyl.

“Cycloalkyl” and “Heterocycloalkyl” by themselves or as part of another substituent 25 refer to cyclic versions of “alkyl” and “heteroalkyl” groups, respectively. For heteroalkyl groups, a heteroatom can occupy the position that is attached to the remainder of the molecule. Typical cycloalkyl groups include, but are not limited to, cyclopropyl; cyclobutyls such as cyclobutanyl and cyclobutenyl; cyclopentyls such as cyclopentanyl and cyclopentenyl; cyclohexyls such as cyclohexanyl and cyclohexenyl; and the like. Typical 30 heterocycloalkyl groups include, but are not limited to, tetrahydrofuranyl (*e.g.*, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, etc.), piperidinyl (*e.g.*, piperidin-1-yl, piperidin-

2-yl, etc.), morpholinyl (e.g., morpholin-3-yl, morpholin-4-yl, etc.), piperazinyl (e.g., piperazin-1-yl, piperazin-2-yl, etc.), and the like.

“Acyclic Heteroatomic Bridge” refers to a divalent bridge in which the backbone atoms are exclusively heteroatoms and/or heteroatomic groups. Typical acyclic 5 heteroatomic bridges include, but are not limited to, -O-, -S-, -S-O-, -NR'-, -PH-, -S(O)-, -S(O)₂-, -S(O) NR'-, -S(O)₂NR'-, and the like, including combinations thereof, where each R' is independently hydrogen or (C1-C8) alkyl.

“Parent Aromatic Ring System” refers to an unsaturated cyclic or polycyclic ring system having a conjugated π electron system. Specifically included within the definition 10 of “parent aromatic ring system” are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, fluorene, indane, indene, phenalene, tetrahydronaphthalene, etc. Typical parent aromatic ring systems include, but are not limited to, aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, 15 fluorene, hexacene, hexaphene, hexalene, indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, tetrahydronaphthalene, triphenylene, trinaphthalene, and the like.

“Aryl” by itself or as part of another substituent refers to a monovalent aromatic 20 hydrocarbon group having the stated number of carbon atoms (*i.e.*, C₆-C₁₅ means from 6 to 15 carbon atoms) derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Typical aryl groups include, but are not limited to, groups derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, 25 *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like, as well as the various hydro isomers thereof. In preferred embodiments, the aryl group is (C₆-C₁₅) aryl, with (C₆-C₁₀) being more typical. Specific exemplary aryls include phenyl and 30 naphthyl.

“Arylaryl” by itself or as part of another substituent refers to a monovalent hydrocarbon group derived by the removal of one hydrogen atom from a single carbon atom of a ring system in which two or more identical or non-identical parent aromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent aromatic ring systems involved. Typical 5 arylaryl groups include, but are not limited to, biphenyl, triphenyl, phenyl-naphthyl, binaphthyl, biphenyl-naphthyl, and the like. Where the number of carbon atoms in an arylaryl group are specified, the numbers refer to the carbon atoms comprising each parent aromatic ring. For example, (C6-C15) arylaryl is an arylaryl group in which each aromatic 10 ring comprises from 6 to 15 carbons, *e.g.*, biphenyl, triphenyl, binaphthyl, phenylnaphthyl, etc. In some embodiments, each parent aromatic ring system of an arylaryl group is independently a (C6-C15) aromatic, more preferably a (C6-C10) aromatic. Specific 15 exemplary arylaryl groups include those in which all of the parent aromatic ring systems are identical, *e.g.*, biphenyl, triphenyl, binaphthyl, trinaphthyl, etc.

“Biaryl” by itself or as part of another substituent refers to an arylaryl group having 15 two identical parent aromatic systems joined directly together by a single bond. Typical biaryl groups include, but are not limited to, biphenyl, binaphthyl, bianthracyl, and the like. In some embodiments, the aromatic ring systems are (C6-C15) aromatic rings, more 20 typically (C6-C10) aromatic rings. A particular exemplary biaryl group is biphenyl.

“Arylalkyl” by itself or as part of another substituent refers to an acyclic alkyl group 20 in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with an aryl group. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenylethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and 25 the like. Where specific alkyl moieties are intended, the nomenclature arylalkanyl, arylalkenyl and/or arylalkynyl is used. In some embodiments, the arylalkyl group is (C7-C21) arylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C1-C6) and the aryl moiety is (C6-C15). In some specific embodiments the arylalkyl 30 group is (C7-C13), *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is (C1-C3) and the aryl moiety is (C6-C10).

“Parent Heteroaromatic Ring System” refers to a parent aromatic ring system in

which one or more carbon atoms are each independently replaced with the same or different heteroatoms or heteroatomic groups. Typical heteroatoms or heteroatomic groups to replace the carbon atoms include, but are not limited to, N, NH, P, O, S, S(O), S(O)₂, Si, etc. Specifically included within the definition of “parent heteroaromatic ring systems” are fused 5 ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, benzodioxan, benzofuran, chromane, chromene, indole, indoline, xanthene, etc. Also included in the definition of “parent heteroaromatic ring system” are those recognized rings that include common substituents, such as, for example, benzopyrone and 1-methyl-1,2,3,4-tetrazole. Specifically excluded 10 from the definition of “parent heteroaromatic ring system” are benzene rings fused to cyclic polyalkylene glycols such as cyclic polyethylene glycols. Typical parent heteroaromatic ring systems include, but are not limited to, acridine, benzimidazole, benzisoxazole, benzodioxan, benzodioxole, benzofuran, benzopyrone, benzothiadiazole, benzothiazole, benzotriazole, benzoxazine, benzoxazole, benzoxazoline, carbazole, β -carboline, chromane, 15 chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, 20 thiophene, triazole, xanthene, and the like.

“Heteroaryl” by itself or as part of another substituent refers to a monovalent heteroaromatic group having the stated number of ring atoms (e.g., “5-14 membered” means from 5 to 14 ring atoms) derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. Typical heteroaryl groups include, but are not 25 limited to, groups derived from acridine, benzimidazole, benzisoxazole, benzodioxan, benzodiazole, benzofuran, benzopyrone, benzothiadiazole, benzothiazole, benzotriazole, benzoxazine, benzoxazole, benzoxazoline, carbazole, β -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, 30 oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole,

pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like, as well as the various hydro isomers thereof. In preferred embodiments, the heteroaryl group is a 5-14 membered heteroaryl, with 5-10 membered heteroaryl being particularly preferred.

5 “Heteroaryl-Heteroaryl” by itself or as part of another substituent refers to a monovalent heteroaromatic group derived by the removal of one hydrogen atom from a single atom of a ring system in which two or more identical or non-identical parent heteroaromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent heteroaromatic ring 10 systems involved. Typical heteroaryl-heteroaryl groups include, but are not limited to, bipyridyl, tripyridyl, pyridylpurinyl, bipurinyl, etc. Where the number of atoms are specified, the numbers refer to the number of atoms comprising each parent heteroaromatic ring systems. For example, 5-15 membered heteroaryl-heteroaryl is a heteroaryl-heteroaryl group in which each parent heteroaromatic ring system comprises from 5 to 15 atoms, *e.g.*, 15 bipyridyl, tripuridyl, etc. In some embodiments, each parent heteroaromatic ring system is independently a 5-15 membered heteroaromatic, more typically a 5-10 membered heteroaromatic. Specific exemplary heteroaryl-heteroaryl groups include those in which all of the parent heteroaromatic ring systems are identical.


20 “Biheteroaryl” by itself or as part of another substituent refers to a heteroaryl-heteroaryl group having two identical parent heteroaromatic ring systems joined directly together by a single bond. Typical biheteroaryl groups include, but are not limited to, bipyridyl, bipurinyl, biquinolinyl, and the like. In some embodiments, the heteroaromatic ring systems are 5-15 membered heteroaromatic rings, more typically 5-10 membered heteroaromatic rings.

25 “Heteroarylalkyl” by itself or as part of another substituent refers to an acyclic alkyl group in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heteroaryl group. Where specific alkyl moieties are intended, the nomenclature heteroarylalkanyl, heteroarylalkenyl and/or heteroarylalkynyl is used. In some embodiments, the heteroarylalkyl group is a 6-21 membered heteroarylalkyl, 30 *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the heteroarylalkyl is (C1-C6) alkyl and the heteroaryl moiety is a 5-15-membered heteroaryl. In some specific exemplary

embodiments, the heteroarylalkyl is a 6-13 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety is (C1-C3) alkyl and the heteroaryl moiety is a 5-10 membered heteroaryl.

“Halogen” or “Halo” by themselves or as part of another substituent, unless 5 otherwise stated, refer to fluoro, chloro, bromo and iodo.

“(C₃-C₇) Spirocycloalkyl” refers to divalent cyclic groups from 3 to 7 carbon atoms having a 3 to 7 membered cycloalkyl ring with a spiro union (the union formed by a single atom which is the only common member of the rings) as exemplified by the following C₃ structure:



10

“Substituted,” when used to modify a specified group or radical, means that one or more hydrogen atoms of the specified group or radical are each, independently of one another, replaced with the same or different substituent(s). Substituent groups useful for substituting for hydrogens on saturated carbon atoms in the specified group or radical 15 include, but are not limited to -R⁶⁰, halo, -O⁻M⁺, =O, -OR⁷⁰, -SR⁷⁰, -SM⁺, =S, -NR⁸⁰R⁸⁰, =NR⁷⁰, =N-OR⁷⁰, trihalomethyl, -CF₃, -CN, -OCN, -SCN, -NO, -NO₂, =N₂, -N₃, -S(O)₂R⁷⁰, -S(O)₂O⁻M⁺, -S(O)₂OR⁷⁰, -OS(O)₂R⁷⁰, -OS(O)₂O⁻M⁺, -OS(O)₂OR⁷⁰, -P(O)(O⁻)₂(M⁺)₂, -P(O)(OR⁷⁰)O⁻M⁺, -P(O)(OR⁷⁰)(OR⁷⁰), -C(O)R⁷⁰, -C(S)R⁷⁰, -C(NR⁷⁰)R⁷⁰, -C(O)O⁻M⁺, -C(O)OR⁷⁰, -C(S)OR⁷⁰, -C(O)NR⁸⁰R⁸⁰, -C(NR⁷⁰)NR⁸⁰R⁸⁰, -OC(O)R⁷⁰, -OC(S)R⁷⁰, 20 -OC(O)O⁻M⁺, -OC(O)OR⁷⁰, -OC(S)OR⁷⁰, -NR⁷⁰C(O)R⁷⁰, -NR⁷⁰C(S)R⁷⁰, -NR⁷⁰C(O)O⁻M⁺, -NR⁷⁰C(O)OR⁷⁰, -NR⁷⁰C(S)OR⁷⁰, -NR⁷⁰C(O)NR⁸⁰R⁸⁰, -NR⁷⁰C(NR⁷⁰)R⁷⁰ and -NR⁷⁰C(NR⁷⁰)NR⁸⁰R⁸⁰, where R⁶⁰ is selected from the group consisting of alkyl, cycloalkyl, heteroalkyl, cycloheteroalkyl, aryl, arylalkyl, heteroaryl and heteroarylalkyl; each R⁷⁰ is independently hydrogen or R⁶⁰; each R⁸⁰ is independently R⁷⁰ or alternatively, the two R⁸⁰s, 25 taken together with the nitrogen atom to which they are bonded, form a 5-, 6- or 7-membered cycloheteroalkyl which may optionally include from 1 to 4 of the same or different additional heteroatoms selected from the group consisting of O, N and S; and each M⁺ is a counter ion with a positive charge, for example, a positive charge independently

selected from K^+ , Na^+ , $^+N(R^{60})_4$, and Li^+ , or two of M^+ , combine to form a divalent counterion, for example a divalent counterion selected from Ca^{2+} , Mg^{2+} , and Ba^{2+} . As specific examples, $-NR^{80}R^{80}$ is meant to include $-NH_2$, $-NH$ -alkyl, N -pyrrolidinyl and N -morpholinyl.

5 Similarly, substituent groups useful for substituting for hydrogens on unsaturated carbon atoms in the specified group or radical include, but are not limited to, $-R^{60}$, halo, $-O^-$ M^+ , $-OR^{70}$, $-SR^{70}$, $-S^-M^+$, $-NR^{80}R^{80}$, trihalomethyl, $-CF_3$, $-CN$, $-OCN$, $-SCN$, $-NO$, $-NO_2$, $-N_3$, $-S(O)_2R^{70}$, $-S(O)_2O^-M^+$, $-S(O)_2OR^{70}$, $-OS(O)_2R^{70}$, $-OS(O)_2O^-M^+$, $-OS(O)_2OR^{70}$, $-P(O)(O^-)_2(M^+)_2$, $-P(O)(OR^{70})O^-M^+$, $-P(O)(OR^{70})(OR^{70})$, $-C(O)R^{70}$, $-C(S)R^{70}$, $-C(NR^{70})R^{70}$, $-C(O)O^-M^+$, $-C(O)OR^{70}$, $-C(S)OR^{70}$, $-C(O)NR^{80}R^{80}$, $-C(NR^{70})NR^{80}R^{80}$, $-OC(O)R^{70}$, $-OC(S)R^{70}$, $-OC(O)O^-M^+$, $-OC(O)OR^{70}$, $-OC(S)OR^{70}$, $-NR^{70}C(O)R^{70}$, $-NR^{70}C(S)R^{70}$, $-NR^{70}C(O)O^-M^+$, $-NR^{70}C(O)OR^{70}$, $-NR^{70}C(S)OR^{70}$, $-NR^{70}C(O)NR^{80}R^{80}$, $-NR^{70}C(NR^{70})R^{70}$ and $-NR^{70}C(NR^{70})NR^{80}R^{80}$, where R^{60} , R^{70} , R^{80} and M^+ are as previously defined.

15 Substituent groups, other than R^p , useful for substituting for hydrogens on nitrogen atoms in heteroalkyl and cycloheteroalkyl groups include, but are not limited to, $-R^{60}$, $-O^-$ M^+ , $-OR^{70}$, $-SR^{70}$, $-S^-M^+$, $-NR^{80}R^{80}$, trihalomethyl, $-CF_3$, $-CN$, $-NO$, $-NO_2$, $-S(O)_2R^{70}$, $-S(O)_2O^-M^+$, $-S(O)_2OR^{70}$, $-OS(O)_2R^{70}$, $-OS(O)_2O^-M^+$, $-OS(O)_2OR^{70}$, $-P(O)(O^-)_2(M^+)_2$, $-P(O)(OR^{70})O^-M^+$, $-P(O)(OR^{70})(OR^{70})$, $-C(O)R^{70}$, $-C(S)R^{70}$, $-C(NR^{70})R^{70}$, $-C(O)OR^{70}$, $-C(S)OR^{70}$, $-C(O)NR^{80}R^{80}$, $-C(NR^{70})NR^{80}R^{80}$, $-OC(O)R^{70}$, $-OC(S)R^{70}$, $-OC(O)OR^{70}$, $-OC(S)OR^{70}$, $-NR^{70}C(O)R^{70}$, $-NR^{70}C(S)R^{70}$, $-NR^{70}C(O)OR^{70}$, $-NR^{70}C(S)OR^{70}$, $-NR^{70}C(O)NR^{80}R^{80}$, $-NR^{70}C(NR^{70})R^{70}$ and $-NR^{70}C(NR^{70})NR^{80}R^{80}$, where R^{60} , R^{70} , R^{80} and M^+ are as previously defined.

20 Substituent groups from the above lists useful for substituting other groups or atoms specified as “substituted” will be apparent to those of skill in the art.

25 “Protecting group” refers to a group of atoms that, when attached to a reactive functional group in a molecule, mask, reduce or prevent the reactivity of the functional group. Typically, a protecting group may be selectively removed as desired during the course of a synthesis. Examples of protecting groups can be found in Greene and Wuts, *Protective Groups in Organic Chemistry*, 3rd Ed., 1999, John Wiley & Sons, NY and Harrison *et al.*, *Compendium of Synthetic Organic Methods*, Vols. 1-8, 1971-1996, John

Wiley & Sons, NY. Representative amino protecting groups include, but are not limited to, formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (“CBZ”), *tert*-butoxycarbonyl (“Boc”), trimethylsilyl (“TMS”), 2-trimethylsilyl-ethanesulfonyl (“TES”), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (“FMOC”), nitroveratryloxycarbonyl (“NVOC”) and the like. Representative hydroxyl protecting groups include, but are not limited to, those where the hydroxyl group is either acylated or alkylated such as benzyl and trityl ethers, as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers (e.g., TMS or TIPPS groups) and allyl ethers.

“Fc Receptor” refers to a member of the family of cell surface molecules that binds the Fc portion (containing the specific constant region) of an immunoglobulin. Each Fc receptor binds immunoglobulins of a specific type. For example the Fc α receptor (“Fc α R”) binds IgA, the Fc ϵ R binds IgE and the Fc γ R binds IgG.

The Fc α R family includes the polymeric Ig receptor involved in epithelial transport of IgA/IgM, the myeloid specific receptor R α RI (also called CD89), the Fc α / μ R and at least two alternative IgA receptors (for a recent review see Monteiro & van de Winkel, 2003, *Annu. Rev. Immunol.*, advanced e-publication). The Fc α RI is expressed on neutrophils, eosinophils, monocytes/macrophages, dendritic cells and kupffer cells. The Fc α RI includes one alpha chain and the FcR gamma homodimer that bears an activation motif (ITAM) in the cytoplasmic domain and phosphorylates Syk kinase.

The Fc ϵ R family includes two types, designated Fc ϵ RI and Fc ϵ RII (also known as CD23). Fc ϵ RI is a high affinity receptor (binds IgE with an affinity of about $10^{10} M^{-1}$) found on mast, basophil and eosinophil cells that anchors monomeric IgE to the cell surface. The Fc ϵ RII possesses one alpha chain, one beta chain and the gamma chain homodimer discussed above. The Fc ϵ RII is a low affinity receptor expressed on mononuclear phagocytes, B lymphocytes, eosinophils and platelets. The Fc ϵ RII comprises a single polypeptide chain and does not include the gamma chain homodimer.

The Fc γ R family includes three types, designated Fc γ RI (also known as CD64), Fc γ RII (also known as CD32) and Fc γ RIII (also known as CD16). Fc γ RI is a high affinity receptor (binds IgG1 with an affinity of $10^8 M^{-1}$) found on mast, basophil, mononuclear, neutrophil, eosinophil, dendritic and phagocyte cells that anchors nomomeric IgG to the cell

surface. The Fc γ RI includes one alpha chain and the gamma chain dimer shared by Fc α RI and Fc ϵ RI.

The Fc γ RII is a low affinity receptor expressed on neutrophils, monocytes, eosinophils, platelets and B lymphocytes. The Fc γ RII includes one alpha chain, and does not include the gamma chain homodimer discussed above.

The Fc γ RIII is a low affinity (binds IgG1 with an affinity of $5 \times 10^5 M^{-1}$) expressed on NK, eosinophil, macrophage, neutrophil and mast cells. It comprises one alpha chain and the gamma homodimer shared by Fc α RI, Fc ϵ RI and Fc γ RI.

Skilled artisans will recognize that the subunit structure and binding properties of these various Fc receptors, as well as the cell types expressing them, are not completely characterized. The above discussion merely reflects the current state-of-the-art regarding these receptors (see, e.g., Immunobiology: The Immune System in Health & Disease, 5th Edition, Janeway et al., Eds, 2001, ISBN 0-8153-3642-x, Figure 9.30 at pp. 371), and is not intended to be limiting with respect to the myriad receptor signaling cascades that can be regulated with the prodrugs described herein.

“Fc Receptor-Mediated Degranulation” or “Fc Receptor-Induced Degranulation” refers to degranulation that proceeds *via* an Fc receptor signal transduction cascade initiated by crosslinking of an Fc receptor.

“IgE-Induced Degranulation” or “Fc ϵ RI-Mediated Degranulation” refers to degranulation that proceeds *via* the IgE receptor signal transduction cascade initiated by crosslinking of Fc ϵ RI-bound IgE. The crosslinking may be induced by an IgE-specific allergen or other multivalent binding agent, such as an anti-IgE antibody. In mast and/or basophil cells, the Fc ϵ RI signaling cascade leading to degranulation may be broken into two stages: upstream and downstream. The upstream stage includes all of the processes that occur prior to calcium ion mobilization. The downstream stage includes calcium ion mobilization and all processes downstream thereof. Compounds that inhibit Fc ϵ RI-mediated degranulation may act at any point along the Fc ϵ RI-mediated signal transduction cascade. Compounds that selectively inhibit upstream Fc ϵ RI-mediated degranulation act to inhibit that portion of the Fc ϵ RI signaling cascade upstream of the point at which calcium ion mobilization is induced. In cell-based assays, compounds that selectively inhibit

upstream Fc ϵ RI-mediated degranulation inhibit degranulation of cells such as mast or basophil cells that are activated or stimulated with an IgE-specific allergen or binding agent (such as an anti-IgE antibody) but do not appreciably inhibit degranulation of cells that are activated or stimulated with degranulating agents that bypass the Fc ϵ RI signaling pathway, 5 such as, for example the calcium ionophores ionomycin and A23187.

“IgG-Induced Degranulation” or “Fc γ RI-Mediated Degranulation” refers to degranulation that proceeds *via* the Fc γ RI signal transduction cascade initiated by crosslinking of Fc γ RI-bound IgG. The crosslinking may be induced by an IgG-specific allergen or another multivalent binding agent, such as an anti-IgG or fragment antibody. 10 Like the Fc ϵ RI signaling cascade, in mast and basophil cells the Fc γ RI signaling cascade also leads to degranulation which may be broken into the same two stages: upstream and downstream. Similar to Fc ϵ RI-mediated degranulation, compounds that selectively inhibit upstream Fc γ RI-mediated degranulation act upstream of the point at which calcium ion 15 mobilization is induced. In cell-based assays, compounds that selectively inhibit upstream Fc γ RI-mediated degranulation inhibit degranulation of cells such as mast or basophil cells that are activated or stimulated with an IgG-specific allergen or binding agent (such as an anti-IgG antibody or fragment) but do not appreciably inhibit degranulation of cells that are activated or stimulated with degranulating agents that bypass the Fc γ RI signaling pathway, such as, for example the calcium ionophores ionomycin and A23187.

20 “Ionophore-Induced Degranulation” or “Ionophore-Mediated Degranulation” refers to degranulation of a cell, such as a mast or basophil cell, that occurs upon exposure to a calcium ionophore such as, for example, ionomycin or A23187.

“Syk Kinase” refers to the well-known 72kDa non-receptor (cytoplasmic) spleen protein tyrosine kinase expressed in B-cells and other hematopoietic cells. Syk kinase 25 includes two consensus Src-homology 2 (SH2) domains in tandem that bind to phosphorylated immunoreceptor tyrosine-based activation motifs (“ITAMs”), a “linker” domain and a catalytic domain (for a review of the structure and function of Syk kinase see Sada *et al.*, 2001, *J. Biochem. (Tokyo)* 130:177-186); see also Turner *et al.*, 2000, *Immunology Today* 21:148-154). Syk kinase has been extensively studied as an effector of 30 B-cell receptor (BCR) signaling (Turner *et al.*, 2000, *supra*). Syk kinase is also critical for tyrosine phosphorylation of multiple proteins which regulate important pathways leading

from immunoreceptors, such as Ca^{2+} mobilization and mitogen-activated protein kinase (MAPK) cascades and degranulation. Syk kinase also plays a critical role in integrin signaling in neutrophils (see, e.g., Mocsai *et al.* 2002, *Immunity* 16:547-558).

As used herein, Syk kinase includes kinases from any species of animal, including 5 but not limited to, homosapiens, simian, bovine, porcine, rodent, etc., recognized as belonging to the Syk family. Specifically included are isoforms, splice variants, allelic variants, mutants, both naturally occurring and man-made. The amino acid sequences of such Syk kinases are well known and available from GENBANK. Specific examples of mRNAs encoding different isoforms of human Syk kinase can be found at GENBANK 10 accession no. gi|21361552|ref|NM_003177.2|, gi|496899|emb|Z29630.1|HSSYKPTK[496899] and gi|15030258|gb|BC011399.1|BC011399[15030258], which are incorporated herein by reference.

Skilled artisans will appreciate that tyrosine kinases belonging to other families may 15 have active sites or binding pockets that are similar in three-dimensional structure to that of Syk. As a consequence of this structural similarity, such kinases, referred to herein as "Syk mimics," are expected to catalyze phosphorylation of substrates phosphorylated by Syk. Thus, it will be appreciated that such Syk mimics, signal transduction cascades in which 20 such Syk mimics play a role, and biological responses effected by such Syk mimics and Syk mimic-dependent signaling cascades may be regulated, and in particular inhibited, with many of the prodrugs described herein.

"Syk-Dependent Signaling Cascade" refers to a signal transduction cascade in which Syk kinase plays a role. Non-limiting examples of such Syk-dependent signaling cascades include the Fc α RI, Fc ϵ RI, Fc γ RI, Fc γ RIII, BCR and integrin signaling cascades.

25 "Autoimmune Disease" refers to those diseases which are commonly associated with the nonanaphylactic hypersensitivity reactions (Type II, Type III and/or Type IV hypersensitivity reactions) that generally result as a consequence of the subject's own humoral and/or cell-mediated immune response to one or more immunogenic substances of endogenous and/or exogenous origin. Such autoimmune diseases are distinguished from 30 diseases associated with the anaphylactic (Type I or IgE-mediated) hypersensitivity

reactions.


“Prodrug” refers to a derivative of an active 2,4-pyrimidinediamine compound (drug) that requires a transformation under the conditions of use, such as within the body, to release the active 2,4-pyrimidinediamine drug or an active metabolite thereof. Prodrugs are 5 frequently, but not necessarily, pharmacologically inactive until converted into the active drug. Prodrugs are typically obtained by masking one or more functional groups in the 2,4-pyrimidinediamine drug believed to be in part required for activity with a progroup (defined below) to form a promoiety which undergoes a transformation, such as cleavage, under the specified conditions of use to release the functional group, and hence the active 2,4-pyrimidinediamine drug. The cleavage of the promoiety may proceed spontaneously, such as by way of a hydrolysis reaction, or it may be catalyzed or induced by another agent, such as by an enzyme, by light, by acid or base, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature. The agent may be endogenous to the conditions of use, such as an enzyme present in the cells to which the prodrug is 10 administered or the acidic conditions of the stomach, or it may be supplied exogenously. 15

A wide variety of progroups, as well as the resultant promoieties, suitable for masking functional groups in the active 2,4-pyrimidinediamines compounds to yield prodrugs are well-known in the art. For example, a hydroxyl functional group may be masked as a sulfonate, ester or carbonate promoiety, which may be hydrolyzed *in vivo* to 20 provide the hydroxyl group. An amino functional group may be masked as an amide, carbamate, imine, urea, phosphenyl, phosphoryl or sulfenyl promoiety, which may be hydrolyzed *in vivo* to provide the amino group. A carboxyl group may be masked as an ester (including silyl esters and thioesters), amide or hydrazide promoiety, which may be hydrolyzed *in vivo* to provide the carboxyl group. Nitrogen protecting groups and nitrogen 25 pro-drugs of the invention may include lower alkyl groups as well as amides, carbamates, etc. Other specific examples of suitable progroups and their respective promoieties will be apparent to those of skill in the art.

“Progroup” refers to a type of protecting group that, when used to mask a functional group within an active 2,4-pyrimidinediamine drug to form a promoiety, converts the drug 30 into a prodrug. Progroups are typically attached to the functional group of the drug *via* bonds that are cleavable under specified conditions of use. Thus, a progroup is that portion

of a promoiety that cleaves to release the functional group under the specified conditions of use. As a specific example, an amide promoiety of the formula $-\text{NH}-\text{C}(\text{O})\text{CH}_3$ comprises the progroup $-\text{C}(\text{O})\text{CH}_3$.

Accordingly, the present invention relates to a compound according to formula I:



or a stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof, wherein:

Y is selected from the group consisting of S, O, SO, SO_2 , and $\text{C}(\text{R}^7)_2$;

each R^{35} is independently selected from the group consisting of hydrogen, (C_1-C_4)alkyl, and halo, or both R^{35} together with the carbon to which they are attached form a carbonyl group;

10 W is selected from the group consisting of $\text{C}=\text{O}$, $\text{C}=\text{S}$, $\text{C}=\text{NH}$, $\text{C}(\text{R}^7)_2$, and NR^{37} ;

Z is $\text{C}=\text{O}$ or NR^{37} , provided that Z and W are not both NR^{37} and provided that when Z is $\text{C}=\text{O}$, W is $\text{C}(\text{R}^7)_2$;

15 X is CH or N;

each R^{31} is independently (C_1-C_4)alkyl or both R^{31} together form a (C_1-C_2)alkylene group optionally substituted with one to two (C_1-C_4)alkyl groups or substituted with one (C_3-C_7) spirocycloalkyl group;

each R^7 is independently hydrogen or (C_1-C_4)alkyl; and

20 R^{37} is hydrogen or methyl optionally substituted with phenyl or pyridyl, wherein said phenyl or pyridyl is optionally substituted with (C_1-C_4)alkoxy.

In one embodiment, Y is O or S. In some aspects, Y is O.

In another embodiment, Y is S or an oxidized form of S, such as a sulfoxide SO or a sulfone SO_2 .

25 In still another embodiment, Y is $\text{C}(\text{R}^7)_2$. In some aspects, Y is $\text{C}(\text{Me})_2$.

In one embodiment, both R^{35} are the same. In some aspects, both R^{35} are methyl. In still other aspects both R^{35} are hydrogen or both are fluoro. In another aspect, both R^{35}

together with the carbon to which they are attached form a carbonyl group.

In one embodiment, W is C=O or C=S. In some aspects, W is C=O.

In another embodiment, W is C(R⁷)₂. In some aspects, W is CH₂.

In still another embodiment, W is NR³⁷. In some aspects, W is NH.


5 In one embodiment, Z is NR³⁷. In some aspects, R³⁷ is hydrogen so that Z is NH. In other aspects, R³⁷ is methyl, 2-pyridylmethyl, or 4-methoxybenzyl.

In another embodiment, Z is C=O.

In one embodiment, X is N.

10 In one embodiment each R³¹ is independently (C₁-C₂) alkyl. In some aspects, both of R³¹ are methyl.

In another embodiment, both R³¹ together form a (C₁-C₂)alkylene group optionally substituted with one to two (C₁-C₄)alkyl groups or substituted with one (C₃-C₇) spirocycloalkyl group. In some embodiments, both of R³¹ combine to form an acetal or ketal carbon. When both R³¹ together form a (C₁-C₂)alkylene group, a bicyclic fused ring system is formed with the phenyl group bearing the -OR³¹ groups. In some embodiments, the bicyclic fused ring system include the following structures wherein each R is (C₁-C₄)alkyl, or two of R combine to form a (C₃-C₇) spirocycloalkyl:



20 In still another embodiment, the present invention relates to a compound, stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof selected from the group consisting of

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine;

N4-[2,2-Dimethyl-3-oxo-benzo[1,4]thiazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;

5 N4-[2,2-Dimethyl-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;

N4-[2,2-Dimethyl-3-oxo-pyrid[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;

N4-[2,2-Difluoro-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-4-(2-pyridylmethyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;

15 N4-(3,4-Dihydro-2H-2,2-dimethyl-5-pyrido[1,4]oxazin-6-yl)-N2-[3,4-dimethoxyphenyl-5-hydroxyphenyl]-5-fluoro-2,4-pyrimidinediamine

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(1,3-(2H)-4,4-dimethylisoquinolinedione-7-yl)-2,4-pyrimidinediamine;

20 (R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;

(R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine;

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(2,2,4-trimethyl-1,1,3-trioxo-benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine; and


25 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(4-methyl-3-oxo-benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine.

Those of skill in the art will appreciate that the 3-hydroxyphenyl-2,4-pyrimidinediamine compounds described herein may include functional groups that can be masked with progroups to create prodrugs. Such prodrugs are usually, but need not be, pharmacologically inactive until converted into their active drug form. For example, ester groups commonly undergo acid-catalyzed hydrolysis to yield the parent carboxylic acid

when exposed to the acidic conditions of the stomach, or base-catalyzed hydrolysis when exposed to the basic conditions of the intestine or blood. Thus, when administered to a subject orally, 3-hydroxyphenyl-2,4-pyrimidinediamines that include ester moieties may be considered prodrugs of their corresponding carboxylic acid, regardless of whether the ester form is pharmacologically active.

In the prodrugs of the invention, any available functional moiety may be masked with a progroup to yield a prodrug. Myriad progroups suitable for masking such functional groups to yield promoietyes that are cleavable under the desired conditions of use are known in the art. All of these progroups, alone or in combinations, may be included in the prodrugs of the invention.

In one embodiment, a prodrug of the invention is a compound of formula I is a having formula II



II

15 wherein:

W is selected from the group consisting of C=O, C=S, C=NH, C(R⁷)₂, and NP¹;

Z is selected from the group consisting of C=O, NR³⁷, and NP¹; provided that when Z is C=O, then W is C(R⁷)₂ or NP¹, and provided that W and Z are not both one of NR³⁷ or NP¹;

20 Y, R³⁵, X, and R³¹ are as previously defined for formula I; and

P¹, P², P³, and P⁴ are independently hydrogen or a progroup R^P, provided that at least one of P¹, P², P³, and P⁴ is a progroup.

In some embodiments of a compound of formula II, P³ is a progroup R^P, wherein P³ and the oxygen to which it is attached form an ester, thioester, carbonate, or carbamate promoiety. In some aspects, the -OP³ promoiety is an ester.

In some embodiments of a compound of formula II, the 2-amino, 4-amino, or Z=N group is bound to a progroup R^P. In some aspects, the Z is N-R^P.

In some of the aforementioned embodiments, the progroup R^P is a phosphorous-containing progroup.

In some of the aforementioned embodiments, the progroup R^P includes a group or moiety that is metabolized under the conditions of use to yield an unstable

5 α -hydroxymethyl, α -aminomethyl or α -thiomethyl intermediate, which then further metabolized *in vivo* to yield the active 3-hydroxyphenyl-2,4- pyrimidinediamine drug. In some embodiments, the progroup includes an α -hydroxyalkyl, α -aminoalkyl or α -thioalkyl moiety, for example an α -hydroxymethyl, α -aminomethyl, α -thiomethyl moiety, that metabolizes under the conditions of use to yield the active 3-hydroxyphenyl-2,4

10 pyrimidinediamine drug. For example, in some embodiments the progroup R^P is of the formula -CR^dR^d-AR³, where each R^d is, independently of the other, selected from hydrogen, cyano, optionally substituted (C1-C20) alkyl, (C1-C20) perfluoroalkyl, optionally substituted (C7-C30) arylalkyl and optionally substituted 6-30 membered heteroarylalkyl, where each optional substituent is, independently of the others, selected from hydrogen, alkyl, aryl, arylalkyl, heteroaryl and heteroalkyl, or, alternatively, the two R^d are taken

15 together with the carbon atom to which they are bonded to form a cycloalkyl containing from 3 to 8 carbon atoms ; A is selected from O, S and NR⁵⁰, where R⁵⁰ is selected from hydrogen, alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl and cycloheteroalkyl, or alternatively is combined with R³, and, together with the nitrogen to which they are

20 attached, form a three to seven membered ring; and R³ represents a group that can be metabolized *in vivo* to yield a group of the formula -CR^dR^d-AH, where R^d and A are as previously defined.

The identity of R³ is not critical, provided that it can be metabolized under the desired conditions of use, for example under the acidic conditions found in the stomach and/or by enzymes found *in vivo*, to yield a group of the formula -CR^dR^d-AH, where A and R^d are as previously defined. Thus, skilled artisans will appreciate that R³ can comprise virtually any known or later-discovered hydroxyl, amine or thiol protecting group. Non-limiting examples of suitable protecting groups can be found, for example, in *Protective Groups in Organic Synthesis*, Greene & Wuts, 2nd Ed., John Wiley & Sons, New York, 1991 (especially pages 10-142 (alcohols, 277-308 (thiols) and 309-405 (amines) the disclosure of which is incorporated herein by reference).

In a specific embodiment, R^3 includes, together with A, an ether, a thioether, a silyl ether, a silyl thioether, an ester, a thioester, an amide, a carbonate, a thiocarbonate, a carbamate, a thiocarbamate, or a urea linkage, $-OCH_2SO_3R$, where R is hydrogen, alkyl, aryl, arylalkyl or a metal salt (e.g., sodium, lithium, potassium); $-GCH_2^+N(R^{51})_3M^-$, where G is absent, $-OPO_3^-$, OSO_3^- or $-CO_2^-$, R^{51} is hydrogen, alkyl, aryl, arylalkyl, cycloheteroalkyl or cycloheteroalkylalkyl and M- is a counterion, usually a halide ion or the like (acetate, sulfate, phosphate, etc.). Specific exemplary embodiments include, but are not limited to, progroups R^P in which R^3 is selected from R^f , $-C(O)R^f$, $-C(O)OR^f$, $-C(O)NR^fR^f$ and $-SiR^fR^fR^f$, where each R^f is, independently of the others, selected from hydrogen, optionally substituted lower alkyl, optionally substituted lower heteroalkyl, optionally substituted lower cycloalkyl, optionally substituted lower heterocycloalkyl, optionally substituted (C6-C10) aryl, optionally substituted 5-10 membered heteroaryl, optionally substituted (C7-C18) arylalkyl and optionally substituted 6-18 membered heteroarylalkyl. In a specific embodiment, each R^f is the same.

The identity of the progroup(s) R^P can be selected to tailor the water-solubility and other properties of the underlying active 3-hydroxyphenyl-2,4-pyrimidinediamine compound to be optimized for a particular mode of administration. It can also be selected to provide for removal at specified organs and/or tissues within the body, such as, for example, in the digestive tract, in blood and/or serum, or via enzymes residing in specific organs, such as the liver.

In some embodiments, progroups R^P that are phosphorous-containing progroups include phosphate moieties that can be cleaved *in vitro* by enzymes such as esterases, lipases and/or phosphatases. Such enzymes are prevalent throughout the body, residing in, for example, the stomach and digestive tract, blood and/or serum, and in virtually all tissues and organs. Such phosphate-containing progroups R^P will generally increase the water-solubility of the underlying active 3-hydroxyphenyl-2,4-pyrimidinediamine compound, making such phosphate-containing prodrugs ideally suited for modes of administration where water-solubility is desirable, such as, for example, oral, buccal, intravenous, intramuscular and ocular modes of administration.

In some embodiments, each phosphate-containing progroup R^P in the prodrug is of the formula $-(CR^dR^d)_y-O-P(O)(OH)(OH)$, or a salt thereof, wherein R^d is as previously

defined and y is an integer ranging from 1 to 3, typically 1 or 2. In one specific embodiment, each R^d is, independently of the others, selected from hydrogen, substituted or unsubstituted lower alkyl, substituted or unsubstituted phenyl, substituted or unsubstituted methyl and substituted or unsubstituted benzyl. In another specific embodiment, each R^d is, 5 independently of the others, selected from hydrogen and unsubstituted lower alkyl. Specific exemplary phosphate-containing progroups R^p include $-CH_2-O-P(O)(OH)(OH)$ and $-CH_2CH_2-O-P(O)(OH)(OH)$ and/or the corresponding salts.

While not intending to be bound by any theory of operation, when y is 1 in the exemplary phosphate-containing progroups R^p , it is believed that the phosphate-containing prodrugs are converted *in vivo* by enzymes such as phosphatases, lipases and/or esterases to the corresponding hydroxymethylamines, which are then further metabolized *in vivo* by the elimination of formaldehyde to yield the active 2,4-pyrimidinediamine drug compound. The phosphate and formaldehyde metabolic by-products are innocuous.

When y is 2 in the exemplary phosphate-containing prodrugs, it is believed that the prodrugs are metabolized to the active 3-hydroxyphenyl-2,4-pyrimidinediamine drug compound *in vivo* by elimination of enol phosphate, which further metabolizes to acetaldehyde and phosphate. The phosphate and acetaldehyde metabolic by-products are innocuous.

Skilled artisans will appreciate that certain types of precursors can be converted *in vivo* to phosphate groups. Such precursors include, by way of example and not limitation, phosphate esters, phosphites and phosphite esters. For example, phosphites can be oxidized *in vivo* to phosphates. Phosphate esters can be hydrolyzed *in vivo* to phosphates. Phosphite esters can be oxidized *in vivo* to phosphate esters, which can in turn be hydrolyzed *in vivo* to phosphates. As a consequence of the ability of these phosphate precursor groups to convert to phosphates *in vivo*, the prodrugs can also include progroups that comprise such phosphate precursors. In some embodiments, the phosphate precursor groups may be directly metabolized to the active 2,4-pyrimidinediamine drug, without first being converted into a phosphate prodrug. In other embodiments, prodrugs comprising progroups that include such phosphate precursors are first metabolized into the corresponding phosphate prodrug, 20 which then metabolizes to the active 3-hydroxyphenyl-2,4-pyrimidinediamine drug *via* a hydroxymethylamine, as discussed above.


In some embodiments, such phosphate precursor groups are phosphate esters. The phosphate esters can be acyclic or cyclic, and can be phosphate triesters or phosphate diesters. Such esters are generally less water-soluble than the corresponding phosphate acid prodrugs and the corresponding active 3-hydroxyphenyl-2,4-pyrimidinediamine 5 compounds, and are therefore typically suitable for modes of delivering prodrugs of active 3-hydroxyphenyl-2,4-pyrimidinediamine compounds where low water-solubility is desired, including, by way of example and not limitation, administration *via* inhalation. The 10 solubility of the prodrug can be specifically tailored for specific modes of administration by appropriate selection of the number and identity(ies) of the esterifying groups in the phosphate ester.

The mechanism by which the phosphate ester group metabolizes to the corresponding phosphate group can be controlled by appropriate selection of the esterifying moieties. For example, it is well-known that certain esters are acid (or base) labile, generating the corresponding phosphate under the acidic conditions found in the stomach 15 and digestive tract. In instances where it is desirable for the phosphate ester prodrug to metabolize to the corresponding phosphate prodrug in the digestive tract (such as, for example, where the prodrugs are administered orally), phosphate ester progroups that are acid-labile can be selected. Other types of phosphate esters are acid and base stable, being converted into the corresponding phosphates *via* enzymes found in certain tissues and 20 organs of the body (see, e.g., the various cyclic phosphate esters described in Erion *et al.*, 2004, J. Am. Chem. Soc. 126:5154-5163, incorporated herein by reference). In instances where it is desirable to convert a phosphate ester prodrug into the corresponding phosphate prodrug within a desired target tissue or site within the body, phosphate esters having the desired metabolic properties can be selected.


25 In some embodiments, each phosphate ester-containing progroup R^P in the prodrug is an acyclic phosphate ester of the formula -(CR^dR^d)_y-O-P(O)(OH)(OR^e) or -(CR^dR^d)_y-O-P(O)(OR^e)(OR^e), or a salt thereof, wherein each R^e is, independently of the others, selected from substituted or unsubstituted lower alkyl, substituted or unsubstituted (C6-C14) aryl (e.g., phenyl, naphthyl, 4-loweralkoxyphenyl, 4-methoxyphenyl), substituted 30 or unsubstituted (C7-C20) arylalkyl (e.g., benzyl, 1-phenylethan-1-yl, 2-phenylethan-1-yl), -(CR^dR^d)_y-OR^f, -(CR^dR^d)_y-O-C(O)R^f, -(CR^dR^d)_y-O-C(O)OR^f, -(CR^dR^d)_y-S-C(O)R^f,

$-(CR^dR^d)_y-S-C(O)OR^f$, $-(CR^dR^d)_y-NH-C(O)R^f$, $-(CR^dR^d)_y-NH-C(O)OR^f$ and $-Si(R^d)_3$, wherein R^d , R^f and y are as defined above. In a specific embodiment, each R^d is selected from hydrogen and unsubstituted lower alkyl and/or each R^e is an unsubstituted lower alkanyl or benzyl. Specific exemplary phosphate ester progroups include, but are not limited to, $-CH_2-O-P(O)(OH)(OR^e)$, $-CH_2CH_2-O-P(O)(OH)(OR^e)$, $-CH_2-O-P(O)(OR^e)(OR^e)$ and $-CH_2CH_2-O-P(O)(OR^e)(OR^e)$, where R^e is selected from lower alkanyl, *i*-propyl and *t*-butyl.

In other embodiments, each phosphate ester-containing progroup R^P is a cyclic phosphate ester of the formula




where each R^g is, independently of the others, selected from hydrogen and lower alkyl; each R^h is, independently of the others, selected from hydrogen, substituted or unsubstituted lower alkyl, substituted or unsubstituted lower cycloheteroalkyl, substituted or unsubstituted (C6-C14) aryl, substituted or unsubstituted (C7-C20) arylalkyl and substituted or unsubstituted 5-14 membered heteroaryl; z is an integer ranging from 0 to 2; and R^d and y are as previously defined. In a specific embodiment, each phosphate ester-containing progroup R^P is a cyclic phosphate ester of the formula




where R^d , R^h and y are as previously defined.

The mechanism by which cyclic phosphate ester prodrugs including such cyclic phosphate ester progroups metabolize *in vivo* to the active drug compound depends, in part, on the identity of the R^h substituent. For example, cyclic phosphate ester progroups in which each R^h is, independently of the others, selected from hydrogen and lower alkyl are cleaved *in vivo* by esterases. Thus, in some embodiments, the cyclic phosphate ester progroups are selected such that they are cleavable *in vivo* by esterases. Specific examples

of such cyclic phosphate ester progroups include, but are not limited to, progroups selected



5 Alternatively, cyclic phosphate ester prodrugs having progroups in which the R^h substituents are substituted or unsubstituted aryl, arylalkyl and heteroaryl groups, are not typically cleaved by esterases, but are instead metabolized to the active prodrug by enzymes, such as cytochrome P₄₅₀ enzymes, that reside in the liver. For example, a series of 10 cyclic phosphate ester nucleotide prodrugs that undergo an oxidative cleavage reaction catalyzed by a cytochrome P₄₅₀ enzyme (CYP) expressed predominantly in the liver are described in Erion *et al.*, 2004, J. Am. Chem. Soc. 126:5154-5163. In some embodiments, the cyclic phosphate ester progroups are selected such that they are cleavable by CYP enzymes expressed in the liver. Specific exemplary embodiments of such cyclic phosphate ester-containing progroups R^p include, but are not limited to, progroups having the formula



15 where R^h is selected from phenyl, 3-chlorophenyl, 4-pyridyl and 4-methoxyphenyl.

As skilled artisans will appreciate, phosphites and phosphite esters can undergo oxidation *in vivo* to yield the corresponding phosphate and phosphate ester analogs. Such reactions can be carried out *in vivo* by, for example, oxidase enzymes, oxoreductase 20 enzymes and other oxidative enzymes. Thus, the phosphorous-containing progroups R^p can also include phosphite and phosphite ester analogs of any of the phosphate and phosphate

ester progroups described above. In some embodiments the phosphorous-containing progroups R^p include, but are not limited to, groups of the formula -(CR^dR^d)_y-O-P(OH)(OH), -(CR^dR^d)_y-O-P(OH)(OR^e) and -(CR^dR^d)_y-O-P(OR^e)(R^e), or salts thereof, where R^d, R^e and y are as previously defined. Specific exemplary embodiments 5 include groups in which each R^d is, independently of the others, selected from hydrogen and unsubstituted lower alkyl and/or each R^e is, independently of the others, selected from unsubstituted lower alkanyl and benzyl. Specific exemplary acyclic phosphite and phosphite-ester progroups include, but are not limited to, -CH₂-O-P(OH)(OH), -CH₂CH₂-O-P(OH)(OH), -CH₂-O-P(OH)(OR^e), and -CH₂CH₂-O-P(OR^e)(OR^e), where each 10 R^e is selected from lower alkanyl, *i*-propyl and *t*-butyl. Specific exemplary cyclic phosphite ester prodrugs include phosphite analogs of the above-described cyclic phosphate ester progroups. Conceptually, prodrug compounds including such phosphite and/or phosphite ester progroups can be thought of as prodrugs of the corresponding phosphate and phosphate ester prodrugs.


15 As mentioned above, it is believed that certain phosphate-containing prodrugs metabolize *in vivo* through the corresponding hydroxymethylamines. Although these hydroxymethylamines metabolize *in vivo* to the corresponding active 3-hydroxyphenyl-2,4-pyrimidinediamine compounds, they are stable at pH 7 and can be prepared and administered as hydroxyalkyl-containing prodrugs. In some embodiments, each 20 hydroxyalkyl-containing progroup R^p of such prodrugs is of the formula -CR^dR^d-OH, where R^d is as previously defined. A specific exemplary hydroxyalkyl-containing progroup R^p is -CH₂OH.

25 In one embodiment, R^p has the formula -(CR^dR^d)_y-O-P(O)(OH)₂, or a salt thereof, where y is an integer ranging from 1 to 3; each R^d is, independently of the others, selected from hydrogen, optionally substituted lower alkyl, optionally substituted (C6-C14) aryl and 30 optionally substituted (C7-C20) arylalkyl; where the optional substituents are, independently of one another, selected from hydroxyl, lower alkoxy, (C6-C14) aryloxy, lower alkoxyalkyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl and halogen, or, alternatively, two R^d bonded to the same carbon atom are taken together with the carbon atom to which they are bonded to form a cycloalkyl group containing from 3 to 8 carbon atoms.

In one embodiment, R^p is selected from -CH₂-O-P(O)(OH)₂ and -CH₂CH₂-O-P(O)(OH)₂ and salts thereof.

In one embodiment, R^p comprises a phosphate ester group.

In one embodiment, R^p is selected from $-(CR^dR^d)y-O-P(O)(OR^e)(OH)$,
5 $-(CR^dR^d)y-O-P(O)(OR^e)(OR^e)$,



and salts thereof, wherein each R^e is, independently of the others, selected from substituted or unsubstituted lower alkyl, substituted or unsubstituted (C6-C14) aryl (e.g., phenyl, naphthyl, 4-loweralkoxyphenyl, 4-methoxyphenyl), substituted or unsubstituted (C7-C20) 10 arylalkyl (e.g., benzyl, 1-phenylethan-1-yl, 2-phenylethan-1-yl), $-(CR^dR^d)_y-OR^f$, $-(CR^dR^d)_y-O-C(O)R^f$, $-(CR^dR^d)_y-O-C(O)OR^f$, $-(CR^dR^d)_y-S-C(O)R^f$, $-(CR^dR^d)_y-S-C(O)OR^f$, $-(CR^dR^d)_y-NH-C(O)R^f$, $-(CR^dR^d)_y-NH-C(O)OR^f$ and $-Si(R^d)_3$, wherein each R^f is, independently of the others, selected from hydrogen, unsubstituted or substituted lower alkyl, substituted or unsubstituted (C6-C14) aryl, and substituted or unsubstituted (C7-C20) 15 arylalkyl; each R^g is, independently of the others, selected from hydrogen and lower alkyl; each R^h is, independently of the others, selected from hydrogen, substituted or unsubstituted lower alkyl, substituted or unsubstituted lower cycloheteroalkyl, substituted or unsubstituted (C6-C14) aryl, substituted or unsubstituted (C7-C20) arylalkyl and substituted or 20 unsubstituted 5-14 membered heteroaryl; z is an integer ranging from 0 to 2; and R^d and y are as previously defined.

In one embodiment, R^p is selected from $-CH_2-O-P(O)(OH)_2$, $-CH_2CH_2-O-P(O)(OH)_2$, $-CH_2OH$ and salts thereof. In some such aspects Z is $N-R^p$.

In another embodiment the present invention provides a composition comprising a compound of formula I or II and a carrier, excipient, or diluent.

25 Those of skill in the art will appreciate that many of the compounds and prodrugs of the invention, as well as the various compound species specifically described and/or illustrated herein, may exhibit the phenomena of tautomerism, conformational isomerism,

geometric isomerism and/or optical isomerism. For example, the compounds and prodrugs of the invention may include one or more chiral centers and/or double bonds and as a consequence may exist as stereoisomers, such as double-bond isomers (i.e., geometric isomers), enantiomers and diasteromers and mixtures thereof, such as racemic mixtures. As another example, the compounds and prodrugs of the invention may exist in several tautomeric forms, including the enol form, the keto form and mixtures thereof. As the various compound names, formulae and compound drawings within the specification and claims can represent only one of the possible tautomeric, conformational isomeric, optical isomeric or geometric isomeric forms, it should be understood that the invention encompasses any tautomeric, conformational isomeric, optical isomeric and/or geometric isomeric forms of the compounds or prodrugs having one or more of the utilities described herein, as well as mixtures of these various different isomeric forms. In cases of limited rotation around the 2,4-pyrimidinediamine core structure, atropisomers are also possible and are also specifically included in the compounds of the invention.

Moreover, skilled artisans will appreciate that when lists of alternative substituents include members which, owing to valency requirements or other reasons, cannot be used to substitute a particular group, the list is intended to be read in context to include those members of the list that are suitable for substituting the particular group. For example, skilled artisans will appreciate that while all of the listed alternatives for R^b can be used to substitute an alkyl group, certain of the alternatives, such as =O, cannot be used to substitute a phenyl group. It is to be understood that only possible combinations of substituent-group pairs are intended.

The compounds and/or prodrugs of the invention may be identified by either their chemical structure or their chemical name. When the chemical structure and the chemical name conflict, the chemical structure is determinative of the identity of the specific compound.


Depending upon the nature of the various substituents, the 2,4-pyrimidinediamine compounds and prodrugs of the invention may be in the form of salts. Such salts include salts suitable for pharmaceutical uses (“pharmaceutically-acceptable salts”), salts suitable for veterinary uses, etc. Such salts may be derived from acids or bases, as is well-known in the art.

In one embodiment, the salt is a pharmaceutically acceptable salt. Generally, pharmaceutically acceptable salts are those salts that retain substantially one or more of the desired pharmacological activities of the parent compound and which are suitable for administration to humans. Pharmaceutically acceptable salts include acid addition salts 5 formed with inorganic acids or organic acids. Inorganic acids suitable for forming pharmaceutically acceptable acid addition salts include, by way of example and not limitation, hydrohalide acids (*e.g.*, hydrochloric acid, hydrobromic acid, hydriodic, etc.), sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids suitable for forming pharmaceutically acceptable acid addition salts include, by way of example and not 10 limitation, acetic acid, trifluoroacetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, oxalic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, palmitic acid, benzoic acid, 3-(4-hydroxybenzoyl) benzoic acid, cinnamic acid, mandelic acid, alkylsulfonic acids (*e.g.*, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic 15 acid, 2-hydroxyethanesulfonic acid, etc.), arylsulfonic acids (*e.g.*, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, cycloalkylsulfonic acids (*e.g.*, camphorsulfonic acid), 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary 20 butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like.

Pharmaceutically acceptable salts also include salts formed when an acidic proton present in the parent compound is either replaced by a metal ion (*e.g.*, an alkali metal ion, an alkaline earth metal ion or an aluminum ion), an ammonium ion or coordinates with an organic base (*e.g.*, ethanolamine, diethanolamine, triethanolamine, N-methylglucamine, morpholine, piperidine, dimethylamine, diethylamine, etc.).

The 2,4-pyrimidinediamine compounds and of the invention, as well as the salts thereof, may also be in the form of hydrates, solvates and N-oxides, as are well-known in the art. In one implementation, this invention provides a compound, or stereoisomer, tautomer, prodrug, solvate, or pharmaceutically acceptable salt thereof, selected from **Table 30 I.**

Table I



| cmpd# | Y | R ³⁵ | W | Z | X | R ³¹ |
|---------|----------------------------------|-----------------|-----------------|-----------------------|----|-----------------|
| R909384 | S | H,H | C=O | NH | CH | Me,Me |
| R909385 | S | Me,Me | C=O | NH | CH | Me,Me |
| R909390 | O | Me,Me | C=O | NH | CH | Me,Me |
| R909391 | O | Me,Me | C=O | NH | N | Me,Me |
| R909402 | O | H,H | C=O | NH | CH | Me,Me |
| R909403 | O | △ | C=O | NH | CH | Me,Me |
| R909404 | O | F,F | C=O | NH | CH | Me,Me |
| R909406 | O | H,H | C=O | | CH | Me,Me |
| R909407 | O | H,H, | C=O | N-CH ₂ C≡N | CH | Me,Me |
| R909408 | S | H,H, | C=O | N-CH ₂ C≡N | CH | Me,Me |
| R935879 | O | H,H | CH ₂ | NH | N | Me,Me |
| R909414 | C(CH ₃) ₂ | =O | NH | C=O | CH | Me,Me |
| R909415 | O | Me,H | C=O | | CH | Me,Me |
| R909416 | S | Me,H | C=O | | CH | Me,Me |
| R909417 | SO ₂ | Me,Me | C=O | N-Me | CH | Me,Me |
| R909418 | S | H,H | C=O | N-Me | CH | Me,Me |


Methods of Synthesis

The compounds and prodrugs of the invention may be synthesized *via* a variety of different synthetic routes using commercially available starting materials and/or starting materials prepared by conventional synthetic methods. Suitable exemplary methods that 5 may be routinely adapted to synthesize the 2,4-pyrimidinediamine compounds and prodrugs of the invention are found in U.S. Patent No. 5,958,935, U.S. Patent Application 10/355,543, filed January 31, 2003 (US Publication US20040029902-A1), WO 03/063794, published August 1, 2003, U.S. Patent Application 10/631,029, filed July 29, 2003 and WO 2004/014382, published February 19, 2004, and U.S. Patent Application Serial 10 No.10/903,870 filed July 30, 2004 the disclosures of which are incorporated herein by reference. All of the compounds of structural formulae (I) and (II) may be prepared by routine adaptation of these methods.

A variety of exemplary synthetic routes that can be used to synthesize the 3-hydroxy-2,4-pyrimidinediamine compounds of the invention are described in Schemes (I)-15 (II), below. These methods may be routinely adapted to synthesize the prodrugs according to structural formulas (III) and (IV).

In one exemplary embodiment, the compounds can be synthesized from substituted or unsubstituted uracils or thiouracils as illustrated in Scheme (I), below:

Scheme (I)



20

In Scheme (I), R³⁵, R³¹, Y, W, Z, and X are as previously defined for structural

formula (I), X' is a halogen (e.g., F, Cl, Br or I) and Q and Q' are each, independently of one another, selected from the group consisting of O and S. Referring to Scheme (I), uracil or thiouracil **2** is dihalogenated at the 2- and 4-positions using standard halogenating agent POX_3 (or other standard halogenating agent) under standard conditions to yield 2,4-bishalo pyrimidine **4**. Typically, in pyrimidine **4**, the halide at the C4 position is more reactive towards nucleophiles than the halide at the C2 position. This differential reactivity can be exploited to synthesize 2,4-pyrimidinediamines according structural formula (I) by first reacting 2,4-bishalopyrimidine **4** with one equivalent of amine **10**, yielding 4N-substituted-2-halo-4-pyrimidineamine **8**, followed by amine **6** to yield a 2,4-pyrimidinediamine according structural formula (I).

Typically, the C4 halide is more reactive towards nucleophiles, as illustrated in the Scheme. However, as will be recognized by skilled artisans, the regioselectivity of the reaction can be controlled by adjusting the solvent and other synthetic conditions (such as temperature), as is well-known in the art.

The reactions depicted in Scheme (I) may proceed more quickly when the reaction mixtures are heated *via* microwave. When heating in this fashion, the following conditions may be used: heat to 175°C - 185°C in ethanol for 5-60 minutes in a Smith Reactor (Personal Chemistry) in a sealed tube (at 20 bar pressure).

The uracil or thiouracil **2** starting materials may be purchased from commercial sources or prepared using standard techniques of organic chemistry. Commercially available uracils and thiouracils that can be used as starting materials in Scheme (I) include, by way of example and not limitation, uracil (Aldrich #13,078-8; CAS Registry 66-22-8); 2-thio-uracil (Aldrich #11,558-4; CAS Registry 141-90-2); 2,4-dithiouracil (Aldrich #15,846-1; CAS Registry 2001-93-6); 5-acetouracil (Chem. Sources Int'l 2000; CAS Registry 6214-65-9); 5-azidouracil; 5-aminouracil (Aldrich #85,528-6; CAS Registry 932-52-5); 5-bromouracil (Aldrich #85,247-3; CAS Registry 51-20-7); 5-(trans-2-bromovinyl)-uracil (Aldrich #45,744-2; CAS Registry 69304-49-0); 5-(trans-2-chlorovinyl)-uracil (CAS Registry 81751-48-2); 5-(trans-2-carboxyvinyl)-uracil; uracil-5-carboxylic acid (2,4-dihydroxypyrimidine-5-carboxylic acid hydrate; Aldrich #27,770-3; CAS Registry 23945-44-0); 5-chlorouracil (Aldrich #22,458-8; CAS Registry 1820-81-1); 5-cyanouracil (Chem. Sources Int'l 2000; CAS Registry 4425-56-3); 5-ethyluracil (Aldrich #23,044-8;

CAS Registry 4212-49-1); 5-ethenyluracil (CAS Registry 37107-81-6); 5-fluorouracil (Aldrich #85,847-1; CAS Registry 51-21-8); 5-iodouracil (Aldrich #85,785-8; CAS Registry 696-07-1); 5-methyluracil (thymine; Aldrich #13,199-7; CAS Registry 65-71-4); 5-nitrouracil (Aldrich #85,276-7; CAS Registry 611-08-5); uracil-5-sulfamic acid (Chem. 5 Sources Int'l 2000; CAS Registry 5435-16-5); 5-(trifluoromethyl)-uracil (Aldrich #22,327-1; CAS Registry 54-20-6); 5-(2,2,2-trifluoroethyl)-uracil (CAS Registry 155143-31-6); 5-(pentafluoroethyl)-uracil (CAS Registry 60007-38-3); 6-aminouracil (Aldrich #A5060-6; CAS Registry 873-83-6) uracil-6-carboxylic acid (orotic acid; Aldrich #0-840-2; CAS Registry 50887-69-9); 6-methyluracil (Aldrich #D11,520-7; CAS Registry 626-48-2); 10 uracil-5-amino-6-carboxylic acid (5-aminoorotic acid; Aldrich #19,121-3; CAS Registry #7164-43-4); 6-amino-5-nitrosouracil (6-amino-2,4-dihydroxy-5-nitrosopyrimidine; Aldrich #27,689-8; CAS Registry 5442-24-0); uracil-5-fluoro-6-carboxylic acid (5-fluoroorotic acid; Aldrich #42,513-3; CAS Registry 00000-00-0); and uracil-5-nitro-6-carboxylic acid (5-nitroorotic acid; Aldrich #18,528-0; CAS Registry 600779-49-9). Additional 5-, 6- and 15 5,6-substituted uracils and/or thiouracils are available from General Intermediates of Canada, Inc., Edmonton, Alberta, CA (www.generalintermediates.com) and/or Interchim, France (www.interchim.com), or may be prepared using standard techniques. Myriad textbook references teaching suitable synthetic methods are provided *infra*.


Amines **6** and **10** may be purchased from commercial sources or, alternatively, may 20 be synthesized utilizing standard techniques. For example, suitable amines may be synthesized from nitro precursors using standard chemistry. Specific exemplary reactions are provided in the Examples section. See also Vogel, 1989, *Practical Organic Chemistry*, Addison Wesley Longman, Ltd. and John Wiley & Sons, Inc.

Skilled artisans will recognize that in some instances, amines **6** (such as the 3-hydroxy moiety) and **10** and/or other substituents on uracil or thiouracil **2**, may include 25 functional groups that require protection during synthesis. The exact identity of any protecting group(s) used will depend upon the identity of the functional group being protected, and will be apparent to those of skill in the art. Guidance for selecting appropriate protecting groups, as well as synthetic strategies for their attachment and 30 removal, may be found, for example, in Greene & Wuts, *Protective Groups in Organic Synthesis*, 3d Edition, John Wiley & Sons, Inc., New York (1999) and the references cited

therein (hereinafter “Greene & Wuts”).

A specific embodiment of Scheme (I) utilizing 5-fluorouracil (Aldrich #32,937-1) as a starting material is illustrated in Scheme (II), below:


Scheme (II)



In Scheme (II), Y, Z, and X are as previously defined for Scheme (I). According to Scheme (II), 5-fluorouracil **3** is halogenated with POCl_3 to yield 2,4-dichloro-5-fluoropyrimidine **5**, which is then reacted with one equivalent of amine **10'** (to yield 2-chloro-N4-substituted-5-fluoro-4-pyrimidineamine **8'**) followed by one or more equivalents of amine **6** to give compounds of formula (II).

Prodrugs according to structural formula (II) may be prepared by routine modification of the above-described methods. Alternatively, such prodrugs may be prepared by reacting a suitably protected 2,4-pyrimidinediamine of structural formula (I) with a suitable progroup. Conditions for carrying out such reactions and for deprotecting the product to yield a prodrug of formulas (III) and (IV) are well-known, and include, for example, those shown in U.S. Patent Application Publication No. 2006-0211657, PCT Publication WO 2006/078846 and in U.S. application Serial No. 11/295,752, filed 6 December 2005, each of which is hereby incorporated by reference in its entirety.

In another embodiment, a prodrug of the formula V



V

wherein Y, Z, and X are as defined for formula I and P³ and the oxygen atom to which it is attached form an ester moiety, is prepared by reacting a 3-hydroxy phenyl compound of formula V wherein P³ is hydrogen with an appropriate acid halide or anhydride and optionally in the presence of a base such as an amine. In another embodiment the compound is reacted with an appropriate acid in the presence of an acid catalyst or a coupling reagent. In some embodiments, the acid catalyst is sulfuric acid or HCl. In other embodiments the coupling reagent is a carbodiimide such as dicyclohexylcarbodiimide or is 1,1'-carbonyldiimidazole.

Myriad references teaching methods useful for synthesizing pyrimidines generally, as well as starting materials described in Schemes (I)-(II), are known in the art. For specific guidance, the reader is referred to Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16* (Weissberger, A., Ed.), 1962, Interscience Publishers, (A Division of John Wiley & Sons), New York ("Brown I"); Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16, Supplement I* (Weissberger, A. and Taylor, E. C., Ed.), 1970, Wiley-Interscience, (A Division of John Wiley & Sons), New York ("Brown II"); Brown, D. J., "The Pyrimidines", in *The Chemistry of Heterocyclic Compounds, Volume 16, Supplement II* (Weissberger, A. and Taylor, E. C., Ed.), 1985, An Interscience Publication (John Wiley & Sons), New York ("Brown III"); Brown, D. J., "The Pyrimidines" in *The Chemistry of Heterocyclic Compounds, Volume 52* (Weissberger, A. and Taylor, E. C., Ed.), 1994, John Wiley & Sons, Inc., New York, pp. 1-1509 ("Brown IV"); Kenner, G. W. and Todd, A., in *Heterocyclic Compounds, Volume 6*, (Elderfield, R. C., Ed.), 1957, John Wiley, New York, Chapter 7 (pyrimidines); Paquette, L. A., *Principles of Modern Heterocyclic Chemistry*, 1968, W. A. Benjamin, Inc., New York, pp. 1 – 401 (uracil synthesis pp. 313, 315;

pyrimidine synthesis pp. 313-316; amino pyrimidine synthesis pp. 315); Joule, J. A., Mills, K. and Smith, G. F., *Heterocyclic Chemistry*, 3rd Edition, 1995, Chapman and Hall, London, UK, pp. 1 – 516; Vorbrüggen, H. and Ruh-Pohlenz, C., *Handbook of Nucleoside Synthesis*, John Wiley & Sons, New York, 2001, pp. 1-631 (protection of pyrimidines by acylation pp. 5 90-91; silylation of pyrimidines pp. 91-93); Joule, J. A., Mills, K. and Smith, G. F., *Heterocyclic Chemistry*, 4th Edition, 2000, Blackwell Science, Ltd, Oxford, UK, pp. 1 – 589; and *Comprehensive Organic Synthesis*, Volumes 1-9 (Trost, B. M. and Fleming, I., Ed.), 1991, Pergamon Press, Oxford, UK.

Inhibition of Fc Receptor Signal Cascades

10 Active 2,4-pyrimidinediamine compounds of the invention inhibit Fc receptor signalling cascades that lead to, among other things, degranulation of cells. As a specific example, the compounds inhibit the Fc ϵ RI and/or Fc γ RI signal cascades that lead to degranulation of immune cells such as neutrophil, eosinophil, mast and/or basophil cells. Both mast and basophil cells play a central role in allergen-induced disorders, including, for 15 example, allergic rhinitis and asthma. Upon exposure allergens, which may be, among other things, pollen or parasites, allergen-specific IgE antibodies are synthesized by B-cells activated by IL-4 (or IL-13) and other messengers to switch to IgE class specific antibody synthesis. These allergen-specific IgEs bind to the high affinity Fc ϵ RI. Upon binding of antigen, the Fc ϵ RI-bound IgEs are cross-linked and the IgE receptor signal transduction 20 pathway is activated, which leads to degranulation of the cells and consequent release and/or synthesis of a host of chemical mediators, including histamine, proteases (e.g., tryptase and chymase), lipid mediators such as leukotrienes (e.g., LTC4), platelet-activating factor (PAF) and prostaglandins (e.g., PGD2) and a series of cytokines, including TNF- α , IL-4, IL-13, IL-5, IL-6, IL-8, GMCSF, VEGF and TGF- β . The release and/or synthesis of 25 these mediators from mast and/or basophil cells accounts for the early and late stage responses induced by allergens, and is directly linked to downstream events that lead to a sustained inflammatory state.

The molecular events in the Fc ϵ RI signal transduction pathway that lead to release of preformed mediators *via* degranulation and release and/or synthesis of other chemical 30 mediators are well-known. Fc ϵ RI is a heterotetrameric receptor composed of an IgE-

binding alpha-subunit, a beta subunit, and two gamma subunits (gamma homodimer). Cross-linking of Fc ϵ RI-bound IgE by multivalent binding agents (including, for example IgE-specific allergens or anti-IgE antibodies or fragments) induces the rapid association and activation of the Src-related kinase Lyn. Lyn phosphorylates immunoreceptor tyrosine-based activation motifs (ITAMS) on the intracellular beta and gamma subunits, which leads to the recruitment of additional Lyn to the beta subunit and Syk kinase to the gamma homodimer. These receptor-associated kinases, which are activated by intra- and intermolecular phosphorylation, phosphorylate other components of the pathway, such as the Btk kinase, LAT, and phospholipase C-gamma PLC-gamma. Activated PLC-gamma initiates pathways that lead to protein kinase C activation and Ca^{2+} mobilization, both of which are required for degranulation. Fc ϵ R1 cross-linking also activates the three major classes of mitogen activated protein (MAP) kinases, *i.e.* ERK1/2, JNK1/2, and p38. Activation of these pathways is important in the transcriptional regulation of proinflammatory mediators, such as TNF- α and IL-6, as well as the lipid mediator leukotriene CA (LTC4).

Although not illustrated, the Fc γ RI signaling cascade is believed to share some common elements with the Fc ϵ RI signaling cascade. Importantly, like Fc ϵ RI, the Fc γ RI includes a gamma homodimer that is phosphorylated and recruits Syk, and like Fc ϵ RI, activation of the Fc γ RI signaling cascade leads to, among other things, degranulation. Other Fc receptors that share the gamma homodimer, and which can be regulated by the active 2,4-pyrimidinediamine compounds include, but are not limited to, Fc α RI and Fc γ RIII.

The ability of the 2,4-pyrimidinediamine compounds of the invention to inhibit Fc receptor signaling cascades may be simply determined or confirmed in *in vitro* assays. Suitable assays for confirming inhibition of Fc ϵ RI-mediated degranulation are provided in the Examples section. In one typical assay, cells capable of undergoing Fc ϵ RI-mediated degranulation, such as mast or basophil cells, are first grown in the presence of IL-4, Stem Cell Factor (SCF), IL-6 and IgE to increase expression of the Fc ϵ RI, exposed to a 2,4-pyrimidinediamine test compound of the invention and stimulated with anti-IgE antibodies (or, alternatively, an IgE-specific allergen). Following incubation, the amount of a chemical mediator or other chemical agent released and/or synthesized as a consequence of activating

the Fc ϵ RI signaling cascade may be quantified using standard techniques and compared to the amount of the mediator or agent released from control cells (*i.e.*, cells that are stimulated but that are not exposed to test compound). The concentration of test compound that yields a 50% reduction in the quantity of the mediator or agent measured as compared to control 5 cells is the IC₅₀ of the test compound. The origin of the mast or basophil cells used in the assay will depend, in part, on the desired use for the compounds and will be apparent to those of skill in the art. For example, if the compounds will be used to treat or prevent a particular disease in humans, a convenient source of mast or basophil cells is a human or other animal which constitutes an accepted or known clinical model for the particular 10 disease. Thus, depending upon the particular application, the mast or basophil cells may be derived from a wide variety of animal sources, ranging from, for example, lower mammals such as mice and rats, to dogs, sheep and other mammals commonly employed in clinical testing, to higher mammals such as monkeys, chimpanzees and apes, to humans. Specific examples of cells suitable for carrying out the *in vitro* assays include, but are not limited to, 15 rodent or human basophil cells, rat basophil leukemia cell lines, primary mouse mast cells (such as bone marrow-derived mouse mast cells “BMMC”) and primary human mast cells isolated from cord blood (“CHMC”) or other tissues such as lung. Methods for isolating and culturing these cell types are well-known or are provided in the Examples section (*see, e.g.*, Demo *et al.*, 1999, Cytometry 36(4):340-348 and copending application Serial No. 20 10/053,355, filed November 8, 2001, the disclosures of which are incorporated herein by reference). Of course, other types of immune cells that degranulate upon activation of the Fc ϵ RI signaling cascade may also be used, including, for example, eosinophils.

As will be recognized by skilled artisans, the mediator or agent quantified is not critical. The only requirement is that it be a mediator or agent released and/or synthesized 25 as a consequence of initiating or activating the Fc receptor signaling cascade. For example, activation of the Fc ϵ RI signaling cascade in mast and/or basophil cells leads to numerous downstream events. For example, activation of the Fc ϵ RI signal cascade leads to the immediate release (*i.e.*, within 1-3 min. following receptor activation) of a variety of preformed chemical mediators and agents *via* degranulation. Thus, in one embodiment, the 30 mediator or agent quantified may be specific to granules (*i.e.*, present in granules but not in the cell cytoplasm generally). Examples of granule-specific mediators or agents that can be

quantified to determine and/or confirm the activity of a 2,4-pyrimidinediamine compound of the invention include, but are not limited to, granule-specific enzymes such as hexosaminidase and tryptase and granule-specific components such as histamine and serotonin. Assays for quantifying such factors are well-known, and in many instances are 5 commercially available. For example, tryptase and/or hexosaminidase release may be quantified by incubating the cells with cleavable substrates that fluoresce upon cleavage and quantifying the amount of fluorescence produced using conventional techniques. Such cleavable fluorogenic substrates are commercially available. For example, the fluorogenic substrates Z-Gly-Pro-Arg-AMC (Z=benzyloxycarbonyl; AMC=7-amino-4-methylcoumarin; 10 BIOMOL Research Laboratories, Inc., Plymouth Meeting, PA 19462, Catalog No. P-142) and Z-Ala-Lys-Arg-AMC (Enzyme Systems Products, a division of ICN Biomedicals, Inc., Livermore, CA 94550, Catalog No. AMC-246) can be used to quantify the amount of tryptase released. The fluorogenic substrate 4-methylumbelliferyl-N-acetyl- β -D-glucosaminide (Sigma, St. Louis, MO, Catalog #69585) can be used to quantify the amount 15 of hexosaminidase released. Histamine release may be quantified using a commercially available enzyme-linked immunosorbent assay (ELISA) such as Immunotech histamine ELISA assay #IM2015 (Beckman-Coulter, Inc.). Specific methods of quantifying the release of tryptase, hexosaminidase and histamine are provided in the Examples section. Any of these assays may be used to determine or confirm the activity of the 2,4- 20 pyrimidinediamine compounds of the invention.

Degranulation is only one of several responses initiated by the Fc ϵ RI signaling cascade. In addition, activation of this signaling pathway leads to the *de novo* synthesis and release of cytokines and chemokines such as IL-4, IL-5, IL-6, TNF- α , IL-13 and MIP1- α , and release of lipid mediators such as leukotrienes (e.g., LTC4), platelet activating factor 25 (PAF) and prostaglandins. Accordingly, the 2,4-pyrimidinediamine compounds of the invention may also be assessed for activity by quantifying the amount of one or more of these mediators released and/or synthesized by activated cells.

Unlike the granule-specific components discussed above, these “late stage” mediators are not released immediately following activation of the Fc ϵ RI signaling cascade. 30 Accordingly, when quantifying these late stage mediators, care should be taken to insure that the activated cell culture is incubated for a time sufficient to result in the synthesis (if

necessary) and release of the mediator being quantified. Generally, PAF and lipid mediators such as leukotriene C4 are released 3-30 min. following Fc ϵ RI activation. The cytokines and other late stage mediators are released approx. 4-8 hrs. following Fc ϵ RI activation. Incubation times suitable for a specific mediator will be apparent to those of skill in the art. Specific guidance and assays are provided in the Examples section.

The amount of a particular late stage mediator released may be quantified using any standard technique. In one embodiment, the amount(s) may be quantified using ELISA assays. ELISA assay kits suitable for quantifying the amount of TNF α , IL-4, IL-5, IL-6 and/or IL-13 released are available from, for example, Biosource International, Inc., 10 Camarillo, CA 93012 (see, e.g., Catalog Nos. KHC3011, KHC0042, KHC0052, KHC0061 and KHC0132). ELISA assay kits suitable for quantifying the amount of leukotriene C4 (LTC4) released from cells are available from Cayman Chemical Co., Ann Arbor, MI 48108 (see, e.g., Catalog No. 520211).

Typically, active 2,4-pyrimidinediamine compounds of the invention will exhibit IC₅₀s with respect to Fc ϵ RI-mediated degranulation and/or mediator release or synthesis of about 20 μ M or lower, as measured in an *in vitro* assay, such as one of the *in vitro* assays described above or in the Examples section. Of course, skilled artisans will appreciate that compounds which exhibit lower IC₅₀s, for example on the order of 10 μ M, 1 μ M, 100 nM, 10 nM, 1 nM, or even lower, are particularly useful.

Skilled artisans will also appreciate that the various mediators discussed above may induce different adverse effects or exhibit different potencies with respect to the same adverse effect. For example, the lipid mediator LTC4 is a potent vasoconstrictor – it is approximately 1000-fold more potent at inducing vasoconstriction than histamine. As another example, in addition to mediating atopic or Type I hypersensitivity reactions, 25 cytokines can also cause tissue remodeling and cell proliferation. Thus, although compounds that inhibit release and/or synthesis of any one of the previously discussed chemical mediators are useful, skilled artisans will appreciate that compounds which inhibit the release and/or synthesis of a plurality, or even all, of the previously described mediators find particular use, as such compounds are useful for ameliorating or avoiding altogether a 30 plurality, or even all, of the adverse effects induced by the particular mediators. For

example, compounds which inhibit the release of all three types of mediators—granule-specific, lipid and cytokine—are useful for treating or preventing immediate Type I hypersensitivity reactions as well as the chronic symptoms associated therewith.

Compounds of the invention capable of inhibiting the release of more than one type of mediator (e.g., granule-specific or late stage) may be identified by determining the IC₅₀ with respect to a mediator representative of each class using the various *in vitro* assays described above (or other equivalent *in vitro* assays). Compounds of the invention which are capable of inhibiting the release of more than one mediator type will typically exhibit an IC₅₀ for each mediator type tested of less than about 20 μ M. For example, a compound which exhibits an IC₅₀ of 1 μ M with respect to histamine release (IC₅₀^{histamine}) and an IC₅₀ of 1 nM with respect to leukotriene LTC4 synthesis and/or release (IC₅₀^{LTC4}) inhibits both immediate (granule-specific) and late stage mediator release. As another specific example, a compound that exhibits an IC₅₀^{tryptase} of 10 μ M, an IC₅₀^{LTC4} of 1 μ M and an IC₅₀^{IL-4} of 1 μ M inhibits immediate (granule-specific), lipid and cytokine mediator release. Although the above specific examples utilize the IC₅₀s of one representative mediator of each class, skilled artisans will appreciate that the IC₅₀s of a plurality, or even all, mediators comprising one or more of the classes may be obtained. The quantity(ies) and identity(ies) of mediators for which IC₅₀ data should be ascertained for a particular compound and application will be apparent to those of skill in the art.

Similar assays may be utilized to confirm inhibition of signal transduction cascades initiated by other Fc receptors, such as Fc α RI, Fc γ RI and/or Fc γ RIII signaling, with routine modification. For example, the ability of the compounds to inhibit Fc γ RI signal transduction may be confirmed in assays similar to those described above, with the exception that the Fc γ RI signaling cascade is activated, for example by incubating the cells with IgG and an IgG-specific allergen or antibody, instead of IgE and an IgE-specific allergen or antibody. Suitable cell types, activating agents and agents to quantify to confirm inhibition of other Fc receptors, such as Fc receptors that comprise a gamma homodimer, will be apparent to those of skill in the art.

One particularly useful class of compounds includes those 2,4-pyrimidinediamine compounds that inhibit the release of immediate granule-specific mediators and late stage

mediators with approximately equivalent IC₅₀s. By approximately equivalent is meant that the IC₅₀s for each mediator type are within about a 10-fold range of one another. Another particularly useful class of compounds includes those 2,4-pyrimidinediamine compounds that inhibit the release of immediate granule-specific mediators, lipid mediators and 5 cytokine mediators with approximately equivalent IC₅₀s. In a specific embodiment, such compounds inhibit the release of the following mediators with approximately equivalent IC₅₀s: histamine, tryptase, hexosaminidase, IL-4, IL-5, IL-6, IL-13, TNF α and LTC4. Such compounds are particularly useful for, among other things, ameliorating or avoiding altogether both the early and late stage responses associated with atopic or immediate Type 10 I hypersensitivity reactions.

Ideally, the ability to inhibit the release of all desired types of mediators will reside in a single compound. However, mixtures of compounds can also be identified that achieve the same result. For example, a first compound which inhibits the release of granule specific mediators may be used in combination with a second compound which inhibits the 15 release and/or synthesis of cytokine mediators.

In addition to the Fc ϵ RI or Fc γ RI degranulation pathways discussed above, degranulation of mast and/or basophil cells can be induced by other agents. For example, ionomycin, a calcium ionophore that bypasses the early Fc ϵ RI or Fc γ RI signal transduction machinery of the cell, directly induces a calcium flux that triggers degranulation. Activated 20 PLC γ initiates pathways that lead to, among other things, calcium ion mobilization and subsequent degranulation. As illustrated, this Ca²⁺ mobilization is triggered late in the Fc ϵ RI signal transduction pathway. As mentioned above, ionomycin directly induces Ca²⁺ mobilization and a Ca²⁺ flux that leads to degranulation. Other ionophores that induce degranulation in this manner include A23187. The ability of granulation-inducing 25 ionophores such as ionomycin to bypass the early stages of the Fc ϵ RI and/or Fc γ RI signaling cascades may be used as a counter screen to identify active compounds of the invention that specifically exert their degranulation-inhibitory activity by blocking or inhibiting the early Fc ϵ RI or Fc γ RI signaling cascades, as discussed above. Compounds which specifically inhibit such early Fc ϵ RI or Fc γ RI-mediated degranulation inhibit not 30 only degranulation and subsequent rapid release of histamine, tryptase and other granule contents, but also inhibit the pro-inflammatory activation pathways causing the release of

TNF α , IL-4, IL-13 and the lipid mediators such as LTC4. Thus, compounds which specifically inhibit such early Fc ϵ RI and/or Fc γ RI-mediated degranulation block or inhibit not only acute atopic or Type I hypersensitivity reactions, but also late responses involving multiple inflammatory mediators.

5 Compounds of the invention that specifically inhibit early Fc ϵ RI and/or Fc γ RI-mediated degranulation are those compounds that inhibit Fc ϵ RI and/or Fc γ RI-mediated degranulation (for example, have an IC₅₀ of less than about 20 μ M with respect to the release of a granule-specific mediator or component as measured in an *in vitro* assay with cells stimulated with an IgE or IgG binding agent) but that do not appreciably inhibit
10 ionophore-induced degranulation. In one embodiment, compounds are considered to not appreciably inhibit ionophore-induced degranulation if they exhibit an IC₅₀ of ionophore-induced degranulation of greater than about 20 μ M, as measured in an *in vitro* assay. Of course, active compounds that exhibit even higher IC₅₀s of ionophore-induced degranulation, or that do not inhibit ionophore-induced degranulation at all, are particularly
15 useful. In another embodiment, compounds are considered to not appreciably inhibit ionophore-induced degranulation if they exhibit a greater than 10-fold difference in their IC₅₀s of Fc ϵ RI and/or Fc γ RI-mediated degranulation and ionophore-induced degranulation, as measured in an *in vitro* assay. Assays suitable for determining the IC₅₀ of ionophore-induced degranulation include any of the previously-described degranulation assays, with
20 the modification that the cells are stimulated or activated with a degranulation-inducing calcium ionophore such as ionomycin or A23187 (A.G. Scientific, San Diego, CA) instead of anti-IgE antibodies or an IgE-specific allergen. Specific assays for assessing the ability of a particular 2,4-pyrimidinediamine compound of the invention to inhibit ionophore-induced degranulation are provided in the Examples section.

25 As will be recognized by skilled artisans, compounds which exhibit a high degree of selectivity of Fc ϵ RI-mediated degranulation find particular use, as such compounds selectively target the Fc ϵ RI cascade and do not interfere with other degranulation mechanisms. Similarly, compounds which exhibit a high degree of selectivity of Fc γ RI-mediated degranulation find particular use, as such compounds selectively target the Fc γ RI cascade and do not interfere with other degranulation mechanisms. Compounds which
30 exhibit a high degree of selectivity are generally 10-fold or more selective for Fc ϵ RI- or

Fc γ RI -mediated degranulation over ionophore-induced degranulation, such as ionomycin-induced degranulation.

Accordingly, the activity of the 2,4-pyrimidinediamine compounds of the invention may also be confirmed in biochemical or cellular assays of Syk kinase activity. In the Fc ϵ RI signaling cascade in mast and/or basophil cells, Syk kinase phosphorylates LAT and PLC-gamma1, which leads to, among other things, degranulation. Any of these activities may be used to confirm the activity of the 2,4-pyrimidinediamine compounds of the invention. In one embodiment, the activity is confirmed by contacting an isolated Syk kinase, or an active fragment thereof with a 2,4-pyrimidinediamine compound in the presence of a Syk kinase substrate (e.g., a synthetic peptide or a protein that is known to be phosphorylated by Syk in a signaling cascade) and assessing whether the Syk kinase phosphorylated the substrate. Alternatively, the assay may be carried out with cells that express a Syk kinase. The cells may express the Syk kinase endogenously or they may be engineered to express a recombinant Syk kinase. The cells may optionally also express the Syk kinase substrate. Cells suitable for performing such confirmation assays, as well as methods of engineering suitable cells will be apparent to those of skill in the art. Specific examples of biochemical and cellular assays suitable for confirming the activity of the 2,4-pyrimidinediamine compounds are provided in the Examples section.

Generally, compounds that are Syk kinase inhibitors will exhibit an IC₅₀ with respect to a Syk kinase activity, such as the ability of Syk kinase to phosphorylate a synthetic or endogenous substrate, in an *in vitro* or cellular assay in the range of about 20 μ M or less. Skilled artisans will appreciate that compounds that exhibit lower IC₅₀s, such as in the range of 10 μ M, 1 μ M, 100 nM, 10 nM, 1 nM, or even lower, are particularly useful.

Uses and Compositions

As previously discussed, the active compounds of the invention inhibit Fc receptor signaling cascades, especially those Fc receptors including a gamma homodimer, such as the Fc ϵ RI and/or Fc γ RI signaling cascades, that lead to, among other things, the release and/or synthesis of chemical mediators from cells, either *via* degranulation or other processes. As also discussed, the active compounds are also potent inhibitors of Syk kinase. As a consequence of these activities, the active compounds of the invention may be used in

a variety of *in vitro*, *in vivo* and *ex vivo* contexts to regulate or inhibit Syk kinase, signaling cascades in which Syk kinase plays a role, Fc receptor signaling cascades, and the biological responses effected by such signaling cascades. For example, in one embodiment, the compounds may be used to inhibit Syk kinase, either *in vitro* or *in vivo*, in virtually any 5 cell type expressing Syk kinase. They may also be used to regulate signal transduction cascades in which Syk kinase plays a role. Such Syk-dependent signal transduction cascades include, but are not limited to, the Fc ϵ RI, Fc γ RI, Fc γ RIII, BCR and integrin signal transduction cascades. The compounds may also be used *in vitro* or *in vivo* to regulate, and in particular inhibit, cellular or biological responses effected by such Syk-dependent signal 10 transduction cascades. Such cellular or biological responses include, but are not limited to, respiratory burst, cellular adhesion, cellular degranulation, cell spreading, cell migration, cell aggregation, phagocytosis, cytokine synthesis and release, cell maturation and Ca²⁺ flux. Importantly, the compounds may be used to inhibit Syk kinase *in vivo* as a therapeutic 15 approach towards the treatment or prevention of diseases mediated, either wholly or in part, by a Syk kinase activity. Non-limiting examples of Syk kinase mediated diseases that may be treated or prevented with the compounds are those discussed in more detail, below.

In another embodiment, the active compounds may be used to regulate or inhibit the Fc receptor signaling cascades and/or Fc ϵ RI- and/or Fc γ RI-mediated degranulation as a therapeutic approach towards the treatment or prevention of diseases characterized by, 20 caused by and/or associated with the release or synthesis of chemical mediators of such Fc receptor signaling cascades or degranulation. Such treatments may be administered to animals in veterinary contexts or to humans. Diseases that are characterized by, caused by or associated with such mediator release, synthesis or degranulation, and that can therefore be treated or prevented with the active compounds include, by way of example and not 25 limitation, atopy or anaphylactic hypersensitivity or allergic reactions, allergies (e.g., allergic conjunctivitis, allergic rhinitis, atopic asthma, atopic dermatitis and food allergies), low grade scarring (e.g., of scleroderma, increased fibrosis, keloids, post-surgical scars, pulmonary fibrosis, vascular spasms, migraine, reperfusion injury and post myocardial infarction), diseases associated with tissue destruction (e.g., of COPD, cardiobronchitis and 30 post myocardial infarction), diseases associated with tissue inflammation (e.g., irritable bowel syndrome, spastic colon and inflammatory bowel disease), inflammation and

scarring.

In addition to the myriad diseases discussed above, cellular and animal empirical data confirm that the 2,4-pyrimidinediamine compounds described herein are also useful for the treatment or prevention of autoimmune diseases, as well as the various symptoms 5 associated with such diseases. The types of autoimmune diseases that may be treated or prevented with the 2,4-pyrimidinediamine compounds generally include those disorders involving tissue injury that occurs as a result of a humoral and/or cell-mediated response to immunogens or antigens of endogenous and/or exogenous origin. Such diseases are frequently referred to as diseases involving the nonanaphylactic (*i.e.*, Type II, Type III 10 and/or Type IV) hypersensitivity reactions.

As discussed previously, Type I hypersensitivity reactions generally result from the release of pharmacologically active substances, such as histamine, from mast and/or basophil cells following contact with a specific exogenous antigen. As mentioned above, such Type I reactions play a role in numerous diseases, including allergic asthma, allergic 15 rhinitis, etc.

Type II hypersensitivity reactions (also referred to as cytotoxic, cytolytic complement-dependent or cell-stimulating hypersensitivity reactions) result when immunoglobulins react with antigenic components of cells or tissue, or with an antigen or hapten that has become intimately coupled to cells or tissue. Diseases that are commonly 20 associated with Type II hypersensitivity reactions include, but are not limited, to autoimmune hemolytic anemia, erythroblastosis fetalis and Goodpasture's disease.

Type III hypersensitivity reactions, (also referred to as toxic complex, soluble complex, or immune complex hypersensitivity reactions) result from the deposition of soluble circulating antigen-immunoglobulin complexes in vessels or in tissues, with 25 accompanying acute inflammatory reactions at the site of immune complex deposition. Non-limiting examples of prototypical Type III reaction diseases include the Arthus reaction, rheumatoid arthritis, serum sickness, systemic lupus erythematosus, certain types of glomerulonephritis, multiple sclerosis and bullous pemphigoid.

Type IV hypersensitivity reactions (frequently called cellular, cell-mediated, 30 delayed, or tuberculin-type hypersensitivity reactions) are caused by sensitized T-

lymphocytes which result from contact with a specific antigen. Non-limiting examples of diseases cited as involving Type IV reactions are contact dermatitis and allograft rejection.

Autoimmune diseases associated with any of the above nonanaphylactic hypersensitivity reactions may be treated or prevented with the 2,4-pyrimidinediamine compounds of the invention. In particular, the methods may be used to treat or prevent those autoimmune diseases frequently characterized as single organ or single cell-type autoimmune disorders including, but not limited to: Hashimoto's thyroiditis, autoimmune hemolytic anemia, autoimmune atrophic gastritis of pernicious anemia, autoimmune encephalomyelitis, autoimmune orchitis, Goodpasture's disease, autoimmune thrombocytopenia, sympathetic ophthalmia, myasthenia gravis, Graves' disease, primary biliary cirrhosis, chronic aggressive hepatitis, ulcerative colitis and membranous glomerulopathy, as well as those autoimmune diseases frequently characterized as involving systemic autoimmune disorder, which include but are not limited to: systemic lupus erythematosus, rheumatoid arthritis, Sjogren's syndrome, Reiter's syndrome, polymyositis-dermatomyositis, systemic sclerosis, polyarteritis nodosa, multiple sclerosis and bullous pemphigoid.

It will be appreciated by skilled artisans that many of the above-listed autoimmune diseases are associated with severe symptoms, the amelioration of which provides significant therapeutic benefit even in instances where the underlying autoimmune disease may not be ameliorated. Many of these symptoms, as well as their underlying disease states, result as a consequence of activating the Fc γ R signaling cascade in monocyte cells. As the 2,4-pyrimidinediamine compounds described herein are potent inhibitors of such Fc γ R signaling in monocytes and other cells, the methods find use in the treatment and/or prevention of myriad adverse symptoms associated with the above-listed autoimmune diseases.

As a specific example, rheumatoid arthritis (RA) typically results in swelling, pain, loss of motion and tenderness of target joints throughout the body. RA is characterized by chronically inflamed synovium that is densely crowded with lymphocytes. The synovial membrane, which is typically one cell layer thick, becomes intensely cellular and assumes a form similar to lymphoid tissue, including dendritic cells, T-, B- and NK cells, macrophages and clusters of plasma cells. This process, as well as a plethora of immunopathological

mechanisms including the formation of antigen-immunoglobulin complexes, eventually result in destruction of the integrity of the joint, resulting in deformity, permanent loss of function and/or bone erosion at or near the joint. The methods may be used to treat or ameliorate any one, several or all of these symptoms of RA. Thus, in the context of RA, the 5 methods are considered to provide therapeutic benefit (discussed more generally, *infra*) when a reduction or amelioration of any of the symptoms commonly associated with RA is achieved, regardless of whether the treatment results in a concomitant treatment of the underlying RA and/or a reduction in the amount of circulating rheumatoid factor (“RF”).

As another specific example, systemic lupus erythematosus (“SLE”) is typically 10 associated with symptoms such as fever, joint pain (arthralgias), arthritis, and serositis (pleurisy or pericarditis). In the context of SLE, the methods are considered to provide therapeutic benefit when a reduction or amelioration of any of the symptoms commonly associated with SLE are achieved, regardless of whether the treatment results in a concomitant treatment of the underlying SLE.

15 As another specific example, multiple sclerosis (“MS”) cripples the patient by disturbing visual acuity; stimulating double vision; disturbing motor functions affecting walking and use of the hands; producing bowel and bladder incontinence; spasticity; and sensory deficits (touch, pain and temperature sensitivity). In the context of MS, the methods are considered to provide therapeutic benefit when an improvement or a reduction 20 in the progression of any one or more of the crippling effects commonly associated with MS is achieved, regardless of whether the treatment results in a concomitant treatment of the underlying MS.

When used to treat or prevent such diseases, the active compounds may be 25 administered singly, as mixtures of one or more active compounds or in mixture or combination with other agents useful for treating such diseases and/or the symptoms associated with such diseases. The active compounds may also be administered in mixture or in combination with agents useful to treat other disorders or maladies, such as steroids, membrane stabilizers, 5LO inhibitors, leukotriene synthesis and receptor inhibitors, inhibitors of IgE isotype switching or IgE synthesis, IgG isotype switching or IgG synthesis, 30 β-agonists, tryptase inhibitors, aspirin, COX inhibitors, methotrexate, anti-TNF drugs, Rituximab, PD4 inhibitors, p38 inhibitors, PDE4 inhibitors, and antihistamines, to name a

few. The active compounds may be administered *per se* in the form of prodrugs or as pharmaceutical compositions, comprising an active compound or prodrug.

Pharmaceutical compositions comprising the active compounds of the invention (or prodrugs thereof) may be manufactured by means of conventional mixing, dissolving, 5 granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilization processes. The compositions may be formulated in conventional manner using one or more physiologically acceptable carriers, diluents, excipients or auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically.

10 The active compound or prodrug may be formulated in the pharmaceutical compositions *per se*, or in the form of a hydrate, solvate, N-oxide or pharmaceutically acceptable salt, as previously described. Typically, such salts are more soluble in aqueous solutions than the corresponding free acids and bases, but salts having lower solubility than the corresponding free acids and bases may also be formed.

15 Pharmaceutical compositions of the invention may take a form suitable for virtually any mode of administration, including, for example, topical, ocular, oral, buccal, systemic, nasal, injection, transdermal, rectal, vaginal, etc., or a form suitable for administration by inhalation or insufflation.

20 For topical administration, the active compound(s) or prodrug(s) may be formulated as solutions, gels, ointments, creams, suspensions, etc. as are well-known in the art.

Systemic formulations include those designed for administration by injection, *e.g.*, subcutaneous, intravenous, intramuscular, intrathecal or intraperitoneal injection, as well as those designed for transdermal, transmucosal oral or pulmonary administration.

25 Useful injectable preparations include sterile suspensions, solutions or emulsions of the active compound(s) in aqueous or oily vehicles. The compositions may also contain formulating agents, such as suspending, stabilizing and/or dispersing agent. The formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or in multidose containers, and may contain added preservatives.

Alternatively, the injectable formulation may be provided in powder form for

reconstitution with a suitable vehicle, including but not limited to sterile pyrogen free water, buffer, dextrose solution, etc., before use. To this end, the active compound(s) may be dried by any art-known technique, such as lyophilization, and reconstituted prior to use.

5 For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are known in the art.

For oral administration, the pharmaceutical compositions may take the form of, for example, lozenges, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, 10 microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulfate). The tablets may be coated by methods well known in the art with, for example, sugars, films or enteric coatings.

15 Liquid preparations for oral administration may take the form of, for example, elixirs, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily 20 esters, ethyl alcohol, cremophoreTM or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, preservatives, flavoring, coloring and sweetening agents as appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound or prodrug, as is well known.

25 For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For rectal and vaginal routes of administration, the active compound(s) may be formulated as solutions (for retention enemas) suppositories or ointments containing conventional suppository bases such as cocoa butter or other glycerides.

For nasal administration or administration by inhalation or insufflation, the active compound(s) or prodrug(s) can be conveniently delivered in the form of an aerosol spray from pressurized packs or a nebulizer with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, fluorocarbons, 5 carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges for use in an inhaler or insufflator (for example capsules and cartridges comprised of gelatin) may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

10 A specific example of an aqueous suspension formulation suitable for nasal administration using commercially-available nasal spray devices includes the following ingredients: active compound or prodrug (0.5-20 mg/ml); benzalkonium chloride (0.1-0.2 mg/mL); polysorbate 80 (TWEEN® 80; 0.5-5 mg/ml); carboxymethylcellulose sodium or microcrystalline cellulose (1-15 mg/ml); phenylethanol (1-4 mg/ml); and dextrose (20-50 15 mg/ml). The pH of the final suspension can be adjusted to range from about pH5 to pH7, with a pH of about pH 5.5 being typical.

20 Another specific example of an aqueous suspension suitable for administration of the compounds *via* inhalation, and in particular for such administration of a compound of the invention, contains 1-20 mg/mL of the compound or prodrug, 0.1-1% (v/v) Polysorbate 80 (TWEEN®80), 50 mM citrate and/or 0.9% sodium chloride.

For ocular administration, the active compound(s) or prodrug(s) may be formulated as a solution, emulsion, suspension, etc. suitable for administration to the eye. A variety of vehicles suitable for administering compounds to the eye are known in the art. Specific non-limiting examples are described in U.S. Patent No. 6,261,547; U.S. Patent No. 25 6,197,934; U.S. Patent No. 6,056,950; U.S. Patent No. 5,800,807; U.S. Patent No. 5,776,445; U.S. Patent No. 5,698,219; U.S. Patent No. 5,521,222; U.S. Patent No. 5,403,841; U.S. Patent No. 5,077,033; U.S. Patent No. 4,882,150; and U.S. Patent No. 4,738,851.

30 For prolonged delivery, the active compound(s) or prodrug(s) can be formulated as a depot preparation for administration by implantation or intramuscular injection. The active

ingredient may be formulated with suitable polymeric or hydrophobic materials (e.g., as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, e.g., as a sparingly soluble salt. Alternatively, transdermal delivery systems manufactured as an adhesive disc or patch which slowly releases the active compound(s) for percutaneous absorption may be used. To this end, permeation enhancers may be used to facilitate transdermal penetration of the active compound(s). Suitable transdermal patches are described in for example, U.S. Patent No. 5,407,713.; U.S. Patent No. 5,352,456; U.S. Patent No. 5,332,213; U.S. Patent No. 5,336,168; U.S. Patent No. 5,290,561; U.S. Patent No. 5,254,346; U.S. Patent No. 5,164,189; U.S. Patent No. 5,163,899; U.S. Patent No. 5,088,977; U.S. Patent No. 5,087,240; U.S. Patent No. 5,008,110; and U.S. Patent No. 4,921,475.

Alternatively, other pharmaceutical delivery systems may be employed. Liposomes and emulsions are well-known examples of delivery vehicles that may be used to deliver active compound(s) or prodrug(s). Certain organic solvents such as dimethylsulfoxide (DMSO) may also be employed, although usually at the cost of greater toxicity.

The pharmaceutical compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active compound(s). The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

20 **Effective Dosages**

The active compound(s) or prodrug(s) of the invention, or compositions thereof, will generally be used in an amount effective to achieve the intended result, for example in an amount effective to treat or prevent the particular disease being treated. The compound(s) may be administered therapeutically to achieve therapeutic benefit or prophylactically to achieve prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated and/or eradication or amelioration of one or more of the symptoms associated with the underlying disorder such that the patient reports an improvement in feeling or condition, notwithstanding that the patient may still be afflicted with the underlying disorder. For example, administration of a compound to a patient suffering from an allergy provides therapeutic benefit not only when the underlying allergic

response is eradicated or ameliorated, but also when the patient reports a decrease in the severity or duration of the symptoms associated with the allergy following exposure to the allergen. As another example, therapeutic benefit in the context of asthma includes an improvement in respiration following the onset of an asthmatic attack, or a reduction in the 5 frequency or severity of asthmatic episodes. Therapeutic benefit also includes halting or slowing the progression of the disease, regardless of whether improvement is realized.

For prophylactic administration, the compound may be administered to a patient at risk of developing one of the previously described diseases. For example, if it is unknown whether a patient is allergic to a particular drug, the compound may be administered prior to 10 administration of the drug to avoid or ameliorate an allergic response to the drug.

Alternatively, prophylactic administration may be applied to avoid the onset of symptoms in a patient diagnosed with the underlying disorder. For example, a compound may be administered to an allergy sufferer prior to expected exposure to the allergen. Compounds may also be administered prophylactically to healthy individuals who are repeatedly 15 exposed to agents known to one of the above-described maladies to prevent the onset of the disorder. For example, a compound may be administered to a healthy individual who is repeatedly exposed to an allergen known to induce allergies, such as latex, in an effort to prevent the individual from developing an allergy. Alternatively, a compound may be administered to a patient suffering from asthma prior to partaking in activities which trigger 20 asthma attacks to lessen the severity of, or avoid altogether, an asthmatic episode.

The amount of compound administered will depend upon a variety of factors, including, for example, the particular indication being treated, the mode of administration, whether the desired benefit is prophylactic or therapeutic, the severity of the indication being treated and the age and weight of the patient, the bioavailability of the particular 25 active compound, etc. Determination of an effective dosage is well within the capabilities of those skilled in the art.

Effective dosages may be estimated initially from *in vitro* assays. For example, an initial dosage for use in animals may be formulated to achieve a circulating blood or serum concentration of active compound that is at or above an IC₅₀ of the particular compound as 30 measured in an *in vitro* assay, such as the *in vitro* CHMC or BMMC and other *in vitro* assays described in the Examples section. Calculating dosages to achieve such circulating

blood or serum concentrations taking into account the bioavailability of the particular compound is well within the capabilities of skilled artisans. For guidance, the reader is referred to Fingl & Woodbury, "General Principles," *In: Goodman and Gilman's The Pharmaceutical Basis of Therapeutics*, Chapter 1, pp. 1-46, latest edition, Pergamon Press, and the references cited therein.

5 Initial dosages can also be estimated from *in vivo* data, such as animal models. Animal models useful for testing the efficacy of compounds to treat or prevent the various diseases described above are well-known in the art. Suitable animal models of hypersensitivity or allergic reactions are described in Foster, 1995, *Allergy* 50(21Suppl):6-9, discussion 34-38 and Tumas *et al.*, 2001, *J. Allergy Clin. Immunol.* 107(6):1025-1033. Suitable animal models of allergic rhinitis are described in Szelenyi *et al.*, 2000, *Arzneimittelforschung* 50(11):1037-42; Kawaguchi *et al.*, 1994, *Clin. Exp. Allergy* 24(3):238-244 and Sugimoto *et al.*, 2000, *Immunopharmacology* 48(1):1-7. Suitable animal models of allergic conjunctivitis are described in Carreras *et al.*, 1993, *Br. J. Ophthalmol.* 77(8):509-514; Saiga *et al.*, 1992, *Ophthalmic Res.* 24(1):45-50; and Kunert *et al.*, 2001, *Invest. Ophthalmol. Vis. Sci.* 42(11):2483-2489. Suitable animal models of systemic mastocytosis are described in O'Keefe *et al.*, 1987, *J. Vet. Intern. Med.* 1(2):75-80 and Bean-Knudsen *et al.*, 1989, *Vet. Pathol.* 26(1):90-92. Suitable animal models of hyper IgE syndrome are described in Claman *et al.*, 1990, *Clin. Immunol. Immunopathol.* 56(1):46-53. 10 Suitable animal models of B-cell lymphoma are described in Hough *et al.*, 1998, *Proc. Natl. Acad. Sci. USA* 95:13853-13858 and Hakim *et al.*, 1996, *J. Immunol.* 157(12):5503-5511. Suitable animal models of atopic disorders such as atopic dermatitis, atopic eczema and atopic asthma are described in Chan *et al.*, 2001, *J. Invest. Dermatol.* 117(4):977-983 and Suto *et al.*, 1999, *Int. Arch. Allergy Immunol.* 120(Suppl 1):70-75. Ordinarily skilled 15 artisans can routinely adapt such information to determine dosages suitable for human administration. Additional suitable animal models are described in the Examples section.

20 Dosage amounts will typically be in the range of from about 0.0001 or 0.001 or 0.01 mg/kg/day to about 100 mg/kg/day, but may be higher or lower, depending upon, among other factors, the activity of the compound, its bioavailability, the mode of administration 25 and various factors discussed above. Dosage amount and interval may be adjusted individually to provide plasma levels of the compound(s) which are sufficient to maintain

therapeutic or prophylactic effect. For example, the compounds may be administered once per week, several times per week (e.g., every other day), once per day or multiple times per day, depending upon, among other things, the mode of administration, the specific indication being treated and the judgment of the prescribing physician. In cases of local 5 administration or selective uptake, such as local topical administration, the effective local concentration of active compound(s) may not be related to plasma concentration. Skilled artisans will be able to optimize effective local dosages without undue experimentation.


Preferably, the compound(s) will provide therapeutic or prophylactic benefit without causing substantial toxicity. Toxicity of the compound(s) may be determined using 10 standard pharmaceutical procedures. The dose ratio between toxic and therapeutic (or prophylactic) effect is the therapeutic index. Compounds(s) that exhibit high therapeutic indices are preferred.

The foregoing and other aspects of the present invention may be better understood in connection with the following representative examples.

15

EXAMPLES

Example 1. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine




A mixture of 40 mg of 2-Chloro-5-fluoro-N4-[3-oxo-benzo[1,4]thiazin-6-yl]-4-pyrimidineamine and 48 mg of 3-Hydroxy-4,5-dimethoxyaniline Hydrochloride salt in 700 ²⁰ uL EtOH was heated in the microwave at 180 °C for 1 hour. The precipitate formed was collected by suction filtration, dried, suspended in deionized water and the pH was adjusted to pH 5 with dilute sodium bicarbonate solution, brine was added and after the suspension was briefly sonicated the solid was collected by suction filtration and dried to yield 25 mg ²⁵ 43 % yield of the desired product 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine. ¹H NMR (DMSO-d6): δ 8.06 (d, 1H, J = 2.7 Hz), 7.58 (s, 1H), 7.40 (dd, 1H, J = 8.4 Hz, J = 1.5 Hz), 7.19 (s, 1H), 6.92 (s, 1H),

6.76 (d, 1H, $J = 1.5$ Hz), 3.60 (s, 3H), 3.58 (s, 3H), 3.43 (s, 2H); purity 92 %; MS (m/e): 444 (MH^+).

Examples 2-13 were prepared according to the procedure in Example 1.

Example 2. N4-[2,2-Dimethyl-3-oxo-benzo[1,4]thiazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine


5



^1H NMR (DMSO-d6): δ 8.07 (d, 1H, $J = 3.3$ Hz), 7.67 (s, 1H), 7.46 (dd, 1H, $J = 9$ Hz, $J = 2.1$ Hz), 7.18 (d, 1H, $J = 9$ Hz), 6.92 (s, 1H), 6.78 (d, 1H, $J = 2.1$ Hz), 3.62 (s, 3H), 3.58 (s, 3H), 1.35 (s, 6H); purity 94 %; MS (m/e): 472 (MH^+).

10


Example 3. N4-[2,2-Dimethyl-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine



^1H NMR (DMSO-d6): δ 8.03 (d, 1H, $J = 3.9$ Hz), 7.38 (dd, 1H, $J = 9$ Hz, $J = 2.1$ Hz), 7.29 (d, 1H, $J = 1.8$ Hz), 6.81 (m, 2H), 6.77 (d, 1H, $J = 2.1$ Hz), 3.59 (s, 3H), 3.58 (s, 3H), 1.38 (s, 6H); purity 95 %; MS (m/e): 455 (MH^+).

Example 4. N4-[2,2-Dimethyl-3-oxo-pyrid[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine


20



¹H NMR (DMSO-d6): δ 8.09 (d, 1H, J = 3.6 Hz), 7.72 (d, 1H, J = 8.1 Hz), 7.31 (d, 1H, J = 8.1 Hz), 6.91 (d, 1H, J = 2.7 Hz), 6.75 (d, 1H, J = 2.7 Hz), 3.63 (s, 3H), 3.58 (s, 3H), 1.41 (s, 6H); purity 96 %; MS (m/e): 457 (MH^+).


5

Example 5. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine




¹H NMR (DMSO-d6): δ 8.02 (d, 1H, J = 3.9 Hz), 7.34 (dd, 1H, J = 9 Hz, J = 2.1 Hz), 7.25 (d, 1H, J = 1.8 Hz), 6.84 (m, 2H), 6.78 (d, 1H, J = 2.1 Hz), 4.52 (s, 2H), 3.57 (s, 3H), 3.56 (s, 3H); purity 97 %; MS (m/e): 428 (MH^+).

10
15
20
Example 6. N4-[2,2-Difluoro-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine




¹H NMR (DMSO-d6): δ 8.15 (d, 1H, J = 3.6 Hz), 7.68 (dd, 1H, J = 9.3 Hz, J = 2.1 Hz), 7.58 (d, 1H, J = 2.1 Hz), 7.25 (d, 1H, J = 9.3 Hz), 6.93 (d, 1H, J = 2.1 Hz), 6.84 (d, 1H, J = 2.7 Hz), 3.64 (s, 3H), 3.63 (s, 3H); purity 95 %; MS (m/e): 464 (MH^+).

Example 7. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-4-(2-pyridylmethyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine




¹H NMR (DMSO-d6): δ 8.47 (δ , 1H, ϑ = 3.9 H ζ), 8.07 (d, 1H, J = 4.2 Hz), 7.70 (m, 1H), 7.38 (m, 2H), 7.26 (m, 2H), 6.96 (d, 1H, J = 8.7 Hz), 6.68 (m, 2H), 5.08 (s, 2H), 4.76 (s, 2H), 3.61 (s, 3H), 3.59 (s, 3H); purity 95 %; MS (m/e): 519 (MH $^+$).

5 **Example 8. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(1,3-(2H)-4,4-dimethylisoquinolinedione-7-yl)-2,4-pyrimidinediamine**




¹H NMR (DMSO-d6): δ 8.36 (m, 1H), 8.10 (m, 2H), 7.56 (d, 1H, J = 9.0 Hz), 6.86 (d, 1H, J = 2.4 Hz), 6.745 (d, 1H, J = 2.7 Hz), 3.58 (s, 3H), 3.54 (s, 3H), 1.51 (s, 6H); purity 10 93 %; MS (m/e): 468 (MH $^+$).

Example 9. (R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine




15 ¹H NMR (DMSO-d6): δ 8.02 (d, 1H, J = 3.2 Hz), 7.44 (m, 2H), 7.06 (m, 2H), 6.92 (m, 2H), 6.76 (m, 3H), 4.86 (s, 2H), 4.79 (q, 1H, J = 7.2 Hz), 3.64 (s, 3H), 3.63 (s, 3H), 3.54 (s, 3H), 1.47 (d, 3H, J = 7.2 Hz); purity 92 %; MS (m/e): 562 (MH $^+$).

Example 10. (R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine



¹H NMR (DMSO-d6): δ 8.06 (d, 1H, J = 3.3 Hz), 7.59 (m, 2H), 7.26 (d, 1H, J = 10.8 Hz), 6.99 (m, 2H), 6.76 (m, 4H), 4.94 (s, 2H), 3.75 (q, 1H, J = 7.2 Hz), 3.63 (s, 3H), 3.58 (s, 3H), 3.55 (s, 3H), 1.36 (d, 3H, J = 7.2 Hz); purity 90 %; MS (m/e): 578 (MH^+).

Example 11. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(2,2,4-trimethyl-1,1,3-trioxo-benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine




10

¹H NMR (DMSO-d6): δ 8.22 (d, 1H, 3.3 Hz), 8.04 (d, 1H, J = 2.1 Hz), 7.92 (dd, 1H, J = 8.4 Hz, J = 2.1 Hz), 7.52 (d, 1H, J = 8.4 Hz), 6.85 (d, 1H, J = 2.1 Hz), 6.71 (d, 1H, J = 2.1 Hz), 3.59 (s, 3H), 3.58 (s, 3H), 3.25 (s, 3H), 1.41 (s, 6H); purity 98 %; MS (m/e): 518 (MH^+).

15


Example 12. 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(4-methyl-3-oxo-benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine



¹H NMR (DMSO-d6): δ 8.09 (d, 1H, 3.6 Hz), 7.71 (d, 1H, J = 1.8 Hz), 7.67 (dd, 1H, J = 8.1 Hz, J = 2.1 Hz), 7.28 (d, 1H, J = 8.1 Hz), 6.83 (d, 1H, J = 1.8 Hz), 6.75 (d, 1H, J

= 2.1 Hz), 3.58 (s, 3H), 3.57 (s, 3H), 3.47 (s, 2H), 3.18 (s, 3H); purity 95 %; MS (m/e): 458 (MH⁺).

5 **Example 13. N4-(3,4-Dihydro-2H-2,2-dimethyl-5-pyrido[1,4]oxazin-6-yl)-N2-[3,4-dimethoxyphenyl-5-hydroxyphenyl]-5-fluoro-2,4-pyrimidinediamine**



10 ¹H NMR (DMSO-d6): δ 8.92 (s, 1H), 8.89 (s, 1H), 8.64 (s, 1H), 8.02 (d, 1H, J = 3.5 Hz), 7.33 (d, 1H, J = 8.2 Hz), 6.92 (d, 1H, J = 2.3 Hz), 6.88 (d, 1H, J = 8.2 Hz), 6.78 (d, 1H, J = 8.2 Hz), 6.56 (s, 1H), 3.63 (s, 3H), 3.59 (s, 3H), 3.12 (d, 1H, J = 2.3 Hz), 1.24 (s, 6H). LCMS: ret. time: 9.36 min.; purity: 97%; MS (m/e): 443 (MH⁺).

Inhibition of FcεRI Receptor-Mediated Degranulation

15 The ability of the 2,4-pyrimidinediamine compounds of the invention to inhibit IgE-induced degranulation is demonstrated in a variety of cellular assays with cultured human mast cells (CHMC) and/or mouse bone marrow derived cells (BMMC). Inhibition of degranulation is measured at both low and high cell density by quantifying the release of the granule specific factors tryptase, histamine and hexosaminidase. Inhibition of release and/or synthesis of lipid mediators is assessed by measuring the release of leukotriene LTC4 and inhibition of release and/or synthesis of cytokines is monitored by quantifying TNF-α, IL-6 and IL-13. Tryptase and hexosaminidase are quantified using fluorogenic substrates as 20 described in their respective examples. Histamine, TNFα, IL-6, IL-13 and LTC4 are quantified using the following commercial ELISA kits: histamine (Immunotech #2015, Beckman Coulter), TNFα (Biosource #KHC3011), IL-6 (Biosource #KMC0061), IL-13 (Biosource #KHC0132) and LTC4 (Cayman Chemical #520211). The protocols of the various assays are provided below.

25 **Culturing of Human Mast and Basophil Cells**

Human mast and basophil cells are cultured from CD34-negative progenitor cells as described below (see also the methods described in copending U.S. application Serial No. 10/053,355, filed November 8, 2001, the disclosure of which is incorporated herein by

reference).

Preparation of STEMPRO-34 Complete Medium

To prepare STEMPRO-34 complete medium (“CM”), 250 mL STEMPRO-34TM serum free medium (“SFM”; GibcoBRL, Catalog No. 10640) was added to a filter flask. To 5 this was added 13 mL STEMPRO-34 Nutrient Supplement (“NS”; GibcoBRL, Catalog No. 10641) (prepared as described in more detail, below). The NS container was rinsed with approximately 10 mL SFM and the rinse added to the filter flask. Following addition of 5 mL L-glutamine (200 mM; Mediatech, Catalog No. MT 25-005-CI and 5 mL 100X penicillin/streptomycin (“pen-strep”; HyClone, Catalog No. SV30010), the volume was 10 brought to 500 mL with SFM and the solution was filtered.

The most variable aspect of preparing the CM is the method by which the NS is thawed and mixed prior to addition to the SFM. The NS should be thawed in a 37° C water bath and swirled, not vortexed or shaken, until it is completely in solution. While swirling, take note whether there are any lipids that are not yet in solution. If lipids are present and 15 the NS is not uniform in appearance, return it to the water bath and repeat the swirling process until it is uniform in appearance. Sometimes this component goes into solution immediately, sometimes after a couple of swirling cycles, and sometimes not at all. If, after a couple of hours, the NS is still not in solution, discard it and thaw a fresh unit. NS that appears non-uniform after thaw should not be used.

20 Expansion of CD34+ Cells

A starting population of CD34-positive (CD34+) cells of relatively small number (1-5 x 10⁶ cells) was expanded to a relatively large number of CD34-negative progenitor cells (about 2-4 x 10⁹ cells) using the culture media and methods described below. The CD34+ cells (from a single donor) were obtained from Allcells (Berkeley, CA). Because there is a 25 degree of variation in the quality and number of CD34+ cells that Allcells typically provides, the newly delivered cells were transferred to a 15 mL conical tube and brought up to 10 mL in CM prior to use.

On day 0, a cell count was performed on the viable (phase-bright) cells and the cells were spun at 1200 rpm to pellet. The cells were resuspended to a density of 275,000 30 cells/mL with CM containing 200 ng/mL recombinant human Stem Cell Factor (“SCF”;

Peprotech, Catalog No. 300-07) and 20 ng/mL human flt-3 ligand (Peprotech, Catalog No. 300-19) (“CM/SCF/flt-3 medium”). On about day 4 or 5, the density of the culture was checked by performing a cell count and the culture was diluted to a density of 275,000 cells/mL with fresh CM/SCF/flt-3 medium. On about day 7, the culture was transferred to a 5 sterile tube and a cell count was performed. The cells were spun at 1200 rpm and resuspended to a density of 275,000 cells/mL with fresh CM/SCF/flt-3 medium.

This cycle was repeated, starting from day 0, a total of 3-5 times over the expansion period.

When the culture is large and being maintained in multiple flasks and is to be 10 resuspended, the contents of all of the flasks are combined into a single container prior to performing a cell count. This ensures that an accurate cell count is achieved and provides for a degree of uniformity of treatment for the entire population. Each flask is checked separately for contamination under the microscope prior to combining to prevent contamination of the entire population.

15 Between days 17-24, the culture can begin to go into decline (*i.e.*, approximately 5-10% of the total number of cells die) and fail to expand as rapidly as before. The cells are then monitored on a daily basis during this time, as complete failure of the culture can take place in as little as 24 hours. Once the decline has begun, the cells are counted, spun down at 850 rpm for 15 minutes, and resuspended at a density of 350,000 cells/mL in 20 CM/SCF/flt-3 medium to induce one or two more divisions out of the culture. The cells are monitored daily to avoid failure of the culture.

When greater than 15% cell death is evident in the progenitor cell culture and some debris is present in the culture, the CD34-negative progenitor cells are ready to be differentiated.

25 **Differentiation of CD34-Negative Progenitor Cells into Mucosal Mast Cells**

A second phase is performed to convert the expanded CD34-negative progenitor cells into differentiated mucosal mast cells. These mucosal cultured human mast cells (“CHMC”) are derived from CD34+ cells isolated from umbilical cord blood and treated to form a proliferated population of CD34-negative progenitor cells, as described above. To 30 produce the CD34-negative progenitor cells, the resuspension cycle for the culture was the

same as that described above, except that the culture was seeded at a density of 425,000 cells/mL and 15% additional media was added on about day four or five without performing a cell count. Also, the cytokine composition of the medium was modified such that it contained SCF (200 ng/mL) and recombinant human IL-6 (200 ng/mL; Peprotech, Catalog 5 No. 200-06 reconstituted to 100 ug/mL in sterile 10 mM acetic acid) (“CM/SCF/IL-6 medium”).

Phases I and II together span approximately 5 weeks. Some death and debris in the culture is evident during weeks 1-3 and there is a period during weeks 2-5 during which a small percentage of the culture is no longer in suspension, but is instead attached to the 10 surface of the culture vessel.

As during Phase I, when the culture is to be resuspended on day seven of each cycle, the contents of all flasks are combined into a single container prior to performing a cell count to ensure uniformity of the entire population. Each flask is checked separately for contamination under the microscope prior to combining to prevent contamination of the 15 entire population.

When the flasks are combined, approximately 75% of the volume is transferred to the communal container, leaving behind about 10 mL or so in the flask. The flask containing the remaining volume was rapped sharply and laterally to dislodge the attached cells. The rapping was repeated at a right angle to the first rap to completely dislodge the 20 cells.

The flask was leaned at a 45 degree angle for a couple of minutes before the remaining volume was transferred to the counting vessel. The cells were spun at 950 rpm for 15 min prior to seeding at 35-50 mL per flask (at a density of 425,000 cells/mL).

Differentiation of CD34-Negative Progenitor Cells into Connective Tissue-Type Mast Cells

A proliferated population of CD34-negative progenitor cells is prepared as above and treated to form a tryptase/chymase positive (connective tissue) phenotype. The methods are performed as described above for mucosal mast cells, but with the substitution of IL-4 for IL-6 in the culture medium. The cells obtained are typical of connective tissue 30 mast cells.

Differentiation of CD34-Negative Progenitor Cells into Basophil Cells

A proliferated population of CD34-negative progenitor cells is prepared as described above and used to form a proliferated population of basophil cells. The CD34-negative cells are treated as described for mucosal mast cells, but with the substitution of IL-3 (at 20-5 50 ng/mL) for IL-6 in the culture medium.

CHMC Low Cell Density IgE Activation: Tryptase and LTC4 Assays

To duplicate 96-well U-bottom plates (Costar 3799) add 65 ul of compound dilutions or control samples that have been prepared in MT [137 mM NaCl, 2.7 mM KCl, 1.8 mM CaCl₂, 1.0 mM MgCl₂, 5.6 mM Glucose, 20 mM Hepes (pH 7.4), 0.1% Bovine 10 Serum Albumin, (Sigma A4503)] containing 2% MeOH and 1% DMSO. Pellet CHMC cells (980 rpm, 10 min) and resuspend in pre-warmed MT. Add 65 ul of cells to each 96-well plate. Depending on the degranulation activity for each particular CHMC donor, load 1000-1500 cells/well. Mix four times followed by a 1 hr incubation at 37°C. During the 1 hr incubation, prepare 6X anti-IgE solution [rabbit anti-human IgE (1 mg/ml, Bethyl 15 Laboratories A80-109A) diluted 1:167 in MT buffer]. Stimulate cells by adding 25 ul of 6X anti-IgE solution to the appropriate plates. Add 25 ul MT to un-stimulated control wells. Mix twice following addition of the anti-IgE. Incubate at 37°C for 30 minutes. During the 30 minute incubation, dilute the 20 mM tryptase substrate stock solution [(Z-Ala-Lys-Arg- 20 AMC'2TFA; Enzyme Systems Products, #AMC-246)] 1:2000 in tryptase assay buffer [0.1 M Hepes (pH 7.5), 10 % w/v Glycerol, 10 uM Heparin (Sigma H-4898) 0.01% NaN₃]. Spin plates at 1000 rpm for 10 min to pellet cells. Transfer 25 ul of supernatant to a 96-well black bottom plate and add 100 ul of freshly diluted tryptase substrate solution to each well. Incubate plates at room temperature for 30 min. Read the optical density of the plates at 355nm/460nm on a spectrophotometric plate reader.

25 Leukotriene C4 (LTC4) is also quantified using an ELISA kit on appropriately diluted supernatant samples (determined empirically for each donor cell population so that the sample measurement falls within the standard curve) following the supplier's instructions.

CHMC High Cell Density IgE Activation: Degranulation (Tryptase, Histamine), Leukotriene (LTC4), and Cytokine (TNF α , IL-13) Assays

Cultured human mast cells (CHMC) are sensitized for 5 days with IL-4 (20 ng/ml), SCF (200 ng/ml), IL-6 (200 ng/ml), and Human IgE (CP 1035K from Cortx Biochem, 100-500ng/ml depending on generation) in CM medium. After sensitizing, cells are counted, pelleted (1000 rpm, 5-10 minutes), and resuspended at 1-2 x10⁶ cells/ml in MT buffer. Add 100 ul of cell suspension to each well and 100 ul of compound dilutions. The final vehicle concentration is 0.5% DMSO. Incubate at 37°C (5% CO₂) for 1 hour. After 1 hour of compound treatment, stimulate cells with 6X anti-IgE. Mix wells with the cells and allow 10 plates to incubate at 37°C (5% CO₂) for one hour. After 1 hour incubation, pellet cells (10 minutes, 1000 RPM) and collect 200 ul per well of the supernatant, being careful not to disturb pellet. Place the supernatant plate on ice. During the 7-hour step (see next) perform tryptase assay on supernatant that had been diluted 1:500. Resuspend cell pellet in 240 ul of CM media containing 0.5% DMSO and corresponding concentration of compound.

15 Incubate CHMC cells for 7 hours at 37°C (5% CO₂). After incubation, pellet cells (1000 RPM, 10 minutes) and collect 225 ul per well and place in -80°C until ready to perform ELISAS. ELISAS are performed on appropriately diluted samples (determined empirically for each donor cell population so that the sample measurement falls within the standard curve) following the supplier's instructions.

20 Inhibition of the Upstream IgE Receptor Cascade

Assays for ionomycin-induced mast cell degranulation are carried out as described for the CHMC Low Density IgE Activation assays, with the exception that during the 1 hour incubation, 6X ionomycin solution [5mM ionomycin (Sigma I-0634) in MeOH (stock) diluted 1:416.7 in MT buffer (2 μ M final)] was prepared and cells were stimulated by 25 adding 25 μ l of the 6X ionomycin solution to the appropriate plates.

Inhibition of Syk Kinase in Biochemical Assays

Compounds are tested for the ability to inhibit Syk kinase catalyzed phosphorylation of a peptide substrate in a biochemical fluorescenced polarization assay with isolated Syk kinase. In this experiment, compounds are diluted to 1% DMSO in kinase buffer (20 mM 30 HEPES, pH 7.4, 5 mM MgCl₂, 2 mM MnCl₂, 1 mM DTT, 0.1 mg/mL acetylated Bovine Gamma Globulin). Compound in 1% DMSO (0.2% DMSO final) is mixed with

ATP/substrate solution at room temperature. Syk kinase (Upstate, Lake Placid NY) is added to a final reaction volume of 20 uL, and the reaction was incubated for 30 minutes at room temperature. Final enzyme reaction conditions were 20 mM HEPES, pH 7.4, 5 mM MgCl₂, 2 mM MnCl₂, 1 mM DTT, 0.1 mg/mL acetylated Bovine Gamma Globulin, 0.125 ng Syk, 4 5 uM ATP, 2.5 uM peptide substrate (biotin-EQEDEPEGDYEEVLE-CONH₂, SynPep Corporation). EDTA (10 mM final)/anti-phosphotyrosine antibody (1X final)/fluorescent phosphopeptide tracer (0.5X final) is added in FP Dilution Buffer to stop the reaction for a total volume of 40 uL according to manufacturer's instructions (PanVera Corporation) The plate is incubated for 30 minutes in the dark at room temperature. Plates are read on a 10 Polarion fluorescence polarization plate reader (Tecan). Data is converted to amount of phosphopeptide present using a calibration curve generated by competition with the phosphopeptide competitor provided in the Tyrosine Kinase Assay Kit, Green (PanVera Corporation).

When tested in the LD Tryptase assay, the 3-hydroxyphenyl-2,4-pyrimidinediamine 15 compounds of Examples 1-13 were all found to have an activity of less than 5uM in the assay, as shown in Table 1 below where A indicates an activity of less than 1 uM and B indicates an activity of less than 5 uM. Moreover, the 2-hydroxyphenyl compounds that were tested relative to their 3,4,5-trimethoxyphenyl counterparts showed between about 10% and 500% improved potency.

Table 1.


| Compound Ex. No. | LD Tryptase, CHMC, IgE, 8pt |
|---------------------|--------------------------------|
| 1 | A |
| 2 | A |
| 3 | A |
| 4 | A |
| 5 | A |
| 6 | A |
| 7 | B |
| 8 | A |
| 9 | B |
| 10 | B |
| 11 | A |
| 12 | A |
| 13 | A |

Although the foregoing invention has been described in some detail to facilitate understanding, it will be apparent that certain changes and modifications may be practiced 5 within the scope of the appended claims. Accordingly, the described embodiments are to be considered as illustrative and not restrictive, and the invention is not to be limited to the details given herein, but may be modified within the scope and equivalents of the appended claims.

All literature and patent references cited throughout the application are incorporated 10 by reference into the application for all purposes.

What is claimed is:

1. A compound according to formula I:



I

or a stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof, wherein:

Y is selected from the group consisting of S, O, SO, SO₂, and C(R⁷)₂;

each R³⁵ is independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, and halo, or both R³⁵ together with the carbon to which they are attached form a carbonyl group;

W is selected from the group consisting of C=O, C=S, C=NH, C(R⁷)₂, and NR³⁷;

Z is C=O or NR³⁷, provided that Z and W are not both NR³⁷ and provided that when Z is C=O, then W is C(R⁷)₂ or NR³⁷;

X is CH or N;

each R³¹ is independently (C₁-C₄)alkyl or both R³¹ together form a (C₁-C₂)alkylene group optionally substituted with one to two (C₁-C₄)alkyl groups or substituted with one (C₃-C₇) spirocycloalkyl group;

each R⁷ is independently hydrogen or (C₁-C₄)alkyl; and

R³⁷ is hydrogen or methyl optionally substituted with phenyl or pyridyl, wherein said phenyl or pyridyl is optionally substituted with (C₁-C₄)alkoxy.

2. A compound according to claim 1 wherein Y is O or S.

3. A compound according to claim 2 wherein Y is O.

4. A compound according to claim 1 wherein both R³⁵ are the same.

5. A compound according to claim 4 wherein both R³⁵ are methyl.

25 6. A compound according to claim 1 wherein W is C=O or C=S.

7. A compound according to claim 6 wherein W is C=O.

8. A compound according to claim 1 wherein Z is NR³⁷.
9. A compound according to claim 8 wherein Z is NH.
10. A compound according to claim 1 wherein X is N.
11. A compound according to claim 1 wherein R³¹ is methyl.
- 5 12. A compound according to claim 1 wherein Z is NR³⁷ and R³⁷ is methyl, 2-pyridylmethyl, or 4-methoxybenzyl.
13. The compound of according to claim 1 selected from the group consisting of 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine;
- 10 N4-[2,2-Dimethyl-3-oxo-benzo[1,4]thiazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;
- 14 N4-[2,2-Dimethyl-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;
- 15 N4-[2,2-Dimethyl-3-oxo-pyrid[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;
- 16 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;
- 17 N4-[2,2-Difluoro-3-oxo-benz[1,4]oxazin-6-yl]-5-fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-2,4-pyrimidinediamine;
- 20 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[3-oxo-4-(2-pyridylmethyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;
- 21 N4-(3,4-Dihydro-2H-2,2-dimethyl-5-pyrido[1,4]oxazin-6-yl)-N2-[3,4-dimethoxyphenyl-5-hydroxyphenyl]-5-fluoro-2,4-pyrimidinediamine;
- 22 5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(1,3-(2H)-4,4-dimethylisoquinolinedione-7-yl)-2,4-pyrimidinediamine;
- 23 (R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benz[1,4]oxazin-6-yl]-2,4-pyrimidinediamine;
- 24 (R/S)-5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-[2-methyl-3-oxo-4-(4-methoxybenzyl)-benzo[1,4]thiazin-6-yl]-2,4-pyrimidinediamine;

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(2,2,4-trimethyl-1,1,3-trioxo-
benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine; and

5-Fluoro-N2-(3-hydroxy-4,5-dimethoxyphenyl)-N4-(4-methyl-3-oxo-
benzo[1,4]thiazin-6-yl)-2,4-pyrimidinediamine; or

5 a stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof.

14. A composition comprising a compound, stereoisomer, salt, hydrate, solvate, N-
oxide, or prodrug thereof according to claim 1 and a carrier, excipient, or diluent.

15. A method of inhibiting the IgE signaling cascade of a cell expressing an IgE
receptor, comprising contacting the cell with an amount of a compound, a stereoisomer, salt,
10 hydrate, solvate, N-oxide, or prodrug thereof according to claim 1 effective to inhibit the
signaling cascade.

16. A method of inhibiting an Fc receptor signal transduction cascade in a subject,
comprising administering to the subject a compound, a stereoisomer, salt, hydrate, solvate,
N-oxide, or prodrug thereof according to claim 1 effective to inhibit the Fc receptor signal
15 transduction cascade.

17. The method of claim 16 in which the Fc receptor is selected from Fc α RI, Fc γ RI,
Fc γ RIII and Fc ε RI.

18. A method of treating a disease characterized by mast or basophil cell degranulation,
comprising administering to a subject suffering from such a disease an amount of a
20 compound, a stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof according to
claim 1 effective to treat the disease.

19. A method according to claim 18, wherein the disease is a disease associated with
tissue inflammation.

20. A method according to claim 19, wherein the disease associated with tissue
25 inflammation is irritable bowel syndrome, spastic colon or inflammatory bowel disease),

21. A method according to claim 18, wherein the disease is mediated by Syk kinase.

22. A method of treating or preventing an autoimmune disease in a subject, and/or one
or more symptoms associated therewith, comprising administering to the subject an amount

of a subject a compound, a stereoisomer, salt, hydrate, solvate, N-oxide, or prodrug thereof according to claim 1 effective to treat or prevent the autoimmune disease.

23. The method of claim 22 in which the autoimmune disease is selected from autoimmune diseases that are frequently designated as single organ or single cell-type

5 autoimmune disorders and autoimmune disease that are frequently designated as involving systemic autoimmune disorder.

24. Use of a compound of any one of claims 1-13 for the manufacture of a medicament.